

Fig. Illustration of the hot smoke airflow in the smoking chamber

Cold smoking

Fish can also be subjected to cold smoking. Temperatures of cold smoking typically do not exceed 30 °C. Thus, cold smoked products are not cooked and typically heavily salted. Compared to the traditional hot smoking, cold smoking runs longer, has a higher yield and retains the original textural properties much better than the hot-smoked ones. Cold smoking of varied fish species has been reported, including rainbow trout.

Liquid smoking

Liquid smoke is smoke condensate that is dissolved in a solvent, such as water or oil (Maga, 1988). Liquid smoke can be used directly on products by dipping or spraying. It is rapid and much easier to achieve a uniform smoke flavour than traditional cold and hot smoking processes, although the flavour and colour from the traditional smoking cannot be exactly duplicated (Varlet et al., 2007). Some potential harmful ingredients (e.g. polycyclic aromatic hydrocarbons, PAHs) in the nature smoke can be separated out and excluded from the liquid smoke (Chen & Lin, 1997). Other advantages of liquid smoke include easy modification, application to food items that traditionally are not smoked, lower operation cost, and less environmental pollution (Abu-Ali & Barringer,

2007). However, the application of liquid smoking may be expensive compared to other methods. Liquid smoking of fish species had been reported on swordfish, salmon and rainbow trout.

Electrostatic smoking

Electrostatic smoking is another rapid way to smoke. In the electrostatic smoking, fish are sent into a tunnel where an electrostatic field is created. Smoke particles are given a positive charge and deposit onto the surface of the fish which are negative charged. Although this procedure will change the composition of the smoke, the efficiency of smoking is still higher than that of the traditional smoking. It can also be operated continuously. The smoke compound ratio in the vapour phase may be modified by the electrostatic field, which results in increased level of carbonyl compounds (Ruiter, 1979). Factors that may influence the electrostatic smoking operation include the skin thickness, presence of scales, and subcutaneous fat amount (Maga, 1988). This operation may present safety problems to employees. Applications of electrostatic smoking have been reported mainly in salmon and herring.




Fig. Schematic diagram of smoke house with basic components.

Hot smoking of fish

Good smoked products can only be obtained from good raw material (Dore, 1993). In addition, control of the smoking procedures plays an equal importance in the production of good products. From raw material preparation to final product storage, smoking includes several operations, such as brining, drying, smoking, packaging and storage.

Brining

This is the stage when the flavours and spices are introduced into the fish. Cleaned fish are submerged under a prepared brine solution for a certain amount of time. A brine time less than 12 hours at 3.3 °C (38 °F) is recommended to minimize the possible spoilage in the fish (Lee, 1977). Salt is an important ingredient to be delivered into the fish tissue at this stage as well as a key hazard analysis and critical control point (HACCP) preventive measure for smoked fish. Not only does it bring the taste but also reduces the water activity (a_w) in the product, so that bacterial growth can be inhibited in the smoked fish.

Of all the bacteria that can exist in fish products, *Clostridium botulinum* is a major concern for vacuum or reduced packaged fish products. *C. botulinum* is a strictly anaerobic, gram positive bacillus bacterium. The vegetative cells and their neurotoxins can be easily destroyed by heat (less than five minutes) at 85 °C. However, their spores are very resistant to heat and can survive for up to 2 hours at 100 °C (Caya, 2001). Thus, prevention of botulism from hot-smoked fish products depends on the destruction of all *C. botulinum* spores or inhibition germination of the spores that may be present in the products.

Water phase salt (WPS) is used to measure the amount of salt in the fish products.

The WPS is calculated as (FDA, 2001):

$$WPS = \frac{\%Salt}{\%Salt + \%Moisture} \times 100$$

The higher the WPS value, the less the availability of the water. When sodium chloride is the only major humectant in the cured food, the relationship between the a_w and WPS can be expressed as (Ross & Dalgaard, 2004):

$$a_w = 1 - 0.0052471 \cdot WPS\% - 0.00012206 \cdot (WPS\%)^2$$

or

$$WPS\% = 8 - 140.07 \cdot (a_w - 0.95) - 405.12 \cdot (a_w - 0.95)^2$$

Current regulations require at least 3.5% WPS in the loin muscle of the vacuum packaged smoke products; at least 3.0% WPS if at least an additional 100 ppm nitrite exists in the vacuum packaged product; air packaged smoked fish products must contain at least 2.5% WPS (FDA, 2001).

Several salting methods are available to deliver the salt into the fish. The most common techniques used by the industry are dry and brine salting. Dry salting is widely used in

low fat fish. Basically, fish are put into layers with dry salt separating each layer. Water removed by salt is allowed to drain away. Periodical reshuffling of the layers may be necessary to make sure all the fish get uniform salting and pressure. Muscle fiber shrinks more during dry salting than brine salting (Sigurgisladottir et al., 2000b). Thus, dry salting of fish typically results in over-dried fish and low yield. A better quality and higher yield is usually obtained from brine salting.

Fish are brine salted by completely being covered in a prepared brine solution for a certain time period. The brine solution can have a salt concentration from relatively low to saturated levels. Brine salting is also used widely for most fatty fish since oxygen cannot oxidize the fish fat easily. Some modern processors inject the brine to speed up the process, therefore lowering the cost and minimizing the chance of fish deterioration. Salt is distributed evenly in the fish when injection brine is used. A higher brine yield can be obtained through injection brine as compared to brine or dry salting. Flavour ingredients can also be incorporated into the injection solution. However, the injecting brine operation has to be carefully controlled to avoid contamination delivered by the needles into the previously sterile flesh. Brine salting is still one of the most widely used salting methods for smoked fish. Efficiency of salt penetration into the fish tissue is affected by several factors, such as species, physiological state of fish (rigor), fish quality (fresh/frozen) fish dimension (thickness), brine concentration, brine time, brine to fish ratio, brine temperature, fat content, texture, etc.

After brining, fish have to be rinsed with clean water to remove the brine solution on its surface because a harsh, salty flavour can develop due to residues of brine solution.

Drying

It is widely known that reducing the water activity (a_w) will result in a reduction of microbial activity. The a_w is defined as:

$$a_w = p / p_0$$

where p is the vapour pressure of the product, and p_0 is the vapour pressure of pure water at the same temperature (Olley, Doe, & Heruwati, 1989).

For ideal solutions (real solutions at low concentrations), water activity can be calculated from the formula:

$$a_w = n_1 / (n_1 + n_2)$$

where n_1 is the number of moles of solvent, and n_2 is the number of moles of the solute.

This relationship may become complex due to the interactions between moisture and the fish tissue and also the relatively high solute concentration involved in cured fish. Drying of the fish can still be simulated with the formula in a way that drying the fish will cause a decrease in n_1 and an increase in n_2 , which finally decreases the a_w .

A certain amount of moisture has to be lost from fish after brining; so that water activity (a_w) can be decreased and a good texture can be obtained at the end of the smoking process. Drying of fish occurs at the early stage of smoking process. An air flow is applied on the fish; so that moisture in the fish tissue can migrate to the surface and leave the fish by evaporation. The temperature, relative humidity and velocity of the air flow are keys to the rate of drying. Drying with a low relative humidity air at high velocity may not drive the moisture out of the fish fast. If the temperature is too high

fish surface may be hardened at the beginning of drying resulting in a blocking layer to the inside moisture migration. The hardened surface may also prevent smoke penetrating into the tissue, which decreases the preservative effects of the smoke. Tissues under the hardened surface will tend to spoil from inside.

Drying at temperatures below 70 to 80 °C was recommended to minimize the damage to protein quality in fish (Opstvedt, 1989). Drying also influences the quality of finished smoked fish product.

Smoking

Smoke is generated from the incomplete combustion of wood at certain temperatures followed by thermal disintegration or pyrolysis of high molecular organic compounds into volatile lower molecular mass (Eyo, 2001). Smoke is composed of two phases: a particulate or dispersed phase and a gaseous or dispersing phase. The major parts of dispersed phase are particles in the droplet form having an average diameter of 0.196 to 0.346 μm (Maga, 1988; Wheaton & Lawson, 1985). These particles are mainly tars, wood resins, and compounds with high or low boiling points. The dispersed phase is the visible part of the smoke. The dispersing phase is responsible for flavouring, colouring, antioxidative, and bacteriostatic roles of the smoke (Hall, 1997). The composition of the dispersing smoke phase is complicated, many of which have yet been identified. More than 200 components have been identified. The most abundant chemicals found in smoke are carbonyls, organic acids, phenols, alcohols, and hydrocarbons.

Quality and composition of the smoke are affected by several factors, such as combustion temperature, wood type, moisture content of wood, air ventilation rate, and wood size.

Cellulose, hemicellulose and lignin are three main components in wood and their contents and compositions vary in different types of wood. Cellulose levels are fairly consistent among different species. Softwoods have higher lignin content than hardwoods. Hardwoods typically contain more hemicellulose than softwoods. Decomposition of hemicellulose happens at the early stage of smoking and produces furan and its derivatives as well as aliphatic carboxylic acids, which drops the pH in the smoked product. Softwoods also contain more resin acids than hardwoods, which typically introduces unpleasant flavor to the fish. Hardwoods, such as hickory, oak, cherry, apple and beech, are preferred in most situations over the softwoods for smoke generation. This is because hardwoods tend to produce more phenols and organic acids which contribute to the flavor and preservation effect of smoking (Hall, 1997).

The amount of air present during the production of smoke also influences the results of wood pyrolysis. Lower temperature and less air produce a smoke with more flavoring and preserving substances. While a higher temperature and more air burn the woods into carbon dioxide and water. Smoke production can be influenced by the size of wood. Wood can be used as chunks, chips or sawdust forms. However, their combustion rates will vary if same ventilation rate is used. Sawdust produces more smoke than chunks or chips due to its self-smoldering effect, which blocks the access of oxygen. Fish is also more likely to be charred with less smoke when chunks or chips are used. Most modern smokers use continuously fed sawdust to maintain a consistent production of smoke.

Although people like the flavour and taste of the smoked product, there are concerns about the negative side of smoked products, which are mainly focused on the

carcinogenic substances found in the smoke: the polynuclear aromatic hydrocarbons (PAHs). PAHs are composed of multiple fused benzene rings. It can be thermally produced by either high temperature pyrolysis or from the incomplete combustion of materials containing carbon and hydrogen. Up to 100 PAHs compounds have been either identified or detected (Maga, 1988). The level of PAHs can be reduced by decreasing the combustion temperature since the PAHs content was found to change linearly from 5 to 20 $\mu\text{g}/100\text{g}$ in temperature range 400 to 1000 $^{\circ}\text{C}$ (Eyo, 2001). Indirect smoking like liquid and electrostatic smoking also significantly reduces the PAHs amount.




Fig. Smoking kiln

Potential hazards associated with smoking of fish

I. Biological hazards

Generally, Cold smoking will typically reduce the level of microorganism by 90 to 99%. But after the cold smoking there is no such steps to eliminate or reduce the level of microorganisms. Typical temperature used for cold smoking is 22-28 $^{\circ}\text{C}$. However, this temperature is not sufficient to eliminate the risk from *Listeria monocytogens*, a gram positive, facultative anaerobic, psychrotropic bacteria causing deadly septicaemia, meningitis, spontaneous abortion, and foetal death in adult human beings. Specific high risk categories like persons with altered immune system, pregnant ladies, old aged persons etc. will be more susceptible to listeriosis followed by accidental inclusion. Comparatively high temperature used in hot-smoking process and long-time of exposure to that temperature (60-70 $^{\circ}\text{C}$ for 2-3 h) can inactivate the *L. monocytogenes* effectively, provided the raw material is not extra-ordinarily contaminated with the bacteria prior to processing. At the same time listericidal process

should be validated to ensure that the treatments are effective and can be applied continuously. But the hot smoked products are susceptible to post-process contaminations from many of the micro-organisms due to improper handling and storage of the products. Sufficient heat treatment, proper hygienic handling and cold chain maintenance during distribution can reduce the risk of biological hazards in smoked fish and fishery products.

Another important biological hazard associated with storage of smoked fish is *Clostridium botulinum*. The toxin produced by *C. botulinum* can lead to botulism, serious illness and death to the consumer. Even a few micrograms of intoxication can lead to ill-health with symptoms like weakness, vertigo, double vision, difficulty in speaking, swallowing and breathing, abdominal swelling, constipation, paralysis and death. The symptoms will start within 18-36 h after consumption of the infected product. By achieving proper salt concentration in processed fish, proper refrigeration during storage and reduced oxygen packaging like Modified Atmosphere Packaging (MAP) and vacuum packaging of the products can prevent the occurrence of *C. botulinum* in smoked fish and fishery products, especially type E and non-proteolytic types B and F. Salt along with smoke effectively prevents the toxin formation from type E, B and F.

In cold smoked fish and fishery products, which undergoes mild heat processing, the presence of spoilage organisms prevents the growth of *C. botulinum* and toxin production. Whereas in hot-smoked products, high temperature application causes damages to spores of *C. botulinum* thus prevents the toxin formation. Same process also prevents the prevalence of spoilage organisms and thus extends the shelf life of the product. Thus, the time- temperature combination for smoking, along with salt concentration plays critical roles in safety and quality aspects of the smoked fish and fishery products.

II. Chemical hazards

1. Polycyclic Aromatic Hydrocarbons (PAHs)

PAHs are large class of organic compounds containing two or more fused aromatic rings made up of carbon and hydrogen atoms. Incomplete combustion (pyrolysis), during smoking can lead to formation and release of PAHs into the smoked product. Some of them are carcinogenic and mutagenic substances causing serious health issues to the consumers. Processing procedures such as smoking, drying, roasting, baking, frying and barbecuing/grilling can lead to formation of PAHs in food items. Many reports indicate that individual PAHs in smoked fish can go up to a level of 200 µg/Kg. Among the 33 PAHs evaluated by the scientific committee on Food (SCF, 2002) of EU, 15 were found to be having mutagenicity/Geno toxicity in somatic cells of experimental animal in-vivo. They are benzo[a]anthracene, benzo[b]-, benzo[j]- and benzo[k]fluoranthene, benzo[ghi]perylene, benzo[a]pyrene, chrysene, cyclopenta[cd]pyrene, dibenz[a,h]anthracene, dibenzo[a,e]-, dibenzo[a,h]-, dibenzo[a,i]-, dibenzo[a,l]pyrene, indeno[1,2,3-cd]pyrene and 5-methylchrysene. The carcinogenic and genotoxic potentials of PAH are largest among the high molecular weight PAH, i.e. compounds with 4 rings or more. Among that benzo[a]pyrene regarded as potentially genotoxic and carcinogenic to humans. They can cause long-term adverse health effects following dietary intake of PAH.

The PAH contamination in smoked products can be significantly reduced by using indirect smoking process instead of direct smoking of the fish. In indirect

smoking, the smoke generated in an external smoking kiln, under controlled conditions, is used for smoking process. The smoke produced can be even, washed before coming into contact with the food material processed. In addition to that, use of lean fish for smoking, and cooking at lower temperature for longer time can also reduce the PAH contamination significantly. If the smoke condensate is used for smoking, usage of smoke condensate from reputed reliable resources approved by competent authority can effectively reduce the occurrence of PAH contamination in the final product. The formation of PAH in smoked fish can be minimised by following Code of Practice for the Reduction of Contamination of Food with Polycyclic Hydrocarbons (PAH) from Smoking and Direct Drying Processes (CAC/RCP 68-2009) given by Codex Alimentarius Commission. EU No.835/2011 specifies that maximum level of benzopyrene, and PAH4 (benzo[a]pyrene + chrysene+ benz[a]anthracene+benzo[b]fluoranthene) should be 2µg/Kg wet weight and 12µg/Kg in meat of smoked fish and fishery products, 5µg/Kg and 30µg/Kg in smoked sprats and 6µg/Kg and 35µg/Kg in smoked bivalve mollusc respectively.

2. Histamine:

Histamine poisoning is associated with Scombroid fishes and other dark meat fishes. The fishes showing potential treats of histamine poisoning are tunas, bonitos, mackerel, mahimahi, carangids, herring etc. These fishes having high content of free histidine, which during spoilage are converted to histamine by bacteria like *Morganellamorgani*, *Klebsiellapnuemoniae* and *Hafniaalvei*. Histamine is heat stable, even cooking or canning cannot destroy it. Presence of other biogenic amines like cadaverine and putrescine will act as potentiators for histamine production. As per Codexstandards, the maximum allowable histamine content in smoked fishes is 200 mg/Kg for species like *Scombridae*, *Clupeidae*, *Engraulidae*, *Coryphaenidae*, *Pomatomidae*, and *Scomberesocidae*. Low temperature storage of fishes right from catch can effectively reduce the production of histamine in fishes.

3. Biotoxins:

Biotoxins causing a number of food borne diseases. The poisoning due to biotoxins are caused by consuming finfish/shell fish containing poisonous tissues with accumulated toxins from plankton they consumed. Paralytic shellfish poisoning (PSP), diarrhetic shellfish poisoning (DSP), amnesic shellfish poisoning (ASP), and neurotoxic shellfish poisoning (NSP) are mostly associated with shellfish species such as oysters, clam and mussels. The control of biotoxin is very difficult. They cannot be destroyed by any of the processing methods like cooking, smoking, drying or salting. Environmental monitoring of plankton and proper depuration process of the bivalves only can reduce the occurrence significantly.

III. Physical Hazards

Presence of parasites like nematodes, cestodes, trematodes and any other extraneous matter can be considered as physical hazards. Particular attention needs to be paid to cold smoked or smoke-flavoured products, which should be frozen before or after smoking if a parasite hazard is present.

IV. Other potential hazards associated with smoking of fish

If wood or plant material is using for smoking of fish, there is a chance of presence of natural toxins, chemicals, paint, or impregnating material in plant or wood used which may result in imparting undesirable odour in processed products. This can be prevented by using sufficiently dried wood or plant material for smoke generation, judicious selection of the species of wood or plant and not using woods having mould or fungus growth for smoking process. Moreover, the material for smoking should be kept in a clean dry place during storage to prevent any kind of contamination, till the usage.
