

Nutritional Changes during Production of High Temperature Short Time (HTST) Air Puffed Potato and Potato-Soy Snacks

A. Nath¹, P. K. Chattopadhyay², Manish Pandey³, R. P. Mishra¹

¹Project Directorate for Farming System Research, Meerut, UP, India

²IIT, Kharagpur, West Bengal, India

³ICAR, Umiam, Meghalaya, India

amitnath2005@gmail.com

Abstract

Two different potato based ready-to-eat (RTE) snacks were prepared using high temperature short time (HTST) air puffing followed by oven toasting and evaluated for its nutritional changes that took place during high temperature short time air puffing and oven toasting process. The nutritional compositions of initial samples before puffing and oven toasting and final products were measured in terms of moisture content (% db), protein (%), fat (%), ash (%), crude fiber (%), carbohydrates (%), ascorbic acid content (mg/100g), nitrogen solubility index (%), protein digestibility (%), trypsin inhibitor activity (%), calorific value (Kcal/100g) and protein energy ratio. Significant increases in protein (56.67%), ash (70.49%), crude fiber (27.2%) and protein energy ratio (63.20%) were recorded in RTE potato-soy snack as compared to RTE potato snack due to the addition of soy flour (10.26%) in the product.

Keywords

Ready-to-Eat; Potato; Soy Flour; Nutrition; High Temperature Short Time Air Puffing; Oven Toasting

Introduction

Consumers in the recent world are becoming increasingly aware of the relationship between diet and health. Thus, the demand for a balanced diet and nutrient composition of food products that addresses specific health benefits is growing steadily. Processing and cooking conditions cause changes in availability of different food nutrients. During production of ready-to-eat (RTE) foods, several changes in nutritional composition take place in processing steps and become very important for process formulation.

The dehydration process, which is generally more widely applied to potatoes than to fruit and vegetables, can be very destructive to L-AA with a loss of 75%

being reported (Mishkin, Saguy, & Karel, 1984). The effect of different combinations of puffing temperature, puffing time, moisture content and air velocity on ascorbic acid content of ready-to eat dehydrated puffed potato cubes was reported by Mukherjee and Chattopadhyay (2007). It reported that the maximum ascorbic acid loss (14.48 mg/100g dry matter) and the minimum (3.12 mg/100g dry matter) were observed at the process conditions of 175°C, 125 s, 45% (wb) and 3.6 m/s and 150°C, 50 s, 35% (wb) and 3.0 m/s respectively. Haase and Weber (2003) observed degradation of ascorbic acid during processing of French fries and potato chips. During processing total losses of ascorbic acid were about 52% for French fries and about 26% for potato chips respectively. Losses of vitamin C during processing depend on the degree of heating (Alam, Singh, & Sawhney, 2010), leaching into the cooking medium, surface area exposed to oxygen and any other factors that facilitate oxidation (Eitenmiller & Laden, 1999).

The effect of temperature on decreasing the activity of soy product trypsin inhibitors is well known and described in the literature (Dublish, Chauhan, & Bains, 1988). Thermal inactivation of trypsin inhibitors of soybean preparations added to meat was reported by Kozłowska, Elkowicz, & Rutkowski (1980) and they found heat treatments at 70 and 100°C for 5 min reduced the trypsin inhibitor activity in the product by 40 and 70% respectively in comparison with unheated samples. By extending the heating time up to 15 min, a further reduction of activity by about 20 and 4% was caused at 70 and 100°C respectively when Dublish et al (1988) studied on nutritional quality of extruded rice, ragi and defatted soy flour blends and observed extrusion cooking resulted in the inactivation of trypsin inhibitor activity and at 65 and 95±2°C the inactivation was 42.6-71.4 and 72.3-100% respectively.

Nitrogen Solubility Index (NSI) is frequently used as indicators of protein functionality and potential end use (Hutton & Campbell, 1977). NSI is considered to be a good index of the potential use of a protein, providing information that is useful in snack foods processing and optimization. Several functional properties are directly related to this type of heat treatment (Mattil, 1971). Heat treatment to the food materials leads to alterations in specific structure of the proteins. In many instances, increase in digestibility of the protein occurs, perhaps by opening of the complex structure which serves as the site for digestive enzyme (Phillips, Chhinnan, & Mendosa, 1983). Bhole (1992) studied on the development of cereal-pulses based convenience food and reported 14.72% and 5.54% increase in nitrogen solubility index and invitro protein digestibility index respectively in the developed product. Quality evaluation of three canteen snacks of nutritative, fat and bacteriological was reported by Goyle, & Dugar (1994). They observed that the energy contents of the snacks were in between 249 and 475 kcal/100g.

There is no literature available on nutritional changes that took place during high temperature short time (HTST) air puffed potato and potato-soy snack production. Therefore, the present study was undertaken to evaluate the nutritional changes that took place during the production of RTE potato and potato-soy snacks.

Materials and Methods

Preparation of Potato Flour

Potato flour was prepared from freshly harvested potatoes (*Kufri chandramukhi*) as per the method described by Nath, Chattopadhyay, & Majumder (2007). The potato flour obtained were packed in air-tight containers and kept in dry and cold place for future use.

Preparation of Soy Flour

Soy beans (*Glycine max* Cv. JS 335) were procured from local market and washed thoroughly with clean water. Immediately after washing and sorting, raw soybean seeds were blanched for 15 min in hot water (100°C). Blanched soybeans were dehulled by manual rubbing. Splits obtained were steamed for 10 min. and then dried at 60°C for 5-6 hours in hot air cabinet drier up to a final moisture content of 8-10% (Khetarpaul, Grewal, Goyal, & Garg, 2004). The dried splits were ground in a domestic grinder (Sumeet Brand) for 15-20

min and passed through the sieve (British Sieve Standards) to yield flour of 50 mesh size. The soy flour thus obtained was packed in air-tight containers for future use.

Sample Preparation

RTE potato and potato-soy snack were prepared from potato flour dough with about 37% (wb) moisture content along with 2% NaCl and 10% soy flour added in total weight of potato flour respectively. Dolly pasta machine (LaMonferrina Make) was used to prepare a rectangular shape (15-20mm length, 10mm width and 1.5mm thickness) dimension chips from the dough. These chips were air puffed in a high temperature short time (HTST) fluidized bed air puffer specially designed and fabricated for the purpose (Nath et al., 2007). These snacks were subjected to oven toasting (Nath & Chattopadhyay, 2007) for obtaining final RTE potato and potato-soy snack. The nutritional compositions of initial samples before puffing and oven toasting and final products were measured in terms of moisture content, protein, fat, ash, crude fiber, carbohydrates (by difference), ascorbic acid content, nitrogen solubility index, protein digestibility, trypsin inhibitor activity, calorific value and protein energy ratio.

Moisture Content

The moisture content of the samples at every stage of the process was determined by hot air oven method as described by Ranganna (1995).

Protein Content

Protein Content was determined by AOAC (1984) method No. 2.049. Protein content was estimated as per equations given below:

$$\text{nitrogen\%} = \frac{[\{\text{Sample titre} - \text{Blank titre}\} \times \text{Normality of HCL} \times 4 \times \text{Vol. made up of the digest} \times 100]}{\text{Aliquot of the digest taken} \times \text{weight of the sample taken}} \quad (1)$$

$$\text{Protein} = \text{Nitrogen\%} \times 6.25 \quad (2)$$

Crude Fiber Content

It was determined according to AOAC (1984) method No. 7.054. Crude fiber content was estimated as per Eq 3 and 4 shown below:

$$\% \text{ crude fibre in ground sample} = C = \frac{(\text{loss in weight on ignition} - \text{loss in weight of asbestos blank}) \times 100}{\text{weight of the sample}} \quad (3)$$

$$\begin{aligned} & \% \text{ crude fibre on derived moisture basis} \\ & = C \times \frac{(100 - \% \text{ moisture disived})}{(100 - \% \text{ moisture in ground sample})} \end{aligned} \quad (4)$$

Ash, Fat and Carbohydrate Content

Ash content was determined according to AOAC (1984). However, fat soluble material in snack food was extracted from an oven-dried sample using a Soxhlet extraction apparatus with solvent hexane. Fat content was estimated as per Eq 5 shown below:

$$\begin{aligned} & \% \text{ Fat content} \\ & = \text{Wt of fat - soluble material} * 100 / \text{Wt of sample} \end{aligned} \quad (5)$$

The carbohydrate content was estimated by subtracting the values of moisture, protein, ash, crude fat and crude fiber from 100.

Ascorbic Acid Content

The ascorbic acid content of the samples was determined by visual titration method using 2, 6-dichlorophenol-indophenol (Ranganna, 1995). The ascorbic acid content of the sample was calculated by the formula:

$$\begin{aligned} & \text{Ascorbic acid content(mg/100g)} \\ & = \frac{\text{Titre} \times \text{Dye factor} \times \text{Volume made up} \times 100}{\text{Aliquot of extract} \times \text{Volume of sample}} \end{aligned} \quad (6)$$

Protein Digestibility Index

In vitro protein digestibility was estimated by enzymatic method of Akesson & Stahmann (1964). Digestibility was calculated using Eq 7 as shown below:

$$\begin{aligned} & \text{Protein digestibility (\%)} \\ & = \frac{N_2 \text{ in supernatant} - N_2 \text{ in sample blank at zero hour}}{N_2 \text{ in sample} - N_2 \text{ in enzyme blank (in TCA)}} \end{aligned} \quad (7)$$

Nitrogen Solubility Index

It was determined according to BIS (1975) method No. 7837. The nitrogen solubility index of the sample was calculated by using Eq 8 and 9 as shown below:

$$\begin{aligned} & \text{Water soluble nitrogen, percent by mass} \\ & = \frac{(V_1 - V_2) \times N \times 0.014 \times 100}{M} \end{aligned} \quad (8)$$

$$\begin{aligned} & \text{Nitrogen solubility index (NSI)} \\ & = \frac{\text{Water soluble nitrogen}}{\text{Total nitrogen}} \times 100 \end{aligned} \quad (9)$$

where, V_1 (volume in ml of alkali for back titration of blank); V_2 (volume in ml of alkali for back titration of sample); N (normality of alkali) and M (mass of sample).

Trypsin Inhibitor Activity

Trypsin inhibitor activity was determined according to BIS (1975) method No. 7837.

Calorific Value

Calorific values (kcal/100g) of both the raw materials and final products were calculated by summing up the multiplication of percent protein, fat and carbohydrate present in these materials by 4.04, 9.0 and 4.04 respectively (Mudambi & Rajagopal, 1983).

Protein Energy Ratio

The energy contributed by the protein to that of the total energy obtained from 100 g of the developed product was calculated by using Eq 10 as shown below:

$$\text{Protein energy ratio} = \frac{\text{Total protein(\%)}}{\text{Total energy (kcal/100g)}} \quad (10)$$

Statistical Analysis

Paired samples T-test were conducted using the statistical software 'SPSS, V-10' to compare the mean values of RTE potato snack with RTE potato-soy snack. The null hypothesis was tested and level of significant difference was also recorded between the products.

Results and Discussion

Preparation of Potato Flour and Soybean Flour

Potato slices were dried at a temperature of 60°C in a hot air cabinet drier for 4 hours in a single layer and obtained flour of 100 mesh sizes with drying ratios of 5.68 and yield of 16.2% with moisture content of 9.45% (db). However, soybean flour of 50 mesh size was obtained from dried soybean (Glycine max Cv. JS 335) splits by grinding in a domestic grinder with final moisture content of 10.1% (db) and flour yield was recorded as 83.6%.

Proximate and Nutritional Compositions of Raw

Materials

Potato flour, soy flour and defatted soy flour were analyzed for their proximate compositions and nutritional properties (Table 1). Potato flour was found to be rich in carbohydrate (75.52%) and ascorbic acid (41.76 mg/100g). It was observed that defatted soy flour was rich in protein (48.34%), ash (6.92%) and crude fiber (4.07%) as compared to full fat soy flour with protein (40.16%), ash (6.34%) and crude fiber (3.81%). Defatted soy flour was found 5.5 times higher in protein energy ratio than as compared to potato

flour. The proximate compositions of potato flour presented in Table 1 were found consistent with the results reported by Kulkarni, Kerure, & Kulkarni (1988) and Marwaha (1997). Tripathi, & Misra (2005) and, Murugkar, & Jha (2009) reported 13-25% oil, 30-50% protein and 14-24% carbohydrate in soybean seed and these results are in agreement with the present findings. On the contrary, Akubor, & Ukwuru (2003) reported higher values of carbohydrates (33%) and fat (18%) in soybean, which might be due to differences in agro-climatic conditions and varietal differences. Other nutritional compositions viz. nitrogen solubility index, protein digestibility index, energy value and protein energy ratio and were recorded as 65.70%, 83.44%, 342.05 kCal/100g and 0.024; 60.25%, 79.15%, 387.20 kCal/100g and 0.104; 62.25%, 81.37%, 365.90 kCal/100g and 0.132 respectively for potato flour, full fat and defatted soy flour (Table 1). However, trypsin inhibitor activities of full fat soy flour and defatted soy flour were recorded as 38.5% and 35.7% respectively which were much below the maximum prescribed limit of 55 units (BIS: 7836-1975 and 7837-1975).

TABLE 1 PROXIMATE AND NUTRITIONAL COMPOSITIONS OF RAW MATERIALS (POTATO FLOUR AND SOY FLOUR)

Parameters	Potato flour	Soy flour	
		full fat	defatted
MC, %, db.	9.5	10.1	9.8
Fat, %, db.	0.46	12.4	1.3
Protein, %, db.	8.1	40.2	48.3
Ash, %, db	3.8	6.3	6.9
Crude fiber, %, db	1.6	3.8	4.1
Carbohydrates, %*	75.5	28.2	39.4
Ascorbic acid, mg/100g, db	41.8	0.00	0.00
Nitrogen solubility index, %	65.7	60.3	62.0
Protein digestibility index, %	83.4	79.2	81.4
Trypsin inhibitor activity, %	-	38.5	35.7
Energy, Kcal/100g	342.1	387.2	365.9
Protein Energy Ratio	0.02	0.10	0.13

* By difference

MC, moisture content

Nutritional Properties of RTE Potato Snack and RTE Potato-Soy Snack during Processing

The proximate composition and nutritional properties of RTE potato snacks and RTE potato-soy snacks (initial and final products) were analyzed and presented in Table 2. During processing the maximum loss of fat, protein, carbohydrates, ascorbic acid content and protein energy ratio were found to be 31.11%, 2.19%, 0.44%, 31.87% and 1.28% respectively for RTE potato snack and 27.69%, 2.25%, 0.53%, 30.12% and 1.05% respectively for RTE potato-soy snack while, the maximum loss of trypsin inhibitor activity was observed to be 63.66% for RTE potato-soy snack. Haase,

& Weber (2003) also observed degradation of ascorbic acid during processing of French fries and potato chips. During processing total losses of ascorbic acid AA were about 52% for French fries and about 26% for potato chips respectively. Losses of vitamin C during processing depend on the degree of heating, leaching into the cooking medium, surface area exposed to oxygen and any other factors that facilitate oxidation (Eitenmiller & Laden, 1999). The effect of temperature (i.e. processing) on decreasing the activity of soy product trypsin inhibitors is well known and described in the literature (Chauhan, Verma, & Bains, 1985 and Dublish et al., 1988). Thermal inactivation of trypsin inhibitors of soybean preparations added to meat was reported by Kozłowska et al. (1980) and they found heat treatments at 70 and 100°C for 5 min reduced the trypsin inhibitor activity in the product by 40 and 70% respectively, in comparison with unheated samples. By extending the heating time up to 15 min, a further reduction of activity by about 20 and 4% was caused at 70 and 100°C respectively when Dublish et al. (1988) studied on nutritional quality of extruded rice, ragi and defatted soy flour blends and observed extrusion cooking resulted in the inactivation of trypsin inhibitor activity and at 65 and 95±2°C the inactivation was 42.6-71.4 and 72.3-100% respectively. However, during processing the maximum increase of ash content, crude fiber, nitrogen solubility index and invitro protein digestibility index were recorded to be 16.99%, 2.42%, 12.69% and 8.59% respectively for RTE potato snack and 11.15%, 5.39%, 11.52% and 8.84% respectively for RTE potato-soy snack. It was observed that due to processing Nitrogen Solubility Index (NSI) and Protein Digestibility Index (PDI) values were also increased and this may due to the release of some protein fraction into the products during processing. NSI are frequently used as indicators of protein functionality and potential end use (Hutton and Campbell 1977). NSI is considered to be a good index of the potential use of a protein, providing information that is useful in snack foods processing and optimization. Several functional properties are directly related to this type of heat treatment (Mattil, 1971). Heat treatment to the food materials leads to alterations in specific structure of the proteins. In many instances, increase in digestibility of the protein occurs, perhaps by opening of the complex structure which serves as the site for digestive enzyme (Phillips et al. 1983). Bhole (1992) studied on the development of cereal-pulses based convenience food and he reported 14.72% and 5.54% increase in nitrogen solubility index and invitro protein digestibility index respectively in the developed product. Quality

evaluation of three canteen snacks of nutritive value, fat and bacteriological was reported by Goyle, & Dugar (1994). They observed that the energy contents of the snacks were in between 249 and 475 kcal/100g. These results are in agreement with the present findings.

Comparison of Nutritional Properties of RTE Potato Snack with RTE Potato-Soy Snack

The mean values of nutritional parameters for RTE potato snack and RTE potato-soy snack and their standard deviations are presented in Table 3. The null hypothesis was tested and there was highly significant difference recorded between the products. High significant difference ($p \leq 0.01$) between RTE potato snack and RTE potato-soy snack were found except for moisture content in which it was non-significant (Table 3). Significant increase in protein (56.67%), ash (70.49%), crude fiber (27.2%) and protein energy ratio (63.20%) of RTE potato-soy snack as compared to RTE potato snack has been recorded due to addition of soy flour (10.26%) in the product. Laul, Bhalerao, &

Mulmuley (1985) studied on the preparation of extruded finger shaped snack foods supplemented with defatted soybean to corn, broken rice and sorghum and they reported that addition of soy flour increased protein content in the finished product to 12% level. Soy flour incorporation in potato based foods like extruded and traditional type can, not only give better colored product, but will make more protein available for nutrition due to lesser formation of mailared product (Kulkarni & Joshi, 1992). These findings are in accordance with the present findings.

Conclusions

Significant changes in nutritional composition between RTE potato and RTE potato-soy snacks were observed during high temperature short time air puffing and oven toasting process. Addition of soy flour (10.26%) caused increase in protein (56.67%), ash (70.49%), crude fiber (27.2%) and protein energy ratio (63.20%). The maximum loss of trypsin inhibitor activity was observed to be 63.66% for RTE potato-soy snack.

TABLE 2 CHANGES IN NUTRITIONAL PROPERTIES OF RTE POTATO AND RTE POTATO-SOY SNACKS DURING PROCESSING

Parameters	RTE potatosnack		Loss, %	Gain, %	RTE potato-soysnack		Loss, %	Gain, %
	Initial	Final			Initial	Final		
MC, %	35.6	3.3	90.7		35.6	3.4	90.6	
Fat, %	0.45	0.31	31.1		0.65	0.47	27.7	
Protein, %	8.2	8.0	2.2		12.9	12.6	2.3	
Ash, %	3.7	4.3		16.9	6.6	7.3		11.2
Crude fiber, %	1.7	1.7		2.4	2.0	2.2		5.4
Carbohydrates, %*	77.5	77.2	0.44		69.4	68.9	0.53	
Ascorbic acid, mg/100g	37.2	25.4	31.9		32.7	22.8	30.1	
Nitrogen solubility index	66.4	74.9		12.7	62.3	69.5		11.5
Protein digestibility index	85.2	92.6		8.6	82.5	89.7		8.8
Trypsin inhibitor activity	-	-			3.6	1.3	63.7	
Energy, Kcal/100g	350.4	347.00	0.96		338.1	333.8	1.3	
Protein Energy Ratio	0.02	0.02	1.3		0.04	0.04	1.1	

* By difference

MC, moisture content; RTE, ready-to-eat

TABLE 3 COMPARISON OF NUTRITIONAL PROPERTIES OF RTE POTATO SNACK WITH RTE POTATO-SOY SNACK

Parameters	Mean \pm SD		%Variation	Standard error of Mean	Sig(2-tailed)
	RTE potato	RTE potato-soy			
MC, %	3.3 \pm 0.047	3.4 \pm 0.049	0.90	0.03	0.441
Fat, %	0.31 \pm 0.023	0.47 \pm 0.016	51.6	0.02	0.000
Protein, %	8.0 \pm 0.093	12.6 \pm 0.053	56.7	0.04	0.000
Ash, %	4.3 \pm 0.041	7.3 \pm 0.098	70.5	0.05	0.000
Crude fiber, %	1.7 \pm 0.070	2.2 \pm 0.040	27.2	0.03	0.000
Carbohydrates, %*	77.2 \pm 0.684	68.9 \pm 0.336	-10.6	0.36	0.000
Ascorbic acid, mg/100g	25.4 \pm 0.186	22.8 \pm 0.186	-10.0	0.14	0.000
Nitrogen solubility index	74.9 \pm 0.380	69.5 \pm 0.509	-7.6	0.25	0.000
Protein digestibility index	92.6 \pm 0.207	89.7 \pm 0.449	-3.0	0.22	0.000
Trypsin inhibitor activity	0	1.3 \pm 0.100		0.04	0.000
Energy, Kcal/100g	347.0 \pm 15.313	333.8 \pm 1.625	-3.8	6.6	0.114
Protein Energy Ratio	0.02 \pm 0.000	0.04 \pm 0.000	63.2	0.00	0.000

* By difference

MC, moisture content; SD, standard deviation; RTE, ready-to-eat

REFERENCES

- Akeson, W.R., & Stahmann, M.A. (1964). A pepsin-pancreatin digest index of protein quality. *Journal of Nutrition*, 83(1), 257-261.
- Akubor, P.I., & Ukwuru, K. Functional properties and biscuit making potential of soybean and cassava flour blends. *Plant Food for Human Nutrition*, 2003, 55(1), 1-12.
- Alam, M.S., Singh, A., & Sawhney, B.K. Response surface optimization of osmotic dehydration process for aonla slices. *Journal of Food Science and Technology*, 2010, 47(1): 47-54.
- AOAC. (1984). Official methods of analysis of the Association of Official Analytical Chemists. 14th edn, Washington, DC.
- Bhole, S.N. (1992). Studies on the development of cereal-pulses based convenience food. Unpublished Ph.D. thesis, PHTC, IIT, Kharagpur (W.B.), India.
- BIS. (1975). Specification for edible full-fat and low fat-soya flour. Optimum requirements, IS: 7836-1975 and 7837-1975. Bureau of Indian Standards, New Delhi.
- Chauhan, G.S., Verma, N.S. & Bains, G.S. (1985). Effect of extrusion cooking on nutritional quality of extruded rice-legume blends. Paper presented in 5th Indian Convention of Food Scientists and Technologists, 03-05 Apr, 1985, New Delhi.
- Dublish, R.K., Chauhan, G.S. & Bains, G.S. (1988). Nutritional quality of extruded rice, ragi and defatted soy flour blends. *Journal of Food Science and Technology*, 25(2), 35-39.
- Eitenmiller, R.R. & Laden, W.O. (1999). *Ascorbic acid*. In: Eitenmiller RR and Laden WO. Edn, Vitamin Analysis for the Health and Food Science. CRC Press, Boca Raton, FL, 226-228.
- Goyle, A. & Dugar, N. (1994). Quality evaluation of three canteen snacks: nutritive, fat and bacteriological. *Indian Food Packer*, 1, 53-58.
- Haase, N.U. & Weber, L. (2003). Ascorbic acid losses during processing of French fries and potato chips. *Journal of Food Engineering*, 56(2), 207-209.
- Hutton, C.W. & Campbell, A.M. (1977). Functional properties of soy concentrate and soy isolate in simple systems-NSI and water absorption. *Journal of Food Science*, 42(4), 454-457.
- Khetarpaul, N., Grewal, R.B., Goyal, R. & Garg, R. (2004). Development of partially defatted soy flour and dhal. *Food Chemistry*, 87, 355-359.
- Kozłowska, H., Elkowicz, K. & Rutkowski, A. (1980). Thermal inactivation of trypsin inhibitors of soybean preparations added to meat. *Meat Science*, 4, 95-102.
- Kulkarni, K.D. & Joshi, K.C. (1992). Potato starch soy-blends: Possible effects of starch properties on few aspects of end products. *Indian Food Packer*, 66, 38-49.
- Kulkarni, K.D., Kerure, Y.E. & Kulkarni, D.N. (1988). Evaluation of potato cultivars for flour production. *Beverage and Food World*, 8, 214-221.
- Laul, M.S., Bhalerao, S.D. & Mulmuley, G.V. (1985). Studies on the preparation of extruded finger shaped snack foods with protein enrichment. *Indian Food Packer*, 32, 68-77.
- Marwaha, R.S. (1997). Processing of potatoes: current status, need, future potential and suitability of Indian varieties - a critical appraisal. *Journal of Food Science and Technology*, 34, 457-471.
- Mattil, K.F. (1971). The functional requirement of proteins in foods. *Journal of American Oil and Chemical Society*, 48, 477-480.
- Mishkin, M., Saguy, I. & Karel, M. (1984). A dynamic test for kinetic models of chemical changes during processing: Ascorbic acid degradation in dehydrated potatoes. *Journal of Food Science*, 49, 1276-1274.
- Mudambi, S.R. & Rajagopal, M.V. (1983). *The nutrients*. In: Fundamentals of Food and Nutrition. Wiley Eastern Ltd., New Delhi.
- Mukherjee, S. & Chattopadhyay, P.K. (2007). Whirling bed blanching of potato cubes and its effects on product quality. *Journal of Food Engineering*, 78, 52-60.
- Murugkar, D.A. & Jha, K. (2009). Effect of sprouting on nutritional and functional characteristics of soybean. *Journal of Food Science and Technology*, 46(3), 240-243.
- Nath, A. & Chattopadhyay, P.K. (2007). Optimization of Oven Toasting for Improving Crispness and Other Quality Attributes of Ready to Eat Potato-Soy Snack Using Response Surface Methodology. *Journal of Food Engineering*, 80, 1282-1292.
- Nath, A., Chattopadhyay, P. K. & Majumder, G.C. (2007). High temperature short time air puffed ready-to-eat (RTE) potato snacks: process parameter optimization. *Journal of Food Engineering*, 80, 770-780.

Phillips, R.D., Chinnan, M.S. & Mendosa, L.G. (1983). Effect of temperature and moisture content on kinetics of TIA, protein in vitro digestibility and nitrogen solubility of cowpea flour. *Journal of Food Science*, 48, 1863.

Ranganna, S. (1995). *Handbook of Analysis and Quality Control*

for Fruits and Vegetable Products. Second edition. Tata McGraw Hill Publishing Co. Ltd. New Delhi.

Tripathi, A. K. & Misra, A. K. (2005). Soybean- a consummate functional food: A review. *Journal of Food Science and Technology*, 42, 111-119.