

**State: Jharkhand**

**Agriculture Contingency Plan for District: Garhwa**

<b>1.0 District Agriculture profile</b>					
<b>1.1</b>	<b>Agro-Climatic/Ecological Zone</b>				
	Agro Ecological Sub Region (ICAR)	Moderately To Gently Sloping ChattisgarhMahanadi Basin, Hot Moist/Dry Subhumid Transitional ESR (11.0)			
	Agro-Climatic Zone (Planning Commission)	EASTERN PLATEAU AND HILLS REGION (VII)			
	Agro Climatic Zone (NARP)	WESTERN PLATEAU ZONE (BI-5)			
	Geographic coordinates of district headquarters	Latitude	Longitude	Altitude	
		23 <sup>0</sup> 34' 11'' – 24 <sup>0</sup> 32' -05'' N	83 <sup>0</sup> .10' 13'' - 83 <sup>0</sup> 56' 38'' E	364 m	
	Name and address of the concerned ZRS/ ZARS/ RARS/ RRS/ RRTTS	Zonal Research Station (Z.R.S.), Chianki, medinigar, Palamau, Pin – 822133 (Birsa Agricultural University, Ranchi) Pin – 834006.			
Mention the KVK located in the district	Krishi Vigyan Kendra, Garhwa, Pin - 822114				
<b>1.2</b>	<b>Rainfall</b>	Normal RF(mm) 2008	Normal Rainy days (number)	Normal Onset ( specify week and month)	Normal Cessation (specify week and month)
	SW monsoon (June-Sep):	780.8	47	3 <sup>rd</sup> week of June	4 <sup>th</sup> week of September
	NE Monsoon(Oct-Dec):	00	00		
	Winter (Jan- March)	53.4	31	-	-
	Summer (Apr-May)	31.6	3	-	-
	Annual Average rain fall 1355	865.8	53	-	-

<b>1.3</b>	<b>Land use pattern of the district</b> (latest statistics)	Geographical area	Cultivable area	Forest area	Land under non-agricultural use	Permanent pastures	Cultivable wasteland	Land under Misc. tree crops and groves	Barren and uncultivable land	Current fallows	Other fallows
	<b>Area ('000 ha)</b>	428.550	100.950	169.790	19.456	2.05704	6.600	2.271	24.727	102.351	

<b>1.4</b>	<b>Major Soils (common names like red sandy loam deep soils (etc.))*</b>	<b>Area ('000 ha)</b>	<b>Percent (%) of total</b>
	Sandy loam	201.419	47
	Red loam (Moram)	184.277	43
	Grey soil	42.855	10

<b>1.5</b>	<b>Agricultural land use</b>	<b>Area ('000 ha)</b>	<b>Cropping intensity %</b>
	Net sown area	100.950	115
	Area sown more than once	35.178	
	Gross cropped area	136.128	

<b>1.6</b>	<b>Irrigation</b>	<b>Area ('000 ha)</b>		
	Net irrigated area	35.180		
	Gross irrigated area	35.180		
	Rainfed area	65.770		
	<b>Sources of Irrigation</b>	<b>Number</b>	<b>Area ('000 ha)</b>	<b>Percentage of total irrigated area</b>
	Canals, Ponds, check dams, lifts		19.740	19.554
	Tanks			7.53
	Open wells		7.6	0.86
	Bore wells, Pump sets		0.870	

Lift irrigation schemes			
Micro-irrigation			
Other sources (please specify)			
Total Irrigated Area		28.210	27.95
Pump sets			
No. of Tractors			
<b>Groundwater availability and use*</b> <b>(Data source: State/Central Ground</b> <b>water Department /Board)</b>	No. of blocks/ Tehsils	(%) area	Quality of water (specify the problem such as high levels of arsenic, fluoride, saline etc)
Over exploited			
Critical			
Semi- critical			
Safe			
Wastewater availability and use			
Ground water quality			
*over-exploited: groundwater utilization > 100%; critical: 90-100%; semi-critical: 70-90%; safe: <70%			

**1.7 Area under major field crops & horticulture (as per latest figures) (Specify year 2009-10 eg., 2008-09)**

1.7	Major field crops cultivated	Area ('000 ha)							
		<i>Kharif</i>			<i>Rabi</i>				
		Irrigated	Rainfed	Total	Irrigated	Rainfed	Total	Summer	Grand total
	Paddy			51.00					
	Maize	1.5	25.6	27.1					
	Wheat				6.875	-	6.875		
	Pigeonpea		14.833	14.833					
	Mustard						4.00		
	Chickpea					3.033	3.033		

Source : Rabi and Kharif report (2009-10) Department of Agriculture, Garhwa.

	Horticulture crops - Fruits	Area ('000 ha)		
		Total	Irrigated	Rainfed
		-	-	-
	Horticulture crops - Vegetables	Total	Irrigated	Rainfed
	Potato	1.5		

	Okra	1.00		
	Chilies	0.670		
	Brinjal	0.640		
	Tomato	0.600	0.600	
	Cauliflower	0.520	0.520	
<b>Medicinal and Aromatic crops -</b>				
<b>Plantation crops -</b>				
<b>Fodder crops -</b>				
	<b>Total fodder crop area</b>			
	<b>Grazing land</b>			
	<b>Sericulture etc</b>			
	<b>Others (specify)</b>			

<b>1.8</b>	<b>Livestock</b>	<b>Male ('000)</b>	<b>Female ('000)</b>	<b>Total ('000)</b>
	Non descriptive Cattle (local low yielding)	193.407 (Bullocks)	397.995	591.402
	Calf			79.173
	Non descriptive Buffaloes (local low yielding)			15.835
	Graded Buffaloes			-
	Goat			159.305

	Sheep				1.131		
	Pig				28.051		
	Commercial dairy farms (Number)						
<b>1.9</b>	<b>Poultry</b>	<b>No. of farms</b>	<b>Total No. of birds ('000)</b>				
	Commercial		344.966				
	Backyard						
<b>1.10</b>	<b>Fisheries (Data source: Chief Planning Officer)</b>						
	<b>A. Capture</b>						
	<b>i) Marine</b> (Data Source: Fisheries Department)	<b>No. of fishermen</b>	<b>Boats</b>		<b>Nets</b>		<b>Storage facilities (Ice plants etc.)</b>
			Mechanized	Non-mechanized	Mechanized (Trawl nets, Gill nets)	Non-mechanized (Shore Seines, Stake & trap nets)	
	<b>ii) Inland</b> (Data Source: Fisheries Department)	<b>No. Farmer owned ponds</b>		<b>No. of Reservoirs</b>	<b>No. of village tanks</b>		
	<b>B. Culture</b>						
		<b>Water Spread Area (ha)</b>		<b>Yield (t/ha)</b>		<b>Production ('000 tons)</b>	
	<b>i) Brackish water</b> (Data Source: MPEDA/ Fisheries Department)						
	<b>ii) Fresh water</b> (Data Source: Fisheries Department)						
	<b>Others</b>						

**1.11 Production and Productivity of major crops** (Average of last 5 years: 2004, 05, 06, 07, 08; specify years)

1.11	Name of crop	Kharif		Rabi		Summer		Total		Crop residue as fodder ('000 tons)
		Production ('000 t)	Productivity (kg/ha)	Production ('000 t)	Productivity (kg/ha)	Production ('000 t)	Productivity (kg/ha)	Production ('000 t)	Productivity (kg/ha)	
<b>Major Field crops (Crops to be identified based on total acreage)</b>										
	Rice	18.58	1279.0					18.58	1279.0	
	Maize	14.24	870.0	3.75	2500.00			17.99		
	Pigeonpea	3.00	460.0					3.00	460.00	
	Wheat			13.75	2000.00			13.75	2000.00	
	Mustard			2.00	500.00			2.00	500.00	
<b>Major Horticultural crops (Crops to be identified based on total acreage) -</b>										

1.12	Sowing window for 5 major field crops (start and end of normal sowing period)	Paddy	Maize	Pigeonpea	Wheat	Mustard
	Kharif- Rainfed	2 <sup>nd</sup> week of June to 3 <sup>rd</sup> week of June	2 <sup>nd</sup> week of June to 1 <sup>st</sup> week of July	1 <sup>st</sup> week of June to 3 <sup>rd</sup> week of June		
	Kharif-Irrigated	1 <sup>st</sup> week of June to 1 <sup>st</sup> week of July	2 <sup>nd</sup> week of June to 4 <sup>th</sup> week of June	2 <sup>nd</sup> week of June to 3 <sup>rd</sup> week of July		
	Rabi- Rainfed				2 <sup>nd</sup> week of October to 4 <sup>th</sup> week of October	1 <sup>st</sup> week of October to 2 <sup>nd</sup> week of October
	Rabi-Irrigated				2 <sup>nd</sup> week of November to 4 <sup>th</sup>	1 <sup>st</sup> week of October to 2 <sup>nd</sup>

					week of November	week of November
--	--	--	--	--	------------------	------------------

1.13	What is the major contingency the district is prone to? (Tick mark)	Regular	Occasional	None
	Drought			
	Flood			
	Cyclone			
	Hail storm			
	Heat wave			
	Cold wave			
	Frost			
	Sea water intrusion			
	Pests and disease outbreak (specify)			
	Others (specify) Late blight in potato			

1.14	Include Digital maps of the district for		
		Location map of district within State as Annexure I	Enclosed: Yes
		Mean annual rainfall as Annexure 2	Enclosed: No
		Soil map as Annexure 3	Enclosed: Yes / No



Annexure 1



## 2.0 Strategies for weather related contingencies

### 2.1 Drought

#### 2.1.1 Rainfed situation

Condition			Suggested Contingency measures		
Early season drought (delayed onset)	Major Farming situation <sup>a</sup>	Normal Crop / Cropping system <sup>b</sup>	Change in crop / cropping system <sup>c</sup> including variety	Agronomic measures <sup>d</sup>	Remarks on Implementation <sup>e</sup>
Delay by 2 weeks (Specify month)*  1 <sup>st</sup> week of July	1 Upland	Maize + Pigeonpea	Maize + Cowpea Maize: Birsa Maize-1	Maize : 60 cm x 20 cm	Supply of seeds under RKVY
		Pigeonpea + Groundnut	Pigeonpea : Bahar, Narendra Arhar, Birsa Arhar-1		
		Maize + Groundnut	Groundnut: AK. 12.24, Birsa Bold		
		Blackgram/ Sesame	Sesame : Kanke safed		
	Finger millet	Finger millet: A 404 (Transplanting)/ Pigeonpea + Castor /Soybean/ Sorghum Soybean : Birsa Soybean safed – 2 Sorghum: CSV – 20	Transplanting Field management by banding	MNREGS & NWDPRA Schemes	
	2 Medium land Medium deep sandy loam	Rice Hybrids	Rice : Sahbhagi, Naveen, IR - 64	Direct sowing through drum seeder Use of pre emergence weedicides e.g. Glyphosate or post emergence Butachlor.	Supply of seeds and implements under RKVY
			Hybrids: PAC. 807, Uday – 111, Abhishek, 27P31		
3 Lowland deep	Rice	Rajendra Mansuri, Rajshree (OP) Sonam,	• Transplanting at		

	clay		Rupali, Mansuri, Vandana, BVD – 109, IR- 64 Local, Birsamati, MTU - 7029	<ul style="list-style-type: none"> <li>closer spacing</li> <li>Field Bunding</li> </ul>	
		Arize – 6444, Advanta – 801	IR – 36, IR - 64		
		PHB – 71 (Hybrid)	PAC – 807, 25P31, Abhishek ,Uday-111 (Hybrids)		MNREGS & NWDPRASchemes
		Blackgram	Urd – Birsa Urd – 1, T – 9, Pant U - 19		
<b>Condition</b>			<b>Suggested Contingency measures</b>		
<b>Early season drought (delayed onset)</b>	<b>Major Farming situation<sup>a</sup></b>	<b>Normal Crop/cropping system<sup>b</sup></b>	<b>Change in crop/cropping system<sup>c</sup></b>	<b>Agronomic measures<sup>d</sup></b>	<b>Remarks on Implementation<sup>e</sup></b>
<b>Delay by 4 weeks (Specify month)</b> 3 <sup>rd</sup> week of July	Rainfed, shallow, acidic, redsoil	Maize + Pigeonpea, Maize + Groundnut Maize + Blackgram	Maize + Cowpea, Maize + Okra, Maize + Ridgeground Maize : Birsa Vikas makka -2  Pigeonpea : Birsa Arhar – 1 ICPH – 2671 Pigeonpea + Fingermillet (A 404)  Soybean : Birsa Soybean Safed- 2  Sorghum: CSV – 20  Sesame : Kanke white, Pragati.	Closer spacing of Pigeon pea and maize	Seeds to be needed available under RKVY and NFSM

	Medium deep sandy loom soil	Rice – IR – 36, IR – 64	Rice : Vandana, Naveen, Sahbhagi, PAC – 807, 27 P 31, Uday - 111, Abhishek (Hybrid)	
	Deep, heavy clay soil	Rice	Naveen, IR – 64, IR – 36, Rupali, PAC – 807, Uday – 111, Abhishek	

Condition	Major Farming situation <sup>a</sup>	Normal Crop/cropping system <sup>b</sup>	Suggested Contingency measures		
			Change in crop/cropping system <sup>c</sup>	Agronomic measures <sup>d</sup>	Remarks on Implementation <sup>e</sup>
<b>Early season drought (delayed onset)</b>  <b>Delay by 6 weeks (Specify month)</b> 1 <sup>st</sup> week of August	Upland	Maize + Pigeonpea	Maize + Cowpea (Swarna Harita/S. Sweta) Maize : Birsa Maize – 2 Cowpea : Swarna Harita, Swarna Sweta		
		Sesame/ Blackgram	Pigeonpea : Birsa Arhar – 1 + Okra		
		Finger millet	Sorghum: CSV – 20		
			Groundnut : Birsa Bold		
			Soybean : Birsa Soyabean safed – 2		
	2. Medium land sandy loom with medium depth	Rice	Rice : Sahabhagi, Naveen (OP) PAC – 807 27P - 31 (Hybrid)		Vegetable seeds supply under RKVY
		Maize	Maize : Birsa Maize – 2, HSPM - 1		
		Vegetable	Vegetable : Tomato, Brinjal, Cowpea		

Condition			Suggested Contingency measures		
Early season drought (delayed onset)	Major Farming situation <sup>a</sup>	Normal Crop/cropping system <sup>b</sup>	Change in crop/cropping system <sup>c</sup>	Agronomic measures <sup>d</sup>	Remarks on Implementation <sup>e</sup>
Delay by 8 weeks (Specify month) 3 <sup>rd</sup> week of August	1. Upland, Rainfed, Redsoil.	Maize + Pigeonpea	Pigeonpea – Birsa Arhar-1		
		Sesame/ Blackgram	Sesame – Kanke safed, Pragati, Krishna		
			Blackgram - Shekhar-2		
			Finger millet - A 404 (Transplanting)		
			Niger – Birsa Niger – 1		
			Horsegram : Birsa Kulthi–1 JNC – 6		
2. Medium land medium deep sandy loom soil	Rice / Pigeonpea +Sesame	Pigeonpea – ICPH – 2671, Bahar Narendra Arhar – 1/ Pigeonpea + Castor/Sesame /Toria – PT 303/ Niger–JNC-6,Birsa Niger–1/Horsegram – Birsa Kulthi–1/ Tomato – Swarna Lalima /Cabbage – Early Kuwari, Pusa Deepali		<ul style="list-style-type: none"> <li>• Close planting (60cm x 20 cm)</li> <li>• Closer spacing and planting of average seedlings</li> <li>• Closer spacing</li> <li>• Heavy dose of NPK</li> </ul>	Seed supply under RKVY
3. Lowland clay soil		Rice : Naveen, Sahbhagi, Vandana, PAC-807, Abhishek, Uday-111	Transplanting of average seedlings of long duration varieties/ hybrids		

Condition	Major Farming situation <sup>a</sup>	Normal Crop/cropping system <sup>b</sup>	Suggested Contingency measures		
			Crop management <sup>c</sup>	Soil nutrient & moisture conservation measures <sup>d</sup>	Remarks on Implementation <sup>e</sup>
Early season drought (Normal onset)					
Normal onset followed by 15-20 days dry spell after sowing leading to poor germination/crop stand etc.	1. Upland	Maize + Pigeonpea	Re sowing Maize (Birsa Vikas Makka-2 Birsa Arhar – 1 Sesame – Kanke safed Groundnut : Birsa Bold	Sprinkler system or irrigation by lifting the water from rivers	Supply of sprinkler system under RKVY
		Maize + Blackgram			
		Pigeonpea + Sesame			
		Maize + Groundnut			Supply of seeds of pulses under NFSM (Pulses)
		Pigeonpea + Groundnut			
Finger millet					
2 Medium deep Medium land	Rice soil	Direct sowing of Naveen, Shabhagi, Vandana.	Irrigation of crop by lifting the water from ponds, wells.	Supply of Pipe – Pumps under RKVY	
3. Low land deep clay soil.	Rice crop		Irrigation of crops by lifting the water from well of ponds		

Condition	Major Farming situation <sup>a</sup>	Normal Crop/cropping system <sup>b</sup>	Suggested Contingency measures		
			Crop management <sup>c</sup>	Soil nutrient & moisture conservation measures <sup>d</sup>	Remarks on Implementation <sup>e</sup>
Mid season drought (long dry spell, consecutive 2 weeks rainless (>2.5 mm) period)					
	Uplands	Maize + Pigeonpea /	1. Supply of life saving	1. Application of	

At vegetative stage		Blackgram / Pigeonpea + Sesame/ Maize + Groundnut/Cowpea / Finger millet	irrigation 2. Weeding cum – hoeing to break capillarity 3. Finger millet has better drought tolerance capacity (Area extension) 4. Weeding and weed mulching of the field	compost to enhance the water holding capacity of soil 2. Judicious land of P for better penetration of root system 3. Weeding and weed mulching of the field 4. Pre sowing application of compost and judicious land of P&K for better water holding and root growth.	
	2. Medium land	Rice – IR – 36, IR – 64, Saryu – 52	IR – 36, IR – 64, Saryu – 52	<ul style="list-style-type: none"> <li>• Life saving irrigation through pumps and sprinkler.</li> <li>• Area extension under Shabhagi Dhan</li> </ul>	Supply of Pumps (Sprinkler) sets under RKVY
	3. Lowland	rice	Low lands mostly covered under hybrids with stand 2 – 3 weeks long stress PHB – 71, A – 801, Arize 6444, Rupali, Sonam	Life saving irrigation through Pumps	

Condition	Major Farming situation <sup>a</sup>	Normal Crop/cropping system <sup>b</sup>	Suggested Contingency measures		
			Crop management <sup>c</sup>	Soil nutrient & moisture conservation measures <sup>d</sup>	Remarks on Implementation <sup>e</sup>
Mid season drought (long dry spell)	1. Upland Shallow red soils	Maize + Pigeonpea Maize + Blackgram Pigeonpea + Sesame Maize + Groundnut Pigeonpea + Groundnut Finger millet	Life saving irrigation through sprinkler system Weed – cum – hoeing and weed mulching	Intercultivation ( soil mulching )  Conservation Furrow	Supply of seeds through D.A.O.  Supply seeds through N.F.S.M.
	2. Medium land medium deep sandy clay loom.	Rice : IR – 36 , IR - 64		Life irrigation by lifting the water from ponds/ wells	Supply of irrigation devices under RKVY.
	3. Low land deep heavy clay soil.	Rice : Varieties and Hybrids Sonam, Rupali, Arize – 6444, PHB – 71		Life saving irrigation through Pumps/Ponds/wells	



Condition	Major Farming situation <sup>a</sup>	Normal Crop/cropping system <sup>b</sup>	Suggested Contingency measures		
			Crop management <sup>c</sup>	Rabi Crop planning <sup>d</sup>	Remarks on Implementation <sup>e</sup>
Terminal drought (Early withdrawal of monsoon)					
	1. Upland shall one red soils	Maize + Pigeonpea / Blackgram/ Groundnut/Cowpea Pigeonpea + Sesame Finger millet	Life saving irrigation Harvesting of pods of Cowpea and Black gram for vegetable purpose and fodder	Niger, Mustard, Chickpea, Linseed.	Supply of Pumps (Sprinkler) sets under RKVY
	2. Medium land	Rice : Varieties	Supply of life saving irrigation lifting the water from ponds wells.	Rye + Wheat, Linseed + Horsegram, Niger, Toria, Chickpea, Vegetable like – tomato, Vegetable pea, Potato, Wheat + Mustard, Lentil	Seeds and planting materials supply under RKVY
	3. Low land	Long duration rice varieties and hybrids.	Life saving irrigation Crop protection measures		Ponds/wells under MNREGS and RKVY

### 2.1.2 Drought - Irrigated situation

Condition	Major Farming situation <sup>f</sup>	Normal Crop/cropping system <sup>g</sup>	Suggested Contingency measures		
			Change in crop/cropping system <sup>h</sup>	Agronomic measures <sup>i</sup>	Remarks on Implementation <sup>j</sup>
Delayed release of water in canals due to low rainfall	Mention source of irrigation, topography (upland/lowland) and soil colour & depth Eg; canal irrigated shallow red soils; tankfed medium deep				

Condition	Suggested Contingency measures				
	Major Farming situation <sup>f</sup>	Normal Crop/cropping system <sup>g</sup>	Change in crop/cropping system <sup>h</sup>	Agronomic measures <sup>i</sup>	Remarks on Implementation <sup>j</sup>
	black soils				
	Medium land sandy/clay looms	Medium land rice – IR – 36, IR – 64.	Aerobic rice var. Naveen, Shabhagi, Vandana.	Limited irrigation by the sprinklers.	Seed through RKVY, NFSM (Pigeonpea)
			Maize + Pigeonpea	Irrigation through drip or alternate row irrigation.	
			Pigeonpea + Sesame		
			Vegetable – Tomato, Chilli		
			Treated Cucurbits		

Condition	Suggested Contingency measures				
	Major Farming situation <sup>f</sup>	Normal Crop/cropping system <sup>g</sup>	Change in crop/cropping system <sup>h</sup>	Agronomic measures <sup>i</sup>	Remarks on Implementation <sup>j</sup>
		Rice	Aerobic rice / Transplanted rice var. Naveen, Shabhagi, Vandana.		
			Pigeonpea + Sesame Maize + Pigeonpea Tomato, Chillies		
	Medium land	Medium maturing rice varieties	Direct sown (Aerobic) Rice var. Naveen Shabhagi, Vandana, IR - 64		

Condition	Suggested Contingency measures				
	Major Farming situation <sup>f</sup>	Normal Crop/cropping system <sup>g</sup>	Change in crop/cropping system <sup>h</sup>	Agronomic measures <sup>i</sup>	Remarks on Implementation <sup>j</sup>
Non release of	NA				

Condition	Suggested Contingency measures				
	Major Farming situation <sup>f</sup>	Normal Crop/cropping system <sup>g</sup>	Change in crop/cropping system <sup>h</sup>	Agronomic measures <sup>i</sup>	Remarks on Implementation <sup>j</sup>
water in canals under delayed onset of monsoon in catchment					
Condition	Suggested Contingency measures				
	Major Farming situation <sup>f</sup>	Normal Crop/cropping system <sup>g</sup>	Change in crop/cropping system <sup>h</sup>	Agronomic measures <sup>i</sup>	Remarks on Implementation <sup>j</sup>
Lack of inflows into tanks due to insufficient /delayed onset of monsoon	NA				

Condition	Suggested Contingency measures				
	Major Farming situation <sup>f</sup>	Normal Crop/cropping system <sup>g</sup>	Change in crop/cropping system <sup>h</sup>	Agronomic measures <sup>i</sup>	Remarks on Implementation <sup>j</sup>
Insufficient groundwater recharge due to low rainfall	NA				

## 2.2 Unusual rains (untimely, unseasonal etc) (for both rainfed and irrigated situations)

Condition	Suggested contingency measure			
Continuous high rainfall in a short span leading to water logging	Vegetative stage <sup>k</sup>	Flowering stage <sup>l</sup>	Crop maturity stage <sup>m</sup>	Post harvest <sup>n</sup>
Rice	Bunding to store water in medium lands	Water restoring	Draining out	Shifting to

		in medium lands		safer place and drying
Pigeon pea	Draining out since cultivated in uplands which is undulated, self drainage occur therefore	Draining out application of plant protection chemicals	Provided drainage	Shifting to safer place and drying
Maize	Provided drainage	Provided drainage	Provided drainage	Shifting to safer place and drying
Wheat	Draining out	Draining out	Draining out	Shifting to safer place and drying
Groundnut	Draining out followed by hoeing the field	Draining out followed by hoeing	Draining out	Shifting to safer place and drying
<b>Horticulture</b>				
Potato	Provided drainage and plant protection		Drainage and diggiout	Shifting to safer place and drying
Brinjal	Provided drainage and plant protection	Provided drainage and Plant Protection	Provided drainage	Shifting to safer place and drying
Tomato	Provided drainage and plant protection	Provided drainage and Plant Protection	Drainage and Picking up fruits	Value addition
Cucurbit	Provided drainage and plant protection	Proper drainage	Drainage and Picking up fruits	Value addition
Chillies	Provided drainage and plant protection	Drainage and Plant protection	Drainage and Picking up fruits	Storage or drying or value addition
<b>Heavy rainfall with high speed</b>				

<b>winds in a short span<sup>2</sup></b>				
Rice	Water restoring in medium land		Immediate drainage and harvesting crop	Drying on safer place
Pigeonpea	Draining and earthing up	Drainage and erecting the Plants	Drainage and erecting the Plants	Drying safer place
Maize	Draining and earthing up	Drainage and erecting the Plants	Drainage and harvesting at physiological maturity	Piling at safe place
Wheat	Drainage	Drainage	Drainage	Drying safer place
Pulses (Rabi)	Drainage and hoeing	Drainage and hoeing	Drainage	Drying safer place
<b>Horticulture</b>				
Potato	Drainage and Plant protection	Drainage and Plant protection	Drainage	Shade drying and store in cold storage
Brinjal	Drainage and Plant protection	Drainage and Plant protection	Drainage	Sail
Tomato	Drainage and Plant protection	Drainage and Plant protection	Drainage	Value addition
Cucurbits	Drainage and Plant protection	Drainage and Plant protection	Drainage	Value addition
Chillies	Drainage and Plant protection	Drainage and Plant protection	Drainage	Preservation and drying
<b>Outbreak of pests and diseases due to unseasonal rains</b>				
Rice	Need based plant protection measure IPM or IDPM for field crop			
Pigeonpea	Need based plant protection measure IPM or IDPM for field crop			
Maize	Need based plant protection measure IPM or IDPM for field crop			

Pulses (Rabi)	Need based plant protection measure IPM or IDPM for field crop			
Rye – Mustard	Need based plant protection measure IPM or IDPM for field crop			
<b>Horticulture</b>				

### 2.3 Floods

Condition	Suggested contingency measure <sup>o</sup>			
	Seedling / nursery stage	Vegetative stage	Reproductive stage	At harvest
Transient water logging/ partial inundation <sup>1</sup>	NA			
<b>Horticulture</b>	NA			
Continuous submergence for more than 2 days <sup>2</sup>				
<b>Horticulture</b>	NA			
Sea water intrusion <sup>3</sup>	NA			

### 2.4 Extreme events: Heat wave / Cold wave/Frost/ Hailstorm /Cyclone

Extreme event type	Suggested contingency measure <sup>r</sup>			
	Seedling / nursery stage	Vegetative stage	Reproductive stage	At harvest
<b>Heat Wave<sup>p</sup> (loo)</b>				
Wheat	-	-	Maintaining soil moisture by irrigation	
Pigeonpea	-	-	Maintaining soil moisture by irrigation	
Rye	-	-	Maintaining soil moisture by irrigation	
Lentil and Chickpea	-	-	Maintaining soil moisture by irrigation	
Rice	Maintaining soil moisture	-	Maintaining soil moisture by irrigation	
<b>Horticulture</b>				

Okra	Moisture management	Maintaining wind breaks and soil moisture	Wind breaks and irrigation	
Chillies	Moisture management	Maintaining wind breaks and soil moisture	Wind breaks and irrigation	
Cucurbits	Moisture management			
<b>Cold wave<sup>a</sup></b>				
Pigeonpea	N.A.	Spray of fungicide and IPM	Before initiation of flower use of IPM Which include fungicide + insecticide	
Pea		Spray of fungicide and IPM	Watering of plant	
Lentil		Spray of fungicide and IPM	Smoking at 3 to 4 am in the morning	
Rabi Maize				
<b>Horticulture</b>				
Potato	-	Proper moisture mgt.		
Tomato	-	Proper moisture mgt.		
Chilli	-	Proper moisture mgt.		
<b>Frost</b>				
<b>Horticulture</b>	NA			
<b>Hailstorm</b>	NA			
<b>Horticulture</b>	NA			
<b>Cyclone</b>	NA			
<b>Horticulture</b>	NA			

## 2.5 Contingent strategies for Livestock, Poultry & Fisheries

### 2.5.1 Livestock

	Suggested contingency measures		
	Before the event <sup>s</sup>	During the event	After the event
<b>Drought</b>			
Feed and fodder availability			
Drinking water			
Health and disease management			
<b>Floods</b>			
Feed and fodder availability			
Drinking water			
Health and disease management			
<b>Cyclone</b>			
Feed and fodder availability			
Drinking water			
Health and disease management			
<b>Heat wave and cold wave</b>			
Shelter/environment management			
Health and disease management			

<sup>s</sup> based on forewarning wherever available



## 2.5.2

## Poultry

	Suggested contingency measures			Convergence/linkages with ongoing programs, if any
	Before the event <sup>a</sup>	During the event	After the event	
<b>Drought</b>				
Shortage of feed ingredients				
Drinking water				
Health and disease management				
<b>Floods</b>				
Shortage of feed ingredients				
Drinking water				
Health and disease management				
<b>Cyclone</b>				
Shortage of feed ingredients				
Drinking water				
Health and disease management				
<b>Heat wave and cold wave</b>				
Shelter/environment management				
Health and disease management				

## 2.5.3

## Fisheries/ Aquaculture

	Suggested contingency measures		
	Before the event <sup>a</sup>	During the event	After the event
<b>1) Drought</b>			
<b>A. Capture</b>			
Marine			
Inland			
(i) Shallow water depth due to insufficient rains/inflow			
(ii) Changes in water quality			
(iii) Any other			
<b>B. Aquaculture</b>			
(i) Shallow water in ponds due to insufficient rains/inflow			
(ii) Impact of salt load build up in ponds / change in water quality			
(iii) Any other			
<b>2) Floods</b>			
<b>A. Capture</b>			
Marine			
Inland			
(i) Average compensation paid due to loss of human life			
(ii) No. of boats / nets/damaged			

(iii) No.of houses damaged			
(iv) Loss of stock			
(v) Changes in water quality			
(vi) Health and diseases			
<b>B. Aquaculture</b>			
(i) Inundation with flood water			
(ii) Water contamination and changes in water quality			
(iii) Health and diseases			
(iv) Loss of stock and inputs (feed, chemicals etc)			
(v) Infrastructure damage (pumps, aerators, huts etc)			
(vi) Any other			
<b>3. Cyclone / Tsunami</b>			
A. Capture			
Marine			
(i) Average compensation paid due to loss of fishermen lives			
(ii) Avg. no. of boats / nets/damaged			
(iii) Avg. no. of houses damaged			
Inland			
B. Aquaculture			

(i) Overflow / flooding of ponds			
(ii) Changes in water quality (fresh water / brackish water ratio)			
(iii) Health and diseases			
(iv) Loss of stock and inputs (feed, chemicals etc)			
(v) Infrastructure damage (pumps, aerators, shelters/huts etc)			
(vi) Any other			
<b>4. Heat wave and cold wave</b>			
<b>A. Capture</b>			
Marine			
Inland			
<b>B. Aquaculture</b>			
(i) Changes in pond environment (water quality)			
(ii) Health and Disease management			
(iii) Any other			

<sup>a</sup> based on forewarning wherever available