



E-ISSN: 2320-7078

P-ISSN: 2349-6800

JEZS 2019; 7(6): 15-18

© 2019 JEZS

Received: 09-09-2019

Accepted: 13-10-2019

**Amit Paschapur**ICAR-Vivekananda Parvatiya  
Krishi Anusandhana Sansthan,  
Almora, Uttarakhand, India**ARNS Subbanna**ICAR-Vivekananda Parvatiya  
Krishi Anusandhana Sansthan,  
Almora, Uttarakhand, India**Ashish Kumar Singh**ICAR-Vivekananda Parvatiya  
Krishi Anusandhana Sansthan,  
Almora, Uttarakhand, India**Jeevan B**ICAR-Vivekananda Parvatiya  
Krishi Anusandhana Sansthan,  
Almora, Uttarakhand, India**J Stanley**ICAR-Vivekananda Parvatiya  
Krishi Anusandhana Sansthan,  
Almora, Uttarakhand, India**KK Mishra**ICAR-Vivekananda Parvatiya  
Krishi Anusandhana Sansthan,  
Almora, Uttarakhand, India**Corresponding Author:****Amit Paschapur**ICAR-Vivekananda Parvatiya  
Krishi Anusandhana Sansthan,  
Almora, Uttarakhand, India

## Baseline susceptibility of major sucking pests of hill crops in north western Himalayan region

**Amit Paschapur, ARNS Subbanna, Ashish Kumar Singh, Jeevan B, J Stanley and KK Mishra**

**Abstract**

Greenhouse whitefly (*Trialeurodes vaporariorum*), green peach aphid (*Myzus persicae*) and cabbage aphid (*Brevicoryne brassicae*) are the three major sucking pests that infect hill crops in North Western Himalayan region and cause severe yield losses. Chemical management tactics are the widely practiced methods for management of these pests in the locality. However, knowledge about the efficacy of different pesticides against these native populations is lacking. In view of this, seven insecticides against whitefly and six against aphids were tested for their bio-efficacy under laboratory conditions. The results showed that, field populations of greenhouse whitefly were highly susceptible to three insecticides; Thiomethaxam > Imidacloprid > Pymetrozine with LC<sub>50</sub> values 12.30, 18.62 and 22.38 ppm respectively. Whereas, LC<sub>50</sub> values of botanical insecticides NSKE and nimbecidine were very high; 524.81ppm and 4365.16ppm respectively, indicating their non-suitability against greenhouse whitefly management. The susceptibility for *B. brassicae*, a major pest of crucifers is in the following order; Thiomethaxam > Imidacloprid > Indoxacarb > Pymetrozine > Nimbecidine > NSKE and Thiomethaxam was the most toxic insecticide with LC<sub>50</sub> and LC<sub>90</sub> values as low as 0.024ppm and 0.25ppm respectively. Green peach aphid (*M. persicae*) was highly susceptible to Thiomethaxam and was followed by Pymetrozine, indoxacarb, Imidacloprid, Nimbecidine and NSKE respectively. The LC<sub>50</sub> values of Thiomethaxam and Pymetrozine were at par with a recorded value of 2.54ppm and 2.57ppm, respectively. The three sucking pests under study are well known for their resistance development mechanisms against several groups of insecticides. But, the level of susceptibility that was recorded in our study indicates presence of susceptible population of these sucking pests in Indian Himalayan region that could help to maintain a refugee or buffer populations against development of insecticide resistant strains.

**Keywords:** Greenhouse whitefly, green peach aphid, cabbage aphid, susceptibility, lethal concentrations, north western Indian Himalayas

**1. Introduction**

Greenhouse whitefly (*Trialeurodes vaporariorum*), green peach aphid (*Myzus persicae*) and cabbage aphid (*Brevicoryne brassicae*) are the three most important sucking pests in hill agriculture (Kim *et al.*, 1986; Hill, 1987) <sup>[1, 2]</sup>, wherein, the former two being polyphagous pests infecting large number of crops in both open field and greenhouse conditions, while the latter being a stenophagous in habit, infecting cruciferous crops. The sucking pests cause damage in three ways *viz.*, reduction in vitality due to loss in cell sap; inhibition in photosynthesis due to sooty mould and transmission of lethal viral diseases (Johnson *et al.*, 1992; Blackman and Eastop, 2000; Castillo *et al.*, 2011; Cavalieri *et al.*, 2014) <sup>[3, 4, 5, 6]</sup>. The well-evolved reproduction (parthenogenesis in aphids and arrhenotokous parthenogenetic reproduction in whiteflies) and survival strategies (pseudopupa/ prepupal stage in whiteflies) in these pests make them superior even under adverse climatic conditions.

Although, several management strategies are available to suppress the pests on crops; efficacy, availability and cost of operations are not comparable with chemical control measures. Besides these are the last resorts to suppress any type of pest populations in order to avoid economic damage. Therefore, integrated pest management systems designed to suppress sucking pests on various crops still include insecticides as an important component (Liu *et al.*, 1993; Toscano *et al.*, 1998) <sup>[7, 8]</sup>. However, indiscriminate use of insecticides leads to the development of resistance to various organophosphates, carbamates and synthetic pyrethroids (Elhag and Horn, 1983; Buitrago *et al.*, 1994; Zheng and Gao, 1995; Denholm and Jespersen, 1998) <sup>[9, 10, 11, 12]</sup>. Intensive research has been carried out in recent years for evaluating insecticides with novel modes of action against sucking pests due to low pace of resistance

development to these insecticides. Although, resistance and resurgence in sucking pests is becoming widespread due to their indiscriminate use (Nauen and Denholm, 2005) [13]. In spite of number of publications re-establishing toxicity for a range of compounds that include imidacloprid and other novel insecticides; the lack of up-to-date resistance and susceptibility monitoring data for *T. vaporariorum*, *B. brassicae* and *M. persicae* is a subject of concern (Bi *et al.*, 2002; Wang *et al.*, 2003) [14, 15].

In hill agriculture, farmers either use botanicals and/or safer novel insecticide molecules, which can manage the insect pests at lower dosages and in turn have least residues in the soil, agriculture produce and environment. Taking this into consideration; seven insecticide molecules popularly used by farmers in the locality to manage the above three sucking pests were selected and their acute toxicities were calculated for field populations of aphids and whiteflies collected from ICAR-VPKAS, experimental farm, Hawalbagh (29.63°N and 79.63°E, 1250 amsl) representing North Western Himalayan region. The median lethal concentrations for all the insecticides were calculated and the most effective among them was recommended to farmers for managing respective insect pests.

## 2. Materials and Methods

**A. Insects:** Adults of greenhouse whitefly (*Trialeurodes vaporariorum*), green peach aphid (*Myzus persicae*) and cabbage aphid (*Brevicoryne brassicae*) infecting tomato, capsicum and cauliflower respectively, were collected from open fields or polyhouses at ICAR-VPKAS, experimental farm, Hawalbagh (29.63°N and 79.63°E, 1250 amsl). The field populations, after confirming the identity, were tested directly without rearing in the laboratory. Apterous adults of aphids and winged adults of whiteflies were used for the study.

**B. Insecticides:** Seven most commonly used insecticide formulations to manage sucking pests in Uttarakhand hills were selected for the study; Acetamiprid (Prime @ 20% SP; Crop chemicals), Imidacloprid (Sacdor @ 17.8%SL; Shivalik crop sciences), Thiomethaxam (Dxstar @ 25%WG; Nagarjun Agrichemicals Ltd), Indoxacarb (King carb @ 14.5%SC; Parijat), Pymetrozine (Simca @ 50%WG; Adama), NSKE (Vanguard 1500@ 1500ppm azadirachtin EC; Agriland) and Nimbicidine (Nimbicidine 0.03%EC @ 300ppm azadirachtin;

Stanes). All the insecticides were purchased from local dealers. All these seven insecticides were tested against whiteflies and six insecticides except acetamiprid were tested against aphids.

**C. Bioassay:** Leaf dip bioassay was conducted for all the insecticides with seven different concentrations ranging from 0.05ppm to 15ppm for aphids and 0.1ppm to 100ppm for whiteflies for each insecticide and each treatment was replicated three times. A control treatment was set up with double distilled water as treatment at each time. Respective host leaf discs of approximately 7.5cm diameter were dipped in insecticide solution for 1 minute and air dried for 5-10 minutes at room temperature (25±2°C). The treated leaves were placed in a Petri dish (9cm diameter) with adaxial side of the leaf facing upwards and with a moist filter paper beneath to maintain humidity. Ten numbers of insects were released with a camel brush on each leaf and care was taken to avoid any physical damage while transferring. The lid with minute perforations was closed and plates were incubated at 25±2°C for 96 hours.

**3. Data analysis:** Mortality of the adults was recorded at every 24 hours of insecticide exposure and was corrected by Abbott's (1925) [16] formula. Insects showing no movements after gentle touch with the camel brush were considered dead. Thus obtained data was subjected to probit analysis (Finney, 1971) [17] using the software package PoloPlus (LeOra Software 2013)

## 4. Results

**Susceptibility of *T. vaporariorum* to insecticides:** Field populations of greenhouse whitefly were highly susceptible to three insecticides; Thiomethaxam> Imidacloprid> Pymetrozine. The LC<sub>50</sub> values for botanical insecticides NSKE and nimbicidine were very high, 524.81ppm and 4365.16ppm respectively. Whereas, LC<sub>50</sub> values for acetamiprid, the most commonly used insecticide in North Western Himalayan region for management of greenhouse whiteflies was 38.02ppm. Indoxacarb, an insecticide used by farmers to manage both American tomato pin worm (*Tuta absoluta*) and *T. vaporariorum* in polyhouses showed LC<sub>50</sub> value of 52.48ppm. The details of the probit analysis along with LC<sub>50</sub> and LC<sub>90</sub> are mentioned in table 1.

**Table 1:** Toxicity of selected insecticides against field populations of *Trialeurodes vaporariorum* collected from ICAR-VPKAS, Experimental farm, Hawalbagh, Almora, Uttarakhand

Insecticide	Linear equation (Y= ax+b)	Slope±SE	$\chi^2$	Df	LC <sub>50</sub> (in ppm)	LC <sub>90</sub> (in ppm)
Acetamiprid	Y=2.51x+1.02	2.51±0.77	0.76	6	38.02	123.03
Imidacloprid	Y =2.73x+1.52	2.73±0.51	0.90	6	18.62	54.92
Thiomethaxam	Y =2.85x+1.87	2.85±0.50	0.91	6	12.30	35.48
Indoxacarb	Y =2.36x+0.93	2.36±0.72	0.76	6	52.48	467.74
Pymetrozine	Y =2.43x+1.71	2.43±0.44	0.90	6	22.38	75.86
NSKE	Y =1.70x+0.38	1.70±0.74	0.62	6	524.81	2951.21
Nimbicidine	Y =1.35x+0.08	1.35±0.80	0.47	6	4365.16	38904.51

**Susceptibility of *B. brassicae* to insecticides:** *B. brassicae* a major aphid pest of crucifers in North western Himalayan region was highly susceptible to all the six tested insecticides including botanical insecticides, NSKE and nimbicidine. The baseline susceptibility decreased in the following order;

Thiomethaxam>Imidacloprid>Indoxacarb>Pymetrozine> Nimbicidine> NSKE. Thiomethaxam was the most toxic insecticide with LC<sub>50</sub> and LC<sub>90</sub> values as low as 0.024ppm and 0.25ppm respectively. The details of the probit analysis along with LC<sub>50</sub> and LC<sub>90</sub> are mentioned in table 2.

**Table 2:** Toxicity of selected insecticides against field populations of *Brevicoryne brassicae* collected from ICAR-VPKAS, Experimental farm, Hawalbagh, Almora, Uttarakhand

Insecticide	Linear equation (Y= ax+b)	Slope±SE	$\chi^2$	Df	LC <sub>50</sub> (in ppm)	LC <sub>90</sub> (in ppm)
Imidacloprid	Y=1.13x+5.12	1.13±0.08	0.96	6	0.78	10.47
Thiomethaxam	Y=1.26x+7.05	1.26±0.23	0.82	6	0.024	0.25
Indoxacarb	Y=1.43x+4.99	1.43±0.32	0.76	6	1.02	7.94
Pymetrozine	Y=1.29x+5.36	1.29±0.26	0.79	6	1.9	8.91
NSKE	Y=1.88x+4.03	1.00±0.37	0.81	6	3.31	15.49
Nimbecidine	Y=1.42x+5.44	1.42±0.22	0.87	6	2.04	13.89

**Susceptibility of *M. persicae* to insecticides:** *M. persicae* a major polyphagous aphid pest infecting more than 100 crops in hill region showed almost uniform susceptibility to all the six insecticides tested. Thiomethaxam being the most toxic insecticide followed by Pymetrozine, indoaxcarb, Imidacloprid, Nimbecidine and NSKE. The LC<sub>50</sub> values for Thiomethaxam and Pymetrozine were on par with each other (2.54ppm and 2.57ppm) respectively. The details of the probit analysis along with LC<sub>50</sub> and LC<sub>90</sub> are mentioned in table 3.

**Table 3:** Toxicity of selected insecticides against field populations of *Myzus persicae* collected from ICAR-VPKAS, Experimental farm, Hawalbagh, Almora, Uttarakhand

Insecticide	Linear equation	Slope±SE	$\chi^2$	Df	LC <sub>50</sub>	LC <sub>90</sub>
Imidacloprid	y=2.69x+2.71	2.69±0.37	0.89	6	7.07	21.38
Thiomethaxam	Y=2.93x+3.81	2.93±0.36	0.91	6	2.54	6.92
Indoxacarb	y=2.37x+3.1	2.37±0.22	0.95	6	6.31	21.87
Pymetrozine	Y=2.95x+3.79	2.95±0.35	0.92	6	2.57	6.92
NSKE	y=2.46x+2.58	2.46±0.38	0.87	6	9.55	31.62
Nimbecidine	Y=2.28x+2.98	2.28±0.21	0.95	6	7.58	28.18

All in all the bioassay results against three major sucking pests of North Western Himalayan region showed that thiomethaxam (Dxstar @ 25%WG; Nagarjun Agrichemicals Ltd) was the most toxic among the seven insecticides tested against both whitefly and two aphid species. The botanical formulations NSKE (Vanguard 1500 @ 1500ppm azadirachtin EC; Agriland) was least toxic to aphid species and Nimbecidine (Nimbecidine 0.03%EC @ 300ppm azadirachtin; Stanes) was least toxic to greenhouse whitefly. All these mortalities were recorded at 24 hours after the treatment, which indicates that the aphid and whitefly populations were highly susceptible to the major insecticides available in the market and adapted by the farmers.

## 5. Discussion

The three sucking pests under study are well examined for their dynamic physiology of developing resistance to major conventional insecticides like organophosphates, carbamates and pyrethroids (Zou and Zheng, 1988; Omer *et al.*, 1992)<sup>[18]</sup><sup>[19]</sup>. *Myzus persicae* is known for overproduction of insecticide-detoxifying carboxylesterases, resulting from structural amplification of genes encoding these enzymes (Foster *et al.*, 2000)<sup>[20]</sup>. Ahmad and Aslam, (2005)<sup>[21]</sup> showed that *Brevicoryne brassicae* populations from Pakistan were resistant to major organophosphates and pyrethroids. Whereas, enhanced production of P450 CYP6CM1 in *T. vaporariorum* has shown resistant to all conventional insecticides along with few neonicotinoids (Imidacloprid) and even pymetrozine (Karatolos *et al.*, 2010)<sup>[22]</sup> indicating the scenario parallel to *Bemisia tabaci*. Among the reports of insecticide resistance development, those populations collected from Indian Himalayas are known to be highly susceptible to novel insecticides (Sood *et al.*, 2006)<sup>[23]</sup> and the present research findings are in agreement with this.

However, botanicals were not of much importance in whitefly management due to higher LC<sub>50</sub> values recorded for NSKE and Nimbecidine respectively. Whereas, the three neonicotinoid insecticides, Imidacloprid, acetamiprid and thiomethaxam were highly toxic and are a welcome new class of chemistry for aphid and whitefly control. Indoxacarb a novel molecule recommended for coleopteran and lepidopteran pest management had significant toxicity to sucking pests (Homoptera) which shows its broad activity spectrum. Pymetrozine a specialised sucking pest control molecule proved to be a promising insecticide molecule and recorded lower LC<sub>50</sub> values that are at par with neonicotinoids.

In conclusion, the insect pest populations of Indian Himalayas are least exposed to pesticide treatments and are susceptible to most of novel molecules. This status of susceptibility has to be maintained by discriminate and wise use of available insecticides to avoid development of resistant populations.

## 6. Acknowledgement

The authors are thankful to The Director, ICAR-VPKAS, Almora for providing all facilities to carry out the research work and for his valuable guidance during the course of work.

## 7. References

- Kim IS, Hwang EY, Kim JH, Lee MH. Studies on host plants, development and distribution within plants of the greenhouse whitefly, *Trialeurodes vaporariorum* (Westwood). Korean Journal of Plant Protection. 1986; 25(4):201-207.
- Hill DS. Agricultural insect pests of temperate regions and their control. Cambridge University Press London, 1987, 659.
- Johnson MW, Caprio LC, Coughlin JA, Tabashnik BE, Rosenheim JA, Welter SC *et al.* Effect of *Trialeurodes vaporariorum* (Homoptera: Aleyrodidae) on yield of fresh market tomatoes. Journal of Economic Entomology. 1992; 85(6):2370-2376.
- Blackman RL, Eastop VF. Aphids on the world's crops: an identification and information guide (No. Ed. 2). John Wiley & Sons Ltd, 2000.
- Castillo NJ, Olive EF, Campos SS. Emerging virus disease transmitted by whiteflies. Annual Review of Phytopathology. 2011; 49:219-248.
- Cavaliere V, Mangli A, Tiberini A, Tomassoli L, Rapisarda C. Rapid identification of *Trialeurodes vaporariorum*, *Bemisia tabaci* (MEAM1 and MED) and tomato-infecting criniviruses in whiteflies and in tomato leaves by real-time reverse transcription-PCR assay. Bulletin of Insectology. 2014; 67(2):219-225.
- Liu TX, Oetting RD, Buntin GD. Population dynamics and distribution of *Trialeurodes vaporariorum* and *Bemisia tabaci* (Homoptera: Aleyrodidae) on poinsettia following applications of three chemical insecticides.

- Journal of Entomological Science. 1993; 28(1):126-135.
8. Toscano NC, Prabhaker N, Zhou S, Ballmer G. Toxicity of Applaud and Knack against silverleaf whiteflies from southern California: Implications for susceptibility monitoring. In: Dugger, P., Richer, D. (Eds.), Proceedings of Beltwide Cotton Research Conference, National Cotton Council of America, Memphis, 1998, 1093-1095.
  9. Elhag EA, Horn DJ. Resistance of greenhouse whitefly (Homoptera: Aleyrodidae) to insecticides in selected Ohio greenhouses. Journal of Economic Entomology. 1983; 76(4):945-948.
  10. Buitrago NA, Cardona C, Acosta A. Niveles de resistencia a insecticidas en *Trialeurodes vaporariorum* (Westwood) (Homoptera: Aleyrodidae), plaga del frijol común. Revista Colombiana de Entomología. 1994; 20(2):109-114.
  11. Zheng BZ, Gao XW. Monitoring insecticide resistance in greenhouse whitefly adults in Beijing, China, 1991-1995. Resistant Pest Management. 1995; 7(2):23-24.
  12. Denholm I, Jespersen JB. Insecticide resistance management in Europe: recent developments and prospects. Pesticide Science. 1998; 52(2):193-195.
  13. Nauen R, Denholm I. Resistance of insect pests to neonicotinoid insecticides: current status and future prospects. Archives of Insect Biochemistry and Physiology. 2005; 58:200-215.
  14. Bi JL, Toscano NC, Ballmer GR. Greenhouse and field evaluation of six novel insecticides against the greenhouse whitefly *Trialeurodes vaporariorum* on strawberries. Crop Protection. 2002; 21:49-55.
  15. Wang KY, Kong XB, Jiang XY, Yi MQ, Liu TX. Susceptibility of immature and adult stages of *Trialeurodes vaporariorum* (Homoptera, Aleyrodidae) to selected insecticides. Journal of Applied Entomology. 2003; 127:527-533.
  16. Abbott WS. Abbott's formula. Journal of Economic Entomology. 1925; 18:267-268.
  17. Finney DJ. Probit analysis, Cambridge University Press. Cambridge, UK, 1971.
  18. Zou YQ, Zheng BZ. The toxicity of some insecticides to greenhouse whitefly (*Trialeurodes vaporariorum* Westwood) and monitoring of resistance. Acta Phytosociologica Sinica. 1988; 15:277-281.
  19. Omer AD, Leigh TF, Granett J. Insecticide resistance in field populations of greenhouse whitefly (Homoptera: Aleyrodidae) in the San Joaquin Valley (California) cotton cropping system. Journal of Economic Entomology. 1992; 85(1):21-27.
  20. Foster SP, Denholm I, Devonshire AL. The ups and downs of insecticide resistance in peach-potato aphids (*Myzus persicae*) in the UK. Crop Protection. 2000; 19:873-879.
  21. Ahmad M, Aslam M. Resistance of cabbage aphid, *Brevicoryne brassicae* (Linnaeus) to endosulfan, organophosphates and synthetic pyrethroids. Pakistan Journal of Zoology. 2005; 37(4), 293.
  22. Karatolos N, Denholm I, Williamson M, Nauen R, Gorman K. Incidence and characterisation of resistance to neonicotinoid insecticides and pymetrozine in the greenhouse whitefly, *Trialeurodes vaporariorum* Westwood (Homoptera: Aleyrodidae). Pest Management Science. 2010; 66(12):1304-1307.
  23. Sood S, Sood AK, Verma KS. Determination of baseline toxicity of some insecticides to greenhouse whitefly, *Trialeurodes vaporariorum* (Westwood) population from North-Western Indian Himalayas. Pest Management in Horticultural Ecosystems. 2006; 12(1):67-70.