



हर कदम, हर डगर
किसानों का हमसफर
भारतीय कृषि अनुसंधान परिषद

AgriSearch with a human touch

**LAND RESOURCE INVENTORY AND SOCIO-ECONOMIC STATUS OF
FARM HOUSEHOLDS FOR WATERSHED PLANNING AND
DEVELOPMENT**

CHINCHOLI KHURD-1 (4D5C2A1c) MICROWATERSHED

Aland Taluk, Gulbarga District, Karnataka

Karnataka Watershed Development Project – II

SUJALA – III

World Bank funded Project



ICAR – NATIONAL BUREAU OF SOIL SURVEY AND LAND USE PLANNING



ICAR - NBSS & LUP



**WATERSHED DEVELOPMENT DEPARTMENT
GOVT. OF KARNATAKA, BANGALORE**



About ICAR - NBSS&LUP

The ICAR-National Bureau of Soil Survey and Land Use Planning (ICAR-NBSS&LUP), Nagpur, a premier Institute of the Indian Council of Agricultural Research (ICAR), was set up during 1976 with the objective to prepare soil resource maps at national, state and district levels and to provide research inputs in soil resource mapping and its applications, land evaluation, land use planning, land resource management, and database management using GIS for optimising land use on different kinds of soils in the country.

The Bureau has been engaged in carrying out soil resource survey, agro-ecological and soil degradation mapping at the country, state and district levels for qualitative assessment and monitoring the soil health towards viable land use planning. The research activities have resulted in identifying the soil potentials and problems, and the various applications of the soil surveys with the ultimate objective of sustainable agricultural development. The Bureau has the mandate to correlate and classify soils of the country and maintain a National Register of all the established soil series. The Institute is also imparting in-service training to staff of the soil survey agencies in the area of soil survey, land evaluation and soil survey interpretations for land use planning. The Bureau in collaboration with Panjabrao Krishi Vidyapeeth, Akola is running post-graduate teaching and research programme in land resource management, leading to M.Sc. and Ph.D. degrees.

Citation: Rajendra Hegde, Ramesh Kumar, S.C., K.V. Niranjana, S. Srinivas, M.Lalitha, B.A. Dhanorkar, R.S. Reddy and S.K. Singh (2019). "Land Resource Inventory and Socio-Economic Status of Farm Households for Watershed Planning and Development of Chincholi Khurd-1 (4D5C2A1c) Microwatershed, Aland Taluk, Gulbarga District, Karnataka", ICAR-NBSS&LUP Sujala MWS Publ.7, ICAR – NBSS & LUP, RC, Bangalore. p.91 & 28.

TO OBTAIN COPIES,

Please write to:

Director, ICAR - NBSS & LUP,

Amaravati Road, NAGPUR - 440 033, India

Phone : (0712) 2500386, 2500664, 2500545 (O)

Telefax : 0712-2522534

E-Mail : director@nbsslup.ernet.in

Website URL : nbsslup.in

Or

Head, Regional Centre, ICAR - NBSS&LUP, Hebbal, Bangalore - 560 024

Phone : (080) 23412242, 23510350 (O)

Telefax : 080-23510350

E-Mail : nbssrcb@gmail.com



**LAND RESOURCE INVENTORY AND SOCIO-ECONOMIC
STATUS OF FARM HOUSEHOLDS FOR WATERSHED
PLANNING AND DEVELOPMENT**

CHINCHOLI KHURD-1 (4D5C2A1c) MICROWATERSHED

Aland Taluk, Gulbarga District, Karnataka

Karnataka Watershed Development Project – II

Sujala-III

World Bank funded Project



**ICAR – NATIONAL BUREAU OF SOIL SURVEY AND LAND USE
PLANNING**



ICAR - NBSS & LUP



**WATERSHED DEVELOPMENT DEPARTMENT, GOVT. OF
KARNATAKA, BANGALORE**



PREFACE

In Karnataka, as in other Indian States, the livelihoods of rural people are intertwined with farming pursuits. The challenges in agriculture are seriously threatening the livelihood of a large number of farmers as they have been practicing farming in contextual factors beyond their control. Climatic factors are the most important ones and have become much more significant in recent times due to rapid climate changes induced by intensive anthropogenic activities affecting our ecosystem in multiple ways. Climate change has become the reality, it is happening and efforts to evolve and demonstrate climate resilient technologies have become essential. Due to the already over stressed scenario of agrarian sector, the climate change is resulting in manifold increase in the complexities, pushing the rural mass to face more and more unpredictable situations. The rising temperatures and unpredictable rainfall patterns are going to test seriously the informed decisions farmers have to make in order to survive in farming and sustain their livelihood.

It is generally recognized that impacts of climate change shall not be uniform across the globe. It is said that impact of climate change is more severe in South Asia. Based on the analysis of meteorological data, it is predicted that in India, there will be upward trend in mean temperature, downward trend in relative humidity, annual rainfall and number of wet days in a year. Also, in general, phenomena like erratic monsoon, spread of tropical diseases, rise in sea levels, changes in availability of fresh water, frequent floods, droughts, heat waves, storms and hurricanes are predicted. Each one of these adverse situations are already being experienced in various parts of India and also at the global level. Decline in agricultural productivity of small and marginal farmers becoming more vulnerable is already witnessed.

In Karnataka, more than 60 per cent of the population live in rural areas and depend on agriculture and allied activities for their livelihood. Though the state has achieved significant progress in increasing the yield of many crops, there is tremendous pressure on the land resources due to the growing and competing demands of various land uses. This is reflected in the alarming rate of land degradation observed. Already more than 50 per cent of the area is affected by various forms of degradation. If this trend continues, the sustainability of the fragile ecosystem will be badly affected. The adverse effects of change in the climatic factors are putting additional stress on the land resources and the farmers dependent on this.

The natural resources (land, water and vegetation) of the state need adequate and constant care and management, backed by site-specific technological interventions and investments particularly by the government. Detailed database pertaining to the nature of

the land resources, their constraints, inherent potentials and suitability for various land based rural enterprises, crops and other uses is a prerequisite for preparing location-specific action plans, which are in tune with the inherent capability of the resources. Any effort to evolve climate resilient technologies has to be based on the baseline scientific database. Then only one can expect effective implementation of climate resilient technologies, monitor the progress, make essential review of the strategy, and finally evaluate the effectiveness of the implemented programs. The information available at present on the land resources of the state are of general nature and useful only for general purpose planning. Since the need of the hour is to have site-specific information suitable for farm level planning and detailed characterization and delineation of the existing land resources of an area into similar management units is the only option.

ICAR-NBSS&LUP, Regional Centre, Bangalore has taken up a project sponsored by the Karnataka Watershed Development Project-II, (Sujala-III), Government of Karnataka funded by the World Bank under Component -1 Land Resource Inventory. This study was taken up to demonstrate the utility of such a database in reviewing, monitoring and evaluating all the land based watershed development programs on a scientific footing. To meet the requirements of various land use planners at grassroots level, the present study on “Land Resource Inventory and Socio-Economic Status of Farm Households for Watershed Planning and Development of Chincholi Khurd-1 Microwatershed, Aland Taluk, Gulbarga District, Karnataka” for integrated development was taken up in collaboration with the State Agricultural Universities, IISC, KRSRAC, KSNDMC as Consortia partners. The project provides detailed land resource information at cadastral level (1:7920 scale) for all the plots and socio-economic status of farm households covering thirty per cent farmers randomly selected representing landed and landless class of farmers in the micro-watershed. The project report with the accompanying maps for the microwatershed will provide required detailed database for evolving effective land use plan, alternative land use options and conservation plans for the planners, administrators, agricultural extension personnel, KVK officials, developmental departments and other land users to manage the land resources in a sustainable manner.

It is hoped that this database will be useful to the planners, administrators and developmental agencies working in the area in not only for formulating location specific developmental schemes but also for their effective monitoring at the village/watershed level.

Nagpur

Date: 10.05..2016

S.K. SINGH

Director, ICAR - NBSS&LUP, Nagpur

Contributors

Dr. RajendraHegde Principal Scientist, Head & Project Leader, Sujala-III Project ICAR-NBSS&LUP, Regional Centre Bangalore	Dr. S.K.Singh Director, ICAR-NBSS&LUP Coordinator, Sujala-III Project Nagpur
Soil Survey, Mapping & Report Preparation	
Dr. B.A. Dhanorkar	Sh. R.S. Reddy
Dr. K.V. Niranjana	Sh. Nagendra, B.R.
	Smt. Chaitra, S.P.
Field Work	
Sh. C.BacheGowda	Sh. Mahesha, D.B.
Sh. Somashekar	Sh. Ashok S. Sindagi
Sh. VenkataGiriyappa	Sh. Veerabhadrapa
Sh. M. Jayaramaiah	Sh. Anand
Sh. Paramesha, K.	Sh. Arun N Kambar
	Sh. Shankarappa, K.
	Sh. Kamalesh K. Avate
	Sh. Sharan Kumar Huppar
	Sh. Yogesh, H.N.
	Sh. Kalaveerachari R. Kammar
GIS Work	
Dr. S.Srinivas	Sh. A.G.Devendra Prasad
Dr. M.Ramesh	Sh. Prakashanaik, M.K.
Sh. D.H.Venkatesh	Sh. AbhijithSastry, N.S.
Smt.K.Sujatha	Sh. Sudip Kumar Suklabaidya
Smt. K.V.Archana	Sh. Mahamad Ali, M.
Sh. N.Maddileti	Sh. Avinash, K.N.
	Sh. Amar Suputhra, S
	Sh. Anudeep, Y.
	Sh. Deepak, M.J.
	Smt. K.Karunya Lakshmi
	Ms. Seema, K.V.
	Ms. A. Rajab Nisha

Laboratory Analysis	
Dr. K.M.Nair	Dr. H.R. Savitha
Smt. ArtiKoyal	Ms. Steffi Peter
Smt. Parvathy, S.	Ms. Thara, V.R
	Ms. Roopa, G.
	Ms. Swati, H.
	Sh. Shantaveera Swami
	Ms. Shwetha, N.K.
	Smt. Ishrat Haji
	Ms. P. PavanKumari
	Ms. Padmaja
	Ms. Veena, M.
Soil & Water Conservation	
Sh. Sunil P. Maske	
Socio-Economic Analysis	
Dr. S.C. Ramesh Kumar	Sh. M. K. Prakashanaik
	Ms. Sowmya K.B
	Sh.Manjunath M
	Sh.Veerabhadraswamy R
	Sh.Lankesh RS
	Sh.Kalaveerachari R Kammar
	Sh.Pradyumma U
	Sh.Yogेशha HN
	Sh.Vijay kumar lamani
	Sh.Arun N Kambar
	Sh.Vinay
	Sh.Basavaraj.Biradar
	Sh.Vinod R
	Sh.Praveenkumar P Achalkar
	Sh.Rajendra D
Watershed Development Department, GoK, Bangalore	
Sh. Rajeev Ranjan IFS Project Director & Commissioner, WDD	Dr. A. Natarajan NRM Consultant, Sujala-III Project
Dr. S.D. Pathak IFS Executive Director & Chief Conservator of Forests, WDD	

PART-A

LAND RESOURCE INVENTORY

Contents

Preface		
Contributors		
Executive Summary		
Chapter 1	Introduction	1
Chapter 2	Geographical Setting	5
2.1	Location and Extent	5
2.2	Geology	6
2.3	Physiography	6
2.4	Drainage	7
2.5	Climate	7
2.6	Natural Vegetation	8
2.7	Land Utilization	9
Chapter 3	Survey Methodology	13
3.1	Base maps	13
3.2	Field Investigation	15
3.3	Laboratory Characterization	16
3.4	Finalization of Soil Map	16
Chapter 4	The Soils	23
4.1	Soils of Basalt Landscape	23
Chapter 5	Interpretation for Land Resource Management	31
5.1	Land Capability Classification	31
5.2	Soil Depth	33
5.3	Surface Soil Texture	34
5.4	Soil Gravelliness	35
5.5	Available Water Capacity	36
5.6	Soil Slope	37
5.7	Soil Erosion	38
Chapter 6	Fertility Status	41
6.1	Soil Reaction (pH)	41
6.2	Electrical Conductivity (EC)	41
6.3	Organic Carbon (OC)	41
6.4	Available Phosphorus	43
6.5	Available Potassium	43
6.6	Available Sulphur	43
6.7	Available Boron	43
6.8	Available Iron	46
6.9	Available Manganese	46
6.10	Available Copper	46

6.11	Available Zinc	46
Chapter 7	Land Suitability for Major Crops	49
7.1	Land suitability for Sorghum	49
7.2	Land suitability for Maize	52
7.3	Land suitability for Red gram	54
7.4	Land suitability for Sunflower	55
7.5	Land suitability for Cotton	57
7.6	Land suitability for Sugarcane	58
7.7	Land suitability for Soybean	60
7.8	Land suitability for Guava	61
7.9	Land suitability for Mango	62
7.10	Land suitability for Sapota	64
7.11	Land suitability for Jackfruit	66
7.12	Land suitability for Jamun	67
7.13	Land Suitability for Musambi	68
7.14	Land Suitability for Lime	69
7.15	Land Suitability for Cashew	71
7.16	Land Suitability for Custard Apple	72
7.17	Land Suitability for Amla	73
7.18	Land Suitability for Tamarind	74
7.19	Land Management Units	75
7.20	Proposed Crop Plan	77
Chapter 8	Soil Health Management	79
Chapter 9	Soil and Water conservation Treatment Plan	83
9.1	Treatment Plan	84
9.2	Recommended Soil and Water Conservation measures	87
9.3	Greening of microwatershed	88
	References	90
	Appendix I	I-VI
	Appendix II	VII-X
	Appendix III	XI-XIII

LIST OF TABLES

2.1	Mean Monthly Rainfall, PET, 1/2 PET at Aland Taluk, Gulbarga District	7
2.2	Land Utilization in Aland Taluk	10
3.1	Differentiating Characteristics used for Identifying Soil Series	16
3.2	Soil Legend	21
7.1	Soil-Site Characteristics of Chincholi Khurd-1 microwatershed	50
7.2	Crop suitability criteria for Sorghum	51
7.3	Crop suitability criteria for Maize	53
7.4	Crop suitability criteria for Red gram	54
7.5	Crop suitability criteria for Sunflower	56
7.6	Crop suitability criteria for Cotton	57
7.7	Crop suitability criteria for Sugarcane	59
7.8	Crop suitability criteria for Guava	61
7.9	Crop suitability criteria for Mango	63
7.10	Crop suitability criteria for Sapota	65
7.11	Crop suitability criteria for Lime	70
7.12	Proposed Crop Plan for Chincholi Khurd-1 Microwatershed	77

LIST OF FIGURES

2.1	Location map of Chincholi Khurd-1 microwatershed	5
2.2	Rock formations in Chincholi Khurd-1 microwatershed	6
2.3	Rainfall distribution in Aland Taluk, Gulbarga District	8
2.4	Natural Vegetation of Chincholi Khurd-1	9
2.5	Current Land use – Chincholi Khurd-1 microwatershed	10
2.6	Location of Wells- Chincholi Khurd-1 microwatershed	11
3.1	Scanned and Digitized Cadastral map of Chincholi Khurd-1 microwatershed	14
3.2	Satellite image of Chincholi Khurd-1 microwatershed	14
3.3	Cadastral map overlaid on IRS PAN+LISS IV merged imagery of Chincholi Khurd-1 microwatershed	15
3.4	Soil Phase or management units of Chincholi Khurd-1 microwatershed	19
5.1	Land Capability Classification of Chincholi Khurd-1 microwatershed	33
5.2	Soil Depth map of Chincholi Khurd-1 microwatershed	34
5.3	Surface Soil Texture map of Chincholi Khurd-1 microwatershed	35
5.4	Soil Gravelliness map of Chincholi Khurd-1 microwatershed	36
5.5	Soil Available Water Capacity map of Chincholi Khurd-1 microwatershed	37
5.6	Soil Slope map of Chincholi Khurd-1 microwatershed	38
5.7	Soil Erosion map of Chincholi Khurd-1 microwatershed	39
6.1	Soil Reaction (pH) map of Chincholi Khurd-1 microwatershed	42
6.2	Electrical Conductivity (EC) map of Chincholi Khurd-1 microwatershed	42
6.3	Soil Organic Carbon (OC) map of Chincholi Khurd-1 microwatershed	44
6.4	Soil Available Phosphorus map of Chincholi Khurd-1 microwatershed	44
6.5	Soil Available Potassium map of Chincholi Khurd-1 microwatershed	45
6.6	Soil Available Sulphur map of Chincholi Khurd-1 microwatershed	45
6.7	Soil Available Boron map of Chincholi Khurd-1 microwatershed	46
6.8	Soil Available Iron map of Chincholi Khurd-1 microwatershed	47
6.9	Soil Available Manganese map of Chincholi Khurd-1 microwatershed	47
6.10	Soil Available Copper map of Chincholi Khurd-1 microwatershed	48

6.11	Soil Available Zinc map of Chincholi Khurd-1 microwatershed	48
7.1	Land Suitability map of Sorghum	52
7.2	Land Suitability map of Maize	53
7.3	Land Suitability map of Red gram	55
7.4	Land Suitability map of Sunflower	56
7.5	Land Suitability map of Cotton	58
7.6	Land Suitability map of Sugarcane	59
7.7	Land Suitability map of Soybean	60
7.8	Land Suitability map of Guava	62
7.9	Land Suitability map of Mango	64
7.10	Land Suitability map of Sapota	66
7.11	Land Suitability map of Jackfruit	67
7.12	Land Suitability map of Jamun	68
7.13	Land Suitability map of Musambi	69
7.14	Land Suitability map of Lime	71
7.15	Land Suitability map of Cashew	72
7.16	Land Suitability map of Custard Apple	73
7.17	Land Suitability map of Amla	74
7.18	Land Suitability map of Tamarind	75
7.19	Land Management Units map of Chincholi Khurd-1 microwatershed	76
9.1	Soil and Water Conservation map of Chincholi Khurd-1 microwatershed	88

EXECUTIVE SUMMARY

The land resource inventory of Chincholi Khurd-1microwatershed was conducted using village cadastral maps and IRS satellite imagery on 1:7920 scale. The false colour composites of IRS imagery were interpreted for physiography and these physiographic delineations were used as base for mapping soils. The soils were studied in several transects and a soil map was prepared with phases of soil series as mapping units. Random checks were made all over the area outside the transects to confirm and validate the soil map unit boundaries. The soil map shows the geographic distribution and extent, characteristics, classification and use potentials of the soils in the microwatershed.

The present study covers an area of 532ha in Alandtaluk of Gulbarga district, Karnataka. The climate is semiarid and categorized as drought prone with an average annual rainfall of 786 mm of which about 595 mm is received during south –west monsoon, 116mm during north-east and the remaining 74 mm during the rest of the year. An area of about 98 per cent is covered by soils, two per cent by waterbodies, settlements and others. The salient findings from the land resource inventory are summarized briefly below.

- ❖ The soils belong to 6 soil series and 20 soil phases (management units) and 5land management units.*
- ❖ The length of crop growing period is about 150days starting from the 3rd week of June to 1st week of October.*
- ❖ From the master soil map, several interpretative and thematic maps like land capability, soil depth, surface soil texture, soil gravelliness, available water capacity, soil slope and soil erosion were generated.*
- ❖ Soil fertility status maps for macro and micronutrients were generated based on the surface soil samples collected at every 250 m grid interval.*
- ❖ Land suitability for growing major agricultural and horticultural crops were assessed and maps showing degree of suitability along with constraints were generated.*
- ❖ About 98 per cent area is suitable for agriculture and two per cent is not suitable for agriculture but well suited for forestry, pasture, agroforestry, silvi-pasture, recreation, installation of wind mills and as habitat for wildlife.*
- ❖ About 19 per cent of the soils are very deep (>150 cm) to deep (100 - 150 cm), 38 per cent are moderately shallow to shallow (25-75 cm) and about 42 per cent are very shallow (<25 cm) soils.*
- ❖ About 98 per cent of the area has clayey soils at the surface.*
- ❖ About 66 per cent of the area has non-gravelly soils, 31 per cent gravelly soils (15-35 % gravel) and one per cent very gravelly (35- 60% gravel) soils.*

- ❖ *About 16 per cent of the area has soils that are very high (>200mm/m) in available water capacity, one per cent medium (100-150 mm/m) and about 81 per cent low (50-100 mm/m) and very low (<50mm/m).*
- ❖ *About 72 per cent of the area has nearly level (0-1%) to very gently sloping (1-3% slope) lands and about 26 per cent area is gently (3-5%) to moderately sloping (5-10%) lands.*
- ❖ *An area of about 16 per cent has soils that are slightly eroded (e1), 69 per cent moderately eroded (e2) and 12 per cent are severely eroded (e3).*
- ❖ *An area of about 82 per cent has soils that are moderately alkaline (pH 7.8 to 8.4), 5 per cent strongly alkaline (pH 8.4 to >9.0) and about 11 per cent slightly alkaline (pH 7.3-7.8).*
- ❖ *The Electrical Conductivity (EC) of the soils are dominantly <2 dsm⁻¹ indicating that most of the soils are non-saline.*
- ❖ *About 62 per cent medium (0.5-0.75%), 30 per cent high (>0.75%) and 7 per cent low (<0.5%) in organic carbon.*
- ❖ *Major area of 86per cent has soils that are low (<23 kg/ha), 10 per cent medium (23-57 kg/ha) and 2 per cent in high (>57 kg/ha)in available phosphorus.*
- ❖ *About 17 per cent medium (145-337 kg/ha) and 81 per cent high (>337 kg/ha) in available potassium.*
- ❖ *Available sulphur is low (<10 ppm) in about 19per cent area, medium (10-20 ppm) in 67 per cent and 12 per cent is high (>20 ppm).*
- ❖ *Available boron is low (<0.5 ppm) in about 84 per cent area and 15 per cent medium (0.5-1.0 ppm).*
- ❖ *About 3 per cent area has soils that are deficient (<4.5 ppm) in available iron and 95per cent is sufficient (>0.6 ppm).*
- ❖ *Available manganese and copper are sufficient in all the soils.*
- ❖ *About 69 per cent area has soils that are deficient (<0.6 ppm) in available zinc and 29per cent is sufficient (>0.6 ppm).*
- ❖ *The land suitability for 18 major crops grown in the microwatershedwere assessed and the areas that are highly suitable (S1) and moderately suitable (S2) are given below. It is however to be noted that a given soil may be suitable for various crops but what specific crop to be grown may be decided by the farmer looking to his capacity to invest on various inputs, marketing infrastructure, price and finally the demand and supply position.*

Land suitability for various crops in the microwatershed

Crop	Suitability Area in ha (%)		Crop	Suitability Area in ha (%)	
	Highly suitable (S1)	Moderately suitable (S2)		Highly suitable (S1)	Moderately suitable (S2)
Sorghum	91 (17)	16 (3)	Sapota	-	-
Maize	-	-	Jackfruit	-	-
Red gram	-	107 (20)	Jamun	-	16 (3)
Sunflower	91 (17)	9 (2)	Musambi	91 (17)	9 (2)
Cotton	91 (17)	16 (3)	Lime	91 (17)	9 (2)
Sugarcane	-	-	Cashew	-	-
Soybean	91(17)	16 (3)	Custard apple	91 (17)	16 (3)
Guava	-	-	Amla	91 (17)	16 (3)
Mango	-	-	Tamarind	-	16 3)

- ❖ *Apart from the individual crop suitability, a proposed crop plan has been prepared for the 5 identified LMUs by considering only the highly and moderately suitable lands for different crops and cropping systems with food, fibre and horticulture crops.*
- ❖ *Maintaining soil-health is vital to crop production and conserve soil and land resource base for maintaining ecological balance and to mitigate climate change. For this, several ameliorative measures have been suggested to these problematic soils like saline/alkali, highly eroded, sandy soils etc.,*
- ❖ *Soil and water conservation treatment plan has been prepared that would help in identifying the sites to be treated and also the type of structures required.*
- ❖ *As part of the greening programme, several tree species have been suggested to be planted in marginal and submarginal lands and also in the hillocks, mounds and ridges.*

INTRODUCTION

Soil being a vital natural resource on whose proper use depends the life supporting systems of a country and the socioeconomic development of its people. Soils provide food, fodder, fibre and fuel for meeting the basic human and animal needs. With the ever increasing growth in human and animal population, the demand on soil for more food and fodder production is on the increase. The area available for agriculture is about 51 per cent of the total geographical area and more than 60 per cent of the people are still dependant on agriculture for their livelihood. However, the capacity of a soil to produce is limited and the limits to the production are set by its intrinsic characteristics, agro climatic setting and use and management. There is, therefore, tremendous pressure on land and water resources, which is causing decline in soil-health and stagnation in productivity. The soils have been degrading at an estimated rate of one million hectares per year and ground water levels have been receding at an alarming rate resulting in decline in the ground water resource. Further, land degradation has emerged as a serious problem which has already affected about 38 lakh ha of cultivated area in the State. Soil erosion alone has degraded about 35 lakh ha. Almost all the uncultivated areas are facing various degrees of degradation, particularly soil erosion; salinity and alkalinity has emerged as a major problem (>3.5 lakh ha) in the irrigated areas of the State. Nutrient depletion and declining factor productivity is common in both rainfed and irrigated areas. The degradation is continuing at an alarming rate and there appears to be no systematic effort among the stakeholders to contain this process. In recent times, an aberration of weather due to climate change phenomenon has added another dimension leading to unpredictable situations to be tackled by the farmers.

In this critical juncture, the challenge before us is not only to increase the productivity per unit area which is steadily declining and showing a fatigue syndrome, but also to prevent or at least reduce the severity of degradation. If the situation is not reversed at the earliest, then the sustainability of the already fragile crop production system and the overall ecosystem will be badly affected in the state.

Added to this, every year there is a significant diversion of farm lands and water resources for non-agricultural purposes. Thus, developing strategies to slow down the degradation process or reclaim the soils to normal condition and ensure sustainability of production system are the major issues today. This demands a systematic appraisal of our soil and land resources with respect to their extent, geographic distribution, characteristics, behaviour and use potential, which is very important for developing an effective land use and cropping systems for augmenting agricultural production on a sustainable basis.

The soil and land resource inventories made so far in Karnataka had limited utility because the surveys were of different types, scales and intensities carried out at different times with specific objectives.

Hence, there is an urgent need to generate detailed site-specific farm level database on various land resources for all the villages/watersheds in a time bound manner that would help to protect the valuable soil and land resources and also to stabilize the farm production.

Therefore, the land resource inventory required for farm level planning is the one which investigates all the parameters which are critical for productivity *viz.*, soils, site characteristics like slope, erosion, gravelliness and stoniness, climate, water, topography, geology, hydrology, vegetation, crops, land use pattern, animal population, socio-economic conditions, infrastructure, marketing facilities and various schemes and developmental works of the government etc. From the data collected at farm level, the specific problems and potentials of the area can be identified and highlighted, conservation measures required for the area can be planned on a scientific footing, suitability of the area for various uses can be worked out and finally viable and sustainable land use options suitable for each and every land holding can be prescribed.

The Land Resource Inventory is basically done for identifying potential and problem areas, developing sustainable land use plans, estimation of surface run off and water harvesting potential, preparation of soil and water conservation plans, land degradation/desertification etc. The Bureau is presently engaged in developing an LRI methodology using high resolution satellite remote sensing data and Digital Elevation Model (DEM) data to prepare Landscape Ecological Units (LEU) map representing agro-ecosystem as a whole. The LEU is preferred over landform as the base map for LRI. LEU is the assemblage of landform, slope and land use. An attempt was made to upscale the soil resource information from 1:250000 and 1:50000 scale to the LEU map in Goa and other states. Here, an attempt is being made to uplink the LRI data generated under Sujala-III Project to the Landscape Ecological Units (LEUs) map. For this, the major physiographic region, *i.e.*, South Deccan Plateau is taken as an example.

The Karnataka state has been divided into three major physiographic divisions, namely the Deccan Plateau, Hill Ranges and Coastal Plain (NATMO, 1980). These divisions have been subdivided into four regions based on their geographic location, namely the South Deccan Plateau, Western Ghats, Eastern Ghats and West Coast Plains.

The South Deccan Plateau locally known as the Karnataka Plateau, covers an area of about 15.8 m ha. The major part of the Plateau is a peneplain in various stages of development and destruction. The Plateau is divided into *Malnad* (Hilly area) and *Maidan* (Plains). *Malnad* is an area of rolling to undulating uplands with many valleys and is a transitional zone between the Western Ghats and the *Maidan*. It covers an area of about 6.2 mha. The *Maidan* has a rolling surface with altitude of 900-1150 m, 600-850 m, 450-550 m and 300-400 m above MSL. The highest surface is in the southwestern part of the state. The lowest surface is in the northeast in the valleys of the Tungabhadra and Hagari rivers.

The northern part of the Plateau is drained by the Krishna river and its tributaries, the Bhima, Malaprabha, Ghataprabha and Tungabhadra and the southern part by the Cauvery river and its tributaries, the Hemavathi, Kabini and Lakshmanthirtha.

The plateau has been divided into five landscapes (Shiva Prasad. *et, al.*, 1998) namely, Granite and gneiss landscape (Dsa), Basalt landscape (Dsb), Laterite landscape (Dsc), Metamorphic landscape (Dsd 1) and Sedimentary landscape (Dsd 2)

The land resource inventory aims to provide site specific database for Chincholi Khurd-1 microwatershed in Aland Taluk, Gulbarga District, Karnataka state for the Karnataka Watershed Development Department. The database was generated by using cadastral map of the village as a base along with high resolution IRS LISS IV and Cartosat-1 merged satellite imagery. Later, an attempt will be made to uplink this LRI data generated at 1:7920 scale under Sujala-III Project to the proposed Landscape Ecological Units (LEUs) map.

The study was organized and executed by the ICAR- National Bureau of Soil Survey and Land Use Planning, Regional Centre, Bangalore under Generation of Land Resource Inventory Data Base Component-1 of the Sujala-III Project funded by the World Bank.

GEOGRAPHICAL SETTING

2.1 Location and Extent

The study area of Chincholi Khurd-1 microwatershed (Khairat subwatershed) is located in the northeastern part of Karnataka in Aland Taluk, Gulbarga District, Karnataka State (Fig.2.1). It comprises of parts of Chincholi Budruk, Padsavali and Sarasamba villages. It lies between $17^{\circ} 35'$ and $17^{\circ} 45'$ North latitude and between $76^{\circ} 10'$ and $77^{\circ} 45'$ east longitude and covers an area of 532 ha. It is about 15 km south of Aland and is surrounded by Khanapur on the southeast, Chincholi Budruk village in the northwest, Nagalgaon on the southwest and Nirgudi on the north.

LOCATION MAP OF CHINCHOLI KHURD-1 MICRO-WATERSHED




Fig.2.1 Location map of Chincholi Khurd-1 microwatershed

2.2 Geology

Major rock formation observed in the microwatershed is Basalt (Fig.2.2) or Deccan Trap. The Deccan Traps cover the whole of Bidar, Parts of Gulbarga, Bijapur and Belgam districts. Eight lava flows have been identified in Karnataka horizontally overlying the older formations. The thickness of the individual flows averages about five meters. It is relatively uniform in petrographic character. The most common types is augite basalt. Dominant colour is grayish green and texture ranges from cryptocrystalline to glassy. The rock is often vesicular and scoriaceous filled up with secondary minerals like coloured agate, quartz, calcite and a large variety of zeolites. The Deccan Traps form an excellent building material and also used as road-metal and railway ballast.



Fig. 2.2 Basalt

2.3 Physiography

Physiographically, the area has been identified as Basalt landscape based on geology. Based on slope and its relief features, the area has been further subdivided into four landforms, viz; mounds/ridges, summits, side slopes and very gently sloping uplands. The elevation ranges from 480 to 566 m. The mounds and ridges are mostly covered by rock outcrops.

2.4 Drainage

The area is drained by several small parallel streams that join Monia nala which further joins Awarja river along its course. Though, it is not a perennial one, during rainy season it carries large quantities of rain water. The microwatershed has only few small tanks which are not capable of storing the water that flows during the rainy season. Due to this, the ground water recharge is very much affected. This is reflected in the failure of many bore wells in the village. If the available rain water is properly harnessed by constructing new tanks and recharge structures at appropriate places in the village, then the drinking and irrigation needs of the entire area can be easily met. The drainage network is parallel to sub parallel and dendritic.

2.5 Climate

The Gulbarga district lies in the northern plains of Karnataka and falls under semiarid tract of the state and is categorized as drought prone with average annual rainfall of 786 mm (Table 2.1). Of the total rainfall, maximum of 595 mm is received during the south-west monsoon period from June to September, the north-east monsoon from October to early December contributes about 116 mm, and the remaining 74 mm during the rest of the year. December is the coldest month with mean daily maximum and minimum temperatures being 29.5°C & 15°C to 10°C respectively. During peak summer, temperature shoots up to 45°C. Relative humidity varies from 26 per cent in summer to 62 per cent in winter. Rainfall distribution is shown in Figure 2.3. The average potential evapotranspiration (PET) is 150 mm and varies from a low of 115 mm in December to 232 mm in the month of May. The PET is always higher than precipitation in all the months except August and September. Generally, the length of crop growing period (LGP) is 150 days and starts from 3rd week of June to third week of November.

Table 2.1 Mean Monthly Rainfall, PET, 1/2 PET at Aland Taluk, Gulbarga District

Sl.No.	Months	Rainfall	PET	1/2 PET
1	January	7.50	126.80	63.40
2	February	3.40	143.90	71.95
3	March	11.30	189.90	94.95
4	April	19.40	209.80	104.90
5	May	32.70	232.20	116.10
6	June	111.00	186.40	93.20
7	July	139.20	152.80	76.40
8	August	172.40	147.60	73.80
9	September	172.30	131.70	65.85
10	October	91.30	145.50	72.75
11	November	19.30	129.80	64.90
12	December	5.80	114.80	57.40
Total		785.6	149.70	




Fig 2.3 Rainfall distribution in Aland Taluk, Gulbarga District

2.6 Natural Vegetation

The natural vegetation is sparse comprising few tree species, shrubs and herbs. The mounds, ridges and boulders occupy very sizeable area which are under thin to moderately thick forest vegetation. Still, there are some remnants of the past forest cover which can be seen in patches in some ridges and hillocks in the microwatershed (Fig. 2.4).

Apart from the continuing deforestation, the presence of large population of goats, sheep and other cattle in the microwatershed is causing vegetative degradation of whatever little vegetation left in the area. The uncontrolled grazing has left no time for the regeneration of the vegetative cover. This leads to the accelerated rate of erosion on the hill slopes resulting in the formation of deep gullies in the foot slopes that eventually result in the heavy siltation of tanks and reservoirs in the microwatershed.



Fig. 2.4 Natural vegetation (Scrub) of Chincholi Khurd-1 microwatershed

2.7 Land Utilization

About 89 per cent area (Table 2.2) in Aland taluk is cultivated at present. An area of about 2 per cent is currently barren. Forests occupy an area of about 2 per cent and the tree cover is in a very poor state. Most of the mounds, ridges and bouldery areas have very poor vegetative cover. Major crops grown in the area are sorghum, maize, cotton, sugarcane, safflower, groundnut, red gram and sapota. While carrying out land resource inventory, the land use/land cover particulars are collected from all the survey numbers and a current land use map of the microwatersheds is prepared. The current land use map prepared shows the arable and non-arable lands, other land uses and different types of crops grown in the area. The current land use map of Chincholi Khurd-1 microwatershed is presented in Fig.2.5.

Table 2.2 Land Utilization in Aland Taluk

Sl. No.	Agricultural land use	Area (ha)	Per cent
1.	Total geographical area	173417	
2.	Total cultivated area	153806	88.69
3.	Area sown more than once	7910	
4.	Trees and grooves	59	0.034
5.	Forest	2854	1.64
6.	Cultivable wasteland	974	0.56
7.	Permanent Pasture land	3469	2.00
8.	Barren land	3142	1.81
9.	Non- Agriculture land	3465	1.99
10.	Currently Fallow lands	5648	3.25




Fig.2.5 Current Land Use – Chincholi Khurd-1 microwatershed

Simultaneously, enumeration of wells (bore wells and open wells) in the microwatershed was made and their location in different survey numbers is located on the cadastral map. Map showing the location of wells and other water bodies in the Chincholi Khurd-1 microwatershed is given (Fig.2.6).




Fig.2.6 Location of Wells – Chincholi Khurd-1 microwatershed

SURVEY METHODOLOGY

The purpose of land resource inventory is to delineate similar areas (soil series and phases), which respond or expected to respond similarly to a given level of management. This was achieved in Chincholi Khurd -1 microwatershed by the detailed study of all the soil characteristics (depth, texture, colour, structure, consistence, coarse fragments, porosity, soil reaction, soil horizons etc.) and site (slope of the land, erosion, drainage, occurrence of rock fragments etc.) followed by grouping of similar areas based on soil-site characteristics into homogeneous (management units) units and showing their extent and geographic distribution on the microwatershed cadastral map. The detailed survey at 1:7920 scale was carried out in 532 ha area. The methodology followed for carrying out land resource inventory was as per the guidelines given in Soil Survey Manual (IARI, 1971; Soil Survey Staff, 2006; Natarajan *et al.*, 2015) which is briefly described below

3.1 Base Maps

The detailed survey of the land resources occurring in the microwatershed was carried out by using digitized cadastral map as a base. The cadastral map shows field boundaries with their survey numbers, location of tanks, streams and other permanent features of the area (Fig. 3.1). Apart from the cadastral map, remote sensing data products from Cartosat-1 and LISS-IV merged at the scale of 1:7920 were used in conjunction with the cadastral map to identify the rock types, the landscapes, landforms and other surface features. The imagery helped in the identification and delineation of boundaries between hills, uplands and lowlands, water bodies, forest and vegetated areas, roads, habitations and other cultural features of the area (Fig. 3.2). The cadastral map was overlaid on the satellite imagery (Fig.3.3) that helps to identify the parcel boundaries and other permanent features. Apart from cadastral maps and images, toposheets of the area (1:50,000 scale) were used for initial traversing, identification of geology and landforms, drainage features, present land use and also for selection of transects in the microwatershed.




Fig 3.1 Scanned and Digitized Cadastral map of Chincholi Khurd-1 microwatershed




Fig.3.2 Satellite Image of Chincholi Khurd-1 microwatershed




Fig.3.3 Cadastral map overlaid on IRS PAN+LISS IV merged imagery of Chincholi Khurd-1 microwatershed

3.2 Field Investigation

Preliminary traverse of the microwatershed was carried out with the help of cadastral map, imagery and toposheets. While traversing, landforms and physiographic units identified were checked and preliminary soil legend was prepared by studying soils at few selected places.

The field boundaries and survey numbers given on the cadastral sheet were located on the ground by following permanent features like roads, cart tracks, nallas, streams, tanks etc., and wherever changes were noticed, they were incorporated on the microwatershed cadastral map. Then, intensive traversing of each physiographic unit like hills, ridges and uplands was carried out. Based on the variability observed on the surface, transects were selected across the slope covering all the landform units in the microwatershed (Natarajan and Dipak Sarkar, 2010).

In the selected transect, soil profiles were located at closely spaced intervals to take care of any change in the land features like break in slope, erosion, gravel, stones etc. In the selected sites, profiles (vertical cut showing the soil layers from surface to the rock) were opened up to 200 cm or to the depth limited by rock or hard substratum and studied in detail for all their morphological and physical characteristics. The soil and site characteristics were recorded for all profile sites on a standard proforma as per the guidelines given in USDA Soil Survey Manual (Soil Survey Staff, 2012). Apart from the transect study, profiles were also studied at random, almost like in a grid pattern, outside the transect areas.

Based on the soil-site characteristics, the soils were grouped into different soil series (soil series is the most homogeneous unit having similar horizons and properties and behaves similarly for a given level of management). Soil depth, texture, colour, kind of horizon and horizon sequence, amount and nature of gravel present, nature of substratum etc, were used as the major differentiating characteristics for identifying soil series occurring in the area. The differentiating characteristics used for identifying the soil series are given in Table 3.1. Based on the above characteristics, 6 soil series were identified in the Chincholi Khurd-1 microwatershed.

Table 3.1 Differentiating Characteristics used for Identifying Soil Series
(Characteristics are of Series Control Section)

SOILS OF BASALT LANDSCAPE							
Sl. no	Soil Series	Depth (cm)	Colour	Text-ure	Gravel (%)	Horizon sequence	Calcar-eousness
1	Margutti (MGT)	<25	10YR3/3,4/3,5/4 7.5YR4/3	c	15-35	Ap- cr	-
2	Novinihala (NHA)	25-50	10YR3/2,3/1,4/2 7.5YR3/4	c	<15	Ap-Bw- cr/R	-
3	Bhimanahalli (BHI)	25-50	10YR3/2,3/3,3/1 7.5YR3/2,4/2	c	15-35	Ap-Bw- cr/R	-
4	Dinsi (DSI)	50-75	10YR3/2,3/3,4/3 3/2	c	<15	Ap-BA- Bss	-
5	Rajnala (RNL)	100- 150	10YR3/2,3/1,4/2,4/3	c	<15	Ap-BA- -Bss-cr	-
6	Mahagaon (MAN)	>150	10YR3/2,3/1	c	<15	Ap-BA- Bss	-

3.3 Laboratory Characterization

Soil samples were collected from representative master profiles for laboratory characterization by following the methods outlined in the Laboratory Manual (Sarma *et al*, 1987). Surface soil samples collected from farmer's fields (84 samples) for fertility status (major and micronutrients) at 250 m grid interval were analyzed in the laboratory. (Katyal and Rattan, 2003) By linking the soil fertility data to the survey numbers through GIS, soil fertility maps were generated for the microwatershed.

3.4 Finalization of Soil Maps

The area under each soil series was further separated into soil phases and their boundaries delineated on the cadastral map based on the variations observed in the texture of the surface soil, slope, erosion, presence of gravel, stoniness etc. A soil phase is a subdivision of soil series based mostly on surface features that affect its use and management.

The soil mapping units are shown on the map (Fig.3.4) in the form of symbols. During the survey about 21 profile pits, few minipits and a few auger bores representing different landforms occurring in the microwatershed were studied. All the profile locations are indicated on the village cadastral map in the form of a triangle. In addition to the profile study, spot observations in the form of minipits, road cuts, terrace cuts etc., were studied to validate the soil boundaries on the soil map. The soil map shows the geographic distribution of 20 mapping units representing 6 soil series occurring in the microwatershed. The soil map unit (soil legend) description is presented in Table 3.2.

The soil phase map (management units) shows the distribution of 20 phases mapped in the microwatershed. Each mapping unit (soil phase) delineated on the map has similar soil and site characteristics. In other words, all the farms or survey numbers included in one phase will have similar management needs and they have to be treated accordingly.

The 20 soil phases identified and mapped in the microwatershed were regrouped into 5 Land Management Units (LMU's) for the purpose of preparing a proposed land use plan for sustained development of the microwatershed. The database (soil phases) generated under LRI was utilized for identifying Land Management Units (LMUs) based on the management needs. One or more than one soil site characteristic having influence on the management have been chosen for identification and delineation of LMUs. For Chincholi Khurd-1 microwatershed, five soil and site characteristics, namely soil depth, soil texture, slope, erosion and gravel content have been considered for defining LMUs. The land management units are expected to behave similarly for a given level of management.




Fig 3.4 Soil phase or management units map of Chincholi Khurd-1 microwatershed

Table 3.2 Soil Legend

Soil map unit No	Soil series	Soil phase	Mapping unit Description	Area in ha (%)
Soils of Basalt Landscape				
	MGT	Marguti soils are very shallow (<25cm), well drained, have very dark grayish brown to dark brown clayey soils occurring on nearly level to moderately sloping uplands		221.15 (41.54)
1		MGTiC3g1	Sandy clay surface, slope 3-5 %, severe erosion, gravelly (15-35%)	51.54 (9.68)
2		MGTiD3g2	Sandy clay surface, slope 5-10 %, severe erosion, very gravelly (35-60 %)	5.10 (0.96)
3		MGTmA1	Clay surface, slope 0-1%, slight erosion	7.71 (1.45)
4		MGTmB1	Clay surface, 1-3 % slope, slight erosion	5.68 (1.07)
5		MGTmB2	Clay surface, 1-3 % slope, moderate erosion	93.02 (17.47)
6		MGTmB2g1	Clay surface, 1-3 % slope, moderate erosion, gravelly (15-35%).	13.87 (2.60)
7		MGTmB3g1	Clay surface, 1-3 % slope, severe erosion, gravelly (15-35%).	5.00 (0.94)
8		MGTmC2	Clay surface, slope 3-5%, moderate erosion.	11.17 (2.10)
9		MGTmC2g1	Clay surface, slope 3-5%, moderate erosion, gravelly (15-35%).	28.06 (5.27)
	NHA	Novinihala soils are shallow (25-50 cm), well drained, have very dark grayish brown to dark brown clayey soils occurring on very gently sloping uplands		89.85 (16.88)
10		NHAmB1	Clay surface, slope 1-3%, slight erosion	7.36 (1.38)
11		NHAmB2	Clay surface, slope 1-3%, moderate erosion	82.49 (15.49)
	BHI	Bhimanahalli soils area shallow (25-50 cm), well drained, have very dark gray to brown clay soils occurring on very gently sloping to gently sloping uplands.		103.90 (19.52)
12		BHImB1	Clay surface, slope 1-3%, slight erosion	46.18 (8.67)
13		BHImB2g1	Clay surface, slope 1-3%, moderate erosion, gravelly (15-35%).	25.76 (4.84)
14		BHImC2g1	Clay surface, 3-5% slope, moderate erosion, gravelly (15-35%).	31.97 (6.00)
	DSI	Dinsi soils are moderately shallow (50-75 cm), moderately well drained, have very dark gray to brown clayey soils occurring on very gently sloping uplands		7.22 (1.36)
15		DSImB2	Clay surface, slope 1-3%, moderate erosion.	7.22 (1.36)

	RNL	Rajnala soils are deep (100-150 cm), moderately well drained, have very dark gray to brown calcareous clay soils occurring on very gently to gently sloping uplands	15.84 (2.98)
16	RNLmB1	Clay surface, slope 1-3 %, slight erosion	6.57 (1.23)
17	RNLmC2g1	Clay surface, slope 3-5%, moderate erosion, gravelly (15-35%).	9.27 (1.74)
	MAN	Mahagaon soils are very deep (>150 cm), moderately well drained, have very dark gray to very dark grayish brown cracking clay soils occurring on nearly level to very gently sloping uplands	84.66 (15.90)
18	MANmA1	Clay surface, 0-1% slope, slight erosion	1.74 (0.33)
19	MANmB1	Clay surface, slope 1-3%, slight erosion	18.17 (3.41)
20	MANmB2	Clay surface, slope 1-3%, moderate erosion	64.75 (12.16)
Miscellaneous Lands			
21	Water body		9.79 (1.84)

THE SOILS

Detailed information pertaining to the nature, extent and distribution of different kinds of soils occurring in Chincholi Khurd-1 microwatershed is provided in this chapter. The microwatershed area has Basaltic landscape. In all, 6 soil series were identified in this landscape. Soil formation is the result of the combined effect of environmental and terrain factors that are reflected in soil morphology. In the Basalt landscape, it is by parent material and climate. A brief description of each of the 6 soil series identified followed by 20 soil phases (management units) mapped under each series are furnished below. The soils in any one map unit differ from place to place in their depth, texture, slope, gravelliness, erosion or any other site characteristics that affect management. The soil phase map can be used for identifying the suitability of areas for growing specific crops or for other alternative uses and also for deciding the type of conservation structures needed. The detailed information on soil and site-characteristics like soil depth, surface soil texture, slope, erosion, gravelliness, AWC, LCC etc, with respect to each of the soil phase identified is given village/survey number wise for the microwatershed in Appendix-I.

4.1 Soils of Basalt Landscape

In this landscape, 6 soil series are identified and mapped. Of these, Margutti (MGT) soil series occupies maximum area of about 221.15 (41.54). The brief description of each series along with the soil phases identified and mapped is given below.

4.1.1 Margutti (MGT) Series: Marguti soils are very shallow (<25cm), well drained, have very dark grayish brown to dark brown clay soils. They have developed from basalt and occur on very gently sloping to moderately sloping uplands.

The total depth of the soil ranges from 10 to 23 cm. The thickness of A horizon ranges from 7 to 18 cm. Its colour is in 10 YR and 7.5 YR hue with value 3 to 4 and chroma 2 to 4. The texture is clay with 15 to 35 per cent gravel. The available water capacity is very low (<50 mm/m). Nine phases were identified:

MGTiC3g1	Sandy clay surface, slope 3-5 %, severe erosion, gravelly (15-35%)
MGTiD3g2	Sandy clay surface, slope 5-10 %, severe erosion, very gravelly (35-60 %)
MGTmA1	Clay surface, slope 0-1%, slight erosion
MGTmB1	Clay surface, 1-3 % slope, slight erosion
MGTmB2	Clay surface, 1-3 % slope, moderate erosion
MGTmB2g1	Clay surface, 1-3 % slope, moderate erosion, gravelly (15-35%)
MGTmB3g1	Clay surface, 1-3 % slope, severe erosion, gravelly (15-35%)
MGTmC2	Clay surface, slope 3-5%, moderate erosion
MGTmC2g1	Clay surface, slope 3-5%, moderate erosion, gravelly (15-35%)



Landscape and Soil Profile characteristics of Margutti (MGT) Series

4.1.2 Novanihala (NHA) Series: Novanihala soils are shallow (25-50 cm), well drained, have very dark grayish brown to dark brown clay soils. They have developed from basalt and occur on very gently sloping uplands.

The thickness of the solum ranges from 27 to 48 cm. The thickness of A horizon ranges from 12 to 20 cm. Its colour is in 7.5 YR and 10 YR hue with value 3 to 4 and chroma 2 to 4. The texture varies from sandy clay to clay with 10 to 20 per cent gravel. The thickness of B horizon ranges from 22 to 37 cm. Its colour is in 10 YR and 7.5 YR hue with value 3 to 4 and chroma 2 to 4. Its texture is clay with gravel content of 10-15 per cent. The available water capacity is low (51-100 mm/m).

Two phases were identified:

NHAmB1	Clay surface, slope 1-3%, slight erosion
NHAmB2	Clay surface, slope 1-3%, moderate erosion




.. Landscape and Soil profile Characteristics of Novanihala (NHA) Series

4.1.3 Bhimanahalli (BHI) Series: Bhimanahalli soils are shallow (25-50 cm), well drained, have very dark gray to brown clay soils. They have developed from basalt and occur on very gently sloping to gently sloping uplands.

The thickness of the solum ranges from 29 to 48 cm. The thickness of A horizon ranges from 15 to 20 cm. Its colour is in 7.5 YR and 10 YR hue with value 3 to 4 and chroma 2 to 4. The texture varies from sandy clay to clay with 15 to 25 per cent gravel. The thickness of B horizon ranges from 23 to 33 cm. Its colour is in 10 YR and 7.5 YR hue with value 3 to 4 and chroma 1 to 3. Its texture is clay with gravel content of 15 to 35 per cent. The available water capacity is very low (<50 mm/m).

Three phases were identified:

BHImB1	Clay surface, slope 1-3%, slight erosion
BHImB2g1	Clay surface, slope 1-3%, moderate erosion, gravelly (15-35%)
BHImC2g1	Clay surface, 3-5% slope, moderate erosion, gravelly (15-35%)



Landscape and Soil profile Characteristics of Bhimanahalli (BHI) Series

4.1.4 Dinsi (DSI) Series: Dinsi soils are moderately shallow (50-75 cm), moderately well drained, have very dark gray to brown clay soils. They have developed from basalt and occur on gently sloping uplands.

The thickness of the solum ranges from 51 to 71 cm. The thickness of A horizon ranges from 9 to 24 cm. Its colour is in 10 YR hue with value 3 and chroma 1 to 3. The texture is clay with 5 to 10 per cent gravel. The thickness of B horizon ranges from 27 to 62 cm. Its colour is in 10 YR hue with value 3 to 4 and chroma 2 to 4. Its texture is clay with gravel content of less than 15 per cent. The available water capacity is medium (101-150 mm/m).

Only one phase was identified:

DSImB2	Clay surface, slope 1-3%, moderate erosion
--------	--



Landscape and Soil profile Characteristics of Dinsi (DSI) Series

4.1.5 Rajnala (RNL) Series: Rajnala soils are deep (100-150 cm), moderately well drained, have very dark gray to brown cracking clay soils. They have developed from basalt and occur on very gently to gently sloping uplands.

The thickness of the solum ranges from 125 to 140 cm. The thickness of A horizon ranges from 14 to 23 cm. Its colour is in 10 YR hue with value 3 to 4 and chroma 1 to 2. The texture is clay with less than 10 per cent gravel. The thickness of B horizon ranges from 85 to 130 cm. Its colour is in 10 YR hue with value 3 to 4 and chroma 1 to 3. Its texture is clay with gravel content of less than 15 per cent. The available water capacity is very high (>200 mm/m).

Two phases were identified

RNLmB1	Clay surface, slope 1-3 %, slight erosion
RNLmC2g1	Clay surface, slope 3-5%, moderate erosion, gravelly (15-35%).



Landscape and Soil profile Characteristics of Rajnala (RNL) Series

4.1.6 Mahagaon (MAN) Series: Mahagaon soils are very deep (>150 cm), moderately well drained, have very dark gray to very dark grayish brown cracking clay soils. They have developed from basalt and occur on nearly level to very gently sloping uplands.

The thickness of the solum ranges from 151 to 170 cm. The thickness of A horizon ranges from 18 to 22 cm. Its colour is in 10 YR and 7.5 YR hue with value 3 to 4 and chroma 1 to 3. The texture is clay with less than 10 per cent gravel. The thickness of B horizon ranges from 130 to 160 cm. Its colour is in 10 YR hue with value 3 to 4 and chroma 1 to 2. Its texture is clay with gravel content of less than 15 per cent. The available water capacity is very high (>200 mm/m).

Two phases were identified

MANmA1	Clay surface, 0-1% slope, slight erosion
MANmB1	Clay surface, slope 1-3%, slight erosion
MANmB2	Clay surface, slope 1-3%, moderate erosion



Landscape and Soil profile Characteristics of Mahagaon (MAN) Series

INTERPRETATION FOR LAND RESOURCE MANAGEMENT

The most important soil and site characteristics that affect the land use and conservation needs of an area are land capability, soil depth, soil texture, coarse fragments, available water capacity, soil slope, soil erosion, soil reaction etc. These are interpreted from the data base gathered through land resource inventory and several thematic maps are generated. These would help in identifying the areas suitable for growing crops and, soil and water conservation measures and structures needed thus helping to maintain good soil health for sustained crop production. The various thematic maps generated are described below.

5.1 Land Capability Classification

Land capability classification is an interpretative grouping of soil map units (soil phases) mainly based on inherent soil characteristics, external land features and environmental factors that limit the use of land for agriculture, pasture, forestry, or other uses on a sustained basis (IARI, 1971). The land and soil characteristics used to group the land resources in an area into various land capability classes, subclasses and units are

Soil characteristics: Soil depth, soil texture, coarse fragments, soil reaction, available water capacity, calcareousness, salinity/alkali *etc.*

Land characteristics: Slope, erosion, drainage, rock outcrops.

Climate: Total rainfall and its distribution, and length of crop growing period.

The Land capability classification system is divided into land capability classes, subclasses and units based on the level of information available. Eight land capability classes are recognized. They are

Class I: The soil map units have few or very few limitations that restrict their use.

Class II: The soil map units have moderate limitations that reduce the choice of crops or that require moderate conservation practices.

Class III: The soil map units have severe limitations that reduce the choice of crops or that require special conservation practices.

Class IV: The soil map units have very severe limitations that reduce the choice of crops or that require very careful management.

Class V: Soils in the mapping units are not likely to erode, but have other limitations that are impractical to remove and as such not suitable for agriculture.

Class VI: The lands have severe limitations that make them generally unsuitable for cultivation.

Class VII: The lands have very severe limitations that make them unsuitable for cultivation.

Class VIII: Soil and other miscellaneous areas that have very severe limitations that nearly preclude their use for any crop production.

The land capability subclasses are recognised based on the dominant limitations observed within a given land capability class. The subclasses are designated by adding a lower case letter like 'e', 'w', 's', or 'c' to the class numeral. The subclass "e" indicates that the main hazard is risk of erosion, "w" indicates drainage or wetness as a limitation for plant growth, "s" indicates shallow soil depth, coarse or heavy textures, calcareousness, salinity/alkalinity or gravelliness and "c" indicates limitation due to climate.

The land capability subclasses have been further subdivided into land capability units based on the kinds of limitations present in each subclass. Ten land capability units are used in grouping the soil map units. They are stony or rocky (0), erosion hazard (slope, erosion) (1), coarse texture (sand, loamy sand, sandy loam) (2), fine texture (cracking clay, silty clay) (3) slowly permeable subsoil (4), coarse underlying material (5), salinity/alkali (6), stagnation, overflow, high ground water table (7), soil depth (8) and fertility problems (9). The capability units thus identified have similar soil and land characteristics that respond similarly to a given level of management. The soils of the microwatershed have been classified upto land capability subclass level only.

The 20 soil map units identified in the Chincholi Khurd -1 microwatershed are grouped under 3 land capability classes and 5 land capability subclasses. About 98 per cent area in the microwatershed is suitable for agriculture (Fig. 5.1)

Good cultivable lands (Class II) cover about 20 per cent area and are distributed in the north and northeastern part of the micowatershed with moderate problems of soil and erosion.

Moderately good cultivable lands (Class III) cover maximum area of about 66 per cent and are distributed in all over the microwatershed with moderate problems of erosion and soil.

The fairly cultivable lands (class IV) cover about 12 per cent area. They have limitations of erosion and soil and are distributed in the northern, northeastern and southern part of the microwatershed.




Fig. 5.1 Land Capability map of Chincholi Khurd-1 microwatershed

5.2 Soil Depth

Soil depth refers to the depth of the soil occurring above the parent material or hard rock. The depth of the soil determines the effective rooting depth for plants and in accordance with soil texture, mineralogy and gravel content, the capacity of the soil column to hold water and nutrient availability. Soil depth is one of the most important soil characteristic that is used in differentiating soils into different soil series. The soil depth classes used in identifying soils in the field are very shallow (<25 cm), shallow (25-50 cm), moderately shallow (50-75 cm), moderately deep (75-100 cm), deep (100-150 cm) and very deep (>150 cm). They were used to classify the soils into different depth classes and a soil depth map was prepared (Fig. 5.2).

Very deep soils (>150 cm) occur in about 85 ha (16%) and are distributed in the northern and central part of the microwatershed. Deep soils (100-150 cm) occur in small area of about 16 ha (3%) and are distributed in the northeastern part of the micro watershed. Moderately shallow (50-75 cm) soils occupy about 7 ha (1%) and are distributed in the southwestern part of the microwatershed.

Maximum area of about 221 ha (41%) is under very shallow (<25 cm) and are distributed in all parts of the microwatershed. Shallow soils (25-50 cm) occupy an area of about 194 ha (36%) in the southeastern, central and western part of the microwatershed. The most productive lands 85 ha (16%) with respect to soil rooting depth where all climatically

adapted annual and perennial crops can be grown are very deep soils (>150 cm depth) occurring in the northern and central part of the microwatershed.

The most problem lands with an area of about 415 ha (77%) having very shallow (<25 cm) and shallow (25-50 cm) rooting depth occur in all parts of the microwatershed. They are not suitable for growing agricultural crops but well suited for pasture, forestry or other recreational purposes. Occasionally, short duration crops may be grown if rainfall is normal.




Fig. 5.2 Soil Depth map of Chincholi Khurd-1 microwatershed

5.3 Surface Soil Texture

Texture is an expression to indicate the coarseness or fineness of the soil as determined by the relative proportion of primary particles of sand, silt and clay. It has a direct bearing on the structure, porosity, adhesion and consistence. The surface layer of a soil to a depth of about 25 cm is the layer that is most used by crops and plants. The surface soil textural class provides a guide to understanding soil-water retention and availability, nutrient holding capacity, infiltration, workability, drainage, physical and chemical behaviour, microbial activity and crop suitability.

Maximum area of 466 ha (88%) has soils that are clayey and are distributed all over the microwatershed and a minor area has soils that are sandy clay 57 ha (11%). They are distributed in the northern and southeastern part of the microwatershed (Fig. 5.3).

The most productive lands (88%) with respect to surface soil texture are the clayey soils that have high potential for soil-water retention and availability, and nutrient retention and availability, but have problems of drainage, infiltration, workability and other physical problems.




Fig. 5.3 Surface Soil Texture map of Chincholi Khurd-1 microwatershed

5.4 Soil Gravelliness

Gravel is the term used for describing coarse fragments between 2 mm and 7.5 cm diameter and stones for those between 7.5 cm and 25 cm. The presence of gravel and stones in soil reduces the volume of soil responsible for moisture and nutrient storage, drainage, infiltration and runoff and hinders plant growth by impeding root growth and seedling emergence, intercultural operations and farm mechanization.

Maximum area has soils that are nongravelly (<15%) covering about 352 ha (66%) and are distributed all over the microwatershed (Fig.5.4).

About 165 ha (31%) of area in the micro watershed has soils that are gravelly (15-35%) and are distributed in central, eastern and western part of the microwatershed followed by small area of soils that are very gravelly (35-60%) covering about 5 ha (1%) that are distributed in the southeastern part of the microwatershed.

The most productive lands with respect to gravelliness are found to be 66 per cent. They are nongravelly with less than 15 per cent gravel and have potential for growing both annual and perennial crops.

The problem soils (<1%) that are very gravelly (35-60%), where only short duration crops can be grown.




Fig. 5.4 Soil Gravelliness map of Chincholi Khurd-1 microwatershed

5.5 Available Water Capacity

The soil available water capacity (AWC) is estimated based on the ability of the soil column to retain water between the tensions of 0.33 and 15 bar in a depth of 100 cm or the entire solum if the soil is shallower. The AWC of the soils (soil series) as estimated by considering the soil texture, mineralogy, soil depth and gravel content (Sehgal *et al.*, 1990) and accordingly the soil map units were grouped into five AWC classes *viz.*, very low (<50 mm/m), low (50-100 mm/m), medium (100-150 mm/m), high (150-200 mm/m) and very high (>200 mm/m) and using these values, an AWC map was prepared (Fig. 5.5).

Major area of about 325 ha (61%) has soils that are very low (<50 mm/m) in available water capacity and are distributed in all parts of the microwatershed. An area of about 106 ha (20%) has soils that are low (51-100 mm/m) in available water capacity and are distributed in the southeastern and western part of the microwatershed. Small area in the microwatershed has soils that are medium (101-150 mm/m) in available water capacity. They occur in small area of about 7 ha (1%) and are distributed in the southwestern part of the microwatershed and followed by soils that are very high (>200 mm/m) in AWC covering about 85 ha (16%) in the microwatershed and are distributed in northern parts of the microwatershed.

An area of about 85 ha (16%) has soils that have very high potential (>200 mm/m) with regard to available water capacity. In these areas, if the rainfall is normal and well

distributed, all climatically adapted long duration annual and perennial crops can be grown. About 325 ha (61%) area in the microwatershed has soils that are problematic with regard to available water capacity. Here, only short or medium duration crops can be grown and the probability of crop failure is very high. These areas are best put to other alternative uses.




Fig. 5.5 Soil Available Water Capacity map of Chincholi Khurd-1 microwatershed

5.6 Soil Slope

Soil slope refers to the inclination of the surface of the land. It is defined by gradient, shape and length, and is an integral feature of any soil as a natural body. Slope is considered important in soil genesis, land use and land development. The length and gradient of slope influences the rate of runoff, infiltration, erosion and deposition. The soil map units were grouped into four slope classes and a slope map was prepared showing the area extent and geographic distribution of different slope classes in the microwatershed (Fig. 5.6).

Major area of the microwatershed falls under very gently sloping (1-3% slope) slope class. It covers an area of about 376 ha (71%) and is distributed in all parts of the microwatershed and an area of about 132 ha (25 %) in the microwatershed falls under gently sloping (3-5%) slope class and is distributed in the northern, northeastern and southeastern part of the microwatershed.

Moderately sloping (5-10% slope) slope class cover a very minor area of about 5 ha (<1%) and is distributed in the southeastern part of the microwatershed. Nearly level lands

(0-1%) slope class covers about 5 ha (12%) and distributed in southwestern parts of the microwatershed.

An area of about 376 ha (72%) in the microwatershed has soils that have high potential in respect of soil slopes. In these areas, all climatically adapted annual and perennial crops can be grown without much soil and water conservation and other land development measures.




Fig. 5.6 Soil Slope map of Chincholi Khurd-1 microwatershed

5.7 Soil Erosion

Soil erosion refers to the wearing away of the earth's surface by the forces of water, wind and ice involving detachment and transport of soil by raindrop impact. It is used for accelerated soil erosion resulting from disturbance of the natural landscape by burning, excessive grazing and indiscriminate felling of forest trees and tillage, all usually by man. The erosion classes showing an estimate of the current erosion status as judged from field observations in the form of rills, gullies or a carpet of gravel on the surface are recorded. Four erosion classes, viz, slight erosion (e1), moderate erosion (e2), severe erosion (e3) and very severe erosion (e4) are recognized. The soil map units were grouped into different erosion classes and soil erosion map prepared. The area extent and their spatial distribution in the microwatershed is given in Figure 5.7.

Soils that are slightly eroded (e1 class) covers about 93 ha (17%) and are distributed in the eastern and western part of the microwatershed.

Soils that are moderately eroded (e2 class) cover about 368 ha (69%) in the microwatershed and are distributed in all parts of the microwatershed. Severely eroded (e3 class) soils cover an area about 62 ha (12%) and are distributed in all parts of the microwatershed. Top priority is to be given to these areas for taking up soil and water conservation and other land development measures followed by moderately eroded lands (69%).




Fig. 5.7 Soil Erosion map of Chincholi Khurd-1 microwatershed

FERTILITY STATUS

Soil fertility plays an important role in increasing crop yield. The adoption of high yielding varieties that require high amounts of nutrients has resulted in deficiency symptoms in crops and plants due to imbalanced fertilization and poor inherent fertility status. Hence, it is necessary to know the fertility (macro and micro nutrients) status of the soils of the watersheds for assessing the kind and amount of fertilizers required for each of the crop intended to be grown. For this purpose, the surface soil samples collected from the grid points (one soil sample at every 250 m interval) all over the microwatershed through land resource inventory in the year 2014 were analysed for pH, EC, organic carbon, available phosphorus and potassium and for micronutrients like zinc, copper, iron and manganese, and secondary nutrient sulphur.

Soil fertility data generated has been assessed and individual maps for all the nutrients for the microwatershed have been prepared. The village/survey number wise fertility data for the microwatershed is given in Appendix-II.

6.1 Soil Reaction (pH)

The soil fertility analysis of the Chincholi Khurd-1 microwatershed for soil reaction (pH) that about 57 ha (11%) area is slightly alkaline (pH 7.3-7.8) and is distributed in the northeastern and southeastern part of the microwatershed. Maximum area of about 437 ha (82%) area is moderately alkaline (pH 7.8-8.4) and is distributed in all parts of the microwatershed. About 28 ha (5%) area (Fig.6.1) is under strongly alkaline (pH 8.4-9.0) and is distributed in the southern, southeastern and southwestern part of the microwatershed.

6.2 Electrical Conductivity (EC)

The Electrical Conductivity of the soils of the entire microwatershed area is $<2 \text{ dSm}^{-1}$ (Fig 6.2) and as such soils are nonsaline.

6.3 Organic Carbon

The soil organic carbon content of the soils in the microwatershed is medium (0.5-0.75%) in 329 ha (62%) area that are distributed in all parts of the microwatershed (Fig.6.3). High ($>0.75\%$) organic carbon content accounts for 157 ha (30%) area in the microwatershed and is distributed in the central, northern and western part of the microwatershed. Low ($<0.5\%$) organic carbon content accounts for a small area of 36 ha (7%) in the microwatershed and is distributed in the southeastern parts of the microwatershed.




Fig.6.1 Soil Reaction (pH) map of Chincholi Khurd-1 microwatershed




Fig.6.2 Electrical Conductivity (EC) map of Chincholi Khurd-1 microwatershed

6.4 Available Phosphorus

The soil fertility analysis revealed that available phosphorus is low (<23 kg/ha) in 459 ha (86%) area (Fig.6.4) and is distributed in all parts of the microwatershed. There is an urgent need to increase the dose of phosphorous for all the crops by 25 per cent over the recommended dose to realize better crop performance. About 53 ha (10%) area in the microwatershed is medium (23-57 kg/ha) and is distributed in the southeastern part of the microwatershed. A small area of about 11 ha (2%) is high (>57 kg/ha) and is distributed in the southeastern part of the microwatershed.

6.5 Available Potassium

Available potassium content is medium (145-337 kg/ha) in 89 ha (17%) area distributed in central, eastern and western part of the microwatershed (Fig.6.5); high available potassium (>337 kg/ ha) content accounts for major area of 433 ha (81%) and is distributed in all parts of the microwatershed.

6.6 Available Sulphur

Available sulphur content is low (<10 ppm) in 99 ha (19%) area and is distributed in the central and western part of the microwatershed. Maximum area of about 359 ha (67%) is medium (10-20 ppm) in available sulphur and is distributed in all parts of the microwatershed (Fig.6.6). Available sulphur is high (>20 ppm) in small area of 65 ha (12%) and is distributed in the eastern parts of the microwatershed.

6.7 Available Boron

Available boron content is low (<0.5 ppm) in major area of about 445 ha (84%) and is distributed in all parts of the microwatershed. About 78 ha (15%) has soils that are medium (0.5-1.0 ppm) in available boron (Fig 6.7) and is distributed in the eastern and western part of the microwatershed. A very minute area of less than 1 ha is high (>1.0 ppm) in available boron.




Fig.6.3 Soil Organic Carbon map of Chincholi Khurd-1 microwatershed




Fig.6.4 Soil available Phosphorus map of Chincholi Khurd-1 microwatershed




Fig.6.5 Soil available Potassium map of Chincholi Khurd-1 microwatershed




Fig.6.6 Soil available Sulphur map of Chincholi Khurd-1 microwatershed




Fig.6.7 Soil available Boron map of Chincholi Khurd-1 microwatershed

6.8 Available Iron

Available iron content is deficient (<4.5 ppm) in a small area of 19 ha (3%) and is distributed in the eastern and western part of the microwatershed. It is sufficient in 504 ha (95%) area (Fig 6.8) and are distributed in all parts of the microwatershed.

6.9 Available Manganese

Available manganese content is sufficient (>1.0 ppm) in the entire microwatershed area (Fig 6.9).

6.10 Available Copper

Available copper content is sufficient (>0.2 ppm) in the entire microwatershed area (Fig 6.10).

6.11 Available Zinc

Available zinc content is deficient (<0.6 ppm) in major area of about 367 ha (69%) and is distributed in all parts of the microwatershed. It is sufficient (>0.6 ppm) in 155 ha (29%) area (Fig 6.11) and is distributed in the northwestern, southeastern and northern part of the microwatershed.




Fig.6.8 Soil available Iron map of Chincholi Khurd-1 microwatershed




Fig.6.9 Soil available Manganese map of Chincholi Khurd-1 microwatershed




Fig.6.10 Soil available Copper map of Chincholi Khurd-1 microwatershed




Fig.6.11 Soil available Zinc map of Chincholi Khurd-1 microwatershed

LAND SUITABILITY FOR MAJOR CROPS

The soil and land resource units (soil phases) of Chincholi Khurd-1 microwatershed were assessed for their suitability for growing food, fibre, fodder and horticulture crops by following the procedure as outlined in FAO, 1976 and 1983. Crop requirements were developed for each of the crop from the available research data, Naidu *et. al.* (2006) and Natarajan *et. al.* (2015). The crop requirements were matched with the soil and land characteristics (Table 7.1) to arrive at the crop suitability. In FAO land suitability classification, two orders are recognized. Order S- Suitable and Order N- Not suitable. The orders have classes, subclasses and units. Order-S has three classes, Class S1- Highly Suitable, Class S2- Moderately Suitable and Class S3- Marginally Suitable. Order N has two classes, N1- Currently not Suitable and N2- Permanently not Suitable. There are no subclasses within the class S1 as they will have very minor or no limitations for crop growth. Classes S2 and S3 are divided into subclasses based on the kinds of limitations encountered. The limitations that affect crop production are ‘c’ for erratic rainfall and its distribution and length of growing period (LGP), ‘e’ for erosion hazard, ‘r’ for rooting condition, ‘t’ for lighter or heavy texture, ‘g’ for gravelliness or stoniness, ‘n’ for nutrient availability, ‘l’ for topography, ‘m’ for moisture availability and ‘w’ for drainage. These limitations are indicated as lower case letters to the class symbol. For example, moderately suitable land with the limitations of soil depth and erosion is designated as S2re. For the microwatershed, the soil mapping units were evaluated and classified up to subclass level only; land suitability units are not worked out.

Using the above criteria, the soil map units of the microwatershed were evaluated and land suitability maps for 18 major annual and perennial crops were prepared. The detailed information on the kind of suitability of each of the soil phase for the crops assessed are given village/ survey number wise for the microwatershed in Appendix-III.

7.1 Land Suitability for Sorghum (*Sorghum bicolor*)

Sorghum is one of the major crops grown in Karnataka in an area of 11.02 lakh ha in northern Karnataka in Bijapur, Gulbarga, Raichur, Bidar, Belgaum, Dharwad and Bellary districts. The crop requirements for growing sorghum (Table 7.2) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and land suitability map for growing sorghum was prepared. The area extent and their geographic distribution of different suitability subclasses in the microwatershed is given in Figure. 7.1.

A small area of about 91 ha (17%) in the microwatershed has soils that are highly suitable (class S1) for growing sorghum crop. They are distributed mainly in the central, northern and small areas in southeastern part of the microwatershed. A very small area of about 16 ha (3%) is moderately suitable (class S2) for growing sorghum and are distributed in southwestern and eastern part of the microwatershed. They have major limitations of erosion and rooting depth.

Table 7.1 Soil-Site Characteristics of Chincholi Khurd-1 microwatershed

Soil Map Units	Climate (P) (mm)	Growing period (Days)	Drainage class	Soil depth (cm)	Soil texture		Gravelliness		AWC (mm/m)	Slope (%)	Erosion	pH	EC	ES P	CEC [Cmol (p ⁺) kg ⁻¹]	BS (%)
					Surf-ace	Sub-surface	Sur-face (%)	Subsurfa-ce (%)								
MGTiC3g1	786	150	WD	<25	sc	c	15-35	15-35	<50	3-5	Severe	6.8	0.3	0.2	46	100
MGTiD3g2	786	150	WD	<25	sc	c	35-60	15-35	<50	5-10	Severe	6.8	0.3	0.2	46	100
MGTmA1	786	150	WD	<25	c	c	-	15-35	<50	0-1	Slight	6.8	0.3	0.2	46	100
MGTmB1	786	150	WD	<25	c	c	-	15-35	<50	1-3	Slight	6.8	0.3	0.2	46	100
MGTmB2	786	150	WD	<25	c	c	-	15-35	<50	1-3	Moderate	6.8	0.3	0.2	46	100
MGTmB2g1	786	150	WD	<25	c	c	15-35	15-35	<50	1-3	Moderate	6.8	0.3	0.2	46	100
MGTmB3g1	786	150	WD	<25	c	c	15-35	15-35	<50	1-3	Severe	6.8	0.3	0.2	46	100
MGTmC2	786	150	WD	<25	c	c	-	15-35	<50	3-5	Moderate	6.8	0.3	0.2	46	100
MGTmC2g1	786	150	WD	<25	c	c	15-35	15-35	<50	3-5	Moderate	6.8	0.3	0.2	46	100
NHAmB1	786	150	WD	25-50	c	c	-	<15	51-100	1-3	Slight	7.2	0.1	0.3	40	100
NHAmB2	786	150	WD	25-50	c	c	-	<15	51-100	1-3	Moderate	7.2	0.1	0.3	40	100
BHImB1	786	150	WD	25-50	c	c	-	15-35	<50	1-3	Slight	7.0	0.1	0.2	28	100
BHImB2g1	786	150	WD	25-50	c	c	15-35	15-35	<50	1-3	Moderate	7.0	0.1	0.2	28	100
BHImC2g1	786	150	WD	25-50	c	c	15-35	15-35	<50	3-5	Moderate	7.0	0.1	0.2	28	100
DSImB2	786	150	MWD	50-75	c	c	-	<15	101-150	1-3	Moderate	7.0	0.1	0.3	62	100
RNLmB1	786	150	MWD	100-150	c	c	-	<15	>200	1-3	Slight	8.4	0.2	0.2	60	100
RNLmC2g1	786	150	MWD	100-150	c	c	15-35	<15	>200	3-5	Moderate	8.4	0.2	0.2	60	100
MANmA1	786	150	MWD	>150	c	c	-	<15	>200	0-1	Slight	8.3	0.2	0.1	58	100
MANmB1	786	150	MWD	>150	c	c	-	<15	>200	1-3	Slight	8.3	0.2	0.1	58	100
MANmB2	786	150	MWD	>150	c	c	-	<15	>200	1-3	moderate	8.3	0.2	0.1	58	100

*Symbols and abbreviations are according to Field Guide for LRI under Sujala-III Project, Karnataka

Marginally suitable lands (class S3) for growing sorghum occupy about 194 ha (36%) and occur in the western, central and southeastern part of the microwatershed. They have severe limitation of rooting depth. Major area of about 221 ha (42%) is not suitable for growing sorghum in the microwatershed and occur in all parts of the microwatershed.

Table 7.2 Crop suitability criteria for Sorghum

Crop requirement		Rating			
Soil –site characteristics	unit	Highly suitable (S1)	Moderately Suitable (S2)	Marginally suitable (S3)	Not suitable (N)
Slope	%	2-3	3-8	8-15	>15
LGP	Days	120-150	120-90	<90	
Soil drainage	class	Well to mod. drained	imperfect	Poorly/excessively	V. poorly
Soil reaction	pH	6.0-8.0	5.5-5.9 8.1-8.5	<5.5 8.6-9.0	>9.0
Surface soil texture	Class	C, cl, sicl, sc	l, sil, sic	Sl, ls	S, fragmental skeletal
Soil depth	Cm	100-75	50-75	30-50	<30
Gravel content	% vol.	5-15	15-30	30-60	>60
Salinity (EC)	dSm ⁻¹	2-4	4-8	8-10	>10
Sodicity (ESP)	%	5-8	8-10	10-15	>15




Fig. 7.1 Land Suitability map of Sorghum

7.2 Land Suitability for Maize (*Zea mays*)

Maize is the most important food crop grown in an area of 13.73 lakh ha in all the districts of Karnataka. The crop requirements for growing maize (Table 7.3) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing maize was generated. The area and geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.2.

The marginally suitable (class S3) lands cover a maximum area of about 302 ha (57%) and occur in all parts of the microwatershed. They have severe limitations of texture and rooting depth. About 221 ha (42%) area is not suitable for growing maize and occur in the southern, eastern and northwestern part of the microwatershed.

Table 7.3 Crop suitability criteria for Maize

Crop requirement		Rating			
Soil –site characteristics	unit	Highly suitable (S1)	Moderately Suitable (S2)	Marginally suitable (S3)	Not suitable (N)
Slope	%	<3	3.5	5-8	
LGP	Days	>100	100-80	60-80	
Soil drainage	class	Well drained	Mod. to imperfectly	Poorly/excessively	V.poorly
Soil reaction	pH	5.5-7.5	7.6-8.5	8.6-9.0	
Surface soil texture	Class	l, cl, scl, sil	Sl, sicl, sic	C(s-s), ls	S,fragmental
Soil depth	Cm	>75	50-75	25-50	<25
Gravel content	% vol.	<15	15-35	35-50	>50
Salinity (EC)	dSm ⁻¹	<1.0	1.0-2.0	2.0-4.0	
Sodicity (ESP)	%	<10	10-15	>15	




Fig. 7.2 Land Suitability map of Maize

7.3 Land Suitability for Red gram (*Cajanus cajan*)

Red gram is one of the major pulse crop grown in an area of 8.23 lakh ha in northern Karnataka in Bijapur, Gulbarga, Raichur, Bidar, Belgaum, Dharwad and Bellary districts. The crop requirements for growing red gram (Table 7.4) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and land suitability map for growing red gram was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed is given in Figure 7.3.

An area of about 107 ha (20%) is moderately suitable (class S2) for red gram and is distributed in the northeastern, northern and southwestern part of the microwatershed. They have major limitations of rooting depth, erosion and texture. Marginally suitable lands (class S3) for growing red gram occupy about 194 ha (36%) and occur in all parts of the microwatershed. They have severe limitations of rooting depth. An area of about 221 ha (42%) is not suitable for growing red gram in the microwatershed and occur in the northwestern, eastern and southern part of the microwatershed.

Table 7.4 Crop suitability criteria for Red gram

Crop requirement		Rating			
Soil characteristics	–site unit	Highly suitable (S1)	Moderately Suitable (S2)	Marginally suitable (S3)	Not suitable (N)
Slope	%	<3	3-5	5-10	>10
LGP	Days	>210	180-210	150-180	<150
Soil drainage	class	Well drained	Mod. to well drained	Imperfectly drained	Poorly drained
Soil reaction	pH	6.5-7.5	5.0-6.5 7.6-8.0	8.0-9.0	>9.0
Surface soil texture	Class	l, scl, sil, cl, sl	sicl, c(m)	sic, ls	S, fragmental
Soil depth	Cm	>100	85-100	40-85	<40
Gravel content	% vol.	<20	20-35	35-60	>60
Salinity (EC)	dSm ⁻¹	<1.0	1.0-2.0	>2.0	
Sodicity (ESP)	%	<10	10-15	>15	

Table 7.5 Crop suitability criteria for Sunflower

Crop requirement		Rating			
Soil characteristics	–site unit	Highly suitable (S1)	Moderately Suitable (S2)	Marginally suitable (S3)	Not suitable (N)
Slope	%	<3	3-5	5-10	>10
LGP	Days	>90	80-90	70-80	<70
Soil drainage	class	Well drained	mod. drained	Well imperfectly drained	Poorly drained
Soil reaction	pH	6.5-8.0	8.1-8.5 5.5-6.4	8.6-9.0; 4.5-5.4	>9.0 <4.5
Surface soil texture	Class	l, cl, sil, sc	Scl, sic, c,	c (>60%), sl	ls, s
Soil depth	Cm	>100	75-100	50-75	<50
Gravel content	% vol.	<15	15-35	35-60	>60
Salinity (EC)	dSm ⁻¹	<1.0	1.0-2.0	>2.0	
Sodicity (ESP)	%	<10	10-15	>15	




Fig. 7.4 Land Suitability map of Sunflower

7.5 Land Suitability for Cotton (*Gossypium hirsutum*)

Cotton is the most important fibre crop grown in the State in about 6.6 lakh ha area in Raichur, Dharwad, Belgaum, Gulbarga, Bijapur, Bidar, Bellary, Chitradurga and Chamarajnagar districts. The crop requirements for growing cotton (Table 7.6) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing cotton was generated and the area extent and geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.5.

Highly suitable (class S1) lands are found to occur in an area of 91 ha (17%) and distributed in the northern and eastern part of the microwatershed.

Moderately suitable (class S2) lands are found to occur in a very minor area of about 16 ha (3%). The soils have moderate limitations of erosion and rooting depth. They are dominantly distributed in the southwestern and eastern part of the microwatershed. The marginally suitable (class S3) lands cover about 194 ha (36%) area in the microwatershed and mainly occur in the central, western and southeastern part of the microwatershed. They have severe limitations of rooting depth. Major area of about 221 ha (42%) area is not suitable for growing cotton and are distributed in all parts of the microwatershed.

Table 7.6 Crop suitability criteria for Cotton

Crop requirement		Rating			
Soil-site characteristics	unit	Highly suitable (S1)	Moderately Suitable (S2)	Marginally suitable (S3)	Not suitable (N)
Slope	%	1-2	2-3	3-5	>5
LGP	Days	180-240	120-180	<120	
Soil drainage	class	Well to moderately well	imperfectly drained	Poor somewhat excessive	Stagnant/excessive
Soil reaction	pH	6.5-7.5	7.6-8.0	8.1-9.0	>9.0 >6.5
Surface soil texture	Class	Sic, c	Sicl, cl	Si, sil, sc, scl, l	S1, s,ls
Soil depth	Cm	100-150	60-100	30-60	<30
Gravel content	% vol.	<5	5-10	10-15	15-35
CaCO ₃ in root zone	%	<3	3-5	5-10	10-20
Salinity (EC)	dSm ⁻¹	2-4	4.0-8.0	8.0-12	>12
Sodicity (ESP)	%	5-10	10-20	20-30	>30




Fig. 7.5 Land Suitability map of Cotton

7.6 Land Suitability for Sugarcane (*Saccharum officinarum*)

Sugarcane is the most important commercial crop grown in 6.7 lakh ha area in Gulbarga, Bijapur, Bagalkot, Bidar, Mysore and Mandya districts. The crop requirements for growing sugarcane (Table 7.7) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing sugarcane was generated and the area extent and geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.6.

The marginally suitable (class S3) lands cover about 108 ha (20%) area in the microwatershed and mainly occur in the northern and northeastern part of the microwatershed. They have severe limitations of texture.

Major area of about 415 ha (78%) is not suitable for growing sugarcane and occur in all parts of the microwatershed.

Table 7.7 Crop suitability criteria for Sugarcane

Land use requirement		Rating			
Soil-site characteristics	unit	Highly suitable (S1)	Moderately Suitable (S2)	Marginally suitable (S3)	Not suitable (N)
Slope	%	<3	3-5	5-8	>8
Soil drainage	class	Well drained	Mod./imperfectly drained	Poorly drained	V.poor/excessively drained
Soil reaction	pH	7.0-8.0	6.0-6.9 8.1-9.0	4.0-5.9 9.1-9.5	<4.0/ >9.5
Surface soil texture	Class	l, cl, sil, sicl	C(m/k), sl	C+(ss)	
Soil depth	cm	>100	100-75	75-50	<50
stoniness	%	<15	15-35	35-50	>50
Salinity (EC)	dSm ⁻¹	<2.0	2.0-4.0	4.0-9.0	>9
Sodicity (ESP)	%	<10	10-15	15-25	>25




Fig. 7.6 Land Suitability map of Sugarcane

7.7 Land Suitability for Soybean (*Glycine max*)

Soybean is the most important pulse and oil seed crop grown in about 1.68 lakh ha area in Bijapur, Raichur, Gulbarga, Dharwad, Belgaum and Bellary all the districts of the state. The crop requirements for growing soybean were matched with the soil-site characteristics and a land suitability map for growing soybean was generated and the area extent and geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.7.

Highly suitable (class S1) lands are found to occur in an area of 91 ha (17%) and are distributed in the northern and eastern part of the microwatershed. Moderately suitable (class S2) lands are found to occur in small area of about 16 ha (3%). The soils have moderate limitations of erosion and rooting depth. They are dominantly distributed in the eastern and southwestern part of the microwatershed.

The marginally suitable (class S3) lands cover about 194 ha (36%) area in the microwatershed and mainly occur in the northern, central, western and southeastern part of the microwatershed. They have severe limitations of rooting depth. Major area of about 221 ha (42%) area is not suitable for growing soybean and occur in all parts of the microwatershed.




Fig. 7.7 Land Suitability map of Soybean

7.8 Land Suitability for Guava (*Psidium guajava*)

Guava is the most important fruit crop grown in the State in Raichur, Dharwad, Belgaum, Gulbarga, Bijapur, Bidar, Bellary, Chitradurga and Chamarajnar districts. The crop requirements for growing guava (Table 7.8) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing guava was generated and the area extent and geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.8.

The marginally suitable (class S3) lands cover a small area of about 23 ha (4%) in the microwatershed and mainly occur in the eastern and southwestern part of the microwatershed. They have severe limitations of texture, slope and rooting depth. Major area of about 500 ha (94%) is not suitable for growing guava and occur in all parts of the microwatershed.

Table 7.8 Crop suitability criteria for Guava

Crop requirement			Rating			
Soil –site characteristics		unit	Highly suitable (S1)	Moderately Suitable (S2)	Marginally suitable (S3)	Not suitable (N)
climate	Temperature in growing season	⁰ C	28-32	33-36 24-27	37-42 20-23	
Soil moisture	Growing period	Days	>150	120-150	90-120	<90
Soil aeration	Soil drainage	class	Well drained	Mod. to imperfectly	poor	Very poor
Nutrient availability	Texture	Class	Scl, l, cl, sil	Sl,sicl,sic.,sc,c	C (<60%)	C (>60%)
	pH	1:2.5	6.0-7.5	7.6-8.0:5.0-5.9	8.1-8.5:4.5-4.9	>8.5:<4.5
	CaCO ₃ in root zone	%	Non calcareous	<10	10-15	>15
Rooting conditions	Soil depth	cm	>100	75-100	50-75	<50
	Gravel content	% vol.	<15	15-35	>35	
Soil toxicity	Salinity	dS/m	<2.0	2.0-4.0	4.0-6.0	
	Sodicity	%	Non sodic	10-15	15-25	>25
Erosion	Slope	%	<3	3-5	5-10	>10




Fig 7.8 Land Suitability map of Guava

7.9 Land Suitability for Mango (*Mangifera indica*)

Mango is the most important fruit crop grown in all the districts of the State. The crop requirements for growing mango (Table 7.9) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing mango was generated and the area and geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.9.

The marginally suitable (class S3) lands cover a very small area of about 16 ha (3%) and mainly occur in the eastern part of the microwatershed. They have severe limitations of texture and slope. Major area of about 507 ha (95%) is not suitable for growing mango and occur in all parts of the microwatershed.

Table 7.9 Crop suitability criteria for Mango

Crop requirement			Rating			
soil-site characteristics		unit	Highly suitable (S1)	Moderately Suitable (S2)	Marginally suitable (S3)	Not suitable (N)
climate	Temp in growing season	⁰ C	28-32	24-27 33-35	36-40	20-24
	Min. temp. before flowering	⁰ C	10-15	15-22	>22	
Soil moisture	Growing period	Days	>180	150-180	120-150	<120
Soil aeration	Soil drainage	class	Well drained	Mod. To imperfectly drained	Poor drained	Very poorly drained
	Water table	M	>3	2.50-3.0	2.5-1.5	<1.5
Nutrient availability	Texture	Classes	Sc, l, sil, cl	Sl, sc, sic, l, c	C (<60%)	C (>60%),
	pH	1:2.5	5.5-7.5	7.6-8.55.0-5.4	8.6-9.0 4.0-4.9	>9.0 <4.0
	OC	%	High	medium	low	
	CaCO ₃ in root zone	%	Non calcareous	<5	5-10	>10
Rooting conditions	Soil depth	cm	>200	125-200	75-125	<75
	Gravel content	% vol	Non gravelly	<15	15-35	>35
Soil toxicity	Salinity	dS/m	Non saline	<2.0	2.0-3.0	>3.0
	Sodicity	%	Non sodic	<10	10-15	>15
Erosion	Slope	%	<3	3-5	5-10	




Fig. 7.9 Land Suitability map of Mango

7.10 Land Suitability for Sapota (*Manilkara zapota*)

Sapota is the most important fruit crop grown in almost all the districts of the state. The crop requirements for growing sapota (Table 7.10) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing sapota was generated and the area extent and geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.10.

The marginally suitable (class S3) lands cover a small area of about 23 ha (4%) and mainly occur in the eastern and southwestern part of the microwatershed. They have severe limitations of rooting depth, texture and slope. Major area of about 500 ha (94%) is not suitable for growing sapota and occur in all parts of the microwatershed.

Table 7.10 Crop suitability criteria for Sapota

Crop requirement			Rating			
Soil –site characteristics		unit	Highly suitable (S1)	Moderately Suitable (S2)	Marginally suitable (S3)	Not suitable (N)
climate	Temperature in growing season	° C	28-32	33-36 24-27	37-42 20-23	>42 <18
Soil moisture	Growing period	Days	>150	120-150	90-120	<120
Soil aeration	Soil drainage	class	Well drained	Moderately well drained	Imperfectly drained	Poorly drained
Nutrient availability	Texture	Class	Scl, l, cl, sil	Sl, sicl, sc	C (<60%)	ls, s, C (>60%)
	pH	1:2.5	6.0-7.5	7.6-8.0 5.0-5.9	8.1-9.0 4.5-4.9	>9.0 <4.5
	CaCO ₃ in root zone	%	Non calcareous	<10	10-15	>15
Rooting conditions	Soil depth	cm	>150	75-150	50-75	<50
	Gravel content	% vol.	Non gravelly	<15	15-35	<35
Soil toxicity	Salinity	dS/m	Non saline	Up to 1.0	1.0-2.0	2.0-4.0
	Sodicity	%	Non sodic	10-15	15-25	>25
Erosion	Slope	%	<3	3-5	5-10	>10




Fig. 7.10 Land Suitability map of Sapota

7.11 Land Suitability for Jackfruit (*Artocarpus heterophyllus*)

Jackfruit is the most important fruit crop grown in almost all the districts of the state. The crop requirements for growing jackfruit were matched with the soil-site characteristics and a land suitability map for growing jackfruit was generated and the area extent and geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.11.

The marginally suitable (class S3) lands cover a small area of about 23 ha (4%) and mainly occur in the eastern and southwestern part of the microwatershed. They have severe limitations of rooting depth, texture and slope. Major area of about 500 ha (94%) is not suitable for growing jackfruit and occur in all parts of the microwatershed.




Fig 7.11 Land Suitability map of Jackfruit

7.12 Land Suitability for Jamun (*Syzygium cumini*)

Jamun is the most important fruit crop grown in almost all the districts of the state. The crop requirements for growing jamun were matched with the soil-site characteristics and a land suitability map for growing jamun was generated and the area extent and geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.12.

Moderately suitable (class S2) lands are found to occur in a very small area of about 16 ha (3%). The soils have moderate limitations of texture and slope. They are dominantly distributed in eastern part of the microwatershed

The marginally suitable (class S3) lands cover about a very minor area of 7 ha (1%) and mainly occur in the southwestern part of the microwatershed. They have severe limitations of rooting depth. Major area of about 500 ha (94%) is not suitable for growing jamun and occur in all parts of the microwatershed.




Fig 7.12 Land Suitability map of Jamun

7.13 Land Suitability for Musambi (*Citrus limetta*)

Musambi is the most important fruit crop grown in almost all the districts of the state. The crop requirements for growing musambi were matched with the soil-site characteristics and a land suitability map for growing musambi was generated and the area extent and geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.13.

Highly suitable (class S1) lands are found to occur in an area of about 91 ha (17%) and are distributed in the northern part of the microwatershed.

Moderately suitable (class S2) lands are found to occur in a very minor area of about 9 ha (2%). The soils have moderate limitations of erosion and slope. They are dominantly distributed in the eastern part of the microwatershed.

The marginally suitable (class S3) lands also cover a very small area of about 7 ha (1%) in the microwatershed and mainly occur in the southwestern part of the microwatershed. They have severe limitations of rooting depth. Major area of about 415 ha (78%) is not suitable for growing musambi and occur in all parts of the microwatershed.




Fig 7.13 Land Suitability map of Musambi

7.14 Land Suitability for Lime (*Citrus sp*)

Lime is the most important fruit crop grown in almost all the districts of the state. The crop requirements for growing Lime (Table 7.11) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing Lime was generated and the area extent and geographic distribution of different suitability subclasses in the microwatershed is given in Figure 7.14.

Highly suitable (class S1) lands are found to occur in an area of about 91 ha (17%) and are distributed in the northern part of the microwatershed. Moderately suitable (class S2) lands are found to occur in a very minor area of about 9 ha (2%). The soils have moderate limitations of slope. They are dominantly distributed in the eastern part of the microwatershed.

The marginally suitable (class S3) lands cover a very small area of about 7 ha (1%) in the microwatershed and occur in the southwestern part of the microwatershed. They have severe limitations of rooting depth. Major area of about 415 ha (78%) is not suitable for growing lime and occur in all parts of the microwatershed.

Table 7.11 Crop suitability criteria for Lime

Land use requirement			Rating			
Soil –site characteristics		unit	Highly suitable (S1)	Moderately Suitable (S2)	Marginally suitable (S3)	Not suitable (N)
Climate	Temp in growing season	⁰ C	28-30	31-35 24-27	36-40 20-23	>40 <20
	Soil moisture	Growing period	Days	240-265	180-240	150-180 <150
Soil aeration	Soil drainage	class	Well drained	Mod. to imperfectly drained	poorly	Very poorly
Nutrient availability	Texture	Class	Scl, l, sicl, cl, s	Sc, sc, c	C (>70%)	S, ls
	pH	1:2.5	6.0-7.5	5.5-6.4/ 7.6-8.0	4.0-5.4 8.1-8.5	<4.0 >8.5
	CaCO ₃ in root zone	%	Non calcareous	Upto 5	5-10	>10
Rooting condition	Soil depth	cm	>150	100-150	50-100	<50
	Gravel content	% vol.	Non gravelly	15-35	35-55	>55
Soil toxicity	Salinity	dS/m	Non saline	Upto 1.0	1.0-2.5	>2.5
	Sodicity	%	Non sodic	5-10	10-15	>15
Erosion	Slope	%	<3	3-5	5-10	




Fig 7.14 Land Suitability map of Lime

7.15 Land Suitability for Cashew (*Anacardium occidentale*)

Cashew is the most important plantation crop grown in almost all the districts. The crop requirements for growing Cashew were matched with the soil-site characteristics and a land suitability map for growing Cashew was generated and the area and geographic distribution of different suitability subclasses in the microwatershed is given in Figure 7.15. The entire area is not suitable for growing cashew in the microwatershed.




Fig 7.15 Land Suitability map of Cashew

7.16 Land Suitability for Custard Apple (*Annona reticulata*)

Custard apple is the most important fruit crop grown in almost all the districts of the state. The crop requirements for growing custard apple were matched with the soil-site characteristics and a land suitability map for growing custard apple was generated and the area extent and geographic distribution of different suitability subclasses in the microwatershed is given in Figure 7.16.

Highly suitable (class S1) lands are found to occur in an area of 91 ha (17%) and are distributed in the northern part of the microwatershed.

Moderately suitable (class S2) lands are found to occur in a small area of about 16 ha (3%). The soils have moderate limitations of slope and rooting depth. They are dominantly distributed in the southwestern and eastern part of the microwatershed.

The marginally suitable (class S3) lands cover about 90 ha (17%) area in the microwatershed and occur in the southeastern and western part of the microwatershed. They have severe limitations of rooting depth. Major area of about 325 ha (61%) is not suitable for growing custard apple and occur in all parts of the microwatershed.




Fig 7.16 Land Suitability map of Custard Apple

7.17 Land Suitability for Amla (*Phyllanthus emblica*)

Amla is the most important fruit crop grown in almost all the districts of the state. The crop requirements for growing amla were matched with the soil-site characteristics and a land suitability map for growing amla was generated and the area extent and geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.17.

Highly suitable (class S1) lands are found to occur in an area of 91 ha (17%) and are distributed in the northern part of the microwatershed.

Moderately suitable (class S2) lands are found to occur in a small area of about 16 ha (3%). The soils have moderate limitations of slope and rooting depth. They are dominantly distributed in the southwestern and eastern part of the microwatershed.

The marginally suitable (class S3) lands cover about 90 ha (17%) area in the microwatershed and occur in the southeastern and western part of the microwatershed. They have severe limitations of rooting depth. Major area of about 325 ha (61%) is not suitable for growing amla and occur in all parts of the microwatershed.




Fig 7.17 Land Suitability map of Amla

7.18 Land Suitability for Tamarind (*Tamarindus indica*)

Tamarind is the most important spice crop raised in almost all the districts of the state. The crop requirements for growing tamarind were matched with the soil-site characteristics and a land suitability map for growing tamarind was generated and the area extent and geographic distribution of different suitability subclasses in the microwatershed is given in Figure 7.18.

Moderately suitable (class S2) lands are found to occur in a very minor area of about 16 ha (3%). The soils have moderate limitations of slope and rooting depth. They are dominantly distributed in the eastern part of the microwatershed. Major area of about 507 ha (95%) is not suitable for growing tamarind and occurs all parts of the microwatershed.




Fig 7.18 Land Suitability map of Tamarind

7.19 Land Management Units (LMUs)

The 20 soil map units identified in Chincholi Khurd-1 microwatershed have been regrouped into 5 Land Management Units (LMU's) for the purpose of preparing Proposed Crop Plan. The map units that have been grouped into 5 land management units along with brief description of soil and site characteristics are given below.

LMUs	Soil map units	Soil and site characteristics
1	MGTiC3g1, MGTiD3g2, MGTmB3g1	Very shallow, black soils with slopes of 1-10%, gravelly to very gravelly (15-60%) and severe erosion
2	MGTmA1, MGTmB1, MGTmB2, MGTmB2g1, MGTmC2, MGTmC2g1	Very shallow, black soils with slopes of <1-5%, gravelly (15-35%) and slight to moderate erosion
3	NHAmB1, NHAmB2, BHImB1, BHImB2g1, BHImC2g1	Shallow, black soils with slopes of 1-5 %, gravelly (15-35%) and slight to moderate erosion
4	DSImB2	Moderately shallow, black soils with slopes of 1-3 % and moderate erosion
5	RNLmB1, RNLmC2g1, MANmA1, MANmB1, MANmB2	Deep to very deep, black soils with slopes of 1-5 %, gravelly (15-35%) and slight to moderate erosion

Land Management Units are grouped based on the similarities in respect of the type of soil, the depth of the soil, the surface soil texture, gravel content, AWC, slope, erosion etc. and a Land Management Units map (Fig.7.19) has been prepared. These Land Management Units are expected to behave similarly for a given level of management.




Fig. 7.19 Land Management Units map - Chincholi Khurd-1 microwatershed

7.20 Proposed Crop Plan for Chincholi Khurd-1 microwatershed

After assessing the land suitability for the 18 crops, a proposed crop plan has been prepared for the 5 identified LMUs by considering only the highly (class S1) and moderately suitable (class S2) lands for each of the eighteen crops. The resultant proposed crop plan is presented below in Table 7.12

Table 7.12 Proposed Crop Plan for Chincholi Khurd - 1 microwatershed

LMUs No	Mapping Units	Survey Number	Field crops	Forestry Crop/Grasses	Horticulture crops (Rainfed Condition)	Horticulture crops with suitable intervention	Recommended Interventions
LMU 1	1, 2, 7 (<25 cm)	Chincholi: 12 Padsavli: 220,268,269, 271, 274	Horse gram	Neem, Glyricydia Silviculture, Agave, Simaroba	-	-	Crescent bunds
LMU 2	3, 4, 5, 6, 8, 9 (<25 cm)	Chincholi: 19,20,21,22, 25,26,27 Padsavli: 211,219,221, 222,223,229/1,229/2, 242/1,242/2,243,267 Sarasamba: 30,31,32,33,34	Horse gram	Neem, Glyricydia Silviculture, Agave, Simaroba	-	-	Crescent bunds

To be continued...

LMU 3	10, 11, 12, 13, 14 (25-50 cm)	Chincholi: 5,6,7,8,9,10, 11,15,16,17,18,28,29 Padsavli: 224,225,244/1, 244/2,245,246/1,246/2,27 0,272,273 Sarasamba: 25,26,28	Bajra, Linseed, Green gram, Black gram, Chick pea	Subabhul, Neem, Teak	Custard apple, Charoli, Ber, Amla Vegetables: Ladies finger, Brinjal, Cowpea, Flowers: Marigold, Chrysanthemum	Custard apple, Charoli, Ber, Amla Vegetables: Onion, Tomato, Brinjal, Chillies, Bhendi Flowers: Marigold, Chrysanthemum	Drip irrigation, suitable soil and water conservations like cultivation on raised beds with mulches and drip
LMU 4	15 (50-75 cm)	Sarasamba: 29	Sorghum, Cotton, Red Gram, Black gram, Green gram, Soybean, Sesame, Sunflower, Safflower Rabi: Sorghum, Chickpea	Subabhul, Neem, Teak	Custard apple, Charoli, Ber, Amla Vegetables: Ladies finger, Brinjal, Cowpea, Flowers: Marigold, Chrysanthemum	Custard apple, Charoli, Ber, Amla, Papaya, Banana, Lime, Citrus Vegetables: Onion, Tomato, Brinjal, Chillies, Bhendi Flowers: Marigold, Chrysanthemum	Graded bunds, Strengthening of field bunds
LMU 5	16, 17, 18, 19, 20 (100-150 & >150 cm)	Chincholi: 3,4,13,14 Padsavli: 4,226,227,228, 230/1,230/2,231,232,233,2 34,235,236,237,238,239,2 40,241	Sorghum, Cotton, Red Gram Black gram, Green gram, Soybean, Sesame, Sunflower, Safflower, Rabi: Sorghum, Chickpea	-	Vegetables: Ladies finger, Brinjal, Cowpea, coriander Field crops: Sorghum, Cotton, Red Gram, Sunflower, Safflower, Perennial component: Guava, Tamarind, Sapota, Lime, Mosambi Flowers: Marigold, Chrysanthemum	Banana, Papaya, Lime. Mosambi, Guava, Tamrind Vegetables: Onion, Tomato, Brinjal, Chillies, Bhendi Flowers: Marigold, Chrysanthemum	Graded bunds, Strengthening of field bunds

SOIL HEALTH MANAGEMENT

8.1 Soil Health

Soil is fundamental to crop production. Without soil, no food could be produced nor would livestock be fed on a large scale. Because it is finite and fragile, soil is a precious resource that requires special care from its users.

Soil health or the capacity of the soil to function is critical to human survival. Soil health has been defined as: “the capacity of the soil to function as a living system without adverse effect on the ecosystem”. Healthy soils maintain a diverse community of soil organisms that help to form beneficial symbiotic associations with plant roots, recycle essential plant nutrients, improve soil structure with positive repercussions for soil, water and nutrient holding capacity and ultimately improve crop production and also contribute to mitigating climate change by maintaining or increasing its carbon content.

Functional interactions of soil biota with organic and inorganic components, air and water determine a soil’s potential to store and release nutrients and water to plants and to promote and sustain plant growth. Thus, maintaining soil health is vital to crop production and conserve soil resource base for sustaining agriculture.

The most important characteristics of a healthy soil are

- Good soil tilth
- Sufficient soil depth
- Good water storage and good drainage
- Adequate supply, but not excess of nutrients
- Large population of beneficial organisms
- Small proportion of plant pathogens and insect pests
- Low weed pressure
- Free of chemicals and toxins that may harm the crop
- Resistance to degradation
- Resilience when unfavourable conditions occur

Characteristics of Chincholi Khurd-1 microwatershed

- The soil phases with sizeable area identified in the microwatershed belonged to the soil series of MGT (221 ha), BHI (104 ha), NHA (90 ha), MAN (86 ha), RNL (16 ha) and DSI (7 ha). As per land capability classification, nearly 98 per cent area comes under arable land category (Class II, III and IV) and two per cent area belongs to nonarable land category. The major limitations identified in the arable lands were soil and erosion.
- On the basis of soil reaction, about 437 ha (82%) of total area is moderately alkaline (pH 7.8-8.4) followed by strongly alkaline (pH 8.4-9.0) 28 ha (5%) and slightly alkaline (pH 7.3-7.8) 57 ha (11%) of the area. Thus, about 98 per cent of the soils are alkaline in reaction.

Soil Health Management

The following actions are required to improve the current land husbandry practices that provide a sound basis for the successful adoption of sustainable crop production system.

Alkaline soils

(Slightly alkaline to moderately alkaline soils)

1. Regular addition of organic manure, green manuring, green leaf manuring, crop residue incorporation and mulching needs to be taken up to improve the soil organic matter status.
2. Application of biofertilizers (Azospirillum, Azotobacter, Rhizobium).
3. Application of 25% of extra N and P (125 % RDN&P).
4. Application of ZnSO₄ – 12.5 kg/ha (once in three years).
5. Application of Boron – 5kg/ha (once in three years).

Neutral soils

1. Regular addition of organic manure, green manuring, green leaf manuring, crop residue incorporation and mulching needs to be taken up to improve the soil organic matter status.
2. Application of biofertilizers, (Azospirillum, Azotobacter, Rhizobium).
3. Application of 100 per cent RDF.
4. Need based micronutrient applications.

Besides the above recommendations, the best transfer of technology options are also to be adopted.

Disseminate information and communicate benefits. Any large scale implementation of soil health management requires that supporting information is made available widely, particularly through channels familiar to farmers and extension workers. Given the very high priority attached to soil health especially by the Central Government on issuing Soil-Health Cards to all the farmers, media outlets like regional, state and national newspapers and radio programs in local languages but also modern information and communication technologies such as cellular phones and the Internet, which can be much more effective in reaching younger farmers.

Inputs for Net Planning and Interventions needed

Net planning in IWMP is focusing on preparation of

1. Soil and Water Conservation plan for each plot or farm.
2. Productivity enhancement measures/ interventions for existing crops/livestock/other farm enterprises.
3. Diversification of farming mainly with perennial horticultural crops and livestock.
4. Improving livelihood opportunities and income generating activities.

In this connection, how various outputs of Sujala-III are of use in addressing these objectives of Net Planning are briefly presented.


- ❖ **Soil Depth:** The depth of a soil decides the amount of moisture and nutrients it can hold, what crops can be taken up or not, depending on the rooting depth and the length of growing period available for raising any crop. Deeper the soil, better for a wide variety of crops. If sufficient depth is not available for growing deep rooted crops, deeper planting pits need to be opened and additional good quality soil brought from outside has to be filled into the planting pits.
- ❖ **Surface soil texture:** Lighter soil texture in the top soil means, better rain water infiltration, less run-off and soil moisture conservation, less capillary rise and less evaporation losses. Lighter surface textured soils are highly suitable for crops like groundnut, root vegetables (carrot, radish, potato etc) but not ideal for crops that need stagnant water like lowland paddy. Heavy textured soils are poor in water infiltration and percolation. They are prone for sheet erosion; such soils can be improved by sand mulching. The technology that is developed by the AICRP-Dryland Agriculture, Vijayapura, Karnataka may be adopted.
- ❖ **Gravelliness:** More gravel content is favorable for run-off harvesting but poor in soil moisture storage and nutrient availability. It is a significant parameter that decides the kind of crop to be raised.
- ❖ **Land Capability Classification:** The land capability map shows the areas suitable and not suitable for agriculture and the major constraints in each of the plot/survey number. Hence, one can decide what kind of enterprise is possible in each of these units. In general, erosion and soil are the major constraints in Chincholi Khurd-1 watershed.
- ❖ **Organic Carbon:** In about 36 ha (7%) area, the OC content is low (<0.5%), about 329 ha (62%) area, the OC content is medium (0.5-0.75%) and in about 157 ha (30%) area it is high (>0.75%). The areas that are low and medium in OC needs to be further improved by applying farmyard manure and rotating crops with cereals and legumes or mixed cropping.
- ❖ **Promoting green manuring:** Growing of green manuring crops costs Rs. 1250/ha (green manuring seeds) and about Rs. 2000/ha towards cultivation that totals to Rs. 3250/- per ha. On the other hand, application of organic manure @ 10 tons/ha costs Rs. 5000/ha. The practice needs to be continued for 2-3 years or more. Nitrogen fertilizer needs to be supplemented by 25% in addition to the recommended level in 65 ha area where OC is less than 0.5-0.75%. For example, for rainfed maize, recommended level is 50 kg N per ha and an additional 12 kg /ha needs to be applied for all the crops grown in these plots.

- ❖ **Available Phosphorus:** In 459 ha (86%), the available phosphorus is low, about 53 ha (10%) area is medium in available phosphorus, Hence for all the crops, 25% additional P-needs to be applied and about 11 ha (2%) area is high in available phosphorus.
 - ❖ **Available Potassium:** Available potassium is medium in 89 ha (17%) area of the microwatershed. Hence, in all these plots, for all crops, an additional 25 % potassium can be applied. It is high in 433 ha (81%) area of the microwatershed.
 - ❖ **Available Sulphur:** Available sulphur is a very critical nutrient for oilseed crops. It is low in 99 ha (19%) area of the microwatershed, medium in 359 ha (67%).These areas need to be applied with magnesium sulphate or gypsum or Factamphos (p) fertilizer (13% sulphur) for 2-3 years for the deficiency to be corrected. About 65 ha (12 %) area has soils that are high in available sulphur.
 - ❖ **Available iron:** It is deficient in 19 ha (3%) area of the microwatershed. To manage iron deficiency, iron sulphate @ 25kg /ha needs to be applied for 2-3 years and sufficient in 504 ha (95 %) area in the microwatershed.
 - ❖ **Available Zinc:** It is deficient in 367 ha (69%) area of the microwatershed. Application of zinc sulphate @25kg/ha is to be applied and sufficient in 155 ha (29%) area in the microwatershed.
 - ❖ **Soil alkalinity:** The microwatershed has 328 ha area with soils that are moderately to strongly alkaline. These areas need application of gypsum and wherever calcium is in excess, iron pyrites and element sulphur can be recommended. Management practices like treating repeatedly with good quality water to drain out the excess salts, growing of salt tolerant crops like Casuarina, Acasia, Neem, Ber etc. are recommended.
- Land Suitability for various crops:** Areas that are highly, moderately and marginally suitable for growing various crops are indicated. Along with the suitability, various constraints that are limiting the productivity are also indicated. For example, in case of cotton, gravel content, rooting depth and salinity/alkalinity are the major constraints in various plots. With suitable management interventions, the productivity can be enhanced. In order to increase water holding capacity of light textured soils, growing of green manure crops and application of organic manure is recommended.

SOIL AND WATER CONSERVATION TREATMENT PLAN

For preparing soil and water conservation treatment plan for Chincholi Khurd-1 microwatershed, the land resource inventory database generated under Sujala-III project has been transformed as information through series of interpretative (thematic) maps using soil phase map as a base. The various thematic maps (1:7920 scale) generated were

- Soil depth
- Surface soil texture
- Available water capacity
- Soil slope
- Soil gravelliness
- Land capability
- Present land use & land cover
- Crop suitability maps
- Rainfall map
- Hydrology
- Water Resources
- Socio-economic data
- Contour plan with existing features- Network of water ways, pothissa boundaries, cut up/ minor terraces etc.
- Cadastral map (1:7920 scale)
- Satellite imagery (1:7920 scale)



Apart from these, Hand Level/ Hydro Marker/ Dumpy Level/ Total Station and Kathedars' List has to be collected.

Steps for Survey and Preparation of Treatment Plan

The boundaries of Land user Groups' and Survey No. boundaries are traced in the field.

Naming of user groups and farmers

- Identification of arable and non arable lands
- Identification of drainage lines and gullies
- Identification of non treatable areas
- Identification of priority areas in the arable lands
- Treatment plan for arable lands
- Location of water harvesting and recharge structures

9.1 Treatment Plan

The treatment plan recommended for arable lands is briefly described below


9.1.1 Arable Land Treatment

A. BUNDING

Steps for Survey and Preparation of Treatment Plan		USER GROUP-1
Cadastral map (1:7920 scale) is enlarged to a scale of 1:2500 scale		
Existing network of waterways, pottissa boundaries, grass belts, natural drainage lines/ watercourse, cut ups/ terraces are marked on the cadastral map to the scale		
Drainage lines are demarcated into		
Small gullies	(up to 5 ha catchment)	
Medium gullies	(5-15 ha catchment)	
Ravines	(15-25 ha catchment) and	
Halla/Nala	(more than 25ha catchment)	

Measurement of Land Slope

Land slope is estimated or determined by the study and interpretation of contours or by measurement in the field using simple instruments like Hand level or hydrometer.



Vertical and Horizontal intervals between bunds as recommended by the Watershed Development

Slope percentage	Vertical interval (m)	Corresponding Horizontal Distance (m)
2 - 3%	0.6	24
3 - 4%	0.9	21
4 - 5%	0.9	21
5 - 6%	1.2	21
6 - 7%	1.2	21

Note: i) The above intervals are maximum.

(ii) Considering the slope class and erosion status (A1....) the intervals have to be decided.

Bund length recording: Considering the contour plan and the existing grass belts/partitions, the bunds are aligned and lengths are measured.

Section of the Bund

Bund section is decided considering the soil texture class and gravelliness class (bg₀, loamy sand, <15% gravel). The recommended Sections for different soils are given below.


Recommended Bund Section

Top width (m)	Base width (m)	Height (m)	Side slope (Z:1;H:V)	Cross section (sq m)	Soil Texture	Remarks
0.3	0.9	0.3	01:01	0.18	Sandy loam	Vegetative bund
0.3	1.2	0.3	1.5:1	0.225	Sandy clay	
0.3	1.2	0.5	0.9:1	0.375	Red gravelly soil	
0.3	1.2	0.6	0.75:1	0.45		
0.3	1.5	0.6	01:01	0.54	Red sandy loam	
0.3	2.1	0.6	1.5:1	0.72	Very shallow black soils	
0.45	2	0.75	01:01	0.92		
0.45	2.4	0.75	1.3:1	1.07	Shallow black soils	
0.6	3.1	0.7	1.78:1	1.29	Medium black soils	
0.5	3	0.85	1.47:1	1.49		

Formation of Trench cum Bund

Dimensions of the Borrow Pits/ Trenches to be excavated (machinery are decided considering the Bund Section).

Details of Borrow Pit dimensions are given below




TRENCH CUM BUND


WATER STORAGE AREA

0.45 Sq.m section

IDEAL FOR HORTICULTURE CROPS



'A' FRAME FOR INTERBUND MANAGEMENT



1. ಸಮಸಾತಳ ಉಳುವಂ

2. ಸಮಸಾತಳ ಬಿತ್ತನೆ/ನಾಟ

Size of Borrow Pits/ Trench recommended for Trench cum Bund (by machinery)

Bund section	Bund length	Earth quantity	Pit				Berm (pit to pit)	Soil depth class
			L(m)	W(m)	D(m)	QUANTITY (m ³)		
m ²	m	m ³	L(m)	W(m)	D(m)	QUANTITY (m ³)	m	
0.375	6	2.25	5.85	0.85	0.45	2.24	0.15	Shallow
0.45	6	2.7	5.4	1.2	0.43	2.79	0.6	Shallow
0.45	6	2.7	5	0.85	0.65	2.76	1	Moderately Shallow
0.54	5.6	3.02	5.5	0.85	0.7	3.27	0.1	Moderately shallow
0.54	5.5	2.97	5	1.2	0.5	3	0.5	Shallow
0.72	6.2	4.46	6	1.2	0.7	5.04	0.2	Moderately shallow
0.72	5.2	3.74	5.1	0.85	0.9	3.9	0.1	Moderately deep

B. Water Ways

- Existing water ways are marked on the cadastral map (1:7920 scale) and their dimensions are recorded.
- Considering the contour plan of the MWS, additional water ways/ modernization of the existing ones can be thought of.
- The design details are given in the Manual.

C. Farm Ponds

Water ways and the catchment area will give an indication on the size of the Farm Pond. Location of the pond can be decided based on the contour plan/ field condition and farmers' need/desire.

D. Diversion Channel

Existing EPT/ CPT are marked on the cadastral map. Looking to the need, these can be modernized or fresh diversion channel can be proposed and runoff from this can be stored in Gokatte/ Recharge ponds.

9.1.2 Non-Arable Land Treatment

Depending on the gravelliness and crops preferred by the farmers, the concerned authorities can decide appropriate treatment plan. The recommended treatments may be

Contour Trench, Staggered Trench, Crescent Bund, Boulder Bund or Pebble Bunds are formed in the field.

9.1.3 Treatment of Natural Water Course/ Drainage Lines

- a) The cadastral map has to be updated as regards the network of drainage lines (gullies/ nalas/ hallas) and existing structures are marked to the scale and storage capacity of the existing water bodies are documented.
- b) The drainage line will be demarcated into Upper Reach, Middle Reach and Lower Reach.
- c) Considering the Catchment, Nala bed and bank conditions, suitable structures are decided.
- d) Number of storage structures (Check dam/ Nala bund/ Percolation tank) will be decided considering the commitments and available runoff in water budgeting and quality of water in the wells and site suitability.
- e) Detailed Levelling Survey using Dumpy Level / Total Station has to be carried out to arrive at the site-specific designs as shown in the Manual.
- f) The location of ground water recharge structures are decided by examining the lineaments and fracture zones from geological maps.
- g) Rainfall intensity data of the nearest Rain gauge station is considered for Hydrologic Designs.
- h) Silt load to the Storage/Recharge structures is reduced by providing vegetative, boulder and earthen checks in the natural water course. Location and design details are given in the Manual.

9.2 Recommended Soil and Water Conservation Measures

The appropriate conservation structures best suited for each of the land parcel/ survey number (Appendix-I) are selected based on the slope per cent, severity of erosion, amount of rainfall, land use and soil type. The different kinds of conservation structures recommended are

1. Graded / Strengthening of bunds
2. Trench cum Bunds (TCB)
3. Trench cum Bunds / Strengthening
4. Crescent Bunds

A map (Fig. 9.1) showing soil and water conservation plan with different kinds of structures recommended has been prepared which shows the spatial distribution and extent of area. An area of about 422 ha (79%) requires trench cum bunding and about 100 ha (16%) area needs graded bunds.

The conservation plan prepared may be presented to all the stakeholders including farmers and after including their suggestions, the conservation plan for the microwatershed may be finalised in a participatory approach.




Fig. 9.1 Soil and Water Conservation Plan map of Chincholi Khurd-1 microwatershed

9.3 Greening of Microwatershed

As part of the greening programme in the watersheds, it is envisaged to plant a variety of horticultural and other tree plants that are edible, economical and produce lot of biomass which helps to restore the ecological balance in the watersheds. The lands that are suitable for greening programme are non-arable lands (land capability classes V, VI and VII) and also the lands that are not suitable or marginally suitable for growing annual and perennial crops. The method of planting these trees is given below.

It is recommended to open pits during the 1st week of March along the contour and heap the dug out soil on the lower side of the slope in order to harness the flowing water and facilitate weathering of soil in the pit. Exposure of soil in the pit also prevents spread of pests and diseases due to scorching sun rays. The pits should be filled with mixture of soil and organic manure during the second week of April and keep ready with sufficiently tall seedlings produced either in poly bags or in root trainer nurseries so that planting can be done during the 2nd or 3rd week of April depending on the rainfall.

The tree species suitable for the area considering rainfall, temperature and adaptability is listed below; waterlogged areas are recommended to be planted with species like Neral (*Sizygium cumini*) and Bamboo. Dry areas are to be planted with species like Honge, Bevu, Seetaphal etc.

Dry Deciduous Species			Temp (°C)	Rainfall (mm)
1.	Bevu	<i>Azadiracta indica</i>	21–32	400 –1,200
2.	Tapasi	<i>Holoptelia integrifolia</i>	20-30	500 - 1000
3.	Seetaphal	<i>Anona Squamosa</i>	20-40	400 - 1000
4.	Honge	<i>Pongamia pinnata</i>	20 -50	500– 2,500
5.	Kamara	<i>Hardwickia binata</i>	25 -35	400 - 1000
6.	Bage	<i>Albezzia lebbek</i>	20 - 45	500 - 1000
7.	Ficus	<i>Ficus bengalensis</i>	20 - 50	500–2,500
8.	Sisso	<i>Dalbargia Sissoo</i>	20 - 50	500 -2000
9.	Ailanthus	<i>Ailanthus excelsa</i>	20 - 50	500 - 1000
10.	Hale	<i>Wrightia tinctoria</i>	25 - 45	500 - 1000
11.	Uded	<i>Steriospermum chelanoides</i>	25 - 45	500 -2000
12.	Dhupa	<i>Boswellia Serrata</i>	20 - 40	500 - 2000
13.	Nelli	<i>Emblica Officinalis</i>	20 - 50	500 -1500
14.	Honne	<i>Pterocarpus marsupium</i>	20 - 40	500 - 2000
Moist Deciduous Species			Temp (°C)	Rainfall (mm)
15.	Teak	<i>Tectona grandis</i>	20 - 50	500-5000
16.	Nandi	<i>Legarstroemia lanceolata</i>	20 - 40	500 - 4000
17.	Honne	<i>Pterocarpus marsupium</i>	20 - 40	500 - 3000
18.	Mathi	<i>Terminalia alata</i>	20 -50	500 - 2000
19.	Shivane	<i>Gmelina arboria</i>	20 -50	500 -2000
20.	Kindal	<i>T.Paniculata</i>	20 - 40	500 - 1500
21.	Beete	<i>Dalbargia latifolia</i>	20 - 40	500 - 1500
22.	Tare	<i>T. belerica</i>	20 - 40	500 - 2000
23.	Bamboo	<i>Bambusa arundinasia</i>	20 - 40	500 - 2500
24.	Bamboo	<i>Dendrocalamus strictus</i>	20 – 40	500 – 2500
25.	Muthuga	<i>Butea monosperma</i>	20 - 40	400 - 1500
26.	Hippe	<i>Madhuca latifolia</i>	20 - 40	500 - 2000
27.	Sandal	<i>Santalum album</i>	20 - 50	400 - 1000
28.	Nelli	<i>Emblica officinalis</i>	20 - 40	500 - 2000
29.	Nerale	<i>Sizyzium cumini</i>	20 - 40	500 - 2000
30.	Dhaman	<i>Grevia tilifolia</i>	20 - 40	500 - 2000
31.	Kaval	<i>Careya arborea</i>	20 - 40	500 - 2000
32.	Harada	<i>Terminalia chebula</i>	20 - 40	500 - 2000

References

1. FAO (1976) Framework for Land Evaluation, Food and Agriculture Organization, Rome.72 pp.
2. FAO (1983) Guidelines for Land Evaluation for Rainfed Agriculture, FAO, Rome, 237 pp.
3. IARI (1971) Soil Survey Manual, All India Soil and Land Use Survey Organization, IARI, New Delhi, 121 pp.
4. Katyal, J.C. and Rattan, R.K. (2003) Secondary and Micronutrients; Research Gap and Future Needs. Fert. News 48 (4); 9-20.
5. Naidu, L.G.K., Ramamurthy, V., Challa, O., Hegde, R. and Krishnan, P. (2006) Manual Soil Site Suitability Criteria for Major Crops, NBSS Publ. No. 129, NBSS &LUP, Nagpur, 118 pp.
6. Natarajan, A. and Dipak Sarkar (2010) Field Guide for Soil Survey, National Bureau of Soil Survey and Land Use Planning (ICAR), Nagpur, India.
7. Natarajan, A., Rajendra Hegde, Raj, J.N. and Shivananda Murthy, H.G. (2015) Implementation Manual for Sujala-III Project, Watershed Development Department, Bengaluru, Karnataka.
8. Sarma, V.A.K., Krishnan, P. and Budihal, S.L. (1987) Laboratory Manual, Tech. Bull. 23, NBSS &LUP, Nagpur.
9. Sehgal, J.L. (1990) Soil Resource Mapping of Different States of India; Why and How?, National Bureau of Soil Survey and Land Use Planning, Nagpur, 49 pp.
10. Shivaprasad, C.R., R.S. Reddy, J. Sehgal and M. Velayuthum (1998) Soils of Karnataka for Optimising Land Use, NBSS Publ. No. 47b, NBSS & LUP, Nagpur, India.
11. Soil Survey Staff (2006) Keys to Soil Taxonomy, Tenth edition, U.S. Department of Agriculture/ NRCS, Washington DC, U.S.A.
12. Soil Survey Staff (2012) Soil Survey Manual, Handbook No. 18, USDA, Washington DC, USA.

Appendix I

Soil Phase Information

VILLAGE	Survey No.	Total Area (ha)	Soils phase	Land Management Unit	Soil Depth	Surface Soil Texture	Soil Gravelliness	AWC	Slope	Soil Erosion	CLU code	WELLS	Land Capability	Conservation Plan
Chincholi (K)	3	6.45	MANmB2	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Borewell,Openwell	Iise	GB/strengt hening of Bunds
Chincholi (K)	4	1.52	MANmB2	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iise	GB/strengt hening of Bunds
Chincholi (K)	5	8.19	BHImC2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Moderate	Greengram (Gg)	Borewell,Borewell,Openwell,Openwell	IIIse	Crescent bund/TCB
Chincholi (K)	6	6.68	BHImC2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Moderate	Greengram+Redgram+Sugarcarne(Gg+Rg+Sc)	Not Available	IIIse	Crescent bund/TCB
Chincholi (K)	7	10.49	BHImC2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Moderate	Redgram+Sugar cane (Rg+Sc)	Checkdam	IIIse	Crescent bund/TCB
Chincholi (K)	8	9.87	BHImB2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIIse	Crescent bund/TCB
Chincholi (K)	9	10.15	BHImB2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram+Redgram+No Crop (Gg+Rg+NC)	Not Available	IIIse	Crescent bund/TCB
Chincholi (K)	10	7.81	BHImB1	LMU-3	Shallow (25-50 cm)	Clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Slight	Redgram+No Crop (Rg+NC)	Not Available	IIIse	Crescent bund/TCB
Chincholi (K)	11	13.2	BHImB1	LMU-3	Shallow (25-50 cm)	Clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Slight	Greengram+Redgram+Sunflower (Gg+Rg+Sf)	Not Available	IIIse	Crescent bund/TCB
Chincholi (K)	12	2.36	MGTiC3g1	LMU-1	Very shallow (<25 cm)	Sandy clay	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Greengram (Gg)	Not Available	IVse	Crescent bund/TCB
Chincholi (K)	13	14.07	MANmB2	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram+Redgram (Gg+Rg)	Openwell	Iise	GB/strengt hening of Bunds
Chincholi (K)	14	6.36	MANmB2	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram (Gg)	Borewell	Iise	GB/strengt hening of Bunds
Chincholi (K)	15	1.42	BHImB1	LMU-3	Shallow (25-50 cm)	Clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Slight	No Crop (NC)	Not Available	IIIse	Crescent bund/TCB
Chincholi (K)	16	10.68	BHImB1	LMU-3	Shallow (25-50 cm)	Clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Slight	Redgram+No Crop (Rg+NC)	Not Available	IIIse	Crescent bund/TCB
Chincholi (K)	17	9.88	BHImB1	LMU-3	Shallow (25-50 cm)	Clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Slight	Greengram+Redgram (Gg+Rg)	Not Available	IIIse	Crescent bund/TCB

VILLAGE	Survey No.	Total Area (ha)	Soils phase	Land Management Unit	Soil Depth	Surface Soil Texture	Soil Gravelliness	AWC	Slope	Soil Erosion	CLU code	WELLS	Land Capabi-lity	Conserv-ation Plan
Chincholi (K)	18	8.25	NHAmB2	LMU-3	Shallow (25-50 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram+Redgram (Gg+Rg)	Not Available	IIIse	Crescent bund/TCB
Chincholi (K)	19	8.53	MGTmB2	LMU-2	Very shallow (<25 cm)	Clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram+Redgram (Gg+Rg)	Openwell	IIIse	Crescent bund/TCB
Chincholi (K)	20	3.97	MGTmB2	LMU-2	Very shallow (<25 cm)	Clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIIse	Crescent bund/TCB
Chincholi (K)	21	5.8	MGTmC2g1	LMU-2	Very shallow (<25 cm)	Clay	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Moderate	Redgram (Rg)	Not Available	IIIse	Crescent bund/TCB
Chincholi (K)	22	4.03	MGTmC2g1	LMU-2	Very shallow (<25 cm)	Clay	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Moderate	Redgram (Rg)	Not Available	IIIse	Crescent bund/TCB
Chincholi (K)	25	0.31	MGTmB2	LMU-2	Very shallow (<25 cm)	Clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIIse	Crescent bund/TCB
Chincholi (K)	26	3.38	MGTmB2	LMU-2	Very shallow (<25 cm)	Clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIIse	Crescent bund/TCB
Chincholi (K)	27	6.9	MGTmB2	LMU-2	Very shallow (<25 cm)	Clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIIse	Crescent bund/TCB
Chincholi (K)	28	6.93	NHAmB2	LMU-3	Shallow (25-50 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram+Redgram (Gg+Rg)	Openwell ,Borewell	IIIse	Crescent bund/TCB
Chincholi (K)	29	7.67	NHAmB1	LMU-3	Shallow (25-50 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Greengram+No Crop (Gg+NC)	Borewell, Borewell	IIIs	Crescent bund/TCB
Chincholi (K)	STREAM	2.71	MANmB2	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Waterbody	Not Available	IIse	GB/strengthening of Bunds
Padasavali	4	0.05	MANmA1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Greengram (Gg)	Not Available	IIs	GB/strengthening of Bunds
Padasavali	7/1	0.32	Waterbody	Others	Others	Others	Others	Others	Others	Others	Greengram (Gg)	Not Available	Others	Others
Padasavali	211	6.26	MGTmC2g1	LMU-2	Very shallow (<25 cm)	Clay	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Moderate	Greengram+Redgram (Gg+Rg)	Not Available	IIIse	Crescent bund/TCB
Padasavali	219	2.83	MGTmB2	LMU-2	Very shallow (<25 cm)	Clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIIse	Crescent bund/TCB
Padasavali	220	7.94	MGTmB3g1	LMU-1	Very shallow (<25 cm)	Clay	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Greengram+Redgram (Gg+Rg)	Not Available	IVse	Crescent bund/TCB

VILLAGE	Survey No.	Total Area (ha)	Soils phase	Land Management Unit	Soil Depth	Surface Soil Texture	Soil Gravelliness	AWC	Slope	Soil Erosion	CLU code	WELLS	Land Capability	Conservation Plan
Padasavali	221	8.22	MGTmB2	LMU-2	Very shallow (<25 cm)	Clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram+Redgram (Gg+Rg)	Not Available	IIIse	Crescent bund/TCB
Padasavali	222	15.09	MGTmB2	LMU-2	Very shallow (<25 cm)	Clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram+Redgram+Sunflower (Gg+Rg+Sf)	Not Available	IIIse	Crescent bund/TCB
Padasavali	223	14.98	MGTmB2	LMU-2	Very shallow (<25 cm)	Clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram+Redgram+No Crop (Gg+Rg+NC)	Not Available	IIIse	Crescent bund/TCB
Padasavali	224	13.38	NHAmB2	LMU-3	Shallow (25-50 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram+Redgram (Gg+Rg)	Not Available	IIIse	Crescent bund/TCB
Padasavali	225	13.4	NHAmB2	LMU-3	Shallow (25-50 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Sugar cane (Rg+Sc)	Not Available	IIIse	Crescent bund/TCB
Padasavali	226	8.6	RNLmB1	LMU-5	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Greengram+Redgram (Gg+Rg)	Checkdam,Borewell,Borewell	IIs	GB/strengthening of Bunds
Padasavali	227	7.96	MANmB1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Redgram+Sugar cane (Rg+Sc)	Borewell	IIs	GB/strengthening of Bunds
Padasavali	228	15.15	MANmB1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Greengram+Redgram+Sugarcane (Gg+Rg+Sc)	Openwell	IIs	GB/strengthening of Bunds
Padasavali	229/1	1.69	MGTmB2	LMU-2	Very shallow (<25 cm)	Clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Sugarcane (Sc)	Openwell,Borewell	IIIse	Crescent bund/TCB
Padasavali	229/2	1.83	MGTmB2	LMU-2	Very shallow (<25 cm)	Clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIIse	Crescent bund/TCB
Padasavali	230/1	2.97	MANmB1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Redgram (Rg)	Borewell,Borewell,Openwell	IIs	GB/strengthening of Bunds
Padasavali	230/2	2.44	MANmB1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Sugarcane (Sc)	Not Available	IIs	GB/strengthening of Bunds
Padasavali	231	3.11	MANmB1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Sugarcane (Sc)	Not Available	IIs	GB/strengthening of Bunds
Padasavali	232	6.26	MANmB2	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram+Redgram (Gg+Rg)	Not Available	IIs	GB/strengthening of Bunds
Padasavali	233	2.36	MANmB2	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIs	GB/strengthening of Bunds
Padasavali	234	6.69	MANmB2	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Sugar cane (Rg+Sc)	Borewell,Openwell,Openwell	IIs	GB/strengthening of Bunds

VILLAGE	Survey No.	Total Area (ha)	Soils phase	Land Management Unit	Soil Depth	Surface Soil Texture	Soil Gravelliness	AWC	Slope	Soil Erosion	CLU code	WELLS	Land Capability	Conservation Plan
Padasavali	235	6.48	MANmB2	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+No Crop (Rg+NC)	Borewell	IIsE	GB/strengthening of Bunds
Padasavali	236	6.22	MANmB2	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram+Redgram (Gg+Rg)	Not Available	IIsE	GB/strengthening of Bunds
Padasavali	237	5.71	MANmB2	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram+Redgram (Gg+Rg)	Not Available	IIsE	GB/strengthening of Bunds
Padasavali	238	5.04	MANmB2	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram+No Crop (Gg+NC)	Openwell	IIsE	GB/strengthening of Bunds
Padasavali	239	7.34	MANmB2	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram+Redgram (Gg+Rg)	Openwell,Borewell	IIsE	GB/strengthening of Bunds
Padasavali	240	7	RNLmC2g1	LMU-5	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Gently sloping (3-5%)	Moderate	Redgram (Rg)	Not Available	IIsE	GB/strengthening of Bunds
Padasavali	241	10.65	RNLmC2g1	LMU-5	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Gently sloping (3-5%)	Moderate	Greengram+Redgram (Gg+Rg)	Checkdam	IIsE	GB/strengthening of Bunds
Padasavali	242/1	1.23	MGTmC2g1	LMU-2	Very shallow (<25 cm)	Clay	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Moderate	Redgram (Rg)	Not Available	IIIsE	Crescent bund/TCB
Padasavali	242/2	2.16	MGTmC2g1	LMU-2	Very shallow (<25 cm)	Clay	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Moderate	Redgram (Rg)	Not Available	IIIsE	Crescent bund/TCB
Padasavali	243	2.94	MGTmC2g1	LMU-2	Very shallow (<25 cm)	Clay	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Moderate	No Crop (NC)	Not Available	IIIsE	Crescent bund/TCB
Padasavali	244/1	2.56	BHImB2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	No Crop (NC)	Not Available	IIIsE	Crescent bund/TCB
Padasavali	244/2	2.98	BHImB2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	No Crop (NC)	Not Available	IIIsE	Crescent bund/TCB
Padasavali	245	7.78	BHImB2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	No Crop (NC)	Not Available	IIIsE	Crescent bund/TCB
Padasavali	246/1	4.5	BHImB1	LMU-3	Shallow (25-50 cm)	Clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Slight	No Crop (NC)	Not Available	IIIsE	Crescent bund/TCB
Padasavali	246/2	4.28	BHImB1	LMU-3	Shallow (25-50 cm)	Clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Slight	No Crop (NC)	Not Available	IIIsE	Crescent bund/TCB
Padasavali	267	2.78	MGTmB2	LMU-2	Very shallow (<25 cm)	Clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Sugarcane (Sc)	Not Available	IIIs	Crescent bund/TCB

VILLAGE	Survey No.	Total Area (ha)	Soils phase	Land Management Unit	Soil Depth	Surface Soil Texture	Soil Gravelliness	AWC	Slope	Soil Erosion	CLU code	WELLS	Land Capability	Conservation Plan
Padasavali	268	8.32	MGTiC3g1	LMU-1	Very shallow (<25 cm)	Sandy clay	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Redgram+Sugarcane (Rg+Sc)	Borewell	IVse	Crescent bund/TCB
Padasavali	269	11.9	MGTiC3g1	LMU-1	Very shallow (<25 cm)	Sandy clay	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Greengram+Redgram (Gg+Rg)	Not Available	IVse	Crescent bund/TCB
Padasavali	270	8.81	NHAmB2	LMU-3	Shallow (25-50 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIIse	Crescent bund/TCB
Padasavali	271	6.47	MGTiC3g1	LMU-1	Very shallow (<25 cm)	Sandy clay	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Sugarcane+No Crop (Sc+NC)	Checkdam	IVse	Crescent bund/TCB
Padasavali	272	4.85	BHImC2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Moderate	Redgram+Sugarcane (Rg+Sc)	Borewell	IIIse	Crescent bund/TCB
Padasavali	273	4.06	NHAmB2	LMU-3	Shallow (25-50 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIIse	Crescent bund/TCB
Padasavali	274	8.96	MGTiC3g1	LMU-1	Very shallow (<25 cm)	Sandy clay	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Greengram+Redgram (Gg+Rg)	Openwell	IVse	Crescent bund/TCB
Padasavali	STREAM	0.71	NHAmB2	LMU-3	Shallow (25-50 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Waterbody	Not Available	IIIse	Crescent bund/TCB
Sarasamba	25	0.31	NHAmB2	LMU-3	Shallow (25-50 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIIse	Crescent bund/TCB
Sarasamba	26	1.06	NHAmB2	LMU-3	Shallow (25-50 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIIse	Crescent bund/TCB
Sarasamba	28	12.4	NHAmB2	LMU-3	Shallow (25-50 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram+Redgram (Gg+Rg)	Not Available	IIIse	Crescent bund/TCB
Sarasamba	29	11.27	DSImB2	LMU-4	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	No Crop (NC)	Not Available	IIs	TCB
Sarasamba	30	7.98	MGTmC2	LMU-2	Very shallow (<25 cm)	Clay	Non gravelly (<15%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Moderate	Greengram+Redgram+No Crop (Gg+Rg+NC)	Not Available	IIIse	Crescent bund/TCB
Sarasamba	31	5.77	MGTmB1	LMU-2	Very shallow (<25 cm)	Clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Slight	Greengram+Redgram+Sugarcane (Gg+Rg+Sc)	Not Available	IIIs	Crescent bund/TCB
Sarasamba	32	10.51	MGTmA1	LMU-2	Very shallow (<25 cm)	Clay	Non gravelly (<15%)	Very low (<50 mm/m)	Nearly level (0-1%)	Slight	Redgram+Sunflower (Rg+Sf)	Openwell	IIIs	Crescent bund/TCB
Sarasamba	33	8.21	MGTmB2	LMU-2	Very shallow (<25 cm)	Clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram+Redgram (Gg+Rg)	Openwell	IIIse	Crescent bund/TCB
Sarasamba	34	9.85	MGTmB2g1	LMU-2	Very shallow (<25 cm)	Clay	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+No Crop (Rg+NC)	Not Available	IIIse	Crescent bund/TCB

VILLAGE	Survey No.	Soil Reaction	EC	Organic Carbon	Available Phosphorus	Available Potassium	Available Sulphur	Available Boron	Available Iron	Available Manganese	Available Copper	Available Zinc
Padasavali	274	Moderately alkaline (pH 7.8 – 8.4)	Non saline (<2 dsm)	Low (< 0.5 %)	Medium (23 - 57 kg/ha)	High (> 337 kg/ha)	High (> 10 ppm)	Medium (0.5 - 1.0 ppm)	Sufficient (> 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)
Padasavali	STRE AM	Strongly alkaline (pH 8.4 – 9.0)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Low (< 23 kg/ha)	High (> 337 kg/ha)	High (> 10 ppm)	Medium (0.5 - 1.0 ppm)	Sufficient (> 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Sarasamba	25	Strongly alkaline (pH 8.4 – 9.0)	Non saline (<2 dsm)	High (>0.75 %)	Low (< 23 kg/ha)	High (> 337 kg/ha)	Medium (10-20 ppm)	Low (< 0.5 ppm)	Sufficient (> 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Sarasamba	26	Strongly alkaline (pH 8.4 – 9.0)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Low (< 23 kg/ha)	High (> 337 kg/ha)	Medium (10-20 ppm)	Low (< 0.5 ppm)	Dwficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Sarasamba	28	Strongly alkaline (pH 8.4 – 9.0)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Low (< 23 kg/ha)	High (> 337 kg/ha)	Medium (10-20 ppm)	Medium (0.5 - 1.0 ppm)	Sufficient (> 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Sarasamba	29	Moderately alkaline (pH 7.8 – 8.4)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Low (< 23 kg/ha)	High (> 337 kg/ha)	Medium (10-20 ppm)	Low (< 0.5 ppm)	Sufficient (> 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Sarasamba	30	Moderately alkaline (pH 7.8 – 8.4)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Low (< 23 kg/ha)	Medium (145 - 337 kg/ha)	Medium (10-20 ppm)	Low (< 0.5 ppm)	Sufficient (> 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Sarasamba	31	Moderately alkaline (pH 7.8 – 8.4)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Low (< 23 kg/ha)	High (> 337 kg/ha)	Low (< 10 ppm)	Low (< 0.5 ppm)	Sufficient (> 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Sarasamba	32	Moderately alkaline (pH 7.8 – 8.4)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Low (< 23 kg/ha)	High (> 337 kg/ha)	Medium (10-20 ppm)	Low (< 0.5 ppm)	Sufficient (> 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Sarasamba	33	Moderately alkaline (pH 7.8 – 8.4)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Low (< 23 kg/ha)	High (> 337 kg/ha)	Medium (10-20 ppm)	Low (< 0.5 ppm)	Sufficient (> 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Sarasamba	34	Moderately alkaline (pH 7.8 – 8.4)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Low (< 23 kg/ha)	High (> 337 kg/ha)	Low (< 10 ppm)	Low (< 0.5 ppm)	Sufficient (> 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)

Appendix III

Soil Suitability Information

VILLAGE	Survey No.	Sorghum	Maize	Red gram	Sunflower	Cotton	Sugarcane	Soyabean
Chincholi (K)	3	S2e	S3t	S2te	S2e	S2e	S3t	S2e
Chincholi (K)	4	S2e	S3t	S2te	S2e	S2e	S3t	S2e
Chincholi (K)	5	S3r	S3r	S3r	N	S3r	N	S3r
Chincholi (K)	6	S3r	S3r	S3r	N	S3r	N	S3r
Chincholi (K)	7	S3r	S3r	S3r	N	S3r	N	S3r
Chincholi (K)	8	S3r	S3r	S3r	N	S3r	N	S3r
Chincholi (K)	9	S3r	S3r	S3r	N	S3r	N	S3r
Chincholi (K)	10	S3r	S3r	S3r	N	S3r	N	S3r
Chincholi (K)	11	S3r	S3r	S3r	N	S3r	N	S3r
Chincholi (K)	12	N	N	N	N	N	N	N
Chincholi (K)	13	S2e	S3t	S2te	S2e	S2e	S3t	S2e
Chincholi (K)	14	S2e	S3t	S2te	S2e	S2e	S3t	S2e
Chincholi (K)	15	S3r	S3r	S3r	N	S3r	N	S3r
Chincholi (K)	16	S3r	S3r	S3r	N	S3r	N	S3r
Chincholi (K)	17	S3r	S3r	S3r	N	S3r	N	S3r
Chincholi (K)	18	S3r	S3rt	S3r	N	S3r	N	S3r
Chincholi (K)	19	N	N	N	N	N	N	N
Chincholi (K)	20	N	N	N	N	N	N	N
Chincholi (K)	21	N	N	N	N	N	N	N
Chincholi (K)	22	N	N	N	N	N	N	N
Chincholi (K)	25	N	N	N	N	N	N	N
Chincholi (K)	26	N	N	N	N	N	N	N
Chincholi (K)	27	N	N	N	N	N	N	N
Chincholi (K)	28	S3r	S3rt	S3r	N	S3r	N	S3r
Chincholi (K)	29	S3r	S3rt	S3r	N	S3r	N	S3r
Chincholi (K)	STREAM	S2e	S3t	S2te	S2e	S2e	S3t	S2e

VILLAGE	Survey No.	Sorghum	Maize	Red gram	Sunflower	Cotton	Sugarcane	Soyabean
Padasavali	4	S1	S3t	S2t	S1	S1	S3t	S1
Padasavali	7/1	Others	Others	Others	Others	Others	Others	Others
Padasavali	211	N	N	N	N	N	N	N
Padasavali	219	N	N	N	N	N	N	N
Padasavali	220	N	N	N	N	N	N	N
Padasavali	221	N	N	N	N	N	N	N
Padasavali	222	N	N	N	N	N	N	N
Padasavali	223	N	N	N	N	N	N	N
Padasavali	224	S3r	S3rt	S3r	N	S3r	N	S3r
Padasavali	225	S3r	S3rt	S3r	N	S3r	N	S3r
Padasavali	226	S1	S3t	S2t	S1	S1	S3t	S1
Padasavali	227	S1	S3t	S2t	S1	S1	S3t	S1
Padasavali	228	S1	S3t	S2t	S1	S1	S3t	S1
Padasavali	229/1	N	N	N	N	N	N	N
Padasavali	229/2	N	N	N	N	N	N	N
Padasavali	230/1	S1	S3t	S2t	S1	S1	S3t	S1
Padasavali	230/2	S1	S3t	S2t	S1	S1	S3t	S1
Padasavali	231	S1	S3t	S2t	S1	S1	S3t	S1
Padasavali	232	S2e	S3t	S2te	S2e	S2e	S3t	S2e
Padasavali	233	S2e	S3t	S2te	S2e	S2e	S3t	S2e
Padasavali	234	S2e	S3t	S2te	S2e	S2e	S3t	S2e
Padasavali	235	S2e	S3t	S2te	S2e	S2e	S3t	S2e
Padasavali	236	S2e	S3t	S2te	S2e	S2e	S3t	S2e
Padasavali	237	S2e	S3t	S2te	S2e	S2e	S3t	S2e
Padasavali	238	S2e	S3t	S2te	S2e	S2e	S3t	S2e
Padasavali	239	S2e	S3t	S2te	S2e	S2e	S3t	S2e
Padasavali	240	S2e	S3t	S2te	S2e	S2e	S3t	S2e
Padasavali	241	S2e	S3t	S2te	S2e	S2e	S3t	S2e
Padasavali	242/1	N	N	N	N	N	N	N

VILLAGE	Survey No.	Sorghum	Maize	Red gram	Sunflower	Cotton	Sugarcane	Soyabean
Padasavali	242/2	N	N	N	N	N	N	N
Padasavali	243	N	N	N	N	N	N	N
Padasavali	244/1	S3r	S3r	S3r	N	S3r	N	S3r
Padasavali	244/2	S3r	S3r	S3r	N	S3r	N	S3r
Padasavali	245	S3r	S3r	S3r	N	S3r	N	S3r
Padasavali	246/1	S3r	S3r	S3r	N	S3r	N	S3r
Padasavali	246/2	S3r	S3r	S3r	N	S3r	N	S3r
Padasavali	267	N	N	N	N	N	N	N
Padasavali	268	N	N	N	N	N	N	N
Padasavali	269	N	N	N	N	N	N	N
Padasavali	270	S3r	S3rt	S3r	N	S3r	N	S3r
Padasavali	271	N	N	N	N	N	N	N
Padasavali	272	S3r	S3r	S3r	N	S3r	N	S3r
Padasavali	273	S3r	S3rt	S3r	N	S3r	N	S3r
Padasavali	274	N	N	N	N	N	N	N
Padasavali	STREAM	S3r	S3rt	S3r	N	S3r	N	S3r
Sarasamba	25	S3r	S3rt	S3r	N	S3r	N	S3r
Sarasamba	26	S3r	S3rt	S3r	N	S3r	N	S3r
Sarasamba	28	S3r	S3rt	S3r	N	S3r	N	S3r
Sarasamba	29	S2r	S3t	S2r	S3r	S2r	S3t	S2r
Sarasamba	30	N	N	N	N	N	N	N
Sarasamba	31	N	N	N	N	N	N	N
Sarasamba	32	N	N	N	N	N	N	N
Sarasamba	33	N	N	N	N	N	N	N
Sarasamba	34	N	N	N	N	N	N	N

PART-B

SOCIO-ECONOMIC STATUS OF FARM HOUSEHOLDS

CONTENTS

1	Executive summary	1-3
2	Introduction	5
3	Methodology	7-11
4	Results and discussions	13-28

LIST OF TABLES

I. Social status		
1	Human population among sample households	13
2	Basic needs of sample households	15
3	Migration details among the sample households	15
II. Economic status		
4	Occupational pattern in sample households	16
5	Domestic assets among samples households	17
6	Farm assets among samples households	17
7	Livestock assets among sample households	18
8	Milk produced and Fodder availability of sample households	18
9	Women empowerment of sample households	18
10	Per capita daily consumption of food among the sample households	19
11	Annual average Income from various sources	20
12	Average annual expenditure of sample farmers	21
13	Distribution of land holding among the sample households	21
14	Land holding among samples households	22
III. Resource use pattern		
15	Number of tree/plants covered in sample farm households	22
16	Present cropping pattern among samples households	22
17	Distribution of soil series in the watershed	23-24
IV. Economic land evaluation		
18	Cropping pattern on major soil series	24
19	Alternative land use options for different size group of farmers (Benefit Cost Ratio)	24
20	Economics Land evaluation and bridging yield gap for different crops	25
21	Estimation of onsite cost of soil erosion	26
22	Ecosystem services of food grain production	27
23	Ecosystem services of fodder production	27
24	Ecosystem services of water supply for crop production	27
25	Farming constraints	28

LIST OF FIGURES

1	Location of study area	8
2	ALPES Framework	9
3	Basic needs of sample households	14
4	Domestic assets among the sample households	16
5	Livestock assets among sample households	17
6	Per capita daily consumption of food among the sample farmers	19
7	Average annual expenditure of sample households	20
8	Estimation of onsite cost of soil erosion	26

EXECUTIVE SUMMARY

Baseline socioeconomic characterisation is prerequisite to prepare action plan for program implementation and to assess the project performance before making any changes in the watershed development program. The baseline provides appropriate policy direction for enhancing productivity and sustainability in agriculture.

Methodology: *Chincholi Khurd-1 micro-watershed (Khairat sub-watershed, Aland taluk, Gulbarga district) is located in between 17°35'–17°37' North latitudes and 76°25'–76°27' East longitudes, covering an area of about 532.43 ha, bounded by Chincholi Budruk, Nirgudi, Khanapur and Nagalogaon villages with length of growing period (LGP) 120-150 days. We used soil resource map as basis for sampling farm households to test the hypothesis that soil quality influence crop selection and conservation investment of farm households. The level of technology adoption and productivity gaps and livelihood patterns were analyses. The cost of soil degradation and ecosystem services were quantified.*

Results: *The socio-economic outputs for Chincholi Khurd-1 micro-watershed (Khairat sub-watershed, Aland taluk, Gulbarga district) are presented here.*

Social Indicators;

- ❖ *Male and female ratio is 54.8 to 45.2 per cent to the total sample population.*
- ❖ *Younger age 18 to 50 years group of population is around 59.5 per cent to the total population.*
- ❖ *Literacy population is around 95.2 per cent.*
- ❖ *Social groups belong to other backward caste (OBC) is around 37.5 and general caste around 62.5 per cent.*
- ❖ *Fire wood is the source of energy for a cooking among 50 per cent.*
- ❖ *About 37.5 per cent of households have a yashaswini health card.*
- ❖ *Majority of farm households (50 %) are having MGNREGA card for rural employment.*
- ❖ *Dependence on ration cards for food grains through public distribution system is around 87.5 per cent.*
- ❖ *Swach bharath program providing closed toilet facilities around 50 per cent of sample households.*
- ❖ *Rural migration to urban centre for employment is prevalent among 2.4 per cent of farm households.*
- ❖ *Women participation in decisions making for agriculture production among the households was found.*

Economic Indicators;

- ❖ *The average land holding is 1.7 ha indicates that majority of farm households are belong to marginal and small farmers. The dry land is total cultivated land area among the sample farmers.*
- ❖ *Agriculture is the main occupation among 40.5 per cent and agriculture is the main and agriculture labour is subsidiary occupation among 42.9 per cent in sample households.*
- ❖ *The average value of domestic assets is around Rs. 57401 per household. Mobile and television are popular media mass communication.*
- ❖ *The average value of farm assets is around Rs. 35400 per household, about 25 per cent of sample farmers having weeder and tractor (12.5 %).*
- ❖ *The average value of livestock is around Rs. 46250 per household; about 75 per cent of household are having livestock.*
- ❖ *The average per capita food consumption is around 802.8 grams (1853.60 kilo calories) against national institute of nutrition (NIN) recommendation at 827 gram. Around 25 per cent of sample households are consuming less than the NIN recommendation.*
- ❖ *The annual average income is around Rs.47453 per household. About 12.5 per cent of farm households are below poverty line.*
- ❖ *The per capita average monthly expenditure is around Rs.2519.*

Environmental Indicators-Ecosystem Services;

- ❖ *The value of ecosystem service helps to support investment to decision on soil and water conservation and in promoting sustainable land use.*
- ❖ *The onsite cost of different soil nutrients lost due to soil erosion is around Rs. 1468 per ha/year. The total cost of annual soil nutrients is around Rs. 767543 per year for the total area of 532.43 ha.*
- ❖ *The average value of ecosystem service for food grain production is around Rs 11369/ ha/year. Per hectare food grain production services is maximum in red gram (Rs. 11537) and sorghum (Rs. 9833).*
- ❖ *The average value of ecosystem service for fodder production is around Rs 1707/ ha/year in sorghum.*
- ❖ *The data on water requirement for producing one quintal of grain is considered for estimating the total value of water required for crop production. The per hectare value of water used and value of water was maximum in red gram (Rs. 55810) and sorghum (Rs. 36725).*

Economic Land Evaluation;

- ❖ *The major cropping pattern is red gram (96.4 %) and sorghum (3.6 %).*
- ❖ *In Chincholi Khurd-1 micro-watershed, major soils are Margutti (MGT) series is having very shallow soil depth cover around 41.5 % of area. On this soil farmers*

are presently growing red gram (93.7 %) and sorghum (6.3 %). Novinahala (NHA) are also having shallow soil depth cover 16.9 % of area, the crop is a red gram. Bhimanahalli (BHI) soil series having shallow soil depth cover around 19.6 % of areas, crop is red gram. Mahagaon (MAN) soil series having very deep soil depth cover around 15.9 % of area crop is red gram.

- ❖ The total cost of cultivation and benefit cost ratio (BCR) in study area for red gram ranges between Rs. 43620/ha in MAN soil (with BCR of 1.00) and Rs.23317/ha in MGT soil (with BCR of 1.53).
- ❖ In sorghum the cost of cultivation in MGT soil is Rs 32338/ha (with BCR of 1.36).
- ❖ The land management practices reported by the farmers are crop rotation, tillage practices, fertilizer application and use of farm yard manure (FYM). Due to higher wages farmer are following labour saving strategies is not prating soil and water conservation measures. Less ownership of livestock limiting application of FYM.
- ❖ It was observed soil quality influences on the type and intensity of land use. More fertilizer applications in deeper soil to maximize returns.

Suggestions;

- ❖ Involving farmers in watershed planning helps in strengthening institutional participation.
- ❖ The per capita food consumption and monthly income is very low. Diversifying income generation activities from crop and livestock production in order to reduce risk related to drought and market prices.
- ❖ Majority of farmers reported that they are not getting timely support/extension services from the concerned development departments.
- ❖ By strengthening agricultural extension for providing timely advice improved technology there is scope to increase in net income of farm households. By adopting recommended package of practices by following the soil test fertiliser recommendation, there is scope to increase yield in red gram (3.6 to 28.5 %), sorghum (57.1 %).

INTRODUCTION

Watershed Development program aim to restore degraded watersheds in rainfed regions to increase their capacity to capture and store rain water, reduce soil erosion, and improved soil nutrients and carbon contents so they can produce greater agricultural yields and other benefits. As majority of rural poor live in these regions and dependent on natural resources for their livelihood and sustenance, improvements in agricultural yields improve human welfare and simultaneously improve national food security.

Sujala–III watershed development project conceptualised and implemented by the Watershed Development Department of Government of Karnataka with tripartite cost-sharing arrangements. The World Bank through International Development Association provided major portion of plan outlay as a loan to Government of India and in turn loan to Government of Karnataka.

The objectives of Sujala-III is to demonstrate more effective watershed management through greater integration of programs related to rain fed agriculture, innovative and science based approaches and strengthened institutions and capacities. The project is implemented in 11 districts of Bidar, Vijayapura, Gulbarga, Yadgir, Koppal, Gadag, Raichur, Davanagere, Tumkur, Chikkamangalur and Chamarajanagar which have been identified by the Watershed Development Department based on rainfall and socio-economic conditions. The project will be implemented over six years and linked with the centrally financed integrated watershed management programme.

Economic evaluations can better guide in watershed planning and implementation, as well as raise awareness of benefits of ecosystem restoration for food security and poverty alleviation program. The present study aims to characterize socio-economic status of farm households, assess the land and water use status, evaluate the economic viability of land use, prioritize farming constraints and suggest the measures for soil and water conservation for sustainable agriculture.

Objectives of the study

1. To characterize socio-economic status of farm households
2. To evaluate the economic viability of land use and land related constraints
3. To estimate the ecosystem service provided by the watershed and
4. To suggest alternatives for sustainable agriculture production.

METHODOLOGY

Study area

Chincholi Khurd-1 micro-watershed located in north-eastern dry zone of Karnataka (Figure 1): The total geographic area of this zone is about 1.76 M ha covering 8 taluks of Gulbarga district and 3 taluks of Raichur. Net cultivated area in the zone is about 1.31 M ha of which about 0.09 M ha are irrigated. The mean elevation of the zone is 300-450 m MSL. The main soil type is deep to very deep soils with small pockets of shallow to medium black soils. The zone is cropped predominantly during Rabi due to insufficient rainfall (465-785 mm). The principal crops of the zone are jowar, bajra, oilseeds, pulses, cotton and sugarcane. It's represented Agro Ecological Sub Region (AESR) 6.2 having LGP 120-150 days.

Chincholi Khurd-1 micro-watershed (Khairat sub-watershed, Aland taluk, Gulbarga district) is located in between 17⁰35'–17⁰37' North latitudes and 76⁰25'–76⁰27' East longitudes, covering an area of about 532.43 ha, bounded by Chincholi Budruk, Nirgudi, Khanapur and Nagalogaon villages.

Sampling Procedure:

In this study we have followed soil variability as criterion for sampling the farm households. In each micro-watershed the survey numbers and associated soil series are listed. Minimum three farm households for each soil series were taken and summed up to arrive at total sample for analysis.

Sources of data and analysis:

For evaluating the specific objectives of the study, primary data was collected from the sample respondents by personal interview method with the help of pre-tested questionnaire. The data on socio-economic characteristics of respondents such as family size and composition, land holdings, asset position, occupational pattern and education level was collected. The present cropping pattern and the level of input use and yields collected during survey. The data collected from the representative farm households were analysed using Automated Land Potential Evaluation System (Figure 2).

LOCATION MAP OF CHINCHOLI KHURD-1 MICRO-WATERSHED




Figure 1: Location of study area

Steps followed in socio-economic assessment

- 1 • After the completion of soil profile study link the cadastral number to the soil profile in the micro watershed.
- 2 • Download the names of the farmers who are owning the land for each cadastral number in the Karnataka BHOOMI Website.
- 3 • Compiling the names of the farmers representing for all the soil profiles studied in the micro watershed for socio-economic Survey.
- 4 • Conducting the socioeconomic survey of selected farm households in the micro watershed .
- 5 • Farm households database created using the Automated Land Potential Evaluation System (ALPES) for analysis of socio economic status for each micro watershed .
- 6 • Synthesis of tables and preparation of report for each micro watershed .




Figure 2: ALPES FRAMEWORK

The sample farmers were post classified in to marginal and small (0.0 to ≤ 2 ha), medium and semi medium (>2 to ≤ 10 ha) and large (>10 ha). The steps involved in estimation of soil potential involve estimation of total cost of cultivation, the yield/gross returns and net income per hectare. The cost of inputs such seed, manure and fertilizer, plant protection chemicals, payment towards human and bullock labour and interest on working capita are included under operational costs. In the case of perennial crops, the cost of establishment was estimated by using actual physical requirements and prevailing market prices. Estimation cost included maintenance cost up to bearing period. The value of main product and by product from the crop enterprise at the market rates were the gross returns of the crop. Net returns were worked out by deducting establishment and maintained cost from gross returns.

Operational Cost = cost of seeds, fertilizers, pesticides. Cost of human and bullock labour, cost of machinery, cost of irrigation water + interest on working capital.

Gross returns = Yield (Quintals/hectare)*Price (Rs/Quintal)

Net returns = Gross returns-Operational cost.


Benefit Cost Ratio = Net returns/Total cost.

Economic suitability classes: once each land use –land area combination has been assigned an economic value by the land evaluation, the question arises as to its ‘suitability’, that is, the degree to which it satisfies the land user. The FAO framework defines two suitability orders: ‘S’(suitable if benefit cost ratio (BCR) >1) and ‘N’(not suitable if (BCR <1), which are divided into five economic suitability classes: ‘S1’(highly suitable if BCR >3), ‘S2’(suitable if BCR >2 and <3), ‘S3’(Marginally suitable if BCR >1 and <2), ‘N1’(Not suitable for economic reasons but physically suitable) and ‘N2’(not suitable for physical reasons). The limit between ‘S3’ and ‘N1’ must be at least at the point of financial feasibility (i.e. net returns, NPV, or IRR >0 and BCR >1). The other limits depend on social factors such as farm size, family size, alternative employment or investment possibilities and wealth expectations; these need to be specified for the soil series.

Economic Valuation of Soil ecosystem services:

The replacement cost approach was followed for estimating the onsite cost of soil erosion, Market price method was followed for estimating the value of food and fodder production. Value transfer methods was followed for estimating the value of water demand by different crops in the micro watershed.

Steps followed in Replacement cost methods for estimation of onsite cost of soil erosion



RESULTS AND DISCUSSIONS

The demographic information shows that the household population dynamics encompasses the socioeconomic status of the farmer. For a rural family, the household size should be optimal to earn a comfortable livelihood through farm and non-farm wage earning. The total number of population in watershed area was 42, out of which 54.8 per cent were males and 45.2 per cent females. Average family size of the households is 5.3. Age is an important factor, which affects the potential employment and mobility status of respondents. The data on age wise distribution of farmers in the sample households indicated that majority of the farmers are coming under the age group of 18 to 30 years (33.3 %) followed by , 0 to 18 years (28.6 %), 30 to 50 years (26.2 %) and more than 50 years (11.9 %). Hence, in the study area in general, the respondents were of young and middle age, indicating thereby that the households had almost settled with whatever livelihood options they were practicing and sample respondents were young by age who could venture into various options of livelihood sources. Data on literacy indicated that 4.8 per cent of respondents were illiterate and 95.2 per cent literate (Table 1).

Table 1: Human population among sample households in Chincholi Khurd 1Microwatershed

Particulars	Units	Value
Total human population in sample HHs	Number	42
Male	% to total Population	54.8
Female	% to total Population	45.2
Average family size	Number	5.3
Age group		
0 to 18 years	% to total Population	28.6
18 to 30 years	% to total Population	33.3
30 to 50 years	% to total Population	26.2
>50 years	% to total Population	11.9
Average age	Age in years	29.4
Education Status		
Illiterates	% to total Population	4.8
Literates	% to total Population	95.2
Primary School (<5 class)	% to total Population	28.6
Middle School (6- 8 class)	% to total Population	11.9
High School (9- 10 class)	% to total Population	19.0
Others	% to total Population	35.7

The ethnic groups among the sample farm households found to be 62.5 per cent belonging to general caste and 37.5 per cent belonging to other backward caste (Table 2 and Figure 3). About 50.0per cent of sample households are using fire wood as source of fuel for cooking. All the sample farmers are having electricity connection. About 37.5 per cent are sample households having health cards. Majority (50 %) are having MNREGA job cards for employment generation. About 87.5 per cent of farm households are having

ration cards for taking food grains from public distribution system. About 50 per cent of farm households are having toilet facilities.




Figure 3: Basic needs of sample households in Chincholi Khurd -1 Micro-watershed

The data collected on the source of drinking water in the study area is presented in Table 2. All the sample respondents are having tube well source for water supply for domestic purpose.

Table 2: Basic needs of sample households in Chincholi Khurd -1 Micro-watershed

Particulars	Units	Value
Social groups		
OBC	% of Households	37.5
General	% of Households	62.5
Types of fuel use for cooking		
Fire wood	% of Households	50.0
Fire wood & Gas	% of Households	25.0
Gas	% of Households	25.0
Energy supply for home		
Electricity	% of Households	100.0
Number of households having Health card		
Yes	% of Households	37.5
No	% of Households	62.5
MGNREGA Card		
Yes	% of Households	50.0
No	% of Households	50.0
Ration Card		
Yes	% of Households	87.5
No	% of Households	12.5
Households with toilet		
Yes	% of Households	50.0
No	% of Households	50.0
Drinking water facilities		
Tube Well	% of Households	100.00

The data on migration in Chincholi Khurd -1 Micro-watershed is given in Table 3. It indicated that around 2.4 per cent of samples households were migrated. The average distance travelled for seeking employment is 100 km.

Table 3: Migration details among the sample households in Chincholi Khurd-1 micro-watershed

Particulars	Value
% of households showing migration	2.4
% of persons migrating	12.5
No. of months migrated in a year	6.0
Average Distance of migration(Km)	100.0
Nature of job (%)	
Job/wage/work	100.0

The occupational pattern (Table 4) among sample households shows that agriculture is the main occupation around 40.5 per cent of farmers followed by subsidiary occupations like agriculture labour(42.9%), private service (4.8%), trade and business

(2.4%), self employed (2.4 %). About 2.4 percent of sample household's main occupation is government services and 4.6 percent of private service.

Table 4: Occupational pattern in sample population in Chincholi Khurd -1 Micro-watershed

Occupation		% to total
Main	Subsidiary	
Agriculture	Agriculture	40.5
	Agriculture labour	42.9
	Private service	4.8
	Trade and business	2.4
	Self employed	2.4
Govt. service		2.4
Private service		4.6
Grand Total		100
Family labour availability		Man days/month
Male		25.0
Female		22.9
Total		47.9




Figure 4: Domestic assets among the sample households in Chincholi Khurd -1 Micro-watershed

The important assets especially with reference to domestic assets were analyzed and are given in Table 5 and Figure 4. The important domestic assets possessed by all categories of farmers are television (100 %) followed by mobile phones (100 %), motorcycle (62.5 %), mixer/grinder (37.5 %), refrigerator (25 %), four wheeler (12.5%)

and landline Phone (12.5 %), radio (12.5 %) and computer/laptop (12.5 %). The average value of domestic assets is around Rs.57401 per households.

Table 5: Domestic assets among the sample households in Chincholi Khurd -1 Micro-watershed

Particulars	% of households	Average value in Rs
Computer/laptop	12.5	30000
Four wheeler	12.5	400000
Landline Phone	12.5	1200
Mixer/grinder	37.5	2000
Mobile Phone	100.0	6963
Motorcycle	62.5	41200
Radio	12.5	7000
Refrigerator	25.0	20000
Television	100.0	8250
Average Value	57401	

The most popularly owned farm equipments were tractor and weeders were only present in the sampled farmers; these were primary implements in agriculture. The per cent of households owned tractor (12.5 %) and weeder (25 %). The average value of farm assets is around Rs 35400 per households (Table 6).

Table 6: Farm assets among samples households in Chincholi Khurd -1 Micro-watershed

Particulars	% of households	Average value in Rs
Tractor	12.5	70000
Weeder	25.0	800
Average Value	35400	




Figure 5: Livestock assets among sample households in Chincholi Khurd -1 micro-watershed

Livestock is an integral component of the conventional farming systems (Table 7 and Figure 5). The highest livestock population is local dry cow was around 33.3 per cent and local milching cow (33.3%), milching buffalos is (16.7 %) and bullocks is (16.7 %). The average livestock value was Rs.46250 per household.

Table 7: Livestock assets among sample households in Chincholi Khurd 1micro-watershed

Particulars	% of livestock population	Average value in Rs
Local Dry Cow	33.3	17500
Local Milching Cow	33.3	17500
Milching Buffalos	16.7	50000
Bullocks	16.7	100000
Average value	46250	

Average milk produced in sample households is 1156 liters/ annum (Table 8). The number livestock having sample farm households was 75 per cent.

Table 8: Milk produced and fodder availability of sample households in Chincholi Khurd -1 Micro-watershed

Particulars	
Name of the Livestock	Ltr./Lactation/animal
Local Milching Cow	1440
Milching Buffalos	800
Average Milk produced	1156
Livestock having households (%)	75
Livestock population (Numbers)	8

A woman participation in decision making in this micro-watershed is presented in Table 9. Among the all farm women taking decision in her family and agriculture related activities and women earning for her family requirement. Around 25.0 per cent of women participation in local organisation activates and about 12.5 per cent of women elected as panchayat member.

Table 9: Women empowerment of sample households in Chincholi Khurd -1 Micro-watershed

Particulars	% to Grand Total	
	Yes	No
Women participation in local organization activities	25.0	75.0
Women elected as panchayat member	12.5	87.5
Women earning for her family requirement	100.0	00.0
Women taking decision in her family and agriculture related activities	100.0	00.0

The food intake in terms of kilo calorie (kcal) per person per day was calculated and presented in the Table 10 and Figure 6. More quantity of cereals is consumed by sample farmers which accounted for 1255.03.kcal per person. The other important food items consumed was pulses 144.48 kcal followed by cooking oil 182.14 kcal, milk 93.42 kcal, vegetables 27.68 kcal, egg 129.02 kcal and meat 21.83 kcal. In the sampled households, farmers were consuming less (1853.6 kcal) than NIN- recommended food requirement (2250 kcal).

Table 10: Per capita daily consumption of food among the sample households in Chincholi Khurd -1 Micro-watershed

Particulars	NIN recommendation (gram/ per day/ person)	Present level of consumption (gram/ per day/ person)	Kilo Calories /day/person
Cereals	396	369.1	1255.03
Pulses	43	42.1	144.48
Milk	200	143.7	93.42
Vegetables	143	115.3	27.68
Cooking Oil	31	32.0	182.14
Egg	0.5	86.0	129.02
Meat	14.2	14.6	21.83
Total	827.7	802.8	1853.60
Threshold of NIN recommendation		827 gram*	2250 Kcal*
% Below NIN		25.0	00.0
% Above NIN		75.0	100

Note: * day/person




Figure 6: Per capita daily consumption of food among the sample households in Chincholi Khurd -1 Micro-watershed

Annual income of the sample HHs: The average annual household income is around Rs 47453.4. Major source of income to the farmers in the study area is from Livestock income (Rs 24120) followed by crop production (Rs. 21917). The income from Non farm income was very low at Rs 1416. The monthly per capita income is Rs. 1583, which is more than the threshold monthly income of Rs 753.2 for considering above poverty line. Due to the fact that erratic rainfall and shortage of water, farmers are

diverting from crop production activities to enable the household for a comfortable livelihood. The incomes from the other aforesaid sources are very meagre (Table 11).

Table 11: Annual average income of HHs from various sources in Chincholi Khurd -1 Micro-watershed

Particulars	Income *
Nonfarm income (Rs)	1416 (12.5)
Livestock income (Rs)	24120 (37.5)
Crop Production (Rs)	21917 (100)
Total Annual Income (Rs)	47453
Average monthly per capita income (Rs)	753
Threshold for Poverty level (Rs 975 per month/person)	
% of households below poverty line	87.5
% of households above poverty line	12.5

* Figure in the parenthesis indicates % of Households




Figure 7: Average annual expenditure of sample HHs in Chincholi Khurd -1 Micro-watershed

The average annual expenditure of farm households indicated that farmers in the study area spend highest on food (Rs. 158675) followed by education, clothing, social function and health. Now a day's education is most important among all of us. In today's

competitive world, education is a necessity for man after food, clothing, and shelter. It is the only fundamental way by which a desired change in the society can happen. The average per capita monthly expenditure is around Rs 2519 and about 12.5 per cent of farm households are below poverty line and 87.5 per cent of farm households are above poverty line (Table 12 and Figure 7).

Table 12: Average annual expenditure of sample HHs in Chincholi Khurd -1 Micro-watershed

Particulars	Value in Rupees	Per cent
Food	47487	29.9
Education	23938	15.1
Clothing	9875	6.2
Social functions	56250	35.4
Health	21125	13.3
Total Expenditure (Rs/year)	158675	100.0
Monthly per capita expenditure (Rs)	2519	

Land holding: Total sample households are total area cultivated by them is 14.2 ha. The average land holding of sample HHs is 1.7 ha. Large number of sample HHs (62.5 %) belong to small size group with an average holding size of 0.9 ha followed by medium farmers (25 %) with an average holding size of 2.3 ha and a large farmer (12.5 %) with an average land holding size of 4.8 ha (Table 13).

Table 13: Distribution of land holding among the sample households in Chincholi Khurd I Micro-watershed

Particulars	Units	Values
Small farmers		
Total land	ha	4.7
Sample size	Per cent	62.5
Average land holding	ha	0.9
Medium farmers		
Total land	ha	4.6
Sample size	Per cent	25.0
Average land holding	ha	2.3
Large farmers		
Total land	ha	4.8
Sample size	Per cent	12.5
Average land holding	ha	4.8
Total sample households		
Total land	ha	14.2
Sample size	Per cent	100.0
Average land holding	ha	1.7

Land use: The total land holding in the Chincholi Khurd -1 Microwatershed is 28.6 ha is rain fed land (Table 14). The average land holding per household is worked out to be 1.78 ha.

Table 14: Land use among samples households in Chincholi Khurd -1 Micro-watershed

Particulars	Per cent	Area in ha
Irrigated land	0.0	0.0
Rain fed Land	100.0	14.3
Fallow Land	0.0	0.0
Total land holding	100.0	14.3
Average land holding	1.78	

In the micro-watershed, the prevalent present land uses under perennial plants are neem trees (89.7 %) and mango tree (10.3 %) (Table 15).

Table 15: Number of trees/plants covered in sample farm households in Chincholi Khurd -1 Micro-watershed

Particulars	Number of Plants/trees	Per cent
Mango	4	10.3
Neem trees	35	89.7
Grand Total	39	100.0

The land use decisions are usually based on experience of farmers, tradition, expected profit, personal preferences, resources and social requirements (Table 16). The present dominant crops grown in dry lands in the study area were by red gram (96.4 %) and sorghum (3.6 %) during Kharif season respectively.

Table 16: Present cropping pattern and cropping intensity in Chincholi Khurd -1 Micro-watershed

Crops	(% to grand total)	
	Kharif	Grand Total
Red gram	96.4	96.44
Sorghum	3.6	3.6
Grand Total	100.0	100.0

Economic land evaluation

The main purpose to characterise the socio-economic systems in the watershed is to identify the existing production constraints and propose the potential/alternate options for agro-technology transfer and for bridging the adoption and yield gap.

In Chincholi Khurd 1micro-watershed, 6 soil series are identified and mapped (Table 17). The distribution of major soil series are Margutti covering an area around 221.15 ha (41.54 %) followed by Novinihala 89.85 ha (83.85 %), Bhimanahalli 103.91 ha (19.51 %), Dinsi 7.22 ha (1.36 %), Rajnala 15.84 ha (2.97%) and Mahagaon 84.66 ha (15.9 %).

Table 17: Distribution of soil series in Chincholi Khurd -1 Micro-watershed

Sl. No	Soil series	Map Description	Area in ha (%)
1	MGT iC3g1	Very shallow, black gravelly clay soils developed from weathered basalt on gently sloping uplands; sandy clay surface on 3-5 % slope, severely eroded, slightly gravelly, 15-35 per cent gravels.	51.54 (9.68)
	MGT iD3g2	Very shallow, black gravelly clay soils developed from weathered basalt on moderately sloping uplands; sandy clay surface on 5-10 % slope, severely eroded, moderately gravelly, 35-60 per cent gravels.	5.10 (0.96)
	MGT mA1	Very shallow, black gravelly clay soils developed from weathered basalt on nearly level uplands; clay surface on 0-1% slope, slightly eroded	7.71 (1.45)
	MGT mB1	Very shallow, black gravelly clay soils developed from weathered basalt on very gently sloping uplands; clay surface on 1-3 % slope, slightly eroded	5.68 (1.07)
	MGT mB2	Very shallow, black gravelly clay soils developed from weathered basalt on very gently sloping uplands; clay surface on 1-3 % slope, moderately eroded	93.02 (17.47)
	MGT mB2g1	Very shallow, black gravelly clay soils developed from weathered basalt on very gently sloping uplands; clay surface on 1-3 % slope, moderately eroded, slightly gravelly, 15-35 per cent gravels.	13.87 (2.60)
	MGT mB3g1	Very shallow, black gravelly clay soils developed from weathered basalt on very gently sloping uplands; clay surface on 1-3 % slope, severely eroded, slightly gravelly, 15-35 per cent gravels.	5.00 (0.94)
	MGT mC2	Very shallow, black gravelly clay soils developed from weathered basalt on gently sloping uplands; clay surface on 3-5 % slope, moderately eroded.	11.17 (2.10)
	MGT mC2g1	Very shallow, black gravelly clay soils developed from weathered basalt on gently sloping uplands; clay surface on 3-5 % slope, moderately eroded, slightly gravelly, 15-35 per cent gravels.	28.06 (5.27)
2	NHA mB1	Shallow, black clayey soils developed from weathered basalt on very gently sloping uplands; clay surface on 1-3% slope, slightly eroded	7.36 (1.38)
	NHA mB2	Shallow, black clayey soils developed from weathered basalt on very gently sloping uplands; clay surface on 1-3% slope, moderately eroded	82.49 (15.49)
3	BHI mB1	Shallow, black clay soils developed from weathered basalt on very gently sloping uplands; clay surface on 1-3% slope, slightly eroded	46.18 (8.67)
	BHI mB2g1	Shallow, black clay soils developed from weathered basalt on very gently sloping uplands; clay surface on 1-3% slope, moderately eroded, slightly gravelly, 15-35 per cent gravels.	25.76 (4.84)
	BHI mC2g1	Shallow, black clay soils developed from weathered basalt on gently sloping uplands; clay surface on 3-5% slope, moderately eroded, slightly gravelly, 15-35 per cent gravels.	31.97 (6.00)
4	DSI mB2	Moderately shallow, black clayey soils developed from weathered basalt on very gently sloping uplands; clay surface on 1-3% slope, moderately eroded.	7.22 (1.36)
5	RNL mB1	Deep, black clayey soils developed from weathered basalt on very gently sloping uplands; clay surface on 1-3 % slope, slightly eroded	6.57 (1.23)

	RNL mC2g1	Deep, black clayey soils developed from weathered basalt on gently sloping uplands; clay surface on 3-5 % slope, moderately eroded, slightly gravelly, 15-35 per cent gravels.	9.27 (1.74)
6	MAN mA1	Very deep, black clayey soils developed from weathered basalt on nearly level uplands; clay surface on 0-1% slope, slightly eroded	1.74 (0.33)
	MAN mB1	Very deep, black clayey soils developed from weathered basalt on very gently sloping uplands; clay surface on 1-3% slope, slightly eroded	18.17 (3.41)
	MAN mB2	Very deep, black clayey soils developed from weathered basalt on very gently sloping uplands; clay surface on 1-3% slope, moderately eroded	64.75 (12.16)
Water body			9.79 (1.84)

Present cropping pattern on different soil series are given in Table 18. Crops grown on Margutti soils are red gram and sorghum. Sorghum is grown on Novinihala soils. Red gram is grown on Bhimanahalli soils. Red gram is grown on Mahagaon soils series.

Table 18: Cropping pattern on major soil series in Chincholi Khurd -1 micro-watershed (Area in per cent)

Soil Series	Soil Depth	Crops	Dry	Grand Total
			Kharif	
MGT	Very Shallow (<25 cm)	Red gram	93.7	93.7
		Sorghum	6.3	6.3
NHA	Shallow (25-50 cm)	Red gram	100	100
BHI	shallow (50-75 cm)	Red gram	100	100
MAN	Very deep (50-75 cm)	Red gram	100	100

Land is used for agricultural use for growing cereals, pulse, oilseeds and commercial crops. The soil/ land potential are measures in terms of physical yield and net income. The alternative land use options for each micro-watershed are given below (Table 19).

Table 19: Alternative land use options for different size group of farmers (Benefit Cost Ratio) in Chincholi Khurd -1 Microwatershed.

Soil Series	Small Farmers	Medium Farmers	Large Farmers
BHI	Red gram (1.66)		
MAN	Red gram (1.00)		
MGT	Red gram (1.21)&Sorghum (1.36)	Red gram (1.90)	Red gram (1.47)
NHA		Red gram (1.65)	

The productivity of different crops grown in Chincholi Khurd -1 micro-watershed under potential yield of the crops is given in Table 20.

The data on cost of cultivation and benefit cost ratio (BCR) of different crops is given in Table 20. The total cost of cultivation in study area for red gram ranges between Rs.43620 /ha in MAN soil (with BCR of 1.00) and Rs. 23317/ha in MGT soil (with BCR

of 1.53) and sorghum the cost of cultivation in MGT soil is Rs.32338/ha (with BCR of 1.36).

Table 20: Economic land evaluation and bridging yield gap for different crops in Chincholi Khurd -1 micro-watershed

Particulars	MGT (<25 cm)		BHI (25-50 cm)	NHA (25-50 cm)	MAN (>150 cm)
	Red gram	Sorghum	Red gram	Red gram	Red gram
Total cost (Rs/ha)	23317	32338	24161	24919	43620
Gross Return (Rs/ha)	34202	43878	40087	41167	43717
Net returns (Rs/ha)	10885	11540	15926	16248	97
BCR	1.53	1.36	1.66	1.65	1.00
Farmers Practices (FP)					
FYM (t/ha)	0.9	4.1	0.9	3.2	4.4
Nitrogen (kg/ha)	78.2	18.3	78.1	25.0	19.9
Phosphorus (kg/ha)	53.9	46.7	56.1	63.9	50.9
Potash (kg/ha)	0.0	0.0	0.0	0.0	0.0
Grain (Qtl/ha)	8.8	12.2	11.6	11.9	11.1
Price of Yield (Rs/Qtl)	3867	3500	3500	3500	4000
Soil test based fertilizer Recommendation (STBR)					
FYM (t/ha)	7.4	7.4	7.4	7.4	7.4
Nitrogen (kg/ha)	24.7	61.1	24.7	24.7	24.7
Phosphorus (kg/ha)	61.8	71.0	61.8	49.4	61.8
Potash (kg/ha)	22.6	39.5	18.5	18.5	18.5
Grain (Qtl/ha)	12.4	28.4	12.4	12.4	12.4
% of Adoption/yield gap (STBR-FP) / (STBR)					
FYM (%)	87.4	45.1	87.7	57.2	40.3
Nitrogen (%)	-216.6	70.1	-216.0	-1.2	19.4
Phosphorus (%)	12.7	34.2	9.1	-29.3	17.6
Potash (%)	100.0	100.0	100.0	100.0	100.0
Grain (%)	28.5	57.1	6.1	3.6	10.4
Value of yield and Fertilizer (Rs)					
Additional Cost (Rs/ha)	6633	5717	6477	3965	3891
Additional Benefits (Rs/ha)	13621	56735	2652	1558	5152
Net change Income (Rs/ha)	6988	51017	-3825	-2406	1261

The data on FYM, Nitrogen, Phosphorus and Potash application by the farmers to different crops and recommended FYM for different crops is given in Table 20. There is a huge gap between FYM application by farmers and recommended FYM in all the crops across the soils. There is a larger yield gap in crops grown across different soil series. Adequate knowledge about recommended package of practices is the pre-requisite for their use in cultivation of crops. It is a fact that, recommended practices are major contributing factors to yield. Inadequate knowledge about recommended practices leads to their improper adoption. Strengthening of extension services by concerned agency is required to increase adoption of recommended cultivation practices and ultimately reducing the gap. By adopting soil-test fertiliser recommendation, there is scope to

increase yield and income to maximum of Rs. 51017 in sorghum and a minimum of Rs. 6988 in red gram cultivation.

Economic valuation of Ecosystem Services (ES) was aimed at combining use and non-use values to determine Total Economic Value (TEV) of ES. Ecosystem Services (ES) were valued based on their annual flow or utilization in common monetary units, Rs/year. The valuation of ES was based on market price in 2017 or market cost approaches whichever is applicable, and in other cases on value or benefit transfer from previous valuation studies.

The onsite cost of different soil nutrients lost due to soil erosion is given in Table 21 and Figure 8. The average value of soil nutrient loss is around Rs 1467.58 per ha/year. The total cost of annual soil nutrients is around Rs 767543 per year for the total area of 532.43 ha.




Figure 8: Estimation of onsite cost of soil erosion in Chincholi Khurd -1 micro-watershed

Table 21: Estimation of onsite cost of soil erosion in Chincholi Khurd -1 micro-watershed

Particulars	Quantity(kg)		Value (Rs)	
	Per ha	Total	Per ha	Total
Organic matter	196.51	102776	1238.03	647492
Phosphorus	0.12	61	5.15	2691
Potash	3.55	1858	71.06	37163
Iron	0.15	77	7.10	3713
Manganese	0.36	186	97.83	51164
Cupper	0.06	33	35.68	18662
Zinc	0.01	5	0.41	212
Sulpher	0.30	157	12.03	6294
Boron	0.01	4	0.29	153
Total	201.07	105159	1467.58	767543

The average value of ecosystem service for food grain production is around Rs 10685/year (Table 22). Per hectare food grain production services is maximum in red gram (Rs 11537) followed by sorghum (Rs 9833).

Table 22: Ecosystem services of food grain production in Chincholi Khurd -1 Micro-watershed

Production items	Crops	Area in ha	Yield (Qtl/ha)	Price (Rs/Qtl)	Gross Returns (Rs/ha)	Cost of Cultivation (Rs/ha)	Net Returns (Rs/ha)
Cereals	Sorghum	0.5	12.0	3500	42171	32338	9833
Pulses	Red gram	13.7	10.0	3729	38224	26687	11537
Average value		14.2	11.0	3615	40198	29513	10685

The average value of ecosystem service for fodder production is around Rs. 1707/ha/year (Table 23). Per hectare fodder production services is maximum in sorghum (Rs 1707).

Table 23: Ecosystem services of fodder production in Chincholi Khurd -1 Micro watershed

Production items	Crops	Area in ha	Yield (Qtl/ha)	Price (Rs/Qtl)	Net Returns (Rs/ha)
Cereals	Sorghum	0.5	2.0	850	1707

The water demand for production of different crops was worked out in arriving at the ecosystem services of water support to crop growth. The data on water requirement for producing one quintal of grain is considered for estimating the total value of water required for crop production. Per hectare value of water used and value of water was maximum (Table 24) in red gram (Rs 55810) and sorghum (Rs 36725).

Table 24: Ecosystem services of water supply in Chincholi Khurd -1 Micro-watershed

Crops	Yield (Qtl/ha)	Virtual water (cubic meter) per ha	Value of Water (Rs/ha)	Water consumption (Cubic meters/Qtl)
Red gram	10.3	5581	55810	544
Sorghum	12	3672	36725	305
Average value	22.3	4626.5	46267.5	424.5

The main farming constraints in Chincholi Khurd 1micro-watershed to be found are less rainfall, non availability fertilizers, high crop pests & diseases, animal pests & diseases, damage of crops by wild animals and non availability of plant protection chemicals. Majority of farmers depend up on money lender of the sources of loan for purpose of crop production. Farmers to sell the agriculture produce through village market and the farmers getting the agriculture related information on newspaper and television. Farmers reported that they are not getting timely support/extension services from the concerned development department (Table 25).

Table 25: Farming constraints related land resources of sample households in Chincholi Khurd -1 Micro-watershed

Sl. No	Particulars	Per cent
1	Less Rainfall	100.0
2	Lack of good quality seeds	25.0
3	Non availability Fertilizers	12.5
4	Animal Pests & Diseases	12.5
5	Lack of transportation	50.0
6	Lack of storage	62.5
7	Damage of crops by Wild Animals	87.5
8	Non availability of Plant Protection Chemicals	100.0
9	Source of loan	
	Bank	75.0
	Village merchants	25.0
9	Market for selling	
	Regulated	12.5
	Village market	87.5
10	Sources of Agri-Technology information	
	Newspaper	12.5
	Television	87.5

The findings of the study would be very much useful to the planners and policy makers of the study area to identify the irrationality in the existing production pattern and to suggest appropriate production plans for efficient utilization of their scarce resources resulting in increased net farm incomes and employment. The study also throws light on future potentialities of increasing net farm income and employment under different situations viz., with existing and recommended technology.