



हर कदम, हर डगर
किसानों का हमसफर
भारतीय कृषि अनुसंधान परिषद

AgriSearch with a human touch

**LAND RESOURCE INVENTORY AND SOCIO-ECONOMIC STATUS OF
FARM HOUSEHOLDS FOR WATERSHED PLANNING AND
DEVELOPMENT**

BETTAGERI-1 (4D4A2Q4c) MICRO WATERSHED

Alawandi Hobli, Koppal Taluk and District, Karnataka

Karnataka Watershed Development Project – II

SUJALA – III

World Bank funded Project



ICAR – NATIONAL BUREAU OF SOIL SURVEY AND LAND USE PLANNING



ICAR - NBSS & LUP



**WATERSHED DEVELOPMENT DEPARTMENT
GOVT. OF KARNATAKA, BANGALORE**



About ICAR - NBSS&LUP

The ICAR-National Bureau of Soil Survey and Land Use Planning (ICAR-NBSS&LUP), Nagpur, a premier Institute of the Indian Council of Agricultural Research (ICAR), was set up during 1976 with the objective to prepare soil resource maps at national, state and district levels and to provide research inputs in soil resource mapping and its applications, land evaluation, land use planning, land resource management, and database management using GIS for optimising land use on different kinds of soils in the country.

The Bureau has been engaged in carrying out soil resource survey, agro-ecological and soil degradation mapping at the country, state and district levels for qualitative assessment and monitoring the soil health towards viable land use planning. The research activities have resulted in identifying the soil potentials and problems, and the various applications of the soil surveys with the ultimate objective of sustainable agricultural development. The Bureau has the mandate to correlate and classify soils of the country and maintain a National Register of all the established soil series. The Institute is also imparting in-service training to staff of the soil survey agencies in the area of soil survey, land evaluation and soil survey interpretations for land use planning. The Bureau in collaboration with Panjabrao Krishi Vidyapeeth, Akola is running post-graduate teaching and research programme in land resource management, leading to M.Sc. and Ph.D. degrees.

Citation: Rajendra Hegde, Ramesh Kumar, S.C., K.V. Niranjana, S. Srinivas, M.Lalitha, B.A. Dhanorkar, R.S. Reddy and S.K. Singh (2019), "Land Resource Inventory and Socio-Economic Status of Farm Households for Watershed Planning and Development of Bettageri-1 (4D4A2Q4c) Microwatershed, Alavandi Hobli, Koppal Taluk & District, Karnataka", ICAR-NBSS&LUP Sujala MWS Publ.142, ICAR – NBSS & LUP, RC, Bangalore. P.101 & 28.

TO OBTAIN COPIES,

Please write to:

Director, ICAR - NBSS & LUP,

Amaravati Road, NAGPUR - 440 033, India

Phone : (0712) 2500386, 2500664, 2500545 (O)

Telefax : 0712-2522534

E-Mail : director@nbsslup.ernet.in

Website URL : nbsslup.in

Or

Head, Regional Centre, ICAR - NBSS&LUP, Hebbal, Bangalore - 560 024

Phone : (080) 23412242, 23510350 (O)

Telefax : 080-23510350

E-Mail : nbssrcb@gmail.com



LAND RESOURCE INVENTORY AND SOCIO-ECONOMIC STATUS OF FARM HOUSEHOLDS FOR WATERSHED PLANNING AND DEVELOPMENT

BETTAGERI-1 (4D4A2Q4c) MICRO WATERSHED

Alawandi Hobli, Koppal Taluk and District, Karnataka

Karnataka Watershed Development Project – II

Sujala-III

World Bank funded Project



ICAR – NATIONAL BUREAU OF SOIL SURVEY AND LAND USE
PLANNING



ICAR - NBSS & LUP



WATERSHED DEVELOPMENT DEPARTMENT, GOVT. OF
KARNATAKA, BANGALORE



PREFACE

In Karnataka, as in other Indian States, the livelihoods of rural people are intertwined with farming pursuits. The challenges in agriculture are seriously threatening the livelihood of a large number of farmers as they have been practicing farming in contextual factors beyond their control. Climatic factors are the most important ones and have become much more significant in recent times due to rapid climate changes induced by intensive anthropogenic activities affecting our ecosystem in multiple ways. Climate change has become the reality, it is happening and efforts to evolve and demonstrate climate resilient technologies have become essential. Due to the already over stressed scenario of agrarian sector, the climate change is resulting in manifold increase in the complexities, pushing the rural mass to face more and more unpredictable situations. The rising temperatures and unpredictable rainfall patterns are going to test seriously the informed decisions farmers have to make in order to survive in farming and sustain their livelihood.

It is generally recognized that impacts of climate change shall not be uniform across the globe. It is said that impact of climate change is more severe in South Asia. Based on the analysis of meteorological data, it is predicted that in India, there will be upward trend in mean temperature, downward trend in relative humidity, annual rainfall and number of wet days in a year. Also, in general, phenomena like erratic monsoon, spread of tropical diseases, rise in sea levels, changes in availability of fresh water, frequent floods, droughts, heat waves, storms and hurricanes are predicted. Each one of these adverse situations are already being experienced in various parts of India and also at the global level. Decline in agricultural productivity of small and marginal farmers becoming more vulnerable is already witnessed.

In Karnataka, more than 60 per cent of the population live in rural areas and depend on agriculture and allied activities for their livelihood. Though the state has achieved significant progress in increasing the yield of many crops, there is tremendous pressure on the land resources due to the growing and competing demands of various land uses. This is reflected in the alarming rate of land degradation observed. Already more than 50 per cent of the area is affected by various forms of degradation. If this trend continues, the sustainability of the fragile ecosystem will be badly affected. The adverse effects of change in the climatic factors are putting additional stress on the land resources and the farmers dependent on this.

The natural resources (land, water and vegetation) of the state need adequate and constant care and management, backed by site-specific technological interventions and investments particularly by the government. Detailed database pertaining to the nature of

the land resources, their constraints, inherent potentials and suitability for various land based rural enterprises, crops and other uses is a prerequisite for preparing location-specific action plans, which are in tune with the inherent capability of the resources. Any effort to evolve climate resilient technologies has to be based on the baseline scientific database. Then only one can expect effective implementation of climate resilient technologies, monitor the progress, make essential review of the strategy, and finally evaluate the effectiveness of the implemented programs. The information available at present on the land resources of the state are of general nature and useful only for general purpose planning. Since the need of the hour is to have site-specific information suitable for farm level planning and detailed characterization and delineation of the existing land resources of an area into similar management units is the only option.

ICAR-NBSS&LUP, Regional Centre, Bangalore has taken up a project sponsored by the Karnataka Watershed Development Project-II, (Sujala-III), Government of Karnataka funded by the World Bank under Component -1 Land Resource Inventory. This study was taken up to demonstrate the utility of such a database in reviewing, monitoring and evaluating all the land based watershed development programs on a scientific footing. To meet the requirements of various land use planners at grassroots level, the present study on “Land Resource Inventory and Socio-Economic Status of Farm Households for Watershed Planning and Development of Bettageri-1 microwatershed in Koppal Taluk and District, Karnataka” for integrated development was taken up in collaboration with the State Agricultural Universities, IISC, KRSAC, KSNDMC as Consortia partners. The project provides detailed land resource information at cadastral level (1:7920 scale) for all the plots and socio-economic status of farm households covering thirty per cent farmers randomly selected representing landed and landless class of farmers in the microwatershed. The project report with the accompanying maps for the microwatershed will provide required detailed database for evolving effective land use plan, alternative land use options and conservation plans for the planners, administrators, agricultural extension personnel, KVK officials, developmental departments and other land users to manage the land resources in a sustainable manner.

It is hoped that this database will be useful to the planners, administrators and developmental agencies working in the area in not only for formulating location specific developmental schemes but also for their effective monitoring at the village/watershed level.

Nagpur

Date:17.12.2018

S.K. SINGH

Director, ICAR - NBSS&LUP, Nagpur

Contributors

Dr. RajendraHegde Principal Scientist, Head & Project Leader, Sujala-III Project ICAR-NBSS&LUP, Regional Centre, Bangalore	Dr. S.K.Singh Director, ICAR-NBSS&LUP Coordinator, Sujala-III Project Nagpur
Soil Survey, Mapping & Report Preparation	
Dr. K.V. Niranjana	Sh. R.S. Reddy
Dr. B.A. Dhanorkar	Smt. Chaitra, S.P.
	Dr. GopaliBardhan
	Sh. Somashekar T.N
	Ms. Arpitha M
	Dr. Mahendra
Field Work	
Sh. C.Bache Gowda	Sh. MayurPatil
Sh. Somashekar	Sh. Arun Kumar, S.
Sh. M. Jayaramaiah	Sh. Sunil Raj
	Sh. Yogesh Kumar, B.
	Sh. Vikas, N.K.
	Sh. Arun Kumar, S.G.
	Sh. UmeshJadiyappaMadolli
	Sh. Praveen Kumar P. Achalkar
	Sh. Veerabhadraswamy
	Sh. Vinay
	Sh. Shankarappa, K.
	Sh. Lankesh, R.S.
	Sh. Appanna B. Hattigoudar
	Sh. Maharudra
GIS Work	
Dr. S.Srinivas	Sh. A.G. Devendra Prasad
Sh. D.H.Venkatesh	Sh. AbhijithSastry, N.S.
Smt.K.Sujatha	Sh. NagendraBabuKolukondu
Smt. K.V.Archana	Sh. Avinash
Sh. N.Maddileti	Sh. Amar Suputhra, S.
	Sh. Deepak M.J.
	Sh. Madappaswamy
	Smt. K.Karunya Lakshmi
	Ms. Seema, K.V.
	Ms. Ramireddy Lakshmi Silpa
	Ms. BhanuRekha, T.
	Ms. RajataBhat
	Ms. Shruthi
	Ms. Suman, S.

Laboratory Analysis	
Dr. M. Lalitha	Ms. Thara, V.R.
Smt. ArtiKoyal	Ms. Roopa, G.
Smt. Parvathy, S.	Ms. Vindhya, N.G.
	Ms. Shwetha N.K.
	Ms. PavanaKumari, P.
	Ms. Leelavathy, K.U.
	Ms. Rashmi, N.
	Ms. Padmaja, S.
	Ms. Veena, M.
	Ms. Chaithrashree B
	Ms. Shwetha N
Socio-economic Analysis	
Dr. Ramesh Kumar, S.C.	Sh. M.K. Prakashanaik
	Ms. Karuna V. Kulkarni
	Mrs. Sowmya A.N
	Sh. Vinod R
	Sh. Basavaraja
	Sh. Vijay Kumar Lamani
	Ms. Sowmya K.B
	Mrs. Prathibha, D.G
	Sh. Rajendra,D
Soil & Water Conservation	
Sh. Sunil P. Maske	
Watershed Development Department, GoK, Bangalore	
Sh. Rajeev Ranjan IFS Project Director & Commissioner, WDD	Dr. A. Natarajan NRM Consultant, Sujala-III Project
Dr. S.D. Pathak IFS Executive Director & Chief Conservator of Forests, WDD	

PART-A

LAND RESOURCE INVENTORY

Contents

Preface		
Contributors		
Executive Summary		
Chapter 1	Introduction	1
Chapter 2	Geographical Setting	3
2.1	Location and Extent	3
2.2	Geology	3
2.3	Physiography	4
2.4	Drainage	5
2.5	Climate	5
2.6	Natural Vegetation	6
2.7	Land Utilization	7
Chapter 3	Survey Methodology	11
3.1	Base maps	11
3.2	Image Interpretation for Physiography	11
3.3	Field Investigation	14
3.4	Soil mapping	16
3.5	Laboratory Characterization	16
3.6	Land Use Classes	16
Chapter 4	The Soils	21
4.1	Soils of Granite Gneiss Landscape	21
4.2	Soils of Alluvial Landscape	24
Chapter 5	Interpretation for Land Resource Management	33
5.1	Land Capability Classification	33
5.2	Soil Depth	35
5.3	Surface Soil Texture	36
5.4	Soil Gravelliness	37
5.5	Available Water Capacity	37
5.6	Soil Slope	39
5.7	Soil Erosion	39
Chapter 6	Fertility Status	41
6.1	Soil Reaction (pH)	41
6.2	Electrical Conductivity (EC)	41
6.3	Organic Carbon (OC)	41
6.4	Available Phosphorus	42
6.5	Available Potassium	44
6.6	Available Sulphur	44
6.7	Available Boron	44

6.8	Available Iron	45
6.9	Available Manganese	45
6.10	Available Copper	45
6.11	Available Zinc	45
Chapter 7	Land Suitability for Major Crops	49
7.1	Land suitability for Sorghum	49
7.2	Land suitability for Maize	52
7.3	Land suitability for Bajra	53
7.4	Land suitability for Groundnut	54
7.5	Land suitability for Sunflower	55
7.6	Land suitability for Chilli	56
7.7	Land suitability for Tomato	58
7.8	Land suitability for Drumstick	59
7.9	Land suitability for Mulberry	60
7.10	Land suitability for Mango	61
7.11	Land suitability for Sapota	63
7.12	Land suitability for Pomegranate	64
7.13	Land Suitability for Guava	65
7.14	Land Suitability for Jackfruit	67
7.15	Land Suitability for Jamun	68
7.16	Land Suitability for Musambi	69
7.17	Land Suitability for Lime	71
7.18	Land Suitability for Cashew	72
7.19	Land Suitability for Custard apple	73
7.20	Land suitability for Amla	75
7.21	Land suitability for Tamarind	76
7.22	Land suitability for Marigold	77
7.23	Land suitability for Chrysanthemum	79
7.24	Land suitability for Jasmine	80
7.25	Land Suitability for Crossandra	81
7.26	Proposed Crop Plan	83
Chapter 8	Soil Health Management	87
Chapter9	Soil and Water conservation Treatment Plan	93
9.1	Treatment Plan	94
9.2	Recommended Soil and Water Conservation measures	97
9.3	Greening of microwatershed	98
	References	101
	Appendix I	I-V
	Appendix II	VII-XI
	Appendix III	XIII-XVII

LIST OF TABLES

2.1	Mean Monthly Rainfall, PET, ½ PET atKoppalTaluk and District	5
2.2	Land Utilization in Koppal District	7
3.1	Differentiating Characteristics used for Identifying Soil Series	15
3.2	Soil map unit description of Bettageri-1microwatershed	19
4.1	Physical and chemical characteristics of soil series identified in Bettageri-1microwatershed	27
7.1	Soil-Site Characteristics of Bettageri-1microwatershed	50
7.2	Land suitability for Sorghum	51
7.3	Land suitability for Maize	52
7.4	Land suitability forbajra	53
7.5	Land suitability for Groundnut	54
7.6	Land suitability for Sunflower	55
7.7	Land suitability for Chilli	57
7.8	Land suitability for Tomato	58
7.9	Land suitability for Drumstick	59
7.10	Land suitability for Mulberry	61
7.11	Land suitability for Mango	62
7.12	Land Suitability for Sapota	63
7.13	Land suitability for Pomegranate	65
7.14	Land suitability for Guava	66
7.15	Land suitability for jackfruit	67
7.16	Land suitability forjamun	68
7.17	Land suitability forMusambi	70
7.18	Land Suitability for Lime	71
7.19	Land suitability for Cashew	73
7.20	Land suitability for custard apple	74
7.21	Land suitability forAmla	75
7.22	Land suitability for Tamarind	76
7.23	Land suitability for Marigold	78
7.24	Land suitability for chrysanthemum	79

7.25	Land suitability for Jasmine	80
7.26	Proposed Crop Plan for Bettageri-1Microwatershed	84

LIST OF FIGURES

2.1	Location map of Bettageri-1Microwatershed	3
2.2a	Granite and granite gneiss rocks	4
2.2b	Alluvium	4
2.3	Rainfall distribution in KoppalTaluk,Koppal District	6
2.4	Natural vegetation of Bettageri-1microwatershed	6
2.5a	Different crops and cropping systems in Bettageri-1 Microwatershed	8
2.5b	Different crops and cropping systems in Bettageri-1 Microwatershed	9
2.6	Current Land use – Bettageri-1 Microwatershed	10
2.7	Location of Wells- Bettageri-1 Microwatershed	10
3.1	Scanned and Digitized Cadastral map ofBettageri-1Microwatershed	13
3.2	Satellite image of Bettageri-1Microwatershed	13
3.3	Cadastral map overlaid on IRS PAN+LISS IV merged imagery of Bettageri-1Microwatershed	14
3.4	Location of profiles in a transect	14
3.5	Soil phase or management units of Bettageri-1 Microwatershed	17
5.1	Land Capability Classification ofBettageri-1Microwatershed	34
5.2	Soil Depth map ofBettageri-1Microwatershed	35
5.3	Surface Soil Texture map ofBettageri-1Microwatershed	36
5.4	Soil Gravelliness map ofBettageri-1Microwatershed	37
5.5	Soil Available Water Capacity map ofBettageri-1Microwatershed	38
5.6	Soil Slope map ofBettageri-1Microwatershed	39
5.7	Soil Erosion map ofBettageri-1Microwatershed	40
6.1	Soil Reaction (pH) map ofBettageri-1Microwatershed	42
6.2	Electrical Conductivity (EC) map ofBettageri-1Microwatershed	43
6.3	Soil Organic Carbon (OC) map ofBettageri-1Microwatershed	43
6.4	Soil Available Phosphorus map ofBettageri-1Microwatershed	44
6.5	Soil Available Potassium map ofBettageri-1Microwatershed	45
6.6	Soil Available Sulphur map ofBettageri-1Microwatershed	46
6.7	Soil Available Boron map ofBettageri-1Microwatershed	46
6.8	Soil Available Iron map ofBettageri-1Microwatershed	47
6.9	Soil Available Manganese map ofBettageri-1Microwatershed	47
6.10	Soil Available Copper map ofBettageri-1Microwatershed	48

6.11	Soil Available Zinc map of Bettageri-1 Microwatershed	48
7.1	Land suitability for Sorghum	51
7.2	Land suitability for Maize	52
7.3	Land suitability for Bajra	53
7.4	Land suitability for Groundnut	55
7.5	Land suitability for Sunflower	56
7.6	Land suitability for Chilli	57
7.7	Land suitability for Tomato	59
7.8	Land suitability for Drumstick	60
7.9	Land suitability for Mulberry	61
7.10	Land suitability for Mango	62
7.11	Land suitability for Sapota	64
7.12	Land suitability for Pomegranate	65
7.13	Land Suitability for Guava	66
7.14	Land Suitability for Jackfruit	68
7.15	Land Suitability for Jamun	69
7.16	Land Suitability for Musambi	70
7.17	Land Suitability for Lime	72
7.18	Land Suitability for Cashew	73
7.19	Land Suitability for Custard apple	74
7.20	Land suitability for Amla	76
7.21	Land suitability for Tamarind	77
7.22	Land suitability for Marigold	78
7.23	Land suitability for Chrysanthemum	79
7.24	Land suitability for Jasmine	81
7.25	Land suitability for Crossandra	81
7.26	Land Management Units map of Bettageri-1 micro watershed	83
9.1	Soil and water conservation map of Bettageri-1 Microwatershed	98

EXECUTIVE SUMMARY

The land resource inventory of Bettageri-1microwatershed was conducted using village cadastral maps and IRS satellite imagery on 1:7920 scale. The false colour composites of IRS imagery were interpreted for physiography and these physiographic delineations were used as base for mapping soils. The soils were studied in several transects and a soil map was prepared with phases of soil series as mapping units. Random checks were made all over the area outside the transects to confirm and validate the soil map unit boundaries. The soil map shows the geographic distribution and extent, characteristics, classification, behavior and use potentials of the soils in the microwatershed.

The present study covers an area of 391ha in Koppal taluk and district, Karnataka. The climate is semiarid and categorized as drought - prone with an average annual rainfall of 662 mm, of which about 424 mm is received during south –west monsoon, 161mm during north-east and the remaining 77 mm during the rest of the year. An area of about 91 per cent is covered by soils, 9 per cent by habitation and water bodies. The salient findings from the land resource inventory are summarized briefly below.

- ❖ The soils belong to 8 soil series and 9 soil phases (management units) and 7 land use classes.*
- ❖ The length of crop growing period is <90 days and starts from 2nd week of August to 2nd week of November.*
- ❖ From the master soil map, several interpretative and thematic maps like land capability, soil depth, surface soil texture, soil gravelliness, available water capacity, soil slope and soil erosion were generated.*
- ❖ Soil fertility status maps for macro and micronutrients were generated based on the surface soil samples collected at every 250 m grid interval.*
- ❖ Land suitability for growing 24 major agricultural and horticultural crops were assessed and maps showing the degree of suitability along with constraints were generated.*
- ❖ Entire area is suitable for agriculture.*
- ❖ About 42 per cent of the soils are shallow to moderately shallow (25-75 cm) and 49 per cent area are moderately deep to deep (75-150 cm).*
- ❖ About 46 per cent area has loamy soils and 45 per cent area has clayey soils at the surface.*
- ❖ About 5 per cent of the area has non-gravelly soils, 44 per cent gravelly soils (15-35 % gravel) and 41 per cent very gravelly (35- 60% gravel) soils.*
- ❖ About 63 per cent area has very low to low (<50-100 mm/m), 23 per cent medium (101-150 mm/m) and 5 per cent very high (>200mm/m) in available water capacity.*
- ❖ Entire area has very gently sloping (1-3%) lands.*
- ❖ About 33 per cent area is slightly eroded and about 58 per cent area is moderately eroded (e2) lands.*

- ❖ *About 6 per cent area is slightly acidic (pH 6.0-6.5) about 18 per cent neutral, 67 per cent area is slightly alkaline (pH 8.4 to 9.0) to very strongly alkaline (pH > 9.0) in soil reaction.*
- ❖ *The Electrical Conductivity (EC) of the soils are dominantly $< 2 \text{ dsm}^{-1}$ indicating that the soils are non-saline.*
- ❖ *Organic carbon is low ($< 0.5\%$) in about 20 per cent, 39 per cent of the soils are medium (0.5-0.75%) and 32 per cent of the soils are high ($> 0.75\%$) in organic carbon.*
- ❖ *About 6 per cent area is low ($< 23 \text{ kg/ha}$), medium (23-57 kg/ha) in major area of 71 per cent and 14 per cent area is high ($> 57 \text{ kg/ha}$) available phosphorus.*
- ❖ *Major area of about 86 per cent is medium (145-337 kg/ha) in available potassium and a small area of about 5 per cent has high ($> 337 \text{ kg/ha}$) in available potassium.*
- ❖ *Available sulphur is low ($< 10 \text{ ppm}$) in 66 per cent area, medium (10-20 ppm) in about 24 per cent area and about < 1 per cent area is high ($> 20 \text{ ppm}$).*
- ❖ *Available boron is low (0.5 ppm) in about 60 per cent area and medium (0.5-1.0 ppm) in 31 per cent area.*
- ❖ *Available iron is sufficient ($> 4.5 \text{ ppm}$) in 40 per cent area and deficient ($< 4.5 \text{ ppm}$) in about 51 per cent area.*
- ❖ *Available zinc is deficient ($< 0.6 \text{ ppm}$) in an area of about 36 per cent and sufficient in 55 per cent area.*
- ❖ *Available manganese and copper are sufficient in all the soils.*
- ❖ *The land suitability for 25 major crops grown in the micro watershed were assessed and the areas that are highly suitable (S1) and moderately suitable (S2) are given below. It is however to be noted that a given soil may be suitable for various crops but what specific crop to be grown may be decided by the farmer looking to his capacity to invest on various inputs, marketing infrastructure, market price and finally the demand and supply position.*

Land suitability for various crops in the microwatershed

Crop	Suitability Area in ha (%)		Crop	Suitability Area in ha (%)	
	Highly suitable (S1)	Moderately suitable (S2)		Highly suitable (S1)	Moderately suitable (S2)
<i>Sorghum</i>	20 (5)	143 (37)	<i>Guava</i>	-	101 (26)
<i>Maize</i>	-	142 (36)	<i>Jackfruit</i>	-	101 (26)
<i>Bajra</i>	88 (23)	54 (14)	<i>Jamun</i>	-	108 (28)
<i>Groundnut</i>	-	142 (36)	<i>Musambi</i>	20 (5)	101 (26)
<i>Sunflower</i>	20 (5)	101 (26)	<i>Lime</i>	20 (5)	101 (26)
<i>Chilli</i>	-	142 (36)	<i>Cashew</i>	-	160 (41)
<i>Tomato</i>	-	142 (36)	<i>Custard apple</i>	108 (28)	127 (32)
<i>Drumstick</i>	-	121 (31)	<i>Amla</i>	88 (23)	147 (37)
<i>Mulberry</i>	-	193 (49)	<i>Tamarind</i>	-	20 (5)
<i>Mango</i>	-	-	<i>Marigold</i>	-	163 (42)
<i>Sapota</i>	-	101 (26)	<i>Chrysanthemum</i>	-	163 (42)
<i>Pomegranate</i>	-	121 (31)	<i>Jasmine</i>	-	143 (37)
			<i>Crossandra</i>	-	145 (36)

Apart from the individual crop suitability, a proposed crop plan has been prepared for the 7 identified LUCs by considering only the highly and moderately suitable lands for different crops and cropping systems with food, fodder, fibre and other horticulture crops.

- ❖ Maintaining soil-health is vital to crop production and conserve soil and land resource base for maintaining ecological balance and to mitigate climate change. For this, several ameliorative measures have been suggested to these problematic soils like saline/alkali, highly eroded, sandy soils etc.,
- ❖ Soil and water conservation treatment plan has been prepared that would help in identifying the sites to be treated and also the type of structures required.
- ❖ As part of the greening programme, several tree species have been suggested to be planted in marginal and submarginal lands, field bunds and also in the hillocks, mounds and ridges. That would help in supplementing the farm income, provide fodder and fuel, and generate lot of biomass which in turn would help in maintaining the ecological balance and contribute to mitigating the climate change.

INTRODUCTION

Soil is a finite natural resource that is central to sustainable agriculture and food security. Over the years, this precious resource is faced with the problems of erosion, salinity, alkalinity, degradation, depletion of nutrients and even decline in availability of land for agriculture. It is a known fact, that it takes thousands of years to form a few centimetres of soil, thus, soil is a precious gift of nature. The area available for agriculture is about 51 per cent of the total geographical area and more than 60 per cent of the people are still dependant on agriculture for their livelihood. However, the capacity of a soil to produce is limited and the limits to the production are set by its intrinsic characteristics, agro climatic setting, and use and management. There is, therefore, tremendous pressure on land and water resources, which is causing decline in soil-health and stagnation in productivity. As much as 121 m ha of land is reportedly degraded which leads to impaired soil quality. It is imperative that steps are urgently taken to check and reverse land degradation without any further loss of time. The improvements in productivity will have to come from sustainable intensification measures that make the most effective use of land and water resources. Soil erosion alone has degraded about 35 lakh ha. Almost all the uncultivated areas are facing various degrees of degradation, particularly soil erosion; salinity and alkalinity has emerged as a major problem in more than 3.5 lakh ha in the irrigated areas of the State. Nutrient depletion and declining factor productivity is common in both rainfed and irrigated areas. The degradation is continuing at an alarming rate and there appears to be no systematic effort among the stakeholders to contain this process. In recent times, an aberration of weather due to climate change phenomenon has added another dimension leading to unpredictable situations to be tackled by the farmers.

In this critical juncture, the challenge before us is not only to increase the productivity per unit area which is steadily declining and showing a fatigue syndrome, but also to prevent or at least reduce the severity of degradation. If the situation is not reversed at the earliest, then the sustainability of the already fragile crop production system and the overall ecosystem will be badly affected in the state.

Added to this, every year there is a significant diversion of farm lands and water resources for non-agricultural purposes. Thus, developing strategies to slow down the degradation process or reclaim the soils to normal condition and ensure sustainability of production system are the major issues today. This demands a systematic appraisal of our soil and land resources with respect to their extent, geographic distribution, characteristics, behaviour and use potential, which is very important for developing an effective land use and cropping systems for augmenting agricultural production on a sustainable basis. The soil and land resource inventories made so far in Karnataka had limited utility because the surveys were of different types, scales and intensities carried out at different times with specific objectives. Hence, there is an urgent need to generate

detailed site-specific farm level database on various land resources for all the villages/watersheds in a time bound manner that would help to protect the valuable soil and land resources and also to stabilize the farm production.

Therefore, the land resource inventory required for farm level planning is the one which investigates all the parameters which are critical for productivity *viz.*, soils, site characteristics like slope, erosion, gravelliness and stoniness, climate, water, topography, geology, hydrology, vegetation, crops, land use pattern, animal population, socio-economic conditions, infrastructure, marketing facilities and various schemes and developmental works of the government etc. From the data collected at farm level, the specific problems and potentials of the area can be identified and highlighted, conservation measures required for the area can be planned on a scientific footing, suitability of the area for various uses can be worked out and finally viable and sustainable land use options suitable for each and every land holding can be prescribed.

The Land Resource Inventory is basically done for identifying potential and problem areas, developing sustainable land use plans, estimation of surface run off and water harvesting potential, preparation of soil and water conservation plans, land degradation/desertification etc. The Bureau is presently engaged in developing an LRI methodology using high resolution satellite remote sensing data and Digital Elevation Model (DEM) data to prepare Landscape Ecological Units (LEU) map representing agro-ecosystem as a whole. The LEU is preferred over landform as the base map for LRI. LEU is the assemblage of landform, slope and land use. An attempt was made to upscale the soil resource information from 1:250000 and 1:50000 scale to the LEU map in Goa and other states.

The land resource inventory aims to provide site specific database for Bettageri-1 microwatershed in Koppal Taluk and District, Karnataka state for the Karnataka Watershed Development Department. The database was generated by using cadastral map of the village as a base along with high resolution IRS LISS IV and Cartosat-1 merged satellite imagery. Later, an attempt will be made to uplink this LRI data generated at 1:7920 scale under Sujala-III Project to the proposed Landscape Ecological Units (LEUs) map.

The study was organized and executed by the ICAR- National Bureau of Soil Survey and Land Use Planning, Regional Centre, Bangalore under Generation of Land Resource Inventory Data Base Component-1 of the Sujala-III Project funded by the World Bank.

GEOGRAPHICAL SETTING

2.1 Location and Extent

The Bettageri-1 Microwatershed is located in the central part of northern Karnataka in Koppal Taluk, Koppal District, Karnataka State (Fig.2.1). It comprises parts of Bettageri village. It lies between $15^{\circ}11'$ and $15^{\circ}13'$ North latitudes and $76^{\circ}01'$ and $76^{\circ}03'$ East longitudes and covers an area of 391 ha. It is about 40 km southwest of Koppal town and is surrounded by Bettageri village on all parts of the microwatershed.




Fig.2.1 Location map of Bettageri-1 Microwatershed

2.2 Geology

Major rock formations observed in the microwatershed are granite gneiss and alluvium (Figs.2.2a and b). Granite gneisses are essentially pink to gray and are coarse to medium grained. They consist primarily of quartz, feldspar, biotite and hornblende. The gray granite gneisses are highly weathered, fractured and fissured upto a depth of about

10 m. Dolerite dykes and quartz veins are common with variable width and found to occur in Gudigeri-3 village. The soil thickness of the alluvium generally is limited to less than a meter, except in river valleys where it is very deep extending to tens of meters. Such soils are transported and represent paleo black soil originally formed at higher elevation, but now occupying river valleys.



Fig.2.2 Granite and granite gneiss rocks



Fig.2.2 b Alluvium

2.3 Physiography

Physiographically, the area has been identified as Granite gneiss and Alluvial landscapes based on geology. The microwatershed area has been further divided into mounds/ridges, summits, side slopes and very gently sloping uplands and nearly level

plains based on slope and its relief features. The elevation ranges from 511 to 565 m in the gently sloping uplands. The mounds and ridges are mostly covered by rock outcrops.

2.4 Drainage

The area is drained by several small seasonal streams that join Hire *halla* and Chenna *halla* along its course. Though, the streams are not perennial, during rainy season they carry large quantities of rain water. The microwatershed has only few small tanks which are not able to store the water flowing during the rainy season. Due to this, the ground water recharge is very much affected in the village. This is reflected in the failure of many bore wells in the village. If the available rain water is properly harnessed by constructing tanks and recharge structures at appropriate places in the villages, then the drinking and irrigation needs of the area can be easily met. The drainage network is dendritic to sub parallel.

2.5 Climate

The district falls under semiarid tract of the state and is categorized as drought - prone with total annual rainfall of 662 mm (Table 2.1) of this maximum of 424 mm precipitation takes place during south–west monsoon period from June to September, north-east monsoon contributes about 161 mm and prevails from October to early December and the remaining 77 mm received during the rest of the year. The winter season is from December to February. During April and May, the temperatures reach up to 45°C and in December and January, the temperatures will go down to 16°C. Rainfall distribution is shown in Figure 2.3. The average Potential Evapo Transpiration (PET) is 145 mm and varies from a low of 101 mm in December to 193 mm in the months of May. The PET is always higher than precipitation in all the months except in the month of September. Generally, the Length of crop Growing Period (LGP) is <90 days and starts from 2nd week of August to 2nd week of November.

Table 2.1 Mean Monthly Rainfall, PET, 1/2 PET at Koppal Taluk and District

Sl.No.	Months	Rainfall	PET	1/2 PET
1	January	1.60	116.70	58.35
2	February	1.50	129.20	64.60
3	March	14.10	169.80	84.90
4	April	18.10	180.60	90.30
5	May	41.60	193.50	96.75
6	June	85.80	167.90	83.95
7	July	72.10	156.20	78.10
8	August	110.50	152.50	76.25
9	September	155.60	138.50	69.25
10	October	116.30	122.30	61.15
11	November	36.00	106.40	53.20
12	December	9.10	101.00	50.50
TOTAL		662.30	144.55	




Fig. 2.3 Rainfall distribution in Koppal Taluk and District

2.6 Natural Vegetation

The natural vegetation is sparse comprising few tree species, shrubs and herbs. The mounds, ridges and boulders occupy sizeable areas which are under thin to moderately thick forest vegetation. Still, there are some remnants of the past forest cover which can be seen in patches in some ridges and hillocks in the microwatershed Fig (2.4).

Apart from the continuing deforestation, the presence of large population of goats, sheep and other cattle in the microwatershed is causing vegetative degradation of whatever little vegetation left in the area. The uncontrolled grazing has left no time for the regeneration of the vegetative cover. This leads to the accelerated rate of erosion on the hill slopes, resulting in the formation of deep gullies in the foot slopes and eventually resulting in the heavy siltation of few tanks and reservoirs in the microwatershed.



Fig 2.4 Natural vegetation of Bettageri-1 microwatershed

2.7 Land Utilization

About 91 per cent area (Table 2.2) in Koppal district is cultivated at present and about 17 per cent of the area is sown more than once. An area of about 3 per cent is currently barren. Forests occupy a small area of about 5 per cent and the tree cover is in a very poor state. Most of the mounds, ridges and bouldery areas have very poor vegetative cover. Major crops grown in the area are sorghum, maize, bajra, cotton, safflower, sunflower, red gram, horse gram, onion, mulberry, pomegranate, sugarcane, bengalgram and groundnut (Fig 2.5). While carrying out land resource inventory, the land use/land cover particulars are collected from all the survey numbers and a current land use map of the microwatershed is prepared. The current land use map prepared shows the arable and non-arable lands, other land uses and different types of crops grown in the area. The current land use map of Bettageri-1 Microwatershed is presented in Fig.2.6. Simultaneously, enumeration of existing wells (bore wells and open wells) and other soil and water conservation structures in the microwatershed is made and their location in different survey numbers is marked on the cadastral map. Map showing the location of wells and other water bodies in Bettageri-1 Microwatershed is given Fig.2.7.

Table 2.2 Land Utilization in Koppal District

Sl.No.	Agricultural land use	Area (ha)	Per cent
1	Total geographical area	552495	
2	Total cultivated area	500542	90.6
3	Area sown more than once	92696	16.8
4	Trees and groves	210	0.04
5	Cropping intensity	-	118
6	Forest	29451	5.33
7	Cultivable wasteland	2568	0.46
8	Permanent Pasture land	14675	2.66
9	Barren land	16627	3.01
10	Non agricultural land	40591	7.35
11	Current fallow	19660	3.56



Fig.2.5 (a) Different crops and cropping systems in Bettageri-1 Microwatershed



Fig.2.5 (b) Different crops and cropping systems in Bettageri-1 Microwatershed




Fig.2.6 Current Land Use – Bettageri-1 Microwatershed




Fig.2.7 Location of wells - Bettageri-1 Microwatershed

SURVEY METHODOLOGY

The purpose of land resource inventory is to delineate similar areas (soil series and phases), which respond or expected to respond similarly to a given level of management. This was achieved in Bettageri-1 Microwatershed by the detailed study of all the soil characteristics (depth, texture, colour, structure, consistence, coarse fragments, porosity, soil reaction, soil horizons etc.) and site (slope, erosion, drainage, occurrence of rock fragments etc.) followed by grouping of similar areas based on soil-site characteristics into homogeneous (management units) units and showing their extent and geographic distribution on the microwatershed cadastral map. The detailed soil survey at 1:7920 scale was carried out in 391 ha area. The methodology followed for carrying out land resource inventory was as per the guidelines given in Soil Survey Manual (IARI, 1971; Soil Survey Staff, 2006; Natarajan *et al.*, 2015) which is briefly described below.

3.1 Base Maps

The detailed survey of the land resources occurring in the microwatershed was carried out by using digitized cadastral map as a base. The cadastral map shows field boundaries with their survey numbers, location of tanks, streams and other permanent features of the area (Fig. 3.1). Apart from the cadastral map, remote sensing data products from Cartosat-1 and LISS IV merged at the scale of 1:7920 were used in conjunction with the cadastral map to identify the geology, landscapes, landforms and other surface features. The imagery helped in the identification and delineation of boundaries between hills, uplands and lowlands, water bodies, forest and vegetated areas, roads, habitations and other cultural features of the area (Fig.3.2).The cadastral map was overlaid on the satellite imagery (Fig.3.3) that helps to identify the parcel boundaries and other permanent features. Apart from cadastral maps and images, toposheets of the area (1:50,000 scale) were used for initial traversing, identification of geology, landscapes and landforms, drainage features, present land use and also for selection of transects in the microwatershed.

3.2 Image Interpretation for Physiography

False Colour Composites (FCC) of Cartosat-I and LISS-IV merged satellite data covering the microwatershed area was visually interpreted using image interpretation elements and all the available collateral data with local knowledge. The delineated physiographic boundaries were transferred on to a cadastral map overlaid on satellite imagery. Physiographically, the area has been identified as granite gneiss and alluvial landscapes and is divided into landforms such as ridges, mounds and uplands based on slope. They were further subdivided into physiographic/ image interpretation units based on image characteristics. The image interpretation legend for Physiography is given below.

Image Interpretation Legend for Physiography

G- Granite gneiss landscape

- G1 Hills/ Ridges/ Mounds
 - G11 Summits
 - G12 Side slopes
 - G121 Side slopes with dark grey tones
- G2 Uplands
 - G21 Summits
 - G22 Gently sloping uplands
 - G221 Gently sloping uplands, yellowish green (eroded)
 - G222 Gently sloping uplands, yellowish white (severely eroded)
 - G23 Very gently sloping uplands
 - G231 Very gently sloping uplands, yellowish green
 - G232 Very gently sloping uplands, medium green and pink
 - G233 Very gently sloping uplands, pink and green (scrub land)
 - G234 Very gently sloping uplands, medium greenish grey
 - G235 Very gently sloping uplands, yellowish white (eroded)
 - G236 Very gently sloping uplands, dark green
 - G237 Very gently sloping uplands, medium pink (coconut garden)
 - G238 Very gently sloping uplands, pink and bluish white (eroded)

DSe Alluvial landscape

DSe 1 Summit

- DSe 11 Nearly level Summit with dark grey tone
- DSe 12 Nearly level Summit with medium grey tone
- DSe 13 Nearly level Summit with whitish grey tone
- DSe 14 Nearly level Summit with whitish tone (Calcareousness)
- DSe 15 Nearly level Summit with pinkish grey tone
- DSe 16 Nearly level Summit with medium pink tone
- DSe 17 Nearly level Summit with bluish white tone
- DSe 18 Nearly level Summit with greenish grey tone

DSe 2 Very gently sloping

- DSe 21 Very gently sloping, whitish tone
- DSe 22 Very gently sloping, greyish pink tone
- DSe 23 Very gently sloping, whitish grey tone
- DSe 24 Very gently sloping, medium grey tone
- DSe 25 Very gently sloping, medium pink tone
- DSe 26 Very gently sloping, dark grey tone
- DSe 27 Very gently sloping, bluish grey tone
- DSe 28 Very gently sloping, greenish grey tone
- DSe 29 Very gently sloping, Pinkish grey




Fig 3.1 Scanned and Digitized Cadastral map of Bettageri-1 Microwatershed




Fig.3.2 Satellite Image of Bettageri-1 Microwatershed




Fig.3.3 Cadastral map overlaid on IRS PAN+LISS IV merged imagery of Bettageri-1 Microwatershed

3.3 Field Investigation

The field boundaries and survey numbers given on the cadastral sheet were located on the ground by following permanent features like roads, cart tracks, *nallas*, streams, tanks etc., and wherever changes were noticed, they were incorporated on the microwatershed cadastral map. Preliminary traverse of the microwatershed was carried out with the help of cadastral map, imagery and toposheets. While traversing, landforms and physiographic units identified were checked and preliminary soil legend was prepared by studying soils at few selected places. Then, intensive traversing of each physiographic unit like hills, ridges, uplands and plains was carried out. Based on the variability observed on the surface, transects (Fig 3.4) were selected across the slope covering all the landform units in the microwatershed (Natarajan and Dipak Sarkar, 2010).




Fig: 3.4. Location of profiles in a transect

In the selected transect, soil profiles (Fig.3.4) were located at closely spaced intervals to take care of any change in the land features like break in slope, erosion, gravel, stones etc. In the selected sites, profiles (vertical cut showing the soil layers from surface to the rock) were opened up to 200 cm or to the depth limited by rock or hard substratum and studied in detail for all their morphological and physical characteristics. The soil and site characteristics were recorded for all profile sites on a standard proforma as per the guidelines given in USDA Soil Survey Manual (Soil Survey Staff, 2012). Apart from the transect study, profiles were also studied at random, almost like in a grid pattern, outside the transect areas to validate the soil map unit boundaries.

Based on the soil characteristics, the soils were grouped into different soil series. Soil series is the most homogeneous unit having similar horizons and properties and behaves similarly for a given level of management. Soil depth, texture, colour, kind of horizon and horizon sequence, amount and nature of gravel present, calcareousness, nature of substratum etc, were used as the major differentiating characteristics for identifying soil series occurring in the area. The differentiating characteristics used for identifying the soil series are given in Table 3.1. Based on the above characteristics, 8 soil series were identified in Bettageri-1 Microwatershed.

**Table 3.1 Differentiating Characteristics used for identifying Soil Series
(Characteristics are of Series Control Section)**

Sl. No	Soil Series	Depth (cm)	Colour (moist)	Texture	Gravel (%)	Horizon sequence	Calcareousness
Soils of Granite and granite gneiss Landscape							
1	Harve (HRV)	25-50	2.5YR3/4,3/6 5YR3/3,4/4,3/4	gscl	>35	Ap-Bt-Cr-	-
2	Mukhadahalli (MKH)	50-75	5YR3/3,3/4,4/3, 5/4,6/6 2.5YR3/4	gscl	>35	Ap-Bt-Cr	-
3	Tigari (TGR)	75-100	5YR 3/3, 4/3, 2.5YR 2/3, 3/3, 3/4	gscl-gsc	15-35	Ap-Bt-Cr	e-es
4	Chikkamegheri (CKM)	75-100	2.5YR2.5/3,3/4, 3/6	sc	-	Ap-Bt-Cr	-
5	Bidanagere (BDG)	75-100	5YR3/3,3/4,4/3,5/ 4 2.5YR3/4	gc	35-60	Ap-Bt-Cr	-
Soils of Alluvial Landscape							
6	Muttal (MTL)	25-50	10YR3/2,3/3,4/2 7.5YR3/2,3/3,6/4	gc	15-35	Ap-Bw-Ck	e-ev
7	Ravanaki (RNK)	50-75	7.5YR3/2,3/3,5/2, 5/3 10YR3/1,3/2,4/1, 4/2, 5/1,6/1	c	<15	Ap-Bw-C	e-ev
8	Gatareddihal (GRH)	100-150	10YR 2/1, 3/1, 2.5Y 4/3, 5/4	c	<15	Ap-Bw-BC-C	es

3.4 Soil Mapping

The area under each soil series was further separated into soil phases and their boundaries delineated on the cadastral map based on the variations observed in the texture of the surface soil, slope, erosion, presence of gravel, stones etc. A soil phase is a subdivision of soil series based mostly on surface features that affect its use and management. The soil mapping units are shown on the map (Fig.3.5) in the form of symbols. During the survey many soil profile pits, few minipits and a few auger bores representing different landforms occurring in the microwatershed were studied. In addition to the profile study, spot observations in the form of minipits, road cuts, terrace cuts etc., were studied to validate the soil boundaries on the soil map. The soil map shows the geographic distribution of 9 mapping units representing 8 soil series occurring in the microwatershed. The soil map unit (soil legend) description is presented in Table 3.2. The soil phase map (management units) shows the distribution of 9 phases mapped in the microwatershed. Each mapping unit (soil phase) delineated on the map has similar soil and site characteristics. In other words, all the farms or survey numbers included in one phase will have similar management needs and have to be treated accordingly.

3.5 Laboratory Characterization

Soil samples for each series were collected from representative master profiles for laboratory characterization by following the methods outlined in the Laboratory Manual (Sarma *et al*, 1987). Surface soil samples collected in the year 2017 from Bettageri-1 farmer's fields (38 samples) for fertility status (major and micronutrients) at 250 m grid interval were analyzed in the laboratory (Katyal and Rattan, 2003). By linking the soil fertility data to the survey numbers through GIS, soil fertility maps were generated using Kriging method for the microwatershed.

3.6 Land Use Classes (LUCs)

The 9 soil phases identified and mapped in the microwatershed were regrouped into 7 Land Use Classes (LUC's) for the purpose of preparing a Proposed Crop Plan for sustained development of the microwatershed. The database (soil phases) generated under LRI was utilized for identifying Land Use Classes (LUC's) based on the management needs. One or more than one soil site characteristic having influence on the management have been chosen for identification and delineation of LUCs. For Bettageri-1 Microwatershed, five soil and site characteristics, namely soil depth, soil texture, slope, erosion and gravel content have been considered for defining LUCs. The land use classes are expected to behave similarly for a given level of management.




Fig 3.5 Soil Phase or Management Units- Bettageri-1 Microwatershed

Table 3.2 Soil map unit description of Bettageri-1 Microwatershed

Soil map unit No*	Soil Series	Soil Phase Symbol	Mapping Unit Description	Area in ha (%)
Soils of granite and granite gneiss landscape				
	HRV		Harve soils are shallow (25-50 cm), well drained, dark red to dark reddish brown, red gravelly loamy soils occurring on nearly level to gently sloping uplands under cultivation	19 (4.88)
24		HRVhB1g2	Sandy clay loam surface, slope 1-3%, slight erosion, very gravelly (35-60%)	19 (4.88)
27		HRVhB2g2	Sandy clay loam surface, slope 1-3%, moderate erosion, very gravelly (35-60%)	0.14 (0.04)
	MKH		Mukhadahalli soils are moderately shallow (50-75 cm), well drained, have dark brown to reddish brown gravelly red sandy clay loam soils occurring on gently very gently to gently sloping uplands under cultivation	41 (10.57)
91		MKHhB2g2	Sandy clay surface, slope 1-3%, moderate erosion, very gravelly (35-60%)	41 (10.57)
	TGR		Tigari soils are moderately deep (75-100 cm), well drained, have reddish brown to dark reddish brown red calcareous gravelly sandy clay loam to sandy clay soils occurring on very gently sloping uplands under cultivation	13 (3.22)
151		TGRiB2g1	Sandy clay surface, slope 1-3%, moderate erosion, gravelly (15-35%)	13 (3.22)
	CKM		Chikkamegheri soils are moderately deep (75-100 cm), well drained, have dark brown to dark reddish brown red sandy clay soils occurring on nearly level to very gently sloping uplands under cultivation	88 (22.53)
174		CKMhB1g1	Sandy clay loam surface, slope 1-3%, slight erosion, gravelly (15-35%)	88 (22.53)
	BDG		Bidanagere soils are moderately deep (75-100 cm), well drained, have dark reddish brown gravelly red sandy clay loam to sandy clay soils occurring on nearly level to gently sloping uplands under cultivation	72 (18.34)
188		BDGhB2g1	Sandy clay loam surface, slope 1-3%, moderate erosion, gravelly (15-35%)	72 (18.34)
Soils of alluvial landscape				
	MTL		Muttal soils are shallow (25-50 cm), well drained, have very dark grayish brown to dark brown, calcareous black gravelly clay soils occurring on gently to very gently sloping uplands under cultivation	101 (25.83)
305		MTLiB2g2	Sandy clay surface, slope 1-3%, moderate erosion, very gravelly (35-60%)	101 (25.83)
	RNK		Ravanaki soils are moderately shallow (50-75 cm), moderately well drained, have dark brown to very dark grayish brown and dark gray, calcareous clayey black soils occurring on very gently sloping uplands under cultivation	1 (0.35)
337		RNKmB2g1	Clay surface, slope 1-3%, moderate erosion, gravelly (15-35%)	1 (0.35)
	GRH		Gatareddihal soils are deep (100-150 cm), moderately well drained, have light olive brown to very dark gray, calcareous black cracking clay soils occurring on nearly level to very gently sloping plains under cultivation	20 (5.09)
371		GRHmB1	Clay surface, slope 1-3%, slight erosion	20 (5.09)
1000		Others	Habitation	36 (9.16)

*Soil map unit numbers are continuous for the taluk, not for the microwatersheds

THE SOILS

Detailed information pertaining to the nature, extent and distribution of different kinds of soils occurring in Bettageri-1 Microwatershed is provided in this chapter. The microwatershed area has been identified as granite gneiss and alluvial landscapes based on geology. In all, 8 soil series are identified. Soil formation is the result of the combined effect of environmental and terrain factors that are reflected in soil morphology. The soil formation is dominantly influenced by the parent material, climate, time and relief.

A brief description of each of the 8 soil series identified followed by 9 soil phases (management units) mapped (Fig. 3.5) are furnished below. The physical and chemical characteristics of soil series identified in Bettageri-1 microwatershed is given in Table 4.1. The soils in any one map unit differ from place to place in their depth, texture, slope, gravelliness, erosion or any other site characteristic that affect management. The soil phase map can be used for identifying the suitability of areas for growing specific crops or for other alternative uses and also for deciding the type of conservation structures needed. The detailed information on soil and site-characteristics like soil depth, surface soil texture, slope, erosion, gravelliness, AWC, LCC etc, with respect to each of the soil phase identified is given village/survey number wise for the microwatershed in Appendix-I.

4.1 Soils of Granite and Granite gneiss landscape

In this landscape, 5 soil series are identified and mapped. Of these, Chikkamegheri (CKM) occupies maximum area of about 88 ha (23%), Bidanagere (BDG) 72 ha (18%), and other occur in a small area. The brief description of each soil series along with the soil phases identified and mapped is given below.

4.1.1 Harve (HRV) Series: Harve soils are shallow (25-50 cm), well drained, have reddish brown to dark red gravelly sandy clay loam soils. They have developed from granite gneiss and occur on very gently to moderately sloping uplands. The Harve series has been tentatively classified as a member of the loamy- skeletal, mixed, isohyperthermic family of Typic Haplustalfs.

The thickness of the solum ranges from 28 to 48 cm. The thickness of A-horizon ranges from 12 to 17 cm. Its colour is in 5YR and 2.5 YR hue with value 3 to 4 and chroma 4 to 6. The texture varies from loamy sand to sandy loam with 20 to 60 per cent gravel. The thickness of B-horizon ranges from 16 to 32 cm. Its colour is in 2.5 YR and 5 YR hue with value 3 to 4 and chroma 4 to 6. Its texture is sandy clay loam with gravel content of more than 35 per cent. The available water capacity is very low (<50 mm/m). Only one soil phase is identified and mapped.



Landscape and soil profile characteristics of Harve (HRV) Series

4.1.2 Mukhadahalli (MKH) Series: Mukhadahalli soils are moderately shallow (50-75 cm), well drained, have dark brown to reddish brown gravelly sandy clay loam soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands. The Mukhadahalli series has been classified as a member of the clayey-skeletal, mixed, isohyperthermic family of Typic Haplustalfs.

The thickness of the solum ranges from 51 to 72 cm. The thickness of A horizon ranges from 12 to 17 cm. Its colour is in 5 YR and 7.5 YR hue with value 3 to 4 and chroma 2 to 4. The texture varies from loamy sand to sandy loam with 20 to 45 per cent gravel. The thickness of B horizon ranges from 40 to 68 cm. Its colour is in 2.5 YR and 5 YR hue with value and chroma 3 to 6. Texture is sandy clay loam with 35 to 50 per cent gravel. The available water capacity is low (51-100 mm/m). Only one soil phase identified and mapped.



Landscape and soil profile characteristics of Mukhadahalli (MKH) Series

4.1.3 Tigari (TGR) Series: Tigari soils are moderately deep (75-100 cm), well drained, have dark reddish brown to reddish brown, calcareous gravelly sandy clay loam to sandy clay soils. They have developed from weathered granite gneiss and occur on very gently sloping uplands under cultivation.

The thickness of the solum ranges from 77 to 100 cm. The thickness of A horizon ranges from 11 to 21 cm. Its colour is in 5 YR and 10 YR hue with value 3 to 4 and chroma 2 to 4. The texture is sandy clay to clay. The thickness of B horizon ranges from 56 to 87 cm. Its colour is in 2.5 YR and 5 YR hue with value and chroma 2 to 4. Its texture ranges from gravelly sandy clay loam to sandy clay and is calcareous. The available water capacity is low (51-100 mm/m).

4.1.4 Chikkamegheri (CKM) Series: Chikkamegheri soils are moderately deep (75-100 cm), well drained, have dark brown to dark reddish brown and red sandy clay soils. They have developed from granite gneiss and occur on nearly level to very gently sloping uplands. The Chikkamegheri series has been classified as a member of the fine, mixed, isohyperthermic family of Rhodic Paleustalfs.

The thickness of the solum ranges from 76 to 100 cm. The thickness of A horizon ranges from 11 to 24 cm. Its colour is in 7.5 YR, 5YR and 2.5 YR hue with value 2 to 4 and chroma 3 to 6. The texture varies from sandy clay loam to sandy clay with 10 to 15 per cent gravel. The thickness of B horizon ranges from 65 to 86 cm. Its colour is in 2.5 YR hue with value 2.5 to 3 and chroma 3 to 6. Its texture is dominantly sandy clay. The available water capacity is medium (101-150 mm/m). Only one soil phase is identified and mapped



Landscape and soil profile characteristics of Chikkamegheri (CKM) Series

4.1.5 Bidanagere (BDG) Series: Bidanagere soils are moderately deep (75-100 cm), well drained, have dark reddish brown gravelly clay soils. They have developed from weathered granite gneiss and occur on very gently sloping uplands under cultivation. The

Bidanagere soil series has been classified as a member of the clayey-skeletal, mixed, isohyperthermic family of Rhodic, Paleustalfs.

The thickness of the solum ranges from 78 to 99 cm. The thickness of A-horizon ranges from 12 to 19 cm. Its colour is in 2.5 YR and 5 YR hue with value 2 to 3 and chroma 3 to 4. The texture varies from sandy clay loam to sandy clay with 10 to 20 per cent gravel. The thickness of B-horizon ranges from 68 to 85 cm. Its colour is in 5 YR and 2.5 YR hue with value 3 to 5 and chroma 3 to 4. Its texture is gravelly clay with gravel content of 35-60 per cent. The available water capacity is very low (<50 mm/m). only one soil phase is identified and mapped.



Landscape Soil Profile Characteristics of Bidanagere (BDG) Series

4.2 Soils of Alluvial Landscape

In this landscape, 3 soil series are identified and mapped. Of these, Muttal (MTL) occupies maximum area of about 101 ha (26%), Gatareddihal (GRH) 20 ha (5%) and others occur in small area. The brief description of each soil series along with the soil phases identified and mapped is given below.

4.2.1 Muttal (MTL) Series: Muttal soils are shallow (25-50 cm), well drained, have dark brown to very dark grayish brown, calcareous gravelly clay soils. They have developed from alluvium and occur on nearly level to very gently sloping uplands. The Muttal series has been classified as a member of the clayey, mixed, isohyperthermic (Calc) family of (Paralithic) Haplustepts.

The thickness of the solum ranges from 30 to 48 cm. The thickness of A horizon ranges from 15 to 18 cm. Its colour is in 7.5 YR and 10 YR hue with value 2 to 3 and chroma 2.5 to 4. The texture varies from sandy clay to clay with 10 to 15 per cent gravel. The thickness of B horizon ranges from 18 to 32 cm. Its colour is in 10 YR and 7.5 YR hue with value 2 to 6 and chroma 2 to 4. Its texture is sandy clay to clay. The available water capacity is low (51-100 mm/m). Only one soil phase was identified and mapped.



Landscape and soil profile characteristics of Muttal (MTL) Series

4.2.2 Ravanaki (RNK) Series: Ravanaki soils are moderately shallow (50-75 cm), well drained, have dark brown to very dark grayish brown, calcareous clayey soils. They have developed from alluvium and occur on nearly level to very gently sloping uplands. The Ravanaki series has been classified as a member of the very fine, smectitic, isohyperthermic (calc) family of Typic Haplustepts.


The thickness of the solum ranges from 50 to 75 cm. The thickness of A horizon ranges from 15 to 20 cm. Its colour is in 7.5 YR and 10 YR hue with value 2 to 3 and chroma 2.5 to 4. The texture varies from sandy clay to clay with 10 to 15 per cent gravel. The thickness of B horizon ranges from 35 to 60 cm. Its colour is in 10 YR and 7.5 YR hue with value 2 to 6 and chroma 2 to 4. Its texture is sandy clay to clay and are calcareous with gravel content of 10 to 20 per cent. The available water capacity is low (51-100 mm/m). one soil phases were identified and mapped.



Landscape and Soil Profile Characteristics of Ravanaki (RNK) Series

4.2.3 Gatareddihal (GRH) Series: Gatareddihal soils are deep (100-150 cm), moderately well drained have black or dark grey to light olive brown calcareous clay soils. They are developed from Alluvium and occur on nearly level to very gently sloping uplands under cultivation. The Gatareddihal series has been classified as a member of the fine, smectitic, isohyperthermic (calc) family of Vertic Haplustepts.

The thickness of the solum ranges from 102 to 149 cm. The thickness of A-horizon ranges from 12 to 19 cm. Its colour is in 7.5 YR, 10 YR hue with value 3 to 4 and chroma 1 to 6. The texture is sandy clay loam to clay. The thickness of B-horizon ranges from 86 to 117 cm. Its colour is in 10 YR and 7.5 YR hue with value 3 and chroma 2 to 6. Texture is clay with less than 15 per cent gravel and are calcareous. The available water capacity is very high (>200 mm/m). Only one soil phase was identified and mapped.



Landscape and soil profile characteristics of Gatareddihal (GRH) Series

Table: 4.1 Physical and Chemical Characteristics of Soil Series identified in Bettageri-1 Microwatershed

Series Name: Mukhadahalli (MKH), Pedon: R-11

Location: 15°22'05.4"N, 76°04'10.3"E, Halageri village, Koppal taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bangalore. **Classification:** Clayey-skeletal, mixed, isohyperthermic Typic Haplustalfs

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-19	Ap	65.71	8.83	25.46	9.27	9.06	14.42	21.52	11.43	70	scl	16.54	8.60
19-32	Bt	55.89	11.13	32.98	6.47	9.18	11.89	19.19	9.18	50	scl	19.24	12.78
32-58	Bt	47.95	10.41	41.63	17.52	3.78	9.13	9.55	7.97	50	sc	24.03	16.02

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO ₃	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP
	Water	CaCl ₂	M KCl				Ca	Mg	K	Na	Total				
				dS m ⁻¹	%	%	cmol kg ⁻¹							%	%
0-19	7.38	-	-	0.09	0.2	0.00	8.97	4.32	0.26	0.22	13.77	14.84	0.58	93	1.49
19-32	7.5	-	-	0.106	0.41	0.00	15.98	3.27	0.16	0.50	19.91	20.88	0.63	95	2.38
32-58	7.46	-	-	0.173	0.49	0.00	19.71	4.53	0.23	1.32	25.79	25.76	0.62	100	5.11

Contd...

Series Name: Chikkamegheri (CKM), Pedon: RM-2

Location: 15°21'40"N, 76°16'43"E, Gudannahalli village, Koppal taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bangalore. **Classification:** Fine, mixed, isohyperthermic, Rhodic Paleustalfs

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-10	Ap	66.80	5.51	27.69	10.14	10.04	20.29	14.75	11.58	-	scl	20.59	7.15
10-25	Bt1	39.52	7.17	53.32	8.75	9.59	7.27	8.43	5.48	-	c	26.96	13.99
25-38	Bt2	42.00	7.16	50.84	13.16	8.74	6.42	8.53	5.16	-	c	26.51	13.42
38-55	Bt3	41.77	10.31	47.92	15.19	8.54	6.33	7.38	4.32	10	c	25.28	14.10
55-70	Bt4	44.03	8.96	47.01	15.72	9.22	6.92	6.81	5.35	20	c	24.30	14.35
70-90	Bt5	56.02	8.46	35.52	11.41	17.07	12.36	10.26	4.92	25	sc	20.59	13.06

Depth (cm)	pH (1:2.5)			E.C. (1:2.5) dS m ⁻¹	O.C. %	CaCO ₃ %	Exchangeable bases					CEC	CEC/Clay	Base saturation %	ESP %
	Water	CaCl ₂	M KCl				Ca	Mg	K	Na	Total				
0-10	7.99	-	-	0.326	0.83	4.44	9.35	4.76	0.28	0.54	14.93	12.50	0.45	119	4.33
10-25	7.36	-	-	0.345	0.99	2.40	10.37	4.84	0.10	1.18	16.48	17.60	0.33	94	6.68
25-38	6.69	-	-	0.477	0.79	0.00	10.25	4.20	0.09	1.61	16.15	16.10	0.32	100	10.01
38-55	6.45	-	-	0.548	0.63	0.00	9.43	2.86	0.10	1.52	13.91	14.80	0.31	94	10.27
55-70	6.35	-	-	0.532	0.71	0.00	9.59	2.79	0.11	1.66	14.16	14.60	0.31	97	11.39
70-90	6.44	-	-	0.613	0.27	0.00	9.58	3.10	0.19	1.87	14.74	14.70	0.41	100	12.69

Contd...

Series: Bidanagere (BDG), **Pedon:** RM-3

Location: 13°22'11"N, 76°38'03"E, (4D3D8G1a), Tharabenahalli village, Chikkanayakanahalli taluk, Tumakuru district

Analysis at: NBSS&LUP, Regional Centre, Bengaluru **Classification:** Clayey-skeletal, mixed, isohyperthermic Rhodic Paleustalfs

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-20	Ap	81.19	11.25	7.56	12.54	15.07	17.90	21.94	13.75	50	ls	-	-
20-35	Bt1	57.45	11.45	31.10	12.76	11.02	10.92	12.45	10.31	50	scl	-	-
35-92	Bt2	44.63	7.85	47.52	12.40	9.61	8.37	7.75	6.51	60	c	-	-

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO ₃	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP			
	Water	CaCl ₂	M KCl				dS m ⁻¹	%	%	Ca	Mg					K	Na	Total
										cmol kg ⁻¹								
0-20	6.24	-	-	0.06	0.60	0.00	1.61	0.26	0.10	0.01	1.98	3.76	0.50	52.56	0.35			
20-35	5.99	-	-	0.02	0.40	0.00	4.25	0.46	0.08	0.28	5.07	8.02	0.26	63.18	3.46			
35-92	6.70	-	-	0.03	0.20	0.00	5.45	0.31	0.10	0.22	6.09	9.90	0.21	61.48	2.24			

Contd...

Series Name: Muttal (MTL), **Pedon:** RM-13

Location: 15°14'30.8"N, 75°56'50.6"E, Gatareddihal village, Koppal taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bangalore. **Classification:** Clayey, mixed, isohyperthermic (calc) (Paralithic) Haplustepts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-20	Ap	39.05	13.74	47.21	3.05	5.05	8.21	14.63	8.11	15-30	c	29.95	17.94
20-34	Bwk	28.77	19.57	51.66	4.81	4.71	4.92	9.09	5.24	10	c	33.44	21.56

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO ₃	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP
	Water	CaCl ₂	M KCl				Ca	Mg	K	Na	Total				
	Water	CaCl ₂	M KCl	dS m ⁻¹	%	%	cmol kg ⁻¹							%	%
0-20	8.27	-	-	0.202	0.79	6.10	-	-	0.62	0.25	-	36.64	0.78	-	0.69
20-34	8.36	-	-	0.177	0.99	23.04	-	-	0.29	0.38	-	39.60	0.77	-	0.96

Contd...

Series Name: Ravanaki (RNK), **Pedon:** RM-20

Location: 15°14'22.7"N, 75°57'45.8"E, Gatareddihal village, Koppal taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bangalore. **Classification:** Very fine, smectitic, isohyperthermic (calc) Typic Haplustepts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-28	Ap	24.43	17.76	57.81	5.30	3.89	3.78	7.14	4.32	20	c	41.40	29.60
28-55	Bw	18.77	15.59	65.64	2.74	3.73	2.85	4.83	4.61	10	c	46.71	35.18
55-80	Bc	12.53	15.43	72.04	2.60	1.92	1.47	3.16	3.39	10	c	56.82	43.73

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO ₃	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP
	Water	CaCl ₂	M KCl				Ca	Mg	K	Na	Total				
	cmol kg ⁻¹														
0-28	8.86	-	-	0.483	0.63	15.48	-	-	0.86	6.27		37.00	0.64	-	16.94
28-55	8.61	-	-	1.4	0.23	13.68	-	-	0.68	12.27		53.20	0.81	-	23.06
55-80	8.35	-	-	4.53	0.91	11.40	-	-	0.75	28.97		54.80	0.76	-	52.86

INTERPRETATION FOR LAND RESOURCE MANAGEMENT

The most important soil and site characteristics that affect the land use and conservation needs of an area are land capability, land irrigability, soil depth, soil texture, coarse fragments, available water capacity, soil slope, soil erosion, soil reaction etc. These are interpreted from the data base generated through land resource inventory and several thematic maps are generated. These would help in identifying the areas suitable for growing crops and, soil and water conservation measures and structures needed thus helping to maintain good soil health for sustained crop production. The various thematic maps generated are described below.

5.1 Land Capability Classification

Land capability classification is an interpretative grouping of soil map units (soil phases) mainly based on inherent soil characteristics, external land features and environmental factors that limit the use of land for agriculture, pasture, forestry or other uses on a sustained basis (IARI, 1971). The land and soil characteristics used to group the land resources in an area into various land capability classes, subclasses and units are

Soil characteristics: Soil depth, soil texture, coarse fragments, soil reaction, available water capacity, calcareousness, salinity/alkali *etc.*

Land characteristics: Slope, erosion, drainage, rock outcrops.

Climate: Total rainfall and its distribution, and length of crop growing period.

The Land Capability Classification system is divided into land capability classes, subclasses and units based on the level of information available. Eight land capability classes are recognized. They are

Class I: They are very good lands that have no limitations or very few limitations that restrict their use.

Class II: They are good lands that have minor limitations and require moderate conservation practices.

Class III: They are moderately good lands that have severe limitations that reduce the choice of crops or that require special conservation practices.

Class IV: They are fairly good lands that have very severe limitations that reduce the choice of crops or that require very careful management.

Class V: Soils in these lands are not likely to erode, but have other limitations like wetness that are impractical to remove and as such not suitable for agriculture, but suitable for pasture or forestry with minor limitations.

Class VI: The lands have severe limitations that make them generally unsuitable for cultivation, but suitable for pasture or forestry with moderate limitations.

Class VII: The lands have very severe limitations that make them unsuitable for cultivation, but suitable for pasture or forestry with major limitations.

Class VIII: Soil and other miscellaneous areas (rock lands) that have very severe limitations that nearly preclude their use for any crop production, but suitable for wildlife, recreation and installation of wind mills.

The land capability subclasses are recognised based on the dominant limitations observed within a given land capability class. The subclasses are designated by adding a lower case letter like ‘e’, ‘w’, ‘s’, or ‘c’ to the class numeral. The subclass “e” indicates that the main hazard is risk of erosion, “w” indicates drainage or wetness as a limitation for plant growth, “s” indicates shallow soil depth, coarse or heavy textures, calcareousness, salinity/alkalinity or gravelliness and “c” indicates limitation due to climate.




Fig. 5.1 Land Capability map of Bettageri-1 Microwatershed

The land capability subclasses have been further subdivided into land capability units based on the kinds of limitations present in each subclass. Ten land capability units are identified in grouping the soil map units. They are stony or rocky (0), erosion hazard (slope, erosion) (1), coarse texture (sand, loamy sand, sandy loam) (2), fine texture (cracking clay, silty clay) (3), slowly permeable subsoil (4), coarse underlying material (5), salinity/alkali (6), stagnation, overflow, high ground water table (7), soil depth (8) and fertility problems (9). The capability units thus identified have similar soil and land characteristics that respond similarly to a given level of management. The soils of the microwatershed have been classified upto land capability subclass level.

The 9 soil map units identified in the Bettageri-1 microwatershed are grouped under two land capability classes and four land capability subclasses (Fig. 5.1).

Entire area is suitable for agriculture. Good lands (Class II) cover an area of 31 per cent and are distributed in the northern and northeastern part of the microwatershed with moderate problems of soil and erosion. Moderately good lands (Class III) cover a major area of about 60 per cent and are distributed in all parts of the microwatershed with moderate problems of soil and erosion.

5.2 Soil Depth

Soil depth refers to the depth of the soil occurring above the parent material or hard rock. The depth of the soil determines the effective rooting depth for plants and in accordance with soil texture, mineralogy and gravel content, the capacity of the soil column to hold water and nutrient availability. Soil depth is one of the most important soil characteristic that is used in differentiating soils into different soil series. The soil depth classes used in identifying soils in the field are very shallow (<25 cm), shallow (25-50 cm), moderately shallow (50-75 cm), moderately deep (75-100 cm), deep (100-150 cm) and very deep (>150 cm). They were used to classify the soils into different depth classes and a soil depth map was generated (Fig. 5.2).

An area of 120 ha (31%) has shallow soils (25-50 cm) and are distributed in the northwestern and southern part of the microwatershed. An area of about 43 ha (11%) is moderately shallow (50-75 cm) and are distributed in the northern and southwestern part of the microwatershed. Moderately deep (75-100 cm) soils occupy a major area of about 172 ha (44%) and occur in all parts of the microwatershed. Deep soils (100-150 cm) cover about 20 ha (5%) in the microwatershed.




Fig. 5.2 Soil Depth map of Bettageri-1 Microwatershed

The problem lands with an area of about 120 ha (31%) having shallow rooting depth are marginally suitable for growing agricultural crops but well suited for pasture, forestry or other recreational purposes. Occasionally, short duration crops may be grown if rainfall is normal. The most productive lands cover about 20 ha (5%) where all climatically adapted long duration crops be grown.

5.3 Surface Soil Texture

Texture is an expression to indicate the coarseness or fineness of the soil as determined by the relative proportion of primary particles of sand, silt and clay. It has a direct bearing on the structure, porosity, adhesion and consistence. The surface layer of a soil to a depth of about 25 cm is the layer that is most used by crops and plants. The surface soil textural class provides a guide to understanding soil-water retention and availability, nutrient holding capacity, infiltration, workability, drainage, physical and chemical behaviour, microbial activity and crop suitability.

An area of about 176 ha (45%) has soils that are clayey texture at the surface (Fig. 5.3) and Major area of about 179 ha (46%) has soils that are loamy at the surface. Both are most productive lands that have high potential for soil-water retention and availability, and nutrient retention and availability, but have problems of drainage, infiltration, workability and other physical problems




Fig. 5.3 Surface Soil Texture map of Bettageri-1 Microwatershed

5.4 Soil Gravelliness

Gravel is the term used for describing coarse fragments between 2 mm and 7.5 cm diameter and stones for those between 7.5 cm and 25 cm. The presence of gravel and stones in soil reduces the volume of soil responsible for moisture and nutrient storage, drainage, infiltration and runoff, and hinders plant growth by impeding root growth and seedling emergence, intercultural operations and farm mechanization.

The soils that are non-gravelly (<15% gravel) cover a small area of about 20 ha (5%) and are distributed in the northern part of the microwatershed, Major area of 174 ha (44%) is covered by gravelly (15-35% gravel) soils and are distributed in the central part of the microwatershed. A small area of about 161 ha (41%) has soils that are very gravelly (35-60% gravel) and are distributed in the southern and northern part of the microwatershed (Fig. 5.4).

The most productive lands with respect to gravelliness are found to be 5%. They are non-gravelly with less than 15 per cent gravel and have potential for growing both annual and perennial crops. The problem soils (41%) that are very gravelly (35-60%) where only short duration crops can be grown.




Fig. 5.4 Soil Gravelliness map of Bettageri-1 Microwatershed

5.5 Available Water Capacity

The soil available water capacity (AWC) is estimated based on the ability of the soil column to retain water between the tensions of 0.33 and 15 bars in a depth of 100 cm or the entire solum if the soil is shallower. The AWC of the soils (soil series) as estimated

by considering the soil texture, mineralogy, soil depth and gravel content (Sehgal *et al.*, 1990) and accordingly the soil map units were grouped into five AWC classes *viz*, very low (<50 mm/m), low (50-100 mm/m), medium (100-150 mm/m), high (150-200 mm/m) and very high (>200 mm/m) and using these values, an AWC map was generated (Fig. 5.5).

Major area of about 132 ha (34%) has soils that are very low (<50 mm) in available water capacity. An area of about 115 ha (29%) has soils that are low (51-100 mm/m) in available water capacity and are distributed in the southwestern and northern part of the microwatershed. An area of about 88 ha (23%) is medium (101-150 mm/m) in available water capacity and are distributed in the eastern part of the microwatershed and small area of about 20 ha (5%) is very high in available water capacity and are distributed in the northern part of the microwatershed.

An area of about 132 ha (34%) in the microwatershed has soils that are problematic with regard to available water capacity. Here, only short duration crops can be grown and the probability of crop failure is very high. These areas are best put to other alternative uses. An area of about 20 ha (5%) has soils that have high potential (>200 mm/m) with regard to available water capacity where all climatically adapted long duration crops can be grown successfully.




Fig. 5.5 Soil Available Water Capacity map of Bettageri-1 Microwatershed

5.6 Soil Slope

Soil slope refers to the inclination of the surface of the land. It is defined by gradient, shape and length, and is an integral feature of any soil as a natural body. Slope is considered important in soil genesis, land use and land development. The length and gradient of slope influences the rate of runoff, infiltration, erosion and deposition. The soil map units were grouped into four slope classes and a slope map was generated showing the area extent and their geographic distribution of different slope classes in the microwatershed (Fig. 5.6).

Entire area falls under very gently sloping (1-3% slope) lands. In all these areas, all climatically adapted annual and perennial crops can be grown without much soil and water conservation and other land development measures.




Fig. 5.6 Soil Slope map of Bettageri-1 Microwatershed

5.7 Soil Erosion

Soil erosion refers to the wearing away of the earth's surface by the forces of water, wind and ice involving detachment and transport of soil by raindrop impact. It is used for accelerated soil erosion resulting from disturbance of the natural landscape by burning, excessive grazing and indiscriminate felling of forest trees and tillage, all usually by man. The erosion classes showing an estimate of the current erosion status as judged from field observations in the form of rills, gullies or a carpet of gravel on the surface are recorded. Four erosion classes, viz, slight erosion (e1), moderate erosion (e2), severe erosion (e3) and very severe erosion (e4) are recognized. The soil map units were

grouped into different erosion classes and a soil erosion map generated. The area extent and their spatial distribution in the microwatershed is given in Figure 5.7.

Major area of 228 ha (58%) has soils that are moderately eroded (e2 class). These are problematic and need appropriate soil and water conservation and other land development measures and an area of about 127 ha (33%) has soils that are slightly eroded (e1 class) and are distributed in the northern and eastern part of the microwatershed.




Fig. 5.7 Soil Erosion map of Bettageri-1 Microwatershed

FERTILITY STATUS

Soil fertility plays an important role in increasing crop yield. The adoption of high yielding varieties that require high amounts of nutrients has resulted in deficiency symptoms in crops and plants due to imbalanced fertilization and poor inherent fertility status, as these areas are characterised by low rainfall and high temperatures. Hence, it is necessary to know the fertility (macro and micro nutrients) status of the soils of the watersheds for assessing the kind and amount of fertilizers required for each of the crop intended to be grown. For this purpose, the surface soil samples collected from the grid points (one soil sample at every 250 m grid interval) all over the microwatershed through land resource inventory in the year 2017 were analysed for pH, EC, organic carbon, available phosphorus and potassium, and for micronutrients like zinc, boron, copper, iron and manganese, and secondary nutrient sulphur.

Soil fertility data generated has been assessed and individual maps for all the nutrients for the microwatershed have been generated by using the Kriging method under GIS. The village/survey number wise fertility data for the microwatershed is given in Appendix-II.

6.1 Soil Reaction (pH)

The soil analysis of the Bettageri-1 microwatershed for soil reaction (pH) showed that a small area of 23 ha (6%) is slightly acid (pH 6.0-6.5) and is distributed in the northwestern part of the microwatershed. 70 ha (18%) area is neutral (pH 6.5-7.3) and occur in the northern and western part. Major area of about 202 ha (52%) soils are slightly to moderately alkaline (pH 7.3-8.4) occur in all parts and 59 ha (15%) area is under strongly alkaline to very strongly alkaline (pH 8.4- >9.0) and is distributed in the northern and southern part of the microwatershed. Major area of about 354 ha (64%) is very strongly alkaline (pH > 9.0) and are distributed in the southern and northern part of the microwatershed (Fig.6.1).

6.2 Electrical Conductivity (EC)

The Electrical Conductivity of the soils of the entire microwatershed area is <2 dSm⁻¹ (Fig 6.2) and as such the soils are non-saline.

6.3 Organic Carbon

The soil organic carbon content (an index of available Nitrogen) of the microwatershed is low (<0.5%) in an area of 79 ha (20%) and is distributed in the northern part of the microwatershed and medium (0.5-0.75%) in the major area of about 152 ha (39%) and is distributed in the central and southern part of the microwatershed. An area of about 124 ha (32%) is high (>0.75%) in organic carbon content and occur in the southern part of the microwatershed (Fig.6.3).

6.4 Available Phosphorus

Small area of about 24 ha (6%) is low (<23 kg/ha) in available phosphorus and distributed in the northern part of the microwatershed. Medium (23-57 kg/ha) in major area of about 277 ha (71%) and distributed in all parts of the microwatershed. There is an urgent need to increase the dose of phosphorous for all the crops by 25 per cent over the recommended dose to realize better crop performance. High (>57 kg/ha) in about 54 ha (14%) area and occur in the central and western part of the microwatershed (Fig 6.4).




Fig.6.1 Soil Reaction (pH) map of Bettageri-1 Microwatershed




Fig.6.2 Electrical Conductivity (EC) map of Bettageri-1 Microwatershed




Fig.6.3 Soil Organic Carbon map of Bettageri-1 Microwatershed




Fig.6.4 Soil Available Phosphorus map of Bettageri-1 Microwatershed

6.5 Available Potassium

Small area of about 18 ha (5%) is high (>337 kg/ha) in available potassium (Fig.6.5). Hence, in these plots, for all the crops, 25% less potassium than recommended may be applied. Medium (145-337 kg/ha) in major area of about 337 ha (86%) and occur in the western part of the microwatershed. Additional 25% potassium may be applied in areas where potassium is medium.

6.6 Available Sulphur

Major area of 259 ha (66%) is low (<10 ppm) in available sulphur and is distributed in all parts of the microwatershed. An area of about 96 ha (24%) is medium (10-20 ppm) in available sulphur and is distributed in the northern and southern part of the microwatershed. A very minor area of about <1 ha (0.17%) is high (>20 ppm) in available sulphur and are distributed in the northern part of the microwatershed (Fig.6.6). The areas that are low and medium in available sulphur need to be applied with magnesium sulphate or gypsum or factomphos (p) fertilizer (13% sulphur) for 2-3 years for the deficiency to be corrected.

6.7 Available Boron

Available boron content is low (<0.5 ppm) in maximum area of 233 ha (60%) in the microwatershed and is distributed in all parts of the microwatershed. An area of about 121 ha (31%) is medium (0.5-1.0 ppm) in available boron and is distributed in the central,

northern and southern part of the microwatershed (Fig.6.7). These areas need to be applied with sodium borate @ 10kg/ha as a soil application or 0.2% borax as foliar spray to correct the deficiency

6.8 Available Iron

Available iron content is deficient (<4.5 ppm) in major area of 200 ha (51%) and distributed in the southern, central and northern part of the microwatershed and sufficient (>4.5 ppm) in about 155 ha (40%) area and occur in the northern, central and southern part of the microwatershed (Fig 6.8).

6.9 Available Manganese

Available manganese content is sufficient (>1.0 ppm) in the entire microwatershed area (Fig 6.9).

6.10 Available Copper

Available copper content is sufficient (>0.2 ppm) in the entire microwatershed area (Fig 6.10).

6.11 Available Zinc

Available zinc content is deficient (<0.6 ppm) in an area 139 ha (36%) and a major area of 215 ha (55%) is sufficient (>0.6 ppm) in available zinc in the microwatershed Fig 6.11).




Fig.6.5 Soil Available Potassium map of Bettageri-1 Microwatershed




Fig.6.6 Soil Available Sulphur map of Bettageri-1 Microwatershed




Fig.6.7 Soil Available Boron map of Bettageri-1 Microwatershed




Fig.6.8 Soil Available Iron map of Bettageri-1 Microwatershed




Fig.6.9 Soil Available Manganese map of Bettageri-1 Microwatershed




Fig.6.10 Soil Available Copper map of Bettageri-1 Microwatershed




Fig.6.11 Soil Available Zinc map of Bettageri-1 Microwatershed

LAND SUITABILITY FOR MAJOR CROPS

The soil and land resource units (soil phases) of Bettageri-1 Microwatershed were assessed for their suitability for growing food, fodder, fibre and other horticulture crops by following the procedure as outlined in FAO, 1976 and 1983. Crop requirements were developed for each of the crop from the available research data and also by referring to Naidu *et. al.* (2006) and Natarajan *et. al* (2015). The crop requirements were matched with the soil and land characteristics (Table 7.1) to arrive at the crop suitability. In FAO land suitability classification, two orders are recognized. Order S- Suitable and Order N- Not suitable. The orders have Classes, subclasses and units. Order-S has three classes, Class S1- Highly Suitable, Class S2- Moderately Suitable and Class S3- Marginally Suitable. Order N has two classes, N1- Currently not Suitable and N2- Permanently not Suitable. There are no subclasses within the Class S1 as they will have very minor or no limitations for crop growth. Classes S2, S3, N1 and N2 are divided into subclasses based on the kinds of limitations encountered. The limitations that affect crop production are ‘c’ for erratic rainfall and its distribution and length of growing period (LGP), ‘e’ for erosion hazard, ‘r’ for rooting condition, ‘t’ for lighter or heavy texture, ‘g’ for gravelliness or stoniness, ‘n’ for nutrient availability, ‘l’ for topography, ‘m’ for moisture availability, ‘z’ for calcareousness and ‘w’ for drainage. These limitations are indicated as lower case letters to the class symbol. For example, moderately suitable lands with the limitations of soil depth and erosion are designated as S2re. For the microwatershed, the soil mapping units were evaluated and classified up to subclass level.

Using the above criteria, the soil map units of the microwatershed were evaluated and land suitability maps for 25 major annual and perennial crops were generated. The detailed information on the kind of suitability of each of the soil phase for the crops assessed are given village/ survey number wise for the microwatershed in Appendix-III.

7.1 Land Suitability for Sorghum (*Sorghum bicolor*)

Sorghum is one of the major crops grown in Karnataka in an area of 10.47 lakh ha in Bijapur, Gulbarga, Raichur, Bidar, Belgaum, Dharwad, and Bellary, Chitradurga, Mysore and Chamarajnaragar districts. The crop requirements for growing sorghum (Table 7.2) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and land suitability map for growing sorghum was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure. 7.1.

Highly suitable (Class S1) land occupying a small area of about 20 ha (5%) for growing sorghum and occur in the northern part of the microwatershed. An area of about 143 ha (37%) is moderately suitable (Class S2) for growing sorghum and are distributed in the northern, eastern and southwestern part of the microwatershed.

Table 7.1 Soil-Site Characteristics of Bettageri-1 Microwatershed

Soil Map Units	Climate (P) (mm)	Growing period (Days)	Drainage Class	Soil depth (cm)	Soil texture		Gravelliness		AWC (mm/m)	Slope (%)	Erosion	pH	EC	ESP	CEC [Cmol (p ⁺)kg ⁻¹]	BS (%)
					Surf-ace	Sub-surface	Sur-face	Sub-surface								
HRVhB1g2	662	<90	WD	25-50	scl	scl	35-60	>35	<50	1-3	slight	-	-	-	-	-
HRVhB2g2	662	<90	WD	25-50	scl	scl	35-60	>35	<50	1-3	moderate	-	-	-	-	-
MKHiB2g2	662	<90	WD	50-75	sc	scl	35-60	>35	51-100	1-3	moderate	7.38	0.09	1.49	14.84	93
TGRiB2g1	662	<90	WD	75-100	sc	scl-sc	15-35	15-35	51-100	1-3	moderate	-	-	-	-	-
CKMhB1g1	662	<90	WD	75-100	scl	sc	15-35	-	101-150	1-3	slight	7.99	0.32	4.33	12.50	94
BDGhB2g1	662	<90	WD	75-100	scl	c	15-35	35-60	<50	1-3	moderate	6.24	0.06	3.76	52.56	0.35
MTLiB2g2	662	<90	WD	25-50	sc	c	35-60	15-35	51-100	1-3	moderate	8.27	0.20	0.69	36.64	-
RNKmB2g1	662	<90	MWD	50-75	c	c	15-35	<15	51-100	1-3	moderate	8.86	0.48	16.94	37	-
GRHmB1	662	<90	MWD	100-150	c	c	-	<15	>200	1-3	slight	8.27	1.11	11.72	31.60	-

*Symbols and abbreviations are according to Field Guide for LRI under Sujala-III

They have minor limitations of gravelliness, rooting depth and calcareousness. Major area of about 192 ha (52%) is marginally suitable (Class S3) for growing sorghum and occur in the western and southern part of the microwatershed with moderate limitations of gravelliness, calcareous and rooting depth.

Table 7.2 Crop suitability criteria for Sorghum

Crop requirement		Rating			
Soil –site characteristics	Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N)
Slope	%	2-3	3-8	8-15	>15
LGP	Days	120-150	120-90	<90	
Soil drainage	Class	Well to mod.Well drained	imperfect	Poorly/excessively	V.poorly
Soil reaction	pH	5.5-8.0	5.0-5.5,7.8-8.4	8.4-9.0	>9.0
Surface soil texture	Class	c, sicl, sc	l, scl, sil, sic, cl,	S1, ls	S, fragmental skeletal
Soil depth	Cm	100-75	50-75	25-50	<25
Gravel content	% vol.	<15	15-35	35-60	>60
Salinity (EC)	dSm ⁻¹	2-4	4-8	8-10	>10
Sodicity (ESP)	%	5-8	8-10	10-15	>15




Fig. 7.1 Land Suitability map of Sorghum

7.2 Land Suitability for Maize (*Zea mays*)

Maize is one of the most important food crop grown in an area of 13.37 lakh ha in almost all the districts of the State. The crop requirements for growing maize (Table 7.3) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing maize was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.2.

Table 7.3 Crop suitability criteria for Maize

Crop requirement		Rating			
Soil-site characteristics	Unit	Highly suitable(S1)	Moderately suitable(S2)	Marginally suitable(S3)	Not suitable (N)
Slope	%	<3	3.5	5-8	
LGP	Days	>100	100-80	60-80	
Soil drainage	Class	Well drained	Mod. to imperfectly	Poorly/excessively	V. poorly
Soil reaction	pH	5.5-7.5	7.6-8.5	8.6-9.0	
Surface soil texture	Class	l,cl, scl, sil,sc	c(s-s),c,sicl, sic	sl ,ls	s,fragmental
Soil depth	Cm	>75	50-75	25-50	<25
Gravel content	% vol.	<15	15-35	35-60	>60
Salinity (EC)	dSm ⁻¹	<1.0	1.0-2.0	2.0-4.0	
Sodicity (ESP)	%	<10	10-15	>15	




Fig. 7.2 Land Suitability map of Maize

An area of about 142 ha (36%) moderately suitable (S2) for growing maize and distributed in the northern, central and southwestern part of the microwatershed with minor limitation of rooting depth, gravelliness, texture and calcareousness. Major area of about 213 ha (55%) has marginally suitable (Class S3) lands. They have moderate

limitations of texture, rooting depth, gravelliness and calcareousness and distributed in all parts of the microwatershed.

7.3 Land Suitability for Bajra (*Pennisetum glaucum*)

Bajra is one of the major food crop grown in an area of 2.34 lakh ha in Karnataka in the northern districts. The crop requirements (Table 7.4) for growing bajra were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and land suitability map for growing bajra was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.3.

Table 7.4 Crop suitability criteria for Bajra

Crop requirement		Rating			
Soil –site characteristics	Unit	Highly suitable(S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable(N)
Slope	%	2-3	3-8	8-15	>15
LGP	Days	120-150	120-90	<90	
Soil drainage	Class	Well to mod.Well drained	imperfect	Poorly/excessively	V.poorly
Soil reaction	pH	5.5-8.0	5.0-5.5,7.8-8.4	8.4-9.0	>9.0
Surface soil texture	Class	C (red), sicl, sc, sl,cl	l, c (black) scl, sil, sic	S1, ls	S, fragmental skeletal
Soil depth	Cm	100-75	50-75	25-50	<25
Gravel content	% vol.	15-35	35-60	60-80	-
Salinity (EC)	dSm ⁻¹	2-4	4-8	8-10	>10
Sodicity (ESP)	%	5-8	8-10	10-15	>15




Fig. 7.3 Land Suitability map of Bajra

Highly suitable (Class S1) land occupying an area of about 88 ha (23%) for growing bajra and occur in the eastern part of the microwatershed. A small area of about 54 ha (14%) is moderately suitable (Class S2) for growing bajra and are distributed in the northwestern and southwestern part of the microwatershed They have minor limitations of gravelliness, rooting depth and calcareousness. Major area of about 213 ha (55%) is marginally suitable (Class S3) for growing bajra and occur in all parts of the microwatershed with moderate limitations of gravelliness, texture, calcareous and rooting depth.

7.4 Land Suitability for Groundnut (*Arachis hypogaea*)

Groundnut is one of the major oilseed crop grown in an area of 6.54 lakh ha in Karnataka in most of the districts either as rainfed or irrigated crop. The crop requirements for growing groundnut (Table 7.5) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing groundnut was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.4.

An area of about 142 ha (36%) is moderately suitable (Class S2) for groundnut and are distributed in the northern, central and southwestern part of the microwatershed. They have minor limitations of texture, rooting depth, gravelliness and calcareousness and marginally suitable (Class S3) lands occupy major area of about 213 ha (55%) and are distributed in all parts of the microwatershed with moderate limitations of texture, rooting depth, gravelliness and calcareousness.

Table 7.5 Crop suitability criteria for Groundnut

Crop requirement		Rating			
Soil-site characteristics	Unit	Highly suitable(S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable(N)
Slope	%	<3	3-5	5-10	>10
LGP	Days	100-125	90-105	75-90	
Soil drainage	Class	Well drained	Mod. Well drained	Imperfectly drained	Poorly drained
Soil reaction	pH	6.0-8.0	8.1-8.5,5.5-5.9	>8.5,<5.5	
Surface soil texture	Class	l, sil, sicl	cl, sc, sic, sl	s, ls,c (>60%)	s, fragmental
Soil depth	Cm	>75	50-75	25-50	<25
Gravel content	% vol.	<35	35-60	>60	
CaCO ₃ in root zone	%	high	Medium	low	
Salinity (EC)	dSm ⁻¹	<2.0	2.0-4.0	4.0-8.0	
Sodicity (ESP)	%	<5	5-10	>10	




Fig. 7.4 Land Suitability map of Groundnut

7.5 Land Suitability for Sunflower (*Helianthus annus*)

Sunflower is one of the most important oilseed crop grown in an area of 3.56 lakh ha in the State in all the districts. The crop requirements for growing sunflower (Table 7.6) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing sunflower was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.5.

Table 7.6 Crop suitability criteria for Sunflower

Crop requirement		Rating			
Soil-site characteristics	Unit	Highly suitable(S1)	Moderately suitable(S2)	Marginally suitable(S3)	Not suitable(N)
Slope	%	<3	3-5	5-10	>10
LGP	Days	>90	80-90	70-80	<70
Soil drainage	class	Well drained	mod. Well drained	imperfectly drained	Poorly drained
Soil reaction	pH	6.5-7.8	7.8-8.4,5.5-6.5	8.4-9.0;5.0-5.5	>9.0,<5.0
Surface soil texture	Class	l, cl, sil, sc	Scl, sic	ls sl	s
Soil depth	Cm	>100	75-100	50-75	<50
Gravel content	% vol.	<15	15-35	35-60	>60
Salinity (EC)	dSm ⁻¹	<1.0	1.0-2.0	>2.0	
Sodicity (ESP)	%	<10	10-15	>15	




Fig. 7.5 Land Suitability map of Sunflower

A small area of about 20 ha (5%) is highly suitable (S1) for growing sunflower and is distributed in the northern part of the microwatershed. An area of about **101 ha (26%)** is moderately suitable (Class S2) for growing sunflower and are distributed in the central and northwestern part of the microwatershed. They have minor limitations of rooting depth and calcareousness. Marginally suitable (Class S3) lands occupy small area of about 114 ha (29%) and are distributed in the northwestern and western part of the microwatershed with moderate limitations of rooting depth, gravelliness and calcareousness and an area of about 120 ha (31%) is not suitable (Class N1) for growing sunflower and occur in the western and southern part of the microwatershed with severe limitations of gravelliness, calcareousness and rooting depth.

7.6 Land Suitability for Chilli (*Capsicum annum L*)

Chilli is one of the major fruit and spice crop grown in an area of 0.42 lakh ha in Karnataka State. The crop requirements for growing chilli (Table 7.7) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing chilli was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.6.

An area of about 142 ha (36%) is moderately suitable (Class S2) for growing chilli and are distributed in the central, northern and southwestern part of the microwatershed. They have minor limitations of rooting depth, gravelliness and calcareousness. Marginally suitable (Class S3) lands occupy major area of about 213 ha (55%) and are

distributed in all parts of the microwatershed with moderate limitations of rooting depth, texture, gravelliness and calcareousness.

Table 7.7 Crop suitability criteria for Chilli

Crop requirement		Rating			
Soil –site characteristics	Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable(S3)	Not suitable(N)
Mean temperature in growing season	°C	20-30	30-35	35-40	>40
Slope	%	<3	3-5	5-10	>10
LGP	Days	>150	120-150	90-120	<90
Soil drainage	Class	Well drained	Moderately drained	Imp./ poor drained /excessively	V.poorly drained
Soil reaction	pH	6.5-7.8,6.0-7.0	7.8-8.4	8.4-9.0,5.0-5.9	>9.0
Surface soil texture	Class	scl, cl, sil	sl, sc, sic,c(m/k)	C(ss), ls, s	
Soil depth	Cm	>75	50-75	25-50	<25
Gravel content	% vol.	<15	15-35	35-60	>60
Salinity (ECe)	dsm ⁻¹	<1.0	1.0-2.0	2.0-4.0	<4
Sodicity (ESP)	%	<5	5-10	10-15	




Fig. 7.6 Land Suitability map of Chilli

7.7 Land Suitability for Tomato (*Solanum lycopersicum*)

Tomato is one of the most important vegetable crop grown in an area of 0.65 lakh ha in almost all the districts of the State. The crop requirements (Table 7.8) for growing tomato were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing tomato was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.7.

An area of about 142 ha (36%) has moderately suitable (Class S2) for growing tomato and are distributed in the central, northern and southwestern part of the microwatershed. They have minor limitations of rooting depth, gravelliness and calcareousness. Marginally suitable (Class S3) lands occupy major area of about 213 ha (55%) and are distributed in all parts of the microwatershed with moderate limitations of rooting depth, texture, gravelliness and calcareousness.

Table 7.8 Crop suitability criteria for Tomato

Crop requirement			Rating			
Soil-site characteristics	Unit		Highly suitable(S1)	Moderately suitable(S2)	Marginally suitable(S3)	Not suitable(N)
Climate	Temperature in growing season	°c	25-28	29-32 20-24	15-19 33-36	<15 >36
Soil moisture	Growing period	Days	>150	120-150	90-120	
Soil aeration	Soil drainage	Class	Well drained	Moderately well drained	Imperfectly drained	Poorly drained
Nutrient availability	Texture	Class	l, sl, cl, scl	Sic,siel,sc,c(m/k)	C (ss), ls	s
	pH	1:2.5	6.0-7.0	5.0-5.9,7.1-8.5	<5;>8.5	
	CaCO ₃ in root zone	%	Non calcareous	Slightly calcareous	Strongly calcareous	
Rooting conditions	Soil depth	Cm	>75	50-75	25-50	<25
	Gravel content	% vol.	<15	15-35	>35	
Soil toxicity	Salinity	ds/m	Non saline	slight	strongly	
	Sodicity (ESP)	%	<10	10-15	>15	-
Erosion	Slope	%	1-3	3-5	5-10	>10




Fig. 7.7 Land Suitability map of Tomato

7.8 Land Suitability for Drumstick (*Moringa oleifera*)

Drumstick is one of the most important vegetable crop grown in 2403 ha area in the state. The crop requirements for growing drumstick (Table 7.9) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing drumstick was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.8.

Table 7.9 Land suitability criteria for Drumstick

Crop requirement			Rating			
Soil-site characteristics		Unit	Highly suitable(S1)	Moderately suitable(S2)	Marginally suitable(S3)	Not suitable(N)
Soil aeration	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V. Poorly drained
Nutrient availability	Texture	Class	Sc, scl, cl, c (red)	Sl, c (black)	ls	S
	pH	1:2.5	5.5-6.5	5-5.5,6.5-7.3	7.8-8.4	>8.4
Rooting conditions	Soil depth	Cm	>100	75-100	50-75	<50
	Gravel content	% vol.	0-35	35-60	60-80	>80
Erosion	Slope	%	0-3	3-10	-	>10




Fig. 7.8 Land Suitability map of Drumstick

Moderately suitable (Class S2) lands occupy an area of about 121 ha (31%) and occur in the northern and northwestern part of the microwatershed. They have minor limitation of rooting depth, texture and calcareousness, An area of 114 ha (29%) is marginally (Class S3) suitable for growing drumstick with moderate limitations of rooting depth, gravelliness and calcareousness and distributed in the western, southwestern and northern part of the microwatershed. An area of about 120 ha (31%) is not suitable (Class N1) and occur in the southern and western part of the microwatershed and have severe limitations of rooting depth, gravelliness and calcareousness.

7.9 Land Suitability for Mulberry (*Morus nigra*)

Mulberry is the important leaf crop grown for rearing silkworms in about 1.66 lakh ha in all the districts of the state. The crop requirements for growing mulberry (Table 7.10) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing mulberry was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.9.

Moderately suitable (Class S2) lands occupy maximum area of about 193 ha (49%) and occur in all parts of the microwatershed. They have minor limitations of texture, gravelliness, rooting depth and calcareousness. Marginally suitable lands cover a small area of about 42 ha (11%) and occur in the northern and southwestern part of the microwatershed. They have moderate limitations of rooting depth, gravelliness and calcareousness. and an area of about 120 ha (31%) is not suitable (Class N1) and occur in

the southern and western part of the microwatershed and have severe limitations of rooting depth, gravelliness and calcareousness.

Table 7.10 Land suitability criteria for Mulberry

Crop requirement			Rating			
Soil-site characteristics		Unit	Highly suitable(S1)	Moderately suitable(S2)	Marginally suitable(S3)	Not suitable(N)
Soil aeration	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V. Poorly drained
Nutrient availability	Texture	Class	Sc, cl, scl	C (red)	C (black), sl, ls	-
	pH	1:2.5				
Rooting conditions	Soil depth	Cm	>100	75-100	50-75	<50
	Gravel content	% vol.	0-35	35-60	60-80	>80
Erosion	Slope	%	0-3	3-5	5-10	>10

Note: Suitability evaluation only for Mulberry leaf not for Silk worm rearing




Fig. 7.9 Land Suitability map of Mulberry

7.10 Land suitability for Mango (*Mangifera indica*)

Mango is one of the most important fruit crop grown in about 1.73 lakh ha in almost all the districts of the State. The crop requirements (Table 7.11) for growing mango were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing mango was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.10.

Table 7.11 Crop suitability criteria for Mango

Crop requirement			Rating			
Soil-site characteristics	Unit		Highly suitable(S1)	Moderately suitable (S2)	Marginally suitable(S3)	Not suitable(N)
Climate	Temp. in growing season	°C	28-32	24-27 33-35	36-40	20-24
	Min. temp. before flowering	°C	10-15	15-22	>22	
Soil moisture	Growing period	Days	>180	150-180	120-150	<120
Soil aeration	Soil drainage	Class	Well drained	Mod. To imp. drained	Poor drained	V.poorly drained
	Water table	M	>3	2.50-3.0	2.5-1.5	<1.5
Nutrient availability	Texture	Class	Sc, l, sil, cl	sc, sic, l, c	C (<60%), ls, sl	C (>60%),
	pH	1:2.5	5.5-7.5	7.6-8.55.0-5.4	8.6-9.04.0-4.9	>9.0<4.0
	OC	%	High	medium	low	
	CaCO ₃ in root zone	%	Non calcareous	<5	5-10	>10
Rooting conditions	Soil depth	cm	>200	125-200	75-125	<75
	Gravel content	% vol	Non-gravelly	<15	15-35	>35
Soil toxicity	Salinity	dS/m	Non saline	<2.0	2.0-3.0	>3.0
	Sodicity	%	Non sodic	<10	10-15	>15
Erosion	Slope	%	<3	3-5	5-10	




Fig. 7.10 Land Suitability map of Mango

There are no highly (Class S1) and moderately (Class S2) suitable lands for growing mango. Marginally suitable (Class S3) lands cover a maximum area of about 193 ha (49%) and occur in all parts of the microwatershed. They have moderate limitations of rooting depth, gravelliness, texture and calcareousness. About 162 ha (42%) is not suitable (Class N1) for growing mango and occur in the northern, southwestern and northwestern part of the microwatershed with severe limitations of texture, calcareousness, gravelliness and rooting depth .

7.11 Land suitability for Sapota (*Manilkara zapota*)

Sapota is one of the most important fruit crop grown in an area of about 29373 ha in almost all the districts of the state. The crop requirements (Table 7.12) for growing sapota were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing sapota was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.11.

About 101 ha (26%) area is moderately suitable (S2) for growing sapota and occur in the eastern and northwestern part of the microwatershed with minor limitations of rooting depth, gravelliness and calcareousness. Marginally suitable (Class S3) lands cover a maximum area of about 134 ha (34%) and occur in the northern and western part of the microwatershed. They have moderate limitations of rooting depth, texture and calcareousness and an area of about 120 ha (31%) is not suitable (Class N1) for growing sapota and occur in the northwestern and northern part of the microwatershed with severe limitations of calcareousness, gravelliness and rooting depth.

Table 7.12 Crop suitability criteria for Sapota

Crop requirement			Rating			
Soil –site characteristics	Unit		Highly suitable(S1)	Moderately suitable(S2)	Marginally suitable(S3)	Not suitable(N)
Climate	Temperature in growing season	° C	28-32	33-36	37-42	>42
				24-27	20-23	<18
Soil moisture	Growing period	Days	>150	120-150	90-120	<120
Soil aeration	Soil drainage	Class	Well drained	Moderately well drained	Imperfectly drained	Poorly drained
Nutrient availability	Texture	Class	Scl, l, cl, sil	Sl, sicl, sc	C (<60%),ls	, s,C (>60%)
	pH	1:2.5	6.0-7.5	7.6-8.0,5.0-5.9	8.1-9.0,4.5-4.9	>9.0,<4.5
	CaCO ₃ in root zone	%	Non calcareous	<10	10-15	>15
Rooting conditions	Soil depth	Cm	>150	75-150	50-75	<50
	Gravel content	% vol.	Non gravelly	<15	15-35	<35
Soil toxicity	Salinity	dS/m	Non saline	Up to 1.0	1.0-2.0	2.0-4.0
	Sodicity	%	Non sodic	10-15	15-25	>25
Erosion	Slope	%	<3	3-5	5-10	>10




Fig. 7.11 Land Suitability map of Sapota

7.12 Land Suitability for Pomegranate (*Punica granatum*)

Pomegranate is one of the commercially grown fruit crop in about 18488 ha in Karnataka mainly in Bijapur, Bagalkot, Koppal, Gadag and Chitradurga districts. The crop requirements for growing pomegranate (Table 7.13) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing pomegranate was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.12.

Moderately suitable (Class S2) lands occupy major area of about 121 ha (31%) and are distributed in the northern and northeastern part of the microwatershed. They have minor limitations of rooting depth, gravelliness, calcareousness and texture. Marginally suitable (Class S3) lands for growing pomegranate occupy an area of about 114 ha (29%) and are distributed in the western and southwestern part of the microwatershed with moderate limitations of rooting depth, gravelliness and calcareousness. An area of about 120 ha (31%) is not suitable (Class N1) for growing pomegranate and occur in the northwestern and southern part of the microwatershed with severe limitations of calcareousness, gravelliness and rooting depth.

Table 7.13 Crop suitability criteria for Pomegranate

Crop requirement			Rating			
Soil –site characteristics		Unit	Highly suitable(S1)	Moderately suitable(S2)	Marginally suitable(S3)	Not suitable(N)
Climate	Temperature in growing season	⁰ C	30-34	35-38 25-29	39-40 15-24	
Soil moisture	Growing period	Days	>150	120-150	90-120	<90
Soil aeration	Soil drainage	Class	Well drained	imperfectly drained		
Nutrient availability	Texture	Class	Sl, scl, l, cl	C, sic, siel	Cl, s, ls	S,fragmental
Rooting conditions	pH	1:2.5	5.5-7.5	7.6-8.5	8.6-9.0	
	Soil depth	Cm	>100	75-100	50-75	<50
	Gravel content	% vol.	nil	15-35	35-60	>60
Soil toxicity	Salinity	dS/m	Nil	<9	>9	<50
	Sodicity	%	nil			
Erosion	Slope	%	<3	3-5	5-10	




Fig. 7.12 Land Suitability map of Pomegranate

7.13 Land Suitability for Guava (*Psidium guajava*)

Guava is one of the most important fruit crop grown in an area of about 6558 ha in almost all the districts of the state. The crop requirements (Table 7.14) for growing guava were matched with the soil-site characteristics (Table 7.1) and a land suitability map for

growing guava was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.13.

Table 7.14 Crop suitability criteria for Guava

Crop requirement			Rating			
Soil –site characteristics	Unit		Highly suitable(S1)	Moderately suitable(S2)	Marginally suitable(S3)	Not suitable(N)
Climate	Temperature in growing season	⁰ C	28-32	33-36 24-27	37-42 20-23	
Soil moisture	Growing period	Days	>150	120-150	90-120	<90
Soil aeration	Soil drainage	Class	Well drained	Mod. to imperfectly	poor	Very poor
Nutrient availability	Texture	Class	Scl, l, cl, sil sc, c (red)	Sl,sicl,sic	C (<60%),ls	C (>60%)
	pH	1:2.5	6.0-7.5	7.6-8.0:5.0-5.9	8.1-8.5:4.5-4.9	>8.5:<4.5
	CaCO ₃ in root zone	%	Non calcareous	<10	10-15	>15
Rooting conditions	Soil depth	Cm	>100	75-100	50-75	<50
	Gravel content	% vol.	<15	15-35	>35	
Soil toxicity	Salinity	dS/m	<2.0	2.0-4.0	4.0-6.0	
	Sodicity	%	Non sodic	10-15	15-25	>25
Erosion	Slope	%	<3	3-5	5-10	>10




Fig. 7.13 Land Suitability map of Guava

About 101 ha (26%) area is moderately suitable (S2) for growing guava and occur in the eastern and northwestern part of the microwatershed with minor limitations of rooting depth, gravelliness and calcareousness. Marginally suitable (Class S3) lands cover a maximum area of about 134 ha (34%) and occur in the northern and western part of the microwatershed. They have moderate limitations of rooting depth, gravelliness, texture and calcareousness An area of about 120 ha (31%) is not suitable (Class N1) for growing guava and occur in the northwestern and northern part of the microwatershed with severe limitations of texture, gravelliness and rooting depth.

7.14 Land Suitability for Jackfruit (*Artocarpus heterophyllus*)

Jackfruit is the most important fruit crop grown in 5368 ha in all the districts of the state. The crop requirements (Table 7.15) for growing jackfruit were matched with the soil-site characteristics and a land suitability map for growing jackfruit was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in figure 7.14.

About 101 ha (26%) area is moderately suitable (S2) for growing jackfruit and occur in the eastern and northwestern part of the microwatershed with minor limitations of rooting depth, gravelliness and calcareousness. Marginally suitable (Class S3) lands cover a maximum area of about 134 ha (%) and occur in the northern and western part of the microwatershed. They have moderate limitations of rooting depth, gravelliness, texture and calcareousness. An area of about 120 ha (31%) is not suitable (Class N1) for growing jackfruit and occur in the northwestern and northern part of the microwatershed with severe limitations of rooting depth, gravelliness and rooting depth

Table 7.15 Crop suitability criteria for Jackfruit

Crop requirement			Rating			
Soil –site characteristics		Unit	Highly suitable(S1)	Moderately suitable(S2)	Marginally suitable (S3)	Not suitable(N)
Soil aeration	Soil drainage	class	well	Mod. well	Poorly	V. Poorly
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	-	sl, ls, c (black)	-
	pH	1:2.5	5.5-7.3	5.0-5.5,7.3-7.8	7.8-8.4	>8.4
Rooting conditions	Soil depth	Cm	>100	75-100	50-75	<50
	Gravel content	% vol.	<15	15-35	35-60	>60
Erosion	Slope	%	0-3	3-5	>5	-




Fig. 7.14 Land Suitability map of Jackfruit

7.15 Land Suitability for Jamun (*Syzygium cumini*)

Jamun is an important fruit crop grown in almost all the districts of the state. The crop requirements (Table 7.16) for growing jamun were matched with the soil-site characteristics and a land suitability map for growing jamun was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.15.

Table 7.16 Crop suitability criteria for Jamun

Crop requirement			Rating			
Soil –site characteristics		Unit	Highly suitable(S1)	Moderately suitable(S2)	Marginally suitable (S3)	Not suitable(N)
Soil aeration	Soil drainage	Class	Well	Mod. well	Poorly	V.Poorly
Nutrient availability	Texture	Class	Scl, cl, sc, C(red)	Sl, C black)	ls	-
	pH	1:2.5	6.0-7.8	5.0-6.0	7.8-8.4	>8.4
Rooting conditions	Soil depth	Cm	>150	100-150	50-100	<50
	Gravel content	% vol.	<15	15-35	35-60	>60
Erosion	Slope	%	0-3	3-5	5-10	>10




Fig. 7.15 Land Suitability map of Jamun

There are no highly suitable (Class S1) lands for growing jamun. An area of about 108 ha (28%) is moderately suitable (Class S2) and occur in the northern and eastern part of the microwatershed. They have minor limitations of rooting depth, texture, gravelliness and calcareousness. The marginally suitable (Class S3) lands cover an area of about 127 ha (32%) and are distributed in the northwestern and western part of the microwatershed with moderate limitations of rooting depth, gravelliness, calcareousness and texture. an area of about 120 ha (31%) is not suitable (Class N1) for growing jamun and are distributed in the northwestern and southern part of the microwatershed with severe limitations of rooting depth, gravelliness and texture.

7.16 Land Suitability for Musambi (*Citrus limetta*)

Musambi is one of the most important fruit crop grown in an area of 5446 ha in almost all the districts of the state. The crop requirements (Table 7.17) for growing musambi were matched with the soil-site characteristics (Table 7.14) and a land suitability map for growing musambi was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.16.

Table 7.17 Crop suitability criteria for Musambi

Crop requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable(S2)	Marginally suitable(S3)	Not suitable(N)
Climate	Temperature in growing season	°C	28-30	31-35 24-27	36-40 20-23	>40 <20
Soil moisture	Growing period	Days	240-265	180-240	150-180	<150
Soil aeration	Soil drainage	Class	Well drained	Mod. to imp. drained	Poorly	Very poorly
Nutrient availability	Texture	Class	Scl,l,siel,cl,s	Sc, sc, c	C(>70%),ls	S
	pH	1:2.5	6.0-7.5	5.5-6.47.6-8.0	4.0-5.4,8.1-8.5	<4.0,>8.5
	CaCO ₃ in root zone	%	Non calcareous	Upto 5	5-10	>10
Rooting conditions	Soil depth	Cm	>150	100-150	50-100	<50
	Gravel content	% vol.	Non gravelly	15-35	35-55	>55
Soil toxicity	Salinity	dS/m	Non saline	Upto 1.0	1.0-2.5	>2.5
	Sodicity	%	Non sodic	5-10	10-15	>15
Erosion	Slope	%	<3	3-5	5-10	




Fig. 7.16 Land Suitability map of Musambi

A small area of about 20 ha (5%) is highly suitable (Class S1) for growing musambi and occur in the northern and northwestern part of the microwatershed. An area of about 101 ha (26%) is moderately suitable (Class S2) for growing musambi and are distributed in the northwestern and eastern part of the microwatershed. They have minor

limitations of rooting depth, gravelliness and calcareousness. An area of about 114 ha (29%) is marginally suitable (Class S3) for growing musambi and are distributed in the southwestern and northwestern part of the microwatershed with moderate limitations of rooting depth, gravelliness and calcareousness. An area of about 120 ha (29%) is not suitable (Class N1) for growing musambi and are distributed in the northwestern and southern part of the microwatershed with severe limitations of rooting depth, gravelliness and calcareousness.

7.17 Land Suitability for Lime (*Citrus sp*)

Lime is one of the most important fruit crop grown in an area of 11752 ha in almost all the districts of the State. The crop requirements (Table 7.18) for growing lime (Table 7.15) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing lime was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.17.

Table 7.18 Crop suitability criteria for Lime

Crop requirement			Rating			
Soil –site characteristics	Unit		Highly suitable (S1)	Moderately suitable(S2)	Marginally suitable(S3)	Not suitable(N)
Climate	Temperature in growing season	°C	28-30	31-35 24-27	36-40 20-23	>40 <20
Soil moisture	Growing period	Days	240-265	180-240	150-180	<150
Soil aeration	Soil drainage	Class	Well drained	Mod. to imp. drained	Poorly	Very poorly
Nutrient availability	Texture	Class	Scl,l,sicl,cl,s	Sc, sc, c	C(>70%), ls	S
	pH	1:2.5	6.0-7.5	5.5-6.4,7.6-8.0	4.0-5.4,8.1-8.5	<4.0,>8.5
	CaCO ₃ in root zone	%	Non calcareous	Upto 5	5-10	>10
Rooting conditions	Soil depth	Cm	>150	100-150	50-100	<50
	Gravel content	% vol.	Non gravelly	15-35	35-55	>55
Soil toxicity	Salinity	dS/m	Non saline	Upto 1.0	1.0-2.5	>2.5
	Sodicity	%	Non sodic	5-10	10-15	>15
Erosion	Slope	%	<3	3-5	5-10	




Fig. 7.17 Land Suitability map of Lime

A small area of about 20 ha (5%) is highly suitable (Class S1) for growing lime and occur in the northern and northwestern part of the microwatershed. An area of about 101 ha (26%) is moderately suitable (Class S2) for growing lime and are distributed in the northwestern and eastern part of the microwatershed. They have minor limitations of rooting depth, gravelliness and calcareousness. An area of about 114 ha (29%) is marginally suitable (Class S3) for growing lime and are distributed in the southwestern and northwestern part of the microwatershed with moderate limitations of rooting depth, gravelliness and calcareousness. An area of about 120 ha (29%) is not suitable (Class N1) for growing lime and are distributed in the northwestern and southern part of the microwatershed with severe limitations of rooting depth, gravelliness and calcareousness.

7.18 Land Suitability for Cashew (*Anacardium occidentale*)

Cashew is one of the most important nut crop grown in an area of 7052 ha in almost all the districts of the State. The crop requirements (Table 7.19) for growing cashew were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing cashew was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.18.

Major area of about 160 ha (41%) is moderately suitable (Class S2) for growing cashew and are distributed in the western, central and eastern part of the microwatershed. They have minor limitations of rooting depth, gravelliness and texture. A small area of about 14 ha (11%) is marginally suitable (Class S3) for growing cashew and are

distributed in the southwestern and northwestern part of the microwatershed with moderate limitations of rooting depth and gravelliness. An area of about 154 ha (39%) is not suitable (Class N1) for growing cashew and are distributed in the northwestern and southern part of the microwatershed with severe limitations of rooting depth, gravelliness, texture and calcareousness.

Table 7.19 Crop suitability criteria for Cashew

Crop requirement			Rating			
Soil –site characteristics		Unit	Highly suitable(S1)	Moderately suitable(S2)	Marginally suitable (S3)	Not suitable(N)
Soil aeration	Soil drainage	Class	Well drained	Mod. well drained	Poorly drained	V. Poorly drainage
Nutrient availability	Texture	Class				
	pH	1:2.5	5.5-6.5	5.0-5.5,6.5-7.3	7.3-7.8	>7.8
Rooting conditions	Soil depth	Cm	>100	75-100	50-75	<50
	Gravel content	% vol.	<15	15-35	35-60	>60
Erosion	Slope	%	0-3	3-10	>10	




Fig. 7.18 Land Suitability map of Cashew

7.19 Land Suitability for Custard Apple (*Annona reticulata*)

Custard apple is one of the most important fruit crop grown in 1426 ha in almost all the districts of the State. The crop requirements (Table 7.20) for growing custard apple were matched with the soil-site characteristics (Table 7.1) and a land suitability map for

growing custard apple was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.19.

Table 7.20 Crop suitability criteria for Custard apple

Crop requirement			Rating			
Soil –site characteristics		Unit	Highly suitable(S1)	Moderately suitable(S2)	Marginally suitable (S3)	Not suitable(N)
Soil aeration	Soil drainage	Class	Well drained	Mod. well drained	Poorly drained	V. Poorly drained
Nutrient availability	Texture	Class	Scl, cl, sc, c (red), c (black)	-	Sl, ls	-
	pH	1:2.5	6.0-7.3	7.3-8.4	5.0-5.5,8.4-9.0	>9.0
Rooting conditions	Soil depth	Cm	>75	50-75	25-50	<25
	Gravel content	% vol.	<15-35	35-60	60-80	-
Erosion	Slope	%	0-3	3-5	>5	-




Fig. 7.19 Land Suitability map of Custard Apple

An area of about 108 ha (28%) is highly suitable (Class S1) for growing custard apple. They are distributed in the northern and eastern part of the microwatershed. Maximum area of about 127 ha (32%) is moderately suitable (Class S2) and occur in northwestern and southwestern part of the microwatershed. They have minor limitations of gravelliness, rooting depth and calcareousness and area of about 120 ha (31%) is

marginally suitable (Class S3) for growing custard apple and are distributed in the northwestern and southern part of the microwatershed with moderate limitations of rooting depth, gravelliness and calcareousness.

7.20 Land Suitability for Amla (*Phyllanthus emblica*)

Amla is the most important fruit and medicinal crop grown in an area of 151 ha and distributed in almost all the districts of the state. The crop requirements for (Table 7.21) growing amla were matched with the soil-site characteristics and a land suitability map for growing amla was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.20.

An area of about 88 ha (23%) is highly suitable (Class S1) for growing amla. They are distributed in the eastern part of the microwatershed. Maximum area of about 147 ha (37%) is moderately suitable (Class S2) and occur in northern and southwestern part of the microwatershed. They have minor limitations of gravelliness, rooting depth, texture and calcareousness and area of about 120 ha (31%) is marginally suitable (Class S3) for growing amla and are distributed in the northwestern and southern part of the microwatershed with moderate limitations of rooting depth, gravelliness, texture and calcareousness.

Table 7.21 Crop suitability criteria for Amla

Crop requirement			Rating			
Soil –site characteristics		Unit	Highly suitable(S1)	Moderately suitable(S2)	Marginally suitable (S3)	Not suitable(N)
Soil aeration	Soil drainage	Class	Well drained	Mod.well drained	Poorly drained	V. Poorly drained
Nutrient availability	Texture	Class	Scl,cl,sc,c(red)	C (black)	ls, sl	-
	pH	1:2.5	5.5-7.3	5.0-5.5	7.8-8.4	>8.4
Rooting conditions	Soil depth	Cm	>75	50-75	25-50	<25
	Gravel content	% vol.	<15-35	35-60	60-80	
Erosion	Slope	%	0-3	3-5	5-10	>10




Fig. 7.20 Land Suitability map of Amla

7.21 Land Suitability for Tamarind (*Tamarindus indica*)

Tamarind is the most important spice crop grown in 14897 ha in all the districts of the state. The crop requirements (Table 7.22) for growing tamarind were matched with the soil-site characteristics and a land suitability map for growing tamarind was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.21.

Table 7.22) Crop suitability criteria for Tamarind

Crop requirement			Rating			
Soil –site characteristics		Unit	Highly suitable(S1)	Moderately suitable(S2)	Marginally suitable (S3)	Not suitable(N)
Soil aeration	Soil drainage	Class	Well drained	Mod.well drained	Poorly drained	V.Poorly drained
Nutrient availability	Texture	Class	Scl,cl,sc,c(red)	Sl, c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0,7.3-7.8	7.8-8.4	>8.4
Rooting conditions	Soil depth	Cm	>150	100-150	75-100	<75
	Gravel content	% vol.	<15	15-35	35-60	60-80
Erosion	Slope	%	0-3	3-5	5-10	>10




Fig. 7.21 Land Suitability map of Tamarind

There are no highly suitable lands (Class S1) for growing tamarind. An area of about 20 ha (5%) is moderately suitable (Class S2) and occurs in the northern part of the microwatershed. They have minor limitations of rooting depth. Major area of about 173 ha (44%) is marginally suitable (Class S3) and occur in the central, eastern and western part of the microwatershed. They have moderate limitations of rooting depth, gravelliness and calcareousness followed by an area of about 162 ha (42%) is not suitable (Class N1) for growing tamarind and are distributed in the northwestern and southern part of the microwatershed. They have severe limitations of rooting depth, gravelliness and calcareousness.

7.22 Land Suitability for Marigold (*Tagetes erecta*)

Marigold is one of the most important flower crop grown in an area of 9108 ha in almost all the districts of the state. The crop requirements (Table 7.23) for growing marigold were matched with the soil-site characteristics and a land suitability map for growing marigold was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.22.

Table 7.23 crop suitability criteria for Marigold

Crop requirement			Rating			
Soil –site characteristics		Unit	Highly suitable(S1)	Moderately Suitable(S2)	Marginally suitable(S3)	Not suitable(N)
climate	Temperature in growing season		18-23	17-15 24-35	35-40 10-14	>40 <10
Soil aeration	Soil drainage	class	Well drained	Moderately well drained	Imperfectly drained	Poorly drained
Nutrient availability	Texture	Class	l,sl,scl,cl, sil	sicl, sc, sic, c	C, ls	s
	pH	1:2.5	7.0-7.5	5.5-5.9,7.6-8.5	<5,>8.5	-
	CaCO ₃ in root zone	%	Non calcareous	Slightly calcareous	Strongly calcareous	-
Rooting conditions	Soil depth	Cm	>75	50-75	25-50	<25
	Gravelcontent	% vol.	<15	15-35	>35	-
Soil toxicity	Salinity	ds/m	Non saline	Slightly	Strongly	-
	Sodicity(ESP)	%	<10	10-15	>15	-
Erosion	Slope	%	1-3	3-5	5-10	-




Fig. 7.22 Land Suitability map of Marigold

An area of about **163 ha (42%)** is moderately suitable (Class S2) for growing marigold and occur in the northern, central and southwestern part of the microwatershed. They have minor limitations of calcareousness, gravelliness, rooting depth and texture and major area of about 192 ha (49%) is marginally suitable (Class S3) for growing marigold and occur in the northwestern, southern and western part of the microwatershed. They have severe limitations of rooting depth, gravelliness and calcareousness.

7.23 Land Suitability for Chrysanthemum (*Chrysanthemum indicum*)

Chrysanthemum is one of the most important flower crop grown in an area of 4978 ha in almost all the districts of the State. The crop requirements (Table 7.24) for growing chrysanthemum were matched with the soil-site characteristics and a land suitability map for growing chrysanthemum was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed is given in Figure 7.23.

Table 7.24 Crop suitability criteria for Chrysanthemum

Crop requirement			Rating			
Soil –site characteristics	Unit		Highly suitable(S1)	Moderately Suitable(S2)	Marginally suitable(S3)	Not suitable(N)
climate	Temperature in growing season		18-23	17-15 24-35	35-40 10-14	>40 <10
Soil aeration	Soil drainage	class	Well drained	Mod. well drained	Imperfectly drained	Poorly drained
Nutrient availability	Texture	Class	l,sl, scl, cl, sil	sicl, sc, sic, c	C , ls	, s
	pH	1:2.5	7.0-7.5	5.5-5.9,7.6-8.5	<5,>8.5	
	CaCO ₃ in root zone	%	Non calcareous	Slightly calcareous	Strongly calcareous	
Rooting conditions	Soil depth	Cm	>75	50-75	25-50	<25
	Gravel content	% vol.	<15	15-35	>35	
Soil toxicity	Salinity	ds/m	Non saline	slightly	strongly	
	Sodicity(ESP)	%	<10	10-15	>15	-
Erosion	Slope	%	1-3	3-5	5-10	




Fig. 7.23 Land Suitability map of Chrysanthemum

An area of about 163 ha (42%) is moderately suitable (Class S2) for growing chrysanthemum and occur in the northern, central and southwestern part of the microwatershed. They have minor limitations of calcareousness, gravelliness, rooting depth and texture and major area of about 192 ha (49%) is marginally suitable (Class S3) for growing chrysanthemum and occur in the northwestern, southern and western part of the microwatershed. They have severe limitations of rooting depth, gravelliness and calcareousness.

7. 24 Land Suitability for Jasmine (*Jasminum sp.*)

Jasmine is the most important flower crop grown in an area of 803 ha in almost all the districts of the State. The crop requirements (Table 7.25) for growing jasmine were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing jasmine was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.24.

An area of about 143 ha (37%) is moderately suitable (Class S2) for growing jasmine and occur in the northern, central and northwestern part of the microwatershed. They have minor limitations of calcareousness, gravelliness, rooting depth and texture and major area of about 212 ha (54%) is marginally suitable (Class S3) for growing jasmine and occur in the northwestern, southern and western part of the microwatershed. They have moderate limitations of texture, rooting depth, gravelliness and calcareousness.

Table 7.25 Land suitability criteria for jasmine (irrigated)

Crop requirement			Rating			
Soil-site characteristics	Unit		Highly suitable(S1)	Moderately suitable(S2)	Marginally suitable(S3)	Not suitable(N)
Climate	Temperature in growing season		18-23	17-15 24-35	35-40 10-14	
Soil aeration	Soil drainage	Class	Well drained	Moderately drained	Imperfectly drained	Poorly drained
Nutrient availability	Texture	Class	Scl, l, scl, cl, sil	si cl, sc, sic, c(m/k)	C(ss), ls,	s
	pH	1:2.5	6.0-7.5	5.5-5.9, 7.6-8.5	<5, >8.5	
	CaCO ₃ in root zone	%	Non calcareous	Slightly calcareous	Strong calcareous	
Rooting conditions	Soil depth	Cm	>75	50-75	25-50	<25
	Gravel content	% vol.	<15	15-35	>35	
Soil toxicity	Salinity	ds/m	Non saline	Slight	Strongly	
	Sodicity	%	Non sodic	Slight	Strongly	
Erosion	Slope	%	1-3	3-5	5-10	




Fig. 7.24 Land Suitability map of Jasmine

7.25 Land Suitability for Crossandra (*Crossandra infundibuliformis*.)

Crossandra is the most important flower crop grown in all the districts of the state. The crop requirements for growing crossandra were matched with the soil-site characteristics and a land suitability map for growing crossandra was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.25.




Fig. 7.25 Land Suitability map of Crossandra

An area of about 142 ha (36%) is moderately suitable (Class S2) for growing crossandra and occur in the northern, eastern, southwestern and northwestern part of the microwatershed. They have minor limitations of calcareousness, gravelliness, rooting depth and texture and major area of about 213 ha (55%) is marginally suitable (Class S3) for growing crossandra and occur in the northwestern, southern and western part of the microwatershed. They have moderate limitations of texture, rooting depth, gravelliness and calcareousness.

7.26 Land Management Units (LMU)

The 9 soil map units identified in Bettageri-1 microwatershed have been grouped into 7 Land Management Units (LMU) for the purpose of preparing a Proposed Crop Plan. Land Management Units are grouped based on the similarities in respect of the type of soil, the depth of the soil, the surface soil texture, gravel content, AWC, slope, erosion etc. and a Land Management Units map (Fig.7.25) has been generated. These Land Management Units are expected to behave similarly for a given level of management.

The map units that have been grouped into seven Land Management Units along with brief description of soil and site characteristics are given below.

LMU	Mapping unit	Soil and site characteristics
1	371.GRHmB1	Deep, black calcareous clay soils with slopes of 1-3%, slight erosion
2	188.BDGhB2g1	Moderately deep, red gravelly clay soils with slopes of 1-3%, moderate erosion, gravelly (15-35%)
3	151.TGRiB2g1 174.CKMhB1g1	Moderately deep, red calcareous to non calcareous sandy clay to sandy clay loam soils with slopes of 1-3%, slight to moderate erosion, gravelly (15-35%)
4	91.MKHiB2g2	Moderately shallow, gravelly red loamy soils with slopes of 1-3%, moderate erosion, very gravelly (35-60%)
5	337.RNKmB2g1	Moderately shallow, black calcareous clay soils with slopes of 1-3%, moderate erosion, gravelly (15-35%)
6	305.MTLiB2g2	Shallow, calcareous black gravelly sandy clay to clay soils with slopes of 1-3%, moderate erosion, very gravelly (35-60%)
7	24.HRVhB1g2 27.HRVhB2g2	Shallow, red gravelly loamy soils with slopes of 1-3%, slight to moderate erosion, very gravelly (35-60%)




Fig 7.25 Land Management Units map of Bettageri-1 microwatershed

7.26 Proposed Crop Plan for Bettageri-1 Microwatershed

After assessing the land suitability for the 25 crops, the proposed crop plan has been prepared for the 7 identified LUCs by considering only the highly (Class S1) and moderately (Class S2) suitable lands for each of the 25 crops. The resultant proposed crop plan is presented in Table 7.26.

Table 7.26 Proposed Crop Plan for Bettageri-1 Microwatershed

Proposed Land use Class	Soil Map Units	Survey Number	Field Crops	Horticulture Crops	Suitable Interventions
1	371.GRHmB1 (Deep, calcareous black clay soils)	Betageri : 10,109,110,111,112, 113, 200,201	Sunflower, Sorghum, Cotton, Bengal gram, Safflower, Linseed, Bajra	Fruit crops: Musambi, Pomegranate, Lime, Tamarind, Jamun, Amla, Custard apple Vegetables: Drumstick, Chilli, Coriander, Bhendi Flowers: Marigold, Chrysanthemum	Application of FYM, Biofertilizers and micronutrients, drip irrigation, Mulching, Suitable soil and water conservation practises
2	188.BDGhB2g1 (Moderately deep, red gravelly sandy clay to sandy clay loam soils)	Betageri : 428,429,433,434,436,437,438,439,440,441,442, 447,448,449,450,451,452,453,454,455,510,513	Groundnut, Red gram, Bajra, Horsegram, Castor	Fruit crops: Musambi, Lime, Jamun, Jackfruit Amla, Custard apple Vegetables: Drumstick	Drip irrigation, mulching, suitable soil and water conservation practises (Crescent Bunding with Catch Pit etc)
3	151.TGRiB2g1 174.CKMhB1g1 (Moderately deep, red calcareous to non calcareous sandy clay to sandy clay loam soils)	Betageri : 345,346,348,349,350,351,352,353,354,355,356, 357,358,359,360,365,366,367,372,373,374,375, 376,589,590,591,592	Sorghum, Bajra, Groundnut, Redgram, Castor	Fruit crops: Pomegranate, Sapota, Jackfruit, Jamun, Lime, Musambi, Amla, Custard apple Vegetables: Drumstick, Tomato, Chilli, Brinjal Flowers: Marigold, Chrysanthemum, Jasmine	Drip irrigation, mulching, suitable soil and water conservation practises (Crescent Bunding with Catch Pit etc)
4	91.MKHiB2g2 (Moderately shallow, gravelly red loamy soils)	Betageri 2,3,4,5,6,8,9,377, 378,379,380,381, 383,384,435,588	Sorghum, Groundnut, Bajra, Castor	Fruit crops: Amla, Cashew, Custard apple	Drip irrigation, mulching, suitable soil and water conservation practises (Crescent Bunding with Catch Pit etc)

5	337.RNKmB2g1 (Moderately shallow, black calcareous clay soils)	Betageri: 313,314	Sorghum, Bajra, Bengal gram, Linseed, Safflower, Coriander	Fruit crops: Amla, Custard apple Flowers: Marigold, Jasmine Chrysanthemum	Application of FYM, Biofertilizers and micronutrients, drip irrigation, mulching, suitable soil and water conservation practises
6	305.MTLiB2g2 (Shallow, calcareous black gravelly sandy clay to clay soils)	Betageri : 312,316,317,318, 319,320,321,322, 323,324,325,326, 327,328,335,336, 337,338,339,340, 341,342,343,344, 347,361,362,363, 364,382	Bengal gram, safflower, linseed, coriander	Agri-Silvi-Pasture: Hybrid Napier, <i>Styloxanthes hamata</i> , <i>Styloxanthes scabra</i>	Sowing across the slope, drip irrigation and mulching is recommended
7	24.HRVhB1g2 27.HRVhB2g2 (Shallow, red gravelly loamy soils)	Betageri: 443,444,445,446, 583,584	Horsegram, Bajra	Agri-Silvi-Pasture: Custard apple, Amla, Hybrid Napier, <i>Styloxanthes hamata</i> , Glyricidia, <i>Styloxanthes scabra</i>	Use of short duration varieties, sowing across the slope and split application of nitrogen fertilizers

SOIL HEALTH MANAGEMENT

8.1 Soil Health

Soil health is basic to plant health and plant health is basic to human health. Soil is fundamental to crop production. Without soil, no food could be produced nor would livestock be fed on a large scale. Because it is finite and fragile, soil is a precious resource that requires special care from its users.

Soil health or the capacity of the soil to function is critical to human survival. Soil health has been defined as: “the capacity of the soil to function as a living system without adverse effect on the ecosystem”. Healthy soils maintain a diverse community of soil organisms that help to form beneficial symbiotic associations with plant roots, recycle essential plant nutrients, improve soil structure with positive repercussions for soil, water and nutrient holding capacity and ultimately improve crop production and also contribute to mitigating climate change by maintaining or increasing its carbon content.

Functional interactions of soil biota with organic and inorganic components, air and water determine a soil’s potential to store and release nutrients, and water to plants and to promote and sustain plant growth. Thus, maintaining soil health is vital to crop production and conserve soil resource base for sustaining agriculture.

The most important characteristics of a healthy soil are

- Good soil tilth
- Sufficient soil depth
- Good water storage and good drainage
- Adequate supply, but not excess of nutrients
- Large population of beneficial organisms
- Small proportion of plant pathogens and insect pests
- Low weed pressure
- Free of chemicals and toxins that may harm the crop
- Resistance to degradation
- Resilience when unfavorable conditions occur

Characteristics of Bettageri-1 Microwatershed

- ❖ The soil phases with sizeable area identified in the microwatershed belonged to the soil series of MTL (101 ha), CKM (88 ha), BDG (72 ha), MKH (41 ha) and other series in a small area.
- ❖ As per land capability classification, entire area in the microwatershed falls under arable land category (Class II and III). The major limitations identified in the arable lands were soil and erosion.
- ❖ On the basis of soil reaction, an area of about 261 ha (67%) under slightly alkaline (pH 7.3-7.8) to very strongly alkaline in reaction (pH >9.0), small area of 23 ha (6%) is under slightly acid (pH 6.0-6.5) and 70 ha (18%) area is neutral in reaction.

Soil Health Management

The following actions are required to improve the current land husbandry practices that provide a sound basis for the successful adoption of sustainable crop production system.

Acid soils

1. Growing of crops suitable for a particular soil pH.
2. Ameliorating the soils through the application of amendments (liming materials).

Liming materials:

1. CaCO_3 (Calcium Carbonate). More than 90% use in India.
2. Dolomite [$\text{Ca Mg} (\text{CO}_3)_2$]
3. Quick lime (Cao)
4. Slaked lime [$\text{Ca} (\text{OH})_2$]

For normal pH and pH-4.8 (35 t/ha) and pH 6.0-7.0 (4 t/ha) lime is required.

Alkaline soils

(Slightly alkaline to strongly alkaline soils)

1. Regular addition of organic manure, green manuring, green leaf manuring, crop residue incorporation and mulching needs to be taken up to improve the soil organic matter status.
2. Application of biofertilizers (Azospirillum, Azotobacter, Rhizobium).
3. Application of 25% extra N and P (125 % RDN&P).
4. Application of ZnSO_4 – 12.5 kg/ha (once in three years).
5. Application of Boron – 5 kg/ha (once in three years).

Neutral soils

1. Regular addition of organic manure, green manuring, green leaf manuring, crop residue incorporation and mulching needs to be taken up to improve the soil organic matter status.
2. Application of biofertilizers, (Azospirillum, Azotobacter, Rhizobium).
3. Application of 100 per cent RDF.
4. Need based micronutrient applications.

Besides the above recommendations, the best transfer of technology options are also to be adopted.

Soil Degradation

Soil erosion is one of the major factor affecting the soil health in the microwatershed. About 228 ha (58%) area is suffering from moderate erosion. These areas need immediate soil and water conservation and, other land development and land husbandry practices for restoring soil health.

Dissemination of Information and Communication of Benefits

Any large scale implementation of soil health management requires that supporting information is made available widely, particularly through channels familiar to farmers and extension workers. Given the very high priority attached to soil health especially by the Central Government on issuing Soil-Health Cards to all the farmers, media outlets like Regional, State and National Newspapers, Radio and Dooradarshan programs in local languages but also modern information and communication technologies such as Cellular phones and the Internet, which can be much more effective in reaching the younger farmers.

Inputs for Net Planning (Saturation Plan) and Interventions needed

Net planning in IWMP is focusing on preparation of

1. Soil and Water Conservation Treatment Plans for each plot or farm.
2. Productivity enhancement measures/ interventions for existing crops/livestock/other farm enterprises.
3. Diversification of farming mainly with perennial horticultural crops and livestock.
4. Improving livelihood opportunities and income generating activities.

In this connection, how various outputs of Sujala-III are of use in addressing these objectives of Net Planning are briefly presented below.

- ❖ **Soil Depth:** The depth of a soil decides the amount of moisture and nutrients it can hold, what crops can be taken up or not, depending on the rooting depth and the length of growing period available for raising any crop. Deeper the soil, better for a wide variety of crops. If sufficient depth is not available for growing deep rooted crops, either choose medium or short duration crops or deeper planting pits need to be opened and additional good quality soil brought from outside has to be filled into the planting pits.
- ❖ **Surface soil texture:** Lighter soil texture in the top soil means, better rain water infiltration, less run-off and soil moisture conservation, less capillary rise and less evaporation losses. Lighter surface textured soils are amenable to good soil tilth and are highly suitable for crops like groundnut, root vegetables (carrot, raddish, potato etc) but not ideal for crops that need stagnant water like lowland paddy. Heavy textured soils are poor in water infiltration and percolation. They are prone for sheet erosion; such soils can be improved by sand mulching. The technology that is developed by the AICRP-Dryland Agriculture, Vijayapura, Karnataka can be adopted.
- ❖ **Gravelliness:** More gravel content is favorable for run-off harvesting but poor in soil moisture storage and nutrient availability. It is a significant parameter that decides the kind of crop to be raised.
- ❖ **Land Capability Classification:** The land capability map shows the areas suitable and not suitable for agriculture and the major constraints in each of the plot/survey number. Hence, one can decide what kind of enterprise is possible in each of these

units. In general, erosion and soil are the major constraints in Bettageri-1 Microwatershed.

- ❖ **Organic Carbon:** The OC content is medium (0.5-0.75%) in an area of about 152 ha (20%), low in an area of (<0.5%) in 79 ha (20%) and high (>0.5%) in about 124 ha (32%). The areas that are low and medium in OC needs to be further improved by applying farmyard manure and rotating crops with cereals and legumes or mixed cropping.
- ❖ **Promoting green manuring:** Growing of green manuring crops costs Rs. 1250/ha (green manuring seeds) and about Rs. 2000/ha towards cultivation that totals to Rs. 3250/- per ha. On the other hand, application of organic manure @ 10 tons/ha costs Rs. 5000/ha. The practice needs to be continued for 2-3 years or more. Nitrogen fertilizer needs to be supplemented by 25% in addition to the recommended level in 79 ha area where OC is less than 0.5% and 152 ha area medium (0.5-0.75%). For example, for rainfed maize, recommended level is 50 kg N per ha and an additional 12 kg /ha needs to be applied for all the crops grown in these plots.
- ❖ **Available Phosphorus:** An area of about 24 ha (6%) is low (<23 kg/ha) and medium in 277 ha (71%) (23-57 kg/ha) area in available phosphorus. Hence for all the crops, 25% additional P-needs to be applied and high (>57 kg/ha) in small area of 54 ha (14%). For all crops 25% less phosphorus needs to be applied in areas where it is high.
- ❖ **Available Potassium:** Available potassium is high in small area of 18 ha (5%) for these areas 25% less potassium needs to be applied and medium (145-337 kg/ha) in major area of 337 ha (86%) the microwatershed. For all crops, 25 % less potassium may be applied in areas where it is medium or low.
- ❖ **Available Sulphur:** Available sulphur is a very critical nutrient for oilseed crops. Available sulphur is low (<10 ppm) in 259 ha (66%) area and medium in an area of about 96 ha (24%) in the microwatershed. These areas need to be applied with magnesium sulphate or gypsum or Factamphos (p) fertilizer (13% sulphur) for 2-3 years for the deficiency to be corrected. It is high in minute area of the microwatershed.
- ❖ **Available Boron:** Major area of about 233 ha (60%) is low (<0.5 ppm) in available boron and an area of 121 ha (31%) is medium (0.5 -1.0 ppm) in available boron content. These areas need to be applied with sodium borate @ 10kg/ha as a soil application or 0.2% borax as foliar spray to correct the deficiency
- ❖ **Available iron:** It is sufficient (>4.5 ppm) in 155 ha (40%) and deficient (<4.5 ppm) in 200 ha (51%) the entire microwatershed for deficiency areas , iron sulphate @ 25 kg/ha needs to be applied for 2-3 years correct the deficiency.
- ❖ **Available Zinc:** It is deficient (<0.6 ppm) in an area of 139 ha (36%). Application of zinc sulphate @ 25kg/ha is to be followed in areas that are deficient in available zinc and 215 ha (55%) area is sufficient (>0.6 ppm) in available zinc.


❖ **Soil alkalinity:** About 261 ha (67%) area in the microwatershed has soils that are slightly to very strongly alkaline. These areas need application of gypsum and wherever calcium is in excess, iron pyrites and element sulphur can be recommended. Management practices like treating repeatedly with good quality water to drain out the excess salts and provision of subsurface drainage and growing of salt tolerant crops like Casuarina, Acasia, Neem, Ber etc, are recommended.

Land Suitability for various crops: Areas that are highly, moderately and marginally suitable for growing various crops are indicated. Along with the suitability, various constraints that are limiting the productivity are also indicated. For example, in case of cotton, gravel content, rooting depth and salinity/alkalinity are the major constraints in various plots. With suitable management interventions, the productivity can be enhanced. In order to increase water holding capacity of light textured soils, growing of green manure crops and application of organic manure is recommended.

SOIL AND WATER CONSERVATION TREATMENT PLAN

For preparing soil and water conservation treatment plan for Bettageri-1 Microwatershed, the land resource inventory database generated under Sujala-III project has been transformed as information through series of interpretative (thematic) maps using soil phase map as a base. The various thematic maps (1:7920 scale) generated were

- Soil depth
 - Surface soil texture
 - Available water capacity
 - Soil slope
 - Soil gravelliness
 - Land capability
 - Present land use and land cover
 - Crop suitability maps
 - Rainfall map
 - Hydrology
 - Water Resources
 - Socio-economic data
 - Contour plan with existing features- network of waterways, pottissa boundaries, cut up/ minor terraces etc.
 - Cadastral map (1:7920 scale)
 - Satellite imagery (1:7920 scale)
- Apart from these, Hand Level/ Hydro Marker/ Dumpy Level/ Total Station and Kathedars' List needs to be collected.



Steps for Survey and Preparation of Treatment Plan

The boundaries of Land User Groups' and Survey No. boundaries are traced in the field.

- Naming of user groups and farmers
- Identification of arable and non arable lands
- Identification of drainage lines and gullies
- Identification of non treatable areas
- Identification of priority areas in the arable lands
- Treatment plan for arable lands
- Location of water harvesting and recharge structures

9.1 Treatment Plan

The treatment plan recommended for arable lands is briefly described below.

9.1.1 Arable Land Treatment

A. BUNDING

Steps for Survey and Preparation of Treatment Plan		USER GROUP-1 CLASSIFICATION OF GULLIES
Cadastral map (1:7920 scale) is enlarged to a scale of 1:2500 scale		
Existing network of waterways, pothissa boundaries, grass belts, natural drainage lines/ watercourse, cut ups/ terraces are marked on the cadastral map to the scale		
Drainage lines are demarcated into		
Small gullies	(up to 5 ha catchment)	
Medium gullies	(5-15 ha catchment)	
Ravines	(15-25 ha catchment) and	
<i>Halla/Nala</i>	(more than 25ha catchment)	

Measurement of Land Slope

Land slope is estimated or determined by the study and interpretation of contours or by measurement in the field using simple instruments like Hand Level or Hydromarker.



Vertical and Horizontal intervals between bunds as recommended by the Watershed Development Department.

Slope percentage	Vertical interval (m)	Corresponding Horizontal Distance (m)
2 - 3%	0.6	24
3 - 4%	0.9	21
4 - 5%	0.9	21
5 - 6%	1.2	21
6 - 7%	1.2	21

Note: i) The above intervals are maximum.

(ii) Considering the slope class and erosion status (A1... A= 0-1% slope, 1= slight erosion.) the intervals have to be decided.

Bund length recording: Considering the contour plan and the existing grass belts/partitions, the bunds are aligned and lengths are measured.

Section of the Bund

Bund section is decided considering the soil texture class and gravelliness class (bg₀b= loamy sand, g₀ = <15% gravel). The recommended sections for different soils are given below.

Recommended Bund Section



Top width (m)	Base width (m)	Height (m)	Side slope (Z:1;H:V)	Cross section (sq m)	Soil Texture	Remarks
0.3	0.9	0.3	01:01	0.18	Sandy loam	Vegetative bund
0.3	1.2	0.3	1.5:1	0.225	Sandy clay	
0.3	1.2	0.5	0.9:1	0.375	Red gravelly soils	
0.3	1.2	0.6	0.75:1	0.45		
0.3	1.5	0.6	01:01	0.54	Red sandy loam	
0.3	2.1	0.6	1.5:1	0.72	Very shallow black clayey soils	
0.45	2	0.75	01:01	0.92		
0.45	2.4	0.75	1.3:1	1.07	Shallow black clayey soils	
0.6	3.1	0.7	1.78:1	1.29	Medium black clayey soils	
0.5	3	0.85	1.47:1	1.49		

Formation of Trench cum Bund

Dimensions of the Borrow Pits/ Trenches to be excavated (machinery are decided considering the Bund Section).

Details of Borrow Pit dimensions are given below

TRENCH CUM BUND





WATER STORAGE AREA

0.45 Sq.m section

IDEAL FOR HORTICULTURE CROPS

'A' FRAME FOR INTERBUND MANAGEMENT



1. ಸಮಸಾರಾತಲ ಉಳುಮೆ
2. ಸಮಸಾರಾತಲ ಬಿತ್ತನೆ/ನಾಟಿ

Size of Borrow Pits/ Trench recommended for Trench cum Bund (by machinery)

Bund section	Bund length	Earth quantity	Pit				Berm (pit to pit)	Soil depth Class
			L(m)	W(m)	D(m)	QUANTITY (m ³)		
m ²	m	m ³					m	
0.375	6	2.25	5.85	0.85	0.45	2.24	0.15	Shallow
0.45	6	2.7	5.4	1.2	0.43	2.79	0.6	Shallow
0.45	6	2.7	5	0.85	0.65	2.76	1	Moderately Shallow
0.54	5.6	3.02	5.5	0.85	0.7	3.27	0.1	Moderately shallow
0.54	5.5	2.97	5	1.2	0.5	3	0.5	Shallow
0.72	6.2	4.46	6	1.2	0.7	5.04	0.2	Moderately shallow
0.72	5.2	3.74	5.1	0.85	0.9	3.9	0.1	Moderately deep

B. Waterways

- Existing waterways are marked on the cadastral map (1:7920 scale) and their dimensions are recorded.
- Considering the contour plan of the MWS, additional waterways/ modernization of the existing ones can be thought of.
- The design details are given in the Manual.

C. Farm Ponds

Waterways and the catchment area will give an indication on the size of the Farm Pond. Location of the pond can be decided based on the contour plan/ field condition and farmers' need/desire.

D. Diversion Channel

Existing EPT/ CPT are marked on the cadastral map. Looking to the need, these can be modernized or fresh diversion channel can be proposed and runoff from this can be stored in *Gokatte/* Recharge ponds.

9.1.2 Non-Arable Land Treatment

Depending on the gravelliness and crops preferred by the farmers, the concerned authorities can decide appropriate treatment plan. The recommended treatments may be Contour Trench, Staggered Trench, Crescent Bund, Boulder Bund or Pebble Bund are formed in the field.

9.1.3 Treatment of Natural Water Course/ Drainage Lines

- a) The cadastral map has to be updated as regards the network of drainage lines (gullies/ *nalas*/ *hallas*) and existing structures are marked to the scale and storage capacity of the existing water bodies are documented.
- b) The drainage line will be demarcated into Upper Reach, Middle Reach and Lower Reach.
- c) Considering the Catchment, *Nala* bed and bank conditions, suitable structures are decided.
- d) Number of storage structures (Check dam/ *Nala* bund/ Percolation tank) will be decided considering the commitments and available runoff in water budgeting and quality of water in the wells and site suitability.
- e) Detailed Leveling Survey using Dumpy Level / Total Station has to be carried out to arrive at the site-specific designs as shown in the Manual.
- f) The location of ground water recharge structures are decided by examining the lineaments and fracture zones from geological maps.
- g) Rainfall intensity data of the nearest Rain Gauge Station is considered for Hydrologic Designs.
- h) Silt load to the Storage/Recharge Structures is reduced by providing vegetative, boulder and earthen checks in the natural water course. Location and design details are given in the Manual.

9.2 Recommended Soil and Water Conservation Measures

The appropriate conservation structures best suited for each of the land parcel/ survey number (Appendix-I) are selected based on the slope per cent, severity of erosion, amount of rainfall, land use and soil type. The different kinds of conservation structures recommended are

1. Graded / Strengthening of Bunds
2. Trench cum Bunds (TCB)
3. Trench cum Bunds / Strengthening
4. Crescent Bunds

A map (Fig. 9.1) showing soil and water conservation plan with different kinds of structures recommended has been prepared which shows the spatial distribution and extent of area. About 233 ha (60%) area is trench cum bunding and 122 ha (31%) area is graded bunding. The conservation plan prepared may be presented to all the stakeholders including farmers and after considering their suggestions, the conservation plan for the microwatershed may be finalised in a participatory approach.




Fig. 9.1 Soil and Water Conservation Plan map of Bettageri-1 Microwatershed

9.3 Greening of Microwatershed

As part of the greening programme in the watersheds, it is envisaged to plant a variety of horticultural and other tree plants that are edible, economical and produce lot of biomass which helps to restore the ecological balance in the watersheds. The lands that are suitable for greening programme are non-arable lands (land capability classes V, VI VII and VIII) and also the lands that are not suitable or marginally suitable for growing annual and perennial crops. The method of planting these trees is given below.

It is recommended to open the pits during the 1st week of March along the contour and heap the dug out soil on the lower side of the slope in order to harness the flowing water and facilitate weathering of soil in the pit. Exposure of soil in the pit also prevents spread of pests and diseases due to scorching sun rays. The pits should be filled with mixture of soil and organic manure during the second week of April and keep ready with sufficiently tall seedlings produced either in poly bags or in root trainer nurseries so that planting can be done during the 2nd or 3rd week of April depending on the rainfall.

The tree species suitable for the area considering rainfall, temperature and adaptability is listed below; waterlogged areas are recommended to be planted with species like Neral (*Syzgium cumini*) and Bamboo. Dry areas are to be planted with species like Honge, Bevu, Seetaphal *etc.*

Dry Deciduous Species			Temp (°C)	Rainfall (mm)
1.	Bevu	<i>Azadiracta indica</i>	21–32	400 –1,200
2.	Tapasi	<i>Holoptelia integrifolia</i>	20-30	500 - 1000
3.	Seetaphal	<i>Anona Squamosa</i>	20-40	400 - 1000
4.	Honge	<i>Pongamia pinnata</i>	20 -50	500–2,500
5.	Kamara	<i>Hardwickia binata</i>	25 -35	400 - 1000
6.	Bage	<i>Albezzia lebbek</i>	20 - 45	500 - 1000
7.	Ficus	<i>Ficus bengalensis</i>	20 - 50	500–2,500
8.	Sisso	<i>Dalbargia Sissoo</i>	20 - 50	500 -2000
9.	Ailanthus	<i>Ailanthus excelsa</i>	20 - 50	500 - 1000
10.	Hale	<i>Wrightia tinctoria</i>	25 - 45	500 - 1000
11.	Uded	<i>Steriospermum chelanoides</i>	25 - 45	500 -2000
12.	Dhupa	<i>Boswella Serrata</i>	20 - 40	500 - 2000
13.	Nelli	<i>Emblia Officinalis</i>	20 - 50	500 -1500
14.	Honne	<i>Pterocarpus marsupium</i>	20 - 40	500 - 2000
Moist Deciduous Species			Temp (°C)	Rainfall (mm)
15.	Teak	<i>Tectona grandis</i>	20 - 50	500-5000
16.	Nandi	<i>Legarstroemia lanceolata</i>	20 - 40	500 - 4000
17.	Honne	<i>Pterocarpus marsupium</i>	20 - 40	500 - 3000
18.	Mathi	<i>Terminalia alata</i>	20 -50	500 - 2000
19.	Shivane	<i>Gmelina arborea</i>	20 -50	500 -2000
20.	Kindal	<i>T.Paniculata</i>	20 - 40	500 - 1500
21.	Beete	<i>Dalbargia latifolia</i>	20 - 40	500 - 1500
22.	Tare	<i>T. belerica</i>	20 - 40	500 - 2000
23.	Bamboo	<i>Bambusa arundinasia</i>	20 - 40	500 - 2500
24.	Bamboo	<i>Dendrocalamus strictus</i>	20 – 40	500 – 2500
25.	Muthuga	<i>Butea monosperma</i>	20 - 40	400 - 1500
26.	Hippe	<i>Madhuca latifolia</i>	20 - 40	500 - 2000
27.	Sandal	<i>Santalum album</i>	20 - 50	400 - 1000
28.	Nelli	<i>Emblia officinalis</i>	20 - 40	500 - 2000
29.	Nerale	<i>Sizyium cumini</i>	20 - 40	500 - 2000
30.	Dhaman	<i>Grevia tilifolia</i>	20 - 40	500 - 2000
31.	Kaval	<i>Careya arborea</i>	20 - 40	500 - 2000
32.	Harada	<i>Terminalia chebula</i>	20 - 40	500 - 2000

References

1. FAO (1976) Framework for Land Evaluation, Food and Agriculture Organization, Rome.72 pp.
2. FAO (1983) Guidelines for Land Evaluation for Rainfed Agriculture, FAO, Rome, 237 pp.
3. IARI (1971) Soil Survey Manual, All India Soil and Land Use Survey Organization, IARI, New Delhi, 121 pp.
4. Katyal, J.C. and Rattan, R.K. (2003) Secondary and Micronutrients; Reaserch Gap and future needs. Fert. News 48 (4); 9-20.
5. Naidu, L.G.K., Ramamurthy, V., Challa, O., Hegde, R. and Krishnan, P. (2006) Manual Soil Site Suitability Criteria for Major Crops, NBSS Publ. No. 129, NBSS &LUP, Nagpur, 118 pp.
6. Natarajan, A. and Dipak Sarkar (2010) Field Guide for Soil Survey, National Bureau of Soil Survey and Land Use Planning (ICAR), Nagpur, India.
7. Natarajan, A., Rajendra Hegde, Raj, J.N. and Shivananda Murthy, H.G. (2015) Implementation Manual for Sujala-III Project, Watershed Development Department, Bengaluru, Karnataka.
8. Sarma, V.A.K., Krishnan, P. and Budihal, S.L. (1987) Laboratory Manual, Tech. Bull. 23, NBSS &LUP, Nagpur.
9. Sehgal, J.L. (1990) Soil Resource Mapping of Different States of India; Why and How?, National Bureau of Soil Survey and Land Use Planning, Nagpur, 49 pp.
10. Shivaprasad, C.R., R.S. Reddy, J. Sehgal and M. Velayuthum (1998) Soils of Karntaka for Optimising Land Use, NBSS Publ. No. 47b, NBSS & LUP, Nagpur, India.
11. Soil Survey Staff (2006) Keys to Soil Taxonomy, Tenth edition, U.S. Department of Agriculture/ NRCS, Washington DC, U.S.A.
12. Soil Survey Staff (2012) Soil Survey Manual, Handbook No. 18, USDA, Washington DC, USA.

Appendix I
Betageri-1 Microwatershed
Soil Phase Information

Village	Survey No.	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	WELLS	Land Capability	Conservation Plan
Betageri	1	11.33	Habitation	Habitation	Habitation	Habitation	Habitation	Habitation	Habitation	Habitation	Habitation	Habitation	Habitation	Habitation
Betageri	2	3.32	MKHiB2g2	LMU-4	Moderately shallow (50-75 cm)	Sandy clay	Very gravelly (35-60%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Bajra (Bj)	Not Available	Illes	Trench cum bunding
Betageri	3	3.29	MKHiB2g2	LMU-4	Moderately shallow (50-75 cm)	Sandy clay	Very gravelly (35-60%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	Illes	Trench cum bunding
Betageri	4	0.83	MKHiB2g2	LMU-4	Moderately shallow (50-75 cm)	Sandy clay	Very gravelly (35-60%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	Illes	Trench cum bunding
Betageri	5	1.23	MKHiB2g2	LMU-4	Moderately shallow (50-75 cm)	Sandy clay	Very gravelly (35-60%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	Illes	Trench cum bunding
Betageri	6	0.01	MKHiB2g2	LMU-4	Moderately shallow (50-75 cm)	Sandy clay	Very gravelly (35-60%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Bajra (Bj)	Not Available	Illes	Trench cum bunding
Betageri	8	0.82	MKHiB2g2	LMU-4	Moderately shallow (50-75 cm)	Sandy clay	Very gravelly (35-60%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	Illes	Trench cum bunding
Betageri	9	0.6	MKHiB2g2	LMU-4	Moderately shallow (50-75 cm)	Sandy clay	Very gravelly (35-60%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow (Cf)	Not Available	Illes	Trench cum bunding
Betageri	10	0	GRHmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	IIs	Graded bunding
Betageri	109	0.01	GRHmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Current fallow (Cf)	Not Available	IIs	Graded bunding
Betageri	110	1.76	GRHmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Bajra (Bj)	Not Available	IIs	Graded bunding
Betageri	111	7.09	GRHmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Current fallow+Fallow land (Cf+Fl)	Not Available	IIs	Graded bunding
Betageri	112	7.42	GRHmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Current fallow (Cf)	Not Available	IIs	Graded bunding
Betageri	113	0	GRHmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Current fallow+Maize (Cf+Mz)	Not Available	IIs	Graded bunding
Betageri	200	0.31	GRHmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Current fallow+Bajra (Cf+Bj)	Not Available	IIs	Graded bunding
Betageri	201	2.18	GRHmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Fallow land (Fl)	Not Available	IIs	Graded bunding
Betageri	265	0.88	Habitation	Habitation	Habitation	Habitation	Habitation	Habitation	Habitation	Habitation	Habitation	Habitation	Habitation	Habitation
Betageri	266	3.09	Habitation	Habitation	Habitation	Habitation	Habitation	Habitation	Habitation	Habitation	Habitation	Habitation	Habitation	Habitation
Betageri	312	0.09	MTLiB2g2	LMU-6	Shallow (25-50 cm)	Sandy clay	Very gravelly (35-60%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Fallow land+Maize (Fl+Mz)	Not Available	Illes	Graded bunding
Betageri	313	0.07	RNKmB2g1	LMU-5	Moderately shallow (50-75 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Maize+Eucalyptus (Mz+Eu)	Not Available	IIs	Graded bunding
Betageri	314	1.1	RNKmB2g1	LMU-5	Moderately shallow (50-75 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIs	Graded bunding
Betageri	316	0.32	MTLiB2g2	LMU-6	Shallow (25-50 cm)	Sandy clay	Very gravelly (35-60%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Bajra+Maize (Bj+Mz)	Not Available	Illes	Graded bunding
Betageri	317	0.44	MTLiB2g2	LMU-6	Shallow (25-50 cm)	Sandy clay	Very gravelly (35-60%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	Illes	Graded bunding

Village	Survey No.	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	WELLS	Land Capability	Conservation Plan
Betageri	318	0.73	MTLiB2g2	LMU-6	Shallow (25-50 cm)	Sandy clay	Very gravelly (35-60%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Bajra (Bj)	Not Available	IIles	Graded bunding
Betageri	319	1.09	MTLiB2g2	LMU-6	Shallow (25-50 cm)	Sandy clay	Very gravelly (35-60%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Sunflower+Maize (Sf+Mz)	Not Available	IIles	Graded bunding
Betageri	320	1.53	MTLiB2g2	LMU-6	Shallow (25-50 cm)	Sandy clay	Very gravelly (35-60%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIles	Graded bunding
Betageri	321	2.97	MTLiB2g2	LMU-6	Shallow (25-50 cm)	Sandy clay	Very gravelly (35-60%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Bajra+Sunflower+Maize (Bj+Sf+Mz)	Not Available	IIles	Graded bunding
Betageri	322	2.28	MTLiB2g2	LMU-6	Shallow (25-50 cm)	Sandy clay	Very gravelly (35-60%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIles	Graded bunding
Betageri	323	5.21	MTLiB2g2	LMU-6	Shallow (25-50 cm)	Sandy clay	Very gravelly (35-60%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Maize+Bajra (Mz+Bj)	Not Available	IIles	Graded bunding
Betageri	324	5.72	MTLiB2g2	LMU-6	Shallow (25-50 cm)	Sandy clay	Very gravelly (35-60%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIles	Graded bunding
Betageri	325	7.14	MTLiB2g2	LMU-6	Shallow (25-50 cm)	Sandy clay	Very gravelly (35-60%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Bajra+Maize (Bj+Mz)	Not Available	IIles	Graded bunding
Betageri	326	3.56	MTLiB2g2	LMU-6	Shallow (25-50 cm)	Sandy clay	Very gravelly (35-60%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Maize+Bajra (Mz+Bj)	Not Available	IIles	Graded bunding
Betageri	327	9.16	MTLiB2g2	LMU-6	Shallow (25-50 cm)	Sandy clay	Very gravelly (35-60%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Maize+Bajra (Mz+Bj)	Not Available	IIles	Graded bunding
Betageri	328	1.57	MTLiB2g2	LMU-6	Shallow (25-50 cm)	Sandy clay	Very gravelly (35-60%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Bajra (Bj)	Not Available	IIles	Graded bunding
Betageri	335	0.13	MTLiB2g2	LMU-6	Shallow (25-50 cm)	Sandy clay	Very gravelly (35-60%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow (Cf)	Not Available	IIles	Graded bunding
Betageri	336	9.4	MTLiB2g2	LMU-6	Shallow (25-50 cm)	Sandy clay	Very gravelly (35-60%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIles	Graded bunding
Betageri	337	5.57	MTLiB2g2	LMU-6	Shallow (25-50 cm)	Sandy clay	Very gravelly (35-60%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Bajra+Maize (Bj+Mz)	Not Available	IIles	Graded bunding
Betageri	338	5.94	MTLiB2g2	LMU-6	Shallow (25-50 cm)	Sandy clay	Very gravelly (35-60%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Maize+Sunflower+Bajra (Mz+Sf+Bj)	Not Available	IIles	Graded bunding
Betageri	339	2.62	MTLiB2g2	LMU-6	Shallow (25-50 cm)	Sandy clay	Very gravelly (35-60%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIles	Graded bunding
Betageri	340	1.72	MTLiB2g2	LMU-6	Shallow (25-50 cm)	Sandy clay	Very gravelly (35-60%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Bajra (Bj)	Not Available	IIles	Graded bunding
Betageri	341	4.82	MTLiB2g2	LMU-6	Shallow (25-50 cm)	Sandy clay	Very gravelly (35-60%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Bajra+Drumstick (Bj+Ds)	Not Available	IIles	Graded bunding
Betageri	342	1.39	MTLiB2g2	LMU-6	Shallow (25-50 cm)	Sandy clay	Very gravelly (35-60%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow (Cf)	Not Available	IIles	Graded bunding
Betageri	343	2.54	MTLiB2g2	LMU-6	Shallow (25-50 cm)	Sandy clay	Very gravelly (35-60%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Bajra (Bj)	Not Available	IIles	Graded bunding
Betageri	344	1.67	MTLiB2g2	LMU-6	Shallow (25-50 cm)	Sandy clay	Very gravelly (35-60%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIles	Graded bunding
Betageri	345	5.25	CKMhB1g1	LMU-3	Moderately deep (75-100 cm)	Sandy clay loam	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Slight	Maize+Groundnut (Mz+Gn)	Not Available	IIs	Trench cum bunding
Betageri	346	4.74	CKMhB1g1	LMU-3	Moderately deep (75-100 cm)	Sandy clay loam	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Slight	Bajra+Maize (Bj+Mz)	Not Available	IIs	Trench cum bunding
Betageri	347	5.82	MTLiB2g2	LMU-6	Shallow (25-50 cm)	Sandy clay	Very gravelly (35-60%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Bajra+Maize (Bj+Mz)	Not Available	IIles	Graded bunding

Village	Survey No.	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	WELLS	Land Capability	Conservation Plan
Betageri	348	5.33	CKMhB1g1	LMU-3	Moderately deep (75-100 cm)	Sandy clay loam	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Slight	Maize+Bajra+Jowar (Mz+Bj+Jw)	Not Available	IIs	Trench cum bunding
Betageri	349	3.47	CKMhB1g1	LMU-3	Moderately deep (75-100 cm)	Sandy clay loam	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Slight	Maize+Bajra (Mz+Bj)	Not Available	IIs	Trench cum bunding
Betageri	350	0.07	CKMhB1g1	LMU-3	Moderately deep (75-100 cm)	Sandy clay loam	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	IIs	Trench cum bunding
Betageri	351	2.49	CKMhB1g1	LMU-3	Moderately deep (75-100 cm)	Sandy clay loam	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	IIs	Trench cum bunding
Betageri	352	2.37	CKMhB1g1	LMU-3	Moderately deep (75-100 cm)	Sandy clay loam	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	IIs	Trench cum bunding
Betageri	353	6.2	CKMhB1g1	LMU-3	Moderately deep (75-100 cm)	Sandy clay loam	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	IIs	Trench cum bunding
Betageri	354	1.48	CKMhB1g1	LMU-3	Moderately deep (75-100 cm)	Sandy clay loam	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Slight	Bajra (Bj)	Not Available	IIs	Trench cum bunding
Betageri	355	3.78	CKMhB1g1	LMU-3	Moderately deep (75-100 cm)	Sandy clay loam	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Slight	Bajra+Maize (Bj+Mz)	Not Available	IIs	Trench cum bunding
Betageri	356	2.35	CKMhB1g1	LMU-3	Moderately deep (75-100 cm)	Sandy clay loam	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Slight	Bajra (Bj)	Not Available	IIs	Trench cum bunding
Betageri	357	1.67	CKMhB1g1	LMU-3	Moderately deep (75-100 cm)	Sandy clay loam	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	IIs	Trench cum bunding
Betageri	358	1.87	CKMhB1g1	LMU-3	Moderately deep (75-100 cm)	Sandy clay loam	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Slight	Bajra (Bj)	Not Available	IIs	Trench cum bunding
Betageri	359	4.24	CKMhB1g1	LMU-3	Moderately deep (75-100 cm)	Sandy clay loam	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Slight	Maize+Bajra (Mz+Bj)	Not Available	IIs	Trench cum bunding
Betageri	360	4.48	CKMhB1g1	LMU-3	Moderately deep (75-100 cm)	Sandy clay loam	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Slight	Maize+Bajra (Mz+Bj)	Not Available	IIs	Trench cum bunding
Betageri	361	3.44	MTLiB2g2	LMU-6	Shallow (25-50 cm)	Sandy clay	Very gravelly (35-60%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIIs	Graded bunding
Betageri	362	3.23	MTLiB2g2	LMU-6	Shallow (25-50 cm)	Sandy clay	Very gravelly (35-60%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIIs	Graded bunding
Betageri	363	4.32	MTLiB2g2	LMU-6	Shallow (25-50 cm)	Sandy clay	Very gravelly (35-60%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIIs	Graded bunding
Betageri	364	4.72	MTLiB2g2	LMU-6	Shallow (25-50 cm)	Sandy clay	Very gravelly (35-60%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIIs	Graded bunding
Betageri	365	7.81	CKMhB1g1	LMU-3	Moderately deep (75-100 cm)	Sandy clay loam	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Slight	Maize+Bajra (Mz+Bj)	Not Available	IIs	Trench cum bunding
Betageri	366	3.31	CKMhB1g1	LMU-3	Moderately deep (75-100 cm)	Sandy clay loam	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Slight	Bajra (Bj)	Not Available	IIs	Trench cum bunding
Betageri	367	2.34	CKMhB1g1	LMU-3	Moderately deep (75-100 cm)	Sandy clay loam	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Slight	Bajra (Bj)	Not Available	IIs	Trench cum bunding
Betageri	368	1.82	Habitation	Habitation	Habitation	Habitation	Habitation	Habitation	Habitation	Habitation	Habitation	Habitation	Habitation	Habitation
Betageri	369	1.11	Habitation	Habitation	Habitation	Habitation	Habitation	Habitation	Habitation	Habitation	Habitation	Habitation	Habitation	Habitation
Betageri	370	0.56	Habitation	Habitation	Habitation	Habitation	Habitation	Habitation	Habitation	Habitation	Habitation	Habitation	Habitation	Habitation
Betageri	371	1.48	Habitation	Habitation	Habitation	Habitation	Habitation	Habitation	Habitation	Habitation	Habitation	Habitation	Habitation	Habitation
Betageri	372	3.63	CKMhB1g1	LMU-3	Moderately deep (75-100 cm)	Sandy clay loam	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	IIs	Trench cum bunding
Betageri	373	2.79	CKMhB1g1	LMU-3	Moderately deep (75-100 cm)	Sandy clay loam	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Slight	Bajra (Bj)	Not Available	IIs	Trench cum bunding

Village	Survey No.	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	WELLS	Land Capability	Conservation Plan
Betageri	374	1.61	CKMhB1g1	LMU-3	Moderately deep (75-100 cm)	Sandy clay loam	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	IIs	Trench cum bunding
Betageri	375	4.37	CKMhB1g1	LMU-3	Moderately deep (75-100 cm)	Sandy clay loam	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Slight	Bajra+Maize (Bj+Mz)	Not Available	IIs	Trench cum bunding
Betageri	376	2.45	CKMhB1g1	LMU-3	Moderately deep (75-100 cm)	Sandy clay loam	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	1 Bore well	IIs	Trench cum bunding
Betageri	377	6.2	MKHiB2g2	LMU-4	Moderately shallow (50-75 cm)	Sandy clay	Very gravelly (35-60%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Maize+Current fallow (Mz+Cf)	Not Available	IIIs	Trench cum bunding
Betageri	378	4.63	MKHiB2g2	LMU-4	Moderately shallow (50-75 cm)	Sandy clay	Very gravelly (35-60%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIIs	Trench cum bunding
Betageri	379	8.47	MKHiB2g2	LMU-4	Moderately shallow (50-75 cm)	Sandy clay	Very gravelly (35-60%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Bajra+Maize (Bj+Mz)	Not Available	IIIs	Trench cum bunding
Betageri	380	2.23	MKHiB2g2	LMU-4	Moderately shallow (50-75 cm)	Sandy clay	Very gravelly (35-60%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Bajra (Bj)	Not Available	IIIs	Trench cum bunding
Betageri	381	1.62	MKHiB2g2	LMU-4	Moderately shallow (50-75 cm)	Sandy clay	Very gravelly (35-60%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Bajra (Bj)	Not Available	IIIs	Trench cum bunding
Betageri	382	4.53	MTLiB2g2	LMU-6	Shallow (25-50 cm)	Sandy clay	Very gravelly (35-60%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Bajra+Current fallow (Bj+Cf)	Not Available	IIIs	Graded bunding
Betageri	383	7.07	MKHiB2g2	LMU-4	Moderately shallow (50-75 cm)	Sandy clay	Very gravelly (35-60%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Bajra+Maize+Current fallow (Bj+Mz+Cf)	Not Available	IIIs	Trench cum bunding
Betageri	384	1.4	MKHiB2g2	LMU-4	Moderately shallow (50-75 cm)	Sandy clay	Very gravelly (35-60%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Maize+Bajra+Current fallow (Mz+Bj+Cf)	Not Available	IIIs	Trench cum bunding
Betageri	428	1.33	BDGhB2g1	LMU-2	Moderately deep (75-100 cm)	Sandy clay loam	Gravelly (15-35%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Bajra (Bj)	Not Available	IIIs	Trench cum bunding
Betageri	429	2.59	BDGhB2g1	LMU-2	Moderately deep (75-100 cm)	Sandy clay loam	Gravelly (15-35%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIIs	Trench cum bunding
Betageri	433	2.91	BDGhB2g1	LMU-2	Moderately deep (75-100 cm)	Sandy clay loam	Gravelly (15-35%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Maize+Bajra (Mz+Bj)	Not Available	IIIs	Trench cum bunding
Betageri	434	2.85	BDGhB2g1	LMU-2	Moderately deep (75-100 cm)	Sandy clay loam	Gravelly (15-35%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Bajra+Current fallow (Bj+Cf)	Not Available	IIIs	Trench cum bunding
Betageri	435	3.17	MKHiB2g2	LMU-4	Moderately shallow (50-75 cm)	Sandy clay	Very gravelly (35-60%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIIs	Trench cum bunding
Betageri	436	3.79	BDGhB2g1	LMU-2	Moderately deep (75-100 cm)	Sandy clay loam	Gravelly (15-35%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut+Maize (Gn+Mz)	Not Available	IIIs	Trench cum bunding
Betageri	437	2.81	BDGhB2g1	LMU-2	Moderately deep (75-100 cm)	Sandy clay loam	Gravelly (15-35%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIIs	Trench cum bunding
Betageri	438	4.51	BDGhB2g1	LMU-2	Moderately deep (75-100 cm)	Sandy clay loam	Gravelly (15-35%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Maize+Bajra (Mz+Bj)	1 Bore well	IIIs	Trench cum bunding
Betageri	439	3.51	BDGhB2g1	LMU-2	Moderately deep (75-100 cm)	Sandy clay loam	Gravelly (15-35%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIIs	Trench cum bunding
Betageri	440	8.48	BDGhB2g1	LMU-2	Moderately deep (75-100 cm)	Sandy clay loam	Gravelly (15-35%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Maize+Current fallow (Mz+Cf)	Not Available	IIIs	Trench cum bunding
Betageri	441	5.04	BDGhB2g1	LMU-2	Moderately deep (75-100 cm)	Sandy clay loam	Gravelly (15-35%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Bajra+Maize+Current fallow (Bj+Mz+Cf)	1 Bore well	IIIs	Trench cum bunding
Betageri	442	3.07	BDGhB2g1	LMU-2	Moderately deep (75-100 cm)	Sandy clay loam	Gravelly (15-35%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Maize+Onion (Mz+On)	1 Bore well	IIIs	Trench cum bunding
Betageri	443	2.97	HRVhB1g2	LMU-7	Shallow (25-50 cm)	Sandy clay loam	Very gravelly (35-60%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Slight	Bajra (Bj)	Not Available	IIIs	Trench cum bunding

Village	Survey No.	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	WELLS	Land Capability	Conservation Plan
Betageri	444	8.77	HRVhB1g2	LMU-7	Shallow (25-50 cm)	Sandy clay loam	Very gravelly (35-60%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Slight	Maize+Groundnut (Mz+Gn)	Not Available	IIIs	Trench cum bunding
Betageri	445	1.55	HRVhB1g2	LMU-7	Shallow (25-50 cm)	Sandy clay loam	Very gravelly (35-60%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Slight	Bajra (Bj)	Not Available	IIIs	Trench cum bunding
Betageri	446	1.07	HRVhB1g2	LMU-7	Shallow (25-50 cm)	Sandy clay loam	Very gravelly (35-60%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Slight	Sunflower (Sf)	Not Available	IIIs	Trench cum bunding
Betageri	447	0.87	BDGhB2g1	LMU-2	Moderately deep (75-100 cm)	Sandy clay loam	Gravelly (15-35%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Maize+Sunflower+Groundnut+Bajra (Mz+Sf+Gn+Bj)	Not Available	IIIes	Trench cum bunding
Betageri	448	1.29	BDGhB2g1	LMU-2	Moderately deep (75-100 cm)	Sandy clay loam	Gravelly (15-35%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIIes	Trench cum bunding
Betageri	449	2.25	BDGhB2g1	LMU-2	Moderately deep (75-100 cm)	Sandy clay loam	Gravelly (15-35%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIIes	Trench cum bunding
Betageri	450	7.89	BDGhB2g1	LMU-2	Moderately deep (75-100 cm)	Sandy clay loam	Gravelly (15-35%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Bajra (Bj)	Not Available	IIIes	Trench cum bunding
Betageri	451	1.84	BDGhB2g1	LMU-2	Moderately deep (75-100 cm)	Sandy clay loam	Gravelly (15-35%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut (Gn)	Not Available	IIIes	Trench cum bunding
Betageri	452	1.7	BDGhB2g1	LMU-2	Moderately deep (75-100 cm)	Sandy clay loam	Gravelly (15-35%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIIes	Trench cum bunding
Betageri	453	4.37	BDGhB2g1	LMU-2	Moderately deep (75-100 cm)	Sandy clay loam	Gravelly (15-35%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Maize+Bajra+Groundnut (Mz+Bj+Gn)	Not Available	IIIes	Trench cum bunding
Betageri	454	2.52	BDGhB2g1	LMU-2	Moderately deep (75-100 cm)	Sandy clay loam	Gravelly (15-35%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Bajra (Bj)	Not Available	IIIes	Trench cum bunding
Betageri	455	4.16	BDGhB2g1	LMU-2	Moderately deep (75-100 cm)	Sandy clay loam	Gravelly (15-35%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIIes	Trench cum bunding
Betageri	510	0.13	BDGhB2g1	LMU-2	Moderately deep (75-100 cm)	Sandy clay loam	Gravelly (15-35%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Bajra (Bj)	Not Available	IIIes	Trench cum bunding
Betageri	513	0.02	BDGhB2g1	LMU-2	Moderately deep (75-100 cm)	Sandy clay loam	Gravelly (15-35%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Bajra (Bj)	Not Available	IIIes	Trench cum bunding
Betageri	583	1.66	HRVhB1g2	LMU-7	Shallow (25-50 cm)	Sandy clay loam	Very gravelly (35-60%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Slight	Bajra+Maize (Bj+Mz)	Not Available	IIIs	Trench cum bunding
Betageri	584	0.88	HRVhB1g2	LMU-7	Shallow (25-50 cm)	Sandy clay loam	Very gravelly (35-60%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Slight	Bajra+Maize (Bj+Mz)	Not Available	IIIs	Trench cum bunding
Betageri	588	0.49	MKHiB2g2	LMU-4	Moderately shallow (50-75 cm)	Sandy clay	Very gravelly (35-60%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIIes	Trench cum bunding
Betageri	589	0.86	TGRiB2g1	LMU-3	Moderately deep (75-100 cm)	Sandy clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIs	Trench cum bunding
Betageri	590	3.15	TGRiB2g1	LMU-3	Moderately deep (75-100 cm)	Sandy clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Bajra (Bj)	Not Available	IIs	Trench cum bunding
Betageri	591	7.2	TGRiB2g1	LMU-3	Moderately deep (75-100 cm)	Sandy clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Maize+Bajra+Cucumber (Mz+Bj+Cu)	1 Bore well	IIs	Trench cum bunding
Betageri	592	2.77	TGRiB2g1	LMU-3	Moderately deep (75-100 cm)	Sandy clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Bajra+Sunflower (Bj+Sf)	Not Available	IIs	Trench cum bunding

Appendix III
Betageri-1 Microwatershed
Soil Suitability Information

Village	Survey No.	Mango	Maize	Sapota	Sorgham	Guava	Cotton	Tamarind	Lime	Bengalgram	Sunflower	Redgram	Amla	Jackfruit	Custard apple	Cashew	Jamun	Musambi	Groundnut	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Jasmine	Crossandra	Drumstick	Mulbery	
Betageri	1	Habit	Habit	Habita	Habit	Habita	Habita	Habita	Habit	Habit	Habit	Habit	Habit	Habit	Habit	Habit	Habit	Habit	Habit	Habit	Habit	Habit	Habit	Habit	Habit	Habit	Habit	Habit	Habit	Habit
Betageri	2	N1r	S2rg	S3rg	S2rg	S3rg	S2rg	N1r	S3rg	S2rt	S3rg	S3rg	S2rg	S3rg	S2rg	S3rg	S3rg	S3rg	S2rg	S2rg	S2rg	S2rg	S2rg	S3rg	S2rg	S2rg	S2rg	S3rg	S3rg	
Betageri	3	N1r	S2rg	S3rg	S2rg	S3rg	S2rg	N1r	S3rg	S2rt	S3rg	S3rg	S2rg	S3rg	S2rg	S3rg	S3rg	S3rg	S2rg	S2rg	S2rg	S2rg	S2rg	S3rg	S2rg	S2rg	S2rg	S3rg	S3rg	
Betageri	4	N1r	S2rg	S3rg	S2rg	S3rg	S2rg	N1r	S3rg	S2rt	S3rg	S3rg	S2rg	S3rg	S2rg	S3rg	S3rg	S3rg	S2rg	S2rg	S2rg	S2rg	S2rg	S3rg	S2rg	S2rg	S2rg	S3rg	S3rg	
Betageri	5	N1r	S2rg	S3rg	S2rg	S3rg	S2rg	N1r	S3rg	S2rt	S3rg	S3rg	S2rg	S3rg	S2rg	S3rg	S3rg	S3rg	S2rg	S2rg	S2rg	S2rg	S2rg	S3rg	S2rg	S2rg	S2rg	S3rg	S3rg	
Betageri	6	N1r	S2rg	S3rg	S2rg	S3rg	S2rg	N1r	S3rg	S2rt	S3rg	S3rg	S2rg	S3rg	S2rg	S3rg	S3rg	S3rg	S2rg	S2rg	S2rg	S2rg	S2rg	S3rg	S2rg	S2rg	S2rg	S3rg	S3rg	
Betageri	8	N1r	S2rg	S3rg	S2rg	S3rg	S2rg	N1r	S3rg	S2rt	S3rg	S3rg	S2rg	S3rg	S2rg	S3rg	S3rg	S3rg	S2rg	S2rg	S2rg	S2rg	S2rg	S3rg	S2rg	S2rg	S2rg	S3rg	S3rg	
Betageri	9	N1r	S2rg	S3rg	S2rg	S3rg	S2rg	N1r	S3rg	S2rt	S3rg	S3rg	S2rg	S3rg	S2rg	S3rg	S3rg	S3rg	S2rg	S2rg	S2rg	S2rg	S2rg	S3rg	S2rg	S2rg	S2rg	S3rg	S3rg	
Betageri	10	S3t	S3t	S3t	S1	S3t	S1	S2r	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S3t	S2t	S2t	S2t	S3t	S3t	S3t	S2t	S2t	
Betageri	109	S3t	S3t	S3t	S1	S3t	S1	S2r	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S3t	S2t	S2t	S2t	S3t	S3t	S3t	S2t	S2t	
Betageri	110	S3t	S3t	S3t	S1	S3t	S1	S2r	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S3t	S2t	S2t	S2t	S3t	S3t	S3t	S2t	S2t	
Betageri	111	S3t	S3t	S3t	S1	S3t	S1	S2r	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S3t	S2t	S2t	S2t	S3t	S3t	S3t	S2t	S2t	
Betageri	112	S3t	S3t	S3t	S1	S3t	S1	S2r	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S3t	S2t	S2t	S2t	S3t	S3t	S3t	S2t	S2t	
Betageri	113	S3t	S3t	S3t	S1	S3t	S1	S2r	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S3t	S2t	S2t	S2t	S3t	S3t	S3t	S2t	S2t	
Betageri	200	S3t	S3t	S3t	S1	S3t	S1	S2r	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S3t	S2t	S2t	S2t	S3t	S3t	S3t	S2t	S2t	
Betageri	201	S3t	S3t	S3t	S1	S3t	S1	S2r	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S3t	S2t	S2t	S2t	S3t	S3t	S3t	S2t	S2t	
Betageri	265	Habit	Habit	Habita	Habit	Habita	Habita	Habita	Habit	Habit	Habit	Habit	Habit	Habit	Habit	Habit	Habit	Habit	Habit	Habit	Habit	Habit	Habit	Habit	Habit	Habit	Habit	Habit	Habit	Habit
Betageri	266	Habit	Habit	Habita	Habit	Habita	Habita	Habita	Habit	Habit	Habit	Habit	Habit	Habit	Habit	Habit	Habit	Habit	Habit	Habit	Habit	Habit	Habit	Habit	Habit	Habit	Habit	Habit	Habit	Habit
Betageri	312	N1rt	S3tz	N1rz	S3rz	N1rt	S3rz	N1rz	N1rz	S3rz	N1rz	N1rz	S3tz	N1rt	S3zg	N1rt	N1rt	N1rz	S3tz	S3rz	S3rz	S3rz	S3rz	N1rz	S3rz	S3rz	S3rz	N1rz	N1rz	
Betageri	313	N1rz	S3tz	S3rz	S2rz	S3tz	S2rz	N1rz	S3rz	S2rz	S3rz	S3rz	S2rz	S3tz	S2rz	N1tz	S3tz	S3rz	S3tz	S3tz	S3tz	S3tz	S2rz	S2rz	S3rz	S3tz	S2rz	S3rz	S3rz	
Betageri	314	N1rz	S3tz	S3rz	S2rz	S3tz	S2rz	N1rz	S3rz	S2rz	S3rz	S3rz	S2rz	S3tz	S2rz	N1tz	S3tz	S3rz	S3tz	S3tz	S3tz	S3tz	S2rz	S2rz	S3rz	S3tz	S2rz	S3rz	S3rz	
Betageri	316	N1rt	S3tz	N1rz	S3rz	N1rt	S3rz	N1rz	N1rz	S3rz	N1rz	N1rz	S3tz	N1rt	S3zg	N1rt	N1rt	N1rz	S3tz	S3rz	S3rz	S3rz	S3rz	N1rz	S3rz	S3rz	S3rz	N1rz	N1rz	
Betageri	317	N1rt	S3tz	N1rz	S3rz	N1rt	S3rz	N1rz	N1rz	S3rz	N1rz	N1rz	S3tz	N1rt	S3zg	N1rt	N1rt	N1rz	S3tz	S3rz	S3rz	S3rz	S3rz	N1rz	S3rz	S3rz	S3rz	N1rz	N1rz	
Betageri	318	N1rt	S3tz	N1rz	S3rz	N1rt	S3rz	N1rz	N1rz	S3rz	N1rz	N1rz	S3tz	N1rt	S3zg	N1rt	N1rt	N1rz	S3tz	S3rz	S3rz	S3rz	S3rz	N1rz	S3rz	S3rz	S3rz	N1rz	N1rz	
Betageri	319	N1rt	S3tz	N1rz	S3rz	N1rt	S3rz	N1rz	N1rz	S3rz	N1rz	N1rz	S3tz	N1rt	S3zg	N1rt	N1rt	N1rz	S3tz	S3rz	S3rz	S3rz	S3rz	N1rz	S3rz	S3rz	S3rz	N1rz	N1rz	

Village	Survey No.	Mango	Maize	Sapota	Sorgham	Guava	Cotton	Tamarind	Lime	Bengalgram	Sunflower	Redgram	Amla	Jackfruit	Custard apple	Cashew	Jamun	Musambi	Groundnut	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Jasmine	Crosandra	Drumstick	Mulbery
Betageri	382	N1rt	S3tz	N1rz	S3rz	N1rt	S3rz	N1rz	N1rz	S3rz	N1rz	N1rz	S3tz	N1rt	S3zg	N1rt	N1rt	N1rz	S3tz	S3rz	S3rz	S3rz	S3rz	N1rz	S3rz	S3rz	S3rz	N1rz	N1rz
Betageri	383	N1r	S2rg	S3rg	S2rg	S3rg	S2rg	N1r	S3rg	S2rt	S3rg	S3rg	S2rg	S3rg	S2rg	S3rg	S3rg	S3rg	S2rg	S2rg	S2rg	S2rg	S2rg	S3rg	S2rg	S2rg	S2rg	S3rg	S3rg
Betageri	384	N1r	S2rg	S3rg	S2rg	S3rg	S2rg	N1r	S3rg	S2rt	S3rg	S3rg	S2rg	S3rg	S2rg	S3rg	S3rg	S3rg	S2rg	S2rg	S2rg	S2rg	S2rg	S3rg	S2rg	S2rg	S2rg	S3rg	S3rg
Betageri	428	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S2g	S3g	S2g	S2rg	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S2g
Betageri	429	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S2g	S3g	S2g	S2rg	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S2g
Betageri	433	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S2g	S3g	S2g	S2rg	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S2g
Betageri	434	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S2g	S3g	S2g	S2rg	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S2g
Betageri	435	N1r	S2rg	S3rg	S2rg	S3rg	S2rg	N1r	S3rg	S2rt	S3rg	S3rg	S2rg	S3rg	S2rg	S3rg	S3rg	S3rg	S2rg	S2rg	S2rg	S2rg	S2rg	S3rg	S2rg	S2rg	S2rg	S3rg	S3rg
Betageri	436	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S2g	S3g	S2g	S2rg	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S2g
Betageri	437	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S2g	S3g	S2g	S2rg	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S2g
Betageri	438	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S2g	S3g	S2g	S2rg	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S2g
Betageri	439	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S2g	S3g	S2g	S2rg	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S2g
Betageri	440	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S2g	S3g	S2g	S2rg	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S2g
Betageri	441	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S2g	S3g	S2g	S2rg	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S2g
Betageri	442	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S2g	S3g	S2g	S2rg	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S2g
Betageri	443	N1rg	S3rg	N1rg	S3rg	N1rg	S3rt	N1rg	N1rg	S3rt	N1rg	N1rg	S3rg	N1rg	S3rg	N1rg	N1rg	N1rg	S3rg	S3rg	S3rg	S3rg	S3rg	N1rg	S3rg	S3rg	S3rg	N1rg	N1rg
Betageri	444	N1rg	S3rg	N1rg	S3rg	N1rg	S3rt	N1rg	N1rg	S3rt	N1rg	N1rg	S3rg	N1rg	S3rg	N1rg	N1rg	N1rg	S3rg	S3rg	S3rg	S3rg	S3rg	N1rg	S3rg	S3rg	S3rg	N1rg	N1rg
Betageri	445	N1rg	S3rg	N1rg	S3rg	N1rg	S3rt	N1rg	N1rg	S3rt	N1rg	N1rg	S3rg	N1rg	S3rg	N1rg	N1rg	N1rg	S3rg	S3rg	S3rg	S3rg	S3rg	N1rg	S3rg	S3rg	S3rg	N1rg	N1rg
Betageri	446	N1rg	S3rg	N1rg	S3rg	N1rg	S3rt	N1rg	N1rg	S3rt	N1rg	N1rg	S3rg	N1rg	S3rg	N1rg	N1rg	N1rg	S3rg	S3rg	S3rg	S3rg	S3rg	N1rg	S3rg	S3rg	S3rg	N1rg	N1rg
Betageri	447	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S2g	S3g	S2g	S2rg	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S2g
Betageri	448	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S2g	S3g	S2g	S2rg	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S2g
Betageri	449	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S2g	S3g	S2g	S2rg	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S2g
Betageri	450	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S2g	S3g	S2g	S2rg	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S2g
Betageri	451	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S2g	S3g	S2g	S2rg	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S2g
Betageri	452	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S2g	S3g	S2g	S2rg	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S2g
Betageri	453	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S2g	S3g	S2g	S2rg	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S2g
Betageri	454	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S2g	S3g	S2g	S2rg	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S2g
Betageri	455	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S2g	S3g	S2g	S2rg	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S2g

Village	Survey No.	Mango	Maize	Sapota	Sorgham	Guava	Cotton	Tamarind	Lime	Bengalgram	Sunflower	Redgram	Amla	Jackfruit	Custard apple	Cashew	Jamun	Musambi	Groundnut	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Jasmine	Crosandra	Drumstick	Mulbery	
Betageri	510	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S2g	S3g	S2g	S2rg	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S2g
Betageri	513	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S2g	S3g	S2g	S2rg	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S2g
Betageri	583	N1rg	S3rg	N1rg	S3rg	N1rg	S3rt	N1rg	N1rg	S3rt	N1rg	N1rg	S3rg	N1rg	S3rg	N1rg	N1rg	N1rg	S3rg	S3rg	S3rg	S3rg	S3rg	N1rg	S3rg	S3rg	S3rg	N1rg	N1rg	N1rg
Betageri	584	N1rg	S3rg	N1rg	S3rg	N1rg	S3rt	N1rg	N1rg	S3rt	N1rg	N1rg	S3rg	N1rg	S3rg	N1rg	N1rg	N1rg	S3rg	S3rg	S3rg	S3rg	S3rg	N1rg	S3rg	S3rg	S3rg	N1rg	N1rg	N1rg
Betageri	588	N1r	S2rg	S3rg	S2rg	S3rg	S2rg	N1r	S3rg	S2rt	S3rg	S3rg	S2rg	S3rg	S2rg	S3rg	S3rg	S3rg	S2rg	S2rg	S2rg	S2rg	S2rg	S3rg	S2rg	S2rg	S2rg	S2rg	S3rg	S3rg
Betageri	589	S3rz	S2zg	S2rz	S2tz	S2rz	S2rz	S3rz	S2rz	S2z	S2rz	S2rz	S2z	S2rz	S2z	N1tz	S3rz	S2rz	S2z	S2zg	S2zg	S2zg	S2zg	S2rz	S2zg	S2zg	S2rz	S2rz	S2z	
Betageri	590	S3rz	S2zg	S2rz	S2tz	S2rz	S2rz	S3rz	S2rz	S2z	S2rz	S2rz	S2z	S2rz	S2z	N1tz	S3rz	S2rz	S2z	S2zg	S2zg	S2zg	S2zg	S2rz	S2zg	S2zg	S2rz	S2rz	S2z	
Betageri	591	S3rz	S2zg	S2rz	S2tz	S2rz	S2rz	S3rz	S2rz	S2z	S2rz	S2rz	S2z	S2rz	S2z	N1tz	S3rz	S2rz	S2z	S2zg	S2zg	S2zg	S2zg	S2rz	S2zg	S2zg	S2rz	S2rz	S2z	
Betageri	592	S3rz	S2zg	S2rz	S2tz	S2rz	S2rz	S3rz	S2rz	S2z	S2rz	S2rz	S2z	S2rz	S2z	N1tz	S3rz	S2rz	S2z	S2zg	S2zg	S2zg	S2zg	S2rz	S2zg	S2zg	S2rz	S2rz	S2z	

PART-B

SOCIO-ECONOMIC STATUS OF FARM HOUSEHOLDS

CONTENTS

1.	Findings of the socio-economic survey	1-2
2.	Introduction	3
3	Methodology	4-5
4	Salient features of the survey	7-24
5	Summary	25-28

LIST OF TABLES

1	Households sampled for socio economic survey	7
2	Population characteristics	7
3	Age wise classification of household members	7
4	Education level of household members	8
5	Occupation of household heads	8
6	Occupation of family members	9
7	Institutional participation of household members	9
8	Type of house owned by households	9
9	Durable assets owned by households	10
10	Average value of durable assets owned by households	10
11	Farm implements owned by households	10
12	Average value of farm implements	11
13	Livestock possession by households	11
14	Average labour availability	11
15	Adequacy of hired labour	11
16	Distribution of land (ha)	12
17	Average land value (Rs./ha)	12
18	Status of bore wells	12
19	Source of irrigation	12
20	Depth of water(Avg in meters)	12
21	Irrigated area (ha)	13
22	Cropping pattern	13
23	Cropping intensity	13
24	Possession of bank account and saving	13
25	Borrowing status	13
26.a	Cost of cultivation of Bajra	14
26.b	Cost of cultivation of Groundnut	15
26.c	Cost of cultivation of Bengal gram	16
26.d	Cost of cultivation of Sorghum	17
26.e	Cost of cultivation of Maize	18
26.f	Cost of cultivation of Sunflower	19

27	Adequacy of fodder	20
28	Annual gross income	20
29	Average annual expenditure	20
30	Horticultural species grown	20
31	Forest species grown	21
32	Marketing of the agricultural produce	21
33	Marketing channels used for sale of agricultural produce	21
34	Mode of transport of agricultural produce	22
35	Incidence of soil and water erosion problems	22
36	Interest shown towards soil testing	22
37	Usage pattern of fuel for domestic use	22
38	Source of drinking water	22
39	Source of light	23
40	Existence of sanitary toilet facility	23
41	Possession of public distribution system (PDS) card	23
42	Participation in NREGA programme	23
43	Adequacy of food items	24
44	Inadequacy of food items	24
45	Farming constraints experienced	24

FINDINGS OF THE SOCIO-ECONOMIC SURVEY

- ❖ *The survey was conducted in Bettageri-1 is located at North latitude 15^o 13' 35.664" and 16^o 11' 29.015" and East longitude 76^o 3' 22.648" and 76^o 2' 6.455" covering an area of about 391.08 ha coming under bettageri village of Koppal taluk.*
- ❖ *Socio-economic analysis indicated that, out of the total sample of 35 respondents, - 5 (14.29%) were marginal, 11(31.43%) were small and 10 (28.57%) were semi medium, 7 (20%) were medium.*
- ❖ *The population characteristics of households indicated that, there were 84 (57.14%) men and 62 (42.18%) were women.*
- ❖ *Majority of the respondents (40.82%) were in the age group of 35-60 years. Education level of the sample households indicated that, majority there were 17.69 per cent illiterates and only 11.56 per cent attained graduation.*
- ❖ *About, 54.29 per cent of household heads practicing agriculture and 31.43 per cent of the household heads were engaged as agricultural labourers.*
- ❖ *Agriculture was the major occupation for 36.73 per cent of the household members.*
- ❖ *In the study area, 91.43 per cent of the households possess katcha house and 2.86 per cent possess pucca house.*
- ❖ *The durable assets owned by the households showed that, 80 per cent possess TV, 37.14 per cent possess mixer grinder and 100 per cent possess mobile phones.*
- ❖ *Farm implements owned by the households indicated that 11.43 per cent of the households possess plough.*
- ❖ *Regarding livestock possession by the households, 17.14 per cent possess local cow and 5.71 per cent possess buffalo respectively.*
- ❖ *The average labour availability in the study area showed that, own men and women labour availability in the micro watershed was 7.71 each, while the hired labour (men) availability was 1.58.*
- ❖ *Further, 8.57 per cent of the households opined that hired labour was inadequate during the agricultural season.*
- ❖ *Out of the total land holding of the sample respondents (66.25 ha), 73.68 per cent of the area is under dry condition and the remaining 26.21 per cent area is irrigated land.*
- ❖ *There were 9 bore wells and 0 dry bore wells among the sampled households.*
- ❖ *Bore well was the major source of irrigation for 25.71 per cent of the households.*
- ❖ *The major crops grown by sample farmers are Bajra, Groundnut, Bengalgram, Sorghum and Maize and cropping intensity was recorded as 98.47 per cent.*
- ❖ *The sample households possessed 5.71 per cent bank account and 5.71 per cent of them have savings in the account.*

- ❖ *About 5.71 per cent of the respondents borrowed credit from various sources.*
- ❖ *The per hectare cost of cultivation for Bajra, Groundnut, Bengalgram, Sorghum and Maize was Rs.28756.92 , 101401.12, 40347.70, 30650.44, and 27142.57 with benefit cost ratio of 1:1.20, 1:0.80, 1:0.90, 1:1.50, and 1:1.60, respectively.*
- ❖ *Further, 28.57 per cent of the households opined that dry fodder was adequate and 28.57 per cent of the households have opined that the green fodder was adequate.*
- ❖ *The average annual gross income of the farmers was Rs. 122345.71 in micro-watershed, of which Rs. 69631.43 comes from agriculture.*
- ❖ *Sampled households have grown horticulture crops has planted 9 coconut trees in the fields and forest species have grown 33 neem trees and 1 banyan trees together in both field and backyard.*
- ❖ *Regarding marketing channels, 94.29 per cent of the households have sold agricultural produce to the local/village merchants, while, 17.14 per cent have sold by Agents/Traders.*
- ❖ *Further, 114.29 per cent of the households have used tractor for the transport of agriculture commodity.*
- ❖ *Majority of the farmers (57.14 %) have experienced soil and water erosion problems in the watershed and 88.57 per cent of the households were interested towards soil testing.*
- ❖ *Firewood connection was the major source of fuel for domestic use for 94.29 per cent of the households and 5.71 per cent households has LPG.*
- ❖ *Piped supply was the major source for drinking water for 91.43 per cent of the households.*
- ❖ *Electricity was the major source of light for 100 per cent of the households.*
- ❖ *In the study area, 40 per cent of the households possess toilet facility.*
- ❖ *Regarding possession of PDS card, 88.57 per cent of the households possessed BPL card and 11.43 per cent do not possess PDS card.*
- ❖ *Cereals (105.71%), pulses (80%), oilseeds (0%) were adequate for consumption.*
- ❖ *Farming constraints experienced by households in the micro watersheds were lower fertility status of the soil (88.57%) wild animal menace on farm field (11.43%), frequent incidence of pest and diseases (80%), inadequacy of irrigation water (28.57%), high cost of fertilizers and plant protection chemicals (80%), high rate of interest on credit (65.71%), low price for the agricultural commodities (77.14%), lack of marketing facilities in the area (60%), inadequate extension services (0%), lack of transport for safe transport of the agricultural produce to the market (11.43%), Less rainfall (8.57%) and Source of Agri-technology information(Newspaper/TV/Mobile) (5.71%).*

INTRODUCTION

Soil and water are the two precious natural resources which are essential for crop production and existence of life on earth. Rainfed agriculture is under severe stress due to various constraints related to agriculture like uneven and erratic distribution of rainfall, indiscriminate use of fertilizers, chemicals and pesticides, adoption of improper land management practices, soil erosion, decline in soil fertility, decline in ground water resources leading to low crop productivity. The area under rainfed agriculture has to be managed effectively using the best available practices to enhance the production of food, fodder and fuel. This is possible if the land resources are characterized at each parcel of land through detailed land resource inventory using the best available techniques of remote sensing, GPS and GIS. The watershed development programs are aimed at the sustainable distribution of its resources and the process of creating and implementing plans, programs, and projects to sustain and enhance watershed functions that affect the plant, animal and human communities within a watershed boundary.

World Bank funded KWDP II, SUJALA III project was implemented in with Broad objective of demonstrating more effective watershed management through greater integration of programmes related to rain-fed agriculture, innovative and science based approaches and strengthen institutional capacities and If successful, it is expected that the systems and tools could be mainstreamed into the overall IWMP in the State of Karnataka and in time, throughout other IWMP operations in India. With this background the socio-economic survey has been carried out with following specific objectives:

1. To understand the demographic features of the households in the micro-watershed
2. To understand the extent of family labour available and additional employment opportunities available within the village.
3. To know the status of assets of households in the micro-watershed for suggesting possible improvements.
4. To study the cropping pattern, cropped area and productivity levels of different households in micro-watershed.
5. To determine the type and extent of livestock owned by different categories of HHs
6. Availability of fodder and level of livestock management.

Scope and importance of survey

Survey helps in identification of different socio-economic and resource use-patterns of farmers at the Micro watershed. Household survey provides demographic features, labor force, and levels of education; land ownership and asset position (including livestock and other household assets) of surveyed households; and cropping patterns, input intensities, and average crop yields from farmers' fields. It also discusses crop utilization and the degree of commercialization of production in the areas; farmers' access to and utilization of credit from formal and informal sources; and the level of adoption and use of soil, water, and pest management technologies.

METHODOLOGY

The description of the methods, components selected for the survey and procedures followed in conducting the baseline survey are furnished under the following heads.

1. Description of the study area

Koppal district is an administrative district in the state of Karnataka in India. In the past Koppal was referred to as 'Kopana Nagara'. Koppal, now a district headquarters is ancient Kopana a major holy place of the Jainas. The district occupies an area of 7,190 km² and has a population of 1,196,089, which 16.58% were urban as of 2001. The Koppal district was formed after split of Raichur district.

Geographers are very particular about the physiography or relief of a region. It plays a very important role in the spatial analysis of agricultural situation of the study area. The undulating topography with black cotton soil shrips, cut across by numerous nalas or streams is the major characteristic feature of the study region. Three physiographic divisions have made considering the local conditions of landforms and crops grown in the district. On the basis of physiography, Koppal district can be divided into three major divisions. They are (a) Koppal & Yelburga plateau, (b) Maidan division, (c) Tungabhadra valley. The district is part of Krishna basin the main streams draining the area are Maskinala, Ilkal-nadi and Hirenala. These are Ephemeral in nature, these come under Tungabhadra sub-basin. The drainage exhibit dentritic to subdentritic with drainage density varies from 1.4 to 7.0kms/sq.km.

According to the 2011 census Koppal district has a population of 1,391,292, roughly equal to the nation of Swaziland or the US state of Hawaii. This gives it a ranking of 350th in India (out of a total of 640). The district has a population density of 250 inhabitants per square kilometre (650/sq mi). Its population growth rate over the decade 2001-2011 was 16.32%.Koppal has a sex ratio of 983 females for every 1000 males, and a literacy rate of 67.28%.

2. Locale of the survey and description of the micro-watershed and

The study was conducted in Bettageri-1 micro-watershed (Bettageri sub-watershed, Koppal taluk & District) is located at North latitude 15⁰ 13' 35.664" and 16⁰ 11' 29.015" and East longitude 76⁰ 3' 22.648" and 76⁰ 2' 6.455" covering an area of about 391.08 ha bounded by under Bettageri Village.

3. Selection of the respondents for the study

The micro-watershed is marked with 320 square meters grids. One farmer from every alternate grid in the micro-watershed was selected for the study and interviewed for socio-economic data. Totally 35 households were interviewed for the survey.

4. The parameters considered for socio-economic survey of households

Two forms of data were collected from the micro-watershed which includes primary data from the farm households and secondary data about the villages under the micro-watershed jurisdiction.

The following parameters were considered for the primary data collection about the socio-economic data of the households, (1) Demographic information, (2) Farm and durable assets owned by households, (3) Livestock possession, (4) Labour availability, (5) Level of migration in the village, Land holding, (7) Cropping pattern, (8) Source of irrigation, (9) Borrowing status, (10) Cost of cultivation of major crops, (11) Economics of subsidiary activities, (12) Fodder availability, (13) Family annual income from different sources, (14) Horticulture and forestry species grown, (15) Additional investment capacity, (16) Marketing practices, (17) Status of soil and water conservation structure, (18) Access to basic needs and (19) Constraints and suggestion.

The following parameters were considered for the secondary data regarding the villages under the micro-watershed jurisdiction, (1) Number of villages in each micro-watershed jurisdiction, (2) Village wise number of households, (3) Geographical area of the villages, (4) Cultivable area including rainfed and irrigated, (5) Number and type of house in each village, (6) Human and livestock population, (7) Facilities in the village such as roads, transport facility for conveyance, drinking water supply, street light and (8) Community based organizations in the villages.

5. Development of interview schedule and data collection

Taking into the consideration the objectives of the survey, an interview schedule was prepared after thorough consultation with the experts in the field of social sciences. A comprehensive interview schedule covering all the major parameters for measuring the socio-economic situation was developed.

6. Tools used to analyze the data

The statistical components such as frequency and percentage were used to analyze the data.

Abbreviations used in the report

LL=Landless

MF=Marginal Farmers

SF=Small farmers

SMF=Semi medium farmers

MDF=Medium farmers

LF=Large Farmers

FINDINGS OF THE SURVEY

This chapter deals with systematic presentation of results of the survey. Keeping in view the objectives, the salient features of the survey are presented under the following headings.

Households sampled for socio-economic survey: The data on households sampled for socio economic survey in Bettageri-1 Micro watershed is presented in Table 1 and it indicated that 35 farmers were sampled in Bettageri-1 micro-watershed among households surveyed 5 (14.3%) were marginal, 11 (31.4 %) were small farmers, 10 (28.6%) were semi-medium and 7 (20%) were large farmers. 2 landless farmers were also interviewed for the survey.

Table 1. Households sampled for socio economic survey in Bettageri-1 micro-watershed

Sl.No.	Particulars	LL (2)		MF (5)		SF (11)		SMF (10)		MDF (7)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Farmers	2	5.71	5	14.3	11	31.4	10	28.6	7	20	35	100

Population characteristics: The population characteristics of households sampled for socio-economic survey in Bettageri-1 Micro watershed is presented in Table 2. The data indicated that, there were 84 (57.14%) men and 62 (42.18%) were women. The average size of the family was 4.5 landless, 4.6 marginal farmers, 4.2 small farmers, 4.1 semi medium farmers and 4 medium farmers.

Table 2. Population characteristics in Bettageri-1 micro-watershed

Sl.No.	Particulars	LL (9)		MF (23)		SF (46)		SMF (41)		MDF (28)		All (147)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Men	6	66.7	14	61	24	52	25	61	15	53.6	84	57.1
2	Women	3	33.3	9	39	21	46	16	39	13	46.4	62	42.2
3	Other	0	0	0	0	1	2.2	0	0	0	0	1	0.68
Total		9	100	23	100	46	100	41	100	28	100	147	100
Average		4.5		4.6		4.2		4.1		4.0		4.2	

Age wise classification of population: The age wise classification of household members in Bettageri-1 Micro watershed is presented in Table 3. The indicated that, 20 (13.61%) of population were 0-15 years of age, 60 (40.82%) were 16-35 years of age, 51(34.69%) were 36-60 years of age and 16 (10.88 %) were above 61 years of age.

Table 3: Age wise classification of members of the household in Bettageri-1 micro-watershed

Sl.No.	Particulars	LL (9)		MF (23)		SF (46)		SMF (41)		MDF (28)		All (147)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	0-15 years of age	0	0	5	21.7	6	13	3	7.32	6	21	20	13.61
2	16-35 years of age	4	44.4	10	43.5	21	45.7	16	39.02	9	32	60	40.82
3	36-60 years of age	4	44.4	6	26.1	15	32.6	15	36.59	11	39	51	34.69
4	> 61 years	1	11.1	2	8.7	4	8.7	7	17.07	2	7.1	16	10.88
Total		9	100	23	100	46	100	41	100	28	100	147	100

Education level of household members: Education level of household members in Bettageri-1 Micro watershed is presented in Table 4. The results indicated that, there were 17.69 per cent of illiterates, 17.69 per cent of them had primary school education, 4.08 per cent middle school education, and 23.81 per cent high school education, 12.93 per cent of them had PUC education, 0.68 per cent of them had Diploma, 11.56 per cent attained graduation, and 5.44 them had other education.

Table 4. Education level of members of the household in Bettageri-1 micro-watershed

Sl.No.	Particulars	LL (9)		MF (23)		SF (46)		SMF (41)		MDF(28)		All (147)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Illiterate	3	33.3	5	21.7	11	23.9	5	12.2	2	7.14	26	17.7
2	Primary School	0	0	7	30.4	6	13	10	24.4	3	10.71	26	17.7
3	Middle School	0	0	0	0	2	4.35	2	4.88	2	7.14	6	4.08
4	High School	4	44.4	3	13	10	21.7	11	26.8	7	25	35	23.8
5	PUC	2	22.2	4	17.4	5	10.9	3	7.32	5	17.86	19	12.9
6	Diploma	0	0	0	0	0	0	1	2.44	0	0	1	0.68
7	ITI	0	0	0	0	5	10.9	2	4.88	0	0	7	4.76
8	Degree	0	0	3	13	4	8.7	6	14.6	4	14.29	17	11.6
9	Masters	0	0	0	0	1	2.17	1	2.44	0	0	2	1.36
Total		9	100	23	100	46	100	41	100	28	100	147	100

Occupation of head of households: The data regarding the occupation of the household heads in Bettageri-1 Micro watershed is presented in Table 5. The results indicate that, 54.29 per cent of households heads were practicing agriculture, 31.43 per cent of the household heads were agricultural Labour and housewife (0%).

Table 5: Occupation of heads of households in Bettageri-1 micro-watershed

Sl.No.	Particulars	LL (2)		MF (5)		SF (11)		SMF (10)		MDF (7)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Agriculture	0	0	4	80	4	36.36	7	70	4	57.1	19	54.29
2	Agricultural Labour	2	100	0	0	4	36.36	2	20	3	42.9	11	31.43
3	Government Service	0	0	0	0	1	9.09	1	10	0	0	2	5.71
4	Private Service	0	0	0	0	1	9.09	0	0	0	0	1	2.86
5	Others	0	0	1	20	1	9.09	0	0	0	0	2	5.71
Total		2	100	5	100	11	100	10	100	7	100	35	100

Occupation of the members of the household: The data regarding the occupation of the household members in Bettageri-1 Micro watershed is presented in Table 6. The results indicate that, agriculture was the major occupation for 36.73 per cent of the household members, 27.21 per cent were agricultural labour, 1.36 per cent were working in government sector, 18.37 per cent were working in pursuing education, 1.36 per cent were involved as housewife and 4.08 per cent were children's.

Table 6: Occupation of members of the household in Bettageri-1 micro-watershed

Sl.No.	Particulars	LL (9)		MF (23)		SF (46)		SMF (41)		MDF (28)		All (147)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Agriculture	0	0	12	52.2	13	28.26	21	51.22	8	29	54	36.7
2	Agricultural Labour	9	100	1	4.35	13	28.26	6	14.63	11	39	40	27.2
3	Household industry	0	0	0	0	1	2.17	0	0	1	3.6	2	1.36
4	Government Service	0	0	0	0	1	2.17	1	2.44	0	0	2	1.36
5	Private Service	0	0	0	0	6	13.04	4	9.76	2	7.1	12	8.16
6	Student	0	0	8	34.8	10	21.74	8	19.51	1	3.6	27	18.4
7	Others	0	0	1	4.35	1	2.17	0	0	0	0	2	1.36
8	Housewife	0	0	0	0	1	2.17	1	2.44	0	0	2	1.36
9	Children	0	0	1	4.35	0	0	0	0	5	18	6	4.08
Total		9	100	23	100	46	100	41	100	28	100	147	100

Institutional Participation of household members: The data regarding the institutional participation of the household members in Bettageri-1 Micro watershed is presented in Table 7. The results show that, out of the total family members in the households 0.68 per cent of them are participating in NGOs and 99.3 per cent of them were not participating in any of the local institutions.

Table 7: Institutional Participation of household member in Bettageri-1 micro-watershed

Sl.No.	Particulars	LL (9)		MF (23)		SF (46)		SMF (41)		MDF (28)		All (147)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	NGOs	0	0	0	0	1	2.17	0	0	0	0	1	0.68
2	No Participation	9	100	23	100	45	97.8	41	100	28	100	146	99.3
Total		9	100	23	100	46	100	41	100	28	100	147	100

Type of house owned: The data regarding the type of house owned by the households in Bettageri-1 Micro watershed is presented in Table 8. The results indicate that, 5.71 percent possess thatched house, 91.43 per cent of the households possess katcha house and 2.86 per cent possess pucca house

Table 8. Type of house owned by households in Bettageri-1 micro-watershed

Sl.No.	Particulars	LL (2)		MF (5)		SF (11)		SMF (10)		MDF (7)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Thatched	1	50	0	0	0	0	1	10	0	0	2	5.71
2	Katcha	1	50	5	100	10	90.91	9	90	7	100	32	91.43
3	Pucca/RCC	0	0	0	0	1	9.09	0	0	0	0	1	2.86
Total		2	100	5	100	11	100	10	100	7	100	35	100

Durable assets owned by the households: The data regarding the Durable Assets owned by the households in Bettageri-1 Micro watershed is presented in Table 9. The results shows that, 80 per cent possess TV, 37.14 per cent possess mixer grinder, 2.86 per cent possess refrigerator, 17.14 per cent possess Bicycle, 45.71 per cent possess motor cycle and 100 per cent possess mobile phones.

Table 9. Durable assets owned by households in Bettageri-1 micro-watershed

Sl.No.	Particulars	LL (2)		MF (5)		SF (11)		SMF (10)		MDF (7)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Radio	1	50	0	0	0	0	0	0	0	0	1	2.86
2	Television	1	50	4	80	11	100	7	70	5	71.4	28	80
3	Mixer/Grinder	0	0	3	60	4	36.4	4	40	2	28.6	13	37.14
4	Refrigerator	0	0	0	0	0	0	1	10	0	0	1	2.86
5	Bicycle	0	0	3	60	2	18.2	1	10	0	0	6	17.14
6	Motor Cycle	0	0	0	0	8	72.7	4	40	4	57.1	16	45.71
7	Mobile Phone	2	100	5	100	11	100	10	100	7	100	35	100

Average value of durable assets: The data regarding the average value of durable assets owned by the households in Bettageri-1 Micro watershed is presented in Table 10. The result shows that, the average value of television was Rs.8303, mixer grinder was Rs.2076, refrigerator was 10000, bicycle was Rs.1816, motor cycle was Rs. 40750, and mobile phone was Rs.2662.

Table 10. Average value of durable assets owned in Bettageri-1 micro-watershed

Average Value (Rs.)

Sl.No.	Particulars	LL (2)	MF (5)	SF (11)	SMF (10)	MDF (7)	All (35)
1	Radio	1000	0	0	0	0	1000
2	Television	9000	7125	7909	9000	9000	8303
3	Mixer/Grinder	0	1666	2500	2000	2000	2076
4	Refrigerator	0	0	0	10000	0	10000
5	Bicycle	0	1633	2000	2000	0	1816
6	Motor Cycle	0	0	40000	39250	43750	40750
7	Mobile Phone	4000	2466	2062	2714	3714	2662

Farm implements owned: The data regarding the farm implements owned by the households in Bettageri-1 Micro watershed is presented in Table 11. About 11.43 per cent of the households possess Bullock Cart, 11.43 per cent possess plough, 48.57 per cent possess Weeder, 8.57 per cent possess tractor and 2.86 per cent possess chaff cutter.

Table 11. Farm implements owned in Bettageri-1 micro-watershed

Sl.No.	Particulars	LL (2)		MF (5)		SF (11)		SMF (10)		MDF (7)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Bullock Cart	0	0	0	0	0	0	1	10	3	42.9	4	11.43
2	Plough	0	0	0	0	0	0	1	10	3	42.9	4	11.43
3	Tractor	0	0	0	0	1	9.09	1	10	1	14.3	3	8.57
4	Weeder	0	0	4	80	6	54.55	4	40	3	42.9	17	48.57
5	Chaff Cutter	0	0	0	0	1	9.09	0	0	0	0	1	2.86
6	Blank	2	100	1	20	4	36.36	5	50	2	28.6	14	40

Average value of farm implements: The data regarding the average value of farm Implements owned by the households in Bettageri-1 Micro watershed is presented in Table

12. The results show that the average value of plough was Rs.671, bullock Cart was Rs.20500, weeder was Rs.54, chaff cutter was Rs.200 and tractor Rs. 366666.

Table 12. Average value of farm implements in Bettageri-1 micro-watershed

Average Value (Rs.)							
Sl.No.	Particulars	LL (2)	MF (5)	SF (11)	SMF (10)	MDF (7)	All (35)
1	Bullock Cart	0	0	0	18000	21333	20500
2	Plough	0	0	0	1500	533	671
3	Tractor	0	0	400000	400000	300000	366666
4	Weeder	0	47	73	44	42	54
5	Chaff Cutter	0	0	200	0	0	200

Livestock possession by the households: The data regarding the Livestock possession by the households in Bettageri-1 Micro watershed is presented in Table 13. The indicate that, 17.14 per cent of the households possess bullocks and local cow and 5.71 per cent possess buffalo.

Table 13. Livestock possession by households in Bettageri-1 micro-watershed

Sl.No.	Particulars	LL (2)		MF (5)		SF (11)		SMF (10)		MDF (7)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Bullock	0	0	1	20	0	0	1	10	4	57.1	6	17.14
2	Local cow	0	0	0	0	2	18.18	3	30	1	14.3	6	17.14
3	Buffalo	0	0	0	0	0	0	0	0	2	28.6	2	5.71

Average Labour availability: The data regarding the average labour availability in Bettageri-1 Micro watershed is presented in Table 14. The indicated that, own labour men available in the micro watershed was 6.59, women available in the micro watershed was 1.12, hired labour (men) available was 1.58 and hired labour (women) available was 7.48.

Table 14. Average labour availability in Bettageri-1 micro-watershed

Sl.No.	Particulars	LL (2)	MF (5)	SF (11)	SMF (10)	MDF (7)	All (35)
1	Hired labour Female	0	5.83	7.73	6.8	5.14	6.59
2	Own Labour Female	0	1.33	1.09	1	1.14	1.12
3	Own labour Male	0	1.6	1.45	1.9	1.29	1.58
4	Hired labour Male	0	7	8.91	6.8	6.57	7.48

Adequacy of hired labour: The data regarding the adequacy of hired labour in Bettageri-1 Micro watershed is presented in Table 15. The results indicate that, 88.57 per cent of the household opined that hired labour was adequate, 8.57 per cent of the household opined that hired labour was Inadequate.

Table 15. Adequacy of hired labour in Bettageri-1 micro-watershed

Sl.No.	Particulars	LL (2)		MF (5)		SF (11)		SMF (10)		MDF (7)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Adequate	0	0	5	100	9	81.8	10	100	7	100	31	88.6
2	Inadequate	0	0	1	20	2	18.2	0	0	0	0	3	8.57

Distribution of land (ha): The data regarding the distribution of land (ha) in Bettageri-1 Micro watershed is presented in Table 16. The results indicate that, 48.81 ha (73.68%) of dry land and 17.36 ha (26.21 %) of irrigated land.

Table 16. Distribution of land (ha) in Bettageri-1 micro-watershed

Sl. No.	Particulars	LL (2)		MF (5)		SF (11)		SMF (10)		MDF (7)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Dry	0	0	3.89	100	15.66	99.51	18.21	82.64	11.05	44.9	48.81	73.68
2	Irrigated	0	0	0	0	0	0	3.82	17.36	13.54	55.1	17.36	26.21
Total		0	100	3.89	100	15.74	100	22.03	100	24.59	100	66.25	100

Average value of land (ha): The data regarding the average land value (Rs./ha) in Bettageri-1 Micro watershed is presented in Table 17. The results show that the average value of dry land was Rs.231415.31 and the average value of irrigated land was Rs.385757.57.

Table 17. Average value of land (ha) in Bettageri-1 micro-watershed

Sl.No.	Particulars	LL (2)	MF (5)	SF (11)	SMF (10)	MDF (7)	All (35)
1	Dry	0	436486.5	255297.2	219604.4	144761.9	231415.3
2	Irrigated	0	0	0	548888.9	339671.2	385757.6

Status of bore wells: The data regarding the status of bore wells in Bettageri-1 Micro watershed is presented in Table 18. The results indicate that, there were 9 functioning bore wells among the sampled households in micro watershed.

Table 18. Status of bore wells in Bettageri-1 micro-watershed

Sl.No	Particulars	LL(2)	MF (5)	SF(11)	SMF(10)	MDF(7)	All (35)
1	Functioning	0	0	0	3	6	9

Source of irrigation: The data regarding the source of irrigation in Bettageri-1 Micro watershed is presented in Table 19. The results that bore well were major source of irrigation for 25.71 per cent of the households.

Table 19. Source of irrigation in Bettageri-1 micro-watershed

Sl.No.	Particulars	LL (2)		MF (5)		SF (11)		SMF (10)		MDF (7)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Bore Well	0	0	0	0	0	0	3	30	6	85.71	9	25.71

Depth of water (Avg. In meters): The data regarding the depth of water in Bettageri-1 Micro watershed is presented in Table 20. The results revealed that, the depth of bore well was 27.43 meter.

Table 20. Depth of water (Avg. In meters) in Bettageri-1 micro-watershed

Sl.No.	Particulars	LL (2)	MF (5)	SF (11)	SMF (10)	MDF (7)	All (35)
1	Bore Well	0	0	0	32	91.44	27.43

Irrigated Area (ha): The data regarding the irrigated area (ha) in Bettageri-1 Micro watershed is presented in Table 21. The results indicate that, the availability of irrigation water was used for kharif crops was 17.37 ha.

Table 21. Irrigated Area (ha) in Bettageri-1 micro-watershed

Sl.No.	Particulars	LL (2)	MF (5)	SF (11)	SMF (10)	MDF (7)	All (35)
1	Kharif	0	0	0	3.83	13.54	17.37
	Total	0	0	0	3.83	13.54	17.37

Cropping pattern: The data regarding the cropping pattern in Bettageri-1 Micro watershed is presented in Table 22. The results indicate that, farmers have grown crop1 (12.17 ha), crop 2(11.59 ha), crop 3 (2.19 ha) crop4 (1.32 ha), crop5 (0.96 ha) and crop 6 (0.81 ha).

Table 22. Cropping pattern in Bettageri-1 micro-watershed

Sl.No.	Particulars	LL (2)	MF (5)	SF (11)	SMF(10)	MDF (7)	All (35)
1	Kharif - Maize	0	0.81	3.69	16.74	11.81	33.06
2	Kharif - Sunflower	0	1.32	3.78	1.3	5.78	12.17
3	Kharif - Bajra	0	0.96	6.82	2.48	1.34	11.59
4	Kharif - Bengal gram	0	0.81	1.38	0	0	2.19
5	Kharif - Sorghum	0	0	0	1.32	0	1.32
6	Rabi - Jowar	0	0.96	0	0	0	0.96
7	Kharif - Groundnut	0	0	0	0	0.81	0.81

Cropping intensity: The data regarding the cropping intensity in Bettageri-1 Micro watershed is presented in Table 23. The results indicate that, the cropping intensity was 98.47 per cent.

Table 23. Cropping intensity (%) in Bettageri-1 micro-watershed

Sl.No.	Particulars	LL (2)	MF (5)	SF (11)	SMF (10)	MDF (7)	All (35)
1	Cropping Intensity	0	100	95.37	99.06	100	98.47

Possession of bank account and savings: The data regarding the possession of bank account and saving in Bettageri-1 micro-watershed is presented in Table 24. The results indicate that, 5.71 cent of the households posses bank account and 5.71 per cent of them have savings.

Table 24. Possession of Bank account and savings in Bettageri-1 micro-watershed

Sl.No.	Particulars	LL (2)		MF (5)		SF (11)		SMF (10)		MDF (7)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Account	0	0	0	0	2	18.18	0	0	0	0	2	5.71
2	Savings	0	0	0	0	2	18.18	0	0	0	0	2	5.71

Borrowing status: The data regarding the borrowing status in Bettageri-1 micro-watershed is presented in Table 25. The results indicate that, 5.71 percent of the sample farmers have borrowed credit from different sources.

Table 25. Borrowing status in Bettageri-1 micro-watershed

Sl.No.	Particulars	LL (2)		MF (5)		SF (11)		SMF (10)		MDF (7)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Credit Aailed	0	0	0	0	2	18.2	0	0	0	0	2	5.71

Cost of Cultivation of Bajra: The data regarding the cost of cultivation (Rs/ha) of Bajra in Bettageri-1 micro watershed is presented in Table 26.a. The results indicate that, the total cost of cultivation (Rs/ha) for Bajra was Rs. 28756.92. The gross income realized by the farmers was Rs. 35301.75. The net income from Bajra cultivation was Rs.6544.83, thus the benefit cost ratio was found to be 1:1.20.

Table 26(a). Cost of Cultivation of Bajra in Bettageri-1 micro-watershed

Sl.No	Particulars	Units	Phy units	Value(Rs.)	% to C3
I	Cost A1				
1	Hired Human Labour	Man days	39.52	6906.73	24.02
2	Bullock	Pairs/day	1.01	606.41	2.11
3	Tractor	Hours	3.92	3122.57	10.86
4	Machinery	Hours	0.93	741	2.58
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	7.21	1224.23	4.26
7	FYM	Quintal	3.8	760.61	2.64
8	Fertilizer + micronutrients	Quintal	6.4	5738.62	19.96
9	Pesticides (PPC)	Kgs / liters	0.91	776.76	2.7
10	Irrigation	Number	0	0	0
12	Msc. Charges (Marketing costs etc)		0	0	0
13	Depreciation charges		0	1433.82	4.99
14	Land revenue and Taxes		0	2.88	0.01
II	Cost B1				
16	Interest on working capital			1020.28	3.55
17	Cost B1 = (Cost A1 + sum of 15 and 16)			22333.92	77.66
III	Cost B2				
18	Rental Value of Land			312.5	1.09
19	Cost B2 = (Cost B1 + Rental value)			22646.42	78.75
IV	Cost C1				
20	Family Human Labour		17.81	3494.11	12.15
21	Cost C1 = (Cost B2 + Family Labour)			26140.53	90.9
V	Cost C2				
22	Risk Premium			2.13	0.01
23	Cost C2 = (Cost C1 + Risk Premium)			26142.65	90.91
VI	Cost C3				
24	Managerial Cost			2614.27	9.09
25	Cost C3 = (Cost C2 + Managerial Cost)			28756.92	100
VII	Economics of the Crop				
a.	Main Product	a) Main Product (q)	22.84	34552.99	
		b) Main Crop Sales Price (Rs.)		1512.5	
	By Product	e) Main Product (q)	1.05	748.76	
		f) Main Crop Sales Price (Rs.)		712.5	
b.	Gross Income (Rs.)			35301.75	
c.	Net Income (Rs.)			6544.83	
d.	Cost per Quintal (Rs./q.)			1258.79	
e.	Benefit Cost Ratio (BC Ratio)			1:1.2	

Cost of Cultivation of Groundnut: The data regarding the cost of cultivation (Rs/ha) of Groundnut in Bettageri-1 micro watershed is presented in Table 26.b. The results indicate that, the total cost of cultivation (Rs/ha) for Groundnut was Rs. 101401.12. The gross income realized by the farmers was Rs. 86450. The net income from Groundnut cultivation was Rs.-14951.12, thus the benefit cost ratio was found to be 1:0.80.

Table 26(b). Cost of Cultivation of Groundnut in Bettageri-1 micro-watershed

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
I	Cost A1				
1	Hired Human Labour	Man days	60.52	11917.75	11.75
2	Bullock	Pairs/day	1.24	741	0.73
3	Tractor	Hours	11.12	8892	8.77
4	Machinery	Hours	4.94	3952	3.9
5	Seed Main Crop	Kgs (Rs.)	247	37050	36.54
7	FYM	Quintal	1.24	247	0.24
8	Fertilizer + micronutrients	Quintal	14.82	12251.2	12.08
9	Pesticides (PPC)	Kgs / liters	0	0	0
10	Irrigation	Number	0	0	0
11	Repairs		0	0	0
12	Msc. Charges (Marketing costs etc)		0	0	0
13	Depreciation charges		0	4.94	0
14	Land revenue and Taxes		0	3.29	0
II	Cost B1				
16	Interest on working capital			5945.9	5.86
17	Cost B1 = (Cost A1 + sum of 15 and 16)			81005.09	79.89
III	Cost B2				
18	Rental Value of Land			0	0
19	Cost B2 = (Cost B1 + Rental value)			81005.09	79.89
IV	Cost C1				
20	Family Human Labour		55.57	11176.75	11.02
21	Cost C1 = (Cost B2 + Family Labour)			92181.84	90.91
V	Cost C2				
22	Risk Premium			1	0
23	Cost C2 = (Cost C1 + Risk Premium)			92182.84	90.91
VI	Cost C3				
24	Managerial Cost			9218.28	9.09
25	Cost C3 = (Cost C2 + Managerial Cost)			101401.12	100
VII	Economics of the Crop				
a.	Main Product	a) Main Product (q)		18.53	74100
		b) Main Crop Sales Price (Rs.)			4000
	By Product	e) Main Product (q)		2.47	12350
		f) Main Crop Sales Price (Rs.)			5000
b.	Gross Income (Rs.)			86450	
c.	Net Income (Rs.)			-14951.12	
d.	Cost per Quintal (Rs./q.)			5473.74	
e.	Benefit Cost Ratio (BC Ratio)			1:0.8	

Cost of Cultivation of Bengalgram: The data regarding the cost of cultivation (Rs/ha) of Bengalgram in Bettageri-1 micro watershed is presented in Table 26.c. The results indicate, the total cost of cultivation (Rs/ha) for Bengalgram was Rs.40347.70. The gross income realized by the farmers was Rs. 36503.33. The net income from Bengalgram cultivation was Rs. -3844.37, thus the benefit cost ratio was found to be 1:0.90.

Table 26(c). Cost of Cultivation of Bengalgram in Bettageri-1 micro-watershed

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
I	Cost A1				
1	Hired Human Labour	Man days	28.33	4818.68	11.94
2	Bullock	Pairs/day	1.96	1176.88	2.92
3	Tractor	Hours	2.32	1859.76	4.61
4	Machinery	Hours	0	0	0
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	73.37	17900.23	44.36
6	Seed Inter Crop	Kgs.	0	0	0
7	FYM	Quintal	0	0	0
8	Fertilizer + micronutrients	Quintal	3.31	3966.53	9.83
9	Pesticides (PPC)	Kgs/liters	0.98	980.74	2.43
10	Irrigation	Number	0	0	0
11	Repairs		0	0	0
13	Depreciation charges		0	3.2	0.01
14	Land revenue and Taxes		0	3.29	0.01
II	Cost B1				
16	Interest on working capital			2741.82	6.8
17	Cost B1 = (Cost A1 + sum of 15 and 16)			33451.14	82.91
III	Cost B2				
18	Rental Value of Land			333.33	0.83
19	Cost B2 = (Cost B1 + Rental value)			33784.47	83.73
IV	Cost C1				
20	Family Human Labour		14.49	2894.26	7.17
21	Cost C1 = (Cost B2 + Family Labour)			36678.73	90.91
V	Cost C2				
22	Risk Premium			1	0
23	Cost C2 = (Cost C1 + Risk Premium)			36679.73	90.91
VI	Cost C3				
24	Managerial Cost			3667.97	9.09
25	Cost C3 = (Cost C2 + Managerial Cost)			40347.7	100
VII	Economics of the Crop				
a.	Main Product	a) Main Product (q)		8.21	35914.89
		b) Main Crop Sales Price (Rs.)			4375
	By Product	e) Main Product (q)		0.98	588.44
		f) Main Crop Sales Price (Rs.)			600
b.	Gross Income (Rs.)			36503.33	
c.	Net Income (Rs.)			-3844.37	
d.	Cost per Quintal (Rs./q.)			4914.99	
e.	Benefit Cost Ratio (BC Ratio)			1:0.9	

Cost of Cultivation of Sorghum: The data regarding the cost of cultivation (Rs/ha) of Sorghum in Bettageri-1 micro watershed is presented in Table 26.d. The results indicate that, the total cost of cultivation (Rs/ha) for Sorghum was Rs. 30650.44. The gross income realized by the farmers was Rs.46816. The net income from Sorghum cultivation was Rs. 16165.56, thus the benefit cost ratio was found to be 1:1.50.

Table 26(d). Cost of Cultivation of Sorghum in Bettageri-1 micro-watershed

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
I	Cost A1				
1	Hired Human Labour	Man days	20.52	2660	8.68
2	Bullock	Pairs/day	2.28	1368	4.46
3	Tractor	Hours	2.28	1824	5.95
4	Machinery	Hours	0	0	0
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	26.6	3724	12.15
7	FYM	Quintal	1.52	304	0.99
8	Fertilizer + micronutrients	Quintal	10.64	10214.4	33.33
9	Pesticides (PPC)	Kgs / liters	0.76	760	2.48
10	Irrigation	Number	1.52	0	0
12	Msc. Charges (Marketing costs etc)		0	0	0
13	Depreciation charges		0	273.6	0.89
14	Land revenue and Taxes		0	3.29	0.01
II	Cost B1				
16	Interest on working capital			1800.41	5.87
17	Cost B1 = (Cost A1 + sum of 15 and 16)			22931.7	74.82
III	Cost B2				
18	Rental Value of Land			333.33	1.09
19	Cost B2 = (Cost B1 + Rental value)			23265.03	75.9
IV	Cost C1				
20	Family Human Labour		35.72	4598	15
21	Cost C1 = (Cost B2 + Family Labour)			27863.03	90.91
V	Cost C2				
22	Risk Premium			1	0
23	Cost C2 = (Cost C1 + Risk Premium)			27864.03	90.91
VI	Cost C3				
24	Managerial Cost			2786.4	9.09
25	Cost C3 = (Cost C2 + Managerial Cost)			30650.44	100
VII	Economics of the Crop				
a.	Main Product	a) Main Product (q)		38	45600
		b) Main Crop Sales Price (Rs.)			1200
	By Product	e) Main Product (q)		1.52	1216
		f) Main Crop Sales Price (Rs.)			800
b.	Gross Income (Rs.)			46816	
c.	Net Income (Rs.)			16165.56	
d.	Cost per Quintal (Rs./q.)			806.59	
e.	Benefit Cost Ratio (BC Ratio)			1:1.5	

Cost of Cultivation of Maize: The data regarding the cost of cultivation (Rs/ha) of Maize in Bettageri-1 micro watershed is presented in Table 26.e. The results indicate that, the total cost of cultivation (Rs/ha) for Maize was Rs.27142.57. The gross income realized by the farmers was Rs. 42359.90. The net income from Maize cultivation was Rs. 15217.33, thus the benefit cost ratio was found to be 1:1.60.

Table 26(e). Cost of Cultivation of Maize in Bettageri-1 micro-watershed

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
I	Cost A1				
1	Hired Human Labour	Man days	31.7	5853.22	21.56
2	Bullock	Pairs/day	0.78	467.09	1.72
3	Tractor	Hours	2.96	2360.59	8.7
4	Machinery	Hours	0.84	675.35	2.49
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	19.93	2391.52	8.81
7	FYM	Quintal	0.98	196.27	0.72
8	Fertilizer + micronutrients	Quintal	7.86	6737.04	24.82
9	Pesticides (PPC)	Kgs/ liters	0.62	588.02	2.17
10	Irrigation	Number	2.41	0	0
11	Repairs		0	0	0
12	Msc. Charges		0	0	0
13	Depreciation charges		0	247.47	0.91
14	Land revenue and Taxes		0	2.91	0.01
II	Cost B1				
16	Interest on working capital			1189.79	4.38
17	Cost B1 = (Cost A1 + sum of 15 and 16)			20709.26	76.3
III	Cost B2				
18	Rental Value of Land			313.73	1.16
19	Cost B2 = (Cost B1 + Rental value)			21022.99	77.45
IV	Cost C1				
20	Family Human Labour		19.85	3650.01	13.45
21	Cost C1 = (Cost B2 + Family Labour)			24673	90.9
V	Cost C2				
22	Risk Premium			2.06	0.01
23	Cost C2 = (Cost C1 + Risk Premium)			24675.06	90.91
VI	Cost C3				
24	Managerial Cost			2467.51	9.09
25	Cost C3 = (Cost C2 + Managerial Cost)			27142.57	100
VII	Economics of the Crop				
a.	Main Product	a) Main Product (q)		32.05	39596.55
		b) Main Crop Sales Price (Rs.)			1235.29
	By Product	e) Main Product (q)		2.3	2763.35
		f) Main Crop Sales Price (Rs.)			1200
b.	Gross Income (Rs.)			42359.9	
c.	Net Income (Rs.)			15217.33	
d.	Cost per Quintal (Rs./q.)			846.77	
e.	Benefit Cost Ratio (BC Ratio)			1:1.6	

Cost of Cultivation of Sunflower: The data regarding the cost of cultivation (Rs/ha) of Sunflower in Bettageri-1 micro watershed is presented in Table 26.f. The results indicate that, the total cost of cultivation (Rs/ha) for Sunflower was Rs. 25646.36. The gross income realized by the farmers was Rs. 93892.32. The net income from Sunflower cultivation was Rs. 68245.96, thus the benefit cost ratio was found to be 1:3.7.

Table 26(f). Cost of Cultivation of Sunflower in Bettageri-1 micro-watershed

Sl.No	Particulars	Units	PhyUnits	Value(Rs.)	% toC3
I	Cost A1				
1	Hired Human Labour	Man days	34.1	6213.96	24.23
2	Bullock	Pairs/day	1.17	704.48	2.75
3	Tractor	Hours	2.36	1891.94	7.38
4	Machinery	Hours	0.79	634.49	2.47
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	5.49	2304.94	8.99
6	Seed Inter Crop	Kgs.	0	0	0
7	FYM	Quintal	1.55	310.25	1.21
8	Fertilizer + micronutrients	Quintal	5.02	4117.87	16.06
9	Pesticides (PPC)	Kgs / liters	0.94	944.14	3.68
10	Irrigation	Number	0	0	0
11	Repairs		0	0	0
12	Msc. Charges (Marketing costs etc)		0	0	0
13	Depreciation charges		0	468.79	1.83
14	Land revenue and Taxes		0	3.29	0.01
II	Cost B1				
16	Interest on working capital			921.38	3.59
17	Cost B1 = (Cost A1 + sum of 15 and 16)			18515.52	72.2
III	Cost B2				
18	Rental Value of Land			333.33	1.3
19	Cost B2 = (Cost B1 + Rental value)			18848.85	73.5
IV	Cost C1				
20	Family Human Labour		25.5	4465.02	17.41
21	Cost C1 = (Cost B2 + Family Labour)			23313.87	90.91
V	Cost C2				
22	Risk Premium			1	0
23	Cost C2 = (Cost C1 + Risk Premium)			23314.87	90.91
VI	Cost C3				
24	Managerial Cost			2331.49	9.09
25	Cost C3 = (Cost C2 + Managerial Cost)			25646.36	100
VII	Economics of the Crop				
a.	Main Product	a) Main Product (q)	11.69	93892.32	
		b) Main Crop Sales Price (Rs.)		8033.33	
b.	Gross Income (Rs.)			93892.32	
c.	Net Income (Rs.)			68245.96	
d.	Cost per Quintal (Rs./q.)			2194.28	
e.	Benefit Cost Ratio (BC Ratio)			1:3.7	

Adequacy of fodder: The data regarding the adequacy of fodder in Bettageri-1 Micro watershed is presented in Table 27. The results indicate that, 28.57 per cent of the households opined that dry fodder was adequate. With respect to green fodder availability, 28.57 percent of them opined it was sufficient.

Table 27. Adequacy of fodder in Bettageri-1 micro-watershed

Sl. No	Particulars	LL(2)		MF(5)		SF (11)		SMF(10)		MDF (7)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Adequate-Dry Fodder	0	0	2	40	2	18.1	3	30	3	42.9	10	28.5
2	Adequate-Green Fodder	0	0	2	40	2	18.1	3	30	3	42.9	10	28.5

Average annual gross income: The data regarding the annual gross income in Bettageri-1 Micro watershed is presented in Table 28. The results indicate that, the farmers have annual gross income of Rs. 122345.71 in micro-watershed, of which Rs. 69631.43 is from agriculture itself.

Table 28. Average annual gross income in Bettageri-1 micro-watershed

(Avg value in Rs.)

Sl.No.	Particulars	LL (2)	MF (5)	SF (11)	SMF (10)	MDF (7)	All (35)
1	Service/salary	0	0	60545.5	45000	14285.7	34742.9
2	Wage	18000	18600	13636.4	18000	18571.4	16828.6
3	Agriculture	0	41800	41786.4	95945	115571	69631.4
4	Dairy Farm	0	0	0	0	5714.29	1142.86
Income(Rs.)		18000	60400	115968	158945	154143	122346

Average annual Expenditure: The data regarding the average annual expenditure in Bettageri-1 Micro watershed is presented in Table 29. The results indicate that, the farmers have annual gross expenditure of Rs. 523906.06 in micro-watershed, of which Rs. 45342.86 is from agriculture itself.

Table 29. Average annual Expenditure in Bettageri-1 micro-watershed

(Avg value in Rs.)

Sl.No.	Particulars	LL (2)	MF (5)	SF (11)	SMF (10)	MDF (7)	All (35)
1	Service/salary	0	0	82500	103333	32500	24857.1
2	Wage	11000	10500	17600	24000	30000	10342.9
3	Agriculture	0	21400	24272.7	61800	85000	45342.9
4	Dairy Farm	0	0	0	0	20000	571.43
Total		11000	31900	124373	189133	167500	523906

Horticulture species grown: The data regarding horticulture species grown in Bettageri-1 Micro watershed is presented in Table 30. The results indicate that, the total number of horticultural trees grown (both field and backyard) coconut (9).

Table 30. Horticulture species grown in Bettageri-1 micro-watershed

Sl.No.	Particulars	LL (2)		MF (5)		SF (11)		SMF (10)		MDF (7)		All (35)	
		F	B	F	B	F	B	F	B	F	B	F	B
1	Coconut	0	0	0	0	0	2	0	1	6	0	6	3

*F= Field B=Back Yard

Forest species grown: The data regarding forest species grown in Bettageri-1 Micro watershed is presented in Table 31. The results indicate that, households have planted 0 Eucalyptus trees, 0 cashew trees, 0 teak trees, 33 neem trees and 1 banyan trees together in both field and backyard.

Table 31. Forest species grown in Bettageri-1 micro-watershed

Sl.No.	Particulars	LL (2)		MF (5)		SF (11)		SMF (10)		MDF (7)		All (35)	
		F	B	F	B	F	B	F	B	F	B	F	B
1	Neem	0	0	4	1	7	1	17	0	3	0	31	2
2	Banyan	0	0	0	0	1	0	0	0	0	0	1	0

*F= Field B=Back Yard

Marketing of agricultural produce: The data regarding marketing of the agricultural produce in Bettageri-1 Micro watershed is presented in Table 32. The results indicated that, 89.50 percent of output of bajra was sold in the market; 70.59 percent of output of Bengal gram was sold in the market; 66.67 percent of output of Groundnut was sold in the market; 37.50 percent of output of jowar was sold in the market and 96.78 percent of output of maize was sold in the market; 90 per cent of output of sorghum was sold in the market; and 100 per cent of output of sunflower was sold in the market.

Table 32. Marketing of agricultural produce in Bettageri-1 micro-watershed

Sl. No	Crops	Output obtained (q)	Output retained (q)	Output sold (q)	Output sold (%)	Avg. Price obtained (Rs/q)
1	Bajra	238	25	213	89	1513
2	Bengalgram	17	5	12	71	4375
3	Groundnut	15	5	10	67	4000
4	Jowar	16	10	6	38	2100
5	Maize	1026	33	993	97	1167
6	Sorghum	50	5	45	90	1200
7	Sunflower	140	0	140	100	8033

Marketing channels used for sale of agricultural produce: The data regarding marketing channels used for sale of agricultural produce in Bettageri-1 Micro watershed is presented in Table 33. The results indicated that, 94.29 cent of the households have sold agricultural produce to the local/village merchants, 2.86 per per cent have sold to Agent/Traders and 17.14 per cent of regulated market.

Table 33. Marketing channels used for sale of agricultural produce in Bettageri-1 micro-watershed

Sl. No	Particulars	LL(2)		MF(5)		SF(11)		SMF(10)		MDF(7)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Agent/Traders	0	0	0	0	0	0	0	0	1	14.3	1	2.86
2	Local/village Merchant	0	0	4	80	9	81.8	10	100	9	129	33	94.29
3	Regulated Market	0	0	2	40	4	36.4	0	0	0	0	6	17.14

Mode of transport of agricultural produce: The data regarding mode of transport of agricultural produce in Bettageri-1 Micro watershed is presented in Table 34. The results indicated that, 114.29 cent of the households have used tractor.

Table 34. Mode of transport of agricultural produce in Bettageri-1 micro-watershed

Sl.No.	Particulars	LL (2)		MF (5)		SF (11)		SMF (10)		MDF (7)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Tractor	0	0	6	120	13	118	11	110	10	143	40	114.3

Incidence of soil and water erosion problems: The data regarding incidence of incidence of soil and water erosion problems in Bettageri-1 Micro watershed is presented in Table 35. The results indicate that, 57.14 per cent of the households have experienced soil and water erosion problems.

Table 35. Incidence of soil and water erosion problems in Bettageri-1 micro-watershed

Sl. No	Particulars	LL (2)		MF(5)		SF(11)		SMF(10)		MDF(7)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Soil and water erosion problems in the farm	0	0	1	20	6	54.6	8	80	5	71	20	57.1

Interest towards soil testing: The data regarding Interest shown towards soil testing in Bettageri-1 Micro watershed is presented in Table 36. The results indicated that, 88.57 per cent of the households were interested towards soil testing.

Table 36. Interest regarding soil testing in Bettageri-1 micro-watershed

Sl.No	Particulars	LL(2)		MF (5)		SF (11)		SMF(10)		MDF(7)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Interest in soil test	0	0	5	100	10	90	10	100	6	85.7	31	88.5

Usage pattern of fuel for domestic use: The data on usage pattern of fuel for domestic use in Bettageri-1 Micro watershed is presented in Table 37. The results indicated that, LPG was the major source of fuel for domestic use for 5.71 per cent of the households followed by firewood (94.29 %).

Table 37. Usage pattern of fuel for domestic use in Bettageri-1 micro-watershed

Sl.No.	Particulars	LL (2)		MF (5)		SF (11)		SMF (10)		MDF (7)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Fire Wood	2	100	5	100	11	100	9	90	6	85.7	33	94.29
2	LPG	0	0	0	0	0	0	1	10	1	14.3	2	5.71

Source of drinking water: The data on source of drinking water in Bettageri-1 Micro watershed is presented in Table 38. The results indicated that, tank supply of water was the major source for drinking water for 0 per cent of the households followed by piped waters supply (91.43 %), bore well water (2.86%) and open well water for (2.86%).

Table 38. Source of drinking water in Bettageri-1 micro-watershed

Sl.No.	Particulars	LL (2)		MF (5)		SF (11)		SMF (10)		MDF (7)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Piped supply	2	100	4	80	9	81.82	10	100	7	100	32	91.43
2	Bore Well	0	0	1	20	0	0	0	0	0	0	1	2.86
3	Open well	0	0	0	0	1	9.09	0	0	0	0	1	2.86

Source of light: The data on source of light in Bettageri-1 Micro watershed is presented in Table 39. The results indicated that, electricity was the major source of light for 100 per cent of the households.

Table 39. Source of light in Bettageri-1 micro-watershed

Sl.No.	Particulars	LL (2)		MF (5)		SF (11)		SMF (10)		MDF (7)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Electricity	2	100	5	100	11	100	10	100	7	100	35	100

Existence of sanitary toilet facility: The data on availability of toilet facility in Bettageri-1 Micro watershed is presented in Table 40. The results indicated that, 40 per cent of the households possess toilets.

Table 40. Existence of sanitary toilet facility in Bettageri-1 micro-watershed

Sl.No	Particulars	LL (2)		MF (5)		SF (11)		SMF(10)		MDF(7)		All(35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Sanitary toilet facility	2	100	1	20	6	54.5	3	30	2	28.6	14	40

Possession of PDS card: The data regarding possession of PDS card in Bettageri-1 Micro watershed is presented in Table 41. The results indicated that, 88.57 per cent of the households possessed BPL card and 11.43 per cent do not possess PDS card.

Table 41. Possession of PDS card in Bettageri-1 micro-watershed

Sl.No	Particulars	LL (2)		MF (5)		SF (11)		SMF(10)		MDF (7)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	BPL	2	100	5	100	10	90.9	8	80	6	85.7	31	88.5
2	Not possessed	0	0	0	0	1	9.09	2	20	1	14.3	4	11.4

Participation in NREGA programme: The data regarding Participation in NREGA programme in Bettageri-1 Micro watershed is presented in Table 42. The results indicated that, only 40 percent of the participate have participated in NREGA programme.

Table 42. Participation in NREGA programme in Bettageri-1 micro-watershed

Sl. No	Particulars	LL(2)		MF(5)		SF(11)		SMF(10)		MDF(7)		All(35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Participation in NREGA programme	0	0	3	60	5	45	6	60	0	0	14	40

Adequacy of food items: The data regarding adequacy of food items in Bettageri-1 Micro watershed is presented in Table 43. The results indicated that, the extent of adequacy of food items for cereals, pulses, and vegetables were 105.71, 80, 88.57 per cent respectively, similarly for milk, Egg and Meat (91.43%).

Inadequacy of food items: The data regarding in adequacy of food items in Bettageri-1 Micro watershed is presented in Table 44. The results indicated that, the extent of in adequacy of food items for pulses, Oilseeds and vegetables were 20, 100, 8.57 per cent respectively, similarly for fruits (100%), milk (5.71%) and egg (8.57%).

Table 43. Adequacy of food items in Bettageri-1 micro-watershed

Sl.No.	Particulars	LL (2)		MF (5)		SF (11)		SMF (10)		MDF (7)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Cereals	2	100	5	100	11	100	10	100	7	100	35	100
2	Pulses	2	100	3	60	7	63.64	9	90	7	100	28	80
3	Vegetables	2	100	4	80	9	81.82	9	90	7	100	31	88.57
4	Milk	2	100	4	80	9	81.82	10	100	7	100	32	91.43
5	Egg	2	100	4	80	9	81.82	10	100	7	100	32	91.43
6	Meat	2	100	4	80	9	81.82	10	100	7	100	32	91.43

Table 44. Inadequacy of food items in Bettageri-1 micro-watershed

Sl.No.	Particulars	LL (2)		MF (5)		SF (11)		SMF (10)		MDF (7)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Pulses	0	0	2	40	4	36.36	1	10	0	0	7	20
2	Oilseed	2	100	5	100	11	100	10	100	7	100	35	100
3	Vegetables	0	0	0	0	2	18.18	1	10	0	0	3	8.57
4	Fruits	2	100	5	100	11	100	10	100	7	100	35	100
5	Milk	0	0	1	20	1	9.09	0	0	0	0	2	5.71
6	Egg	0	0	1	20	2	18.18	0	0	0	0	3	8.57

Farming constraints: The data regarding farming constraints experienced by households in Bettageri-1 Micro watershed is presented in Table 45. The results indicated that, lower fertility status of the soil was the constraint experienced by (88.57 %) per cent of the households, wild animal menace on farm field (11.43%), frequent incidence of pest and diseases (80%), inadequacy of irrigation water (28.57%), high cost of fertilizers and plant protection chemicals (80%), high rate of interest on credit (65.71%), low price for the agricultural commodities (77.14 %), lack of marketing facilities in the area (60%), lack of transport for safe transport of the agricultural produce to the market (11.43%), less rainfall (8.57%), source of agri-technology information (5.71%).

Table 45. Farming constraints experienced in Bettageri-1 micro-watershed

Sl.	Particulars	LL (2)		MF(5)		SF (11)		SMF(10)		MDF(7)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Lower fertility status of the soil	0	0	4	80	10	90.9	10	100	7	100	31	88.5
2	Wild animal menace on farm field	0	0	2	40	1	9.09	1	10	0	0	4	11.4
3	Frequent incidence of pest and diseases	0	0	4	80	8	72.7	9	90	7	100	28	80
4	Inadequacy of irrigation water	0	0	0	0	2	18.1	3	30	5	71.43	10	28.5
5	High cost of Fertilizers and plant protection chemicals	0	0	4	80	9	81.8	9	90	6	85.71	28	80
6	High rate of interest on credit	0	0	4	80	8	72.7	7	70	4	57.14	23	65.7
7	Low price for the agricultural commodities	0	0	3	60	8	72.7	9	90	7	100	27	77.1
8	Lack of marketing facilities in the area	0	0	4	80	4	36.3	8	80	5	71.43	21	60
9	Lack of transport for safe transport of the Agril produce to the market.	0	0	1	20	2	18.1	0	0	1	14.29	4	11.4
10	Less rainfall	0	0	1	20	2	18.1	0	0	0	0	3	8.57
11	Source of Agri-technology information	0	0	1	20	1	9.09	0	0	0	0	2	5.71

SUMMARY AND IMPLICATIONS

In order to assess the socio-economic condition of the farmers in the watershed 35 households located in the micro watershed were interviewed for the survey. The study was conducted in Bettageri-1 micro-watershed (Bettageri sub-watershed, Koppal taluk & District) is located at North latitude 15⁰ 13' 35.664" and 16⁰ 11' 29.015" and East longitude 76⁰ 3' 22.648" and 76⁰ 2' 6.455" covering an area of about 391.08 ha bounded by under Bettageri Village.

Socio-economic analysis indicated that, out of the total sample of 35 respondents, 5 (14.29%) were marginal, 11(31.43%) were small and 10 (28.57%) were semi medium, 7 (20%) were medium. The population characteristics of households indicated that, there were 84 (57.14%) men and 62 (42.18%) were women. Majority of the respondents (40.82%) were in the age group of 35-60 years. Education level of the sample households indicated that, majority there were 17.69 per cent illiterates and only 11.56 per cent attained graduation. About, 54.29 per cent of household heads practicing agriculture and 31.43 per cent of the household heads were engaged as agricultural labourers. Agriculture was the major occupation for 36.73 per cent of the household members.

In the study area, 91.43 per cent of the households possess katcha house and 2.86 per cent possess pucca house. The durable assets owned by the households showed that, 80 per cent possess TV, 37.14 per cent possess mixer grinder and 100 per cent possess mobile phones. Farm implements owned by the households indicated that 11.43 per cent of the households possess plough. Regarding livestock possession by the households, 17.14 per cent possess local cow and 5.71 per cent possess buffalo respectively.

The average labour availability in the study area showed that, own men and women labour availability in the micro watershed was 7.71 each, while the hired labour (men) availability was 1.58. Further, 8.57 per cent of the households opined that hired labour was inadequate during the agricultural season. Out of the total land holding of the sample respondents (66.25 ha), 73.68 per cent of the area is under dry condition and the remaining 26.21 per cent area is irrigated land. There were 9 bore wells the sampled households. Bore well was the major source of irrigation for 25.71 per cent of the households.

The major crops grown by sample farmers are Bajra, Groundnut, Bengalgram, Sorghum and Maize and cropping intensity was recorded as 98.47 per cent.

The sample households possessed 5.71 per cent bank account and 5.71 per cent of them have savings in the account. About 5.71 per cent of the respondents borrowed credit from various sources.

The per hectare cost of cultivation for Bajra, Groundnut, Bengalgram, Sorghum and Maize was Rs.28756.92 , 101401.12, 40347.70, 30650.44 and 27142.57 with benefit cost ratio of 1:1.20, 1:0.80, 1:0.90, 1:1.50 and 1:1.60 respectively.

Further, 28.57 per cent of the households opined that dry fodder was adequate and 28.57 per cent of the households have opined that the green fodder was adequate.

The average annual gross income of the farmers was Rs. 122345.71 in micro-watershed, of which Rs. 69631.43 comes from agriculture.

Sampled households have grown horticulture crops has planted 9 coconut trees in the fields and forest species have grown 33 neem trees and 1 banyan trees together in both field and backyard.

Regarding marketing channels, 94.29 per cent of the households have sold agricultural produce to the local/village merchants, while, 17.14 per cent have sold by Agents/Traders. Further, 114.29 per cent of the households have used tractor for the transport of agriculture commodity.

Majority of the farmers (57.14 %) have experienced soil and water erosion problems in the watershed and 88.57 per cent of the households were interested towards soil testing.

Firewood connection was the major source of fuel for domestic use for 94.29 per cent of the households and 5.71 per cent households has LPG. Piped supply was the major source for drinking water for 91.43 per cent of the households. Electricity was the major source of light for 100 per cent of the households. In the study area, 40 per cent of the households possess toilet facility. Regarding possession of PDS card, 88.57 per cent of the households possessed BPL card and 11.43 per cent do not possess PDS card. Cereals (105.71%), pulses (80%), oilseeds (0%) were adequate for consumption.

Farming constraints experienced by households in the micro watersheds were lower fertility status of the soil (88.57%) wild animal menace on farm field (11.43%), frequent incidence of pest and diseases (80%), inadequacy of irrigation water (28.57%), high cost of fertilizers and plant protection chemicals (80%), high rate of interest on credit (65.71%), low price for the agricultural commodities (77.14%), lack of marketing facilities in the area (60%), inadequate extension services (0%), lack of transport for safe transport of the agricultural produce to the market (11.43%), Less rainfall (8.57%) and Source of Agri-technology information(Newspaper/TV/Mobile) (5.71%).

Implications of the survey

- ✓ Result indicated that, there were 17.69 per cent were illiterate hence, extension methodologies such as demonstration, street play, drama, video shows will be effective in dissemination of the technologies in the micro watershed.
- ✓ The data indicate that, 91.43 per cent of the households possess katcha house. Hence, the development department while implementing the watershed plan should focus on agriculture to enhance the productivity of major crops in the area to increase the income of the farmers.

- ✓ Results indicated that the local institutional participation of the household members in the micro watershed is minimal hence, activities like membership campaign, awareness creation about the benefits of membership in local institutions and strengths of organized groups must be conveyed.
- ✓ Majority of the households in the watershed have experience in use of mobile phones, and television hence, these mass media can be effectively utilized for transfer of technology as well as for information dissemination.
- ✓ The farm machinery/implement possession in the micro watershed was found to be minimum the reasons may lack of knowledge or lack of financial ability which can be addressed through training on use of different farm implements, providing information on different sources of finance for purchase of farm implements.
- ✓ The possession of livestock such as crossbred cow found is less hence, farmers must be made aware of the benefits of crossbred cow in increased milk production.
- ✓ The possession of livestock such as sheep, goat and poultry was found to be low hence, farmers may be informed the role of subsidiary enterprises in enhancing the income and information on financial support for subsidiary activities.
- ✓ The data indicate that, job/work was the reason for all the migrants hence, farmers may be trained on profitable agriculture or self employment such as animal husbandry, plate making, sheep rearing, goat rearing, rabbit rearing with suitable information on sources of financial support.
- ✓ The results indicate that there was a change in quality of life due to migration hence, the developmental departments should take actions to arrest migration and to improve the quality of the life in rural areas.
- ✓ Households possess 48.81ha (73.68 %) of dry land and 17.36ha (26.21 %) of irrigated land hence, the availability of the dryland agricultural technologies such as short duration crops, high yielding drought resistance crop varieties, drip irrigation technology and subsidy information will be helpful for the farmers to enhance the productivity of land and as well as farmers income.
- ✓ Few of the bore well in micro watershed found non functional hence, farmers may be trained on possibility of bore well rejuvenation.
- ✓ Open well was major source of irrigation for 0 per cent of the households. hence, in order to increase the area under irrigation as well as to increase the water use efficiency farmers may trained on drip irrigation and provide the information on subsidy for drip irrigation equipment's along with the information on different agencies which provides the financial assistance for drip irrigation.
- ✓ Farmers have grown horticulture crops has 9 coconut trees in the fields. Hence, production technologies related to these crops can be made available to the farmers for better adoption.

- ✓ The cropping intensity in the micro watershed was found to be (98.47 %) hence, care must be taken by the implementing agency to bring uncultivated land into cultivation through suitable measures.
- ✓ Many of the household members have borrowed loan from cooperative banks which has higher rate of interest hence, farmers may be sensitized on the different sources of credit with lesser interest rate such SHGs etc.
- ✓ The results indicated the non availability of both green and dry fodder throughout the year hence, fodder development activities can be taken up in the micro watershed.
- ✓ The average annual gross income of the households Rs.69631.43 from agriculture and Rs. 16828.57 from wages and. Agriculture was found to be the major source of income for households hence; the development activities should focus on productivity enhancement, marketing arrangements and agricultural technology dissemination to have a direct impact on the farmers.
- ✓ The cultivation of forest species is found minimal hence; information and production technology related to agro-forestry and integrated farming system.
- ✓ The data indicated that, 57.14 per cent of the households have experienced soil and water erosion problems. Hence, those farmers who reported the soil and water erosion problems may be given attention while implementation of the watershed development plan.
- ✓ The data indicated that, 88.57 per cent of the households have interest in soil testing hence, farmers must be provided with the information on various institutions which are involved in soil testing for the benefit of the farmers.
- ✓ Except summer ploughing the adoption of other soil and water conservation structures is minimum hence, the farmers in the micro watershed should be sensitized on the use of different conservation structures for soil water conservation.
- ✓ Cereals and pulses found be adequate for per cent of the households respectively hence, farm households and the farm women must be trained on importance of balanced nutrition and role of vegetable, milk, egg, meat in balanced diet.
- ✓ Lower fertility status of the soil (88.57%), wild animal menace on farm field (11.43%), frequent incidence of pest and diseases (80%), high cost of fertilizers and plant protection chemicals (80%), high rate of interest on credit (65.71%), low price for the agricultural commodities (77.14%), lack of marketing facilities in the area (60%), inadequate extension services (0%), lack of transport for safe transport of the agricultural produce to the market (11.43%) were the major farming constraints experienced hence, these constraints must be addressed immediately for the welfare of the farmers. Awareness to be created among the farmers to approach nearest KVKs/RSKs and other developmental departments for technical and for subsidized inputs and utilize the well established regulated markets, approaching the contract firms, direct markets to avoid the involvement of middlemen.