



हर कदम, हर डगर
किसानों का हमसफर
भारतीय कृषि अनुसंधान परिषद

AgriSearch with a human touch

**LAND RESOURCE INVENTORY AND SOCIO-ECONOMIC STATUS OF
FARM HOUSEHOLDS FOR WATERSHED PLANNING AND
DEVELOPMENT**

CHANDRAKI-1 (4D2D6E1b) MICROWATERSHED

Yadgir Taluk & District, Karnataka

Karnataka Watershed Development Project – II

SUJALA – III

World Bank funded Project



THE WORLD BANK



ICAR – NATIONAL BUREAU OF SOIL SURVEY AND LAND USE PLANNING



ICAR - NBSS & LUP

**WATERSHED DEVELOPMENT DEPARTMENT
GOVT. OF KARNATAKA, BANGALORE**



About ICAR - NBSS&LUP

The ICAR-National Bureau of Soil Survey and Land Use Planning (ICAR-NBSS&LUP), Nagpur, a premier Institute of the Indian Council of Agricultural Research (ICAR), was set up during 1976 with the objective to prepare soil resource maps at national, state and district levels and to provide research inputs in soil resource mapping and its applications, land evaluation, land use planning, land resource management, and database management using GIS for optimising land use on different kinds of soils in the country.

The Bureau has been engaged in carrying out soil resource survey, agro-ecological and soil degradation mapping at the country, state and district levels for qualitative assessment and monitoring the soil health towards viable land use planning. The research activities have resulted in identifying the soil potentials and problems, and the various applications of the soil surveys with the ultimate objective of sustainable agricultural development. The Bureau has the mandate to correlate and classify soils of the country and maintain a National Register of all the established soil series. The Institute is also imparting in-service training to staff of the soil survey agencies in the area of soil survey, land evaluation and soil survey interpretations for land use planning. The Bureau in collaboration with Panjabrao Krishi Vidyapeeth, Akola is running post-graduate teaching and research programme in land resource management, leading to M.Sc. and Ph.D. degrees.

Citation: Rajendra Hegde, Ramesh kumar S.C. B.A. Dhanorkar, K.V. Niranjana, S. Srinivas, M.Lalitha, R.S. Reddy and S.K. Singh (2019). Land resource inventory and socio-economic status of farm households for watershed planning and development of Chandraki-1 (6E1b) Microwatershed, Yadgir Taluk and District, Karnataka”, NBSS&LUPSujala MWS Publ .587, ICAR – NBSS & LUP, RC, Bangalore. p.131 & 35.

TO OBTAIN COPIES,

Please write to:

Director, ICAR - NBSS & LUP,

Amaravati Road, NAGPUR - 440 033, India

Phone : (0712) 2500386, 2500664, 2500545 (O)

Telefax : 0712-2522534

E-Mail : director@nbsslup.ernet.in

Website URL : nbsslup.in

Or

Head, Regional Centre, ICAR - NBSS&LUP, Hebbal, Bangalore - 560 024

Phone : (080) 23412242, 23510350 (O)

Telefax : 080-23510350

E-Mail : nbssrcb@gmail.com



**LAND RESOURCE INVENTORY AND SOCIO-ECONOMIC
STATUS OF FARM HOUSEHOLDS FOR WATERSHED
PLANNING AND DEVELOPMENT
CHANDRAKI-1 (4D2D6E1b) MICROWATERSHED**

Yadgir Taluk & District, Karnataka

Karnataka Watershed Development Project – II

Sujala-III

World Bank funded Project



THE WORLD BANK



**ICAR – NATIONAL BUREAU OF SOIL SURVEY AND LAND USE
PLANNING**



**WATERSHED DEVELOPMENT DEPARTMENT, GOVT. OF
KARNATAKA, BANGALORE**



PREFACE

In Karnataka, as in other Indian States, the livelihoods of rural people are intertwined with farming pursuits. The challenges in agriculture are seriously threatening the livelihood of a large number of farmers as they have been practicing farming in contextual factors beyond their control. Climatic factors are the most important ones and have become much more significant in recent times due to rapid climate changes induced by intensive anthropogenic activities affecting our ecosystem in multiple ways. Climate change has become the reality, it is happening and efforts to evolve and demonstrate climate resilient technologies have become essential. Due to the already over stressed scenario of agrarian sector, the climate change is resulting in manifold increase in the complexities, pushing the rural mass to face more and more unpredictable situations. The rising temperatures and unpredictable rainfall patterns are going to test seriously the informed decisions farmers have to make in order to survive in farming and sustain their livelihood.

It is generally recognized that impacts of climate change shall not be uniform across the globe. It is said that impact of climate change is more severe in South Asia. Based on the analysis of meteorological data, it is predicted that in India, there will be upward trend in mean temperature, downward trend in relative humidity, annual rainfall and number of wet days in a year. Also, in general, phenomena like erratic monsoon, spread of tropical diseases, rise in sea levels, changes in availability of fresh water, frequent floods, droughts, heat waves, storms and hurricanes are predicted. Each one of these adverse situations are already being experienced in various parts of India and also at the global level. Decline in agricultural productivity of small and marginal farmers becoming more vulnerable is already witnessed.

In Karnataka, more than 60 per cent of the population live in rural areas and depend on agriculture and allied activities for their livelihood. Though the state has achieved significant progress in increasing the yield of many crops, there is tremendous pressure on the land resources due to the growing and competing demands of various land uses. This is reflected in the alarming rate of land degradation observed. Already more than 50 per cent of the area is affected by various forms of degradation. If this trend continues, the sustainability of the fragile ecosystem will be badly affected. The adverse effects of change in the climatic factors are putting additional stress on the land resources and the farmers dependent on this.

The natural resources (land, water and vegetation) of the state need adequate and constant care and management, backed by site-specific technological interventions and investments particularly by the government. Detailed database pertaining to the nature of the land resources, their constraints, inherent potentials and suitability for various land

based rural enterprises, crops and other uses is a prerequisite for preparing location-specific action plans, which are in tune with the inherent capability of the resources. Any effort to evolve climate resilient technologies has to be based on the baseline scientific database. Then only one can expect effective implementation of climate resilient technologies, monitor the progress, make essential review of the strategy, and finally evaluate the effectiveness of the implemented programs. The information available at present on the land resources of the state are of general nature and useful only for general purpose planning. Since the need of the hour is to have site-specific information suitable for farm level planning and detailed characterization and delineation of the existing land resources of an area into similar management units is the only option.

ICAR-NBSS&LUP, Regional Centre, Bangalore has taken up a project sponsored by the Karnataka Watershed Development Project-II, (Sujala-III), Government of Karnataka funded by the World Bank under Component-1 Land Resource Inventory. This study was taken up to demonstrate the utility of such a database in reviewing, monitoring and evaluating all the land based watershed development programs on a scientific footing. To meet the requirements of various land use planners at grassroots level, the present study on “Land Resource Inventory and Socio-Economic Status of Farm Households for Watershed Planning and Development of Chandaraki-1 Microwatershed, Yadgir Taluk & District, Karnataka” for integrated development was taken up in collaboration with the State Agricultural Universities, IISC, KRSRAC, KSNDMC as Consortia partners. The project provides detailed land resource information at cadastral level (1:7920 scale) for all the plots and socio-economic status of farm households covering thirty per cent farmers randomly selected representing landed and landless class of farmers in the microwatershed. The project report with the accompanying maps for the Microwatershed will provide required detailed database for evolving effective land use plan, alternative land use options and conservation plans for the planners, administrators, agricultural extension personnel, KVK officials, developmental departments and other land users to manage the land resources in a sustainable manner.

It is hoped that this database will be useful to the planners, administrators and developmental agencies working in the area in not only for formulating location specific developmental schemes but also for their effective monitoring at the village/watershed level.

Nagpur

Date: 15-12-2019

S.K. SINGH

Director, ICAR - NBSS&LUP, Nagpur

Contributors

Dr. Rajendra Hegde Principal Scientist, Head & Project Leader, Sujala-III Project ICAR-NBSS&LUP, Regional Centre, Bangalore	Dr. S.K.Singh Director, ICAR-NBSS&LUP Coordinator, Sujala-III Project Nagpur
Soil Survey, Mapping & Report Preparation	
Dr. B.A. Dhanorkar	Sh. R.S. Reddy
Dr. K.V. Niranjana	Mr. Somashekar T N
	Smt. Chaitra, S.P.
	Dr. Gopali bardhan
	Ms. Arpitha
	Dr. Mahendra Kumar, M.B.
Field Work	
Sh. C.BacheGowda	Sh. Mahesh, D.B.
Sh. Somashekar	Sh. Ashok S Sindagi
Sh. M. Jayaramaiah	Sh. Veerabhadrapa B.
Sh. Paramesha, K.	Sh. Shankarappa
Sh. B. M. Narayana Reddy	Sh. Anand
	Sh. Arun N Kambar.
	Sh Kamalesh Awate
	Sh. Sharaan Kumar Huppar
	Sh. Yogesh H.N.
	Sh. Kalaveerachari R Kammar
GIS Work	
Dr. S.Srinivas	Sh. A.G.Devendra Prasad
Sh. D.H.Venkatesh	Sh. Prakashanaik, M.K.
Smt.K.Sujatha	Sh. Abhijith Sastry, N.S.
Smt. K.V.Archana	Sh. Sudip Kumar Suklabaidya
Sh. N. Maddileti	Sh. Avinash, K.N.
	Sh. Amar Suputhra, S
	Sh. Deepak, M.J.
	Smt. K.Karunya Lakshmi
	Ms. Seema, K.V.
	Ms. A. Rajab Nisha
Laboratory Analysis	
Dr. K.M.Nair	Ms. Steffi Peter
Smt. Arti Koyal	Ms. Thara, V.R
Smt. Parvathy	Ms. Roopa, G.
	Ms. Swati, H.
	Sh. Shantaveera Swami
	Ms. Shwetha, N.K.

	Smt. Ishrat Haji
	Ms. P. Pavan Kumari
	Ms. Padmaja
	Ms. Veena, M.
Soil & Water Conservation	
Sh. Sunil P. Maske	
Socio-Economic Analysis	
Dr. S.C. Ramesh Kumar	Sh. M. K. Prakashanaik
	Ms. Sowmya K.B
	Sh.Manjunath M
	Sh.Veerabhadraswamy R
	Sh.Lankesh RS
	Sh.Kalaveerachari R Kammar
	Sh.Pradyumma U
	Sh.Yogेशha HN
	Sh.Vijay kumar lamani
	Sh.Arun N Kambar
	Sh.Vinay
	Sh.Basavaraj.Biradar
	Sh.Vinod R
	Sh.Praveenkumar P Achalkar
	Sh.Rajendra D
Watershed Development Department, GoK, Bangalore	
Sh. Rajeev Ranjan IFS Project Director & Commissioner, WDD	Dr. A. Natarajan NRM Consultant, Sujala-III Project
Dr. S.D. Pathak IFS Executive Director & Chief Conservator of Forests, WDD	

PART-A

LAND RESOURCE INVENTORY

Contents

Preface		
Contributors		
Executive Summary		
Chapter 1	Introduction	1
Chapter 2	Geographical Setting	3
2.1	Location and Extent	3
2.2	Geology	3
2.3	Physiography	4
2.4	Drainage	4
2.5	Climate	4
2.6	Natural Vegetation	6
2.7	Land Utilization	6
Chapter 3	Survey Methodology	9
3.1	Base maps	9
3.2	Image Interpretation for Physiography	9
3.3	Field Investigation	12
3.4	Soil Mapping	14
3.5	Land Management Units (LMU's)	14
3.6	Laboratory Characterization	15
Chapter 4	The Soils	19
4.1	Soils of granite gneiss landscape	19
Chapter 5	Interpretation for Land Resource Management	35
5.1	Land Capability Classification	35
5.2	Soil Depth	37
5.3	Surface Soil Texture	38
5.4	Soil Gravelliness	39
5.5	Available Water Capacity	40
5.6	Soil Slope	41
5.7	Soil Erosion	42
Chapter 6	Fertility Status	45
6.1	Soil Reaction (pH)	45
6.2	Electrical Conductivity (EC)	45
6.3	Organic Carbon (OC)	45
6.4	Available Phosphorus	47
6.5	Available Potassium	47
6.6	Available Sulphur	47
6.7	Available Boron	48
6.8	Available Iron	48
6.9	Available Manganese	48
6.10	Available Copper	48
6.11	Available Zinc	48

Chapter 7	Land Suitability for Major Crops	53
7.1	Land suitability for Sorghum	53
7.2	Land suitability for Maize	54
7.3	Land suitability for Bajra	55
7.4	Land suitability for Groundnut	56
7.5	Land suitability for Sunflower	57
7.6	Land suitability for Redgram	58
7.7	Land suitability for Bengal gram	59
7.8	Land suitability for Cotton	60
7.9	Land suitability for Chilli	61
7.10	Land suitability for Tomato	62
7.11	Land suitability for Brinjal	63
7.12	Land suitability for Onion	64
7.13	Land suitability for Bhendi	65
7.14	Land suitability for Drumstick	66
7.15	Land suitability for Mango	67
7.16	Land suitability for Guava	68
7.17	Land suitability for Sapota	69
7.18	Land Suitability for Pomegranate	70
7.19	Land Suitability for Musambi	71
7.20	Land Suitability for Lime	72
7.21	Land Suitability for Amla	73
7.22	Land Suitability for Cashew	74
7.23	Land Suitability for Jackfruit	75
7.24	Land Suitability for Jamun	76
7.25	Land Suitability for Custard apple	77
7.26	Land Suitability for Tamarind	78
7.27	Land Suitability for Mulberry	79
7.28	Land Suitability for Marigold	80
7.29	Land Suitability for Chrysanthemum	81
7.30	Land use classes	113
7.31	Proposed Crop Plan	114
Chapter 8	Soil Health Management	117
Chapter 9	Soil and Water conservation Treatment Plan	123
9.1	Treatment Plan	124
9.2	Recommended Soil and Water Conservation measures	127
9.3	Greening of Microwatershed	128
	References	131
	Appendix I	I-X
	Appendix II	XI-XX
	Appendix III	XXI-XXXII

LIST OF TABLES

2.1	Mean Monthly Rainfall, PET, 1/2 PET at Yadgir Taluk & District	5
2.2	Land Utilization in Yadgir district	7
3.1	Differentiating Characteristics used for Identifying Soil Series	14
3.2	Soil map unit description of Chandaraki-1Microwatershed	15
4.1	Physical and Chemical Characteristics of Soil Series identified in Chandaraki-1microwatershed	25
7.1	Soil-Site Characteristics of Chandaraki-1Microwatershed	83
7.2	Land suitability criteria for Sorghum	84
7.3	Land suitability criteria for Maize	85
7.4	Land suitability criteria for Bajra	86
7.5	Land suitability criteria for Groundnut	87
7.6	Land suitability criteria for Sunflower	88
7.7	Land suitability criteria for Redgram	89
7.8	Land suitability criteria for Bengal gram	90
7.9	Land suitability criteria for Cotton	91
7.10	Land suitability criteria for Chilli	92
7.11	Land suitability criteria for Tomato	93
7.12	Land suitability criteria for Brinjal	94
7.13	Land suitability criteria for Onion	95
7.14	Land suitability criteria for Bhenidi	96
7.15	Land suitability criteria for Drumstick	97
7.16	Land suitability criteria for Mango	98
7.17	Land suitability criteria for Guava	99
7.18	Land suitability criteria for Sapota	100
7.19	Land suitability criteria for Pomegranate	101
7.20	Land suitability criteria for Musambi	102
7.21	Land suitability criteria for Lime	103
7.22	Land suitability criteria for Amla	104
7.23	Land suitability criteria for Cashew	105
7.24	Land suitability criteria for Jackfruit	106
7.25	Land suitability criteria for Jamun	107
7.26	Land suitability criteria for Custard apple	108
7.27	Land suitability criteria for Tamarind	109

7.28	Land suitability criteria for Mulberry	110
7.29	Land suitability criteria for Marigold	111
7.30	Land suitability criteria for Chrysanthemum	112
7.31	Proposed Crop Plan for Chandaraki-1 Microwatershed	115

LIST OF FIGURES

2.1	Location map of Chandaraki-1 Microwatershed	3
2.2 a	Granite and granite gneiss rock formation	4
2.3	Rainfall distribution in Yadgir Taluk & District	5
2.4	Natural vegetation of Chandaraki-1 Microwatershed	6
2.5	Current Land use map of Chandaraki-1 Microwatershed	7
2.6 a & b	Different crops and cropping systems in Chandaraki-1 Microwatershed	8
3.1	Scanned and Digitized Cadastral map of Chandaraki-1 Microwatershed	11
3.2	Satellite image of Chandaraki-1 Microwatershed	11
3.3	Cadastral map overlaid on IRS PAN+LISS IV merged imagery of Chandaraki-1 Microwatershed	12
3.4	Location of profiles in a transect	13
3.5	Soil phase or management units of Chandaraki-1 Microwatershed	17
5.1	Land Capability Classification map of Chandaraki-1 Microwatershed	37
5.2	Soil Depth map of Chandaraki-1 Microwatershed	38
5.3	Surface Soil Texture map of Chandaraki-1 Microwatershed	39
5.4	Soil Gravelliness map of Chandaraki-1 Microwatershed	40
5.5	Soil Available Water Capacity map of Chandaraki-1 Microwatershed	41
5.6	Soil Slope map of Chandaraki-1 Microwatershed	42
5.7	Soil Erosion map of Chandaraki-1 Microwatershed	43
6.1	Soil Reaction (pH) map of Chandaraki-1 Microwatershed	46
6.2	Electrical Conductivity (EC) map of Chandaraki-1 Microwatershed	46
6.3	Soil Organic Carbon (OC) map of Chandaraki-1 Microwatershed	47
6.4	Soil Available Phosphorus map of Chandaraki-1 Microwatershed	48
6.5	Soil Available Potassium map of Chandaraki-1 Microwatershed	49
6.6	Soil Available Sulphur map of Chandaraki-1 Microwatershed	49
6.7	Soil Available Boron map of Chandaraki-1 Microwatershed	50
6.8	Soil Available Iron map of Chandaraki-1 Microwatershed	50
6.9	Soil Available Manganese map of Chandaraki-1 Microwatershed	51
6.10	Soil Available Copper map of Chandaraki-1 Microwatershed	51
6.11	Soil Available Zinc map of Chandaraki-1 Microwatershed	52
7.1	Land suitability for Sorghum	54
7.2	Land suitability for Maize	55

7.3	Land suitability for Bajra	56
7.4	Land suitability for Groundnut	57
7.5	Land suitability for Sunflower	58
7.6	Land suitability for Redgram	59
7.7	Land suitability for Bengal gram	60
7.8	Land suitability for Cotton	61
7.9	Land suitability for Chilli	62
7.10	Land suitability for Tomato	63
7.11	Land Suitability map of Brinjal	64
7.12	Land Suitability map of Onion	65
7.13	Land Suitability map of Bhendi	66
7.14	Land suitable for Drumstick	67
7.15	Land suitability for Mango	68
7.16	Land suitability for Guava	69
7.17	Land suitability for Sapota	70
7.18	Land suitability for Pomegranate	71
7.19	Land suitability for Musambi	72
7.20	Land suitability for Lime	73
7.21	Land suitability for Amla	74
7.22	Land suitability for Cashew	75
7.23	Land suitability for Jackfruit	76
7.24	Land suitability for Jamun	77
7.25	Land suitability for Custard apple	78
7.26	Land suitability for Tamarind	79
7.27	Land suitability for Mulberry	80
7.28	Land suitability for Marigold	81
7.29	Land suitability for Chrysanthemum	82
7.30	Land use classes map of Chandaraki-1Microwatershed	114
9.1	Soil and water conservation map of Chandaraki-1Microwatershed	128

EXECUTIVE SUMMARY

The land resource inventory of Chandaraki-1 Microwatershed was conducted using village cadastral maps and IRS satellite imagery on 1:7920 scale. The false colour composites of IRS imagery were interpreted for physiography and the physiographic delineations were used as base for mapping soils. The soils were studied in several transects and a soil map was prepared with phases of soil series as mapping units. Random checks were made all over the area outside the transects to confirm and validate the soil map unit boundaries. The soil map shows the geographic distribution and extent, characteristics, classification, behavior and use potentials of the soils in the microwatershed.

The present study covers an area of 608 ha in Yadgir taluk & district, Karnataka. The climate is semiarid and categorized as drought-prone with an average annual rainfall of 866 mm, of which about 652 mm is received during south-west monsoon, 138 mm during north-east and the remaining 76 mm during the rest of the year. An area of 505 ha in the microwatershed is covered by soils, 1 ha is covered by rock outcrops and 101 ha by others (habitation and water body). The salient findings from the land resource inventory are summarized briefly below.

- ❖ The soils belong to 9 soil series and 3 soil phases (management units) and 4 land management units.*
- ❖ The length of crop growing period is about 120-150 days starting from 1st week of June to 4th week of October.*
- ❖ From the master soil map, several interpretative and thematic maps like land capability, soil depth, surface soil texture, soil gravelliness, available water capacity, soil slope and soil erosion were generated.*
- ❖ Soil fertility status maps for macro and micronutrients were generated based on the surface soil samples collected at every 320 m grid interval.*
- ❖ Land suitability for growing 29 major agricultural and horticultural crops was assessed and maps showing the degree of suitability along with constraints were generated.*
- ❖ About 505 ha area in the microwatershed is suitable for agriculture.*
- ❖ About <1 per cent of area is very shallow (<25 cm), 45 per cent of area is shallow (25-50 cm), 1 per cent of area of the microwatershed has soils that are moderately shallow (50-75 cm), 13 per cent of area of the microwatershed has soils that are moderately deep (75-100 cm), 22 per cent of area is deep (100 - 150 cm) and 2 per cent of area is very deep (>150 cm).*
- ❖ About <1 per cent area in the microwatershed has sandy soils, 4 per cent loamy soils and 79 per cent clayey soils at the surface.*
- ❖ About 64 per cent area in the microwatershed is non gravelly (<15%) and 20 per cent is gravelly (15-35%).*
- ❖ About 24 per cent area of the microwatershed is very high (>200 mm/m) in available water capacity, 13 per cent is low (51-100 mm/m), <1 per cent is medium (101-150 mm/m) and 46 per cent area is very low (<50 mm/m).*

- ❖ *Entire area in the microwatershed is under very gently sloping (1-3% slope) lands.*
- ❖ *Maximum area of about 83 per cent is moderately (e2) eroded and <1 per cent area is severely (e3) eroded.*
- ❖ *An area of about 29 per cent is slightly acid (pH 6.0-6.5), 4 per cent is moderately acid (pH 5.5-6.0), 31 per cent is neutral (pH 6.5-7.3), 10 per cent is slightly alkaline (pH 7.3-7.8), 6 per cent moderately alkaline (pH 7.8-8.4) and 4 per cent is strongly alkaline (pH 8.4-9.0) in soil reaction.*
- ❖ *The Electrical Conductivity (EC) in the entire area of the microwatershed is <2dsm¹ indicating that the soils are non-saline.*
- ❖ *Entire microwatershed area is high (>0.75%) in organic carbon content of the soil.*
- ❖ *About 35 per cent of area is medium (23-57 kg/ha) in available phosphorus content of the soil, 38 per cent of area is low (<23 kg/ha) and 10 per cent of area is high (>57 kg/ha) in the microwatershed.*
- ❖ *Available potassium content is medium (145-337 kg/ha) in 45 per cent of area and high (>337 kg/ha) in 38 per cent of area in the microwatershed.*
- ❖ *Available sulphur is low (<10 ppm) in an entire area of the microwatershed.*
- ❖ *Available boron is high (>1.0 ppm) in an area of 18 per cent and medium (0.5-0.1 ppm) in an area of 65 per cent in the microwatershed.*
- ❖ *Available iron is sufficient (>4.5 ppm) in the entire area of the microwatershed.*
- ❖ *Available manganese and copper are sufficient in all the soils of the microwatershed.*
- ❖ *Available zinc is deficient (<0.6 ppm) in an area of 23 per cent and sufficient (>0.6 ppm) in an area of 61 per cent of the microwatershed.*
- ❖ *The land suitability for 29 major crops grown in the microwatershed were assessed and the areas that are highly suitable (S1) and moderately suitable (S2) are given below. It is however to be noted that a given soil may be suitable for various crops but what specific crop to be grown may be decided by the farmer looking to his capacity to invest on various inputs, marketing infrastructure, market price and finally the demand and supply position.*

Land suitability for various crops in the Microwatershed

Crop	Suitability Area in ha (%)		Crop	Suitability Area in ha (%)	
	Highly suitable (S1)	Moderately suitable (S2)		Highly suitable (S1)	Moderately suitable (S2)
<i>Sorghum</i>	135 (22)	94 (15)	<i>Guava</i>	-	77 (13)
<i>Maize</i>	77 (13)	151 (25)	<i>Sapota</i>	-	77 (13)
<i>Bajra</i>	77 (13)	151 (25)	<i>Pomegranate</i>	-	221 (36)
<i>Groundnut</i>	74 (12)	10 (2)	<i>Musambi</i>	131 (22)	90 (15)
<i>Sunflower</i>	131 (22)	90 (15)	<i>Lime</i>	131 (22)	90 (15)
<i>Redgram</i>	-	221 (36)	<i>Amla</i>	77 (13)	138 (23)
<i>Bengal gram</i>	131 (22)	13 (2)	<i>Cashew</i>	-	-
<i>Cotton</i>	131 (22)	16 (3)	<i>Jackfruit</i>	-	77 (13)
<i>Chilli</i>	77 (13)	151 (25)	<i>Jamun</i>	-	131 (22)
<i>Tomato</i>	77 (13)	7 (1)	<i>Custard apple</i>	205 (34)	23 (4)
<i>Brinjal</i>	77 (13)	7 (1)	<i>Tamarind</i>	-	131 (22)
<i>Onion</i>	77 (13)	7 (1)	<i>Mulberry</i>	-	77 (13)
<i>Bhendi</i>	77 (13)	151 (25)	<i>Marigold</i>	77 (13)	151 (25)
<i>Drumstick</i>	-	208 (34)	<i>Chrysanthemum</i>	77 (13)	151 (25)
<i>Mango</i>	-	-			

- ❖ *Apart from the individual crop suitability, a proposed crop plan has been prepared for the identified LMUs by considering only the highly and moderately suitable lands for different crops and cropping systems with food, fodder, fibre and other horticulture crops.*
- ❖ *Maintaining soil-health is vital for crop production and conserve soil and land resource base for maintaining ecological balance and to mitigate climate change. For this, several ameliorative measures have been suggested for these problematic soils like saline/alkali, highly eroded, sandy soils etc.,*
- ❖ *Soil and water conservation treatment plan has been prepared that would help in identifying the sites to be treated and also the type of structures required.*
- ❖ *As part of the greening programme, several tree species have been suggested to be planted in marginal and submarginal lands, field bunds and also in the hillocks, mounds and ridges. This would help in not only supplementing the farm income but also provide fodder and fuel and generate lot of biomass which would help in maintaining an ecological balance and also contribute to mitigating the climate change.*



INTRODUCTION

Soil being a vital natural resource on whose proper use depends the life supporting systems of a country and the socioeconomic development of its people. Soils provide food, fodder, fibre and fuel for meeting the basic human and animal needs. With the ever increasing growth in human and animal population, the demand on soil for more food and fodder production is on the increase. The area available for agriculture is about 51 per cent of the total geographical area and more than 60 per cent of the people are still dependant on agriculture for their livelihood. However, the capacity of a soil to produce is limited and the limits to the production are set by its intrinsic characteristics, agro-climatic setting, and, use and management. There is, therefore, tremendous pressure on land and water resources, which is causing decline in soil-health and stagnation in productivity. The soils have been degrading at an estimated rate of one million hectares per year and ground water levels have been receding at an alarming rate resulting in decline in the ground water resource. Further, land degradation has emerged as a serious problem which has already affected about 38 lakh ha of cultivated area in the State. Soil erosion alone has degraded about 35 lakh ha. Almost all the uncultivated areas are facing various degrees of degradation, particularly soil erosion; salinity and alkalinity has emerged as a major problem in more than 3.5 lakh ha in the irrigated areas of the State. Nutrient depletion and declining factor productivity is common in both rainfed and irrigated areas. The degradation is continuing at an alarming rate and there appears to be no systematic effort among the stakeholders to contain this process. In recent times, an aberration of weather due to climate change phenomenon has added another dimension leading to unpredictable situation to be tackled by the farmers.

In this critical juncture, the challenge before us is not only to increase the productivity per unit area which is steadily declining and showing a fatigue syndrome, but also to prevent or at least reduce the severity of degradation. If the situation is not reversed at the earliest, then the sustainability of the already fragile crop production system and the overall ecosystem will be badly affected in the state. Added to this, every year there is a significant diversion of farm lands and water resources for non-agricultural purposes. Thus, developing strategies to slow down the degradation process or reclaim the soils to normal condition and ensure sustainability of production system are the major issues today. This demands a systematic appraisal of our soil and land resources with respect to their extent, geographic distribution, characteristics, behaviour and use potential, which is very important for developing an effective land use and cropping systems for augmenting agricultural production on a sustainable basis.

The soil and land resource inventories made so far in Karnataka had limited utility because the surveys were of different types, scales and intensities carried out at different times with specific objectives. Hence, there is an urgent need to generate detailed site-

specific farm level database on various land resources for all the villages/watersheds in a time bound manner that would help to protect the valuable soil and land resources and also to stabilize the farm production. Therefore, the land resource inventory required for farm level planning is the one which investigates all the parameters which are critical for productivity *viz.*, soils, site characteristics like slope, erosion, gravelliness and stoniness, climate, water, topography, geology, hydrology, vegetation, crops, land use pattern, animal population, socio-economic conditions, infrastructure, marketing facilities and various schemes and developmental works of the government etc. From the data collected at farm level, the specific problems and potentials of the area can be identified and highlighted, conservation measures required for the area can be planned on a scientific footing, suitability of the area for various uses can be worked out and finally viable and sustainable land use options suitable for each and every land holding can be prescribed.

The Land Resource Inventory is basically done for identifying potential and problem areas, developing sustainable land use plans, estimation of surface run off and water harvesting potential, preparation of soil and water conservation plans, land degradation/desertification etc. The Bureau is presently engaged in developing an LRI methodology using high resolution satellite remote sensing data and Digital Elevation Model (DEM) data to prepare Landscape Ecological Units (LEU) map representing agro-ecosystem as a whole. The LEU is preferred over landform as the base map for LRI. LEU is the assemblage of landform, slope and land use. An attempt has already been made to upscale the soil resource information from 1:250000 and 1:50000 scale to the LEU map in Goa and in some other states.

The land resource inventory aims to provide site-specific database for Chandaraki-1 microwatershed in Yadgir Taluk and Yadgir District, Karnataka State for the Karnataka Watershed Development Department. The database was generated by using cadastral map of the village as a base along with high resolution IRS LISS IV and Cartosat-1 merged satellite imagery. Later, an attempt will be made to uplink this LRI data generated at 1:7920 scale under Sujala-III Project to the proposed Landscape Ecological Units (LEUs) map.

The study was organized and executed by the ICAR- National Bureau of Soil Survey and Land Use Planning, Regional Centre, Bangalore under Generation of Land Resource Inventory Data Base Component-1 of the Sujala-III Project funded by the World Bank.

GEOGRAPHICAL SETTING

2.1 Location and Extent

The Chandaraki-1 microwatershed is located in the northern part of Karnataka in Yadgir Taluk & District, Karnataka State (Fig.2.1). It comprises parts of Chandaraki and Madhepalli villages. It lies between $16^{\circ} 50'$ and $16^{\circ} 52'$ North latitudes and $77^{\circ} 26'$ and $77^{\circ} 27'$ East longitudes covering an area of about 607 ha. It is about 45 km southeast of Yadgir town and is surrounded by Madhepalli on the south and Chandaraki on the east, west and northern side.




Fig.2.1 Location map of Chandaraki-1 Microwatershed

2.2 Geology

Major rock formations observed in the microwatershed are granite gneiss (Figs.2.2). They are essentially pink to gray and are coarse to medium grained. They consist primarily of quartz, feldspar, biotite and hornblende. The gray granite gneisses are highly weathered, fractured and fissured upto a depth of about 10 m. Dolerite dykes and quartz

veins are common with variable width and found to occur in Chandaraki-1 microwatershed.



Fig.2.2 Granite and granite gneiss rocks

2.3 Physiography

Physiographically, the area has been identified as granite gneiss landscape based on geology. The area has been further subdivided into five landforms, *viz*; mounds/ridges, summits, side slopes and very gently sloping uplands, plains and valleys based on slope and its relief features. The elevation ranges from 541-648 m above MSL. The mounds and ridges are mostly covered by rock outcrops.

2.4 Drainage

The area is drained by several parallel streams like Bori, Amerja and Kanga which finally join the river Bhima along its course. Though, they are not perennial, during rainy season they carry large quantities of rain water. The microwatershed has only few small tanks which are not capable of storing the water that flows during the rainy season. Due to this, the ground water recharge is very much affected. This is reflected in the failure of many bore wells in the villages. If the available rain water is properly harnessed by constructing new tanks and recharge structures at appropriate places in the villages, then the drinking and irrigation needs of the area can be easily met. The drainage network is parallel to sub parallel and dendritic.

2.5 Climate

The Yadgir district lies in the northern plains of Karnataka and falls under semiarid tract of the state and is categorized as drought- prone with total annual rainfall of 866 mm (Table 2.1). Of the total rainfall, maximum of 652 mm is received during the south–west monsoon period from June to September, the north-east monsoon from October to early December contributes about 138 mm and the remaining 76 mm during the rest of the year. The summer season starts during the middle of February and

continues up to the first week of June. The period from December to the middle of February is the coldest season. December is the coldest month with mean daily maximum and minimum temperatures being 29.5⁰C and 10⁰C respectively. During peak summer, temperature shoots up to 45⁰C. Relative humidity varies from 26% in summer to 62% in winter. Rainfall distribution is shown in Figure 2.3. The average Potential Evapo-Transpiration (PET) is 141 mm and varies from a low of 81 mm in December to 199 mm in the month of May. The PET is always higher than precipitation in all the months except July, August and September. Generally, the Length of crop Growing Period (LGP) is 120-150 days and starts from 1st week of June to 4th week of October.

Table 2.1 Mean Monthly Rainfall, PET, 1/2 PET at Yadgir Taluk, Yadgir District

Sl. No.	Months	Rainfall	PET	1/2 PET
1	January	4.30	86.0	43.0
2	February	2.30	125.5	62.7
3	March	15.10	166.0	83.0
4	April	18.50	179.8	89.9
5	May	36.0	198.8	97.9
6	June	118.0	175.1	87.5
7	July	171.80	156.3	78.1
8	August	182.9	150.3	75.1
9	September	179.7	142.0	71.0
10	October	105.3	138.5	69.2
11	November	26.4	97.60	48.6
12	December	6.0	80.90	40.4
Total		866.3		




Fig 2.3 Rainfall distribution in Yadgir Taluk, Yadgir District

2.6 Natural Vegetation

The natural vegetation is sparse comprising few tree species, shrubs and herbs. The mounds, ridges and boulders occupy very sizeable area which is under thin to moderately thick forest vegetation. Still, there are some remnants of the past forest cover which can be seen in patches in some ridges and hillocks in the microwatershed (Fig 2.4).

Apart from the continuing deforestation, the presence of large population of goats, sheep and other cattle in the microwatershed is causing vegetative degradation of whatever little vegetation left in the area. The uncontrolled grazing has left no time for the regeneration of the vegetative cover. This leads to the accelerated rate of erosion on the hill slopes resulting in the formation of deep gullies in the foot slopes that eventually result in the heavy siltation of tanks and reservoirs in the microwatershed.



Fig 2.4 Natural vegetation of Chandaraki-1 microwatershed

2.7 Land Utilization

About 72 per cent area (Table 2.2) in Yadgir district is cultivated at present. An area of about 2 per cent is permanently under pasture, 20 per cent under current fallows and 6 per cent under non-agricultural land, and 5 per cent under currently barren. Forests occupy an area of about 7 per cent and the tree cover is in a very poor state. Most of the mounds, ridges and bouldery areas have very poor vegetative cover. Major crops grown in the area are sorghum, maize, cotton, sunflower, groundnut, Bengal gram, red gram and paddy. The cropping intensity is 120 per cent in the taluk. While carrying out land resource inventory, the land use/land cover particulars are collected from all the survey numbers and a current land use map of the microwatershed is prepared. The current land use map prepared shows the arable and non-arable lands, other land uses and different types of crops grown in the area. The current land use map of Chandaraki-1

microwatershed is presented in Fig.2.5. The different crops and cropping systems adopted in the microwatershed is presented in the Figures 2.6.

Table 2.2 Land Utilization in Yadgir District

Sl. No.	Agricultural land use	Area (ha)	Per cent
1.	Total geographical area	516088	-
2.	Total cultivated area	373617	72.4
3.	Area sown more than once	74081	14.3
4.	Cropping intensity	-	119.8
5.	Trees and grooves	737	0.14
6.	Forest	33773	6.54
7.	Cultivable wasteland	2385	0.46
8.	Permanent Pasture land	11755	2.28
9.	Barren land	27954	5.41
10.	Non- Agriculture land	29623	5.73
11.	Current Fallows	105212	20.4




Fig.2.5 Current Land Use map of Chandaraki-1 Microwatershed



Fig 2.6 Different Crops and Cropping Systems in Chandaraki-1 Microwatershed

SURVEY METHODOLOGY

The purpose of land resource inventory is to delineate similar areas (soil series and phases), which respond or expected to respond similarly for a given level of management. This was achieved in Chandaraki-1 microwatershed by the detailed study of all the soil characteristics (depth, texture, colour, structure, consistence, coarse fragments, porosity, soil reaction, soil horizons etc.) and site characteristics (slope of the land, erosion, drainage, occurrence of rock fragments etc.) followed by grouping of similar areas based on soil-site characteristics into homogeneous (management units) units, and showing the area extent and their geographic distribution on the microwatershed cadastral map. The detailed survey at 1:7920 scale was carried out in an area of 608 ha. The methodology followed for carrying out land resource inventory was as per the guidelines given in Soil Survey Manual (IARI, 1971; Soil Survey Staff, 2006; Natarajan *et al.*, 2015) which is briefly described below.

3.1 Base Maps

The detailed survey of the land resources occurring in the microwatershed was carried out by using digitized cadastral map and satellite imagery as base supplied by KRSRSAC. The cadastral map shows field boundaries with their survey numbers, location of tanks, streams and other permanent features of the area (Fig. 3.1). Apart from the cadastral map, remote sensing data products from Cartosat-1 and LISS IV merged at the scale of 1:7920 were used in conjunction with the cadastral map to identify the landscapes, landforms and other surface features. The imagery helped in the identification and delineation of boundaries between hills, uplands and lowlands, water bodies, forest and vegetated areas, roads, habitations and other cultural features of the area (Fig. 3.2). The cadastral map was overlaid on the satellite imagery (Fig. 3.3) that helps to identify the parcel boundaries and other permanent features. Apart from cadastral maps and images, toposheets of the area (1:50,000 scale) were also used for initial traversing, identification of geology and landforms, drainage features, present land use and also for selection of transects in the microwatershed.

3.2 Image Interpretation for Physiography

False Colour Composites (FCCs) of Cartosat-I and LISS-IV merged satellite data covering microwatershed area was visually interpreted using image interpretation elements and all the available collateral data with local knowledge. The delineated physiographic boundaries were transferred on to a cadastral map overlaid on satellite imagery. Physiographically, the area has been identified as granite gneiss landscape. It was divided into five landforms, *viz.*; ridges and mounds, gently and very gently sloping uplands and lowlands based on slope and image characteristics. They were further

subdivided into physiographic/image interpretation units based on image characteristics. The image interpretation legend for physiography is given below.

Image Interpretation Legend for Physiography

G- Granite Gneiss Landscape

G1	Hills/ Ridges/ Mounds
G11	Summits
G12	Side slopes
G121	Side slopes with dark grey tones
G2	Uplands
G21	Summits
G22	Gently sloping uplands
G221	Gently sloping uplands, yellowish green (eroded)
G222	Gently sloping uplands, yellowish white (severely eroded)
G23	Very gently sloping uplands
G231	Very gently sloping uplands, yellowish green
G232	Very gently sloping uplands, medium green and pink
G233	Very gently sloping uplands, pink and green (scrub land)
G234	Very gently sloping uplands, medium greenish grey
G235	Very gently sloping uplands, yellowish white (eroded)
G236	Very gently sloping uplands, dark green
G237	Very gently sloping uplands, medium pink (coconut garden)
G238	Very gently sloping uplands, pink and bluish white (eroded)
G24	Valleys/ lowlands
G241	Valleys, pink tones
G242	Valleys gray mixed with pink tones




Fig 3.1 Scanned and Digitized Cadastral map of Chandaraki-1 Microwatershed




Fig.3.2 Satellite Image of Chandaraki-1 Microwatershed




Fig.3.3 Cadastral map overlaid on IRS PAN+LISS IV merged imagery of Chandaraki-1 Microwatershed

3.3 Field Investigation

The field boundaries and survey numbers given on the cadastral sheet were located on the ground by following permanent features like roads, cart tracks, *nallas*, streams, tanks etc., and wherever changes were noticed, they were incorporated on the microwatershed cadastral map. Preliminary traverse of the microwatershed was carried out with the help of cadastral map, imagery and toposheets. While traversing, landforms and physiographic units identified were checked and preliminary soil legend was prepared by studying soils at few selected places. Then, intensive traversing of each physiographic unit like hills, ridges, uplands and valleys was carried out. Based on the variability observed on the surface, transects (Fig. 3.4) were selected across the slope covering all the landform units in the microwatershed (Natarajan and Dipak Sarkar, 2010).




Fig: 3.4. Location of profiles in a transect

In the selected transect, soil profiles were located (Fig. 3.4) at closely spaced intervals to take care of any change in the land features like break in slope, erosion, gravel, stones etc. In the selected sites, soil profiles (vertical cut showing the soil layers from surface to the rock) were opened upto 200 cm or to the depth limited by rock or hard substratum and studied in detail for all their morphological and physical characteristics. The soil and site characteristics were recorded for all profile sites on a standard proforma as per the guidelines given in USDA Soil Survey Manual (Soil Survey Staff, 2012). Apart from the transect study, profiles were also studied at random, almost like in a grid pattern, outside the transect areas.

Based on the soil characteristics, the soils were grouped into different soil series. Soil series is the most homogeneous unit having similar horizons and properties and behaves similarly for a given level of management. Soil depth, texture, colour, kind of horizon and horizon sequence, calcareousness, amount and nature of gravel present, nature of substratum etc, were used as the major differentiating characteristics for identifying soil series occurring in the area. The differentiating characteristics used for identifying the soil series are given in Table 3.1. Based on the above characteristics, 9 soil series were identified in the Chandaraki-1 microwatershed.

Table 3.1 Differentiating Characteristics used for identifying Soil Series
(Characteristics are of Series Control Section)

Soils of Granite gneiss Landscape							
Sl. no	Soil Series	Depth (cm)	Colour (moist)	Texture	Gravel (%)	Horizon sequence	Calcareousness
1	BDP (Baddeppalli)	<25	7.5YR 3/2,3/4 5YR 3/4	scl	-	Ap-Ac	es
2	KKR (Kakalawar)	<25	7.5YR 4/3 10YR 6/3	sl	10-15	Ap-AC	-
3	BDL (Badiyala)	25-50	7.5YR 2.5/3,2.5/2,3/3 10YR 3/4,4/3	sl	-	Ap-Bw	e
4	HTK (Hattikuni)	25-50	10YR4/6,4/4 7.5YR4/4,3/3	sl	10-25	Ap-AC	-
5	JNK (Jinkera)	50-75	10YR3/1,3/2 7.5YR3/4	scl	-	Ap-Bw	e
6	SHT (Shettalli)	75-100	10YR 3/1	gsc	15-35	Ap-Bw	e
7	HSL (Hosalli)	75-100	10YR 5/4,4/4,4/6	sc	-	Ap-Bw	e
8	BGD (Belagundi)	100-150	10YR 5/4,4/4 7.5YR4/4	c	-	Ap-AB- Bss	es
9	BMN (Bhimanahalli)	>150	10YR 3/1	c	-	Ap-Bss	es

3.4 Soil Mapping

The area under each soil series was further separated into soil phases and their boundaries delineated on the cadastral map based on the variations observed in the texture of the surface soil, slope, erosion, presence of gravel, stoniness etc. A soil phase is a subdivision of soil series based mostly on surface features that affect its use and management. The soil mapping units are shown on the map (Fig.3.5) in the form of symbols. During the survey many soil profile pits, few minipits and a few auger bores representing different landforms occurring in the microwatershed were studied. In addition to the profile study, spot observations in the form of minipits, road cuts, terrace cuts etc., were studied to validate the soil boundaries on the soil map. The soil map shows the geographic distribution of 13 mapping units representing 9 soil series occurring in the microwatershed. The soil map unit (soil legend) description is presented in Table 3.2. The soil phase map (management units) shows the distribution of 13 soil phases mapped in the microwatershed. Each mapping unit (soil phase) delineated on the map has similar soil and site characteristics. In other words, all the farms or survey numbers included in one soil phase will have similar management needs and have to be treated accordingly.

3.5 Land Management Units (LMU's)

The 13 soil phases identified and mapped in the microwatershed were grouped into 4 Land Management Units (LMU's) for the purpose of preparing a Proposed Crop Plan for sustained development of the microwatershed. The database (soil phases)

generated under LRI was utilized for identifying Land Management Units (LMU's) based on the management needs. One or more than one soil site characteristic having influence on the management have been chosen for identification and delineation of LMUs. For Chandaraki-1 microwatershed, five soil and site characteristics, namely soil depth, soil texture, slope, erosion and gravel content have been considered for defining LMUs. The Land Management Units are expected to behave similarly for a given level of management.

3.6 Laboratory Characterization

Soil samples for each soil series were collected from representative master profiles for laboratory characterization by following the methods outlined in the Laboratory Manual (Sarma *et al*, 1987). Surface soil samples collected from farmer's fields for fertility status (major and micronutrients) at 320 m grid interval in the year 2018 were analyzed in the laboratory (Katyral and Rattan, 2003). By linking the soil fertility data to the survey numbers through GIS, soil fertility maps were generated by using Kriging method for the microwatershed.

Table 3.2 Soil map unit description of Chandaraki-1 Microwatershed

*Soil map unit No.	Soil Series	Soil Phase	Mapping Unit Description	Area in ha(%)
Soils of Granite and Granite Gneiss Landscape				
	BDP		Baddeppalli soils are very shallow (<25 cm), well drained, have dark brown to dark reddish brown, calcareous sandy clay loam soils occurring on very gently sloping uplands under cultivation	2 (0.3)
119		BDPiB3	Sandy clay surface, slope 1-3%, severe erosion	2 (0.3)
	KKR		Kakalawar soils are very shallow (<25 cm), well drained, have dark brown sandy loam soils occurring on very gently sloping uplands under cultivation	0.17 (0.03)
153		KKRbB2g1	Loamy sand surface, slope 1-3%, moderate erosion, gravelly (15-35%)	0.17(0.03)
	BDL		Badiyala soils are shallow (25-50 cm), well drained, have dark brown to very dark brown and dark yellowish brown, slightly calcareous sandy loam soils occurring on very gently to gently sloping uplands under cultivation	269 (44.98)
4		BDLhB2	Sandy clay loam surface, slope 1-3%, moderate erosion	0.24 (0.04)
162		BDLhB2g1	Sandy clay loam surface, slope 1-3%, moderate erosion, gravelly (15-35%)	20 (3.29)
5		BDLiB2	Sandy clay surface, slope 1-3%, moderate erosion	249 (40.95)
	HTK		Hattikuni soils are shallow (25-50 cm), well drained, have dark yellowish brown sandy loam soils	7 (1.08)

*Soil map unit No.	Soil Series	Soil Phase	Mapping Unit Description	Area in ha(%)
			occurring on very gently sloping uplands under cultivation	
165		HTKcB2	Sandy loam surface, slope 1-3%, moderate erosion	7 (1.08)
	JNK		Jinkera soils are moderately shallow (50-75 cm), well drained, have dark brown to very dark grayish brown, slightly calcareous sandy clay loam soils occurring on very gently sloping uplands under cultivation	7 (1.12)
23		JNKiB2g1	Sandy clay surface, slope 1-3%, moderate erosion, gravelly (15-35%)	6 (1.03)
24		JNKiB3g1	Sandy clay surface, slope 1-3%, severe erosion, gravelly (15-35%)	1 (0.09)
	SHT		Shettalli soils are moderately deep (75-100 cm), well drained, have very dark gray, slightly calcareous gravelly sandy clay loam soils occurring on very gently sloping uplands under cultivation	74 (12.11)
112		SHTmB2	Clay surface, slope 1-3%, moderate erosion	74 (12.11)
	HSL		Hosalli soils are moderately deep (75-100 cm), moderately well drained, have yellowish brown to dark yellowish brown, slightly calcareous sandy clay soils occurring on very gently sloping uplands under cultivation	3 (0.53)
33		HSLiB2	Sandy clay surface, slope 1-3%, moderate erosion	3 (0.53)
	BGD		Belagundi soils are deep (100-150 cm) well drained, have brown to dark yellowish brown, calcareous cracking clay soils occurring on very gently sloping uplands under cultivation	132 (21.61)
115		BGDmB2	Clay surface, slope 1-3%, moderate erosion	40 (6.53)
151		BGDmB2g1	Clay surface, slope 1-3%, moderate erosion, gravelly (15-35%)	92 (15.08)
	BMN		Bhimanahalli soils are very deep (>150 cm), moderately well drained, have very dark gray, calcareous cracking clay black soils occurring on very gently sloping uplands under cultivation	13 (2.08)
62		BMNmB2	Clay surface, slope 1-3%, moderate erosion	13 (2.08)
999		Rock outcrops	Rock lands, both massive and bouldery with little or no soil	1 (0.19)
1000		Others	Habitation and waterbody	101 (16.65)




Fig 3.5 Soil Phase or Management Units - Chandaraki-1 Microwatershed

THE SOILS

Detailed information pertaining to the nature, extent and their distribution of different kinds of soils occurring in Chandaraki-1 microwatershed is provided in this chapter. The microwatershed area has been identified as granite gneiss landscape based on geology. In all, 15 soil series are identified. Soil formation is the result of the combined effect of environmental and terrain factors that are reflected in soil morphology. In the granite gneiss landscape, it is by parent material, relief and climate.

A brief description of each of the 15 soil series identified followed by 23 soil phases (management units) mapped are furnished below. The physical and chemical characteristics of soil series identified in Chandaraki-1 microwatershed are given in Table 4.1 along with soil classification. The soils in any one map unit differ from place to place in their depth, texture, slope, gravelliness, erosion or any other site characteristic that affect management. The soil phase map can be used for identifying the suitability of areas for growing specific crops or for other alternative uses and also for deciding the type of conservation structures needed. The detailed information on soil and site-characteristics like soil depth, surface soil texture, slope, erosion, gravelliness, AWC, LCC etc, with respect to each of the soil phase identified is given village/survey number wise for the microwatershed in Appendix-I.

4.1 Soils of granite gneiss landscape

In this landscape, 9 soil series are identified and mapped. Of these, MDG series occupies a maximum area of 130 ha (17%) followed by BDP 99 ha (13%), BDL 93 ha (12%), VNK 83 ha (11%), ANR 77 ha (10%), SHT 73 ha (9%), YLR 47 ha (6%), YDR 32 ha (4%), MDR 30 (4%), HTK 7 ha (<1%), DSB 3 ha (<1%), VKS 3 ha (<1%), SBR 2 ha (<1), JNK 0.47 ha (<1%) and BMN 0.18 ha (<1). Brief description of each series identified and number of soil phases mapped is given below.

4.1.1 Baddeppalli (BDP) Series: Baddeppalli soils are very shallow (<25cm), well drained, have dark brown to dark reddish brown, calcareous sandy clay loam soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation. The Baddeppalli series has been classified as a member of the loamy, mixed (calcareous), isohyperthermic family of Lithic Ustorthents.

The thickness of the soil is less than 25 cm. Its colour is in 7.5 YR and 5 YR hue with value 3 and chroma 2 to 4. The texture varies from sandy clay loam to sandy clay and is calcareous. The available water capacity is very low (<50 mm/m). Four phases were identified and mapped.



Landscape and Soil Profile characteristics of Baddeppalli (BDP) Series

4.1.3 Badiyala (BDL) Series: Badiyala soils are shallow (25-50 cm), well drained, have very dark brown to dark yellow brown and dark brown, slightly calcareous sandy loam soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation. The Badiyala series has been classified as a member of the coarse-loamy, mixed, isohyperthermic family of Fluventic Haplustepts.

The thickness of the solum ranges from 28 to 50 cm. The thickness of A horizon ranges from 4 to 12 cm. Its colour is in 10YR hue with value 3 to 4 and chroma 3 to 4. The texture is loamy sand, sandy clay loam and sandy clay. The thickness of B horizon ranges from 27 to 45 cm. Its colour is in 10 YR and 7.5 YR hue with value 2 to 4 and chroma 3 to 4. Its texture is sandy loam to sandy clay loam and is slightly calcareous. The available water capacity is very low (<50mm/m). Only one phase was identified and mapped.



Landscape and Soil Profile characteristics of Badiyala (BDL) Series

4.1.4 Hattikuni (HTK) Series: Hattikuni soils are shallow (25-50 cm), well drained, have dark brown to dark yellowish brown sandy loam soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation. The Hattikuni series has been classified as a member of the mixed, isohyperthermic family of Lithic Ustipsamments.

The thickness of the soil ranges from 36 to 50 cm. The thickness of A horizon ranges from 8 to 12 cm. Its colour is in 10YR and 7.5 YR hue with value 3 to 4 and chroma 4 to 6. The texture varies from loamy sand to sandy loam. The thickness of subsurface horizon ranges from 28 to 42 cm. Its colour is in 10 YR and 7.5 YR hue with value 3 to 4 and chroma 4 to 6. Its texture varies from loamy sand to sand and sandy loam. The available water capacity is very low (<50 mm/m). Only one phase was identified and mapped.



Landscape and Soil Profile characteristics of Hattikuni (HTK) Series

4.1.5 Jinkera (JNK) Series: Jinkera soils are moderately shallow (50-75 cm), well drained, have very dark gray to very dark grayish brown and dark brown, slightly calcareous sandy clay loam soils. They are developed from weathered granite gneiss and occur on very gently sloping uplands under cultivation. The Jinkera series has been classified as a member of the fine-loamy, mixed, isohyperthermic family of Typic Haplustepts.

The thickness of the solum ranges from 51-75 cm. Thickness of A horizon ranges from 6 to 11 cm. Its colour is in hue 10 YR and 7.5 YR with value and chroma of 3 to 4. The texture varies from sandy loam to sandy clay. The thickness of B horizon ranges from 53 to 66 cm. Its colour is in 10 YR and 7.5 YR hue with value and chroma of 2 to 4. The texture varies from sandy clay loam to sandy clay and is slightly calcareous. The available water capacity is low (51-100 mm/m). Only one phase was identified and mapped.



Landscape and Soil Profile characteristics of Jinkera (JNK) Series

4.1.6 Shettalli (SHT) Series: Shettalli soils are moderately deep (75-100 cm), well drained, have very dark gray, slightly calcareous gravelly sandy clay soils. They are developed from weathered granite gneiss and occur on very gently sloping uplands under cultivation. The Shettalli series has been classified as a member of the fine, mixed, isohyperthermic family of Typic Haplustepts.

The thickness of the solum ranges from 78 to 100 cm. The thickness of A horizon ranges from 7 to 12 cm. Its colour is in hue 7.5 YR with value and chroma of 3 to 4. Its texture varies from sandy loam to sandy clay with 20 per cent gravel. The thickness of B horizon ranges from 68 to 92 cm. Its colour is in hue 7.5 YR with value 2 to 4 and chroma 1 to 3. Its texture is sandy clay with 15-35 per cent gravel and is slightly calcareous. The available water capacity is low (51-100 mm/m). Two phases were identified and mapped.



Landscape and Soil Profile characteristics of Shettalli (SHT) Series

4.1.7 Hosalli (HSL) Series: Hosalli soils are moderately deep (75-100 cm), moderately well drained, have dark yellowish brown to yellowish brown, slightly calcareous sandy clay soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation. The Hosalli series has been classified as a member of the fine, mixed, isohyperthermic family of Typic Haplustepts.

The thickness of the solum ranges from 76 to 100 cm. The thickness of A horizon ranges from 6 to 15 cm. Its colour is in hue 10 YR and 7.5 YR with value 3 to 5 and chroma 2 to 4. Its texture varies from loamy sand to sandy loam and sandy clay loam. The thickness of B horizon ranges from 62 to 93 cm. Its colour is in hue 10 YR with value 3 to 4 and chroma 2 to 4. Its texture varies from sandy clay loam to sandy clay and clay and is slightly calcareous. The available water capacity is medium (101-150 mm/m). Two phases were identified and mapped.



Landscape and Soil Profile characteristics of Hosalli (HSL) Series

4.1.8 Belagundi (BGD) Series: Belagundi soils are deep (100-150 cm), moderately well drained, have dark yellowish brown to yellowish brown and dark brown calcareous cracking clay soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation. The Belagundi series has been classified as a member of the very fine, smectitic (calcareous), isohyperthermic family of Typic Haplusterts.

The thickness of the solum ranges from 100 to 145 cm. The thickness of A horizon ranges from 5 to 12 cm. Its colour is in 10 YR and 5 YR hue with value 5 and chroma 2 to 4. The texture varies from sandy to loamy sand. The thickness of B horizon ranges from 95 to 135 cm. Its colour is in 10 YR and 7.5 YR hue with value 4 to 5 and chroma 4. Texture is sandy clay to clay and is slightly calcareous. The available water capacity is very high (>200 mm/m). Only one phase was identified and mapped.



Landscape and Soil Profile characteristics of Belagundi (BGD) Series

4.1.9 Bhimanahalli (BMN) Series: Bhimanahalli soils are very deep (>150 cm), moderately well drained, have very dark gray calcareous cracking clay soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation. The Bhimanahalli series has been classified as a member of the fine, smectitic (calcareous), isohyperthermic family of Typic Haplusterts.

The thickness of the solum is more than 150 cm. The thickness of A horizon ranges from 6 to 13 cm. Its colour is in 10 YR hue with value 3 to 4 and chroma 1 to 2 with clay texture. The thickness of B horizon ranges from 163 to 176 cm. Its colour is in 10 YR hue with value 2 to 4 and chroma 1. Its texture is clay and is calcareous. The available water capacity is very high (>200 mm/m). Only one phase was identified and mapped.



Landscape and Soil Profile characteristics of Bhimanahalli (BMN) Series

Table: 4.1 Physical and Chemical Characteristics of Soil Series identified in Chandaraki-1 microwatershed

Soil Series: Baddeppalli (BDP) **Pedon:** R-11

Location: 16°43'84.4"N 77°14'06.4"E, Halagera village, Yadgir hobli, Yadgir taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bengaluru **Classification:** Loamy, mixed (calcareous), isohyperthermic Lithic Ustorthents

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-16	Ap	58.67	17.02	24.31	19.03	13.74	9.62	10.57	5.71	<15	scl	16.19	8.18

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO ₃	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP
	Water	CaCl ₂	M KCl				Ca	Mg	K	Na	Total				
0-16	8.58	-	-	0.262	1.60	7.67	-	-	0.24	0.06	-	18.10	0.74	100	0.35

Contd...

Soil Series: Kakalawar (KKR), Pedon: R-7

Location: 16°50'25.9"N 77°15'97.1"E, Yampada village, Gurumitkal hobli, Yadgir taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bengaluru **Classification:** Mixed, isohyperthermic, Lithic Ustipsamments

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-22	Ap	83.81	10.37	5.82	17.31	20.65	17.91	5.67	22.27	10-20	ls	9.77	4.65

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO ₃	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP
	Water	CaCl ₂	M KCl				Ca	Mg	K	Na	Total				
				dS m ⁻¹	%	%	cmol kg ⁻¹					%	%		
0-22	5.85	-	-	0.027	0.19	-	0.72	0.21	0.62	0.03	1.58	2.6	0.45	60.90	1.17

Contd...

Soil Series: Badiyala (BDL) **Pedon:** R-5

Location: 16°37'10.0"N 77°20'21.5", Gudalagunta village, Balichakra hobli, Yadgir taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bengaluru **Classification:** Coarse-loamy, mixed, isohyperthermic Fluventic Haplustepts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-12	Ap	87.13	7.04	5.83	10.03	24.32	23.61	23.51	5.67	<15	ls	6.27	2.44
12-28	Bw1	64.63	13.30	22.07	6.74	13.07	22.30	17.01	5.50	<15	scl	16.34	7.83
28-50	BC	73.11	12.02	14.87	3.93	16.03	26.89	18.41	7.86	<15	sl	12.94	5.47

Depth (cm)	pH (1:2.5)			E.C. (1:2.5) dS m ⁻¹	O.C. %	CaCO ₃ %	Exchangeable bases					CEC	CEC/Clay	Base saturation %	ESP %
	Water	CaCl ₂	M KCl				Ca	Mg	K	Na	Total				
0-12	6.20	-	-	0.074	1.00	0.00	2.80	0.98	0.14	0.01	3.92	4.20	0.72	93	0.20
12-28	9.04	-	-	0.253	0.80	3.20	-	-	0.16	0.69	-	16.90	0.77	100	4.09
28-50	9.41	-	-	0.364	1.10	3.60	-	-	0.16	1.39	-	11.10	0.75	100	12.52

Contd...

Soil Series: Hattikuni (HTK), Pedon: R-7

Location: 16°50'46.5"N 77°10'16.4"E, Yaddalli village, Hattikuni hobli, Yadgir taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bengaluru **Classification:** Mixed, isohyperthermic Lithic Ustipsamments

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-12	Ap	90.89	5.62	3.49	8.50	13.46	29.86	29.55	9.51	20	s	7.73	3.16
12-22	A1	89.97	6.53	3.50	7.19	13.48	29.48	29.79	10.03	20	s	8.00	3.05
22-45	A2	87.20	6.43	6.38	11.09	14.42	31.55	7.16	22.98	40	ls	7.67	3.96

Depth (cm)	pH (1:2.5)			E.C. (1:2.5) dS m ⁻¹	O.C. %	CaCO ₃ %	Exchangeable bases					CEC	CEC/Clay	Base saturation %	ESP %
	Water	CaCl ₂	M KCl				Ca	Mg	K	Na	Total				
							cmol kg ⁻¹								
0-12	6.81	-	-	0.062	0.07	-	2.35	0.50	0.16	0.01	3.02	3.0	0.86	100	0.38
12.0-22	6.80	-	-	0.050	0.21	-	1.67	0.30	0.09	0.01	2.07	2.4	0.69	86.30	0.45
22-45	6.85	-	-	0.044	0.19	-	1.82	0.42	0.10	0.06	2.40	2.6	0.41	92.41	2.17

Contd...

Soil Series: Jinkera (JNK) **Pedon:** R-1

Location: 16°45'13.5"N 77°10'59.8"E, Varkanahalli village, Yadgir hobli, Yadgir taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bengaluru **Classification:** Fine-loamy, mixed, isohyperthermic Typic Haplustepts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-15	Ap	66.84	13.62	19.54	12.15	21.22	11.23	12.56	9.68	10	sl	14.42	7.70
15-38	Bw1	59.08	12.11	28.81	12.53	12.42	17.85	8.77	7.52	20	scl	18.21	12.23
38-52	Bw2	68.21	11.68	20.11	17.90	21.81	10.60	10.80	7.10	10	scl	14.54	8.96

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO ₃	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP
	Water	CaCl ₂	M KCl				Ca	Mg	K	Na	Total				
				dS m ⁻¹	%	%	cmol kg ⁻¹						%	%	
0-15	8.42	-	-	0.148	0.70	0.65	-	-	0.15	0.03	-	14.50	0.74	100	0.18
15-38	8.38	-	-	0.226	0.31	2.21	-	-	0.09	0.23	-	21.70	0.75	100	1.05
38-52	8.40	-	-	0.195	0.25	1.17	-	-	0.07	0.19	-	15.90	0.79	100	1.23

Contd...

Soil Series: Shettalli (SHT) **Pedon:** R-14

Location: 16°47'21.1"N 77°04'91.1"E, Thumakura village, Yadgir hobli, Yadgir taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bengaluru **Classification:** Fine, mixed, isohyperthermic Typic Haplustepts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-14	Ap	74.39	10.89	14.73	5.64	8.30	21.00	28.89	10.55	50	sl	12.58	4.51
14-35	Bw1	54.37	14.73	30.90	3.58	5.90	15.38	21.71	7.80	25	scl	20.37	10.92
35-63	Bw2	41.16	20.63	38.21	1.71	1.71	10.61	13.61	13.50	30	cl	24.34	15.03
63-83	Bw3	36.96	21.52	41.51	4.31	5.28	8.94	12.39	6.03	35	c	24.76	16.17

Depth (cm)	pH (1:2.5)			E.C. (1:2.5) dS m ⁻¹	O.C. %	CaCO ₃ %	Exchangeable bases					CEC	CEC/Clay	Base saturation %	ESP %
	Water	CaCl ₂	M KCl				Ca	Mg	K	Na	Total				
0-14	7.26	-	-	0.199	0.91	0.13	-	-	0.28	0.09	-	10.60	0.72	100	0.86
14-35	7.05	-	-	0.051	0.80	1.17	-	-	0.12	0.09	-	18.20	0.59	100	0.48
35-63	7.67	-	-	0.238	0.70	2.86	-	-	0.14	0.16	-	24.40	0.64	100	0.64
63-83	8.67	-	-	0.142	0.20	12.48	-	-	0.13	0.23	-	27.40	0.66	100	0.84

Contd...

Soil Series: Hosalli (HSL) **Pedon:** R-3

Location: 16°46'60.3"N 77°05'47.6"E, Mudhanala village, Yadgir hobli, Yadgir taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bengaluru **Classification:** Fine, mixed, isohyperthermic Typic Haplustepts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-10	Ap	88.43	5.15	6.42	5.69	6.40	36.04	27.31	12.99	-	s	7.40	2.74
10-30	Bw1	58.47	7.24	34.29	4.26	9.37	19.91	19.28	5.64	-	scl	19.07	11.57
30-50	Bw2	51.43	12.67	35.90	3.49	8.89	16.72	15.87	6.46	<15	sc	21.64	12.44
50-90	Bw3	49.89	13.64	36.47	2.43	2.96	20.61	16.17	7.72	<15	sc	21.12	12.95

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO ₃	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP			
	Water	CaCl ₂	M KCl				dS m ⁻¹	%	%	Ca	Mg					K	Na	Total
										cmol kg ⁻¹								
0-10	7.16	-	-	0.117	0.48	0.00	2.83	1.50	0.15	0.29	4.76	4.90	0.76	97	5.94			
10-30	6.91	-	-	0.040	0.36	0.00	10.64	5.43	0.10	0.26	16.43	17.80	0.52	92	1.47			
30-50	8.17	-	-	0.182	0.24	1.43	-	-	0.12	0.22	-	19.90	0.55	100	1.08			
50-90	8.60	-	-	0.148	0.20	4.29	-	-	0.13	0.16	-	19.70	0.54	100	0.81			

Contd...

Soil Series: Belagundi (BGD) **Pedon:** T₁/P₂

Location: 16°31'65.3"N 77°20'84.9"E, Kadechoora village, Sydhapura hobli, Yadgir taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bengaluru **Classification:** Very fine, smectitic (calcareous), isohyperthermic Typic Haplusterts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-13	Ap	14.90	17.83	67.27	0.77	2.10	2.65	5.96	3.42	-	c	43.97	29.27
13-40	AB	13.07	18.32	68.61	0.80	2.05	2.61	4.20	3.41	-	c	41.23	30.48
40-80	Bss1	11.68	17.18	71.13	0.80	2.06	2.29	3.32	3.21	-	c	46.72	32.41
80-113	Bss2	12.17	16.53	71.30	1.95	1.61	3.21	2.41	2.99	-	c	46.87	35.13

Depth (cm)	pH (1:2.5)			E.C. (1:2.5) dS m ⁻¹	O.C. %	CaCO ₃ %	Exchangeable bases					CEC	CEC/Clay	Base saturation %	ESP %
	Water	CaCl ₂	M KCl				Ca	Mg	K	Na	Total				
0-13	7.85	-	-	0.253	0.87	5.20	-	-	0.67	0.17	-	65.90	0.98	100	0.26
13-40	8.11	-	-	0.172	0.74	4.29	-	-	0.31	0.16	-	66.70	0.97	100	0.23
40-80	8.44	-	-	0.205	0.58	5.59	-	-	0.20	0.27	-	66.30	0.93	100	0.40
80-113	8.82	-	-	0.201	0.39	10.14	-	-	0.19	0.17	-	63.80	0.89	100	0.27

Contd...

Soil Series: Bhimanahalli (BMN) **Pedon:** R-3

Location: 16°31'82.4"N 77°12'70.8"E, Bheemanahalli village, Sydhapura hobli, Yadgir taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bengaluru **Classification:** Fine, smectitic (calcareous), isohyperthermic Typic Haplusterts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-8	Ap	20.34	19.94	59.72	2.68	5.03	3.75	5.25	3.64	-	c	50.19	33.49
8-40	Bss1	19.61	22.76	57.62	1.94	2.59	5.28	4.96	4.85	-	c	43.22	29.05
40-70	Bss2	21.25	17.65	61.10	3.02	5.26	3.91	5.48	3.58	-	c	44.30	30.25
70-120	Bss3	19.08	22.29	58.63	1.75	5.04	3.84	5.15	3.29	-	c	43.26	30.31
120-170	Bss4	11.11	20.44	68.45	2.04	1.93	1.70	2.83	2.61	-	c	51.33	33.51

Depth (cm)	pH (1:2.5)			E.C. (1:2.5) dS m ⁻¹	O.C. %	CaCO ₃ %	Exchangeable bases					CEC	CEC/Clay	Base saturation %	ESP %
	Water	CaCl ₂	M KCl				Ca	Mg	K	Na	Total				
0-8	8.2	-	-	0.284	0.72	4.94	-	-	1.20	0.34	-	52.70	0.88	100	0.65
8-40	8.44	-	-	0.139	0.40	7.28	-	-	0.30	0.48	-	52.06	0.90	100	0.93
40-70	8.32	-	-	0.202	0.40	6.37	-	-	0.18	0.40	-	52.52	0.86	100	0.77
70-120	9.3	-	-	0.282	0.36	6.89	-	-	0.27	0.38	-	50.97	0.87	100	0.75
120-170	8.47	-	-	0.305	0.37	8.19	-	-	0.28	0.91	-	58.19	0.85	100	1.57

INTERPRETATION FOR LAND RESOURCE MANAGEMENT

The most important soil and site characteristics that affect the land use and conservation needs of an area are land capability, land irrigability, soil depth, soil texture, coarse fragments, available water capacity, soil slope, soil erosion, soil reaction etc. These are interpreted from the data base generated through land resource inventory and several thematic maps are generated. These would help in identifying the areas suitable for growing crops and, soil and water conservation measures and structures needed thus helping to maintain good soil health for sustained crop production. The various interpretative and thematic maps generated are described below.

5.1 Land Capability Classification

Land capability classification is an interpretative grouping of soil map units (soil phases) mainly based on inherent soil characteristics, external land features and environmental factors that limit the use of land for agriculture, pasture, forestry, or other uses on a sustained basis (IARI, 1971). The land and soil characteristics used to group the land resources in an area into various land capability classes, subclasses and units are

Soil Characteristics: Depth, texture, gravelliness, calcareousness.

Land characteristics: Slope, erosion, drainage, rock outcrops.

Climate: Total rainfall and its distribution, and length of crop growing period.

The land capability classification system is divided into land capability classes, subclasses and units based on the level of information available. Eight land capability classes are recognized. They are

Class I: They are very good lands that have no limitations or very few limitations that restrict their use.

Class II: They are good lands that have minor limitations and require moderate conservation practices.

Class III: They are moderately good lands that have moderate limitations that reduce the choice of crops or that require special conservation practices.

Class IV: They are fairly good lands that have very severe limitations that reduce the choice of crops or that require very careful management.

Class V: Soils in these lands are not likely to erode, but have other limitations like wetness that are impractical to remove and as such not suitable for agriculture, but suitable for pasture or forestry with minor limitations.

Class VI: The lands have severe limitations that make them generally unsuitable for cultivation, but suitable for pasture or forestry with moderate limitations.

Class VII: The lands have very severe limitations that make them unsuitable for cultivation, but suitable for pasture or forestry with major limitations.

Class VIII: Soil and other miscellaneous areas (rock lands) that have very severe limitations that nearly preclude their use for any crop production, but suitable for wildlife, recreation and installation of wind mills.

The land capability subclasses are recognised based on the dominant limitations observed within a given land capability class. The subclasses are designated by adding a lower case letter like ‘e’, ‘w’, ‘s’, or ‘c’ to the class numeral. The subclass “e” indicates that the main hazard is risk of erosion, “w” indicates drainage or wetness as a limitation for plant growth, “s” indicates shallow soil depth, coarse or heavy textures, calcareousness, salinity/alkalinity or gravelliness and “c” indicates limitation due to climate.

The land capability subclasses have been further subdivided into land capability units based on the kinds of limitations present in each subclass. Ten land capability units are used in grouping the soil map units. They are stony or rocky (0), erosion hazard (slope, erosion) (1), coarse texture (sand, loamy sand, sandy loam) (2), fine texture (cracking clay, silty clay) (3), slowly permeable subsoil (4), coarse underlying material (5), salinity/alkali (6), stagnation, overflow, high ground water table (7), soil depth (8) and fertility problems (9). The capability units thus identified have similar soil and land characteristics that respond similarly to a given level of management. The soils of the microwatershed have been classified upto land capability subclass level.

The 13 soil map units identified in Chandaraki-1 microwatershed are grouped under 3 land capability classes and 3 land capability subclasses. An area of about 505 ha (83%) in the microwatershed is suitable for agriculture. About 101 ha (17%) area is covered by others (water body & habitation), and 1 ha (<1%) is under rock outcrops (Fig. 5.1).

Good lands (Class II) cover an area of about 37 per cent and are distributed in the southern, western, southwestern, northwestern, northeastern and eastern part of the microwatershed with minor problems of soil and erosion. Moderately good lands (Class III) cover an area of about 45 per cent and are distributed in the northern, northwestern, southern, southeastern and central part of the microwatershed with moderate problems of soil and erosion. Fairly good (Class IV) lands occur in an area of about <1 per cent of the microwatershed and are distributed in the southern part of the microwatershed with very severe problems of soil and erosion.




Fig. 5.1 Land Capability map of Chandaraki-1 Microwatershed

5.2 Soil Depth

Soil depth refers to the depth of the soil occurring above the parent material or hard rock. The depth of the soil determines the effective rooting depth for plants and in accordance with soil texture, mineralogy and gravel content, the capacity of the soil column to hold water and nutrient availability. Soil depth is one of the most important soil characteristic that is used in differentiating soils into different soil series. The soil depth classes used in identifying soils in the field are very shallow (<25 cm), shallow (25-50 cm), moderately shallow (50-75 cm), moderately deep (75-100 cm), deep (100-150 cm) and very deep (>150 cm). They were used to classify the soils into different depth classes and a soil depth map was generated. The area extent and their geographical distribution in the microwatershed is given in Fig. 5.2.

Very shallow (<25 cm) soils occur in an area of 2 ha (<1%) and are distributed in the southern part of the microwatershed. Shallow (25-50 cm) soils occur in an area of 276 ha (45%) and are distributed in the central, northern, southeastern, northwestern and southern part of the microwatershed. Moderately shallow (50-75 cm) soils occur in an area of 7 ha (1%) and are distributed in the southern part of the microwatershed. Moderately deep (75-100 cm) soils occur in an area of 77 ha (13%) and are distributed in the northwestern, southwestern and western part of the microwatershed. Deep (100-150 cm) soils cover an area of 131 ha (22%) and are distributed in the northeastern, southeastern, northwestern and eastern part of the microwatershed. Very deep (>150 cm)

soils cover an area of 13 ha (2%) and are distributed in the northwestern part of the microwatershed.

The most productive lands covering 144 ha (24%) with respect to soil rooting depth where all climatically adapted annual and perennial crops can be grown are deep to very deep (100- >150 cm depth) soils. The problem soils occupy an area of 278 ha (46%) where only short duration crops can be grown occasionally and the probability of crop failure is very high.




Fig. 5.2 Soil Depth map of Chandaraki-1 Microwatershed

5.3 Surface Soil Texture

Texture is an expression to indicate the coarseness or fineness of the soil as determined by the relative proportion of primary particles of sand, silt and clay. It has a direct bearing on the structure, porosity, adhesion and consistence. The surface layer of a soil to a depth of about 25 cm is the layer that is most used by crops and plants. The surface soil textural class provides a guide to understanding soil-water retention and availability, nutrient holding capacity, infiltration, workability, drainage, physical and chemical behaviour, microbial activity and crop suitability. The textural classes used for LRI were used to classify and a surface soil texture map was generated. The area extent and their geographical distribution in the microwatershed is shown in Figure 5.3.

An area of 0.17 ha (<1%) of the microwatershed has sandy soils at the surface and are distributed in the southern part. An area of 27 ha (4%) of the microwatershed has loamy soils at the surface and are distributed in the southern part. An area of about 479 ha

(79%) of the microwatershed has soils that are clayey and are distributed in the major part. Both loamy and clay soils have high potential for soil-water retention and availability, and nutrient retention and availability, but clayey soils have more problems of drainage, infiltration, workability and other physical problems. Problem soils have limitations of moisture and nutrient availability but are suited for root or tuber crops.




Fig. 5.3 Surface Soil Texture map of Chandaraki-1 Microwatershed

5.4 Soil Gravelliness

Gravel is the term used for describing coarse fragments between 2 mm and 7.5 cm diameter and stones for those between 7.5 cm and 25 cm. The presence of gravel and stones in soil reduces the volume of soil responsible for moisture and nutrient storage, drainage, infiltration and runoff, and hinders plant growth by impeding root growth and seedling emergence, inter-cultural operations and farm mechanization. The gravelliness classes used in LRI were used to classify the soils and using these classes a gravelliness map was generated. The area extent and their geographic distribution in the microwatershed is shown in Figure 5.4.

Non gravelly (<15%) soils cover an area of 387 ha (64%) and are distributed in the major part of the microwatershed. These are the most productive soils, where all climatically adapted short and long duration crops can be grown. Gravelly (15-35%) soils occur in an area of 119 ha (20%) and distributed in the northeastern, eastern, southeastern and southern part of the microwatershed. These lands are low in moisture holding

capacity and hence growing of short duration crops is ideal with best management practice.




Fig. 5.4 Soil Gravelliness map of Chandaraki-1 Microwatershed

5.5 Available Water Capacity

The soil available water capacity (AWC) is estimated based on the ability of the soil column to retain water between the tensions of 0.33 and 15 bar in a depth of 100 cm or the entire solum if the soil is shallower. The AWC of the soils (soil series) as estimated by considering the soil texture, mineralogy, soil depth and gravel content (Sehgal *et al.*, 1990) and accordingly the soil map units were grouped into five AWC classes *viz*, very low (<50 mm/m), low (50-100 mm/m), medium (100-150 mm/m), high (150-200 mm/m) and very high (>200 mm/m) and using these values, an AWC map was generated. The area extent and their geographic distribution of different AWC classes in the microwatershed is given in Figure 5.5.

An area of about 278 ha (46%) in the microwatershed has soils that are very low (<50 mm/m) in available water capacity and is distributed in the northern, central, northwestern, southern and southeastern part of the microwatershed. An area of about 80 ha (13%) is low (51-100 mm/m) in available water capacity and are distributed in the western, northwestern and southwestern part of the microwatershed. An area of about 3 ha (<1%) is medium (101-150 mm/m) in available water capacity and are distributed in the southern part of the microwatershed Very high (>200 mm/m) in 144 ha (24%) and are

distributed in the eastern, southeastern, northwestern and northeastern part of the microwatershed.

An area of about 358 ha (59%) in the microwatershed has soils that are problematic with regard to available water capacity. Here, only short duration crops can be grown and probability of the crop failure is very high. These areas are best put to other alternative uses. An area of 144 ha (24%) are potential areas with regard to AWC where all climatically adapted annual and perennial crops can be grown.




Fig. 5.5 Soil Available Water Capacity map of Chandaraki-1 Microwatershed

5.6 Soil Slope

Soil slope refers to the inclination of the surface of the land. It is defined by gradient, shape and length, and is an integral feature of any soil as a natural body. Slope is considered important in soil genesis, land use and land development. The length and gradient of slope influences the rate of runoff, infiltration, erosion and deposition. The soil map units were grouped into single slope class and a slope map was generated showing the area extent and their geographic distribution in the microwatershed (Fig. 5.6).

Entire area in the microwatershed is under very gently sloping (1-3% slope) lands. In these areas, all climatically adapted annual and perennial crops can be grown without much soil and water conservation and other land development measures.




Fig. 5.6 Soil Slope map of Chandaraki-1 Microwatershed

5.7 Soil Erosion

Soil erosion refers to the wearing away of the earth's surface by the forces of water, wind and ice involving detachment and transport of soil by raindrop impact. It is used for accelerated soil erosion resulting from disturbance of the natural landscape by burning, excessive grazing and indiscriminate felling of forest trees and tillage, all usually by man. The erosion classes showing an estimate of the current erosion status as judged from field observations in the form of rills, gullies or a carpet of gravel on the surface are recorded. Four erosion classes, viz, slight erosion (e1), moderate erosion (e2), severe erosion (e3) and very severe erosion (e4) are recognized. The soil map units were grouped into different erosion classes and a soil erosion map generated. The area extent and their spatial distribution in the microwatershed is given in Figure 5.7.

Moderately eroded (e2 class) soils cover a maximum area of 503 ha (83%) and are distributed in the major part of the microwatershed. Severely eroded (e3) soils cover an area of 2 ha (<1%) and are distributed in the southern part of the microwatershed

Entire area of the microwatershed is problematic because of moderate and severe erosion. For these areas, taking up of soil and water conservation and other land development measures are needed.




Fig. 5.7 Soil Erosion map of Chandaraki-1 Microwatershed

FERTILITY STATUS

Soil fertility plays an important role in increasing crop yield. The adoption of high yielding varieties that require high amounts of nutrients has resulted in deficiency symptoms in crops and plants due to imbalanced fertilization and poor inherent fertility status as these areas are characterised by low rainfall and high temperatures. Hence, it is necessary to know the fertility (macro and micro nutrients) status of the soils of the watersheds for assessing the kind and amount of fertilizers required for each of the crop intended to be grown. For this purpose, the surface soil samples collected from the grid points (one soil sample at every 320 m interval) all over the microwatershed through land resource inventory in the year 2018 were analysed for pH, EC, organic carbon, available phosphorus and potassium, and for micronutrients like zinc, boron, copper, iron and manganese, and secondary nutrient sulphur.

Soil fertility data generated has been assessed and individual maps for all the nutrients for the microwatershed have been generated using Kriging method under GIS. The village/survey number wise fertility data for the microwatershed is given in Appendix-II.

6.1 Soil Reaction (pH)

The soil analysis of the Chandaraki-1 microwatershed for soil reaction (pH) showed that an area of about 174 ha (29%) is slightly acid (pH 6.0-6.5) and are distributed in the northern, central, southern, northeastern and eastern part. An area of about 26 ha (4%) is moderately acid (pH 5.5-6.0) and are distributed in the eastern part. An area of about 186 ha (31%) is neutral (pH 6.5-7.3) and are distributed in the northern, southern, central, southeastern, northeastern and western part. An area of about 61 ha (10%) is slightly alkaline (pH 7.3-7.8) and are distributed in the northern, western, central and northwestern part. An area of about 33 ha (6%) are moderately alkaline (pH 7.8-8.4) and are distributed in the western, northwestern and central part. An area of 25 ha (4%) area is strongly alkaline (pH 8.4-9.0) and are distributed in the northwestern part of the microwatershed (Fig. 6.1)

6.2 Electrical Conductivity (EC)

The Electrical Conductivity in the entire area of the microwatershed is <2 dS/m (Fig. 6.2) and as such the soils are non saline.

6.3 Organic Carbon

The soil organic carbon content (an index of available Nitrogen) is high ($>0.75\%$) in the entire area of the microwatershed (Fig. 6.3).




Fig.6.1 Soil Reaction (pH) map of Chandaraki-1 Microwatershed




Fig.6.2 Electrical Conductivity (EC) map of Chandaraki-1 Microwatershed




Fig.6.3 Soil Organic Carbon map of Chandaraki-1 Microwatershed

6.4 Available Phosphorus

Available phosphorus content is medium (23-57 kg/ha) in an area of 214 ha (35%) and distributed in the eastern, central, northern, western, southeastern and southwestern part of the microwatershed. High (>57 kg/ha) in an area of 63 ha (10%) and are distributed in the southern and western part and low (<23 kg/ha) in an area of 229 ha (38%) and are distributed in the southeastern, northwestern, central eastern, northeastern and northern part of the microwatershed (Fig. 6.4).

6.5 Available Potassium

Available potassium content is medium (145-337 kg/ha) in an area of 273 ha (45%) and are distributed in the northwestern, central, eastern, southeastern, southern and southwestern part of the microwatershed and high (>337 kg/ha) in an area of 232 ha (38%) and are distributed in the western, central, eastern, northern, northeastern and southeastern part of the microwatershed (Fig. 6.5)

6.6 Available Sulphur

Available sulphur content is low (<10 ppm) in the entire area of the microwatershed (Fig. 6.6).




Fig.6.4 Soil Available Phosphorus map of Chandaraki-1 Microwatershed

6.7 Available Boron

Available boron content is high (>1.0 ppm) in an area of 109 ha (18%) and are distributed in the western, southwestern and southern part of the microwatershed and medium (0.5-1.0 ppm) in an area of 396 ha (65%) and are distributed in the major part of the microwatershed (Fig. 6.7).

6.8 Available Iron

Available iron content is sufficient (>4.5 ppm) in the entire area of the microwatershed (Fig 6.8).

6.9 Available Manganese

Available manganese content is sufficient (>1.0 ppm) in the entire microwatershed area (Fig 6.9).

6.10 Available Copper

Available copper content is sufficient (>0.2 ppm) in the entire microwatershed area (Fig 6.10).

6.11 Available Zinc

Available zinc content is deficient (<0.6 ppm) in an area of 137 ha (23%) and are distributed in the southern, central, southeastern and southwestern part and sufficient (>0.6 ppm) in an area of 368 ha (61%) and are distributed in the major part of the microwatershed (Fig 6.11).




Fig.6.5 Soil Available Potassium map of Chandaraki-1 Microwatershed




Fig.6.6 Soil Available Sulphur map of Chandaraki-1 Microwatershed




Fig.6.7 Soil Available Boron map of Chandaraki-1 Microwatershed




Fig.6.8 Soil Available Iron map of Chandaraki-1 Microwatershed




Fig.6.9 Soil Available Manganese map of Chandaraki-1 Microwatershed




Fig.6.10 Soil Available Copper map of Chandaraki-1 Microwatershed




Fig.6.11 Soil Available Zinc map of Chandaraki-1 Microwatershed

LAND SUITABILITY FOR MAJOR CROPS

The soil and land resource units (soil phases) of Chandaraki-1 microwatershed were assessed for their suitability for growing food, fodder, fibre and other horticulture crops by following the procedure as outlined in FAO, 1976 and 1983. Crop requirements were developed for each of the crop from the available research data and also by referring to Naidu *et. al.* (2006) and Natarajan *et. al* (2015). The soil and land characteristics were matched with the crop requirement to arrive at the crop suitability. The soil and land characteristics (Table 7.1) table and crop requirement tables (Tables 7.2 to 7.30) are given at the end of the chapter. In FAO land suitability classification, two orders are recognized. Order S-Suitable and Order N-Not suitable. The orders have classes, subclasses and units. Order-S has three classes, Class S1-Highly Suitable, Class S2-Moderately Suitable and Class S3- Marginally Suitable. Order N has two classes, N1-Currently not Suitable and N2- Permanently not Suitable. There are no subclasses within the Class S1 as they will have very minor or no limitations for crop growth. Classes S2, S3, N1 and N2 are divided into subclasses based on the kinds of limitations encountered. The limitations that affect crop production are ‘c’ for erratic rainfall and its distribution and length of growing period (LGP), ‘e’ for erosion hazard, ‘r’ for rooting condition, ‘t’ for lighter or heavy texture, ‘g’ for gravelliness or stoniness, ‘n’ for nutrient availability, ‘l’ for topography, ‘m’ for moisture availability, ‘w’ for drainage, ‘s’ for sodium and ‘z’ for calcareousness. These limitations are indicated as lower case letters to the Class symbol. For example, moderately suitable lands with the limitations of soil depth and erosion are designated as S2re. For the microwatershed, the soil mapping units were evaluated and classified up to subclass level.

Using the above criteria, the soil map units of the microwatershed were evaluated and land suitability maps for 29 major agricultural and horticultural crops were generated. The detailed information on the kind of suitability of each of the soil phase for the crops assessed are given village/ survey number wise for the microwatershed in Appendix-III.

7.1 Land Suitability for Sorghum (*Sorghum bicolor*)

Sorghum is one of the major food crop grown in Karnataka in an area of 10.47 lakh ha in Bijapur, Gulbarga, Raichur, Bidar, Belgaum, Dharwad, Bellary, Chitradurga, Mysore and Tumakuru districts. The crop requirements for growing sorghum (Table 7.2) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing sorghum was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.1.

Highly suitable (Class S1) lands for growing sorghum occur in an area of 135 ha (22%) and are distributed in the northeastern, eastern, southeastern and northwestern part

of the microwatershed. An area of about 94 ha (15%) is moderately suitable (Class S2) for growing sorghum and is distributed in the northern, southern, western, northwestern and southwestern part of the microwatershed with minor limitations of rooting depth, calcareousness and texture. An area of about 276 ha (45%) is marginally suitable (Class S3) for growing sorghum and is distributed in the central, eastern, northern, northwestern, southeastern and southern part of the microwatershed with moderate limitations of rooting depth and texture. Currently not suitable (Class N1) lands occur in an area of 2 ha (<1%) and are distributed in the southern part of the microwatershed with severe limitation of rooting depth.




Fig. 7.1 Land Suitability map of Sorghum

7.2 Land Suitability for Maize (*Zea mays*)

Maize is one of the most important food crop grown in an area of 13.37 lakh ha in almost all the districts of the State. The crop requirements for growing maize (Table 7.3) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing maize was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.2.

Highly suitable (Class S1) lands for growing maize occur in an area of 77 ha (13%) and are distributed in the western and northwestern part of the microwatershed. An area of about 151 ha (25%) is moderately suitable (Class S2) for growing maize and is distributed in the northeastern, eastern, southern, northwestern and southeastern part of

the microwatershed with minor limitations of rooting depth, calcareousness and texture. An area of about 276 ha (45%) is marginally suitable (Class S3) for growing maize and is distributed in the central, eastern, northern, northwestern, southeastern and southern part of the microwatershed with moderate limitations of rooting depth and texture. Currently not suitable (Class N1) lands occur in an area of 2 ha (<1%) and are distributed in the southern part of the microwatershed with severe limitation of rooting depth.




Fig. 7.2 Land Suitability map of Maize

7.3 Land Suitability for Bajra (*Pennisetum glaucum*)

Bajra is one of the most important millet crop grown in an area of 2.34 lakh ha in the northern districts of Karnataka state. The crop requirements for growing bajra (Table 7.4) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing bajra was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.3.

Highly suitable (Class S1) lands for growing bajra occur in an area of 77 ha (13%) and are distributed in the western and northwestern part of the microwatershed. An area of about 151 ha (25%) is moderately suitable (Class S2) for growing bajra and is distributed in the northeastern, eastern, southern, northwestern and southeastern part of the microwatershed with minor limitations of rooting depth, calcareousness and texture. An area of about 276 ha (45%) is marginally suitable (Class S3) for growing bajra and is distributed in the central, eastern, northern, northwestern, southeastern and southern part

of the microwatershed with moderate limitations of rooting depth and texture. Currently not suitable (Class N1) lands occur in an area of 2 ha (<1%) and are distributed in the southern part of the microwatershed with severe limitation of rooting depth.




Fig. 7.3 Land Suitability map of Bajra

7.4 Land Suitability for Groundnut (*Arachis hypogaea*)

Groundnut is one of the major oilseed crop grown in an area of 6.54 lakh ha in Karnataka in most of the districts either as rainfed or irrigated crop. The crop requirements for growing groundnut (Table 7.5) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing groundnut was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.4.

Highly suitable (Class S1) lands for growing groundnut occur in an area of 74 ha (12%) and are distributed in the western and northwestern part of the microwatershed. An area of about 10 ha (2%) is moderately suitable (Class S2) for growing groundnut and is distributed in the southern and southwestern part of the microwatershed with minor limitations of rooting depth and texture. An area of about 420 ha (69%) is marginally suitable (Class S3) for growing groundnut and is distributed in the major part of the microwatershed with moderate limitations of rooting depth, calcareousness and texture. Currently not suitable (Class N1) lands occur in an area of 2 ha (<1%) and are distributed in the southern part of the microwatershed with severe limitation of rooting depth.




Fig. 7.4 Land Suitability map of Groundnut

7.5 Land Suitability for Sunflower (*Helianthus annuus*)

Sunflower is one of the most important oilseed crop grown in an area of 4.1 lakh ha in the State in all the districts. The crop requirements for growing sunflower (Table 7.6) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing sunflower was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.5.

Highly suitable (Class S1) lands for growing sunflower occur in an area of 131 ha (22%) and are distributed in the northeastern, eastern, southeastern and northwestern part of the microwatershed. An area of about 90 ha (15%) is moderately suitable (Class S2) for growing sunflower and is distributed in the northern, southern, western, northwestern and southwestern part of the microwatershed with minor limitations of rooting depth, calcareousness and texture. An area of about 7 ha (1%) is marginally suitable (Class S3) and is distributed in the southern part of the microwatershed with moderate limitation of rooting depth. Currently not suitable (Class N1) lands occur in an area of 278 ha (46%) and are distributed in the southern, southeastern, central, northern and northwestern part of the microwatershed with severe limitation of rooting depth.




Fig. 7.5 Land Suitability map of Sunflower

7.6 Land Suitability for Red gram (*Cajanus Cajan*)

Redgram is one of the most important pulse crop grown in an area of 7.28 lakh ha in almost all the districts of the State. The crop requirements for growing red gram (Table 7.7) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing redgram was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.6.

An area of about 221 ha (36%) is moderately suitable (Class S2) for redgram and are distributed in the northeastern, western, southwestern, northwestern, southeastern and eastern part of the microwatershed. They have minor limitations of rooting depth, calcareousness and texture. An area of about 7 ha (1%) is marginally suitable (Class S3) and is distributed in the southern part of the microwatershed with moderate limitation of rooting depth. Currently not suitable (Class N1) lands occur in an area of 278 ha (46%) and are distributed in the southern, southeastern, central, northern and northwestern part of the microwatershed with severe limitation of rooting depth.




Fig. 7.6 Land Suitability map of Redgram

7.7 Land Suitability for Bengal gram (*Cicer aerativum*)

Bengal gram is one of the most important pulse crop grown in about 9.39 lakh ha area in Bijapur, Raichur, Kalaburgi, Dharwad, Belgaum and Bellary districts. The crop requirements for growing Bengal gram (Table 7.8) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing Bengal gram was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.7.

Highly suitable (Class S1) lands for growing bengalgram occur in an area of 131 ha (22%) and are distributed in the northeastern, eastern, southeastern and northwestern part of the microwatershed. An area of about 13 ha (2%) is moderately suitable (Class S2) for growing bengalgram and is distributed in the northwestern part of the microwatershed with minor limitations of rooting depth, calcareousness and texture. An area of about 84 ha (14%) is marginally suitable (Class S3) and is distributed in the southern, northern and western part of the microwatershed with moderate limitation of texture. Currently not suitable (Class N1) lands occur in an area of 278 ha (46%) and are distributed in the southern, southeastern, central, northern and northwestern part of the microwatershed with severe limitations of rooting depth and texture.




Fig. 7.7 Land Suitability map of Bengal gram

7.8 Land Suitability for Cotton (*Gossypium hirsutum*)

Cotton is one of the most important fibre crop grown in the State in about 8.75 lakh ha area in Raichur, Dharwad, Belgaum, Kalaburgi, Bijapur, Bidar, Bellary, Chitradurga and Chamarajnagar districts. The crop requirements for growing cotton (Table 7.9) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing cotton was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.8.

Highly suitable (Class S1) lands for growing cotton occur in an area of 131 ha (22%) and are distributed in the northeastern, eastern, southeastern and northwestern part of the microwatershed. An area of about 16 ha (3%) is moderately suitable (Class S2) for growing cotton and is distributed in the northwestern and southwestern part of the microwatershed with minor limitations of rooting depth and calcareousness. An area of about 80 ha (13%) is marginally suitable (Class S3) and is distributed in the southern, northern and western part of the microwatershed with moderate limitation of texture. Currently not suitable (Class N1) lands occur in an area of 278 ha (46%) and are distributed in the southern, southeastern, central, northern and northwestern part of the microwatershed with severe limitations of rooting depth and texture.




Fig. 7.8 Land Suitability map of Cotton

7.9 Land Suitability for Chilli (*Capsicum annuum*)

Chilli is one of the most important spice crop grown in about 0.42 lakh ha in Karnataka State. The crop requirements for growing chilli (Table 7.10) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing chilli was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.9.

Highly suitable (Class S1) lands for growing chilli occur in an area of 77 ha (13%) and are distributed in the western and northwestern part of the microwatershed. An area of about 151 ha (25%) is moderately suitable (Class S2) for growing chilli and is distributed in the northeastern, eastern, southern, northwestern and southeastern part of the microwatershed with minor limitations of rooting depth, calcareousness and texture. An area of about 276 ha (45%) is marginally suitable (Class S3) for growing chilli and is distributed in the central, eastern, northern, northwestern, southeastern and southern part of the microwatershed with moderate limitation of rooting depth. Currently not suitable (Class N1) lands occur in an area of 2 ha (<1%) and are distributed in the southern part of the microwatershed with severe limitation of rooting depth.




Fig 7.9 Land Suitability map of Chilli

7.10 Land Suitability for Tomato (*Lycopersicon esculentum*)

Tomato is one of the most important vegetable crop grown in about 0.61 lakh ha covering almost all the district of the state. The crop requirements for growing tomato (Table 7.11) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing tomato was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.10.

Highly suitable (Class S1) lands for growing tomato occur in an area of 77 ha (12%) and are distributed in the western and northwestern part of the microwatershed. An area of about 7 ha (1%) is moderately suitable (Class S2) for growing tomato and is distributed in the southern and southwestern part of the microwatershed with minor limitation of rooting depth. An area of about 420 ha (69%) is marginally suitable (Class S3) for growing tomato and is distributed in the major part of the microwatershed with moderate limitations of rooting depth and texture. Currently not suitable (Class N1) lands occur in an area of 2 ha (<1%) and are distributed in the southern part of the microwatershed with severe limitation of rooting depth.




Fig 7.10 Land Suitability map of Tomato

7.11 Land Suitability for Brinjal (*Solanum melongena*)

Brinjal is one of the most important vegetable crop grown in the state. The crop requirements for growing brinjal (Table 7.12) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing brinjal was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.11.

Highly suitable (Class S1) lands for growing brinjal occur in an area of 77 ha (12%) and are distributed in the western and northwestern part of the microwatershed. An area of about 7 ha (1%) is moderately suitable (Class S2) for growing brinjal and is distributed in the southern and southwestern part of the microwatershed with minor limitation of rooting depth. An area of about 420 ha (69%) is marginally suitable (Class S3) for growing brinjal and is distributed in the major part of the microwatershed with moderate limitations of rooting depth and texture. Currently not suitable (Class N1) lands occur in an area of 2 ha (<1%) and are distributed in the southern part of the microwatershed with severe limitation of rooting depth.




Fig 7.11 Land Suitability map of Brinjal

7.12 Land Suitability for Onion (*Allium cepa L.*)

Onion is one of the most important vegetable crop grown in the state. The crop requirements for growing onion (Table 7.13) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing onion was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.12.

Highly suitable (Class S1) lands for growing onion occur in an area of 77 ha (12%) and are distributed in the western and northwestern part of the microwatershed. An area of about 7 ha (1%) is moderately suitable (Class S2) for growing onion and is distributed in the southern and southwestern part of the microwatershed with minor limitation of rooting depth. An area of about 420 ha (69%) is marginally suitable (Class S3) for growing onion and is distributed in the major part of the microwatershed with moderate limitations of rooting depth and texture. Currently not suitable (Class N1) lands occur in an area of 2 ha (<1%) and are distributed in the southern part of the microwatershed with severe limitation of rooting depth.




Fig 7.12 Land Suitability map of Onion

7.13 Land Suitability for Bhendi (*Abelmoschus esculentus*)

Bhendi is one of the most important vegetable crop grown in the state. The crop requirements for growing bhendi (Table 7.14) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing bhendi was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.13.

Highly suitable (Class S1) lands for growing bhendi occur in an area of 77 ha (13%) and are distributed in the western and northwestern part of the microwatershed. An area of about 151 ha (25%) is moderately suitable (Class S2) for growing bhendi and is distributed in the northeastern, eastern, southern, northwestern and southeastern part of the microwatershed with minor limitations of rooting depth, calcareousness and texture. An area of about 276 ha (45%) is marginally suitable (Class S3) for growing bhendi and is distributed in the central, eastern, northern, northwestern, southeastern and southern part of the microwatershed with moderate limitation of rooting depth. Currently not suitable (Class N1) lands occur in an area of 2 ha (<1%) and are distributed in the southern part of the microwatershed with severe limitation of rooting depth.




Fig 7.13 Land Suitability map of Bhendi

7.14 Land Suitability for Drumstick (*Moringa oleifera*)

Drumstick is one of the most important vegetable crop grown in about 2403 ha in the state. The crop requirements for growing drumstick (Table 7.15) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing drumstick was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.14.

An area of about 208 ha (34%) is moderately suitable (Class S2) for growing drumstick and is distributed in the western, eastern, southeastern, southwestern, northern and northwestern part of the microwatershed with minor limitations of rooting depth and texture. An area of about 20 ha (3%) is marginally suitable (Class S3) and is distributed in the northwestern and southern part of the microwatershed with moderate limitations of rooting depth and calcareousness. Currently not suitable (Class N1) lands occur in an area of 278 ha (46%) and are distributed in the southern, southeastern, central, northern and northwestern part of the microwatershed with severe limitation of rooting depth.




Fig 7.14 Land Suitability map of Drumstick

7.15 Land Suitability for Mango (*Mangifera indica*)

Mango is one of the most important fruit crop grown in an area of 1.73 lakh ha in almost all the districts of the State. The crop requirements (Table 7.16) for growing mango were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing mango was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed is given in Figure 7.15.

An area of about 221 ha (36%) is marginally suitable (Class S3) and is distributed in the southwestern, western, eastern, northeastern, northern and northwestern part of the microwatershed with moderate limitations of texture and rooting depth. Currently not suitable (Class N1) lands occur in an area of 284 ha (47%) and are distributed in the northern, northwestern, southern, southeastern and central part of the microwatershed with severe limitation of rooting depth.




Fig. 7.15 Land Suitability map of Mango

7.16 Land Suitability for Guava (*Psidium guajava*)

Guava is one of the most important fruit crop grown in an area of 6558 ha in almost all the districts of the State. The crop requirements (Table 7.17) for growing guava were matched with the soil-site characteristics (7.1) and a land suitability map for growing guava was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.16.

An area of about 77 ha (13%) is moderately suitable (Class S2) for growing guava and is distributed in the western, southwestern and northwestern part of the microwatershed with minor limitation of rooting depth. An area of about 151 ha (25%) is marginally suitable (Class S3) and is distributed in the northeastern, eastern, southeastern, northwestern and southern part of the microwatershed with moderate limitations of rooting depth, texture and calcareousness. Currently not suitable (Class N1) lands occur in an area of 278 ha (46%) and are distributed in the southern, southeastern, central, northern and northwestern part of the microwatershed with severe limitation of rooting depth.




Fig. 7.16 Land Suitability map of Guava

7.17 Land Suitability for Sapota (*Manilkara zapota*)

Sapota is one of the most important fruit crop grown in an area of 29373 ha in almost all the districts of the State. The crop requirements (Table 7.18) for growing sapota were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing sapota was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.17.

An area of about 77 ha (13%) is moderately suitable (Class S2) for growing sapota and is distributed in the western, southwestern and northwestern part of the microwatershed with minor limitation of rooting depth. An area of about 151 ha (25%) is marginally suitable (Class S3) and is distributed in the northeastern, eastern, southeastern, northwestern and southern part of the microwatershed with moderate limitations of rooting depth and texture. Currently not suitable (Class N1) lands occur in an area of 278 ha (46%) and are distributed in the southern, southeastern, central, northern and northwestern part of the microwatershed with severe limitation of rooting depth.




Fig. 7.17 Land Suitability map of Sapota

7.18 Land Suitability for Pomegranate (*Punica granatum*)

Pomegranate is one of the most important fruit crop commercially grown in about 18488 ha in Karnataka, mainly in Bijapur, Bagalkot, Koppal, Gadag and Chitradurga districts. The crop requirements for growing pomegranate (Table 7.19) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing pomegranate was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.18.

An area of about 221 ha (36%) is moderately suitable (Class S2) for pomegranate and are distributed in the northeastern, western, southwestern, northwestern, southeastern and eastern part of the microwatershed. They have minor limitations of rooting depth, calcareousness and texture. An area of about 7 ha (1%) is marginally suitable (Class S3) and is distributed in the southern part of the microwatershed with moderate limitation of rooting depth. Currently not suitable (Class N1) lands occur in an area of 278 ha (46%) and are distributed in the southern, southeastern, central, northern and northwestern part of the microwatershed with severe limitation of rooting depth.




Fig 7.18 Land Suitability map of Pomegranate

7.19 Land Suitability for Musambi (*Citrus limetta*)

Musambi is one of the important fruit crop grown in an area of 3446 ha in almost all the districts of the State. The crop requirements for growing musambi (Table 7.20) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing musambi was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.19.

Highly suitable (Class S1) lands for growing musambi occur in an area of 131 ha (22%) and are distributed in the northeastern, eastern, southeastern and northwestern part of the microwatershed. An area of about 90 ha (15%) is moderately suitable (Class S2) for growing musambi and is distributed in the northern, southern, western, northwestern and southwestern part of the microwatershed with minor limitations of calcareousness and rooting depth. An area of about 7 ha (1%) is marginally suitable (Class S3) and is distributed in the southern part of the microwatershed with moderate limitation of rooting depth. Currently not suitable (Class N1) lands occur in an area of 278 ha (46%) and are distributed in the southern, southeastern, central, northern and northwestern part of the microwatershed with severe limitation of rooting depth.




Fig. 7.19 Land Suitability map of Musambi

7.20 Land Suitability for Lime (*Citrus sp*)

Lime is one of the most important fruit crop grown in an area of 0.11 lakh ha in almost all the districts of the State. The crop requirements for growing lime (Table 7.21) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing lime was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7. 20.

Highly suitable (Class S1) lands for growing lime occur in an area of 131 ha (22%) and are distributed in the northeastern, eastern, southeastern and northwestern part of the microwatershed. An area of about 90 ha (15%) is moderately suitable (Class S2) for growing lime and is distributed in the northern, southern, western, northwestern and southwestern part of the microwatershed with minor limitations of calcareousness and rooting depth. An area of about 7 ha (1%) is marginally suitable (Class S3) and is distributed in the southern part of the microwatershed with moderate limitation of rooting depth. Currently not suitable (Class N1) lands occur in an area of 278 ha (46%) and are distributed in the southern, southeastern, central, northern and northwestern part of the microwatershed with severe limitation of rooting depth.




Fig. 7.20 Land Suitability map of Lime

7.21 Land Suitability for Amla (*Phyllanthus emblica*)

Amla is one of the medicinal fruit crop grown in almost all the districts of the State. The crop requirements for growing amla (Table 7.22) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing amla was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.21.

Highly suitable (Class S1) lands for growing amla occur in an area of 77 ha (13%) and are distributed in the western and northwestern part of the microwatershed. An area of about 138 ha (23%) is moderately suitable (Class S2) for growing amla and is distributed in the northeastern, eastern, southern, northwestern and southeastern part of the microwatershed with minor limitations of rooting depth and texture. An area of about 289 ha (47%) is marginally suitable (Class S3) for growing amla and is distributed in the central, eastern, northern, northwestern, southeastern and southern part of the microwatershed with moderate limitations of rooting depth, calcareousness and texture. Currently not suitable (Class N1) lands occur in an area of 2 ha (<1%) and are distributed in the southern part of the microwatershed with severe limitation of rooting depth.




Fig. 7.21 Land Suitability map of Amla

7.22 Land Suitability for Cashew (*Anacardium occidentale*)

Cashew is one of the most important plantation nut crop grown in an area of 0.7 lakh ha in almost all the districts of the State. The crop requirements for growing cashew (Table 7.23) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing cashew was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.22.

An area of about 77 ha (13%) is marginally suitable (Class S3) and is distributed in the northwestern, western and southwestern part of the microwatershed with moderate limitation of nutrient availability. Currently not suitable (Class N1) lands occur in an area of 429 ha (71%) and are distributed in the major part of the microwatershed with severe limitations of rooting depth, texture and nutrient availability.




Fig. 7.22 Land Suitability map of Cashew

7. 23 Land Suitability for Jackfruit (*Artocarpus heterophyllus*)

Jackfruit is one of the most important fruit crop grown in an area of 5368 ha in almost all the districts of the State. The crop requirements for growing jackfruit (Table 7.24) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing jackfruit was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.23.

An area of about 77 ha (13%) is moderately suitable (Class S2) for growing jackfruit and is distributed in the western, southwestern and northwestern part of the microwatershed with minor limitation of rooting depth. An area of about 151 ha (25%) is marginally suitable (Class S3) and is distributed in the northeastern, eastern, southeastern, northwestern and southern part of the microwatershed with moderate limitations of rooting depth, calcareousness and texture. Currently not suitable (Class N1) lands occur in an area of 278 ha (46%) and are distributed in the southern, southeastern, central, northern and northwestern part of the microwatershed with severe limitation of rooting depth.




Fig. 7.23 Land Suitability map of Jackfruit

7.24 Land Suitability for Jamun (*Syzygium cumini*)

Jamun is an important fruit crop grown in almost all the districts of the State. The crop requirements for growing jamun (Table 25) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing jamun was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.24.

An area of about 131 ha (22%) is moderately suitable (Class S2) for growing jamun and is distributed in the eastern, southeastern, northern and northwestern part of the microwatershed with minor limitations of rooting depth and texture. An area of about 97 ha (16%) is marginally suitable (Class S3) and is distributed in the northwestern, western, southwestern and southern part of the microwatershed with moderate limitations of rooting depth and calcareousness. Currently not suitable (Class N1) lands occur in an area of 278 ha (46%) and are distributed in the southern, southeastern, central, northern and northwestern part of the microwatershed with severe limitation of rooting depth.




Fig. 7.24 Land Suitability map of Jamun

7.25 Land Suitability for Custard Apple (*Annona reticulata*)

Custard apple is one of the most important fruit crop grown in almost all the districts of the State. The crop requirements for growing custard apple (Table 7.26) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing custard apple was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.25.

Highly suitable lands (Class S1) for growing custard apple occupy an area of about 205 ha (34%) and are distributed in the northwestern, northeastern, southeastern, eastern, northern and western part of the microwatershed. An area of about 23 ha (4%) is moderately suitable (Class S2) for growing custard apple and are distributed in the northwestern, southern and southwestern part of the microwatershed. They have minor limitations of rooting depth and calcareousness. Marginally suitable lands (Class S3) for growing custard apple occupy an area of about 276 ha (45%) and are distributed in the central, northern, southern, northwestern, northeastern and southwestern part of the microwatershed with moderate limitations of texture and rooting depth. Currently not suitable (Class N1) lands occur in an area of 2 ha (<1%) and are distributed in the southern part of the microwatershed with severe limitation of rooting depth.




Fig. 7.25 Land Suitability map of Custard Apple

7.26 Land Suitability for Tamarind (*Tamarindus indica*)

Tamarind is one of the most important spice crop grown in almost all the districts of the state. The crop requirements for growing tamarind (Table 7.27) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing tamarind was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.26.

An area of about 131 ha (22%) is moderately suitable (Class S2) for growing tamarind and is distributed in the eastern, southeastern, northern and northwestern part of the microwatershed with minor limitations of rooting depth and texture. An area of about 90 ha (13%) is marginally suitable (Class S3) and is distributed in the northwestern, western, southwestern and southern part of the microwatershed with moderate limitations of rooting depth and calcareousness. Currently not suitable (Class N1) lands occur in an area of 284 ha (47%) and are distributed in the southern, southeastern, central, northern and northwestern part of the microwatershed with severe limitation of rooting depth.




Fig. 7.26 Land Suitability map of Tamarind

7.27 Land Suitability for Mulberry (*Morus nigra*)

Mulberry is one of the important leaf crop grown for rearing silk worms in about 1.6 lakh ha area in all the districts of the state. The crop requirements for growing mulberry (Table 7.28) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing mulberry was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.27.

An area of about 77 ha (13%) is moderately suitable (Class S2) for growing mulberry and is distributed in the western, southwestern and northwestern part of the microwatershed with minor limitation of rooting depth. An area of about 151 ha (25%) is marginally suitable (Class S3) and is distributed in the northeastern, eastern, southeastern, northwestern and southern part of the microwatershed with moderate limitations of rooting depth, calcareousness and texture. Currently not suitable (Class N1) lands occur in an area of 278 ha (46%) and are distributed in the southern, southeastern, central, northern and northwestern part of the microwatershed with severe limitation of rooting depth.




Fig 7.27 Land Suitability map of Mulberry

7.28 Land Suitability for Marigold (*Tagetes sps.*)

Marigold is one of the most important flower crop grown in an area of 9108 ha in almost all the districts of the State. The crop requirements (Table 7.29) for growing marigold were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing marigold was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.28.

Highly suitable (Class S1) lands for growing marigold occur in an area of 77 ha (13%) and are distributed in the western and northwestern part of the microwatershed. An area of about 151 ha (25%) is moderately suitable (Class S2) for growing marigold and is distributed in the northeastern, eastern, southern, northwestern and southeastern part of the microwatershed with minor limitations of rooting depth, calcareousness and texture. An area of about 276 ha (45%) is marginally suitable (Class S3) for growing marigold and is distributed in the central, eastern, northern, northwestern, southeastern and southern part of the microwatershed with moderate limitation of rooting depth. Currently not suitable (Class N1) lands occur in an area of 2 ha (<1%) and are distributed in the southern part of the microwatershed with severe limitation of rooting depth.




Fig. 7.28 Land Suitability map of Marigold

7.29 Land Suitability for Chrysanthemum (*Dendranthema grandiflora*)

Chrysanthemum is one of the most important flower crop grown in an area of 4978 ha in almost all the districts of the State. The crop requirements (Table 7.30) for growing chrysanthemum were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing chrysanthemum was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.29.

Highly suitable (Class S1) lands for growing chrysanthemum occur in an area of 77 ha (13%) and are distributed in the western and northwestern part of the microwatershed. An area of about 151 ha (25%) is moderately suitable (Class S2) for growing chrysanthemum and is distributed in the northeastern, eastern, southern, northwestern and southeastern part of the microwatershed with minor limitations of rooting depth, calcareousness and texture. An area of about 276 ha (45%) is marginally suitable (Class S3) for growing chrysanthemum and is distributed in the central, eastern, northern, northwestern, southeastern and southern part of the microwatershed with moderate limitation of rooting depth. Currently not suitable (Class N1) lands occur in an area of 2 ha (<1%) and are distributed in the southern part of the microwatershed with severe limitation of rooting depth.




Fig. 7.29 Land Suitability map of Chrysanthemum

Table 7.1 Soil-Site Characteristics of Chandaraki-1 Microwatershed

Soil Map Units	Climate (P) (mm)	Growing period (Days)	Drain-age Class	Soil depth (cm)	Soil texture		Gravelliness		AWC (mm/m)	Slope (%)	Erosion	pH	EC (dSm ⁻¹)	ESP (%)	CEC [Cmol (p ⁺)kg ⁻¹]	BS (%)
					Sur-face	Sub-surface	Surface (%)	Sub-surface (%)								
BDPiB3	866	150	WD	<25	sc	scl	<15	<15	<50	1-3	severe	8.58	0.262	0.35	18.10	100
KKRbB2g1	866	150	WD	<25	ls	sl	15-35	10-15	<50	1-3	moderate	-	5.82	-	9.77	0-22
BDLhB2	866	150	WD	25-50	scl	sl	<15	<15	<50	1-3	moderate	6.20	0.074	0.20	4.20	93
BDLhB2g1	866	150	WD	25-50	scl	sl	15-35	<15	<50	1-3	moderate	6.20	0.074	0.20	4.20	93
BDLiB2	866	150	WD	25-50	sc	sl	<15	<15	<50	1-3	moderate	6.20	0.074	0.20	4.20	93
HTKcB2	866	150	WD	25-50	sl	sl	<15	10-25	<50	1-3	moderate	6.81	0.062	0.38	3	101
JNKiB2g1	866	150	WD	50-75	sc	scl	15-35	<15	51-100	1-3	moderate	8.42	0.148	0.18	14.50	100
JNKiB3g1	866	150	WD	50-75	sc	scl	15-35	<15	51-100	1-3	severe	8.42	0.148	0.18	14.50	100
SHTmB2	866	150	WD	75-100	c	scl	15-35	<15	51-100	1-3	moderate	7.26	0.199	0.86	10.60	100
HSLiB2	866	150	MW	75-100	sc	sc	<15	<15	101-150	1-3	moderate	7.16	0.117	5.94	4.90	97
BGDmB2	866	150	MW	100-150	c	c	<15	<15	>200	1-3	moderate	7.85	0.253	0.26	65.90	100
BGDdB2g1	866	150	MW	100-150	ls	c	15-35	<15	>200	1-3	moderate	7.85	0.253	0.26	65.90	100
BMNmB2	866	150	MW	>150	c	c	<15	<15	>200	1-3	moderate	8.2	0.284	0.65	52.70	100

*Symbols and abbreviations are according to Field Guide for LRI under Sujala-III Project, Karnataka

Table 7.2 Land suitability criteria for Sorghum

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	26–30	30–34; 24–26	34–40; 20–24	>40; <20
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, c (red), c (black)	scl, cl	ls, sl	-
	pH	1:2.5	5.5-7.8	5.0-5.5 7.8-9.0	>9.0	-
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	10-15
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

Table 7.3 Land suitability criteria for Maize

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	30-34	35-38 26-30	38-40 26-20	
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc	c (red), c (black)	ls, sl	-
	pH	1:2.5	5.5-7.8	5.0-5.5 7.8-9.0	>9.0	-
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	-
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

Table 7.4 Land suitability criteria for Bajra

Land use requirement		Rating				
Soil –site characteristics	Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)	
Climatic regime	Mean temperature in growing season	°C	28-32	33-38 24-27	39-40 20-23	<20
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm	500-750	400-500	200-400	<200
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl, scl, cl,sc,c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.8	5.0-5.5 7.8-9.0	5.5-6.0 >9.0	
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	15-35	35-60	>60	
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	1-3	3-5	5-10	>10

Table 7.5 Land suitability criteria for Groundnut

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	24–33	22–24; 33–35	20–22; 35–40	<20; >40
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. Well drained	Poorly drained	Very Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl	sl,cl, sc	c (red), c (black), ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<35	35-60	>60	
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.6 Land suitability criteria for Sunflower

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	24–30	30–34; 20–24	34–38; 16–20	>38; <16
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	mod. Well drained	-	Poorly to very drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	cl, sc,c (red), c (black)	scl	ls, sl	-
	pH	1:2.5	6.5-7.8	7.8-8.4 5.5-6.5	8.4-9.0; 5.0-5.5	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.7 Land suitability criteria for Redgram

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	30-35(G) 20-25(AV) 15-18 (F&PS) 35-40(M)	25-30(G) 20-25 (AV) 12-15 (F&PS) 30-35(M)	20-25(G) 15-20(AV) 10-12 (F&PS) 25-30(M)	< 20 <15 <10 <25
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. Well drained	Poorly drained	Very Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, c (red)	c (black),sl, scl, cl	ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-9.0	5.0-5.5 >9.0	-
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-50	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<1.0	1.0-2.0	>2.0	
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.8 Land suitability criteria for Bengal gram

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	20–25	25–30; 15–20	30–35; 10–15	>35; <10
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. Well drained	Poorly drained	Very Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	c(black)	-	c (red), scl, cl, sc	ls, sl
	pH	1:2.5	6.0-7.8	5.0-6.0 7.8-9.0	>9.0	-
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	-
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.9 Land suitability criteria for Cotton

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	22-32	>32	<19	-
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well to moderately well	Poorly drained/Some what excessively drained	-	very poorly/excessively drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, c (red,black)	cl	scl	ls, sl
	pH	1:2.5	6.5-7.8	7.8-8.4	5.5-6.5 8.4->9.0	<5.5
	CEC	C mol (p+)Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	50-100	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	<3	3-5	-	>5

Table 7.10 Land suitability criteria for Chilli

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	25-32	33-35 20-25	35-38 <20	>38
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc	c (black), sl	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.11 Land suitability criteria for Tomato

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	25-28	29-32 20-24	15-19 33-36	<15 >36
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl, scl, cl, sc, c (red)	-	ls, c(black)	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.12 Land suitability criteria for Brinjal

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	Well drained	Moderately well drained	Poorly drained	V. Poorly drained
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class				
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl, scl, cl, sc c (red)	-	ls, c (black)	-
	pH	1:2.5	6.0-7.3	7.3-8.4 5.0-6.0	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	>60
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.13 Land suitability criteria for Onion

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	20-30	30-35	35-40	>40
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately /imperfectly	-	Poorly to V poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl,scl,cl,sc,c (red)	-	c (Black),ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<1.0	1.0-2.0	2.0-4.0	<4
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.14 Land suitability criteria for Bhendi

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	25-28	29-32 20-24	15-19 33-36	<15 >36
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Imperfectly drained	Poorly to very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl,sc, c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.15 Land suitability criteria for Drumstick

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, scl, cl, c (red)	sl, c (black)	ls	s
	pH	1:2.5	6.0-7.3	5.0-5.5 7.3-7.8	5.5-6.0 7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<35	35-60	60-80	>80
Soil toxicity	Salinity (EC saturation extract)	ds/m				
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-10	-	>10

Table 7.16 Land suitability criteria for Mango

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-32	24-27 33-35	36-40	20-24
	Min temp. before flowering	°C	10-15	15-22	>22	-
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration	Days				
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V. Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	-	ls, sl, c (black)	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>150	100-150	75-100	<75
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.17 Land suitability criteria for Guava

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-32	33-36 24-27	37-42 20-23	
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	sl	c (black), ls	-
	pH	1:2.5	6.0-7.8	5.0-6.0	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.18 Land suitability criteria for Sapota

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-32	33-36 24-27	37-42 20-23	>42 <18
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	-	Poorly to very drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	sl	ls, c (black)	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.19 Land suitability criteria for Pomegranate

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	30-34	35-38 25-29	39-40 15-24	
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl,cl, sc, c (red)	c (black),sl	ls	-
	pH	1:2.5	5.5-7.8	7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.20 Land suitability criteria for Musambi

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-30	31-35 24-27	36-40 20-23	>40 <20
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately drained	poorly	Very poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c	sl	ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.21 Land suitability criteria for Lime

Land use requirement		Rating				
Soil –site characteristics	Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)	
Climatic regime	Mean temperature in growing season	°C	28-30	31-35 24-27	36-40 20-23	>40 <20
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately drained	poorly	Very poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c	sl	ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.22 Land suitability criteria for Amla

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. well drained	Poorly drained	V. Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	c (black)	ls, sl	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15-35	35-60	60-80	-
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

Table 7.23 Land suitability criteria for Cashew

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	32 to 34	28 to 32; 34 to 38	24 to 28; 38 to 40	<20; >40
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	-	sl, ls	c (black)
	pH	1:2.5	5.5-6.5	5.0-5.5 6.5-7.3	7.3-7.8	>7.8
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-10	>10	-

Table 7.24 Land suitability criteria for Jackfruit

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. well	Poorly	V. Poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	-	sl, ls, c (black)	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	>60
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10-

Table 7.25 Land suitability criteria for Jamun

Land use requirement		Rating				
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well	Mod. well	Poorly	V.Poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c(red)	sl, c (black)	ls	-
	pH	1:2.5	6.0-7.8	5.0-6.0	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>150	100-150	50-100	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	>60
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

Table 7.26 Land suitability criteria for Custard apple

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	Scl, cl, sc, c (red), c (black)	-	Sl, ls	-
	pH	1:2.5	6.0-7.3	5.5-6.0 7.3-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15-35	35-60	60-80	-
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	>5	-

Table 7.27 Land suitability criteria for Tamarind

Land use requirement		Rating				
Soil –site characteristics	Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)	
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod.well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl,sc, c (red)	sl, c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>150	100-150	75-100	<75
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

Table 7.28 Land suitability criteria for Mulberry

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	24–28	22–24; 28–32	32–38; 22–18	>38; <18
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V. Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, cl, scl	c (red)	c (black), sl, ls	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.8-8.4	7.3-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	0-35	35-60	60-80	>80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

Table 7.29 Land suitability criteria for Marigold

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	18-23	17-15 24-35	35-40 10-14	>40 <10
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl,scl, cl, sc, c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%				
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.30 Land suitability criteria for Chrysanthemum

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	18-23	17-15 24-35	35-40 10-14	>40 <10
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl,scl, cl, sc, c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%				
Erosion hazard	Slope	%	<3	3-5	5-10	>10

7.30 Land Management Units (LMUs)

The 13 soil map units identified in Chandaraki-1 microwatershed have been grouped into 4 Land Management Units (LMU's) for the purpose of preparing a Proposed Crop Plan. Land Management Units are grouped based on the similarities in respect of the type of soil, the depth of the soil, the surface soil texture, gravel content, AWC, slope, erosion etc. and a Land Management Units map (Fig. 7.30) has been generated. These Land Management Units are expected to behave similarly for a given level of management.

The 13 map units that have been grouped into 4 Land Management Units along with brief description of soil and site characteristics are given below.

LMU	Soil map units	Soil and site characteristics
1	115.BGDmB2 151.BGDmB2g1 62.BMNmB2 33.HSLiB2 112.SHTmB2	Moderately deep to very deep (75 to >150), black calcareous to non calcareous clay soils, 1-3% slopes, non gravelly to gravelly (<15-35%), moderate erosion.
2	23.JNKiB2g1 24.JNKiB3g1	Moderately shallow (50 to 75 cm), sandy clay loam soils, 1-3% slopes, gravelly (15-35%), moderate to severe erosion.
3	4.BDLhB2 162.BDLhB2g1 5.BDLiB2 165.HTKcB2	Shallow (25 to 50 cm), sandy clay loam to sandy loam soils, 1-3% slopes, non gravelly to gravelly (<15-35%), moderate erosion.
4	119.BDPiB3 153.KKRbB2g1	Very shallow to shallow soils (<25 cm), 1-3% slopes, non gravelly to gravelly (<15-35%), moderate to severe erosion.




Fig. 7.30 Land Management Units Map- Chandaraki-1 Microwatershed

7.31 Proposed Crop Plan for Chandaraki-1 Microwatershed

After assessing the land suitability for the 29 crops, the Proposed Crop Plan has been prepared for the 4 identified LMUs by considering only the highly (Class S1) and moderately (Class S2) suitable lands for each of the 29 crops. The resultant proposed crop plan is presented below in Table 7.31.

Table 7.31 Proposed Crop Plan for Chandaraki-1 Microwatershed

LMU	Soil Map Units	Survey Number	Field Crops/ Commercial crops	Horticulture Crops (Rainfed/Irrigated)	Suitable Interventions
1	115.BGDmB2 151.BGDmB2g1 62.BMNB2 33.HSLiB2 112.SHTmB2 (Moderately deep to very deep, black calcareous to non calcareous clay soils)	Chandraki: 1,3,5,6,7,8,9,10,11,12,13,14,71,83,84,92,93,94,95,96,97,99,101,118,119,120,121,122,123,124,126,128,135,155,158,160,161,162,164,165,166,167,168,169,170/2,170/3,171,173/1,173/2,175,178,179,180,181,182,183,184,185,186,187,189,190,191,192,193,194,197,198,199,200,201,202,203,204,205,206,207,208,209,210,211,212,213,217,230,232,233,234,235,236,237,238,239,240,498,499/1,499/2,500,501,502,521,542,543,545,546,547,550,551/1,551/2,551/3, 552,553,554 Madhepalli : 72,73,75,76,77,82	Maize, Sorghum, Sunflower, Cotton, Redgram, Bengalgram, Bajra	Fruit crops: Lime, Musambi, Custard apple, Pomegranate Vegetables: Chilli, Bhendi Flowers: Marigold, Chrysanthemum	Application of FYM, Biofertilizers and micronutrients, drip irrigation, mulching, suitable soil and water conservation practices
2	23.JNKiB2g1 24.JNKiB3g1 (Moderately shallow, sandy clay loam soils)	Madhepalli : 12,13	Maize, Sorghum Groundnut, Bajra	Fruit crops: Amla, Custard apple Vegetables: Tomato, Chilli, Brinjal, Bhendi, Onion Flowers: Marigold, Chrysanthemum	Application of FYM, Biofertilizers and micronutrients, drip irrigation, mulching, suitable soil and water conservation practices
3	4.BDLhB2 162.BDLhB2g1 5.BDLiB2 165.HTKcB2 (Shallow, sandy clay loam to sandy loam soils)	Chandraki: 2,4,15,16,72/2,72/3,88/1,88/2,89,90,91,98,100,102,103,105,106,107,108,109,110,112,113,114,115,116,117/1,117/10,117/11,117/12,117/13,117/14,117/15,117/2,117/3,117/4,117/5,117/6,117/7,117/8,117/9,125,127,129,130,131,132,141,142,143/1,143/2,144,146,151/1,151/2,195,503,541,548,549,555 Madhepalli: 1,10,2,23,3,4,5,6,7,74,78,79,8,80, 81,83,84,85	-	Agri-Silvi-Pasture: Hybrid Napier, <i>Styloxanthes hamata</i> , <i>Styloxanthes scabra</i>	Use of short duration varieties, sowing across the slope, drip irrigation is recommended
4	119.BDPiB3 153.KKRbB2g1 (Very shallow, sandy clay loam to sandy loam soils)	Madhepalli: 9	-	Hybrid Napier, <i>Styloxanthes hamata</i> , <i>Styloxanthes scabra</i>	Use of short duration varieties, sowing across the slope, drip irrigation is recommended

SOIL HEALTH MANAGEMENT

8.1 Soil Health

Soil health is basic to plant health and plant health is basic to human and bovine health. Soil is fundamental to crop production. Without soil, no food could be produced nor would livestock be fed on a large scale. Because it is finite and fragile, soil is a precious resource that requires special care from its users.

Soil health or the capacity of the soil to function is critical to human survival. Soil health has been defined as: “the capacity of the soil to function as a living system without adverse effect on the ecosystem”. Healthy soils maintain a diverse community of soil organisms that help to form beneficial symbiotic associations with plant roots, recycle essential plant nutrients, improve soil structure with positive repercussions for soil, water and nutrient holding capacity and ultimately improve crop production and also contribute to mitigating climate change by maintaining or increasing its carbon content.

Functional interactions of soil biota with organic and inorganic components, air and water determine a soil’s potential to store and release nutrients and water to plants and to promote and sustain plant growth. Thus, maintaining soil health is vital to crop production and conserve soil resource base for sustaining agriculture.

The most important characteristics of a healthy soil are

- Good soil tilth
- Sufficient soil depth
- Good water storage and good drainage
- Adequate supply, but not excess of nutrients
- Large population of beneficial organisms
- Small proportion of plant pathogens and insect pests
- Low weed pressure
- Free of chemicals and toxins that may harm the crop
- Resistance to degradation
- Resilience when unfavorable conditions occur

Characteristics of Chandaraki-1 Microwatershed

- ❖ The soil phases identified in the microwatershed belonged to the soil series of MDG series occupies a maximum area of 130 ha (17%) followed by BDP 99 ha (13%), BDL 93 ha (12%), VNK 83 ha (11%), ANR 77 ha (10%), SHT 73 ha (9%), YLR 47 ha (6%), YDR 32 ha (4%), MDR 30 (4%), HTK 7 ha (<1%), DSB 3 ha (<1%), VKS 3 ha (<1%), SBR 2 ha (<1), JNK 0.47 ha (<1%) and BMN 0.18 ha (<1).
- ❖ As per land capability classification entire area of the microwatershed falls under arable land category (Class II, III & IV). The major limitations identified in the arable lands were soil and erosion.

- ❖ On the basis of soil reaction, an area of about 29 per cent is slightly acid (pH 6.0-6.5). An area of about 4 per cent is moderately acid (pH 5.5-6.0). An area of about 31 per cent is neutral (pH 6.5-7.3). An area of about 10 per cent is slightly alkaline (pH 7.3-7.8). An area of about 6 per cent is moderately alkaline (pH 7.8-8.4). An area of 4 per cent is strongly alkaline (pH 8.4-9.0) in the microwatershed.

Soil Health Management

The following actions are required to improve the current land husbandry practices that provide a sound basis for the successful adoption of sustainable crop production system.

Acid soils

Acid soils occur in 200 ha area in the microwatershed.

1. Growing of crops suitable for a particular soil pH.
2. Amelioration of soils through the application of amendments (liming materials).

Liming materials:

1. CaCO_3 (Calcium Carbonate).
2. Dolomite [$\text{Ca Mg} (\text{CO}_3)_2$]
3. Quick lime (Cao)
4. Slaked lime [$\text{Ca} (\text{OH})_2$]

For normal pH and pH 4.8 (35 t/ha) and pH 6.0-7.0 (4 t/ha) lime is required.

Alkaline soils

Alkaline soils occur in 119 ha area in the microwatershed.

1. Regular addition of organic manure, green manuring, green leaf manuring, crop residue incorporation and mulching needs to be taken up to improve the soil organic matter status.
2. Application of biofertilizers (Azospirillum, Azotobacter, Rhizobium).
3. Application of 25% extra N and P (125 % RDN&P).
4. Application of ZnSO_4 – 12.5 kg/ha (once in three years).
5. Application of Boron – 5kg/ha (once in three years).

Neutral soils

Neutral soils occur in 186 ha area of the microwatershed.

1. Regular addition of organic manure, green manuring, green leaf manuring, crop residue incorporation and mulching needs to be taken up to improve the soil organic matter status.
2. Application of biofertilizers, (Azospirillum, Azotobacter, Rhizobium).
3. Application of 100 per cent RDF.
4. Need based micronutrient applications.

Besides the above recommendations, the best transfer of technology options are also to be adopted.

Soil Degradation

Soil erosion is one of the major factor affecting the soil health in the microwatershed. Entire microwatershed area is suffering from moderate and severe erosion. In areas of moderate and severe erosion immediate soil and water conservation and, other land development and land husbandry practices are required for restoring soil health.

Dissemination of Information and Communication of Benefits

Any large scale implementation of soil health management requires that supporting information is made available widely, particularly through channels familiar to farmers and extension workers. Given the very high priority attached to soil-health especially by the Central Government on issuing Soil-Health Cards to all the farmers, media outlets like Regional, State and National Newspapers, Radio and Dooradarshan programs in local languages but also modern information and communication technologies such as Cellular phones and the Internet, which can be much more effective in reaching the younger farmers.

Inputs for Net Planning (Saturation Plan) and Interventions needed

Net planning (Saturation Plan) in IWMP is focusing on preparation of

1. Soil and Water Conservation Plan for each plot or farm.
2. Productivity enhancement measures/ interventions for existing crops/livestock/other farm enterprises.
3. Diversification of farming mainly with perennial horticultural crops and livestock.
4. Improving livelihood opportunities and income generating activities.

In this connection, how various outputs of Sujala-III are of use in addressing these objectives of Net Planning (Saturation Plan) are briefly presented below.

- ❖ **Soil Depth:** The depth of a soil decides the amount of moisture and nutrients it can hold, what crops can be taken up or not, depending on the rooting depth and the length of growing period available for raising any crop. Deeper the soil, better for a wide variety of crops. If sufficient depth is not available for growing deep rooted crops, either choose medium or short duration crops or deeper planting pits need to be opened and additional good quality soil brought from outside has to be filled into the planting pits.
- ❖ **Surface Soil Texture:** Lighter soil texture in the top soil means, better rain water infiltration, less run-off and soil moisture conservation, less capillary rise and less evaporation losses. Lighter surface textured soils are amenable to good soil tilth and are highly suitable for crops like groundnut, root vegetables (carrot, radish, potato etc) but not ideal for crops that need stagnant water like lowland paddy. Heavy

textured soils are poor in water infiltration and percolation. They are prone for sheet erosion; such soils can be improved by sand mulching. The technology that is developed by the AICRP-Dryland Agriculture, Vijayapura, Karnataka can be adopted.


- ❖ **Gravelliness:** More gravel content is favorable for run-off harvesting but poor in soil moisture storage and nutrient availability. It is a significant parameter that decides the kind of crop to be raised.
- ❖ **Land Capability Classification:** The land capability map shows the areas suitable and not suitable for agriculture and the major constraints in each of the plot/survey number. Hence, one can decide what kind of enterprise is possible in each of these units. In general, erosion and soil are the major constraints in Chandaraki-1 microwatershed.
- ❖ **Organic Carbon:** The OC content (an index of available Nitrogen) is high (>0.75%) in an entire area of the microwatershed.
- ❖ **Promoting green manuring:** Growing of green manuring crops costs Rs. 1250/ha (green manuring seeds) and about Rs. 2000/ha towards cultivation that totals to Rs. 3250/- per ha. On the other hand, application of organic manure @ 10 tons/ha costs Rs. 5000/ha. The practice needs to be continued for 2-3 years or more.
- ❖ **Available Phosphorus:** Available Phosphorus is medium (23-57 kg/ha) in an area of 214 ha (35%) in the microwatershed. Low (<23 kg/ha) in an area of 229 ha (38%) and high (>57 kg/ha) in an area of 63 ha (10%). In medium and low areas, for all the crops 25% additional P needs to be applied.
- ❖ **Available Potassium:** Available potassium is medium (145-337 kg/ha) in an area of 273 ha (45%) and high (>337 kg/ha) in an area of 232 ha (38%) in the microwatershed. All the plots, where available potassium is medium, for all the crops, additional 25% potassium may be applied.
- ❖ **Available Sulphur:** Available sulphur is a very critical nutrient for oilseed crops. It is low in an entire area of the microwatershed. Low areas need to be applied with magnesium sulphate or gypsum or Factamphos (p) fertilizer (13% sulphur) for 2-3 years for the deficiency to be corrected.
- ❖ **Available Boron:** An area of 109 ha (18%) is high (>0.1 ppm) in available boron and medium (0.5-1.0 ppm) in an area of 396 ha (65%). Application of sodium tetra borate @ 10 kg/ha as soil application or 0.2 % borax as foliar spray is recommended for medium areas.
- ❖ **Available Iron:** Entire area is sufficient (>4.5 ppm) in available iron content in the microwatershed.
- ❖ **Available Manganese:** All the soils in the microwatershed are sufficient (>1.0 ppm) in available manganese.
- ❖ **Available Copper:** All the soils in the microwatershed are sufficient (>0.2 ppm) in available copper.

- ❖ **Available Zinc:** An area of 137 ha (23%) is deficient (<0.6 ppm) in available zinc content of the microwatershed and 368 ha (61%) area is sufficient (>0.6 ppm). Application of zinc sulphate @25 kg/ha is recommended for zinc deficient areas.
- ❖ **Soil Alkalinity:** Alkaline soils occur 119 ha area in the microwatershed. Alkaline soils need application of gypsum and wherever calcium is in excess, iron pyrites and element sulphur can be recommended. Management practices like treating repeatedly with good quality water to drain out the excess salts and provision of subsurface drainage and growing of salt tolerant crops like Casuarina, Acacia, Neem, Ber etc, are recommended.
- ❖ **Land Suitability for various crops:** Areas that are highly, moderately and marginally suitable and not suitable for growing various crops are indicated. Along with the suitability, various constraints that are limiting the productivity are also indicated. For example, in case of cotton, gravel content, rooting depth and salinity/alkalinity are the major constraints in various plots. With suitable management interventions, the productivity can be enhanced. In order to increase the water holding capacity of light textured soils, growing of green manure crops and application of organic manure is recommended.

SOIL AND WATER CONSERVATION TREATMENT PLAN

For preparing soil and water conservation treatment plan for Chandaraki-1 microwatershed, the land resource inventory database generated under Sujala-III project has been transformed as information through series of interpretative (thematic) maps using soil phase map as a base. The various thematic maps (1:7920 scale) generated were

- Soil depth
- Surface soil texture
- Available water capacity
- Soil slope
- Soil gravelliness
- Land capability
- Present land use and land cover
- Crop suitability
- Rainfall
- Hydrology
- Water Resources
- Socio-economic data
- Contour plan with existing features- network of waterways, pothissa boundaries, cut up/ minor terraces etc.
- Cadastral map (1:7920 scale)
- Satellite imagery (1:7920 scale)



Apart from these, Hand Level/ Hydro Marker/ Dumpy Level/ Total Station and Kathedars' List to be collected.

Steps for Survey and Preparation of Treatment Plan

The boundaries of Land User Groups' and Survey No. boundaries are traced in the field.

- Naming of user groups and farmers
- Identification of arable and non arable lands
- Identification of drainage lines and gullies
- Identification of non treatable areas
- Identification of priority areas in the arable lands
- Treatment plan for arable lands
- Location of water harvesting and recharge structures

9.1 Treatment Plan

The treatment plan recommended for arable lands is briefly described below


9.1.1 Arable Land Treatment

A. BUNDING

Steps for Survey and Preparation of Treatment Plan		USER GROUP-1 CLASSIFICATION OF GULLIES
<ul style="list-style-type: none"> • Cadastral map (1:7920 scale) is enlarged to a scale of 1:2500 scale • Existing network of waterways, pothissa boundaries, grass belts, natural drainage lines/ watercourse, cut ups/ terraces are marked on the cadastral map to the scale • Drainage lines are demarcated into 		
Small gullies	(up to 5 ha catchment)	
Medium gullies	(5-15 ha catchment)	
Ravines	(15-25 ha catchment) and	
<i>Halla/Nala</i>	(more than 25ha catchment)	

Measurement of Land Slope

Land slope is estimated or determined by the study and interpretation of contours or by measurement in the field using simple instruments like Hand Level or Hydromarker.



Vertical and Horizontal intervals between bunds as recommended by the Watershed Development Department.

Slope percentage	Vertical interval (m)	Corresponding Horizontal Distance (m)
2 - 3%	0.6	24
3 - 4%	0.9	21
4 - 5%	0.9	21
5 - 6%	1.2	21
6 - 7%	1.2	21

Note: (i) The above intervals are maximum.

(ii) Considering the slope class and erosion status (A1... A=0-1 % slope, 1= slight erosion) the intervals have to be decided.

Bund length recording: Considering the contour plan and the existing grass belts/partitions, the bunds are aligned and lengths are measured.

Section of the Bund

Bund section is decided considering the soil texture class and gravelliness class (bg₀... b=loamy sand, g₀ = <15% gravel). The recommended Sections for different soils are given below.

Recommended Bund Section

Top width (m)	Base width (m)	Height (m)	Side slope (Z:1;H:V)	Cross section (sq m)	Soil Texture	Remarks
0.3	0.9	0.3	01:01	0.18	Sandy loam	Vegetative bund
0.3	1.2	0.3	1.5:1	0.225	Sandy clay	
0.3	1.2	0.5	0.9:1	0.375	Red gravelly soils	
0.3	1.2	0.6	0.75:1	0.45		
0.3	1.5	0.6	01:01	0.54	Red sandy loam	
0.3	2.1	0.6	1.5:1	0.72	Very shallow black clayey soils	
0.45	2	0.75	01:01	0.92		
0.45	2.4	0.75	1.3:1	1.07	Shallow black clayey soils	
0.6	3.1	0.7	1.78:1	1.29	Medium black clayey soils	
0.5	3	0.85	1.47:1	1.49		

Formation of Trench cum Bund

Dimensions of the Borrow Pits/Trenches to be excavated (machinery are decided considering the Bund Section).

Details of Borrow Pit dimensions are given below:

TRENCH CUM BUND

'A' FRAME FOR INTERBUND MANAGEMENT

Size of Borrow Pits/ Trench recommended for Trench cum Bund (by machinery)

Bund section	Bund length	Earth quantity	Pit				Berm (pit to pit)	Soil depth class
			L(m)	W(m)	D(m)	Quantity (m ³)		
m ²	m	m ³					m	
0.375	6	2.25	5.85	0.85	0.45	2.24	0.15	Shallow
0.45	6	2.7	5.4	1.2	0.43	2.79	0.6	Shallow
0.45	6	2.7	5	0.85	0.65	2.76	1	Moderately Shallow
0.54	5.6	3.02	5.5	0.85	0.7	3.27	0.1	Moderately shallow
0.54	5.5	2.97	5	1.2	0.5	3	0.5	Shallow
0.72	6.2	4.46	6	1.2	0.7	5.04	0.2	Moderately shallow
0.72	5.2	3.74	5.1	0.85	0.9	3.9	0.1	Moderately deep

B. Water Ways

- Existing waterways are marked on the cadastral map (1:7920 scale) and their dimensions are recorded.
- Considering the contour plan of the MWS, additional waterways/ modernization of the existing ones can be thought of.
- The design details are given in the Manual.

C. Farm Ponds

Waterways and the catchment area will give an indication on the size of the Farm Pond. Location of the pond can be decided based on the contour plan/ field condition and farmers' need/desire.

D. Diversion Channel

Existing EPT/ CPT are marked on the cadastral map. Looking to the need, these can be modernized or fresh diversion channel can be proposed and runoff from this can be stored in *Gokatte*/ Recharge ponds.

9.1.2 Non-Arable Land Treatment

Depending on the gravelliness and crops preferred by the farmers, the concerned authorities can decide appropriate treatment plan. The recommended treatments may be Contour Trench, Staggered Trench, Crescent Bund, Boulder Bund or Pebble Bund.

9.1.3 Treatment of Natural Water Course/ Drainage Lines

- a) The cadastral map has to be updated as regards the network of drainage lines (gullies/*nalas/hallas*) and existing structures are marked to the scale and storage capacity of the existing water bodies are documented.
- b) The drainage line will be demarcated into Upper Reach, Middle Reach and Lower Reach.
- c) Considering the Catchment, *Nala* bed and bank conditions, suitable structures are decided.
- d) Number of storage structures (Check dam/*Nala* bund/ Percolation tank) will be decided considering the commitments and available runoff from water budgeting and quality of water in the wells and site suitability.
- e) Detailed Leveling Survey using Dumpy Level / Total Station has to be carried out to arrive at the site-specific designs as shown in the Manual.
- f) The location of ground water recharge structures are decided by examining the lineaments and fracture zones from geological maps.
- g) Rainfall intensity data of the nearest Rain Gauge Station is considered for Hydrologic Designs.
- h) Silt load to the Storage/Recharge Structures is reduced by providing vegetative, boulder and earthen checks in the natural water course. Location and design details are given in the Manual.

9.2 Recommended Soil and Water Conservation Measures

The appropriate conservation structures best suited for each of the land parcel/ survey number (Appendix-I) are selected based on the slope per cent, severity of erosion, amount of rainfall, land use and soil type. The different kinds of conservation structures recommended are:

1. Graded / Strengthening of Bunds
2. Trench cum Bunds (TCB)
3. Trench cum Bunds / Strengthening
4. Crescent Bunds

A map (Fig. 9.1) showing soil and water conservation plan with different kinds of structures recommended has been prepared which shows the spatial distribution and extent of area. An area of about 503 ha (83%) needs Graded Bunding and an area of 2 ha (<1%) needs Trench cum Bunding in the microwatershed.

The conservation plan prepared may be presented to all the stakeholders including farmers and after considering their suggestions, the conservation plan for the microwatershed may be finalised in a participatory approach.




Fig. 9.1 Soil and Water Conservation Plan map of Chandaraki-1 Microwatershed

9.3 Greening of Microwatershed

As part of the greening programme in the watersheds, it is envisaged to plant a variety of horticultural and other tree plants that are edible, economical and produce lot of biomass which helps to restore the ecological balance in the watersheds. The lands that are suitable for greening programme are non-arable lands (land capability classes V, VI VII and VIII) and also the lands that are not suitable or marginally suitable and field bunds for growing annual and perennial crops. The method of planting these trees is given below.

It is recommended to open pits during the 1st week of March along the contour and heap the dugout soil on the lower side of the slope in order to harness the flowing water and facilitate weathering of soil in the pit. Exposure of soil in the pit also prevents spread of pests and diseases due to scorching sun rays. The pits should be filled with mixture of soil and organic manure during the second week of April and keep ready with sufficiently tall seedlings produced either in poly bags or in root trainer nurseries so that planting can be done during the 2nd or 3rd week of April depending on the rainfall.

The tree species suitable for the area considering rainfall, temperature and adaptability is listed below; waterlogged areas are recommended to be planted with species like Nerale (*Syzgium cumini*) and Bamboo. Dry areas are to be planted with species like Honge, Bevu, Seetaphal etc.

Dry Deciduous Species			Temp (°C)	Rainfall (mm)
1.	Bevu	<i>Azadiracta indica</i>	21–32	400 –1,200
2.	Tapasi	<i>Holoptelia integrifolia</i>	20-30	500 - 1000
3.	Seetaphal	<i>Anona Squamosa</i>	20-40	400 - 1000
4.	Honge	<i>Pongamia pinnata</i>	20 -50	500– 2,500
5.	Kamara	<i>Hardwickia binata</i>	25 -35	400 - 1000
6.	Bage	<i>Albezzia lebbek</i>	20 - 45	500 - 1000
7.	Ficus	<i>Ficus bengalensis</i>	20 - 50	500–2,500
8.	Sisso	<i>Dalbargia Sissoo</i>	20 - 50	500 -2000
9.	Ailanthus	<i>Ailanthus excelsa</i>	20 - 50	500 - 1000
10.	Hale	<i>Wrightia tinctoria</i>	25 - 45	500 - 1000
11.	Uded	<i>Steriospermum chelanoides</i>	25 - 45	500 -2000
12.	Dhupa	<i>Boswella Serrata</i>	20 - 40	500 - 2000
13.	Nelli	<i>Emblca Officinalis</i>	20 - 50	500 -1500
14.	Honne	<i>Pterocarpus marsupium</i>	20 - 40	500 - 2000
Moist Deciduous Species			Temp (°C)	Rainfall (mm)
15.	Teak	<i>Tectona grandis</i>	20 - 50	500-5000
16.	Nandi	<i>Legarstroemia lanceolata</i>	20 - 40	500 - 4000
17.	Honne	<i>Pterocarpus marsupium</i>	20 - 40	500 - 3000
18.	Mathi	<i>Terminalia alata</i>	20 -50	500 - 2000
19.	Shivane	<i>Gmelina arboria</i>	20 -50	500 -2000
20.	Kindal	<i>T.Paniculata</i>	20 - 40	500 - 1500
21.	Beete	<i>Dalbargia latifolia</i>	20 - 40	500 - 1500
22.	Tare	<i>T. belerica</i>	20 - 40	500 - 2000
23.	Bamboo	<i>Bambusa arundinasia</i>	20 - 40	500 - 2500
24.	Bamboo	<i>Dendrocalamus strictus</i>	20 – 40	500 – 2500
25.	Muthuga	<i>Butea monosperma</i>	20 - 40	400 - 1500
26.	Hippe	<i>Madhuca latifolia</i>	20 - 40	500 - 2000
27.	Sandal	<i>Santalum album</i>	20 - 50	400 - 1000
28.	Nelli	<i>Emblca officinalis</i>	20 - 40	500 - 2000
29.	Nerale	<i>Sizyzium cumini</i>	20 - 40	500 - 2000
30.	Dhaman	<i>Grevia tilifolia</i>	20 - 40	500 - 2000
31.	Kaval	<i>Careya arborea</i>	20 - 40	500 - 2000
32.	Harada	<i>Terminalia chebula</i>	20 - 40	500 - 2000

References

1. FAO (1976) Framework for Land Evaluation, Food and Agriculture Organization, Rome. 72 pp.
2. FAO (1983) Guidelines for Land Evaluation for Rainfed Agriculture, FAO, Rome, 237 pp.
3. IARI (1971) Soil Survey Manual, All India Soil and Land Use Survey Organization, IARI, New Delhi, 121 pp.
4. Katyal, J.C. and Rattan, R.K. (2003) Secondary and Micronutrients; Research Gap and Future Needs. Fert. News 48 (4); 9-20.
5. Naidu, L.G.K., Ramamurthy, V., Challa, O., Hegde, R. and Krishnan, P. (2006) Manual Soil Site Suitability Criteria for Major Crops, NBSS Publ. No. 129, NBSS & LUP, Nagpur, 118 pp.
6. Natarajan, A. and Dipak Sarkar (2010) Field Guide for Soil Survey, National Bureau of Soil Survey and Land Use Planning (ICAR), Nagpur, India.
7. Natarajan, A., Rajendra Hegde, Raj, J.N. and Shivananda Murthy, H.G. (2015) Implementation Manual for Sujala-III Project, Watershed Development Department, Bengaluru, Karnataka.
8. Sarma, V.A.K., Krishnan, P. and Budihal, S.L. (1987) Laboratory Manual, Tech. Bull. 23, NBSS & LUP, Nagpur.
9. Sehgal, J.L. (1990) Soil Resource Mapping of Different States of India; Why and How?, National Bureau of Soil Survey and Land Use Planning, Nagpur, 49 pp.
10. Shivaprasad, C.R., R.S. Reddy, J. Sehgal and M. Velayuthum (1998) Soils of Karnataka for Optimizing Land Use, NBSS Publ. No. 47b, NBSS & LUP, Nagpur, India.
11. Soil Survey Staff (2006) Keys to Soil Taxonomy, Tenth edition, U.S. Department of Agriculture/ NRCS, Washington DC, U.S.A.
12. Soil Survey Staff (2012) Soil Survey Manual, Handbook No. 18, USDA, Washington DC, USA.

Appendix I
Chandraki-1 (6E1b) Microwatershed
Soil Phase Information

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Madhepalli	1	5.67	BDLhB2g1	LMU-3	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut+Redgram (Gn+Rg)	Not Available	IIes	Graded bunding
Madhepalli	2	3.78	BDLhB2g1	LMU-3	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Madhepalli	3	4.58	BDLhB2g1	LMU-3	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut (Gn)	Not Available	IIes	Graded bunding
Madhepalli	4	6.12	HTKcB2	LMU-3	Shallow (25-50 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut (Gn)	Not Available	IIes	Graded bunding
Madhepalli	5	0.96	HTKcB2	LMU-3	Shallow (25-50 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Madhepalli	6	0.64	HTKcB2	LMU-3	Shallow (25-50 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut (Gn)	Not Available	IIes	Graded bunding
Madhepalli	7	0.32	HTKcB2	LMU-3	Shallow (25-50 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut (Gn)	Not Available	IIes	Graded bunding
Madhepalli	8	0.37	BDLiB2	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut (Gn)	Not Available	IIes	Graded bunding
Madhepalli	9	2.26	BDPiB3	LMU-4	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Redgram (Rg)	Not Available	IVes	Trench cum bunding
Madhepalli	10	2.72	BDLiB2	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut (Gn)	Not Available	IIes	Graded bunding
Madhepalli	11	0.39	RO	RO	RO	RO	RO	RO	RO	RO	Groundnut+Redgram (Gn+Rg)	Not Available	RO	RO
Madhepalli	12	4.86	JNKiB2g1	LMU-2	Moderately shallow (50-75 cm)	Sandy clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton+Redgram (Ct+Rg)	Not Available	IIes	Graded bunding
Madhepalli	13	2.69	JNKiB2g1	LMU-2	Moderately shallow (50-75 cm)	Sandy clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton+Redgram (Ct+Rg)	Not Available	IIes	Graded bunding
Madhepalli	23	3.11	BDLhB2g1	LMU-3	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut (Gn)	Not Available	IIes	Graded bunding
Madhepalli	24	0.0008	Habitation	Others	Others	Others	Others	Others	Others	Others	Groundnut+Redgram+Habitation (Gn+Rg+Hb)	Not Available	Others	Others
Madhepalli	57	0.001	Habitation	Others	Others	Others	Others	Others	Others	Others	NA	Not Available	Others	Others
Madhepalli	70	0.02	HSLiB2	LMU-1	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	NA	Not Available	IIes	Graded bunding
Madhepalli	71	0.74	Habitation	Others	Others	Others	Others	Others	Others	Others	Habitation (Hb)	Not Available	Others	Others
Madhepalli	72	0.91	HSLiB2	LMU-1	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IIes	Graded bunding
Madhepalli	73	0.25	HSLiB2	LMU-1	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IIes	Graded bunding
Madhepalli	74	0.65	BDLhB2g1	LMU-3	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IIes	Graded bunding
Madhepalli	75	0.92	BDLhB2g1	LMU-3	Shallow (25-50 cm)	Sandy clay	Gravelly	Very low (<50 mm/m)	Very gently	Moderate	Paddy (Pd)	Not	IIes	Graded

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
						loam	(15-35%)	mm/m)	sloping (1-3%)			Available		bunding
Madhepalli	76	2.57	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Madhepalli	77	0.98	HSLiB2	LMU-1	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Madhepalli	78	0.73	BDLhB2g1	LMU-3	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IIles	Graded bunding
Madhepalli	79	6.4	BDLiB2	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIles	Graded bunding
Madhepalli	80	0.42	BDLhB2g1	LMU-3	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut (Gn)	Not Available	IIles	Graded bunding
Madhepalli	81	0.44	BDLhB2g1	LMU-3	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut (Gn)	Not Available	IIles	Graded bunding
Madhepalli	82	0.59	HSLiB2	LMU-1	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Madhepalli	83	0.31	BDLhB2g1	LMU-3	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIles	Graded bunding
Madhepalli	84	1.01	BDLhB2g1	LMU-3	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IIles	Graded bunding
Madhepalli	85	0.27	BDLhB2g1	LMU-3	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Habitation (Hb)	Not Available	IIles	Graded bunding
Chandraki	1	4.81	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Chandraki	2	0.41	BDLiB2	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIles	Graded bunding
Chandraki	3	2.91	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	1 Bore Well	Iles	Graded bunding
Chandraki	4	1.55	BDLiB2	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	1 Bore Well	IIles	Graded bunding
Chandraki	5	5.97	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Forest (Fo)	Not Available	Iles	Graded bunding
Chandraki	6	0.13	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iles	Graded bunding
Chandraki	7	0.33	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iles	Graded bunding
Chandraki	8	0.63	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Acacia (Aca)	Not Available	Iles	Graded bunding
Chandraki	9	0.68	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Chandraki	10	0.6	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Chandraki	11	0.52	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Chandraki	12	0.29	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Chandraki	13	0.25	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Chandraki	14	0.01	SHTmB2	LMU-1	Moderately deep	Clay	Non gravelly	Low (51-100	Very gently	Moderate	Paddy (Pd)	Not	Iles	Graded

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
					(75-100 cm)		(<15%)	mm/m)	sloping (1-3%)			Available		bunding
Chandraki	16	2.53	BDLiB2	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton+Redgram (Ct+Rg)	Not Available	IIes	Graded bunding
Chandraki	22	1.1	Waterbody	Others	Others	Others	Others	Others	Others	Others	Scrub land (Sl)	Not Available	Others	Others
Chandraki	71	0.06	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iies	Graded bunding
Chandraki	72/2	0.16	BDLiB2	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Chandraki	72/3	0.66	BDLiB2	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Chandraki	83	0.11	BGDmB2g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Blackgram (Rg+Bm)	Not Available	Iies	Graded bunding
Chandraki	84	0.001	BGDmB2g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iies	Graded bunding
Chandraki	88/1	0.15	BDLiB2	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Chandraki	88/2	0.03	BDLiB2	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIes	Graded bunding
Chandraki	89	5.37	BDLiB2	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Chandraki	90	5.91	BDLiB2	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Chandraki	91	2.71	BDLiB2	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Chandraki	92	4.77	BGDmB2g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iies	Graded bunding
Chandraki	93	3.57	BGDmB2g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Blackgram (Rg+Bm)	Not Available	Iies	Graded bunding
Chandraki	94	4.19	BGDmB2g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iies	Graded bunding
Chandraki	95	6.15	BGDmB2g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iies	Graded bunding
Chandraki	96	4.87	BGDmB2g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iies	Graded bunding
Chandraki	97	4.45	BGDmB2g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iies	Graded bunding
Chandraki	98	4.21	BDLiB2	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Blackgram (Rg+Bm)	Not Available	IIes	Graded bunding
Chandraki	99	5.86	BGDmB2g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iies	Graded bunding
Chandraki	100	6.76	BDLiB2	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Chandraki	101	7.91	BGDmB2g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Blackgram (Rg+Bm)	Not Available	Iies	Graded bunding
Chandraki	102	5.56	BDLiB2	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Chandraki	103	4.41	BDLiB2	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly	Very low (<50	Very gently	Moderate	Redgram (Rg)	Not	IIes	Graded

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Graveliness (<15%)	Available Water Capacity (mm/m)	Slope (1-3%)	Soil Erosion	Current Land Use	Wells Available	Land Capability	Conservation Plan
Chandraki	104	2.74	Waterbody	Others	Others	Others	Others	Others	Others	Others	NA	Not Available	Others	Others
Chandraki	105	4.53	BDLiB2	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIles	Graded bunding
Chandraki	106	0.85	BDLiB2	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	IIles	Graded bunding
Chandraki	107	7	BDLiB2	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIles	Graded bunding
Chandraki	108	7.14	BDLiB2	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIles	Graded bunding
Chandraki	109	6.15	BDLiB2	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIles	Graded bunding
Chandraki	110	2.92	BDLiB2	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIles	Graded bunding
Chandraki	111	6.26	Waterbody	Others	Others	Others	Others	Others	Others	Others	NA	Not Available	Others	Others
Chandraki	112	2.53	BDLiB2	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Scrub land (Sl)	Not Available	IIles	Graded bunding
Chandraki	113	1.33	BDLiB2	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIles	Graded bunding
Chandraki	114	9.67	BDLiB2	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Forest (Fo)	Not Available	IIles	Graded bunding
Chandraki	115	8.2	BDLiB2	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Forest (Fo)	Not Available	IIles	Graded bunding
Chandraki	116	5.35	BDLiB2	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Forest (Fo)	Not Available	IIles	Graded bunding
Chandraki	117/1	49.92	BDLiB2	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Forest (Fo)	Not Available	IIles	Graded bunding
Chandraki	117/10	2.71	BDLiB2	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Forest (Fo)	Not Available	IIles	Graded bunding
Chandraki	117/11	3.71	BDLiB2	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Forest (Fo)	Not Available	IIles	Graded bunding
Chandraki	117/12	2.48	BDLiB2	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Forest (Fo)	Not Available	IIles	Graded bunding
Chandraki	117/13	2.39	BDLiB2	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIles	Graded bunding
Chandraki	117/14	1.87	BDLiB2	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Scrub land (Sl)	Not Available	IIles	Graded bunding
Chandraki	117/15	1.29	BDLiB2	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIles	Graded bunding
Chandraki	117/2	3.35	BDLiB2	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIles	Graded bunding
Chandraki	117/3	1.54	BDLiB2	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIles	Graded bunding
Chandraki	117/4	3.48	BDLiB2	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIles	Graded bunding
Chandraki	117/5	3.7	BDLiB2	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly	Very low (<50	Very gently	Moderate	Redgram (Rg)	Not	IIles	Graded

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness (<15%)	Available Water Capacity (mm/m)	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
									sloping (1-3%)			Available		bunding
Chandraki	117/6	1.64	BDLiB2	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIes	Graded bunding
Chandraki	117/7	1.07	BDLiB2	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIes	Graded bunding
Chandraki	117/8	3.17	BDLiB2	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Chandraki	117/9	3.07	BDLiB2	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Forest (Rg+Fo)	Not Available	IIes	Graded bunding
Chandraki	118	3.77	BGDmB2g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iies	Graded bunding
Chandraki	119	8.64	BGDmB2g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iies	Graded bunding
Chandraki	120	6.79	BGDmB2g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Blackgram (Rg+Bm)	Not Available	Iies	Graded bunding
Chandraki	121	6.07	BGDmB2g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Blackgram (Rg+Bm)	Not Available	Iies	Graded bunding
Chandraki	122	5.37	BGDmB2g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton+Redgram (Ct+Rg)	Not Available	Iies	Graded bunding
Chandraki	123	5.22	BGDmB2g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iies	Graded bunding
Chandraki	124	4.88	BGDmB2g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Forest (Rg+Fo)	Not Available	Iies	Graded bunding
Chandraki	125	3.98	BDLiB2	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Chandraki	126	6.05	BGDmB2g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iies	Graded bunding
Chandraki	127	4.75	BDLiB2	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Chandraki	128	3.66	BGDmB2g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iies	Graded bunding
Chandraki	129	3.05	BDLiB2	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Chandraki	130	3.92	BDLiB2	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Chandraki	131	6.3	BDLiB2	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Chandraki	132	0.31	BDLiB2	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIes	Graded bunding
Chandraki	133	5.64	Waterbody	Others	Others	Others	Others	Others	Others	Others	Scrub land (Sl)	Not Available	Others	Others
Chandraki	134	18.19	Waterbody	Others	Others	Others	Others	Others	Others	Others	NA	Not Available	Others	Others
Chandraki	135	2.87	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iies	Graded bunding
Chandraki	136	2.47	Waterbody	Others	Others	Others	Others	Others	Others	Others	Waterbody (Wb)	Not Available	Others	Others
Chandraki	137	2.86	Waterbody	Others	Others	Others	Others	Others	Others	Others	Waterbody (Wb)	Not	Others	Others

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
												Available		
Chandraki	138	7.64	Waterbody	Others	Others	Others	Others	Others	Others	Others	Waterbody (Wb)	Not Available	Others	Others
Chandraki	139	3.81	Waterbody	Others	Others	Others	Others	Others	Others	Others	Acacia (Aca)	Not Available	Others	Others
Chandraki	140	4.56	Waterbody	Others	Others	Others	Others	Others	Others	Others	Acacia (Aca)	Not Available	Others	Others
Chandraki	141	1.78	BDLiB2	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Chandraki	142	0.66	BDLiB2	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Acacia (Aca)	Not Available	IIes	Graded bunding
Chandraki	143/1	0.71	BDLiB2	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Acacia (Aca)	Not Available	IIes	Graded bunding
Chandraki	143/2	0.58	BDLiB2	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIes	Graded bunding
Chandraki	144	2.51	BDLiB2	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Chandraki	145/1	0.79	Waterbody	Others	Others	Others	Others	Others	Others	Others	Acacia (Aca)	Not Available	Others	Others
Chandraki	145/2	0.29	Waterbody	Others	Others	Others	Others	Others	Others	Others	Acacia (Aca)	Not Available	Others	Others
Chandraki	146	0.14	Waterbody	Others	Others	Others	Others	Others	Others	Others	Acacia (Aca)	Not Available	Others	Others
Chandraki	147	0.66	Waterbody	Others	Others	Others	Others	Others	Others	Others	Acacia (Aca)	Not Available	Others	Others
Chandraki	148	0.47	Waterbody	Others	Others	Others	Others	Others	Others	Others	Acacia (Aca)	Not Available	Others	Others
Chandraki	149	1.09	Waterbody	Others	Others	Others	Others	Others	Others	Others	Acacia (Aca)	Not Available	Others	Others
Chandraki	150	0.91	Waterbody	Others	Others	Others	Others	Others	Others	Others	Acacia (Aca)	Not Available	Others	Others
Chandraki	151/1	0.62	BDLiB2	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Chandraki	151/2	0.61	BDLiB2	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Chandraki	151/3	2.97	Waterbody	Others	Others	Others	Others	Others	Others	Others	Acacia (Aca)	Not Available	Others	Others
Chandraki	152	0.41	Waterbody	Others	Others	Others	Others	Others	Others	Others	Acacia (Aca)	Not Available	Others	Others
Chandraki	153	0.33	Waterbody	Others	Others	Others	Others	Others	Others	Others	Acacia (Aca)	Not Available	Others	Others
Chandraki	154	0.38	Waterbody	Others	Others	Others	Others	Others	Others	Others	Acacia (Aca)	1 Bore Well	Others	Others
Chandraki	155	0.96	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Acacia (Aca)	Not Available	IIes	Graded bunding
Chandraki	156	0.6	Waterbody	Others	Others	Others	Others	Others	Others	Others	Acacia (Aca)	Not Available	Others	Others

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Graveliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Chandraki	157	0.95	Waterbody	Others	Others	Others	Others	Others	Others	Others	Acacia (Aca)	Not Available	Others	Others
Chandraki	158	0.84	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iles	Graded bunding
Chandraki	159	0.58	Waterbody	Others	Others	Others	Others	Others	Others	Others	Not Available (NA)	Not Available	Others	Others
Chandraki	160	0.58	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Chandraki	161	0.8	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iles	Graded bunding
Chandraki	162	0.73	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iles	Graded bunding
Chandraki	163	0.68	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Habitation (Hb)	Not Available	Iles	Graded bunding
Chandraki	164	1.09	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Chandraki	165	0.49	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Chandraki	166	0.56	Waterbody	Others	Others	Others	Others	Others	Others	Others	Not Available (NA)	Not Available	Others	Others
Chandraki	167	0.49	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iles	Graded bunding
Chandraki	168	0.16	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iles	Graded bunding
Chandraki	169	1.08	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iles	Graded bunding
Chandraki	170/1	1.45	Habitation	Others	Others	Others	Others	Others	Others	Others	Habitation (Hb)	Not Available	Others	Others
Chandraki	170/2	0.46	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Habitation (Hb)	Not Available	Iles	Graded bunding
Chandraki	170/3	0.52	Habitation	Others	Others	Others	Others	Others	Others	Others	Habitation (Hb)	Not Available	Others	Others
Chandraki	170/4	0.22	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Habitation (Hb)	Not Available	Iles	Graded bunding
Chandraki	171	1.56	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Habitation (Hb)	Not Available	Iles	Graded bunding
Chandraki	172	3.09	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Habitation (Hb)	Not Available	Iles	Graded bunding
Chandraki	173/1	0.32	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Chandraki	173/2	4.3	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Scrub land (Sl)	Not Available	Iles	Graded bunding
Chandraki	174	0.3	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Chandraki	175	0.003	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Chandraki	178	0.02	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Chandraki	179	0.62	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Chandraki	180	0.23	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iles	Graded bunding
Chandraki	181	0.89	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iles	Graded bunding
Chandraki	182	0.49	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iles	Graded bunding
Chandraki	183	0.79	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Chandraki	184	0.86	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Chandraki	185	0.74	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Chandraki	186	0.67	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Chandraki	187	0.95	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Chandraki	188	0.81	Waterbody	Others	Others	Others	Others	Others	Others	Others	Not Available (NA)	Not Available	Others	Others
Chandraki	189	4.03	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iles	Graded bunding
Chandraki	190	0.33	Waterbody	Others	Others	Others	Others	Others	Others	Others	Not Available (NA)	Not Available	Others	Others
Chandraki	191	0.23	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iles	Graded bunding
Chandraki	192	2.04	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Chandraki	193	1.57	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Chandraki	194	1.36	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iles	Graded bunding
Chandraki	195	0.66	BDLiB2	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IIles	Graded bunding
Chandraki	196	0.65	Waterbody	Others	Others	Others	Others	Others	Others	Others	Paddy (Pd)	Not Available	Others	Others
Chandraki	197	0.56	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Chandraki	198	1.13	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Chandraki	199	0.86	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Chandraki	200	0.86	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Chandraki	201	0.66	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Chandraki	202	4.15	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	NA	Not Available	Iles	Graded bunding

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Graveliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Chandraki	203	0.35	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Chandraki	204	0.56	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Chandraki	205	0.77	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Chandraki	206	0.95	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Chandraki	207	0.81	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Chandraki	208	0.85	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Chandraki	209	0.48	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iles	Graded bunding
Chandraki	210	0.4	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Chandraki	211	0.91	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Chandraki	212	0.91	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Chandraki	213	0.25	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Chandraki	217	0.26	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Chandraki	230	0.26	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Chandraki	232	0.06	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Chandraki	233	0.77	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Chandraki	234	0.77	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Chandraki	235	0.91	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Chandraki	236	0.52	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Chandraki	237	0.45	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Chandraki	238	0.12	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Chandraki	239	0.04	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Chandraki	240	0.03	SHTmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Chandraki	315/3	0.42	Habitation	Others	Others	Others	Others	Others	Others	Others	Habitation (Hb)	Not Available	Others	Others
Chandraki	498	4.19	BGDmB2	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Graveliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Chandraki	499/1	1.3	BGDmB2	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Chandraki	499/2	1.28	BGDmB2	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Chandraki	500	7.68	BGDmB2	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Iles	Graded bunding
Chandraki	501	5.5	BGDmB2	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Blackgram (Rg+Bm)	Not Available	Iles	Graded bunding
Chandraki	502	2.48	BGDmB2	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Blackgram (Rg+Bm)	Not Available	Iles	Graded bunding
Chandraki	503	2.21	BDLiB2	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Blackgram (Rg+Bm)	Not Available	IIles	Graded bunding
Chandraki	521	0.83	BMNmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Chandraki	541	0.77	BDLiB2	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIles	Graded bunding
Chandraki	542	3.79	BMNmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Chandraki	543	2.74	BMNmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Chandraki	545	1.99	BMNmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Chandraki	546	6.64	BMNmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Blackgram (Rg+Bm)	Not Available	Iles	Graded bunding
Chandraki	547	6.64	BGDmB2	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Blackgram (Bm)	Not Available	Iles	Graded bunding
Chandraki	548	4.11	BDLiB2	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIles	Graded bunding
Chandraki	549	3.59	BDLiB2	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Paddy (Rg+Pd)	2 Bore Well	IIles	Graded bunding
Chandraki	550	4.18	BGDmB2	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Chandraki	551/1	1.74	BGDmB2	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Iles	Graded bunding
Chandraki	551/2	1.05	BGDmB2	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Chandraki	551/3	0.85	BGDmB2	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Chandraki	552	2.88	BGDmB2	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Chandraki	553	2.28	BGDmB2	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Chandraki	554	1.84	BGDmB2	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Habitation (Hb)	Not Available	Iles	Graded bunding
Chandraki	555	1.89	BDLiB2	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Scrub land (Sl)	Not Available	IIles	Graded bunding

Village	Survey No	Soil Reaction	Salinity	Organic Carbon	Available Phosphorus	Available Potassium	Available Sulphur	Available Boron	Available Iron	Available Manganese	Available Copper	Available Zinc
		(pH 7.8 - 8.4)	(<2 dsm)	0.75 %)	kg/ha)	337 kg/ha)	ppm)	1.0 ppm)	(>4.5 ppm)	1.0 ppm)	0.2 ppm)	0.6 ppm)
Chandraki	546	Strongly alkaline (pH 8.4 - 9.0)	Non saline (<2 dsm)	High (> 0.75 %)	Low (< 23 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	Medium (0.5 - 1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)
Chandraki	547	Strongly alkaline (pH 8.4 - 9.0)	Non saline (<2 dsm)	High (> 0.75 %)	Low (< 23 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	Medium (0.5 - 1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)
Chandraki	548	Strongly alkaline (pH 8.4 - 9.0)	Non saline (<2 dsm)	High (> 0.75 %)	Low (< 23 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	Medium (0.5 - 1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)
Chandraki	549	Strongly alkaline (pH 8.4 - 9.0)	Non saline (<2 dsm)	High (> 0.75 %)	Low (< 23 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	Medium (0.5 - 1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Chandraki	550	Moderately alkaline (pH 7.8 - 8.4)	Non saline (<2 dsm)	High (> 0.75 %)	Low (< 23 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	Medium (0.5 - 1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)
Chandraki	551/1	Moderately alkaline (pH 7.8 - 8.4)	Non saline (<2 dsm)	High (> 0.75 %)	Low (< 23 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	Medium (0.5 - 1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)
Chandraki	551/2	Moderately alkaline (pH 7.8 - 8.4)	Non saline (<2 dsm)	High (> 0.75 %)	Low (< 23 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	Medium (0.5 - 1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)
Chandraki	551/3	Moderately alkaline (pH 7.8 - 8.4)	Non saline (<2 dsm)	High (> 0.75 %)	Low (< 23 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	Medium (0.5 - 1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)
Chandraki	552	Slightly alkaline (pH 7.3 - 7.8)	Non saline (<2 dsm)	High (> 0.75 %)	Low (< 23 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	Medium (0.5 - 1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)
Chandraki	553	Slightly alkaline (pH 7.3 - 7.8)	Non saline (<2 dsm)	High (> 0.75 %)	Low (< 23 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	Medium (0.5 - 1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)
Chandraki	554	Slightly alkaline (pH 7.3 - 7.8)	Non saline (<2 dsm)	High (> 0.75 %)	Low (< 23 kg/ha)	High (> 337 kg/ha)	Low (<10 ppm)	Medium (0.5 - 1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)
Chandraki	555	Slightly acid (pH 6.0 - 6.5)	Non saline (<2 dsm)	High (> 0.75 %)	High (> 57 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	High (> 1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)

Appendix III
Chandraki-1 (6E1b) Microwatershed
Soil Suitability Information

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry	
Madhepalli	1	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Madhepalli	2	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Madhepalli	3	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Madhepalli	4	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1r	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Madhepalli	5	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1r	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Madhepalli	6	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1r	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Madhepalli	7	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1r	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Madhepalli	8	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Madhepalli	9	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Madhepalli	10	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Madhepalli	11	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0
Madhepalli	12	N1r	S2r	S3r	S2rt	S3r	S3t	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	N1n	S3r	S3r	S2r	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Madhepalli	13	N1r	S2r	S3r	S2rt	S3r	S3t	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	N1n	S3r	S3r	S2r	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Madhepalli	23	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Madhepalli	24	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Madhepalli	57	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Madhepalli	70	S3r	S1	S2r	S1	S2r	S2r	S3r	S2r	S3t	S2r	S2r	S1	S2r	S2r	S3n	S3r	S2r	S2t	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S1	S2r	S2r
Madhepalli	71	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Madhepalli	72	S3r	S1	S2r	S1	S2r	S2r	S3r	S2r	S3t	S2r	S2r	S1	S2r	S2r	S3n	S3r	S2r	S2t	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r	

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry
Madhepalli	73	S3r	S1	S2r	S1	S2r	S2r	S3r	S2r	S3t	S2r	S2r	S1	S2r	S2r	S3n	S3r	S2r	S2t	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r
Madhepalli	74	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r
Madhepalli	75	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r
Madhepalli	76	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r
Madhepalli	77	S3r	S1	S2r	S1	S2r	S2r	S3r	S2r	S3t	S2r	S2r	S1	S2r	S2r	S3n	S3r	S2r	S2t	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r
Madhepalli	78	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r
Madhepalli	79	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r
Madhepalli	80	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r
Madhepalli	81	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r
Madhepalli	82	S3r	S1	S2r	S1	S2r	S2r	S3r	S2r	S3t	S2r	S2r	S1	S2r	S2r	S3n	S3r	S2r	S2t	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r
Madhepalli	83	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r
Madhepalli	84	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r
Madhepalli	85	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r
Chandra ki	1	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r
Chandra ki	2	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r
Chandra ki	3	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r
Chandra ki	4	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r
Chandra ki	5	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r
Chandra ki	6	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r
Chandra ki	7	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r
Chandra ki	8	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r
Chandra	9	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry
ki																														
Chandra ki	10	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r
Chandra ki	11	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r
Chandra ki	12	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r
Chandra ki	13	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r
Chandra ki	14	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r
Chandra ki	16	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r
Chandra ki	22	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs
Chandra ki	71	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r
Chandra ki	72/2	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r
Chandra ki	72/3	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r
Chandra ki	83	S3t	S2t	S3t	S1	S3t	S1	S2rt	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S2t	S3t	S2t	S2t	S2t	S2t	S3t	S2t	S2t	S3t
Chandra ki	84	S3t	S2t	S3t	S1	S3t	S1	S2rt	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S2t	S3t	S2t	S2t	S2t	S2t	S3t	S2t	S2t	S3t
Chandra ki	88/1	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r
Chandra ki	88/2	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r
Chandra ki	89	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r
Chandra ki	90	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r
Chandra ki	91	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r
Chandra ki	92	S3t	S2t	S3t	S1	S3t	S1	S2rt	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S2t	S3t	S2t	S2t	S2t	S2t	S3t	S2t	S2t	S3t
Chandra ki	93	S3t	S2t	S3t	S1	S3t	S1	S2rt	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S2t	S3t	S2t	S2t	S2t	S2t	S3t	S2t	S2t	S3t
Chandra ki	94	S3t	S2t	S3t	S1	S3t	S1	S2rt	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S2t	S3t	S2t	S2t	S2t	S2t	S3t	S2t	S2t	S3t
Chandra ki	95	S3t	S2t	S3t	S1	S3t	S1	S2rt	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S2t	S3t	S2t	S2t	S2t	S2t	S3t	S2t	S2t	S3t

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry	
Chandra ki	96	S3t	S2t	S3t	S1	S3t	S1	S2rt	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S2t	S3t	S2t	S2t	S2t	S2t	S3t	S2t	S2t	S3t	
Chandra ki	97	S3t	S2t	S3t	S1	S3t	S1	S2rt	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S2t	S3t	S2t	S2t	S2t	S2t	S3t	S2t	S2t	S3t	
Chandra ki	98	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	N1r	
Chandra ki	99	S3t	S2t	S3t	S1	S3t	S1	S2rt	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S2t	S3t	S2t	S2t	S2t	S2t	S3t	S2t	S2t	S3t	
Chandra ki	100	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	N1r	
Chandra ki	101	S3t	S2t	S3t	S1	S3t	S1	S2rt	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S2t	S3t	S2t	S2t	S2t	S2t	S3t	S2t	S2t	S3t	
Chandra ki	102	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	N1r	
Chandra ki	103	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	N1r	
Chandra ki	104	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs
Chandra ki	105	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	N1r	
Chandra ki	106	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	N1r	
Chandra ki	107	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	N1r	
Chandra ki	108	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	N1r	
Chandra ki	109	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	N1r	
Chandra ki	110	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	N1r	
Chandra ki	111	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs
Chandra ki	112	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	N1r	
Chandra ki	113	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	N1r	
Chandra ki	114	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	N1r	
Chandra ki	115	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	N1r	
Chandra ki	116	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	N1r	
Chandra	117/	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	N1r	

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry	
ki	1																														
Chandra ki	117/10	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Chandra ki	117/11	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Chandra ki	117/12	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Chandra ki	117/13	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Chandra ki	117/14	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Chandra ki	117/15	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Chandra ki	117/2	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Chandra ki	117/3	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Chandra ki	117/4	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Chandra ki	117/5	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Chandra ki	117/6	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Chandra ki	117/7	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Chandra ki	117/8	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Chandra ki	117/9	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Chandra ki	118	S3t	S2t	S3t	S1	S3t	S1	S2rt	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S2t	S3t	S2t	S2t	S2t	S2t	S3t	S2t	S2t	S3t	
Chandra ki	119	S3t	S2t	S3t	S1	S3t	S1	S2rt	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S2t	S3t	S2t	S2t	S2t	S2t	S3t	S2t	S2t	S3t	
Chandra ki	120	S3t	S2t	S3t	S1	S3t	S1	S2rt	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S2t	S3t	S2t	S2t	S2t	S2t	S3t	S2t	S2t	S3t	
Chandra ki	121	S3t	S2t	S3t	S1	S3t	S1	S2rt	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S2t	S3t	S2t	S2t	S2t	S2t	S3t	S2t	S2t	S3t	
Chandra ki	122	S3t	S2t	S3t	S1	S3t	S1	S2rt	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S2t	S3t	S2t	S2t	S2t	S2t	S3t	S2t	S2t	S3t	
Chandra ki	123	S3t	S2t	S3t	S1	S3t	S1	S2rt	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S2t	S3t	S2t	S2t	S2t	S2t	S3t	S2t	S2t	S3t	
Chandra ki	124	S3t	S2t	S3t	S1	S3t	S1	S2rt	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S2t	S3t	S2t	S2t	S2t	S2t	S3t	S2t	S2t	S3t	

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry
Chandra ki	125	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	S3r	N1r	N1r
Chandra ki	126	S3t	S2t	S3t	S1	S3t	S1	S2rt	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S2t	S3t	S2t	S2t	S2t	S2t	S3t	S2t	S2t	S3t
Chandra ki	127	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	S3r	N1r	N1r
Chandra ki	128	S3t	S2t	S3t	S1	S3t	S1	S2rt	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S2t	S3t	S2t	S2t	S2t	S2t	S3t	S2t	S2t	S3t
Chandra ki	129	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	S3r	N1r	N1r
Chandra ki	130	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	S3r	N1r	N1r
Chandra ki	131	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	S3r	N1r	N1r
Chandra ki	132	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	S3r	N1r	N1r
Chandra ki	133	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs
Chandra ki	134	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs
Chandra ki	135	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r
Chandra ki	136	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs
Chandra ki	137	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs
Chandra ki	138	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs
Chandra ki	139	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs
Chandra ki	140	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs
Chandra ki	141	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	S3r	N1r	N1r
Chandra ki	142	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	S3r	N1r	N1r
Chandra ki	143/1	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	S3r	N1r	N1r
Chandra ki	143/2	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	S3r	N1r	N1r
Chandra ki	144	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	S3r	N1r	N1r
Chandra	145/	Othe	Othe	Othe	Othe	Othe	Othe	Othe	Othe	Othe	Othe	Othe	Othe	Othe	Othe	Othe	Othe	Othe	Othe	Othe	Othe	Othe	Othe	Othe	Othe	Othe	Othe	Othe	Othe	Othe

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry	
Chandra ki	164	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r	
Chandra ki	165	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r	
Chandra ki	166	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs
Chandra ki	167	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r	
Chandra ki	168	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r	
Chandra ki	169	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r	
Chandra ki	170/1	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs
Chandra ki	170/2	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r	
Chandra ki	170/3	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs
Chandra ki	170/4	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r	
Chandra ki	171	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r	
Chandra ki	172	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r	
Chandra ki	173/1	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r	
Chandra ki	173/2	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r	
Chandra ki	174	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r	
Chandra ki	175	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r	
Chandra ki	178	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r	
Chandra ki	179	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r	
Chandra ki	180	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r	
Chandra ki	181	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r	
Chandra ki	182	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r	
Chandra	183	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r	

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry
ki																														
Chandra ki	184	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r
Chandra ki	185	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r
Chandra ki	186	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r
Chandra ki	187	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r
Chandra ki	188	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs
Chandra ki	189	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r
Chandra ki	190	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs
Chandra ki	191	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r
Chandra ki	192	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r
Chandra ki	193	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r
Chandra ki	194	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r
Chandra ki	195	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r
Chandra ki	196	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs
Chandra ki	197	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r
Chandra ki	198	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r
Chandra ki	199	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r
Chandra ki	200	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r
Chandra ki	201	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r
Chandra ki	202	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r
Chandra ki	203	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r
Chandra ki	204	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry	
Chandra ki	205	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r	
Chandra ki	206	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r	
Chandra ki	207	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r	
Chandra ki	208	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r	
Chandra ki	209	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r	
Chandra ki	210	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r	
Chandra ki	211	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r	
Chandra ki	212	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r	
Chandra ki	213	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r	
Chandra ki	217	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r	
Chandra ki	230	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r	
Chandra ki	232	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r	
Chandra ki	233	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r	
Chandra ki	234	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r	
Chandra ki	235	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r	
Chandra ki	236	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r	
Chandra ki	237	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r	
Chandra ki	238	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r	
Chandra ki	239	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r	
Chandra ki	240	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r	
Chandra ki	315/3	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs
Chandra	498	S3t	S2t	S3t	S1	S3t	S1	S2rt	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S2t	S3t	S2t	S2t	S2t	S2t	S3t	S2t	S2t	S3t	

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry
ki																														
Chandra ki	499/1	S3t	S2t	S3t	S1	S3t	S1	S2rt	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S2t	S3t	S2t	S2t	S2t	S2t	S3t	S2t	S2t	S3t
Chandra ki	499/2	S3t	S2t	S3t	S1	S3t	S1	S2rt	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S2t	S3t	S2t	S2t	S2t	S2t	S3t	S2t	S2t	S3t
Chandra ki	500	S3t	S2t	S3t	S1	S3t	S1	S2rt	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S2t	S3t	S2t	S2t	S2t	S2t	S3t	S2t	S2t	S3t
Chandra ki	501	S3t	S2t	S3t	S1	S3t	S1	S2rt	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S2t	S3t	S2t	S2t	S2t	S2t	S3t	S2t	S2t	S3t
Chandra ki	502	S3t	S2t	S3t	S1	S3t	S1	S2rt	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S2t	S3t	S2t	S2t	S2t	S2t	S3t	S2t	S2t	S3t
Chandra ki	503	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r
Chandra ki	521	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz
Chandra ki	541	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r
Chandra ki	542	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz
Chandra ki	543	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz
Chandra ki	545	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz
Chandra ki	546	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz
Chandra ki	547	S3t	S2t	S3t	S1	S3t	S1	S2rt	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S2t	S3t	S2t	S2t	S2t	S2t	S3t	S2t	S2t	S3t
Chandra ki	548	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r
Chandra ki	549	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r
Chandra ki	550	S3t	S2t	S3t	S1	S3t	S1	S2rt	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S2t	S3t	S2t	S2t	S2t	S2t	S3t	S2t	S2t	S3t
Chandra ki	551/1	S3t	S2t	S3t	S1	S3t	S1	S2rt	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S2t	S3t	S2t	S2t	S2t	S2t	S3t	S2t	S2t	S3t
Chandra ki	551/2	S3t	S2t	S3t	S1	S3t	S1	S2rt	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S2t	S3t	S2t	S2t	S2t	S2t	S3t	S2t	S2t	S3t
Chandra ki	551/3	S3t	S2t	S3t	S1	S3t	S1	S2rt	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S2t	S3t	S2t	S2t	S2t	S2t	S3t	S2t	S2t	S3t
Chandra ki	552	S3t	S2t	S3t	S1	S3t	S1	S2rt	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S2t	S3t	S2t	S2t	S2t	S2t	S3t	S2t	S2t	S3t
Chandra ki	553	S3t	S2t	S3t	S1	S3t	S1	S2rt	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S2t	S3t	S2t	S2t	S2t	S2t	S3t	S2t	S2t	S3t

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry
Chandra ki	554	S3t	S2t	S3t	S1	S3t	S1	S2rt	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S2t	S3t	S2t	S2t	S2t	S2t	S3t	S2t	S2t	S3t
Chandra ki	555	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r

RO-Rock outcrops

PART-B

SOCIO-ECONOMIC STATUS OF FARM HOUSEHOLDS

CONTENTS

1	Findings of the socio-economic survey	1-3
2	Introduction	5-6
3	Methodology	7-8
4	Salient features of the survey	9-29
5	Summary	31-35

LIST OF TABLES

1	Households sampled for socio economic survey	9
2	Population characteristics	9
3	Age wise classification of household members	9
4	Education level of household members	10
5	Occupation of household heads	10
6	Occupation of family members	10
7	Institutional participation of household members	11
8	Type of house owned by households	11
9	Durable assets owned by households	11
10	Average value of durable assets owned by households	11
11	Farm implements owned by households	12
12	Average value of farm implements	12
13	Livestock possession by households	12
14	Average labour availability	13
15	Adequacy of hired labour	13
16	Migration among the households	13
17	Average distance and duration of migration	13
18	Purpose of migration	14
19	Distribution of land (ha)	14
20	Average land value (Rs./ha)	14
21	Status of Open wells	14
22	Source of irrigation	15
23	Depth of water(Avg in meters)	15
24	Cropping pattern	15
25	Cropping intensity	15
26	Possession of bank account and saving	15
27	Borrowing status	16
28	Source of credit	16
29	Avg. Credit amount	16
30	Purpose of credit borrowed - Institutional Credit	16
31	Purpose of credit borrowed Non - Institutional Credit	17
32	Repayment status of credit borrowed from institutional Source	17

33	Repayment status of credit borrowed from Non-institutional Source	17
34.a	Cost of cultivation of Red gram	18
34.b	Cost of cultivation of Cotton	19
34.c	Cost of cultivation of Bengal gram	20
34.d	Cost of cultivation of Jowar	21
34.e	Cost of cultivation of Paddy	22
35	Adequacy of fodder	23
36	Average Annual gross income	23
37	Average annual expenditure	23
38	Horticulture species grown	23
39	Forest species grown	24
40	Average additional investment capacity	24
41	Source of funds for additional investment	24
42	Marketing of the agricultural produce	25
43	Marketing channels used for sale of agricultural produce	25
44	Mode of transport of agricultural produce	25
45	Incidence of soil and water erosion problems	25
46	Interest shown towards soil testing	26
47	Soil and water conservation practices and structures	26
48	Status soil and water conservation structures	26
49	Agencies involved in the soil and water conservation structures	26
50	Usage pattern of fuel for domestic use	27
51	Source of drinking water	27
52	Source of light	27
53	Existence of sanitary toilet facility	27
54	Possession of public distribution system (PDS) card	27
55	Participation in NREGA programme	28
56	Adequacy of food items	28
57	Inadequacy of food items	28
58	Farming constraints experienced	29

FINDINGS OF THE SOCIO-ECONOMIC SURVEY

- ❖ *The survey was conducted in Chandaraki-1 is located at North latitude 16° 52' 29.368" and 16° 50' 46.558" and East longitude 77° 27' 47.027" and 77° 26' 17.986" covering an area of about 603.07 ha coming under Chandaraki and Madhepalli villages of Yadagiri taluk.*
- ❖ *Socio-economic analysis of Chandaraki-1 micro watersheds of Chandaraki sub-watershed, Yadgiri taluk & District indicated that, out of the total sample of 36 farmers were sampled in Chandaraki-1 micro-watershed among households surveyed 14 (38.89%) were marginal, 7 (19.44%) were small, 10 (27.78 %) were semi medium and 3 (8.33 %) were medium large farmers. 2 landless farmers were also interviewed for the survey.*
- ❖ *The population characteristics of households indicated that, there were 89 (55.97%) men and 70 (44.03 %) were women. The average population of landless was 2.5, marginal farmers were 4.6, small farmers were 4.3, semi medium farmers were 4.8 and medium farmers were 4.*
- ❖ *Majority of the respondents (38.36%) were in the age group of 16-35 years.*
- ❖ *Education level of the sample households indicated that, there were 44.65 per cent of illiterates, 17.61 per cent of them had primary school education, 13.84 per cent high school education, 8.18 per cent of them had PUC education, 0.63 per cent of them had Diploma, 9.43 per cent attained graduation and 3.14 them had other education.*
- ❖ *About, 80.56 per cent of household heads practicing agriculture and 2.78 per cent of the household heads were engaged as agricultural labourers.*
- ❖ *Agriculture was the major occupation for 80.56 per cent of the household members.*
- ❖ *In the study area, 72.22 per cent of the households possess katcha house and 8.33 per cent possess pucca house.*
- ❖ *The durable assets owned by the households showed that, 72.22 per cent possess TV, 80.56 per cent possess mixer grinder, 94.44 per cent possess mobile phones and 30.56 per cent possess motor cycles.*
- ❖ *Farm implements owned by the households indicated that, 8.33 per cent of the households possess Bullock Cart, 36.11 per cent possess plough and 77.78 per cent possess Weeder, 8.33 per cent possess tractor, 5.56 per cent possess Harvester and 36.11 per cent possess Thresher.*
- ❖ *Regarding livestock possession by the households, 2.78 per cent possess local cow and 5.56 per cent possess buffalo.*
- ❖ *The average labour availability in the study area showed that, own men and women labour availability in the micro watershed was 14.06 each, while the hired labour (men) availability was 1.22.*

- ❖ *Further, 5.56 per cent of the households opined that hired labour was inadequate during the agricultural season.*
- ❖ *In the study area, about 0.63 per cent of the respondents migrated from the micro watershed in search of jobs with an average distance of 850.00 kms for about 4.00 months.*
- ❖ *Out of the total land holding of the sample respondents 94.31 per cent (59.84 ha) of the area is under dry condition and the remaining 4.24 per cent area is irrigated land.*
- ❖ *Open well was the major source of irrigation for 5.56 per cent of the households.*
- ❖ *The major crops grown by sample farmers are Red gram, Cotton, Bengal gram, Jowar and Paddy and cropping intensity was recorded as 100.00 per cent.*
- ❖ *Out of the sample households 94.44 percent possessed bank account.*
- ❖ *About 94.44 per cent of the respondents borrowed credit from various sources*
- ❖ *Among the credit borrowed by households, 13.64 per cent have borrowed loan from commercial banks and 86.36 per cent from co-operative/Grameena bank.*
- ❖ *Majority of the respondents (95.45%) have borrowed loan for agriculture purpose.*
- ❖ *The per hectare cost of cultivation for Red gram, Cotton, Bengal gram, Jowar and Paddy was Rs.48868.88, 44389.25, 18089.82, 26988.16 and 23946.94 with benefit cost ratio of 1:0.90, 1: 1.00, 1: 3.80, 1: 0.50 and 1:0.70 respectively.*
- ❖ *Further, 25.00 per cent of the households opined that dry fodder was adequate and 8.33 per cent of the households have opined that the green fodder was adequate.*
- ❖ *The average annual gross income of the farmers was Rs. 76055.56 in micro-watershed, of which Rs. 41027.78 comes from agriculture.*
- ❖ *Sampled households have grown 8 horticulture trees and 100 forestry trees together in the fields and back yards.*
- ❖ *Households have an average investment capacity of Rs. 12361.11 for land development.*
- ❖ *Source of funds for additional investment is concerned, 88.89 per cent depends on own funds and 2.78 per cent depends on bank loan for land development activities.*
- ❖ *Regarding marketing channels, 116.67 per cent of the households have sold agricultural produce to the local/village merchants.*
- ❖ *Further, 116.67 per cent of the households have used tractor for the transport of agriculture commodity.*
- ❖ *Majority of the farmers (88.89%) have experienced soil and water erosion problems in the watershed and 94.44 per cent of the households were interested towards soil testing.*

- ❖ *Fire was the major source of fuel for domestic use for 88.89 per cent of the households and 11.11 per cent households has LPG connection.*
- ❖ *Piped supply was the major source for drinking water for 97.22 per cent of the households.*
- ❖ *Electricity was the major source of light for 100.00 per cent of the households.*
- ❖ *In the study area, 30.56 per cent of the households possess toilet facility.*
- ❖ *Regarding possession of PDS card, 100.00 per cent of the households possessed BPL card.*
- ❖ *Households opined that, the requirement of cereals (86.11%), pulses (83.33%) and oilseeds (5.56%) are adequate for consumption.*
- ❖ *Farming constraints experienced by households in the micro watersheds were lower fertility status of the soil (91.67%) wild animal menace on farm field (55.56%), frequent incidence of pest and diseases (55.56%), inadequacy of irrigation water (72.22%), high cost of fertilizers and plant protection chemicals (63.89%), high rate of interest on credit (66.67%), low price for the agricultural commodities (69.44%), lack of marketing facilities in the area (52.78%), inadequate extension services (41.67%), lack of transport for safe transport of the agricultural produce to the market (36.11%), Less rainfall (2.78%) and Source of Agri-technology information (Newspaper/ TV/Mobile) (2.78%).*

INTRODUCTION

Soil and water are the two precious natural resources which are essential for crop production and existence of life on earth. Rainfed agriculture is under severe stress due to various constraints related to agriculture like uneven and erratic distribution of rainfall, indiscriminate use of fertilizers, chemicals and pesticides, adoption of improper land management practices, soil erosion, decline in soil fertility, decline in ground water resources leading to low crop productivity. The area under rainfed agriculture has to be managed effectively using the best available practices to enhance the production of food, fodder and fuel. This is possible if the land resources are characterized at each parcel of land through detailed land resource inventory using the best available techniques of remote sensing, GPS and GIS. The watershed development programs are aimed at the sustainable distribution of its resources and the process of creating and implementing plans, programs, and projects to sustain and enhance watershed functions that affect the plant, animal and human communities within a watershed boundary.

World Bank funded KWDP II, SUJALA III project was implemented in with Broad objective of demonstrating more effective watershed management through greater integration of programmes related to rain-fed agriculture, innovative and science based approaches and strengthen institutional capacities and If successful, it is expected that the systems and tools could be mainstreamed into the overall IWMP in the State of Karnataka and in time, throughout other IWMP operations in India. With this background the socio-economic survey has been carried out with following specific objectives:

1. To understand the demographic features of the households in the micro-watershed
2. To understand the extent of family labour available and additional employment opportunities available within the village.
3. To know the status of assets of households in the micro-watershed for suggesting possible improvements.
4. To study the cropping pattern, cropped area and productivity levels of different households in micro-watershed.
5. To determine the type and extent of livestock owned by different categories of HHs
6. Availability of fodder and level of livestock management.

Scope and importance of survey

Survey helps in identification of different socio-economic and resource use-patterns of farmers at the Micro watershed. Household survey provides demographic features, labour force, and levels of education; land ownership and asset position (including livestock and other household assets) of surveyed households; and

cropping patterns, input intensities, and average crop yields from farmers' fields. It also discusses crop utilization and the degree of commercialization of production in the areas; farmers' access to and utilization of credit from formal and informal sources; and the level of adoption and use of soil, water, and pest management technologies.

METHODOLOGY

The description of the methods, components selected for the survey and procedures followed in conducting the baseline survey are furnished under the following heads.

1. Description of the study area

Yadgir District is one of the 30 districts of Karnataka state in southern India. This district was carved out from the erstwhile Gulbarga district as the 30th district of Karnataka on 10 April 2010. Yadgir town is the administrative headquarters of the district. The district comprises of 3 taluks namely, Shahapur, Yadgiri and Shorapur (There are 16 hoblies, 117 Gram Panchayats, 4 Municipalities, 8 Towns/ Urban agglomeration and 487 inhabited & 32 un-inhabited villages The district occupies an area of 5,160.88 km².

Yadgir district is the second smallest district in the state, area wise is very rich in cultural traditions. The vast stretch of fertile black soil of the district is known for bumper red gram and jowar crops. The district is a "Daal bowl" of the state. The district is also known for cluster of cement industries and a distinct stone popularly known as "Malakheda Stone". Two main rivers, Krishna and Bhima, and a few tributaries flow in this region. Krishna and Bhima Rivers drain the district. They constitute the two major river basins of the district. Kagna and Amarja are the two sub-basins of Bhima River, which occur within the geographical area of the district

According to the 2011 census Yadgir district has a population of 1, 172,985, roughly equal to the nation of Timor-Leste or the US state of Rhode Island. This gives it a ranking of 404th in India (out of a total of 640). The district has a population density of 224 inhabitants per square kilometre (580/sq mi). Its population growth rate over the decade 2001-2011 was 22.67%. Yadgir has a sex ratio of 984 females for every 1000 males, and a literacy rate of 52.36%.

2. Locale of the survey and description of the micro-watershed and

The study was conducted in Chandaraki-1 micro-watershed (Chandaraki sub-watershed, Yadgiri taluk & District) is located at North latitude 16^o 52' 29.368" and 16^o 50' 46.558" and East longitude 77^o 27' 47.027" and 77^o 26' 17.986" covering an area of about 603.07 ha bounded by under Chandaraki and Madhepalli Villages.

3. Selection of the respondents for the study

The micro-watershed is marked with 320 square meters grids. One farmer from every alternate grid in the micro-watershed was selected for the study and interviewed for socio-economic data. Totally 36 households were interviewed for the survey.

4. The parameters considered for socio-economic survey of households

Two forms of data were collected from the micro-watershed which includes primary data from the farm households and secondary data about the villages under the micro-watershed jurisdiction.

The following parameters were considered for the primary data collection about the socio-economic data of the households, (1) Demographic information, (2) Farm and durable assets owned by households, (3) Livestock possession, (4) Labour availability, (5) Level of migration in the village, Land holding, (7) Cropping pattern, (8) Source of irrigation, (9) Borrowing status, (10) Cost of cultivation of major crops, (11) Economics of subsidiary activities, (12) Fodder availability, (13) Family annual income from different sources, (14) Horticulture and forestry species grown, (15) Additional investment capacity, (16) Marketing practices, (17) Status of soil and water conservation structure, (18) Access to basic needs and (19) Constraints and suggestion.

The following parameters were considered for the secondary data regarding the villages under the micro-watershed jurisdiction, (1) Number of villages in each micro-watershed jurisdiction, (2) Village wise number of households, (3) Geographical area of the villages, (4) Cultivable area including rainfed and irrigated, (5) Number and type of house in each village, (6) Human and livestock population, (7) Facilities in the village such as roads, transport facility for conveyance, drinking water supply, street light and (8) Community based organizations in the villages.

5. Development of interview schedule and data collection

Taking into the consideration the objectives of the survey, an interview schedule was prepared after thorough consultation with the experts in the field of social sciences. A comprehensive interview schedule covering all the major parameters for measuring the socio-economic situation was developed.

6. Tools used to analyze the data

The statistical components such as frequency and percentage were used to analyze the data.

Abbreviations used in the report

LL=Landless

MF=Marginal Farmers

SF=Small farmers

SMF=Semi medium farmers

MDF=Medium farmers

LF=Large Farmers

FINDINGS OF THE SURVEY

This chapter deals with systematic presentation of results of the survey. Keeping in view the objectives, the salient features of the survey are presented under the following headings.

Households sampled for socio-economic survey: The data on households sampled for socio economic survey in Chandaraki-1 Micro watershed is presented in Table 1 and it indicated that 36 farmers were sampled in Chandaraki-1 micro-watershed among households surveyed 14 (38.89%) were marginal, 7 (19.44%) were small, 10 (27.78 %) were semi medium and 3 (8.33 %) were medium large farmers. 2 landless farmers were also interviewed for the survey.

Table 1. Households sampled for socio economic survey in Chandaraki-1 micro-watershed

Sl. No.	Particulars	LL (2)		MF (14)		SF (7)		SMF (10)		MDF (3)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Farmers	2	5.56	14	38.9	7	19.4	10	27.8	3	8.33	36	100

Population characteristics: The population characteristics of households sampled for socio-economic survey in Chandaraki-1 Micro watershed is presented in Table 2. The data indicated that, there were 89 (55.97%) men and 70 (44.03%) were women. The average population of landless was 2.5, marginal farmers were 4.6, small farmers were 4.3, semi medium farmers were 4.8 and medium farmers were 4.

Table 2. Population characteristics in Chandaraki-1 micro-watershed

Sl. No.	Particulars	LL (5)		MF (64)		SF (30)		SMF (48)		MDF (12)		All (159)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Men	4	80	36	56	17	57	26	54.2	6	50	89	56
2	Women	1	20	28	44	13	43	22	45.8	6	50	70	44
Total		5	100	64	100	30	100	48	100	12	100	159	100
Average		2.5		4.6		4.3		4.8		4.0		4.4	

Age wise classification of population: The age wise classification of household members in Chandaraki-1 Micro watershed is presented in Table 3. The indicated that, 35 (22.01%) of population were 0-15 years of age, 61 (38.36%) were 16-35 years of age, 48(30.19%) were 36-60 years of age and 15 (9.43 %) were above 61 years of age.

Table 3: Age wise classification of members of the household in Chandaraki-1 micro-watershed

Sl.No.	Particulars	LL (5)		MF (64)		SF (30)		SMF (48)		MDF (12)		All (159)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	0-15 years of age	2	40	16	25	7	23.3	6	12.5	4	33	35	22.01
2	16-35 years of age	2	40	30	46.9	10	33.3	17	35.42	2	17	61	38.36
3	36-60 years of age	1	20	13	20.3	11	36.7	17	35.42	6	50	48	30.19
4	> 61 years	0	0	5	7.81	2	6.67	8	16.67	0	0	15	9.43
Total		5	100	64	100	30	100	48	100	12	100	159	100

Education level of household members: Education level of household members in Chandaraki-1 Micro watershed is presented in Table 4. The results indicated that, there were 44.65 per cent of illiterates, 17.61 per cent of them had primary school education, 13.84 per cent high school education, 8.18 per cent of them had PUC education, 0.63 per cent of them had Diploma, 9.43 per cent attained graduation and 3.14 them had other education.

Table 4. Education level of members of the household in Chandaraki-1 micro-watershed

Sl. No.	Particulars	LL (5)		MF (64)		SF (30)		SMF (48)		MDF (12)		All (159)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Illiterate	3	60	32	50	12	40	20	41.7	4	33.33	71	44.7
2	Primary School	2	40	14	21.9	4	13.3	6	12.5	2	16.67	28	17.6
3	High School	0	0	7	10.9	3	10	10	20.8	2	16.67	22	13.8
4	PUC	0	0	5	7.81	2	6.67	4	8.33	2	16.67	13	8.18
5	Diploma	0	0	1	1.56	0	0	0	0	0	0	1	0.63
6	ITI	0	0	0	0	2	6.67	2	4.17	0	0	4	2.52
7	Degree	0	0	3	4.69	5	16.7	5	10.4	2	16.67	15	9.43
8	Others	0	0	2	3.13	2	6.67	1	2.08	0	0	5	3.14
Total		5	100	64	100	30	100	48	100	12	100	159	100

Occupation of head of households: The data regarding the occupation of the household heads in Chandaraki-1 Micro watershed is presented in Table 5. The results indicate that, 80.56 per cent of households heads were practicing agriculture and 2.78 per cent of the household heads were agricultural Labour.

Table 5: Occupation of heads of households in Chandaraki-1 micro-watershed

Sl.No.	Particulars	LL (2)		MF (14)		SF (7)		SMF (10)		MDF (3)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Agriculture	2	100	12	86	6	85.71	8	80	1	33.3	29	80.56
2	Agricultural Labour	0	0	1	7.1	0	0	0	0	0	0	1	2.78
3	Government Service	0	0	0	0	0	0	1	10	0	0	1	2.78
4	Private Service	0	0	1	7.1	1	14.29	0	0	1	33.3	3	8.33
Total		2	100	14	100	7	100	9	100	2	100	34	100

Table 6: Occupation of members of the household in Chandaraki-1 micro-watershed

Sl.No.	Particulars	LL (5)		MF (64)		SF (30)		SMF (48)		MDF (12)		All (159)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Agriculture	2	100	12	85.7	6	85.71	8	80	1	33	29	80.6
2	Agricultural Labour	0	0	1	7.14	0	0	0	0	0	0	1	2.78
3	Government Service	0	0	0	0	0	0	1	10	0	0	1	2.78
4	Private Service	0	0	1	7.14	1	14.29	0	0	1	33	3	8.33
Total		2	100	14	100	7	100	9	100	2	100	34	100

Occupation of the members of the household: The data regarding the occupation of the household members in Chandaraki-1 Micro watershed is presented in Table 6. The results indicate that, agriculture was the major occupation for 80.56 per cent of

the household members, 2.78 per cent were agricultural labour and 2.78 per cent were working in government sector.

Institutional Participation of household members: The data regarding the institutional participation of the household members in Chandaraki-1 Micro watershed is presented in Table 7. The results show that, out of the total family members in the households 0.63 per cent of them are participating in NGOs.

Table 7: Institutional Participation of household member in Chandaraki-1 micro-watershed

Sl.No.	Particulars	LL (5)		MF (64)		SF (30)		SMF (48)		MDF (12)		All (159)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	NGOs	0	0	0	0	1	3.33	0	0	0	0	1	0.63
2	No Participation	5	100	64	100	29	96.7	48	100	12	100	158	99.4
Total		5	100	64	100	30	100	48	100	12	100	159	100

Type of house owned: The data regarding the type of house owned by the households in Chandaraki-1 Micro watershed is presented in Table 8. The results indicate that, 19.44 percent possess thatched house, 72.22 per cent of the households possess katcha house and 8.33 per cent possess pacca house.

Table 8. Type of house owned by households in Chandaraki-1 micro-watershed

Sl.No.	Particulars	LL (5)		MF (64)		SF (30)		SMF (48)		MDF (12)		All (159)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Thatched	1	50	3	21	0	0	3	30	0	0	7	19.44
2	Katcha	1	50	11	79	6	85.71	6	60	2	67	26	72.22
3	Pucca/RCC	0	0	0	0	1	14.29	1	10	1	33	3	8.33
Total		2	100	14	100	7	100	10	100	3	100	36	100

Durable assets owned by the households: The data regarding the Durable Assets owned by the households in Chandaraki-1 Micro watershed is presented in Table 9. The results show that, 72.22 per cent possess TV, 80.56 per cent possess mixer grinder, 30.56 per cent possess motor cycle and 94.44 per cent possess mobile phones.

Table 9. Durable assets owned by households in Chandaraki-1 micro-watershed

Sl.No.	Particulars	LL (2)		MF (14)		SF (7)		SMF (10)		MDF (3)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Television	1	50	11	79	4	57.1	8	80	2	66.7	26	72.22
2	Mixer/Grinder	0	0	11	79	7	100	8	80	3	100	29	80.56
3	Motor Cycle	0	0	4	29	4	57.1	1	10	2	66.7	11	30.56
4	Mobile Phone	2	100	12	86	7	100	10	100	3	100	34	94.44

Table 10. Average value of durable assets owned in Chandaraki-1 micro-watershed
Average Value (Rs.)

Sl.No.	Particulars	LL (2)	MF (14)	SF (7)	SMF (10)	MDF (3)	All (36)
1	Television	5,000.00	4,727.00	6,000.00	4,875.00	6,000.00	5,076.00
2	Mixer/Grinder	0.00	1,845.00	1,771.00	1,430.00	1,533.00	1,664.00
3	Motor Cycle	0.00	47,500.00	55,000.00	60,000.00	50,000.00	51,818.00
4	Mobile Phone	2,250.00	2,062.00	1,692.00	2,352.00	2,666.00	2,134.00

Average value of durable assets: The data regarding the average value of durable assets owned by the households in Chandaraki-1 Micro watershed is presented in Table 10. The result shows that, the average value of television was Rs. 5,076.00, Mixer/Grinder was Rs. 1,664.00, motor cycle was Rs. 51,818.00 and Rs.2,134.00.

Farm implements owned: The data regarding the farm implements owned by the households in Chandaraki-1 Micro watershed is presented in Table 11. About 8.33 per cent of the households possess Bullock Cart, 36.11 per cent possess plough and 77.78 per cent possess Weeder, 8.33 per cent possess tractor, 5.56 per cent possess Harvester and 36.11 per cent possess Thresher.

Table 11. Farm implements owned in Chandaraki-1 micro-watershed

Sl.No.	Particulars	LL (2)		MF (14)		SF (7)		SMF (10)		MDF (3)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Bullock Cart	0	0	2	14.3	0	0	1	10	0	0	0	8.33
2	Plough	0	0	6	42.9	2	28.57	3	30	2	66.7	0	36.11
3	Tractor	0	0	1	7.14	1	14.29	0	0	1	33.3	0	8.33
4	Weeder	0	0	13	92.9	6	85.71	7	70	2	66.7	0	77.78
5	Harvester	0	0	0	0	1	14.29	1	10	0	0	0	5.56
6	Thresher	0	0	4	28.6	4	57.14	3	30	2	66.7	0	36.11
7	Blank	2	100	1	7.14	0	0	2	20	0	0	0	13.89

Average value of farm implements: The data regarding the average value of farm Implements owned by the households in Chandaraki-1 Micro watershed is presented in Table 12. The results show that the average value of plough was Rs.1664.00, bullock Cart was Rs.13933.00, weeder was Rs.135.00, Harvester was Rs. 4333, tractor was Rs. 800000 and Thresher Rs. 496.

Table 12. Average value of farm implements in Chandaraki-1 micro-watershed

Sl.No.	Particulars	Average Value (Rs.)					
		LL (2)	MF (14)	SF (7)	SMF (10)	MDF(3)	All (36)
1	Bullock Cart	0	20000	0	1800	0	13933
2	Plough	0	1685	2000	1833	1000	1664
3	Tractor	0	800000	800000	0	800000	800000
4	Weeder	0	124	172	125	175	135
5	Harvester	0	0	10000	1500	0	4333
6	Thresher	0	393	600	436	666	496

Table 13. Livestock possession by households in Chandaraki-1 micro-watershed

Sl.No.	Particulars	LL (2)		MF (14)		SF (7)		SMF (10)		MDF (3)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Bullock	0	0	2	14	2	28.57	1	10	1	33.3	6	16.67
2	Local cow	0	0	1	7.1	0	0	0	0	0	0	1	2.78
3	Crossbred cow	0	0	0	0	0	0	1	10	0	0	1	2.78
4	Buffalo	0	0	0	0	1	14.29	1	10	0	0	2	5.56
5	Poultry birds	0	0	0	0	0	0	2	20	0	0	2	5.56
6	blank	2	100	11	79	5	71.43	5	50	2	66.7	25	69.44

Livestock possession by the households: The data regarding the Livestock possession by the households in Chandaraki-1 Micro watershed is presented in Table 13. The indicate that, 16.67 per cent of the households possess bullocks, 2.78 per cent possess local cow, 5.56 per cent possess buffalo, 2.78 per cent possess crossbred cow and 5.56 per cent were poultry birds.

Average Labour availability: The data regarding the average labour availability in Chandaraki-1 Micro watershed is presented in Table 14. The indicated that, own labour men available in the micro watershed was 1.22, women available in the micro watershed was 1.17, hired labour (men) available was 15.03 and hired labour (women) available was 12.89.

Table 14. Average labour availability in Chandaraki-1 micro-watershed

Sl.No.	Particulars	LL (2)	MF (14)	SF (7)	SMF (10)	MDF (3)	All (36)
		N	N	N	N	N	N
1	Hired labour Female	3	6.43	16.43	16.8	28.3	12.89
2	Own Labour Female	1	1.07	1	1.5	1	1.17
3	Own labour Male	1	1.21	1	1.5	1	1.22
4	Hired labour Male	3	8	16.43	22.3	28.3	15.03

Adequacy of hired labour: The data regarding the adequacy of hired labour in Chandaraki-1 Micro watershed is presented in Table 15. The results indicate that, 94.44 per cent of the household opined that hired labour was adequate and 5.56 per cent of the household opined that hired labour was Inadequate.

Table 15. Adequacy of hired labour in Chandaraki-1 micro-watershed

Sl.No.	Particulars	LL (2)		MF (14)		SF (7)		SMF (10)		MDF (3)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Adequate	0	0	14	100	7	100	10	100	3	100	34	94.4
2	Inadequate	2	100	0	0	0	0	0	0	0	0	2	5.56

Migration among the households: The data regarding the migration (Table 16) indicate that, 0.63 percent of the population was being migrated from the micro watershed.

Table 16. Migration among the households in Chandaraki-1 micro-watershed

Sl.No.	Particulars	LL (5)		MF (64)		SF (30)		SMF (48)		MDF (12)		All (159)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Migration	0	0.00	1	1.56	0	0.00	0	0.00	0	0.00	1	0.63

Table 17. Average distance and duration of migration in Chandaraki-1 micro-watershed

Sl.No.	Particulars	LL (0)	MF (1)	SF (0)	SMF (0)	MDF (0)	All (1)
		N	N	N	N	N	N
1	Avg. Distance (kms)	0	850	0	0	0	850
2	Avg. Duration (months)	0	4	0	0	0	4

Average distance and duration of migration: The data regarding the average distance and duration of migration (Table 17) indicate that, people migrated to a distance of 850 kms on an average for 4 months.

Purpose of migration: The data regarding the purpose of migration (Table 18) indicate that, 100.00 percent of them went for the purpose of job/wage/work.

Table 18. Purpose of migration by members of households in Chandaraki-1 micro-watershed

Sl. No.	Particulars	LL (0)		MF (1)		SF (0)		SMF (0)		MDF (0)		All (1)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Job/wage/work	0	0	1	100	0	0	0	0	0	0	1	100
Total		0	100	1	100	0	100	0	100	0	100	1	100

Distribution of land (ha): The data regarding the distribution of land (ha) in Chandaraki-1 Micro watershed is presented in Table 19. The results indicate that, 56.43 ha (94.31%) of dry land and 2.54 ha (4.24 %) of irrigated land.

Table 19. Distribution of land (ha) in Chandaraki-1 micro-watershed

Sl.No.	Particulars	LL (2)		MF (14)		SF (7)		SMF (10)		MDF (3)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Dry	0	0	8.19	90.39	10.3	100	26.94	100	11	81.3	56.43	94.31
2	Irrigated	0	0	0	0	0	0	0	0	2.54	18.8	2.54	4.24
3	Permanent Fallow	0	0	0.87	9.61	0	0	0	0	0	0	0.87	1.45
Total		0	100	9.06	100	10.3	100	26.94	100	13.53	100	59.84	100

Average value of land (ha): The data regarding the average land value (Rs./ha) in Chandaraki-1 Micro watershed is presented in Table 20. The results show that the average value of dry land was Rs.313532.70, and the average value of irrigated land was Rs.157575.76.

Table 20. Average value of land (ha) in Chandaraki-1 micro-watershed

Sl.No.	Particulars	LL (2)	MF (14)	SF (7)	SMF (10)	MDF (3)	All (36)
		N	N	N	N	N	N
1	Dry	0	634898.7	281343.3	322754.6	81818.18	313532.7
2	Irrigated	0	0	0	0	157575.8	157575.8
3	Permanent Fallow	0	459534.9	0	0	0	459534.9

Status of open wells: The data regarding the status of open wells in Chandaraki-1 Micro watershed is presented in Table 21. The results indicate that, there were 2 De-functioning open wells and 2 functioning open wells among the sampled households in micro watershed.

Table 21. Status of open wells in Chandaraki-1 micro-watershed

Sl. No.	Particulars	LL (2)	MF (14)	SF (7)	SMF (10)	MDF (3)	All (36)
		N	N	N	N	N	N
1	De-functioning	0	0	0	2	0	2
2	Functioning	0	0	0	2	0	2

Source of irrigation: The data regarding the source of irrigation in Chandaraki-1 Micro watershed is presented in Table 22. The results that open well were major source of irrigation for 5.56 per cent of the households.

Table 22. Source of irrigation in Chandaraki-1 micro-watershed

Sl.No.	Particulars	LL (2)		MF (14)		SF (7)		SMF (10)		MDF (3)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Open Well	0	0	0	0	0	0	2	20	0	0	2	5.56

Depth of water (Avg. In meters): The data regarding the depth of water in Chandaraki-1 Micro watershed is presented in Table 23. The results revealed that, the depth of open well was 1.69 meter.

Table 23. Depth of water (Avg. In meters) in Chandaraki-1 micro-watershed

Sl.No.	Particulars	LL (2)	MF (14)	SF (7)	SMF (10)	MDF (3)	All (36)
		N	N	N	N	N	N
1	Open Well	0	0	0	6.1	0	1.69

Cropping pattern: The data regarding the cropping pattern in Chandaraki-1 Micro watershed is presented in Table 24. The results indicate that, farmers have grown Red gram (48.33 ha), Cotton (4.81 ha), Jowar (3.46 ha), Paddy (1.21 ha) and Bangalore brinjal (0.81 ha)

Table 24. Cropping pattern in Chandaraki-1 micro-watershed

Sl.No.	Particulars	LL (2)	MF (14)	SF (7)	SMF (10)	MDF (3)	All (36)
1	Kharif - Red gram	0	7.64	9.03	21.94	9.72	48.33
2	Kharif - Cotton	0	0	2.25	2.55	0	4.81
3	Kharif - Jowar	0	0.55	0	0.44	2.47	3.46
4	Kharif - Paddy	0	0	0	0	1.21	1.21
5	Kharif - brinjal	0	0	0	0.81	0	0.81
Total		0	8.19	11.28	25.74	13.4	58.62

Cropping intensity: The data regarding the cropping intensity in Chandaraki-1 Micro watershed is presented in Table 25. The results indicate that, the cropping intensity was 100.00 per cent.

Table 25. Cropping intensity (%) in Chandaraki-1 micro-watershed

Sl.No.	Particulars	LL (2)	MF (14)	SF (7)	SMF (10)	MDF (3)	All (36)
1	Cropping Intensity	0	100	100	100	100	100

Possession of bank account and savings: The data regarding the possession of bank account and saving in Chandaraki-1 micro-watershed is presented in Table 26. The results indicate that, 94.44 cent of the households posses bank account.

Table 26. Possession of Bank account and savings in Chandaraki-1 micro-watershed

Sl. No.	Particulars	LL (2)		MF (14)		SF (7)		SMF (10)		MDF (3)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Account	0	0	14	100	7	100	10	100	3	100	34	94.44

Borrowing status: The data regarding the borrowing status in Chandaraki-1 micro-watershed is presented in Table 27. The results indicate that, 94.44 percent of the sample farmers have borrowed credit from different sources.

Table 27. Borrowing status in Chandaraki-1 micro-watershed

Sl.No.	Particulars	LL (2)		MF (14)		SF (7)		SMF (10)		MDF (3)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Credit Availed	0	0	14	100	7	100	10	100	3	100	34	94.44

Source of credit: The data regarding the source of credit availed by households in Chandaraki-1 micro-watershed is presented in Table 28. The results show that, 13.64 per cent have borrowed loan from commercial banks, 4.55 per cent have borrowed loan from Friends/Relatives and 86.36 per cent have borrowed loan from Grameena Bank.

Table 28. Source of credit borrowed by households in Chandaraki-1 micro-watershed

Sl.No.	Particulars	LL (0)		MF (9)		SF (3)		SMF (7)		MDF (3)		All (22)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Commercial Bank	0	0	0	0	0	0	2	29	1	33.33	3	13.64
2	Friends/Relatives	0	0	0	0	0	0	1	14	0	0	1	4.545
3	Grameena Bank	0	0	9	100	3	100	5	71	2	66.67	19	86.36

Avg. Credit amount: The data regarding the avg. Credit amount in Chandaraki-1 micro-watershed is presented in Table 29. The results show that, farmers have borrowed Avg. Credit of Rs.35611.11 from different sources.

Table 29. Avg. Credit amount in Chandaraki-1 micro-watershed

Sl.No.	Particulars	LL (0)		MF (9)		SF (3)		SMF (7)		MDF (3)		All (8)	
		N	N	N	N	N	N	N	N	N	N		
1	Average Credit	0	80000	51666.7	40000	42000	35611.1						

Purpose of credit borrowed (institutional Source): The data regarding the purpose of credit borrowed - Institutional Credit in Chandaraki-1 micro-watershed is presented in Table 30. The results indicate that, 95.45 per cent of the households have borrowed loan for agriculture and animal husbandry (4.55%).

Table 30. Purpose of credit borrowed (institutional Source) by households in Chandaraki-1 micro-watershed

SN	Particulars	LL (0)		MF (9)		SF (3)		SMF (7)		MDF (3)		All (22)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Agriculture production	0	0	8	88.9	3	100	7	100	3	100	21	95.5
2	Animal husbandry	0	0	1	11.1	0	0	0	0	0	0	1	4.55

Purpose of credit borrowed (Private Source): The data regarding the purpose of credit borrowed – Private Source in Chandaraki-1 micro-watershed is presented in Table 31. The results indicate that, 100.00 per cent of the households have borrowed loan for agriculture.

Table 31. Purpose of credit borrowed (Private Source) by households in Chandaraki-1 micro-watershed

Sl.No.	Particulars	LL (0)		MF (0)		SF (0)		SMF (1)		MDF (0)		All (1)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Agriculture production	0	0	0	0	0	0	1	100	0	0	1	100

Repayment status of household (institutional Source): The data regarding the repayment status of credit borrowed from institutional Source by households in Chandaraki-1 micro watershed is presented in Table 32. The results indicate that, 100.00 per cent have unpaid.

Table 32. Repayment status of household (institutional Source) in Chandaraki-1 micro-watershed

Sl.No.	Particulars	LL (0)		MF (9)		SF (3)		SMF (7)		MDF (3)		All (22)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Un paid	0	0	9	100	3	100	7	100	3	100	22	100

Repayment status of household (Private Source): The data regarding the repayment status of credit borrowed from private sources by households in Chandaraki-1 micro watershed is presented in Table 33. The results indicate that, 100 per cent of the households has unpaid.

Table 33. Repayment status of household (Private Source) in Chandaraki-1 micro-watershed

Sl.No.	Particulars	LL (0)		MF (0)		SF (0)		SMF (1)		MDF (0)		All (1)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Un paid	0	0	0	0	0	0	1	100	0	0	1	100

Cost of Cultivation of Red gram: The data regarding the cost of cultivation (Rs/ha) of Red gram in Chandaraki-1 micro watershed is presented in Table 34.a. The results indicate that, the total cost of cultivation (Rs/ha) for Red gram was Rs. 48868.88. The gross income realized by the farmers was Rs. 42065.03. The net income from Red gram cultivation was Rs.-6803.84, thus the benefit cost ratio was found to be 1:0.90.

Table 34(a). Cost of Cultivation of Red gram in Chandaraki-1 micro-watershed

Sl.No	Particulars	Units	Phy Units	Value (Rs.)	% to C3
I	Cost A1				
1	Hired Human Labour	Man days	44.57	7782.37	15.92
2	Bullock	Pairs/day	2.55	1703.95	3.49
3	Tractor	Hours	4.03	3023.48	6.19
4	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	17.36	1437.21	2.94
5	FYM	Quintal	4.8	11407.37	23.34
6	Fertilizer + micronutrients	Quintal	8.88	7494.48	15.34
7	Pesticides (PPC)	Kgs / liters	1.8	1482.19	3.03
8	Irrigation	Number	0.82	0	0
9	Depreciation charges		0	1341.64	2.75
II	Cost B1				
10	Interest on working capital			2619.75	5.36
11	Cost B1 = (Cost A1 + sum of 15 and 16)			38292.42	78.36
III	Cost B2				
12	Rental Value of Land			217.17	0.44
13	Cost B2 = (Cost B1 + Rental value)			38509.6	78.8
IV	Cost C1				
14	Family Human Labour		25.22	5906.65	12.09
15	Cost C1 = (Cost B2 + Family Labour)			44416.25	90.89
V	Cost C2				
16	Risk Premium			10	0.02
17	Cost C2 = (Cost C1 + Risk Premium)			44426.25	90.91
VI	Cost C3				
18	Managerial Cost			4442.63	9.09
19	Cost C3 = (Cost C2 + Managerial Cost)			48868.88	100
VII	Economics of the Crop				
a.	Main Product	a) Main Product (q)		7.2	42065.03
		b) Main Crop Sales Price (Rs.)			5838.71
b.	Gross Income (Rs.)			42065.03	
c.	Net Income (Rs.)			-6803.84	
d.	Cost per Quintal (Rs./q.)			6783.1	
e.	Benefit Cost Ratio (BC Ratio)			1:0.9	

Cost of Cultivation of Cotton: The data regarding the cost of cultivation (Rs/ha) of Cotton in Chandaraki-1 micro watershed is presented in Table 34.b. The results indicate that, the total cost of cultivation (Rs/ha) for Cotton was Rs. 44389.25. The gross income realized by the farmers was Rs. 44332.27. The net income from Cotton cultivation was Rs.-56.97, thus the benefit cost ratio was found to be 1:1.00.

Table 34(b). Cost of Cultivation of Cotton in Chandaraki-1 micro-watershed

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
I	Cost A1				
1	Hired Human Labour	Man days	39.13	6671.79	15.03
2	Bullock	Pairs/day	2.91	1837.34	4.14
3	Tractor	Hours	1.89	1496.73	3.37
4	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	4.95	7811.81	17.6
5	FYM	Quintal	4.94	5928	13.35
6	Fertilizer + micronutrients	Quintal	6.38	5590.78	12.59
7	Pesticides (PPC)	Kgs / liters	2.07	1255.54	2.83
8	Depreciation charges		0	1907.4	4.3
II	Cost B1				
9	Interest on working capital			2471.54	5.57
10	Cost B1 = (Cost A1 + sum of 15 and 16)			34970.93	78.78
III	Cost B2				
11	Rental Value of Land			166.67	0.38
12	Cost B2 = (Cost B1 + Rental value)			35137.59	79.16
IV	Cost C1				
13	Family Human Labour		23.77	5206.27	11.73
14	Cost C1 = (Cost B2 + Family Labour)			40343.86	90.89
V	Cost C2				
15	Risk Premium			10	0.02
16	Cost C2 = (Cost C1 + Risk Premium)			40353.86	90.91
VI	Cost C3				
17	Managerial Cost			4035.39	9.09
18	Cost C3 = (Cost C2 + Managerial Cost)			44389.25	100
VII	Economics of the Crop				
a.	Main Product	a) Main Product (q)	8.53	44332.27	
		b) Main Crop Sales Price (Rs.)		5200	
b.	Gross Income (Rs.)			44332.27	
c.	Net Income (Rs.)			-56.97	
d.	Cost per Quintal (Rs./q.)			5206.68	
e.	Benefit Cost Ratio (BC Ratio)			1:1	

Cost of Cultivation of Bengal gram: The data regarding the cost of cultivation (Rs/ha) of Bengal gram in Chandaraki-1 micro watershed is presented in Table 34.c. The results indicate, the total cost of cultivation (Rs/ha) for Bengal gram was Rs.18089.82. The gross income realized by the farmers was Rs. 69160.00. The net income from Bengal gram cultivation was Rs. 51070.18, thus the benefit cost ratio was found to be 1:3.80.

Table 34(c). Cost of Cultivation of Bengal gram in Chandaraki-1 micro-watershed

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
I	Cost A1				
1	Hired Human Labour	Man days	29.64	5001.75	27.65
2	Bullock	Pairs/day	2.47	1482	8.19
3	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	12.35	926.25	5.12
4	Fertilizer + micronutrients	Quintal	3.71	3211	17.75
5	Pesticides (PPC)	Kgs / liters	1.24	617.5	3.41
6	Depreciation charges		0	12.35	0.07
II	Cost B1				
7	Interest on working capital			571.77	3.16
8	Cost B1 = (Cost A1 + sum of 15 and 16)			11822.62	65.36
III	Cost B2				
9	Rental Value of Land			166.67	0.92
10	Cost B2 = (Cost B1 + Rental value)			11989.29	66.28
IV	Cost C1				
11	Family Human Labour		19.76	4446	24.58
12	Cost C1 = (Cost B2 + Family Labour)			16435.29	90.85
V	Cost C2				
13	Risk Premium			10	0.06
14	Cost C2 = (Cost C1 + Risk Premium)			16445.29	90.91
VI	Cost C3				
15	Managerial Cost			1644.53	9.09
16	Cost C3 = (Cost C2 + Managerial Cost)			18089.82	100
VII	Economics of the Crop				
a.	Main Product	a) Main Product (q)		8.65	69160
		b) Main Crop Sales Price (Rs.)			8000
b.	Gross Income (Rs.)			69160	
c.	Net Income (Rs.)			51070.18	
d.	Cost per Quintal (Rs./q.)			2092.52	
e.	Benefit Cost Ratio (BC Ratio)			1:3.8	

Cost of Cultivation of Jowar: The data regarding the cost of cultivation (Rs/ha) of Jowar in Chandaraki-1 micro watershed is presented in Table 34.d. The results indicate that, the total cost of cultivation (Rs/ha) for Jowar was Rs. 26988.16. The gross income realized by the farmers was Rs.14670.55. The net income from Jowar cultivation was Rs. -12317.61, thus the benefit cost ratio was found to be 1:0.50.

Table 34(d). Cost of Cultivation of Jowar in Chandaraki-1 micro-watershed

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
I	Cost A1				
1	Hired Human Labour	Man days	46.85	8101.11	30.02
2	Bullock	Pairs/day	1.48	915.12	3.39
3	Tractor	Hours	1.44	1148.88	4.26
4	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	13.37	302.92	1.12
5	Fertilizer + micronutrients	Quintal	6.17	5348.24	19.82
6	Pesticides (PPC)	Kgs / liters	2.11	1592.2	5.9
7	Depreciation charges		0	2183.89	8.09
II	Cost B1				
8	Interest on working capital			870.4	3.23
9	Cost B1 = (Cost A1 + sum of 15 and 16)			20462.77	75.82
III	Cost B2				
10	Rental Value of Land			166.67	0.62
11	Cost B2 = (Cost B1 + Rental value)			20629.44	76.44
IV	Cost C1				
12	Family Human Labour		15.61	3895.25	14.43
13	Cost C1 = (Cost B2 + Family Labour)			24524.69	90.87
V	Cost C2				
14	Risk Premium			10	0.04
15	Cost C2 = (Cost C1 + Risk Premium)			24534.69	90.91
VI	Cost C3				
16	Managerial Cost			2453.47	9.09
17	Cost C3 = (Cost C2 + Managerial Cost)			26988.16	100
VII	Economics of the Crop				
a.	Main Product	a) Main Product (q)	7.34	14670.55	
		b) Main Crop Sales Price (Rs.)		2000	
b.	Gross Income (Rs.)			14670.55	
c.	Net Income (Rs.)			-12317.61	
d.	Cost per Quintal (Rs./q.)			3679.23	
e.	Benefit Cost Ratio (BC Ratio)			1:0.5	

Cost of Cultivation of Paddy: The data regarding the cost of cultivation (Rs/ha) of Paddy in Chandaraki-1 micro watershed is presented in Table 34.e. The results indicate that, the total cost of cultivation (Rs/ha) for Paddy was Rs.23946.94. The gross income realized by the farmers was Rs. 17784.00. The net income from Paddy cultivation was Rs. -6162.94, thus the benefit cost ratio was found to be 1:0.70.

Table 34(e). Cost of Cultivation of Paddy in Chandaraki-1 micro-watershed

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
I	Cost A1				
1	Hired Human Labour	Man days	33.76	6298.5	26.3
2	Bullock	Pairs/day	1.65	1152.67	4.81
3	Tractor	Hours	0	0	0
4	Machinery	Hours	0	0	0
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	4.94	172.9	0.72
6	Seed Inter Crop	Kgs.	0	0	0
7	FYM	Quintal	1.65	1976	8.25
8	Fertilizer + micronutrients	Quintal	7.41	6422	26.82
9	Pesticides (PPC)	Kgs / liters	1.65	823.33	3.44
10	Irrigation	Number	0.82	0	0
11	Repairs		0	0	0
12	Msc. Charges (Marketing costs etc)		0	0	0
13	Depreciation charges		0	37.87	0.16
14	Land revenue and Taxes		0	0	0
II	Cost B1				
16	Interest on working capital			1128.51	4.71
17	Cost B1 = (Cost A1 + sum of 15 and 16)			18011.78	75.22
III	Cost B2				
18	Rental Value of Land			166.67	0.7
19	Cost B2 = (Cost B1 + Rental value)			18178.45	75.91
IV	Cost C1				
20	Family Human Labour		15.64	3581.5	14.96
21	Cost C1 = (Cost B2 + Family Labour)			21759.95	90.87
V	Cost C2				
22	Risk Premium			10	0.04
23	Cost C2 = (Cost C1 + Risk Premium)			21769.95	90.91
VI	Cost C3				
24	Managerial Cost			2176.99	9.09
25	Cost C3 = (Cost C2 + Managerial Cost)			23946.94	100
VII	Economics of the Crop				
a.	Main Product	a) Main Product (q)		9.88	17784
		b) Main Crop Sales Price (Rs.)			1800
b.	Gross Income (Rs.)			17784	
c.	Net Income (Rs.)			-6162.94	
d.	Cost per Quintal (Rs./q.)			2423.78	
e.	Benefit Cost Ratio (BC Ratio)			1:0.7	

Adequacy of fodder: The data regarding the adequacy of fodder in Chandaraki-1 Micro watershed is presented in Table 35. The results indicate that, 25.00 per cent of the households opined that dry fodder was adequate and 8.33 per cent of them opined green fodder was sufficient.

Table 35. Adequacy of fodder in Chandaraki-1 micro-watershed

Sl.No.	Particulars	LL (2)		MF (14)		SF (7)		SMF (10)		MDF (3)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Adequate-Dry Fodder	0	0	3	21.43	2	28.57	3	30	1	33.3	9	25
2	Adequate-Green Fodder	0	0	1	7.14	0	0	2	20	0	0	3	8.33

Average annual gross income: The data regarding the annual gross income in Chandaraki-1 Micro watershed is presented in Table 36. The results indicate that, the farmers have annual gross income of Rs. 76055.56 in micro-watershed, of which Rs. 41027.78 is from agriculture itself.

Table 36. Average annual gross income in Chandaraki-1 micro-watershed

Sl.No.	Particulars	LL (2)	MF (14)	SF (7)	SMF (10)	MDF (3)	All (36)
		Rs.	Rs.	Rs.	Rs.	Rs.	Rs.
1	Service/salary	0	0	0	40000	0	11111.1
2	Wage	40500	26071.4	25000	14800	26666.7	23583.3
3	Agriculture	0	23500	47857.1	55300	86666.7	41027.8
4	Dairy Farm	0	0	1428.57	200	0	333.33
Income(Rs.)		40500	49571.4	74285.7	110300	113333	76055.6

Average annual Expenditure: The data regarding the average annual expenditure in Chandaraki-1 Micro watershed is presented in Table 37. The results indicate that, the farmers have annual gross expenditure of Rs. 495400.18 in micro-watershed, of which Rs. 26638.89 is from agriculture itself.

Table 37. Average annual Expenditure in Chandaraki-1 micro-watershed

Sl.No.	Particulars	LL (2)	MF (14)	SF (7)	SMF (10)	MDF (3)	All (36)
		Rs.	Rs.	Rs.	Rs.	Rs.	Rs.
1	Service/salary	0	0	0	300000	0	8333.33
2	Wage	17500	11384.6	11333.3	6750	2000	8583.33
3	Agriculture	0	15384.6	27714.3	37500	63333.3	26638.9
4	Dairy Farm	0	0	2000	500	0	69.44
Total		17500	26769.2	41047.6	344750	65333.3	495400

Table 38. Horticulture species grown in Chandaraki-1 micro-watershed

Sl.No.	Particulars	LL (2)		MF (14)		SF (7)		SMF (10)		MDF (3)		All (36)	
		F	B	F	B	F	B	F	B	F	B	F	B
1	Mango	0	0	0	0	8	0	0	0	0	0	8	0

*F= Field B=Back Yard

Horticulture species grown: The data regarding horticulture species grown in Chandaraki-1 Micro watershed is presented in Table 38. The results indicate that, the

total number of horticultural trees grown (both field and backyard) by the sampled households were Mango (8).

Forest species grown: The data regarding forest species grown in Chandaraki-1 Micro watershed is presented in Table 39. The results indicate that, households have planted 10 teak trees, 68 neem trees, 11 pongamia trees, 10 acacia trees and 1 banyan trees together in both field and backyard.

Table 39. Forest species grown in Chandaraki-1 micro-watershed

Sl.No.	Particulars	LL (2)		MF (14)		SF (7)		SMF (10)		MDF (3)		All (36)	
		F	B	F	B	F	B	F	B	F	B	F	B
1	Teak	0	0	4	0	3	3	0	0	0	0	7	3
2	Neem	0	0	15	7	11	3	22	2	8	0	56	12
3	Pongamia	0	0	0	0	10	0	1	0	0	0	11	0
4	Acacia	0	0	3	1	2	0	0	0	4	0	9	1
5	Banyan	0	0	0	0	0	0	0	0	1	0	1	0

*F= Field B=Back Yard

Average additional investment capacity: The data regarding average additional investment capacity in Chandaraki-1 Micro watershed is presented in Table 40. The results indicate that, households have an average investment capacity of Rs. 12361.11 for land development, Rs.2750.00 for adoption of improved crop production and Rs.277.78 adoption of improved livestock management.

Table 40. Average additional investment capacity of households in Chandaraki-1 micro-watershed

Sl. No.	Particulars	LL (2)	MF (14)	SF (7)	SMF (10)	MDF (3)	All (36)
		Rs.	Rs.	Rs.	Rs.	Rs.	Rs.
1	Land development	0	8928.57	15285.7	15100	20666.7	12361.1
2	Improved crop production	0	2928.57	1857.14	3000	5000	2750
3	Improved livestock management	0	357.14	0	500	0	277.78

Source of funds for additional investment: The data regarding source of funds for additional investment in Chandaraki-1 Micro watershed is presented in Table 41. The results indicate that, the sources of finance raised from bank as a loan and from own sources for land development was 2.78 and 88.89 per cent.

Table 41. Source of funds for additional investment in Chandaraki-1 micro-watershed

Sl.No	Item	Land development		Improved crop production		Improved livestock management	
		N	%	N	%	N	%
1	Bank Funds	1	2.78	0	0	0	0
2	Own Funds	32	88.89	20	55.56	2	5.56

Marketing of agricultural produce: The data regarding marketing of the agricultural produce in Chandaraki-1 Micro watershed is presented in Table 42. The results indicated that, 85.71 per cent of output of Bengal gram was sold in the market

with average price of Rs. 8000.00; 97.44 percent of output of Cotton was sold in the market with average price of Rs. 5200.00; 78.26 percent of output of Jowar was sold in the market with average price of Rs. 2000.00; 75.00 percent of output of Paddy was sold in the market with average price of Rs. 1800.00 and 86.49 percent of output of Red gram was sold in the market with average price of Rs. 5656.25.

Table 42. Marketing of agricultural produce in Chandaraki-1 micro-watershed

Sl. No	Crops	Output obtained (q)	Output retained (q)	Output sold (q)	Output sold (%)	Avg. Price obtained (Rs/q)
1	Bengal gram	7	1	6	86	8000
2	Cotton	39	1	38	97	5200
3	Jowar	23	5	18	78	2000
4	Paddy	12	3	9	75	1800
5	Red gram	259	35	224	86	5656

Marketing channels used for sale of agricultural produce: The data regarding marketing channels used for sale of agricultural produce in Chandaraki-1 Micro watershed is presented in Table 43. The results indicated that, 116.67 cent of the households have sold agricultural produce to the local/village merchants, 0.00 per per cent have sold to Agent/Traders, 0.00 per cent of regulated market, 0.00 per cent of cooperative marketing society, 0.00 per cent of contract marketing arrangement and 0.00 per cent of outside the state.

Table 43. Marketing channels used for sale of agricultural produce in Chandaraki-1 micro-watershed

Sl.No.	Particulars	LL (2)		MF (14)		SF (7)		SMF (10)		MDF (3)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Local/village Merchant	0	0	14	100	7	100	10	100	3	100	36	100

Mode of transport of agricultural produce: The data regarding mode of transport of agricultural produce in Chandaraki-1 Micro watershed is presented in Table 44. The results indicated that, 100 cent of the households have used tractor for the transport of agriculture commodity.

Table 44. Mode of transport of agricultural produce in Chandaraki-1 micro-watershed

Sl.No.	Particulars	LL (2)		MF (14)		SF (7)		SMF (10)		MDF (3)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Tractor	0	0	14	100	7	100	10	100	3	100	36	100

Table 45. Incidence of soil and water erosion problems in Chandaraki-1 micro-watershed

Sl.No.	Particulars	LL (2)		MF (14)		SF (7)		SMF (10)		MDF (3)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Soil and water erosion problems in the farm	0	0	12	86	7	100	10	100	3	100	32	88.89

Incidence of soil and water erosion problems: The data regarding incidence of incidence of soil and water erosion problems in Chandaraki-1 Micro watershed is

presented in Table 45. The results indicate that, 88.89 per cent of the households have experienced soil and water erosion problems.

Interest towards soil testing: The data regarding Interest shown towards soil testing in Chandaraki-1 Micro watershed is presented in Table 46. The results indicated that, 94.44 per cent of the households were interested towards soil testing.

Table 46. Interest regarding soil testing in Chandaraki-1 micro-watershed

Sl.No.	Particulars	LL (2)		MF (14)		SF (7)		SMF (10)		MDF (3)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Interest in soil test	0	0	14	100	7	100	10	100	3	100	34	94.44

Soil and water conservation practices and structures adopted: The data regarding soil and water conservation practices and structures adopted in Chandaraki-1 Micro watershed is presented in Table 47. The results indicated that 2.78 per cent of farmers practicing Farm Pond as soil and water conservation practice.

Table 47. Soil and water conservation practices and structures adopted in Chandaraki-1 micro-watershed

Sl.No.	Particulars	LL (2)		MF (14)		SF (7)		SMF (10)		MDF (3)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Farm Pond	0	0	0	0	1	14	0	0	0	0	1	2.78

Status of soil and water conservation structures: The data regarding status soil and water conservation structures adopted in Chandaraki-1 Micro watershed is presented in Table 48. The results indicated that, the households have adopted Farm Pond as a soil and water conservation structures out of which 100.00 per cent was in good condition.

Table 48. Status of soil and water conservation structures in Chandaraki-1 micro-watershed

Sl.No	Item	Good	
		N	%
1	Farm Pond	1	100

Agencies involved in the soil and water conservation structures: The data regarding Agencies involved in the soil and water conservation structures adopted in Chandaraki-1 Micro watershed is presented in Table 49. The results indicated that, 2.78 per cent were done by Govt.

Table 49. Agencies involved in the soil and water conservation structures in Chandaraki-1 micro-watershed

Sl.No.	Particulars	LL (2)		MF (14)		SF (7)		SMF (10)		MDF (3)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Govt.	0	0	0	0	1	14.29	0	0	0	0	1	2.78

Usage pattern of fuel for domestic use: The data on usage pattern of fuel for domestic use in Chandaraki-1 Micro watershed is presented in Table 50. The results

indicated that, firewood was the major source of fuel for domestic use for 88.89 per cent of the households followed by LPG (11.11%).

Table 50. Usage pattern of fuel for domestic use in Chandaraki-1 micro-watershed

Sl.No.	Particulars	LL (2)		MF (14)		SF (7)		SMF (10)		MDF (3)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Fire Wood	2	100	13	92.9	6	85.7	9	90	2	66.7	32	88.89
2	LPG	0	0	1	7.14	1	14.3	1	10	1	33.3	4	11.11

Source of drinking water: The data on source of drinking water in Chandaraki-1 Micro watershed is presented in Table 51. The results indicated that, piped waters supply was the major source for drinking water for 97.22 per cent of the households followed by bore well water (2.78%).

Table 51. Source of drinking water in Chandaraki-1 micro-watershed

Sl.No.	Particulars	LL (2)		MF (14)		SF (7)		SMF (10)		MDF (3)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Piped supply	2	100	13	92.9	7	100	10	100	3	100	35	97.22
2	Bore Well	0	0	1	7.14	0	0	0	0	0	0	1	2.78

Source of light: The data on source of light in Chandaraki-1 Micro watershed is presented in Table 52. The results indicated that, electricity was the major source of light for 100.00 per cent of the households.

Table 52. Source of light in Chandaraki-1 micro-watershed

Sl.No.	Particulars	LL (2)		MF (14)		SF (7)		SMF (10)		MDF (3)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Electricity	2	100	14	100	7	100	10	100	3	100	36	100

Existence of sanitary toilet facility: The data on availability of toilet facility in Chandaraki-1 Micro watershed is presented in Table 53. The results indicated that, 30.56 per cent of the households possess toilets.

Table 53. Existence of sanitary toilet facility in Chandaraki-1 micro-watershed

Sl.No.	Particulars	LL (2)		MF (14)		SF (7)		SMF (10)		MDF (3)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Sanitary toilet facility	1	50	6	43	2	28.57	1	10	1	33.3	11	30.6

Possession of PDS card: The data regarding possession of PDS card in Chandaraki-1 Micro watershed is presented in Table 54. The results indicated that, 100.00 per cent of the households possessed BPL card.

Table 54. Possession of PDS card in Chandaraki-1 micro-watershed

Sl.No.	Particulars	LL (2)		MF (14)		SF (7)		SMF (10)		MDF (3)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	BPL	2	100	14	100	7	100	10	100	3	100	36	100

Participation in NREGA programme: The data regarding Participation in NREGA programme in Chandaraki-1 Micro watershed is presented in Table 55. The results

indicated that, only 38.89 percent of the households have participated in NREGA programme.

Table 55. Participation in NREGA programme in Chandaraki-1 micro-watershed

Sl. No.	Particulars	LL (2)		MF (14)		SF (7)		SMF (10)		MDF (3)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Participation in NREGA programme	0	0	3	21.4	6	85.7	4	40	1	33	14	38.9

Adequacy of food items: The data regarding adequacy of food items in Chandaraki-1 Micro watershed is presented in Table 56. The results indicated that, the extent of adequacy of food items for cereals, pulses, Oilseeds and vegetables were 86.11, 83.33, 5.56, 27.78 per cent respectively, similarly for Fruits (5.56%), milk (91.67%), Egg (16.67%) and Meat (16.67%).

Table 56. Adequacy of food items in Chandaraki-1 micro-watershed

Sl.No.	Particulars	LL (2)		MF (14)		SF (7)		SMF (10)		MDF (3)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Cereals	2	100	12	85.7	7	100	8	80	2	66.67	31	86.11
2	Pulses	1	50	12	85.7	7	100	8	80	2	66.67	30	83.33
3	Oilseed	0	0	0	0	2	28.57	0	0	0	0	2	5.56
4	Vegetables	0	0	5	35.7	0	0	4	40	1	33.33	10	27.78
5	Fruits	0	0	0	0	0	0	2	20	0	0	2	5.56
6	Milk	1	50	13	92.9	7	100	10	100	2	66.67	33	91.67
7	Egg	0	0	3	21.4	0	0	3	30	0	0	6	16.67
8	Meat	0	0	3	21.4	0	0	3	30	0	0	6	16.67

Inadequacy of food items: The data regarding in adequacy of food items in Chandaraki-1 Micro watershed is presented in Table 57. The results indicated that, the extent of in adequacy of food items for cereals, pulses, Oilseeds and vegetables were 13.89, 16.67, 38.89, 13.89 and 25.00 per cent respectively, similarly for fruits (36.11%), milk (2.78%), egg (27.78%) and meat (25.00%).

Table 57. Inadequacy of food items in Chandaraki-1 micro-watershed

Sl.No.	Particulars	LL (2)		MF (14)		SF (7)		SMF (10)		MDF (3)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Cereals	0	0	2	14.3	0	0	2	20	1	33.33	5	13.89
2	Pulses	1	50	2	14.3	0	0	2	20	1	33.33	6	16.67
3	Oilseed	2	100	6	42.9	0	0	5	50	1	33.33	14	38.89
4	Vegetables	2	100	1	7.14	2	28.57	0	0	0	0	5	13.89
5	Fruits	1	50	5	35.7	2	28.57	4	40	1	33.33	13	36.11
6	Milk	1	50	0	0	0	0	0	0	0	0	1	2.78
7	Egg	2	100	2	14.3	2	28.57	3	30	1	33.33	10	27.78
8	Meat	2	100	2	14.3	1	14.29	3	30	1	33.33	9	25

Farming constraints: The data regarding farming constraints experienced by households in Chandaraki-1 Micro watershed is presented in Table 58. The results indicated that, lower fertility status of the soil was the constraint experienced by

(91.67 %) per cent of the households, wild animal menace on farm field (55.56%), frequent incidence of pest and diseases (55.56%), inadequacy of irrigation water (72.22%), high cost of fertilizers and plant protection chemicals (63.89%), high rate of interest on credit (66.67%), low price for the agricultural commodities (69.44 %), lack of marketing facilities in the area (52.78%), inadequate extension services (41.67 %), lack of transport for safe transport of the agricultural produce to the market (36.11%), less rainfall (2.78%), source of agri-technology information (Newspaper/Tv/Mobile) (2.78%).

Table 58. Farming constraints experienced in Chandaraki-1 micro-watershed

SN	Particulars	MF (14)		SF (7)		SMF (10)		MDF (3)		All (36)	
		N	%	N	%	N	%	N	%	N	%
1	Lower fertility status of the soil	13	92.86	7	100	10	100	3	100	33	91.67
2	Wild animal menace on farm field	7	50	3	42.86	7	70	2	66.67	20	55.56
3	Frequent incidence of pest and diseases	9	64.29	3	42.86	6	60	1	33.33	20	55.56
4	Inadequacy of irrigation water	11	78.57	6	85.71	7	70	2	66.67	26	72.22
5	High cost of Fertilizers and plant protection chemicals	6	42.86	6	85.71	9	90	2	66.67	23	63.89
6	High rate of interest on credit	12	85.71	1	14.29	8	80	2	66.67	24	66.67
7	Low price for the agricultural commodities	12	85.71	5	71.43	7	70	1	33.33	25	69.44
8	Lack of marketing facilities in the area	7	50	4	57.14	7	70	1	33.33	19	52.78
9	Inadequate extension services	6	42.86	3	42.86	5	50	1	33.33	15	41.67
10	Lack of transport for safe transport of the Agril produce to the market.	5	35.71	2	28.57	4	40	1	33.33	13	36.11
11	Less rainfall	0	0	0	0	0	0	0	0	1	2.78
12	Source of Agri-technology information	0	0	0	0	0	0	0	0	1	2.78

SUMMARY AND IMPLICATIONS

In order to assess the socio-economic condition of the farmers in the watershed 36 households located in the micro watershed were interviewed for the survey. The study was conducted in Chandaraki-1 micro-watershed (Chandaraki sub-watershed, Yadgiri taluk & District) is located at North latitude 16⁰ 52' 29.368" and 16⁰ 50' 46.558" and East longitude 77⁰ 27' 47.027" and 77⁰ 26' 17.986" covering an area of about 603.07 ha bounded by under Chandraki and Madhepalli Villages.

Socio-economic analysis of Chandaraki-1 micro watersheds of Chandaraki sub-watershed, Yadgiri taluk & District indicated that, out of the total sample of 36 farmers were sampled in Chandaraki-1 micro-watershed among households surveyed 14 (38.89%) were marginal, 7 (19.44%) were small, 10 (27.78 %) were semi medium and 3 (8.33 %) were medium large farmers. 2 landless farmers were also interviewed for the survey. The population characteristics of households indicated that, there were 89 (55.97%) men and 70 (44.03 %) were women. The average population of landless was 2.5, marginal farmers were 4.6, small farmers were 4.3, semi medium farmers were 4.8 and medium farmers were 4. Majority of the respondents (38.36%) were in the age group of 16-35 years.

Education level of the sample households indicated that, there were 44.65 per cent of illiterates, 17.61 per cent of them had primary school education, 13.84 per cent high school education, 8.18 per cent of them had PUC education, 0.63 per cent of them had Diploma, 9.43 per cent attained graduation and 3.14 them had other education. About, 80.56 per cent of household heads practicing agriculture and 2.78 per cent of the household heads were engaged as agricultural labourers.

Agriculture was the major occupation for 80.56 per cent of the household members. In the study area, 72.22 per cent of the households possess katcha house and 8.33 per cent possess pucca house. The durable assets owned by the households showed that, 72.22 per cent possess TV, 80.56 per cent possess mixer grinder, 94.44 per cent possess mobile phones and 30.56 per cent possess motor cycles.

Farm implements owned by the households indicated that, 8.33 per cent of the households possess Bullock Cart, 36.11 per cent possess plough and 77.78 per cent possess Weeder, 8.33 per cent possess tractor, 5.56 per cent possess Harvester and 36.11 per cent possess Thresher. Regarding livestock possession by the households, 2.78 per cent possess local cow and 5.56 per cent possess buffalo.

The average labour availability in the study area showed that, own men and women labour availability in the micro watershed was 14.06 each, while the hired labour (men) availability was 1.22. Further, 5.56 per cent of the households opined that hired labour was inadequate during the agricultural season.

In the study area, about 0.63 per cent of the respondents migrated from the micro watershed in search of jobs with an average distance of 850.00 kms for about 4.00 months. Out of the total land holding of the sample respondents 94.31 per cent (59.84 ha) of the area is under dry condition and the remaining 4.24 per cent area is irrigated land.

Open well was the major source of irrigation for 5.56 per cent of the households. The major crops grown by sample farmers are Red gram, Cotton, Bengal gram, Jowar and Paddy and cropping intensity was recorded as 100.00 per cent. Out of the sample households 94.44 percent possessed bank account. About 94.44 per cent of the respondents borrowed credit from various sources. Among the credit borrowed by households, 13.64 per cent have borrowed loan from commercial banks and 86.36 per cent from co-operative/Grameena bank. Majority of the respondents (95.45%) have borrowed loan for agriculture purpose.

The per hectare cost of cultivation for Red gram, Cotton, Bengal gram, Jowar and Paddy was Rs.48868.88 , 44389.25, 18089.82, 26988.16, and 23946.94 with benefit cost ratio of 1:0.90, 1: 1.00, 1: 3.80, 1: 0.50, and 1:0.70 , respectively. Further, 25.00 per cent of the households opined that dry fodder was adequate and 8.33 per cent of the households have opined that the green fodder was adequate.

The average annual gross income of the farmers was Rs. 76055.56 in micro-watershed, of which Rs. 41027.78 comes from agriculture. Sampled households have grown 8 horticulture trees and 100 forestry trees together in the fields and back yards. Households have an average investment capacity of Rs. 12361.11 for land development. Source of funds for additional investment is concerned, 88.89 per cent depends on own funds and 2.78 per cent depends on bank loan for land development activities.

Regarding marketing channels, 116.67 per cent of the households have sold agricultural produce to the local/village merchants. Further, 116.67 per cent of the households have used tractor for the transport of agriculture commodity. Majority of the farmers (88.89%) have experienced soil and water erosion problems in the watershed and 94.44 per cent of the households were interested towards soil testing.

Firewood was the major source of fuel for domestic use for 88.89 per cent of the households and 11.11 per cent households has LPG connection. Piped supply was the major source for drinking water for 97.22 per cent of the households. Electricity was the major source of light for 100.00 per cent of the households. In the study area, 30.56 per cent of the households possess toilet facility. Regarding possession of PDS card, 100.00 per cent of the households possessed BPL card. Households opined that, the requirement of cereals (86.11%), pulses (83.33%) and oilseeds (5.56%) are adequate for consumption.

Farming constraints experienced by households in the micro watersheds were lower fertility status of the soil (91.67%) wild animal menace on farm field (55.56%), frequent incidence of pest and diseases (55.56%), inadequacy of irrigation water (72.22%), high cost of fertilizers and plant protection chemicals (63.89%), high rate of interest on credit (66.67%), low price for the agricultural commodities (69.44%), lack of marketing facilities in the area (52.78%), inadequate extension services (41.67%), lack of transport for safe transport of the agricultural produce to the market (36.11%), Less rainfall (2.78%) and Source of Agri-technology information (Newspaper/ TV/Mobile) (2.78%).

Implications of the survey

- ✓ Result indicated that, there were 44.65 per cent were illiterate hence, extension methodologies such as demonstration, street play, drama, video shows will be effective in dissemination of the technologies in the micro watershed.
- ✓ The data indicate that, 72.22 per cent of the households possess katcha house. Hence, the development department while implementing the watershed plan should focus on agriculture to enhance the productivity of major crops in the area to increase the income of the farmers.
- ✓ Results indicated that the local institutional participation of the household members in the micro watershed is minimal hence, activities like membership campaign, awareness creation about the benefits of membership in local institutions and strengths of organized groups must be conveyed.
- ✓ Majority of the households in the watershed have experience in use of mobile phones, and television hence, these mass media can be effectively utilized for transfer of technology as well as for information dissemination.
- ✓ The farm machinery/implement possession in the micro watershed was found to be minimum the reasons may lack of knowledge or lack of financial ability which can be addressed through training on use of different farm implements, providing information on different sources of finance for purchase of farm implements.
- ✓ The possession of livestock such as crossbred cow found is less hence, farmers must be made aware of the benefits of crossbred cow in increased milk production.
- ✓ The possession of livestock such as sheep, goat and poultry was found to be low hence, farmers may be informed the role of subsidiary enterprises in enhancing the income and information on financial support for subsidiary activities.
- ✓ The data indicate that, job/work was the reason for all the migrants hence, farmers may be trained on profitable agriculture or self employment such as animal husbandry, plate making, sheep rearing, goat rearing, rabbit rearing with suitable information on sources of financial support.

- ✓ The results indicate that there was a change in quality of life due to migration hence, the developmental departments should take actions to arrest migration and to improve the quality of the life in rural areas.
- ✓ Households possess 56.43ha (94.31 %) of dry land and 2.54ha (4.24 %) of irrigated land hence, the availability of the dry land agricultural technologies such as short duration crops, high yielding drought resistance crop varieties, drip irrigation technology and subsidy information will be helpful for the farmers to enhance the productivity of land and as well as farmers income.
- ✓ Few of the bore well in micro watershed found non functional hence, farmers may be trained on possibility of bore well rejuvenation.
- ✓ Bore well was major source of irrigation for 12.0 per cent of the households. Hence, in order to increase the area under irrigation as well as to increase the water use efficiency farmers may trained on drip irrigation and provides the information on subsidy for drip irrigation equipment's along with the information on different agencies which provides the financial assistance for drip irrigation.
- ✓ The cropping intensity in the micro watershed was found to be (100.00 %) hence, care must be taken by the implementing agency to bring uncultivated land into cultivation through suitable measures.
- ✓ Many of the household members have borrowed loan from cooperative banks which has higher rate of interest hence, farmers may be sensitized on the different sources of credit with lesser interest rate such SHGs etc.
- ✓ The results indicated the non availability of both green and dry fodder throughout the year hence, fodder development activities can be taken up in the micro watershed.
- ✓ The average annual gross income of the households Rs.41027.78 from agriculture and Rs. 23583.33 from wages and. Agriculture was found to be the major source of income for households hence; the development activities should focus on productivity enhancement, marketing arrangements and agricultural technology dissemination to have a direct impact on the farmers.
- ✓ The cultivation of forest species is found minimal hence, information and production technology related to agro-forestry and integrated farming system.
- ✓ The data indicated that, 88.89 per cent of the households have experienced soil and water erosion problems. Hence, those farmers who reported the soil and water erosion problems may be given attention while implementation of the watershed development plan.
- ✓ The data indicated that, 94.44 per cent of the households have interest in soil testing hence, farmers must be provided with the information on various institutions which are involved in soil testing for the benefit of the farmers.
- ✓ Except summer ploughing the adoption of other soil and water conservation structures is minimum hence, the farmers in the micro watershed should be

sensitized on the use of different conservation structures for soil water conservation.

- ✓ Cereals and pulses found be adequate for per cent of the households respectively hence, farm households and the farm women must be trained on importance of balanced nutrition and role of vegetable, milk, egg, meat in balanced diet.
- ✓ Lower fertility status of the soil (91.67%), wild animal menace on farm field (55.56%), frequent incidence of pest and diseases (55.56%), high cost of fertilizers and plant protection chemicals (63.89%), high rate of interest on credit (66.67%), low price for the agricultural commodities (69.44%), lack of marketing facilities in the area (52.78%), inadequate extension services (41.67%), lack of transport for safe transport of the agricultural produce to the market (36.11%) were the major farming constraints experienced hence, these constraints must be addressed immediately for the welfare of the farmers. Awareness to be created among the farmers to approach nearest KVKs/RSKs and other developmental departments for technical and for subsidized inputs and utilize the well established regulated markets, approaching the contract firms, direct markets to avoid the involvement of middlemen.