



हर कदम, हर डगर
किसानों का हमसफर
भारतीय कृषि अनुसंधान परिषद

Agrisearch with a human touch

**LAND RESOURCE INVENTORY AND SOCIO-ECONOMIC STATUS OF
FARM HOUSEHOLDS FOR WATERSHED PLANNING AND
DEVELOPMENT**

YAGAPUR TANDA-1 (4D5B2H1c) MICROWATERSHED

Yadgir Taluk & District, Karnataka

Karnataka Watershed Development Project – II

SUJALA – III

World Bank funded Project



THE WORLD BANK



ICAR – NATIONAL BUREAU OF SOIL SURVEY AND LAND USE PLANNING



ICAR - NBSS & LUP

**WATERSHED DEVELOPMENT DEPARTMENT
GOVT. OF KARNATAKA, BANGALORE**



About ICAR - NBSS&LUP

The ICAR-National Bureau of Soil Survey and Land Use Planning (ICAR-NBSS&LUP), Nagpur, a premier Institute of the Indian Council of Agricultural Research (ICAR), was set up during 1976 with the objective to prepare soil resource maps at national, state and district levels and to provide research inputs in soil resource mapping and its applications, land evaluation, land use planning, land resource management, and database management using GIS for optimising land use on different kinds of soils in the country.

The Bureau has been engaged in carrying out soil resource survey, agro-ecological and soil degradation mapping at the country, state and district levels for qualitative assessment and monitoring the soil health towards viable land use planning. The research activities have resulted in identifying the soil potentials and problems, and the various applications of the soil surveys with the ultimate objective of sustainable agricultural development. The Bureau has the mandate to correlate and classify soils of the country and maintain a National Register of all the established soil series. The Institute is also imparting in-service training to staff of the soil survey agencies in the area of soil survey, land evaluation and soil survey interpretations for land use planning. The Bureau in collaboration with Panjabrao Krishi Vidyapeeth, Akola is running post-graduate teaching and research programme in land resource management, leading to M.Sc. and Ph.D. degrees.

Citation: Rajendra Hegde, Ramesh kumar S.C. B.A. Dhanorkar, K.V. Niranjana, S. Srinivas, M.Lalitha, R.S. Reddy and S.K. Singh (2019). Land resource inventory and socio-economic status of farm households for watershed planning and development of Yagapur Tanda-1 (4D5B2H1c) Microwatershed, Yadgir Taluk and District, Karnataka”, ICAR-NBSS&LUP Sujala MWS Publ.579, ICAR – NBSS & LUP, RC, Bangalore. p.140 & 32.

TO OBTAIN COPIES,

Please write to:

Director, ICAR - NBSS & LUP,

Amaravati Road, NAGPUR - 440 033, India

Phone : (0712) 2500386, 2500664, 2500545 (O)

Telefax : 0712-2522534

E-Mail : director@nbsslup.ernet.in

Website URL : nbsslup.in

Or

Head, Regional Centre, ICAR - NBSS&LUP, Hebbal, Bangalore - 560 024

Phone : (080) 23412242, 23510350 (O)

Telefax : 080-23510350

E-Mail : nbssrcb@gmail.com



**LAND RESOURCE INVENTORY AND SOCIO-ECONOMIC
STATUS OF FARM HOUSEHOLDS FOR WATERSHED
PLANNING AND DEVELOPMENT**

YAGAPUR TANDA-1 (4D5B2H1c) MICROWATERSHED

Yadgir Taluk & District, Karnataka

Karnataka Watershed Development Project – II

Sujala-III

World Bank funded Project



THE WORLD BANK



**ICAR – NATIONAL BUREAU OF SOIL SURVEY AND LAND USE
PLANNING**



**WATERSHED DEVELOPMENT DEPARTMENT, GOVT. OF
KARNATAKA, BANGALORE**



PREFACE

In Karnataka, as in other Indian States, the livelihoods of rural people are intertwined with farming pursuits. The challenges in agriculture are seriously threatening the livelihood of a large number of farmers as they have been practicing farming in contextual factors beyond their control. Climatic factors are the most important ones and have become much more significant in recent times due to rapid climate changes induced by intensive anthropogenic activities affecting our ecosystem in multiple ways. Climate change has become the reality, it is happening and efforts to evolve and demonstrate climate resilient technologies have become essential. Due to the already over stressed scenario of agrarian sector, the climate change is resulting in manifold increase in the complexities, pushing the rural mass to face more and more unpredictable situations. The rising temperatures and unpredictable rainfall patterns are going to test seriously the informed decisions farmers have to make in order to survive in farming and sustain their livelihood.

It is generally recognized that impacts of climate change shall not be uniform across the globe. It is said that impact of climate change is more severe in South Asia. Based on the analysis of meteorological data, it is predicted that in India, there will be upward trend in mean temperature, downward trend in relative humidity, annual rainfall and number of wet days in a year. Also, in general, phenomena like erratic monsoon, spread of tropical diseases, rise in sea levels, changes in availability of fresh water, frequent floods, droughts, heat waves, storms and hurricanes are predicted. Each one of these adverse situations are already being experienced in various parts of India and also at the global level. Decline in agricultural productivity of small and marginal farmers becoming more vulnerable is already witnessed.

In Karnataka, more than 60 per cent of the population live in rural areas and depend on agriculture and allied activities for their livelihood. Though the state has achieved significant progress in increasing the yield of many crops, there is tremendous pressure on the land resources due to the growing and competing demands of various land uses. This is reflected in the alarming rate of land degradation observed. Already more than 50 per cent of the area is affected by various forms of degradation. If this trend continues, the sustainability of the fragile ecosystem will be badly affected. The adverse effects of change in the climatic factors are putting additional stress on the land resources and the farmers dependent on this.

The natural resources (land, water and vegetation) of the state need adequate and constant care and management, backed by site-specific technological interventions and investments particularly by the government. Detailed database pertaining to the nature of the land resources, their constraints, inherent potentials and suitability for various land

based rural enterprises, crops and other uses is a prerequisite for preparing location-specific action plans, which are in tune with the inherent capability of the resources. Any effort to evolve climate resilient technologies has to be based on the baseline scientific database. Then only one can expect effective implementation of climate resilient technologies, monitor the progress, make essential review of the strategy, and finally evaluate the effectiveness of the implemented programs. The information available at present on the land resources of the state are of general nature and useful only for general purpose planning. Since the need of the hour is to have site-specific information suitable for farm level planning and detailed characterization and delineation of the existing land resources of an area into similar management units is the only option.

ICAR-NBSS&LUP, Regional Centre, Bangalore has taken up a project sponsored by the Karnataka Watershed Development Project-II, (Sujala-III), Government of Karnataka funded by the World Bank under Component-1 Land Resource Inventory. This study was taken up to demonstrate the utility of such a database in reviewing, monitoring and evaluating all the land based watershed development programs on a scientific footing. To meet the requirements of various land use planners at grassroots level, the present study on Land Resource Inventory and Socio-Economic Status of Farm Households for Watershed Planning and Development of for Yagapur Tanda-1 microwatershed in Yadgir Taluk and District, Karnataka” for integrated development was taken up in collaboration with the State Agricultural Universities, IISC, KRSRAC, KSNDMC as Consortia partners. The project provides detailed land resource information at cadastral level (1:7920 scale) for all the plots and socio-economic status of farm households covering thirty per cent farmers randomly selected representing landed and landless class of farmers in the micro-watershed. The project report with the accompanying maps for the microwatershed will provide required detailed database for evolving effective land use plan, alternative land use options and conservation plans for the planners, administrators, agricultural extension personnel, KVK officials, developmental departments and other land users to manage the land resources in a sustainable manner.

It is hoped that this database will be useful to the planners, administrators and developmental agencies working in the area in not only for formulating location specific developmental schemes but also for their effective monitoring at the village/watershed level.

Nagpur
Date:23-11-2019

S.K. SINGH
Director, ICAR - NBSS&LUP, Nagpur

Contributors

Dr. Rajendra Hegde Principal Scientist, Head & Project Leader, Sujala-III Project ICAR-NBSS&LUP, Regional Centre, Bangalore	Dr. S.K.Singh Director, ICAR-NBSS&LUP Coordinator, Sujala-III Project Nagpur
Soil Survey, Mapping & Report Preparation	
Dr. B.A. Dhanorkar	Sh. R.S. Reddy
Dr. K.V. Niranjana	Sh. Somashekar T N
	Smt. Chaitra, S.P.
	Dr. Gopali bardhan
Field Work	
Sh. C.BacheGowda	Sh. Mahesh, D.B.
Sh. Somashekar	Sh. Ashok S Sindagi
Sh. M. Jayaramaiah	Sh. Veerabhadrapa B.
Sh. Paramesha, K.	Sh. Shankarappa
Sh. B. M. Narayana Reddy	Sh. Anand
	Sh. Arun N Kambar.
	Sh Kamalesh Awate
	Sh. Sharaan Kumar Huppar
	Sh. Yogesh H.N.
	Sh. Kalaveerachari R Kammar
GIS Work	
Dr. S.Srinivas	Sh. A.G.Devendra Prasad
Sh. D.H.Venkatesh	Sh. Prakashanaik, M.K.
Smt.K.Sujatha	Sh. Abhijith Sastry, N.S.
Smt. K.V.Archana	Sh. Sudip Kumar Suklabaidya
Sh. N. Maddileti	Sh. Avinash, K.N.
	Sh. Amar Suputhra, S
	Sh. Deepak, M.J.
	Smt. K.Karunya Lakshmi
	Ms. Seema, K.V.
	Ms. A. Rajab Nisha

Laboratory Analysis	
Dr. K.M.Nair	Ms. Steffi Peter
Smt. Arti Koyal	Ms. Thara, V.R
Smt. Parvathy	Ms. Roopa, G.
	Ms. Swati, H.
	Sh. Shantaveera Swami
	Ms. Shwetha, N.K.
	Smt. Ishrat Haji
	Ms. P. Pavan Kumari
	Ms. Padmaja
	Ms. Veena, M.
Socio-Economic Analysis	
Dr. S.C. Ramesh Kumar	Sh. M.K. Prakashanaik
	Ms. Karuna V. Kulkarni
	Mrs. Sowmya A.N
	Sh. Vinod R
	Sh. Basavaraja
	Sh. Vijay Kumar Lamani
	Ms. Sowmya K.B
	Mrs. Prathibha, D.G
	Sh. Rajendra,D
Soil & Water Conservation	
Sh. Sunil P. Maske	
Watershed Development Department, GoK, Bangalore	
Sh. Rajeev Ranjan IFS Project Director & Commissioner, WDD	Dr. A. Natarajan NRM Consultant, Sujala-III Project
Dr. S.D. Pathak IFS Executive Director & Chief Conservator of Forests, WDD	

PART-A

LAND RESOURCE INVENTORY

Contents

Preface		
Contributors		
Executive Summary		
Chapter 1	Introduction	1
Chapter 2	Geographical Setting	3
2.1	Location and Extent	3
2.2	Geology	3
2.3	Physiography	4
2.4	Drainage	4
2.5	Climate	4
2.6	Natural Vegetation	6
2.7	Land Utilization	6
Chapter 3	Survey Methodology	11
3.1	Base maps	11
3.2	Image Interpretation for Physiography	11
3.3	Field Investigation	14
3.4	Soil Mapping	16
3.5	Land Management Units	17
3.6	Laboratory Characterization	17
Chapter 4	The Soils	23
4.1	Soils of granite gneiss landscape	23
Chapter 5	Interpretation for Land Resource Management	47
5.1	Land Capability Classification	47
5.2	Soil Depth	49
5.3	Surface Soil Texture	50
5.4	Soil Gravelliness	52
5.5	Available Water Capacity	53
5.6	Soil Slope	54
5.7	Soil Erosion	55
Chapter 6	Fertility Status	56
6.1	Soil Reaction (pH)	56
6.2	Electrical Conductivity (EC)	56
6.3	Organic Carbon (OC)	56
6.4	Available Phosphorus	58
6.5	Available Potassium	58
6.6	Available Sulphur	58
6.7	Available Boron	58
6.8	Available Iron	59
6.9	Available Manganese	59
6.10	Available Copper	59
6.11	Available Zinc	63

Chapter 7	Land Suitability for Major Crops	64
7.1	Land suitability for Sorghum	64
7.2	Land suitability for Maize	65
7.3	Land suitability for Bajra	66
7.4	Land suitability for Groundnut	67
7.5	Land suitability for Sunflower	68
7.6	Land suitability for Redgram	69
7.7	Land suitability for Bengal gram	70
7.8	Land suitability for Cotton	71
7.9	Land suitability for Chilli	72
7.10	Land suitability for Tomato	73
7.11	Land suitability for Brinjal	74
7.12	Land suitability for Onion	75
7.13	Land suitability for Bhendi	76
7.14	Land suitability for Drumstick	77
7.15	Land suitability for Mango	78
7.16	Land suitability for Guava	79
7.17	Land suitability for Sapota	80
7.18	Land Suitability for Pomegranate	81
7.19	Land Suitability for Musambi	82
7.20	Land Suitability for Lime	83
7.21	Land Suitability for Amla	84
7.22	Land Suitability for Cashew	85
7.23	Land Suitability for Jackfruit	86
7.24	Land Suitability for Jamun	87
7.25	Land Suitability for Custard apple	88
7.26	Land Suitability for Tamarind	89
7.27	Land Suitability for Mulberry	90
7.28	Land Suitability for Marigold	91
7.29	Land Suitability for Chrysanthemum	92
7.30	Land Management Units (LMUs)	124
7.31	Proposed Crop Plan for Yagapur Tanda-1 Microwatershed	125
Chapter 8	Soil Health Management	128
Chapter 9	Soil and Water conservation Treatment Plan	132
9.1	Treatment Plan	132
9.2	Recommended Soil and Water Conservation measures	136
9.3	Greening of Microwatershed	137
	References	140
	Appendix I	I-VIII
	Appendix II	IX-XV
	Appendix III	XVI-XXI

LIST OF TABLES

2.1	Mean Monthly Rainfall, PET, 1/2 PET at Yadgir Taluk & District	5
2.2	Land Utilization in Yadgir district	7
3.1	Differentiating Characteristics used for Identifying Soil Series	16
3.2	Soil map unit description of Yagapur Tanda-1 Microwatershed	17
4.1	Physical and Chemical Characteristics of Soil Series identified in Yagapur Tanda-1 microwatershed	33
7.1	Soil-Site Characteristics of Yagapur Tanda-1 Microwatershed	94
7.2	Crop suitability for Sorghum	95
7.3	Crop suitability for Maize	96
7.4	Crop suitability for Bajra	97
7.5	Crop suitability for Groundnut	98
7.6	Crop suitability for Sunflower	99
7.7	Crop suitability for Redgram	100
7.8	Crop suitability for Bengal gram	101
7.9	Crop suitability for Cotton	102
7.10	Crop suitability for Chilli	103
7.11	Crop suitability for Tomato	104
7.12	Crop suitability for Brinjal	105
7.13	Crop suitability for Onion	106
7.14	Crop suitability for Bhendi	107
7.15	Crop suitability for Drumstick	108
7.16	Crop suitability for Mango	109
7.17	Crop suitability for Guava	110
7.18	Crop suitability for Sapota	111
7.19	Crop suitability for Pomegranate	112
7.20	Crop suitability for Musambi	113
7.21	Crop suitability for Lime	114
7.22	Crop suitability for Amla	115
7.23	Crop suitability for Cashew	116
7.24	Crop suitability for Jackfruit	117
7.25	Crop suitability for Jamun	118
7.26	Crop suitability for Custard apple	119

7.27	Crop suitability for Tamarind	120
7.28	Crop suitability for Mulberry	121
7.29	Crop suitability for Marigold	122
7.30	Crop suitability for Chrysanthemum	123
7.31	Proposed Crop Plan for Yagapur Tanda-1 Microwatershed	126

LIST OF FIGURES

2.1	Location map of Yagapur Tanda-1 Microwatershed	3
2.2	Granite and granite gneiss rock formation	4
2.3	Rainfall distribution in Yadgir Taluk & District	5
2.4	Natural vegetation of Yagapur Tanda-1 Microwatershed	6
2.5	Current Land use map of Yagapur Tanda-1 Microwatershed	7
2.6	Location of wells map of Yagapur Tanda-1 Microwatershed.	8
2.7 a	Different crops and cropping systems in Yagapur Tanda-1 Microwatershed	8
2.7 b	Different crops and cropping systems in Yagapur Tanda-1 Microwatershed	9
3.1	Scanned and Digitized Cadastral map of Yagapur Tanda-1 Microwatershed	13
3.2	Satellite image of Yagapur Tanda-1 Microwatershed	13
3.3	Cadastral map overlaid on IRS PAN+LISS IV merged imagery of Yagapur Tanda-1 Microwatershed	14
3.4	Location of profiles in a transect	15
3.5	Soil phase or management units of Yagapur Tanda-1 Microwatershed	21
5.1	Land Capability Classification map of Yagapur Tanda-1 Microwatershed	49
5.2	Soil Depth map of Yagapur Tanda-1 Microwatershed	50
5.3	Surface Soil Texture map of Yagapur Tanda-1 Microwatershed	51
5.4	Soil Gravelliness map of Yagapur Tanda-1 Microwatershed	52
5.5	Soil Available Water Capacity map Yagapur Tanda-1 Microwatershed	53
5.6	Soil Slope map of Yagapur Tanda-1 Microwatershed	54
5.7	Soil Erosion map of Yagapur Tanda-1 Microwatershed	55
6.1	Soil Reaction (pH) map of Yagapur Tanda-1 Microwatershed	57
6.2	Electrical Conductivity (EC) map of Yagapur Tanda-1 Microwatershed	57
6.3	Soil Organic Carbon (OC) map of Yagapur Tanda-1 Microwatershed	58
6.4	Soil Available Phosphorus map of Yagapur Tanda-1 Microwatershed	59
6.5	Soil Available Potassium map of Yagapur Tanda-1 Microwatershed	60
6.6	Soil Available Sulphur map of Yagapur Tanda-1 Microwatershed	60
6.7	Soil Available Boron map of Yagapur Tanda-1 Microwatershed	61
6.8	Soil Available Iron map of Yagapur Tanda-1 Microwatershed	61
6.9	Soil Available Manganese map of Yagapur Tanda-1 Microwatershed	62

6.10	Soil Available Copper map of Yagapur Tanda-1 Microwatershed	62
6.11	Soil Available Zinc map of Yagapur Tanda-1 Microwatershed	63
7.1	Land suitability for Sorghum	65
7.2	Land suitability for Maize	66
7.3	Land suitability for Bajra	67
7.4	Land suitability for Groundnut	68
7.5	Land suitability for Sunflower	69
7.6	Land suitability for Redgram	70
7.7	Land suitability for Bengal gram	71
7.8	Land suitability for Cotton	72
7.9	Land suitability for Chilli	73
7.10	Land suitability for Tomato	74
7.11	Land suitable for Brinjal	75
7.12	Land suitable for Onion	76
7.13	Land suitable for Bhendi	77
7.14	Land suitable for Drumstick	78
7.15	Land suitability for Mango	79
7.16	Land suitability for Guava	80
7.17	Land suitability for Sapota	81
7.18	Land suitability for Pomegranate	82
7.19	Land suitability for Musambi	70
7.20	Land suitability for Lime	71
7.21	Land suitability for Amla	72
7.22	Land suitability for Cashew	73
7.23	Land suitability for Jackfruit	75
7.24	Land suitability for Jamun	76
7.25	Land suitability for Custard apple	77
7.26	Land suitability for Tamarind	78
7.27	Land suitability for Mulberry	79
7.28	Land suitability for Marigold	81
7.29	Land suitability for Chrysanthemum	82
7.30	Land management units map of Yagapur Tanda-1 Microwatershed	84
9.1	Soil and water conservation plan map of Yagapur Tanda-1 Microwatershed	100

EXECUTIVE SUMMARY

The land resource inventory of Yagapur Tanda-I Microwatershed was conducted using village cadastral maps and IRS satellite imagery on 1:7920 scale. The false colour composites of IRS imagery were interpreted for physiography and the physiographic delineations were used as base for mapping soils. The soils were studied in several transects and a soil map was prepared with phases of soil series as mapping units. Random checks were made all over the area outside the transects to confirm and validate the soil map unit boundaries. The soil map shows the geographic distribution and extent, characteristics, classification, behavior and use potentials of the soils in the microwatershed.

The present study covers an area of 720 ha in Yadgir taluk & district, Karnataka. The climate is semiarid and categorized as drought-prone with an average annual rainfall of 866 mm, of which about 652 mm is received during south-west monsoon, 138 mm during north-east and the remaining 76 mm during the rest of the year. An area of 498 ha (69%) ha in the microwatershed is covered by soils, about 199 ha (28%) by rock outcrops and about 23 ha (3%) by others (Habitation and water body). The salient findings from the land resource inventory are summarized briefly below.

- ❖ *The soils belong to 14 soil series and 22 soil phases (management units) and 9 land management units.*
- ❖ *The length of crop growing period is about 120-150 days starting from 1st week of June to 4th week of October.*
- ❖ *From the master soil map, several interpretative and thematic maps like land capability, soil depth, surface soil texture, soil gravelliness, available water capacity, soil slope and soil erosion were generated.*
- ❖ *Soil fertility status maps for macro and micronutrients were generated based on the surface soil samples collected at every 320 m grid interval.*
- ❖ *Land suitability for growing 29 major agricultural and horticultural crops was assessed and maps showing the degree of suitability along with constraints were generated.*
- ❖ *An area of about 69 per cent is suitable for agriculture in the microwatershed.*
- ❖ *About 28 per cent area of the microwatershed has soils that are deep to very deep (100- >150 cm), 5 per cent soils are moderately deep (75-100 cm), 9 per cent soils are moderately shallow (50-75 cm) and about 27 per cent soils are very shallow and shallow (<25-50 cm) in the microwatershed.*
- ❖ *About 8 percent soils are sandy, 16 percent soils are loamy and 45 per cent is clayey soils at the surface.*
- ❖ *An area of about 54 per cent is non gravelly (<15%) soils, about 15 per cent soils are gravelly (15-35%) and about <1 per cent soils are very gravelly (35-60%) in the microwatershed.*
- ❖ *About 18 per cent area of the microwatershed is very high (>200 mm/m) in available water capacity. About 6 per cent area of the microwatershed is medium (101-150 mm/m) and about 45 per cent soils are low (51-100 mm/m) and very low (<51mm/m) in available water capacity.*

- ❖ *An area of about 67 percent is very gently sloping (1-3% slope) lands and about 3 per cent soils are nearly level (0-1% slope) lands in the microwatershed.*
- ❖ *An area of about 66 per cent is moderately (e2) eroded and about 3 per cent are slightly (e1) eroded lands in the microwatershed.*
- ❖ *An area of about 3 percent is slightly to moderately acid (pH 5.5- 6.5), about 12 percent is neutral (6.5-7.3), about 24 percent is slightly alkaline (ph 7.3-7.8), about 37 per cent is moderately alkaline (pH 7.8-8.4) in the microwatershed.*
- ❖ *The Electrical Conductivity (EC) of the soils in the entire cultivated area of the microwatershed is $<2 \text{ ds}^{\text{m}^{-1}}$ indicating that the soils are non-saline.*
- ❖ *Organic carbon content is high ($>0.75\%$) in the entire cultivated area of the microwatershed.*
- ❖ *An area of about 6 percent is low ($<23 \text{ kg/ha}$), about 22 percent is medium (23-57 kg/ha) and 41 percent soils are high ($>57 \text{ kg/ha}$) in available phosphorus.*
- ❖ *An area of about 38 per cent is medium (145-337 kg/ha) and 31 per cent is high ($>337 \text{ kg/ha}$) in available potassium in the microwatershed.*
- ❖ *Available sulphur is high ($>20 \text{ ppm}$) in the entire cultivated area of the microwatershed.*
- ❖ *Available boron is low ($<0.5 \text{ ppm}$) in an area of about 60 per cent and medium (0.5-1.0 ppm) in about 9 per cent soils.*
- ❖ *Available iron content is sufficient ($>4.5 \text{ ppm}$) in the entire cultivated area of the microwatershed.*
- ❖ *Available manganese and copper are sufficient in all the soils of the microwatershed.*
- ❖ *Available zinc content is deficient ($<0.6 \text{ ppm}$) in the entire cultivated area of the microwatershed.*
- ❖ *The land suitability for 29 major crops grown in the microwatershed were assessed and the areas that are highly suitable (S1) and moderately suitable (S2) are given below. It is however to be noted that a given soil may be suitable for various crops but what specific crop to be grown may be decided by the farmer looking to his capacity to invest on various inputs, marketing infrastructure, market price and finally the demand and supply position.*

Land suitability for various crops in the Microwatershed

Crop	Suitability Area in ha (%)		Crop	Suitability Area in ha (%)	
	Highly suitable (S1)	Moderately suitable (S2)		Highly suitable (S1)	Moderately suitable (S2)
<i>Sorghum</i>	23(3)	113(16)	<i>Guava</i>	16(2)	11(1)
<i>Maize</i>	20(3)	116(16)	<i>Sapota</i>	16(2)	11(1)
<i>Bajra</i>	26(4)	110(15)	<i>Pomegranate</i>	16(2)	52(7)
<i>Groundnut</i>	22(3)	4(<1)	<i>Musambi</i>	35(3)	33(5)
<i>Sunflower</i>	19(3)	49(7)	<i>Lime</i>	35(3)	33(5)
<i>Redgram</i>	-	74(10)	<i>Amla</i>	26(4)	82(11)
<i>Bengal gram</i>	19(3)	22(3)	<i>Cashew</i>	-	23(3)
<i>Cotton</i>	19(3)	89(12)	<i>Jackfruit</i>	16(2)	11(1)
<i>Chilli</i>	20(3)	110(15)	<i>Jamun</i>	16(2)	19(3)
<i>Tomato</i>	20(3)	69(10)	<i>Custard apple</i>	41(6)	99(12)
<i>Brinjal</i>	20(3)	69(10)	<i>Tamarind</i>	16(2)	19(3)
<i>Onion</i>	20(3)	69(10)	<i>Mulberry</i>	16(2)	11(1)
<i>Bhendi</i>	20(3)	110(15)	<i>Marigold</i>	20(3)	110(15)
<i>Drumstick</i>	16(2)	30(4)	<i>Chrysanthemum</i>	20(3)	110(15)
<i>Mango</i>	16(2)	-			

- ❖ *Apart from the individual crop suitability, a proposed crop plan has been prepared for the identified LMUs by considering only the highly and moderately suitable lands for different crops and cropping systems with food, fodder, fiber and horticulture crops.*
- ❖ *Maintaining soil-health is vital to crop production and conserve soil and land resource base for maintaining ecological balance and to mitigate climate change. For this, several ameliorative measures have been suggested to these problematic soils like saline/alkali, highly eroded, sandy soils etc.*
- ❖ *Soil and water conservation treatment plan has been prepared that would help in identifying the sites to be treated and also the type of structures required.*
- ❖ *As part of the greening programme, several tree species have been suggested to be planted in marginal and submarginal lands, field bunds and also in the hillocks, mounds and ridges. This would help in not only supplementing the farm income but also provide fodder and fuel to generate lot of biomass which would help in maintaining an ecological balance and also contribute to mitigating the climate change.*

INTRODUCTION

Land is a scarce resource and basic unit for any material production. It can support the needs of the growing population, provided they use the land in a rational and judicious manner. But what is happening in many areas of the state is a cause for concern to everyone involved in the management of land resources at the grassroots level. The area available for agriculture is about 51 per cent of the total geographical area and more than 60 per cent of the people are still dependant on agriculture for their livelihood. The limited land area is under severe stress and strain due to increasing population pressure and competing demands of various land uses. Due to this, every year there is significant diversion of farm lands and water resources for non-agricultural purposes. Apart from this, due to lack of interest in farmers for farming, large tracts of cultivable lands are turning into fallows in many areas and this trend is continuing at an alarming rate.

Further, land degradation has emerged as a serious problem which has already affected about 38 lakh ha of cultivated area in the state. Soil erosion alone has degraded about 35 lakh ha. Almost all the uncultivated areas are facing various degrees of degradation, particularly soil erosion. Salinity and alkalinity has emerged as a major problem in more than 3.5 lakh ha in the irrigated areas of the state. Nutrient depletion and declining factor productivity is common in both rainfed and irrigated areas. The degradation is continuing at an alarming rate and there appears to be no systematic effort among the stakeholders to contain this process. In recent times, an aberration of weather due to climate change phenomenon has added another dimension leading to unpredictable situations to be tackled by the farmers.

In this critical juncture, the challenge before us is not only to increase the productivity per unit area which is steadily declining and showing a fatigue syndrome, but also to prevent or at least reduce the severity of degradation. If the situation is not reversed at the earliest, then the sustainability of the already fragile crop production system and the overall ecosystem will be badly affected in the state. The continued neglect and unscientific use of the resources for a long time has led to the situation observed at present in the state. It is a known fact and established beyond doubt by many studies in the past that the cause for all kinds of degradation is the neglect and irrational use of the land resources. Hence, there is an urgent need to generate a detailed site-specific farm level database on various land resources for all the villages/watersheds in a time bound manner that would help to protect the valuable soil and land resources and also to stabilize the farm production.

Therefore, the land resource inventory required for farm level planning is the one which investigates not only the surface but also consider the other parameters which are critical for productivity *viz.*, soils, climate, water, minerals and rocks, topography, geology, hydrology, vegetation, crops, land use pattern, animal population, socio-

economic conditions, infrastructure, marketing facilities and various schemes and developmental works of the government etc. From the data collected at farm level, the specific problems and potentials of the area can be identified and highlighted, conservation measures required for the area can be planned on a scientific footing, suitability of the area for various uses can be worked out and finally viable and sustainable land use options suitable for each and every land holding can be prescribed.

The Land Resource Inventory is basically done for identifying the potential and problem areas, developing sustainable land use plans, estimation of surface run off and water harvesting potential, preparation of soil and water conservation plans, land degradation/desertification etc. The Bureau is presently engaged in developing an LRI methodology using high resolution satellite remote sensing data and Digital Elevation Model (DEM) data to prepare Landscape Ecological Units (LEU) map representing agro-ecosystem as a whole. The LEU is preferred over landform as the base map for LRI. LEU is the assemblage of landform, slope and land use. An attempt has already been made to upscale the soil resource information from 1:250000 and 1:50000 scale to the LEU map in Goa and other states.

The land resource inventory aims to provide site specific database for Yagapur Tanda-1 microwatershed in Yadgir Taluk & District, Karnataka State for the Karnataka Watershed Development Department. The database was generated by using cadastral map of the village as a base along with high resolution IRS LISS IV and Cartosat-1 merged satellite imagery. Later, an attempt will be made to uplink this LRI data generated at 1:7920 scale under Sujala-III Project to the proposed Landscape Ecological Units (LEUs) map.

The study was organized and executed by the ICAR- National Bureau of Soil Survey and Land Use Planning, Regional Centre, Bangalore under Generation of Land Resource Inventory Data Base Component-1 of the Sujala-III Project funded by the World Bank.

GEOGRAPHICAL SETTING

2.1 Location and Extent

The Yagapur Tanda-1 microwatershed is located in the northern part of Karnataka in Yadgir Taluk & District, Karnataka State (Fig.2.1). It comprises parts of Shivanagara and Yagapura villages. It lies between $16^{\circ} 54'$ and $16^{\circ} 56'$ North latitudes and $77^{\circ} 5'$ and $77^{\circ} 7'$ East longitudes, covering an area of about 720 ha. It is in the northwestern side of Yadgir town and is surrounded by Shivanagara on the north and northwest and Yagapura on the west, east and southern of the microwatershed.




Fig.2.1 Location map of Yagapur Tanda-1 Microwatershed

2.2 Geology

Major rock formations observed in the microwatershed are granite gneiss (Figs.2.2a). Granite gneisses are essentially pink to gray and are coarse to medium grained. They consist primarily of quartz, feldspar, biotite and hornblende. The gray granite gneisses are

highly weathered, fractured and fissured upto a depth of about 10 m. Dolerite dykes and quartz veins are common with variable width and found to occur in Yagapur Tanda-1 microwatershed.



Fig.2.2a Granite and granite gneiss rocks

2.3 Physiography

Physiographically, the area has been identified as granite gneiss based on geology. The area has been further subdivided into five landforms, viz; mounds/ridges, summits, side slopes and very gently sloping uplands, plains and valleys based on slope and its relief features. The elevation ranges from 395-548 m above MSL. The mounds and ridges are mostly covered by rock outcrops.

2.4 Drainage

The area is drained by several parallel streams like Bori, Amerja and Kanga which finally join the river Bhima along its course. Though, they are not perennial, during rainy season they carry large quantities of rain water. The microwatershed has only few small tanks which are not capable of storing the water that flows during the rainy season. Due to this, the ground water recharge is very much affected. This is reflected in the failure of many bore wells in the villages. If the available rain water is properly harnessed by constructing new tanks and recharge structures at appropriate places in the villages, then the drinking and irrigation needs of the area can be easily met. The drainage network is parallel to sub parallel and dendritic.

2.5 Climate

The Yadgir district lies in the northern plains of Karnataka and falls under semiarid tract of the state and is categorized as drought- prone with total annual rainfall of 866 mm (Table 2.1). Of the total rainfall, maximum of 652 mm is received during the south–west monsoon period from June to September; the north-east monsoon from

October to early December contributes about 138 mm and the remaining 76 mm during the rest of the year. The summer season starts during the middle of February and continues up to the first week of June. The period from December to the middle of February is the coldest season. December is the coldest month with mean daily maximum and minimum temperatures being 29.5⁰C and 10⁰C respectively. During peak summer, temperature shoots up to 45⁰C. Relative humidity varies from 26% in summer to 62% in winter. Rainfall distribution is shown in Figure 2.3. The average Potential Evapo-Transpiration (PET) is 141 mm and varies from a low of 81 mm in December to 199 mm in the month of May. The PET is always higher than precipitation in all the months except end of June to end of September. Generally, the Length of crop Growing Period (LGP) is 120-150 days and starts from 1st week of June to 4th week of October.

Table 2.1 Mean Monthly Rainfall, PET, 1/2 PET at Yadgir Taluk, Yadgir District

Sl. No.	Months	Rainfall	PET	1/2 PET
1	January	4.30	86.0	43.0
2	February	2.30	125.5	62.7
3	March	15.10	166.0	83.0
4	April	18.50	179.8	89.9
5	May	36.0	198.8	97.9
6	June	118.0	175.1	87.5
7	July	171.80	156.3	78.1
8	August	182.9	150.3	75.1
9	September	179.7	142.0	71.0
10	October	105.3	138.5	69.2
11	November	26.4	97.60	48.6
12	December	6.0	80.90	40.4
Total		866.3		




Fig 2.3 Rainfall distribution in Yadgir Taluk, Yadgir District

2.6 Natural Vegetation

The natural vegetation is sparse comprising few tree species, shrubs and herbs. The mounds, ridges and boulders occupy very sizeable area which is under thin to moderately thick forest vegetation. Still, there are some remnants of the past forest cover which can be seen in patches in some ridges and hillocks in the microwatershed (Fig 2.4).

Apart from the continuing deforestation, the presence of large population of goats, sheep and other cattle in the microwatershed is causing vegetative degradation of whatever little vegetation left in the area. The uncontrolled grazing has left no time for the regeneration of the vegetative cover. This leads to the accelerated rate of erosion on the hill slopes resulting in the formation of deep gullies in the foot slopes that eventually result in the heavy siltation of tanks and reservoirs in the microwatershed.



Fig 2.4 Natural vegetation of Yagapur Tanda-1 Microwatershed

2.7 Land Utilization

About 72 per cent area (Table 2.2) in Yadgir district is cultivated at present. An area of about 2 per cent is permanently under pasture, 20 per cent under current fallows and 6 per cent under non-agricultural land and 5 per cent under currently barren. Forests occupy an area of about 7 per cent and the tree cover is in a very poor state. Most of the mounds, ridges and bouldery areas have very poor vegetative cover. Major crops grown in the area are sorghum, maize, cotton, sunflower, groundnut, red gram, mango, pomegranate, marigold and sapota. While carrying out land resource inventory, the land use/land cover particulars are collected from all the survey numbers and a current land use map of the microwatershed is prepared. The current land use map prepared shows the arable and non-arable lands, other land uses and different types of crops grown in the area. The current land use map of Yagapur Tanda-1 microwatershed is presented in Fig.2.5. The location of wells and conservation structures in the Yagapur Tanda-1

microwatershed is shown in Fig. 2.6. The different crops and cropping systems adopted in the microwatershed are presented in Figures 2.7 a & b.

Table 2.2 Land Utilization in Yadgir District

Sl. No.	Agricultural land use	Area (ha)	Per cent
1	Total geographical area	516088	-
2	Total cultivated area	373617	72.4
3	Area sown more than once	74081	14.3
4	Cropping intensity	-	119.8
5	Trees and grooves	737	0.14
6	Forest	33773	6.54
7	Cultivable wasteland	2385	0.46
8	Permanent Pasture land	11755	2.28
9	Barren land	27954	5.41
10	Non- Agriculture land	29623	5.73
11	Current Fallows	105212	20.4




Fig.2.5 Current Land Use map of Yagapur Tanda-1 Microwatershed




Fig.2.6 Location of wells and conservation structures map of Yagapur Tanda-1 Microwatershed.




Fig. 2.7 a. Different Crops and Cropping Systems in Yagapur Tanda-1 Microwatershed



Sunflower



Groundnut



Mango



Pomegranate



Cotton



Sorghum

Fig. 2.7 b. Different Crops and Cropping Systems in Yagapur Tanda-1 Microwatershed

SURVEY METHODOLOGY

The purpose of land resource inventory is to delineate similar areas (soil series and phases), which respond or expected to respond similarly to a given level of management. This was achieved in Yagapur Tanda-1 microwatershed by the detailed study of all the soil characteristics (depth, texture, colour, structure, consistence, coarse fragments, porosity, soil reaction, soil horizons etc.) and site characteristics (slope of the land, erosion, drainage, occurrence of rock fragments etc.) followed by grouping of similar areas based on soil-site characteristics into homogeneous (management units) units, and showing the area extent and their geographic distribution on the microwatershed cadastral map. The detailed survey at 1:7920 scale was carried out in an area of 720 ha. The methodology followed for carrying out land resource inventory was as per the guidelines given in Soil Survey Manual (IARI, 1971; Soil Survey Staff, 2006; Natarajan *et al.*, 2015) which is briefly described below.

3.1 Base Maps

The detailed survey of the land resources occurring in the microwatershed was carried out by using digitized cadastral and IRS satellite imagery map as base supplied by KSRSAC. The cadastral map shows field boundaries with their survey numbers, location of tanks, streams and other permanent features of the area (Fig. 3.1). Apart from the cadastral map, remote sensing data products from Cartosat-1 and LISS IV merged at the scale of 1:7920 were used in conjunction with the cadastral map to identify the landscapes, landforms and other surface features. The imagery helped in the identification and delineation of boundaries between hills, uplands and lowlands, water bodies, forest and vegetated areas, roads, habitations and other cultural features of the area (Fig. 3.2). The cadastral map was overlaid on the satellite imagery (Fig. 3.3) that helps to identify the parcel boundaries and other permanent features. Apart from cadastral maps and images, toposheets of the area (1:50,000 scale) were also used for initial traversing, identification of geology and landforms, drainage features, present land use and also for selection of transects in the microwatershed.

3.2 Image Interpretation for Physiography

False Colour Composites (FCCs) of Cartosat-I and LISS-IV merged satellite data covering microwatershed area was visually interpreted using image interpretation elements and all the available collateral data with local knowledge. The delineated physiographic boundaries were transferred on to a cadastral map overlaid on satellite imagery. Physiographically, the area has been identified as granite gneiss landscape. It was divided into five landforms, *viz.*; ridges and mounds, gently and very gently sloping uplands and lowlands based on slope and image characteristics. They were further subdivided into physiographic/image interpretation units based on image characteristics. The image interpretation legend for physiography is given below.

Image Interpretation Legend for Physiography

G- Granite Gneiss Landscape

G1	Hills/ Ridges/ Mounds
G11	Summits
G12	Side slopes
G121	Side slopes with dark grey tones
G2	Uplands
G21	Summits
G22	Gently sloping uplands
G221	Gently sloping uplands, yellowish green (eroded)
G222	Gently sloping uplands, yellowish white (severely eroded)
G23	Very gently sloping uplands
G231	Very gently sloping uplands, yellowish green
G232	Very gently sloping uplands, medium green and pink
G233	Very gently sloping uplands, pink and green (scrub land)
G234	Very gently sloping uplands, medium greenish grey
G235	Very gently sloping uplands, yellowish white (eroded)
G236	Very gently sloping uplands, dark green
G237	Very gently sloping uplands, medium pink (coconut garden)
G238	Very gently sloping uplands, pink and bluish white (eroded)
G3	Valleys/ lowlands
G31	Valleys, pink tones
G32	Valleys gray mixed with pink tones




Fig 3.1 Scanned and Digitized Cadastral map of Yagapur Tanda-1 Microwatershed




Fig.3.2 Satellite Image of Yagapur Tanda-1 Microwatershed




Fig.3.3 Cadastral map overlaid on IRS PAN+LISS IV merged imagery of Yagapur Tanda-1 Microwatershed

3.3 Field Investigation

The field boundaries and survey numbers given on the cadastral sheet were located on the ground by following permanent features like roads, cart tracks, *nallas*, streams, tanks etc., and wherever changes were noticed, they were incorporated on the microwatershed cadastral map. Preliminary traverse of the microwatershed was carried out with the help of cadastral map, imagery and toposheets. While traversing, landforms and physiographic units identified were checked and preliminary soil legend was prepared by studying soils at few selected places. Then, intensive traversing of each physiographic unit like hills, ridges, uplands and valleys was carried out. Based on the variability observed on the surface, transects (Fig. 3.4) were selected across the slope covering all the landform units in the microwatershed (Natarajan and Dipak Sarkar, 2010).




Fig: 3.4. Location of profiles in a transect

In the selected transect, soil profiles were located (Fig. 3.4) at closely spaced intervals to take care of any change in the land features like break in slope, erosion, gravel, stones etc. In the selected sites, profiles (vertical cut showing the soil layers from surface to the rock) were opened upto 200 cm or to the depth limited by rock or hard substratum and studied in detail for all their morphological and physical characteristics. The soil and site characteristics were recorded for all profile sites on a standard proforma as per the guidelines given in USDA Soil Survey Manual (Soil Survey Staff, 2012). Apart from the transect study, profiles were also studied at random, almost like in a grid pattern, outside the transect areas.

Based on the soil characteristics, the soils were grouped into different soil series. Soil series is the most homogeneous unit having similar horizons and properties and behaves similarly for a given level of management. Soil depth, texture, colour, kind of horizon and horizon sequence, calcareousness, amount and nature of gravel present, nature of substratum etc, were used as the major differentiating characteristics for identifying soil series occurring in the area. The differentiating characteristics used for identifying the soil series are given in Table 3.1. Based on the above characteristics, 14 soil series were identified in the Yagapur Tanda-1 microwatershed.

Table 3.1 Differentiating Characteristics used for identifying Soil Series
(Characteristics are of Series Control Section)

Soils of Granite gneiss Landscape							
Sl. no	Soil Series	Depth (cm)	Colour (moist)	Texture	Gravel (%)	Horizon sequence	Calcareousness
1	KKR (Kakalawar)	<25	7.5YR 4/3 10YR 6/3	sl	10-15	Ap-AC	-
2	HTK (Hattikuni)	25-50	10YR 4/6, 4/4 7.5YR 4/4, 3/3	sl	10-25	Ap-AC	-
3	BDL (Badiyala)	25-50	7.5YR2.5/3,2.5/ 3/3,10YR 3/4,4/3	sl	<15	Ap-Bw	e
4	VNK (Vanakanahalli)	25-50	2.5YR 3/4	sc	<15	Ap-Bt-Cr	-
5	YLR (Yalleri)	50-75	2.5YR 3/4,4/4 5YR 3/4 7.5 YR4/4	gc	15-35	Ap-Bt	-
6	HSL (Hosalli)	75-100	10YR 5/4,4/4,4/6	sc	<15	Ap-Bw	e
7	GWD (Gowdagera)	75-100	10YR 3/1,3/2,4/2	scl	<15	Ap-Bw	es
8	BLC (Balichakra)	75-100	2.5YR 5/3,2.5/4, 5YR 4/3, 3/3	scl	<15	Ap-Bt	-
9	YDR (Yadgir)	100-150	10YR 4/3,4/4	sl	<15	Ap-Ac	-
10	BGD (Belagundi)	100-150	10 YR 5/4,4/4 7.5YR 4/4	c	<15	Ap-Bss	es
11	ANR (Anur)	100-150	10YR 4/3,4/1	c	<15	Ap-Bw	es
12	MDG (Mundargi)	100-150	10YR 4/4,3/3 7.5YR 4/4	scl	<15	Ap-Bw	-
13	BMN (Bhimanahalli)	>150	10YR 3/1	c	<15	Ap-Bss	es
14	BMD (Bomaraladoddi)	>150	5YR 3/3,4/1,4/3,4/6	scl	-	Ap-Bt	e

3.4 Soil Mapping

The area under each soil series was further separated into soil phases and their boundaries delineated on the cadastral map based on the variations observed in the texture of the surface soil, slope, erosion, presence of gravel, stoniness etc. A soil phase is a subdivision of soil series based mostly on surface features that affect its use and management. The soil mapping units are shown on the map (Fig.3.5) in the form of symbols. During the survey many profile pits, few minipits and a few auger bores representing different landforms occurring in the microwatershed were studied. In addition to the profile study, spot observations in the form of minipits, road cuts, terrace cuts etc., were studied to validate the soil boundaries on the soil map.

The soil map shows the geographic distribution of 22 mapping units representing 14 soil series occurring in the microwatershed. The soil map unit (soil legend) description

is presented in Table 3.2. The soil phase map (management units) shows the distribution of 22 soil phases mapped in the microwatershed. Each mapping unit (soil phase) delineated on the map has similar soil and site characteristics. In other words, all the farms or survey numbers included in one phase will have similar management needs and have to be treated accordingly.

3.5 Land Management Units

The 22 soil phases identified and mapped in the microwatershed were grouped into 9 Land Management Units (LMU's) for the purpose of preparing a Proposed Crop Plan for sustained development of the microwatershed. The database (soil phases) generated under LRI was utilized for identifying Land Management Units (LMU's) based on the management needs. One or more than one soil site characteristic having influence on the management have been chosen for identification and delineation of LMUs. For Yagapur Tanda-1 microwatershed, five soil and site characteristics, namely soil depth, soil texture, slope, erosion and gravel content have been considered for defining LMUs. The land use classes are expected to behave similarly for a given level of management.

3.6 Laboratory Characterization

Soil samples were collected from representative master profiles for laboratory characterization by following the methods outlined in the Laboratory Manual (Sarma *et al*, 1987). Surface soil samples collected from farmer's fields for fertility status (major and micronutrients) at 320 m grid interval in the year 2018 were analyzed in the laboratory (Katyal and Rattan, 2003). By linking the soil fertility data to the survey numbers through GIS, soil fertility maps were generated by using Kriging method for the microwatershed.

Table 3.2 Soil map unit description of Yagapur Tanda-1 Microwatershed

*Soil map unit No.	Soil Series	Soil Phase	Mapping Unit Description	Area in ha (%)
Soils of Granite and granite gneiss Landscape				
	KKR		Kakalawar soils are very shallow (<25 cm), well drained, have dark brown sandy loam soils occurring on very gently sloping uplands under cultivation	10 (1.4)
153		KKRbB2g1	Loamy sand surface, slope 1-3%, moderate erosion, gravelly (15-35%)	2 (0.29)
175		KKRcB2	Sandy loam surface, slope 1-3%, moderate erosion	8 (1.11)
	HTK		Hattikuni soils are shallow (25-50 cm), well drained, have dark yellowish brown sandy loam soils occurring on very gently sloping uplands under cultivation	7 (0.91)
165		HTKcB2	Sandy loam surface, slope 1-3%, moderate erosion	7 (0.91)
	BDL		Badiyala soils are shallow (25-50 cm), well drained, have dark brown to very dark brown and dark yellowish brown,	71 (9.86)

*Soil map unit No.	Soil Series	Soil Phase	Mapping Unit Description	Area in ha (%)
			slightly calcareous sandy loam soils occurring on very gently to gently sloping uplands under cultivation	
174		BDLcB2g2	Sandy loam surface, slope 1-3%, moderate erosion, very gravelly (35-60%)	3 (0.47)
4		BDLhB2	Sandy clay loam surface, slope 1-3%, moderate erosion	20 (2.75)
162		BDLhB2g1	Sandy clay loam, slope 1-3%, moderate erosion, gravelly (15-35%)	26 (3.6)
5		BDLiB2	Sandy clay surface, slope 1-3%, moderate erosion	22 (3.04)
	VNK		Vanakanahalli soils are shallow (25-50 cm), well drained, have dark reddish brown, sandy clay red soils occurring on very gently to moderately sloping uplands under cultivation	109 (15.15)
8		VNKbB2g1	Loamy sand surface, slope 1-3%, moderate erosion, gravelly (15-35%)	53 (7.38)
10		VNKiB2	Sandy clay surface, slope 1-3%, moderate erosion	37 (5.16)
109		VNKmB2g1	Clay surface, slope 1-3%, moderate erosion, gravelly (15-35%)	19 (2.61)
	YLR		Yalleri soils are moderately shallow (50-75 cm), well drained, have brown to reddish brown and dark reddish brown, gravelly clay red soils occurring on very gently to gently sloping uplands under cultivation	63 (8.79)
31		YLRiB2	Sandy clay surface, slope 1-3%, moderate erosion	63 (8.79)
	HSL		Hosalli soils are moderately deep (75-100 cm), moderately well drained, have yellowish brown to dark yellowish brown, slightly calcareous sandy clay soils occurring on very gently sloping uplands under cultivation	4 (0.56)
111		HSLbB2	Loamy sand surface, slope 1-3%, moderate erosion	4 (0.56)
	GWD		Gowdagera soils are moderately deep (75-100 cm), moderately well drained, have dark grayish brown to very dark grayish brown, calcareous, sodic sandy clay loam soils occurring on very gently sloping uplands under cultivation	24 (3.3)
127		GWDmB2	Clay surface, slope 1-3%, moderate erosion	24 (3.3)
	BLC		Balichakra soils are moderately deep (75-100 cm), well drained, have reddish brown to dark reddish brown, sandy clay red soils occurring on very gently sloping uplands under cultivation	7 (0.92)
38		BLCiB2	Sandy clay surface, slope 1-3%, moderate erosion	7 (0.92)
	YDR		Yadgir soils are deep (100-150 cm), well drained, have brown to dark yellowish brown and olive brown, sodic	57 (7.93)

*Soil map unit No.	Soil Series	Soil Phase	Mapping Unit Description	Area in ha (%)
			sandy loam soils occurring on very gently sloping uplands under cultivation	
42		YDRcB2	Sandy loam surface, slope 1-3%, moderate erosion	41 (5.74)
43		YDRiB2	Sandy clay surface, slope 1-3%, moderate erosion	16 (2.19)
	BGD		Belagundi soils are deep (100-150 cm) well drained, have brown to dark yellowish brown, calcareous cracking clay soils occurring on very gently sloping uplands under cultivation	19 (2.62)
177		BGDia1	Sandy clay surface, slope 0-1%, slight erosion	19 (2.62)
	ANR		Anur soils are deep (100-150 cm), moderately well drained, have dark gray to dark brown, calcareous, sodic clay soils occurring on very gently to gently sloping uplands under cultivation	84 (11.6)
55		ANRiB2	Sandy clay surface, slope 1-3%, moderate erosion	84 (11.6)
	MDG		Mundargi soils are deep (100-150 cm), well drained, have brown to dark yellowish brown, sandy clay loam soils occurring on very gently sloping uplands under cultivation	6 (0.85)
58		MDGiB2	Sandy clay surface, slope 1-3%, moderate erosion	6 (0.85)
	BMN		Bhimanahalli soils are very deep (>150 cm), moderately well drained, have very dark gray, calcareous cracking clay black soils occurring on very gently sloping uplands under cultivation	22 (3.07)
62		BMNmB2	Clay surface, slope 1-3%, moderate erosion	22 (3.07)
	BMD		Bomraddoddi soils are very deep (>150 cm), well drained, have dark reddish brown to dark grey, reddish brown, dark brown and yellowish red, slightly calcareous sandy clay loam soils occurring on nearly level to very gently sloping uplands under cultivation.	16 (2.18)
64		BMDcB2	Sandy loam surface, slope 1-3%, moderate erosion	10 (1.37)
65		BMDiB2g1	Sandy clay surface, slope 1-3%, moderate erosion, gravelly (15-35%)	6 (0.81)
999		Rock outcrops	Rock lands, both massive and bouldery with little or no soil	199 (27.66)
1000		Others	Habitation	23 (3.21)




Fig 3.5 Soil Phase or Management Units - Yagapur Tanda-1 Microwatershed

THE SOILS

Detailed information pertaining to the nature, extent and their distribution of different kinds of soils occurring in Yagapur Tanda-1 microwatershed is provided in this chapter. The microwatershed area has been identified as granite gneiss landscape based on geology. In all, 14 soil series are identified. Soil formation is the result of the combined effect of environmental and terrain factors that are reflected in soil morphology. In the granite gneiss landscape, it is by parent material, relief and climate.

A brief description of each of the 14 soil series identified followed by 22 soil phases (management units) mapped under each series are furnished below. The physical and chemical characteristics of soil series identified in Yagapur Tanda-1 microwatershed are given in Table 4.1 along with soil classification. The soils in any one map unit differ from place to place in their depth, texture, slope, gravelliness, erosion or any other site characteristic that affect management. The soil phase map can be used for identifying the suitability of areas for growing specific crops or for other alternative uses and also for deciding the type of conservation structures needed. The detailed information on soil and site-characteristics like soil depth, surface soil texture, slope, erosion, gravelliness, AWC, LCC etc, with respect to each of the soil phase identified is given village/survey number wise for the microwatershed in Appendix-I.

4.1 Soils of granite gneiss landscape

In this landscape, 14 soil series are identified and mapped. VNK series occupies maximum area of 109 ha (15%) followed by ANR 84 ha (12%), BDL 71 ha (10%), YLR 63 ha (9%), YDR 53 ha (8%), GWD 24 ha (3%), BMN 22 ha (3%), BGD 19 ha (3%), BMD 16 ha (2%), KKR 10 ha (1%), BLC 7 ha (<1%), HTK 7 ha (<1%), MDG 6 ha (<1%) and HSL 4 ha (<1%). Brief description of each series identified and number of soil phases mapped is given below.

4.1.1 Kakalawar (KKR) Series: Kakalawar soils are very shallow (<25cm), well drained, have dark brown to light brown, sandy loam soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation. The Kakalawar series has been classified as a member of the mixed, isohyperthermic family of Lithic Ustipsamments.

The thickness of the soil is less than 25 cm. Its colour is in 10 YR and 7.5 YR hue with value 4 to 6 and chroma 3 to 4. The texture varies from loamy sand to sand. The available water capacity is very low (<50 mm/m). Two soil phase were identified and mapped.



Landscape and Soil Profile characteristics of Kakalawar (KKR) Series

4.1.2 Hattikuni (HTK) Series: Hattikuni soils are shallow (25-50 cm), well drained, have dark brown to dark yellowish brown sandy loam soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation. The Hattikuni series has been classified as a member of the mixed, isohyperthermic family of Lithic Ustipsamments.

The thickness of the soil ranges from 36 to 50 cm. The thickness of A horizon ranges from 8 to 12 cm. Its colour is in 10YR and 7.5 YR hue with value 3 to 4 and chroma 4 to 6. The texture varies from loamy sand to sandy loam. The thickness of subsurface horizon ranges from 28 to 42 cm. Its colour is in 10 YR and 7.5 YR hue with value 3 to 4 and chroma 4 to 6. Its texture varies from loamy sand to sand and sandy loam. The available water capacity is very low (<50 mm/m). One soil phase was identified and mapped.



Landscape and Soil Profile characteristics of Hattikuni (HTK) Series

4.1.3 Badiyala (BDL) Series: Badiyala soils are shallow (25-50 cm), well drained, have very dark brown to dark yellow brown and dark brown, slightly calcareous sandy loam soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation. The Badiyala series has been classified as a member of the coarse-loamy, mixed, isohyperthermic family of Fluventic Haplustepts.

The thickness of the solum ranges from 28 to 50 cm. The thickness of A horizon ranges from 4 to 12 cm. Its colour is in 10YR hue with value 3 to 4 and chroma 3 to 4. The texture is loamy sand, sandy clay loam and sandy clay. The thickness of B horizon ranges from 27 to 45 cm. Its colour is in 10 YR and 7.5 YR hue with value 2 to 4 and chroma 3 to 4. Its texture is sandy loam to sandy clay loam and is slightly calcareous. The available water capacity is very low (<50mm/m). Four soil phases were identified and mapped.



Landscape and Soil Profile characteristics of Badiyala (BDL) Series

4.1.4 Vanakanahalli (VNK) Series: Vanakanahalli soils are shallow (25-50 cm), well drained, have dark reddish brown sandy clay red soils. They have developed from weathered granite gneiss and occur on very gently sloping uplands under cultivation. The Vanakanahalli series has been classified as a member of the clayey, mixed isohyperthermic family of (Paralthic) Haplustalfs.

The thickness of the solum ranges from 25 to 49 cm. The thickness of A horizon ranges from 7 to 16 cm. Its colour is in 2.5 YR and 5 YR with value 3 and chroma 2 to 4. The texture is sandy loam to sandy clay loam and sandy clay. The thickness of B horizon ranges from 20 to 40 cm. Its colour is in 2.5 YR and 5 YR hue with value 3 to 4 and chroma 3 to 4. Its texture is sandy clay. The available water capacity is very low (<50 mm/m). Three soil phases were identified and mapped.



Landscape and Soil Profile characteristics of Vanakanahalli (VNK) Series

4.1.5 Yalleri (YLR) Series: Yalleri soils are moderately shallow (50-75 cm), well drained, have very dark reddish brown to dark brown, gravelly clay soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation. The Yalleri series has been classified as a member of the fine, mixed, isohyperthermic family of Typic Paleustalfs.

The thickness of the solum ranges from 50 to 74 cm. The thickness of A horizon ranges from 10 to 13 cm. Its colour is in 7.5 YR and 5 YR hue with value and chroma 2 to 4. The texture is sandy loam, loamy sand, and sandy clay loam. The thickness of B horizon ranges from 45 to 64 cm. Its colour is in 7.5 YR and 5 YR hue with value 2 to 4 and chroma 2 to 4. Its texture is clay with gravel content of 15-35 per cent. The available water capacity is low (51-100 mm/m). One soil phase was identified and mapped.



Landscape and Soil Profile characteristics of Yalleri (YLR) Series

4.1.6 Hosalli (HSL) Series: Hosalli soils are moderately deep (75-100 cm), moderately well drained, have dark yellowish brown to yellowish brown, slightly calcareous sandy clay soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation. The Hosalli series has been classified as a member of the fine, mixed, isohyperthermic family of Typic Haplustepts.

The thickness of the solum ranges from 76 to 100 cm. The thickness of A horizon ranges from 6 to 15 cm. Its colour is in hue 10 YR and 7.5 YR with value 3 to 5 and chroma 2 to 4. Its texture varies from loamy sand to sandy loam and sandy clay loam. The thickness of B horizon ranges from 62 to 93 cm. Its colour is in hue 10 YR with value 3 to 4 and chroma 2 to 4. Its texture varies from sandy clay loam to sandy clay and clay and is slightly calcareous. The available water capacity is medium (101-150 mm/m). One soil phase was identified and mapped.



Landscape and Soil Profile characteristics of Hosalli (HSL) Series

4.1.7 Gowdagera (GWD) Series: Gowdagera soils are moderately deep (75-100 cm), well drained, have very dark gray to dark grayish brown, calcareous sodic sandy clay loam soils. They are developed from weathered granite gneiss and occur on very gently sloping uplands under cultivation. The Gowdagera series has been classified as a member of the fine-loamy, mixed (calcareous), isohyperthermic family of Typic Haplustepts.

The thickness of the solum ranges from 76 to 100 cm. The thickness of A horizon ranges from 8 to 16 cm. Its colour is in hue 10 YR with value 3 to 4 and chroma 2 to 4. Its texture varies from sandy loam to sandy clay loam. The thickness of B horizon ranges from 61 to 91 cm. Its colour is in hue 10 YR with value 2 to 4 and chroma 1 to 4. Its texture is sandy clay loam to sandy clay and is calcareous sodic soils. The available water capacity is medium (101-150 mm/m). One soil phase was identified and mapped.



Landscape and Soil Profile characteristics of Gowdagera (GWD) Series

4.1.8 Balichakra (BLC) Series: Balichakra soils are moderately deep (75-100 cm), well drained, have dark reddish brown to reddish brown, sandy clay loam soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation. The Balichakra series has been classified as a member of the fine-loamy, mixed, isohyperthermic family of Typic Haplustalfs.

The thickness of the solum ranges from 80 to 100 cm. The thickness of A horizon ranges from 10 to 16 cm. Its colour is in hue 5 YR with value and chroma of 3 to 4. Its texture varies from sandy clay loam and sandy clay. The thickness of B horizon ranges from 70 to 88 cm. Its colour is in hue 2.5 YR and 5 YR with value 3 to 5 and chroma 3 to 4. Its texture is sandy clay loam to sandy clay. The available water capacity is medium (101-150 mm/m). One soil phase was identified and mapped.



Landscape and Soil Profile characteristics of Balichakra (BLC) Series

4.1.9 Yadgir (YDR) Series: Yadgir soils are deep (100-150 cm), well drained, have very dark yellowish brown to light olive brown, sodic sandy loam soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation. The Yadgir series has been classified as a member of the coarse-loamy, mixed, isohyperthermic family of Fuluventic Haplustepts.

The thickness of the soil ranges from 105 to 145 cm. The thickness of A horizon ranges from 6 to 10 cm. Its colour is in 10 YR hue with value 4 and chroma 3. The texture is loamy sand. The thickness of subsurface horizons ranges from 95 to 130 cm. Its colour is in 10 YR and 2.5 Y hue with value 4 to 5 and chroma 3 to 4. Texture is sandy loam and sandy clay loam and is sodic soils. The available water capacity is low (51-100 mm/m). Two soil phases was identified and mapped.



Landscape and Soil Profile characteristics of Yadgir (YDR) Series

4.1.10 Belagundi (BGD) Series: Belagundi soils are deep (100-150 cm), moderately well drained, have dark yellowish brown to yellowish brown and dark brown calcareous cracking clay soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation. The Belagundi series has been classified as a member of the very fine, smectitic (calcareous), isohyperthermic family of Typic Haplusterts.

The thickness of the solum ranges from 100 to 145 cm. The thickness of A horizon ranges from 5 to 12 cm. Its colour is in 10 YR and 5 YR hue with value 5 and chroma 2 to 4. The texture varies from sandy to loamy sand. The thickness of B horizon ranges from 95 to 135 cm. Its colour is in 10 YR and 7.5 YR hue with value 4 to 5 and chroma 4. Texture is sandy clay to clay and is slightly calcareous. The available water capacity is very high (>200 mm/m). One soil phase was identified and mapped.



Landscape and Soil Profile characteristics of Belagundi (BGD) Series

4.1.11 Anur (ANR) Series: Anur soils are deep (100-150 cm), moderately well drained, have dark gray to dark brown, calcareous sodic clay soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation. The Anur series has been classified as a member of the fine, mixed (calcareous), isohyperthermic family of Typic Haplustepts.

The thickness of the solum ranges from 102 to 148 cm. The thickness of A-horizon ranges from 9 to 17 cm. Its colour is in 10 YR hue with value 3 to 4 and chroma 2 to 4. The texture ranges from loamy sand to sandy clay loam and sandy clay and are calcareous. The thickness of B horizon ranges from 102 to 135 cm. Its colour is in 10 YR hue with value 3 to 5 and chroma 1 to 6. Texture is sandy clay loam to sandy clay and clay and is calcareous sodic soils. The available water capacity is very high (>200 mm/m). One soil phase was identified and mapped.



Landscape and Soil Profile characteristics of Anur (ANR) Series

4.1.12 Mundargi (MDG) Series: Mundargi soils are deep (100-150 cm), well drained, have dark brown to dark yellowish brown, sandy clay loam soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation. The Mundargi series has been classified as a member of the fine-loamy, mixed, isohyperthermic family of Fluventic Haplustepts.

The thickness of the solum ranges from 100 to 149 cm. The thickness of A horizon ranges from 8 to 20 cm. Its colour is in 10 YR hue with value 3 and chroma 1 to 4. The texture ranges from sandy loam to sandy clay loam and sandy clay. The thickness of B horizon ranges from 105 to 140 cm. Its colour is in 10 YR hue with value 3 to 4 and chroma 2 to 4. The texture varies from sandy loam to sandy clay loam and sandy clay. The available water capacity is very high (>200 mm/m). One soil phase was identified and mapped.



Landscape and Soil Profile characteristics of Mundargi (MDG) Series

4.1.13 Bhimanahalli (BMN) Series: Bhimanahalli soils are very deep (>150 cm), moderately well drained, have very dark gray, calcareous cracking clay soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation. The Bhimanahalli series has been classified as a member of the fine, smectitic (calcareous), isohyperthermic family of Typic Haplusterts.

The thickness of the solum is more than 150 cm. The thickness of A horizon ranges from 6 to 13 cm. Its colour is in 10 YR hue with value 3 to 4 and chroma 1 to 2 with clay texture. The thickness of B horizon ranges from 163 to 176 cm. Its colour is in 10 YR hue with value 2 to 4 and chroma 1. Its texture is clay and is calcareous. The available water capacity is very high (>200 mm/m). One soil phase was identified and mapped.



Landscape and Soil Profile characteristics of Bhimanahalli (BMN) Series

4.1.14 Bomraldoddi (BMD) Series: Bomraldoddi soils are very deep (>150 cm), well drained, have dark reddish brown to dark grey, reddish brown, dark brown and yellowish red, slightly calcareous sandy clay loam soils. They are developed from weathered granite gneiss and occur on nearly level to very gently sloping uplands under cultivation. The bomraldoddi series has been classified as a member of the fine loamy mixed, isohyperthermic family of Rhodic Paleustalfs.

The thickness of the solum is more than 150 cm. The thickness of A horizon ranges from 11 to 17 cm. Its colour is in 5 YR hue with value 3 to 4 and chroma 1 to 5. Texture varies from sandy loam to sandy clay loam. The thickness of B horizon is more than 150 cm. Its colour is in hue 5 YR with value 4 and chroma 1 to 6. Texture varies from sandy clay loam to sandy clay and is slightly calcareous. The available water capacity is high (151-200 mm/m). Two soil phases were identified and mapped.

Table: 4.1 Physical and Chemical Characteristics of Soil Series identified in Yagapur Tanda-1 microwatershed

Soil Series: Kakalawar (KKR), Pedon: R-7

Location: 16°50'25.9"N 77°15'97.1"E, Yampada village, Gurumitkal hobli, Yadgir taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bengaluru **Classification:** Mixed, isohyperthermic Lithic Ustipsamments

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-22	Ap	83.81	10.37	5.82	17.31	20.65	17.91	5.67	22.27	10-20	ls	9.77	4.65

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO ₃	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP						
	Water	CaCl ₂	M KCl				dS m ⁻¹	%	%	Ca	Mg					K	Na	Total	cmol kg ⁻¹	%	%
0-22	5.85	-	-	0.027	0.19	-	0.72	0.21	0.62	0.03	1.58	2.6	0.45	60.90	1.17						

Contd...

Soil Series: Hattikuni (HTK), Pedon: R-7

Location: 16°50'46.5"N 77°10'16.4"E, Yaddalli village, Hattikuni hobli, Yadgir taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bengaluru **Classification:** Mixed, isohyperthermic Lithic Ustipsamments

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-12	Ap	90.89	5.62	3.49	8.50	13.46	29.86	29.55	9.51	20	s	7.73	3.16
12-22	A1	89.97	6.53	3.50	7.19	13.48	29.48	29.79	10.03	20	s	8.00	3.05
22-45	A2	87.20	6.43	6.38	11.09	14.42	31.55	7.16	22.98	40	ls	7.67	3.96

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO ₃	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP			
	Water	CaCl ₂	M KCl				dS m ⁻¹	%	%	Ca	Mg					K	Na	Total
										cmol kg ⁻¹								
0-12	6.81	-	-	0.062	0.07	-	2.35	0.50	0.16	0.01	3.02	3.0	0.86	100	0.38			
12.0-22	6.80	-	-	0.050	0.21	-	1.67	0.30	0.09	0.01	2.07	2.4	0.69	86.30	0.45			
22-45	6.85	-	-	0.044	0.19	-	1.82	0.42	0.10	0.06	2.40	2.6	0.41	92.41	2.17			

Contd...

Soil Series: Badiyala (BDL) **Pedon:** R-5

Location: 16°37'10.0"N 77°20'21.5", Gudalagunta village, Balichakra hobli, Yadgir taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bengaluru **Classification:** Coarse-loamy, mixed, isohyperthermic Fluventic Haplustepts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-12	Ap	87.13	7.04	5.83	10.03	24.32	23.61	23.51	5.67	<15	ls	6.27	2.44
12-28	Bw	64.63	13.30	22.07	6.74	13.07	22.30	17.01	5.50	<15	scl	16.34	7.83
28-50	BC	73.11	12.02	14.87	3.93	16.03	26.89	18.41	7.86	<15	sl	12.94	5.47

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO ₃	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP			
	Water	CaCl ₂	M KCl				dS m ⁻¹	%	%	Ca	Mg					K	Na	Total
										cmol kg ⁻¹								
0-12	6.20	-	-	0.074	1.00	0.00	2.80	0.98	0.14	0.01	3.92	4.20	0.72	93	0.20			
12-28	9.04	-	-	0.253	0.80	3.20	-	-	0.16	0.69	-	16.90	0.77	100	4.09			
28-50	9.41	-	-	0.364	1.10	3.60	-	-	0.16	1.39	-	11.10	0.75	100	12.52			

Contd...

Soil Series: Vanakanahalli (VNK) **Pedon:** R-15

Location: 16°43'49.5"N 77°17'17.9"E, Yaleri village, Balichakra hobli, Yadgiri taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bengaluru **Classification:** Clayey, mixed isohyperthermic (Paralthic) Haplustalfs

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-18	Ap	82.61	8.09	9.30	6.77	8.59	21.13	34.58	11.53	-	ls	8.85	3.53
18-61	Bt	54.51	8.73	36.77	4.93	6.18	14.15	20.75	8.49	-	sc	18.88	11.63

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO ₃	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP			
	Water	CaCl ₂	M KCl				dS m ⁻¹	%	%	Ca	Mg					K	Na	Total
0-18	5.37	-	-	0.11	0.60	0.00	2.96	1.45	0.13	0.14	4.68	6.27	0.67	75	2.22			
18-61	4.71	-	-	0.05	0.81	0.00	5.56	2.24	0.10	0.05	7.95	13.31	0.36	60	0.38			

Contd...

Soil Series: Yalleri (YLR) **Pedon:** R-16

Location: 16°32'54.3"N 77°22'71.2"E, Duppalli village, Sydhapura hobli, Yadgir taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bengaluru **Classification:** Fine, mixed, isohyperthermic Typic Paleustalfs

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-5	Ap	81.69	5.44	12.87	6.10	8.65	33.88	21.57	11.50	-	sl	8.60	3.37
5-34	Bt1	38.78	6.73	54.49	3.38	9.91	12.42	8.93	4.14	-	c	25.33	15.82
34-75	Bt2	40.35	2.90	56.75	12.91	6.83	10.30	7.48	2.82	35-60	c	24.49	16.20

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO ₃	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP			
	Water	CaCl ₂	M KCl				dS m ⁻¹	%	%	Ca	Mg					K	Na	Total
										cmol kg ⁻¹								
0-5	6.91	-	-	0.069	0.70	0.00	5.29	1.37	0.28	0.03	6.96	6.90	0.54	100	0.45			
5-34	7.05	-	-	0.053	0.62	0.00	16.43	3.89	0.26	0.09	20.67	21.60	0.40	96	0.42			
34-75	7.25	-	-	0.058	0.59	0.00	15.22	3.46	0.25	0.14	19.06	19.90	0.35	96	0.69			

Contd...

Soil Series: Hosalli (HSL) **Pedon:** R-3

Location: 16°46'60.3"N 77°05'47.6"E, Mudhanala village, Yadgir hobli, Yadgir taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bengaluru **Classification:** Fine, mixed, isohyperthermic Typic Haplustepts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-10	Ap	88.43	5.15	6.42	5.69	6.40	36.04	27.31	12.99	-	s	7.40	2.74
10-30	Bw1	58.47	7.24	34.29	4.26	9.37	19.91	19.28	5.64	-	scl	19.07	11.57
30-50	Bw2	51.43	12.67	35.90	3.49	8.89	16.72	15.87	6.46	<15	sc	21.64	12.44
50-90	Bw3	49.89	13.64	36.47	2.43	2.96	20.61	16.17	7.72	<15	sc	21.12	12.95

Depth (cm)	pH (1:2.5)			E.C. (1:2.5) dS m ⁻¹	O.C. %	CaCO ₃ %	Exchangeable bases					CEC	CEC/Clay	Base saturation %	ESP %
	Water	CaCl ₂	M KCl				Ca	Mg	K	Na	Total				
							cmol kg ⁻¹								
0-10	7.16	-	-	0.117	0.48	0.00	2.83	1.50	0.15	0.29	4.76	4.90	0.76	97	5.94
10-30	6.91	-	-	0.040	0.36	0.00	10.64	5.43	0.10	0.26	16.43	17.80	0.52	92	1.47
30-50	8.17	-	-	0.182	0.24	1.43	-	-	0.12	0.22	-	19.90	0.55	100	1.08
50-90	8.60	-	-	0.148	0.20	4.29	-	-	0.13	0.16	-	19.70	0.54	100	0.81

Contd...

Soil Series: Gowdagera (GWD) **Pedon:** R-13

Location: 16°38'24.4"N 77°21'24.0"E, Madhawara village, Balichakara hobli, Yadgir taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bengaluru , **Classification:** Fine-loamy, mixed (calcareous), isohyperthermic Typic Haplustepts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-18	Ap	79.61	13.94	6.45	14.17	17.53	23.65	17.02	7.24	-	ls	11.36	3.86
18-42	BW1	69.09	10.58	21.06	10.54	16.58	22.01	14.43	5.53	-	scl	31.62	12.30
42-81	Bw2	51.37	13.51	35.60	7.59	10.55	16.24	11.60	5.38	-	sc	67.57	26.89

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO ₃	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP			
	Water	CaCl ₂	M KCl				dS m ⁻¹	%	%	Ca	Mg					K	Na	Total
										cmol kg ⁻¹								
0-18	9.89	-	-	0.74	0.66	1.20	-	-	0.18	3.63	-	8.35	1.29	100	17.40			
18-42	10.82	-	-	1.60	0.27	5.76	-	-	0.19	19.23	-	15.84	0.75	100	40.17			
42-81	10.83	-	-	2.30	0.27	7.80	-	-	0.40	26.71	-	26.54	0.75	100	40.27			

Contd...

Soil Series: Balichakra (BLC) **Pedon:** T1/P2

Location: 16°33'25.0"N 77°20'52.3"E, Sowrashtralli village, Sydhapura hobli, Yadgir taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bengaluru **Classification:** Fine-loamy, mixed, isohyperthermic Typic Haplustalfs

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-8	Ap	65.46	8.38	26.16	12.51	18.72	18.82	10.44	4.96	-	scl	15.15	8.63
8-19	BA	63.48	8.16	28.36	12.80	15.84	17.21	12.49	5.14	-	scl	16.45	8.81
19-40	Bt	52.64	11.58	35.79	13.19	13.19	14.35	8.23	3.69	-	sc	21.49	10.36
40-75	BC	55.14	10.71	34.15	14.10	14.42	14.63	7.53	4.45	-	scl	17.77	8.99

Depth (cm)	pH (1:2.5)			E.C. (1:2.5) dS m ⁻¹	O.C. %	CaCO ₃ %	Exchangeable bases					CEC	CEC/Clay	Base saturation %	ESP %
	Water	CaCl ₂	M KCl				Ca	Mg	K	Na	Total				
0-8	6.75	-	-	0.19	0.72	0.00	12.18	3.10	0.43	0.22	15.92	16.80	0.64	95	1.31
8-19	7.23	-	-	0.12	0.68	0.84	11.37	2.50	0.23	0.18	14.28	14.77	0.52	97	1.24
19-40	7.13	-	-	0.08	0.50	0.48	13.80	2.82	0.18	0.09	16.89	17.66	0.49	96	0.51
40-75	7.07	-	-	0.07	0.35	0.84	13.00	2.90	0.17	0.10	16.16	17.55	0.51	92	0.57

Contd...

Soil Series: Yadgir (YDR) **Pedon:** R-5

Location: 16°35'43.6"N 77°17'06.4"E, Kanikal village, Balichakra hobli, Yadgir taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bengaluru **Classification:** Coarse-loamy, mixed, isohyperthermic Fulventic Haplustepts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-14	Ap	73.39	11.31	15.30	6.76	20.27	24.87	15.66	5.83	-	sl	12.14	7.22
14-43	A2	86.59	8.77	4.64	23.19	26.92	14.11	15.22	7.16	-	ls	6.97	2.68
43-89	Bw1	80.41	3.75	15.84	8.06	13.47	36.73	15.71	6.43	-	sl	22.84	10.18
89-110	Bw2	63.55	5.40	31.05	8.10	23.05	19.00	9.87	3.53	15-35	scl	38.46	17.70

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO ₃	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP			
	Water	CaCl ₂	M KCl				dS m ⁻¹	%	%	Ca	Mg					K	Na	Total
0-14	9.47	-	-	0.371	0.32	1.30	14.71	4.28	0.38	1.54	20.91	12.70	0.83	165	4.86			
14-43	7.25	-	-	0.114	0.56	0.00	2.29	0.86	0.07	0.03	3.25	3.40	0.73	96	0.31			
43-89	10.30	-	-	0.820	0.16	0.52	1.70	0.98	0.15	6.62	9.45	8.61	0.54	110	30.77			
89-110	10.80	-	-	1.440	0.12	0.91	1.02	2.00	0.29	14.43	17.74	16.17	0.52	110	35.688			

Contd...

Soil Series: Belagundi (BGD) **Pedon:** T₁/P₂

Location: 16°31'65.3"N 77°20'84.9"E, Kadechoora village, Sydhapura hobli, Yadgir taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bengaluru, **Classification:** Very fine, smectitic (calcareous), isohyperthermic Typic Haplusterts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-13	Ap	14.90	17.83	67.27	0.77	2.10	2.65	5.96	3.42	-	c	43.97	29.27
13-40	AB	13.07	18.32	68.61	0.80	2.05	2.61	4.20	3.41	-	c	41.23	30.48
40-80	Bss1	11.68	17.18	71.13	0.80	2.06	2.29	3.32	3.21	-	c	46.72	32.41
80-113	Bss2	12.17	16.53	71.30	1.95	1.61	3.21	2.41	2.99	-	c	46.87	35.13

Depth (cm)	pH (1:2.5)			E.C. (1:2.5) dS m ⁻¹	O.C. %	CaCO ₃ %	Exchangeable bases					CEC	CEC/Clay	Base saturation %	ESP %
	Water	CaCl ₂	M KCl				Ca	Mg	K	Na	Total				
0-13	7.85	-	-	0.253	0.87	5.20	-	-	0.67	0.17	-	65.90	0.98	100	0.26
13-40	8.11	-	-	0.172	0.74	4.29	-	-	0.31	0.16	-	66.70	0.97	100	0.23
40-80	8.44	-	-	0.205	0.58	5.59	-	-	0.20	0.27	-	66.30	0.93	100	0.40
80-113	8.82	-	-	0.201	0.39	10.14	-	-	0.19	0.17	-	63.80	0.89	100	0.27

Contd...

Soil Series: Anur (ANR) **Pedon:** R-15

Location: 16°32'45.0"N 77°23'57.4"E, Duppalli village, Sydhapura hobli, Yadgir taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bengaluru **Classification:** Fine, mixed (calcareous), isohyperthermic Typic Haplustepts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-18	Ap	64.60	13.44	21.96	7.33	10.42	18.68	20.12	8.05	<15	scl	16.59	7.96
18-49	Bw1	56.66	12.19	31.15	4.73	9.80	18.66	17.02	6.45	-	scl	33.38	13.51
49-95	Bw2	39.94	17.81	42.25	3.09	3.30	15.44	10.65	7.45	<15	c	44.68	25.23
95-123	Bw3	30.65	17.58	51.77	1.50	5.57	10.18	9.65	3.75	<15	c	54.94	32.07

Depth (cm)	pH (1:2.5)			E.C. (1:2.5) dS m ⁻¹	O.C. %	CaCO ₃ %	Exchangeable bases					CEC	CEC/Clay	Base saturation %	ESP %
	Water	CaCl ₂	M KCl				Ca	Mg	K	Na	Total				
0-18	10.17	-	-	0.365	0.48	6.11	-	-	0.25	3.52	-	19.90	0.91	100	7.08
18-49	10.32	-	-	1.38	0.30	6.76	-	-	0.21	16.03	-	24.60	0.79	100	26.07
49-95	10.08	-	-	2.55	0.17	6.11	-	-	0.33	21.49	-	32.60	0.77	100	26.36
95-123	9.92	-	-	2.56	0.12	7.93	-	-	0.51	26.03	-	36.00	0.70	100	28.92

Contd...

Soil Series: Mundargi (MDG) **Pedon:** R-2

Location: 16°46'82.4"N 77°04'85.2"E, Thumakura village, Yadgir hobli, Yadgir taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bengaluru **Classification:** Fine-loamy, mixed, isohyperthermic Fluventic Haplustepts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-9	Ap	81.23	12.97	5.80	4.84	10.19	14.83	37.94	13.42	<15	ls	11.75	3.31
9-20	A2	76.82	16.19	6.98	4.96	10.12	20.75	27.53	13.46	-	ls	14.52	3.99
20-46	Bw1	42.43	17.43	40.15	2.26	5.59	11.49	14.93	8.16	-	c	34.90	21.14
46-90	Bw2	54.51	16.56	28.93	4.72	5.03	19.92	16.67	8.18	-	scl	36.73	18.88
90-110	Bw3	53.69	11.00	35.30	9.57	9.89	16.23	13.01	4.99	-	sc	38.72	20.53

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO ₃	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP						
	Water	CaCl ₂	M KCl				dS m ⁻¹	%	%	Ca	Mg					K	Na	Total	cmol kg ⁻¹	%	%
0-9	8.2	-	-	0.399	0.44	0.78	-	-	0.16	0.38	-	4.90	0.84	100	3.08						
9-20	8.44	-	-	0.075	0.29	1.82	-	-	0.05	0.35	-	4.90	0.70	100	2.88						
20-46	9.39	-	-	0.451	0.32	2.73	-	-	0.12	5.22	-	20.77	0.52	100	10.06						
46-90	9.75	-	-	0.616	0.24	3.25	-	-	0.12	5.72	-	16.56	0.57	100	13.82						
90-110	9.72	-	-	0.725	0.24	3.64	-	-	0.14	6.84	-	19.76	0.56	100	13.836						

Contd...

Soil Series: Bhimanahalli (BMN) **Pedon:** R-3

Location: 16°31'82.4"N 77°12'70.8"E, Bheemanahalli village, Sydhapura hobli, Yadgir taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bengaluru **Classification:** Fine, smectitic (calcareous), isohyperthermic Typic Haplusterts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-8	Ap	20.34	19.94	59.72	2.68	5.03	3.75	5.25	3.64	-	c	50.19	33.49
8-40	Bss1	19.61	22.76	57.62	1.94	2.59	5.28	4.96	4.85	-	c	43.22	29.05
40-70	Bss2	21.25	17.65	61.10	3.02	5.26	3.91	5.48	3.58	-	c	44.30	30.25
70-120	Bss3	19.08	22.29	58.63	1.75	5.04	3.84	5.15	3.29	-	c	43.26	30.31
120-170	Bss4	11.11	20.44	68.45	2.04	1.93	1.70	2.83	2.61	-	c	51.33	33.51

Depth (cm)	pH (1:2.5)			E.C. (1:2.5) dS m ⁻¹	O.C. %	CaCO ₃ %	Exchangeable bases					CEC	CEC/Clay	Base saturation %	ESP %
	Water	CaCl ₂	M KCl				Ca	Mg	K	Na	Total				
0-8	8.2	-	-	0.284	0.72	4.94	-	-	1.20	0.34	-	52.70	0.88	100	0.65
8-40	8.44	-	-	0.139	0.40	7.28	-	-	0.30	0.48	-	52.06	0.90	100	0.93
40-70	8.32	-	-	0.202	0.40	6.37	-	-	0.18	0.40	-	52.52	0.86	100	0.77
70-120	9.3	-	-	0.282	0.36	6.89	-	-	0.27	0.38	-	50.97	0.87	100	0.75
120-170	8.47	-	-	0.305	0.37	8.19	-	-	0.28	0.91	-	58.19	0.85	100	1.57

INTERPRETATION FOR LAND RESOURCE MANAGEMENT

The most important soil and site characteristics that affect the land use and conservation needs of an area are land capability, soil depth, soil texture, coarse fragments, available water capacity, soil slope, soil erosion, soil reaction etc. These are interpreted from the data base generated through land resource inventory and several thematic maps are generated. These would help in identifying the areas suitable for growing crops and, soil and water conservation measures and structures needed thus helping to maintain good soil health for sustained crop production. The various interpretative and thematic maps generated are described below.

5.1 Land Capability Classification

Land capability classification is an interpretative grouping of soil map units (soil phases) mainly based on inherent soil characteristics, external land features and environmental factors that limit the use of land for agriculture, pasture, forestry, or other uses on a sustained basis (IARI, 1971). The land and soil characteristics used to group the land resources in an area into various land capability classes, subclasses and units are

Soil Characteristics: Depth, texture, gravelliness, calcareousness.

Land characteristics: Slope, erosion, drainage, rock outcrops.

Climate: Total rainfall and its distribution, and length of crop growing period.

The Land capability classification system is divided into land capability classes, subclasses and units based on the level of information available. Eight land capability classes are recognized. They are

Class I: They are very good lands that have no limitations or very few limitations that restrict their use.

Class II: They are good lands that have minor limitations and require moderate conservation practices.

Class III: They are moderately good lands that have moderate limitations that reduce the choice of crops or that require special conservation practices.

Class IV: They are fairly good lands that have very severe limitations that reduce the choice of crops or that require very careful management.

Class V: Soils in these lands are not likely to erode, but have other limitations like wetness that are impractical to remove and as such not suitable for agriculture, but suitable for pasture or forestry with minor limitations.

Class VI: The lands have severe limitations that make them generally unsuitable for cultivation, but suitable for pasture or forestry with moderate limitations.

Class VII: The lands have very severe limitations that make them unsuitable for cultivation, but suitable for pasture or forestry with major limitations.

Class VIII: Soil and other miscellaneous areas (rock lands) that have very severe limitations that nearly preclude their use for any crop production, but suitable for wildlife, recreation and installation of wind mills.

The land capability subclasses are recognised based on the dominant limitations observed within a given land capability class. The subclasses are designated by adding a lower case letter like 'e', 'w', 's', or 'c' to the class numeral. The subclass "e" indicates that the main hazard is risk of erosion, "w" indicates drainage or wetness as a limitation for plant growth, "s" indicates shallow soil depth, coarse or heavy textures, calcareousness, salinity/alkalinity or gravelliness and "c" indicates limitation due to climate.

The land capability subclasses have been further subdivided into land capability units based on the kinds of limitations present in each subclass. Ten land capability units are used in grouping the soil map units. They are stony or rocky (0), erosion hazard (slope, erosion) (1), coarse texture (sand, loamy sand, sandy loam) (2), fine texture (cracking clay, silty clay) (3), slowly permeable subsoil (4), coarse underlying material (5), salinity/alkali (6), stagnation, overflow, high ground water table (7), soil depth (8) and fertility problems (9). The capability units thus identified have similar soil and land characteristics that respond similarly to a given level of management. The soils of the microwatershed have been classified upto land capability subclass level.

The 22 soil map units identified in the Yagapur Tanda-1 microwatershed are grouped under 3 land capability classes and 4 subclasses. An area of about 498 ha (69%) in the microwatershed is suitable for agriculture, about 199 ha (28%) covered by rock outcrops and about 23 ha (3%) covered by others in the microwatershed. (Fig. 5.1).




Fig. 5.1 Land Capability map of Yagapur Tanda-1 Microwatershed

Good lands (Class II) cover an area of 137 ha (19%) and are distributed in the northern, northeastern, northwestern, western, southern and eastern part of the microwatershed. They have minor limitations of soil and erosion. Moderately good lands (Class III) cover an area of 187 ha (26%) and are distributed in the major part of the cultivated area. They have moderate limitations of soil and erosion. Fairly good lands (Class IV) cover an area of about 174 ha (24%) and are distributed in the central, southern and eastern part of the microwatershed. They have very severe limitations of soil and erosion.

5.2 Soil Depth

Soil depth refers to the depth of the soil occurring above the parent material or hard rock. The depth of the soil determines the effective rooting depth for plants and in accordance with soil texture, mineralogy and gravel content, the capacity of the soil column to hold water and nutrient availability. Soil depth is one of the most important soil characteristic that is used in differentiating soils into different soil series. The soil depth classes used in identifying soils in the field are very shallow (<25 cm), shallow (25-50 cm), moderately shallow (50-75 cm), moderately deep (75-100 cm), deep (100-150 cm) and very deep (>150 cm). They were used to classify the soils into different depth classes and a soil depth map was generated. The area extent and their geographical distribution in the microwatershed is given in Fig. 5.2.




Fig. 5.2 Soil Depth map of Yagapur Tanda-1 Microwatershed

Very shallow (<25 cm) soils cover an area of 10 ha (1%) and are distributed in the eastern and southern part of the microwatershed. Shallow (25-50 cm) soils cover an area of 187 ha (26%) and are distributed in the major part of the microwatershed. Moderately shallow (50-75 cm) soils cover an area of 63 ha (9%) and are distributed in the northeastern and northern part of the microwatershed. Moderately deep (75-100 cm) soils cover an area of 34 ha (5%) and are distributed in the northwestern, western and southern part of the microwatershed. Deep (100-150 cm) soils cover an area of 166 ha (23%) and are distributed in central, eastern and northern part of the microwatershed. Very deep (>150 cm) soils cover an area of 38 ha (5%) and are distributed in eastern part of the microwatershed.

The most productive lands 204 ha (28%) with respect to soil rooting depth where all climatically adapted annual and perennial crops can be grown are deep to very deep (100 - >150 cm) soils. Problem soils cover about 197 ha (27%) area where short duration crops can be grown and probability of crop failure is high.

5.3 Surface Soil Texture

Texture is an expression to indicate the coarseness or fineness of the soil as determined by the relative proportion of primary particles of sand, silt and clay. It has a direct bearing on the structure, porosity, adhesion and consistence. The surface layer of a soil to a depth of about 25 cm is the layer that is most used by crops and plants. The surface soil textural class provides a guide to understanding soil-water retention and

availability, nutrient holding capacity, infiltration, workability, drainage, physical and chemical behaviour, microbial activity and crop suitability. The textural classes used for LRI were used to classify and a surface soil texture map was generated. The area extent and their geographical distribution in the microwatershed is shown in Figure 5.3.




Fig. 5.3 Surface Soil Texture map of Yagapur Tanda-1 Microwatershed

An area of 59 ha (8%) has soils that are sandy at the surface and occur in the eastern, western and southern part of the microwatershed. An area of 115 ha (16%) has soils that are loamy at the surface and occur in the eastern, western, northern and southern part of the microwatershed. An area of 324 ha (45%) has soils that are clayey at the surface and occur in the major part of the microwatershed.

An area of 439 ha (61%) in the microwatershed is most productive with respect to surface soil texture. The clayey soils (45%) have high potential for soil-water retention and availability, and nutrient retention and availability, but have more problems of drainage, infiltration, workability and other physical problems. The other productive lands are loamy soils (16%) which also have high potential for soil-water retention and nutrient availability but have no drainage or other physical problems. The sandy soils (8%) are problematic but productive for root and tuber crops, but these soils have the major limitation of moisture and nutrient retention capacity, hence frequent and shallow irrigation with balanced fertilizer application is to be followed in order to get better crop yields.

5.4 Soil Gravelliness

Gravel is the term used for describing coarse fragments between 2 mm and 7.5 cm diameter and stones for those between 7.5 cm and 25 cm. The presence of gravel and stones in soil reduces the volume of soil responsible for moisture and nutrient storage, drainage, infiltration and runoff, and hinders plant growth by impeding root growth and seedling emergence, intercultural operations and farm mechanization. The gravelliness classes used in LRI were used to classify the soils and using these classes a gravelliness map was generated. The area extent and their geographic distribution in the microwatershed is shown in Figure 5.4.




Fig. 5.4 Soil Gravelliness map of Yagapur Tanda-1 Microwatershed

An area of about 389 ha (54%) is non gravelly (<15%), and are distributed in the major part of the microwatershed. About 106 ha (15%) is gravelly (15-35%) soils, and are distributed in the central, eastern, western and southern part of the microwatershed and about 3 ha (<1%) is very gravelly (35-60%) soils, and are distributed in the eastern part of the microwatershed

The most productive soils (54%) that are non gravelly (<15%), where all climatically adapted long duration crops can be grown. Problem soils cover an area of (<1%) in the microwatershed.

5.5 Available Water Capacity

The soil available water capacity (AWC) is estimated based on the ability of the soil column to retain water between the tensions of 0.33 and 15 bar in a depth of 100 cm or the entire solum if the soil is shallower. The AWC of the soils (soil series) as estimated by considering the soil texture, mineralogy, soil depth and gravel content (Sehgal *et al.*, 1990) and accordingly the soil map units were grouped into five AWC classes *viz.*, very low (<50 mm/m), low (50-100 mm/m), medium (100-150 mm/m), high (150-200 mm/m) and very high (>200 mm/m) and using these values, an AWC map was generated. The area extent and their geographic distribution of different AWC classes in the microwatershed is given in Figure 5.5.




Fig. 5.5 Soil Available Water Capacity map of Yagapur Tanda-1 Microwatershed

An area of about 197 ha (27%) and 127 ha (18%) are very low (<50 mm) and low (51-100 mm/m) in available water capacity and are distributed in the major part of the microwatershed. About 43 ha (6%) is medium (101-150 mm/m) in available water capacity and are distributed in the western part of the microwatershed and about 131 ha (18%) is very high (>200 mm/m) in available water capacity and are distributed in the central, southern and northwestern part of the microwatershed.

An area of 324 ha (45%) in the microwatershed is problematic with regard to available water capacity. Here, only short duration crops can be grown and the probability of crop failure is very high. These areas are best put to other alternative uses. An area of

131 ha (18%) is potential, where all climatically adapted long duration crops can be grown.

5.6 Soil Slope

Soil slope refers to the inclination of the surface of the land. It is defined by gradient, shape and length, and is an integral feature of any soil as a natural body. Slope is considered important in soil genesis, land use and land development. The length and gradient of slope influences the rate of runoff, infiltration, erosion and deposition. The soil map units were grouped into two slope classes and a slope map was generated showing the area extent and their geographic distribution in the microwatershed (Fig. 5.6).

An area of about 479 ha (67%) falls under very gently sloping (1-3% slope) lands and are distributed in the major part of the microwatershed and about 19 ha (3%) falls under nearly level (0-1% slope) lands and are distributed in the northwestern part of the microwatershed.




Fig. 5.6 Soil Slope map of Yagapur Tanda-1 Microwatershed

Entire cultivated area of about 498 ha (69%) in the microwatershed is high potential in respect of soil slopes. In these areas, all climatically adapted annual and perennial crops can be grown without much soil and water conservation and other land development measures.

5.7 Soil Erosion

Soil erosion refers to the wearing away of the earth's surface by the forces of water, wind and ice involving detachment and transport of soil by raindrop impact. It is used for accelerated soil erosion resulting from disturbance of the natural landscape by burning, excessive grazing and indiscriminate felling of forest trees and tillage, all usually by man. The erosion classes showing an estimate of the current erosion status as judged from field observations in the form of rills, gullies or a carpet of gravel on the surface are recorded. Four erosion classes, viz, slight erosion (e1), moderate erosion (e2), severe erosion (e3) and very severe erosion (e4) are recognized. The soil map units were grouped into different erosion classes and a soil erosion map generated. The area extent and their spatial distribution in the microwatershed is given in Figure 5.7.

Soils that are slightly eroded (e1 class) cover an area of 19 ha (3%) and are distributed in the northern part of the microwatershed. Soils that are moderately eroded (e2 class) cover a maximum area of 479 ha (67%) and are distributed in the major part of the microwatershed.

Maximum area of about 479 ha (67%) in the microwatershed is problematic because of moderate erosion. For these areas, taking up soil and water conservation and other land development measures are needed.




Fig. 5.7 Soil Erosion map of Yagapur Tanda-1 Microwatershed

FERTILITY STATUS

Soil fertility plays an important role in increasing crop yield. The adoption of high yielding varieties that require high amounts of nutrients has resulted in deficiency symptoms in crops and plants due to imbalanced fertilization and poor inherent fertility status as these areas are characterised by low rainfall and high temperatures. Hence, it is necessary to know the fertility (macro and micro nutrients) status of the soils of the watersheds for assessing the kind and amount of fertilizers required for each of the crop intended to be grown. For this purpose, the surface soil samples collected from the grid points (one soil sample at every 320 m interval) all over the microwatershed through land resource inventory in the year 2018 were analysed for pH, EC, organic carbon, available phosphorus and potassium, and for micronutrients like zinc, boron, copper, iron and manganese, and secondary nutrient sulphur.

Soil fertility data generated has been assessed and individual maps for all the nutrients for the microwatershed have been generated using Kriging method under GIS. The village/survey number wise fertility data for the microwatershed is given in Appendix-II.

6.1 Soil Reaction (pH)

The soil analysis of the Yagapur Tanda-1 microwatershed for soil reaction (pH) showed that an area of about 25 ha (3%) is slightly to moderately acid (pH 5.5- 6.5) and are distributed in the northern and northwestern part of the microwatershed. An area of about 109 ha (12%) is neutral (6.5-7.3) and are distributed in the central, western, northeastern, northwestern part of the microwatershed. An area of about 220 ha (24%) is slightly alkaline (ph 7.3-7.8) and are distributed in the central, western, northern, northeastern and southern part of the microwatershed. An area of about 343 ha (37%) is moderately alkaline (ph 7.8-8.4) and are distributed in the major part of the microwatershed (fig.6.1). In all, major area of about 563 ha is alkaline, 109 ha is under neutral soils and 25 ha is under acidic.

6.2 Electrical Conductivity (EC)

The Electrical Conductivity of the soils in the microwatershed area is $<2 \text{ dS m}^{-1}$ (Fig 6.2) and as such the soils are non-saline.

6.3 Organic Carbon

The soil organic carbon content (an index of available Nitrogen) in the soils of the microwatershed is high ($>0.75\%$) in the entire cultivated area of the microwatershed. (Fig. 6.3).




Fig.6.1 Soil Reaction (pH) map of Yagapur Tanda-1 Microwatershed




Fig.6.2 Electrical Conductivity (EC) map of Yagapur Tanda-1 Microwatershed




Fig.6.3 Soil Organic Carbon map of Yagapur Tanda-1 Microwatershed

6.4 Available Phosphorus

Available phosphorus content is high (>57 kg/ha) covering an area of about 296 ha (41%) and occur in the major part of the microwatershed. Medium (23-57 kg/ha) in an area of about 157 ha (22%) and occur in the southern, eastern, northern and northeastern part of the microwatershed and low (<23 kg/ha) in an area of about 45 ha (6%) and occur in the northwestern part of the microwatershed (Fig. 6.4).

6.5 Available Potassium

Available potassium content is medium (145-337 kg/ha) in an area of 277 ha (38%) and are distributed in the major part of the microwatershed and high (>337 kg/ha) in an area of 221 ha (31%) and are distributed in the central, western, eastern, northeastern and northern and northwestern part of the microwatershed the microwatershed (Fig.6.5).

6.6 Available Sulphur

Available sulphur content is low (<10 ppm) in the entire cultivated area of the microwatershed. (Fig. 6.6).

6.7 Available Boron

Available boron content is medium (0.5-1.0 ppm) covering an area of 64 ha (9%) and are distributed in the eastern, northeastern and northern part of the microwatershed and about 434 ha (60%) is low (<0.5 ppm) in available boron and are distributed in the major part of the microwatershed (Fig. 6.7).

6.8 Available Iron

Available iron content is sufficient (>4.5 ppm) in entire cultivated area of the microwatershed (Fig 6.8).

6.9 Available Manganese

Available manganese content is sufficient (>1.0 ppm) in the entire cultivated area of the microwatershed (Fig 6.9).

6.10 Available Copper

Available copper content is sufficient (>0.2 ppm) in the entire cultivated area of the microwatershed (Fig 6.10).




Fig.6.4 Soil Available Phosphorus map of Yagapur Tanda-1 Microwatershed




Fig.6.5 Soil Available Potassium map of Yagapur Tanda-1 Microwatershed




Fig.6.6 Soil Available Sulphur map of Yagapur Tanda-1 Microwatershed




Fig.6.7 Soil Available Boron map of Yagapur Tanda-1 Microwatershed




Fig.6.8 Soil Available Iron map of Yagapur Tanda-1 Microwatershed




Fig.6.9 Soil Available Manganese map of Yagapur Tanda-1 Microwatershed




Fig.6.10 Soil Available Copper map of Yagapur Tanda-1 Microwatershed

6.11 Available Zinc

Available zinc content is deficient (<0.6 ppm) in the entire cultivated area of the microwatershed (Fig 6.11).




Fig.6.11 Soil Available Zinc map of Yagapur Tanda-1 Microwatershed

LAND SUITABILITY FOR MAJOR CROPS

The soil and land resource units (soil phases) of Yagapur Tanda-1 microwatershed were assessed for their suitability for growing food, fodder, fibre and other horticulture crops by following the procedure as outlined in FAO, 1976 and 1983. Crop requirements were developed for each of the crop from the available research data and also by referring to Naidu *et. al.* (2006) and Natarajan *et. al.* (2015). The soil and land characteristics were matched with the crop requirement to arrive at the crop suitability. The soil and land characteristics (Table 7.1) and crop requirement (Table 7.2 to 7.30) are given at the end of the chapter. In FAO land suitability classification, two orders are recognized. Order S-Suitable and Order N-Not suitable. The orders have classes, subclasses and units. Order S has three classes, Class S1-Highly Suitable, Class S2-Moderately Suitable and Class S3- Marginally Suitable. Order N has two classes, N1-Currently not Suitable and N2-Permanently not Suitable. There are no subclasses within the Class S1 as they will have very minor or no limitations for crop growth. Classes S2, S3 and N1 are divided into subclasses based on the kinds of limitations encountered. The limitations that affect crop production are 'c' for erratic rainfall and its distribution and length of growing period (LGP), 'e' for erosion hazard, 'r' for rooting condition, 't' for lighter or heavy texture, 'g' for gravelliness or stoniness, 'n' for nutrient availability, 'l' for topography, 'm' for moisture availability, 'w' for drainage and 'z' for calcareousness. These limitations are indicated as lower case letters to the Class symbol. For example, moderately suitable lands with the limitations of soil depth and erosion are designated as S2re. For the microwatershed, the soil mapping units were evaluated and classified up to subclass level.

Using the above criteria, the soil map units of the microwatershed were evaluated and land suitability maps for 29 major annual and perennial crops were generated. The detailed information on the kind of suitability of each of the soil phase for the crops assessed are given village/ survey number wise for the microwatershed in Appendix-IV.

7.1 Land Suitability for Sorghum (*Sorghum bicolor*)

Sorghum is one of the major food crop grown in Karnataka in an area of 10.47 lakh ha in Bijapur, Gulbarga, Raichur, Bidar, Belgaum, Dharwad, Bellary, Chitradurga, Mysore and Tumakuru districts. The crop requirements for growing sorghum (Table 7.2) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing sorghum was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.1.

Highly (Class S1) suitable lands for growing sorghum occur in an area of 23 ha (3%) and are distributed in the western and northwestern part of the microwatershed. An area of about 113 ha (16%) is moderately suitable (Class S2) for growing sorghum and

are distributed in the northern, northeastern, northwestern, eastern and southern part of the microwatershed. They have minor limitations of nutrient availability, rooting depth, texture, gravelliness and calcareousness. About 352 ha (49%) is marginally suitable (Class S3) for growing sorghum and are distributed in the major part of the microwatershed with moderate limitations of rooting depth, texture, nutrient availability and calcareousness. About 10 ha (1%) is currently not suitable (Class N1) for growing sorghum and are distributed in the southern and eastern part of the microwatershed with severe limitation of rooting depth.




Fig. 7.1 Land Suitability map of Sorghum

7.2 Land Suitability for Maize (*Zea mays*)

Maize is one of the most important food crop grown in an area of 13.37 lakh ha in almost all the districts of the State. The crop requirements for growing maize (Table 7.3) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing maize was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.2.

An area of about 20 ha (3%) is highly suitable (Class S1) for growing maize and are distributed in the western and northwestern part of the microwatershed. An area of about 116 ha (16%) is moderately suitable (Class S2) for growing maize and are distributed in the northern, northwestern, northeastern, southern and eastern part of the microwatershed. They have minor limitations of nutrient availability, rooting depth, texture and calcareousness. About 352 ha (49%) is marginally suitable (Class S3) for

growing maize and are distributed in the major part of the microwatershed with moderate limitations of rooting depth, texture, nutrient availability and calcareousness. About 10 ha (1%) is currently not suitable (Class N1) for growing maize and is distributed in the southeastern and eastern part of the microwatershed with severe limitation of rooting depth.




Fig. 7.2 Land Suitability map of Maize

7.3 Land Suitability for Bajra (*Pennisetum glaucum*)

Bajra is one of the most important millet crop grown in an area of 2.34 lakh ha in the northern districts of Karnataka state. The crop requirements for growing bajra (Table 7.4) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing bajra was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.3.

An area of about 26 ha (4%) is highly suitable (Class S1) for growing bajra and are distributed in the western and northwestern part of the microwatershed. An area of about 110 ha (15%) is moderately suitable (Class S2) for growing bajra and are distributed in the northern, northwestern, northeastern, southern and southeastern part of the microwatershed. They have minor limitations of nutrient availability, rooting depth, texture and calcareousness. About 352 ha (49%) is marginally suitable (Class S3) for growing bajra and are distributed in the major part of the microwatershed with moderate limitations of rooting depth, texture, nutrient availability and calcareousness. About 10 ha (1%) is currently not suitable (Class N1) for growing bajra and is distributed in the

southeastern and eastern part of the microwatershed with severe limitation of rooting depth.




Fig. 7.3 Land Suitability map of Bajra

7.4 Land Suitability for Groundnut (*Arachis hypogaea*)

Groundnut is one of the major oilseed crop grown in an area of 6.54 lakh ha in Karnataka in most of the districts either as rainfed or irrigated crop. The crop requirements for growing groundnut (Table 7.5) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing groundnut was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.4.

An area of about 22 ha (3%) is highly suitable (Class S1) for growing groundnut and are distributed in the western and northwestern part of the microwatershed. An area of about 4 ha (<1%) is moderately suitable (Class S2) for growing bajra and are distributed in the western part of the microwatershed. They have minor limitation of texture. About 291 ha (41%) is marginally suitable (Class S3) for growing bajra and are distributed in the major part of the microwatershed with moderate limitations of rooting depth, texture, nutrient availability and calcareousness. About 174 ha (24%) is currently not suitable (Class N1) for growing groundnut and is distributed in the southeastern, southern and central part of the microwatershed with severe limitation of rooting depth.




Fig. 7.4 Land Suitability map of Groundnut

7.5 Land Suitability for Sunflower (*Helianthus annus*)

Sunflower is one of the most important oilseed crop grown in an area of 3.56 lakh ha in the State in all the districts. The crop requirements for growing sunflower (Table 7.6) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing sunflower was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.5.

Highly (Class S1) suitable lands for growing sunflower occur in an area of 19 ha (3%) and are distributed in the northwestern part of the microwatershed. An area of about 49 ha (7%) is moderately suitable (Class S2) for growing sunflower and are distributed in the southeastern, southwestern, western and northwestern part of the microwatershed. They have minor limitations of rooting depth, texture, gravelliness and calcareousness. An area of about 69 ha (10%) is marginally suitable (Class S3) for growing sunflower and are distributed in the northeastern, northern, northwestern and southwestern part of the microwatershed. They have moderate limitations of nutrient availability and rooting depth. About 361 ha (50%) is currently not suitable (Class N1) for growing sunflower and are distributed in major part of the microwatershed with severe limitations of nutrient availability and rooting depth.




Fig. 7.5 Land Suitability map of Sunflower

7.6 Land Suitability for Red gram (*Cajanus Cajan*)

Redgram is one of the most important pulse crop grown in an area of 7.28 lakh ha in almost all the districts of the State. The crop requirements for growing red gram (Table 7.7) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing redgram was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.6.

An area of about 74 ha (10%) is moderately suitable (Class S2) for redgram and are distributed in the northwestern, western, southwestern and southeastern part of the microwatershed. They have minor limitations of texture, calcareousness, graveliness, rooting depth and nutrient availability. An area of about 228 ha (32%) is marginally suitable (Class S3) for growing redgram and are distributed in the major part of the microwatershed. They have moderate limitations of nutrient availability, rooting depth and calcareousness. About 197 ha (27%) is currently not suitable (Class N1) for growing redgram and are distributed in the central, northern, eastern, western, southern part of the microwatershed with severe limitation of rooting depth.




Fig. 7.6 Land Suitability map of Redgram

7.7 Land Suitability for Bengal gram (*Cicer aerativum*)

Bengal gram is one of the most important pulse crop grown in about 9.39 lakh ha area in Bijapur, Raichur, Kalaburgi, Dharwad, Belgaum and Bellary districts. The crop requirements for growing Bengal gram (Table 7.8) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing Bengal gram was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.7.

Highly (Class S1) suitable lands for growing bengalgram occur in an area of 19 ha (3%) and are distributed in the northwestern part of the microwatershed. An area of about 22 ha (3%) is moderately suitable (Class S2) for growing bengalgram and are distributed in the southeastern part of the microwatershed. They have minor limitation of calcareousness. An area of about 313 ha (43%) is marginally suitable (Class S3) for growing bengalgram and are distributed in the major part of the microwatershed. They have moderate limitations of nutrient availability, texture, calcareousness and rooting depth. About 145 ha (20%) is currently not suitable (Class N1) for growing bengalgram and are distributed in eastern, western, northern and southeastern part of the microwatershed with severe limitations of texture and rooting depth.




Fig. 7.7 Land Suitability map of Bengal gram.

7.8 Land Suitability for Cotton (*Gossypium hirsutum*)

Cotton is one of the most important fibre crop grown in the State in about 8.75 lakh ha area in Raichur, Dharwad, Belgaum, Kalaburgi, Bijapur, Bidar, Bellary, Chitradurga and Chamarajnagar districts. The crop requirements for growing cotton (Table 7.9) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing cotton was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.8.

Highly (Class S1) suitable lands for growing cotton occur in an area of 19 ha (3%) and are distributed in the northwestern part of the microwatershed. An area of about 89 ha (12%) is moderately suitable (Class S2) for growing cotton and are distributed in the southeastern, northeastern, northwestern, northern and western part of the microwatershed. They have minor limitations of rooting depth and calcareousness. An area of about 245 ha (34%) is marginally suitable (Class S3) for growing cotton and are distributed in the major part of the microwatershed. They have moderate limitations of nutrient availability, texture, calcareousness and rooting depth. About 145 ha (20%) is currently not suitable (Class N1) for growing cotton and are distributed in eastern, western, northern and southeastern part of the microwatershed with severe limitations of texture and rooting depth.




Fig. 7.8 Land Suitability map of Cotton

7.9 Land Suitability for Chilli (*Capsicum annum*)

Chilli is one of the most important spice crop grown in about 0.42 lakh ha in Karnataka State. The crop requirements for growing chilli (Table 7.10) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing chilli was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.9.

Highly (Class S1) suitable lands for growing chilli occur in an area of 20 ha (3%) and are distributed in the western, northwestern and southwestern part of the microwatershed. An area of about 110 ha (15%) is moderately suitable (Class S2) for growing chilli and are distributed in the southeastern, northeastern, northern and northwestern part of the microwatershed. They have minor limitations of texture, rooting depth, gravelliness and calcareousness. About 193 ha (27%) is marginally suitable (Class S3) for growing chilli and are distributed in the major part of the microwatershed with moderate limitations of nutrient availability and rooting depth. About 174 ha (24%) is currently not suitable (Class N1) for growing chilli and are distributed in the central, southwestern, southeastern and southern part of the microwatershed with severe limitations of nutrient availability and rooting depth.




Fig 7.9 Land Suitability map of Chilli

7.10 Land Suitability for Tomato (*Lycopersicon esculentum*)

Tomato is one of the most important vegetable crop grown in about 0.61 lakh ha covering almost all the district of the state. The crop requirements for growing tomato (Table 7.11) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing tomato was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.10.

Highly (Class S1) suitable lands for growing tomato occur in an area of 20 ha (3%) and are distributed in the western, northwestern and southwestern part of the microwatershed. An area of about 69 ha (10%) is moderately suitable (Class S2) for growing tomato and are distributed in the northeastern, northern and northwestern part of the microwatershed. They have minor limitations of rooting depth and gravelliness. About 234 ha (32%) is marginally suitable (Class S3) for growing tomato and are distributed in the major part of the microwatershed with moderate limitations of nutrient availability, texture and rooting depth. About 174 ha (24%) is currently not suitable (Class N1) for growing tomato and are distributed in the central, southwestern, southeastern and southern part of the microwatershed with severe limitations of nutrient availability and rooting depth.




Fig 7.10 Land Suitability map of Tomato

7.11 Land Suitability for Brinjal (*Solanum melongena*)

Brinjal is one of the most important vegetable crop grown in the state. The crop requirements for growing brinjal (Table 7.12) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing Brinjal was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.11.

Highly (Class S1) suitable lands for growing brinjal occur in an area of 20 ha (3%) and are distributed in the western, northwestern and southwestern part of the microwatershed. An area of about 69 ha (10%) is moderately suitable (Class S2) for growing brinjal and are distributed in the northeastern, northern and northwestern part of the microwatershed. They have minor limitations of rooting depth and gravelliness. About 234 ha (32%) is marginally suitable (Class S3) for growing brinjal and are distributed in the major part of the microwatershed with moderate limitations of nutrient availability, texture and rooting depth. About 174 ha (24%) is currently not suitable (Class N1) for growing brinjal and are distributed in the central, southwestern, southeastern and southern part of the microwatershed with severe limitations of nutrient availability and rooting depth.




Fig 7.11 Land Suitability map of Brinjal

7.12 Land Suitability for Onion (*Allium cepa* L.,)

Onion is one of the most important vegetable crop grown in the state. The crop requirements for growing onion (Table 7.13) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing onion was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.12.

Highly (Class S1) suitable lands for growing onion occur in an area of 20 ha (3%) and are distributed in the western, northwestern and southwestern part of the microwatershed. An area of about 69 ha (10%) is moderately suitable (Class S2) for growing onion and are distributed in the northeastern, northern and northwestern part of the microwatershed. They have minor limitations of rooting depth and gravelliness. About 228 ha (32%) is marginally suitable (Class S3) for growing onion and are distributed in the major part of the microwatershed with moderate limitations of texture and rooting depth. About 181 ha (25%) is currently not suitable (Class N1) for growing onion and are distributed in the central, southeastern and southern part of the microwatershed with severe limitations of nutrient availability and rooting depth.




Fig 7.12 Land Suitability map of Onion

7.13 Land Suitability for Bhendi (*Abelmoschus esculentus*)

Bhendi is one of the most important vegetable crop grown in the state. The crop requirements for growing bhendi (Table 7.14) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing bhendi was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.13.

Highly (Class S1) suitable lands for growing bhendi occur in an area of 20 ha (3%) and are distributed in the western, northwestern and southwestern part of the microwatershed. An area of about 110 ha (15%) is moderately suitable (Class S2) for growing bhendi and are distributed in the southeastern, northeastern, northern and northwestern part of the microwatershed. They have minor limitations of texture, rooting depth, gravelliness and calcareousness. About 193 ha (27%) is marginally suitable (Class S3) for growing bhendi and are distributed in the major part of the microwatershed with moderate limitations of nutrient availability and rooting depth. About 174 ha (24%) is currently not suitable (Class N1) for growing bhendi and are distributed in the central, southwestern, southeastern and southern part of the microwatershed with severe limitations of nutrient availability and rooting depth.




Fig 7.13 Land Suitability map of Bhendi

7.14 Land Suitability for Drumstick (*Moringa oleifera*)

Drumstick is one of the most important vegetable crop grown in about 2403 ha in the state. The crop requirements for growing drumstick (Table 7.15) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing drumstick was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.14.

Highly (Class S1) suitable lands for growing drumstick occur in an area of 16 ha (2%) and are distributed in the western, northwestern and southwestern part of the microwatershed. An area of about 30 ha (4%) is moderately suitable (Class S2) for growing drumstick and are distributed in the northwestern, western and southwestern part of the microwatershed. They have minor limitations of texture and rooting depth. About 85 ha (12%) is marginally suitable (Class S3) for growing drumstick and are distributed in the southeastern, northeastern, northern and northwestern part of the microwatershed with moderate limitations of calcareousness and rooting depth. About 368 ha (51%) is currently not suitable (Class N1) for growing drumstick and are distributed in the major part of the microwatershed with severe limitations of nutrient availability and rooting depth.




Fig 7.14 Land Suitability map of Drumstick

7.15 Land Suitability for Mango (*Mangifera indica*)

Mango is one of the most important fruit crop grown in an area of 1.73 lakh ha in almost all the districts of the State. The crop requirements (Table 7.16) for growing mango were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing mango was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed is given in Figure 7.15.

Highly (Class S1) suitable lands for growing mango occur in an area of 16 ha (2%) and are distributed in the western part of the microwatershed. About 58 ha (8%) is marginally suitable (Class S3) for growing mango and are distributed in the southeastern, western, northwestern and southwestern part of the microwatershed with moderate limitations of texture, nutrient availability and rooting depth. About 424 ha (59%) is currently not suitable (Class N1) for growing mango and are distributed in the major part of the microwatershed with severe limitations of nutrient availability and rooting depth.




Fig. 7.15 Land Suitability map of Mango

7.16 Land Suitability for Guava (*Psidium guajava*)

Guava is one of the most important fruit crop grown in an area of 0.06 lakh ha in almost all the districts of the State. The crop requirements (Table 7.17) for growing guava were matched with the soil-site characteristics (7.1) and a land suitability map for growing guava was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.16.

Highly (Class S1) suitable lands for growing guava occur in an area of 16 ha (2%) and are distributed in the western part of the microwatershed. An area of about 11 ha (1%) is moderately suitable (Class S2) for growing guava and are distributed in the northwestern, western and southwestern part of the microwatershed. They have minor limitation of rooting depth. About 104 ha (14%) is marginally suitable (Class S3) for growing drumstick and are distributed in the southeastern, northeastern, northern and northwestern part of the microwatershed with moderate limitations of calcareousness, texture and rooting depth. About 368 ha (51%) is currently not suitable (Class N1) for growing guava and are distributed in the major part of the microwatershed with severe limitations of nutrient availability and rooting depth.




Fig. 7.16 Land Suitability map of Guava

7.17 Land Suitability for Sapota (*Manilkara zapota*)

Sapota is one of the most important fruit crop grown in an area of 29373 ha in almost all the districts of the State. The crop requirements (Table 7.18) for growing sapota were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing sapota was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.17.

Highly (Class S1) suitable lands for growing sapota occur in an area of 16 ha (2%) and are distributed in the western part of the microwatershed. An area of about 11 ha (1%) is moderately suitable (Class S2) for growing sapota and are distributed in the northwestern, western and southwestern part of the microwatershed. They have minor limitation of rooting depth. About 110 ha (15%) is marginally suitable (Class S3) for growing sapota and are distributed in the southeastern, northeastern, northern and northwestern part of the microwatershed with moderate limitations of nutrient availability, texture and rooting depth. About 316 ha (50%) is currently not suitable (Class N1) for growing sapota and are distributed in the major part of the microwatershed with severe limitations of nutrient availability and rooting depth.




Fig. 7.17 Land Suitability map of Sapota

7.18 Land Suitability for Pomegranate (*Punica granatum*)

Pomegranate is one of the most important fruit crop commercially grown in about 18488 ha in Karnataka, mainly in Bijapur, Bagalkot, Koppal, Gadag and Chitradurga districts. The crop requirements for growing pomegranate (Table 7.19) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing pomegranate was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.18.

Highly (Class S1) suitable lands for growing pomegranate occur in an area of 16 ha (2%) and are distributed in the western part of the microwatershed. An area of about 52 ha (7%) is moderately suitable (Class S2) for growing pomegranate and is distributed in the northwestern, western, southeastern and southwestern part of the microwatershed. They have minor limitations of rooting depth, texture and calcareousness. About 69 ha (10%) is marginally suitable (Class S3) for growing pomegranate and are distributed in the northeastern, northern, southwestern and northwestern part of the microwatershed with moderate limitations of calcareousness, texture and rooting depth. About 361 ha (50%) is currently not suitable (Class N1) for growing pomegranate and are distributed in the major part of the microwatershed with severe limitations of nutrient availability and rooting depth.




Fig 7.18 Land Suitability map of Pomegranate

7.19 Land Suitability for Musambi (*Citrus limetta*)

Musambi is one of the important fruit crop grown in an area of 3446 ha in almost all the districts of the State. The crop requirements for growing musambi (Table 7.20) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing musambi was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.19.

Highly (Class S1) suitable lands for growing musambi occur in an area of 35 ha (3%) and are distributed in the western and southwestern part of the microwatershed. An area of about 33 ha (5%) is moderately suitable (Class S2) for growing musambi and are distributed in the southeastern, western and northwestern part of the microwatershed. They have minor limitations of rooting depth and calcareousness. An area of about 69 ha (10%) is marginally suitable (Class S3) for growing musambi and are distributed in the northeastern, northern, northwestern and southwestern part of the microwatershed. They have moderate limitations of rooting depth and nutrient availability. About 361 ha (50%) is currently not suitable (Class N1) for growing musambi and are distributed in the major part of the microwatershed with severe limitations of nutrient availability and rooting depth.




Fig. 7.19 Land Suitability map of Musambi

7.20 Land Suitability for Lime (*Citrus sp*)

Lime is one of the most important fruit crop grown in an area of 0.11 lakh ha in almost all the districts of the State. The crop requirements for growing lime (Table 7.21) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing lime was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7. 20.

Highly (Class S1) suitable lands for growing lime occur in an area of 35 ha (3%) and are distributed in the western and southwestern part of the microwatershed. An area of about 33 ha (5%) is moderately suitable (Class S2) for growing lime and are distributed in the southeastern, western and northwestern part of the microwatershed. They have minor limitations of rooting depth and calcareousness. An area of about 69 ha (10%) is marginally suitable (Class S3) for growing lime and are distributed in the northeastern, northern, northwestern and southwestern part of the microwatershed. They have moderate limitations of rooting depth and nutrient availability. About 361 ha (50%) is currently not suitable (Class N1) for growing lime and are distributed in the major part of the microwatershed with severe limitations of nutrient availability and rooting depth.




Fig. 7.20 Land Suitability map of Lime

7.21 Land Suitability for Amla (*Phyllanthus emblica*)

Amla is one of the medicinal fruit crop grown in almost all the districts of the State. The crop requirements for growing amla (Table 7.22) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing amla was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.21.

Highly (Class S1) suitable lands for growing amla occur in an area of 26 ha (4%) and are distributed in the western and northwestern part of the microwatershed. An area of about 82 ha (11%) is moderately suitable (Class S2) for growing amla and are distributed in the northeastern, northern and northwestern part of the microwatershed. They have minor limitation of rooting depth, calcareousness and texture. About 209 ha (29%) is marginally suitable (Class S3) for growing amla and are distributed in the major part of the microwatershed with moderate limitations of rooting depth, texture and calcareousness. About 181 ha (25%) is currently not suitable (Class N1) for growing amla and are distributed in the central, southern, southwestern and southeastern and eastern part of the microwatershed with severe limitations of rooting depth and nutrient availability.




Fig. 7.21 Land Suitability map of Amla

7.22 Land Suitability for Cashew (*Anacardium occidentale*)

Cashew is one of the most important plantation nut crop grown in an area of 0.7 lakh ha in almost all the districts of the State. The crop requirements for growing cashew (Table 7.23) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing cashew was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.22.

An area of about 23 ha (3%) is moderately suitable (Class S2) for growing cashew and are distributed in the northwestern part of the microwatershed. They have minor limitation of rooting depth and nutrient availability. About 67 ha (9%) is marginally suitable (Class S3) for growing cashew and are distributed in the northeastern, northern, western and northwestern part of the microwatershed with moderate limitations of rooting depth and nutrient availability. About 409 ha (57%) is currently not suitable (Class N1) for growing cashew and are distributed in the major part of the microwatershed with severe limitations of rooting depth and nutrient availability.




Fig. 7.22 Land Suitability map of Cashew

7. 23 Land Suitability for Jackfruit (*Artocarpus heterophyllus*)

Jackfruit is one of the most important fruit crop grown in an area of 5368 ha in almost all the districts of the State. The crop requirements for growing jackfruit (Table 7.24) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing jackfruit was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.23.

Highly (Class S1) suitable lands for growing jackfruit occur in an area of 16 ha (2%) and are distributed in the western part of the microwatershed. An area of about 11 ha (1%) is moderately suitable (Class S2) for growing jackfruit and are distributed in the western and northwestern part of the microwatershed. They have minor limitation of rooting depth. About 104 ha (14%) is marginally suitable (Class S3) for growing jackfruit and are distributed in the northeastern, northern, northwestern and southeastern part of the microwatershed with moderate limitations of rooting depth, texture and calcareousness. About 368 ha (51%) is currently not suitable (Class N1) for growing jackfruit and are distributed in the major part of the microwatershed with severe limitations of nutrient availability and rooting depth.




Fig. 7.23 Land Suitability map of Jackfruit

7.24 Land Suitability for Jamun (*Syzygium cumini*)

Jamun is an important fruit crop grown in almost all the districts of the State. The crop requirements for growing jamun (Table 25) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing jamun was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.24.

Highly (Class S1) suitable lands for growing jamun occur in an area of 16 ha (2%) and are distributed in the western part of the microwatershed. An area of about 19 ha (3%) is moderately suitable (Class S2) for growing jamun and are distributed in the northwestern part of the microwatershed. They have minor limitations of rooting depth and texture. About 96 ha (13%) is marginally suitable (Class S3) for growing jamun and are distributed in the northeastern, western, northern, northwestern and southeastern part of the microwatershed with moderate limitations of rooting depth and calcareousness. About 368 ha (51%) is currently not suitable (Class N1) for growing jamun and are distributed in the major part of the microwatershed with severe limitations of nutrient availability and rooting depth.




Fig. 7.24 Land Suitability map of Jamun

7.25 Land Suitability for Custard Apple (*Annona reticulata*)

Custard apple is one of the most important fruit crop grown in almost all the districts of the State. The crop requirements for growing custard apple (Table 7.26) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing custard apple was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.25.

Highly (Class S1) suitable lands for growing custard apple occur in an area of 41 ha (6%) and are distributed in the northwestern part of the microwatershed. An area of about 89 ha (12%) is moderately suitable (Class S2) for growing custard apple and are distributed in the southeastern, northeastern, northern, northwestern, western part of the microwatershed. They have minor limitations of rooting depth and calcareousness. About 193 ha (27%) is marginally suitable (Class S3) for growing custard apple and are distributed in the major part of the microwatershed with moderate limitations of nutrient availability, rooting depth and texture. About 174 ha (24%) is currently not suitable (Class N1) for growing custard apple and are distributed in the central, eastern, northern, southern and northwestern part of the microwatershed with severe limitations of nutrient availability and rooting depth.




Fig. 7.25 Land Suitability map of Custard Apple

7.26 Land Suitability for Tamarind (*Tamarindus indica*)

Tamarind is one of the most important spice crop grown in almost all the districts of the state. The crop requirements for growing tamarind (Table 7.27) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing tamarind was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Fig. 7.26.

Highly (Class S1) suitable lands for growing tamarind occur in an area of 16 ha (2%) and are distributed in the western part of the microwatershed. An area of about 19 ha (3%) is moderately suitable (Class S2) for growing tamarind and are distributed in the northwestern part of the microwatershed. They have minor limitations of rooting depth and texture. About 33 ha (5%) is marginally suitable (Class S3) for growing tamarind and are distributed in the southeastern, northwestern and western part of the microwatershed with moderate limitations of nutrient availability and rooting depth. About 431 ha (60%) is currently not suitable (Class N1) for growing tamarind and are distributed in the major part of the microwatershed with severe limitations of nutrient availability and rooting depth.




Fig. 7.26 Land Suitability map of Tamarind

7.27 Land Suitability for Mulberry (*Morus nigra*)

Mulberry is the important leaf crop grown for rearing of silkworms in about 1.6 lakh ha area in all the districts of the state. The crop requirements for growing mulberry (Table 7.28) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing mulberry was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.27.

Highly (Class S1) suitable lands for growing mulberry occur in an area of 16 ha (2%) and are distributed in the northwestern part of the microwatershed. An area of about 11 ha (1%) is moderately suitable (Class S2) for growing mulberry and is distributed in the western and northwestern part of the microwatershed. They have minor limitation of rooting depth. About 104 ha (14%) is marginally suitable (Class S3) for growing mulberry and are distributed in the northern, northwestern, northeastern and southeastern part of the microwatershed with moderate limitations of texture, calcareousness and rooting depth. About 368 ha (51%) is currently not suitable (Class N1) for growing mulberry and are distributed in the major part of the microwatershed with severe limitations of nutrient availability and rooting depth.




Fig 7.27 Land Suitability map of Mulberry

7.28 Land Suitability for Marigold (*Tagetes sps.*)

Marigold is one of the most important flower crop grown in an area of 9108 ha in almost all the districts of the State. The crop requirements (Table 7.29) for growing marigold were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing marigold was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.28.

Highly (Class S1) suitable lands for growing marigold occur in an area of 20 ha (3%) and are distributed in the western, northwestern and southwestern part of the microwatershed. An area of about 110 ha (15%) is moderately suitable (Class S2) for growing marigold and are distributed in the southeastern, northeastern, northern and northwestern part of the microwatershed. They have minor limitations of texture, rooting depth, gravelliness and calcareousness. About 193 ha (27%) is marginally suitable (Class S3) for growing marigold and are distributed in the major part of the microwatershed with moderate limitations of nutrient availability and rooting depth. About 174 ha (24%) is currently not suitable (Class N1) for growing marigold and are distributed in the central, southwestern, southeastern and southern part of the microwatershed with severe limitations of nutrient availability and rooting depth.




Fig. 7.28 Land Suitability map of Marigold

7.29 Land Suitability for Chrysanthemum (*Dendranthema grandiflora*)

Chrysanthemum is one of the most important flower crop grown in an area of 4978 ha in almost all the districts of the State. The crop requirements (Table 7.30) for growing chrysanthemum were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing chrysanthemum was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.29.

Highly (Class S1) suitable lands for growing chrysanthemum occur in an area of 20 ha (3%) and are distributed in the western, northwestern and southwestern part of the microwatershed. An area of about 110 ha (15%) is moderately suitable (Class S2) for growing chrysanthemum and are distributed in the southeastern, northeastern, northern and northwestern part of the microwatershed. They have minor limitations of texture, rooting depth, gravelliness and calcareousness. About 193 ha (27%) is marginally suitable (Class S3) for growing chrysanthemum and are distributed in the major part of the microwatershed with moderate limitations of nutrient availability and rooting depth. About 174 ha (24%) is currently not suitable (Class N1) for growing chrysanthemum and are distributed in the central, southwestern, southeastern and southern part of the microwatershed with severe limitations of nutrient availability and rooting depth.




Fig. 7.29 Land Suitability map of Chrysanthemum

Table 7.1 Soil-Site Characteristics of Yagapur Tanda-1 Microwatershed

Soil Map Units	Climate (P) (mm)	Growing period (Days)	Drainage Class	Soil depth (cm)	Soil texture		Gravelliness		AWC (mm/m)	Slope (%)	Erosion	pH	EC (dSm ⁻¹)	ESP (%)	CEC [Cmol (p ⁺)kg ⁻¹]	BS (%)
					Surface	Sub-surface	Surface (%)	Sub-surface (%)								
KKRbB2g1	866	150	WD	<25	ls	sl	15-35	10-15	<50	1-3	moderate	5.85	0.027	1.17	2.6	60.90
KKRcB2	866	150	WD	<25	sl	sl	<15	10-15	<50	1-3	moderate	5.85	0.027	1.17	2.6	60.90
HTKcB2	866	150	WD	25-50	sl	sl	<15	10-25	<50	1-3	moderate	6.81	0.062	0.38	3.0	100
BDLcB2g2	866	150	WD	25-50	sl	sl	35-60	<15	<50	1-3	moderate	6.20	0.074	0.20	4.20	93
BDLhB2	866	150	WD	25-50	scl	sl	<15	<15	<50	1-3	moderate	6.20	0.074	0.20	4.20	93
BDLhB2g1	866	150	WD	25-50	scl	sl	15-35	<15	<50	1-3	moderate	6.20	0.074	0.20	4.20	93
BDLiB2	866	150	WD	25-50	sc	sl	<15	<15	<50	1-3	moderate	6.20	0.074	0.20	4.20	93
VNKbB2g1	866	150	WD	25-50	ls	sc	15-35	<15	<50	1-3	moderate	5.37	0.11	2.22	6.27	75
VNKiB2	866	150	WD	25-50	sc	sc	<15	<15	<50	1-3	moderate	5.37	0.11	2.22	6.27	75
VNKmB2g1	866	150	WD	25-50	c	sc	15-35	<15	<50	1-3	moderate	5.37	0.11	2.22	6.27	75
YLRiB2	866	150	WD	50-75	sc	c	<15	15-35	51-100	1-3	moderate	6.91	0.069	0.45	6.90	100
HSLbB2	866	150	MWD	75-100	ls	sc	<15	<15	101-150	1-3	moderate	7.16	0.117	5.94	4.90	97
GWDmB2	866	150	MWD	75-100	c	scl	<15	<15	101-150	1-3	moderate	9.89	0.74	17.40	8.35	100
BLCiB2	866	150	W	75-100	sc	scl	<15	<15	101-150	1-3	moderate	6.75	0.19	1.31	16.80	95
YDRcB2	866	150	WD	100-150	sl	sl	15-35	<15	51-100	1-3	moderate	9.47	0.371	4.86	12.70	165
YDRiB2	866	150	WD	100-150	sc	sl	15-35	<15	51-100	1-3	moderate	9.47	0.371	4.86	12.70	165
BGDia1	866	150	MWD	100-150	sc	c	<15	<15	>200	0-1	slight	7.85	0.253	0.26	65.90	100
ANRiB2	866	150	MWD	100-150	sc	c	<15	<15	>200	1-3	moderate	10.17	0.365	7.08	19.90	100
MDGiB2	866	150	WD	100-150	sc	scl	<15	<15	>200	1-3	moderate	8.20	0.399	3.08	4.90	100
BMNmB2	866	150	MWD	>150	c	c	<15	<15	>200	1-3	moderate	8.20	0.284	0.65	52.70	100
BMDcB2	866	150	WD	>150	sl	scl	<15	-	151-200	1-3	moderate	-	-	-	-	-
BMDiB2g1	866	150	WD	>150	sc	scl	15-35	-	151-200	1-3	moderate	-	-	-	-	-

*Symbols and abbreviations are according to Field Guide for LRI under Sujala-III Project, Karnataka

Table 7.2 Land suitability criteria for Sorghum

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	26–30	30–34; 24–26	34–40; 20–24	>40; <20
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, c (red), c (black)	scl, cl	ls, sl	-
	pH	1:2.5	5.5-7.8	5.0-5.5 7.8-9.0	>9.0	-
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	10-15
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

Table 7.3 Land suitability criteria for Maize

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	30-34	35-38 26-30	38-40 26-20	
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	Mm				
	Rainfall in growing season	Mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc	c (red), c (black)	ls, sl	-
	pH	1:2.5	5.5-7.8	5.0-5.5 7.8-9.0	>9.0	-
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	Cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	-
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

Table 7.4 Land suitability criteria for Bajra

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-32	33-38 24-27	39-40 20-23	<20
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm	500-750	400-500	200-400	<200
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl, scl, cl,sc,c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.8	5.0-5.5 7.8-9.0	5.5-6.0 >9.0	
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	15-35	35-60	>60	
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	1-3	3-5	5-10	>10

Table 7.5 Land suitability criteria for Groundnut

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	24–33	22–24; 33–35	20–22; 35–40	<20; >40
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	Mm				
	Rainfall in growing season	Mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. Well drained	Poorly drained	Very Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl	sl,cl, sc	c (red), c (black), ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	Cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<35	35-60	>60	
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.6 Land suitability criteria for Sunflower

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	24–30	30–34; 20–24	34–38; 16–20	>38; <16
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	mod. Well drained	-	Poorly to very drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	cl, sc,c (red), c (black)	scl	ls, sl	-
	pH	1:2.5	6.5-7.8	7.8-8.4 5.5-6.5	8.4-9.0; 5.0-5.5	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.7 Land suitability criteria for Redgram

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	30-35(G) 20-25(AV) 15-18 (F&PS) 35-40(M)	25-30(G) 20-25 (AV) 12-15 (F&PS) 30-35(M)	20-25(G) 15-20(AV) 10-12 (F&PS) 25-30(M)	< 20 <15 <10 <25
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	Mm				
	Rainfall in growing season	Mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. Well drained	Poorly drained	Very Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, c (red)	c (black),sl, scl, cl	ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-9.0	5.0-5.5 >9.0	-
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	Cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-50	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<1.0	1.0-2.0	>2.0	
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.8 Land suitability criteria for Bengal gram

Land use requirement		Rating				
Soil –site characteristics	Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)	
Climatic regime	Mean temperature in growing season	°C	20–25	25–30; 15–20	30–35; 10–15	>35; <10
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. Well drained	Poorly drained	Very Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	c(black)	-	c (red), scl, cl, sc	ls, sl
	pH	1:2.5	6.0-7.8	5.0-6.0 7.8-9.0	>9.0	-
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
Rooting conditions	OC	%				
	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
Soil toxicity	Coarse fragments	Vol %	<15	15-35	35-60	60-80
	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
Erosion hazard	Sodicity (ESP)	%	5-10	10-15	>15	-
	Slope	%	<3	3-5	5-10	>10

Table 7.9 Land suitability criteria for Cotton

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	22-32	>32	<19	-
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well to moderately well	Poorly drained/Some what excessively drained	-	very poorly/excessively drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, c (red,black)	cl	scl	ls, sl
	pH	1:2.5	6.5-7.8	7.8-8.4	5.5-6.5 8.4->9.0	<5.5
	CEC	C mol (p+)Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	50-100	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	<3	3-5	-	>5

Table 7.10 Land suitability criteria for Chili

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	25-32	33-35 20-25	35-38 <20	>38
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc	c (black), sl	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.11 Land suitability criteria for Tomato

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	25-28	29-32 20-24	15-19 33-36	<15 >36
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl, scl, cl, sc, c (red)	-	ls, c(black)	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.12 Land suitability criteria for Brinjal

Land use requirement		Rating				
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	Well drained	Moderately well drained	Poorly drained	V. Poorly drained
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class				
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl, scl, cl, sc c (red)	-	ls, c (black)	-
	pH	1:2.5	6.0-7.3	7.3-8.4 5.0-6.0	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	>60
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.13 Land suitability criteria for Onion

Land use requirement		Rating				
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	20-30	30-35	35-40	>40
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately /imperfectly	-	Poorly to V poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl,scl,cl,sc,c (red)	-	c (Black),ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<1.0	1.0-2.0	2.0-4.0	<4
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.14 Land suitability criteria for Bhendi

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	25-28	29-32 20-24	15-19 33-36	<15 >36
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Imperfectly drained	Poorly to very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl,sc, c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.15 Land suitability criteria for Drumstick

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, scl, cl, c (red)	sl, c (black)	ls	S
	pH	1:2.5	6.0-7.3	5.0-5.5 7.3-7.8	5.5-6.0 7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<35	35-60	60-80	>80
Soil toxicity	Salinity (EC saturation extract)	ds/m				
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-10	-	>10

Table 7.16 Land suitability criteria for Mango

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-32	24-27 33-35	36-40	20-24
	Min temp. before flowering	°C	10-15	15-22	>22	-
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration	Days				
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V. Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	-	ls, sl, c (black)	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>150	100-150	75-100	<75
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.17 Land suitability criteria for Guava

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-32	33-36 24-27	37-42 20-23	
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	sl	c (black), ls	-
	pH	1:2.5	6.0-7.8	5.0-6.0	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.18 Land suitability criteria for Sapota

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-32	33-36 24-27	37-42 20-23	>42 <18
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	-	Poorly to very drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	sl	ls, c (black)	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.19 Land suitability criteria for Pomegranate

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	30-34	35-38 25-29	39-40 15-24	
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl,cl, sc, c (red)	c (black),sl	ls	-
	pH	1:2.5	5.5-7.8	7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.20 Land suitability criteria for Musambi

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-30	31-35 24-27	36-40 20-23	>40 <20
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately drained	poorly	Very poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c	sl	ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.21 Land suitability criteria for Lime

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-30	31-35 24-27	36-40 20-23	>40 <20
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately drained	poorly	Very poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c	sl	ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.22 Land suitability criteria for Amla

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. well drained	Poorly drained	V. Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	c (black)	ls, sl	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15-35	35-60	60-80	-
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

Table 7.23 Land suitability criteria for Cashew

Land use requirement		Rating				
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	32 to 34	28 to 32; 34 to 38	24 to 28; 38 to 40	<20; >40
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	-	sl, ls	c (black)
	pH	1:2.5	5.5-6.5	5.0-5.5 6.5-7.3	7.3-7.8	>7.8
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-10	>10	-

Table 7.24 Land suitability criteria for Jackfruit

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. well	Poorly	V. Poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	-	sl, ls, c (black)	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	>60
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10-

Table 7.25 Land suitability criteria for Jamun

Land use requirement		Rating				
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well	Mod. well	Poorly	V.Poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c(red)	sl, c (black)	ls	-
	pH	1:2.5	6.0-7.8	5.0-6.0	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>150	100-150	50-100	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	>60
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

Table 7.26 Land suitability criteria for Custard apple

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	Scl, cl, sc, c (red), c (black)	-	Sl, ls	-
	pH	1:2.5	6.0-7.3	5.5-6.0 7.3-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15-35	35-60	60-80	-
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	>5	-

Table 7.27 Land suitability criteria for Tamarind

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod.well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl,sc, c (red)	sl, c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>150	100-150	75-100	<75
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

Table 7.28 Land suitability criteria for Mulberry

Land use requirement		Rating				
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	24–28	22–24; 28–32	32–38; 22–18	>38; <18
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V. Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, cl, scl	c (red)	c (black), sl, ls	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.8-8.4	7.3-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	0-35	35-60	60-80	>80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

Table 7.29 Land suitability criteria for Marigold

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	18-23	17-15 24-35	35-40 10-14	>40 <10
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl,scl, cl, sc, c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%				
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.30 Land suitability criteria for Chrysanthemum

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	18-23	17-15 24-35	35-40 10-14	>40 <10
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl,scl, cl, sc, c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%				
Erosion hazard	Slope	%	<3	3-5	5-10	>10

7.30 Land Management Units (LMUs)

The 22 soil map units identified in Yagapur Tanda-1 microwatershed have been grouped into 9 Land Management Units (LMUs) for the purpose of preparing a Proposed Crop Plan. Land Management Units are grouped based on the similarities in respect of the type of soil, the depth of the soil, the surface soil texture, gravel content, AWC, slope, erosion etc. and a Land Management Units map (Fig. 7.30) has been generated. These Land Management Units are expected to behave similarly for a given level of management.




Fig. 7.30 Land Management Units Map Yagapur Tanda-1 Microwatershed

The map units that have been grouped into 9 Land Management Units along with brief description of soil and site characteristics are given below.

LMU	Soil map units	Soil and site characteristics
1	64.BMDcB2 65.BMDiB2g1	Very deep, sandy clay loam soils (>150cm), 1-3 % slopes, non-gravelly to gravelly (<15-35%), moderate erosion.
2	62.BMNmB2 177.BGDIA1 111.HSLbB2	Moderately deep to very deep, black calcareous to non calcareous clay soils (75 - >150 cm), 1-3 % slopes, non-gravelly (<15%), slight to moderate erosion.
3	55.ANRiB2 127.GWDmB2	Moderately deep to deep, sodic soils (75- 150 cm), 1- 3% slopes, non- gravelly (<15%), moderate erosion.
4	58.MDGiB2	Deep, sandy clay loam and strongly alkaline soils (75->150 cm), 1-3 % slopes, non-gravelly (<15%), moderate erosion.
5	42.YDRcB2 43.YDRiB2	Deep sodic soils (100- 150 cm), 1-3 % slopes, non-gravelly (<15%), moderate erosion
6	38.BLCiB2	Moderately deep, red sandy clay soils (75- 100 cm), 1-3 % slopes, non-gravelly (<15%), moderate erosion
7	31.YLRiB2	Moderately shallow, red clay soils (50- 75 cm), 1-3 % slopes, non-gravelly (<15%), moderate erosion.
8	174.BDLcB2g2 4.BDLhB2 162.BDLhB2g1 5.BDLiB2 165.HTKcB2 8.VNKbB2g1 10.VNKiB2 109.VNKmB2g1	Shallow, soils (25- 50 cm), 1-3 % slopes, non-gravelly to gravelly (<15-35%), moderate erosion.
9	153.KKRbB2g1 175.KKRcB2	Very shallow, soils (<25 cm), 1-3 % slopes, non-gravelly to gravelly (<15-35%), moderate erosion.

7.31 Proposed Crop Plan for Yagapur Tanda-1 Microwatershed

After assessing the land suitability for the 29 crops, the Proposed Crop Plan has been prepared for the 9 identified LMUs by considering only the highly (Class S1) and moderately (Class S2) suitable lands for each of the 29 crops. The resultant proposed crop plan is presented below in Table 7.31.

Table 7.31 Proposed Crop Plan for Yagapur Tanda-1 Microwatershed

LMU	Soil Map Units	Survey Number	Field Crops/ Commercial crops	Horticulture Crops (Rainfed/Irrigated)	Suitable Interventions
1	64.BMDcB2 65.BMDiB2g1	Yagapura : 84,85	Sunflower, Sorghum, Maize, Groundnut, Red gram, Bajra	Fruit crops: Musambi, Sapota, Tamarind, Pomegranate, Amla, Custard apple, Guava, Jackfruit, Jamun, Lime Vegetables: Tomato, Onion, Bhendi, Chilli, Brinjal, Drumstick, Coriander Flowers: Marigold, Chrysanthemum	Application of FYM, Biofertilizers and micronutrients, drip irrigation, mulching, suitable soil and water conservation practices
2	62.BMNmB2 177.BGDia1 111.HSLbB2	Shivanagara: 14,15,16, 17, 40 Yagapura: 29,3,67,71,8 9,208,209,212,218,219, 220,221,222, 223,224	Maize, Sorghum, Sunflower, Cotton, Red gram, Bengalgram, Bajra	Fruit crops: Lime, Musambi, Custard apple, Pomegranate Vegetables: Chilli, Bhendi Flowers: Marigold, Chrysanthemum	Application of FYM, Biofertilizers and micronutrients, drip irrigation, mulching, suitable soil and water conservation practices
3	55.ANRiB2 127.GWDmB2	Yagapura: 44,45,46,49, 50,53,54,55,56,59,74,75 ,76,77,78,79,80,81,82,8 3,184,187,188,189,198, 199,200,201,202,203,20 5,206,207,211	-	Agri-Silvi-Pasture: Ber, Aonla, Acacia sp. Dhaincha, Rhodes grass, Para grass ,Bermuda grass	Application of gypsum, iron pyrites and elemental sulphur. Addition of farm yard manure, green manures and providing subsurface drainage
4	58.MDGiB2	Yagapura: 68,168,169,1 71,178, 181,182	Sorghum, Maize, Bajra	Agri-Silvi-Pasture: Ber, Aonla, Acacia sp. Dhaincha, Rhodes grass, Para grass ,Bermuda grass	Application of FYM, Biofertilizers and micronutrients, drip irrigation, mulching, suitable soil and water conservation practices
5	42.YDRcB2 43.YDRiB2	Yagapura: ,28,40,41,42, 43,47,48,179,180,185,1 86,190,191	-	Agri-Silvi-Pasture: Ber, Aonla, Acacia sp. Dhaincha, Rhodes grass, Para grass ,Bermuda grass	Application of gypsum, iron pyrites and elemental sulphur. Addition of farm yard manure, green

LMU	Soil Map Units	Survey Number	Field Crops/ Commercial crops	Horticulture Crops (Rainfed/Irrigated)	Suitable Interventions
					manures and providing subsurface drainage
6	38.BLCiB2	Shivanagara :5,6,13,41	Sunflower, Sorghum, Maize, Groundnut, Red gram, Bajra	Fruit crops: Mango, Musambi, Sapota, Tamarind, Pomegranate, Amla, Custard apple, Guava, Jackfruit, Jamun, Lime Vegetables: Tomato, Onion, Bhendi, Chilli, Brinjal, Drumstick, Coriander Flowers: Marigold, Chrysanthemum	Application of FYM, Biofertilizers and micronutrients, drip irrigation, mulching, suitable soil and water conservation practices
7	31.YLRiB2	Shivanagara:3,4,18,19,2,20,22,23,28,38,54,57,58,59,60	Maize, Sorghum, Cotton, Bajra	Fruit crops: Amla, Custard apple Vegetables: Tomato, Onion, Bhendi, Chilli, Brinjal Flowers: Marigold, Chrysanthemum	Drip irrigation, mulching, suitable soil and water conservation practices (Crescent Bunding with Catch Pit etc)
8	174.BDLcB2g2 4.BDLhB2 162.BDLhB2g1 5.BDLiB2 165.HTKcB2 8.VNKbB2g1 10.VNKiB2 109.VNKmB2g1	Shivanagara : 39 Yagapura:30,31,32,33,34,35,36,38,39,51,52,57,58,60,61,62,63,64,65,66,72,73,86,88,167,183,193,194,195,197,210,214,215,216	-	Agri-Silvi-Pasture: Custard apple, Hybrid Napier, <i>Styloxanthes hamata</i> , <i>Glyricidia</i> , <i>Styloxanthes scabra</i>	Use of short duration varieties, sowing across the slope and split application of nitrogen fertilizers
9	153.KKRbB2g1 175.KKRcB2	Yagapura : 2,213,217	-	Hybrid Napier, <i>Styloxanthes hamata</i> , <i>Styloxanthes scabra</i>	Use of short duration varieties, sowing across the slope, drip irrigation is recommended

SOIL HEALTH MANAGEMENT

8.1 Soil Health

Soil health is basic to plant health and plant health is basic to human and bovine health. Soil is fundamental to crop production. Without soil, no food could be produced nor would livestock be fed on a large scale. Because it is finite and fragile, soil is a precious resource that requires special care from its users.

Soil health or the capacity of the soil to function is critical to human survival. Soil health has been defined as: “the capacity of the soil to function as a living system without adverse effect on the ecosystem”. Healthy soils maintain a diverse community of soil organisms that help to form beneficial symbiotic associations with plant roots, recycle essential plant nutrients, improve soil structure with positive repercussions for soil, water and nutrient holding capacity and ultimately improve crop production and also contribute to mitigating climate change by maintaining or increasing its carbon content.

Functional interactions of soil biota with organic and inorganic components, air and water determine a soil’s potential to store and release nutrients and water to plants and to promote and sustain plant growth. Thus, maintaining soil health is vital to crop production and conserve soil resource base for sustaining agriculture.

The most important characteristics of a healthy soil are

- Good soil tilth
- Sufficient soil depth
- Good water storage and good drainage
- Adequate supply, but not excess of nutrients
- Large population of beneficial organisms
- Small proportion of plant pathogens and insect pests
- Low weed pressure
- Free of chemicals and toxins that may harm the crop
- Resistance to degradation
- Resilience when unfavorable conditions occur

Characteristics of Yagapur Tanda-1 Microwatershed

- ❖ The soil phases identified in the microwatershed belonged to different soil series, VNK series occupies maximum area of 109 ha (15%) followed by ANR 84 ha (12%), BDL 71 ha (10%), YLR 63 ha (9%), YDR 53 ha (8%), GWD 24 ha (3%), BMN 22 ha (3%), BGD 19 ha (3%), BMD 16 ha (2%), KKR 10 ha (1%), BLC 7 ha (<1%), HTK 7 ha (<1%), MDG 6 ha (<1%) and HSL 4 ha (<1%).
- ❖ As per land capability classification entire area of 720 ha in the microwatershed falls under arable land category (Class II, III & IV). The major limitations identified in the arable lands were soil and erosion.
- ❖ On the basis of soil reaction an area of about 25 ha (3%) is slightly to moderately acid (pH 5.5- 6.5). About 109 ha (12%) is neutral (6.5-7.3). About 220 ha (24%) is slightly

alkaline (pH 7.3-7.8). An area of about 343 ha (37%) is moderately alkaline (pH 7.8-8.4) and are distributed in the major part of the microwatershed.

Soil Health Management

The following actions are required to improve the current land husbandry practices that provide a sound basis for the successful adoption of sustainable crop production system.

Acid soils

An area of 25 ha is under acid soils

1. Growing of crops suitable for particular soil pH.
2. Ameliorating the soils through the application of amendments (liming materials).

Liming materials:

1. CaCO_3 (Calcium Carbonate).
2. Dolomite [$\text{Ca Mg} (\text{CO}_3)_2$]
3. Quick lime (Cao)
4. Slaked lime [$\text{Ca} (\text{OH})_2$]

For normal pH and pH 4.8 (35 t/ha) and pH 6.0-7.0 (4 t/ha) lime is required.

Alkaline soils

Moderately and slightly alkaline soils cover an area of about 220 ha in the microwatershed

1. Regular addition of organic manure, green manuring, green leaf manuring, crop residue incorporation and mulching needs to be taken up to improve the soil organic matter status.
2. Application of Biofertilizers (Azospirillum, Azotobacter, Rhizobium).
3. Application of 25% extra N and P (125 % RDN&P).
4. Application of ZnSO_4 – 12.5 kg/ha (once in three years).
5. Application of Boron – 5kg/ha (once in three years).

Neutral soils

An area of about 109 ha is under neutral soils.

1. Regular addition of organic manure, green manuring, green leaf manuring, crop residue incorporation and mulching needs to be taken up to improve the soil organic matter status.
2. Application of Biofertilizers, (Azospirillum, Azotobacter, Rhizobium).
3. Application of 100 per cent RDF.
4. Need based micronutrient applications.

Besides the above recommendations, the best transfer of technology options are also to be adopted.

Soil Degradation

Soil erosion is one of the major factors affecting the soil health in the microwatershed. Out of total 720 ha area in the microwatershed, an area of about 479 ha

(67%) is under moderate erosion. These areas need immediate soil and water conservation and, other land development and land husbandry practices for restoring soil health.

Dissemination of Information and Communication of Benefits

Any large scale implementation of soil health management requires that supporting information is made available widely, particularly through channels familiar to farmers and extension workers. Given the very high priority attached to soil-health especially by the Central Government on issuing Soil-Health Cards to all the farmers, media outlets like Regional, State and National Newspapers, Radio and Dooradarshan programs in local languages but also modern information and communication technologies such as Cellular phones and the Internet, which can be much more effective in reaching the younger farmers.

Inputs for Net Planning (Saturation Plan) and Interventions needed

Net planning (Saturation Plan) in IWMP is focusing on preparation of

1. Soil and Water Conservation Plan for each plot or farm.
2. Productivity enhancement measures/ interventions for existing crops/livestock/other farm enterprises.
3. Diversification of farming mainly with perennial horticultural crops and livestock.
4. Improving livelihood opportunities and income generating activities.

In this connection, how various outputs of Sujala-III are of use in addressing these objectives of Net Planning (Saturation Plan) are briefly presented below.


- ❖ **Soil Depth:** The depth of a soil decides the amount of moisture and nutrients it can hold, what crops can be taken up or not, depending on the rooting depth and the length of growing period available for raising any crop. Deeper the soil, better for a wide variety of crops. If sufficient depth is not available for growing deep rooted crops, either choose medium or short duration crops or deeper planting pits need to be opened and additional good quality soil brought from outside has to be filled into the planting pits.
- ❖ **Surface Soil Texture:** Lighter soil texture in the top soil means, better rain water infiltration, less run-off and soil moisture conservation, less capillary rise and less evaporation losses. Lighter surface textured soils are amenable to good soil tilth and are highly suitable for crops like groundnut, root vegetables (carrot, raddish, potato etc) but not ideal for crops that need stagnant water like lowland paddy. Heavy textured soils are poor in water infiltration and percolation. They are prone for sheet erosion; such soils can be improved by sand mulching. The technology that is developed by the AICRP-Dryland Agriculture, Vijayapura, Karnataka can be adopted.
- ❖ **Gravelliness:** More gravel content is favorable for run-off harvesting but poor in soil moisture storage and nutrient availability. It is a significant parameter that decides the kind of crop to be raised.

- ❖ **Land Capability Classification:** The land capability map shows the areas suitable and not suitable for agriculture and the major constraints in each of the plot/survey number. Hence, one can decide what kind of enterprise is possible in each of these units. In general, erosion and soil are the major constraints in Yagapur Tanda-1 microwatershed.
- ❖ **Organic Carbon:** The OC content (an index of available Nitrogen) is high (>0.75%) in the entire cultivated area of the microwatershed.
- ❖ **Available Phosphorus:** Available Phosphorus is high (>57 kg/ha) covering an area of about 296 ha (41%), medium (23-57 kg/ha) in an area of about 157 ha (22%) and low (<23 kg/ha) in an area of about 45 ha (6%). For all the crops 25% additional P needs to be applied where available P is medium and low.
- ❖ **Available Potassium:** Available potassium content is medium (145-337 kg/ha) in an area of 277 ha (38%) and high (>337 kg/ha) in an area of 221 ha (31%). All the plots, where available potassium is medium, additional 25% potassium may be applied.
- ❖ **Available Sulphur:** Available sulphur content is a very critical nutrient for oilseed crops. It is high (>20 ppm) in the entire cultivated area of the microwatershed.
- ❖ **Available Boron:** Available boron content is medium (0.5-1.0 ppm) covering an area of 64 ha (9%) and about 434 ha (60%) is low (<0.5 ppm) in available boron. For these low and medium areas, application of sodium borate @ 10 kg/ha as soil application or 0.2 % borax as foliar spray is recommended.
- ❖ **Available Iron:** Available iron content is sufficient (>4.5 ppm) in the entire cultivated area of the microwatershed.
- ❖ **Available Manganese:** Entire cultivated area in the microwatershed is sufficient in the available manganese content.
- ❖ **Available Copper:** Entire cultivated area in the microwatershed is sufficient in available copper content.
- ❖ **Available Zinc:** Available zinc content is deficient (<0.6 ppm) in the entire cultivated area of the microwatershed. Application of zinc sulphate @25 kg/ha is recommended for the deficient areas.
- ❖ **Land Suitability for various crops:** Areas that are highly, moderately and marginally suitable for growing various crops are indicated. Along with the suitability, various constraints that are limiting the productivity are also indicated. For example, in case of cotton, gravel content, rooting depth and salinity/alkalinity are the major constraints in various plots. With suitable management interventions, the productivity can be enhanced. In order to increase the water holding capacity of light textured soils, growing of green manure crops and application of organic manure is recommended.

SOIL AND WATER CONSERVATION TREATMENT PLAN

For preparing soil and water conservation treatment plan for Yagapur Tanda-1 microwatershed, the land resource inventory database generated under Sujala-III project has been transformed as information through series of interpretative (thematic) maps using soil phase map as a base. The various thematic maps (1:7920 scale) generated were

- Soil depth
- Surface soil texture
- Available water capacity
- Soil slope
- Soil gravelliness
- Land capability
- Present land use and land cover
- Crop suitability
- Rainfall
- Hydrology
- Water Resources
- Socio-economic data
- Contour plan with existing features- network of waterways, pothissa boundaries, cut up/ minor terraces etc.
- Cadastral map (1:7920 scale)
- Satellite imagery (1:7920 scale)



Apart from these, Hand Level/ Hydro Marker/ Dumpy Level/ Total Station and Kathedars' List to be collected.

Steps for Survey and Preparation of Treatment Plan

The boundaries of Land User Groups' and Survey No. boundaries are traced in the field.

- Naming of user groups and farmers
- Identification of arable and non arable lands
- Identification of drainage lines and gullies
- Identification of non treatable areas
- Identification of priority areas in the arable lands
- Treatment plan for arable lands
- Location of water harvesting and recharge structures

9.1 Treatment Plan

The treatment plan recommended for arable lands is briefly described below

9.1.1 Arable Land Treatment


A. BUNDING

Steps for Survey and Preparation of Treatment Plan	
<ul style="list-style-type: none"> • Cadastral map (1:7920 scale) is enlarged to a scale of 1:2500 scale • Existing network of waterways, pothissa boundaries, grass belts, natural drainage lines/ watercourse, cut ups/ terraces are marked on the cadastral map to the scale • Drainage lines are demarcated into 	
Small gullies	(up to 5 ha catchment)
Medium gullies	(5-15 ha catchment)
Ravines	(15-25 ha catchment) and
<i>Halla/Nala</i>	(more than 25ha catchment)

USER GROUP-1

CLASSIFICATION OF GULLIES

ಕೊರಕಲಿನ ವರ್ಗೀಕರಣ



Measurement of Land Slope

Land slope is estimated or determined by the study and interpretation of contours or by measurement in the field using simple instruments like Hand Level or Hydromarker.



Vertical and Horizontal intervals between bunds as recommended by the Watershed Development Department.

Slope percentage	Vertical interval (m)	Corresponding Horizontal Distance (m)
2 - 3%	0.6	24
3 - 4%	0.9	21
4 - 5%	0.9	21
5 - 6%	1.2	21
6 - 7%	1.2	21

Note: (i) The above intervals are maximum.

(ii) Considering the slope class and erosion status (A1... A=0-1 % slope, 1= slight erosion) the intervals have to be decided.

Bund length recording: Considering the contour plan and the existing grass belts/partitions, the bunds are aligned and lengths are measured.

Section of the Bund

Bund section is decided considering the soil texture class and gravelliness class (bg₀... b=loamy sand, g₀ = <15% gravel). The recommended Sections for different soils are given below.

Recommended Bund Section



Top width (m)	Base width (m)	Height (m)	Side slope (Z:1;H:V)	Cross section (sq m)	Soil Texture	Remarks
0.3	0.9	0.3	01:01	0.18	Sandy loam	Vegetative bund
0.3	1.2	0.3	1.5:1	0.225	Sandy clay	
0.3	1.2	0.5	0.9:1	0.375	Red gravelly soils	
0.3	1.2	0.6	0.75:1	0.45		
0.3	1.5	0.6	01:01	0.54	Red sandy loam	
0.3	2.1	0.6	1.5:1	0.72	Very shallow black soils	
0.45	2	0.75	01:01	0.92		
0.45	2.4	0.75	1.3:1	1.07	Shallow black soils	
0.6	3.1	0.7	1.78:1	1.29	Medium black soils	
0.5	3	0.85	1.47:1	1.49		

Formation of Trench cum Bund

Dimensions of the Borrow Pits/Trenches to be excavated (machinery are decided considering the Bund Section).

Details of Borrow Pit dimensions are given below:



TRENCH CUM BUND

WATER STORAGE AREA

IDEAL FOR HORTICULTURE CROPS

'A' FRAME FOR INTERBUND MANAGEMENT

1. ಸಮಾನಾತಲ ಉಳುವಂ
2. ಸಮಾನಾತಲ ಬಿತ್ತನೆ/ನಾಟಿ

Size of Borrow Pits/ Trench recommended for Trench cum Bund (by machinery)

Bund section	Bund length	Earth quantity	Pit				Berm (pit to pit)	Soil depth class
			L(m)	W(m)	D(m)	Quantity (m ³)		
m ²	M	m ³					m	
0.375	6	2.25	5.85	0.85	0.45	2.24	0.15	Shallow
0.45	6	2.7	5.4	1.2	0.43	2.79	0.6	Shallow
0.45	6	2.7	5	0.85	0.65	2.76	1	Moderately Shallow
0.54	5.6	3.02	5.5	0.85	0.7	3.27	0.1	Moderately shallow
0.54	5.5	2.97	5	1.2	0.5	3	0.5	Shallow
0.72	6.2	4.46	6	1.2	0.7	5.04	0.2	Moderately shallow
0.72	5.2	3.74	5.1	0.85	0.9	3.9	0.1	Moderately deep

B. Water Ways

- Existing waterways are marked on the cadastral map (1:7920 scale) and their dimensions are recorded.
- Considering the contour plan of the MWS, additional waterways/ modernization of the existing ones can be thought of.
- The design details are given in the Manual.

C. Farm Ponds

Waterways and the catchment area will give an indication on the size of the Farm Pond. Location of the pond can be decided based on the contour plan/ field condition and farmers' need/desire.

D. Diversion Channel

Existing EPT/ CPT are marked on the cadastral map. Looking to the need, these can be modernized or fresh diversion channel can be proposed and runoff from this can be stored in Gokatte/ Recharge ponds.

9.1.2 Non-Arable Land Treatment

Depending on the gravelliness and crops preferred by the farmers, the concerned authorities can decide appropriate treatment plan. The recommended treatments may be Contour Trench, Staggered Trench, Crescent Bund, Boulder Bund or Pebble Bund.

9.1.3 Treatment of Natural Water Course/ Drainage Lines

- a) The cadastral map has to be updated as regards the network of drainage lines (gullies/*nalas*/*hallas*) and existing structures are marked to the scale and storage capacity of the existing water bodies are documented.
- b) The drainage line will be demarcated into Upper Reach, Middle Reach and Lower Reach.
- c) Considering the Catchment, *Nala* bed and bank conditions, suitable structures are decided.
- d) Number of storage structures (Check dam/ *Nala* bund/ Percolation tank) will be decided considering the commitments and available runoff from water budgeting and quality of water in the wells and site suitability.
- e) Detailed Leveling Survey using Dumpy Level / Total Station has to be carried out to arrive at the site-specific designs as shown in the Manual.
- f) The location of ground water recharge structures are decided by examining the lineaments and fracture zones from geological maps.
- g) Rainfall intensity data of the nearest Rain Gauge Station is considered for Hydrologic Designs.
- h) Silt load to the Storage/Recharge structures is reduced by providing vegetative, boulder and earthen checks in the natural water course. Location and design details are given in the Manual.

9.2 Recommended Soil and Water Conservation Measures

The appropriate conservation structures best suited for each of the land parcel/ survey number (Appendix-I) are selected based on the slope per cent, severity of erosion, amount of rainfall, land use and soil type. The different kinds of conservation structures recommended are:

1. Graded / Strengthening of Bunds
2. Trench cum Bunds (TCB)
3. Trench cum Bunds / Strengthening
4. Crescent Bunds

A map (Fig. 9.1) showing soil and water conservation plan with different kinds of structures recommended has been prepared which shows the spatial distribution and extent of area. An area of about 195 ha (27%) requires trench cum bunding, 284 ha (39%) requires Graded bunding and about 19 ha (3%) needs strengthening of existing bunds.

The conservation plan prepared may be presented to all the stakeholders including farmers and after considering their suggestions, the conservation plan for the microwatershed may be finalised in a participatory approach.




Fig. 9.1 Soil and Water Conservation Plan map of Yagapur Tanda-1 Microwatershed

9.3 Greening of Microwatershed

As part of the greening programme in the watersheds, it is envisaged to plant a variety of horticultural and other tree plants that are edible, economical and produce lot of biomass which helps to restore the ecological balance in the watersheds. The lands that are suitable for greening programme are non-arable lands (land capability classes V, VI VII and VIII) and also the lands that are not suitable or marginally suitable and field bunds for growing edible and economical annual and perennial crops. The method of planting these trees is given below.

It is recommended to open pits during the 1st week of March along the contour and heap the dug out soil on the lower side of the slope in order to harness the flowing water and facilitate weathering of soil in the pit. Exposure of soil in the pit also prevents spread of pests and diseases due to scorching sun rays. The pits should be filled with mixture of soil and organic manure during the second week of April and keep ready with sufficiently tall seedlings produced either in poly bags or in root trainer nurseries so that planting can be done during the 2nd or 3rd week of April depending on the rainfall.

The tree species suitable for the area considering rainfall, temperature and adaptability is listed below; waterlogged areas are recommended to be planted with species like Nerale (*Syzgium cumini*) and Bamboo. Dry areas are to be planted with species like Honge, Bevu, Seetaphal *etc.*

Dry Deciduous Species			Temp (°C)	Rainfall (mm)
1.	Bevu	<i>Azadiracta indica</i>	21–32	400 –1,200
2.	Tapasi	<i>Holoptelia integrifolia</i>	20-30	500 - 1000
3.	Seetaphal	<i>Anona Squamosa</i>	20-40	400 - 1000
4.	Honge	<i>Pongamia pinnata</i>	20 -50	500– 2,500
5.	Kamara	<i>Hardwickia binata</i>	25 -35	400 - 1000
6.	Bage	<i>Albezzia lebbek</i>	20 - 45	500 - 1000
7.	Ficus	<i>Ficus bengalensis</i>	20 - 50	500–2,500
8.	Sisso	<i>Dalbargia Sissoo</i>	20 - 50	500 -2000
9.	Ailanthus	<i>Ailanthus excelsa</i>	20 - 50	500 - 1000
10.	Hale	<i>Wrightia tinctoria</i>	25 - 45	500 - 1000
11.	Uded	<i>Steriospermum chelanoides</i>	25 - 45	500 -2000
12.	Dhupa	<i>Boswellia Serrata</i>	20 - 40	500 - 2000
13.	Nelli	<i>Embllica Officinalis</i>	20 - 50	500 -1500
14.	Honne	<i>Pterocarpus marsupium</i>	20 - 40	500 - 2000
Moist Deciduous Species			Temp (°C)	Rainfall (mm)
15.	Teak	<i>Tectona grandis</i>	20 - 50	500-5000
16.	Nandi	<i>Legarstroemia lanceolata</i>	20 - 40	500 - 4000
17.	Honne	<i>Pterocarpus marsupium</i>	20 - 40	500 - 3000
18.	Mathi	<i>Terminalia alata</i>	20 -50	500 - 2000
19.	Shivane	<i>Gmelina arborea</i>	20 -50	500 -2000
20.	Kindal	<i>T.Paniculata</i>	20 - 40	500 - 1500
21.	Beete	<i>Dalbargia latifolia</i>	20 - 40	500 - 1500
22.	Tare	<i>T. belerica</i>	20 - 40	500 - 2000
23.	Bamboo	<i>Bambusa arundinasia</i>	20 - 40	500 - 2500
24.	Bamboo	<i>Dendrocalamus strictus</i>	20 – 40	500 – 2500
25.	Muthuga	<i>Butea monosperma</i>	20 - 40	400 - 1500
26.	Hippe	<i>Madhuca latifolia</i>	20 - 40	500 - 2000
27.	Sandal	<i>Santalum album</i>	20 - 50	400 - 1000
28.	Nelli	<i>Embllica officinalis</i>	20 - 40	500 - 2000
29.	Nerale	<i>Syzizium cumini</i>	20 - 40	500 - 2000
30.	Dhaman	<i>Grevia tilifolia</i>	20 - 40	500 - 2000
31.	Kaval	<i>Careya arborea</i>	20 - 40	500 - 2000
32.	Harada	<i>Terminalia chebula</i>	20 - 40	500 - 2000

References

1. FAO (1976) Framework for Land Evaluation, Food and Agriculture Organization, Rome.72 pp.
2. FAO (1983) Guidelines for Land Evaluation for Rainfed Agriculture, FAO, Rome, 237 pp.
3. IARI (1971) Soil Survey Manual, All India Soil and Land Use Survey Organization, IARI, New Delhi, 121 pp.
4. Katyal, J.C. and Rattan, R.K. (2003) Secondary and Micronutrients; Research Gap and Future Needs. Fert. News 48 (4); 9-20.
5. Naidu, L.G.K., Ramamurthy, V., Challa, O., Hegde, R. and Krishnan, P. (2006) Manual Soil Site Suitability Criteria for Major Crops, NBSS Publ. No. 129, NBSS & LUP, Nagpur, 118 pp.
6. Natarajan, A. and Dipak Sarkar (2010) Field Guide for Soil Survey, National Bureau of Soil Survey and Land Use Planning (ICAR), Nagpur, India.
7. Natarajan, A., Rajendra Hegde, Raj, J.N. and Shivananda Murthy, H.G. (2015) Implementation Manual for Sujala-III Project, Watershed Development Department, Bengaluru, Karnataka.
8. Sarma, V.A.K., Krishnan, P. and Budihal, S.L. (1987) Laboratory Manual, Tech. Bull. 23, NBSS &LUP, Nagpur.
9. Sehgal, J.L. (1990) Soil Resource Mapping of Different States of India; Why and How?, National Bureau of Soil Survey and Land Use Planning, Nagpur, 49 pp.
10. Shivaprasad, C.R., R.S. Reddy, J. Sehgal and M. Velayuthum (1998) Soils of Karnataka for Optimizing Land Use, NBSS Publ. No. 47b, NBSS & LUP, Nagpur, India.
11. Soil Survey Staff (2006) Keys to Soil Taxonomy, Tenth edition, U.S. Department of Agriculture/ NRCS, Washington DC, U.S.A.
12. Soil Survey Staff (2012) Soil Survey Manual, Handbook No. 18, USDA, Washington DC, USA.

Appendix I
Yagapur tanda-1 (2H1c) Microwatershed
Soil Phase Information

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Shivanagara	1	2.66	RO	RO	RO	RO	RO	RO	RO	RO	Not Available (NA)	Not Available	RO	RO
Shivanagara	2	2.58	YLRiB2	LMU-7	Moderately shallow (50-75 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Iies	Trench cum bunding
Shivanagara	3	4.23	YLRiB2	LMU-7	Moderately shallow (50-75 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton+Redgram (Ct+Rg)	Not Available	Iies	Trench cum bunding
Shivanagara	4	7.13	YLRiB2	LMU-7	Moderately shallow (50-75 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Iies	Trench cum bunding
Shivanagara	5	2.05	BLCiB2	LMU-6	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Iies	Trench cum bunding
Shivanagara	6	0.09	BLCiB2	LMU-6	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Iies	Trench cum bunding
Shivanagara	13	0.71	BLCiB2	LMU-6	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Iies	Trench cum bunding
Shivanagara	14	5.46	BGDia1	LMU-2	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Cotton (Ct)	Not Available	Iis	Strengthening of existing bunds
Shivanagara	15	6.06	BGDia1	LMU-2	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Cotton (Ct)	Not Available	Iis	Strengthening of existing bunds
Shivanagara	16	6.15	BGDia1	LMU-2	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Cotton (Ct)	Not Available	Iis	Strengthening of existing bunds
Shivanagara	17	5.32	BGDia1	LMU-2	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Cotton (Ct)	Not Available	Iis	Strengthening of existing bunds
Shivanagara	18	5.06	YLRiB2	LMU-7	Moderately shallow (50-75 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Iies	Trench cum bunding
Shivanagara	19	4.98	YLRiB2	LMU-7	Moderately shallow (50-75 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Iies	Trench cum bunding
Shivanagara	20	1.72	YLRiB2	LMU-7	Moderately shallow (50-75 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Iies	Trench cum bunding
Shivanagara	21	1.86	RO	RO	RO	RO	RO	RO	RO	RO	Cotton (Ct)	Not Available	RO	RO
Shivanagara	22	5.77	YLRiB2	LMU-7	Moderately shallow (50-75 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Iies	Trench cum bunding
Shivanagara	23	4.54	YLRiB2	LMU-7	Moderately shallow (50-75 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Iies	Trench cum bunding
Shivanagara	28	0.26	YLRiB2	LMU-7	Moderately shallow (50-75 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iies	Trench cum bunding
Shivanagara	38	0.84	YLRiB2	LMU-7	Moderately shallow (50-75 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Iies	Trench cum bunding
Shivanagara	39	4.25	HTKcB2	LMU-8	Shallow (25-50 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	IIies	Graded bunding
Shivanagara	40	4.97	BGDia1	LMU-2	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Cotton (Ct)	Not Available	Iis	Strengthening of existing bunds
Shivanagara	41	0.89	BLCiB2	LMU-6	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Iies	Trench cum bunding

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Shivanagara	54	8.23	YLRiB2	LMU-7	Moderately shallow (50-75 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Ies	Trench cum bunding
Shivanagara	55	4.09	RO	RO	RO	RO	RO	RO	RO	RO	Not Available (NA)	Not Available	RO	RO
Shivanagara	56	5.57	RO	RO	RO	RO	RO	RO	RO	RO	Hill	Not Available	RO	RO
Shivanagara	57	6.24	YLRiB2	LMU-7	Moderately shallow (50-75 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Ies	Trench cum bunding
Shivanagara	58	0.16	YLRiB2	LMU-7	Moderately shallow (50-75 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Ies	Trench cum bunding
Shivanagara	59	0.73	YLRiB2	LMU-7	Moderately shallow (50-75 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Ies	Trench cum bunding
Shivanagara	60	2.99	YLRiB2	LMU-7	Moderately shallow (50-75 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Ies	Trench cum bunding
Shivanagara	61	2.91	RO	RO	RO	RO	RO	RO	RO	RO	Hill	Not Available	RO	RO
Shivanagara	62	3.63	RO	RO	RO	RO	RO	RO	RO	RO	Hill	Not Available	RO	RO
Shivanagara	63	38.37	RO	RO	RO	RO	RO	RO	RO	RO	Hill	Not Available	RO	RO
Shivanagara	64	5.7	RO	RO	RO	RO	RO	RO	RO	RO	Hill	Not Available	RO	RO
Yagapura	1	0.54	Habitatio n	Others	Others	Others	Others	Others	Others	Others	Redgram (Rg)	Not Available	Others	Others
Yagapura	2	0.38	KKRbB2g 1	LMU-9	Very shallow (<25 cm)	Loamy sand	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IVes	Graded bunding
Yagapura	3	0.9	BMNmB2	LMU-2	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Ies	Graded bunding
Yagapura	28	9.49	YDRcB2	LMU-5	Deep (100-150 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IVes	Graded bunding
Yagapura	29	5.62	BMNmB2	LMU-2	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	Ies	Graded bunding
Yagapura	30	4.18	BDLhB2g 1	LMU-8	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	IIes	Graded bunding
Yagapura	31	1.96	BDLhB2g 1	LMU-8	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Yagapura	32	6.61	BDLhB2g 1	LMU-8	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	IIes	Graded bunding
Yagapura	33	8.02	BDLhB2g 1	LMU-8	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	IIes	Graded bunding
Yagapura	34	7.26	VNKiB2	LMU-8	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	IIes	Trench cum bunding
Yagapura	35	3.77	VNKiB2	LMU-8	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Trench cum bunding
Yagapura	36	2.74	VNKiB2	LMU-8	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	IIes	Trench cum bunding
Yagapura	37	4.99	Habitatio n	Others	Others	Others	Others	Others	Others	Others	Not Available (NA)	Not Available	Others	Others

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Yagapura	38	2.84	BDLhB2g 1	LMU-8	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Yagapura	39	2.43	BDLhB2g 1	LMU-8	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Yagapura	40	7	YDRcB2	LMU-5	Deep (100-150 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	IVes	Graded bunding
Yagapura	41	7.83	YDRcB2	LMU-5	Deep (100-150 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IVes	Graded bunding
Yagapura	42	1.68	YDRcB2	LMU-5	Deep (100-150 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IVes	Graded bunding
Yagapura	43	2.87	YDRcB2	LMU-5	Deep (100-150 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	IVes	Graded bunding
Yagapura	44	4.79	ANRiB2	LMU-3	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Jowar (Rg+Jw)	Not Available	IVes	Graded bunding
Yagapura	45	7.8	ANRiB2	LMU-3	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	IVes	Graded bunding
Yagapura	46	5.44	ANRiB2	LMU-3	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IVes	Graded bunding
Yagapura	47	4.89	YDRcB2	LMU-5	Deep (100-150 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IVes	Graded bunding
Yagapura	48	5.67	YDRcB2	LMU-5	Deep (100-150 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IVes	Graded bunding
Yagapura	49	8.94	ANRiB2	LMU-3	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IVes	Graded bunding
Yagapura	50	7.1	ANRiB2	LMU-3	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IVes	Graded bunding
Yagapura	51	4.22	VNKmB2g 1	LMU-8	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Trench cum bunding
Yagapura	52	2.93	VNKmB2g 1	LMU-8	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	IIes	Trench cum bunding
Yagapura	53	6.36	ANRiB2	LMU-3	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IVes	Graded bunding
Yagapura	54	0.71	ANRiB2	LMU-3	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	IVes	Graded bunding
Yagapura	55	1.08	ANRiB2	LMU-3	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	IVes	Graded bunding
Yagapura	56	3.95	ANRiB2	LMU-3	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	IVes	Graded bunding
Yagapura	57	3.32	VNKiB2	LMU-8	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	IIes	Trench cum bunding
Yagapura	58	5.44	VNKiB2	LMU-8	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Trench cum bunding
Yagapura	59	0.63	ANRiB2	LMU-3	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IVes	Graded bunding
Yagapura	60	4.81	BDLiB2	LMU-8	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	IIes	Graded bunding
Yagapura	61	0.45	BDLiB2	LMU-8	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Yagapura	62	0.92	BDLiB2	LMU-8	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Yagapura	63	0.79	BDLiB2	LMU-8	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Yagapura	64	0.93	BDLiB2	LMU-8	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Yagapura	65	7.14	BDLiB2	LMU-8	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Yagapura	66	8.78	BDLiB2	LMU-8	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Yagapura	67	2.22	HSLbB2	LMU-2	Moderately deep (75-100 cm)	Loamy sand	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Ies	Graded bunding
Yagapura	68	0.06	MDGiB2	LMU-4	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Ies	Graded bunding
Yagapura	71	1.1	HSLbB2	LMU-2	Moderately deep (75-100 cm)	Loamy sand	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Ies	Graded bunding
Yagapura	72	11.68	BDLhB2	LMU-8	Shallow (25-50 cm)	Sandy clay loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Jowar (Rg+Jw)	1 Bore well	IIes	Graded bunding
Yagapura	73	6.8	BDLiB2	LMU-8	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	IIes	Graded bunding
Yagapura	74	1.27	ANRiB2	LMU-3	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	IVes	Graded bunding
Yagapura	75	1.11	ANRiB2	LMU-3	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	IVes	Graded bunding
Yagapura	76	0.78	ANRiB2	LMU-3	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	IVes	Graded bunding
Yagapura	77	1.3	ANRiB2	LMU-3	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	IVes	Graded bunding
Yagapura	78	0.45	ANRiB2	LMU-3	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IVes	Graded bunding
Yagapura	79	2.03	ANRiB2	LMU-3	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IVes	Graded bunding
Yagapura	80	6.2	ANRiB2	LMU-3	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IVes	Graded bunding
Yagapura	81	8.77	ANRiB2	LMU-3	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IVes	Graded bunding
Yagapura	82	13.22	ANRiB2	LMU-3	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton+Redgram (Ct+Rg)	1 Chek Dam	IVes	Graded bunding
Yagapura	83	3.95	ANRiB2	LMU-3	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Hill	Not Available	IVes	Graded bunding
Yagapura	84	6.84	BMDcB2	LMU-1	Very deep (>150 cm)	Sandy loam	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Ies	Trench cum bunding
Yagapura	85	9.82	BMDcB2	LMU-1	Very deep (>150 cm)	Sandy loam	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Scrub land+Redgram (Sl+Rg)	1 Bore well	Ies	Trench cum bunding
Yagapura	86	3.12	VNkbB2g 1	LMU-8	Shallow (25-50 cm)	Loamy sand	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIes	Trench cum bunding
Yagapura	87	7.88	Habitat ion	Others	Others	Others	Others	Others	Others	Others	Not Available (NA)	Not Available	Others	Others

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Yagapura	88	7.68	BDLhB2	LMU-8	Shallow (25-50 cm)	Sandy clay loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Yagapura	89	0.67	HSLbB2	LMU-2	Moderately deep (75-100 cm)	Loamy sand	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Ies	Graded bunding
Yagapura	100	1.11	RO	RO	RO	RO	RO	RO	RO	RO	Hill	Not Available	RO	RO
Yagapura	165	111.26	RO	RO	RO	RO	RO	RO	RO	RO	Hill	Not Available	RO	RO
Yagapura	167	2.48	VNKbB2g1	LMU-8	Shallow (25-50 cm)	Loamy sand	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Hill	Not Available	IIIes	Trench cum bunding
Yagapura	168	0.89	MDGiB2	LMU-4	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	Ies	Graded bunding
Yagapura	169	0.08	MDGiB2	LMU-4	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Ies	Graded bunding
Yagapura	171	0.31	MDGiB2	LMU-4	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	Ies	Graded bunding
Yagapura	178	0.76	MDGiB2	LMU-4	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Ies	Graded bunding
Yagapura	179	0.12	YDRiB2	LMU-5	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IVes	Graded bunding
Yagapura	180	0.08	YDRiB2	LMU-5	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IVes	Graded bunding
Yagapura	181	0.32	MDGiB2	LMU-4	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Ies	Graded bunding
Yagapura	182	0.57	MDGiB2	LMU-4	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Ies	Graded bunding
Yagapura	183	2.77	BDLiB2	LMU-8	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIIes	Graded bunding
Yagapura	184	8.67	ANRiB2	LMU-3	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Jowar (Rg+Jw)	1 Bore well	IVes	Graded bunding
Yagapura	185	0.6	YDRiB2	LMU-5	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IVes	Graded bunding
Yagapura	186	4.47	YDRiB2	LMU-5	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IVes	Graded bunding
Yagapura	187	0.74	ANRiB2	LMU-3	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	IVes	Graded bunding
Yagapura	188	0.98	ANRiB2	LMU-3	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IVes	Graded bunding
Yagapura	189	0.74	ANRiB2	LMU-3	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IVes	Graded bunding
Yagapura	190	6.84	YDRiB2	LMU-5	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	IVes	Graded bunding
Yagapura	191	5.39	YDRiB2	LMU-5	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	IVes	Graded bunding
Yagapura	192	6.27	Habitatio n	Others	Others	Others	Others	Others	Others	Others	Not Available (NA)	Not Available	Others	Others
Yagapura	193	10.83	VNKbB2g1	LMU-8	Shallow (25-50 cm)	Loamy sand	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Fallo w land (Rg+Fl)	Not Available	IIIes	Trench cum bunding

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Yagapura	194	1.86	VNKbB2g 1	LMU-8	Shallow (25-50 cm)	Loamy sand	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Trench cum bunding
Yagapura	195	6.68	VNKbB2g 1	LMU-8	Shallow (25-50 cm)	Loamy sand	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Trench cum bunding
Yagapura	196	2.35	Habitatio n	Others	Others	Others	Others	Others	Others	Others	Not Available (NA)	Not Available	Others	Others
Yagapura	197	7.12	VNKbB2g 1	LMU-8	Shallow (25-50 cm)	Loamy sand	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Trench cum bunding
Yagapura	198	2.41	GWDmB2	LMU-3	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IVes	Graded bunding
Yagapura	199	0.3	GWDmB2	LMU-3	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IVes	Graded bunding
Yagapura	200	0.47	GWDmB2	LMU-3	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IVes	Graded bunding
Yagapura	201	6.05	GWDmB2	LMU-3	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IVes	Graded bunding
Yagapura	202	3.46	GWDmB2	LMU-3	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IVes	Graded bunding
Yagapura	203	0.26	GWDmB2	LMU-3	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IVes	Graded bunding
Yagapura	204	2.37	Habitatio n	Others	Others	Others	Others	Others	Others	Others	Not Available (NA)	Not Available	Others	Others
Yagapura	205	4.82	GWDmB2	LMU-3	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IVes	Graded bunding
Yagapura	206	1.25	GWDmB2	LMU-3	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IVes	Graded bunding
Yagapura	207	0.62	GWDmB2	LMU-3	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IVes	Graded bunding
Yagapura	208	1.42	BMNmB2	LMU-2	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Yagapura	209	0.79	BMNmB2	LMU-2	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iles	Graded bunding
Yagapura	210	4.59	VNKiB2	LMU-8	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Trench cum bunding
Yagapura	211	3.32	GWDmB2	LMU-3	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IVes	Graded bunding
Yagapura	212	1.14	BMNmB2	LMU-2	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Yagapura	213	2.04	KKRcB2	LMU-9	Very shallow (<25 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IVes	Graded bunding
Yagapura	214	2.13	VNKbB2g 1	LMU-8	Shallow (25-50 cm)	Loamy sand	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIes	Trench cum bunding
Yagapura	215	3.82	VNKbB2g 1	LMU-8	Shallow (25-50 cm)	Loamy sand	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Trench cum bunding
Yagapura	216	2.18	VNKbB2g 1	LMU-8	Shallow (25-50 cm)	Loamy sand	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Trench cum bunding
Yagapura	217	4.65	KKRcB2	LMU-9	Very shallow (<25 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IVes	Graded bunding

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Yagapura	218	5.22	BMNmB2	LMU-2	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	1 Bore well	Iles	Graded bunding
Yagapura	219	0.33	BMNmB2	LMU-2	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Yagapura	220	0.39	BMNmB2	LMU-2	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Yagapura	221	0.71	BMNmB2	LMU-2	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Yagapura	222	3.16	BMNmB2	LMU-2	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Yagapura	223	1.22	BMNmB2	LMU-2	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Yagapura	224	1	BMNmB2	LMU-2	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding

Village	Survey Number	Soil Reaction	Salinity	Organic Carbon	Available Phosphorus	Available Potassium	Available Sulphur	Available Boron	Available Iron	Available Manganese	Available Copper	Available Zinc
Yagapura	218	Moderately alkaline (pH 7.8 - 8.4)	Non saline (<2 dsm)	High (> 0.75 %)	High (> 57 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Yagapura	219	Slightly alkaline (pH 7.3 - 7.8)	Non saline (<2 dsm)	High (> 0.75 %)	High (> 57 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Yagapura	220	Moderately alkaline (pH 7.8 - 8.4)	Non saline (<2 dsm)	High (> 0.75 %)	High (> 57 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Yagapura	221	Moderately alkaline (pH 7.8 - 8.4)	Non saline (<2 dsm)	High (> 0.75 %)	High (> 57 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Yagapura	222	Moderately alkaline (pH 7.8 - 8.4)	Non saline (<2 dsm)	High (> 0.75 %)	High (> 57 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Yagapura	223	Moderately alkaline (pH 7.8 - 8.4)	Non saline (<2 dsm)	High (> 0.75 %)	High (> 57 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Yagapura	224	Moderately alkaline (pH 7.8 - 8.4)	Non saline (<2 dsm)	High (> 0.75 %)	High (> 57 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)

Appendix III
Yagapur tanda-1 (2H1c) Microwatershed
Soil Suitability Information

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry	
Shivanagara	1	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0
Shivanagara	2	N1r	S2rt	S3r	S2r	S3r	S2r	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	S3r	S3r	S3r	S3t	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Shivanagara	3	N1r	S2rt	S3r	S2r	S3r	S2r	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	S3r	S3r	S3r	S3t	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Shivanagara	4	N1r	S2rt	S3r	S2r	S3r	S2r	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	S3r	S3r	S3r	S3t	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Shivanagara	5	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S2rn	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r	
Shivanagara	6	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S2rn	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r	
Shivanagara	13	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S2rn	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r	
Shivanagara	14	S3t	S2t	S3t	S1	S3t	S1	S2rt	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S2t	S3t	S2t	S2t	S2t	S2t	S3t	S2t	S2t	S3t	
Shivanagara	15	S3t	S2t	S3t	S1	S3t	S1	S2rt	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S2t	S3t	S2t	S2t	S2t	S2t	S3t	S2t	S2t	S3t	
Shivanagara	16	S3t	S2t	S3t	S1	S3t	S1	S2rt	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S2t	S3t	S2t	S2t	S2t	S2t	S3t	S2t	S2t	S3t	
Shivanagara	17	S3t	S2t	S3t	S1	S3t	S1	S2rt	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S2t	S3t	S2t	S2t	S2t	S2t	S3t	S2t	S2t	S3t	
Shivanagara	18	N1r	S2rt	S3r	S2r	S3r	S2r	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	S3r	S3r	S3r	S3t	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Shivanagara	19	N1r	S2rt	S3r	S2r	S3r	S2r	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	S3r	S3r	S3r	S3t	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Shivanagara	20	N1r	S2rt	S3r	S2r	S3r	S2r	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	S3r	S3r	S3r	S3t	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Shivanagara	21	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0
Shivanagara	22	N1r	S2rt	S3r	S2r	S3r	S2r	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	S3r	S3r	S3r	S3t	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Shivanagara	23	N1r	S2rt	S3r	S2r	S3r	S2r	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	S3r	S3r	S3r	S3t	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Shivanagara	28	N1r	S2rt	S3r	S2r	S3r	S2r	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	S3r	S3r	S3r	S3t	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Shivanagara	38	N1r	S2rt	S3r	S2r	S3r	S2r	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	S3r	S3r	S3r	S3t	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Shivanagara	39	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1r	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Shivanagara	40	S3t	S2t	S3t	S1	S3t	S1	S2rt	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S2t	S3t	S2t	S2t	S2t	S2t	S3t	S2t	S2t	S3t	
Shivanagara	41	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S2rn	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r	
Shivanagara	54	N1r	S2rt	S3r	S2r	S3r	S2r	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	S3r	S3r	S3r	S3t	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Shivanagara	55	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brimjal	Bhendi	Drumstick	Mulberry	
Shivanagara	56	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	
Shivanagara	57	N1r	S2rt	S3r	S2r	S3r	S2r	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	S3r	S3r	S3r	S3t	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Shivanagara	58	N1r	S2rt	S3r	S2r	S3r	S2r	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	S3r	S3r	S3r	S3t	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Shivanagara	59	N1r	S2rt	S3r	S2r	S3r	S2r	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	S3r	S3r	S3r	S3t	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Shivanagara	60	N1r	S2rt	S3r	S2r	S3r	S2r	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	S3r	S3r	S3r	S3t	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Shivanagara	61	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO
Shivanagara	62	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO
Shivanagara	63	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO
Shivanagara	64	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO
Yagapura	1	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Yagapura	2	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Yagapura	3	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz	
Yagapura	28	N1n	S3nt	N1n	S3nt	N1n	N1t	N1n	N1n	N1t	N1n	S3n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n	
Yagapura	29	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz	
Yagapura	30	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Yagapura	31	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Yagapura	32	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Yagapura	33	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Yagapura	34	N1r	S3r	N1r	S3r	N1r	S3r	N1r	N1r	S3rt	N1r	N1r	S3r	N1r	S3r	N1r	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Yagapura	35	N1r	S3r	N1r	S3r	N1r	S3r	N1r	N1r	S3rt	N1r	N1r	S3r	N1r	S3r	N1r	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Yagapura	36	N1r	S3r	N1r	S3r	N1r	S3r	N1r	N1r	S3rt	N1r	N1r	S3r	N1r	S3r	N1r	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Yagapura	37	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Yagapura	38	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Yagapura	39	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Yagapura	40	N1n	S3nt	N1n	S3nt	N1n	N1t	N1n	N1n	N1t	N1n	S3n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n	
Yagapura	41	N1n	S3nt	N1n	S3nt	N1n	N1t	N1n	N1n	N1t	N1n	S3n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n	
Yagapura	42	N1n	S3nt	N1n	S3nt	N1n	N1t	N1n	N1n	N1t	N1n	S3n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n	

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brijjal	Bhendi	Drumstick	Mulberry	
Yagapura	43	N1n	S3nt	N1n	S3nt	N1n	N1t	N1n	N1n	N1t	N1n	S3n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n	
Yagapura	44	N1n	S3n	N1n	S3n	N1n	S3n	N1n	N1n	S3n	N1n	S3n	N1n	N1n	N1n	N1tn	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n
Yagapura	45	N1n	S3n	N1n	S3n	N1n	S3n	N1n	N1n	S3n	N1n	S3n	N1n	N1n	N1n	N1tn	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n
Yagapura	46	N1n	S3n	N1n	S3n	N1n	S3n	N1n	N1n	S3n	N1n	S3n	N1n	N1n	N1n	N1tn	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n
Yagapura	47	N1n	S3nt	N1n	S3nt	N1n	N1t	N1n	N1n	N1t	N1n	S3n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n
Yagapura	48	N1n	S3nt	N1n	S3nt	N1n	N1t	N1n	N1n	N1t	N1n	S3n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n
Yagapura	49	N1n	S3n	N1n	S3n	N1n	S3n	N1n	N1n	S3n	N1n	S3n	N1n	N1n	N1n	N1tn	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n
Yagapura	50	N1n	S3n	N1n	S3n	N1n	S3n	N1n	N1n	S3n	N1n	S3n	N1n	N1n	N1n	N1tn	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n
Yagapura	51	N1r	S3r	N1r	S3r	N1r	S3r	N1r	N1r	S3rt	N1r	N1r	S3r	N1r	S3r	N1r	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r
Yagapura	52	N1r	S3r	N1r	S3r	N1r	S3r	N1r	N1r	S3rt	N1r	N1r	S3r	N1r	S3r	N1r	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r
Yagapura	53	N1n	S3n	N1n	S3n	N1n	S3n	N1n	N1n	S3n	N1n	S3n	N1n	N1n	N1n	N1tn	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n
Yagapura	54	N1n	S3n	N1n	S3n	N1n	S3n	N1n	N1n	S3n	N1n	S3n	N1n	N1n	N1n	N1tn	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n
Yagapura	55	N1n	S3n	N1n	S3n	N1n	S3n	N1n	N1n	S3n	N1n	S3n	N1n	N1n	N1n	N1tn	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n
Yagapura	56	N1n	S3n	N1n	S3n	N1n	S3n	N1n	N1n	S3n	N1n	S3n	N1n	N1n	N1n	N1tn	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n
Yagapura	57	N1r	S3r	N1r	S3r	N1r	S3r	N1r	N1r	S3rt	N1r	N1r	S3r	N1r	S3r	N1r	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r
Yagapura	58	N1r	S3r	N1r	S3r	N1r	S3r	N1r	N1r	S3rt	N1r	N1r	S3r	N1r	S3r	N1r	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r
Yagapura	59	N1n	S3n	N1n	S3n	N1n	S3n	N1n	N1n	S3n	N1n	S3n	N1n	N1n	N1n	N1tn	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n
Yagapura	60	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r
Yagapura	61	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r
Yagapura	62	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r
Yagapura	63	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r
Yagapura	64	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r
Yagapura	65	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r
Yagapura	66	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r
Yagapura	67	S3r	S1	S2r	S1	S2r	S2r	S3r	S2r	S3t	S2r	S2r	S1	S2r	S2r	S3n	S3r	S2r	S2t	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S1	S2r	S2r
Yagapura	68	S3n	S2n	S3n	S2tn	N1n	S3tn	N1n	S3n	S3t	S3n	S2tn	N1n	N1n	S3n	N1n	N1n	S3n	S3n	N1n	S3n	S3n	S3n	S3n	S3n	S2n	S3n	S3n	S3n	N1n	N1n
Yagapura	71	S3r	S1	S2r	S1	S2r	S2r	S3r	S2r	S3t	S2r	S2r	S1	S2r	S2r	S3n	S3r	S2r	S2t	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S1	S2r	S2r

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brimjal	Bhendi	Drumstick	Mulberry
Yagapura	72	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r
Yagapura	73	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r
Yagapura	74	N1n	S3n	N1n	S3n	N1n	S3n	N1n	N1n	S3n	N1n	S3n	N1n	N1n	N1n	N1tn	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n
Yagapura	75	N1n	S3n	N1n	S3n	N1n	S3n	N1n	N1n	S3n	N1n	S3n	N1n	N1n	N1n	N1tn	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n
Yagapura	76	N1n	S3n	N1n	S3n	N1n	S3n	N1n	N1n	S3n	N1n	S3n	N1n	N1n	N1n	N1tn	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n
Yagapura	77	N1n	S3n	N1n	S3n	N1n	S3n	N1n	N1n	S3n	N1n	S3n	N1n	N1n	N1n	N1tn	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n
Yagapura	78	N1n	S3n	N1n	S3n	N1n	S3n	N1n	N1n	S3n	N1n	S3n	N1n	N1n	N1n	N1tn	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n
Yagapura	79	N1n	S3n	N1n	S3n	N1n	S3n	N1n	N1n	S3n	N1n	S3n	N1n	N1n	N1n	N1tn	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n
Yagapura	80	N1n	S3n	N1n	S3n	N1n	S3n	N1n	N1n	S3n	N1n	S3n	N1n	N1n	N1n	N1tn	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n
Yagapura	81	N1n	S3n	N1n	S3n	N1n	S3n	N1n	N1n	S3n	N1n	S3n	N1n	N1n	N1n	N1tn	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n
Yagapura	82	N1n	S3n	N1n	S3n	N1n	S3n	N1n	N1n	S3n	N1n	S3n	N1n	N1n	N1n	N1tn	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n
Yagapura	83	N1n	S3n	N1n	S3n	N1n	S3n	N1n	N1n	S3n	N1n	S3n	N1n	N1n	N1n	N1tn	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n
Yagapura	84	S1	S1	S1	S2t	S1	S3t	S1	S1	S3t	S2t	S2t	S1	S1	S1	S2n	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1
Yagapura	85	S1	S1	S1	S2t	S1	S3t	S1	S1	S3t	S2t	S2t	S1	S1	S1	S2n	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1
Yagapura	86	N1r	S3r	N1r	S3r	N1r	S3r	N1r	N1r	S3rt	N1r	N1r	S3r	N1r	S3r	N1r	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r
Yagapura	87	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Yagapura	88	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r
Yagapura	89	S3r	S1	S2r	S1	S2r	S2r	S3r	S2r	S3t	S2r	S2r	S1	S2r	S2r	S3n	S3r	S2r	S2t	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r
Yagapura	100	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO
Yagapura	165	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO
Yagapura	167	N1r	S3r	N1r	S3r	N1r	S3r	N1r	N1r	S3rt	N1r	N1r	S3r	N1r	S3r	N1r	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r
Yagapura	168	S3n	S2n	S3n	S2tn	N1n	S3tn	N1n	S3n	S3t	S3n	S2tn	N1n	N1n	S3n	N1n	N1n	S3n	S3n	N1n	S3n	S3n	S3n	S3n	S3n	S2n	S3n	S3n	N1n	N1n
Yagapura	169	S3n	S2n	S3n	S2tn	N1n	S3tn	N1n	S3n	S3t	S3n	S2tn	N1n	N1n	S3n	N1n	N1n	S3n	S3n	N1n	S3n	S3n	S3n	S3n	S3n	S2n	S3n	S3n	N1n	N1n
Yagapura	171	S3n	S2n	S3n	S2tn	N1n	S3tn	N1n	S3n	S3t	S3n	S2tn	N1n	N1n	S3n	N1n	N1n	S3n	S3n	N1n	S3n	S3n	S3n	S3n	S3n	S2n	S3n	S3n	N1n	N1n
Yagapura	178	S3n	S2n	S3n	S2tn	N1n	S3tn	N1n	S3n	S3t	S3n	S2tn	N1n	N1n	S3n	N1n	N1n	S3n	S3n	N1n	S3n	S3n	S3n	S3n	S3n	S2n	S3n	S3n	N1n	N1n
Yagapura	179	N1n	S3nt	N1n	S3nt	N1n	N1t	N1n	N1n	N1t	N1n	S3n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n
Yagapura	180	N1n	S3nt	N1n	S3nt	N1n	N1t	N1n	N1n	N1t	N1n	S3n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brimjal	Bhendi	Drumstick	Mulberry	
Yagapura	181	S3n	S2n	S3n	S2tn	N1n	S3tn	N1n	S3n	S3t	S3n	S2tn	N1n	N1n	S3n	N1n	N1n	S3n	S3n	N1n	S3n	S3n	S3n	S3n	S3n	S2n	S3n	S3n	N1n	N1n	
Yagapura	182	S3n	S2n	S3n	S2tn	N1n	S3tn	N1n	S3n	S3t	S3n	S2tn	N1n	N1n	S3n	N1n	N1n	S3n	S3n	N1n	S3n	S3n	S3n	S3n	S3n	S2n	S3n	S3n	N1n	N1n	
Yagapura	183	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Yagapura	184	N1n	S3n	N1n	S3n	N1n	S3n	N1n	N1n	S3n	N1n	S3n	N1n	N1n	N1n	N1tn	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n	
Yagapura	185	N1n	S3nt	N1n	S3nt	N1n	N1t	N1n	N1n	N1t	N1n	S3n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n	
Yagapura	186	N1n	S3nt	N1n	S3nt	N1n	N1t	N1n	N1n	N1t	N1n	S3n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n	
Yagapura	187	N1n	S3n	N1n	S3n	N1n	S3n	N1n	N1n	S3n	N1n	S3n	N1n	N1n	N1n	N1tn	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n	
Yagapura	188	N1n	S3n	N1n	S3n	N1n	S3n	N1n	N1n	S3n	N1n	S3n	N1n	N1n	N1n	N1tn	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n	
Yagapura	189	N1n	S3n	N1n	S3n	N1n	S3n	N1n	N1n	S3n	N1n	S3n	N1n	N1n	N1n	N1tn	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n	
Yagapura	190	N1n	S3nt	N1n	S3nt	N1n	N1t	N1n	N1n	N1t	N1n	S3n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n	
Yagapura	191	N1n	S3nt	N1n	S3nt	N1n	N1t	N1n	N1n	N1t	N1n	S3n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n	
Yagapura	192	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Yagapura	193	N1r	S3r	N1r	S3r	N1r	S3r	N1r	N1r	S3rt	N1r	N1r	S3r	N1r	S3r	N1r	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Yagapura	194	N1r	S3r	N1r	S3r	N1r	S3r	N1r	N1r	S3rt	N1r	N1r	S3r	N1r	S3r	N1r	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Yagapura	195	N1r	S3r	N1r	S3r	N1r	S3r	N1r	N1r	S3rt	N1r	N1r	S3r	N1r	S3r	N1r	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Yagapura	196	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Yagapura	197	N1r	S3r	N1r	S3r	N1r	S3r	N1r	N1r	S3rt	N1r	N1r	S3r	N1r	S3r	N1r	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Yagapura	198	N1n	S3nz	N1n	S3nz	N1n	S3nz	N1n	N1n	S3nz	N1n	S3nz	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3nz	N1n	N1n	N1n	N1n	
Yagapura	199	N1n	S3nz	N1n	S3nz	N1n	S3nz	N1n	N1n	S3nz	N1n	S3nz	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3nz	N1n	N1n	N1n	N1n	
Yagapura	200	N1n	S3nz	N1n	S3nz	N1n	S3nz	N1n	N1n	S3nz	N1n	S3nz	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3nz	N1n	N1n	N1n	N1n	
Yagapura	201	N1n	S3nz	N1n	S3nz	N1n	S3nz	N1n	N1n	S3nz	N1n	S3nz	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3nz	N1n	N1n	N1n	N1n	
Yagapura	202	N1n	S3nz	N1n	S3nz	N1n	S3nz	N1n	N1n	S3nz	N1n	S3nz	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3nz	N1n	N1n	N1n	N1n	
Yagapura	203	N1n	S3nz	N1n	S3nz	N1n	S3nz	N1n	N1n	S3nz	N1n	S3nz	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3nz	N1n	N1n	N1n	N1n	
Yagapura	204	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Yagapura	205	N1n	S3nz	N1n	S3nz	N1n	S3nz	N1n	N1n	S3nz	N1n	S3nz	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3nz	N1n	N1n	N1n	N1n	
Yagapura	206	N1n	S3nz	N1n	S3nz	N1n	S3nz	N1n	N1n	S3nz	N1n	S3nz	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3nz	N1n	N1n	N1n	N1n	
Yagapura	207	N1n	S3nz	N1n	S3nz	N1n	S3nz	N1n	N1n	S3nz	N1n	S3nz	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3nz	N1n	N1n	N1n	N1n	

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry	
Yagapura	208	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz	
Yagapura	209	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz	
Yagapura	210	N1r	S3r	N1r	S3r	N1r	S3r	N1r	N1r	S3rt	N1r	N1r	S3r	N1r	S3r	N1r	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Yagapura	211	N1n	S3nz	N1n	S3nz	N1n	S3nz	N1n	N1n	S3nz	N1n	S3nz	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3nz	N1n	N1n	N1n	N1n	
Yagapura	212	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz	
Yagapura	213	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Yagapura	214	N1r	S3r	N1r	S3r	N1r	S3r	N1r	N1r	S3rt	N1r	N1r	S3r	N1r	S3r	N1r	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Yagapura	215	N1r	S3r	N1r	S3r	N1r	S3r	N1r	N1r	S3rt	N1r	N1r	S3r	N1r	S3r	N1r	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Yagapura	216	N1r	S3r	N1r	S3r	N1r	S3r	N1r	N1r	S3rt	N1r	N1r	S3r	N1r	S3r	N1r	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Yagapura	217	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Yagapura	218	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz	
Yagapura	219	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz	
Yagapura	220	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz	
Yagapura	221	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz	
Yagapura	222	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz	
Yagapura	223	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz	
Yagapura	224	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz	

RO-Rock outcrops

PART-B

SOCIO-ECONOMIC STATUS OF FARM HOUSEHOLDS

CONTENTS

1	Findings of the socio-economic survey	1-3
2	Introduction	5
3	Methodology	7-8
4	Salient features of the survey	9-27
5	Summary	29-32

LIST OF TABLES

1	Households sampled for socio economic survey	9
2	Population characteristics	9
3	Age wise classification of household members	9
4	Education level of household members	10
5	Occupation of household heads	10
6	Occupation of family members	10
7	Institutional participation of household members	11
8	Type of house owned by households	11
9	Durable assets owned by households	11
10	Average value of durable assets owned by households	11
11	Farm implements owned by households	12
12	Average value of farm implements	12
13	Livestock possession by households	13
14	Average labour availability	13
15	Adequacy of hired labour	13
16	Distribution of land (ha)	13
17	Average land value (Rs./ha)	13
18	Status of bore wells	14
19	Source of irrigation	14
20	Depth of water(Avg in meters)	14
21	Irrigated area (ha)	14
22	Cropping pattern	14
23	Cropping intensity	15
24	Possession of bank account and saving	15
25	Borrowing status	15
26	Source of credit	15
27	Avg. credit borrowed	15
28	Purpose of credit borrowed from institutional sources	16
29	Purpose of credit borrowed from Non-institutional sources	16
30	Repayment status of household from institutional sources	16
31	Repayment status of household from Non-institutional sources	16
32.a	Cost of cultivation of Red gram	17

32.b	Cost of cultivation of Cotton	18
32.c	Cost of cultivation of Jowar	19
32.d	Cost of cultivation of Maize	20
32.e	Cost of cultivation of Groundnut	21
33	Adequacy of fodder	22
34	Annual gross income	22
35	Average annual expenditure	22
36	Horticultural species grown	22
37	Forest species grown	23
38	Average additional investment capacity	23
39	Source of funds for additional investment	23
40	Marketing of the agricultural produce	23
41	Marketing channels used for sale of agricultural produce	24
42	Mode of transport of agricultural produce	24
43	Incidence of soil and water erosion problems	24
44	Interest shown towards soil testing	24
45	Usage pattern of fuel for domestic use	25
46	Source of drinking water	25
47	Source of light	25
48	Existence of sanitary toilet facility	25
49	Possession of public distribution system (PDS) card	26
50	Participation in NREGA programme	26
51	Adequacy of food items	26
52	Inadequacy of food items	26
53	Farming constraints experienced	27

FINDINGS OF THE SOCIO-ECONOMIC SURVEY

- ❖ *The survey was conducted in Yagapur Tanda-1 is located at North latitude 16° 56' 28.758" and 16° 54' 29.936" and East longitude 77° 7' 1.416" and 77° 5' 27.747" covering an area of about 725.94 ha coming under Yagapura Village of Chithapura taluk.*
- ❖ *Socio-economic analysis of Yagapur Tanda-1 micro watersheds of Yaragal sub-watershed, Chithapura taluk & Kalaburagi District indicated that, out of the total sample of 34 farmers were sampled in Yagapur Tanda-1 micro-watershed among households surveyed 12 (35.29%) were marginal, 11 (32.35%) were small and 7 (20.59 %) were semi medium farmers. 4 landless farmers were also interviewed for the survey.*
- ❖ *The population characteristics of households indicated that, there were 90 (52.94%) men and 80 (47.06 %) were women. The average population of landless was 4.8, marginal farmers were 4.7, small farmers were 5.6 and semi medium farmers were 4.7.*
- ❖ *Majority of the respondents (44.12%) were in the age group of 16-35 years.*
- ❖ *Education level of the sample households indicated that, there were 62.94 per cent illiterates, 37.05 per cent pre university education and 1.18 per cent attained graduation.*
- ❖ *About, 97.06 per cent of household heads practicing agriculture and 2.94 per cent of the household heads were engaged as agricultural labourers.*
- ❖ *Agriculture was the major occupation for 62.94 per cent of the household members.*
- ❖ *In the study area, 85.29 per cent of the households possess katcha house and 8.82 per cent possess pucca house.*
- ❖ *The durable assets owned by the households showed that, 79.41 per cent possess TV, 41.18 per cent possess mixer grinder, 100.00 per cent possess mobile phones and 5.88 per cent possess motor cycles.*
- ❖ *Farm implements owned by the households indicated that, 23.53 per cent of the households possess plough, 2.94 per cent possess tractor, 14.71 per cent possess bullock cart and 8.82 per cent possess sprayer.*
- ❖ *Regarding livestock possession by the households, 2.94 per cent possess local cow.*
- ❖ *The average labour availability in the study area showed that, own labour men available in the micro watershed was 1.73, women available in the micro watershed was 1.67, hired labour (men) available was 12 and hired labour (women) available was 10.37.*
- ❖ *Out of the total land holding of the sample respondents 97.14 per cent (42.49 ha) of the area is under dry condition and the remaining 2.86 per cent area is irrigated land.*

- ❖ *There were 1.00 live bore wells and 1.00 dry bore wells among the sampled households.*
- ❖ *Bore well was the major source of irrigation for 2.94 per cent of the households.*
- ❖ *The major crops grown by sample farmers are Red gram, Cotton, Jowar, Maize and Groundnut and cropping intensity was recorded as 100.00 per cent.*
- ❖ *Out of the sample households 94.12 percent possessed bank account and 35.29 per cent of them have savings in the account.*
- ❖ *About 100.00 per cent of the respondents borrowed credit from various sources.*
- ❖ *Among the credit borrowed by households, 100.00 per cent from co-operative/Grameena bank.*
- ❖ *Majority of the respondents (100.00%) have borrowed loan for agriculture purpose.*
- ❖ *The per hectare cost of cultivation for Red gram, Cotton, Jowar, Maize and Groundnut was Rs.28112.10 , 30197.34, 25900.72, 12693.85 and 51094.17 with benefit cost ratio of 1:1.20, 1: 1.22, 1: 0.80, 1: 2.90 and 1:0.70 respectively.*
- ❖ *Further, 20.59 per cent of the households opined that dry fodder was adequate and 5.88 per cent of the households have opined that the green fodder was adequate.*
- ❖ *The average annual gross income of the farmers was Rs. 64638.24 in micro-watershed, of which Rs. 37991.18 comes from agriculture.*
- ❖ *Sampled households have grown 3 horticulture trees and 85 forestry trees together in the fields and back yards.*
- ❖ *Households have an average investment capacity of Rs. 1911.76 for land development.*
- ❖ *Source of funds for additional investment is concerned, 47.06 per cent depends on bank loan for land development activities.*
- ❖ *Regarding marketing channels, 88.24 per cent of the households have sold agricultural produce to the local/village merchants.*
- ❖ *Further, 88.24 per cent of the households have used tractor for the transport of agriculture commodity.*
- ❖ *Majority of the farmers (55.88%) have experienced soil and water erosion problems in the watershed and 88.24 per cent of the households were interested towards soil testing.*
- ❖ *Fire was the major source of fuel for domestic use for 73.53 per cent of the households and 26.47 per cent households has LPG connection.*
- ❖ *Piped supply was the major source for drinking water for 58.82 per cent of the households.*
- ❖ *Electricity was the major source of light for 100.00 per cent of the households.*
- ❖ *In the study area, 47.06 per cent of the households possess toilet facility.*
- ❖ *Regarding possession of PDS card, 100.00 per cent of the households possessed BPL card.*

- ❖ *Households opined that, the requirement of cereals (61.76%), pulses (67.65%) and oilseeds (23.53%) are adequate for consumption.*
- ❖ *Farming constraints experienced by households in the micro watersheds were lower fertility status of the soil (94.12%) wild animal menace on farm field (91.18%), frequent incidence of pest and diseases (61.76%), inadequacy of irrigation water (88.24%), high cost of fertilizers and plant protection chemicals (76.47%), high rate of interest on credit (76.47%), low price for the agricultural commodities (64.71%), lack of marketing facilities in the area (38.24%), inadequate extension services (67.65%) and lack of transport for safe transport of the agricultural produce to the market (67.65%).*

INTRODUCTION

Soil and water are the two precious natural resources which are essential for crop production and existence of life on earth. Rainfed agriculture is under severe stress due to various constraints related to agriculture like uneven and erratic distribution of rainfall, indiscriminate use of fertilizers, chemicals and pesticides, adoption of improper land management practices, soil erosion, decline in soil fertility, decline in ground water resources leading to low crop productivity. The area under rainfed agriculture has to be managed effectively using the best available practices to enhance the production of food, fodder and fuel. This is possible if the land resources are characterized at each parcel of land through detailed land resource inventory using the best available techniques of remote sensing, GPS and GIS. The watershed development programs are aimed at the sustainable distribution of its resources and the process of creating and implementing plans, programs, and projects to sustain and enhance watershed functions that affect the plant, animal and human communities within a watershed boundary.

World Bank funded KWDP II, SUJALA III project was implemented in with Broad objective of demonstrating more effective watershed management through greater integration of programmes related to rain-fed agriculture, innovative and science based approaches and strengthen institutional capacities and If successful, it is expected that the systems and tools could be mainstreamed into the overall IWMP in the State of Karnataka and in time, throughout other IWMP operations in India. With this background the socio-economic survey has been carried out with following specific objectives:

1. To understand the demographic features of the households in the micro-watershed
2. To understand the extent of family labour available and additional employment opportunities available within the village.
3. To know the status of assets of households in the micro-watershed for suggesting possible improvements.
4. To study the cropping pattern, cropped area and productivity levels of different households in micro-watershed.
5. To determine the type and extent of livestock owned by different categories of HHs
6. Availability of fodder and level of livestock management.

Scope and importance of survey

Survey helps in identification of different socio-economic and resource use-patterns of farmers at the Micro watershed. Household survey provides demographic features, labour force, and levels of education; land ownership and asset position (including livestock and other household assets) of surveyed households; and cropping patterns, input intensities, and average crop yields from farmers' fields. It also discusses crop utilization and the degree of commercialization of production in the areas; farmers' access to and utilization of credit from formal and informal sources; and the level of adoption and use of soil, water, and pest management technologies.

METHODOLOGY

The description of the methods, components selected for the survey and procedures followed in conducting the baseline survey are furnished under the following heads.

1. Description of the study area

Kalaburagi district is one of the three districts that were transferred from Hyderabad State to Karnataka state at the time of re-organization of the state in 1956. The district is one among the 30 districts of Karnataka State. It is located in the Northern part of the state and lies between 76°.04' and 77°.42 east longitude, and 17°.12' and 17°.46' north latitude, covering an area of 10,951 km². It is bounded on the west by Bijapur district of Karnataka and Sholapur district of Maharashtra, on the west by RangaReddy and Medak district of Telegana State, on the north by Bidar district and Osmanabad district of Maharashtra and on the south by Yadgir district of Karnataka. Kalaburagi is famous for toordal Pigeon pea and the limestone deposits are more in Kalaburagi District. As per Census 2011, Kalaburagi City is an Urban Agglomeration coming under category of Class I UAs/Towns.

The District was under the rule of Nijam s of Hyderabad before independence. The district has a rich background of knowledge and culture. The existence of university at Nagai in Chitapur, Vignaneeshwaras Mitakshara, Nrupatungas Kavirajmarg and the religious and social revolution led by Shivsharanas and the Sufi saint Banda Nawaz are all evidence of it. However, due to erratic rainfall and continuous occurrence of droughts in the 19th century the life of the people was never smooth and secure. Further during the Nizams period, the district could not develop due to the negligence and inefficient administration.

Kalaburagi is situated in Deccan Plateau located at 17.33°N 76.83°E and the general elevation ranges from 300 to 750 meters above mean sea level. Two main rivers, Krishna and Bhima, flow in the district. Black soil is predominant soil type in the district. The district has a large number of tanks which, in addition to the rivers, irrigate the land. The Upper Krishna Project is major irrigation venture in the district. Bajra, toor, sugarcane, groundnut, sunflower, sesame, castor bean, black gram, jowar, wheat, cotton, ragi, Bengal gram, and linseed are grown in this district.

According to the 2011 census Kalaburagi district has a population of 2,564,892. The district has a population density of 233 inhabitants per square kilometre (600/sq mi). Kalaburagi has a sex ratio of 962 females for every 1000 males, and a literacy rate of 65.65%.

2. Locale of the survey and description of the micro-watershed and

The study was conducted in Yagapur Tanda-1 micro-watershed (Yaragal sub-watershed, Chithapura taluk & Kalaburagi District) is located at North latitude 16⁰ 56'

28.758" and 16⁰ 54' 29.936" and East longitude 77⁰ 7' 1.416" and 77⁰ 5' 27.747" covering an area of about 725.94 ha bounded by under Yagapura Village.

3. Selection of the respondents for the study

The micro-watershed is marked with 320 square meters grids. One farmer from every alternate grid in the micro-watershed was selected for the study and interviewed for socio-economic data. Totally 34 households were interviewed for the survey.

4. The parameters considered for socio-economic survey of households

Two forms of data were collected from the micro-watershed which includes primary data from the farm households and secondary data about the villages under the micro-watershed jurisdiction.

The following parameters were considered for the primary data collection about the socio-economic data of the households, (1) Demographic information, (2) Farm and durable assets owned by households, (3) Livestock possession, (4) Labour availability, (5) Level of migration in the village, Land holding, (7) Cropping pattern, (8) Source of irrigation, (9) Borrowing status, (10) Cost of cultivation of major crops, (11) Economics of subsidiary activities, (12) Fodder availability, (13) Family annual income from different sources, (14) Horticulture and forestry species grown, (15) Additional investment capacity, (16) Marketing practices, (17) Status of soil and water conservation structure, (18) Access to basic needs and (19) Constraints and suggestion.

The following parameters were considered for the secondary data regarding the villages under the micro-watershed jurisdiction, (1) Number of villages in each micro-watershed jurisdiction, (2) Village wise number of households, (3) Geographical area of the villages, (4) Cultivable area including rainfed and irrigated, (5) Number and type of house in each village, (6) Human and livestock population, (7) Facilities in the village such as roads, transport facility for conveyance, drinking water supply, street light and (8) Community based organizations in the villages.

5. Development of interview schedule and data collection

Taking into the consideration the objectives of the survey, an interview schedule was prepared after thorough consultation with the experts in the field of social sciences. A comprehensive interview schedule covering all the major parameters for measuring the socio-economic situation was developed.

6. Tools used to analyze the data

The statistical components such as frequency and percentage were used to analyze the data.

Abbreviations used in the report

LL=Landless

MF=Marginal Farmers

SF=Small farmers

SMF=Semi medium farmers

MDF=Medium farmers

LF=Large Farmers

FINDINGS OF THE SURVEY

This chapter deals with systematic presentation of results of the survey. Keeping in view the objectives, the salient features of the survey are presented under the following headings.

Households sampled for socio-economic survey: The data on households sampled for socio economic survey in Yagapur Tanda-1 Micro watershed is presented in Table 1 and it indicated that 34 farmers were sampled in Yagapur Tanda-1 micro-watershed among households surveyed 12 (35.29%) were marginal, 11 (32.35%) were small and 7 (20.59 %) were semi medium farmers. 4 landless farmers were also interviewed for the survey.

Table 1. Households sampled for socio economic survey in Yagapur Tanda-1 micro-watershed

Sl.No.	Particulars	LL (4)		MF (12)		SF (11)		SMF (7)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Farmers	4	11.8	12	35.3	11	32.4	7	20.6	34	100

Population characteristics: The population characteristics of households sampled for socio-economic survey in Yagapur Tanda-1 Micro watershed is presented in Table 2. The data indicated that, there were 90 (52.94%) men and 80 (47.06%) were women. The average population of landless was 4.8, marginal farmers were 4.7, small farmers were 5.6 and semi medium farmers were 4.7.

Table 2. Population characteristics in Yagapur Tanda-1 micro-watershed

Sl.No.	Particulars	LL (19)		MF (56)		SF (62)		SMF (33)		All (170)	
		N	%	N	%	N	%	N	%	N	%
1	Men	10	52.6	29	52	32	52	19	57.6	90	52.9
2	Women	9	47.4	27	48	30	48	14	42.4	80	47.1
Total		19	100	56	100	62	100	33	100	170	100
Average		4.8		4.7		5.6		4.7		5.0	

Age wise classification of population: The age wise classification of household members in Yagapur Tanda-1 Micro watershed is presented in Table 3. The indicated that, 40 (23.53%) of population were 0-15 years of age, 75 (44.12%) were 16-35 years of age, 43(25.29%) were 36-60 years of age and 12 (7.06 %) were above 61 years of age.

Table 3: Age wise classification of members of the household in Yagapur Tanda-1 micro-watershed

Sl.No.	Particulars	LL (19)		MF (56)		SF (62)		SMF (33)		All (170)	
		N	%	N	%	N	%	N	%	N	%
1	0-15 years of age	4	21.1	9	16.1	15	24.2	12	36.36	40	23.53
2	16-35 years of age	8	42.1	30	53.6	27	43.6	10	30.3	75	44.12
3	36-60 years of age	7	36.8	14	25	15	24.2	7	21.21	43	25.29
4	> 61 years	0	0	3	5.36	5	8.06	4	12.12	12	7.06
Total		19	100	56	100	62	100	33	100	170	100

Education level of household members: Education level of household members in Yagapur Tanda-1 Micro watershed is presented in Table 4. The results indicated that, there were 62.94 per cent of illiterates, 20.00 per cent of them had primary school education, 6.47 per cent high school education, 5.29 per cent of them had PUC education, 1.18 per cent attained graduation and 4.12 them had other education.

Table 4. Education level of members of the household in Yagapur Tanda-1 micro-watershed

Sl.No.	Particulars	LL (19)		MF (56)		SF (62)		SMF (33)		All (170)	
		N	%	N	%	N	%	N	%	N	%
1	Illiterate	13	68.4	42	75	38	61.3	14	42.4	107	62.9
2	Primary School	4	21.1	7	12.5	13	21	10	30.3	34	20
3	High School	2	10.5	0	0	6	9.68	3	9.09	11	6.47
4	PUC	0	0	2	3.57	3	4.84	4	12.1	9	5.29
5	Degree	0	0	0	0	0	0	2	6.06	2	1.18
6	Others	0	0	5	8.93	2	3.23	0	0	7	4.12
Total		19	100	56	100	62	100	33	100	170	100

Occupation of head of households: The data regarding the occupation of the household heads in Yagapur Tanda-1 Micro watershed is presented in Table 5. The results indicate that, 97.06 per cent of households heads were practicing agriculture and 2.94 per cent of the household heads were agricultural.

Table 5: Occupation of heads of households in Yagapur Tanda-1 micro-watershed

Sl.No.	Particulars	LL (4)		MF (12)		SF (11)		SMF (7)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Agriculture	4	100	12	100	10	90.91	7	100	33	97.06
2	Agricultural Labour	1	25	0	0	0	0	0	0	1	2.94
Total		5	100	12	100	10	100	7	100	34	100

Occupation of the members of the household: The data regarding the occupation of the household members in Yagapur Tanda-1 Micro watershed is presented in Table 6. The results indicate that, agriculture was the major occupation for 62.94 per cent of the household members, 2.35 per cent were agricultural labour, 6.47 per cent were general labour, 0.59 per cent were working in government sector, 21.76 per cent were working in pursuing education and 4.12 per cent were childrens.

Table 6: Occupation of members of the household in Yagapur Tanda-1 micro-watershed

Sl.No.	Particulars	LL (19)		MF (56)		SF (62)		SMF (33)		All (170)	
		N	%	N	%	N	%	N	%	N	%
1	Agriculture	9	47.4	39	69.6	39	62.9	20	60.61	107	62.9
2	Agricultural Labour	4	21.1	0	0	0	0	0	0	4	2.35
3	General Labour	1	5.26	6	10.7	4	6.45	0	0	11	6.47
4	Government Service	0	0	1	1.79	0	0	0	0	1	0.59
5	Student	3	15.8	4	7.14	17	27.42	13	39.39	37	21.8
6	Others	2	10.5	1	1.79	0	0	0	0	3	1.76
7	Children	0	0	5	8.93	2	3.23	0	0	7	4.12
Total		19	100	56	100	62	100	33	100	170	100

Institutional Participation of household members: The data regarding the institutional participation of the household members in Yagapur Tanda-1 Micro watershed is presented in Table 7. The results show that, out of the total family members in the households 100 per cent of them were not participating in any of the institutions.

Table 7: Institutional Participation of household member in Yagapur Tanda-1 micro-watershed

Sl.No.	Particulars	LL (19)		MF (56)		SF (62)		SMF (33)		All (170)	
		N	%	N	%	N	%	N	%	N	%
1	No Participation	19	100	56	100	62	100	33	100	170	100
	Total	19	100	56	100	62	100	33	100	170	100

Type of house owned: The data regarding the type of house owned by the households in Yagapur Tanda-1 Micro watershed is presented in Table 8. The results indicate that, 5.88 percent possess thatched house, 85.29 per cent of the households possess katcha house and 8.82 per cent possess pacca house.

Table 8. Type of house owned by households in Yagapur Tanda-1 micro-watershed

Sl.No.	Particulars	LL (4)		MF (12)		SF (11)		SMF (7)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Thatched	1	25	1	8.3	0	0	0	0	2	5.88
2	Katcha	2	50	10	83	10	90.91	7	100	29	85.29
3	Pucca/RCC	1	25	1	8.3	1	9.09	0	0	3	8.82
	Total	4	100	12	100	11	100	7	100	34	100

Durable assets owned by the households: The data regarding the Durable Assets owned by the households in Yagapur Tanda-1 Micro watershed is presented in Table 9. The results shows that, 79.41 per cent possess TV, 41.18 per cent possess mixer grinder, 26.47 per cent possess Bicycle, 5.88 per cent possess motor cycle and 100.00 per cent possess mobile phones.

Table 9. Durable assets owned by households in Yagapur Tanda-1 micro-watershed

Sl.No.	Particulars	LL (4)		MF (12)		SF (11)		SMF (7)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Television	3	75	9	75	10	90.9	5	71	27	79.41
2	Mixer/Grinder	0	0	6	50	6	54.6	2	29	14	41.18
3	Bicycle	0	0	2	17	6	54.6	1	14	9	26.47
4	Motor Cycle	0	0	0	0	2	18.2	0	0	2	5.88
5	Mobile Phone	4	100	12	100	11	100	7	100	34	100

Table 10. Average value of durable assets owned in Yagapur Tanda-1 micro-watershed

Sl.No.	Particulars	LL (4)	MF (12)	SF (11)	SMF (7)	All (34)
1	Television	3666	3000	3000	4000	3259
2	Mixer/Grinder	0	966	1000	900	971
3	Bicycle	0	1000	1000	1000	1000
4	Motor Cycle	0	0	35000	0	35000
5	Mobile Phone	1020	1152	1063	1810	1220

Average value of durable assets: The data regarding the average value of durable assets owned by the households in Yagapur Tanda-1 Micro watershed is presented in Table 10. The result shows that, the average value of television was Rs.3259.00, mixer grinder was Rs.971.00, bicycle was Rs.1000.00, motor cycle was Rs. 35000.00 and mobile phone was Rs.1220.00.

Farm implements owned: The data regarding the farm implements owned by the households in Yagapur Tanda-1 Micro watershed is presented in Table 11. About 14.71 per cent of the households possess Bullock Cart, 23.53 per cent possess plough and 8.82 per cent possess Sprayer, 41.18 per cent possess Weeder and 2.94 per cent possess tractor.

Table 11. Farm implements owned in Yagapur Tanda-1 micro-watershed

Sl.No.	Particulars	LL (4)		MF (12)		SF (11)		SMF (7)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Bullock Cart	0	0	0	0	3	27.27	2	28.6	5	14.71
2	Plough	0	0	0	0	6	54.55	2	28.6	8	23.53
3	Tractor	0	0	1	8.33	0	0	0	0	1	2.94
4	Sprayer	0	0	1	8.33	2	18.18	0	0	3	8.82
5	Weeder	1	25	5	41.7	6	54.55	2	28.6	14	41.18
6	Harvester	0	0	1	8.33	0	0	0	0	1	2.94
7	Thresher	0	0	1	8.33	0	0	0	0	1	2.94
8	Chaff Cutter	0	0	0	0	3	27.27	0	0	3	8.82
9	Blank	3	75	7	58.3	2	18.18	4	57.1	16	47.06
10	Earth remover/Duster	0	0	1	8.33	0	0	0	0	1	2.94

Average value of farm implements: The data regarding the average value of farm Implements owned by the households in Yagapur Tanda-1 Micro watershed is presented in Table 12. The results show that the average value of plough was Rs.1000.00, bullock Cart was Rs.33600.00, sprayer was Rs.4666.00, weeder was Rs.80.00 and tractor Rs. 300000.

Table 12. Average value of farm implements in Yagapur Tanda-1 micro-watershed
Average Value (Rs.)

Sl.No.	Particulars	LL (4)	MF (12)	SF (11)	SMF (7)	All (34)
1	Bullock Cart	0	0	33333	34000	33600
2	Plough	0	0	866	2000	1000
3	Tractor	0	300000	0	0	300000
4	Sprayer	0	5000	4500	0	4666
5	Weeder	25	25	150	37	80
6	Harvester	0	5500	0	0	5500
7	Thresher	0	6000	0	0	6000
8	Chaff Cutter	0	0	3000	0	3000
9	Earth remover/Duster	0	55000	0	0	55000

Livestock possession by the households: The data regarding the Livestock possession by the households in Yagapur Tanda-1 Micro watershed is presented in Table 13. The results indicate that, 29.41 per cent of the households possess bullocks and 2.94 per cent possess local cow.

Table 13. Livestock possession by households in Yagapur Tanda-1 micro-watershed

Sl.No.	Particulars	LL (4)		MF (12)		SF (11)		SMF (7)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Bullock	0	0	0	0	7	63.64	3	43	10	29.41
2	Local cow	0	0	0	0	1	9.09	0	0	1	2.94
3	blank	4	100	12	100	4	36.36	4	57	24	70.59

Average Labour availability: The data regarding the average labour availability in Yagapur Tanda-1 Micro watershed is presented in Table 14. The indicated that, own labour men available in the micro watershed was 1.73, women available in the micro watershed was 1.67, hired labour (men) available was 12 and hired labour (women) available was 10.37.

Table 14. Average labour availability in Yagapur Tanda-1 micro-watershed

Sl.No.	Particulars	LL (4)	MF (12)	SF (11)	SMF (7)	All (34)
		N	N	N	N	N
1	Hired labour Female	0	6.25	13.64	12.29	10.37
2	Own Labour Female	0	2	1.64	1.14	1.67
3	Own labour Male	0	2	1.73	1.29	1.73
4	Hired labour Male	0	8.33	15.82	12.29	12

Adequacy of hired labour: The data regarding the adequacy of hired labour in Yagapur Tanda-1 Micro watershed is presented in Table 15. The results indicate that, 88.24 per cent of the household opined that hired labour was adequate.

Table 15. Adequacy of hired labour in Yagapur Tanda-1 micro-watershed

Sl.No.	Particulars	LL (4)		MF (12)		SF (11)		SMF (7)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Adequate	0	0	12	100	11	100	7	100	30	88.2

Distribution of land (ha): The data regarding the distribution of land (ha) in Yagapur Tanda-1 Micro watershed is presented in Table 16. The results indicate that, 41.28 ha (97.14%) of dry land and 1.21 ha (2.86 %) of irrigated land.

Table 16. Distribution of land (ha) in Yagapur Tanda-1 micro-watershed

Sl.No.	Particulars	LL (4)		MF (12)		SF (11)		SMF (7)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Dry	0	0	8.44	100	16.81	100	16.03	92.96	41.28	97.14
2	Irrigated	0	0	0	0	0	0	1.21	7.04	1.21	2.86
	Total	0	100	8.44	100	16.81	100	17.25	100	42.49	100

Table 17. Average value of land (ha) in Yagapur Tanda-1 micro-watershed

Sl.No.	Particulars	LL (4)	MF (12)	SF (11)	SMF (7)	All (34)
		N	N	N	N	N
1	Dry	0	627865.7	273585.4	199495.2	317225.5
2	Irrigated	0	0	0	494000	494000

Average value of land (ha): The data regarding the average land value (Rs./ha) in Yagapur Tanda-1 Micro watershed is presented in Table 17. The results show that the

average value of dry land was Rs.317225.49 and the average value of irrigated land was Rs.494000.00.

Status of bore wells: The data regarding the status of bore wells in Yagapur Tanda-1 Micro watershed is presented in Table 18. The results indicate that, there were 1 De-functioning bore wells and 1 functioning bore wells among the sampled households in micro watershed.

Table 18. Status of bore wells in Yagapur Tanda-1 micro-watershed

Sl.No.	Particulars	LL (4)	MF (12)	SF (11)	SMF (7)	All (34)
		N	N	N	N	N
1	De-functioning	0	0	1	0	1
2	Functioning	0	0	1	0	1

Source of irrigation: The data regarding the source of irrigation in Yagapur Tanda-1 Micro watershed is presented in Table 19. The results that bore well were major source of irrigation for 2.94 per cent of the households.

Table 19. Source of irrigation in Yagapur Tanda-1 micro-watershed

Sl.No.	Particulars	LL (4)		MF (12)		SF (11)		SMF (7)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Bore Well	0	0	0	0	1	9.09	0	0	1	2.94

Depth of water (Avg. In meters): The data regarding the depth of water in Yagapur Tanda-1 Micro watershed is presented in Table 20. The results revealed that, the depth of bore well was 0.90 meter.

Table 20. Depth of water (Avg. In meters) in Yagapur Tanda-1 micro-watershed

Sl.No.	Particulars	LL (4)	MF (12)	SF (11)	SMF (7)	All (34)
		N	N	N	N	N
1	Bore Well	0	0	2.77	0	0.9

Irrigated Area (ha): The data regarding the irrigated area (ha) in Yagapur Tanda-1 Micro watershed is presented in Table 21. The results indicate that, the availability of irrigation water was used for 2.15 ha for perennial crop.

Table 21. Irrigated Area (ha) in Yagapur Tanda-1 micro-watershed

Sl.No.	Particulars	LL (4)	MF (12)	SF (11)	SMF (7)	All (34)
1	Perennial Crops	0	0	2.15	0	2.15

Table 22. Cropping pattern in Yagapur Tanda-1 micro-watershed

Sl.No.	Particulars	LL (4)	MF (12)	SF (11)	SMF (7)	All (34)
1	Kharif - Red gram (togari)	0	2.99	9.29	6.88	19.17
2	Kharif - Groundnut	0	2.49	7.5	8.27	18.26
3	Kharif - Maize	0	0	0	2.11	2.11
4	Kharif - Cotton	0	1.73	0	0	1.73
5	Kharif - Jowar	0	1.26	0	0	1.26
	Total	0	8.47	16.79	17.26	42.51

Cropping pattern: The data regarding the cropping pattern in Yagapur Tanda-1 Micro watershed is presented in Table 22. The results indicate that, farmers have grown Red

gram (19.17 ha), Groundnut (18.26 ha), Maize (2.11 ha), Cotton (1.73 ha) and Jowar (1.26 ha).

Cropping intensity: The data regarding the cropping intensity in Yagapur Tanda-1 Micro watershed is presented in Table 23. The results indicate that, the cropping intensity was 100.00 per cent.

Table 23. Cropping intensity (%) in Yagapur Tanda-1 micro-watershed

Sl.No.	Particulars	LL (4)	MF (12)	SF (11)	SMF (7)	All (34)
1	Cropping Intensity	0	100	100	100	100

Possession of bank account and savings: The data regarding the possession of bank account and saving in Yagapur Tanda-1 micro-watershed is presented in Table 24. The results indicate that, 94.12 cent of the households posses bank account and 35.29 per cent of them have savings.

Table 24. Possession of Bank account and savings in Yagapur Tanda-1 micro-watershed

Sl.No.	Particulars	LL (4)		MF (12)		SF (11)		SMF (7)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Account	3	75	12	100	10	90.91	7	100	32	94.12
2	Savings	1	25	5	41.67	3	27.27	3	42.86	12	35.29

Borrowing status: The data regarding the borrowing status in Yagapur Tanda-1 micro-watershed is presented in Table 25. The results indicate that, 100.00 percent of the sample farmers have borrowed credit from different sources.

Table 25. Borrowing status in Yagapur Tanda-1 micro-watershed

Sl.No.	Particulars	LL (4)		MF (12)		SF (11)		SMF (7)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Credit Availed	3	75	12	100	12	109	7	100	34	100

Source of credit: The data regarding the source of credit availed by households in Yagapur Tanda-1 micro-watershed is presented in Table 26. The results show that, 100.00 per cent have borrowed loan from Grameena Bank and 30.00 per cent have borrowed loan from money lender.

Table 26. Source of credit borrowed by households in Yagapur Tanda-1 micro-watershed

Sl.No.	Particulars	LL (0)		MF (3)		SF (6)		SMF (1)		All (10)	
		N	%	N	%	N	%	N	%	N	%
1	Grameena Bank	0	0	3	100	6	100	1	100	10	100
2	Money Lender	0	0	2	66.7	1	16.7	0	0	3	30

Table 27. Avg. Credit amount in Yagapur Tanda-1 micro-watershed

Sl.No.	Particulars	LL (0)	MF (3)	SF (6)	SMF (1)	All (10)
		N	N	N	N	N
1	Average Credit	0	25000	35000	50000	18333.3

Avg. Credit amount: The data regarding the avg. Credit amount in Yagapur Tanda-1 micro-watershed is presented in Table 27. The results show that, farmers have borrowed Avg. Credit of Rs.18333.33 from different sources.

Purpose of credit borrowed (institutional Source): The data regarding the purpose of credit borrowed - Institutional Credit in Yagapur Tanda-1 micro-watershed is presented in Table 28. The results indicate that, 100.00 per cent of the households have borrowed loan for agriculture.

Table 28. Purpose of credit borrowed (institutional Source) by households in Yagapur Tanda-1 micro-watershed

SN	Particulars	LL (0)		MF (3)		SF (6)		SMF (1)		All (10)	
		N	%	N	%	N	%	N	%	N	%
1	Agriculture production	0	0	3	100	6	100	1	100	10	100

Purpose of credit borrowed (Private Source): The data regarding the purpose of credit borrowed – Private Source in Yagapur Tanda-1 micro-watershed is presented in Table 29. The results indicate that, 100.00 per cent of the households have borrowed loan for agriculture.

Table 29. Purpose of credit borrowed (Private Source) by households in Yagapur Tanda-1 micro-watershed

Sl.No.	Particulars	LL (0)		MF (2)		SF (1)		SMF (0)		All (3)	
		N	%	N	%	N	%	N	%	N	%
1	Agriculture production	0	0	2	100	1	100	0	0	3	100

Repayment status of household (institutional Source): The data regarding the repayment status of credit borrowed from institutional Source by households in Yagapur Tanda-1 micro watershed is presented in Table 30. The results indicate that, 100.00 per cent have unpaid.

Table 30. Repayment status of household (institutional Source) in Yagapur Tanda-1 micro-watershed

Sl.No.	Particulars	LL (0)		MF (3)		SF (6)		SMF (1)		All (10)
		N	%	N	%	N	%	N	%	%
1	Un paid	0	0	3	100	6	100	1	100	100

Repayment status of household (Private Source): The data regarding the repayment status of credit borrowed from private sources by households in Yagapur Tanda-1 micro watershed is presented in Table 31. The results indicate that, 100 per cent of the households have unpaid.

Table 31. Repayment status of household (Private Source) in Yagapur Tanda-1 micro-watershed

Sl.No.	Particulars	LL (0)		MF (2)		SF (1)		SMF (0)		All (3)	
		N	%	N	%	N	%	N	%	N	%
1	Un paid	0	0	2	100	1	100	0	0	3	100

Cost of Cultivation of Red gram: The data regarding the cost of cultivation (Rs/ha) of Red gram in Yagapur Tanda-1 micro watershed is presented in Table 32.a. The results indicate that, the total cost of cultivation (Rs/ha) for Red gram was Rs. 28112.10. The gross income realized by the farmers was Rs. 33036.05. The net income from Red gram cultivation was Rs.4923.95, thus the benefit cost ratio was found to be 1:1.20.

Table 32(a). Cost of Cultivation of Red gram in Yagapur Tanda-1 micro-watershed

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
I	Cost A1				
1	Hired Human Labour	Man days	22.63	4526.32	16.1
2	Bullock	Pairs/day	0.74	560.2	1.99
3	Tractor	Hours	1.85	1490.9	5.3
4	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	16.33	1387.7	4.94
5	FYM	Quintal	2.57	5042.92	17.94
6	Fertilizer + micronutrients	Quintal	3.64	5296.68	18.84
7	Pesticides (PPC)	Kgs / liters	0.99	1226.62	4.36
8	Depreciation charges		0	121.93	0.43
II	Cost B1				
9	Interest on working capital			1555.59	5.53
10	Cost B1 = (Cost A1 + sum of 15 and 16)			21208.85	75.44
III	Cost B2				
11	Rental Value of Land			166.67	0.59
12	Cost B2 = (Cost B1 + Rental value)			21375.51	76.04
IV	Cost C1				
13	Family Human Labour		15.91	4171.63	14.84
14	Cost C1 = (Cost B2 + Family Labour)			25547.15	90.88
V	Cost C2				
15	Risk Premium			9.31	0.03
16	Cost C2 = (Cost C1 + Risk Premium)			25556.45	90.91
VI	Cost C3				
17	Managerial Cost			2555.65	9.09
18	Cost C3 = (Cost C2 + Managerial Cost)			28112.1	100
VII	Economics of the Crop				
a.	Main Product	a) Main Product (q)		6.89	33036.05
		b) Main Crop Sales Price (Rs.)			4796.15
b.	Gross Income (Rs.)				33036.05
c.	Net Income (Rs.)				4923.95
d.	Cost per Quintal (Rs./q.)				4081.3
e.	Benefit Cost Ratio (BC Ratio)				1:1.2

Cost of Cultivation of Cotton: The data regarding the cost of cultivation (Rs/ha) of Cotton in Yagapur Tanda-1 micro watershed is presented in Table 32.b. The results indicate that, the total cost of cultivation (Rs/ha) for Cotton was Rs. 30197.34. The gross income realized by the farmers was Rs. 36875.90. The net income from Cotton cultivation was Rs.6678.56, thus the benefit cost ratio was found to be 1:1.22.

Table 32(b). Cost of Cultivation of Cotton in Yagapur Tanda-1 micro-watershed

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
I	Cost A1				
1	Hired Human Labour	Man days	33.21	5330.63	17.65
2	Bullock	Pairs/day	0	0	0
3	Tractor	Hours	2.32	1626.17	5.39
4	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	2.87	4587.45	15.19
5	Fertilizer + micronutrients	Quintal	4.65	3833.12	12.69
6	Pesticides (PPC)	Kgs / liters	2.32	5734.32	18.99
7	Depreciation charges		0	0.02	0
II	Cost B1				
8	Interest on working capital			1699.79	5.63
9	Cost B1 = (Cost A1 + sum of 15 and 16)			22811.51	75.54
III	Cost B2				
10	Rental Value of Land			166.67	0.55
11	Cost B2 = (Cost B1 + Rental value)			22978.18	76.09
IV	Cost C1				
12	Family Human Labour		18.36	4463.95	14.78
13	Cost C1 = (Cost B2 + Family Labour)			27442.13	90.88
V	Cost C2				
14	Risk Premium			10	0.03
15	Cost C2 = (Cost C1 + Risk Premium)			27452.13	90.91
VI	Cost C3				
16	Managerial Cost			2745.21	9.09
17	Cost C3 = (Cost C2 + Managerial Cost)			30197.34	100
VII	Economics of the Crop				
a.	Main Product	a) Main Product (q)	9.22	36875.9	
		b) Main Crop Sales Price (Rs.)		4000	
b.	Gross Income (Rs.)			36875.9	
c.	Net Income (Rs.)			6678.56	
d.	Cost per Quintal (Rs./q.)			3275.56	
e.	Benefit Cost Ratio (BC Ratio)			1:1.22	

Cost of Cultivation of Jowar: The data regarding the cost of cultivation (Rs/ha) of Jowar in Yagapur Tanda-1 micro watershed is presented in Table 32.c. The results indicate, the total cost of cultivation (Rs/ha) for Jowar was Rs.25900.72. The gross income realized by the farmers was Rs. 20868.90. The net income from Jowar cultivation was Rs. -5031.82, thus the benefit cost ratio was found to be 1:0.80.

Table 32(c). Cost of Cultivation of Jowar in Yagapur Tanda-1 micro-watershed

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
I	Cost A1				
1	Hired Human Labour	Man days	49.01	9231.66	35.64
2	Bullock	Pairs/day	0.61	605.39	2.34
3	Tractor	Hours	2.38	1663.12	6.42
4	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	11.88	415.78	1.61
5	FYM	Quintal	1.21	2421.57	9.35
6	Fertilizer + micronutrients	Quintal	2.94	2648.88	10.23
7	Pesticides (PPC)	Kgs / liters	1.21	1210.78	4.67
8	Depreciation charges		0	0.04	0
II	Cost B1				
9	Interest on working capital			804.84	3.11
10	Cost B1 = (Cost A1 + sum of 15 and 16)			19002.05	73.36
III	Cost B2				
11	Rental Value of Land			166.67	0.64
12	Cost B2 = (Cost B1 + Rental value)			19168.72	74.01
IV	Cost C1				
13	Family Human Labour		17.89	4367.39	16.86
14	Cost C1 = (Cost B2 + Family Labour)			23536.11	90.87
V	Cost C2				
15	Risk Premium			10	0.04
16	Cost C2 = (Cost C1 + Risk Premium)			23546.11	90.91
VI	Cost C3				
17	Managerial Cost			2354.61	9.09
18	Cost C3 = (Cost C2 + Managerial Cost)			25900.72	100
VII	Economics of the Crop				
a.	Main Product	a) Main Product (q)	11.93	20868.9	
		b) Main Crop Sales Price (Rs.)		1750	
b.	Gross Income (Rs.)			20868.9	
c.	Net Income (Rs.)			-5031.82	
d.	Cost per Quintal (Rs./q.)			2171.95	
e.	Benefit Cost Ratio (BC Ratio)			1:0.8	

Cost of Cultivation of Maize: The data regarding the cost of cultivation (Rs/ha) of Maize in Yagapur Tanda-1 micro watershed is presented in Table 32.d. The results indicate that, the total cost of cultivation (Rs/ha) for Maize was Rs. 12693.85. The gross income realized by the farmers was Rs.37050.00. The net income from Maize cultivation was Rs. 24356.15, thus the benefit cost ratio was found to be 1:2.90.

Table 32(d). Cost of Cultivation of Maize in Yagapur Tanda-1 micro-watershed

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
I	Cost A1				
1	Hired Human Labour	Man days	21.85	4655	36.67
2	Bullock	Pairs/day	0	0	0
3	Tractor	Hours	2.38	1781.25	14.03
4	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	7.13	249.38	1.96
5	Fertilizer + micronutrients	Quintal	2.85	2850	22.45
6	Pesticides (PPC)	Kgs / liters	0.48	237.5	1.87
7	Depreciation charges		0	0.95	0.01
II	Cost B1				
16	Interest on working capital			401.63	3.16
17	Cost B1 = (Cost A1 + sum of 15 and 16)			10175.7	80.16
III	Cost B2				
18	Rental Value of Land			166.67	1.31
19	Cost B2 = (Cost B1 + Rental value)			10342.37	81.48
IV	Cost C1				
20	Family Human Labour		4.75	1187.5	9.35
21	Cost C1 = (Cost B2 + Family Labour)			11529.87	90.83
V	Cost C2				
22	Risk Premium			10	0.08
23	Cost C2 = (Cost C1 + Risk Premium)			11539.87	90.91
VI	Cost C3				
24	Managerial Cost			1153.99	9.09
25	Cost C3 = (Cost C2 + Managerial Cost)			12693.85	100
VII	Economics of the Crop				
a.	Main Product	a) Main Product (q)	14.25	37050	
		b) Main Crop Sales Price (Rs.)		2600	
b.	Gross Income (Rs.)			37050	
c.	Net Income (Rs.)			24356.15	
d.	Cost per Quintal (Rs./q.)			890.8	
e.	Benefit Cost Ratio (BC Ratio)			1:2.9	

Cost of Cultivation of Groundnut: The data regarding the cost of cultivation (Rs/ha) of Groundnut in Yagapur Tanda-1 micro watershed is presented in Table 32.e. The results indicate that, the total cost of cultivation (Rs/ha) for Groundnut was Rs.51094.17. The gross income realized by the farmers was Rs. 36455.60. The net income from Groundnut cultivation was Rs. -14638.57, thus the benefit cost ratio was found to be 1:0.70.

Table 32(e). Cost of Cultivation of Groundnut in Yagapur Tanda-1 micro-watershed

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
I	Cost A1				
1	Hired Human Labour	Man days	30.46	6138.89	12.01
2	Bullock	Pairs/day	1.2	1059.06	2.07
3	Tractor	Hours	1.95	1542.75	3.02
4	Machinery	Hours	0	0	0
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	173.94	13407.51	26.24
6	Seed Inter Crop	Kgs.	0	0	0
7	FYM	Quintal	2.45	9710.61	19.01
8	Fertilizer + micronutrients	Quintal	4.63	3973.05	7.78
9	Pesticides (PPC)	Kgs / liters	1.07	802.36	1.57
10	Irrigation	Number	0	0	0
11	Repairs		0	0	0
12	Msc. Charges (Marketing costs etc)		0	0	0
13	Depreciation charges		0	856.04	1.68
14	Land revenue and Taxes		0	0	0
II	Cost B1				
16	Interest on working capital			3348.42	6.55
17	Cost B1 = (Cost A1 + sum of 15 and 16)			40838.69	79.93
III	Cost B2				
18	Rental Value of Land			166.67	0.33
19	Cost B2 = (Cost B1 + Rental value)			41005.36	80.25
IV	Cost C1				
20	Family Human Labour		21.92	5433.89	10.64
21	Cost C1 = (Cost B2 + Family Labour)			46439.25	90.89
V	Cost C2				
22	Risk Premium			10	0.02
23	Cost C2 = (Cost C1 + Risk Premium)			46449.25	90.91
VI	Cost C3				
24	Managerial Cost			4644.92	9.09
25	Cost C3 = (Cost C2 + Managerial Cost)			51094.17	100
VII	Economics of the Crop				
a.	Main Product	a) Main Product (q)	7.91	36455.6	
		b) Main Crop Sales Price (Rs.)		4608.33	
b.	Gross Income (Rs.)			36455.6	
c.	Net Income (Rs.)			-14638.57	
d.	Cost per Quintal (Rs./q.)			6458.79	
e.	Benefit Cost Ratio (BC Ratio)			1:0.7	

Adequacy of fodder: The data regarding the adequacy of fodder in Yagapur Tanda-1 Micro watershed is presented in Table 33. The results indicate that, 20.59 per cent of the households opined that dry fodder was adequate and 2.94 per cent of them opined dry fodder was inadequate. With respect to green fodder availability, 5.88 percent of them opined it was sufficient.

Table 33. Adequacy of fodder in Yagapur Tanda-1 micro-watershed

Sl.No.	Particulars	LL (4)		MF (12)		SF (11)		SMF (7)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Adequate-Dry Fodder	0	0	0	0	4	36.36	3	42.9	7	20.59
2	Inadequate-Dry Fodder	0	0	0	0	1	9.09	0	0	1	2.94
3	Adequate-Green Fodder	0	0	0	0	2	18.18	0	0	2	5.88

Average annual gross income: The data regarding the annual gross income in Yagapur Tanda-1 Micro watershed is presented in Table 34. The results indicate that, the farmers have annual gross income of Rs. 64638.24 in micro-watershed, of which Rs. 37991.18 is from agriculture itself.

Table 34. Average annual gross income in Yagapur Tanda-1 micro-watershed

Sl.No.	Particulars	LL (4)	MF (12)	SF (11)	SMF (7)	All (34)
		Rs.	Rs.	Rs.	Rs.	Rs.
1	Wage	53750	29500	22454.6	12857.1	26647.1
2	Agriculture	0	28058.3	54545.5	50714.3	37991.2
Income(Rs.)		53750	57558.3	77000	63571.4	64638.2

Average annual Expenditure: The data regarding the average annual expenditure in Yagapur Tanda-1 Micro watershed is presented in Table 35. The results indicate that, the farmers have annual gross expenditure of Rs. 180944.81 in micro-watershed, of which Rs. 31205.88 is from agriculture itself.

Table 35. Average annual Expenditure in Yagapur Tanda-1 micro-watershed

Sl.No.	Particulars	LL (4)	MF (12)	SF (11)	SMF (7)	All (34)
		Rs.	Rs.	Rs.	Rs.	Rs.
1	Wage	18750	3285.71	0	45000	4205.88
2	Agriculture	0	23181.8	42727.3	48000	31205.9
Total		18750	26467.5	42727.3	93000	180945

Horticulture species grown: The data regarding horticulture species grown in Yagapur Tanda-1 Micro watershed is presented in Table 36. The results indicate that, the total number of horticultural trees grown (both field and backyard) by the sampled households were coconut (2) and Lemon (1).

Table 36. Horticulture species grown in Yagapur Tanda-1 micro-watershed

Sl.No.	Particulars	LL (4)		MF (12)		SF (11)		SMF (7)		All (34)	
		F	B	F	B	F	B	F	B	F	B
1	Coconut	0	0	0	0	2	0	0	0	2	0
2	Lemon	0	0	0	0	1	0	0	0	1	0

*F= Field B=Back Yard

Forest species grown: The data regarding forest species grown in Yagapur Tanda-1 Micro watershed is presented in Table 37. The results indicate that, households have planted 2 teak trees, 66 neem trees, 6 tamarind trees, 10 acacia trees and 1 banyan trees together in both field and backyard.

Table 37. Forest species grown in Yagapur Tanda-1 micro-watershed

Sl.No.	Particulars	LL (4)		MF (12)		SF (11)		SMF (7)		All (34)	
		F	B	F	B	F	B	F	B	F	B
1	Neem	0	0	12	0	40	2	12	0	64	2
2	Teak	0	0	0	0	2	0	0	0	2	0
3	Tamarind	0	0	0	0	6	0	0	0	6	0
4	Acacia	0	0	1	0	2	0	7	0	10	0
5	Banyan	0	0	0	0	1	0	0	0	1	0

*F= Field B=Back Yard

Average additional investment capacity: The data regarding average additional investment capacity in Yagapur Tanda-1 Micro watershed is presented in Table 38. The results indicate that, households have an average investment capacity of Rs. 1911.76 for land development and Rs.1411.76 for adoption of improved livestock breeds.

Table 38. Average additional investment capacity of households in Yagapur Tanda-1 micro-watershed

Sl.No.	Particulars	LL (4)	MF (12)	SF (11)	SMF (7)	All (34)
		Rs.	Rs.	Rs.	Rs.	Rs.
1	Land development	0	1333.33	1090.91	5285.71	1911.76
2	Improved crop production	0	1333.33	909.09	3142.86	1411.76

Source of funds for additional investment: The data regarding source of funds for additional investment in Yagapur Tanda-1 Micro watershed is presented in Table 39. The results indicate that, the sources of finance raised from own sources for land development were 47.06.

Table 39. Source of funds for additional investment in Yagapur Tanda-1 micro-watershed

Sl.No	Item	Land development		Irrigation facility		Improved crop production	
		N	%	N	%	N	%
1	Own funds	16	47.06	0	0	8	23.53

Table 40. Marketing of agricultural produce in Yagapur Tanda-1 micro-watershed

Sl.No	Crops	Output obtained (q)	Output retained (q)	Output sold (q)	Output sold (%)	Avg. Price obtained (Rs/q)
1	Cotton	16	0	16	100	4000
2	Groundnut	127	8	119	94	4608
3	Jowar	16	4	12	75	1750
4	Maize	30	0	30	100	2600
5	Red gram	118	8	110	93	4796

Marketing of agricultural produce: The data regarding marketing of the agricultural produce in Yagapur Tanda-1 Micro watershed is presented in Table 40. The results indicated that, 100.00 percent of output of Cotton was sold in the market with average price of Rs. 4000.00; 93.70 per cent of output of Groundnut was sold in the market with average price of Rs. 4608.33; 75.00 per cent of output of Jowar was sold in the market with average price of Rs. 1750.00 and 100.00 per cent of output of Maize was sold in the market with average price of Rs. 2600.00.

Marketing channels used for sale of agricultural produce: The data regarding marketing channels used for sale of agricultural produce in Yagapur Tanda-1 Micro watershed is presented in Table 41. The results indicated that, 88.24 cent of the households have sold agricultural produce to the local/village merchants.

Table 41. Marketing channels used for sale of agricultural produce in Yagapur Tanda-1 micro-watershed 5

Sl.No.	Particulars	LL (4)		MF (12)		SF (11)		SMF (7)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Local/village Merchant	0	0	12	100	11	100	7	100	30	88.24

Mode of transport of agricultural produce: The data regarding mode of transport of agricultural produce in Yagapur Tanda-1 Micro watershed is presented in Table 42. The results indicated that, 88.24 cent of the households have used tractor for the transport of agriculture commodity.

Table 42. Mode of transport of agricultural produce in Yagapur Tanda-1 micro-watershed

Sl.No.	Particulars	LL (4)		MF (12)		SF (11)		SMF (7)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Tractor	0	0	12	100	11	100	7	100	30	88.24

Incidence of soil and water erosion problems: The data regarding incidence of incidence of soil and water erosion problems in Yagapur Tanda-1 Micro watershed is presented in Table 43. The results indicate that, 55.88 per cent of the households have experienced soil and water erosion problems.

Table 43. Incidence of soil and water erosion problems in Yagapur Tanda-1 micro-watershed

Sl.No	Particulars	LL (4)		MF (12)		SF (11)		SMF (7)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Soil and water erosion problems in the farm	0	0	7	58	5	45.5	7	100	19	55.88

Table 44. Interest regarding soil testing in Yagapur Tanda-1 micro-watershed

Sl.No.	Particulars	LL (4)		MF (12)		SF (11)		SMF (7)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Interest in soil test	0	0	12	100	11	100	7	100	30	88.24

Interest towards soil testing: The data regarding Interest shown towards soil testing in Yagapur Tanda-1 Micro watershed is presented in Table 44. The results indicated that, 88.24 per cent of the households were interested towards soil testing.

Usage pattern of fuel for domestic use: The data on usage pattern of fuel for domestic use in Yagapur Tanda-1 Micro watershed is presented in Table 45. The results indicated that, firewood was the major source of fuel for domestic use for 73.53 per cent of the households followed by LPG (26.47%) and Biogas (2.94 %).

Table 45. Usage pattern of fuel for domestic use in Yagapur Tanda-1 micro-watershed

Sl.No.	Particulars	LL (4)		MF (12)		SF (11)		SMF (7)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Fire Wood	3	75	9	75	7	63.6	6	85.7	25	73.53
2	Biogas	0	0	1	8.33	0	0	0	0	1	2.94
3	LPG	1	25	2	16.7	4	36.4	2	28.6	9	26.47

Source of drinking water: The data on source of drinking water in Yagapur Tanda-1 Micro watershed is presented in Table 46. The results indicated that, piped waters supply was the major source for drinking water for 58.82 per cent of the households followed by bore well water (41.18%).

Table 46. Source of drinking water in Yagapur Tanda-1 micro-watershed

Sl.No.	Particulars	LL (4)		MF (12)		SF (11)		SMF (7)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Piped supply	1	25	7	58.3	6	54.55	6	85.7	20	58.82
2	Bore Well	3	75	5	41.7	5	45.45	1	14.3	14	41.18

Source of light: The data on source of light in Yagapur Tanda-1 Micro watershed is presented in Table 47. The results indicated that, electricity was the major source of light for 100.00 per cent of the households.

Table 47. Source of light in Yagapur Tanda-1 micro-watershed

Sl.No.	Particulars	LL (4)		MF (12)		SF (11)		SMF (7)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Electricity	4	100	12	100	11	100	7	100	34	100

Existence of sanitary toilet facility: The data on availability of toilet facility in Yagapur Tanda-1 Micro watershed is presented in Table 48. The results indicated that, 47.06 per cent of the households possess toilets.

Table 48. Existence of sanitary toilet facility in Yagapur Tanda-1 micro-watershed

Sl.No.	Particulars	LL (4)		MF (12)		SF (11)		SMF (7)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Sanitary toilet facility	3	75	4	33	5	45.45	4	57	16	47.1

Possession of PDS card: The data regarding possession of PDS card in Yagapur Tanda-1 Micro watershed is presented in Table 49. The results indicated that, 100.00 per cent of the households possessed BPL card.

Table 49. Possession of PDS card in Yagapur Tanda-1 micro-watershed

Sl.No.	Particulars	LL (4)		MF (12)		SF (11)		SMF (7)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	BPL	4	100	12	100	11	100	7	100	34	100

Participation in NREGA programme: The data regarding Participation in NREGA programme in Yagapur Tanda-1 Micro watershed is presented in Table 50. The results indicated that, only 58.82 percent of the households have participated in NREGA programme.

Table 50. Participation in NREGA programme in Yagapur Tanda-1 micro-watershed

Sl.No.	Particulars	LL (4)		MF (12)		SF (11)		SMF (7)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Participation in NREGA programme	2	50	7	58.3	7	63.6	4	57.1	20	58.8

Adequacy of food items: The data regarding adequacy of food items in Yagapur Tanda-1 Micro watershed is presented in Table 51. The results indicated that, the extent of adequacy of food items for cereals, pulses, Oilseeds and vegetables were 61.76, 67.65, 23.53, 58.82 per cent respectively, similarly for Fruits (38.24%), milk (67.65%) and Egg (5.88%).

Table 51. Adequacy of food items in Yagapur Tanda-1 micro-watershed

Sl.No.	Particulars	LL (4)		MF (12)		SF (11)		SMF (7)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Cereals	2	50	8	66.7	8	72.73	3	42.9	21	61.76
2	Pulses	2	50	8	66.7	8	72.73	5	71.4	23	67.65
3	Oilseed	0	0	3	25	2	18.18	3	42.9	8	23.53
4	Vegetables	2	50	7	58.3	5	45.45	6	85.7	20	58.82
5	Fruits	1	25	5	41.7	6	54.55	1	14.3	13	38.24
6	Milk	2	50	9	75	9	81.82	3	42.9	23	67.65
7	Egg	0	0	0	0	1	9.09	1	14.3	2	5.88

Inadequacy of food items: The data regarding in adequacy of food items in Yagapur Tanda-1 Micro watershed is presented in Table 52. The results indicated that, the extent of in adequacy of food items for cereals, pulses, Oilseeds and vegetables were 44.12, 41.18, 76.47, 38.24 and 76.47 per cent respectively, similarly for fruits (38.24%), milk (29.41%), egg (70.59%) and meat (76.47%).

Table 52. Inadequacy of food items in Yagapur Tanda-1 micro-watershed

Sl.No.	Particulars	LL (4)		MF (12)		SF (11)		SMF (7)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Cereals	3	75	4	33.3	4	36.36	4	57.1	15	44.12
2	Pulses	2	50	4	33.3	4	36.36	4	57.1	14	41.18
3	Oilseed	3	75	9	75	10	90.91	4	57.1	26	76.47
4	Vegetables	1	25	5	41.7	6	54.55	1	14.3	13	38.24
5	Fruits	2	50	4	33.3	4	36.36	3	42.9	13	38.24
6	Milk	1	25	3	25	2	18.18	4	57.1	10	29.41
7	Egg	3	75	9	75	9	81.82	3	42.9	24	70.59
8	Meat	3	75	9	75	10	90.91	4	57.1	26	76.47

Farming constraints: The data regarding farming constraints experienced by households in Yagapur Tanda-1 Micro watershed is presented in Table 53. The results indicated that, lower fertility status of the soil was the constraint experienced by (94.12 %) per cent of the households, wild animal menace on farm field (91.18%), frequent incidence of pest and diseases (61.76%), inadequacy of irrigation water (88.24%), high cost of fertilizers and plant protection chemicals (76.47%), high rate of interest on credit (76.47%), low price for the agricultural commodities (64.71 %), lack of marketing facilities in the area (38.24%), inadequate extension services (67.65 %) and lack of transport for safe transport of the agricultural produce to the market (67.65%).

Table 53. Farming constraints experienced in Yagapur Tanda-1 micro-watershed

SN	Particulars	LL (4)		MF (12)		SF (11)		SMF (7)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Lower fertility status of the soil	0	0	12	100	12	109.09	7	100	32	94.12
2	Wild animal menace on farm field	0	0	12	100	11	100	7	100	31	91.18
3	Frequent incidence of pest and diseases	0	0	8	66.67	8	72.73	4	57.14	21	61.76
4	Inadequacy of irrigation water	0	0	12	100	11	100	7	100	30	88.24
5	High cost of Fertilizers and plant protection chemicals	0	0	9	75	10	90.91	7	100	26	76.47
6	High rate of interest on credit	0	0	9	75	11	100	6	85.71	26	76.47
7	Low price for the agricultural commodities	0	0	9	75	9	81.82	4	57.14	22	64.71
8	Lack of marketing facilities in the area	0	0	5	41.67	6	54.55	2	28.57	13	38.24
9	Inadequate extension services	0	0	9	75	10	90.91	4	57.14	23	67.65
10	Lack of transport for safe transport of the Agril produce to the market.	0	0	9	75	10	90.91	4	57.14	23	67.65

SUMMARY AND IMPLICATIONS

In order to assess the socio-economic condition of the farmers in the watershed 34 households located in the micro watershed were interviewed for the survey. The study was conducted in Yagapur Tanda-1 micro-watershed (Yaragal sub-watershed, Chithapura taluk & Kalaburagi District) is located at North latitude $16^{\circ} 56' 28.758''$ and $16^{\circ} 54' 29.936''$ and East longitude $77^{\circ} 7' 1.416''$ and $77^{\circ} 5' 27.747''$ covering an area of about 725.94 ha bounded by under Yagapura Village.

Socio-economic analysis of Yagapur Tanda-1 micro watersheds of Yaragal sub-watershed, Chithapura taluk & Kalaburagi District indicated that, out of the total sample of 34 farmers were sampled in Yagapur Tanda-1 micro-watershed among households surveyed 12 (35.29%) were marginal, 11 (32.35%) were small and 7 (20.59 %) were semi medium farmers. 4 landless farmers were also interviewed for the survey. The population characteristics of households indicated that, there were 90 (52.94%) men and 80 (47.06 %) were women. The average population of landless was 4.8, marginal farmers were 4.7, small farmers were 5.6 and semi medium farmers were 4.7. Majority of the respondents (44.12%) were in the age group of 16-35 years.

Education level of the sample households indicated that, there were 62.94 per cent illiterates, 37.05 per cent pre university education and 1.18 per cent attained graduation. About, 97.06 per cent of household heads practicing agriculture and 2.94 per cent of the household heads were engaged as agricultural labourers. Agriculture was the major occupation for 62.94 per cent of the household members.

In the study area, 85.29 per cent of the households possess katcha house and 8.82 per cent possess pucca house. The durable assets owned by the households showed that, 79.41 per cent possess TV, 41.18 per cent possess mixer grinder, 100.00 per cent possess mobile phones and 5.88 per cent possess motor cycles. Farm implements owned by the households indicated that, 23.53 per cent of the households possess plough, 2.94 per cent possess tractor, 14.71 per cent possess bullock cart and 8.82 per cent possess sprayer. Regarding livestock possession by the households, 2.94 per cent possess local cow. The average labour availability in the study area showed that, own labour men available in the micro watershed was 1.73, women available in the micro watershed was 1.67, hired labour (men) available was 12 and hired labour (women) available was 10.37.

Out of the total land holding of the sample respondents 97.14 per cent (42.49 ha) of the area is under dry condition and the remaining 2.86 per cent area is irrigated land. There were 1.00 live bore wells and 1.00 dry bore wells among the sampled households. Bore well was the major source of irrigation for 2.94 per cent of the households. The major crops grown by sample farmers are Red gram, Cotton, Jowar, Maize and Groundnut and cropping intensity was recorded as 100.00 per cent. Out of the sample

households 94.12 percent possessed bank account and 35.29 per cent of them have savings in the account.

About 100.00 per cent of the respondents borrowed credit from various sources. Among the credit borrowed by households, 100.00 per cent from co-operative/Grameena bank. Majority of the respondents (100.00%) have borrowed loan for agriculture purpose. The per hectare cost of cultivation for Red gram, Cotton, Jowar, Maize and Groundnut was Rs.28112.10 , 30197.34, 25900.72, 12693.85 and 51094.17 with benefit cost ratio of 1:1.20, 1: 1.22, 1: 0.80, 1: 2.90 and 1:0.70 respectively. Further, 20.59 per cent of the households opined that dry fodder was adequate and 5.88 per cent of the households have opined that the green fodder was adequate.

The average annual gross income of the farmers was Rs. 64638.24 in micro-watershed, of which Rs. 37991.18 comes from agriculture. Sampled households have grown 3 horticulture trees and 85 forestry trees together in the fields and back yards. Households have an average investment capacity of Rs. 1911.76 for land development. Source of funds for additional investment is concerned, 47.06 per cent depends on bank loan for land development activities.

Regarding marketing channels, 88.24 per cent of the households have sold agricultural produce to the local/village merchants. Further, 88.24 per cent of the households have used tractor for the transport of agriculture commodity. Majority of the farmers (55.88%) have experienced soil and water erosion problems in the watershed and 88.24 per cent of the households were interested towards soil testing. Fire was the major source of fuel for domestic use for 73.53 per cent of the households and 26.47 per cent households has LPG connection. Piped supply was the major source for drinking water for 58.82 per cent of the households. Electricity was the major source of light for 100.00 per cent of the households.

In the study area, 47.06 per cent of the households possess toilet facility. Regarding possession of PDS card, 100.00 per cent of the households possessed BPL card. Households opined that, the requirement of cereals (61.76%), pulses (67.65%) and oilseeds (23.53%) are adequate for consumption.

Farming constraints experienced by households in the micro watersheds were lower fertility status of the soil (94.12%) wild animal menace on farm field (91.18%), frequent incidence of pest and diseases (61.76%), inadequacy of irrigation water (88.24%), high cost of fertilizers and plant protection chemicals (76.47%), high rate of interest on credit (76.47%), low price for the agricultural commodities (64.71%), lack of marketing facilities in the area (38.24%), inadequate extension services (67.65%) and lack of transport for safe transport of the agricultural produce to the market (67.65%).

Implications of the survey

- ✓ Result indicated that, there were 62.94 per cent were illiterate hence, extension methodologies such as demonstration, street play, drama, video shows will be effective in dissemination of the technologies in the micro watershed.
- ✓ The data indicate that, 85.29 per cent of the households possess katcha house. Hence, the development department while implementing the watershed plan should focus on agriculture to enhance the productivity of major crops in the area to increase the income of the farmers.
- ✓ Results indicated that the local institutional participation of the household members in the micro watershed is minimal hence, activities like membership campaign, awareness creation about the benefits of membership in local institutions and strengths of organized groups must be conveyed.
- ✓ Majority of the households in the watershed have experience in use of mobile phones, and television hence, these mass media can be effectively utilized for transfer of technology as well as for information dissemination.
- ✓ The farm machinery/implement possession in the micro watershed was found to be minimum the reasons may lack of knowledge or lack of financial ability which can be addressed through training on use of different farm implements, providing information on different sources of finance for purchase of farm implements.
- ✓ The possession of livestock such as crossbred cow found is less hence, farmers must be made aware of the benefits of crossbred cow in increased milk production.
- ✓ The possession of livestock such as sheep, goat and poultry was found to be low hence, farmers may be informed the role of subsidiary enterprises in enhancing the income and information on financial support for subsidiary activities.
- ✓ The results indicate that there was a change in quality of life due to migration hence, the developmental departments should take actions to arrest migration and to improve the quality of the life in rural areas.
- ✓ Households possess 41.28ha (97.14 %) of dry land and 1.21ha (2.86 %) of irrigated land hence, the availability of the dryland agricultural technologies such as short duration crops, high yielding drought resistance crop varieties, drip irrigation technology and subsidy information will be helpful for the farmers to enhance the productivity of land and as well as farmers income.
- ✓ Few of the bore well in micro watershed found non functional hence, farmers may be trained on possibility of bore well rejuvenation.
- ✓ Bore well was major source of irrigation for 2.94 per cent of the households. hence, in order to increase the area under irrigation as well as to increase the water use efficiency farmers may trained on drip irrigation and provide the information on subsidy for drip irrigation equipment's along with the information on different agencies which provides the financial assistance for drip irrigation.

- ✓ The cropping intensity in the micro watershed was found to be (100.00 %) hence, care must be taken by the implementing agency to bring uncultivated land into cultivation through suitable measures.
- ✓ Many of the household members have borrowed loan from cooperative banks which has higher rate of interest hence, farmers may be sensitized on the different sources of credit with lesser interest rate such SHGs etc.
- ✓ The results indicated the non availability of both green and dry fodder throughout the year hence, fodder development activities can be taken up in the micro watershed.
- ✓ The average annual gross income of the households Rs.37991.18 from agriculture, and Rs. 26647.06 from wages. Agriculture was found to be the major source of income for households hence; the development activities should focus on productivity enhancement, marketing arrangements and agricultural technology dissemination to have a direct impact on the farmers.
- ✓ The cultivation of forest species is found minimal hence, information and production technology related to agro-forestry and integrated farming system.
- ✓ The data indicated that, 55.88 per cent of the households have experienced soil and water erosion problems. Hence, those farmers who reported the soil and water erosion problems may be given attention while implementation of the watershed development plan.
- ✓ The data indicated that, 88.24 per cent of the households have interest in soil testing hence, farmers must be provided with the information on various institutions which are involved in soil testing for the benefit of the farmers.
- ✓ Except summer ploughing the adoption of other soil and water conservation structures is minimum hence, the farmers in the micro watershed should be sensitized on the use of different conservation structures for soil water conservation.
- ✓ Cereals and pulses found be adequate for per cent of the households respectively hence, farm households and the farm women must be trained on importance of balanced nutrition and role of vegetable, milk, egg, meat in balanced diet.
- ✓ Lower fertility status of the soil (94.12%), wild animal menace on farm field (91.18%), frequent incidence of pest and diseases (61.76%), high cost of fertilizers and plant protection chemicals (76.47%), high rate of interest on credit (76.47%), low price for the agricultural commodities (64.71%), lack of marketing facilities in the area (38.24%), inadequate extension services (67.65%), lack of transport for safe transport of the agricultural produce to the market (67.65%) were the major farming constraints experienced hence, these constraints must be addressed immediately for the welfare of the farmers. Awareness to be created among the farmers to approach nearest KVKs/RSKs and other developmental departments for technical and for subsidized inputs and utilize the well established regulated markets, approaching the contract firms, direct markets to avoid the involvement of middlemen.