



हर कदम, हर डगर  
किसानों का हमसफर  
भारतीय कृषि अनुसंधान परिषद

*Agrisearch with a human touch*

**LAND RESOURCE INVENTORY AND SOCIO-ECONOMIC STATUS OF  
FARM HOUSEHOLDS FOR WATERSHED PLANNING AND  
DEVELOPMENT**

**HITNAL (4D3A9D2d) MICRO WATERSHED**

**Koppal Taluk and District, Karnataka**

**Karnataka Watershed Development Project – II**

**SUJALA – III**

**World Bank funded Project**



**THE WORLD BANK**



**ICAR – NATIONAL BUREAU OF SOIL SURVEY AND LAND USE PLANNING**



**WATERSHED DEVELOPMENT DEPARTMENT  
GOVT. OF KARNATAKA, BANGALORE**



ICAR - NBSS & LUP



## **About ICAR - NBSS&LUP**

The ICAR-National Bureau of Soil Survey and Land Use Planning (ICAR-NBSS&LUP), Nagpur, a premier Institute of the Indian Council of Agricultural Research (ICAR), was set up during 1976 with the objective to prepare soil resource maps at national, state and district levels and to provide research inputs in soil resource mapping and its applications, land evaluation, land use planning, land resource management, and database management using GIS for optimising land use on different kinds of soils in the country.

The Bureau has been engaged in carrying out soil resource survey, agro-ecological and soil degradation mapping at the country, state and district levels for qualitative assessment and monitoring the soil health towards viable land use planning. The research activities have resulted in identifying the soil potentials and problems, and the various applications of the soil surveys with the ultimate objective of sustainable agricultural development. The Bureau has the mandate to correlate and classify soils of the country and maintain a National Register of all the established soil series. The Institute is also imparting in-service training to staff of the soil survey agencies in the area of soil survey, land evaluation and soil survey interpretations for land use planning. The Bureau in collaboration with Panjabrao Krishi Vidyapeeth, Akola is running post-graduate teaching and research programme in land resource management, leading to M.Sc. and Ph.D. degrees.

**Citation:** Rajendra Hegde, Ramesh Kumar, S.C., K.V. Niranjana, S. Srinivas, M.Lalitha, B.A. Dhanorkar, R.S. Reddy and S.K. Singh (2019). "Land resource inventory and socio-economic status of farm households for watershed planning and development of Hitnal (4D3A9D2d) Microwatershed, Koppal Taluk and District, Karnataka", ICAR-NBSS&LUP Sujala MWS Publ.566 ICAR – NBSS & LUP, RC, Bangalore. p.129 & 29.

### **TO OBTAIN COPIES,**

**Please write to:**

**Director, ICAR - NBSS & LUP,**

Amaravati Road, NAGPUR - 440 033, India

Phone : (0712) 2500386, 2500664, 2500545 (O)

Telefax : 0712-2522534

E-Mail : [director@nbsslup.ernet.in](mailto:director@nbsslup.ernet.in)

Website URL : [nbsslup.in](http://nbsslup.in)

Or

**Head, Regional Centre, ICAR - NBSS&LUP, Hebbal, Bangalore - 560 024**

Phone : (080) 23412242, 23510350 (O)

Telefax : 080-23510350

E-Mail : [nbssrcb@gmail.com](mailto:nbssrcb@gmail.com)



# LAND RESOURCE INVENTORY AND SOCIO-ECONOMIC STATUS OF FARM HOUSEHOLDS FOR WATERSHED PLANNING AND DEVELOPMENT

HITNAL (4D3A9D2d) MICRO WATERSHED

Koppal Taluk and District, Karnataka

Karnataka Watershed Development Project – II

Sujala-III

World Bank funded Project



THE WORLD BANK



ICAR – NATIONAL BUREAU OF SOIL SURVEY AND LAND USE  
PLANNING



ICAR - NBSS & LUP



WATERSHED DEVELOPMENT DEPARTMENT, GOVT. OF  
KARNATAKA, BANGALORE





## PREFACE

In Karnataka, as in other Indian States, the livelihoods of rural people are intertwined with farming pursuits. The challenges in agriculture are seriously threatening the livelihood of a large number of farmers as they have been practicing farming in contextual factors beyond their control. Climatic factors are the most important ones and have become much more significant in recent times due to rapid climate changes induced by intensive anthropogenic activities affecting our ecosystem in multiple ways. Climate change has become the reality, it is happening and efforts to evolve and demonstrate climate resilient technologies have become essential. Due to the already over stressed scenario of agrarian sector, the climate change is resulting in manifold increase in the complexities, pushing the rural mass to face more and more unpredictable situations. The rising temperatures and unpredictable rainfall patterns are going to test seriously the informed decisions farmers have to make in order to survive in farming and sustain their livelihood.

It is generally recognized that impacts of climate change shall not be uniform across the globe. It is said that impact of climate change is more severe in South Asia. Based on the analysis of meteorological data, it is predicted that in India, there will be upward trend in mean temperature, downward trend in relative humidity, annual rainfall and number of wet days in a year. Also, in general, phenomena like erratic monsoon, spread of tropical diseases, rise in sea levels, changes in availability of fresh water, frequent floods, droughts, heat waves, storms and hurricanes are predicted. Each one of these adverse situations are already being experienced in various parts of India and also at the global level. Decline in agricultural productivity of small and marginal farmers becoming more vulnerable is already witnessed.

In Karnataka, more than 60 per cent of the population live in rural areas and depend on agriculture and allied activities for their livelihood. Though the state has achieved significant progress in increasing the yield of many crops, there is tremendous pressure on the land resources due to the growing and competing demands of various land uses. This is reflected in the alarming rate of land degradation observed. Already more than 50 per cent of the area is affected by various forms of degradation. If this trend continues, the sustainability of the fragile ecosystem will be badly affected. The adverse effects of change in the climatic factors are putting additional stress on the land resources and the farmers dependent on this.

The natural resources (land, water and vegetation) of the state need adequate and constant care and management, backed by site-specific technological interventions and investments particularly by the government. Detailed database pertaining to the nature of

the land resources, their constraints, inherent potentials and suitability for various land based rural enterprises, crops and other uses is a prerequisite for preparing location-specific action plans, which are in tune with the inherent capability of the resources. Any effort to evolve climate resilient technologies has to be based on the baseline scientific database. Then only one can expect effective implementation of climate resilient technologies, monitor the progress, make essential review of the strategy, and finally evaluate the effectiveness of the implemented programs. The information available at present on the land resources of the state are of general nature and useful only for general purpose planning. Since the need of the hour is to have site-specific information suitable for farm level planning and detailed characterization and delineation of the existing land resources of an area into similar management units is the only option.

ICAR-NBSS&LUP, Regional Centre, Bangalore has taken up a project sponsored by the Karnataka Watershed Development Project-II, (Sujala-III), Government of Karnataka funded by the World Bank under Component -1 Land Resource Inventory. This study was taken up to demonstrate the utility of such a database in reviewing, monitoring and evaluating all the land based watershed development programs on a scientific footing. To meet the requirements of various land use planners at grassroots level, the present study on “Land Resource Inventory and Socio-Economic Status of Farm Households for Watershed Planning and Development of for Hitnal microwatershed in Koppal Taluk and District, Karnataka” for integrated development was taken up in collaboration with the State Agricultural Universities, IISC, KRSRAC, KSNDMC as Consortia partners. The project provides detailed land resource information at cadastral level (1:7920 scale) for all the plots and socio-economic status of farm households covering thirty per cent farmers randomly selected representing landed and landless class of farmers in the micro-watershed. The project report with the accompanying maps for the microwatershed will provide required detailed database for evolving effective land use plan, alternative land use options and conservation plans for the planners, administrators, agricultural extension personnel, KVK officials, developmental departments and other land users to manage the land resources in a sustainable manner.

It is hoped that this database will be useful to the planners, administrators and developmental agencies working in the area in not only for formulating location specific developmental schemes but also for their effective monitoring at the village/watershed level.

Nagpur

Date:19-11-2019

**S.K. SINGH**

Director, ICAR - NBSS&LUP , Nagpur

## Contributors

<b>Dr. Rajendra Hegde</b> Principal Scientist, Head & Project Leader, Sujala-III Project ICAR-NBSS&LUP, Regional Centre, Bangalore	<b>Dr. S.K.Singh</b> Director, ICAR-NBSS&LUP Coordinator, Sujala-III Project Nagpur
<b>Soil Survey, Mapping &amp; Report Preparation</b>	
Dr. K.V. Niranjana	Sh. R.S. Reddy
Dr. B.A. Dhanorkar	Smt. Chaitra, S.P.
	Dr. Gopali Bardhan
	Mr. Somashekar T.N
	Ms. Arpitha G.M
	Dr. Mahendra kumar M.B
<b>Field Work</b>	
Sh. C. Bache Gowda	Sh. Mayur Patil
Sh. Somashekar	Sh. Arun Kumar, S.
Sh. M. Jayaramaiah	Sh. Sunil Raj
	Sh. Yogesh Kumar, B.
	Sh. Vikas, N.K.
	Sh. Arun Kumar, S.G.
	Sh. Umesh Jadiyahappa Madolli
	Sh. Praveen Kumar P. Achalkar
	Sh. Veerabhadraswamy
	Sh. Vinay
	Sh. Shankarappa, K.
	Sh. Lankesh, R.S.
	Sh. Appanna B. Hattigoudar
	Sh. Maharudra
<b>GIS Work</b>	
Dr. S.Srinivas	Sh. A.G.Devendra Prasad
Sh. D.H.Venkatesh	Sh. Abhijith Sastry, N.S.
Smt. K.Sujatha	Smt. Shyla, B.
Smt. K.V.Archana	Smt. Swetha ,K.
Sh. N.Maddileti	Ms. Vidya, P.C.
	Sh. Deepak, M.J.
	Smt. K.Karunya Lakshmi
	Ms. Seema, K.V.

<b>Laboratory Analysis</b>	
Dr. M. Lalitha	Sh. Vindhya, N.G.
Smt. Arti Koyal	Ms. P. Pavanakumari, P.
Smt. Parvathy, S.	Ms. Rashmi, N.
	Ms. Leelavathy, K.U.
	Smt. Usha Kiran, G.
<b>Socio-Economic Analysis</b>	
Dr. S.C. Ramesh Kumar	Sh. M.K. Prakashanaik
	Ms. Karuna V. Kulkarni
	Mrs. Sowmya A.N
	Sh. Vinod R
	Sh. Basavaraja
	Sh. Vijay Kumar Lamani
	Ms. Sowmya K.B
	Mrs. Prathibha, D.G
	Sh. Rajendra,D
<b>Soil &amp; Water Conservation</b>	
Sh. Sunil P. Maske	
<b>Watershed Development Department, GoK, Bangalore</b>	
Sh. Prabhash Chandra Ray, IFS Project Director & Commissioner, WDD	Dr. A. Natarajan NRM Consultant, Sujala-III Project
Sh. A. Padmaya Naik, Director (In-Charge) Executive Director, KWDP-II, Sujala-III, WDD	



# **PART-A**

## **LAND RESOURCE INVENTORY**



## Contents

Preface		
Contributors		
Executive Summary		
Chapter 1	Introduction	1
Chapter 2	Geographical Setting	3
2.1	Location and Extent	3
2.2	Geology	3
2.3	Physiography	4
2.4	Drainage	5
2.5	Climate	5
2.6	Natural Vegetation	6
2.7	Land Utilization	7
Chapter 3	Survey Methodology	9
3.1	Base maps	9
3.2	Image Interpretation for Physiography	9
3.3	Field Investigation	12
3.4	Soil mapping	14
3.5	Land Management Units	14
3.6	Laboratory Characterization	14
Chapter 4	The Soils	19
4.1	Soils of Granite Gneiss Landscape	19
4.2	Soils of Alluvial Landscape	22
Chapter 5	Interpretation for Land Resource Management	29
5.1	Land Capability Classification	29
5.2	Soil Depth	31
5.3	Surface Soil Texture	32
5.4	Soil Gravelliness	33
5.5	Available Water Capacity	34
5.6	Soil Slope	35
5.7	Soil Erosion	36
Chapter 6	Fertility Status	39
6.1	Soil Reaction (pH)	39
6.2	Electrical Conductivity (EC)	39
6.3	Organic Carbon (OC)	39
6.4	Available Phosphorus	41
6.5	Available Potassium	41
6.6	Available Sulphur	41
6.7	Available Boron	42
6.8	Available Iron	42
6.9	Available Manganese	42
6.10	Available Copper	42
6.11	Available Zinc	42

Chapter 7	Land Suitability for Major Crops	47
7.1	Land suitability for Sorghum	47
7.2	Land suitability for Maize	48
7.3	Land suitability for Bajra	49
7.4	Land suitability for Redgram	50
7.5	Land suitability for Bengalgram	51
7.6	Land suitability for Groundnut	52
7.7	Land suitability for Sunflower	53
7.8	Land suitability for Cotton	54
7.9	Land suitability for Chilli	55
7.10	Land suitability for Tomato	56
7.11	Land suitability for Brinjal	57
7.12	Land suitability for Onion	58
7.13	Land suitability for Bhendi	59
7.14	Land suitability for Drumstick	60
7.15	Land suitability for Mulberry	61
7.16	Land suitability for Mango	62
7.17	Land Suitability for Sapota	63
7.18	Land suitability for Pomegranate	64
7.19	Land suitability for Guava	65
7.20	Land Suitability for Jackfruit	66
7.21	Land Suitability for Jamun	67
7.22	Land Suitability for Musambi	68
7.23	Land Suitability for Lime	69
7.24	Land Suitability for Cashew	70
7.25	Land Suitability for Custard apple	71
7.26	Land suitability for Amla	72
7.27	Land suitability for Tamarind	73
7.28	Land suitability for Marigold	74
7.29	Land suitability for Chrysanthemum	75
7.30	Land suitability for Jasmine	76
7.31	Land suitability for Crossandra	77
7.32	Land Management Units	111
7.33	Proposed Crop Plan	112
Chapter 8	Soil Health Management	115
Chapter 9	Soil and Water conservation Treatment Plan	121
9.1	Treatment Plan	121
9.2	Recommended Soil and Water Conservation measures	125
9.3	Greening of microwatershed	126
	References	129
	Appendix I	I-VIII
	Appendix II	IX-XIV
	Appendix III	XV-XX

## LIST OF TABLES

2.1	Mean Monthly Rainfall, PET, 1/2 PET at Koppal Taluk and District	5
2.2	Land Utilization in Koppal District	7
3.1	Differentiating Characteristics used for Identifying Soil Series	13
3.2	Soil map unit description of Hitnal microwatershed	15
4.1	Physical and chemical characteristics of soil series identified in Hitnal microwatershed	24
7.1	Soil-Site Characteristics of Hitnal microwatershed	79
7.2	Land suitability for Sorghum	80
7.3	Land suitability for Maize	81
7.4	Land suitability for Bajra	82
7.5	Land suitability for Redgram	83
7.6	Land suitability for Bengalgram	84
7.7	Land suitability for Groundnut	85
7.8	Land suitability for Sunflower	86
7.9	Land suitability for Cotton	87
7.10	Land suitability for Chilli	88
7.11	Land suitability for Tomato	89
7.12	Land suitability for Brinjal	90
7.13	Land suitability for Onion	91
7.14	Land suitability for Bhendi	92
7.15	Land suitability for Drumstick	93
7.16	Land suitability for Mulberry	94
7.17	Land suitability for Mango	95
7.18	Land Suitability for Sapota	96
7.19	Land suitability for Pomegranate	97
7.20	Land suitability for Guava	98
7.21	Land suitability for Jackfruit	99
7.22	Land suitability for Jamun	100
7.23	Land Suitability for Musambi	101
7.24	Land Suitability for Lime	102

7.25	Land Suitability for Cashew	103
7.26	Land Suitability for Custard apple	104
7.27	Land Suitability for Amla	105
7.28	Land Suitability for Tamarind	106
7.29	Land Suitability for Marigold	107
7.30	Land Suitability for Chrysanthemum	108
7.31	Land suitability for Jasmine	109
7.32	Land suitability for Crossandra	110
7.33	Proposed Crop Plan for Hitnal Microwatershed	113

## LIST OF FIGURES

2.1	Location map of Hitnal Microwatershed	3
2.2a	Granite and granite gneiss rocks	4
2.2b	Alluvial rocks	4
2.3	Rainfall distribution in Koppal Taluk, Koppal District	6
2.4	Natural vegetation of Hitnal microwatershed	6
3.1	Scanned and Digitized Cadastral map of Hitnal Microwatershed	11
3.2	Satellite image of Hitnal Microwatershed	11
3.3	Cadastral map overlaid on IRS PAN+LISS IV merged imagery of Hitnal Microwatershed	12
3.4	Location of profiles in a transect	13
3.5	Soil phase or management units of Hitnal Microwatershed	17
5.1	Land Capability Classification of Hitnal Microwatershed	30
5.2	Soil Depth map of Hitnal Microwatershed	31
5.3	Surface Soil Texture map of Hitnal Microwatershed	33
5.4	Soil Gravelliness map of Hitnal Microwatershed	34
5.5	Soil Available Water Capacity map of Hitnal Microwatershed	35
5.6	Soil Slope map of Hitnal Microwatershed	36
5.7	Soil Erosion map of Hitnal Microwatershed	37
6.1	Soil Reaction (pH) map of Hitnal Microwatershed	40
6.2	Electrical Conductivity (EC) map of Hitnal Microwatershed	40
6.3	Soil Organic Carbon (OC) map of Hitnal Microwatershed	41
6.4	Soil Available Phosphorus map of Hitnal Microwatershed	43
6.5	Soil Available Potassium map of Hitnal Microwatershed	43
6.6	Soil Available Sulphur map of Hitnal Microwatershed	44
6.7	Soil Available Boron map of Hitnal Microwatershed	44
6.8	Soil Available Iron map of Hitnal Microwatershed	45
6.9	Soil Available Manganese map of Hitnal Microwatershed	45
6.10	Soil Available Copper map of Hitnal Microwatershed	46
6.11	Soil Available Zinc map of Hitnal Microwatershed	46
7.1	Land suitability map of Sorghum	48
7.2	Land suitability map of Maize	49
7.3	Land suitability map of Bajra	50

7.4	Land suitability map of Redgram	51
7.5	Land suitability map of Bengalgram	52
7.6	Land suitability map of Groundnut	53
7.7	Land suitability map of Sunflower	54
7.8	Land suitability map of Cotton	55
7.9	Land suitability map of Chilli	56
7.10	Land suitability map of Tomato	57
7.11	Land suitability map of Brinjal	58
7.12	Land suitability map of Onion	59
7.13	Land suitability map of Bhendi	60
7.14	Land suitability map of Drumstick	61
7.15	Land suitability map of Mulberry	62
7.16	Land suitability map of Mango	63
7.17	Land Suitability map of Sapota	64
7.18	Land suitability for Pomegranate	65
7.19	Land suitability map of Guava	66
7.20	Land Suitability map of Jackfruit	67
7.21	Land Suitability map of Jamun	68
7.22	Land Suitability map of Musambi	69
7.23	Land Suitability map of Lime	70
7.24	Land Suitability map of Cashew	71
7.25	Land Suitability map of Custard apple	72
7.26	Land suitability map of Amla	73
7.27	Land suitability map of Tamarind	74
7.28	Land suitability map of Marigold	75
7.29	Land suitability map of Chrysanthemum	76
7.30	Land suitability map of Jasmine	77
7.31	Land suitability map of Crossandra	78
7.32	Land Management Units map of Hitnal microwatershed	111
9.1	Soil and water conservation map of Hitnal microwatershed	126



## **EXECUTIVE SUMMARY**

*The land resource inventory of Hitnal microwatershed was conducted using village cadastral maps and IRS satellite imagery on 1:7920 scale. The false colour composites of IRS imagery were interpreted for physiography and these physiographic delineations were used as base for mapping soils. The soils were studied in several transects and a soil map was prepared with phases of soil series as mapping units. Random checks were made all over the area outside the transects to confirm and validate the soil map unit boundaries. The soil map shows the geographic distribution and extent, characteristics, classification, behavior and use potentials of the soils in the Microwatershed.*

*The present study covers an area of 311 ha in Koppal taluk and district, Karnataka. The climate is semiarid and categorized as drought - prone with an average annual rainfall of 662 mm, of which about 424 mm is received during south –west monsoon, 161 mm during north-east and the remaining 77 mm during the rest of the year. An area of about 34 per cent is covered by soil, 8 per cent by rockout crops, 3 per cent by Mining/ Industrial area, 12 per cent by water bodies, settlements, <1 per cent by Railway and 42 per cent of the area is not surveyed. The salient findings from the land resource inventory are summarized briefly below.*

- ❖ *The soils belong to 5 soil series and 7 soil phases (management units) and 4 land management units.*
- ❖ *The length of crop growing period is <90 days and starts from 2<sup>nd</sup> week of August to 2<sup>nd</sup> week of November.*
- ❖ *From the master soil map, several interpretative and thematic maps like land capability, soil depth, surface soil texture, soil gravelliness, available water capacity, soil slope and soil erosion were generated.*
- ❖ *Soil fertility status maps for macro and micronutrients were generated based on the surface soil samples collected at every 320 m grid interval.*
- ❖ *Land suitability for growing 31 major agricultural and horticultural crops were assessed and maps showing the degree of suitability along with constraints were generated.*
- ❖ *Entire area is suitable for agriculture.*
- ❖ *About 15 per cent of the soils are moderately shallow (50-75 cm), 19 per cent moderately deep (75- 100 cm) and 1 per cent is very deep (100->150cm) soils.*
- ❖ *About 33 per cent loamy (sandy loam and sandy clay loam) and 2 per cent has clayey (sandy clay and clay) soils at the surface.*
- ❖ *About 1 per cent of the area has non-gravelly (<15%) and 34 per cent gravelly (15-35%) soils.*

- ❖ *With respect to available water capacity 15 per cent of the area has very low (<50mm/m), 19 per cent of the area has low (51-100 mm/m) and 1 per cent very high (>200mm/m) in available water capacity.*
- ❖ *An area of about 1 per cent has nearly level (0-1%) and 34 per cent has very gently sloping (1-3%) lands.*
- ❖ *An area of about 1 per cent is slightly eroded (e1) and 34 per cent is moderately eroded (e2).*
- ❖ *An area of about 8 per cent is neutral (pH 6.5 to 7.3), 5 per cent is slightly alkaline (pH 7.3 to 7.8), 21 per cent moderately alkaline (pH 7.8 to 8.4), <1 per cent is strongly alkaline (pH 8.4-9.0) and <1 per cent is very strongly alkaline (pH>9.0).*
- ❖ *The Electrical Conductivity (EC) of the soils are <2 dsm<sup>-1</sup> indicating that soils are non saline.*
- ❖ *Organic carbon is medium (0.5-0.75%) in 11 per cent and high (>0.75%) in 25 per cent area of the soils.*
- ❖ *Available phosphorus is low (<23 kg/ha) in 6 per cent, medium (<23 kg/ha) in 18 per cent and high (>57 kg/ha) in 11 per cent area of the soils.*
- ❖ *Available potassium is low (<145 kg/ha) in 7 per cent, medium (145-337 kg/ha) in 27 per cent and high (>337 kg/ha) in 1 per cent area of the soils.*
- ❖ *Available sulphur is low (<10 ppm) in 5 per cent, medium (10-20 ppm) in 17 per cent and high (>20 ppm) in 13 per cent area of the soils.*
- ❖ *Available boron is low (<0.5 ppm) in 30 per cent, medium (0.5-1.0 ppm) in 4 per cent and high (>1.0 ppm) in 1 per cent area of the microwatershed.*
- ❖ *Available iron is deficient (<4.5ppm) in 17 per cent and sufficient (>4.5 ppm) in 19 per cent of the area.*
- ❖ *Available zinc is deficient (<0.6 ppm) in 7 per cent and sufficient (>0.6 ppm) in 29 per cent area of the microwatershed.*
- ❖ *Available manganese and copper are sufficient in the entire area.*
- ❖ *The land suitability for 31 major agricultural and horticultural crops grown in the microwatershed was assessed and the areas that are highly suitable (class S1) and moderately suitable (class S2) are given below. It is however to be noted that a given soil may be suitable for various crops but what specific crop to be grown may be decided by the farmer looking to his capacity to invest on various inputs, marketing infrastructure, market price, and finally the demand and supply position.*

**Land suitability for various crops in the microwatershed**

Crop	Suitability Area in ha (%)		Crop	Suitability Area in ha (%)	
	Highly suitable (S1)	Moderately suitable (S2)		Highly suitable (S1)	Moderately suitable (S2)
Sorghum	4(1)	59(19)	Sapota	-	59(19)
Maize	-	63(20)	Pomegranate	-	63(20)
Bajra	-	110(35)	Guava	-	59(19)
Redgram	-	63(20)	Jackfruit	-	59(19)
Bengal gram	4(1)	-	Jamun	-	4(1)
Groundnut	-	106(34)	Musambi	4(1)	59(19)
Sunflower	4(1)	59(19)	Lime	4(1)	59(19)
Cotton	4(1)	59(19)	Cashew	-	59(19)
Chilli	-	63(20)	Custard apple	4(1)	106(34)
Tomato	-	59(19)	Amla	-	110(35)
Brinjal	-	59(19)	Tamarind	-	4(1)
Onion	-	59(19)	Marigold	-	63(20)
Bhendi	-	63(20)	Chrysanthemum	-	63 (20)
Drumstick	-	63(20)	Jasmine	-	59(19)
Mulberry	-	59(19)	Crossandra	-	59(19)
Mango	-	-	-	-	-

- ❖ *Apart from the individual crop suitability, a proposed crop plan has been prepared for the 4 identified LMUs by considering only the highly and moderately suitable lands for different crops and cropping systems with food, fodder, fibre and other horticulture crops.*
- ❖ *Maintaining soil-health is vital for crop production and conserve soil and land resource base for maintaining ecological balance and to mitigate climate change. For this, several ameliorative measures have been suggested to these problematic soils like saline/alkali, highly eroded, sandy soils etc.,*
- ❖ *Soil and water conservation and drainage line treatment plans have been prepared that would help in identifying the sites to be treated and also the type of structures required.*
- ❖ *As part of the greening programme, several tree species have been suggested to be planted in marginal and submarginal lands, field bunds and also in the hillocks, mounds and ridges. That would help in supplementing the farm income, provide fodder and fuel, and generate lot of biomass which in turn would help in maintaining the ecological balance and contribute to mitigating the climate change.*



## **INTRODUCTION**

Land is a scarce resource and basic unit for any material production. It can support the needs of the growing population, provided they use the land in a rational and judicious manner. But what is happening in many areas of the state is a cause for concern to everyone involved in the management of land resources at the grassroots level. The area available for agriculture is about 51 per cent of the total area and more than 60 per cent of the people are still dependant on agriculture for their livelihood. The limited land area is under severe stress and strain due to increasing population pressure and competing demands of various land uses. Due to this, every year there is significant diversion of farm lands and water resources for non-agricultural purposes. Apart from this, due to lack of interest in farmers for farming, large tracts of cultivable lands are turning into fallows in many areas and this trend is continuing at an alarming rate.

Further, land degradation has emerged as a serious problem which has already affected about 38 lakh ha of cultivated area in the state. Soil erosion alone has degraded about 35 lakh ha. Almost all the uncultivated areas are facing various degrees of degradation, particularly soil erosion. Salinity and alkalinity has emerged as a major problem in more than 3.5 lakh ha in the irrigated areas of the state. Nutrient depletion and declining factor productivity is common in both rainfed and irrigated areas. The degradation is continuing at an alarming rate and there appears to be no systematic effort among the stakeholders to contain this process. In recent times, an aberration of weather due to climate change phenomenon has added another dimension leading to unpredictable situations to be tackled by the farmers.

In this critical juncture, the challenge before us is not only to increase the productivity per unit area which is steadily declining and showing a fatigue syndrome, but also to prevent or at least reduce the severity of degradation. If the situation is not reversed at the earliest, then the sustainability of the already fragile crop production system and the overall ecosystem will be badly affected in the state. The continued neglect and unscientific use of the resources for a long time has led to the situation observed at present in the state. It is a known fact and established beyond doubt by many studies in the past that the cause for all kinds of degradation is the neglect and irrational use of the land resources. Hence, there is urgent need to generate a detailed site-specific farm level database on various land resources for all the villages/watersheds in a time bound manner that would help to protect the valuable soil and land resources and also to stabilize the farm production.

Therefore, the land resource inventory required for farm level planning is the one which investigates not only the surface but also consider the other parameters which are critical for productivity *viz.*, soils, climate, water, minerals and rocks, topography, geology, hydrology, vegetation, crops, land use pattern, animal population, socio-

economic conditions, infrastructure, marketing facilities and various schemes and developmental works of the government etc. From the data collected at farm level, the specific problems and potentials of the area can be identified and highlighted, conservation measures required for the area can be planned on a scientific footing, suitability of the area for various uses can be worked out and finally viable and sustainable land use options suitable for each and every land holding can be prescribed.

The Land Resource Inventory is basically done for identifying potential and problem areas, developing sustainable land use plans, estimation of surface run off and water harvesting potential, preparation of soil and water conservation plans, land degradation/desertification etc. The Bureau is presently engaged in developing an LRI methodology using high resolution satellite remote sensing data and Digital Elevation Model (DEM) data to prepare Landscape Ecological Units (LEU) map representing agro-ecosystem as a whole. The LEU is preferred over landform as the base map for LRI. LEU is the assemblage of landform, slope and land use. An attempt was made to upscale the soil resource information from 1:250000 and 1:50000 scale to the LEU map in Goa and other states.

The land resource inventory aims to provide site-specific database for Hitnal Microwatershed in Koppal Taluk, Koppal District, Karnataka State for the Karnataka Watershed Development Department. The database was generated by using cadastral map of the village as a base along with high resolution IRS LISS IV and Cartosat-1 merged satellite imagery. Later, an attempt will be made to uplink this LRI data generated at 1:7920 scale under Sujala-III Project to the proposed Landscape Ecological Units (LEUs) map.

The study was organized and executed by the ICAR- National Bureau of Soil Survey and Land Use Planning, Regional Centre, Bangalore under Generation of Land Resource Inventory Data Base Component-1 of the Sujala-III Project funded by the World Bank.

## GEOGRAPHICAL SETTING

### 2.1 Location and Extent

The Hitnal micro-watershed is located in the central part of Karnataka in Koppal taluk and district (Fig 2.1). It lies between  $15^{\circ} 26'$  and  $15^{\circ} 28'$  North latitudes and  $76^{\circ} 19'$  and  $76^{\circ} 21'$  East longitudes, and covers an area of about 311 ha. It comprises parts of Chikkakasanakandi, Bevinahalli, Hitnala, Agalakeri and Shapura villages. It is about 23 km from Koppal town and is bounded by Bevinahalli on the west, Chikkakasanakandi on the south, Shapura and Agalakeri on the north and Hitnala on the east and western side of the microwatershed.




Fig.2.1 Location map of Hitnal Microwatershed

### 2.2 Geology

Major rock formations observed in the microwatershed are granite gneiss and alluvium (Fig.2.2 a and b). Granite gneisses are essentially pink to gray and are coarse to

medium grained. They consist primarily of quartz, feldspar, biotite and hornblende. The gray granite gneisses are highly weathered, fractured and fissured upto a depth of about 10 m. Dolerite dykes and quartz veins are common with variable width and found to occur in Hitnal village. The thickness of the alluvium generally is limited to less than a meter, except in river valleys where it is very deep extending to tens of meters. Such soils are transported and represent paleo black soils originally formed at higher elevation, but now occupying river valleys.



Fig.2.2 a Granite and granite gneiss rocks



Fig.2.2 b Alluvium

### 2.3 Physiography

Physiographically, the area has been identified as Granite gneiss and Alluvial landscapes based on geology. The microwatershed area has been further divided into



mounds/ridges, summits, side slopes and very gently sloping uplands and nearly level plains based on slope and its relief features. The elevation ranges from 482 to 511 m in the gently sloping uplands. The mounds and ridges are mostly covered by rock outcrops.

## 2.4 Drainage

The area is drained by several small seasonal streams that join Hire *halla* and Chenna *halla* along its course. Though, the streams are not perennial, during rainy season they carry large quantities of rain water. The microwatershed has only few small tanks which are not able to store the water flowing during the rainy season. Due to this, the ground water recharge is very much affected in the villages. This is reflected in the failure of many bore wells in the villages. If the available rain water is properly harnessed by constructing tanks and recharge structures at appropriate places in the villages, then the drinking and irrigation needs of the area can be easily met. The drainage network is dendritic to sub parallel.

## 2.5 Climate

The district falls under semiarid tract of the state and is categorized as drought - prone with total annual rainfall of 662 mm (Table 2.1). Of this, a maximum of 424 mm precipitation is received during south–west monsoon period from June to September, north-east monsoon contributes about 161 mm and prevails from October to early December and the remaining 77 mm is received during the rest of the year. The winter season is from December to February. During April and May, the temperatures reach up to 45°C and in December and January, the temperatures will go down to 16°C. Rainfall distribution is shown in Figure 2.3. The average Potential Evapo Transpiration (PET) is 145 mm and varies from a low of 101 mm in December to 193 mm in the month of May. The PET is always higher than precipitation in all the months except in the month of September. Generally, the Length of crop Growing Period (LGP) is <90 days and starts from 2<sup>nd</sup> week of August to 2<sup>nd</sup> week of November.

**Table 2.1 Mean Monthly Rainfall, PET, 1/2 PET at Koppal Taluk and District**

Sl. no.	Months	Rainfall	PET	1/2 PET
1	January	1.60	116.70	58.35
2	February	1.50	129.20	64.60
3	March	14.10	169.80	84.90
4	April	18.10	180.60	90.30
5	May	41.60	193.50	96.75
6	June	85.80	167.90	83.95
7	July	72.10	156.20	78.10
8	August	110.50	152.50	76.25
9	September	155.60	138.50	69.25
10	October	116.30	122.30	61.15
11	November	36.00	106.40	53.20
12	December	9.10	101.00	50.50
<b>TOTAL</b>		<b>662.30</b>	<b>144.55</b>	




Fig. 2.3 Rainfall distribution in Koppal Taluk and District

## 2.6 Natural Vegetation

The natural vegetation is sparse comprising few tree species, shrubs and herbs. The mounds, ridges and boulders occupy sizeable areas which are under thin to moderately thick forest vegetation. Still, there are some remnants of the past forest cover which can be seen in patches in some ridges and hillocks in the microwatershed (Fig 2.4).

Apart from the continuing deforestation, the presence of large population of goats, sheep and other cattle in the microwatershed is causing vegetative degradation of whatever little vegetation left in the area. The uncontrolled grazing has left no time for the regeneration of the vegetative cover. This leads to the accelerated rate of erosion on the hill slopes, resulting in the formation of deep gullies in the foot slopes and eventually resulting in the heavy siltation of few tanks and reservoirs in the microwatershed.



Fig 2.4 Natural vegetation of Hitnal Microwatershed

## 2.7 Land Utilization

About 91 per cent area (Table 2.2) in Koppal district is cultivated at present and about 17 per cent of the area is sown more than once. An area of about 3 per cent is currently barren. Forests occupy a small area of about 5 per cent and the tree cover is in a very poor state. Most of the mounds, ridges and boulder areas have very poor vegetative cover.

**Table 2.2 Land Utilization in Koppal District**

Sl. No.	Agricultural land use	Area ( ha)	Per cent
1	Total geographical area	552495	
2	Total cultivated area	500542	90.6
3	Area sown more than once	92696	16.8
4	Trees and groves	210	0.04
5	Cropping intensity	-	118
6	Forest	29451	5.33
7	Cultivable wasteland	2568	0.46
8	Permanent Pasture land	14675	2.66
9	Barren land	16627	3.01
10	Non agricultural land	40591	7.35
11	Current fallow	19660	3.56



## SURVEY METHODOLOGY

The purpose of land resource inventory is to delineate similar areas (soil series and phases), which respond or expected to respond similarly for a given level of management. This was achieved in Hitnal microwatershed by the detailed study of all the soil characteristics (depth, texture, colour, structure, consistence, coarse fragments, porosity, soil reaction, soil horizons etc.) and site characteristics (slope, erosion, drainage, occurrence of rock fragments etc.) followed by grouping of similar areas based on soil-site characteristics into homogeneous (management units) units and showing their extent and geographic distribution on the microwatershed cadastral map. The detailed soil survey at 1:7920 scale was carried out in 311 ha area. The methodology followed for carrying out land resource inventory was as per the guidelines given in Soil Survey Manual (IARI, 1971; Soil Survey Staff, 2006; Natarajan *et al.*, 2015) which is briefly described below.

### 3.1 Base Maps

The detailed survey of the land resources occurring in the microwatershed was carried out by using digitized cadastral map and satellite imagery as base supplied by the KRSRSAC. The cadastral map shows field boundaries with their survey numbers, location of tanks, streams and other permanent features of the area (Fig. 3.1). Apart from the cadastral map, remote sensing data products from Cartosat-1 and LISS IV merged at the scale of 1:7920 were used in conjunction with the cadastral map to identify the geology, landscapes, landforms and other surface features. The imagery helped in the identification and delineation of boundaries between hills, uplands and lowlands, water bodies, forest and vegetated areas, roads, habitations and other cultural features of the area (Fig.3.2). The cadastral map was overlaid on the satellite imagery (Fig.3.3) that helps to identify the parcel boundaries and other permanent features. Apart from cadastral maps and images, toposheets of the area (1:50,000 scale) were used for initial traversing, identification of geology, landscapes and landforms, drainage features, present land use and also for selection of transects in the microwatershed.

### 3.2 Image Interpretation for Physiography

False Colour Composites (FCC) of Cartosat-I and LISS-IV merged satellite data covering the microwatershed area was visually interpreted using image interpretation elements and all the available collateral data with local knowledge. The delineated physiographic boundaries were transferred on to a cadastral map overlaid on satellite imagery. Physiographically, the area has been identified as granite gneiss and alluvial landscapes and is divided into landforms such as ridges, mounds and uplands based on slope. They were further subdivided into physiographic/ image interpretation units based on image characteristics. The image interpretation legend for physiography is given below.

## **Image Interpretation Legend for Physiography**

### **G- Granite gneiss landscape**

- G1 Hills/ Ridges/ Mounds
  - G11 Summits
  - G12 Side slopes
    - G121 Side slopes with dark grey tones
- G2 Uplands
  - G21 Summits
  - G22 Gently sloping uplands
    - G221 Gently sloping uplands, yellowish green (eroded)
    - G222 Gently sloping uplands, yellowish white (severely eroded)
  - G23 Very gently sloping uplands
    - G231 Very gently sloping uplands, yellowish green
    - G232 Very gently sloping uplands, medium green and pink
    - G233 Very gently sloping uplands, pink and green (scrub land)
    - G234 Very gently sloping uplands, medium greenish grey
    - G235 Very gently sloping uplands, yellowish white (eroded)
    - G236 Very gently sloping uplands, dark green
    - G237 Very gently sloping uplands, medium pink (coconut garden)
    - G238 Very gently sloping uplands, pink and bluish white (eroded)

### **DSe -Alluvial landscape**

#### **DSe 1 Summit**

- DSe 11 Nearly level Summit with dark grey tone
- DSe 12 Nearly level Summit with medium grey tone
- DSe 13 Nearly level Summit with whitish grey tone
- DSe 14 Nearly level Summit with whitish tone (Calcareousness)
- DSe 15 Nearly level Summit with pinkish grey tone
- DSe 16 Nearly level Summit with medium pink tone
- DSe 17 Nearly level Summit with bluish white tone
- DSe 18 Nearly level Summit with greenish grey tone

#### **DSe 2 Very gently sloping**

- DSe 21 Very gently sloping, whitish tone
- DSe 22 Very gently sloping, greyish pink tone
- DSe 23 Very gently sloping, whitish grey tone
- DSe 24 Very gently sloping, medium grey tone
- DSe 25 Very gently sloping, medium pink tone
- DSe 26 Very gently sloping, dark grey tone
- DSe 27 Very gently sloping, bluish grey tone
- DSe 28 Very gently sloping, greenish grey tone
- DSe 29 Very gently sloping, Pinkish grey




Fig 3.1 Scanned and Digitized Cadastral map of Hitnal Microwatershed




Fig.3.2 Satellite Image of Hitnal Microwatershed




Fig.3.3 Cadastral map overlaid on IRS PAN+LISS IV merged imagery of Hitnal Microwatershed

### 3.3 Field Investigation

The field boundaries and survey numbers given on the cadastral sheet were located on the ground by following permanent features like roads, cart tracks, *nallas*, streams, tanks etc., and wherever changes were noticed, they were incorporated on the microwatershed cadastral map. Preliminary traverse of the microwatershed was carried out with the help of cadastral map, imagery and toposheets. While traversing, landforms and physiographic units identified were checked and preliminary soil legend was prepared by studying soils at few selected places. Then, intensive traversing of each physiographic unit like hills, ridges, uplands and plains was carried out. Based on the variability observed on the surface, transects (Fig 3.4) were selected across the slope covering all the landform units in the microwatershed (Natarajan and Dipak Sarkar, 2010).






Fig: 3.4. Location of profiles in a transect

In the selected transect, soil profiles (Fig.3.4) were located at closely spaced intervals to take care of any change in the land features like break in slope, erosion, gravel, stones etc. In the selected sites, profiles (vertical cut showing the soil layers from surface to the rock) were opened upto 200 cm or to the depth limited by rock or hard substratum and studied in detail for all their morphological and physical characteristics. The soil and site characteristics were recorded for all profile sites on a standard proforma as per the guidelines given in USDA Soil Survey Manual (Soil Survey Staff, 2012). Apart from the transect study, profiles were also studied at random, almost like in a grid pattern, outside the transect areas to validate the soil map unit boundaries.

Based on the soil characteristics, the soils were grouped into different soil series. Soil series is the most homogeneous unit having similar horizons and properties and behaves similarly for a given level of management. Soil depth, texture, colour, kind of horizon and horizon sequence, amount and nature of gravel present, calcareousness, nature of substratum etc, were used as the major differentiating characteristics for identifying soil series occurring in the area. The differentiating characteristics used for identifying the soil series are given in Table 3.1. Based on the above characteristics, 5 soil series were identified in Hitnal microwatershed.

**Table 3.1 Differentiating Characteristics used for identifying Soil Series  
(Characteristics are of Series Control Section)**

Soils of Granite Gneiss Landscape							
Sl. No	Soil Series	Depth (cm)	Colour (moist)	Texture	Gravel (%)	Horizon sequence	Calcareousness
1	<b>Lakkur (LKR)</b>	50-75	2.5YR 2.5/3, 2.5/4, 3/4, 3/6	gsc	40-60	Ap-Bt- Bc-Cr	-
2	<b>Mukhadahalli (MKH)</b>	50-75	5YR3/3,3/4,4/3, 5/4,6/6 2.5YR3/4	gsc	>35	Ap-Bt-Cr	-
3	<b>Hatti (HTI)</b>	50-75	5 YR 3/3, 3/4,	gsc	15-35	Ap-Bt-Cr	-
4	<b>Hooradhahalli</b>	75-100	2.5YR2.5/4,3/4,	gsc-gc	>35	Ap-Bt-Cr	-

	(HDH)		3/6				
<b>Soils of Alluvial Landscape</b>							
5	<b>Kadagathur (KDT)</b>	>150	10 YR 3/1, 3/2, 3/3, 7.5YR 3/3, 3/4	sc-c	-	Ap-Bw	-

### 3.4 Soil Mapping

The area under each soil series was further separated into soil phases and their boundaries delineated on the cadastral map based on the variations observed in the texture of the surface soil, slope, erosion, presence of gravel, stoniness etc. A soil phase is a subdivision of soil series based mostly on surface features that affect its use and management. The soil mapping units are shown on the map (Fig.3.5) in the form of symbols. During the survey many soil profile pits, few mini pits and a few auger bores representing different landforms occurring in the microwatershed were studied. In addition to the profile study, spot observations in the form of mini pits, road cuts, terrace cuts etc., were studied to validate the soil boundaries on the soil map.

The soil map shows the geographic distribution of 7 mapping units representing 5 soil series occurring in the microwatershed. The soil map unit (soil legend) description is presented in Table 3.2. The soil phase map (management units) shows the distribution of 7 phases mapped in the microwatershed. Each mapping unit (soil phase) delineated on the map has similar soil and site characteristics. In other words, all the farms or survey numbers included in one soil phase will have similar management needs and have to be treated accordingly.

### 3.5 Land Management Units

The 7 soil phases identified and mapped in the microwatershed were regrouped into four Land Management Units (LMU's) for the purpose of preparing a Proposed Crop Plan for sustained development of the microwatershed. The database (soil phases) generated under LRI was utilized for identifying Land Management Units (LMU's) based on the management needs. One or more than one soil site characteristic having influence on the management have been chosen for identification and delineation of LMUs. For Hitnal microwatershed, five soil and site characteristics, namely the soil depth, soil texture, slope, erosion and gravel content have been considered for defining LMUs. The land management units are expected to behave similarly for a given level of management.

### 3.6 Laboratory Characterization

Soil samples for each soil series soil were collected from representative master profiles for laboratory characterization by following the methods outlined in the Laboratory Manual (Sarma *et al*, 1987). Surface soil samples collected in the year 2018 from farmer's fields in Hitnal microwatershed for fertility status (major and micronutrients) at 320 m grid interval were analyzed in the laboratory (Katyal and Rattan,

2003). By linking the soil fertility data to the survey numbers through GIS, soil fertility maps were generated using Kriging method for the microwatershed.

**Table 3.2 Soil map unit description of Hitnal Microwatershed**

Soil map unit No*	Soil Series	Soil Phase Symbol	Mapping Unit Description	Area in ha (%)
<b>Soils of Granite gneiss Landscape</b>				
	LKR		Lakkur soils are moderately shallow (50-75 cm), well drained, have dark reddish brown to dark red, red gravelly sandy clay soils occurring on very gently to moderately sloping uplands under cultivation	<b>32 (10.14)</b>
43		LKRcB2g1	Sandy loam surface, slope 1-3%, moderate erosion, gravelly (15-35%)	27 (8.66)
452		LKRhB2g1	Sandy clay loam surface, slope 1-3%, moderate erosion, gravelly (15-35%)	5 (1.48)
	MKH		Mukhadahalli soils are moderately shallow (50-75 cm), well drained, have dark brown to reddish brown, red gravelly sandy clay soils occurring on gently very gently to gently sloping uplands under cultivation	<b>16 (4.99)</b>
85		MKHhB2g1	Sandy clay loam surface, slope 1-3%, moderate erosion, gravelly (15-35%)	16 (4.99)
	HTI		Hatti soils are moderately shallow (50-75 cm), well drained, have dark reddish brown, red gravelly sandy clay soils occurring on nearly level to very gently sloping uplands under cultivation	<b>0.31 (0.1)</b>
100		HTIiB2	Sandy clay surface, slope 1-3%, moderate erosion	0.31(0.1)
	HDH		Hooradhahalli soils are moderately deep (75-100 cm), well drained, have dark red to dark reddish brown, red gravelly sandy clay to clay soils occurring on nearly level to moderately sloping uplands under cultivation	<b>59 (18.98)</b>
111		HDHcB2g1	Sandy loam surface, slope 1-3%, moderate erosion, gravelly (15-35%)	57 (18.33)
128		HDHiB2g1	Sandy clay surface, slope 1-3%, moderate erosion, gravelly (15-35%)	2 (0.65)
<b>Soils of Alluvial Landscape</b>				
	KDT		Kadagathur soils are very deep (>150 cm), moderately well drained, have dark brown to very dark grayish brown, sandy clay to clay black soils occurring on nearly level to very gently sloping plains under cultivation	<b>4 (1.33)</b>
403		KDTmA1	Clay surface, slope 0-1%, slight erosion	4 (1.33)
992		Railway		<b>2 (0.65)</b>
993		Not Surveyed		<b>129 (41.53)</b>
994		Mining/Industrial	Mining/Industrial area	<b>9 (2.96)</b>
999		Rock outcrops	Rock lands, both massive & bouldery with little or no soil	<b>24 (7.7)</b>
1000		Others	Habitation and Waterbody	<b>36 (11.61)</b>

\*Soil map unit numbers are continuous for the taluk, not the microwatersheds






Fig 3.5 Soil Phase or Management Units- Hitnal Microwatershed



## THE SOILS

Detailed information pertaining to the nature, extent and distribution of different kinds of soils occurring in Hitnal microwatershed is provided in this chapter. The microwatershed area has been identified as granite gneiss and alluvial landscapes based on geology. In all, 5 soil series were identified. Soil formation is the result of the combined effect of environmental and terrain factors that are reflected in soil morphology. The soil formation is dominantly influenced by the parent material, climate, time and relief.

A brief description of each of the 5 soil series identified followed by 7 soil phases (management units) mapped (Fig. 3.5) are furnished below. The physical and chemical characteristics of soil series identified in Hitnal microwatershed are given in Table 4.1 along with soil classification. The soils in any one map unit differ from place to place in their depth, texture, slope, gravelliness, erosion or any other site characteristic that affect management. The soil phase map can be used for identifying the suitability of areas for growing specific crops or for other alternative uses and also for deciding the type of conservation structures needed. The detailed information on soil and site-characteristics like soil depth, surface soil texture, slope, erosion, gravelliness, AWC, LCC etc, with respect to each of the soil phase identified is given village/survey number wise for the microwatershed in Appendix-I.

### 4.1 Soils of Granite gneiss Landscape

In this landscape, four soil series were identified and mapped. Of these series, Hooradhahalli (HDH) series occupies a maximum area of 59 ha (19 %) and others occupy minor area. The brief description of the soil series along with the soil phases identified and mapped is given below.

**4.1.1 Lakkur (LKR) Series:** Lakkur soils are moderately shallow (50-75cm), well drained, have reddish brown to dark red gravelly sandy clay soils. They have developed from weathered granite gneiss and occur on nearly level to very gently and gently sloping uplands. The Lakkur series has been classified as a member of the clayey-skeletal, mixed, isohyperthermic family of Typic Rhodustalfs.

The thickness of the solum ranges from 51 to 74 cm. The thickness of A horizon ranges from 12 to 18 cm. Its colour is in 5YR and 2.5 YR hue with value 3 to 4 and chroma 4 to 6. The texture varies from loamy sand to sandy clay loam with 15 to 50 per cent gravel. The thickness of B horizon ranges from 39 to 58 cm. Its colour is in 2.5 YR hue with value 3 to 4 and chroma 4 to 6. Texture is sandy clay with 40 to 60 per cent gravel. The available water capacity is low (50-100 mm/m). Two soil phases were identified and mapped.



Landscape and soil profile characteristics of Lakkur (LKR) Series

**4.1.2 Mukhadahalli (MKH) Series:** Mukhadahalli soils are moderately shallow (50-75 cm), well drained, have dark brown to reddish brown gravelly sandy clay soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands. The Mukhadahalli series has been classified as a member of the clayey-skeletal, mixed, isohyperthermic family of Typic Haplustalfs.

The thickness of the solum ranges from 51 to 72 cm. The thickness of A horizon ranges from 12 to 17 cm. Its colour is in 5 YR and 7.5 YR hue with value 3 to 4 and chroma 2 to 4. The texture varies from loamy sand to sandy loam with 20 to 45 per cent gravel. The thickness of B horizon ranges from 40 to 68 cm. Its colour is in 2.5 YR and 5 YR hue with value and chroma 3 to 6. Texture is sandy clay loam to sandy clay with 35 to 50 per cent gravel. The available water capacity is low (50-100 mm/m). One soil phase was identified and mapped.



Landscape and soil profile characteristics of Mukhadahalli (MKH) Series



**4.1.3 Hatti (HTI) Series:** Hatti soils are moderately shallow (50-75cm), well drained, have dark reddish brown, gravelly sandy clay soils. They are developed from weathered granite gneiss and occur on very gently sloping uplands under cultivation. The Hatti series has been classified as a member of the fine, mixed, isohyperthermic Typic Paleustalfs.

The thickness of the solum ranges from 57 to 74 cm. The thickness of A horizon ranges from 16 to 20 cm. Its colour is in 5 YR hue with value and chroma 3 to 4. The texture varies from sandy loam to sandy clay loam and sandy clay with 15 to 60 per cent gravel. The thickness of B horizon ranges from 45 to 56 cm. Its colour is in 5 YR hue with value 3 and chroma 3 to 4. Texture is sandy clay with 15 to 35 per cent gravel. The available water capacity is low (50-100 mm/m). One soil phase was identified and mapped.



Landscape and soil profile characteristics of Hatti (HTI) Series

**4.1.4 Hooradhahalli (HDH) Series:** Hooradhahalli soils are moderately deep (75-100 cm), well drained, have red to dark red and reddish brown gravelly sandy clay to clay soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands. The Hooradhahalli series has been classified as a member of the clayey-skeletal, mixed, isohyperthermic family of Rhodic Paleustalfs.

The thickness of the solum ranges from 76 to 100 cm. The thickness of A horizon ranges from 11 to 19 cm. Its colour is in 5 YR and 2.5 YR hue with value 3 to 4 and chroma 3 to 6. The texture varies from loamy sand to sandy clay with 15 to 50 per cent gravel. The thickness of B horizon varies from 65 to 83 cm. Its colour is in 2.5 YR hue with value 2.5 to 3 and chroma 4 to 6. Texture is sandy clay to clay with 35 to 50 per cent gravel. The available water capacity is low (50-100mm/m). Two soil phases were identified and mapped.



Landscape and soil profile characteristics of Hooradhahalli (HDH) Series

#### 4.2 Soils of Alluvial Landscape

In this landscape, only one soil series was identified and mapped. The brief description of the soil series along with the soil phases identified and mapped is given below.

**4.2.1 Kadagathur (KDT) Series:** Kadagathur soils are very deep (>150 cm), moderately well drained, have dark brown to very dark grayish brown, sandy clay to clay soils. They have developed from alluvium and occur on nearly level to very gently sloping plains under cultivation. The Kadagathur series has been classified as a member of the fine, mixed, isohyperthermic family of Fluventic Haplustepts.

The thickness of the solum is more than 150 cm. The thickness of A horizon ranges from 8 to 14 cm. Its colour is in 10 YR hue with value 3 and chroma 4. The texture varies is sandy loam. The thickness of B horizon is more than 150 cm. Its colour is in 10 YR and 7.5 YR hue with value 3 and chroma 1 to 4. Its texture is sandy clay to clay. The available water capacity is very high (>200 mm/m). One soil phase was identified and mapped.



Landscape and soil profile characteristics of Kadagathur (KDT) Series

**Table: 4.1 Physical and Chemical Characteristics of Soil Series identified in Hitnal microwatershed**

**Soil Series:** Lakkur (LKR), **Pedon:** RM-8.

**Location:** 15°04'26.3"N, 75°37'84.1"E, (4D4A3I1f), Belhatti village, Shirahatti taluk, Gadag district

**Analysis at:** NBSS&LUP, Regional Centre, Bengaluru **Classification:** Clayey-skeletal, mixed, isohyperthermic Typic Rhodustalfs

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-21	Ap	74.00	8.34	17.66	9.62	11.57	15.76	23.13	13.92	20	sl	-	-
21-35	Bt	54.37	10.48	35.14	16.33	8.64	9.69	11.59	8.11	40	sc	-	-
35-56	Bc	48.37	13.46	38.17	10.96	7.69	9.17	11.28	9.27	60	sc	-	-

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO <sub>3</sub>	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP						
	Water	CaCl <sub>2</sub>	M KCl				dS m <sup>-1</sup>	%	%	Ca	Mg					K	Na	Total	cmol kg <sup>-1</sup>	%	%
0-21	8.18	-	-	0.30	0.56	0.94	-	-	0.31	0.55	0.86	12.19	0.69	100.00	4.51						
21-35	8.17	-	-	0.30	0.52	1.29	-	-	0.19	0.84	1.03	22.18	0.63	100.00	3.79						
35-56	7.95	-	-	0.46	0.48	1.99	-	-	0.24	0.58	0.82	22.94	0.60	100.00	2.53						

*Contd...*

**Series Name:** Mukahadahalli (MKH), **Pedon:** R-11

**Location:** 15°22'05.4"N, 76°04'10.3"E, Halageri village, Koppal Taluk and District

**Analysis at:** NBSS&LUP, Regional Centre, Bangalore. **Classification:** Clayey-skeletal, mixed, isohyperthermic Typic Haplustalfs

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-19	Ap	65.71	8.83	25.46	9.27	9.06	14.42	21.52	11.43	70	scl	16.54	8.60
19-32	Bt	55.89	11.13	32.98	6.47	9.18	11.89	19.19	9.18	50	scl	19.24	12.78
32-58	Bt	47.95	10.41	41.63	17.52	3.78	9.13	9.55	7.97	50	sc	24.03	16.02

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO <sub>3</sub>	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP			
	Water	CaCl <sub>2</sub>	M KCl				dS m <sup>-1</sup>	%	%	Ca	Mg					K	Na	Total
										cmol kg <sup>-1</sup>								
0-19	7.38	-	-	0.09	0.2	0.00	8.97	4.32	0.26	0.22	13.77	14.84	0.58	93	1.49			
19-32	7.5	-	-	0.106	0.41	0.00	15.98	3.27	0.16	0.50	19.91	20.88	0.63	95	2.38			
32-58	7.46	-	-	0.173	0.49	0.00	19.71	4.53	0.23	1.32	25.79	25.76	0.62	100	5.11			

Contd...

**Series Name:** Hatti (HTI), **Pedon:** R-20

**Location:** 15°21'45"N, 76°03'06" E Lakshmapura village Koppal taluk and district

**Analysis at:** NBSS&LUP, Regional Centre, Bangalore. **Classification:** Fine, mixed, isohyperthermic Typic Paleustalfs

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-16	Ap	65.33	12.19	22.48	13.79	11.32	13.37	18.31	8.54	15-20	scl	16.83	5.49
16-41	Bt1	41.54	14.04	44.42	6.47	6.26	9.50	13.36	5.95	15-20	c	27.26	16.64
41-64	Bt2	48.71	8.48	42.81	26.06	7.55	5.38	6.31	3.41	55-60	sc	27.22	12.63

Depth (cm)	pH (1:2.5)			E.C. (1:2.5) dS m <sup>-1</sup>	O.C. %	CaCO <sub>3</sub> %	Exchangeable bases					CEC	CEC/Clay	Base saturation %	ESP %
	Water	CaCl <sub>2</sub>	M KCl				Ca	Mg	K	Na	Total				
0-16	7.11			0.109	0.92		21.06	8.23	0.39	0.06	29.74	20.19	0.90	147	0.30
16-41	7.54			0.220	0.92		21.93	8.47	0.23	0.27	30.90	31.31	0.70	99	0.85
41-64	7.82			0.168	0.55		19.43	7.09	0.31	0.47	27.30	26.57	0.62	103	1.77

Contd...

**Soil Series:** Hooradhahalli (HDH), **Pedon:** RM-69

**Location:** 13°24'31"N, 76°33'41"E, (4D3D8G2d), Hesarahalli village, Chikkanayakanahalli taluk, Tumukura district

**Analysis at:** NBSS&LUP, Regional Centre, Bengaluru **Classification:** Clayey-skeletal, mixed isohyperthermic Rhodic Paleustalfs

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-18	Ap	72.56	15.17	12.27	4.57	8.33	17.38	23.88	18.39	35	sl	-	-
18-33	Bt1	56.29	10.75	32.96	7.88	10.24	13.41	14.43	10.34	55	scl	-	-
33-58	Bt2	46.66	10.79	42.55	10.79	9.87	8.43	9.04	8.53	55	sc	-	-
58-90	Bt3	43.09	13.63	43.27	9.90	8.25	7.32	8.76	8.87	45	c	-	-

Depth (cm)	pH (1:2.5)			E.C. (1:2.5) dS m <sup>-1</sup>	O.C. %	CaCO <sub>3</sub> %	Exchangeable bases					CEC	CEC/Clay	Base saturation %	ESP %
	Water	CaCl <sub>2</sub>	M KCl				Ca	Mg	K	Na	Total				
0-18	6.54	-	-	0.07	0.60	0.00	2.68	1.38	0.44	0.42	4.91	5.84	0.48	84.07	7.11
18-33	5.90	-	-	0.07	0.52	0.00	3.99	1.27	0.09	0.37	5.71	8.61	0.26	66.32	4.29
33-58	6.16	-	-	0.07	0.44	0.00	4.92	1.67	0.08	0.55	7.22	10.00	0.24	72.23	5.50
58-90	6.39	-	-	0.06	0.40	0.00	4.30	2.02	0.08	0.46	6.87	9.21	0.21	74.61	5.05

Contd...

**Series Name:** Kadagathur (KDT), **Pedon :** R-7

**Location:** 15°26'48"N, 76°09'51" E Budashettynala village, Koppal taluk and district

**Analysis at:** NBSS&LUP, Regional Centre, Bangalore. **Classification:** Fine, mixed, isohyperthermic Fluventic Haplustepts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-12	Ap	75.90	8.77	15.33	17.33	18.36	14.36	15.90	9.95	-	sl	10.66	5.33
12-37	A2	62.54	11.35	26.11	8.46	20.54	13.31	12.07	8.15	-	scl	15.61	8.22
37-71	Bw1	52.73	10.51	36.77	6.08	18.24	12.47	9.01	6.92	-	sc	19.66	11.21
71-93	Bw2	33.26	22.65	44.09	3.13	12.53	7.78	5.18	4.64	-	c	30.08	17.34
93-118	Bw3	31.01	24.57	44.42	2.04	10.41	8.26	6.01	4.29	-	c	34.92	18.16
118-170	Bw4	38.31	18.73	42.96	2.99	14.62	10.35	6.30	4.06	-	c	46.06	19.59

Depth (cm)	pH (1:2.5)			E.C. (1:2.5) dS m <sup>-1</sup>	O.C. %	CaCO <sub>3</sub> %	Exchangeable bases					CEC	CEC/Clay	Base saturation %	ESP %
	Water	CaCl <sub>2</sub>	M KCl				Ca	Mg	K	Na	Total				
0-12	6.95	-	-	0.17	1.28	0.39	9.17	2.76	0.10	0.08	12.11	12.10	0.79	100.09	0.65
12-37	7.55	-	-	0.17	0.40	0.40	8.36	4.51	0.08	0.40	13.35	13.30	0.51	100.37	3.02
37-71	7.60	-	-	0.21	0.44	0.39	10.67	8.19	0.10	0.74	19.70	19.10	0.52	103.12	3.88
71-93	8.26	-	-	0.28	0.72	1.56	14.97	12.13	0.12	3.07	30.29	29.40	0.67	103.01	10.45
93-118	8.44	-	-	0.58	0.68	1.17	13.32	10.77	0.13	4.76	28.98	28.50	0.64	101.68	12.40
118-170	9.06	-	-	0.64	0.44	1.17	8.92	8.14	0.23	12.32	29.61	28.60	0.67	103.53	37.27



## INTERPRETATION FOR LAND RESOURCE MANAGEMENT

The most important soil and site characteristics that affect the land use and conservation needs of an area are land capability, land irrigability, soil depth, soil texture, coarse fragments, available water capacity, soil slope, soil erosion, soil reaction etc. These are interpreted from the data base generated through land resource inventory and several thematic maps are generated. These would help in identifying the areas suitable for growing crops and, soil and water conservation measures and structures needed thus helping to maintain good soil health for sustained crop production. The various thematic maps generated are described below.

### 5.1 Land Capability Classification

Land capability classification is an interpretative grouping of soil map units (soil phases) mainly based on inherent soil characteristics, external land features and environmental factors that limit the use of land for agriculture, pasture, forestry, or other uses on a sustained basis (IARI, 1971). The land and soil characteristics used to group the land resources in an area into various land capability classes, subclasses and units are

*Soil characteristics:* Soil depth, soil texture, coarse fragments, soil reaction, available water capacity, calcareousness, salinity/alkali *etc.*

*Land characteristics:* Slope, erosion, drainage, rock outcrops.

*Climate:* Total rainfall and its distribution, and length of crop growing period.

The Land Capability Classification system is divided into land capability classes, subclasses and units based on the level of information available. Eight land capability classes are recognized. They are

*Class I:* They are very good lands that have no limitations or very few limitations that restrict their use.

*Class II:* They are good lands that have minor limitations and require moderate conservation practices.

*Class III:* They are moderately good lands that have severe limitations that reduce the choice of crops or that require special conservation practices.

*Class IV:* They are fairly good lands that have very severe limitations that reduce the choice of crops or that require very careful management.

*Class V:* Soils in these lands are not likely to erode, but have other limitations like wetness that are impractical to remove and as such not suitable for agriculture, but suitable for pasture or forestry with minor limitations.

*Class VI:* The lands have severe limitations that make them generally unsuitable for cultivation, but suitable for pasture or forestry with moderate limitations.

*Class VII:* The lands have very severe limitations that make them unsuitable for cultivation, but suitable for pasture or forestry with major limitations.

*Class VIII:* Soil and other miscellaneous areas (rock lands) that have very severe limitations that nearly preclude their use for any crop production, but suitable for wildlife, recreation and installation of wind mills.

The land capability subclasses are recognized based on the dominant limitations observed within a given land capability class. The subclasses are designated by adding a lower case letter like ‘e’, ‘w’, ‘s’, or ‘c’ to the class numeral. The subclass “e” indicates that the main hazard is risk of erosion, “w” indicates drainage or wetness as a limitation for plant growth, “s” indicates shallow soil depth, coarse or heavy textures, calcareousness, salinity/alkalinity or gravelliness and “c” indicates limitation due to climate.

The land capability subclasses have been further subdivided into land capability units based on the kinds of limitations present in each subclass. Ten land capability units are used in grouping the soil map units. They are stony or rocky (0), erosion hazard (slope, erosion) (1), coarse texture (sand, loamy sand, sandy loam) (2), fine texture (cracking clay, silty clay) (3), slowly permeable subsoil (4), coarse underlying material (5), salinity/alkali (6), stagnation, overflow, high ground water table (7), soil depth (8) and fertility problems (9). The capability units thus identified have similar soil and land characteristics that respond similarly to a given level of management. The soils of the microwatershed have been classified upto land capability subclass level.

The 7 soil map units identified in the Hitnal Microwatershed are grouped under two land capability classes and three land capability subclasses (Fig. 5.1).




Fig. 5.1 Land Capability map of Hitnal Microwatershed

Entire cultivated area in the microwatershed is suitable for agriculture. Good lands (Class II) cover an area of about 63 ha (20%) and distributed in the major part of the microwatershed with minor problems of soil and erosion. Moderately good lands (Class III) occupy an area of about 47 ha (15%) and distributed in the southern and central part of the microwatershed with severe limitations of soil and erosion. An area of about 129 ha (42%) is not surveyed, 9 ha (3%) is covered by Mining/Industrial area, 2 ha (<1%) is under railway crops, 24 ha (8%) is under rock out crops and 36 ha (12%) is covered by habitation and water body.

## 5.2 Soil Depth

Soil depth refers to the depth of the soil occurring above the parent material or hard rock. The depth of the soil determines the effective rooting depth for plants and in accordance with soil texture, mineralogy and gravel content, the capacity of the soil column to hold water and nutrient availability. Soil depth is one of the most important soil characteristic that is used in differentiating soils into different soil series. The soil depth classes used in identifying soils in the field are very shallow (<25 cm), shallow (25-50 cm), moderately shallow (50-75 cm), moderately deep (75-100 cm), deep (100-150 cm) and very deep (>150 cm). They were used to classify the soils into different depth classes and a soil depth map was generated (Fig. 5.2). The area extent and their geographical distribution in the microwatershed is given in Fig. 5.2.




Fig. 5.2 Soil Depth map of Hitnal Microwatershed

Moderately shallow (50-75 cm) soils cover an area of about 47 ha (15%) and distributed in the southern and central part of the microwatershed. An area of about 59 ha (19%) is moderately deep soils (75-100 cm) and distributed in the central part of the microwatershed. Very deep (>150 cm) soils occupy an area of about 4 ha (1%) and distributed in the northern part of the microwatershed.

The most productive lands cover about 4 ha (1%) where all climatically adopted long duration crops be grown.

### **5.3 Surface Soil Texture**

Texture is an expression to indicate the coarseness or fineness of the soil as determined by the relative proportion of primary particles of sand, silt and clay. It has a direct bearing on the structure, porosity, adhesion and consistence. The surface layer of a soil to a depth of about 25 cm is the layer that is most used by crops and plants. The surface soil textural class provides a guide to understanding soil-water retention and availability, nutrient holding capacity, infiltration, workability, drainage, physical and chemical behavior, microbial activity and crop suitability. The textural classes used for LRI were used to classify and a surface soil texture map was generated. The area extent and their geographical distribution in the microwatershed is shown in Fig 5.3.

An area of about 104 ha (33%) is loamy (sandy loam and sandy clay loam) at the surface and distributed in the major part of the microwatershed. Clayey (sandy clay and clay) soils cover about 6 ha (2%) and are distributed in the northern and southern part of the microwatershed.

The most productive lands with respect to surface soil texture are clayey soils that (2%) have high potential for soil-water retention and availability and nutrient retention and availability, but have more problems of drainage, infiltration, workability and other physical problems. The other productive lands are loamy (33%) soils which also have high potential for soil- water retention and nutrient availability but have no drainage or other physical problems.




Fig. 5.3 Surface Soil Texture map of Hitnal Microwatershed

#### 5.4 Soil Gravelliness

Gravel is the term used for describing coarse fragments between 2 mm and 7.5 cm diameter and stones for those between 7.5 cm and 25 cm. The presence of gravel and stones in soil reduces the volume of soil responsible for moisture and nutrient storage, drainage, infiltration and runoff, and hinders plant growth by impeding root growth and seedling emergence, intercultural operations and farm mechanization. The gravelliness classes used in LRI were used to classify the soils and using these classes a gravelliness map was generated. The area extent and their geographic distribution in the microwatershed is shown in Fig. 5.4.

The soils that are non-gravelly (<15% gravel) cover an area of about 4 ha (1%) and distributed in the northern part of the microwatershed. Maximum area of about 106 ha (34 %) is covered by gravelly (15-35% gravel) soils and are distributed in the major part of the microwatershed (Fig. 5.4).

The most productive lands with respect to gravelliness are found to be one per cent. They are non-gravelly with less than 15 per cent gravel and have potential for growing both annual and perennial crops.




Fig. 5.4 Soil Gravelliness map of Hitnal Microwatershed

### 5.5 Available Water Capacity

The soil available water capacity (AWC) is estimated based on the ability of the soil column to retain water between the tensions of 0.33 and 15 bar in a depth of 100 cm or the entire solum if the soil is shallower. The AWC of the soils (soil series) as estimated by considering the soil texture, mineralogy, soil depth and gravel content (Sehgal *et al.*, 1990) and accordingly the soil map units were grouped into five AWC classes *viz*, very low (<50 mm/m), low (50-100 mm/m), medium (100-150 mm/m), high (150-200 mm/m) and very high (>200 mm/m) and using these values, an AWC map was generated. The area extent and their geographic distribution of different AWC classes in the microwatershed is shown in Fig. 5.5.

An area of about 47 ha (15%) in the microwatershed has soils that are very low (<50 mm/m) in available water capacity and are distributed in the southern and central part of the microwatershed. An area of about 59 ha (19%) has soils that are low (51 to 100 mm/m) in available water capacity and are distributed in the central part of the microwatershed. An area of about 4 ha (1 %) is very high (200 mm/min) in available water capacity and distributed in the northern part of the microwatershed.

An area of about 106 ha (34%) in the microwatershed has soils that are problematic with regard to available water capacity. Here, only short duration crops can be grown and the probability of crop failure is very high. These areas are best put to other alternative uses. An area of about 4 ha (1%) has soils that have high potential (>200

mm/m) with regard to available water capacity where all climatically adapted long duration crops can be grown successfully.




Fig. 5.5 Soil Available Water Capacity map of Hitnal Microwatershed

## 5.6 Soil Slope

Soil slope refers to the inclination of the surface of the land. It is defined by gradient, shape and length, and is an integral feature of any soil as a natural body. Slope is considered important in soil genesis, land use and land development. The length and gradient of slope influences the rate of runoff, infiltration, erosion and deposition. The soil map units were grouped into two slope classes and a slope map was generated showing the area extent and their geographic distribution of different slope classes in the microwatershed (Fig. 5.6).

Nearly level (0-1%) lands cover an area of about 4 ha (1%) and distributed in the northern part of the microwatershed. Very gently sloping (1-3%) lands cover a maximum area of about 106 ha (34%) and distributed in the major part of the microwatershed. In all these areas, all climatically adapted annual and perennial crops can be grown without much soil and water conservation and other land development measures.




Fig. 5.6 Soil Slope map of Hitnal Microwatershed

## 5.7 Soil Erosion

Soil erosion refers to the wearing away of the earth's surface by the forces of water, wind and ice involving detachment and transport of soil by raindrop impact. It is used for accelerated soil erosion resulting from disturbance of the natural landscape by burning, excessive grazing and indiscriminate felling of forest trees and tillage, all usually by man. The erosion classes showing an estimate of the current erosion status as judged from field observations in the form of rills, gullies or a carpet of gravel on the surface are recorded. Four erosion classes, viz, slight erosion (e1), moderate erosion (e2), severe erosion (e3) and very severe erosion (e4) are recognized. The soil map units were grouped into different erosion classes and a soil erosion map generated. The area extent and their spatial distribution in the microwatershed is given in Figure 5.7.

Slightly eroded lands cover an area of about 4 ha (1 %) and distributed in the northern part of the microwatershed. Maximum area of about 106 ha (34 %) is moderately eroded (e2 class) and distributed in the major part of the microwatershed. Moderately eroded lands are problematic and need appropriate soil and water conservation and other land development measures.






Fig. 5.7 Soil Erosion map of Hitnal Microwatershed



## **FERTILITY STATUS**

Soil fertility plays an important role in increasing crop yield. The adoption of high yielding varieties that require high amounts of nutrients has resulted in deficiency symptoms in crops and plants due to imbalanced fertilization and poor inherent fertility status, as these areas are characterized by low rainfall and high temperatures. Hence, it is necessary to know the fertility (macro and micro nutrients) status of the soils of the watersheds for assessing the kind and amount of fertilizers required for each of the crop intended to be grown. For this purpose, the surface soil samples collected from the grid points (one soil sample at every 320 m grid interval) all over the microwatershed through land resource inventory in the year 2018 were analyzed for pH, EC, organic carbon, available phosphorus and potassium, and for micronutrients like zinc, boron, copper, iron and manganese, and secondary nutrient sulphur.

Soil fertility data generated has been assessed and individual maps for all the nutrients for the microwatershed have been generated by using the Kriging method under GIS. The village/survey number wise fertility data for the microwatershed is given in Appendix-II.

### **6.1 Soil Reaction (pH)**

The soil analysis of the Hitnal microwatershed for soil reaction (pH) showed that neutral soils (pH 6.5-7.3) cover an area of about 25 ha (8%) and distributed in the southern part of the microwatershed. Slightly to moderately alkaline soils (pH 7.3-8.4) cover an area of about 81 ha (26%) and distributed in the major part of the microwatershed. Strongly to very strongly alkaline (pH 8.4->9.0) soils cover an area of about 4 ha (1%) and distributed in the northern part of the microwatershed (Fig.6.1). An area of about 25 ha (8%) is neutral and 85 ha (27%) is alkaline in reaction.

### **6.2 Electrical Conductivity (EC)**

The Electrical Conductivity of the soils of the entire microwatershed area is  $<2 \text{ dSm}^{-1}$  (Fig 6.2) and as such the soils are non-saline.

### **6.3 Organic Carbon**

An area of about 34 ha (11%) is medium (0.5-0.75%) and distributed in the western and central part of the microwatershed. An area of about 77 ha (25%) is high ( $>0.75\%$ ) and distributed in the major part of the microwatershed (Fig.6.3).




Fig.6.1 Soil Reaction (pH) map of Hitnal Microwatershed




Fig.6.2 Electrical Conductivity (EC) map of Hitnal Microwatershed




Fig.6.3 Soil Organic Carbon map of Hitnal Microwatershed

#### 6.4 Available Phosphorus

An area of about 18 ha (6%) is low (<23 kg/ha) and distributed in the western part of the microwatershed. An area of about 57 ha (18%) is medium (23-57 kg/ha) in available phosphorus and distributed in the major part of the microwatershed. An area of about 35 ha (11%) is high (>57 kg/ha) and distributed in the northern part of the microwatershed. The areas with high phosphorus content reduce 25 per cent from the recommended dose to avoid the excess application of fertilizer. Apply additional 25% phosphorus in areas where it is low and medium (Fig 6.4).

#### 6.5 Available Potassium

Available potassium is low (<145 kg ha) in 22 ha (7%) and distributed in the southern part of the microwatershed. Maximum area of about 84 ha (27%) is medium (145-337 kg/ha) and distributed in the major part of the microwatershed. An area of about 4 ha (1%) is high (>337 kg/ha) and distributed in the northern part of the microwatershed. The areas with high potassium content reduce 25 per cent from the recommended dose to avoid the excess application of fertilizer. Apply additional 25% potassium in areas where it is low and medium (Fig 6.5).

#### 6.6 Available Sulphur

Soil analysis of available sulphur content in Hitnal microwatershed showed that an area of about 17 ha (5%) is low and distributed in the northern and southern part of the

microwatershed. An area of about 54 ha (17%) is medium (10-20 ppm) in available sulphur content and distributed in the major part of the microwatershed. An area of about 40 ha (13%) is high (>20 ppm) and distributed in the central part of the microwatershed (Fig.6.6). The areas that are low and medium in available sulphur need to be applied with magnesium sulphate or gypsum or factomphos (p) fertilizer (13% sulphur) for 2-3 years for the deficiency to be corrected.

### **6.7 Available Boron**

An area of about 92 ha (30%) is low (< 0.5ppm) in available boron and distributed in the major part of the microwatershed. An area of about 14 ha (4%) is medium (0.5-1.0 ppm) and distributed in the western part of the microwatershed. An area of about 4 ha (1%) is high (>1.0 ppm) and distributed in the northern part of the microwatershed (Fig.6.7).

### **6.8 Available Iron**

Available iron content in the soils of the Hitnal microwatershed is deficient (<4.5 ppm) in an area of about 52 ha (17%) and distributed in the central part of the microwatershed. An area of about 59 ha (19%) showed sufficiency (>4.5 ppm) with respect to iron content and distributed in the northern and southern part of the microwatershed (Fig 6.8).

### **6.9 Available Manganese**

Available manganese content is sufficient (>1.0 ppm) in the entire microwatershed area (Fig 6.9).

### **6.10 Available Copper**

Available copper content is sufficient (>0.2 ppm) in the entire microwatershed area (Fig 6.10).

### **6.11 Available Zinc**

Available zinc content is deficient (<0.6 ppm) in 21 ha (7 %) and distributed in the western and central part of the microwatershed. Maximum area of about 89 ha (29%) is sufficient (>0.6 ppm) and distributed in the major part of the microwatershed (Fig 6.11).




Fig.6.4 Soil Available Phosphorus map of Hitnal Microwatershed




Fig.6.5 Soil Available Potassium map of Hitnal Microwatershed




Fig.6.6 Soil Available Sulphur map of Hitnal Microwatershed




Fig.6.7 Soil Available Boron map of Hitnal Microwatershed






Fig.6.8 Soil Available Iron map of Hitnal Microwatershed




Fig.6.9 Soil Available Manganese map of Hitnal Microwatershed




Fig.6.10 Soil Available Copper map of Hitnal Microwatershed




Fig.6.11 Soil Available Zinc map of Hitnal Microwatershed

## LAND SUITABILITY FOR MAJOR CROPS

The soil and land resource units (soil phases) of Hitnal Microwatershed were assessed for their suitability for growing food, fodder, fibre and other horticulture crops by following the procedure as outlined in FAO, 1976 and 1983. Crop requirements were developed for each of the crop from the available research data and also by referring to Naidu *et. al.* (2006) and Natarajan *et. al.* (2015). The soil and land characteristics were matched with the crop requirements to arrive at the crop suitability. The soil and land characteristics table (Table 7.1) were matched with the crop requirements (Tables 7.2-7.32) to arrive at the crop suitability and the crop requirement tables are given at the end of the chapter. In FAO land suitability classification, two orders are recognized. Order S- Suitable and Order N- Not suitable. The orders have classes, subclasses and units. Order S has three classes, Class S1- Highly Suitable, Class S2- Moderately Suitable and Class S3- Marginally Suitable. Order N has two Classes, N1- Currently not Suitable and N2- Permanently not Suitable. There are no subclasses within the Class S1 as they will have very minor or no limitations for crop growth. Classes S2, S3 and N1 are divided into subclasses based on the kinds of limitations encountered. The limitations that affect crop production are 'c' for erratic rainfall and its distribution and length of growing period (LGP), 'e' for erosion hazard, 'r' for rooting condition, 't' for lighter or heavy texture, 'g' for gravelliness or stoniness, 'n' for nutrient availability, 'l' for topography, 'm' for moisture availability, 's' for sodium 'z' for calcareousness and 'w' for drainage. These limitations are indicated as lower case letters to the class symbol. For example, moderately suitable lands with the limitations of soil depth and erosion are designated as S2re. For the microwatershed, the soil mapping units were evaluated and classified up to subclass level.

Using the above criteria, the soil map units of the microwatershed were evaluated and land suitability maps for 31 major agricultural and horticultural crops were generated. The detailed information on the kind of suitability of each of the soil phase for the crops assessed are given village/ survey number wise for the microwatershed in Appendix-III.

### 7.1 Land Suitability for Sorghum (*Sorghum bicolor*)

Sorghum is one of the major food crop grown in Karnataka in an area of 10.47 lakh ha in Bijapur, Gulbarga, Raichur, Bidar, Belgaum, Dharwad, Bellary, Chitradurga, Mysore and Chamarajnagar districts. The crop requirements for growing sorghum (Table 7.2) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing sorghum was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.1.

Highly suitable (Class S1) lands occupy an area of about 4 (1%) for growing sorghum and occur in the northern part of the microwatershed. A maximum area of about

59 ha (19%) is moderately suitable (Class S2) for growing sorghum and distributed in the major part of the microwatershed with minor limitations of gravelliness and rooting depth. An area of about 47 ha (15%) is marginally suitable for growing sorghum and distributed in the southern and central part of the microwatershed. They have moderate limitation of gravelliness.




Fig. 7.1 Land Suitability map of Sorghum

## 7.2 Land Suitability for Maize (*Zea mays*)

Maize is one of the most important food crop grown in an area of 13.37 lakh ha in almost all the districts of the State. The crop requirements for growing maize (Table 7.3) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing maize was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.2.

Maximum area of about 63 ha (20%) is moderately suitable (Class S2) for growing maize and distributed in the major part of the microwatershed with minor limitations of gravelliness, rooting depth and texture. An area of about 47 ha (15%) is marginally suitable for growing maize and distributed in the central and southern part of the microwatershed. They have moderate limitation of gravelliness.




Fig. 7.2 Land Suitability map of Maize

### 7.3 Land Suitability for Bajra (*Pennisetum glaucum*)

Bajra is one of the major food crop grown in an area of 2.34 lakh ha in Karnataka in the northern districts. The crop requirements (Table 7.4) for growing bajra were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing bajra was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.3.

Entire cultivated area of 110 ha(25%) in the microwatershed is moderately suitable (Class S2) for growing bajra with minor limitations of texture, rooting depth and gravelliness.




Fig. 7.3 Land Suitability map of Bajra

#### 7.4 Land Suitability for Redgram (*Cajanus cajan*)

Redgram is one of the most important pulse crop grown in an area of 7.28 lakh ha in almost all the districts of the State. The crop requirements for growing redgram (Table 7.5) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing redgram was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.4.

Maximum area of about 63 ha (20%) is moderately suitable (Class S2) for growing redgram and distributed in the major part of the microwatershed. They have minor limitations of gravelliness, rooting depth and texture. Marginally suitable lands (Class S3) occupy an area of about 47 ha (15%) and occur in the southern and central part of the microwatershed. They have moderate limitations of rooting depth and gravelliness.




Fig. 7.4 Land Suitability map of Redgram

### 7.5 Land Suitability for Bengal gram (*Cicer arietinum*)

Bengal gram is one of the major pulse crop grown in an area of 9.39 lakh ha in northern Karnataka in Bijapur, Gulbarga, Raichur, Bidar, Belgaum, Dharwad and Bellary districts. The crop requirements for growing Bengal gram (Table 7.6) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing Bengal gram was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.5.

Highly suitable (Class S1) lands occupy an area of about 4 ha (1%) for growing bengal gram and occur in the northern part of the microwatershed. Marginally suitable (Class S3) lands cover an area of about 106 ha (34%) and are distributed in the major part of the microwatershed. They have moderate limitations of gravelliness and texture.




Fig. 7.5 Land Suitability map of Bengal gram

## 7.6 Land Suitability for Groundnut (*Arachis hypogaea*)

Groundnut is one of the major oilseed crop grown in an area of 6.54 lakh ha in Karnataka in most of the districts either as rainfed or irrigated crop. The crop requirements for growing groundnut (Table 7.7) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing groundnut was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.6.

An area of about 106 ha (34%) is moderately suitable (Class S2) for growing groundnut and distributed in the major part of the microwatershed. They have minor limitations of gravelliness, rooting depth and texture. An area of about 4 ha (1%) is marginally suitable (Class S3) for growing groundnut and are distributed in the northern part of the microwatershed with moderate limitation of texture.






Fig. 7.6 Land Suitability map of Groundnut

### 7.7 Land Suitability for Sunflower (*Helianthus annuus*)

Sunflower is one of the most important oilseed crop grown in an area of 3.56 lakh ha in the State in all the districts. The crop requirements for growing sunflower (Table 7.8) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing sunflower was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.7.

An area of about 4 ha (1%) is highly suitable (Class S1) for growing sunflower and are distributed in the northern part of the microwatershed. An area of about 59 ha (19%) is moderately suitable (Class S2) and are distributed in the major part of the microwatershed. They have minor limitations of gravelliness and rooting depth. Marginally suitable (Class S3) lands occupy an area of about 47 ha (15%) and are distributed in the central and southern part of the microwatershed with moderate limitations of rooting depth and gravelliness.




Fig. 7.7 Land Suitability map of Sunflower

### 7.8 Land Suitability for Cotton (*Gossypium hirsutum*)

Cotton is one of the most important fibre crop grown in the State in about 8.75 lakh ha area in Raichur, Dharwad, Belgaum, Gulbarga, Bijapur, Bidar, Bellary, Chitradurga and Chamarajnarag districts. The crop requirements for growing cotton (Table 7.9) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing cotton was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.8.

Highly suitable (Class S1) lands occupy an area of about 4 ha (1%) for growing cotton and occur in the northern part of the microwatershed. Maximum area of about 59 ha (19%) is moderately suitable (Class S2) and are distributed in the major part of the microwatershed. They have minor limitations of rooting depth and gravelliness. Marginally suitable (Class S3) lands occupy an area of about 48 ha (15%) and are distributed in the northern and central part of the microwatershed with moderate limitations of texture and gravelliness.




Fig. 7.8 Land Suitability map of Cotton

### 7.9 Land Suitability for Chilli (*Capsicum annum L*)

Chilli is one of the most important spice crop grown in an area of 0.42 lakh ha in Karnataka State. The crop requirements for growing chilli (Table 7.10) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing chilli was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.9.

An area of about 63 ha (20%) is moderately suitable (Class S2) and are distributed in the major part of the microwatershed. They have minor limitations of texture, rooting depth and gravelliness. Marginally suitable (Class S3) lands cover an area of about 47 ha (15%) and distributed in the central and southern part of the microwatershed. They have moderate limitation of gravelliness.




Fig. 7.9 Land Suitability map of Chilli

### 7.10 Land Suitability for Tomato (*Solanum lycopersicum*)

Tomato is one of the most important vegetable crop grown in an area of 0.65 lakh ha in almost all the districts of the State. The crop requirements (Table 7.11) for growing tomato were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing tomato was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.10.

An area of about 59 ha (19%) is moderately suitable (Class S2) and are distributed in the central and southern part of the microwatershed. They have minor limitations of rooting depth and gravelliness. Marginally suitable (Class S3) lands cover an area of about 51 ha (16%) and distributed in the northern, central and southern part of the microwatershed. They have moderate limitations of gravelliness and texture.




Fig. 7.10 Land Suitability map of Tomato

### 7.11 Land Suitability for Brinjal (*Solanum melongena*)

Brinjal is one of the most important vegetable crop grown in the state. The crop requirements for growing brinjal (Table 7.12) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing brinjal was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.11.

An area of about 59 ha (19%) is moderately suitable (Class S2) for growing Brinjal and distributed in the southern and central part of the microwatershed with minor limitations of rooting depth and gravelliness. Marginally suitable (Class S3) lands cover an area of about 51 ha (16%) and occur in the northern, central and eastern part of the microwatershed with moderate limitations of gravelliness and texture.




Fig 7.11 Land Suitability map of Brinjal

### 7.12 Land Suitability for Onion (*Allium cepa L.*)

Onion is one of the most important vegetable crop grown in the state. The crop requirements for growing onion (Table 7.13) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing onion was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.12.

An area of about 59 ha (19%) is moderately suitable (Class S2) for growing Onion and distributed in the southern and central part of the microwatershed with minor limitations of rooting depth and gravelliness. Marginally suitable (Class S3) lands cover an area of about 51 ha (16%) and occur in the northern, central and eastern part of the microwatershed with moderate limitations of gravelliness and texture.




Fig 7.12 Land Suitability map of Onion

### 7.13 Land Suitability for Bhendi (*Abelmoschus esculentus*)

Bhendi is one of the most important vegetable crop grown in the state. The crop requirements for growing bhendi (Table 7.14) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing bhendi was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.13.

Moderately suitable (Class S1) lands occupy an area of about 63 ha (20%) for growing Bhendi and occur in the major part of the microwatershed with minor limitations of gravelliness, texture and rooting depth. An area of about 47 ha (15%) is marginally suitable (Class S3) for growing Bhendi and distributed in the southern and central part of the microwatershed with moderate limitation of gravelliness.




Fig 7.13 Land Suitability map of Bhendi

#### 7.14 Land Suitability for Drumstick (*Moringa oleifera*)

Drumstick is one of the most important vegetable crop grown in 2403 ha area in the state. The crop requirements for growing drumstick (Table 7.15) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing drumstick was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.14.

Moderately suitable (Class S2) lands cover an area of about 63 ha (20%) and are distributed in the central and southern part of the microwatershed. They have minor limitations of rooting depth, texture and gravelliness. Marginally suitable (Class S3) lands cover an area of about 47 ha (15%) and occur in the southern and central part of the microwatershed. They have moderate limitations of gravelliness and rooting depth






Fig. 7.14 Land Suitability map of Drumstick

### 7.15 Land Suitability for Mulberry (*Morus nigra*)

Mulberry is the most important leaf crop grown for rearing silkworms in about 1.66 lakh ha in all the districts of the state. The crop requirements for growing mulberry (Table 7.16) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing mulberry was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.15.

An area of about 59 ha (19%) is moderately suitable (Class S2) for growing mulberry and distributed in the central part of the microwatershed. They have minor limitations of gravelliness and rooting depth. Marginally suitable (Class S3) lands cover an area of about 51 ha (16%) and occur in the southern, central and northern part of the microwatershed. They have moderate limitations of rooting depth, texture and gravelliness.




Fig. 7.15 Land Suitability map of Mulberry

### 7.16 Land Suitability for Mango (*Mangifera indica*)

Mango is one of the most important fruit crop grown in about 1.73 lakh ha in almost all the districts of the State. The crop requirements (Table 7.17) for growing mango were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing mango was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.16.

Marginally suitable (Class S3) lands cover an area of about 63 ha (20%) and occur in the major part of the microwatershed. They have moderate limitations of rooting depth and texture. Area currently not suitable (Class N1) for growing mango cover about 47 ha (15%) and distributed in the southern and central part of the microwatershed with severe limitation of rooting depth.




Fig. 7.16 Land Suitability map of Mango

### 7.17 Land Suitability for Sapota (*Manilkara zapota*)

Sapota is one of the most important fruit crop grown in an area of about 29373 ha in almost all the districts of the state. The crop requirements (Table 7.18) for growing sapota were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing sapota was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.17.

An area of about 59 ha (19%) is moderately suitable (Class S2) for growing sapota and distributed in the southern and central part of the microwatershed. They have minor limitations of rooting depth and gravelliness. Marginally suitable (Class S3) lands cover an area of about 51 ha (16%) and occur in the southern, central and northern part of the microwatershed. They have moderate limitations of rooting depth, gravelliness and texture.




Fig. 7.17 Land Suitability map of Sapota

### 7.18 Land Suitability for Pomegranate (*Punica granatum*)

Pomegranate is one of the commercially grown fruit crop in about 18488 ha in Karnataka mainly in Bijapur, Bagalkot, Koppal, Gadag and Chitradurga districts. The crop requirements for growing pomegranate (Table 7.19) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing pomegranate was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.18.

Moderately suitable (Class S2) lands occupy an area of about 63 ha (20%) and are distributed in the major part of the microwatershed. They have minor limitations of rooting depth, gravelliness and texture. Marginally suitable (Class S3) lands for growing pomegranate occupy an area of about 47 ha (15%) and are distributed in the southern and central part of the microwatershed with moderate limitations of rooting depth and gravelliness.




Fig. 7.18 Land Suitability map of Pomegranate

### 7.19 Land Suitability for Guava (*Psidium guajava*)

Guava is one of the most important fruit crop grown in an area of about 6558 ha in almost all the districts of the state. The crop requirements (Table 7.20) for growing guava were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing guava was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.19.

Moderately suitable (Class S2) lands occupy an area of about 59 ha (19%) and are distributed in the central and southern part of the microwatershed. They have minor limitations of rooting depth and gravelliness. Marginally suitable (Class S3) lands for growing guava occupy an area of about 51 ha (16%) and are distributed in the central northern and southern part of the microwatershed with moderate limitations of gravelliness, texture and rooting depth.




Fig. 7.19 Land Suitability map of Guava

## 7.20 Land Suitability for Jackfruit (*Artocarpus heterophyllus*)

Jackfruit is one of the most important fruit crop grown in 5368 ha in all the districts of the state. The crop requirements (Table.7.21) for growing jackfruit were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing jackfruit was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in figure 7.20.

Moderately suitable (Class S2) lands occupy an area of about 59 ha (19%) and are distributed in the central and southern part of the microwatershed. They have minor limitations of rooting depth and gravelliness. Marginally suitable (Class S3) lands for growing jackfruit occupy an area of about 51 ha (16%) and are distributed in the central northern and southern part of the microwatershed with moderate limitations of gravelliness, texture and rooting depth.




Fig. 7.20 Land Suitability map of Jackfruit

### 7.21 Land Suitability for Jamun (*Syzygium cumini*)

Jamun is an important fruit crop grown in almost all the districts of the state. The crop requirements (Table 7.22) for growing jamun were matched with the soil-site characteristics and a land suitability map for growing jamun was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.21.

Moderately suitable (Class S2) lands occupy an area of about 4 ha (1%) and distributed in the northern part of the microwatershed. They have minor limitation of texture. Marginally suitable (Class S3) lands cover an area of about 106 ha (34%) and are distributed in the major part of the microwatershed with moderate limitations of rooting depth and gravelliness.




Fig. 7.21 Land Suitability map of Jamun

## 7.22 Land Suitability for Musambi (*Citrus limetta*)

Musambi is one of the most important fruit crop grown in an area of 5446 ha in almost all the districts of the state. The crop requirements (Table 7.23) for growing musambi were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing musambi was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.22.

An area of about 4 ha (1%) is highly suitable (Class S1) for growing musambi and are distributed in the northern part of the microwatershed. An area of about 59 ha (19%) is moderately suitable (Class S2) and occur in the southern and central part of the microwatershed. They have minor limitations of gravelliness and rooting depth. An area of about 47 ha (15%) is marginally suitable (Class S3) for growing musambi and are distributed in the central and southern part of the microwatershed with moderate limitations of gravelliness and rooting depth.






Fig. 7.22 Land Suitability map of Musambi

### 7.23 Land Suitability for Lime (*Citrus sp*)

Lime is one of the most important fruit crop grown in an area of 11752 ha in almost all the districts of the State. The crop requirements for growing lime (Table 7.24) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing lime was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.23.

An area of about 4 ha (1%) is highly suitable (Class S1) for growing lime and are distributed in the northern part of the microwatershed. An area of about 59 ha (19%) is moderately suitable (Class S2) and occur in the southern and central part of the microwatershed. They have minor limitations of gravelliness and rooting depth. An area of about 47 ha (15%) is marginally suitable (Class S3) for growing lime and are distributed in the central and southern part of the microwatershed with moderate limitations of gravelliness and rooting depth.




Fig. 7.23 Land Suitability map of Lime

### 7.24 Land Suitability for Cashew (*Anacardium occidentale*)

Cashew is one of the most important nut crop grown in an area of 7052 ha in almost all the districts of the State. The crop requirements for growing cashew (Table 7.25) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing cashew was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.24.

An area of about 59 ha (19%) is moderately suitable (Class S2) and occur in the central part of the microwatershed. They have minor limitations of gravelliness and rooting depth. An area of about 47 ha (15%) is marginally suitable (Class S3) for growing cashew and are distributed in the southern and central part of the microwatershed with moderate limitations of gravelliness and rooting depth. An area of about 4 ha (1%) is currently not suitable (Class N1) for growing cashew and distributed in the northern part of the microwatershed with severe limitation of texture.




Fig. 7.24 Land Suitability map of Cashew

### 7.25 Land Suitability for Custard Apple (*Annona reticulata*)

Custard apple is one of the most important fruit crop grown in 1426 ha in almost all the districts of the State. The crop requirements( Table 7.26) for growing custard apple were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing custard apple was generated .The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.25.

An area of about 4 ha (1%) is highly suitable (Class S1) for growing custard apple and are distributed in the northern part of the microwatershed. Moderately suitable (Class S2) lands cover an area of about 106 ha (34%) and occur in the major part of the microwatershed. They have minor limitations of rooting depth and gravelliness.




Fig. 7.25 Land Suitability map of Custard Apple

### 7.26 Land Suitability for Amla (*Phyllanthus emblica*)

Amla is one of the most important fruit and medicinal crop grown in an area of 151 ha and distributed in almost all the districts of the state. The crop requirements (Table 7.27) for growing amla were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing amla was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.26.

Moderately suitable (Class S2) lands cover an entire area of the microwatershed. They have minor limitations of rooting depth, texture and gravelliness.




Fig. 7.26 Land Suitability map of Amla

### 7.27 Land Suitability for Tamarind (*Tamarindus indica*)

Tamarind is one of the most important spice crop grown in 14897 ha in all the districts of the state. The crop requirements (Table 7.28) for growing tamarind were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing tamarind was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.27.

An area of about 4 ha (1%) is moderately suitable (Class S2) and occur in the northern part of the microwatershed. They have minor limitation of texture. An area of about 59 ha (19%) is marginally suitable (Class S3) for growing tamarind and are distributed in the central part of the microwatershed with moderate limitation of rooting depth. An area of about 47 ha (15%) is currently not suitable (Class N1) for growing tamarind and distributed in the southern and central part of the microwatershed with severe limitation of rooting depth.




Fig. 7.27 Land Suitability map of Tamarind

### 7.28 Land Suitability for Marigold (*Tagetes erecta*)

Marigold is one of the most important flower crop grown in an area of 9108 ha in almost all the districts of the state. The crop requirements (Table 7.29) for growing marigold were matched with the soil-site characteristics and a land suitability map for growing marigold was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.28.

Maximum area of about 63 ha (20%) is moderately suitable (Class S2) and occur in the central and northern part of the microwatershed. They have minor limitations of gravelliness, rooting depth and texture. An area of about 47 ha (15%) is marginally suitable (Class S3) for growing marigold and are distributed in the southern and central part of the microwatershed with moderate limitation of gravelliness.




Fig. 7.28 Land Suitability map of Marigold

### 7.29 Land Suitability for Chrysanthemum (*Chrysanthemum indicum*)

Chrysanthemum is one of the most important flower crop grown in an area of 4978 ha in almost all the districts of the State. The crop requirements (Table 7.30) for growing chrysanthemum were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing chrysanthemum was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed is given in Figure 7.29.

Maximum area of about 63 ha (20%) is moderately suitable (Class S2) and occur in the central and northern part of the microwatershed. They have minor limitations of gravelliness, rooting depth and texture. An area of about 47 ha (15%) is marginally suitable (Class S3) for growing chrysanthemum and are distributed in the southern and central part of the microwatershed with moderate limitation of gravelliness.




Fig. 7.29 Land Suitability map of Chrysanthemum

### 7. 30 Land Suitability for Jasmine (*Jasminum sp.*)

Jasmine is one of the most important flower crop grown in an area of 803 ha in almost all the districts of the State. The crop requirements (Table 7.31) for growing jasmine were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing jasmine was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.30.

An area of about 59 ha (19%) is moderately suitable (Class S2) and occur in the central and southern part of the microwatershed. They have minor limitations of gravelliness and rooting depth. An area of about 51 ha (16%) is marginally suitable (Class S3) for growing jasmine and are distributed in the northern, central and southern part of the microwatershed with moderate limitations of gravelliness and texture.






Fig. 7.30 Land Suitability map of Jasmine

### 7. 31 Land Suitability for Crossandra (*Crossandra infundibuliformis*)

Crossandra is one of the most important flower crop grown in almost all the districts of the State. The crop requirements (Table 7.32) for growing crossandra were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing crossandra was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.31.

An area of about 59 ha (19%) is moderately suitable (Class S2) and occur in the central and southern part of the microwatershed. They have minor limitations of gravelliness and rooting depth. An area of about 51 ha (16%) is marginally suitable (Class S3) for growing crossandra and are distributed in the northern, central and southern part of the microwatershed with moderate limitations of gravelliness and texture.




Fig. 7.31 Land Suitability map of Crossandra

**Table 7.1 Soil-Site Characteristics of Hitnal Microwatershed**

Soil Map Units	Climate (P) (mm)	Growing period (Days)	Drainage Class	Soil depth (cm)	Soil texture		Gravelliness		AWC (mm/m)	Slope (%)	Erosion	pH	EC (dSm <sup>-1</sup> )	ESP	CEC [Cmol (p <sup>+</sup> )kg <sup>-1</sup> ]	BS (%)
					Surf-ace	Sub-surface	Sur-face	Sub-surface								
LKRcB2g1	662	<90	WD	50-75	sl	gsc	15-35	40-60	51-100	1-3	moderate	8.18	0.30	4.51	12.19	100
LKRhB2g1	662	<90	WD	50-75	scl	gsc	15-35	40-60	51-100	1-3	moderate	8.18	0.30	4.51	12.19	100
MKHhB2g1	662	<90	WD	50-75	scl	gsc	15-35	>35	51-100	1-3	moderate	7.38	0.09	1.49	14.84	93
HTliB2	662	<90	WD	50-75	sc	gsc	-	15-35	51-100	1-3	moderate	7.11	0.10	0.30	0.90	147
HDHcB2g1	662	<90	WD	75-100	sl	gsc-gc	15-35	>35	51-100	1-3	moderate	6.54	0.07	7.11	5.84	84.7
HDHiB2g1	662	<90	WD	75-100	sc	gsc-gc	15-35	>35	51-100	1-3	moderate	6.54	0.07	7.11	5.84	84.7
KDTmA1	662	<90	MWD	>150	c	sc-c	-	-	>200	0-1	slight	6.95	0.17	0.65	12.10	100

**Table 7.2 Land suitability criteria for Sorghum**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime 1	Mean temperature in growing season	°C	26–30	30–34; 24–26	34–40; 20–24	>40; <20
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristics					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V. poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, c (red), c (black)	scl, cl	ls, sl	-
	pH	1:2.5	5.5-7.8	5.0-5.5 7.8-9.0	>9.0	-
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	10-15
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

**Table 7.3 Land suitability criteria for Maize**

Land use requirement		Rating				
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	30-34	35-38 26-30	38-40 26-20	
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc	c (red), c (black)	ls, sl	-
	pH	1:2.5	5.5-7.8	5.0-5.5 7.8-9.0	>9.0	-
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	-
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

**Table 7.4 Land suitability criteria for Bajra**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-32	33-38 24-27	39-40 20-23	<20
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm	500-750	400-500	200-400	<200
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	Sl, scl, cl,sc,c (red)	C (black)	ls	-
	pH	1:2.5	6.0-7.8	5.0-5.5 7.8-9.0	5.5-6.0 >9.0	
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	15-35	35-60	>60	
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	1-3	3-5	5-10	>10

**Table 7.5 Land suitability criteria for Red gram**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	30-35(G) 20-25(AV) 15-18 (F&PS) 35-40(M)	25-30(G) 20-25 (AV) 12-15 (F&PS) 30-35(M)	20-25(G) 15-20(AV) 10-12 (F&PS) 25-30(M)	< 20 <15 <10 <25
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. Well drained	Poorly drained	Very Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, c (red)	c (black),sl, scl, cl	ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-9.0	5.0-5.5 >9.0	-
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-50	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<1.0	1.0-2.0	>2.0	
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.6 Land suitability criteria for Bengal gram**

Land use requirement			Rating			
Soil –site characteristics	Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)	
Climatic regime	Mean temperature in growing season	°C	20–25	25–30; 15–20	30–35; 10–15	>35; <10
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. Well drained	Poorly drained	Very Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	C (black)	-	c (red), scl, cl, sc	ls, sl
	pH	1:2.5	6.0-7.8	5.0-6.0 7.8-9.0	>9.0	-
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	-
Erosion hazard	Slope	%	<3	3-5	5-10	>10



**Table 7.7 Land suitability criteria for Groundnut**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	24–33	22–24; 33–35	20–22; 35–40	<20; >40
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. Well drained	Poorly drained	Very Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl	sl,cl, sc	c (red), c (black), ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<35	35-60	>60	
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.8 Land suitability criteria for Sunflower**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	24–30	30–34; 20–24	34–38; 16–20	>38; <16
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	mod. Well drained	-	Poorly to very drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	cl, sc,c (red), c (black)	scl	ls, sl	-
	pH	1:2.5	6.5-7.8	7.8-8.4 5.5-6.5	8.4-9.0; 5.0-5.5	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.9 Land suitability criteria for Cotton**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	22-32	>32	<19	-
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well to moderately well	Poorly drained/Some what excessively drained	-	very poorly/excessively drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, c (red,black)	cl	scl	ls, sl
	pH	1:2.5	6.5-7.8	7.8-8.4	5.5-6.5 8.4->9.0	<5.5
	CEC	C mol (p+)Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	50-100	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	<3	3-5	-	>5

**Table 7.10 Land suitability criteria for Chili**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	25-32	33-35 20-25	35-38 <20	>38
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc	c (black), sl	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.11 Land suitability criteria for Tomato**

Land use requirement		Rating				
Soil –site characteristics	Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)	
Climatic regime	Mean temperature in growing season	°C	25-28	29-32 20-24	15-19 33-36	<15 >36
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl, scl, cl, sc, c (red)	-	ls, c(black)	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.12 Land suitability criteria for Brinjal**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	Well drained	Moderately well drained	Poorly drained	V. Poorly drained
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class				
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl, scl, cl, sc c (red)	-	ls, c (black)	-
	pH	1:2.5	6.0-7.3	7.3-8.4 5.0-6.0	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	>60
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.13 Land suitability criteria for Onion**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	20-30	30-35	35-40	>40
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately /imperfectly	-	Poorly to V poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl,scl,cl,sc,c (red)	-	c (Black),ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<1.0	1.0-2.0	2.0-4.0	<4
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.14 Land suitability criteria for Bhendi**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	25-28	29-32 20-24	15-19 33-36	<15 >36
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Imperfectly drained	Poorly to very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl,sc, c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10



**Table 7.15 Land suitability criteria for Drumstick**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, scl, cl, c (red)	sl, c (black)	ls	s
	pH	1:2.5	6.0-7.3	5.0-5.5 7.3-7.8	5.5-6.0 7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<35	35-60	60-80	>80
Soil toxicity	Salinity (EC saturation extract)	dS/m				
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-10	-	>10

**Table 7.16 Land suitability criteria for Mulberry**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	24–28	22–24; 28–32	32–38; 22–18	>38; <18
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V. Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, cl, scl	c (red)	c (black), sl, ls	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.8-8.4	7.3-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	0-35	35-60	60-80	>80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

**Note:** Suitability evaluation only for Mulberry leaf not for Silk worm rearing

**Table 7.17 Land suitability criteria for Mango**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-32	24-27 33-35	36-40	20-24
	Min temp. before flowering	°C	10-15	15-22	>22	-
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration	Days				
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	-	ls, sl, c (black)	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>150	100-150	75-100	<75
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.18 Land suitability criteria for Sapota**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-32	33-36 24-27	37-42 20-23	>42 <18
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	-	Poorly to very drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	sl	ls, c (black)	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.19 Land suitability criteria for Pomegranate**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	30-34	35-38 25-29	39-40 15-24	
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl,cl, sc, c (red)	c (black),sl	ls	-
	pH	1:2.5	5.5-7.8	7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.20 Land suitability criteria for Guava**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-32	33-36 24-27	37-42 20-23	
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	sl	c (black), ls	-
	pH	1:2.5	6.0-7.8	5.0-6.0	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.21 Land suitability criteria for Jackfruit**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. well	Poorly	V. Poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	-	sl, ls, c (black)	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	>60
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10-

**Table 7.22 Land suitability criteria for Jamun**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well	Mod. well	Poorly	V.Poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c(red)	sl, c (black)	ls	-
	pH	1:2.5	6.0-7.8	5.0-6.0	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>150	100-150	50-100	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	>60
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10



**Table 7.23 Land suitability criteria for Musambi**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-30	31-35 24-27	36-40 20-23	>40 <20
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately drained	poorly	Very poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c	sl	ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.24 Land suitability criteria for Lime**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-30	31-35 24-27	36-40 20-23	>40 <20
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately drained	poorly	Very poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c	sl	ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.25 Land suitability criteria for Cashew**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	32 to 34	28 to 32; 34 to 38	24 to 28; 38 to 40	<20; >40
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	-	sl, ls	c (black)
	pH	1:2.5	5.5-6.5	5.0-5.5 6.5-7.3	7.3-7.8	>7.8
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-10	>10	-

**Table 7.26 Land suitability criteria for Custard apple**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	Scl, cl, sc, c (red), c (black)	-	Sl, ls	-
	pH	1:2.5	6.0-7.3	5.5-6.0 7.3-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15-35	35-60	60-80	-
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	>5	-

**Table 7.27 Land suitability criteria for Amla**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod.well drained	Poorly drained	V. Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	c (black)	ls, sl	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15-35	35-60	60-80	-
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

**Table 7.28 Land suitability criteria for Tamarind**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod.well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl,sc, c (red)	sl, c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>150	100-150	75-100	<75
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

**Table 7.29 Land suitability criteria for Marigold**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	18-23	17-15 24-35	35-40 10-14	>40 <10
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl,scl, cl, sc, c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%				
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.30 Land suitability criteria for Chrysanthemum**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	18-23	17-15 24-35	35-40 10-14	>40 <10
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl,scl, cl, sc, c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%				
Erosion hazard	Slope	%	<3	3-5	5-10	>10



**Table 7.31 Land suitability criteria for Jasmine (irrigated)**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	18-23	17-15 24-35	35-40 10-14	-
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	sl	ls, c (black)	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%				
Erosion hazard	Slope	%	<3	3-5	5-10	>10

### 7.32 Land suitability criteria for Crossandra

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	-	Poorly to very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c(red)	sl,	c (black),ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%				
Erosion hazard	Slope	%	<3	3-5	5-10	>10

### 7.32 Land Management Units (LMUs)

The 7 soil map units identified in Hitnal Microwatershed have been grouped into four Land Management Units (LMUs) for the purpose of preparing a Proposed Crop Plan. Land Management Units are grouped based on the similarities in respect of the type of soil, the depth of the soil, the surface soil texture, gravel content, AWC, slope, erosion etc. and a Land Management Units map (Fig.7.32) has been generated. These Land Management Units are expected to behave similarly for a given level of management.

The map units that have been grouped into four Land Management Units along with brief description of soil and site characteristics are given below.

LMU	Mapping unit	Soil and site characteristics
1	KDTmA1	Very deep, black clay soils with slopes of 0-1%, slight erosion
2	HDHcB2g1, HDHiB2g1	Moderately deep, gravelly red sandy clay to clay soils with slopes of 1-3%, moderate erosion, gravelly (15-35%)
3	LKRcB2g1, LKRhB2g1, MKHhB2g1	Moderately shallow, red gravelly loamy soils with slopes of 1-3%, moderate erosion, gravelly (15-35%)
4	HTTiB2	Moderately shallow, red gravelly sandy clay soils with slopes of 1-3%, moderate erosion




Fig 7.32 Land Management Units map of Hitnal microwatershed

### **7.33 Proposed Crop Plan for Hitnal Microwatershed**

After assessing the land suitability for the 31 crops, the proposed crop plan has been prepared for the four identified LMUs by considering only the highly (Class S1) and moderately (Class S2) suitable lands for each of the 31 crops. The resultant proposed crop plan is presented in Table 7.33.

**Table 7.33 Proposed Crop Plan for Hitnal Microwatershed**

LMU	Soil Map Units	Survey Number	Field Crops/ Commercial crops	Horticulture Crops (Rainfed/Irrigated )	Suitable Interventions
1	403.KDTmA1 (Very deep, black clay soils)	<b>Hitnala</b> : 238	Maize, Sorghum, Sunflower, Cotton, Bengal gram, Safflower, Linseed, Bajra , Soybean	<b>Fruit crops:</b> Sapota, Pomegranate, Jamun, Lime, Musambi, Tamarind, Amla, Custard apple <b>Vegetables:</b> Drumstick, Chilli, Coriander, Tomato, Bhendi <b>Flowers:</b> Marigold, Chrysanthemum, Crossandra, Jasmine	Application of FYM, Bio-fertilizers and micronutrients, drip irrigation, mulching, suitable soil and water conservation practices
2	111.HDHcB2g1 128.HDHhB2g1 (moderately deep, gravelly red sandy clay to clay soils)	<b>Bevinahalli:</b> 199,200,46 <b>Chikkakasanakandi:</b> 1, 2,3,37,39,46 <b>Hitnala:</b> 165,195,196,1 97,198,199,202,203,205	Maize, Sorghum, Sunflower, Groundnut, Bajra, Cotton, Red gram	<b>Fruit crops</b> : Sapota, Pomegranate, Amla, Cashew, Guava, Custard apple, Jack fruit, Jamun, Lime, Musambi <b>Vegetables:</b> Tomato, Chilli, Drumstick, Onion, Bhendi, Brinjal, Curry leaves <b>Flowers:</b> Marigold, Chrysanthemum, Jasmine, Crossandra	Drip irrigation, mulching, suitable soil and water conservation practices (Crescent Bunding with Catch Pit <i>etc</i> )
3	43.LKRcB2g1 452.LKRhB2g1 85.MKHhB2g1 (Moderately shallow, red gravelly loamy soils)	<b>Chikkakasanakandi:</b> 3 3,36,38 <b>Hitnala:</b> 193,200,201,2 04,206,209,210,213,214 ,215,224,225,226,228,2 29,230	Bajra, Groundnut, Horse gram, Castor	<b>Fruit crops</b> : Amla, Custard apple <b>Vegetables:</b> Curry leaves <b>Flowers:</b> Marigold, Chrysanthemum	Drip irrigation, mulching, suitable soil and water conservation practices (Crescent Bunding with Catch Pit <i>etc</i> )
4	100.HTLiB2 (Moderately shallow, red gravelly sandy clay soils)	<b>Shapura</b> :43,44	Maize, Sorghum, Groundnut, Bajra, Cotton, Horse gram, Castor	<b>Fruit crops</b> : Amla, Custard apple <b>Vegetables:</b> Tomato, Chilli, Onion, Bhendi, Brinjal ,Curry leaves <b>Flowers:</b> Marigold, Chrysanthemum, Jasmine, Crossandra	Drip irrigation, mulching, suitable soil and water conservation practices (Crescent Bunding with Catch Pit <i>etc</i> )



## SOIL HEALTH MANAGEMENT

### 8.1 Soil Health

Soil health is basic to plant health and plant health is basic to human and bovine health. Soil is fundamental to crop production. Without soil, no food could be produced nor would livestock be fed on a large scale. Because it is finite and fragile, soil is a precious resource that requires special care from its users.

Soil health or the capacity of the soil to function is critical to human survival. Soil health has been defined as: “the capacity of the soil to function as a living system without adverse effect on the ecosystem”. Healthy soils maintain a diverse community of soil organisms that help to form beneficial symbiotic associations with plant roots, recycle essential plant nutrients, improve soil structure with positive repercussions for soil, water and nutrient holding capacity and ultimately improve crop production and also contribute to mitigating climate change by maintaining or increasing its carbon content.

Functional interactions of soil biota with organic and inorganic components, air and water determine a soil’s potential to store and release nutrients, and water to plants and to promote and sustain plant growth. Thus, maintaining soil health is vital to crop production and conserve soil resource base for sustaining agriculture.

#### **The most important characteristics of a healthy soil are**

- Good soil tilth
- Sufficient soil depth
- Good water storage and good drainage
- Adequate supply, but not excess of nutrients
- Large population of beneficial organisms
- Small proportion of plant pathogens and insect pests
- Low weed pressure
- Free of chemicals and toxins that may harm the crop
- Resistance to degradation
- Resilience when unfavourable conditions occur

#### **Characteristics of Hitnal Microwatershed**

- ❖ The soil phases with sizeable area identified in the microwatershed belonged to the soil series of HDH (59 ha), LKR (32 ha), MKH (16 ha), KDT (4 ha) and HTI (<1 ha).
- ❖ As per land capability classification, entire area in the microwatershed falls under arable land category (Class II and III). The major limitations identified in the arable lands were soil and erosion.
- ❖ On the basis of soil reaction, an area of about 25 ha (8%) is neutral (pH 6.5-7.3), 16 ha (5%) is slightly alkaline (pH 7.3-7.8), 65 ha (21%) is moderately alkaline (pH 7.8-

8.4), 1 ha (<1%) is strongly alkaline (pH 8.4-9.0) and 3 ha (<1%) is very strongly alkaline (pH >9.0) in reaction.

### **Soil Health Management**

The following actions are required to improve the current land husbandry practices that provide a sound basis for the successful adoption of sustainable crop production system.

#### **Alkaline soils**

An area of about 85 ha (28%) is under alkaline soils. The following actions are recommended.

1. Regular addition of organic manure, green manuring, green leaf manuring, crop residue incorporation and mulching needs to be taken up to improve the soil organic matter status.
2. Application of biofertilizers (Azospirillum, Azotobacter, Rhizobium).
3. Application of 25% extra N and P (125 % RDN&P).
4. Application of ZnSO<sub>4</sub> – 12.5 kg/ha (once in three years).
5. Application of Boron – 5 kg/ha (once in three years).

#### **Neutral soils**

Neutral soils cover about 25 ha (8%) and the following actions are recommended.

1. Regular addition of organic manure, green manuring, green leaf manuring, crop residue incorporation and mulching needs to be taken up to improve the soil organic matter status.
2. Application of biofertilizers, (Azospirillum, Azotobacter, Rhizobium).
3. Application of 100 per cent RDF.
4. Need based micronutrient applications.

#### **Soil Degradation**

Soil erosion is one of the major factor affecting the soil health in the microwatershed. An area of about 106 ha (34%) is under moderate erosion. The areas with moderate erosion need immediate soil and water conservation and, other land development and land husbandry practices for restoring soil health.

#### **Dissemination of Information and Communication of Benefits**

Any large scale implementation of soil health management requires that supporting information is made available widely, particularly through channels familiar to farmers and extension workers. Given the very high priority attached to soil health especially by the Central Government on issuing Soil-Health Cards to all the farmers, media outlets like Regional, State and National Newspapers, Radio and Dooradarshan programs in local languages but also modern information and communication technologies such as Cellular phones and the Internet, which can be much more effective in reaching the younger farmers.



## **Inputs for Net Planning (Saturation Plan) and Interventions needed**

Net planning in IWMP is focusing on preparation of

1. Soil and Water Conservation Treatment Plans for each plot or farm.
2. Productivity enhancement measures/ interventions for existing crops/livestock/other farm enterprises.
3. Diversification of farming mainly with perennial horticultural crops and livestock.
4. Improving livelihood opportunities and income generating activities.

In this connection, how various outputs of Sujala-III are of use in addressing these objectives of Net Planning are briefly presented below.

- ❖ **Soil Depth:** The depth of a soil decides the amount of moisture and nutrients it can hold, what crops can be taken up or not, depending on the rooting depth and the length of growing period available for raising any crop. Deeper the soil, better for a wide variety of crops. If sufficient depth is not available for growing deep rooted crops, either choose medium or short duration crops or deeper planting pits need to be opened and additional good quality soil brought from outside has to be filled into the planting pits.
- ❖ **Surface Soil Texture:** Lighter soil texture in the top soil means, better rain water infiltration, less run-off and soil moisture conservation, less capillary rise and less evaporation losses. Lighter surface textured soils are amenable to good soil tilth and are highly suitable for crops like groundnut, root vegetables (carrot, radish, potato etc) but not ideal for crops that need stagnant water like lowland paddy. Heavy textured soils are poor in water infiltration and percolation. They are prone for sheet erosion; such soils can be improved by sand mulching. The technology that is developed by the AICRP-Dryland Agriculture, Vijayapura, Karnataka can be adopted.
- ❖ **Gravelliness:** More gravel content is favorable for run-off harvesting but poor in soil moisture storage and nutrient availability. It is a significant parameter that decides the kind of crop to be raised.
- ❖ **Land Capability Classification:** The land capability map shows the areas suitable and not suitable for agriculture and the major constraints in each of the plot/survey number. Hence, one can decide what kind of enterprise is possible in each of these units. In general, erosion and soil are the major constraints in Hitnal Microwatershed.
- ❖ **Organic Carbon:** An area of about 34 ha (11%) is medium (0.5-0.75%) and 77 ha (25%) is high (>0.75%). The areas that are low and medium in OC needs to be further improved by applying farmyard manure and rotating crops with cereals and legumes or mixed cropping.
- ❖ **Promoting green manuring:** Growing of green manuring crops costs Rs. 1250/ha (green manuring seeds) and about Rs. 2000/ha towards cultivation that totals to Rs. 3250/- per ha. On the other hand, application of organic manure @ 10 tons/ha costs Rs. 5000/ha. The practice needs to be continued for 2-3 years or more. Nitrogen

fertilizer needs to be supplemented by 25% in addition to the recommended level in 34 ha area where OC is less than 0.75 per cent. For example, for rainfed maize, recommended level is 50 kg N per ha and an additional 12 kg /ha needs to be applied for all the crops grown in these plots.

- ❖ **Available Phosphorus:** Available phosphorus is low (<23 kg/ha) in 18 ha (6%), medium (23-57 kg/ha) in 57 ha (18%) and high (>57 kg/ha) in 35 ha (11%) area of the microwatershed. The areas with high phosphorus content reduce 25% from the RDF to avoid the excess application of fertilizer and apply additional 25% phosphorus in areas where it is low and medium.
- ❖ **Available Potassium:** Available potassium is low (<145 kg/ha) in 22 ha (7%), medium (145-337 kg/ha) in 84 ha (27%) and high (>337 kg/ha) in 4 ha (1%) area of the microwatershed. The areas with high potassium content reduce 25% from the RDF to avoid the excess application of fertilizer and apply additional 25% potassium in areas where it is medium.
- ❖ **Available Sulphur:** Available sulphur is a very critical nutrient for oilseed crops. Available sulphur is low (<10 ppm) in 17 ha (5%), medium in 54 ha (17%) and high (>20 ppm) in 40 ha (13%) area of the microwatershed. Areas with low and medium in available sulphur need to be applied with magnesium sulphate or gypsum or Factamphos (p) fertilizer (13% sulphur) for 2-3 years for the deficiency to be corrected.
- ❖ **Available Iron:** It is deficient (<4.5 ppm) in 52 ha (17 %) and sufficient (>4.5 ppm) in 59 ha (19 %) area of the microwatershed. To manage iron deficiency iron sulphate @ 25 kg/ha needs to be applied for 2-3 years.
- ❖ **Available Zinc:** It is deficient (<0.6 ppm) in 21 ha (7%) and sufficient (>0.6 ppm) in 89 ha (29%) area of the microwatershed. Application of zinc sulphate @ 25kg/ha is to be followed in areas that are deficient in available zinc.
- ❖ **Available Boron:** Available boron is low in (<0.5ppm) 92 ha (30%), medium (0.5-1.0 ppm) in 14 ha (4%) and high (>1.0 ppm) in 4 ha (1%) area of the microwatershed. The areas with low and medium in boron content need to be applied with sodium borate @ 10kg/ha as soil application or 0.2% borax as foliar spray to correct the deficiency.
- ❖ **Available Manganese:** It is sufficient in the entire area of the microwatershed.
- ❖ **Available Copper:** It is sufficient in the entire area of the microwatershed.
- ❖ **Soil Alkalinity:** An area of about 85 ha (28%) in the microwatershed has soils that are slightly to very strongly alkaline. These areas need application of gypsum and wherever calcium is in excess, iron pyrites and element sulphur can be recommended. Management practices like treating repeatedly with good quality water to drain out the excess salts and provision of subsurface drainage and growing of salt tolerant crops like Casuarina, Acacia, Neem, Ber etc, are recommended.


- ❖ **Land Suitability for various crops:** Areas that are highly, moderately and marginally suitable and not suitable for growing various crops are indicated. Along with the suitability, various constraints that are limiting the productivity are also indicated. For example, in case of cotton, gravel content, rooting depth and salinity/alkalinity are the major constraints in various plots. With suitable management interventions, the productivity can be enhanced. In order to increase water holding capacity of light textured soils, growing of green manure crops and application of organic manure is recommended.



## SOIL AND WATER CONSERVATION TREATMENT PLAN

For preparing soil and water conservation treatment plan for Hitnal Microwatershed, the land resource inventory database generated under Sujala-III project has been transformed as information through series of interpretative (thematic) maps using soil phase map as a base. The various thematic maps (1:7920 scale) generated were

- Soil depth
- Surface soil texture
- Available water capacity
- Soil slope
- Soil gravelliness
- Land capability
- Present land use and land cover
- Crop suitability maps
- Rainfall map
- Hydrology
- Water Resources
- Socio-economic data
- Contour plan with existing features- network of waterways, pottissa boundaries, cut up/ minor terraces etc.
- Cadastral map (1:7920 scale)
- Satellite imagery (1:7920 scale)



Apart from these, Hand Level/ Hydro Marker/ Dumpy Level/ Total Station and Kathedars' List to be collected.

### Steps for Survey and Preparation of Treatment Plan

The boundaries of Land User Groups' and Survey No. boundaries are traced in the field.

- Naming of user groups and farmers
- Identification of arable and non arable lands
- Identification of drainage lines and gullies
- Identification of non treatable areas
- Identification of priority areas in the arable lands
- Treatment plan for arable lands
- Location of water harvesting and recharge structures

### 9.1 Treatment Plan

The treatment plan recommended for arable lands is briefly described below.



### 9.1.1 Arable Land Treatment

#### A. BUNDING

Steps for Survey and Preparation of Treatment Plan		USER GROUP-1
Cadastral map (1:7920 scale) is enlarged to a scale of 1:2500 scale		<p><b>CLASSIFICATION OF GULLIES</b></p> <p>ಕೊರಕಾಲಿನ ವರ್ಗೀಕರಣ</p> <ul style="list-style-type: none"> <li>ಮೇಲ್ಭಾಗ (UPPER REACH) 15 Ha.</li> <li>ಮಧ್ಯಭಾಗ (MIDDLE REACH) 15+10=25 ಹೆ.</li> <li>ಕೆಳಭಾಗ (LOWER REACH) 25 ಹೆಚ್ಚರಗಿನ ಅಧಿಕ</li> </ul> <p>POINT OF CONCENTRATION</p>
Existing network of waterways, pothissa boundaries, grass belts, natural drainage lines/ watercourse, cut ups/ terraces are marked on the cadastral map to the scale		
Drainage lines are demarcated into		
Small gullies	(up to 5 ha catchment)	
Medium gullies	(5-15 ha catchment)	
Ravines	(15-25 ha catchment) and	
Halla/Nala	(more than 25ha catchment)	

#### Measurement of Land Slope

Land slope is estimated or determined by the study and interpretation of contours or by measurement in the field using simple instruments like Hand Level or Hydromarker.



Vertical and Horizontal intervals between bunds as recommended by the Watershed Development Department.

Slope percentage	Vertical interval (m)	Corresponding Horizontal Distance (m)
2 - 3%	0.6	24
3 - 4%	0.9	21
4 - 5%	0.9	21
5 - 6%	1.2	21
6 - 7%	1.2	21

**Note:** i) The above intervals are maximum.

(ii) Considering the slope class and erosion status (A1... A= 0-1% slope, 1= slight erosion) the intervals have to be decided.

**Bund length recording:** Considering the contour plan and the existing grass belts/partitions, the bunds are aligned and lengths are measured.

**Section of the Bund**

Bund section is decided considering the soil texture class and gravelliness class (bg<sub>0</sub> .....b = loamy sand, g<sub>0</sub> = <15% gravel). The recommended sections for different soils are given below.

**Recommended Bund Section**



Top width (m)	Base width (m)	Height (m)	Side slope (Z:1;H:V)	Cross section (sq m)	Soil Texture	Remarks
0.3	0.9	0.3	01:01	0.18	Sandy loam	Vegetative bund
0.3	1.2	0.3	1.5:1	0.225	Sandy clay	
0.3	1.2	0.5	0.9:1	0.375	Red gravelly soils	
0.3	1.2	0.6	0.75:1	0.45		
0.3	1.5	0.6	01:01	0.54	Red sandy loam	
0.3	2.1	0.6	1.5:1	0.72	Very shallow clayey black soils	
0.45	2	0.75	01:01	0.92		
0.45	2.4	0.75	1.3:1	1.07	Shallow clayey black soils	
0.6	3.1	0.7	1.78:1	1.29	Medium clayey black soils	
0.5	3	0.85	1.47:1	1.49		

**Formation of Trench cum Bund**

Dimensions of the Borrow Pits/ Trenches to be excavated (machinery are decided considering the Bund Section).


Details of Borrow Pit dimensions are given below

**TRENCH CUM BUND**

IDEAL FOR HORTICULTURE CROPS

**'A' FRAME FOR INTERBUND MANAGEMENT**



1. ಸಮಾನಾಕಳ ಉಳುವುದು
2. ಸಮಾನಾಕಳ ಬಿತ್ತನೆ/ನಾಟಿ

### Size of Borrow Pits/ Trench recommended for Trench cum Bund (by machinery)

Bund section	Bund length	Earth quantity	Pit				Berm (pit to pit)	Soil depth Class
			L(m)	W(m)	D(m)	Quantity (m <sup>3</sup> )		
0.375	6	2.25	5.85	0.85	0.45	2.24	0.15	Shallow
0.45	6	2.7	5.4	1.2	0.43	2.79	0.6	Shallow
0.45	6	2.7	5	0.85	0.65	2.76	1	Moderately Shallow
0.54	5.6	3.02	5.5	0.85	0.7	3.27	0.1	Moderately shallow
0.54	5.5	2.97	5	1.2	0.5	3	0.5	Shallow
0.72	6.2	4.46	6	1.2	0.7	5.04	0.2	Moderately shallow
0.72	5.2	3.74	5.1	0.85	0.9	3.9	0.1	Moderately deep

### B. Waterways

- a) Existing waterways are marked on the cadastral map (1:7920 scale) and their dimensions are recorded.
- b) Considering the contour plan of the MWS, additional waterways/ modernization of the existing ones can be thought of.
- c) The design details are given in the Manual.

### C. Farm Ponds

Waterways and the catchment area will give an indication on the size of the Farm Pond. Location of the pond can be decided based on the contour plan/ field condition and farmers' need/desire.

### D. Diversion Channel

Existing EPT/ CPT are marked on the cadastral map. Looking to the need, these can be modernized or fresh diversion channel can be proposed and runoff from this can be stored in *Gokatte/ Recharge Ponds*.

### 9.1.2 Non-Arable Land Treatment

Depending on the gravelliness and crops preferred by the farmers, the concerned authorities can decide appropriate treatment plan. The recommended treatments may be Contour Trench, Staggered Trench, Crescent Bund, Boulder Bund or Pebble Bund.



### **9.1.3 Treatment of Natural Water Course/ Drainage Lines**

- a) The cadastral map has to be updated as regards the network of drainage lines (gullies/ *nalas/ hallas*) and existing structures are marked to the scale and storage capacity of the existing water bodies are documented.
- b) The drainage line will be demarcated into Upper Reach, Middle Reach and Lower Reach.
- c) Considering the Catchment, *Nala* bed and bank conditions, suitable structures are decided.
- d) Number of storage structures (Check dam/ *Nala* bund/ Percolation tank) will be decided considering the commitments and available runoff in water budgeting and quality of water in the wells and site suitability.
- e) Detailed Levelling Survey using Dumpy Level / Total Station has to be carried out to arrive at the site-specific designs as shown in the Manual.
- f) The location of ground water recharge structures are decided by examining the lineaments and fracture zones from geological maps.
- g) Rainfall intensity data of the nearest Rain Gauge Station is considered for Hydrologic Designs.
- h) Silt load to the Storage/Recharge Structures is reduced by providing vegetative, boulder and earthen checks in the natural water course. Location and design details are given in the Manual.

### **9.2 Recommended Soil and Water Conservation Measures**

The appropriate conservation structures best suited for each of the land parcel/ survey number (Appendix-I) are selected based on the slope per cent, severity of erosion, amount of rainfall, land use and soil type. The different kinds of conservation structures recommended are

1. Graded / Strengthening of Bunds
2. Trench cum Bunds (TCB)
3. Trench cum Bunds / Strengthening
4. Crescent Bunds

A map (Fig. 9.1) showing soil and water conservation plan with different kinds of structures recommended has been prepared which shows the spatial distribution and extent of area. A maximum area of about 106 ha (34%) needs trench cum bunding and an area of about 4 ha (1%) requires strengthening of existing bunds/ bunding. The conservation plan prepared may be presented to all the stakeholders including farmers and after considering their suggestions, the conservation plan for the microwatershed may be finalized in a participatory approach.




Fig. 9.1 Soil and Water Conservation Plan map of Hitnal Microwatershed

### 9.3 Greening of Microwatershed

As part of the greening programme in the watersheds, it is envisaged to plant a variety of horticultural and other tree plants that are edible, economical and produce lot of biomass which helps to restore the ecological balance in the watersheds. The lands that are suitable for greening programme are non-arable lands (land capability classes V, VI VII and VIII) and also the lands that are not suitable or marginally suitable for growing annual and perennial crops. The method of planting these trees is given below.

It is recommended to open the pits during the 1<sup>st</sup> week of March along the contour and heap the dugout soil on the lower side of the slope in order to harness the flowing water and facilitate weathering of soil in the pit. Exposure of soil in the pit also prevents spread of pests and diseases due to scorching sun rays. The pits should be filled with mixture of soil and organic manure during the second week of April and keep ready with sufficiently tall seedlings produced either in poly bags or in root trainer nurseries so that planting can be done during the 2<sup>nd</sup> or 3<sup>rd</sup> week of April depending on the rainfall.

The tree species suitable for the area considering rainfall, temperature and adaptability is listed below; waterlogged areas are recommended to be planted with species like Neral (*Syzgium cumini*) and Bamboo. Dry areas are to be planted with species like Honge, Bevu, Seetaphal *etc.*

<b>Dry Deciduous Species</b>			<b>Temp (°C)</b>	<b>Rainfall (mm)</b>
1.	Bevu	<i>Azadiracta indica</i>	21–32	400 –1,200
2.	Tapasi	<i>Holoptelia integrifolia</i>	20-30	500 - 1000
3.	Seetaphal	<i>Anona Squamosa</i>	20-40	400 - 1000
4.	Honge	<i>Pongamia pinnata</i>	20 -50	500–2,500
5.	Kamara	<i>Hardwickia binata</i>	25 -35	400 - 1000
6.	Bage	<i>Albezzia lebbek</i>	20 - 45	500 - 1000
7.	Ficus	<i>Ficus bengalensis</i>	20 - 50	500–2,500
8.	Sisso	<i>Dalbargia Sissoo</i>	20 - 50	500 -2000
9.	Ailanthus	<i>Ailanthus excelsa</i>	20 - 50	500 - 1000
10.	Hale	<i>Wrightia tinctoria</i>	25 - 45	500 - 1000
11.	Uded	<i>Steriospermum chelanoides</i>	25 - 45	500 -2000
12.	Dhupa	<i>Boswella Serrata</i>	20 - 40	500 - 2000
13.	Nelli	<i>Emblca Officinalis</i>	20 - 50	500 -1500
14.	Honne	<i>Pterocarpus marsupium</i>	20 - 40	500 - 2000
<b>Moist Deciduous Species</b>			<b>Temp (°C)</b>	<b>Rainfall (mm)</b>
15.	Teak	<i>Tectona grandis</i>	20 - 50	500-5000
16.	Nandi	<i>Legarstroemia lanceolata</i>	20 - 40	500 - 4000
17.	Honne	<i>Pterocarpus marsupium</i>	20 - 40	500 - 3000
18.	Mathi	<i>Terminalia alata</i>	20 -50	500 - 2000
19.	Shivane	<i>Gmelina arborea</i>	20 -50	500 -2000
20.	Kindal	<i>T.Paniculata</i>	20 - 40	500 - 1500
21.	Beete	<i>Dalbargia latifolia</i>	20 - 40	500 - 1500
22.	Tare	<i>T. belerica</i>	20 - 40	500 - 2000
23.	Bamboo	<i>Bambusa arundinasia</i>	20 - 40	500 - 2500
24.	Bamboo	<i>Dendrocalamus strictus</i>	20 – 40	500 – 2500
25.	Muthuga	<i>Butea monosperma</i>	20 - 40	400 - 1500
26.	Hippe	<i>Madhuca latifolia</i>	20 - 40	500 - 2000
27.	Sandal	<i>Santalum album</i>	20 - 50	400 - 1000
28.	Nelli	<i>Emblca officinalis</i>	20 - 40	500 - 2000
29.	Nerale	<i>Sizyzium cumini</i>	20 - 40	500 - 2000
30.	Dhaman	<i>Grevia tilifolia</i>	20 - 40	500 - 2000
31.	Kaval	<i>Careya arborea</i>	20 - 40	500 - 2000
32.	Harada	<i>Terminalia chebula</i>	20 - 40	500 - 2000



## References

1. FAO (1976) Framework for Land Evaluation, Food and Agriculture Organization, Rome.72 pp.
2. FAO (1983) Guidelines for Land Evaluation for Rainfed Agriculture, FAO, Rome, 237 pp.
3. IARI (1971) Soil Survey Manual, All India Soil and Land Use Survey Organization, IARI, New Delhi, 121 pp.
4. Katyal, J.C. and Rattan, R.K. (2003) Secondary and Micronutrients; Research Gap and future needs. Fert. News 48 (4); 9-20.
5. Naidu, L.G.K., Ramamurthy, V., Challa, O., Hegde, R. and Krishnan, P. (2006) Manual Soil Site Suitability Criteria for Major Crops, NBSS Publ. No. 129, NBSS &LUP, Nagpur, 118 pp.
6. Natarajan, A. and Dipak Sarkar (2010) Field Guide for Soil Survey, National Bureau of Soil Survey and Land Use Planning (ICAR), Nagpur, India.
7. Natarajan, A., Rajendra Hegde, Raj, J.N. and Shivananda Murthy, H.G. (2015) Implementation Manual for Sujala-III Project, Watershed Development Department, Bengaluru, Karnataka.
8. Sarma, V.A.K., Krishnan, P. and Budihal, S.L. (1987) Laboratory Manual, Tech. Bull. 23, NBSS &LUP, Nagpur.
9. Sehgal, J.L. (1990) Soil Resource Mapping of Different States of India; Why and How? National Bureau of Soil Survey and Land Use Planning, Nagpur, 49 pp.
10. Shivaprasad, C.R., R.S. Reddy, J. Sehgal and M. Velayuthum (1998) Soils of Karnataka for Optimizing Land Use, NBSS Publ. No. 47b, NBSS & LUP, Nagpur, India.
11. Soil Survey Staff (2006) Keys to Soil Taxonomy, Tenth edition, U.S. Department of Agriculture/ NRCS, Washington DC, U.S.A.
12. Soil Survey Staff (2012) Soil Survey Manual, Handbook No. 18, USDA, Washington DC, USA.



**Appendix I**  
**Hitnal (9D2d) Microwatershed**  
**Soil Phase Information**

Village	Survey Number	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Graveliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	WELLS	Land Capability	Conservation Plan
Chikkakasa nakandi	1	4.66	HDHcB2g 1	LMU-2	Moderately deep (75-100 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iies	Trench cum bunding
Chikkakasa nakandi	2	4.14	HDHcB2g 1	LMU-2	Moderately deep (75-100 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iies	Trench cum bunding
Chikkakasa nakandi	3	0.03	HDHcB2g 1	LMU-2	Moderately deep (75-100 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iies	Trench cum bunding
Chikkakasa nakandi	4	3.72	Habitatio n	Others	Others	Others	Others	Others	Others	Others	Not Available (NA)	Not Available	Others	Others
Chikkakasa nakandi	5	3.71	Habitatio n	Others	Others	Others	Others	Others	Others	Others	Not Available (NA)	Not Available	Others	Others
Chikkakasa nakandi	6	2.7	Habitatio n	Others	Others	Others	Others	Others	Others	Others	Not Available (NA)	Not Available	Others	Others
Chikkakasa nakandi	7	2.21	Habitatio n	Others	Others	Others	Others	Others	Others	Others	Not Available (NA)	Not Available	Others	Others
Chikkakasa nakandi	8	0.17	Habitatio n	Others	Others	Others	Others	Others	Others	Others	Not Available (NA)	Not Available	Others	Others
Chikkakasa nakandi	9	2.22	Habitatio n	Others	Others	Others	Others	Others	Others	Others	Not Available (NA)	Not Available	Others	Others
Chikkakasa nakandi	10	1.51	Habitatio n	Others	Others	Others	Others	Others	Others	Others	Not Available (NA)	Not Available	Others	Others
Chikkakasa nakandi	12	0.01	RO	RO	RO	RO	RO	RO	RO	RO	Not Available (NA)	Not Available	RO	RO
Chikkakasa nakandi	25	27.61	RO	RO	RO	RO	RO	RO	RO	RO	Not Available (NA)	Not Available	RO	RO
Chikkakasa nakandi	33	7.12	MKHhB2g 1	LMU-3	Moderately shallow (50-75 cm)	Sandy clay loam	Gravelly (15-35%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIies	Trench cum bunding
Chikkakasa nakandi	34	4.66	MI	MI	MI	MI	MI	MI	MI	MI	Not Available (NA)	Not Available	MI	MI
Chikkakasa nakandi	35	0.75	MI	MI	MI	MI	MI	MI	MI	MI	Not Available (NA)	Not Available	MI	MI
Chikkakasa nakandi	36	2.98	MKHhB2g 1	LMU-3	Moderately shallow (50-75 cm)	Sandy clay loam	Gravelly (15-35%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIies	Trench cum bunding
Chikkakasa nakandi	37	4.11	HDHcB2g 1	LMU-2	Moderately deep (75-100 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iies	Trench cum bunding
Chikkakasa nakandi	38	3.78	MKHhB2g 1	LMU-3	Moderately shallow (50-75 cm)	Sandy clay loam	Gravelly (15-35%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIies	Trench cum bunding
Chikkakasa nakandi	39	8.36	HDHcB2g 1	LMU-2	Moderately deep (75-100 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iies	Trench cum bunding
Hitnala	1	0.1	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	2	0.43	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	3	0.98	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS

Village	Survey Number	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	WELLS	Land Capability	Conservation Plan
Hitnala	4	1.04	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	5	1.52	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	6	0.83	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	7	1.65	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	8	0.38	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	9	0.65	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	10	0.46	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	11	1.52	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	12	1.37	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	13	1.24	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	14	1.52	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	15	1.08	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	16	0.53	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	17	1.07	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	18	0.64	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	19	1.45	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	20	0.96	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	21	3.5	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	22	4.92	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	23	0.93	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	24	0.77	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	25	1.16	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	26	0.67	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	27	1.77	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS



Village	Survey Number	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	WELLS	Land Capability	Conservation Plan
Hitnala	28	1.64	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	29	1.39	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	30	0.81	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	31	1.79	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	32	0.9	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	33	0.99	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	34	1.71	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	35	0.4	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	36	0.23	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	37	0.0002	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	38	0.2	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	39	2.17	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	40	1.75	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	41	0.26	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	42	0.97	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	43	1.33	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	44	1.8	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	45	1.15	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	46	1.17	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	47	1.4	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	48	1.32	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	49	2.28	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	50	0.49	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	51	1.3	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS

Village	Survey Number	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	WELLS	Land Capability	Conservation Plan
Hitnala	52	0.86	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	53	0.78	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	54	0.32	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	55	0.33	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	56	0.06	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	65	0.002	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	68	0.45	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	69	0.01	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	70	0.21	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	71	0.37	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	72	0.08	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	164	0.16	Habitatio n	Others	Others	Others	Others	Others	Others	Others	Not Available (NA)	Not Available	Others	Others
Hitnala	165	2.24	HDHcB2g 1	LMU-2	Moderately deep (75-100 cm)	Sandy loam	Gravelly (15- 35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iies	Trench cum bundling
Hitnala	188	0.92	Habitatio n	Others	Others	Others	Others	Others	Others	Others	Not Available (NA)	Not Available	Others	Others
Hitnala	189	0.68	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	190	1.82	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	191	0.98	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	192	0.35	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	193	4.09	LKRcB2g1	LMU-3	Moderately shallow (50-75 cm)	Sandy loam	Gravelly (15- 35%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIies	Trench cum bundling
Hitnala	194	6.33	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	195	5.86	HDHcB2g 1	LMU-2	Moderately deep (75-100 cm)	Sandy loam	Gravelly (15- 35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iies	Trench cum bundling
Hitnala	196	7.48	HDHcB2g 1	LMU-2	Moderately deep (75-100 cm)	Sandy loam	Gravelly (15- 35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iies	Trench cum bundling
Hitnala	197	4.07	HDHcB2g 1	LMU-2	Moderately deep (75-100 cm)	Sandy loam	Gravelly (15- 35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iies	Trench cum bundling
Hitnala	198	4.15	HDHcB2g 1	LMU-2	Moderately deep (75-100 cm)	Sandy loam	Gravelly (15- 35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iies	Trench cum bundling

Village	Survey Number	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Graveliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	WELLS	Land Capability	Conservation Plan
Hitnala	199	4.4	HDHcB2g1	LMU-2	Moderately deep (75-100 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iles	Trench cum bunding
Hitnala	200	1.76	LKRcB2g1	LMU-3	Moderately shallow (50-75 cm)	Sandy loam	Gravelly (15-35%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIles	Trench cum bunding
Hitnala	201	1.16	LKRcB2g1	LMU-3	Moderately shallow (50-75 cm)	Sandy loam	Gravelly (15-35%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIles	Trench cum bunding
Hitnala	202	2.15	HDHcB2g1	LMU-2	Moderately deep (75-100 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iles	Trench cum bunding
Hitnala	203	5.54	HDHcB2g1	LMU-2	Moderately deep (75-100 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iles	Trench cum bunding
Hitnala	204	6.66	LKRcB2g1	LMU-3	Moderately shallow (50-75 cm)	Sandy loam	Gravelly (15-35%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIles	Trench cum bunding
Hitnala	205	3.34	HDHcB2g1	LMU-2	Moderately deep (75-100 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iles	Trench cum bunding
Hitnala	206	0.13	LKRcB2g1	LMU-3	Moderately shallow (50-75 cm)	Sandy loam	Gravelly (15-35%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIles	Trench cum bunding
Hitnala	209	0.33	LKRhB2g1	LMU-3	Moderately shallow (50-75 cm)	Sandy clay loam	Gravelly (15-35%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIles	Trench cum bunding
Hitnala	210	1.29	LKRhB2g1	LMU-3	Moderately shallow (50-75 cm)	Sandy clay loam	Gravelly (15-35%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIles	Trench cum bunding
Hitnala	211	3.5	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	212	2.24	NS	NS	NS	NS	NS	NS	NS	NS	NS	Not Available	NS	NS
Hitnala	213	2.15	LKRcB2g1	LMU-3	Moderately shallow (50-75 cm)	Sandy loam	Gravelly (15-35%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIles	Trench cum bunding
Hitnala	214	0.42	LKRcB2g1	LMU-3	Moderately shallow (50-75 cm)	Sandy loam	Gravelly (15-35%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIles	Trench cum bunding
Hitnala	215	0.93	LKRcB2g1	LMU-3	Moderately shallow (50-75 cm)	Sandy loam	Gravelly (15-35%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIles	Trench cum bunding
Hitnala	216	0.82	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	217	1.21	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	218	4.43	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	219	2.52	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	220	1.76	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	221	0.23	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	222	1.61	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	223	2.32	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	224	2.9	LKRcB2g1	LMU-3	Moderately shallow (50-75 cm)	Sandy loam	Gravelly (15-35%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIles	Trench cum bunding

Village	Survey Number	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	WELLS	Land Capability	Conservation Plan
Hitnala	225	0.85	LKRcB2g1	LMU-3	Moderately shallow (50-75 cm)	Sandy loam	Gravelly (15-35%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIes	Trench cum bunding
Hitnala	226	0.27	LKRcB2g1	LMU-3	Moderately shallow (50-75 cm)	Sandy loam	Gravelly (15-35%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIes	Trench cum bunding
Hitnala	228	0.06	LKRcB2g1	LMU-3	Moderately shallow (50-75 cm)	Sandy loam	Gravelly (15-35%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIes	Trench cum bunding
Hitnala	229	1.22	LKRcB2g1	LMU-3	Moderately shallow (50-75 cm)	Sandy loam	Gravelly (15-35%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIes	Trench cum bunding
Hitnala	230	1.59	LKRcB2g1	LMU-3	Moderately shallow (50-75 cm)	Sandy loam	Gravelly (15-35%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIes	Trench cum bunding
Hitnala	238	3.06	KDTmA1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Not Available (NA)	Not Available	IIs	Strengthening of existing bunds/bunding
Hitnala	241	5.97	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	242	0.84	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	243	1.01	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	244	0.9	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	245	0.5	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Hitnala	246	0.87	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Shapura	43	0.2	HTIIB2	LMU-4	Moderately shallow (50-75 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIs	Trench cum bunding
Shapura	44	0.16	HTIIB2	LMU-4	Moderately shallow (50-75 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIs	Trench cum bunding
Agalakeri	230	0.84	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Agalakeri	231	0.34	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Agalakeri	232	0.16	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Agalakeri	233	0.77	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Agalakeri	234	0.94	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Agalakeri	235	0.7	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Agalakeri	236	0.65	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Agalakeri	237	0.87	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Agalakeri	238	0.003	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS

Village	Survey Number	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	WELLS	Land Capability	Conservation Plan
Agalakeri	239	0.01	NS	NS	NS	NS	NS	NS	NS	NS	Not Available (NA)	Not Available	NS	NS
Bevinahalli	46	0.33	HDHcB2g 1	LMU-2	Moderately deep (75-100 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Not available (NA)	Not Available	Iies	Trench cum bunding



**Appendix II**  
**Hitnal (9D2d) Microwatershed**  
**Soil Fertility Information**

Village	Survey No	Soil Reaction	Salinity	Organic Carbon	Available Phosphorus	Available Potassium	Available Sulphur	Available Boron	Available Iron	Available Manganese	Available Copper	Available Zinc
Chikkakasa nakandi	1	Slightly alkaline (pH 7.3 - 7.8)	Non saline (<2 dsm)	High (> 0.75 %)	High (> 57 kg/ha)	Medium (145 - 337 kg/ha)	Medium (10 - 20 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)
Chikkakasa nakandi	2	Slightly alkaline (pH 7.3 - 7.8)	Non saline (<2 dsm)	High (> 0.75 %)	High (> 57 kg/ha)	Low (<145 kg/ha)	Medium (10 - 20 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Deficient (< 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)
Chikkakasa nakandi	3	Slightly alkaline (pH 7.3 - 7.8)	Non saline (<2 dsm)	High (> 0.75 %)	High (> 57 kg/ha)	Low (<145 kg/ha)	Medium (10 - 20 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)
Chikkakasa nakandi	4	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Chikkakasa nakandi	5	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Chikkakasa nakandi	6	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Chikkakasa nakandi	7	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Chikkakasa nakandi	8	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Chikkakasa nakandi	9	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Chikkakasa nakandi	10	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Chikkakasa nakandi	12	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO
Chikkakasa nakandi	25	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO
Chikkakasa nakandi	33	Neutral (pH 6.5 - 7.3)	Non saline (<2 dsm)	High (> 0.75 %)	High (> 57 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)
Chikkakasa nakandi	34	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI
Chikkakasa nakandi	35	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI
Chikkakasa nakandi	36	Neutral (pH 6.5 - 7.3)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	Medium (10 - 20 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)
Chikkakasa nakandi	37	Neutral (pH 6.5 - 7.3)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 - 57 kg/ha)	Low (<145 kg/ha)	Medium (10 - 20 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)
Chikkakasa nakandi	38	Neutral (pH 6.5 - 7.3)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 - 57 kg/ha)	Low (<145 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)
Chikkakasa nakandi	39	Slightly alkaline (pH 7.3 - 7.8)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 - 57 kg/ha)	Low (<145 kg/ha)	Medium (10 - 20 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)
Hitnala	1	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	2	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	3	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	4	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS

Village	Survey No	Soil Reaction	Salinity	Organic Carbon	Available Phosphorus	Available Potassium	Available Sulphur	Available Boron	Available Iron	Available Manganese	Available Copper	Available Zinc
Hitnala	5	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	6	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	7	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	8	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	9	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	10	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	11	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	12	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	13	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	14	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	15	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	16	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	17	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	18	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	19	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	20	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	21	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	22	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	23	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	24	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	25	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	26	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	27	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	28	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	29	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	30	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	31	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	32	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	33	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	34	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS



Village	Survey No	Soil Reaction	Salinity	Organic Carbon	Available Phosphorus	Available Potassium	Available Sulphur	Available Boron	Available Iron	Available Manganese	Available Copper	Available Zinc
Hitnala	35	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	36	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	37	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	38	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	39	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	40	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	41	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	42	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	43	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	44	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	45	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	46	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	47	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	48	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	49	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	50	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	51	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	52	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	53	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	54	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	55	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	56	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	65	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	68	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	69	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	70	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	71	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	72	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	164	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Hitnala	165	Moderately alkaline (pH 7.8 - 8.4)	Non saline (<2 dsm )	High (> 0.75 %)	High (> 57 kg/ha)	Medium (145 - 337 kg/ha)	Medium (10 - 20 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)



Village	Survey No	Soil Reaction	Salinity	Organic Carbon	Available Phosphorus	Available Potassium	Available Sulphur	Available Boron	Available Iron	Available Manganese	Available Copper	Available Zinc
Hitnala	216	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	217	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	218	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	219	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	220	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	221	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	222	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	223	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	224	Moderately alkaline (pH 7.8 - 8.4)	Non saline (<2 dsm )	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	Medium (10 - 20 ppm)	Medium (0.5 - 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)
Hitnala	225	Moderately alkaline (pH 7.8 - 8.4)	Non saline (<2 dsm )	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	Medium (10 - 20 ppm)	Low (< 0.5 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)
Hitnala	226	Moderately alkaline (pH 7.8 - 8.4)	Non saline (<2 dsm )	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	Medium (10 - 20 ppm)	Low (< 0.5 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)
Hitnala	228	Moderately alkaline (pH 7.8 - 8.4)	Non saline (<2 dsm )	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	Medium (10 - 20 ppm)	Low (< 0.5 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)
Hitnala	229	Moderately alkaline (pH 7.8 - 8.4)	Non saline (<2 dsm )	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	Medium (10 - 20 ppm)	Medium (0.5 - 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Hitnala	230	Moderately alkaline (pH 7.8 - 8.4)	Non saline (<2 dsm )	Medium (0.5 - 0.75 %)	Low (< 23 kg/ha)	Medium (145 - 337 kg/ha)	Medium (10 - 20 ppm)	Medium (0.5 - 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Hitnala	238	Very strongly alkaline (pH > 9.0)	Non saline (<2 dsm )	High (> 0.75 %)	High (> 57 kg/ha)	High (> 337 kg/ha)	Low (<10 ppm)	High (> 1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)
Hitnala	241	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	242	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	243	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	244	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	245	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Hitnala	246	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Shapura	43	Strongly alkaline (pH 8.4 - 9.0)	Non saline (<2 dsm )	High (> 0.75 %)	High (> 57 kg/ha)	High (> 337 kg/ha)	Low (<10 ppm)	High (> 1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)
Shapura	44	Strongly alkaline (pH 8.4 - 9.0)	Non saline (<2 dsm )	High (> 0.75 %)	High (> 57 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	High (> 1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)
Agalakeri	230	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Agalakeri	231	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Agalakeri	232	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Agalakeri	233	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Agalakeri	234	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS

Village	Survey No	Soil Reaction	Salinity	Organic Carbon	Available Phosphorus	Available Potassium	Available Sulphur	Available Boron	Available Iron	Available Manganese	Available Copper	Available Zinc
Agalakeri	235	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Agalakeri	236	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Agalakeri	237	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Agalakeri	238	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Agalakeri	239	NS	NS	NS	NS	Quarry	NS	NS	NS	NS	NS	NS
Bevinahalli	46	Slightly alkaline (pH 7.3 - 7.8)	Non saline (<2 dsm )	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	Low (<145 kg/ha)	High (> 20 ppm)	Low (< 0.5 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)

**Appendix III**  
**Hitnal (9D2d) Microwatershed**  
**Soil Suitability Information**

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Jasmine	Bhendi	Brinjal	Crossandra	Drumstick	Mulberry	Onion	
Chikkakasa nakandi	1	S3r	S2g	S2rg	S2g	S2rg	S2rg	S3r	S2rg	S3t	S2rg	S2rg	S2g	S2rg	S2g	S2rg	S3r	S2rg	S2gt	S2g	S2g	S2g	S2rg	S2g	S2g	S2g	S2g	S2g	S2g	S2rg	S2rg	S2g	
Chikkakasa nakandi	2	S3r	S2g	S2rg	S2g	S2rg	S2rg	S3r	S2rg	S3t	S2rg	S2rg	S2g	S2rg	S2g	S2rg	S3r	S2rg	S2gt	S2g	S2g	S2g	S2g	S2rg	S2g	S2g	S2g	S2g	S2g	S2rg	S2rg	S2g	
Chikkakasa nakandi	3	S3r	S2g	S2rg	S2g	S2rg	S2rg	S3r	S2rg	S3t	S2rg	S2rg	S2g	S2rg	S2g	S2rg	S3r	S2rg	S2gt	S2g	S2g	S2g	S2g	S2rg	S2g	S2g	S2g	S2g	S2g	S2rg	S2rg	S2g	
Chikkakasa nakandi	4	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	
Chikkakasa nakandi	5	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	
Chikkakasa nakandi	6	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	
Chikkakasa nakandi	7	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	
Chikkakasa nakandi	8	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	
Chikkakasa nakandi	9	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	
Chikkakasa nakandi	10	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	
Chikkakasa nakandi	12	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	
Chikkakasa nakandi	25	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO
Chikkakasa nakandi	33	N1r	S3g	S3rg	S3g	S3rg	S3gt	N1r	S3rg	S3gt	S3rg	S3rg	S2rg	S3rg	S2rg	S3rg	S3rg	S3rg	S2rg	S3g	S3g	S3g	S3g	S3rg	S2rg	S3g	S3g	S3g	S3g	S3g	S3rg	S3rg	S3g
Chikkakasa nakandi	34	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI
Chikkakasa nakandi	35	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI
Chikkakasa nakandi	36	N1r	S3g	S3rg	S3g	S3rg	S3gt	N1r	S3rg	S3gt	S3rg	S3rg	S2rg	S3rg	S2rg	S3rg	S3rg	S3rg	S2rg	S3g	S3g	S3g	S3g	S3rg	S2rg	S3g	S3g	S3g	S3g	S3g	S3rg	S3rg	S3g
Chikkakasa nakandi	37	S3r	S2g	S2rg	S2g	S2rg	S2rg	S3r	S2rg	S3t	S2rg	S2rg	S2g	S2rg	S2g	S2rg	S3r	S2rg	S2gt	S2g	S2g	S2g	S2g	S2rg	S2g	S2g	S2g	S2g	S2g	S2g	S2rg	S2rg	S2g
Chikkakasa nakandi	38	N1r	S3g	S3rg	S3g	S3rg	S3gt	N1r	S3rg	S3gt	S3rg	S3rg	S2rg	S3rg	S2rg	S3rg	S3rg	S3rg	S2rg	S3g	S3g	S3g	S3g	S3rg	S2rg	S3g	S3g	S3g	S3g	S3g	S3rg	S3rg	S3g
Chikkakasa nakandi	39	S3r	S2g	S2rg	S2g	S2rg	S2rg	S3r	S2rg	S3t	S2rg	S2rg	S2g	S2rg	S2g	S2rg	S3r	S2rg	S2gt	S2g	S2g	S2g	S2g	S2rg	S2g	S2g	S2g	S2g	S2g	S2rg	S2rg	S2g	
Hitnala	1	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Chilly	Tomato	Margold	Chrysanthemum	Pomegranate	Bajra	Jasmine	Bhendi	Brinjal	Crossandra	Drumstick	Mulberry	Onion	
Hitnala	2	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	
Hitnala	3	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	4	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	5	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	6	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	7	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	8	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	9	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	10	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	11	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	12	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	13	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	14	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	15	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	16	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	17	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	18	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	19	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	20	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	21	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	22	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	23	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	24	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	25	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	26	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	27	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	28	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Jasmine	Bhendi	Brinjal	Crossandra	Drumstick	Mulberry	Onion	
Hitnala	29	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	
Hitnala	30	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	31	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	32	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	33	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	34	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	35	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	36	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	37	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	38	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	39	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	40	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	41	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	42	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	43	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	44	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	45	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	46	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	47	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	48	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	49	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	50	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	51	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	52	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	53	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	54	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	55	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Jasmine	Bhendi	Brinjal	Crossandra	Drumstick	Mulberry	Onion	
Hitnala	56	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	
Hitnala	65	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	68	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	69	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	70	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	71	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	72	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	164	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs
Hitnala	165	S3r	S2g	S2rg	S2g	S2rg	S2rg	S3r	S2rg	S3t	S2rg	S2rg	S2g	S2rg	S2g	S2rg	S3r	S2rg	S2gt	S2g	S2g	S2g	S2g	S2rg	S2g	S2g	S2g	S2g	S2g	S2rg	S2rg	S2g	S2g
Hitnala	188	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs
Hitnala	189	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	190	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	191	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	192	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	193	N1r	S3g	S3rg	S3g	S3rg	S3g	N1r	S3rg	S3gt	S3rg	S3rg	S2rg	S3rg	S2rg	S3rg	S3rg	S3rg	S2rg	S3g	S3g	S3g	S3g	S3rg	S2rg	S3g	S3g	S3g	S3g	S3rg	S3rg	S3g	S3g
Hitnala	194	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	195	S3r	S2g	S2rg	S2g	S2rg	S2rg	S3r	S2rg	S3t	S2rg	S2rg	S2g	S2rg	S2g	S2rg	S3r	S2rg	S2gt	S2g	S2g	S2g	S2g	S2rg	S2g	S2g	S2g	S2g	S2g	S2rg	S2rg	S2g	S2g
Hitnala	196	S3r	S2g	S2rg	S2g	S2rg	S2rg	S3r	S2rg	S3t	S2rg	S2rg	S2g	S2rg	S2g	S2rg	S3r	S2rg	S2gt	S2g	S2g	S2g	S2g	S2rg	S2g	S2g	S2g	S2g	S2g	S2rg	S2rg	S2g	S2g
Hitnala	197	S3r	S2g	S2rg	S2g	S2rg	S2rg	S3r	S2rg	S3t	S2rg	S2rg	S2g	S2rg	S2g	S2rg	S3r	S2rg	S2gt	S2g	S2g	S2g	S2g	S2rg	S2g	S2g	S2g	S2g	S2g	S2rg	S2rg	S2g	S2g
Hitnala	198	S3r	S2g	S2rg	S2g	S2rg	S2rg	S3r	S2rg	S3t	S2rg	S2rg	S2g	S2rg	S2g	S2rg	S3r	S2rg	S2gt	S2g	S2g	S2g	S2g	S2rg	S2g	S2g	S2g	S2g	S2g	S2rg	S2rg	S2g	S2g
Hitnala	199	S3r	S2g	S2rg	S2g	S2rg	S2rg	S3r	S2rg	S3t	S2rg	S2rg	S2g	S2rg	S2g	S2rg	S3r	S2rg	S2gt	S2g	S2g	S2g	S2g	S2rg	S2g	S2g	S2g	S2g	S2g	S2rg	S2rg	S2g	S2g
Hitnala	200	N1r	S3g	S3rg	S3g	S3rg	S3g	N1r	S3rg	S3gt	S3rg	S3rg	S2rg	S3rg	S2rg	S3rg	S3rg	S3rg	S2rg	S3g	S3g	S3g	S3g	S3rg	S2rg	S3g	S3g	S3g	S3g	S3rg	S3rg	S3g	S3g
Hitnala	201	N1r	S3g	S3rg	S3g	S3rg	S3g	N1r	S3rg	S3gt	S3rg	S3rg	S2rg	S3rg	S2rg	S3rg	S3rg	S3rg	S2rg	S3g	S3g	S3g	S3g	S3rg	S2rg	S3g	S3g	S3g	S3g	S3rg	S3rg	S3g	S3g
Hitnala	202	S3r	S2g	S2rg	S2g	S2rg	S2rg	S3r	S2rg	S3t	S2rg	S2rg	S2g	S2rg	S2g	S2rg	S3r	S2rg	S2gt	S2g	S2g	S2g	S2g	S2rg	S2g	S2g	S2g	S2g	S2g	S2rg	S2rg	S2g	S2g
Hitnala	203	S3r	S2g	S2rg	S2g	S2rg	S2rg	S3r	S2rg	S3t	S2rg	S2rg	S2g	S2rg	S2g	S2rg	S3r	S2rg	S2gt	S2g	S2g	S2g	S2g	S2rg	S2g	S2g	S2g	S2g	S2g	S2rg	S2rg	S2g	S2g
Hitnala	204	N1r	S3g	S3rg	S3g	S3rg	S3g	N1r	S3rg	S3gt	S3rg	S3rg	S2rg	S3rg	S2rg	S3rg	S3rg	S3rg	S2rg	S3g	S3g	S3g	S3g	S3rg	S2rg	S3g	S3g	S3g	S3g	S3rg	S3rg	S3g	S3g



Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Chilly	Tomato	Margold	Chrysanthemum	Pomegranate	Bajra	Jasmine	Bhendi	Brinjal	Crossandra	Drumstick	Mulberry	Onion	
Hitnala	205	S3r	S2g	S2rg	S2g	S2rg	S2rg	S3r	S2rg	S3t	S2rg	S2rg	S2g	S2rg	S2g	S2rg	S3r	S2rg	S2gt	S2g	S2g	S2g	S2g	S2rg	S2g	S2g	S2g	S2g	S2g	S2rg	S2rg	S2g	
Hitnala	206	N1r	S3g	S3rg	S3g	S3rg	S3g	N1r	S3rg	S3gt	S3rg	S3rg	S2rg	S3rg	S2rg	S3rg	S3rg	S3rg	S2rg	S3g	S3g	S3g	S3g	S3rg	S2rg	S3g	S3g	S3g	S3g	S3g	S3rg	S3rg	S3g
Hitnala	209	N1r	S3g	S3rg	S3g	S3rg	S3g	N1r	S3rg	S3gt	S3rg	S3rg	S2rg	S3rg	S2rg	S3rg	S3rg	S3rg	S2rg	S3g	S3g	S3g	S3g	S3rg	S2rg	S3g	S3g	S3g	S3g	S3g	S3rg	S3rg	S3g
Hitnala	210	N1r	S3g	S3rg	S3g	S3rg	S3g	N1r	S3rg	S3gt	S3rg	S3rg	S2rg	S3rg	S2rg	S3rg	S3rg	S3rg	S2rg	S3g	S3g	S3g	S3g	S3rg	S2rg	S3g	S3g	S3g	S3g	S3g	S3rg	S3rg	S3g
Hitnala	211	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	212	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	213	N1r	S3g	S3rg	S3g	S3rg	S3g	N1r	S3rg	S3gt	S3rg	S3rg	S2rg	S3rg	S2rg	S3rg	S3rg	S3rg	S2rg	S3g	S3g	S3g	S3g	S3rg	S2rg	S3g	S3g	S3g	S3g	S3rg	S3rg	S3g	
Hitnala	214	N1r	S3g	S3rg	S3g	S3rg	S3g	N1r	S3rg	S3gt	S3rg	S3rg	S2rg	S3rg	S2rg	S3rg	S3rg	S3rg	S2rg	S3g	S3g	S3g	S3g	S3rg	S2rg	S3g	S3g	S3g	S3g	S3rg	S3rg	S3g	
Hitnala	215	N1r	S3g	S3rg	S3g	S3rg	S3g	N1r	S3rg	S3gt	S3rg	S3rg	S2rg	S3rg	S2rg	S3rg	S3rg	S3rg	S2rg	S3g	S3g	S3g	S3g	S3rg	S2rg	S3g	S3g	S3g	S3g	S3rg	S3rg	S3g	
Hitnala	216	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	217	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	218	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	219	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	220	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	221	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	222	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	223	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	224	N1r	S3g	S3rg	S3g	S3rg	S3g	N1r	S3rg	S3gt	S3rg	S3rg	S2rg	S3rg	S2rg	S3rg	S3rg	S3rg	S2rg	S3g	S3g	S3g	S3g	S3rg	S2rg	S3g	S3g	S3g	S3g	S3rg	S3rg	S3g	
Hitnala	225	N1r	S3g	S3rg	S3g	S3rg	S3g	N1r	S3rg	S3gt	S3rg	S3rg	S2rg	S3rg	S2rg	S3rg	S3rg	S3rg	S2rg	S3g	S3g	S3g	S3g	S3rg	S2rg	S3g	S3g	S3g	S3g	S3rg	S3rg	S3g	
Hitnala	226	N1r	S3g	S3rg	S3g	S3rg	S3g	N1r	S3rg	S3gt	S3rg	S3rg	S2rg	S3rg	S2rg	S3rg	S3rg	S3rg	S2rg	S3g	S3g	S3g	S3g	S3rg	S2rg	S3g	S3g	S3g	S3g	S3rg	S3rg	S3g	
Hitnala	228	N1r	S3g	S3rg	S3g	S3rg	S3g	N1r	S3rg	S3gt	S3rg	S3rg	S2rg	S3rg	S2rg	S3rg	S3rg	S3rg	S2rg	S3g	S3g	S3g	S3g	S3rg	S2rg	S3g	S3g	S3g	S3g	S3rg	S3rg	S3g	
Hitnala	229	N1r	S3g	S3rg	S3g	S3rg	S3g	N1r	S3rg	S3gt	S3rg	S3rg	S2rg	S3rg	S2rg	S3rg	S3rg	S3rg	S2rg	S3g	S3g	S3g	S3g	S3rg	S2rg	S3g	S3g	S3g	S3g	S3rg	S3rg	S3g	
Hitnala	230	N1r	S3g	S3rg	S3g	S3rg	S3g	N1r	S3rg	S3gt	S3rg	S3rg	S2rg	S3rg	S2rg	S3rg	S3rg	S3rg	S2rg	S3g	S3g	S3g	S3g	S3rg	S2rg	S3g	S3g	S3g	S3g	S3rg	S3rg	S3g	
Hitnala	238	S3t	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2t	S1	S3t	S2t	S3t	S2t	S2t	S2t	S2t	S3t	S2t	S3t	S3t	S2t	S3t	S3t	
Hitnala	241	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	242	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Hitnala	243	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Jasmine	Bhendi	Brinjal	Crossandra	Drumstick	Mulberry	Onion				
Hitnala	244	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS			
Hitnala	245	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS		
Hitnala	246	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	
Shapura	43	N1r	S2r	S3r	S2r	S3r	S2r	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	S3r	S3r	S3r	S2rt	S2rg	S2rg	S2rg	S2rg	S3r	S2r	S2rg	S2rg	S2rg	S2rg	S3r	S3r	S2rg	S2rg			
Shapura	44	N1r	S2r	S3r	S2r	S3r	S2r	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	S3r	S3r	S3r	S2rt	S2rg	S2rg	S2rg	S2rg	S3r	S2r	S2rg	S2rg	S2rg	S2rg	S3r	S3r	S2rg	S2rg			
Agalakeri	230	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS		
Agalakeri	231	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	
Agalakeri	232	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	
Agalakeri	233	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	
Agalakeri	234	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Agalakeri	235	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Agalakeri	236	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Agalakeri	237	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Agalakeri	238	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Agalakeri	239	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Bevinahalli	46	S3r	S2g	S2rg	S2g	S2rg	S2rg	S3r	S2rg	S3t	S2rg	S2rg	S2g	S2rg	S2g	S2rg	S3r	S2rg	S2gt	S2g	S2g	S2g	S2g	S2rg	S2g	S2g	S2g	S2g	S2g	S2g	S2g	S2rg	S2rg	S2g	S2g	

RO-Rock outcrops, MI-Mining/industry, NS- Not surveyed

# **PART-B**

**SOCIO-ECONOMIC STATUS OF FARM HOUSEHOLDS**



## CONTENTS

1	Findings of the socio-economic survey	1-3
2	Introduction	5
3	Methodology	7-8
4	Salient features of the survey	9-24
5	Summary	25-29



## LIST OF TABLES

1	Households sampled for socio economic survey	9
2	Population characteristics	9
3	Age wise classification of household members	9
4	Education level of household members	10
5	Occupation of household heads	10
6	Occupation of family members	10
7	Institutional participation of household members	11
8	Type of house owned by households	11
9	Durable assets owned by households	11
10	Average value of durable assets owned by households	11
11	Farm implements owned by households	12
12	Average value of farm implements	12
13	Livestock possession by households	12
14	Average labour availability	13
15	Adequacy of hired labour	13
16	Distribution of land (ha)	13
17	Average land value (Rs./ha)	13
18	Status of bore wells	14
19	Source of irrigation	14
20	Depth of water (Avg in meters)	14
21	Irrigated area (ha)	14
22	Cropping pattern	14
23	Cropping intensity	15
24	Possession of bank account and saving	15
25	Borrowing status	15
26	Source of credit	15
27	Avg. credit borrowed	15
28	Purpose of credit borrowed from institutional sources	16
29	Repayment status of household from institutional sources	16
30	Opinion regarding institutional sources of credit	16
31. a	Cost of cultivation of Maize	17

31. b	Cost of cultivation of Bajra	18
31. c	Cost of cultivation of Paddy	19
32	Adequacy of fodder	20
33	Annual gross income	20
34	Average annual expenditure	20
35	Horticultural species grown	20
36	Forest species grown	21
37	Marketing of the agricultural produce	21
38	Marketing channels used for sale of agricultural produce	21
39	Mode of transport of agricultural produce	21
40	Incidence of soil and water erosion problems	22
41	Interest shown towards soil testing	22
42	Usage pattern of fuel for domestic use	22
43	Source of drinking water	22
44	Source of light	22
45	Existence of sanitary toilet facility	22
46	Possession of public distribution system (PDS) card	23
47	Participation in NREGA programme	23
48	Adequacy of food items	23
49	Inadequacy of food items	23
50	Response on market surplus of food items	24
51	Farming constraints experienced	24



**FINDINGS OF THE SOCIO-ECONOMIC SURVEY**

- ❖ *The survey was conducted in Hitnal is located at North latitude 15<sup>0</sup> 20' 35.836'' and 15<sup>0</sup> 19' 2.884'' and East longitude 76<sup>0</sup> 19' 29.484'' and 76<sup>0</sup> 17' 1.088'' covering an area of about 351.12 ha coming under Hitnal and Chikkasanakandi villages of Koppal taluk.*
- ❖ *Socio-economic analysis of Hitnal micro watersheds of Shahpura sub-watershed, Koppala taluk & District indicated that, out of the total sample of 34 total respondents, 10 (29.41 %) were marginal, 6 (17.65%) were small, 9 (26.47 %) were Semi medium and 4 (11.76 %) were medium farmers.*
- ❖ *The population characteristics of households indicated that, there were 84 (56.00%) men and 66 (44.00 %) were women.*
- ❖ *Majority of the respondents (48.00%) were in the age group of 16-35 years.*
- ❖ *Education level of the sample households indicated that, there were 47.33 per cent of illiterates, 18.67 per cent of them had primary school education, 8.67 per cent middle school education, and 7.33 per cent high school education, 11.33 per cent of them had PUC education, 0.67 per cent of them had Diploma, 2.00 per cent attained graduation.*
- ❖ *About, 91.18 per cent of household heads practicing agriculture and 8.82 per cent of the household heads were engaged as agricultural labourers.*
- ❖ *Agriculture was the major occupation for 21.33 per cent of the household members.*
- ❖ *In the study area, 88.24 per cent of the households possess katcha house and 8.82 per cent possess pucca house.*
- ❖ *The durable assets owned by the households showed that, 70.59 per cent possess TV, 32.35 per cent possess mixer grinder, 85.29 per cent possess mobile phones and 50.00 per cent possess motor cycles.*
- ❖ *Farm implements owned by the households indicated that, 14.71 per cent of the households possess plough, 5.88 per cent possess tractor, 5.88 per cent possess bullock cart and 2.94 per cent possess sprayer.*
- ❖ *Regarding livestock possession by the households, 17.65 per cent possess local cow and 2.94 per cent possess buffalo.*
- ❖ *The average labour availability in the study area showed that, own men and women labour availability in the micro watershed was 21.71 each, while the hired labour (men) availability was 1.76.*
- ❖ *Further, 88.24 per cent of the households opined that hired labour was inadequate during the agricultural season.*
- ❖ *Out of the total land holding of the sample respondents 67.15 per cent (45.93 ha) of the area is under dry condition and the remaining 32.85 per cent area is irrigated land.*

- ❖ *There were 5.00 live bore wells and 3.00 dry bore wells among the sampled households.*
- ❖ *Bore/open well was the major source of irrigation for 20.59 per cent of the households.*
- ❖ *The major crops grown by sample farmers are Maize, Bajra, Paddy, 0 and 0 and cropping intensity was recorded as 100.00 per cent.*
- ❖ *Out of the sample households 85.29 percent possessed bank account and 85.29 per cent of them have savings in the account.*
- ❖ *About 85.29 per cent of the respondents borrowed credit from various sources.*
- ❖ *Among the credit borrowed by households, 3.45 per cent have borrowed loan from commercial banks.*
- ❖ *Majority of the respondents (100.00%) have borrowed loan for agriculture purpose.*
- ❖ *Regarding the opinion on institutional sources of credit, 100.00 per cent of the households opined that credit helped to perform timely agricultural operations.*
- ❖ *The per hectare cost of cultivation for Maize, Bajra, and Paddy was Rs.22744.14 , 24612.64 and 46877.36 with benefit cost ratio of 1:1.50, 1: 1.02 and 1: 1.20 respectively.*
- ❖ *Further, 23.53 per cent of the households opined that dry fodder was adequate.*
- ❖ *The average annual gross income of the farmers was Rs. 60641.18 in micro-watershed, of which Rs. 46611.76 comes from agriculture.*
- ❖ *Sampled households have grown 1 horticulture trees and 4 forestry trees together in the fields and back yards.*
- ❖ *Regarding marketing channels, 11.76 per cent of the households have sold agricultural produce to the local/village merchants, while, 70.59 per cent have sold in regulated markets.*
- ❖ *Further, 82.35 per cent of the households have used tractor for the transport of agriculture commodity.*
- ❖ *Majority of the farmers (85.29%) have experienced soil and water erosion problems in the watershed and 82.35 per cent of the households were interested towards soil testing.*
- ❖ *Fire was the major source of fuel for domestic use for 100.00 per cent of the households.*
- ❖ *Piped supply was the major source for drinking water for 100.00 per cent of the households.*
- ❖ *Electricity was the major source of light for 100.00 per cent of the households.*
- ❖ *In the study area, 100.00 per cent of the households possess toilet facility.*
- ❖ *Regarding possession of PDS card, 100.00 per cent of the households possessed BPL card.*

- ❖ *Households opined that, the requirement of cereals (91.18%), pulses (88.24%) and oilseeds (17.65%) are adequate for consumption.*
- ❖ *Farming constraints experienced by households in the micro watersheds were lower fertility status of the soil (85.29%) wild animal menace on farm field (76.47%), frequent incidence of pest and diseases (55.88%), inadequacy of irrigation water (20.59%), high cost of fertilizers and plant protection chemicals (38.24%), high rate of interest on credit (8.82%), low price for the agricultural commodities (14.71%), lack of marketing facilities in the area (11.76%), inadequate extension services (14.71%), lack of transport for safe transport of the agricultural produce to the market (38.24%), Less rainfall (32.35%) and Source of Agri-technology information (Newspaper/ TV/Mobile) (38.24%).*



## INTRODUCTION

Soil and water are the two precious natural resources which are essential for crop production and existence of life on earth. Rainfed agriculture is under severe stress due to various constraints related to agriculture like uneven and erratic distribution of rainfall, indiscriminate use of fertilizers, chemicals and pesticides, adoption of improper land management practices, soil erosion, decline in soil fertility, decline in ground water resources leading to low crop productivity. The area under rainfed agriculture has to be managed effectively using the best available practices to enhance the production of food, fodder and fuel. This is possible if the land resources are characterized at each parcel of land through detailed land resource inventory using the best available techniques of remote sensing, GPS and GIS. The watershed development programs are aimed at the sustainable distribution of its resources and the process of creating and implementing plans, programs, and projects to sustain and enhance watershed functions that affect the plant, animal and human communities within a watershed boundary.

World Bank funded KWDP II, SUJALA III project was implemented in with Broad objective of demonstrating more effective watershed management through greater integration of programmes related to rain-fed agriculture, innovative and science based approaches and strengthen institutional capacities and If successful, it is expected that the systems and tools could be mainstreamed into the overall IWMP in the State of Karnataka and in time, throughout other IWMP operations in India. With this background the socio-economic survey has been carried out with following specific objectives:

1. To understand the demographic features of the households in the micro-watershed
2. To understand the extent of family labour available and additional employment opportunities available within the village.
3. To know the status of assets of households in the micro-watershed for suggesting possible improvements.
4. To study the cropping pattern, cropped area and productivity levels of different households in micro-watershed.
5. To determine the type and extent of livestock owned by different categories of HHs
6. Availability of fodder and level of livestock management.

### **Scope and importance of survey**

Survey helps in identification of different socio-economic and resource use-patterns of farmers at the Micro watershed. Household survey provides demographic features, labour force, and levels of education; land ownership and asset position (including livestock and other household assets) of surveyed households; and cropping patterns, input intensities, and average crop yields from farmers' fields. It also discusses crop utilization and the degree of commercialization of production in the areas; farmers' access to and utilization of credit from formal and informal sources; and the level of adoption and use of soil, water, and pest management technologies.



## METHODOLOGY

The description of the methods, components selected for the survey and procedures followed in conducting the baseline survey are furnished under the following heads.

### 1. Description of the study area

Koppal district is an administrative district in the state of Karnataka in India. In the past Koppal was referred to as 'Kopana Nagara'. Koppal, now a district headquarters is ancient Kopana a major holy place of the Jainas. The district occupies an area of 7,190 km<sup>2</sup> and has a population of 1,196,089, which 16.58% were urban as of 2001. The Koppal district was formed after split of Raichur district.

Geographers are very particular about the physiography or relief of a region. It plays a very important role in the spatial analysis of agricultural situation of the study area. The undulating topography with black cotton soil shrips, cut across by numerous nalas or streams is the major characteristic feature of the study region. Three physiographic divisions have made considering the local conditions of landforms and crops grown in the district. On the basis of physiography, Koppal district can be divided into three major divisions. They are (a) Koppal & Yelburga plateau, (b) Maidan division, (c) Tungabhadra valley. The district is part of Krishna basin the main streams draining the area are Maskinala, Ilkal-nadi and Hirenala. These are Ephemeral in nature, these come under Tungabhadra sub-basin. The drainage exhibit dentritic to subdentritic with drainage density varies from 1.4 to 7.0kms/sq.km.

According to the 2011 census Koppal district has a population of 1,391,292, roughly equal to the nation of Swaziland or the US state of Hawaii. This gives it a ranking of 350th in India (out of a total of 640). The district has a population density of 250 inhabitants per square kilometre (650/sq mi). Its population growth rate over the decade 2001-2011 was 16.32%.Koppal has a sex ratio of 983 females for every 1000 males, and a literacy rate of 67.28%.

### 2. Locale of the survey and description of the micro-watershed and

The study was conducted in Hitnal micro-watershed (Shahpura sub-watershed, Koppala taluk & District) is located at North latitude 15<sup>0</sup> 20' 35.836" and 15<sup>0</sup> 19' 2.884" and East longitude 76<sup>0</sup> 19' 29.484" and 76<sup>0</sup> 17' 1.088" covering an area of about 351.12 ha bounded by under Hitnal and Chikkasanakandi Villages.

### 3. Selection of the respondents for the study

The micro-watershed is marked with 320 square meters grids. One farmer from every alternate grid in the micro-watershed was selected for the study and interviewed for socio-economic data. Totally 34 households were interviewed for the survey.

#### **4. The parameters considered for socio-economic survey of households**

Two forms of data were collected from the micro-watershed which includes primary data from the farm households and secondary data about the villages under the micro-watershed jurisdiction.

The following parameters were considered for the primary data collection about the socio-economic data of the households, (1) Demographic information, (2) Farm and durable assets owned by households, (3) Livestock possession, (4) Labour availability, (5) Level of migration in the village, Land holding, (7) Cropping pattern, (8) Source of irrigation, (9) Borrowing status, (10) Cost of cultivation of major crops, (11) Economics of subsidiary activities, (12) Fodder availability, (13) Family annual income from different sources, (14) Horticulture and forestry species grown, (15) Additional investment capacity, (16) Marketing practices, (17) Status of soil and water conservation structure, (18) Access to basic needs and (19) Constraints and suggestion.

The following parameters were considered for the secondary data regarding the villages under the micro-watershed jurisdiction, (1) Number of villages in each micro-watershed jurisdiction, (2) Village wise number of households, (3) Geographical area of the villages, (4) Cultivable area including rainfed and irrigated, (5) Number and type of house in each village, (6) Human and livestock population, (7) Facilities in the village such as roads, transport facility for conveyance, drinking water supply, street light and (8) Community based organizations in the villages.

#### **5. Development of interview schedule and data collection**

Taking into the consideration the objectives of the survey, an interview schedule was prepared after thorough consultation with the experts in the field of social sciences. A comprehensive interview schedule covering all the major parameters for measuring the socio-economic situation was developed.

#### **6. Tools used to analyze the data**

The statistical components such as frequency and percentage were used to analyze the data.

##### **Abbreviations used in the report**

LL=Landless

MF=Marginal Farmers

SF=Small farmers

SMF=Semi medium farmers

MDF=Medium farmers

LF=Large Farmers



## FINDINGS OF THE SURVEY

This chapter deals with systematic presentation of results of the survey. Keeping in view the objectives, the salient features of the survey are presented under the following headings.

**Households sampled for socio-economic survey:** The data on households sampled for socio economic survey in Hitnal Micro watershed is presented in Table 1 and it indicated that 34 farmers were sampled in Hitnal micro-watershed among households surveyed 10 (29.41%) were marginal, 6 (17.65%) were small, 9 (26.47 %) were semi medium, 4 (11.76 %) were medium farmers. 5 landless farmers were also interviewed for the survey.

**Table 1. Households sampled for socio economic survey in Hitnal micro-watershed**

Sl.No.	Particulars	LL (5)		MF (10)		SF (6)		SMF (9)		MDF (4)		All (34)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Farmers	5	14.7	10	29.4	6	17.7	9	26.5	4	11.8	34	100

**Population characteristics:** The population characteristics of households sampled for socio-economic survey in Hitnal Micro watershed is presented in Table 2. The data indicated that, there were 84 (56.00%) men and 66 (44.00%) were women.

**Table 2. Population characteristics in Hitnal micro-watershed**

Sl.N.	Particulars	LL (17)		MF (51)		SF (26)		SMF (35)		MDF (21)		All (150)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Men	8	47.1	31	61	15	58	20	57.1	10	47.6	84	56
2	Women	9	52.9	20	39	11	42	15	42.9	11	52.4	66	44
Total		17	100	51	100	26	100	35	100	21	100	150	100
Average		3.4		5.1		4.3		3.9		5.3		4.4	

**Age wise classification of population:** The age wise classification of household members in Hitnal Micro watershed is presented in Table 3. The indicated that, 14 (9.33%) of population were 0-15 years of age, 72 (48.00%) were 16-35 years of age, 53(35.33%) were 36-60 years of age and 11 (7.33 %) were above 61 years of age.

**Table 3: Age wise classification of members of the household in Hitnal micro-watershed**

Sl.No.	Particulars	LL (17)		MF (51)		SF (26)		SMF (35)		MDF (21)		All (150)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	0-15 years of age	1	5.88	7	13.7	3	11.5	1	2.86	2	9.5	14	9.33
2	16-35 years of age	10	58.8	24	47.1	12	46.2	16	45.71	10	48	72	48
3	36-60 years of age	5	29.4	18	35.3	8	30.8	16	45.71	6	29	53	35.33
4	> 61 years	1	5.88	2	3.92	3	11.5	2	5.71	3	14	11	7.33
Total		17	100	51	100	26	100	35	100	21	100	150	100

**Education level of household members:** Education level of household members in Hitnal Micro watershed is presented in Table 4. The results indicated that, there were 47.33 per cent of illiterates, 18.67 per cent of them had primary school education, 8.67 per cent middle school education, and 7.33 per cent high school education, 11.33 per cent

of them had PUC education, 0.67 per cent of them had Diploma, 2.00 per cent attained graduation and 1.33 them had other education.

**Table 4. Education level of members of the household in Hitnal micro-watershed**

Sl.No.	Particulars	LL (17)		MF (51)		SF (26)		SMF (35)		MDF (21)		All (150)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Illiterate	8	47.1	21	41.2	14	53.9	15	42.9	13	61.9	71	47.3
2	Primary School	3	17.7	15	29.4	2	7.69	6	17.1	2	9.52	28	18.7
3	Middle School	0	0	4	7.84	3	11.5	5	14.3	1	4.76	13	8.67
4	High School	1	5.88	1	1.96	1	3.85	5	14.3	3	14.29	11	7.33
5	PUC	5	29.4	5	9.8	5	19.2	1	2.86	1	4.76	17	11.3
6	Diploma	0	0	1	1.96	0	0	0	0	0	0	1	0.67
7	ITI	0	0	3	5.88	0	0	1	2.86	0	0	4	2.67
8	Degree	0	0	0	0	1	3.85	1	2.86	1	4.76	3	2
9	Others	0	0	1	1.96	0	0	1	2.86	0	0	2	1.33
Total		17	100	51	100	26	100	35	100	21	100	150	100

**Occupation of head of households:** The data regarding the occupation of the household heads in Hitnal Micro watershed is presented in Table 5. The results indicate that, 91.18 per cent of households heads were practicing agriculture, 8.82 per cent of the household heads were agricultural Labour.

**Table 5: Occupation of heads of households in Hitnal micro-watershed**

Sl.No.	Particulars	LL (5)		MF (10)		SF (6)		SMF (9)		MDF (4)		All (34)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Agriculture	2	40	10	100	6	100	9	100	4	100	31	91.18
2	Agricultural Labour	3	60	0	0	0	0	0	0	0	0	3	8.82
Total		5	100	10	100	6	100	9	100	4	100	34	100

**Occupation of the members of the household:** The data regarding the occupation of the household members in Hitnal Micro watershed is presented in Table 6. The results indicate that, agriculture was the major occupation for 21.33 per cent of the household members, 72.00 per cent were agricultural labour, 5.33 per cent were working in pursuing education and 0.67 per cent were childrens.

**Table 6: Occupation of members of the household in Hitnal micro-watershed**

Sl.No.	Particulars	LL (17)		MF (51)		SF (26)		SMF (35)		MDF (21)		All (150)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Agriculture	2	11.8	10	19.6	7	26.92	9	25.71	4	19	32	21.3
2	Agricultural Labour	15	88.2	37	72.6	17	65.38	23	65.71	16	76	108	72
3	Private Service	0	0	0	0	0	0	1	2.86	0	0	1	0.67
4	Student	0	0	3	5.88	2	7.69	2	5.71	1	4.8	8	5.33
5	Children	0	0	1	1.96	0	0	0	0	0	0	1	0.67
Total		17	100	51	100	26	100	35	100	21	100	150	100

**Institutional Participation of household members:** The data regarding the institutional participation of the household members in Hitnal Micro watershed is presented in Table 7. The results show that, out of the total family members in the households were not participating in any of the institutions.

**Table 7: Institutional Participation of household member in Hitnal micro-watershed**

Sl.No.	Particulars	LL (17)		MF (51)		SF (26)		SMF (35)		MDF (21)		All (150)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	No Participation	17	100	51	100	26	100	35	100	21	100	150	100
	Total	17	100	51	100	26	100	35	100	21	100	150	100

**Type of house owned:** The data regarding the type of house owned by the households in Hitnal Micro watershed is presented in Table 8. The results indicate that, 2.94 percent possess thatched house, 88.24 per cent of the households possess katcha house and 8.82 per cent possess pucca house.

**Table 8. Type of house owned by households in Hitnal micro-watershed**

Sl.No.	Particulars	LL (5)		MF (10)		SF (6)		SMF (9)		MDF (4)		All (34)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Thatched	0	0	1	10	0	0	0	0	0	0	1	2.94
2	Katcha	5	100	8	80	5	83.33	8	88.9	4	100	30	88.24
3	Pucca/RCC	0	0	1	10	1	16.67	1	11.1	0	0	3	8.82
	Total	5	100	10	100	6	100	9	100	4	100	34	100

**Durable assets owned by the households:** The data regarding the Durable Assets owned by the households in Hitnal Micro watershed is presented in Table 9. The results shows that, 70.59 per cent possess TV, 32.35 per cent possess mixer grinder, 5.88 per cent possess Bicycle, 50.00 per cent possess motor cycle and 85.29 per cent possess mobile phones.

**Table 9. Durable assets owned by households in Hitnal micro-watershed**

Sl.No.	Particulars	LL (5)		MF (10)		SF (6)		SMF (9)		MDF (4)		All (34)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Television	0	0	8	80	5	83.3	8	89	3	75	24	70.59
2	Mixer/Grinder	0	0	5	50	3	50	2	22	1	25	11	32.35
3	Bicycle	0	0	2	20	0	0	0	0	0	0	2	5.88
4	Motor Cycle	0	0	4	40	6	100	4	44	3	75	17	50
5	Mobile Phone	2	40	9	90	5	83.3	9	100	4	100	29	85.29
6	Blank	3	60	0	0	0	0	0	0	0	0	3	8.82

**Table 10. Average value of durable assets owned in Hitnal micro-watershed**

Sl.No.	Particulars	Average Value (Rs.)					
		LL (5)	MF (10)	SF (6)	SMF (9)	MDF (4)	All (34)
1	Television	0	8062	4400	12250	4833	8291
2	Mixer/Grinder	0	1700	1900	1800	2000	1800
3	Bicycle	0	1500	0	0	0	1500
4	Motor Cycle	0	45000	34333	47500	46666	42117
5	Mobile Phone	2500	2125	2300	1961	1220	1989

**Average value of durable assets:** The data regarding the average value of durable assets owned by the households in Hitnal Micro watershed is presented in Table 10. The result shows that, the average value of television was Rs.8291.00, mixer grinder was

Rs.1800.00, bicycle was Rs.1500.00, motor cycle was Rs. 42117.00 and mobile phone was Rs.1989.00.

**Farm implements owned:** The data regarding the farm implements owned by the households in Hitnal Micro watershed is presented in Table 11. About 5.88 per cent of the households possess Bullock Cart, 14.71 per cent possess plough, 2.94 per cent possess Sprayer, 41.18 per cent possess Weeder and 5.88 per cent possess tractor.

**Table 11. Farm implements owned in Hitnal micro-watershed**

Sl.No.	Particulars	LL (5)		MF (10)		SF (6)		SMF (9)		MDF (4)		All (34)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Bullock Cart	0	0	1	10	0	0	1	11.1	0	0	2	5.88
2	Plough	0	0	3	30	1	16.67	1	11.1	0	0	5	14.71
3	Irrigation Pump	0	0	0	0	1	16.67	0	0	0	0	1	2.94
4	Tractor	0	0	0	0	2	33.33	0	0	0	0	2	5.88
5	Sprayer	0	0	0	0	1	16.67	0	0	0	0	1	2.94
6	Weeder	0	0	5	50	3	50	4	44.4	2	50	14	41.18
7	Thresher	0	0	2	20	1	16.67	2	22.2	2	50	7	20.59
8	Blank	5	100	5	50	1	16.67	5	55.6	2	50	18	52.94

**Average value of farm implements:** The data regarding the average value of farm Implements owned by the households in Hitnal Micro watershed is presented in Table 12. The results show that the average value of plough was Rs.5258.00, bullock Cart was Rs.9500.00, seed/fertilizer drill was Rs.15000.00, sprayer and weeder was Rs.126 and tractor was Rs. 450000.

**Table 12. Average value of farm implements in Hitnal micro-watershed**  
**Average Value (Rs.)**

Sl.No.	Particulars	LL (5)	MF (10)	SF (6)	SMF (9)	MDF (4)	All (34)
1	Bullock Cart	0	18000	0	1000	0	9500
2	Plough	0	9600	375	2000	0	5258
3	Irrigation Pump	0	0	2000	0	0	2000
4	Tractor	0	0	450000	0	0	450000
5	Sprayer	0	0	15000	0	0	15000
6	Weeder	0	97	88	178	166	126
7	Thresher	0	325	200	387	325	316

**Table 13. Livestock possession by households in Hitnal micro-watershed**

Sl.No.	Particulars	LL (5)		MF (10)		SF (6)		SMF (9)		MDF (4)		All (34)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Bullock	0	0	1	10	1	16.67	1	11	1	25	4	11.76
2	Local cow	0	0	1	10	2	33.33	1	11	2	50	6	17.65
3	Crossbred cow	0	0	0	0	0	0	1	11	0	0	1	2.94
4	Buffalo	0	0	1	10	0	0	0	0	0	0	1	2.94
5	Sheep	0	0	0	0	1	16.67	0	0	0	0	1	2.94
6	blank	5	100	8	80	3	50	7	78	2	50	25	73.53

**Livestock possession by the households:** The data regarding the Livestock possession by the households in Hitnal Micro watershed is presented in Table 13. The indicate that,

11.76 per cent of the households possess bullocks, 17.65 per cent possess local cow, 2.94 per cent possess buffalo, 2.94 per cent possess crossbred cow and 2.94 per cent possess sheep.

**Average Labour availability:** The data regarding the average labour availability in Hitnal Micro watershed is presented in Table 14. The indicated that, own labour men available in the micro watershed was 20.52, women available in the micro watershed was 1.19, hired labour (men) available was 1.76 and hired labour (women) available was 19.97.

**Table 14. Average labour availability in Hitnal micro-watershed**

Sl.No.	Particulars	LL (5)		MF (10)		SF (6)		SMF (9)		MDF (4)		All (34)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Hired labour Female	0	0	24.6	81.1	26	73.1	21	94.3	16.3	0	30.84	20.52
2	Own Labour Female	0	0	1.33	18.9	1.33	26.9	1.33	5.7	1.25	10.66	15.09	1.19
3	Own labour Male	0	0	1.8	1.92	1.67	2.3	2.3	1.21	1.75	1.75	1.76	1.76
4	Hired labour Male	0	0	24.5	71.3	24	67.1	21	58.3	15	41.7	19.97	19.97

**Adequacy of hired labour:** The data regarding the adequacy of hired labour in Hitnal Micro watershed is presented in Table 15. The results indicate that, 8.82 per cent of the household opined that hired labour was adequate, 88.24 per cent of the household opined that hired labour was Inadequate.

**Table 15. Adequacy of hired labour in Hitnal micro-watershed**

Sl.No.	Particulars	LL (5)		MF (10)		SF (6)		SMF (9)		MDF (4)		All (34)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Adequate	0	0	1	10	1	16.7	0	0	1	25	3	8.82
2	Inadequate	3	60	9	90	5	83.3	10	111	3	75	30	88.2

**Distribution of land (ha):** The data regarding the distribution of land (ha) in Hitnal Micro watershed is presented in Table 16. The results indicate that, 30.84 ha (67.15%) of dry land and 15.09 ha (32.85 %) of irrigated land.

**Table 16. Distribution of land (ha) in Hitnal micro-watershed**

Sl.No.	Particulars	LL (5)		MF (10)		SF (6)		SMF (9)		MDF (4)		All (34)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Dry	0	0	5.56	81.1	5.21	73.1	20.07	94.3	0	0	30.84	67.15
2	Irrigated	0	0	1.3	18.9	1.92	26.9	1.21	5.7	10.66	100	15.09	32.85
Total		0	100	6.85	100	7.13	100	21.29	100	10.66	100	45.93	100

**Average value of land (ha):** The data regarding the average land value (Rs./ha) in Hitnal Micro watershed is presented in Table 17. The results show that the average value of dry land was Rs.259283.56, and the average value of irrigated land was Rs.351153.43.

**Table 17. Average value of land (ha) in Hitnal micro-watershed**

Sl.No.	Particulars	LL (5)		MF (10)		SF (6)		SMF (9)		MDF (4)		All (34)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Dry	0	0	521704.3	81.1	249301.2	73.1	189233.9	94.3	0	0	259283.6	67.15
2	Irrigated	0	0	1157812	18.9	677426.2	26.9	411666.7	5.7	187547.5	100	351153.4	32.85

**Status of bore wells:** The data regarding the status of bore wells in Hitnal Micro watershed is presented in Table 18. The results indicate that, there were 3 De-functioning bore wells and 5 functioning bore wells among the sampled households in micro watershed.

**Table 18. Status of bore wells in Hitnal micro-watershed**

Sl.No.	Particulars	LL (5)	MF (10)	SF (6)	SMF (9)	MDF (4)	All (34)
		N	N	N	N	N	N
1	De-functioning	0	0	2	0	1	3
2	Functioning	0	1	1	0	3	5

**Source of irrigation:** The data regarding the source of irrigation in Hitnal Micro watershed is presented in Table 19. The results that bore well were major source of irrigation for 14.71 per cent of the households followed by the canal and Tank was 2.94 per cent of the households.

**Table 19. Source of irrigation in Hitnal micro-watershed**

Sl.No.	Particulars	LL (5)		MF (10)		SF (6)		SMF (9)		MDF (4)		All (34)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Bore Well	0	0	1	10	1	16.67	0	0	3	75	5	14.71
2	Canal	0	0	0	0	0	0	0	0	1	25	1	2.94
3	Tank	0	0	1	10	0	0	0	0	0	0	1	2.94

**Depth of water (Avg. In meters):** The data regarding the depth of water in Hitnal Micro watershed is presented in Table 20. The results revealed that, the depth of bore well was 9.50 meter.

**Table 20. Depth of water (Avg. In meters) in Hitnal micro-watershed**

Sl.N.	Particulars	LL (5)	MF (10)	SF (6)	SMF (9)	MDF (4)	LF (0)	All (34)
		N	N	N	N	N	N	N
1	Bore Well	0	6.1	16.26	0	41.15	0	9.5
2	Tank	0	0.04	0	0	0	0	0.01

**Irrigated Area (ha):** The data regarding the irrigated area (ha) in Hitnal Micro watershed is presented in Table 21. The results indicate that, the availability of irrigation water was used for kharif crops was 9.63 ha.

**Table 21. Irrigated Area (ha) in Hitnal micro-watershed**

Sl.No.	Particulars	LL (5)	MF (10)	SF (6)	SMF (9)	MDF (4)	All (34)
1	Kharif	0	0.89	0.5	0	8.23	9.63
Total		0	0.89	0.5	0	8.23	9.63

**Table 22. Cropping pattern in Hitnal micro-watershed**

Sl.No.	Particulars	LL (5)	MF (10)	SF (6)	SMF (9)	MDF (4)	All (34)
1	Kharif - Maize	0	3.05	3.44	15.07	2.02	23.58
2	Kharif - Paddy	0	1.3	1.92	1.21	6.21	10.64
3	Kharif - Bajra	0	2.51	1.77	3.24	2.43	9.95
Total		0	6.85	7.13	19.52	10.66	44.17

**Cropping pattern:** The data regarding the cropping pattern in Hitnal Micro watershed is presented in Table 22. The results indicate that, farmers have grown maize (23.58 ha), Paddy (10.64 ha) and bajra (9.95 ha).

**Cropping intensity:** The data regarding the cropping intensity in Hitnal Micro watershed is presented in Table 23. The results indicate that, the cropping intensity was 100.00 per cent.

**Table 23. Cropping intensity (%) in Hitnal micro-watershed**

Sl.No.	Particulars	LL (5)	MF (10)	SF (6)	SMF (9)	MDF (4)	All (34)
1	Cropping Intensity	0	100	100	100	100	100

**Possession of bank account and savings:** The data regarding the possession of bank account and saving in Hitnal micro-watershed is presented in Table 24. The results indicate that, 85.29 cent of the households posses bank account and 85.29 per cent of them have savings.

**Table 24. Possession of Bank account and savings in Hitnal micro-watershed**

Sl.No.	Particulars	LL (5)		MF (10)		SF (6)		SMF (9)		MDF (4)		All (34)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Account	0	0	10	100	6	100	9	100	4	100	29	85.29
2	Savings	0	0	10	100	6	100	9	100	4	100	29	85.29

**Borrowing status:** The data regarding the borrowing status in Hitnal micro-watershed is presented in Table 25. The results indicate that, 85.29 percent of the sample farmers have borrowed credit from different sources.

**Table 25. Borrowing status in Hitnal micro-watershed**

Sl.No.	Particulars	LL (5)		MF (10)		SF (6)		SMF (9)		MDF (4)		All (34)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Credit Availed	0	0	10	100	6	100	9	100	4	100	29	85.29

**Source of credit:** The data regarding the source of credit availed by households in Hitnal micro-watershed is presented in Table 26. The result shows that, 3.45 per cent have borrowed loan from commercial banks.

**Table 26. Source of credit borrowed by households in Hitnal micro-watershed**

Sl.No.	Particulars	LL (0)		MF (10)		SF (6)		SMF (9)		MDF (4)		All (29)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Commercial Bank	0	0	0	0	0	0	1	11	0	0	1	3.45

**Avg. Credit amount:** The data regarding the avg. Credit amount in Hitnal micro-watershed is presented in Table 27. The results show that, farmers have borrowed Avg. Credit of Rs.1724.14 from different sources.

**Table 27. Avg. Credit amount in Hitnal micro-watershed**

Sl.No.	Particulars	LL (0)		MF (10)		SF (6)		SMF (9)		MDF (4)		All (29)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Average Credit	0		0		0		5555.56		0		1724.14	

**Purpose of credit borrowed (institutional Source):** The data regarding the purpose of credit borrowed - Institutional Credit in Hitnal micro-watershed is presented in Table 28.

The results indicate that, 100.00 per cent of the households have borrowed loan for agriculture.

**Table 28. Purpose of credit borrowed (institutional Source) by households in Hitnal micro-watershed**

SN	Particulars	SMF (1)		All (1)	
		N	%	N	%
1	Agriculture production	1	100	1	100

**Repayment status of household (institutional Source):** The data regarding the repayment status of credit borrowed from institutional Source by households in Hitnal micro watershed is presented in Table 29. The results indicate that, 100.00 per cent have unpaid.

**Table 29. Repayment status of household (institutional Source) in Hitnal micro-watershed**

Sl.No.	Particulars	SMF (1)		All (1)	
		N	%	N	%
1	Un paid	1	100	1	100

**Opinion regarding institutional sources of credit:** The data regarding the opinion on institutional sources of credit in Hitnal micro watershed is presented in Table 30. The results indicate that, 100.00 per cent of the households opined that credit helped to perform timely agricultural operations.

**Table 30. Opinion regarding institutional sources of credit in Hitnal micro-watershed**

Sl.No.	Particulars	SMF (1)		All (1)	
		N	%	N	%
1	Helped to perform timely agricultural operations	1	100	1	100



**Cost of Cultivation of Maize:** The data regarding the cost of cultivation (Rs/ha) of Maize in Hitnal micro watershed is presented in Table 31.a. The results indicate that, the total cost of cultivation (Rs/ha) for Maize was Rs. 22744.14. The gross income realized by the farmers was Rs. 35001.59. The net income from Maize cultivation was Rs.12257.45, thus the benefit cost ratio was found to be 1:1.50.

**Table 31(a). Cost of Cultivation of Maize in Hitnal micro-watershed**

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
<b>I</b>	<b>Cost A1</b>				
1	Hired Human Labour	Man days	20.51	5131.08	22.56
2	Bullock	Pairs/day	2.16	1187.1	5.22
3	Tractor	Hours	2.01	1504.44	6.61
4	Machinery	Hours	0	0	0
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	17.18	1952.76	8.59
6	Seed Inter Crop	Kgs.	0	0	0
7	FYM	Quintal	4.94	988	4.34
8	Fertilizer + micronutrients	Quintal	3.28	2351.43	10.34
9	Pesticides (PPC)	Kgs / liters	1.58	1953.51	8.59
10	Irrigation	Number	0	0	0
11	Repairs		0	0	0
12	Msc. Charges (Marketing costs etc)		0	0	0
13	Depreciation charges		0	372.01	1.64
14	Land revenue and Taxes		0	0	0
<b>II</b>	<b>Cost B1</b>				
16	Interest on working capital			870.68	3.83
17	<b>Cost B1 = (Cost A1 + sum of 15 and 16)</b>			16311	71.72
<b>III</b>	<b>Cost B2</b>				
18	Rental Value of Land			178.57	0.79
19	<b>Cost B2 = (Cost B1 + Rental value)</b>			16489.57	72.5
<b>IV</b>	<b>Cost C1</b>				
20	Family Human Labour		15.57	4176.92	18.36
21	<b>Cost C1 = (Cost B2 + Family Labour)</b>			20666.49	90.87
<b>V</b>	<b>Cost C2</b>				
22	Risk Premium			10	0.04
23	<b>Cost C2 = (Cost C1 + Risk Premium)</b>			20676.49	90.91
<b>VI</b>	<b>Cost C3</b>				
24	Managerial Cost			2067.65	9.09
25	<b>Cost C3 = (Cost C2 + Managerial Cost)</b>			22744.14	100
<b>VII</b>	<b>Economics of the Crop</b>				
a.	Main Product	a) Main Product (q)	28.04	33545.88	
		b) Main Crop Sales Price (Rs.)		1196.43	
	By Product	e) Main Product (q)	18.53	1455.71	
		f) Main Crop Sales Price (Rs.)		78.57	
b.	Gross Income (Rs.)			35001.59	
c.	Net Income (Rs.)			12257.45	
d.	Cost per Quintal (Rs./q.)			811.18	
e.	Benefit Cost Ratio (BC Ratio)			1:1.5	

**Cost of Cultivation of Bajra:** The data regarding the cost of cultivation (Rs/ha) of Bajra in Hitnal micro watershed is presented in Table 31.b. The results indicate that, the total cost of cultivation (Rs/ha) for Bajra was Rs. 24612.64. The gross income realized by the farmers was Rs. 25185.15. The net income from Bajra cultivation was Rs.572.51, thus the benefit cost ratio was found to be 1:1.02.

**Table 31(b). Cost of Cultivation of Bajra in Hitnal micro-watershed**

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
<b>I</b>	<b>Cost A1</b>				
1	Hired Human Labour	Man days	34.79	9005.01	36.59
2	Bullock	Pairs/day	1.73	952.06	3.87
3	Tractor	Hours	1.83	1370.13	5.57
4	Machinery	Hours	0	0	0
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	13.38	1704.68	6.93
6	Seed Inter Crop	Kgs.	0	0	0
7	FYM	Quintal	0	0	0
8	Fertilizer + micronutrients	Quintal	3.25	2299.98	9.34
9	Pesticides (PPC)	Kgs / liters	1.57	2093.63	8.51
10	Irrigation	Number	0	0	0
11	Repairs		0	0	0
12	Msc. Charges (Marketing costs etc)		0	0	0
13	Depreciation charges		0	7.07	0.03
14	Land revenue and Taxes		0	0	0
<b>II</b>	<b>Cost B1</b>				
16	Interest on working capital			732.99	2.98
17	<b>Cost B1 = (Cost A1 + sum of 15 and 16)</b>			18165.56	73.81
<b>III</b>	<b>Cost B2</b>				
18	Rental Value of Land			138.89	0.56
19	<b>Cost B2 = (Cost B1 + Rental value)</b>			18304.45	74.37
<b>IV</b>	<b>Cost C1</b>				
20	Family Human Labour		15.22	4060.68	16.5
21	<b>Cost C1 = (Cost B2 + Family Labour)</b>			22365.13	90.87
<b>V</b>	<b>Cost C2</b>				
22	Risk Premium			10	0.04
23	<b>Cost C2 = (Cost C1 + Risk Premium)</b>			22375.13	90.91
<b>VI</b>	<b>Cost C3</b>				
24	Managerial Cost			2237.51	9.09
25	<b>Cost C3 = (Cost C2 + Managerial Cost)</b>			24612.64	100
<b>VII</b>	<b>Economics of the Crop</b>				
a.	Main Product	a) Main Product (q)	20.25	23292.38	
		b) Main Crop Sales Price (Rs.)		1150	
	By Product	e) Main Product (q)	18.93	1892.77	
		f) Main Crop Sales Price (Rs.)		100	
b.	Gross Income (Rs.)			25185.15	
c.	Net Income (Rs.)			572.51	
d.	Cost per Quintal (Rs./q.)			1215.18	
e.	Benefit Cost Ratio (BC Ratio)			1:1.02	

**Cost of Cultivation of Paddy:** The data regarding the cost of cultivation (Rs/ha) of Paddy in Hitnal micro watershed is presented in Table 31.c. The results indicate, the total cost of cultivation (Rs/ha) for Paddy was Rs.46877.36. The gross income realized by the farmers was Rs. 56959.99. The net income from Paddy cultivation was Rs. 10082.63, thus the benefit cost ratio was found to be 1:1.20.

**Table 31(c). Cost of Cultivation of Paddy in Hitnal micro-watershed**

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
<b>I</b>	<b>Cost A1</b>				
1	Hired Human Labour	Man days	57.31	14810.75	31.59
2	Bullock	Pairs/day	4.84	2663.52	5.68
3	Tractor	Hours	2.67	2000.24	4.27
4	Machinery	Hours	0	0	0
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	41.56	1870.02	3.99
6	Seed Inter Crop	Kgs.	0	0	0
7	FYM	Quintal	13.93	2786.06	5.94
8	Fertilizer + micronutrients	Quintal	6.21	4512.66	9.63
9	Pesticides (PPC)	Kgs / liters	3.06	3534.09	7.54
10	Irrigation	Number	11.02	0	0
11	Repairs		0	0	0
12	Msc. Charges (Marketing costs etc)		0	0	0
13	Depreciation charges		0	1455.32	3.1
14	Land revenue and Taxes		0	0	0
<b>II</b>	<b>Cost B1</b>				
16	Interest on working capital			1525.54	3.25
17	<b>Cost B1 = (Cost A1 + sum of 15 and 16)</b>			35158.21	75
<b>III</b>	<b>Cost B2</b>				
18	Rental Value of Land			185.19	0.4
19	<b>Cost B2 = (Cost B1 + Rental value)</b>			35343.39	75.4
<b>IV</b>	<b>Cost C1</b>				
20	Family Human Labour		28.4	7262.39	15.49
21	<b>Cost C1 = (Cost B2 + Family Labour)</b>			42605.78	90.89
<b>V</b>	<b>Cost C2</b>				
22	Risk Premium			10	0.02
23	<b>Cost C2 = (Cost C1 + Risk Premium)</b>			42615.78	90.91
<b>VI</b>	<b>Cost C3</b>				
24	Managerial Cost			4261.58	9.09
25	<b>Cost C3 = (Cost C2 + Managerial Cost)</b>			46877.36	100
<b>VII</b>	<b>Economics of the Crop</b>				
a.	Main Product	a) Main Product (q)	44.71	54894.08	
		b) Main Crop Sales Price (Rs.)		1227.78	
	By Product	e) Main Product (q)	23.24	2065.91	
		f) Main Crop Sales Price (Rs.)		88.89	
b.	Gross Income (Rs.)			56959.99	
c.	Net Income (Rs.)			10082.63	
d.	Cost per Quintal (Rs./q.)			1048.47	
e.	Benefit Cost Ratio (BC Ratio)			1:1.2	

**Adequacy of fodder:** The data regarding the adequacy of fodder in Hitnal Micro watershed is presented in Table 32. The results indicate that, 23.53 per cent of the households opined that dry fodder was adequate.

**Table 32. Adequacy of fodder in Hitnal micro-watershed**

Sl.No.	Particulars	LL (5)		MF (10)		SF (6)		SMF (9)		MDF (4)		All (34)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Adequate-Dry Fodder	0	0	3	30	2	33.33	1	11.1	2	50	8	23.53

**Average annual gross income:** The data regarding the annual gross income in Hitnal Micro watershed is presented in Table 33. The results indicate that, the farmers have annual gross income of Rs. 60641.18 in micro-watershed, of which Rs. 46611.76 is from agriculture itself.

**Table 33. Average annual gross income in Hitnal micro-watershed**

Sl.No.	Particulars	LL (5)	MF (10)	SF (6)	SMF (9)	MDF (4)	All (34)
		Rs.	Rs.	Rs.	Rs.	Rs.	Rs.
1	Service/salary	0	0	0	20000	0	5294.12
2	Wage	20000	7000	0	8333.33	6250	7941.18
3	Agriculture	4000	23800	65333.3	60644.4	97250	46611.8
4	Dairy Farm	0	0	0	3000	0	794.12
Income(Rs.)		24000	30800	65333.3	91977.8	103500	60641.2

**Average annual Expenditure:** The data regarding the average annual expenditure in Hitnal Micro watershed is presented in Table 34. The results indicate that, the farmers have annual gross expenditure of Rs. 242266.67 in micro-watershed, of which Rs. 18794.12 is from agriculture itself.

**Table 34. Average annual Expenditure in Hitnal micro-watershed**

Sl.No.	Particulars	LL (5)	MF (10)	SF (6)	SMF (9)	MDF (4)	All (34)
		Rs.	Rs.	Rs.	Rs.	Rs.	Rs.
1	Service/salary	0	0	0	80000	0	2352.94
2	Wage	9500	9333.33	0	8333.33	12000	3029.41
3	Agriculture	5000	9100	23000	23500	42500	18794.1
4	Dairy Farm	0	0	0	20000	0	588.24
Total		14500	18433.3	23000	131833	54500	242267

**Horticulture species grown:** The data regarding horticulture species grown in Hitnal Micro watershed is presented in Table 35. The results indicate that, the total number of horticultural trees grown (both field and backyard) by the sampled households were Mango (1).

**Table 35. Horticulture species grown in Hitnal micro-watershed**

Sl.No.	Particulars	LL (5)		MF (10)		SF (6)		SMF (9)		MDF (4)		All (34)	
		F	B	F	B	F	B	F	B	F	B	F	B
1	Mango	0	0	0	0	0	0	0	0	1	0	1	0

\*F= Field B=Back Yard

**Forest species grown:** The data regarding forest species grown in Hitnal Micro watershed is presented in Table 36. The results indicate that, households have planted 4 neem trees together in both field and backyard.

**Table 36. Forest species grown in Hitnal micro-watershed**

Sl.No.	Particulars	LL (5)		MF (10)		SF (6)		SMF (9)		MDF (4)		All (34)	
		F	B	F	B	F	B	F	B	F	B	F	B
1	Neem	0	0	0	0	0	0	4	0	0	0	4	0

\*F= Field B=Back Yard

**Marketing of agricultural produce:** The data regarding marketing of the agricultural produce in Hitnal Micro watershed is presented in Table 37. The results indicated that, 86.63 percent of output of Bajra was sold in the market with average price of Rs. 1150.00; 86.03 percent of output of maize was sold in the market with average price of Rs. 1288.46 and 72.10 percent of output of paddy was sold in the market with average price of Rs. 1227.78;

**Table 37. Marketing of agricultural produce in Hitnal micro-watershed**

Sl.No	Crops	Output obtained (q)	Output retained (q)	Output sold (q)	Output sold (%)	Avg. Price obtained (Rs/q)
1	Bajra	187	25	162	87	1150
2	Maize	630	88	542	86	1288
3	Paddy	466	130	336	72	1228

**Marketing channels used for sale of agricultural produce:** The data regarding marketing channels used for sale of agricultural produce in Hitnal Micro watershed is presented in Table 38. The results indicated that, 11.76 cent of the households have sold agricultural produce to the local/village merchants and 70.59 per cent of regulated market.

**Table 38. Marketing channels used for sale of agricultural produce in Hitnal micro-watershed**

Sl.No.	Particulars	LL (5)		MF (10)		SF (6)		SMF (9)		MDF (4)		All (34)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Local/village Merchant	0	0	0	0	1	16.7	1	11.1	2	50	4	11.76
2	Regulated Market	0	0	9	90	5	83.3	8	88.9	2	50	24	70.59

**Mode of transport of agricultural produce:** The data regarding mode of transport of agricultural produce in Hitnal Micro watershed is presented in Table 39. The results indicated that, 82.35 cent of the households have used tractor.

**Table 39. Mode of transport of agricultural produce in Hitnal micro-watershed**

Sl.No.	Particulars	LL (5)		MF (10)		SF (6)		SMF (9)		MDF (4)		All (34)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Tractor	0	0	9	90	6	100	9	100	4	100	28	82.35

**Incidence of soil and water erosion problems:** The data regarding incidence of incidence of soil and water erosion problems in Hitnal Micro watershed is presented in Table 40. The results indicate that, 85.29 per cent of the households have experienced soil and water erosion problems.

**Table 40. Incidence of soil and water erosion problems in Hitnal micro-watershed**

Sl.No.	Particulars	LL (5)		MF (10)		SF (6)		SMF (9)		MDF (4)		All (34)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Soil and water erosion problems in the farm	0	0	10	100	6	100	9	100	4	100	29	85.29

**Interest towards soil testing:** The data regarding Interest shown towards soil testing in Hitnal Micro watershed is presented in Table 41. The results indicated that, 82.35 per cent of the households were interested towards soil testing.

**Table 41. Interest regarding soil testing in Hitnal micro-watershed**

Sl.No.	Particulars	LL (5)		MF (10)		SF (6)		SMF (9)		MDF (4)		All (34)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Interest in soil test	0	0	9	90	6	100	9	100	4	100	28	82.35

**Usage pattern of fuel for domestic use:** The data on usage pattern of fuel for domestic use in Hitnal Micro watershed is presented in Table 42. The results indicated that, firewood was the major source of fuel for domestic use for 100.00 per cent of the households.

**Table 42. Usage pattern of fuel for domestic use in Hitnal micro-watershed**

Sl.No.	Particulars	LL (5)		MF (10)		SF (6)		SMF (9)		MDF (4)		All (34)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Fire Wood	5	100	10	100	6	100	9	100	4	100	34	100

**Source of drinking water:** The data on source of drinking water in Hitnal Micro watershed is presented in Table 43. The results indicated that, piped waters supply was the major source for drinking water for 100.00 per cent of the households.

**Table 43. Source of drinking water in Hitnal micro-watershed**

Sl.No.	Particulars	LL (5)		MF (10)		SF (6)		SMF (9)		MDF (4)		All (34)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Piped supply	5	100	10	100	6	100	9	100	4	100	34	100

**Source of light:** The data on source of light in Hitnal Micro watershed is presented in Table 44. The results indicated that, electricity was the major source of light for 100.00 per cent of the households.

**Table 44. Source of light in Hitnal micro-watershed**

Sl.No.	Particulars	LL (5)		MF (10)		SF (6)		SMF (9)		MDF (4)		All (34)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Electricity	5	100	10	100	6	100	9	100	4	100	34	100

**Existence of sanitary toilet facility:** The data on availability of toilet facility in Hitnal Micro watershed is presented in Table 45. The results indicated that, 100.00 per cent of the households possess toilets.

**Table 45. Existence of sanitary toilet facility in Hitnal micro-watershed**

Sl.No.	Particulars	LL (5)		MF (10)		SF (6)		SMF (9)		MDF (4)		All (34)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Sanitary toilet facility	5	100	10	100	6	100	9	100	4	100	34	100

**Possession of PDS card:** The data regarding possession of PDS card in Hitnal Micro watershed is presented in Table 46. The results indicated that, 100.00 per cent of the households possessed BPL card.

**Table 46. Possession of PDS card in Hitnal micro-watershed**

Sl.No.	Particulars	LL (5)		MF (10)		SF (6)		SMF (9)		MDF (4)		All (34)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	BPL	5	100	10	100	6	100	9	100	4	100	34	100

**Participation in NREGA programme:** The data regarding Participation in NREGA programme in Hitnal Micro watershed is presented in Table 47. The results indicated that, only 2.94 percent of the participate have participated in NREGA programme.

**Table 47. Participation in NREGA programme in Hitnal micro-watershed**

Sl.No.	Particulars	LL (5)		MF (10)		SF (6)		SMF (9)		MDF (4)		All (34)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Participation in NREGA programme	0	0	1	10	0	0	0	0	0	0	1	2.94

**Adequacy of food items:** The data regarding adequacy of food items in Hitnal Micro watershed is presented in Table 48. The results indicated that, the extent of adequacy of food items for cereals, pulses, Oilseeds and vegetables were 91.18, 88.24, 17.65, 11.76 per cent respectively, similarly for Fruits (17.65%), milk (14.71%), Egg (5.88%), and Meat (5.88%).

**Table 48. Adequacy of food items in Hitnal micro-watershed**

Sl.No.	Particulars	LL (5)		MF (10)		SF (6)		SMF (9)		MDF (4)		All (34)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Cereals	3	60	10	100	6	100	8	88.9	4	100	31	91.18
2	Pulses	3	60	10	100	5	83.33	8	88.9	4	100	30	88.24
3	Oilseed	1	20	3	30	1	16.67	0	0	1	25	6	17.65
4	Vegetables	0	0	2	20	0	0	1	11.1	1	25	4	11.76
5	Fruits	0	0	3	30	1	16.67	1	11.1	1	25	6	17.65
6	Milk	0	0	2	20	0	0	2	22.2	1	25	5	14.71
7	Egg	0	0	0	0	1	16.67	0	0	1	25	2	5.88
8	Meat	0	0	0	0	0	0	1	11.1	1	25	2	5.88

**Table 49. Inadequacy of food items in Hitnal micro-watershed**

Sl.No.	Particulars	LL (5)		MF (10)		SF (6)		SMF (9)		MDF (4)		All (34)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Cereals	2	40	0	0	0	0	0	0	0	0	2	5.88
2	Pulses	2	40	0	0	1	16.67	0	0	0	0	3	8.82
3	Oilseed	4	80	7	70	5	83.33	9	100	3	75	28	82.35
4	Vegetables	5	100	6	60	6	100	8	88.9	3	75	28	82.35
5	Fruits	5	100	6	60	5	83.33	7	77.8	2	50	25	73.53
6	Milk	4	80	8	80	5	83.33	5	55.6	2	50	24	70.59
7	Egg	5	100	9	90	5	83.33	9	100	4	100	32	94.12
8	Meat	5	100	10	100	6	100	8	88.9	3	75	32	94.12

**Inadequacy of food items:** The data regarding in adequacy of food items in Hitnal Micro watershed is presented in Table 49. The results indicated that, the extent of in adequacy of food items for cereals, pulses, Oilseeds and vegetables were 5.88, 8.82, 82.35, 82.35 and 94.12 per cent respectively, similarly for fruits (73.53%), milk (70.59%), egg (94.12%) and meat (94.12%).

**Response on market surplus of food items:** The data regarding adequacy of food items in Hitnal Micro watershed is presented in Table 50. The results indicated that, the extent of adequacy of food items for cereals and pulses were 2.94 and 2.94per cent respectively.

**Table 50. Response on market surplus of food items in Hitnal micro-watershed**

Sl.No.	Particulars	LL (5)		MF (10)		SF (6)		SMF (9)		MDF (4)		All (34)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Cereals	0	0	0	0	0	0	1	11.1	0	0	1	2.94
2	Pulses	0	0	0	0	0	0	1	11.1	0	0	1	2.94

**Farming constraints:** The data regarding farming constraints experienced by households in Hitnal Micro watershed is presented in Table 51. The results indicated that, lower fertility status of the soil was the constraint experienced by (85.29 %) per cent of the households, wild animal menace on farm field (76.47%), frequent incidence of pest and diseases (55.88%), inadequacy of irrigation water (20.59%), high cost of fertilizers and plant protection chemicals (38.24%), high rate of interest on credit (8.82%), low price for the agricultural commodities (14.71 %), lack of marketing facilities in the area (11.76%), inadequate extension services (14.71 %), lack of transport for safe transport of the agricultural produce to the market (38.24%), less rainfall (32.35%), source of agri-technology information (Newspaper/Tv/Mobile) (38.24%).

**Table 51. Farming constraints experienced in Hitnal micro-watershed**

SN	Particulars	LL (5)		MF (10)		SF (6)		SMF (9)		MDF (4)		All (34)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Lower fertility status of the soil	0	0	10	100	6	100	9	100	4	100	29	85.29
2	Wild animal menace on farm field	0	0	8	80	6	100	9	100	3	75	26	76.47
3	Frequent incidence of pest and diseases	0	0	7	70	4	66.67	6	66.67	2	50	19	55.88
4	Inadequacy of irrigation water	0	0	2	20	1	16.67	1	11.11	3	75	7	20.59
5	High cost of Fertilizers and plant protection chemicals	0	0	5	50	3	50	4	44.44	1	25	13	38.24
6	High rate of interest on credit	0	0	1	10	1	16.67	1	11.11	0	0	3	8.82
7	Low price for the agricultural commodities	0	0	2	20	1	16.67	0	0	2	50	5	14.71
8	Lack of marketing facilities in the area	0	0	0	0	1	16.67	3	33.33	0	0	4	11.76
9	Inadequate extension services	0	0	1	10	1	16.67	2	22.22	1	25	5	14.71
10	Lack of transport for safe transport of the Agril produce to the market.	0	0	8	80	1	16.67	1	11.11	3	75	13	38.24
11	Less rainfall	0	0	4	40	2	33.33	5	55.56	0	0	11	32.35
12	Source of Agri-technology information	0	0	4	40	2	33.33	5	55.56	2	50	13	38.24



## SUMMARY AND IMPLICATIONS

In order to assess the socio-economic condition of the farmers in the watershed 34 households located in the micro watershed were interviewed for the survey. The study was conducted in Hitnal micro-watershed (Shahpura sub-watershed, Koppala taluk & District) is located at North latitude  $15^{\circ} 20' 35.836''$  and  $15^{\circ} 19' 2.884''$  and East longitude  $76^{\circ} 19' 29.484''$  and  $76^{\circ} 17' 1.088''$  covering an area of about 351.12 ha bounded by under Hitnal and Chikkasanakandi Villages.

Socio-economic analysis indicated that, out of the total sample of 34 respondents, 10 (29.41%) were marginal, 6 (17.65%) were small and 9 (26.47%) were semi medium, 4 (11.76%) were medium farmers. The population characteristics of households indicated that, there were 84 (56.00%) men and 66 (44.00%) were women. Majority of the respondents (48.00%) were in the age group of 35-60 years. Education level of the sample households indicated that, majority there were 47.33 per cent of illiterates, 18.67 per cent of them had primary school education, 8.67 per cent middle school education, and 7.33 per cent high school education, 11.33 per cent of them had PUC education, 0.67 per cent of them had Diploma, 2.00 per cent attained graduation. About, 91.18 per cent of household heads practicing agriculture and 8.82 per cent of the household heads were engaged as agricultural labourers. Agriculture was the major occupation for 21.33 per cent of the household members.

In the study area, 88.24 per cent of the households possess katcha house and 8.82 per cent possess pucca house. The durable assets owned by the households showed that, 70.59 per cent possess TV, 32.35 per cent possess mixer grinder and 85.29 per cent possess mobile phones. Farm implements owned by the households indicated that, 14.71 per cent of the households possess plough and only 2.94 per cent sprayer. Regarding livestock possession by the households, 17.65 per cent possess local cow and 2.94 per cent possess buffalo respectively.

The average labour availability in the study area showed that, own men and women labour availability in the micro watershed was 21.71 each, while the hired labour (men) availability was 1.76. Further, 88.24 per cent of the households opined that hired labour was inadequate during the agricultural season.

Out of the total land holding of the sample respondents (45.93 ha), 67.15 per cent of the area is under dry condition and the remaining 32.85 per cent area is irrigated land. There were 5.00 bore wells and 3.00 dry bore wells among the sampled households. Bore well was the major source of irrigation for 14.71 per cent of the households. The major crops grown by sample farmers are Maize, Bajra and Paddy and cropping intensity was recorded as 100.00 per cent.

The sample households possessed 85.29 per cent bank account and 85.29 per cent of them have savings in the account. About 85.29 per cent of the respondents borrowed credit from various sources. Among the credit borrowed by households, 3.45 per cent have borrowed loan from commercial banks. Majority of the respondents (100.00 %) have borrowed loan for agriculture purpose. Regarding the opinion on institutional sources of credit, 100.00 per cent of the households opined that credit helped to perform timely agricultural operations.

The per hectare cost of cultivation for Maize, Bajra and Paddy was Rs.22744.14 , 24612.64 and 46877.36 with benefit cost ratio of 1:1.50, 1: 1.02 and 1: 1.20 respectively.

Further, 23.53 per cent of the households opined that dry fodder was adequate.

The average annual gross income of the farmers was Rs. 60641.18 in micro-watershed, of which Rs. 46611.76 comes from agriculture.

Sampled households have grown mango trees in the fields, Further, Cashew, Lemon, Coconut, Guava, Jamun trees were also planted in the farm fields. None of the households shown interest to cultivate horticultural crops.

Regarding marketing channels, 11.76 per cent of the households have sold agricultural produce to the local/village merchants, while, 70.59 per cent have sold by Agents/Traders. Further, 82.35 per cent of the households have used tractor for the transport of agriculture commodity.

Majority of the farmers (85.29 %) have experienced soil and water erosion problems in the watershed and 82.35 per cent of the households were interested towards soil testing.

Firewood connection was the major source of fuel for domestic use for 100.00 per cent of the households. Piped supply was the major source for drinking water for 100.00 per cent of the households. Electricity was the major source of light for 100.00 per cent of the households. In the study area, 100.00 per cent of the households possess toilet facility. Regarding possession of PDS card, 100.00 per cent of the households possessed BPL card. Cereals (91.18%), pulses (88.24%), oilseeds (17.65%) were adequate for consumption.

Farming constraints experienced by households in the micro watersheds were lower fertility status of the soil (85.29%) wild animal menace on farm field (76.47%), frequent incidence of pest and diseases (55.88%), inadequacy of irrigation water (20.59%), high cost of fertilizers and plant protection chemicals (38.24%), high rate of interest on credit (8.82%), low price for the agricultural commodities (14.71%), lack of marketing facilities in the area (11.76%), inadequate extension services (14.71%), lack of transport for safe transport of the agricultural produce to the market (38.24%), Less

rainfall (32.35%) and Source of Agri-technology information (Newspaper/TV/Mobile) (38.24%).

### **Implications of the survey**

- ✓ Result indicated that, there were 47.33 per cent were illiterate hence, extension methodologies such as demonstration, street play, drama, video shows will be effective in dissemination of the technologies in the micro watershed.
- ✓ The data indicate that, 88.24 per cent of the households possess katcha house. Hence, the development department while implementing the watershed plan should focus on agriculture to enhance the productivity of major crops in the area to increase the income of the farmers.
- ✓ Results indicated that the local institutional participation of the household members in the micro watershed is minimal hence, activities like membership campaign, awareness creation about the benefits of membership in local institutions and strengths of organized groups must be conveyed.
- ✓ Majority of the households in the watershed have experience in use of mobile phones, and television hence, these mass media can be effectively utilized for transfer of technology as well as for information dissemination.
- ✓ The farm machinery/implement possession in the micro watershed was found to be minimum the reasons may lack of knowledge or lack of financial ability which can be addressed through training on use of different farm implements, providing information on different sources of finance for purchase of farm implements.
- ✓ The possession of livestock such as crossbred cow found is less hence, farmers must be made aware of the benefits of crossbred cow in increased milk production.
- ✓ The possession of livestock such as sheep, goat and poultry was found to be low hence, farmers may be informed the role of subsidiary enterprises in enhancing the income and information on financial support for subsidiary activities.
- ✓ The data indicate that, job/work was the reason for all the migrants hence, farmers may be trained on profitable agriculture or self employment such as animal husbandry, plate making, sheep rearing, goat rearing, rabbit rearing with suitable information on sources of financial support.
- ✓ The results indicate that there was a change in quality of life due to migration hence, the developmental departments should take actions to arrest migration and to improve the quality of the life in rural areas.
- ✓ Households possess 30.84ha (67.15 %) of dry land and 15.09ha (32.85 %) of irrigated land hence, the availability of the dryland agricultural technologies such as short duration crops, high yielding drought resistance crop varieties, drip irrigation technology and subsidy information will be helpful for the farmers to enhance the productivity of land and as well as farmers income.

- ✓ Few of the bore well in micro watershed found non functional hence, farmers may be trained on possibility of bore well rejuvenation.
- ✓ Bore well was major source of irrigation for 14.71 per cent of the households. hence, in order to increase the area under irrigation as well as to increase the water use efficiency farmers may trained on drip irrigation and provide the information on subsidy for drip irrigation equipment's along with the information on different agencies which provides the financial assistance for drip irrigation.
- ✓ Farmers have grown 1 mango trees in the fields. Hence, production technologies related to these crops can be made available to the farmers for better adoption.
- ✓ The cropping intensity in the micro watershed was found to be (100.00 %) hence, care must be taken by the implementing agency to bring uncultivated land into cultivation through suitable measures.
- ✓ Many of the household members have borrowed loan from cooperative banks which has higher rate of interest hence, farmers may be sensitized on the different sources of credit with lesser interest rate such SHGs etc.
- ✓ The results indicated the non availability of both green and dry fodder throughout the year hence, fodder development activities can be taken up in the micro watershed.
- ✓ The average annual gross income of the households Rs.46611.76 from agriculture and Rs. 7941.18 from wages and. Agriculture was found to be the major source of income for households hence; the development activities should focus on productivity enhancement, marketing arrangements and agricultural technology dissemination to have a direct impact on the farmers.
- ✓ The cultivation of forest species is found minimal hence, information and production technology related to agro-forestry and integrated farming system.
- ✓ The data indicated that, 85.29 per cent of the households have experienced soil and water erosion problems. Hence, those farmers who reported the soil and water erosion problems may be given attention while implementation of the watershed development plan.
- ✓ The data indicated that, 82.35 per cent of the households have interest in soil testing hence, farmers must be provided with the information on various institutions which are involved in soil testing for the benefit of the farmers.
- ✓ Except summer ploughing the adoption of other soil and water conservation structures is minimum hence, the farmers in the micro watershed should be sensitized on the use of different conservation structures for soil water conservation.
- ✓ Cereals and pulses found be adequate for per cent of the households respectively hence, farm households and the farm women must be trained on importance of balanced nutrition and role of vegetable, milk, egg, meat in balanced diet.
- ✓ Lower fertility status of the soil (85.29%), wild animal menace on farm field (76.47%), frequent incidence of pest and diseases (55.88%), high cost of fertilizers and plant protection chemicals (38.24%), high rate of interest on credit (8.82%), low

price for the agricultural commodities (14.71%), lack of marketing facilities in the area (11.76%), inadequate extension services (14.71%), lack of transport for safe transport of the agricultural produce to the market (38.24%) were the major farming constraints experienced hence, these constraints must be addressed immediately for the welfare of the farmers. Awareness to be created among the farmers to approach nearest KVKs/RSKs and other developmental departments for technical and for subsidized inputs and utilize the well established regulated markets, approaching the contract firms, direct markets to avoid the involvement of middlemen.