



हर कदम, हर डगर  
किसानों का हमसफर  
भारतीय कृषि अनुसंधान परिषद

*AgriSearch with a human touch*

**LAND RESOURCE INVENTORY AND SOCIO-ECONOMIC STATUS OF  
FARM HOUSEHOLDS FOR WATERSHED PLANNING AND  
DEVELOPMENT**

**HANDRAHALU-2 (4D4A1V2c) MICRO WATERSHED**

**Alavandi Hobli, Koppal Taluk and District, Karnataka**

**Karnataka Watershed Development Project – II**

**SUJALA – III**

**World Bank funded Project**



**ICAR – NATIONAL BUREAU OF SOIL SURVEY AND LAND USE PLANNING**



ICAR - NBSS & LUP



**WATERSHED DEVELOPMENT DEPARTMENT  
GOVT. OF KARNATAKA, BANGALORE**



## **About ICAR - NBSS&LUP**

The ICAR-National Bureau of Soil Survey and Land Use Planning (ICAR-NBSS&LUP), Nagpur, a premier Institute of the Indian Council of Agricultural Research (ICAR), was set up during 1976 with the objective to prepare soil resource maps at national, state and district levels and to provide research inputs in soil resource mapping and its applications, land evaluation, land use planning, land resource management, and database management using GIS for optimising land use on different kinds of soils in the country.

The Bureau has been engaged in carrying out soil resource survey, agro-ecological and soil degradation mapping at the country, state and district levels for qualitative assessment and monitoring the soil health towards viable land use planning. The research activities have resulted in identifying the soil potentials and problems, and the various applications of the soil surveys with the ultimate objective of sustainable agricultural development. The Bureau has the mandate to correlate and classify soils of the country and maintain a National Register of all the established soil series. The Institute is also imparting in-service training to staff of the soil survey agencies in the area of soil survey, land evaluation and soil survey interpretations for land use planning. The Bureau in collaboration with Panjabrao Krishi Vidyapeeth, Akola is running post-graduate teaching and research programme in land resource management, leading to M.Sc. and Ph.D. degrees.

**Citation:** Rajendra Hegde, Ramesh Kumar, S.C., K.V. Niranjana, S. Srinivas, M.Lalitha, B.A. Dhanorkar, R.S. Reddy and S.K. Singh (2019). "Land Resource Inventory and Socio-Economic Status of Farm Households for Watershed Planning and Development of Handrahalu-2 (4D4A1V2c) Microwatershed, Alavandi Hobli, Koppal Taluk and District, Karnataka", ICAR-NBSS&LUP Sujala MWS Publ.382, ICAR – NBSS & LUP, RC, Bangalore. p.139 & 31.

### **TO OBTAIN COPIES,**

**Please write to:**

**Director, ICAR - NBSS & LUP,**

Amaravati Road, NAGPUR - 440 033, India

Phone : (0712) 2500386, 2500664, 2500545 (O)

Telefax : 0712-2522534

E-Mail : [director@nbsslup.ernet.in](mailto:director@nbsslup.ernet.in)

Website URL : [nbsslup.in](http://nbsslup.in)

Or

**Head, Regional Centre, ICAR - NBSS&LUP, Hebbal, Bangalore - 560 024**

Phone : (080) 23412242, 23510350 (O)

Telefax : 080-23510350

E-Mail : [nbssrcb@gmail.com](mailto:nbssrcb@gmail.com)

ICAR-NBSS&LUP Sujala MWS Publ.382



हर कदम, हर डगर  
किसानों का हमसफर  
भारतीय कृषि अनुसंधान परिषद

*Agri*search with a human touch

# LAND RESOURCE INVENTORY AND SOCIO-ECONOMIC STATUS OF FARM HOUSEHOLDS FOR WATERSHED PLANNING AND DEVELOPMENT

## HANDRAHALU-2 (4D4A1V2c) MICRO WATERSHED

Alavandi Hobli, Koppal Taluk and District, Karnataka

### Karnataka Watershed Development Project – II

### Sujala-III

World Bank funded Project



ICAR – NATIONAL BUREAU OF SOIL SURVEY AND LAND USE  
PLANNING



ICAR - NBSS & LUP



WATERSHED DEVELOPMENT DEPARTMENT, GOVT. OF  
KARNATAKA, BANGALORE





## PREFACE

In Karnataka, as in other Indian States, the livelihoods of rural people are intertwined with farming pursuits. The challenges in agriculture are seriously threatening the livelihood of a large number of farmers as they have been practicing farming in contextual factors beyond their control. Climatic factors are the most important ones and have become much more significant in recent times due to rapid climate changes induced by intensive anthropogenic activities affecting our ecosystem in multiple ways. Climate change has become the reality, it is happening and efforts to evolve and demonstrate climate resilient technologies have become essential. Due to the already over stressed scenario of agrarian sector, the climate change is resulting in manifold increase in the complexities, pushing the rural mass to face more and more unpredictable situations. The rising temperatures and unpredictable rainfall patterns are going to test seriously the informed decisions farmers have to make in order to survive in farming and sustain their livelihood.

It is generally recognized that impacts of climate change shall not be uniform across the globe. It is said that impact of climate change is more severe in South Asia. Based on the analysis of meteorological data, it is predicted that in India, there will be upward trend in mean temperature, downward trend in relative humidity, annual rainfall and number of wet days in a year. Also, in general, phenomena like erratic monsoon, spread of tropical diseases, rise in sea levels, changes in availability of fresh water, frequent floods, droughts, heat waves, storms and hurricanes are predicted. Each one of these adverse situations are already being experienced in various parts of India and also at the global level. Decline in agricultural productivity of small and marginal farmers becoming more vulnerable is already witnessed.

In Karnataka, more than 60 per cent of the population live in rural areas and depend on agriculture and allied activities for their livelihood. Though the state has achieved significant progress in increasing the yield of many crops, there is tremendous pressure on the land resources due to the growing and competing demands of various land uses. This is reflected in the alarming rate of land degradation observed. Already more than 50 per cent of the area is affected by various forms of degradation. If this trend continues, the sustainability of the fragile ecosystem will be badly affected. The adverse effects of change in the climatic factors are putting additional stress on the land resources and the farmers dependent on this.

The natural resources (land, water and vegetation) of the state need adequate and constant care and management, backed by site-specific technological interventions and investments particularly by the government. Detailed database pertaining to the nature of

the land resources, their constraints, inherent potentials and suitability for various land based rural enterprises, crops and other uses is a prerequisite for preparing location-specific action plans, which are in tune with the inherent capability of the resources. Any effort to evolve climate resilient technologies has to be based on the baseline scientific database. Then only one can expect effective implementation of climate resilient technologies, monitor the progress, make essential review of the strategy, and finally evaluate the effectiveness of the implemented programs. The information available at present on the land resources of the state are of general nature and useful only for general purpose planning. Since the need of the hour is to have site-specific information suitable for farm level planning and detailed characterization and delineation of the existing land resources of an area into similar management units is the only option.

ICAR-NBSS&LUP, Regional Centre, Bangalore has taken up a project sponsored by the Karnataka Watershed Development Project-II, (Sujala-III), Government of Karnataka funded by the World Bank under Component -1 Land Resource Inventory. This study was taken up to demonstrate the utility of such a database in reviewing, monitoring and evaluating all the land based watershed development programs on a scientific footing. To meet the requirements of various land use planners at grassroots level, the present study on “Land Resource Inventory and Socio-Economic Status of Farm Households for Watershed Planning and Development of for Handrahalu-2 microwatershed in Koppal Taluk, Koppal District, Karnataka” for integrated development was taken up in collaboration with the State Agricultural Universities, IISC, KRSRAC, KSNDMC as Consortia partners. The project provides detailed land resource information at cadastral level (1:7920 scale) for all the plots and socio-economic status of farm households covering thirty per cent farmers randomly selected representing landed and landless class of farmers in the micro-watershed. The project report with the accompanying maps for the microwatershed will provide required detailed database for evolving effective land use plan, alternative land use options and conservation plans for the planners, administrators, agricultural extension personnel, KVK officials, developmental departments and other land users to manage the land resources in a sustainable manner.

It is hoped that this database will be useful to the planners, administrators and developmental agencies working in the area in not only for formulating location specific developmental schemes but also for their effective monitoring at the village/watershed level.

Nagpur

Date:26-09-2019

**S.K. SINGH**

Director, ICAR - NBSS&LUP Nagpur

## Contributors

<b>Dr. Rajendra Hegde</b> Principal Scientist, Head & Project Leader, Sujala-III Project ICAR-NBSS&LUP, Regional Centre, Bangalore	<b>Dr. S.K.Singh</b> Director, ICAR-NBSS&LUP Coordinator, Sujala-III Project Nagpur
<b>Soil Survey, Mapping &amp; Report Preparation</b>	
Dr. K.V. Niranjana	Sh. R.S. Reddy
Dr. B.A. Dhanorkar	Dr. Gopali Bardhan
	Smt. Chaitra, S.P.
	Sh. Somashekar T.N
	Ms. Arpitha, G.M.
	Dr. Mahendra kumar M.B
	Sh. Tirupati Meti
<b>Field Work</b>	
Sh. C. Bache Gowda	Sh. Mayur Patil
Sh. Somashekar	Sh. Arun Kumar, S.
Sh. M. Jayaramaiah	Sh. Sunil Raj
	Sh. Yogesh Kumar, B.
	Sh. Vikas, N.K.
	Sh. Arun Kumar, S.G.
	Sh. Umesh Jadiyahappa Madolli
	Sh. Praveen Kumar P. Achalkar
	Sh. Veerabhadraswamy
	Sh. Vinay
	Sh. Shankarappa, K.
	Sh. Lankesh, R.S.
	Sh. Appanna B. Hattigoudar
	Sh. Maharudra
<b>GIS Work</b>	
Dr. S.Srinivas	Sh. A.G. Devendra Prasad
Sh. D. H.Venkatesh	Sh. Abhijith Sastry, N.S.
Smt. K.Sujatha	Smt. Shyla, B.
Smt. K. V. Archana	Smt. Swetha ,K.
Sh. N. Maddileti	Ms. Vidya, P.C.
	Sh. Deepak, M.J.
	Smt. K.Karunya Lakshmi
	Ms. Seema, K.V.

<b>Laboratory Analysis</b>	
Dr. M. Lalitha	Sh. Vindhya, N.G.
Smt. Arti Koyal	Ms. P. Pavanakumari, P.
Smt. Parvathy, S.	Ms. Rashmi, N.
	Ms. Leelavathy, K.U.
	Smt. Usha Kiran, G.
<b>Socio-Economic Analysis</b>	
Dr. S.C. Ramesh Kumar	Sh. M.K. Prakashanaik
	Ms. Karuna V. Kulkarni
	Mrs. Sowmya A.N
	Sh. Vinod R
	Sh. Basavaraja
	Sh. Vijay Kumar Lamani
	Ms. Sowmya K.B
	Mrs. Prathibha, D.G
	Sh. Rajendra,D
<b>Soil &amp; Water Conservation</b>	
Sh. Sunil P. Maske	
<b>Watershed Development Department, GOK, Bangalore</b>	
Sh. Rajeev Ranjan IFS Project Director & Commissioner, WDD	Dr. A. Natarajan NRM Consultant, Sujala-III Project
Dr. S.D. Pathak IFS Executive Director & Chief Conservator of Forests, WDD	



# **PART-A**

## **LAND RESOURCE INVENTORY**



## Contents

Preface		
Contributors		
Executive Summary		
Chapter 1	Introduction	1
Chapter 2	Geographical Setting	3
2.1	Location and Extent	3
2.2	Geology	3
2.3	Physiography	4
2.4	Drainage	5
2.5	Climate	5
2.6	Natural Vegetation	6
2.7	Land Utilization	7
Chapter 3	Survey Methodology	11
3.1	Base maps	11
3.2	Image Interpretation for Physiography	11
3.3	Field Investigation	14
3.4	Soil mapping	16
3.5	Laboratory Characterization	17
3.6	Land management units	19
Chapter 4	The Soils	23
4.1	Soils of Granite gneiss	23
4.2	Alluvial Landscape and Lowland	24
Chapter 5	Interpretation for Land Resource Management	41
5.1	Land Capability Classification	41
5.2	Soil Depth	43
5.3	Surface Soil Texture	44
5.4	Soil Gravelliness	45
5.5	Available Water Capacity	46
5.6	Soil Slope	47
5.7	Soil Erosion	48
Chapter 6	Fertility Status	49
6.1	Soil Reaction (pH)	51
6.2	Electrical Conductivity (EC)	51
6.3	Organic Carbon (OC)	51
6.4	Available Phosphorus	54
6.5	Available Potassium	54
6.6	Available Sulphur	54
6.7	Available Boron	54
6.8	Available Iron	54
6.9	Available Manganese	54
6.10	Available Copper	54
6.11	Available Zinc	58
Chapter 7	Land Suitability for Major Crops	59

7.1	Land suitability for Sorghum	59
7.2	Land suitability for Maize	60
7.3	Land suitability for Bajra	61
7.4	Land suitability for Groundnut	62
7.5	Land suitability for Sunflower	63
7.6	Land suitability for Red gram	64
7.7	Land suitability for Bengalgram	65
7.8	Land suitability for Cotton	66
7.9	Land suitability for Chilli	67
7.10	Land suitability for Tomato	68
7.11	Land suitability for Brinjal	69
7.12	Land suitability for onion	70
7.13	Land suitability for Bhindi	71
7.14	Land suitability for Drumstick	72
7.15	Land suitability for Mango	73
7.16	Land suitability for Guava	74
7.17	Land suitability for Sapota	75
7.18	Land suitability for Pomegranate	76
7.19	Land suitability for Musambi	77
7.20	Land suitability for Lime	78
7.21	Land suitability for Amla	79
7.22	Land suitability for Cashew	80
7.23	Land suitability for Jackfruit	81
7.24	Land Suitability for Jamun	82
7.25	Land Suitability for Custard apple	83
7.26	Land Suitability for Tamarind	84
7.27	Land Suitability for Mulberry	85
7.28	Land Suitability for Marigold	86
7.29	Land suitability for Chrysanthemum	87
7.30	Land suitability for Jasmine	88
7.31	Land suitability for Crossandra	89
7.32	Land management units	123
7.32	Proposed Crop Plan	124
Chapter 8	Soil Health Management	127
Chapter 9	Soil and Water conservation Treatment Plan	131
9.1	Treatment Plan	132
9.2	Recommended Soil and Water Conservation measures	135
9.3	Greening of microwatershed	136
	References	139
	Appendix I	I-VIII
	Appendix II	IX-XV
	Appendix III	XVII-XXII

## LIST OF TABLES

2.1	Mean Monthly Rainfall, PET, 1/2 PET at Koppal Taluk and District	5
2.2	Land Utilization in Koppal District	7
3.1	Differentiating Characteristics used for Identifying Soil Series	16
3.2	Soil map unit description of Handrahalu-2 microwatershed	17
4.1	Physical and chemical characteristics of soil series identified in Handrahalu-2 microwatershed	31
7.1	Soil-Site Characteristics of Handrahalu-2 microwatershed	91
7.2	Land suitability for Sorghum	92
7.3	Land suitability for Maize	93
7.4	Land suitability for Bajra	94
7.5	Land suitability for Groundnut	95
7.6	Land suitability for Sunflower	96
7.7	Land suitability for Red gram	97
7.8	Land suitability for Bengalgram	98
7.9	Land suitability for Cotton	99
7.10	Land suitability for Chilli	100
7.11	Land suitability for Tomato	101
7.12	Land suitability for Brinjal	102
7.13	Land suitability for onion	103
7.14	Land suitability for Bhindi	104
7.15	Land suitability for Drumstick	105
7.16	Land suitability for Mango	106
7.17	Land suitability for Guava	107
7.18	Land suitability for Sapota	108
7.19	Land suitability for Pomegranate	109
7.20	Land suitability for Musambi	110
7.21	Land suitability for Lime	111
7.22	Land suitability for Amla	112
7.23	Land suitability for Cashew	113

7.24	Land suitability for Jackfruit	114
7.25	Land Suitability for Jamun	115
7.26	Land Suitability for Custard apple	116
7.27	Land Suitability for Tamarind	117
7.28	Land Suitability for Mulberry	118
7.29	Land Suitability for Marigold	119
7.30	Land suitability for Chrysanthemum	120
7.31	Land suitability for Jasmine	121
7.32	Land suitability for Crossandra	122
7.33	Proposed Crop Plan for Handrahalu-2 microwatershed	125

## LIST OF FIGURES

2.1	Location map of Handrahalu-2 microwatershed	3
2.2a	Granite and granite gneiss rocks	4
2.2b	Alluvial rocks	4
2.3	Rainfall distribution in Koppal Taluk, Koppal District	6
2.4	Natural vegetation of Handrahalu-2 microwatershed	6
2.5a	Different crops and cropping systems in Handrahalu-2 microwatershed	8
2.5b	Different crops and cropping systems in Handrahalu-2 microwatershed	9
2.6	Current Land use – Handrahalu-2 microwatershed	10
2.7	Location of Wells- Handrahalu-2 microwatershed	10
3.1	Scanned and Digitized Cadastral map of Handrahalu-2 microwatershed	13
3.2	Satellite image of Handrahalu-2 microwatershed	13
3.3	Cadastral map overlaid on IRS PAN+LISS IV merged imagery of Handrahalu-2 microwatershed	14
3.4	Location of profiles in a transect	15
3.5	Soil phase or management units of Handrahalu-2 microwatershed	21
5.1	Land Capability Classification of Handrahalu-2 microwatershed	42
5.2	Soil Depth map of Handrahalu-2 microwatershed	44
5.3	Surface Soil Texture map of Handrahalu-2 microwatershed	45
5.4	Soil Gravelliness map of Handrahalu-2 microwatershed	46
5.5	Soil Available Water Capacity map of Handrahalu-2 microwatershed	47
5.6	Soil Slope map of Handrahalu-2 microwatershed	48
5.7	Soil Erosion map of Handrahalu-2 microwatershed	49
6.1	Soil Reaction (pH) map of Handrahalu-2 microwatershed	52
6.2	Electrical Conductivity (EC) map of Handrahalu-2 microwatershed	52
6.3	Soil Organic Carbon (OC) map of Handrahalu-2 microwatershed	53
6.4	Soil Available Phosphorus map of Handrahalu-2 microwatershed	53
6.5	Soil Available Potassium map of Handrahalu-2 microwatershed	55
6.6	Soil Available Sulphur map of Handrahalu-2 microwatershed	55
6.7	Soil Available Boron map of Handrahalu-2 microwatershed	56
6.8	Soil Available Iron map of Handrahalu-2 microwatershed	56
6.9	Soil Available Manganese map of Handrahalu-2 microwatershed	57
6.10	Soil Available Copper map of Handrahalu-2 microwatershed	57
6.11	Soil Available Zinc map of Handrahalu-2 microwatershed	58

7.1	Land suitability for Sorghum	60
7.2	Land suitability for Maize	61
7.3	Land suitability for Bajra	62
7.4	Land suitability for Groundnut	63
7.5	Land suitability for Sunflower	64
7.6	Land suitability for Red gram	65
7.7	Land suitability for Bengalgram	66
7.8	Land suitability for Cotton	67
7.9	Land suitability for Chilli	68
7.10	Land suitability for Tomato	69
7.11	Land suitability for Brinjal	70
7.12	Land suitability for onion	71
7.13	Land suitability for Bhindi	72
7.14	Land suitability for Drumstick	73
7.15	Land suitability for Mango	74
7.16	Land suitability for Guava	75
7.17	Land suitability for Sapota	76
7.18	Land suitability for Pomegranate	77
7.19	Land suitability for Musambi	78
7.20	Land suitability for Lime	79
7.21	Land suitability for Amla	80
7.22	Land suitability for Cashew	81
7.23	Land suitability for Jackfruit	82
7.24	Land Suitability for Jamun	83
7.25	Land Suitability for Custard apple	84
7.26	Land Suitability for Tamarind	85
7.27	Land Suitability for Mulberry	86
7.28	Land Suitability for Marigold	87
7.29	Land suitability for Chrysanthemum	88
7.30	Land suitability for Jasmine	89
7.31	Land suitability for Crossandra	90
7.32	Land Management Units of Handrahalu-2 microwatershed	124
9.2	Soil and water conservation map of Handrahalu-2 microwatershed	136



## **EXECUTIVE SUMMARY**

*The land resource inventory of Handrahalu-2 microwatershed was conducted using village cadastral maps and IRS satellite imagery on 1:7920 scale. The false colour composites of IRS imagery were interpreted for physiography and these physiographic delineations were used as base for mapping soils. The soils were studied in several transects and a soil map was prepared with phases of soil series as mapping units. Random checks were made all over the area outside the transects to confirm and validate the soil map unit boundaries. The soil map shows the geographic distribution and extent, characteristics, classification, behavior and use potentials of the soils in the microwatershed.*

*The present study covers an area of 462 ha in Koppal taluk and district, Karnataka. The climate is semiarid and categorized as drought - prone with an average annual rainfall of 662 mm, of which about 424 mm is received during south-west monsoon, 161 mm during north-east and the remaining 77 mm during the rest of the year. An area of about 96 per cent is covered by soils and 4 per cent by others. The salient findings from the land resource inventory are summarized briefly below.*

- ❖ The soils belong to 10 soil series and 18 soil phases (management units) and 4 land management units.*
- ❖ The length of crop growing period is <90 days and starts from 2<sup>nd</sup> week of August to 2<sup>nd</sup> week of November.*
- ❖ From the master soil map, several interpretative and thematic maps like land capability, soil depth, surface soil texture, soil gravelliness, available water capacity, soil slope and soil erosion were generated.*
- ❖ Soil fertility status maps for macro and micronutrients were generated based on the surface soil samples collected at every 320 m grid interval.*
- ❖ Land suitability for growing 31 major agricultural and horticultural crops were assessed and maps showing the degree of suitability along with constraints were generated.*
- ❖ Entire area is suitable for agriculture.*
- ❖ About 2 per cent of the soils are very shallow (<25 cm), 9 per cent of the soils are shallow (25-50 cm), 12 per cent of the soils are moderately shallow (50-75 cm), 38 per cent of the soils are moderately deep (75-100 cm), 22 per cent soils are deep (100-150 cm) and 12 per cent area has very deep (>150 cm) soils.*
- ❖ Entire area of about 96 per cent area has clayey soils at the surface.*
- ❖ About 49 per cent of the area has non-gravelly (<15%) soils, 44 per cent gravelly (15-35 % gravel) and 3 per cent very gravelly (35-60%) soils.*
- ❖ About 2 per cent are very low (<50 mm/m), 22 per cent low (51-100 mm/m), 38 per cent medium (101-150 mm/m) and 34 per cent very high (>200 mm/m) in available water capacity.*

- ❖ *An area of about 7 per cent has nearly level sloping (0-1%) and 90 per cent has very gently sloping (1-3%) lands.*
- ❖ *An area of about 43 per cent has soils that are slightly eroded (e1) and 53 per cent moderately eroded (e2) lands.*
- ❖ *An area of about 2 per cent are moderately alkaline (pH 7.8-8.4), 70 per cent strongly alkaline (pH 8.4-9.0) and 25 per cent are very strongly alkaline (pH>9.0) in soil reaction.*
- ❖ *The Electrical Conductivity (EC) of the soils is  $<2 \text{ dS m}^{-1}$  and as such the soils are non-saline.*
- ❖ *Organic carbon is low ( $<0.5\%$ ) in about 20 per cent, medium (0.5-0.75%) in about 76 per cent and 1 per cent is high ( $>0.75\%$ ) of the soils in organic carbon.*
- ❖ *Available phosphorus is low ( $<23 \text{ kg/ha}$ ) in about 23 per cent and medium (23-57 kg/ha) in 74 per cent in the microwatershed.*
- ❖ *About 2 per cent of the soils are medium (145-337 kg/ha) and 95 per cent soils are high ( $>337 \text{ kg/ha}$ ) in available potassium content.*
- ❖ *Available sulphur is medium (10-20 ppm) in 21 per cent and 75 per cent are high ( $>20 \text{ ppm}$ ) in the microwatershed.*
- ❖ *Available boron is low (0.5 ppm) in about 66 per cent area, 31 per cent area is medium (0.5-1.0 ppm) in soils.*
- ❖ *Available iron is sufficient ( $>4.5 \text{ ppm}$ ) in entire area of about 96 per cent.*
- ❖ *Available zinc is deficient ( $<0.6 \text{ ppm}$ ) in 4 per cent and sufficient ( $>0.6 \text{ ppm}$ ) in about 92 per cent area.*
- ❖ *Available manganese and copper are sufficient in all the soils.*
- ❖ *The land suitability for 31 major agricultural and horticultural crops grown in the microwatershed were assessed and the areas that are highly suitable (S1) and moderately suitable (S2) are given below. It is however to be noted that a given soil may be suitable for various crops but what specific crop to be grown may be decided by the farmer looking to his capacity to invest on various inputs, marketing infrastructure, market price and finally the demand and supply position.*

### Land suitability for various crops in the microwatershed

Crop	Suitability Area in ha (%)		Crop	Suitability Area in ha (%)	
	Highly suitable (S1)	Moderately suitable (S2)		Highly suitable (S1)	Moderately suitable (S2)
<i>Sorghum</i>	212 (46)	176 (38)	<i>Sapota</i>	-	-
<i>Maize</i>	-	393 (85)	<i>Pomegranate</i>	-	336 (73)
<i>Bajra</i>		388 (84)	<i>Musambi</i>	51 (11)	285 (62)
<i>Groundnut</i>	-	-	<i>Lime</i>	51 (11)	285 (62)
<i>Sunflower</i>	51 (11)	280 (61)	<i>Amla</i>	-	393 (85)
<i>Red gram</i>	-	315 (68)	<i>Cashew</i>	-	-
<i>Bengalgram</i>	230 (50)	159 (34)	<i>Jackfruit</i>	-	-
<i>Cotton</i>	230 (50)	159 (34)	<i>Jamun</i>	-	138 (33)
<i>Chilli</i>	-	-	<i>Custard apple</i>	230 (50)	163 (35)
<i>Tomato</i>	-	-	<i>Tamarind</i>	-	159 (34)
<i>Brinjal</i>	-	394 (85)	<i>Mulberry</i>	-	300 (65)
<i>Onion</i>	-	-	<i>Marigold</i>	-	388 (84)
<i>Bhendi</i>	-	394 (85)	<i>Chrysanthemum</i>	-	388 (84)
<i>Drumstick</i>	-	336 (73)	<i>Jasmine</i>	-	57 (12)
<i>Mango</i>	-	-	<i>Crossandra</i>	-	85 (18)
<i>Guava</i>	-	-			

- ❖ *Apart from the individual crop suitability, a proposed crop plan has been prepared for the 4 identified LMUs by considering only the highly and moderately suitable lands for different crops and cropping systems with food, fodder, fibre and other horticulture crops that helps in maintaining productivity and ecological balance in the microwatershed.*
- ❖ *Maintaining soil-health is vital for crop production and conserve soil and land resource base for maintaining ecological balance and to mitigate climate change. For this, several ameliorative measures have been suggested for these problematic soils like saline/alkali, highly eroded, sandy soils etc.*
- ❖ *Soil and water conservation treatment plan has been prepared that would help in identifying the sites to be treated and also the type of structures required.*
- ❖ *As part of the greening programme, several tree species have been suggested to be planted in marginal and submarginal lands, field bunds and also in the hillocks, mounds and ridges. That would help in supplementing the farm income, provide fodder and fuel, and generate lot of biomass which in turn would help in maintaining the ecological balance and contribute to mitigating the climate change.*



## INTRODUCTION

Land is a scarce resource and basic unit for any material production. It can support the needs of the growing population, provided they use the land in a rational and judicious manner. But what is happening in many areas of the state is a cause for concern to everyone involved in the management of land resources at the grassroots level. The area available for agriculture is about 51 per cent of the total area and more than 60 per cent of the people are still dependant on agriculture for their livelihood. The limited land area is under severe stress and strain due to increasing population pressure and competing demands of various land uses. Due to this, every year there is significant diversion of farm lands and water resources for non-agricultural purposes. Apart from this, due to lack of interest in farmers for farming, large tracts of cultivable lands are turning into fallows in many areas and this trend is continuing at an alarming rate.

Further, land degradation has emerged as a serious problem which has already affected about 38 lakh ha of cultivated area in the state. Soil erosion alone has degraded about 35 lakh ha. Almost all the uncultivated areas are facing various degrees of degradation, particularly soil erosion. Salinity and alkalinity has emerged as a major problem in more than 3.5 lakh ha in the irrigated areas of the state. Nutrient depletion and declining factor productivity is common in both rainfed and irrigated areas. The degradation is continuing at an alarming rate and there appears to be no systematic effort among the stakeholders to contain this process. In recent times, an aberration of weather due to climate change phenomenon has added another dimension leading to unpredictable situations to be tackled by the farmers.

In this critical juncture, the challenge before us is not only to increase the productivity per unit area which is steadily declining and showing a fatigue syndrome, but also to prevent or at least reduce the severity of degradation. If the situation is not reversed at the earliest, then the sustainability of the already fragile crop production system and the overall ecosystem will be badly affected in the state.

The continued neglect and unscientific use of the resources for a long time has led to the situation observed at present in the state. It is a known fact and established beyond doubt by many studies in the past that the cause for all kinds of degradation is the neglect and irrational use of the land resources. Hence, there is urgent need to generate a detailed site-specific farm level database on various land resources for all the villages/watersheds in a time bound manner that would help to protect the valuable soil and land resources and also to stabilize the farm production.

Therefore, the land resource inventory required for farm level planning is the one which investigates not only the surface but also consider the other parameters which are critical for productivity *viz.*, soils, climate, water, minerals and rocks, topography, geology, hydrology, vegetation, crops, land use pattern, animal population, socio-economic conditions, infrastructure, marketing facilities and various schemes and

developmental works of the government etc. From the data collected at farm level, the specific problems and potentials of the area can be identified and highlighted, conservation measures required for the area can be planned on a scientific footing, suitability of the area for various uses can be worked out and finally viable and sustainable land use options suitable for each and every land holding can be prescribed.

The Land Resource Inventory is basically done for identifying potential and problem areas, developing sustainable land use plans, estimation of surface run off and water harvesting potential, preparation of soil and water conservation plans, land degradation/desertification etc. The Bureau is presently engaged in developing an LRI methodology using high resolution satellite remote sensing data and Digital Elevation Model (DEM) data to prepare Landscape Ecological Units (LEU) map representing agro-ecosystem as a whole. The LEU is preferred over landform as the base map for LRI. LEU is the assemblage of landform, slope and land use. An attempt was made to upscale the soil resource information from 1:250000 and 1:50000 scale to the LEU map in Goa and other states.

The land resource inventory aims to provide site specific database for Handrahalu-2 Microwatershed in Koppal Taluk, Koppal District, Karnataka State for the Karnataka Watershed Development Department. The database was generated by using cadastral map of the village as a base along with high resolution IRS LISS IV and Cartosat-1 merged satellite imagery. Later, an attempt will be made to uplink this LRI data generated at 1:7920 scale under Sujala-III Project to the proposed Landscape Ecological Units (LEUs) map.

The study was organized and executed by the ICAR- National Bureau of Soil Survey and Land Use Planning, Regional Centre, Bangalore under Generation of Land Resource Inventory Data Base Component-1 of the Sujala-III Project funded by the World Bank.

## GEOGRAPHICAL SETTING

### 2.1 Location and Extent

The Handrahalu-2 Microwatershed is located in the central part of northern Karnataka in Koppal Taluk, Koppal District, Karnataka State (Fig. 2.1). It comprises of parts of Hanavala, Handrala, Chithepura, Hireshindhogi and Narasapura villages. It lies between  $15^{\circ}17'$  –  $15^{\circ}19'$  North latitudes and  $76^{\circ}1'$  –  $76^{\circ}3'$  East longitudes and covers an area of 462 ha. It is surrounded by Chithapura and Hanavala villages on the north, Handrala on the east, Hanavala and Handrala on the west and Hireshindhogi villages on the on the southern side.




Fig. 2.1 Location map of Handrahalu-2 Microwatershed

### 2.2 Geology

Major rock formations observed in the microwatershed are granite gneiss and alluvium (Figs. 2.2a and b). Granite gneisses are essentially pink to gray and are coarse to medium grained. They consist primarily of quartz, feldspar, biotite and hornblende. The

gray granite gneisses are highly weathered, fractured and fissured upto a depth of about 10 m. Dolerite dykes and quartz veins are common with variable width and found to occur in the village. The thickness of the alluvium generally is limited to less than a meter, except in river valleys where it is very deep extending to tens of meters. Such soils are transported and represent palaeo black soils originally formed at higher elevation, but now occupying river valleys.



Fig. 2.2 a Granite and granite gneiss rocks



Fig. 2.2 b Alluvium

### **2.3 Physiography**

Physiographically, the area has been identified as granite gneiss and alluvial landscapes based on geology. The microwatershed area has been further divided into summits, very gently sloping uplands and nearly level plains based on slope and its relief features. The elevation ranges from 502 to 523 m in the gently sloping uplands.



## 2.4 Drainage

The area is drained by several small seasonal streams that join Hire *halla* and Chenna *halla* along its course. Though, the streams are not perennial, during rainy season they carry large quantities of rain water. The microwatershed has only few small tanks which are not able to store the water flowing during the rainy season. Due to this, the ground water recharge is very much affected in the villages. This is reflected in the failure of many bore wells in the villages. If the available rain water is properly harnessed by constructing tanks and recharge structures at appropriate places in the villages, then the drinking and irrigation needs of the area can be easily met. The drainage network is dendritic to sub parallel.

## 2.5 Climate

The district falls under semiarid tract of the state and is categorized as drought-prone with an average annual rainfall of 662 mm (Table 2.1). Maximum of 424 mm precipitation takes place during the south-west monsoon period from June to September, north-east monsoon contributes about 161 mm and prevails from October to early December and the remaining 77 mm takes place during the rest of the year. The winter season is from December to February. During April and May, the temperatures reach up to 45 °C and in December and January, the temperatures will go down to 16 °C. Rainfall distribution is shown in Figure 2.3. The average Potential Evapo Transpiration (PET) is 145 mm and varies from a low of 101 mm in December and 193 mm in the month of May. The PET is always higher than precipitation in all the months except in the month of September. Generally, the Length of crop Growing Period (LGP) is <90 days and starts from 2<sup>nd</sup> week of August to 2<sup>nd</sup> week of November.

**Table 2.1 Mean Monthly Rainfall, PET, 1/2 PET at Koppal Taluk and District**

Sl. No.	Months	Rainfall	PET	1/2 PET
1	January	1.60	116.70	58.35
2	February	1.50	129.20	64.60
3	March	14.10	169.80	84.90
4	April	18.10	180.60	90.30
5	May	41.60	193.50	96.75
6	June	85.80	167.90	83.95
7	July	72.10	156.20	78.10
8	August	110.50	152.50	76.25
9	September	155.60	138.50	69.25
10	October	116.30	122.30	61.15
11	November	36.00	106.40	53.20
12	December	9.10	101.00	50.50
	<b>TOTAL</b>	<b>662.30</b>	<b>144.55</b>	




Fig. 2.3 Rainfall distribution in Koppal Taluk and District

### 2.6 Natural Vegetation

The natural vegetation is sparse comprising few tree species, shrubs and herbs. The mounds, ridges and boulders occupy sizeable areas which are under thin to moderately thick forest vegetation. Still, there are some remnants of the past forest cover which can be seen in patches in some ridges and hillocks in the microwatershed (Fig 2.4).

Apart from the continuing deforestation, the presence of large population of goats, sheep and other cattle in the microwatershed is causing vegetative degradation of whatever little vegetation left in the area. The uncontrolled grazing has left no time for the regeneration of the vegetative cover. This leads to the accelerated rate of erosion on the hill slopes, resulting in the formation of deep gullies in the foot slopes and eventually resulting in the heavy siltation of few tanks and reservoirs in the microwatershed.



Fig 2.4 Natural vegetation of Handrahalu-2 Microwatershed

## 2.7 Land Utilization

About 91 per cent area (Table 2.2) in Koppal district is cultivated at present and about 16 per cent of the area is sown more than once. The cropping intensity is 118 per cent. An area of about 3 per cent is currently barren. Forests occupy a small area of about 5 per cent and the tree cover is in a very poor state. Most of the mounds, ridges and bouldery areas have very poor vegetative cover. Major crops grown in the area are sorghum, maize, bajra, cotton, safflower, sunflower, red gram, horse gram, onion, mulberry, pomegranate, sugarcane, bengalgram and groundnut (Fig 2.5). While carrying out land resource inventory, the land use/land cover particulars are collected from all the survey numbers and a current land use map of the microwatershed is prepared. The current land use map prepared shows the arable and non-arable lands, other land uses and different types of crops grown in the area. The current land use map of Handrahalu-2 Microwatershed is presented in Fig. 2.6. Simultaneously, enumeration of existing wells (bore wells and open wells) and other soil and water conservation structures in the microwatershed is made and their location in different survey numbers is marked on the cadastral map. Map showing the location of wells and other water bodies in Handrahalu-2 Microwatershed is given Fig. 2.7.

**Table 2.2 Land Utilization in Koppal District**

Sl. no.	Agricultural land use	Area ( ha)	Per cent
1	Total geographical area	552495	
2	Total cultivated area	500542	90.6
3	Area sown more than once	92696	16.8
4	Trees and groves	210	0.04
5	Cropping intensity	-	118
6	Forest	29451	5.33
7	Cultivable wasteland	2568	0.46
8	Permanent Pasture land	14675	2.66
9	Barren land	16627	3.01
10	Non agricultural land	40591	7.35
11	Current fallow	19660	3.56



Fig. 2.5 (a) Different crops and cropping systems in Handrahalu-2 Microwatershed



Fig. 2.5 (b) Different crops and cropping systems in Handrahalu-2 Microwatershed




Fig. 2.6 Current Land Use – Handrahalu-2 Microwatershed




Fig. 2.7 Location of wells-Handrahalu-2 Microwatershed

## SURVEY METHODOLOGY

The purpose of land resource inventory is to delineate similar areas (soil series and phases), which respond or expected to respond similarly for a given level of management. This was achieved in Handrahalu-2 Microwatershed by the detailed study of all the soil characteristics (depth, texture, colour, structure, consistence, coarse fragments, porosity, soil reaction, soil horizons etc.) and site characteristics (slope, erosion, drainage, occurrence of rock fragments etc.) followed by grouping of similar areas based on soil-site characteristics into homogeneous (management units) units and showing their extent and geographic distribution on the microwatershed cadastral map. The detailed soil survey at 1:7920 scale was carried out in 462 ha area. The methodology followed for carrying out land resource inventory was as per the guidelines given in Soil Survey Manual (IARI, 1971; Soil Survey Staff, 2006; Natarajan *et al.*, 2015) which is briefly described below.

### 3.1 Base Maps

The detailed survey of the land resources occurring in the microwatershed was carried out by using digitized cadastral map and satellite imagery as base supplied by the KRSRSAC. The cadastral map shows field boundaries with their survey numbers, location of tanks, streams and other permanent features of the area (Fig. 3.1). Apart from the cadastral map, remote sensing data products from Cartosat-1 and LISS IV merged at the scale of 1:7920 were used in conjunction with the cadastral map to identify the geology, landscapes, landforms and other surface features. The imagery helped in the identification and delineation of boundaries between hills, uplands and lowlands, water bodies, forest and vegetated areas, roads, habitations and other cultural features of the area (Fig. 3.2). The cadastral map was overlaid on the satellite imagery (Fig. 3.3) that helps to identify the parcel boundaries and other permanent features. Apart from cadastral maps and images, toposheets of the area (1:50,000 scale) were used for initial traversing, identification of geology, landscapes and landforms, drainage features, present land use and also for selection of transects in the microwatershed.

### 3.2 Image Interpretation for Physiography

False Colour Composites (FCC) of Cartosat-I and LISS-IV merged satellite data covering the microwatershed area was visually interpreted using image interpretation elements and all the available collateral data with local knowledge. The delineated physiographic boundaries were transferred on to a cadastral map overlaid on satellite imagery. Physiographically, the area has been identified as granite gneiss and alluvial landscapes and is divided into landforms such as ridges, mounds and uplands based on slope. They were further subdivided into physiographic/ image interpretation units based on image characteristics. The image interpretation legend for physiography is given below.

## **Image Interpretation Legend for Physiography**

### **G- Granite gneiss landscape**

G1	Hills/ Ridges/ Mounds
G11	Summits
G12	Side slopes
G121	Side slopes with dark grey tones
G2	Uplands
G21	Summits
G22	Gently sloping uplands
G221	Gently sloping uplands, yellowish green (eroded)
G222	Gently sloping uplands, yellowish white (severely eroded)
G23	Very gently sloping uplands
G231	Very gently sloping uplands, yellowish green
G232	Very gently sloping uplands, medium green and pink
G233	Very gently sloping uplands, pink and green (scrub land)
G234	Very gently sloping uplands, medium greenish grey
G235	Very gently sloping uplands, yellowish white (eroded)
G236	Very gently sloping uplands, dark green
G237	Very gently sloping uplands, medium pink (coconut garden)
G238	Very gently sloping uplands, pink and bluish white (eroded)

### **DSe Alluvial landscape**

#### **Dse 1 Summit**

- Dse 11 Nearly level Summit with dark grey tone
- Dse 12 Nearly level Summit with medium grey tone
- Dse 13 Nearly level Summit with whitish grey tone
- Dse 14 Nearly level Summit with whitish tone (Calcareousness)
- Dse 15 Nearly level Summit with pinkish grey tone
- Dse 16 Nearly level Summit with medium pink tone
- Dse 17 Nearly level Summit with bluish white tone
- Dse 18 Nearly level Summit with greenish grey tone

#### **Dse 2 Very gently sloping**

- Dse 21 Very gently sloping, whitish tone
- Dse 22 Very gently sloping, greyish pink tone
- Dse 23 Very gently sloping, whitish grey tone
- Dse 24 Very gently sloping, medium grey tone
- Dse 25 Very gently sloping, medium pink tone
- Dse 26 Very gently sloping, dark grey tone
- Dse 27 Very gently sloping, bluish grey tone
- Dse 28 Very gently sloping, greenish grey tone
- Dse 29 Very gently sloping, Pinkish grey

#### **Dsa 25 – Nearly Level Lands**

- Dsa 251- Nearly level, Grayish green tone
- Dsa 252- Nearly level, Bluish grey tone
- Dsa 253- Nearly level, Light green tone
- Dsa 254- Nearly level, Medium green tone
- Dsa 255- Nearly level, Greenish pink tone
- Dsa 256- Nearly level, Whitish green
- Dsa 257- Nearly level, Pink tone
- Dsa 258- Nearly level, Whitish grey tone
- Dsa 259- Nearly level, Grayish Pink






Fig. 3.1 Scanned and Digitized Cadastral map of Handrahalu-2 Microwatershed




Fig. 3.2 Satellite Image of Handrahalu-2 Microwatershed




Fig. 3.3 Cadastral map overlaid on IRS PAN+LISS IV merged imagery of Handrahalu-2 Microwatershed

### 3.3 Field Investigation

The field boundaries and survey numbers given on the cadastral sheet were located on the ground by following permanent features like roads, cart tracks, *nallas*, streams, tanks etc., and wherever changes were noticed, they were incorporated on the microwatershed cadastral map. Preliminary traverse of the microwatershed was carried out with the help of cadastral map, imagery and toposheets. While traversing, landforms and physiographic units identified were checked and preliminary soil legend was prepared by studying soils at few selected places. Then, intensive traversing of each physiographic unit like hills, ridges, uplands and plains was carried out. Based on the variability observed on the surface, transects (Fig. 3.4) were selected across the slope covering all the landform units in the microwatershed (Natarajan and Dipak Sarkar, 2010).




Fig. 3.4 Location of profiles in a transect

In the selected transect, soil profiles (Fig. 3.4) were located at closely spaced intervals to take care of any change in the land features like break in slope, erosion, gravel, stones etc. In the selected sites, soil profiles (vertical cut showing the soil layers from surface to the rock) were opened upto 200 cm or to the depth limited by rock or hard substratum and studied in detail for all their morphological and physical characteristics. The soil and site characteristics were recorded for all profile sites on a standard proforma as per the guidelines given in USDA Soil Survey Manual (Soil Survey Staff, 2012). Apart from the transect study, profiles were also studied at random, almost like in a grid pattern, outside the transect areas to validate the soil map unit boundaries.

Based on the soil characteristics, the soils were grouped into different soil series. Soil series is the most homogeneous unit having similar horizons and properties and behaves similarly for a given level of management. Soil depth, texture, colour, kind of horizon and horizon sequence, amount and nature of gravel present, calcareousness, nature of substratum etc, were used as the major differentiating characteristics for identifying soil series occurring in the area. The differentiating characteristics used for identifying the soil series are given in Table 3.1. Based on the above characteristics, 10 soil series were identified in Handrahalu-2 Microwatershed.

**Table 3.1 Differentiating Characteristics used for identifying Soil Series  
(Characteristics are of Series Control Section)**

Soils of Granite gneiss Landscape							
Sl. No	Soil Series	Depth (cm)	Colour (moist)	Texture	Gravel (%)	Horizon sequence	Calcareousness
1	Belagatti (BGT)	<25	10YR3/1,3/2,4/2	gc	>35	Ap-Crk	es
Soils of Alluvial Landscape							
2	Muttal (MTL)	25-50	10YR3/2,3/3,4/2 7.5YR3/2,3/3,6/4	gc	15-35	Ap-Bw-Ck	e-ev
3	Ravanaki (RNK)	50-75	7.5YR3/2,3/3,5/2,5/3 10YR3/1,3/2,4/1,4/2,5/1,6/1	c	<15	Ap-Bw-Cr	e-ev
4	Dambarahalli (DRL)	75-100	10YR2/1,3/1,4/3	c	<15	Ap-Bss-Ck	e-es
5	Narasapura (NSP)	75-100	10YR3/1,3/2,4/2	c		Ap-Bw-Cr	e-es
6	Gatareddihal (GRH)	100-150	10YR 2/1,3/1 2.5YR4/3,5/4	c	<15	Ap-Bss-BC-C	es
7	Handrala (HDL)	100-150	10YR 2/1,3/1,4/1	c	-	Ap-Bss-Ck	es
8	Kadagathur (KDT)	>150	10YR3/1,3/2,3/3, 7.5YR3/3,3/4	sc-c	-	Ap-Bw	-
9	Murlapur (MLR)	>150	10YR2/1,2/2,3/1,3/2,4/1	c	10-20	Ap-Bss	e-es
10	Bardur (BDR)	>150	10YR2/1,3/1,3/2	c	<15	Ap-Bss	es

### 3.4 Soil Mapping

The area under each soil series was further separated into 18 soil phases and their boundaries delineated on the cadastral map based on the variations observed in the texture of the surface soil, slope, erosion, presence of gravel, stoniness etc. A soil phase is a subdivision of soil series based mostly on surface features that affect its use and management.

The soil mapping units are shown on the soil map (Fig. 3.5) in the form of symbols. During the survey many soil profile pits, few minipits and a few auger bores representing different landforms occurring in the microwatershed were studied. In addition to the profile study, spot observations in the form of minipits, road cuts, terrace cuts etc., were studied to validate the soil boundaries on the soil map. The soil map shows the geographic distribution of 18 mapping units representing 10 soil series occurring in the microwatershed. The soil map unit (soil legend) description is presented in Table 3.2. The soil phase map (management units) shows the distribution of 18 soil phases mapped in the microwatershed. Each mapping unit (soil phase) delineated on the map has similar soil and site characteristics. In other words, all the farms or survey numbers included in one phase will have similar management needs and have to be treated accordingly.

### 3.5 Laboratory Characterization

Soil samples for each soil series were collected from representative master profiles for laboratory characterization by following the methods outlined in the Laboratory Manual (Sarma *et al*, 1987). Surface soil samples collected in the year 2017 from Handrahalu-2 farmer's fields (42 samples) for fertility status (major and micronutrients) at 320 m grid interval were analyzed in the laboratory (Katyal and Rattan, 2003). By linking the soil fertility data to the survey numbers through GIS, soil fertility maps were generated using Kriging method for the microwatershed.

**Table 3.2 Soil map unit description of Handrahalu-2 Microwatershed**

Soil map unit No*	Soil Series	Soil Phase Symbol	Mapping Unit Description	Area in ha (%)
<b>Soils of Granite and Granite gneiss landscape</b>				
	BGT		Belagatti soils are very shallow (<25 cm), well drained, have very dark gray to very dark grayish brown, calcareous gravelly clay black soils occurring on very gently to gently sloping uplands under cultivation	<b>8 (1.74)</b>
8		BGTmB1g2	Clay surface, slope 1-3%, slight erosion, very gravelly (35-60%)	8 (1.74)
<b>Soils of Alluvial landscape</b>				
	MTL		Muttal soils are shallow (25-50 cm), well drained, have very dark grayish brown to dark brown, calcareous black clay soils occurring on nearly level to gently sloping plains under cultivation	<b>43 (9.34)</b>
311		MTLmB2g1	Clay surface, slope 1-3%, moderate erosion, gravelly (15-35%)	43 (9.34)
	RNK		Ravanaki soils are moderately shallow (50-75 cm), moderately well drained, have dark brown to very dark grayish brown and dark gray, calcareous clay black soils occurring on nearly level to very gently sloping plains under cultivation	<b>57 (12.4)</b>
333		RNKmB1	Clay surface, slope 1-3%, slight erosion	25 (5.49)
334		RNKmB1g1	Clay surface, slope 1-3%, slight erosion, gravelly (15-35%)	31 (6.74)
337		RNKmB2g1	Clay surface, slope 1-3%, moderate erosion, gravelly (15-35%)	1 (0.17)
	DRL		Dambarahalli soils are moderately deep (75-100 cm), moderately well drained, have dark brown to very dark gray, calcareous black cracking clay soils occurring on nearly level to very gently sloping plains under cultivation	<b>16 (3.52)</b>
348		DRLmB1	Clay surface, slope 1-3%, slight erosion	16 (3.52)
	NSP		Narasapura soils are moderately deep (75-100 cm), moderately well drained, have dark grayish brown to very dark grayish brown and very dark gray, black calcareous cracking clay soils occurring on nearly level to very	<b>161 (34.8)</b>

Soil map unit No*	Soil Series	Soil Phase Symbol	Mapping Unit Description	Area in ha (%)
			gently sloping plains under cultivation	
360		NSPmB1	Clay surface, slope 1-3%, slight erosion	39 (8.37)
361		NSPmB1g1	Clay surface, slope 1-3%, slight erosion, gravelly (15-35%)	7 (1.46)
362		NSPmB2	Clay surface, slope 1-3%, moderate erosion	65 (14.1)
363		NSPmB2g1	Clay surface, slope 1-3%, moderate erosion, gravelly (15-35%)	50 (10.87)
	GRH		Gatareddihal soils are deep (100-150 cm), moderately well drained, have light olive brown to very dark gray, calcareous black cracking clay soils occurring on nearly level to very gently sloping plains under cultivation	<b>54</b> <b>(11.76)</b>
372		GRHmB1g1	Clay surface, slope 1-3%, slight erosion, gravelly (15-35%)	12 (2.56)
374		GRHmB2g1	Clay surface, slope 1-3%, moderate erosion, gravelly (15-35%)	42 (9.2)
	HDL		Handrala soils are deep (100-150 cm), moderately well drained, have dark gray to very dark gray, black calcareous cracking clay soils occurring on very gently sloping plains under cultivation	<b>48</b> <b>(10.32)</b>
380		HDLmB1	Clay surface, slope 1-3%, slight erosion	27 (5.85)
382		HDLmB2	Clay surface, slope 1-3%, moderate erosion	3 (0.63)
383		HDLmB2g1	Clay surface, slope 1-3%, moderate erosion, gravelly (15-35%)	18 (3.84)
	KDT		Kadagathur soils are very deep (>150 cm), moderately well drained, have dark brown to very dark grayish brown, sandy clay to clay black soils occurring on nearly level to very gently sloping plains under cultivation	<b>21</b> <b>(4.64)</b>
405		KDTmB2	Clay surface, slope 1-3%, moderate erosion	21 (4.64)
	MLR		Murlapur soils are very deep (>150 cm), moderately well drained, have very dark grayish brown to very dark gray, calcareous black cracking clay soils occurring on nearly level to very gently sloping plains under cultivation	<b>31</b> <b>(6.68)</b>
411		MLRmA1	Clay surface, slope 0-1%, slight erosion	31 (6.68)
	BDR		Bardur soils are very deep (>150 cm), moderately well drained, have very dark grayish brown to very dark gray, black cracking calcareous clay soils occurring on nearly level to very gently sloping plains under cultivation	<b>5 (1.01)</b>
432		BDRmB1g2	Clay surface, slope 1-3%, slight erosion, very gravelly (35-60%)	5 (1.01)
1000	Others		Habitaion and waterbody	18 (3.79)

\*Soil map unit numbers are continuous for the taluk, not the microwatersheds

### **3.6 Land Management Units**

The 18 soil phases identified and mapped in the microwatershed were regrouped into 4 Land Management units (LMU's) for the purpose of preparing a Proposed Crop Plan for sustained development of the microwatershed. The database (soil phases) generated under LRI was utilized for identifying Land Management units (LMU's) based on the management needs. One or more than one soil site characteristic having influence on the management have been chosen for identification and delineation of LMUs. For Handrahalu-2 Microwatershed, five soil and site characteristics, namely soil depth, soil texture, slope, erosion and gravel content have been considered for defining LMUs. The land use classes are expected to behave similarly for a given level of management.








Fig 3.5 Soil Phase or Management Units-Handrahalu-2 Microwatershed



## THE SOILS

Detailed information pertaining to the nature, extent and distribution of different kinds of soils occurring in Handrahalu-2 Microwatershed is provided in this chapter. The microwatershed area has been identified as granite gneiss and alluvial landscapes based on geology. In all, 10 soil series are identified. Soil formation is the result of the combined effect of environmental and terrain factors that are reflected in soil morphology. The soil formation is dominantly influenced by the parent material, climate, time and relief.

A brief description of each of the 10 soil series identified followed by 18 soil phases (management units) mapped (Fig. 3.4) are furnished below. The physical and chemical characteristics of soil series identified in Handrahalu-2 Microwatershed are given in Table 4.1. The soils in any one map unit differ from place to place in their depth, texture, slope, gravelliness, erosion or any other site characteristic that affect management. The soil phase map can be used for identifying the suitability of areas for growing specific crops or for other alternative uses and also for deciding the type of conservation structures needed. The detailed information on soil and site-characteristics like soil depth, surface soil texture, slope, erosion, gravelliness, AWC, LCC etc, with respect to each of the soil phase identified is given village/survey number wise for the microwatershed in Appendix-I.

### **4.1 Soils of granite and granite gneiss landscape**

In this landscape, only one soil series are identified and mapped. Of these, Belagatti (BGT) series occupies an area of 8 ha (2%) in the microwatershed. The brief description of each soil series along with the soil phases identified and mapped is given below.

**4.1.1 Belagatti (BGT) Series:** Belagatti soils are very shallow (< 25 cm), well drained, have dark gray to dark grayish brown gravelly clay soils. They have developed from granite gneiss and occur on very gently sloping uplands. The Belagatti series has been classified as a member of the clayey mixed, isohyperthermic family of Lithic Ustorthents.

The thickness of the soil is less than 25 cm. Its colour is in 10 YR hue with value 3 to 4 and chroma 1 to 2. The texture is clay with more than 35 per cent gravel and the available water capacity is very low (<50 mm/m). Only one phase was identified and mapped.



Landscape and soil profile characteristics of Belagatti (BGT) Series

## 4.2 Soils of Alluvial landscape

In this landscape, nine soil series have been identified and mapped. Of these, Narasapura (NSP) series occupies maximum area of 161 ha (35%), Ravanaki (RNK) 57 ha (12%), Gatareddihal (GRH) 54 ha (12%), Handrala (HDL) 48 ha (10%), Muttal (MTL) 43 ha (9%), Murlapur (MLR) 31 ha (7%), Kadagathur (KDT) 21 ha (5%), Dambarahalli (DRL) 16 ha (4%) and Bardur (BDR) occupy an area of about 5 ha (1%) in the microwatershed. The brief description of soil series along with the soil phases identified and mapped is given below.

**4.2.1 Muttal (MTL) Series:** Muttal soils are shallow (25-50 cm), well drained, have dark brown to very dark grayish brown, gravelly clay soils. They have developed from alluvium and occur on nearly level to very gently sloping plains. The Muttal series has been classified as a member of the clayey, mixed, isohyperthermic (calc) family of (Paralithic) Haplustepts.

The thickness of the solum ranges from 30 to 50 cm. The thickness of A horizon ranges from 15 to 18 cm. Its colour is in 7.5 YR and 10 YR hue with value 2 to 3 and chroma 2.5 to 4. The texture varies from sandy clay to clay with 10 to 15 per cent gravel. The thickness of B horizon ranges from 18 to 32 cm. Its colour is in 10 YR and 7.5 YR hue with value 2 to 6 and chroma 2 to 4. Its texture is sandy clay to clay. The available water capacity is low (51-100 mm/m). Only one phase was identified and mapped.



Landscape and soil profile characteristics of Muttal (MTL) Series

**4.1.3 Ravanaki (RNK) Series:** Ravanaki soils are moderately shallow (50-75 cm), well drained, have dark brown to very dark grayish brown, calcareous sodic clay soils. They have developed from alluvium and occur on nearly level to very gently sloping plains. The Ravanaki series has been classified as a member of the very fine, smectitic, isohyperthermic (calc) family of Typic Haplustepts.

The thickness of the solum ranges from 50 to 75 cm. The thickness of A horizon ranges from 15 to 20 cm. Its colour is in 7.5 YR and 10 YR hue with value 2 to 3 and chroma 2.5 to 4. The texture varies from sandy clay to clay with 10 to 15 per cent gravel. The thickness of B horizon ranges from 35 to 60 cm. Its colour is in 10 YR and 7.5 YR hue with value 2 to 6 and chroma 2 to 4. Its texture is sandy clay to clay with gravel content of 10 to 20 per cent. The available water capacity is low (51-100 mm/m). Three phases were identified and mapped.



Landscape and soil Profile Characteristics of Ravanaki (RNK) Series

**4.1.4 Dambarahalli (DRL) Series:** Dambarahalli soils are moderately deep (75-100 cm), moderately well drained, have black and very dark gray to dark brown calcareous cracking clay soils. They have developed from alluvium and occur on very gently to gently sloping plains under cultivation. The Dambarahalli series has been classified as a member of the very fine, smectitic (calc) isohyperthermic family of Typic Haplusterts.

The thickness of the solum ranges from 75 to 99 cm. The thickness of A horizon ranges from 13 to 24 cm. Its colour is in 10 YR hue with value 3 to 4 and chroma 1 to 2. The texture is clay. The thickness of B horizon ranges from 54 to 85 cm. Its colour is in 10 YR hue with value 2 to 4 and chroma 1 to 3. Its texture is clay and are calcareous. The available water capacity is high (151-200 mm/m). Two soil phases were identified and mapped. Only one phase was identified and mapped.



Landscape and soil profile characteristics of Dambarahalli (DRL) Series.

**4.1.5 Narsapura (NSP) series:** Narsapura soils are moderately deep (75-100 cm), moderately well drained, have dark grayish brown to very dark grayish brown and very dark gray, black calcareous cracking clay soils. They have developed from alluvium and occur on very gently sloping uplands. The Narsapura series has been classified as a member of the very fine, smectitic, isohyperthermic (calc) family of Typic Haplustepts.


The thickness of the solum is 76 to 98 cm. The thickness of A horizon ranges from 15 to 19 cm. Its colour is in 10 YR hue with value 3 and chroma 1 to 2. The texture is clay with no gravel. The thickness of B horizon ranges from 57 to 83 cm. Its colour is in 10 YR hue with value 3 to 5 and chroma 1 to 3. Its texture is clay and are calcareous. The available water capacity is medium (101-150 mm/m). Four phases were identified and mapped.



Landscape and soil profile characteristics of Narsapura (NSP) series

**4.1.6 Gatareddihal (GRH) Series:** Gatareddihal soils are deep (100-150 cm), moderately well drained have black or dark grey to light olive brown, calcareous sodic clay soils. They are developed from alluvium and occur on nearly level to very gently sloping plains under cultivation. The Gatareddihal series has been classified as a member of the very fine, smectitic, isohyperthermic family of Sodic Haplusterts.

The thickness of the solum ranges from 102 to 149 cm. The thickness of A-horizon ranges from 12 to 19 cm. Its colour is in 7.5 YR, 10 YR hue with value 3 to 4 and chroma 1 to 6. The texture is sandy clay loam to clay. The thickness of B-horizon ranges from 86 to 117 cm. Its colour is in 10 YR and 7.5 YR hue with value 3 and chroma 2 to 6. Texture is clay with less than 15 per cent gravel. The available water capacity is very high (>200 mm/m). Two phases were identified and mapped.



Landscape and soil profile characteristics of Gatareddihal (GRH) Series

**4.1.7 Handrala (HDL) Series:** Handrala soils are deep (100-150 cm), moderately well drained, have black, very dark brown to dark gray cracking clay soils. They are developed from alluvium and occur on very gently to gently sloping plains. The Handrala series has been classified as a member of the very fine, smectitic, isohyperthermic (calc) family of Typic Haplusterts.

The thickness of the solum ranges from 102 to 149 cm. The thickness of A horizon ranges from 14 to 26 cm. Its colour is in 10 YR hue with value 3 and chroma 1. The texture is clay. The thickness of B horizon ranges from 103 to 127 cm. Its colour is in 10 YR hue with value 2 to 4 and chroma 1 to 2. Texture is dominantly clay. The available water capacity is very high (>200 mm/m). Three phases were identified and mapped.




Landscape and soil Profile Characteristics of Handrala (HDL) Series

**4.1.8 Kadagathur (KDT) Series:** Kadagathur soils are very deep (>150 cm), moderately well drained, have dark brown to very dark grayish brown sandy clay to clay soils. They have developed from weathered alluvium and occur on nearly level to very gently sloping plains under cultivation. The Kadagathur series has been classified as a member of the fine, mixed, isohyperthermic family of Fluventic Haplustepts.

The thickness of the solum is more than 150 cm. The thickness of A horizon ranges from 8 to 14 cm. Its colour is in 10 YR hue with value 3 and chroma 4. The texture varies is sandy loam. The thickness of B horizon is more than 150 cm. Its colour is in 10 YR and 7.5 YR hue with value 3 and chroma 1 to 4. Its texture is sandy clay to clay. The available water capacity is very high (>200 mm/m). Only one phase was identified and mapped.





Landscape and soil profile characteristics of Kadagathur (KDT) Series

**4.1.9 Murlapur (MLR) series:** Murlapur soils are very deep (>150 cm), moderately well drained, have very dark grayish brown to very dark gray, calcareous black cracking clay soils. They have developed from alluvium and occur on nearly level to very gently sloping uplands. The murlapur series has been classified as a member of the very fine, smectitic, isohyperthermic (Calc) family of Typic Haplusterts.

The thickness of the solum is >150 cm. The thickness of A horizon ranges from 20 to 25 cm. Its colour is in 10 YR hue with value 3 and chroma 1. The texture is clay with no gravel. The thickness of B horizon ranges from 150 to 190 cm. Its colour is in 10 YR hue with value 3 to 4 and chroma 1 to 2. Its texture is clay. The available water capacity is very high (>200 mm/m). Only one phase was identified and mapped.



Landscape and soil profile characteristics of Murlapur (MLR) series

**4.1.10 Bardur (BDR) Series:** Bardur soils are very deep (>150 cm), moderately well drained, have very dark grayish brown to very dark gray, black calcareous cracking clay soils. They are developed from alluvium and occur on nearly level to very gently sloping plains under cultivation. The Bardur series has been classified as a member of the very fine, smectitic, isohyperthermic (calcareous) family of Typic Haplusterts.

The thickness of the solum is more than 150 cm. The thickness of A horizon ranges from 15 to 19 cm. Its colour is in 10 YR hue with value 2 and chroma 1 with clay texture. The thickness of B horizon ranges from 146 to 180 cm. Its colour is in 10 YR hue with value 2 to 3 and chroma 1 to 2. Its texture is clay and is calcareous with less than 15 per cent gravel. The available water capacity is very high (>200 mm/m). Only one phase was identified and mapped.



Landscape and soil profile characteristics of Bardur (BDR) Series

**Table: 4.1 Physical and Chemical Characteristics of Soil Series identified in Handrahalu-2 Microwatershed**

**Series Name:** Belagatti (BGT), **Pedon:** A2/RM-5

**Location:** 15°19'10.8"N, 75°57'48.1"E, Kavalura village, Koppal Taluk and District

**Analysis at:** NBSS&LUP, Regional Centre, Bangalore. **Classification:** Clayey mixed, isohyperthermic Lithic Ustorthents

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-23	Ap	36.14	20.34	43.52	10.87	6.93	5.97	8.42	3.94	40	c	29.53	17.97

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO <sub>3</sub>	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP
	Water	CaCl <sub>2</sub>	M KCl	dS m <sup>-1</sup>	%	%	Ca	Mg	K	Na	Total				
							cmol kg <sup>-1</sup>								
0-23	8.4			0.157	0.12	18.24			0.73	0.50		44.84	1.03		1.11

*Contd...*

**Series Name:** Muttal (MTL), **Pedon:** RM-13

**Location:** 15°14'30.8"N, 75°56'50.6"E, Gatareddihalla village, Koppal Taluk and District

**Analysis at:** NBSS&LUP, Regional Centre, Bangalore. **Classification:** Clayey, mixed, isohyperthermic (calc) (paralithic) Haplustepts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-20	Ap	39.05	13.74	47.21	3.05	5.05	8.21	14.63	8.11	15-30	c	29.95	17.94
20-34	Bwk	28.77	19.57	51.66	4.81	4.71	4.92	9.09	5.24	10	c	33.44	21.56

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO <sub>3</sub>	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP
	Water	CaCl <sub>2</sub>	M KCl				Ca	Mg	K	Na	Total				
				dS m <sup>-1</sup>	%	%	cmol kg <sup>-1</sup>						%	%	
0-20	8.27	-	-	0.202	0.79	6.10	-	-	0.62	0.25	-	36.64	0.78	-	0.69
20-34	8.36	-	-	0.177	0.99	23.04	-	-	0.29	0.38	-	39.60	0.77	-	0.96

*Contd...*

**Series Name:** Ravanaki (RNK), **Pedon:** RM-20

**Location:** 15°14'22.7"N, 75°57'45.8"E, Gatareddihalla village, Koppal Taluk and District

**Analysis at:** NBSS&LUP, Regional Centre, Bangalore. **Classification:** Very fine, smectitic, isohyperthermic (calc) Typic Haplustepts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-28	Ap	24.43	17.76	57.81	5.30	3.89	3.78	7.14	4.32	20	c	41.40	29.60
28-55	Bw	18.77	15.59	65.64	2.74	3.73	2.85	4.83	4.61	10	c	46.71	35.18

Depth (cm)	pH (1:2.5)			E.C. (1:2.5) dS m <sup>-1</sup>	O.C. %	CaCO <sub>3</sub> %	Exchangeable bases					CEC	CEC/Clay	Base saturation %	ESP %
	Water	CaCl <sub>2</sub>	M KCl				Ca	Mg	K	Na	Total				
							cmol kg <sup>-1</sup>								
0-28	8.86	-	-	0.483	0.63	15.48	-	-	0.86	6.27	-	37.00	0.64	-	6.78
28-55	8.61	-	-	1.4	0.23	13.68	-	-	0.68	12.27	-	53.20	0.81	-	9.22

*Contd...*

**Series Name:** Dombarahalli (DRL)

**Pedon:** R-8

**Location:** 15°13'96.2"N, 75°57'48.6" E Rangunathanahalli village, Koppal Taluk and District

**Analysis at:** NBSS&LUP, Regional Centre, Bangalore.

**Classification:** Very fine, smectitic, isohyperthermic (calc) Typic Haplusterts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-15	Ap	28.25	19.48	52.27	4.76	4.44	4.87	8.23	5.95	-	c	39.86	27.20
15-27	BA1	21.55	20.00	58.45	3.76	2.76	3.43	6.30	5.30	-	c	46.35	34.84
27-45	Bss1	14.86	20.89	64.25	2.46	2.23	2.23	3.91	4.02	-	c	57.99	41.06
45-80	Bss2	10.42	19.04	70.54	1.74	1.97	1.27	2.78	2.66	-	c	66.36	36.24

Depth (cm)	pH (1:2.5)			E.C. (1:2.5) dS m <sup>-1</sup>	O.C. %	CaCO <sub>3</sub> %	Exchangeable bases					CEC	CEC/Clay	Base saturation %	ESP %
	Water	CaCl <sub>2</sub>	M KCl				Ca	Mg	K	Na	Total				
0-15	8.78	-	-	0.42	0.32	12.35	-	-	0.59	4.25	-	49.70	0.95	100.00	5.62
15-27	9.03	-	-	0.61	0.30	12.48	-	-	0.30	8.96	-	57.23	0.98	100.00	10.07
27-45	9.10	-	-	0.67	0.34	11.70	-	-	0.25	11.85	-	60.71	0.95	100.00	14.05
45-80	9.18	-	-	0.86	0.32	13.39	-	-	0.27	15.40	-	63.33	0.90	100.00	18.45

Contd...

**Series Name:** Narsapura (NSP), **Pedon:** A2/RM-2

**Location:** 15°19'86.9"N, 75°57'86.1"E, Kavalura village, Koppal Taluk and District

**Analysis at:** NBSS&LUP, Regional Centre, Bangalore.

**Classification:** Very fine, smectitic, isohyperthermic (Calc) Vertic Haplustepts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-29	Ap	31.32	16.52	52.16	5.51	5.40	5.51	9.83	5.08	10	c	38.86	27.64
29-52	Bw1	13.30	22.08	64.62	2.52	2.41	2.41	3.67	2.29	05	c	49.88	40.05
52-77	BW2	13.22	17.39	69.40	3.56	2.41	1.95	2.76	2.53	05	c	51.33	41.55

Depth (cm)	pH (1:2.5)			E.C. (1:2.5) dS m <sup>-1</sup>	O.C. %	CaCO <sub>3</sub> %	Exchangeable bases					CEC	CEC/Clay	Base saturation %	ESP %
	Water	CaCl <sub>2</sub>	M KCl				Ca	Mg	K	Na	Total				
0-29	9.16	-	-	0.615	0.23	9.36	-	-	0.72	10.98	-	51.09	0.98	-	8.60
29-52	8.69	-	-	2.01	0.5	8.64	-	-	0.55	24.42	-	60.63	0.94	-	16.11
52-77	8.52	-	-	2.68	0.46	7.68	-	-	0.50	25.65	-	60.74	0.88	-	16.90

Contd...

**Series Name:** Gatareddihal (GRH) Pedon: R-7

**Location:** 15°14'20.8"N, 76°04'28.4" E Gudlanur village, Koppal Taluk and District

**Analysis at:** NBSS&LUP, Regional Centre, Bangalore.

**Classification:** Very fine, smectitic, isohyperthermic Sodic Haplusterts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-18	Ap	20.07	19.71	60.23	1.76	3.75	3.64	3.42	7.50	-	c	41.70	29.56
18-51	Bss1	15.11	17.47	67.42	3.16	3.04	2.25	3.38	3.27	-	c	59.43	38.52
51-80	Bss2	13.19	18.74	68.07	1.80	2.93	2.37	3.04	3.04	-	c	60.69	40.91
80-107	Bss3	17.54	19.50	62.96	2.46	4.13	3.24	4.25	3.46	-	c	57.25	37.31
107-131	BC	9.42	17.48	73.10	1.48	1.82	1.36	1.93	2.84	-	c	64.62	43.98

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO <sub>3</sub>	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP			
	Water	CaCl <sub>2</sub>	M KCl				dS m <sup>-1</sup>	%	%	Ca	Mg					K	Na	Total
										cmol kg <sup>-1</sup>						%	%	
0-18	9.08	-	-	0.23	0.33	6.89	-	-	0.70	6.36	-	63.21	1.05	100.00	7.11			
18-51	9.19	-	-	0.61	0.49	9.10	-	-	0.54	14.20	-	66.05	0.98	100.00	15.98			
51-80	9.27	-	-	0.56	0.29	9.36	-	-	0.49	14.75	-	65.63	0.96	100.00	17.07			
80-107	9.28	-	-	0.57	0.39	9.62	-	-	0.44	14.64	-	63.95	1.02	100.00	17.49			
107-131	9.04	-	-	1.08	0.31	8.32	-	-	0.52	16.40	-	68.36	0.94	100.00	17.30			

Contd...



**Series Name:** Handrala (HDL), **Pedon:** A2/RM-1

**Location:** 15°19'69.8"N, 75°58'00"E, Kavalura village, Koppal Taluk and District

**Analysis at:** NBSS&LUP, Regional Centre, Bangalore. **Classification:** Very fine, smectitic, isohyperthermic (calc) Typic Haplusterts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-25	Ap	21.68	16.62	61.70	4.42	3.98	3.43	5.64	4.20	10	c	41.36	31.27
25-50	Bss1	14.93	15.76	69.32	2.64	2.53	2.99	3.33	3.44	05	c	48.92	39.19
50-82	Bss2	23.11	16.60	60.29	4.51	3.61	6.31	4.74	3.95	05	c	42.46	33.85
82-117	Bss3	10.50	18.38	71.12	1.98	1.98	1.63	2.57	2.33	05	c	52.95	42.82

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO <sub>3</sub>	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP			
	Water	CaCl <sub>2</sub>	M KCl				dS m <sup>-1</sup>	%	%	Ca	Mg					K	Na	Total
										cmol kg <sup>-1</sup>						%	%	
0-25	9.06	-	-	0.371	0.16	4.80	-	-	0.80	7.93	-	62.33	1.01	-	5.09			
25-50	9.09	-	-	0.719	0.2	7.20	-	-	0.42	14.94	-	67.10	0.97	-	8.90			
50-82	9.28	-	-	0.47	0.19	9.36	-	-	0.47	11.59	-	60.21	1.00	-	7.70			
82-117	8.76	-	-	1.55	0.36	8.64	-	-	0.11	2.28	-	25.33	0.36	-	3.61			

Contd...

**Series Name:** Kadagathur (KDT)

**Pedon :** R-7

**Location:** 15<sup>0</sup>26'48"N, 76<sup>0</sup>09'51" E Budashettynala village, Koppal Taluk and District

**Analysis at:** NBSS&LUP, Regional Centre, Bangalore.

**Classification:** Fine, mixed, isohyperthermic Fluventic Haplustepts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-12	Ap	75.90	8.77	15.33	17.33	18.36	14.36	15.90	9.95	-	sl	10.66	5.33
12-37	A2	62.54	11.35	26.11	8.46	20.54	13.31	12.07	8.15	-	scl	15.61	8.22
37-71	Bw1	52.73	10.51	36.77	6.08	18.24	12.47	9.01	6.92	-	sc	19.66	11.21
71-93	Bw2	33.26	22.65	44.09	3.13	12.53	7.78	5.18	4.64	-	c	30.08	17.34
93-118	Bw3	31.01	24.57	44.42	2.04	10.41	8.26	6.01	4.29	-	c	34.92	18.16
118-170	Bw4	38.31	18.73	42.96	2.99	14.62	10.35	6.30	4.06	-	c	46.06	19.59

Depth (cm)	pH (1:2.5)			E.C. (1:2.5) dS m <sup>-1</sup>	O.C. %	CaCO <sub>3</sub> %	Exchangeable bases					CEC	CEC/Clay	Base saturation %	ESP %
	Water	CaCl <sub>2</sub>	M KCl				Ca	Mg	K	Na	Total				
0-12	6.95	-	-	0.17	1.28	0.39	9.17	2.76	0.10	0.08	12.11	12.10	0.79	100.09	0.65
12-37	7.55	-	-	0.17	0.40	0.40	8.36	4.51	0.08	0.40	13.35	13.30	0.51	100.37	3.02
37-71	7.60	-	-	0.21	0.44	0.39	10.67	8.19	0.10	0.74	19.70	19.10	0.52	103.12	3.88
71-93	8.26	-	-	0.28	0.72	1.56	14.97	12.13	0.12	3.07	30.29	29.40	0.67	103.01	10.45
93-118	8.44	-	-	0.58	0.68	1.17	13.32	10.77	0.13	4.76	28.98	28.50	0.64	101.68	12.40
118-170	9.06	-	-	0.64	0.44	1.17	8.92	8.14	0.23	12.32	29.61	28.60	0.67	103.53	37.27

Contd...

**Series Name:** Murlapur (MLR), **Pedon:** R-A1/16

**Location:** 15°19'42.9"N, 75°55'84.7"E, Kavalura village, Koppal Taluk and District

**Analysis at:** NBSS&LUP, Regional Centre, Bangalore. **Classification:** Very fine, smectitic, isohyperthermic (calc) Typic Haplusterts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-30	Ap	27.97	13.96	58.07	4.22	4.77	6.66	8.10	4.22	10	c	36.24	25.90
30-53	BA	26.34	17.48	56.17	4.17	5.05	6.04	7.24	3.84	05	c	38.55	28.98
53-83	Bss1	19.35	19.55	61.10	3.13	3.91	4.03	5.48	2.80	05	c	44.48	33.69
83-105	Bss2	16.63	17.47	65.90	2.70	3.93	2.92	3.93	3.15	<5	c	50.55	38.11
105-160	Bss3	14.69	20.34	64.97	0.79	2.26	4.07	4.18	3.39	<5	c	51.54	40.19

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO <sub>3</sub>	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP						
	Water	CaCl <sub>2</sub>	M KCl				dS m <sup>-1</sup>	%	%	Ca	Mg					K	Na	Total	cmol kg <sup>-1</sup>	%	%
0-30	9.19	-	-	0.313	0.57	10.08	-	-	0.64	5.67	-	42.08	0.72	-	5.39						
30-53	9.22	-	-	0.449	0.24	13.08	-	-	0.35	8.23	-	41.02	0.73	-	8.02						
53-83	9.17	-	-	0.377	0.82	16.92	-	-	0.39	14.28	-	51.20	0.84	-	11.16						
83-105	9.18	-	-	0.477	0.61	15.48	-	-	0.35	13.19	-	53.11	0.81	-	9.94						
105-160	9.01	-	-	1.17	0.24	16.92	-	-	0.43	19.61	-	53.95	0.83	-	14.54						

Contd...

**Series Name:** Bardur (BDR), **Pedon:** R-4

**Location:** 15°14'31.7"N, 76°01'19.1"E, Moranali village, Koppal Taluk and District

**Analysis at:** NBSS&LUP, Regional Centre, Bangalore. **Classification:** Very fine, smectitic, isohyperthermic (calc) Typic Haplusterts.

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-25	Ap	21.78	22.78	55.44	2.17	3.68	4.44	6.61	4.88	-	c	36.78	26.95
25-53	BA	18.62	18.56	62.82	2.23	4.24	3.46	5.24	3.46	-	c	41.25	29.87
53-90	Bss1	15.87	18.60	65.53	2.23	1.34	4.25	3.91	4.13	-	c	44.73	33.64
90-126	Bss2	13.66	20.02	66.32	1.68	2.80	2.35	3.70	3.14	-	c	49.24	38.37
126-152	Bss3	11.64	20.79	67.57	1.69	1.81	1.81	3.50	2.82	-	c	53.50	41.90
152-210	Bss4	11.38	22.78	65.42	2.16	2.16	1.93	3.07	2.05	-	c	51.53	39.64

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO <sub>3</sub>	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP
	Water	CaCl <sub>2</sub>	M KCl				Ca	Mg	K	Na	Total				
				dS m <sup>-1</sup>	%	%	cmol kg <sup>-1</sup>					%	%		
0-25	8.73	-	22.78	0.203	0.24	5.76	-	-	0.65	4.43	-	40.56	0.73	-	4.37
25-53	9.17	-	18.56	0.295	0.45	4.92	-	-	0.32	10.47	-	74.70	1.19	-	5.61
53-90	9.27	-	18.60	0.388	0.66	6.00	-	-	0.24	10.49	-	76.20	1.16	-	5.51
90-126	9.22	-	20.02	0.608	0.57	5.88	-	-	0.21	15.93	-	77.20	1.16	-	8.25
126-152	9.21	-	20.79	0.936	0.33	6.60	-	-	0.37	20.88	-	80.90	1.20	-	10.32
152-210	9.03	-	23.21	1.47	0.33	8.16	-	-	0.24	15.34	-	73.10	1.12	-	8.39

## INTERPRETATION FOR LAND RESOURCE MANAGEMENT

The most important soil and site characteristics that affect the land use and conservation needs of an area are land capability, land irrigability, soil depth, soil texture, coarse fragments, available water capacity, soil slope, soil erosion, soil reaction etc. These are interpreted from the data base generated through land resource inventory and several thematic maps are generated. These would help in identifying the areas suitable for growing crops and, soil and water conservation measures and structures needed thus helping to maintain good soil health for sustained crop production. The various thematic maps generated are described below.

### 5.1 Land Capability Classification

Land capability classification is an interpretative grouping of soil map units (soil phases) mainly based on inherent soil characteristics, external land features and environmental factors that limit the use of land for agriculture, pasture, forestry or other uses on a sustained basis (IARI, 1971). The land and soil characteristics used to group the land resources in an area into various land capability classes, subclasses and units are

*Soil characteristics:* Soil depth, soil texture, coarse fragments, soil reaction, available water capacity, calcareousness, salinity/alkali *etc.*

*Land characteristics:* Slope, erosion, drainage, rock outcrops.

*Climate:* Total rainfall and its distribution, and length of crop growing period.

The land capability classification system is divided into land capability classes, subclasses and units based on the level of information available. Eight land capability classes are recognized. They are

*Class I:* They are very good lands that have no limitations or very few limitations that restrict their use.

*Class II:* They are good lands that have minor limitations and require moderate conservation practices.

*Class III:* They are moderately good lands that have severe limitations that reduce the choice of crops or that require special conservation practices.

*Class IV:* They are fairly good lands that have very severe limitations that reduce the choice of crops or that require very careful management.

*Class V:* Soils in these lands are not likely to erode, but have other limitations like wetness that are impractical to remove and as such not suitable for agriculture, but suitable for pasture or forestry with minor limitations.

*Class VI:* The lands have severe limitations that make them generally unsuitable for cultivation, but suitable for pasture or forestry with moderate limitations.

*Class VII:* The lands have very severe limitations that make them unsuitable for cultivation, but suitable for pasture or forestry with major limitations.

*Class VIII:* Soil and other miscellaneous areas (rock lands) that have very severe limitations that nearly preclude their use for any crop production, but suitable for wildlife, recreation and installation of wind mills.

The land capability subclasses are recognised based on the dominant limitations observed within a given land capability class. The subclasses are designated by adding a lower case letter like ‘e’, ‘w’, ‘s’, or ‘c’ to the class numeral. The subclass “e” indicates that the main hazard is risk of erosion, “w” indicates drainage or wetness as a limitation for plant growth, “s” indicates shallow soil depth, coarse or heavy textures, calcareousness, salinity/alkalinity or graveliness and “c” indicates limitation due to climate.

The land capability subclasses have been further subdivided into land capability units based on the kinds of limitations present in each subclass. Ten land capability units are used in grouping the soil map units. They are stony or rocky (0), erosion hazard (slope, erosion) (1), coarse texture (sand, loamy sand, sandy loam) (2), fine texture (cracking clay, silty clay) (3), slowly permeable subsoil (4), coarse underlying material (5), salinity/alkali (6), stagnation, overflow, high ground water table (7), soil depth (8) and fertility problems (9). The capability units thus identified have similar soil and land characteristics that respond similarly to a given level of management. The soils of the microwatershed have been classified upto land capability subclass level.

The 18 soil map units identified in the Handrahalu-2 Microwatershed are grouped under three land capability classes and four land capability subclasses (Fig. 5.1).




Fig. 5.1 Land Capability map of Handrahalu-2 Microwatershed

Entire area of the microwatershed is suitable for agriculture. An area of 393 ha (85%) are good lands (Class II) that have minor limitations and require moderate conservation practices and are distributed in all parts of the microwatershed. Moderately good lands (Class III) cover an area of 43 ha (9%) and are distributed in the northern and northeastern part of the microwatershed with moderate problems of soil that require special conservation practices. An area of 8 ha (2%) is fairly good lands (Class IV) and are distributed in the western part of the microwatershed. The other miscellaneous areas cover about 4 per cent that have very severe limitations that preclude them for any crop productivity, but well suited for wildlife, recreation and installation of wind mills.

## **5.2 Soil Depth**

Soil depth refers to the depth of the soil occurring above the parent material or hard rock. The depth of the soil determines the effective rooting depth for plants and in accordance with soil texture, mineralogy and gravel content, the capacity of the soil column to hold water and nutrient availability. Soil depth is one of the most important soil characteristic that is used in differentiating soils into different soil series. The soil depth classes used in identifying soils in the field are very shallow (<25 cm), shallow (25-50 cm), moderately shallow (50-75 cm), moderately deep (75-100 cm), deep (100-150 cm) and very deep (>150 cm). They were used to classify the soils into different depth classes and a soil depth map was generated (Fig. 5.2).

An area of 8 ha (2%) is very shallow (<25 cm) and are distributed in the western part of the microwatershed. Shallow (25-50 cm) soils occupy an area of 43 ha (9%) and are distributed in the northern and northeastern part of the microwatershed. An area of 57 ha (12%) is moderately shallow (50-75 cm) and are distributed in the western part of the microwatershed. Moderately deep soils (75-100 cm) occupy a maximum area of 177 ha (38%) and occur in the major part of the microwatershed. Deep (100-150 cm) to very deep (>150 cm) soils occupy an area of 159 ha (34%) and are distributed in the northwestern, central, eastern and southern part of the microwatershed.

The most problem lands with an area of about 8 ha (2%) having very shallow (<25 cm) rooting depth. They are suitable for growing short duration agricultural crops but well suited for pasture, forestry or other recreational purposes. The most productive lands cover a minor area about 159 ha (34%) where all climatically adapted long duration crops be grown.




Fig. 5.2 Soil Depth map of Handrahalu-2 Microwatershed

### 5.3 Surface Soil Texture

Texture is an expression to indicate the coarseness or fineness of the soil as determined by the relative proportion of primary particles of sand, silt and clay. It has a direct bearing on the structure, porosity, adhesion and consistence. The surface layer of a soil to a depth of about 25 cm is the layer that is most used by crops and plants. The surface soil textural class provides a guide to understanding soil-water retention and availability, nutrient holding capacity, infiltration, workability, drainage, physical and chemical behaviour, microbial activity and crop suitability. The textural classes used for LRI were used to classify and a surface soil texture map was generated (fig. 5.3). The area extent and their spatial distribution in the microwatershed is shown in figure 5.3.

An entire area of about 444 ha (96%) has soils that are clayey soils at the surface and are distributed in all parts of the microwatershed (Fig. 5.3).

The most productive lands 444 ha (96%) with respect to surface soil texture are the clayey soils that have high potential for soil-water retention and availability, and nutrient retention and availability, but have more problems of drainage, infiltration, workability and other physical problems.






Fig. 5.3 Surface Soil Texture map of Handrahalu-2 Microwatershed

#### 5.4 Soil Gravelliness

Gravel is the term used for describing coarse fragments between 2 mm and 7.5 cm diameter and stones for those between 7.5 cm and 25 cm. The presence of gravel and stones in soil reduces the volume of soil responsible for moisture and nutrient storage, drainage, infiltration and runoff, and hinders plant growth by impeding root growth and seedling emergence, intercultural operations and farm mechanization. The gravelliness classes used in LRI were used to classify the soils and using these classes a gravelliness map was generated. The area extent and their spatial distribution in the microwatershed is given in figure 5.4.

The soils that are non-gravelly (<15% gravel) cover a maximum area of 228 ha (49%) and are distributed in the major part of the microwatershed. An area of 204 ha (44%) is covered by gravelly (15-35% gravel) soils and are distributed in the northern, eastern, western and central part of the microwatershed. An area of about 13 ha (3%) is very gravelly (35-60%) and are distributed in the eastern and western part of the microwatershed (Fig. 5.4).

The most productive lands with respect to gravelliness are found to be 49%. They are non-gravelly with less than 15 per cent gravel and have potential for growing both annual and perennial crops. The problem soils that are gravelly (15-35%) to very gravelly (35-60%) cover 217 ha (47%) where only short or medium duration crops can be grown.




Fig. 5.4 Soil Gravelliness map of Handrahalu-2 Microwatershed

### 5.5 Available Water Capacity

The soil available water capacity (AWC) is estimated based on the ability of the soil column to retain water between the tensions of 0.33 and 15 bar in a depth of 100 cm or the entire solum if the soil is shallower. The AWC of the soils (soil series) as estimated by considering the soil texture, mineralogy, soil depth and gravel content (Sehgal *et al.*, 1990) and accordingly the soil map units were grouped into five AWC classes *viz*, very low (<50 mm/m), low (50-100 mm/m), medium (100-150 mm/m), high (150-200 mm/m) and very high (>200 mm/m) and using these values, an AWC map was generated (Fig. 5.5), showing the area extent and their spatial distribution in the microwatershed.

An area of about 8 ha (2%) are very low (<50 mm/m) in available water capacity and are distributed in the western part of the microwatershed. An area of about 100 ha (22%) has soils that are low (51-100 mm/m) in available water capacity and are distributed in the northern and western part of the microwatershed. Soils with medium available water capacity (101-150 mm/m) occupy a maximum area of 177 ha (38%) and are distributed in the major part of the microwatershed. An area of about 159 ha (34%) is very high (>200 mm/m) in available water capacity and are distributed in the eastern, northwestern, central and southern part of the microwatershed.

An area of about 8 ha (2%) in the microwatershed has soils that are problematic with regard to available water capacity. Here, only short duration crops can be grown and the probability of crop failure is very high. These areas are best put to other alternative

uses. The potential soils with respect to AWC cover about 159 ha (34%) that have very high AWC, where all climatically adapted long duration crops can be grown.




Fig. 5.5 Soil Available Water Capacity map of Handrahalu-2 Microwatershed

## 5.6 Soil Slope

Soil slope refers to the inclination of the surface of the land. It is defined by gradient, shape and length, and is an integral feature of any soil as a natural body. Slope is considered important in soil genesis, land use and land development. The length and gradient of slope influences the rate of runoff, infiltration, erosion and deposition. The soil map units were grouped into four slope classes and a slope map was generated showing the area extent and their geographic distribution of different slope classes in the microwatershed (Fig. 5.6).

An area of 31 ha (7%) is nearly level (0-1% slope) and are distributed in the southern part of the microwatershed. Maximum area of about 413 ha (90%) falls under very gently sloping (1-3% slope) and are distributed in all parts of the microwatershed.




Fig. 5.6 Soil Slope map of Handrahalu-2 Microwatershed

## 5.7 Soil Erosion

Soil erosion refers to the wearing away of the earth's surface by the forces of water, wind and ice involving detachment and transport of soil by raindrop impact. It is used for accelerated soil erosion resulting from disturbance of the natural landscape by burning, excessive grazing and indiscriminate felling of forest trees and tillage, all usually by man. The erosion classes showing an estimate of the current erosion status as judged from field observations in the form of rills, gullies or a carpet of gravel on the surface are recorded. Four erosion classes, viz, slight erosion (e1), moderate erosion (e2), severe erosion (e3) and very severe erosion (e4) are recognized. The soil map units were grouped into different erosion classes and a soil erosion map generated. The area extent and their spatial distribution in the microwatershed is given in Figure 5.7.

Soils that are slightly eroded (e1 Class) occupy an area of about 200 ha (43%) and are distributed in the northern, eastern, western and southern part of the microwatershed. Moderately eroded (e2 Class) soils cover a maximum area of 244 ha (53%) and are distributed in the major part of the microwatershed.

An area of about 200 ha (43%) in the microwatershed is problematic because of moderate erosion. These areas need soil and water conservation and other land development measures for restoring the soil health.




Fig. 5.7 Soil Erosion map of Handrahalu-2 Microwatershed



## **FERTILITY STATUS**

Soil fertility plays an important role in increasing crop yield. The adoption of high yielding varieties that require high amounts of nutrients has resulted in deficiency symptoms in crops and plants due to imbalanced fertilization and poor inherent fertility status, as these areas are characterised by low rainfall and high temperatures. Hence, it is necessary to know the fertility (macro and micro nutrients) status of the soils of the watersheds for assessing the kind and amount of fertilizers required for each of the crop intended to be grown. For this purpose, the surface soil samples collected from the grid points (one soil sample at every 320 m grid interval) all over the microwatershed through land resource inventory in the year 2017 were analysed for pH, EC, organic carbon, available phosphorus and potassium and for micronutrients like zinc, boron, copper, iron and manganese, and secondary nutrient sulphur.

Soil fertility data generated has been assessed and individual maps for all the nutrients for the microwatershed have been generated by using the Kriging method under GIS. The village/survey number wise fertility data for the microwatershed is given in Appendix-II.

### **6.1 Soil Reaction (pH)**

The soil analysis of the Handrahalu-2 Microwatershed for soil reaction (pH) showed that an area of 8 ha (2%) is moderately alkaline (pH 7.8-8.4) and are distributed in the southwestern part of the microwatershed. Strongly alkaline (pH 8.4-9.0) soils cover an area of 322 ha (70%) and are distributed in the major part of the microwatershed. An area of 115 ha (25%) is very strongly alkaline (pH >9.0) and are distributed in the eastern and northern part of the microwatershed. Thus, major soils in the microwatershed are alkaline in reaction covering 444 ha.

### **6.2 Electrical Conductivity (EC)**

The Electrical Conductivity of the soils is <2 dS m<sup>-1</sup> in the entire microwatershed (Fig. 6.2) area and as such the soils are nonsaline.

### **6.3 Organic Carbon**

The soil organic carbon content (an index of available Nitrogen) in the soils of the microwatershed is low (<0.5%) covering an area of 93 ha (20%) and are distributed in the northern part of the microwatershed. Medium (0.5-0.75%) in organic carbon covering an area of 349 ha (76%) and is distributed in the major part of the microwatershed (Fig. 6.3). An area of 2 ha (<1%) is high (>0.75%) and are distributed in the southern part of the microwatershed.




Fig. 6.1 Soil Reaction (pH) map of Handrahalu-2 Microwatershed




Fig. 6.2 Electrical Conductivity (EC) map of Handrahalu-2 Microwatershed






Fig. 6.3 Soil Organic Carbon map of Handrahalu-2 Microwatershed




Fig. 6.4 Soil Available Phosphorus map of Handrahalu-2 Microwatershed

#### **6.4 Available Phosphorus**

An area of 105 ha (23%) is low (<23 kg/ha) and are distributed in the southern and western part of the microwatershed. Maximum cultivated area of about 340 ha (74%) is medium (<23 kg/ha) in available phosphorus and is distributed in all parts of the microwatershed (Fig. 6.4).

#### **6.5 Available Potassium**

An area of about 8 ha (2%) is medium (145-337 kg/ha) in available potassium and are distributed in the northern part of the microwatershed. Maximum area of 437 ha (95%) is high (>337 kg/ha) and are distributed in all parts of the microwatershed (Fig. 6.5).

#### **6.6 Available Sulphur**

Soils that are medium in available sulphur content (10-20 ppm) cover an area of 96 ha (21%) and are distributed in the southern, western and southeastern part of the microwatershed (Fig. 6.6). The areas that are medium in available sulphur need to be applied with magnesium sulphate or gypsum or factomphos (p) fertilizer (13% sulphur) for 2-3 years for the deficiency to be corrected. It is high (>20 ppm) in a maximum area of 348 ha (75%) and are distributed in the major part of the microwatershed.

#### **6.7 Available Boron**

Available boron content is low (<0.5 ppm) in a maximum area of 303 ha (66%) and are distributed in the major part of the microwatershed. An area of about 141 ha (31%) is medium (0.5-1.0 ppm) in available boron and are distributed in the northeastern, central and southehrn part of the microwatershed (Fig. 6.7). It is high (>1.0 ppm) in a minor area of (<1%) in the microwatershed.

#### **6.8 Available Iron**

Available iron content is sufficient (>4.5 ppm) in an entire area of 444 ha (96%) of the microwatershed (Fig. 6.8).

#### **6.9 Available Manganese**

Available manganese content is sufficient (>1.0 ppm) in the entire microwatershed area (Fig. 6.9).

#### **6.10 Available Copper**

Available copper content is sufficient (>0.2 ppm) in the entire microwatershed area (Fig. 6.10).




Fig. 6.5 Soil Available Potassium map of Handrahalu-2 Microwatershed




Fig. 6.6 Soil Available Sulphur map of Handrahalu-2 Microwatershed




Fig. 6.7 Soil Available Boron map of Handrahalu-2 Microwatershed




Fig. 6.8 Soil Available Iron map of Handrahalu-2 Microwatershed




Fig. 6.9 Soil Available Manganese map of Handrahalu-2 Microwatershed




Fig. 6.10 Soil Available Copper map of Handrahalu-2 Microwatershed

## 6.11 Available Zinc

Available zinc content is sufficient ( $>0.6$  ppm) in an area of 20 ha (4%) and are distributed in the southern part of the microwatershed. Maximum area of 424 ha (92%) is deficient ( $<0.6$  ppm) and are distributed in the major part of the microwatershed (Fig. 6.11).




Fig. 6.11 Soil Available Zinc map of Handrahalu-2 Microwatershed

## LAND SUITABILITY FOR MAJOR CROPS

The soil and land resource units (soil phases) of Handrahalu-2 Microwatershed were assessed for their suitability for growing food, fodder, fibre and other horticulture crops by following the procedure as outlined in FAO, 1976 and 1983. Crop requirements were developed for each of the crop from the available research data and also by referring to Naidu *et. al.* (2006) and Natarajan *et. al* (2015). The crop requirements (Table 7.2 to 7.33) were matched with the soil and land characteristics (Table 7.1) to arrive at the crop suitability. In FAO land suitability classification, two orders are recognized. Order S-Suitable and Order N-Not suitable. The orders have classes, subclasses and units. Order S has three classes, Class S1-Highly Suitable, Class S2-Moderately Suitable and Class S3-Marginally Suitable. Order N has two Classes, N1-Currently not Suitable and N2-Permanently not Suitable. There are no subclasses within the Class S1 as they will have very minor or no limitations for crop growth. Classes S2, S3, N1 and N2 are divided into subclasses based on the kinds of limitations encountered. The limitations that affect crop production are 'c' for erratic rainfall and its distribution and length of growing period (LGP), 'e' for erosion hazard, 'r' for rooting condition, 't' for lighter or heavy texture, 'g' for gravelliness or stoniness, 'n' for nutrient availability, 'l' for topography, 'm' for moisture availability, 'z' for calcareousness 's' for sodium and 'w' for drainage. These limitations are indicated as lower case letters to the class symbol. For example, moderately suitable lands with the limitations of soil depth and erosion are designated as S2re. For the microwatershed, the soil mapping units were evaluated and classified up to subclass level.

Using the above criteria, the soil map units of the microwatershed were evaluated and land suitability maps for 31 major agricultural and horticultural crops were generated. The detailed information on the kind of suitability of each of the soil phase for the crops assessed are given village/ survey number wise for the microwatershed in Appendix-III.

### 7.1 Land Suitability for Sorghum (*Sorghum bicolor*)

Sorghum is one of the major food crop grown in Karnataka in an area of 10.47 lakh ha in Bijapur, Gulbarga, Raichur, Bidar, Belgaum, Dharwad, Bellary, Chitradurga, Mysore and Chamarajnagar districts. The crop requirements for growing sorghum (Table 7.2) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing sorghum was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure. 7.1.

Maximum area of 212 ha (46%) is highly suitable (Class S1) lands for growing sorghum and are distributed in the major part of the microwatershed. An area of 176 ha (38%) is moderately suitable (Class S2) and are distributed in the northern, eastern, northwestern and southern part of the microwatershed. They have minor limitations of

gravelliness, calcareousness, nutrient availability and rooting condition. An area of about 48 ha (10%) is marginally suitable (Class S3) for growing sorghum and are distributed in the northern and northeastern part of the microwatershed with moderate limitations of gravelliness, calcareousness and rooting condition. An area of 8 ha (2%) is currently not suitable (Class N1) and are distributed in the western part of the microwatershed with severe limitations of rooting condition and gravelliness.




Fig. 7.1 Land Suitability map of Sorghum

## 7.2 Land Suitability for Maize (*Zea mays*)

Maize is one of the most important food crop grown in an area of 13.37 lakh ha in almost all the districts of the State. The crop requirements for growing maize (Table 7.3) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing maize was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.2.

There are no highly suitable (Class S1) lands for growing maize in the microwatershed. Maximum area of 393 ha (85%) is moderately suitable (Class S2) for growing maize and are distributed in the major part of the microwatershed with minor limitations of rooting condition and texture. Marginally suitable (Class S3) lands cover an area of 43 ha (9%) and are distributed in the northern and northeastern part of the microwatershed. They have moderate limitations of calcareousness and texture. Currently suitable lands cover an area of 8 ha (2%) and are distributed in the western part of the microwatershed with severe limitations of rooting condition and gravelliness.






Fig. 7.2 Land Suitability map of Maize

### 7.3 Land Suitability for Bajra (*Pennisetum glaucum*)

Bajra is one of the major food crop grown in an area of 2.34 lakh ha in the northern districts of the Karnataka State. The crop requirements for growing bajra (Table 7.4) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing bajra was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.3.

There are no highly suitable (Class S1) lands for growing bajra and in the microwatershed. Maximum area of 388 ha (84%) is moderately suitable (Class S2) and are distributed in the major part of the microwatershed with minor limitations of texture and calcareousness. Marginally suitable (Class S3) lands cover an area of 48 ha (10%) and are distributed in the northern and northeastern part of the microwatershed. They have moderate limitations of gravelliness, texture, calcareousness and rooting depth. Currently suitable lands cover an area of 8 ha (2%) and are distributed in the western part of the microwatershed with severe limitations of rooting condition and gravelliness.




Fig. 7.3 Land Suitability map of Bajra

#### 7.4 Land Suitability for Groundnut (*Arachis hypogaea*)

Groundnut is one of the major oilseed crop grown in an area of 6.54 lakh ha in Karnataka in most of the districts either as rainfed or irrigated crop. The crop requirements for growing groundnut (Table 7.5) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing groundnut was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.4.

There are no highly (Class S1) and moderately (Class S2) suitable lands for growing groundnut in the microwatershed. Maximum area of 436 ha (94%) is marginally suitable (Class S3) for groundnut and are distributed in the major part of the microwatershed. They have moderate limitations of gravelliness, calcareousness and texture. An area of 8 ha (2%) is currently nor suitable (Class N1) for growing groundnut and are distributed in the western part of the microwatershed with severe limitations of rooting condition and gravelliness.




Fig. 7.4 Land Suitability map of Groundnut

### 7.5 Land Suitability for Sunflower (*Helianthus annuus*)

Sunflower is one of the most important oilseed crop grown in an area of 3.56 lakh ha in the State in all the districts. The crop requirements for growing sunflower (Table 7.6) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing sunflower was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.5.

An area of 51 ha (11%) is highly suitable (Class S1) lands for growing sunflower and are distributed in the western and central part of the microwatershed. Maximum area of 280 ha (61%) is moderately suitable (Class S2) and are distributed in the major part of the microwatershed. They have minor limitations of gravelliness, calcareousness and rooting condition. An area of 62 ha (13%) is marginally suitable (Class S3) for growing sunflower and distributed in the northern, western and northeastern part of the microwatershed with moderate limitations of rooting condition, calcareousness and gravelliness. Currently not suitable (Class N1) lands cover an area of 51 ha (11%) and are distributed in the northern, northeastern and western part of the microwatershed with severe limitations of rooting condition, calcareousness and gravelliness.




Fig. 7.5 Land Suitability map of Sunflower

## 7.6 Land Suitability for Red gram (*Cajanus cajan*)

Redgram is one of the most important pulse crop grown in an area of 7.28 lakh ha in almost all the districts of the State. The crop requirements for growing redgram (Table 7.7) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing redgram was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.6.

There are no highly suitable (Class S1) lands growing redgram in the microwatershed. Moderately suitable (Class S2) lands occupy a maximum area of 315 ha (68%) and are distributed in the major part of the microwatershed with minor limitations of gravelliness, texture and calcareousness. Marginally suitable (Class S3) lands cover an area of 79 ha (17%) and are distributed in the northern, western and eastern part of the microwatershed. They have moderate limitations of gravelliness, calcareousness, texture and rooting condition. Currently not suitable (Class N1) lands cover an area of 51 ha (11%) for growing redgram and are distributed in the western and northeastern part of the microwatershed with severe limitations of gravelliness, calcareousness and rooting condition.




Fig. 7.6 Land Suitability map of Redgram

### 7.7 Land Suitability for Bengalgram (*Cicer arietinum*)

Bengalgram is one of the major pulse crop grown in an area of 9.39 lakh ha in northern Karnataka in Bijapur, Gulbarga, Raichur, Bidar, Belgaum, Dharwad and Bellary districts. The crop requirements for growing Bengalgram (Table 7.8) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing Bengalgram was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.7.

Maximum area of 230 ha (50%) is highly suitable (Class S1) lands available for growing bengalgram and are distributed in the major part of the microwatershed. Moderately suitable lands (Class S2) occupy an area of 159 ha (34%) and are distributed in the northern, northwestern, western, southern and eastern part of the microwatershed with minor limitations of gravelliness, rooting condition and calcareousness. Marginally suitable (Class S3) lands cover an area of 48 ha (10%) and are distributed in the northern and northeastern part of the microwatershed. They have moderate limitations of rooting condition, texture, calcareousness and gravelliness. An area of 8 ha (2%) is currently not suitable (Class N1) and are distributed in the western part of the microwatershed with severe limitations of rooting condition and gravelliness.




Fig. 7.7 Land Suitability map of Bengalgram

### 7.8 Land Suitability for Cotton (*Gossypium hirsutum*)

Cotton is one of the most important fibre crop grown in the State in about 8.75 lakh ha area in Raichur, Dharwad, Belgaum, Gulbarga, Bijapur, Bidar, Bellary, Chitradurga and Chamarajnagar districts. The crop requirements for growing cotton (Table 7.9) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing cotton was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.8.

Maximum area of 230 ha (50%) is highly suitable (Class S1) lands for growing cotton and are distributed in the major part of the microwatershed. Moderately suitable (Class S2) lands occupy an area of 159 ha (34%) and are distributed in the northern, eastern, northwestern, western and southern part of the microwatershed. They have minor limitations of rooting condition, gravelliness and calcareousness. Marginally suitable (Class S3) lands cover an area of 48 ha (10%) and are distributed in the northeastern and northern part of the microwatershed. They have moderate limitations of gravelliness, calcareousness, texture and rooting condition. Currently suitable land occur in an area of 8 ha (2%) and are distributed in the western part of the microwatershed with severe limitations of gravelliness and rooting condition.




Fig. 7.8 Land Suitability map of Cotton

### 7.9 Land Suitability for Chilli (*Capsicum annum L*)

Chilli is one of the most important commercial spice crop grown in an area of 0.89 lakh ha in all the districts of Karnataka State. The crop requirements for growing chilli (Table 7.10) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing chilli was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.9.

There are no highly (Class S1) and moderately (Class S2) suitable lands for growing chilli in the microwatershed. Marginally suitable (Class S3) lands cover a maximum area of about 436 ha (94%) and are distributed in all parts of the microwatershed. They have moderate limitations of gravelliness, texture, calcareousness and rooting condition. An area of 8 ha (2%) is currently not suitable (Class N1) and are distributed in the western part of the microwatershed with severe limitations of rooting condition and gravelliness.




Fig. 7.9 Land Suitability map of Chilli

### 7.10 Land Suitability for Tomato (*Solanum lycopersicum*)

Tomato is one of the most important vegetable crop grown in an area of 0.65 lakh ha in almost all the districts of the State. The crop requirements (Table 7.11) for growing tomato were matched with the soil-site characteristics (Table 7.11) and a land suitability map for growing tomato was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.10.

There are no highly (Class S1) and moderately (Class S2) suitable lands for growing tomato in the microwatershed. Marginally suitable (Class S3) lands occupy a maximum area of 436 ha (94%) and are distributed in all parts of the microwatershed with moderate limitations of gravelliness, rooting condition, texture and calcareousness. An area of 8 ha (2%) is currently not suitable (Class N1) and are distributed in the western part of the microwatershed with severe limitations of rooting condition and gravelliness.






Fig. 7.10 Land Suitability map of Tomato

### 7.11 Land Suitability for Brinjal (*Solanum melongena*)

Brinjal is one of the most important vegetable crop grown in all the districts. The crop requirements for growing brinjal (Table 7.12) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing brinjal was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.11.

There are no highly suitable (Class S1) for growing brinjal in the microwatershed. Maximum area of about 394 ha (85%) is moderately suitable (Class S2) for growing brinjal and are distributed in the major part of the microwatershed with minor limitations of texture, calcareousness, gravelliness and rooting condition. Marginally suitable lands (Class S3) occur in an area of 43 ha (9%) and are distributed in the northern and northeastern part of the microwatershed with moderate limitation of rooting condition. Currently suitable (Class N1) lands occur in an area of 8 ha (2%) and are distributed in the western part of the microwatershed with severe limitation of rooting condition.




Fig. 7.11 Land Suitability map of Brinjal

## 7.12 Land Suitability for Onion (*Allium cepa*)

Onion is one of the most important vegetable crop grown in Raichur, Dharwad, Belgaum, Gulbarga, Bijapur, Bidar, Bellary, Chitradurga and Tumakuru districts. The crop requirements for growing onion (Table 7.13) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing Onion was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.12.

There are no highly (Class S1) and moderately (Class S2) suitable for growing onion in the microwatershed. Marginally suitable lands (Class S3) for growing onion occupy an area of 437 ha (94%) and are distributed in all parts of the microwatershed with moderate limitations of rooting depth, gravelliness, calcareousness and texture. Currently suitable (Class N1) lands occur in an area of 8 ha (2%) and are distributed in the western part of the microwatershed with severe limitation of rooting condition.




Fig. 7.12 Land Suitability map of Onion

### 7.13 Land Suitability for Bhendi (*Abelmoschus esculentus*)

Bhendi is one of the most important vegetable crop grown in all the districts. The crop requirements for growing bhendi (Table 7.14) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing bhendi was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.13.

There are no highly suitable (Class S1) for growing bhendi in the microwatershed. Maximum area of about 394 ha (85%) is moderately suitable (Class S2) and are distributed in the major part of the microwatershed with minor limitations of texture, calcareousness, gravelliness and rooting depth. Marginally suitable lands (Class S3) occur in an area of 43 ha (9%) and are distributed in the northern and northeastern part of the microwatershed with moderate limitation of rooting condition. An area of 8 ha (2%) is currently not suitable (Class N1) and are distributed in the western part of the microwatershed with severe limitations of rooting condition.




Fig. 7.13 Land Suitability map of Bhendi

#### 7.14 Land Suitability for Drumstick (*Moringa oleifera*)

Drumstick is one of the most important vegetable crop grown in 2403 ha area in the State. The crop requirements for growing drumstick (Table 7.15) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing drumstick was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.14.

There are no highly suitable (Class S1) lands for growing drumstick in the microwatershed. Maximum area of 336 ha (73%) is moderately suitable (Class S2) and are distributed in the major part of the microwatershed. They have minor limitations of gravelliness, texture, rooting condition and calcareousness. Marginally suitable (Class S3) lands cover an area of 57 ha (12%) and are distributed in the western part of the microwatershed. They have moderate limitations of calcareousness and rooting condition. Currently not suitable (Class N1) lands cover an area of 51 ha (11%) and are distributed in the northern, western and northeastern part of the microwatershed with severe limitations of rooting condition, calcareousness and gravelliness.




Fig. 7.14 Land Suitability map of Drumstick

### 7.15 Land Suitability for Mango (*Mangifera indica*)

Mango is one of the most important fruit crop grown in about 1.73 lakh ha in almost all the districts of the State. The crop requirements (Table 7.16) for growing mango were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing mango was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.15.

There are no highly (Class S1) and moderately (Class S2) suitable lands for growing mango in the microwatershed. Marginally suitable (Class S3) lands cover a maximum area of 336 ha (73%) and are distributed in the major part of the microwatershed. They have moderate limitations of gravelliness, rooting condition, texture and calcareousness. An area of 108 ha (23%) is currently not suitable (Class N1) for growing mango and occur in the northern, western and northeastern part of the microwatershed with severe limitations of gravelliness, texture, calcareousness and rooting condition.




Fig. 7.15 Land Suitability map of Mango

### 7.16 Land suitability for Guava (*Psidium guajava*)

Guava is one of the most important fruit crop grown in an area of about 0.64 lakh ha in almost all the districts of the State. The crop requirements (Table 7.17) for growing guava were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing guava was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.16.

There are no highly (Class S1) suitable and moderately (Class S2) suitable lands for growing guava in the microwatershed. Marginally suitable (Class S3) lands cover a maximum area of 393 ha (85%) and are distributed in all parts of the microwatershed. They have moderate limitations of gravelliness, texture and calcareousness. An area of about 51 ha (11%) area is currently not suitable (Class N1) for growing guava and occur in the northern, western and northeastern part of the microwatershed with severe limitations of rooting condition, texture and gravelliness.




Fig. 7.16 Land Suitability map of Guava

### 7.17 Land Suitability for Sapota (*Manilkara zapota*)

Sapota is one of the most important fruit crop grown in an area of about 29373 ha in almost all the districts of the State. The crop requirements (Table 7.18) for growing sapota were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing sapota was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.17.

There are no highly (Class S1) and moderately (Class S2) suitable lands for growing sapota in the microwatershed. Maximum area of about 393 ha (85%) is marginally suitable (Class S3) and are distributed in the major part of the microwatershed. They have minor limitations of gravelliness, texture, calcareousness and rooting condition. An area of 51 ha (11%) is currently not suitable (Class N1) for growing sapota and occur in the northern, western and northeastern part of the microwatershed with severe limitations of gravelliness, calcareousness and rooting condition.




Fig. 7.17 Land Suitability map of Sapota

### 7.18 Land Suitability for Pomegranate (*Punica granatum*)

Pomegranate is one of the commercially grown fruit crop in about 18488 ha in Karnataka mainly in Bijapur, Bagalkot, Koppal, Gadag and Chitradurga districts. The crop requirements for growing pomegranate (Table 7.19) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing pomegranate was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.18.

There are no highly suitable (Class S1) lands for growing pomegranate in the microwatershed. Moderately suitable (Class S2) lands occupy a maximum area of 336 ha (73%) and are distributed in the major part of the microwatershed. They have minor limitations of texture, rooting condition, gravelliness and calcareousness. An area of 57 ha (12%) is marginally suitable (Class S3) for growing pomegranate and are distributed in the northern and western part of the microwatershed. They have moderate limitations of rooting condition and calcareousness. An area of 51 ha (11%) is currently not suitable (Class N1) for growing pomegranate and are distributed in the northern, northeastern and western part of the microwatershed with severe limitations of gravelliness, calcareousness and rooting condition.






Fig. 7.18 Land Suitability map of Pomegranate

### 7.19 Land Suitability for Musambi (*Citrus limetta*)

Musambi is one of the most important fruit crop grown in almost all the districts of the State. The crop requirements for growing musambi (Table 7.20) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing musambi was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.19.

An area of 51 ha (11%) is highly suitable (Class S1) lands for growing musambi and are distributed in the western and central part of the microwatershed. Maximum area of 285 ha (62%) is moderately suitable (Class S2) and are distributed in the major part of the microwatershed. They have minor limitations of calcareousness, rooting condition and gravelliness. Marginally suitable (Class S3) lands occur in an area of 57 ha (12%) and are distributed in the northern and western part of the microwatershed with moderate limitations of rooting condition and gravelliness. An area of 51 ha (11%) is currently not suitable (Class N1) for growing musambi and are distributed in the northeastern and western part of the microwatershed. They have severe limitations of gravelliness, calcareousness and rooting condition.




Fig. 7.19 Land Suitability map of Musambi

## 7.20 Land Suitability for Lime (*Citrus sp*)

Lime is one of the most important fruit crop grown in an area of 0.11 lakh ha in almost all the districts of the State. The crop requirements for growing lime (Table 7.21) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing lime was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.20.

An area of 51 ha (11%) is highly suitable (Class S1) lands for growing lime and are distributed in the western and central part of the microwatershed. Maximum area of 285 ha (62%) is moderately suitable (Class S2) and are distributed in the major part of the microwatershed. They have minor limitations of calcareousness, rooting condition and gravelliness. Marginally suitable (Class S3) lands occur in an area of 57 ha (12%) for growing lime and distributed in the northern and western part of the microwatershed with moderate limitations of rooting condition and calcareousness. An area of 51 ha (11%) is currently not suitable (Class N1) for growing lime and are distributed in the northern, western, northeastern part of the microwatershed with severe limitations of gravelliness, calcareousness and rooting condition.




Fig. 7.20 Land Suitability map of Lime

### 7.21 Land Suitability for Amla (*Phyllanthus emblica*)

Amla is one of the most important fruit and medicinal crop grown in 151 ha area and distributed in almost all the districts of the State. The crop requirements for growing amla (Table 7.22) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing amla was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.21.

There are no highly suitable (Class S1) lands for growing amla in the microwatershed. An area of 393 ha (85%) has soils that are moderately suitable (Class S2) and are distributed in the major part of the microwatershed. They have minor limitations of rooting condition, gravelliness, texture and calcareousness. The marginally suitable (Class S3) lands cover an area of 43 ha (9%) and occur in the northern and northeastern part of the microwatershed with moderate limitations of calcareousness and texture. An area of 8 ha (2%) is currently not suitable (Class N1) and are distributed in the western part of the microwatershed with severe limitations of rooting condition and gravelliness.




Fig. 7.21 Land Suitability map of Amla

## 7.22 Land Suitability for Cashew (*Anacardium occidentale*)

Cashew is one of the most important nut crop grown in an area of 1.24 lakh ha in almost all the districts of the State. The crop requirements for growing cashew (Table 7.23) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing cashew was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.22.

There are no highly (Class S1), moderately (Class S2) and marginally (Class S3) suitable for growing cashew in the microwatershed. Maximum area of about 444 ha (96%) is currently not suitable (Class N1) for growing cashew and are distributed in the entire part of the microwatershed with severe limitations of texture, rooting condition, calcareousness and gravelliness.




Fig. 7.22 Land Suitability map of Cashew

### 7.23 Land Suitability for Jackfruit (*Artocarpus heterophyllus*)

Jackfruit is one of the most important fruit crop grown in 5368 ha in all the districts of the State. The crop requirements for growing jackfruit (Table 7.24) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing jackfruit was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in figure 7.23.

There are no highly (Class S1) and moderately (Class S2) lands suitable for growing jackfruit in the microwatershed. Maximum area of about 393 ha (85%) is marginally suitable (Class S3) and are distributed in all parts of the microwatershed. They have moderate limitations of gravelliness, texture and calcareousness. An area of 51 ha (11%) is currently not suitable (Class N1) and occur in the northern, western, and northeastern part of the microwatershed with severe limitations of gravelliness, texture and rooting condition.




Fig. 7.23 Land Suitability map of Jackfruit

#### 7.24 Land Suitability for Jamun (*Syzygium cumini*)

Jamun is an important fruit crop grown in almost all the districts of the State. The crop requirements for growing jamun (Table 7.25) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing jamun was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.24.

There are no highly suitable (Class S1) lands for growing jamun in the microwatershed. An area of 138 ha (33%) is moderately suitable (Class S2) and occur in the northwestern, central, eastern and southern part of the microwatershed. They have minor limitations of rooting condition, texture and calcareousness. Marginally suitable (Class S3) lands cover an area of 239 ha (52%) and are distributed in the major part of the microwatershed with moderate limitations of rooting condition, calcareousness, texture and gravelliness. An area of 51 ha (11%) is currently not suitable (Class N1) for growing jamun and are distributed in the northern, western and northeastern part of the microwatershed with severe limitations of gravelliness, texture and rooting condition.




Fig. 7.24 Land Suitability map of Jamun

### 7.25 Land Suitability for Custard Apple (*Annona reticulata*)

Custard apple is one of the most important fruit crop grown in almost all the districts of the State. The crop requirements for growing custard apple (Table 7.26) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing custard apple was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.25.

Maximum area of 230 ha (50%) is highly (Class S1) suitable lands for growing custard apple and are distributed in the major part of the microwatershed. An area of 163 ha (35%) is moderately suitable (Class S2) and are distributed in the northern, eastern, northwestern, western and southern part of the microwatershed. They have minor limitations of gravelliness, rooting condition, texture and calcareousness. An area of 43 ha (9%) is marginally suitable (Class S3) for growing custard apple and are distributed in the northern and northeastern part of the microwatershed with moderate limitations of gravelliness and calcareousness. An area of 8 ha (2%) is currently not suitable (Class N1) and are distributed in the western part of the microwatershed with severe limitations of rooting condition and gravelliness.




Fig. 7.25 Land Suitability map of Custard Apple

## 7.26 Land Suitability for Tamarind (*Tamarindus indica*)

Tamarind is one of the most important spice crop grown in 14897 ha in all the districts of the State. The crop requirements for growing tamarind (Table 7.27) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing tamarind was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.26.

There are no highly (Class S1) suitable lands for growing tamarind in the microwatershed. An area of 159 ha (34%) is moderately suitable (Class S2) and occur in the northern, northwestern, eastern, central and southern part of the microwatershed. They have minor limitations of rooting condition, texture, gravelliness and calcareousness. Maximum area of 177 ha (38%) is marginally suitable (Class S3) and occur in the major part of the microwatershed with moderate limitations of gravelliness, rooting condition and calcareousness. An area of 108 ha (23%) is currently not suitable (Class N1) and are distributed in the northern, western and northeastern part of the microwatershed with severe limitations of rooting condition, calcareousness and gravelliness.






Fig. 7.26 Land Suitability map of Tamarind

### 7.27 Land Suitability for Mulberry (*Morus nigra*)

Mulberry is the most important leaf crop grown for rearing silkworms in about 1.66 lakh ha in all the districts of the State. The crop requirements for growing mulberry (Table 7.28) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing mulberry was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.27.

There are no highly suitable (Class S1) lands for growing mulberry in the microwatershed. Moderately suitable (Class S2) lands occupy a maximum area of 300 ha (65%) and are distributed in the major part of the microwatershed. They have minor limitations of calcareousness, gravelliness, drainage and texture. Marginally suitable (Class S3) lands cover an area of 93 ha (20%) and are distributed in the northern, northeastern, western and southern part of the microwatershed. They have moderate limitations of rooting condition, calcareousness, and texture. An area of 51 ha (11%) is currently not suitable (Class N1) and are distributed in the northern, western and northeastern part of the microwatershed with severe limitations of rooting depth, calcareousness and gravelliness.




Fig. 7.27 Land Suitability map of Mulberry

## 7.28 Land Suitability for Marigold (*Tagetes erecta*)

Marigold is one of the most important flower crop grown in an area of 1858 ha in almost all the districts of the State. The crop requirements for growing marigold (Table 7.29) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing marigold was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.28.

There are no highly suitable (Class S1) lands for growing marigold in the microwatershed. An area of 388 ha (84%) is moderately suitable (Class S2) for growing marigold and are distributed in the major part of the microwatershed. They have minor limitations of texture, gravelliness, rooting condition, drainage and calcareousness. An area of 48 ha (10%) is marginally suitable (Class S3) for growing marigold and are distributed in the northern and northeastern part of the microwatershed. They have moderate limitations of gravelliness, calcareousness, texture and rooting condition. An area of 8 ha (2%) is currently not suitable (Class N1) and are distributed in the northern and western part of the microwatershed with severe limitations of rooting condition and gravelliness.




Fig. 7.28 Land Suitability map of Marigold

### 7.29 Land Suitability for Chrysanthemum (*Chrysanthemum indicum*)

Chrysanthemum is one of the most important flower crop grown in an area of 4978 ha in almost all the districts of the State. The crop requirements for growing chrysanthemum (Table 7.30) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing chrysanthemum was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed is given in Figure 7.29.

There are no highly suitable (Class S1) lands for growing chrysanthemum in the microwatershed. Maximum area of 388 ha (84%) is moderately suitable (Class S2) for growing chrysanthemum and are distributed in the major part of the microwatershed. They have minor limitations of calcareousness, gravelliness, rooting condition, drainage and texture. An area of 48 ha (10%) is marginally suitable (Class S3) and occur in the northern and northeastern part of the microwatershed. They have moderate limitations of gravelliness, rooting condition, texture and calcareousness. Currently not suitable (Class N1) lands occur in an area of 8 ha (2%) and are distributed in the western part of the microwatershed with severe limitations of rooting condition and gravelliness.




Fig. 7.29 Land Suitability map of Chrysanthemum

### 7. 30 Land Suitability for Jasmine (*Jasminum sp.*)

Jasmine is one of the most important flower crop grown in an area of 6146 ha in almost all the districts of the State. The crop requirements (Table 7.31) for growing jasmine were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing jasmine was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.30.

There are no highly suitable lands (Class S1) for growing jasmine in the microwatershed. An area of 57 ha (12%) is moderately suitable (Class S2) and occur in the northern and western part of the microwatershed. They have minor limitations of rooting condition and calcareousness. A maximum area of 379 ha (82%) is marginally suitable (Class S3) for growing jasmine and are distributed in the major part of the microwatershed. They have moderate limitations of gravelliness, texture, rooting condition, drainage and calcareousness. Currently suitable (Class N1) lands occur in an area of 8 ha (2%) and are distributed in the western part of the microwatershed with severe limitations of rooting condition and gravelliness.




Fig. 7.30 Land Suitability map of Jasmine

### 7.31 Land Suitability for Crossandra (*Crossandra in fundibuliformis*)

Crossandra is one of the most important flower crop grown in an area of 6146 ha in almost all the districts of the State. The crop requirements (Table 7.32) for growing crossandra were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing crossandra was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.31.

There are no highly suitable lands (Class S1) for growing crossandra in the microwatershed. An area of 85 ha (18%) is moderately suitable (Class S2) for growing crossandra and occur in the western, central, southern and eastern part of the microwatershed. They have minor limitations of texture and calcareousness. A maximum area of 350 ha (76%) is marginally suitable (Class S3) and are distributed in the major part of the microwatershed. They have moderate limitations of gravelliness, rooting condition, texture and calcareousness. An area of 8 ha (2%) is currently not suitable (Class N1) and are distributed in the western part of the microwatershed with severe limitations of rooting condition and gravelliness.




Fig. 7.31 Land Suitability map of Crossandra

**Table 7.1 Soil-Site Characteristics of Handrahalu-2 Microwatershed**

Soil Map Units	Climate (P) (mm)	Growing period (Days)	Drainage Class	Soil depth (cm)	Soil texture		Gravelliness		AWC (mm/m)	Slope (%)	Erosion	pH	EC	ESP	CEC [Cmol (p <sup>+</sup> ) kg <sup>-1</sup> ]	BS (%)
					Surface	Sub-surface	Surface	Sub-surface								
BGTmB1g2	662	90	WD	<25	c	gc	35-60	>35	<50	1-3	Slight	8.4	0.15	1.11	44.84	-
MTLmB2g1	662	90	WD	25-50	c	gc	15-35	15-35	51-100	1-3	Moderate	8.27	0.20	0.69	36.64	-
RNKmB1	662	90	WD	50-75	c	c	-	<15	51-100	1-3	Slight	8.86	0.48	6.78	37.0	-
RNKmB1g1	662	90	WD	50-75	c	c	15-35	<15	51-100	1-3	Slight	8.86	0.48	6.78	37.0	-
RNKmB2g1	662	90	WD	50-75	c	c	15-35	<15	51-100	1-3	Moderate	8.86	0.48	6.78	37.0	-
DRLmB1	662	90	WD	75-100	c	c	-	<15	151-200	1-3	Slight	8.78	0.42	5.62	49.70	100
NSPmB1	662	90	WD	75-100	c	c	-	-	101-150	1-3	Slight	9.16	0.61	8.60	51.09	-
NSPmB1g1	662	90	WD	75-100	c	c	15-35	-	101-150	1-3	Slight	9.16	0.61	8.60	51.09	-
NSPmB2	662	90	WD	75-100	c	c	-	-	101-150	1-3	Moderate	9.16	0.61	8.60	51.09	-
NSPmB2g1	662	90	WD	75-100	c	c	15-35	-	101-150	1-3	Moderate	9.16	0.61	8.60	51.09	-
GRHmB1g1	662	90	WD	100-150	c	c	15-35	<15	>200	1-3	Slight	9.08	0.23	7.11	63.21	100
GRHmB2g1	662	90	WD	100-150	c	c	15-35	<15	>200	1-3	Moderate	9.08	0.23	7.11	63.21	100
HDLmB1	662	90	WD	100-150	c	c	-	-	>200	1-3	Slight	9.06	0.37	5.09	62.33	-
HDLmB2	662	90	WD	100-150	c	c	-	-	>200	1-3	Moderate	9.06	0.37	5.09	62.33	-
HDLmB2g1	662	90	WD	100-150	c	c	15-35	-	>200	1-3	Moderate	9.06	0.37	5.09	62.33	-
KDTmB2	662	90	WD	>150	c	sc-c	-	-	>200	1-3	Moderate	6.95	0.17	0.65	12.10	100
MLRmA1	662	90	WD	>150	c	c	-	10-20	>200	0-1	Slight	9.19	0.31	5.39	42.08	-
BDRmB1g2	662	90	WD	>150	c	c	35-60	<15	>200	1-3	Slight	8.73	0.20	4.37	40.56	-

\*Symbols and abbreviations are according to Field Guide for LRI under Sujala-III Project, Karnataka

**Table 7.2 Land suitability criteria for Sorghum**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime 1	Mean temperature in growing season	°C	26–30	30–34; 24–26	34–40; 20–24	>40; <20
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristics					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, c (red), c (black)	sc1, cl	ls, sl	-
	pH	1:2.5	5.5-7.8	5.0-5.5 7.8-9.0	>9.0	-
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	10-15
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	0-3	3-5	5-10	>10



**Table 7.3 Land suitability criteria for Maize**

Land use requirement		Rating				
Soil –site characteristics	Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)	
Climatic regime	Mean temperature in growing season	°C	30-34	35-38 26-30	38-40 26-20	
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc	c (red), c (black)	ls, sl	-
	pH	1:2.5	5.5-7.8	5.0-5.5 7.8-9.0	>9.0	-
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	-
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

**Table 7.4 Land suitability criteria for Bajra**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-32	33-38 24-27	39-40 20-23	<20
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm	500-750	400-500	200-400	<200
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl, scl, cl,sc,c (red)	C (black)	ls	-
	pH	1:2.5	6.0-7.8	5.0-5.5 7.8-9.0	5.5-6.0 >9.0	
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	15-35	35-60	>60	
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	1-3	3-5	5-10	>10

**Table 7.5 Land suitability criteria for Groundnut**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	24–33	22–24; 33–35	20–22; 35–40	<20; >40
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. Well drained	Poorly drained	Very Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl	sl,cl, sc	c (red), c (black), ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<35	35-60	>60	
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.6 Land suitability criteria for Sunflower**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	24–30	30–34; 20–24	34–38; 16–20	>38; <16
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	mod. Well drained	-	Poorly to very drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	cl, sc,c (red), c (black)	scl	ls, sl	-
	pH	1:2.5	6.5-7.8	7.8-8.4 5.5-6.5	8.4-9.0; 5.0-5.5	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.7 Land suitability criteria for Red gram**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	30-35(G) 20-25(AV) 15-18 (F&PS) 35-40(M)	25-30(G) 20-25 (AV) 12-15 (F&PS) 30-35(M)	20-25(G) 15-20(AV) 10-12 (F&PS) 25-30(M)	< 20 <15 <10 <25
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. Well drained	Poorly drained	Very Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, c (red)	c (black),sl, scl, cl	ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-9.0	5.0-5.5 >9.0	-
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-50	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<1.0	1.0-2.0	>2.0	
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.8 Land suitability criteria for Bengal gram**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	20–25	25–30; 15–20	30–35; 10–15	>35; <10
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. Well drained	Poorly drained	Very Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	C (black)	-	c (red), scl, cl, sc	ls, sl
	pH	1:2.5	6.0-7.8	5.0-6.0 7.8-9.0	>9.0	-
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	-
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.9 Land suitability criteria for Cotton**

Land use requirement		Rating				
Soil –site characteristics	Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)	
Climatic regime	Mean temperature in growing season	°C	22-32	>32	<19	-
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well to moderately well	Poorly drained/Some what excessively drained	-	very poorly/excessively drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, c (red,black)	cl	scl	ls, sl
	pH	1:2.5	6.5-7.8	7.8-8.4	5.5-6.5 8.4->9.0	<5.5
	CEC	C mol (p+)Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	50-100	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	<3	3-5	-	>5

**Table 7.10 Land suitability criteria for Chilli**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	25-32	33-35 20-25	35-38 <20	>38
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc	c (black), sl	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10



**Table 7.11 Land suitability criteria for Tomato**

Land use requirement		Rating				
Soil –site characteristics	Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)	
Climatic regime	Mean temperature in growing season	°C	25-28	29-32 20-24	15-19 33-36	<15 >36
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl, scl, cl, sc, c (red)	-	ls, c(black)	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.12 Land suitability criteria for Brinjal**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	Well drained	Moderately well drained	Poorly drained	V. Poorly drained
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class				
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl, scl, cl, sc c (red)	-	ls, c (black)	-
	pH	1:2.5	6.0-7.3	7.3-8.4 5.0-6.0	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	>60
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.13 Land suitability criteria for Onion**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	20-30	30-35	35-40	>40
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately/imperfectly	-	Poorly to V poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl,scl,cl,sc,c (red)	-	c (Black),ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<1.0	1.0-2.0	2.0-4.0	<4
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.14 Land suitability criteria for Bhendi**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	25-28	29-32 20-24	15-19 33-36	<15 >36
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Imperfectly drained	Poorly to very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl,sc, c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.15 Land suitability criteria for Drumstick**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, scl, cl, c (red)	sl, c (black)	ls	s
	pH	1:2.5	6.0-7.3	5.0-5.5 7.3-7.8	5.5-6.0 7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<35	35-60	60-80	>80
Soil toxicity	Salinity (EC saturation extract)	dS/m				
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-10	-	>10

**Table 7.16 Land suitability criteria for Mango**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-32	24-27 33-35	36-40	20-24
	Min temp. before flowering	°C	10-15	15-22	>22	-
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration	Days				
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	-	ls, sl, c (black)	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>150	100-150	75-100	<75
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.17 Land suitability criteria for Guava**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-32	33-36 24-27	37-42 20-23	
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	sl	c (black), ls	-
	pH	1:2.5	6.0-7.8	5.0-6.0	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.18 Land suitability criteria for Sapota**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-32	33-36 24-27	37-42 20-23	>42 <18
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	-	Poorly to very drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	sl	ls, c (black)	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10



**Table 7.19 Land suitability criteria for Pomegranate**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	30-34	35-38 25-29	39-40 15-24	
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl,cl, sc, c (red)	c (black),sl	ls	-
	pH	1:2.5	5.5-7.8	7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.20 Land suitability criteria for Musambi**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-30	31-35 24-27	36-40 20-23	>40 <20
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately drained	poorly	Very poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c	sl	ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.21 Land suitability criteria for Lime**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-30	31-35 24-27	36-40 20-23	>40 <20
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately drained	poorly	Very poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c	sl	ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.22 Land suitability criteria for Amla**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. well drained	Poorly drained	V. Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	c (black)	ls, sl	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15-35	35-60	60-80	-
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

**Table 7.23 Land suitability criteria for Cashew**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	32 to 34	28 to 32; 34 to 38	24 to 28; 38 to 40	<20; >40
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	-	sl, ls	c (black)
	pH	1:2.5	5.5-6.5	5.0-5.5 6.5-7.3	7.3-7.8	>7.8
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-10	>10	-

**Table 7.24 Land suitability criteria for Jackfruit**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. well	Poorly	V. Poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl, cl, sc, c (red)	-	sl, ls, c (black)	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	>60
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10-

**Table 7.25 Land suitability criteria for Jamun**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well	Mod. well	Poorly	V.Poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c(red)	sl, c (black)	ls	-
	pH	1:2.5	6.0-7.8	5.0-6.0	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>150	100-150	50-100	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	>60
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

**Table 7.26 Land suitability criteria for Custard apple**

Land use requirement		Rating				
Soil –site characteristics	Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)	
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	Scl, cl, sc, c (red), c (black)	-	Sl, ls	-
	pH	1:2.5	6.0-7.3	5.5-6.0 7.3-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15-35	35-60	60-80	-
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	>5	-



**Table 7.27 Land suitability criteria for Tamarind**

Land use requirement		Rating				
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod.well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl,sc, c (red)	sl, c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>150	100-150	75-100	<75
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

**Table 7.28 Land suitability criteria for Mulberry**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	24–28	22–24; 28–32	32–38; 22–18	>38; <18
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V. Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, cl, scl	c (red)	c (black), sl, ls	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.8-8.4	7.3-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	0-35	35-60	60-80	>80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

**Note:** Suitability evaluation only for Mulberry leaf not for Silk worm rearing

**Table 7.29 Land suitability criteria for Marigold**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	18-23	17-15 24-35	35-40 10-14	>40 <10
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl,scl, cl, sc, c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%				
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.30 Land suitability criteria for Chrysanthemum**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	18-23	17-15 24-35	35-40 10-14	>40 <10
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl,scl, cl, sc, c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%				
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.31 Land suitability criteria for jasmine (irrigated)**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	18-23	17-15 24-35	35-40 10-14	-
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	sl	ls, c (black)	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%				
Erosion hazard	Slope	%	<3	3-5	5-10	>10

### 7.32 Land suitability criteria for Crossandra

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	-	Poorly to very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c(red)	sl,	c (black),ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%				
Erosion hazard	Slope	%	<3	3-5	5-10	>10

### 7.32 Land Management Units (LMUs)

The 18 soil map units identified in Handrahalu-2 Microwatershed have been grouped into 4 Land Management Units (LMUs) for the purpose of preparing a Proposed Crop Plan. Land Management Units are grouped based on the similarities in respect of the type of soil, the depth of the soil, the surface soil texture, gravel content, AWC, slope, erosion etc. and a Land Management Units map (Fig. 7.32) has been generated. These Land Management Units are expected to behave similarly for a given level of management.

The map units that have been grouped into four Land Management Units along with brief description of soil and site characteristics are given below.

LMU	Soil map unit number	Mapping unit	Soil and site characteristics
1	348, 360, 361, 362, 363, 372, 374, 380, 382, 383, 405, 411, 432	DRLmB1, NSPmB1, NSPmB1g1, NSPmB2, NSPmB2g1, GRHmB1g1, GRHmB2g1, HDLmB1, HDLmB2, HDLmB2g1, KDTmB2, MLRmA1, BDRmB1g2	Moderately deep to very deep calcareous to non calcareous clay soils
2	333, 334, 337	RNKmB1, RNKmB1g1, RNKmB1g1	Moderately shallow, black calcareous clay soils
3	311	MTLmB2g1	Shallow, black calcareous clay soils
4	8	BGTmB1g2	Very shallow, calcareous gravelly black clay soils




Fig 7.32 Land Management Units map of Handrahalu-2 Microwatershed

### 7.33 Proposed Crop Plan for Handrahalu-2 Microwatershed

After assessing the land suitability for the 31 crops, the proposed crop plan has been prepared for the 4 identified LMUs by considering only the highly (Class S1) and moderately (Class S2) suitable lands for each of the 31 crops. The resultant proposed crop plan is presented in Table 7.33.



**Table 7.33 Proposed Crop Plan for Handrahalu-2 Microwatershed**

LMU	Soil Map Units	Survey Number	Soil characters	Field Crops	Horticulture Crops	Suitable Interventions
LMU 1 336 ha (73%)	348.DRLmB1 360.NSPmB1 361.NSPmB1g1 362.NSPmB2 363.NSPmB2g1 372.GRHmB1g1 374.GRHmB2g1 380.HDLmB1 382.HDLmB2 383.HDLmB2g1 405.KDTmB2 411.MLRmA1 432.BDRmB1g2	<b>Hanavala:</b> 58,69,70,71,72,73, 74,75,76,77,78,79,80,83,84,85,86,8 7, 88,89,90,102,103, 104 <b>Handrala:</b> 1,2,3,4,5,6,7,8,9,66,67,6 8,69,70,78,79,80,81,82,83,84,85,86 ,87,88,89,90,91,92,93,94,95,96,97, 98,99,100,102,107,148,150,153,15 4,155,156,158,159,160,162,163,16 4,165,166,167,168,169,170,171,17 2,173,174,175,176,177,178,179,18 0,181 <b>Hireshindhogi:</b> 273,274,275,276,2 77,278,279,280,281,284,285,286,2 87,288,289,290,293,294,295 <b>Narasapura:</b> 33	Moderately deep to very deep calcareous to non calcareous clay soils	Maize, Sorghum, Sunflower, Cotton, Bengal gram, Safflower, Linseed, Bajra , Soybean	<b>Fruit crops:</b> Mango, Sapota, Pomegranate, Jamun, Lime, Musambi, Tamarind, Amla, Custard apple <b>Vegetables:</b> Drumstick, Chilli, Coriander, Tomato, Bhendi <b>Flowers:</b> Marigold, Chrysanthemum, Crossandra, Jasmine	Application of FYM, Biofertilizers and micronutrients, drip irrigation, mulching, suitable soil and water conservation practices
LMU 2 57 ha (12%)	333.RNKmB1 334.RNKmB1g1 337.RNKmB2g1	<b>Handrala:</b> 101,103,104,105,106, 108,109,110,111,112, 113,114,152	Moderately shallow, black calcareous clay soils	Sorghum, Bajra, Bengal gram, linseed, Safflower, Coriander	<b>Fruit crops:</b> Amla, Custard apple <b>Flower crops:</b> Marigold, Jasmine Chrysanthemum	Application of FYM, Biofertilizers and micronutrients, drip irrigation, mulching, suitable soil and water conservation practiCes
LMU 3 43 ha (9%)	311.MTLmB2g1	<b>Hanavala:</b> 56,57,59,60,61,62,63, 64,65,66,67,68	Shallow, black calcareous clay soils	Bengal gram	<b>Agri-Silvi-Pasture:</b> Hybrid Napier, <i>Styloxanthes hamata</i> , <i>Styloxanthes scabra</i>	Sowing across the slope, drip irrigation and mulching is recommended
LMU 4 8 ha (2%)	8.BGTmB1g2	<b>Handrala :</b> 147,157	Very shallow, calcareous gravelly black clay soils	Bengal gram	<b>Agri-Silvi-Pasture:</b> Hybrid Napier, <i>Styloxanthes hamata</i> , <i>Styloxanthes scabra</i>	Use of short duration varieties, sowing across the slope



## SOIL HEALTH MANAGEMENT

### 8.1 Soil Health

Soil health is basic to plant health and plant health is basic to human and bovine health. Soil is fundamental to crop production. Without soil, no food could be produced nor would livestock be fed on a large scale. Because it is finite and fragile, soil is a precious resource that requires special care from its users.

Soil health or the capacity of the soil to function is critical to human survival. Soil health has been defined as: “the capacity of the soil to function as a living system without adverse effect on the ecosystem”. Healthy soils maintain a diverse community of soil organisms that help to form beneficial symbiotic associations with plant roots, recycle essential plant nutrients, improve soil structure with positive repercussions for soil, water and nutrient holding capacity and ultimately improve crop production and also contribute to mitigating climate change by maintaining or increasing its carbon content.

Functional interactions of soil biota with organic and inorganic components, air and water determine a soil’s potential to store and release nutrients, and water to plants and to promote and sustain plant growth. Thus, maintaining soil health is vital to crop production and conserve soil resource base for sustaining agriculture.

#### **The most important characteristics of a healthy soil are**

- Good soil tilth
- Sufficient soil depth
- Good water storage and good drainage
- Adequate supply, but not excess of nutrients
- Large population of beneficial organisms
- Small proportion of plant pathogens and insect pests
- Low weed pressure
- Free of chemicals and toxins that may harm the crop
- Resistance to degradation
- Resilience when unfavourable conditions occur

#### **Characteristics of Handrahalu-2 Microwatershed**

- ❖ The soil phases with sizeable area identified in the microwatershed belonged to the soil series of Narasapura (NSP) 161 ha (35%), Ravanaki (RNK) 57 ha (12%), Gatareddihal (GRH) 54 ha (12%), Handrala (HDL) 48 ha (10%), Muttal (MTL) 43 ha (9%), Murlapur (MLR) 31 ha (7%), Kadagathur (KDT) 21 ha (5%), Dambarahalli (DRL) 16 ha (4%), Belagatti (BGT) 8 ha (2%) and Bardur (BDR) occupy an area of about 5 ha (1%) in the microwatershed.

- ❖ As per land capability classification, entire area in the microwatershed falls under arable land category (Class II, III & IV). The major limitations identified in the arable lands were soil and erosion.
- ❖ On the basis of soil reaction, an area of about 8 ha (2%) is moderately alkaline (pH 7.8-8.4), 322 ha (70%) is strongly alkaline (pH 8.4-9.0) and about 115 ha (25%) is very strongly alkaline (pH >9.0) in the microwatershed. Entire area in the microwatershed is moderately to very strongly alkaline in reaction.

### **Soil Health Management**

The following actions are required to improve the current land husbandry practices that provide a sound basis for the successful adoption of sustainable crop production system.

#### **Alkaline soils**

Slightly to very strongly alkaline soils cover an area of 497 ha.

1. Regular addition of organic manure, green manuring, green leaf manuring, crop residue incorporation and mulching needs to be taken up to improve the soil organic matter status.
2. Application of biofertilizers (Azospirillum, Azatobacter, Rhizobium).
3. Application of 25% extra N and P (125 % RDN&P).
4. Application of ZnSO<sub>4</sub> – 12.5 kg/ha (once in three years).
5. Application of Boron – 5 kg/ha (once in three years).

#### **Soil Degradation**

Soil erosion is one of the major factor affecting the soil health in the microwatershed. Out of total 482 ha area in the microwatershed, an area of about 200 ha (43%) is suffering from slight erosion and 244 ha (53%) is suffering from moderate erosion. The areas suffering from moderate erosion need immediate soil and water conservation and, other land development and land husbandry practices for restoring soil health.

#### **Dissemination of Information and Communication of Benefits**

Any large scale implementation of soil health management requires that supporting information is made available widely, particularly through channels familiar to farmers and extension workers. Given the very high priority attached to soil health especially by the Central Government on issuing Soil-Health Cards to all the farmers, media outlets like Regional, State and National Newspapers, Radio and Dooradarshan programs in local languages but also modern information and communication technologies such as Cellular phones and the Internet, which can be much more effective in reaching the younger farmers.

#### **Inputs for Net Planning (Saturation Plan) and Interventions needed**

Net planning (Saturation plan) in IWMP is focusing on preparation of

1. Soil and Water Conservation Treatment Plans for each plot or farm.

2. Productivity enhancement measures/ interventions for existing crops/livestock/other farm enterprises.
3. Diversification of farming mainly with perennial horticultural crops and livestock.
4. Improving livelihood opportunities and income generating activities.

In this connection, how various outputs of Sujala-III are of use in addressing these objectives of Net Planning are briefly presented below.

- ❖ **Soil Depth:** The depth of a soil decides the amount of moisture and nutrients it can hold, what crops can be taken up or not, depending on the rooting depth and the length of growing period available for raising any crop. Deeper the soil, better for a wide variety of crops. If sufficient depth is not available for growing deep rooted crops, either choose medium or short duration crops or deeper planting pits need to be opened and additional good quality soil brought from outside has to be filled into the planting pits.
- ❖ **Surface Soil Texture:** Lighter soil texture in the top soil means, better rain water infiltration, less run-off and soil moisture conservation, less capillary rise and less evaporation losses. Lighter surface textured soils are amenable to good soil tilth and are highly suitable for crops like groundnut, root vegetables (carrot, radish, potato etc) but not ideal for crops that need stagnant water like lowland paddy. Heavy textured soils are poor in water infiltration and percolation. They are prone for sheet erosion; such soils can be improved by sand mulching. The technology that is developed by the AICRP-Dryland Agriculture, Vijayapura, Karnataka can be adopted.
- ❖ **Gravelliness:** More gravel content is favorable for run-off harvesting but poor in soil moisture storage and nutrient availability. It is a significant parameter that decides the kind of crop to be raised.
- ❖ **Land Capability Classification:** The land capability map shows the areas suitable and not suitable for agriculture and the major constraints in each of the plot/survey number. Hence, one can decide what kind of enterprise is possible in each of these units. In general, erosion and soil are the major constraints in Handrahalu-2 Microwatershed.
- ❖ **Organic Carbon:** The OC content (an index of available Nitrogen) is low (<0.5%) in 93 ha (20%) and medium (0.5-0.75%) in 349 ha (76%). The areas that are low and medium in OC needs to be further improved by applying farmyard manure and rotating crops with cereals and legumes or mixed cropping. It is high (>0.75%) in an area of 2 ha (1%) in the microwatershed.
- ❖ **Promoting green manuring:** Growing of green manuring crops costs Rs. 1250/ha (green manuring seeds) and about Rs. 2000/ha towards cultivation that totals to Rs. 3250/- per ha. On the other hand, application of organic manure @ 10 tons/ha costs Rs. 5000/ha. The practice needs to be continued for 2-3 years or more. Nitrogen fertilizer needs to be supplemented by 25% in addition to the recommended level in 442 ha area where OC is low and medium. For example, for rainfed maize, recommended level is 50 kg N per ha and an additional 12 kg /ha needs to be applied for all the crops grown in these plots.


- ❖ **Available Phosphorus:** An area of about 105 ha (23%) is low (<23 kg/ha) and 340 ha (74%) is medium (23-57 kg/ha) in available phosphorus. Hence for all the crops, 25% additional P-needs to be applied where it is low and medium.
- ❖ **Available Potassium:** Available potassium medium (145-337 kg/ha) in 8 ha (1%) and high (>337 kg/ha) in 437 (95%) in the microwatershed. Additional 25% potassium needs to be applied in areas where it is medium.
- ❖ **Available Sulphur:** Available sulphur is a very critical nutrient for oilseed crops. Available sulphur is medium (10-20 ppm) in 96 ha (21%) and high (>20 ppm) is about 348 ha (75%) in the microwatershed. These areas need to be applied with magnesium sulphate or gypsum or Factamphos (p) fertilizer (13% sulphur) for 2-3 years for the deficiency to be corrected.
- ❖ **Available Boron:** An area of about 303 ha (66%) is low (<0.5 ppm) and 141 ha (31%) is medium (0.5-1.0 ppm) in available boron content. The areas that are low and medium need to be applied with sodium borate @ 10 kg/ha as soil application or 0.2% borax as foliar spray to correct the deficiency.
- ❖ **Available iron:** It is sufficient (>4.5 ppm) in 444 ha (96%) area in the microwatershed.
- ❖ **Available manganese:** Entire area in the microwatershed is sufficient (>1.0 ppm) in available manganese.
- ❖ **Available copper:** Entire area is sufficient (>0.2 ppm) in available copper in the microwatershed.
- ❖ **Available Zinc:** It is deficient (<0.6 ppm) in 424 ha (92%) and sufficient (>0.6 ppm) in 20 ha (4%) area in the microwatershed. Application of zinc sulphate @ 25 kg/ha is to be followed in areas that are deficient in available zinc.
- ❖ **Soil alkalinity:** The microwatershed has 445 ha (96%) soils that are moderately to very strongly alkaline. These areas need application of gypsum and wherever calcium is in excess, iron pyrites and element sulphur can be recommended. Management practices like treating repeatedly with good quality water to drain out the excess salts and provision of subsurface drainage and growing of salt tolerant crops like Casuarina, Acacia, Neem, Ber etc, are recommended.

**Land suitability for various crops:** Areas that are highly, moderately and marginally suitable and not suitable for growing various crops are indicated. Along with the suitability, various constraints that are limiting the productivity are also indicated. For example, in case of cotton, gravel content, rooting depth and salinity/alkalinity are the major constraints in various plots. With suitable management interventions, the productivity can be enhanced. In order to increase water holding capacity of light textured soils, growing of green manure crops and application of organic manure is recommended.

## SOIL AND WATER CONSERVATION TREATMENT PLAN

For preparing soil and water conservation treatment plan for Handrahalu-2 Microwatershed, the land resource inventory database generated under Sujala-III project has been transformed as information through series of interpretative (thematic) maps using soil phase map as a base. The various thematic maps (1:7920 scale) generated were

- Soil depth
  - Surface soil texture
  - Available water capacity
  - Soil slope
  - Soil gravelliness
  - Land capability
  - Present land use and land cover
  - Crop suitability maps
  - Rainfall map
  - Hydrology
  - Water Resources
  - Socio-economic data
  - Contour plan with existing features- network of waterways, pottissa boundaries, cut up/ minor terraces etc.
  - Cadastral map (1:7920 scale)
  - Satellite imagery (1:7920 scale)
- Apart from these, Hand Level/ Hydro Marker/ Dumpy Level/ Total Station and Kathedars' List to be collected.



### Steps for Survey and Preparation of Treatment Plan

The boundaries of Land User Groups' and Survey No. boundaries are traced in the field.

- Naming of user groups and farmers
- Identification of arable and non arable lands
- Identification of drainage lines and gullies
- Identification of non treatable areas
- Identification of priority areas in the arable lands
- Treatment plan for arable lands
- Location of water harvesting and recharge structures

## 9.1 Treatment Plan

The treatment plan recommended for arable lands is briefly described below.



### 9.1.1 Arable Land Treatment

#### A. BUNDING

Steps for Survey and Preparation of Treatment Plan		<b>USER GROUP-1</b> <b>CLASSIFICATION OF GULLIES</b> 
Cadastral map (1:7920 scale) is enlarged to a scale of 1:2500 scale Existing network of waterways, pottissa boundaries, grass belts, natural drainage lines/watercourse, cut ups/ terraces are marked on the cadastral map to the scale Drainage lines are demarcated into		
Small gullies	(up to 5 ha catchment)	UPPER REACH MIDDLE REACH LOWER REACH POINT OF CONCENTRATION
Medium gullies	(5-15 ha catchment)	
Ravines	(15-25 ha catchment) and	
Halla/Nala	(more than 25ha catchment)	

#### Measurement of Land Slope

Land slope is estimated or determined by the study and interpretation of contours or by measurement in the field using simple instruments like Hand level or Hydromarker.



Vertical and Horizontal intervals between bunds as recommended by the Watershed Development Department.

Slope percentage	Vertical interval (m)	Corresponding Horizontal Distance (m)
2 - 3%	0.6	24
3 - 4%	0.9	21
4 - 5%	0.9	21
5 - 6%	1.2	21
6 - 7%	1.2	21



**Note:** i) The above intervals are maximum.

(ii) Considering the slope class and erosion status (A1... A= 0-1% slope, 1= slight erosion) the intervals have to be decided.

**Bund length recording:** Considering the contour plan and the existing grass belts/partitions, the bunds are aligned and lengths are measured.

### Section of the Bund

Bund section is decided considering the soil texture class and gravelliness class ( $bg_0 \dots b = \text{loamy sand}, g_0 = <15\% \text{ gravel}$ ). The recommended sections for different soils are given below.

#### Recommended Bund Section



Top width (m)	Base width (m)	Height (m)	Side slope (Z:1;H:V)	Cross section (sq m)	Soil Texture	Remarks
0.3	0.9	0.3	01:01	0.18	Sandy loam	Vegetative bund
0.3	1.2	0.3	1.5:1	0.225	Sandy clay	
0.3	1.2	0.5	0.9:1	0.375	Red gravelly soils	
0.3	1.2	0.6	0.75:1	0.45		
0.3	1.5	0.6	01:01	0.54	Red sandy loam	
0.3	2.1	0.6	1.5:1	0.72	Very shallow black clayey soils	
0.45	2	0.75	01:01	0.92		
0.45	2.4	0.75	1.3:1	1.07	Shallow black clayey soils	
0.6	3.1	0.7	1.78:1	1.29	Medium black clayey soils	
0.5	3	0.85	1.47:1	1.49		

### Formation of Trench cum Bund

Dimensions of the Borrow Pits/ Trenches to be excavated (machinery are decided considering the Bund Section).


Details of Borrow Pit dimensions are given below

#### TRENCH CUM BUND

IDEAL FOR HORTICULTURE CROPS

#### 'A' FRAME FOR INTERBUND MANAGEMENT



1. ಸಮಸಾಹಕಳೆ ಉಳಿಸುವೆ
2. ಸಮಸಾಹಕಳೆ ಬಿತ್ತನೆ/ನಾಟಿ

### Size of Borrow Pits/ Trench recommended for Trench cum Bund (by machinery)

Bund section	Bund length	Earth quantity	Pit				Berm (pit to pit)	Soil depth Class
			L(m)	W(m)	D(m)	QUANTITY (m <sup>3</sup> )		
m <sup>2</sup>	m	m <sup>3</sup>	L(m)	W(m)	D(m)	QUANTITY (m <sup>3</sup> )	m	
0.375	6	2.25	5.85	0.85	0.45	2.24	0.15	Shallow
0.45	6	2.7	5.4	1.2	0.43	2.79	0.6	Shallow
0.45	6	2.7	5	0.85	0.65	2.76	1	Moderately Shallow
0.54	5.6	3.02	5.5	0.85	0.7	3.27	0.1	Moderately shallow
0.54	5.5	2.97	5	1.2	0.5	3	0.5	Shallow
0.72	6.2	4.46	6	1.2	0.7	5.04	0.2	Moderately shallow
0.72	5.2	3.74	5.1	0.85	0.9	3.9	0.1	Moderately deep

#### B. Waterways

- Existing waterways are marked on the cadastral map (1:7920 scale) and their dimensions are recorded.
- Considering the contour plan of the MWS, additional waterways/ modernization of the existing ones can be thought of.
- The design details are given in the Manual.

#### C. Farm Ponds

Waterways and the catchment area will give an indication on the size of the Farm Pond. Location of the pond can be decided based on the contour plan/ field condition and farmers' need/desire.

#### D. Diversion channel

Existing EPT/ CPT are marked on the cadastral map. Looking to the need, these can be modernized or fresh diversion channel can be proposed and runoff from this can be stored in *Gokatte/* Recharge ponds.

#### 9.1.2 Non-Arable Land Treatment

Depending on the gravelliness and crops preferred by the farmers, the concerned authorities can decide appropriate treatment plan. The recommended treatments may be Contour Trench, Staggered Trench, Crescent Bund, Boulder Bund or Pebble Bund.

### **9.1.3 Treatment of Natural Water Course/ Drainage Lines**

- a) The cadastral map has to be updated as regards the network of drainage lines (gullies/*nalas/hallas*) and existing structures are marked to the scale and storage capacity of the existing water bodies are documented.
- b) The drainage line will be demarcated into Upper Reach, Middle Reach and Lower Reach.
- c) Considering the Catchment, *Nala* bed and bank conditions, suitable structures are decided.
- d) Number of storage structures (Check dam/ *Nala* bund/ Percolation tank) will be decided considering the commitments and available runoff in water budgeting and quality of water in the wells and site suitability.
- e) Detailed Leveling Survey using Dumpy Level / Total Station has to be carried out to arrive at the site-specific designs as shown in the Manual.
- f) The location of ground water recharge structures are decided by examining the lineaments and fracture zones from geological maps.
- g) Rainfall intensity data of the nearest Rain Gauge Station is considered for Hydrologic Designs.
- h) Silt load to the Storage/Recharge Structures is reduced by providing vegetative, boulder and earthen checks in the natural water course. Location and design details are given in the Manual.

### **9.2 Recommended Soil and Water Conservation Measures**

The appropriate conservation structures best suited for each of the land parcel/ survey number (Appendix-I) are selected based on the slope per cent, severity of erosion, amount of rainfall, land use and soil type. The different kinds of conservation structures recommended are

1. Graded / Strengthening of Bunds
2. Trench cum Bunds (TCB)
3. Trench cum Bunds / Strengthening
4. Crescent Bunds

A map (Fig. 9.2) showing soil and water conservation plan with different kinds of structures recommended has been prepared which shows the spatial distribution and extent of area. A maximum area of about 413 ha (90%) area requires Graded Bunding and about 31 ha (7%) area required Strengthening of existing bunds/bunding in the microwatershed. The conservation plan prepared may be presented to all the stakeholders including farmers and after including their suggestions, the conservation plan for the microwatershed may be finalised in a participatory approach.




Fig. 9.2 Soil and Water Conservation Plan map of Handrahalu-2 Microwatershed

### Greening of Microwatershed

As part of the greening programme in the watersheds, it is envisaged to plant a variety of horticultural and other tree plants that are edible, economical and produce lot of biomass which helps to restore the ecological balance in the watersheds. The lands that are suitable for greening programme are non-arable lands (land capability classes V, VI VII and VIII) and also the lands that are not suitable or marginally suitable for growing annual and perennial crops. The method of planting these trees is given below.

It is recommended to open the pits during the 1<sup>st</sup> week of March along the contour and heap the dug out soil on the lower side of the slope in order to harness the flowing water and facilitate weathering of soil in the pit. Exposure of soil in the pit also prevents spread of pests and diseases due to scorching sun rays. The pits should be filled with mixture of soil and organic manure during the second week of April and keep ready with sufficiently tall seedlings produced either in poly bags or in root trainer nurseries so that planting can be done during the 2<sup>nd</sup> or 3<sup>rd</sup> week of April depending on the rainfall.

The tree species suitable for the area considering rainfall, temperature and adaptability is listed below; waterlogged areas are recommended to be planted with species like Neral (*Syzgium cumini*) and Bamboo. Dry areas are to be planted with species like Honge, Bevu, Seetaphal *etc.*

<b>Dry Deciduous Species</b>			<b>Temp (°C)</b>	<b>Rainfall (mm)</b>
1.	Bevu	<i>Azadiracta indica</i>	21–32	400 –1,200
2.	Tapasi	<i>Holoptelia integrifolia</i>	20-30	500 - 1000
3.	Seetaphal	<i>Anona Squamosa</i>	20-40	400 - 1000
4.	Honge	<i>Pongamia pinnata</i>	20 -50	500–2,500
5.	Kamara	<i>Hardwickia binata</i>	25 -35	400 - 1000
6.	Bage	<i>Albezzia lebbek</i>	20 - 45	500 - 1000
7.	Ficus	<i>Ficus bengalensis</i>	20 - 50	500–2,500
8.	Sisso	<i>Dalbargia Sissoo</i>	20 - 50	500 -2000
9.	Ailanthus	<i>Ailanthus excelsa</i>	20 - 50	500 - 1000
10.	Hale	<i>Wrightia tinctoria</i>	25 - 45	500 - 1000
11.	Uded	<i>Steriospermum chelanoides</i>	25 - 45	500 -2000
12.	Dhupa	<i>Boswellia Serrata</i>	20 - 40	500 - 2000
13.	Nelli	<i>Emblica Officinalis</i>	20 - 50	500 -1500
14.	Honne	<i>Pterocarpus marsupium</i>	20 - 40	500 - 2000
<b>Moist Deciduous Species</b>			<b>Temp (°C)</b>	<b>Rainfall (mm)</b>
15.	Teak	<i>Tectona grandis</i>	20 - 50	500-5000
16.	Nandi	<i>Legarstroemia lanceolata</i>	20 - 40	500 - 4000
17.	Honne	<i>Pterocarpus marsupium</i>	20 - 40	500 - 3000
18.	Mathi	<i>Terminalia alata</i>	20 -50	500 - 2000
19.	Shivane	<i>Gmelina arboria</i>	20 -50	500 -2000
20.	Kindal	<i>T.Paniculata</i>	20 - 40	500 - 1500
21.	Beete	<i>Dalbargia latifolia</i>	20 - 40	500 - 1500
22.	Tare	<i>T. belerica</i>	20 - 40	500 - 2000
23.	Bamboo	<i>Bambusa arundinasia</i>	20 - 40	500 - 2500
24.	Bamboo	<i>Dendrocalamus strictus</i>	20 – 40	500 – 2500
25.	Muthuga	<i>Butea monosperma</i>	20 - 40	400 - 1500
26.	Hippe	<i>Madhuca latifolia</i>	20 - 40	500 - 2000
27.	Sandal	<i>Santalum album</i>	20 - 50	400 - 1000
28.	Nelli	<i>Emblica officinalis</i>	20 - 40	500 - 2000
29.	Nerale	<i>Sizyzium cumini</i>	20 - 40	500 - 2000
30.	Dhaman	<i>Grevia tilifolia</i>	20 - 40	500 - 2000
31.	Kaval	<i>Careya arborea</i>	20 - 40	500 - 2000
32.	Harada	<i>Terminalia chebula</i>	20 - 40	500 - 2000



## References

1. FAO (1976) Framework for Land Evaluation, Food and Agriculture Organization, Rome.72 pp.
2. FAO (1983) Guidelines for Land Evaluation for Rainfed Agriculture, FAO, Rome, 237 pp.
3. IARI (1971) Soil Survey Manual, All India Soil and Land Use Survey Organization, IARI, New Delhi, 121 pp.
4. Katyal, J.C. and Rattan, R.K. (2003) Secondary and Micronutrients; Research Gap and future needs. Fert. News 48 (4); 9-20.
5. Naidu, L.G.K., Ramamurthy, V., Challa, O., Hegde, R. and Krishnan, P. (2006) Manual Soil Site Suitability Criteria for Major Crops, NBSS Publ. No. 129, NBSS &LUP, Nagpur, 118 pp.
6. Natarajan, A. and Dipak Sarkar (2010) Field Guide for Soil Survey, National Bureau of Soil Survey and Land Use Planning (ICAR), Nagpur, India.
7. Natarajan, A., Rajendra Hegde, Raj, J.N. and Shivananda Murthy, H.G. (2015) Implementation Manual for Sujala-III Project, Watershed Development Department, Bengaluru, Karnataka.
8. Sarma, V.A.K., Krishnan, P. and Budihal, S.L. (1987) Laboratory Manual, Tech. Bull. 23, NBSS &LUP, Nagpur.
9. Sehgal, J.L. (1990) Soil Resource Mapping of Different States of India; Why and How?, National Bureau of Soil Survey and Land Use Planning, Nagpur, 49 pp.
10. Shivaprasad, C.R., R.S. Reddy, J. Sehgal and M. Velayuthum (1998) Soils of Karnataka for Optimizing Land Use, NBSS Publ. No. 47b, NBSS & LUP, Nagpur, India.
11. Soil Survey Staff (2006) Keys to Soil Taxonomy, Tenth edition, U.S. Department of Agriculture/ NRCS, Washington DC, U.S.A.
12. Soil Survey Staff (2012) Soil Survey Manual, Handbook No. 18, USDA, Washington DC, USA.





**Appendix-I**  
**Handrahalu-2 (1V2c) Microwatershed**  
**Soil Phase Information**

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Chithapura	51	2.28	NSPmB1	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Slight	Not Available	Not Available	IIs	Graded bunding
Chithapura	52	6.39	GRHmB2g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available	Not Available	IIses	Graded bunding
Chithapura	53	0.02	GRHmB2g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available	Not Available	IIses	Graded bunding
Chithapura	57	0.38	GRHmB2g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available	Not Available	IIses	Graded bunding
Hanavala	56	0.62	MTLmB2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Habitation	Not Available	IIIses	Graded bunding
Hanavala	57	6.96	MTLmB2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Sunflower+Fallow land+Redgram (Sf+Fl+Rg)	Not Available	IIIses	Graded bunding
Hanavala	58	6.93	NSPmB1g1	LMU-1	Moderately deep (75-100 cm)	Clay	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	IIs	Graded bunding
Hanavala	59	6.86	MTLmB2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Maize+Sunflower (Mz+Sf)	Not Available	IIIses	Graded bunding
Hanavala	60	1.24	MTLmB2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIIses	Graded bunding
Hanavala	61	7.19	MTLmB2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	1 Borewell	IIIses	Graded bunding
Hanavala	62	0.2	MTLmB2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIIses	Graded bunding
Hanavala	63	3.85	MTLmB2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	1 Borewell	IIIses	Graded bunding
Hanavala	64	0.73	MTLmB2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIIses	Graded bunding
Hanavala	65	0.44	MTLmB2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	1 Borewell	IIIses	Graded bunding
Hanavala	66	1.01	MTLmB2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIIses	Graded bunding
Hanavala	67	2.55	MTLmB2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	1 Borewell	IIIses	Graded bunding
Hanavala	68	4.19	MTLmB2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	1 Borewell	IIIses	Graded bunding
Hanavala	69	3.39	NSPmB1	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Slight	Bajra+Sunflower (Bj+Sf)	Not Available	IIs	Graded bunding
Hanavala	70	7.44	GRHmB2g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Maize+Sunflower (Mz+Sf)	1 Borewell	IIses	Graded bunding
Hanavala	71	3.59	GRHmB2g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Maize+Fallow land (Mz+Fl)	Not Available	IIses	Graded bunding
Hanavala	72	2.99	GRHmB2g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIses	Graded bunding
Hanavala	73	2.62	GRHmB2g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIses	Graded bunding

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Hanavala	74	2.85	GRHmB2g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Chilli+Onion (Ch+On)	Not Available	Iles	Graded bunding
Hanavala	75	2	KDTmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	2 Borewell	Iles	Graded bunding
Hanavala	76	1.18	KDTmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Onion (On)	1 Borewell	Iles	Graded bunding
Hanavala	77	0.19	KDTmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Chilli (Ch)	Not Available	Iles	Graded bunding
Hanavala	78	10.16	KDTmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Maize+Sunflower+Chilli (Mz+Sf+Ch)	Not Available	Iles	Graded bunding
Hanavala	79	4.43	GRHmB2g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	Iles	Graded bunding
Hanavala	80	4.97	KDTmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Maize+Sunflower+Onion (Mz+Sf+On)	Not Available	Iles	Graded bunding
Hanavala	83	1.92	GRHmB2g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	Iles	Graded bunding
Hanavala	84	7.2	GRHmB2g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	Iles	Graded bunding
Hanavala	85	8.12	GRHmB2g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Sunflower+Onion+Chilli (Sf+On+Ch)	Not Available	Iles	Graded bunding
Hanavala	86	8.31	NSPmB1	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Slight	Sunflower+Bajra+Maize (Sf+Bj+Mz)	Not Available	Ils	Graded bunding
Hanavala	87	6.81	NSPmB1	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Slight	Maize+Chilli (Mz+Ch)	1 Borewell	Ils	Graded bunding
Hanavala	88	7.73	NSPmB1	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	Ils	Graded bunding
Hanavala	89	5.67	NSPmB1	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Slight	Sunflower+Maize+Chilli (Sf+Mz+Ch)	Not Available	Ils	Graded bunding
Hanavala	90	0.46	NSPmB1g1	LMU-1	Moderately deep (75-100 cm)	Clay	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Slight	Onion+Maize+Sunflower (On+Mz+Sf)	Not Available	Ils	Graded bunding
Hanavala	102	0.27	NSPmB1	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Slight	Maize+Bengalgram+Sunflower (Mz+Bg+Sf)	Not Available	Ils	Graded bunding
Hanavala	103	2.75	NSPmB1	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Slight	Bengalgram+Maize+Fallow land (Bg+Mz+Fl)	Not Available	Ils	Graded bunding
Hanavala	104	4.73	NSPmB1	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Slight	Sunflower+Onion+Maize (Sf+On+Mz)	1 Farm pond	Ils	Graded bunding
Handrala	1	0.09	GRHmB1g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Habitation	Not Available	Ils	Graded bunding
Handrala	2	0.09	GRHmB1g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Habitation	Not Available	Ils	Graded bunding
Handrala	3	0.67	BDRmB1g2	LMU-1	Very deep (>150 cm)	Clay	Very gravelly (35-60%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Habitation	Not Available	Ils	Graded bunding
Handrala	4	0.77	BDRmB1g2	LMU-1	Very deep (>150 cm)	Clay	Very gravelly (35-60%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Fallow land (Fl)	Not Available	Ils	Graded bunding
Handrala	5	0.82	BDRmB1g2	LMU-1	Very deep (>150 cm)	Clay	Very gravelly (35-60%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Fallow land (Fl)	Not Available	Ils	Graded bunding
Handrala	6	1.35	BDRmB1g2	LMU-1	Very deep (>150 cm)	Clay	Very gravelly	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Habitation	Not Available	Ils	Graded bunding

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth cm)	Surface Soil Texture	Soil Gravelliness (35-60%)	Available Water Capacity mm/m)	Slope sloping (1-3%)	Soil Erosion	Current Land Use	Wells Available	Land Capability IIs	Conservation Plan bundling
Handrala	7	0.77	BDRmB1g2	LMU-1	Very deep (>150 cm)	Clay	Very gravelly (35-60%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Habitation	Not Available	IIs	Graded bundling
Handrala	8	0.78	BDRmB1g2	LMU-1	Very deep (>150 cm)	Clay	Very gravelly (35-60%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Fallow land (Fl)	Not Available	IIs	Graded bundling
Handrala	9	0.01	BDRmB1g2	LMU-1	Very deep (>150 cm)	Clay	Very gravelly (35-60%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Fallow land (Fl)	Not Available	IIs	Graded bundling
Handrala	66	0.18	NSPmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Maize+Onion (Mz+On)	Not Available	IIs	Graded bundling
Handrala	67	2.15	NSPmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Sunflower (Sf)	Not Available	IIs	Graded bundling
Handrala	68	4.85	NSPmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Sunflower+Onion (Sf+On)	Not Available	IIs	Graded bundling
Handrala	69	2.1	NSPmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIs	Graded bundling
Handrala	70	2.43	NSPmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Maize+Sunflower (Mz+Sf)	1 Borewell	IIs	Graded bundling
Handrala	78	2.46	GRHmB1g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Fallow land (Fl)	1 Borewell	IIs	Graded bundling
Handrala	79	1.41	NSPmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	1 Borewell	IIs	Graded bundling
Handrala	80	1.96	GRHmB1g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Onion (On)	Not Available	IIs	Graded bundling
Handrala	81	5.87	NSPmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Onion+Maize (On+Mz)	1 Borewell	IIs	Graded bundling
Handrala	82	2.48	NSPmB2g1	LMU-1	Moderately deep (75-100 cm)	Clay	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIs	Graded bundling
Handrala	83	2.94	NSPmB2g1	LMU-1	Moderately deep (75-100 cm)	Clay	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Maize+Onion (Mz+On)	Not Available	IIs	Graded bundling
Handrala	84	3.28	NSPmB2g1	LMU-1	Moderately deep (75-100 cm)	Clay	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIs	Graded bundling
Handrala	85	1.94	NSPmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut (Gn)	Not Available	IIs	Graded bundling
Handrala	86	6.69	NSPmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Onion+Maize (On+Mz)	Not Available	IIs	Graded bundling
Handrala	87	5.36	NSPmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Maize+Bengalgram (Mz+Bg)	Not Available	IIs	Graded bundling
Handrala	88	3.4	NSPmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIs	Graded bundling
Handrala	89	2.55	NSPmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Sunflower (Sf)	Not Available	IIs	Graded bundling
Handrala	90	6.81	NSPmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIs	Graded bundling
Handrala	91	3.76	NSPmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Maize+Sunflower (Mz+Sf)	Not Available	IIs	Graded bundling
Handrala	92	5.25	NSPmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Maize+Onion (Mz+On)	Not Available	IIs	Graded bundling

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Handrala	93	8.11	HDLmB2g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Maize+Onion (Mz+On)	Not Available	Iles	Graded bunding
Handrala	94	6.96	NSPmB2g1	LMU-1	Moderately deep (75-100 cm)	Clay	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Sunflower+Maize (Sf+Mz)	Not Available	Iles	Graded bunding
Handrala	95	5.52	NSPmB2g1	LMU-1	Moderately deep (75-100 cm)	Clay	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	Iles	Graded bunding
Handrala	96	2.41	NSPmB2g1	LMU-1	Moderately deep (75-100 cm)	Clay	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Maize+Onion (Mz+On)	1 Borewell	Iles	Graded bunding
Handrala	97	1.76	NSPmB2g1	LMU-1	Moderately deep (75-100 cm)	Clay	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	Iles	Graded bunding
Handrala	98	1.47	NSPmB2g1	LMU-1	Moderately deep (75-100 cm)	Clay	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	Iles	Graded bunding
Handrala	99	7.97	NSPmB2g1	LMU-1	Moderately deep (75-100 cm)	Clay	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Maize+Onion (Mz+On)	1 Borewell	Iles	Graded bunding
Handrala	100	5.64	NSPmB2g1	LMU-1	Moderately deep (75-100 cm)	Clay	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Maize+Onion+Red gram (Mz+On+Rg)	Not Available	Iles	Graded bunding
Handrala	101	4.92	RNKmB1g1	LMU-2	Moderately shallow (50-75 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Maize+Redgram (Mz+Rg)	Not Available	IIs	Graded bunding
Handrala	102	4.36	HDLmB2g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Maize+Onion (Mz+On)	Not Available	Iles	Graded bunding
Handrala	103	5.06	RNKmB1g1	LMU-2	Moderately shallow (50-75 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Bajra+Maize (Bj+Mz)	1 Borewell	IIs	Graded bunding
Handrala	104	7.94	RNKmB1g1	LMU-2	Moderately shallow (50-75 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Current fallow land+Maize (Cfl+Mz)	Not Available	IIs	Graded bunding
Handrala	105	1.97	RNKmB1g1	LMU-2	Moderately shallow (50-75 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Bajra (Bj)	Not Available	IIs	Graded bunding
Handrala	106	3.74	RNKmB1g1	LMU-2	Moderately shallow (50-75 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Bajra+Sunflower (Bj+Sf)	Not Available	IIs	Graded bunding
Handrala	107	7.29	HDLmB2g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Bengalgram+Maize (Bg+Mz)	Not Available	Iles	Graded bunding
Handrala	108	3.25	RNKmB1	LMU-2	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Bengalgram+Maize (Bg+Mz)	Not Available	IIs	Graded bunding
Handrala	109	6.93	RNKmB1	LMU-2	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Maize+Redgram (Mz+Rg)	Not Available	IIs	Graded bunding
Handrala	110	5.07	RNKmB1g1	LMU-2	Moderately shallow (50-75 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Maize+Fallow land (Mz+Fl)	Not Available	IIs	Graded bunding
Handrala	111	0.2	RNKmB1	LMU-2	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Redgram+Maize (Rg+Mz)	Not Available	IIs	Graded bunding
Handrala	112	1.52	RNKmB1	LMU-2	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	IIs	Graded bunding
Handrala	113	5.66	RNKmB1	LMU-2	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Maize+Redgram (Mz+Rg)	Not Available	IIs	Graded bunding
Handrala	114	0.14	RNKmB1	LMU-2	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Maize+Onion (Mz+On)	Not Available	IIs	Graded bunding
Handrala	147	3.48	BGTmB1g2	LMU-4	Very shallow (<25 cm)	Clay	Very gravelly (35-60%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Slight	Onion+Redgram (On+Rg)	Not Available	IVs	Graded bunding
Handrala	148	0.97	NSPmB2g1	LMU-1	Moderately deep (75-100 cm)	Clay	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Fallow land+Maze (Fl+Mz)	Not Available	Iles	Graded bunding

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Handrala	150	0.49	NSPmB2g1	LMU-1	Moderately deep (75-100 cm)	Clay	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Fallow land (Fl)	Not Available	Iies	Graded bunding
Handrala	152	0.14	RNKmB1	LMU-2	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Maize+Onion (Mz+On)	Not Available	Iis	Graded bunding
Handrala	153	4.35	NSPmB2g1	LMU-1	Moderately deep (75-100 cm)	Clay	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	Iies	Graded bunding
Handrala	154	0.65	NSPmB2g1	LMU-1	Moderately deep (75-100 cm)	Clay	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iies	Graded bunding
Handrala	155	1.4	NSPmB2g1	LMU-1	Moderately deep (75-100 cm)	Clay	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Sunflower (Sf)	Not Available	Iies	Graded bunding
Handrala	156	0.06	NSPmB2g1	LMU-1	Moderately deep (75-100 cm)	Clay	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Sunflower (Sf)	Not Available	Iies	Graded bunding
Handrala	157	4.96	BGTmB1g2	LMU-4	Very shallow (<25 cm)	Clay	Very gravelly (35-60%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Slight	Fallow land (Fl)	Not Available	IVs	Graded bunding
Handrala	158	1.54	HDLmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Fallow land (Fl)	1 Borewell	Iis	Graded bunding
Handrala	159	4.3	HDLmB2	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	2 Borewell	Iies	Graded bunding
Handrala	160	0.97	HDLmB2	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	Iies	Graded bunding
Handrala	161	1.15	Waterbody	Others	Others	Others	Others	Others	Others	Others	Maize (Mz)	Not Available	Others	Others
Handrala	162	0.4	HDLmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	Iis	Graded bunding
Handrala	163	5.88	HDLmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Maize+Redgram+Onion (Mz+Rg+On)	1 Farm pond, 1 Borewell	Iis	Graded bunding
Handrala	164	6.67	HDLmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Onion+Maize+Fallow land (On+Mz+Fl)	1 Borewell	Iis	Graded bunding
Handrala	165	3.24	HDLmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Maize+Onion (Mz+On)	Not Available	Iis	Graded bunding
Handrala	166	4.13	HDLmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	Iis	Graded bunding
Handrala	167	1.63	HDLmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	1 Borewell	Iis	Graded bunding
Handrala	168	3.16	HDLmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Maize+Onion (Mz+On)	Not Available	Iis	Graded bunding
Handrala	169	4.49	DRLmB1	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Slight	Maize+Onion (Mz+On)	1 Borewell	Iis	Graded bunding
Handrala	170	1.53	DRLmB1	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	Iis	Graded bunding
Handrala	171	1.59	NSPmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	Iies	Graded bunding
Handrala	172	1.18	DRLmB1	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Slight	Sunflower (Sf)	Not Available	Iis	Graded bunding
Handrala	173	2.04	DRLmB1	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Slight	Sunflower (Sf)	Not Available	Iis	Graded bunding
Handrala	174	1.67	DRLmB1	LMU-1	Moderately deep	Clay	Non gravelly	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Slight	Sunflower (Sf)	Not Available	Iis	Graded bunding

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
					(75-100 cm)		(<15%)	mm/m)	sloping (1-3%)			Available		bunding
Handrala	175	1.06	DRLmB1	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Slight	Sunflower (Sf)	Not Available	IIs	Graded bunding
Handrala	176	0.11	NSPmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIs	Graded bunding
Handrala	177	1.76	DRLmB1	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	1 Borewell	IIs	Graded bunding
Handrala	178	1.42	DRLmB1	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	IIs	Graded bunding
Handrala	179	2.24	DRLmB1	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	IIs	Graded bunding
Handrala	180	0.36	DRLmB1	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Slight	Fallow land (Fl)	Not Available	IIs	Graded bunding
Handrala	181	2.83	DRLmB1	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Slight	Fallow land (Fl)	Not Available	IIs	Graded bunding
Hireshindh ogi	273	1.08	MLRmA1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Maize (Mz)	Not Available	IIs	Graded bunding
Hireshindh ogi	274	4.3	MLRmA1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Sunflower (Sf)	Not Available	IIs	Graded bunding
Hireshindh ogi	275	1.89	MLRmA1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Sunflower (Sf)	Not Available	IIs	Graded bunding
Hireshindh ogi	276	8.92	MLRmA1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Maize (Mz)	Not Available	IIs	Graded bunding
Hireshindh ogi	277	6.84	MLRmA1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Maize (Mz)	Not Available	IIs	Graded bunding
Hireshindh ogi	278	2.45	MLRmA1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Bengalgram (Bg)	Not Available	IIs	Graded bunding
Hireshindh ogi	279	0.83	MLRmA1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Maize (Mz)	Not Available	IIs	Graded bunding
Hireshindh ogi	280	0.38	MLRmA1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Maize (Mz)	Not Available	IIs	Graded bunding
Hireshindh ogi	281	0.12	MLRmA1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Cotton (Ct)	Not Available	IIs	Graded bunding
Hireshindh ogi	284	0.07	MLRmA1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Maize (Mz)	Not Available	IIs	Graded bunding
Hireshindh ogi	285	3.94	MLRmA1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Maize (Mz)	Not Available	IIs	Graded bunding
Hireshindh ogi	286	3.03	MLRmA1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Maize (Mz)	Not Available	IIs	Graded bunding
Hireshindh ogi	287	0.88	MLRmA1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Maize (Mz)	Not Available	IIs	Graded bunding
Hireshindh ogi	288	1.02	MLRmA1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Maize (Mz)	Not Available	IIs	Graded bunding
Hireshindh ogi	289	1.82	MLRmA1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Maize (Mz)	Not Available	IIs	Graded bunding
Hireshindh ogi	290	0.09	MLRmA1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Maize (Mz)	Not Available	IIs	Graded bunding

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Hireshindh ogi	293	0.56	MLRmA1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Maize (Mz)	Not Available	IIs	Graded bunding
Hireshindh ogi	294	1.01	NSPmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIs	Graded bunding
Hireshindh ogi	295	1.63	NSPmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIs	Graded bunding
Narasapura	33	0.28	NSPmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Maize+Current fallow land (Mz+Cf)	Not Available	IIs	Graded bunding

















Village	Survey No	Soil Reaction	Salinity	Organic Carbon	Available Phosphorus	Available Potassium	Available Sulphur	Available Boron	Available Iron	Available Manganese	Available Copper	Available Zinc
Hireshindh ogi	295	Strongly alkaline (pH 8.4 - 9.0)	Non saline (<2 dsm )	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	High (> 337 kg/ha)	Medium (10 - 20 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Narasapura	33	Strongly alkaline (pH 8.4 - 9.0)	Non saline (<2 dsm )	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	High (> 337 kg/ha)	Medium (10 - 20 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)





**Appendix III**  
**Handrahalu-2 (1V2c) Microwatershed**  
**Soil Suitability Information**

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Jasmine	Bhendi	Brinjal	Crossandra	Drumstick	Mulberry	Onion
Chithapura	51	S3rt	S2tz	S3rt	S1	S3t	S1	S3r	S2r	S1	S2r	S2t	S3t	S1	N1t	S3rt	S2r	S3t	S3t	S3t	S2t	S2t	S2rt	S2t	S3t	S2tz	S2tz	S3t	S2rt	S2t	S3tz	
Chithapura	52	S3t	S2t	S3t	S2g	S3t	S2g	S2rg	S2g	S2g	S2g	S2gt	S2gt	S3t	S2g	N1t	S2rt	S2g	S3t	S3t	S3t	S2tg	S2tg	S2tg	S2t	S3t	S2t	S2t	S3t	S2tg	S2gt	S3t
Chithapura	53	S3t	S2t	S3t	S2g	S3t	S2g	S2rg	S2g	S2g	S2g	S2gt	S2gt	S3t	S2g	N1t	S2rt	S2g	S3t	S3t	S3t	S2tg	S2tg	S2tg	S2t	S3t	S2t	S2t	S3t	S2tg	S2gt	S3t
Chithapura	57	S3t	S2t	S3t	S2g	S3t	S2g	S2rg	S2g	S2g	S2g	S2gt	S2gt	S3t	S2g	N1t	S2rt	S2g	S3t	S3t	S3t	S2tg	S2tg	S2tg	S2t	S3t	S2t	S2t	S3t	S2tg	S2gt	S3t
Hanavala	56	N1rt	S3tz	N1rz	S3rz	N1rt	S3rz	N1rz	N1rz	S3rz	N1rz	N1rz	S3tz	N1rt	S3zg	N1rt	N1rt	N1rz	S3tz	S3rz	S3rz	S3rz	S3rz	N1rz	S3rz	S3rz	S3r	S3r	S3rz	N1rz	N1rz	S3rt
Hanavala	57	N1rt	S3tz	N1rz	S3rz	N1rt	S3rz	N1rz	N1rz	S3rz	N1rz	N1rz	S3tz	N1rt	S3zg	N1rt	N1rt	N1rz	S3tz	S3rz	S3rz	S3rz	S3rz	N1rz	S3rz	S3rz	S3r	S3r	S3rz	N1rz	N1rz	S3rt
Hanavala	58	S3rt	S2tz	S3rt	S1	S3t	S1	S3r	S2rg	S1	S2rg	S2t	S2gt	S3t	S1	N1t	S3rt	S2rg	S3t	S3t	S3t	S2tg	S2tg	S2rt	S2t	S3t	S2tz	S2tz	S3t	S2rt	S2gt	S3tz
Hanavala	59	N1rt	S3tz	N1rz	S3rz	N1rt	S3rz	N1rz	N1rz	S3rz	N1rz	N1rz	S3tz	N1rt	S3zg	N1rt	N1rt	N1rz	S3tz	S3rz	S3rz	S3rz	S3rz	N1rz	S3rz	S3rz	S3r	S3r	S3rz	N1rz	N1rz	S3rt
Hanavala	60	N1rt	S3tz	N1rz	S3rz	N1rt	S3rz	N1rz	N1rz	S3rz	N1rz	N1rz	S3tz	N1rt	S3zg	N1rt	N1rt	N1rz	S3tz	S3rz	S3rz	S3rz	S3rz	N1rz	S3rz	S3rz	S3r	S3r	S3rz	N1rz	N1rz	S3rt
Hanavala	61	N1rt	S3tz	N1rz	S3rz	N1rt	S3rz	N1rz	N1rz	S3rz	N1rz	N1rz	S3tz	N1rt	S3zg	N1rt	N1rt	N1rz	S3tz	S3rz	S3rz	S3rz	S3rz	N1rz	S3rz	S3rz	S3r	S3r	S3rz	N1rz	N1rz	S3rt
Hanavala	62	N1rt	S3tz	N1rz	S3rz	N1rt	S3rz	N1rz	N1rz	S3rz	N1rz	N1rz	S3tz	N1rt	S3zg	N1rt	N1rt	N1rz	S3tz	S3rz	S3rz	S3rz	S3rz	N1rz	S3rz	S3rz	S3r	S3r	S3rz	N1rz	N1rz	S3rt
Hanavala	63	N1rt	S3tz	N1rz	S3rz	N1rt	S3rz	N1rz	N1rz	S3rz	N1rz	N1rz	S3tz	N1rt	S3zg	N1rt	N1rt	N1rz	S3tz	S3rz	S3rz	S3rz	S3rz	N1rz	S3rz	S3rz	S3r	S3r	S3rz	N1rz	N1rz	S3rt
Hanavala	64	N1rt	S3tz	N1rz	S3rz	N1rt	S3rz	N1rz	N1rz	S3rz	N1rz	N1rz	S3tz	N1rt	S3zg	N1rt	N1rt	N1rz	S3tz	S3rz	S3rz	S3rz	S3rz	N1rz	S3rz	S3rz	S3r	S3r	S3rz	N1rz	N1rz	S3rt
Hanavala	65	N1rt	S3tz	N1rz	S3rz	N1rt	S3rz	N1rz	N1rz	S3rz	N1rz	N1rz	S3tz	N1rt	S3zg	N1rt	N1rt	N1rz	S3tz	S3rz	S3rz	S3rz	S3rz	N1rz	S3rz	S3rz	S3r	S3r	S3rz	N1rz	N1rz	S3rt
Hanavala	66	N1rt	S3tz	N1rz	S3rz	N1rt	S3rz	N1rz	N1rz	S3rz	N1rz	N1rz	S3tz	N1rt	S3zg	N1rt	N1rt	N1rz	S3tz	S3rz	S3rz	S3rz	S3rz	N1rz	S3rz	S3rz	S3r	S3r	S3rz	N1rz	N1rz	S3rt
Hanavala	67	N1rt	S3tz	N1rz	S3rz	N1rt	S3rz	N1rz	N1rz	S3rz	N1rz	N1rz	S3tz	N1rt	S3zg	N1rt	N1rt	N1rz	S3tz	S3rz	S3rz	S3rz	S3rz	N1rz	S3rz	S3rz	S3r	S3r	S3rz	N1rz	N1rz	S3rt
Hanavala	68	N1rt	S3tz	N1rz	S3rz	N1rt	S3rz	N1rz	N1rz	S3rz	N1rz	N1rz	S3tz	N1rt	S3zg	N1rt	N1rt	N1rz	S3tz	S3rz	S3rz	S3rz	S3rz	N1rz	S3rz	S3rz	S3r	S3r	S3rz	N1rz	N1rz	S3rt
Hanavala	69	S3rt	S2tz	S3rt	S1	S3t	S1	S3r	S2r	S1	S2r	S2t	S3t	S1	N1t	S3rt	S2r	S3t	S3t	S3t	S2t	S2t	S2rt	S2t	S3t	S2tz	S2tz	S3t	S2rt	S2t	S3tz	
Hanavala	70	S3t	S2t	S3t	S2g	S3t	S2g	S2rg	S2g	S2g	S2g	S2gt	S2gt	S3t	S2g	N1t	S2rt	S2g	S3t	S3t	S3t	S2tg	S2tg	S2tg	S2t	S3t	S2t	S2t	S3t	S2tg	S2gt	S3t
Hanavala	71	S3t	S2t	S3t	S2g	S3t	S2g	S2rg	S2g	S2g	S2g	S2gt	S2gt	S3t	S2g	N1t	S2rt	S2g	S3t	S3t	S3t	S2tg	S2tg	S2tg	S2t	S3t	S2t	S2t	S3t	S2tg	S2gt	S3t
Hanavala	72	S3t	S2t	S3t	S2g	S3t	S2g	S2rg	S2g	S2g	S2g	S2gt	S2gt	S3t	S2g	N1t	S2rt	S2g	S3t	S3t	S3t	S2tg	S2tg	S2tg	S2t	S3t	S2t	S2t	S3t	S2tg	S2gt	S3t
Hanavala	73	S3t	S2t	S3t	S2g	S3t	S2g	S2rg	S2g	S2g	S2g	S2gt	S2gt	S3t	S2g	N1t	S2rt	S2g	S3t	S3t	S3t	S2tg	S2tg	S2tg	S2t	S3t	S2t	S2t	S3t	S2tg	S2gt	S3t
Hanavala	74	S3t	S2t	S3t	S2g	S3t	S2g	S2rg	S2g	S2g	S2g	S2gt	S2gt	S3t	S2g	N1t	S2rt	S2g	S3t	S3t	S3t	S2tg	S2tg	S2tg	S2t	S3t	S2t	S2t	S3t	S2tg	S2gt	S3t
Hanavala	75	S3t	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2t	S1	S3t	S3t	S3t	S2t	S2t	S2t	S2t	S3t	S2t	S2t	S2t	S2t	S2t	S3t

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Jasmine	Bhendi	Brinjal	Crossandra	Drumstick	Mulberry	Onion
Hanavala	76	S3t	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2t	S1	S3t	S3t	S3t	S2t	S2t	S2t	S2t	S3t	S2t	S2t	S2t	S2t	S2t	S3t
Hanavala	77	S3t	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2t	S1	S3t	S3t	S3t	S2t	S2t	S2t	S2t	S3t	S2t	S2t	S2t	S2t	S2t	S3t
Hanavala	78	S3t	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2t	S1	S3t	S3t	S3t	S2t	S2t	S2t	S2t	S3t	S2t	S2t	S2t	S2t	S2t	S3t
Hanavala	79	S3t	S2t	S3t	S2g	S3t	S2g	S2rg	S2g	S2g	S2g	S2gt	S2gt	S3t	S2g	N1t	S2rt	S2g	S3t	S3t	S3t	S2tg	S2tg	S2tg	S2t	S3t	S2t	S2t	S3t	S2tg	S2gt	S3t
Hanavala	80	S3t	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2t	S1	S3t	S3t	S3t	S2t	S2t	S2t	S2t	S3t	S2t	S2t	S2t	S2t	S2t	S3t
Hanavala	83	S3t	S2t	S3t	S2g	S3t	S2g	S2rg	S2g	S2g	S2g	S2gt	S2gt	S3t	S2g	N1t	S2rt	S2g	S3t	S3t	S3t	S2tg	S2tg	S2tg	S2t	S3t	S2t	S2t	S3t	S2tg	S2gt	S3t
Hanavala	84	S3t	S2t	S3t	S2g	S3t	S2g	S2rg	S2g	S2g	S2g	S2gt	S2gt	S3t	S2g	N1t	S2rt	S2g	S3t	S3t	S3t	S2tg	S2tg	S2tg	S2t	S3t	S2t	S2t	S3t	S2tg	S2gt	S3t
Hanavala	85	S3t	S2t	S3t	S2g	S3t	S2g	S2rg	S2g	S2g	S2g	S2gt	S2gt	S3t	S2g	N1t	S2rt	S2g	S3t	S3t	S3t	S2tg	S2tg	S2tg	S2t	S3t	S2t	S2t	S3t	S2tg	S2gt	S3t
Hanavala	86	S3rt	S2tz	S3rt	S1	S3t	S1	S3r	S2r	S1	S2r	S2t	S2t	S3t	S1	N1t	S3rt	S2r	S3t	S3t	S3t	S2t	S2t	S2rt	S2t	S3t	S2tz	S2tz	S3t	S2rt	S2t	S3tz
Hanavala	87	S3rt	S2tz	S3rt	S1	S3t	S1	S3r	S2r	S1	S2r	S2t	S2t	S3t	S1	N1t	S3rt	S2r	S3t	S3t	S3t	S2t	S2t	S2rt	S2t	S3t	S2tz	S2tz	S3t	S2rt	S2t	S3tz
Hanavala	88	S3rt	S2tz	S3rt	S1	S3t	S1	S3r	S2r	S1	S2r	S2t	S2t	S3t	S1	N1t	S3rt	S2r	S3t	S3t	S3t	S2t	S2t	S2rt	S2t	S3t	S2tz	S2tz	S3t	S2rt	S2t	S3tz
Hanavala	89	S3rt	S2tz	S3rt	S1	S3t	S1	S3r	S2r	S1	S2r	S2t	S2t	S3t	S1	N1t	S3rt	S2r	S3t	S3t	S3t	S2t	S2t	S2rt	S2t	S3t	S2tz	S2tz	S3t	S2rt	S2t	S3tz
Hanavala	90	S3rt	S2tz	S3rt	S1	S3t	S1	S3r	S2rg	S1	S2rg	S2t	S2gt	S3t	S1	N1t	S3rt	S2rg	S3t	S3t	S3t	S2tg	S2tg	S2rt	S2t	S3t	S2tz	S2tz	S3t	S2rt	S2gt	S3tz
Hanavala	102	S3rt	S2tz	S3rt	S1	S3t	S1	S3r	S2r	S1	S2r	S2t	S2t	S3t	S1	N1t	S3rt	S2r	S3t	S3t	S3t	S2t	S2t	S2rt	S2t	S3t	S2tz	S2tz	S3t	S2rt	S2t	S3tz
Hanavala	103	S3rt	S2tz	S3rt	S1	S3t	S1	S3r	S2r	S1	S2r	S2t	S2t	S3t	S1	N1t	S3rt	S2r	S3t	S3t	S3t	S2t	S2t	S2rt	S2t	S3t	S2tz	S2tz	S3t	S2rt	S2t	S3tz
Hanavala	104	S3rt	S2tz	S3rt	S1	S3t	S1	S3r	S2r	S1	S2r	S2t	S2t	S3t	S1	N1t	S3rt	S2r	S3t	S3t	S3t	S2t	S2t	S2rt	S2t	S3t	S2tz	S2tz	S3t	S2rt	S2t	S3tz
Handrala	1	S3t	S2t	S3t	S2g	S3t	S2g	S2rg	S2g	S2g	S2g	S2gt	S2gt	S3t	S2g	N1t	S2rt	S2g	S3t	S3t	S3t	S2tg	S2tg	S2tg	S2t	S3t	S2t	S2t	S3tg	S2tg	S2gt	S3t
Handrala	2	S3t	S2t	S3t	S2g	S3t	S2g	S2rg	S2g	S2g	S2g	S2gt	S2gt	S3t	S2g	N1t	S2rt	S2g	S3t	S3t	S3t	S2tg	S2tg	S2tg	S2t	S3t	S2t	S2t	S3tg	S2tg	S2gt	S3t
Handrala	3	S3tg	S2t	S3tg	S3g	S3tg	S3tg	S2tg	S2g	S3tg	S3g	S3tg	S2gt	S3tg	S2gt	N1tg	S3gt	S2g	S3tg	S3tg	S3tg	S3tg	S3tg	S2tg	S3tg	S3tg	S2tg	S2tg	S3tg	S2tg	S3t	S3tg
Handrala	4	S3tg	S2t	S3tg	S3g	S3tg	S3tg	S2tg	S2g	S3tg	S3g	S3tg	S2gt	S3tg	S2gt	N1tg	S3gt	S2g	S3tg	S3tg	S3tg	S3tg	S3tg	S2tg	S3tg	S3tg	S2tg	S2tg	S3tg	S2tg	S3t	S3tg
Handrala	5	S3tg	S2t	S3tg	S3g	S3tg	S3tg	S2tg	S2g	S3tg	S3g	S3tg	S2gt	S3tg	S2gt	N1tg	S3gt	S2g	S3tg	S3tg	S3tg	S3tg	S3tg	S2tg	S3tg	S3tg	S2tg	S2tg	S3tg	S2tg	S3t	S3tg
Handrala	6	S3tg	S2t	S3tg	S3g	S3tg	S3tg	S2tg	S2g	S3tg	S3g	S3tg	S2gt	S3tg	S2gt	N1tg	S3gt	S2g	S3tg	S3tg	S3tg	S3tg	S3tg	S2tg	S3tg	S3tg	S2tg	S2tg	S3tg	S2tg	S3t	S3tg
Handrala	7	S3tg	S2t	S3tg	S3g	S3tg	S3tg	S2tg	S2g	S3tg	S3g	S3tg	S2gt	S3tg	S2gt	N1tg	S3gt	S2g	S3tg	S3tg	S3tg	S3tg	S3tg	S2tg	S3tg	S3tg	S2tg	S2tg	S3tg	S2tg	S3t	S3tg
Handrala	8	S3tg	S2t	S3tg	S3g	S3tg	S3tg	S2tg	S2g	S3tg	S3g	S3tg	S2gt	S3tg	S2gt	N1tg	S3gt	S2g	S3tg	S3tg	S3tg	S3tg	S3tg	S2tg	S3tg	S3tg	S2tg	S2tg	S3tg	S2tg	S3t	S3tg
Handrala	9	S3tg	S2t	S3tg	S3g	S3tg	S3tg	S2tg	S2g	S3tg	S3g	S3tg	S2gt	S3tg	S2gt	N1tg	S3gt	S2g	S3tg	S3tg	S3tg	S3tg	S3tg	S2tg	S3tg	S3tg	S2tg	S2tg	S3tg	S2tg	S3t	S3tg
Handrala	66	S3rt	S2tz	S3rt	S1	S3t	S1	S3r	S2r	S1	S2r	S2t	S2t	S3t	S1	N1t	S3rt	S2r	S3t	S3t	S3tw	S2tw	S2rt	S2t	S3tw	S2tz	S2tz	S3t	S2rt	S2tw	S3tz	

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Jasmine	Bhendi	Brinjal	Crossandra	Drumstick	Mulberry	Onion
Handrala	67	S3rt	S2tz	S3rt	S1	S3t	S1	S3r	S2r	S1	S2r	S2t	S2t	S3t	S1	N1t	S3rt	S2r	S3t	S3t	S3tw	S2tw	S2tw	S2rt	S2t	S3tw	S2tz	S2tz	S3t	S2rt	S2tw	S3tz
Handrala	68	S3rt	S2tz	S3rt	S1	S3t	S1	S3r	S2r	S1	S2r	S2t	S2t	S3t	S1	N1t	S3rt	S2r	S3t	S3t	S3tw	S2tw	S2tw	S2rt	S2t	S3tw	S2tz	S2tz	S3t	S2rt	S2tw	S3tz
Handrala	69	S3rt	S2tz	S3rt	S1	S3t	S1	S3r	S2r	S1	S2r	S2t	S2t	S3t	S1	N1t	S3rt	S2r	S3t	S3t	S3tw	S2tw	S2tw	S2rt	S2t	S3tw	S2tz	S2tz	S3t	S2rt	S2tw	S3tz
Handrala	70	S3rt	S2tz	S3rt	S1	S3t	S1	S3r	S2r	S1	S2r	S2t	S2t	S3t	S1	N1t	S3rt	S2r	S3t	S3t	S3tw	S2tw	S2tw	S2rt	S2t	S3tw	S2tz	S2tz	S3t	S2rt	S2tw	S3tz
Handrala	78	S3t	S2t	S3t	S2g	S3t	S2g	S2rg	S2g	S2g	S2g	S2gt	S2gt	S3t	S2g	N1t	S2rt	S2g	S3t	S3t	S3t	S2tg	S2tg	S2tg	S2t	S3t	S2t	S2t	S3tg	S2tg	S2gt	S3t
Handrala	79	S3rt	S2tz	S3rt	S1	S3t	S1	S3r	S2r	S1	S2r	S2t	S2t	S3t	S1	N1t	S3rt	S2r	S3t	S3t	S3tw	S2tw	S2tw	S2rt	S2t	S3tw	S2tz	S2tz	S3t	S2rt	S2tw	S3tz
Handrala	80	S3t	S2t	S3t	S2g	S3t	S2g	S2rg	S2g	S2g	S2g	S2gt	S2gt	S3t	S2g	N1t	S2rt	S2g	S3t	S3t	S3t	S2tg	S2tg	S2tg	S2t	S3t	S2t	S2t	S3tg	S2tg	S2gt	S3t
Handrala	81	S3rt	S2tz	S3rt	S1	S3t	S1	S3r	S2r	S1	S2r	S2t	S2t	S3t	S1	N1t	S3rt	S2r	S3t	S3t	S3tw	S2tw	S2tw	S2rt	S2t	S3tw	S2tz	S2tz	S3t	S2rt	S2tw	S3tz
Handrala	82	S3rt	S2tz	S3rt	S1	S3t	S1	S3rg	S2r	S1	S2r	S2t	S2gt	S3t	S1	N1t	S3rt	S2r	S3t	S3t	S3tw	S2tw	S2tw	S2rt	S2t	S3tw	S2tz	S2tz	S3t	S2rt	S2tw	S3tz
Handrala	83	S3rt	S2tz	S3rt	S1	S3t	S1	S3rg	S2r	S1	S2r	S2t	S2gt	S3t	S1	N1t	S3rt	S2r	S3t	S3t	S3tw	S2tw	S2tw	S2rt	S2t	S3tw	S2tz	S2tz	S3t	S2rt	S2tw	S3tz
Handrala	84	S3rt	S2tz	S3rt	S1	S3t	S1	S3rg	S2r	S1	S2r	S2t	S2gt	S3t	S1	N1t	S3rt	S2r	S3t	S3t	S3tw	S2tw	S2tw	S2rt	S2t	S3tw	S2tz	S2tz	S3t	S2rt	S2tw	S3tz
Handrala	85	S3rt	S2tz	S3rt	S1	S3t	S1	S3r	S2r	S1	S2r	S2t	S2t	S3t	S1	N1t	S3rt	S2r	S3t	S3t	S3tw	S2tw	S2tw	S2rt	S2t	S3tw	S2tz	S2tz	S3t	S2rt	S2tw	S3tz
Handrala	86	S3rt	S2tz	S3rt	S1	S3t	S1	S3r	S2r	S1	S2r	S2t	S2t	S3t	S1	N1t	S3rt	S2r	S3t	S3t	S3tw	S2tw	S2tw	S2rt	S2t	S3tw	S2tz	S2tz	S3t	S2rt	S2tw	S3tz
Handrala	87	S3rt	S2tz	S3rt	S1	S3t	S1	S3r	S2r	S1	S2r	S2t	S2t	S3t	S1	N1t	S3rt	S2r	S3t	S3t	S3tw	S2tw	S2tw	S2rt	S2t	S3tw	S2tz	S2tz	S3t	S2rt	S2tw	S3tz
Handrala	88	S3rt	S2tz	S3rt	S1	S3t	S1	S3r	S2r	S1	S2r	S2t	S2t	S3t	S1	N1t	S3rt	S2r	S3t	S3t	S3tw	S2tw	S2tw	S2rt	S2t	S3tw	S2tz	S2tz	S3t	S2rt	S2tw	S3tz
Handrala	89	S3rt	S2tz	S3rt	S1	S3t	S1	S3r	S2r	S1	S2r	S2t	S2t	S3t	S1	N1t	S3rt	S2r	S3t	S3t	S3tw	S2tw	S2tw	S2rt	S2t	S3tw	S2tz	S2tz	S3t	S2rt	S2tw	S3tz
Handrala	90	S3rt	S2tz	S3rt	S1	S3t	S1	S3r	S2r	S1	S2r	S2t	S2t	S3t	S1	N1t	S3rt	S2r	S3t	S3t	S3tw	S2tw	S2tw	S2rt	S2t	S3tw	S2tz	S2tz	S3t	S2rt	S2tw	S3tz
Handrala	91	S3rt	S2tz	S3rt	S1	S3t	S1	S3r	S2r	S1	S2r	S2t	S2t	S3t	S1	N1t	S3rt	S2r	S3t	S3t	S3tw	S2tw	S2tw	S2rt	S2t	S3tw	S2tz	S2tz	S3t	S2rt	S2tw	S3tz
Handrala	92	S3rt	S2tz	S3rt	S1	S3t	S1	S3r	S2r	S1	S2r	S2t	S2t	S3t	S1	N1t	S3rt	S2r	S3t	S3t	S3tw	S2tw	S2tw	S2rt	S2t	S3tw	S2tz	S2tz	S3t	S2rt	S2tw	S3tz
Handrala	93	S3t	S2t	S3t	S2g	S3t	S1	S2rt	S2g	S1	S2g	S2t	S2t	S3t	S1	N1t	S2rt	S2g	S3tg	S3t	S3t	S2tg	S2tg	S2tg	S2t	S3t	S2tz	S2tz	S2t	S2tg	S2gt	S3tz
Handrala	94	S3rt	S2tz	S3rt	S1	S3t	S1	S3rg	S2r	S1	S2r	S2t	S2gt	S3t	S1	N1t	S3rt	S2r	S3t	S3t	S3tw	S2tw	S2tw	S2rt	S2t	S3tw	S2tz	S2tz	S3t	S2rt	S2tw	S3tz
Handrala	95	S3rt	S2tz	S3rt	S1	S3t	S1	S3rg	S2r	S1	S2r	S2t	S2gt	S3t	S1	N1t	S3rt	S2r	S3t	S3t	S3tw	S2tw	S2tw	S2rt	S2t	S3tw	S2tz	S2tz	S3t	S2rt	S2tw	S3tz
Handrala	96	S3rt	S2tz	S3rt	S1	S3t	S1	S3rg	S2r	S1	S2r	S2t	S2gt	S3t	S1	N1t	S3rt	S2r	S3t	S3t	S3tw	S2tw	S2tw	S2rt	S2t	S3tw	S2tz	S2tz	S3t	S2rt	S2tw	S3tz
Handrala	97	S3rt	S2tz	S3rt	S1	S3t	S1	S3rg	S2r	S1	S2r	S2t	S2gt	S3t	S1	N1t	S3rt	S2r	S3t	S3t	S3tw	S2tw	S2tw	S2rt	S2t	S3tw	S2tz	S2tz	S3t	S2rt	S2tw	S3tz
Handrala	98	S3rt	S2tz	S3rt	S1	S3t	S1	S3rg	S2r	S1	S2r	S2t	S2gt	S3t	S1	N1t	S3rt	S2r	S3t	S3t	S3tw	S2tw	S2tw	S2rt	S2t	S3tw	S2tz	S2tz	S3t	S2rt	S2tw	S3tz
Handrala	99	S3rt	S2tz	S3rt	S1	S3t	S1	S3rg	S2r	S1	S2r	S2t	S2gt	S3t	S1	N1t	S3rt	S2r	S3t	S3t	S3tw	S2tw	S2tw	S2rt	S2t	S3tw	S2tz	S2tz	S3t	S2rt	S2tw	S3tz

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Jasmine	Bhendi	Brinjal	Crossandra	Drumstick	Mulberry	Onion
Handrala	100	S3rt	S2tz	S3rt	S1	S3t	S1	S3rg	S2r	S1	S2r	S2t	S2gt	S3t	S1	N1t	S3rt	S2r	S3t	S3t	S3tw	S2tw	S2rt	S2t	S3tw	S2tz	S2tz	S3t	S2rt	S2tw	S3tz	
Handrala	101	N1rz	S2tz	S3rz	S2rz	S3tz	S2rz	N1rz	S3rz	S2rz	S3rz	S3rz	S2rz	S3tz	S2rz	N1tz	S3tz	S3rz	S3tz	S3tz	S3tz	S2rz	S2rz	S3rz	S2tz	S2rz	S2rt	S2rt	S3rz	S3rz	S3rz	S3tz
Handrala	102	S3t	S2t	S3t	S2g	S3t	S1	S2rt	S2g	S1	S2g	S2t	S2t	S3t	S1	N1t	S2rt	S2g	S3tg	S3t	S3t	S2tg	S2tg	S2tg	S2t	S3t	S2tz	S2tz	S2t	S2tg	S2gt	S3tz
Handrala	103	N1rz	S2tz	S3rz	S2rz	S3tz	S2rz	N1rz	S3rz	S2rz	S3rz	S3rz	S2rz	S3tz	S2rz	N1tz	S3tz	S3rz	S3tz	S3tz	S3tz	S2rz	S2rz	S3rz	S2tz	S2rz	S2rt	S2rt	S3rz	S3rz	S3rz	S3tz
Handrala	104	N1rz	S2tz	S3rz	S2rz	S3tz	S2rz	N1rz	S3rz	S2rz	S3rz	S3rz	S2rz	S3tz	S2rz	N1tz	S3tz	S3rz	S3tz	S3tz	S3tz	S2rz	S2rz	S3rz	S2tz	S2rz	S2rt	S2rt	S3rz	S3rz	S3rz	S3tz
Handrala	105	N1rz	S2tz	S3rz	S2rz	S3tz	S2rz	N1rz	S3rz	S2rz	S3rz	S3rz	S2rz	S3tz	S2rz	N1tz	S3tz	S3rz	S3tz	S3tz	S3tz	S2rz	S2rz	S3rz	S2tz	S2rz	S2rt	S2rt	S3rz	S3rz	S3rz	S3tz
Handrala	106	N1rz	S2tz	S3rz	S2rz	S3tz	S2rz	N1rz	S3rz	S2rz	S3rz	S3rz	S2rz	S3tz	S2rz	N1tz	S3tz	S3rz	S3tz	S3tz	S3tz	S2rz	S2rz	S3rz	S2tz	S2rz	S2rt	S2rt	S3rz	S3rz	S3rz	S3tz
Handrala	107	S3t	S2t	S3t	S2g	S3t	S1	S2rt	S2g	S1	S2g	S2t	S2t	S3t	S1	N1t	S2rt	S2g	S3tg	S3t	S3t	S2tg	S2tg	S2tg	S2t	S3t	S2tz	S2tz	S2t	S2tg	S2gt	S3tz
Handrala	108	N1rz	S2tz	S3rz	S2rz	S3tz	S2rz	N1rz	S3rz	S2rz	S3rz	S3rz	S2rz	S3tz	S2rz	N1tz	S3tz	S3rz	S3tz	S3tz	S3tz	S2rz	S2rz	S3rz	S2tz	S2rz	S2rt	S2rt	S3rz	S3rz	S3rz	S3tz
Handrala	109	N1rz	S2tz	S3rz	S2rz	S3tz	S2rz	N1rz	S3rz	S2rz	S3rz	S3rz	S2rz	S3tz	S2rz	N1tz	S3tz	S3rz	S3tz	S3tz	S3tz	S2rz	S2rz	S3rz	S2tz	S2rz	S2rt	S2rt	S3rz	S3rz	S3rz	S3tz
Handrala	110	N1rz	S2tz	S3rz	S2rz	S3tz	S2rz	N1rz	S3rz	S2rz	S3rz	S3rz	S2rz	S3tz	S2rz	N1tz	S3tz	S3rz	S3tz	S3tz	S3tz	S2rz	S2rz	S3rz	S2tz	S2rz	S2rt	S2rt	S3rz	S3rz	S3rz	S3tz
Handrala	111	N1rz	S2tz	S3rz	S2rz	S3tz	S2rz	N1rz	S3rz	S2rz	S3rz	S3rz	S2rz	S3tz	S2rz	N1tz	S3tz	S3rz	S3tz	S3tz	S3tz	S2rz	S2rz	S3rz	S2tz	S2rz	S2rt	S2rt	S3rz	S3rz	S3rz	S3tz
Handrala	112	N1rz	S2tz	S3rz	S2rz	S3tz	S2rz	N1rz	S3rz	S2rz	S3rz	S3rz	S2rz	S3tz	S2rz	N1tz	S3tz	S3rz	S3tz	S3tz	S3tz	S2rz	S2rz	S3rz	S2tz	S2rz	S2rt	S2rt	S3rz	S3rz	S3rz	S3tz
Handrala	113	N1rz	S2tz	S3rz	S2rz	S3tz	S2rz	N1rz	S3rz	S2rz	S3rz	S3rz	S2rz	S3tz	S2rz	N1tz	S3tz	S3rz	S3tz	S3tz	S3tz	S2rz	S2rz	S3rz	S2tz	S2rz	S2rt	S2rt	S3rz	S3rz	S3rz	S3tz
Handrala	114	N1rz	S2tz	S3rz	S2rz	S3tz	S2rz	N1rz	S3rz	S2rz	S3rz	S3rz	S2rz	S3tz	S2rz	N1tz	S3tz	S3rz	S3tz	S3tz	S3tz	S2rz	S2rz	S3rz	S2tz	S2rz	S2rt	S2rt	S3rz	S3rz	S3rz	S3tz
Handrala	147	N1r g	N1r g	N1r g	N1r g	N1r g	N1r g	N1r g	N1r g	N1r g	N1r g	N1r g	N1r g	N1r g	N1r g	N1r g	N1r g	N1r g	N1r g	N1r g	N1rg	N1rg	N1rg	N1rg	N1rg	N1r	N1r	N1rg	N1rg	N1rg	N1r	
Handrala	148	S3rt	S2tz	S3rt	S1	S3t	S1	S3rg	S2r	S1	S2r	S2t	S2gt	S3t	S1	N1t	S3rt	S2r	S3t	S3t	S3tw	S2tw	S2rt	S2t	S3tw	S2tz	S2tz	S3t	S2rt	S2tw	S3tz	
Handrala	150	S3rt	S2tz	S3rt	S1	S3t	S1	S3rg	S2r	S1	S2r	S2t	S2gt	S3t	S1	N1t	S3rt	S2r	S3t	S3t	S3tw	S2tw	S2rt	S2t	S3tw	S2tz	S2tz	S3t	S2rt	S2tw	S3tz	
Handrala	152	N1rz	S2tz	S3rz	S2rz	S3tz	S2rz	N1rz	S3rz	S2rz	S3rz	S3rz	S2rz	S3tz	S2rz	N1tz	S3tz	S3rz	S3tz	S3tz	S3tz	S2rz	S2rz	S3rz	S2tz	S2rz	S2rt	S2rt	S3rz	S3rz	S3rz	S3tz
Handrala	153	S3rt	S2tz	S3rt	S1	S3t	S1	S3rg	S2r	S1	S2r	S2t	S2gt	S3t	S1	N1t	S3rt	S2r	S3t	S3t	S3tw	S2tw	S2rt	S2t	S3tw	S2tz	S2tz	S3t	S2rt	S2tw	S3tz	
Handrala	154	S3rt	S2tz	S3rt	S1	S3t	S1	S3rg	S2r	S1	S2r	S2t	S2gt	S3t	S1	N1t	S3rt	S2r	S3t	S3t	S3tw	S2tw	S2rt	S2t	S3tw	S2tz	S2tz	S3t	S2rt	S2tw	S3tz	
Handrala	155	S3rt	S2tz	S3rt	S1	S3t	S1	S3rg	S2r	S1	S2r	S2t	S2gt	S3t	S1	N1t	S3rt	S2r	S3t	S3t	S3tw	S2tw	S2rt	S2t	S3tw	S2tz	S2tz	S3t	S2rt	S2tw	S3tz	
Handrala	156	S3rt	S2tz	S3rt	S1	S3t	S1	S3rg	S2r	S1	S2r	S2t	S2gt	S3t	S1	N1t	S3rt	S2r	S3t	S3t	S3tw	S2tw	S2rt	S2t	S3tw	S2tz	S2tz	S3t	S2rt	S2tw	S3tz	
Handrala	157	N1r g	N1r g	N1r g	N1r g	N1r g	N1r g	N1r g	N1r g	N1r g	N1r g	N1r g	N1r g	N1r g	N1r g	N1r g	N1r g	N1r g	N1r g	N1r g	N1rg	N1rg	N1rg	N1rg	N1rg	N1r	N1r	N1rg	N1rg	N1rg	N1r	
Handrala	158	S3t	S2t	S3t	S1	S3t	S1	S2rt	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S3t	S2t	S2t	S2t	S2t	S3t	S2tz	S2tz	S2t	S2t	S2t	S3tz
Handrala	159	S3t	S2t	S3t	S1	S3t	S1	S2rt	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S3t	S2t	S2t	S2t	S2t	S3t	S2tz	S2tz	S2t	S2t	S2t	S3tz

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Jasmine	Bhendi	Brinjal	Crossandra	Drumstick	Mulberry	Onion	
Handrala	160	S3t	S2t	S3t	S1	S3t	S1	S2rt	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S3t	S2t	S2t	S2t	S2t	S3t	S2tz	S2tz	S2t	S2t	S2t	S3tz	
Handrala	161	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs
Handrala	162	S3t	S2t	S3t	S1	S3t	S1	S2rt	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S3t	S2t	S2t	S2t	S2t	S3t	S2tz	S2tz	S2t	S2t	S2t	S3tz	
Handrala	163	S3t	S2t	S3t	S1	S3t	S1	S2rt	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S3t	S2t	S2t	S2t	S2t	S3t	S2tz	S2tz	S2t	S2t	S2t	S3tz	
Handrala	164	S3t	S2t	S3t	S1	S3t	S1	S2rt	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S3t	S2t	S2t	S2t	S2t	S3t	S2tz	S2tz	S2t	S2t	S2t	S3tz	
Handrala	165	S3t	S2t	S3t	S1	S3t	S1	S2rt	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S3t	S2t	S2t	S2t	S2t	S3t	S2tz	S2tz	S2t	S2t	S2t	S3tz	
Handrala	166	S3t	S2t	S3t	S1	S3t	S1	S2rt	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S3t	S2t	S2t	S2t	S2t	S3t	S2tz	S2tz	S2t	S2t	S2t	S3tz	
Handrala	167	S3t	S2t	S3t	S1	S3t	S1	S2rt	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S3t	S2t	S2t	S2t	S2t	S3t	S2tz	S2tz	S2t	S2t	S2t	S3tz	
Handrala	168	S3t	S2t	S3t	S1	S3t	S1	S2rt	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S3t	S2t	S2t	S2t	S2t	S3t	S2tz	S2tz	S2t	S2t	S2t	S3tz	
Handrala	169	S3rz	S2tz	S3tz	S2nz	S3tz	S2rz	S3rz	S2rz	S2rz	S2rz	S3rz	S2tz	S3tz	S2z	N1tz	S3rz	S2rz	S3tz	S3tz	S3tz	S2tz	S2tz	S2rt	S2tz	S3tz	S2tz	S2tz	S2z	S2rz	S2tz	S3tz	
Handrala	170	S3rz	S2tz	S3tz	S2nz	S3tz	S2rz	S3rz	S2rz	S2rz	S2rz	S3rz	S2tz	S3tz	S2z	N1tz	S3rz	S2rz	S3tz	S3tz	S3tz	S2tz	S2tz	S2rt	S2tz	S3tz	S2tz	S2tz	S2z	S2rz	S2tz	S3tz	
Handrala	171	S3rt	S2tz	S3rt	S1	S3t	S1	S3r	S2r	S1	S2r	S2t	S2t	S3t	S1	N1t	S3rt	S2r	S3t	S3t	S3t w	S2t w	S2t w	S2rt	S2t	S3t w	S2tz	S2tz	S3t	S2rt	S2t w	S3tz	
Handrala	172	S3rz	S2tz	S3tz	S2nz	S3tz	S2rz	S3rz	S2rz	S2rz	S2rz	S3rz	S2tz	S3tz	S2z	N1tz	S3rz	S2rz	S3tz	S3tz	S3tz	S2tz	S2tz	S2rt	S2tz	S3tz	S2tz	S2tz	S2z	S2rz	S2tz	S3tz	
Handrala	173	S3rz	S2tz	S3tz	S2nz	S3tz	S2rz	S3rz	S2rz	S2rz	S2rz	S3rz	S2tz	S3tz	S2z	N1tz	S3rz	S2rz	S3tz	S3tz	S3tz	S2tz	S2tz	S2rt	S2tz	S3tz	S2tz	S2tz	S2z	S2rz	S2tz	S3tz	
Handrala	174	S3rz	S2tz	S3tz	S2nz	S3tz	S2rz	S3rz	S2rz	S2rz	S2rz	S3rz	S2tz	S3tz	S2z	N1tz	S3rz	S2rz	S3tz	S3tz	S3tz	S2tz	S2tz	S2rt	S2tz	S3tz	S2tz	S2tz	S2z	S2rz	S2tz	S3tz	
Handrala	175	S3rz	S2tz	S3tz	S2nz	S3tz	S2rz	S3rz	S2rz	S2rz	S2rz	S3rz	S2tz	S3tz	S2z	N1tz	S3rz	S2rz	S3tz	S3tz	S3tz	S2tz	S2tz	S2rt	S2tz	S3tz	S2tz	S2tz	S2z	S2rz	S2tz	S3tz	
Handrala	176	S3rt	S2tz	S3rt	S1	S3t	S1	S3r	S2r	S1	S2r	S2t	S2t	S3t	S1	N1t	S3rt	S2r	S3t	S3t	S3tw	S2tw	S2tw	S2rt	S2t	S3tw	S2tz	S2tz	S3t	S2rt	S2tw	S3tz	
Handrala	177	S3rz	S2tz	S3tz	S2nz	S3tz	S2rz	S3rz	S2rz	S2rz	S2rz	S3rz	S2tz	S3tz	S2z	N1tz	S3rz	S2rz	S3tz	S3tz	S3tz	S2tz	S2tz	S2rt	S2tz	S3tz	S2tz	S2tz	S2z	S2rz	S2tz	S3tz	
Handrala	178	S3rz	S2tz	S3tz	S2nz	S3tz	S2rz	S3rz	S2rz	S2rz	S2rz	S3rz	S2tz	S3tz	S2z	N1tz	S3rz	S2rz	S3tz	S3tz	S3tz	S2tz	S2tz	S2rt	S2tz	S3tz	S2tz	S2tz	S2z	S2rz	S2tz	S3tz	
Handrala	179	S3rz	S2tz	S3tz	S2nz	S3tz	S2rz	S3rz	S2rz	S2rz	S2rz	S3rz	S2tz	S3tz	S2z	N1tz	S3rz	S2rz	S3tz	S3tz	S3tz	S2tz	S2tz	S2rt	S2tz	S3tz	S2tz	S2tz	S2z	S2rz	S2tz	S3tz	
Handrala	180	S3rz	S2tz	S3tz	S2nz	S3tz	S2rz	S3rz	S2rz	S2rz	S2rz	S3rz	S2tz	S3tz	S2z	N1tz	S3rz	S2rz	S3tz	S3tz	S3tz	S2tz	S2tz	S2rt	S2tz	S3tz	S2tz	S2tz	S2z	S2rz	S2tz	S3tz	
Handrala	181	S3rz	S2tz	S3tz	S2nz	S3tz	S2rz	S3rz	S2rz	S2rz	S2rz	S3rz	S2tz	S3tz	S2z	N1tz	S3rz	S2rz	S3tz	S3tz	S3tz	S2tz	S2tz	S2rt	S2tz	S3tz	S2tz	S2tz	S2z	S2rz	S2tz	S3tz	
Hireshindho gi	273	S3tz	S2tz	S3tz	S2z	S3tz	S2z	S2tz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2tz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S3tz	S3tz	
Hireshindho gi	274	S3tz	S2tz	S3tz	S2z	S3tz	S2z	S2tz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2tz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S3tz	S3tz	
Hireshindho gi	275	S3tz	S2tz	S3tz	S2z	S3tz	S2z	S2tz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2tz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S3tz	S3tz	

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Jasmine	Bhendi	Brinjal	Crossandra	Drumstick	Mulberry	Onion
Hireshindho gi	276	S3tz	S2tz	S3tz	S2z	S3tz	S2z	S2tz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2tz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S3tz	S3tz
Hireshindho gi	277	S3tz	S2tz	S3tz	S2z	S3tz	S2z	S2tz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2tz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S3tz	S3tz
Hireshindho gi	278	S3tz	S2tz	S3tz	S2z	S3tz	S2z	S2tz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2tz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S3tz	S3tz
Hireshindho gi	279	S3tz	S2tz	S3tz	S2z	S3tz	S2z	S2tz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2tz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S3tz	S3tz
Hireshindho gi	280	S3tz	S2tz	S3tz	S2z	S3tz	S2z	S2tz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2tz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S3tz	S3tz
Hireshindho gi	281	S3tz	S2tz	S3tz	S2z	S3tz	S2z	S2tz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2tz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S3tz	S3tz
Hireshindho gi	284	S3tz	S2tz	S3tz	S2z	S3tz	S2z	S2tz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2tz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S3tz	S3tz
Hireshindho gi	285	S3tz	S2tz	S3tz	S2z	S3tz	S2z	S2tz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2tz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S3tz	S3tz
Hireshindho gi	286	S3tz	S2tz	S3tz	S2z	S3tz	S2z	S2tz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2tz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S3tz	S3tz
Hireshindho gi	287	S3tz	S2tz	S3tz	S2z	S3tz	S2z	S2tz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2tz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S3tz	S3tz
Hireshindho gi	288	S3tz	S2tz	S3tz	S2z	S3tz	S2z	S2tz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2tz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S3tz	S3tz
Hireshindho gi	289	S3tz	S2tz	S3tz	S2z	S3tz	S2z	S2tz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2tz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S3tz	S3tz
Hireshindho gi	290	S3tz	S2tz	S3tz	S2z	S3tz	S2z	S2tz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2tz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S3tz	S3tz
Hireshindho gi	293	S3tz	S2tz	S3tz	S2z	S3tz	S2z	S2tz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2tz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S3tz	S3tz
Hireshindho gi	294	S3rt	S2tz	S3rt	S1	S3t	S1	S3r	S2r	S1	S2r	S2t	S3t	S1	N1t	S3rt	S2r	S3t	S3t	S3tw	S2tw	S2tw	S2rt	S2t	S3tw	S2tz	S2tz	S3t	S2rt	S2tw	S3tz	
Hireshindho gi	295	S3rt	S2tz	S3rt	S1	S3t	S1	S3r	S2r	S1	S2r	S2t	S3t	S1	N1t	S3rt	S2r	S3t	S3t	S3tw	S2tw	S2tw	S2rt	S2t	S3tw	S2tz	S2tz	S3t	S2rt	S2tw	S3tz	
Narasapura	33	S3rt	S2tz	S3rt	S1	S3t	S1	S3r	S2r	S1	S2r	S2t	S3t	S1	N1t	S3rt	S2r	S3t	S3t	S3tw	S2tw	S2tw	S2rt	S2t	S3tw	S2tz	S2tz	S3t	S2rt	S2tw	S3tz	

# **PART-B**

**SOCIO-ECONOMIC STATUS OF FARM HOUSEHOLDS**





## CONTENTS

1.	Findings of the socio-economic survey	1-3
2.	Introduction	5
3	Methodology	7-8
4	Salient features of the survey	9-26
5	Summary	27-31



## LIST OF TABLES

1	Households sampled for socio economic survey	9
2	Population characteristics	9
3	Age wise classification of household members	9
4	Education level of household members	10
5	Occupation of household heads	10
6	Occupation of family members	10
7	Institutional participation of household members	11
8	Type of house owned by households	11
9	Durable assets owned by households	11
10	Average value of durable assets owned by households	12
11	Farm implements owned by households	12
12	Average value of farm implements	12
13	Livestock possession by households	12
14	Average labour availability	13
15	Adequacy of hired labour	13
16	Distribution of land (ha)	13
17	Average land value (Rs./ha)	13
18	Status of bore wells	14
19	Source of irrigation	14
20	Depth of water(Avg in meters)	14
21	Irrigated area (ha)	14
22	Cropping pattern	14
23	Cropping intensity	14
24.a	Cost of cultivation of Maize	15
24.b	Cost of cultivation of Sunflower	16
24.c	Cost of cultivation of Green gram	17
24.d	Cost of cultivation of Cotton	18
24.e	Cost of cultivation of Onion	19
25	Adequacy of fodder	20
26	Annual gross income	20
27	Average annual expenditure	20

28	Forest species grown	21
29	Average additional investment capacity	21
30	Source of funds for additional investment	21
31	Marketing of the agricultural produce	22
32	Marketing channels used for sale of agricultural produce	22
33	Mode of transport of agricultural produce	22
34	Incidence of soil and water erosion problems	23
35	Interest shown towards soil testing	23
36	Soil and water conservation practices and structures	23
37	Status soil and water conservation structures	23
38	Agencies involved in the soil and water conservation structures	23
39	Usage pattern of fuel for domestic use	24
40	Source of drinking water	24
41	Source of light	24
42	Existence of sanitary toilet facility	24
43	Possession of public distribution system (PDS) card	25
44	Participation in NREGA programme	25
45	Adequacy of food items	25
46	Inadequacy of food items	25
47	Farming constraints experienced	26

**FINDINGS OF THE SOCIO-ECONOMIC SURVEY**

- ❖ *The survey was conducted in Handrahalu-2 is located at North latitude 15<sup>o</sup> 19' 52.722" and 15<sup>o</sup> 17' 50.924" and East longitude 76<sup>o</sup> 3' 8.831" and 76<sup>o</sup> 1' 41.664" covering an area of about 450.01 ha coming under Handrala, Hirehindogi and Hanavala Villages of Koppal taluk.*
- ❖ *Socio-economic analysis of Handrahalu-2 micro watersheds of Tadakal sub-watershed, Koppala taluk & District indicated that, out of the total sample of 34 farmers were sampled in Handrahalu-2 micro-watershed among households surveyed 6 (17.65%) were marginal, 10 (29.41%) were small and 13 (38.24 %) were semi medium farmers. 5 landless farmers were also interviewed for the survey.*
- ❖ *The population characteristics of households indicated that, there were 92 (57.14%) men and 69 (42.86 %) were women. The average population of landless was 4.8, marginal farmers were 4.8, small farmers were 5 and semi medium farmers were 4.5.*
- ❖ *Majority of the respondents (37.27%) were in the age group of 16-35 years.*
- ❖ *Education level of the sample households indicated that, there were 37.27 per cent illiterates, 60.87 per cent pre university education and 2.48 per cent attained graduation.*
- ❖ *About, 97.06 per cent of household heads practicing agriculture.*
- ❖ *Agriculture was the major occupation for 59.63 per cent of the household members.*
- ❖ *In the study area, 79.41 per cent of the households possess katcha house and 2.94 per cent possess pucca house.*
- ❖ *The durable assets owned by the households showed that, 70.59 per cent possess TV, 23.53 per cent possess mixer grinder, 94.12 per cent possess mobile phones and 20.59 per cent possess motor cycles.*
- ❖ *Farm implements owned by the households indicated that, 14.71 per cent of the households possess plough, 2.94 per cent possess tractor, 20.59 per cent possess bullock cart.*
- ❖ *Regarding livestock possession by the households, 17.65 per cent possess local cow.*
- ❖ *The average labour availability in the study area showed that, own labour men available in the micro watershed was 1.68, women available in the micro watershed was 1.38, hired labour (men) available was 10 and hired labour (women) available was 8.24.*
- ❖ *Further, 17.65 per cent of the households opined that hired labour was inadequate during the agricultural season.*

- ❖ *Out of the total land holding of the sample respondents 65.19 per cent (39.67 ha) of the area is under dry condition and the remaining 34.81 per cent area is irrigated land.*
- ❖ *There were 16.00 live bore wells and 15.00 dry bore wells among the sampled households.*
- ❖ *Bore well was the major source of irrigation for 47.06 per cent of the households.*
- ❖ *The major crops grown by sample farmers are Maize, Sunflower, Green gram, Cotton and Onion and cropping intensity was recorded as 91.83 per cent.*
- ❖ *Out of the sample households 73.53 percent possessed bank account.*
- ❖ *Majority of the respondents (100.00%) have borrowed loan for agriculture purpose.*
- ❖ *Regarding the opinion on institutional sources of credit, 69.23 per cent of the households opined that credit helped to perform timely agricultural operations.*
- ❖ *The per hectare cost of cultivation for Maize, Sunflower, Green gram, Cotton and Onion was Rs.34869.68 , 44381.39, 28905.04, 28381.13 and 42875.32 with benefit cost ratio of 1:1.10, 1: 3.60, 1: 1.30, 1: 2.10 and 1:4.80 respectively.*
- ❖ *Further, 2.94 per cent of the households opined that dry fodder was adequate and 2.94 per cent of the households have opined that the green fodder was adequate.*
- ❖ *The average annual gross income of the farmers was Rs. 68723.53 in micro-watershed, of which Rs. 60282.35 comes from agriculture.*
- ❖ *Sampled households have grown 1 horticulture trees and 48 forestry trees together in the fields and back yards.*
- ❖ *Households have an average investment capacity of Rs. 3705.88 for land development and Rs. 941.18 for irrigation facility.*
- ❖ *Source of funds for additional investment is concerned, 55.88 per cent depends on own funds and 5.88 per cent depends on bank loan for land development activities.*
- ❖ *Regarding marketing channels, 61.76 per cent of the households have sold agricultural produce to the local/village merchants, while, 2.94 per cent have sold in regulated markets.*
- ❖ *Further, 67.65 per cent of the households have used tractor for the transport of agriculture commodity.*
- ❖ *Majority of the farmers (61.76%) have experienced soil and water erosion problems in the watershed and 76.47 per cent of the households were interested towards soil testing.*
- ❖ *Fire was the major source of fuel for domestic use for 79.41 per cent of the households and 23.53 per cent households has LPG connection.*
- ❖ *Piped supply was the major source for drinking water for 73.53 per cent of the households.*
- ❖ *Electricity was the major source of light for 100.00 per cent of the households.*

- ❖ *In the study area, 32.35 per cent of the households possess toilet facility.*
- ❖ *Regarding possession of PDS card, 94.12 per cent of the households possessed BPL card, 2.94 per cent of the household's possessed APL card and 2.94 per cent of the household's were not having ration cards.*
- ❖ *Households opined that, the requirement of cereals (73.53%), pulses (67.65%) and oilseeds (14.71%) are adequate for consumption.*
- ❖ *Farming constraints experienced by households in the micro watersheds were lower fertility status of the soil (61.76%) wild animal menace on farm field (76.47%), frequent incidence of pest and diseases (35.29%), inadequacy of irrigation water (50.00%), high cost of fertilizers and plant protection chemicals (73.53%), high rate of interest on credit (61.76%), low price for the agricultural commodities (47.06%), lack of marketing facilities in the area (61.76%), inadequate extension services (38.24%), lack of transport for safe transport of the agricultural produce to the market (47.06%), Less rainfall (8.82%) and Source of Agri-technology information (Newspaper/ TV/Mobile) (8.82%).*





## INTRODUCTION

Soil and water are the two precious natural resources which are essential for crop production and existence of life on earth. Rainfed agriculture is under severe stress due to various constraints related to agriculture like uneven and erratic distribution of rainfall, indiscriminate use of fertilizers, chemicals and pesticides, adoption of improper land management practices, soil erosion, decline in soil fertility, decline in ground water resources leading to low crop productivity. The area under rainfed agriculture has to be managed effectively using the best available practices to enhance the production of food, fodder and fuel. This is possible if the land resources are characterized at each parcel of land through detailed land resource inventory using the best available techniques of remote sensing, GPS and GIS. The watershed development programs are aimed at the sustainable distribution of its resources and the process of creating and implementing plans, programs, and projects to sustain and enhance watershed functions that affect the plant, animal and human communities within a watershed boundary.

World Bank funded KWDP II, SUJALA III project was implemented in with Broad objective of demonstrating more effective watershed management through greater integration of programmes related to rain-fed agriculture, innovative and science based approaches and strengthen institutional capacities and If successful, it is expected that the systems and tools could be mainstreamed into the overall IWMP in the State of Karnataka and in time, throughout other IWMP operations in India. With this background the socio-economic survey has been carried out with following specific objectives:

1. To understand the demographic features of the households in the micro-watershed
2. To understand the extent of family labour available and additional employment opportunities available within the village.
3. To know the status of assets of households in the micro-watershed for suggesting possible improvements.
4. To study the cropping pattern, cropped area and productivity levels of different households in micro-watershed.
5. To determine the type and extent of livestock owned by different categories of HHs
6. Availability of fodder and level of livestock management.

### **Scope and importance of survey**

Survey helps in identification of different socio-economic and resource use-patterns of farmers at the Micro watershed. Household survey provides demographic features, labor force, and levels of education; land ownership and asset position (including livestock and other household assets) of surveyed households; and cropping patterns, input intensities, and average crop yields from farmers' fields. It also discusses crop utilization and the degree of commercialization of production in the areas; farmers' access to and utilization of credit from formal and informal sources; and the level of adoption and use of soil, water, and pest management technologies.



## METHODOLOGY

The description of the methods, components selected for the survey and procedures followed in conducting the baseline survey are furnished under the following heads.

### 1. Description of the study area

Koppal district is an administrative district in the state of Karnataka in India. In the past Koppal was referred to as 'Kopana Nagara'. Koppal, now a district headquarters is ancient Kopana a major holy place of the Jains. The district occupies an area of 7,190 km<sup>2</sup> and has a population of 1,196,089, which 16.58% were urban as of 2001. The Koppal district was formed after split of Raichur district.

Geographers are very particular about the physiography or relief of a region. It plays a very important role in the spatial analysis of agricultural situation of the study area. The undulating topography with black cotton soil shrips, cut across by numerous nalas or streams is the major characteristic feature of the study region. Three physiographic divisions have made considering the local conditions of landforms and crops grown in the district. On the basis of physiography, Koppal district can be divided into three major divisions. They are (a) Koppal & Yelburga plateau, (b) Maidan division, (c) Tungabhadra valley. The district is part of Krishna basin the main streams draining the area are Maskinala, Ilkal-nadi and Hirenala. These are Ephemeral in nature, these come under Tungabhadra sub-basin. The drainage exhibit dentritic to subdentritic with drainage density varies from 1.4 to 7.0kms/sq.km.

According to the 2011 census Koppal district has a population of 1,391,292, roughly equal to the nation of Swaziland or the US state of Hawaii. This gives it a ranking of 350th in India (out of a total of 640). The district has a population density of 250 inhabitants per square kilometre (650/sq mi). Its population growth rate over the decade 2001-2011 was 16.32%. Koppal has a sex ratio of 983 females for every 1000 males, and a literacy rate of 67.28%.

### 2. Locale of the survey and description of the micro-watershed and

The study was conducted in Handrahalu-2 micro-watershed (Tadakal sub-watershed, Koppala taluk & District) is located at North latitude 15<sup>0</sup> 19' 52.722" and 15<sup>0</sup> 17' 50.924" and East longitude 76<sup>0</sup> 3' 8.831" and 76<sup>0</sup> 1' 41.664" covering an area of about 450.01 ha bounded by under Handrala, Hireshindogi and Hanavala Villages.

### 3. Selection of the respondents for the study

The micro-watershed is marked with 320 square meters grids. One farmer from every alternate grid in the micro-watershed was selected for the study and interviewed for socio-economic data. Totally 34 households were interviewed for the survey.

#### **4. The parameters considered for socio-economic survey of households**

Two forms of data were collected from the micro-watershed which includes primary data from the farm households and secondary data about the villages under the micro-watershed jurisdiction.

The following parameters were considered for the primary data collection about the socio-economic data of the households, (1) Demographic information, (2) Farm and durable assets owned by households, (3) Livestock possession, (4) Labour availability, (5) Level of migration in the village, Land holding, (7) Cropping pattern, (8) Source of irrigation, (9) Borrowing status, (10) Cost of cultivation of major crops, (11) Economics of subsidiary activities, (12) Fodder availability, (13) Family annual income from different sources, (14) Horticulture and forestry species grown, (15) Additional investment capacity, (16) Marketing practices, (17) Status of soil and water conservation structure, (18) Access to basic needs and (19) Constraints and suggestion.

The following parameters were considered for the secondary data regarding the villages under the micro-watershed jurisdiction, (1) Number of villages in each micro-watershed jurisdiction, (2) Village wise number of households, (3) Geographical area of the villages, (4) Cultivable area including rainfed and irrigated, (5) Number and type of house in each village, (6) Human and livestock population, (7) Facilities in the village such as roads, transport facility for conveyance, drinking water supply, street light and (8) Community based organizations in the villages.

#### **5. Development of interview schedule and data collection**

Taking into the consideration the objectives of the survey, an interview schedule was prepared after thorough consultation with the experts in the field of social sciences. A comprehensive interview schedule covering all the major parameters for measuring the socio-economic situation was developed.

#### **6. Tools used to analyze the data**

The statistical components such as frequency and percentage were used to analyze the data.

#### **Abbreviations used in the report**

LL=Landless

MF=Marginal Farmers

SF=Small farmers

SMF=Semi medium farmers

MDF=Medium farmers

LF=Large Farmers

## FINDINGS OF THE SURVEY

This chapter deals with systematic presentation of results of the survey. Keeping in view the objectives, the salient features of the survey are presented under the following headings.

**Households sampled for socio-economic survey:** The data on households sampled for socio economic survey in Handrahalu-2 Micro watershed is presented in Table 1 and it indicated that 34 farmers were sampled in Handrahalu-2 micro-watershed among households surveyed 6 (17.65%) were marginal, 10 (29.41%) were small and 13 (38.24 %) were semi medium farmers. 5 landless farmers were also interviewed for the survey.

**Table 1. Households sampled for socio economic survey in Handrahalu-2 micro-watershed**

Sl.No.	Particulars	LL (5)		MF (6)		SF (10)		SMF (13)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Farmers	5	14.7	6	17.7	10	29.4	13	38.2	34	100

**Population characteristics:** The population characteristics of households sampled for socio-economic survey in Handrahalu-2 Micro watershed is presented in Table 2. The data indicated that, there were 92 (57.14%) men and 69 (42.86%) were women. The average population of landless was 4.8, marginal farmers were 4.8, small farmers were 5 and semi medium farmers were 4.5.

**Table 2. Population characteristics in Handrahalu-2 micro-watershed**

Sl.No.	Particulars	LL (24)		MF (29)		SF (50)		SMF (58)		All (161)	
		N	%	N	%	N	%	N	%	N	%
1	Men	15	62.5	18	62	27	54	32	55.2	92	57.1
2	Women	9	37.5	11	38	23	46	26	44.8	69	42.9
Total		24	100	29	100	50	100	58	100	161	100
Average		4.8		4.8		5.0		4.5		4.7	

**Age wise classification of population:** The age wise classification of household members in Handrahalu-2 Micro watershed is presented in Table 3. The indicated that, 46 (28.57%) of population were 0-15 years of age, 60 (37.27%) were 16-35 years of age, 45(27.95%) were 36-60 years of age and 10 (6.21 %) were above 61 years of age.

**Table 3: Age wise classification of members of the household in Handrahalu-2 micro-watershed**

Sl.No.	Particulars	LL (24)		MF (29)		SF (50)		SMF (58)		All (161)	
		N	%	N	%	N	%	N	%	N	%
1	0-15 years of age	9	37.5	10	34.5	15	30	12	20.69	46	28.57
2	16-35 years of age	9	37.5	11	37.9	19	38	21	36.21	60	37.27
3	36-60 years of age	5	20.8	7	24.1	11	22	22	37.93	45	27.95
4	> 61 years	1	4.17	1	3.45	5	10	3	5.17	10	6.21
Total		24	100	29	100	50	100	58	100	161	100

**Education level of household members:** Education level of household members in Handrahalu-2 Micro watershed is presented in Table 4. The results indicated that, there

were 37.27 per cent of illiterates, 18.63 per cent of them had primary school education, 16.15 per cent middle school education, 17.39 per cent high school education, 4.35 per cent of them had PUC education, 2.48 per cent attained graduation and 1.24 them had other education.

**Table 4. Education level of members of the household in Handrahalu-2 micro-watershed**

Sl.No.	Particulars	LL (24)		MF (29)		SF (50)		SMF (58)		All (161)	
		N	%	N	%	N	%	N	%	N	%
1	Illiterate	13	54.2	7	24.1	19	38	21	36.2	60	37.3
2	Primary School	1	4.17	6	20.7	15	30	8	13.8	30	18.6
3	Middle School	5	20.8	4	13.8	5	10	12	20.7	26	16.2
4	High School	3	12.5	7	24.1	6	12	12	20.7	28	17.4
5	PUC	1	4.17	0	0	1	2	5	8.62	7	4.35
6	ITI	0	0	2	6.9	1	2	0	0	3	1.86
7	Degree	0	0	1	3.45	3	6	0	0	4	2.48
8	Masters	1	4.17	0	0	0	0	0	0	1	0.62
9	Others	0	0	2	6.9	0	0	0	0	2	1.24
Total		24	100	29	100	50	100	58	100	161	100

**Occupation of head of households:** The data regarding the occupation of the household heads in Handrahalu-2 Micro watershed is presented in Table 5. The results indicate that, 97.06 per cent of households heads were practicing agriculture.

**Table 5: Occupation of heads of households in Handrahalu-2 micro-watershed**

Sl.No.	Particulars	LL (5)		MF (6)		SF (10)		SMF (13)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Agriculture	5	100	6	100	9	90	13	100	33	97.06
2	Others	0	0	0	0	1	10	0	0	1	2.94
Total		5	100	6	100	10	100	13	100	34	100

**Table 6: Occupation of members of the household in Handrahalu-2 micro-watershed**

Sl.No.	Particulars	LL (24)		MF (29)		SF (50)		SMF (58)		All (161)	
		N	%	N	%	N	%	N	%	N	%
1	Agriculture	13	54.2	14	48.3	30	60	39	67.24	96	59.6
2	Agricultural Labour	0	0	0	0	0	0	1	1.72	1	0.62
3	General Labour	0	0	0	0	0	0	1	1.72	1	0.62
4	Government Service	0	0	1	3.45	0	0	0	0	1	0.62
5	Private Service	1	4.17	2	6.9	0	0	0	0	3	1.86
6	Student	3	12.5	9	31	16	32	16	27.59	44	27.3
7	Others	3	12.5	0	0	4	8	1	1.72	8	4.97
8	Housewife	0	0	1	3.45	0	0	0	0	1	0.62
9	Children	4	16.7	2	6.9	0	0	0	0	6	3.73
Total		24	100	29	100	50	100	58	100	161	100

**Occupation of the members of the household:** The data regarding the occupation of the household members in Handrahalu-2 Micro watershed is presented in Table 6. The results indicate that, agriculture was the major occupation for 59.63 per cent of the household members, 0.62 per cent were agricultural labour, 0.62 per cent were general labour, 0.62

per cent were working in government sector, 27.33 per cent were working in pursuing education, 0.62 per cent were involved as housewife and 3.73 per cent were children.

**Institutional Participation of household members:** The data regarding the institutional participation of the household members in Handrahalu-2 Micro watershed is presented in Table 7. The results show that, out of the total family members in the households 100 per cent of them were not participating in any of the institutions.

**Table 7: Institutional Participation of household member in Handrahalu-2 micro-watershed**

Sl.No.	Particulars	LL (24)		MF (29)		SF (50)		SMF (58)		All (161)	
		N	%	N	%	N	%	N	%	N	%
1	No Participation	24	100	29	100	50	100	58	100	161	100
	Total	24	100	29	100	50	100	58	100	161	100

**Type of house owned:** The data regarding the type of house owned by the households in Handrahalu-2 Micro watershed is presented in Table 8. The results indicate that, 8.82 percent possess thatched house, 79.41 per cent of the households possess katcha house, 2.94 per cent possess pacca house and 8.82 percent possess semi pacca house.

**Table 8. Type of house owned by households in Handrahalu-2 micro-watershed**

Sl.No.	Particulars	LL (5)		MF (6)		SF (10)		SMF (13)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Thatched	3	60	0	0	0	0	0	0	3	8.82
2	Katcha	1	20	5	83	9	90	12	92.3	27	79.41
3	Pucca/RCC	1	20	0	0	0	0	0	0	1	2.94
4	Semi pacca	0	0	1	17	1	10	1	7.69	3	8.82
	Total	5	100	6	100	10	100	13	100	34	100

**Durable assets owned by the households:** The data regarding the Durable Assets owned by the households in Handrahalu-2 Micro watershed is presented in Table 9. The results shows that, 70.59 per cent possess TV, 23.53 per cent possess mixer grinder, 8.82 per cent possess Bicycle, 20.59 per cent possess motor cycle and 94.12 per cent possess mobile phones.

**Table 9. Durable assets owned by households in Handrahalu-2 micro-watershed**

Sl.No.	Particulars	LL (5)		MF (6)		SF (10)		SMF (13)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Radio	0	0	0	0	1	10	0	0	1	2.94
2	Television	5	100	4	67	5	50	10	77	24	70.59
3	Mixer/Grinder	0	0	3	50	4	40	1	7.7	8	23.53
4	Bicycle	0	0	0	0	1	10	2	15	3	8.82
5	Motor Cycle	1	20	2	33	3	30	1	7.7	7	20.59
6	Mobile Phone	5	100	6	100	10	100	11	85	32	94.12
7	Blank	0	0	0	0	0	0	1	7.7	1	2.94

**Average value of durable assets:** The data regarding the average value of durable assets owned by the households in Handrahalu-2 Micro watershed is presented in Table 10. The result shows that, the average value of television was Rs.6875.00, mixer grinder was

Rs.1150.00, bicycle was Rs.18000.00, motor cycle was Rs. 37285.00 and mobile phone was Rs.4617.00.

**Table 10. Average value of durable assets owned in Handrahalu-2 micro-watershed**  
Average Value (Rs.)

Sl.No.	Particulars	LL (5)	MF (6)	SF (10)	SMF (13)	All (34)
1	Radio	0	0	2000	0	2000
2	Television	7000	6000	5800	7700	6875
3	Mixer/Grinder	0	1633	825	1000	1150
4	Bicycle	0	0	2000	26000	18000
5	Motor Cycle	60000	41500	23333	48000	37285
6	Mobile Phone	5400	8571	3227	3136	4617

**Farm implements owned:** The data regarding the farm implements owned by the households in Handrahalu-2 Micro watershed is presented in Table 11. About 20.59 per cent of the households possess Bullock Cart, 14.71 per cent possess plough, 29.41 per cent possess Weeder and 2.94 per cent possess tractor.

**Table 11. Farm implements owned in Handrahalu-2 micro-watershed**

Sl.No.	Particulars	LL (5)		MF (6)		SF (10)		SMF (13)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Bullock Cart	0	0	3	50	1	10	3	23.1	7	20.59
2	Plough	0	0	3	50	1	10	1	7.69	5	14.71
3	Tractor	0	0	0	0	0	0	1	7.69	1	2.94
4	Weeder	0	0	3	50	3	30	4	30.8	10	29.41
5	Blank	5	100	3	50	7	70	6	46.2	21	61.76

**Average value of farm implements:** The data regarding the average value of farm Implements owned by the households in Handrahalu-2 Micro watershed is presented in Table 12. The results show that the average value of plough was Rs.2500.00, bullock Cart was Rs.27857.00, weeder was Rs.82.00 and tractor Rs. 300000.

**Table 12. Average value of farm implements in Handrahalu-2 micro-watershed**  
Average Value (Rs.)

Sl.No.	Particulars	LL (5)	MF (6)	SF (10)	SMF (13)	All (34)
1	Bullock Cart	0	31666	30000	23333	27857
2	Plough	0	2666	2000	2500	2500
3	Tractor	0	0	0	300000	300000
4	Weeder	0	65	75	100	82

**Table 13. Livestock possession by households in Handrahalu-2 micro-watershed**

Sl.No.	Particulars	LL (5)		MF (6)		SF (10)		SMF (13)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Bullock	0	0	4	67	1	10	4	31	9	26.47
2	Local cow	0	0	2	33	1	10	3	23	6	17.65
3	Sheep	0	0	0	0	0	0	1	7.7	1	2.94
4	Goat	0	0	0	0	1	10	0	0	1	2.94
5	blank	5	100	1	17	8	80	8	62	22	64.71

**Livestock possession by the households:** The data regarding the Livestock possession by the households in Handrahalu-2 Micro watershed is presented in Table 13. The results



indicate that, 26.47 per cent of the households possess bullocks, 17.65 per cent possess local cow, 2.94 per cent possess sheep and 2.94 per cent possess goat.

**Average Labour availability:** The data regarding the average labour availability in Handrahalu-2 Micro watershed is presented in Table 14. The indicated that, own labour men available in the micro watershed was 1.68, women available in the micro watershed was 1.38, hired labour (men) available was 10 and hired labour (women) available was 8.24.

**Table 14. Average labour availability in Handrahalu-2 micro-watershed**

Sl.No.	Particulars	LL (5)	MF (6)	SF (10)	SMF (13)	All (34)
		N	N	N	N	N
1	Hired labour Female	1	6.17	8.3	11.92	8.24
2	Own Labour Female	1	1	1.4	1.69	1.38
3	Own labour Male	1	1.33	1.9	1.92	1.68
4	Hired labour Male	1	8	10.2	14.23	10

**Adequacy of hired labour:** The data regarding the adequacy of hired labour in Handrahalu-2 Micro watershed is presented in Table 15. The results indicate that, 82.35 per cent of the household opined that hired labour was adequate and 17.65 per cent of the household opined that hired labour was Inadequate.

**Table 15. Adequacy of hired labour in Handrahalu-2 micro-watershed**

Sl.No.	Particulars	LL (5)		MF (6)		SF (10)		SMF (13)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Adequate	0	0	6	100	9	90	13	100	28	82.4
2	Inadequate	5	100	0	0	1	10	0	0	6	17.7

**Distribution of land (ha):** The data regarding the distribution of land (ha) in Handrahalu-2 Micro watershed is presented in Table 16. The results indicate that, 25.86 ha (65.19%) of dry land and 13.81 ha (34.81 %) of irrigated land.

**Table 16. Distribution of land (ha) in Handrahalu-2 micro-watershed**

Sl.No.	Particulars	LL (5)		MF (6)		SF (10)		SMF (13)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Dry	0	0	3.55	88.6	9.8	82.24	12.51	52.68	25.86	65.19
2	Irrigated	0	0	0.46	11.4	2.12	17.76	11.23	47.32	13.81	34.81
Total		0	100	4.01	100	11.91	100	23.74	100	39.67	100

**Average value of land (ha):** The results (Table 17) show that the average value of dry land was Rs.237336.46 and the average value of irrigated land was Rs.390190.51.

**Table 17. Average value of land (ha) in Handrahalu-2 micro-watershed**

Sl.No.	Particulars	LL (5)	MF (6)	SF (10)	SMF (13)	All (34)
		N	N	N	N	N
1	Dry	0	450113.9	275464.7	147033.3	237336.5
2	Irrigated	0	87433.63	873709.4	311419.3	390190.5

**Status of bore wells:** The data regarding the status of bore wells in Handrahalu-2 Micro watershed is presented in Table 18. The results indicate that, there were 15 De-

functioning bore wells and 16 functioning bore wells among the sampled households in micro watershed.

**Table 18. Status of bore wells in Handrahalu-2 micro-watershed**

Sl.No.	Particulars	LL (5)	MF (6)	SF (10)	SMF (13)	All (34)
		N	N	N	N	N
1	De-functioning	0	2	3	10	15
2	Functioning	0	2	3	11	16

**Source of irrigation:** The results (Table 19) indicate that bore well for 47.06 per cent of the households.

**Table 19. Source of irrigation in Handrahalu-2 micro-watershed**

Sl.No.	Particulars	LL (5)		MF (6)		SF (10)		SMF (13)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Bore Well	0	0	2	33.3	3	30	11	84.6	16	47.06

**Depth of water (Avg. In meters):** The results (Table 20) revealed that, the depth of bore well was 25.10 meter.

**Table 20. Depth of water (Avg. In meters) in Handrahalu-2 micro-watershed**

Sl.No.	Particulars	LL (5)	MF (6)	SF (10)	SMF (13)	All (34)
		N	N	N	N	N
1	Bore Well	0	18.8	19.51	41.97	25.1

**Irrigated Area (ha):** The results (Table 21) indicate that, the availability of irrigation water was used for kharif crops was 14.60 ha.

**Table 21. Irrigated Area (ha) in Handrahalu-2 micro-watershed**

Sl.No.	Particulars	LL (5)	MF (6)	SF (10)	SMF (13)	All (34)
1	Kharif	0	1.27	2.11	11.22	14.6
	Total	0	1.27	2.11	11.22	14.6

**Cropping pattern:** The results (Table 22) indicate that, farmers have grown Maize (12.39 ha), Sunflower (11.33 ha), Green gram (4.62 ha), Cotton (2.83 ha), Onion (2.43 ha), Bajra (2.15 ha) and Chick pea (2.15 ha).

**Table 22. Cropping pattern in Handrahalu-2 micro-watershed**

Sl.No.	Particulars	LL (5)	MF (6)	SF (10)	SMF (13)	All (34)
1	Kharif - Maize	0	0.56	3.08	8.75	12.39
2	Kharif - Sunflower	0	2.08	4.33	4.92	11.33
3	Kharif - Green gram	0	0	0.87	3.74	4.62
4	Kharif - Cotton	0	0	0.81	2.02	2.83
5	Kharif - Onion	0	0	1.21	1.21	2.43
6	Kharif - Bajra	0	0.93	1.21	0	2.15
7	Kharif - Chick pea	0	0	0	2.15	2.15

**Cropping intensity:** The results (Table 23) indicate that, the cropping intensity was 91.83 per cent.

**Table 23. Cropping intensity (%) in Handrahalu-2 micro-watershed**

Sl.No.	Particulars	LL (5)	MF (6)	SF (10)	SMF (13)	All (34)
1	Cropping Intensity	0	98.59	100	87.64	91.83

**Cost of Cultivation of Maize:** The data regarding the cost of cultivation (Rs/ha) of Maize in Handrahalu-2 micro watershed is presented in Table 24.a. The results indicate that, the total cost of cultivation (Rs/ha) for Maize was Rs. 34869.68. The gross income realized by the farmers was Rs. 39629.06. The net income from Maize cultivation was Rs.4759.38, thus the benefit cost ratio was found to be 1:1.10.

**Table 24(a). Cost of Cultivation of Maize in Handrahalu-2 micro-watershed**

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
<b>I</b>	<b>Cost A1</b>				
1	Hired Human Labour	Man days	29.84	4775.84	13.7
2	Bullock	Pairs/day	0.56	484	1.39
3	Tractor	Hours	3.31	3642.78	10.45
4	Machinery	Hours	0.39	316.77	0.91
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	12.77	2237.18	6.42
6	FYM	Quintal	3.77	6577.46	18.86
7	Fertilizer + micronutrients	Quintal	7.05	6138.65	17.6
8	Pesticides (PPC)	Kgs / liters	0.98	966.92	2.77
9	Depreciation charges		0	561.73	1.61
10	Land revenue and Taxes		0	0.33	0
<b>II</b>	<b>Cost B1</b>				
11	Interest on working capital			1911.52	5.48
12	<b>Cost B1 = (Cost A1 + sum of 15 and 16)</b>			27613.17	79.19
<b>III</b>	<b>Cost B2</b>				
13	Rental Value of Land			166.67	0.48
14	<b>Cost B2 = (Cost B1 + Rental value)</b>			27779.84	79.67
<b>IV</b>	<b>Cost C1</b>				
15	Family Human Labour		19.12	3910.76	11.22
16	<b>Cost C1 = (Cost B2 + Family Labour)</b>			31690.6	90.88
<b>V</b>	<b>Cost C2</b>				
17	Risk Premium			9.1	0.03
18	<b>Cost C2 = (Cost C1 + Risk Premium)</b>			31699.7	90.91
<b>VI</b>	<b>Cost C3</b>				
19	Managerial Cost			3169.97	9.09
20	<b>Cost C3 = (Cost C2 + Managerial Cost)</b>			34869.68	100
<b>VII</b>	<b>Economics of the Crop</b>				
a.	Main Product	a) Main Product (q)		30.47	36257.27
		b) Main Crop Sales Price (Rs.)			1190
	By Product	e) Main Product (q)		8.03	3371.78
		f) Main Crop Sales Price (Rs.)			420
b.	Gross Income (Rs.)			39629.06	
c.	Net Income (Rs.)			4759.38	
d.	Cost per Quintal (Rs./q.)			1144.46	
e.	Benefit Cost Ratio (BC Ratio)			1:1.1	

**Cost of Cultivation of Sunflower:** The data regarding the cost of cultivation (Rs/ha) of Sunflower in Handrahalu-2 micro watershed is presented in Table 24.b. The results indicate that, the total cost of cultivation (Rs/ha) for Sunflower was Rs. 44381.39. The gross income realized by the farmers was Rs. 157515.89. The net income from Sunflower cultivation was Rs.113134.50, thus the benefit cost ratio was found to be 1:3.60.

**Table 24(b). Cost of Cultivation of Sunflower in Handrahalu-2 micro-watershed**

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
<b>I</b>	<b>Cost A1</b>				
1	Hired Human Labour	Man days	39.52	6597.9	14.87
2	Bullock	Pairs/day	0.73	675.46	1.52
3	Tractor	Hours	3.27	3369.29	7.59
4	Machinery	Hours	1.71	2087.15	4.7
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	5.96	1356.64	3.06
6	FYM	Quintal	3.86	11586.67	26.11
7	Fertilizer + micronutrients	Quintal	8.02	6852.04	15.44
8	Pesticides (PPC)	Kgs / liters	1.13	1124.28	2.53
9	Irrigation	Number	0.82	0	0
10	Depreciation charges		0	244.39	0.55
11	Land revenue and Taxes		0	0.27	0
<b>II</b>	<b>Cost B1</b>				
12	Interest on working capital			2511.47	5.66
13	<b>Cost B1 = (Cost A1 + sum of 15 and 16)</b>			36405.58	82.03
<b>III</b>	<b>Cost B2</b>				
14	Rental Value of Land			125	0.28
15	<b>Cost B2 = (Cost B1 + Rental value)</b>			36530.58	82.31
<b>IV</b>	<b>Cost C1</b>				
16	Family Human Labour		17.65	3806.89	8.58
17	<b>Cost C1 = (Cost B2 + Family Labour)</b>			40337.47	90.89
<b>V</b>	<b>Cost C2</b>				
18	Risk Premium			9.25	0.02
19	<b>Cost C2 = (Cost C1 + Risk Premium)</b>			40346.72	90.91
<b>VI</b>	<b>Cost C3</b>				
20	Managerial Cost			4034.67	9.09
21	<b>Cost C3 = (Cost C2 + Managerial Cost)</b>			44381.39	100
<b>VII</b>	<b>Economics of the Crop</b>				
a.	Main Product	a) Main Product (q)		53.37	157005.32
		b) Main Crop Sales Price (Rs.)			2941.67
	By Product	c) Main Product (q)		3.6	510.56
		d) Main Crop Sales Price (Rs.)			141.67
b.	Gross Income (Rs.)			157515.89	
c.	Net Income (Rs.)			113134.5	
d.	Cost per Quintal (Rs./q.)			831.53	
e.	Benefit Cost Ratio (BC Ratio)			1:3.6	

**Cost of Cultivation of Green gram:** The data regarding the cost of cultivation (Rs/ha) of Green gram in Handrahalu-2 micro watershed is presented in Table 24.c. The results indicate, the total cost of cultivation (Rs/ha) for Green gram was Rs.28905.04. The gross income realized by the farmers was Rs. 36022.10. The net income from Green gram cultivation was Rs. 7117.06, thus the benefit cost ratio was found to be 1:1.30.

**Table 24(c). Cost of Cultivation of Green gram in Handrahalu-2 micro-watershed**

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
<b>I</b>	<b>Cost A1</b>				
1	Hired Human Labour	Man days	24.48	3158.81	10.93
2	Bullock	Pairs/day	0.13	80.11	0.28
3	Tractor	Hours	2.12	2905.32	10.05
4	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	5.24	851.15	2.94
5	Depreciation charges		0	1.08	0
6	Land revenue and Taxes		0	1.65	0.01
<b>II</b>	<b>Cost B1</b>				
7	Interest on working capital			1891.75	6.54
8	<b>Cost B1 = (Cost A1 + sum of 15 and 16)</b>			23797.79	82.33
<b>III</b>	<b>Cost B2</b>				
9	Rental Value of Land			250	0.86
10	<b>Cost B2 = (Cost B1 + Rental value)</b>			24047.79	83.2
<b>IV</b>	<b>Cost C1</b>				
11	Family Human Labour		12.34	2224.02	7.69
12	<b>Cost C1 = (Cost B2 + Family Labour)</b>			26271.81	90.89
<b>V</b>	<b>Cost C2</b>				
13	Risk Premium			5.5	0.02
14	<b>Cost C2 = (Cost C1 + Risk Premium)</b>			26277.31	90.91
<b>VI</b>	<b>Cost C3</b>				
15	Managerial Cost			2627.73	9.09
16	<b>Cost C3 = (Cost C2 + Managerial Cost)</b>			28905.04	100
<b>VII</b>	<b>Economics of the Crop</b>				
a.	Main Product	a) Main Product (q)	8.28	36022.1	
		b) Main Crop Sales Price (Rs.)		4350	
b.	Gross Income (Rs.)			36022.1	
c.	Net Income (Rs.)			7117.06	
d.	Cost per Quintal (Rs./q.)			3490.55	
e.	Benefit Cost Ratio (BC Ratio)			1:1.3	

**Cost of Cultivation of Cotton:** The data regarding the cost of cultivation (Rs/ha) of Cotton in Handrahalu-2 micro watershed is presented in Table 24.d. The results indicate that, the total cost of cultivation (Rs/ha) for Cotton was Rs. 28381.13. The gross income realized by the farmers was Rs.59354.10. The net income from Cotton cultivation was Rs. 30972.97, thus the benefit cost ratio was found to be 1:2.10.

**Table 24(d). Cost of Cultivation of Cotton in Handrahalu-2 micro-watershed**

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
<b>I</b>	<b>Cost A1</b>				
1	Hired Human Labour	Man days	26.31	4094.03	14.43
2	Bullock	Pairs/day	0.86	864.5	3.05
3	Tractor	Hours	4.94	5434	19.15
4	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	3.4	305.66	1.08
5	FYM	Quintal	1.73	5187	18.28
6	Fertilizer + micronutrients	Quintal	5.19	4633.72	16.33
7	Pesticides (PPC)	Kgs / liters	0.86	864.5	3.05
8	Irrigation	Number	1.24	0	0
9	Depreciation charges		0	0.02	0
<b>II</b>	<b>Cost B1</b>				
10	Interest on working capital			1320.11	4.65
11	<b>Cost B1 = (Cost A1 + sum of 15 and 16)</b>			22703.53	80
<b>III</b>	<b>Cost B2</b>				
12	Rental Value of Land			0	0
13	<b>Cost B2 = (Cost B1 + Rental value)</b>			22703.53	80
<b>IV</b>	<b>Cost C1</b>				
14	Family Human Labour		15.07	3087.5	10.88
15	<b>Cost C1 = (Cost B2 + Family Labour)</b>			25791.03	90.87
<b>V</b>	<b>Cost C2</b>				
16	Risk Premium			10	0.04
17	<b>Cost C2 = (Cost C1 + Risk Premium)</b>			25801.03	90.91
<b>VI</b>	<b>Cost C3</b>				
18	Managerial Cost			2580.1	9.09
19	<b>Cost C3 = (Cost C2 + Managerial Cost)</b>			28381.13	100
<b>VII</b>	<b>Economics of the Crop</b>				
a.	Main Product	a) Main Product (q)	11.73	58662.5	
		b) Main Crop Sales Price (Rs.)		5000	
	By Product	e) Main Product (q)	1.73	691.6	
		f) Main Crop Sales Price (Rs.)		400	
b.	Gross Income (Rs.)			59354.1	
c.	Net Income (Rs.)			30972.97	
d.	Cost per Quintal (Rs./q.)			2419.02	
e.	Benefit Cost Ratio (BC Ratio)			1:2.1	

**Cost of Cultivation of Onion:** The data regarding the cost of cultivation (Rs/ha) of Onion in Handrahalu-2 micro watershed is presented in Table 24.e. The results indicate that, the total cost of cultivation (Rs/ha) for Onion was Rs.42875.32. The gross income realized by the farmers was Rs. 207297.04. The net income from Onion cultivation was Rs. 164421.71, thus the benefit cost ratio was found to be 1:4.80.

**Table 24(e). Cost of Cultivation of Onion in Handrahalu-2 micro-watershed**

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
<b>I</b>	<b>Cost A1</b>				
1	Hired Human Labour	Man days	25.39	3962.29	9.24
2	Bullock	Pairs/day	0	0	0
3	Tractor	Hours	3.71	4507.75	10.51
4	Machinery	Hours	0	0	0
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	4.12	790.4	1.84
6	Seed Inter Crop	Kgs.	0	0	0
7	FYM	Quintal	4.39	13173.33	30.72
8	Fertilizer + micronutrients	Quintal	8.65	8296.46	19.35
9	Pesticides (PPC)	Kgs / liters	1.85	1852.5	4.32
10	Irrigation	Number	0	0	0
11	Repairs		0	0	0
12	Msc. Charges (Marketing costs etc)		0	0	0
13	Depreciation charges		0	3.33	0.01
14	Land revenue and Taxes		0	0	0
<b>II</b>	<b>Cost B1</b>				
16	Interest on working capital			2894.72	6.75
17	<b>Cost B1 = (Cost A1 + sum of 15 and 16)</b>			35480.79	82.75
<b>III</b>	<b>Cost B2</b>				
18	Rental Value of Land			111.11	0.26
19	<b>Cost B2 = (Cost B1 + Rental value)</b>			35591.9	83.01
<b>IV</b>	<b>Cost C1</b>				
20	Family Human Labour		21.41	3375.67	7.87
21	<b>Cost C1 = (Cost B2 + Family Labour)</b>			38967.57	90.89
<b>V</b>	<b>Cost C2</b>				
22	Risk Premium			10	0.02
23	<b>Cost C2 = (Cost C1 + Risk Premium)</b>			38977.57	90.91
<b>VI</b>	<b>Cost C3</b>				
24	Managerial Cost			3897.76	9.09
25	<b>Cost C3 = (Cost C2 + Managerial Cost)</b>			42875.32	100
<b>VII</b>	<b>Economics of the Crop</b>				
a.	Main Product	a) Main Product (q)	141.34	207297.04	
		b) Main Crop Sales Price (Rs.)		1466.67	
b.	Gross Income (Rs.)			207297.04	
c.	Net Income (Rs.)			164421.71	
d.	Cost per Quintal (Rs./q.)			303.35	
e.	Benefit Cost Ratio (BC Ratio)			1:4.8	

**Adequacy of fodder:** The data regarding the adequacy of fodder in Handrahalu-2 Micro watershed is presented in Table 25. The results indicate that, 2.94 per cent of the households opined that dry fodder was adequate and 11.76 per cent of them opined dry fodder was inadequate. With respect to green fodder availability, 2.94 percent of them opined it was sufficient and 11.76 percent of them opined it was insufficient.

**Table 25. Adequacy of fodder in Handrahalu-2 micro-watershed**

Sl.No.	Particulars	LL (5)		MF (6)		SF (10)		SMF (13)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Adequate-Dry Fodder	0	0	1	16.67	0	0	0	0	1	2.94
2	Inadequate-Dry Fodder	0	0	1	16.67	1	10	2	15.4	4	11.76
3	Adequate-Green Fodder	0	0	1	16.67	0	0	0	0	1	2.94
4	Inadequate-Green Fodder	0	0	1	16.67	1	10	2	15.4	4	11.76

**Average annual gross income:** The data regarding the annual gross income in Handrahalu-2 Micro watershed is presented in Table 26. The results indicate that, the farmers have annual gross income of Rs. 68723.53 in micro-watershed, of which Rs. 60282.35 is from agriculture itself.

**Table 26. Average annual gross income in Handrahalu-2 micro-watershed**

Sl.No.	Particulars	LL (5)	MF (6)	SF (10)	SMF (13)	All (34)
		Rs.	Rs.	Rs.	Rs.	Rs.
1	Service/salary	0	20000	0	0	3529.41
2	Wage	0	1333.33	8400	1923.08	3441.18
3	Agriculture	0	53000	64700	83430.8	60282.4
4	Dairy Farm	0	2500	0	2692.31	1470.59
Income(Rs.)		0	76833.3	73100	88046.2	68723.5

**Average annual Expenditure:** The data regarding the average annual expenditure in Handrahalu-2 Micro watershed is presented in Table 27. The results indicate that, the farmers have annual gross expenditure of Rs. 197041.03 in micro-watershed, of which Rs. 34558.82 is from agriculture itself.

**Table 27. Average annual Expenditure in Handrahalu-2 micro-watershed**

Sl.No.	Particulars	LL (5)	MF (6)	SF (10)	SMF (13)	All (34)
		Rs.	Rs.	Rs.	Rs.	Rs.
1	Service/salary	0	50000	0	0	1470.59
2	Wage	0	1000	6400	15000	1411.76
3	Agriculture	0	25833.3	40500	47307.7	34558.8
4	Dairy Farm	0	5000	0	6000	500
Total		0	81833.3	46900	68307.7	197041

**Forest species grown:** The data regarding forest species grown in Handrahalu-2 Micro watershed is presented in Table 28. The results indicate that, households have planted 1 teak trees, 36 neem trees, 4 tamarind trees, 1 acacia trees and 6 banyan trees together in both field and backyard.



**Table 28. Forest species grown in Handrahalu-2 micro-watershed**

Sl.No.	Particulars	LL (5)		MF (6)		SF (10)		SMF (13)		All (34)	
		F	B	F	B	F	B	F	B	F	B
1	Teak	0	0	1	0	0	0	0	0	1	0
2	Neem	0	0	2	0	19	0	15	0	36	0
3	Tamarind	0	0	1	0	2	0	1	0	4	0
4	Acacia	0	0	0	0	1	0	0	0	1	0
5	Banyan	0	0	0	0	3	0	3	0	6	0

\*F= Field B=Back Yard

**Average additional investment capacity:** The data regarding average additional investment capacity in Handrahalu-2 Micro watershed is presented in Table 29. The results indicate that, households have an average investment capacity of Rs. 3705.88 for land development, Rs. 941.18 for creation of irrigation facility, Rs.2235.29 for adoption of improved livestock breeds and Rs.1794.12 adoption of improved crop production activities.

**Table 29. Average additional investment capacity of households in Handrahalu-2 micro-watershed**

Sl.No.	Particulars	LL (5)	MF (6)	SF (10)	SMF (13)	All (34)
		Rs.	Rs.	Rs.	Rs.	Rs.
1	Land development	0	4166.67	3400	5153.85	3705.88
2	Irrigation facility	0	0	100	2384.62	941.18
3	Improved crop production	0	3000	1600	3230.77	2235.29
4	Improved livestock management	0	2833.33	800	2769.23	1794.12

**Source of funds for additional investment:** The data regarding source of funds for additional investment in Handrahalu-2 Micro watershed is presented in Table 30. The results indicate that, the sources of finance raised from bank as a loan and from own sources for land development were 5.88 and 55.88 per cent, for irrigation facility was 5.88 and 941.18 per cent.

**Table 30. Source of funds for additional investment in Handrahalu-2 micro-watershed**

Sl.No	Item	Land development		Irrigation facility		Improved crop production		Improved livestock management	
		N	%	N	%	N	%	N	%
1	Loan from bank	2	5.88	2	5.88	1	2.94	0	0
2	Own funds	19	55.88	2	5.88	14	41.18	10	29.4

**Marketing of agricultural produce:** The data regarding marketing of the agricultural produce in Handrahalu-2 Micro watershed is presented in Table 31. The results indicated that, 100.00 per cent of output of Bajra was sold in the market with average price of Rs. 1400.00; 100.00 per cent of output of Bengal gram was sold in the market with average price of Rs. 2000.00; 100.00 per cent of output of Cotton was sold in the market with average price of Rs. 5000.00; 100.00 per cent of output of Green gram was sold in the

market with average price of Rs. 4350.00 and 100.00 per cent of output of Jowar was sold in the market with average price of Rs. 1600.00.

**Table 31. Marketing of agricultural produce in Handrahalu-2 micro-watershed**

Sl.No	Crops	Output obtained (q)	Output retained (q)	Output sold (q)	Output sold (%)	Avg. Price obtained (Rs/q)
1	Bajra	63	0	63	100	1400
2	Bengal gram (Kadale)	7	0	7	100	2000
3	Cotton	25	0	25	100	5000
4	Green gram	29	0	29	100	4350
5	Jowar	28	0	28	100	1600
6	Maize	355	25	330	93	1322
7	Onion	275	0	275	100	1467
8	Sorghum	22	0	22	100	2600

**Marketing channels used for sale of agricultural produce:** The data regarding marketing channels used for sale of agricultural produce in Handrahalu-2 Micro watershed is presented in Table 32. The results indicated that, 61.76 cent of the households have sold agricultural produce to the local/village merchants, 38.24 per cent have sold to Agent/Traders and 2.94 per cent of regulated market.

**Table 32. Marketing channels used for sale of agricultural produce in Handrahalu-2 micro-watershed**

Sl.No.	Particulars	LL (5)		MF (6)		SF (10)		SMF (13)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Agent/Traders	0	0	0	0	7	70	6	46.2	13	38.24
2	Local/village Merchant	0	0	7	117	4	40	10	76.9	21	61.76
3	Regulated Market	0	0	0	0	1	10	0	0	1	2.94

**Mode of transport of agricultural produce:** The data regarding mode of transport of agricultural produce in Handrahalu-2 Micro watershed is presented in Table 33. The results indicated that, 67.65 cent of the households have used tractor and 14.71 per cent have used Cart for the transport of agriculture commodity.

**Table 33. Mode of transport of agricultural produce in Handrahalu-2 micro-watershed**

Sl.No.	Particulars	LL (5)		MF (6)		SF (10)		SMF (13)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Cart	0	0	1	17	2	20	2	15.4	5	14.71
2	Tractor	0	0	4	67	7	70	12	92.3	23	67.65
3	Truck	0	0	2	33	3	30	2	15.4	7	20.59

**Incidence of soil and water erosion problems:** The data regarding incidence of incidence of soil and water erosion problems in Handrahalu-2 Micro watershed is presented in Table 34. The results indicate that, 61.76 per cent of the households have experienced soil and water erosion problems.

**Table 34. Incidence of soil and water erosion problems in Handrahalu-2 micro-watershed**

Sl.No.	Particulars	LL (5)		MF (6)		SF (10)		SMF (13)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Soil and water erosion problems in the farm	0	0	5	83	5	50	11	85	21	61.76

**Interest towards soil testing:** The data regarding Interest shown towards soil testing in Handrahalu-2 Micro watershed is presented in Table 35. The results indicated that, 76.47 per cent of the households were interested towards soil testing.

**Table 35. Interest regarding soil testing in Handrahalu-2 micro-watershed**

Sl.No.	Particulars	LL (5)		MF (6)		SF (10)		SMF (13)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Interest in soil test	0	0	5	83	9	90	12	92	26	76.47

**Soil and water conservation practices and structures adopted:** The data regarding soil and water conservation practices and structures adopted in Handrahalu-2 Micro watershed is presented in Table 36. The results indicated that 100 per cent of farmers practicing summer ploughing as soil and water conservation practice.

**Table 36. Soil and water conservation practices and structures adopted in Handrahalu-2 micro-watershed**

Sl.No.	Particulars	LL (5)		MF (6)		SF (10)		SMF (13)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Field Bunding	0	0	2	33	0	0	2	15.4	4	11.76

**Status of soil and water conservation structures:** The data regarding status soil and water conservation structures adopted in Handrahalu-2 Micro watershed is presented in Table 37. The results indicated that, the households have adopted field bunding as a soil and water conservation structures out of which 100.00 per cent was slightly damaged.

**Table 37. Status of soil and water conservation structures in Handrahalu-2 micro-watershed**

Sl. No	Item	Good		Slightly Damaged		Severely Damaged		Full Replacement Required	
		N	%	N	%	N	%	N	%
1	Field Bunding	0	0	4	100	0	0	0	0

**Table 38. Agencies involved in the soil and water conservation structures in Handrahalu-2 micro-watershed**

Sl.No.	Particulars	LL (5)		MF (6)		SF (10)		SMF (13)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Own	0	0	2	33	0	0	0	0	2	5.88
2	Other	0	0	0	0	0	0	2	15	2	5.88

**Agencies involved in the soil and water conservation structures:** The data regarding Agencies involved in the soil and water conservation structures adopted in Handrahalu-2

Micro watershed is presented in Table 38. The results indicated that, 5.88 per cent of the households have adopted by their own.

**Usage pattern of fuel for domestic use:** The data on usage pattern of fuel for domestic use in Handrahalu-2 Micro watershed is presented in Table 39. The results indicated that, firewood was the major source of fuel for domestic use for 79.41 per cent of the households followed by LPG (23.53%).

**Table 39. Usage pattern of fuel for domestic use in Handrahalu-2 micro-watershed**

Sl.No.	Particulars	LL (5)		MF (6)		SF (10)		SMF (13)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Fire Wood	1	20	6	100	9	90	11	84.6	27	79.41
2	LPG	4	80	0	0	1	10	3	23.1	8	23.53

**Source of drinking water:** The data on source of drinking water in Handrahalu-2 Micro watershed is presented in Table 40. The results indicated that, piped waters supply was the major source for drinking water for 73.53 per cent of the households followed by bore well water (26.47%).

**Table 40. Source of drinking water in Handrahalu-2 micro-watershed**

Sl.No.	Particulars	LL (5)		MF (6)		SF (10)		SMF (13)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Piped supply	3	60	3	50	10	100	9	69.2	25	73.53
2	Bore Well	2	40	3	50	0	0	4	30.8	9	26.47

**Source of light:** The data on source of light in Handrahalu-2 Micro watershed is presented in Table 41. The results indicated that, electricity was the major source of light for 100.00 per cent of the households.

**Table 41. Source of light in Handrahalu-2 micro-watershed**

Sl.No.	Particulars	LL (5)		MF (6)		SF (10)		SMF (13)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Electricity	5	100	6	100	10	100	13	100	34	100

**Existence of sanitary toilet facility:** The data on availability of toilet facility in Handrahalu-2 Micro watershed is presented in Table 42. The results indicated that, 32.35 per cent of the households possess toilets.

**Table 42. Existence of sanitary toilet facility in Handrahalu-2 micro-watershed**

Sl.No.	Particulars	LL (5)		MF (6)		SF (10)		SMF (13)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Sanitary toilet facility	5	100	2	33	1	10	3	23	11	32.4

**Possession of PDS card:** The data regarding possession of PDS card in Handrahalu-2 Micro watershed is presented in Table 43. The results indicated that, 94.12 per cent of the households possessed BPL card, 2.94 per cent possessed APL card and 2.94 per cent do not possess PDS card.

**Table 43. Possession of PDS card in Handrahalu-2 micro-watershed**

Sl.No.	Particulars	LL (5)		MF (6)		SF (10)		SMF (13)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	APL	0	0	0	0	0	0	1	7.7	1	2.94
2	BPL	5	100	6	100	9	90	12	92	32	94.12
3	Not Possessed	0	0	0	0	1	10	0	0	1	2.94

**Participation in NREGA programme:** The data regarding Participation in NREGA programme in Handrahalu-2 Micro watershed is presented in Table 44. The results indicated that, only 41.18 percent of the households have participated in NREGA programme.

**Table 44. Participation in NREGA programme in Handrahalu-2 micro-watershed**

Sl.No.	Particulars	LL (5)		MF (6)		SF (10)		SMF (13)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Participation in NREGA programme	0	0	1	16.7	5	50	8	61.5	14	41.2

**Adequacy of food items:** The data regarding adequacy of food items in Handrahalu-2 Micro watershed is presented in Table 45. The results indicated that, the extent of adequacy of food items for cereals, pulses, Oilseeds and vegetables were 73.53, 67.65, 14.71, 58.82 per cent respectively, similarly for Fruits (23.53%), milk (79.41%), Egg (26.47%), and Meat (23.53%).

**Table 45. Adequacy of food items in Handrahalu-2 micro-watershed**

Sl.No.	Particulars	LL (5)		MF (6)		SF (10)		SMF (13)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Cereals	5	100	5	83.3	5	50	10	76.9	25	73.53
2	Pulses	5	100	4	66.7	5	50	9	69.2	23	67.65
3	Oilseed	0	0	2	33.3	2	20	1	7.69	5	14.71
4	Vegetables	3	60	4	66.7	4	40	9	69.2	20	58.82
5	Fruits	0	0	2	33.3	2	20	4	30.8	8	23.53
6	Milk	5	100	6	100	4	40	12	92.3	27	79.41
7	Egg	1	20	2	33.3	2	20	4	30.8	9	26.47
8	Meat	1	20	2	33.3	3	30	2	15.4	8	23.53

**Table 46. Inadequacy of food items in Handrahalu-2 micro-watershed**

Sl.No.	Particulars	LL (5)		MF (6)		SF (10)		SMF (13)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Cereals	0	0	1	16.7	1	10	3	23.1	5	14.71
2	Pulses	0	0	2	33.3	1	10	3	23.1	6	17.65
3	Oilseed	5	100	4	66.7	4	40	14	108	27	79.41
4	Vegetables	2	40	2	33.3	2	20	4	30.8	10	29.41
5	Fruits	5	100	4	66.7	3	30	8	61.5	20	58.82
6	Milk	0	0	0	0	2	20	0	0	2	5.88
7	Egg	4	80	3	50	3	30	9	69.2	19	55.88
8	Meat	4	80	3	50	2	20	9	69.2	18	52.94

**Inadequacy of food items:** The data regarding in adequacy of food items in Handrahalu-2 Micro watershed is presented in Table 46. The results indicated that, the extent of in

adequacy of food items for cereals, pulses, Oilseeds and vegetables were 14.71, 17.65, 79.41, 29.41 and 52.94 per cent respectively, similarly for fruits (58.82%), milk (5.88%), egg (55.88%) and meat (52.94%).

**Farming constraints:** The data regarding farming constraints experienced by households in Handrahalu-2 Micro watershed is presented in Table 47. The results indicated that, lower fertility status of the soil was the constraint experienced by (61.76 %) per cent of the households, wild animal menace on farm field (76.47%), frequent incidence of pest and diseases (35.29%), inadequacy of irrigation water (50.00%), high cost of fertilizers and plant protection chemicals (73.53%), high rate of interest on credit (61.76%), low price for the agricultural commodities (47.06 %), lack of marketing facilities in the area (61.76%), inadequate extension services (38.24 %), lack of transport for safe transport of the agricultural produce to the market (47.06%), less rainfall (8.82%), source of agri-technology information (Newspaper/Tv/Mobile) (8.82%).

**Table 47. Farming constraints experienced in Handrahalu-2 micro-watershed**

SN	Particulars	LL (5)		MF (6)		SF (10)		SMF (13)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Lower fertility status of the soil	0	0	5	83.33	5	50	11	84.62	21	61.76
2	Wild animal menace on farm field	0	0	5	83.33	9	90	12	92.31	26	76.47
3	Frequent incidence of pest and diseases	0	0	3	50	5	50	4	30.77	12	35.29
4	Inadequacy of irrigation water	0	0	3	50	6	60	8	61.54	17	50
5	High cost of Fertilizers and plant protection chemicals	0	0	4	66.67	10	100	11	84.62	25	73.53
6	High rate of interest on credit	0	0	3	50	9	90	9	69.23	21	61.76
7	Low price for the agricultural commodities	0	0	4	66.67	5	50	7	53.85	16	47.06
8	Lack of marketing facilities in the area	0	0	4	66.67	8	80	9	69.23	21	61.76
9	Inadequate extension services	0	0	1	16.67	7	70	5	38.46	13	38.24
10	Lack of transport for safe transport of the Agril produce to the market.	0	0	5	83.33	4	40	7	53.85	16	47.06
11	Less rainfall	0	0	1	16.67	1	10	1	7.69	3	8.82
12	Source of Agri-technology information	0	0	0	0	0	0	3	23.08	3	8.82

**SUMMARY AND IMPLICATIONS**

In order to assess the socio-economic condition of the farmers in the watershed 34 households located in the micro watershed were interviewed for the survey. The study was conducted in Handrahalu-2 micro-watershed (Tadakal sub-watershed, Koppala taluk & District) is located at North latitude 15° 19' 52.722" and 15° 17' 50.924" and East longitude 76° 3' 8.831" and 76° 1' 41.664" covering an area of about 450.01 ha bounded by under Handrala, Hireshindogi and Hanavala Villages.

Socio-economic analysis of Handrahalu-2 micro watersheds of Tadakal sub-watershed, Koppala taluk & District indicated that, out of the total sample of 34 farmers were sampled in Handrahalu-2 micro-watershed among households surveyed 6 (17.65%) were marginal, 10 (29.41%) were small and 13 (38.24 %) were semi medium farmers. 5 landless farmers were also interviewed for the survey. The population characteristics of households indicated that, there were 92 (57.14%) men and 69 (42.86 %) were women. The average population of landless was 4.8, marginal farmers were 4.8, small farmers were 5 and semi medium farmers were 4.5. Majority of the respondents (37.27%) were in the age group of 16-35 years.

Education level of the sample households indicated that, there were 37.27 per cent illiterates, 60.87 per cent pre university education and 2.48 per cent attained graduation. About, 97.06 per cent of household heads practicing agriculture. Agriculture was the major occupation for 59.63 per cent of the household members.

In the study area, 79.41 per cent of the households possess katcha house and 2.94 per cent possess pucca house. The durable assets owned by the households showed that, 70.59 per cent possess TV, 23.53 per cent possess mixer grinder, 94.12 per cent possess mobile phones and 20.59 per cent possess motor cycles. Farm implements owned by the households indicated that, 14.71 per cent of the households possess plough, 2.94 per cent possess tractor, 20.59 per cent possess bullock cart.

Regarding livestock possession by the households, 17.65 per cent possess local cow. The average labour availability in the study area showed that, own labour men available in the micro watershed was 1.68, women available in the micro watershed was 1.38, hired labour (men) available was 10 and hired labour (women) available was 8.24. Further, 17.65 per cent of the households opined that hired labour was inadequate during the agricultural season.

Out of the total land holding of the sample respondents 65.19 per cent (39.67 ha) of the area is under dry condition and the remaining 34.81 per cent area is irrigated land. There were 16.00 live bore wells and 15.00 dry bore wells among the sampled households. Bore well was the major source of irrigation for 47.06 per cent of the

households. The major crops grown by sample farmers are Maize, Sunflower, Green gram, Cotton and Onion and cropping intensity was recorded as 91.83 per cent.

Out of the sample households 73.53 percent possessed bank account. Majority of the respondents (100.00%) have borrowed loan for agriculture purpose. Regarding the opinion on institutional sources of credit, 69.23 per cent of the households opined that credit helped to perform timely agricultural operations.

The per hectare cost of cultivation for Maize, Sunflower, Green gram, Cotton and Onion was Rs.34869.68 , 44381.39, 28905.04, 28381.13 and 42875.32 with benefit cost ratio of 1:1.10, 1: 3.60, 1: 1.30, 1: 2.10 and 1:4.80 respectively. Further, 2.94 per cent of the households opined that dry fodder was adequate and 2.94 per cent of the households have opined that the green fodder was adequate.

The average annual gross income of the farmers was Rs. 68723.53 in micro-watershed, of which Rs. 60282.35 comes from agriculture. Sampled households have grown 1 horticulture trees and 48 forestry trees together in the fields and back yards. Households have an average investment capacity of Rs. 3705.88 for land development and Rs. 941.18 for irrigation facility. Source of funds for additional investment is concerned, 55.88 per cent depends on own funds and 5.88 per cent depends on bank loan for land development activities.

Regarding marketing channels, 61.76 per cent of the households have sold agricultural produce to the local/village merchants, while, 2.94 per cent have sold in regulated markets. Further, 67.65 per cent of the households have used tractor for the transport of agriculture commodity. Majority of the farmers (61.76%) have experienced soil and water erosion problems in the watershed and 76.47 per cent of the households were interested towards soil testing.

Firewood was the major source of fuel for domestic use for 79.41 per cent of the households and 23.53 per cent households has LPG connection. Piped supply was the major source for drinking water for 73.53 per cent of the households. Electricity was the major source of light for 100.00 per cent of the households. In the study area, 32.35 per cent of the households possess toilet facility.

Regarding possession of PDS card, 94.12 per cent of the households possessed BPL card, 2.94 per cent of the household's possessed APL card and 2.94 per cent of the household's were not having ration cards. Households opined that, the requirement of cereals (73.53%), pulses (67.65%) and oilseeds (14.71%) are adequate for consumption.

Farming constraints experienced by households in the micro watersheds were lower fertility status of the soil (61.76%) wild animal menace on farm field (76.47%), frequent incidence of pest and diseases (35.29%), inadequacy of irrigation water (50.00%), high cost of fertilizers and plant protection chemicals (73.53%), high rate of



interest on credit (61.76%), low price for the agricultural commodities (47.06%), lack of marketing facilities in the area (61.76%), inadequate extension services (38.24%), lack of transport for safe transport of the agricultural produce to the market (47.06%), Less rainfall (8.82%) and Source of Agri-technology information (Newspaper/ TV/Mobile) (8.82%).

### **Implications of the survey**

- ✓ Result indicated that, there were 37.27 per cent were illiterate hence, extension methodologies such as demonstration, street play, drama, video shows will be effective in dissemination of the technologies in the micro watershed.
- ✓ The data indicate that, 79.41 per cent of the households possess katcha house. Hence, the development department while implementing the watershed plan should focus on agriculture to enhance the productivity of major crops in the area to increase the income of the farmers.
- ✓ Results indicated that the local institutional participation of the household members in the micro watershed is minimal hence, activities like membership campaign, awareness creation about the benefits of membership in local institutions and strengths of organized groups must be conveyed.
- ✓ Majority of the households in the watershed have experience in use of mobile phones, and television hence, these mass media can be effectively utilized for transfer of technology as well as for information dissemination.
- ✓ The farm machinery/implement possession in the micro watershed was found to be minimum the reasons may lack of knowledge or lack of financial ability which can be addressed through training on use of different farm implements, providing information on different sources of finance for purchase of farm implements.
- ✓ The possession of livestock such as crossbred cow found is less hence, farmers must be made aware of the benefits of crossbred cow in increased milk production.
- ✓ The possession of livestock such as sheep, goat and poultry was found to be low hence, farmers may be informed the role of subsidiary enterprises in enhancing the income and information on financial support for subsidiary activities.
- ✓ The data indicate that, job/work was the reason for all the migrants hence, farmers may be trained on profitable agriculture or self employment such as animal husbandry, plate making, sheep rearing, goat rearing, rabbit rearing with suitable information on sources of financial support.
- ✓ The results indicate that there was a change in quality of life due to migration hence, the developmental departments should take actions to arrest migration and to improve the quality of the life in rural areas.
- ✓ Households possess 25.86ha (65.19 %) of dry land and 13.81ha (34.81 %) of irrigated land hence, the availability of the dry land agricultural technologies such as short duration crops, high yielding drought resistance crop varieties, drip irrigation

technology and subsidy information will be helpful for the farmers to enhance the productivity of land and as well as farmers income.

- ✓ Few of the bore well in micro watershed found non functional hence, farmers may be trained on possibility of bore well rejuvenation.
- ✓ Bore well was major source of irrigation for 47.06 per cent of the households. hence, in order to increase the area under irrigation as well as to increase the water use efficiency farmers may trained on drip irrigation and provide the information on subsidy for drip irrigation equipment's along with the information on different agencies which provides the financial assistance for drip irrigation.
- ✓ The cropping intensity in the micro watershed was found to be (91.83 %) hence, care must be taken by the implementing agency to bring uncultivated land into cultivation through suitable measures.
- ✓ Many of the household members have borrowed loan from cooperative banks which has higher rate of interest hence, farmers may be sensitized on the different sources of credit with lesser interest rate such SHGs etc.
- ✓ The results indicated the non availability of both green and dry fodder throughout the year hence, fodder development activities can be taken up in the micro watershed.
- ✓ The average annual gross income of the households Rs.60282.35 from agriculture, and Rs. 3441.18 from wages. Agriculture was found to be the major source of income for households hence; the development activities should focus on productivity enhancement, marketing arrangements and agricultural technology dissemination to have a direct impact on the farmers.
- ✓ The cultivation of forest species is found minimal hence, information and production technology related to agro-forestry and integrated farming system.
- ✓ The data indicated that, 61.76 per cent of the households have experienced soil and water erosion problems. Hence, those farmers who reported the soil and water erosion problems may be given attention while implementation of the watershed development plan.
- ✓ The data indicated that, 76.47 per cent of the households have interest in soil testing hence, farmers must be provided with the information on various institutions which are involved in soil testing for the benefit of the farmers.
- ✓ Except summer ploughing the adoption of other soil and water conservation structures is minimum hence, the farmers in the micro watershed should be sensitized on the use of different conservation structures for soil water conservation.
- ✓ Cereals and pulses found be adequate for per cent of the households respectively hence, farm households and the farm women must be trained on importance of balanced nutrition and role of vegetable, milk, egg, meat in balanced diet.
- ✓ Lower fertility status of the soil (61.76%), wild animal menace on farm field (76.47%), frequent incidence of pest and diseases (35.29%), high cost of fertilizers and plant protection chemicals (73.53%), high rate of interest on credit (61.76%),

low price for the agricultural commodities (47.06%), lack of marketing facilities in the area (61.76%), inadequate extension services (38.24%), lack of transport for safe transport of the agricultural produce to the market (47.06%) were the major farming constraints experienced hence, these constraints must be addressed immediately for the welfare of the farmers. Awareness to be created among the farmers to approach nearest KVKs/RSKs and other developmental departments for technical and for subsidized inputs and utilize the well established regulated markets, approaching the contract firms, direct markets to avoid the involvement of middlemen.