



हर कदम, हर डगर
किसानों का हमसफर
भारतीय कृषि अनुसंधान परिषद

AgriSearch with a human touch

**LAND RESOURCE INVENTORY AND SOCIO-ECONOMIC STATUS OF
FARM HOUSEHOLDS FOR WATERSHED PLANNING AND
DEVELOPMENT**

YADGIR RF-1 (4D5B1F2a) MICROWATERSHED

Gurumitkal Hobli, Yadgir Taluk and District, Karnataka

Karnataka Watershed Development Project – II

SUJALA – III

World Bank funded Project



ICAR – NATIONAL BUREAU OF SOIL SURVEY AND LAND USE PLANNING



ICAR - NBSS & LUP



**WATERSHED DEVELOPMENT DEPARTMENT
GOVT. OF KARNATAKA, BANGALORE**



About ICAR - NBSS&LUP

The ICAR-National Bureau of Soil Survey and Land Use Planning (ICAR-NBSS&LUP), Nagpur, a premier Institute of the Indian Council of Agricultural Research (ICAR), was set up during 1976 with the objective to prepare soil resource maps at national, state and district levels and to provide research inputs in soil resource mapping and its applications, land evaluation, land use planning, land resource management, and database management using GIS for optimising land use on different kinds of soils in the country.

The Bureau has been engaged in carrying out soil resource survey, agro-ecological and soil degradation mapping at the country, state and district levels for qualitative assessment and monitoring the soil health towards viable land use planning. The research activities have resulted in identifying the soil potentials and problems, and the various applications of the soil surveys with the ultimate objective of sustainable agricultural development. The Bureau has the mandate to correlate and classify soils of the country and maintain a National Register of all the established soil series. The Institute is also imparting in-service training to staff of the soil survey agencies in the area of soil survey, land evaluation and soil survey interpretations for land use planning. The Bureau in collaboration with Panjabrao Krishi Vidyapeeth, Akola is running post-graduate teaching and research programme in land resource management, leading to M.Sc. and Ph.D. degrees.

Citation: Rajendra Hegde, Ramesh Kumar, S.C., B.A. Dhanorkar, S. Srinivas, M.Lalitha, K.V. Niranjana, R.S. Reddy and S.K. Singh (2019). "Land Resource Inventory and Socio-Economic Status of Farm Households for Watershed Planning and Development of Yadgir Rf-1 (4D5B1F2a) Microwatershed, Gurumitkal Hobli, Yadgir Taluk and District, Karnataka", ICAR-NBSS&LUP Sujala MWS Publ.324, ICAR – NBSS & LUP, RC, Bangalore. p.129 & 30.

TO OBTAIN COPIES,

Please write to:

Director, ICAR - NBSS & LUP,

Amaravati Road, NAGPUR - 440 033, India

Phone : (0712) 2500386, 2500664, 2500545 (O)

Telefax : 0712-2522534

E-Mail : director@nbsslup.ernet.in

Website URL : nbsslup.in

Or

Head, Regional Centre, ICAR - NBSS&LUP, Hebbal, Bangalore - 560 024

Phone : (080) 23412242, 23510350 (O)

Telefax : 080-23510350

E-Mail : nbssrcb@gmail.com

ICAR-NBSS&LUP Sujala MWS Publ.324



हर कदम, हर डगर
किसानों का हमसफर
भारतीय कृषि अनुसंधान परिषद

Agriseach with a human touch

LAND RESOURCE INVENTORY AND SOCIO-ECONOMIC STATUS OF FARM HOUSEHOLDS FOR WATERSHED PLANNING AND DEVELOPMENT

YADGIR RF-1 (4D5B1F2a) MICROWATERSHED

Gurumitkal Hobli, Yadgir Taluk and District, Karnataka

Karnataka Watershed Development Project – II

Sujala-III

World Bank funded Project



**ICAR – NATIONAL BUREAU OF SOIL SURVEY AND LAND USE
PLANNING**



ICAR - NBSS & LUP



**WATERSHED DEVELOPMENT DEPARTMENT, GOVT. OF
KARNATAKA, BANGALORE**



PREFACE

In Karnataka, as in other Indian States, the livelihoods of rural people are intertwined with farming pursuits. The challenges in agriculture are seriously threatening the livelihood of a large number of farmers as they have been practicing farming in contextual factors beyond their control. Climatic factors are the most important ones and have become much more significant in recent times due to rapid climate changes induced by intensive anthropogenic activities affecting our ecosystem in multiple ways. Climate change has become the reality, it is happening and efforts to evolve and demonstrate climate resilient technologies have become essential. Due to the already over stressed scenario of agrarian sector, the climate change is resulting in manifold increase in the complexities, pushing the rural mass to face more and more unpredictable situations. The rising temperatures and unpredictable rainfall patterns are going to test seriously the informed decisions farmers have to make in order to survive in farming and sustain their livelihood.

It is generally recognized that impacts of climate change shall not be uniform across the globe. It is said that impact of climate change is more severe in South Asia. Based on the analysis of meteorological data, it is predicted that in India, there will be upward trend in mean temperature, downward trend in relative humidity, annual rainfall and number of wet days in a year. Also, in general, phenomena like erratic monsoon, spread of tropical diseases, rise in sea levels, changes in availability of fresh water, frequent floods, droughts, heat waves, storms and hurricanes are predicted. Each one of these adverse situations are already being experienced in various parts of India and also at the global level. Decline in agricultural productivity of small and marginal farmers becoming more vulnerable is already witnessed.

In Karnataka, more than 60 per cent of the population live in rural areas and depend on agriculture and allied activities for their livelihood. Though the state has achieved significant progress in increasing the yield of many crops, there is tremendous pressure on the land resources due to the growing and competing demands of various land uses. This is reflected in the alarming rate of land degradation observed. Already more than 50 per cent of the area is affected by various forms of degradation. If this trend continues, the sustainability of the fragile ecosystem will be badly affected. The adverse effects of change in the climatic factors are putting additional stress on the land resources and the farmers dependent on this.

The natural resources (land, water and vegetation) of the state need adequate and constant care and management, backed by site-specific technological interventions and investments particularly by the government. Detailed database pertaining to the nature of the land resources, their constraints, inherent potentials and suitability for various land

based rural enterprises, crops and other uses is a prerequisite for preparing location-specific action plans, which are in tune with the inherent capability of the resources. Any effort to evolve climate resilient technologies has to be based on the baseline scientific database. Then only one can expect effective implementation of climate resilient technologies, monitor the progress, make essential review of the strategy, and finally evaluate the effectiveness of the implemented programs. The information available at present on the land resources of the state are of general nature and useful only for general purpose planning. Since the need of the hour is to have site-specific information suitable for farm level planning and detailed characterization and delineation of the existing land resources of an area into similar management units is the only option.

ICAR-NBSS&LUP, Regional Centre, Bangalore has taken up a project sponsored by the Karnataka Watershed Development Project-II, (Sujala-III), Government of Karnataka funded by the World Bank under Component-1 Land Resource Inventory. This study was taken up to demonstrate the utility of such a database in reviewing, monitoring and evaluating all the land based watershed development programs on a scientific footing. To meet the requirements of various land use planners at grassroots level, the present study on “Land Resource Inventory and Socio-Economic Status of Farm Households for Watershed Planning and Development of YadgirRF1 Microwatershed, Yadgir Taluk and District, Karnataka” for integrated development was taken up in collaboration with the State Agricultural Universities, IISC, KRSRAC, KSNDMC as Consortia partners. The project provides detailed land resource information at cadastral level (1:7920 scale) for all the plots and socio-economic status of farm households covering thirty per cent farmers randomly selected representing landed and landless class of farmers in the microwatershed. The project report with the accompanying maps for the Microwatershed will provide required detailed database for evolving effective land use plan, alternative land use options and conservation plans for the planners, administrators, agricultural extension personnel, KVK officials, developmental departments and other land users to manage the land resources in a sustainable manner.

It is hoped that this database will be useful to the planners, administrators and developmental agencies working in the area in not only for formulating location specific developmental schemes but also for their effective monitoring at the village/watershed level.

Nagpur

Date: 21-08-2019

S.K. SINGH

Director, ICAR - NBSS&LUP Nagpur

Contributors

Dr. Rajendra Hegde Principal Scientist, Head & Project Leader, Sujala-III Project ICAR-NBSS&LUP, Regional Centre, Bangalore	Dr. S.K.Singh Director, ICAR-NBSS&LUP Coordinator, Sujala-III Project Nagpur
Soil Survey, Mapping & Report Preparation	
Dr. B.A. Dhanorkar	Sh. R.S. Reddy
Dr. K.V. Niranjana	Mr. Somashekar T N
	Smt. Chaitra, S.P.
	Dr. Gopali bardhan
	Ms. Arpitha
	Dr. Mahendra Kumar, M.B.
Field Work	
Sh. C.BacheGowda	Sh. Mahesh, D.B.
Sh. Somashekar	Sh. Ashok S Sindagi
Sh. M. Jayaramaiah	Sh. Veerabhadrapa B.
Sh. Paramesha, K.	Sh. Shankarappa
Sh. B. M. Narayana Reddy	Sh. Anand
	Sh. Arun N Kambar.
	Sh Kamalesh Awate
	Sh. Sharaan Kumar Huppar
	Sh. Yogesh H.N.
	Sh. Kalaveerachari R Kammar
GIS Work	
Dr. S.Srinivas	Sh. A.G.Devendra Prasad
Sh. D.H.Venkatesh	Sh. Prakashanaik, M.K.
Smt.K.Sujatha	Sh. Abhijith Sastry, N.S.
Smt. K.V.Archana	Sh. Sudip Kumar Suklabaidya
Sh. N. Maddileti	Sh. Avinash, K.N.
	Sh. Amar Suputhra, S
	Sh. Deepak, M.J.
	Smt. K.Karunya Lakshmi
	Ms. Seema, K.V.
	Ms. A. Rajab Nisha

Laboratory Analysis	
Dr. K.M.Nair	Ms. Steffi Peter
Smt. Arti Koyal	Ms. Thara, V.R
Smt. Parvathy	Ms. Roopa, G.
	Ms. Swati, H.
	Sh. Shantaveera Swami
	Ms. Shwetha, N.K.
	Smt. Ishrat Haji
	Ms. P. Pavan Kumari
	Ms. Padmaja
	Ms. Veena, M.
Socio-Economic Analysis	
Dr. S.C. Ramesh Kumar	Sh. M.K. Prakashanaik
	Mrs. Sowmya A.N
	Ms. Karuna V Kulkarni
	Sh. Vijay Kumar
	Sh. Pradyumna
	Ms. Sowmya K.B
	Mrs. Prathibha, D.G
	Sh. Rajendra,D
Soil & Water Conservation	
Sh. Sunil P. Maske	
Watershed Development Department, GoK, Bangalore	
Sh. Rajeev Ranjan IFS Project Director & Commissioner, WDD	Dr. A. Natarajan NRM Consultant, Sujala-III Project
Dr. S.D. Pathak IFS Executive Director & Chief Conservator of Forests, WDD	

PART-A

LAND RESOURCE INVENTORY

Contents

Preface		
Contributors		
Executive Summary		
Chapter 1	Introduction	1
Chapter 2	Geographical Setting	3
2.1	Location and Extent	3
2.2	Geology	3
2.3	Physiography	4
2.4	Drainage	4
2.5	Climate	4
2.6	Natural Vegetation	6
2.7	Land Utilization	6
Chapter 3	Survey Methodology	11
3.1	Base maps	11
3.2	Image Interpretation for Physiography	11
3.3	Field Investigation	14
3.4	Soil Mapping	16
3.5	Land Management Units	16
3.5	Laboratory Characterization	17
Chapter 4	The Soils	21
4.1	Soils of granite gneiss landscape	21
Chapter 5	Interpretation for Land Resource Management	33
5.1	Land Capability Classification	33
5.2	Soil Depth	35
5.3	Surface Soil Texture	36
5.4	Soil Gravelliness	37
5.5	Available Water Capacity	38
5.6	Soil Slope	39
5.7	Soil Erosion	40
Chapter 6	Fertility Status	43
6.1	Soil Reaction (pH)	43
6.2	Electrical Conductivity (EC)	43
6.3	Organic Carbon (OC)	43
6.4	Available Phosphorus	45
6.5	Available Potassium	45
6.6	Available Sulphur	45
6.7	Available Boron	46
6.8	Available Iron	46
6.9	Available Manganese	46
6.10	Available Copper	46
6.11	Available Zinc	50

Chapter 7	Land Suitability for Major Crops	51
7.1	Land suitability for Sorghum	51
7.2	Land suitability for Maize	52
7.3	Land suitability for Bajra	53
7.4	Land suitability for Groundnut	54
7.5	Land suitability for Sunflower	55
7.6	Land suitability for Redgram	56
7.7	Land suitability for Bengal gram	57
7.8	Land suitability for Cotton	58
7.9	Land suitability for Chilli	59
7.10	Land suitability for Tomato	60
7.11	Land suitability for Brinjal	61
7.12	Land suitability for Onion	62
7.13	Land suitability for Bhendi	63
7.14	Land suitability for Drumstick	64
7.15	Land suitability for Mango	65
7.16	Land suitability for Guava	66
7.17	Land suitability for Sapota	67
7.18	Land Suitability for Pomegranate	68
7.19	Land Suitability for Musambi	69
7.20	Land Suitability for Lime	70
7.21	Land Suitability for Amla	71
7.22	Land Suitability for Cashew	72
7.23	Land Suitability for Jackfruit	73
7.24	Land Suitability for Jamun	74
7.25	Land Suitability for Custard apple	75
7.26	Land Suitability for Tamarind	76
7.27	Land Suitability for Mulberry	77
7.28	Land Suitability for Marigold	78
7.29	Land Suitability for Chrysanthemum	79
7.30	Land management units	111
7.31	Proposed Crop Plan	112
Chapter 8	Soil Health Management	115
Chapter 9	Soil and Water conservation Treatment Plan	121
9.1	Treatment Plan	122
9.2	Recommended Soil and Water Conservation measures	125
9.3	Greening of Microwatershed	126
	References	129
	Appendix I	I-VIII
	Appendix II	XI-XVI
	Appendix III	XVII-XXII

LIST OF TABLES

2.1	Mean Monthly Rainfall, PET, 1/2 PET at Yadgir Taluk & District	5
2.2	Land Utilization in Yadgir district	7
3.1	Differentiating Characteristics used for Identifying Soil Series	16
3.2	Soil map unit description of YadgirRF1Microwatershed	17
4.1	Physical and Chemical Characteristics of Soil Series identified in YadgirRF1microwatershed	26
7.1	Soil-Site Characteristics of YadgirRF1Microwatershed	81
7.2	Land suitability criteria for Sorghum	82
7.3	Land suitability criteria for Maize	83
7.4	Land suitability criteria for Bajra	84
7.5	Land suitability criteria for Groundnut	85
7.6	Land suitability criteria for Sunflower	86
7.7	Land suitability criteria for Redgram	87
7.8	Land suitability criteria for Bengal gram	88
7.9	Land suitability criteria for Cotton	89
7.10	Land suitability criteria for Chilli	90
7.11	Land suitability criteria for Tomato	91
7.12	Land suitability criteria for Brinjal	92
7.13	Land suitability criteria for Onion	93
7.14	Land suitability criteria for Bhendi	94
7.15	Land suitability criteria for Drumstick	95
7.16	Land suitability criteria for Mango	96
7.17	Land suitability criteria for Guava	97
7.18	Land suitability criteria for Sapota	98
7.19	Land suitability criteria for Pomegranate	99
7.20	Land suitability criteria for Musambi	100
7.21	Land suitability criteria for Lime	101
7.22	Land suitability criteria for Amla	102
7.23	Land suitability criteria for Cashew	103
7.24	Land suitability criteria for Jackfruit	104
7.25	Land suitability criteria for Jamun	105
7.26	Land suitability criteria for Custard apple	106

7.27	Land suitability criteria for Tamarind	107
7.28	Land suitability criteria for Mulberry	108
7.29	Land suitability criteria for Marigold	109
7.30	Land suitability criteria for Chrysanthemum	110
7.31	Proposed Crop Plan for YadgirRF1Microwatershed	113

LIST OF FIGURES

2.1	Location map of YadgirRF1 Microwatershed	3
2.2 a	Granite and granite gneiss rock formation	4
2.3	Rainfall distribution in Yadgir Taluk & District	5
2.4	Natural vegetation of YadgirRF1Microwatershed	6
2.5	Current Land use map of YadgirRF1Microwatershed	7
2.6	Different crops and cropping systems in YadgirRF1Microwatershed	8
2.7	Location of Wells in YadgirRF1 microwatershed	9
3.1	Scanned and Digitized Cadastral map of YadgirRF1Microwatershed	13
3.2	Satellite image of YadgirRF1Microwatershed	13
3.3	Cadastral map overlaid on IRS PAN+LISS IV merged imagery of YadgirRF1Microwatershed	14
3.4	Location of profiles in a transect	15
3.5	Soil phase or management units of YadgirRF1Microwatershed	19
5.1	Land Capability Classification map of YadgirRF1Microwatershed	35
5.2	Soil Depth map of YadgirRF1Microwatershed	36
5.3	Surface Soil Texture map of YadgirRF1Microwatershed	37
5.4	Soil Gravelliness map of YadgirRF1Microwatershed	38
5.5	Soil Available Water Capacity map of YadgirRF1Microwatershed	39
5.6	Soil Slope map of YadgirRF1Microwatershed	40
5.7	Soil Erosion map of YadgirRF1Microwatershed	41
6.1	Soil Reaction (pH) map of YadgirRF1Microwatershed	44
6.2	Electrical Conductivity (EC) map of YadgirRF1Microwatershed	44
6.3	Soil Organic Carbon (OC) map of YadgirRF1Microwatershed	45
6.4	Soil Available Phosphorus map of YadgirRF1Microwatershed	46
6.5	Soil Available Potassium map of YadgirRF1Microwatershed	47
6.6	Soil Available Sulphur map of YadgirRF1Microwatershed	47
6.7	Soil Available Boron map of YadgirRF1Microwatershed	48
6.8	Soil Available Iron map of YadgirRF1Microwatershed	48
6.9	Soil Available Manganese map of YadgirRF1Microwatershed	49
6.10	Soil Available Copper map of YadgirRF1Microwatershed	49
6.11	Soil Available Zinc map of YadgirRF1Microwatershed	50

7.1	Land suitability for Sorghum	52
7.2	Land suitability for Maize	53
7.3	Land suitability for Bajra	54
7.4	Land suitability for Groundnut	55
7.5	Land suitability for Sunflower	56
7.6	Land suitability for Redgram	57
7.7	Land suitability for Bengal gram	58
7.8	Land suitability for Cotton	59
7.9	Land suitability for Chilli	60
7.10	Land suitability for Tomato	61
7.11	Land suitability for Brinjal	62
7.12	Land suitability for Onion	63
7.13	Land suitability for Bhendi	64
7.14	Land suitable for Drumstick	65
7.15	Land suitability for Mango	66
7.16	Land suitability for Guava	67
7.17	Land suitability for Sapota	68
7.18	Land suitability for Pomegranate	69
7.19	Land suitability for Musambi	70
7.20	Land suitability for Lime	71
7.21	Land suitability for Amla	72
7.22	Land suitability for Cashew	73
7.23	Land suitability for Jackfruit	74
7.24	Land suitability for Jamun	75
7.25	Land suitability for Custard apple	76
7.26	Land suitability for Tamarind	77
7.27	Land suitability for Mulberry	78
7.28	Land suitability for Marigold	79
7.29	Land suitability for Chrysanthemum	80
7.30	Land management units map of YadgirRF1Microwatershed	112
9.1	Soil and water conservation map of YadgirRF1Microwatershed	126

EXECUTIVE SUMMARY

The land resource inventory of Yadgir RFIMicrowatershed was conducted using village cadastral maps and IRS satellite imagery on 1:7920 scale. The false colour composites of IRS imagery were interpreted for physiography and the physiographic delineations were used as base for mapping soils. The soils were studied in several transects and a soil map was prepared with phases of soil series as mapping units. Random checks were made all over the area outside the transects to confirm and validate the soil map unit boundaries. The soil map shows the geographic distribution and extent, characteristics, classification, behavior and use potentials of the soils in the microwatershed.

The present study covers an area of 490 ha in Yadgir taluk & district, Karnataka. The climate is semiarid and categorized as drought-prone with an average annual rainfall of 866 mm, of which about 652 mm is received during south-west monsoon, 138 mm during north-east and the remaining 76 mm during the rest of the year. An area of 485 ha in the microwatershed is covered by soils, 0.25 ha by rock outcrops and 5 ha by others (habitation and water body). The salient findings from the land resource inventory are summarized briefly below.

- ❖ The soils belong to 6 soil series and 12 soil phases (management units) and 3 land management units.*
- ❖ The length of crop growing period is about 120-150 days starting from 1st week of June to 4th week of October.*
- ❖ From the master soil map, several interpretative and thematic maps like land capability, soil depth, surface soil texture, soil gravelliness, available water capacity, soil slope and soil erosion were generated.*
- ❖ Soil fertility status maps for macro and micronutrients were generated based on the surface soil samples collected at every 320 m grid interval.*
- ❖ Land suitability for growing 29 major agricultural and horticultural crops was assessed and maps showing the degree of suitability along with constraints were generated.*
- ❖ Entire area in the microwatershed is suitable for agriculture.*
- ❖ About 32 per cent area is very shallow (<25 cm), 42 per cent area is shallow (25-50 cm), 15 per cent area of the microwatershed has soils that are moderately shallow (50-75 cm) and 10 per cent area is moderately deep (75 - 100 cm).*
- ❖ About 41 per cent area in the microwatershed has sandy soils, 48 per cent loamy soils and 9 per cent clayey soils at the surface.*
- ❖ Maximum of 74 per cent area in the microwatershed is gravelly (15-35%) and 25 per cent is non gravelly (<15%).*
- ❖ About 10 per cent is medium (101-150 mm/m) in available water capacity and 89 is very low (<50 mm/m).*

- ❖ *About 63 per cent area in the microwatershed has very gently sloping (1-3% slope) lands and 36 per cent has gently sloping (3-5% slope) lands.*
- ❖ *An area of about 86 per cent is moderately (e2) eroded and 13 per cent area is severely (e3) eroded.*
- ❖ *An area of about 46 per cent is slightly acid (pH 6.0-6.5) in soil reaction and 53 per cent area is neutral (pH 6.5-7.3).*
- ❖ *The Electrical Conductivity (EC) of entire soils of the microwatershed is dominantly $<2 \text{ dsm}^{-1}$ indicating that the soils are non-saline.*
- ❖ *About 20 per cent of the soils are medium (0.5-0.75%) in organic carbon content and 79 per cent area is high ($>0.75\%$).*
- ❖ *Entire area is medium (23-57 kg/ha) in available phosphorus.*
- ❖ *About 70 per cent is medium (145-337 kg/ha) in available potassium and 29 per cent is low ($<145 \text{ kg/ha}$).*
- ❖ *Available sulphur is medium (10-20 ppm) in an entire area of about 99 per cent.*
- ❖ *Available boron is low ($<0.5 \text{ ppm}$) in an area of about 89 per cent and medium (0.5-1.0 ppm) in 10 per cent area of the microwatershed.*
- ❖ *Available iron is sufficient ($>4.5 \text{ ppm}$) in the entire area of the microwatershed.*
- ❖ *Available manganese and copper are sufficient in all the soils of the microwatershed.*
- ❖ *Available zinc is deficient ($<0.6 \text{ ppm}$) in an area of 12 per cent and sufficient ($>0.6 \text{ ppm}$) in 87 per cent area of the microwatershed.*
- ❖ *The land suitability for 29 major crops grown in the microwatershed were assessed and the areas that are highly suitable (S1) and moderately suitable (S2) are given below. It is however to be noted that a given soil may be suitable for various crops but what specific crop to be grown may be decided by the farmer looking to his capacity to invest on various inputs, marketing infrastructure, market price and finally the demand and supply position.*

Land suitability for various crops in the Microwatershed

Crop	Suitability Area in ha (%)		Crop	Suitability Area in ha (%)	
	Highly suitable (S1)	Moderately suitable (S2)		Highly suitable (S1)	Moderately suitable (S2)
<i>Sorghum</i>	29 (6)	20 (4)	<i>Guava</i>	-	-
<i>Maize</i>	-	20 (4)	<i>Sapota</i>	-	-
<i>Bajra</i>	29 (6)	20 (4)	<i>Pomegranate</i>	-	50 (10)
<i>Groundnut</i>	-	-	<i>Musambi</i>	-	50 (10)
<i>Sunflower</i>	1-	50 (10)	<i>Lime</i>	-	50 (10)
<i>Redgram</i>	-	50 (10)	<i>Amla</i>	-	49 (10)
<i>Bengal gram</i>	29 (6)	20 (4)	<i>Cashew</i>	-	-
<i>Cotton</i>	-	50 (10)	<i>Jackfruit</i>	-	-
<i>Chilli</i>	-	50 (10)	<i>Jamun</i>	-	-
<i>Tomato</i>	-	-	<i>Custard apple</i>	20 (4)	29 (6)
<i>Brinjal</i>	50 (10)	-	<i>Tamarind</i>	-	-
<i>Onion</i>	50 (10)	-	<i>Mulberry</i>	-	-
<i>Bhendi</i>	50 (10)	-	<i>Marigold</i>	-	50 (10)
<i>Drumstick</i>	-	50 (10)	<i>Chrysanthemum</i>	-	50 (10)
<i>Mango</i>	-	-			

- ❖ *Apart from the individual crop suitability, a proposed crop plan has been prepared for the identified LMUs by considering only the highly and moderately suitable lands for different crops and cropping systems with food, fodder, fibre and other horticulture crops.*
- ❖ *Maintaining soil-health is vital for crop production and conserve soil and land resource base for maintaining ecological balance and to mitigate climate change. For this, several ameliorative measures have been suggested for these problematic soils like saline/alkali, highly eroded, sandy soils etc.,*
- ❖ *Soil and water conservation treatment plan has been prepared that would help in identifying the sites to be treated and also the type of structures required.*
- ❖ *As part of the greening programme, several tree species have been suggested to be planted in marginal and submarginal lands, field bunds and also in the hillocks, mounds and ridges. This would help in not only supplementing the farm income but also provide fodder and fuel and generate lot of biomass which would help in maintaining an ecological balance and also contribute to mitigating the climate change.*

INTRODUCTION

Soil being a vital natural resource on whose proper use depends the life supporting systems of a country and the socioeconomic development of its people. Soils provide food, fodder, fibre and fuel for meeting the basic human and animal needs. With the ever increasing growth in human and animal population, the demand on soil for more food and fodder production is on the increase. The area available for agriculture is about 51 per cent of the total geographical area and more than 60 per cent of the people are still dependant on agriculture for their livelihood. However, the capacity of a soil to produce is limited and the limits to the production are set by its intrinsic characteristics, agro-climatic setting, and, use and management. There is, therefore, tremendous pressure on land and water resources, which is causing decline in soil-health and stagnation in productivity. The soils have been degrading at an estimated rate of one million hectares per year and ground water levels have been receding at an alarming rate resulting in decline in the ground water resource. Further, land degradation has emerged as a serious problem which has already affected about 38 lakh ha of cultivated area in the State. Soil erosion alone has degraded about 35 lakh ha. Almost all the uncultivated areas are facing various degrees of degradation, particularly soil erosion; salinity and alkalinity has emerged as a major problem in more than 3.5 lakh ha in the irrigated areas of the State. Nutrient depletion and declining factor productivity is common in both rainfed and irrigated areas. The degradation is continuing at an alarming rate and there appears to be no systematic effort among the stakeholders to contain this process. In recent times, an aberration of weather due to climate change phenomenon has added another dimension leading to unpredictable situation to be tackled by the farmers.

In this critical juncture, the challenge before us is not only to increase the productivity per unit area which is steadily declining and showing a fatigue syndrome, but also to prevent or at least reduce the severity of degradation. If the situation is not reversed at the earliest, then the sustainability of the already fragile crop production system and the overall ecosystem will be badly affected in the state. Added to this, every year there is a significant diversion of farm lands and water resources for non-agricultural purposes. Thus, developing strategies to slow down the degradation process or reclaim the soils to normal condition and ensure sustainability of production system are the major issues today. This demands a systematic appraisal of our soil and land resources with respect to their extent, geographic distribution, characteristics, behaviour and use potential, which is very important for developing an effective land use and cropping systems for augmenting agricultural production on a sustainable basis.

The soil and land resource inventories made so far in Karnataka had limited utility because the surveys were of different types, scales and intensities carried out at different times with specific objectives. Hence, there is an urgent need to generate detailed site-

specific farm level database on various land resources for all the villages/watersheds in a time bound manner that would help to protect the valuable soil and land resources and also to stabilize the farm production. Therefore, the land resource inventory required for farm level planning is the one which investigates all the parameters which are critical for productivity *viz.*, soils, site characteristics like slope, erosion, gravelliness and stoniness, climate, water, topography, geology, hydrology, vegetation, crops, land use pattern, animal population, socio-economic conditions, infrastructure, marketing facilities and various schemes and developmental works of the government etc. From the data collected at farm level, the specific problems and potentials of the area can be identified and highlighted, conservation measures required for the area can be planned on a scientific footing, suitability of the area for various uses can be worked out and finally viable and sustainable land use options suitable for each and every land holding can be prescribed.

The Land Resource Inventory is basically done for identifying potential and problem areas, developing sustainable land use plans, estimation of surface run off and water harvesting potential, preparation of soil and water conservation plans, land degradation/desertification etc. The Bureau is presently engaged in developing an LRI methodology using high resolution satellite remote sensing data and Digital Elevation Model (DEM) data to prepare Landscape Ecological Units (LEU) map representing agro-ecosystem as a whole. The LEU is preferred over landform as the base map for LRI. LEU is the assemblage of landform, slope and land use. An attempt has already been made to upscale the soil resource information from 1:250000 and 1:50000 scale to the LEU map in Goa and in some other states.

The land resource inventory aims to provide site-specific database for Yadgir RF1microwatershed in Yadgir Taluk and Yadgir District, Karnataka State for the Karnataka Watershed Development Department. The database was generated by using cadastral map of the village as a base along with high resolution IRS LISS IV and Cartosat-1 merged satellite imagery. Later, an attempt will be made to uplink this LRI data generated at 1:7920 scale under Sujala-III Project to the proposed Landscape Ecological Units (LEUs) map.

The study was organized and executed by the ICAR- National Bureau of Soil Survey and Land Use Planning, Regional Centre, Bangalore under Generation of Land Resource Inventory Data Base Component-1 of the Sujala-III Project funded by the World Bank.

GEOGRAPHICAL SETTING

2.1 Location and Extent

The Yadgir RF1 microwatershed is located in the northern part of Karnataka in Yadgir Taluk & District, Karnataka State (Fig.2.1). It comprises parts of Gajarakota and Yampada villages. It lies between $16^{\circ} 50'$ and $16^{\circ} 52'$ North latitudes and $77^{\circ} 15'$ and $77^{\circ} 17'$ East longitudes covering an area of about 490 ha. It is about 18 km southeast of Yadgir town and is surrounded by Gajarakota on the north and northeast and Yampada village on the southeast, south and west and Hattikuni hobli on the western side.




Fig.2.1 Location map of Yadgir RF1 Microwatershed

2.2 Geology

Major rock formations observed in the microwatershed are granite gneiss (Figs.2.2). They are essentially pink to gray and are coarse to medium grained. They consist primarily of quartz, feldspar, biotite and hornblende. The gray granite gneisses are highly

weathered, fractured and fissured upto a depth of about 10 m. Dolerite dykes and quartz veins are common with variable width and found to occur in Yadgir RF1 microwatershed.



Fig.2.2 Granite and granite gneiss rocks

2.3 Physiography

Physiographically, the area has been identified as granite gneiss landscape based on geology. The area has been further subdivided into five landforms, viz; mounds/ridges, summits, side slopes and very gently sloping uplands, plains and valleys based on slope and its relief features. The elevation ranges from 300-450 m above MSL. The mounds and ridges are mostly covered by rock outcrops.

2.4 Drainage

The area is drained by several parallel streams like Bori, Amerja and Kanga which finally join the river Bhima along its course. Though, they are not perennial, during rainy season they carry large quantities of rain water. The microwatershed has only few small tanks which are not capable of storing the water that flows during the rainy season. Due to this, the ground water recharge is very much affected. This is reflected in the failure of many bore wells in the villages. If the available rain water is properly harnessed by constructing new tanks and recharge structures at appropriate places in the villages, then the drinking and irrigation needs of the area can be easily met. The drainage network is parallel to sub parallel and dendritic.

2.5 Climate

The Yadgir district lies in the northern plains of Karnataka and falls under semiarid tract of the state and is categorized as drought-prone with total annual rainfall of 866 mm (Table 2.1). Of the total rainfall, maximum of 652 mm is received during the south-west monsoon period from June to September, the north-east monsoon from October to early December contributes about 138 mm and the remaining 76 mm during

the rest of the year. The summer season starts during the middle of February and continues up to the first week of June. The period from December to the middle of February is the coldest season. December is the coldest month with mean daily maximum and minimum temperatures being 29.5⁰C and 10⁰C respectively. During peak summer, temperature shoots up to 45⁰C. Relative humidity varies from 26% in summer to 62% in winter. Rainfall distribution is shown in Figure 2.3. The average Potential Evapo-Transpiration (PET) is 141 mm and varies from a low of 81 mm in December to 199 mm in the month of May. The PET is always higher than precipitation in all the months except July, August and September. Generally, the Length of crop Growing Period (LGP) is 120-150 days and starts from 1st week of June to 4th week of October.

Table 2.1 Mean Monthly Rainfall, PET, 1/2 PET at Yadgir Taluk, Yadgir District

Sl. No.	Months	Rainfall	PET	1/2 PET
1	January	4.30	86.0	43.0
2	February	2.30	125.5	62.7
3	March	15.10	166.0	83.0
4	April	18.50	179.8	89.9
5	May	36.0	198.8	97.9
6	June	118.0	175.1	87.5
7	July	171.80	156.3	78.1
8	August	182.9	150.3	75.1
9	September	179.7	142.0	71.0
10	October	105.3	138.5	69.2
11	November	26.4	97.60	48.6
12	December	6.0	80.90	40.4
Total		866.3		




Fig 2.3 Rainfall distribution in Yadgir Taluk, Yadgir District

2.6 Natural Vegetation

The natural vegetation is sparse comprising few tree species, shrubs and herbs. The mounds, ridges and boulders occupy very sizeable area which is under thin to moderately thick forest vegetation. Still, there are some remnants of the past forest cover which can be seen in patches in some ridges and hillocks in the microwatershed (Fig 2.4).

Apart from the continuing deforestation, the presence of large population of goats, sheep and other cattle in the microwatershed is causing vegetative degradation of whatever little vegetation left in the area. The uncontrolled grazing has left no time for the regeneration of the vegetative cover. This leads to the accelerated rate of erosion on the hill slopes resulting in the formation of deep gullies in the foot slopes that eventually result in the heavy siltation of tanks and reservoirs in the microwatershed.



Fig 2.4 Natural vegetation of Ramasamudram-2 microwatershed

2.7 Land Utilization

About 72 per cent area (Table 2.2) in Yadgir district is cultivated at present. An area of about 2 per cent is permanently under pasture, 20 per cent under current fallows and 6 per cent under non-agricultural land, and 5 per cent under currently barren. Forests occupy an area of about 7 per cent and the tree cover is in a very poor state. Most of the mounds, ridges and bouldery areas have very poor vegetative cover. Major crops grown in the area are sorghum, maize, cotton, sunflower, groundnut, red gram and paddy. The cropping intensity is 120 per cent in the taluk. While carrying out land resource inventory, the land use/land cover particulars are collected from all the survey numbers and a current land use map of the microwatershed is prepared. The current land use map prepared shows the arable and non-arable lands, other land uses and different types of crops grown in the area. The current land use map of Yadgir RF1 microwatershed is presented in

Fig.2.5. The different crops and cropping systems adopted in the microwatershed is presented in the Figures 2.6. The location of wells and conservation structures in the Yadgir RF1 microwatershed is given in Fig.2.7.

Table 2.2 Land Utilization in Yadgir District

Sl. No.	Agricultural land use	Area (ha)	Per cent
1.	Total geographical area	516088	-
2.	Total cultivated area	373617	72.4
3.	Area sown more than once	74081	14.3
4.	Cropping intensity	-	119.8
5.	Trees and grooves	737	0.14
6.	Forest	33773	6.54
7.	Cultivable wasteland	2385	0.46
8.	Permanent Pasture land	11755	2.28
9.	Barren land	27954	5.41
10.	Non- Agriculture land	29623	5.73
11.	Current Fallows	105212	20.4




Fig.2.5 Current Land Use map of Yadgir RF1Microwatershed




Fig 2.6 Different Crops and Cropping Systems in Yadgir RF1 Microwatershed




Fig 2.7 Location of wells and conservation structures in Yadgir RF1 Microwatershed

SURVEY METHODOLOGY

The purpose of land resource inventory is to delineate similar areas (soil series and phases), which respond or expected to respond similarly for a given level of management. This was achieved in Yadgir RF1 microwatershed by the detailed study of all the soil characteristics (depth, texture, colour, structure, consistence, coarse fragments, porosity, soil reaction, soil horizons etc.) and site characteristics (slope of the land, erosion, drainage, occurrence of rock fragments etc.) followed by grouping of similar areas based on soil-site characteristics into homogeneous (management units) units, and showing the area extent and their geographic distribution on the microwatershed cadastral map. The detailed survey at 1:7920 scale was carried out in an area of 490 ha. The methodology followed for carrying out land resource inventory was as per the guidelines given in Soil Survey Manual (IARI, 1971; Soil Survey Staff, 2006; Natarajan *et al.*, 2015) which is briefly described below.

3.1 Base Maps

The detailed survey of the land resources occurring in the microwatershed was carried out by using digitized cadastral map and satellite imagery as base supplied by KSRSAC. The cadastral map shows field boundaries with their survey numbers, location of tanks, streams and other permanent features of the area (Fig. 3.1). Apart from the cadastral map, remote sensing data products from Cartosat-1 and LISS IV merged at the scale of 1:7920 were used in conjunction with the cadastral map to identify the landscapes, landforms and other surface features. The imagery helped in the identification and delineation of boundaries between hills, uplands and lowlands, water bodies, forest and vegetated areas, roads, habitations and other cultural features of the area (Fig. 3.2). The cadastral map was overlaid on the satellite imagery (Fig. 3.3) that helps to identify the parcel boundaries and other permanent features. Apart from cadastral maps and images, toposheets of the area (1:50,000 scale) were also used for initial traversing, identification of geology and landforms, drainage features, present land use and also for selection of transects in the microwatershed.

3.2 Image Interpretation for Physiography

False Colour Composites (FCCs) of Cartosat-I and LISS-IV merged satellite data covering microwatershed area was visually interpreted using image interpretation elements and all the available collateral data with local knowledge. The delineated physiographic boundaries were transferred on to a cadastral map overlaid on satellite imagery. Physiographically, the area has been identified as granite gneiss landscape. It was divided into five landforms, *viz.*; ridges and mounds, gently and very gently sloping uplands and lowlands based on slope and image characteristics. They were further

subdivided into physiographic/image interpretation units based on image characteristics. The image interpretation legend for physiography is given below.

Image Interpretation Legend for Physiography

G- Granite Gneiss Landscape

G1	Hills/ Ridges/ Mounds
G11	Summits
G12	Side slopes
G121	Side slopes with dark grey tones
G2	Uplands
G21	Summits
G22	Gently sloping uplands
G221	Gently sloping uplands, yellowish green (eroded)
G222	Gently sloping uplands, yellowish white (severely eroded)
G23	Very gently sloping uplands
G231	Very gently sloping uplands, yellowish green
G232	Very gently sloping uplands, medium green and pink
G233	Very gently sloping uplands, pink and green (scrub land)
G234	Very gently sloping uplands, medium greenish grey
G235	Very gently sloping uplands, yellowish white (eroded)
G236	Very gently sloping uplands, dark green
G237	Very gently sloping uplands, medium pink (coconut garden)
G238	Very gently sloping uplands, pink and bluish white (eroded)
G24	Valleys/ lowlands
G241	Valleys, pink tones
G242	Valleys gray mixed with pink tones




Fig 3.1 Scanned and Digitized Cadastral map of Yadgir RF1 Microwatershed




Fig.3.2 Satellite Image of Yadgir RF1 Microwatershed




Fig.3.3 Cadastral map overlaid on IRS PAN+LISS IV merged imagery of Yadgir RF1Microwatershed

3.3 Field Investigation

The field boundaries and survey numbers given on the cadastral sheet were located on the ground by following permanent features like roads, cart tracks, *nallas*, streams, tanks etc., and wherever changes were noticed, they were incorporated on the microwatershed cadastral map. Preliminary traverse of the microwatershed was carried out with the help of cadastral map, imagery and toposheets. While traversing, landforms and physiographic units identified were checked and preliminary soil legend was prepared by studying soils at few selected places. Then, intensive traversing of each physiographic unit like hills, ridges, uplands and valleys was carried out. Based on the variability observed on the surface, transects (Fig. 3.4) were selected across the slope covering all the landform units in the microwatershed (Natarajan and Dipak Sarkar, 2010).




Fig: 3.4. Location of profiles in a transect

In the selected transect, soil profiles were located (Fig. 3.4) at closely spaced intervals to take care of any change in the land features like break in slope, erosion, gravel, stones etc. In the selected sites, soil profiles (vertical cut showing the soil layers from surface to the rock) were opened upto 200 cm or to the depth limited by rock or hard substratum and studied in detail for all their morphological and physical characteristics. The soil and site characteristics were recorded for all profile sites on a standard proforma as per the guidelines given in USDA Soil Survey Manual (Soil Survey Staff, 2012). Apart from the transect study, profiles were also studied at random, almost like in a grid pattern, outside the transect areas.

Based on the soil characteristics, the soils were grouped into different soil series. Soil series is the most homogeneous unit having similar horizons and properties and behaves similarly for a given level of management. Soil depth, texture, colour, kind of horizon and horizon sequence, calcareousness, amount and nature of gravel present, nature of substratum etc, were used as the major differentiating characteristics for identifying soil series occurring in the area. The differentiating characteristics used for identifying the soil series are given in Table 3.1. Based on the above characteristics, 6 soil series were identified in the Yadgir RF1 microwatershed.

Table 3.1 Differentiating Characteristics used for identifying Soil Series
(Characteristics are of Series Control Section)

Soils of Granite gneiss Landscape							
Sl. no	Soil Series	Depth (cm)	Colour (moist)	Texture	Gravel (%)	Horizon sequence	Calcareousness
1	BDP (Baddeppalli)	<25	7.5YR 3/2,3/4 5YR 3/4	scl	-	Ap-Ac	es
2	HTK (Hattikuni)	25-50	10YR4/6,4/4 7.5YR34/4,3/3	sl	10-25	Ap-Ac	-
3	KKR (Kakalwar)	<25	7.5 YR 4/3, 10 YR 6/3	sl	10-25	Ap-Ac	-
4	BDL (Badiyala)	25-50	7.5 YR 2.5/3,2.5/2,3/3 10YR 3/4,4/3	sl	<15	Ap-Bw	e
5	SBR (Sambra)	50-75	10YR 7/1 7.5YR 7/4	ls-s	-	Ap-AC	-
6	GWD (Gowdgera)	75-100	10 YR 3/1,3/2.4/2	scl	-	Ap-Bw	es

3.4 Soil Mapping

The area under each soil series was further separated into soil phases and their boundaries delineated on the cadastral map based on the variations observed in the texture of the surface soil, slope, erosion, presence of gravel, stoniness etc. A soil phase is a subdivision of soil series based mostly on surface features that affect its use and management. The soil mapping units are shown on the map (Fig.3.5) in the form of symbols. During the survey many soil profile pits, few minipits and a few auger bores representing different landforms occurring in the microwatershed were studied. In addition to the profile study, spot observations in the form of minipits, road cuts, terrace cuts etc., were studied to validate the soil boundaries on the soil map. The soil map shows the geographic distribution of 12 mapping units representing 6 soil series occurring in the microwatershed. The soil map unit (soil legend) description is presented in Table 3.2. The soil phase map (management units) shows the distribution of 12 soil phases mapped in the microwatershed. Each mapping unit (soil phase) delineated on the map has similar soil and site characteristics. In other words, all the farms or survey numbers included in one soil phase will have similar management needs and have to be treated accordingly.

3.5 Land Management Units (LMU's)

The 12 soil phases identified and mapped in the microwatershed were grouped into 3 Land Management Units (LMU's) for the purpose of preparing a Proposed Crop Plan for sustained development of the microwatershed. The database (soil phases) generated under LRI was utilized for identifying Land Management Units (LMU's) based on the management needs. One or more than one soil site characteristic having influence on the management have been chosen for identification and delineation of LMUs. For Yadgir RF1 microwatershed, five soil and site characteristics, namely soil depth, soil

texture, slope, erosion and gravel content have been considered for defining LMUs. The Land Management Units are expected to behave similarly for a given level of management.

3.6 Laboratory Characterization

Soil samples for each soil series were collected from representative master profiles for laboratory characterization by following the methods outlined in the Laboratory Manual (Sarma *et al*, 1987). Surface soil samples collected from farmer's fields (46 samples) for fertility status (major and micronutrients) at 320 m grid interval in the year 2017 were analyzed in the laboratory (Katyal and Rattan, 2003). By linking the soil fertility data to the survey numbers through GIS, soil fertility maps were generated by using Kriging method for the microwatershed.

Table 3.2 Soil map unit description of Yadgir RF1Microwatershed

*Soil map unit No.	Soil Series	Soil Phase	Mapping Unit Description	Area in ha (%)
Soils of Granite and Granite Gneiss Landscape				
	BDP		Baddeppalli soils are very shallow (<25 cm), well drained, have dark brown to dark reddish brown, calcareous sandy clay loam soils occurring on very gently sloping uplands under cultivation	20(4.01)
119		BDPiB3	Sandy clay surface, slope 1-3%, severe erosion	17 (3.46)
120		BDPhB2	Sandy clay loam surface, slope 1-3%, moderate erosion	3 (0.55)
	HTK		Hattikuni soils are shallow (25-50 cm), well drained, have dark yellowish brown sandy loam soils occurring on very gently sloping uplands under cultivation	177(36.1)
113		HTKcC2g1	Sandy loam surface, slope 3-5%, moderate erosion, gravelly (15-35%)	128 (26.09)
161		HTKbB2g1	Loam sandy surface, slope 1-3%, moderate erosion, gravelly (15-35%)	49 (10.01)
	KKR		Kakalawar soils are very shallow (<25 cm), well drained, have dark brown sandy loam soils occurring on very gently sloping uplands under cultivation	139(28.39)
153		KKRbB2g1	Loam sandy surface, slope 1-3%, moderate erosion, gravelly (15-35%)	137 (28.03)
175		KKRcB2	Sandy loam surface, slope 1-5%, moderate erosion	2 (0.36)
	BDL		Badiyala soils are shallow (25-50 cm), well drained, have dark brown to very dark brown and dark yellowish brown, slightly calcareous sandy loam soils occurring on very gently to gently sloping uplands under cultivation	27(5.42)
2		BDLbB2	Loam sandy surface, slope 1-3%,	16 (3.28)

			moderate erosion	
4		BDLhB2	Sandy clay loam surface, slope 1-3%, moderate erosion	11 (2.14)
	SBR	Sambara soils are moderately shallow (50-75 cm), somewhat excessively drained, have light gray to pink, loamy sand soils occurring on very gently to gently sloping uplands under cultivation		72(14.85)
11		SBRcB2	Sandy loam surface, slope 1-3%, moderate erosion	24 (4.96)
12		SBRcC3g1	Sandy loam surface, slope 3-5%, severe erosion, gravelly (15-35%)	48 (9.89)
	GWD	Gowdagera soils are moderately deep (75-100 cm), moderately well drained, have dark grayish brown to very dark grayish brown, sodic calcareous sandy clay loam soils occurring on very gently sloping uplands under cultivation		49(10.13)
34		GWDcB2	Sandy loam surface, slope 1-3%, moderate erosion	20 (4.12)
127		GWDmB2	Clay surface, slope 1-3%, moderate erosion	29 (6.01)
999		Rock lands	both massive and bouldery with little or no soil	0.25 (0.05)
1000		Others	Habitation and Water body	5 (1.04)




Fig 3.5 Soil Phase or Management Units - Yadgir RF1Microwatershed

THE SOILS

Detailed information pertaining to the nature, extent and their distribution of different kinds of soils occurring in Yadgir RF1 microwatershed is provided in this chapter. The microwatershed area has been identified as granite gneiss landscape based on geology. In all, 6 soil series are identified. Soil formation is the result of the combined effect of environmental and terrain factors that are reflected in soil morphology. In the granite gneiss landscape, it is by parent material, relief and climate.

A brief description of each of the 6 soil series identified followed by 12 soil phases (management units) mapped are furnished below. The physical and chemical characteristics of soil series identified in Yadgir RF1 microwatershed are given in Table 4.1 along with soil classification. The soils in any one map unit differ from place to place in their depth, texture, slope, gravelliness, erosion or any other site characteristic that affect management. The soil phase map can be used for identifying the suitability of areas for growing specific crops or for other alternative uses and also for deciding the type of conservation structures needed. The detailed information on soil and site-characteristics like soil depth, surface soil texture, slope, erosion, gravelliness, AWC, LCC etc, with respect to each of the soil phase identified is given village/survey number wise for the microwatershed in Appendix-I.

4.1 Soils of granite gneiss landscape

In this landscape, 6 soil series are identified and mapped. Of these, HTK series occupies a maximum area of 177 ha (36%) followed by KKR 139 ha (28%), SBR 72 ha (15%), GWD 49 ha (10%), BDL 27 ha (5%) and BDP 20 ha (4%). Brief description of each series identified and number of soil phases mapped is given below.

4.1.1 Baddeppalli (BDP) Series: Baddeppalli soils are very shallow (<25cm), well drained, have dark brown to dark reddish brown, calcareous sandy clay loam soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation. The Baddeppalli series has been classified as a member of the loamy, mixed, (calcareous) isohyperthermic family of Lithic Ustorthents.

The thickness of the soil is less than 25 cm. Its colour is in 7.5 YR and 5 YR hue with value 3 and chroma 2 to 4. The texture varies from sandy clay loam to sandy clay and is calcareous. The available water capacity is very low (<50 mm/m). Two phases were identified and mapped.



Landscape and Soil Profile characteristics of Baddeppalli (BDP) Series

4.1.2 Hattikuni (HTK) Series: Hattikuni soils are shallow (25-50 cm), well drained, have dark brown to dark yellowish brown sandy loam soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation. The Hattikuni series has been classified as a member of the mixed, isohyperthermic family of Lithic Ustipsamments.

The thickness of the soil ranges from 36 to 50 cm. The thickness of A horizon ranges from 8 to 12 cm. Its colour is in 10YR and 7.5 YR hue with value 3 to 4 and chroma 4 to 6. The texture varies from loamy sand to sandy loam. The thickness of subsurface horizon ranges from 28 to 42 cm. Its colour is in 10 YR and 7.5 YR hue with value 3 to 4 and chroma 4 to 6. Its texture varies from loamy sand to sand and sandy loam. The available water capacity is very low (<50 mm/m). Two phases were identified and mapped.



Landscape and Soil Profile characteristics of Hattikuni (HTK) Series

4.1.3 Kakalawar (KKR) Series: Kakalawar soils are very shallow (<25cm), well drained, have dark brown to light brown, sandy loam soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation. The Kakalawar series has been classified as a member of the mixed, isohyperthermic family of Lithic Ustipsamments.

The thickness of the soil is less than 25 cm. Its colour is in 10 YR and 7.5 YR hue with value 4 to 6 and chroma 3 to 4. The texture varies from loamy sand to sand. The available water capacity is very low (<50 mm/m). Two phases were identified and mapped.



Landscape and Soil Profile characteristics of Kakalawar (KKR) Series

4.1.4 Badiyala (BDL) Series: Badiyala soils are shallow (25-50 cm), well drained, have very dark brown, dark yellow brown and dark brown, slightly calcareous sandy loam soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation. The Badiyala series has been classified as a member of the coarse-loamy, mixed, isohyperthermic family of Fluventic Haplustepts.

The thickness of the solum ranges from 28 to 50 cm. The thickness of A horizon ranges from 4 to 12 cm. Its colour is in 10YR hue with value 3 to 4 and chroma 3 to 4. The texture is loamy sand, sandy clay loam and sandy clay. The thickness of B horizon ranges from 27 to 45 cm. Its colour is in 10 YR and 7.5 YR hue with value 2 to 4 and chroma 3 to 4. Its texture is sandy loam to sandy clay loam and is slightly calcareous. The available water capacity is very low (<50mm/m). Two phases were identified and mapped.



Landscape and Soil Profile characteristics of Badiyala (BDL) Series

4.1.5 Sambara (SBR) Series: Sambara soils are moderately shallow (50-75 cm), somewhat excessively drained, have light grey to reddish yellow, loamy sand soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation. The Sambara series has been classified as a member of the mixed, isohyperthermic family of Typic Ustipsamments.

The thickness of the soil ranges from 52-75 cm. Thickness of A horizon ranges from 8 to 23 cm. Its colour is in hue 10 YR and 7.5 YR with value 3 and chroma 1 to 4. The texture varies from loamy sand to sandy loam. The thickness of subsurface horizons ranges from 41 to 66 cm. Its colour is in 10 YR and 7.5 YR hue with value 3 to 5 and chroma 1 to 4. The texture is loamy sand. The available water capacity is very low (<50 mm/m). Two phases were identified and mapped.



Landscape and Soil Profile characteristics of Sambara (SBR) Series

4.1.6 Gowdagera (GWD) Series: Gowdagera soils are moderately deep (75-100 cm), moderately well drained, very dark gray to dark grayish brown, calcareous sodic sandy clay loam soils. They are developed from weathered granite gneiss and occur on very gently sloping uplands under cultivation. The Gowdagera series has been classified as a member of the fine-loamy, mixed, calcareous, isohyperthermic family of Typic Haplustepts.

The thickness of the solum ranges from 76 to 100 cm. The thickness of A horizon ranges from 8 to 16 cm. Its colour is in hue 10 YR with value 3 to 4 and chroma 2 to 4. Its texture varies from sandy loam to sandy clay loam. The thickness of B horizon ranges from 61 to 91 cm. Its colour is in hue 10 YR with value 2 to 4 and chroma 1 to 4. Its texture is sandy clay loam to sandy clay and is calcareous sodic soils. The available water capacity is medium (101-150 mm/m). Two phases were identified and mapped.



Landscape and Soil Profile characteristics of Gowdagera (GWD) Series

Table: 4.1 Physical and Chemical Characteristics of Soil Series identified in Yadgir RF1microwatershed

Soil Series: Baddeppalli (BDP) **Pedon:** R-11

Location: 16°43'84.4"N 77°14'06.4"E, Halagera village, Yadgir hobli, Yadgir taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bengaluru **Classification:** Loamy, mixed (calcareous), isohyperthermic, Lithic Ustorthents

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-16	Ap	58.67	17.02	24.31	19.03	13.74	9.62	10.57	5.71	<15	scl	16.19	8.18

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO ₃	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP
	Water	CaCl ₂	M KCl				Ca	Mg	K	Na	Total				
				dS m ⁻¹	%	%	cmol kg ⁻¹					%	%		
0-16	8.58	-	-	0.262	1.60	7.67	-	-	0.24	0.06	-	18.10	0.74	100	0.35

Contd...

Soil Series: Hattikuni (HTK), Pedon: R-7

Location: 16°50'46.5"N 77°10'16.4"E, Yaddalli village, Hattikuni hobli, Yadgir taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bengaluru **Classification:** Mixed, isohyperthermic, Lithic Ustipsamments

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-12	Ap	90.89	5.62	3.49	8.50	13.46	29.86	29.55	9.51	20	s	7.73	3.16
12-22	A1	89.97	6.53	3.50	7.19	13.48	29.48	29.79	10.03	20	s	8.00	3.05
22-45	A2	87.20	6.43	6.38	11.09	14.42	31.55	7.16	22.98	40	ls	7.67	3.96

Depth (cm)	pH (1:2.5)			E.C. (1:2.5) dS m ⁻¹	O.C. %	CaCO ₃ %	Exchangeable bases					CEC	CEC/Clay	Base saturation %	ESP %
	Water	CaCl ₂	M KCl				Ca	Mg	K	Na	Total				
0-12	6.81	-	-	0.062	0.07	-	2.35	0.50	0.16	0.01	3.02	3.0	0.86	100	0.38
12.0-22	6.80	-	-	0.050	0.21	-	1.67	0.30	0.09	0.01	2.07	2.4	0.69	86.30	0.45
22-45	6.85	-	-	0.044	0.19	-	1.82	0.42	0.10	0.06	2.40	2.6	0.41	92.41	2.17

Contd...

Soil Series: Kakalawar (KKR), Pedon: R-7

Location: 16°50'25.9"N 77°15'97.1"E, Yampada village, Gurumitkal hobli, Yadgir taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bengaluru **Classification:** Mixed, isohyperthermic, Lithic Ustipsamments

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-22	Ap	83.81	10.37	5.82	17.31	20.65	17.91	5.67	22.27	10-20	ls	9.77	4.65

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO ₃	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP
	Water	CaCl ₂	M KCl				Ca	Mg	K	Na	Total				
				dS m ⁻¹	%	%	cmol kg ⁻¹					%	%		
0-22	5.85	-	-	0.027	0.19	-	0.72	0.21	0.62	0.03	1.58	2.6	0.45	60.90	1.17

Contd...

Soil Series: Badiyala (BDL) **Pedon:** R-5

Location: 16°37'10.0"N 77°20'21.5", Gudalagunta village, Balichakra hobli, Yadgir taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bengaluru **Classification:** Coarse-loamy, mixed, isohyperthermic Fluventic Haplustepts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-12	Ap	87.13	7.04	5.83	10.03	24.32	23.61	23.51	5.67	<15	ls	6.27	2.44
12-28	Bw1	64.63	13.30	22.07	6.74	13.07	22.30	17.01	5.50	<15	scl	16.34	7.83
28-52	BC	73.11	12.02	14.87	3.93	16.03	26.89	18.41	7.86	<15	sl	12.94	5.47

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO ₃	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP
	Water	CaCl ₂	M KCl	dS m ⁻¹	%	%	Ca	Mg	K	Na	Total				
							cmol kg ⁻¹								
0-12	6.20	-	-	0.074	1.00	0.00	2.80	0.98	0.14	0.01	3.92	4.20	0.72	93	0.20
12-28	9.04	-	-	0.253	0.80	3.20	-	-	0.16	0.69	-	16.90	0.77	100	4.09
28-52	9.41	-	-	0.364	1.10	3.60	-	-	0.16	1.39	-	11.10	0.75	100	12.52

Contd...

Soil Series: Sambara (SBR) **Pedon:** R-10

Location: 16°42'04.5"N 77°14'35.3"E, Jinatera village, Balichakra hobli, Yadgir taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bengaluru **Classification:** Mixed, isohyperthermic Typic Ustipsamments

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-9	Ap	81.90	8.22	9.88	23.76	14.05	23.76	10.62	9.71	-	ls	9.45	2.69
9-17	C1	84.08	6.59	9.33	21.30	20.69	17.65	17.65	6.80	-	ls	7.84	2.65
17-60	C2	86.86	6.17	6.98	11.53	21.54	25.08	23.46	5.26	-	ls	5.48	2.62
60-78	C3	87.27	6.92	5.81	15.05	20.91	26.36	19.29	5.66	-	ls	5.19	2.81

Depth (cm)	pH (1:2.5)			E.C. (1:2.5) dS m ⁻¹	O.C. %	CaCO ₃ %	Exchangeable bases					CEC	CEC/Clay	Base saturation %	ESP %
	Water	CaCl ₂	M KCl				Ca	Mg	K	Na	Total				
0-9	8.24	-	-	0.145	0.61	0.91	-	-	0.12	0.09	-	7.50	0.76	100	1.15
9-17	8.21	-	-	0.068	0.57	0.39	-	-	0.06	0.12	-	6.70	0.72	100	1.82
17-60	8.47	-	-	0.080	0.38	0.48	-	-	0.03	0.17	-	2.70	0.39	100	6.34
60-78	8.50	-	-	0.081	0.30	0.52	-	-	0.03	0.17	-	2.70	0.46	100	6.43

Contd...

Soil Series: Gowdagera (GWD) **Pedon:** R-13

Location: 16°38'24.4"N 77°21'24.0"E, Madhawara village, Balichakara hobli, Yadgir taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bengaluru **Classification:** Fine-loamy, mixed (calcareous) isohyperthermic Typic Haplustepts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-18	Ap	79.61	13.94	6.45	14.17	17.53	23.65	17.02	7.24	-	ls	11.36	3.86
18-42	BW1	69.09	10.58	21.06	10.54	16.58	22.01	14.43	5.53	-	scl	31.62	12.30
42-81	Bw2	51.37	13.51	35.60	7.59	10.55	16.24	11.60	5.38	-	sc	67.57	26.89

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO ₃	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP
	Water	CaCl ₂	M KCl	dS m ⁻¹	%	%	Ca	Mg	K	Na	Total				
							cmol kg ⁻¹								
0-18	9.89	-	-	0.74	0.66	1.20	-	-	0.18	3.63	-	8.35	1.29	100	17.40
18-42	10.82	-	-	1.60	0.27	5.76	-	-	0.19	19.23	-	15.84	0.75	100	40.17
42-81	10.83	-	-	2.30	0.27	7.80	-	-	0.40	26.71	-	26.54	0.75	100	40.27

INTERPRETATION FOR LAND RESOURCE MANAGEMENT

The most important soil and site characteristics that affect the land use and conservation needs of an area are land capability, land irrigability, soil depth, soil texture, coarse fragments, available water capacity, soil slope, soil erosion, soil reaction etc. These are interpreted from the data base generated through land resource inventory and several thematic maps are generated. These would help in identifying the areas suitable for growing crops and, soil and water conservation measures and structures needed thus helping to maintain good soil health for sustained crop production. The various interpretative and thematic maps generated are described below.

5.1 Land Capability Classification

Land capability classification is an interpretative grouping of soil map units (soil phases) mainly based on inherent soil characteristics, external land features and environmental factors that limit the use of land for agriculture, pasture, forestry, or other uses on a sustained basis (IARI, 1971). The land and soil characteristics used to group the land resources in an area into various land capability classes, subclasses and units are

Soil Characteristics: Depth, texture, gravelliness, calcareousness.

Land characteristics: Slope, erosion, drainage, rock outcrops.

Climate: Total rainfall and its distribution, and length of crop growing period.

The land capability classification system is divided into land capability classes, subclasses and units based on the level of information available. Eight land capability classes are recognized. They are

Class I: They are very good lands that have no limitations or very few limitations that restrict their use.

Class II: They are good lands that have minor limitations and require moderate conservation practices.

Class III: They are moderately good lands that have moderate limitations that reduce the choice of crops or that require special conservation practices.

Class IV: They are fairly good lands that have very severe limitations that reduce the choice of crops or that require very careful management.

Class V: Soils in these lands are not likely to erode, but have other limitations like wetness that are impractical to remove and as such not suitable for agriculture, but suitable for pasture or forestry with minor limitations.

Class VI: The lands have severe limitations that make them generally unsuitable for cultivation, but suitable for pasture or forestry with moderate limitations.

Class VII: The lands have very severe limitations that make them unsuitable for cultivation, but suitable for pasture or forestry with major limitations.

Class VIII: Soil and other miscellaneous areas (rock lands) that have very severe limitations that nearly preclude their use for any crop production, but suitable for wildlife, recreation and installation of wind mills.

The land capability subclasses are recognised based on the dominant limitations observed within a given land capability class. The subclasses are designated by adding a lower case letter like 'e', 'w', 's', or 'c' to the class numeral. The subclass "e" indicates that the main hazard is risk of erosion, "w" indicates drainage or wetness as a limitation for plant growth, "s" indicates shallow soil depth, coarse or heavy textures, calcareousness, salinity/alkalinity or gravelliness and "c" indicates limitation due to climate.

The land capability subclasses have been further subdivided into land capability units based on the kinds of limitations present in each subclass. Ten land capability units are used in grouping the soil map units. They are stony or rocky (0), erosion hazard (slope, erosion) (1), coarse texture (sand, loamy sand, sandy loam) (2), fine texture (cracking clay, silty clay) (3), slowly permeable subsoil (4), coarse underlying material (5), salinity/alkali (6), stagnation, overflow, high ground water table (7), soil depth (8) and fertility problems (9). The capability units thus identified have similar soil and land characteristics that respond similarly to a given level of management. The soils of the microwatershed have been classified upto land capability subclass level.

The 12 soil map units identified in Yadgir RF1 microwatershed are grouped under 3 land capability classes and 4 land capability subclasses. An entire area of 410 ha (68%) in the microwatershed is suitable for agriculture. About <1 ha (<1%) area is having rock outcrops and about 5 ha (1%) is covered by others (water body & habitation) (Fig. 5.1).

Good cultivable lands (Class II) cover an area of about 10 per cent and are distributed in the southern, southeastern and southwestern part of the microwatershed with minor problems of soil and erosion. Moderately good cultivable lands (Class III) cover an area of about 42 per cent and are distributed in the central, northeastern, southeastern and southern part of the microwatershed with moderate problems of soil and erosion. Fairly good cultivable lands (Class IV) occur in 47 per cent area of the microwatershed and they have severe problem of soil and erosion and distributed in the major part of the microwatershed.




Fig. 5.1 Land Capability map of Yadgir RF1 Microwatershed

5.2 Soil Depth

Soil depth refers to the depth of the soil occurring above the parent material or hard rock. The depth of the soil determines the effective rooting depth for plants and in accordance with soil texture, mineralogy and gravel content, the capacity of the soil column to hold water and nutrient availability. Soil depth is one of the most important soil characteristic that is used in differentiating soils into different soil series. The soil depth classes used in identifying soils in the field are very shallow (<25 cm), shallow (25-50 cm), moderately shallow (50-75 cm), moderately deep (75-100 cm), deep (100-150 cm) and very deep (>150 cm). They were used to classify the soils into different depth classes and a soil depth map was generated. The area extent and their geographical distribution in the microwatershed is given in Fig. 5.2.

Very shallow (<25 cm) soils occur in an area of 159 ha (32%) and are distributed in the southwestern, southeastern, northern, northeastern and northwestern part of the microwatershed. Shallow (25-50 cm) soils occur in a small area of 204 ha (42%) and are distributed in the major part of the microwatershed. Moderately shallow (50-75 cm) soils occur in an area of 73 ha (15%) and are distributed in the southern, eastern and central part of the microwatershed. Moderately deep (75-100 cm) soils occur in an area of 50 ha (10%) and are distributed in the northern and southeastern part of the microwatershed.

The problem soils occupy only 363 ha (74%) area where only short duration crops can be grown occasionally and the probability of crop failure is very high.




Fig. 5.2 Soil Depth map of Yadgir RF1 Microwatershed

5.3 Surface Soil Texture

Texture is an expression to indicate the coarseness or fineness of the soil as determined by the relative proportion of primary particles of sand, silt and clay. It has a direct bearing on the structure, porosity, adhesion and consistence. The surface layer of a soil to a depth of about 25 cm is the layer that is most used by crops and plants. The surface soil textural class provides a guide to understanding soil-water retention and availability, nutrient holding capacity, infiltration, workability, drainage, physical and chemical behaviour, microbial activity and crop suitability. The textural classes used for LRI were used to classify and a surface soil texture map was generated. The area extent and their geographical distribution in the microwatershed is shown in Figure 5.3.

An area of about 203 ha (41%) of the microwatershed has sandy soils at the surface and are distributed in all parts except central and southern part. Maximum area of 236 ha (48%) of the microwatershed has soils that are loamy and are distributed in the major part. An area of 46 ha (9%) of the microwatershed has soils that are clayey and are distributed in the northeastern, eastern and southern part. Both soils have high potential for soil-water retention and availability, and nutrient retention and availability, but clay soils have more problems of drainage, infiltration, workability and other physical problems. Problem soils cover 203 ha that are sandy: here only tuber crops can be grown which requires frequent irrigation and addition of manures and fertilizers.




Fig. 5.3 Surface Soil Texture map of Yadgir RF1 Microwatershed

5.4 Soil Gravelliness

Gravel is the term used for describing coarse fragments between 2 mm and 7.5 cm diameter and stones for those between 7.5 cm and 25 cm. The presence of gravel and stones in soil reduces the volume of soil responsible for moisture and nutrient storage, drainage, infiltration and runoff, and hinders plant growth by impeding root growth and seedling emergence, intercultural operations and farm mechanization. The gravelliness classes used in LRI were used to classify the soils and using these classes a gravelliness map was generated. The area extent and their geographic distribution in the microwatershed is shown in Figure 5.4.

Non gravelly (<15%) soils cover maximum area of 122 ha (25%) and distributed in the southeastern, northern and northeastern and eastern part of the microwatershed. These are the most productive soils, where all climatically adapted short and long duration crops can be grown. Gravelly (15-35%) soils occur in an area of 363 ha (74%) and distributed in the major part of the microwatershed; these lands are low in moisture holding capacity and hence growing of short duration crops is ideal with best management practice.




Fig. 5.4 Soil Gravelliness map of Yadgir RF1 Microwatershed

5.5 Available Water Capacity

The soil available water capacity (AWC) is estimated based on the ability of the soil column to retain water between the tensions of 0.33 and 15 bar in a depth of 100 cm or the entire solum if the soil is shallower. The AWC of the soils (soil series) as estimated by considering the soil texture, mineralogy, soil depth and gravel content (Sehgal *et al.*, 1990) and accordingly the soil map units were grouped into five AWC classes *viz*, very low (<50 mm/m), low (50-100 mm/m), medium (100-150 mm/m), high (150-200 mm/m) and very high (>200 mm/m) and using these values, an AWC map was generated. The area extent and their geographic distribution of different AWC classes in the microwatershed is given in Figure 5.5.

Maximum area of about 435 ha (89%) in the microwatershed has soils that are very low (<50 mm/m) in available water capacity and is distributed in the major part of the microwatershed. An area of about 50 ha (10%) is medium (51-100 mm/m) in available water capacity and are distributed in the northern, southern and southeastern part of the microwatershed.

An area of about 435 ha (89%) in the microwatershed has soils that are problematic with regard to available water capacity. Here, only short duration crops can be grown and probability of the crop failure is very high. These areas are best put to other alternative uses.




Fig. 5.5 Soil Available Water Capacity map of Yadgir RF1 Microwatershed

5.6 Soil Slope

Soil slope refers to the inclination of the surface of the land. It is defined by gradient, shape and length, and is an integral feature of any soil as a natural body. Slope is considered important in soil genesis, land use and land development. The length and gradient of slope influences the rate of runoff, infiltration, erosion and deposition. The soil map units were grouped into single slope class and a slope map was generated showing the area extent and their geographic distribution in the microwatershed (Fig. 5.6).

An area of about 308 ha (63%) of the microwatershed falls under very gently sloping (1-3% slope) lands, thus these areas have high potential in respect of soil slopes. In these areas, all climatically adapted annual and perennial crops can be grown without much soil and water conservation and other land development measures. Gently sloping (3-5%) lands occur in 176 ha (36%) and are distributed in the central and southern part of the microwatershed. In these areas the soil and water conservation measures should be adopted in order to increase the productivity of soils.




Fig. 5.6 Soil Slope map of Yadgir RF1 Microwatershed

5.7 Soil Erosion

Soil erosion refers to the wearing away of the earth's surface by the forces of water, wind and ice involving detachment and transport of soil by raindrop impact. It is used for accelerated soil erosion resulting from disturbance of the natural landscape by burning, excessive grazing and indiscriminate felling of forest trees and tillage, all usually by man. The erosion classes showing an estimate of the current erosion status as judged from field observations in the form of rills, gullies or a carpet of gravel on the surface are recorded. Four erosion classes, viz, slight erosion (e1), moderate erosion (e2), severe erosion (e3) and very severe erosion (e4) are recognized. The soil map units were grouped into different erosion classes and a soil erosion map generated. The area extent and their spatial distribution in the microwatershed is given in Figure 5.7.

Soils that are moderately eroded (e2 class) cover a maximum area of 419 ha (86%) and are distributed in the major part of the microwatershed. Severely eroded soils (e3 class) cover an area of 65 ha (13%) and are distributed in the eastern, western and southern part of the microwatershed.

Entire area of the microwatershed is problematic because of moderate and severe erosion. For these areas, taking up of soil and water conservation and other land development measures are needed.




Fig. 5.7 Soil Erosion map of Yadgir RF1 Microwatershed

FERTILITY STATUS

Soil fertility plays an important role in increasing crop yield. The adoption of high yielding varieties that require high amounts of nutrients has resulted in deficiency symptoms in crops and plants due to imbalanced fertilization and poor inherent fertility status as these areas are characterised by low rainfall and high temperatures. Hence, it is necessary to know the fertility (macro and micro nutrients) status of the soils of the watersheds for assessing the kind and amount of fertilizers required for each of the crop intended to be grown. For this purpose, the surface soil samples collected from the grid points (one soil sample at every 320 m interval) all over the microwatershed through land resource inventory in the year 2017 were analysed for pH, EC, organic carbon, available phosphorus and potassium, and for micronutrients like zinc, boron, copper, iron and manganese, and secondary nutrient sulphur.

Soil fertility data generated has been assessed and individual maps for all the nutrients for the microwatershed have been generated using Kriging method under GIS. The village/survey number wise fertility data for the microwatershed is given in Appendix-II.

6.1 Soil Reaction (pH)

The soil analysis of the Yadgir RF1 microwatershed for soil reaction (pH) showed that an area of 227 ha (46%) is slightly acid (pH 6.0-6.5) and are distributed in the central, southeastern, eastern and western part of the microwatershed. An area of 258 ha (53%) is neutral (pH 6.5-7.3) and are distributed in the southwestern, southeastern, northwestern, northeastern, northern and southern part of the microwatershed (Fig. 6.1).

6.2 Electrical Conductivity (EC)

The Electrical Conductivity in entire area of the microwatershed is non saline (<2 dS/m) (Fig. 6.2) and as such the soils are non-saline.

6.3 Organic Carbon

The soil organic carbon content (an index of available Nitrogen) in the soils of the microwatershed is high (>0.75 %) in 388 ha (79%) and are distributed in the major part of the microwatershed. Medium (0.5-0.75%) in about 97 ha (20%) and are distributed in the western and southwestern part of the microwatershed (Fig. 6.3).




Fig.6.1 Soil Reaction (pH) map of Yadgir RF1 Microwatershed




Fig.6.2 Electrical Conductivity (EC) map of Yadgir RF1 Microwatershed




Fig.6.3 Soil Organic Carbon map of Yadgir RF1 Microwatershed

6.4 Available Phosphorus

Available phosphorus content is medium (23-57 kg/ha) in an entire area of about 485 ha (99%) and occur in the major part of the microwatershed (Fig. 6.4).

6.5 Available Potassium

Available potassium content is medium (145-337 kg/ha) in maximum area of about 343 ha (70%) and are distributed in the major part of the microwatershed. Low (<145 kg/ha) in an area of 142 ha (29%) and are distributed in the southwestern, central northeastern and southwestern part of the microwatershed (Fig. 6.5).

6.6 Available Sulphur

Available sulphur content is medium (10-20 ppm) in the entire area of the microwatershed (Fig. 6.6).

6.7 Available Boron

Available boron content is low (<0.5 ppm) in 436 ha (89%) and are distributed in all parts of the microwatershed. Medium (0.5-1.0ppm) in an area of 49 ha (10%) and are distributed in the southeastern and eastern part of the microwatershed (Fig. 6.7).

6.8 Available Iron

Available iron content is sufficient (>4.5 ppm) in the entire area of the microwatershed (Fig 6.8).

6.9 Available Manganese

Available manganese content is sufficient (>1.0 ppm) in the entire microwatershed area (Fig 6.9).

6.10 Available Copper

Available copper content is sufficient (>0.2 ppm) in the entire microwatershed area (Fig 6.10).




Fig.6.4 Soil Available Phosphorus map of Yadgir RF1Microwatershed




Fig.6.5 Soil Available Potassium map of Yadgir RF1 Microwatershed




Fig.6.6 Soil Available Sulphur map of Yadgir RF1 Microwatershed




Fig.6.7 Soil Available Boron map of Yadgir RF1Microwatershed




Fig.6.8 Soil Available Iron map of Yadgir RF1Microwatershed




Fig.6.9 Soil Available Manganese map of Yadgir RF1Microwatershed




Fig.6.10 Soil Available Copper map of Yadgir RF1Microwatershed

6.11 Available Zinc

Available zinc content is deficient (<0.6 ppm) in an area of 60 ha (12%) and is distributed in the western and southwestern part of the microwatershed. Maximum area of about 425 ha (87%) is sufficient (>0.6 ppm) and is distributed in all parts of the microwatershed (Fig 6.11).




Fig.6.11 Soil Available Zinc map of Yadgir RF1 Microwatershed

LAND SUITABILITY FOR MAJOR CROPS

The soil and land resource units (soil phases) of Yadgir RF1 microwatershed were assessed for their suitability for growing food, fodder, fibre and other horticulture crops by following the procedure as outlined in FAO, 1976 and 1983. Crop requirements were developed for each of the crop from the available research data and also by referring to Naidu *et. al.* (2006) and Natarajan *et. al.* (2015). The soil and land characteristics were matched with the crop requirement to arrive at the crop suitability. The soil and land characteristics table and crop requirement tables are given at the end. In FAO land suitability classification, two orders are recognized. Order S-Suitable and Order N-Not suitable. The orders have classes, subclasses and units. Order-S has three classes, Class S1-Highly Suitable, Class S2-Moderately Suitable and Class S3- Marginally Suitable. Order N has two classes, N1-Currently not Suitable and N2- Permanently not Suitable. There are no subclasses within the Class S1 as they will have very minor or no limitations for crop growth. Classes S2, S3, N1 and N2 are divided into subclasses based on the kinds of limitations encountered. The limitations that affect crop production are ‘c’ for erratic rainfall and its distribution and length of growing period (LGP), ‘e’ for erosion hazard, ‘r’ for rooting condition, ‘t’ for lighter or heavy texture, ‘g’ for gravelliness or stoniness, ‘n’ for nutrient availability, ‘l’ for topography, ‘m’ for moisture availability, ‘w’ for drainage, ‘s’ for sodium and ‘z’ for calcareousness. These limitations are indicated as lower case letters to the Class symbol. For example, moderately suitable lands with the limitations of soil depth and erosion are designated as S2re. For the microwatershed, the soil mapping units were evaluated and classified up to subclass level.

Using the above criteria, the soil map units of the microwatershed were evaluated and land suitability maps for 29 major agricultural and horticultural crops were generated. The detailed information on the kind of suitability of each of the soil phase for the crops assessed are given village/ survey number wise for the microwatershed in Appendix-III.

7.1 Land Suitability for Sorghum (*Sorghum bicolor*)

Sorghum is one of the major crop grown in Karnataka in an area of 10.47 lakh ha in Bijapur, Gulbarga, Raichur, Bidar, Belgaum, Dharwad, Bellary, Chitradurga, Mysore and Tumakuru districts. The crop requirements for growing sorghum (Table 7.2) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing sorghum was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.1.

Highly suitable (Class S1) lands for growing sorghum occur in an area of 29 ha (6%) and are distributed in the southeastern and southern part of the microwatershed. An area of about 20 ha (4%) is moderately suitable (Class S2) for growing sorghum and are distributed in the northern part of the microwatershed. They have minor limitations of

calcareousness and drainage. An area of about 277 ha (56%) is marginally suitable (Class S3) for growing sorghum and is distributed in the central, northeastern, southern, southeastern, eastern and western part of the microwatershed with moderate limitations of rooting depth and texture. Currently not suitable (Class N1) lands occur in an area of 159 ha (32%) and are distributed in the major part of the microwatershed with severe limitation of rooting depth.




Fig. 7.1 Land Suitability map of Sorghum

7.2 Land Suitability for Maize (*Zea mays*)

Maize is one of the most important food crop grown in an area of 13.37 lakh ha in almost all the districts of the State. The crop requirements for growing maize (Table 7.3) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing maize was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.2.

No highly suitable (Class S1) lands available for growing maize in the microwatershed. Moderately suitable (Class S2) lands occur in an area of 20 ha (4%) and are distributed in the northern part of the microwatershed with minor limitations of texture and calcareousness. Marginally suitable lands (Class S3) for growing maize occupy a maximum area of 306 ha (62%) and occur in all parts of the microwatershed. They have moderate limitations of rooting depth, calcareousness and texture. Currently not suitable (Class N1) lands occur in a maximum area of 159 ha (32%) and are

distributed in the northwestern, northeastern, western, southwestern and southeastern part of the microwatershed with severe limitation of rooting depth.




Fig. 7.2 Land Suitability map of Maize

7.3 Land Suitability for Bajra (*Pennisetum glaucum*)

Bajra is one of the most important millet crop grown in an area of 2.34 lakh ha in the northern districts of Karnataka state. The crop requirements for growing bajra (Table 7.4) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing bajra was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.3.

Highly suitable (Class S1) lands for growing bajra occur in an area of 29 ha (6%) and are distributed in the southern and southeastern part of the microwatershed. An area of about 20 ha (4%) is moderately suitable (Class S2) for growing bajra and are distributed in the northern part of the microwatershed. They have minor limitations of calcareousness and drainage. Marginally suitable lands (Class S3) occupy a maximum area of 277 ha (56%) and are distributed in the major part of the microwatershed. They have moderate limitations of rooting depth and texture. Currently not suitable (Class N1) lands occur in an area of 159 ha (32%) and are distributed in the northwestern, northeastern, western, southwestern and southeastern part of the microwatershed with severe limitation of rooting depth.




Fig. 7.3 Land Suitability map of Bajra

7.4 Land Suitability for Groundnut (*Arachis hypogaea*)

Groundnut is one of the major oilseed crop grown in an area of 6.54 lakh ha in Karnataka in most of the districts either as rainfed or irrigated crop. The crop requirements for growing groundnut (Table 7.5) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing groundnut was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.4.

There are no highly suitable (Class S1) lands available for growing groundnut in the microwatershed. Marginally suitable lands (Class S3) for growing groundnut occupy a maximum area of about 327 ha (67%) with moderate limitations of texture, calcareousness and rooting depth and are distributed in the major part of the microwatershed. Currently not suitable (Class N1) lands occur in an area of 159 ha (32%) and are distributed in the northwestern, northeastern, western, southwestern and southeastern part of the microwatershed with severe limitation of rooting depth.




Fig. 7.4 Land Suitability map of Groundnut

7.5 Land Suitability for Sunflower (*Helianthus annus*)

Sunflower is one of the most important oilseed crop grown in an area of 4.1 lakh ha in the State in all the districts. The crop requirements for growing sunflower (Table 7.6) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing sunflower was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.5.

An area of about 50 ha (10%) is moderately suitable (Class S2) for sunflower and are distributed in the northern, southeastern and southern part of the microwatershed. They have minor limitations of rooting depth and calcareousness. An area of about 73 ha (15%) is marginally suitable (Class S3) and is distributed in the southern, eastern and western part of the microwatershed with moderate limitation of texture. Currently not suitable (Class N1) lands occur in a maximum area of 362 ha (74%) and are distributed in all parts of the microwatershed with severe limitation of rooting depth.




Fig. 7.5 Land Suitability map of Sunflower

7.6 Land suitability for Red gram (*Cajanus Cajan*)

Redgram is one of the most important pulse crop grown in an area of 7.28 lakh ha in almost all the districts of the State. The crop requirements for growing red gram (Table 7.7) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing redgram was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.6.

No highly suitable (Class S1) lands are available for growing redgram in the microwatershed. An area of about 50 ha (10%) is moderately suitable (Class S2) for growing redgram and are distributed in the northern, southern and southeastern part of the microwatershed. They have minor limitations of rooting depth and calcareousness. Marginally suitable lands (Class S3) for growing redgram occupy an area of about 100 ha (20%) and occur in the southern, southeastern, western and northeastern part of the microwatershed. They have moderate limitations of rooting depth and texture. Currently not suitable (Class N1) lands occur in a maximum area of 336 ha (68%) and are distributed in all parts of the microwatershed with severe limitation of rooting depth.




Fig. 7.6 Land Suitability map of Redgram

7.7 Land Suitability for Bengal gram (*Cicer aerativum*)

Bengal gram is one of the most important pulse crop grown in about 9.39 lakh ha area in Bijapur, Raichur, Kalaburgi, Dharwad, Belgaum and Bellary districts. The crop requirements for growing Bengal gram (Table 7.8) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing Bengal gram was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.7.

Highly (Class S1) suitable lands for growing Bengal gram occupy an area of 29 ha (6%) and are distributed in the southern and southeastern part of the microwatershed. An area of about 20 ha (4%) is moderately suitable (Class S2) for growing Bengal gram and are distributed in the northern part of the microwatershed. They have minor limitations of calcareousness and drainage. Marginally suitable lands (Class S3) occupy an area of about 27 ha (5%) and are distributed in the northeastern and southeastern part of the microwatershed. They have moderate limitation of rooting depth. Currently not suitable (Class N1) lands occur in an area of 409 ha (83%) and are distributed in all parts of the microwatershed with severe limitations of rooting depth and texture.




Fig. 7.7 Land Suitability map of Bengal gram

7.8 Land Suitability for Cotton (*Gossypium hirsutum*)

Cotton is one of the most important fibre crop grown in the State in about 8.75 lakh ha area in Raichur, Dharwad, Belgaum, Kalaburgi, Bijapur, Bidar, Bellary, Chitradurga and Chamarajnagar districts. The crop requirements for growing cotton (Table 7.9) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing cotton was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.8.

Moderately suitable (Class S2) lands are found to occur in an area of about 50 ha (10%). These soils have minor limitations of rooting depth and calcareousness. They are distributed in the northern, southern and southeastern part of the microwatershed. Marginally suitable (Class S3) lands for growing cotton occur in an area of 27 ha (5%) with moderate limitation of rooting depth and are distributed in the northeastern and southeastern part the microwatershed. Currently not suitable (Class N1) lands occur in an area of 409 ha (83%) and are distributed in all parts of the microwatershed with severe limitations of rooting depth and texture.




Fig. 7.8 Land Suitability map of Cotton

7.9 Land Suitability for Chilli (*Capsicum annuum*)

Chilli is one of the most important spice crop grown in about 0.42 lakh ha in Karnataka State. The crop requirements for growing chilli (Table 7.10) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing chilli was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.9.

An area of about 50 ha (10%) is moderately suitable (Class S2) for growing chilli and are distributed in the northern, southern and southeastern part of the microwatershed. They have minor limitations of texture and calcareousness. Marginally suitable lands (Class S3) occupy a maximum area of 277 ha (56%) and are distributed in the major part of the microwatershed. They have moderate limitations of rooting depth and texture. Currently not suitable (Class N1) lands occur in an area of 159 ha (32%) and are distributed in all parts of the microwatershed with severe limitation of rooting depth.




Fig 7.9 Land Suitability map of Chilli

7.10 Land Suitability for Tomato (*Lycopersicon esculentum*)

Tomato is one of the most important vegetable crop grown in about 0.61 lakh ha covering almost all the district of the state. The crop requirements for growing tomato (Table 7.11) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing tomato was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.10.

Maximum area of 327 ha (67%) is marginally suitable for tomato (Class S3) and is distributed in the major part of the microwatershed. They have moderate limitations of rooting depth, drainage and texture. Currently not suitable (Class N1) lands occur in an area of 159 ha (32%) and are distributed in all parts of the microwatershed with severe limitations of rooting depth.




Fig 7.10 Land Suitability map of Tomato

7.11 Land Suitability for Brinjal (*Solanum melongena*)

Brinjal is one of the most important vegetable crop grown in the state. The crop requirements for growing brinjal (Table 7.12) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing brinjal was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.11.

Highly (Class S1) suitable lands for growing brinjal occur in an area of 50 ha (10%) and are distributed in the northern, southeastern and southern part of the microwatershed. Maximum area of 277 ha (56%) is marginally suitable (Class S3) and is distributed in the major part of the microwatershed with moderate limitations of rooting depth and texture. Currently not suitable (Class N1) lands occur in an area of 159 ha (32%) and are distributed in all parts of the microwatershed with severe limitation of rooting depth.




Fig 7.11 Land Suitability map of Brinjal

7.12 Land Suitability for Onion (*Allium cepa L.*)

Onion is one of the most important vegetable crop grown in the state. The crop requirements for growing onion (Table 7.13) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing onion was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.12.

Highly (Class S1) suitable lands for growing onion occur in an area of 50 ha (10%) and are distributed in the northern, southern and southeastern part of the microwatershed. Maximum area of 277 ha (56%) is marginally suitable and is distributed in the major part of the microwatershed with moderate limitations of rooting depth and texture. Currently not suitable (Class N1) lands occur in an area of 159 ha (32%) and are distributed in all parts of the microwatershed with severe limitation of rooting depth.




Fig 7.12 Land Suitability map of Onion

7.13 Land Suitability for Bhendi (*Abelmoschus esculentus*)

Bhendi is one of the most important vegetable crop grown in the state. The crop requirements for growing bhendi (Table 7.14) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing bhendi was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.13.

Highly (Class S1) suitable lands for growing bhendi occur in a maximum area of 50 ha (10%) and are distributed in the northern, southern and southeastern part of the microwatershed. Maximum area of 277 ha (56%) is marginally suitable and is distributed in the major part of the microwatershed with moderate limitations of rooting depth and texture. Currently not suitable (Class N1) lands occur in an area of 159 ha (32%) and are distributed in all parts of the microwatershed with severe limitation of rooting depth.




Fig 7.13 Land Suitability map of Bhendi

7.14 Land Suitability for Drumstick (*Moringa oleifera*)

Drumstick is one of the most important vegetable crop grown in about 2403 ha in the state. The crop requirements for growing drumstick (Table 7.15) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing drumstick was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.14.

There are no highly (Class S1) suitable lands available for growing drumstick in the microwatershed. An area of about 50 ha (10%) is moderately suitable (Class S2) for drumstick and is distributed in the northern, southern and southeastern part of the microwatershed. They have minor limitations of rooting depth and calcareousness. An area of 73 ha (15%) is marginally suitable and is distributed in the southern, southeastern and southwestern part of the microwatershed with moderate limitation of texture. Currently not suitable (Class N1) lands occur in a maximum area of 363 ha (74%) and are distributed in all parts of the microwatershed with severe limitations of rooting depth and texture.




Fig 7.14 Land Suitability map of Drumstick

7.15 Land suitability for Mango (*Mangifera indica*)

Mango is one of the most important fruit crop grown in an area of 1.73 lakh ha in almost all the districts of the State. The crop requirements (Table 7.16) for growing mango were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing mango was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed is given in Figure 7.15.

An area of 50 ha (10%) is marginally suitable (Class S3) for growing mango with moderate limitations of calcareousness and rooting depth and are distributed in the northern, southern and southeastern part of the microwatershed. Maximum area of about 435 ha (89%) is currently not suitable (Class N1) for growing mango and are distributed in the major part of the microwatershed with severe limitation of rooting depth.




Fig. 7.15 Land Suitability map of Mango

7.16 Land suitability for Guava (*Psidium guajava*)

Guava is one of the most important fruit crop grown in an area of 6558 ha in almost all the districts of the State. The crop requirements (Table 7.17) for growing guava were matched with the soil-site characteristics (7.1) and a land suitability map for growing guava was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.16.

Marginally suitable (Class S3) lands cover an area of about 123 ha (25%) and are distributed in the northern, southern, southeastern and southwestern part of the microwatershed. They have moderate limitations of rooting depth, calcareousness and texture. Maximum area of about 363 ha (74%) is currently not suitable (N) for growing guava and occur in the major part of the microwatershed with severe limitations of rooting depth and texture.




Fig. 7.16 Land Suitability map of Guava

7.17 Land suitability for Sapota (*Manilkara zapota*)

Sapota is one of the most important fruit crop grown in an area of 29373 ha in almost all the districts of the State. The crop requirements (Table 7.18) for growing sapota were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing sapota was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.17.

An area of about 123 ha (25%) is marginally suitable (Class S3) for growing sapota and are distributed in the northern, southern, southeastern and southwestern part of the microwatershed. They have moderate limitations of texture, rooting depth and calcareousness. Maximum area of 362 ha (74%) is currently not suitable (Class N1) for growing sapota and occur in the major part of the microwatershed with severe limitation of rooting depth.




Fig. 7.17 Land Suitability map of Sapota

7.18 Land Suitability for Pomegranate (*Punica granatum*)

Pomegranate is one of the most important fruit crop commercially grown in about 18488 ha in Karnataka, mainly in Bijapur, Bagalkot, Koppal, Gadag and Chitradurga districts. The crop requirements for growing pomegranate (Table 7.19) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing pomegranate was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.18.

No highly (Class S1) suitable lands available for growing pomegranate in the microwatershed. An area of about 50 ha (10%) is moderately suitable (Class S2) for growing pomegranate and is distributed in the northern, southern and southeastern part of the microwatershed. They have minor limitations of rooting depth and calcareousness. An area of 73 ha (15%) is marginally suitable for pomegranate and is distributed in the southern, southwestern and southeastern part of the microwatershed with moderate limitations of rooting depth and texture. Maximum area of about 362 ha (74%) is currently not suitable (Class N1) for growing pomegranate and is distributed in the major part of the microwatershed with severe limitation of rooting depth.




Fig 7.18 Land Suitability map of Pomegranate

7.19 Land Suitability for Musambi (*Citrus limetta*)

Musambi is one of the important fruit crop grown in an area of 3446 ha in almost all the districts of the State. The crop requirements for growing musambi (Table 7.20) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing musambi was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.19.

An area of about 50 ha (10%) is moderately suitable (Class S2) for growing Musambi and are distributed in the northern, southern and southeastern part of the microwatershed. They have minor limitations of calcareousness and rooting depth. An area of about 73 ha (15%) is marginally suitable and is distributed in the southern, southeastern and southwestern part of the microwatershed with moderate limitation of rooting depth and texture. Currently not suitable (Class N1) lands occur in a maximum area of 362 ha (74%) and are distributed in the major part of the microwatershed with severe limitation of rooting depth.




Fig. 7.19 Land Suitability map of Musambi

7.20 Land Suitability for Lime (*Citrus sp*)

Lime is one of the most important fruit crop grown in an area of 0.11 lakh ha in almost all the districts of the State. The crop requirements for growing lime (Table 7.21) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing lime was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7. 20.

An area of about 50 ha (10%) is moderately suitable (Class S2) for growing lime and are distributed in the northern, southern and southeastern part of the microwatershed. They have minor limitations of calcareousness and rooting depth. An area of about 73 ha (15%) is marginally suitable and is distributed in the southern, southeastern and southwestern part of the microwatershed with moderate limitations of rooting depth and texture. Currently not suitable (Class N1) lands occur in a maximum area of 362 ha (74%) and are distributed in the major part of the microwatershed with severe limitation of rooting depth.




Fig. 7.20 Land Suitability map of Lime

7.21 Land Suitability for Amla (*Phyllanthus emblica*)

Amla is one of the medicinal fruit crop grown in almost all the districts of the State. The crop requirements for growing amla (Table 7.22) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing amla was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.21.

An area of about 49 ha (10%) has soils that are moderately suitable (Class S2) for growing Amla with minor limitations of calcareousness and drainage and are distributed in the northern, southeastern and southern part of the microwatershed. An area of 277 ha (56%) is marginally suitable (Class S3) for growing amla with moderate limitations of texture and rooting depth and is distributed in the major part of the microwatershed. Currently not suitable (Class N1) lands occur in an area of 159 ha (32%) and are distributed in the northern, northeastern, northwestern, western, southwestern and southeastern part of the microwatershed with severe limitation of rooting depth.




Fig. 7.21 Land Suitability map of Amla

7.22 Land Suitability for Cashew (*Anacardium occidentale*)

Cashew is one of the most important plantation nut crop grown in an area of 0.7 lakh ha in almost all the districts of the State. The crop requirements for growing cashew (Table 7.23) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing cashew was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.22.

An area of 73 ha (15%) is marginally suitable (Class S3) for cashew and is distributed in the southern, southeastern and southwestern part of the microwatershed with severe limitations of rooting depth and texture. Currently not suitable (Class N1) lands occur in an area of 413 ha (84%) and are distributed in all parts of the microwatershed with severe limitations of rooting depth and texture.




Fig. 7.22 Land Suitability map of Cashew

7. 23 Land Suitability for Jackfruit (*Artocarpus heterophyllus*)

Jackfruit is one of the most important fruit crop grown in an area of 5368 ha in almost all the districts of the State. The crop requirements for growing jackfruit (Table 7.24) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing jackfruit was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.23.

Marginally suitable (Class S3) lands occupy an area of about 123 ha (25%) and are distributed in the northern, southern, southeastern and western part of the microwatershed. They have moderate limitations of texture, rooting depth and calcareousness. Maximum area of about 363 ha (74%) is currently not suitable (Class N1) and is distributed in the major part of the microwatershed with severe limitations of rooting depth and texture.




Fig. 7.23 Land Suitability map of Jackfruit

7.24 Land Suitability for Jamun (*Syzygium cumini*)

Jamun is an important fruit crop grown in almost all the districts of the State. The crop requirements for growing jamun (Table 25) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing jamun was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.24.

Marginally suitable (Class S3) lands occupy an area of about 123 ha (25%) and are distributed in the northern, southern, southeastern and western part of the microwatershed. They have moderate limitations of texture, rooting depth and calcareousness. Maximum area of about 363 ha (74%) is currently not suitable (Class N1) and is distributed in the major part of the microwatershed with severe limitations of rooting depth and texture.




Fig. 7.24 Land Suitability map of Jamun

7.25 Land Suitability for Custard Apple (*Annona reticulata*)

Custard apple is one of the most important fruit crop grown in almost all the districts of the State. The crop requirements for growing custard apple (Table 7.26) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing custard apple was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.25.

Highly suitable (Class S1) lands for growing custard apple occur in an area of 20 ha (4%) and are distributed in the northern part of the microwatershed. An area of about 29 ha (6%) is moderately suitable (Class S2) for growing custard apple and is distributed in the southern and southeastern part of the microwatershed with minor limitation of calcareousness. Marginally suitable (Class S3) lands occur in an area of 277 ha (56%) and are distributed in all parts of the microwatershed with moderate limitations of texture and rooting depth. Currently not suitable (Class N1) lands occur in an area of 159 ha (32%) and are distributed in the northern, northeastern, northwestern, western, southwestern and southeastern part of the microwatershed with severe limitation of rooting depth.




Fig. 7.25 Land Suitability map of Custard Apple

7.26 Land Suitability for Tamarind (*Tamarindus indica*)

Tamarind is one of the most important spice crop grown in almost all the districts of the state. The crop requirements for growing tamarind (Table 7.27) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing tamarind was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.26.

Marginally suitable (Class S3) lands for growing Tamarind occupy an area of about 50 ha (10%) and are distributed in the northern, southeastern and southern part of the microwatershed. They have moderate limitations of rooting depth and calcareousness. Currently not suitable (Class N1) lands occur in an area of 436 ha (89%) and are distributed in the major part of the microwatershed with severe limitations of rooting depth and texture.




Fig. 7.26 Land Suitability map of Tamarind

7.27 Land Suitability for Mulberry (*Morus nigra*)

Mulberry is one of the important leaf crop grown for rearing silk worms in about 1.6 lakh ha area in all the districts of the state. The crop requirements for growing mulberry (Table 7.28) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing mulberry was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.27.

Marginally suitable (Class S3) lands occur in a maximum area of 123 ha (25%) and are distributed in the northern, southern, southeastern and western part of the microwatershed with moderate limitations of texture and calcareousness. Currently not suitable lands (Class N1) occupy an area of about 363 ha (74%) and distributed in the major part of the microwatershed. They have severe limitations of rooting depth and texture.




Fig 7.27 Land Suitability map of Mulberry

7.28 Land suitability for Marigold (*Tagetes sps.*)

Marigold is one of the most important flower crop grown in an area of 9108 ha in almost all the districts of the State. The crop requirements (Table 7.29) for growing marigold were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing marigold was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.28.

An area of about 50 ha (10%) is moderately suitable (Class S2) for growing Marigold and are distributed in the northern, southern and southeastern part of the microwatershed. They have minor limitations of texture, rooting depth and calcareousness. Marginally suitable (Class S3) lands occupy an area of about 277 ha (56%) and are distributed in the major part of the microwatershed. They have moderate limitations of rooting depth and texture. Currently not suitable (Class N1) lands occur in an area of 159 ha (32%) and are distributed in the northern, northeastern, northwestern, western, southwestern and southeastern part of the microwatershed with severe limitation of rooting depth.




Fig. 7.28 Land Suitability map of Marigold

7.29 Land Suitability for Chrysanthemum (*Dendranthema grandiflora*)

Chrysanthemum is one of the most important flower crop grown in an area of 4978 ha in almost all the districts of the State. The crop requirements (Table 7.30) for growing chrysanthemum were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing chrysanthemum was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.29.

An area of about 50 ha (10%) is moderately suitable (Class S2) for growing chrysanthemum and are distributed in the northern, southern and southeastern part of the microwatershed. They have minor limitations of texture, rooting depth and calcareousness. Marginally suitable (Class S3) lands occupy an area of about 277 ha (56%) and are distributed in the major part of the microwatershed. They have moderate limitations of rooting depth and texture. Currently not suitable (Class N1) lands occur in an area of 159 ha (32%) and are distributed in the northern, northeastern, northwestern, western, southwestern and southeastern part of the microwatershed with severe limitation of rooting depth.




Fig. 7.29 Land Suitability map of Chrysanthemum

Table 7.1 Soil-Site Characteristics of Yadgir RF1Microwatershed

Soil Map Units	Climate (P) (mm)	Growing period (Days)	Drainage Class	Soil depth (cm)	Soil texture		Gravelliness		AWC (mm/m)	Slope (%)	Erosion	pH	EC (dSm ⁻¹)	ESP (%)	CEC [Cmol (p ⁺)kg ⁻¹]	BS (%)
					Surface	Sub-surface	Surface (%)	Sub-surface (%)								
BDPiB3	866	150	WD	<25	sc	scl	<15	<15	<50	1-3	severe	8.58	0.26	0.35	18.10	100
BDPhB2	866	150	WD	<25	scl	scl	<15	<15	<50	1-3	moderate	8.58	0.26	0.35	18.10	100
HTKcC2g1	866	150	WD	25-50	sl	sl	15-35	10-25	<50	3-5	moderate	6.81	0.06	0.38	3	92
HTKbB2g1	866	150	WD	25-50	ls	sl	15-35	10-25	<50	1-3	moderate	6.81	0.06	0.38	3	92
KKRbB2g1	866	150	WD	25-50	ls	sl	15-35	10-15	<50	1-3	moderate	5.85	0.03	1.17	2.6	61
KKRcB2	866	150	WD	25-50	sl	sl	<15	10-15	<50	1-3	moderate	5.85	0.03	1.17	2.6	61
BDLbB2	866	150	WD	25-50	ls	sl	<15	<15	<50	1-3	moderate	6.20	0.074	0.20	4.20	93
BDLhB2	866	150	WD	25-50	scl	sl	<15	<15	<50	1-3	moderate	6.20	0.074	0.20	4.20	93
SBRcB2	866	150	SED	50-75	sl	ls	<15	<15	<50	1-3	moderate	8.24	0.14	1.15	7.50	100
SBRcC3g1	866	150	SED	50-75	sl	ls	15-35	<15	<50	3-5	severe	8.24	0.14	1.15	7.50	100
GWDcB2	866	150	MWD	75-100	sl	scl	<15	<15	101-150	1-3	moderate	9.89	0.74	17.40	8.35	100
GWDmB2	866	150	MMD	75-100	c	scl	<15	<15	101-150	1-3	moderate	9.89	0.74	17.40	8.35	100

*Symbols and abbreviations are according to Field Guide for LRI under Sujala-III Project, Karnataka

Table 7.2 Land suitability criteria for Sorghum

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	26–30	30–34; 24–26	34–40; 20–24	>40; <20
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, c (red), c (black)	scl, cl	ls, sl	-
	pH	1:2.5	5.5-7.8	5.0-5.5 7.8-9.0	>9.0	-
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	10-15
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

Table 7.3 Land suitability criteria for Maize

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	30-34	35-38 26-30	38-40 26-20	
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc	c (red), c (black)	ls, sl	-
	pH	1:2.5	5.5-7.8	5.0-5.5 7.8-9.0	>9.0	-
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	-
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

Table 7.4 Land suitability criteria for Bajra

Land use requirement		Rating				
Soil –site characteristics	Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)	
Climatic regime	Mean temperature in growing season	°C	28-32	33-38 24-27	39-40 20-23	<20
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm	500-750	400-500	200-400	<200
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl, scl, cl,sc,c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.8	5.0-5.5 7.8-9.0	5.5-6.0 >9.0	
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	15-35	35-60	>60	
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	1-3	3-5	5-10	>10

Table 7.5 Land suitability criteria for Groundnut

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	24–33	22–24; 33–35	20–22; 35–40	<20; >40
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. Well drained	Poorly drained	Very Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl	sl,cl, sc	c (red), c (black), ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<35	35-60	>60	
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.6 Land suitability criteria for Sunflower

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	24–30	30–34; 20–24	34–38; 16–20	>38; <16
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	mod. Well drained	-	Poorly to very drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	cl, sc,c (red), c (black)	scl	ls, sl	-
	pH	1:2.5	6.5-7.8	7.8-8.4 5.5-6.5	8.4-9.0; 5.0-5.5	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.7 Land suitability criteria for Redgram

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	30-35(G) 20-25(AV) 15-18 (F&PS) 35-40(M)	25-30(G) 20-25 (AV) 12-15 (F&PS) 30-35(M)	20-25(G) 15-20(AV) 10-12 (F&PS) 25-30(M)	< 20 <15 <10 <25
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. Well drained	Poorly drained	Very Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, c (red)	c (black),sl, scl, cl	ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-9.0	5.0-5.5 >9.0	-
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-50	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<1.0	1.0-2.0	>2.0	
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.8 Land suitability criteria for Bengal gram

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	20–25	25–30; 15–20	30–35; 10–15	>35; <10
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. Well drained	Poorly drained	Very Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	c(black)	-	c (red), scl, cl, sc	ls, sl
	pH	1:2.5	6.0-7.8	5.0-6.0 7.8-9.0	>9.0	-
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	-
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.9 Land suitability criteria for Cotton

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	22-32	>32	<19	-
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well to moderately well	Poorly drained/Some what excessively drained	-	very poorly/excessively drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, c (red,black)	cl	scl	ls, sl
	pH	1:2.5	6.5-7.8	7.8-8.4	5.5-6.5 8.4->9.0	<5.5
	CEC	C mol (p+)Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	50-100	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	<3	3-5	-	>5

Table 7.10 Land suitability criteria for Chilli

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	25-32	33-35 20-25	35-38 <20	>38
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc	c (black), sl	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.11 Land suitability criteria for Tomato

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	25-28	29-32 20-24	15-19 33-36	<15 >36
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl, scl, cl, sc, c (red)	-	ls, c(black)	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.12 Land suitability criteria for Brinjal

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	Well drained	Moderately well drained	Poorly drained	V. Poorly drained
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class				
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl, scl, cl, sc c (red)	-	ls, c (black)	-
	pH	1:2.5	6.0-7.3	7.3-8.4 5.0-6.0	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	>60
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.13 Land suitability criteria for Onion

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	20-30	30-35	35-40	>40
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately /imperfectly	-	Poorly to V poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl,scl,cl,sc,c (red)	-	c (Black),ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<1.0	1.0-2.0	2.0-4.0	<4
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.14 Land suitability criteria for Bhendi

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	25-28	29-32 20-24	15-19 33-36	<15 >36
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Imperfectly drained	Poorly to very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl,sc, c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.15 Land suitability criteria for Drumstick

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, scl, cl, c (red)	sl, c (black)	ls	s
	pH	1:2.5	6.0-7.3	5.0-5.5 7.3-7.8	5.5-6.0 7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<35	35-60	60-80	>80
Soil toxicity	Salinity (EC saturation extract)	ds/m				
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-10	-	>10

Table 7.16 Land suitability criteria for Mango

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-32	24-27 33-35	36-40	20-24
	Min temp. before flowering	°C	10-15	15-22	>22	-
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration	Days				
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V. Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	-	ls, sl, c (black)	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>150	100-150	75-100	<75
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.17 Land suitability criteria for Guava

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-32	33-36 24-27	37-42 20-23	
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	sl	c (black), ls	-
	pH	1:2.5	6.0-7.8	5.0-6.0	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.18 Land suitability criteria for Sapota

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-32	33-36 24-27	37-42 20-23	>42 <18
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	-	Poorly to very drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	sl	ls, c (black)	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.19 Land suitability criteria for Pomegranate

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	30-34	35-38 25-29	39-40 15-24	
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl,cl, sc, c (red)	c (black),sl	ls	-
	pH	1:2.5	5.5-7.8	7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.20 Land suitability criteria for Musambi

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-30	31-35 24-27	36-40 20-23	>40 <20
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately drained	poorly	Very poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c	sl	ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.21 Land suitability criteria for Lime

Land use requirement		Rating				
Soil –site characteristics	Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)	
Climatic regime	Mean temperature in growing season	°C	28-30	31-35 24-27	36-40 20-23	>40 <20
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately drained	poorly	Very poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c	sl	ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.22 Land suitability criteria for Amla

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. well drained	Poorly drained	V. Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	c (black)	ls, sl	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15-35	35-60	60-80	-
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

Table 7.23 Land suitability criteria for Cashew

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	32 to 34	28 to 32; 34 to 38	24 to 28; 38 to 40	<20; >40
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	-	sl, ls	c (black)
	pH	1:2.5	5.5-6.5	5.0-5.5 6.5-7.3	7.3-7.8	>7.8
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-10	>10	-

Table 7.24 Land suitability criteria for Jackfruit

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. well	Poorly	V. Poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	-	sl, ls, c (black)	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	>60
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10-

Table 7.25 Land suitability criteria for Jamun

Land use requirement		Rating				
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well	Mod. well	Poorly	V.Poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c(red)	sl, c (black)	ls	-
	pH	1:2.5	6.0-7.8	5.0-6.0	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>150	100-150	50-100	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	>60
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

Table 7.26 Land suitability criteria for Custard apple

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	Scl, cl, sc, c (red), c (black)	-	Sl, ls	-
	pH	1:2.5	6.0-7.3	5.5-6.0 7.3-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15-35	35-60	60-80	-
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	>5	-

Table 7.27 Land suitability criteria for Tamarind

Land use requirement		Rating				
Soil –site characteristics	Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)	
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod.well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl,sc, c (red)	sl, c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>150	100-150	75-100	<75
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

Table 7.28 Land suitability criteria for Mulberry

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	24–28	22–24; 28–32	32–38; 22–18	>38; <18
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V. Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, cl, scl	c (red)	c (black), sl, ls	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.8-8.4	7.3-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	0-35	35-60	60-80	>80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

Table 7.29 Land suitability criteria for Marigold

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	18-23	17-15 24-35	35-40 10-14	>40 <10
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl,scl, cl, sc, c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%				
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.30 Land suitability criteria for Chrysanthemum

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	18-23	17-15 24-35	35-40 10-14	>40 <10
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl,scl, cl, sc, c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%				
Erosion hazard	Slope	%	<3	3-5	5-10	>10

7.30 Land Management Units (LMUs)

The 12 soil map units identified in Yadgir RF1 microwatershed have been grouped into 3 Land Management Units (LMU's) for the purpose of preparing a Proposed Crop Plan. Land Management Units are grouped based on the similarities in respect of the type of soil, the depth of the soil, the surface soil texture, gravel content, AWC, slope, erosion etc. and a Land Management Units map (Fig. 7.30) has been generated. These Land Management Units are expected to behave similarly for a given level of management.

The 12 map units that have been grouped into 3 Land Management Units along with brief description of soil and site characteristics are given below.

LMU	Soil map units	Soil and site characteristics
1	34.GWDcB2 127.GWDmB2	Moderately deep (75 to 100 cm), calcareous, sandy clay loam soils, 1-3% slopes, non gravelly (15%), moderate erosion.
2	11.SBRcB2 12.SBRcC3g1	Moderately shallow (50 to 75 cm), loamy sand soils, 1-5% slopes, non gravelly to gravelly (<15-35%), moderate to severe erosion.
3	2.BDLbB2 4.BDLhB2 113.HTKcC2g1 119.BDPiB3 120.BDPhB2 153.KKRbB2g1 161.HTKbB2g1 175.KKRcB2	Very shallow to shallow (<25 to 50 cm), sandy loam soils, 1-5% slopes, non gravelly to gravelly (<15-35%), moderate to severe erosion.




Fig. 7.30 Land Management Units Map- Yadgir RF1Microwatershed

7.31 Proposed Crop Plan for Yadgir RF1Microwatershed

After assessing the land suitability for the 29 crops, the Proposed Crop Plan has been prepared for the 3 identified LMUs by considering only the highly (Class S1) and moderately (Class S2) suitable lands for each of the 29 crops. The resultant proposed crop plan is presented below in Table 7.31.

Table 7.31 Proposed Crop Plan for Yadgir RF1Microwatershed

LMU	Soil Map Units	Survey Number	Field Crops/Commercial crops	Horticulture Crops (Rainfed/Irrigated)	Suitable Interventions
1	34.GWDcB2 127.GWDmB2 (Moderately deep, calcareous sandy clay loam soils)	Gajarakota: 696,697,703,704,705 Yampada: 179,180,216,217,259,260,263,264,265,266	Sunflower, Sorghum, Maize, Soybean, Cotton, Safflower, Linseed, Bajra	Fruit crops: , Pomegranate, Sapota, Guava, Lime, Musambi, Amla, Custard apple Vegetables: Cluster bean, Chilli, Drumstick, Bhendi, Coriander Flowers: Marigold, Chrysanthemum	Application of FYM, Biofertilizers and micronutrients, drip irrigation, mulching, suitable soil and water conservation practices
2	11.SBRcB2 12.SBRcC3g1 (Moderately shallow, loamy sand soils)	Yampada: 202,210,211,212,213,218,219,220,221,258,261,262,267,268,269,270,273,287,296,297,298,299,300,301,302,303,304,308,309,310,311,312,313,314	- Tuber crops	Agri- silvi- Pasture: Hybrid napier, <i>Styloxanthes hamata</i> , <i>Styloxanthes scabra</i>	Use of short duration varieties, sowing across the slope and split application of nitrogenous fertilizers
3	2.BDLbB2 4.BDLhB2 113.HTKcC2g1 119.BDPiB3 120.BDPhB2 153.KKRbB2g1 161.HTKbB2g1 175.KKRcB2 (Shallow to very shallow, sandy loam soils)	Gajarakota: 685,686,687,688,689,690,691,692,693,694,695,698,699,700,706,707,708,709,814 Yampada: 145,155,158,159,160,161,171,173,174,175,176,177,178,181,209,214,222,223,247,248,249,271,272,274,275,276,277,278,279,280,281,286,289,292,293,294,295,307,315,316,317,318,319,320,321,322,323,324,325,326,327,328,329,330,331,332,333,334,335,336,337,338,339,342,344,345,346,347,348,349,350,351,352,353,354,355,356,357,358,359,360,361,362,363	-	Agri-Silvi-Pasture: Hybrid Napier, <i>Styloxanthes hamata</i> , <i>Styloxanthes scabra</i>	Use of short duration varieties, sowing across the slope, drip irrigation and mulching is recommended

SOIL HEALTH MANAGEMENT

8.1 Soil Health

Soil health is basic to plant health and plant health is basic to human and bovine health. Soil is fundamental to crop production. Without soil, no food could be produced nor would livestock be fed on a large scale. Because it is finite and fragile, soil is a precious resource that requires special care from its users.

Soil health or the capacity of the soil to function is critical to human survival. Soil health has been defined as: “the capacity of the soil to function as a living system without adverse effect on the ecosystem”. Healthy soils maintain a diverse community of soil organisms that help to form beneficial symbiotic associations with plant roots, recycle essential plant nutrients, improve soil structure with positive repercussions for soil, water and nutrient holding capacity and ultimately improve crop production and also contribute to mitigating climate change by maintaining or increasing its carbon content.

Functional interactions of soil biota with organic and inorganic components, air and water determine a soil’s potential to store and release nutrients and water to plants and to promote and sustain plant growth. Thus, maintaining soil health is vital to crop production and conserve soil resource base for sustaining agriculture.

The most important characteristics of a healthy soil are

- Good soil tilth
- Sufficient soil depth
- Good water storage and good drainage
- Adequate supply, but not excess of nutrients
- Large population of beneficial organisms
- Small proportion of plant pathogens and insect pests
- Low weed pressure
- Free of chemicals and toxins that may harm the crop
- Resistance to degradation
- Resilience when unfavorable conditions occur

Characteristics of Yadgir RF1 Microwatershed

- ❖ The soil phases identified in the microwatershed belonged to the soil series of HTK 177 ha (36%), KKR 139 ha (28%), SBR 72 ha (15%), GWD 49 ha (10%), BDL 27 (5%) and BDP 20 ha (4%).
- ❖ As per land capability classification entire area of the microwatershed falls under arable land category (Class II, III & IV). The major limitations identified in the arable lands were soil and erosion.
- ❖ On the basis of soil reaction, about 227 ha (46%) area is slightly acid and 258 ha (53%) is neutral in soil reaction.

Soil Health Management

The following actions are required to improve the current land husbandry practices that provide a sound basis for the successful adoption of sustainable crop production system.

Acid soils

Acid soils cover about 227 ha area in the microwatershed.

1. Growing of crops suitable for a particular soil pH.
2. Amelioration of soils through the application of amendments (liming materials).

Liming materials:

1. CaCO_3 (Calcium Carbonate).
2. Dolomite [$\text{Ca Mg} (\text{CO}_3)_2$]
3. Quick lime (Cao)
4. Slaked lime [$\text{Ca} (\text{OH})_2$]

For normal pH and pH 4.8 (35 t/ha) and pH 6.0-7.0 (4 t/ha) lime is required.

Alkaline soils

Slightly Alkaline soils are not found in the microwatershed.

1. Regular addition of organic manure, green manuring, green leaf manuring, crop residue incorporation and mulching needs to be taken up to improve the soil organic matter status.
2. Application of biofertilizers (Azospirillum, Azatobacter, Rhizobium).
3. Application of 25% extra N and P (125 % RDN&P).
4. Application of ZnSO_4 – 12.5 kg/ha (once in three years).
5. Application of Boron – 5kg/ha (once in three years).

Neutral soils

Neutral soils occur in 258 ha area in the microwatershed.

1. Regular addition of organic manure, green manuring, green leaf manuring, crop residue incorporation and mulching needs to be taken up to improve the soil organic matter status.
2. Application of biofertilizers, (Azospirillum, Azotobacter, Rhizobium).
3. Application of 100 per cent RDF.
4. Need based micronutrient applications.

Besides the above recommendations, the best transfer of technology options are also to be adopted.

Soil Degradation

Soil erosion is one of the major factor affecting the soil health in the microwatershed. Out of total 490 ha area in the microwatershed, about 419 ha (86%) is suffering from moderate erosion and 65 ha (13%) from severe erosion. In areas of

moderate and severe erosion immediate soil and water conservation and, other land development and land husbandry practices are required for restoring soil health.

Dissemination of Information and Communication of Benefits

Any large scale implementation of soil health management requires that supporting information is made available widely, particularly through channels familiar to farmers and extension workers. Given the very high priority attached to soil-health especially by the Central Government on issuing Soil-Health Cards to all the farmers, media outlets like Regional, State and National Newspapers, Radio and Dooradarshan programs in local languages but also modern information and communication technologies such as Cellular phones and the Internet, which can be much more effective in reaching the younger farmers.

Inputs for Net Planning (Saturation Plan) and Interventions needed

Net planning (Saturation Plan) in IWMP is focusing on preparation of

1. Soil and Water Conservation Plan for each plot or farm.
2. Productivity enhancement measures/ interventions for existing crops/livestock/other farm enterprises.
3. Diversification of farming mainly with perennial horticultural crops and livestock.
4. Improving livelihood opportunities and income generating activities.

In this connection, how various outputs of Sujala-III are of use in addressing these objectives of Net Planning (Saturation Plan) are briefly presented below.

- ❖ **Soil Depth:** The depth of a soil decides the amount of moisture and nutrients it can hold, what crops can be taken up or not, depending on the rooting depth and the length of growing period available for raising any crop. Deeper the soil, better for a wide variety of crops. If sufficient depth is not available for growing deep rooted crops, either choose medium or short duration crops or deeper planting pits need to be opened and additional good quality soil brought from outside has to be filled into the planting pits.
- ❖ **Surface Soil Texture:** Lighter soil texture in the top soil means, better rain water infiltration, less run-off and soil moisture conservation, less capillary rise and less evaporation losses. Lighter surface textured soils are amenable to good soil tilth and are highly suitable for crops like groundnut, root vegetables (carrot, raddish, potato etc) but not ideal for crops that need stagnant water like lowland paddy. Heavy textured soils are poor in water infiltration and percolation. They are prone for sheet erosion; such soils can be improved by sand mulching. The technology that is developed by the AICRP-Dryland Agriculture, Vijayapura, Karnataka can be adopted.
- ❖ **Gravelliness:** More gravel content is favorable for run-off harvesting but poor in soil moisture storage and nutrient availability. It is a significant parameter that decides the kind of crop to be raised.

- ❖ **Land Capability Classification:** The land capability map shows the areas suitable and not suitable for agriculture and the major constraints in each of the plot/survey number. Hence, one can decide what kind of enterprise is possible in each of these units. In general, erosion and soil are the major constraints in Yadgir RF1 microwatershed.
- ❖ **Organic Carbon:** The OC content (an index of available Nitrogen) is high (>0.75%) in an area of 388 ha (79%) and medium (0.5-0.75%) in 97 ha (20%) of the area. The areas that are medium and low in OC needs to be further improved by applying farmyard manure and crop rotation with cereals and legumes or mixed cropping.
- ❖ **Promoting green manuring:** Growing of green manuring crops costs Rs. 1250/ha (green manuring seeds) and about Rs. 2000/ha towards cultivation that totals to Rs. 3250/- per ha. On the other hand, application of organic manure @ 10 tons/ha costs Rs. 5000/ha. The practice needs to be continued for 2-3 years or more. Nitrogen fertilizer needs to be supplemented by 25% in addition to the recommended level in 97 ha area where OC is medium (<0.5 - 0.75%). For example, for rainfed maize, recommended level is 50 kg N per ha and an additional 12 kg /ha needs to be applied for all the crops grown in these plots.
- ❖ **Available Phosphorus:** Available Phosphorus is medium (23-57 kg/ha) in an entire area of the microwatershed. In medium areas, for all the crops 25% additional P needs to be applied.
- ❖ **Available Potassium:** Available potassium is medium (145-337 kg/ha) in an area of 343 ha (70%) of the microwatershed and low (<145 kg/ha) in 142 ha (29%). All the plots, where available potassium is low and medium, for all the crops, additional 25% potassium may be applied.
- ❖ **Available Sulphur:** Available sulphur is a very critical nutrient for oilseed crops. It is medium in 485 ha (99%). Medium areas need to be applied with magnesium sulphate or gypsum or Factamphos (p) fertilizer (13% sulphur) for 2-3 years for the deficiency to be corrected.
- ❖ **Available Boron:** Maximum area of 436 ha (89%) is low (<0.5 ppm) and 49 ha (10%) is medium (0.5-1.0 ppm) in available boron. Application of sodium tetra borate @ 10 kg/ha as soil application or 0.2 % borax as foliar spray is recommended for low and medium areas.
- ❖ **Available Iron:** Entire area of the microwatershed is sufficient (>4.5 ppm) in available iron content.
- ❖ **Available Manganese:** All the soils in the microwatershed are sufficient (>1.0 ppm) in available manganese.
- ❖ **Available Copper:** All the soils in the microwatershed are sufficient (>0.2 ppm) in available copper.


- ❖ **Available Zinc:** An area of about 60 ha (12%) is deficient (<0.6 ppm) and an area of 425 ha (87%) is sufficient in available zinc content. Application of zinc sulphate @25 kg/ha is recommended for deficient areas.
- ❖ **Soil Alkalinity:** No alkaline area in the microwatershed. Soils that are slightly to very strongly alkaline in these area need application of gypsum and wherever calcium is in excess, iron pyrites and element sulphur can be recommended. Management practices like treating repeatedly with good quality water to drain out the excess salts and provision of subsurface drainage and growing of salt tolerant crops like Casuarina, Acacia, Neem, Ber etc, are recommended.

Land Suitability for various crops: Areas that are highly, moderately and marginally suitable and not suitable for growing various crops are indicated. Along with the suitability, various constraints that are limiting the productivity are also indicated. For example, in case of cotton, gravel content, rooting depth and salinity/alkalinity are the major constraints in various plots. With suitable management interventions, the productivity can be enhanced. In order to increase the water holding capacity of light textured soils, growing of green manure crops and application of organic manure is recommended.

SOIL AND WATER CONSERVATION TREATMENT PLAN

For preparing soil and water conservation treatment plan for Yadgir RF1 microwatershed, the land resource inventory database generated under Sujala-III project has been transformed as information through series of interpretative (thematic) maps using soil phase map as a base. The various thematic maps (1:7920 scale) generated were

- Soil depth
- Surface soil texture
- Available water capacity
- Soil slope
- Soil gravelliness
- Land capability
- Present land use and land cover
- Crop suitability
- Rainfall
- Hydrology
- Water Resources
- Socio-economic data
- Contour plan with existing features- network of waterways, pottissa boundaries, cut up/ minor terraces etc.
- Cadastral map (1:7920 scale)
- Satellite imagery (1:7920 scale)



Apart from these, Hand Level/ Hydro Marker/ Dumpy Level/ Total Station and Kathedars' List to be collected.

Steps for Survey and Preparation of Treatment Plan

The boundaries of Land User Groups' and Survey No. boundaries are traced in the field.

- Naming of user groups and farmers
- Identification of arable and non arable lands
- Identification of drainage lines and gullies
- Identification of non treatable areas
- Identification of priority areas in the arable lands
- Treatment plan for arable lands
- Location of water harvesting and recharge structures

9.1 Treatment Plan

The treatment plan recommended for arable lands is briefly described below


9.1.1 Arable Land Treatment

A. BUNDING

Steps for Survey and Preparation of Treatment Plan		USER GROUP-1 CLASSIFICATION OF GULLIES
<ul style="list-style-type: none"> • Cadastral map (1:7920 scale) is enlarged to a scale of 1:2500 scale • Existing network of waterways, pothissa boundaries, grass belts, natural drainage lines/ watercourse, cut ups/ terraces are marked on the cadastral map to the scale • Drainage lines are demarcated into 		
Small gullies	(up to 5 ha catchment)	
Medium gullies	(5-15 ha catchment)	
Ravines	(15-25 ha catchment) and	
<i>Halla/Nala</i>	(more than 25ha catchment)	

Measurement of Land Slope

Land slope is estimated or determined by the study and interpretation of contours or by measurement in the field using simple instruments like Hand Level or Hydromarker.



Vertical and Horizontal intervals between bunds as recommended by the Watershed Development Department.

Slope percentage	Vertical interval (m)	Corresponding Horizontal Distance (m)
2 - 3%	0.6	24
3 - 4%	0.9	21
4 - 5%	0.9	21
5 - 6%	1.2	21
6 - 7%	1.2	21

Note: (i) The above intervals are maximum.

(ii) Considering the slope class and erosion status (A1... A=0-1 % slope, 1= slight erosion) the intervals have to be decided.

Bund length recording: Considering the contour plan and the existing grass belts/partitions, the bunds are aligned and lengths are measured.

Section of the Bund

Bund section is decided considering the soil texture class and gravelliness class (bg₀... b=loamy sand, g₀ = <15% gravel). The recommended Sections for different soils are given below.

Recommended Bund Section

Top width (m)	Base width (m)	Height (m)	Side slope (Z:1;H:V)	Cross section (sq m)	Soil Texture	Remarks
0.3	0.9	0.3	01:01	0.18	Sandy loam	Vegetative bund
0.3	1.2	0.3	1.5:1	0.225	Sandy clay	
0.3	1.2	0.5	0.9:1	0.375	Red gravelly soils	
0.3	1.2	0.6	0.75:1	0.45		
0.3	1.5	0.6	01:01	0.54	Red sandy loam	
0.3	2.1	0.6	1.5:1	0.72	Very shallow black clayey soils	
0.45	2	0.75	01:01	0.92		
0.45	2.4	0.75	1.3:1	1.07	Shallow black clayey soils	
0.6	3.1	0.7	1.78:1	1.29	Medium black clayey soils	
0.5	3	0.85	1.47:1	1.49		

Formation of Trench cum Bund

Dimensions of the Borrow Pits/Trenches to be excavated (machinery are decided considering the Bund Section).

Details of Borrow Pit dimensions are given below:

TRENCH CUM BUND

'A' FRAME FOR INTERBUND MANAGEMENT

Size of Borrow Pits/ Trench recommended for Trench cum Bund (by machinery)

Bund section	Bund length	Earth quantity	Pit				Berm (pit to pit)	Soil depth class
			L(m)	W(m)	D(m)	Quantity (m ³)		
m ²	m	m ³					m	
0.375	6	2.25	5.85	0.85	0.45	2.24	0.15	Shallow
0.45	6	2.7	5.4	1.2	0.43	2.79	0.6	Shallow
0.45	6	2.7	5	0.85	0.65	2.76	1	Moderately Shallow
0.54	5.6	3.02	5.5	0.85	0.7	3.27	0.1	Moderately shallow
0.54	5.5	2.97	5	1.2	0.5	3	0.5	Shallow
0.72	6.2	4.46	6	1.2	0.7	5.04	0.2	Moderately shallow
0.72	5.2	3.74	5.1	0.85	0.9	3.9	0.1	Moderately deep

B. Water Ways

- Existing waterways are marked on the cadastral map (1:7920 scale) and their dimensions are recorded.
- Considering the contour plan of the MWS, additional waterways/ modernization of the existing ones can be thought of.
- The design details are given in the Manual.

C. Farm Ponds

Waterways and the catchment area will give an indication on the size of the Farm Pond. Location of the pond can be decided based on the contour plan/ field condition and farmers' need/desire.

D. Diversion Channel

Existing EPT/ CPT are marked on the cadastral map. Looking to the need, these can be modernized or fresh diversion channel can be proposed and runoff from this can be stored in *Gokatte*/ Recharge ponds.

9.1.2 Non-Arable Land Treatment

Depending on the gravelliness and crops preferred by the farmers, the concerned authorities can decide appropriate treatment plan. The recommended treatments may be Contour Trench, Staggered Trench, Crescent Bund, Boulder Bund or Pebble Bund.

9.1.3 Treatment of Natural Water Course/ Drainage Lines

- a) The cadastral map has to be updated as regards the network of drainage lines (gullies/*nalas/hallas*) and existing structures are marked to the scale and storage capacity of the existing water bodies are documented.
- b) The drainage line will be demarcated into Upper Reach, Middle Reach and Lower Reach.
- c) Considering the Catchment, *Nala* bed and bank conditions, suitable structures are decided.
- d) Number of storage structures (Check dam/*Nala* bund/ Percolation tank) will be decided considering the commitments and available runoff from water budgeting and quality of water in the wells and site suitability.
- e) Detailed Leveling Survey using Dumpy Level / Total Station has to be carried out to arrive at the site-specific designs as shown in the Manual.
- f) The location of ground water recharge structures are decided by examining the lineaments and fracture zones from geological maps.
- g) Rainfall intensity data of the nearest Rain Gauge Station is considered for Hydrologic Designs.
- h) Silt load to the Storage/Recharge Structures is reduced by providing vegetative, boulder and earthen checks in the natural water course. Location and design details are given in the Manual.

9.2 Recommended Soil and Water Conservation Measures

The appropriate conservation structures best suited for each of the land parcel/ survey number (Appendix-I) are selected based on the slope per cent, severity of erosion, amount of rainfall, land use and soil type. The different kinds of conservation structures recommended are:

1. Graded / Strengthening of Bunds
2. Trench cum Bunds (TCB)
3. Trench cum Bunds / Strengthening
4. Crescent Bunds

A map (Fig. 9.1) showing soil and water conservation plan with different kinds of structures recommended has been prepared which shows the spatial distribution and extent of area. An area of about 20 ha (4%) needs Trench Cum Bunding. Maximum area of about 465 ha (95%) needs Graded Bunding.

The conservation plan prepared may be presented to all the stakeholders including farmers and after considering their suggestions, the conservation plan for the microwatershed may be finalised in a participatory approach.




Fig. 9.1 Soil and Water Conservation Plan map of Yadgir RF1 Microwatershed

9.3 Greening of Microwatershed

As part of the greening programme in the watersheds, it is envisaged to plant a variety of horticultural and other tree plants that are edible, economical and produce lot of biomass which helps to restore the ecological balance in the watersheds. The lands that are suitable for greening programme are non-arable lands (land capability classes V, VI VII and VIII) and also the lands that are not suitable or marginally suitable and field bunds for growing annual and perennial crops. The method of planting these trees is given below.

It is recommended to open pits during the 1st week of March along the contour and heap the dugout soil on the lower side of the slope in order to harness the flowing water and facilitate weathering of soil in the pit. Exposure of soil in the pit also prevents spread of pests and diseases due to scorching sun rays. The pits should be filled with mixture of soil and organic manure during the second week of April and keep ready with sufficiently tall seedlings produced either in poly bags or in root trainer nurseries so that planting can be done during the 2nd or 3rd week of April depending on the rainfall.

The tree species suitable for the area considering rainfall, temperature and adaptability is listed below; waterlogged areas are recommended to be planted with species like Nerale (*Syzgium cumini*) and Bamboo. Dry areas are to be planted with species like Honge, Bevu, Seetaphal etc.

Dry Deciduous Species			Temp (°C)	Rainfall (mm)
1.	Bevu	<i>Azadiracta indica</i>	21–32	400 –1,200
2.	Tapasi	<i>Holoptelia integrifolia</i>	20-30	500 - 1000
3.	Seetaphal	<i>Anona Squamosa</i>	20-40	400 - 1000
4.	Honge	<i>Pongamia pinnata</i>	20 -50	500– 2,500
5.	Kamara	<i>Hardwickia binata</i>	25 -35	400 - 1000
6.	Bage	<i>Albezzia lebbek</i>	20 - 45	500 - 1000
7.	Ficus	<i>Ficus bengalensis</i>	20 - 50	500–2,500
8.	Sisso	<i>Dalbargia Sissoo</i>	20 - 50	500 -2000
9.	Ailanthus	<i>Ailanthus excelsa</i>	20 - 50	500 - 1000
10.	Hale	<i>Wrightia tinctoria</i>	25 - 45	500 - 1000
11.	Uded	<i>Steriospermum chelanoides</i>	25 - 45	500 -2000
12.	Dhupa	<i>Boswella Serrata</i>	20 - 40	500 - 2000
13.	Nelli	<i>Embllica Officinalis</i>	20 - 50	500 -1500
14.	Honne	<i>Pterocarpus marsupium</i>	20 - 40	500 - 2000
Moist Deciduous Species			Temp (°C)	Rainfall (mm)
15.	Teak	<i>Tectona grandis</i>	20 - 50	500-5000
16.	Nandi	<i>Legarstroemia lanceolata</i>	20 - 40	500 - 4000
17.	Honne	<i>Pterocarpus marsupium</i>	20 - 40	500 - 3000
18.	Mathi	<i>Terminalia alata</i>	20 -50	500 - 2000
19.	Shivane	<i>Gmelina arborea</i>	20 -50	500 -2000
20.	Kindal	<i>T.Paniculata</i>	20 - 40	500 - 1500
21.	Beete	<i>Dalbargia latifolia</i>	20 - 40	500 - 1500
22.	Tare	<i>T. belerica</i>	20 - 40	500 - 2000
23.	Bamboo	<i>Bambusa arundinasia</i>	20 - 40	500 - 2500
24.	Bamboo	<i>Dendrocalamus strictus</i>	20 – 40	500 – 2500
25.	Muthuga	<i>Butea monosperma</i>	20 - 40	400 - 1500
26.	Hippe	<i>Madhuca latifolia</i>	20 - 40	500 - 2000
27.	Sandal	<i>Santalum album</i>	20 - 50	400 - 1000
28.	Nelli	<i>Embllica officinalis</i>	20 - 40	500 - 2000
29.	Nerale	<i>Sizygium cumini</i>	20 - 40	500 - 2000
30.	Dhaman	<i>Grevia tilifolia</i>	20 - 40	500 - 2000
31.	Kaval	<i>Careya arborea</i>	20 - 40	500 - 2000
32.	Harada	<i>Terminalia chebula</i>	20 - 40	500 - 2000

References

1. FAO (1976) Framework for Land Evaluation, Food and Agriculture Organization, Rome. 72 pp.
2. FAO (1983) Guidelines for Land Evaluation for Rainfed Agriculture, FAO, Rome, 237 pp.
3. IARI (1971) Soil Survey Manual, All India Soil and Land Use Survey Organization, IARI, New Delhi, 121 pp.
4. Katyal, J.C. and Rattan, R.K. (2003) Secondary and Micronutrients; Research Gap and Future Needs. Fert. News 48 (4); 9-20.
5. Naidu, L.G.K., Ramamurthy, V., Challa, O., Hegde, R. and Krishnan, P. (2006) Manual Soil Site Suitability Criteria for Major Crops, NBSS Publ. No. 129, NBSS & LUP, Nagpur, 118 pp.
6. Natarajan, A. and Dipak Sarkar (2010) Field Guide for Soil Survey, National Bureau of Soil Survey and Land Use Planning (ICAR), Nagpur, India.
7. Natarajan, A., Rajendra Hegde, Raj, J.N. and Shivananda Murthy, H.G. (2015) Implementation Manual for Sujala-III Project, Watershed Development Department, Bengaluru, Karnataka.
8. Sarma, V.A.K., Krishnan, P. and Budihal, S.L. (1987) Laboratory Manual, Tech. Bull. 23, NBSS & LUP, Nagpur.
9. Sehgal, J.L. (1990) Soil Resource Mapping of Different States of India; Why and How?, National Bureau of Soil Survey and Land Use Planning, Nagpur, 49 pp.
10. Shivaprasad, C.R., R.S. Reddy, J. Sehgal and M. Velayuthum (1998) Soils of Karnataka for Optimizing Land Use, NBSS Publ. No. 47b, NBSS & LUP, Nagpur, India.
11. Soil Survey Staff (2006) Keys to Soil Taxonomy, Tenth edition, U.S. Department of Agriculture/ NRCS, Washington DC, U.S.A.
12. Soil Survey Staff (2012) Soil Survey Manual, Handbook No. 18, USDA, Washington DC, USA.

Appendix-I
Yadgir Rf-1 (1F2a) Microwatershed
Soil Phase Information

Village	SY NO	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Gajarakota	685	0.49	KKRbB2g1	LMU-3	Very shallow (<25 cm)	Loamy sand	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Scrubland (Sl)	Not Available	IVes	Graded bunding
Gajarakota	686	2.8	KKRbB2g1	LMU-3	Very shallow (<25 cm)	Loamy sand	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IVes	Graded bunding
Gajarakota	687	3.71	KKRbB2g1	LMU-3	Very shallow (<25 cm)	Loamy sand	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IVes	Graded bunding
Gajarakota	688	0.59	KKRbB2g1	LMU-3	Very shallow (<25 cm)	Loamy sand	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IVes	Graded bunding
Gajarakota	689	8.98	KKRbB2g1	LMU-3	Very shallow (<25 cm)	Loamy sand	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut+Redgram (Gn+Rg)	Not Available	IVes	Graded bunding
Gajarakota	690	8.12	BDLbB2	LMU-3	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	1 Borewell	IIes	Graded bunding
Gajarakota	691	43.44	KKRbB2g1	LMU-3	Very shallow (<25 cm)	Loamy sand	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Hill+Redgram (Hill+Rg)	Not Available	IVes	Graded bunding
Gajarakota	692	5.59	KKRbB2g1	LMU-3	Very shallow (<25 cm)	Loamy sand	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IVes	Graded bunding
Gajarakota	693	6.02	KKRbB2g1	LMU-3	Very shallow (<25 cm)	Loamy sand	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Scrubland (Sl)	Not Available	IVes	Graded bunding
Gajarakota	694	9.2	KKRbB2g1	LMU-3	Very shallow (<25 cm)	Loamy sand	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut (Gn)	Not Available	IVes	Graded bunding
Gajarakota	695	4.17	BDLbB2	LMU-3	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Gajarakota	696	4.44	GWDcB2	LMU-1	Moderately deep (75-100 cm)	Sandy loam	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Gajarakota	697	4.16	GWDcB2	LMU-1	Moderately deep (75-100 cm)	Sandy loam	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Gajarakota	698	2.41	KKRbB2g1	LMU-3	Very shallow (<25 cm)	Loamy sand	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IVes	Graded bunding
Gajarakota	699	8.39	KKRbB2g1	LMU-3	Very shallow (<25 cm)	Loamy sand	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IVes	Graded bunding
Gajarakota	700	5.68	KKRbB2g1	LMU-3	Very shallow (<25 cm)	Loamy sand	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IVes	Graded bunding
Gajarakota	703	2.92	GWDcB2	LMU-1	Moderately deep (75-100 cm)	Sandy loam	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Gajarakota	704	5.28	GWDcB2	LMU-1	Moderately deep (75-100 cm)	Sandy loam	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Gajarakota	705	3.76	GWDcB2	LMU-1	Moderately deep (75-100 cm)	Sandy loam	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Gajarakota	706	2.96	KKRbB2g1	LMU-3	Very shallow (<25 cm)	Loamy sand	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IVes	Graded bunding
Gajarakota	707	4.15	KKRbB2g1	LMU-3	Very shallow (<25 cm)	Loamy sand	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut (Gn)	Not Available	IVes	Graded bunding
Gajarakota	708	1.45	BDPhB2	LMU-3	Very shallow (<25 cm)	Sandy clay loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IVes	Trench cum bunding

Village	SY NO	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Gajarakota	709	0.03	KKRbB2g1	LMU-3	Very shallow (<25 cm)	Loamy sand	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut+Redgram (Gn+Rg)	Not Available	Ives	Graded bunding
Gajarakota	814	0.25	KKRbB2g1	LMU-3	Very shallow (<25 cm)	Loamy sand	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Ives	Graded bunding
Yampada	145	0.02	KKRbB2g1	LMU-3	Very shallow (<25 cm)	Loamy sand	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Ives	Graded bunding
Yampada	155	0.48	HTKbB2g1	LMU-3	Shallow (25-50 cm)	Loamy sand	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Yampada	158	1.24	KKRbB2g1	LMU-3	Very shallow (<25 cm)	Loamy sand	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut (Gn)	Not Available	Ives	Graded bunding
Yampada	159	0.49	KKRbB2g1	LMU-3	Very shallow (<25 cm)	Loamy sand	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	Ives	Graded bunding
Yampada	160	4.58	KKRbB2g1	LMU-3	Very shallow (<25 cm)	Loamy sand	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut (Gn)	Not Available	Ives	Graded bunding
Yampada	161	0.32	KKRbB2g1	LMU-3	Very shallow (<25 cm)	Loamy sand	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	Ives	Graded bunding
Yampada	171	0.29	KKRcB2	LMU-3	Very shallow (<25 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Ives	Graded bunding
Yampada	173	1.84	HTKbB2g1	LMU-3	Shallow (25-50 cm)	Loamy sand	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut (Gn)	Not Available	IIes	Graded bunding
Yampada	174	8.04	HTKbB2g1	LMU-3	Shallow (25-50 cm)	Loamy sand	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut (Gn)	Not Available	IIes	Graded bunding
Yampada	175	6.68	HTKbB2g1	LMU-3	Shallow (25-50 cm)	Loamy sand	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar+Redgram (Jw+Rg)	Not Available	IIes	Graded bunding
Yampada	176	0.32	HTKbB2g1	LMU-3	Shallow (25-50 cm)	Loamy sand	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	IIes	Graded bunding
Yampada	177	4.14	HTKbB2g1	LMU-3	Shallow (25-50 cm)	Loamy sand	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut (Gn)	Not Available	IIes	Graded bunding
Yampada	178	3.69	HTKbB2g1	LMU-3	Shallow (25-50 cm)	Loamy sand	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut (Gn)	1 Borewell	IIes	Graded bunding
Yampada	179	4.42	GWDmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut+Redgram (Gn+Rg)	Not Available	Iles	Graded bunding
Yampada	180	0.2	GWDmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut+Redgram (Gn+Rg)	Not Available	Iles	Graded bunding
Yampada	181	2.87	HTKbB2g1	LMU-3	Shallow (25-50 cm)	Loamy sand	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut+Redgram (Gn+Rg)	Not Available	IIes	Graded bunding
Yampada	202	0.02	SBRcB2	LMU-2	Moderately shallow (50-75 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IVs	Graded bunding
Yampada	209	1.01	BDLhB2	LMU-3	Shallow (25-50 cm)	Sandy clay loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIes	Graded bunding
Yampada	210	0.99	SBRcB2	LMU-2	Moderately shallow (50-75 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IVs	Graded bunding
Yampada	211	0.73	SBRcB2	LMU-2	Moderately shallow (50-75 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IVs	Graded bunding
Yampada	212	0.74	SBRcB2	LMU-2	Moderately shallow (50-75 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IVs	Graded bunding
Yampada	213	1.26	SBRcB2	LMU-2	Moderately shallow (50-75 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut (Gn)	Not Available	IVs	Graded bunding

Village	SY NO	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Yampada	214	4.83	BDLhB2	LMU-3	Shallow (25-50 cm)	Sandy clay loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut (Gn)	Not Available	IIes	Graded bunding
Yampada	216	3.08	GWDmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Pad dy (Rg+Pd)	Not Available	Iles	Graded bunding
Yampada	217	2.24	GWDmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Yampada	218	0.75	SBRcB2	LMU-2	Moderately shallow (50-75 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut (Gn)	Not Available	IVs	Graded bunding
Yampada	219	0.18	SBRcB2	LMU-2	Moderately shallow (50-75 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut (Gn)	Not Available	IVs	Graded bunding
Yampada	220	1.06	SBRcB2	LMU-2	Moderately shallow (50-75 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut (Gn)	Not Available	IVs	Graded bunding
Yampada	221	1.67	SBRcB2	LMU-2	Moderately shallow (50-75 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut (Gn)	Not Available	IVs	Graded bunding
Yampada	222	6.95	HTKbB2g1	LMU-3	Shallow (25-50 cm)	Loamy sand	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut+Jowar (Gn+Jw)	1 Openwell	IIes	Graded bunding
Yampada	223	3.69	HTKbB2g1	LMU-3	Shallow (25-50 cm)	Loamy sand	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut (Gn)	Not Available	IIes	Graded bunding
Yampada	247	3.8	HTKbB2g1	LMU-3	Shallow (25-50 cm)	Loamy sand	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut (Gn)	Not Available	IIes	Graded bunding
Yampada	248	5.26	HTKbB2g1	LMU-3	Shallow (25-50 cm)	Loamy sand	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut (Gn)	Not Available	IIes	Graded bunding
Yampada	249	0.14	HTKbB2g1	LMU-3	Shallow (25-50 cm)	Loamy sand	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	IIes	Graded bunding
Yampada	258	7.57	SBRcB2	LMU-2	Moderately shallow (50-75 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut (Gn)	Not Available	IVs	Graded bunding
Yampada	259	5.92	GWDmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut (Gn)	Not Available	Iles	Graded bunding
Yampada	260	5.56	GWDmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	Iles	Graded bunding
Yampada	261	4.52	SBRcC3g1	LMU-2	Moderately shallow (50-75 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Redgram (Rg)	Not Available	IVes	Graded bunding
Yampada	262	0.81	SBRcC3g1	LMU-2	Moderately shallow (50-75 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Groundnut (Gn)	Not Available	IVes	Graded bunding
Yampada	263	0.81	GWDmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut (Gn)	Not Available	Iles	Graded bunding
Yampada	264	2.93	GWDmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut (Gn)	Not Available	Iles	Graded bunding
Yampada	265	0.52	GWDmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut (Gn)	Not Available	Iles	Graded bunding
Yampada	266	0.88	GWDmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut (Gn)	Not Available	Iles	Graded bunding
Yampada	267	1.2	SBRcC3g1	LMU-2	Moderately shallow (50-75 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Groundnut (Gn)	Not Available	IVes	Graded bunding
Yampada	268	0.88	SBRcC3g1	LMU-2	Moderately shallow (50-75 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Groundnut (Gn)	Not Available	IVes	Graded bunding
Yampada	269	0.45	SBRcC3g1	LMU-2	Moderately shallow (50-75 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Redgram (Rg)	Not Available	IVes	Graded bunding

Village	SY NO	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Yampada	270	5.48	SBRcC3g1	LMU-2	Moderately shallow (50-75 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Groundnut (Gn)	Not Available	IVes	Graded bunding
Yampada	271	3.43	HTKcC2g1	LMU-3	Shallow (25-50 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Moderate	Groundnut (Gn)	Not Available	IIes	Graded bunding
Yampada	272	5.28	HTKcC2g1	LMU-3	Shallow (25-50 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Moderate	Jowar (Jw)	Not Available	IIes	Graded bunding
Yampada	273	0.19	SBRcC3g1	LMU-2	Moderately shallow (50-75 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Redgram (Rg)	Not Available	IVes	Graded bunding
Yampada	274	1.41	KKRbB2g1	LMU-3	Very shallow (<25 cm)	Loamy sand	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut (Gn)	Not Available	IVes	Graded bunding
Yampada	275	1.25	KKRbB2g1	LMU-3	Very shallow (<25 cm)	Loamy sand	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut (Gn)	Not Available	IVes	Graded bunding
Yampada	276	7.07	KKRbB2g1	LMU-3	Very shallow (<25 cm)	Loamy sand	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut+Redgram (Gn+Rg)	Not Available	IVes	Graded bunding
Yampada	277	1.17	KKRbB2g1	LMU-3	Very shallow (<25 cm)	Loamy sand	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut (Gn)	Not Available	IVes	Graded bunding
Yampada	278	0.6	HTKcC2g1	LMU-3	Shallow (25-50 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Moderate	Groundnut (Gn)	Not Available	IIes	Graded bunding
Yampada	279	8.54	HTKcC2g1	LMU-3	Shallow (25-50 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Moderate	Groundnut (Gn)	Not Available	IIes	Graded bunding
Yampada	280	0.57	KKRbB2g1	LMU-3	Very shallow (<25 cm)	Loamy sand	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IVes	Graded bunding
Yampada	281	3.44	KKRbB2g1	LMU-3	Very shallow (<25 cm)	Loamy sand	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IVes	Graded bunding
Yampada	286	0.96	KKRbB2g1	LMU-3	Very shallow (<25 cm)	Loamy sand	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut (Gn)	Not Available	IVes	Graded bunding
Yampada	287	1.22	SBRcC3g1	LMU-2	Moderately shallow (50-75 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Groundnut+Redgram (Gn+Rg)	Not Available	IVes	Graded bunding
Yampada	289	3.88	HTKcC2g1	LMU-3	Shallow (25-50 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Moderate	Groundnut (Gn)	Not Available	IIes	Graded bunding
Yampada	292	1.42	KKRbB2g1	LMU-3	Very shallow (<25 cm)	Loamy sand	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IVes	Graded bunding
Yampada	293	4.3	HTKcC2g1	LMU-3	Shallow (25-50 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Moderate	Redgram+Scrubland (Rg+Sl)	Not Available	IIes	Graded bunding
Yampada	294	1.92	KKRbB2g1	LMU-3	Very shallow (<25 cm)	Loamy sand	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IVes	Graded bunding
Yampada	295	5.99	HTKcC2g1	LMU-3	Shallow (25-50 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Moderate	Groundnut (Gn)	Not Available	IIes	Graded bunding
Yampada	296	1.07	SBRcC3g1	LMU-2	Moderately shallow (50-75 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Groundnut (Gn)	Not Available	IVes	Graded bunding
Yampada	297	6.28	SBRcC3g1	LMU-2	Moderately shallow (50-75 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Groundnut (Gn)	Not Available	IVes	Graded bunding
Yampada	298	3.12	SBRcC3g1	LMU-2	Moderately shallow (50-75 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Groundnut (Gn)	Not Available	IVes	Graded bunding
Yampada	299	0.97	SBRcC3g1	LMU-2	Moderately shallow (50-75 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Not Available (NA)	Not Available	IVes	Graded bunding
Yampada	300	2.85	SBRcC3g1	LMU-2	Moderately shallow (50-75 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Redgram (Rg)	Not Available	IVes	Graded bunding

Village	SY NO	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Yampada	301	3.12	SBRcC3g1	LMU-2	Moderately shallow (50-75 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Not Available (NA)	Not Available	Ives	Graded bunding
Yampada	302	1.78	SBRcC3g1	LMU-2	Moderately shallow (50-75 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Scrubland (Sl)	Not Available	Ives	Graded bunding
Yampada	303	0.72	SBRcC3g1	LMU-2	Moderately shallow (50-75 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Not Available (NA)	Not Available	Ives	Graded bunding
Yampada	304	0.88	SBRcC3g1	LMU-2	Moderately shallow (50-75 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Not Available (NA)	Not Available	Ives	Graded bunding
Yampada	305	1.03	Habitation	Others	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others
Yampada	306	0.17	Habitation	Others	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others
Yampada	307	0.55	HTKcC2g1	LMU-3	Shallow (25-50 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Moderate	Habitation	Not Available	IIes	Graded bunding
Yampada	308	1.24	SBRcC3g1	LMU-2	Moderately shallow (50-75 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Not Available (NA)	Not Available	Ives	Graded bunding
Yampada	309	0.59	SBRcC3g1	LMU-2	Moderately shallow (50-75 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Not Available (NA)	Not Available	Ives	Graded bunding
Yampada	310	1.74	SBRcC3g1	LMU-2	Moderately shallow (50-75 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Not Available (NA)	Not Available	Ives	Graded bunding
Yampada	311	1.49	SBRcC3g1	LMU-2	Moderately shallow (50-75 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Not Available (NA)	Not Available	Ives	Graded bunding
Yampada	312	2.22	SBRcC3g1	LMU-2	Moderately shallow (50-75 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Not Available (NA)	Not Available	Ives	Graded bunding
Yampada	313	1.35	SBRcC3g1	LMU-2	Moderately shallow (50-75 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Not Available (NA)	Not Available	Ives	Graded bunding
Yampada	314	2.51	SBRcC3g1	LMU-2	Moderately shallow (50-75 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Groundnut (Gn)	Not Available	Ives	Graded bunding
Yampada	315	0.57	HTKcC2g1	LMU-3	Shallow (25-50 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Moderate	Groundnut (Gn)	1 Borewell	IIes	Graded bunding
Yampada	316	1.08	HTKcC2g1	LMU-3	Shallow (25-50 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Moderate	Groundnut (Gn)	Not Available	IIes	Graded bunding
Yampada	317	0.47	HTKcC2g1	LMU-3	Shallow (25-50 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Moderate	Groundnut (Gn)	Not Available	IIes	Graded bunding
Yampada	318	0.16	HTKcC2g1	LMU-3	Shallow (25-50 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Yampada	319	0.65	HTKcC2g1	LMU-3	Shallow (25-50 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Yampada	320	0.77	HTKcC2g1	LMU-3	Shallow (25-50 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Yampada	321	0.78	HTKcC2g1	LMU-3	Shallow (25-50 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Yampada	322	0.69	HTKcC2g1	LMU-3	Shallow (25-50 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Yampada	323	0.25	HTKcC2g1	LMU-3	Shallow (25-50 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Yampada	324	5.91	HTKcC2g1	LMU-3	Shallow (25-50 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding

Village	SY NO	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Yampada	325	6.79	HTKbB2g1	LMU-3	Shallow (25-50 cm)	Loamy sand	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Yampada	326	6.68	HTKcC2g1	LMU-3	Shallow (25-50 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Moderate	Groundnut (Gn)	Not Available	IIes	Graded bunding
Yampada	327	0.55	HTKcC2g1	LMU-3	Shallow (25-50 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Moderate	Groundnut (Gn)	Not Available	IIes	Graded bunding
Yampada	328	6.11	HTKcC2g1	LMU-3	Shallow (25-50 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Yampada	329	6.54	HTKcC2g1	LMU-3	Shallow (25-50 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Moderate	Groundnut (Gn)	Not Available	IIes	Graded bunding
Yampada	330	3.72	HTKcC2g1	LMU-3	Shallow (25-50 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Moderate	Groundnut (Gn)	Not Available	IIes	Graded bunding
Yampada	331	3.72	KKRbB2g1	LMU-3	Very shallow (<25 cm)	Loamy sand	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut (Gn)	Not Available	IVes	Graded bunding
Yampada	332	7.89	HTKcC2g1	LMU-3	Shallow (25-50 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Moderate	Redgram (Rg)	1 Borewell	IIes	Graded bunding
Yampada	333	4.8	HTKcC2g1	LMU-3	Shallow (25-50 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Moderate	Groundnut (Gn)	Not Available	IIes	Graded bunding
Yampada	334	3.53	HTKcC2g1	LMU-3	Shallow (25-50 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Moderate	Scrubland (Sl)	Not Available	IIes	Graded bunding
Yampada	335	0.96	HTKcC2g1	LMU-3	Shallow (25-50 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Moderate	Scrubland (Sl)	Not Available	IIes	Graded bunding
Yampada	336	0.71	HTKcC2g1	LMU-3	Shallow (25-50 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Moderate	Scrubland (Sl)	Not Available	IIes	Graded bunding
Yampada	337	4.75	HTKcC2g1	LMU-3	Shallow (25-50 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Moderate	Groundnut (Gn)	Not Available	IIes	Graded bunding
Yampada	338	3.16	HTKcC2g1	LMU-3	Shallow (25-50 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Moderate	Groundnut (Gn)	Not Available	IIes	Graded bunding
Yampada	339	0.55	HTKcC2g1	LMU-3	Shallow (25-50 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Moderate	Groundnut (Gn)	Not Available	IIes	Graded bunding
Yampada	342	5.9	KKRbB2g1	LMU-3	Very shallow (<25 cm)	Loamy sand	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut (Gn)	Not Available	IVes	Graded bunding
Yampada	344	7.12	HTKcC2g1	LMU-3	Shallow (25-50 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Moderate	Groundnut (Gn)	Not Available	IIes	Graded bunding
Yampada	345	3.86	HTKcC2g1	LMU-3	Shallow (25-50 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Moderate	Groundnut (Gn)	Not Available	IIes	Graded bunding
Yampada	346	5.74	HTKcC2g1	LMU-3	Shallow (25-50 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Moderate	Groundnut (Gn)	Not Available	IIes	Graded bunding
Yampada	347	0.29	HTKcC2g1	LMU-3	Shallow (25-50 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Yampada	348	5.34	HTKcC2g1	LMU-3	Shallow (25-50 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Yampada	349	0.85	HTKcC2g1	LMU-3	Shallow (25-50 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Moderate	Groundnut (Gn)	Not Available	IIes	Graded bunding
Yampada	350	0.17	HTKcC2g1	LMU-3	Shallow (25-50 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Moderate	Groundnut (Gn)	Not Available	IIes	Graded bunding
Yampada	351	0.63	HTKcC2g1	LMU-3	Shallow (25-50 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Moderate	Groundnut (Gn)	Not Available	IIes	Graded bunding

Village	SY NO	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Yampada	352	1.18	HTKcC2g1	LMU-3	Shallow (25-50 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Moderate	Groundnut (Gn)	Not Available	IIIes	Graded bunding
Yampada	353	0.98	HTKcC2g1	LMU-3	Shallow (25-50 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Moderate	Groundnut (Gn)	Not Available	IIIes	Graded bunding
Yampada	354	6.9	HTKcC2g1	LMU-3	Shallow (25-50 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Moderate	Redgram (Rg)	Not Available	IIIes	Graded bunding
Yampada	355	3.33	HTKcC2g1	LMU-3	Shallow (25-50 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Moderate	Groundnut (Gn)	Not Available	IIIes	Graded bunding
Yampada	356	2.19	KKRbB2g1	LMU-3	Very shallow (<25 cm)	Loamy sand	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IVes	Graded bunding
Yampada	357	2.77	BDLhB2	LMU-3	Shallow (25-50 cm)	Sandy clay loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Scrubland (Sl)	Not Available	IIIes	Graded bunding
Yampada	358	3.79	KKRbB2g1	LMU-3	Very shallow (<25 cm)	Loamy sand	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Scrubland (Sl)	Not Available	IVes	Graded bunding
Yampada	359	2.89	HTKcC2g1	LMU-3	Shallow (25-50 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Moderate	Redgram (Rg)	Not Available	IIIes	Graded bunding
Yampada	360	0.36	HTKcC2g1	LMU-3	Shallow (25-50 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Moderate	Redgram (Rg)	Not Available	IIIes	Graded bunding
Yampada	361	0.23	HTKcC2g1	LMU-3	Shallow (25-50 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Moderate	Redgram (Rg)	Not Available	IIIes	Graded bunding
Yampada	362	6.42	HTKcC2g1	LMU-3	Shallow (25-50 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Moderate	Redgram (Rg)	Not Available	IIIes	Graded bunding
Yampada	363	0.91	HTKcC2g1	LMU-3	Shallow (25-50 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Moderate	Redgram (Rg)	Not Available	IIIes	Graded bunding
Yampada	364	0.49	Habitation	Others	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others

Village	Survey No	Soil Reaction	Salinity	Organic Carbon	Available Phosphorus	Available Potassium	Available Sulphur	Available Boron	Available Iron	Available Manganese	Available Copper	Available Zinc
Yampada	353	Slightly acid (pH 6.0 - 6.5)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	Medium (10 - 20 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)
Yampada	354	Slightly acid (pH 6.0 - 6.5)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	Medium (10 - 20 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)
Yampada	355	Slightly acid (pH 6.0 - 6.5)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	Medium (10 - 20 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)
Yampada	356	Slightly acid (pH 6.0 - 6.5)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	Medium (10 - 20 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)
Yampada	357	Slightly acid (pH 6.0 - 6.5)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	Medium (10 - 20 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)
Yampada	358	Slightly acid (pH 6.0 - 6.5)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	Medium (10 - 20 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)
Yampada	359	Slightly acid (pH 6.0 - 6.5)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	Medium (10 - 20 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)
Yampada	360	Slightly acid (pH 6.0 - 6.5)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	Medium (10 - 20 ppm)	Medium (0.5 - 1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)
Yampada	361	Slightly acid (pH 6.0 - 6.5)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 - 57 kg/ha)	Low (<145 kg/ha)	Medium (10 - 20 ppm)	Medium (0.5 - 1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)
Yampada	362	Slightly acid (pH 6.0 - 6.5)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	Medium (10 - 20 ppm)	Medium (0.5 - 1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)
Yampada	363	Slightly acid (pH 6.0 - 6.5)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	Medium (10 - 20 ppm)	Medium (0.5 - 1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)
Yampada	364	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others

Appendix III
Yadgir Rf-1 (1F2a) Microwatershed
Soil Suitability Information

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry
Gajarakota	685	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Gajarakota	686	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Gajarakota	687	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Gajarakota	688	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Gajarakota	689	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Gajarakota	690	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt
Gajarakota	691	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Gajarakota	692	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Gajarakota	693	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Gajarakota	694	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Gajarakota	695	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt
Gajarakota	696	S3rz	S2tz	S3tz	S2zw	S3tz	S2rz	S3rz	S2rz	S2zw	S2rz	S2rz	S2zw	S3tz	S1	N1t	S3rz	S2rz	S3tz	S1	S2tz	S3tz	S2tz	S2tz	S2rz	S2zw	S1	S1	S2rz	S3tz
Gajarakota	697	S3rz	S2tz	S3tz	S2zw	S3tz	S2rz	S3rz	S2rz	S2zw	S2rz	S2rz	S2zw	S3tz	S1	N1t	S3rz	S2rz	S3tz	S1	S2tz	S3tz	S2tz	S2tz	S2rz	S2zw	S1	S1	S2rz	S3tz
Gajarakota	698	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Gajarakota	699	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Gajarakota	700	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Gajarakota	703	S3rz	S2tz	S3tz	S2zw	S3tz	S2rz	S3rz	S2rz	S2zw	S2rz	S2rz	S2zw	S3tz	S1	N1t	S3rz	S2rz	S3tz	S1	S2tz	S3tz	S2tz	S2tz	S2rz	S2zw	S1	S1	S2rz	S3tz
Gajarakota	704	S3rz	S2tz	S3tz	S2zw	S3tz	S2rz	S3rz	S2rz	S2zw	S2rz	S2rz	S2zw	S3tz	S1	N1t	S3rz	S2rz	S3tz	S1	S2tz	S3tz	S2tz	S2tz	S2rz	S2zw	S1	S1	S2rz	S3tz
Gajarakota	705	S3rz	S2tz	S3tz	S2zw	S3tz	S2rz	S3rz	S2rz	S2zw	S2rz	S2rz	S2zw	S3tz	S1	N1t	S3rz	S2rz	S3tz	S1	S2tz	S3tz	S2tz	S2tz	S2rz	S2zw	S1	S1	S2rz	S3tz
Gajarakota	706	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Gajarakota	707	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Gajarakota	708	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Gajarakota	709	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Gajarakota	814	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry	
Yampada	145	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	
Yampada	155	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Yampada	158	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Yampada	159	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Yampada	160	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Yampada	161	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Yampada	171	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Yampada	173	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Yampada	174	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Yampada	175	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Yampada	176	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Yampada	177	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Yampada	178	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Yampada	179	S3rz	S3tz	S3tz	S1	S3tz	S2rz	S3rz	S2rz	S1	S2rz	S2rz	S2z	S3tz	S2z	N1t	S3rz	S2rz	S3tz	S1	S2tz	S3tz	S2tz	S2tz	S2rz	S1	S1	S1	S2rz	S3tz	
Yampada	180	S3rz	S3tz	S3tz	S1	S3tz	S2rz	S3rz	S2rz	S1	S2rz	S2rz	S2z	S3tz	S2z	N1t	S3rz	S2rz	S3tz	S1	S2tz	S3tz	S2tz	S2tz	S2rz	S1	S1	S1	S2rz	S3tz	
Yampada	181	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Yampada	202	N1r	S3t	S3rt	S3t	S3rt	N1t	N1r	S3rt	N1t	S3t	S3t	S3t	S3rt	S3t	S3rt	S3rt	S3rt	S3t	S3t	S3t	S3t	S3t	S3t	S3rt	S3t	S3t	S3t	S3t	S3t	
Yampada	209	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	
Yampada	210	N1r	S3t	S3rt	S3t	S3rt	N1t	N1r	S3rt	N1t	S3t	S3t	S3t	S3rt	S3t	S3rt	S3rt	S3rt	S3t	S3t	S3t	S3t	S3t	S3t	S3rt	S3t	S3t	S3t	S3t	S3t	
Yampada	211	N1r	S3t	S3rt	S3t	S3rt	N1t	N1r	S3rt	N1t	S3t	S3t	S3t	S3rt	S3t	S3rt	S3rt	S3rt	S3t	S3t	S3t	S3t	S3t	S3t	S3rt	S3t	S3t	S3t	S3t	S3t	
Yampada	212	N1r	S3t	S3rt	S3t	S3rt	N1t	N1r	S3rt	N1t	S3t	S3t	S3t	S3rt	S3t	S3rt	S3rt	S3rt	S3t	S3t	S3t	S3t	S3t	S3t	S3rt	S3t	S3t	S3t	S3t	S3t	
Yampada	213	N1r	S3t	S3rt	S3t	S3rt	N1t	N1r	S3rt	N1t	S3t	S3t	S3t	S3rt	S3t	S3rt	S3rt	S3rt	S3t	S3t	S3t	S3t	S3t	S3t	S3rt	S3t	S3t	S3t	S3t	S3t	
Yampada	214	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	
Yampada	216	S3rz	S3tz	S3tz	S1	S3tz	S2rz	S3rz	S2rz	S1	S2rz	S2rz	S2z	S3tz	S2z	N1t	S3rz	S2rz	S3tz	S1	S2tz	S3tz	S2tz	S2tz	S2rz	S1	S1	S1	S2rz	S3tz	
Yampada	217	S3rz	S3tz	S3tz	S1	S3tz	S2rz	S3rz	S2rz	S1	S2rz	S2rz	S2z	S3tz	S2z	N1t	S3rz	S2rz	S3tz	S1	S2tz	S3tz	S2tz	S2tz	S2rz	S1	S1	S1	S2rz	S3tz	
Yampada	218	N1r	S3t	S3rt	S3t	S3rt	N1t	N1r	S3rt	N1t	S3t	S3t	S3t	S3rt	S3t	S3rt	S3rt	S3rt	S3t	S3t	S3t	S3t	S3t	S3t	S3rt	S3t	S3t	S3t	S3t	S3t	
Yampada	219	N1r	S3t	S3rt	S3t	S3rt	N1t	N1r	S3rt	N1t	S3t	S3t	S3t	S3rt	S3t	S3rt	S3rt	S3rt	S3t	S3t	S3t	S3t	S3t	S3t	S3rt	S3t	S3t	S3t	S3t	S3t	

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry		
Yampada	220	N1r	S3t	S3rt	S3t	S3rt	N1t	N1r	S3rt	N1t	S3t	S3t	S3t	S3rt	S3t	S3rt	S3rt	S3rt	S3t	S3t	S3t	S3t	S3t	S3t	S3rt	S3t	S3t	S3t	S3t	S3t		
Yampada	221	N1r	S3t	S3rt	S3t	S3rt	N1t	N1r	S3rt	N1t	S3t	S3t	S3t	S3rt	S3t	S3rt	S3rt	S3rt	S3t	S3t	S3t	S3t	S3t	S3t	S3rt	S3t	S3t	S3t	S3t	S3t	S3t	
Yampada	222	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r		
Yampada	223	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r		
Yampada	247	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r		
Yampada	248	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r		
Yampada	249	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r		
Yampada	258	N1r	S3t	S3rt	S3t	S3rt	N1t	N1r	S3rt	N1t	S3t	S3t	S3t	S3rt	S3t	S3rt	S3rt	S3rt	S3t	S3t	S3t	S3t	S3t	S3t	S3rt	S3t	S3t	S3t	S3t	S3t	S3t	
Yampada	259	S3rz	S3tz	S3tz	S1	S3tz	S2rz	S3rz	S2rz	S1	S2rz	S2rz	S2z	S3tz	S2z	N1t	S3rz	S2rz	S3tz	S1	S2tz	S3tz	S2tz	S2tz	S2rz	S1	S1	S1	S1	S2rz	S3tz	
Yampada	260	S3rz	S3tz	S3tz	S1	S3tz	S2rz	S3rz	S2rz	S1	S2rz	S2rz	S2z	S3tz	S2z	N1t	S3rz	S2rz	S3tz	S1	S2tz	S3tz	S2tz	S2tz	S2rz	S1	S1	S1	S1	S2rz	S3tz	
Yampada	261	N1r	S3t	S3rt	S3t	S3rt	N1t	N1r	S3rt	N1t	S3t	S3t	S3t	S3rt	S3t	S3rt	S3rt	S3rt	S3t	S3t	S3t	S3t	S3t	S3t	S3rt	S3t	S3t	S3t	S3t	S3t	S3t	
Yampada	262	N1r	S3t	S3rt	S3t	S3rt	N1t	N1r	S3rt	N1t	S3t	S3t	S3t	S3rt	S3t	S3rt	S3rt	S3rt	S3t	S3t	S3t	S3t	S3t	S3t	S3rt	S3t	S3t	S3t	S3t	S3t	S3t	
Yampada	263	S3rz	S3tz	S3tz	S1	S3tz	S2rz	S3rz	S2rz	S1	S2rz	S2rz	S2z	S3tz	S2z	N1t	S3rz	S2rz	S3tz	S1	S2tz	S3tz	S2tz	S2tz	S2rz	S1	S1	S1	S1	S2rz	S3tz	
Yampada	264	S3rz	S3tz	S3tz	S1	S3tz	S2rz	S3rz	S2rz	S1	S2rz	S2rz	S2z	S3tz	S2z	N1t	S3rz	S2rz	S3tz	S1	S2tz	S3tz	S2tz	S2tz	S2rz	S1	S1	S1	S1	S2rz	S3tz	
Yampada	265	S3rz	S3tz	S3tz	S1	S3tz	S2rz	S3rz	S2rz	S1	S2rz	S2rz	S2z	S3tz	S2z	N1t	S3rz	S2rz	S3tz	S1	S2tz	S3tz	S2tz	S2tz	S2rz	S1	S1	S1	S1	S2rz	S3tz	
Yampada	266	S3rz	S3tz	S3tz	S1	S3tz	S2rz	S3rz	S2rz	S1	S2rz	S2rz	S2z	S3tz	S2z	N1t	S3rz	S2rz	S3tz	S1	S2tz	S3tz	S2tz	S2tz	S2rz	S1	S1	S1	S1	S2rz	S3tz	
Yampada	267	N1r	S3t	S3rt	S3t	S3rt	N1t	N1r	S3rt	N1t	S3t	S3t	S3t	S3rt	S3t	S3rt	S3rt	S3rt	S3t	S3t	S3t	S3t	S3t	S3t	S3rt	S3t	S3t	S3t	S3t	S3t	S3t	
Yampada	268	N1r	S3t	S3rt	S3t	S3rt	N1t	N1r	S3rt	N1t	S3t	S3t	S3t	S3rt	S3t	S3rt	S3rt	S3rt	S3t	S3t	S3t	S3t	S3t	S3t	S3rt	S3t	S3t	S3t	S3t	S3t	S3t	
Yampada	269	N1r	S3t	S3rt	S3t	S3rt	N1t	N1r	S3rt	N1t	S3t	S3t	S3t	S3rt	S3t	S3rt	S3rt	S3rt	S3t	S3t	S3t	S3t	S3t	S3t	S3rt	S3t	S3t	S3t	S3t	S3t	S3t	
Yampada	270	N1r	S3t	S3rt	S3t	S3rt	N1t	N1r	S3rt	N1t	S3t	S3t	S3t	S3rt	S3t	S3rt	S3rt	S3rt	S3t	S3t	S3t	S3t	S3t	S3t	S3rt	S3t	S3t	S3t	S3t	S3t	S3t	
Yampada	271	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r		
Yampada	272	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r		
Yampada	273	N1r	S3t	S3rt	S3t	S3rt	N1t	N1r	S3rt	N1t	S3t	S3t	S3t	S3rt	S3t	S3rt	S3rt	S3rt	S3t	S3t	S3t	S3t	S3t	S3t	S3rt	S3t	S3t	S3t	S3t	S3t	S3t	
Yampada	274	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Yampada	275	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Yampada	276	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Yampada	277	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry	
Yampada	278	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Yampada	279	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Yampada	280	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Yampada	281	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Yampada	286	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Yampada	287	N1r	S3t	S3rt	S3t	S3rt	N1t	N1r	S3rt	N1t	S3t	S3t	S3t	S3rt	S3t	S3rt	S3rt	S3rt	S3t	S3t	S3t	S3t	S3t	S3t	S3rt	S3t	S3t	S3t	S3t	S3t	
Yampada	289	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Yampada	292	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Yampada	293	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Yampada	294	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Yampada	295	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Yampada	296	N1r	S3t	S3rt	S3t	S3rt	N1t	N1r	S3rt	N1t	S3t	S3t	S3t	S3rt	S3t	S3rt	S3rt	S3rt	S3t	S3t	S3t	S3t	S3t	S3t	S3rt	S3t	S3t	S3t	S3t	S3t	
Yampada	297	N1r	S3t	S3rt	S3t	S3rt	N1t	N1r	S3rt	N1t	S3t	S3t	S3t	S3rt	S3t	S3rt	S3rt	S3rt	S3t	S3t	S3t	S3t	S3t	S3t	S3rt	S3t	S3t	S3t	S3t	S3t	
Yampada	298	N1r	S3t	S3rt	S3t	S3rt	N1t	N1r	S3rt	N1t	S3t	S3t	S3t	S3rt	S3t	S3rt	S3rt	S3rt	S3t	S3t	S3t	S3t	S3t	S3t	S3rt	S3t	S3t	S3t	S3t	S3t	
Yampada	299	N1r	S3t	S3rt	S3t	S3rt	N1t	N1r	S3rt	N1t	S3t	S3t	S3t	S3rt	S3t	S3rt	S3rt	S3rt	S3t	S3t	S3t	S3t	S3t	S3t	S3rt	S3t	S3t	S3t	S3t	S3t	
Yampada	300	N1r	S3t	S3rt	S3t	S3rt	N1t	N1r	S3rt	N1t	S3t	S3t	S3t	S3rt	S3t	S3rt	S3rt	S3rt	S3t	S3t	S3t	S3t	S3t	S3t	S3rt	S3t	S3t	S3t	S3t	S3t	
Yampada	301	N1r	S3t	S3rt	S3t	S3rt	N1t	N1r	S3rt	N1t	S3t	S3t	S3t	S3rt	S3t	S3rt	S3rt	S3rt	S3t	S3t	S3t	S3t	S3t	S3t	S3rt	S3t	S3t	S3t	S3t	S3t	
Yampada	302	N1r	S3t	S3rt	S3t	S3rt	N1t	N1r	S3rt	N1t	S3t	S3t	S3t	S3rt	S3t	S3rt	S3rt	S3rt	S3t	S3t	S3t	S3t	S3t	S3t	S3rt	S3t	S3t	S3t	S3t	S3t	
Yampada	303	N1r	S3t	S3rt	S3t	S3rt	N1t	N1r	S3rt	N1t	S3t	S3t	S3t	S3rt	S3t	S3rt	S3rt	S3rt	S3t	S3t	S3t	S3t	S3t	S3t	S3rt	S3t	S3t	S3t	S3t	S3t	
Yampada	304	N1r	S3t	S3rt	S3t	S3rt	N1t	N1r	S3rt	N1t	S3t	S3t	S3t	S3rt	S3t	S3rt	S3rt	S3rt	S3t	S3t	S3t	S3t	S3t	S3t	S3rt	S3t	S3t	S3t	S3t	S3t	
Yampada	305	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	
Yampada	306	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	
Yampada	307	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Yampada	308	N1r	S3t	S3rt	S3t	S3rt	N1t	N1r	S3rt	N1t	S3t	S3t	S3t	S3rt	S3t	S3rt	S3rt	S3rt	S3t	S3t	S3t	S3t	S3t	S3t	S3rt	S3t	S3t	S3t	S3t	S3t	
Yampada	309	N1r	S3t	S3rt	S3t	S3rt	N1t	N1r	S3rt	N1t	S3t	S3t	S3t	S3rt	S3t	S3rt	S3rt	S3rt	S3t	S3t	S3t	S3t	S3t	S3t	S3rt	S3t	S3t	S3t	S3t	S3t	
Yampada	310	N1r	S3t	S3rt	S3t	S3rt	N1t	N1r	S3rt	N1t	S3t	S3t	S3t	S3rt	S3t	S3rt	S3rt	S3rt	S3t	S3t	S3t	S3t	S3t	S3t	S3rt	S3t	S3t	S3t	S3t	S3t	

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry	
Yampada	311	N1r	S3t	S3rt	S3t	S3rt	N1t	N1r	S3rt	N1t	S3t	S3t	S3t	S3rt	S3t	S3rt	S3rt	S3rt	S3t	S3t	S3t	S3t	S3t	S3t	S3rt	S3t	S3t	S3t	S3t	S3t	
Yampada	312	N1r	S3t	S3rt	S3t	S3rt	N1t	N1r	S3rt	N1t	S3t	S3t	S3t	S3rt	S3t	S3rt	S3rt	S3rt	S3t	S3t	S3t	S3t	S3t	S3t	S3rt	S3t	S3t	S3t	S3t	S3t	
Yampada	313	N1r	S3t	S3rt	S3t	S3rt	N1t	N1r	S3rt	N1t	S3t	S3t	S3t	S3rt	S3t	S3rt	S3rt	S3rt	S3t	S3t	S3t	S3t	S3t	S3t	S3rt	S3t	S3t	S3t	S3t	S3t	
Yampada	314	N1r	S3t	S3rt	S3t	S3rt	N1t	N1r	S3rt	N1t	S3t	S3t	S3t	S3rt	S3t	S3rt	S3rt	S3rt	S3t	S3t	S3t	S3t	S3t	S3t	S3rt	S3t	S3t	S3t	S3t	S3t	
Yampada	315	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Yampada	316	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Yampada	317	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Yampada	318	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Yampada	319	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Yampada	320	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Yampada	321	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Yampada	322	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Yampada	323	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Yampada	324	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Yampada	325	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Yampada	326	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Yampada	327	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Yampada	328	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Yampada	329	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Yampada	330	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Yampada	331	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Yampada	332	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Yampada	333	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Yampada	334	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Yampada	335	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Yampada	336	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Yampada	337	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry	
Yampada	338	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Yampada	339	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Yampada	342	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Yampada	344	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Yampada	345	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Yampada	346	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Yampada	347	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Yampada	348	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Yampada	349	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Yampada	350	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Yampada	351	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Yampada	352	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Yampada	353	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Yampada	354	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Yampada	355	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Yampada	356	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Yampada	357	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	
Yampada	358	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Yampada	359	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Yampada	360	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Yampada	361	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Yampada	362	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Yampada	363	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Yampada	364	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	

PART-B

SOCIO-ECONOMIC STATUS OF FARM HOUSEHOLDS

CONTENTS

1.	Salient findings of the survey	1-4
2.	Introduction	5
3	Methodology	7
4	Salient features of the survey	9-25
5	Summary	27-30

LIST OF TABLES

1	Households sampled for socio economic survey	9
2	Population characteristics	9
3	Age wise classification of household members	9
4	Education level of household members	10
5	Occupation of household heads	10
6	Occupation of family members	11
7	Institutional participation of household members	11
8	Type of house owned by households	11
9	Durable assets owned by households	11
10	Average value of durable assets owned by households	12
11	Farm implements owned by households	12
12	Average value of farm implements	12
13	Livestock possession by households	13
14	Average labour availability	13
15	Adequacy of hired labour	13
16	Distribution of land (ha)	14
17	Average land value (Rs./ha)	14
18	Status of bore wells	14
19	Source of irrigation	15
20	Depth of water	15
21	Irrigated area (ha)	15
22	Cropping pattern	15
23	Cropping intensity	15
24	Cost of cultivation of Redgram	16
25	Cost of cultivation of Greengram	17
26	Cost of cultivation of Paddy	18
27	Cost of cultivation of Groundnut	19
28	Adequacy of fodder	20
29	Annual gross income	20
30	Average annual expenditure	20
31	Horticulture species grown	21

32	Forest species grown	21
33	Average Additional investment capacity	21
34	Source of additional investment	21
35	Marketing of the agricultural produce	22
36	Marketing channels used for sale of agricultural produce	22
37	Mode of transport of agricultural produce	22
38	Incidence of soil and water erosion problems	22
39	Interest towards soil testing	23
40	Usage pattern of fuel for domestic use	23
41	Source of drinking water	23
42	Source of light	23
43	Existence of sanitary toilet facility	23
44	Possession of public distribution system(PDS) card	24
45	Participation in NREGA programme	24
46	Adequacy of food items	24
47	Response on inadequacy of food items	24
48	Farming constraints experienced	25

SALIENT FINDINGS OF THE SURVEY

- ❖ *The data indicated that there were 111(55.78%) men and 88 (44.22%) women among the sampled households.*
- ❖ *The average family size of landless farmers' was 4.6, marginal farmers' was 5.3, small farmers' was 5.45 and semi medium farmers' was 7.8.*
- ❖ *The data indicated that, 32 (16.08%) people were in 0-15 years of age, 91 (45.73%) were in 16-35 years of age, 44 (22.11%) were in 36-60 years of age and 32 (16.08%) were above 61 years of age.*
- ❖ *The results indicated that Yadgir Rf-1 had 51.76 per cent illiterates, 0.50 per cent of them had functional illiterate, diploma, ITI and masters, 12.06 per cent of them had primary school, 6.03 per cent of them had middle school and degree education, 11.56 per cent of them had high school education and 4.52 per cent of them had PUC education.*
- ❖ *The results indicate that agriculture was the major occupation for 25.13 per cent of the household members, 30.65 per cent were agricultural labourers, 1.51 per cent were in general labour, 4.02 per cent were private service, 4.52 per cent were trade and business, 12.56 per cent were students, 15.08 per cent were housewives and 6.53 per cent were children.*
- ❖ *The results show that, 100 per cent of the population in the micro watershed has not participated in any local institutions.*
- ❖ *The results indicate that 77.14 per cent of the households possess katcha house and 22.86 per cent of them possess pucca/RCC house.*
- ❖ *The results show that 80 per cent of the households possess TV, 11.43 per cent of the households possess mixer/grinder and bicycle, 2.86 per cent of the households possess refrigerator, 20 per cent of the households possess motor cycle and 100 per cent of the households possess mobile phones.*
- ❖ *The results show that the average value of television was Rs. 8,785, mixer/grinder was Rs.2000, refrigerator was Rs. 8,000, bicycle was Rs. 2,325, motor cycle was Rs.50,714 and mobile phone was Rs.2,787.*
- ❖ *About 2.86 per cent of the households possess bullock cart, 42.86 per cent of them possess plough, 2.86 per cent of them possess seed/fertilizer drill and 40 per cent of them possess weeder.*
- ❖ *The results show that the average value of bullock cart was Rs. 19,000, plough was Rs. 2,426, seed/fertilizer drill was Rs. 2,000 and the average value of weeder was Rs.51.*
- ❖ *The results indicate that, 34.29 per cent of the households possess bullocks, 31.43 per cent of the households possess local cow, 11.43 per cent of the households possess buffalo and 5.71 per cent of the households possess goat and poultry birds.*

- ❖ *The results indicate that, average own labour men available in the micro watershed was 2.31, average own labour (women) available was 1.66, average hired labour (men) available was 14.59 and average hired labour (women) available was 11.22.*
- ❖ *The results indicate that, 91.43 per cent of the households opined that the hired labour was adequate.*
- ❖ *The results indicate that, households of the Yadgir Rf-1 micro-watershed possess 32.95 ha (93.24%) of dry land, 1.98 ha (5.61%) of irrigated land and 0.40 ha (1.15%) of permanent fallow land. Marginal farmers possess 9.17 ha (95.13%) of dry land, 0.06 ha (1.92%) and 0.4 ha (4.20%) of permanent fallow land. Small farmers possess 11.52 ha (85.73%) of dry land and 1.92 ha (14.27%) of irrigated land. Semi medium farmers possess 12.26 ha (100%) of dry land.*
- ❖ *The results indicate that, the average value of dry land was Rs. 646,088.67; the average value of irrigated land was Rs. 1,310,612.24 and the average value of permanent fallow land was Rs. 1,976,000. In case of marginal famers, the average land value was Rs. 1,144,022.94 for dry land, the average value of irrigated land was Rs. 6,175,000.14 and the average value of permanent fallow land was Rs. 1,976,000. In case of small famers, the average land value was Rs. 563,926.94 for dry land and Rs. 1,146,413.50 for irrigated land. In case of semi medium famers, the average land value was Rs. 350,643.78 for dry land.*
- ❖ *The results indicate that, there were 4 functioning bore wells in the micro watershed.*
- ❖ *The results indicate that, bore well was the major irrigation source in the micro water shed for 11.43 per cent of the farmers.*
- ❖ *The results indicate that, the depth of bore well was found to be 12.19 meters.*
- ❖ *The results indicate that, marginal and small farmers had an irrigated area of 0.06 ha and 1.92 ha respectively.*
- ❖ *The results indicate that, farmers have grown red gram (31.5 ha), groundnut (2.44 ha), paddy (0.61 ha) and green gram (4.02 ha). Marginal farmers have grown red gram, paddy and green gram. Small farmers have grown red gram, groundnut and paddy. Semi medium farmers have grown red gram and groundnut.*
- ❖ *The results indicate that, the cropping intensity in Yadgir Rf-1 micro-watershed was found to be 99.20 per cent.*
- ❖ *The results indicate that, the total cost of cultivation for red gram was Rs. 31285.33. The gross income realized by the farmers was Rs. 61970.38. The net income from red gram cultivation was Rs. 30685.06. Thus the benefit cost ratio was found to be 1:1.98.*
- ❖ *The total cost of cultivation for green gram was Rs. 70223.42. The gross income realized by the farmers was Rs. 61750. The net income from green gram cultivation was Rs. -8473.42. Thus the benefit cost ratio was found to be 1:0.88.*

- ❖ *The total cost of cultivation for paddy was Rs. 292727.65. The gross income realized by the farmers was Rs. 310942.70. The net income from paddy cultivation was Rs. 18215.05. Thus the benefit cost ratio was found to be 1:1.06.*
- ❖ *The total cost of cultivation for groundnut was Rs. 69276.29. The gross income realized by the farmers was Rs. 100829.94. The net income from groundnut cultivation was Rs. 31553.65. Thus the benefit cost ratio was found to be 1:1.46.*
- ❖ *The results indicate that, 45.71 per cent of the households opined that dry fodder was adequate and 48.57 per cent of the households opined that green fodder was adequate.*
- ❖ *The results indicate that the annual gross income was Rs. 108,333.33 for landless farmers, for marginal farmers it was Rs. 105,785, for small farmers it was Rs. 179,718.18 and for semi medium farmers it was Rs. 191,928.*
- ❖ *The results indicate that the average annual expenditure is Rs. 18,655.77. For landless households it was Rs. 45,000, for marginal farmers it was Rs. 9,227.20, for small farmers it was Rs. 16,506.06 and for semi medium farmers it was Rs. 37,750.*
- ❖ *The results indicate that, sampled households have grown 2 coconut and 17 custard apple and mango tree in their field and also 2 coconut, custard apple and guava trees in their backyard.*
- ❖ *The results indicate that, households have planted 65 neem and 2 tamarind trees in their field and also 26 neem trees in their backyard.*
- ❖ *The results indicated that, households have an average investment capacity of Rs. 3,428.57 for land development, Rs. 15,285.71 for irrigation facility and Rs. 285.71 for improved crop production.*
- ❖ *The results indicated that loan from bank was the source of additional investment for 14.29 per cent for land development, 28.57 for irrigation facility and for 2.86 per cent of irrigation facility.*
- ❖ *The results indicated that, green gram was sold to the extent of 60 per cent, groundnut was sold to the extent of 71.43 per cent, paddy was sold to the extent of 26.83 per cent and red gram to the extent of 61.98 per cent.*
- ❖ *The results indicated that, about 82.86 per cent of the farmers sold their produce to local/village merchants and 11.43 per cent of the farmers sold their produce to regulated markets.*
- ❖ *The results indicated that, 94.29 per cent of the households have used tractor as a mode of transportation.*
- ❖ *The results indicated that, 42.86 per cent of the households have experienced soil and water erosion problems in the farm.*
- ❖ *The results indicated that, 85.71 per cent have shown interest in soil test.*
- ❖ *The results indicated that, 97.14 per cent of the households used firewood and 2.86 per cent of them used LPG as a source of fuel.*

- ❖ *The results indicated that, piped supply was the major source of drinking water for 100 per cent of the households in the micro watershed.*
- ❖ *Electricity was the major source of light for 100 per cent of the households in micro watershed.*
- ❖ *The results indicated that, 57.14 per cent of the households possess sanitary toilet facility.*
- ❖ *The results indicated that, 100 per cent of the sampled households possessed BPL cards.*
- ❖ *The results indicated that, 82.86 per cent of the households participated in NREGA programme.*
- ❖ *The results indicated that, cereals, milk, egg and meat were adequate for 100 per cent of the households, pulses were adequate for 94.29 per cent, oilseeds were adequate for 5.71 per cent, vegetables were adequate for 71.43 per cent and fruits were adequate for 17.14 per cent.*
- ❖ *The results indicated that, pulses were inadequate for 5.71 per cent of the households, oilseeds were inadequate for 94.29 per cent, vegetables were inadequate for 51.43 per cent and fruits were inadequate for 82.86 per cent of the households.*
- ❖ *The results indicated that, lower fertility status of the soil, frequent incidence of pest and diseases, inadequacy of irrigation water and high cost of fertilizers and plant protection chemicals was the constraint experienced by 91.43 per cent of the households, wild animal menace on farm field and high rate of interest on credit (88.57%), low price for the agricultural commodities (11.43%), lack of marketing facilities in the area and inadequate extension services (2.86%).*

INTRODUCTION

Soil and water are the two precious natural resources which are essential for crop production and existence of life on earth. Rainfed agriculture is under severe stress due to various constraints related to agriculture like uneven and erratic distribution of rainfall, indiscriminate use of fertilizers, chemicals and pesticides, adoption of improper land management practices, soil erosion, decline in soil fertility, decline in ground water resources leading to low crop productivity. The area under rainfed agriculture has to be managed effectively using the best available practices to enhance the production of food, fodder and fuel. This is possible if the land resources are characterized at each parcel of land through detailed land resource inventory using the best available techniques of remote sensing, GPS and GIS. The watershed development programs are aimed at the sustainable distribution of its resources and the process of creating and implementing plans, programs, and projects to sustain and enhance watershed functions that affect the plant, animal and human communities within a watershed boundary.

World Bank funded KWDP II, SUJALA III project was implemented in with Broad objective of demonstrating more effective watershed management through greater integration of programmes related to rain-fed agriculture, innovative and science based approaches and strengthen institutional capacities and If successful, it is expected that the systems and tools could be mainstreamed into the overall IWMP in the State of Karnataka and in time, throughout other IWMP operations in India. With this background the socio-economic survey has been carried out with following specific objectives:

1. To understand the demographic features of the households in the micro-watershed
2. To understand the extent of family labour available and additional employment opportunities available within the village.
3. To know the status of assets of households in the micro-watershed for suggesting possible improvements.
4. To study the cropping pattern, cropped area and productivity levels of different households in micro-watershed.
5. To determine the type and extent of livestock owned by different categories of HHs
6. Availability of fodder and level of livestock management.

Scope and importance of survey

Survey helps in identification of different socio-economic and resource use-patterns of farmers at the Micro watershed. Household survey provides demographic features, labour force, and levels of education; land ownership and asset position (including livestock and other household assets) of surveyed households; and cropping patterns, input intensities, and average crop yields from farmers' fields. It also discusses crop utilization and the degree of commercialization of production in the areas; farmers' access to and utilization of credit from formal and informal sources; and the level of adoption and use of soil, water, and pest management technologies.

METHODOLOGY

The description of the methods, components selected for the survey and procedures followed in conducting the baseline survey are furnished under the following heads.

Description of the study area

Yadgir District is one of the 30 districts of Karnataka state in southern India. This district was carved out from the erstwhile Gulbarga district as the 30th district of Karnataka on 10 April 2010. Yadgir town is the administrative headquarters of the district. The district comprises of 3 taluks namely, Shahapur, Yadgiri and Shorapur (There are 16 hoblies, 117 Gram Panchayats, 4 Municipalities, 8 Towns/ Urban agglomeration and 487 inhabited & 32 un-inhabited villages The district occupies an area of 5,160.88 km².

Yadgiri district is the second smallest district in the state, area wise is very rich in cultural traditions. The vast stretch of fertile black soil of the district is known for bumper red gram and jowar crops. The district is a "Daal bowl" of the state. The district is also known for cluster of cement industries and a distinct stone popularly known as "Malakheda Stone". Two main rivers, Krishna and Bhima, and a few tributaries flow in this region. Krishna and Bhima Rivers drain the district. They constitute the two major river basins of the district. Kagna and Amarja are the two sub - basins of Bhima River, which occur within the geographical area of the district

According to the 2011 census Yadgiri district has a population of 1, 172,985, roughly equal to the nation of Timor-Leste or the US state of Rhode Island. This gives it a ranking of 404th in India (out of a total of 640). The district has a population density of 224 inhabitants per square kilometre (580/sq mi). Its population growth rate over the decade 2001-2011 was 22.67%. Yadgir has a sex ratio of 984 females for every 1000 males, and a literacy rate of 52.36%.

Description of the micro watershed

Yadgir Rf-1 micro-watershed in Bewanahalli sub-watershed (Yadgiri taluk and district) is located in between 16°52'12.613" to 16° 50'13.113" North latitudes and 77° 17'2.044" to 77°15'35.439" East longitudes, covering an area of about 489.94 ha, bounded by Gajarkota and Yampada villages.

Methodology followed in assessing socio-economic status of households

In order to assess the socio-economic condition of the farmers in the watershed a comprehensive questionnaire was prepared. Major components such as demographic conditions, migration details, food consumption and family expenditure pattern, material possession, land holding, land use management, cropping pattern, cost of cultivation of crops, livestock management. The statistical components such as frequency and percentage were used to analyze the data. About 35 households located in the micro-watershed were interviewed for the survey.

SALIENT FEATURES OF THE SURVEY

Households sampled for socio-economic survey: The data on households sampled for socio economic survey in Yadgir Rf-1 micro-watershed is presented in Table 1 and it indicated that 35 farmers were sampled in Yadgir Rf-1 micro-watershed among them 3 (8.57%) were landless, 16 (45.71%) were marginal farmers, 11 (31.43%) were small farmers and 5 (14.29%) were semi medium farmers.

Table 1: Households sampled for socio economic survey in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (3)		MF (16)		SF (11)		SMF (5)		All (35)	
		N	%	N	%	N	%	N	%	N	%
1	Farmers	3	8.57	16	45.71	11	31.43	5	14.29	35	100

Population characteristics: The population characteristics of households sampled for socio-economic survey in Yadgir Rf-1 micro-watershed is presented in Table 2. The data indicated that there were 111(55.78%) men and 88 (44.22%) women among the sampled households. The average family size of landless farmers' was 4.6, marginal farmers' was 5.3, small farmers' was 5.45 and semi medium farmers' was 7.8.

Table 2: Population characteristics of Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (14)		MF (86)		SF (60)		SMF (39)		All (199)	
		N	%	N	%	N	%	N	%	N	%
1	Men	8	57.14	47	54.65	32	53.33	24	61.54	111	55.78
2	Women	6	42.86	39	45.35	28	46.67	15	38.46	88	44.22
	Total	14	100	86	100	60	100	39	100	199	100
	Average		4.6		5.3		5.45		7.8		5.6

Age wise classification of population: The age wise classification of household members in Yadgir Rf-1 micro-watershed is presented in Table 3. The data indicated that, 32 (16.08%) people were in 0-15 years of age, 91 (45.73%) were in 16-35 years of age, 44 (22.11%) were in 36-60 years of age and 32 (16.08%) were above 61 years of age.

Table 3: Age wise classification of household members in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (14)		MF (86)		SF (60)		SMF (39)		All (199)	
		N	%	N	%	N	%	N	%	N	%
1	0-15 years of age	0	0	18	20.93	9	15	5	12.82	32	16.08
2	16-35 years of age	8	57.14	38	44.19	27	45	18	46.15	91	45.73
3	36-60 years of age	4	28.57	18	20.93	14	23.33	8	20.51	44	22.11
4	> 61 years	2	14.29	12	13.95	10	16.67	8	20.51	32	16.08
	Total	14	100	86	100	60	100	39	100	199	100

Education level of household members: Education level of household members in Yadgir Rf-1 micro-watershed is presented in Table 4. The results indicated that Yadgir Rf-1 had 51.76 per cent illiterates, 0.50 per cent of them had functional illiterate, diploma,

ITI and masters, 12.06 per cent of them had primary school, 6.03 per cent of them had middle school and degree education, 11.56 per cent of them had high school education and 4.52 per cent of them had PUC education

Table 4. Education level of household members in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (14)		MF (86)		SF (60)		SMF (39)		All (199)	
		N	%	N	%	N	%	N	%	N	%
1	Illiterate	8	57.14	38	44.19	26	43.33	31	79.49	103	51.76
2	Functional Literate	0	0	0	0	1	1.67	0	0	1	0.50
3	Primary School	0	0	13	15.12	8	13.33	3	7.69	24	12.06
4	Middle School	2	14.29	6	6.98	4	6.67	0	0	12	6.03
5	High School	2	14.29	14	16.28	5	8.33	2	5.13	23	11.56
6	PUC	1	7.14	2	2.33	5	8.33	1	2.56	9	4.52
7	Diploma	0	0	0	0	1	1.67	0	0	1	0.50
8	ITI	0	0	1	1.16	0	0	0	0	1	0.50
9	Degree	1	7.14	5	5.81	5	8.33	1	2.56	12	6.03
10	Masters	0	0	0	0	1	1.67	0	0	1	0.50
11	Others	0	0	7	8.14	4	6.67	1	2.56	12	6.03
Total		14	100	86	100	60	100	39	100	199	100

Occupation of household heads: The data regarding the occupation of the household heads in Yadgir Rf-1 micro-watershed is presented in Table 5. The results indicate that, 40 per cent of household heads were practicing agriculture, 42.86 per cent of the household heads were agricultural labourers, 5.71 per cent of the household heads were general labour and housewives, 2.86 cent of the household heads were in private service and 8.57 per cent of them were trade and business.

Table 5: Occupation of household heads in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (3)		MF (16)		SF (11)		SMF (5)		All (35)	
		N	%	N	%	N	%	N	%	N	%
1	Agriculture	0	0	8	50	4	36.36	2	40	14	40
2	Agricultural Labour	2	66.67	5	31.25	4	36.36	4	80	15	42.86
3	General Labour	0	0	2	12.50	0	0	0	0	2	5.71
4	Private Service	0	0	0	0	1	9.09	0	0	1	2.86
5	Trade & Business	1	33.33	1	6.25	1	9.09	0	0	3	8.57
6	Housewife	0	0	1	6.25	1	9.09	0	0	2	5.71
Total		3	100	17	100	11	100	6	100	37	100

Occupation of the household members: The data regarding the occupation of the household members in Yadgir Rf-1 micro-watershed is presented in Table 6. The results indicate that agriculture was the major occupation for 25.13 per cent of the household members, 30.65 per cent were agricultural labourers, 1.51 per cent were in general labour, 4.02 per cent were private service, 4.52 per cent were trade and business, 12.56 per cent were students, 15.08 per cent were housewives and 6.53 per cent were children.

Table 6: Occupation of family members in Yadgir Rf-1 micro-watershed

Sl. No.	Particulars	LL (14)		MF (86)		SF (60)		SMF (39)		All (199)	
		N	%	N	%	N	%	N	%	N	%
1	Agriculture	0	0	28	32.56	19	31.67	3	7.69	50	25.13
2	Agricultural Labour	6	42.86	19	22.09	16	26.67	20	51.28	61	30.65
3	General Labour	0	0	3	3.49	0	0	0	0	3	1.51
4	Private Service	0	0	2	2.33	5	8.33	1	2.56	8	4.02
5	Trade & Business	6	42.86	2	2.33	1	1.67	0	0	9	4.52
6	Student	2	14.29	15	17.44	6	10	2	5.13	25	12.56
7	Children	0	0	7	8.14	5	8.33	1	2.56	13	6.53
8	Housewife	0	0	10	11.63	8	13.33	12	30.77	30	15.08
Total		14	100	86	100	60	100	39	100	199	100

Institutional participation of the household members: The data regarding the institutional participation of the household members in Yadgir Rf-1 micro-watershed is presented in Table 7. The results show that, 100 per cent of the population in the micro watershed has not participated in any local institutions.

Table 7. Institutional Participation of household members in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (14)		MF (86)		SF (60)		SMF (39)		All (199)	
		N	%	N	%	N	%	N	%	N	%
1	No Participation	14	100	86	100	60	100	39	100	199	100
Total		14	100	86	100	60	100	39	100	199	100

Type of house owned: The data regarding the type of house owned by the households in Yadgir Rf-1 micro-watershed is presented in Table 8. The results indicate that 77.14 per cent of the households possess katcha house and 22.86 per cent of them possess pucca/RCC house.

Table 8. Type of house owned by households in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (3)		MF (16)		SF (11)		SMF (5)		All (35)	
		N	%	N	%	N	%	N	%	N	%
1	Katcha	3	100	12	75	8	72.73	4	80	27	77.14
2	Pucca/RCC	0	0	4	25	3	27.27	1	20	8	22.86
Total		3	100	16	100	11	100	5	100	35	100

Table 9. Durable Assets owned by households in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (3)		MF (16)		SF (11)		SMF (5)		All (35)	
		N	%	N	%	N	%	N	%	N	%
1	Television	3	100	12	75	9	81.82	4	80	28	80
2	Mixer/Grinder	0	0	3	18.75	1	9.09	0	0	4	11.43
3	Refrigerator	0	0	0	0	1	9.09	0	0	1	2.86
4	Bicycle	2	66.67	0	0	2	18.18	0	0	4	11.43
5	Motor Cycle	0	0	3	18.75	3	27.27	1	20	7	20
6	Mobile Phone	3	100	16	100	11	100	5	100	35	100

Durable Assets owned by the households: The data regarding the Durable Assets owned by the households in Yadgir Rf-1 micro-watershed is presented in Table 9. The

results show that 80 per cent of the households possess TV, 11.43 per cent of the households possess mixer/grinder and bicycle, 2.86 per cent of the households possess refrigerator, 20 per cent of the households possess motor cycle and 100 per cent of the households possess mobile phones.

Average value of durable assets: The data regarding the average value of durable assets owned by the households in Yadgir Rf-1 micro-watershed is presented in Table 10. The results show that the average value of television was Rs. 8,785, mixer/grinder was Rs.2000, refrigerator was Rs. 8,000, bicycle was Rs. 2,325, motor cycle was Rs.50,714 and mobile phone was Rs.2,787.

Table 10. Average value of durable assets owned by households in Yadgir Rf-1 micro-watershed
Average value (Rs.)

Sl.No.	Particulars	LL (3)	MF (16)	SF (11)	SMF (5)	All (35)
1	Television	9,000	8,583	8,888	9,000	8,785
2	Mixer/Grinder	0	2,000	2,000	0	2,000
3	Refrigerator	0	0	8,000	0	8,000
4	Bicycle	1,900	0	2,750	0	2,325
5	Motor Cycle	0	41,666	61,666	45,000	50,714
6	Mobile Phone	2,000	2,694	2,972	2,892	2,787

Farm Implements owned: The data regarding the farm implements owned by the households in Yadgir Rf-1 micro-watershed is presented in Table 11. About 2.86 per cent of the households possess bullock cart, 42.86 per cent of them possess plough, 2.86 per cent of them possess seed/fertilizer drill and 40 per cent of them possess weeder.

Table 11. Farm Implements owned by households in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (3)		MF (16)		SF (11)		SMF (5)		All (35)	
		N	%	N	%	N	%	N	%	N	%
1	Bullock Cart	0	0	1	6.25	0	0	0	0	1	2.86
2	Plough	0	0	6	37.50	4	36.36	5	100	15	42.86
3	Seed/Fertilizer Drill	0	0	1	6.25	0	0	0	0	1	2.86
4	Weeder	0	0	5	31.25	4	36.36	5	100	14	40
5	Blank	3	100	10	62.50	7	63.64	0	0	20	57.14

Table 12. Average value of farm implements owned by households in Yadgir Rf-1 micro-watershed
Average Value (Rs.)

Sl.No.	Particulars	LL (3)	MF (16)	SF (11)	SMF (5)	All (35)
1	Bullock Cart	0	19,000	0	0	19,000
2	Plough	0	3,833	1,500	1,480	2,426
3	Seed/Fertilizer Drill	0	2,000	0	0	2,000
4	Weeder	0	52	50	50	51

Average value of farm implements: The data regarding the average value of farm Implements owned by the households in Yadgir Rf-1 micro-watershed is presented in Table 12. The results show that the average value of bullock cart was Rs. 19,000, plough

was Rs. 2,426, seed/fertilizer drill was Rs. 2,000 and the average value of weeder was Rs.51.

Livestock possession by the households: The data regarding the Livestock possession by the households in Yadgir Rf-1 micro-watershed is presented in Table 13. The results indicate that, 34.29 per cent of the households possess bullocks, 31.43 per cent of the households possess local cow, 11.43 per cent of the households possess buffalo and 5.71 per cent of the households possess goat and poultry birds.

Table 13. Livestock possession by households in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (3)		MF (16)		SF (11)		SMF (5)		All (35)	
		N	%	N	%	N	%	N	%	N	%
1	Bullock	0	0	5	31.25	4	36.36	3	60	12	34.29
2	Local cow	0	0	5	31.25	3	27.27	3	60	11	31.43
3	Buffalo	0	0	2	12.50	1	9.09	1	20	4	11.43
4	Goat	0	0	1	6.25	0	0	1	20	2	5.71
5	Poultry birds	0	0	0	0	1	9.09	1	20	2	5.71

Average Labour availability: The data regarding the average labour availability in Yadgir Rf-1 micro-watershed is presented in Table 14. The results indicate that, average own labour men available in the micro watershed was 2.31, average own labour (women) available was 1.66, average hired labour (men) available was 14.59 and average hired labour (women) available was 11.22.

In case of marginal farmers, average own labour men available was 2, average own labour (women) was 1.38, average hired labour (men) was 14.25 and average hired labour (women) available was 9.69. In case of small farmers, average own labour men available was 2, average own labour (women) was 1.27, average hired labour (men) was 14.45 and average hired labour (women) available was 12.64. In case of semi medium farmers, average own labour men available was 4, average own labour (women) was 1.27, average hired labour (men) was 14.45 and average hired labour (women) available was 12.64.

Table 14. Average Labour availability in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (3)	MF (16)	SF (11)	SMF (5)	All (35)
1	Hired labour Female	0	9.69	12.64	13	11.22
2	Own Labour Female	0	1.38	1.27	3.40	1.66
3	Own labour Male	0	2	2	4	2.31
4	Hired labour Male	0	14.25	14.45	16	14.59

Table 15. Adequacy of Hired Labour in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (3)		MF (16)		SF (11)		SMF (5)		All (35)	
		N	%	N	%	N	%	N	%	N	%
1	Adequate	0	0	16	100	11	100	5	100	32	91.43

Adequacy of Hired Labour: The data regarding the adequacy of hired labour in Yadgir Rf-1 micro-watershed is presented in Table 15. The results indicate that, 91.43 per cent of the households opined that the hired labour was adequate.

Distribution of land (ha): The data regarding the distribution of land (ha) in Yadgir Rf-1 micro-watershed is presented in Table 16. The results indicate that, households of the Yadgir Rf-1 micro-watershed possess 32.95 ha (93.24%) of dry land, 1.98 ha (5.61%) of irrigated land and 0.40 ha (1.15%) of permanent fallow land. Marginal farmers possess 9.17 ha (95.13%) of dry land, 0.06 ha (1.92%) and 0.4 ha (4.20%) of permanent fallow land. Small farmers possess 11.52 ha (85.73%) of dry land and 1.92 ha (14.27%) of irrigated land. Semi medium farmers possess 12.26 ha (100%) of dry land.

Table 16. Distribution of land (Ha) in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (3)		MF (16)		SF (11)		SMF (5)		All (35)	
		ha	%	ha	%	ha	%	ha	%	ha	%
1	Dry	0	0	9.17	95.13	11.52	85.73	12.26	100	32.95	93.24
2	Irrigated	0	0	0.06	0.67	1.92	14.27	0	0	1.98	5.61
3	Permanent Fallow	0	0	0.40	4.20	0	0	0	0	0.40	1.15
Total		0	100	9.64	100	13.44	100	12.26	100	35.34	100

Average land value (Rs./ha): The data regarding the average land value (Rs./ha) in Yadgir Rf-1 micro-watershed is presented in Table 17. The results indicate that, the average value of dry land was Rs. 646,088.67; the average value of irrigated land was Rs. 1,310,612.24 and the average value of permanent fallow land was Rs. 1,976,000. In case of marginal famers, the average land value was Rs. 1,144,022.94 for dry land, the average value of irrigated land was Rs. 6,175,000.14 and the average value of permanent fallow land was Rs. 1,976,000. In case of small famers, the average land value was Rs. 563,926.94 for dry land and Rs. 1,146,413.50 for irrigated land. In case of semi medium famers, the average land value was Rs. 350,643.78 for dry land.

Table 17. Average land value (Rs./ha) in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (3)	MF (16)	SF (11)	SMF (5)	All (35)
1	Dry	0	1,144,022.94	563,926.94	350,643.78	646,088.67
2	Irrigated	0	6,175,000.14	1,146,413.50	0	1,310,612.24
3	Permanent Fallow	0	1,976,000	0	0	1,976,000

Status of bore wells: The data regarding the status of bore wells in Yadgir Rf-1 micro-watershed is presented in Table 18. The results indicate that, there were 4 functioning bore wells in the micro watershed.

Table 18. Status of bore wells in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (3)	MF (16)	SF (11)	SMF (5)	All (35)
1	Functioning	0	1	3	0	4

Source of irrigation: The data regarding the source of irrigation in Yadgir Rf-1 micro-watershed is presented in Table 19. The results indicate that, bore well was the major irrigation source in the micro water shed for 11.43 per cent of the farmers.

Table 19. Source of irrigation in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (3)		MF (16)		SF (11)		SMF (5)		All (35)	
		N	%	N	%	N	%	N	%	N	%
1	Bore Well	0	0	1	6.25	3	27.27	0	0	4	11.43

Depth of water (Avg in meters): The data regarding the depth of water in Yadgir Rf-1 micro-watershed is presented in Table 20. The results indicate that, the depth of bore well was found to be 12.19 meters.

Table 20. Depth of water (Avg in meters) in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (3)	MF (16)	SF (11)	SMF (5)	All (35)
1	Bore Well	0	6.67	29.09	0	12.19

Irrigated Area (ha): The data regarding the irrigated area (ha) in Yadgir Rf-1 micro-watershed is presented in Table 21. The results indicate that, marginal and small farmers had an irrigated area of 0.06 ha and 1.92 ha respectively.

Table 21. Irrigated Area (ha) in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (3)	MF (16)	SF (11)	SMF (5)	All (35)
1	Kharif	0	0.06	1.92	0	1.98

Cropping pattern: The data regarding the cropping pattern in Yadgir Rf-1 micro-watershed is presented in Table 22. The results indicate that, farmers have grown red gram (31.5 ha), groundnut (2.44 ha), paddy (0.61 ha) and green gram (4.02 ha). Marginal farmers have grown red gram, paddy and green gram. Small farmers have grown red gram, groundnut and paddy. Semi medium farmers have grown red gram and groundnut.

Table 22. Cropping pattern in Yadgir Rf-1 micro-watershed (Area in ha)

Sl.No.	Particulars	LL (3)	MF (16)	SF (11)	SMF (5)	All (35)
1	Kharif - Red gram	0	8.77	11.3	11.43	31.5
2	Kharif - Groundnut	0	0	1.63	0.81	2.44
3	Kharif - Paddy	0	0.06	0.55	0	0.61
4	Kharif - Greengram	0	0.4	0	0	0.4
	Total	0	9.24	13.47	12.24	34.96

Cropping intensity: The data regarding the cropping intensity in Yadgir Rf-1 micro-watershed is presented in Table 23. The results indicate that, the cropping intensity in Yadgir Rf-1 micro-watershed was found to be 99.20 per cent.

Table 23. Cropping intensity (%) in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (3)	MF (16)	SF (11)	SMF (5)	All (35)
1	Cropping Intensity	0	100	97.94	100	99.20

Cost of cultivation of Red gram: The data regarding the cost of cultivation of red gram in Yadgir Rf-1 micro-watershed is presented in Table 27. The results indicate that, the total cost of cultivation for red gram was Rs. 31285.33. The gross income realized by the farmers was Rs. 61970.38. The net income from red gram cultivation was Rs. 30685.06. Thus the benefit cost ratio was found to be 1:1.98.

Table 27. Cost of Cultivation of red gram in Yadgir Rf-1 micro-watershed

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
I	Cost A1				
1	Hired Human Labour	Man days	31.75	5181.16	16.56
2	Bullock	Pairs/day	8.01	4808.97	15.37
3	Tractor	Hours	4.02	3214.93	10.28
4	Machinery	Hours	0.05	41.82	0.13
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	6.84	820.63	2.62
7	FYM	Quintal	2.28	455.36	1.46
8	Fertilizer + micronutrients	Quintal	5.35	4252.96	13.59
9	Pesticides (PPC)	Kgs /liters	1.22	1215.73	3.89
10	Irrigation	Number	0	0	0
11	Repairs		0	0	0
12	Msc. Charges (Marketing costs etc)		0	0	0
13	Depreciation charges		0	73.81	0.24
14	Land revenue and Taxes		0	3.29	0.01
II	Cost B1				
16	Interest on working capital			809.48	2.59
17	Cost B1 = (Cost A1 + sum of 15 and 16)			20878.13	66.73
III	Cost B2				
18	Rental Value of Land			407.41	1.30
19	Cost B2 = (Cost B1 + Rental value)			21285.54	68.04
IV	Cost C1				
20	Family Human Labour		29.65	7154.67	22.87
21	Cost C1 = (Cost B2 + Family Labour)			28440.21	90.91
V	Cost C2				
22	Risk Premium			1	0
23	Cost C2 = (Cost C1 + Risk Premium)			28441.21	90.91
VI	Cost C3				
24	Managerial Cost			2844.12	9.09
25	Cost C3 = (Cost C2 + Managerial Cost)			31285.33	100
VII	Economics of the Crop				
a.	Main Product	a) Main Product (q)		9.41	43289.75
		b) Main Crop Sales Price (Rs.)			4600
	By Product	e) Main Product (q)		7.20	18680.64
		f) Main Crop Sales Price (Rs.)			2596.30
b.	Gross Income (Rs.)			61970.38	
c.	Net Income (Rs.)			30685.06	
d.	Cost per Quintal (Rs./q.)			3324.40	
e.	Benefit Cost Ratio (BC Ratio)			1:1.98	

Cost of Cultivation of Green gram: The data regarding the cost of cultivation of green gram in Yadgir Rf-1 micro-watershed is presented in Table 28. The results indicate that, the total cost of cultivation for green gram was Rs. 70223.42. The gross income realized by the farmers was Rs. 61750. The net income from green gram cultivation was Rs. - 8473.42. Thus the benefit cost ratio was found to be 1:0.88.

Table 28. Cost of Cultivation of green gram in Yadgir Rf-1 micro-watershed

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
I	Cost A1				
1	Hired Human Labour	Man days	74.10	12967.50	18.47
2	Bullock	Pairs/day	9.88	5928	8.44
3	Tractor	Hours	9.88	7904	11.26
4	Machinery	Hours	0	0	0
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	7.41	889.20	1.27
6	Seed Inter Crop	Kgs.	0	0	0
7	FYM	Quintal	0	0	0
8	Fertilizer + micronutrients	Quintal	19.76	23712	33.77
9	Pesticides (PPC)	Kgs / liters	2.47	2470	3.52
10	Irrigation	Number	0	0	0
11	Repairs		0	0	0
12	Msc. Charges (Marketing costs etc)		0	0	0
13	Depreciation charges		0	83.98	0.12
14	Land revenue and Taxes		0	3.29	0
II	Cost B1				
16	Interest on working capital			3248.66	4.63
17	Cost B1 = (Cost A1 + sum of 15 and 16)			57206.64	81.46
III	Cost B2				
18	Rental Value of Land			333.33	0.47
19	Cost B2 = (Cost B1 + Rental value)			57539.97	81.94
IV	Cost C1				
20	Family Human Labour		27.17	6298.50	8.97
21	Cost C1 = (Cost B2 + Family Labour)			63838.47	90.91
V	Cost C2				
22	Risk Premium			1	0
23	Cost C2 = (Cost C1 + Risk Premium)			63839.47	90.91
VI	Cost C3				
24	Managerial Cost			6383.95	9.09
25	Cost C3 = (Cost C2 + Managerial Cost)			70223.42	100
VII	Economics of the Crop				
a.	Main Product	a) Main Product (q)	12.35	61750	
		b) Main Crop Sales Price (Rs.)		5000	
b.	Gross Income (Rs.)			61750	
c.	Net Income (Rs.)			-8473.42	
d.	Cost per Quintal (Rs./q.)			5686.11	
e.	Benefit Cost Ratio (BC Ratio)			1:0.88	

Cost of Cultivation of paddy: The data regarding the cost of cultivation of paddy in Yadgir Rf-1 micro-watershed is presented in Table 29. The results indicate that, the total cost of cultivation for paddy was Rs. 292727.65. The gross income realized by the farmers was Rs. 310942.70. The net income from paddy cultivation was Rs. 18215.05. Thus the benefit cost ratio was found to be 1:1.06.

Table 29. Cost of Cultivation of paddy in Yadgir Rf-1 micro-watershed

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
I	Cost A1				
1	Hired Human Labour	Man days	364.04	70343.54	24.03
2	Bullock	Pairs/day	3.66	2195.56	0.75
3	Tractor	Hours	18.18	14545.56	4.97
4	Machinery	Hours	0	0	0
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	200.12	60606.48	20.70
6	Seed Inter Crop	Kgs.	0	0	0
7	FYM	Quintal	17.27	3453.43	1.18
8	Fertilizer + micronutrients	Quintal	65.92	50040.37	17.09
9	Pesticides (PPC)	Kgs / liters	8.63	8633.56	2.95
10	Irrigation	Number	56.78	0	0
11	Repairs		0	0	0
12	Msc. Charges (Marketing costs etc)		0	0	0
13	Depreciation charges		0	0.17	0
14	Land revenue and Taxes		0	3.29	0
II	Cost B1				
16	Interest on working capital			14728.18	5.03
17	Cost B1 = (Cost A1 + sum of 15 and 16)			224550.15	76.71
III	Cost B2				
18	Rental Value of Land			166.67	0.06
19	Cost B2 = (Cost B1 + Rental value)			224716.81	76.77
IV	Cost C1				
20	Family Human Labour		176.85	41398.23	14.14
21	Cost C1 =(Cost B2 + family Labour)			266115.04	90.91
V	Cost C2				
22	Risk Premium			1	0
23	Cost C2 = (Cost C1 +Risk Premium)			266116.04	90.91
VI	Cost C3				
24	Managerial Cost			26611.60	9.09
25	Cost C3 = (Cost C2 + managerial Cost)			292727.65	100
VII	Economics of the Crop				
a.	Main Product	a) Main Product (q)		139.57	258197.91
		b) Main Crop Sales Price (Rs.)			1850
	By Product	e) Main Product (q)		35.16	52744.79
		f) MainCropSalesPrice(Rs.)			1500
b.	Gross Income (Rs.)			310942.70	
c.	Net Income (Rs.)			18215.05	
d.	Cost per Quintal (Rs./q.)			2097.41	
e.	Benefit Cost Ratio (BC Ratio)			1:1.06	

Cost of cultivation of Groundnut: The data regarding the cost of cultivation of groundnut in Yadgir Rf-1 micro-watershed is presented in Table 31. The results indicate that, the total cost of cultivation for groundnut was Rs. 69276.29. The gross income realized by the farmers was Rs. 100829.94. The net income from groundnut cultivation was Rs. 31553.65. Thus the benefit cost ratio was found to be 1:1.46.

Table 31. Cost of Cultivation of groundnut in Yadgir Rf-1 micro-watershed

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
I	Cost A1				
1	Hired Human Labour	Man days	26.94	5907.42	8.53
2	Bullock	Pairs/day	12.28	7366.20	10.63
3	Tractor	Hours	6.96	5569.46	8.04
4	Machinery	Hours	0	0	0
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	112.60	16889.99	24.38
7	FYM	Quintal	2.46	491.57	0.71
8	Fertilizer + micronutrients	Quintal	8.16	6715.68	9.69
9	Pesticides (PPC)	Kgs / liters	1.23	1228.92	1.77
10	Irrigation	Number	6.08	0	0
11	Repairs		0	0	0
12	Msc. Charges (Marketing costs etc)		0	0	0
13	Depreciation charges		0	40.96	0.06
14	Land revenue and Taxes		0	3.29	0
II	Cost B1				
16	Interest on working capital			3039.26	4.39
17	Cost B1 = (Cost A1 + sum of 15 and 16)			47252.75	68.21
III	Cost B2				
18	Rental Value of Land			333.33	0.48
19	Cost B2 = (Cost B1 + Rental value)			47586.08	68.69
IV	Cost C1				
20	Family Human Labour		48.46	15391.37	22.22
21	Cost C1 = (Cost B2 + Family Labour)			62977.45	90.91
V	Cost C2				
22	Risk Premium			1	0
23	Cost C2 = (Cost C1 + Risk Premium)			62978.45	90.91
VI	Cost C3				
24	Managerial Cost			6297.84	9.09
25	Cost C3 = (Cost C2 + Managerial Cost)			69276.29	100
VII	Economics of the Crop				
a.	Main Product	a) Main Product (q)		17.22	84363.28
		b) Main Crop Sales Price(Rs.)			4900
	By Product	e) Main Product (q)		16.47	16466.67
		f) Main Crop Sales Price (Rs.)			1000
b.	Gross Income (Rs.)			100829.94	
c.	Net Income (Rs.)			31553.65	
d.	Cost per Quintal (Rs./q.)			4023.72	
e.	Benefit Cost Ratio (BC Ratio)			1:1.46	

Adequacy of fodder: The data regarding the adequacy of fodder in Yadgir Rf-1 micro-watershed is presented in Table 32. The results indicate that, 45.71 per cent of the households opined that dry fodder was adequate and 48.57 per cent of the households opined that green fodder was adequate.

Table 32. Adequacy of fodder in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (3)		MF (16)		SF (11)		SMF (5)		All (35)	
		N	%	N	%	N	%	N	%	N	%
1	Adequate-Dry Fodder	0	0	6	37.50	5	45.45	5	100	16	45.71
2	Adequate-Green Fodder	0	0	7	43.75	5	45.45	5	100	17	48.57

Annual gross income: The data regarding the annual gross income in Yadgir Rf-1 micro-watershed is presented in Table 33. The results indicate that the annual gross income was Rs. 108,333.33 for landless farmers, for marginal farmers it was Rs. 105,785, for small farmers it was Rs. 179,718.18 and for semi medium farmers it was Rs. 191,928.

Table 33. Annual gross income in Yadgir Rf-1 micro-watershed (Avg value in Rs.)

Sl.No.	Particulars	LL (3)	MF (16)	SF (11)	SMF (5)	All (35)
1	Service/salary	0	0	40,909.09	30,000	17,142.86
2	Business	66,666.67	6,250	18,181.82	0	14,285.71
3	Wage	41,666.67	56,750	56,818.18	50,000	54,514.29
4	Agriculture	0	34,475	61,845.45	97,240	49,088.57
5	Dairy Farm	0	5,185	1,963.64	7,488	4,057.14
6	Goat Farming	0	3,125	0	7,200	2,457.14
Income(Rs.)		108,333.33	105,785	179,718.18	191,928	141,545.71

Average annual expenditure: The data regarding the average annual expenditure in Yadgir Rf-1 micro-watershed is presented in Table 34. The results indicate that the average annual expenditure is Rs. 18,655.77. For landless households it was Rs. 45,000, for marginal farmers it was Rs. 9,227.20, for small farmers it was Rs. 16,506.06 and for semi medium farmers it was Rs. 37,750.

Table 34. Average annual expenditure in Yadgir Rf-1 micro-watershed

(Avg value in Rs.)

Sl.No.	Particulars	LL (3)	MF (16)	SF (11)	SMF (5)	All (35)
1	Service/salary	0	0	73,333.33	85,000	8,714.29
2	Business	100,000	65,000	45,000	0	7,285.71
3	Wage	35,000	34,928.57	25,800	25,600	27,000
4	Agriculture	0	19,126.67	33,500	52,400	26,211.43
5	Dairy Farm	0	8,580	3,933.33	5,750	2,220
6	Goat Farming	0	20,000	0	20,000	1,142.86
Total		135,000	147,635.24	181,566.67	188,750	652,951.90
Average		45,000	9,227.20	16,506.06	37,750	18,655.77

Horticulture species grown: The data regarding horticulture species grown in Yadgir Rf-1 micro-watershed is presented in Table 35. The results indicate that, sampled households have grown 2 coconut and 17 custard apple and mango tree in their field and also 2 coconut, custard apple and guava trees in their backyard.

Table 35. Horticulture species grown in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (3)		MF (16)		SF (11)		SMF (5)		All (35)	
		F	B	F	B	F	B	F	B	F	B
1	Coconut	0	0	1	0	0	0	1	2	2	2
2	Custard apple	0	0	15	2	2	0	0	0	17	2
3	Guava	0	0	0	0	0	2	0	0	0	2
4	Mango	0	0	12	0	0	0	5	0	17	0

*F= Field B=Back Yard

Forest species grown: The data regarding forest species grown in Yadgir Rf-1 micro-watershed is presented in Table 36. The results indicate that, households have planted 65 neem and 2 tamarind trees in their field and also 26 neem trees in their backyard.

Table 36: Forest species grown in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (3)		MF (16)		SF (11)		SMF (5)		All (35)	
		F	B	F	B	F	B	F	B	F	B
1	Neem	0	0	15	5	16	12	34	9	65	26
2	Tamarind	0	0	0	0	2	0	0	0	2	0

*F= Field B=Back Yard

Average Additional investment capacity: The data regarding average additional investment capacity in Yadgir Rf-1 micro-watershed is presented in Table 37. The results indicated that, households have an average investment capacity of Rs. 3,428.57 for land development, Rs. 15,285.71 for irrigation facility and Rs. 285.71 for improved crop production.

Table 37: Source of funds for additional investment capacity in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (3)	MF (16)	SF (11)	SMF (5)	All (35)
1	Land development	0	5,000	3,636.36	0	3,428.57
2	Irrigation facility	0	12,500	18,181.82	27,000	15,285.71
3	Improved crop production	0	625	0	0	285.71

Source of additional investment: The data regarding source of funds for additional investment in Yadgir Rf-1 micro-watershed is presented in Table 38. The results indicated that loan from bank was the source of additional investment for 14.29 per cent for land development, 28.57 for irrigation facility and for 2.86 per cent of irrigation facility.

Table 38: Source of funds for additional investment capacity in Yadgir Rf-1 micro-watershed

Sl.No	Item	Land development		Irrigation facility		Improved crop production	
		N	%	N	%	N	%
1	Loan from bank	5	14.29	10	28.57	1	2.86

Marketing of the agricultural produce: The data regarding marketing of the agricultural produce in Yadgir Rf-1 micro-watershed is presented in Table 39. The results indicated that, green gram was sold to the extent of 60 per cent, groundnut was sold to the

extent of 71.43 per cent, paddy was sold to the extent of 26.83 per cent and red gram to the extent of 61.98 per cent.

Table 39. Marketing of the agricultural produce in Yadgir Rf-1 micro-watershed

Sl.No	Crops	Output obtained (q)	Output retained (q)	Output sold (q)	Output sold (%)	Avg. Price obtained (Rs/q)
1	Green gram	5	2	3	60	5000.0
2	Groundnut	42	12	30	71.43	4900.0
3	Paddy	41	30	11	26.83	1850.0
4	Red gram	263	100	163	61.98	4600.0

Marketing Channels used for sale of agricultural produce: The data regarding marketing channels used for sale of agricultural produce in Yadgir Rf-1 micro-watershed is presented in Table 40. The results indicated that, about 82.86 per cent of the farmers sold their produce to local/village merchants and 11.43 per cent of the farmers sold their produce to regulated markets.

Table 40. Marketing Channels used for sale of agricultural produce in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (3)		MF (16)		SF (11)		SMF (5)		All (35)	
		N	%	N	%	N	%	N	%	N	%
1	Local/village Merchant	0	0	14	87.50	12	109.09	3	60	29	82.86
2	Regulated Market	0	0	1	6.25	0	0	3	60	4	11.43

Mode of transport of agricultural produce: The data regarding mode of transport of agricultural produce in Yadgir Rf-1 micro-watershed is presented in Table 41. The results indicated that, 94.29 per cent of the households have used tractor as a mode of transportation.

Table 41. Mode of transport of agricultural produce in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (3)		MF (16)		SF (11)		SMF (5)		All (35)	
		N	%	N	%	N	%	N	%	N	%
1	Tractor	0	0	15	93.75	12	109.09	6	120	33	94.29

Incidence of soil and water erosion problems: The data regarding incidence of soil and water erosion problems in Yadgir Rf-1 micro-watershed is presented in Table 42. The results indicated that, 42.86 per cent of the households have experienced soil and water erosion problems in the farm.

Table 42. Incidence of soil and water erosion problems in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (3)		MF (16)		SF (11)		SMF (5)		All (35)	
		N	%	N	%	N	%	N	%	N	%
1	Soil and water erosion problems in the farm	0	0	8	50	5	45.45	2	40	15	42.86

Interest shown towards soil testing: The data regarding Interest shown towards soil testing in Yadgir Rf-1 micro-watershed is presented in Table 43. The results indicated that, 85.71 per cent have shown interest in soil test.

Table 43. Interest shown towards soil testing in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (3)		MF (16)		SF (11)		SMF (5)		All (35)	
		N	%	N	%	N	%	N	%	N	%
1	Interest in soil test	0	0	16	100	10	90.91	4	80	30	85.71

Usage pattern of fuel for domestic use: The data regarding usage pattern of fuel for domestic use in Yadgir Rf-1 micro-watershed is presented in Table 44. The results indicated that, 97.14 per cent of the households used firewood and 2.86 per cent of them used LPG as a source of fuel.

Table 44. Usage pattern of fuel for domestic use in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (3)		MF (16)		SF (11)		SMF (5)		All (35)	
		N	%	N	%	N	%	N	%	N	%
1	Fire Wood	3	100	15	93.75	11	100	5	100	34	97.14
2	LPG	0	0	1	6.25	0	0	0	0	1	2.86

Source of drinking water: The data regarding source of drinking water in Yadgir Rf-1 micro-watershed is presented in Table 45. The results indicated that, piped supply was the major source of drinking water for 100 per cent of the households in the micro watershed.

Table 45. Source of drinking water in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (3)		MF (16)		SF (11)		SMF (5)		All (35)	
		N	%	N	%	N	%	N	%	N	%
1	Piped supply	3	100	16	100	11	100	5	100	35	100

Source of light: The data regarding source of light in Yadgir Rf-1 micro-watershed is presented in Table 46. The results indicated that, Electricity was the major source of light for 100 per cent of the households in micro watershed.

Table 46. Source of light in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (3)		MF (16)		SF (11)		SMF (5)		All (35)	
		N	%	N	%	N	%	N	%	N	%
1	Electricity	3	100	16	100	11	100	5	100	35	100

Existence of Sanitary toilet facility: The data regarding existence of sanitary toilet facility in Yadgir Rf-1 micro-watershed is presented in Table 47. The results indicated that, 57.14 per cent of the households possess sanitary toilet facility.

Table 47. Existence of Sanitary toilet facility in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (3)		MF (16)		SF (11)		SMF (5)		All (35)	
		N	%	N	%	N	%	N	%	N	%
1	Sanitary toilet facility	2	66.67	10	62.50	5	45.45	3	60	20	57.14

Possession of PDS card: The data regarding possession of PDS card in Yadgir Rf-1 micro-watershed is presented in Table 48. The results indicated that, 100 per cent of the sampled households possessed BPL cards.

Table 48. Possession of PDS card in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (3)		MF (16)		SF (11)		SMF (5)		All (35)	
		N	%	N	%	N	%	N	%	N	%
1	BPL	3	100	16	100	11	100	5	100	35	100

Participation in NREGA program: The data regarding participation in NREGA programme in Yadgir Rf-1 micro-watershed is presented in Table 49. The results indicated that, 82.86 per cent of the households participated in NREGA programme.

Table 49. Participation in NREGA programme in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (3)		MF (16)		SF (11)		SMF (5)		All (35)	
		N	%	N	%	N	%	N	%	N	%
1	Participation in NREGA programme	3	100	14	87.50	9	81.82	3	60	29	82.86

Adequacy of food items: The data regarding adequacy of food items in Yadgir Rf-1 micro-watershed is presented in Table 50. The results indicated that, cereals, milk, egg and meat were adequate for 100 per cent of the households, pulses were adequate for 94.29 per cent, oilseeds were adequate for 5.71 per cent, vegetables were adequate for 71.43 per cent and fruits were adequate for 17.14 per cent.

Table 50. Adequacy of food items in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (3)		MF (16)		SF (11)		SMF (5)		All (35)	
		N	%	N	%	N	%	N	%	N	%
1	Cereals	3	100	16	100	11	100	5	100	35	100
2	Pulses	3	100	14	87.50	11	100	5	100	33	94.29
3	Oilseed	1	33.33	0	0	0	0	1	20	2	5.71
4	Vegetables	2	66.67	10	62.50	5	45.45	1	20	18	51.43
5	Fruits	0	0	3	18.75	3	27.27	0	0	6	17.14
6	Milk	3	100	16	100	11	100	5	100	35	100
7	Egg	3	100	16	100	11	100	5	100	35	100
8	Meat	3	100	16	100	11	100	5	100	35	100

Response on Inadequacy of food items: The data regarding inadequacy of food items in Yadgir Rf-1 micro-watershed is presented in Table 51. The results indicated that, pulses were inadequate for 5.71 per cent of the households, oilseeds were inadequate for 94.29 per cent, vegetables were inadequate for 51.43 per cent and fruits were inadequate for 82.86 per cent of the households.

Table 51. Response on Inadequacy of food items in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (3)		MF (16)		SF (11)		SMF (5)		All (35)	
		N	%	N	%	N	%	N	%	N	%
1	Pulses	0	0	2	12.50	0	0	0	0	2	5.71
2	Oilseed	2	66.67	16	100	11	100	4	80	33	94.29
3	Vegetables	1	33.33	7	43.75	6	54.55	4	80	18	51.43
4	Fruits	3	100	13	81.25	8	72.73	5	100	29	82.86

Farming constraints: The data regarding farming constraints experienced by households in Yadgir Rf-1 micro-watershed is presented in Table 52. The results indicated that, lower fertility status of the soil, frequent incidence of pest and diseases, inadequacy of irrigation

water and high cost of fertilizers and plant protection chemicals was the constraint experienced by 91.43 per cent of the households, wild animal menace on farm field and high rate of interest on credit (88.57%), low price for the agricultural commodities (11.43%), lack of marketing facilities in the area and inadequate extension services (2.86%).

Table 52. Farming constraints Experienced in Yadgir Rf-1 micro-watershed

Sl. No.	Particulars	MF (16)		SF (11)		SMF (5)		All (35)	
		N	%	N	%	N	%	N	%
1	Lower fertility status of the soil	16	100	11	100	5	100	32	91.43
2	Wild animal menace on farm field	15	93.75	11	100	5	100	31	88.57
3	Frequent incidence of pest and diseases	16	100	11	100	5	100	32	91.43
4	Inadequacy of irrigation water	16	100	11	100	5	100	32	91.43
5	High cost of Fertilizers and plant protection chemicals	16	100	11	100	5	100	32	91.43
6	High rate of interest on credit	16	100	10	90.91	5	100	31	88.57
7	Low price for the agricultural commodities	1	6.25	2	18.18	1	20	4	11.43
8	Lack of marketing facilities in the area	1	6.25	0	0	0	0	1	2.86
9	Inadequate extension services	1	6.25	0	0	0	0	1	2.86

SUMMARY

In order to assess the socio-economic condition of the farmers in the watershed a comprehensive questionnaire was prepared. Major components such as demographic conditions, migration details, food consumption and family expenditure pattern, material possession, land holding, land use management, cropping pattern, cost of cultivation of crops, livestock management. The statistical components such as frequency and percentage were used to analyze the data. About 35 households located in the micro watershed were interviewed for the survey.

The data indicated that there were 111(55.78%) men and 88 (44.22%) women among the sampled households. The average family size of landless farmers' was 4.6, marginal farmers' was 5.3, small farmers' was 5.45 and semi medium farmers' was 7.8. The data indicated that, 32 (16.08%) people were in 0-15 years of age, 91 (45.73%) were in 16-35 years of age, 44 (22.11%) were in 36-60 years of age and 32 (16.08%) were above 61 years of age.

The results indicated that Yadgir Rf-1 had 51.76 per cent illiterates, 0.50 per cent of them had functional illiterate, diploma, ITI and masters, 12.06 per cent of them had primary school, 6.03 per cent of them had middle school and degree education, 11.56 per cent of them had high school education and 4.52 per cent of them had PUC education.

The results indicate that agriculture was the major occupation for 25.13 per cent of the household members, 30.65 per cent were agricultural labourers, 1.51 per cent were in general labour, 4.02 per cent were private service, 4.52 per cent were trade and business, 12.56 per cent were students, 15.08 per cent were housewives and 6.53 per cent were children.

The results show that, 100 per cent of the population in the micro watershed has not participated in any local institutions. The results indicate that 77.14 per cent of the households possess katcha house and 22.86 per cent of them possess pucca/RCC house.

The results show that 80 per cent of the households possess TV, 11.43 per cent of the households possess mixer/grinder and bicycle, 2.86 per cent of the households possess refrigerator, 20 per cent of the households possess motor cycle and 100 per cent of the households possess mobile phones. The results show that the average value of television was Rs. 8,785, mixer/grinder was Rs.2000, refrigerator was Rs. 8,000, bicycle was Rs. 2,325, motor cycle was Rs.50,714 and mobile phone was Rs.2,787.

About 2.86 per cent of the households possess bullock cart, 42.86 per cent of them possess plough, 2.86 per cent of them possess seed/fertilizer drill and 40 per cent of them possess weeder. The results show that the average value of bullock cart was Rs. 19,000,

plough was Rs. 2,426, seed/fertilizer drill was Rs. 2,000 and the average value of weeder was Rs.51.

The results indicate that, 34.29 per cent of the households possess bullocks, 31.43 per cent of the households possess local cow, 11.43 per cent of the households possess buffalo, 5.71 per cent of the households possess goat and poultry birds.

The results indicate that, average own labour men available in the micro watershed was 2.31, average own labour (women) available was 1.66, average hired labour (men) available was 14.59 and average hired labour (women) available was 11.22. The results indicate that, 91.43 per cent of the households opined that the hired labour was adequate.

The results indicate that, households of the Yadgir Rf-1 micro-watershed possess 32.95 ha (93.24%) of dry land, 1.98 ha (5.61%) of irrigated land and 0.40 ha (1.15%) of permanent fallow land. Marginal farmers possess 9.17 ha (95.13%) of dry land, 0.06 ha (1.92%) and 0.4 ha (4.20%) of permanent fallow land. Small farmers possess 11.52 ha (85.73%) of dry land and 1.92 ha (14.27%) of irrigated land. Semi medium farmers possess 12.26 ha (100%) of dry land.

The results indicate that, the average value of dry land was Rs. 646,088.67; the average value of irrigated land was Rs. 1,310,612.24 and the average value of permanent fallow land was Rs. 1,976,000. In case of marginal famers, the average land value was Rs. 1,144,022.94 for dry land, the average value of irrigated land was Rs. 6,175,000.14 and the average value of permanent fallow land was Rs. 1,976,000. In case of small famers, the average land value was Rs. 563,926.94 for dry land and Rs. 1,146,413.50 for irrigated land. In case of semi medium famers, the average land value was Rs. 350,643.78 for dry land.

The results indicate that, there were 4 functioning bore wells in the micro watershed. The results indicate that, bore well was the major irrigation source in the micro water shed for 11.43 per cent of the farmers. The results indicate that, the depth of bore well was found to be 12.19 meters.

The results indicate that, marginal and small farmers had an irrigated area of 0.06 ha and 1.92 ha respectively. The results indicate that, farmers have grown red gram (31.5 ha), groundnut (2.44 ha), paddy (0.61 ha) and green gram (4.02 ha). Marginal farmers have grown red gram, paddy and green gram. Small farmers have grown red gram, groundnut and paddy. Semi medium farmers have grown red gram and groundnut. The results indicate that, the cropping intensity in Yadgir Rf-1 micro-watershed was found to be 99.20 per cent.

The results indicate that, the total cost of cultivation for red gram was Rs. 31285.33. The gross income realized by the farmers was Rs. 61970.38. The net income from red gram cultivation was Rs. 30685.06. Thus the benefit cost ratio was found to be 1:1.98. The total cost of cultivation for green gram was Rs. 70223.42. The gross income realized by the farmers was Rs. 61750. The net income from green gram cultivation was Rs. -8473.42. Thus the benefit cost ratio was found to be 1:0.88. The total cost of cultivation for paddy was Rs. 292727.65. The gross income realized by the farmers was Rs. 310942.70. The net income from paddy cultivation was Rs. 18215.05. Thus the benefit cost ratio was found to be 1:1.06. The total cost of cultivation for groundnut was Rs. 69276.29. The gross income realized by the farmers was Rs. 100829.94. The net income from groundnut cultivation was Rs. 31553.65. Thus the benefit cost ratio was found to be 1:1.46.

The results indicate that, 45.71 per cent of the households opined that dry fodder was adequate and 48.57 per cent of the households opined that green fodder was adequate.

The results indicate that the annual gross income was Rs. 108,333.33 for landless farmers, for marginal farmers it was Rs. 105,785, for small farmers it was Rs. 179,718.18 and for semi medium farmers it was Rs. 191,928. The results indicate that the average annual expenditure is Rs. 18,655.77. For landless households it was Rs. 45,000, for marginal farmers it was Rs. 9,227.20, for small farmers it was Rs. 16,506.06 and for semi medium farmers it was Rs. 37,750.

The results indicate that, sampled households have grown 2 coconut and 17 custard apple and mango tree in their field and also 2 coconut, custard apple and guava trees in their backyard. The results indicate that, households have planted 65 neem and 2 tamarind trees in their field and also 26 neem trees in their backyard.

The results indicated that, households have an average investment capacity of Rs. 3,428.57 for land development, Rs. 15,285.71 for irrigation facility and Rs. 285.71 for improved crop production. The results indicated that loan from bank was the source of additional investment for 14.29 per cent for land development, 28.57 for irrigation facility and for 2.86 per cent of irrigation facility.

The results indicated that, green gram was sold to the extent of 60 per cent, groundnut was sold to the extent of 71.43 per cent, paddy was sold to the extent of 26.83 per cent and red gram to the extent of 61.98 per cent.

The results indicated that, about 82.86 per cent of the farmers sold their produce to local/village merchants and 11.43 per cent of the farmers sold their produce to regulated markets. The results indicated that, 94.29 per cent of the households have used tractor as a mode of transportation.

The results indicated that, 42.86 per cent of the households have experienced soil and water erosion problems in the farm. The results indicated that, 85.71 per cent have shown interest in soil test.

The results indicated that, 97.14 per cent of the households used firewood and 2.86 per cent of them used LPG as a source of fuel. The results indicated that, piped supply was the major source of drinking water for 100 per cent of the households in the micro watershed.

Electricity was the major source of light for 100 per cent of the households in micro watershed. The results indicated that, 57.14 per cent of the households possess sanitary toilet facility. The results indicated that, 100 per cent of the sampled households possessed BPL cards. The results indicated that, 82.86 per cent of the households participated in NREGA programme.

The results indicated that, cereals, milk, egg and meat were adequate for 100 per cent of the households, pulses were adequate for 94.29 per cent, oilseeds were adequate for 5.71 per cent, vegetables were adequate for 71.43 per cent and fruits were adequate for 17.14 per cent.

The results indicated that, pulses were inadequate for 5.71 per cent of the households, oilseeds were inadequate for 94.29 per cent, vegetables were inadequate for 51.43 per cent and fruits were inadequate for 82.86 per cent of the households.

The results indicated that, lower fertility status of the soil, frequent incidence of pest and diseases, inadequacy of irrigation water and high cost of fertilizers and plant protection chemicals was the constraint experienced by 91.43 per cent of the households, wild animal menace on farm field and high rate of interest on credit (88.57%), low price for the agricultural commodities (11.43%), lack of marketing facilities in the area and inadequate extension services (2.86%).