



हर कदम, हर डगर
किसानों का हमसफर
भारतीय कृषि अनुसंधान परिषद

Agrisearch with a human touch

LAND RESOURCE INVENTORY AND SOCIO-ECONOMIC STATUS OF FARM HOUSEHOLDS FOR WATERSHED PLANNING AND DEVELOPMENT

BACHANAHALLI-1 (4B3E2E1e) MICROWATERSHED

Gundlupet Taluk, Chamarajanagara District, Karnataka

Karnataka Watershed Development Project – II

SUJALA – III

World Bank funded Project



ICAR – NATIONAL BUREAU OF SOIL SURVEY AND LAND USE PLANNING



WATERSHED DEVELOPMENT DEPARTMENT
GOVT. OF KARNATAKA, BANGALORE



About ICAR - NBSS&LUP

The ICAR-National Bureau of Soil Survey and Land Use Planning (ICAR-NBSS&LUP), Nagpur, a premier Institute of the Indian Council of Agricultural Research (ICAR), was set up during 1976 with the objective to prepare soil resource maps at national, state and district levels and to provide research inputs in soil resource mapping and its applications, land evaluation, land use planning, land resource management, and database management using GIS for optimising land use on different kinds of soils in the country.

The Bureau has been engaged in carrying out soil resource survey, agro-ecological and soil degradation mapping at the country, state and district levels for qualitative assessment and monitoring the soil health towards viable land use planning. The research activities have resulted in identifying the soil potentials and problems, and the various applications of the soil surveys with the ultimate objective of sustainable agricultural development. The Bureau has the mandate to correlate and classify soils of the country and maintain a National Register of all the established soil series. The Institute is also imparting in-service training to staff of the soil survey agencies in the area of soil survey, land evaluation and soil survey interpretations for land use planning. The Bureau in collaboration with Panjabrao Krishi Vidyapeeth, Akola is running post-graduate teaching and research programme in land resource management, leading to M.Sc. and Ph.D. degrees.

Citation: Rajendra Hegde, Ramesh Kumar, S.C., K.V. Niranjana, S. Srinivas, M.Lalitha, B.A. Dhanorkar, R.S. Reddy and S.K. Singh (2019). "Land Resource Inventory and Socio-Economic Status of Farm Households for Watershed Planning and Development of Bachanahalli-1 (4B3E2E1e) Microwatershed, Gundlupet Taluk and Chamrajangar District, Karnataka", ICAR-NBSS&LUP Sujala MWS Publ.217, ICAR – NBSS & LUP, RC, Bangalore. p.107 & 31.

TO OBTAIN COPIES,

Please write to:

Director, ICAR - NBSS & LUP,

Amaravati Road, NAGPUR - 440 033, India

Phone : (0712) 2500386, 2500664, 2500545 (O)

Telefax : 0712-2522534

E-Mail : director@nbsslup.ernet.in

Website URL : nbsslup.in

Or

Head, Regional Centre, ICAR - NBSS&LUP, Hebbal, Bangalore - 560 024

Phone : (080) 23412242, 23510350 (O)

Telefax : 080-23510350

E-Mail : nbssrcb@gmail.com



LAND RESOURCE INVENTORY AND SOCIO-ECONOMIC STATUS OF FARM HOUSEHOLDS FOR WATERSHED PLANNING AND DEVELOPMENT

BACHANAHALLI-1 (4B3E2E1e) MICRO WATERSHED

Gundlupet Taluk, Chamarajanagara District, Karnataka

Karnataka Watershed Development Project – II Sujala-III

World Bank funded Project



ICAR – NATIONAL BUREAU OF SOIL SURVEY AND LAND USE
PLANNING



ICAR - NBSS & LUP



WATERSHED DEVELOPMENT DEPARTMENT, GOVT. OF
KARNATAKA, BANGALORE



PREFACE

In Karnataka, as in other Indian States, the livelihoods of rural people are intertwined with farming pursuits. The challenges in agriculture are seriously threatening the livelihood of a large number of farmers as they have been practicing farming in contextual factors beyond their control. Climatic factors are the most important ones and have become much more significant in recent times due to rapid climate changes induced by intensive anthropogenic activities affecting our ecosystem in multiple ways. Climate change has become the reality, it is happening and efforts to evolve and demonstrate climate resilient technologies have become essential. Due to the already over stressed scenario of agrarian sector, the climate change is resulting in manifold increase in the complexities, pushing the rural mass to face more and more unpredictable situations. The rising temperatures and unpredictable rainfall patterns are going to test seriously the informed decisions farmers have to make in order to survive in farming and sustain their livelihood.

It is generally recognized that impacts of climate change shall not be uniform across the globe. It is said that impact of climate change is more severe in South Asia. Based on the analysis of meteorological data, it is predicted that in India, there will be upward trend in mean temperature, downward trend in relative humidity, annual rainfall and number of wet days in a year. Also, in general, phenomena like erratic monsoon, spread of tropical diseases, rise in sea levels, changes in availability of fresh water, frequent floods, droughts, heat waves, storms and hurricanes are predicted. Each one of these adverse situations are already being experienced in various parts of India and also at the global level. Decline in agricultural productivity of small and marginal farmers becoming more vulnerable is already witnessed.

In Karnataka, more than 60 per cent of the population live in rural areas and depend on agriculture and allied activities for their livelihood. Though the state has achieved significant progress in increasing the yield of many crops, there is tremendous pressure on the land resources due to the growing and competing demands of various land uses. This is reflected in the alarming rate of land degradation observed. Already more than 50 per cent of the area is affected by various forms of degradation. If this trend continues, the sustainability of the fragile ecosystem will be badly affected. The adverse effects of change in the climatic factors are putting additional stress on the land resources and the farmers dependent on this.

The natural resources (land, water and vegetation) of the state need adequate and constant care and management, backed by site-specific technological interventions and investments particularly by the government. Detailed database pertaining to the nature of the land resources, their constraints, inherent potentials and suitability for various land

based rural enterprises, crops and other uses is a prerequisite for preparing location-specific action plans, which are in tune with the inherent capability of the resources. Any effort to evolve climate resilient technologies has to be based on the baseline scientific database. Then only one can expect effective implementation of climate resilient technologies, monitor the progress, make essential review of the strategy, and finally evaluate the effectiveness of the implemented programs. The information available at present on the land resources of the state are of general nature and useful only for general purpose planning. Since the need of the hour is to have site-specific information suitable for farm level planning and detailed characterization and delineation of the existing land resources of an area into similar management units is the only option.

ICAR-NBSS&LUP, Regional Centre, Bangalore has taken up a project sponsored by the Karnataka Watershed Development Project-II, (Sujala-III), Government of Karnataka funded by the World Bank under Component-1 Land Resource Inventory. This study was taken up to demonstrate the utility of such a database in reviewing, monitoring and evaluating all the land based watershed development programs on a scientific footing. To meet the requirements of various land use planners at grassroots level, the present study on “Land Resource Inventory and Socio-Economic Status of Farm Households for Watershed Planning and Development of Bachanahalli-1 Microwatershed, Gundlupet Taluk and Chamarajanagar District, Karnataka” for integrated development was taken up in collaboration with the State Agricultural Universities, IISC, KRSRAC, KSNDMC as Consortia partners. The project provides detailed land resource information at cadastral level (1:7920 scale) for all the plots and socio-economic status of farm households covering thirty per cent farmers randomly selected representing landed and landless class of farmers in the microwatershed. The project report with the accompanying maps for the Microwatershed will provide required detailed database for evolving effective land use plan, alternative land use options and conservation plans for the planners, administrators, agricultural extension personnel, KVK officials, developmental departments and other land users to manage the land resources in a sustainable manner.

It is hoped that this database will be useful to the planners, administrators and developmental agencies working in the area in not only for formulating location specific developmental schemes but also for their effective monitoring at the village/watershed level.

Nagpur

Date: 29.12.2018

S.K. SINGH

Director, ICAR - NBSS&LUP, Nagpur

Contributors

Dr. Rajendra Hegde Principal Scientist, Head & Project Leader, Sujala-III Project ICAR-NBSS&LUP, Regional Centre, Bangalore	Dr. S.K.Singh Director, ICAR-NBSS&LUP Coordinator, Sujala-III Project Nagpur
Soil Survey, Mapping & Report Preparation	
Dr. K.V. Niranjana	Sh. R.S. Reddy
Dr. B.A. Dhanorkar	Sh. Nagendra, B.R.
Field Work	
Sh. C.Bache Gowda	Sh. Sandesh Shastri
Sh. Somashekar	Sh. Rajeev, G.S.
Sh. Venkata Giriappa	Sh. Balasubramanyam, M.G.
Sh. M. Jayaramaiah	Sh. Vijaya Kumar
Sh. Paramesha, K.	Sh. Mayur Patil
	Sh. Kamalesh K. Avate
GIS Work	
Dr. S.Srinivas	Sh. A.G. Devendra Prasad
Sh. D.H.Venkatesh	Sh. Prakashanaik, M.K.
Smt.K.Sujatha	Sh. Abhijith Sastry, N.S.
Smt. K.V.Archana	Sh. Mahamad Ali, M.
Sh. N.Maddileti	Sh. Amar Suputhra, S.
	Sh. Avinash, K.N.
	Sh. Anudeep, Y.
	Sh. Sudip Kumar Suklabaidya
	Smt. K.Karunya Lakshmi
	Ms. Seema, K.V.
	Ms. A. Rajab Nisha
Laboratory Analysis	
Dr. K.M.Nair	Smt. Savitha, H.R.
Smt. Arti Koyal	Smt. Steffi Peter
Smt. Parvathy, S.	Smt. Thara, V.R.
	Smt. Roopa, G.
	Ms. Shwetha, N.K.

	Smt. Ishrat Haji
	Ms. Pavana Kumari, P.
	Sh. Shanthaveeraswamy, H.M.
	Sh. Ramesh, K.
	Ms. Padmaja, S.
	Ms. Veena, M.
Soil & Water Conservation	
Sh. Sunil P. Maske	
Socio-Economic Analysis	
Dr. S.C. Ramesh Kumar	Sh. M. K. Prakashanaik
	Ms. Sowmya K.B
	Sh.Manjunath M
	Sh.Veerabhadraswamy R
	Sh.Lankesh RS
	Sh.Kalaveerachari R Kammar
	Sh.Pradyumma U
	Sh.Yogesha HN
	Sh.Vijay kumar lamani
	Sh.Arun N Kambar
	Sh.Vinay
	Sh.Basavaraj.Biradar
	Sh.Vinod R
	Sh.Praveenkumar P Achalkar
	Sh.Rajendra D
Watershed Development Department, GoK, Bangalore	
Sh. Rajeev Ranjan IFS Project Director & Commissioner, WDD	Dr. A. Natarajan NRM Consultant, Sujala-III Project
Dr. S.D. Pathak IFS Executive Director & Chief Conservator of Forests, WDD	

PART-A

LAND RESOURCE INVENTORY

Contents

Preface	
Contributors	
Executive Summary	
Chapter 1	Introduction 1
Chapter 2	Geographical Setting 3
2.1	Location and Extent 3
2.2	Geology 3
2.3	Physiography 4
2.4	Drainage 5
2.5	Climate 5
2.6	Natural Vegetation 6
2.7	Land Utilization 7
Chapter 3	Survey Methodology 11
3.1	Base maps 11
3.2	Image Interpretation for Physiography 11
3.3	Field Investigation 14
3.4	Soil Mapping 16
3.5	Laboratory Characterization 16
3.6	Land Management Units 18
Chapter 4	The Soils 21
4.1	Soils of Granite gneiss Landscape 21
Chapter 5	Interpretation for Land Resource Management 35
5.1	Land Capability Classification 35
5.2	Soil Depth 37
5.3	Surface Soil Texture 38
5.4	Soil Gravelliness 39
5.5	Available Water Capacity 40
5.6	Soil Slope 41
5.7	Soil Erosion 42
Chapter 6	Fertility Status 45
6.1	Soil Reaction (pH) 45
6.2	Electrical Conductivity (EC) 45
6.3	Organic Carbon (OC) 45
6.4	Available Phosphorus 47
6.5	Available Potassium 47
6.6	Available Sulphur 47
6.7	Available Boron 47
6.8	Available Iron 50
6.9	Available Manganese 50
6.10	Available Copper 50

6.11	Available Zinc	50
Chapter 7	Land Suitability for Major Crops	53
7.1	Land suitability for Sorghum	53
7.2	Land suitability for Maize	56
7.3	Land suitability for Redgram	57
7.4	Land suitability for Horsegram	58
7.5	Land suitability for Fieldbean	60
7.6	Land suitability for Groundnut	61
7.7	Land suitability for Sunflower	62
7.8	Land suitability for Cotton	64
7.9	Land suitability for Onion	65
7.10	Land suitability for Potato	66
7.11	Land suitability for Beans	68
7.12	Land suitability for Beetroot	69
7.13	Land suitability for Mango	70
7.14	Land suitability for Sapota	72
7.15	Land suitability for Guava	73
7.16	Land suitability for Banana	75
7.17	Land suitability for Jackfruit	76
7.18	Land Suitability for Jamun	77
7.19	Land Suitability for Musambi	78
7.20	Land Suitability for Lime	79
7.21	Land Suitability for Cashew	81
7.22	Land Suitability for Custard Apple	81
7.23	Land Suitability for Amla	82
7.24	Land Suitability for Tamarind	83
7.25	Land suitability for Marigold	84
7.26	Land suitability for Chrysanthemum	86
7.27	Land suitability for Turmeric	87
7.28	Land Management Units	89
7.29	Proposed Crop Plan	90
Chapter 8	Soil Health Management	93
Chapter 9	Soil and Water conservation Treatment Plan	99
9.1	Treatment Plan	99
9.2	Recommended Soil and Water Conservation measures	103
9.3	Greening of Microwatershed	104
	References	107
	Appendix I	i-vi
	Appendix II	vii-xii
	Appendix III	xiii-xvi

LIST OF TABLES

2.1	Mean Monthly Rainfall, PET, 1/2 PET at Gundlupet Taluk, Chamarajanagar District	5
2.2	Land Utilization in Gundlupet District	7
3.1	Differentiating Characteristics used for Identifying Soil Series	16
3.2	Soil map unit description of Bachanahalli-1 Microwatershed	17
4.1	The physical and chemical characteristics of soil series identified in Bachanahalli-1 microwatershed	27
7.1	Soil-Site Characteristics of Bachanahalli-1 Microwatershed	54
7.2	Crop suitability criteria for Sorghum	55
7.3	Crop suitability criteria for Maize	57
7.4	Land suitability for Red gram	58
7.5	Land suitability for Horse gram	59
7.6	Land suitability for Field bean	60
7.7	Crop suitability criteria for Groundnut	62
7.8	Crop suitability criteria for Sunflower	63
7.9	Crop suitability criteria for Cotton	64
7.10	Land suitability for Onion	66
7.11	Land suitability for Potato	67
7.12	Crop suitability criteria for Mango	71
7.13	Crop suitability criteria for Sapota	72
7.14	Crop suitability criteria for Guava	74
7.15	Crop suitability criteria for Banana	75
7.16	Crop suitability criteria for Lime	80
7.17	Crop suitability criteria for Marigold	85
7.18	Crop suitability criteria for Chrysanthemum	86
7.19	Crop suitability criteria for Turmeric	88
7.20	Proposed Crop Plan for Bachanahalli-1 Microwatershed	91

LIST OF FIGURES

2.1	Location map of Bachanahalli-1 Microwatershed	3
2.2a	Granite and granite gneiss rocks	4
2.2b	Granite rocks	4
2.3	Rainfall distribution in Gundlupet Taluk, Chamarajanagar District	6
2.4	Natural vegetation of Shivapura-1 Microwatershed	6
2.5a	Different crops and cropping systems in Shivapura-1 Microwatershed	8
2.5b	Different crops and cropping systems in Shivapura-1 Microwatershed	9
2.6	Current Land use – Bachanahalli-1 Microwatershed	9
2.7	Location of Wells- Kalagatahundi-2Microwatershed	10
3.1	Scanned and Digitized Cadastral map ofBachanahalli-1 Microwatershed	13
3.2	Satellite image of Bachanahalli-1 Microwatershed	13
3.3	Cadastral map overlaid on IRS PAN+LISS IV merged imagery of Bachanahalli-1 Microwatershed	14
3.4	Location of profiles in a transect	15
3.5	Soil phase or management units of Bachanahalli-1 Microwatershed	19
5.1	Land Capability Classification of Bachanahalli-1 Microwatershed	37
5.2	Soil Depth map of Bachanahalli-1 Microwatershed	38
5.3	Surface Soil Texture map of Bachanahalli-1 Microwatershed	39
5.4	Soil Gravelliness map of Bachanahalli-1 Microwatershed	40
5.5	Soil Available Water Capacity map of Bachanahalli-1 Microwatershed	41
5.6	Soil Slope map of Bachanahalli-1 Microwatershed	42
5.7	Soil Erosion map of Bachanahalli-1 Microwatershed	43
6.1	Soil Reaction (pH) map of Bachanahalli-1 Microwatershed	46
6.2	Electrical Conductivity (EC) map of Bachanahalli-1 Microwatershed	46
6.3	Soil Organic Carbon (OC) map of Bachanahalli-1 Microwatershed	47
6.4	Soil Available Phosphorus map of Bachanahalli-1 Microwatershed	48
6.5	Soil Available Potassium map of Bachanahalli-1 Microwatershed	48
6.6	Soil Available Sulphur map of Bachanahalli-1 Microwatershed	49
6.7	Soil Available Boron map of Bachanahalli-1 Microwatershed	49
6.8	Soil Available Iron map of Bachanahalli-1 Microwatershed	50
6.9	Soil Available Manganese map of Bachanahalli-1 Microwatershed	51
6.10	Soil Available Copper map of Bachanahalli-1 Microwatershed	51

6.11	Soil Available Zinc map of Bachanahalli-1 Microwatershed	52
7.1	Land suitability for Sorghum	55
7.2	Land suitability for Maize	57
7.3	Land suitability for Redgram	58
7.4	Land suitability for Horsegram	59
7.5	Land suitability for Fieldbean	61
7.6	Land suitability for Groundnut	62
7.7	Land suitability for Sunflower	63
7.8	Land suitability for Cotton	65
7.9	Land suitability for Onion	66
7.10	Land suitability for Potato	68
7.11	Land suitability for Beans	69
7.12	Land suitability for Beetroot	70
7.13	Land suitability for Mango	71
7.14	Land suitability for Sapota	73
7.15	Land suitability for Guava	74
7.16	Land suitability for Banana	76
7.17	Land suitability for Jackfruit	77
7.18	Land Suitability for Jamun	78
7.19	Land Suitability for Musambi	79
7.20	Land Suitability for Lime	80
7.21	Land Suitability for Cashew	81
7.22	Land Suitability for Custard Apple	82
7.23	Land Suitability for Amla	83
7.24	Land Suitability for Tamarind	84
7.25	Land suitability for Marigold	85
7.26	Land suitability for Chrysanthemum	87
7.27	Land suitability for Turmeric	88
7.28	Land Management Units map of Bachanahalli-1 Microwatershed	90
9.1	Soil and water conservation map of Bachanahalli-1 Microwatershed	104

EXECUTIVE SUMMARY

The land resource inventory of Bachanahalli-1 Microwatershed was conducted using village cadastral maps and IRS satellite imagery on 1:7920 scale. The false colour composites of IRS imagery were interpreted for physiography and the physiographic delineations were used as base for mapping soils. The soils were studied in several transects and a soil map was prepared with phases of soil series as mapping units. Random checks were made all over the area outside the transects to confirm and validate the soil map unit boundaries. The soil map shows the geographic distribution and extent, characteristics, classification and use potentials of the soils in the microwatershed.

The present study covers an area of 567 ha in Gundlupet taluk of Chamarajanagar district, Karnataka. The climate is semiarid and categorized as drought prone with an average annual rainfall of 734 mm of which about 254 mm is received during south-west monsoon, 268 mm during north-east and the remaining 212 mm during the rest of the year. An area of about 66 per cent is covered by soils, 31 per cent by forest and 2 per cent by others. The salient findings from the land resource inventory are summarized briefly below.

- ❖ *The soils belong to 8 soil series and 20 soil phases (management units) and 7 land management units.*
- ❖ *The length of crop growing period is about 150 days starting from the 3rd week of June to 3rd week of November.*
- ❖ *From the master soil map, several interpretative and thematic maps like land capability, soil depth, surface soil texture, soil gravelliness, available water capacity, soil slope and soil erosion were generated.*
- ❖ *Soil fertility status maps for macro and micronutrients were generated based on the surface soil samples collected at every 320 m grid interval.*
- ❖ *Land suitability for growing major agricultural and horticultural crops were assessed and maps showing the degree of suitability along with constraints were generated.*
- ❖ *About 66 per cent area is suitable for agriculture.*
- ❖ *About 10 per cent soils are shallow (25-50 cm), 2 per cent of soils are moderately shallow (50-75 cm), 22 per cent of the soils are moderately deep (75-100 cm) and 33 per cent soils are deep (100-150 cm) to very deep (>150 cm).*
- ❖ *About 17 per cent of the area has clayey soils at the surface and 49 per cent area has loamy soils.*
- ❖ *About 35 per cent of the area has non-gravelly soils and 31 per cent gravelly soils (15-35 % gravel) soils.*
- ❖ *About 13 per cent has soils that are very low (<50 mm/m), 20 per cent has low (51-100 mm/m) in available water capacity and an area of 33 per cent has very high (>200 mm/m) available water capacity.*

- ❖ *About 50 per cent of the area has very gently sloping (1-3% slope) lands and 16 per cent is under gently sloping (3-5%) lands.*
- ❖ *An area of about 37 per cent has soils that are slightly eroded (e1) and 29 per cent soils are moderately eroded (e2).*
- ❖ *An area of about 35 per cent has soils that are moderately acidic (pH 5.5-6.0) to slightly acidic (pH 6.0-6.5); 22 per cent area has neutral (pH 6.5-7.3) and an area of about 9 per cent has soils that are slightly alkaline (pH 7.3 to 7.8) to moderately alkaline (pH 7.8 to 8.4).*
- ❖ *The Electrical Conductivity (EC) of the soils are dominantly $<2 \text{ dSm}^{-1}$ indicating 65 per cent soils are non-saline and 2 per cent are low ($2-4 \text{ dSm}^{-1}$).*
- ❖ *About 10 per cent of the soils are low ($<0.5\%$), 50 per cent of the soils are medium ($0.5-0.75\%$) and 6 per cent of the soils are high ($>0.75\%$) in organic carbon.*
- ❖ *Entire area of the microwatershed about 66 per cent of the soils are medium (23-57 kg/ha) in available phosphorus.*
- ❖ *About 7 per cent of the soils are medium (145-337 kg/ha) and 60 per cent are high ($>337 \text{ kg/ha}$) in available potassium.*
- ❖ *Available sulphur is low ($<10 \text{ ppm}$) in entire area of about 66 per cent.*
- ❖ *Available boron is low ($<0.5 \text{ ppm}$) in entire area of about 17 per cent.*
- ❖ *Available iron is sufficient ($>4.5 \text{ ppm}$) in entire area of about 66 per cent.*
- ❖ *Available manganese and copper are sufficient in all the soils of the microwatershed.*
- ❖ *Available zinc is deficient ($<0.6 \text{ ppm}$) in 61 per cent and sufficient ($>0.6 \text{ ppm}$) in about 5 per cent area.*
- ❖ *The land suitability for 27 major crops grown in the microwatershed were assessed and the areas that are highly suitable (S1) and moderately suitable (S2) are given below. It is however to be noted that a given soil may be suitable for various crops but what specific crop to be grown may be decided by the farmer looking to his capacity to invest on various inputs, marketing infrastructure, price and finally the demand and supply position.*

Land suitability for various crops in the Microwatershed

Crop	Suitability Area in ha (%)		Crop	Suitability Area in ha (%)	
	Highly suitable(S1)	Moderately suitable(S2)		Highly suitable(S1)	Moderately suitable(S2)
<i>Sorghum</i>	247 (44)	41 (7)	<i>Sapota</i>	66 (12)	122 (22)
<i>Maize</i>	158 (28)	10 (2)	<i>Guava</i>	67 (12)	122 (22)
<i>Red gram</i>	158 (28)	120 (21)	<i>Banana</i>	66 (12)	211 (37)
<i>Horse gram</i>	158 (28)	10 (2)	<i>Jackfruit</i>	66 (12)	1 (<1)
<i>Field bean</i>	66 (12)	221 (39)	<i>Jamun</i>	186 (33)	1 (<1)
<i>Groundnut</i>	90 (16)	77 (14)	<i>Musambi</i>	186 (33)	1 (<1)
<i>Sunflower</i>	36 (6)	242 (43)	<i>Lime</i>	186 (33)	1 (<1)
<i>Cotton</i>	155 (27)	101 (18)	<i>Cashew</i>	66 (12)	123 (22)
<i>Onion</i>	66 (12)	221 (30)	<i>Custard apple</i>	187 (33)	132 (23)
<i>Potato</i>	66 (12)	101 (18)	<i>Amla</i>	187 (33)	132 (23)
<i>Beans</i>	66 (12)	221 (39)	<i>Tamarind</i>	186 (33)	1 (<1)
<i>Beetroot</i>	66 (12)	101 (18)	<i>Marigold</i>	158 (28)	130 (23)
<i>Turmeric</i>	66 (12)	101 (18)	<i>Chrysanthemum</i>	66 (12)	221 (39)
<i>Mango</i>	66 (12)	1 (<1)			

Apart from the individual crop suitability, a proposed crop plan has been prepared for the 7 identified LMUs by considering only the highly and moderately suitable lands for different crops and cropping systems with food, fibre and horticulture crops.

- ❖ *Maintaining soil-health is vital to crop production and conserve soil and land resource base for maintaining ecological balance and to mitigate climate change. For this, several ameliorative measures have been suggested to these problematic soils like saline/alkali, highly eroded, sandy soils etc.,*
- ❖ *Soil and water conservation treatment plan has been prepared that would help in identifying the sites to be treated and also the type of structures required.*
- ❖ *As part of the greening programme, several tree species have been suggested to be planted in marginal and submarginal lands and also in the hillocks, mounds and ridges.*

INTRODUCTION

Soil is a finite natural resource that is central to sustainable agriculture and food security. Over the years, this precious resource is faced with the problems of erosion, salinity, alkalinity, degradation, depletion of nutrients and even decline in availability of land for agriculture. It is a known fact, that it takes thousands of years to form a few centimetres of soil, thus, soil is a precious gift of nature. The area available for agriculture is about 51 per cent of the total geographical area and more than 60 per cent of the people are still dependant on agriculture for their livelihood. However, the capacity of a soil to produce is limited and the limits to the production are set by its intrinsic characteristics, agroclimatic setting, and use and management. There is therefore, tremendous pressure on land and water resources, which is causing decline in soil-health and stagnation in productivity. As much as 121 m ha of land is reportedly degraded which leads to impaired soil quality. It is imperative that steps are urgently taken to check and reverse land degradation without any further loss of time. The improvements in productivity will have to come from sustainable intensification measures that make the most effective use of land and water resources. Soil erosion alone has degraded about 35 lakh ha. Almost all the areas are facing various degrees of degradation, particularly soil erosion. Salinity and alkalinity has emerged as a major problem (>3.5 lakh ha) in the irrigated areas of the State. Nutrient depletion and declining factor productivity is common in both rainfed and irrigated areas. The degradation is continuing at an alarming rate and there appears to be no systematic effort among the stakeholders to contain this process. In recent times, an aberration of weather due to climate change phenomenon has added another dimension leading to unpredictable situations to be tackled by the farmers.

In this critical juncture, the challenge before us is not only to increase the productivity per unit area which is steadily declining and showing a fatigue syndrome, but also to prevent or at least reduce the severity of degradation. If the situation is not reversed at the earliest, then the sustainability of the already fragile crop production system and the overall ecosystem will be badly affected in the state. Added to this, every year there is a significant diversion of farm lands and water resources for non-agricultural purposes. Thus, developing strategies to slow down the degradation process or reclaim the soils to normal condition and ensure sustainability of production system are the major issues today. This demands a systematic appraisal of our soil and land resources with respect to their extent, geographic distribution, characteristics, behaviour and usepotential, which is very important for developing an effective land use and cropping systems for augmenting agricultural production on a sustainable basis.

The soil and land resource inventories made so far in Karnataka had limited utility because the surveys were of different types, scales and intensities carried out at different times with specific objectives. Hence, there is an urgent need to generate detailed site-specific farm level database on various land resources for all the villages/watersheds in a

time bound manner that would help to protect the valuable soil and land resources and also to stabilize farm production. Therefore, the land resource inventory required for farm level planning is the one which investigates all the parameters which are critical for productivity viz., soils, site characteristics (slope, erosion, gravelliness and stoniness) climate, water, topography, geology, hydrology, vegetation, crops, land use pattern, animal population, socio-economic conditions, infrastructure, marketing facilities and various schemes and developmental works of the government etc. From the data collected at farm level, the specific problems and potentials of the area can be identified and highlighted, conservation measures required for the area can be planned on a scientific footing, suitability of the area for various uses can be worked out and finally viable and sustainable land use options suitable for each and every land holding can be prescribed.

The Land Resource Inventory is basically done for identifying potential and problem areas, developing sustainable land use plans, estimation of surface run off and water harvesting potential, preparation of soil and water conservation plans, land degradation/desertification etc. The Bureau is presently engaged in developing an LRI methodology using high resolution satellite remote sensing data and Digital Elevation Model (DEM) data to prepare Landscape Ecological Units (LEU) map representing agro-ecosystem as a whole. The LEU is preferred over landform as the base for LRI. LEU is the assemblage of landform, slope and land use. An attempt has been made to upscale the soil resource information from 1:250000 and 1:50000 scale to the LEU map in Goa and other states. An attempt will be made later to uplink the LRI data generated under Sujala-III Project to the Landscape Ecological Units (LEUs) map. For this, the major physiographic region, *i.e.*, South Deccan Plateau will be taken as an example.

The land resource inventory data and maps presented here aims to provide site specific database for Bachanahalli-1 microwatershed in Gundlupet Taluk, Chamarajanagar District, Karnataka state for the Karnataka Watershed Development Department. The database was generated by using cadastral map of the village as a base along with high resolution IRS LISS IV and Cartosat-1 merged satellite imagery. Later, an attempt will be made to uplink this LRI data generated at 1:7920 scale under Sujala-III Project to the proposed Landscape Ecological Units (LEUs) map.

The study was organized and executed by the ICAR- National Bureau of Soil Survey and Land Use Planning, Regional Centre, Bangalore under Generation of Land Resource Inventory Data Base Component-1 of the Sujala-III Project funded by the World Bank.

GEOGRAPHICAL SETTING

2.1 Location and Extent

The Bachanahalli-1 microwatershed (Gopalpur subwatershed) is located in the southern part of Karnataka in Gundlupet Taluk, Chamarajanagara District, Karnataka State (Fig. 2.1). It comprises of Bachanahalli, Berambadi and maddur villages. It lies between $11^{\circ}44'$ to $11^{\circ}47'$ North latitudes and $76^{\circ}35'$ to $76^{\circ}38'$ East longitudes and covers an area of 584 ha. It is about 20 km south of Gundlupet and is surrounded by Maddur village on the north, Lakkipur village on the east and Berambadi State Forest in the southwest and southern side.




Fig. 2.1 Location map of Bachanahalli-1 Microwatershed

2.2 Geology

Major rock formations observed in the micowatershed are of Archaean age and comprise of (Fig. 2.2a and 2.2b) granite and gneiss. They are essentially pink to gray

granite gneisses. The rocks are coarse to medium grained. They consist primarily of quartz, feldspar, biotite and hornblende. The gray granite gneisses are highly weathered, fractured and fissured upto a depth of about 10 m. Dolerite dykes and quartz veins are common with variable width and found to occur in the microwatershed.



Fig. 2.2a Granite and granite gneiss rocks



Fig. 2.2b Granite rocks

2.3 Physiography

Physiographically, the area has been broadly divided into one landscape based on granite gneiss geology. The microwatershed area has been further divided into mounds/ridges, summits, side slopes and very gently sloping uplands based on slope and relief features. The mounds and ridges are mostly covered by rock outcrops.

2.4 Drainage

There are no perennial rivers flowing in Gundlupet taluk. However, the area is drained by several small seasonal streams like Gundlu hole along its course. Though, it is not a perennial one, during rainy season, it carries large quantities of rain water. The microwatershed has only a few small tanks which are not capable of storing water that flows during the rainy season. Due to this, the ground water recharge is very much affected in the villages. This is reflected in the failure of many bore wells in the villages. If the available rain water is properly harnessed by constructing tanks and recharge structures at appropriate places in the villages, then the drinking and irrigation needs of the entire area can be easily met. The drainage network is dendritic to sub parallel.

2.5 Climate

The district falls under semiarid tract and is categorized as drought-prone with average annual rainfall of 734 mm (Table 2.1). Of the total rainfall, a maximum of 254 mm is received during south–west monsoon period from June to September, north-east monsoon from October to early December contributes maximum about 268 mm and the remaining 212 mm is received during the rest of the year. The winter season is from December to February. During April and May, the temperatures reach up to 42°C and in December and January, the temperatures will go down to 16°C. Rainfall distribution is shown in Figure 2.3. The average Potential Evapotranspiration (PET) is 128 mm and varies from a low of 106mm in November to 165 mm in the month of March. The PET is always higher than precipitation in all the months except in the last week of September to first week of November. Generally, the length of crop growing period (LGP) is 150 days and starts from 3rd week of June to third week of November.

Table 2.1 Mean Monthly Rainfall, PET, 1/2 PET in Gundlupet Taluk, Chamarajanagara District

Sl. no.	Months	Rainfall	PET	1/2 PET
1	JAN	0.80	129.10	64.55
2	FEB	6.80	133.80	66.90
3	MAR	26.90	164.90	82.45
4	APR	73.60	153.80	76.90
5	MAY	103.90	147.20	73.60
6	JUN	56.00	124.60	62.30
7	JUL	50.40	116.40	58.20
8	AUG	55.80	117.10	58.55
9	SEP	92.00	116.80	58.40
10	OCT	164.10	111.10	55.55
11	NOV	80.50	106.20	53.10
12	DEC	23.50	109.90	54.95
	Total	734.30	127.57	




Fig 2.3 Rainfall distribution in Gundlupet Taluk, Chamarajanagara District

2.6 Natural Vegetation

Forests occupy about 32 per cent area in Gundlupet taluk. The major area of these forests are found in Bandipur National Park and Himavad Gopalaswamy Betta. The rest of the area in the taluk has sparse natural vegetation comprising few tree species, shrubs and herbs. The mounds, ridges and boulders occupy very sizeable areas which are under thin to moderately thick forest vegetation. Still, there are some remnants of the past forest cover which can be seen in patches in some ridges and hillocks in the microwatershed.



Fig. 2.4 Natural vegetation of Bachanahalli-1 Microwatershed

Apart from the continuing deforestation, the presence of large population of goats, sheep and other cattle in the microwatershed is causing vegetative degradation of

whatever little vegetation is left in the area. The uncontrolled grazing has left no time for the regeneration of the vegetative cover. This leads to the accelerated rate of erosion on the hill slopes, resulting in the formation of deep gullies in the foot slopes and eventually resulting in the heavy siltation of few tanks and reservoirs in the microwatershed.

2.7 Land Utilization

About 48 per cent area (Table 2.2) in Gundlupet taluk is cultivated at present. An area of about 6 per cent is currently barren. Forests occupy an area of about 32 per cent and the tree cover is in a very poor state except in Bandipura National Park and Gopalaswamy Betta. Most of the mounds, ridges and bouldery areas have very poor vegetative cover. Major crops grown in the area are sorghum, maize, cotton, onion, sugarcane, groundnut, red gram, horse gram and sapota (Fig. 2.5 a, b & c). While carrying out land resource inventory, the land use/land cover particulars are collected from all the survey numbers and a current land use map of the microwatershed is generated. The current land use map generated shows the arable and non-arable lands, other land uses and different types of crops grown in the area (Fig.2.6). Simultaneously, enumeration of wells (bore wells and open wells) and existing conservation structures in the microwatershed are made and their location in different survey numbers is located on the cadastral map. Map showing the location of wells and other water bodies in Bachanahalli-1 microwatershed is given in Figure 2.7.

Table 2.2 Land Utilization in Gundlupet Taluk

Sl. No.	Agricultural land use	Area (ha)	Per cent
1.	Total geographical area	140607	
2.	Total cultivated area	67339	47.84
3.	Area sown more than once	13532	
4.	Cropping intensity	-	120.09
5.	Trees and grooves	3485	2.47
6.	Forest	44859	31.98
7.	Cultivable wasteland	3265	2.32
8.	Permanent Pasture land	10287	7.31
9.	Barren land	7988	5.68
10.	Non- Agriculture land	3384	2.40




Fig. 2.5.a. Different crops and cropping systems in Bachanahalli -I Microwatershed



Fig. 2.5.b. Different crops and cropping systems in Bachanahalli-1 Microwatershed




Fig. 2.6 Current Land Use map of Bachanahalli-1 Microwatershed




Fig. 2.7 Location of Wells map of Bachanahalli-1 Microwatershed

SURVEY METHODOLOGY

The purpose of land resource inventory is to delineate similar areas (soil series and phases), which respond or expected to respond similarly to a given level of management. This was achieved in Bachanahalli-1 microwatershed by the detailed study of all the soil characteristics (depth, texture, colour, structure, consistence, coarse fragments, porosity, soil reaction, soil horizons etc.), and site characteristics (slope, erosion, drainage, occurrence of rock fragments etc.) and followed by grouping of similar areas based on soil-site characteristics into homogeneous (management units) units, and showing their extent and geographic distribution on the microwatershed cadastral map. The detailed soil survey at 1:7920 scale was carried out in an area of 567 ha. The methodology followed for carrying out land resource inventory was as per the guidelines given in Soil Survey Manual (IARI, 1971; Soil Survey Staff, 2006; Natarajan *et al.*, 2015) which is briefly described below.

3.1 Base Maps

The detailed survey of the land resources occurring in the microwatershed was carried out by using digitized cadastral map as a base supplied by the KSRSAC. The cadastral map shows field boundaries with their survey numbers, location of tanks, streams and other permanent features of the area (Fig. 3.1). Apart from the cadastral map, remote sensing data products from Cartosat-1 and LISS IV merged at the scale of 1:7920 were used in conjunction with the cadastral map to identify the landscapes, landforms and other surface features. The imagery helped in the identification and delineation of boundaries between hills, uplands and lowlands, water bodies, forest and vegetated areas, roads, habitations and other cultural features of the area (Fig. 3.2). The cadastral map was overlaid on the satellite imagery (Fig. 3.3) that helps to identify the parcel boundaries and other permanent features. Apart from cadastral maps and images, toposheets of the area (1:50,000 scale) were also used for initial traversing, identification of geology and landforms, drainage features, present land use and also for selection of transects in the microwatershed.

3.2 Image Interpretation for Physiography

False Colour Composites (FCC) of Cartosat-I and LISS-IV merged satellite data covering the microwatershed area was visually interpreted using image interpretation elements and all the available collateral data with local knowledge. The delineated physiographic boundaries were transferred on to a cadastral map overlaid on satellite imagery. Physiographically, the area has been identified as granite gneiss landscape. They were divided into landforms such as ridges, mounds and uplands based on slope and other relief features. They were further subdivided into physiographic/ image interpretation

units based on image characteristics. The image interpretation legend for physiography is given below.

Image Interpretation Legend for Physiography

G - Granite gneiss landform

G1	Hills/ Ridges/ Mounds
G11	Summits
G12	Side slopes
G121	Side slopes with dark grey tones
G2	Uplands
G21	Summits
G22	Gently sloping uplands
G221	Gently sloping uplands, yellowish green (eroded)
G222	Gently sloping uplands, yellowish white (severely eroded)
G23	Very gently sloping uplands
G231	Very gently sloping uplands, yellowish green
G232	Very gently sloping uplands, medium green and pink
G233	Very gently sloping uplands, pink and green (scrub land)
G234	Very gently sloping uplands, medium greenish grey
G235	Very gently sloping uplands, yellowish white (eroded)
G236	Very gently sloping uplands, dark green
G237	Very gently sloping uplands, medium pink (coconut garden)
G238	Very gently sloping uplands, pink and bluish white (eroded)
G24	Valleys/ lowlands
G241	Valleys, pink tones
G242	Valleys gray mixed with pink tones




Fig 3.1 Scanned and Digitized Cadastral map of Bachanahalli-1 Microwatershed




Fig.3.2 Satellite Image of Bachanahalli-1 Microwatershed




Fig. 3.3 Cadastral map overlaid on IRS PAN+LISS IV merged imagery of Bachanahalli-1 Microwatershed

3.3 Field Investigation

The field boundaries and survey numbers given on the cadastral sheet were located on the ground by following permanent features like roads, cart tracks, *nallas*, streams, tanks etc., and wherever changes were noticed, they were incorporated on the microwatershed cadastral map. Preliminary traverse of the microwatershed was carried out with the help of cadastral map, imagery and toposheets. While traversing, landforms and physiographic units identified were checked and preliminary soil legend was generated by studying soils at few selected places. Then, intensive traversing of each physiographic unit like hills, ridges and uplands was carried out. Based on the variability observed on the surface, transects (Fig. 3.4) were selected across the slope covering all the landform units in the microwatershed (Natarajan and Dipak Sarkar, 2010).




Fig. 3.4 Location of profiles in a transect

In the selected transect (Fig. 3.4), soil profiles were located at closely spaced intervals to take care of any change in the land features like break in slope, erosion, gravel, stones etc. In the selected sites, soil profiles (vertical cut showing the soil layers from surface to the rock) were opened up to 200 cm or to the depth limited by rock or hard substratum and studied in detail for all their morphological and physical characteristics. The soil and site characteristics were recorded for all profile sites on a standard proforma as per the guidelines given in USDA Soil Survey Manual (Soil Survey Staff, 2012). Apart from the transect study, profiles were also studied at random, almost like in a grid pattern, outside the transect areas to validate the soil map unit boundaries.

Based on the soil characteristics, the soils were grouped into different soil series. Soil series is the most homogeneous unit having similar horizons and properties and behaves similarly for a given level of management. Soil depth, texture, colour, kind of horizon and horizon sequence, amount and nature of gravel present, nature of substratum etc, were used as the major differentiating characteristics for identifying soil series occurring in the area. The differentiating characteristics used for identifying the soil series are given in Table 3.1. Based on the above characteristics, 4 soil series were identified in the Bachanahalli-1 microwatershed.

Table 3.1 Differentiating Characteristics used for identifying Soil Series
(Characteristics are of Series Control Section)

Soils of Granite gneiss Landscape						
Sl. no	Soil Series	Depth (cm)	Colour (moist)	Texture	Gravel (%)	Horizon sequence
1	Annurkeri (ARK)	>150	2.5YR2.5/2,3/2, 2.5/3,3/3,2.5/4,3/4	sc-c	<15	Ap-Bt
2	Beemanabeedu (BMB)	>150	10YR2/1,2/2,3/1, 3/2,4/1	sc-c	-	Ap-Bw
3	Berambadi (BMD)	25-50	10YR3/3,4/2, 7.5YR3/2,3/3,	scl	<15	Ap-Bw-Cr
4	Gopalapura (GPR)	75-100	2.5YR3/2, 3/3 5YR3/3,4/3,	gscl-sc	15-35	AP-Bt-Cr
5	Hundipura (HDR)	25-50	2.5YR2.5/4, 5YR3/2	sl	<15	Ap-A2-Cr
6	Honnegaudanahalli (HGH)	>150	7.5YR2.5/2,2.5/3, 3/3,2.5/4,3/4	scl-c	<15	Ap-Bw
7	Hullipura (HPR)	50-75	7.5YR2.5YR2.5/2, 3/2	gsl-sc	15-35	AP-Bw-Cr
8	Kallipura (KLP)	100-150	2.5YR2.5/3,2.5/4, 3/4	scl-c	<15	AP-Bt-Cr

3.4 Soil Mapping

The area under each soil series was further separated into 20 soil phases and their boundaries delineated on the cadastral map based on the variations observed in the texture of the surface soil, slope, erosion, presence of gravel, stoniness etc. A soil phase is a subdivision of soil series based mostly on surface features that affect its use and management.

The soil mapping units are shown on the map (Fig. 3.5) in the form of symbols. During the survey of 24 soil profile pits, few minipits and a few auger bores representing different landforms occurring in the microwatershed were studied. All the profile locations are indicated on the village cadastral map in the form of a triangle. In addition to the profile study, spot observations in the form of minipits, road cuts, terrace cuts etc., were studied to validate the soil boundaries on the soil map. The soil map shows the geographic distribution of 20 mapping units representing 4 soil series occurring in the microwatershed. The soil map unit (soil legend) description is presented in Table 3.2. The soil phase map (management units) shows the distribution of 20 phases mapped in the microwatershed. Each mapping unit (soil phase) delineated on the map has similar soil and site characteristics. In other words, all the farms or survey numbers included in one phase will have similar management needs and they have to be treated accordingly.

3.5 Laboratory Characterization

Soil samples were collected from representative master profiles for laboratory characterization by following the methods outlined in the Laboratory Manual (Sarma *et al*, 1987). Surface soil samples collected from farmer's fields (60 samples) for fertility

status (major and micronutrients) at 320 m grid interval were analyzed in the laboratory (Katyal and Rattan, 2003). By linking the soil fertility data to the survey numbers through GIS, soil fertility maps were generated using kriging method for the microwatershed.

Table 3.2 Soil map unit description of Bachanahalli -1 Microwatershed

Soil No	Soil Series	Soil Phase	Mapping Unit Description	Area in ha (%)
	ARK	Annurkeri soils are very deep (>150 cm), well drained, have dark reddish brown to very dusky red sandy clay to clay soils occurring on very gently sloping uplands under cultivation.		35.71 (6.30)
1		ARKhB1	Sandy clay loam surface, slope 1-3%, slight erosion	18.30(3.23)
2		ARKhB2g1	Sandy clay loam surface, slope 1-3%, moderate erosion, gravelly (15-35%)	17.41(3.07)
	BMB	Beemanabeedu soils are very deep (>150 cm), moderately well drained, have very dark greyish brown to dark grey and very dark brown clayey soils occurring on very gently sloping lowlands under cultivation		119.68 (21.12)
3		BMBhB2	Sandy clay loam surface, slope 1-3%, moderate erosion	37.91(6.69)
4		BMBiB1	Sandy clay surface, slope 1-3%, slight erosion	9.77(1.72)
5		BMBmB1	Clay surface, slope 1-3%, slight erosion	72.0 (12.71)
	BMD	Berambadi soils are shallow (25-50 cm), well drained, dark brown to dark greyish brown clayey soils occurring on very gently to gently sloping uplands under cultivation		51.41 (9.08)
6		BMDhB1	Sandy clay loam surface, slope 1-3%, slight erosion	8.19 (1.45)
7		BMDhB1g1	Sandy clay loam surface, slope 1-3%, slight erosion, gravelly (15-35%)	21.63 (3.82)
8		BMDhC2g1	Sandy clay loam surface, slope 3-5%, moderate erosion, gravelly (15-35%)	21.59 (3.81)
	GPR	Gopalapura soils are moderately deep (75-100 cm), well drained, have dark brown to dark reddish brown and reddish brown gravelly sandy clay loam to sandy clay soils occurring on very gently to gently sloping uplands under cultivation		122.10 (21.55)
9		GPRcB1	Sandy loam surface, slope 1-3%, slight erosion	20.07 (3.54)
10		GPRcC2g1	Sandy loam surface, slope 3-5%, moderate erosion, gravelly (15-35%)	70.02 (12.36)
11		GPRhB1g1	Sandy clay loam surface, slope 1-3%, slight erosion, gravelly (15-35%)	22.64 (4.00)
12		GPRiB1	Sandy clay surface, slope 1-3%, slight erosion	9.37 (1.65)
	HDR	Hindupur soils are shallow (25-50 cm), well drained, have dark reddish brown to dusky red sandy clay loam to sandy clay soils occurring on very gently sloping uplands under		4.78 (0.84)

		cultivation	
13		HDRhB1g1 Sandy clay loam surface, slope 1-3%, slight erosion, gravelly (15-35%)	4.78 (0.84)
	HGH	Honnegaudanahalli soils are very deep (>150 cm), well drained, have very dark brown to brown and dark reddish brown sandy clay loam soils occurring on very gently sloping uplands under cultivation	30.57 (5.39)
14		HGHcB2 Sandy loam surface, slope 1-3%, moderate erosion	6.68 (1.18)
15		HGHhB1 Sandy clay loam surface, 1-3% slope, slight erosion	10.16 (1.79)
16		HGHhB1g1 Sandy clay loam surface, slope 1-3%, slight erosion, gravelly (15-35%)	10.37 (1.83)
17		HGHhB2 Sandy clay loam surface, slope 1-3%, moderate erosion	3.36 (0.59)
	HPR	Hullipura soils are moderately shallow (50-75 cm), well drained, have dark brown to very dark brown gravelly sandy clay loam to sandy clay soils occurring on very gently sloping uplands under cultivation	9.53 (1.68)
18		HPRcB1 Sandy loam surface, slope 1-3%, slight erosion	3.70 (0.65)
19		HPRiB2g1 Sandy Clay surface, slope 1-3%, moderate erosion, gravelly (15-35%)	5.83 (1.03)
	KLP	Kallipura soils are deep (100-150 cm), well drained, have dark reddish brown to dark red gravelly sandy clay loam to sandy clay soils occurring on very gently sloping uplands under cultivation	1.15 (0.20)
20		KLPhB2 Sandy clay loam, slope 1-3%, moderate erosion	1.15 (0.20)
21		Forest	178.29 (31.47)
22		Others Habitation and waterbodies	13.33 (2.35)

3.6 Land Management Units

The 20 soil phases identified and mapped in the microwatershed were regrouped into 7 Land Management Units (LMU's) for the purpose of preparing a proposed crop plan for sustained development of the microwatershed. The database (soil phases) generated under LRI was utilized for identifying Land Management Units (LMUs) based on the management needs. One or more than one soil site characteristic having influence on the management have been chosen for identification and delineation of LMUs. For Bachanahalli-1 microwatershed, five soil and site characteristics, namely soil depth, soil texture, slope, erosion and gravel content have been considered for defining LMUs. The land management units are expected to behave similarly for a given level of management.

THE SOILS

Detailed information pertaining to the nature, extent and distribution of different kinds of soils occurring in Bachanahalli-1 microwatershed is provided in this chapter. The microwatershed area has been identified as granite gneiss landscape based on geology. In all, 8 soil series were identified in different landforms. Soil formation is the result of the combined effect of environmental and terrain factors that are reflected in soil morphology. In the Granite gneiss landform, it is by parent material, relief and climate.

A brief description of each of the 8 soil series identified followed by 20 soil phases (management units) mapped (Fig. 3.5) are furnished below. The physical and chemical characteristics of soil series identified in Bachanahalli-1 microwatershed along with soil classification are given in Table 4.1. The soils in any one map unit differ from place to place in their depth, texture, slope, gravelliness, erosion or any other site characteristics that affect management. The soil phase map can be used for identifying the suitability of areas for growing specific crops or for other alternative uses and also for deciding the type of conservation structures needed. The detailed information on soil and site-characteristics like soil depth, surface soil texture, slope, erosion, gravelliness, AWC, LCC etc, with respect to each of the soil phase identified is given village/survey number wise for the microwatershed in Appendix-I.

4.1 Soils of Granite gneiss Landscape

In this landscape, 8 soil series are identified and mapped. Brief description of each series and their phases identified are given below. Of these, Gopalapura (GPR) soil series occupies maximum area of about 122.10 ha (22%) followed by Beemanabeedu (BMB) soil series 119.68 ha (21%), Berambadi (BMD) soil series 51.41 ha (9%), Annurkeri (ARK) soil series 35.71 ha (6%), Honnegaudanahalli (HGH) soil series 30.57 ha (5%), Hullipura (HPR) soil series 9.53 ha (2%), Hindupur (HDR) soil series 4.78 ha (1%) and Kallipura (KLP) soil series occupies minor area of about 1.15 ha (1%).

4.1.1 Annurkeri (ARK) Series: Annurkeri soils are very deep (>150 cm), well drained, have dark reddish brown to very dusky red sandy clay to sandy clay loam soils. They are developed from weathered granite gneiss and occur on very gently sloping uplands. The Annurkeri series has been classified as a member of the fine-loamy, mixed, isohyperthermic family of Rhodic Paleustalfs.

The thickness of the solum ranges from 150 to 200 cm. The thickness of A horizon ranges from 11 to 18 cm. Its colour is in 5YR and 2.5 YR hue with value 3 and chroma 3 to 4. The texture varies from sandy clay to sandy clay loam with 10 to 15 per cent gravel. The thickness of B horizon is more than 150 cm. Its colour is in 2.5 YR hue with value 2.5 to 3 and chroma 2 to 4. Texture is dominantly sandy clay to clay with less than 15 per cent gravel. The available water capacity is very high (>200 mm/m). Two phases were identified and mapped.



Landscape and soil profile characteristics of Annurkeri (ARK) Series

4.1.2 Beemanabeedu (BMB) Series: Beemanabeedu soils are very deep (>150 cm), moderately well drained, have very dark greyish brown to dark grey and very dark brown sandy clay to clay soils. They are developed from weathered granite gneiss and occur on very gently sloping lowlands. The Beemanabeedu series has been classified as a member of the fine, mixed, isohyperthermic family of Typic Haplustepts.

The thickness of the solum ranges from 150 to 200 cm. The thickness of A horizon ranges from 12 to 17 cm. Its colour is in 10 YR and 7.5 YR hue with value 2.5 to 4 and chroma 2 to 4. The texture varies from sandy clay to clay with less than 15 per cent gravel. The thickness of B horizon is more than 150 cm. Its colour is in 10 YR hue with value 2 to 4 and chroma 1 to 2. Texture is sandy clay to clay with less than 15 per cent gravel. The available water capacity is very high (>200 mm/m). Two phases were identified and mapped.



Landscape and soil profile characteristics of Beemanabeedu (BMB) Series

4.1.3 Berambadi (BMD) Series: Berambadi soils are shallow (25-50 cm), well drained, very dark gray to dark brown sandy clay loam soils. They have developed from granite gneiss and occur on very gently to moderately sloping uplands under cultivation.

The thickness of the solum ranges from 30 to 50 cm. The thickness of A-horizon ranges from 10 to 17 cm. Its colour is in 10 YR hue with value 3 to 4 and chroma 3. The texture varies from sandy loam to sandy clay with 15 to 25 per cent gravel. The thickness of B horizon ranges from 30 to 40 cm. Its colour is in 10YR and 7.5YR hue with value 3 to 4 and chroma 2 to 4. Its texture is sandy clay loam with gravel content of 10 to 15 per cent. The available water capacity is low (50-100 mm/m).

4.1.4 Gopalapura (GPR) Series: Gopalapura soils are moderately deep (75-100 cm), well drained, have dark brown to dark reddish brown and reddish brown sandy clay loam to sandy clay soils. They have developed from granite gneiss and occur on very gently to gently sloping uplands.

The thickness of the solum ranges from 73 to 97 cm. The thickness of A horizon ranges from 12 to 18 cm. Its colour is in 2.5 YR and 7.5 YR hue with value 3 and chroma 2. The texture varies from gravelly sandy clay to sandy clay loam with 10-25 per cent gravel. The thickness of B horizon ranges from 66 to 79 cm. Its colour is in 2.5 YR and 5 YR hue with value 3 to 4 and chroma 2 to 3. Its texture is gravelly sandy clay loam to gravelly sandy clay with gravel content of 15-35 per cent. The available water capacity is low (50-100 mm/m). Two phases were identified and mapped.



Landscape and soil profile characteristics of Gopalapura (GPR) Series

4.1.5 Hundipura (HDR) Series: Hundipura soils are shallow (25-50 cm), well drained, have dark reddish brown to dusky red sandy loam soils. They have developed from granite gneiss and occur on very gently to moderately sloping uplands. The Hundipura series has been classified as a member of the sandy, mixed, isohyperthermic family of (paralithic) Ustorthents.

The thickness of the soil ranges from 35 to 46 cm. The thickness of A horizon ranges from 7 to 18 cm. Its colour is in 7.5 YR and 5 YR hue with value 3 and chroma 3 to 4. The texture varies from loamy sand to sandy clay loam with 10 to 20 per cent gravel. The thickness of B horizon ranges from 19 to 31 cm. Its colour is in 2.5 YR and 5 YR hue with value 2.5 to 3 and chroma 2 to 4. Its texture is sandy loam with gravel content of < 15 per cent. The available water capacity is very low (<50 mm/m). Three phases were identified and mapped.



Landscape and soil profile characteristics of Hundipura (HDR) Series

4.1.6 Honnegaudanahalli (HGH) Series: Honnegaudanahalli soils are very deep (>150 cm), well drained, have very dark brown to brown and dark reddish brown sandy clay loam to clay soils. They are developed from weathered granite gneiss and occur on very gently sloping upland under cultivation. The Honnegaudanahalli series has been classified as a member of the fine loamy over clayey, mixed, isohyperthermic family of Typic Haplustepts.

The thickness of the solum ranges from 150-200 cm. The thickness of A horizon ranges from 14 to 19 cm. Its colour is in 7.5 YR hue with value 2.5 to 4 and chroma 2 to 6. The texture varies from sandy loam to clay with 10 to 15 per cent gravel. The thickness of B horizon is more than 150 cm. Its colour is in 7.5 YR hue with value 2.5 to 3 and chroma 2 to 4. Texture is sandy clay loam to clay with 35 to 50 per cent gravel. The available water capacity is very high (>200 mm/m). Three phases were identified and mapped.



Landscape and soil profile characteristics of Honnegaudanahalli (HGH) Series

4.1.7 Hullipura (HPR) Series: Hullipura soils are moderately shallow (50-75 cm), well drained, have dark brown to very dark brown sandy clay loam soils. They have developed from granite gneiss and occur on very gently to gently sloping uplands. The Hullipura series has been classified as a member of the fine-loamy, mixed, isohyperthermic family of Typic Haplustepts.

The thickness of the solum ranges from 51 to 71 cm. The thickness of A horizon ranges from 13 to 18 cm. Its colour is in 7.5YR and 10 YR hue with value 2.5 to 3 and chroma 2 to 4. The texture varies from gravelly sandy loam to gravelly clay with 15 to 25 per cent gravel. The thickness of B horizon ranges from 38 to 52 cm. Its colour is in 2.5 YR and 7.5 YR hue with value 2.5 to 3 and chroma 2. Its texture is gravelly sandy clay loam with gravel content of 15 to 35 per cent. The available water capacity is low (50-100 mm/m). Five phases were identified and mapped.



Landscape and soil profile characteristics of Hullipura (HPR) Series

4.1.8 Kallipura (KLP) Series: Kallipura soils are moderately shallow (50 to 75 cm), well drained, have brown to very dark brown and dark reddish brown sandy loam to clay loam soils. They have developed from granite gneiss and occur on nearly level to gently sloping uplands.

The thickness of the solum ranges from 54 to 75 cm. The thickness of A horizon ranges from 11 to 19 cm. Its colour is in 7.5 YR, 5YR and 2.5 YR hue with value 2.5 to 4 and chroma 2 to 6. The texture varies from gravelly sandy loam to gravelly clay loam with 10 to 20 per cent gravel. The thickness of B horizon ranges from 43 to 60 cm. Its colour is in 2.5 YR hue with value 3 and chroma 4 to 6. Its texture is gravelly sandy clay loam to gravelly sandy clay with gravel content of 15 to 35 per cent. The available water capacity is medium (100 mm/m).



Landscape and soil profile characteristics of Kallipura (KLP) Series

Table 4.1 The physical and chemical characteristics of soil series identified in Bachanahalli-1 microwatershed

Soil Series: Annurkeri (ARK), **Pedon:** RM-83

Location: 11°43'17.6"N, 76°38'74" (4B3E2G2c) Siddhapura village, Gundlupet Taluk and Chamarajanagara District

Analysis at: NBSS&LUP, Regional Centre, Bangalore. **Classification:** Fine-loamy, mixed, isohyperthermic Rhodic Paleustalfs

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-25	Ap	79.05	5.82	15.13	3.64	15.38	21.46	25.61	12.96	<5	sl	-	-
25-46	A2	78.83	7.26	13.91	2.42	6.35	25.60	29.84	14.62	<5	sl	-	-
46-77	Bt1	55.15	5.36	39.49	2.60	16.34	14.36	13.94	7.91	<5	sc	-	-
77-102	Bt2	52.89	8.50	38.61	3.00	10.54	18.90	12.71	7.75	<5	sc	-	-
102-126	Bt3	48.90	7.24	43.86	1.15	3.97	18.18	15.26	10.34	<5	sc	-	-
126-156	Bt4	47.75	10.71	41.54	3.25	10.16	12.36	13.93	8.06	<5	sc	-	-
156-180	Bt5	45.27	11.65	43.08	3.82	9.68	12.07	11.97	7.70	<5	sc	-	-

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO ₃	Exchangeable bases					CEC	CEC/ Clay	Base saturation	ESP
							Ca	Mg	K	Na	Total				
	Water	CaCl ₂	M KCl	dS m ⁻¹	%	%	cmol kg ⁻¹						%	%	
0-25	7.17	-	-	0.16	0.51	0.00	-	-	0.74	0.13	-	4.79	0.32	100	2.71
25-46	7.09	-	-	0.11	0.47	0.00	-	-	0.37	0.06	-	5.81	0.42	100	1.03
46-77	7.33	-	-	0.06	0.47	0.00	-	-	0.27	0.07	-	15.96	0.40	100	0.44
77-102	7.21	-	-	0.07	0.55	0.00	-	-	0.21	0.09	-	15.62	0.40	100	0.58
102-126	7.22	-	-	0.07	0.47	0.00	-	-	0.23	0.22	-	15.39	0.35	100	1.43
126-156	7.27	-	-	0.07	0.43	0.00	-	-	0.56	0.51	-	14.93	0.36	100	3.42
156-180	7.33	-	-	0.09	0.39	0.00	-	-	0.42	0.51	-	14.82	0.34	100	3.44

Contd...

Soil Series: Bheemanabeedu (BMB), **Pedon:** RM-34

Location: 11°43'45.9"N, 76°37'18.6"E, (4B3E2F1c) Honnegaudanahalli village, Gundlupet taluk, Chamarajanagara district

Analysis at: NBSS&LUP, Regional Centre, Bangalore. **Classification:** Fine, mixed, isohyperthermic Typic Haplustepts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-13	Ap	63.25	17.32	19.42	3.89	9.31	13.61	19.14	17.30	5	sc	-	-
13-32	Bw1	61.95	18.00	20.05	3.18	12.21	13.54	18.56	14.46	-	sc	-	-
32-53	Bw2	54.42	20.16	25.42	3.89	9.79	12.63	14.95	13.16	-	sc	-	-
53-76	Bw3	40.54	20.40	39.05	2.40	8.25	9.82	11.91	8.15	-	c	-	-
76-114	Bw4	35.54	17.90	46.56	2.00	7.36	8.94	9.88	7.36	-	c	-	-
114-146	Bw5	28.31	19.25	52.44	2.55	5.73	6.68	6.89	6.47	-	c	-	-
146-180	Bw6	19.43	27.91	52.66	1.61	2.35	3.84	6.06	5.57	-	c	-	-

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO ₃	Exchangeable bases					CEC	CEC/ Clay	Base saturation	ESP
	Water	CaCl ₂	M KCl				Ca	Mg	K	Na	Total				
				dS m ⁻¹	%	%	cmol kg ⁻¹							%	%
0-13	7.59	-	-	0.48	0.78	0.59	-	-	1.04	0.17	-	10.83	0.56	100	1.57
13-32	7.94	-	-	0.18	0.66	0.35	-	-	0.62	0.26	-	11.74	0.59	100	2.21
32-53	8.11	-	-	0.13	0.58	0.76	-	-	0.41	0.59	-	14.71	0.58	100	4.01
53-76	8.34	-	-	0.12	0.54	0.71	-	-	0.49	0.92	-	25.31	0.65	100	3.63
76-114	8.21	-	-	0.20	0.50	1.59	-	-	0.64	0.82	-	28.39	0.61	100	2.89
114-146	8.37	-	-	0.27	0.46	2.47	-	-	0.51	0.65	-	30.55	0.58	100	2.13
146-180	8.46	-	-	0.27	0.35	5.28	-	-	0.57	0.94	-	35.34	0.67	100	2.66

Contd...

Soil Series: Berambadi (BMD), **Pedon:** RM-125

Location: 11°43'17"N, 76°41' 41.4"E, (4B3E2J2a) Hundipur village, Gundlupet Taluk, Chamarajanagara District

Analysis at: NBSS&LUP, Regional Centre, Bangalore. **Classification:** Fine-loamy, mixed, isohyperthermic, Typic Haplustepts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-12	Ap	64.81	10.27	24.92	5.78	13.11	18.99	17.54	9.39	15	scl	-	-
12-30	A1	63.68	11.19	25.13	5.52	12.59	16.86	19.98	8.74	<10	scl	-	-
30-43	Bw	55.49	10.87	33.65	17.55	10.76	11.29	10.14	5.75	<10	scl	-	-

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO ₃	Exchangeable bases					CEC	CEC/ Clay	Base saturation	ESP
	Water	CaCl ₂	M KCl				Ca	Mg	K	Na	Total				
				dS m ⁻¹	%	%	cmol kg ⁻¹							%	%
0-12	8.28	-	-	0.23	0.74	1.02	-	-	0.73	0.15	-	18.13	0.73	100	0.83
12-30	8.61	-	-	0.18	0.70	1.14	-	-	0.98	0.54	-	19.61	0.78	100	2.75
30-43	8.68	-	-	0.20	0.51	2.28	-	-	0.48	0.58	-	19.15	0.57	1003	3.03

Contd...

Soil Series: Hundipura (HDR), **Pedon:** TR-15/P1

Location: 11°42'25.8"N, 76°41'26"E, (4B3E2J2a) Hundipura village, Gundlupet Taluk and Chamarajanagara District

Analysis at: NBSS&LUP, Regional Centre, Bangalore **Classification:** Sandy, mixed, isohyperthermic, (paralithic) Ustorthents

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-17	Ap	76.89	9.00	14.11	11.25	13.60	19.12	22.39	10.53	15	sl	-	-
17-37	A2	75.31	11.73	12.96	9.02	11.27	18.24	23.36	13.42	10	sl	-	-

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO ₃	Exchangeable bases					CEC	CEC/ Clay	Base saturation	ESP
	Water	CaCl ₂	M KCl				Ca	Mg	K	Na	Total				
				dS m ⁻¹	%	%	cmol kg ⁻¹							%	%
0-17	8.28	-	-	0.18	0.50	1.06	-	-	0.63	0.08	-	12.54	0.89	100	0.64
17-37	8.53	-	-	0.10	0.43	0.65	-	-	0.38	0.04	-	10.03	0.77	100	0.40

Contd...

Soil Series: Honnegaudanahalli (HGH), **Pedon:** RM-33

Location: 11°45'42.5"N, 76°36'46.2"E, (4B3E2F1c) Honnegaudanahalli village, Gundlupet Taluk and Chamarajanagara District

Analysis at: NBSS&LUP, Regional Centre, Bangalore.

Classification: Fine-loamy over clayey, mixed, isohyperthermic, Typic Haplustepts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-18	Ap	66.23	11.55	22.23	2.95	10.89	18.41	21.36	12.61	10	scl	-	-
18-30	Bw1	62.83	16.68	20.49	3.77	10.90	15.99	21.59	10.59	-	scl	-	-
30-52	Bw2	59.33	16.64	24.03	2.14	10.91	13.56	18.76	13.97	-	scl	-	-
52-78	Bw3	65.74	11.85	22.41	3.69	13.13	16.00	19.49	13.44	-	scl	-	-
78-100	Bw4	68.91	10.58	20.51	3.16	6.73	25.08	21.61	12.33	-	scl	-	-
100-148	Bw5	30.06	14.12	55.82	1.57	9.81	6.78	6.58	5.32	-	c	-	-
148-160	Bw6	22.82	29.70	47.48	1.49	4.13	6.54	5.39	5.28	-	c	-	-
160-190	Bw7	24.83	28.46	46.71	0.80	2.86	8.47	6.18	6.52	-	c	-	-

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO ₃	Exchangeable bases					CEC	CEC/ Clay	Base saturation	ESP
	Ca	Mg	K				Na	Total							
	Water	CaCl ₂	M KCl	dS m ⁻¹	%	%	cmol kg ⁻¹						%	%	
0-18	7.37	-	-	0.18	0.5	0.00	-	-	0.74	0.13	-	8.66	0.39	100	1.50
18-30	7.55	-	-	0.11	0.46	0.00	-	-	0.37	0.06	-	8.66	0.42	100	0.69
30-52	7.78	-	-	0.08	0.39	0.00	-	-	0.27	0.07	-	10.03	0.42	100	0.70
52-78	7.98	-	-	0.08	0.39	0.00	-	-	0.21	0.09	-	8.55	0.38	100	1.05
78-100	8.01	-	-	0.05	0.31	0.00	-	-	0.23	0.22	-	8.32	0.41	100	2.64
100-148	7.86	-	-	0.09	0.43	7.20	-	-	0.56	0.51	-	25.65	0.46	100	1.99
148-160	8.47	-	-	0.18	0.35	5.88	-	-	0.42	0.51	-	23.37	0.49	100	2.18
160-190	8.55	-	-	0.17	0.27	6.48	-	-	0.45	0.56	-	23.57	0.50	100	2.38

Contd...

Soil Series: Hullipura (HPR), **Pedon:** RM-126

Location: 11°43'21.8"N, 76°41'28.1"E, (4B3E2J2a) Hundipur village, Gundlupet Taluk, Chamarajanagara District

Analysis at: NBSS&LUP, Regional Centre, Bangalore **Classification:** Fine-loamy, mixed, isohyperthermic, Typic Haplustepts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-15	Ap	47.38	17.57	35.05	7.38	9.95	8.98	10.80	10.27	15	sc	-	-
15-28	Bw1	49.95	18.96	31.10	7.21	10.29	9.97	12.51	9.97	15	scl	-	-
28-55	Bc	70.38	11.41	18.21	6.23	13.18	14.20	21.45	15.32	20	sl	-	-

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO ₃	Exchangeable bases					CEC	CEC/ Clay	Base saturation	ESP
	Water	CaCl ₂	M KCl				Ca	Mg	K	Na	Total				
				dS m ⁻¹	%	%	cmol kg ⁻¹							%	%
0-15	8.36	-	-	0.36	0.60	2.34	-	-	0.93	0.43	-	24.17	0.69	100	1.78
15-28	8.36	-	-	0.37	0.72	3.30	-	-	1.10	0.61	-	6.16	0.20	100	9.90
28-55	8.73	-	-	0.23	0.24	0.84	-	-	0.41	0.85	-	16.30	0.90	100	5.21

Contd...

Soil Series: Kallipura (KLP), **Pedon:** RM-52

Location: 11°43'45.9"N, 76°37'18.6"E, (4B3E2F2c) Kallipura village, Gundlupet Taluk and Chamarajanagara District

Analysis at: NBSS&LUP, Regional Centre, Bangalore. **Classification:** Fine, mixed, isohyperthermic Typic Rhodustalfs

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-28	Ap	65.05	10.57	24.38	5.17	13.86	17.79	18.20	10.03	10	scl	-	-
28-48	Bt1	61.05	10.50	28.45	2.53	9.37	19.47	20.74	8.95	-	scl	-	-
48-75	Bt2	58.11	10.79	31.10	2.01	12.51	18.45	18.88	6.26	-	scl	-	-
75-97	Bt3	47.69	6.51	45.80	4.74	11.73	12.92	13.02	5.27	-	sc	-	-
97-120	Bt4	37.80	10.25	51.95	4.57	9.45	9.14	9.66	4.98	-	c	-	-
120-136	Bc	53.11	5.08	41.81	20.55	6.95	11.49	9.48	4.64	40	sc	-	-

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO ₃	Exchangeable bases					CEC	CEC/ Clay	Base saturation	ESP
	Water	CaCl ₂	M KCl				Ca	Mg	K	Na	Total				
				dS m ⁻¹	%	%	cmol kg ⁻¹							%	%
0-28	6.07	-	-	0.14	0.70	0.00	-	-	0.59	0.26	-	9.23	0.38	100	2.82
28-48	6.45	-	-	0.04	0.58	0.00	-	-	0.25	0.19	-	9.92	0.35	100	1.92
48-75	6.47	-	-	0.04	0.54	0.00	-	-	0.24	0.13	-	9.80	0.32	100	1.33
75-97	6.50	-	-	0.06	0.58	0.00	-	-	0.26	0.17	-	13.22	0.29	100	1.29
97-120	6.51	-	-	0.05	0.50	0.00	-	-	0.36	0.32	-	15.50	0.30	100	2.06
120-136	6.88	-	-	0.04	0.46	0.00	-	-	0.24	0.17	-	14.82	0.35	100	1.15

INTERPRETATION FOR LAND RESOURCE MANAGEMENT

The most important soil and site characteristics that affect land use and conservation needs of an area are land capability, land irrigability, soil depth, texture, coarse fragments, available water capacity, soil slope, soil erosion, soil reaction etc., are interpreted from the data base generated through the land resource inventory and several thematic maps are generated. These would help in identifying the areas suitable for growing crops and conservation structures needed thus helping to maintain good soil health for sustained crop production. The various interpretative and theme maps generated are described below.

5.1 Land Capability Classification

Land capability classification is an interpretative grouping of soil map units (soil phases) mainly based on inherent soil characteristics, external land features and environmental factors that limit the use of land for agriculture, pasture, forestry, or other uses on a sustained basis (IARI, 1971). The land and soil characteristics used to group the land resources in an area into various land capability classes, subclasses and units are

Soil characteristics: Soil depth, texture, gravelliness, soil reaction, available water capacity, calcareousness, salinity/alkali etc.

Land characteristics: Slope, erosion, drainage, rock outcrops.

Climate: Total rainfall and its distribution, and length of crop growing period.

The Land capability classification system is divided into land capability classes, subclasses and units based on the level of information available. Eight land capability classes are recognized. They are

Class I: The soil map units have few or very few limitations that restrict their use.

Class II: The soil map units have moderate limitations that reduce the choice of crops or that require moderate conservations practices.

Class III: The soil map units have severe limitations that reduce the choice of crops or that require special conservation practices.

Class IV: The soil map units have severe limitations that reduce the choice of crops or that require very careful management.

Class V: Soils in the mapping units are not likely to erode, but have other limitations that are impractical to remove and as such not suitable for agriculture.

Class VI: The lands have severe limitations that make them generally unsuitable for cultivation.

Class VII: The lands have very severe limitations that make them unsuitable for cultivation.

Class VIII: Soil and other miscellaneous areas that have very severe limitations that nearly preclude their use for any crop production.

The land capability subclasses are recognised based on the dominant limitations observed within the given capability class. The subclasses are designated by adding a lower case letter like 'e', 'w', 's', or 'c' to the class numeral. The subclass "e" indicates that the main hazard is risk of erosion, "w" indicates drainage or wetness as a limitation for plant growth, "s" indicates shallow soil depth, coarse or heavy textures, calcareousness, salinity/alkaline or gravelliness and "c" indicates limitation due to climate.

The land capability subclasses have been further subdivided into land capability units based on the kinds of limitations present in each subclass. Ten land capability units are used in grouping the soil map units. They are stony or rocky (0), erosion hazard (slope, erosion) (1), coarse texture (sand, loamy sand, sandy loam (2), fine texture (cracking clay, silty clay), (3) slowly permeable subsoil (4), coarse underlying material (5), salinity/alkali (6), stagnation, overflow, high ground water table (7), soil depth (8) and fertility problems (9). The capability units have similar soil and land characteristics that respond similarly to a given level of management. The soils of the microwatershed have been classified upto land capability subclass level.

The 20 soil map units identified in the Bachanahalli-1 microwatershed are grouped under 2 land capability classes and 6 land capability subclasses (Fig. 5.1).

Of the lands suitable for agriculture, about 37 percent are good cultivable lands (Class II) with minor limitations of soil characteristics and erosion and distributed in the northern, northeastern, northwestern, central, western, southwestern, southern and southeastern parts of the microwatershed. About 29 per cent are moderately good cultivable lands (Class III) with moderate limitations of soil characteristics, erosion and excess of water limitations are distributed in northern, western, central, eastern, southern and northeastern parts of the microwatershed.




Fig. 5.1 Land Capability map of Bachanahalli-1 Microwatershed

5.2 Soil Depth

Soil depth refers to the depth of the soil occurring above the parent material or hard rock. The depth of the soil determines the effective rooting depth for plants and in accordance with soil texture, mineralogy and gravel content, the capacity of the soil column to hold water and nutrient availability. Soil depth is one of the most important soil characteristic that is used in differentiating soils into different soil series. The soil depth classes used in identifying soils in the field are very shallow (<25 cm), shallow (25-50 cm), moderately shallow (50-75 cm), moderately deep (75-100 cm), deep (100-150 cm) and very deep (>150 cm). They were used to classify the soils into different depth classes and a soil depth map was generated. The area extent and their geographical distribution in the microwatershed is shown in Figure 5.2.

Shallow (25-50 cm) soils occupy about 56 ha (10%) area and are distributed in the northeastern, central and southeastern parts of the microwatershed. Moderately shallow (50-75 cm) occupy about 10 ha (2%) in the northern and northwestern parts of the microwatershed. About 122 ha (22%) area has soils that are moderately deep (75-100 cm) soils and occur in the northeastern, eastern, southern and southeastern parts of the microwatershed. Deep (100-150 cm) soils occupy an area about 1 ha (<1%) and are distributed in northwestern parts of the microwatershed. An area of about 186 ha (33%) occupied by the very deep (>150 cm) soils and are distributed in northern, northwest, western, central, eastern and southeastern parts of the microwatershed.

The most problematic lands 56 ha (10%) having shallow (25-50 cm) rooting depth. They are suitable for growing short duration agricultural crops but well suited for pasture, forestry or other recreational purposes. The most productive lands cover about 186 ha (33%) where all climatically adapted long duration crops be grown.




Fig. 5.1 Soil Depth map of Bachanahalli-1Microwatershed

5.3 Surface Soil Texture

Texture is an expression to indicate the coarseness or fineness of the soil as determined by the relative proportion of primary particles of sand, silt and clay. It has a direct bearing on the structure, porosity, adhesion and consistence. The surface layer of a soil to a depth of about 25 cm is the layer that is most used by crops and plants. The surface soil textural class provides a guide to understanding soil-water retention and availability, nutrient holding capacity, infiltration, workability, drainage, physical and chemical behaviour, microbial activity and crop suitability. The textural classes used for LRI were used to classify the soils and a surface soil texture map was generated. The area extent and their geographical distribution in the microwatershed is shown in Figure 5.3.

An area of about 277 ha (49%) has soils that are loamy at the surface and are distributed in the major part of the microwatershed. An area of 97 ha (17%) has soils that are clayey at the surface and are distributed in the northern, western and eastern parts of the microwatershed (Fig. 5.3).

The productive lands 97 ha (17%) with respect to surface soil texture are the clayey soils that have high potential for soil-water retention and availability, and nutrient retention and availability, but have problems of drainage, infiltration, workability and other physical

problems. The other most productive lands (49%) are loamy soils which also have high potential for AWC, nutrient availability but have no drainage or other physical problems.




Fig. 5.3 Surface Soil Texture map of Bachanahalli-1 Microwatershed

5.4 Soil Gravelliness

Gravel is the term used for describing coarse fragments between 2 mm and 7.5 cm diameter and stones for those between 7.5 cm and 25 cm. The presence of gravel and stones in soil reduces the volume of soil responsible for moisture and nutrient storage, drainage, infiltration and runoff and hinders plant growth by impeding root growth and seedling emergence, intercultural operations and farm mechanization. The gravelliness classes used in LRI were used to classify the soils and using these classes a gravelliness map was generated. The area extent and their geographic distribution in the microwatershed is shown in Figure 5.4.

The soils that are non gravelly (<15%) cover a maximum area of about 201 ha (35%) and are distributed in the northern, western, central and eastern parts of the microwatershed (Fig. 5.4). An area in the microwatershed has soils that are gravelly (15-35%) covering about 174 ha (31%) and are distributed in the northwestern, northeastern, central and all southern parts of the microwatershed.

The most productive lands with respect to gravelliness are found to be 35 per cent. They are non-gravelly with less than 15 per cent gravel and have potential for growing both annual and perennial crops. The problem soils (31%) are those that are gravelly (15-35%) where only short duration crops can be grown.




Fig. 5.4 Soil Gravelliness map of Bachanahalli-1Microwatershed

5.5 Available Water Capacity

The soil available water capacity (AWC) is estimated based on the ability of the soil column to retain water between the tensions of 0.33 and 15 bar in a depth of 100 cm or the entire solum if the soil is shallower. The AWC of the soils (soil series) as estimated by considering the soil texture, mineralogy, soil depth and gravel content (Sehgal *et al.*, 1990) and accordingly the soil map units were grouped into five AWC classes *viz*, very low (<50 mm/m), low (50-100 mm/m), medium (100-150 mm/m), high (150-200 mm/m) and very high (>200 mm/m) and using these values, an AWC map was generated. The area extent and their geographic distribution of different AWC classes in the microwatershed is given in Figure 5.5.

An area of about 76 ha (13%) has soils that are Very low (<50 mm/m) in available water capacity and are distributed in the northeastern and southeastern parts of the microwatershed. About 113 ha (20%) area is low in available water capacity (51-100 mm/m) and are distributed in the northern, northwestern, southwestern, southern, northeastern and southeastern parts of the microwatershed. A maximum area of about 186 ha (33%) has soils that are very high (>200 mm/m) in available water capacity and are distributed in north, northwest, west and southeast, northeastern parts of the microwatershed.

About 186 ha (33%) area in the microwatershed has soils that are most productive with regard to available water capacity. This is most suitable to long duration crops. An area of about 76 ha (13%) are problematic with respect to available water capacity. Here,

only short or medium duration crops can be grown and the probability of crop failure is very high. These areas are best put to other alternative uses.




Fig. 5.5 Soil Available Water Capacity map of Bachanahalli-1 Microwatershed

5.6 Soil Slope

Soil slope refers to the inclination of the surface of the land. It is defined by gradient, shape and length, and is an integral feature of any soil as a natural body. Slope is considered important in soil genesis, land use and land development. The length and gradient of slope influences the rate of runoff, infiltration, erosion and deposition. The soil map units were grouped into four slope classes and a slope map was generated showing the area extent and their geographic distribution in the microwatershed (Fig. 5.6).

The major area of the microwatershed falls under very gently sloping (1-3% slope) class. It covers an area of about 283 ha (50%) and is distributed in the major part of the microwatershed. An area of about 92 ha (16%) comes under gently sloping (3-5 %) class and are distributed in all southern parts of the microwatershed.

An area of about 283 ha (50%) in the microwatershed has soils that have high potential in respect of soil slopes. In these areas, all climatically adapted annual and perennial crops can be grown without much soil and water conservation and other land development measures.




Fig. 5.6 Soil Slope map of Bachanahalli-1 Microwatershed

5.7 Soil Erosion

Soil erosion refers to the wearing away of the earth's surface by the forces of water, wind and ice involving detachment and transport of soil by raindrop impact. It is used for accelerated soil erosion resulting from disturbance of the natural landscape by burning, excessive grazing and indiscriminate felling of forest trees and tillage, all usually by man. The erosion classes showing an estimate of the current erosion status as judged from the field observations in the form of rills, gullies or a carpet of gravel on the surface are recorded. Four erosion classes, viz, slight erosion (e1), moderate erosion (e2), severe erosion (e3) and very severe erosion (e4) are recognized. The soil map units were grouped into different erosion classes and soil erosion map generated. The area extent and their spatial distribution in the microwatershed is given in Figure 5.7.

Soils that are slight (e1 class) cover maximum area of about 211 ha (37%) in the microwatershed. They are distributed in northern, northeastern, eastern, central and southeastern parts of the microwatershed. An area of about 164 ha (29%) comes under moderate erosion (e2 class) and are distributed in the western, southern and southeastern parts of the microwatershed.

An area of about 164 ha (29%) in the microwatershed is problematic because of moderate erosion. These areas need soil and water conservation and other land development measures for restoring the soil health.




Fig. 5.7 Soil Erosion map of Bachanahalli-1 Microwatershed

FERTILITY STATUS

Soil fertility plays an important role in increasing crop yield. The adoption of high yielding varieties that require high amounts of nutrients has resulted in deficiency symptoms in crops and plants due to imbalanced fertilization and poor inherent fertility status. Hence, it is necessary to know the fertility (macro and micro nutrients) status of the soils of the watersheds for assessing the kind and amount of fertilizers required for each of the crop intended to be grown. For this purpose, the surface soil samples collected from the grid points (one soil sample at every 320 m grid interval) all over the microwatershed through land resource inventory in the year 2014 were analysed for pH, EC, organic carbon, available phosphorus and potassium, and for micronutrients like zinc, boron, copper, iron and manganese, and secondary nutrient sulphur.

Soil fertility data generated has been assessed and individual maps for all the nutrients for the microwatershed have been generated using kriging method under GIS. The village/survey number wise fertility data for the microwatershed is given in Appendix-II.

6.1 Soil Reaction (pH)

The soil analysis of the Bachanahalli-1 microwatershed for soil reaction (pH) showed that about 34 ha (6 %) area is moderately acid (pH 5.5-6.0) and is distributed in the sentral, eastern and southern parts of the microwatershed. About 163 ha (29%) area is slightly acid (pH 6.0-6.5) and is distributed in the major part of the microwatershed. An area of about 125 ha (22%) is neutral (pH 6.5-7.3) and distributed in northern, western, central and northeastern parts of the microwatershed. Slightly alkaline (pH 7.3-7.8) to moderately alkaline (pH 7.8-8.4) (covers an area about 53 ha (9%) and are distributed in northern parts of the microwatershed (Fig. 6.1).

6.2 Electrical Conductivity (EC)

The Electrical Conductivity of the non saline soils ($<2 \text{ dSm}^{-1}$) covers an area of about 365 ha (65 %) and are distributed in major parts of the microwatershed. An area of about 9 ha (2 %) is in low ($2-4 \text{ dSm}^{-1}$) and are distributed in eastern parts of the microwatershed (Fig 6.2).

6.3 Organic Carbon

The soil organic carbon content of the microwatershed area is low ($<0.5\%$) covering about 54 ha (10%) and are distributed in the central and eastern parts of the microwatershed (Fig. 6.3). An area of about 285 ha (50%) soils are medium in organic carbon ($0.5-0.75\%$) and are distributed in the major part of the microwatershed. High ($>0.75\%$) in organic carbon covers an area of about 36 ha (6%) and are distributed in the northwestern and southern parts of the microwatershed.




Fig.6.1 Soil Reaction (pH) map of Bachanahalli -1 Microwatershed




Fig.6.2 Electrical Conductivity (EC) map of Bachanahalli-1 Microwatershed




Fig. 6.3 Soil Organic Carbon map of Bachanahalli-1 Microwatershed

6.4 Available Phosphorus

The soil analysis revealed that the entire area of available phosphorus is Medium (23-57 kg/ha) in an area of 375 ha (66.18 %) of the microwatershed (Fig. 6.4).

6.5 Available Potassium

Available potassium content is medium (145-337 kg/ha) in 37 ha (7%) area and is distributed in the southwestern parts of the microwatershed (Fig. 6.5). High available potassium (>337 kg/ ha) content accounts for 338 ha (60%) and is distributed in major part of the microwatershed.

6.6 Available Sulphur

Available sulphur content is low (<10 ppm) in 375 ha (66%) area in the microwatershed and is distributed in the entire parts of the microwatershed (Fig. 6.6).

6.7 Available Boron

Available boron content is low (<0.5 ppm) in 375 ha (66 %) area and is distributed in the all parts of the microwatershed (Fig. 6.7).




Fig. 6.4 Soil available Phosphorus map of Bachanahalli-1 Microwatershed




Fig. 6.5 Soil available Potassium map of Bachanahalli-1 Microwatershed




Fig. 6.6 Soil available Sulphur map of Bachanahalli-1 Microwatershed




Fig. 6.7 Soil available Boron map of Bachanahalli-1 Microwatershed

6.8 Available Iron

Available iron content is sufficient (>4.5 ppm) in 375 ha (66%) area and is distributed in entire parts of the microwatershed (Fig. 6.8).

6.9 Available Manganese

Available manganese content is sufficient (>1.0 ppm) in the entire microwatershed area (Fig 6.9).

6.10 Available Copper

Available copper content is sufficient (>0.2 ppm) in the entire microwatershed area (Fig 6.10).

6.11 Available Zinc

Available zinc content is deficient (<0.6 ppm) in 347 ha (61%) area and is distributed in the major part of the microwatershed (Fig 6.11). An area of about 28 ha (5%) area is sufficient in available zinc (>0.6 ppm) and is distributed in northeastern and northwestern parts of the microwatershed.




Fig. 6.8 Soil available Iron map of Bachanahalli-1Microwatershed




Fig. 6.9 Soil available Manganese map of Bachanahalli-1Microwatershed




Fig. 6.10 Soil available Copper map of Bachanahalli-1Microwatershed




Fig. 6.11 Soil available Zinc map of Bachanahalli-1 Microwatershed

LAND SUITABILITY FOR MAJOR CROPS

The soil and land resource units (soil phases) of Bachanahalli-1 microwatershed were assessed for their suitability for growing food, fibre, fodder and horticulture crops by following the procedure as outlined in FAO, 1976 and 1983. Crop requirements were developed for each of the crop from the available research data, Naidu *et. al.* (2006) and Natarajan *et. al* (2015). The crop requirements were matched with the soil and land characteristics (Table 7.1) to arrive at the crop suitability. In FAO land suitability classification, two orders are recognized. Order S-Suitable and Order N-Not suitable. The orders have classes, subclasses and units. Order-S has three classes, Class S1-Highly Suitable, Class S2-Moderately Suitable and Class S3-Marginally Suitable. Order N has two classes, N1-Currently not Suitable and N2-Permanently not Suitable. There are no subclasses within the class S1 as they will have very minor or no limitations for crop growth. Classes S2 and S3 are divided into subclasses based on the kinds of limitations encountered. The limitations that affect crop production are ‘c’ for erratic rainfall and its distribution and length of growing period (LGP), ‘e’ for erosion hazard, ‘r’ for rooting condition, ‘t’ for lighter or heavy texture, ‘g’ for gravelliness or stoniness, ‘n’ for nutrient availability, ‘l’ for topography, ‘m’ for moisture availability ‘z’ for calcareousness, ‘s’ for sodium and ‘w’ for drainage. These limitations are indicated as lower case letters to the class symbol. For example, moderately suitable land with the limitations of soil depth and erosion is designated as S2re. For the microwatershed, the soil mapping units were evaluated and classified up to subclass level.

Using the above criteria, the soil map units of the microwatershed were evaluated and land suitability maps for 27 major agricultural and horticultural crops were generated. The detailed information on the kind of suitability of each of the soil phase for the crops assessed are given village/ survey number wise for the microwatershed in Appendix-III.

7.1 Land Suitability for Sorghum (*Sorghum bicolor*)

Sorghum is one of the major crops grown in Karnataka in an area of 11.02 lakh ha in northern Karnataka in Bijapur, Kalaburgi, Raichur, Bidar, Belgaum, Dharwad and Bellary districts. The crop requirements for growing sorghum (Table 7.2) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and land suitability map for growing sorghum was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed is given in Figure 7.1.

An area of about 247 ha (44%) is highly suitable (Class S1) for growing sorghum and are distributed in the major part of the microwatershed. About 41 ha (7%) in the microwatershed has soils that are moderately suitable (Class S2) for growing sorghum and are distributed mainly in the northern, northwestern, eastern parts of the microwatershed. They have minor limitations of rooting depth and texture.

Table 7.1 Soil-Site Characteristics of Bachanahalli-1 Microwatershed

Soil Map Units	Climate (P)(mm)	Growing period(Days)	Drainage class	Soil depth (cm)	Soil texture		Gravelliness		AWC (mm/m)	Slope (%)	Erosion	pH	EC	ESP	CEC [Cmol (p ⁺)kg ⁻¹]	BS (%)
					Surface	Sub-surface	Surface (%)	Sub surface (%)								
ARKhB1	754	120-150	WD	>150	scl	sc-c	-	<15	>200	1-3	slight	7.17	0.16	2.71	4.79	100
ARKhB2g1	754	120-150	WD	>150	scl	sc-c	15-35	<15	>200	1-3	Moderate	7.17	0.16	2.71	4.79	100
BMBhB2	754	120-150	MWD	>150	scl	sc-c	-	-	>200	1-3	Moderate	7.59	0.48	1.57	10.83	100
BMBiB1	754	120-150	MWD	>150	sc	sc-c	-	-	>200	1-3	slight	7.59	0.48	1.57	10.83	100
BMBmB1	754	120-150	MWD	>150	c	sc-c	-	-	>200	1-3	slight	7.59	0.48	1.57	10.83	100
BMDhB1	754	120-150	WD	25-50	scl	scl	-	<15	-	1-3	slight	8.28	0.23	0.83	18.13	100
BMDhB1g1	754	120-150	WD	25-50	scl	scl	15-35	<15	-	1-3	moderate	8.28	0.23	0.83	18.13	100
BMDhC2g1	754	120-150	WD	25-50	scl	scl	15-35	<15	-	3-5	slight	8.28	0.23	0.83	18.13	100
GPRcB1	754	120-150	WD	75-100	sl	gscl-sc	-	15-35	50-100	1-3	slight	-	-	-	-	-
GPRcC2g1	754	120-150	WD	75-100	sl	gscl-sc	15-35	15-35	50-100	3-5	moderate	-	-	-	-	-
GPRhB1g1	754	120-150	WD	75-100	scl	gscl-sc	15-35	15-35	50-100	1-3	slight	-	-	-	-	-
GPRiB1	754	120-150	WD	75-100	sc	gscl-sc	-	15-35	50-100	1-3	slight	-	-	-	-	-
HDRhB1g1	754	120-150	WD	25-50	scl	sl	15-35	<15	<50	1-3	slight	8.28	0.18	0.64	12.54	100
HGHcB2	754	120-150	WD	>150	sl	scl-c	-	<15	>200	1-3	moderate	7.37	0.18	1.50	8.66	100
HGHhB1	754	120-150	WD	>150	scl	scl-c	-	<15	>200	1-3	slight	7.37	0.18	1.50	8.66	100
HGHhB1g1	754	120-150	WD	>150	scl	scl-c	15-35	<15	>200	1-3	slight	7.37	0.18	1.50	8.66	100
HGHhB2	754	120-150	WD	>150	scl	scl-c	-	<15	>200	1-3	moderate	7.37	0.18	1.50	8.66	100
HPRcB1	754	120-150	WD	50-75	sl	gsl-sc	-	15-35	50-100	1-3	slight	8.36	0.36	1.78	24.17	100
HPRiB2g1	754	120-150	WD	50-75	sc	gsl-sc	15-35	15-35	50-100	1-3	moderate	8.36	0.36	1.78	24.17	100
KLPhB2	754	120-150	WD	100-150	scl	scl-c	-	<15	100	1-3	moderate	6.07	0.14	2.82	9.23	100

*Symbols and abbreviations are according to Field Guide for LRI under Sujala-III Project, Karnatak

Marginally suitable lands (Class S3) for growing sorghum occupy an area of about 88 ha (16%) and are distributed in northeastern, central and eastern parts of the microwatershed. They have moderately limitation of rooting condition.

Table 7.2 Land suitability criteria for Sorghum

Crop requirement		Rating			
Soil –site characteristics	Unit	Highly suitable (S1)	Moderately Suitable (S2)	Marginally suitable (S3)	Not suitable (N)
Slope	%	2-3	3-8	8-15	>15
LGP	Days	120-150	120-90	<90	
Soil drainage	Class	Well to mod. Well drained	imperfect	Poorly/ excessively	V. poorly
Soil reaction	pH	6.0-8.0	5.5-5.9 8.1-8.5	<5.5 8.6-9.0	>9.0
Surface soil texture	Class	c, cl, sicl, sc	l, sil, sic	sl, ls	s, fragmental skeletal
Soil depth	cm	100-75	50-75	30-50	<30
Gravel content	% vol.	5-15	15-30	30-60	>60
Salinity (EC)	dSm ⁻¹	2-4	4-8	8-10	>10
Sodicity (ESP)	%	5-8	8-10	10-15	>15




Fig. 7.1 Land suitability map of Sorghum

7.2 Land Suitability for Maize (*Zea mays*)

Maize is the most important food crop grown in an area of 13.73 lakh ha in all the district of the state. The crop requirements for growing maize (Table 7.3) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing maize was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.2.

An area of about 158 ha (28%) in the microwatershed has soils that are highly suitable (Class S1) for growing maize crop and are distributed in the northwestern, southwestern, southern and eastern parts of the microwatershed. About 10 ha (2%) has soils that are moderately suitable (Class S2) for growing maize and are distributed in the northern and western parts of the microwatershed. They have minor limitations of rooting depth. The marginally suitable (Class S3) lands cover a maximum area of about 208 ha (62%) and occur in major part of the microwatershed. They have moderate limitations of rooting depth, texture and drainage.

Table 7.3 Land suitability criteria for Maize

Crop requirement		Rating			
Soil-site characteristics	Unit	Highly suitable(S1)	Moderately Suitable(S2)	Marginally suitable(S3)	Not suitable(N)
Slope	%	<3	3.5	5-8	
LGP	Days	>100	100-80	60-80	
Soil drainage	class	Well drained	Mod. to imperfectly	Poorly/excessively	V. poorly
Soil reaction	pH	5.5-7.5	7.6-8.5	8.6-9.0	
Surface soil texture	Class	l, cl, scl, sil	sl, siel, sic	c(s-s), ls	s, fragmental
Soil depth	cm	>75	50-75	25-50	<25
Gravel content	% vol.	<15	15-35	35-50	>50
Salinity (EC)	dSm ⁻¹	<1.0	1.0-2.0	2.0-4.0	
Sodicity (ESP)	%	<10	10-15	>15	




Fig. 7.2 Land suitability map of Maize

7.3 Land Suitability for Red gram (*Cajanus cajan*)

Red gram is one of the major pulse crop grown in an area of 8.23 lakh ha mainly in northern Karnataka in Bijapur, Kalaburgi, Raichur, Bidar, Belgaum, Dharwad and Bellary districts. The crop requirements for growing red gram (Table 7.4) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing red gram was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed is given in Figure 7.3.

The highly suitable (Class S1) lands for growing red gram occupy about 158 ha (28%) and are distributed in the major part of the microwatershed. An area of about 120 ha (21%) is moderately suitable (Class S2) and are distributed in northern, central and western parts of the microwatershed. They have minor limitation of drainage. Marginally suitable lands (Class S3) occupy a small area of about 10 ha (2%) and occur in the northern and western parts of the microwatershed. They have moderate limitation of rooting depth. An area of about 88 ha (16%) is currently not suitable for cultivation and are distributed in central, northeastern and northwestern parts of the microwatershed with severe limitations of rooting condition and texture.

Table 7.4 Land suitability criteria for Red gram

Crop requirement		Rating			
Soil –site characteristics	Unit	Highly suitable(S1)	Moderately Suitable(S2)	Marginally suitable(S3)	Not suitable(N)
Slope	%	<3	3-5	5-10	>10
LGP	Days	>210	180-210	150-180	<150
Soil drainage	class	Well drained	Mod. to well drained	Imperfectly drained	Poorly drained
Soil reaction	pH	6.5-7.5	5.0-6.5;7.6-8.0	8.0-9.0	>9.0
Surface soil texture	Class	l, scl, sil, cl, sl	sicl, sic, c(m)	ls	s,fragmental
Soil depth	cm	>100	85-100	40-85	<40
Gravel content	% vol.	<20	20-35	35-60	>60
Salinity (EC)	dSm ⁻¹	<1.0	1.0-2.0	>2.0	
Sodicity (ESP)	%	<10	10-15	>15	




Fig. 7.3 Land suitability map of Red gram

7.4 Land Suitability for Horse gram (*Macrotyloma uniflorum*)

Horse gram is the most important pulse crop grown in an area of 1.8 lakh ha in almost all the districts of the State. The crop requirements for growing horse gram (Table 7.5) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing horse gram was generated and the area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.4.

The highly suitable (Class S1) for growing horse gram occupy about 158 ha (28%) and are distributed in the northwestern, southwestern, southern and eastern parts of the microwatershed. An area of about 130 ha (23%) is moderately suitable (Class S2) for horse gram and are distributed in northern, western and central parts of the microwatershed. They have minor limitations of rooting depth and drainage. An area of about 88 ha (16%) is marginally suitable (Class S3) with moderate limitation of rooting condition and are distributed in the central, northeastern and southwestern parts of the microwatershed.

Table 7.5 Land suitability criteria for Horse gram

Crop requirement		Rating			
Soil –site characteristics	Unit	Highly suitable (S1)	Moderately Suitable (S2)	Marginally suitable (S3)	Not suitable(N)
Slope	%	<3	3-5	5-10	>10
LGP	Days				
Soil drainage	class	Well drained/ mod. well drained	imperfectly drained	Poorly drained	Very Poorly drained
Soil reaction	pH	6.0-8.5	8.5-9.0;5.5-5.9	9.1-9.5;5.0-5.4	>9.5
Surface soil texture	Class	l, sl, scl, cl, sc	ls, sic, sicl, c, ls	Heavy clays (>60%)	-
Soil depth	cm	50-75	25-50	<25	-
CaCO ₃ in root zone	% vol.	<15	15-25	25-30	>30
Salinity (ECe)	dSm ⁻¹	<1.0	1.0-2.0	>2.0	
Sodicity (ESP)	%	<10	10-15	>15	-




Fig. 7.4 Land suitability map of Horse gram

7.5 Land Suitability for Field bean (*Dolichos lablab*)

Field Bean is the most important pulse crop grown in an area of 0.68 lakh ha in almost all the districts of the State. The crop requirements (Table 7.4) for growing field bean were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing field bean was generated. The area extent and their geographical distribution of different suitability subclasses in different microwatersheds is given in Figure 7.5.

The area of about 66 ha (12%) is highly suitable (Class S1) for field bean and are distributed in the northwestern and eastern parts of the microwatershed. Maximum area of about 221 ha (40%) is moderately suitable (Class S2) and are distributed in the major part of the microwatershed. They have minor limitations of rooting depth, gravelliness and drainage. There is an area of about 88 ha (16%) in marginally suitable (Class S3) for field bean and are distributed in eastern and central parts of the microwatershed with moderate limitation of rooting condition.

Table 7.6 Land suitability criteria for Field Bean

Crop requirement		Rating			
Soil –site characteristics	Unit	Highly suitable(S1)	Moderately Suitable(S2)	Marginally suitable(S3)	Not suitable (N)
Slope	%	<3	3-5	5-10	>10
LGP	Days	>120	90-120	70-90	<70
Soil drainage	class	Well drained/mod. well drained	imperfectly drained	Poorly drained	Very Poorly drained
Soil reaction	pH	6.0-8.5	8.5-9.0;5.5-5.9	9.1-9.5;5.0-5.4	>9.5
Sub Surface soil texture	Class	l, sl, scl, cl, sc	sic, sicl, c	Heavy clays (>60%), ls	s
Soil depth	cm	>75	50-75	25-50	<25
CaCO ₃ in root zone	% vol.	<15	15-35	35-50	>50
Salinity (EC)	dsm ⁻¹	<1.0	1.0-2.0	>2.0	
Sodicity (ESP)	%	<10	10-15	15-20	>20




Fig. 7.5 Land suitability map of Field bean

7.6 Land Suitability for Groundnut (*Arachis hypogaea*)

Groundnut is one of the major oilseed crop grown in an area of 6.5 lakh ha in Karnataka in most of the districts either as rainfed or irrigated crop. The crop requirements for growing groundnut (Table 7.7) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and land suitability map for growing groundnut was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.6.

The highly suitable (Class S1) lands available for growing Groundnut is about 90 ha (16%) and are distributed in central and southern parts of the microwatershed. An area of 77 ha (14%) is moderately suitable (Class S2) for growing groundnut and are distributed in the northern, northwestern and eastern parts of the microwatershed. They have minor limitations of rooting depth and texture. Marginally suitable lands (class S3) for growing groundnut occupy area of about 208 ha (37%) and are distributed in the major part of the microwatershed. They have moderate limitations of rooting conditions, drainage and texture.

Table 7.7 Land suitability criteria for Groundnut

Crop requirement		Rating			
Soil-site characteristics	Unit	Highly suitable(S1)	Moderately Suitable(S2)	Marginally suitable(S3)	Not suitable (N)
Slope	%	<3	3-5	5-10	>10
LGP	Days	100-125	90-105	75-90	
Soil drainage	class	Well drained	mod. rained	imperfectly drained	Poorly drained
Soil reaction	pH	6.0-8.0	8.1-8.5;5.5-5.9	>8.5,<5.5	
Surface soil texture	Class	l, cl, sil, sc, sicl	sc, sic, c,	s, ls, sl,c (>60%)	s, fragmental
Soil depth	cm	>75	50-75	25-50	<25
Gravel content	% vol.	<35	35-50	>50	
CaCO ₃ in root zone	%	high	Medium	low	
Salinity (EC)	dSm ⁻¹	<2.0	2.0-4.0	4.0-8.0	
Sodicity (ESP)	%	<5	5-10	>10	




Fig. 7.6 Land suitability map of Ground nut

7.7 Land Suitability for Sunflower (*Helianthus annus*)

Sunflower is the most important oilseed crop grown in an area of 4.1 lakh ha in the State in all the districts. The crop requirements for growing sunflower (Table 7.8) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing sunflower was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.7.

An area of 36 ha (6%) is highly suitable (Class S1) lands for growing sunflower in the microwatershed and are distributed in eastern and northwestern parts of the microwatershed. The moderately suitable (Class S2) lands are found to occur in maximum area of about 242 ha (43%) and are distributed in the major part of the microwatershed with minor limitations of gravelliness, texture and drainage. The marginally suitable (Class S3) lands cover about 10 ha (2%) area and occur in the northern and northwestern parts of the microwatershed. They have moderate limitation of rooting depth. An area of about 88 ha (16%) is not suitable (Class N) for cultivation and are distributed in eastern and central parts of the microwatershed with severe limitations of rooting condition and texture.

Table 7.8 Land suitability criteria for Sunflower

Crop requirement		Rating			
Soil-site characteristics	Unit	Highly suitable(S1)	Moderately Suitable (S2)	Marginally suitable (S3)	Not suitable N)
Slope	%	<3	3-5	5-10	>10
LGP	Days	>90	80-90	70-80	<70
Soil drainage	class	Well drained	mod. well drained	Imper. drained	Poorly drained
Soil reaction	pH	6.5-8.0	8.1-8.5;5.5-6.4	8.6-9.0;4.5-5.4	>9.0 <4.5
Surface soil texture	Class	l, cl, sil, sc	scl, sic, c,	c (>60%), sl	ls, s
Soil depth	cm	>100	75-100	50-75	<50
Gravel content	% vol.	<15	15-35	35-60	>60
Salinity (EC)	dSm ⁻¹	<1.0	1.0-2.0	>2.0	
Sodicity (ESP)	%	<10	10-15	>15	




Fig. 7.7 Land suitability map of Sunflower

7.8 Land Suitability for Cotton (*Gossypium hirsutum*)

Cotton is the most important fibre crop grown in the State in about 6.6 lakh ha area in Raichur, Dharwad, Belgaum, Kalaburgi, Bijapur, Bidar, Bellary, Chitradurga and Chamarajnagar districts. The crop requirements for growing cotton (Table 7.9) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing cotton was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.8.

The highly suitable (Class S1) lands available for growing cotton is about 155 ha (27%) area in the microwatershed. They are distributed in the northern, western and eastern parts of the microwatershed. An area of about 101 ha (18%) has soils that are moderately suitable (Class S2) for growing cotton with minor limitations of rooting depth and texture. They are distributed in the northern, western and southern parts of the microwatershed. An area of about 119 ha (21%) is in marginally suitable (Class S3) with minor limitations of rooting depth and texture. They are distributed in western, central and eastern parts of the microwatershed.

Table 7.9 Land suitability criteria for Cotton

Crop requirement		Rating			
Soil-site characteristics	Unit	Highly suitable(S1)	Moderately Suitable (S2)	Marginally suitable(S3)	Not suitable (N)
Slope	%	1-2	2-3	3-5	>5
LGP	Days	180-240	120-180	<120	
Soil drainage	class	Well to moderately well	imperfectly drained	Poor somewhat excessive	Stagnant/ excessive
Soil reaction	pH	6.5-7.5	7.6-8.0	8.1-9.0	>9.0 ;>6.5
Surface soil texture	Class	sic, c	sicl, cl	si, sil, sc, scl, l	sl, s,ls
Soil depth	cm	100-150	60-100	30-60	<30
Gravel content	% vol.	<5	5-10	10-15	15-35
CaCO ₃ in root zone	%	<3	3-5	5-10	10-20
Salinity (EC)	dSm ⁻¹	2-4	4.0-8.0	8.0-12	>12
Sodicity (ESP)	%	5-10	10-20	20-30	>30




Fig. 7.8 Land suitability map of Cotton

7.9 Land Suitability for Onion (*Allium cepa*)

Onion is the most important vegetable crop grown in an area in Raichur, Dharwad, Belgaum, Gulbarga, Bijapur, Bidar, Bellary, Chitradurga and Chamarajnagar districts. The crop requirements for growing Onion (Table 7.10) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing Onion was generated and the area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.9.

The area of about 66 ha (12%) is highly suitable (Class S1) lands available for growing onion and are distributed in northwest and eastern parts of the microwatershed. An area of about 221 ha (40%) has soils that are moderately suitable (Class S2) with minor limitations of rooting depth, gravelliness, drainage and texture. They are distributed in the major part of the microwatershed. An area of 88 ha (16%) is marginally suitable (Class 3) for onion and are distributed in central and eastern parts of the microwatershed with moderate limitations of texture and rooting conditions.

Table 7.10 Land suitability criteria for Onion

Crop requirement		Rating			
Soil –site characteristics	Unit	Highly suitable(S1)	Moderately Suitable (S2)	Marginally suitable(S3)	Not suitable(N)
Mean temperature in growing season	⁰ C	20-30	30-35	35-40	>40
Slope	%	<3	3-5	5-10	>10
Soil drainage	class	Well drained	Moderately /imperfectly	Poor drained	Very poorly drained
Soil reaction	pH	6.5-7.3	7.3-7.8;5.0-5.4	7.8-8.4;<5.0	>8.4
Surface soil texture	Class	scl, sil, sl	sc, sicl, c (red soil)	sc, c (black soil)	ls
Soil depth	cm	>75	50-75	25-50	<25
Gravel content	% vol.	<15	15-35	35-60	60-80
Salinity (ECe)	dsm ⁻¹	<1.0	1.0-2.0	2.0-4.0	<4
Sodicity (ESP)	%	<5	5-10	10-15	>15




Fig. 7.9Land suitability map of Onion

7.10 Land Suitability for Potato (*Solanum tuberosum*)

Potato is the most important vegetable crop grown in Raichur, Dharwad, Belgaum, Chikkaballapur, Kolar, Gulbarga, Bijapur, Bidar, Bellary, Chitradurga and Chamarajnagar districts. The crop requirements for growing Potato (Table 7.11) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for

growing Potato was generated and the area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.10.

The highly suitable (Class S1) cover in an area of 66 ha (12%) for potato and are distributed in northwestern and eastern parts of the microwatershed. An area of about 101 ha (18%) is moderately suitable (Class S2) and are distributed in the northern, western and southern parts of the microwatershed. They have minor limitations of rooting depth and gravelliness. Marginally suitable lands (Class S3) for growing potato occupy an area of about 208 ha (37%) and occur in northern, western, southeastern and northeastern parts of the microwatershed. They have moderate limitations of rooting conditions, texture and drainage.

Table 7.11 Land suitability criteria for Potato

Crop requirement			Rating			
Soil –site characteristics		Unit	Highly suitable(S1)	Moderately Suitable (S2)	Marginally suitable(S3)	Not suitable (N)
Slope	Hills	%	<5	5-10	10-15	>15
	Plains	%	<3	3-5	5-8	>8
Mean temperature in growing season		⁰ C	16-25	26-30 13-15	31-32 10-12	>32 <10
Soil drainage		class	Well drained	Mod /imperfectly	Poor drained	V. poorly drained
Soil reaction		pH	5.5-6.5	6.6-8.2;5.0-5.4	>8.2;<5.0	-
Surface soil texture		Class	scl, sil	s, sil	s	
Soil depth		cm	75-100	50-75	25-50	<25
Stoniness		%	0-10	10-15	15-35	>35
Salinity (ECe)		dsm ⁻¹	<1.0	1.0-2.0	2.0-4.0	>4.0
Sodicity (ESP)		%	<10	10-15	>15	-




Fig. 7.10 Land suitability map of Potato

7.11 Land Suitability for Beans (*Phaseolus vulgaris*)

Beans is the most important vegetable crop grown in almost all the districts of Karnataka. The crop requirements for growing beans were matched with the soil–site characteristics and a land suitability map for growing beans was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.11.

The highly suitable (Class S1) occur in an area of 66 ha (12%) and are distributed in northwestern and eastern parts of the microwatershed. The moderately suitable (Class S2) lands are found to occur in maximum area of about 221 ha (40%) are distributed in the major part of the microwatershed. They have minor limitations of gravelliness, texture, rooting depth and drainage. An area of about 88 ha (16%) is in marginally suitable (Class S3) and are distributed in the eastern and central parts of the microwatershed. They have moderate limitations of rooting condition and texture.




Fig. 7.11 Land suitability map of Beans

7.12 Land Suitability for Beetroot (*Beta vulgaris*)

Beetroot is the most important vegetable crop grown in almost all the districts of Karnataka. The crop requirement for growing beetroot were matched with the soil site characteristics and a land suitability map for growing beetroot was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.12.

An area of 66 ha (12%) is highly suitable (Class S1) and are distributed in northwestern and eastern parts of the microwatershed. The moderately suitable (Class S2) lands are found to occur in an area of about 101 ha (18%) and are distributed in the northern, western and southern parts of the microwatershed. They have minor limitations of gravelliness and rooting depth. The marginally suitable (Class S3) lands cover an area of about 208 ha (37%) and occur in the major part of the microwatershed. They have moderate limitations of rooting condition, texture and drainage.




Fig. 7.12 Land suitability map of Beetroot

7.13 Land suitability for Mango (*Mangifera indica*)

Mango is the most important fruit crop grown in almost all the districts of the State. The crop requirements (Table 7.13) for growing mango were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing mango was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.13.

The highly suitable (Class S1) occur in an area of 66 ha (12 %) and are distributed in northwestern and eastern parts of the microwatershed. The moderately suitable (Class S2) lands are found to occur in a minor area of about 1 ha (<1%) and are distributed in the western parts of the microwatershed. They have minor limitation of rooting depth. An area of about 123 ha (21%) is in marginally suitable (Class S3) for mango and are distributed in southern, eastern and northeastern parts of the microwatershed with moderate limitations of rooting conditions and texture. Maximum area of about 185 ha (33%) is currently not suitable (Class N) for growing mango and are distributed in major part of the microwatershed.

Table 7.12 Land suitability criteria for Mango

Crop requirement			Rating			
soil-site characteristics		Unit	Highly suitable(S1)	Moderately Suitable(S2)	Marginally suitable(S3)	Not suitable(N)
climate	Temp in growing season	⁰ C	28-32	24-27 33-35	36-40	20-24
	Min. temp. before flowering	⁰ C	10-15	15-22	>22	
Soil moisture	Growing period	Days	>180	150-180	120-150	<120
Soil aeration	Soil drainage	class	Well drained	Mod. To imperfectly drained	Poor drained	Very poorly drained
	Water table	M	>3	2.50-3.0	2.5-1.5	<1.5
Nutrient availability	Texture	Class	sc, l, sil, cl	sl, sc, sic, l, c	c (<60%)	c (>60%),
	pH	1:2.5	5.5-7.5	7.6-8.55.0-5.4	8.6-9.04.0-4.9	>9.0<4.0
	OC	%	High	medium	low	
	CaCO ₃ in root zone	%	Non calcareous	<5	5-10	>10
Rooting conditions	Soil depth	cm	>200	125-200	75-125	<75
	Gravel content	% vol	Non gravelly	<15	15-35	>35
Soil toxicity	Salinity	dS/m	Non saline	<2.0	2.0-3.0	>3.0
	Sodicity	%	Non sodic	<10	10-15	>15
Erosion	Slope	%	<3	3-5	5-10	




Fig. 7.13 Land suitability map of Mango

7.14 Land suitability for Sapota (*Manilkara zapota*)

Sapota is the most important fruit crop grown in an area of 3.11 lakh ha in almost all the districts of the State. The crop requirements (Table 7.14) for growing sapota were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing sapota was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.14.

The highly suitable (Class S1) occur in an area of 66 ha (12%) and are distributed in northwestern and eastern parts of the microwatershed. The moderately suitable (class S2) lands are found to occur in an area of about 122 ha (22%) and are distributed in the southern, and eastern parts of the microwatershed. They have minor limitation of rooting conditions. The marginally suitable (class S3) lands cover a major area of about 131 ha (23%) and are distributed in the major part of the microwatershed. They have moderate limitations of texture and drainage. Currently not suitable (Class N) area is 56 ha (10 %) for sapota and are distributed in northeastern, central and southeastern parts of the microwatershed.

Table 7.13 Land suitability criteria for Sapota

Crop requirement			Rating			
Soil –site characteristics		Unit	Highly suitable(S1)	Moderately Suitable (S2)	Marginally suitable (S3)	Not suitable(N)
Climate	Temp. in growing season	⁰ C	28-32	33-36 24-27	37-42 20-23	>42 <18
Soil moisture	Growing period	Days	>150	120-150	90-120	<120
Soil aeration	Soil drainage	class	Well drained	Moderately well drained	Imperfectly drained	Poorly drained
Nutrient availability	Texture	Class	scl, l, cl, sil	sl, sicl, sc	c (<60%)	ls, s, c (>60%)
	pH	1:2.5	6.0-7.5	7.6-8.0;5.0-5.9	8.1-9.0;4.5-4.9	>9.0;<4.5
	CaCO ₃ in root zone	%	Non calcareous	<10	10-15	>15
Rooting conditions	Soil depth	cm	>150	75-150	50-75	<50
	Gravel content	% vol.	Non gravelly	<15	15-35	<35
Soil toxicity	Salinity	dS/m	Non saline	Up to 1.0	1.0-2.0	2.0-4.0
	Sodicity	%	Non sodic	10-15	15-25	>25
Erosion	Slope	%	<3	3-5	5-10	>10




Fig. 7.14 Land suitability map of Sapota

7.15 Land suitability for Guava (*Psidium guajava*)

Guava is the most important fruit crop grown in an area of 0.64 lakh ha in almost all the districts of the State. The crop requirements (Table 7.15) for growing guava were matched with the soil-site characteristics (7.1) and a land suitability map for growing guava was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.15.

The highly suitable (Class S1) cover in an area of 67 ha (12 %) and are distributed in northwestern and eastern parts of the microwatershed. The moderately suitable (Class S2) lands are found to occur in an area of about 122 ha (22%) and are distributed in the southern and eastern parts of the microwatershed. They have minor limitation of rooting conditions. The marginally suitable (Class S3) lands cover a small area of about 10 ha (2%) and are distributed in the northern and western parts of the microwatershed. They have moderate limitation of rooting depth. About 176 ha (31%) area is not suitable (Class N) for growing guava and are distributed in the major part of the microwatershed.

Table 7.14 Land suitability criteria for Guava

Crop requirement			Rating			
Soil –site characteristics	Unit		Highly suitable(S1)	Moderately Suitable (S2)	Marginally suitable (S3)	Not suitable(N)
Climate	Temperature in growing season	⁰ C	28-32	33-36 24-27	37-42 20-23	
Soil moisture	Growing period	Days	>150	120-150	90-120	<90
Soil aeration	Soil drainage	Class	Well drained	Mod. to imperfectly	poor	Very poor
Nutrient availability	Texture	Class	scl, l, cl, sil	sl,sicl,sic.,sc,c	c (<60%)	c (>60%)
	pH	1:2.5	6.0-7.5	7.6-8.0:5.0-5.9	8.1-8.5:4.5-4.9	>8.5:<4.5
	CaCO ₃ in root zone	%	Non calcareous	<10	10-15	>15
Rooting conditions	Soil depth	cm	>100	75-100	50-75	<50
	Gravel content	% vol.	<15	15-35	>35	
Soil toxicity	Salinity	dS/m	<2.0	2.0-4.0	4.0-6.0	
	Sodicity	%	Non sodic	10-15	15-25	>25
Erosion	Slope	%	<3	3-5	5-10	>10




Fig. 7.15 Land suitability map of Guava

7.16 Land Suitability for Banana (*Musa paradisiaca*)

Banana is one of the major fruit crop grown in an area of 1.02 lakh ha in Karnataka State. The crop requirements for growing banana (Table 7.16) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing banana was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.16.

The highly suitable (Class S1) occur in an area of 66 ha (12%) and are distributed in northwestern and eastern parts of the microwatershed. The moderately suitable (Class S2) lands are found to occur in a maximum area of about 211 ha (28%) and are distributed in the major part of the microwatershed. They have minor limitations of drainage, texture and gravelliness. The marginally suitable (Class S3) lands cover a major area of about 10 ha (2 %) and are distributed in the northern and western parts of the microwatershed. They have moderate limitation of rooting depth. An area of about 88 ha (15 %) is currently not suitable (Class N) for banana and are distributed in the northeastern, central and southeastern parts of the microwatershed with severe limitations of rooting condition and texture.

Table 7.15 Land suitability criteria for Banana

Crop requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately Suitable (S2)	Marginally suitable (S3)	Not suitable(N)
Climate	Temperature in growing season	⁰ C	26-33	34-36 24-25	37-38	>38
Soil aeration	Soil drainage	Class	Well drained	Moderately to imper. drained	Poorly drained	V.poorly drained
Nutrient availability	Texture	Class	l,cl, scl,sil	sicl,sc,c(<45%)	c (>45%),sic, sl	ls, s
	pH	1:2.5	6.5-7.0	7.1-8.5;5.5-6.4	>8.5;<5.5	
Rooting conditions	Soil depth	cm	>125	76-125	50-75	<50
	Stoniness	%	<10	10-15	15-35	>35
Soil toxicity	Salinity	dS/m	<1.0	1-2	>2	
	Sodicity	%	<5	5-10	10-15	>15
Erosion	Slope	%	<3	3-5	5-15	>15




Fig. 7.16 Land suitability map of Banana

7.17 Land Suitability for Jackfruit (*Artocarpus heterophyllus*)

Jackfruit is the most important fruit crop grown in almost all the districts of the State. The crop requirements for growing jackfruit were matched with the soil-site characteristics and a land suitability map for growing jackfruit was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.17.

An area of 66 ha (12 %) is highly suitable (Class S1) for growing jackfruits and are distributed in northwestern and eastern parts of the microwatershed. The moderately suitable (Class S2) lands are found to occur in a minor area of about 1 ha (<1%) are distributed in the northwestern parts of the microwatershed. They have minor limitation of rooting depth. The marginally suitable (Class S3) lands cover an area of about 132 ha (23 %) and are distributed in the southern, northern, northwestern, northeastern and eastern parts of the microwatershed. They have moderate limitation of rooting depth. About 176 ha (31 %) area is currently not suitable (Class N) for growing jackfruit and are distributed in the major part of the microwatershed.




Fig. 7.17 Land suitability map of Jackfruit

7.18 Land Suitability for Jamun (*Syzygium cumini*)

Jamun is the most important fruit crop grown in almost all the districts of the state. The crop requirements for growing jamun were matched with the soil-site characteristics and a land suitability map for growing jamun was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.18.

An area of 186 ha (33 %) is highly suitable (Class S1) for growing jamun and are distributed in the major part of the microwatershed. The moderately suitable (Class S2) lands are found to occur in a minor area of about 1 ha (<1%) and are distributed in the western parts of the microwatershed. They have minor limitation of rooting depth. The marginally suitable (Class S3) lands cover an area of about 126 ha (22%) and are distributed in the northern, southern and eastern parts of the microwatershed. They have moderate limitation of rooting depth. About 62 ha (11 %) area is currently not suitable (Class N) for growing jamun and are distributed in the western, central, southeastern and northeastern parts of the microwatershed.




Fig. 7.18 Land suitability map of Jamun

7.19 Land Suitability for Musambi (*Citrus limetta*)

Musambi is the most important fruit crop grown in almost all the districts of the state. The crop requirements for growing musambi were matched with the soil-site characteristics and a land suitability map for growing musambi was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.19.

The highly suitable (Class S1) lands cover about 186 ha (33%) and are distributed in the major part of the microwatershed. The moderately suitable (Class S2) lands are found to occur in a minor area of about 1 ha (1%) and are distributed in the western parts of the microwatershed. They have minor limitation of rooting depth. The marginally suitable (Class S3) lands cover an area of about 132 ha (23%) and are distributed in the northern, western, southern, northeastern and southeastern parts of the microwatershed. They have moderate limitation of rooting depth. About 56 ha (10%) area is currently not suitable (Class N) for growing musambi and are distributed in the northeastern, central and southeastern parts of the microwatershed.




Fig. 7.19 Land suitability map of Musambi

7.20 Land Suitability for Lime (*Citrus sp*)

Lime is the most important fruit crop grown in an area of 0.11 lakh ha in almost all the districts of the state. The crop requirements for growing lime (Table 7.17) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing lime was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed is given in Figure 7.20.

The highly suitable (Class S1) lands for growing lime cover an area of about 186 ha (33%) and are distributed in the major part of the microwatershed. The moderately suitable (Class S2) lands are found to occur in a minor area of about 1 ha (<1%) and are distributed in the western parts of the microwatershed. They have minor limitation of rooting depth. The marginally suitable (Class S3) lands cover a major area of about 132 ha (23%) and are distributed in the northern, western, southern and eastern part of the microwatershed. They have moderate limitation of rooting depth. About 56 ha (10%) area is currently not suitable (Class N) for growing lime and are distributed in the northeastern, central and southeastern parts of the microwatershed.

Table 7.16 Land suitability criteria for lime

Crop requirement			Rating			
Soil –site characteristics		Unit	Highly suitable(S1)	Moderately suitable(S2)	Marginally suitable(S3)	Not suitable(N)
Climate	Temp. in growing season	⁰ C	28-30	31-35 24-27	36-40 20-23	>40 <20
Soil moisture	Growing period	Days	240-265	180-240	150-180	<150
Soil aeration	Soil drainage	Class	Well drained	Mod. to imperfectly drained	poorly	Very poorly
Nutrient availability	Texture	Class	scl, l, sicl, cl, s	sc, sc, c	c (>70%)	s, ls
	pH	1:2.5	6.0-7.5	5.5-6.4 7.6-8.0	4.0-5.4 8.1-8.5	<4.0 >8.5
	CaCO ₃ in root zone	%	Non calcareous	Upto 5	5-10	>10
Rooting conditions	Soil depth	cm	>150	100-150	50-100	<50
	Gravel content	% vol.	Non gravelly	15-35	35-55	>55
Soil toxicity	Salinity	dS/m	Non saline	Upto 1.0	1.0-2.5	>2.5
	Sodicity	%	Non sodic	5-10	10-15	>15
Erosion	Slope	%	<3	3-5	5-10	




Fig. 7.20 Land suitability map of Lime

7.21 Land Suitability for Cashew (*Anacardium occidentale*)

Cashew is the most important plantation crop grown in an area of 1.24 lakh ha in almost all the districts. The crop requirements for growing Cashew were matched with the soil-site characteristics and a land suitability map for growing Cashew was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed is given in Figure 7.21.

An area of 66 ha (12 %) is highly suitable (Class S1) for cashew and are distributed in northwestern and eastern parts of the microwatershed. The moderately suitable (Class S2) lands are found to occur in an area of about 123 ha (22%) and are distributed in the western, southern and eastern parts of the microwatershed. They have minor limitation of rooting depth. The marginally suitable (Class S3) lands cover a minor area of about 10 ha (2%) and are distributed in the northern and western parts of the microwatershed. They have moderate limitation of rooting depth. About 176 ha (31%) area is currently not suitable (Class N) for growing cashew and are distributed in the major part of the microwatershed




Fig. 7.21 Land suitability map of Cashew

7.22 Land Suitability for Custard Apple (*Annona reticulata*)

Custard apple is the most important fruit crop grown in almost all the districts of the state. The crop requirements for growing custard apple were matched with the soil-site characteristics and a land suitability map for growing custard apple was generated.

The area extent and their geographic distribution of different suitability subclasses in the microwatershed is given in Figure 7.21.

An area of 187 ha (33%) is highly suitable (Class S1) lands for growing custard apple and are distributed in the major part of the microwatershed. The moderately suitable (Class S2) lands are found to occur in an area of about 132 ha (23%) and are distributed in the northern, western, southern, southeastern and northeastern parts of the microwatershed. They have minor limitation of rooting depth. The marginally suitable (Class S3) lands cover a major area of about 56 ha (10%) and are distributed in the central and eastern parts of the microwatershed with severe limitation of rooting condition.




Fig. 7.22 Land suitability map of Custard Apple

7.23 Land Suitability for Amla (*Phyllanthus emblica*)

Amla is the most important medicinal fruit crop grown in almost all the districts of the state. The crop requirements for growing amla were matched with the soil-site characteristics and a land suitability map for growing amla was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.23.

The highly suitable (Class S1) lands for growing amla cover about 187 ha (33%) and are distributed in the major part of the microwatershed. The moderately suitable (Class S2) lands are found to occur in an area of about 132 ha (23%) and are distributed in the northern, western, southern and eastern parts of the microwatershed. They have minor limitation of rooting depth. The marginally suitable (Class S3) lands cover an area

of about 56 ha (10%) and are distributed in the central, southeastern and northeastern parts of the microwatershed with severe limitation of rooting depth.




Fig. 7.23 Land suitability map of Amla

7.24 Land Suitability for Tamarind (*Tamarindus indica*)

Tamarind is the most important spice crop raised in all the districts of the state. The crop requirements for growing tamarind were matched with the soil-site characteristics and a land suitability map for growing tamarind was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed is given in Figure 7.24.

The highly suitable (Class S1) lands for growing Tamarind cover about 186 ha (33%) and are distributed in the major part of the microwatershed. The moderately suitable (Class S2) lands are found to occur in a minor area of about 1 ha (<1%) and are distributed in the northwestern parts of the microwatershed. They have minor limitation of rooting depth. The marginally suitable (Class S3) lands cover an area of about 132 ha (23%) and are distributed in the northern, western, southern and eastern parts of the microwatershed. They have moderate limitation of rooting depth. An area of 56 ha (10%) is currently not suitable (Class N) for growing tamarind and are distributed in the central, southeastern and northeastern parts of the microwatershed.




Fig. 7.24 Land suitability map of Tamarind

7.25 Land Suitability for Marigold (*Tagetes erecta*)

Marigold is the most important flower crop grown in an area of 1858 ha in almost all the districts of the State. The crop requirements (Table 7.18) for growing marigold were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing marigold was prepared. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.25.

The highly suitable (Class S1) lands for growing marigold cover about 158 ha (28%) and are distributed in the northwestern, eastern and southern parts of the microwatershed. The moderately suitable (Class S2) lands are found to occur in an area of about 130 ha (23%) and are distributed in northern, central and western parts of the microwatershed. They have minor limitations of rooting depth, texture and drainage. An area of about 88 ha (16%) is in marginally suitable (Class S3) and are distributed in the central, southeastern and northeastern parts of the microwatershed and moderate limitations of rooting condition and texture.

Table 7.17 Land suitability criteria for Marigold

Crop requirement			Rating			
Soil site characteristics	Unit		Highly suitable(S1)	Moderately suitable(S2)	Marginally suitable(S3)	Not suitable(N)
climate	Temperature in growing season	⁰ C	18-23	17-15 24-35	35-40 10-14	>40 <10
Soil aeration	Soil drainage	Class	Well drained	Moderately well drained	Imperfectly drained	Poorly drained
Nutrient availability	Texture	Class	l, sl, scl, cl, sil	sicl, sc, sic, c	c	ls, s
	pH	1:2.5	7.0-7.5	5.5-5.9;7.6-8.5	<5;>8.5	-
	CaCO ₃ in root zone	%	Non calcareous	Slightly calcareous	Strongly calcareous	-
Rooting conditions	Soil depth	cm	>75	50-75	25-50	<25
	Gravel content	% vol.	<15	15-35	>35	-
Soil toxicity	Salinity	ds/m	Non saline	Slightly	Strongly	-
	Sodicity(ESP)	%	<10	10-15	>15	-
Erosion	Slope	%	1-3	3-5	5-10	-




Fig. 7.25 Land suitability map of Marigold

7.26 Land Suitability for Chrysanthemum (*Chrysanthemum indicum*)

Chrysanthemum is the most important flower crop grown in an area of 803 ha in almost all the districts of the State. The crop requirements (7.19) for growing chrysanthemum were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing chrysanthemum was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.26.

An area of 66 ha (12%) is highly suitable (Class S1) lands for growing Chrysanthemum and are distributed in the northwestern and eastern parts of the microwatershed. The moderately suitable (Class S2) lands are found to occur in an area of about 221 ha (40%) and are distributed in the major part of the microwatershed. They have minor limitations of rooting depth, texture, drainage and gravelliness. An area of about 88 ha (41%) is in marginally suitable (Class S3) and are distributed in the central, southeastern and northeastern parts of the microwatershed with moderate limitations of rooting condition and texture.

Table 7.18 Land suitability criteria for Chrysanthemum

Crop requirement			Rating			
Soil site characteristics		Unit	Highly suitable(S1)	Moderately Suitable (S2)	Marginally suitable(S3)	Not suitable(N)
Climate	Temperature in growing season		18-23	17-15 24-35	35-40 10-14	>40 <10
Soil aeration	Soil drainage	Class	Well drained	Moderately well drained	Imperfectly drained	Poorly drained
Nutrient availability	Texture	Class	l ,sl, scl, cl, sil	sicl, sc, sic, c	c	ls, s
	pH	1:2.5	7.0-7.5	5.5-5.9;7.6-8.5	<5;>8.5	
	CaCO ₃ in root zone	%	Non calcareous	Slightly calcareous	Strongly calcareous	
Rooting conditions	Soil depth	cm	>75	50-75	25-50	<25
	Gravel content	% vol.	<15	15-35	>35	
Soil toxicity	Salinity	ds/m	Non saline	slightly	strongly	
	Sodicity(ESP)	%	<10	10-15	>15	-
Erosion	Slope	%	1-3	3-5	5-10	




Fig.26 Land suitability map of Chrysanthemum

7.27 Land Suitability for Turmeric (*Curcuma longa*)

Turmeric is the most important spice crop grown in an area of 1.39 lakh ha in almost all the districts of the State. The crop requirements for growing turmeric (Table 7.12) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing turmeric was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.27.

An area of 66 ha (12%) is highly suitable (Class S1) area of turmeric and are distributed in northwestern, southeast and northeastern parts of the microwatershed. The moderately suitable (Class S2) lands are found to occur in an area of about 101 ha (18%) and are distributed in the northern, southern and western parts of the microwatershed. They have minor limitations of gravelliness and rooting depth. The marginally suitable (Class S3) lands cover an area of about 208 ha (37%) and occur in the major part of the microwatershed. They have moderate limitations of rooting conditions, texture and drainage.

Table 7.19 Land suitability criteria for Turmeric

Crop requirement			Rating			
Soil –site characteristics		Unit	Highly suitable(S1)	Moderately suitable(S2)	Marginally suitable(S3)	Not suitable(N)
Climate	Temperature in growing season	⁰ C	28-32	20-27 33-37	10-19 38-40	<10 >40
Soil aeration	Soil drainage	Class	Well drained	Mod. well drained	Imperfectly drained	Poorly drained
Nutrient availability	Texture	Class	l, cl, scl, sl	sc, sic, sicl	c(40-60%), ls	Stony heavy clay>60%
	pH	1:2.5				
	Available nutrient status (NPK)	Fertility rating class	high	medium	low	
Rooting conditions	Soil depth	cm	>75	50-75	25-50	<25
Erosion	Slope	%	<3	3-8	8-15	>15 mm




Fig. 7.27 Land suitability map of Turmeric

7.28 Land Management Units (LMUs)

The 27 soil map units identified in Bachanahalli-1 microwatershed have been regrouped into 7 Land Management Units (LMU's) for the purpose of preparing Proposed Crop Plan. Land Management Units are grouped based on the similarities in respect of the type of soil, the depth of the soil, the surface soil texture, gravel content, AWC, slope, erosion etc. and a Land Management Units map (Fig. 7.28) has been generated. These Land Management Units are expected to behave similarly for a given level of management.

The map units that have been grouped into 7 land management units along with brief description of soil and site characteristics are given below.

LMU	Soil map units number	Mapping unit	Soil and site characteristics
1	1, 2, 14, 15, 16, 17	ARKhB1,ARKhB2g1, HGHcB2,HGHhB1 HGHhB1g1,HGHhB2	very deep (>150 cm),dark reddish brown to very dusky red sandy clay to clay soils andvery darkbrown and dark reddish brown sandy clay loam soils
2	3, 4, 5	BMBhB2, BMBiB1 BMBmB1	very deep (>150 cm),very dark greyish brown to dark grey and very darkbrown clayey soils
3	20	KLPhB2	deep (100-150 cm), dark reddish brown to dark red gravelly sandy clay loam to sandy clay soils
4	9, 10, 11, 12	GPRcB1,GPRcC2g1 GPRhB1g1 GPRiB1	moderately deep (75-100 cm),have dark brown to dark reddish brown and reddish brown gravelly sandy clay loam to sandy clay soils
5	18, 19	HPRcB1 HPRiB2g1	moderately shallow (50-75 cm),dark brown to very dark brown gravelly sandy clay loam to sandy clay soils
6	6, 7, 8	BMDhB1 BMDhB1g1 BMDhC2g1	shallow (25-50 cm),dark brown to dark greyish brown clayey soils
7	13	HDRhB1g1	shallow (25-50 cm),dark reddish brown to dusky red sandy clay loam to sandy clay soils




Fig. 7.28 Land management units map- Bachanahalli-1Microwatershed

7.29 Proposed Crop Plan for Bachanahalli -1 Microwatershed

After assessing the land suitability for the 27 crops, the proposed crop plan has been generated for the 3 identified LMUs by considering only the highly (class S1) and moderately (class S2) suitable lands for each of the 27 crops. The resultant proposed crop plan is presented below in Table 7.20

Table 7.20 Proposed Crop Plan for Bachanahalli-1 Microwatershed

LMU	Mapping Units	Survey Number	Field crops/Forestry	Suitable Horticulture crops under irrigation	Horticulture crops with interventions	Suitable interventions
LMU 1 66 ha (12%)	1, 2, 14, 15, 16, 17 (>150 cm)	Bachanahalli: 1,8,9,11,15,28,3,40,41,42,5,50, Berambadi: 40,41,42,49,50,62,63,64,65,66,67,68,69,70,72,75,77,78, 79, 86,87,355,	Maize, Sorghum, Redgram, Cotton, Sugarcane, Sunflower, Multiple crop rotation: Redgram+Maize Redgram+Groundnut Pulses+Ragi Pulses+Sorghum	Mango, Sapota, Guava, Lime, Turmeric, Banana, Beans, Tomato, Bhendi	Flower crops: Marigold, Chrysanthemum Perennial components: Mango, Sapota, Custard apple, Amla, Lime, Musambi Vegetables: Chilli, Bhendi	Drip irrigation, Mulching, crop suitable conservation practices
LMU 2 120 ha (21%)	3, 4, 5 (>150 cm)	Bachanahalli: 6,7,10,12,13,14,16, 29,37 Berambadi: 53,54,55,60,61,73,123,315,324,325,326,327,328,340,341,346,350,366, 367,370, Maddur: 33,56,62,63,64,66, 70	Sorghum, Redgram, Cotton, Sunflower, Sugarcane Multiple crop rotation: Redgram+Fodder sorghum Pulses+Sorghum	Vegetables: Beetroot, Banana, Lime, Tomato, Beans, Bhendi	Flower crops: Marigold, Chrysanthemum Perennial components: Custard apple, Amla, Lime, Vegetables: Chilli, Bhendi	Drip irrigation, Mulching, crop suitable conservation practices
LMU 3 1 ha (<1%)	20 (100-150 cm)	Bachanahalli: 43	Maize, Sorghum, Redgram, Cotton, Sugarcane, Sunflower Multiple crop rotation: Redgram+Maize Redgram+Groundnut Pulses+Sorghum	Vegetables: Tomato, Beetroot, Potato, Beans, Bhendi, Turmeric	Perennial components: Sapota, lime Flower crops: Marigold, Chrysanthemum Vegetables: Chilli, Bhendi	Drip irrigation, Mulching, Crop suitable conservation practices
LMU 4 122 ha (22%)	9, 10, 11, 12 (75-100 cm)	Berambadi: 34,35,36,37,38,39,43,57,58,76,88,89,124,125,126,127,130,31	Maize, Sorghum, Cotton, Ragi, Sunflower, Pulses+Sorghum	Fieldbean, Beetroot, Onion,	Perennial components: Sapota, Guava Flower crops:	Drip irrigation, Mulching, Crop suitable

		6,317,320,321,322,323,342,343,344,356,357,358,359,362,363,364		Turmeric, Tomato	Marigold, Chrysanthemum Vegetables: Bhendi, Chilli	conservation practices
LMU 5 10 ha (2%)	18, 19 (50-75 cm)	Bachanahalli: 23,24,53 Maddur: 61,69	Ragi, Groundnut, Maize, Sorghum, Cotton, Pulses+Sorghum	Fieldbean, Beetroot, Onion, Turmeric, Tomato, Banana	Custard apple, Ber, Aonla Flower crops: Marigold, Chrysanthemum, Gaillardia Vegetables: Bhendi, Cluster bean	Drip irrigation, Mulching, Crop suitable conservation practices
LMU 6 52 ha (9%)	6, 7, 8 (25-50 cm)	Berambadi: 51,52,56,128,129,131,132,318,319,347,368,	Horsegram, Bengal gram	Custard apple, Amla	Custard apple, Ber	Drip irrigation, Mulching, Crop suitable conservation practices
LMU 7 5 ha (1%)	13 (25-50 cm)	Berambadi: 90,91	Groundnut, Horsegram	Custard apple, Amla	Custard apple, Ber	Drip irrigation, Mulching, Crop suitable conservation practices

SOIL HEALTH MANAGEMENT

8.1 Soil Health

Soil is fundamental to crop production. Without soil, no food could be produced nor would livestock be fed on a large scale. Because it is finite and fragile, soil is a precious resource that requires special care from its users.

Soil health or the capacity of the soil to function is critical to human survival. Soil health has been defined as: “the capacity of the soil to function as a living system without adverse effect on the ecosystem”. Healthy soils maintain a diverse community of soil organisms that help to form beneficial symbiotic associations with plant roots, recycle essential plant nutrients, improve soil structure with positive repercussions for soil, water and nutrient holding capacity and ultimately improve crop production and also contribute to mitigating climate change by maintaining or increasing its carbon content.

Functional interactions of soil biota with organic and inorganic components, air and water determine a soil’s potential to store and release nutrients and water to plants and to promote and sustain plant growth. Thus, maintaining soil health is vital to crop production and conserve soil resource base for sustaining agriculture.

The most important characteristics of a healthy soil are

- Good soil tilth
- Sufficient soil depth
- Good water storage and good drainage
- Adequate supply, but not excess of nutrients
- Large population of beneficial organisms
- Small proportion of plant pathogens and insect pests
- Low weed pressure
- Free of chemicals and toxins that may harm the crop
- Resistance to degradation
- Resilience when unfavourable conditions occur

Characteristics of Bachanahalli-1 Microwatershed

- ❖ The soil phases identified in the microwatershed belonged to the soil series of ARK (36 ha) BMB (120 ha), BMD (51 ha), GPR (122 ha), HDR (5 ha), HGH (31 ha), HPR (10 ha) and KLP (1 ha).
- ❖ As per land capability classification, about 66 per cent area in the microwatershed falls under arable land category (Class II and III). The major limitations identified in the arable and non arable lands were soil, wetness and erosion.
- ❖ On the basis of soil reaction, an area of about 197 ha (35%) is moderately to slightly acid (pH 6.0-6.5), about 125 ha (22%) is under neutral (pH 6.5-7.3) and about 55 ha (9%) is slightly to moderately alkaline (pH 7.3- 8.4).

Soil Health Management

The following actions are required to improve the current land husbandry practices that provide a sound basis for the successful adoption of sustainable crop production system.

Acid soils

Acid soils occupy an area of about 197 ha in the microwatershed. The following measures recommended for reclaiming acid soils

1. Growing of crops suitable for a particular soil pH.
2. Ameliorating the soils through the application of amendments (liming materials).

Liming materials:

1. CaCO_3 (Calcium Carbonate). More than 90% use in India.
2. Dolomite [$\text{Ca Mg} (\text{CO}_3)_2$]
3. Quick lime (CaO)
4. Slaked lime [$\text{Ca} (\text{OH})_2$]

For normal pH and pH-4.8 (35 t/ha) and pH 6 .0-7.0 (4 t/ha) lime is required

Alkaline soils

(Slightly alkaline to moderately alkaline soils)

1. Regular addition of organic manure, green manuring, green leaf manuring, crop residue incorporation and mulching needs to be taken up to improve the soil organic matter status.
2. Application of biofertilizers (Azospirillum, Azotobacter, Rhizobium).
3. Application of 25% extra N and P (125 % RDN&P).
4. Application of ZnSO_4 – 12.5 kg/ha (once in three years).
5. Application of Boron – 5kg/ha (once in three years).

Neutral soils

1. Regular addition of organic manure, green manuring, green leaf manuring, crop residue incorporation and mulching needs to be taken up to improve the soil organic matter status.
2. Application of biofertilizers, (Azospirillum, Azotobacter, Rhizobium).
3. Application of 100 per cent RDF.
4. Need based micronutrient applications.

Besides the above recommendations, the best transfer of technology options are also to be adopted.

Soil Degradation

Soil erosion is one of the major factor affecting the soil health in the microwatershed. Out of total 567 ha area in the microwatershed, about 164 ha is suffering

from moderate erosion. These areas need immediate soil and water conservation and, other land development and land husbandry practices for restoring soil health.

Dissemination of Information and Communicate of Benefits

Any large scale implementation of soil health management requires that supporting information is made available widely, particularly through channels familiar to farmers and extension workers. Given the very high priority attached to soil health especially by the Central Government on issuing Soil-Health Cards to all the farmers, media outlets like regional, state and national newspapers, Radio and Dooradarshan programs in local languages but also modern information and communication technologies such as cellular phones and the Internet, which can be much more effective in reaching younger farmers.

Inputs for Net Planning and Interventions needed

Net planning in IWMP is focusing on preparation of

1. Soil and Water Conservation Plans for each plot or farm.
2. Productivity enhancement measures/ interventions for existing crops/livestock/other farm enterprises.
3. Diversification of farming mainly with perennial horticultural crops and livestock.
4. Improving livelihood opportunities and income generating activities.

In this connection, how various outputs of Sujala-III are of use in addressing these objectives of Net Planning are briefly presented below.

- ❖ **Soil Depth:** The depth of a soil decides the amount of moisture and nutrients it can hold, what crops can be taken up or not, depending on the rooting depth and the length of growing period available for raising any crop. Deeper the soil, better for a wide variety of crops. If sufficient depth is not available for growing deep rooted crops, either choose medium or short duration crops or deeper planting pits need to be opened and additional good quality soil brought from outside has to be filled into the planting pits.
- ❖ **Surface soil texture:** Lighter soil texture in the top soil means, better rain water infiltration, less run-off and soil moisture conservation, less capillary rise and less evaporation losses. Lighter surface textured soils are amenable to good soil tilth and are highly suitable for crops like groundnut, root vegetables (carrot, radish, potato etc) but not ideal for crops that need stagnant water like lowland paddy. Heavy textured soils are poor in water infiltration and percolation. They are prone for sheet erosion; such soils can be improved by sand mulching. The technology that is developed by the AICRP-Dryland Agriculture, Vijayapura, Karnataka can be adopted.
- ❖ **Gravelliness:** More gravel content is favorable for run-off harvesting but poor in soil moisture storage and nutrient availability. It is a significant parameter that decides the kind of crop to be raised.

- ❖ **Land Capability Classification:** The land capability map shows the areas suitable and not suitable for agriculture and the major constraints in each of the plot/survey number. Hence, one can decide what kind of enterprise is possible in each of these units. In general, erosion and soil are the major constraints in Bachanahalli-1 microwatershed.
- ❖ **Organic Carbon:** The OC content is low (<5%) in about 54 ha (10%), medium (0.5-0.75%) in about 285 ha (50%) area, High (>0.75%) in about 36 ha (6%). The areas that are low and medium in OC need to be further improved by applying farmyard manure and rotating crops with cereals and legumes or mixed cropping.
- ❖ **Promoting green manuring:** Growing of green manuring crops costs Rs. 1250/ha (green manuring seeds) and about Rs. 2000/ha towards cultivation that totals to Rs. 3250/- per ha. On the other hand, application of organic manure @ 10 tons/ha costs Rs. 5000/ha. The practice needs to be continued for 2-3 years or more. Nitrogen fertilizer needs to be supplemented by 25% in addition to the recommended level in 335 ha area where OC is medium (0.5-0.75%). For example, for rainfed maize, recommended level is 50 kg N per ha and an additional 12 kg /ha needs to be applied for all the crops grown in these plots.
- ❖ **Available Phosphorus:** In 375 ha (66%) area, the available phosphorus is medium (23-57 kg/ha). Hence for all the crops, 25% additional P-needs to be applied, where it is medium.
- ❖ **Available Potassium:** Available potassium is medium (145-337 kg/ha) in 37 ha (7%) area of the microwatershed, about 338 ha (60%) area is high (<337 kg/ha) in available potassium and maximum area is high (>337 kg/ha) in available potassium. Hence, in all those plots where available potassium is medium, for all the crops, additional 25 % potassium may be applied.
- ❖ **Available Sulphur:** Available sulphur is a very critical nutrient for oilseed crops. It is low (<10 ppm) in a maximum of 375 ha (66%) area of the microwatershed. These areas need to be applied with magnesium sulphate or gypsum or Factamphos (p) fertilizer (13% sulphur) for 2-3 years for the deficiency to be corrected.
- ❖ **Available Boron:** An area of about 375 ha (66%) is low (<0.5 ppm) in available boron content. The areas with low and medium in boron content need to be applied with sodium borate @ 10kg/ha as a soil application or 0.2% borax as foliar spray to correct the deficiency.
- ❖ **Available Iron:** It is sufficient (>4.5 ppm) in entire area of the microwatershed. To manage iron deficiency, iron sulphate @ 25kg /ha needs to be applied for 2-3 years.
- ❖ **Available manganese:** It is sufficient in the entire area of the microwatershed.
- ❖ **Available copper:** It is sufficient in the entire area of the microwatershed.
- ❖ **Available Zinc:** It is deficient (<0.6 ppm) in 347 ha (61%) area in the microwatershed and an area about 28 ha (5%) is available zinc in sufficient (>0.6 ppm). Application of zinc sulphate @25kg/ha is to be recommended.
- ❖ **Soil alkalinity:** The microwatershed has 53 ha (9%) area with soils that are slightly to moderately alkaline. These areas need application of gypsum and wherever calcium is in


excess, iron pyrites and element sulphur can be recommended. Management practices like treating repeatedly with good quality water to drain out the excess salts and provision of subsurface drainage and growing of salt tolerant crops like Casuarina, Acacia, Neem, Ber etc, are recommended.

Land Suitability for various crops: Areas that are highly, moderately and marginally suitable for growing various crops are indicated. Along with the suitability, various constraints that are limiting the productivity are also indicated. For example, in case of cotton, gravel content, rooting depth and salinity/alkalinity are the major constraints in various plots. With suitable management interventions, the productivity can be enhanced. In order to increase water holding capacity of light textured soils, growing of green manure crops and application of organic manure is recommended.

SOIL AND WATER CONSERVATION TREATMENT PLAN

For preparing soil and water conservation treatment plan for Bachanahalli-1 microwatershed, the land resource inventory database generated under Sujala-III project has been transformed as information through series of interpretative (thematic) maps using soil phase map as a base. The various thematic maps (1:7920 scale) generated were

- Soil depth
 - Surface soil texture
 - Soil gravelliness
 - Available water capacity
 - Soil slope
 - Soil erosion
 - Land capability
 - Present land use /land cover
 - Crop suitability
 - Rainfall
 - Hydrology
 - Water Resources
 - Socio-economic data
 - Contour plan with existing features- Network of waterways, pothissa boundaries, cut up/ minor terraces etc.
 - Cadastral map (1:7920 scale)
 - Satellite imagery (1:7920 scale)
- Apart from these, Hand Level/ Hydro Marker/ Dumpy Level/ Total Station and Kathedars' List has to be collected.



Steps for Survey and Preparation of Treatment Plan

The boundaries of Land User Groups' and Survey No. boundaries are traced in the field.

- Naming of user groups and farmers
- Identification of arable and non arable lands
- Identification of drainage lines and gullies
- Identification of non treatable areas
- Identification of priority areas in the arable lands
- Treatment plan for arable lands
- Location of water harvesting and recharge structures

9.1 Treatment Plan

The treatment plan recommended for arable lands is briefly described below.

9.1.1 Arable Land Treatment

A. BUNDING

Steps for Survey and Preparation of Treatment Plan		USER GROUP-1	
Cadastral map (1:7920 scale) is enlarged to a scale of 1:2500 scale Existing network of waterways, pothissa boundaries, grass belts, natural drainage lines/ watercourse, cut ups/ terraces are marked on the cadastral map to the scale Drainage lines are demarcated into		<p>CLASSIFICATION OF GULLIES</p> <p>ಕೊರಕಲಿನ ವರ್ಗೀಕರಣ</p> <p>UPPER REACH 15 Ha.</p> <p>MIDDLE REACH 15+10=25 ಹೆ.</p> <p>LOWER REACH 25 ಹೆಕ್ಕರ ಗಿಂತ ಅಧಿಕ</p> <p>POINT OF CONCENTRATION</p>	
Small gullies	(up to 5 ha catchment)		
Medium gullies	(5-15 ha catchment)		
Ravines	(15-25 ha catchment) and		
Halla/Nala	(more than 25ha catchment)		

Measurement of Land Slope

Land slope is estimated or determined by the study and interpretation of contours or by measurement in the field using simple instruments like Hand level or Hydromarker.

		<p>HYDRO MARKER (WATER TUBE)</p> <p>FALL: 1.5 - 0.6 = 0.9 m.</p>
--	--	---

Vertical and Horizontal intervals between bunds as recommended by the Watershed Development Department.

Slope percentage	Vertical interval (m)	Corresponding Horizontal Distance (m)
2 - 3%	0.6	24
3 - 4%	0.9	21
4 - 5%	0.9	21
5 - 6%	1.2	21
6 - 7%	1.2	21

Note: i) The above intervals are maximum.

(ii) Considering the slope class and erosion status ($A_1 \dots A=0-1\%$, 1= slight erosion)) the intervals have to be decided.

Bund length recording: Considering the contour plan and the existing grass belts/partitions, the bunds are aligned and lengths are measured.

Section of the Bund

Bund section is decided considering the soil texture class and gravelliness class (bg_0-b = loamy sand, $g_0 \leq 15\%$ gravel). The recommended Sections for different soils are given below.

Recommended Bund Section


Top width(m)	Base width(m)	Height (m)	Side slope (Z:1;H:V)	Cross section (sq m)	Soil Texture	Remarks
0.3	0.9	0.3	01:01	0.18	Sandy loam	Vegetative bund
0.3	1.2	0.3	1.5:1	0.225	Sandy clay	
0.3	1.2	0.5	0.9:1	0.375	Red gravelly soils	
0.3	1.2	0.6	0.75:1	0.45		
0.3	1.5	0.6	01:01	0.54	Red sandy loam	
0.3	2.1	0.6	1.5:1	0.72	Very shallow black soils	
0.45	2	0.75	01:01	0.92		
0.45	2.4	0.75	1.3:1	1.07	Shallow black soils	
0.6	3.1	0.7	1.78:1	1.29	Medium black soils	
0.5	3	0.85	1.47:1	1.49		

Formation of Trench cum Bund

Dimensions of the Borrow Pits/ Trenches to be excavated (machinery are decided considering the Bund Section).


Details of Borrow Pit dimensions are given below

TRENCH CUM BUND



IDEAL FOR HORTICULTURE CROPS

'A' FRAME FOR INTERBUND MANAGEMENT



1. ಸಮಸಾಹಿತ್ಯ ಉಳಿಸುವುದು
2. ಸಮಸಾಹಿತ್ಯ ಬಿಡುಗಡೆ/ನಾಶ

Size of Borrow Pits/ Trench recommended for Trench cum Bund (by machinery)

Bund section	Bund length	Earth quantity	Pit				Berm (pit to pit)	Soil depth class
m ²	m	m ³	L(m)	W(m)	D(m)	Quantity (m ³)	m	
0.375	6	2.25	5.85	0.85	0.45	2.24	0.15	Shallow
0.45	6	2.7	5.4	1.2	0.43	2.79	0.6	Shallow
0.45	6	2.7	5	0.85	0.65	2.76	1	Moderately Shallow
0.54	5.6	3.02	5.5	0.85	0.7	3.27	0.1	Moderately shallow
0.54	5.5	2.97	5	1.2	0.5	3	0.5	Shallow
0.72	6.2	4.46	6	1.2	0.7	5.04	0.2	Moderately shallow
0.72	5.2	3.74	5.1	0.85	0.9	3.9	0.1	Moderately deep

B. Waterways

1. Existing waterways are marked on the cadastral map (1:7920 scale) and their dimensions are recorded.
2. Considering the contour plan of the MWS, additional waterways/ modernization of the existing ones can be thought of.
3. The design details are given in the Manual.

C. Farm Ponds

Waterways and the catchment area will give an indication on the size of the Farm Pond. Location of the pond can be decided based on the contour plan/ field condition and farmers' need/desire.

D. Diversion channel

Existing EPT/ CPT are marked on the cadastral map. Looking to the need, these can be modernized or fresh diversion channel can be proposed and runoff from this can be stored in *Gokatte/* Recharge ponds.

9.1.2 Non-Arable Land Treatment

Depending on the gravelliness and crops preferred by the farmers, the concerned authorities can decide appropriate treatment plan. The recommended treatments may be Contour Trench, Staggered Trench, Crescent Bund, Boulder Bund or Pebble Bunds are formed in the field.

9.1.3 Treatment of Natural Water Course/ Drainage Lines

- a) The cadastral map has to be updated as regards the network of drainage lines (gullies/ *nalas*/ *hallas*) and existing structures are marked to the scale and storage capacity of the existing water bodies are documented.
- b) The drainage line will be demarcated into Upper Reach, Middle Reach and Lower Reach.
- c) Considering the Catchment, *Nala* bed and bank conditions, suitable structures are decided.
- d) Number of storage structures (Check dam/ *Nala* bund/ Percolation tank) will be decided considering the commitments and available runoff in water budgeting and quality of water in the wells and site suitability.
- e) Detailed Leveling Survey using Dumpy Level / Total Station has to be carried out to arrive at the site-specific designs as shown in the Manual.
- f) The location of ground water recharge structures are decided by examining the lineaments and fracture zones from geological maps.
- g) Rainfall intensity data of the nearest Rain gauge station is considered for Hydrologic Designs.
- h) Silt load to the Storage/Recharge structures is reduced by providing vegetative, boulder and earthen checks in the natural water course. Location and design details are given in the Manual.

9.2 Recommended Soil and Water Conservation Measures

The appropriate conservation structures best suited for each of the land parcel/ survey number (Appendix-I) are selected based on the slope per cent, severity of erosion, amount of rainfall, land use and soil type. The different kinds of conservation structures recommended are

1. Graded / Strengthening of bunds
2. Trench cum Bunds (TCB)
3. Trench cum Bunds / Strengthening
4. Crescent Bunds

A map (Fig. 9.1) showing soil and water conservation plan with different kinds of structures recommended has been generated which shows the spatial distribution and extent of area. A maximum area of about 255 ha (45%) requires Trench cum bunding and about 120 ha (21%) area needs Graded bunds or strengthening of existing bunds. The conservation plan generated may be presented to all the stakeholders including farmers and after including their suggestions, the conservation plan for the microwatershed may be finalised in a participatory approach.




Fig. 9.1 Soil and water conservation plan map of Bachanahalli-1 Microwatershed

9.3 Greening of Microwatershed

As part of the greening programme in the watersheds, it is envisaged to plant a variety of horticultural and other tree plants that are edible, economical and produce lot of biomass which helps to restore the ecological balance in the watersheds. The lands that are suitable for greening programme are non-arable lands (land capability classes V, VI and VII) and also the lands that are not suitable or marginally suitable for growing annual and perennial crops. The method of planting these trees is given below.

It is recommended to open pits during the 1st week of March along the contour and heap the dug out soil on the lower side of the slope in order to harness the flowing water and facilitate weathering of soil in the pit. Exposure of soil in the pit also prevents spread of pests and diseases due to scorching sun rays. The pits should be filled with mixture of soil and organic manure during the second week of April and keep ready with sufficiently tall seedlings produced either in poly bags or in root trainer nurseries so that planting can be done during the 2nd or 3rd week of April depending on the rainfall.

The tree species suitable for the area considering rainfall, temperature and adaptability is listed below; waterlogged areas are recommended to be planted with species like Neral (*Sizyziumcumini*) and Bamboo. Dry areas are to be planted with species like Honge, Bevu, Seetapha*etc.*

Dry Deciduous Species			Temp (°C)	Rainfall(mm)
1.	Bevu	<i>Azadiracta indica</i>	21–32	400 –1,200
2.	Tapasi	<i>Holoptelia integrifolia</i>	20-30	500 - 1000
3.	Seetaphal	<i>Anona Squamosa</i>	20-40	400 - 1000
4.	Honge	<i>Pongamia pinnata</i>	20 -50	500–2,500
5.	Kamara	<i>Hardwickia binata</i>	25 -35	400 - 1000
6.	Bage	<i>Albezzia lebbek</i>	20 - 45	500 - 1000
7.	Ficus	<i>Ficus bengalensis</i>	20 - 50	500–2,500
8.	Sisso	<i>Dalbargia Sissoo</i>	20 - 50	500 -2000
9.	Ailanthus	<i>Ailanthus excelsa</i>	20 - 50	500 - 1000
10.	Hale	<i>Wrightia tinctoria</i>	25 - 45	500 - 1000
11.	Uded	<i>Steriospermum chelanoides</i>	25 - 45	500 -2000
12.	Dhupa	<i>Boswellia Serrata</i>	20 - 40	500 - 2000
13.	Nelli	<i>Emblica Officinalis</i>	20 - 50	500 -1500
14.	Honne	<i>Pterocarpus marsupium</i>	20 - 40	500 - 2000
Moist Deciduous Species				
15.	Teak	<i>Tectona grandis</i>	20 - 50	500-5000
16.	Nandi	<i>Legarstroemia lanceolata</i>	20 - 40	500 - 4000
17.	Honne	<i>Pterocarpus marsupium</i>	20 - 40	500 - 3000
18.	Mathi	<i>Terminalia alata</i>	20 -50	500 - 2000
19.	Shivane	<i>Gmelina arboria</i>	20 -50	500 -2000
20.	Kindal	<i>T.Paniculata</i>	20 - 40	500 - 1500
21.	Beete	<i>Dalbargia latifolia</i>	20 - 40	500 - 1500
22.	Tare	<i>T. belerica</i>	20 - 40	500 - 2000
23.	Bamboo	<i>Bambusa arundinasia</i>	20 - 40	500 - 2500
24.	Bamboo	<i>Dendrocalamus strictus</i>	20 – 40	500 – 2500
25.	Muthuga	<i>Butea monosperma</i>	20 - 40	400 - 1500
26.	Hippe	<i>Madhuca latifolia</i>	20 - 40	500 - 2000
27.	Sandal	<i>Santalum album</i>	20 - 50	400 - 1000
28.	Nelli	<i>Emblica officinalis</i>	20 - 40	500 - 2000
29.	Nerale	<i>Sizyzium cumini</i>	20 - 40	500 - 2000
30.	Dhaman	<i>Grevia tilifolia</i>	20 - 40	500 - 2000
31.	Kaval	<i>Careya arborea</i>	20 - 40	500 - 2000
32.	Harada	<i>Terminalia chebula</i>	20 - 40	500 - 2000

References

1. FAO (1976) Framework for Land Evaluation, Food and Agriculture Organization, Rome. 72 pp.
2. FAO (1983) Guidelines for Land Evaluation for Rainfed Agriculture, FAO, Rome, 237 pp.
3. IARI (1971) Soil Survey Manual, All India Soil and Land Use Survey Organization, IARI, New Delhi, 121 pp.
4. Katyal, J.C. and Rattan, R.K. (2003) Secondary and Micronutrients; Research Gap and future needs. Fert. News 48 (4); 9-20.
5. Naidu, L.G.K., Ramamurthy, V., Challa, O., Hegde, R. and Krishnan, P. (2006) Manual Soil Site Suitability Criteria for Major Crops, NBSS Publ. No. 129, NBSS &LUP, Nagpur, 118 pp.
6. Natarajan, A. and Dipak Sarkar (2010) Field Guide for Soil Survey, National Bureau of Soil Survey and Land Use Planning (ICAR), Nagpur, India.
7. Natarajan, A., Rajendra Hegde, Raj, J.N. and Shivananda Murthy, H.G. (2015) Implementation Manual for Sujala-III Project, Watershed Development Department, Bengaluru, Karnataka.
8. Sarma, V.A.K., Krishnan, P. and Budihal, S.L. (1987) Laboratory Manual, Tech. Bull. 23, NBSS &LUP, Nagpur.
9. Sehgal, J.L. (1990) Soil Resource Mapping of Different States of India; Why and How?, National Bureau of Soil Survey and Land Use Planning, Nagpur, 49 pp.
10. Shivaprasad, C.R., R.S. Reddy, J. Sehgal and M. Velayuthum (1998) Soils of Karnataka for Optimizing Land Use, NBSS Publ. No. 47b, NBSS & LUP, Nagpur, India.
11. Soil Survey Staff (2006) Keys to Soil Taxonomy, Tenth edition, U.S. Department of Agriculture/ NRCS, Washington DC, U.S.A.
12. Soil Survey Staff (2012) Soil Survey Manual, Handbook No. 18, USDA, Washington DC, USA.

Appendix I
Bachanahalli -1_2E1e appendix
Soil Phase Information

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	WELLS	Land Capability	Conservation Plan
Bachanahalli	1	2.26	ARKhB2g1	LMU-1	Very deep (>150 cm)	Sandy clay loam	Gravelly (15-35%)	very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Marigold+Banana (Mg+Ba)	Not Available	Iles	Trench cum bunding
Bachanahalli	3	2.19	ARKhB2g1	LMU-1	Very deep (>150 cm)	Sandy clay loam	Gravelly (15-35%)	very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	Iles	Trench cum bunding
Bachanahalli	5	1.73	ARKhB2g1	LMU-1	Very deep (>150 cm)	Sandy clay loam	Gravelly (15-35%)	very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	Iles	Trench cum bunding
Bachanahalli	6	3.22	BMBmB1	LMU-2	Very deep (>150 cm)	Clay	Non gravelly (<15%)	very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Banana+Sorghum (Ba+Sg)	Not Available	IIIw	Graded bunding
Bachanahalli	7	1.71	BMBmB1	LMU-2	Very deep (>150 cm)	Clay	Non gravelly (<15%)	very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Marigold+Turmeric (Mg+Tu)	Not Available	IIIw	Graded bunding
Bachanahalli	8	2.29	ARKhB2g1	LMU-1	Very deep (>150 cm)	Sandy clay loam	Gravelly (15-35%)	very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Maize+Jowar (Mz+Jw)	Not Available	Iles	Trench cum bunding
Bachanahalli	9	1.3	ARKhB2g1	LMU-1	Very deep (>150 cm)	Sandy clay loam	Gravelly (15-35%)	very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	Iles	Trench cum bunding
Bachanahalli	10	0.7	BMBmB1	LMU-2	Very deep (>150 cm)	Clay	Non gravelly (<15%)	very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	IIIw	Graded bunding
Bachanahalli	11	1.85	ARKhB2g1	LMU-1	Very deep (>150 cm)	Sandy clay loam	Gravelly (15-35%)	very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Marigold (Mg)	Not Available	Iles	Trench cum bunding
Bachanahalli	12	2.47	BMBmB1	LMU-2	Very deep (>150 cm)	Clay	Non gravelly (<15%)	very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Marigold+Jowar (Mg+Jw)	Not Available	IIIw	Graded bunding
Bachanahalli	13	2.05	BMBmB1	LMU-2	Very deep (>150 cm)	Clay	Non gravelly (<15%)	very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Marigold+Maize (Mg+Mz)	Not Available	IIIw	Graded bunding
Bachanahalli	14	2.02	BMBmB1	LMU-2	Very deep (>150 cm)	Clay	Non gravelly (<15%)	very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Turmeric+Garlic (Tu+Gc)	Not Available	IIIw	Graded bunding
Bachanahalli	15	1.61	HGHcB2	LMU-1	Very deep (>150 cm)	Sandy loam	Non gravelly (<15%)	very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Marigold (Mg)	Not Available	Ils	Trench cum bunding
Bachanahalli	16	3.69	BMBmB1	LMU-2	Very deep (>150 cm)	Clay	Non gravelly (<15%)	very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Marigold+Maize (Mg+Mz)	1 Borewell	IIIw	Graded bunding
Bachanahalli	23	2.06	HPRIb2g1	LMU-5	Moderately shallow (50-75 cm)	Sandy clay	Gravelly (15-35%)	low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Teak+Current fallow (Te+Cf)	Not Available	Iles	Trench cum bunding
Bachanahalli	24	1.42	HPRIb2g1	LMU-5	Moderately shallow (50-75 cm)	Sandy clay	Gravelly (15-35%)	low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Teak+Current fallow (Te+Cf)	Not Available	Iles	Trench cum bunding
Bachanahalli	28	2.47	ARKhB2g1	LMU-1	Very deep (>150 cm)	Sandy clay loam	Gravelly (15-35%)	very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Marigold+Maize (Mg+Mz)	1 Borewell	Iles	Trench cum bunding
Bachanahalli	29	2.2	BMBmB1	LMU-2	Very deep (>150 cm)	Clay	Non gravelly (<15%)	very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Turmeric+Garlic (Tu+Gc)	1 Borewell	IIIw	Graded bunding
Bachanahalli	37	0.99	BMBmB1	LMU-2	Very deep (>150 cm)	Clay	Non gravelly (<15%)	very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Not Available (NA)	Not Available	IIIw	Graded bunding
Bachanahalli	40	1.41	HGHcB2	LMU-1	Very deep (>150 cm)	Sandy loam	Non gravelly (<15%)	very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Marigold (Mg)	Not Available	Ils	Trench cum bunding
Bachanahalli	41	1.15	HGHcB2	LMU-1	Very deep (>150 cm)	Sandy loam	Non gravelly (<15%)	very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Marigold (Mg)	Not Available	Ils	Trench cum bunding
Bachanahalli	42	1.17	HGHcB2	LMU-1	Very deep (>150 cm)	Sandy loam	Non gravelly (<15%)	very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Marigold (Mg)	Not Available	Ils	Trench cum bunding

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	WELLS	Land Capability	Conservation Plan
Bachanahalli	43	1.27	KLPhB2	LMU-3	Deep (100-150 cm)	Sandy clay loam	Non gravelly (<15%)	low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Turmeric+Teak (Tu+Te)	Not Available	Iles	Trench cum bunding
Bachanahalli	50	1.36	HGHcB2	LMU-1	Very deep (>150 cm)	Sandy loam	Non gravelly (<15%)	very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Marigold+Jowar (Mg+Jw)	Not Available	IIs	Trench cum bunding
Bachanahalli	53	2.13	HPRI2g1	LMU-5	Moderately shallow (50-75 cm)	Sandy clay	Gravelly (15-35%)	low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Teak+Current fallow (Te+Cf)	Not Available	Iles	Trench cum bunding
Berambadi State Forest	1	0.74	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest (F)	Not Available	Forest	Forest
Berambadi District Forest	XX	64.68	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest (F)	Not Available	Forest	Forest
Berambadi	NA	9.87	BMBmB1	LMU-2	Very deep (>150 cm)	Clay	Non gravelly (<15%)	very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Marigold+Maize+Garlic+Turmeric (Mg+Mz+Gc+Tu)	5 Borewell	IIIw	Graded bunding
Berambadi	settlement	5.24	Habitation	Others	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others
Berambadi	TANK	1.07	Waterbody	Others	Others	Others	Others	Others	Others	Others	Waterbody	Not Available	Others	Others
Berambadi	XX	2.38	BMBiB1	LMU-2	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Not Available (NA)	Not Available	IIw	Graded bunding
Berambadi	34	0.01	GPRiB1	LMU-4	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Marigold (Mg)	Not Available	IIs	Trench cum bunding
Berambadi	35	1.66	GPRiB1	LMU-4	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Marigold (Mg)	Not Available	IIs	Trench cum bunding
Berambadi	36	1.91	GPRiB1	LMU-4	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Marigold (Mg)	Not Available	IIs	Trench cum bunding
Berambadi	37	1.7	GPRiB1	LMU-4	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Marigold (Mg)	Not Available	IIs	Trench cum bunding
Berambadi	38	1.87	GPRiB1	LMU-4	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Marigold (Mg)	Not Available	IIs	Trench cum bunding
Berambadi	39	1.31	GPRiB1	LMU-4	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Marigold+Maize (Mg+Mz)	Not Available	IIs	Trench cum bunding
Berambadi	40	3.26	HGHhB1	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Marigold+Jowar (Mg+Jw)	Not Available	IIs	Trench cum bunding
Berambadi	41	2	HGHhB1	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Marigold+Jowar (Mg+Jw)	Not Available	IIs	Trench cum bunding
Berambadi	42	2.05	HGHhB2	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Marigold+Jowar (Mg+Jw)	Not Available	Iles	Trench cum bunding
Berambadi	43	2.22	GPRhB1g1	LMU-4	Moderately deep (75-100 cm)	Sandy clay loam	Gravelly (15-35%)	low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Marigold+Maize (Mg+Mz)	Not Available	IIs	Trench cum bunding
Berambadi	48	71.63	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Marigold+Jowar (Mg+Jw)	Not Available	Forest	Forest
Berambadi	49	1.77	HGHhB1	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Turmeric+Ginger (Tu+Gr)	Not Available	IIs	Trench cum bunding
Berambadi	50	1.37	HGHhB1	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Ginger (Gr)	Not Available	IIs	Trench cum bunding
Berambadi	52	1.73	BMDhB1	LMU-6	Shallow (25-50 cm)	Sandy clay loam	Non gravelly (<15%)	very low (<50 mm/m)	Very gently sloping (1-3%)	Slight	Marigold+Maize (Mg+Mz)	Not Available	IIIs	Trench cum bunding
Berambadi	53	1.33	BMBiB1	LMU-2	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Marigold (Mg)	2 Borewell	IIw	Graded bunding
Berambadi	54	1.04	BMBiB1	LMU-2	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Marigold+Maize (Mg+Mz)	Not Available	IIw	Graded bunding

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	WELLS	Land Capability	Conservation Plan
Berambadi	55	2.33	BMBiB1	LMU-2	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Marigold+Maize (Mg+Mz)	Not Available	IIw	Graded bunding
Berambadi	56	2.73	BMDhB1	LMU-6	Shallow (25-50 cm)	Sandy clay loam	Non gravelly (<15%)	very low (<50 mm/m)	Very gently sloping (1-3%)	Slight	Marigold+Ginger (Mg+Gr)	1 Borewell	IIIs	Trench cum bunding
Berambadi	57	2.04	GPRIb1	LMU-4	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Marigold (Mg)	Not Available	IIs	Trench cum bunding
Berambadi	58	0.56	GPRIb1	LMU-4	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Not Available (NA)	Not Available	IIs	Trench cum bunding
Berambadi	59_TA NK	4.29	Habitation	Others	Others	Others	Others	Others	Others	Others	Waterbody	Not Available	Others	Others
Berambadi	60	0.4	BMBmB1	LMU-2	Very deep (>150 cm)	Clay	Non gravelly (<15%)	very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Not Available (NA)	Not Available	IIIw	Graded bunding
Berambadi	61	2.45	BMBmB1	LMU-2	Very deep (>150 cm)	Clay	Non gravelly (<15%)	very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Not Available (NA)	Not Available	IIIw	Graded bunding
Berambadi	62	1.7	ARKhB1	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Marigold+Jowar (Mg+Jw)	Not Available	IIs	Trench cum bunding
Berambadi	63	1.01	ARKhB1	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Marigold+Jowar (Mg+Jw)	Not Available	IIs	Trench cum bunding
Berambadi	64	2.16	ARKhB1	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Marigold+Maize (Mg+Mz)	Not Available	IIs	Trench cum bunding
Berambadi	65	2.3	ARKhB1	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Marigold+Current fallow (Mg+Cf)	Not Available	IIs	Trench cum bunding
Berambadi	66	1.82	ARKhB1	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Not Available (NA)	Not Available	IIs	Trench cum bunding
Berambadi	67	0.63	ARKhB1	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Marigold (Mg)	Not Available	IIs	Trench cum bunding
Berambadi	68	2	ARKhB1	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Marigold (Mg)	Not Available	IIs	Trench cum bunding
Berambadi	69	2.01	ARKhB1	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Marigold+Banana (Mg+Ba)	Not Available	IIs	Trench cum bunding
Berambadi	70	2.32	ARKhB1	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Marigold+Jowar (Mg+Jw)	Not Available	IIs	Trench cum bunding
Berambadi	72	1.56	ARKhB1	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Marigold (Mg)	Not Available	IIs	Trench cum bunding
Berambadi	73	1.63	BMBmB1	LMU-2	Very deep (>150 cm)	Clay	Non gravelly (<15%)	very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Marigold+Ragi (Mg+Ra)	Not Available	IIIw	Graded bunding
Berambadi	75	2.49	HGHhB1g1	LMU-1	Very deep (>150 cm)	Sandy clay loam	Gravelly (15-35%)	very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Marigold (Mg)	Not Available	IIs	Trench cum bunding
Berambadi	76	1.04	GPRhB1g1	LMU-4	Moderately deep (75-100 cm)	Sandy clay loam	Gravelly (15-35%)	low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Marigold (Mg)	Not Available	IIs	Trench cum bunding
Berambadi	77	0.45	HGHhB1g1	LMU-1	Very deep (>150 cm)	Sandy clay loam	Gravelly (15-35%)	very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Marigold (Mg)	Not Available	IIs	Trench cum bunding
Berambadi	78	0.74	HGHhB1g1	LMU-1	Very deep (>150 cm)	Sandy clay loam	Gravelly (15-35%)	very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Marigold (Mg)	Not Available	IIs	Trench cum bunding
Berambadi	79	2.39	HGHhB1g1	LMU-1	Very deep (>150 cm)	Sandy clay loam	Gravelly (15-35%)	very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Marigold+Jowar (Mg+Jw)	Not Available	IIs	Trench cum bunding
Berambadi	86	2.48	HGHhB1g1	LMU-1	Very deep (>150 cm)	Sandy clay loam	Gravelly (15-35%)	very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Marigold+Jowar (Mg+Jw)	Not Available	IIs	Trench cum bunding

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	WELLS	Land Capability	Conservation Plan
Berambadi	87	1.42	HGHhB1g1	LMU-1	Very deep (>150 cm)	Sandy clay loam	Gravelly (15-35%)	very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Marigold+Jowar (Mg+Jw)	Not Available	IIs	Trench cum bunding
Berambadi	88	1.61	GPRhB1g1	LMU-4	Moderately deep (75-100 cm)	Sandy clay loam	Gravelly (15-35%)	low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Marigold (Mg)	1 Borewell	IIs	Trench cum bunding
Berambadi	89	2.39	GPRhB1g1	LMU-4	Moderately deep (75-100 cm)	Sandy clay loam	Gravelly (15-35%)	low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Marigold+Jowar (Mg+Jw)	Not Available	IIs	Trench cum bunding
Berambadi	90	2.08	HDRhB1g1	LMU-7	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	very low (<50 mm/m)	Very gently sloping (1-3%)	Slight	Marigold+Jowar (Mg+Jw)	Not Available	IIIs	Trench cum bunding
Berambadi	91	2.68	HDRhB1g1	LMU-7	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	very low (<50 mm/m)	Very gently sloping (1-3%)	Slight	Marigold+Jowar (Mg+Jw)	Not Available	IIIs	Trench cum bunding
Berambadi	118	2.11	Habitation	Others	Others	Others	Others	Others	Others	Others	Not Available (NA)	Not Available	Others	Others
Berambadi	122	102.79	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Marigold+Forest (Mg+F)	1 Borewell	Forest	Forest
Berambadi	123	78.12	BMBhB2	LMU-2	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Marigold+Jowar+Banana+Turmeric (Mg+Jw+Ba+Tu)	Not Available	IIIw	Graded bunding
Berambadi	124	1.49	GPRcB1	LMU-4	Moderately deep (75-100 cm)	Sandy loam	Non gravelly (<15%)	very low (<50 mm/m)	Very gently sloping (1-3%)	Slight	Marigold+Maize (Mg+Mz)	Not Available	IIs	Trench cum bunding
Berambadi	125	2.83	GPRcB1	LMU-4	Moderately deep (75-100 cm)	Sandy loam	Non gravelly (<15%)	very low (<50 mm/m)	Very gently sloping (1-3%)	Slight	Marigold+Jowar (Mg+Jw)	Not Available	IIs	Trench cum bunding
Berambadi	126	1.61	GPRcB1	LMU-4	Moderately deep (75-100 cm)	Sandy loam	Non gravelly (<15%)	very low (<50 mm/m)	Very gently sloping (1-3%)	Slight	Marigold (Mg)	Not Available	IIs	Trench cum bunding
Berambadi	127	2.12	GPRcB1	LMU-4	Moderately deep (75-100 cm)	Sandy loam	Non gravelly (<15%)	very low (<50 mm/m)	Very gently sloping (1-3%)	Slight	Marigold (Mg)	Not Available	IIs	Trench cum bunding
Berambadi	128	1.33	BMDhB1g1	LMU-6	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	very low (<50 mm/m)	Very gently sloping (1-3%)	Slight	Marigold (Mg)	Not Available	IIIs	Trench cum bunding
Berambadi	129	3.25	BMDhB1g1	LMU-6	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	very low (<50 mm/m)	Very gently sloping (1-3%)	Slight	Marigold+Jowar (Mg+Jw)	Not Available	IIIs	Trench cum bunding
Berambadi	130	1.26	GPRcB1	LMU-4	Moderately deep (75-100 cm)	Sandy loam	Non gravelly (<15%)	very low (<50 mm/m)	Very gently sloping (1-3%)	Slight	Turmeric+Garlic (Tu+Gc)	1 Borewell	IIs	Trench cum bunding
Berambadi	131	2.28	BMDhB1g1	LMU-6	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	very low (<50 mm/m)	Very gently sloping (1-3%)	Slight	Marigold (Mg)	Not Available	IIIs	Trench cum bunding
Berambadi	132	2.83	BMDhB1g1	LMU-6	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	very low (<50 mm/m)	Very gently sloping (1-3%)	Slight	Marigold+Jowar (Mg+Jw)	Not Available	IIIs	Trench cum bunding
Berambadi	315	1.98	BMBmB1	LMU-2	Very deep (>150 cm)	Clay	Non gravelly (<15%)	very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Banana (Ba)	Not Available	IIIw	Graded bunding
Berambadi	316	1.8	GPRcB1	LMU-4	Moderately deep (75-100 cm)	Sandy loam	Non gravelly (<15%)	very low (<50 mm/m)	Very gently sloping (1-3%)	Slight	Marigold (Mg)	Not Available	IIs	Trench cum bunding
Berambadi	317	2.72	GPRcB1	LMU-4	Moderately deep (75-100 cm)	Sandy loam	Non gravelly (<15%)	very low (<50 mm/m)	Very gently sloping (1-3%)	Slight	Marigold (Mg)	Not Available	IIs	Trench cum bunding
Berambadi	318	1.62	BMDhB1g1	LMU-6	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	very low (<50 mm/m)	Very gently sloping (1-3%)	Slight	Marigold (Mg)	Not Available	IIIs	Trench cum bunding
Berambadi	319	1.82	BMDhB1g1	LMU-6	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	very low (<50 mm/m)	Very gently sloping (1-3%)	Slight	Marigold (Mg)	Not Available	IIIs	Trench cum bunding
Berambadi	320	1.39	GPRhB1g1	LMU-4	Moderately deep (75-100 cm)	Sandy clay loam	Gravelly (15-35%)	low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Marigold+Banana (Mg+Ba)	Not Available	IIs	Trench cum bunding
Berambadi	321	2.7	GPRhB1g1	LMU-4	Moderately deep (75-100 cm)	Sandy clay loam	Gravelly (15-35%)	low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Marigold+Jowar (Mg+Jw)	Not Available	IIs	Trench cum bunding

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	WELLS	Land Capability	Conservation Plan
Berambadi	322	1.55	GPRhB1g1	LMU-4	Moderately deep (75-100 cm)	Sandy clay loam	Gravelly (15-35%)	low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Marigold+Jowar (Mg+Jw)	Not Available	IIs	Trench cum bunding
Berambadi	323	1.99	GPRhB1g1	LMU-4	Moderately deep (75-100 cm)	Sandy clay loam	Gravelly (15-35%)	low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Turmeric+Garlic (Tu+Gc)	Not Available	IIs	Trench cum bunding
Berambadi	324	2.19	BMBmB1	LMU-2	Very deep (>150 cm)	Clay	Non gravelly (<15%)	very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Marigold+Maize (Mg+Mz)	2 Borewell	IIIw	Graded bunding
Berambadi	325	2.07	BMBmB1	LMU-2	Very deep (>150 cm)	Clay	Non gravelly (<15%)	very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Maize+Turmeric (Mz+Tu)	Not Available	IIIw	Graded bunding
Berambadi	326	2.09	BMBmB1	LMU-2	Very deep (>150 cm)	Clay	Non gravelly (<15%)	very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Banana+Turmeric (Ba+Tu)	1 Borewell	IIIw	Graded bunding
Berambadi	327	2.31	BMBmB1	LMU-2	Very deep (>150 cm)	Clay	Non gravelly (<15%)	very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Banana+Garlic (Ba+Gc)	Not Available	IIIw	Graded bunding
Berambadi	328	3	BMBmB1	LMU-2	Very deep (>150 cm)	Clay	Non gravelly (<15%)	very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Marigold+Maize (Mg+Mz)	Not Available	IIIw	Graded bunding
Berambadi	340	3.13	BMBmB1	LMU-2	Very deep (>150 cm)	Clay	Non gravelly (<15%)	very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Not Available (NA)	Not Available	IIIw	Graded bunding
Berambadi	341	3.23	BMBmB1	LMU-2	Very deep (>150 cm)	Clay	Non gravelly (<15%)	very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Marigold+Maize (Mg+Mz)	Not Available	IIIw	Graded bunding
Berambadi	342	2.25	GPRcB1	LMU-4	Moderately deep (75-100 cm)	Sandy loam	Non gravelly (<15%)	very low (<50 mm/m)	Very gently sloping (1-3%)	Slight	Marigold (Mg)	Not Available	IIs	Trench cum bunding
Berambadi	343	2.51	GPRcC2g1	LMU-4	Moderately deep (75-100 cm)	Sandy loam	Gravelly (15-35%)	low (51-100 mm/m)	Gently sloping (3-5%)	Moderate	Marigold (Mg)	Not Available	IIs	Trench cum bunding
Berambadi	344	2.14	GPRcB1	LMU-4	Moderately deep (75-100 cm)	Sandy loam	Non gravelly (<15%)	very low (<50 mm/m)	Very gently sloping (1-3%)	Slight	Marigold+Maize (Mg+Mz)	Not Available	IIs	Trench cum bunding
Berambadi	346	2.07	BMBmB1	LMU-2	Very deep (>150 cm)	Clay	Non gravelly (<15%)	very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Marigold+Jowar (Mg+Jw)	Not Available	IIIw	Graded bunding
Berambadi	347	1.98	BMDhB1g1	LMU-6	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	very low (<50 mm/m)	Very gently sloping (1-3%)	Slight	Marigold+Jowar (Mg+Jw)	Not Available	IIIs	Trench cum bunding
Berambadi	350	2.72	BMBmB1	LMU-2	Very deep (>150 cm)	Clay	Non gravelly (<15%)	very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Marigold+Maize (Mg+Mz)	Not Available	IIIw	Graded bunding
Berambadi	355	2.26	HGHhB1	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Turmeric+Garlic (Tu+Gc)	1 Borewell	IIs	Trench cum bunding
Berambadi	356	1.37	GPRhB1g1	LMU-4	Moderately deep (75-100 cm)	Sandy clay loam	Gravelly (15-35%)	low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Marigold+Maize (Mg+Mz)	Not Available	IIs	Trench cum bunding
Berambadi	357	0.99	GPRhB1g1	LMU-4	Moderately deep (75-100 cm)	Sandy clay loam	Gravelly (15-35%)	low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Marigold (Mg)	Not Available	IIs	Trench cum bunding
Berambadi	358	0.8	GPRhB1g1	LMU-4	Moderately deep (75-100 cm)	Sandy clay loam	Gravelly (15-35%)	low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Marigold (Mg)	Not Available	IIs	Trench cum bunding
Berambadi	359	1.13	GPRhB1g1	LMU-4	Moderately deep (75-100 cm)	Sandy clay loam	Gravelly (15-35%)	low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Marigold (Mg)	Not Available	IIs	Trench cum bunding
Berambadi	362	0.93	GPRhB1g1	LMU-4	Moderately deep (75-100 cm)	Sandy clay loam	Gravelly (15-35%)	low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Marigold (Mg)	Not Available	IIs	Trench cum bunding
Berambadi	363	0.62	GPRhB1g1	LMU-4	Moderately deep (75-100 cm)	Sandy clay loam	Gravelly (15-35%)	low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Marigold (Mg)	Not Available	IIs	Trench cum bunding
Berambadi	364	1.25	GPRhB1g1	LMU-4	Moderately deep (75-100 cm)	Sandy clay loam	Gravelly (15-35%)	low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Marigold (Mg)	Not Available	IIs	Trench cum bunding
Berambadi	366	3.03	BMBmB1	LMU-2	Very deep (>150 cm)	Clay	Non gravelly (<15%)	very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Marigold+Maize (Mg+Mz)	Not Available	IIIw	Graded bunding

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	WELLS	Land Capability	Conservation Plan
Berambadi	367	1.43	BMBhB2	LMU-2	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Marigold+Jowar (Mg+Jw)	Not Available	IIIw	Graded bunding
Berambadi	368	1.67	BMDhB1g1	LMU-6	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	very low (<50 mm/m)	Very gently sloping (1-3%)	Slight	Marigold (Mg)	Not Available	IIIs	Trench cum bunding
Berambadi	370	1.84	BMBmB1	LMU-2	Very deep (>150 cm)	Clay	Non gravelly (<15%)	very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Marigold (Mg)	Not Available	IIIw	Graded bunding
Maddur	33	0.69	BMBmB1	LMU-2	Very deep (>150 cm)	Clay	Non gravelly (<15%)	very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Not Available (NA)	Not Available	IIIw	Graded bunding
Maddur	56	2.34	BMBmB1	LMU-2	Very deep (>150 cm)	Clay	Non gravelly (<15%)	very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Marigold+Maize (Mg+Mz)	Not Available	IIIw	Graded bunding
Maddur	61	1.6	HPRcB1	LMU-5	Moderately shallow (50-75 cm)	Sandy loam	Non gravelly (<15%)	low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Marigold+Maize (Mg+Mz)	Not Available	IIs	Trench cum bunding
Maddur	62	0.98	BMBmB1	LMU-2	Very deep (>150 cm)	Clay	Non gravelly (<15%)	very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	IIIw	Graded bunding
Maddur	63	1.66	BMBmB1	LMU-2	Very deep (>150 cm)	Clay	Non gravelly (<15%)	very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Marigold (Mg)	Not Available	IIIw	Graded bunding
Maddur	64	0.92	BMBmB1	LMU-2	Very deep (>150 cm)	Clay	Non gravelly (<15%)	very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Marigold (Mg)	Not Available	IIIw	Graded bunding
Maddur	66	0.32	BMBmB1	LMU-2	Very deep (>150 cm)	Clay	Non gravelly (<15%)	very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Not Available (NA)	Not Available	IIIw	Graded bunding
Maddur	69	1.1	HPRcB1	LMU-5	Moderately shallow (50-75 cm)	Sandy loam	Non gravelly (<15%)	low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Banana (Ba)	Not Available	IIs	Trench cum bunding
Maddur	70	1.32	BMBmB1	LMU-2	Very deep (>150 cm)	Clay	Non gravelly (<15%)	very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Marigold (Mg)	Not Available	IIIw	Graded bunding

Appendix II

Bachanahalli -1_2E1e appendix

Soil Fertility Information

[illegible]

Village	Survey No	Soil Reaction	Salinity	Organic Carbon	Available Phosphorus	Available Potassium	Available Sulphur	Available Boron	Available Iron	Available Manganese	Available Copper	Available Zinc
Berambadi	367	Neutral (pH 6.5-7.3)	Non saline (<2 dsm)	Low (<0.5 %)	Medium (23-57 kg/ha)	High (>337 kg/ha)	Low (<10 ppm)	Low (<0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Berambadi	368	Neutral (pH 6.5-7.3)	Non saline (<2 dsm)	Medium (0.5-0.75 %)	Medium (23-57 kg/ha)	High (>337 kg/ha)	Low (<10 ppm)	Low (<0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Berambadi	370	Neutral (pH 6.5-7.3)	Non saline (<2 dsm)	Low (<0.5 %)	Medium (23-57 kg/ha)	High (>337 kg/ha)	Low (<10 ppm)	Low (<0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Maddur	33	Moderately alkaline (pH 7.8-8.4)	Non saline (<2 dsm)	High (>0.75 %)	Medium (23-57 kg/ha)	High (>337 kg/ha)	Low (<10 ppm)	Low (<0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Maddur	56	Moderately alkaline (pH 7.8-8.4)	Non saline (<2 dsm)	High (>0.75 %)	Medium (23-57 kg/ha)	High (>337 kg/ha)	Low (<10 ppm)	Low (<0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Maddur	61	Slightly alkaline (pH 7.3-7.8)	Non saline (<2 dsm)	Medium (0.5-0.75 %)	Medium (23-57 kg/ha)	High (>337 kg/ha)	Low (<10 ppm)	Low (<0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Maddur	62	Moderately alkaline (pH 7.8-8.4)	Non saline (<2 dsm)	Medium (0.5-0.75 %)	Medium (23-57 kg/ha)	High (>337 kg/ha)	Low (<10 ppm)	Low (<0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Maddur	63	Slightly alkaline (pH 7.3-7.8)	Non saline (<2 dsm)	Medium (0.5-0.75 %)	Medium (23-57 kg/ha)	High (>337 kg/ha)	Low (<10 ppm)	Low (<0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Maddur	64	Slightly alkaline (pH 7.3-7.8)	Non saline (<2 dsm)	Medium (0.5-0.75 %)	Medium (23-57 kg/ha)	High (>337 kg/ha)	Low (<10 ppm)	Low (<0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Maddur	66	Moderately alkaline (pH 7.8-8.4)	Non saline (<2 dsm)	High (>0.75 %)	Medium (23-57 kg/ha)	Medium (145-337 kg/ha)	Low (<10 ppm)	Low (<0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Maddur	69	Slightly alkaline (pH 7.3-7.8)	Non saline (<2 dsm)	Medium (0.5-0.75 %)	Medium (23-57 kg/ha)	High (>337 kg/ha)	Low (<10 ppm)	Low (<0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Maddur	70	Slightly alkaline (pH 7.3-7.8)	Non saline (<2 dsm)	Medium (0.5-0.75 %)	Medium (23-57 kg/ha)	High (>337 kg/ha)	Low (<10 ppm)	Low (<0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)

Appendix III

Bachanahalli -1_2E1e Appendix

Soil Suitability Information

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Marigold	Chrysanthemum	Banana	Horse gram	Field-bean	Turmeric	Beetroot	Potato	Beans	
Bachanahalli	1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	
Bachanahalli	3	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	
Bachanahalli	5	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	
Bachanahalli	6	N	S3wt	S3wt	S1	N	S1	S1	S1	S2w	S2w	S1	N	S1	N	S1	S1	S3wt	S2wt	S2wt	S2wt	S2wt	S2wt	S2w	S2w	S3wt	S3wt	S3wt	S2wt
Bachanahalli	7	N	S3wt	S3wt	S1	N	S1	S1	S1	S2w	S2w	S1	N	S1	N	S1	S1	S3wt	S2wt	S2wt	S2wt	S2wt	S2wt	S2w	S2w	S3wt	S3wt	S3wt	S2wt
Bachanahalli	8	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	
Bachanahalli	9	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	
Bachanahalli	10	N	S3wt	S3wt	S1	N	S1	S1	S1	S2w	S2w	S1	N	S1	N	S1	S1	S3wt	S2wt	S2wt	S2wt	S2wt	S2wt	S2w	S2w	S3wt	S3wt	S3wt	S2wt
Bachanahalli	11	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	
Bachanahalli	12	N	S3wt	S3wt	S1	N	S1	S1	S1	S2w	S2w	S1	N	S1	N	S1	S1	S3wt	S2wt	S2wt	S2wt	S2wt	S2wt	S2w	S2w	S3wt	S3wt	S3wt	S2wt
Bachanahalli	13	N	S3wt	S3wt	S1	N	S1	S1	S1	S2w	S2w	S1	N	S1	N	S1	S1	S3wt	S2wt	S2wt	S2wt	S2wt	S2wt	S2w	S2w	S3wt	S3wt	S3wt	S2wt
Bachanahalli	14	N	S3wt	S3wt	S1	N	S1	S1	S1	S2w	S2w	S1	N	S1	N	S1	S1	S3wt	S2wt	S2wt	S2wt	S2wt	S2wt	S2w	S2w	S3wt	S3wt	S3wt	S2wt
Bachanahalli	15	S1	S1	S1	S2t	S1	S3t	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	
Bachanahalli	16	N	S3wt	S3wt	S1	N	S1	S1	S1	S2w	S2w	S1	N	S1	N	S1	S1	S3wt	S2wt	S2wt	S2wt	S2wt	S2wt	S2w	S2w	S3wt	S3wt	S3wt	S2wt
Bachanahalli	23	N	S2r	S3r	S2r	S3r	S2r	S3r	S3r	S3r	S3r	S2r	S3r	S2r	S3r	N	S3r	S2rt	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S2r	S2r	S2r
Bachanahalli	24	N	S2r	S3r	S2r	S3r	S2r	S3r	S3r	S3r	S3r	S2r	S3r	S2r	S3r	N	S3r	S2rt	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S2r	S2r	S2r
Bachanahalli	28	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	
Bachanahalli	29	N	S3wt	S3wt	S1	N	S1	S1	S1	S2w	S2w	S1	N	S1	N	S1	S1	S3wt	S2wt	S2wt	S2wt	S2wt	S2wt	S2w	S2w	S3wt	S3wt	S3wt	S2wt
Bachanahalli	37	N	S3wt	S3wt	S1	N	S1	S1	S1	S2w	S2w	S1	N	S1	N	S1	S1	S3wt	S2wt	S2wt	S2wt	S2wt	S2wt	S2w	S2w	S3wt	S3wt	S3wt	S2wt
Bachanahalli	40	S1	S1	S1	S2t	S1	S3t	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	
Bachanahalli	41	S1	S1	S1	S2t	S1	S3t	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	
Bachanahalli	42	S1	S1	S1	S2t	S1	S3t	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	
Bachanahalli	43	S2r	S1	S3r	S1	S1	S2t	S2r	S2r	S2gt	S1	S1	S2r	S1	S2r	S2r	S2r	S2t	S2g	S1	S2g	S2g	S1	S2g	S2g	S2g	S2g	S2g	
Bachanahalli	50	S1	S1	S1	S2t	S1	S3t	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	
Bachanahalli	53	N	S2r	S3r	S2r	S3r	S2r	S3r	S3r	S3r	S3r	S2r	S3r	S2r	S3r	N	S3r	S2rt	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S2r	S2r	

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Marigold	Chrysanthemum	Banana	Horse gram	Field-bean	Turmeric	Beetroot	Potato	Beans
Berambadi State Forest	1	Forest	forest	forest	forest	forest	forest	forest	forest	forest	forest	forest	forest	forest	forest	forest	forest	forest	forest	forest	forest	forest	forest	forest	forest	forest	forest	forest
Berambadi District Forest	XX	Forest	forest	forest	forest	forest	forest	forest	forest	forest	forest	forest	forest	forest	forest	forest	forest	forest	forest	forest	forest	forest	forest	forest	forest	forest	forest	forest
Berambadi	NA	N	S3wt	S3wt	S1	N	S1	S1	S1	S2w	S2w	S1	N	S1	N	S1	S1	S3wt	S2wt	S2wt	S2wt	S2wt	S2w	S2w	S3wt	S3wt	S3wt	S2wt
Berambadi	Settlement	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Berambadi	Tank	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Berambadi	XX	N	S3wt	S3wt	S1	N	S1	S1	S1	S2w	S2w	S1	N	S1	N	S1	S1	S3wt	S2wt	S2wt	S2wt	S2wt	S2w	S2w	S3wt	S3wt	S3wt	S2wt
Berambadi	34	S3rt	S3r	S2r	S3r	S2r	S3r	S3r	S3r	Nr	Nr	S2r	S3r	S2r	S2r	S3r	S3r	S3r	S3r	S3r	S3r	Nr	S3r	S3r	S3r	S3r	S3r	S3r
Berambadi	35	S3rt	S3r	S2r	S3r	S2r	S3r	S3r	S3r	Nr	Nr	S2r	S3r	S2r	S2r	S3r	S3r	S3r	S3r	S3r	S3r	Nr	S3r	S3r	S3r	S3r	S3r	S3r
Berambadi	36	S3rt	S3r	S2r	S3r	S2r	S3r	S3r	S3r	Nr	Nr	S2r	S3r	S2r	S2r	S3r	S3r	S3r	S3r	S3r	S3r	Nr	S3r	S3r	S3r	S3r	S3r	S3r
Berambadi	37	S3rt	S3r	S2r	S3r	S2r	S3r	S3r	S3r	Nr	Nr	S2r	S3r	S2r	S2r	S3r	S3r	S3r	S3r	S3r	S3r	Nr	S3r	S3r	S3r	S3r	S3r	S3r
Berambadi	38	S3rt	S3r	S2r	S3r	S2r	S3r	S3r	S3r	Nr	Nr	S2r	S3r	S2r	S2r	S3r	S3r	S3r	S3r	S3r	S3r	Nr	S3r	S3r	S3r	S3r	S3r	S3r
Berambadi	39	S3rt	S3r	S2r	S3r	S2r	S3r	S3r	S3r	Nr	Nr	S2r	S3r	S2r	S2r	S3r	S3r	S3r	S3r	S3r	S3r	Nr	S3r	S3r	S3r	S3r	S3r	S3r
Berambadi	40	S1	S1	S1	S2t	S1	S3t	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1
Berambadi	41	S1	S1	S1	S2t	S1	S3t	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1
Berambadi	42	S1	S1	S1	S2t	S1	S3t	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1
Berambadi	43	S3rt	S3r	S2r	S3r	S2r	S3r	S3r	S3r	Nr	Nr	S2r	S3r	S2r	S2r	S3r	S3r	S3r	S3r	S3r	S3r	Nr	S3r	S3r	S3r	S3r	S3r	S3r
Berambadi	48	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest
Berambadi	49	S1	S1	S1	S2t	S1	S3t	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1
Berambadi	50	S1	S1	S1	S2t	S1	S3t	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1
Berambadi	51	N	S3rt	N	S3r	N	S3r	N	N	Nrt	Nrt	S3r	N	S3r	N	N	N	S3rt	S3rt	S3rt	S3rt	Nrt	S3r	S3r	S3rt	S3rt	S3rt	S3rt
Berambadi	52	N	S3rt	N	S3r	N	S3r	N	N	Nrt	Nrt	S3r	N	S3r	N	N	N	S3rt	S3rt	S3rt	S3rt	Nrt	S3r	S3r	S3rt	S3rt	S3rt	S3rt
Berambadi	53	N	S3wt	S3wt	S1	N	S1	S1	S1	S2w	S2w	S1	N	S1	N	S1	S1	S3wt	S2wt	S2wt	S2wt	S2wt	S2w	S2w	S3wt	S3wt	S3wt	S2wt
Berambadi	54	N	S3wt	S3wt	S1	N	S1	S1	S1	S2w	S2w	S1	N	S1	N	S1	S1	S3wt	S2wt	S2wt	S2wt	S2wt	S2w	S2w	S3wt	S3wt	S3wt	S2wt
Berambadi	55	N	S3wt	S3wt	S1	N	S1	S1	S1	S2w	S2w	S1	N	S1	N	S1	S1	S3wt	S2wt	S2wt	S2wt	S2wt	S2w	S2w	S3wt	S3wt	S3wt	S2wt
Berambadi	56	N	S3rt	N	S3r	N	S3r	N	N	Nrt	Nrt	S3r	N	S3r	N	N	N	S3rt	S3rt	S3rt	S3rt	Nrt	S3r	S3r	S3rt	S3rt	S3rt	S3rt
Berambadi	57	S3rt	S3r	S2r	S3r	S2r	S3r	S3r	S3r	Nr	Nr	S2r	S3r	S2r	S2r	S3r	S3r	S3r	S3r	S3r	S3r	Nr	S3r	S3r	S3r	S3r	S3r	S3r
Berambadi	58	S3rt	S3r	S2r	S3r	S2r	S3r	S3r	S3r	Nr	Nr	S2r	S3r	S2r	S2r	S3r	S3r	S3r	S3r	S3r	S3r	Nr	S3r	S3r	S3r	S3r	S3r	S3r

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Marigold	Chrysanthemum	Banana	Horse gram	Field-bean	Turmeric	Beetroot	Potato	Beans
Berambadi	59_TAN K	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Berambadi	60	N	S3wt	S3wt	S1	N	S1	S1	S1	S2w	S2w	S1	N	S1	N	S1	S1	S3wt	S2wt	S2wt	S2wt	S2wt	S2w	S2w	S3wt	S3wt	S3wt	S2wt
Berambadi	61	N	S3wt	S3wt	S1	N	S1	S1	S1	S2w	S2w	S1	N	S1	N	S1	S1	S3wt	S2wt	S2wt	S2wt	S2wt	S2w	S2w	S3wt	S3wt	S3wt	S2wt
Berambadi	62	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1
Berambadi	63	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1
Berambadi	64	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1
Berambadi	65	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1
Berambadi	66	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1
Berambadi	67	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1
Berambadi	68	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1
Berambadi	69	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1
Berambadi	70	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1
Berambadi	72	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1
Berambadi	73	N	S3wt	S3wt	S1	N	S1	S1	S1	S2w	S2w	S1	N	S1	N	S1	S1	S3wt	S2wt	S2wt	S2wt	S2wt	S2w	S2w	S3wt	S3wt	S3wt	S2wt
Berambadi	75	S1	S1	S1	S2t	S1	S3t	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1
Berambadi	76	S3rt	S3r	S2r	S3r	S2r	S3r	S3r	S3r	Nr	Nr	S2r	S3r	S2r	S2r	S3r	S3r	S3r	S3r	S3r	S3r	Nr	S3r	S3r	S3r	S3r	S3r	S3r
Berambadi	77	S1	S1	S1	S2t	S1	S3t	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1
Berambadi	78	S1	S1	S1	S2t	S1	S3t	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1
Berambadi	79	S1	S1	S1	S2t	S1	S3t	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1
Berambadi	86	S1	S1	S1	S2t	S1	S3t	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1
Berambadi	87	S1	S1	S1	S2t	S1	S3t	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1
Berambadi	88	S3rt	S3r	S2r	S3r	S2r	S3r	S3r	S3r	Nr	Nr	S2r	S3r	S2r	S2r	S3r	S3r	S3r	S3r	S3r	S3r	Nr	S3r	S3r	S3r	S3r	S3r	S3r
Berambadi	89	S3rt	S3r	S2r	S3r	S2r	S3r	S3r	S3r	Nr	Nr	S2r	S3r	S2r	S2r	S3r	S3r	S3r	S3r	S3r	S3r	Nr	S3r	S3r	S3r	S3r	S3r	S3r
Berambadi	90	N	S3r	N	S3r	N	S3r	N	N	Nr	Nr	S3r	N	S3r	N	N	N	S3r	S3r	S3r	S3r	Nr	S3r	S3r	S3r	S3r	S3r	S3r
Berambadi	91	N	S3r	N	S3r	N	S3r	N	N	Nr	Nr	S3r	N	S3r	N	N	N	S3r	S3r	S3r	S3r	Nr	S3r	S3r	S3r	S3r	S3r	S3r
Berambadi	118	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Berambadi	122	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Marigold	Chrysanthemum	Banana	Horse gram	Field-bean	Turmeric	Beetroot	Potato	Beans
Berambadi	12s3	N	S3wt	S3wt	S1	N	S1	S1	S1	S2w	S2w	S1	N	S1	N	S1	S1	S3wt	S2wt	S2wt	S2wt	S2wt	S2w	S2w	S3wt	S3wt	S3wt	S2wt
Berambadi	124	S3r	S1	S2r	S1	S2r	S2t	S3r	S3r	S2gt	S1	S2r	S3r	S2r	S2r	S3r	S3r	S1	S2g	S1	S2g	S2g	S1	S2g	S2g	S2g	S2g	S2g
Berambadi	125	S3r	S1	S2r	S1	S2r	S2t	S3r	S3r	S2gt	S1	S2r	S3r	S2r	S2r	S3r	S3r	S1	S2g	S1	S2g	S2g	S1	S2g	S2g	S2g	S2g	S2g
Berambadi	126	S3r	S1	S2r	S1	S2r	S2t	S3r	S3r	S2gt	S1	S2r	S3r	S2r	S2r	S3r	S3r	S1	S2g	S1	S2g	S2g	S1	S2g	S2g	S2g	S2g	S2g
Berambadi	127	S3r	S1	S2r	S1	S2r	S2t	S3r	S3r	S2gt	S1	S2r	S3r	S2r	S2r	S3r	S3r	S1	S2g	S1	S2g	S2g	S1	S2g	S2g	S2g	S2g	S2g
Berambadi	128	N	S3rt	N	S3r	N	S3r	N	N	Nrt	Nrt	S3r	N	S3r	N	N	N	S3rt	S3rt	S3rt	S3rt	Nrt	S3r	S3r	S3rt	S3rt	S3rt	S3rt
Berambadi	129	N	S3rt	N	S3r	N	S3r	N	N	Nrt	Nrt	S3r	N	S3r	N	N	N	S3rt	S3rt	S3rt	S3rt	Nrt	S3r	S3r	S3rt	S3rt	S3rt	S3rt
Berambadi	130	S3r	S1	S2r	S1	S2r	S2t	S3r	S3r	S2gt	S1	S2r	S3r	S2r	S2r	S3r	S3r	S1	S2g	S1	S2g	S2g	S1	S2g	S2g	S2g	S2g	S2g
Berambadi	131	N	S3rt	N	S3r	N	S3r	N	N	Nrt	Nrt	S3r	N	S3r	N	N	N	S3rt	S3rt	S3rt	S3rt	Nrt	S3r	S3r	S3rt	S3rt	S3rt	S3rt
Berambadi	132	N	S3rt	N	S3r	N	S3r	N	N	Nrt	Nrt	S3r	N	S3r	N	N	N	S3rt	S3rt	S3rt	S3rt	Nrt	S3r	S3r	S3rt	S3rt	S3rt	S3rt
Berambadi	315	N	S3wt	S3wt	S1	N	S1	S1	S1	S2w	S2w	S1	N	S1	N	S1	S1	S3wt	S2wt	S2wt	S2wt	S2wt	S2w	S2w	S3wt	S3wt	S3wt	S2wt
Berambadi	316	S3r	S1	S2r	S1	S2r	S2t	S3r	S3r	S2gt	S1	S2r	S3r	S2r	S2r	S3r	S3r	S1	S2g	S1	S2g	S2g	S1	S2g	S2g	S2g	S2g	S2g
Berambadi	317	S3r	S1	S2r	S1	S2r	S2t	S3r	S3r	S2gt	S1	S2r	S3r	S2r	S2r	S3r	S3r	S1	S2g	S1	S2g	S2g	S1	S2g	S2g	S2g	S2g	S2g
Berambadi	318	N	S3rt	N	S3r	N	S3r	N	N	Nrt	Nrt	S3r	N	S3r	N	N	N	S3rt	S3rt	S3rt	S3rt	Nrt	S3r	S3r	S3rt	S3rt	S3rt	S3rt
Berambadi	319	N	S3rt	N	S3r	N	S3r	N	N	Nrt	Nrt	S3r	N	S3r	N	N	N	S3rt	S3rt	S3rt	S3rt	Nrt	S3r	S3r	S3rt	S3rt	S3rt	S3rt
Berambadi	320	S3rt	S3r	S2r	S3r	S2r	S3r	S3r	S3r	Nr	Nr	S2r	S3r	S2r	S2r	S3r	S3r	S3r	S3r	S3r	S3r	Nr	S3r	S3r	S3r	S3r	S3r	S3r
Berambadi	321	S3rt	S3r	S2r	S3r	S2r	S3r	S3r	S3r	Nr	Nr	S2r	S3r	S2r	S2r	S3r	S3r	S3r	S3r	S3r	S3r	Nr	S3r	S3r	S3r	S3r	S3r	S3r
Berambadi	322	S3rt	S3r	S2r	S3r	S2r	S3r	S3r	S3r	Nr	Nr	S2r	S3r	S2r	S2r	S3r	S3r	S3r	S3r	S3r	S3r	Nr	S3r	S3r	S3r	S3r	S3r	S3r
Berambadi	323	S3rt	S3r	S2r	S3r	S2r	S3r	S3r	S3r	Nr	Nr	S2r	S3r	S2r	S2r	S3r	S3r	S3r	S3r	S3r	S3r	Nr	S3r	S3r	S3r	S3r	S3r	S3r
Berambadi	324	N	S3wt	S3wt	S1	N	S1	S1	S1	S2w	S2w	S1	N	S1	N	S1	S1	S3wt	S2wt	S2wt	S2wt	S2wt	S2w	S2w	S3wt	S3wt	S3wt	S2wt
Berambadi	325	N	S3wt	S3wt	S1	N	S1	S1	S1	S2w	S2w	S1	N	S1	N	S1	S1	S3wt	S2wt	S2wt	S2wt	S2wt	S2w	S2w	S3wt	S3wt	S3wt	S2wt
Berambadi	326	N	S3wt	S3wt	S1	N	S1	S1	S1	S2w	S2w	S1	N	S1	N	S1	S1	S3wt	S2wt	S2wt	S2wt	S2wt	S2w	S2w	S3wt	S3wt	S3wt	S2wt
Berambadi	327	N	S3wt	S3wt	S1	N	S1	S1	S1	S2w	S2w	S1	N	S1	N	S1	S1	S3wt	S2wt	S2wt	S2wt	S2wt	S2w	S2w	S3wt	S3wt	S3wt	S2wt
Berambadi	328	N	S3wt	S3wt	S1	N	S1	S1	S1	S2w	S2w	S1	N	S1	N	S1	S1	S3wt	S2wt	S2wt	S2wt	S2wt	S2w	S2w	S3wt	S3wt	S3wt	S2wt
Berambadi	340	N	S3wt	S3wt	S1	N	S1	S1	S1	S2w	S2w	S1	N	S1	N	S1	S1	S3wt	S2wt	S2wt	S2wt	S2wt	S2w	S2w	S3wt	S3wt	S3wt	S2wt
Berambadi	341	N	S3wt	S3wt	S1	N	S1	S1	S1	S2w	S2w	S1	N	S1	N	S1	S1	S3wt	S2wt	S2wt	S2wt	S2wt	S2w	S2w	S3wt	S3wt	S3wt	S2wt
Berambadi	342	S3r	S1	S2r	S1	S2r	S2t	S3r	S3r	S2gt	S1	S2r	S3r	S2r	S2r	S3r	S3r	S1	S2g	S1	S2g	S2g	S1	S2g	S2g	S2g	S2g	S2g
Berambadi	343	S3r	S1	S2r	S1	S2r	S2t	S3r	S3r	S2gt	S1	S2r	S3r	S2r	S2r	S3r	S3r	S1	S2g	S1	S2g	S2g	S1	S2g	S2g	S2g	S2g	S2g

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Marigold	Chrysanthemum	Banana	Horse gram	Field-bean	Turmeric	Beetroot	Potato	Beans
Berambadi	344	S3r	S1	S2r	S1	S2r	S2t	S3r	S3r	S2gt	S1	S2r	S3r	S2r	S2r	S3r	S3r	S1	S2g	S1	S2g	S2g	S1	S2g	S2g	S2g	S2g	S2g
Berambadi	346	N	S3wt	S3wt	S1	N	S1	S1	S1	S2w	S2w	S1	N	S1	N	S1	S1	S3wt	S2wt	S2wt	S2wt	S2wt	S2w	S2w	S3wt	S3wt	S3wt	S2wt
Berambadi	347	N	S3rt	N	S3r	N	S3r	N	N	Nrt	Nrt	S3r	N	S3r	N	N	N	S3rt	S3rt	S3rt	S3rt	Nrt	S3r	S3r	S3rt	S3rt	S3rt	S3rt
Berambadi	350	N	S3wt	S3wt	S1	N	S1	S1	S1	S2w	S2w	S1	N	S1	N	S1	S1	S3wt	S2wt	S2wt	S2wt	S2wt	S2w	S2w	S3wt	S3wt	S3wt	S2wt
Berambadi	355	S1	S1	S1	S2t	S1	S3t	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1
Berambadi	356	S3rt	S3r	S2r	S3r	S2r	S3r	S3r	S3r	Nr	Nr	S2r	S3r	S2r	S2r	S3r	S3r	S3r	S3r	S3r	S3r	S3r	Nr	S3r	S3r	S3r	S3r	S3r
Berambadi	357	S3rt	S3r	S2r	S3r	S2r	S3r	S3r	S3r	Nr	Nr	S2r	S3r	S2r	S2r	S3r	S3r	S3r	S3r	S3r	S3r	S3r	Nr	S3r	S3r	S3r	S3r	S3r
Berambadi	358	S3rt	S3r	S2r	S3r	S2r	S3r	S3r	S3r	Nr	Nr	S2r	S3r	S2r	S2r	S3r	S3r	S3r	S3r	S3r	S3r	S3r	Nr	S3r	S3r	S3r	S3r	S3r
Berambadi	359	S3rt	S3r	S2r	S3r	S2r	S3r	S3r	S3r	Nr	Nr	S2r	S3r	S2r	S2r	S3r	S3r	S3r	S3r	S3r	S3r	S3r	Nr	S3r	S3r	S3r	S3r	S3r
Berambadi	362	S3rt	S3r	S2r	S3r	S2r	S3r	S3r	S3r	Nr	Nr	S2r	S3r	S2r	S2r	S3r	S3r	S3r	S3r	S3r	S3r	S3r	Nr	S3r	S3r	S3r	S3r	S3r
Berambadi	363	S3rt	S3r	S2r	S3r	S2r	S3r	S3r	S3r	Nr	Nr	S2r	S3r	S2r	S2r	S3r	S3r	S3r	S3r	S3r	S3r	S3r	Nr	S3r	S3r	S3r	S3r	S3r
Berambadi	364	S3rt	S3r	S2r	S3r	S2r	S3r	S3r	S3r	Nr	Nr	S2r	S3r	S2r	S2r	S3r	S3r	S3r	S3r	S3r	S3r	S3r	Nr	S3r	S3r	S3r	S3r	S3r
Berambadi	366	N	S3wt	S3wt	S1	N	S1	S1	S1	S2w	S2w	S1	N	S1	N	S1	S1	S3wt	S2wt	S2wt	S2wt	S2wt	S2w	S2w	S3wt	S3wt	S3wt	S2wt
Berambadi	367	N	S3wt	S3wt	S1	N	S1	S1	S1	S2w	S2w	S1	N	S1	N	S1	S1	S3wt	S2wt	S2wt	S2wt	S2wt	S2w	S2w	S3wt	S3wt	S3wt	S2wt
Berambadi	368	N	S3rt	N	S3r	N	S3r	N	N	Nrt	Nrt	S3r	N	S3r	N	N	N	S3rt	S3rt	S3rt	S3rt	Nrt	S3r	S3r	S3rt	S3rt	S3rt	S3rt
Berambadi	370	N	S3wt	S3wt	S1	N	S1	S1	S1	S2w	S2w	S1	N	S1	N	S1	S1	S3wt	S2wt	S2wt	S2wt	S2wt	S2w	S2w	S3wt	S3wt	S3wt	S2wt
Maddur	33	N	S3wt	S3wt	S1	N	S1	S1	S1	S2w	S2w	S1	N	S1	N	S1	S1	S3wt	S2wt	S2wt	S2wt	S2wt	S2w	S2w	S3wt	S3wt	S3wt	S2wt
Maddur	56	N	S3wt	S3wt	S1	N	S1	S1	S1	S2w	S2w	S1	N	S1	N	S1	S1	S3wt	S2wt	S2wt	S2wt	S2wt	S2w	S2w	S3wt	S3wt	S3wt	S2wt
Maddur	61	N	S2r	S3r	S2r	S3r	S2r	S3r	S3r	S3r	S3r	S2r	S3r	S2r	S3r	S3r	S3r	S2rt	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S2r	S2r	S2r
Maddur	62	N	S3wt	S3wt	S1	N	S1	S1	S1	S2w	S2w	S1	N	S1	N	S1	S1	S3wt	S2wt	S2wt	S2wt	S2wt	S2w	S2w	S3wt	S3wt	S3wt	S2wt
Maddur	63	N	S3wt	S3wt	S1	N	S1	S1	S1	S2w	S2w	S1	N	S1	N	S1	S1	S3wt	S2wt	S2wt	S2wt	S2wt	S2w	S2w	S3wt	S3wt	S3wt	S2wt
Maddur	64	N	S3wt	S3wt	S1	N	S1	S1	S1	S2w	S2w	S1	N	S1	N	S1	S1	S3wt	S2wt	S2wt	S2wt	S2wt	S2w	S2w	S3wt	S3wt	S3wt	S2wt
Maddur	66	N	S3wt	S3wt	S1	N	S1	S1	S1	S2w	S2w	S1	N	S1	N	S1	S1	S3wt	S2wt	S2wt	S2wt	S2wt	S2w	S2w	S3wt	S3wt	S3wt	S2wt
Maddur	69	N	S2r	S3r	S2r	S3r	S2r	S3r	S3r	S3r	S3r	S2r	S3r	S2r	S3r	S3r	S3r	S2rt	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S2r	S2r	S2r
Maddur	70	N	S3wt	S3wt	S1	N	S1	S1	S1	S2w	S2w	S1	N	S1	N	S1	S1	S3wt	S2wt	S2wt	S2wt	S2wt	S2w	S2w	S3wt	S3wt	S3wt	S2wt

PART-B

SOCIO-ECONOMIC STATUS OF FARM HOUSEHOLDS

CONTENTS

1.	Executive summary	1-4
2.	Introduction	5
3.	Methodology	7-11
5.	Results and discussions	13-31

LIST OF TABLES

I. Social status		
1	Human population among sample households	13
2	Basic needs of sample households	14
3	Institutional participation among the sample population	15
II. Economic status		
4	Occupational pattern in sample households	16
5	Domestic assets among samples households	16
6	Farm assets among samples households	17
7	Livestock assets among sample households	18
8	Milk produced and Fodder availability of sample households	19
9	Women empowerment of sample households	19
10	Per capita daily consumption of food among the sample farmers	20
11	Annual average Income from various sources	21
12	Average annual expenditure of sample farmers	21
13	Land use among samples households	22
III. Resource use pattern		
14	Number of tree/plants covered in sample farm households	22
15	Present cropping pattern among samples households	23
16	Distribution of soil series in the watershed	24
IV. Economic land evaluation		
17	Cropping pattern on major soil series	25
18	Alternative land use options for different size group of farmers (Benefit Cost Ratio)	25
19	Economics Land evaluation and bridging yield gap for different crops	26
20	Estimation of onsite cost of soil erosion	28
21	Ecosystem services of food production	29
22	Ecosystem services of fodder production	30
23	Ecosystem services of water supply for crop production	30
24	Farming constraints	31

LIST OF FIGURES

1	Location of study area	8
2	ALPES Framework	9
3	Basic needs of sample households	15
4	Domestic assets among the sample households	17
5	Farm assets among samples households	17
6	Livestock assets among sample households	18
7	Per capita daily consumption of food among the sample farmers	20
8	Average annual expenditure of sample households	22
9	Present cropping pattern	23
10	Estimation of onsite cost of soil erosion	28
11	Ecosystem services of food production	29
12	Ecosystem services of water supply	31

EXECUTIVE SUMMARY

Baseline socioeconomic characterisation is prerequisite to prepare action plan for program implementation and to assess the project performance before making any changes in the watershed development program. The baseline provides appropriate policy direction for enhancing productivity and sustainability in agriculture.

Methodology: *Bachanahalli-1 micro-watershed (Gopalapur sub-watershed, Gundlupet taluk, Chamarajanagar district) is located in between 11°44' – 11°46' North latitudes and 76°31' – 76°34' East longitudes, covering an area of about 567 ha, bounded by Maddur, Berambadi villages and Berambadi State Forest with length of growing period (LGP) 120-150 days. We used soil resource map as basis for sampling farm households to test the hypothesis that soil quality influence crop selection, and conservation investment of farm households. The level of technology adoption and productivity gaps and livelihood patterns were analyses. The cost of soil degradation and ecosystem services were quantified.*

Results: *The socio-economic outputs for the Bachanahalli-1 micro-watershed (Gopalapur sub-watershed, Gundlupet taluk, Chamarajanagar district) are presented here.*

Social Indicators;

- ❖ *Male and female ratio is 53.4 to 46.5 per cent to the total sample population.*
- ❖ *Younger age 18 to 50 years group of population is 53.5 around per cent to the total population.*
- ❖ *Literacy population is around 65.2 per cent.*
- ❖ *Social groups belong to scheduled caste (SC) around 20 per cent.*
- ❖ *Liquefied petroleum gas (LPG) is the source of energy for a cooking among 90.0 per cent.*
- ❖ *About 80.0 per cent of households have a yashaswini health card.*
- ❖ *Farm households are having MGNREGA card only 20 per cent for rural employment.*
- ❖ *Dependence on ration cards for food grains through public distribution system among all sample households.*
- ❖ *Swachh bharath program providing closed toilet facilities around 70 per cent of sample households.*
- ❖ *Institutional participation is only 16.2 per cent of sample households.*
- ❖ *Rural migration to urban centre for employment is prevalent among 0.5 per cent of farm households.*
- ❖ *Women participation in decisions making are around 33.3 per cent of households.*

Economic Indicators;

- ❖ *The average land holding is 1.58 ha indicates that majority of farm households are belong to small and medium farmers. The dry land of 87.0 per cent and irrigated land 12.9 per cent of total cultivated land area among the sample farmers.*
- ❖ *Agriculture is the main occupation among 9.3 per cent and agriculture is the main non agriculture labour is subsidiary occupation for 83.7 per cent of sample households.*
- ❖ *The average value of domestic assets is around Rs. 12143. per household. Mobile and television are popular media mass communication.*
- ❖ *The average value of farm assets is around Rs. 31311 per household; Among all cent of sample farmers own plough and tractor (20 %).*
- ❖ *The average value of livestock is around Rs. 18500 per household; about 45.4 per cent of household are having livestock.*
- ❖ *The average per capita food consumption is around 802.12 grams (2019.3 kilo calories) against national institute of nutrition (NIN) recommendation at 827 gram. Around 60 per cent of sample households are consuming less than the NIN recommendation.*
- ❖ *The annual average income is around Rs. 50208 per household. About 70.0 per cent of farm households are below poverty line.*
- ❖ *The per capita monthly average expenditure is around Rs.1392.*

Environmental Indicators-Ecosystem Services;

- ❖ *The value of ecosystem service helps to support investment to decision on soil and water conservation and in promoting sustainable land use.*
- ❖ *The onsite cost of different soil nutrients lost due to soil erosion is around Rs.765 per ha/year. The total cost of annual soil nutrients is around Rs.287031 per year for the total area of 567 ha.*
- ❖ *The average value of ecosystem service for food grain production is around Rs 38998/ ha/year. Per hectare food grain production services is maximum in turmeric (Rs.290477) followed by garlic (Rs. 95513), cotton (Rs. 26050), maize (Rs. 20493), bengal gram (Rs. 9922), onion (Rs. 7611), marigold (Rs. 3975), groundnut (Rs. 2855), sunflower (Rs. 2659) and sorghum of negative net returns.*
- ❖ *The average value of ecosystem service for fodder production is around Rs. 1539/ ha/year. Per hectare fodder production services is maximum in groundnut (Rs. 2413) followed by ragi (Rs. 1900), sorghum (Rs. 1482), maize (Rs. 1240), horse gram (Rs. 659).*
- ❖ *The data on water requirement for producing one quintal of grain is considered for estimating the total value of water required for crop production. The per*

hectare value of water used and value of water was maximum in turmeric (Rs. 69577) followed by cotton (Rs. 57375), bengal gram (Rs. 50667), sorghum (Rs. 45171), maize (Rs. 36553), onion (Rs. 26874), garlic (Rs. 21822), groundnut (Rs. 19176) and ragi (Rs. 13931).

Economic Land Evaluation;

- ❖ The major cropping pattern is areca nut (31.1 %) followed by cotton (17.3 %), maize (15.9 %), horse gram (9.8 %), marigold (9.0 %), groundnut (5.9 %), onion (2.8 %), garlic (2.8 %), sorghum (2.8 %) and turmeric (2.8 %).
- ❖ In Bachanahalli 1 micro-watershed, major soil are soil of alluvial landscape of Hundipura (HDH) series is having shallow soil depth cover around 4.78 ha (0.84 %) of area. On this soil farmers are presently growing cotton (57.4 %), maize (21.3 %), and turmeric (21.3). Annurkeri (ARK) is also having shallow soil depth cover 35.71 per cent of area, the crops are horse gram (50 %) and marigold (50 %). Honnegoudanahalli (HGH) soil series having deep soil depth cover around 30.57 % of areas, crops are Ragi. Beramabadi (BMB) soil series having shallow soil depth cover around 51.41 ha (9.08 %) of area, crops are groundnut and Gopalapura (GPR) soil series are having moderately deep soil depth cover around 122.1 ha (21.5 %) of area. The major crops grown are cotton (21.7 %), sunflower (16.2 %), bengal gram (6.1 %), garlic (6 %), onion (6.0%), marigold (6.1) and maize (3.9).
- ❖ The total cost of cultivation and benefit cost ratio (BCR) in study area for coconut ranges between Rs.62778/ha in GPR soil (with BCR of 1.22) and Rs.26977/ha in HDR soil (with BCR of 2.94).
- ❖ In maize the cost of cultivation range between Rs 51457/ha in HDR soil (with of 2.14) and Rs.20555/ha in GPR soil (with BCR of 1.41).
- ❖ In marigold the cost of cultivation range between Rs. 65497/ha in GPR soil (with BCR of 1.04) and Rs. 35720/ha in ARK soil (with BCR of 1.18).
- ❖ In turmeric the cost of cultivation is Rs.129423/ha in HDR soil (with BCR of 3.24).
- ❖ In bengal gram the cost of cultivation is Rs 32631/ha in GPR soil (with BCR of 1.3).
- In garlic the cost cultivation is Rs. 200887/ha in GPR soil (with BCR of 1.48).
- In sorghum the cost cultivation is Rs. 41512/ha in GPR soil (with BCR of 1.04).
- In sunflower the cost of cultivation is Rs. 24235/ha in GPR soil (with BCR of 1.11).
- In horse gram the cost cultivation is Rs. 11981/ ha ARK soil (with BCR of 1.18).
- In groundnut the cost cultivation is Rs. 26785/ha in BMB soil (with BCR of 1.20) and ragi the cost cultivation is Rs. 30251/ha in HGH in soil (with BCR of 1.19).

- *The land management practices reported by the farmers are crop rotation, tillage practices, fertilizer application and use of farm yard manure (FYM). Due to higher wages farmer are following labour saving strategies is not prating soil and water conservation measures. Less ownership of livestock limiting application of FYM.*
- *It was observed soil quality influences on the type and intensity of land use. More fertilizer applications on deeper soil to maximize returns.*

Suggestions;

- *Involving farmers in watershed planning helps in strengthening institutional participation.*
- *The per capita food consumption and monthly income is very low. Diversifying income generation activities from crop and livestock production in order to reduce risk related to drought and market prices.*
- *Majority of farmers reported that they are not getting timely support/extension services from the concerned development departments.*
- *By strengthening agricultural extension for providing timely advice improved technology there is scope to increase in net income of farm households.*
- *By adopting recommended package of practices by following the soil test fertiliser recommendation, there is scope to increase yield in coconut (9 to 21 %), maize (35 to 79 %), marigold (0 to 21.3 %), bengal gram (49.9 %), garlic (52.6%), groundnut (59.6 %), horse gram (32.5%), onion (59.6 %), ragi (62.6%), sorghum (47.2%) and sunflower (60.6 %).*

INTRODUCTION

Watershed Development program aim to restore degraded watersheds in rainfed regions to increase their capacity to capture and store rain water, reduce soil erosion, and improved soil nutrients and carbon contents so they can produce greater agricultural yields and other benefits. As majority of rural poor live in these regions and dependent on natural resources for their livelihood and sustenance, improvements in agricultural yields improve human welfare and simultaneously improve national food security.

Sujala-III watershed development project conceptualised and implemented by the Watershed Development Department of Government of Karnataka with tripartite cost-sharing arrangements. The World Bank through International Development Association provided major portion of plan outlay as a loan to Government of India and in turn loan to Government of Karnataka.

The objectives of Sujala-III is to demonstrate more effective watershed management through greater integration of programs related to rain fed agriculture, innovative and science based approaches and strengthened institutions and capacities. The project is implemented in 11 districts of Bidar, Vijayapura, Gulbarga, Yadgir, Koppal, Gadag, Raichur, Davanagere, Tumkur, Chikkamangalur and Chamarajanagar which have been identified by the Watershed Development Department based on rainfall and socio-economic conditions. The project will be implemented over six years and linked with the centrally financed integrated watershed management programme.

Economic evaluations can better guide in watershed planning and implementation, as well as raise awareness of benefits of ecosystem restoration for food security and poverty alleviation program. The present study aims to characterize socio-economic status of farm households, assess the land and water use status, evaluate the economic viability of land use, prioritize farming constraints and suggest the measures for soil and water conservation for sustainable agriculture.

Objectives of the study

1. To characterize socio-economic status of farm households
2. To evaluate the economic viability of land use and land related constraints
3. To estimate the ecosystem service provided by the watershed and
4. To suggest alternatives for sustainable agriculture production.

METHODOLOGY

Study area

Bachanahalli-1 micro-watershed is located in Southern Dry Zone of Karnataka (Figure 1). It has a total geographical area of 1.56 M ha with 0.74 M ha under cultivation of which 0.22 M ha is irrigated. The mean elevation ranges from 450 to 900 m MSL; most part of the zone is situated at 800-900m. The major soils are red loams with pockets of black soils in Kollegal, Yalandur and T.N. Pura taluks of Mysore district. The average annual rainfall ranges from 670 to 890 mm, of which about 55 to 75 per cent is received during the kharif season. The major crops grown are rice, ragi, sugarcane, pulses and minor millets. It's represented Agro Ecological Region (AER) – 3 having LGP 120-150 days.

Bachanahalli-1 micro-watershed (Gopalapur sub-watershed, Gundlupet taluk, Chamaraajanagar district) is located in between $11^{\circ}44'$ – $11^{\circ}46'$ North latitudes and $76^{\circ}31'$ – $76^{\circ}34'$ East longitudes, covering an area of about 567 ha, bounded by Maddur, Berambadi villages and Berambadi State Forest.

Sampling Procedure:

In this study we have followed soil variability as criterion for sampling the farm households. In each micro-watershed the survey numbers and associated soil series are listed. Minimum three farm households for each soil series were taken and summed up to arrive at total sample for analysis.

Sources of data and analysis:

For evaluating the specific objectives of the study, primary data was collected from the sample respondents by personal interview method with the help of pre-tested questionnaire. The data on socio-economic characteristics of respondents such as family size and composition, land holdings, asset position, occupational pattern and education level was collected. The present cropping pattern and the level of input use and yields collected during survey. The data collected from the representative farm households were analysed using Automated Land Potential Evaluation System (Figure 2).




Figure 1: Location of study area

Steps followed in socio-economic assessment

- 1 • After the completion of soil profile study link the cadastral number to the soil profile in the micro watershed.
- 2 • Download the names of the farmers who are owning the land for each cadastral number in the Karnataka BHOOMI Website.
- 3 • Compiling the names of the farmers representing for all the soil profiles studied in the micro watershed for socio-economic Survey.
- 4 • Conducting the socioeconomic survey of selected farm households in the micro watershed .
- 5 • Farm households database created using the Automated Land Potential Evaluation System (ALPES) for analysis of socio economic status for each micro watershed .
- 6 • Synthesis of tables and preparation of report for each micro watershed .


KOPPAL-Kavalur Village 1-213

Home Create External Data Database Tools

» Main

AUTOMATED LAND POTENTIAL EVALUATION SYSTEM

ALPES



FARM HOUSE HOLD DIGITAL LIBRARY

ADD DATA
EDIT DATA
ANALYSIS
EXIT

Form View

Home Create External Data Database Tools

» Main Report

Livelihood Patterns

I. SOCIAL STATUS

Table 1. Spatial distribution of sample

Table 3. Total human population in sample HHs

Average Family size

Age groups

Education status

Table 4. Drinking water facilities

Table 5. Institutional participation

Average age of HHs

Farming Constraints

Farmers awareness of climate change

Reasons

ICT Use for agril information

II. ECONOMIC STATUS

Table 6: Migration Details

Nature of Job migrates

Table 7. Domestic assets

Table 8 Farm assets

Table 9 Livestock assets

Table 10 Food consumption in KiloCalories

Grams

Table 11 Annual average income (Rs)

Table 12 Crop production income

Table 13 Average annual Expenditure

Poverty estimation

Priority for crop selection

Kind of Government support

Socio-economic Indicators

III. RESOURCE USE PATTERN

Table 14 Land holding sample HHs

Table 15 Land use among sample HHs

Table 16 Cropping pattern

IV. ECONOMIC LAND EVALUATION

Table 17. Cropping pattern on diff soil series

Table 18. Economics different crops in Soil Series

Yield gap analysis of crops

Optimum land use planning

Input-Output coefficients

Ecosystem

Food Provision

Fodder Provision

Livestock Economics

Water consumption budgeting

Back

Record: 1 of 1 No Filter Search

Form View

Figure 2: ALPES FRAMEWORK

The sample farmers were post classified in to marginal and small (0.0 to ≤ 2 ha), medium and semi medium (>2 to ≤ 10 ha) and large (>10 ha). The steps involved in estimation of soil potential involve estimation of total cost of cultivation, the yield/gross returns and net income per hectare. The cost of inputs such seed, manure and fertilizer, plant protection chemicals, payment towards human and bullock labour and interest on working capita are included under operational costs. In the case of perennial crops, the cost of establishment was estimated by using actual physical requirements and prevailing market prices. Estimation cost included maintenance cost up to bearing period. The value of main product and by product from the crop enterprise at the market rates were the gross returns of the crop. Net returns were worked out by deducting establishment and maintained cost from gross returns.

Operational Cost = cost of seeds, fertilizers, pesticides. Cost of human and bullock labour, cost of machinery, cost of irrigation water + interest on working capital.

Gross returns = Yield (Quintals/hectare)*Price (Rs/Quintal)

Net returns = Gross returns-Operational cost.


Benefit Cost Ratio = Net returns/Total cost.

Economic suitability classes: once each land use –land area combination has been assigned an economic value by the land evaluation, the question arises as to its ‘suitability’, that is, the degree to which it satisfies the land user. The FAO framework defines two suitability orders: ‘S’(suitable if benefit cost ratio (BCR) >1) and ‘N’(not suitable if (BCR <1), which are divided into five economic suitability classes: ‘S1’(highly suitable if BCR >3), ‘S2’(suitable if BCR >2 and <3), ‘S3’(Marginally suitable if BCR >1 and <2), ‘N1’(Not suitable for economic reasons but physically suitable) and ‘N2’(not suitable for physical reasons). The limit between ‘S3’ and ‘N1’ must be at least at the point of financial feasibility (i.e. net returns, NPV, or IRR >0 and BCR >1). The other limits depend on social factors such as farm size, family size, alternative employment or investment possibilities and wealth expectations; these need to be specified for the Soil series.

Economic Valuation of Soil ecosystem services:

The replacement cost approach was followed for estimating the onsite cost of soil erosion, Market price method was followed for estimating the value of food and fodder production. Value transfer methods was followed for estimating the value of water demand by different crops in the micro watershed.

Steps followed in Replacement cost methods for estimation of onsite cost of soil erosion



RESULTS AND DISCUSSIONS

The demographic information shows that the household population dynamics encompasses the socioeconomic status of the farmer. For a rural family, the household size should be optimal to earn a comfortable livelihood through farm and non-farm wage earning. The total number of population in watershed area was 43, out of which 53.4 per cent were males and 46.5 per cent females. Average family size of the households is 4.3. Age is an important factor, which affects the potential employment and mobility status of respondents. The data on age wise distribution of farmers in the sample households indicated that majority of the farmers are coming under the age group of 18 to 30 years (27.9 %) followed by 0 to 18 years (16.2 %), 30 to 50 years (23.2 %) and more than 50 years (32.5 %). Hence, in the study area in general, the respondents were of young and middle age, indicating thereby that the households had almost settled with whatever livelihood options they were practicing and sample respondents were young by age who could venture into various options of livelihood sources. Data on literacy indicated that 65.2 per cent of respondents were illiterate and 34.8 per cent literate (Table 1).

Table 1: Human population among sample households in Bachanahalli 1 Microwatershed

Particulars	Units	Value
Total human population in sample HHs	Number	43.0
Male	% to total Population	53.4
Female	% to total Population	46.5
Average family size	Number	4.3
Age group		
0 to 18 years	% to total Population	12.5
18 to 30 years	% to total Population	16.0
30 to 50 years	% to total Population	37.5
>50 years	% to total Population	33.9
Average age	Age in years	41.5
Education Status		
Illiterates	% to total Population	34.8
Literates	% to total Population	65.2
Primary School (<5 class)	% to total Population	11.6
Middle School (6- 8 class)	% to total Population	16.2
High School (9- 10 class)	% to total Population	20.9
Others	% to total Population	16.2

The ethnic groups among the sample farm households found to be 20.0 per cent belonging to other backward caste (OBC) followed by 20.0 per cent belonging to scheduled caste (SC) and followed by 60 per cent general caste (Table 2 and Figure 3).

About 10.0 per cent of sample households are using fire wood as source of fuel for cooking and LPG gas (90.0%). All the sample farmers are having electricity connection. About 80.0 per cent are sample households having health cards. Majority (20 %) are having MNREGA job cards for employment generation. Among all farm households are having ration cards for taking food grains from public distribution system. About 70.0 per cent of farm households are having toilet facilities.

Table 2: Basic needs of sample households in Bachanahalli 1 Micro watershed

Particulars	Units	Value
Social groups		
OBC	% of Households	20.0
SC	% of Households	20.0
General	% of Households	60.0
Types of fuel use for cooking		
Fire wood	% of Households	10.0
Gas	% of Households	90.0
Energy supply for home		
Electricity	% of Households	100
Number of households having Health card		
Yes	% of Households	80.0
No	% of Households	20.0
MGNREGA Card		
Yes	% of Households	20.0
No	% of Households	80.0
Ration Card		
Yes	% of Households	100.0
No	% of Households	0.0
Households with toilet		
Yes	% of Households	70.0
No	% of Households	30.0
Drinking water facilities		
Tube well	% of Households	100.0

The data collected on the source of drinking water in the study area is presented in Table 2. Majority of the sample respondents are having Tube well source for water supply for domestic purpose (100 %).

Only 16.3 per cent of the farmers are participating in community based organizations (Table 3). Among them majority were participating in self help group organization (9.4 %) followed by users group (6.9 %) and like Sri Dharmasthala Swasahaya Sangha, Stri Shakhti Sangha.




Figure 3: Basic needs of sample households in Bachanahalli 1 Microwatershed

Table 3: Institutional participation among the sample population in Bachanahalli 1 Microwatershed

Particulars	Units	Value
No. Of people participating	% to total	16.3
Self help groups(SHG's)	% to total	9.4
Users groups	% to total	6.9
No. Of people not participating	% to total	83.7

The occupational pattern (Table 4) among sample households shows that agriculture is the main occupation is 9.3 per cent and agriculture is the main occupation and agriculture labour is subsidiary occupation was around 83.7 per cent of farmers.

Table 4: Occupational pattern in sample population in Bachanahalli 1 Microwatershed

Occupation		% to total
Main	Subsidiary	
Agriculture	Agriculture	9.3
	Agriculture Labour	83.7
Studying		7.0
Grand Total		100.0
Family labour availability		Man days/month
Male		38.9
Female		31.1
Total		70.0

The important assets especially with reference to domestic assets were analyzed and are given in Table 5 and Figure 4. The important domestic assets possessed by all categories of farmers are mobile phones (80.0 %) followed by television (80.0 %), mixer/grinder (70.0 %), bicycle (30.0 %) and motorcycle (20.0 %). The average value of domestic assets is around Rs 12143 per households.

Table 5: Domestic assets among the sample households in Bachanahalli 1 Microwatershed

Particulars	% of households	Average value in Rs
Bicycle	30.0	800
Mixer/grinder	70.0	2228
Mobile Phone	80.0	7000
Motor cycle	20.0	44000
Television	90.0	6668
Average value	12143	

The most popularly owned farm equipments were sickles, plough, cattle shed; pump sets, chaff cutter, bullock cart, sprayer and thresher. Plough and sickle were commonly present in all the sampled farmers; these were primary implements in agriculture. The per cent of households owned plough (100 %), weeder (10 %), bullock cart (20 %), sprayer (20 %), harvester (20 %), tractor (20 %) and drip/sprinkler (10 %) was found highest among the sample farmers. The average value of farm assets is around Rs 31311 per households (Table 6 and Figure 5).




Figure 4: Domestic assets among the sample households in Bachanahalli 1 Microwatershed

Table 6: Farm assets among samples households in Bachanahalli 1 Microwatershed

Particulars	% of households	Average value in Rs
Bullock cart	20	22500
Drip/sprinkler	10	2700
Harvester	20	1550
Plough	100	1833
Sprayer	20	3500
Tractor	20	500000
Weeder	10	208
Average value	31311	




Figure5: Farm assets among samples households in Bachanahalli 1 Microwatershed

Livestock is an integral component of the conventional farming systems (Table 7 and Figure 6). The highest livestock population is bullocks were around 40 per cent followed by local dry cow (20 %), local milching cow (20 %), and crossbred milching cow (20 %). The average livestock value was Rs 18500 per livestock.

Table 7: Livestock assets among sample households in Bachanahalli 1 Micro-watershed

Particulars	% of livestock population	Average value in Rs
Bullocks	40.0	10000
Crossbred Milching Cow	20.0	3000
Local Dry Cow	20.0	42500
Local Milching Cow	20.0	
Average value	18500	




Figure 6: Livestock assets among sample households in Bachanahalli 1 micro-watershed

Average milk produced in sample households is 900 liters/ annum. Among the farm households, sorghum, bajra and maize are the main crops for domestic food and fodder for animals. About 1498.6 kg /ha of average fodder is available per season for the livestock feeding (Table 8).

Table 8: Milk produced and fodder availability of sample households in Bachanahalli 1 Microwatershed

Particulars	
Name of the Livestock	Ltr./Lactation/animal
Crossbred Milching Cow	1440
Local Milching Cow	360
Average Milk produced	900
Fodder produces	Fodder yield (kg/ha.)
Sorghum	1666
Groundnut	1162
Horsegram	1666
Average fodder availability	1498
Livestock having households (%)	45.4
Livestock population (Numbers)	6

A woman participation in decision making in this micro-watershed is presented in Table 9. About 40.0 per cent of women participation in local organisation activities, 10 per cent of women elected as panchayat member and 50.0 per cent of women taking decision in her family and agriculture related activities.

Table 9: Women empowerment of sample households in Bachanahalli 1 Microwatershed

Particulars	% to Grand Total	
	Yes	No
Women participation in local organization activities	40.0	60.0
Women elected as panchayat member	10.0	90.0
Women earning for her family requirement	0.0	100
Women taking decision in her family and agriculture related activities	50.0	50.0

The food intake in terms of kilo calorie (kcal) per person per day was calculated and presented in the Table 10 and Figure 7. More quantity of cereals are consumed by sample farmers which accounted for 1412.1 kcal per person. The other important food items consumed was pulses 181.7 kcal followed by cooking oil 177 kcal, milk 106.3 kcal, vegetables 12.7 kcal, egg 110.4 kcal and meat 18.9 kcal. In the sampled households, farmers were consuming less (2019.3 kcal) than NIN- recommended food requirement (2250 kcal).

Table 10: Per capita daily consumption of food among the sample households in Bachanahalli 1 Microwatershed

Particulars	NIN recommendation (gram/ per day/ person)	Present level of consumption (gram/ per day/ person)	Kilo Calories /day/person
Cereals	396.0	415.3	1412.1
Pulses	43.0	52.9	181.7
Milk	200.0	163.5	106.3
Vegetables	143.0	52.9	12.7
Cooking Oil	31.0	31.0	177.0
Egg	0.5	73.6	110.4
Meat	14.2	12.6	18.9
Total	827.7	802.12	2019.3
Threshold of NIN recommendation		827 gram*	2250 Kcal*
% Below NIN		60.0	70.0
% Above NIN		40.0	30.0

Note: * day/person




Figure 7: Per capita daily consumption of food among the sample households in Bachanahalli 1 Microwatershed

Annual income of the sample HHs: The average annual household income is around Rs 50208. Major source of income to the farmers in the study area is from crop production (Rs 30673) followed by livestock (Rs. 19535). The income from livestock income was very low at Rs 19535. The monthly per capita income is Rs.973, which is

less than the threshold monthly income of Rs 975 for considering above poverty line. Due to the fact that erratic rainfall and shortage of water, farmers are diverting from crop production activities to enable the household for a comfortable livelihood. The incomes from the other aforesaid sources are very meagre (Table 11).

Table 11: Annual average income of HHs from various sources in Bachanahalli 1 Microwatershed

Particulars	Income *
Nonfarm income	0 (0)
Livestock income (Rs)	19535 (20)
Crop Production (Rs)	30673(100)
Total Annual Income (Rs)	50208
Average monthly per capita income (Rs)	973
Threshold for Poverty level (Rs 975 per month/person)	
% of households below poverty line	70.0
% of households above poverty line	30.0

* Figure in the parenthesis indicates % of Households

The average annual expenditure of farm households indicated that farmers in the study area spend highest on food (Rs. 39804) followed by education, clothing, social function and health. Now a day's education is most important among all of us. In today's competitive world, education is a necessity for man after food, clothing, and shelter. It is the only fundamental way by which a desired change in the society can happen. The average per capita monthly expenditure is around Rs 1392 and about 70 per cent of farm households are below poverty line (Table 12 and Figure 8).

Table 12: Average annual expenditure of sample HHs in Bachanahalli 1 Microwatershed

Particulars	Value in Rupees	Per cent
Food	39804	55.4
Education	13000	18.0
Clothing	5020	6.9
Social functions	4400	6.1
Health	9600	13.3
Total Expenditure (Rs/year)	71824	100
Monthly per capita expenditure (Rs)	1392	




Figure 8: Average annual expenditure of sample HHs in Bachanahalli 1 Microwatershed

Land use: The total land holding in the Bachanahalli 1 micro-watershed is 15.8 ha (Table 13). Of which 13.8 ha is rain fed land and 2.0 ha is irrigated land. The average land holding per household is worked out to be 1.58 ha.

Table 13: Land use among samples households in Bachanahalli 1 Micro watershed

Particulars	Per cent	Area in ha
Irrigated land	12.9	2.0
Rainfed Land	87.0	13.8
Fallow Land	0	0.0
Total land holding	15.8	15.8
Average land holding	1.58	

In the micro-watershed, the prevalent present land uses under perennial plants are neem trees (40.0 %) followed by teak tree (30.0%), jack fruit (10.0 %) and people tree (arali) (2.0 %) (Table 14).

Table 14: Number of trees/plants covered in sample farm households in Bachanahalli 1 Microwatershed

Particulars	Number of Plants/trees	Per cent
Neem trees	4	40.0
Teak	3	30
Peepole tree(Arali)	2	20.0
Jack Fruit	1	10.0
Grand Total	10	100.0

The land use decisions are usually based on experience of farmers, tradition, expected profit, personal preferences, resources and social requirements.

The present dominant crops grown in dry lands in the study area were by areca nut (27.5 %) followed by cotton (17.3 %), marigold (9.0 %), groundnut (5.9 %), garlic (2.8 %), maize (2.8 %), onion (2.8), sorghum (2.8) and turmeric (2.8) which are taken during kharif and maize (13.1 %), horse gram (9.8 %) and areca nut (3.6 %) during Rabi season respectively. The cropping intensity was 113.6 per cent (Table 15 and Figure 9).

Table 15: Present cropping pattern and cropping intensity in Bachanahalli 1 Microwatershed

Crops	Kharif	Rabi	Grand Total
Arecanut	27.5	3.6	31.1
Cotton	17.3	0.0	17.3
Garlic	2.8	0.0	2.8
Groundnut	5.9	0.0	5.9
Horsegram	0.0	9.8	9.8
Maize	2.8	13.1	15.9
Marigold	9.0	0.0	9.0
Onion	2.8	0.0	2.8
Sorghum	2.8	0.0	2.8
Turmeric	2.8	0.0	2.8
Grand Total	73.5	26.5	100
Cropping Intensity (%)	135.9		




Figure 9: Present cropping pattern in Bachanahalli 1 Microwatershed

Economic land evaluation

The main purpose of economic land evaluation in the watershed is to identify the existing production constraints and propose the potential/alternate options for agro-technology transfer and for bridging the adoption and yield gap.

In Bachanahalli 1 micro-watershed, 8 soil series are identified and mapped (Table 16). The distribution of major soil series are Annurkeri covering an area around 35.71 ha (6.30%) followed by Beemanabeedu 119.6 ha (21.1 %), Berambadi 51.4 ha (9.0 %), Gopalaura 122.1ha (21.5 %), Hundipura 4.78 ha (0.84%), Honnegaudanahalli 30.5 ha (5.3 %), Hullipura 9.53 ha (1.68 %), Kallipura 1.1 ha (0.2 %).

Table 16: Distribution of soil series in Bachanahalli 1 Microwatershed

Soil No	Soil Series	Mapping Unit Description	Area in ha (%)
1	ARK	Annurkeri soils are very deep (>150 cm), well drained, have dark reddish brown to very dusky red sandy clay to clay soils occurring on very gently sloping uplands under cultivation.	35.71 (6.30)
2	BMB	Beemanabeedu soils are very deep (>150 cm), moderately well drained, have very dark greyish brown to dark grey and very dark brown clayey soils occurring on very gently sloping lowlands under cultivation	119.68 (21.12)
3	BMD	Berambadi soils are shallow (25-50 cm), well drained, dark brown to dark greyish brown clayey soils occurring on very gently to gently sloping uplands under cultivation	51.41 (9.08)
4	GPR	Gopalapura soils are moderately deep (75-100 cm), well drained, have dark brown to dark reddish brown and reddish brown gravelly sandy clay loam to sandy clay soils occurring on very gently to gently sloping uplands under cultivation	122.10 (21.55)
5	HDR	Hindupur soils are shallow (25-50 cm), well drained, have dark reddish brown to dusky red sandy clay loam to sandy clay soils occurring on very gently sloping uplands under cultivation	4.78 (0.84)
6	HGH	Honnegaudanahalli soils are very deep (>150 cm), well drained, have very dark brown to brown and dark reddish brown sandy clay loam soils occurring on very gently sloping uplands under cultivation	30.57 (5.39)
7	HPR	Hullipura soils are moderately shallow (50-75 cm), well drained, have dark brown to very dark brown gravelly sandy clay loam to sandy clay soils occurring on very gently sloping uplands under cultivation	9.53 (1.68)
8	KLP	Kallipura soils are deep (100-150 cm), well drained, have dark reddish brown to dark red gravelly sandy clay loam to sandy clay soils occurring on very gently sloping uplands under cultivation	1.15 (0.20)
9		Forest	178.29 (31.47)
10		Others	13.33 (2.35)

Present cropping pattern on different soil series are given in Table 17. Crops grown on Hundipura soils are maize, cotton and Turmeric. Horsegram and marigold on Annurkeri soils is grown. Ragi on Honnegoudanahalli soils is grown. Groundnut on Beemanabeedu soil Bengal gram, cotton, maize, sunflower, garlic, marigold, onion and sorghum on Gopalpura soils are grow.

Table 17: Cropping pattern on major soil series in Bachanahalli 1 micro-watershed
(Area in per cent)

Soil Series	Soil Depth	Crops	Dry		Irrigated	Grand Total
			Kharif	Rabi	Kharif	
HDR	Shallow (25-50 cm)	Cotton	57.4	0.0	0.0	57.4
		Maize	0.0	0.0	21.3	21.3
		Turmeric	0.0	0.0	21.3	21.3
GPR	Moderately deep (75-100 cm)	Bengal gram	0.0	6.1	0.0	6.1
		Cotton	21.7	0.0	0.0	21.7
		Garlic	0.0	0.0	6.0	6.0
		Maize	15.6	16.2	0.0	31.9
		Marigold	6.1	0.0	0.0	6.1
		Onion	0.0	0.0	6.0	6.0
		Sorghum	6.0	0.0	0.0	6.0
		Sunflower	16.2	0.0	0.0	16.2
HGH	Very deep (>150 cm)	Ragi	100.0	0.0	0.0	100.0
BMB	Very deep (>150 cm)	Groundnut	100.0	0.0	0.0	100.0
ARK	Very deep (>150 cm)	Horse gram	0.0	50.0	0.0	50.0
		Marigold	50.0	0.0	0.0	50.0

Land is used for agricultural use for growing cereals, pulse, oilseeds and commercial crops. The soil/ land potential are measures in terms of physical yield and net income. The alternative land use options for each micro-watershed are given below (Table 18)

Table 18: Alternative land use options for different size group of farmers (Benefit Cost Ratio) in Bachanahalli 1 Micro watershed.

Soil Series	Small farmers	Large farmers
HDR	Cotton (2.94), maize (2.14) and Turmeric (3.24)	
GPR	cotton (1.11), Bengal gram (1.30), Maize (1.33), Garlic (1.48), Marigold(1.05), Onion (1.04), sorghum (1.04), sunflower (1.11)	Cotton (1.36), Marigold (1.50)
ARK	Horse gram (1.43), Marigold (1.18)	
HGH	Ragi (1.19).	
BMB	Groundnut (1.20).	

The productivity of different crops grown in Bachanahalli 1 micro-watershed under potential yield of the crops is given in Table 19.

Table 19: Economic land evaluation and bridging yield gap for different crops in Bachanahalli 1 micro-watershed

Particulars	HDR (25-50 cm)			GPR (75-100 cm)								ARK (>150 cm)		BMB (>150cm)	HGH (150 cm)
	Cot ton	Maize	Tur meric	Bengal gram	Cot ton	Garlic	Maize	Mari gold	Onion	Sorg hum	Sun flower	Horse gram	Mari gold	Groud nut	Ragi
Total cost (Rs/ha)	26977	51457	129423	32631	62778	200887	20555	65497	170229	41512	24236	11981	35720	26785	30251
Gross Return (Rs/ha)	79314	110162	419900	42552	75335	296400	28949	68475	177840	42978	26896	17125	42264	32053	36100
Net returns (Rs/ha)	52337	58705	290477	9922	12557	95513	8393	2978	7611	1466	2659	5144	6544	5267	5849
BCR	2.94	2.14	3.24	1.30	1.22	1.48	1.41	1.05	1.04	1.04	1.11	1.43	1.18	1.20	1.19
Farmers Practices (FP)															
FYM (t/ha)	0.0	0.0	15.0	0.0	1.0	5.0	0.0	0.0	5.0	5.0	0.9	0.0	0.0	0.0	3.8
Nitrogen (kg/ha)	38.0	160.0	160.0	22.3	50.6	72.5	50.2	22.3	72.5	51.3	50.5	35.6	35.6	84.9	61.5
Phosphorus (kg/ha)	42.6	115.0	115.0	56.9	85.5	102.5	107.4	56.9	102.5	57.5	101.4	25.6	25.6	80.2	44.2
Potash (kg/ha)	27.8	75.0	75.0	74.3	51.9	75.0	14.4	74.3	75.0	75.0	0.0	33.3	33.3	69.8	57.7
Grain (Qtl/ha)	15.7	55.0	42.5	7.4	13.8	37.5	17.9	99.0	100.0	15.0	6.5	6.7	77.8	7.0	11.5
Price of Yield (Rs/Qtl)	5100	2000	10000	5800	5550	8000	1575	700	1800	2800	4200	2500	550	4300	3000
Soil test based fertilizer Recommendation (STBR)															
FYM (t/ha)	12.4	8.6	24.7	7.4	12.4	24.7	8.6	19.8	29.6	7.4	6.6	0.0	19.8	8.6	8.6
Nitrogen (kg/ha)	148.2	123.5	148.2	18.5	148.2	123.5	123.5	222.3	123.5	81.5	55.2	30.9	277.9	30.9	74.1
Phosphorus (kg/ha)	74.1	61.8	123.5	37.1	74.1	61.8	61.8	59.3	74.1	56.8	59.3	37.1	59.3	61.8	43.2
Potash (kg/ha)	55.6	24.1	185.3	27.8	55.6	46.3	24.1	44.5	92.6	29.6	27.8	18.5	44.5	23.2	33.3
Grain (Qtl/ha)	17.3	84.0	24.7	14.8	17.3	79.0	84.0	98.8	247.0	28.4	16.5	9.9	98.8	17.3	30.9
% of Adoption/yield gap (STBR-FP) / (STBR)															
FYM (%)	100.0	100.0	39.3	100.0	92.2	79.8	100.0	100.0	83.1	32.5	85.9	0	100.0	100.0	55.5
Nitrogen (%)	74.4	-29.6	-8.0	-20.3	65.8	41.3	59.3	90.0	41.3	37.1	8.5	-15.2	87.2	-174.9	17.0
Phosphorus (%)	42.5	-86.2	6.9	-53.7	-15.4	-66.0	-74.0	4.0	-38.3	-1.2	-71.0	31.0	56.9	-29.9	-2.3
Potash (%)	50.0	-211.4	59.5	-167.2	6.6	-61.9	40.1	-67.0	0.0	-153.0	100.0	-79.9	0.0	-201.3	-73.0
Grain (%)	9.0	34.5	-72.1	49.9	20.5	52.6	78.7	-0.2	59.5	47.2	60.6	32.5	21.3	59.6	62.6
Value of yield and Fertilizer (Rs)															
Additional Cost (Rs/ha)	15615	4846	12137	5561	12132	17945	7708	21668	24355	1836	4420	153	24374	6251	4418
Additional Benefits (Rs/ha)	7901	57960	-178000	42887	19647	332320	104055	-147	264600	37534	41938	8033	11562	44347	58010
Net change Income (Rs/ha)	-7714	53114	-190137	37326	7515	314375	96348	-21815	240245	35698	37518	7880	-12812	38096	53591

The data on cost of cultivation and benefit cost ratio (BCR) of different crops is given in Tables 19. The total cost of cultivation in study area for coconut ranges between Rs.62778/ha in GPR soil (with BCR of 1.22) and Rs.26977/ha in HDR soil (with BCR of 2.94), maize range between Rs 51457/ha in HDR soil (with of 2.14) and Rs.20555/ha in GPR soil (with BCR of 1.41), marigold range between Rs. 65497/ha in GPR soil (with BCR of 1.04) and Rs. 35720/ha in ARK soil (with BCR of 1.18), turmeric cost of cultivation is Rs.129423/ha in HDR soil (with BCR of 3.24), bengal gram cost of cultivation is Rs 32631/ha in GPR soil (with BCR of 1.30), garlic cost cultivation is 200887/ ha in GPR soil (with BCR of 1.48), onion cost cultivation is 170229 ha in GPR in soil (with BCR of 1.1.04), sorghum cost cultivation is 41512/ ha in GPR soil (with BCR of 1.04) sunflower cost of cultivation is Rs 24235/ha in GPR soil (with BCR of 1.11), horse gram cost cultivation is 11981/ha in ARK soil (with BCR of 1.18), ground nut cost cultivation is 26785/ha in BMB soil (with BCR of 1.20) and ragi cost cultivation is 30251 ha in HGH in soil (with BCR of 1.19).


The data on FYM, Nitrogen, Phosphorus and Potash application by the farmers to different crops and recommended FYM for different crops is given in Table 19. There is a huge gap between FYM application by farmers and recommended FYM in all the crops across the soils. There is a larger yield gap in crops grown across different soil series. Adequate knowledge about recommended package of practices is the pre-requisite for their use in cultivation of crops. It is a fact that, recommended practices are major contributing factors to yield. Inadequate knowledge about recommended practices leads to their improper adoption. Strengthening of extension services by concerned agency is required to increase adoption of recommended cultivation practices and ultimately reducing the gap. By adopting soil-test fertiliser recommendation, there is scope to increase yield and income to a maximum of Rs 314375 in garlic and a minimum of Rs 7515 in cotton cultivation.

Economic valuation of Ecosystem Services (ES) was aimed at combining use and non-use values to determine Total Economic Value (TEV) of ES. Ecosystem Services (ES) were valued based on their annual flow or utilization in common monetary units, Rs/year. The valuation of ES was based on market price in 2017 or market cost approaches whichever is applicable, and in other cases on value or benefit transfer from previous valuation studies.

The onsite cost of different soil nutrients lost due to soil erosion is given in Table 20 and Figure 10. The average value of soil nutrient loss is around Rs 765.42 per ha/year. The total cost of annual soil nutrients is around Rs 287031 per year for the total area of 567 ha.

Table 20: Estimation of onsite cost of soil erosion in Bachanahalli 1 micro-watershed

Particulars	Quantity(kg)		Value (Rs)	
	Per ha	Total	Per ha	Total
Organic matter	102.50	38438	645.75	242158
Phosphorus	0.18	68	8.01	3002
Potash	1.61	603	32.14	12053
Iron	0.21	79	10.15	3808
Manganese	0.21	78	57.43	21537
Copper	0.02	6	9.17	3437
Zinc	0.00	1	0.13	49
Sulphur	0.06	23	2.48	930
Boron	0.00	1	0.15	57
Total	104.80	39298	765.42	287031

**Figure 10: Estimation of onsite cost of soil erosion in Bachanahalli 1 micro-watershed**

The average value of ecosystem service for food production is around Rs 38998/ha/year (Table 21 and Figure 11). Per ha food production services is maximum in turmeric (Rs 290477) followed by garlic (Rs 95513), cotton (Rs 26050), maize (Rs 20493), bengal gram Rs 9922), onion (Rs 7611), horse gram (Rs 4486), marigold (Rs 3975), ragi (Rs 3949), groundnut (Rs 2855) and sunflower (Rs 2659).

Table 21: Ecosystem services of food grain production in Bachanahalli 1 Microwatershed

Production items	Crops	Area in ha	Yield (Qtl/ha)	Price (Rs/Qtl)	Gross Returns (Rs/ha)	Cost of Cultivation (Rs/ha)	Net Returns (Rs/ha)
Cereals	Maize	2.6	29.9	1717	51349	30856	20493
	Ragi	0.5	11.4	3000	34200	30251	3949
	Sorghum	0.4	14.8	2800	41496	41512	-16
Pulses	Bengal gram	0.4	7.3	5800	42552	32631	9922
	Horse gram	0.9	6.6	2500	16467	11981	4486
Oil seeds	Groundnut	0.9	6.9	4300	29640	26785	2855
	Sunflower	1.1	6.4	4200	26896	24236	2659
Vegetables	Onion	0.4	98.8	1800	177840	170229	7611
Commercial Crops	Cotton	2.6	14.2	5400	76899	50844	26055
	Marigold	1.3	87.3	625	54583	50609	3975
Spice crops	Garlic	0.4	37.1	8000	296400	200887	95513
	Turmeric	0.4	42	10000	419900	129423	290477
Average value		11.9	30.2	4179	105685	66687	38998




Figure 11: Ecosystem services of food grains production in Bachanahalli 1 Microwatershed

The average value of ecosystem service for fodder production is around Rs 1539/ha/year (Table 22). Per ha fodder production services is maximum in groundnut (Rs 2413) followed by ragi (Rs1900), sorghum (Rs 1482), maize (Rs 1240) and horse gram (Rs 659).

Table 22: Ecosystem services of fodder production in Bachanahalli 1 Micro watershed

Production items	Crops	Area in ha	Yield (Qtl/ha)	Price (Rs/Qtl)	Net Returns (Rs/ha)
Cereals	Maize	2.6	2.1	600	1240
	Ragi	0.5	1.9	1000	1900
	Sorghum	0.4	2.5	600	1482
Pulses	Horse gram	0.9	1.1	600	659
Oil seeds	Groundnut	0.9	1.1	2100	2413
Average value		5.3	1.7	980	1539

The water demand for production of different crops was worked out in arriving at the ecosystem services of water support to crop growth. The data on water requirement for producing one quintal of grain is considered for estimating the total value of water required for crop production. The per hectare value of water used and value of water was maximum (Table 23 and Figure 12) in turmeric (Rs 69577) followed by cotton (Rs 57375), bengal gram (Rs 50667), sorghum (Rs 45171), maize (Rs 36553), onion (Rs 26874), garlic (Rs 21822), sunflower (Rs 21555), groundnut (Rs 19176) and ragi (Rs 13931).

Table 23: Ecosystem services of water supply in Bachanahalli 1 Micro watershed

Crops	Yield (Qtl/ha)	Virtual water (cubic meter) per ha	Value of Water (Rs/ha)	Water consumption (Cubic meters/Qtl)
Bengalgram	7.3	5067	50667	691
Cotton	14.2	5738	57375	403
Garlic	37.1	2182	21822	59
Groundnut	6.9	1918	19176	278
Horsegram	6.6	2027	20274	308
Maize	29.9	3655	36553	122
Onion	98.8	2687	26874	27
ragi	11.4	1393	13931	122
Sorghum	14.8	4517	45171	305
Sunflower	6.4	2155	21555	337
Turmeric	42.0	6958	69577	166
Grand Total	25.0	3482	34816	256

The main farming constraints in Bachanahalli 1 micro-watershed to be found are less rainfall, damage of crops by wild animals, non availability of plant protection chemicals, non availability fertilizers, lack of transportation, lack of good quality seeds and high crop pests & diseases. Majority of farmers depend up on money lender of the sources of loan for purpose of crop production. Farmers to sell the agriculture produce through village market and the farmers getting the agriculture related information on

newspaper and television. Farmers reported that they are not getting timely support/extension services from the concerned development department (Table 24).




Figure 12: Ecosystem services of water supply in Bachanahalli 1 Microwatershed

Table 24: Farming constraints related land resources of sample households in Bachanahalli 1 Microwatershed

Sl. No	Particulars	Per cent
1	Less Rainfall	80.0
2	Lack of good quality seeds	10.0
3	Non availability Fertilizers	20.0
4	High Crop Pests & Diseases	10.0
5	Lack of transportation	20.0
7	Damage of crops by Wild Animals	80.0
8	Non availability of Plant Protection Chemicals	70.0
9	Source of loan	
	Money Leander	100.0
10	Market for selling	
	Village market	100.0
11	Sources of Agri-Technology information	
	Newspaper	100.0

The findings of the study would be very much useful to the planners and policy makers of the study area to identify the irrationality in the existing production pattern and to suggest appropriate production plans for efficient utilization of their scarce resources resulting in increased net farm incomes and employment. The study also throws light on future potentialities of increasing net farm income and employment under different situations viz., with existing and recommended technology.