




Contents lists available at ScienceDirect

Journal of Environmental Management

journal homepage: <http://www.elsevier.com/locate/jenvman>

Review

Soil salinity under climate change: Challenges for sustainable agriculture and food security

Raj Mukhopadhyay^a, Binoy Sarkar^b, Hanuman Sahay Jat^{a,**}, Parbodh Chander Sharma^{a,*}, Nanthi S. Bolan^{c,d}^a ICAR- Central Soil Salinity Research Institute, Karnal, Haryana, 132001, India^b Lancaster Environment Centre, Lancaster University, Lancaster, LA1 4YQ, United Kingdom^c Global Centre for Environmental Remediation, University of Newcastle, Callaghan, NSW, 2308, Australia^d Cooperative Research Centre for High Performance Soils, Callaghan, NSW, 2308, Australia

ARTICLE INFO

Keywords:

Salt-affected soil
Climate change
Soil reclamation
Environmental quality
Farmers' livelihood
Sustainability

ABSTRACT

Soil salinity is one of the major and widespread challenges in the recent era that hinders global food security and environmental sustainability. Worsening the situation, the harmful impacts of climate change accelerate the development of soil salinity, potentially spreading the problem in the near future to currently unaffected regions. This paper aims to synthesise information from published literature about the extent, development mechanisms, and current mitigation strategies for tackling soil salinity, highlighting the opportunities and challenges under climate change situations. Mitigation approaches such as application of amendments, cultivation of tolerant genotypes, suitable irrigation, drainage and land use strategies, conservation agriculture, phytoremediation, and bioremediation techniques have successfully tackled the soil salinity issue, and offered associated benefits of soil carbon sequestration, and conservation and recycling of natural resources. These management practices further improve the socio-economic conditions of the rural farming community in salt-affected areas. We also discuss emerging reclamation strategies such as saline aquaculture integrated with sub surface drainage, tolerant microorganisms integrated with tolerant plant genotypes, integrated agro-farming systems that warrant future research attention to restore the agricultural sustainability and global food security under climate change scenarios.

1. Introduction

Soil salinity is one of the largest global challenges in the arid and semi-arid regions that severely affects agricultural production (El hasini et al., 2019). Soil salinity already covers 20% of total cultivated, and 33% of the irrigated agricultural lands worldwide (Srivastava and Kumar, 2015) and expected to increase at a faster rate than now by the year 2050 (Central Soil Salinity Research Institute (CSSRI), 2014). For example, the percentage of saline soils in Bangladesh had increased from less than 1% in 1990 to 33% in 2015 mainly due to sea water intrusion in the coastal areas resulting from excessive extraction of ground water sources (Rahman et al., 2018). The salt stress in soil is becoming prominent also due to other anthropogenic activities (e.g., over-application of groundwater and synthetic fertilizers) of the ever-increasing global population pressure (United Nations, 2011).

Recently, the alarming impact of climate change on the build-up of soil salinity has attracted widespread research attention. The rise in atmospheric greenhouse gases' (GHGs) concentrations and the consequent increase in air temperature and decline in relative humidity together with extreme events of rainfall are probable indicators of climate change that have huge impact on the pace of soil salinity development (IPCC, 2013; Haj-Amor and Bouri, 2019). The climate change could accelerate salt water intrusion into fertile soils due to sea level rise, and excess groundwater extraction in the dry regions of the world could also increase soil and groundwater salinity (Dasgupta et al., 2015). It is estimated that about 600 million people living in the coastal zones throughout the world could be affected by salinization (Wheeler, 2011; Dasgupta et al., 2015). Numerous studies reporting the impact of climate change on crop yield indicate positive (e.g., wheat yield increased with increased CO₂ concentration under optimal temperature) or negative (e.g., 3.8% drop in maize yield during 1980–2008) impacts,

* Corresponding author.

** Corresponding author.

E-mail addresses: hsjat_agron@yahoo.com (H.S. Jat), pcsharma.knl@gmail.com (P.C. Sharma).<https://doi.org/10.1016/j.jenvman.2020.111736>

Received 1 June 2020; Received in revised form 19 November 2020; Accepted 21 November 2020

0301-4797/© 2020 Elsevier Ltd. All rights reserved.

Abbreviations

CSSRI	Central Soil Salinity Research Institute	CL	Corn cropland
GHGs	Greenhouse gases	AF	Alfalfa forage
IPCC	Intergovernmental Panel on Climate Change	AG	Monoculture <i>Lyemus chinensis</i> grassland
FAO	Food and Agriculture Organization	AG + M	Monoculture <i>Lyemus chinensis</i> grassland for hay (mowing)
CSMPs	Climate smart management practices	MBC	Microbial biomass carbon
CA	Conservation agriculture	RG	Regrowth grassland
EC _e	Electrical conductivity of the saturated paste extract	ROM	Rice-okra-mentha
EC _{iw}	Electrical conductivity of irrigation water	PTR	Puddled transplanted rice
ESP	Exchangeable sodium percentage	US\$	United States Dollar
Mha	Million hectare	FP	Farm pond
ILO	International Labour Organization	RF	Deep furrow and high ridge
PTEs	Potentially toxic elements	PCF	Paddy cum fish
FGD	Flue gas desulphurization	CSCA	Climate smart conservation agriculture
RDN	Recommended dose of N	Sc	Scenario
RDF	Recommended dose of fertilizer	RWMS	Rice-wheat-mungbean system
OC	Organic carbon	MW	Maize-wheat
SOC	Soil organic carbon	ZT	Zero tillage
MSW	Municipal solid waste	R _m	Residue mulch
RW	Rice-wheat	PGPR	Plant growth promoting rhizobacteria
mmol	millimole concentration	ACC	1-aminocyclopropane-1-carboxylate deaminase
SAR	Sodium adsorption ratio	IAA	Indole acetic acid
DSR	Direct seeded rice	VAM	Vesicular arbuscular mycorrhizae
SSD	Sub surface drainage	EPS	Extra-cellular polymeric substances
		B: C	Benefit-cost ratio
		IRR	Internal rate of return

which could be balanced equally up to 2030 worldwide, but after that a clear dominance of the negative impact on crop yields will be visible (The Food and Agriculture Organization (FAO), 2017). Moreover, about 40 million people would be at risk due to malnourishment if the current pace of climate change continues (FAO, 2017). Therefore, climate change is likely to become one of the primary obstacles to sustainable agriculture and global food security (Corwin, 2020).

Rising air temperature, extreme events of rainfall, weather conditions (e.g., prolonged droughts and floods), changing soil fertility and health, and new pest infestations coupled with increasing salt-affected areas are major factors contributing to stagnant agricultural growth (Corwin, 2020). Climate-smart agriculture is considered a pragmatic approach to ensuring food security in the challenging environment (Jat et al., 2019). The climate smart management practices (CSMPs) include site-specific reclamation management strategies (e.g., amendments, irrigation and drainage), conservation agriculture (CA) and use of stress-tolerant genotypes. These practices may deliver co-benefits in the forms of reduced GHG emission, and enhance soil carbon sequestration and ecosystem services. Therefore, CSMPs are the need of the hour to tackle soil salinity under current and future climatic conditions.

Most of the literature is concentrated on desalination of brine water using reverse osmosis, and electro-remediation (Werber et al., 2017). These technologies have been proven effective for domestic water supply, but are expensive for irrigation of agricultural crops. Very limited literature is available concerning the complete management practice packages of soil salinity (Saifullah et al., 2018; Meena et al., 2019), especially under the varying climatic scenarios. A review paper portraying our current status of knowledge about the soil salinity management strategies under climate change scenarios aiming for food security is thus very important. In this article, we therefore aim to provide a holistic overview of global status of salt-affected soils, its relationship with climate change and food security along with various successful cost-effective climate smart reclamation strategies for salt-affected soils. We critically discuss about amendments, irrigation and drainage strategies, CA, land use patterns, bioremediation and phytoremediation approaches for the management of soil salinity. Further, we shed lights on the environmental and economic implications

of those strategies and suggest future research directions.

2. Methodology adopted for the review

A large number of published reports (n = 140) covering salt-affected soils were collected to make an initial assessment on salt-affected soil genesis, classification, extent of distribution, mechanisms of salt stress and mitigation strategies on the basis of the topic and hypothesis (Khan, 2019). We covered literature reviews based on information available from [sciencedirect.com](https://www.sciencedirect.com), [springer.com](https://www.springer.com), [wiley.com](https://www.wiley.com), FAO reports, CSSRI technical bulletins, Scopus and Google Scholar databases using relevant keywords such as salinity, sodicity, reclamation, salt tolerance, and climate sustainability. After systematic review and content analysis, we identified the key management practices and key challenges associated with certain reclamation strategies. Finally, we reached to the conclusions and future research recommendations focussing on the hypothesis and objective of this review.

3. Soil salinity and its global extent

Soil salinity is classified based on pH of saturated soil paste, electrical conductivity of saturated paste extract (EC_e) and exchangeable sodium percentage (ESP) (Richards, 1954). Salt-affected soil includes saline soil, sodic soil and saline-sodic soil. Saline soils have pH < 8.5, EC_e > 4 dS/m and ESP < 15 containing soluble salts of Cl⁻ and SO₄²⁻ of Na⁺, Ca²⁺, and Mg²⁺. On the contrary, sodic soils have pH > 8.5, EC_e < 4 dS/m and ESP > 15 containing CO₃²⁻ and HCO₃⁻ salts of Na⁺, Ca²⁺ and Mg²⁺ (Richards, 1954). Saline-sodic soil shows the characteristics of both saline and sodic soils. These soils are characterized by pH > 8.5, EC_e > 4 dS/m and ESP > 15 and contain a mixture of Cl⁻, SO₄²⁻, CO₃²⁻ and HCO₃⁻ salts of Na⁺, Ca²⁺, and Mg²⁺. Salt-affected soils are distributed in 954 million hectare (Mha) area of 120 countries of the world, and contribute to 7–8% productivity loss (Meena et al., 2019). Among these, Australia shares the highest area of salt-affected soils constituting more than 50% of the sodic soils worldwide (Table 1) (Shahid et al., 2018).

India is currently having 121 Mha of degraded land out of which 6.73 Mha area is covered by salt-affected soil (NAAS, 2012). In India, 2.96

Table 1
Continent-wise area distribution of salt-affected soils (in Mha).

Continents	Area under saline soil (Mha)	Area under sodic soil (Mha)	Total salt-affected area (Mha)	Sharing of the total global salt-affected area (%)	Reference
North America	6.19	9.56	15.75	1.69	Fageria et al. (2011)
Mexico and Central America	1.97	0.00	1.97	0.21	Fageria et al. (2011)
South America	69.41	59.57	128.98	13.84	Fageria et al. (2011)
Africa	53.49	26.95	80.44	8.63	Fageria et al. (2011)
Australia and New Zealand	17.36	339.97	357.33	38.35	Fageria et al. (2011)
Europe	7.8	22.9	30.7	3.31	Shahid et al. (2018)
Asia	194.7	121.9	316.5	33.97	Shahid et al. (2018)
Total	350.92	580.85	931.67	100	Shahid et al. (2018)

Mha is saline, and 3.77 Mha is sodic soil (Tripathi, 2011). Thus, India's 2% of the total geographic area is salt-affected, which poses a potential threat to the country's sustainable agriculture and food security.

4. Mechanisms of salt stress in plants

Soil salinity results in increased EC_e , poor soil structure and low soil water potential (ψ_w). The development of salt stress in plants could be

described in two ways: (i) osmotic phase, and (ii) ionic phase (Fig. 1) (Munns and Tester, 2008; Sirault et al., 2009). Osmotic phase takes place within few minutes of salt accumulation in root zone. Stomata closure, increase in leaf temperature and inhibited shoot elongation are the fundamental indicators of plants during osmotic phase because of low soil water potential and thick inner wall of the guard cells. Conversely, the ionic phase starts after few minutes to few hours in different cases of salt input which involves salt accumulation in shoots over a long period of time and leads to leaf senescence and premature abscission.

5. Interrelationship between soil salinity, climate change and food security

Climate change refers to long-term changes in weather conditions and climate systems. As a result, the global air temperature has increased by 1.5 °C above the pre-industrialization level, and the rise in CO₂ concentration in the atmosphere has gone up 20 $\mu\text{mol/mol}$ per decade since 2000, and now it has reached >400 $\mu\text{mol/mol}$ (Corwin, 2020). Consequently, following drastic changes have already been observed, which greatly influence the development of soil salinity (Corwin, 2020):

- Increase in the frequency of extreme weather conditions such as rise in air temperature, evaporation rate, excessive rainfall and heat stress.
- Global warming due to increased concentration of GHGs (e.g., CO₂, N₂O, CH₄) which trap the heat within the atmosphere.
- Spatial and temporal variability of rainfall distribution leads to changes in soil moisture contents.
- Increase in precipitation leads to soil erosion, groundwater recharge, infiltration and storage, whereas rise in temperature promotes the transpiration and moisture depletion from the soil profile.
- Rise in sea-level and sea water intrusion in the coastal areas limit their application for irrigation. It is projected that 130 million




Fig. 1. Plant's response to salinity stress (Munns and Tester, 2008; Sirault et al., 2009).

people will be inundated by the rise in sea level within 120 years (Chen and Mueller, 2018).

Besides, excessive use of mineral fertilizers and groundwater during the post green revolution era added neutral soluble salts to the soil, which in turn contributed to salinity build up.

The livelihood of 40–50% people in Asia is highly dependent on agricultural practices, while the corresponding value is 66% for Africa (ILO, 2007). Reclamation of salt-affected soils can potentially contribute to increased production of millions of tonnes of food grains worldwide. Thus, a complete package of climate smart technologies for reclamation of salt-affected soils is the need of the hour. Some of these reclamation approaches, which may reduce the area under salt-affected soil, helping to maintain agricultural sustainability and global food security, are discussed in the subsequent sections.

6. Soil salinity mitigation approaches for sustainable agriculture and food security

Useful techniques for reclaiming salt-affected soils in affected countries along with their major cropping systems are given in Table 2. In addition to various organic and inorganic amendments, applications of microorganisms, halophytes, tree species, land use pattern change, CA, and innovative irrigation and drainage strategies have been employed to reclaim salt-affected soils worldwide.

6.1. Amendments

Organic amendments such as biochar, compost of municipal solid wastes (MSW) and inorganic amendments that are rich in Ca (e.g., fly ash, gypsum, phosphogypsum), and zeolites have been used to reclaim sodic soils (Singh et al., 2018a; Mishra et al., 2019). Application of the


Table 2
Country-wise salt reclamation strategies with major cropping systems in the world.

Type of salt-related problem	Countries	Popular methods of reclamation	Major cropping system	Reference
Sodicity	Australia	Gypsum application	Wheat-pulses	Stevens and Pitt (2012)
Salinity and sodicity	India	Sub surface drainage, salt tolerant genotypes and gypsum application	Rice-wheat	CSSRI, 2014; Tiwari and Goel (2015)
Salinity and sodicity	China	Sub surface drainage, scraping out of salts and gypsum application	Rice-rice	Li et al. (2014)
Salinity and sodicity	United States & Mexico	Salt flushing, drainage, gypsum and organic amendment applications	Corn-wheat	MacMillan and Marciak (2001)
Salinity and sodicity	Pakistan	Sub surface drainage, amendments, salt-tolerant genotypes and gypsum application	Rice-wheat	Ahmad (2014)
Salinity	Egypt	Sub surface drainage	Cotton-wheat	El-Agha et al. (2019)
Salinity	Iraq	Sub surface drainage	Wheat-oil seeds/legume	Tiwari and Goel (2015)
Salinity	Iran	Drainage	Wheat-vegetables	Tiwari and Goel (2015)
Salinity	Israel	Drip irrigation	Wheat-vegetables	Girma and Abdulahi (2015)

above amendment materials improve the soil bulk density, aggregate stability, hydraulic conductivity, and lower down the pH, EC, ESP of salt-affected soils (Mishra et al., 2019; Sundha et al., 2020). In addition, improved soil biological properties (e.g., soil enzymatic activities, microbial population, and microbial biomass N and P contents) are observed due to beneficial effects of amendments. A schematic diagram is presented in Fig. 2 to elucidate the influences of various amendments on soil properties.

6.1.1. Gypsum

Gypsum is the most effective amendment for reclaiming sodic soil due to its wider availability, and substantial Ca^{2+} supply capacity. Ca^{2+} replaces Na^+ from the soil colloids, and leaches NaSO_4 deeper into the soil profile (Eq. (1)) (Singh et al., 2018a). A combination of gypsum and mineral langbeinite (rich in Mg^{2+} and K^+) resulted in significant reduction of soil exchangeable Na^+ and sodium adsorption ratio (SAR), and improved soil saturated hydraulic conductivity (Aydemir and Najjar, 2005). Naturally available mined gypsum is of poor quality, and its availability to agriculture is limited due to excess mining within the cement industry. Phosphogypsum, a by-product of phosphate fertilizer production, is used as an alternative to gypsum to minimize the use of mineral gypsum (Singh et al., 2018a). However, phosphogypsum may contain traces of potentially toxic elements (PTEs) (e.g., Cd), and its solubility is less than gypsum. Flue gas desulphurization (FGD) gypsum, a by-product of FGD process involving the capture of sulphur gases in coal-fired power stations, is also a rich source of Ca^{2+} , hence can be used to replace Na^+ with Ca^{2+} from soil exchange sites (Seshadri et al., 2013).



6.1.2. Compost

Composts such as green waste compost, green manure compost, and municipal solid waste compost were reported to increase soil salinity initially, but decreased it substantially in the later stage. El hasini et al. (2019) used a green waste compost (mixture of melon rind and olive pomace), sugarcane compost and gypsum to reclaim saline soils. They reported that the combined application of organic amendment (green waste compost) and gypsum (3.8 mg/g soil) increased the soil EC initially during 100 days due to the presence of dissolved salts in the compost and limited flushing. However, the EC was reduced to 2.80 from 16.65 dS/m after 120 days due to replacement of Na^+ with Ca^{2+} in soil exchange sites and solute leaching. Similarly, vermicompost at the rate of 10 t/ha in combination with 100% recommended dose of N (RDN) decreased the soil bulk density, pH, EC, ESP and soil solution Na^+ content of a degraded sodic soil by 2.0, 4.2, 26.5, 42.8, and 56.6%, respectively, and increased the soil organic carbon (SOC) content by 34.6% over control (Singh et al., 2019). The properties of degraded sodic soils could be improved considerably due to the decomposition of organic residues by enhanced microbial activity, microflora populations and displacement of excess Na^+ by Ca^{2+} in the soil exchange sites (Wang et al., 2014).

6.1.3. Municipal solid waste compost

Municipal solid waste (MSW) compost received wide acceptance for reclaiming sodic soils, and improved soil physicochemical properties when it was combined with gypsum, mineral fertilizers and other inorganic amendments (Sundha et al., 2020). Singh et al. (2018a) reported that application of gypsum (at the rate of 25% of the gypsum requirement) in combination with on-farm MSW compost at the rate of 10 t/ha reduced soil ESP and bulk density by 14 and 11%, respectively, and increased infiltration rate, SOC content and available N by 54, 10 and 13%, respectively, over recommended dose of gypsum. Similarly, a combined application of MSW (at the rate of 10 t/ha) along with 75% RDN improved the dehydrogenase activity in a sodic soil to the tune of




Fig. 2. Beneficial effects of different types of amendments on soil properties in salt-affected soils.

9.3–47.3% due to enhanced intra and extracellular enzyme secretions (Singh et al., 2019). Thus, to reclaim sodic soils, MSW compost could minimize the requirement of mineral gypsum. A similar study conducted by Meena et al. (2016) in saline soil showed that soil nutrient availability (N, P and K) was improved by the application of rice, wheat straw compost in combination with 50% recommended dose of fertilizer (RDF) instead of MSW compost. The N, P and K availability was increased by 14, 17 and 9%, respectively, after pearl millet harvest, likely due to the slow release of nutrients from the compost in the degraded soil. However, the combined use of chemical fertilizers and compost might decrease the organic P fraction in the soil, likely due to the increased microbial activities in the rhizosphere (Chang Hoon et al., 2004). In contrast, Meena et al. (2018) reported that a combined application of MSW compost (at the rate of 8 t/ha) and 50% RDF increased all fractions of P in soil under a mustard and pearl millet cropping system, likely due to the addition of organic P through the compost. However, MSW added some amounts of PTEs to the soil, especially lead (Pb^{2+}) and chromium (Cr^{6+}) depending upon the raw materials of the MSW compost. The measured PTEs levels were well below the critical level for inducing soil pollution (Meena et al., 2019). Therefore, a holistic improvement of salt-affected soil properties (physicochemical and biological) could be achieved via MSW applications with or without fertilizers, or other amendments. However, care should be taken for the selection and application rates of MSW compost so that it does not pose the risk of a secondary pollution in reclaimed soils.

6.1.4. Biochar

Biochar has become a popular amendment for soil carbon sequestration, soil health improvement and reclamation of salt-affected soils. Depending upon the nature of feedstock and preparation conditions, biochar can contain a substantial amount of N, P, K, and micronutrients (Sun et al., 2017; Purakayastha et al., 2019). Additionally, N-losses via NH_3 volatilization and denitrification can be substantially reduced by biochar application, due to NH_4 adsorption inside biochar pore spaces (Mandal et al., 2016). However, high application rates of biochar can increase the volatilization loss of N due to high pH of biochar applied to soil (Sun et al., 2017). The availability of P in sodic soil can be enhanced by decreasing the pH by 0.3 unit, blocking the clay adsorption sites by

dissolved organic carbon, and releasing organic acids for P mobilization in soil (Lashari et al., 2013). A wood-based biochar prepared from hardwood feedstock and pyrolyzed at 450 °C showed acidic pH (5.6) which could be used for reclamation of sodic soil (Shaheen et al., 2018). Acidic wood biochar (pH = 3.25; pyrolysis temperature 650 °C) might also be used to reduce the pH of sodic soil (Qi et al., 2018). In contrast, biochar application could increase the soil pH too, which would decrease P availability due to precipitation of insoluble P compounds (Xu et al., 2016). The K availability was reported to be increased by 44% after biochar application in salt-affected soils depending upon soil types and biochar properties (Lin et al., 2015). Therefore, the selection of biochar feedstock and pyrolysis temperature are important factors to be considered before applying this amendment to salt-affected soils. Soil physical properties (e.g., bulk density, porosity, and water holding capacity) were also improved considerably due to porous nature of biochar (Burrell et al., 2016). Amini et al. (2016) noticed significant effect of biochar (acidic vs alkaline) on soil physical properties such as saturated hydraulic conductivity and aggregate stability in a saline-sodic soil. The organic molecules present in biochar helped to bind polyvalent cations and clay particles for improving the aggregation of particles in degraded salt-affected soils. The soil structure and SOC content were found to be improved in salt-affected soil by biochar application with a simultaneous lowering of ESP (Amini et al., 2016). Similarly, soil EC_e decreased by 42% following a combined application of poultry manure and biochar (Lashari et al., 2015). It is inferred that acidic biochar could be a potential amendment for the reclamation of sodic soils.


6.1.5. Fly ash

Fly ash is a combustion product of coal industry having potential role in reclaiming sodic soils when applied jointly with gypsum and green manure (Mishra et al., 2019). Fly ash, applied in combination with manure and gypsum (25% gypsum requirement), showed significantly higher rice yield than the control treatment that consisted of the application of fly ash only (2.5% fly ash on mass basis) (Mishra et al., 2019). The beneficial properties of fly ash, such as high Ca^{2+} content, low bulk density, and the presence of substantial amounts of sesquioxides, improved the physicochemical properties of the degraded soil (Shirale et al., 2017). In contrast, fly ash application in soil under rice

crop did not significantly reduce the soil pH, but reduced 68.4% exchangeable Na^+ content of the soil (Lal et al., 2012). Fly ash derived from coal combustion is generally alkaline because of the addition of lime for capturing sulphur gases. However, FGD gypsum was reported to improve the N uptake of corn plants when applied together with N-fertilizers in the soil (Seshadri et al., 2013). Therefore, fly ash products could be a potential amendment for the reclamation of sodic soils due to their high Ca^{2+} supplying capacity.

6.1.6. Zeolites

Zeolites, a hydrated crystalline aluminosilicate of alkali and alkaline earth metals, can be used to reclaim saline soils. Zeolites such as clinoptilolite, erionite and heulandites are used as soil conditioners to improve soil properties and nutrient availability (Manjaiah et al., 2019). It was observed that Ca-rich clinoptilolite substantially improved the crop yield and quality of a saline soil due to adsorption of Na^+ and Cl^- in the mineral cavities (Noori et al., 2006). Similarly, 5% zeolite application to a saline soil increased Ca^{2+} concentration, and micronutrient Fe^{2+} and Mn^{2+} by 19 and 10%, respectively (Al-Busaidi et al., 2008). The mechanisms governing the salt removal process by zeolites are mainly ion exchange, adsorption, and salt storage (Wen et al., 2018). The possible ion exchange reaction of zeolites in salt-affected soil is shown in Eq. (2).



6.2. Irrigation and drainage strategies

Saline groundwater in the arid and semi-arid regions of the world has become a major challenge for water management to ensure agricultural sustainability (Yao et al., 2012). Almost 43% of world's irrigated area is groundwater dependent (Minhas et al., 2019), out of which India (39 Mha) and China (19 Mha) share the maximum area under groundwater irrigation. Therefore, strategic water management techniques are required to tackle saline groundwater during crop production.

6.2.1. Irrigation techniques

South-Asian countries mainly focus on rice-wheat (RW) system to ensure food security, and this cropping system is resource intensive, viz. it entails high requirement of irrigation water (200–250 cm/yr), synthetic fertilizers, energy, and labour. A constant depletion of groundwater aquifers and salinity build up are becoming pronounced in South Asia. Irrigation with saline water was reported to increase salinity up to 12.2 dS/m, and SAR up to 20 (Yadav et al., 2007). Consequently, significant negative changes occurred in soil properties including reduced saturated hydraulic conductivity, aggregate stability and increased dispersion and run-off (Mandal et al., 2008). Hence, judicious application of irrigation water along with smart crop management practices is an essential approach to avoid salinity build up and tackle water scarcity in dry land areas (Minhas et al., 2020a). Saline and good quality available water have been used in a mixture to avoid the water salinity during pearl millet production and to compare yield of two varieties AVKB-19 and ICMV-15111 under saline soil (Makrana et al., 2019). The AVKB-19 showed 16.26% higher grain yield than ICMV-15111 when good quality water was mixed with saline water for irrigation. Bed planting of crops, straw mulching, and micro irrigation (e.g., sprinkler, drip (surface and subsurface) irrigation) are known to save irrigation water, and improve N-use efficiency and grain yield. For example, drip irrigation with residue retention and raised bed planting in maize and wheat showed 13.7 and 23.1% higher yield, respectively, than furrow irrigation system without residue retention (Sandhu et al., 2019). The drip system (with residue retention) saved 88 and 168 mm of water, and increased water productivity by 66 and 259%, respectively, in CA-based wheat and

maize compared to the conventional irrigation system. Moreover, sub-surface drip irrigation in combination with CA saved 48–53 and 42–53% of the irrigation water in rice and wheat, respectively, and saved overall 20% N input (Sidhu et al., 2019). Apart from drip system, irrigation techniques using sprinkler and low energy water application device saved 30–40% water compared to surface irrigation, and recorded 4.4 t/ha rice yield which was at par with maximum yield with surface irrigation in a sodic soil (Singh et al., 2018b). Kumar et al. (2019a) proposed novel matrix potential based irrigation strategies for direct seeded rice (DSR) which could save substantial amount of water, and sustain the productivity of salt tolerant basmati rice (CSR30). They suggested that irrigations at or below -30 kPa (field capacity) during initial 90 days of rice growth, and at -15 kPa during rest of the growth period exhibited similar yield under traditional irrigation method. Thus, the use of saline water in conjugation with fresh water, sub-surface drip irrigation, and DSR techniques might effectively reduce the salinity effects, and tackle water scarcity in semi-arid and arid regions.

6.2.2. Drainage strategies and groundwater recharge

Agricultural drainage is important to remove excess water due to high precipitation, eliminate dissolved soluble salts from the soil profile, and maintain the groundwater table. However, due to lack of care, waterlogged salinity remains a severe problem in many productive areas of Australia, Middle East, United States, and Asia (Emadodin et al., 2012). Sub-surface drainage (SSD) was identified as one of the most important techniques to remove salts from 1.5 m depth (Nijland et al., 2005). Two types of SSD can be useful for the reclamation of saline soils: (i) horizontal sub-surface drainage which works up to 1.5–2.0 m depth of root zone, and involves a network of drain consisted of the main drain, lateral drains and collectors (Nijland et al., 2005), and (ii) vertical sub-surface drainage which is related to pumping of excess water by tube well (Bos, 2001). Srinivasulu et al. (2005) reported SSD installation (pipe drain) in 8 ha saline waterlogged area of Prakasam district of India (water table depth 0–3.7 m; EC_e : 1.3–18.6 dS/m) with 30 and 60 m drain spacing. Around 0.20–0.35 m lowering of water table depth, and 50.4 tonnes of salt accumulation in the pipes occurred during 1999–2002. Yields were increased by 50–100% in most of the crops due to installation of SSD, which is really a profitable solution to the farmers of salt-affected areas (Kamra, 2013). Bhattacharya (2007) reported that investments on surface drainage too are economically viable with yield benefits ranging from 20 to 28% in sugarcane, 20–25% in paddy, 32% in gram, and 50% in Indian bean.

Intensively irrigated agriculture suffers severely from long-term increase of salinity in the groundwater derived from irrigated permeable soils in arid and semi-arid regions (Foster et al., 2018). Groundwater recharge through preparation of artificial recharge structures using rainwater was found useful to minimize the salt load in surface water and groundwater. Groundwater salinity of semi-arid area of Karnal district, India, was reduced to 1.33 from 1.36 dS/m within one year while alkalinity reduced from 6.63 to 1.33 meq/L with 3.16 m rise in groundwater table due to groundwater recharge events (Narjary et al., 2014). Similarly, 1.26 and 0.65 m rise in water table during monsoon through an artificial recharge structure was observed in Nirmana and Kutba village of Uttar Pradesh, India, with a reduced salinity level of the groundwater (e.g., EC declined from 0.90 to 0.24 dS/m in Kutba village) (Kumar et al., 2019b). Furthermore, the alkalinity of groundwater was reduced by 2–3 meq/L in Kaithal district of Haryana, India, which minimized crop damage from 35 to 40% in open field to 5–15% under groundwater recharge structure with a benefit to cost ratio of 1.93 (Kumar et al., 2020). Horriche and Benabdallah (2020) reported that an artificial recharge structure with 1500 m^3 /day recharge rate minimized salt load by 5.7 g/L. Hence, groundwater recharge in saline areas is one of the feasible solutions for increasing the level of groundwater table and improving the groundwater quality.

6.3. Salt tolerant genotypes

Salt tolerant crops are an important tool for sustaining productivity in salt-affected regions. Plant salt tolerance has been demonstrated through specificity in ion accumulation and better partitioning of accumulated ions within plant cells and tissues. For example, wheat showed salt tolerance via: (i) salt exclusion, (ii) osmotic tolerance, and (iii) tissue tolerance (Munns et al., 2016). In addition, the salinity tolerance of wheat showed dependence on forms of N-fertilizers (NH_4^+ preferred) indicating that $\text{NO}_3\text{-N}$ could be harmful (Elgharably et al., 2010). In case of barley, salt exclusion and osmotic tolerance were the main operative mechanisms both under hydroponic and saline soil growth conditions (Tavakkoli et al., 2012). Krishnamurthy et al. (2016) evaluated 131 rice accessions at normal (1.2 dS/m) and highly saline (10 dS/m) irrigation water, and found that root and shoot lengths were decreased by 52 and 50%, respectively, due to high Na^+/K^+ ratio in plants. They identified three accessions namely, IC 545004, IC 545486 and IC 545215 which were suitable as parent donors under saline conditions. Ravikiran et al. (2017) experimented with 192 genotypes of rice at normal (1.2 dS/m) and high salinity (12 dS/m) hydroponic conditions. They concluded that CST 7-1 and Arvattelu genotypes could be novel sources of seedling stage salinity tolerance among all the experimental genotypes. A comparison study between rice varieties CSR10 and MI48 was conducted, and CSR10 expressed lower Na^+/K^+ ratio in shoots than MI48 (Singh et al., 2018c). For mustard, a set of 97 salt tolerant genotypes were planted in highly saline soil (EC_e : 10.7 dS/m). The erucic acid content in mustard was increased under salinity by 12.2%, while oil, protein and crude fibre contents were decreased by 5.78, 29.31, and 20.45% (Singh et al., 2014). Tomato genotypes were also evaluated in salt solution (0, 1.0 and 3.0% NaCl salt concentration in Hoagland solution). Important traits of tomato genotypes such as % germination, and root shoot dry weight responded negatively towards high salinity for most of the tomato genotypes except Sel-7 and Arka Vikas (Singh et al., 2012). Sugarcane was also found salt tolerant (varieties-Co 6806, Co 7717 and Co 8208) in coastal saline areas of India, and but experienced major yield loss (~40%) under massive salt water intrusion in coastal areas (Balasundaram, 2004).

6.4. Land use management

Conversion of barren land to crop lands naturally improve the soil nutrients status and organic carbon inputs (Yu et al., 2018). Therefore, changes in land use systems of salt-affected soil through cropping systems, agroforestry, and fruit crops would improve the soil quality. Five land use systems comprised of corn cropland (CL), alfalfa perennial forage (AF), monoculture *Lyemus chinensis* grassland (AG), monoculture *L. chinensis* grassland for hay (Mowing) (AG + M) and successional regrowth grassland (RG) were evaluated in salt-affected soil of north-eastern China (Yu et al., 2019), and it was revealed that SOC, total N, total P and total K contents of surface soils were increased by 40.42, 17.66, 15.71 and 11.5%, respectively, due to addition of organic litter inputs from forage crops. Rice based cropping system showed promising impact for improving soil physicochemical properties and carbon content in the salt-affected canal command area of Indo-Gangetic Plain (Bhardwaj et al., 2019). Results suggested that rice-okra-mentha (ROM) and RW systems displayed a decreasing trend of ESP to the tune of 37 and 35.5%, respectively, and an increasing trend of carbon stock (4–70%) compared to barren sodic land due to addition of organic matter. Likewise, *Jatropha* (*Jatropha curcas* L.) cultivation as an intercrop between sweet basmati rice and *Matricaria* was effective in increasing soil microbial biomass carbon (MBC) (+24.68% over control-no intercrop), microbial activity and improving soil properties in a degraded sodic soil of northern India. The above practice received 5288.4 US\$/ha as economic return via *Jatropha* intercropping with the sweet basmati rice-*Matricaria* cropping system because of high yield and market value of sweet basmati rice and *Matricaria* flowers (Singh et al.,

2016). Another study on the influence of different land uses (sorghum, paddy, forest, wetland, wasteland and meadow) on microbial community structure in saline-sodic soil suggested that bacterial abundance was maximum under wetland (1.03×10^9 copies/g dry weight of soil), whereas fungal population was maximum (5.83×10^6 copies/g dry weight of soil) under forest ecosystem, due to drying and wetting phenomenon in the former (Feng et al., 2019). In case of GHG emission, emission of CO_2 decreased (lowest in bare land) with increasing salinity, which might be due to the reduction of activity of heterotrophic microorganisms. The N_2O production increased with increasing salinity under *Tamarix chinensis* and *Phragmites australis* ecosystems due to increased ion concentrations, less solubility of N_2O , and low N_2O reductase activity (Zhang et al., 2018). However, N application increased the CO_2 emission under *Suaeda salsa* and *P. australis* ecosystems, while N_2O emission was large under vegetative cover because of increased microbial activity (denitrifying bacteria) (Zhang et al., 2019).

Crop establishment methods such as rapeseed grown after non-puddled transplanted rice ((non-PTR) and dry-seeded rice) provided 25–44% higher rice yield than rapeseed grown afterpuddled transplanted rice (PTR), while yields of maize were 8–13% higher when grown after either dry-seeded rice or non-PTR than after PTR in coastal salt-affected areas of West Bengal, India (Sarangi et al., 2019). In addition, maize was found more profitable (US\$301–405/ha) than rapeseed (US\$ 5–113/ha) in coastal salt-affected soils of India. Impacts of different land shaping techniques such as farm pond (FP), deep furrow and high ridge (RF), and paddy cum fish (PCF) systems were studied for rainwater harvesting to improve the productivity of coastal saline soils of Sundarbans, India (Mandal et al., 2019). These areas receive 2.7 times higher annual rainfall than crop evapotranspiration. The estimated runoff was 19.5, 29.1 and 27.75% of the annual rainfall in FP, RF and PCF systems, respectively, whereas in rice-fallow system it was 34.6% of the annual rainfall. Additionally, the calculated water footprints were much higher in the rice-fallow and rice-rice ecosystems than the individual land shaping system, indicating an increased cropping intensity and high net farm income, during the summer and reduced salinity and waterlogging during the rainy season.

6.5. Climate smart conservation agriculture

Climate smart conservation agriculture (CSCA) works on the principles of conservation agriculture along with precise input management practices, i.e., nutrient, water, genotypes, labour, and pesticides. Conservation agriculture maintains three basic principles: (i) residue retention, (ii) minimum tillage, and (iii) crop diversification in order to enhance the productivity of agri-food systems through better adaptation to climate change (Aryal et al., 2016). A short description of CSCA based agro-ecosystem is presented in Fig. 3.

The CSCA is becoming a potential mitigation strategy for soil salinity under climate change in Asian countries, especially in north western parts of India. Jat et al. (2018) evaluated four scenarios (Sc), namely conventional RW cropping system (Sc1), partial CA-based rice-wheat-mungbean system (RWMS) (Sc2), CA-based RWMS (Sc3), and CA-based maize-wheat-mungbean (Sc4) system to assess soil properties and carbon storage under CA-based treatments in a reclaimed sodic soil. Sc2 indicated remarkably lower soil bulk density (1.52 Mg/m^3) compared to the conventional system (1.67 Mg/m^3), while Sc3 and Sc4 showed greater organic carbon (OC) content (0.75 and 0.77% OC content under Sc3 and Sc4, respectively), and N availability (33 and 68% higher under Sc3 and Sc4, respectively) compared to Sc1. The SOC pools at surface soil showed the following order: Sc4 > Sc3 > Sc2 > Sc1 (Jat et al., 2019). Around 34% higher soil respiration was noticed under different CA-based scenarios. The predominant phyla of soil bacteria in all scenarios were *Proteobacteria*, *Acidobacteria*, *Actinobacteria*, and *Bacteroidetes*, accounting for >70% of the identified phyla. The bacterial diversity was prominent under all CA-based maize-wheat-mungbean cropping systems (Choudhary et al., 2018a). Besides, lignocellulose




Fig. 3. Conceptual framework of agro-ecosystems based on climate smart conservation agriculture (adapted from Lal, 2015).

degrading fungal species such as *Aspergillus flavus*, *Aspaergillus terreus*, *Penicillium pinophilum* and *Alternaria alternate* were present for residue decomposition (Choudhary et al., 2016). In this connection, maize-wheat (MW) system with zero tillage (ZT) and residue mulch (R_m) (MW/ZT + R_m) recorded 208, 263, 210 and 48% improvement in MBC, microbial biomass N, dehydrogenase activity, and alkaline phosphatase activity, while RW system under RW/ZT + R_m showed 83, 81, 44 and 13% improvement, respectively, as compared to RW/conventional tillage without residue mulch (Choudhary et al., 2018b, 2018c). Therefore, the CSCA practices hold future potential to tackle soil salinity under climate change scenarios.

6.6. Phytoremediation

Phytoremediation is a well-known technique that uses plant species to accumulate salts in order to reduce their soil concentration. Mainly three kinds of approaches of phytoremediation are available to minimize soil salinity: (i) agroforestry, (ii) biodrainage, and (iii) halophytic plants.

6.6.1. Agroforestry

Salt-tolerant fruits, fodder, and tree species could survive under low water requiring saline irrigation in arid and semi-arid regions (Minhas et al., 2020b). Dagar et al. (2015a) used salt-tolerant fruit crops, namely *Carissa carandas*, *Embllica officinalis*, and *Aegle marmelos* along with companion crops such as *Hordeum vulgare*, *Brassica juncea*, *Cyamopsis tetragonoloba*, and *Pennisetum typhoides* in inter-row spaces, irrigated with (EC_{iw} : low = 4–5; high = 8.5–10.0 dS/m) saline water. Results suggested that *C. carandas* with *P. typhoides* and *H. vulgare* performed the best with saline water irrigation in sandy calcareous soil. Recently, Dagar and Yadav (2017) reported that fruit crops such as gooseberry (*E. officinalis*), ber (*Zizyphus mauritiana*) and sapota (*Achras zapota*) could tolerate ESP up to 60 in sodic soil. Similarly, among agroforestry trees, *Prosopis juliflora* was considered suitable (pH > 10) for sodic soils

followed by *Tamarix articulata* and *Acacia nilotica* (Dagar et al., 2001). However, *P. juliflora*, *T. articulata*, and *Salvadora persica* could be raised successfully in saline soils at up to EC_e 30–40 dS/m (Dagar and Yadav, 2017). In another experiment, alfalfa (*Medicago sativa*) was evaluated for five years, and it reduced salinity considerably and added high C and N in soil to improve the soil quality across the profile (Cao et al., 2012). Biomass mulching, root exclusion and salt removal through alfalfa shoots could be the mechanisms for reduced soil salinity. Medicinal aromatic plants like *Glycyrrhiza glabra* was also found suitable in alkali soil with high economic values in terms of net returns (2.4–6.1 t/ha forage, and 6.0–7.9 t/ha root biomass, per annum) (Dagar et al., 2015b).

6.6.2. Biodrainage

Biodrainage is a technique of using tree species that reduce the water table by transpiration, mainly in waterlogged areas. Ram et al. (2011) evaluated the performance of clonal Eucalyptus (*Eucalyptus tereticornis*) planted on field boundaries in a waterlogged soil, and found that groundwater table was lowered down by 2 m after 5 years. Till date, Eucalyptus has been the most efficient species for lowering down of water table in canal command areas (Dagar et al., 2016). Dagar et al. (2016) revealed that Eucalyptus lowered water table by 43.0 cm in 1 m × 1 m, 38.5 cm in 1 m × 2 m, and 31.5 cm in 1 m × 3 m spacings during the fourth year of plantation compared with no tree plantation. In addition, a recent study showed that Eucalyptus stored carbon to the tune of 21.2–22.8 Mg/ha in surface soil under agri-silviculture system (i. e., intercropping of timber and fuel wood species, and/or fruit and other useful trees with vegetables and other crops in a common space, at the same time) (Kumar et al., 2019c). However, it was recommended that high water demanding trees such as Eucalyptus (*E. tereticornis*), Populus (*Populus deltoides*) along with Panicum (*Panicum maximum*), and Leptochloa (*Leptochloa fusca*) could manage the waterlogged saline soil (Dagar, 2014).

6.6.3. Halophytes

Halophytes refer to the plants which can grow and adapt under saline conditions. Halophytes of the genera such as *Pandanus*, *Pongamia*, *Panicum*, *Plantago*, *Porterasia*, *Prosopis*, *Rhizophora*, *Salicornia*, and *Salvadora* are popular for the reclamation of salt-affected soils. Halophytes are classified according to their salt tolerance or exclusion, which is shown in Table 3 (Grigore and Toma, 2017).

Three key mechanisms of salt tolerance by the halophytes include: (i) avoidance, (ii) evasion, and (iii) tolerance (Batanouny, 1993; Hayat et al., 2019). Cultivation of *Atriplex halimus* decreased EC from 39.2 to 26.5 dS/m of saline-sodic soil, and from 6.2 to 4.9 dS/m of saline soil (Abdul-Kareem and Nazzal, 2013). Ravindran et al. (2007) reported that the halophytes (*Suaeda maritima* and *Sesuvium portulacastrum*) reduced EC_e and SAR of a saline soil (EC_e > 4 dS/m and SAR >13) to a level of normal soil due to accumulation of large salt amounts in the plant tissues. Zahran and Abdel Wahid (1982) reported that *Juncus rigidus* reduced the soil EC from 33 to 22 dS/m during its growth cycle. In addition, mangroves having aerial roots were found useful and economic in coastal saline soil reclamation, protecting the coastal areas from tide and providing a common habitat for saline aquaculture with shrimps (Dagar and Yadav, 2017). Popular mangrove species include *Aegialitis rotundifolia*, *Aegiceras corniculatum*, *Avicennia marina*, *Avicennia officinalis*, and *Bruguera gymnorrhiza* which are habitat of Sundarban, and Andaman Islands in India (Dagar and Yadav, 2017). Therefore, halophytes offer an economic and ecological solutions for management of waterlogged saline soils.

6.7. Bioremediation

Bioremediation involves the use of various microorganisms or microbial consortium to reclaim salt-affected soil. The soil microbes include plant growth promoting rhizobacteria (PGPR), bacteria, mycorrhiza, and cyanobacteria able to reclaim salt-affected soil by producing various hormones and beneficial substances that enhance soil quality and plant growth.

6.7.1. Plant growth promoting rhizobacteria

Various PGPR showed their impact on salt-affected soil remediation by improving plant growth. The main mechanisms include production of 1-aminocyclopropane-1-carboxylate (ACC) deaminase, indole acetic acid (IAA), and exopolysaccharides secretion to enhance crop growth (Singh, 2015). A list of PGPR along with their mechanisms on agricultural crops is given in Table 4 (modified from Choudhary et al., 2019).

6.7.2. Salt tolerant bacteria

Salt-tolerant bacteria generally show high requirement of salts, and exist in highly saline environment to regulate high osmotic pressure. The genera include *Ammoniphilus*, *Arthrobacter*, *Azospirillum*, *Bacillus*,

Table 3

Classification of halophyte plants used for the reclamation of salt-affected soils.

Class	Definition	Example
Euhalophytes	Plants that can accumulate salts and grow in saline condition having low respiration rate and salt permeable cell cytoplasm. They show succulence due to accumulation of salts and high osmotic potential.	<i>Salicornia europaea</i> , <i>S. maritima</i> , <i>Salosa soda</i> and <i>Halocnemum strobilaceum</i>
Cryno-halophytes	Plants that can grow in low to high salinity and excrete salts through salt glands in the leaves.	<i>Statice gmelini</i> and <i>Tamarix gallica</i>
Glyco-halophytes	Plants that have no capacity to salt permeability through cytoplasm but have limited capacity to grow in salts. These are mainly freshwater plants.	<i>Artemisia maritima</i> .

Table 4

List of plant growth promoting rhizobacteria and their mechanisms on agricultural crops in salt-affected soils.

Microorganisms	Associated mechanisms	Crops	Reference
<i>Enterobacter sp.</i>	ACC deaminase	Okra	Habib et al. (2016)
<i>Pseudomonas sp.</i>	ACC deaminase	Tomato	Win et al. (2018)
<i>Pseudomonas sp.</i>	ACC deaminase	Wheat	Nadeem et al. (2013)
<i>Klebsiella sp.</i>	IAA, organic acids	Oat	Sapre et al. (2018)
<i>Azotobacter sp.</i>	IAA, N-fixation	Maize	Rojas-Tapias et al. (2012)
<i>Bacillus amyloliquefaciens</i>	Gibberalic acid, abscisic acid	Soybean	Kim et al. (2017)
<i>Enterobacter cloacae</i>	IAA, ACC deaminase	Canola	Li et al. (2017)
<i>Enterobacter sp.</i>	ACC deaminase	Wheat	Sarkar et al. (2018)
<i>Pseudomonas fluorescens</i> and <i>Azospirillum brasilense</i>	Phosphate solubilisation, auxin production	Wheat	Kadmiri et al. (2018)
<i>Pseudomonas fluorescens</i> and <i>Pseudomonas putida</i>	ACC deaminase	Canola	Jalili et al. (2009)
<i>Pseudomonas fluorescens</i>	IAA, siderophore	Black gram	Yasin et al. (2018)
<i>Pseudomonas fluorescens</i>	IAA	Maize	Zerrouk et al. (2016)
<i>Alcaligenes sp.</i> and <i>Bacillus sp.</i>	ACC deaminase	Rice	Bal et al. (2013)
<i>Enterobacter aerogenes</i> and <i>Pseudomonas aeruginosa</i>	ACC deaminase	Alfalfa	Liu et al. (2018)
<i>Bacillales</i>	ACC deaminase, siderophore production	Rice	Zhang et al. (2018)

Brevibacillus, and *Brevibacterium* which produce IAA, gibberellic acid, and other organic acids that can solubilize and transform nutrients present in soils. Some salt-tolerant endophytes have been reported to show similar mechanisms to PGPR for salt tolerance (Thijs et al., 2014). Research revealed that *Bacillus foraminis* and *Bacillus gibsonii* tolerated up to 7.5% NaCl (Arora et al., 2014). In addition, *Pseudomonas fluorescens*, and *Bacillus subtilis* were found successful in NH₃ production, while phosphate solubilisation was significantly higher under isolated *Acinetobacter baumannii* and *P. fluorescens* (Arora and Vanza, 2017). However, future researches should focus on salt-tolerant bacteria and their application to agricultural fields to avail success in this potential area.

6.7.3. Mycorrhiza

Mycorrhiza is the symbiotic association between roots of higher plants and fungi. Mycorrhiza is known for mobilization and solubilisation of nutrients even under saline environment. Reports are available on the beneficial impact of mycorrhiza such as improved mobility and availability of nutrients (Zn²⁺, Cu²⁺, P) in soils (Chang et al., 2018). Vesicular arbuscular mycorrhizae (VAM) generally helps in solubilisation of phosphate and P supply to plant roots in salt-affected soils because phosphate remains in precipitated forms due to presence of Ca²⁺- and Mg²⁺-based carbonate salts (Zhu et al., 2016). In addition, high K⁺/Na⁺ ratio maintenance by VAM fungi indicated their salt-tolerant mechanisms in saline soils (Wu et al., 2005). Contrasting reports are also available where VAM colonization in wheat was reported to be decreased by increasing salt concentration in the medium (Zhu et al., 2016). Other mechanism related to mycorrhizal action to salt-tolerance is the control of abscisic acid accumulation under osmotic stress (Auge et al., 2015). However, further investigations are needed to understand the molecular mechanisms of mycorrhiza under salt stress conditions.

6.7.4. Cyanobacteria

Cyanobacteria are gram negative, prokaryote, autotrophic, and blue-green bacteria. They can survive in extreme environments including under highly saline condition, and improve soil quality (Rossi et al., 2017). Mostly cyanobacteria were utilized as biofertilizer. It is a challenge to remove salts from soil using cyanobacteria in terms of quantity, but the use of cyanobacteria in association with salt-tolerant plants helps to increase the quantity of removed salts (Jesus et al., 2015). The main mechanisms employed by cyanobacteria are N-fixation, high biomass production, and extra-cellular polymeric substances (EPS) production that help the microorganisms to survive under salt stress conditions. The cyanobacterial genera which were used in different pot and field studies include *Anabaena*, *Nostoc*, *Calothrix*, and *Spirulina* (Li et al., 2019). Soil pH and EC were decreased (Singh and Singh, 2015), while soil fertility and soil enzyme activities were improved using *Nostoc ellipsosporum* HH-205 and *Nostoc punctiforme* HH-206 under saline soil of India (Nisha et al., 2018). Additionally, *Anabaena laxa* RPAN8 showed 21-times higher acetylene reducing activity under salt stress condition, which was an indication of N-fixation (Babu et al., 2015). Similarly, the intracellular trehalose content of *Anabaena fertilissima* increased significantly under 250 mmol NaCl concentration (Swapnil and Rai, 2018). Besides, the IAA concentrations were found to be 20.05 and 27.17 µg/mL under 250 mmol NaCl in *Nostoc carneum* TUBT04 and *Nostoc commune* TUBT05, respectively (Chittapun et al., 2018). Gibberellins and cytokinins were also identified in *Nostoc kihlmani* and *Anabaena cylindrical*. However, potential application of cyanobacteria at field levels should be explored in integration with manure, biochar, and salt tolerant plants to evolve a green remediation technology for salt-affected soils.

7. Socio-economic and environmental impact

The monetary and environmental loss due to salinity, and the implication of proper management practices in the improvement of farmers' livelihood requires special attention, as described in the following sub-sections.

7.1. Socio-economic impact

Soil salinity caused crop growth and yield reductions which led to 27.3 billion US\$ economic loss globally, and 1.2 billion US\$ in India alone (Qadir et al., 2014). Among the mitigation technologies, SSD offered positive impact on farmers' socio-economic conditions. Manually installed SSD provided a benefit-cost (B: C) ratio of 1.26 with viable internal net return of 13.3% (Kamra et al., 2019). Reports revealed an approximate 40–50% yield increase in soybean-wheat cropping system compared to control (without SSD), which resulted in a B: C ratio of 2.6, and an internal rate of return (IRR) of 28% in Rajasthan, India (Sewa et al., 2000). Even after 20 years of SSD installation, 15–20% additional economic benefit was obtained by the farmers compared to the sites without SSD (Tejawat, 2015). Furthermore, the socio-economic analysis of SSD indicated B: C of 1.5, IRR of 20%, and employment generation of 128 man-days/ha every year (Datta et al., 2000; Sewa et al., 2000). Salt-tolerant genotypes played an outstanding role in the development of economy. Sharma (2010) reported that salt-tolerant rice, wheat and mustard varieties developed by CSSRI showed an estimated value of total annual produce around 4384, 46 and 69 million US\$, respectively (calculated on the basis of the minimum support price rates of these crops in India for the years 2009 & 2010). The rice yield was increased to the tune of 1 t/ha where there was an existing practice of cultivating marginal salt-tolerant varieties (Sharma, 2010). The use of salt-tolerant genotypes reduced gypsum application by 10–15 t/ha for the reclamation of sodic soils (Mandal et al., 2018). Gypsum application to sodic soil gave positive impact with net value of 3771 US\$/ha, B: C ratio of 2.47, IRR of 67% (Tripathi, 2011). The agricultural income generated from the reclaimed sodic soils was around 3410 million US\$ in India (Mandal

et al., 2018), which provided an opportunity of 2.8 million man days of jobs per annum. In coastal saline area of India, implementation of different land-shaping models along with best natural management practices resulted in enhanced farmers' income, allowing net returns from around 76 US\$/ha for wet rice to 1935 US\$/ha for wet rice-fish-vegetables (Mandal et al., 2018). Among agroforestry systems, *G. glabra*, a high value medicinal plant, produced 8000–10000 US\$/ha in terms of root and biomass of high medicinal values (Dagar et al., 2015b).

7.2. Environmental impact

Climate change has many adverse environmental impacts on soil and groundwater salinity. Therefore, climate smart salinity management practices could provide different paths to alleviate salinity and its environmental impacts. The emission of CH₄ from salt-affected soils could be reduced by about 28–68% through biochar application, while the N₂O emission could be reduced by about 50% through manure application (Begum et al., 2019; Nguyen et al., 2020). Application of MSW compost to soils for the reclamation of salinity and sodicity might be harmful as the amendment might contain multiple PTEs (e.g., Cd²⁺, Cr⁶⁺, Pb²⁺, Zn²⁺, and Cu²⁺) (Meena et al., 2019). Therefore, MSW compost should be checked before soil application. Studies on groundwater salinity and toxicology due to climate change and salt leaching during the reclamation process also need research priorities.

8. Future research directions

Following are the thrust research areas that need worldwide future attention to sustain productivity of salt-affected soils under climate change conditions:

- (i) Climate change impact on root zone salinity, solute movement at different depths, and the impact on soil properties under various agro-ecological regions of the world need immediate research attention using hydro-salinity modelling approaches (Corwin, 2020).
- (ii) Sub-surface drainage integrated with inland saline aquaculture is a new area of research to achieve sustainable management of salt-affected soils. Experiments in this field should be conducted considering the salt-load, soil nutrients, and carbon loss. Their environmental impacts should then be assessed over long period (Castellano et al., 2019).
- (iii) Halophytes have worldwide potential for salt tolerance (Hayat et al., 2019). The genes of halophytic plants could be transferred to crop genotypes in order to improve the salt-tolerance capacity of crops, especially for coastal saline areas.
- (iv) Naturally available gypsum is scarce and of poor quality. Therefore, development of suitable alternative amendments to gypsum is the need of the hour.
- (v) Functionalized biochar has enormous potential to manage sodic soils. Investigations on the impact of functionalized biochar on CaCO₃ dissolution in sodic soils, and on the associated soil properties should be pursued in the near future.
- (vi) Salt-tolerant plant genotypes can cope up with soil salinity (Genc et al., 2019). Microbiological interventions with cyanobacteria on salt-tolerant plant roots and their integration with various organic amendments are a potential area of salt remediation warranting further research development (Rossi et al., 2017).
- (vii) Integration of land shaping techniques with multi-enterprise agro-farming system is already performing well in coastal saline areas of West Bengal, India, and needs future research attention in coastal saline areas worldwide.

9. Conclusions

Climate change could accelerate the pace of soil salinity development all over the world and primarily in the arid and semi-arid regions. The different mitigation technologies, such as amendments (gypsum, biochar, MSW, zeolites), salt-tolerant genotypes, sub-surface drainage in waterlogged saline areas, micro-irrigation techniques (drip system), climate smart conservation agriculture, land shaping techniques, agroforestry, and microorganisms, have the capacity to reclaim salt-affected soils. These technologies can improve the physicochemical (pH, EC, bulk density, available soil nutrients) and biological properties (enzyme activities, MBC) of salt-affected soils worldwide, allowing to achieve improved soil health and crop productivity. The mitigation approaches are environmentally sound, and socio-economically viable to the farmers across various agro-ecological regions, and can be adopted according to the bio-physical and socio-economic conditions of the farming communities. Incessant research activities in this area (including a worldwide practice of the above technologies) would boost the global agricultural sustainability and food security in salt-affected regions.

Credit author contribution statement

RM prepared the original draft. BS and NSB reviewed, edited and improved the draft. HSJ and PCS conceptualized and supervised the work, and reviewed and edited the draft.

Declaration of competing interest

The authors declare no conflict of interest.

Acknowledgements

The authors acknowledge the Prioritization, Monitoring and Evaluation cell of ICAR-CSSRI, Karnal, for internal review of the manuscript (Review Article/127/2020).

References

- Abdul-Kareem, A.W., Nazzal, K.E., 2013. Phytoremediation of salt-affected soils at Al-Jazeera northern irrigation project/Nineveh/Iraq. *Mesopotamia J. Agric.* 41, 294–298.
- Ahmad, I., 2014. *The Role of Cation Channels in Abiotic Stress Resistance in Rice*. University of York, York.
- Al-Busaidi, A., Yamamoto, T., Inoue, M., Egrinya Eneji, A., Mori, Y., Irshad, M., 2008. Effects of zeolite on soil nutrients and growth of Barley following irrigation with saline water. *J. Plant Nutr.* 31, 1159–1173.
- Amini, S., Ghadiri, H., Chen, C., Marschner, P., 2016. Salt-affected soils, reclamation, carbon dynamics, and biochar: a review. *J. Soils Sediments* 16, 939–953.
- Arora, S., Vanza, M., 2017. Microbial approach for bioremediation of saline and sodic soils. In: Arora, S., Singh, A.K., Singh, Y.P. (Eds.), *Bioremediation of Salt Affected Soils: an Indian Perspective*. Springer International Publishing, pp. 87–100.
- Arora, S., Patel, P., Vanza, M., Rao, G.G., 2014. Isolation and characterization of endophytic bacteria colonizing halophyte and other salt tolerant plant species from coastal Gujarat. *Afr. J. Microbiol. Res.* 8, 1779–1788.
- Aryal, J.P., Sapkota, T.B., Stirling, C.M., Jat, M.L., Jat, H.S., Rai, M., Mittal, S., Sutaliya, J.M., 2016. Conservation agriculture-based wheat production better copes with extreme climate events than conventional tillage-based systems: a case of untimely excess rainfall in Haryana, India. *Agric. Water Manag.* 233, 325–335.
- Auge, R.M., Toler, H.D., Saxton, A.M., 2015. Arbuscular mycorrhizal symbiosis alters stomatal conductance of host plants more under drought than under amply watered conditions: a meta-analysis. *Mycorrhiza* 25, 13–24.
- Aydemir, S., Najjar, N.F., 2005. Application of two amendments (gypsum and langbeinite) to reclaim sodic soil using sodic irrigation water. *Soil Res.* 43, 547–553.
- Babu, S., Prasanna, R., Bidyarani, N., Singh, R., 2015. Analysing the colonisation of inoculated cyanobacteria in wheat plants using biochemical and molecular tools. *J. Appl. Phycol.* 27, 327–338.
- Bal, H.B., Nayak, L., Das, S., Adhya, T.K., 2013. Isolation of ACC deaminase producing PGPR from rice rhizosphere and evaluating their plant growth promoting activity under salt stress. *Plant Soil* 366, 93–105.
- Balasundaram, N., 2004. *Sugarcane Management in Saline Soils*. Extension Publication No : 81. Sugarcane Breeding Institute, Coimbatore, India.

- Batanouny, K.H., 1993. Adaptation of plants to saline conditions in arid regions. In: Lieth, H., Masoom, A. Al (Eds.), *Towards the Rational Use of High Salinity Tolerant Plants*. Kluwer Academic Publishers, pp. 387–401.
- Begum, K., Kuhnert, M., Yeluripati, J.B., Ogle, S.M., Parton, W.J., Williams, S.A., Pan, G., Cheng, K., Ali, M.A., Smith, P., 2019. Modelling greenhouse gas emissions and mitigation potentials in fertilized paddy rice fields in Bangladesh. *Geoderma* 341, 206–215.
- Bhardwaj, A.K., Mishra, V.K., Singh, A.K., Arora, S., Srivastava, S., Singh, Y.P., Sharma, D.K., 2019. Soil salinity and land use-land cover interactions with soil carbon in a salt-affected irrigation canal command of Indo-Gangetic plain. *Catena* 180, 392–400.
- Bhattacharya, A.K., 2007. *Integrated Water Management Drainage: Report of Emeritus Scientist Scheme*. Division of Agricultural Education ICAR, New Delhi, p. 51.
- Bos, M.G., 2001. Selecting the drainage method for agricultural land. *Irrigat. Drain. Syst.* 15, 269–279.
- Burrell, L.D., Zehetner, F., Rampazzo, N., Wimmer, B., Soja, G., 2016. Long-term effects of biochar on soil physical properties. *Geoderma* 282, 96–102.
- Cao, J., Li, X., Kong, X., Zed, R., Dong, L., 2012. Using alfalfa (*Medicago sativa*) to ameliorate salt-affected soils in Yingda irrigation district in Northwest China. *Acta Ecol. Sin.* 32, 68–73.
- Castellano, M.J., Archontoulis, S.V., Helmers, M.J., Poffenbarger, H.J., Six, J., 2019. Sustainable intensification of agricultural drainage. *Nat Sustain* 2, 914–921. <https://doi.org/10.1038/s41893-019-0393-0>.
- Central Soil Salinity Research Institute (CSSRI), 2014. *Vision 2050. Pragmatic Assessment of the Agricultural Production and Food Demand Scenario of India by the Year 2050*. Central Soil Salinity Research Institute, Karnal, India.
- Chang Hoon, L., Lee, I., PilJoo, K., 2004. Effects of long-term fertilization on organic phosphorus fraction in paddy soil. *J. Soil Sci. Plant Nutr.* 50, 485–491.
- Chang, W., Sui, X., Fan, X.X., Jia, T.T., Song, F.Q., 2018. Arbuscular mycorrhizal symbiosis modulates antioxidant response and ion distribution in salt-stressed *Elaeagnus angustifolia* seedlings. *Front. Microbiol.* 9, 652.
- Chen, J., Mueller, V., 2018. Coastal climate change, soil salinity and human migration in Bangladesh. *Nat. Clim. Change*. <https://doi.org/10.1038/s41558-018-0313-8>.
- Chittapun, S., Limbipichai, S., Amnuaysin, N., Boonkerd, R., Charoensook, M., 2018. Effects of using cyanobacteria and fertilizer on growth and yield of rice, Pathum Thani I: a pot experiment. *J. Appl. Phycol.* 30, 79–85.
- Choudhary, M., Chandra, P., Arora, S., 2019. In: Dagar, J.C., Yadav, R.K., Sharma, P.C. (Eds.), *Research Developments in Saline Agriculture*. Springer Publishers, pp. 203–235.
- Choudhary, M., Sharma, P.C., Jat, H.S., Dash, A., Rajashekar, B., McDonald, A.J., Jat, M. L., 2018a. Soil bacterial diversity under conservation agriculture-based cereal systems in Indo-Gangetic Plains. *3 Biotech* 8, 304–315.
- Choudhary, M., Datta, A., Jat, H.S., Yadav, A.K., Gathala, M.K., Sapkota, T.B., Das, A.K., Sharma, P.C., Jat, M.L., Singh, R., Ladha, J.K., 2018b. Changes in soil biology under conservation agriculture based sustainable intensification of cereal systems in indo-Gangetic Plains. *Geoderma* 313, 193–204.
- Choudhary, M., Jat, H.S., Datta, A., Yadav, A.K., Sapkota, T.B., Mondal, S., Meena, R.P., Sharma, P.C., Jat, M.L., 2018c. Sustainable intensification influences soil quality, biota, and productivity in cereal-based agroecosystems. *Appl. Soil Ecol.* 126, 189–198.
- Choudhary, M., Sharma, P.C., Jat, H.S., Nehra, V., McDonald, A.J., Garg, N., 2016. Crop residue degradation by fungi isolated from conservation agriculture fields under rice-wheat system of North-West India. *Int. J. Recycl. Org. Waste Agric.* 5, 349–360.
- Corwin, D.L., 2020. Climate change impacts on soil salinity in agricultural areas. *Eur. J. Soil Sci.* <https://doi.org/10.1111/ejss.13010>.
- Dagar, J.C., 2014. Greening salty and waterlogged lands through agroforestry systems. In: Dagar, J.C., Singh, A.K., Arunachalam, A. (Eds.), *Agroforestry Systems in India: Livelihood Security and Environmental Services- Advances in Agroforestry*. Springer Publishers, pp. 333–344.
- Dagar, J.C., Yadav, R.K., 2017. Climate resilient approaches for enhancing productivity of saline agriculture. *J. Soil Salinity Water Qual.* 9, 9–29.
- Dagar, J.C., Lal, K., Ram, J., Kumar, M., Chaudhari, S.K., Yadav, R.K., Ahamad, S., Singh, G., Kaur, A., 2016. Eucalyptus geometry in agroforestry on waterlogged saline soils influences plant and soil traits in North-West India. *Agr. Ecosyst Environ* 233, 33–42.
- Dagar, J.C., Singh, G., Singh, N.T., 2001. Evaluation of forest and fruit trees used for rehabilitation of semi-arid alkali/sodic soils in India. *Arid Land Res. Manag.* 15, 115–133.
- Dagar, J.C., Yadav, R.K., Tomar, O.S., Minhas, P.S., Yadav, G., Lal, K., 2015a. Fruit-based agroforestry systems for saline water-irrigated semi-arid hyperthermic camborthids soils of north-west India. *Agrofor. Syst.* 90 (6), 1123–1132.
- Dagar, J.C., Yadav, R.K., Dar, S.R., Ahamad, S., 2015b. Liquorice (*Glycyrrhiza glabra*): a potential salt-tolerant highly remunerative medicinal crop which also helps in remediation of alkali soils. *Curr. Sci.* 108, 1683–1688.
- Dasgupta, S., Hossain, M.M., Huq, M., Wheeler, D., 2015. Climate change and soil salinity: the case of coastal Bangladesh. *Ambio*. <https://doi.org/10.1007/s13280-015-0681-5>.
- Datta, K.K., De Jong, C., Singh, O.P., 2000. Reclaiming salt-affected land through drainage in Haryana, India: a financial analysis. *Agric. Water Manag.* 46, 55–71.
- El hasini, S., Halima, I.O., Azzouzi, M.E., Douaik, A., Azim, K., Zouahri, A., 2019. Organic and inorganic remediation of soils affected by salinity in the seabka of sed el mesjoune – marrakech (Morocco). *Soil Tillage Res.* 193, 153–160.
- El-Agha, D., Molle, F., Rap, E., El Bilay, M., El-Hassan, W.A., 2019. Drainage water salinity and quality across nested scales in the Nile Delta of Egypt. *Environ. Sci. Pollut. Res.* <https://doi.org/10.1007/s11356-019-07154-y>.

- Elgharably, A., Marschner, P., Rengasamy, P., 2010. Wheat growth in a saline sandy loam soil as affected by N form and application rate. *Plant Soil* 328, 303–312.
- Emadodin, I., Narita, D., Bork, H.R., 2012. Soil degradation and agricultural sustainability: an overview from Iran. *Environ. Dev. Sustain.* 14, 611–625.
- Fageria, N.K., Gheyi, H.R., Moreira, A., 2011. Nutrient bioavailability in salt affected soils. *J. Plant Nutr.* 34, 945–962.
- Feng, H., Wang, S., Gao, Z., Wang, Z., Ren, X., Hu, S., Pan, H., 2019. Effect of land use on the composition of bacterial and fungal communities in saline-sodic soils. *Land Degrad. Dev.* 30, 1851–1860.
- Food and Agriculture organization of the United Nations (FAO), 2017. The state of food and agriculture: climate change, agriculture and food security. In: 40th Conference Session. Rome, pp. 1–14.
- Foster, S., Pulido-Bosch, A., Vallejos, A., Molina, L., Llop, A., MacDonald, A.M., 2018. Impact of irrigated agriculture on groundwater-recharge salinity: a major sustainability concern in semi-arid regions. *Hydrogeol. J.* 26, 2781–2791.
- Genc, Y., Taylor, J., Lyons, G., Li, Y., Cheong, J., Appelbee, M., Oldach, K., Sutton, T., 2019. Bread wheat with high salinity and sodicity tolerance. *Front. Plant Sci.* 10, 1280. <https://doi.org/10.3389/fpls.2019.01280>.
- Girma, M., Abdulahi, J., 2015. Irrigation system in Israel: a review. *Int. J. Water Resour. Environ. Eng.* 7, 29–37.
- Grigore, M., Toma, C., 2017. Definition and classification of halophytes. In: Grigore, M., Toma, C. (Eds.), *Anatomical Adaptations of Halophytes*. Springer Publishers, pp. 3–28.
- Habib, S.H., Kausar, H., Saud, H.M., 2016. Plant growth-promoting rhizobacteria enhance salinity stress tolerance in okra through ROS-scavenging enzymes. *BioMed Res. Int.* <https://doi.org/10.1155/2016/6284547>.
- Haj-Amor, Z., Bouri, S., 2019. Use of HYDRUS-1D-GIS tool for evaluating effects of climate changes on soil salinization and irrigation management. *Arch. Agron Soil Sci.* <https://doi.org/10.1080/03650340.2019.1608438>.
- Hayat, K., Bundschuh, J., Jan, F., Menhas, S., Hayat, S., Haq, F., Shah, M.A., Chaudhary, H.A., Ullah, A., Zhang, D., Zhou, Y., Zhou, P., 2019. Combating soil salinity with combining saline agriculture and phytomanagement with salt-accumulating plants. *Crit. Rev. Environ. Sci. Technol.* <https://doi.org/10.1080/10643389.2019.1646087>.
- Horriche, F.J., Benabdallah, S., 2020. Assessing aquifer water level and salinity for a managed artificial recharge site using reclaimed water. *Water* 12, 341–352.
- ILO (International Labour Organization), 2007. Employment by sector. In: Key Indicators of the Labour Market (KILM), fifth ed. www.ilo.org/public/english/employment/s trat/kilm/download/kilm04.pdf
- IPCC (Intergovernmental Panel on Climate Change), 2013. Summary for policymakers. In: Stocker, T.F., Qin, D., Plattner, G.K., Tignor, M., Allen, S.K., Boschung, J., Nauels, A., Xia, Y., Bex, V., Midgley, P.M. (Eds.), *Climate Change 2013: The Physical Science Basis. Contribution of Working Group to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press, Cambridge New York (CA, NY), p. 1535.
- Jalili, F., Khavazi, K., Pazira, E., Nejadi, A., Rahmani, H.A., Sadaghiani, H.R., Miransari, M., 2009. Isolation and characterization of ACC deaminase-producing fluorescent Pseudomonads, to alleviate salinity stress on canola (*Brassica napus* L.) growth. *J. Plant Physiol.* 166, 667–674.
- Jat, H.S., Datta, A., Choudhary, M., Sharma, P.C., Yadav, A.K., Choudhary, V., Gathala, M.K., Jat, M.L., McDonald, A., 2019. Climate Smart Agriculture practices improve soil organic carbon pools, biological properties and crop productivity in cereal-based systems of North-West India. *Catena*. <https://doi.org/10.1016/j.catena.2019.05.005>.
- Jat, H.S., Datta, A., Sharma, P.C., Kumar, V., Yadav, A.K., Choudhary, M., Choudhary, V., Gathala, M.K., Sharma, D.K., Jat, M.L., Yaduvanshi, N.P.S., Singh, G., McDonald, A., 2018. Assessing soil properties and nutrient availability under conservation agriculture practices in a reclaimed sodic soil in cereal-based systems of north-West India. *Arch. Agron Soil Sci.* 64, 531–545.
- Jesus, J.M., Danko, A.S., Fiuzza, A., Borges, M.-T., 2015. Phytoremediation of salt-affected soils: a review of processes, applicability, and the impact of climate change. *Environ. Sci. Pollut. Res.* 22, 6511–6525.
- Kadmiri, I.M., Chaouqui, L., Azaroual, S.E., Sijilmassi, B., Yaakoubi, K., Wahby, I., 2018. Phosphate solubilizing and auxin-producing Rhizobacteria promote plant growth under saline conditions. *Arabian J. Sci. Eng.* 43, 3403–3415.
- Kamra, S.K., 2013. Role of farmers' participation for effective management of groundwater recharge structures in Haryana. In: Proceedings Workshop on "Roadmap for Sustainable Groundwater Resources in Punjab and Haryana" Organized by CGWB North West Region, February 27, Chandigarh, India, pp. 88–99.
- Kamra, S.K., Kumar, S., Kumar, N., Dagar, J.C., 2019. Engineering and biological approaches for drainage of irrigated lands. In: Dagar, J.C., Yadav, R.K., Sharma, P.C. (Eds.), *Research Developments in Saline Agriculture*. Springer, pp. 537–577.
- Khan, I., 2019. Power generation expansion plan and sustainability in a developing country: a multi-criteria decision analysis. *J. Clean. Prod.* <https://doi.org/10.1016/j.jclepro.2019.02.161>.
- Kim, M.J., Radhakrishnan, R., Kang, S.M., You, Y.H., Jeong, E.J., Kim, J.G., Lee, I.J., 2017. Plant growth promoting effect of *Bacillus amyloliquefaciens* H-2-5 on crop plants and influence on physiological changes in soybean under soil salinity. *Physiol. Mol. Biol. Plants* 23, 571–580.
- Krishnamurthy, S.L., Sharma, P.C., Sharma, S.K., Batra, V., Kumari, V., Rao, L.V.S., 2016. Effect of salinity and use of stress indices of morphological and physiological traits at the seedling stage in rice. *Ind J. Exp Bot.* 54, 843–850.
- Kumar, S., Narjary, B., Kumar, K., Jat, H.S., Kamra, S.K., Yadav, R.K., 2019a. Developing soil matrix potential based irrigation strategies of direct seeded rice for improving yield and water productivity. *Agric. Water Manag.* 215, 8–15.
- Kumar, S., Pathan, A.L., Vivekanand, Kamra, S.K., 2019b. Evaluating efficacy of recharge structures in augmenting groundwater resources in Muzaffarnagar, Uttar Pradesh. *J. Agric. Eng.* 56, 1–8.
- Kumar, P., Mishra, A.K., Kumar, M., Chaudhari, S.K., Singh, R., Singh, K., Rai, P., Sharma, D.K., 2019c. Biomass production and carbon sequestration of *Eucalyptus tereticornis* plantation in reclaimed sodic soils of north-west India. *Indian J. Agric. Sci.* 89, 1091–1095.
- Kumar, S., Raju, R., Sheoran, P., Sharma, R., Yadav, R.K., Singh, R.K., Sharma, P.C., Chahal, V.P., 2020. Techno-economic evaluation of recharge structure as localized drainage option for sustainable crop production in sodic agro-ecosystems. *Indian J. Agric. Sci.* 90, 212–219.
- Lal, K., Chhabra, R., Mongia, A.D., Meena, R.L., Yadav, R.K., 2012. Release and uptake of potassium and sodium with fly ash application in rice on reclaimed alkali soil. *J. Indian Soc. Soil Sci.* 60, 1–6.
- Lal, R., 2015. Sequestering carbon and increasing productivity by conservation agriculture. *J. Soil Water Conserv.* 70, 55A–62A.
- Lashari, M.S., Liu, Y., Li, L., Pan, W., Fu, J., Pan, G., Zheng, J., Zheng, J., Zhang, X., Yu, X., 2013. Effects of amendment of biochar-manure compost in conjunction with pyrolytic solution on soil quality and wheat yield of a salt-stressed cropland from Central China Great Plain. *Field Crop. Res.* 144, 113–118.
- Lashari, M.S., Ye, Y., Ji, H., Li, L., Kibue, G.W., Lu, H., Zheng, J., Pan, G., 2015. Biochar-manure compost in conjunction with pyrolytic solution alleviated salt stress and improved leaf bioactivity of maize in a saline soil from central China: a 2-year field experiment. *J. Sci. Food Agric.* 95, 1321–1327.
- Li, H., Lei, P., Pang, X., Li, S., Xu, H., Xu, Z., Feng, X., 2017. Enhanced tolerance to salt stress in canola (*Brassica napus* L.) seedlings inoculated with the halotolerant *Enterobacter cloacae* HSNJ4. *Appl. Soil Ecol.* 119, 26–34.
- Li, H., Zhao, Q., Huang, H., 2019. Current status and challenges of salt-affected soil remediation by cyanobacteria. *Sci. Total Environ.* 666, 258–272.
- Li, J., Pu, L., Han, M., Zhu, M., Zhang, R., Xiang, Y., 2014. Soil salinization research in China: advances and prospects. *J. Geogr. Sci.* 24, 943–960.
- Lin, X.W., Xie, Z.B., Zheng, J.Y., Liu, Q., Bei, Q.C., Zhu, J.G., 2015. Effects of biochar application on greenhouse gas emissions, carbon sequestration and crop growth in coastal saline soil. *Eur. J. Soil Sci.* 66, 329–338.
- Liu, J., Tang, L., Gao, H., Zhang, M., Guo, C., 2018. Enhancement of alfalfa yield and quality by plant growth-promoting rhizobacteria under saline-alkali conditions. *J. Sci. Food Agric.* <https://doi.org/10.1002/jsfa.9185>.
- MacMillan, R.A., Marciak, L.C., 2001. Procedures Manual for Watershed-Based Salinity Management: A Comprehensive Manual for Assessing and Addressing Salinity on a Watershed Basis. Alberta Agriculture, Food and Rural Development, Conservation and Development Branch, Edmonton, Alberta, p. 54.
- Makrana, G., Kumar, A., Yadav, R.K., Kumar, R., Soni, P.G., Lata, C., Sheoran, P., 2019. Effect of saline water irrigations on physiological, biochemical and yield attributes of dual purpose pearl millet (*Pennisetum glaucum* L.) varieties. *Indian J. Agric. Sci.* 89, 624–633.
- Mandal, S., Raju, R., Kumar, A., Kumar, P., Sharma, P.C., 2018. Current status of research, technology response and policy needs of salt-affected soils in India—A review. *J. Indian Soc. Coastal Agric. Res.* 36, 40–53.
- Mandal, S., Thangarajan, R., Bolan, N.S., Sarkar, B., Khan, N., Ok, Y.S., Naidu, R., 2016. Biochar-induced concomitant decrease in ammonia volatilization and increase in nitrogen use efficiency by wheat. *Chemosphere* 142, 120–127.
- Mandal, U.K., Bhardwaj, A.K., Warrington, D.N., Goldstein, D., Bar Tal, A., Levy, G.J., 2008. Changes in soil hydraulic conductivity, runoff, and soil loss due to irrigation with different types of saline-sodic water. *Geoderma* 144, 509–516.
- Mandal, U.K., Burman, D., Bhardwaj, A.K., Nayak, D.B., Samui, A., Mahanta, K.K., Lama, T.D., Maji, B., Mondal, S., Rout, S., Sarangi, S.K., 2019. Waterlogging and coastal salinity management through land shaping and cropping intensification in climatically vulnerable Indian Sundarbans. *Agric. Water Manag.* 216, 12–26.
- Manjajiah, K.M., Mukhopadhyay, R., Paul, R., Datta, S.C., Kumararaja, P., Sarkar, B., 2019. Clay minerals and zeolites for environmentally sustainable agriculture. In: Mercurio, M., Sarkar, B., Langella, A. (Eds.), *Modified Clay and Zeolite Nanocomposite Materials*. Elsevier Publishers, pp. 309–329.
- Meena, M.D., Joshi, P.K., Narjary, B., Sheoran, P., Jat, H.S., Chinchmalpure, Anil, R., Yadav, R.K., Sharma, D.K., 2016. Effects of municipal solid waste, rice straw composts and mineral fertilizers on biological and chemical properties of a saline soil and yields of a mustard-pearl millet cropping system. *Soil Res.* 54, 958–966.
- Meena, M.D., Narjary, B., Sheoran, P., Jat, H.S., Joshi, P.K., Chinchmalpure, A.R., Yadav, G., Yadav, R.K., Meena, M.K., 2018. Changes of phosphorus fractions in saline soil amended with municipal solid waste compost and mineral fertilizers in a mustard-pearl millet cropping system. *Catena* 160, 32–40.
- Meena, M.D., Yadav, R.K., Narjary, B., Yadav, G., Jat, H.S., Sheoran, P., Meena, M.K., Antil, R.S., Meena, B.L., Singh, H.V., Meena, V.S., Rai, P.K., Ghosh, A., Moharana, P. C., 2019. Municipal solid waste (MSW): strategies to improve salt affected soil sustainability: a review. *Waste Manag.* 84, 38–53.
- Minhas, P.S., Qadir, M., Yadav, R.K., 2019. Groundwater irrigation induced soil sodification and response options. *Agric. Water Manag.* 215, 74–85.
- Minhas, P.S., Ramos, T.B., Ben-Gal, A., Pereira, L.S., 2020a. Coping with salinity in irrigated agriculture: crop evapotranspiration and water management issues. *Agric. Water Manag.* 227, 105832.
- Minhas, P.S., Yadav, R.K., Bali, A., 2020b. Perspectives on reviving waterlogged and saline soils through plantation forestry. *Agric. Water Manag.* 232, 106063.
- Mishra, V.K., Jha, S.K., Damodaran, T., Singh, Y.P., Srivastava, S., Sharma, D.K., Prasad, J., 2019. Feasibility of coal combustion fly ash alone and in combination with gypsum and green manure for reclamation of degraded sodic soils of the Indo-Gangetic Plains: a mechanism evaluation. *Land Degrad. Dev.* 30, 1300–1312.

- Munns, R., Tester, M., 2008. Mechanisms of salinity tolerance. *Annu. Rev. Plant Biol.* 59, 651–681.
- Munns, R., James, R.A., Gilliam, M., Flowers, T.J., Colmer, T.D., 2016. Tissue tolerance, an essential but elusive trait for salt-tolerant crops. *Funct. Plant Biol.* 43, 1103–1113.
- Nadeem, S.M., Zahir, Z.A., Naveed, M., Nawaz, S., 2013. Mitigation of salinity-induced negative impact on the growth and yield of wheat by plant growth-promoting rhizobacteria in naturally saline conditions. *Ann. Microbiol.* 63, 225–232.
- Narjary, B., Kumar, S.K., Kamra, S.K., Bundela, D.S., Sharma, D.K., 2014. Impact of rainfall variability on groundwater resources and opportunities of artificial recharge structure to reduce its exploitation in fresh groundwater zones of Haryana. *Curr. Sci.* 107, 1305–1312.
- National Academy of Agricultural Sciences (NAAS), 2012. Management of Crop Residues in the Context of Conservation Agriculture; Policy Paper No. 58. National Academy of Agricultural Sciences, New Delhi, India, p. 12.
- Nguyen, B.T., Trinh, N.N., Bach, Q.V., 2020. Methane emissions and associated microbial activities from paddy salt-affected soil as influenced by biochar and cow manure addition. *Appl. Soil Ecol.* 152, 103531.
- Nijland, H.J., Croon, F.W., Ritzema, H.P., 2005. Subsurface Drainage Practices: Guidelines for the Implementation, Operation and Maintenance of Subsurface Pipe Drainage Systems. Wageningen Alterra ILRI Publication No 60, p. 608.
- Nisha, R., Kiran, B., Kaushik, A., Kaushik, C.P., 2018. Bioremediation of salt affected soils using cyanobacteria in terms of physical structure, nutrient status and microbial activity. *Int. J. Environ. Sci. Technol.* 15, 571–580.
- Noori, M., Zendehele, M., Ahmadi, A., 2006. Using natural zeolite for the improvement of soil salinity and crop yield. *Toxicol. Environ. Chem.* 88, 77–84.
- Purakayastha, T.J., Bera, T., Bhaduri, D., Sarkar, B., Mandal, S., Wade, P., Kumari, S., Biswas, S., Menon, M., Pathak, H., Tsang, D.C.W., 2019. A review on biochar modulated soil condition improvements and nutrient dynamics concerning crop yields: pathways to climate change mitigation and global food security. *Chemosphere* 227, 345–365.
- Qadir, M., Quillerou, E., Nangia, V., Murtaza, G., Singh, M., Thomad, R.J., Drecshel, P., Noble, A.D., 2014. Economics of salt induced land degradation and restoration. *Nat. Resour. Forum* 38, 282–295.
- Qi, F., Lamb, D., Naidu, R., Bolan, N.S., Yan, Y., Ok, Y.S., Rahman, M.M., Choppala, G., 2018. Cadmium solubility and bioavailability in soils amended with acidic and neutral biochar. *Sci. Total Environ.* 610–611, 1457–1466.
- Rahman, M.S., Di, L., Yu, E.G., Tang, J., Lin, L., Zhang, C., Yu, Z., Gaigalas, J., 2018. Impact of climate change on soil salinity: a remote sensing based investigation in coastal Bangladesh. In: The Seventh International Conference on Agro-Geoinformatics, August 6–9, China. <https://doi.org/10.1109/Agro-Geoinformatics.2018.8476036>.
- Ram, J., Dagar, J.C., Lal, K., Singh, G., Toky, O.P., Tanwar, R.S., Dar, S.R., Mukesh, K., 2011. Biodrainage to combat water logging, increase farm productivity and sequester carbon in canal command area of north-west India. *Curr. Sci.* 100, 1673–1680.
- Ravikiran, K.T., Krishnamurthy, S.L., Warraich, A.S., Sharma, P.C., 2017. Diversity and haplotypes of rice genotypes for seedling stage salinity tolerance analyzed through morpho-physiological and SSR markers. *Field Crop. Res.* 220, 10–18.
- Ravindran, K.C., Venkatesan, K., Balakrishnan, V., Chellappan, K.P., Balasubramanian, T., 2007. Restoration of saline land by halophytes for Indian soils. *Soil Biol. Biochem.* 39, 2661–2664.
- Richards, L.A. (Ed.), 1954. *Diagnosis and Improvement of Saline Alkali Soils*. USDA Handbook No. 60, Washington, DC, USA.
- Rojas-Tapias, D., Moreno-Galván, A., Pardo-Díaz, S., Obando, M., Rivera, D., Bonilla, R., 2012. Effect of inoculation with plant growth-promoting bacteria (PGPB) on amelioration of saline stress in maize (*Zea mays*). *Appl. Soil Ecol.* 61, 264–272.
- Rossi, F., Li, H., Liu, Y., De Philippis, R., 2017. Cyanobacterial inoculation (cyanobacterisation): perspectives for the development of a standardized multifunctional technology for soil fertilization and desertification reversal. *Earth Sci. Rev.* 171, 28–43.
- Saifullah, Dahlawi, S., Naem, A., Rengel, Z., Naidu, R., 2018. Biochar application for the remediation of salt-affected soils: challenges and opportunities. *Sci. Total Environ.* 625, 320–335.
- Sandhu, O.S., Gupta, R.K., Thind, H.S., Jat, M.L., Sidhu, H.S., Singh, Y., 2019. Drip irrigation and nitrogen management for improving crop yields, nitrogen use efficiency and water productivity of maize-wheat system on permanent beds in north-west India. *Agric. Water Manag.* 219, 19–26.
- Sapre, S., Gontia-Mishra, I., Tiwari, S., 2018. *Klebsiella* sp. confers enhanced tolerance to salinity and plant growth promotion in oat seedlings (*Avena sativa*). *Microbiol. Res.* 206, 25–32.
- Sarang, S.K., Singh, S., Kumar, V., Srivastava, A.K., Sharma, P.C., Johnson, D.E., 2019. Tillage and crop establishment options for enhancing the productivity, profitability, and resource use efficiency of rice-rabi systems of the salt-affected coastal lowlands of eastern India. *Field Crop. Res.* <https://doi.org/10.1016/j.fcr.2019.03.016>.
- Sarkar, A., Ghosh, P.K., Pramanik, K., Mitra, S., Soren, T., Pandey, S., Mondal, M.H., Maiti, T.K., 2018. A halotolerant *Enterobacter* sp. displaying ACC deaminase activity promotes rice seedling growth under salt stress. *Res. Microbiol.* 169, 20–32.
- Seshadri, B., Bolan, N.S., Naidu, R., Wang, H., Sajwan, K., 2013. Clean coal technology combustion products: properties, agricultural and environmental applications, and risk management. *Adv. Agron.* 119, 309–370.
- Sewa, Ram, Rao, K.V.G.K., Viswanatha, N.A., 2000. Impact of subsurface drainage in management of saline soils of the Chambal command. In: 8th ICID International Drainage Workshop, January 31st – February 4th, New Delhi, pp. 97–104.
- Shaheen, S.M., Niazi, N.K., Hassan, N.E.E., Bibi, I., Wang, H., Tsang, D.C.W., Ok, Y.S., Bolan, N., Rinklebe, J., 2018. Wood-based biochar for the removal of potentially toxic elements in water and wastewater: a critical review. *Int. Mater. Rev.* <https://doi.org/10.1080/09506608.2018.1473096>.
- Shahid, S.A., Zaman, M., Heng, L., 2018. Soil salinity: historical perspectives and a world overview of the problem. In: Shahid, S.A., Zaman, M., Heng, L. (Eds.), *Guideline for Salinity Assessment, Mitigation and Adaptation Using Nuclear and Related Techniques*. Springer, pp. 43–53.
- Sharma, S.K., 2010. Salt tolerant crop varieties developed at CSSRI Karnal, India: a success story, pp. 1–13. www.PlantStress.com.
- Shirale, O.A., Kharche, V.K., Zadode, R.S., Meena, B.P., Rajendiran, S., 2017. Soil biological properties and carbon dynamics subsequent to organic amendments addition in sodic black soils. *Arch. Agron Soil Sci.* 63, 2023–2034.
- Sidhu, H.S., Jat, M.L., Singh, Y., Sidhu, R.K., Gupta, N., Singh, P., Jat, H.S., Gerard, B., 2019. Sub-surface drip fertigation with conservation agriculture in a rice-wheat system: a breakthrough for addressing water and nitrogen use efficiency. *Agric. Water Manag.* 216, 273–283.
- Singh, J., Sharma, P.C., Sharma, S.K., Rai, M., 2014. Assessing effect of salinity on oil quality parameters of Indian mustard (*Brassica juncea* L. Czern & Coss) using Fourier Transform Near-Infrared Reflectance (FT-NIR) spectroscopy. *Grasas Aceites* 65, e009. <https://doi.org/10.3989/gya.063413>.
- Singh, Y.P., Arora, S., Mishra, V.K., Dixit, H., Gupta, R.K., 2018a. Effect of organic and inorganic amendments on amelioration of sodic soil and sustaining rice (*Oryza sativa*)-wheat (*Triticum aestivum*) productivity. *Indian J. Agric. Sci.* 88, 1455–1462.
- Singh, A., Arora, S., Singh, Y.P., Verma, C.L., Bhardwaj, A.K., Sharma, N., 2018b. Water use in rice crop through different methods of irrigation in a sodic soil. *Paddy Water Environ.* 16, 587–593.
- Singh, V., Singh, A.P., Bhadoria, J., Giri, J., Singh, J., Vineeth, T.V., Sharma, P.C., 2018c. Differential expression of salt-responsive genes to salinity stress in salt-tolerant and salt-sensitive rice (*Oryza sativa* L.) at seedling stage. *Protoplasma* 255, 1667–1681.
- Singh, J.S., 2015. Plant-microbe interactions: a viable tool for agricultural sustainability. *Appl. Soil Ecol.* 92, 45–46.
- Singh, J., Sastry, E.V.D., Singh, V., 2012. Effect of salinity on tomato (*Lycopersicon esculentum* Mill.) during seed germination stage. *Physiol. Mol. Biol. Plants* 18, 45–50.
- Singh, V., Singh, D.V., 2015. Cyanobacteria mediated changes and its impact on bioremediation of saline-alkaline soils. *Bangladesh J. Bot.* 44, 653–658.
- Singh, Y.P., Arora, S., Mishra, V.K., Dixit, H., Gupta, R.K., 2019. Plant and soil responses to the combined application of organic amendments and inorganic fertilizers in degraded sodic soils of Indo-Gangetic Plains. *Commun. Soil Sci. Plant Anal.* <https://doi.org/10.1080/00103624.2019.1671446>.
- Singh, Y.P., Mishra, V.K., Sharma, D.K., Singh, G., Arora, S., Dixit, H., Cerdà, A., 2016. Harnessing productivity potential and rehabilitation of degraded sodic lands through Jatropha based intercropping systems. *Agric. Ecosyst. Environ.* 233, 121–129.
- Sirault, X.R.R., James, R.A., Furbank, R.T., 2009. A new screening method for osmotic component of salinity tolerance in cereals using infrared thermography. *Funct. Plant Biol.* 36, 970–977.
- Srinivasulu, A., Satyanarayana, T.V., Kumar, H.V.H., 2005. Subsurface drainage in a pilot area in Nagarjuna Sagar right canal command, India. *Irrigat. Drain. Syst.* 19, 61–70.
- Srivastava, P., Kumar, R., 2015. Soil salinity: a serious environmental issue and plant growth promoting bacteria as one of the tools for its alleviation. *Saudi J. Biol. Sci.* 22, 123–131.
- Stevens, R., Pitt, T., 2012. Guidelines for managing soil salinity in groundwater irrigated vineyards. In: *Sustainable Irrigation Management Update*. <http://www.fao.org/ct/act/sustainable-agriculture-platform-pilot-website/soil-salinity-management/en/>.
- Sun, H., Lu, H., Chu, L., Shao, H., Shi, W., 2017. Biochar applied with appropriate rates can reduce N leaching, keep N retention and not increase NH₃ volatilization in a coastal saline soil. *Sci. Total Environ.* 575, 820–825.
- Sundha, P., Basak, N., Rai, A.K., Yadav, R.K., Sharma, P.C., Sharma, D.K., 2020. Can conjunctive use of gypsum, city waste composts and marginal quality water rehabilitate saline-sodic soils? *Soil Tillage Res.* 200, 104608. <https://doi.org/10.1016/j.still.2020.104608>.
- Swapnil, P., Rai, A.K., 2018. Physiological responses to salt stress of salt-adapted and directly salt (NaCl and NaCl + Na₂SO₄ mixture)-stressed cyanobacterium *Anabaena fertilissima*. *Protoplasma* 255, 963–976.
- Tavakkoli, E., Fatehi, F., Rengasamy, P., McDonald, G.K., 2012. A comparison of hydroponic and soil-based screening methods to identify salt tolerance in the field in barley. *J. Exp. Bot.* 63, 3853–3868.
- Tejawat, C.M., 2015. Large scale reclamation of waterlogged saline soils in Chambal Command area, Rajasthan. In: *Proceedings Workshop on Waterlogging and Soil Salinity in Irrigated Agriculture*, September 3–4, Chandigarh, pp. 125–137.
- Thijs, S., Dillewijn, P.W., Sillen, W., Truysens, S., Holtappels, M., Haen, J.D., 2014. Exploring the rhizospheric and endophytic bacterial communities of *Acer pseudoplatanus* growing on a TNT-contaminated soil: towards the development of a rhizocompetent TNT detoxifying plant growth promoting consortium. *Plant Soil* 385, 15–36.
- Tiwari, P., Goel, A., 2015. An overview of impact of subsurface drainage project studies on salinity management in developing countries. *Appl. Water Sci.* 7, 569–580.
- Tripathi, R.S., 2011. Socio-economic impact of reclaiming salt affected lands in India. *J. Soil Salinity Water Qual* 3, 110–126.
- United Nations, 2011. World population prospects: the 2010 revision. In: *Comprehensive Tables*. Department of Economic and Social Affairs, Population Division, vol. 1. United Nations, New York.
- Wang, L., Sun, X., Li, S., Zhang, T., Zhang, W., Zhai, P., 2014. Application of organic amendments to a coastal saline soil in North China: effects on soil physical and chemical properties and tree growth. *PLoS One* 9, 1–9.
- Wen, J., Dong, H., Zeng, G., 2018. Application of zeolite in removing salinity/sodicity from wastewater: a review of mechanisms, challenges and opportunities. *J. Clean. Prod.* <https://doi.org/10.1016/j.jclepro.2018.06.270>.

- Werber, J.R., Deshmukh, A., Elimelech, M., 2017. Can batch or semi-batch processes save energy in reverse-osmosis desalination? *Desalination* 402, 109–122.
- Wheeler, D., 2011. Quantifying Vulnerability to Climate Change: Implications for Adaptation Assistance. Centre for Global Development, Washington. Working Paper No. 240.
- Win, K.T., Tanaka, F., Okazaki, K., Ohwaki, Y., 2018. The ACC deaminase expressing endophyte *Pseudomonas* spp. Enhances NaCl stress tolerance by reducing stress-related ethylene production, resulting in improved growth, photosynthetic performance, and ionic balance in tomato plants. *Plant Physiol. Biochem.* 127, 599–607.
- Wu, Y.Y., Chen, Q.J., Chen, M., Chen, J., Wang, X.C., 2005. Salt-tolerant transgenic perennial ryegrass (*Lolium perenne* L.) obtained by *Agrobacterium tumefaciens*-mediated transformation of the vacuolar Na⁺/H⁺ antiporter gene. *Plant Sci.* 169, 65–73.
- Xu, G., Zhang, Y., Sun, J., Shao, H., 2016. Negative interactive effects between biochar and phosphorus fertilization on phosphorus availability and plant yield in saline sodic soil. *Sci. Total Environ.* 568, 910–915.
- Yadav, R.K., Singh, S.P., Lal, D., Kumar, A., 2007. Fodder production and soil health with conjunctive use of saline and good quality water in ustipsamments of a semi-arid region. *Land Degrad. Dev.* 18, 153–161.
- Yao, F., Huang, J., Cui, K., Nie, L., Xiang, J., Liu, X., Wu, W., Chen, M., Peng, S., 2012. Agronomic performance of high-yielding rice variety grown under alternate wetting and drying irrigation. *Field Crop. Res.* 126, 16–22.
- Yasin, N.A., Khan, W.U., Ahmad, S.R., Ali, A., Ahmad, A., Akram, W., 2018. Imperative roles of halotolerant plant growth-promoting rhizobacteria and kinetin in improving salt tolerance and growth of black gram (*Phaseolus mungo*). *Environ. Sci. Pollut. Res.* 25, 4491–4505.
- Yu, P., Liu, S., Xu, Q., Fan, G., Huang, Y., Zhou, D., 2019. Response of soil nutrients and stoichiometric ratios to short-term land use conversions in a salt-affected region, northeastern China. *Ecol. Eng.* 129, 22–28.
- Yu, P.J., Liu, S.W., Zhang, L., Li, Q., Zhou, D.W., 2018. Selecting the minimum data set and quantitative soil quality indexing of alkaline soils under different land uses in northeastern China. *Sci. Total Environ.* 616–617, 564–571.
- Zahran, M.A., Abdel Wahid, A.A., 1982. Contributions to the ecology of halophytes. *Tasks Veg. Sci.* 2, 235–257.
- Zerrouk, I.Z., Benchabane, M., Khelifi, L., Yokawa, K., Ludwig-Müller, J., Baluska, F., 2016. A *Pseudomonas* strain isolated from date-palm rhizospheres improves root growth and promotes root formation in maize exposed to salt and aluminum stress. *J. Plant Physiol.* 191, 111–119.
- Zhang, L., Shao, H., Wang, B., Zhang, L., Qin, X., 2019. Effects of nitrogen and phosphorus on the production of carbon dioxide and nitrous oxide in salt-affected soils under different vegetation communities. *Atmos. Environ.* <https://doi.org/10.1016/j.atmosenv.2019.02.024>.
- Zhang, L.H., Song, L.P., Wang, B.C., Shao, H.B., Zhang, L.W., Qin, X.C., 2018. Co-effects of salinity and moisture on CO₂ and N₂O emissions of laboratory-incubated salt-affected soils from different vegetation types. *Geoderma* 332, 109–120.
- Zhu, X., Song, F., Liu, S., Liu, F., 2016. Role of arbuscular mycorrhiza in alleviating salinity stress in wheat (*Triticum aestivum* L.) grown under ambient and elevated CO₂. *J. Agron. Crop Sci.* 202, 486–496.