



हर कदम, हर डगर
किसानों का हमसफर
भारतीय कृषि अनुसंधान परिषद

Agrisearch with a human touch

**LAND RESOURCE INVENTORY AND SOCIO-ECONOMIC STATUS OF
FARM HOUSEHOLDS FOR WATERSHED PLANNING AND
DEVELOPMENT**

NEREGALLU-3 (4D4A1T2c) MICROWATERSHED

Koppal Taluk and District, Karnataka

Karnataka Watershed Development Project – II

SUJALA – III

World Bank funded Project



THE WORLD BANK



ICAR – NATIONAL BUREAU OF SOIL SURVEY AND LAND USE PLANNING



ICAR - NBSS & LUP

**WATERSHED DEVELOPMENT DEPARTMENT
GOVT. OF KARNATAKA, BANGALORE**



About ICAR - NBSS&LUP

The ICAR-National Bureau of Soil Survey and Land Use Planning (ICAR-NBSS&LUP), Nagpur, a premier Institute of the Indian Council of Agricultural Research (ICAR), was set up during 1976 with the objective to prepare soil resource maps at national, state and district levels and to provide research inputs in soil resource mapping and its applications, land evaluation, land use planning, land resource management, and database management using GIS for optimising land use on different kinds of soils in the country.

The Bureau has been engaged in carrying out soil resource survey, agro-ecological and soil degradation mapping at the country, state and district levels for qualitative assessment and monitoring the soil health towards viable land use planning. The research activities have resulted in identifying the soil potentials and problems, and the various applications of the soil surveys with the ultimate objective of sustainable agricultural development. The Bureau has the mandate to correlate and classify soils of the country and maintain a National Register of all the established soil series. The Institute is also imparting in-service training to staff of the soil survey agencies in the area of soil survey, land evaluation and soil survey interpretations for land use planning. The Bureau in collaboration with Panjabrao Krishi Vidyapeeth, Akola is running post-graduate teaching and research programme in land resource management, leading to M.Sc. and Ph.D. degrees.

Citation: Rajendra Hegde, Ramesh Kumar, S.C., K.V. Niranjana, S. Srinivas, M.Lalitha, B.A. Dhanorkar, R.S. Reddy and S.K. Singh (2019). "Land resource inventory and socio-economic status of farm households for watershed planning and development of Naregallu-3 (4D4A1T2c) Microwatershed, Koppal Taluk and District, Karnataka", ICAR-NBSS&LUP Sujala MWS Publ .434, ICAR – NBSS & LUP, RC, Bangalore. p.145 &37

TO OBTAIN COPIES,

Please write to:

Director, ICAR - NBSS & LUP,

Amaravati Road, NAGPUR - 440 033, India

Phone : (0712) 2500386, 2500664, 2500545 (O)

Telefax : 0712-2522534

E-Mail : director@nbsslup.ernet.in

Website URL : nbsslup.in

Or

Head, Regional Centre, ICAR - NBSS&LUP, Hebbal, Bangalore - 560 024

Phone : (080) 23412242, 23510350 (O)

Telefax : 080-23510350

E-Mail : nbssrcb@gmail.com



**LAND RESOURCE INVENTORY AND SOCIO-ECONOMIC
STATUS OF FARM HOUSEHOLDS FOR WATERSHED
PLANNING AND DEVELOPMENT**

NEREGALLU-3 (4D4A1T2c) MICROWATERSHED

Koppal Taluk and District, Karnataka

Karnataka Watershed Development Project – II

Sujala-III

World Bank funded Project



**ICAR – NATIONAL BUREAU OF SOIL SURVEY AND LAND
USE PLANNING**



ICAR - NBSS & LUP



**WATERSHED DEVELOPMENT DEPARTMENT, GOVT. OF
KARNATAKA, BANGALORE**



PREFACE

In Karnataka, as in other Indian States, the livelihoods of rural people are intertwined with farming pursuits. The challenges in agriculture are seriously threatening the livelihood of a large number of farmers as they have been practicing farming in contextual factors beyond their control. Climatic factors are the most important ones and have become much more significant in recent times due to rapid climate changes induced by intensive anthropogenic activities affecting our ecosystem in multiple ways. Climate change has become the reality, it is happening and efforts to evolve and demonstrate climate resilient technologies have become essential. Due to the already over stressed scenario of agrarian sector, the climate change is resulting in manifold increase in the complexities, pushing the rural mass to face more and more unpredictable situations. The rising temperatures and unpredictable rainfall patterns are going to test seriously the informed decisions farmers have to make in order to survive in farming and sustain their livelihood.

It is generally recognized that impacts of climate change shall not be uniform across the globe. It is said that impact of climate change is more severe in South Asia. Based on the analysis of meteorological data, it is predicted that in India, there will be upward trend in mean temperature, downward trend in relative humidity, annual rainfall and number of wet days in a year. Also, in general, phenomena like erratic monsoon, spread of tropical diseases, rise in sea levels, changes in availability of fresh water, frequent floods, droughts, heat waves, storms and hurricanes are predicted. Each one of these adverse situations are already being experienced in various parts of India and also at the global level. Decline in agricultural productivity of small and marginal farmers becoming more vulnerable is already witnessed.

In Karnataka, more than 60 per cent of the population live in rural areas and depend on agriculture and allied activities for their livelihood. Though the state has achieved significant progress in increasing the yield of many crops, there is tremendous pressure on the land resources due to the growing and competing demands of various land uses. This is reflected in the alarming rate of land degradation observed. Already more than 50 per cent of the area is affected by various forms of degradation. If this trend continues, the sustainability of the fragile ecosystem will be badly affected. The adverse effects of change in the climatic factors are putting additional stress on the land resources and the farmers dependent on this.

The natural resources (land, water and vegetation) of the state need adequate and constant care and management, backed by site-specific technological interventions and investments particularly by the government. Detailed database pertaining to the nature of the land resources, their constraints, inherent potentials and suitability for various land

based rural enterprises, crops and other uses is a prerequisite for preparing location-specific action plans, which are in tune with the inherent capability of the resources. Any effort to evolve climate resilient technologies has to be based on the baseline scientific database. Then only one can expect effective implementation of climate resilient technologies, monitor the progress, make essential review of the strategy, and finally evaluate the effectiveness of the implemented programs. The information available at present on the land resources of the state are of general nature and useful only for general purpose planning. Since the need of the hour is to have site-specific information suitable for farm level planning and detailed characterization and delineation of the existing land resources of an area into similar management units is the only option.

ICAR-NBSS&LUP, Regional Centre, Bangalore has taken up a project sponsored by the Karnataka Watershed Development Project-II, (Sujala-III), Government of Karnataka funded by the World Bank under Component -1 Land Resource Inventory. This study was taken up to demonstrate the utility of such a database in reviewing, monitoring and evaluating all the land based watershed development programs on a scientific footing. To meet the requirements of various land use planners at grassroots level, the present study on “Land Resource Inventory and Socio-Economic Status of Farm Households for Watershed Planning and Development of for Neregallu-3 microwatershed in Koppal Taluk, Koppal District, Karnataka” for integrated development was taken up in collaboration with the State Agricultural Universities, IISC, KSRSAC, KSNDMC as Consortia partners. The project provides detailed land resource information at cadastral level (1:7920 scale) for all the plots and socio-economic status of farm households covering thirty per cent farmers randomly selected representing landed and landless class of farmers in the micro-watershed. The project report with the accompanying maps for the microwatershed will provide required detailed database for evolving effective land use plan, alternative land use options and conservation plans for the planners, administrators, agricultural extension personnel, KVK officials, developmental departments and other land users to manage the land resources in a sustainable manner.

It is hoped that this database will be useful to the planners, administrators and developmental agencies working in the area in not only for formulating location specific developmental schemes but also for their effective monitoring at the village/watershed level.

Nagpur

Date:21-10-2019

S.K. SINGH

Director, ICAR - NBSS&LUP,Nagpur

Contributors

Dr. Rajendra Hegde Principal Scientist, Head & Project Leader, Sujala-III Project ICAR-NBSS&LUP, Regional Centre, Bangalore	Dr. S.K.Singh Director, ICAR-NBSS&LUP Coordinator, Sujala-III Project Nagpur
Soil Survey, Mapping & Report Preparation	
Dr. K.V. Niranjana	Sh. R.S. Reddy
Dr. B.A. Dhanorkar	Smt. Chaitra, S.P.
	Dr. Gopali Bardhan
	Sh. Somashekar T.N
	Ms. Arpitha G.M
	Dr. Mahendra kumar M.B
	Sh. Tirupati Meti
Field Work	
Sh. C. Bache Gowda	Sh. Mayur Patil
Sh. Somashekar	Sh. Arun Kumar, S.
Sh. M. Jayaramaiah	Sh. Sunil Raj
	Sh. Yogesh Kumar, B.
	Sh. Vikas, N.K.
	Sh. Arun Kumar, S.G.
	Sh. Umesh Jادیappa Madolli
	Sh. Praveen Kumar P. Achalkar
	Sh. Veerabhadraswamy
	Sh. Vinay
	Sh. Shankarappa, K.
	Sh. Lankesh, R.S.
	Sh. Appanna B. Hattigoudar
	Sh. Maharudra
GIS Work	
Dr. S.Srinivas	Sh. A.G.Devendra Prasad
Sh. D.H.Venkatesh	Sh. Abhijith Sastry, N.S.
Smt. K.Sujatha	Smt. Shyla, B.
Smt. K.V.Archana	Smt. Swetha ,K.
Sh. N.Maddileti	Ms. Vidya, P.C.
	Sh. Deepak, M.J.
	Smt. K.Karunya Lakshmi
	Ms. Seema, K.V.

Laboratory Analysis	
Dr. M. Lalitha	Sh. Vindhya, N.G.
Smt. Arti Koyal	Ms. P. Pavanakumari, P.
Smt. Parvathy, S.	Ms. Rashmi, N.
	Ms. Leelavathy, K.U.
	Smt. Usha Kiran, G.
Socio-Economic Analysis	
Dr. S.C. Ramesh Kumar	Sh. M.K. Prakashanaik
	Ms. Shraddha Hegde
	Mrs. Sowmya A N
	Sh. Vijay Kumar
	Sh. Pradyumna
	Ms. Sowmya K.B
	Mrs. Prathibha, D.G
	Sh. Rajendra,D
Soil & Water Conservation	
Sh. Sunil P. Maske	
Watershed Development Department, GoK, Bangalore	
Sh. Rajeev Ranjan IFS Project Director & Commissioner, WDD	Dr. A. Natarajan NRM Consultant, Sujala-III Project
Dr. S.D. Pathak IFS Executive Director & Chief Conservator of Forests, WDD	

PART-A

LAND RESOURCE INVENTORY

Contents

Preface		
Contributors		
Executive Summary		
Chapter 1	Introduction	1
Chapter 2	Geographical Setting	3
2.1	Location and Extent	3
2.2	Geology	3
2.3	Physiography	4
2.4	Drainage	5
2.5	Climate	5
2.6	Natural Vegetation	6
2.7	Land Utilization	7
Chapter 3	Survey Methodology	11
3.1	Base maps	11
3.2	Image Interpretation for Physiography	11
3.3	Field Investigation	14
3.4	Soil mapping	16
3.5	Laboratory Characterization	17
3.6	Land Management Units	17
Chapter 4	The Soils	23
4.1	Soils of Granite Gneiss Landscape	23
4.2	Soils of Alluvial Landscape	25
Chapter 5	Interpretation for Land Resource Management	43
5.1	Land Capability Classification	43
5.2	Soil Depth	45
5.3	Surface Soil Texture	46
5.4	Soil Gravelliness	47
5.5	Available Water Capacity	48
5.6	Soil Slope	49
5.7	Soil Erosion	50
Chapter 6	Fertility Status	53
6.1	Soil Reaction (pH)	53
6.2	Electrical Conductivity (EC)	53
6.3	Organic Carbon (OC)	53
6.4	Available Phosphorus	53
6.5	Available Potassium	54
6.6	Available Sulphur	54
6.7	Available Boron	57
6.8	Available Iron	57
6.9	Available Manganese	57
6.10	Available Copper	57
6.11	Available Zinc	58

Chapter 7	Land Suitability for Major Crops	62
7.1	Land Suitability for Sorghum	62
7.2	Land Suitability for Maize	63
7.3	Land Suitability for Bajra	64
7.4	Land Suitability for Groundnut	65
7.5	Land Suitability for Sunflower	66
7.6	Land Suitability for Cotton	67
7.7	Land Suitability for Red gram	68
7.8	Land Suitability for Bengal gram	69
7.9	Land Suitability for Chilli	70
7.10	Land Suitability for Tomato	71
7.11	Land Suitability for Brinjal	72
7.12	Land Suitability for Onion	73
7.13	Land Suitability for Bhendi	74
7.14	Land Suitability for Drumstick	75
7.15	Land Suitability for Mulberry	76
7.16	Land Suitability for Mango	77
7.17	Land Suitability for Sapota	78
7.18	Land Suitability for Pomegranate	79
7.19	Land Suitability for Guava	80
7.20	Land Suitability for Jackfruit	81
7.21	Land Suitability for Jamun	82
7.22	Land Suitability for Musambi	83
7.23	Land Suitability for Lime	84
7.24	Land Suitability for Cashew	85
7.25	Land Suitability for Custard apple	86
7.26	Land Suitability for Amla	87
7.27	Land Suitability for Tamarind	88
7.28	Land Suitability for Marigold	89
7.29	Land Suitability for Chrysanthemum	90
7.30	Land Suitability for Jasmine	91
7.31	Land Suitability for Crossandra	92
7.32	Land Management Units (LMUs)	126
7.33	Proposed Crop Plan for Naregallu-3 Microwatershed	127
Chapter 8	Soil Health Management	130
8.1	Soil health	130
Chapter 9	Soil and Water conservation Treatment Plan	136
9.1	Treatment Plan	136
9.2	Recommended Soil and Water Conservation measures	140
9.3	Greening of microwatershed	141
	References	144
	Appendix I	I-VI
	Appendix II	VII-XII
	Appendix III	XIII-XVII

LIST OF TABLES

2.1	Mean Monthly Rainfall, PET, ½ PET at Koppal Taluk and District	5
2.2	Land Utilization in Koppal District	7
3.1	Differentiating Characteristics used for Identifying Soil Series	16
3.2	Soil map unit description of Naregallu-3 microwatershed	17
4.1	Physical and chemical characteristics of soil series identified in Naregallu-3 microwatershed	32
7.1	Soil-Site Characteristics of Naregallu-3 microwatershed	94
7.2	Land suitability criteria for Sorghum	95
7.3	Land suitability criteria for Maize	96
7.4	Land suitability criteria for Bajra	97
7.5	Land suitability criteria for Groundnut	98
7.6	Land suitability criteria for Sunflower	99
7.7	Land suitability criteria for Cotton	100
7.8	Land suitability criteria for Red gram	101
7.9	Land suitability criteria for Bengal gram	102
7.10	Land suitability criteria for Chilli	103
7.11	Land suitability criteria for Tomato	104
7.12	Land suitability criteria for Brinjal	105
7.13	Land suitability criteria for Onion	106
7.14	Land suitability criteria for Bhendi	107
7.15	Land suitability criteria for Drumstick	108
7.16	Land suitability criteria for Mulberry	109
7.17	Land suitability criteria for Mango	110
7.18	Land suitability criteria for Sapota	111
7.19	Land suitability criteria for Pomegranate	112
7.20	Land suitability criteria for Guava	113
7.21	Land suitability criteria for Jackfruit	114
7.22	Land suitability criteria for Jamun	115
7.23	Land suitability criteria for Musambi	116
7.24	Land suitability criteria for Lime	117
7.25	Land suitability criteria for Cashew	118
7.26	Land suitability criteria for Custard apple	119
7.27	Land suitability criteria for Amla	120
7.28	Land suitability criteria for Tamarind	121
7.29	Land suitability criteria for Marigold	122

7.30	Land suitability criteria for Chrysanthemum	123
7.31	Land suitability criteria for Jasmine	124
7.32	Land suitability criteria for Crossandra	125
7.33	Proposed Crop Plan for Naregallu-3 Microwatershed	128

LIST OF FIGURES

2.1	Location map of Naregallu-3 Microwatershed	3
2.2a	Granite and granite gneiss rocks	4
2.2b	Alluvium	4
2.3	Rainfall distribution in Koppal Taluk and District	6
2.4	Natural vegetation of Naregallu-3 Microwatershed	6
2.5	Different crops and cropping systems in Naregallu-3 Microwatershed	8
2.6	Current Land use – Naregallu-3 Microwatershed	9
2.7	Location of Wells and Conservation structures-Naregallu-3 Microwatershed	9
3.1	Scanned and Digitized Cadastral map of Naregallu-3 Microwatershed	13
3.2	Satellite image of Naregallu-3 Microwatershed	13
3.3	Cadastral map overlaid on IRS PAN+LISS IV merged imagery of Naregallu-3 Microwatershed	14
3.4	Location of profiles in a transect	15
3.5	Soil phase or management units-Naregallu-3 Microwatershed	21
5.1	Land Capability Classification of Naregallu-3 Microwatershed	45
5.2	Soil Depth map of Naregallu-3 Microwatershed	46
5.3	Surface Soil Texture map of Naregallu-3 Microwatershed	47
5.4	Soil Gravelliness map of Naregallu-3 Microwatershed	48
5.5	Soil Available Water Capacity map of Naregallu-3 Microwatershed	49
5.6	Soil Slope map of Naregallu-3 Microwatershed	50
5.7	Soil Erosion map of Naregallu-3 Microwatershed	51
6.1	Soil Reaction (pH) map of Naregallu-3 Microwatershed	54
6.2	Electrical Conductivity (EC) map of Naregallu-3 Microwatershed	55
6.3	Soil Organic Carbon (OC) map of Naregallu-3 Microwatershed	55
6.4	Soil Available Phosphorus map of Naregallu-3 Microwatershed	56
6.5	Soil Available Potassium map of Naregallu-3 Microwatershed	56
6.6	Soil Available Sulphur map of Naregallu-3 Microwatershed	57
6.7	Soil Available Boron map of Naregallu-3 Microwatershed	58
6.8	Soil Available Iron map of Naregallu-3 Microwatershed	59
6.9	Soil Available Manganese map of Naregallu-3 Microwatershed	59
6.10	Soil Available Copper map of Naregallu-3 Microwatershed	60
6.11	Soil Available Zinc map of Naregallu-3 Microwatershed	61
7.1	Land suitability map of Sorghum	63
7.2	Land suitability map of Maize	64
7.3	Land suitability map of Bajra	65
7.4	Land suitability map of Groundnut	66
7.5	Land suitability map of Sunflower	67

7.6	Land suitability map of Cotton	68
7.7	Land suitability map of Redgram	69
7.8	Land suitability map of Bengal gram	70
7.9	Land suitability map of Chilli	71
7.10	Land suitability map of Tomato	72
7.11	Land suitability map of Brinjal	73
7.12	Land suitability map of Onion	74
7.13	Land suitability map of Bhendi	75
7.14	Land suitability map of Drumstick	76
7.15	Land suitability map of Mulberry	77
7.16	Land suitability map of Mango	78
7.17	Land suitability map of Sapota	79
7.18	Land suitability map of Pomegranate	80
7.19	Land suitability map of Guava	81
7.20	Land suitability map of Jackfruit	82
7.21	Land suitability map of Jamun	83
7.22	Land suitability map of Musambi	84
7.23	Land suitability map of Lime	85
7.24	Land suitability map of Cashew	86
7.25	Land suitability map of Custard apple	87
7.26	Land suitability map of Amla	88
7.27	Land suitability map of Tamarind	89
7.28	Land suitability map of Marigold	90
7.29	Land suitability map of Chrysanthemum	91
7.30	Land suitability map of Jasmine	92
7.31	Land suitability map of Crossandra	93
7.32	Land Management Unit map of Naregallu-3 microwatershed	127
9.1	Soil and water conservation Plan map of Naregallu-3 Microwatershed	141

EXECUTIVE SUMMARY

The land resource inventory of Naregallu-3 microwatershed was conducted using village cadastral maps and IRS satellite imagery on 1:7920 scale. The false colour composites of IRS imagery were interpreted for physiography and these physiographic delineations were used as base for mapping soils. The soils were studied in several transects and a soil map was prepared with phases of soil series as mapping units. Random checks were made all over the area outside the transects to confirm and validate the soil map unit boundaries. The soil map shows the geographic distribution and extent, characteristics, classification, behavior and use potentials of the soils in the microwatershed.

The present study covers an area of 551 ha in Koppal taluk and district, Karnataka. The climate is semiarid and categorized as drought - prone with an average annual rainfall of 662 mm, of which about 424 mm is received during south-west monsoon, 161 mm during north-east and the remaining 77 mm during the rest of the year.

An area of 99 per cent is covered by soils and 1 per cent is by habitation and settlements. The salient findings from the land resource inventory are summarized briefly below.

- ❖ The soils belong to 11 soil series and 19 soil phases (management units) and 4 Land Management Units.*
- ❖ The length of crop growing period is <90 days and starts from 2nd week of August to 2nd week of November.*
- ❖ From the master soil map, several interpretative and thematic maps like land capability, soil depth, surface soil texture, soil gravelliness, available water capacity, soil slope and soil erosion were generated.*
- ❖ Soil fertility status maps for macro and micronutrients were generated based on the surface soil samples collected at every 320 m grid interval.*
- ❖ Land suitability for growing 31 major agricultural and horticultural crops were assessed and maps showing the degree of suitability along with constraints were generated.*
- ❖ An area of about 99 per cent is suitable for agriculture.*
- ❖ About 20 per cent of the soils are shallow to moderately shallow (25-75 cm), 58 per cent of the soils are moderately deep to deep (75-150 cm) and 21 per cent soils are very deep (>150 cm).*
- ❖ Entire cultivated area in the microwatershed has clayey soils at the surface.*
- ❖ About 89 per cent area has non-gravelly (<15% gravel) soils and 10 per cent has gravelly (15-35% gravel) soils.*
- ❖ About 20 per cent area is low (51-100 mm/m), 27 per cent area is medium to high (101-200 mm/m) and 52 per cent area is very high (>200 mm/m) in available water capacity.*

- ❖ *About 8 per cent area of the microwatershed has nearly level (0-1% slope) lands and 90 per cent area of the microwatershed has very gently sloping (1-3% slope) lands.*
- ❖ *An area of about 51 per cent area is moderately (e2) eroded and about 48 per cent area is slightly (e1) eroded.*
- ❖ *An area of about 37 per cent soils are strongly alkaline (pH 8.4-9.0) and 61 per cent soil are very strongly alkaline (pH >9.0) in soil reaction.*
- ❖ *The Electrical Conductivity (EC) of the soils in the entire cultivated area of the microwatershed is dominantly <math> < 2 \text{ dsm}^{-1}</math> indicating that the soils are non-saline.*
- ❖ *Organic carbon is medium (0.5-0.75%) in 23 per cent area and low (<math> < 0.5\%</math>) in 76 per cent area.*
- ❖ *An area of about 13 per cent is medium (23-57 kg/ha) and 86 per cent is low (<math> < 23 \text{ kg/ha}</math>) in available phosphorus.*
- ❖ *An area of about 43 per cent is medium (145-337 kg/ha) and 56 per cent is high (>337 kg/ha) in available potassium.*
- ❖ *Entire cultivated area in the microwatershed is low (<math> < 10 \text{ ppm}</math>) in available sulphur content.*
- ❖ *An area of about 49 per cent is low (<math> < 0.5 \text{ ppm}</math>), 44 per cent is medium (0.5-1.0 ppm) and 6 per cent is high (>1.0 ppm) in available boron content.*
- ❖ *An area of about 22 per cent is sufficient (>4.5 ppm) and 76 per cent is deficient (<math> < 4.5 \text{ ppm}</math>) in available iron content.*
- ❖ *Entire cultivated area of the microwatershed is sufficient (>1.0 ppm) in available manganese content.*
- ❖ *Entire cultivated area of the microwatershed is sufficient (>0.2 ppm) in available copper content.*
- ❖ *Entire cultivated area of the microwatershed is deficient (<math> < 0.6 \text{ ppm}</math>) in available zinc content.*
- ❖ *The land suitability for 31 major crops grown in the microwatershed was assessed and the areas that are highly suitable (S1) and moderately suitable (S2) are given below. It is however to be noted that a given soil may be suitable for various crops but what specific crop to be grown may be decided by the farmer looking to his capacity to invest on various inputs, marketing infrastructure, market price and finally the demand and supply position.*

Land suitability for various crops in the microwatershed

Crop	Suitability Area in ha (%)		Crop	Suitability Area in ha (%)	
	Highly suitable (S1)	Moderately suitable (S2)		Highly suitable (S1)	Moderately suitable (S2)
Sorghum	207(38)	284(52)	Sapota	34(6))	-
Maize	33(6)	459(83)	Pomegranate	34(6))	400(73)
Bajra	34(6)	457(83)	Guava	34(6))	-
Groundnut	34(6)	-	Jackfruit	34(6))	-
Sunflower	172(31)	262(48)	Jamun	34(6))	310(56)
Cotton	175(32)	318(57)	Musambi	172(31)	262(48)
Red gram	34(6)	345(63)	Lime	172(31)	262(48)
Bengalgram	206(37)	286(52)	Cashew	34(6))	-
Chilli	34(6))	-	Custard apple	207(38)	285(52)
Tomato	34(6)	-	Amla	34(6)	458(83)
Brinjal	33(6)	459(83)	Tamarind	34(6))	310(56)
Onion	33(6)	2(<1)	Marigold	34(6)	458(83)
Bhendi	33(6)	459(83)	Chrysanthemum	34(6)	458(83)
Drumstick	34(6))	400(73)	Jasmine	34(6)	57(10)
Mulberry	34(6))	345(62)	Crossandra	34(6)	166(30)
Mango	34(6))	116(21)			

- ❖ Apart from the individual crop suitability, a proposed crop plan has been prepared for the 4 identified LMUs by considering only the highly and moderately suitable lands for different crops and cropping systems with food, fodder, fibre and other horticulture crops.
- ❖ Maintaining soil-health is vital for crop production and conserves soil and land resource base for maintaining ecological balance and to mitigate climate change. For this, several ameliorative measures have been suggested for these problematic soils like saline/alkali, highly eroded, sandy soils etc.,
- ❖ Soil and water conservation treatment plan has been prepared that would help in identifying the sites to be treated and also the type of structures required.
- ❖ As part of the greening programme, several tree species have been suggested to be planted in marginal and submarginal lands, field bunds and also in the hillocks, mounds and ridges. That would help in supplementing the farm income, provide fodder and fuel, and generate lot of biomass which in turn would help in maintaining the ecological balance and contribute to mitigating the climate change.

INTRODUCTION

Soil is a finite natural resource that is central to sustainable agriculture and food security. Over the years, this precious resource is faced with the problems of erosion, salinity, alkalinity, degradation, depletion of nutrients and even decline in availability of land for agriculture. It is a known fact, that it takes thousands of years to form a few centimetres of soil, thus, soil is a precious gift of nature. The area available for agriculture is about 51 per cent of the total geographical area and more than 60 per cent of the people are still dependant on agriculture for their livelihood. However, the capacity of a soil to produce is limited and the limits to the production are set by its intrinsic characteristics, agro-climatic setting, and use and management. There is, therefore, tremendous pressure on land and water resources, which is causing decline in soil-health and stagnation in productivity. As much as 121 m ha of land is reportedly degraded which leads to impaired soil quality. It is imperative that steps are urgently taken to check and reverse land degradation without any further loss of time. The improvements in productivity will have to come from sustainable intensification measures that make the most effective use of land and water resources. Soil erosion alone has degraded about 35 lakh ha. Almost all the uncultivated areas are facing various degrees of degradation, particularly soil erosion; salinity and alkalinity has emerged as a major problem in more than 3.5 lakh ha in the irrigated areas of the state. Nutrient depletion and declining factor productivity is common in both rainfed and irrigated areas. The degradation is continuing at an alarming rate and there appears to be no systematic effort among the stakeholders to contain this process. In recent times, an aberration of weather due to climate change phenomenon has added another dimension leading to unpredictable situations to be tackled by the farmers. In this critical juncture, the challenge before us is not only to increase the productivity per unit area which is steadily declining and showing a fatigue syndrome, but also to prevent or at least reduce the severity of degradation. If the situation is not reversed at the earliest, then the sustainability of the already fragile crop production system and the overall ecosystem will be badly affected in the state.

Added to this, every year there is a significant diversion of farm lands and water resources for non-agricultural purposes. Thus, developing strategies to slow down the degradation process or reclaim the soils to normal condition and ensure sustainability of production system are the major issues today. This demands a systematic appraisal of our soil and land resources with respect to their extent, geographic distribution, characteristics, behaviour and uses potential, which is very important for developing an effective land use and cropping systems for augmenting agricultural production on a sustainable basis. The soil and land resource inventories made so far in Karnataka had limited utility because the surveys were of different types, scales and intensities carried out at different times with specific objectives. Hence, there is an urgent need to generate

detailed site-specific farm level database on various land resources for all the villages/watersheds in a time bound manner that would help to protect the valuable soil and land resources and also to stabilize the farm production.

Therefore, the land resource inventory required for farm level planning is the one which investigates all the parameters which are critical for productivity *viz.*, soils, site characteristics like slope, erosion, gravelliness and stoniness, climate, water, topography, geology, hydrology, vegetation, crops, land use pattern, animal population, socio-economic conditions, infrastructure, marketing facilities and various schemes and developmental works of the government etc. From the data collected at farm level, the specific problems and potentials of the area can be identified and highlighted, conservation measures required for the area can be planned on a scientific footing, suitability of the area for various uses can be worked out and finally viable and sustainable land use options suitable for each and every land holding can be prescribed.

The Land Resource Inventory is basically done for identifying potential and problem areas, developing sustainable land use plans, estimation of surface run off and water harvesting potential, preparation of soil and water conservation plans, land degradation/desertification etc. The Bureau is presently engaged in developing an LRI methodology using high resolution satellite remote sensing data and Digital Elevation Model (DEM) data to prepare Landscape Ecological Units (LEU) map representing agro-ecosystem as a whole. The LEU is preferred over landform as the base map for LRI. LEU is the assemblage of landform, slope and land use. An attempt was made to upscale the soil resource information from 1:250000 and 1:50000 scale to the LEU map in Goa and other states.

The land resource inventory aims to provide site-specific database for Naregallu-3 microwatershed in Koppal Taluk and District, Karnataka State for the Karnataka Watershed Development Department. The database was generated by using cadastral map of the village as a base along with high resolution IRS LISS IV and Cartosat-1 merged satellite imagery. Later, an attempt will be made to uplink this LRI data generated at 1:7920 scales under Sujala-III Project to the proposed Landscape Ecological Units (LEUs) map.

The study was organized and executed by the ICAR- National Bureau of Soil Survey and Land Use Planning, Regional Centre, Bangalore under Generation of Land Resource Inventory Data Base Component-1 of the Sujala-III Project funded by the World Bank.

GEOGRAPHICAL SETTING

2.1 Location and Extent

The Naregallu-3 Microwatershed is located in the central part of northern Karnataka in Koppal Taluk, Koppal District, Karnataka State (Fig.2.1). It comprises parts of Madhinura and Neregalla villages. It lies between $15^{\circ}23'$ – $15^{\circ}25'$ North latitudes and $76^{\circ}5'$ – $76^{\circ}8'$ East longitudes and covers an area of 551 ha. It is about 11 km from Koppal town and is surrounded by Neregalla village on the east, southeast and south, and Madhinura village on the west, northwest and northern side of the microwatershed.




Fig.2.1 Location map of Naregallu-3 Microwatershed

2.2 Geology

Major rock formation observed in the microwatershed are granite gneiss and alluvium (Figs.2.2 a & b). Granite gneisses are essentially pink to gray and are coarse to medium grained. They consist primarily of quartz, feldspar, biotite and hornblende. The

gray granite gneisses are highly weathered, fractured and fissured upto a depth of about 10 m. Dolerite dykes and quartz veins are common with variable width and found to occur in Bettageri village. The soil thickness of the alluvium generally is limited to less than a meter, except in river valleys where it is very deep extending to tens of meters. Such soils are transported and represent palaeo black soils originally formed at higher elevation, but now occupying river valleys.



Fig.2.2a Granite and granite gneiss rocks



Fig.2.2b Alluvium

2.3 Physiography

Physiographically, the area has been identified as Granite gneiss and Alluvial landscapes based on geology. The microwatershed area has been further divided into mounds/ridges, summits, side slopes and very gently sloping uplands and nearly level plains based on slope and its relief features. The elevation ranges from 527-554 m in the gently sloping uplands. The mounds and ridges are mostly covered by rock outcrops.

2.4 Drainage

The area is drained by several small seasonal streams that join Hire *halla* and Chenna *halla* along its course. Though, the streams are not perennial, during rainy season they carry large quantities of rain water. The microwatershed has only few small tanks which are not able to store the water flowing during the rainy season. Due to this, the ground water recharge is very much affected in the villages. This is reflected in the failure of many bore wells in the villages. If the available rain water is properly harnessed by constructing tanks and recharge structures at appropriate places in the villages, then the drinking and irrigation needs of the area can be easily met. The drainage network is dendritic to sub parallel.

2.5 Climate

The district falls under semiarid tract of the state and is categorized as drought - prone with total annual rainfall of 662 mm (Table 2.1) Of this, a maximum of 424 mm precipitation takes place during south–west monsoon period from June to September, north-east monsoon contributes about 161 mm and prevails from October to early December and the remaining 77 mm received during the rest of the year. The winter season is from December to February. During April and May, the temperatures reach up to 45°C and in December and January, the temperatures will go down to 16°C. Rainfall distribution is shown in Figure 2.3. The average Potential Evapo Transpiration (PET) is 145 mm and varies from a low of 101 mm in December and 193 mm in the months of May. The PET is always higher than precipitation in all the months except in the month of September. Generally, the Length of crop Growing Period (LGP) is <90 days and starts from 2nd week of August to 2nd week of November.

Table 2.1 Mean Monthly Rainfall, PET, 1/2 PET at Koppal Taluk and District

Sl. No.	Months	Rainfall	PET	1/2 PET
1	January	1.60	116.70	58.35
2	February	1.50	129.20	64.60
3	March	14.10	169.80	84.90
4	April	18.10	180.60	90.30
5	May	41.60	193.50	96.75
6	June	85.80	167.90	83.95
7	July	72.10	156.20	78.10
8	August	110.50	152.50	76.25
9	September	155.60	138.50	69.25
10	October	116.30	122.30	61.15
11	November	36.00	106.40	53.20
12	December	9.10	101.00	50.50
TOTAL		662.30	144.55	




Fig. 2.3 Rainfall distribution in Koppal Taluk and District

2.6 Natural Vegetation

The natural vegetation is sparse comprising few tree species, shrubs and herbs. The mounds, ridges and boulders occupy sizeable areas which are under thin to moderately thick forest vegetation. Still, there are some remnants of the past forest cover which can be seen in patches in some ridges and hillocks in the microwatershed (Fig 2.4).

Apart from the continuing deforestation, the presence of large population of goats, sheep and other cattle in the microwatershed is causing vegetative degradation of whatever little vegetation left in the area. The uncontrolled grazing has left no time for the regeneration of the vegetative cover. This leads to the accelerated rate of erosion on the hill slopes, resulting in the formation of deep gullies in the foot slopes and eventually resulting in the heavy siltation of few tanks and reservoirs in the microwatershed.



Fig 2.4 Natural vegetation of Naregallu-3 microwatershed

2.7 Land Utilization

About 91 per cent area (Table 2.2) in Koppal district is cultivated at present and about 17 per cent of the area is sown more than once. An area of about 3 per cent is currently barren. Forests occupy a small area of about 5 per cent and the tree cover is in a very poor state. Most of the mounds, ridges and bouldery areas have very poor vegetative cover. Major crops grown in the area are sorghum, maize, bajra, cotton, safflower, sunflower, red gram, horse gram, onion, mulberry, pomegranate, sugarcane, Bengalgram, marigold and groundnut (Fig 2.5). While carrying out land resource inventory, the land use/land cover particulars are collected from all the survey numbers and a current land use map of the microwatershed is prepared. The current land use map prepared shows the arable and non-arable lands, other land uses and different types of crops grown in the area. The current land use map of Naregallu-3 Microwatershed is presented in Fig.2.6. Simultaneously, enumeration of existing wells (bore wells and open wells) and other soil and water conservation structures in the microwatershed is made and their location in different survey numbers is marked on the cadastral map. Map showing the location of wells and conservation structures in Naregallu-3 Microwatershed is given Fig.2.7.

Table 2.2 Land Utilization in Koppal District

Sl. No.	Agricultural land use	Area (ha)	Per cent
1	Total geographical area	552495	
2	Total cultivated area	500542	90.6
3	Area sown more than once	92696	16.8
4	Trees and groves	210	0.04
5	Cropping intensity	-	118
6	Forest	29451	5.33
7	Cultivable wasteland	2568	0.46
8	Permanent Pasture land	14675	2.66
9	Barren land	16627	3.01
10	Non agricultural land	40591	7.35
11	Current fallow	19660	3.56



Groundnut



Sunflower



Cotton



Red gram



Onion



Marigold

Fig.2.5 Different crops and cropping systems in Naregallu-3 Microwatershed




Fig.2.6 Current Land Use – Naregallu-3 Microwatershed




Fig.2.7 Location of wells and conservation structures- Naregallu-3 Microwatershed

SURVEY METHODOLOGY

The purpose of land resource inventory is to delineate similar areas (soil series and phases), which respond or expected to respond similarly for a given level of management. This was achieved in Naregallu-3 Microwatershed by the detailed study of all the soil characteristics (depth, texture, colour, structure, consistence, coarse fragments, porosity, soil reaction, soil horizons etc.) and site characteristics (slope, erosion, drainage, occurrence of rock fragments etc.) followed by grouping of similar areas based on soil-site characteristics into homogeneous (management units) units and showing their extent and geographic distribution on the microwatershed cadastral map. The detailed soil survey at 1:7920 scale was carried out in 551 ha area. The methodology followed for carrying out land resource inventory was as per the guidelines given in Soil Survey Manual (IARI, 1971; Soil Survey Staff, 2006; Natarajan *et al.*, 2015) which is briefly described below.

3.1 Base Maps

The detailed survey of the land resources occurring in the microwatershed was carried out by using digitized cadastral map and satellite imagery as base supplied by the KRSRSAC. The cadastral map shows field boundaries with their survey numbers, location of tanks, streams and other permanent features of the area (Fig. 3.1). Apart from the cadastral map, remote sensing data products from Cartosat-1 and LISS IV merged at the scale of 1:7920 were used in conjunction with the cadastral map to identify the geology, landscapes, landforms and other surface features. The imagery helped in the identification and delineation of boundaries between hills, uplands and lowlands, water bodies, forest and vegetated areas, roads, habitations and other cultural features of the area (Fig.3.2).The cadastral map was overlaid on the satellite imagery (Fig.3.3) that helps to identify the parcel boundaries and other permanent features. Apart from cadastral maps and images, toposheets of the area (1:50,000 scale) were used for initial traversing, identification of geology, landscapes and landforms, drainage features, present land use and also for selection of transects in the microwatershed.

3.2 Image Interpretation for Physiography

False Colour Composites (FCC) of Cartosat-I and LISS-IV merged satellite data covering the microwatershed area was visually interpreted using image interpretation elements and all the available collateral data with local knowledge. The delineated physiographic boundaries were transferred on to a cadastral map overlaid on satellite imagery. Physiographically, the area has been identified as granite gneiss and alluvial landscapes and is divided into landforms such as uplands, summits and very gently sloping based on slope. They were further subdivided into physiographic/ image

interpretation units based on image characteristics. The image interpretation legend for physiography is given below.

Image Interpretation Legend for Physiography

G- Granite gneiss landscape

- G1 Hills/ Ridges/ Mounds
 - G11 Summits
 - G12 Side slopes
 - G121 Side slopes with dark grey tones
- G2 Uplands
 - G21 Summits
 - G22 Gently sloping uplands
 - G221 Gently sloping uplands, yellowish green (eroded)
 - G222 Gently sloping uplands, yellowish white (severely eroded)
 - G23 Very gently sloping uplands
 - G231 Very gently sloping uplands, yellowish green
 - G232 Very gently sloping uplands, medium green and pink
 - G233 Very gently sloping uplands, pink and green (scrub land)
 - G234 Very gently sloping uplands, medium greenish grey
 - G235 Very gently sloping uplands, yellowish white (eroded)
 - G236 Very gently sloping uplands, dark green
 - G237 Very gently sloping uplands, medium pink (coconut garden)
 - G238 Very gently sloping uplands, pink and bluish white (eroded)

DSe Alluvial landscape

DSe 1 Summit

- DSe 11 Nearly level Summit with dark grey tone
- DSe 12 Nearly level Summit with medium grey tone
- DSe 13 Nearly level Summit with whitish grey tone
- DSe 14 Nearly level Summit with whitish tone (Calcareousness)
- DSe 15 Nearly level Summit with pinkish grey tone
- DSe 16 Nearly level Summit with medium pink tone
- DSe 17 Nearly level Summit with bluish white tone
- DSe 18 Nearly level Summit with greenish grey tone

DSe 2 Very gently sloping

- DSe 21 Very gently sloping, whitish tone
- DSe 22 Very gently sloping, greyish pink tone
- DSe 23 Very gently sloping, whitish grey tone
- DSe 24 Very gently sloping, medium grey tone
- DSe 25 Very gently sloping, medium pink tone
- DSe 26 Very gently sloping, dark grey tone
- DSe 27 Very gently sloping, bluish grey tone
- DSe 28 Very gently sloping, greenish grey tone
- DSe 29 Very gently sloping, Pinkish grey




Fig 3.1 Scanned and Digitized Cadastral map of Naregallu-3 Microwatershed




Fig.3.2 Satellite Image of Naregallu-3 Microwatershed




Fig.3.3 Cadastral map overlaid on IRS PAN+LISS IV merged imagery of Naregallu-3 Microwatershed

3.3 Field Investigation

The field boundaries and survey numbers given on the cadastral sheet were located on the ground by following permanent features like roads, cart tracks, *nallas*, streams, tanks etc., and wherever changes were noticed, they were incorporated on the microwatershed cadastral map. Preliminary traverse of the microwatershed was carried out with the help of cadastral map, imagery and toposheets. While traversing, landforms and physiographic units identified were checked and preliminary soil legend was prepared by studying soils at few selected places. Then, intensive traversing of each physiographic unit like uplands and plains was carried out. Based on the variability observed on the surface, transects (Fig 3.4) were selected across the slope covering all the landform units in the microwatershed (Natarajan and Dipak Sarkar, 2010).




Fig: 3.4. Location of profiles in a transect

In the selected transect, soil profiles (Fig.3.4) were located at closely spaced intervals to take care of any change in the land features like break in slope, erosion, gravel, stones etc. In the selected sites, profiles (vertical cut showing the soil layers from surface to the rock) were opened up to 200 cm or to the depth limited by rock or hard substratum and studied in detail for all their morphological and physical characteristics. The soil and site characteristics were recorded for all profile sites on a standard proforma as per the guidelines given in USDA Soil Survey Manual (Soil Survey Staff, 2012). Apart from the transect study, profiles were also studied at random, almost like in a grid pattern, outside the transect areas to validate the soil map unit boundaries.

Based on the soil characteristics, the soils were grouped into different soil series. Soil series is the most homogeneous unit having similar horizons and properties and behaves similarly for a given level of management. Soil depth, texture, colour, kind of horizon and horizon sequence, amount and nature of gravel present, calcareousness, nature of substratum etc, were used as the major differentiating characteristics for identifying soil series occurring in the area. The differentiating characteristics used for identifying the soil series are given in Table 3.1. Based on the above characteristics, 11 soil series were identified in Naregallu-3 Microwatershed.

**Table 3.1 Differentiating Characteristics used for identifying Soil Series
(Characteristics are of Series Control Section)**

Sl. No.	Soil Series	Depth (cm)	Colour (moist)	Texture	Gravel (%)	Horizon sequence	Calcareousness
SOILS OF GRANITE GNEISS LANDSCAPE							
1	Muradi (MRD)	>150	2.5YR3/6,4/6,5/6, 5/8	scl	<15	Ap-Bt	-
2	Ranatur (RTR)	>150	2.5YR2.5/3,2.5/4,3/3, 4/6	c	<15	Ap-Bt	-
SOILS OF ALLUVIAL LANDSCAPE							
3	Muttal (MTL)	25-50	10YR3/2,3/3,4/2 7.5YR3/2,3/3,6/4	gc	15-35	Ap-Bw-Ck	e-ev
4	Ravanaki (RNK)	50-75	7.5YR3/2,3/3,5/2,5/3 10YR3/1,3/2,4/1,4/2, 5/1,6/1	c	<15	Ap-Bw-Cr	e-ev
5	Dambarahalli (DRL)	75-100	10YR 2/1, 3/1, 4/3	c	<15	Ap-Bss	e-es
6	Narasapura (NSP)	75-100	10 YR 3/1, 3/2, 4/2,	c	<15	Ap-Bw-Cr	e-es
7	Gatareddihal (GRH)	100-150	10YR2/1,3/1 2.5Y 4/3, 5/4	c	<15	Ap-Bss-Bck-Cr	es
8	Handrala (HDL)	100-150	10 YR 2/1, 3/1,4/1,	c	<15	Ap-Bss-Ck	es
9	Kavalur (KVR)	100-150	10 YR 2/2, 3/1, 3/2, 3/3, 4/4	c	<15	Ap-Bss-Bck-Cr	es-ev
10	Budagumpa (BGP)	>150	7.5YR3/2,5/1 10YR4/1,4/4	c	<15	Ap-Bw	es
11	Bardur (BDR)	>150	10YR 2/1, 3/1, 3/2	c	<15	Ap-Bss	es

3.4 Soil Mapping

The area under each soil series was further separated into soil phases and their boundaries delineated on the cadastral map based on the variations observed in the texture of the surface soil, slope, erosion, presence of gravel, stoniness etc. A soil phase is a subdivision of soil series based mostly on surface features that affect its use and management. The soil mapping units are shown on the map (Fig.3.5) in the form of symbols. During the survey many soil profile pits, few minipits and a few auger bores representing different landforms occurring in the microwatershed were studied. In addition to the profile study, spot observations in the form of minipits, road cuts, terrace cuts etc., were studied to validate the soil boundaries on the soil map. The soil map shows the geographic distribution and area extent of 19 mapping units representing 11 soil series occurring in the microwatershed. The soil map unit (soil legend) description is presented in Table 3.2. The soil phase map (management units) shows the distribution of 19 phases mapped in the microwatershed. Each mapping unit (soil phase) delineated on the map has similar soil and site characteristics. In other words, all the farms or survey numbers

included in one phase will have similar management needs and have to be treated accordingly.

3.5 Laboratory Characterization

Soil samples for each series were collected from representative master profiles for laboratory characterization by following the methods outlined in the Laboratory Manual (Sarma *et al*, 1987). Surface soil samples collected in the year 2018 from Naregallu-3 farmer's fields (51 samples) for fertility status (major and micronutrients) at 320 m grid interval were analyzed in the laboratory (Katyal and Rattan, 2003). By linking the soil fertility data to the survey numbers through GIS, soil fertility maps were generated using Kriging method for the microwatershed.

3.6 Land Management Units (LMUs)

The 19 soil phases identified and mapped in the microwatershed were regrouped into 4 Land Management Units (LMU's) for the purpose of preparing a Proposed Crop Plan for sustained development of the microwatershed. The database (soil phases) generated under LRI was utilized for identifying Land Management Units (LMU's) based on the management needs. One or more than one soil site characteristic having influence on the management have been chosen for identification and delineation of LMUs. For Naregallu-3 Microwatershed, five soil and site characteristics, namely soil depth, soil texture, slope erosion and gravel content have been considered for defining LMUs. The land management units are expected to behave similarly for a given level of management.

Table 3.2 Soil map unit description of Naregallu-3 Microwatershed

Soil map unit No*	Soil Series	Soil Phase Symbol	Mapping Unit Description	Area in ha (%)
SOILS OF GRANITE AND GRANITE GNEISS LANDSCAPE				
	MRD		Muradi soils are very deep (>150 cm), well drained, have red to dark red, sandy clay loam soils occurring on nearly level to gently sloping uplands under cultivation	33(5.91)
283		MRDiB2	Sandy clay surface, slope 1-3%, moderate erosion	33(5.91)
	RTR		Ranatur soils are very deep (>150 cm), well drained, have dark reddish brown to dark red, clay soils occurring on nearly level to very gently sloping uplands under cultivation	2(0.29)
288		RTRiB2	Sandy clay surface, slope 1-3%, moderate erosion	2(0.29)
SOILS OF ALLUVIAL LANDSCAPE				
	MTL		Muttal soils are shallow (25-50 cm), well drained, have very dark grayish brown to dark brown, calcareous, black gravelly clay soils occurring on nearly level to gently sloping plains under cultivation	53(9.56)
301		MTLmB2	Clay surface, slope 1-3%, moderate erosion	53(9.56)
	RNK		Ravanaki soils are moderately shallow (50-75 cm),	57(10.3)

Soil map unit No*	Soil Series	Soil Phase Symbol	Mapping Unit Description	Area in ha (%)
			moderately well drained, have dark brown to very dark grayish brown and dark gray, sodic, calcareous clay black soils occurring on nearly level to very gently sloping plains under cultivation	
333		RNKmB1	Clay surface, slope 1-3%, slight erosion	36(6.45)
336		RNKmB2	Clay surface, slope 1-3%, moderate erosion	21(3.85)
	DRL		Dambarahalli soils are moderately deep (75-100 cm), moderately well drained, have dark brown to very dark gray, calcareous, black cracking clay soils occurring on nearly level to very gently sloping plains under cultivation	55(10.02)
350		DRLmB2	Clay surface, slope 1-3%, moderate erosion	55(10.02)
	NSP		Narasapura soils are moderately deep (75-100 cm), moderately well drained, have dark grayish brown to very dark grayish brown and very dark gray, sodic, calcareous, black cracking clay soils occurring on nearly level to very gently sloping plains under cultivation	34(6.3)
359		NSPmA1	Clay surface, slope 0-1% slight erosion	9 (1.7)
362		NSPmB2	Clay surface, slope 1-3%, moderate erosion	25(4.6)
	GRH		Gatareddihal soils are deep (100-150 cm), moderately well drained, have light olive brown to very dark gray, sodic, calcareous, black cracking clay soils occurring on nearly level to very gently sloping plains under cultivation	58(10.49)
371		GRHmB1	Clay surface, slope 1-3%, slight erosion	9(1.68)
373		GRHmB2	Clay surface, slope 1-3%, moderate erosion	18(3.24)
374		GRHmB2g1	Clay surface, slope 1-3%, moderate erosion, gravelly (15-35%)	31(5.57)
	HDL		Handrala soils are deep (100-150 cm), moderately well drained, have dark gray to very dark gray, calcareous, black cracking clay soils occurring on very gently sloping plains under cultivation	55(10)
382		HDLmB2	Clay surface, slope 1-3%, moderate erosion	44(7.93)
378		HDLmA1	Clay surface, slope 0-1% ,slight erosion	11(2.07)
	KVR		Kavalur soils are deep (100-150 cm), moderately well drained, have dark yellowish brown to very dark grayish brown, calcareous black cracking clay soils occurring on nearly level to very gently sloping plains under cultivation	116 (21.04)
384		KVRiB2	Sandy clay surface, slope 1-3%, moderate erosion	1(0.14)
388		KVRmB1	Clay surface, slope 1-3%, slight erosion	90(16.37)
389		KVRmB1g1	Clay surface, slope 1-3%, slight erosion, gravelly (15-35%)	25(4.53)
	BGP		Budagumpa soils are very deep (>150 cm), moderately well drained, have dark yellowish brown to dark brown and dark gray, sodic, calcareous, black clay soils occurring on nearly level to very gently sloping plains under cultivation	25(4.6)

Soil map unit No*	Soil Series	Soil Phase Symbol	Mapping Unit Description	Area in ha (%)
396		BGPmB1	Clay surface, slope 1-3%, slight erosion	25(4.6)
	BDR		Bardur soils are very deep (>150 cm), moderately well drained, have very dark grayish brown to very dark gray, calcareous, black cracking clay soils occurring on nearly level to very gently sloping plains under cultivation	56(10.18)
428		BDRmA1	Clay surface, slope 0-1%, slight erosion	26(4.71)
430		BDRmB1	Clay surface, slope 1-3%, slight erosion	30(5.47)
1000		Others	Water body	7(1.3)

*Soil map unit numbers are continuous for the taluk, not for the microwatershed




Fig 3.5 Soil Phase or Management Units - Naregallu-3 Microwatershed

THE SOILS

Detailed information pertaining to the nature, extent and distribution of different kinds of soils occurring in Naregallu-3 Microwatershed is provided in this chapter. The microwatershed area has been identified as granite gneiss and alluvial landscapes based on geology. In all, 11 soil series are identified. Soil formation is the result of the combined effect of environmental and terrain factors that are reflected in soil morphology. The soil formation is dominantly influenced by the parent material, climate, time and relief.

A brief description of each of the 11 soil series identified followed by 19 soil phases (management units) mapped (Fig. 3.5) are furnished below. The physical and chemical characteristics of soil series identified in Naregallu-3 microwatershed are given in Table 4.1 along with soil classification. The soils in any one map unit differ from place to place in their depth, texture, slope, gravelliness, erosion or any other site characteristic that affect management. The soil phase map can be used for identifying the suitability of areas for growing specific crops or for other alternative uses and also for deciding the type of conservation structures needed. The detailed information on soil and site-characteristics like soil depth, surface soil texture, slope, erosion, gravelliness, AWC, LCC etc, with respect to each of the soil phase identified is given village/survey number wise for the microwatershed in Appendix-I.

4.1 Soils of Granite gneiss landscape

In this landscape, 2 soil series are identified and mapped. Of these, Muradi (MRD) series occupies major area 33 ha (6%) and Ranatur (RTR) 2 ha (<1%). The brief description of each soil series along with the soil phases identified and mapped is given below.

4.1.1 Muradi (MRD) Series: Muradi soils are very deep (>150 cm), well drained, have red to dark red, sandy clay loam soils. They have developed from weathered granite gneiss and occur on very gently sloping uplands under cultivation. The Muradi series has been classified as a member of the fine-loamy, mixed, isohyperthermic family of Typic Paleustalfs.

The thickness of the solum is more than 150 cm. The thickness of A-horizon ranges from 16 to 26 cm. Its colour is in 2.5 YR hue with value 3 and chroma 4. The texture is Sandy loam. The thickness of B horizon ranges from 126 to 160 cm. Its colour is in 2.5 YR hue with value 3 to 5 and chroma 6 to 8. Its texture is sand clay loam to sandy clay. The available water capacity is medium (101-150 mm/m). Only one soil phase was identified and mapped.



Landscape Soil Profile Characteristics of Muradi (MRD) Series

4.1.2 Ranatur (RTR) Series: Ranatur soils are very deep (> 150 cm), well drained, have dark reddish brown to dark red, clay soils. They have developed from weathered granite gneiss and occur on very gently sloping uplands. The Ranatur series has been classified as a member of the fine, mixed, isohyperthermic family of Rhodic Paleustalfs.

The thickness of the solum is more than 150 cm. The thickness of A-horizon ranges from 8 to 14 cm. Its colour is in 5 YR and 2.5 YR hue with value 2.5 to 4 and chroma 3 to 6. The texture varies from sandy loam to sand clay. The thickness of B horizon is more than 150 cm. Its colour is in 2.5 YR hue with value 2.5 to 3 and chroma 3 to 6. Its texture is clay. The available water capacity is high (150-200 mm/m). Only one soil phase was identified and mapped.



Landscape and soil profile characteristics of Ranatur (RTR) Series

4.2 Soils of Alluvial landscape

In this landscape, 9 soil series were identified and mapped. Of these, Kavalur (KVR) series occupies major area 116 ha (21%) followed by Gatareddihal (GRH) 58 ha (10%), Ravanaki (RNK) 57 ha (10%), Bardur (BDR) 56 ha (10%), Dambarahalli (DRL) 55 ha (10%), Handrala (HDL) 55 ha (10%), Muttal (MTL) 53 ha (10%), Narasapura (NSP) 34 ha (6%) and Budagumpa (BGP) 25 ha (5%). The brief description along with the soil phases identified and mapped is given below.

4.2.1 Muttal (MTL) Series: Muttal soils are shallow (25-50 cm), well drained, have dark brown to very dark grayish brown, calcareous gravelly clay soils. They have developed from alluvium and occur on nearly level to very gently sloping uplands. The Muttal series has been classified as a member of the clayey, mixed (calc), isohyperthermic family of (Paralithic) Haplustepts.

The thickness of the solum ranges from 30 to 50 cm. The thickness of A-horizon ranges from 15 to 18 cm. Its colour is in 7.5 YR and 10 YR hue with value 2 to 3 and chroma 2.5 to 4. The texture varies from sandy clay to clay with 10 to 15 per cent gravel. The thickness of B horizon ranges from 18 to 32 cm. Its colour is in 10 YR and 7.5 YR hue with value 2 to 6 and chroma 2 to 4. Its texture is sandy clay to clay. The available water capacity is low (51-100 mm/m). Only one soil phase was identified and mapped.



Landscape and soil profile characteristics of Muttal (MTL) Series

4.2.2 Ravanaki (RNK) Series: Ravanaki soils are moderately shallow (50-75 cm), well drained, have dark brown to very dark grayish brown, sodic, calcareous clay soils. They have developed from alluvium and occur on nearly level to very gently sloping uplands. The Ravanaki series has been classified as a member of the very-fine, smectitic (calc), isohyperthermic family of Typic Haplustepts.

The thickness of the solum ranges from 50 to 75 cm. The thickness of A-horizon ranges from 15 to 20 cm. Its colour is in 7.5 YR and 10 YR hue with value 2 to 3 and

chroma 2.5 to 4. The texture varies from sandy clay to clay with 10 to 15 per cent gravel. The thickness of B horizon ranges from 35 to 60 cm. Its colour is in 10 YR and 7.5 YR hue with value 2 to 6 and chroma 2 to 4. Its texture is sandy clay to clay with gravel content of 10 to 20 per cent. The available water capacity is low (51-100 mm/m). Two soil phases were identified and mapped.



Landscape and Soil Profile Characteristics of Ravanaki (RNK) Series

4.2.3 Dambarahalli (DRL) Series: Dambarahalli soils are moderately deep (75-100 cm), moderately well drained, have black and very dark gray to dark brown, calcareous cracking clay soils. They have developed from alluvium and occur on very gently to gently sloping uplands under cultivation. The Dambarahalli series has been classified as a member of the very-fine, smectitic (calc), isohyperthermic family of Typic Haplusterts.

The thickness of the solum ranges from 75 to 99 cm. The thickness of A-horizon ranges from 13 to 24 cm. Its colour is in 10 YR hue with value 3 to 4 and chroma 1 to 2. The texture is clay. The thickness of B horizon ranges from 54 to 85 cm. Its colour is in 10 YR hue with value 2 to 4 and chroma 1 to 3. Its texture is clay and is calcareous. The available water capacity is high (151-200 mm/m). Only one soil phase was identified and mapped.



Landscape and soil profile characteristics of Dambarahalli (DRL) Series.

4.2.4 Narsapura (NSP) series: Narsapura soils are moderately deep (75-100 cm), moderately well drained, have dark grayish brown to very dark grayish brown and very dark gray, sodic, calcareous black cracking clay soils. They have developed from alluvium and occur on very gently sloping uplands. The Narsapura series has been classified as a member of the very-fine, smectitic (calc), isohyperthermic family of Vertic Haplustepts.


The thickness of the solum is 76 to 98 cm. The thickness of A-horizon ranges from 15 to 19 cm. Its colour is in 10 YR hue with value 3 and chroma 1 to 2. The texture is clay with no gravel. The thickness of B horizon ranges from 57 to 83 cm. Its colour is in 10 YR hue with value 3 to 5 and chroma 1 to 3. Its texture is clay and is calcareous. The available water capacity is medium (101-150 mm/m). Two soil phases were identified and mapped.



Landscape and soil profile characteristics of Narsapura (NSP) series

4.2.5 Gatareddihal (GRH) Series: Gatareddihal soils are deep (100-150 cm), moderately well drained, have black or dark grey to light olive brown, sodic, calcareous cracking clay soils. They are developed from alluvium and occur on nearly level to very gently sloping uplands under cultivation. The Gatareddihal soil series has been classified as a member of the very-fine, smectitic (calc), isohyperthermic family of Sodic Haplusterts.

The thickness of the solum ranges from 102 to 149 cm. The thickness of A-horizon ranges from 12 to 19 cm. Its colour is in 7.5 YR, 10 YR hue with value 3 to 4 and chroma 1 to 6. The texture is sandy clay loam to clay. The thickness of B-horizon ranges from 86 to 117 cm. Its colour is in 10 YR and 7.5 YR hue with value 3 and chroma 2 to 6. Texture is clay with less than 15 per cent gravel. The available water capacity is very high (>200 mm/m). Three soil phases were identified and mapped.



Landscape and soil profile characteristics of Gatareddihal (GRH) Series

4.2.6 Handrala (HDL) Series: Handrala soils are deep (100-150 cm), moderately well drained, have black, very dark brown to dark gray, calcareous cracking clay soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands. The Handrala series has been classified as a member of the very-fine, smectitic (calc), isohyperthermic family of Typic Haplusterts.

The thickness of the solum ranges from 102 to 149 cm. The thickness of A-horizon ranges from 14 to 26 cm. Its colour is in 10 YR hue with value 3 and chroma 1. The texture is clay. The thickness of B horizon ranges from 103 to 127 cm. Its colour is in 10 YR hue with value 2 to 4 and chroma 1 to 2. Texture is dominantly clay. The available water capacity is very high (>200 mm/m). Two soil phases were identified and mapped.



Landscape and Soil Profile Characteristics of Handrala (HDL) Series

4.2.7 Kavalur (KVR) series: Kavalur soils are deep (100-150 cm), moderately well drained, have dark yellowish brown to very dark brown and very dark gray, calcareous black cracking clay soils. They have developed from alluvium and occur on very gently sloping uplands. The Kavalur series has been classified as a member of the fine, smectitic (calc), isohyperthermic family of Typic Haplusterts.

The thickness of the solum is 113 to 143 cm. The thickness of A-horizon ranges from 9 to 24 cm. Its colour is in 10 YR hue with value 3 and chroma 1. The texture is clay with no gravel. The thickness of B horizon ranges from 89 to 134 cm. Its colour is in 10 YR hue with value 3 and chroma 1. Its texture is clay. The available water capacity is very high (>200 mm/m). Three soil phases were identified and mapped.



Landscape and soil profile characteristics of Kavalur (KVR) series

4.2.8 Budagumpa (BGP) Series: Budagumpa soils are very deep (>150 cm), well drained, have black sodic, calcareous clay soils. They have developed from alluvium and occur on very gently sloping uplands under cultivation. The Budagumpa series has been classified as a member of the fine, mixed (calc), isohyperthermic family of Typic Haplustepts.


The thickness of the solum is more than 150 cm. The thickness of A-horizon ranges from 16 to 26 cm. Its colour is in 7.5 YR and 10 YR hue with value 2 to 3 and chroma 2 to 4. The texture varies from sandy clay to clay with 5 to 10 per cent gravel. The thickness of B horizon ranges from 130 to 160 cm. Its colour is in 10 YR and 7.5 YR hue with value 3 to 5 and chroma 1 to 4. Its texture is clay with gravel content of <15 per cent. These soils are calcareous that increase with depth. The available water capacity is very high (>200 mm/m). Only one soil phase was identified and mapped.



Landscape and Soil Profile Characteristics of Budagumpa (BGP) Series

4.2.9 Bardur (BDR) Series: Bardur soils are very deep (>150 cm), moderately well drained, have very dark grayish brown to very dark gray, calcareous black cracking clay soils. They are developed from alluvium and occur on nearly level to very gently sloping plains under cultivation. The Bardur series has been classified as a member of the very-fine, smectitic (calcareous), isohyperthermic family of Typic Haplusterts.

The thickness of the solum is more than 150 cm. The thickness of A-horizon ranges from 15 to 19 cm. Its colour is in 10 YR hue with value 2 and chroma 1 with clay texture. The thickness of B horizon ranges from 146 to 180 cm. Its colour is in 10 YR hue with value 2 to 3 and chroma 1 to 2. Its texture is clay and is calcareous with less than 15 per cent gravel. The available water capacity is very high (>200 mm/m). Two soil phases were identified and mapped.



Landscape and soil profile characteristics of Bardur (BDR) Series

Table: 4.1 Physical and Chemical Characteristics of Soil Series identified in Naregallu-3 Microwatershed

Soil Series: Ranatur (RTR), **Pedon:** RM-87

Location: 13°21'49.0"N, 76°38'06"E, (4B3D4L2a), J C Pura village, Chikkanayakanahalli taluk, Tumakuru district

Analysis at: NBSS&LUP, Regional Centre, Bengaluru **Classification:** Fine, mixed, isohyperthermic Rhodic Paleustalfs

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-17	Ap	84.16	9.46	6.38	2.22	18.57	26.14	24.32	12.92	-	ls	-	-
17-47	Bt1	51.14	8.30	40.56	1.66	13.49	14.52	13.59	7.88	-	sc	-	-
47-89	Bt2	51.99	11.01	37.00	1.94	13.99	15.32	13.18	7.56	-	sc	-	-
89-123	Bt3	51.58	9.07	39.35	3.47	14.50	14.61	11.64	7.35	-	sc	-	-
123-152	Bt4	47.89	8.88	43.23	2.27	12.36	14.21	11.12	7.93	-	sc	-	-
152-198	Bt5	43.37	13.17	43.45	2.48	9.83	13.25	10.87	6.94	-	c	-	-

Depth (cm)	pH (1:2.5)			E.C. (1:2.5) dS m ⁻¹	O.C. %	CaCO ₃ %	Exchangeable bases					CEC	CEC/Clay	Base saturation %	ESP %
	Water	CaCl ₂	M KCl				Ca	Mg	K	Na	Total				
0-17	5.08	-	-	0.03	0.52	0.00	3.68	0.72	0.06	0.19	4.65	9.21	1.44	50.50	2.06
17-47	6.28	-	-	0.03	0.48	0.00	3.93	0.72	0.08	0.07	4.80	7.92	0.20	60.59	0.94
47-89	6.42	-	-	0.03	0.40	0.00	4.40	0.74	0.08	0.06	5.28	7.52	0.20	70.15	0.79
89-123	6.50	-	-	0.02	0.32	0.00	4.44	0.76	0.09	0.07	5.36	7.82	0.20	68.58	0.93
123-152	6.52	-	-	0.02	0.28	0.00	4.40	0.71	0.09	0.07	5.26	8.22	0.19	64.00	0.81
152-198	7.09	-	-	0.02	0.24	0.00	6.10	0.98	0.10	0.20	7.38	9.60	0.22	76.89	2.09

Contd....

Series Name: Muttal (MTL), **Pedon:** RM-13

Location: 15°14'30.8"N, 75°56'50.6"E, Gatareddihalla village, Koppal Taluk and District

Analysis at: NBSS&LUP, Regional Centre, Bangalore. **Classification:** Clayey, mixed (calc), isohyperthermic (Paralithic) Haplustepts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-20	Ap	39.05	13.74	47.21	3.05	5.05	8.21	14.63	8.11	15-30	c	29.95	17.94
20-34	Bwk	28.77	19.57	51.66	4.81	4.71	4.92	9.09	5.24	10	c	33.44	21.56

Depth (cm)	pH (1:2.5)			E.C. (1:2.5) dS m ⁻¹	O.C. %	CaCO ₃ %	Exchangeable bases					CEC	CEC/Clay	Base saturation %	ESP %
	Water	CaCl ₂	M KCl				Ca	Mg	K	Na	Total				
							cmol kg ⁻¹								
0-20	8.27	-	-	0.202	0.79	6.10	-	-	0.62	0.25	-	36.64	0.78	-	0.69
20-34	8.36	-	-	0.177	0.99	23.04	-	-	0.29	0.38	-	39.60	0.77	-	0.96

Contd....

Series Name: Ravanaki (RNK), **Pedon:** RM-20

Location: 15°14'22.7"N, 75°57'45.8"E, Gatareddihalla village, Koppal Taluk and District

Analysis at: NBSS&LUP, Regional Centre, Bangalore. **Classification:** Very-fine, smectitic (calc), isohyperthermic Typic Haplustepts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-28	Ap	24.43	17.76	57.81	5.30	3.89	3.78	7.14	4.32	20	c	41.40	29.60
28-55	Bw	18.77	15.59	65.64	2.74	3.73	2.85	4.83	4.61	10	c	46.71	35.18

Depth (cm)	pH (1:2.5)			E.C. (1:2.5) dS m ⁻¹	O.C. %	CaCO ₃ %	Exchangeable bases					CEC	CEC/Clay	Base saturation %	ESP %
	Water	CaCl ₂	M KCl				Ca	Mg	K	Na	Total				
							cmol kg ⁻¹								
0-28	8.86	-	-	0.483	0.63	15.48	-	-	0.86	6.27	-	37.00	0.64	-	6.78
28-55	8.61	-	-	1.4	0.23	13.68	-	-	0.68	12.27	-	53.20	0.81	-	9.22

Contd....

Series Name: Dombarahalli (DRL)

Pedon: R-8

Location: 15°13'96.2"N, 75°57'48.6" E Rangunathanahalli village, Koppal Taluk and District

Analysis at: NBSS&LUP, Regional Centre, Bangalore.

Classification: Very-fine, smectitic (calc), isohyperthermic Typic Haplusterts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-15	Ap	28.25	19.48	52.27	4.76	4.44	4.87	8.23	5.95	-	c	39.86	27.20
15-27	BA1	21.55	20.00	58.45	3.76	2.76	3.43	6.30	5.30	-	c	46.35	34.84
27-45	Bss1	14.86	20.89	64.25	2.46	2.23	2.23	3.91	4.02	-	c	57.99	41.06
45-80	Bss2	10.42	19.04	70.54	1.74	1.97	1.27	2.78	2.66	-	c	66.36	36.24

Depth (cm)	pH (1:2.5)			E.C. (1:2.5) dS m ⁻¹	O.C. %	CaCO ₃ %	Exchangeable bases					CEC	CEC/Clay	Base saturation %	ESP %
	Water	CaCl ₂	M KCl				Ca	Mg	K	Na	Total				
0-15	8.78	-	-	0.42	0.32	12.35	-	-	0.59	4.25	-	49.70	0.95	100.00	5.62
15-27	9.03	-	-	0.61	0.30	12.48	-	-	0.30	8.96	-	57.23	0.98	100.00	10.07
27-45	9.10	-	-	0.67	0.34	11.70	-	-	0.25	11.85	-	60.71	0.95	100.00	14.05
45-80	9.18	-	-	0.86	0.32	13.39	-	-	0.27	15.40	-	63.33	0.90	100.00	18.45

Contd....

Series Name: Narsapura (NSP), **Pedon:** A2/RM-2

Location: 15°19'86.9"N, 75°57'86.1"E, Kavalura village, Koppal Taluk and District

Analysis at: NBSS&LUP, Regional Centre, Bangalore. **Classification:** Very-fine, smectitic (calc), isohyperthermic Vertic Haplustepts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-29	Ap	31.32	16.52	52.16	5.51	5.40	5.51	9.83	5.08	10	c	38.86	27.64
29-52	Bw1	13.30	22.08	64.62	2.52	2.41	2.41	3.67	2.29	05	c	49.88	40.05
52-77	BW2	13.22	17.39	69.40	3.56	2.41	1.95	2.76	2.53	05	c	51.33	41.55

Depth (cm)	pH (1:2.5)			E.C. (1:2.5) dS m ⁻¹	O.C. %	CaCO ₃ %	Exchangeable bases					CEC	CEC/Clay	Base saturation %	ESP %
	Water	CaCl ₂	M KCl				Ca	Mg	K	Na	Total				
0-29	9.16	-	-	0.615	0.23	9.36	-	-	0.72	10.98	-	51.09	0.98	-	8.60
29-52	8.69	-	-	2.01	0.5	8.64	-	-	0.55	24.42	-	60.63	0.94	-	16.11
52-77	8.52	-	-	2.68	0.46	7.68	-	-	0.50	25.65	-	60.74	0.88	-	16.90

Contd....

Series Name: Gatareddihal (GRH) Pedon: R-7

Location: 15°14'20.8"N, 76°04'28.4" E Gudlanur village, Koppal Taluk and District

Analysis at: NBSS&LUP, Regional Centre, Bangalore.

Classification: Very-fine, smectitic (calc), isohyperthermic Sodic Haplusterts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-18	Ap	20.07	19.71	60.23	1.76	3.75	3.64	3.42	7.50	-	c	41.70	29.56
18-51	Bss1	15.11	17.47	67.42	3.16	3.04	2.25	3.38	3.27	-	c	59.43	38.52
51-80	Bss2	13.19	18.74	68.07	1.80	2.93	2.37	3.04	3.04	-	c	60.69	40.91
80-107	Bss3	17.54	19.50	62.96	2.46	4.13	3.24	4.25	3.46	-	c	57.25	37.31
107-131	BC	9.42	17.48	73.10	1.48	1.82	1.36	1.93	2.84	-	c	64.62	43.98

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO ₃	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP			
	Water	CaCl ₂	M KCl				dS m ⁻¹	%	%	Ca	Mg					K	Na	Total
										cmol kg ⁻¹						%	%	
0-18	9.08	-	-	0.23	0.33	6.89	-	-	0.70	6.36	-	63.21	1.05	100.00	7.11			
18-51	9.19	-	-	0.61	0.49	9.10	-	-	0.54	14.20	-	66.05	0.98	100.00	15.98			
51-80	9.27	-	-	0.56	0.29	9.36	-	-	0.49	14.75	-	65.63	0.96	100.00	17.07			
80-107	9.28	-	-	0.57	0.39	9.62	-	-	0.44	14.64	-	63.95	1.02	100.00	17.49			
107-131	9.04	-	-	1.08	0.31	8.32	-	-	0.52	16.40	-	68.36	0.94	100.00	17.30			

Contd....

Series Name: Handrala (HDL), **Pedon:** A2/RM-1

Location: 15°19'69.8"N, 75°58'00"E, Kavalura village, Koppal Taluk and District

Analysis at: NBSS&LUP, Regional Centre, Bangalore.

Classification: Very-fine, smectitic (calc), isohyperthermic Typic Haplusterts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-25	Ap	21.68	16.62	61.70	4.42	3.98	3.43	5.64	4.20	10	c	41.36	31.27
25-50	Bss1	14.93	15.76	69.32	2.64	2.53	2.99	3.33	3.44	05	c	48.92	39.19
50-82	Bss2	23.11	16.60	60.29	4.51	3.61	6.31	4.74	3.95	05	c	42.46	33.85
82-117	Bss3	10.50	18.38	71.12	1.98	1.98	1.63	2.57	2.33	05	c	52.95	42.82

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO ₃	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP			
	Water	CaCl ₂	M KCl				dS m ⁻¹	%	%	Ca	Mg					K	Na	Total
										cmol kg ⁻¹						%	%	
0-25	9.06	-	-	0.371	0.16	4.80	-	-	0.80	7.93	-	62.33	1.01	-	5.09			
25-50	9.09	-	-	0.719	0.2	7.20	-	-	0.42	14.94	-	67.10	0.97	-	8.90			
50-82	9.28	-	-	0.47	0.19	9.36	-	-	0.47	11.59	-	60.21	1.00	-	7.70			
82-117	8.76	-	-	1.55	0.36	8.64	-	-	0.11	2.28	-	25.33	0.36	-	3.61			

Contd....

Series Name: Kavalura (KVR), **Pedon:** A2/RM-9

Location: 15°18'86.8"N, 75°56'56.3"E, Kavalura village, Koppal Taluk and District

Analysis at: NBSS&LUP, Regional Centre, Bangalore. **Classification:** Fine, smectitic (calc), isohyperthermic Typic Haplusterts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-24	Ap	36.18	17.80	46.02	7.04	7.47	6.62	9.28	5.76	10	c	28.20	18.75
24-50	Bss1	38.79	15.36	45.85	6.25	6.25	9.70	10.67	5.93	05	c	27.16	18.81
50-85	Bss2	36.80	14.66	48.54	9.63	8.23	7.03	7.58	4.33	<5	c	30.16	22.17
85-124	Bss3	22.66	17.24	60.09	4.18	3.85	5.28	5.06	4.29	<5	c	40.34	31.42

Depth (cm)	pH (1:2.5)			E.C. (1:2.5) dS m ⁻¹	O.C. %	CaCO ₃ %	Exchangeable bases					CEC	CEC/Clay	Base saturation %	ESP %
	Water	CaCl ₂	M KCl				Ca	Mg	K	Na	Total				
0-24	8.4	-	-	0.265	0.2	8.04	-	-	0.97	0.65		43.25	0.94		0.60
24-50	9.27	-	-	0.23	0.37	8.04	-	-	0.31	3.21		41.66	0.91		3.08
50-85	9.44	-	-	0.297	0.41	8.64	-	-	0.35	6.43		43.99	0.91		5.85
85-124	9.37	-	-	0.46	0.41	11.40	-	-	0.42	7.99		51.09	0.85		6.26

Contd....

Series Name: Budagumpa (BGP), **Pedon:** R-21

Location: 15°23'45"N, 76°08'52"E Neregalla village, Koppal Taluk and District

Analysis at: NBSS&LUP, Regional Centre, Bangalore. **Classification:** Fine, mixed (calc), isohyperthermic Typic Haplustepts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-16	Ap	58.30	18.10	23.60	6.34	11.75	11.66	17.44	11.10	-	scl	18.24	10.29
16-38	Bw1	44.26	18.39	37.36	4.71	9.79	9.32	12.24	8.19	-	cl	32.99	18.12
38-68	Bw2	37.84	24.91	37.25	3.66	7.51	8.45	10.89	7.32	-	cl	39.50	22.32
68-83	Bw3	19.17	19.89	60.93	0.87	3.47	3.85	6.07	4.91	-	c	47.27	28.52
83-107	Bw4	14.76	23.22	62.02	0.63	2.41	3.25	4.61	3.87	-	c	46.10	29.36
107-131	Bw5	11.86	17.75	70.39	0.85	2.73	2.45	3.20	2.64	-	c	50.52	28.09
131-160	Bw6	14.48	18.21	67.31	2.23	2.50	2.59	3.84	3.31	-	c	59.14	28.35

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO ₃	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP			
	Water	CaCl ₂	M KCl				dS m ⁻¹	%	%	Ca	Mg					K	Na	Total
0-16	9.20	-	-	0.27	0.51	6.24	-	-	0.42	3.11	-	19.60	0.83	100.00	3.84			
16-38	9.29	-	-	0.88	0.35	5.98	-	-	0.17	9.36	-	28.40	0.76	100.00	15.38			
38-68	8.95	-	-	2.37	0.31	4.81	-	-	0.31	24.10	-	34.90	0.94	100.00	42.65			
68-83	8.65	-	-	4.28	0.33	4.42	-	-	0.39	27.95	-	45.10	0.74	100.00	25.94			
83-107	8.10	-	-	9.50	0.30	3.38	-	-	0.44	31.29	-	44.10	0.71	100.00	12.82			
107-131	8.16	-	-	9.32	0.22	2.73	-	-	0.63	37.86	-	47.20	0.67	100.00	20.37			
131-160	8.49	-	-	5.29	0.19	3.51	-	-	0.60	34.82	-	43.70	0.65	100.00	48.66			

Contd....

Series Name: Bardur (BDR), **Pedon:** R-4

Location: 15°14'31.7"N, 76°01'19.1"E, Moranali village, Koppal Taluk and District

Analysis at: NBSS&LUP, Regional Centre, Bangalore. **Classification:** Very-fine, smectitic (calcareous), isohyperthermic Typic Haplusterts.

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-25	Ap	21.78	22.78	55.44	2.17	3.68	4.44	6.61	4.88	-	c	36.78	26.95
25-53	BA	18.62	18.56	62.82	2.23	4.24	3.46	5.24	3.46	-	c	41.25	29.87
53-90	Bss1	15.87	18.60	65.53	2.23	1.34	4.25	3.91	4.13	-	c	44.73	33.64
90-126	Bss2	13.66	20.02	66.32	1.68	2.80	2.35	3.70	3.14	-	c	49.24	38.37
126-152	Bss3	11.64	20.79	67.57	1.69	1.81	1.81	3.50	2.82	-	c	53.50	41.90
152-210	Bss4	11.38	22.78	65.42	2.16	2.16	1.93	3.07	2.05	-	c	51.53	39.64

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO ₃	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP						
	Water	CaCl ₂	M KCl				dS m ⁻¹	%	%	Ca	Mg					K	Na	Total	cmol kg ⁻¹	%	%
0-25	8.73	-	22.78	0.203	0.24	5.76	-	-	0.65	4.43	-	40.56	0.73	-	4.37						
25-53	9.17	-	18.56	0.295	0.45	4.92	-	-	0.32	10.47	-	74.70	1.19	-	5.61						
53-90	9.27	-	18.60	0.388	0.66	6.00	-	-	0.24	10.49	-	76.20	1.16	-	5.51						
90-126	9.22	-	20.02	0.608	0.57	5.88	-	-	0.21	15.93	-	77.20	1.16	-	8.25						
126-152	9.21	-	20.79	0.936	0.33	6.60	-	-	0.37	20.88	-	80.90	1.20	-	10.32						
152-210	9.03	-	23.21	1.47	0.33	8.16	-	-	0.24	15.34	-	73.10	1.12	-	8.39						

INTERPRETATION FOR LAND RESOURCE MANAGEMENT

The most important soil and site characteristics that affect the land use and conservation needs of an area are land capability, land irrigability, soil depth, soil texture, coarse fragments, available water capacity, soil slope, soil erosion, soil reaction etc. These are interpreted from the data base generated through land resource inventory and several thematic maps are generated. These would help in identifying the areas suitable for growing crops and, soil and water conservation measures and structures needed thus helping to maintain good soil health for sustained crop production. The various thematic maps generated are described below.

5.1 Land Capability Classification

Land capability classification is an interpretative grouping of soil map units (soil phases) mainly based on inherent soil characteristics, external land features and environmental factors that limit the use of land for agriculture, pasture, forestry or other uses on a sustained basis (IARI, 1971). The land and soil characteristics used to group the land resources in an area into various land capability classes, subclasses and units are

Soil characteristics: Soil depth, soil texture, coarse fragments, soil reaction, available water capacity, calcareousness, salinity/alkali *etc.*

Land characteristics: Slope, erosion, drainage and rock-outcrops.

Climate: Total rainfall and its distribution, and length of crop growing period.

The Land Capability Classification system is divided into land capability classes, subclasses and units based on the level of information available. Eight land capability classes are recognized. They are

Class I: They are very good lands that have no limitations or very few limitations that restrict their use.

Class II: They are good lands that have minor limitations and require moderate conservation practices.

Class III: They are moderately good lands that have severe limitations that reduce the choice of crops or that require special conservation practices.

Class IV: They are fairly good lands that have very severe limitations that reduce the choice of crops or that require very careful management.

Class V: Soils in these lands are not likely to erode, but have other limitations like wetness that are impractical to remove and as such not suitable for agriculture, but suitable for pasture or forestry with minor limitations.

Class VI: The lands have severe limitations that make them generally unsuitable for cultivation, but suitable for pasture or forestry with moderate limitations.

Class VII: The lands have very severe limitations that make them unsuitable for cultivation, but suitable for pasture or forestry with major limitations.

Class VIII: Soil and other miscellaneous areas (rock lands) that have very severe limitations that nearly preclude their use for any crop production, but suitable for wildlife, recreation and installation of wind mills.

The land capability subclasses are recognised based on the dominant limitations observed within a given land capability class. The subclasses are designated by adding a lower case letter like ‘e’, ‘w’, ‘s’, or ‘c’ to the class numeral. The subclass “e” indicates that the main hazard is risk of erosion, “w” indicates drainage or wetness as a limitation for plant growth, “s” indicates shallow soil depth, coarse or heavy textures, calcareousness, salinity/alkali or gravelliness and “c” indicates limitation due to climate.

The land capability subclasses have been further subdivided into land capability units based on the kinds of limitations present in each subclass. Ten land capability units are identified in grouping the soil map units. They are stony or rocky (0), erosion hazard (slope, erosion) (1), coarse texture (sand, loamy sand, sandy loam) (2), fine texture (cracking clay, silty clay) (3), slowly permeable subsoil (4), coarse underlying material (5), salinity/alkali (6), stagnation, overflow, high ground water table (7), soil depth (8) and fertility problems (9). The capability units thus identified have similar soil and land characteristics that respond similarly to a given level of management. The soils of the microwatershed have been classified upto land capability subclass level.

The 19 soil map units identified in the Naregallu-3 microwatershed are grouped under 2 Land capability classes and 5 land capability subclasses (Fig. 5.1). Entire cultivated area of about 544 ha (99%) is suitable for agriculture. An area of about 7 ha (1%) is under habitation and settlements.

Maximum area of about 490 ha (89%) is good lands (Class II) with minor problems of soil and erosion and distributed in the major part of the microwatershed. An area about 54 ha (10%) is moderately good lands (Class III) with moderate limitations of soil and erosion and distributed in the southeastern and northwestern part of the microwatershed.




Fig. 5.1 Land Capability map of Naregallu-3 Microwatershed

5.2 Soil Depth

Soil depth refers to the depth of the soil occurring above the parent material or hard rock. The depth of the soil determines the effective rooting depth for plants and in accordance with soil texture, mineralogy and gravel content, the capacity of the soil column to hold water and nutrient availability. Soil depth is one of the most important soil characteristic that is used in differentiating soils into different soil series. The soil depth classes used in identifying soils in the field are very shallow (<25 cm), shallow (25-50 cm), moderately shallow (50-75 cm), moderately deep (75-100 cm), deep (100-150 cm) and very deep (>150 cm). They were used to classify the soils into different depth classes and a soil depth map was generated (Fig. 5.2).

An area of about 53 ha (10%) is under shallow (25-50 cm) soils and distributed in the southeastern and northwestern part of the microwatershed. Moderately shallow (50-75 cm) soils cover an area of about 57 ha (10%) and occur in the northern, southwestern, western and southeastern part of the microwatershed. Moderately deep (75-100 cm) soils occupy an area of about 90 ha (16%) and occur in the central, northern, southern and northwestern part of the microwatershed. Deep (100-150 cm) soils cover a major area of about 229 ha (42%) and distributed in the central, eastern, southeastern, southern, southwestern, western and northwestern part of the microwatershed. Very deep (>150 cm) soils occupy an area of 116 ha (21%) and occur in the central, southern, eastern and western part of the microwatershed.

The most productive lands cover a major area about 345 ha (63%) where all climatically adapted long duration crops can be grown. The problem soils cover about 53 ha (10%) area where only short duration crops can be grown and the probability of crop failure is high.




Fig. 5.2 Soil Depth map of Naregallu-3 Microwatershed

5.3 Surface Soil Texture

Texture is an expression to indicate the coarseness or fineness of the soil as determined by the relative proportion of primary particles of sand, silt and clay. It has a direct bearing on the structure, porosity, adhesion and consistence. The surface layer of a soil to a depth of about 25 cm is the layer that is most used by crops and plants. The surface soil textural class provides a guide to understanding soil-water retention and availability, nutrient holding capacity, infiltration, workability, drainage, physical and chemical behaviour, microbial activity and crop suitability. The textural classes used for LRI were used to classify and a surface soil texture map showing sandy, loamy and clayey at the surface was generated. The area extent and their geographical distribution in the microwatershed is shown in Fig.5.3.

Entire cultivated area in the microwatershed has soils that are clayey at the surface.

Entire cultivated area has most productive lands with respect to surface soil texture where they are clayey soils. They have high potential for soil-water retention and

availability, and nutrient retention and availability, but have more problems of drainage, infiltration, workability and other physical problems.




Fig. 5.3 Surface Soil Texture map of Naregallu-3 Microwatershed

5.4 Soil Gravelliness

Gravel is the term used for describing coarse fragments between 2 mm and 7.5 cm diameter and stones for those between 7.5 cm and 25 cm. The presence of gravel and stones in soil reduces the volume of soil responsible for moisture and nutrient storage, drainage, infiltration and runoff, and hinders plant growth by impeding root growth and seedling emergence, intercultural operations and farm mechanization. The gravelliness classes used in LRI were used to classify the soils and using these classes, a gravelliness map was generated. The area extent and their spatial distribution in the microwatershed is shown in Fig.5.4.

Maximum area of about 488 ha (89%) has non gravelly (<15%) soils and occur in the major part of the microwatershed. An area of about 56 ha (10%) has gravelly (15-35%) soils and distributed in the eastern part of the microwatershed.

An area of about 488 ha (89%) are most productive lands with respect to gravelliness. They are non-gravelly with less than 15 per cent gravel and have potential for growing both annual and perennial crops. The problem lands cover about 56 ha (10%) that are gravelly where only medium or short duration crops can be grown.




Fig. 5.4 Soil Gravelliness map of Naregallu-3 Microwatershed

5.5 Available Water Capacity

The soil available water capacity (AWC) is estimated based on the ability of the soil column to retain water between the tensions of 0.33 and 15 bar in a depth of 100 cm or the entire solum if the soil is shallower. The AWC of the soils (soil series) as estimated by considering the soil texture, mineralogy, soil depth and gravel content (Sehgal *et al.*, 1990) and accordingly the soil map units were grouped into five AWC classes *viz.*, very low (<50 mm/m), low (50-100 mm/m), medium (100-150 mm/m), high (150-200 mm/m) and very high (>200 mm/m) and using these values, an AWC map was generated (Fig. 5.5).

An area of about 109 ha (20%) has soils that are low (51-100 mm/m) in available water capacity and distributed in the northern, southeastern, southwestern, western and northwestern part of the microwatershed. An area of about 149 ha (27%) is medium to high (101-200 mm/m) in available water capacity and occur in the central, southern, northern, northwestern and eastern part of the microwatershed. Maximum area of about 285 ha (52%) is very high (>200 mm/m) in available water capacity and occur in the major part of the microwatershed.

An area of about 109 ha (20%) in the microwatershed has soils that are problematic with regard to available water capacity. Here, only short duration crops can be grown and the probability of crop failure is very high. These areas are best put to other alternative uses. Maximum area of about 344 ha (63%) has soils that have very high

potential (151->200 mm/m) with regard to available water capacity where all climatically adapted long duration crops can be grown successfully.




Fig. 5.5 Soil Available Water Capacity map of Naregallu-3 Microwatershed

5.6 Soil Slope

Soil slope refers to the inclination of the surface of the land. It is defined by gradient, shape and length, and is an integral feature of any soil as a natural body. Slope is considered important in soil genesis, land use and land development. The length and gradient of slope influences the rate of runoff, infiltration, erosion and deposition. The soil map units were grouped into different slope classes and a slope map was generated showing the area extent and their geographic distribution of different slope classes in the microwatershed (Fig. 5.6).

An area of about 47 ha (8%) falls under nearly level (0-1% slope) lands and distributed in the northern and western part of the microwatershed. Maximum area of about 497 ha (90%) falls under very gently sloping (1-3% slope) lands and distributed in the major part of the microwatershed.

Entire cultivated area in the microwatershed has soils that have high potential in respect of soil slopes. In these areas, all climatically adapted annual and perennial crops can be grown without much soil and water conservation and other land development measures.




Fig. 5.6 Soil Slope map of Naregallu-3 Microwatershed

5.7 Soil Erosion

Soil erosion refers to the wearing away of the earth's surface by the forces of water, wind and ice involving detachment and transport of soil by raindrop impact. It is used for accelerated soil erosion resulting from disturbance of the natural landscape by burning, excessive grazing and indiscriminate felling of forest trees and tillage, all usually by man. The erosion classes showing an estimate of the current erosion status as judged from field observations in the form of rills, gullies or a carpet of gravel on the surface are recorded. Four erosion classes, viz, slight erosion (e1), moderate erosion (e2), severe erosion (e3) and very severe erosion (e4) are recognized. The soil map units were grouped into different erosion classes and a soil erosion map generated. The area extent and their spatial distribution in the microwatershed is given in Figure 5.7.

Soils that are slightly eroded (e1 class) cover an area of 262 ha (48%) and distributed in the central, northern, northeastern, eastern, southeastern, southern, southwestern and western part of the microwatershed. Soils that are moderately eroded (e2 class) cover a major area of 282 ha (51%) and distributed in the central, northern, northeastern, eastern, southeastern, southern, northwestern and western part of the microwatershed.

Maximum area of about 282 ha (51%) in the microwatershed is problematic because of moderate erosion. For these areas, taking up soil and water conservation and other land development measures are needed.




Fig. 5.7 Soil Erosion map of Naregallu-3 Microwatershed

FERTILITY STATUS

Soil fertility plays an important role in increasing crop yield. The adoption of high yielding varieties that require high amounts of nutrients has resulted in deficiency symptoms in crops and plants due to imbalanced fertilization and poor inherent fertility status, as these areas are characterised by low rainfall and high temperatures. Hence, it is necessary to know the fertility (macro and micro nutrients) status of the soils of the watersheds for assessing the kind and amount of fertilizers required for each of the crop intended to be grown. For this purpose, the surface soil samples collected from the grid points (one soil sample at every 320 m grid interval) all over the microwatershed through land resource inventory in the year 2018 were analysed for pH, EC, organic carbon, available phosphorus and potassium, and for micronutrients like zinc, boron, copper, iron and manganese, and secondary nutrient sulphur.

Soil fertility data generated has been assessed and individual maps for all the nutrients for the microwatershed have been generated using the Kriging method under GIS. The village/survey number wise fertility data for the microwatershed is given in Appendix-II.

6.1 Soil Reaction (pH)

The soil analysis of the Naregallu-3 microwatershed for soil reaction (pH) showed that an area of about 206 ha (37%) is under strongly alkaline (pH 8.4-9.0) in soil reaction and distributed in the southern, southeastern, eastern, northeastern and northwestern part of the microwatershed. Maximum area of about 337 ha (61%) is under very strongly alkaline (pH >9.0) in soil reaction and occur in the major part of the microwatershed (Fig.6.1). Thus, entire cultivated area of the microwatershed is under alkaline condition.

6.2 Electrical Conductivity (EC)

The Electrical Conductivity of the soils of the entire microwatershed area is $<2 \text{ dS m}^{-1}$ (Fig 6.2) and as such the soils are non-saline.

6.3 Organic Carbon (OC)

The soil organic carbon content (an index of available Nitrogen) of the microwatershed is medium (0.5-0.75%) in an area of about 124 ha (23%) and occur in the southern, southeastern and eastern part of the microwatershed. Maximum area of about 420 ha (76%) is low ($<0.5\%$) in organic carbon and distributed in the major part of the microwatershed (Fig.6.3).

6.4 Available Phosphorus

An area of about 71 ha (13%) is medium (23-57 kg/ha) in available phosphorus and distributed in the eastern part of the microwatershed. Low ($<23 \text{ kg/ha}$) in available

phosphorus cover a major area of about 472 ha (86%) and distributed in the major part of the microwatershed (Fig 6.4).

6.5 Available Potassium

An area of about 237 ha (43%) is medium (145-337 kg/ha) and distributed in the central, northern, eastern, western and northwestern part of the microwatershed. Maximum area of about 307 ha (56%) is high (>337 kg/ha) and distributed in the major part of the microwatershed (Fig.6.5).

6.6 Available Sulphur

Available sulphur content is low (<10 ppm) in the entire cultivated area of the microwatershed (Fig.6.6).




Fig.6.1 Soil Reaction (pH) map of Naregallu-3 Microwatershed




Fig.6.2 Electrical Conductivity (EC) map of Naregallu-3 Microwatershed




Fig.6.3 Soil Organic Carbon (OC) map of Naregallu-3 Microwatershed




Fig.6.4 Soil Available Phosphorus map of Naregallu-3 Microwatershed




Fig.6.5 Soil Available Potassium map of Naregallu-3 Microwatershed




Fig.6.6 Soil Available Sulphur map of Naregallu-3 Microwatershed

6.7 Available Boron

Available boron is low (<0.5 ppm) in a major area of about 268 ha (49%) and distributed in the northeastern, northern, northwestern and western part of the microwatershed. An area of about 245 ha (44%) is medium (0.5-1.0 ppm) and occur in the central, southwestern, southern, southeastern and eastern part of the microwatershed. An area of about 31 ha (6%) is high (>1.0 ppm) and distributed in the southeastern part of the microwatershed (Fig.6.7).

6.8 Available Iron

Available iron content is sufficient (>4.5 ppm) in an area of about 124 ha (22%) and distributed in the western, southeastern and eastern part of the microwatershed. Maximum area of about 420 ha (76%) is deficient (<4.5 ppm) and distributed in the major part of the microwatershed (Fig 6.8).

6.9 Available Manganese

Available manganese content is sufficient (>1.0 ppm) in the entire cultivated area of the microwatershed (Fig 6.9).

6.10 Available Copper

Available copper content is sufficient (>0.2 ppm) in the entire cultivated microwatershed area (Fig 6.10).

6.11 Available Zinc

Available zinc content is deficient (<0.6 ppm) in the entire cultivated area of the microwatershed (Fig 6.11).




Fig.6.7 Soil Available Boron map of Naregallu-3 Microwatershed




Fig.6.8 Soil Available Iron map of Naregallu-3 Microwatershed




Fig.6.9 Soil Available Manganese map of Naregallu-3 Microwatershed




Fig.6.10 Soil Available Copper map of Naregallu-3 Microwatershed




Fig.6.11 Soil Available Zinc map of Naregallu-3 Microwatershed

LAND SUITABILITY FOR MAJOR CROPS

The soil and land resource units (soil phases) of Naregallu-3 Microwatershed were assessed for their suitability for growing food, fodder, fibre and other horticulture crops by following the procedure as outlined in FAO, 1976 and 1983. Crop requirements were developed for each of the crop from the available research data and also by referring to Naidu *et. al.* (2006) and Natarajan *et. al* (2015). The soil and land characteristics (Table 7.1) were matched with the crop requirements (Tables 7.2 to 7.32) to arrive at the crop suitability. The soil and land characteristics table and crop requirements tables are given at the end of the chapter. In FAO land suitability classification, two orders are recognized. Order S-Suitable and Order N- Not suitable. The orders have Classes, subclasses and units. Order-S has three classes, Class S1-Highly Suitable, Class S2-Moderately Suitable and Class S3-Marginally Suitable. Order N has two classes, N1-Currently not Suitable and N2-Permanently not Suitable. There are no subclasses within the Class S1 as they will have very minor or no limitations for crop growth. Classes S2, S3, N1 and N2 are divided into subclasses based on the kinds of limitations encountered. The limitations that affect crop production are ‘c’ for erratic rainfall and its distribution and length of growing period (LGP), ‘e’ for erosion hazard, ‘r’ for rooting condition, ‘t’ for lighter or heavy texture, ‘g’ for gravelliness or stoniness, ‘n’ for nutrient availability, ‘l’ for topography, ‘m’ for moisture availability, ‘z’ for calcareousness and ‘w’ for drainage. These limitations are indicated as lower case letters to the class symbol. For example, moderately suitable lands with the limitations of soil depth and erosion are designated as S2re. For the microwatershed, the soil mapping units were evaluated and classified up to subclass level.

Using the above criteria, the soil map units of the microwatershed were evaluated and land suitability maps for 31 major agricultural and horticultural crops were generated. The detailed information on the kind of suitability of each of the soil phase for the crops assessed are given village/ survey number wise for the microwatershed in Appendix-III

7.1 Land Suitability for Sorghum (*Sorghum bicolor*)

Sorghum is one of the major food crop grown in Karnataka in an area of 10.47 lakh ha in Bijapur, Gulbarga, Raichur, Bidar, Belgaum, Dharwad, Bellary, Chitradurga, Mysore and Chamarajnagar districts. The crop requirements for growing sorghum (Table 7.2) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and land a suitability map for growing sorghum was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure. 7.1.

An area of about 207 ha (38%) area is highly suitable (Class S1) for growing sorghum and occur in the western, northwestern, northern, northeastern and eastern part of the microwatershed. Maximum area of about 284 ha (52%) is moderately suitable

(Class S2) for growing sorghum and distributed in the major part of the microwatershed. They have minor limitations of rooting depth, gravelliness, calcareousness and nutrient availability. An area of about 53 ha (10%) is marginally suitable (Class S3) for growing sorghum and occur in the northwestern and southeastern part of the microwatershed with moderate limitations of rooting depth and calcareousness.




Fig. 7.1 Land Suitability map of Sorghum

7.2 Land Suitability for Maize (*Zea mays*)

Maize is one of the most important food crop grown in an area of 13.37 lakh ha in almost all the districts of the State. The crop requirements for growing maize (Table 7.3) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing maize was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.2.

An area of about 33 ha (6%) is highly suitable (Class S1) for growing maize and occur in the eastern part of the microwatershed. Maximum area of about 459 ha (83%) is moderately suitable (Class S2) for growing maize and distributed in the major part of the microwatershed with minor limitations of texture and calcareousness. An area of about 53 ha (10%) is marginally suitable (Class S3) for growing maize and occur in the northwestern and southeastern part of the microwatershed with moderate limitations of texture and calcareousness.




Fig. 7.2 Land Suitability map of Maize

7.3 Land Suitability for Bajra (*Pennisetum glaucum*)

Bajra is one of the major food crop grown in an area of 2.34 lakh ha in Karnataka in the northern districts. The crop requirements (Table 7.4) for growing bajra were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and land suitability map for growing bajra was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.3.

An area of about 34 ha (6%) is highly suitable (Class S1) for growing bajra and distributed in the eastern part of the microwatershed. Maximum area of about 457 ha (83%) is moderately suitable (Class S2) for growing bajra and distributed in the major part of the microwatershed with minor limitations of texture and calcareousness. An area of about 53 ha (10%) is marginally suitable (Class S3) for growing bajra and distributed in the northwestern and southeastern part of the microwatershed with moderate limitations of rooting depth and calcareousness.




Fig. 7.3 Land Suitability map of Bajra

7.4 Land Suitability for Groundnut (*Arachis hypogaea*)

Groundnut is one of the major oilseed crop grown in an area of 6.54 lakh ha in Karnataka in most of the districts either as rainfed or irrigated crop. The crop requirements for growing groundnut (Table 7.5) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing groundnut was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.4.

An area of about 34 ha (6%) is highly suitable (Class S1) for growing groundnut and distributed in the eastern part of the microwatershed. Maximum area of about 510 ha (93%) is marginally suitable (Class S3) for growing groundnut and distributed in the major part of the microwatershed with moderate limitations of calcareousness and texture.




Fig. 7.4 Land Suitability map of Groundnut

7.5 Land Suitability for Sunflower (*Helianthus annuus*)

Sunflower is one of the most important oilseed crop grown in an area of 3.56 lakh ha in the State in all the districts. The crop requirements for growing sunflower (Table 7.6) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing sunflower was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.5.

An area of about 172 ha (31%) is highly suitable (Class S1) for growing sunflower and distributed in the northeastern, eastern, western and northwestern part of the microwatershed. Maximum area of about 262 ha (48%) is moderately suitable (Class S2) for growing sunflower and distributed in the central, southern, western, northwestern, northern, eastern and southeastern part of the microwatershed with minor limitations of gravelliness, rooting depth and calcareousness. An area of about 57 ha (10%) is marginally suitable (Class S3) for growing sunflower and occur in the northern, southeastern, southwestern and western part of the microwatershed with moderate limitations of calcareousness and rooting depth. An area of about 53 ha (10%) is currently not suitable (Class N1) for growing sunflower and occur in the northwestern and southeastern part of the microwatershed with severe limitations of rooting depth and calcareousness.




Fig. 7.5 Land Suitability map of Sunflower

7.6 Land Suitability for Cotton (*Gossypium hirsutum*)

Cotton is one of the most important fibre crop grown in the state in about 8.75 lakh ha area in Raichur, Dharwad, Belgaum, Kalaburagi, Bijapur, Bidar, Bellary, Chitradurga and Chamarajnar districts. The crop requirements for growing cotton (Table 7.7) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing cotton was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.6.

An area of about 175 ha (32%) is highly (Class S1) suitable for growing cotton and occur in the western, northwestern, northern and northeastern part of the microwatershed. Maximum area of about 318 ha (57%) is moderately suitable (Class S2) for growing cotton and distributed in the major part of the microwatershed with minor limitations of gravelliness, texture, calcareousness and rooting depth. An area of about 53 ha (10%) is marginally suitable (Class S3) for growing cotton and occur in the northwestern and southeastern part of the microwatershed with moderate limitations of rooting depth and calcareousness.




Fig. 7.6 Land Suitability map of Cotton

7.7 Land Suitability for Red gram (*Cajanus cajana*)

Red gram is one of the major pulse crop grown in an area of 7.28 lakh ha mainly in northern Karnataka in Bijapur, Kalaburagi, Raichur, Bidar, Belgaum, Dharwad and Bellary districts. The crop requirements for growing red gram (Table 7.8) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing red gram was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed is given in Figure 7.7.

An area of about 34 ha (6%) is highly suitable (Class S1) for growing red gram and occur in the eastern part of the microwatershed. Maximum area of about 345 ha (63%) is moderately suitable (Class S2) for growing red gram and occur in the major part of the microwatershed. They have minor limitations of texture, calcareousness and gravelliness. An area of about 112 ha (20%) is marginally suitable (Class S3) for growing red gram and distributed in the central, northern, southeastern, southern, southwestern, western and northwestern part of the microwatershed with moderate limitations of rooting depth and calcareousness. An area of about 53 ha (10%) is currently not suitable (Class N1) for growing red gram and occur in the northwestern and southeastern part of the microwatershed with severe limitations of rooting depth and calcareousness.




Fig. 7.7 Land Suitability map of Red gram

7.8 Land Suitability for Bengal gram (*Cicer aerativum*)

Bengal gram is one of the most important pulse crop grown in about 9.39 lakh ha area in Bijapur, Raichur, Kalaburagi, Dharwad, Belgaum and Bellary districts. The crop requirements for growing Bengal gram (Table 7.9) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing Bengal gram was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.8.

Highly suitable (Class S1) lands for growing Bengal gram occur in an area of 206 ha (37%) and distributed in the northern, northeastern, eastern, western and northwestern part of the microwatershed. Maximum area of about 286 ha (52%) is moderately suitable (Class S2) for growing Bengal gram and distributed in the major part of the microwatershed. They have minor limitations of rooting depth, gravelliness, calcareousness and texture. An area of about 53 ha (10%) is marginally suitable (Class S3) for growing Bengal gram and occur in the northwestern and southeastern part of the microwatershed with moderate limitations of rooting depth and calcareousness.




Fig. 7.8 Land Suitability map of Bengal gram

7.9 Land Suitability for Chilli (*Capsicum annum L*)

Chilli is one of the major spice crop grown in an area of 0.42 lakh ha in Karnataka State. The crop requirements for growing chilli (Table 7.10) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing chilli was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.9.

An area of about 34 ha (6%) is highly suitable (Class S1) for growing chilli and distributed in the eastern part of the microwatershed. Maximum area of about 510 ha (93%) is marginally suitable (Class S3) for growing chilli and occur in the major part of the microwatershed with moderate limitations of calcareousness, rooting depth and texture.




Fig. 7.9 Land Suitability map of Chilli

7.10 Land Suitability for Tomato (*Solanum lycopersicum*)

Tomato is one of the most important vegetable crop grown in an area of 0.65 lakh ha in almost all the districts of the State. The crop requirements (Table 7.11) for growing tomato were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing tomato was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.10.

An area of about 34 ha (6%) is highly suitable (Class S1) for growing tomato and distributed in the eastern part of the microwatershed. Major area of about 509 ha (93%) is marginally suitable (Class S3) for growing tomato and occur in the major part of the microwatershed with moderate limitations of calcareousness, rooting depth and texture.




Fig. 7.10 Land Suitability map of Tomato

7.11 Land Suitability for Brinjal (*Solanum melongena*)

Brinjal is one of the most important vegetable crop grown in the state. The crop requirements for growing brinjal (Table 7.12) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing Brinjal was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.11.

Highly suitable (Class S1) lands for growing brinjal occur in an area of 33 ha (6%) and distributed in the eastern part of the microwatershed. Maximum area of about 459 ha (83%) is moderately suitable (Class S2) for brinjal and distributed in the major part of the microwatershed. They have minor limitations of rooting depth, calcareousness and texture. An area about of 53 ha (10%) is marginally suitable (Class S3) and distributed in the southeastern and northwestern part of the microwatershed with moderate limitation of rooting depth.




Fig 7.11 Land Suitability map of Brinjal

7.12 Land Suitability for Onion (*Allium cepa L.*)

Onion is one of the most important vegetable crop grown in the state. The crop requirements for growing onion (Table 7.13) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing onion was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.12.

Highly suitable (Class S1) lands for growing onion occur in an area of 33 ha (6%) and distributed in the eastern part of the microwatershed. An area of about 2 ha (<1%) is moderately suitable (Class S2) for onion and distributed in the eastern part of the microwatershed. They have minor limitation of texture. Maximum area of about 510 ha (93%) is marginally suitable (Class S3) and distributed in the major part of the microwatershed with moderate limitations of rooting depth, texture and calcareousness.




Fig 7.12 Land Suitability map of Onion

7.13 Land Suitability for Bhendi (*Abelmoschus esculentus*)

Bhendi is one of the most important vegetable crop grown in the state. The crop requirements for growing bhendi (Table 7.14) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing bhendi was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.13.

Highly suitable (Class S1) lands for growing bhendi occur in area of 33 ha (6%) and distributed in the eastern part of the microwatershed. Maximum area of about 459 ha (83%) is moderately suitable (Class S2) for bhendi and distributed in the major part of the microwatershed. They have minor limitations of rooting depth, calcareousness and texture. An area about of 53 ha (10%) is marginally suitable (Class S3) and distributed in the northwestern and southeastern part of the microwatershed with moderate limitation of rooting depth.




Fig 7.13 Land Suitability map of Bhendi

7.14 Land Suitability for Drumstick (*Moringa oleifera*)

Drumstick is one of the most important vegetable crop grown in 2403 ha area in the state. The crop requirements for growing drumstick (Table 7.15) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing drumstick was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.14.

An area of about 34 ha (6%) is highly suitable (Class S1) for growing drumstick and distributed in the eastern part of the microwatershed. Maximum area of 400 ha (73%) is moderately suitable (Class S2) for growing drumstick and distributed in the major part of the microwatershed with minor limitations of texture, rooting depth, calcareousness and gravelliness. An area of about 57 ha (10%) is marginally suitable (Class S3) for growing drumstick and occur in the northern, southeastern, southwestern and western part of the microwatershed with moderate limitations of rooting depth and calcareousness. An area of about 53 ha (10%) is currently not suitable (Class N1) for growing drumstick and occur in the northwestern and southeastern part of the microwatershed with severe limitations of rooting depth and calcareousness.




Fig. 7.14 Land Suitability map of Drumstick

7.15 Land Suitability for Mulberry (*Morus nigra*)

Mulberry is the most important leaf crop grown for rearing silkworms in about 1.66 lakh ha in all the districts of the state. The crop requirements for growing mulberry (Table 7.16) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing mulberry was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.15.

Highly suitable (Class S1) lands for growing mulberry occur in an area of 34 ha (6%) and distributed in the eastern part of the microwatershed. Moderately suitable (Class S2) lands occupy a major area of about 345 ha (62%) and occur in the major part of the microwatershed. They have minor limitations of texture, calcareousness, drainage and gravelliness. Marginally suitable (Class S3) lands cover an area of about 113 ha (20%) and occur in the northern, southeastern, southwestern and western part of the microwatershed. They have moderate limitations of rooting depth, texture and calcareousness. An area of about 53 ha (10%) is currently not suitable (Class N1) for growing mulberry and occur in the northwestern and southeastern part of the microwatershed with severe limitations of rooting depth and calcareousness.




Fig. 7.15 Land Suitability map of Mulberry

7.16 Land Suitability for Mango (*Mangifera indica*)

Mango is one of the most important fruit crop grown in about 1.73 lakh ha in almost all the districts of the State. The crop requirements (Table 7.17) for growing mango were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing mango was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.16.

An area of about 34 ha (6%) is highly suitable (Class S1) for growing mango and distributed in the eastern part of the microwatershed. Moderately suitable (Class S2) lands occupy an area of about 116 ha (21%) and occur in the western, southern, southeastern and eastern part of the microwatershed. They have minor limitations of rooting depth and calcareousness. Marginally suitable (Class S3) lands cover a major area of about 284 ha (52%) and occur in the major part of the microwatershed. They have moderate limitations of texture, rooting depth and calcareousness. An area of about 110 ha (20%) is currently not suitable (Class N1) for growing mango and occur in the southwestern, northwestern, northern, western and southeastern part of the microwatershed with severe limitations of calcareousness, texture and rooting depth.




Fig. 7.16 Land Suitability map of Mango

7.17 Land Suitability for Sapota (*Manilkara zapota*)

Sapota is one of the most important fruit crop grown in an area of about 29373 ha in almost all the districts of the state. The crop requirements (Table 7.18) for growing sapota were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing sapota was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.17.

An area of about 34 ha (6%) is highly suitable (Class S1) for growing sapota and distributed in the eastern part of the microwatershed. Major area of about 458 ha (83%) is marginally (Class S3) suitable for growing sapota and occur in the major part of the microwatershed with moderate limitations of texture, rooting depth and calcareousness. An area of about 53 ha (10%) is currently not suitable (Class N1) for growing sapota and occur in the northwestern and southeastern part of the microwatershed with severe limitations of rooting depth and calcareousness.




Fig. 7.17 Land Suitability map of Sapota

7.18 Land Suitability for Pomegranate (*Punica granatum*)

Pomegranate is one of the commercially grown fruit crop in about 18488 ha in Karnataka mainly in Bijapur, Bagalkot, Koppal, Gadag and Chitradurga districts. The crop requirements for growing pomegranate (Table 7.19) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing pomegranate was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.18.

Highly suitable (Class S1) lands for growing pomegranate occur in an area of about 34 ha (6%) and distributed in the eastern part of the microwatershed. Maximum area of about 400 ha (73%) is moderately suitable (Class S2) for growing pomegranate and occur in the major part of the microwatershed with minor limitations of texture, rooting depth, calcareousness and gravelliness. Marginally suitable (Class S3) lands cover a major area of about 57 ha (10%) and occur in the northern, southeastern, southwestern and western part of the microwatershed. They have moderate limitations of rooting depth and calcareousness. An area of about 53 ha (10%) is currently not suitable (Class N1) for growing pomegranate and occur in the northwestern and southeastern part of the microwatershed with severe limitations of rooting depth and calcareousness.




Fig. 7.18 Land Suitability map of Pomegranate

7.19 Land Suitability for Guava (*Psidium guajava*)

Guava is one of the most important fruit crop grown in an area of about 6558 ha in almost all the districts of the state. The crop requirements (Table 7.20) for growing guava were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing guava was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.19.

An area of about 34 ha (6%) is highly suitable (Class S1) for growing guava and distributed in the eastern part of the microwatershed. Maximum area of 457 ha (83%) is marginally (Class S3) suitable for growing guava and occur in the major part of the microwatershed with moderate limitations of texture and calcareousness. An area of about 53 ha (10%) is currently not suitable (Class N1) for growing guava and occur in the northwestern and southeastern part of the microwatershed with severe limitations of rooting depth and texture.




Fig. 7.19 Land Suitability map of Guava

7.20 Land Suitability for Jackfruit (*Artocarpus heterophyllus*)

Jackfruit is one of the most important fruit crop grown in 5368 ha in all the districts of the state. The crop requirements (Table 7.21) for growing jackfruit were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing jackfruit was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in figure 7.20.

Highly suitable (Class S1) lands for growing jackfruit occur in an area of about 34 ha (6%) and distributed in the eastern part of the microwatershed. Major area of about 457 ha (83%) is marginally (Class S3) suitable for growing jackfruit and occur in the major part of the microwatershed with moderate limitations of texture and calcareousness. An area of about 53 ha (10%) is currently not suitable (Class N1) for growing jackfruit and occur in the northwestern and southeastern part of the microwatershed with severe limitations of rooting depth and texture.




Fig. 7.20 Land Suitability map of Jackfruit

7.21 Land Suitability for Jamun (*Syzygium cumini*)

Jamun is one of the important fruit crop grown in almost all the districts of the state. The crop requirements (Table 7.22) for growing jamun were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing jamun was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.21.

An area of about 34 ha (6%) is highly suitable (Class S1) for growing jamun and distributed in the eastern part of the microwatershed. Maximum area of 310 ha (56%) is moderately suitable (Class S2) for growing jamun and occur in the major part of the microwatershed with minor limitations of texture, rooting depth and calcareousness. Marginally suitable (Class S3) lands cover an area of about 147 ha (27%) and occur in the central, northwestern, northern, southeastern, southern, southwestern and western part of the microwatershed. They have moderate limitations of rooting depth, texture and calcareousness. An area of about 53 ha (10%) is currently not suitable (Class N1) for growing jamun and occur in the northwestern and southeastern part of the microwatershed with severe limitations of rooting depth and texture.




Fig. 7.21 Land Suitability map of Jamun

7.22 Land Suitability for Musambi (*Citrus limetta*)

Musambi is one of the most important fruit crop grown in an area of 5446 ha in almost all the districts of the state. The crop requirements (Table 7.23) for growing musambi were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing musambi was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.22.

Highly suitable (Class S1) lands for growing musambi cover an area of about 172 ha (31%) and occur in the eastern, northeastern, western and northwestern part of the microwatershed. Maximum area of about 262 ha (48%) is moderately suitable (Class S2) for growing musambi and occur in the central, northern, eastern, southeastern, southern, western and northwestern part of the microwatershed with minor limitations of rooting depth, calcareousness and gravelliness. Marginally suitable (Class S3) lands cover an area of about 57 ha (10%) and occur in the northern, southeastern, southwestern and western part of the microwatershed. They have moderate limitations of rooting depth and calcareousness. An area of about 53 ha (10%) is currently not suitable (Class N1) for growing musambi and occur in the northwestern and southeastern part of the microwatershed with severe limitations of rooting depth and calcareousness.




Fig. 7.22 Land Suitability map of Musambi

7.23 Land Suitability for Lime (*Citrus sp*)

Lime is one of the most important fruit crop grown in an area of 11752 ha in almost all the districts of the State. The crop requirements (Table 7.24) for growing lime (Table 7.15) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing lime was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.23.

An area of about 172 ha (31%) is highly suitable (Class S1) for growing lime and occurs in the eastern, northeastern, western and northwestern part of the microwatershed. Maximum area of about 262 ha (48%) is moderately suitable (Class S2) for growing lime and occur in the central, northern, eastern, southeastern, southern, western and northwestern part of the microwatershed with minor limitations of rooting depth, calcareousness and gravelliness. Marginally suitable (Class S3) lands cover an area of about 57 ha (10%) and occur in the northern, southeastern, southwestern and western part of the microwatershed. They have moderate limitations of rooting depth and calcareousness. An area of about 53 ha (10%) is currently not suitable (Class N1) for growing lime and occur in the northwestern and southeastern part of the microwatershed with severe limitations of rooting depth and calcareousness.




Fig. 7.23 Land Suitability map of Lime

7.24 Land Suitability for Cashew (*Anacardium occidentale*)

Cashew is one of the most important nut crop grown in an area of 7052 ha in almost all the districts of the State. The crop requirements (Table 7.25) for growing cashew were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing cashew was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.24.

Highly suitable (Class S1) lands for growing cashew cover an area of about 34 ha (6%) and occur in the eastern part of the microwatershed. Currently not suitable (Class N1) lands cover a major area of about 510 ha (93%) and distributed in the major part of the microwatershed with severe limitations of texture, rooting depth and calcareousness.




Fig. 7.24 Land Suitability map of Cashew

7.25 Land Suitability for Custard Apple (*Annona reticulata*)

Custard apple is one of the most important fruit crop grown in 1426 ha in almost all the districts of the State. The crop requirements (Table 7.26) for growing custard apple were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing custard apple was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.25.

An area of about 207 ha (38%) is highly suitable (Class S1) for growing custard apple and occur in the northern, northeastern, eastern, western and northwestern part of the microwatershed. Major area of about 285 ha (52%) is moderately suitable (Class S2) for growing custard apple and occur in the major part of the microwatershed with minor limitations of rooting depth, calcareousness and gravelliness. Marginally suitable (Class S3) lands cover an area of about 53 ha (10%) for growing custard apple and occur in the northwestern and southeastern part of the microwatershed. They have moderate limitations of calcareousness and gravelliness.




Fig. 7.25 Land Suitability map of Custard Apple

7.26 Land Suitability for Amla (*Phyllanthus emblica*)

Amla is one of the most important fruit and medicinal crop grown in an area of 151 ha and distributed in almost all the districts of the state. The crop requirements for (Table 7.27) growing amla were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing amla was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.26.

Highly suitable (Class S1) lands for growing amla cover an area of about 34 ha (6%) and occur in the eastern part of the microwatershed. Major area of about 458 ha (83%) is moderately suitable (Class S2) for growing amla and occur in the major part of the microwatershed with minor limitations of rooting depth, calcareousness, texture and gravelliness. An area of about 53 ha (10%) is marginally suitable (Class S3) for growing amla and occur in the northwestern and southeastern part of the microwatershed with moderate limitations of calcareousness and texture.




Fig. 7.26 Land Suitability map of Amla

7.27 Land Suitability for Tamarind (*Tamarindus indica*)

Tamarind is one of the most important spice crop grown in 14897 ha in all the districts of the state. The crop requirements (Table 7.28) for growing tamarind were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing tamarind was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.27.

An area of about 34 ha (6%) is highly suitable (Class S1) for growing tamarind and occur in the eastern part of the microwatershed. Maximum area of about 310 ha (56%) is moderately suitable (Class S2) for growing tamarind and occur in the major part of the microwatershed with minor limitations of rooting depth, gravelliness, calcareousness and texture. Marginally suitable (Class S3) lands cover an area of 90 ha (16%) for growing tamarind and occur in the central, northern, southern and southwestern part of the microwatershed. They have moderate limitations of rooting depth and calcareousness. An area of about 109 ha (20%) is currently not suitable (Class N1) for growing tamarind and distributed in the northern, southeastern, southwestern, western and northwestern part of the microwatershed. They have severe limitations of rooting depth and calcareousness.




Fig. 7.27 Land Suitability map of Tamarind

7.28 Land Suitability for Marigold (*Tagetes erecta*)

Marigold is one of the most important flower crop grown in an area of 9108 ha in almost all the districts of the state. The crop requirements (Table 7.29) for growing marigold were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing marigold was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.28.

Highly suitable (Class S1) lands for growing marigold cover an area of about 34 ha (6%) and occur in the eastern part of the microwatershed. Maximum area of about 458 ha (83%) is moderately suitable (Class S2) for growing marigold and distributed in the major part of the microwatershed with minor limitations of texture, rooting depth, calcareousness, drainage and gravelliness. An area of about 53 ha (10%) is marginally suitable (Class S3) for growing marigold and occur in the northwestern and southeastern part of the microwatershed with moderate limitations of rooting depth and calcareousness.




Fig. 7.28 Land Suitability map of Marigold

7.29 Land Suitability for Chrysanthemum (*Chrysanthemum indicum*)

Chrysanthemum is one of the most important flower crop grown in an area of 4978 ha in almost all the districts of the State. The crop requirements (Table 7.30) for growing chrysanthemum were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing chrysanthemum was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed is given in Figure 7.29.

An area of about 34 ha (6%) is highly suitable (Class S1) for growing chrysanthemum and occur in the eastern part of the microwatershed. Maximum area of about 458 ha (83%) is moderately suitable (Class S2) for growing chrysanthemum and distributed in the major part of the microwatershed with minor limitations of texture, rooting depth, calcareousness, drainage and gravelliness. An area of about 53 ha (10%) is marginally suitable (Class S3) for growing chrysanthemum and occur in the northwestern and southeastern part of the microwatershed with moderate limitations of rooting depth and calcareousness.




Fig. 7.29 Land Suitability map of Chrysanthemum

7.30 Land Suitability for Jasmine (*Jasminum sp.*)

Jasmine is one of the most important flower crop grown in an area of 803 ha in almost all the districts of the State. The crop requirements (Table 7.31) for growing jasmine were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing jasmine was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.30.

Highly suitable (Class S1) lands for growing jasmine cover an area of about 34 ha (6%) and occur in the eastern part of the microwatershed. An area of about 57 ha (10%) is moderately suitable (Class S2) for growing jasmine and occur in the northern, southeastern, southwestern and western part of the microwatershed. They have minor limitations of calcareousness and rooting depth. Major area of about 458 ha (83%) is marginally suitable (Class S3) for growing jasmine and occur in the major part of the microwatershed. They have moderate limitations of rooting depth, texture, drainage and calcareousness.




Fig. 7.30 Land Suitability map of Jasmine

7. 31 Land Suitability for Crossandra (*Crossandra infundibuliformis*.)

Crossandra is one of the most important flower crop grown in all the districts of the state. The crop requirements (Table 7.32) for growing crossandra were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing crossandra was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.31.

An area of about 34 ha (6%) is highly suitable (Class S1) for growing crossandra and occur in the eastern part of the microwatershed. An area of about 166 ha (30%) is moderately suitable (Class S2) for growing crossandra and occur in the central, southern, northeastern, western and northwestern part of the microwatershed. They have minor limitations of texture and calcareousness. Major area of about 343 ha (62%) is marginally suitable (Class S3) for growing crossandra and occur in the major part of the microwatershed. They have moderate limitations of rooting depth, calcareousness and texture.




Fig. 7.31 Land Suitability map of Crossandra

Table 7.1 Soil-Site Characteristics of Naregallu-3 Microwatershed

Soil Map Units	Climate (P) (mm)	Growing period (Days)	Drainage Class	Soil depth (cm)	Soil texture		Gravelliness		AWC (mm/m)	Slope (%)	Erosion	pH	EC	ESP	CEC [Cmol (p+)kg-1]	BS (%)
					Surf-ace	Sub-surface	Sur-face	Sub-surface								
MRDiB2	662	<90	WD	>150	sc	scl	<15	<15	101-150	1-3	Moderate	-	-	-	-	-
RTRiB2	662	<90	WD	>150	sc	c	<15	<15	151-200	1-3	Moderate	5.08	0.03	2.06	9.21	50.50
MTLmB2	662	<90	WD	25-50	c	gc	<15	15-35	51-100	1-3	Moderate	8.27	0.202	0.69	36.64	-
RNKmB1	662	<90	MWD	50-75	c	c	<15	<15	51-100	1-3	Slight	8.86	0.483	6.78	37.00	-
RNKmB2	662	<90	MWD	50-75	c	c	<15	<15	51-100	1-3	Moderate	8.86	0.483	6.78	37.00	-
DRLmB2	662	<90	MWD	75-100	c	c	<15	<15	151-200	1-3	Moderate	8.78	0.42	5.62	49.70	100
NSPmA1	662	<90	MWD	75-100	c	c	<15	<15	101-150	0-1	Slight	9.16	0.615	8.60	51.09	-
NSPmB2	662	<90	MWD	75-100	c	c	<15	<15	101-150	1-3	Moderate	9.16	0.615	8.60	51.09	-
GRHmB1	662	<90	MWD	100-150	c	c	<15	<15	>200	1-3	Slight	9.08	0.23	7.11	63.21	100
GRHmB2	662	<90	MWD	100-150	c	c	<15	<15	>200	1-3	Moderate	9.08	0.23	7.11	63.21	100
GRHmB2g1	662	<90	MWD	100-150	c	c	15-35	<15	>200	1-3	Moderate	9.08	0.23	7.11	63.21	100
HDLmB2	662	<90	MWD	100-150	c	c	<15	<15	>200	1-3	Moderate	9.06	0.371	5.09	62.33	-
HDLmA1	662	<90	MWD	100-150	c	c	<15	<15	>200	0-1	Slight	9.06	0.371	5.09	62.33	-
KVRiB2	662	<90	MWD	100-150	sc	c	<15	<15	>200	1-3	Moderate	8.4	0.265	0.60	43.25	-
KVRmB1	662	<90	MWD	100-150	c	c	<15	<15	>200	1-3	Slight	8.4	0.265	0.60	43.25	-
KVRmB1g1	662	<90	MWD	100-150	c	c	15-35	<15	>200	1-3	Slight	8.4	0.265	0.60	43.25	-
BGPmB1	662	<90	MWD	>150	c	c	<15	<15	>200	1-3	Slight	9.2	0.27	3.84	19.60	100
BDRmA1	662	<90	MWD	>150	c	c	<15	<15	>200	0-1	Slight	8.73	0.203	4.37	40.56	-
BDRmB1	662	<90	MWD	>150	c	c	<15	<15	>200	1-3	Slight	8.73	0.203	4.37	40.56	-

*Symbols and abbreviations are according to Field Guide for LRI under Sujala-III

Table 7.2 Land suitability criteria for Sorghum

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	26–30	30–34; 24–26	34–40; 20–24	>40; <20
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristics					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, c (red), c (black)	scl, cl	ls, sl	-
	pH	1:2.5	5.5-7.8	5.0-5.5 7.8-9.0	>9.0	-
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	10-15
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

Table 7.3 Land suitability criteria for Maize

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	30-34	35-38 26-30	38-40 26-20	
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc	c (red), c (black)	ls, sl	-
	pH	1:2.5	5.5-7.8	5.0-5.5 7.8-9.0	>9.0	-
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	-
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

Table 7.4 Land suitability criteria for Bajra

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-32	33-38 24-27	39-40 20-23	<20
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm	500-750	400-500	200-400	<200
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	Sl, scl, cl,sc,c (red)	C (black)	ls	-
	pH	1:2.5	6.0-7.8	5.0-5.5 7.8-9.0	5.5-6.0 >9.0	
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	15-35	35-60	>60	
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	1-3	3-5	5-10	>10

Table 7.5 Land suitability criteria for Groundnut

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	24–33	22–24; 33–35	20–22; 35–40	<20; >40
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. Well drained	Poorly drained	Very Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl	sl,cl, sc	c (red), c (black), ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<35	35-60	>60	
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.6 Land suitability criteria for Sunflower

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	24–30	30–34; 20–24	34–38; 16–20	>38; <16
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	mod. Well drained	-	Poorly to very drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	cl, sc,c (red), c (black)	scl	ls, sl	-
	pH	1:2.5	6.5-7.8	7.8-8.4 5.5-6.5	8.4-9.0; 5.0-5.5	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.7 Land suitability criteria for Cotton

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	22-32	>32	<19	-
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well to moderately well	Poorly drained/Somewhat excessively drained	-	very poorly/excessively drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, c (red,black)	cl	scl	ls, sl
	pH	1:2.5	6.5-7.8	7.8-8.4	5.5-6.5 8.4->9.0	<5.5
	CEC	C mol (p+)Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	50-100	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	<3	3-5	-	>5

Table 7.8 Land suitability criteria for Red gram

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	30-35(G) 20-25(AV) 15-18 (F&PS) 35-40(M)	25-30(G) 20-25 (AV) 12-15 (F&PS) 30-35(M)	20-25(G) 15-20(AV) 10-12 (F&PS) 25-30(M)	< 20 <15 <10 <25
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. Well drained	Poorly drained	Very Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, c (red)	c (black),sl, scl, cl	ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-9.0	5.0-5.5 >9.0	-
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-50	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<1.0	1.0-2.0	>2.0	
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.9 Land suitability criteria for Bengal gram

Land use requirement			Rating			
Soil –site characteristics	Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)	
Climatic regime	Mean temperature in growing season	°C	20–25	25–30; 15–20	30–35; 10–15	>35; <10
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. Well drained	Poorly drained	Very Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	C (black)	-	c (red), scl, cl, sc	ls, sl
	pH	1:2.5	6.0-7.8	5.0-6.0 7.8-9.0	>9.0	-
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	-
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.10 Land suitability criteria for Chilli

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	25-32	33-35 20-25	35-38 <20	>38
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl, cl, sc	c (black), sl	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.11 Land suitability criteria for Tomato

Land use requirement		Rating				
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	25-28	29-32 20-24	15-19 33-36	<15 >36
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl, scl, cl, sc, c (red)	-	ls, c(black)	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.12 Land suitability criteria for Brinjal

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	Well drained	Moderately well drained	Poorly drained	V. Poorly drained
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class				
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl, scl, cl, sc c (red)	-	ls, c (black)	-
	pH	1:2.5	6.0-7.3	7.3-8.4 5.0-6.0	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	>60
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.13 Land suitability criteria for Onion

Land use requirement		Rating				
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	20-30	30-35	35-40	>40
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately /imperfectly	-	Poorly to V poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl,scl,cl,sc,c (red)	-	c (Black),ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<1.0	1.0-2.0	2.0-4.0	<4
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.14 Land suitability criteria for Bhendi

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	25-28	29-32 20-24	15-19 33-36	<15 >36
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Imperfectly drained	Poorly to very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl,sc, c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.15 Land suitability criteria for Drumstick

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, scl, cl, c (red)	sl, c (black)	ls	s
	pH	1:2.5	6.0-7.3	5.0-5.5 7.3-7.8	5.5-6.0 7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<35	35-60	60-80	>80
Soil toxicity	Salinity (EC saturation extract)	dS/m				
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-10	-	>10

Table 7.16 Land suitability criteria for Mulberry

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	24–28	22–24; 28–32	32–38; 22–18	>38; <18
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V. Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, cl, scl	c (red)	c (black), sl, ls	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.8-8.4	7.3-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	0-35	35-60	60-80	>80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

Note: Suitability evaluation only for Mulberry leaf not for Silk worm rearing

Table 7.17 Land suitability criteria for Mango

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-32	24-27 33-35	36-40	20-24
	Min temp. before flowering	°C	10-15	15-22	>22	-
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration	Days				
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	-	ls, sl, c (black)	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>150	100-150	75-100	<75
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.18 Land suitability criteria for Sapota

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-32	33-36 24-27	37-42 20-23	>42 <18
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	-	Poorly to very drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	sl	ls, c (black)	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.19 Land suitability criteria for Pomegranate

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	30-34	35-38 25-29	39-40 15-24	
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl,cl, sc, c (red)	c (black),sl	ls	-
	pH	1:2.5	5.5-7.8	7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.20 Land suitability criteria for Guava

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-32	33-36 24-27	37-42 20-23	
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	sl	c (black), ls	-
	pH	1:2.5	6.0-7.8	5.0-6.0	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.21 Land suitability criteria for Jackfruit

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. well	Poorly	V. Poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	-	sl, ls, c (black)	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	>60
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10-

Table 7.22 Land suitability criteria for Jamun

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well	Mod. well	Poorly	V.Poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c(red)	sl, c (black)	ls	-
	pH	1:2.5	6.0-7.8	5.0-6.0	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>150	100-150	50-100	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	>60
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

Table 7.23 Land suitability criteria for Musambi

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-30	31-35 24-27	36-40 20-23	>40 <20
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately drained	poorly	Very poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c	sl	ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.24 Land suitability criteria for Lime

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-30	31-35 24-27	36-40 20-23	>40 <20
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately drained	poorly	Very poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c	sl	ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.25 Land suitability criteria for Cashew

Land use requirement		Rating				
Soil –site characteristics	Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)	
Climatic regime	Mean temperature in growing season	°C	32 to 34	28 to 32; 34 to 38	24 to 28; 38 to 40	<20; >40
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	-	sl, ls	c (black)
	pH	1:2.5	5.5-6.5	5.0-5.5 6.5-7.3	7.3-7.8	>7.8
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-10	>10	-

Table 7.26 Land suitability criteria for Custard apple

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	Scl, cl, sc, c (red), c (black)	-	Sl, ls	-
	pH	1:2.5	6.0-7.3	5.5-6.0 7.3-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15-35	35-60	60-80	-
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	>5	-

Table 7.27 Land suitability criteria for Amla

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod.well drained	Poorly drained	V. Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	c (black)	ls, sl	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15-35	35-60	60-80	-
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

Table 7.28 Land suitability criteria for Tamarind

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod.well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl,sc, c (red)	sl, c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>150	100-150	75-100	<75
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

Table 7.29 Land suitability criteria for Marigold

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	18-23	17-15 24-35	35-40 10-14	>40 <10
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl,scl, cl, sc, c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%				
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.30 Land suitability criteria for Chrysanthemum

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	18-23	17-15 24-35	35-40 10-14	>40 <10
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl,scl, cl, sc, c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%				
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.31 Land suitability criteria for Jasmine (irrigated)

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	18-23	17-15 24-35	35-40 10-14	-
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	sl	ls, c (black)	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%				
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.32 Land suitability criteria for Crossandra

Land use requirement		Rating				
Soil –site characteristics	Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)	
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	-	Poorly to very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c(red)	sl,	c (black),ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%				
Erosion hazard	Slope	%	<3	3-5	5-10	>10

7.32 Land Management Units (LMUs)

The 19 soil map units identified in Naregallu-3 microwatershed have been grouped into 4 Land Management Units (LMUs) for the purpose of preparing a Proposed Crop Plan. Land Management Units are grouped based on the similarities in respect of the type of soil, the depth of the soil, the surface soil texture, gravel content, AWC, slope, erosion etc. and a Land Management Unit map (Fig.7.32) has been generated. These Land Management Units are expected to behave similarly for a given level of management.

The map units that have been grouped into 4 Land Management Units along with brief description of soil and site characteristics are given below.

LMUs	Mapping unit	Soil and site characteristics
1	428.BDRmA1 430.BDRmB1 396.BGPmB1 371.GRHmB1 373.GRHmB2 374.GRHmB2g1 382.HDLmB2 378.HDLmA1 384.KVRiB2 388.KVRmB1 389.KVRmB1g1 350.DRLmB2 359.NSPmA1 362.NSPmB2	Moderately deep to very deep black calcareous clay soils, 0-3% slope, slight to moderate erosion, non-gravelly to gravelly (<15-35%).
2	283.MRDIB2 288.RTRiB2	Very deep, red sandy clay to clay soils, 1-3% slope, moderate erosion, non-gravelly (<15%).
3	333.RNKmB1 336.RNKmB2	Moderately shallow, black calcareous clay soils, 1-3% slope, slight to moderate erosion, non-gravelly (<15%).
4	310.MTLmB2	Shallow, black calcareous clay soils, 1-3% slope, moderate erosion, non-gravelly (<15%).




Fig 7.32 Land Management Units map of Naregallu-3 microwatershed

7.33 Proposed Crop Plan for Naregallu-3 Microwatershed

After assessing the land suitability for the 31 crops, the proposed crop plan has been prepared for the 4 identified LMUs by considering only the highly (Class S1) and moderately (Class S2) suitable lands for each of the 31 crops. The resultant proposed crop plan is presented in Table 7.33.

Table 7.33 Proposed Crop Plan for Naregallu-3 Microwatershed

LMU	Soil Map Units	Survey Number	Soil and site characteristics	Field Crops	Horticulture Crops	Suitable Interventions
1	428.BDRmA1 430.BDRmB1 396.BGPmB1 371.GRHmB1 373.GRHmB2 374.GRHmB2g1 382.HDLmB2 378.HDLmA1 384.KVRiB2 388.KVRmB1 389.KVRmB1g1 350.DRLmB2 359.NSPmA1 362.NSPmB2	Madhinura: 71,72,73,74,75,76,77,78,79,80,83,84,85,86,87,109,111,113,117,118,119,120,122,123,124,125,128,130,131,132,133,135,136,138,139,140,141,142,143,144,146,148,149 Neregalla: 80,81,117,118,119,120,125,126,127,128,169,173,175,176,179,180,181,182,183,184,185,186,187,188,189,190,191,192,193,194,195,198,199,200,201,202,208,209,210,211,212,213,214,215,216,217,218,219,222,223,224,225,226	Moderately deep to very deep black calcareous clay soils, 0-3% slope, slight to moderate erosion, non-gravelly to gravelly (<15-35%).	Maize, Sorghum, Sunflower, Cotton, Bengal gram, Safflower, Linseed, Bajra , Soybean	Fruit crops: Sapota, Pomegranate, Jamun, Lime, Musambi, Tamarind, Amla, Custard apple Vegetables: Drumstick, Chilli, Coriander, Tomato, Bhendi Flowers: Marigold, Chrysanthemum, Crossandra, Jasmine	Application of FYM, Biofertilizers and micronutrients, drip irrigation, mulching, suitable soil and water conservation practices
2	283.MRDIB2 288.RTRiB2	Neregalla: 77,78,79,203,204,205,206,207,220, 221	Very deep, red sandy clay to clay soils, 1-3% slope, moderate erosion, non-gravelly (<15%).	Maize, Sorghum, Sunflower, Bajra, Finger millet, Groundnut, Red gram, Cowpea, Field bean, Castor, Mulberry	Fruit crops: Mango, Pomegranate, Guava, Sapota, Jackfruit, Jamun, Tamarind, Lime, Musambi, Amla, Custard apple, Cashew Vegetable crops: Drumstick, Tomato, Bhendi, Chilli, Brinjal, Onion, Curry leaves Flower crops: Marigold,	Drip irrigation, mulching, suitable soil and water conservation practices (Crescent Bunding with Catch Pit etc)

LMU	Soil Map Units	Survey Number	Soil and site characteristics	Field Crops	Horticulture Crops	Suitable Interventions
					Chrysanthemum, Jasmine, Crossandra	
3	333.RNKmB1 336.RNKmB2	Madhinura: 81,82,121 Neregalla: 164,167,168,170, 171,172,177,178, 196,197	Moderately shallow, black calcareous clay soils, 1-3% slope, slight to moderate erosion, non-gravelly (<15%).	Sorghum, Bajra, Bengal gram, linseed, Safflower, Coriander	Fruit crops: Amla, Custard apple Flower crops: Marigold, Jasmine Chrysanthemum	Application of FYM, Biofertilizers and micronutrients, drip irrigation, mulching, suitable soil and water conservation practices
4	310.MTLmB2	Madhinura: 114,115,116,134 Neregalla: 116,121,122,123	Shallow, black calcareous clay soils, 1-3% slope, moderate erosion, non-gravelly (<15%).	Bengal gram	Agri-Silvi-Pasture: Hybrid Napier, <i>Styloxanthes hamata</i> , <i>Styloxanthes scabra</i>	Use of short duration varieties, sowing across the slope

SOIL HEALTH MANAGEMENT

8.1 Soil Health

Soil health is basic to plant health and plant health is basic to human and bovine health. Soil is fundamental to crop production. Without soil, no food could be produced nor would livestock be fed on a large scale. Because it is finite and fragile, soil is a precious resource that requires special care from its users.

Soil health or the capacity of the soil to function is critical to human survival. Soil health has been defined as: “the capacity of the soil to function as a living system without adverse effect on the ecosystem”. Healthy soils maintain a diverse community of soil organisms that help to form beneficial symbiotic associations with plant roots, recycle essential plant nutrients, improve soil structure with positive repercussions for soil, water and nutrient holding capacity and ultimately improve crop production and also contribute to mitigating climate change by maintaining or increasing its carbon content.

Functional interactions of soil biota with organic and inorganic components, air and water determine a soil’s potential to store and release nutrients, and water to plants and to promote and sustain plant growth. Thus, maintaining soil health is vital to crop production and conserve soil resource base for sustaining agriculture.

The most important characteristics of a healthy soil are

- Good soil tilth
- Sufficient soil depth
- Good water storage and good drainage
- Adequate supply, but not excess of nutrients
- Large population of beneficial organisms
- Small proportion of plant pathogens and insect pests
- Low weed pressure
- Free of chemicals and toxins that may harm the crop
- Resistance to degradation
- Resilience when unfavorable conditions occur

Characteristics of Naregallu-3 Microwatershed

- ❖ The soil phases with sizeable area identified in the microwatershed belonged to the soil series of Kavalur (KVR) series occupies major area of 116 ha (21%) followed by Gatareddihal (GRH) 58 ha (10%), Ravanaki (RNK) 57 ha (10%), Bardur (BDR) 56 ha (10%), Dambarahalli (DRL) 55 ha (10%), Handrala (HDL) 55 ha (10%), Muttal (MTL) 53 ha (10%), Narasapura (NSP) 34 ha (6%), Muradi (MRD) 33 ha (6%), Budagumpa (BGP) 25 ha (5%) and Ranatur (RTR) 2 ha (<1%).
- ❖ As per land capability classification, maximum area of about 490 ha (89%) in the microwatershed falls under good lands (Class II) with minor limitations of soil and

erosion. An area of about 54 ha (10%) is under moderately good lands (Class III) with severe limitations of soil and erosion.

- ❖ On the basis of soil reaction, an area of about 206 ha (37%) are strongly alkaline (pH 8.4-9.0) and 337 ha (61%) are very strongly alkaline (pH >9.0) in soil reaction.

Soil Health Management

The following actions are required to improve the current land husbandry practices that provide a sound basis for the successful adoption of sustainable crop production system.

Alkaline soils

Entire area in the microwatershed has comes under alkaline condition (Strongly alkaline to very strongly alkaline).

1. Regular addition of organic manure, green manuring, green leaf manuring, crop residue incorporation and mulching needs to be taken up to improve the soil organic matter status.
2. Application of Biofertilizers (Azospirillum, Azotobacter, Rhizobium).
3. Application of 25% extra N and P (125 % RDN&P).
4. Application of ZnSO₄ – 12.5 kg/ha (once in three years).
5. Application of Boron – 5 kg/ha (once in three years).

Besides the above recommendations, the best transfer of technology options are also to be adopted.

Soil Degradation

Soil erosion is one of the major factor affecting the soil health in the microwatershed. An area of about 282 ha (51%) is suffering from moderate erosion. These areas need immediate soil and water conservation and, other land development and land husbandry practices for restoring soil health.

Dissemination of Information and Communication of Benefits

Any large scale implementation of soil health management requires that supporting information is made available widely, particularly through channels familiar to farmers and extension workers. Given the very high priority attached to soil health especially by the Central Government on issuing Soil-Health Cards to all the farmers, media outlets like Regional, State and National Newspapers, Radio and Dooradarshan programs in local languages but also modern information and communication technologies such as Cellular phones and the Internet, which can be much more effective in reaching the younger farmers.

Inputs for Net Planning (Saturation Plan) and Interventions needed

Net planning in IWMP is focusing on preparation of

1. Soil and Water Conservation Treatment Plans for each plot or farm.
2. Productivity enhancement measures/ interventions for existing crops/livestock/other farm enterprises.
3. Diversification of farming mainly with perennial horticultural crops and livestock.
4. Improving livelihood opportunities and income generating activities.

In this connection, how various outputs of Sujala-III are of use in addressing these objectives of Net Planning are briefly presented below.

- ❖ **Soil Depth:** The depth of a soil decides the amount of moisture and nutrients it can hold, what crops can be taken up or not, depending on the rooting depth and the length of growing period available for raising any crop. Deeper the soil, better for a wide variety of crops. If sufficient depth is not available for growing deep rooted crops, either choose medium or short duration crops or deeper planting pits need to be opened and additional good quality soil brought from outside has to be filled into the planting pits.
- ❖ **Surface Soil Texture:** Lighter soil texture in the top soil means, better rain water infiltration, less run-off and soil moisture conservation, less capillary rise and less evaporation losses. Lighter surface textured soils are amenable to good soil tilth and are highly suitable for crops like groundnut, root vegetables (carrot, radish, potato etc) but not ideal for crops that need stagnant water like lowland paddy. Heavy textured soils are poor in water infiltration and percolation. They are prone for sheet erosion; such soils can be improved by sand mulching. The technology that is developed by the AICRP-Dryland Agriculture, Vijayapura, Karnataka can be adopted.
- ❖ **Gravelliness:** More gravel content is favorable for run-off harvesting but poor in soil moisture storage and nutrient availability. It is a significant parameter that decides the kind of crop to be raised.
- ❖ **Land Capability Classification:** The land capability map shows the areas suitable and not suitable for agriculture and the major constraints in each of the plot/survey number. Hence, one can decide what kind of enterprise is possible in each of these units. In general, erosion and soil are the major constraints in Naregallu-3 Microwatershed.
- ❖ **Organic Carbon:** The OC content is low (0.5%) in 420 ha (76%) and medium (0.5-0.75%) in 124 ha (23%). These areas needs to be further improved by applying farmyard manure and rotating crops with cereals and legumes or mixed cropping.
- ❖ **Promoting Green Manuring:** Growing of green manuring crops costs Rs. 1250/ha (green manuring seeds) and about Rs. 2000/ha towards cultivation that totals to Rs. 3250/- per ha. On the other hand, application of organic manure @ 10 tons/ha costs Rs. 5000/ha. The practice needs to be continued for 2-3 years or more. Nitrogen fertilizer needs to be supplemented by 25% in addition to the recommended level in entire area where OC is low to medium (<0.5-0.75%). For example, for rainfed maize,

recommended level is 50 kg N per ha and an additional 12 kg /ha needs to be applied for all the crops grown in these plots.


- ❖ **Available Phosphorus:** An area of about 71 ha (13%) is medium (23-57 kg/ha) and 472 ha (86%) is low (>23 kg/ha) in available phosphorus content. Hence, for all the plots, where available phosphorus is low to medium, for all the crops, 25% additional P-needs to be applied
- ❖ **Available Potassium:** Available potassium content is medium (145-337 kg/ha) in 237 ha (43%) and high in 307 ha (56%) of the microwatershed. All the plots, where available potassium is medium, for all the crops, additional 25% of potassium may be applied.
- ❖ **Available Sulphur:** Available sulphur is a very critical nutrient for oilseed crops, Entire cultivated area of the microwatershed is low (<10 ppm) in available sulphur content. Hence these areas need to be applied with magnesium sulphate or gypsum or Factamphos (p) fertilizer (13% of sulphur) for 2-3 years for the deficiency to be corrected.
- ❖ **Available Boron:** An area of about 268 ha (49%) is low (<0.5 ppm), 245 ha (44%) is medium (0.5-1.0 ppm) and 31 ha (6%) is high (>1.0 ppm) in the available boron content. Low to medium (<0.5-1.0 ppm) areas need to be applied with sodium borate @ 10kg/ha as soil application or 0.2% borax as foliar spray to correct the deficiency.
- ❖ **Available Iron:** Available iron content is deficient (<4.5 ppm) in 420 ha (76%) and sufficient (>4.5 ppm) in 124 ha (22%) area of the microwatershed. For deficient areas, iron sulphate @ 25 kg/ha needs to be applied for 2-3 years to correct the deficiency.
- ❖ **Available Manganese:** Entire cultivated area of the microwatershed is sufficient (>1.0 ppm) in the available manganese content.
- ❖ **Available Copper:** Entire cultivated area of the microwatershed is sufficient (>0.2 ppm) in the available copper content.
- ❖ **Available Zinc:** Entire cultivated area of the microwatershed is deficient (<0.6 ppm) in the available zinc content. For deficient areas, application of zinc sulphate @ 25kg/ha is recommended.
- ❖ **Soil Alkalinity:** Entire area in the microwatershed has soils that are strongly alkaline to very strongly alkaline. These areas need application of gypsum and wherever calcium is in excess, iron pyrites and element sulphur can be recommended. Management practices like treating repeatedly with good quality water to drain out the excess salts and provision of subsurface drainage and growing of salt tolerant crops like Casuarina, Acasia, Neem, Ber etc, are recommended.

- ❖ **Land Suitability for various crops:** Areas that are highly, moderately and marginally suitable and not suitable for growing various crops are indicated. Along with the suitability, various constraints that are limiting the productivity are also indicated. For example, in case of cotton, gravel content, rooting depth and salinity/alkalinity are the major constraints in various plots. With suitable management interventions, the productivity can be enhanced. In order to increase water holding capacity of light textured soils, growing of green manure crops and application of organic manure is recommended.

SOIL AND WATER CONSERVATION TREATMENT PLAN

For preparing soil and water conservation treatment plan for Naregallu-3 Microwatershed, the land resource inventory database generated under Sujala-III project has been transformed as information through series of interpretative (thematic) maps using soil phase map as a base. The various thematic maps (1:7920 scale) generated were

- Soil depth
- Surface soil texture
- Available water capacity
- Soil slope
- Soil gravelliness
- Land capability
- Present land use and land cover
- Crop suitability maps
- Rainfall map
- Hydrology
- Water Resources
- Socio-economic data
- Contour plan with existing features- network of waterways, pothissa boundaries, cut up/ minor terraces etc.
- Cadastral map (1:7920 scale)
- Satellite imagery (1:7920 scale)



Apart from these, Hand Level/ Hydro Marker/ Dumpy Level/ Total Station and Kathedars' List needs to be collected.

Steps for Survey and Preparation of Treatment Plan

The boundaries of Land User Groups' and Survey No. boundaries are traced in the field.

- Naming of user groups and farmers
- Identification of arable and non arable lands
- Identification of drainage lines and gullies
- Identification of non treatable areas
- Identification of priority areas in the arable lands
- Treatment plan for arable lands
- Location of water harvesting and recharge structures

9.1 Treatment Plan

The treatment plan recommended for arable lands is briefly described below.


9.1.1 Arable Land Treatment

A. BUNDING

Steps for Survey and Preparation of Treatment Plan		USER GROUP-1
Cadastral map (1:7920 scale) is enlarged to a scale of 1:2500 scale Existing network of waterways, pothissa boundaries, grass belts, natural drainage lines/ watercourse, cut ups/ terraces are marked on the cadastral map to the scale Drainage lines are demarcated into		
Small gullies	(up to 5 ha catchment)	
Medium gullies	(5-15 ha catchment)	
Ravines	(15-25 ha catchment) and	
Halla/Nala	(more than 25ha catchment)	

Measurement of Land Slope

Land slope is estimated or determined by the study and interpretation of contours or by measurement in the field using simple instruments like Hand Level or Hydromarker.



Vertical and Horizontal intervals between bunds as recommended by the Watershed Development Department.

Slope percentage	Vertical interval (m)	Corresponding Horizontal Distance (m)
2 - 3%	0.6	24
3 - 4%	0.9	21
4 - 5%	0.9	21
5 - 6%	1.2	21
6 - 7%	1.2	21

Note: i) The above intervals are maximum.

(ii) Considering the slope class and erosion status (A1... A= 0-1% slope, 1= slight erosion.) the intervals have to be decided.

Bund length recording: Considering the contour plan and the existing grass belts/partitions, the bunds are aligned and lengths are measured.

Section of the Bund

Bund section is decided considering the soil texture class and gravelliness class (bg0b= loamy sand, g0 = <15% gravel). The recommended sections for different soils are given below.

Recommended Bund Section

Top width (m)	Base width (m)	Height (m)	Side slope (Z:1;H:V)	Cross section (sq m)	Soil Texture	Remarks
0.3	0.9	0.3	01:01	0.18	Sandy loam	Vegetative bund
0.3	1.2	0.3	1.5:1	0.225	Sandy clay	
0.3	1.2	0.5	0.9:1	0.375	Red gravelly soils	
0.3	1.2	0.6	0.75:1	0.45		
0.3	1.5	0.6	01:01	0.54	Red sandy loam	
0.3	2.1	0.6	1.5:1	0.72	Very shallow black clayey soils	
0.45	2	0.75	01:01	0.92		
0.45	2.4	0.75	1.3:1	1.07	Shallow black clayey soils	
0.6	3.1	0.7	1.78:1	1.29	Medium black clayey soils	
0.5	3	0.85	1.47:1	1.49		

Formation of Trench cum Bund

Dimensions of the Borrow Pits/ Trenches to be excavated (machinery are decided considering the Bund Section).

Details of Borrow Pit dimensions are given below

TRENCH CUM BUND

IDEAL FOR HORTICULTURE CROPS

'A' FRAME FOR INTERBUND MANAGEMENT

ಸಮಸಾರಕಳು ರೇಖೆ

1. ಸಮಸಾರಕಳು ಉಳಿಸುವುದು
2. ಸಮಸಾರಕಳು ಬಿಡುವುದು/ನಾಟಿ

Size of Borrow Pits/ Trench recommended for Trench cum Bund (by machinery)

Bund section	Bund length	Earth quantity	Pit				Berm (pit to pit)	Soil depth Class
			L(m)	W(m)	D(m)	Quantity (m ³)		
0.375	6	2.25	5.85	0.85	0.45	2.24	0.15	Shallow
0.45	6	2.7	5.4	1.2	0.43	2.79	0.6	Shallow
0.45	6	2.7	5	0.85	0.65	2.76	1	Moderately Shallow
0.54	5.6	3.02	5.5	0.85	0.7	3.27	0.1	Moderately shallow
0.54	5.5	2.97	5	1.2	0.5	3	0.5	Shallow
0.72	6.2	4.46	6	1.2	0.7	5.04	0.2	Moderately shallow
0.72	5.2	3.74	5.1	0.85	0.9	3.9	0.1	Moderately deep

B. Waterways

- Existing waterways are marked on the cadastral map (1:7920 scale) and their dimensions are recorded.
- Considering the contour plan of the MWS, additional waterways/ modernization of the existing ones can be thought of.
- The design details are given in the Manual.

C. Farm Ponds

Waterways and the catchment area will give an indication on the size of the Farm Pond. Location of the pond can be decided based on the contour plan/ field condition and farmers' need/desire.

D. Diversion Channel

Existing EPT/ CPT are marked on the cadastral map. Looking to the need, these can be modernized or fresh diversion channel can be proposed and runoff from this can be stored in *Gokatte/* Recharge ponds.

9.1.2 Non-Arable Land Treatment

Depending on the gravelliness and crops preferred by the farmers, the concerned authorities can decide appropriate treatment plan. The recommended treatments may be Contour Trench, Staggered Trench, Crescent Bund, Boulder Bund or Pebble Bund.

9.1.3 Treatment of Natural Water Course/ Drainage Lines

- a) The cadastral map has to be updated as regards the network of drainage lines (gullies/ *nalas*/ *hallas*) and existing structures are marked to the scale and storage capacity of the existing water bodies are documented.
- b) The drainage line will be demarcated into Upper Reach, Middle Reach and Lower Reach.
- c) Considering the Catchment, *Nala* bed and bank conditions, suitable structures are decided.
- d) Number of storage structures (Check dam/ *Nala* bund/ Percolation tank) will be decided considering the commitments and available runoff in water budgeting and quality of water in the wells and site suitability.
- e) Detailed Leveling Survey using Dumpy Level / Total Station has to be carried out to arrive at the site-specific designs as shown in the Manual.
- f) The location of ground water recharge structures are decided by examining the lineaments and fracture zones from geological maps.
- g) Rainfall intensity data of the nearest Rain Gauge Station is considered for Hydrologic Designs.
- h) Silt load to the Storage/Recharge Structures is reduced by providing vegetative, boulder and earthen checks in the natural water course. Location and design details are given in the Manual.

9.2 Recommended Soil and Water Conservation Measures

The appropriate conservation structures best suited for each of the land parcel/survey number (Appendix-I) are selected based on the slope per cent, severity of erosion, amount of rainfall, land use and soil type. The different kinds of conservation structures recommended are

1. Graded / Strengthening of Bunds
2. Trench cum Bunds (TCB)
3. Trench cum Bunds / Strengthening
4. Crescent Bunds

A map (Fig. 9.1) showing soil and water conservation plan with different kinds of structures recommended has been prepared which shows the spatial distribution and extent of area. An area of about 34 ha (6%) needs Trench cum Bunding, 463 ha (84%) needs Graded Bunding and 47 ha (8%) needs strengthening of existing bunds.

The conservation plan prepared may be presented to all the stakeholders including farmers and after considering their suggestions, the conservation plan for the microwatershed may be finalised in a participatory approach.




Fig. 9.1 Soil and Water Conservation Plan map of Naregallu-3 Microwatershed

9.3 Greening of Microwatershed

As part of the greening programme in the watersheds, it is envisaged to plant a variety of horticultural and other tree plants that are edible, economical and produce lot of biomass which helps to restore the ecological balance in the watersheds. The lands that are suitable for greening programme are non-arable lands (land capability classes V, VI VII and VIII) and also the lands that are not suitable or marginally suitable for growing annual and perennial crops. The method of planting these trees is given below.

It is recommended to open the pits during the 1st week of March along the contour and heap the dug-out soil on the lower side of the slope in order to harness the flowing water and facilitate weathering of soil in the pit. Exposure of soil in the pit also prevents spread of pests and diseases due to scorching sun rays. The pits should be filled with mixture of soil and organic manure during the second week of April and keep ready with sufficiently tall seedlings produced either in poly bags or in root trainer nurseries so that planting can be done during the 2nd or 3rd week of April depending on the rainfall.

The tree species suitable for the area considering rainfall, temperature and adaptability is listed below; waterlogged areas are recommended to be planted with species like Neral (*Syzgium cumini*) and Bamboo. Dry areas are to be planted with species like Honge, Bevu, Seetaphal *etc.*

Dry Deciduous Species			Temp (°C)	Rainfall (mm)
1.	Bevu	<i>Azadiracta indica</i>	21-32	400 -1,200
2.	Tapasi	<i>Holoptelia integrifolia</i>	20-30	500 - 1000
3.	Seetaphal	<i>Anona Squamosa</i>	20-40	400 - 1000
4.	Honge	<i>Pongamia pinnata</i>	20 -50	500-2,500
5.	Kamara	<i>Hardwickia binata</i>	25 -35	400 - 1000
6.	Bage	<i>Albezzia lebbek</i>	20 - 45	500 - 1000
7.	Ficus	<i>Ficus bengalensis</i>	20 - 50	500-2,500
8.	Sisso	<i>Dalbargia Sissoo</i>	20 - 50	500 -2000
9.	Ailanthus	<i>Ailanthus excelsa</i>	20 - 50	500 - 1000
10.	Hale	<i>Wrightia tinctoria</i>	25 - 45	500 - 1000
11.	Uded	<i>Steriospermum chelanoides</i>	25 - 45	500 -2000
12.	Dhupa	<i>Boswellia Serrata</i>	20 - 40	500 - 2000
13.	Nelli	<i>Emblica Officinalis</i>	20 - 50	500 -1500
14.	Honne	<i>Pterocarpus marsupium</i>	20 - 40	500 - 2000
Moist Deciduous Species			Temp (°C)	Rainfall (mm)
15.	Teak	<i>Tectona grandis</i>	20 - 50	500-5000
16.	Nandi	<i>Legarstroemia lanceolata</i>	20 - 40	500 - 4000
17.	Honne	<i>Pterocarpus marsupium</i>	20 - 40	500 - 3000
18.	Mathi	<i>Terminalia alata</i>	20 -50	500 - 2000
19.	Shivane	<i>Gmelina arborea</i>	20 -50	500 -2000
20.	Kindal	<i>T.Paniculata</i>	20 - 40	500 - 1500
21.	Beete	<i>Dalbargia latifolia</i>	20 - 40	500 - 1500
22.	Tare	<i>T. belerica</i>	20 - 40	500 - 2000
23.	Bamboo	<i>Bambusa arundinasia</i>	20 - 40	500 - 2500
24.	Bamboo	<i>Dendrocalamus strictus</i>	20 - 40	500 - 2500
25.	Muthuga	<i>Butea monosperma</i>	20 - 40	400 - 1500
26.	Hippe	<i>Madhuca latifolia</i>	20 - 40	500 - 2000
27.	Sandal	<i>Santalum album</i>	20 - 50	400 - 1000
28.	Nelli	<i>Emblica officinalis</i>	20 - 40	500 - 2000
29.	Nerale	<i>Sizyium cumini</i>	20 - 40	500 - 2000
30.	Dhaman	<i>Grevia tilifolia</i>	20 - 40	500 - 2000
31.	Kaval	<i>Careya arborea</i>	20 - 40	500 - 2000
32.	Harada	<i>Terminalia chebula</i>	20 - 40	500 - 2000

References

1. FAO (1976) Framework for Land Evaluation, Food and Agriculture Organization, Rome.72 pp.
2. FAO (1983) Guidelines for Land Evaluation for Rainfed Agriculture, FAO, Rome, 237 pp.
3. IARI (1971) Soil Survey Manual, All India Soil and Land Use Survey Organization, IARI, New Delhi, 121 pp.
4. Katyal, J.C. and Rattan, R.K. (2003) Secondary and Micronutrients; Research Gap and future needs. Fert. News 48 (4); 9-20.
5. Naidu, L.G.K., Ramamurthy, V., Challa, O., Hegde, R. and Krishnan, P. (2006) Manual Soil Site Suitability Criteria for Major Crops, NBSS Publ. No. 129, NBSS &LUP, Nagpur, 118 pp.
6. Natarajan, A. and Dipak Sarkar (2010) Field Guide for Soil Survey, National Bureau of Soil Survey and Land Use Planning (ICAR), Nagpur, India.
7. Natarajan, A., Rajendra Hegde, Raj, J.N. and Shivananda Murthy, H.G. (2015) Implementation Manual for Sujala-III Project, Watershed Development Department, Bengaluru, Karnataka.
8. Sarma, V.A.K., Krishnan, P. and Budihal, S.L. (1987) Laboratory Manual, Tech. Bull. 23, NBSS &LUP, Nagpur.
9. Sehgal, J.L. (1990) Soil Resource Mapping of Different States of India; Why and How?, National Bureau of Soil Survey and Land Use Planning, Nagpur, 49 pp.
10. Shivaprasad, C.R., R.S. Reddy, J. Sehgal and M. Velayuthum (1998) Soils of Karnataka for Optimizing Land Use, NBSS Publ. No. 47b, NBSS & LUP, Nagpur, India.
11. Soil Survey Staff (2006) Keys to Soil Taxonomy, Tenth edition, U.S. Department of Agriculture/ NRCS, Washington DC, U.S.A.
12. Soil Survey Staff (2012) Soil Survey Manual, Handbook No. 18, USDA, Washington DC, USA.

Appendix I
Neregallu-3 (IT1c) Microwatershed
Soil Phase Information

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Madhin ura	71	0.04	HDLmA1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Bengalgram (Bg)	Not Available	IIs	Graded bunding
Madhin ura	72	0.72	HDLmA1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Ragi (Ra)	Not Available	IIs	Graded bunding
Madhin ura	73	1.48	HDLmA1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Sunflower (Sf)	Not Available	IIs	Graded bunding
Madhin ura	74	0.98	HDLmA1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Maize (Mz)	Not Available	IIs	Graded bunding
Madhin ura	75	0.76	HDLmA1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Bengalgram (Bg)	Not Available	IIs	Graded bunding
Madhin ura	76	4.95	HDLmA1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Bengalgram+Sunflower (Bg+Sf)	Not Available	IIs	Graded bunding
Madhin ura	77	0.1	HDLmA1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Bengalgram (Bg)	Not Available	IIs	Graded bunding
Madhin ura	78	2.31	NSPmA1	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Nearly level (0-1%)	Slight	Sunflower (Sf)	Not Available	IIs	Graded bunding
Madhin ura	79	6.14	NSPmA1	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Nearly level (0-1%)	Slight	Bengalgram (Bg)	Not Available	IIs	Graded bunding
Madhin ura	80	8.7	NSPmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Ragi (Ra)	Not Available	IIs	Graded bunding
Madhin ura	81	1.52	RNKmB2	LMU-3	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIs	Graded bunding
Madhin ura	82	4.86	RNKmB2	LMU-3	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Sunflower (Sf)	Not Available	IIs	Graded bunding
Madhin ura	83	8.26	NSPmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Bengalgram+Sunflower (Bg+Sf)	Not Available	IIs	Graded bunding
Madhin ura	84	6.74	NSPmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Sunflower+Wheat (Sf+Wh)	Not Available	IIs	Graded bunding
Madhin ura	85	3.52	DRLmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Bengalgram+Sunflower (Bg+Sf)	Not Available	IIs	Graded bunding
Madhin ura	86	3.33	NSPmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Sunflower (Sf)	Not Available	IIs	Graded bunding
Madhin ura	87	1.66	NSPmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	IIs	Graded bunding
Madhin ura	109	0.55	DRLmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Bengalgram (Bg)	Not Available	IIs	Graded bunding
Madhin ura	111	0.37	HDLmB2	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Bengalgram (Bg)	Not Available	IIs	Graded bunding
Madhin ura	113	3.82	DRLmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Bengalgram (Bg)	Not Available	IIs	Graded bunding
Madhin ura	114	21	MTLmB2	LMU-4	Shallow (25-50 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Bengalgram+Sunflower (Bg+Sf)	1 Farm Pond	IIIs	Graded bunding
Madhin ura	115	9.71	MTLmB2	LMU-4	Shallow (25-50 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Bengalgram+Jowar+Sunflower (Bg+Jw+Sf)	Not Available	IIIs	Graded bunding

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Madhinura	116	15.11	MTLmB2	LMU-4	Shallow (25-50 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Bengalgram+Maize (Bg+Mz)	Not Available	IIIes	Graded bunding
Madhinura	117	8.32	HDLmB2	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Bengalgram+Sunflower (Bg+Sf)	Not Available	Iles	Graded bunding
Madhinura	118	6.99	HDLmB2	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Sunflower (Sf)	Not Available	Iles	Graded bunding
Madhinura	119	8.96	HDLmB2	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Bengalgram+Sunflower (Bg+Sf)	Not Available	Iles	Graded bunding
Madhinura	120	4.91	HDLmB2	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Bengalgram (Bg)	Not Available	Iles	Graded bunding
Madhinura	121	7.01	RNKmB2	LMU-3	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Fallow land (Fl)	Not Available	Iles	Graded bunding
Madhinura	122	3.79	BDRmA1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Fallow land (Fl)	Not Available	IIs	Graded bunding
Madhinura	123	6.9	BDRmA1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Fallow land (Fl)	Not Available	IIs	Graded bunding
Madhinura	124	4.15	BDRmA1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Bengalgram (Bg)	Not Available	IIs	Graded bunding
Madhinura	125	0.05	BDRmA1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Bengalgram (Bg)	Not Available	IIs	Graded bunding
Madhinura	128	0.49	BDRmA1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Bengalgram+Marigold+Jowar (Bg+Mg+Jw)	Not Available	IIs	Graded bunding
Madhinura	130	5.98	BDRmA1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Bengalgram (Bg)	Not Available	IIs	Graded bunding
Madhinura	131	8.58	BDRmB1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Bengalgram+Sunflower (Bg+Sf)	Not Available	IIs	Graded bunding
Madhinura	132	6.96	BDRmB1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Bengalgram (Bg)	Not Available	IIs	Graded bunding
Madhinura	133	5.89	BDRmB1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Bengalgram (Bg)	Not Available	IIs	Graded bunding
Madhinura	134	1.43	MTLmB2	LMU-4	Shallow (25-50 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Bengalgram (Bg)	Not Available	IIIes	Graded bunding
Madhinura	135	1.97	HDLmB2	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Bengalgram (Bg)	Not Available	Iles	Graded bunding
Madhinura	136	5.12	HDLmB2	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Bengalgram (Bg)	Not Available	Iles	Graded bunding
Madhinura	138	5.23	HDLmB2	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Sunflower (Sf)	Not Available	Iles	Graded bunding
Madhinura	139	4.95	BDRmB1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Sunflower (Sf)	Not Available	IIs	Graded bunding
Madhinura	140	0.7	BDRmB1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Bengalgram (Bg)	Not Available	IIs	Graded bunding
Madhinura	141	2.33	BDRmB1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Bengalgram (Bg)	Not Available	IIs	Graded bunding
Madhinura	142	7.37	BDRmB1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Bengalgram (Bg)	Not Available	IIs	Graded bunding
Madhinura	143	1.8	BDRmA1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Bengalgram (Bg)	Not Available	IIs	Graded bunding

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Madhin ura	144	0.75	HDLmA1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Bengalgram (Bg)	Not Available	IIs	Graded bunding
Madhin ura	146	2.46	BDRmB1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Bengalgram (Bg)	Not Available	IIs	Graded bunding
Madhin ura	148	0.53	BDRmB1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Bengalgram (Bg)	Not Available	IIs	Graded bunding
Madhin ura	149	0.11	BDRmB1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Bengalgram (Bg)	Not Available	IIs	Graded bunding
Neregalla	77	0.16	RTRiB2	LMU-2	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	High (151-200 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	Ile	Trench cum bunding
Neregalla	78	0.87	RTRiB2	LMU-2	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	High (151-200 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Ile	Trench cum bunding
Neregalla	79	3.94	MRDiB2	LMU-2	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	High (151-200 mm/m)	Very gently sloping (1-3%)	Moderate	Fallow land+Current fallow (Fl+Cf)	1 Borewell	Ile	Trench cum bunding
Neregalla	80	0.51	KVRiB2	LMU-1	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIIe	Graded bunding
Neregalla	81	0.4	KVRiB2	LMU-1	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIIe	Graded bunding
Neregalla	116	0	MTLmB2	LMU-4	Shallow (25-50 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIIes	Graded bunding
Neregalla	117	0.2	KVRmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	IIs	Graded bunding
Neregalla	118	0.42	KVRmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	IIs	Graded bunding
Neregalla	119	2.87	KVRmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	IIs	Graded bunding
Neregalla	120	2.86	KVRmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Maize+Fallow land (Mz+Fl)	Not Available	IIs	Graded bunding
Neregalla	121	4.62	MTLmB2	LMU-4	Shallow (25-50 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Maize+Redgram (Mz+Rg)	Not Available	IIIes	Graded bunding
Neregalla	122	3.42	MTLmB2	LMU-4	Shallow (25-50 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Maize+Redgram (Mz+Rg)	Not Available	IIIes	Graded bunding
Neregalla	123	0.03	MTLmB2	LMU-4	Shallow (25-50 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIIes	Graded bunding
Neregalla	125	2.52	KVRmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	IIs	Graded bunding
Neregalla	126	2.31	DRLmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Neregalla	127	0.89	DRLmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	Iles	Graded bunding
Neregalla	128	0.2	DRLmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	Iles	Graded bunding
Neregalla	164	0.03	RNKmB2	LMU-3	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Fallow land+Sunflower (Fl+Sf)	Not Available	Iles	Graded bunding
Neregalla	167	1.65	RNKmB2	LMU-3	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow (Cf)	Not Available	Iles	Graded bunding
Neregalla	168	5.22	RNKmB1	LMU-3	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Fallow land+Sunflower (Fl+Sf)	Not Available	IIs	Graded bunding

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Neregalla	169	5.52	KVRmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Current fallow+Sunflower (Cf+Sf)	Not Available	IIs	Graded bunding
Neregalla	170	4.04	RNKmB1	LMU-3	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Bengalgram+Sunflower (Bg+Sf)	Not Available	IIs	Graded bunding
Neregalla	171	6.71	RNKmB1	LMU-3	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Current fallow+Sunflower+Bengalgram (Cf+Sf+Bg)	Not Available	IIs	Graded bunding
Neregalla	172	7.16	RNKmB1	LMU-3	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Current fallow+Maize (Cf+Mz)	1 Borewell	IIs	Graded bunding
Neregalla	173	0.55	KVRmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Current fallow+Sunflower (Cf+Sf)	Not Available	IIs	Graded bunding
Neregalla	175	2.57	KVRmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Bengalgram+Current fallow (Bg+Cf)	Not Available	IIs	Graded bunding
Neregalla	176	1.53	KVRmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	IIs	Graded bunding
Neregalla	177	2.34	RNKmB1	LMU-3	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	IIs	Graded bunding
Neregalla	178	1.4	RNKmB1	LMU-3	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	IIs	Graded bunding
Neregalla	179	9.25	KVRmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Current fallow+Pearl millet+Bengalgram (Cf+Pm+Bg)	Not Available	IIs	Graded bunding
Neregalla	180	5.1	KVRmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Bengalgram (Bg)	Not Available	IIs	Graded bunding
Neregalla	181	6.38	KVRmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Bengalgram+Redgram (Bg+Rg)	Not Available	IIs	Graded bunding
Neregalla	182	3.6	KVRmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Maize+Redgram (Mz+Rg)	Not Available	IIs	Graded bunding
Neregalla	183	3.91	KVRmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Maize+Redgram (Mz+Rg)	Not Available	IIs	Graded bunding
Neregalla	184	6.41	BGPmB1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	High (151-200 mm/m)	Very gently sloping (1-3%)	Slight	Fallowland+Maize+Sunflower (Fl+Mz+Sf)	Not Available	IIs	Graded bunding
Neregalla	185	4.15	BGPmB1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	High (151-200 mm/m)	Very gently sloping (1-3%)	Slight	Sunflower (Sf)	Not Available	IIs	Graded bunding
Neregalla	186	4.46	KVRmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	IIs	Graded bunding
Neregalla	187	4.79	BGPmB1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	High (151-200 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	IIs	Graded bunding
Neregalla	188	6.22	DRLmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Fallowland+Maize+Redgram (Fl+Mz+Rg)	Not Available	IIs	Graded bunding
Neregalla	189	5.86	DRLmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Maize+Redgram (Mz+Rg)	Not Available	IIs	Graded bunding
Neregalla	190	7.9	DRLmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow (Cf)	Not Available	IIs	Graded bunding
Neregalla	191	0.88	DRLmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIs	Graded bunding
Neregalla	192	8.38	KVRmB1g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Fallow land+Sunflower (Fl+Sf)	Not Available	IIs	Graded bunding

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Neregalla	193	5.61	DRLmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Fallowland+Maize+Sunflower (Fl+Mz+Sf)	Not Available	Iles	Graded bunding
Neregalla	194	9.09	BGPmB1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	High (151-200 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	Ils	Graded bunding
Neregalla	195	6.49	DRLmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Sunflower (Sf)	Not Available	Iles	Graded bunding
Neregalla	196	8.02	RNKmB1	LMU-3	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	Ils	Graded bunding
Neregalla	197	0.75	RNKmB1	LMU-3	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	Ils	Graded bunding
Neregalla	198	2.39	KVRmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	Ils	Graded bunding
Neregalla	199	7.21	KVRmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	Ils	Graded bunding
Neregalla	200	5.34	KVRmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	Ils	Graded bunding
Neregalla	201	2.98	KVRmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	Ils	Graded bunding
Neregalla	202	3.12	KVRmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	Ils	Graded bunding
Neregalla	203	6.19	MRDiB2	LMU-2	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	High (151-200 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	1 Borewell	Ile	Trench cum bunding
Neregalla	204	2.06	MRDiB2	LMU-2	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	High (151-200 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	Ile	Trench cum bunding
Neregalla	205	4.11	MRDiB2	LMU-2	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	High (151-200 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	Ile	Trench cum bunding
Neregalla	206	0.54	MRDiB2	LMU-2	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	High (151-200 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	Ile	Trench cum bunding
Neregalla	207	7.87	MRDiB2	LMU-2	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	High (151-200 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	Ile	Trench cum bunding
Neregalla	208	7.25	KVRmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	Ils	Graded bunding
Neregalla	209	0.22	KVRmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	Ils	Graded bunding
Neregalla	210	5.84	GRHmB2g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Sunflower (Sf)	Not Available	Iles	Graded bunding
Neregalla	211	5.03	KVRmB1g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	Ils	Graded bunding
Neregalla	212	12.26	GRHmB2	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Maize+Pearl millet+Redgram (Mz+Pm+Rg)	Not Available	Iles	Graded bunding
Neregalla	213	4.07	GRHmB2	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow+Pearl millet+Sunflower (Cf+Pm+Sf)	Not Available	Iles	Graded bunding
Neregalla	214	5.59	GRHmB2	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow+Maize (Cf+Mz)	Not Available	Iles	Graded bunding
Neregalla	215	4.45	GRHmB2	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	Iles	Graded bunding

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Neregalla	216	10.05	KVRmB1g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Sunflower (Sf)	Not Available	IIs	Graded bunding
Neregalla	217	2.58	GRHmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	IIs	Graded bunding
Neregalla	218	7.23	GRHmB2g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIs	Graded bunding
Neregalla	219	5.8	GRHmB2g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Maize+Redgram (Mz+Rg)	Not Available	IIs	Graded bunding
Neregalla	220	5.68	MRDiB2	LMU-2	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	High (151-200 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	Ile	Trench cum bunding
Neregalla	221	5.5	MRDiB2	LMU-2	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	High (151-200 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	Ile	Trench cum bunding
Neregalla	222	7.11	GRHmB2g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	1 Borewell	IIs	Graded bunding
Neregalla	223	5.56	GRHmB2g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIs	Graded bunding
Neregalla	224	0.24	GRHmB2g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIs	Graded bunding
Neregalla	225	6.92	GRHmB2g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIs	Graded bunding

Village	Survey No	Soil Reaction	Salinity	Organic Carbon	Available Phosphorus	Available Potassium	Available Sulphur	Available Boron	Available Iron	Available Manganese	Available Copper	Available Zinc
Neregalla	218	Strongly alkaline (pH 8.4 - 9.0)	Non saline (<2 dsm)	Low (< 0.5 %)	Low (< 23 kg/ha)	High (> 337 kg/ha)	Low (<10 ppm)	Medium (0.5 - 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Neregalla	219	Strongly alkaline (pH 8.4 - 9.0)	Non saline (<2 dsm)	Low (< 0.5 %)	Medium (23 - 57 kg/ha)	High (> 337 kg/ha)	Low (<10 ppm)	Medium (0.5 - 1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Neregalla	220	Strongly alkaline (pH 8.4 - 9.0)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	High (> 337 kg/ha)	Low (<10 ppm)	Medium (0.5 - 1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Neregalla	221	Very strongly alkaline (pH > 9.0)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	High (> 337 kg/ha)	Low (<10 ppm)	Medium (0.5 - 1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Neregalla	222	Strongly alkaline (pH 8.4 - 9.0)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	High (> 337 kg/ha)	Low (<10 ppm)	Medium (0.5 - 1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Neregalla	223	Strongly alkaline (pH 8.4 - 9.0)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	High (> 337 kg/ha)	Low (<10 ppm)	Medium (0.5 - 1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Neregalla	224	Strongly alkaline (pH 8.4 - 9.0)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	High (> 337 kg/ha)	Low (<10 ppm)	Medium (0.5 - 1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Neregalla	225	Strongly alkaline (pH 8.4 - 9.0)	Non saline (<2 dsm)	Low (< 0.5 %)	Medium (23 - 57 kg/ha)	High (> 337 kg/ha)	Low (<10 ppm)	Medium (0.5 - 1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)

Appendix III
Neregallu-3 (1T1c) Microwatershed
Soil Suitability Information

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Jasmine	Bhendi	Brinjal	Crossandra	Drumstick	Mulberry	Onion	
Madhinura	71	S3t	S2t	S3t	S1	S3t	S1	S2rt	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S3t	S2t	S2t	S2t	S2t	S3t	S2tz	S2tz	S2t	S2t	S2t	S3tz	
Madhinura	72	S3t	S2t	S3t	S1	S3t	S1	S2rt	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S3t	S2t	S2t	S2t	S2t	S2t	S3t	S2tz	S2tz	S2t	S2t	S2t	S3tz
Madhinura	73	S3t	S2t	S3t	S1	S3t	S1	S2rt	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S3t	S2t	S2t	S2t	S2t	S2t	S3t	S2tz	S2tz	S2t	S2t	S2t	S3tz
Madhinura	74	S3t	S2t	S3t	S1	S3t	S1	S2rt	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S3t	S2t	S2t	S2t	S2t	S2t	S3t	S2tz	S2tz	S2t	S2t	S2t	S3tz
Madhinura	75	S3t	S2t	S3t	S1	S3t	S1	S2rt	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S3t	S2t	S2t	S2t	S2t	S2t	S3t	S2tz	S2tz	S2t	S2t	S2t	S3tz
Madhinura	76	S3t	S2t	S3t	S1	S3t	S1	S2rt	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S3t	S2t	S2t	S2t	S2t	S2t	S3t	S2tz	S2tz	S2t	S2t	S2t	S3tz
Madhinura	77	S3t	S2t	S3t	S1	S3t	S1	S2rt	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S3t	S2t	S2t	S2t	S2t	S2t	S3t	S2tz	S2tz	S2t	S2t	S2t	S3tz
Madhinura	78	S3rt	S2tz	S3rt	S1	S3t	S1	S3r	S2r	S1	S2r	S2t	S2t	S3t	S1	N1t	S3rt	S2r	S3t	S3t	S3t	S2t	S2t	S2rt	S2t	S3t	S2tz	S2tz	S3t	S2rt	S2t	S3tz	
Madhinura	79	S3rt	S2tz	S3rt	S1	S3t	S1	S3r	S2r	S1	S2r	S2t	S2t	S3t	S1	N1t	S3rt	S2r	S3t	S3t	S3t	S2t	S2t	S2rt	S2t	S3t	S2tz	S2tz	S3t	S2rt	S2t	S3tz	
Madhinura	80	S3rt	S2tz	S3rt	S1	S3t	S1	S3r	S2r	S1	S2r	S2t	S2t	S3t	S1	N1t	S3rt	S2r	S3t	S3t	S3tw	S2tw	S2tw	S2rt	S2t	S3tw	S2tz	S2tz	S3t	S2rt	S2tw	S3tz	
Madhinura	81	N1rz	S2tz	S3rz	S2rz	S3tz	S2rz	N1rz	S3rz	S2rz	S3rz	S3rz	S2rz	S3tz	S2rz	N1tz	S3tz	S3rz	S3tz	S3tz	S3tz	S2rz	S2rz	S3rz	S2tz	S2rz	S2rt	S2rt	S3rz	S3rz	S3rz	S3tz	
Madhinura	82	N1rz	S2tz	S3rz	S2rz	S3tz	S2rz	N1rz	S3rz	S2rz	S3rz	S3rz	S2rz	S3tz	S2rz	N1tz	S3tz	S3rz	S3tz	S3tz	S3tz	S2rz	S2rz	S3rz	S2tz	S2rz	S2rt	S2rt	S3rz	S3rz	S3rz	S3tz	
Madhinura	83	S3rt	S2tz	S3rt	S1	S3t	S1	S3r	S2r	S1	S2r	S2t	S2t	S3t	S1	N1t	S3rt	S2r	S3t	S3t	S3tw	S2tw	S2tw	S2rt	S2t	S3tw	S2tz	S2tz	S3t	S2rt	S2tw	S3tz	
Madhinura	84	S3rt	S2tz	S3rt	S1	S3t	S1	S3r	S2r	S1	S2r	S2t	S2t	S3t	S1	N1t	S3rt	S2r	S3t	S3t	S3tw	S2tw	S2tw	S2rt	S2t	S3tw	S2tz	S2tz	S3t	S2rt	S2tw	S3tz	
Madhinura	85	S3rz	S2tz	S3tz	S2nz	S3tz	S2rz	S3rz	S2rz	S2rz	S2rz	S3rz	S2tz	S3tz	S2z	N1tz	S3rz	S2rz	S3tz	S3tz	S3tz	S2tz	S2tz	S2rt	S2tz	S3tz	S2tz	S2tz	S2z	S2rz	S2tz	S3tz	
Madhinura	86	S3rt	S2tz	S3rt	S1	S3t	S1	S3r	S2r	S1	S2r	S2t	S2t	S3t	S1	N1t	S3rt	S2r	S3t	S3t	S3tw	S2tw	S2tw	S2rt	S2t	S3tw	S2tz	S2tz	S3t	S2rt	S2tw	S3tz	
Madhinura	87	S3rt	S2tz	S3rt	S1	S3t	S1	S3r	S2r	S1	S2r	S2t	S2t	S3t	S1	N1t	S3rt	S2r	S3t	S3t	S3tw	S2tw	S2tw	S2rt	S2t	S3tw	S2tz	S2tz	S3t	S2rt	S2tw	S3tz	
Madhinura	109	S3rz	S2tz	S3tz	S2nz	S3tz	S2rz	S3rz	S2rz	S2rz	S2rz	S3rz	S2tz	S3tz	S2z	N1tz	S3rz	S2rz	S3tz	S3tz	S3tz	S2tz	S2tz	S2rt	S2tz	S3tz	S2tz	S2tz	S2z	S2rz	S2tz	S3tz	
Madhinura	111	S3t	S2t	S3t	S1	S3t	S1	S2rt	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S3t	S2t	S2t	S2t	S2t	S2t	S3t	S2tz	S2tz	S2t	S2t	S2t	S3tz
Madhinura	113	S3rz	S2tz	S3tz	S2nz	S3tz	S2rz	S3rz	S2rz	S2rz	S2rz	S3rz	S2tz	S3tz	S2z	N1tz	S3rz	S2rz	S3tz	S3tz	S3tz	S2tz	S2tz	S2rt	S2tz	S3tz	S2tz	S2tz	S2z	S2rz	S2tz	S3tz	
Madhinura	114	N1rt	S3tz	N1rz	S3rz	N1rt	S3rz	N1rz	N1rz	S3rz	N1rz	N1rz	S3tz	N1rt	S3zg	N1rt	N1rt	N1rz	S3tz	S3rz	S3rz	S3rz	S3rz	N1rz	S3rz	S3rz	S3r	S3r	S3rz	N1rz	N1rz	S3rt	
Madhinura	115	N1rt	S3tz	N1rz	S3rz	N1rt	S3rz	N1rz	N1rz	S3rz	N1rz	N1rz	S3tz	N1rt	S3zg	N1rt	N1rt	N1rz	S3tz	S3rz	S3rz	S3rz	S3rz	N1rz	S3rz	S3rz	S3r	S3r	S3rz	N1rz	N1rz	S3rt	
Madhinura	116	N1rt	S3tz	N1rz	S3rz	N1rt	S3rz	N1rz	N1rz	S3rz	N1rz	N1rz	S3tz	N1rt	S3zg	N1rt	N1rt	N1rz	S3tz	S3rz	S3rz	S3rz	S3rz	N1rz	S3rz	S3rz	S3r	S3r	S3rz	N1rz	N1rz	S3rt	
Madhinura	117	S3t	S2t	S3t	S1	S3t	S1	S2rt	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S3t	S2t	S2t	S2t	S2t	S2t	S3t	S2tz	S2tz	S2t	S2t	S2t	S3tz

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Jasmine	Bhendi	Brinjal	Crossandra	Drumstick	Mulberry	Onion	
Madhinura	118	S3t	S2t	S3t	S1	S3t	S1	S2rt	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S3t	S2t	S2t	S2t	S2t	S3t	S2tz	S2tz	S2t	S2t	S2t	S3tz	
Madhinura	119	S3t	S2t	S3t	S1	S3t	S1	S2rt	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S3t	S2t	S2t	S2t	S2t	S3t	S2tz	S2tz	S2t	S2t	S2t	S3tz	
Madhinura	120	S3t	S2t	S3t	S1	S3t	S1	S2rt	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S3t	S2t	S2t	S2t	S2t	S3t	S2tz	S2tz	S2t	S2t	S2t	S3tz	
Madhinura	121	N1rz	S2tz	S3rz	S2rz	S3tz	S2rz	N1rz	S3rz	S2rz	S3rz	S3rz	S2rz	S3tz	S2rz	N1tz	S3tz	S3rz	S3tz	S3tz	S3tz	S2rz	S2rz	S3rz	S2tz	S2rz	S2rt	S2rt	S3rz	S3rz	S3rz	S3tz	
Madhinura	122	S3t	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2t	S1	S3t	S3t	S3t	S2t	S2t	S2t	S2t	S3t	S2t	S2t	S2t	S2t	S3t	S3t	
Madhinura	123	S3t	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2t	S1	S3t	S3t	S3t	S2t	S2t	S2t	S2t	S3t	S2t	S2t	S2t	S2t	S3t	S3t	
Madhinura	124	S3t	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2t	S1	S3t	S3t	S3t	S2t	S2t	S2t	S2t	S3t	S2t	S2t	S2t	S2t	S3t	S3t	
Madhinura	125	S3t	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2t	S1	S3t	S3t	S3t	S2t	S2t	S2t	S2t	S3t	S2t	S2t	S2t	S2t	S3t	S3t	
Madhinura	128	S3t	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2t	S1	S3t	S3t	S3t	S2t	S2t	S2t	S2t	S3t	S2t	S2t	S2t	S2t	S3t	S3t	
Madhinura	130	S3t	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2t	S1	S3t	S3t	S3t	S2t	S2t	S2t	S2t	S3t	S2t	S2t	S2t	S2t	S3t	S3t	
Madhinura	131	S3t	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2t	S1	S3t	S3t	S3t	S2t	S2t	S2t	S2t	S3t	S2t	S2t	S2t	S2t	S3t	S3t	
Madhinura	132	S3t	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2t	S1	S3t	S3t	S3t	S2t	S2t	S2t	S2t	S3t	S2t	S2t	S2t	S2t	S3t	S3t	
Madhinura	133	S3t	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2t	S1	S3t	S3t	S3t	S2t	S2t	S2t	S2t	S3t	S2t	S2t	S2t	S2t	S3t	S3t	
Madhinura	134	N1rt	S3tz	N1rz	S3rz	N1rt	S3rz	N1rz	N1rz	S3rz	N1rz	N1rz	S3tz	N1rt	S3zg	N1rt	N1rt	N1rz	S3tz	S3rz	S3rz	S3rz	S3rz	S3rz	N1rz	S3rz	S3rz	S3r	S3r	S3rz	N1rz	N1rz	S3rt
Madhinura	135	S3t	S2t	S3t	S1	S3t	S1	S2rt	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S3t	S2t	S2t	S2t	S2t	S3t	S2tz	S2tz	S2t	S2t	S2t	S3tz	
Madhinura	136	S3t	S2t	S3t	S1	S3t	S1	S2rt	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S3t	S2t	S2t	S2t	S2t	S3t	S2tz	S2tz	S2t	S2t	S2t	S3tz	
Madhinura	138	S3t	S2t	S3t	S1	S3t	S1	S2rt	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S3t	S2t	S2t	S2t	S2t	S3t	S2tz	S2tz	S2t	S2t	S2t	S3tz	
Madhinura	139	S3t	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2t	S1	S3t	S3t	S3t	S2t	S2t	S2t	S2t	S3t	S2t	S2t	S2t	S2t	S3t	S3t	
Madhinura	140	S3t	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2t	S1	S3t	S3t	S3t	S2t	S2t	S2t	S2t	S3t	S2t	S2t	S2t	S2t	S3t	S3t	
Madhinura	141	S3t	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2t	S1	S3t	S3t	S3t	S2t	S2t	S2t	S2t	S3t	S2t	S2t	S2t	S2t	S3t	S3t	
Madhinura	142	S3t	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2t	S1	S3t	S3t	S3t	S2t	S2t	S2t	S2t	S3t	S2t	S2t	S2t	S2t	S3t	S3t	
Madhinura	143	S3t	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2t	S1	S3t	S3t	S3t	S2t	S2t	S2t	S2t	S3t	S2t	S2t	S2t	S2t	S3t	S3t	
Madhinura	144	S3t	S2t	S3t	S1	S3t	S1	S2rt	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S3t	S2t	S2t	S2t	S2t	S3t	S2tz	S2tz	S2t	S2t	S2t	S3tz	
Madhinura	146	S3t	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2t	S1	S3t	S3t	S3t	S2t	S2t	S2t	S2t	S3t	S2t	S2t	S2t	S2t	S3t	S3t	
Madhinura	148	S3t	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2t	S1	S3t	S3t	S3t	S2t	S2t	S2t	S2t	S3t	S2t	S2t	S2t	S2t	S3t	S3t	
Madhinura	149	S3t	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2t	S1	S3t	S3t	S3t	S2t	S2t	S2t	S2t	S3t	S2t	S2t	S2t	S2t	S3t	S3t	
Neregalla	77	S1	S2t	S1	S1	S1	S1	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S2t	S2t	S1	S1	S1	S2t

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Jasmine	Bhendi	Brinjal	Crossandra	Drumstick	Mulberry	Onion	
Neregalla	78	S1	S2t	S1	S1	S1	S1	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S2t
Neregalla	79	S1	S1	S1	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1
Neregalla	80	S2rz	S2tz	S3tz	S2z	S3tz	S2zg	S2rz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2rz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz
Neregalla	81	S2rz	S2tz	S3tz	S2z	S3tz	S2zg	S2rz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2rz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz
Neregalla	116	N1rt	S3tz	N1rz	S3rz	N1rt	S3rz	N1rz	N1rz	S3rz	N1rz	N1rz	S3tz	N1rt	S3zg	N1rt	N1rt	N1rz	S3tz	S3rz	S3rz	S3rz	S3rz	N1rz	S3rz	S3rz	S3r	S3r	S3rz	N1rz	N1rz	S3rt	
Neregalla	117	S2rz	S2tz	S3tz	S2z	S3tz	S2z	S2rz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2rz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz
Neregalla	118	S2rz	S2tz	S3tz	S2z	S3tz	S2z	S2rz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2rz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz
Neregalla	119	S2rz	S2tz	S3tz	S2z	S3tz	S2z	S2rz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2rz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz
Neregalla	120	S2rz	S2tz	S3tz	S2z	S3tz	S2z	S2rz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2rz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz
Neregalla	121	N1rt	S3tz	N1rz	S3rz	N1rt	S3rz	N1rz	N1rz	S3rz	N1rz	N1rz	S3tz	N1rt	S3zg	N1rt	N1rt	N1rz	S3tz	S3rz	S3rz	S3rz	S3rz	N1rz	S3rz	S3rz	S3r	S3r	S3rz	N1rz	N1rz	S3rt	
Neregalla	122	N1rt	S3tz	N1rz	S3rz	N1rt	S3rz	N1rz	N1rz	S3rz	N1rz	N1rz	S3tz	N1rt	S3zg	N1rt	N1rt	N1rz	S3tz	S3rz	S3rz	S3rz	S3rz	N1rz	S3rz	S3rz	S3r	S3r	S3rz	N1rz	N1rz	S3rt	
Neregalla	123	N1rt	S3tz	N1rz	S3rz	N1rt	S3rz	N1rz	N1rz	S3rz	N1rz	N1rz	S3tz	N1rt	S3zg	N1rt	N1rt	N1rz	S3tz	S3rz	S3rz	S3rz	S3rz	N1rz	S3rz	S3rz	S3r	S3r	S3rz	N1rz	N1rz	S3rt	
Neregalla	125	S2rz	S2tz	S3tz	S2z	S3tz	S2z	S2rz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2rz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz
Neregalla	126	S3rz	S2tz	S3tz	S2nz	S3tz	S2rz	S3rz	S2rz	S2rz	S2rz	S3rz	S2tz	S3tz	S2z	N1tz	S3rz	S2rz	S3tz	S3tz	S3tz	S2tz	S2tz	S2rt	S2tz	S3tz	S2tz	S2tz	S2z	S2rz	S2tz	S3tz	
Neregalla	127	S3rz	S2tz	S3tz	S2nz	S3tz	S2rz	S3rz	S2rz	S2rz	S2rz	S3rz	S2tz	S3tz	S2z	N1tz	S3rz	S2rz	S3tz	S3tz	S3tz	S2tz	S2tz	S2rt	S2tz	S3tz	S2tz	S2tz	S2z	S2rz	S2tz	S3tz	
Neregalla	128	S3rz	S2tz	S3tz	S2nz	S3tz	S2rz	S3rz	S2rz	S2rz	S2rz	S3rz	S2tz	S3tz	S2z	N1tz	S3rz	S2rz	S3tz	S3tz	S3tz	S2tz	S2tz	S2rt	S2tz	S3tz	S2tz	S2tz	S2z	S2rz	S2tz	S3tz	
Neregalla	164	N1rz	S2tz	S3rz	S2rz	S3tz	S2rz	N1rz	S3rz	S2rz	S3rz	S3rz	S2rz	S3tz	S2rz	N1tz	S3tz	S3rz	S3tz	S3tz	S3tz	S2rz	S2rz	S3rz	S2tz	S2rz	S2rt	S2rt	S3rz	S3rz	S3rz	S3tz	
Neregalla	167	N1rz	S2tz	S3rz	S2rz	S3tz	S2rz	N1rz	S3rz	S2rz	S3rz	S3rz	S2rz	S3tz	S2rz	N1tz	S3tz	S3rz	S3tz	S3tz	S3tz	S2rz	S2rz	S3rz	S2tz	S2rz	S2rt	S2rt	S3rz	S3rz	S3rz	S3tz	
Neregalla	168	N1rz	S2tz	S3rz	S2rz	S3tz	S2rz	N1rz	S3rz	S2rz	S3rz	S3rz	S2rz	S3tz	S2rz	N1tz	S3tz	S3rz	S3tz	S3tz	S3tz	S2rz	S2rz	S3rz	S2tz	S2rz	S2rt	S2rt	S3rz	S3rz	S3rz	S3tz	
Neregalla	169	S2rz	S2tz	S3tz	S2z	S3tz	S2z	S2rz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2rz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz
Neregalla	170	N1rz	S2tz	S3rz	S2rz	S3tz	S2rz	N1rz	S3rz	S2rz	S3rz	S3rz	S2rz	S3tz	S2rz	N1tz	S3tz	S3rz	S3tz	S3tz	S3tz	S2rz	S2rz	S3rz	S2tz	S2rz	S2rt	S2rt	S3rz	S3rz	S3rz	S3tz	
Neregalla	171	N1rz	S2tz	S3rz	S2rz	S3tz	S2rz	N1rz	S3rz	S2rz	S3rz	S3rz	S2rz	S3tz	S2rz	N1tz	S3tz	S3rz	S3tz	S3tz	S3tz	S2rz	S2rz	S3rz	S2tz	S2rz	S2rt	S2rt	S3rz	S3rz	S3rz	S3tz	
Neregalla	172	N1rz	S2tz	S3rz	S2rz	S3tz	S2rz	N1rz	S3rz	S2rz	S3rz	S3rz	S2rz	S3tz	S2rz	N1tz	S3tz	S3rz	S3tz	S3tz	S3tz	S2rz	S2rz	S3rz	S2tz	S2rz	S2rt	S2rt	S3rz	S3rz	S3rz	S3tz	
Neregalla	173	S2rz	S2tz	S3tz	S2z	S3tz	S2z	S2rz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2rz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz
Neregalla	175	S2rz	S2tz	S3tz	S2z	S3tz	S2z	S2rz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2rz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz
Neregalla	176	S2rz	S2tz	S3tz	S2z	S3tz	S2z	S2rz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2rz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz
Neregalla	177	N1rz	S2tz	S3rz	S2rz	S3tz	S2rz	N1rz	S3rz	S2rz	S3rz	S3rz	S2rz	S3tz	S2rz	N1tz	S3tz	S3rz	S3tz	S3tz	S3tz	S2rz	S2rz	S3rz	S2tz	S2rz	S2rt	S2rt	S3rz	S3rz	S3rz	S3tz	

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Jasmine	Bhendi	Brinjal	Crossandra	Drumstick	Mulberry	Onion
Neregalla	178	N1rz	S2tz	S3rz	S2rz	S3tz	S2rz	N1rz	S3rz	S2rz	S3rz	S3rz	S2rz	S3tz	S2rz	N1tz	S3tz	S3rz	S3tz	S3tz	S3tz	S2rz	S2rz	S3rz	S2tz	S2rz	S2rt	S2rt	S3rz	S3rz	S3rz	S3tz
Neregalla	179	S2rz	S2tz	S3tz	S2z	S3tz	S2z	S2rz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2rz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz
Neregalla	180	S2rz	S2tz	S3tz	S2z	S3tz	S2z	S2rz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2rz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz
Neregalla	181	S2rz	S2tz	S3tz	S2z	S3tz	S2z	S2rz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2rz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz
Neregalla	182	S2rz	S2tz	S3tz	S2z	S3tz	S2z	S2rz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2rz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz
Neregalla	183	S2rz	S2tz	S3tz	S2z	S3tz	S2z	S2rz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2rz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz
Neregalla	184	S3tz	S2tz	S3tz	S2z	S3tz	S2z	S2tz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2tz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz
Neregalla	185	S3tz	S2tz	S3tz	S2z	S3tz	S2z	S2tz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2tz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz
Neregalla	186	S2rz	S2tz	S3tz	S2z	S3tz	S2z	S2rz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2rz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz
Neregalla	187	S3tz	S2tz	S3tz	S2z	S3tz	S2z	S2tz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2tz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz
Neregalla	188	S3rz	S2tz	S3tz	S2nz	S3tz	S2rz	S3rz	S2rz	S2rz	S2rz	S3rz	S2tz	S3tz	S2z	N1tz	S3rz	S2rz	S3tz	S3tz	S3tz	S2tz	S2tz	S2rt	S2tz	S3tz	S2tz	S2tz	S2z	S2rz	S2tz	S3tz
Neregalla	189	S3rz	S2tz	S3tz	S2nz	S3tz	S2rz	S3rz	S2rz	S2rz	S2rz	S3rz	S2tz	S3tz	S2z	N1tz	S3rz	S2rz	S3tz	S3tz	S3tz	S2tz	S2tz	S2rt	S2tz	S3tz	S2tz	S2tz	S2z	S2rz	S2tz	S3tz
Neregalla	190	S3rz	S2tz	S3tz	S2nz	S3tz	S2rz	S3rz	S2rz	S2rz	S2rz	S3rz	S2tz	S3tz	S2z	N1tz	S3rz	S2rz	S3tz	S3tz	S3tz	S2tz	S2tz	S2rt	S2tz	S3tz	S2tz	S2tz	S2z	S2rz	S2tz	S3tz
Neregalla	191	S3rz	S2tz	S3tz	S2nz	S3tz	S2rz	S3rz	S2rz	S2rz	S2rz	S3rz	S2tz	S3tz	S2z	N1tz	S3rz	S2rz	S3tz	S3tz	S3tz	S2tz	S2tz	S2rt	S2tz	S3tz	S2tz	S2tz	S2z	S2rz	S2tz	S3tz
Neregalla	192	S2rz	S2tz	S3tz	S2zg	S3tz	S2zg	S2rz	S2z	S2gz	S2zg	S2tz	S2tz	S3tz	S2z	N1tz	S2rz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz
Neregalla	193	S3rz	S2tz	S3tz	S2nz	S3tz	S2rz	S3rz	S2rz	S2rz	S2rz	S3rz	S2tz	S3tz	S2z	N1tz	S3rz	S2rz	S3tz	S3tz	S3tz	S2tz	S2tz	S2rt	S2tz	S3tz	S2tz	S2tz	S2z	S2rz	S2tz	S3tz
Neregalla	194	S3tz	S2tz	S3tz	S2z	S3tz	S2z	S2tz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2tz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz
Neregalla	195	S3rz	S2tz	S3tz	S2nz	S3tz	S2rz	S3rz	S2rz	S2rz	S2rz	S3rz	S2tz	S3tz	S2z	N1tz	S3rz	S2rz	S3tz	S3tz	S3tz	S2tz	S2tz	S2rt	S2tz	S3tz	S2tz	S2tz	S2z	S2rz	S2tz	S3tz
Neregalla	196	N1rz	S2tz	S3rz	S2rz	S3tz	S2rz	N1rz	S3rz	S2rz	S3rz	S3rz	S2rz	S3tz	S2rz	N1tz	S3tz	S3rz	S3tz	S3tz	S3tz	S2rz	S2rz	S3rz	S2tz	S2rz	S2rt	S2rt	S3rz	S3rz	S3rz	S3tz
Neregalla	197	N1rz	S2tz	S3rz	S2rz	S3tz	S2rz	N1rz	S3rz	S2rz	S3rz	S3rz	S2rz	S3tz	S2rz	N1tz	S3tz	S3rz	S3tz	S3tz	S3tz	S2rz	S2rz	S3rz	S2tz	S2rz	S2rt	S2rt	S3rz	S3rz	S3rz	S3tz
Neregalla	198	S2rz	S2tz	S3tz	S2z	S3tz	S2z	S2rz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2rz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz
Neregalla	199	S2rz	S2tz	S3tz	S2z	S3tz	S2z	S2rz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2rz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz
Neregalla	200	S2rz	S2tz	S3tz	S2z	S3tz	S2z	S2rz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2rz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz
Neregalla	201	S2rz	S2tz	S3tz	S2z	S3tz	S2z	S2rz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2rz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz
Neregalla	202	S2rz	S2tz	S3tz	S2z	S3tz	S2z	S2rz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2rz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz
Neregalla	203	S1	S1	S1	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1
Neregalla	204	S1	S1	S1	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Jasmine	Bhendi	Brinjal	Crossandra	Drumstick	Mulberry	Onion	
Neregalla	205	S1	S1	S1	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	
Neregalla	206	S1	S1	S1	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	
Neregalla	207	S1	S1	S1	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	
Neregalla	208	S2rz	S2tz	S3tz	S2z	S3tz	S2z	S2rz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2rz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz
Neregalla	209	S2rz	S2tz	S3tz	S2z	S3tz	S2z	S2rz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2rz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz
Neregalla	210	S3t	S2t	S3t	S2g	S3t	S2g	S2rg	S2g	S2g	S2g	S2gt	S2gt	S3t	S2g	N1t	S2rt	S2g	S3t	S3t	S3t	S2tg	S2tg	S2tg	S2t	S3t	S2t	S2t	S3t	S2tg	S2gt	S3t	
Neregalla	211	S2rz	S2tz	S3tz	S2zg	S3tz	S2zg	S2rz	S2z	S2gz	S2zg	S2tz	S2tz	S3tz	S2z	N1tz	S2rz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz
Neregalla	212	S3t	S2t	S3t	S1	S3t	S1	S2r	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S3t	S2t	S2t	S2t	S2t	S2t	S3t	S2t	S2t	S3t	S2t	S2t	S3t
Neregalla	213	S3t	S2t	S3t	S1	S3t	S1	S2r	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S3t	S2t	S2t	S2t	S2t	S2t	S3t	S2t	S2t	S3t	S2t	S2t	S3t
Neregalla	214	S3t	S2t	S3t	S1	S3t	S1	S2r	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S3t	S2t	S2t	S2t	S2t	S2t	S3t	S2t	S2t	S3t	S2t	S2t	S3t
Neregalla	215	S3t	S2t	S3t	S1	S3t	S1	S2r	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S3t	S2t	S2t	S2t	S2t	S2t	S3t	S2t	S2t	S3t	S2t	S2t	S3t
Neregalla	216	S2rz	S2tz	S3tz	S2zg	S3tz	S2zg	S2rz	S2z	S2gz	S2zg	S2tz	S2tz	S3tz	S2z	N1tz	S2rz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz
Neregalla	217	S3t	S2t	S3t	S1	S3t	S1	S2r	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S3t	S2t	S2t	S2t	S2t	S2t	S3t	S2t	S2t	S3t	S2t	S2t	S3t
Neregalla	218	S3t	S2t	S3t	S2g	S3t	S2g	S2rg	S2g	S2g	S2g	S2gt	S2gt	S3t	S2g	N1t	S2rt	S2g	S3t	S3t	S3t	S2tg	S2tg	S2tg	S2t	S3t	S2t	S2t	S3t	S2tg	S2gt	S3t	
Neregalla	219	S3t	S2t	S3t	S2g	S3t	S2g	S2rg	S2g	S2g	S2g	S2gt	S2gt	S3t	S2g	N1t	S2rt	S2g	S3t	S3t	S3t	S2tg	S2tg	S2tg	S2t	S3t	S2t	S2t	S3t	S2tg	S2gt	S3t	
Neregalla	220	S1	S1	S1	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1
Neregalla	221	S1	S1	S1	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1
Neregalla	222	S3t	S2t	S3t	S2g	S3t	S2g	S2rg	S2g	S2g	S2g	S2gt	S2gt	S3t	S2g	N1t	S2rt	S2g	S3t	S3t	S3t	S2tg	S2tg	S2tg	S2t	S3t	S2t	S2t	S3t	S2tg	S2gt	S3t	
Neregalla	223	S3t	S2t	S3t	S2g	S3t	S2g	S2rg	S2g	S2g	S2g	S2gt	S2gt	S3t	S2g	N1t	S2rt	S2g	S3t	S3t	S3t	S2tg	S2tg	S2tg	S2t	S3t	S2t	S2t	S3t	S2tg	S2gt	S3t	
Neregalla	224	S3t	S2t	S3t	S2g	S3t	S2g	S2rg	S2g	S2g	S2g	S2gt	S2gt	S3t	S2g	N1t	S2rt	S2g	S3t	S3t	S3t	S2tg	S2tg	S2tg	S2t	S3t	S2t	S2t	S3t	S2tg	S2gt	S3t	
Neregalla	225	S3t	S2t	S3t	S2g	S3t	S2g	S2rg	S2g	S2g	S2g	S2gt	S2gt	S3t	S2g	N1t	S2rt	S2g	S3t	S3t	S3t	S2tg	S2tg	S2tg	S2t	S3t	S2t	S2t	S3t	S2tg	S2gt	S3t	

PART-B

SOCIO-ECONOMIC STATUS OF FARM HOUSEHOLDS

CONTENTS

1.	Salient findings of the survey	1-5
2.	Introduction	7
3	Methodology	9
4	Salient features of the survey	11-31
5	Summary	33-37

LIST OF TABLES

1	Households sampled for socio economic survey	11
2	Population characteristics	11
3	Age wise classification of household members	11
4	Education level of household members	12
5	Occupation of household heads	12
6	Occupation of family members	12
7	Institutional participation of household members	13
8	Type of house owned by households	13
9	Durable assets owned by households	13
10	Average value of durable assets owned by households	13
11	Farm implements owned by households	14
12	Average value of farm implements	14
13	Livestock possession by households	14
14	Average labour availability	15
15	Adequacy of hired labour	15
16	Distribution of land (ha)	15
17	Average land value (Rs./ha)	16
18	Status of bore wells	16
19	Status of open wells	16
20	Source of irrigation	16
21	Depth of water	16
22	Irrigated area (ha)	17
23	Cropping pattern	17
24	Cropping intensity	17
25	Possession of Bank account and savings	17
26	Borrowing status	18
27	Source of credit availed	18
28	Average Credit amount	18
29	Purpose of credit borrowed – Institutional Credit	18
30	Purpose of credit borrowed – Private Credit	19
31	Repayment status of households- Institutional credit	19

36	Repayment status of households- Private credit	19
37	Opinion on institutional sources of credit	19
38	Opinion on non- institutional sources of credit	19
39	Cost of cultivation of Maize	20
40	Cost of cultivation of Chilly	21
41	Cost of cultivation of Sorghum	22
42	Cost of cultivation of Bengal gram	23
43	Cost of cultivation of Sunflower	24
44	Cost of cultivation of Paddy	25
45	Adequacy of fodder	26
46	Annual gross income	26
47	Average annual expenditure	26
48	Horticulture species grown	27
49	Forest species grown	27
50	Marketing of the agricultural produce	27
51	Marketing channels used for sale of agricultural produce	28
52	Mode of transport of agricultural produce	28
53	Incidence of soil and water erosion problems	28
54	Interest towards soil testing	28
55	Usage pattern of fuel for domestic use	28
56	Source of drinking water	29
57	Source of light	29
58	Existence of sanitary toilet facility	29
59	Possession of public distribution system(PDS) card	29
60	Participation in NREGA programme	29
61	Adequacy of food items	30
62	Response on inadequacy of food items	30
63	Response on market surplus of food items	31
64	Farming constraints experienced	31

SALIENT FINDINGS OF THE SURVEY

- ❖ *The data indicated that there were 94 (56.29%) men and 73 (43.71%) women among the sampled households.*
- ❖ *The average family size of landless farmers' was 3, marginal farmers' was 4.2, small farmers' was 3.83, semi medium farmers' was 5 and medium farmers' was 4.*
- ❖ *The data indicated that, 18 (10.78%) people were in 0-15 years of age, 69 (41.32%) were in 16-35 years of age, 64 (38.32%) were in 36-60 years of age and 16 (9.58%) were above 61 years of age.*
- ❖ *The results indicated that Naregallu-3 had 36.53 per cent illiterates, 35.33 per cent of them had primary school education, 4.19 per cent of them had middle school education, 13.77 per cent of them had high school education, 7.19 per cent of them had PUC education and 1.20 per cent had diploma and ITI.*
- ❖ *The results indicate that, 97.5 per cent of household heads were practicing agriculture and 2.5 per cent of the household heads were agricultural labourers.*
- ❖ *The results indicate that agriculture was the major occupation for 23.95 per cent of the household members, 61.68 per cent were agricultural labourers and 14.37 per cent were student.*
- ❖ *The results show that, 100 per cent of the population in the micro watershed has not participated in any local institutions.*
- ❖ *The results indicate that 5 per cent of the households possess thatched house and 95 per cent of the households possess katcha houses.*
- ❖ *The results show that 70 per cent of the households possess TV, 17.5 per cent of them possess mixer/grinder, 7.50 per cent of them possess bicycle, 35 per cent of the households possess motor cycle, 2.50 per cent of them possess auto and landline phone and 90 per cent of the households possess mobile phones.*
- ❖ *The results show that the average value of television was Rs. 6,732, mixer grinder was Rs. 1,700, bicycle was 1,175, motor cycle was Rs. 32,357, auto was 300,000, landline was 4,000 and mobile phone was Rs. 2,492.*
- ❖ *About 12.50 per cent of the households possess bullock cart, 17.50 per cent of them possess plough, 5 per cent of them possess seed/ fertilizer drill, 2.5 per cent of them possess sprayer, 12.50 per cent of them possess weeder, and 17.50 per cent of them possess chaff cutter. T*
- ❖ *he results show that the average value of bullock cart was Rs. 17,060, plough was Rs. 2800, seed/fertilizer drill was Rs. 8,000, sprayer was Rs. 2,000, weeder was Rs. 21 and the average value of chaff cutter was Rs. 441.*
- ❖ *The results indicate that, 22.50 per cent of the households possess bullocks, 7.50 per cent of the households possess local cow, 2.50 per cent possess buffalo and goat.*

- ❖ *The results indicate that, average own labour men available in the micro watershed was 1.40, average own labour (women) available was 1.85, average hired labour (men) available was 6.25 and average hired labour (women) available was 6.4.*
- ❖ *The results indicate that 100 per cent of the households opined that the hired labour was inadequate.*
- ❖ *The results indicate that, households of the Naregallu-3 micro-watershed possess 29.15 ha (59.87%) of dry land and 19.54 ha (40.13%) of irrigated land. Marginal farmers possess 5.84 ha (86.61%) of dry land and 0.90 ha (13.39%) of irrigated land. Small farmers possess 14.26 ha (84.63%) of dry land and 2.59 ha (15.37%) of irrigated land. Semi medium farmers possess 9.05 ha (40.66%) of dry land and 13.21 ha (59.34%) of irrigated land. Medium farmers possess 2.83 ha (100%) of irrigated land.*
- ❖ *The results indicate that, the average value of dry land was Rs. 264,006.11 and the average value of irrigated land was Rs. 450,207.13. In case of marginal famers, the average land value was Rs. 530,630.63 for dry land and Rs. 1,218,385.64 for irrigated land. In case of small famers, the average land value was Rs. 238,308.74 for dry land and Rs. 694,687.49 for irrigated land. In case of semi medium famers, the average land value was Rs. 132,498.88 for dry land and Rs. 400,949.47 for irrigated land. In case of medium farmers, the average land value was Rs. 211,714.29 for irrigated land.*
- ❖ *The results indicate that, there were 11 functioning and 6 de-functioning bore wells in the micro watershed.*
- ❖ *The results indicate that, there was 1 functioning open well in the micro watershed.*
- ❖ *The results indicate that, bore well was the major irrigation source in the micro water shed for 27.50 per cent of the farmers and open well was the major source of irrigation for 2.50 per cent of the farmers.*
- ❖ *The results indicate that, the depth of bore well was found to be 13.72 meters and the depth of open well was found to be 3.05 meters.*
- ❖ *The results indicate that marginal, small and semi medium farmers had an irrigated area of 1.78 ha, 2.55 ha and 11.92 ha respectively.*
- ❖ *The results indicate that, farmers have grown maize (31.79 ha), sunflower (4.91 ha), sorghum (4.61 ha), Bengal gram (2.14 ha), paddy (1.62 ha), chilly and jowar (1.21 ha). Marginal farmers have grown maize and Bengal gram, while small farmers have grown maize, sunflower, sorghum, Bengal gram paddy and jowar. Semi medium farmers have grown maize, sunflower, sorghum, paddy and chilly. Medium farmers have grown maize.*
- ❖ *The results indicate that, the cropping intensity in Naregallu-3 micro-watershed was found to be 88.20 per cent.*
- ❖ *The results indicate that, 55 per cent of the households have bank account and savings.*

- ❖ *The results indicate that, 55 per cent of the households have availed credit from different sources.*
- ❖ *The results indicate that, 81.82 per cent of the households have borrowed from commercial bank, 4.55 per cent of them borrowed from friends/relatives, grameena bank and money lender, 9.09 per cent of the households borrowed from SHGs and CBOs. The results indicate that, the average credit amount borrowed by households in micro-watershed was Rs, 59,409.09.*
- ❖ *The results indicate that, 100 per cent of the households borrowed from institutional sources for the purpose of agricultural production.*
- ❖ *The results indicate that, 100 per cent of the household's availed credit for the purpose of agricultural production.*
- ❖ *The results indicated that 100 per cent of the households did not repay their loan borrowed from institutional sources.*
- ❖ *The results indicated that 100 per cent of the households did not repay their loan borrowed from private sources.*
- ❖ *The results indicate that, around 100 per cent opined that the loan amount borrowed from institutional sources helped to perform timely agricultural operations.*
- ❖ *The results indicate that, 100 per cent of the households opined that the credit borrowed from private credit helped to perform timely agricultural operations.*
- ❖ *The results indicate that, the total cost of cultivation for maize was Rs. 33037.55. The gross income realized by the farmers was Rs. 45231.00. The net income from maize cultivation was Rs. 12193.46. Thus the benefit cost ratio was found to be 1:1.37.*
- ❖ *The results indicate that, the total cost of cultivation for chilly was Rs. 57632.23. The gross income realized by the farmers was Rs. 89969.75. The net income from chilly cultivation was Rs. 32337.52. Thus the benefit cost ratio was found to be 1:1.56.*
- ❖ *The results indicate that, the total cost of cultivation for chilly was Rs. 57632.23. The gross income realized by the farmers was Rs. 89969.75. The net income from chilly cultivation was Rs. 32337.52. Thus the benefit cost ratio was found to be 1:1.56.*
- ❖ *The results indicate that, the total cost of cultivation for chilly was Rs. 57632.23. The gross income realized by the farmers was Rs. 89969.75. The net income from chilly cultivation was Rs. 32337.52. Thus the benefit cost ratio was found to be 1:1.56.*
- ❖ *The results indicate that, the total cost of cultivation for sunflower was Rs. 28263.78. The gross income realized by the farmers was Rs. 88272.38. The net income from sunflower cultivation was Rs. 60008.60. Thus the benefit cost ratio was found to be 1:3.12.*

- ❖ *The results indicate that, the total cost of cultivation for sunflower was Rs. 28263.78. The gross income realized by the farmers was Rs. 88272.38. The net income from sunflower cultivation was Rs. 60008.60. Thus the benefit cost ratio was found to be 1:3.12.*
- ❖ *The results indicate that, 7.50 per cent of the households opined that dry fodder was adequate, green fodder was adequate for 2.50 per cent of the households and 20 per cent of the households opined that dry fodder was in adequate.*
- ❖ *The results indicate that the annual gross income was Rs. 68,000 for landless farmers, for marginal farmers it was Rs. 64,250, for small farmers it was Rs. 79,483.33, for semi medium farmers it was Rs. 98,566.67 and for medium farmers it was Rs. 160,000. The results indicate that the average annual expenditure is Rs. 5,928.42. For landless households it was Rs. 4,200, for marginal farmers it was Rs. 4,682, for small farmers it was Rs. 5,366.67, for semi medium farmers it was Rs. 5,409.72 and for medium farmers it was Rs. 40,000.*
- ❖ *The results indicate that, sampled households have grown 37 coconut tree trees in their field.*
- ❖ *The results indicate that, households have planted 1 teak 14 neem, 6 tamarind, 9 acacia and 4 banyan trees in their field.*
- ❖ *The results indicated that, Bengal gram was sold to the extent 52.63 per cent, chilly was sold to the extent of 90 per cent, maize was sold to the extent of 96.01 per cent, paddy was sold to the extent of 52 per cent, sorghum was sold to the extent of 54.55 per cent and sunflower were sold to the extent of 100 per cent.*
- ❖ *The results indicated that, about 2.50 per cent of the farmers sold their produce to local/village merchants, 82.50 per cent of the farmers sold their produce to regulated market and 12.50 per cent of them sold their produce through contract marketing arrangement.*
- ❖ *The results indicated that, 95 per cent of the households used tractor and 2.50 per cent of them used flight as a mode of transportation for their agricultural produce.*
- ❖ *The results indicated that, 20 per cent of the households have experienced soil and water erosion problems in the farm.*
- ❖ *The results indicated that, 67.50 per cent have shown interest in soil test.*
- ❖ *The results indicated that, 95 per cent of the households used firewood and 2.5 per cent of the households used LPG as a source of fuel.*
- ❖ *The results indicated that, piped supply was the major source of drinking water for 45 per cent of the households and bore well was the source of drinking water for 55 per cent of the households in micro watershed.*
- ❖ *Electricity was the major source of light for 100 per cent of the households in micro watershed.*
- ❖ *The results indicated that, 42.50 per cent of the households possess sanitary toilet facility.*

- ❖ *The results indicated that, 2.50 per cent of the sampled households possessed APL card and 95 per cent of the sampled households possessed BPL card.*
- ❖ *The results indicated that, 62.50 per cent of the households participated in NREGA programme.*
- ❖ *The results indicated that, cereals were adequate for 90 per cent of the households, pulses were adequate for 72.50 per cent, oilseeds were adequate for 17.50 per cent, vegetables and egg were adequate for 7.50 per cent, fruits were adequate for 12.50 per cent, milk was adequate for 5 per cent and meat were adequate for 2.50 per cent.*
- ❖ *The results indicated that, cereals were in adequate for 10 per cent of the households, pulses were inadequate for 27.50 per cent, oilseeds were inadequate for 65 per cent, vegetables were inadequate for 70 per cent, fruits were inadequate for 67.50 per cent, milk was inadequate for 82.50 per cent, eggs were inadequate for 90 per cent and meat was inadequate for 75 per cent of the households.*
- ❖ *The results indicated that, oilseeds were market surplus for 15 per cent of the households, vegetables were market surplus for 20 per cent of the households, fruits were market surplus for 7.50 per cent of the households and meat were market surplus for 2.50 per cent of the households.*
- ❖ *The results indicated that, lower fertility status of the soil was the constraint experienced by 72.50 per cent of the households, wild animal menace on farm field (75%), frequent incidence of pest and diseases (57.50%), inadequacy of irrigation water (45%), high cost of fertilizers and plant protection chemicals (30%), high rate of interest on credit and lack of marketing facilities in the area (20%), low price for the agricultural commodities (22.5%), inadequate extension services (10%), lack of transport for safe transport of the agricultural produce to the market (25%), less rainfall (60%) and Source of Agri-technology information (37.5%).*

INTRODUCTION

Soil and water are the two precious natural resources which are essential for crop production and existence of life on earth. Rainfed agriculture is under severe stress due to various constraints related to agriculture like uneven and erratic distribution of rainfall, indiscriminate use of fertilizers, chemicals and pesticides, adoption of improper land management practices, soil erosion, decline in soil fertility, decline in ground water resources leading to low crop productivity. The area under rainfed agriculture has to be managed effectively using the best available practices to enhance the production of food, fodder and fuel. This is possible if the land resources are characterized at each parcel of land through detailed land resource inventory using the best available techniques of remote sensing, GPS and GIS. The watershed development programs are aimed at the sustainable distribution of its resources and the process of creating and implementing plans, programs, and projects to sustain and enhance watershed functions that affect the plant, animal and human communities within a watershed boundary.

World Bank funded KWDP II, SUJALA III project was implemented in with Broad objective of demonstrating more effective watershed management through greater integration of programmes related to rain-fed agriculture, innovative and science based approaches and strengthen institutional capacities and If successful, it is expected that the systems and tools could be mainstreamed into the overall IWMP in the State of Karnataka and in time, throughout other IWMP operations in India. With this background the socio-economic survey has been carried out with following specific objectives:

1. To understand the demographic features of the households in the micro-watershed
2. To understand the extent of family labour available and additional employment opportunities available within the village.
3. To know the status of assets of households in the micro-watershed for suggesting possible improvements.
4. To study the cropping pattern, cropped area and productivity levels of different households in micro-watershed.
5. To determine the type and extent of livestock owned by different categories of HHs
6. Availability of fodder and level of livestock management.

Scope and importance of survey

Survey helps in identification of different socio-economic and resource use-patterns of farmers at the Micro watershed. Household survey provides demographic features, labour force, and levels of education; land ownership and asset position (including livestock and other household assets) of surveyed households; and cropping patterns, input intensities, and average crop yields from farmers' fields. It also discusses crop utilization and the degree of commercialization of production in the areas; farmers' access to and utilization of credit from formal and informal sources; and the level of adoption and use of soil, water, and pest management technologies.

METHODOLOGY

The description of the methods, components selected for the survey and procedures followed in conducting the baseline survey are furnished under the following heads.

Description of the study area

Koppal district is an administrative district in the state of Karnataka in India. In the past Koppal was referred to as 'Kopana Nagara'. Koppal, now a district headquarters is ancient Kopana a major holy place of the Jainas. The district occupies an area of 7,190 km² and has a population of 1,196,089, which 16.58% were urban as of 2001. The Koppal district was formed after split of Raichur district.

Geographers are very particular about the physiography or relief of a region. It plays a very important role in the spatial analysis of agricultural situation of the study area. The undulating topography with black cotton soil shrips, cut across by numerous nalas or streams is the major characteristic feature of the study region. Three physiographic divisions have made considering the local conditions of landforms and crops grown in the district. On the basis of physiography, Koppal district can be divided into three major divisions. They are (a) Koppal & Yelburga plateau, (b) Maidan division, (c) Tungabhadra valley. The district is part of Krishna basin the main streams draining the area are Maskinala, Ilkal-nadi and Hirenala. These are Ephemeral in nature, these come under Tungabhadra sub-basin. The drainage exhibit dentritic to subdentritic with drainage density varies from 1.4 to 7.0kms/sq.km.

According to the 2011 census Koppal district has a population of 1,391,292, roughly equal to the nation of Swaziland or the US state of Hawaii. This gives it a ranking of 350th in India (out of a total of 640). The district has a population density of 250 inhabitants per square kilometre (650/sq mi). Its population growth rate over the decade 2001-2011 was 16.32%. Koppal has a sex ratio of 983 females for every 1000 males, and a literacy rate of 67.28%.

Description of the micro watershed

Naregallu-3 micro-watershed in Haligeri sub-watershed (Koppal taluk and district) is located in between 15^o25'2.211" to 15^o 23'57.159" North latitudes and 76^o 8'27.869" to 76^o5'59.853" East longitudes, covering an area of about 551.24 ha, bounded by Naregalla and Madhinura villages.

Methodology followed in assessing socio-economic status of households

In order to assess the socio-economic condition of the farmers in the watershed a comprehensive questionnaire was prepared. Major components such as demographic conditions, migration details, food consumption and family expenditure pattern, material possession, land holding, land use management, cropping pattern, cost of cultivation of crops, livestock management. The statistical components such as frequency and percentage were used to analyze the data. About 40 households located in the micro-watershed were interviewed for the survey.

SALIENT FEATURES OF THE SURVEY

Households Sampled For Socio-Economic Survey: The data on households sampled for socio economic survey in Naregallu-3 micro-watershed is presented in Table 1 and it indicated that 40 farmers were sampled in Naregallu-3 micro-watershed among them 5 (12.50%) were landless, 10 (25%) were marginal farmers, 12 (30%) were small farmers and semi medium farmers and 1 (2.50%) were medium farmers.

Table 1: Households sampled for socio economic survey in Naregallu-3 micro-watershed

Sl.No.	Particulars	LL (5)		MF (10)		SF (12)		SMF (12)		MDF (1)		All (40)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Farmers	5	12.50	10	25.00	12	30.00	12	30.00	1	2.50	40	100.00

Population characteristics: The population characteristics of households sampled for socio-economic survey in Naregallu-3 micro-watershed is presented in Table 2. The data indicated that there were 94 (56.29%) men and 73 (43.71%) women among the sampled households. The average family size of landless farmers' was 3, marginal farmers' was 4.2, small farmers' was 3.83, semi medium farmers' was 5 and medium farmers' was 4.

Table 2: Population characteristics of Naregallu-3 micro-watershed

Sl.No.	Particulars	LL (15)		MF (42)		SF (46)		SMF (60)		MDF (4)		All (167)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Men	8	53.33	24	57.14	26	56.52	34	56.67	2	50.00	94	56.29
2	Women	7	46.67	18	42.86	20	43.48	26	43.33	2	50.00	73	43.71
Total		15	100.00	42	100.00	46	100.00	60	100.00	4	100.00	167	100.00
Average		3.00		4.20		3.83		5.00		4.00		4.18	

Age wise classification of population: The age wise classification of household members in Naregallu-3 micro-watershed is presented in Table 3. The data indicated that, 18 (10.78%) people were in 0-15 years of age, 69 (41.32%) were in 16-35 years of age, 64 (38.32%) were in 36-60 years of age and 16 (9.58%) were above 61 years of age.

Table 3: Age wise classification of household members in Naregallu-3 micro-watershed

Sl.No.	Particulars	LL (15)		MF (42)		SF (46)		SMF (60)		MDF (4)		All (167)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	0-15 years of age	3	20.00	3	7.14	2	4.35	10	16.67	0	0.00	18	10.78
2	16-35 years of age	4	26.67	20	47.62	25	54.35	20	33.33	0	0.00	69	41.32
3	36-60 years of age	5	33.33	16	38.10	14	30.43	26	43.33	3	75.00	64	38.32
4	> 61 years	3	20.00	3	7.14	5	10.87	4	6.67	1	25.00	16	9.58
Total		15	100.00	42	100.00	46	100.00	60	100.00	4	100.00	167	100.00

Education level of household members: Education level of household members in Naregallu-3 micro-watershed is presented in Table 4. The results indicated that Naregallu-3 had 36.53 per cent illiterates, 35.33 per cent of them had primary school education, 4.19 per cent of them had middle school education, 13.77 per cent of them had

high school education, 7.19 per cent of them had PUC education and 1.20 per cent had diploma and ITI.

Table 4. Education level of household members in Naregallu-3 micro-watershed

Sl.No.	Particulars	LL (15)		MF (42)		SF (46)		SMF (60)		MDF (4)		All (167)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Illiterate	3	20.00	11	26.19	21	45.65	24	40.00	2	50.00	61	36.53
2	Primary School	8	53.33	16	38.10	11	23.91	22	36.67	2	50.00	59	35.33
3	Middle School	1	6.67	4	9.52	0	0.00	2	3.33	0	0.00	7	4.19
4	High School	2	13.33	6	14.29	9	19.57	6	10.00	0	0.00	23	13.77
5	PUC	1	6.67	2	4.76	5	10.87	4	6.67	0	0.00	12	7.19
6	ITI	0	0.00	1	2.38	0	0.00	1	1.67	0	0.00	2	1.20
7	Degree	0	0.00	1	2.38	0	0.00	1	1.67	0	0.00	2	1.20
8	Others	0	0.00	1	2.38	0	0.00	0	0.00	0	0.00	1	0.60
Total		15	100.00	42	100.00	46	100.00	60	100.00	4	100.00	167	100.00

Occupation of household heads: The data regarding the occupation of the household heads in Naregallu-3 micro-watershed is presented in Table 5. The results indicate that, 97.5 per cent of household heads were practicing agriculture and 2.5 per cent of the household heads were agricultural labourers.

Table 5: Occupation of household heads in Naregallu-3 micro-watershed

Sl. No.	Particulars	LL (5)		MF (10)		SF (12)		SMF (12)		MDF (1)		All (40)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Agriculture	4	80.00	10	100.00	12	100.00	12	100.00	1	100.00	39	97.50
2	Agricultural Labour	1	20.00	0	0.00	0	0.00	0	0.00	0	0.00	1	2.50
Total		5	100.00	10	100.00	12	100.00	12	100.00	1	100.00	40	100.00

Occupation of the household members: The data regarding the occupation of the household members in Naregallu-3 micro-watershed is presented in Table 6. The results indicate that agriculture was the major occupation for 23.95 per cent of the household members, 61.68 per cent were agricultural labourers and 14.37 per cent were student.

Table 6: Occupation of family members in Naregallu-3 micro-watershed

Sl.No.	Particulars	LL (15)		MF (42)		SF (46)		SMF (60)		MDF (4)		All (167)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Agriculture	4	26.67	10	23.81	12	26.09	13	21.67	1	25.00	40	23.95
2	Agricultural Labour	8	53.33	28	66.67	28	60.87	36	60.00	3	75.00	103	61.68
3	Student	3	20.00	4	9.52	6	13.04	11	18.33	0	0.00	24	14.37
Total		15	100.00	42	100.00	46	100.00	60	100.00	4	100.00	167	100.00

Institutional participation of the household members: The data regarding the institutional participation of the household members in Naregallu-3 micro-watershed is presented in Table 7. The results show that, 100 per cent of the population in the micro watershed has not participated in any local institutions.

Table 7. Institutional Participation of household members in Naregallu-3 micro-watershed

Sl.No.	Particulars	LL (15)		MF (42)		SF (46)		SMF (60)		MDF (4)		All (167)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	No Participation	15	100.00	42	100.00	46	100.00	60	100.00	4	100.00	167	100.00
	Total	15	100.00	42	100.00	46	100.00	60	100.00	4	100.00	167	100.00

Type of house owned: The data regarding the type of house owned by the households in Naregallu-3 micro-watershed is presented in Table 8. The results indicate that 5 per cent of the households possess thatched house and 95 per cent of the households possess katcha houses.

Table 8. Type of house owned by households in Naregallu-3 micro-watershed

Sl.No.	Particulars	LL (5)		MF (10)		SF (12)		SMF (12)		MDF (1)		All (40)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Thatched	2	40.00	0	0.00	0	0.00	0	0.00	0	0.00	2	5.00
2	Katcha	3	60.00	10	100.00	12	100.00	12	100.00	1	100.00	38	95.00
	Total	5	100.00	10	100.00	12	100.00	12	100.00	1	100.00	40	100.00

Durable Assets owned by the households: The data regarding the Durable Assets owned by the households in Naregallu-3 micro-watershed is presented in Table 9. The results show that 70 per cent of the households possess TV, 17.5 per cent of them possess mixer/grinder, 7.50 per cent of them possess bicycle, 35 per cent of the households possess motor cycle, 2.50 per cent of them possess auto and landline phone and 90 per cent of the households possess mobile phones.

Table 9. Durable Assets owned by households in Naregallu-3 micro-watershed

Sl.No.	Particulars	LL (5)		MF (10)		SF (12)		SMF (12)		MDF (1)		All (40)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Television	2	40.00	8	80.00	10	83.33	8	66.67	0	0.00	28	70.00
2	Mixer/Grinder	0	0.00	5	50.00	1	8.33	1	8.33	0	0.00	7	17.50
3	Bicycle	0	0.00	3	30.00	0	0.00	0	0.00	0	0.00	3	7.50
4	Motor Cycle	0	0.00	6	60.00	4	33.33	4	33.33	0	0.00	14	35.00
5	Auto	0	0.00	0	0.00	1	8.33	0	0.00	0	0.00	1	2.50
6	Landline Phone	0	0.00	0	0.00	0	0.00	1	8.33	0	0.00	1	2.50
7	Mobile Phone	3	60.00	9	90.00	12	100.00	11	91.67	1	100.00	36	90.00
8	Blank	2	40.00	1	10.00	0	0.00	0	0.00	0	0.00	3	7.50

Table 10. Average value of durable assets owned by households in Naregallu-3 micro-watershed

Average value (Rs.)

Sl.No.	Particulars	LL (5)	MF (10)	SF (12)	SMF (12)	MDF (1)	All (40)
1	Television	7,500.00	5,312.00	6,500.00	8,250.00	0.00	6,732.00
2	Mixer/Grinder	0.00	1,840.00	700.00	2,000.00	0.00	1,700.00
3	Bicycle	0.00	1,175.00	0.00	0.00	0.00	1,175.00
4	Motor Cycle	0.00	33,833.00	28,750.00	33,750.00	0.00	32,357.00
5	Auto	0.00	0.00	300,000.00	0.00	0.00	300,000.00
6	Landline Phone	0.00	0.00	0.00	4,000.00	0.00	4,000.00
7	Mobile Phone	2,833.00	2,291.00	2,442.00	2,458.00	5,000.00	2,492.00

Average value of durable assets: The data regarding the average value of durable assets owned by the households in Naregallu-3 micro-watershed is presented in Table 10. The results show that the average value of television was Rs. 6,732, mixer grinder was Rs. 1,700, bicycle was 1,175, motor cycle was Rs. 32,357, auto was 300,000, landline was 4,000 and mobile phone was Rs.2,492.

Farm Implements owned: The data regarding the farm implements owned by the households in Naregallu-3 micro-watershed is presented in Table 11. About 12.50 per cent of the households possess bullock cart, 17.50 per cent of them possess plough, 5 per cent of them possess seed/ fertilizer drill, 2.5 per cent of them possess sprayer, 12.50 per cent of them possess weeder, and 17.50 per cent of them possess chaff cutter.

Table 11. Farm Implements owned by households in Naregallu-3 micro-watershed

Sl.No.	Particulars	LL (5)		MF (10)		SF (12)		SMF (12)		MDF (1)		All (40)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Bullock Cart	0	0.00	1	10.00	1	8.33	2	16.67	1	100.00	5	12.50
2	Plough	0	0.00	2	20.00	2	16.67	2	16.67	1	100.00	7	17.50
3	Seed/Fertilizer Drill	0	0.00	1	10.00	0	0.00	1	8.33	0	0.00	2	5.00
4	Sprayer	0	0.00	0	0.00	0	0.00	1	8.33	0	0.00	1	2.50
5	Weeder	0	0.00	2	20.00	2	16.67	1	8.33	0	0.00	5	12.50
6	Chaff Cutter	0	0.00	2	20.00	2	16.67	2	16.67	1	100.00	7	17.50
7	Blank	5	100.00	6	60.00	9	75.00	8	66.67	0	0.00	28	70.00

Average value of farm implements: The data regarding the average value of farm Implements owned by the households in Naregallu-3 micro-watershed is presented in Table 12. The results show that the average value of bullock cart was Rs. 17,060, plough was Rs. 2800, seed/fertilizer drill was Rs. 8,000, sprayer was Rs. 2,000, weeder was Rs. 21 and the average value of chaff cutter was Rs. 441.

Table 12. Average value of farm implements owned by households in Naregallu-3 micro-watershed

Sl.No.	Particulars	MF (10)	SF (12)	SMF (12)	MDF (1)	All (40)
1	Bullock Cart	20,000.00	1,300.00	22,000.00	20,000.00	17,060.00
2	Plough	1,000.00	1,000.00	666.00	500.00	800.00
3	Seed/Fertilizer Drill	15,000.00	0.00	1,000.00	0.00	8,000.00
4	Sprayer	0.00	0.00	2,000.00	0.00	2,000.00
5	Weeder	27.00	15.00	25.00	0.00	21.00
6	Chaff Cutter	180.00	1,110.00	165.00	180.00	441.00

Table 13. Livestock possession by households in Naregallu-3 micro-watershed

Sl.No.	Particulars	LL (5)		MF (10)		SF (12)		SMF (12)		MDF (1)		All (40)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Bullock	0	0.00	3	30.00	2	16.67	3	25.00	1	100.00	9	22.50
2	Local cow	0	0.00	1	10.00	0	0.00	1	8.33	1	100.00	3	7.50
3	Buffalo	0	0.00	0	0.00	1	8.33	0	0.00	0	0.00	1	2.50
4	Goat	0	0.00	0	0.00	1	8.33	0	0.00	0	0.00	1	2.50
5	blank	5	100.00	7	70.00	8	66.67	8	66.67	0	0.00	28	70.00

Livestock possession by the households: The data regarding the Livestock possession by the households in Naregallu-3 micro-watershed is presented in Table 13. The results indicate that, 22.50 per cent of the households possess bullocks, 7.50 per cent of the households possess local cow, 2.50 per cent possess buffalo and goat.

Average Labour availability: The data regarding the average labour availability in Naregallu-3 micro-watershed is presented in Table 14. The results indicate that, average own labour men available in the micro watershed was 1.40, average own labour (women) available was 1.85, average hired labour (men) available was 6.25 and average hired labour (women) available was 6.4.

Table 14. Average Labour availability in Naregallu-3 micro-watershed

Sl.No.	Particulars	MF (10)	SF (12)	SMF (12)	MDF (1)	All (40)
1	Hired labour Female	6.10	6.25	8.75	5.00	6.40
2	Own Labour Female	1.20	1.50	1.83	2.00	1.40
3	Own labour Male	2.40	1.75	2.00	2.00	1.85
4	Hired labour Male	6.10	5.75	8.75	5.00	6.25

Adequacy of Hired Labour: The data regarding the adequacy of hired labour in Naregallu-3 micro-watershed is presented in Table 15. The results indicate that 100 per cent of the households opined that the hired labour was inadequate.

Table 15. Adequacy of Hired Labour in Naregallu-3 micro-watershed

Sl.No.	Particulars	LL (5)		MF (10)		SF (12)		SMF (12)		MDF (1)		All (40)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Inadequate	5	100.00	10	100.00	12	100.00	12	100.00	1	100.00	40	100.00

Distribution of land (ha): The data regarding the distribution of land (ha) in Naregallu-3 micro-watershed is presented in Table 16. The results indicate that, households of the Naregallu-3 micro-watershed possess 29.15 ha (59.87%) of dry land and 19.54 ha (40.13%) of irrigated land. Marginal farmers possess 5.84 ha (86.61%) of dry land and 0.90 ha (13.39%) of irrigated land. Small farmers possess 14.26 ha (84.63%) of dry land and 2.59 ha (15.37%) of irrigated land. Semi medium farmers possess 9.05 ha (40.66%) of dry land and 13.21 ha (59.34%) of irrigated land. Medium farmers possess 2.83 ha (100%) of irrigated land.

Table 16. Distribution of land (Ha) in Naregallu-3 micro-watershed

Sl.No.	Particulars	MF (10)		SF (12)		SMF (12)		MDF (1)		All (40)	
		ha	%	ha	%	ha	%	ha	%	ha	%
1	Dry	5.84	86.61	14.26	84.63	9.05	40.66	0.00	0.00	29.15	59.87
2	Irrigated	0.90	13.39	2.59	15.37	13.21	59.34	2.83	100.00	19.54	40.13
Total		6.74	100.00	16.85	100.00	22.27	100.00	2.83	100.00	48.69	100.00

Average land value (Rs./ha): The data regarding the average land value (Rs./ha) in Naregallu-3 micro-watershed is presented in Table 17. The results indicate that, the average value of dry land was Rs. 264,006.11 and the average value of irrigated land was Rs. 450,207.13. In case of marginal famers, the average land value was Rs. 530,630.63

for dry land and Rs. 1,218,385.64 for irrigated land. In case of small famers, the average land value was Rs. 238,308.74 for dry land and Rs. 694,687.49 for irrigated land. In case of semi medium famers, the average land value was Rs. 132,498.88 for dry land and Rs. 400,949.47 for irrigated land. In case of medium farmers, the average land value was Rs. 211,714.29 for irrigated land.

Table 17. Average land value (Rs./ha) in Naregallu-3 micro-watershed

Sl.No.	Particulars	MF (10)	SF (12)	SMF (12)	MDF (1)	All (40)
1	Dry	530,630.63	238,308.74	132,498.88	0.00	264,006.11
2	Irrigated	1,218,385.64	694,687.49	400,949.47	211,714.29	450,207.13

Status of bore wells: The data regarding the status of bore wells in Naregallu-3 micro-watershed is presented in Table 18. The results indicate that, there were 11 functioning and 6 de-functioning bore wells in the micro watershed.

Table 18. Status of bore wells in Naregallu-3 micro-watershed

Sl.No.	Particulars	LL (5)	MF (10)	SF (12)	SMF (12)	MDF (1)	All (40)
1	De-functioning	0	1	2	3	0	6
2	Functioning	0	1	3	7	0	11

Status of open wells: The data regarding the status of open wells in Naregallu-3 micro-watershed is presented in Table 19. The results indicate that, there was 1 functioning open well in the micro watershed.

Table 19. Status of open wells in Naregallu-3 micro-watershed

Sl.No.	Particulars	LL (5)	MF (10)	SF (12)	SMF (12)	MDF (1)	All (40)
1	Functioning	0	1	0	0	0	1

Source of irrigation: The data regarding the source of irrigation in Naregallu-3 micro-watershed is presented in Table 20. The results indicate that, bore well was the major irrigation source in the micro water shed for 27.50 per cent of the farmers and open well was the major source of irrigation for 2.50 per cent of the farmers.

Table 20. Source of irrigation in Naregallu-3 micro-watershed

Sl.No.	Particulars	MF (10)		SF (12)		SMF (12)		MDF (1)		All (40)	
		N	%	N	%	N	%	N	%	N	%
1	Bore Well	1	10.00	3	25.00	7	58.33	0	0.00	11	27.50
2	Open Well	1	10.00	0	0.00	0	0.00	0	0.00	1	2.50

Depth of water (Avg in meters): The data regarding the depth of water in Naregallu-3 micro-watershed is presented in Table 21. The results indicate that, the depth of bore well was found to be 13.72 meters and the depth of open well was found to be 3.05 meters.

Table 21. Depth of water (Avg in meters) in Naregallu-3 micro-watershed

Sl.No.	Particulars	MF (10)	SF (12)	SMF (12)	MDF (1)	All (40)
1	Bore Well	4.57	18.16	23.75	0.00	13.72
2	Open Well	12.19	0.00	0.00	0.00	3.05

Irrigated Area (ha): The data regarding the irrigated area (ha) in Naregallu-3 micro-watershed is presented in Table 22. The results indicate that marginal, small and semi medium farmers had an irrigated area of 1.78 ha, 2.55 ha and 11.92 ha respectively.

Table 22. Irrigated Area (ha) in Naregallu-3 micro-watershed

Sl.No.	Particulars	MF (10)	SF (12)	SMF (12)	MDF (1)	All (40)
1	Kharif	1.78	2.55	11.92	0.00	16.25

Cropping pattern: The data regarding the cropping pattern in Naregallu-3 micro-watershed is presented in Table 23. The results indicate that, farmers have grown maize (31.79 ha), sunflower (4.91 ha), sorghum (4.61 ha), Bengal gram (2.14 ha), paddy (1.62 ha), chilly and jowar (1.21 ha). Marginal farmers have grown maize and Bengal gram, while small farmers have grown maize, sunflower, sorghum, Bengal gram paddy and jowar. Semi medium farmers have grown maize, sunflower, sorghum, paddy and chilly. Medium farmers have grown maize.

Table 23. Cropping pattern in Naregallu-3 micro-watershed (Area in ha)

Sl.No.	Particulars	MF (10)	SF (12)	SMF (12)	MDF (1)	All (40)
1	Kharif - Maize	7.07	7.13	14.76	2.83	31.79
2	Kharif - Sunflower	0	2.48	2.43	0	4.91
3	Kharif - Sorghum	0	3.39	1.21	0	4.61
4	Kharif - Bengal gram	0	1.65	0	0	1.65
5	Kharif - Paddy	0	0.81	0.81	0	1.62
6	Kharif - Chilly	0	0	1.21	0	1.21
7	Rabi - Jowar	0	1.21	0	0	1.21
8	Rabi - Bengal gram	0.49	0	0	0	0.49
Total		7.55	16.68	20.43	2.83	47.49

Cropping intensity: The data regarding the cropping intensity in Naregallu-3 micro-watershed is presented in Table 24. The results indicate that, the cropping intensity in Naregallu-3 micro-watershed was found to be 88.20 per cent.

Table 24. Cropping intensity (%) in Naregallu-3 micro-watershed

Sl.No.	Particulars	MF (10)	SF (12)	SMF (12)	MDF (1)	All (40)
1	Cropping Intensity	100.00	90.67	81.50	100.00	88.20

Possession of Bank account and savings: The data regarding the possession of bank account and saving in Naregallu-3 micro-watershed is presented in Table 25. The results indicate that, 55 per cent of the households have bank account and savings.

Table 25. Possession of Bank account and savings in Naregallu-3 micro-watershed

Sl.No.	Particulars	MF (10)		SF (12)		SMF (12)		MDF (1)		All (40)	
		N	%	N	%	N	%	N	%	N	%
1	Account	6	60.00	6	50.00	9	75.00	1	100.00	22	55.00
2	Savings	6	60.00	6	50.00	9	75.00	1	100.00	22	55.00

Borrowing status: The data regarding the borrowing status in Naregallu-3 micro-watershed is presented in Table 26. The results indicate that, 55 per cent of the households have availed credit from different sources.

Table 26. Borrowing status in Naregallu-3 micro-watershed

Sl.No.	Particulars	MF (10)		SF (12)		SMF (12)		MDF (1)		All (40)	
		N	%	N	%	N	%	N	%	N	%
1	Credit Availed	6	60.00	6	50.00	9	75.00	1	100.00	22	55.00

Source of credit availed by households: The data regarding the Source of credit availed in Naregallu-3 micro-watershed is presented in Table 27. The results indicate that, 81.82 per cent of the households have borrowed from commercial bank, 4.55 per cent of them borrowed from friends/ relatives, grameena bank and money lender, 9.09 per cent of the households borrowed from SHGs and CBOs.

Table 27. Source of credit availed by household in Naregallu-3 micro-watershed

Sl.No.	Particulars	MF (6)		SF (6)		SMF (9)		MDF (1)		All (22)	
		N	%	N	%	N	%	N	%	N	%
1	Commercial Bank	4	66.67	4	66.67	9	100.00	1	100.00	18	81.82
2	Friends/Relatives	1	16.67	0	0.00	0	0.00	0	0.00	1	4.55
3	Grameena Bank	0	0.00	1	16.67	0	0.00	0	0.00	1	4.55
4	Money Lender	0	0.00	1	16.67	0	0.00	0	0.00	1	4.55
5	SHGs/CBOs	0	0.00	1	16.67	1	11.11	0	0.00	2	9.09

Avg. Credit amount: The data regarding the avg. Credit amount in Naregallu-3 micro-watershed is presented in Table 28. The results indicate that, the average credit amount borrowed by households in micro-watershed was Rs, 59,409.09.

Table 28. Avg. Credit amount by household in Naregallu-3 micro-watershed

Sl.No.	Particulars	MF (6)	SF (6)	SMF (9)	MDF (1)	All (22)
1	Average Credit	29,166.67	94,166.67	60,777.78	20,000.00	59,409.09

Purpose of credit borrowed - Institutional Credit: The data regarding the purpose of credit borrowed - Institutional Credit in Naregallu-3 micro-watershed is presented in Table 29. The results indicate that, 100 per cent of the households borrowed from institutional sources for the purpose of agricultural production.

Table 29. Purpose of credit borrowed - Institutional Credit by household in Naregallu-3 micro-watershed

Sl.No.	Particulars	MF (4)		SF (5)		SMF (9)		MDF (1)		All (19)	
		N	%	N	%	N	%	N	%	N	%
1	Agriculture production	4	100.00	5	100.00	9	100.00	1	100.00	19	100.00

Purpose of credit borrowed - Private Credit: The data regarding the purpose of credit borrowed - Institutional Credit in Naregallu-3 micro-watershed is presented in Table 30. The results indicate that, 100 per cent of the households availed credit for the purpose of agricultural production.

Table 30. Purpose of credit borrowed - Private Credit by household in Naregallu-3 micro-watershed

Sl.No.	Particulars	MF (1)		SF (2)		SMF (1)		All (4)	
		N	%	N	%	N	%	N	%
1	Agriculture production	1	100.00	2	100.00	1	100.00	4	100.00

Repayment status of households – Institutional: The data regarding the repayment status of credit borrowed from institutional sources by households in Naregallu-3 micro watershed is presented in Table 31. The results indicated that 100 per cent of the households did not repay their loan borrowed from institutional sources.

Table 31. Repayment status of households – Institutional Credit in Naregallu-3 micro-watershed

Sl.No.	Particulars	MF (4)		SF (5)		SMF (9)		MDF (1)		All (19)	
		N	%	N	%	N	%	N	%	N	%
1	Un paid	4	100.00	5	100.00	9	100.00	1	100.00	19	100.00

Repayment status of households – Private: The data regarding the repayment status of credit borrowed from private sources by households in Naregallu-3 micro watershed is presented in Table 32. Results indicated that 100 per cent of the households did not repay their loan borrowed from private sources.

Table 32. Repayment status of households (private sources) in Naregallu-3 micro watershed

Sl.No.	Particulars	MF (1)		SF (2)		SMF (1)		All (4)	
		N	%	N	%	N	%	N	%
1	Un paid	1	100.00	2	100.00	1	100.00	4	100.00

Opinion on institutional sources of credit: The data regarding the opinion on institutional sources of credit in Naregallu-3 micro watershed is presented in Table 33. The results indicate that, around 100 per cent opined that the loan amount borrowed from institutional sources helped to perform timely agricultural operations.

Table 33. Opinion on institutional sources of credit in Naregallu-3 micro watershed

Sl.No.	Particulars	MF (4)		SF (5)		SMF (9)		MDF (1)		All (19)	
		N	%	N	%	N	%	N	%	N	%
1	Helped to perform timely agricultural operations	4	100.00	5	100.00	9	100.00	1	100.00	19	100.00

Opinion on non-institutional sources of credit: The data regarding the opinion on non-institutional sources of credit in Naregallu-3 micro watershed is presented in Table 34. The results indicate that, 100 per cent of the households opined that the credit borrowed from private credit helped to perform timely agricultural operations.

Table 34. Opinion on non-institutional sources of credit in Naregallu-3 micro watershed

Sl.No.	Particulars	MF (1)		SF (2)		SMF (1)		All (4)	
		N	%	N	%	N	%	N	%
1	Helped to perform timely agricultural operations	1	100.00	2	100.00	1	100.00	4	100.00

Cost of cultivation of Maize: The data regarding the cost of cultivation of maize in Naregallu-3 micro-watershed is presented in Table 35. The results indicate that, the total cost of cultivation for maize was Rs. 33037.55. The gross income realized by the farmers was Rs. 45231.00. The net income from maize cultivation was Rs. 12193.46. Thus the benefit cost ratio was found to be 1:1.37.

Table 35. Cost of Cultivation of maize in Naregallu-3 micro-watershed

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
I	Cost A1				
1	Hired Human Labour	Man days	35.19	8090.16	24.49
2	Bullock	Pairs/day	1.09	602.22	1.82
3	Tractor	Hours	2.00	1497.52	4.53
4	Machinery	Hours	0.62	370.19	1.12
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	25.39	3046.97	9.22
6	Seed Inter Crop	Kgs.	0.00	0.00	0.00
7	FYM	Quintal	8.75	2378.59	7.20
8	Fertilizer + micronutrients	Quintal	3.34	2976.68	9.01
9	Pesticides (PPC)	Kgs / liters	1.56	1972.32	5.97
10	Irrigation	Number	7.05	0.00	0.00
11	Repairs		0.00	0.00	0.00
12	Msc. Charges (Marketing costs etc)		0.00	0.00	0.00
13	Depreciation charges		0.00	112.56	0.34
14	Land revenue and Taxes		0.00	0.00	0.00
II	Cost B1				
16	Interest on working capital			1246.15	3.77
17	Cost B1 = (Cost A1 + sum of 15 and 16)			22293.37	67.48
III	Cost B2				
18	Rental Value of Land			317.33	0.96
19	Cost B2 = (Cost B1 + Rental value)			22610.70	68.44
IV	Cost C1				
20	Family Human Labour		29.09	7413.43	22.44
21	Cost C1 = (Cost B2 + Family Labour)			30024.13	90.88
V	Cost C2				
22	Risk Premium			10.00	0.03
23	Cost C2 = (Cost C1 + Risk Premium)			30034.13	90.91
VI	Cost C3				
24	Managerial Cost			3003.41	9.09
25	Cost C3 = (Cost C2 + Managerial Cost)			33037.55	100.00
VII	Economics of the Crop				
a.	Main Product	a) Main Product (q)	31.80	38288.12	
		b) Main Crop Sales Price (Rs.)		1204.00	
	By Product	e) Main Product (q)	18.87	6942.88	
		f) Main Crop Sales Price (Rs.)		368.00	
b.	Gross Income (Rs.)			45231.00	
c.	Net Income (Rs.)			12193.46	
d.	Cost per Quintal (Rs./q.)			1038.89	
e.	Benefit Cost Ratio (BC Ratio)			1:1.37	

Cost of Cultivation of Chilly: The data regarding the cost of cultivation of chilly in Naregallu-3 micro-watershed is presented in Table 36. The results indicate that, the total cost of cultivation for chilly was Rs. 57632.23. The gross income realized by the farmers was Rs. 89969.75. The net income from chilly cultivation was Rs. 32337.52. Thus the benefit cost ratio was found to be 1:1.56.

Table 36. Cost of Cultivation of Chilly in Naregallu-3 micro-watershed

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
I	Cost A1				
1	Hired Human Labour	Man days	45.23	9887.72	17.16
2	Bullock	Pairs/day	0.00	0.00	0.00
3	Tractor	Hours	3.24	2431.41	4.22
4	Machinery	Hours	2.01	1204.12	2.09
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	3.24	4013.75	6.96
7	FYM	Quintal	49.40	9880.00	17.14
8	Fertilizer + micronutrients	Quintal	2.78	2825.06	4.90
9	Pesticides (PPC)	Kgs / liters	1.54	1852.50	3.21
10	Irrigation	Number	7.41	0.00	0.00
11	Repairs		0.00	0.00	0.00
12	Msc. Charges (Marketing costs etc)		0.00	0.00	0.00
13	Depreciation charges		0.00	0.03	0.00
14	Land revenue and Taxes		0.00	0.00	0.00
II	Cost B1				
16	Interest on working capital			2229.76	3.87
17	Cost B1 = (Cost A1 + sum of 15 and 16)			34324.35	59.56
III	Cost B2				
18	Rental Value of Land			583.33	1.01
19	Cost B2 = (Cost B1 + Rental value)			34907.69	60.57
IV	Cost C1				
20	Family Human Labour		70.32	17475.25	30.32
21	Cost C1 = (Cost B2 + Family Labour)			52382.94	90.89
V	Cost C2				
22	Risk Premium			10.00	0.02
23	Cost C2 = (Cost C1 + Risk Premium)			52392.94	90.91
VI	Cost C3				
24	Managerial Cost			5239.29	9.09
25	Cost C3 = (Cost C2 + Managerial Cost)			57632.23	100.00
VII	Economics of the Crop				
a.	Main Product	a) Main Product (q)		14.51	89969.75
		b) Main Crop Sales Price (Rs.)			6200.00
b.	Gross Income (Rs.)			89969.75	
c.	Net Income (Rs.)			32337.52	
d.	Cost per Quintal (Rs./q.)			3971.56	
e.	Benefit Cost Ratio (BC Ratio)			1:1.56	

Cost of Cultivation of Sorghum: The data regarding the cost of cultivation of sorghum in Naregallu-3 micro-watershed is presented in Table 37. The results indicate that, the total cost of cultivation for sorghum was Rs. 22927.34. The gross income realized by the farmers was Rs. 33038.15. The net income from sorghum cultivation was Rs. 10110.81. Thus the benefit cost ratio was found to be 1:1.44.

Table 37. Cost of Cultivation of sorghum in Naregallu-3 micro-watershed

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
I	Cost A1				
1	Hired Human Labour	Man days	31.28	6637.21	28.95
2	Bullock	Pairs/day	0.67	367.93	1.60
3	Tractor	Hours	1.43	1068.99	4.66
4	Machinery	Hours	0.77	463.13	2.02
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	7.93	793.45	3.46
7	FYM	Quintal	3.91	782.17	3.41
8	Fertilizer + micronutrients	Quintal	3.81	4281.90	18.68
9	Pesticides (PPC)	Kgs / liters	0.75	820.59	3.58
10	Irrigation	Number	0.00	0.00	0.00
11	Repairs		0.00	0.00	0.00
12	Msc. Charges (Marketing costs etc)		0.00	0.00	0.00
13	Depreciation charges		0.00	22.49	0.10
14	Land revenue and Taxes		0.00	0.00	0.00
II	Cost B1				
16	Interest on working capital			802.57	3.50
17	Cost B1 = (Cost A1 + sum of 15 and 16)			16040.41	69.96
III	Cost B2				
18	Rental Value of Land			125.00	0.55
19	Cost B2 = (Cost B1 + Rental value)			16165.41	70.51
IV	Cost C1				
20	Family Human Labour		19.16	4667.62	20.36
21	Cost C1 = (Cost B2 + Family Labour)			20833.03	90.87
V	Cost C2				
22	Risk Premium			10.00	0.04
23	Cost C2 = (Cost C1 + Risk Premium)			20843.03	90.91
VI	Cost C3				
24	Managerial Cost			2084.30	9.09
25	Cost C3 = (Cost C2 + Managerial Cost)			22927.34	100.00
VII	Economics of the Crop				
a.	Main Product	a) Main Product (q)		18.16	28835.49
		b) Main Crop Sales Price (Rs.)			1587.50
	By Product	e) Main Product (q)		15.28	4202.67
		f) Main Crop Sales Price (Rs.)			275.00
b.	Gross Income (Rs.)			33038.15	
c.	Net Income (Rs.)			10110.81	
d.	Cost per Quintal (Rs./q.)			1262.23	
e.	Benefit Cost Ratio (BC Ratio)			1:1.44	

Cost of cultivation of Bengal gram: The data regarding the cost of cultivation of Bengal gram in Naregallu-3 micro-watershed is presented in Table 38. The results indicate that, the total cost of cultivation for Bengal gram was Rs. 57703.23. The gross income realized by the farmers was Rs. 29624.86. The net income from Bengal gram cultivation was Rs. -28078.36. Thus the benefit cost ratio was found to be 1:0.51.

Table 38. Cost of Cultivation of Bengal gram in Naregallu-3 micro-watershed

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
I	Cost A1				
1	Hired Human Labour	Man days	72.22	15806.79	27.39
2	Bullock	Pairs/day	0.30	166.48	0.29
3	Tractor	Hours	4.00	2996.69	5.19
4	Machinery	Hours	0.00	0.00	0.00
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	82.64	8263.60	14.32
7	FYM	Quintal	5.75	10412.74	18.05
8	Fertilizer + micronutrients	Quintal	2.24	3844.24	6.66
9	Pesticides (PPC)	Kgs / liters	1.33	3580.89	6.21
10	Irrigation	Number	0.00	0.00	0.00
11	Repairs		0.00	0.00	0.00
12	Msc. Charges (Marketing costs etc)		0.00	0.00	0.00
13	Depreciation charges		0.00	1.70	0.00
14	Land revenue and Taxes		0.00	0.00	0.00
II	Cost B1				
16	Interest on working capital			3133.38	5.43
17	Cost B1 = (Cost A1 + sum of 15 and 16)			48206.52	83.54
III	Cost B2				
18	Rental Value of Land			166.67	0.29
19	Cost B2 = (Cost B1 + Rental value)			48373.19	83.83
IV	Cost C1				
20	Family Human Labour		15.56	4074.29	7.06
21	Cost C1 = (Cost B2 + Family Labour)			52447.48	90.89
V	Cost C2				
22	Risk Premium			10.00	0.02
23	Cost C2 = (Cost C1 + Risk Premium)			52457.48	90.91
VI	Cost C3				
24	Managerial Cost			5245.75	9.09
25	Cost C3 = (Cost C2 + Managerial Cost)			57703.23	100.00
VII	Economics of the Crop				
a.	Main Product	a) Main Product (q)	7.93	27360.70	
		b) Main Crop Sales Price (Rs.)		3450.00	
	By Product	e) Main Product (q)	2.06	2264.17	
		f) Main Crop Sales Price (Rs.)		1100.00	
b.	Gross Income (Rs.)			29624.86	
c.	Net Income (Rs.)			-28078.36	
d.	Cost per Quintal (Rs./q.)			7275.99	
e.	Benefit Cost Ratio (BC Ratio)			1:0.51	

Cost of cultivation of Sunflower: The data regarding the cost of cultivation of sunflower in Naregallu-3 micro-watershed is presented in Table 39. The results indicate that, the total cost of cultivation for sunflower was Rs. 28263.78. The gross income realized by the farmers was Rs. 88272.38. The net income from sunflower cultivation was Rs. 60008.60. Thus the benefit cost ratio was found to be 1:3.12.

Table 39. Cost of Cultivation of sunflower in Naregallu-3 micro-watershed

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
I	Cost A1				
1	Hired Human Labour	Man days	41.42	9909.37	35.06
2	Bullock	Pairs/day	1.85	1018.88	3.60
3	Tractor	Hours	1.84	1378.08	4.88
4	Machinery	Hours	0.00	0.00	0.00
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	9.22	1970.88	6.97
7	FYM	Quintal	0.00	0.00	0.00
8	Fertilizer + micronutrients	Quintal	3.07	2949.69	10.44
9	Pesticides (PPC)	Kgs / liters	1.64	2130.63	7.54
10	Irrigation	Number	6.18	0.00	0.00
11	Repairs		0.00	0.00	0.00
12	Msc. Charges (Marketing costs etc)		0.00	0.00	0.00
13	Depreciation charges		0.00	155.31	0.55
14	Land revenue and Taxes		0.00	0.00	0.00
II	Cost B1				
16	Interest on working capital			847.34	3.00
17	Cost B1 = (Cost A1 + sum of 15 and 16)			20360.18	72.04
III	Cost B2				
18	Rental Value of Land			208.33	0.74
19	Cost B2 = (Cost B1 + Rental value)			20568.51	72.77
IV	Cost C1				
20	Family Human Labour		20.19	5115.84	18.10
21	Cost C1 = (Cost B2 + Family Labour)			25684.35	90.87
V	Cost C2				
22	Risk Premium			10.00	0.04
23	Cost C2 = (Cost C1 + Risk Premium)			25694.35	90.91
VI	Cost C3				
24	Managerial Cost			2569.43	9.09
25	Cost C3 = (Cost C2 + Managerial Cost)			28263.78	100.00
VII	Economics of the Crop				
a.	Main Product	a) Main Product (q)		15.69	88233.78
		b) Main Crop Sales Price (Rs.)			5625.00
	By Product	e) Main Product (q)		1.54	38.59
		f) Main Crop Sales Price (Rs.)			25.00
b.	Gross Income (Rs.)			88272.38	
c.	Net Income (Rs.)			60008.60	
d.	Cost per Quintal (Rs./q.)			1801.85	
e.	Benefit Cost Ratio (BC Ratio)			1:3.12	

Cost of cultivation of Paddy: The data regarding the cost of cultivation of paddy in Naregallu-3 micro-watershed is presented in Table 40. The results indicate that, the total cost of cultivation for paddy was Rs. 41985.86. The gross income realized by the farmers was Rs. 51252.50. The net income from paddy cultivation was Rs. 9266.64. Thus the benefit cost ratio was found to be 1:1.22.

Table 40. Cost of Cultivation of paddy in Naregallu-3 micro-watershed

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
I	Cost A1				
1	Hired Human Labour	Man days	41.37	10374.00	24.71
2	Bullock	Pairs/day	0.00	0.00	0.00
3	Tractor	Hours	2.47	1852.50	4.41
4	Machinery	Hours	0.62	370.50	0.88
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	74.10	1852.50	4.41
7	FYM	Quintal	24.70	4940.00	11.77
8	Fertilizer + micronutrients	Quintal	1.85	1667.25	3.97
9	Pesticides (PPC)	Kgs / liters	1.85	2470.00	5.88
10	Irrigation	Number	6.18	0.00	0.00
11	Repairs		0.00	0.00	0.00
12	Msc. Charges (Marketing costs etc)		0.00	0.00	0.00
13	Depreciation charges		0.00	0.02	0.00
14	Land revenue and Taxes		0.00	0.00	0.00
II	Cost B1				
16	Interest on working capital			1312.77	3.13
17	Cost B1 = (Cost A1 + sum of 15 and 16)			24839.54	59.16
III	Cost B2				
18	Rental Value of Land			166.67	0.40
19	Cost B2 = (Cost B1 + Rental value)			25006.21	59.56
IV	Cost C1				
20	Family Human Labour		51.25	13152.75	31.33
21	Cost C1 = (Cost B2 + Family Labour)			38158.96	90.89
V	Cost C2				
22	Risk Premium			10.00	0.02
23	Cost C2 = (Cost C1 + Risk Premium)			38168.96	90.91
VI	Cost C3				
24	Managerial Cost			3816.90	9.09
25	Cost C3 = (Cost C2 + Managerial Cost)			41985.86	100.00
VII	Economics of the Crop				
a.	Main Product	a) Main Product (q)		30.88	50943.75
		b) Main Crop Sales Price (Rs.)			1650.00
	By Product	e) Main Product (q)		6.18	308.75
		f) Main Crop Sales Price (Rs.)			50.00
b.	Gross Income (Rs.)			51252.50	
c.	Net Income (Rs.)			9266.64	
d.	Cost per Quintal (Rs./q.)			1359.87	
e.	Benefit Cost Ratio (BC Ratio)			1:1.22	

Adequacy of fodder: The data regarding the adequacy of fodder in Naregallu-3 micro-watershed is presented in Table 41. The results indicate that, 7.50 per cent of the households opined that dry fodder was adequate, green fodder was adequate for 2.50 per cent of the households and 20 per cent of the households opined that dry fodder was in adequate.

Table 41. Adequacy of fodder in Naregallu-3 micro-watershed

Sl.No.	Particulars	MF (10)		SF (12)		SMF (12)		MDF (1)		All (40)	
		N	%	N	%	N	%	N	%	N	%
1	Adequate-Dry Fodder	0	0.00	1	8.33	2	16.67	0	0.00	3	7.50
2	Inadequate-Dry Fodder	1	10.00	3	25.00	3	25.00	1	100.00	8	20.00
3	Adequate-Green Fodder	0	0.00	1	8.33	0	0.00	0	0.00	1	2.50

Annual gross income: The data regarding the annual gross income in Naregallu-3 micro-watershed is presented in Table 42. The results indicate that the annual gross income was Rs. 68,000 for landless farmers, for marginal farmers it was Rs.64,250, for small farmers it was Rs.79,483.33, for semi medium farmers it was Rs. 98,566.67 and for medium farmers it was Rs. 160,000.

Table 42. Annual gross income in Naregallu-3 micro-watershed

(Avg value in Rs.)

Sl.No.	Particulars	LL (5)	MF (10)	SF (12)	SMF (12)	MDF (1)	All (40)
1	Service/salary	0.00	6,000.00	0.00	4,166.67	0.00	2,750.00
2	Business	0.00	0.00	7,083.33	0.00	0.00	2,125.00
3	Wage	68,000.00	24,500.00	24,166.67	18,333.33	40,000.00	28,375.00
4	Agriculture	0.00	30,750.00	48,233.33	76,066.67	120,000.00	47,977.50
5	Non Farm income	0.00	3,000.00	0.00	0.00	0.00	750.00
	Income(Rs.)	68,000.00	64,250.00	79,483.33	98,566.67	160,000.00	81,977.50

Average annual expenditure: The data regarding the average annual expenditure in Naregallu-3 micro-watershed is presented in Table 43. The results indicate that the average annual expenditure is Rs. 5,928.42. For landless households it was Rs. 4,200, for marginal farmers it was Rs. 4,682, for small farmers it was Rs. 5,366.67, for semi medium farmers it was Rs. 5,409.72 and for medium farmers it was Rs. 40,000.

Table 43. Average annual expenditure in Naregallu-3 micro-watershed

(Avg value in Rs.)

Sl.No.	Particulars	LL (5)	MF (10)	SF (12)	SMF (12)	MDF (1)	All (40)
1	Service/salary	0.00	20,000.00	0.00	20,000.00	0.00	1,000.00
2	Business	0.00	0.00	26,500.00	0.00	0.00	1,325.00
3	Wage	21,000.00	16,000.00	16,400.00	14,500.00	10,000.00	9,100.00
4	Agriculture	0.00	10,820.00	21,500.00	30,416.67	30,000.00	19,030.00
	Total	21,000.00	46,820.00	64,400.00	64,916.67	40,000.00	237,136.67
	Average	4,200.00	4,682.00	5,366.67	5,409.72	40,000.00	5,928.42

Horticulture species grown: The data regarding horticulture species grown in Naregallu-3 micro-watershed is presented in Table 44. The results indicate that, sampled households have grown 37 coconut tree trees in their field.

Table 44. Horticulture species grown in Naregallu-3 micro-watershed

Sl.No.	Particulars	MF (10)		SF (12)		SMF (12)		MDF (1)		All (40)	
		F	B	F	B	F	B	F	B	F	B
1	Coconut	0	0	2	0	35	0	0	0	37	0

*F= Field B=Back Yard

Forest species grown: The data regarding forest species grown in Naregallu-3 micro-watershed is presented in Table 45. The results indicate that, households have planted 1 teak 14 neem, 6 tamarind, 9 acacia and 4 banyan trees in their field.

Table 45: Forest species grown in Naregallu-3 micro-watershed

Sl.No.	Particulars	MF (10)		SF (12)		SMF (12)		MDF (1)		All (40)	
		F	B	F	B	F	B	F	B	F	B
1	Teak	0	0	1	1	0	0	0	0	1	1
2	Neem	6	0	5	0	3	0	0	0	14	0
3	Tamarind	4	0	0	0	2	0	0	0	6	0
4	Acacia	9	0	0	0	0	0	0	0	9	0
5	Banyan	1	0	1	0	2	0	0	0	4	0

*F= Field B=Back Yard

Marketing of the agricultural produce: The data regarding marketing of the agricultural produce in Naregallu-3 micro-watershed is presented in Table 46. The results indicated that, Bengal gram was sold to the extent 52.63 per cent, chilly was sold to the extent of 90 per cent, maize was sold to the extent of 96.01 per cent, paddy was sold to the extent of 52 per cent, sorghum was sold to the extent of 54.55 per cent and sunflower were sold to the extent of 100 per cent.

Table 46. Marketing of the agricultural produce in Naregallu-3 micro-watershed

Sl.No	Crops	Output obtained (q)	Output retained (q)	Output sold (q)	Output sold (%)	Avg. Price obtained (Rs/q)
1	Bengal gram	19	9	10	52.63	3450.0
2	Chilly	20	2	18	90.00	6200.0
3	Maize	953	38	915	96.01	1204.0
4	Paddy	50	24	26	52.00	1650.0
5	Sorghum	110	50	60	54.55	2116.67
6	Sunflower	65	0	65	100.00	5625.0

Marketing Channels used for sale of agricultural produce: The data regarding marketing channels used for sale of agricultural produce in Naregallu-3 micro-watershed is presented in Table 47. The results indicated that, about 2.50 per cent of the farmers sold their produce to local/village merchants, 82.50 per cent of the farmers sold their produce to regulated market and 12.50 per cent of them sold their produce through contract marketing arrangement.

Table 47. Marketing Channels used for sale of agricultural produce in Naregallu-3 micro-watershed

Sl.No.	Particulars	MF (10)		SF (12)		SMF (12)		MDF (1)		All (40)	
		N	%	N	%	N	%	N	%	N	%
1	Local/village Merchant	0	0.00	1	8.33	0	0.00	0	0.00	1	2.50
2	Regulated Market	10	100.00	9	75.00	13	108.33	1	100.00	33	82.50
3	Cooperative marketing Society	0	0.00	3	25.00	2	16.67	0	0.00	5	12.50

Mode of transport of agricultural produce: The data regarding mode of transport of agricultural produce in Naregallu-3 micro-watershed is presented in Table 48. The results indicated that, 95 per cent of the households used tractor and 2.50 per cent of them used flight as a mode of transportation for their agricultural produce.

Table 48. Mode of transport of agricultural produce in Naregallu-3 micro-watershed

Sl.No.	Particulars	LL (5)		MF (10)		SF (12)		SMF (12)		MDF (1)		All (40)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Tractor	0	0.00	10	100.00	12	100.00	15	125.00	1	100.00	38	95.00
2	Flight	0	0.00	0	0.00	1	8.33	0	0.00	0	0.00	1	2.50

Incidence of soil and water erosion problems: The data regarding incidence of soil and water erosion problems in Naregallu-3 micro-watershed is presented in Table 49. The results indicated that, 20 per cent of the households have experienced soil and water erosion problems in the farm.

Table 49. Incidence of soil and water erosion problems in Naregallu-3 micro-watershed

Sl. No.	Particulars	MF (10)		SF (12)		SMF (12)		MDF (1)		All (40)	
		N	%	N	%	N	%	N	%	N	%
1	Soil and water erosion problems in the farm	1	10.00	4	33.33	3	25.00	0	0.00	8	20.00

Interest shown towards soil testing: The data regarding Interest shown towards soil testing in Naregallu-3 micro-watershed is presented in Table 50. The results indicated that, 67.50 per cent have shown interest in soil test.

Table 50. Interest shown towards soil testing in Naregallu-3 micro-watershed

Sl.No.	Particulars	MF (10)		SF (12)		SMF (12)		MDF (1)		All (40)	
		N	%	N	%	N	%	N	%	N	%
1	Interest in soil test	7	70.00	9	75.00	10	83.33	1	100.00	27	67.50

Table 51. Usage pattern of fuel for domestic use in Naregallu-3 micro-watershed

Sl.No.	Particulars	LL (5)		MF (10)		SF (12)		SMF (12)		MDF (1)		All (40)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Fire Wood	4	80.00	9	90.00	12	100.00	12	100.00	1	100.00	38	95.00
2	LPG	0	0.00	1	10.00	0	0.00	0	0.00	0	0.00	1	2.50

Usage pattern of fuel for domestic use: The data regarding usage pattern of fuel for domestic use in Naregallu-3 micro-watershed is presented in Table 51. The results

indicated that, 95 per cent of the households used firewood and 2.5 per cent of the households used LPG as a source of fuel.

Source of drinking water: The data regarding source of drinking water in Naregallu-3 micro-watershed is presented in Table 52. The results indicated that, piped supply was the major source of drinking water for 45 per cent of the households and bore well was the source of drinking water for 55 per cent of the households in micro watershed.

Table 52. Source of drinking water in Naregallu-3 micro-watershed

Sl.No.	Particulars	LL (5)		MF (10)		SF (12)		SMF (12)		MDF (1)		All (40)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Piped supply	4	80.00	2	20.00	7	58.33	5	41.67	0	0.00	18	45.00
2	Bore Well	1	20.00	8	80.00	5	41.67	7	58.33	1	100.00	22	55.00

Source of light: The data regarding source of light in Naregallu-3 micro-watershed is presented in Table 53. The results indicated that, Electricity was the major source of light for 100 per cent of the households in micro watershed.

Table 53. Source of light in Naregallu-3 micro-watershed

Sl.No.	Particulars	LL (5)		MF (10)		SF (12)		SMF (12)		MDF (1)		All (40)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Electricity	5	100.00	10	100.00	12	100.00	12	100.00	1	100.00	40	100.00

Existence of Sanitary toilet facility: The data regarding existence of sanitary toilet facility in Naregallu-3 micro-watershed is presented in Table 54. The results indicated that, 42.50 per cent of the households possess sanitary toilet facility.

Table 54. Existence of Sanitary toilet facility in Naregallu-3 micro-watershed

Sl.No.	Particulars	LL (5)		MF (10)		SF (12)		SMF (12)		MDF (1)		All (40)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Sanitary toilet facility	2	40.00	4	40.00	6	50.00	4	33.33	1	100.00	17	42.50

Possession of PDS card: The data regarding possession of PDS card in Naregallu-3 micro-watershed is presented in Table 55. The results indicated that, 2.50 per cent of the sampled households possessed APL card and 95 per cent of the sampled households possessed BPL card.

Table 55. Possession of PDS card in Naregallu-3 micro-watershed

Sl.No.	Particulars	LL (5)		MF (10)		SF (12)		SMF (12)		MDF (1)		All (40)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	APL	0	0.00	1	10.00	0	0.00	0	0.00	0	0.00	1	2.50
2	BPL	5	100.00	9	90.00	12	100.00	11	91.67	1	100.00	38	95.00

Table 56. Participation in NREGA programme in Naregallu-3 micro-watershed

Sl. No.	Particulars	LL (5)		MF (10)		SF (12)		SMF (12)		MDF (1)		All (40)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Participation in NREGA programme	5	100	4	40.00	8	66.67	7	58.33	1	100	25	62.50

Participation in NREGA program: The data regarding participation in NREGA programme in Naregallu-3 micro-watershed is presented in Table 56. The results indicated that, 62.50 per cent of the households participated in NREGA programme.

Adequacy of food items: The data regarding adequacy of food items in Naregallu-3 micro-watershed is presented in Table 57. The results indicated that, cereals were adequate for 90 per cent of the households, pulses were adequate for 72.50 per cent, oilseeds were adequate for 17.50 per cent, vegetables and egg were adequate for 7.50 per cent, fruits were adequate for 12.50 per cent, milk was adequate for 5 per cent and meat were adequate for 2.50 per cent.

Table 57. Adequacy of food items in Naregallu-3 micro-watershed

Sl.No.	Particulars	LL (5)		MF (10)		SF (12)		SMF (12)		MDF (1)		All (40)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Cereals	1	20.00	10	100.00	12	100.00	12	100.00	1	100.00	36	90.00
2	Pulses	1	20.00	7	70.00	11	91.67	9	75.00	1	100.00	29	72.50
3	Oilseed	0	0.00	1	10.00	1	8.33	5	41.67	0	0.00	7	17.50
4	Vegetables	0	0.00	1	10.00	1	8.33	0	0.00	1	100.00	3	7.50
5	Fruits	2	40.00	1	10.00	1	8.33	1	8.33	0	0.00	5	12.50
6	Milk	0	0.00	0	0.00	1	8.33	1	8.33	0	0.00	2	5.00
7	Egg	1	20.00	0	0.00	1	8.33	1	8.33	0	0.00	3	7.50
8	Meat	1	20.00	0	0.00	0	0.00	0	0.00	0	0.00	1	2.50

Response on Inadequacy of food items: The data regarding inadequacy of food items in Naregallu-3 micro-watershed is presented in Table 58. The results indicated that, cereals were in adequate for 10 per cent of the households, pulses were inadequate for 27.50 per cent, oilseeds were inadequate for 65 per cent, vegetables were inadequate for 70 per cent, fruits were inadequate for 67.50 per cent, milk was inadequate for 82.50 per cent, eggs were inadequate for 90 per cent and meat was inadequate for 75 per cent of the households.

Table 58. Response on Inadequacy of food items in Naregallu-3 micro-watershed

Sl.No.	Particulars	LL (5)		MF (10)		SF (12)		SMF (12)		MDF (1)		All (40)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Cereals	4	80.00	0	0.00	0	0.00	0	0.00	0	0.00	4	10.00
2	Pulses	4	80.00	3	30.00	1	8.33	3	25.00	0	0.00	11	27.50
3	Oilseed	5	100.00	6	60.00	8	66.67	6	50.00	1	100.00	26	65.00
4	Vegetables	4	80.00	6	60.00	8	66.67	10	83.33	0	0.00	28	70.00
5	Fruits	4	80.00	6	60.00	8	66.67	8	66.67	1	100.00	27	67.50
6	Milk	3	60.00	9	90.00	10	83.33	10	83.33	1	100.00	33	82.50
7	Egg	4	80.00	10	100.00	11	91.67	10	83.33	1	100.00	36	90.00
8	Meat	4	80.00	7	70.00	9	75.00	9	75.00	1	100.00	30	75.00

Response on Market surplus of food items: The data regarding market surplus of food items in Naregallu-3 micro-watershed is presented in Table 59. The results indicated that, oilseeds were market surplus for 15 per cent of the households, vegetables were market

surplus for 20 per cent of the households, fruits were market surplus for 7.50 per cent of the households and meat were market surplus for 2.50 per cent of the households.

Table 59. Response on Market surplus of food items in Naregallu-3 micro-watershed

Sl.No.	Particulars	LL (5)		MF (10)		SF (12)		SMF (12)		MDF (1)		All (40)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Oilseed	0	0.00	3	30.00	2	16.67	1	8.33	0	0.00	6	15.00
2	Vegetables	0	0.00	3	30.00	2	16.67	3	25.00	0	0.00	8	20.00
3	Fruits	0	0.00	1	10.00	2	16.67	0	0.00	0	0.00	3	7.50
4	Meat	0	0.00	0	0.00	0	0.00	1	8.33	0	0.00	1	2.50

Farming constraints: The data regarding farming constraints experienced by households in Naregallu-3 micro-watershed is presented in Table 60. The results indicated that, lower fertility status of the soil was the constraint experienced by 72.50 per cent of the households, wild animal menace on farm field (75%), frequent incidence of pest and diseases (57.50%), inadequacy of irrigation water (45%), high cost of fertilizers and plant protection chemicals (30%), high rate of interest on credit and lack of marketing facilities in the area (20%), low price for the agricultural commodities (22.5%), inadequate extension services (10%), lack of transport for safe transport of the agricultural produce to the market (25%), less rainfall (60%) and Source of Agri-technology information (37.5%).

Table 60. Farming constraints Experienced in Naregallu-3 micro-watershed

Sl. No.	Particulars	MF (10)		SF (12)		SMF (12)		MDF (1)		All (40)	
		N	%	N	%	N	%	N	%	N	%
1	Lower fertility status of the soil	7	70	10	83.33	10	83.33	1	100	29	72.50
2	Wild animal menace on farm field	7	70	10	83.33	11	91.67	1	100	30	75
3	Frequent incidence of pest and diseases	8	80	8	66.67	7	58.33	0	0	23	57.50
4	Inadequacy of irrigation water	7	70	5	41.67	4	33.33	1	100	18	45
5	High cost of Fertilizers and plant protection chemicals	2	20	6	50	4	33.33	0	0	12	30
6	High rate of interest on credit	4	40	2	16.67	2	16.67	0	0	8	20
7	Low price for the agricultural commodities	3	30	2	16.67	2	16.67	1	100	9	22.50
8	Lack of marketing facilities in the area	1	10	3	25	4	33.33	0	0	8	20
9	Inadequate extension services	0	0	3	25	1	8.33	0	0	4	10
10	Lack of transport for safe transport of the Agril produce to the market.	1	10	5	41.67	3	25	1	100	10	25
11	Less rainfall	9	90	8	66.67	6	50	0	0	24	60
12	Source of Agri-technology information	4	40	5	41.67	6	50	0	0	15	37.50

SUMMARY

In order to assess the socio-economic condition of the farmers in the watershed a comprehensive questionnaire was prepared. Major components such as demographic conditions, migration details, food consumption and family expenditure pattern, material possession, land holding, land use management, cropping pattern, cost of cultivation of crops, livestock management. The statistical components such as frequency and percentage were used to analyze the data. About 40 households located in the micro watershed were interviewed for the survey.

The data indicated that there were 94 (56.29%) men and 73 (43.71%) women among the sampled households. The average family size of landless farmers' was 3, marginal farmers' was 4.2, small farmers' was 3.83, semi medium farmers' was 5 and medium farmers' was 4. The data indicated that, 18 (10.78%) people were in 0-15 years of age, 69 (41.32%) were in 16-35 years of age, 64 (38.32%) were in 36-60 years of age and 16 (9.58%) were above 61 years of age.

The results indicated that Naregallu-3 had 36.53 per cent illiterates, 35.33 per cent of them had primary school education, 4.19 per cent of them had middle school education, 13.77 per cent of them had high school education, 7.19 per cent of them had PUC education and 1.20 per cent had diploma and ITI.

The results indicate that, 97.5 per cent of household heads were practicing agriculture and 2.5 per cent of the household heads were agricultural labourers. The results indicate that agriculture was the major occupation for 23.95 per cent of the household members, 61.68 per cent were agricultural labourers and 14.37 per cent were student.

The results show that, 100 per cent of the population in the micro watershed has not participated in any local institutions. The results indicate that 5 per cent of the households possess thatched house and 95 per cent of the households possess katcha houses.

The results show that 70 per cent of the households possess TV, 17.5 per cent of them possess mixer/grinder, 7.50 per cent of them possess bicycle, 35 per cent of the households possess motor cycle, 2.50 per cent of them possess auto and landline phone and 90 per cent of the households possess mobile phones. The results show that the average value of television was Rs. 6,732, mixer grinder was Rs. 1,700, bicycle was 1,175, motor cycle was Rs. 32,357, auto was 300,000, landline was 4,000 and mobile phone was Rs. 2,492.

About 12.50 per cent of the households possess bullock cart, 17.50 per cent of them possess plough, 5 per cent of them possess seed/ fertilizer drill, 2.5 per cent of them possess sprayer, 12.50 per cent of them possess weeder, and 17.50 per cent of them

possess chaff cutter. The results show that the average value of bullock cart was Rs. 17,060, plough was Rs. 2800, seed/fertilizer drill was Rs. 8,000, sprayer was Rs. 2,000, weeder was Rs. 21 and the average value of chaff cutter was Rs. 441.

The results indicate that, 22.50 per cent of the households possess bullocks, 7.50 per cent of the households possess local cow, 2.50 per cent possess buffalo and goat.

The results indicate that, average own labour men available in the micro watershed was 1.40, average own labour (women) available was 1.85, average hired labour (men) available was 6.25 and average hired labour (women) available was 6.4. The results indicate that 100 per cent of the households opined that the hired labour was inadequate.

The results indicate that, households of the Naregallu-3 micro-watershed possess 29.15 ha (59.87%) of dry land and 19.54 ha (40.13%) of irrigated land. Marginal farmers possess 5.84 ha (86.61%) of dry land and 0.90 ha (13.39%) of irrigated land. Small farmers possess 14.26 ha (84.63%) of dry land and 2.59 ha (15.37%) of irrigated land. Semi medium farmers possess 9.05 ha (40.66%) of dry land and 13.21 ha (59.34%) of irrigated land. Medium farmers possess 2.83 ha (100%) of irrigated land.

The results indicate that, the average value of dry land was Rs. 264,006.11 and the average value of irrigated land was Rs. 450,207.13. In case of marginal famers, the average land value was Rs. 530,630.63 for dry land and Rs. 1,218,385.64 for irrigated land. In case of small famers, the average land value was Rs. 238,308.74 for dry land and Rs. 694,687.49 for irrigated land. In case of semi medium famers, the average land value was Rs. 132,498.88 for dry land and Rs. 400,949.47 for irrigated land. In case of medium farmers, the average land value was Rs. 211,714.29 for irrigated land.

The results indicate that, there were 11 functioning and 6 de-functioning bore wells in the micro watershed. The results indicate that, there was 1 functioning open well in the micro watershed.

The results indicate that, bore well was the major irrigation source in the micro water shed for 27.50 per cent of the farmers and open well was the major source of irrigation for 2.50 per cent of the farmers. The results indicate that, the depth of bore well was found to be 13.72 meters and the depth of open well was found to be 3.05 meters.

The results indicate that marginal, small and semi medium farmers had an irrigated area of 1.78 ha, 2.55 ha and 11.92 ha respectively. The results indicate that, farmers have grown maize (31.79 ha), sunflower (4.91 ha), sorghum (4.61 ha), Bengal gram (2.14 ha), paddy (1.62 ha), chilly and jowar (1.21 ha). Marginal farmers have grown maize and Bengal gram, while small farmers have grown maize, sunflower, sorghum, Bengal gram paddy and jowar. Semi medium farmers have grown maize, sunflower, sorghum, paddy and chilly. Medium farmers have grown maize. The results indicate that, the cropping intensity in Naregallu-3 micro-watershed was found to be 88.20 per cent.

The results indicate that, 55 per cent of the households have bank account and savings. The results indicate that, 55 per cent of the households have availed credit from different sources. The results indicate that, 81.82 per cent of the households have borrowed from commercial bank, 4.55 per cent of them borrowed from friends/ relatives, grameena bank and money lender, 9.09 per cent of the households borrowed from SHGs and CBOs. The results indicate that, the average credit amount borrowed by households in micro-watershed was Rs, 59,409.09. The results indicate that, 100 per cent of the households borrowed from institutional sources for the purpose of agricultural production. The results indicate that, 100 per cent of the household's availed credit for the purpose of agricultural production. The results indicated that 100 per cent of the households did not repay their loan borrowed from institutional sources. The results indicated that 100 per cent of the households did not repay their loan borrowed from private sources. The results indicate that, around 100 per cent opined that the loan amount borrowed from institutional sources helped to perform timely agricultural operations. The results indicate that, 100 per cent of the households opined that the credit borrowed from private credit helped to perform timely agricultural operations.

The results indicate that, the total cost of cultivation for maize was Rs. 33037.55. The gross income realized by the farmers was Rs. 45231.00. The net income from maize cultivation was Rs. 12193.46. Thus the benefit cost ratio was found to be 1:1.37. The results indicate that, the total cost of cultivation for chilly was Rs. 57632.23. The gross income realized by the farmers was Rs. 89969.75. The net income from chilly cultivation was Rs. 32337.52. Thus the benefit cost ratio was found to be 1:1.56. The results indicate that, the total cost of cultivation for chilly was Rs. 57632.23. The gross income realized by the farmers was Rs. 89969.75. The net income from chilly cultivation was Rs. 32337.52. Thus the benefit cost ratio was found to be 1:1.56. The results indicate that, the total cost of cultivation for sunflower was Rs. 28263.78. The gross income realized by the farmers was Rs. 88272.38. The net income from sunflower cultivation was Rs. 60008.60. Thus the benefit cost ratio was found to be 1:3.12. The results indicate that, the total cost of cultivation for sunflower was Rs. 28263.78. The gross income realized by the farmers was Rs. 88272.38. The net income from sunflower cultivation was Rs. 60008.60. Thus the benefit cost ratio was found to be 1:3.12.

The results indicate that, 7.50 per cent of the households opined that dry fodder was adequate, green fodder was adequate for 2.50 per cent of the households and 20 per cent of the households opined that dry fodder was in adequate.

The results indicate that the annual gross income was Rs. 68,000 for landless farmers, for marginal farmers it was Rs.64,250, for small farmers it was Rs.79,483.33, for

semi medium farmers it was Rs. 98,566.67 and for medium farmers it was Rs. 160,000. The results indicate that the average annual expenditure is Rs. 5,928.42. For landless households it was Rs. 4,200, for marginal farmers it was Rs. 4,682, for small farmers it was Rs. 5,366.67, for semi medium farmers it was Rs. 5,409.72 and for medium farmers it was Rs. 40,000.

The results indicate that, sampled households have grown 37 coconut tree trees in their field. The results indicate that, households have planted 1 teak 14 neem, 6 tamarind, 9 acacia and 4 banyan trees in their field.

The results indicated that, Bengal gram was sold to the extent 52.63 per cent, chilly was sold to the extent of 90 per cent, maize was sold to the extent of 96.01 per cent, paddy was sold to the extent of 52 per cent, sorghum was sold to the extent of 54.55 per cent and sunflower were sold to the extent of 100 per cent.

The results indicated that, about 2.50 per cent of the farmers sold their produce to local/village merchants, 82.50 per cent of the farmers sold their produce to regulated market and 12.50 per cent of them sold their produce through contract marketing arrangement. The results indicated that, 95 per cent of the households used tractor and 2.50 per cent of them used flight as a mode of transportation for their agricultural produce.

The results indicated that, 20 per cent of the households have experienced soil and water erosion problems in the farm. The results indicated that, 67.50 per cent have shown interest in soil test.

The results indicated that, 95 per cent of the households used firewood and 2.5 per cent of the households used LPG as a source of fuel. The results indicated that, piped supply was the major source of drinking water for 45 per cent of the households and bore well was the source of drinking water for 55 per cent of the households in micro watershed.

Electricity was the major source of light for 100 per cent of the households in micro watershed. The results indicated that, 42.50 per cent of the households possess sanitary toilet facility. The results indicated that, 2.50 per cent of the sampled households possessed APL card and 95 per cent of the sampled households possessed BPL card. The results indicated that, 62.50 per cent of the households participated in NREGA programme.

The results indicated that, cereals were adequate for 90 per cent of the households, pulses were adequate for 72.50 per cent, oilseeds were adequate for 17.50 per cent, vegetables and egg were adequate for 7.50 per cent, fruits were adequate for 12.50 per cent, milk was adequate for 5 per cent and meat were adequate for 2.50 per cent.

The results indicated that, cereals were in adequate for 10 per cent of the households, pulses were inadequate for 27.50 per cent, oilseeds were inadequate for 65 per cent, vegetables were inadequate for 70 per cent, fruits were inadequate for 67.50 per cent, milk was inadequate for 82.50 per cent, eggs were inadequate for 90 per cent and meat was inadequate for 75 per cent of the households.

The results indicated that, oilseeds were market surplus for 15 per cent of the households, vegetables were market surplus for 20 per cent of the households, fruits were market surplus for 7.50 per cent of the households and meat were market surplus for 2.50 per cent of the households.

The results indicated that, lower fertility status of the soil was the constraint experienced by 72.50 per cent of the households, wild animal menace on farm field (75%), frequent incidence of pest and diseases (57.50%), inadequacy of irrigation water (45%), high cost of fertilizers and plant protection chemicals (30%), high rate of interest on credit and lack of marketing facilities in the area (20%), low price for the agricultural commodities (22.5%), inadequate extension services (10%), lack of transport for safe transport of the agricultural produce to the market (25%), less rainfall (60%) and Source of Agri-technology information (37.5%).