

**State: BIHAR**

**Agriculture Contingency Plan for District: KAIMUR (Bhabhua)**

<b>1.0 District Agriculture profile</b>				
<b>1.1</b>	<b>Agro-Climatic/Ecological Zone</b>			
	Agro Ecological Sub Region (ICAR)	Northern Plain, Hot Subhumib (Dry) Eco-Region (9.2)		
	Agro-Climatic Zone (Planning Commission)	Middle Gangetic Plain Region (IV)		
	Agro Climatic Zone (NARP)	South Bihar Alluvial Plain Zone (BI-3)		
	List all the districts falling under the NARP Zone>(*>50% area falling in the zone)	Zone-III (Rohtas, Bhojpur, Buxer, Bhabua, Arwal, Patna, Nalanda, Nawada, Sheikhpura, Jahanabad, Aurangabad, Gaya, Munger, Bhagalpur, Banka, Jamui, Lakhisarai)		
	Geographic coordinates of district headquarters	Latitude	Longitude	Altitude
		25-26 <sup>0</sup> N	83-84 <sup>0</sup> E	1800 ft
	Name and address of the concerned ZRS/ ZARS/ RARS/ RRS/ RRTTS	ZARS, Irrigation Research Centre, Bikramganj, Dist.-Rohtas		
	Mention the KVK located in the district with address	Vanvasi Krishi Vigyan Kendra, Vill.+P.O.-Adhaura, Distt.-Kaimur, Bihar, Pin-821116		
Name and address of the nearest Agromet Field Unit (AMFU, IMD) for agro-advisories in the Zone	Rajendra Agricultural University, Pusa, Samastipur			

<b>1.2</b>	<b>Rainfall</b>	<b>Normal RF(mm)</b>	<b>Normal Onset</b>	<b>Normal Cessation</b>
	SW monsoon (June-Sep)	1004.4	3 <sup>rd</sup> week of July	3 <sup>rd</sup> week of September
	NE Monsoon(Oct-Dec)	67.8	1 <sup>st</sup> week of October	
	Winter (Jan-Feb)	46.2		
	Summer (March -May)	45.9		
	Annual	1164.3		

<b>1.3</b>	<b>Land use pattern of the district</b> (latest statistics)	Geographical area	Cultivable area	Forest area	Land under non-agricultural use	Permanent pastures	Cultivable wasteland	Land under Misc. tree crops and groves	Barren and uncultivable land	Current fallows	Other fallows
	Area ('000 ha)	340.4	176.7	96.1	33.2	3.8	14.7	1.5	2.6	8.3	3.1

<b>1.4</b>	<b>Major Soils</b>	<b>Area ('000 ha)</b>	<b>Percent (%) of total</b>
	1. Clay loam soils	80.5	45.6
	2. Sandy loam soils	54.5	30.8
	3. Red Laterite soils	41.6	23.5

<b>1.5</b>	<b>Agricultural land use</b>	<b>Area ('000 ha)</b>	<b>Cropping intensity %</b>
	Net sown area	176.7	136.4
	Area sown more than once	64.3	
	Gross cropped area	241.0	

<b>1.6</b>	<b>Irrigation</b>	<b>Area ('000 ha)</b>		
	Net irrigated area	71.7		
	Gross irrigated area	81.6		
	Rainfed area	95.1		
	<b>Sources of Irrigation</b>	<b>Number</b>	<b>Area ('000 ha)</b>	<b>Percentage of total irrigated area</b>
	Canals	01	55.7	68.3
	Tanks	270	1.0	1.3
	Open wells	6505	0.058	0.07
	Bore wells	18095	9.8	12.1
	Lift irrigation schemes	34	1.01	1.23

	Micro-irrigation	22	0.7	0.85
	Other sources (please specify)		2.45	3.0
	Total Irrigated Area			
	Pump sets	13180		
	No. of Tractors	2275		
	<b>Groundwater availability and use* (Data source: State/Central Ground water Department /Board)</b>	No. of blocks/ Tehsils	(%) area	Quality of water (specify the problem such as high levels of arsenic, fluoride, saline etc)
	Over exploited			
	Critical			
	Semi- critical	06		
	Safe	05		
	Wastewater availability and use			
	Ground water quality			
*over-exploited: groundwater utilization > 100%; critical: 90-100%; semi-critical: 70-90%; safe: <70%				

**1.7 Area under major field crops & horticulture (as per figures of 2009-10)**

1.7	Major field crops cultivated	Area ('000 ha)							
		<i>Kharif</i>			<i>Rabi</i>			Summer	Grand total
		Irrigated	Rainfed	Total	Irrigated	Rainfed	Total		
	Rice			81.7					81.7
	Wheat						68.4		68.4
	Lentil						4.8		4.8
	Chickpea						6.4		6.4
	Linseed						1.4		1.4
	Mustard						1.4		1.4
	Maize			0.079					0.079
	Greengram							0.75	0.75

	Horticulture crops - Fruits	Area ('000 ha) 2010-11		
		Total	Irrigated	Rainfed
	Mango	3.37		
	Guava	1.36		
	Aonla	0.04		

	Lemon	0.27		
	Banana	0.21		
	<b>Horticulture crops - Vegetables</b>	<b>Total</b>	<b>Irrigated</b>	<b>Rainfed</b>
	Potato	4.18		
	Onion	0.88		
	Tomato	0.58		
	Cauliflower	0.77		
	Cabbage	0.44		
	Brinjal	0.69		
	Okra	0.85		
	Chilli	0.40		
	<b>Medicinal and Aromatic crops</b>	<b>Total (year 2009-10)</b>	<b>Irrigated</b>	<b>Rainfed</b>
	Japanese Mint	0.005	0.005	
	Satawar	0.012	0.010	0.002
	Tuberose	0.003	0.003	
	<b>Plantation crops</b>	<b>Total</b>	<b>Irrigated</b>	<b>Rainfed</b>
	Siris	50.5		50.5
	<b>Fodder crops</b>	<b>Total</b>	<b>Irrigated</b>	<b>Rainfed</b>
	Berseem	1.050	1.050	
	Oat	0.005	0.005	
	<b>Total fodder crop area</b>			
	<b>Grazing land</b>	3.8	0.029	3.829
	<b>Sericulture etc</b>	0.001		0.001

<b>1.8</b>	<b>Livestock</b>	<b>Male ('000)</b>	<b>Female ('000)</b>	<b>Total ('000)</b>
	Non descriptive Cattle (local low yielding)	130.5	64.6	195.1
	Improved cattle	5	6	11
	Crossbred cattle	0.042	3.5	3.5
	Non descriptive Buffaloes (local low yielding)	5	80	85
	Descript Buffaloes		25	25

	Goat	40	50	90			
	Sheep	18	17.5	35.5			
	Others (Camel, Pig, Yak etc.)	4	5	9			
	Commercial dairy farms (Number)			0.035			
<b>1.9</b>	<b>Poultry</b>	<b>No. of farms</b>	<b>Total No. of birds ('000)</b>				
	Commercial	150	125				
	Backyard	1500	15				
<b>1.10</b>	<b>Fisheries</b> (Data source: Chief Planning Officer)						
	<b>A. Capture</b>						
	<b>i) Marine</b> (Data Source: Fisheries Department)	<b>No. of fishermen</b>	<b>Boats</b>		<b>Nets</b>		<b>Storage facilities (Ice plants etc.)</b>
			Mechanized	Non-mechanized	Mechanized (Trawl nets, Gill nets)	Non-mechanized (Shore Seines, Stake & trap nets)	
	<b>ii) Inland</b> (Data Source: Fisheries Department)	<b>No. Farmer owned ponds</b>		<b>No. of Reservoirs</b>		<b>No. of village tanks</b>	
		300		40		40	
	<b>B. Culture</b>						
			<b>Water Spread Area (ha)</b>	<b>Yield (t/ha)</b>	<b>Production ('000 tons)</b>		
	<b>i) Brackish water</b> (Data Source: MPEDA/ Fisheries Department)						
	<b>ii) Fresh water</b> (Data Source: Fisheries Department)		2469	3.2	2469		

### 1.11 Production and Productivity of major crops (2006 - 2010)

1.11	Name of crop	Kharif		Rabi		Summer		Total		Crop residue as fodder ('000 tons)
		Production ('000 t)	Productivity (kg/ha)	Production ('000 t)	Productivity (kg/ha)	Production ('000 t)	Productivity (kg/ha)	Production ('000 t)	Productivity (kg/ha)	
<b>Major Field crops (Crops identified based on total acreage)</b>										
	Rice	382.9	3450					382.9	3450	400

	Wheat			199.7	2570			199.7	2570	200
	Lentil			7.9	885			7.9	885	8
	Chickpea			9.2	990			9.2	990	9
	Linseed			3.7	775			3.7	775	
<b>Major Horticultural crops (Crops identified based on total acreage)</b>										
	Potato			115.6	18500			115.6	18500	
	Tomato	39.9	10500					39.9	10500	
	Brinjal	24.8	10800					24.8	10800	
	Okra	15.1	10100					15.1	10100	
	Cucurbits	10.3	10300					10.3	10300	

<b>1.12</b>	<b>Sowing window for 5 major field crops</b> (start and end of normal sowing period)	<b>Rice</b>	<b>Wheat</b>	<b>Chickpea</b>	<b>Linseed</b>	<b>Potato</b>
	Kharif- Rainfed	4 <sup>th</sup> week of June – 2 <sup>nd</sup> week of July	-	-	-	-
	Kharif-Irrigated	4 <sup>th</sup> week of May – 3 <sup>rd</sup> week of June	-	-	-	-
	Rabi- Rainfed	-	2 <sup>nd</sup> week of October - 4 <sup>th</sup> week of October	2 <sup>nd</sup> week of October - 4 <sup>th</sup> week of October	2 <sup>nd</sup> week of October - 3 <sup>rd</sup> week of October	-
	Rabi-Irrigated	-	2 <sup>nd</sup> week of November- 2 <sup>nd</sup> week of December	4 <sup>th</sup> week of October-2 <sup>nd</sup> week of November	3 <sup>rd</sup> week of October - 2 <sup>nd</sup> week of November	4 <sup>th</sup> week of October - 2 <sup>nd</sup> week of November

<b>1.13</b>	<b>What is the major contingency the district is prone to? (Tick mark)</b>	<b>Regular</b>	<b>Occasional</b>	<b>None</b>
	Drought	√		
	Flood		√	
	Cyclone			√
	Hail storm			√
	Heat wave		√	
	Cold wave		√	
Frost		√		

	Sea water intrusion			√
	Pests and disease outbreak		√	

<b>1.14</b>	<b>Include Digital maps of the district for</b>	Location map of district within State as Annexure I	Enclosed : Yes
		Mean annual rainfall as Annexure 2	Enclosed: Yes
		Soil map as Annexure 3	Enclosed : Yes

## ANNEXURE-I

### Agro climatic Zones of Bihar




Source: [krishi.bih.nic.in](http://krishi.bih.nic.in)




### Annexure-II

Mean annual rainfall (mm)



### Annexure-III



Source: NBSS&LUP, Kolkata

## 2.0 Strategies for weather related contingencies

### 2.1 Drought

#### 2.1.1 Rainfed situation

Condition	Major Farming situation	Normal Crop / Cropping system	Suggested Contingency measures		
			Change in crop / cropping system including variety	Agronomic measures	Remarks on Implementation
Early season drought (delayed onset) Delay by 2 weeks 1 <sup>st</sup> week of August	Upland Shallow red soils	Pigeonpea-Fallow	Pigeonpea-ML-13, Narendra Arhar-1, P-9	-	-
		Fallow-Wheat/ Chickpea Blackgram/ Greengram/ Lentil /Linseed	Blackgram Blackgram-Pant Urd-19, Narendra Urd	-	
		Rice-Lentil/Chickpea	Early Rice-Lentil/ Chickpea Prefer medium to long duration varieties	<ul style="list-style-type: none"> <li>• Adopt normal package of practices</li> <li>• Direct seeding of drought tolerant varieties in dry soil in June/ July with pre emergence herbicide application under sufficient soil moisture conditions.</li> </ul>	
	Medium land Fine loamy soils	Rice-Wheat	Rice-Wheat Rice-Rajendra Bhagwati, Sugandha-2/3, Sita, Sarju-52, PRH-10, P-6444, Wheat-K-307, CBW-38, PBW-343	<ul style="list-style-type: none"> <li>• Raise staggered community nursery preferably with medium duration varieties in mid and lowlands</li> </ul>	
	Lowland Clay loamy soils	Rice-Wheat	Rice-Wheat Rice-R. Mahsoori-1, MTU-7029, Sugandha		

Condition	Major Farming situation	Normal Crop/cropping system	Suggested Contingency measures		
			Change in crop/cropping system	Agronomic measures	Remarks on Implementation
Early season drought (delayed onset) Delay by 4 weeks	Upland Shallow red soils	Rice-Wheat	Short duration Rice-Wheat Rice-Turanta, vandana,	<ul style="list-style-type: none"> <li>• Direct seeding of Rice</li> <li>• Application of fertilizers</li> </ul>	Seed from KVK, Adhaura, RAU,

3 <sup>rd</sup> week of August			Prabhat, Pusa 2-21,	especially phosphorous and potash to be ensured under late transplanted conditions	Pusa, BRBN, BHU, NSC
		Fallow-Wheat/ Lentil/ Chickpea /Mustard/ Linseed	Dhaincha (Green manuring)-Wheat/ Lentil/Chickpea/ Mustard/ Linseed  Dhaincha-Punjab Dhaicha-1, Jahirabad-2, Local		
	Medium land Fine loamy soils	Rice-Wheat	Maize-Wheat Maize-Shaktiman-1,2,3,4, 5Suwan, Devaki, Ganga-11	- Apply full basal dose of NPK - Planting of Maize through dibbling method	
		Rice-Wheat	Short duration Rice-Wheat Short duration Rice- R.Bhagwati, Prabhat, PRH-10, R. Subhasini	<ul style="list-style-type: none"> <li>• Mat nursery (dapog method)/ Community nursery can be raised for quick availability of young seedlings for transplanting of medium duration varieties by first fortnight of August</li> <li>• Direct seedling of Rice</li> <li>• Raise staggered community nursery preferably with medium duration varieties in mid and lowlands</li> </ul>	
	Lowland Clay loamy soils	Rice-Wheat	Medium duration Rice-Wheat Rice-R. Sweta, Sita, Rajshree		

Condition	Major Farming situation	Normal Crop/cropping system	Suggested Contingency measures		
			Change in crop/cropping system	Agronomic measures	Remarks on Implementation
Early season drought (delayed onset)	Upland	Fallow-Lenti/1 Chickpea	Niger-Lentil	Balanced use of fertilizer	Seed from KVK,

1 <sup>st</sup> week of September	Shallow red soils	/Linseed /Mustard	Niger-JNC-1, JNC-6, BNS-1 & BNS-2		Adhaura, RAU, Pusa, BRBN, BHU, NSC
		Pigeonpea-Fallow	Pigeonpea-Fallow Pigeonpea-M-13, P-9	Adopt seed rate @20 Kg/ha	
	Medium land Fine loamy soils	Rice-Wheat	Rice-Wheat Rice-Hira, Turanta	<ul style="list-style-type: none"> <li>• Direct seeding of Rice</li> <li>• Application of fertilizers especially phosphorous and potash to be ensured under late transplanted conditions</li> <li>• Life saving irrigation</li> </ul>	
		Rice-Wheat	Blackgram/ Greengram- Wheat Blackgram -Pant U-19&31, T-9 Greengram -HUM-12, HUM16	-	
		Rice-Wheat	Tomato-Wheat Tomato-Kahsi Amrit, Swarna Vaibhav, Swarn Lalima, DVRT-2	-	
Lowland Clay loamy soils	Rice-Wheat	Short duration Rice-Wheat Rice-Hira, Turanta, Vandana	<ul style="list-style-type: none"> <li>• Application of organic manure and vermi compost initially for Rice and other crops.</li> <li>• Fodder varieties of Jowar, Maize, Bajra in combination with legumes (cowpea and horsegram) can be taken up wherever feasible to meet the fodder requirements in deficit rainfall districts</li> </ul>		

Condition	Major Farming situation	Normal Crop/cropping system	Suggested Contingency measures		
			Change in crop/cropping system	Agronomic measures	Remarks on Implementation
Early season drought (delayed onset)	Upland Shallow red soils	Fallow-Wheat	Blackgram-Mustard Blackgram -T-9, Pant U-19, Pant	-Life saving irrigation	Seed from KVK, Adhaura, RAU,

3 <sup>rd</sup> week of September			Urd-31, Naveen	- Light irrigation at critical stage	Pusa, BRBN, BHU, NSC
		Fallow-Lentil/ Chickpea	Blackgram/ Greengram-Lentil/Chickpea Blackgram -T-9, Pant U-19, Pant Urd-31		
	Medium land Fine loamy soils	Fallow-Wheat	Sep. Pigeonpea-Fallow Sep. Pigeon pea- P-9, Sharad, Narendra Arhar-1		
		Fallow-Lentil/ Chickpea	Blackgram/ Greengram-Lentil/Chickpea Blackgram -T-9, Pant U-19, Pant Urd-31		
		Fallow-Wheat	<b>Tomato-Maize</b> Tomato-Kahsi Amrit, Swarna Vaibhav, Swarn Lalima, DVRT-2		

Condition	Major Farming situation	Normal Crop/cropping system	Suggested Contingency measures		
			Crop management	Soil nutrient & moisture conservation measures	Remarks on Implementation
Early season drought (Normal onset)					
Normal onset followed by 15-20 days dry spell after sowing leading to poor germination/ crop stand etc.	Upland Shallow red soils	Rice-Lentil/Chickpea	<ul style="list-style-type: none"> <li>• Re-transplanting of short duration Rice in case of heavy loss</li> <li>• Gap filling</li> <li>• Life saving irrigation</li> </ul>	<ul style="list-style-type: none"> <li>• Mulching through mechanical weeding</li> </ul>	Seed from RAU Pusa, NSC, TDC
	Medium land Fine loamy soils	Rice-Wheat			
	Lowland Clay loamy soils	Rice-Wheat			

Condition	Major Farming	Normal Crop/cropping	Suggested Contingency measures		
			Crop management	Soil nutrient &	Remarks on
Mid season					

<b>drought (long dry spell, consecutive 2 weeks rainless (&gt;2.5 mm) period)</b>	<b>situation</b>	<b>system</b>		<b>moisture conservation measues</b>	<b>Implementation</b>
At vegetative stage	Upland Shallow red soils	Rice-Lentil/Chickpea	<ul style="list-style-type: none"> <li>• Gap filling</li> <li>• Postpone the top dressing</li> </ul>	<ul style="list-style-type: none"> <li>• Mulching</li> <li>• Life saving irrigation</li> </ul>	
	Lowland Clay loamy soils	Rice-Wheat			

<b>Condition</b>			<b>Suggested Contingency measures</b>		
<b>Mid season drought (long dry spell)</b>	<b>Major Farming situation</b>	<b>Normal Crop/cropping system</b>	<b>Crop management</b>	<b>Soil nutrient &amp; moisture conservation measues</b>	<b>Remarks on Implementation</b>
At flowering/ fruiting stage	Upland Shallow red soils	Rice-Lentil/Gram/Linseed	-	<ul style="list-style-type: none"> <li>• Mulching</li> <li>• Life saving irrigation</li> </ul>	
	Midland Fine loamy soils	Rice-Wheat	-		
	Lowland Clay loamy soils	Rice-Wheat	-		
<b>Condition</b>			<b>Suggested Contingency measures</b>		
<b>Terminal drought (Early withdrawal of monsoon)</b>	<b>Major Farming situation</b>	<b>Normal Crop/cropping system</b>	<b>Crop management</b>	<b>Rabi Crop planning</b>	<b>Remarks on Implementation</b>
	Upland Shallow red soils	Rice-Lentil/Chickpea	<ul style="list-style-type: none"> <li>• Mulching</li> <li>• Foliar application of 2% MOP</li> </ul>	<ul style="list-style-type: none"> <li>• Open the Furrow during evening and left the furrow open overnight &amp; plank in the next morning before sunrise for growing of early Rabi crops like Toria, Lentil &amp; Chickpea</li> </ul>	
	Midland Fine loamy soils	Rice-Wheat			
	Lowland Clay loamy soils	Rice-Wheat			

### 2.1.2 Drought - Irrigated situation

Condition			Suggested Contingency measures		
	Major Farming situation	Normal Crop/cropping system	Change in crop/cropping system	Agronomic measures	Remarks on Implementation
Delayed release of water in canals due to low rainfall	Midland Fine loamy soils	Rice-Wheat	Rice-Wheat Medium duration Rice: R. Bhagwati, R. Sweta, Sarju-52	<ul style="list-style-type: none"> <li>Life saving irrigation</li> <li>Mulching</li> </ul>	
	Lowland Clay loamy soils	Rice-Wheat			

Condition			Suggested Contingency measures		
	Major Farming situation	Normal Crop/cropping system	Change in crop/cropping system	Agronomic measures	Remarks on Implementation
Limited release of water in canals due to low rainfall	Midland Fine loamy soils	Rice-Wheat	Rice-Wheat Medium duration Rice: R. Bhagwati, R. Sweta, Sarju-52	<ul style="list-style-type: none"> <li>Life saving irrigation</li> <li>Mulching</li> </ul>	
	Lowland Clay loamy soils	Rice-Wheat			

Condition			Suggested Contingency measures		
	Major Farming situation	Normal Crop/cropping system	Change in crop/cropping system	Agronomic measures	Remarks on Implementation
Non release of water in canals under delayed onset of monsoon in catchment	Midland& Lowland	Rice-Wheat	Blackgram/Greengram-Wheat  Black gram:T-9, Pant U-19, Pant U-30 Green gram :HUM-12, HUM-16	Life saving irrigation	

Condition			Suggested Contingency measures		
	Major Farming situation	Normal Crop/cropping system	Change in crop/cropping system	Agronomic measures	Remarks on Implementation



Condition	Suggested Contingency measures			
	Major Farming situation	Normal Crop/cropping system	Change in crop/cropping system	Agronomic measures
Lack of inflows into tanks due to insufficient /delayed onset of monsoon	Not Applicable			

Condition	Suggested Contingency measures				
	Major Farming situation	Normal Crop/cropping system	Change in crop/cropping system	Agronomic measures	Remarks on Implementation
Insufficient groundwater recharge due to low rainfall		Rice-Wheat	Pigeonpea-Fallow Pigeonpea: P-9, Mal-13, N. Pigeon pea-1	Sprinkler irrigation system	
		Rice-Wheat	Short duration Rice-Mustard Rice-Turanta, Vandana, Prabhat	<ul style="list-style-type: none"> <li>Organic manure &amp; Vermi-compost application</li> <li>Direct seeding of rice</li> <li>Sprinkler irrigation system</li> </ul>	
			Short duration Rice-Lentil  Rice-Turanta, Vandana, Prabhat		

## 2.2 Unusual rains (untimely, unseasonal etc) (for both rainfed and irrigated situations)

Condition	Suggested contingency measure			
	Vegetative stage	Flowering stage	Crop maturity stage	Post harvest
Continuous high rainfall in a short span leading to water logging				
Rice	<ul style="list-style-type: none"> <li>Drainage management</li> <li>Gap filling</li> </ul>	<ul style="list-style-type: none"> <li>Drainage management</li> <li>Subsequently crop if totally damaged i.e. Toria</li> </ul>	<ul style="list-style-type: none"> <li>Drainage management</li> <li>Subsequent crop if totally damaged</li> <li>Harvest at physiological maturity</li> </ul>	<ul style="list-style-type: none"> <li>i) Storage at safer place</li> <li>ii) Moisture level should be 9-10%</li> </ul>
Pigeonpea	<ul style="list-style-type: none"> <li>Drainage management</li> <li>September sowing if Kharif</li> </ul>	<ul style="list-style-type: none"> <li>Drainage management</li> <li>Alternative Rabi crops (OLS &amp;</li> </ul>	<ul style="list-style-type: none"> <li>Drainage management</li> <li>Harvest at physiological</li> </ul>	Storage at safer place

	Pigeonpea is completely damaged • Gap filling if needed	PLS) if totally damaged	maturity	
<b>Heavy rainfall with high speed winds in a short span<sup>2</sup></b>				
Rice	• Drainage management • Gap filling if needed	• Drainage management • Subsequent crop if totally damaged i.e. Toria	• Drainage management • Subsequent crop if totally damaged	Storage at safer place
Pigeonpea	• Drainage management • September sowing if Kharif Arhar is completely damaged • Gap filling if needed	• Drainage management • Alternative Rabi crops (OLS & PLS) if totally damaged	• Drainage management • Harvest at physiological maturity	Storage at safer place
<b>Outbreak of pests and diseases due to unseasonal rains</b>				
Rice	❖ Seedling treatment with granular insecticide – Cartap hydrochloride or phorate 10G or carbofuran 3G. ❖ Maintain shallow water in nursery beds ❖ Providing good drainage.	❖ Use copper fungicides against Bacterial leaf blight. ❖ Split application of N fertilizer (3-4 times)	❖ Harvest at physiological maturity	Proper drying and safe storage
Pigeonpea	Pigeonpea	❖ Provide drainage ❖ Seed treatment with 1 g carbendizim +2g thiram/kg seed.	Provide drainage	Provide drainage

### 2.3 Floods

Condition	Suggested contingency measure <sup>o</sup>			
	Seedling / nursery stage	Vegetative stage	Reproductive stage	At harvest
Transient water logging/partial inundation <sup>1</sup>				
Rice	• Drainage management	• Drainage management	• Drainage management	Storage at safer place

	<ul style="list-style-type: none"> <li>• Re transplanting through Dapog nursery if completely damaged</li> <li>• Gap filling</li> </ul>	<ul style="list-style-type: none"> <li>• Alternative crops if totally damaged</li> <li>• Gap filling</li> <li>• 40-45 days old seedlings may be used for transplanting</li> <li>• Kharuhan (double transplanting)</li> </ul>	<ul style="list-style-type: none"> <li>• Harvest at physiological maturity</li> <li>• Lentil as paira crop can be taken</li> </ul>	
<b>Continuous submergence for more than 2 days<sup>2</sup></b>				
Rice	<ul style="list-style-type: none"> <li>• Gap filling, if needed</li> <li>• Re-sowing if damaged after receding of flood</li> </ul>	<ul style="list-style-type: none"> <li>• Replanting through Kharuhan (double transplanting) by 3-4 seedlings per hill</li> <li>• Short duration rice variety</li> </ul>	<ul style="list-style-type: none"> <li>• Toria/Late wheat if completely damaged</li> </ul>	Storage at safer place

#### 2.4 Extreme events: Heat wave / Cold wave/Frost/ Hailstorm /Cyclone

Extreme event type	Suggested contingency measure <sup>r</sup>			
	Seedling / nursery stage	Vegetative stage	Reproductive stage	At harvest
<b>Heat wave</b>				
Pigeonpea			Provide irrigation	
Wheat			Provide irrigation (Terminal heat)	
<b>Horticulture</b>				
Mango	Provide irrigation	Provide irrigation	Provide irrigation	
Papaya	Provide irrigation	Provide irrigation	Provide irrigation	
<b>Cold wave</b>				
Wheat		Provide irrigation, , mulching		
Mustard		Provide irrigation, , mulching		
Pulses		Provide irrigation, , mulching		
<b>Frost</b>				
Wheat		Provide irrigation, , mulching		
Pigeonpea		Provide irrigation, , mulching		
Lentil		Provide irrigation, , mulching		
<b>Horticulture</b>				
Tomato & Potato		Earth up to 15cm ht. Provide		Harvest in dry

		irrigation, , mulching		weather
<b>Hailstorm</b>	Not Applicable			
<b>Cyclone</b>	Not Applicable			

## 2.5 Contingent strategies for Livestock, Poultry & Fisheries

### 2.5.1 Livestock

	Suggested contingency measures		
	Before the event <sup>s</sup>	During the event	After the event
<b>Drought</b>			
Feed and fodder availability	<ol style="list-style-type: none"> <li>1. Cultivation of fodder tree</li> <li>2. Storage of Improved Quality Fodder</li> <li>3. Conservation &amp; Storage of <ul style="list-style-type: none"> <li>• Feed &amp; Fodder</li> <li>• Hay &amp; Silage: — Preserve the fodder in the form of hay from Berseem &amp; other grasses as well as silage from (a) Maize- harvesting at well developed cob. (b) Jowar - at flowering stage. (c) Oat (d) Hybrid Napier – 40-45 day old. (e) Potato leaves mixing with wheat straw in ratio of 7:1 and should be supplemented with 3% molasses.</li> </ul>           Hay: – <ul style="list-style-type: none"> <li>• Berseem/Lucerne and other grasses.</li> <li>• Bales of hay and other dry fodder should be stored in dry places and covered with asbestos sheet or polythene sheet.</li> </ul> </li> <li>4. Development &amp; storage of: – (a) Complete Feed Block (CFB) (b) Urea-Molasses-Mineral-Block (U.M.M.B)</li> <li>5. Development of Fodder Bank</li> </ol>	<ol style="list-style-type: none"> <li>1. Feeding of Complete Feed Block</li> <li>2. Feeding of Urea-Molasses-Mineral-Block &amp; Fodder</li> <li>3. Feeding of stored Hay/Silage/Improved Quality Fodder</li> <li>4. Feeding of Tree leaves some of which are as follows: <ol style="list-style-type: none"> <li>1. Bamboo leaves</li> <li>2. Neem</li> <li>3. Bargad</li> <li>4. Peepal</li> <li>5. Seesam</li> <li>6. Subabul</li> </ol> </li> <li>5. Azolla feeding with concentrates</li> </ol>	Production of forage crops <ol style="list-style-type: none"> <li>1. Balanced feeding of Animal supported with little higher concentrate mixture</li> <li>2. Jowar/Cowpea</li> <li>3. Maize in September</li> <li>4. Berseem in Nov.-Dec.</li> <li>5. Napier / Para grass</li> </ol>

Drinking water	Repairing of water storage tank, tube-well, hand pump, well etc for water availability	<ol style="list-style-type: none"> <li>1) To ensure drinking water (electrolyte, Gur, Salt added water) to avoid dehydration in animals</li> <li>2) 2) To provide anti-stress drugs in drinking water to build up resistance to animals</li> </ol>	After drought adliv water and medicines should be given to animals to prevent disease in rainy season
Health and disease management	<p>Veterinary Preparedness with Medicines, Vaccines and provision for mobile ambulatory van.</p> <ul style="list-style-type: none"> <li>• Vaccination</li> </ul> <p>Mass vaccination should be conducted by a team of Department staff with proper maintenance of detailed Inoculation Register.</p> <p>Vaccines to be used for different animals</p> <p>Cattle and Buffalo</p> <p>Hemorrhagic Septicemia Vaccine</p> <p>Black Quarter Vaccine</p> <p>FMD Vaccine</p> <p>Anthrax Vaccine as per endemicity.</p> <p>Sheep and Goat</p> <p>Hemorrhagic Septicemia Vaccine</p> <p>PPR Vaccine</p> <p>FMD Vaccine</p> <p>Goat pox Vaccine</p> <p>Enterotoxemia Vaccine</p> <p>Anthrax Vaccine as per endemicity</p> <p>Pigs</p> <p>Hemorrhagic Septicemia Vaccine</p> <p>PPR Vaccine</p> <p>FMD Vaccine</p> <p>Goat pox Vaccine</p> <p>Enterotoxemia Vaccine</p> <p>Anthrax Vaccine as per endemicity.</p> <p>Dogs</p>	<p>Health camp and treatment</p> <p>Diseases that can occur during drought by drinking contaminated water should be given special attention and accordingly medicines should be available in the health camp for the following mentioned diseases.</p> <p>Salmonella spp.</p> <p>Escherichia coli</p> <p>Giardiasis</p> <p>Amoebiasis</p> <p>Rotavirus</p> <p>Leptospirosis</p> <p>Scabies</p> <p>Black leg</p> <p>Malignant Edema</p> <p>Foot rot</p> <p>Anthrax</p> <p>Botulism</p> <p>Tetanus</p> <p>Red water</p> <p>Black disease</p> <p>Entertoxemia</p> <p>Liver fluke</p> <p>Amphistomiasis</p> <p>Brooders pnemonia</p> <p>Treatment of Non infectious</p>	<p>Sanitation, deworming, treatment, health camps Culling of Sick animals and disposal of carcass</p> <p>Maintenance of Sanitation:</p> <ol style="list-style-type: none"> <li>1) Well ventilated animal shed should be created</li> <li>2) Proper disposal of urine and cow-dung to avoid contamination</li> <li>3) Disinfect the premises by application of bleaching powder</li> </ol> <p>De-worming :</p> <p>To control ticks infestation in animals</p> <p>Health Camp after the drought: Protection of livestock from out breaking and communicable diseases be made. Health camps are to be organised in drought affected areas to restore the normal breeding capability of breedable population as well as to restore the normal health of livestock .</p>

	<p>Rabies Vaccine</p> <p>List of life saving Medicines Corticosteroids Nikethamide Antibloat Adrenaline Antihistaminic Antidotes for common poisoning Antisnake venom Broad spectrum antibiotics Anti-inflammatory Antipyretic and Analgesics Fluids and Electrolytes</p> <ul style="list-style-type: none"> <li>• Mobile Veterinary Clinics</li> </ul> <p>Mobile Veterinary Clinics should be kept ready at Veterinary Hospital or Veterinary Camps so that immediate treatment of injured and affected animals may be done.</p> <p>For this MVC must have adequate drugs like antibiotic, analgesic, dewormer, ointment, antisnake venom and emergency health care facilities along with trained personnel.</p> <p>A good no. of mobile clinic teams should be planned consisting dedicated and experienced technical workers with allotment of area of operation.</p> <p>The teams should be kept in readiness having required stock of medicines and equipment to work in any adverse situation.</p> <p>A telephone directory should be maintained at the District level by collecting the telephone nos. of Vets, Para-Vets, NGOs / youth clubs / societies, volunteers etc. to collect feedback and plan the activities during the emergency.</p>	<p>Arrangement should be made for the treatment of drowning and traumatic injuries, aspiration pneumonia, lameness and other surgical cases in the health camp.</p> <p>Disinfection of livestock premises</p>	
Flood			
Cyclone			

Heat wave and cold wave	
-------------------------	--

<sup>s</sup> based on forewarning wherever available

### 2.5.2 Poultry

	Suggested contingency measures			Convergence/linkages with ongoing programs, if any
	Before the event <sup>a</sup>	During the event	After the event	
<b>Drought</b>				
Shortage of feed ingredients				
Drinking water				
Health and disease management	Vaccines to be used Mareks disease vaccine RDV (F1 & R2B), FPV, IBRV & IBDV	Vaccines to be used Mareks disease vaccine RDV (F1 & R2B), FPV, IBRV & IBDV	Vaccines to be used Mareks disease vaccine RDV (F1 & R2B), FPV, IBRV & IBDV	
<b>Floods</b>				
<b>Cyclone</b>				
<b>Heat wave and cold wave</b>				

<sup>a</sup> based on forewarning wherever available

### 2.5.3 Fisheries/ Aquaculture

	Suggested contingency measures		
	Before the event <sup>a</sup>	During the event	After the event
<b>1) Drought</b>			
A. Capture			
<b>B. Aquaculture</b>			
(i) Shallow water in ponds due to insufficient rains/inflow	(i) Thinning of population (ii) Arrangement of water supply from external resource	(i) Partial harvesting (ii) Addition of water (iii) Stocking of air breathing fishes	(i) Maintenances of remaining stock till favorable condition achieved (ii) If not feasible, total harvesting or transfer of fishes may be done. (iii) Preparation of the pond for next crop.

(ii) Impact of salt load build up in ponds / change in water quality	(i) Regular monitoring of water quality parameter. (ii) Arrangement of aeration (iii) Addition of water from external resource	(i) Arrangement of aeration. (ii) Addition of water (iii) Monitoring of water quality (iv) Reduction of manuring according to water level.	
<b>2. Floods</b>			
<b>3. Cyclone / Tsunami</b>			
<b>4. Heat wave and cold wave</b>			

<sup>a</sup> based on forewarning wherever available