

ISSN 0974-0775
NAAS Rating : 4.38

GREEN FARMING

International Journal of Applied Agricultural & Horticultural Sciences

Volume 9 ❖ Number 2 ❖ March-April 2018 ❖ Bimonthly



UGC Approved Jr.No. 45500 Global Impact Factor : 0.468 www.greenfarming.in



GREEN FARMING

(International Journal of Applied Agricultural & Horticultural Sciences)

(Abbr. - Green Farming Int. J.)

EDITORIAL ADVISORY PANEL (HONY.)

Dr. P.M. SALIMATH

Hon'able Vice-Chancellor,
University of Agricultural Sciences, Raichur (Karnataka)

Dr. D.L. MAHESWAR

Hon'able Vice-Chancellor, Univ. of Horticultural Sciences,
Bagalkot, Navanagar, Bagalkot (Karnataka)

Dr. AJAY KUMAR SINGH

Hon'able Vice-Chancellor, Bihar Agricultural University, Sabour,
Bhagalpur (Bihar)

Dr. M.B. CHETTI

Assistant Director General (HRD), Education Division
Indian Council of Agricultural Research, Krishi Bhawan, New Delhi

Dr. RAJEEV K. VARSHNEY

Research Programme Director - Genetic Gains
International Crops Research Institute for the Semi-Arid Tropics
(ICRISAT), Patancheru, Hyderabad (Telangana)

Dr. ASHOK K. PATRA

Director, Indian Institute of Soil Science (ICAR)
Nabi Bagh, Berasia Road, Bhopal (M.P.)

Dr. K.K. SINGH

Director, Central Institute of Agricultural Engineering (ICAR)
Nabi Bagh, Berasia Road, Bhopal (M.P.)

Dr. J.C. TARAFDAR

National Fellow & Principal Scientist (Soil Chem/ Microbiology)
Central Arid Zone Research Institute (ICAR) Jodhpur (Rajasthan)

Dr. K.L. SHARMA

Principal Scientist & National Fellow, Central Research Institute
for Dryland Agriculture (ICAR), Hyderabad (A.P.)

Dr. PRAMOD W. RAMTEKE

Prof. & Dean, Post Graduate Studies, S.H. Institute of Agriculture
Technology & Sciences (Deemed University), Allahabad (U.P.)

Dr. N.C. PATEL

Hon'able Vice-Chancellor
Anand Agricultural University, Anand (Gujarat)

Dr. NARENDRA SINGH RATHORE

Dy. Director General Education, Division of Education,
Krishi Anusandhan Bhawan - II, ICAR, New Delhi

Dr. MYER MULA

Dean, College of Agriculture Science and Technology, Pampanga State
Agricultural University, Magalang, Pampanga, Philippines

Dr. H.S. DHALIWAL

Dean, College of Agriculture
Punjab Agricultural University, Ludhiana (Punjab)

Dr. K.S. RANA

Ex. Head & Principal Scientist, Division of Agronomy,
Indian Agricultural Research Institute (IARI), Pusa, New Delhi

Dr. V.M. BHALE

Dean, P.G.I., College of Agriculture,
Dr. P.D. Krishi Vidyapeeth, Krishinagar, Akola (Maharashtra)

Dr. SURENDRA S. SIWACH

Director of Research and Dean, College of Agriculture
CCS Haryana Agricultural University, Hisar (Haryana)

Dr. M.K. ARVADIA

Principal & Dean, N.M. College of Agriculture,
Navsari Agricultural University, Navsari (Gujarat)

Dr. SRINIVASAN RAMASAMY

Entomologist, AVRDC - The World Vegetable Center,
P.B. Box. 42, Shanhua, Tainan (Chinese) Taipei (China)

Dr. A. NOOR

Ex. Professor (Entom.) Head & ZMT - Plant Protection, Agri. Research
Station (Raj. Agri. Univ), Mandore, Jodhpur, Rajasthan (Hony. Secret.)

Editor

Dr. A. HUSSAIN

Editorial Asstt.

K. SINGH

Chief Editor (Hony.)

Dr. A. NOOR

Publisher, Printer & Owned by : Dr. Anwar Hussain, and Published at White House, 78-A, Bank Colony Road, Near Lal Bangla, Raikabagh, JODHPUR - 342 006 (Rajasthan) INDIA. **Printed at :** Bhandari Offset, JODHPUR - 342 005 (Rajasthan) INDIA, Editor - Dr. Anwar Hussain

Registration No. RAJENG/2010/32178

NAAS Rating - 4.38 (2018)

ISSN 0974-0775

Citation : Abstracting & Indexing in Indian Science Abstracts (NISCAIR) & CABI International; Indexing in Agricultural & Biological Abstracts etc.

Disclaimer : Copy right reserved with the publisher. No part of this journal can be reproduced, stored or transmitted in any form or by any means without the written permission of the publisher. The publisher/editor does not assume any responsibility for opinions offered by the authors published in the Journal. The views expressed in the articles by the authors are their own. All disputes are subject to the exclusive jurisdiction of competent courts in Jodhpur only.

The publisher and authors declare that they have compiled this document carefully. However, for any technical shortcomings, they hold no responsibility whatsoever. They assume no liability whatsoever for any damage resulting from the use of this document or its contents. They do not watch any claim for made by the advertisement of products and services.

Note : The editor does not accept responsibility for returning unsolicited publication material. The publisher and the editor are not responsible for any delay, whatsoever, in publication/delivery of the periodicals to the subscribers due to unforeseen circumstances or postal delay/ lost.

All Correspondence to : Editorial Office, GREEN FARMING, White House, 78-A, Bank Colony Road, Near Lal Bangla, Raikabagh, JODHPUR - 342 006 (Rajasthan) India.



CONTENTS

Previous issue :
Vol. 9, No. 1, pp. 1-190

Review Paper

Organic farming in Eastern India : Relevance, opportunities and challenges

- **AJOY KUMAR SINGH, SRINIVASARAGHAVAN A. and SHRIDHAR PATIL** 191

Research Papers

Heterosis studies for yield and its components under aerobic condition in hybrid rice (*Oryza sativa* L.)

- **SUMITRA UPENDI and N.K. MOTIRAMANI** 198

Combining ability and pooled score studies on baby corn traits in maize (*Zea mays* L.)

- **LASYA BHAT and S.K. DESHPANDE** 206

Characterization of maize genotypes for baby corn traits

- **LASYA BHAT and S.K. DESHPANDE** 212

Genetic variability, character association, path coefficient in forage sorghum [*Sorghum bicolor* (L.) M.]

- **H.I. DAMOR, H.P. PARMAR, D.P. GOHIL and A.A. PATEL** 218

Genetic variability studies for yield and quality traits in rice (*Oryza sativa* L.)

- **KAVURI KALPANA, S. THIRUMENI, V. VENGADESSAN and MOHAMED YASSIN G.** 224

Character association studies for early vigour traits and yield components in direct sown rice (*Oryza sativa* L.)

- **M. SRILAKSHMI CHAITANYA, B. KRISHNA VENI, M. LAL AHAMED, M. GIRIJA RANI and K. JAYA LALITHA** 228

Genetic divergence studies in mungbean (*Vigna radiata* L. Wilczek) genotypes

- **NAVEEN KUMAR MARKAM, S.K. NAIR, ISHU KUMAR KHUTE and PREETI PAINKRA** 231

Genetic variability, heritability and genetic advance in mungbean (*Vigna radiata* L. Wilczek) genotypes

- **SRAVAN KUMAR MALLI, G.R. LAVANYA and B.S.K. NIKHIL** 235

Assessment of genetic diversity and variability in ridge gourd for growth and yield parameters

- **MANOJ Y.B., LAKSHMANA D., SHASHIKALA S. KOLAKAR and CHANDANA B.C.** 239

Studies on character association and genetic variability for important traits in ridge gourd (*Luffa acutangula*)

- **MANOJ Y.B., LAKSHMANA D., GANAPATHI M. and CHANDANA B.C.** 244

Heritability and correlation studies in wild apple genotypes from different locations in Kashmir Valley

- **K.M. BHAT, SHAZIYA HASSAN, H.U. REHMAN, A.H. PANDITH and M.A. MIR** 248

Studies on heterosis for yield attributing traits in cucumber (*Cucumis sativus* L.)

- **PRASHANT R. NAIK, NAGARAJAPPA ADIVAPPAR, V. SRINIVASA, S. GANGAPRASAD and SHRIDHAR P. HERLE** 253

Estimation of genetic variability, heritability, genetic advance, correlations and path analysis in advanced mutant breeding lines of sesame (*Sesamum indicum* L.)

- **MAHALAXMI K. PATIL and R. LOKESHA** 257

Contd.

Relationship of organic and inorganic fertilization with soil fertility and crop yield under maize-wheat cropping system	
— ASMATULLAH DURANI, B.S. BRAR, KHUWAJA SAFIULLAH, AMINULLAH YOUSAFZAI and HASHMATULLAH DURRANI 261
Influence of time of sowing and weed management on weed flora, nutrient content, uptake and productivity of summer blackgram (<i>Vigna mungo</i> L.)	
— S.K. PHAJAGE, V.C. RAJ and A.S. CHAVAN 270
Ameliorative effect of different tree species on soil profile development of salt affected soil	
— RAVENDRA SINGH, S.G. RAJPUT, H.P. CHAUDHARY, R.R. VERMA, NIRMAL KUMAR KATIYAR and YOGESH KUMAR 277
Performance of cultivars in direct seeded rice with various seeding rates and levels of micronutrients under Tunga Bhadra irrigation during summer	
— SHUBHA G.V., CHITTAPUR B.M., MANJUNATH B., REKHA R.G. and MUNIYAPPA N. 284
Effect of sodium silicate on the growth and physiological attributes of tomato	
— CHANCHAL MALHOTRA, RITITHAPAR KAPOOR, DEEPAK GANJEWALA and N.B. SINGH 290
Demand driven nutrient management for enhanced nutrient use efficiency and productivity in rice and wheat under rice-wheat cropping system	
— A. QURESHI, D.K. SINGH, P.C. PANDEY, P.K. CHANDRAKAR and P.S. MARKAM 296
Effect of plastic mulch and supplemental irrigation on growth and yield of black gram (<i>Vigna mungo</i> L.)	
— CHANDRAKANT Y. MAHALE, S.R. BHAKAR, S.S. LAKHAWAT, M. KOTHARI and B.G. CHHIPA 301
Effect of foliar nutrition on the growth and yield of <i>Bt</i> cotton (<i>Gossypium hirsutum</i> L.) under irrigated condition	
— V.L. KIKANI, A.R. PATEL, A.M. POLARA and D.K. DAVARA 306
Fibre yield, quality parameter, light use efficiency and economics of linseed as influenced by intercropping system of safflower and linseed under rainfed condition	
— KARALE GANGADHAR, S.A. BIRADAR, B.K. DESAI, AJITHKUMAR K. and RAJANNA 310
Morphological and Micro-anatomical variation in different ploids of mulberry genotypes (<i>Morus</i> sp.)	
— R. RAVI KUMARA, H.L. RAMESH, M.R. SWAMY GOWDA and P. SUDHAKAR 314
Effect of different land use systems on soil properties in red soils of Jambadahalla subwatershed, Chikkamagaluru district	
— NETHRAVATHI B., T.S. VAGEESH and Y. VISHWANATH SHETTY 318
Response of <i>Bt</i> cotton (<i>Gossypium hirsutum</i> L.) to foliar application of gibberellic acid under rainfed conditions	
— M.D. GIRI, C.P. JAYBHAYE, D.G. KANWADE and J.R. WADKAR 323
Response of okra (<i>Abelmoschus esculentus</i> L.) growth, yield attributes and economics on irrigation levels and mulching	
— M.M. SIPPO, P.K. SINGH, S.R. BHAKAR and S.S. LAKHAWAT 327
Influence of tillage practices and nutrient sources on yield, quality of sweet corn and post-harvest soil fertility	
— V.N. SHETYE, R.C. TIWARI, V.V. SAGVEKAR, S.S. KADAM and A.P. CHAVAN 331
Soil fertility status under <i>Bt.</i> cotton growing vertisols of Dharwad district	
— RAMYA S.H., N.S. HEBSUR, RAVINDRA V.M. and S.S. ANGADI 335

Image analysis for size distribution of <i>kodo</i> and <i>kutki</i> millet	
— P.K. NISHAD, S.D. DESHPANDE and N. KOTWALIWALE 339
Impact of hexanal vapour treatment on the shelf-life and post-harvest quality of banana var. Poovan	
— ASHWINI T., S. GANAPATHY and K.S. SUBRAMANIAN 344
Reflexes of biofungicide (<i>Trichoderma harzianum</i> R.) on seed mycoflora & seed germination of some medicinal plant species	
— SATISH DUBEY, R.K.S. TIWARI and SANTRAM SAHU 348
Efficacy of <i>Cissus quadrangularis</i> L. extract against wilt of tomato (<i>Lycopersicon esculentum</i> L.) caused by <i>Fusarium oxysporum</i> f. sp. lycopersici	
— M.R. SWAMY GOWDA, H.M. MAHESH, M.S. SHARADA and P. SUDHAKAR 353
Molecular characterization of <i>Fusarium pallidoroseum</i> isolates by using RAPD technique	
— T.T. BARIA and R.K. PATIL 357
Effect of bactericides and fungicides in management of citrus canker in Kagzi lime under nursery condition	
— CHETHANKUMAR K., KIRANKUMAR K.C. and PRASANNA KUMAR M.K. 361
Diversity of insect pests and natural enemies in <i>Bt</i> and non- <i>Bt</i> cotton	
— S. ANUSHA and R. SWAMINATHAN 365
Influence of cultivation methods on incidence of rice pests	
— D. SUDHA RANI, CH. NAGA SATYASRI, M.N. VENKATESH and K. ANAND KUMAR 370
Some physical properties of quinoa (<i>Chenopodium quinoa</i> Willd.)	
— SAGAR CHAVAN, SURBHI SHARMA, VINAY PATIL and NARENDRA KUMAR JAIN 375
Effect of harvesting of fruits at different stage of maturity on nutritional quality of Kinnow under Hot Arid Zone of North Western India	
— JULIUS UCHOI, R.S. SINGH, R. BHARGAVA, HARE KRISHNA, VENKATESAN K., ABHISHEK KUMAR and ANIL PATIDAR 379
Development and performance evaluation of tractor operated tuber rhizomes digger	
— POOJA N. WAJIRE, M.V. JADHAV, C.M. PAREEK, S.H. THAKARE and V.P. KHAMBALKAR 383
Development of crop water stress index for onion (<i>Allium cepa</i> L.) crop using infrared thermometry	
— AMITKUMAR L. DAVARIYA, S.R. BHAKAR, M. KOTHARI and R.S. SINGH 387
Study on energy use pattern in groundnut (<i>Arachis hypogaea</i> L.) production and processing in Rajasthan	
— DIKSHA SRIVASTAVA, A.N. MATHUR and MRUNAL K. SHIRSHAT 390

Short Communication

Evaluation of the physico-chemical parameters of underutilized <i>Allium</i> species (<i>Allium hookeri</i> Thw.) in Northeast region of India	
— M. BILASHINI DEVI, S.R. ASSUMI, H.D. TALANG, A.K. JHA and NISHA THAKUR 393

Strategic Vision Message : 44

Agripreneurship : An exciting challenge towards economic development

— MYER G. MULA	
-----------------------	--



Effect of harvesting of fruits at different stage of maturity on nutritional quality of Kinnow under Hot Arid Zone of North western India

JULIUS UCHOI^{b1*}, R.S. SINGH^{a2}, R. BHARGAVA^{a3}, HARE KRISHNA^{a4}, VENKATESAN K.^{c5},
ABHISHEK KUMAR^{b6} and ANIL PATIDAR^{b7}

^a ICAR-Central Institute for Arid Horticulture, Bikaner - 334 006 (Rajasthan)

^b ICAR-Central Arid Zone Research Institute, Regional Research Station, Jaisalmer - 345 001 (Rajasthan)

^c ICAR-Central Island Agricultural Research Institute, Portblair - 744 101 (Andaman & Nicobar Island)

Received : 20 January 2018 ; Revised accepted : 20 March 2018

ABSTRACT

Kinnow fruits were collected from the 15 year old orchard of ICAR-Central Institute for Arid Horticulture (CIAH), Bikaner at three different time of harvesting such as immature (early), fully mature (mid) and over-mature (late) as treatments with five replications under Completely randomized design. Observations made on fifteen different parameters, which include seven physical and eight biochemical traits. Assessment of nutritional quality of Kinnow revealed the impact of each stages of fruit maturity on the quality aspect of this citrus species. Important nutritional quality parameters viz TSS (11.8°Brix), vitamin –C (49.0mg/100ml), total sugar (8.1%), reducing sugar (4.9%), non-reducing sugar (4.1%) are found to increase as the fruits progress towards maturity. Only antioxidant activity (2.1mMTE/L) and total phenolic content (2060.4µg/100g) revealed phenomenal changes decreasing at mid maturity and again increase in the later stage of full maturity. Although the significant difference obtained in physical and biochemical changes between the mid and late maturity stage, the maximum attainment of fruit nutritional quality at the late maturity or full ripe stages of the mandarin suggesting the right stage of consumption of Kinnow fruits.

Key words : *Citrus*, *Kinnow*, *Nutritional quality*, *Stage of consumption*.

INTRODUCTION

Citrus fruits are most popular in many parts of the world due to its distinctive flavor, taste, and aroma as well as multiple health benefits associated with it. The major phenolic compounds detected in different citrus fruits are categorized as flavonoids and phenolic acids (Balasundram *et al.*, 2006). Among the fruits, several species of citrus possess good amount of antioxidant properties. Kinnow (*Citrus reticulata* Blanco), an important introduced citrus crop, is widely popular and cultivated in arid region of western India. Mandarins are rich in ascorbic acid and calcium. They are a great source of vitamin C. The chemical composition of the mandarin is as

given in **Table 1**. The presence of vitamins and antioxidants within the fruit provide a wide scope in combating the nutritional deficiency in ever growing population of the region. There exist an immense potential to tap and assess the various physical and biochemical activity of Kinnow mandarin in India at various stages of ripening, as quantifying and determining the amount of antioxidant activity at various stages of fruit ripening would help in determining the most suitable stage of its consumption. Therefore the present study was conducted on assessment of physical and biochemical activity of Kinnow mandarin in ICAR-Central Institute for Arid Horticulture, Bikaner, Rajasthan at 3 different months from November, 2014 (*i.e.* beginning of colour changing stage) to January, 2015 (*i.e.* full ripen stage).

1,5,6,7 Scientist *(juliusuas@gmail.com), 2,3 Principal Scientist

4 Senior Scientist

Table 1. Chemical composition of Mandarin (Per 100 g of edible portion)

Chemical composition	
Moisture	82.6 - 90.2 g
Protein	0.61 - 0.215 g
Fat	0.05 - 0.32 g
Fiber	0.3 - 0.7 g
Ash	0.29 - 0.54 g
Calcium	25.0 - 46.8 mg
Phosphorus	11.7 - 23.4 mg
Iron	0.17 - 0.62 mg
Carotene	0.013 - 0.175 mg
Thiamine	0.048 - 0.128 mg
Riboflavin	0.014 - 0.041 mg
Niacin	0.199 - 0.38 mg
Ascorbic acid	13.3 - 54.4 mg

Source : Morton, J. 1987

MATERIALS AND METHODS

The experimental material 'Kinnow' fruits were collected from the 15 year old orchard of ICAR-Central Institute for Arid Horticulture (CIAH), Bikaner situated at latitude 28°10' and longitude 73°35'. Experiment were conducted using three different time of harvesting such as immature (early), fully mature (mid) and over-mature (late) as treatments with five replications under completely randomized design. Ten fruits from five randomly selected trees pooled together and divided into five lots with ten fruits each. Each lot was considered as a replication. Observations made on fifteen different parameters of Kinnow fruits, which include seven physical [fruit weight, length, width, volume dispersed, specific gravity, peel weight and juice recovery] and eight biochemical traits.

TSS (Brix) was determined by hand-refractometer. The titratable acidity (%), total and reducing sugar content were estimated by using the methods described in A.O.A.C. (1990), Dubois *et al.* (1951) and Nelsons modification of Somogyi method (Somogyi, 1952), respectively. Total phenol and antioxidant content of Kinnow juice were measured on the basis of Folin–Ciocalteu colorimetric method using gallic acid solution (Singleton *et al.*, 1999) and CUPRAC method using UV-VIS spectrophotometer at 450 nm (Meng *et al.*, 2011), respectively. Data were subjected to one way ANOVA using SPSS software. The means of significant parameters, among the treatment were compared using critical difference (CD) tests at probability level of 5%.

RESULTS AND DISCUSSION

In the present study, changes in seven physical and eight biochemical traits were observed in the selected fruits of Kinnow mandarin at three maturity stages (early, mid and late) (Fig. 1) and it revealed the presence of significant differences (5% level) in three physical and four biochemical traits among




Fig. 1. A view of the Kinnow fruits at three different stage of harvesting (Early, mid and late)

Kinnow fruits collected at different stages (Table 2 and 3). Although, differences in physical changes occurred between Kinnow fruits harvested at early and late maturity stage with respect to fruit weight (g), length (cm), width (cm) and volume dispersed (cc) [as (152.0 and 170.4), (5.8 and 6.3), (7.1 and 7.3) and (146.0 and 183.0), respectively], very little difference noted in these characters between mid and late stage. Decreasing trend was observed in specific gravity of fruits from early to mid and mid to late stage of maturity as 1.1 to 1.0 and 1.0 to 0.9, respectively. But the peel weight (g) of fruits increased from 36.6 at early stage to 82.9 at mid stage of maturity and then, decreased (77.9) in late stage of maturity.

In case of juice recovery of Kinnow, late and middle stage of maturity showed high juice recovery percentage as 51.8 and 41.2%, respectively with significant difference at 5% level as compared to early stage (35.9%). In general, juice contents in citrus fruits increase towards maturity and then decrease when fruits become over-mature. These contents remain variable for the entire period of fruit development. Decrease in juice

Table 2. Physical parameters of the Kinnow fruits at three different stage of harvesting

	Different stages of fruit			CD
	Early	Mid	Late	
Fruit weight (g)	152.0	172.4	170.4	35.14
Fruit length (cm)	5.8	6.2	6.3	0.44
Fruit width (cm)	7.1	7.3	7.3	0.52
Volume dispersed (cc)	146.0	180.0	183.0	40.37
Specific gravity	1.1	1.0	0.9	0.08*
Peel weight (g)	36.6	82.9	77.9	11.18*
Juice recovery (%)	35.9	41.2	51.8	7.31*

* Significant at 5% level

Table 3. Biochemical activities of Kinnow fruits at three different stage of harvesting

	Different stages of fruit			CD
	Early	Mid	Late	
Acidity (%)	1.7	0.4	0.3	0.10*
TSS (Brix)	10.3	11.0	11.8	1.15
Total sugar (%)	6.5	7.3	8.1	0.23
Reducing Sugar (%)	3.6	4.5	4.9	1.91
Non reducing sugar (%)	2.9	3.4	4.0	0.26*
Vit-C (mg/100ml)	46.3	48.0	49.0	15.30
Total phenol ($\mu\text{g}/100\text{g}$)	991.4	179.4	2060.4	50.68*
Antioxidant activity (mMTE/L)	0.3	1.0	2.1	0.05*

* Significant at 5% level

content reveals quality decline (Riaz *et al.*, 2015). Whereas in present study, the juice content of Kinnow increased in over-mature stage which means that there was improvement in quality. It might be due to the effect of weather condition of area such as rainfall, high humidity or cultural operation (irrigation management) before fruit collection (Cruse *et al.*, 1982) and it might also be picking season of fruit (Gillfillan *et al.*, 1972).

Acidity in citrus fruits is one of the quality traits and it should be lower in sweet oranges for better quality (Nawaz *et al.*, 2012). In general, acidity decreased as fruits approach towards maturity and then increased as fruit become over-mature (Grewal *et al.*, 2000; Anwar *et al.*, 1999). In present study, there was an initial rapid rise (1.7) of titratable acids at early stage of maturity, followed by dip in mid and late (0.4 and 0.3, respectively) stage of maturity due to dilution effect, because of increase in size of fruit mainly water content. Trends of total soluble solids and acidity from early to late maturity stage of Kinnow fruits obtained in our study followed the chemical changes occurring in citrus fruits during growth and maturation of it (Syverstsen *et al.*, 1983; Cohen, 1988; Mcanenev *et al.*, 1995; Yamanishi, 1995; Yamanashi and Hasegawa, 1995; Yakushiji *et al.*, 1996). TSS is good quality index of Kinnow fruits and sugars constitute 85% of it. Normally, it increases as fruit become mature and then decreases as over-mature and a similar trend (at early, mid and late stage of maturity as 10.3, 11.0 and 11.8 brix respectively) observed in our study. This rise might be due to increase in sucrose contents, which hydrolyzed to simple sugars during maturity of fruits. Its content varies on basis of position of fruits on tree (Syverstsen *et al.*, 1983; Fallahi and Moon, 1989).

TSS is inversely related with titratable acidity (%), as the fruit attain towards ripening the amount of total soluble solids and other sugar compound increases while the organic acids present in the fruit juice decreases. The increase in sugars accumulation may be due to lower pH which enhanced the sink strength of the fruits (Hockema and Atxeberria, 2001).

Sugars are the major components of citrus juice's soluble solids and sweetness of citrus juice is intrinsic to its sugar composition. Sucrose is present in the largest amounts in citrus juice followed by fructose and glucose (Kelebek & Selli, 2011). The increasing trend observed in total sugar content of Kinnow fruits till maturity (6.5 at early stage to 8.1% at late stage of maturity) in present study also reported in earlier work (Miller and Evans, 1997). Potential factors responsible for variability in sugar content of Kinnow fruits are: variety, rootstock, geographical location, weather and cultural practices. The vitamin-C (ascorbic acid) content showed increasing trend without significant differences at early to late stage of maturity from 46.3 to 49.0 mg/ 100ml. This finding was contrary to the work reported by Danielle *et al.* (2014), where they reported that in citrus vitamin-C concentration decreases with ripening and vitamin-C was found to be at the highest during matured stage than the green stage because the fruit juice volume and fruit size increases in mature fruit. The changes in ascorbic acid can be attributed by effect of processing, storage time, variety, exposure to light and other factors (Gillfillan *et al.*, 1972).

The highest concentrations found in Citrus sp. correspond to flavanone glycosides, followed by flavones, flavonols and the fully polymethoxylated flavones (PMFs) (Kawaii *et al.*, 1999). Hesperidin, narirutin, naringin, eriocitrin and neohesperidin are the major flavanone glycosides (Peterson *et al.*, 2006). Vitamin C is considered to be one of the most important nutrients found in citrus fruit, and it is an important water-soluble antioxidant that plays a crucial role in the suppression of superoxide radicals (Kaur and Kapoor, 2001). In citrus, vitamin-C and phenolic compound constitute the major portion of antioxidants. In present study, the amount phenol content ($\mu\text{g}/100\text{g}$) and anti-oxidant activity (mMTE L⁻¹) observed as 991.4 to 2060.4 and 0.3 to 2.1 at early and late stage of maturity, respectively.

Evidence of increased antioxidant activity in ripe fruits is contradictory to an earlier work of Rekha *et al.* (2012), wherein they reported that as the fruit advances towards ripening, there is a reduction in phenolic and other antioxidant contents as compare to unripe ones. The antioxidant activity may be contributed by integral effect of both ascorbic acid and phenolic compounds present in the citrus juice and also with the possibility of other factors such as the type of citrus cultivar, stage of maturity and the methodology selected for analyzing antioxidants (Juihaimi and Ghafoor, 2013). Being antioxidant rich fruit, consumption and the right stage of harvesting of Kinnow to obtain maximum benefit of this antioxidant identified in present study could provide a wide scope in combating the nutritional deficiency in ever growing population of the region. Although the significant difference obtained in physical and biochemical changes between the mid and late maturity stage,

the maximum attainment of fruit nutritional quality at the late maturity or full ripe stages of the mandarin suggesting the right stage of consumption of Kinnow fruits.

REFERENCES

- A O A C. 1990. *Official Methods of Analysis*. 15th Edition, Association of Official Agricultural Chemist. Washington, D. C., pp. 16.
- Anwar SA, Ahmed B, Sarfarz M, Hussain K M, Bhatti and Saqib M. 1999. Effect of picking time on physical and chemical characteristics of sweet orange. *J. Agric. Bio.* **1** : 59-61.
- Balasundram N, Sundram K and Samman S. 2006. Phenolic compounds in plants and agri-industrial by-products: Antioxidant activity, occurrence and potential uses. *Food Chem.* **99** : 191-203.
- Cohen E. 1988. The Chemical composition and sensory flavor quality of 'mineola' tangerines. Part. 1 : Effects of fruit size and within-tree position. *J. Food Sci Tech.* **35** : 455-457.
- Cruse R R, Wiegler C L and Swanson W A. 1982. The effect of rainfall and irrigation management on citrus juice quality in Texas. *J. American Soc. Hort. Sci.* **107** : 767-770.
- Danielle dos Santo Bonfim de Castro, Lana de Souza Rosa., Ellen Mayra da Silva Menezes and Teodoro A J. 2014. Comparative evaluation of organic and conventional farming on chemical quality parameters and antioxidant activity in fruits. *African J. Biotech.* **13** (18) : 1883-1890.
- Dubois M, Gilies K, Hammlton J K, Robers P A and Smith F A. 1951. A calorimetric method for the determination of sugars related substances. *Ann. Chem.* **28** : 350-356.
- Fallahi E and Moon J W. 1989. Fruit quality and mineral nutrient from exposed versus internal canopy of four citrus varieties. *J. Plant Nutr.* **12** : 523-534.
- Gillfillan I M, Holtzhasusen L C, Koekemar N and Du Plessis J A. 1972. Picking season changes in Valencia quality factor. *Citrus Grow and Sub-trop Fruit J.* **454** : 11-15.
- Grewal A G, Hafiz I A, Chaudhary A H, Khan M I and Chaudhary M I. 2000. Quality estimation during marketing of kinnow and feutrell's early. *Int. J. Agric Biol.* **2** : 328-330.
- Hockema B R and Exteberria E. 2001. Metabolic contributors to drought enhanced accumulation of sugars and acids in oranges. *J. American Soc. Hort. Sci.* **126** : 599-605.
- Juhaimi F Y A and Ghafoor. 2013. Bioactive compounds, antioxidant and physio-chemical properties of juice from lemon, mandarin and orange fruits cultivated in Saudi Arabia. *Pak. J. Bot.* **45** (4) : 1193-1196.
- Kaur C and Kapoor H C. 2001. Antioxidants in fruits and vegetables - The millennium's health. *International Journal of Food Science and Technology.* **36** : 703-725.
- Kawaii S, Tomono Y, Katase E, Ogawa K and Yano M. 1999. Quantitation of flavonoid constituents in citrus fruits. *Journal of Agricultural and Food Chemistry.* **47** : 3565-3571.
- Kelebek H and Selli S. 2011. Determination of volatile, phenolic, organic acid and sugar components in Turkish cv. Dortyol (*Citrus sinensis* L. Osbeck) orange juice. *Journal of the Science of Food and Agriculture.* **91** : 1855-1862.
- Mcaneney K J, Richardson A C, Astill M S and Anderson P A. 1995. The inscrutable mandarin. *Agric for Meteorol.* **75** : 71-84.
- Meng J, Fang Y, Zhang A, Chen S, Xu T, Ren Z, Han G, Liu J, Zhang Z and Wang H. 2011. Phenolic content and antioxidant capacity of Chinese raisins produced in Xinjiang Province. *Food Res Int.* **44** : 2830-2836.
- Miller N J and Rice Evans C A. 1997. Cinnamates and hydroxyl benzoates in the diet: antioxidant activity assessed using the ABTS radical cation. *British Food J.* **99** : 75-61.
- Morton J F. 1987. (*in*) *Fruits of Warm Climates*. Florida Flair Books, Miami. pp. 550.
- Nawaz MA, Ahmed W, Maqbool M, Saleem BA, Hussain Z, Aziz M and Shafique A. 2012. Characteristics of some potential cultivars for diversification of citrus industry of Pakistan. *Int. J. Agric. Appl. Sci.* **4** : 58-62.
- Peterson J J, Beecher G R, Bhagwat S A, Dwyer J T, Gebhardt S E and Haytowitz D B. 2006. Flavanones in grapefruit, lemons, and limes: A compilation and review of the data from the analytical literature. *J. Food Compos. Anal.* **19** : S74-S80.
- Rekha C, Poornima G, Manasa M, Abhipsa V, Pavithra Devi J, Vijay Kumar H T and Prashith Kekuda T R. 2012. Ascorbic acid, total phenol content and antioxidant activity of fresh juices of four ripe and unripe citrus fruits. *Chem Sci Trans.* **1** (2) : 303-310.
- Riaz H, Begum A, Raza S A, Khan Z M U, Yousaf H and Tariq A. 2015. Antimicrobial property and phytochemical study of ginger found in local area of Punjab, Pakistan. *Int. Curr. Pharm. J.* **4** (7) : 405-409
- Singleton V L, Orthofer R and Lamuela Reventos R M. 1999. Analysis of total phenols and other oxidation substrates and antioxidants by means of Folin-Ciocalteu reagent. *Meth. Enzymol.* **299** : 152-178.
- Somogyi M. 1952. Notes on sugar determination. *J. Biol. Chem.* **195** (1) : 19-23.
- Syvetsen W B and Smith M L. 1983. Environmental stress and seasonal changes in proline concentration in kinnow of citrus tree tissues and juice. *J. American Soc. Hort.* **3** : 113-114.
- Yakushiji H, Nonami H, Fukuyama T, Ono S, Takagi N and Hashimoto Y. 1996. Sugar accumulation enhanced by osmo regulation in Satsuma mandarin fruit. *J. American Soc Hort Sci.* **121** : 466-472.
- Yamanishi O K and Hasegawa K. 1995. Trunk strangulation responses to the detrimental effect of heavy shade on fruit size and quality of Tosa Buntan' pummel. *J. Hort Sci.* **70** : 875-887.
- Yamanishi O K. 1995. Trunk strangulation and oranges. winter heating effects on fruit size, internal quality. *J. Hort. Sci.* **70** : 65-74.

