

## Short Note

Vipin Chaudhary\* and Rakesh S. Tripathi

**First record of Little Indian field mouse, *Mus booduga* (Gray 1837) (Rodentia: Muridae), from cold arid region of Leh-Ladakh, Jammu and Kashmir, India**<https://doi.org/10.1515/mammalia-2017-0066>

Received June 2, 2017; accepted March 9, 2018; previously published online April 20, 2018

**Abstract:** *Mus booduga* (Gray 1837) is a mesic rodent, recorded for the first time from the cold arid region of Leh-Ladakh, Jammu and Kashmir, India. The species was collected from crop fields and nearby areas from different altitudes ranging from 3187 to 3768 m above mean sea level. The paper describes the external measurements and diagnostic features of *M. booduga* collected from Leh-Ladakh region. Earlier records had shown its altitudinal distribution from 250 to 3695 m above mean sea level from Uttarakhand in the Himalayan region, however in the present study the mouse was collected up to an altitude of 3768 m which is the highest elevational record of this species for the Himalayan range.

**Keywords:** altitude; distribution; Leh; *Mus booduga*; trapping.

Rodents are one of the most diverse groups of mammals with 2277 species recorded worldwide, whereas 103 species have been reported from India (Wilson and Reeder 2005, Pradhan and Talmale 2009). The Little Indian field mouse, *Mus booduga* (Gray 1837) (= *Leggada booduga* Gray, Charlesworth's Mag. Nat. Hist. 1:586) is one of the smallest and widely distributed rodent species in India. Perusal of extant literature revealed that the Little Indian field mouse is distributed in East Pakistan, India, South Nepal, Sri Lanka and Bangladesh (Blanford 1891, Ellerman 1961, Chesemore 1970, Roberts 1977, Posamentier 1989, Aplin et al. 2016). A recent molecular phylogeny by Shimada concluded that the Myanmar *Mus* previously attributed to *M. booduga* were representing a different clade (Shimada et al. 2010) and these authors retracted the distribution

range of the species in the East to Bangladesh. In India *M. booduga* is reported from almost every part of country viz., from Arunachal Pradesh in the east to Rajasthan and Gujarat in the west and from Jammu and Kashmir in the north to Tamil Nadu and Andaman and Nicobar Islands in the south (Hinton and Lindsay 1926, Ellerman 1961, Biswas and Tiwari 1966, Prakash et al. 1971, Sharma and Sharma 1976, Mandal 1981, Chakraborty 1983, Agrawal 2000, Idris et al. 2003, Birah et al. 2012).

The earliest record of rodents of British India by Blanford (1891) did not mention occurrence of *Mus booduga* from the Himalayan region, however Wroughton (1914) for the first time reported it from the Kumaon region in Nainital and Almora Districts in Uttarakhand at altitudes varying from 250 to 2155 m above mean sea level. Lindsay (1926) has also reported it from Gopalpur (c 2770 m) (Kangra Valley, Himachal Pradesh). Agrawal (1980) and Mandal (1981) reported its occurrence from the eastern offshoot of the Himalayas in North Eastern Hill regions including Arunachal Pradesh. Mandal (1984) reported this species from Nelang village in Uttarkashi district (Uttarakhand) which, till the present report, was the highest altitudinal range (c 3695 m) of distribution of *M. booduga* in the Himalayas. Regarding its distribution in Jammu and Kashmir State (India), the species has been mainly reported from Jammu sector (c 315 m) by Varma (1968); Punch, Udhamapur, Jhajar Kotli and Baderwah by Sharma and Sharma (1976) and Chakraborty (1983).

However, the present study, for the first time reports its occurrence at high altitudes in cold arid regions of Leh-Ladakh (Jammu and Kashmir). In the field surveys undertaken during June–September, 2014–2016, *Mus booduga* was collected from crop fields and field stores in rural areas and from stores and godowns in urban areas of Leh-Ladakh district (Jammu and Kashmir) at an altitudinal ranges from 3187 to 3768 m above mean sea level. This field mouse was the only rodent species recorded from crop fields and field stores, however in urban areas of Leh the species inhabited the stores and godowns along with Turkish rat, *Rattus pyctoris* (Hodgson 1845) (Table 1). During the study period, 44 specimens of *M. booduga* were collected using live Sherman traps. The males (27)

\*Corresponding author: Vipin Chaudhary, All India Network Project on Vertebrate Pest Management, ICAR-Central Arid Zone Research Institute, Jodhpur-342003, India, e-mail: vipin\_cima@yahoo.com

Rakesh S. Tripathi: All India Network Project on Vertebrate Pest Management, ICAR-Central Arid Zone Research Institute, Jodhpur-342003, India

**Table 1:** Collection localities of *Mus booduga* in Leh-Ladakh region (Jammu and Kashmir).

| Villages       | Elevation (m) | Latitude         | Longitude        | General habitat   |
|----------------|---------------|------------------|------------------|---|
| Stakmo         | 3768          | 34° 2' 8.02" N   | 77° 43' 16.49" E | Crop fields ( <i>Hordeum vulgare</i> , <i>Triticum aestivum</i> , <i>Phragmites australis</i> ) |
| Sankar         | 3572          | 34° 10' 26.58" N | 77° 35' 7.61" E  | Crop fields ( <i>Hordeum vulgare</i> , <i>Triticum aestivum</i> , <i>Phragmites australis</i> ) |
| Saboo          | 3550          | 34° 7' 58.36" N  | 77° 38' 0.83" E  | Crop fields ( <i>Hordeum vulgare</i> , <i>Triticum aestivum</i> , <i>Phragmites australis</i> ) |
| Stakna         | 3288          | 34° 0' 18.13" N  | 77° 41' 6.54" E  | Crop fields ( <i>Hordeum vulgare</i> , <i>Triticum aestivum</i> , <i>Phragmites australis</i> ) |
| Thiksey        | 3259          | 34° 3' 24.57" N  | 77° 40' 1.06" E  | Crop fields ( <i>Hordeum vulgare</i> , <i>Triticum aestivum</i> , <i>Phragmites australis</i> ) |
| Chuchot Gongma | 3258          | 34° 2' 8.02" N   | 77° 43' 49" E    | Crop fields ( <i>Hordeum vulgare</i> , <i>Triticum aestivum</i> , <i>Phragmites australis</i> ) |
| Shey           | 3250          | 34° 4' 23.45" N  | 77° 38' 21.45" E | Crop fields ( <i>Hordeum vulgare</i> , <i>Triticum aestivum</i> , <i>Phragmites australis</i> ) |
| Choglamsar     | 3236          | 34° 6' 42.36" N  | 77° 35' 15.83" E | Crop fields ( <i>Hordeum vulgare</i> , <i>Triticum aestivum</i> , <i>Phragmites australis</i> ) |
| Phey           | 3187          | 34° 8' 0.55" N   | 77° 27' 57.22" E | Crop fields ( <i>Hordeum vulgare</i> , <i>Triticum aestivum</i> , <i>Phragmites australis</i> ) |
| Leh city       | 3407          | 34° 9' 9.30" N   | 77° 34' 37.39" E | Godowns (stored rice and wheat grain)   |

outnumbered the females (17) in the collection (sex ratio: 1.59:1.0). All the trapped animals were released to the site of capture after recording their body weight (g), external measurements (mm) and diagnostic characteristics. However, one female specimen was sent to Zoological Survey of India (ZSI) for confirmation of identification. The specimen was identified as *M. booduga* (Gray 1837) and deposited as National Zoological Collection vide registration No. V/3582, at Desert Regional station (DRS), ZSI, Jodhpur (Rajasthan), India. We provide here a description and measurements of its diagnostic characteristics.

The external measurements of the specimens are as follows: (mean  $\pm$  standard error of mean): head body (HB): ♂ 69.88  $\pm$  7.11 mm ♀ 73.23  $\pm$  4.49 mm; tail length (TL): ♂ 69.48  $\pm$  8.03 mm ♀ 73.29  $\pm$  5.87 mm; hind foot (Hf): ♂ 16.52  $\pm$  1.01 mm ♀ 16.47  $\pm$  1.12 mm; ear (E): ♂ 10.41  $\pm$  1.05 mm ♀ 11.05  $\pm$  1.20 mm.

The cranial measurements obtained for specimen V/3582 (one female) are: occipitonasal length (onl): 17.5 mm; condylobasal length (cbl): 16.9 mm; length of nasals (nasl): 6.5 mm; palatal length (pal): 10.7 mm; length of maxillary tooth row (mtr): 3.8 mm; length of diastema (dial): 3.1 mm; length of tympanic bulla (bl): 3.2 mm; length of anterior palatal foramina (apf): 3.2 mm; zygomatic width (zw): 10.3 mm; interorbital width (iw): 3.1 mm; mandibular length (ml): 7.2 mm.

Other characteristics: the body weight of *Mus booduga* in the present collection showed no significant variation between sexes. The pool data on body weight ranged between 15 and 25 g (mean  $\pm$  standard error of mean: 17.06  $\pm$  0.68 g). TL was 100% or slightly less than HB length. Dorsal surface was dark brown with soft fur and the ventral surface white in colour with a bicoloured tail (dark above and light below). Forelegs and hind feet were with four and five toes, respectively, all bearing claws.

All these characteristics and measurements fit within the original diagnostic characteristics of *Mus booduga* except that the HB of the newly collected specimens is slightly larger than the mean HB reported in the literature (i.e. mean HB: 69 and 73 mm in present collection against 56–63 mm as reported by Agrawal 2000). But for the other external characteristics the specimens fit within the range variation of the species. The new collected specimens of *M. booduga* from a cold arid region differed in the colour of their dorsum (dark brown) compared to those inhabiting hot arid regions viz., Bikaner, Jodhpur, Pali, Sirohi in Rajasthan (light brown or sandy colour), however the venter in both the collections were white.

Mandal (1984) reported the highest altitudinal record of *Mus booduga* from Nelang (c 3695 m) of the Himalayan region in Uttar Kashi District, Uttarakhand. However, in the present study, the mice have been recorded from altitudes ranging from 3187 to 3768 m above mean sea level, which may be regarded as the highest altitudinal distribution record of mice in the Himalayan region. From the observations it may be concluded that croplands and field stores provide an ideal habitat for Little Indian field mouse as it supplies energy rich food and safe shelter for survival in cold arid regions.

**Acknowledgements:** The authors are thankful to the Director of ICAR-Central Arid Zone Research Institute (CAZRI), Jodhpur for providing facilities and support in conducting the study. We are also thankful to Mr. Ramesh Chand Meena, and Mr. Surjeet Singh, Senior Technical Officers, CAZRI, Jodhpur for their technical support during the course of study. We would like to thank Dr. S.S. Talmale, Assistant Zoologist, Zoological Survey of India (ZSI), Jabalpur, India for the identification of specimen, and Dr. Sanjeev Kumar, Scientist D, ZSI, Jodhpur, India for registration of voucher specimen.

## References

- Agrawal, V.C. 1980. Murids of northeastern India. Studies on rodents and their control. The Meghalaya Science Society, Shillong. pp. 22.
- Agrawal, V.C. 2000. Taxonomic studies on Indian Muridae and Hystricidae (Mammalia Rodentia). Occas. Pap. No. 180. Zoological Survey of India, Calcutta. pp. 186.
- Aplin, K., S. Molur and P.O. Nameer. 2016. *Mus booduga* (errata version published in 2017). The IUCN Red List of Threatened Species 2016: e.T13953A115115984. <http://dx.doi.org/10.2305/IUCN.UK.2016-3.RLTS.T13953A22403517.en>.
- Birah, A., R.S. Tripathi and A.M.K. Mohan Rao. 2012. New report of little Indian Field mouse, *Mus booduga* (Gray) from Andaman and Nicobar Islands. J. Plant. Crops 40: 149–151.
- Biswas, B. and K.K. Tiwari. 1966. Taxonomy and distribution of Indian rodents. In: (D.W. Parrack, ed.) Proceedings of the Indian Rodent Symposium hosted by Johns Hopkins University, CMRT, and USAID, Calcutta. pp. 9–23.
- Blanford, W.T. 1891. The fauna of British India including Ceylon and Burma: Mammalia. Taylor and Francis, London. pp. 617.
- Chakraborty, S. 1983. Contribution to the knowledge of the mammalian fauna of Jammu and Kashmir, India. Occas. Pap. No. 38. Zoological Survey of India, Calcutta. pp. 129.
- Chesemore, D.L. 1970. Notes on the mammals of southern Nepal. J. Mammal. 51: 162–166.
- Ellerman, J.R. 1961. The Fauna of India including Pakistan, Burma and Ceylon. Mammalia, Vol. 3 (Rodentia), Part I and II. Govt. of India, Delhi. pp. 884.
- Hinton, M.A.C. and H.M. Lindsay. 1926. Bombay Natural History Society's Mammal Survey of India, Burma and Ceylon. Report No. 41 (Assam and Mishmi Hills). J. Bomb. Nat. Hist. Soc. 31: 383–403.
- Idris, M., R.S. Tripathi, V. Chaudhary and B.D. Rana. 2003. Incidence of Indian field mouse, *Mus booduga* in Thar Desert. In: (P. Narain, S. Kathju, M.P. Singh and Praveen-Kumar eds.) Human impact on desert ecosystem. Arid Zone Research Association of India and Scientific Publisher, Jodhpur. pp. 343–347.
- Lindsay, H.M. 1926. Bombay natural history society's mammal survey of India, Burma and Ceylon. Report No. 44 (Kangra & Chamba). J. Bomb. Nat. Hist. Soc. 31: 597–606.
- Mandal, A.K. 1981. Report on the occurrence of the Little Indian Field Mouse, *Mus booduga booduga* (Gray) in Arunachal Pradesh, India. Bull. Zool. Surv. India 3: 271–272.
- Mandal, A.K. 1984. A note on the distribution of the little Indian field mouse, *Mus booduga* (Gray) (Rodentia; Muridae), with an altitudinal record. Bull. Zool. Surv. India 6: 301–303.
- Posamentier, H. 1989. Rodents in agriculture. A review of findings in Bangladesh. Deutsche Gesellschaft für Technische Zusammenarbeit (GT 2) GM bh. Eschborn. pp. 107.
- Pradhan, M.S. and S.S. Talmale. 2009. List of valid Rodent taxa (Class: Mammalia, Order: Rodentia) from Indian Subcontinent region including Myanmar. Occas. Pap. No. 297. Zoological Survey of India, Kolkata. pp. 239.
- Prakash, I., A.P. Jain and B.D. Rana. 1971. New records of rodents from the Rajasthan desert. J. Bombay Nat. Hist. Soc. 68: 447–450.
- Roberts, T.J. 1977. The Mammals of Pakistan. Ernest Benn Ltd., London & Tonbridge. pp. 361.
- Sharma, B.D. and T. Sharma. 1976. Some mammals records for Jammu Province (J & K State). Indian Poul. Rev. 8: 15–17.
- Shimada, T., K.P. Aplin and H. Suzuki. 2010. *Mus lepidoides* (Muridae, Rodentia) of Central Burma is a distinct species of potentially great evolutionary and biogeographic significance. Zool. Sci. 27: 449–459.
- Varma, R.N. 1968. A study of ectoparasites of mammals in the Jammu-Sialkot and Lahore sectors. Armed Forces Med. J. 23: 71–82.
- Wilson, D.E. and D.A.M. Reeder. 2005. Mammal Species of the World. A taxonomic and geographic reference, 3rd ed. Johns Hopkins University Press, Baltimore, Maryland. pp. 2142.
- Wroughton, R.C. 1914. Bombay Natural History Society's Mammal Survey of India, Burma and Ceylon. Report No. 15 (Kumaon, U. P.). J. Bomb. Nat. Hist. Soc. 23: 282–301.