



Inheritance of moderate pod-reticulation in groundnut (*Arachis hypogaea* L.)

M. Y. Samdur, P. Paria, P. Manivel, T. Radhakrishnan, R. K. Mathur and H. K. Gor

National Research Centre for Groundnut, Post Box No. 5, Junagadh 362 001

(Received: January 2002; Revised: July 2002; Accepted: August 2002)

Pod characteristics like reticulation, constriction, beak, pod size etc. are important traits associated with identification of varieites in groundnut. The reports in literature show that prominent reticulation of the pods was reported to be dominant to absence of pod reticulation [1-4]. Badami [1] hypothesized at least four factors for deep reticulation whereas Manoharan and Ramalingam [4] reported two factors for the same. Using three crosses involving parents with pods with moderate reticulation and with smooth pod, Radhakrishnan et al. (personal communication) proposed a digenic inhibitory gene action system for pod reticulation.

The cultivar GG 2, an advanced breeding line PBS 12143 and the germplasm line NCAc 343, all having pods with moderate reticulation, were used as the female parents in this study. The cultivar JL 24 used as the male parent, has smooth pods (absence of reticulation). The F₁'s were raised in medium black calcareous soil (vertisol) during *kharif* 1999 at National Research Centre for Groundnut, Junagadh. The F₁ plants were harvested individually and were raised in plant to progeny rows in F₂ during *kharif* 2000. Pods were critically observed for presence or absence of reticulation (score 0 for no reticulation and 5 for moderate reticulation on 0 to 9 scale) as per the descriptors for groundnut [5]. The goodness of fit between the observed and expected segregation pattern in different crosses

in F₂'s was tested by using χ^2 test (Table 1).

In all the crosses F₁'s produced smooth pods. It is evident from the F₂ segregation data of these crosses that the segregation of smooth pod vs. reticulation fit well to the 13:3 ratio in all the three F₂'s. Out of a total of 259 progenies over the families, 207 produced smooth pods and rest 52 produced reticulated pods. Thus, overall segregation fitted well to ratio 13 smooth pods: 3 reticulated pods. Therefore, it is evident that the reticulation of pods in the parents studied is under the control of digenic inhibitory gene action. Patil [2] found monogenic inheritance of pod reticulation in groundnut and proposed the gene symbol *rp* for the locus. However, the present study suggests the presence of an inhibitor gene in the control of reticulation. Hence, the gene symbol *rp* proposed for reticulation [2] locus may be retained and the symbol of the inhibitory gene for the character may be proposed as *I'/i'*. Thus, the common male parent (JL 24) used in this study having genotype of *I' I' rrp* had smooth pod. Similarly the genotypes of the three reticulated parents used were *I' I' RpRp*.

References

1. **Badami V. K.** 1928. *Arachis hypogaea* (groundnut). PhD thesis, Cambridge University. Cited in H. Hunter and H. M. Leeke (1933). Recent Advances in Agric. Plant Breeding, Blakiston, Philadelphia.
2. **Patil V. H.** 1965. Genetic studies in groundnut (*Arachis hypogaea* L.). MSc (Ag) thesis, Poona University, India.
3. **Murthy T. G. K., Tiwari S. P. and Reddy P. S.** 1988. A linkage group for genes governing pod characters in groundnut. *Euphytica*. **39**: 43-46.
4. **Manoharan V. and Ramalingam R. S.** 1992. Inheritance of testa colour and pod reticulation in groundnut. *Madras Agric. J.* **79**: 646-648.
5. **IBPGR and ICRISAT.** 1992. Descriptors for groundnut, International Board for Plant Genetic Resources, Rome, Italy; International Crops Research Institute for Semi-Arid Tropics, Patancheru, India.

Table 1. F₂ segregation of pod reticulation in groundnut

| Cross | F ₁ | F ₂ progeny | | | χ^2 (13:3) | P |
|-------------------|----------------|------------------------|------------------------|-------|--------------------|-----------|
| | | Smooth | Moderately reticulated | Total | | |
| GG 2 × JL 24 | Smooth | 126 | 32 | 158 | 0.234 | 0.50-0.95 |
| PBS 12143 × JL 24 | Smooth | 37 | 14 | 51 | 2.530 | 0.10-0.20 |
| NC Ac 343 × JL 24 | Smooth | 44 | 6 | 50 | 1.490 | 0.20-0.30 |
| Total | | 207 | 52 | 259 | 0.299 | > 0.95 |
| Heterogeneity | | | | | 3.955 | 0.10-0.20 |