



Journal of Experimental Biology and Agricultural Sciences

<http://www.jebas.org>

ISSN No. 2320 – 8694

EFFECT OF *Morinda citrifolia* IN GROWTH, PRODUCTION AND IMMUNOMODULATORY PROPERTIES IN LIVESTOCK AND POULTRY: A REVIEW

Jai Sunder*, Tamilvannan Sujatha and Anandamoy Kundu

Division of Animal Science, ICAR-Central Inland Agricultural Research Institute, Port Blair, A and N Islands 744105

Received – April 18, 2016; Revision – May 02, 2016; Accepted – May 21, 2016

Available Online – May 25, 2016

DOI: [http://dx.doi.org/10.18006/2016.4\(3S\).249.265](http://dx.doi.org/10.18006/2016.4(3S).249.265)

KEYWORDS

Morinda citrifolia

Livestock

Poultry

Growth

Production

Immunomodulatory
Properties

ABSTRACT

Morinda citrifolia L. is commonly known as Noni and has been found to have wide range of medicinal properties. It is usually found in the coastal region in many countries including Andaman and Nicobar Islands and belongs to the family *Rubiaceae*. This small evergreen tree is widely grown and well adapted to the tropics and can grow in fertile, acidic, alkaline and saline affected soils. It tolerates high soil salinity and brackish water stagnation. All the components of this plant have high demand in case of alternative medicines and herbal medicines. Due to its high demand and as a source of revenue generation the detail study on its nutritional benefits and therapeutic values are essential for its commercial exploitation. More than 200 nutraceutical compounds have been identified from the plant. *Morinda citrifolia* is reported to have broad spectrum biological activities such as antimicrobial, immunomodulatory, antioxidant wound healing etc. Apart from the *in-vitro* scientific validation of the activities and *in-vivo* trial in some lab animal model, the plant has been used for livestock and poultry health and production. A lot of reviews have been written on the different uses of Noni, however, scientific review on the use of this plant on the growth, production, immunomodulator and other pharmacological activities of *M. citrifolia* in livestock and poultry has not been compiled. Therefore this review discusses the compilation of the work done on the use of *M. citrifolia* in livestock and poultry.

* Corresponding author

E-mail: jaisunder@rediffmail.com (Jai Sunder)

Peer review under responsibility of Journal of Experimental Biology and Agricultural Sciences.

Production and Hosting by Horizon Publisher India [HPI]
(<http://www.horizonpublisherindia.in/>).
All rights reserved.

All the article published by Journal of Experimental Biology and Agricultural Sciences is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License Based on a work at www.jebas.org.



1 Introduction

Livestock farming is considered to be a profitable complementary enterprise in agriculture and constitutes an important activity for accelerating the rural economy. They are being reared under small open ranging system to large scale intensive system in our country. This sector has been growing at steady pace; however, during the recent years due to indiscriminate usages of antibiotics in livestock and poultry the problem of multidrug resistance has evolved (Woolhouse et al., 2015). The high dosage of antibiotics, hormones and chemical derivatives for increase and sustain livestock production has increased the problem of its residual effects in their products thus causing a serious health concern (Landers et al., 2012). The organic farming has changed the scenario of the present agricultural production system; similarly the concept of organic livestock production system has been evolved. The use of traditional medicine and alternative medicine has been practiced by the farmers for sustainable production (Galav et al., 2013; Luseba & Tshisikhawe, 2013).

A large number of medicinal plants with known medicinal properties are available and is being used by the farmers (Chinsembu et al., 2014; Verma, 2014; Luseba & Tshisikhawe, 2015). *Morinda citrifolia* is also known as Noni, belongs to the family Rubiaceae and is mostly available in the coastal region (Nelson, 2006). It is commonly known as Indian mulberry, Ba Ji Tian, Nono or Nonu, Cheese fruit and Nhau in various countries throughout the world (Bruggnecate, 1992; Whistler, 1992; Solomon, 1999; Chan-Blanco et al., 2006). In these islands it is commonly known as Lorang, Burmaphal, Pongee phal and Surangi by the tribal. *M. citrifolia* has a rich history in India, where it has been used for tens of centuries in the system of medicine known as Ayurveda. This small evergreen tree or sprint (10-20 ft) is native to India and also distributed to south-eastern Asia to Australia and now has tropical distribution widely adapted to the tropics (Dixon et al., 1999; Ross, 2001). It can grow in challenged environments viz, saline, acidic and alkaline soils. *M. citrifolia* has been known for its wide range of medicinal properties (Younos et al., 1990; Bruggnecate, 1992; Hiramatsu et al., 1993; Hirazumi et al., 1996; Solomon, 1999; Brown, 2012; Assi et al., 2015). Reports are available on the scientific studies of this plant and wide health benefits. The plant is also suitable for saline affected lands, owing to its saline resistant properties the plant can be grown as an alternative crop for the saline affected areas and its high demand both at national and international market, the studies on its phytochemicals and its effect on production and immune response is utmost importance.

Polynesians used the whole plant for treatment of various kinds of illness as herbal remedies (Earle, 2001). Various reports are available for use of this plant for treatment of illness such as diabetes, blood pressure, cancers, arthritis, poor digestion etc. (Singh et al., 1984; Bruggnecate, 1992; Solomon, 1999; Whistler, 1992; Wang et al., 2002; Brown, 2012; Lee et al., 2012; Fletcher et al., 2013; Saminathan et al., 2013a;

Saminathan et al., 2014b; Sharma et al., 2016). However, there is less information on the scientific validation of the *Morinda* fruits viz. treatment of colds and influenza reported by Solomon, 1999. Allen (1873) reported some information on the ethno botanical properties, which is considered to be one of the earliest report on the medicinal use of morinda.

The history of use of *M. citrifolia* for livestock and poultry is very limited. Due to its strong smell and taste many animals do not consume the product and avoid contact with the fruit and seeds. The residents of South Pacific islands have noted health benefits for themselves and their animals by ingesting the *M. citrifolia* fruit (Whistler, 1985). Some animals such as pigs consume the fruit in its natural state (Fugh-Berman, 2003). Most animals have difficulty in consuming and digesting whole fruit. The available literatures revealed that the leaves are used for livestock fodder (eg. Niue, India) and to feed silk worms (India). Over the year's research have been mainly carried out on use of fruit and leaf extract. Studies have been carried out on anthelmintic property (Raj, 1975; Morton, 1992), anti-inflammatory (Basar et al., 2010; Fletcher et al., 2013), antitumor potential (Hirazumi & Furusawa, 1999; Furusawa, 2002; Brown, 2012; Jinhua et al., 2013; Saminathan et al., 2014a), utilization of its juice extract waste in diets of goats (Aregheore, 2005), enhanced mononuclear phagocytic activity of gram-negative bacteria in calves fed with the fruit puree (Schafer et al., 2008). Researchers have been carried out on the volatile components of ripe fruits and their effects on drosophila (Farine et al., 1996). Studies on the effect of its fruit and leaf extract in the total serum protein, creatinine and albumen level of poultry have been carried out (Sunder et al., 2011a). However, the scientific evidence of its use in animal model and its effect on production system is very rare (Fugh-Berman, 2003; Sunder et al., 2007; Sunder et al., 2011b; Sunder et al., 2011b c). In the present review, information on the available literature on the use of *M. citrifolia* in livestock and poultry has been documented.

2 Growth and production promoting properties in poultry

The growth performance ability of the *M. citrifolia* fruit was tested in Nicobari fowl; an indigenous poultry bird of Andaman & Nicobar Islands, India and broiler (Sunder et al., 2011b). Broilers were given fresh noni fruit juice (1.5 ml/bird/day). The performance of the morinda fed group was found more than the control birds in terms of body weight gain, feed conversion ratio and feed efficiency. The overall performance index of Morinda fed group was found to be superior (93.6±16.15) than other groups. Egg production was also recorded to be high in the morinda fed group (95.24%) than in control (83.11 %). Similarly, highest dressing % was obtained in the Morinda fed group (69.05%) than in control (68.38%). The highest body weight gain during the 3rd and 4th month of age was observed due to the growing phase of the bird during which the bird attained more body weight gain than compared to the other phase of the growth. The reports of Morinda on the growth was reported by Sunder et al. (2007) in

broiler birds, however, no reports are available on the effect on the egg production in the Nicoabri fowl.

Studies on feeding of *M. citrifolia* fruit extract to Japanese quail showed better body weight gain, feed conversion ratio and performance index in Morinda fed group than in control group. The overall results revealed that higher body weight gain in Morinda fed group (162.11 ± 0.06 g) compared to control group (153.005 ± 1.05 g) for 0 to 7 weeks of age. FCR of the Morinda fed group (5.99 ± 0.17) was recorded better than control group (6.18 ± 0.16). The feed efficiency of Morinda fed group (0.22 ± 0.05) was also found to be higher ($p < 0.05$) than control group (0.20 ± 0.08). The hen day egg production was recorded better in the Morinda fed group when compared to the control group. The overall analysis of the growth and production performance of both the groups revealed that the *M. citrifolia* crude fruit extract fed @ 5% daily enhanced the body weight gain and the egg production performance of the Japanese quail (Sunder et al., 2013d).

3 Growth, production and immunomodulatory properties in livestock

Ethno veterinary application of noni fruit puree has been studied in the livestock. Brooks et al. (2009) demonstrated that feeding of noni puree enhanced the immunity of neonatal calves and potentially long term health especially in the preweaning stage. In earlier findings, Brooks et al. (2009) reported that supplementation of juice from *M. citrifolia* enhanced the activation of CD4+ and CD8+ T cells in neonatal calves. Bactericidal effect of noni was evaluated via an ex vivo whole blood bactericidal assay by Schäfer et al. (2008). Result showed that noni fed group showed significantly more killing power at day 14 when compared to control calves. Advantages of supplementing Morinda Max have been demonstrated in newly received cattle. Yancey et al. (2013) studied the growth performance effect of feeding of noni in cattle. They have reported that feeding of pulp of *M. citrifolia* to cow at the level of up to 2% in the diet increased body weight gain with better feed conversion ratio. The gain was increased with the increase in concentration of Noni pulp in the diet. Ponce et al. (2011) found that feeding of Noni extract to calf increased the growth performance as well as the immunomodulatory properties. The immune enhancing effect including the antibacterial, anti-inflammatory, anti cancer and anti oxidant effect of noni has been validated (Pawlus et al., 2005; Kusirisin et al., 2009; Nitteranon et al., 2011). Presence of Iridoid and polysaccharide fractions of noni has been shown to induce the release of several immune mediators, many of which have beneficial stimulatory effects and may help in the maturation of the neonatal immune system (Bui et al., 2006; Hirazumi & Furusawa, 1999; Deng et al., 2010).

The growth promotion effect of *M. citrifolia* juice may be due to its rich nutrients value which contain all the essential amino acids, minerals, vitamins and other nutrients which are required for the growing cells (Chunhieng, 2003) It is very rich in proxeronine which is believed to be a precursor to

xeronine which helps in activation of xeroninase (Heinicke, 2001). As the Morinda fruit contain several amino acid, vitamins, minerals, co-enzymes, carbohydrates and alkaloids which directly or indirectly help in metabolism of the nutrients and help in overall growth of the cell and tissues (Chan Blanco et al., 2006; Takashima et al., 2007; Assi et al., 2015). More than 200 nutraceutical compounds have been identified from the plant (Solomon, 1999; Singh, 2012). Leaf and fruit of this plant are reported to be used as feed for livestock and poultry (Fugh-Berman, 2003). However, further investigation on several bioactive compounds present in the *M. citrifolia* will help in understanding the actual mechanism in detail and its use in livestock and poultry as a source of vitamins and minerals supplement for higher growth, production and immunity.

3.1 Anticholesterol properties

Cholesterol is very important compounds used for many physiological functions; however, the unwanted increase in the level of bad cholesterol is the predisposing factors for many of the diseases. Lot of works have been carried out to lower the level of cholesterol in chicken meat by using additives, feed supplements, etc. (Chowdhury et al., 2002; Yalcin et al., 2006; Aydin et al., 2008; Azeke & Ekpo, 2009; Saki et al., 2014). Considering the importance of the cholesterol content the study was conducted to see the effect of *M. citrifolia* fruit and leaf extract supplementation on the blood cholesterol level. A group of 100 poultry birds were fed 5% crude fruit extract and leaf extract of *M. citrifolia* daily along with the normal basal ration. Studies on broilers and Nicobari fowl revealed that feeding of fruit extract (5%) and leaf extract (5%) daily along with the normal basal ration lowered the blood cholesterol level. The cholesterol level in the Morinda fruit extract fed group at 4th week (179 ± 12.3) and at 6th week (201 ± 9.4) was found to be lower than control group (185 ± 11.4) and (233 ± 10.5), respectively. The cholesterol level in the birds fed with leaves extract was found to be slightly higher than fruit extract. The cholesterol level in the Morinda leaf extract fed group at 4th week were (190 ± 11.1) and at 6th week were (210 ± 5.4) was again found to be lower than the control group (222 ± 12.3) and (248 ± 8.1) respectively. The result revealed that the feeding of *M. citrifolia* extract daily in the poultry ration has lowered the level of cholesterol in the blood serum (Sunder et al., 2011a). Reports suggest that the Noni has phytochemicals and beta-sitosterol, a plant sterol with potential for anti-cholesterol activity (Palu et al., 2006; Wang et al., 2006). Research on lowering the cholesterol level in chicken meat and egg is going on worldwide. Research showed that cholesterol content may vary in eggs and blood and lot of work is being carried out on chicken eggs and meat either through the use of additives, dietary fibre, polyunsaturated fatty acids supplementation etc (Barroeta, 2007; Dalkilic et al., 2009). Recent study showed that feeding of morinda fruit extracts to calves reduced the level of total cholesterol, triglycerides, LDL-cholesterol in the morinda fed group than in the control group (Anantharaj et al., 2016).

3.2 Effect on blood biochemical profile of poultry

Concentration of serum protein is useful in monitoring various disease status. It increases during dehydration, multiple myeloma and chronic liver diseases, and decreases in renal diseases and liver failure. Creatinine is a waste product formed in the muscle from the high energy storage compound creatinine phosphate. It is useful indicator of renal function and increases in various renal diseases. Albumin is a plasma protein synthesized in liver from amino acids absorbed from ileum. It increases during dehydration and stasis during venipuncture. It decreases during excess protein loss and decreases synthesis due to dietary, hepatic disease or malabsorption (Babatunde et al., 1992). Lot of reports are available on the effect of feeding of herbal plants and its extracts on the blood biochemical profile of poultry, (Langhout, 2000; Kamel, 2001; Elagib et al., 2012; Hosseinzadeh et al., 2014; Kant et al., 2014). However, very few researches have been carried out to see the effect of feeding of noni in the blood biochemical profile of poultry. Sunder et al., 2011a found that feeding of noni decreases the level of total protein, serum creatinine and albumin in the poultry. There is no report available on the effect of *M. citrifolia* in blood biochemical parameters of poultry, however, similar to this study, Polat et al., 2011 found that feeding of rosemary reduces the blood cholesterol level and increases the creatinine level. Creatinine is a chemical waste molecule that is generated from muscle metabolism. The lower values indicates that no muscular wastage which might have been possibly cause by inadequacy of protein in animals. In the present study also the level of blood protein was found to be low. However, Ghazalah & Ali (2008) found that creatinine levels were all reduced by dietary rosemary leaves compared to controls. It is useful indicator of renal function and increase in various renal diseases. Albumin is synthesized in the liver from amino acids absorbed from the ileum. It increases during dehydration and decreases during excess protein loss. The level of total serum protein, albumin and creatinine was found to be better in the *Morinda* fed group. The effect of *M. citrifolia* on various biochemical parameters may be useful in monitoring physiological status and disease status as well as therapeutic evaluation of the products. Research showed that the feeding of *M. citrifolia* juice enhanced the immune response in poultry (Sunder et al., 2007).

3.3 Antimastitis properties in cattle

Mastitis is a serious problem of the dairy cows and estimated loss due to direct and indirect losses has been estimated to the tune of \$35 billion annually (Mubarack et al., 2011). Infection of the cow's udder and the mastitis is considered as one of the major constraints for growth of dairy industry worldwide (Sasidhar et al., 2002; Osteras & Solverod, 2009). Due to development of multidrug resistance bacteria the use of herbal based preparation for treatment of mastitis and other diseases have been reported in livestock (Virmani et al., 2010; Kalayou et al., 2012; Mir et al., 2014; Zeedan et al., 2014). Several plants have been reported to be used for treatment of mastitis

viz. *Capsicum annuum*, *Lepidium sativum*, *Allium sativum*, *Sesamum indicum*, *Citrus limon*, *Zingiber officinale*, *Citrullus colocynthis*, *Curcuma longa*, *Amomum subulatum*, *Sesamum indicum*, *Cuminum cyminum*, *Rosa indica*, *Centratherrum anthelmisticum*, *Triticum aestivum*, *Nigella sativa* and *Peganum harmala* (Dilshad et al., 2009). In many of the reports, the antibacterial activity of the plant extracts were described against the organisms responsible for mastitis, however, Sunder et al. (2013a) reported the treatment of mastitis by feeding of *M. citrifolia* in dairy cows. Sunder et al. (2013a) observed that oral feeding of noni fruit juice (100 ml) to dairy cows is effective for treatment of mastitis. In the mastitis milk, level of electrical conductivity increases due to changes in ionic concentration which resulted in the decrease in the level of lactose and K^+ and increase in the level of Na^+ and Cl^- (Kitchen, 1981; Kaşikçi et al., 2012). Sunder et al. (2013a) found that feeding of *M. citrifolia* decreased the electrical conductivity from 5.5 mho to less than 5.0 mho, which resulted in decrease in the milk pH (7.0 to 6.6) thus reducing the total microbial load in the milk. Level of microbial load was also decreased from 5.13×10^8 to 3.54×10^8 . Among the normal cow no change was observed in the pH of cow's milk with respect to different teat of the cow (Batavani et al., 2007). In case of mastitis Electrical conductivity will vary in the range of 5.58 mS. It is concluded that the feeding of *M. citrifolia* reduced the milk pH, electrical conductivity, microbial load in the mastitis affected milk.

4 *M. citrifolia* as feed supplement

Plants are being used as a source of feed and feed additives in animal feeding since time immemorial. During the recent years due to ban of most of the antibiotic growth promoters use of medicinal plants as a feed additives, supplements for the livestock and poultry have been increased (Charis, 2000; Tipu et al., 2006; Mirzaei-Aghsaghali, 2012; Eevuri & Putturu, 2013; Mirzaei & Venkatesh, 2012). Use of *M. citrifolia* fruit as a feed ingredient was studied in Japanese quail (Sunder et al., 2013b,d). *M. citrifolia* fruits were cut into small pieces and sun dried. The dried fruits were grounded to make it in the form of small granules. Fruit granules were added in the quail ration up to 15% by replacing maize, rice bran and wheat by 5% each. The body weight at 2nd, 3rd and 5th week of age was higher significantly ($p < 0.05$) in the *Morinda* fed group compared to the control group. Feed conversion ratio and egg production was found to be better in *Morinda* fed group. The overall production performance revealed that at the end of 100 days of egg production the *Morinda* fed group showed 24% more egg production than the control group. The study showed that *M. citrifolia* can be used as feed ingredient in Japanese quail ration to the tune of 15% on dry matter basis. In another study, Japanese quail were fed with fruit granules of *M. citrifolia*, 20% (w/w) as replacement of the normal concentrate mixture in the ration (Sunder et al., 2013 b). Results revealed the higher body weight gain in *Morinda* group (109.4 ± 7.22) than in control group (106.8 ± 6.65) at 5th week of age. Egg production was also found better in *Morinda* group (59.34 ± 12.31) than in control group (56.80 ± 10.71). It is

concluded that part-replacement of quail ration with dried ripe fruit granules of *M. citrifolia* (20%, w/w) would be cost-efficient in quails with no apparent side effects. This is the first report of use of *M. citrifolia* fruit granules as feed supplement in the poultry. There is no report available on the use of *M. citrifolia* fruit pulp as a feed for poultry. However, some reports are available on the use of leaf and fruit of this plant as feed for livestock and poultry (Fugh-Berman, 2003). Use of other medicinal plants viz. Curcuma longa, turmeric, Ocimum sanctum, tulsi, Aloe vera, were also reported to enhanced the body weight gain, better FCR, feed efficiency in poultry (Kumar et al., 2005; Durrani et al., 2006; Lanjewar, 2008; Al-Kassie et al., 2011; Eevuri & Putturu, 2013). The proximate analysis of the noni fruit pulp revealed that the crude protein content is only 5.8%, however, it is very rich in all amino acids, micro and macro minerals and vitamins which are essential for the vital functioning of the cells/tissues for growth and production. The *M. citrifolia* fruit is very rich in the nutraceutical compounds and contains rich amounts of minerals like K, Ca, Mg, Fe, Cu and Mn (Singh et al., 2008). High egg production and growth promoting effect may be due to the nutraceutical compounds and minerals present in the fruit which might have played important role in enhancing the growth and production. The earlier reports with the use of *M. citrifolia* fruit juice revealed that the supplementation of *M. citrifolia* juice @ 5% enhanced body weight gain in broiler and Japanese quail (Sunder et al., 2007, Sunder et al., 2011a). Morinda fruit contain several amino acids, vitamins, minerals, co-enzymes, carbohydrates and alkaloids which directly or indirectly help in metabolism of the nutrients and help in overall growth of the cell and tissues hence in the Morinda fed group the overall performance was better than in control group. The studies on evaluation and utilization of Noni juice extract waste (NJEW) in goat's diet revealed that the unusual taste and low degradability of essential nutrients may be the factors limiting the use of NJEW in ruminant diets. Therefore, research on its palatability quality is required so that the extracts and the fibres may be effectively utilized as a source of livestock food (Aregheore, 2005)

5 *M. citrifolia* as Immunomodulator

Immune system of poultry generally benefits from the medicinal plants and herbs which are rich in flavonoids, vitamin C and carotenoids. During the last decade there has been substantial increase in the use of medicinal herbs as feed supplement, immunostimulant and growth promoters. Many plants are available which are very rich in these compounds including the *M. citrifolia*. There are several reports available with the use of medicinal herbs for immunostimulants in the poultry (Kumari et al., 1994; Emadi & Kermanshahi, 2007; Durrani et al., 2008; Lee et al., 2010; Mahima et al., 2012; Mirzaei-Aghsaghali, 2012; Dhama et al., 2015). Phytochemical analysis of the *M. citrifolia* revealed the presence of several compounds viz. carbohydrates, gums and mucilages, proteins, amino acids, fats and oils, anthraquinone glycosides, coumarin glycosides, flavonoids, alkaloids, tannins, phenolic compounds and citric acid which are

responsible for several activities including the immunostimulation (Nayak et al., 2012). Studies showed that *M. citrifolia* stimulated the immune system for anti tumor, anticancer, antioxidant activities both in-vivo and in-vitro (Hirazumi & Furusawa, 1999; Furusawa & Hirazumi, 2003; Palu et al., 2006; Nayak & Mengi, 2010; Mufidah et al., 2013; Assi et al., 2015; Krishnaiah et al., 2015). However, very few reports are available on the use of *M. citrifolia* as immunostimulant in livestock and poultry (Sunder et al., 2007; Lohani, 2010; Sunder et al., 2011a; Benjamin et al., 2014). Studies showed that feeding 5% crude leaf extract of *M. citrifolia* to Nicobari fowl enhanced B cell mediated immune response. Furusawa et al. (2003) studied the antitumor activity in allogeneic mice and found that immunomodulatory property is due to a polysaccharide rich substance present in the fruit. Sunder et al. (2007) also studied the immunomodulatory properties of *M. citrifolia* in poultry. This was the first report of immunomodulatory properties of *M. citrifolia* in poultry. They found that feeding of *M. citrifolia* fruit juice @ 5% in water enhanced both humoral (B cell mediated) and cellular (T cell mediated) immunity in broilers. The humoral immune response of the Morinda group was significantly better than control group ($P < 0.05$). The peak response was observed at first week post inoculation (PI) in Morinda group (1.48 ± 0.18) compared to control group (0.82 ± 0.1). The direct challenge test of IBDV in the Morinda fed birds showed protection against the infection as only 6.6% mortality was recorded in this group compared to 25% in control.

Humoral immunity and cell mediated immunity of *M. citrifolia* was studied by Sunder et al. (2011b) and found that supplementation of 1.5 ml of fruit juice enhanced B cell and T cell mediated immunity in Nicobari fowl. Literatures also indicated that Noni increases the defences and reinforces the immune system of the body, neutralize its function in all the cells and regenerates the affected cells (Heinicke, 1985). Thus it helps in preventing the diseases such as IBDV infection in the present study as it increases the immune response as observed. Reports also suggests the effect of *M. citrifolia* in inducing the release of interferons, interleukins and nitric oxide (Hirazumi & Furusawa, 1999).

6 Synergistic effect of Morinda with other medicinal plant and probiotic

During the last decade lot of works have been carried out in search of alternative to antibiotics which could be safely used as growth promoter, antimicrobial without any side effect or residual compounds in the end products. Probiotics are single or mixed group of bacteria when administered to the host showed many beneficial effects viz. growth promotion, enhancing nutrient uptake from the intestine, reducing the harmful microorganism, increases the immunity (Kabir, 2009; Brisbin et al., 2010). Reports are available on the use of lactobacillus as growth promoter and probiotics in livestock and poultry (Ibrahim et al., 2005; Salarmoini & Fooladi, 2011; Zamanzad-Ghavidel et al., 2011). Similarly, beneficial properties of medicinal plants have also been reported in the

the livestock and poultry by many workers (Mishra et al., 2008; Javed et al., 2009; Narimani-Rad et al., 2011). Use of probiotics and prebiotics together showed beneficial effect showed that However, Sunder et al. (2012) and Sunder et al. (2015) studied the use of combination of *Morinda* and lactobacillus in poultry. They reported that combination of *M. citrifolia* and lactobacillus showed synergistic effect in terms of body weight gain, immunomodulatory properties, and reduction in gut microbial count and feed efficiency. This is the only report available on the combined use of *M. citrifolia* and lactobacillus in poultry.

Since the ban of antibiotics as growth promoters in poultry, the use of lactobacillus and herbal based nutraceuticals compounds have been increased. Role of nutraceuticals in improving the gut health and growth performance of poultry have been described by many workers (Muir et al., 2000; Yang et al., 2009; Zamanzad-Ghavidel et al., 2011; Adil & Magray, 2012; Das et al., 2012; Fallah et al., 2013; Sugiharto, 2015). However, Sunder et al. (2015) studied the feeding of noni and lactobacillus in broiler which was not studied earlier. They found that feeding of lactobacillus and noni juice showed synergistic effect and enhanced the reduction in gut coliform load. Antimicrobial activity of *M. citrifolia* was also reported by some workers and they found that the activity is mainly due to presence of terpenoid compounds, phenolic compounds such as acubin, alizarin, acopoletin, anthraquinones in the noni fruit (Jin et al., 1998; Lavanya & Brahmaaprakash, 2011; Narimani-Rad et al., 2011; Salarinoi & Fooladi, 2011).

Histology study of the chicken gut after feeding with noni fruit and lactobacillus showed significant changes in crypt depth and villi height, which is considered to be the main site for development of immune response and nutrient uptake (Sunder et al., 2014a). Similar to the finding of use of noni and lactobacillus, use of herbal based feed supplement as a growth promoter and effect on the gut function has also been reported by some workers (Hashemi et al., 2009; Lavinia et al., 2009; Abdulkarim et al., 2013; Kanduri et al., 2013).

7 *Morinda citrifolia* and *Andrographis paniculata* on expression of toll-like receptors

Kalmegh (*Andrographis paniculata*) a promising medicinal plant has been scientifically validated to exhibit functions such as antiinflammation, immunomodulator (Sheeja & Kuttan, 2007; Abu-Ghefreh et al., 2009; Wang et al., 2010; Shen et al., 2013; Gao & Wang, 2016). Toll like receptors (TLRs) are innate immune receptors and induce fast and appropriate host defence reaction against pathogens. TLRs recognise the conserved microbial patterns such as flagellin, LPS, peptidoglycan in an efficient and non self reactive manner to initiate pro inflammatory cytokines. Role of TLR in the immunomodulation has been demonstrated and at least 10 TLRs have been identified in chickens, including TLR1A, 1B, 2A, 2B, 3, 4, 5, 7, 15 and 21 (Barjesteh et al., 2013; St Paul et al., 2013). Sunder et al. (2014b) studied that supplementation of Noni and Kalmegh influenced the expression levels of TLR-

2, TLR-3, TLR-4, TLR-5, TLR-15 and TLR-21 significantly ($P < 0.05$). The gene expression level of TLR-3, TLR-4 and TLR-5 was found more which showed the immunomodulatory properties of herbal extract. The increased expression of TLR-3, TLR-4 and TLR-5 may be due to the effects of phytochemicals on these TLR signal transduction pathway. Andrographolide has been found to induce the APK and PI3K signaling pathway which thereby involved in macrophage activation (Rao, 2001; Fukao & Koyasu, 2003; Wang et al., 2010). Park et al. (2009) also found that production of interferon is mainly due to the presence of quercetin in the noni fruit. This was also supported with the finding of Tanabe et al., 2003 and Tohyama et al., 2005. They also found that IFN- γ plays important role in expression of TLR-3, TLR-4 and TLR-5. The high level of TLR-3, TLR-4 and TLR-5 and decreased TLR-7 gene indicated the antiviral and antibacterial activities.

In another study the effect of feeding of *M. citrifolia* fruit to broiler was done and expression of TLR-4, TLR-5, IL-8, and IL-12 was found to be more while the TLR-7 and IL-6 level was lowered. The high level of interleukins and TLR may be responsible for antiviral and antibacterial activities in the noni fruits.

8 Grommune tonic for poultry

Feeding of Grommune, a noni based herbal tonic showed more body weight gain, better FCR and enhanced B cell mediated immunity in broilers. The dose was standardized and found that feeding of Grommune @ 15 ml per bird up to 4 week and 30 ml per bird up to 8th week of age improved the body weight, feed conversion ratio and immune competency status of broilers (Sunder et al., 2014c).

9 Morical supplement for Japanese quail

Sunder et al. (2013b,c) studied the feeding of different concentration of morical, a herbal based *M. citrifolia* feed supplement in the Japanese quail. They found that supplementation of 4% (w/w) of morical in feed showed more annual egg production (238.5) compared to control (215.4). They also observed that egg shell thickness, high Ca content in egg shell, egg yolk content, total weight of the egg increased with the increasing concentration of morical supplementation up to 8% in the feed.

10 Anticancer Activity

Antitumor activity of the noni fruit was reported due to the presence of a polysaccharide rich substance (Hirazumi et al., 1994). Reports also suggests the role of *M. citrifolia* as anticancer and antitumor properties (Liu et al., 2001; Wang et al., 2001; Wang & Su, 2001; Wong, 2004; Issell et al., 2009; Brown, 2012; Saminathan et al., 2013a; Wu et al., 2015). Later, a lot of studies were carried out to find out the compounds responsible for the antitumor and anticancer activities directly or indirectly (Hiramatsu et al., 1993; Hisawa et al., 1999; Hirazumi & Furusawa, 1999; Mathivanan et al.,

2005). Studies on effect of Noni-ppt showed improvement in the survival time and curative effect in treatment of cancer. Noni-ppt administration significantly prolonged the survival duration of inbred Lewis lung tumor-bearing mice (Hirazumi et al., 1996). Further studies on Noni-ppt suggested that it suppresses the tumor growth through release of TNF- α , IL-1 β , IL-10, IL-12 p70, IFN- γ , and nitric oxide (Hirazumi & Furusawa, 1999). Some individual compounds from Noni juice were reported to function as ras inhibitors and thus suppressed the ras-expressing tumors (Wang et al., 1999).

Liu et al. (2001) found that the antitumor potential of noni fruit due to the presence of glycosides and asperulosidic acid extracted from the noni fruit on the AP-1 transactivation and cell transformation in mouse epidermal JB6 cells. They found that both the compound suppressed the cell transformation and AP-1 activity.

Wang & Su (2001) found that supplementation of noni juice to rats helped in prevention of DMBA-DNA adduct formation by 30% in heart, 41% in lung, 42% in liver, and 80% in kidney. Arpornsuwan & Punjanon (2006) tested the methanol extract from *M. citrifolia* fruits for cytotoxicity activity on the MTT assay. The appearance of cytotoxic changes after exposure to the extract was in a concentration dependent manner. The most sensitive cell line was baby hamster which showed median lethal concentration of 2.5 mg/ml followed by Vero cell (3.0 mg/ml) and human laryngeal carcinoma cell (5.0 mg/ml) respectively.

In another study, Thani et al. (2010) found the cytotoxic activity of noni leaves against the KB cell line while Lv et al. (2011) found the anticancer potential in the root extracts. Gupta & Patel (2013) also demonstrated that combination of Noni and cisplatin were able to induce the anticancerous activity through the p53 and Bax proteins (pro-apoptotic) up regulating pathways and Bcl-2 (anti-apoptotic gene), survivin and Bcl-XL proteins down regulating mechanisms. Saminathan et al. (2013b) showed the antitumor activity of noni fruit against mammary tumours in rats. The frequency of the tumor incidence was found to be significantly decreased in the noni fruit treated group and only benign tumour was observed while in the control group malignant tumour was observed.

11 Hypotensive activity

The first report of the hypotensive potential of the noni fruit was available in 1927, when Davison (1927) found that hot water extract of noni roots lowered the blood pressure of dog, the study was supported by the finding of Youngken et al., 1960; Youngken, 1958; Gilani et al., 2010. Later, Moorthy & Reddy (1970) and Yamaguchi et al. (2002) also reported the hypotensive effect of ethanol extract of noni dogs and rats respectively. Asahina et al., 1994 showed the diuretic effect of noni fruit juice.

12 Anti-inflammatory activity

The first scientific validation of the anti-inflammatory activity of noni juice was reported by Zhang et al. (1994). They studied the inhibitory activity of noni juice on COX-2 and COX-1 activities and compared with traditional anti-inflammatory drugs such as aspirin, indomethacin and celebrex. They found that activity of noni in COX-2 inhibition was on par with the celebrex. Later, Wang et al. (2002) also studied the anti-inflammatory activity of noni juice and found that feeding of 10% noni juice in drinking water for 12 days decreases the inflammatory foci in an acute liver injury model in female rats induced by the CCl₄. McKoy et al. (2002) showed that feeding of 200 mg of *M. citrifolia* prevented the formation of paw edema in the rats. This anti-inflammatory effect may be due to the inhibition of B2 receptor mediated mechanism of bradykinin

Study showed that damnacanthol isolated from the *M. citrifolia* roots mediates anti-inflammatory activity through H1 receptor (Okusada et al., 2011). Dussossoy et al. (2011) reported that the anti-inflammatory activity of the noni is due to the presence of several polyphenols, flavonoids, phenolics compounds, irridoids and ascorbic acid. These compounds probably act through the NO and PGE2 pathways by directly inhibiting the cyclooxygenase COX-1 and COX-2 activities.

Palu et al. (2006) also described the use of noni seed oil as anti-inflammatory activity in skin. Noni seed oil inhibited the COX-2 and 5-LOX enzymes in a concentration dependent manner and found that it is safe for topical use for skin care applications and is non-comedogenic.

13 Analgesic effect

The analgesic potential of *M. citrifolia* was reported by Younos et al. (1990). They evaluated the root extracts of *M. citrifolia* in mice and found that it showed a dose-related analgesic activity in the writhing and hot plate tests in mice. The extract did not show any toxic effect and further, on administration of higher dose, it decreased all the behavioural parameters and induces sleeping which is suggestive of sedative properties of the *M. citrifolia*. The analgesic and tranquilizing properties of the noni fruit was also reported by Joseph Betz (1997) and Wang et al. (2002). They also observed the dose dependent analgesic properties of the *M. citrifolia* in mice.

Punjanon & Nandhasri (2005) evaluated the different dose concentration of noni fruit extract viz. 1, 2, 3 and 4 g in mice and compared with the morphine sulphate. They found that 4 g/kg-1 concentration of noni fruit extract showed similar effect as produced by the morphine sulphate in inhibition of abdominal constriction induced by acetic acid.

This proves that *M. citrifolia* is having analgesic effect; however, detail studies are necessary for the identification of the chemical compounds and to study the mechanism of action. The analgesic efficacy of alcoholic extracts of Noni fruits was also demonstrated in the acetic acid induced writhing test (Okusada et al., 2011). In another study, Basar et al. (2010) demonstrated the analgesic activity of alcohol extract of noni fruit in reducing the pain and arthritis.

14 Antimicrobial Activity

Many reports are available on the antimicrobial activity of *M. citrifolia*. Atkinson, 1956 reported that the antibacterial effect of noni is due to the presence of acubin, L-asperuloside, alizarin and anthraquinone. Reports also suggested that these compounds are responsible for antibacterial activity against *Ps. Aeruginosa*, *Pr. Morgaii*, *B. subtilis*, *S. aureus*, *E. coli*, *Shigella* and *Salmonella* as well as treatment of skin infection, cold fever and other bacterial infection (Bushnell et al., 1950; Tabrah & Eveleth, 1966; Leach et al., 1988; Locher et al., 1995). Duncan et al. (1998) showed that scopoletin, a compound available in the noni is responsible for antibacterial activity against *E. coli* and control of serious illness and even death. Umezawa (1992) demonstrated that anti HIV activity in the noni is due to the presence of a compound i.e 1-methoxy-2-formyl-3-hydroxyanthraquinone which suppressed the cytopathic effect of HIV infected cells.

Broad spectrum antibacterial activity of various solvent extracts of *M. citrifolia* have been reported against Gram positive and Gram negative microorganisms (Wei et al., 2008; Jayaraman et al., 2008; Selvam et al., 2009; Kumar et al., 2010; Usha et al., 2010; Sunder et al., 2012; West et al., 2012).

The first report of the use of noni against tuberculosis was demonstrated by Saludes et al. (2002). They demonstrated that bactericidal activity of the noni leaf extracts was 89% compared to 97 % with rifampicin. The antifungal activity of *M. citrifolia* was demonstrated by Banerjee et al. (2006). They studied the antifungal activity in-vitro and found that *M. citrifolia* inhibited the growth of *C. albicans* in-vitro. They also found that the same extract also showed inhibitory activity against *Apergillus nidulans* spores. Sunder et al. (2012) have studied the wide spectrum antibacterial and antifungal activity of various parts of the *M. citrifolia* extracts. They have found that the methanol, ethanol, ethyl acetate, chloroform, acetone extracts of leaf, stem bark, fruit and seed showed broad spectrum antibacterial and antifungal activity in – vitro.

Conclusion and future perspectives

Farmers in several countries use medicinal plants in the maintenance and conservation of the healthcare of livestock. The last two decades have seen tremendous interest in the area of medicinal and aromatic plants. The role of plant derived drugs have been emphasized both national and international level. Based on the findings of the several researches on the

use of *Morinda citrifolia* for livestock health and production, it is concluded that the *M. citrifolia* may be used as alone or in combination with probiotics and other medicinal plants for growth, production, immunomodulator, antioxidant and many other properties in livestock and poultry. A lot of compounds with nutraceutical values have been identified from the *Morinda citrifolia*. Although lot of products are available in the market for human use, however, there are plenty of scope for development of noni based herbal products for livestock health and production. The antibacterial potential of the plant should be explored and studied in detail to develop the drug against multidrug resistance bacteria mainly for the tuberculosis, malaria, HIV and other diseases. The feeding of noni fruit has been found to exhibit very good antioxidant, anti-cholesterol and growth and immunomodulatory property in livestock and poultry. Study on the palatability of the fruit should be carried out to develop the cheap and best technology which is available at the farmer doorstep for the livestock and poultry. Reports suggested that chicken, duck and pig consumes the raw fruit of noni. However, the palatability of the fruit should be improved for feeding as such to the other livestock which could save the production losses and post harvest losses. The high mineral richness of the fruit and leaf should be studied in detail to study the efficacy of mineral supplement in poultry and large animals.

Since the ban of antibiotics as growth promoters by the European Union in 2006, lot of compounds, products etc have been studied as alternative to the antibiotics for probiotic, prebiotic, growth promotion effects in livestock and poultry. The presence of rich nutraceutical compounds in the noni might be useful for exploring this plant as an alternative to the antibiotic in poultry without giving any side effect. *Morinda citrifolia* trees are widely grown in coastal forest areas of A&N Islands. Owing to its high nutritive value for medicinal importance and having national and international market, increasing demand, there is a possibility for emerging as one of the most remunerative fruit crops to the island farmers. Recently, it has undergone a revival in Andaman and Nicolas Island as interest in plant with nutraceutical properties has increased. Noni plant is distributed in almost all parts of the Island. It can be found near the coast, in open lands and Grassland, in gulches and in distributed forest of the dryer areas.

It tolerates high soil salinity and brackish water stagnation. All the components of this plant have high demand in case of alternative medicines and herbal medicines. Due to its high demand and as a source of revenue generation the detail study on its effect on the livestock health and production should be carried out for its commercial exploitation.

Conflict of interest

Authors would hereby like to declare that there is no conflict of interests that could possibly arise.

References

- Abdulkarim A, Amad, Wendler KR, Zentek J (2013) Effects of a phytogetic feed additive on growth performance, selected blood criteria and jejunal morphology in broiler chickens. *Emirates Journal of Food and Agriculture* 25: 549–554.
- Abu-Ghefreh AA, Canatan H, Ezeamuzie CI (2009) In vitro and in vivo anti-inflammatory effects of andrographolide. *International Journal of Immunopharmacology* 9: 313–318.
- Adil S, Magray SN (2012) Impact and manipulation of gut microflora in poultry: a review. *Journal of Animal and Veterinary Advances* 11: 873–877.
- Al-Kassie GAM, Mohseen AM, Abd-Al-Jaleel RA (2011) Modification of productive performance and physiological aspects of broilers on the addition of a mixture of cumin and turmeric to the diet. *Research Opinions in Animal & Veterinary Sciences* 1: 31-34.
- Allen WH (1873) Some information on the ethnobotanical properties of Noni (*Morinda citrifolia*). In: *The useful plants of India*.
- Anantharaj A, Jeyakumar S, Sathya MM, Sunder J (2016) Biochemical and Antioxidant effects in crossbred calves fed with *Morinda citrifolia*. *Journal of Applied Animal Research*, DOI: 10.1080/09712119.2016.1151794 Online 21 march 2016.
- Aregheore EM (2005) Evaluation and utilization of Noni (*Morinda citrifolia*) juice extract waste in complete diets of goats. *Livestock Research for Rural Development* 17: 39.
- Arpornsuwan T, Punjanon T (2006) Tumor cell-selective antiproliferative effect of the extract from *Morindacitrifolia* fruits. *Phytotherapy Research* 20: 515–551.
- Asahina AY, Ebesu JSM, Ichinotsubo D, Tongson J, Hokama Y (1994) Effect of okadaic acid (OA) and Noni fruit extraction in the synthesis of tumor necrosis factor- α (TNF- α) by peripheral blood mononuclear (PBN) cells in vitro. *The Proceedings of the International Symposium of Ciguatera and Marine Natural Products* 197-205.
- Assi RA, Darwisa Y, Abdulbaqia IM, Khana AA, Vuanghaob, L, Lagharia MH (2015) *Morinda citrifolia* (Noni): A comprehensive review on its industrial uses, pharmacological activities, and clinical trials. *Arabian Journal of Chemistry*. DOI:10.1016/j.arabjc.2015.06.018.
- Atkinson N (1956) Antibacterial substances from flowering plants. 3. Antibacterial activity of dried Australian plants by rapid direct plate test. *Australian Journal of Experimental Biology* 34: 17–26.
- Aydin R, Karaman M, Cicek T, Yardibi H (2008) Black cumin (*Nigella sativa* L.) supplementation into the diet of the laying hen positively influences egg yield parameters, shell quality, and decreases egg cholesterol. *Poultry Science* 87:2590-2595.
- Azeke MA, Ekpo KE (2009) Egg yolk cholesterol lowering effects of garlic and tea. *Journal of Medicinal Plant Research* 3:1113–1117.
- Babatunde GM, Fajimi AO, Oyejide AO (1992) Rubber seed oil versus palm oil in broiler chicken diets. Effects on performance, nutrient digestibility, haematology and carcass characteristics. *Animal feed Science and Technology* 35 : 133–146.
- Banerjee S, Johnson AD, Csiszar K, Wansley DL, McGeady P (2006) An extract of *Morindacitrifolia* interferes with the serum-induced formation of filamentous structures in *Candida albicans* and inhibits germination of *Aspergillus nidulans*. *American Journal of Clinical Medicine* 34:503- 509.
- Barjesteh N, Hodgins DC, St Paul M, Quinteiro-Filho WM, DePass C, Monteiro MA, Shariff S (2013) Induction of chicken cytokine responses in vivo and in vitro by lipooligosaccharide of *Campylobacter jejuni* HS:10. *Veterinary Microbiology* 164: 122–130.
- Barroeta AC (2007) Nutritive value of poultry meat: relationship between vitamin E and PUFA. *World's Poultry Science Journal* 63: 277-284.
- Basar S, Uhlenhut K, Högger P, Schöne F, Westendorf J (2010) Analgesic and antiinflammatory activity of *Morinda citrifolia* L. (Noni) fruit. *Phytotherapy Research* 24:38-42.
- Batavani RA, Asri S, Naebzadeh H (2007) The effect of subclinical mastitis on milk composition in dairy cows. *Iranian Journal of Veterinary Research* 8: 205-211.
- Benjamin J, Richard D, Godbee G (2014) *Morinda citrifolia* based formulations for regulating T cell immunomodulation in neonatal stock animals. US Patent No US 8652546 B2
- Brisbin JT, Gong J, Parvizi P, Sharif S (2010) Effects of lactobacilli on cytokine expression by chicken spleen and caecal tonsil cells. *Clinical and Vaccine Immunology* 17: 1337–1343.
- Brooks V, Schafer J, Sharp M, Xu P, Cai J, Keuler J, Godbee NS, Peek RG, Schultz SF, Suresh RD, Darien M (2009) Effects of *Morindacitrifolia* (Noni) on CD4+ and CD8+ T-cell activation in neonatal calves. *Professional Animal Scientists* 20: 202-206.
- Brown AC (2012) Anticancer activity of *Morindacitrifolia* (Noni) fruit: a review. *Phytotherapy Research* 26 :1427-1440.
- Bruggnecate JT (1992). Native plants can heal your wounds. *Honolulu Star–Bulletin Local News*, Feb 2.

- Bui AKT, Basic A, Pettolino F (2006) Polysaccharide composition of the fruit juice of *Morinda citrifolia* (Noni). *Phytochemistry* 67:1271–1275.
- Bushnell OA, Fukuda M, Makinodian T (1950). The antibacterial properties of some plants found in Hawaii. *Pacific Science* 4: 167-183.
- Chan-Blanco Y, Fabrice V, Perez AM, Reynes M, Jean-Marc B, Brat P (2006) The noni fruit (*Morinda citrifolia* L.): a review of agricultural research, nutritional and therapeutic properties *Journal of Food Composition and Analysis* 19 : 645–654.
- Charis K (2000) A novel look at a classical approach of plant extracts. *Feed Mix (special issue on Nutraceuticals)*: 19-21.
- Chinsembu KC, Negumbo J, Likand OM, Mbangi A (2014) An ethnobotanical study of medicinal plants used to treat livestock diseases in Onayena and Katima Mulilo, Namibia. *South African Journal of Botany* 94:101-107.
- Chowdhury SR, Chowdhury SD, Smith TK (2002) Effects of dietary garlic on cholesterol metabolism in laying hens. *Poultry Science* 81:1856–1862.
- Chunhieng MT (2003) Developpement de nouveaux aliments santé tropical: application a
- Dalkilic B, Ciftci M, Guler T, Cerci IH, Ertas ON, Guvenc M (2009) Influence of dietary cinnamon oil supplementation on fatty acid composition of liver and abdominal fat in broiler chicken. *Journal of Applied Animal Research* 35: 173-176.
- Das L, Bhaumik E, Raychaudhuri U, Chakraborty R (2012) Role of nutraceuticals in human health. *Journal of Food Science and Technology* 49:173–183.
- Davison C (1927) Hawaiian medicine. *The Queen's Hospital Bulletin with Palama Clinic Section* 4: 2-5.
- Deng S, West BJ, Palu AK, Jensen CJ (2010) Determination and comparative analysis of major iridoids in different parts and cultivation sources of *Morinda citrifolia*. *Phytochemical Analysis* 22:26–30.
- Dhama K, Shyma K, Latheef, Mani S, Samad HA, Karthik K, Tiwari R, Khan RU, Alagawany M, Mayada R, Farag, Alam GM, Laudadio V, Tufarelli V (2015). Multiple Beneficial Applications and Modes of Action of Herbs in Poultry Health and Production-A Review. *International Journal of Pharmacology* 11: 152-176.
- Dilshad SMR, Rehman NU, Ahmad N, Iqbal A (2009) Documentation of ethnoveterinary practices for mastitis in dairy animals in Pakistan. *Pakistan Veterinary Journal* 30: 167-71.
- Dixon AR, McMillen H, Etkin NL (1999) Ferment This: The transformation of Noni, a traditional Polynesian medicine (*Morinda citrifolia*, Rubiaceae). *Economic Botany* 53:51-68.
- Duncan SH, Flint HJ, Stewart CS (1998) Inhibitory activity of gut bacteria against *Escherichia coli* 0157 mediated by dietary plant metabolites. *FEMS Microbiology Letter* 164: 283-258.
- Durrani FR, Ismail M, Sultan A, Suhail SM, Naila C, Durrani Z (2006) Effect of different levels of feed added turmeric (*Curcuma longa*) on the performance of broiler chicks. *Journal of Agricultural Biology Science* 1:9-11.
- Durrani FR, Sanaullah N, Chand Z, Durrani, Akhtar S (2008) Using aqueous extract of aloe gel as anticoccidial and immunostimulant agent in broiler production. *Sarhad Journal of Agriculture* 24 : 665-669.
- Dussosoy E, Brat P, Bony E, Boudard F, Poucheret P, Mertz C, Giaimis J, Michel A (2011) Characterization, anti-oxidative and anti-inflammatory effects of Costa Rican noni juice (*Morinda citrifolia*). *Journal of Ethnopharmacology* 133: 108–115.
- Earle JE (2001) *Plantas Medicinales en el Tro'pico Hu'edo*. Editorial Guayaca', San Jose'.
- Eevuri TR, Putturu R (2013) Use of certain herbal preparations in broiler feeds - A review. *Veterinary World* 6:172-179. DOI: 10.5455/vetworld.2013.172-179.
- Elagib HAA, Nabiela EM, Abbass SA, Ginawi TAN (2012) Effect of Natural Spices on Plasma Proteins in Broiler Chicks. *Journal of Nutrition and Food Sciences* 2:152. DOI:10.4172/2155-9600.1000152.
- Emadi M, Kermanshahi H (2007) Effect of turmeric rhizome powder on immunity responses of broiler chickens. *Medwell Journal of Animal and Veterinary Advances* 6: 833-836.
- Fallah R, Kiani A, Azarfar A (2013) A review of the role of five kinds of alternatives to in feed antibiotics in broiler production. *Journal of Veterinary Medicine and Animal Health* 5: 317–321.
- Farine JP, Lega, L, Moreteau B, Jean-Luc Le Quere (1996) Volatile components of ripe fruits of *Morinda citrifolia* and their effects on *Drosophila*. *Phytochemistry* 14:433-438.
- Fletcher HM, Dawkins J, Rattray C, Wharfe G, Reid M, Gordon-Strachan G (2013) *Morinda citrifolia* (Noni) as an Anti-Inflammatory Treatment in Women with Primary Dysmenorrhoea: A Randomised Double-Blind Placebo-Controlled Trial. *Obstetrics and Gynaecology International*. DOI: 10.1155/2013/195454.

- Fugh-Berman A (2003) The 5 minute herbal and dietary supplement consult. Philadelphia Lippincott pp 236-237.
- Fukao T, Koyasu S (2003) PI3K and negative regulation of TLR signaling. Trends in Immunology 24: 358–63.
- Furusawa E (2002) Anti-cancer activity of Noni fruit juice against tumors in mice. In: Nelson SC (Ed.), Proceedings of the 2002 Hawai'i Noni Conference, University of Hawaii at Manoa, College of Tropical Agriculture and Human Resources.
- Furusawa E, Hirazumi A (2003) Story S, Jensen J. Antitumor potential of polysaccharide-rich substance from the fruit juice of *Morinda citrifolia* (Noni) on sarcoma 180 ascites tumor in mice. Phytotherapy Research 17: 1158-1162.
- Furusawa E, Hirazumi A, Story SJ, Jensen J (2003) Antitumor potential of a polysaccharide-rich substance from the fruit juice of *Morindacitrifolia* (Noni) on sarcoma 180 ascites tumour in mice. Phytotherapy Research 17: 1158-1164.
- Gao H, Wang J (2016) Andrographolide inhibits multiple myeloma cells by inhibiting the TLR4/NF- κ B signaling pathway. Molecular Medicine Reports 13 :1827-1832.
- Galav P, Anitha J, Kateva SS (2013) Ethnoveterinary medicines used by tribals of Tadgarh Raoli wildlife sanctuary, Rajasthan. Indian Journal of Traditional Knowledge 12: 56-61.
- Ghazalah AA, Ali AM (2008) Rosemary leaves as a dietary supplement for growth in broiler chickens. International Journal of Poultry Science 7: 234-239.
- Gilani AH, Mandukhail SUR, Iqbal J, Yasinzai M, Aziz N, Khan A, Najeeb-ur-Rehman (2010) Antispasmodic and vasodilator activities of *Morinda citrifolia* root extract are mediated through blockade of voltage dependent calcium channels. BMC Complementary and Alternative Medicine 10.1186/1472-6882-10-2.
- Gupta RK, Patel AK (2013) Do the health claims made for *Morinda citrifolia* (Noni) harmonize with current scientific knowledge and evaluation of its biological effects. Asian Pacific Journal of Cancer Prevention 14: 4495-4499.
- Hashemi SR, Zulkifli I, Hair-Bejo M, Loh TC and Davoodi H (2009). Morphological Changes of the Intestinal in Broiler Chickens Fed Herbal Plant and Acidifier. Academia.edu
- Heinicke R (1985) The pharmacologically active ingredient of Noni. Bulletin of the National Tropical Botanical Garden.
- Heinicke R (2001) The Xerone system: a new cellular mechanism that explains the health.
- Hiramatsu T, Imoto M, Koyano T, Umezawa K (1993) Induction of normal phenotypes in ras-transformed cells by damnacanthol from *Morindacitrifolia*. Cancer Letter 30: 161–166.
- Hirazumi A, Furusawa E (1999) An immunomodulatory polysaccharide rich substance from the fruit juice of *Morinda citrifolia* (noni) with antitumour activity. Phytotherapy Research 13: 380-387.
- Hirazumi A, Furusawa E, Chou SC, Hokama Y (1996). Immunomodulation contributes to the anticancer activity of *Morindacitrifolia* (noni) fruit juice. Proceeding of the Western Pharmacology Society 39: 7-9.
- Hirazumi A, Furusawa E, Chou SC, Hokama Y (1994) Anticancer activity of *Morinda citrifolia* (noni) on intraperitoneally implanted Lewis lung carcinoma in syngeneic mice. Proceeding of the Western Pharmacology Society 37: 145-146.
- Hisawa T, Arase Y, Chen Z, Kita K, Umezawa K, Ito H, Suzuki N (1999) Stimulation of ultraviolet-induced apoptosis of human fibroblast UVr-1 cells by tyrosine kinase inhibitors. FEBS Letter 444: 173-176.
- Hosseinzadeh H, Qotbi AAA, Seidavi A, Norris D, Brown D (2014) Effects of Different Levels of Coriander (*Coriandrum sativum*) Seed Powder and Extract on Serum Biochemical Parameters, Microbiota, and Immunity in Broiler Chicks. The Scientific World Journal (2014), Article ID 628979, 11 pages <http://dx.doi.org/10.1155/2014/628979>.
- Ibrahim KA, Mahmoud A, Elhalim H (2005) Comparison of the efficacies of commercial probiotics on growth performance, carcass characteristics and some plasma constituents of broiler chicks. Suez Canal Veterinary Medicine Journal 8: 1-18.
- Issell BF, Gotay CC, Pagano I, Franke AA (2009) Using quality of life measures in a phase I clinical trial of noni in patients with advanced cancer to select a phase II dose. Journal of Dietary Supplements 6 : 347–359.
- Javed M, Durrani FR, Hafeez A, Khan RU, Ahmad I (2009). Effect of aqueous extract of plant mixture on carcass quality of broiler chicks. Journal of Agricultural and Biological Science 4: 37-40.
- Jayaraman SK, Manoharan MS, Illanchezian S (2008) Antibacterial, antifungal and tumor cell suppression potential of *Morinda citrifolia* fruit extracts. International Journal of Integrative Biology 3:44-49.
- Jin LZ, Ho YW, Abdullah N, Jalaludin S (1998) Growth performance, intestinal microbial populations and serum cholesterol of broilers fed diets containing *Lactobacillus* cultures. Poultry Science 77: 1259-1265.

- Jinhua Li, Lleng-Chee C, Marisa W, Wong DKW, Xianzhong Y, Yanzhang, W (2013) Antitumor activity of fermented Noni exudates and its fractions. *Molecular and Clinical Oncology* 1: 161–164.
- Joseph Betz (1997) Pride Publishing, Noni: Polynesia's natural pharmacy, Pp 13.
- Kabir SML (2009) The role of probiotics in the poultry industry. *International Journal of Molecular Sciences* 10: 3531–3546.
- Kalayou S, Haileselassie M, Gebre-egziabher G, Tiku'e T, Sahle S, Taddele H, Ghezu M (2012) In-vitro antimicrobial activity screening of some ethnoveterinary medicinal plants traditionally used against mastitis, wound and gastrointestinal tract complication in Tigray Region, Ethiopia. *Asian Pacific Journal of Tropical Biomedicine* 2 : 516–522.
- Kamel C (2001) Tracing modes of action and the roles of plant extracts in non-ruminants. In: Garnsworthy PC; Wiseman J (Eds.), in *Recent Advances in Animal Nutrition*, Nottingham University Press, Nottingham, UK, Pp. 135–150.
- Kanduri AB, Saxena MJ, Ravikanth K, Maini S and Kokane SS (2013) Effect of dietary replacement of antibiotic growth promoter with herbal growth promoter on performance of broiler poultry birds. *International Journal of Applied Pharmaceutical Science Bio and Medical Science* 2 : 12–15.
- Kant S, Ali N, Chandra G, Siddique RA , Imran M (2014) Effect of shatavari and vitamin E on hemato-biochemical profile of broilers during winter season. *Veterinary World* 7: 948-951.
- Kaşıkcı, G, Cetin O, Bingol EB, Gunduz MC (2012) Relations between electrical conductivity, somatic cell count, California mastitis test and some quality parameters in the diagnosis of subclinical mastitis in dairy cows. *Turkish Journal of Veterinary & Animal Sciences* 36: 49-55.
- Kitchen BJ (1981) Review of the progress of dairy science: bovine mastitis: milk compositional changes and related diagnostic tests. *Journal of Dairy Research* 48: 167-172.
- Krishnaiah D, Bono A, Sarbatly R, Anisuzzaman SM (2015) Antioxidant activity and total phenolic content of an isolated *Morinda citrifolia* L. methanolic extract from polyethersulphone (pes) membrane separator. *Journal of King Saud University – Engineering Sciences* 27: 63–67.
- Kumar KT, Panda DS, Nanda UN, Khuntia S (2010) Evaluation of antibacterial, antifungal and anthelmintic activity of *Morinda citrifolia* L. (Noni). *International Journal of Pharm Tech Research* 2: 1030-1032.
- Kumar M, Choudhary RS, Vaishnav JK (2005) Effect of supplemental prebiotic, probiotic and turmeric in diet on the performance of broiler chicks during summer. *Indian journal of Poultry Science* 40:137-141.
- Kumari P, Gupta MK, Ranjan R, Singh KK, Yadava R (1994) *Curcuma longa* as feed additive in broiler birds and its pathophysiological effects. *National Toxicology Program Technical Report Series* 435:1-288.
- Kusirisin W, Srichairatanakoo S, Lertrakarnnon P, Lailerd N, Suttajit M, Jaikang C, Chaiyasut C (2009) Antioxidative activity, polyphenolic content and anti-glycation effect of some Thai medicinal plants traditionally used in diabetic patients. *Medicinal Chemistry* 5:139–147.
- Landers TF, Cohen B, Wittum TE, Larson EL (2012) A Review of Antibiotic Use in Food Animals: Perspective, Policy, and Potential. *Public Health Reports* 127: 4–22.
- Langhout P (2000) New additives for broiler chickens. *World Poultry* 16: 22–27.
- Lanjewar RD, Zanzad AA, Ramteke BN, Deshmukh GB (2008) Effect of dietary supplementation of tulsi (*O. sanctum*) leaf powder on the growth performance and serum lipid profile in broilers. *Indian Journal of Animal Nutrition* 25:395-97.
- Lavanya G, Brahma Prakash GP (2011) Phytochemical screening and antimicrobial activity of compounds from selected medicinal and aromatic plants. *International Journal of Science and Nature* 2: 287-291.
- Lavinia S, Dumitrescu G, Drinceanu D, Stef D (2009) The effect of medicinal plants and plant extracted oils on broiler duodenum morphology and immunological profile of broiler. *Romanian Biotechnology Letter* 9: 1906-1914.
- Leach AJ, Leach DN, Leach GJ (1988) Antibacterial activity of some medicinal plants of Papua New Guinea. *Science New Guinea* 14 : 1-7.
- Lee SH, Lillehoj HS, Jang SI, Kim DK, Ionescu C, Bravo D (2010) Effect of dietary *Curcuma*, *capsicum* and *lentinus* on enhancing local immunity against *Eimeria acervulina* infection. *Journal of Poultry Science* 47: 89-95.
- Lee SY, Park SL, Hwang JT, Yi SH, Nam YD, Lim SI (2012) Antidiabetic Effect of *Morinda citrifolia* (Noni) Fermented by Cheonggukjang in KK-Ay Diabetic Mice. *Evidence Based Complementary and Alternative Medicine*. DOI: 10.1155/2012/163280.
- Liu G, Bode A, Ma WY, Sang S, Ho CT, Dong D (2001) Two novel glycosides from the fruits of *Morinda citrifolia* (noni) inhibit AP-1 transactivation and cell transformation in the mouse epidermal JB6 cell line. *Cancer Research* 61: 5749–5756.

- Locher CP, Burch MT, Mower HF, Berestecky H, Davis H, Van Polel B, Lasure A, Vander BDA, Vlieti-Nick AJ (1995) Anti-microbial activity and anti-complement activity of extracts obtained from selected Hawaiian medicinal plants. *Journal of Ethnopharmacology* 49:23-32.
- Lohani M (2010) Immunomodulatory Properties of Noni (*Morindacitrifolia*). Thesis paper presented to the Graduate School of Clemson University in partial fulfillment of the requirements for the Degree of Master of Science in Animal and Veterinary Science. http://etd.lib.clemson.edu/documents/1285619469/LOHANI_clemson_0050M_10649.pdf on 12 October 2013.
- Luseba D, Tshisikhawe MP (2013) Medicinal plants used in the treatment of livestock diseases in Vhembe region, Limpopo province, South Africa. *Journal of Medicinal Plants Research* 7: 593-601.
- Lv L, Chen H, Ho CT, Sang S (2011) Chemical components of the roots of Noni (*Morinda citrifolia*) and their cytotoxic effects. *Fitoterapia* 82: 704-708.
- Mahima, Rahal A, Deb R, Shyma K, Latheef, Samad HA, Tiwari R, Verma AK, Kumar A, Dhama K (2012) Immunomodulatory and Therapeutic Potentials of Herbal, Traditional/Indigenous and Ethnoveterinary Medicines. *Pakistan Journal of Biological Sciences* 15: 754-774.
- Mathivanan N, Surendiran G, Srinivasan K, Sagadevan E, Malarvizhi K (2005) Review on the current scenario of Noni research: Taxonomy, distribution, chemistry, medicinal and therapeutic values of *Morinda citrifolia*. *International Journal of Noni Research* 1: 1-16.
- Mckoy ML, Thomas EA, Simon OR (2002) Preliminary investigation of the anti-inflammatory properties of an aqueous extract from *Morinda citrifolia* (Noni). *Proceeding of the Western Pharmacology Society* 45: 76-78.
- Mir AQ, Bansal BK, Gupta DK (2014) Subclinical mastitis in machine milked dairy farms in Punjab: prevalence, distribution of bacteria and current antibiogram. *Veterinary World* 7: 291-294.
- Mirzaei F, Venkatesh KRH (2012) Efficacy of phyto medicines as supplement in feeding practices on ruminant's performance: a review. *Global Journal of Research in Medicinal Plants and indigenous Medicine* 1: 391-403.
- Mirzaei-Aghsaghali A (2012) Importance of medical herbs in animal feeding: A review. *Annals of Biological Research* 3 : 918-923.
- Mishra KP, Chand AS, Karan D, Ganju L, Sawhney RC (2008) Effect of Seabuckthorn (*Hippophae rhamnoides*) flavone on immune system: An in-vitro approach. *Phytotherapy Research* 22: 1490-1495.
- Moorthy NK, Reddy GS (1970) Preliminary phytochemical and pharmacological study of *Morindacitrifolia*, Linn. *Antiseptic* 67: 167-71.
- Morton JF (1992) The ocean-going Noni, or Indian mulberry (*Morindacitrifolia*, Rubiaceae) and some of its 'colorful' relatives. *Economic Botany* 46: 241-56.
- Mubarack HM, Doss A, Dhanabalan R, Venkataswamy R (2011) Activity of Some Selected Medicinal Plant Extracts Against Bovine Mastitis Pathogens. *Journal of Animal and Veterinary Advances* 10: 738-741.
- Muir WI, Bryden WL, Husband AJ (2000) Immunity, vaccination and the avian intestinal tract. *Development and Comparative Immunology* 24 : 325-342.
- Mufidah Z, Rifa'i, M, Rahayu S (2013) Immunomodulators activity of noni (*Morinda citrifolia* L.) fruit extract in mice infected with *Staphylococcus aureus*. *Jurnal Veteriner* 14: 501-510.
- Narimani-Rad M, Nobakht A, Shahryar HA, Kamani J, Lotfi A (2011) Influence of dietary supplemented medicinal plants mixture (ziziphora, oregano and peppermint) on performance and carcass characterization of broiler chickens. *Journal of Medicinal Plant Research* 5: 5626-5629.
- Nayak S, Chintamaneni M, Mengi S (2012) Anti-infective potential of *Morinda citrifolia* fruit extract. *International Journal of Pharmacy and Pharmaceutical Sciences* 3 : 81-83.
- Nayak S, Mengi S (2010) Immunostimulant activity of noni (*Morinda citrifolia*) on T and B lymphocytes. *Pharmaceutical Biology* 48 : 724-731.
- Nelson SC (2006) *Morindacitrifolia* (noni), ver. 4. In: Elevitch CR (Ed.), *Species Profiles for Pacific Island Agroforestry*. Permanent Agriculture Resources (PAR), Hôlualoa, Hawaii.
- Nitteranon V, Zhang G, Darien BJ, Parkin K (2011) Isolation and synergism of in vitro anti-inflammatory and quinone reductase (QR) inducing agents from the fruits of *Morinda citrifolia* (noni). *Food Research International* 44:2271-2277.
- Okusada K, Nakamoto K, Nishida M, Fujita-Hamabe W, Kamiya K, Mizushima Y, Shogo TS (2011) The antinociceptive and anti-inflammatory action of the CHCl₃-soluble phase and its main active component, damnacanthol, isolated from the root of *Morinda citrifolia*. *Biological and Pharmaceutical Bulletin* 34: 103-107. DOI: <http://doi.org/10.1248/bpb.34.103>.
- Osteras O, Solverod L (2009) Norwegian mastitis control programme. *Irish Veterinary Journal* 62 : 26-33.
- Palu AK, Brett JW, Jarakae J, Bing-Nan Z (2006) *Morindacitrifolia* L. Noni has cholesterol lowering

potential. Folk Botanical Wisdom: Towards Global Markets. June 5-9.

Park HJ, Lee C M, Jung ID, Lee J S, Jeong YI, Chang JH, Chun SH, Kim MJ, Choi W, Ahn SC, Shin YK, Yeom SR, Park YM (2009) Quercetin regulates Th1/Th2 balance in a murine model of asthma. *International Immunopharmacology* 9: 261-267.

Pawlus AD, Su B, Keller WJ, Kinghorn AD (2005) An anthraquinone with potent quinone reductase-inducing activity and other constituents of the fruits of *Morinda citrifolia* (noni). *Journal of Natural Products* 68:1720–1722.

Polat U, Yesilbag D, Eren M (2011) Serum Biochemical Profile of Broiler Chickens Fed Diets Containing Rosemary and Rosemary Volatile Oil. *Journal of Biological and Environmental Science* 5: 23-30.

Ponce CH, Ballou MA, Godbee RG, Hulbert LE, DiLorenzo N, Quinn MJ, Smith DR, Galyean ML (2011) Case study: Effects of *Morinda citrifolia* extract on performance and morbidity of newly received beef heifers. *Professional Animal Scientist* 27: 269–275. DOI: [http://dx.doi.org/10.15232/S1080-7446\(15\)30484-8](http://dx.doi.org/10.15232/S1080-7446(15)30484-8).

Punjanon T, Nand hasri P (2005) Analgesic effect of the alcoholic extract from the fruits of *Morinda citrifolia* 4: 678.

Raj RK (1975) Screening of indigenous plants for anthelmintic action against human *Ascaris Lumbricoides*: Part-II. *Indian Journal of Physiology and Pharmacology* 19: 47-49.

Rao KM (2001) MAP kinase activation in macrophages. *Journal of Leukocyte Biology* 69: 3–10.

Ross IA (2001) *Medical Plants of the World Chemical Constituents, Traditional and Modern Medical Uses*. Humana Press, New Jersey.

Saki AA, Aliarabi H, Siyar SAH, Salari J, Hashemi M (2014) Effect of a phyto-genic feed additive on performance, ovarian morphology, serum lipid parameters and egg sensory quality in laying hen. *Veterinary Research Forum* 5: 287-293.

Salarmoini M, Fooladi MH (2011) Efficacy of *Lactobacillus acidophilus* as probiotic to improve broiler chicks performance. *Journal of Agricultural Science and Technology* 13: 165-172.

Saludes JP, Garson MJ, Franzblau SG, Aguinaldo AM (2002) Antitubercular constituents from the hexane fraction of *Morinda citrifolia* Linn. (Rubiaceae). *Phytotherapy Research* 16: 683-685.

Saminathan M, Rai RB, Dhama K, Ranganath GJ, Palanivelu M, Rajesh G, Kumar SS (2013a) Evaluation of chemopreventive effect of *Morinda citrifolia* fruit juice in N-

Methyl-N-Nitrosourea (NMU) induced mammary tumours in rats. *Proceedings of the 27th Annual Conference on National Symposium and Indian Association of Veterinary Microbiologists, Immunologists*, December 13-15, 2013, Uttar Pradesh, India, pp: 20-21.

Saminathan M, Rai RB, Dhama K, Tiwari R, Chakraborty S, Amarpal, Ranganath GJ, Kannan K (2013b) Systematic Review on Anticancer Potential and other Health Beneficial Pharmacological Activities of Novel Medicinal Plant *Morinda citrifolia* (Noni). *International Journal of Pharmacology* 9: 462-492.

Saminathan M, Rai RB, Dhama K, Jangir BL, Suresh S, Ranganath GJ, Sophia I, Karuppanasamy K, Barathiraja S, Gopalakrishnan A (2014a). Effect of *Morinda citrifolia* (Noni) Fruit Juice on Antioxidant, Hematological and Biochemical Parameters in N-Methyl-N-Nitrosourea (NMU) Induced Mammary Carcinogenesis in Sprague-Dawley Rats. *International Journal of Pharmacology* 10 : 109-119.

Saminathan M, Rai RB, Murugesan V, Kumar SS, Baqir M, Lalruatfela PL (2014b) Induction of Cataract by N-methyl-N-nitrosourea (NMU) and Anti-cataract Activity of *Morinda citrifolia* Fruit Juice and Ethanolic Leaf Extract in Rats. *Indian Veterinary Journal* 91: 18-20.

Sasidhar PVK, Reddy YR, Sudhakar RA (2002) Economics of mastitis. *Indian Journal of Animal Sciences* 72: 439-440.

Schäfer M, Sharp P, Brooks VJ, Xu J, Cai J, Keuler NS, Peek SF, Godbee RG, Schultz RD, Darien BJ (2008) Enhanced bactericidal activity against *Escherichia coli* in calves fed *Morinda citrifolia* (Noni) puree. *Journal of Veterinary Internal Medicine* 22 :499-502.

Sharma K, Pachauri SD, Khandelwal K, Ahmad H, Arya A, Biala P, Agrawal S, Pandey RR, Srivastava A, Srivastav A, Saxena JK, Dwivedi AK (2016) Anticancer Effects of Extracts from the Fruit of *Morinda citrifolia* (Noni) in Breast Cancer Cell Lines. *Drug Research* 66 :141-147.

Sheeja K, Kuttan G (2007) Modulation of natural killer cell activity, antibody-dependent cellular cytotoxicity, and antibody-dependent complement-mediated cytotoxicity by andrographolide in normal and Ehrlich ascites carcinoma-bearing mice. *Integrative Cancer Therapies* 6: 66–73.

Shen T, Yang WS, Yi Y, Sung GH, Rhee MH, Poo H, Kim MY, Kim KW, Kim JH, Cho JY (2013) AP-1/IRF-3 Targeted Anti-Inflammatory Activity of Andrographolide Isolated from *Andrographis paniculata*. Evidence-Based Complementary and Alternative Medicine DOI: [org/10.1155/2013/210736](http://dx.doi.org/10.1155/2013/210736).

Selvam P, Raj K, Vimisha V, Harikrishnan R, Sarija KS, Umalekshmi R (2009) Antimicrobial Activity of Fruit Extracts of *Morinda citrifolia*. *Journal of Applied Chemical Research* 10: 61-63.

- Singh DR (2012) *Morinda citrifolia* L. (Noni): a review of the scientific validation for its nutritional and therapeutic properties. *Journal of Diabetes and Endocrinology* 6 : 77–91.
- Singh DR, Sunder J, Srivastava RC (2008) Peptide and mineral profile of *Morinda citrifolia* fruits and leaves. *International Journal of Noni Research* 2 : 72-78.
- Singh Y, Ikahehifo T, PanuveM, Slatter C (1984) Folk medicine in Tonga. A study on the use of herbal medicines for obstetric and gynecological conditions and disorders. *Journal of Ethnopharmacology* 12: 305-25.
- Solomon N (1999) The tropical fruit with 101 medicinal uses, NONI juice. 2nd ed. Woodland Publishing.
- St Paul M, Brisbin JT, Abdul-Careem MF, Sharif S (2013) Immunostimulatory properties of Toll-like receptor ligands in chickens. *Veterinary Immunology and Immunopathology* 152: 191–199.
- Sugiharto S (2015) Role of nutraceuticals in gut health and growth performance of poultry. *Journal of the Saudi Society of Agricultural Sciences*. doi:10.1016/j.jssas.2014.06.001.
- Sunder J, Kundu A, Singh DR, Jeyakumar S, Srivastava RC (2011a) Effect of feeding of *Morindacitrifolia* fruit juice on growth, production and immune response in Nicobari fowl. *Indian Journal of Animal Science* 81: 68–71.
- Sunder J, De AK, Jeyakumar S, Kundu A (2013a) Effect of feeding of *Morinda citrifolia* fruit juice on the biophysical parameters of healthy as well as mastitis-affected cow milk, *Journal of Applied Animal Research* 41: 29-33.
- Sunder J, Jeyakumar S, Sujatha T, Kundu A (2013b) Effect of feeding of Morical: a herbal based supplement on production and egg quality in Japanese quails. *Advances in Animal and Veterinary Sciences* 1: 157-160.
- Sunder J, Jeyakumar S, Sujatha T, Kundu A (2013c) Effect of feeding of morical (*M.citrifolia*) based herbal supplement on production and egg quality in Nicobari fowl. *Journal of Medicinal Plant Research* 7: 2999-3002.
- Sunder J, Kundu A, Singh DR, Sujatha T, Jeyakumar S (2013d) Production performance of Japanese quails on diets supplemented with dried fruit granules of *Morinda citrifolia*. *Animal Science Reporter* 7 :133-138.
- Sunder J, Jeyakumar S, Sujatha T, Kundu A (2014c) Grommune : *Morindacitrifolia* based herbal tonic for growth and immunity for commercial broilers. *Journal of Applied Animal Research* 43:2, 137-140.
- Sunder J, Rai RB, Yasmeen J, Kundu A, Jeyakumar S (2007) Immunomodulatory effect of *Morinda citrifolia* in poultry. *Indian Journal of Animal Sciences* 77 : 1126-1128.
- Sunder J, Singh DR, Jeyakumar S, Kundu A, Srivastava RC (2012) Antimicrobial activity of *Morinda citrifolia* solvent extracts. *Indian Veterinary Journal* 89: 9-11.
- Sunder J, Singh DR, Kundu A, Jeyakumar S (2011b) Effect of feeding of *Morinda citrifolia* extract on blood biochemical profile of Nicobari fowl and broiler. *Animal Science Reporter* 5 :141-146.
- Sunder J, Singh DR, Kundu A, Jeyakumar S, Verma SK (2011c) Supplementation of *Morindacitrifolia* extract on growth, production and immune response of Japanese quail. *Indian Journal of Veterinary Research* 20 (2): 61-65.
- Sunder J, Sujatha T, Kundu A (2015) Efficacy of feeding of *Morinda citrifolia* fruit juice and *Lactobacillus acidophilus* in broiler. *Asian Journal of Animal and Veterinary Advances* 10 : 352-359.
- Sunder J, Sujatha T, Pazhanivel N, Kundu A, Kundu MS (2014a) Effect of *M. citrifolia* fruit juice and *Lactobacillus acidophilus* on broiler duodenal morphology. *Advances in Animal and Veterinary Sciences* 2 : 28-30
- Sunder J, Sujatha T, Raja A, Kundu A, Kundu MS (2014b) Immunomodulatory effect of *Morindacitrifolia* and *Rorippa paniculata* on Expression of Toll-Like Receptors in Nicobari fowl. Abstract presented in the presented in XXVIII annual convention and international conference on Challenges and opportunities in animal health at the face of globalization and climate change held at DUVASU, Mathura from 30 October to 1 November 2014.
- Tabrah FL, Eveleth BM (1966) Evaluation of the effectiveness of ancient Hawaiian medicine. *Hawaii Medical Journal* 25: 223–230.
- Takashima J, Ikeda Y, Komiyama K, Hayashi M, Kishida A, Ohsaki A (2007) New constituents from the leaves of *Morinda citrifolia*. *Chemical and Pharmaceutical Bulletin* 55: 343–345.
- Tanabe M, Taniguchi M K, Takeuchi K, Takeda M, Ayata M, Ogura H, Matsumoto, M, Seya T (2003) Mechanism of up-regulation of human Toll-like receptor-3 secondary to infection of measles virus-attenuated strains. *Biochemical and Biophysical Research Communications* 311: 39-48.
- Thani W, Vallisuta O, Siripong P, Ruangwises N (2010) Anti-proliferative and antioxidative activities of Thai noni/*Yor (Morinda citrifolia* Linn.) leaf extract. *Southeast Asian Journal of Tropical Medicine and Public Health* 41: 482-489.
- Tipu MA, Akhtar MS, Anjum MI, Raja L (2006) New dimension of medicinal plants as animal feed. *Pakistan Veterinary Journal* 26 : 144-148.
- Tohyama M, Dai X, Sayama K, Yamasaki K, Shirakata Y, Hanakawa Y, Tokumaru, S, Yahata Y, Yang L, Nagai H,

- immunity of epidermal keratinocytes. *Biochemical and Biophysical Research Communications* 335: 505–511.
- Umezawa K (1992) Isolation of 1-methoxy-2-formyl-3-hydroxyanthraquinone from *M. citrifolia* and neoplasm inhibitors containing the same. *Japan Kokai Tokyo Koho JP 06 87, 736 (94-87, 736) Appl 92*: 264-311.
- Usha R, Sashidharan S, Palaniswamy M (2010) Antimicrobial Activity of a Rarely Known Species, *Morinda citrifolia* L. *Ethnobotanical Leaflets* 14:306-311.
- Verma RK (2014) An ethnobotanical study of plants used for the treatment of livestock diseases in Tikamgarh District of Bundelkhand, Central India. *Asian Pacific Journal of Tropical Biomedicine* 4 : 460–467.
- Virmani M, Garg SL, Virmani N, Batra SK (2010) Activity extracts of *Synyzium aromaticum* against microbes of veterinary importance. *Indian Journal of Animal Science* 80: 284-288.
- Wang M, Kikuzaki H, Csiszar K, Boyd CD, Maunakea A, Fong SF, Ghai G, Rosen RT, Nakatani N, Ho CT (1999) Novel trisaccharide fatty acid ester identified from the fruits of *Morinda citrifolia* (noni) *Journal of Agricultural and Food Chemistry* 47 : 4880–4882.
- Wang MY, Su C (2001) Cancer preventive effect of *Morinda citrifolia* (Noni),” *Annals of the New York Academy of Sciences* 952: 161–168.
- Wang MY, Su S, Nowicki AD, Jensen J, Anderson G (2001) *Morinda citrifolia* and cancer prevention. *Journal of Nutrition* 131 : 3151–3152.
- Wang MY, West BJ, Jensen CJ, Nowicki D, Su C, Palu AK, and erson G (2002) *Morindacitriifolia* (Noni): a literature review and recent advances in Noni research *Acta Pharmacologica Sinica* 23 : 1127–1141.
- Wang W, Wang J, Dong S, Liu C, Italiani P, Sun S, Xu J, Boraschi D, Ma S, Qu D (2010) Immunomodulatory activity of andrographolide on macrophage activation and specific antibody response. *Acta Pharmacologica Sinica* 31: 191–201
- Wang, GJ, Chu LY, Zhou MY, Chen WM (2006) Effects of preparation conditions on the microstructure of porous microcapsule membrane with straight open pores. *Journal of Membrane Science* 284: 301–312.
- Wei LS, Musa N, Sengm CT, Wee W, Mohd-Shazili NA (2008) Antimicrobial properties of tropical plants against 12 pathogenic bacteria isolated from aquatic organisms. *African Journal of Biotechnology* 7 : 2275-2278.
- activity of an iridoid rich extract from *Morinda citrifolia* fruit. *Current Research Journal of Biological Sciences* 4 : 52-54.
- Whistler WA (1985) Traditional and herbal medicine in the cook island s. *Journal of Ethnopharmacology* 13: 239-280.
- Whistler WA (1992) *Polynesian Plant Introductions*. Cox PA, Banack SA (Eds), Islands, Plants, and Polynesians. Portland , OR: Dioscorides Press 41-66.
- Woolhouse M, Ward M, Bunnik BV, Farrar J (2015) Antimicrobial resistance in humans, livestock and the wider environment. *Philosophical Transaction of the Royal Society B*. DOI: 10.1098/rstb.2014.0083.
- Wong DKW (2004) Are immune responses pivotal to cancer patient's long term survival? Two clinical case-study reports on the effects of *Morinda citrifolia* (Noni). *Hawaii Medical Journal* 63 : 182–184.
- Wu Y, Girmay S, Silva VMD, Perry B, Hu X, Tan GT (2015) The Role of Endophytic Fungi in the Anticancer Activity of *Morinda citrifolia* Linn. (Noni). *Evidence-Based Complementary and Alternative Medicine*. doi.org/10.1155/2015/393960.
- Yalcin S, Onbasilar EE, Reisli Z, Yalçın S (2006) Effect of garlic powder on the performance, egg traits and blood parameters of laying hens. *Journal of the Science of Food & Agriculture* 86:1336–1339.
- Yamaguchi S, Ohnishi J, Sogawa M, Maru I, Ohta Y, Tsukada Y (2002) Inhibition of angiotensin I converting enzyme by noni (*Morinda citrifolia*) juice. *Japanese Society for Food Science and Technology* 49: 624-627.
- Yancey JWS, Apple JK, Kegley EB, Goodbee RG (2013) Effects of *Morinda citrifolia* (Noni) pulp on growth performance and stress responses of growing cattle. *The Professional Animal Scientist* 29: 420-425.
- Yang Y, Iji PA, Choct M (2009) Dietary modulation of gut microflora in broiler chickens: a review of the role of six kinds of alternatives to in-feed antibiotics. *World Poultry Science Journal* 65: 97–114.
- Youngken HW (1958) A study of the root of *Morinda citrifolia* Linn, I. *Journal of American Pharmacology Association* 47: 162-5.
- Youngken HW, Jenkins HJ, Butler CL (1960) Studies on *Morinda citrifolia* L. *Journal of American Pharmacology Association* 49: 271-273.
- Younos C, Rolland A, Fleurentin J, Lanhers MC Miss LR, Mortier F (1990) Analgesic and behavioural effects of *Morindacitriifolia*. *Planta Medica* 56: 430-434.

Zamanzad-Ghavidel S, Adl KN, Sis NM, Aharizad S, Mirzaei-Aghsaghali A, Mohammadian M, Siadati SA (2011) Effects of Lactobacillus-based probiotic on growth performance, mortality rate and carcass yield in broiler chickens. *Annals of Biological Research* 2: 325-331.

Zeedan GSG, Abdalhamed AM, Abdeen E, Ottai ME, Abdel-Shafy S (2014) Evaluation of antibacterial effect of some Sinai

medicinal plant extracts on bacteria isolated from bovine mastitis. *Veterinary World* 7: 991-998.

Zhang LD, ZhangYL, Xu SH, Zhou G, Jin SB (1994) Traditional Chinesemedicine typing of affective disorders and treatment. *American Journal of Clinical Medicine* 22: 321-327.