

27th Year

MOST CIRCULATED & APPRECIATED MAGAZINE IN INDIA



Dr. RAJEEV RANJAN, I.A.S
Secretary, Department of Fisheries
Article on Ornamental fisheries sector

visit our website
www.aquastarmagazine.com

Aqua Star

Estd : 1994

RNI Regd. No. 67662/97 Regd. Magazine by the Govt. of India

DECEMBER-2020

A MONTHLY MAGAZINE IN THE SERVICE OF FISHERIES & AQUACULTURE



Trout Nutrition, opens state-of-the-art facility in Telangana



Registered with Coastal Aquaculture Authority (CAA) as Antibiotic-free inputs.

Quality Inputs - Quality Management... Towards Sustainable Aquaculture!!



SRIBS BIOTECHNIQS PVT. LTD.

302, Wing-A, Cello Triumph, I.B.Patel Road, Off Western Express Highway, Goregaon East, Mumbai 400063, Maharashtra, India. Phone: +9122 2686 1441/ 2685 1442
Email: info@sribsbio.in • Web: www.sribsbio.in

SRIBS sustainability simplified®

INDEX

★ త్వరలో అమలులోకి రానున్న ఫిష్ ఫీడ్ యాక్ట్	15
★ ఆక్వాసాగులో యాంటిబయోటిక్స్ వాడకూడదు	19
★ రాయ్లసాగులో అత్యాధునిక టెస్ట్లు తప్పనిసరి డాలర్ల పంటకు పరిక్షలే కీలకం	20
★ మత్స్యకారులకు ఇక మంచిదే	23
★ నాణ్యమైన ఆక్వా ఉత్పత్తులే లక్ష్యం..	24

సాంకేతిక సమాచారం

★ మంచినీటి చేపల మరియు వేనామి సాగులో మేత యాజమాన్య పద్ధతులు	27
★ మంచినీటి చేపల బ్రీడింగ్, సర్ప్లీ యాజమాన్యంలో ప్రామాణిక పద్ధతులు	31
★ ప్రేరేపిత ప్రజననము (ఇండ్యూస్ట్ బ్రీడింగ్) - ప్రామాణిక పద్ధతులు	36
★ ఘనంగా మత్స్యకార దినోత్సవం	50
★ కారమేను చేపపిల్లల ఉత్పత్తిలో ప్రామాణిక నిర్వహణా పద్ధతులు	53



Focus Article

★ Ornamental fisheries sector in India towards a new dawn under Pradhan Mantri Matsya Sampada Yojana (PMMSY) By Dr. Rajeev Ranjan IAS	58
--	----

Article

1) Impact of Harmful Algal Bloom (HAB) in Aquatic system	71
2) Paper based packaging material coated with microbial polyhydroxy alkanoates: An approach towards 100% biodegradability	75
3) Climate Change Impacts on Freshwater Resources, Biodiversity and Fisheries in India: A Snapshot	81

National News

★ Inland fish culture has huge potential to grow under fisheries industries CII FISHMART89	89
★ WORLD FISHERIES DAY 2020	90
★ CMFRI Extends Support & Opens Livefish Counter	93
★ ICAR-CIBA Conducted a National webinar on the Most sought after Biofloc Farming....	94
★ Trow Nutrition, foray into India with a state-of-the-art facility in Telangana!	98
★ National Award Winner WORLD FISHERIES DAY - 2020	102
★ Vaisakhi Bio-Marine Bags Best Shrimp Hatchery Award	109

International News

★ World Aquaculture 2020	110
★ RDO Blue In-Situ Offers Optical DO Monitoring Probe Customized for Aquaculture	110
★ World Sustainability Organization Collaborates with The Good Food...	113
★ ICC Brazil among the four Best Companies in the Animal Nutrition sector in Brazil	114

Paper based packaging material coated with microbial polyhydroxyalkanoates: An approach towards 100% biodegradability

Sreelakshmi K.R., Murugadas. V, Remya S., Rehana Raj and Mohan C.O, ICAR-Central Institute of Fisheries Technology, Matsyapuri P.O., Cochin. *Corresponding author: ^aScientist, Fish processing division, ICAR-Central Institute of Fisheries Technology, Matsyapuri P.O., Cochin. Email: sreelakshmi.k.ramanathan@gmail.com.

Polyhydroxyalkanoates (PHAs) are the polyesters produced by microorganisms under limiting conditions of nitrogen, phosphorus, oxygen or magnesium in the presence of excess of carbon. Polyhydroxybutyrate (PHB) was the first discovered form of PHA in 1926 and is from *Bacillus Megaterium*. Today, more than 300 microorganisms are known to synthesis and accumulate PHA intracellularly. Also genetically modified microbes and plants are being reported for the production of PHA. PHA can be of short chain length (3±5 carbon atoms) or medium chain length (6±14 carbon atoms).

Polyhydroxyalkanoates are found to be fully biodegradable. The biodegradability can be ensured based on marine biodegradability, industrially compostability, home compostability, soil biodegradability and anaerobic digestibility. Among the plastics, only thermoplastic starch based and PHA based compounds are fully degradable. Also compared to thermoplastic starch the production of PHA can be done using substrates which cannot initiate a competition for human or animal food. The biodegradability and biocompatibility are the major characteristics of PHA which makes it stands in prime position among other plastics. PHA acts as a potential candidate to replace the petroleum based plastics but the cost of production and lower yield matters. Now a days researchers are focussed to increase the yield of PHA production using waste substrates for microbial culture and also on the utilisation of different wastes



Sreelakshmi K.R.

like municipal waste, sewage, agricultural waste etc. so as to reduce the cost of production. Fish oil was also reported to be useful for the production of PHA (Table. 1).

Paper, as a packaging material is safe for the environment because of its fully biodegradable property. The global production of paper is mainly constituted by use of wood (92%) and only 8% is produced using agricultural wastes (Fahmy et al. 2017). Paper is composed of long chain molecules of cellulose, having a disrupted crystalline structure with amorphous regions. The water vapour barrier properties of paper is low because of this porous nature and also the OH groups contributes to hydrophilic nature. Hence when used as packaging for food, it easily absorbs water and loses the mechanical strength.

Inorder to improve the barrier properties, other materials such as plastic, aluminium etc. are used. This inturn adds to the mechanical properties also. Coatings with ethyl vinyl alcohol, poly olefins, rubber latex, fluorocarbon etc. are given to increase the gas barrier properties. But the biodegradability of paper will be lost because of these synthetic treatments. The use of natural polymers can be an alternative.

Paper coatings with PHA

The properties of PHA depends on the constituted monomers. This in turn is affected by the substrate used, conditions of culture and the microorganism. The hydrophobic character of PHA

Table. 1. Reports on utilisation of fish oil for production of PHA

Substrate	Microorganism	Yield (% of dry cell waste)	Reference
Waste fish oil, glycerol and fish sauce	<i>Salinivibrio species</i> M318	51.5	Van Thuoc et al, 2019
Alaska pollack oil	<i>P. oleovorans</i> NRRL B-778	27	Ashby and Solaiman, 2008
	<i>P. oleovorans</i> NRRL B-14682	6	
	<i>P. oleovorans</i> NRRL B-14683	52	
	<i>P. resinovorans</i> NRRL B-2649	53	
	<i>P. corrugata</i> 388	43	
	<i>P. putida</i> KT2442	39	
Waste fish oil (15g/L), Gum Arabic (2.5g/L), Cell growth medium	<i>Cupravidus necator</i>	75	Kaesavan, 2014

makes it resistant to hydrolytic degradation. They have good barrier property against oxygen and water vapour, good UV resistance and poor resistance towards acids and bases.

Different techniques can be applied for coating PHA over paper. PHB dissolved in chloroform can be used to make a coating using solvent casting method. The barrier properties against moisture, surface roughness and tensile strength increase with increase in concentration of PHB used. Next method is compression moulding which gave similar properties as in solvent casting. The additional step of acetylation of paper board increased the adhesion of PHB over paper and the characteristics were improved. The PHB coating was found to be a good substitute for tetra pak packaging. The third method is extrusion coating. The adhesion of PHB/V extrusion coating on paper is to be improved by pretreatment of the paper with acrylic based primer (Rastogi and Samyn, 2015).

The PHB was found to be suitable for paper sizing than PHBV and a temperature of 160°C gave improved film formation by PHB melting. The PHB can give better water barrier properties but lower oil resistance than PLA coatings. Because of increased microbial activity, the paper coated with PHA was found to be more biodegradable than the biopolymer alone (Masood, 2017).

Disadvantage of PHA as a coating for paper

The major drawback of PHA is its thermal instability which causes difficulties in processing.

The melting temperature of PHA varies depending on the monomers and copolymers. Usually the melting point starts from 120°C and can go beyond 300°C.



SHARAT INDUSTRIES LIMITED

No. 17, (Old No.30) LUZ Avenue, Myloapore, Chennai-600004. India

Phone : 044-24994663, 24997867/88, Fax : 044-24995893.

e-mail : sharatpremium@dataone.in

: NELLORE OFFICE :

16/375-A-2, Ramalingapuram, Nellore-524003 (A.P.) Phone : 0861-2344857/
2344859, 2331727/2329831, Fax : 0861-2325583. email : saratnlr@hotmail.com

Farm, Hatchery & Feed Mill : Venkannapalem Village,

Thotapalli, Gudur Mandal, Nellore - 524 002, ☎ : 0861-2387633, 2387733

Processing Plant : Mahalakshmpuram Village,

T.P.Gudur Mandalam, Nellore Dist. ☎ : 0861-24994663 / 24997867

Corp. Off: Flat No.4, III Floor, Pallavi Apartments, No.57/11 HDFC Bank Compound,
First Main Road, RA Puram, Chennai - 600 028. ☎ : 044-24994663 / 24997867