



## Prevalence of *Rhipicephalus sanguineus* on dogs in Chennai, Tamil Nadu

C.Soundararajan, K.Nagarajan<sup>2</sup>, S.Selvakumar<sup>2</sup>, K.C.Rajesh Kanna<sup>3</sup>, M.Arul Prakash<sup>4</sup> and R.Anandaraja<sup>5</sup>

Department of Veterinary Parasitology, Madras Veterinary College,

Tamil Nadu Veterinary and Animal Sciences University (TANUVAS), Chennai – 600 007, TN.

---

### Abstract

Dog ticks are the most widespread ticks in the world and are well-recognized vectors of many pathogens affecting dogs and occasionally humans. A total of 352 dogs of different breeds were examined for the presence of ticks from June 2009 to May 2011 in Chennai, Tamil Nadu. The collected ticks were identified as *Rhipicephalus sanguineus*. The overall prevalence of ticks was 58.52%. Among the seasons, highest prevalence of *R. sanguineus* was observed in northeast monsoon (34.46%) followed by southwest monsoon (30.10%), summer (23.79%) and lowest level in winter (11.65%). Adult dogs were heavily infested with *R. sanguineus* ticks than the puppies (67.96% vs 32.04%). Male dogs were highly infested (74.76%) than the female dogs (25.24%). Among the breeds, Spitz was highly infested (28.16%) followed by mongrels (24.76%), Labrador (17.48%), German shepherd (12.44%) and Doberman (4.37%). Other breeds were infested with very low levels ranging from 0.49% to 2.43%. Hairy breeds (Spitz, Labrador, German shepherd, Lhasa Apso, Border collie, Golden Retriever and Pomeranian) were heavily infested with ticks (61.17%) than the non-hairy breeds (38.83%).

**Key words:** Tick, *Rhipicephalus sanguineus*, Prevalence, Dog, Chennai, Tamil Nadu

---

### Introduction

*Rhipicephalus sanguineus* is an endophilic, monotropic, three-host tick species and the most widespread tick in the world. It is a well-recognized vector of many pathogens affecting dogs and occasionally humans (Pegram et al., 1987; Dantas-Torres, 2010). It acts as a vector for transmission of *Ehrlichia canis*, *Babesia vogeli* and *Hepatozoon canis* in canines and *Rickettsia conorii* and *R. rickettsia* in human (Otranto et al., 2012). Dogs are the main hosts of both adult and pre-adult parasitic stages of the tick. Adult *R. sanguineus* can also feed on many other mammals, such as goats

(Valls, 2008), sheep (Okoli et al., 2006; Bui et al., 2012), cattle, buffaloes (Chhillar et al., 2014), horses, ungulates, wild carnivores, cats (Filippova, 1997; Ogden et al., 2000), hares and hedgehogs. Pre-adult ticks were found on shrews and many rodents (Walker et al., 2000). It also occurs in human beings (Mentz et al., 2016). This paper reports the prevalence of *R. sanguineus* on different breeds of dogs in Chennai, Tamil Nadu.

### Materials and Methods

A total of 352 tick infested dogs (German Shepherd, Doberman, Dalmatian, Dachshund, Great Dane, Golden Retriever, Labrador, Lhasa Apso, Pug, Boxer, Border collie, Spitz, Pomeranian, Rottweiler and Mongrel) were examined at small animal private clinics, Perambur, Chennai, Tamil Nadu from June 2009 to May 2011 to know the seasonal prevalence of ticks in Chennai, Tamil Nadu. The ticks were collected and examined after clearing them with 10% sodium hydroxide (Sloss, 1970) and identified based on morphological features by

---

Corresponding author. Email: drsoundarpara@gmail.com

<sup>1</sup>Department of Veterinary Pathology,

<sup>2</sup>Department of Clinical Medicine and Jurisprudence, Madras Veterinary College, TANUVAS, Chennai – 600007, TN.

<sup>3</sup>Veterinary Dispensary, Arakandanallur, Villupuram district, TN.

<sup>4</sup>University Research Farm, TANUVAS, Madhavaram, Chennai - 600 051, Tamil Nadu, India.

<sup>5</sup>ICAR-Central Institute of Brackishwater Aquaculture, Chennai - 600028, Tamil Nadu, India.



microscopical examination (Soulsby, 1982). Statistical analysis for the data was carried out using SPSS tools, Version 2017.

**Results and Discussion**

The overall prevalence of *R.sanguineus* was 58.52% (Table 1). Nithikathkul *et al.*(2005)and Otranto *et al.* (2005) reported 80% and 39.4% prevalence of *R. sanguineus* on dogs in north-eastern Thailand and Italy, respectively. Contrary to this, Goldberg *et al.* (2002) and Dantas-Torres *et al.* (2009) reported lowest prevalence of 3.8% and 5.4% *R. sanguineus* infestation on dogs of North West Georgia (USA) and North eastern Brazil respectively.

Among the breeds, Spitz was highly infested (28.16%) with *R. sanguineus* followed by

mongrels (24.76%), Labrador (17.48%), German shepherd(12.44%) and Doberman (4.37 %).Dantas-Torres *et al.* (2004) observed the higher prevalence of ticks in free ranging dogs than the house-trained dogs. Soares *et al.* (2006) reported that the tick prevalence was more in house living dogs with grassy yard when compared to the dogs kept in apartments.Okoli *et al.* (2006) reported that *R. sanguineus* infestation was more in Alsatian than mongrel.Louly *et al.* (2009) reported English Cocker Spaniel was more susceptible than other breeds. In our study, hairy breeds (Spitz, Labrador, German shepherd, Lhasa Apso, Border collie, Golden Retriever and Pomeranian) were found to be heavily infested with ticks (61.17 %) than the non-hairy breeds (38.83%). Tick infestation was significantly ( $P<0.01$ ) higher in hairy breed than the non-hairy breed (Table 1)

**Table 1 Breed wise prevalence of *R.sanguineus* on dog**

Sl. No.	Breed	Skin	Season				Age		Sex		Overall %
			S	SWM	NEM	W	Puppy	Adult	M	F	
1	Boxer	Non-hairy	33.33 (1)	0.00(0)	67.67 (2)	0.00 (0)	0.00 (0)	100 (3)	66.67 (2)	33.33 (1)	1.46 (3)
2	Dalmatian	Non-hairy	0.00 (0)	100 (1)	0.00 (0)	0.00 (0)	0.00 (0)	100 (1)	100 (1)	0.00 (0)	0.49 (1)
3	Dachshund	Non-hairy	33.33 (1)	0.00 (0)	66.67 (2)	0.00 (0)	0.00 (0)	100 (3)	100 (3)	0.00 (0)	1.46 (3)
4	Doberman	Non-hairy	55.56 (5)	0.00 (0)	22.22 (2)	22.22 (2)	44.44 (4)	55.56 (5)	88.89 (8)	11.11 (1)	4.37 (9)
5	Great Dane	Non-hairy	0.00 (0)	33.33 (1)	66.67 (2)	0.00 (0)	66.67 (2)	33.33 (1)	66.67 (2)	33.33 (1)	1.46 (3)
6	Rottweiler	Non-hairy	0.00 (0)	40.00 (2)	40.00(2)	20.00 (1)	20.00 (1)	80.00 (4)	80.00 (4)	20.00 (1)	2.43 (5)
7	Pug	Non-hairy	0.00 (0)	0.00 (0)	80.00 (4)	20.00 (1)	40.00 (2)	60.00 (3)	80.00 (4)	10.00 (1)	2.43 (5)
8	Mongrel	Non-hairy	27.45 (14)	25.50 (13)	7.25 (19)	9.80 (5)	33.33 (17)	66.67 (34)	82.35 (42)	17.65 (9)	24.76 (51)
9	Border collie	Hairy	0.00 (0)	0.00 (0)	100 (1)	0.00 (0)	0.00 (0)	100 (1)	100 (1)	0.00 (0)	0.49 (1)
10	Golden Retriever	Hairy	0.00 (0)	0.00 (0)	100 (1)	0.00 (0)	100 (1)	0.00 (0)	100 (1)	0.00 (0)	0.49 (1)
11	German Shepard	Hairy	4.00 (1)	60.00 (15)	28.00 (7)	8.00 (2)	40.00 (10)	60.00 (15)	64.00 (16)	36.00 (9)	12.14 (25)
12	Lhasa Apso	Hairy	0.00 (0)	75.00 (3)	25.00 (1)	0.00 (0)	25.00 (1)	75.00 (3)	75.00 (3)	25.00 (1)	1.94 (4)
13	Labrador	Hairy	30.56 (11)	33.33 (12)	25.00 (9)	11.11 (4)	30.55 (11)	66.44 (25)	80.56 (29)	19.44 (7)	17.48 (36)
14	Pomeranian	Hairy	0.00 (0)	0.00 (0)	100 (1)	0.00 (0)	0.00 (0)	100 (1)	100 (1)	0.00 (0)	0.49 (1)


15	Spitz	Hairy	27.59 (16)	25.86 (15)	31.03 (18)	15.52 (9)	29.31 (17)	70.69 (41)	63.79 (37)	36.21 (21)	28.16 (58)
Total			23.79 (49)	30.10 (62)	34.46 (71)	11.65 (24)	32.04 (66)	67.96 (140)	77.76 (154)	25.24 (52)	58.52 (206)

S-Summer; SWM-Southwest monsoon, NEM-Northeast monsoon; W-Winter

Dog breed	Seasons			Age			Sex	
	Summer**	South-west monsoon**	North-east monsoon**	Winter*	Puppy**	Adult**	Male**	Female**
Non-hairy	10.5±0.5 <sup>a</sup>	8.5±0.5 <sup>a</sup>	16.5±0.5 <sup>a</sup>	4.5±0.5 <sup>a</sup>	13.0±1.0 <sup>a</sup>	27.0±1.0 <sup>a</sup>	33.0±2.0 <sup>a</sup>	7.0±1.0 <sup>a</sup>
Hairy	14.0±0.0 <sup>b</sup>	22.5±0.5 <sup>b</sup>	19.0±0.0 <sup>b</sup>	7.5±0.5 <sup>b</sup>	20.0±1.0 <sup>b</sup>	43.0±1.0 <sup>b</sup>	44.0±1.0 <sup>b</sup>	19.0±1.0 <sup>b</sup>

\*P<0.05; \*\*P<0.01 – Values bearing different superscript in a column differ significantly.

Among the seasons, highest prevalence of *R.sanguineus* was observed in north-east monsoon (34.46%) followed by southwest monsoon (30.10%), summer (23.79%) and lowest in winter (11.65%). Highest prevalence of ticks was observed during monsoon months (64.56%) than the summer and winter seasons (Figure. 1). On the contrary, Okoli et al. (2006) has reported that *R. sanguineus* occurred during the early rainy season in the month of April and May. Chhillar et al. (2014) observed the occurrence of *R.sanguineus* throughout the year and reduced in winter in case of cattle and buffaloes.



**Figure 1** Season wise prevalence of *R.sanguineus* on dogs

Among the age groups, adult dogs were heavily infested with *R.sanguineus* ticks than the puppies (67.96 % vs 32.04%). In contrast, Tinoco-Gracia et al. (2009) reported higher prevalence in younger dogs. Among the sex, male dogs were highly infested (74.76%) than the female dogs (25.24 %) which is in similarity with report



**Figure 2.** Heavy tick infestation on a dog



**Figure 3.** Heavy tick infestation on ear of dog

of Silveira et al. (2009). Sites of attachment were mostly on the ears, withers, face and perineum, especially under the heavy fur of the mane and in the inter-digital spaces (Figure. 2 and 3) which is similar to that of Okoli et al. (2006).



## References

- Biu, A.A., Abdul Kadir, M.A. and Isadu, T.H., 2012. Effects of temperature and relative humidity on the egg laying pattern of *Rhipicephalus sanguineus* (Koch, 1844) infesting sheep in semi-arid region of Nigeria. *Sokoto J. Vet. Sci.*, 10: 18-20.
- Chhillar, S., Chhilar, J.S. and Kaur, H., 2014. Investigations on some hard ticks (acari:ixodidae) infesting domestic buffalo and cattle from Haryana, India. *J. Entomol. Zool. Stud.*, 2: 99-104.
- Dantas-Torres, F., Figueredo, L.A. and Faustino, M.A.G., 2004. Ectoparasitos de caesprovenientes de alguns municípios da Região Metropolitana do Recife, Pernambuco, Brasil. *Rev Bras Parasitol. Vet.*, 13:151-154.
- Dantas-Torres, F., Melo, M.F., Figueredo, L.A. and Brandao-Filho, S.P., 2009. Ectoparasite infestation on rural dogs in the municipality of Sao Vicente Ferrer, Pernambuco, Northeastern Brazil. *Rev. Bras. Parasitol. Vet.*, 18:75-77.
- Dantas-Torres, F., 2010. 1. Biology and ecology of the brown dog tick, *Rhipicephalus sanguineus* -Review. *Parasit. Vectors*, 3:26.
- Filippova, N.A., 1997: Ixodid Ticks of the Subfamily Amblyomminae. *Arachnoidea 4 (5)*. Fauna of Russia and neighbouring countries. Nauka Publishing House, St. Petersburg, Russia, 436 Pp.
- Goldberg, M., Recha, Y. and Durden, L.A., 2002. Ticks parasitizing dogs in northwestern Georgia. *J. Med. Entomol.*, 39:112-114.
- Louly, C.C.B., Soares, S., Silveira, D., Neto, O., Silva, A. and Borges, L., 2009. Differences in the susceptibility of two breeds of dogs, English cocker spaniel and beagle, to *Rhipicephalus sanguineus* (Acari: Ixodidae). *Int. J. Acarol.*, 35:25-32.
- Mentz, M.B., Trombka, M., Silva, G.I. and Silva, C.E., 2016. *Rhipicephalus sanguineus* (Acari: Ixodidae) biting a human being in Porto Alegre city, Rio Grande Do Sul, Brazil. *Rev. Inst. Med. Trop. Sao Paulo.*, 58:35
- Nithikathkul, C., Polseela, R., Iamsa-ard, J., Wongsawad, C. and Jittapalpong, S., 2005. A study of ectoparasites of *Canis lupus familiaris* in Mueang district, KhonKaen, Thailand. *Southeast Asian J. Trop. Med. Public Health.*, 36:149-151.
- Ogden, N.H., Cripps, P., Davidson, C.C., Owen, G., Parry, J.M., Timms, B.J. and Forbes A.B., 2000. The Ixodid tick species attaching to domestic dogs and cats in Great Britain and Ireland. *Med. Vet. Entomol.*, 14: 332-338.
- Okoli, I.C., Okoli, C.G. and Opara, M., 2006. Environmental and multi-host infestation of the brown dog tick, *Rhipicephalus sanguineus* in Owerri, south-east Nigeria - A Case Report. *Vet. Arhiv.*, 76:93-100.
- Otranto, D., Dantas-Torres, F., Tarallo, V.D., Ramos, R.A., Stanneck, D., Baneth, G. and de Caprariis, D., 2012. Apparent tick paralysis by *Rhipicephalus sanguineus* (Acari: Ixodidae) in dogs. *Vet. Parasitol.*, 188:325-329.
- Otranto, D., Lia, R.P., Cantacessi, C., Galli, G., Paradies, P., Mallia, E. and Capelli, G., 2005. Efficacy of a combination of imidacloprid 10% permethrin 50% versus fipronil 10% (S)-methoprene 12%, against ticks in naturally infected dogs. *Vet. Parasitol.*, 130:293-304.
- Pegram, R.G., Keirans, J.E., Clifford, C.M., and Walker, J.B., 1987. Clarification of the *Rhipicephalus sanguineus* group (Acari, Ixodoidea, Ixodidae). Ii. *R. sanguineus* (Latreille, 1806) and related species. *Syst. Parasitol.* 10, 27-44.
- Silveira, J.A., Passos, L.M. and Ribeiro, M.F., 2009. Population dynamics of *Rhipicephalus sanguineus* (Latreille, 1806) in Belo Horizonte, Minas Gerais state, Brazil. *Vet. Parasitol.* 161:270-275.
- Sloss, M.W., 1970. In: *Veterinary Clinical parasitology*, 4th ed. The Iowa State University, Ames. 181 p.
- Soares, A.O., Souza, A.D., Feliciano, E.A., Rodrigues, A.F., D'Agosto, M. and Daemon, E., 2006. Evaluation of ectoparasites and hemoparasites in dogs kept in apartments and houses with yards in the city of Juiz de Fora, Minas Gerais, Brazil. *Rev. Bras. Parasitol. Vet.*, 15:13-16.
- Soulsby, E.J.L., 1982. *Helmiths, arthropods and protozoa of domesticated animals*. 7th Ed., Bailliere and Tindall. London.



- Tinoco-Gracia, L., Quiroz-Romero, H., Quintero-Martínez, M.T., RenteríaEvangelista, T.B., González-Medina, Y., Barreras-Serrano, A., Hori-Oshima, S., Moro, M.H. andVinasco, J. 2009. Prevalence of *Rhipicephalussanguineus* ticks on dogs in a region on the Mexico-USA border. *Vet. Rec.*, 164:59-61.
- Valls, A., 2008.Species identification of *Rhipicephalus sanguineus*. *J. Investig. Allergol.Clin.Immunol.*18: 232.
- Walker, J.B., Keirans, J.E. andHorak, I.G., 2000. The genus *Rhipicephalus* (Acari: Ixodidae). A Guide to the brown ticks of the world.Cambridge University Press, Cambridge, UK, 643 Pp.