

## Evaluation of biodegradable agricultural substrates for mass production of entomopathogenic fungi

S K Sain\*, N Sathyanarayana\*\* and P Jeyakumar<sup>1</sup>

National Institute of Plant Health Management, Rajendranagar, Hyderabad -500 030, Telangana, India.

<sup>1</sup>ICAR-Indian Institute of Rice Research, Rajendranagar, Hyderabad -500 030, Telangana, India.

E mail : sain.skumar@gmail.com

The indiscriminate use of chemical pesticides is resulting in pest resurgence, pesticide resistance and health hazards as well as negatively affecting natural beneficial insects and the crop ecosystems in recent years. Thus the biological control becomes a safer alternative to chemical pesticides. Biopesticides based on bacteria, viruses, entomopathogenic fungi and nematodes often have considerable scope as insect-pest management agents (Noris *et al.*, 2002). Entomopathogenic fungi cause lethal infections, perpetuate in target insects and regulate insects and mite population in nature by epizootics (Carruthers and Soper, 1987; Ferron, 1987; McCoy *et al.*, 1988). Their characteristics, namely, safety to non-target organisms, ease and economic feasibility for *in vitro* mass multiplication, make them one of the preferred options for pest management. Among the fungal biopesticides, *Beauveria bassiana*, *Metarhizium anisopliae*, *Nomuraea rileyi* and *Lecanicillium lecanii* (Zimm.) Zare and Gams have been reported to be promising under Indian conditions. Various strains and formulations of *M. anisopliae* have been shown to infect 100 foliar and other soil dwelling insect pest species belonging to different genera (Butt *et al.*, 2001; Kim *et al.*, 2001; Horaczek and Viernstein, 2004; Bruck *et al.*, 2005; Fisher and Bruck, 2008; Jin *et al.*, 2009; Ansari *et al.*, 2009 and 2010).

*L. lecanii* (formerly *Verticillium lecanii* (Zimm.) Viegas), known as a "white-halo" fungus, belongs to the class Sordariomycetes and reproduces asexually (anamorphic stage) by conidia without fruiting bodies (Hajek, 1997; Zare and Gams, 2001). The market for biocontrol agents (BCAs) has been growing continuously over the last few decades due to improved awareness on the adverse environmental impacts of chemical pesticides. Production of adequate quantities of a good quality inoculum of any biocontrol organism is a prerequisite for receiving excellent results of a biocontrol programme. Production of entomopathogenic fungi has been successfully accomplished in many synthetic and semisynthetic media. Carbon and nitrogen are the most vital nutrients required for growth and sporulation (Campbell *et al.*, 1983). Natural media, which

are invariably rich in carbon and nitrogen, were proved to support the growth and sporulation of fungi. Broken sorghum grain is reported to produce lesser conidia of *L. lecanii* than cooked rice and bran per gram substrate (Feng *et al.*, 2000; Lakshmi *et al.*, 2001). Malt extract peptone medium yielded more conidia/cm<sup>2</sup> of *L. lecanii*, compared to dextrose peptone yeast extract agar medium (Kamp and Bidochka, 2002). Similarly, the biphasic state production by using molasses yeast broth and rice grains produced the spore count of 1.70g/100g (Derakhshan *et al.*, 2008) and in potato dextrose broth and rice after 10 days, it ranges from 0.23 to 1.75 × 10<sup>9</sup> spores/ml (Nirmala *et al.*, 2006). Though different methods of mass production of entomopathogenic fungi are available.

The development of cheaper method of mass production using locally available materials will be highly useful. Thus, the present study was undertaken to evaluate agriculture products like grains of rice, wheat, ragi, sorghum, pearl millet and maize alone and in combination with carbon and nitrogen sources *i.e.*, 2% jaggery and 0.5% yeast, to find out economically viable substrate for the mass production of *Metarhizium anisopliae* and *L. lecanii* with good number of spores.

The present study was carried out at Center for Biological Control (CBC), National Institute of Plant Health Management (NIPHM) during 2012 and 2013. Laboratory experiments were conducted to test the effect of different agricultural products in solid and enriched solid agricultural substrates on mass production of entomopathogenic fungi (EPFs). The cadavers of the insects or the small hoppers that appeared to be infected by fungi were collected in screw cap vials (18 × 4 mm) during survey and brought to the laboratory and entomopathogens were isolated on specific media. Mycosed insects collected from the fields were surface sterilized with 5 per cent sodium hypochlorite for few seconds and then thoroughly rinsed with sterile distilled water several times. The excess water was removed by keeping the diseased insects on Whatman filter paper

\*Present address: ICAR-Central Institute for Cotton Research Regional Station, Sirsa -125 055, Haryana, India

\*\*Present address: Directorate of Plant Protection Quarantine & Storage, Old CGO Complex, NH-IV, Faridabad- 121001, Haryana, India.

No. 1. In a sterile Petri dish, the diseased-insect specimens were cut into small pieces with the help of sterile blade and the bits were aseptically transferred to a culture plate containing selective medium [sabourand maltose agar enriched with 1% yeast extract (SMYA)] with the help of sterile inoculation needle. The inoculated Petri plates were kept in BOD at  $25 \pm 1$  °C under constant observation for the growth and development of microorganisms. Diseased larvae were also kept on moist filter paper in Petri dish for mycelial growth and sporulation. The diseased larvae either showed whitish mycelial surface growth as characteristic of *L. lecanii* or slight darker green colour growth on the rice plant hoppers as characteristic of *M. anisopliae*. After 5 days of incubation, the microorganisms were sub-cultured for purification. Preliminary identification of fungi was made based on the morphological character with the help of the Atlas of entomopathogenic fungi (MaCoy *et al.*, 1988; Humber, 1997). The identified fungi *M. anisopliae* Ma-11 (Accession No. NFCT 5805.13) and *Lecanicillium lecanii* (Accession No. NFCT 5806.13) are available at CBC-NIPHM. All the cultures were maintained on SMYA and potato dextrose agar (PDA) slants. A loopfull of inoculum from sub cultured plates of *M. anisopliae* and *L. lecanii* was transferred to PDA slants and maintained as pure culture. For laboratory studies, both fungi were cultured on PDA medium, separately. After sporulation, conidia from the medium were harvested by washing them thoroughly with sterilized water containing Tween-80 (0.2%) for immediate use. Otherwise, spores were harvested with the help of a small sterile metal spatula. Harvested conidia were air dried under laminar air flow and stored in small air tight screw cap vials (10 cm with 2.5 cm diameter) in refrigerator at 4 °C for further studies.

Whole grains and their by-products of rice, wheat, sorghum, pearl millet, finger millet, maize, barley, black gram seed bran, rice bran were collected from the local market and used for mass multiplication and estimating the sporulation of *M. anisopliae* and *L. lecanii* at 28 °C. Each sample of 100 g grains was taken, washed well and equal amount of distilled water (w/v) (pH 7.0) was added and packed separately in individual autoclavable polypropylene bags [Hi Media 7" (B) x 11" (H) size]. The bags were plugged using non-absorbent cotton with the support of a piece of 1.5 inch PVC pipe at the opening of the bag and sterilized by autoclaving at 121 °C and 15 lbs pressure for 20 minutes (Mazumdar *et al.*, 1995). Each bag was considered as a replicate and three bags were used for each treatment. In addition to the set of plain grain media, another set of all grain media amended with 0.5% yeast and 2% Sugarcane "Jaggery" (65-85% sucrose) separately

for nutrient enrichment was taken. For enrichment of these grain media, 5 g yeast and 20 g sugarcane "jaggery" were mixed in 1000 ml distilled water. Hundred gram grains of each crop were cleaned, washed well transferred separately in 7" (B) x 11" (H) autoclavable bags (Hi Media) and 100 ml of yeast and sugarcane "jaggery" solution was added to each bag separately and sterilized as mentioned earlier. Three replications were maintained for each grain medium. After, autoclaving the bags were allowed to cool down upto the lukewarm temperature and 1 ml of freshly prepared conidial suspension of *M. anisopliae* and *L. lecanii* (10-days old) was inoculated in each bags, separately under aseptic conditions. The inoculated bags were incubated in BOD incubator at  $25 \pm 1$  °C separately for 10 days. To avoid clumping, after every three-days of inoculation, the bags were shaken vigorously to separate the grain and to break the mycelial mat. Ten days after incubation, grain along with fungal biomass of each sample was transferred in sterilized porcelain trays separately and dried. The biomass was dried to constant weight in hot air oven at 60-65 °C for 12 hours. The determination of dry mycelial weight done separately by weighing the biomass along with fungal mycelium and conidia (Cochrane, 1958).

After 10 days of incubation, 1g homogenous dried grain media sample drawn from each replicate of uniformly sporulating bags was transferred to 10 ml sterilized distilled water containing Tween 80 (0.2%) solution in test tubes. The test tubes were shaken in mechanical shaker for 10 min. The suspension was filtered through double layered muslin cloth and then filtered through Whatman filter paper No.1. The filtered suspension was subjected to the serial dilution up to 12 times. A drop of conidial suspension of *M. anisopliae* and *L. lecanii* was placed on separate hemocytometer. The number of conidia of entomopathogenic fungi was counted under phase contrast microscope at higher resolution in the middle square (V) of Neubauer hemocytometer. The conidial count was determined to find out the number of conidia in 1 g of the dry biomass obtained from each grain media separately using the formula: Conidia per ml = the average count per square ( $1 \text{ mm}^2$ )  $\times$  the dilution factor  $\times 10^4$ . Colony forming units (cfu) were estimated by plating technique. From the filtered stock solution, further dilutions were made to upto  $10^9$  dilution. The study was carried out in a randomized complete block design (RCBD) with three replications. The biomass and conidial production of both the fungi from different substrates were subjected to analysis of variance (ANOVA). The means were separated using least significant difference (LSD) and differences between treatments were considered significant at probability level  $P = 0.05$ .

The results presented in Table 1 revealed that among the grain media tested, the dry weight of *M. anisopliae* per 100 g media was significantly higher in pearl millet (47.7 g) followed by finger millet (45.5 g) and maize (44.2 g) than other grain media at 10 days after inoculation as compared to other grain media. However, there was no significant difference ( $P=0.05$ ) among these three grain media in terms of dry biomass weight of *M. anisopliae*. The significantly highest dry biomass of *L. lecanii* in sorghum (62.6 g) followed by wheat and maize grain media, 60.0 and 58.6 g, respectively than others. There was no significant difference between wheat and maize in terms of dry biomass weight. Compared to the other grain media the spore production of *M. anisopliae* was significantly higher on wheat ( $11.6 \times 10^{10}$ ) and rice ( $11.5 \times 10^{10}$ ) followed by sorghum ( $10.4 \times 10^{10}$ ) than others. The spore count of *L. lecanii* was highest on sorghum and pearl millet enriched with yeast and "jaggery" media ( $11.4 \times 10^{10}$  and  $11.2 \times 10^{10}$ ) should not be superscript, respectively, but not significantly different from each other.

In case of enrichment of these grain media with 0.5% yeast and 2% sugarcane "jaggery" (65-85% sucrose), there was a significant boost in dry biomass production as well as spore count. In case of *M. anisopliae*, enriched pearl millet jaggery and finger millet jaggery media recorded the highest dry weight per 100 g (54.6 and 53.1, respectively) but both were at par with finger millet and pearl millet was statistically insignificant. In case of *L. lecanii*, enriched sorghum jaggery medium resulted in the highest dry biomass yield (66.6 g per 100 g). This was followed by sorghum alone, enriched maize and wheat media (62.6, 61.4 and 61.1, respectively). There was no statistically significant difference among maize, wheat, and sorghum enriched media and jaggery and sorghum alone. The spore count of *M. anisopliae* was recorded highest on enriched rice jaggery and wheat jaggery media ( $12.8 \times 10^{10}$  and  $12.7 \times 10^{10}$  per gram of dry biomass, respectively). But the statistical comparison between the rice and wheat media was insignificant. However, the spore count of

**Table 1. Effect of various grain media alone and or enriched with jaggery plus yeast (JY) on biomass and spore production of *Metarhizium anisopliae* and *Lecanicillium lecanii***

Media	Spore count $\times 10^{10}$ /substrate alone		Fresh weight (g)		Dry weight (g)	
	<i>M. anisopliae</i>	<i>L. lecanii</i>	<i>M. anisopliae</i>	<i>L. lecanii</i>	<i>M. anisopliae</i>	<i>L. lecanii</i>
Rice	11.5	10.2	81.5	82.0	34.9	44.9
Wheat	11.6	9.4	64.1	86.8	33.1	60.0
Sorghum	10.4	11.4	62.1	89.3	42.5	62.6
Pearl millet	9.8	11.2	91.5	90.6	47.7	42.0
Finger millet	10.2	10.4	89.3	88.2	45.5	41.2
Matze	9.5	10.3	94.8	93.8	44.2	58.6
Barley	10.2	10.5	75.9	85.1	27.4	41.0
Black gram seed bran	10.2	9.5	59.6	65.3	41.1	43.1
Rice bran	7.2	6.5	60.1	63.9	42.4	45.2
Rice +JY	12.8	12.7	83.3	83.6	35.5	49.7
Wheat +JY	12.7	11.2	83.5	88.5	36.2	61.1
Sorghum+ JY	11.6	12.6	63.0	91.4	45.6	66.6
Pearl millet + JY	11.1	11.8	92.8	91.0	54.6	45.8
Finger millet +JY	10.6	11.2	91.8	87.9	53.1	44.3
Maize + JY	10.7	11.4	95.7	95.6	46.8	61.4
Barley +JY	11.4	11.8	78.4	86.3	31.3	43.6
Black gram seed bran +JY	11.3	10.9	64.3	68.2	42.5	44.9
Rice bran +JY	8.4	7.8	65.8	67.9	43.9	46.7
SEM $\pm$	0.995	1.033	3.233	3.778	2.271	2.657
CD ( $P = 0.05$ )	2.022	2.099	6.569	6.864	4.615	5.403




Figure 1. Total c.f.u. ( $1 \times 10^{12}/100$  g) of *Metarhizium anisopliae* recorded on different media alone and enriched with jaggery plus yeast (JY)

*L. lecanii* was recorded highest but at par on rice and sorghum medium enriched with yeast plus jaggery ( $12.7 \times 10^{10}$  and  $12.6 \times 10^{10}$ , respectively). Overall, selection of medium was done based on total spore production in terms of total c.f.u. from the harvested biomass. The highest number of c.f.u. of *M. anisopliae* was recorded in pearl millet medium enriched with yeast plus jaggery ( $5.6 \times 10^{12}$  per 100 g); however, the highest c.f.u. ( $8.4 \times 10^{12}/100$  g) of *L. lecanii* was recorded in sorghum medium enriched with yeast plus jaggery. Similarly, sorghum was reported to be ideal for the mass production of *L. lecanii* ( $11.31 \times 10^{10}$  spores/100 g) while, pearl millet was also found as suitable medium for the spore production ( $10.17 \times 10^{10}$  spores/100g) (Sahayaraj *et al.*, 2008) and the spore production can further be maximized by incorporation of molasses at 4% concentration (Derakhshan *et al.*, 2008). Broken sorghum grains are reported to produce  $15 \times 10^8$  conidia /g (Lakshmi *et al.*, 2001), which is in confirmation with the findings of the current study. While Feng *et al.* (2000) found cooked rice and bran substrate to produce higher spores of  $1.5$  and  $1.4 \times 10^9$  spores /g for *L. lecanii*. Grains are cheap, easily available and act as best nutritive media for the mass multiplication of many micro and macro organisms. According to Ibrahim and Low (1993) and Sharma *et al.* (2002), rice was found to be the suitable medium for the mass culture of deuteromycete fungi. The above findings are closely in agreement with Bharati *et al.* (2007), who recorded considerably low spore yields in agro wastes as compared to food grains and Sahayaraj *et al.* (2008) who have observed that sorghum recorded maximum spore production of *Paecilomyces fumosoroseus* and *L. lecanii*. Dangar *et al.* (1991) also observed similar findings in *M. anisopliae*. In Brazil,




Figure 2. Total c.f.u. ( $1 \times 10^{12}/100$  g) *Lecanicillium lecanii* recorded on different media alone and enriched with jaggery plus yeast (JY)

*M. anisopliae* has been found to produce conidial yields of 5-15 times higher using rice bran/rice husk substrate mixtures than yields usually obtained from rice grains, with viabilities of higher than 85% (Dorta *et al.*, 1990). Sahayaraj *et al.* (2008) found that among other grain media sorghum recorded maximum spore production of *P. fumosoroseus* and *L. lecanii*. Puzari *et al.* (1997) further reported that rice husk supplemented with 2% dextrose solution recorded more sporulation of *M. anisopliae*.

In conclusion, the tested fungi were able to grow on a wide variety of agriculture products and by products of solid state and this can be useful to farmers to culture these fungi easily. Moreover, the grain media enriched with low cost carbon and nitrogen sources could further enhance the fungal biomass production. The pearl millet medium enriched with yeast plus jaggery showed significantly high growth rate of *M. anisopliae* in terms of total c.f.u production ( $8.4 \times 10^{12}$  per 100 gram) as compared to other media. However, comparatively sorghum grain media enriched with jaggery plus yeast showed significantly enhanced growth rate of *L. lecanii* in terms of total c.f.u production ( $8.4 \times 10^{12}/100$  g). For each new strain and species of entomopathogenic fungi, the optimal conditions and the source of nutrition required for large scale production may differ from those found previously and need to be tested to produce a high level of infective inoculum at minimal cost. The grain media can be a very good low cost and a suitable alternative for small scale industry and farm level mass multiplication of these beneficial microbial fungi. This method can ultimately serve as an economic way of making quality biopesticides easily available to the farmers.

## References

- Ansari M A, Evans M and Butt T M 2009. Identification of pathogenic strains of entomopathogenic nematodes and fungi for wireworm control. *Crop Protection* **28** : 269-272.
- Ansari M A, Carpenter S and Butt T M 2010. Susceptibility of *Culicoides* biting midges larvae to the entomopathogenic fungus, *Metarhizium ansopliae*. prospects for Bluetongue Vector control. *Acta Tropica* **113** : 1-6.
- Bharati T, Kulkarni J H, Krishnaraj P U and Alagawadi A R 2007. Evaluation of Food Grains and Agro Wastes for Sporulation of *Metarhizium ansopliae* (Ma2). *Karnataka Journal of Agricultural Sciences* **20** : 424-425.
- Bruck D J, Snelling J E, Dreves A J and Jaronski S T 2005. Laboratory bioassays of entomopathogenic fungi for control of *Della radicum* (L.) larvae. *Journal of Invertebrate Pathology* **89** : 179-183.
- Butt T M, Jackson C W and Murugan W 2001. Fungi as biocontrol agents, progress, problems and potentials. CAB International, Wallingford, UK. Pp 240-242.
- Campbell R K B, Arnes G L, Cartwright B O and Eikenbary R D 1983. Growth and sporulation of *Beauveria bassiana* and *Metarhizium ansopliae* in a basal medium containing various carbohydrate sources. *Journal of Invertebrate Pathology* **41** : 117-121.
- Carruthers R I and Soper R S 1987. Fungal diseases. Epizootiology of Insect Diseases (eds. Fuxa J R and Tanada Y) New John Wiley and Sons, New York, USA. Pp. 357-416.
- Cochrane V M 1958. Physiology of fungi. John Wiley and Sons Inc., New York. 524 pp.
- Dangar T E, Geetha L and Pillai G B 1991. Mass production of the entomopathogenic fungi *Metarhizium ansopliae* in coconut water. *Journal of Plantation Crops* **19** : 54-69.
- Derakhshan A, Rabindra R J, Ramanujam B and Rahimi M 2008. Evaluation of different media and methods of cultivation on the production and viability of entomopathogenic fungi *Verticillium lecanii* (Zimm.) Viegas. *Pakistan Journal of Biological Sciences* **11** : 1506-1509.
- Dorta B, Bosch A, Arcas J A and Ertola R J 1990. High level of sporulation of *Metarhizium ansopliae* in a medium containing by-products. *Applied Microbiology and Biotechnology* **33** : 712-715.
- Feng K C, Liu B L and Tzeng Y M 2000. *Verticillium lecanii* spore production in solid-state and liquid-state fermentations. *Bioprocess and Biosystems Engineering* **23** : 25-29.
- Ferron P 1987. Biological control of insect pests by entomopathogenic fungi. *Annual Review of Entomology* **23** : 409-442.
- Fisher J R and Bruck D J 2008. Biology and control of root weevils on berry and nursery crops in Oregon. *Acta Horticulturae* **777** : 345-351.
- Hajek A E 1997. Ecology of terrestrial fungal entomopathogens. *Advances in Microbial Ecology* **15** : 193-249.
- Horaczek A and Viernstein H 2004. Comparison of three commonly used drying technologies with respect to activity and longevity of areal conidia of *Beauveria brongniartii* and *Metarhizium ansopliae*. *Biological Control* **31** : 65-71.
- Humber R A 1997. Manual of Techniques in Insect Pathology. Academic Pree, London. Pp 153-155.
- Ibrahim Y B and Low W 1993. Potential of mass production and field efficacy of isolates of the entomopathogenic fungi *Beauveria bassiana* and *Paecilomyces fumosoroseus* on *Plutella xylostella*. *Journal of Invertebrate Pathology* **39** : 222-232.
- Jin X, Uguine T A, Chen Jian and Streett D A 2009. Method for determining the best hydrophilic-lipophilic balance (HLB) number for a compatible non-ionic surfactant in formulation development for aerial conidia of *Metarhizium ansopliae* (Hypocreales: Clavicipitaceae). *Biocontrol Science and Technology* **19** : 341-347.
- Kamp A M and Bidochka M J 2002. Conidium production by insect production fungi on commercially available agars. *Letters in Applied Microbiology* **35** : 74-75.
- Kim J J, Goettel M S and Gillespie D R 2001. Potential of *Lecanicillium* species for dual microbial control of aphids and cucumber powdery mildew fungus, *Sphaerotheca fuliginea*. *Biological Control* **40** : 327-332.
- Lakshmi S M, Alagammai P L and Jayaraj K 2001. Studies on mass culturing of the entomopathogen white halo fungus *Verticillium lecanii* on three grain media and its inefficacy on *Helicoverpa armigera*. Microbials. Insect Pest Management (eds. Igbachimuthu S, Sen S), Oxford and IBH publishing Pvt Ltd, New Delhi. Pp 23-27.

- Mazumdar D, Hallsworth Magan, Puzari K C and Hazarika L K 1995. Mass production of *Beauveria bassiana* and its potentiality on rice hispa. *Indian Phytopathology* **48** : 275-278.
- McCoy C W, Samson R A and Boucias D G 1988. Entomogenous fungi. Handbook of Natural Pesticides, Microbial Insecticides, Part A. Entomogenous Protozoa and Fungi (eds. Ignoffo C M and Mandava N B). Boca, Raton, Fla: Mr ic Press. Vol. 5.
- Nirmala R, Ramanujam B, Rabindra R J and Rao N S 2006. Growth parameters of some isolates of entomofungal pathogens and production of dust-free spores on rice medium. *Journal of Biological Control* **19** : 129-133.
- Noris R F, Chen E P S and Kogn M 2002. Concepts integrated pest management. Premise Hall of India Private Limited, New Delhi.
- Puzari K C, Sharma D K and Saranka L K 1997. Media of mass production of *Beauveria bassiana*. *Journal of Biological Control* **11** : 96-100.
- SahayaraJ K and Karthick Raja Namasivayam S 2008. Mass production of entomopathogenic fungi using agricultural products and by products. *African Journal of Biotechnology* **7** : 1907-1910.
- Sharma S P, Gupta R B L and Yadava C P S 2002. Selection of a Suitable Medium for Mass Multiplication of Entomofungal pathogens. *Indian Journal of Entomology* **14** : 255-261.
- Zare R and Gams W A 2001. Revision of *Verticillium* section Prostrata. IV. The genera *Lecanicillium* and *Simplicillium* gen. nov. *Nova Hedwigia* **73** : 1-50.

Received : 12-02-2016

Revised : 15-09-2017

Accepted : 27-01-2018