

Cornous Publications

e-ISSN: 2583 - 0147



AgroScience Today

(An Online Magazine)

Volume 2 Issue 12

December 2021



Gauri Jairath

Scientist
ICAR-IVRI Regional Station
Palampur, Himachal Pradesh
India -176 061

Devi Gopinath

Scientist
ICAR-IVRI Regional Station
Palampur, Himachal Pradesh
India -176 061

Ajayta Rialch

Scientist
ICAR-IVRI Regional Station
Palampur, Himachal Pradesh
India -176 061

Gorakh Mal

ICAR-IVRI Regional Station
Palampur, Himachal Pradesh
India -176 061

Birbal Singh

Principal Scientist
ICAR-IVRI Regional Station
Palampur, Himachal Pradesh
India -176 061

Rinku Sharma

Senior Scientist
ICAR-IVRI Regional Station
Palampur, Himachal Pradesh
India -176 061

Corresponding Author

Gauri Jairath
gaurilpt@gmail.com

Plant - Based Antibiotic Adjuvants - A Roadmap to Combat Antibiotic Resistance

Multidrug-resistant bacteria have now emerged as a global threat due to the emergence of new or re-emergence of old pathogens. This alarming situation demands either to find new/modified effective antibiotics or alternatives to reduce the overall load of infectious diseases both in the veterinary and public domain. Antibiotic adjuvants are non-antibiotic compounds that potentiate the existing antibiotic activity. The Himalayan region being a big basket and hot spot of biodiversity, inhabiting a diverse flora of medicinal plants/herbs can be evaluated for antibiotic adjuvants. These adjuvants work either by blocking the main bacterial resistance mechanisms or enhancing the antimicrobial action of the drug. The adjuvants assist in polypharmacy on bacteria by de-energizing the efflux pump channel of bacteria to escape antibiotic action and by allowing antibiotics to have greater internal access. Thus, the plant-based antibiotic adjuvants have the potency to re-empower the existing antibiotics, restoring their activities against target pathogen that too with minimal side effects/toxicity being natural in origin.

INTRODUCTION

In the global scenario, developing countries like India have been surveyed as a major consumption unit for antibiotics with the highest consumption rate of 7.5 units per person. In 2010, the total antibiotic consumption of India was highest at a rate of 12.9 x 10⁹ units followed by China (10.0 x10⁹ units) and the US (6.8 x10⁹ units) [Laxminarayan and Chaudhury, 2016]. It has been further reported that BRICS countries (Brazil, Russia, India, China, and South Africa) are the main consumers of antibiotics and have contributed almost 76% to an

overall increase in consumption during the last decade out of which India shares 23% (Van Boeckel et al., 2014). Further, as per reports of Washington-based Center of Disease Dynamics, Economics and Policy (CDEEP), the percentage change in total use of antibiotics has been about 47.40% between 2010 and 2020. Due to the non-judicial use of antibiotics, antibiotic resistance has emerged as a major issue of concern. Consequently, multidrug-resistant (MDR) bacteria have now emerged as a global threat in the 21st century, as doctors are running short of potent antibiotics against them, resulting in 13 million deaths every year due to the emergence of new or re-emergence of old pathogens (Kourtesi et al., 2013). This alarming situation demands either to find new/modified effective antibiotics or the alternative one to reduce the overall load of microbes both in the veterinary and public domain.

Antibiotic adjuvants have the properties to enhance the efficacy of existing antibiotics. The concept of antibiotic adjuvant can be clearly understood with the example of clavulanic acid in combination with amoxicillin which is a widely used medicine both for humans as well as for veterinary practices. Though the clavulanic acid exhibits weak antibacterial activity, but inhibits β -lactamases and potentiates the antibacterial activities of amoxicillin. (Gibbons et al., 2003). Antibiotic adjuvants from *Buxus sempervirens* and *Cytisus striatus* (isoflavonoids) have been recently identified by Abreu et al. (2016; 2017) as antibiotic adjuvants against MRSA. Similarly, triterpenoids from *Taraxacum officinale* have been reported to be effective against uropathogenic bacteria (Díaz et al., 2018).

Several plant metabolites are being investigated for their antibacterial activity and their mechanism of action has been studied (Gibbons et al., 2004; Kang et al., 2011; Abreu et al., 2012). But the pharmaceutical/clinical applications of plant-derived antimicrobials are very few, despite having antibacterial efficacy due to weak or moderate activity with minimal inhibitory concentrations (MICs) typically in the range of 100–1000 $\mu\text{g}/\text{mL}$ (Tegos et al., 2002). Therefore, a therapeutic contribution can be enhanced by purifying and identifying the bioactive compounds responsible for antibacterial activity to harness their probable use as antibiotic adjuvants (Abreu et al., 2017).

Since, Indian Himalayan region especially, North Western Himalayan is a big basket and hot spot of biodiversity, inhabiting a diverse flora of medicinal plants/herbs (Samant et al., 2007) is a promising source of antimicrobial bioactive compounds owing to their phenolics, flavonoids, isoflavonoids content and many more unidentified compounds (Vidyarthi et al., 2013; Sharma et al., 2014; Thakur et al., 2016). The defense system of plants can be tapped to potentiate the antibacterial effects of available antibiotics and for their possible to act as antibiotic adjuvants. WHO has published a list of bacteria for which there is an urgent need of new antibiotics among which *Escherichia coli* and *Staphylococcus aureus* are on the priority list (WHO, 2017). Nowadays, antibiotics like ciprofloxacin (fluoroquinolone), tetracycline, cefixime (3rd generation cephalosporins), ampicillin (β -lactam) have limited use, due to increased bacterial tolerance. The efficacy of such antibiotics can be enhanced in combination with antibiotic adjuvants. For example, isoflavonoids from *Cytisus striatus* with strong efflux inhibitory activity have been proven to be possible adjuvants for erythromycin against MRSA (Abreu et al., 2017). Many mechanisms like active efflux of antibiotics by bacteria mediated by efflux pumps have been suggested to be a key mechanism in bacterial resistance to many antibiotics. The identification of a probable mechanism of action against bacteria is the most important step to bring them to use in the therapeutic/pharmaceutical industry. Thus, modification of existing antibiotics by combining them with potential antibiotic adjuvants could be a possible way to tackle the resistant bacterial strains of human and animal origin.

CONCLUSION

The current scenario of emerging antibiotic resistance and re-emerging diseases has necessitated the finding of ameliorative measures to reduce the infectious load and combat antibiotic resistance. The Himalayan bioresources may be explored for antibiotic adjuvants to boost antibiotic pharmacy with minimal negative effects. Antibiotic adjuvants being potentiators may enhance the existing efficacy of the antibiotics and make them effective against resistant microbes.

REFERENCES

Abreu, A. C., Coqueiro, A., Sultan, A. R., Lemmens, N., Kim, H. K., Verpoorte, R., Wamel, W. J. B., Simões, M. and Choi, Y. H. (2017). Looking to nature for a new

- concept in antimicrobial treatments: isoflavonoids from *Cytisus striatus* as antibiotic adjuvants against MRSA. *Scientific Reports* 7(1): 3777 (article no.) doi:10.1038/s41598-017-03716-7.
- Abreu, A. C., McBain, A. J. and Simoes, M. (2012). Plants as sources of new antimicrobials and resistance-modifying agents. *Natural Product Reports* 29: 1007–1021.
- Abreu, A. C., Paulet, D., Coqueiro, A., Malheiro, J., Borges, A., Saavedra, M. J., Choi, Y. H. and Simões, M. (2016). Antibiotic adjuvants from *Buxus sempervirens* to promote effective treatment of drug-resistant *Staphylococcus aureus* biofilms. *RSC Advances* 6(97): 95000–95009. doi:10.1039/c6ra21137b.
- Díaz, K., Espinoza, L., Madrid, A., Pizarro, L. and Chamy, R. (2018). Isolation and Identification of compounds from bioactive extracts of *Taraxacum officinale* Weber ex F. H. Wigg. (Dandelion) as a potential source of antibacterial agents. *Evidence-Based Complementary and Alternative Medicine* Article ID 2706417: 1–8. doi:10.1155/2018/2706417.
- Gibbons, S., Moser, E. and Kaatz, G. W. (2004). *Catechin gallates* inhibit multidrug resistance (MDR) in *Staphylococcus aureus*. *Planta Medicine* 70: 1240–1242 (2004).
- Gibbons, S., Oluwatuyi, M. and Kaatz, G. W. (2003). A novel inhibitor of multidrug efflux pumps in *Staphylococcus aureus*. *Journal of Antimicrobial Chemotherapy* 51: 13–17.
- Kang, H. K., Kim, H. Y. and Cha, J. D. (2011). Synergistic effects between silibinin and antibiotics on methicillin-resistant *Staphylococcus aureus* isolated from clinical specimens. *Biotechnology Journal* 6: 1397–1408.
- Kourtesi, C. Ball, A. R., Huang, Y. Y., Jachak, S. M., Vera, D. M., Khondkar, P., Gibbons, S., Hamblin, M. R. and Tegos, G. P. (2013). Microbial efflux systems and inhibitors: approaches to drug discovery and the challenge of clinical implementation. *Open Microbiology Journal* 7: 34–52.
- Laxminarayan, R. and Chaudhury R. R. (2016) Antibiotic Resistance in India: Drivers and Opportunities for Action. *Plos Med* 13(3): e1001974. doi: 10.1371/journal.pmed.1001974
- Samant, S. S., Pant, S., Singh, M., Lal, M., Singh, A., Sharma, A. and Bhandari, S. (2007). Medicinal plants in Himachal Pradesh, North Western Himalaya, India. *International Journal of Biodiversity Science and Management* 3: 234–251.
- Sharma, V., Rishma and Kaur, K. (2014). Identification and characterization of medicinally important plants of Kangra valley with synergistic effects of traditional antibiotics against microbial infections. *The Journal of Phytopharmacology* 3(2): 102–112.
- Tegos, G., Stermitz, F. R., Lomovskaya, O. and Lewis, K. (2002). Multidrug pump inhibitors uncover remarkable activity of plant antimicrobials. *Antimicrobial Agents and Chemotherapy* 46: 3133–3141.
- Thakur, M., Asrani, R. K., Thakur, S., Sharma, P. K., Patil, R. D., Lal, B. and Parkash, O. (2016). Observations on traditional usage of ethnomedicinal plants in humans and animals of Kangra and Chamba districts of Himachal Pradesh in North-Western Himalaya, India. *Journal of Ethnopharmacology* 191: 280–300. doi:10.1016/j.jep.2016.06.033.
- Van Boeckel, T. P., Gandra, S., Ashok, A., Caudron, Q., Grenfell, B. T., Levin, S. A. and Laxminarayan, R. (2014). Global antibiotic consumption 2000 to 2010: An analysis of national pharmaceutical sales data. *Lancet Infectious Diseases* 14(8): 742–50. doi: 10.1016/S1473-3099(14)70780-7.
- Vidyarthi, S., Samant, S. S. and Sharma, P. (2013). Traditional and indigenous uses of medicinal plants by local residents in Himachal Pradesh, North Western Himalaya, India. *International Journal of Biodiversity Science, Ecosystem Services and Management*, 9(3): 185–200. doi:10.1080/21513732.2013.823113
- WHO. (2017). WHO publishes list of bacteria for which new antibiotics are urgently needed. Press release February 27. <https://www.who.int/en/news-room/detail/27-02-2017-who-publishes-list-of-bacteria-for-which-new-antibiotics-are-urgently-needed>.