



## Review

## Role of MAPK/MNK1 signaling in virus replication

Ram Kumar<sup>a,b</sup>, Nitin Khandelwal<sup>a</sup>, Riyesh Thachamvally<sup>a</sup>, Bhupendra Nath Tripathi<sup>a</sup>, Sanjay Barua<sup>a</sup>, Sudhir Kumar Kashyap<sup>b</sup>, Sunil Maherchandani<sup>b</sup>, Naveen Kumar<sup>a,\*</sup>

<sup>a</sup> Virology Laboratory, National Centre for Veterinary Type Cultures, ICAR-National Research Centre on Equines, Hisar, Haryana 125001, India

<sup>b</sup> Department of Veterinary Microbiology and Biotechnology, Rajasthan University of Veterinary and Animal Sciences, Bikaner, Rajasthan 334001, India



## ARTICLE INFO

## Keywords:

Signaling pathway  
MAPK  
MNK1  
Virus replication  
Antiviral drugs

## ABSTRACT

Viruses are obligate intracellular parasites; they heavily depend on the host cell machinery to effectively replicate and produce new progeny virus particles. Following viral infection, diverse cell signaling pathways are initiated by the cells, with the major goal of establishing an antiviral state. However, viruses have been shown to exploit cellular signaling pathways for their own effective replication. Genome-wide siRNA screens have also identified numerous host factors that either support (proviral) or inhibit (antiviral) virus replication. Some of the host factors might be dispensable for the host but may be critical for virus replication; therefore such cellular factors may serve as targets for development of antiviral therapeutics. Mitogen activated protein kinase (MAPK) is a major cell signaling pathway that is known to be activated by diverse group of viruses. MAPK interacting kinase 1 (MNK1) has been shown to regulate both cap-dependent and internal ribosomal entry sites (IRES)-mediated mRNA translation. In this review we have discuss the role of MAPK in virus replication, particularly the role of MNK1 in replication and translation of viral genome.

## 1. Introduction

Cell signaling is a part of a complex system of communication through which living cells interact with the neighbouring cells and extracellular environment (Denef, 2008). Cells are equipped with glycoproteins or glycolipid receptors on the plasma membrane through which they respond to changes in their immediate environment. When a complementary ligand (signaling molecule) binds to the receptor, it initiates a chain of events within the cell called signal transduction, ultimately resulting into a response. This ability of the cells to respond to their microenvironment solely determines growth, development, tissue injury/repair, cell homeostasis and immunity. A cell can communicate signals to neighbouring cells in multiple ways, which include: direct transfer of ions/small molecules through pores in the membrane, endocrine signaling that utilizes hormones, paracrine signaling by secreting chemicals into the common intercellular space, autocrine signaling to alter its own extracellular environment, juxtacrine signaling by making physical contact with adjacent cells and synaptic signaling (nervous system) (Brucher and Jamall, 2014).

Cell signaling starts with an external stimuli (signals); most cell signals are chemical in nature but mechanical stimuli are also possible (Cargnello and Roux, 2011). Cells employ numerous intracellular signaling pathways for transmitting information within the cell. However, a cross talk among these cell signaling pathways is quite possible (Vert

and Chory, 2011). The signaling pathways may be activated in response to external stimuli or via the messengers (ligand/information/metabolic) that are generated within the cell following insults. The information in signaling pathways is conveyed either through protein-protein interactions or via diffusible elements usually referred to as second messengers (Cargnello and Roux, 2011).


## 2. MAPK signaling

Mitogen activated protein kinase (MAPK) is an important cell signaling pathway that converts extracellular stimuli into a wide range of cellular responses (Cargnello and Roux, 2011). MAPK signaling is activated by a variety of chemical and physical stimuli such as cytokine, hormone, growth factors, pathogens (including viruses), osmotic stress, heat, oxidative stress and microtubule disorganization (Boulton et al., 1990; Raman et al., 2007; Rose et al., 2010; Widmann et al., 1999). MAPK pathway regulates gene expression, mitosis, cell survival, apoptosis, metabolism and cell differentiation (Roux and Blenis, 2004; Shaul and Seger, 2007). The information in MAPK pathway is transmitted via protein-protein interactions. By phosphorylating serine and threonine residues in diverse groups of proteins, MAPKs convert a variety of extracellular signals into a multitude of cellular responses (Ceballos-Olvera et al., 2010; Shi et al., 2013).

MAPK signaling toolkit is composed of a cell surface receptor that

\* Corresponding author.

E-mail addresses: [naveenkumar.icar@gmail.com](mailto:naveenkumar.icar@gmail.com), [Naveen.Kumar1@icar.gov.in](mailto:Naveen.Kumar1@icar.gov.in) (N. Kumar).



**Fig. 1. MAPK/MNK signaling.** MAPK/MNK signaling pathway is activated by binding of the ligands to their cognate receptors (RTKs/GPCRs). In RTKs, ligand binding induces dimerization and autophosphorylation at its Ser/Thr residues (intracytoplasmic domain). With the help of adaptor proteins- SOS/Grb2, activated RTKs convert Ras-GDP to Ras-GTP. In GPCRs, ligand binding induces structural changes in heterotrimeric subunits. These structural changes trigger generation of secondary messenger, which eventually converts Ras-GDP to Ras-GTP. Upon formation of Ras-GTP, further information in the pathway is conveyed by three sequentially acting evolutionary conserved proteins of MAPK-ERK family namely Raf (MAPKKKs), MEK1/2 (MAPKKs) and ERK1/2 (MAPK). Besides ERK signaling, Ras-GTP also activates those MAPKKKs, which are required for phosphorylation of MEK4/7 and MEK3/6 to trigger JNK and p38 pathway respectively. The ERK1/2 and p38 activation ultimately results in phosphorylation of MNK1. Activated MNK1 regulates functions of several potential downstream substrates. eIF4E and eIF4 G, which help in formation of 5' cap-dependent mRNA translation initiation complex (eIF4 F). hnRNPA1 helps in pre-mRNA processing. PSF involves in post translational processing as well as translocation of mRNA. cPLA2, which regulates production of eicosanoids, a second messenger, plays an important role in immunity and inflammation. Spry2 is associated with interferon production and negative regulation of ERK/MAPK pathway. Besides, MNK1 also participates in regulation of cap-independent mRNA translation, which involves activation of SRPK, mediated via mTOR. By activating transcriptional factors such as c-Jun, c-Fos, STAT1, ATF1/2/6, Elk1, Ets1, p53 and NF-κB, JNK and p38 induce inflammatory response.

receives extracellular signals in the form of chemical or mechanical ligand, a transducer that converts extracellular stimuli into intracellular signals. MAPK module has three evolutionary conserved, sequentially acting proteins namely, MAPK kinase kinase (MAPKKK), MAPK kinase (MAPKK) and MAPK (Fig. 1). Their downstream targets include MAPK scaffolding protein and MAPK target protein (effectors protein) (Cargnello and Roux, 2011). To be enzymatically active, these proteins must undergo phosphorylation (Robbins et al., 1993).

Extracellular signal-regulated kinase (ERK, also known as p42/44 MAPK), Janus kinase (JNK, also known as stress activated protein kinase-1, SAPK1) and p38 MAP kinase (also known as SAPK2/RK) are three major MAPK pathways in mammals (Gaur et al., 2011; Ludwig et al., 2006). ERK1 and ERK2 are key transducers of proliferation

signals and are often activated by mitogens. JNKs and p38 poorly respond to mitogens but are strongly activated under cellular stress (Cargnello and Roux, 2011). Following activation, these cytosolic proteins (MAPK downstream molecules) either remain in cytoplasm or may translocate into the nucleus to activate numerous proteins and/or transcription factors (Cuadrado and Nebreda, 2010; Li et al., 2016) (Fig. 1). These regulatory molecules not only transmit information to the target effectors but also cross talk with other parallel signaling pathways. These mechanisms allow amplification of the signals and generate a threshold for activation of diverse pathways (Cargnello and Roux, 2011; Maroni et al., 2004).

In order to manipulate cellular functions in its favor, viruses interact with MAPK family members- ERK, p38 and JNK. MAPK pathway is

**Table 1**  
Role of MAPK signaling in virus replication.

Signaling pathway	Cellular mediator	Impact on virus replication	References	
RTK family	Abl	Positively regulates transport and release of poxviruses	(Reeves et al., 2005)	
	EGFR	Facilitates vaccinia virus spread	(Langhammer et al., 2011)	
	VEGF	Facilitates ORFV replication and formation of pock-like lesions	(Wise et al., 1999)	
		Facilitates HSV-1 induced stromal keratitis	(Sharma et al., 2011)	
	Src kinase	Positively regulates assembly, maturation and egress of WNV	(Hirsch et al., 2005)	
	NGFR	Positively regulates assembly and maturation of DENV	(Chu and Yang, 2007)	
	PDGFR	Positively regulates RNA synthesis, vRNP export and release (budding) of IAV	(Kumar et al., 2011a)	
		Facilitates KSHV tumor progression	(Cao et al., 2015; Sturzl et al., 1992)	
GPCRs	GPCRs	Promotes membrane fusion and cell to cell spread in hHSV6 (mediated via viral U51 protein)	(Zhen et al., 2005)	
		Positively regulates replication and reactivation of gamma herpesvirus latency (mediated via viral CXCR2 protein)	(Lee et al., 2003)	
Ras/Akt	N-Ras	Positively regulates flavivirus entry and RNA synthesis	(Le Sommer et al., 2012)	
		Promotes HCV replication by facilitating cell survival and establishment of persistent viral infection	(Mannova and Beretta, 2005)	
	Akt	Promotes clathrin-independent endocytosis and internalization of IAV	(Fujioka et al., 2011)	
		Positively regulates PCV2 replication by inhibiting premature apoptosis	(Wei et al., 2012)	
		By expressing anti-apoptotic genes, positively regulates cowpox virus and vaccinia virus replication	(Soares et al., 2009)	
	mTORC2	Inhibits endocytic uptake of the influenza viruses	(Denisova et al., 2014)	
		Positively regulates adenovirus replication (mediated via viral proteins that mimic nutrient/growth signals regulating mTOR pathway)	(O'Shea et al., 2005)	
Ras/Raf/MEK/ERK	Ras	Positively regulates reovirus replication and spread	(Shmulevitz et al., 2010)	
	Ras-GTP	Positively regulates HBV replication and transcription (mediated via HBV transcriptional transactivating protein HBx)	(Benn and Schneider, 1994; Tang et al., 2005)	
	Cleavage of RasGAP	Positively regulates enterovirus replication	(Huber et al., 1999)	
	Raf	Positively regulates parvovirus transport across nuclear membrane as well as capsid assembly	(Riolobos et al., 2010)	
		Positively regulates synthesis and release of HIV-1	(Flory et al., 1998)	
	RKIP	Modulation of immune response to NDV	(Yin et al., 2015)	
	MEK	Facilitates IAV nuclear export	(Ludwig et al., 2004; Pleschka et al., 2001)	
			Positively regulates coronavirus genome synthesis	(Cai et al., 2007)
			Positively regulates BDV spread to neighboring cells	(Planz et al., 2001)
			Positively regulates replication of human neurotropic polyomavirus, JC	(Ravichandran et al., 2007)
	ERK1/2	Positively regulates RNA/protein synthesis in astroviruses	(Moser and Schultz-Cherry, 2008)	
		Positively regulates viral protein synthesis in alphaviruses	(Voss et al., 2014)	
		Positively regulates CVB3 replication	(Lim et al., 2005)	
		Positively regulates HCV genome synthesis	(Gretton et al., 2010)	
		Facilitates JUNV replication	(Rodriguez et al., 2014)	
		Positively regulates avian leukosis virus replication and virus-induced tumorigenesis	(Dai et al., 2016)	
Ras/p38	p38	Regulates RNA translation of encephalomyocarditis virus	(Hirasawa et al., 2003)	
Ras/JNK	JNK	viral protein and RNA synthesis of avian and human pandemic influenza A viruses	(Nacken et al., 2012)	
		Facilitates HSV-1 replication	(McLean and Bachenheimer, 1999)	
MNK1/eIF4E (cap-dependent translation)	Phosphorylation of eIF4E	Positively regulates HSV-1 genome and protein synthesis	(Walsh and Mohr, 2004)	
		Facilitates IAV protein synthesis	(Kleijn et al., 1996)	
MNK1/SRPK (cap-independent translation)	mTORC1-mediated hypophosphorylation of 4E-BP1	Facilitates CHIKV protein synthesis	(Joubert et al., 2015)	
	Inhibition of mTOR/AKT signaling	Facilitates HSV-1 replication and translation	(Chuluunbaatar et al., 2010)	
	PCB2-mediated recruitment of PIC	Positively regulates poliovirus replication	(Brown et al., 2014b)	
		Positively regulates poliovirus genome synthesis	(Chase et al., 2014)	
		Facilitates HCV replication through binding with 5'UTR	(Wang et al., 2011)	
		PCB2/SRp20 interaction is essential to effectively translate pirconavirus mRNA	(Bedard et al., 2007).	
	SRPK	Facilitates poliovirus cap-independent translation	(Brown et al., 2014a)	
	eIF4 G-MNK1 interaction	Negatively regulates adenovirus replication	(Cuesta et al., 2004)	
	Cleavage of eIF4 G	Facilitates IRES type 1 mediated enterovirus protein synthesis	(Thompson and Sarnow, 2003)	
PKR	eIF2 $\alpha$	Regulation of immune response against viral infections	(Bergmann et al., 2000; Weber et al., 2006)	
	PP2 A	Negatively regulates DNA synthesis in simian virus 40	(Cegielska et al., 1994)	

activated by a wide variety of viruses (Table 1). Depending on the nature of virus, MAPK signaling may either support or down regulate virus replication (Rodriguez et al., 2014). Moreover, MAPK signaling activated by virus independent sources can also be misused by the viruses to facilitate their own replication. For instance, phorbol-12-myristate-13-acetate (PMA) and new born calf serum activate MAPK; arenaviruses use this enhanced activity to promote its own replication (Rodriguez et al., 2014). Such widespread exploitation of MAPK pathway by viruses suggests that it may be used as a target to develop broad-spectrum antiviral drugs.

Depending on the nature of the virus involved, MAPK signaling may regulate single or multiple steps of virus replication (Andrade et al., 2004). For example, Ebola virus entry (Han and Harty, 2007), assembly and budding/release of HIV-1 (Hemonnot et al., 2004), replication/transcription of HCV (Pei et al., 2012), IAV protein synthesis (Gaur et al., 2011) and reactivation of KSHV latency (Jham and Montaner, 2010; Xie et al., 2008). Besides, it also regulates immune response (Gaur et al., 2011) and apoptosis (Bian et al., 2011) in virus infected cells.

Both live as well as inactivated virus can activate MAPK signaling pathway at a similar level (Rodriguez et al., 2014). Moreover, some viral secretory proteins can also lead to sustained activation of MAPK. For example, vaccinia virus growth factor (VGF) [secretary polypeptide from vaccinia virus homologous to epidermal growth factor (EGF) and transforming growth factor (TGF)] can effectively stimulate ERK1/2 (Andrade et al., 2004).

### 2.1. Cell surface receptor for MAPK signaling

Cell-surface receptors are specialized integral membrane proteins that constitute a major route by which signals are conveyed from extracellular to the intracellular environment of the cells. Cell surface receptors sense the extracellular signals by binding to their cognate ligands and are of three types, ligand-gated ion channel receptor (Leite and Cascio, 2001), enzyme-linked receptor and G protein-coupled receptor (GPCR) (Jordan and Devi, 1999). In MAPK signaling, receptor tyrosine kinase (RTKs, enzyme linked receptors) and GPCRs are the major classes of cell surface receptors (Widmann et al., 1999), though integrins may also activate MAPK signaling (Widmann et al., 1999). Besides MAPK signaling, RTKs and GPCRs may also be implicated in regulation of phosphoinositide 3-kinase (PI3K)-Akt signaling pathway (Rajagopalan, 2010).

#### 2.1.1. RTKs

RTKs are the high-affinity cell surface receptors for many growth factors, cytokines and hormones (Lemmon and Schlessinger, 2010). Besides being key regulator of normal cellular processes, RTKs also play a critical role in the development and progression of several types of cancer (Bennasroune et al., 2004). Mutations in RTKs lead to the dysregulation of a series of signaling cascades that may eventually develop disease syndromes. (Lemmon and Schlessinger, 2010). RTKs are member of the large family of proteins that contain a trans-membrane domain, an extracellular N terminal region, and an intracellular C terminal region. The cytoplasmic C terminal region of RTKs is highly conserved, encompassing kinase activity and catalyses receptor autophosphorylation including tyrosine phosphorylation of RTK substrates (Segaliny et al., 2015). Tyrosine kinase that does not possess any trans-membrane domains also exists and is called as non-receptor tyrosine kinase. Currently, over 20 different RTK classes have been identified viz; epidermal growth factor receptor (EGFR), insulin receptor family, platelet derived growth factor receptor (PDGFR), vascular endothelial growth factor receptor (VEGFR), fibroblast growth factor receptor (FGFR), cholecystokinin receptors (CCKR), nerve growth factor receptor (NGFR), hepatocyte growth factor receptor (HGFR), ephrin receptor, AXL receptors, tyrosine kinase with immunoglobulin-like and EGF-like domains 1 (TIE1), related to receptor tyrosine kinase receptor

(RYKR), discoidin domain receptor, 1 (DDR1), rearranged during transfection (RET), ROS receptor family, leukocyte receptor tyrosine kinase (LTK), receptor tyrosine kinase-like orphan receptor (ROR), muscle-specific kinase receptors (MuSK), lemur tyrosine kinase (LMR), and an undetermined class of RTKs (Segaliny et al., 2015).

Ligand binding induces receptor dimerization and autophosphorylation (Fig. 1). Each RTK has several auto-phosphorylation sites, which recruit different Src homology 2 domains (SH2) and phosphotyrosine binding domain (PTB) containing proteins that have intrinsic enzymatic activity. Binding of SH2 and PTB proteins with auto-phosphorylated intra-cytoplasmic domain of RTK leads to initiation of signal transduction pathways (Lemmon and Schlessinger, 2010). Besides SH2 and PTB, occasionally, adaptor proteins such as src homology 3 domain containing (SH3) and growth factor receptor-bound protein 2 (Grb2), which have no intrinsic enzymatic activity, can also initiate MAPK signaling by recruiting son of sevenless (SOS) protein (Gobert Gosse et al., 2005) (Fig. 1). Taken together, ligand binding to RTK induces RTK-Grb2-SOS-Ras/MAPK signaling (Lemmon and Schlessinger, 2010).

Besides MAPK, RTKs may also activate PI3K/Akt signaling (Cargnello and Roux, 2011; Cattaneo et al., 2014; Schlessinger, 2000) (Fig. 1). Src family of RTKs have been shown to support assembly and maturation of dengue virus (DENV) and West Nile virus (WNV) (Chu and Yang, 2007; Hirsch et al., 2005) and, entry of HIV-1 (Tokunaga et al., 1998) and HSV-1 (Qie et al., 1999). RTKs have also been shown to regulate multiple steps of influenza A virus (IAV) replication, which include RNA synthesis, Crm1-dependent nuclear export and, release of viral particles from infected cells through a pathway that is modulated by the lipid biosynthesis enzyme farnesyl diphosphate synthase (FPPS) (Kumar et al., 2011a).

#### 2.1.2. GPCRs

GPCRs lack intrinsic enzymatic (kinases) activity and are coupled to heterotrimeric G proteins, which consist of  $G\alpha$ ,  $G\beta$  and  $G\gamma$  subunits. Ligand binding induces conformational changes in GPCRs, during which the heterotrimeric G-proteins dissociate in GTP-bound  $G\alpha$  and  $G\beta\gamma$  subunits. These dissociated subunits of GPCRs regulate enzymatic activity of several key enzymes including second messengers such as adenylatecyclase, phospholipase C (PLC) which in turn regulate cellular functions by triggering different signaling pathways (Cattaneo et al., 2014; Neves et al., 2002). GPCRs have been shown to promote membrane fusion and cell-to-cell spread of human herpesvirus 6 (Zhen et al., 2005) as well as entry and RNA synthesis in flaviviruses (Le Sommer et al., 2012). Kaposi's sarcoma-associated herpesvirus (KSHV) oncogenes are associated with dysregulation of angiogenesis, which is mediated via GPCRs (Jham and Montaner, 2010).

### 2.2. Ras (transducer)

Upon activation, RTKs and GPCRs induce activation of its downstream signaling component-Rat sarcoma (Ras) which functions as GDP/GTP-regulated molecular switch (Biou and Cherfils, 2004). Ras is a 21-kDa protein belongs to the superfamily of small guanosine triphosphatases (GTPases) and is structurally related to  $G\alpha$  subunit of heterotrimeric G protein. In the inactive state, it binds to the nucleotide guanosine diphosphate (GDP) while in the activated state, binds to guanosine triphosphate (GTP). Ras interacts with and transmits signals to the downstream effector molecules (MAPKKKs/MAPKKs) associated with ERK/JNK/p38 pathway (Rajalingam et al., 2007). The process of exchanging the bound nucleotide is facilitated by guanine nucleotide exchange factors (GEFs) (Li et al., 2016) and GTPase activating proteins (GAPs) (Wennerberg et al., 2005).

Ras has an intrinsic GTPase activity thereby allowing spontaneous conversion of Ras-GTP to Ras-GDP to become inactive (Campbell et al., 1998; Wennerberg et al., 2005) (Fig. 1). Ras-GTP activates rapidly accelerated fibrosarcoma (Raf) which serves as first part of ERK/MAPK module. Ras GTPase-activating proteins (Ras-GAP) are the family of



regulatory proteins that bind to activated G proteins and stimulate their GTPase activity, eventually resulting in termination of the signaling event. By stimulating intrinsic Ras GTPase and consequent hydrolysis of GTP to GDP, Ras-GAP negatively regulates Ras-GTP (Bollag and McCormick, 1991; Trahey et al., 1988). Viral infections induce Ras-GAP cleavage which eventually triggers activation of Ras pathway. For example, Coxsackie virus B 3 (CVB3) infection leads to cleavage of Ras-GAP to produce Ras-GTP (activate state) which is required for efficient virus replication (Huber et al., 1999; Luo et al., 2002). Inhibition of MAPK/ERK kinase (MEK), which regulates cleavage of Ras-GAP, results in reduced CVB3 replication (Huber et al., 1999; Luo et al., 2002) (Table 1).

Ras activation depends on GEFs such as SOS and cell division cycle 25 (cdc25) phosphatase. Autophosphorylation of different intracytosolic domains of RTKs (Margolis and Skolnik, 1994) or conformational changes in GPCRs (Della Rocca et al., 1997) activate Ras by recruiting GEFs. Mutations in Ras may lead to persistent activation of Ras (Chappell et al., 2011; Li et al., 2016). Besides GEFs, some adaptor proteins also support MAPK activation. For example, growth factor receptor-bound protein 2 (Grb2), an adaptor protein, recruits SOS and triggers activation of MAPK and PI3K/Akt pathways (Fig. 1). MAPK activation results in transcription of genes involved in cell growth and differentiation (Li et al., 2016; Luo et al., 2002) whereas PI3K/Akt pathway regulates cytoskeleton integrity and apoptosis. Ras has four subtypes viz; Ha-Ras, N-Ras, KiRas 4A and KiRas 4B of which KiRas strongly activates MAPK pathway (Li et al., 2016; Yan et al., 1998) whereas HaRas more strongly activates PI3K/Akt pathway (Li et al., 2016).

### 2.3. Raf/MEK/ERK pathway

ERK pathway plays multiple pivotal roles in regulating cell fate (Cagnol and Chambard, 2010; Roux and Blenis, 2004; Shaul and Seger, 2007) and cell cycle progression, thereby serving as a reasonable target for the modulation of viral multiplication. To initiate ERK signaling, Raf interacts with MEK1/2 which in turn activates ERK.

#### 2.3.1. Raf (MAPKKK)

Raf (MAPKKK) serves as first part of ERK/MAPK module and holds a pivotal position in the MAPK signaling pathway (Hamden et al., 2005). Three kinases in Raf family viz; Raf-1, B-Raf, and A-Raf, are related with retroviral oncogenes (Roskoski, 2010). Among these, A-Raf serves as a main upstream activator of the ERK pathway (Fig. 1). Activation of Raf kinase involves interaction with Ras-GTPases (Bondeva et al., 2002; Roskoski, 2012). Besides MEK, Raf has a large number of other substrates such as JAK, PKA, ribosomal acidic P proteins (Rap), Src family kinase, Rac (also known as Ras-related C3 botulinum toxin substrate 1), PI3K, and phosphoinositide-dependent protein kinase-1 (PDK1) (Maroni et al., 2004; Zebisch and Troppmair, 2006).

Raf phosphorylates its downstream targets via MEK/ERK pathway (Cseh et al., 2014; Li et al., 2016). Among the Raf family members, only B-Raf (mutated) is able to persistently activate Raf/MEK/ERK pathway (Garnett and Marais, 2004). Mutated B-Raf can also activate MEK/ERK pathway as well as activating other Raf family members (Li et al., 2016; Rushworth et al., 2006). Furthermore, Ras, but not Raf has an intrinsic GTPase activity, thereby mutations at critical catalytic sites do not result in persistent Raf activation. Three Raf family members may induce MEK activation with varying biological effects in the order of B-Raf > Raf-1 > A-Raf (Alessi et al., 1994; Li et al., 2016). The Raf signaling was shown to regulate calcium/calmodulin-mediated budding of Ebola virus-like particle (VLP) (Han and Harty, 2007). Drugs targeting the Raf-MAPK signaling pathways represent a promising new class of antiviral agents to treat severe acute respiratory syndrome coronavirus (SARS-CoV) and poxvirus infections (Fauci and Challberg, 2005; Mizutani et al., 2004).

#### 2.3.2. MEK (MAPKK)

MAPKK (MEK1/2) are tyrosine and serine / threonine dual specificity protein kinases that are activated by Raf-mediated phosphorylation and are second component of ERK/MAPK module. MEK is the most characterized substrate for Raf (Hamden et al., 2005). The interaction between MAPKK and its downstream target MAPK is highly specific. Among the three types of MAPKs, p42/p44 (ERK) is phosphorylated solely by MEK 1 and 2, p38 MAPK is selectively activated by MKK3 and MKK6, while JNK is activated by MKK7 and MKK4. However, MKK4 may also activate p38 MAPK upon over expression (Maroni et al., 2004). Activated MEK has been shown to positively regulate borna disease virus (BDV) spread to neighbouring cells (Planz et al., 2001), RNA synthesis of the coronavirus (Cai et al., 2007) and nuclear export of influenza viruses (Ludwig et al., 2004; Pleschka et al., 2001).

#### 2.3.3. ERK

ERK is a key molecule in ERK/MAPK module that is solely activated by dual specificity MEK 1 and 2 (Plotnikov et al., 2011). In order to initiate ERK signaling, Raf interacts with MEK1/2 which in turn activates ERK. ERK1 and ERK2 are two isoforms with 83% amino acid homology and are expressed in diverse tissue types, which can be activated by a large number of extracellular and intracellular stimuli (Shi et al., 2013). Activated ERK can phosphorylate downstream kinases in the cytoplasm, at cell membrane and in the nucleus, thus diversifying the signaling cascade. ERK1/2 has more than 160 downstream target molecules (Li et al., 2016; McCubrey et al., 2007) that may phosphorylate different transcription factors, such as Ets-1, c-Jun, cMyc, CREB, NFκB, Elk1, c-Fos, c-Jun and STAT1 (Nakano et al., 1998; Shi et al., 2013; Steelman et al., 2004). NF-κB and STAT1 are rapidly activated in response to viral infections and cytokines, thus modulating the expression of genes involved in proliferation, apoptosis and immune response (Kumar et al., 2008; Levy et al., 2011). Some of the downstream targets such as ribosomal S6 kinase (RSK) family members are solely activated by ERK (Chambard et al., 2007; Roux and Blenis, 2004). RSK is shown to regulate KSHV lytic replication by promoting viral protein translation (Kuang et al., 2011).

Overexpression of the dominant negative mutants of Raf/ERK2 and MEK was shown to block influenza virus (Pleschka et al., 2001) and HCV replication respectively (Gretton et al., 2010). Likewise, overexpression of an active form of the Raf catalytic domain enhances parvovirus (Riolobos et al., 2010) and HIV-1 (Flory et al., 1998) replication. These evidences suggest that enhanced signaling activity may be misused by the viruses for their effective replication. For example, IAV interacts with ERK1/2 that allows effective trafficking of viral ribonucleoprotein (vRNP) (Pleschka et al., 2001), viral genome synthesis (Gaur et al., 2011) and induction of immune response, mediated via RANTES (regulated on activation, normal T cell expressed and secreted) (Kujime et al., 2000). Adenovirus type 7 activates Raf/MEK/ERK pathway that results in induction of IL-8 which plays an important role in viral pathogenesis (Alcorn et al., 2001). ERK2 is involved in regulating HIV-1 virion assembly and release by phosphorylating p6Gag (Hemonnot et al., 2004; Snyder et al., 2010). In HCV infection, MAPK/ERK pathway regulates viral RNA translation by directing the viral genome towards IRES (Pei et al., 2012). By regulating expression of lytic gene (switching viral latency from latent to lytic phase), MAPK pathway also facilitates Kaposi's sarcoma-associated herpesvirus (KSHV) replication (Xie et al., 2008).

Most common viral infections induce biphasic activation of ERK1/2 in the target cells. First peak of ERK1/2 activation occurs immediately following infection, suggesting that signaling is initiated by binding of the virus to its cognate receptors/co-receptor complex. For instance, engagement of HIV-1 p56lck and/or p59fyn with RTKs leads to ERK activation (Shenoy-Scaria et al., 1992). Only replication-competent (but not UV-irradiated/inactivated) viruses are capable of triggering late ERK activation thereby suggesting that high level of late-phase ERK activation is dependent on accumulation of viral gene/gene products;

the late phase ERK activation is associated with expression of HCV core protein (Shenoy-Scaria et al., 1992) and the HIV-1 Tat protein (Rusnati et al., 2001).

#### 2.4. JNK pathway

In mammals, JNKs are encoded by three distinct genes (*JNK1*, *JNK2* and *JNK3*). JNK1 and JNK2 are ubiquitously expressed, while JNK3 is limited to brain, heart and testis (Yan et al., 2010). During JNK activation, initial signaling pathway starts with the activation of several MAPKKs including TAK1, MEKK1/4, MLK2/3, and ASK1, which phosphorylate MEK4 or MEK7 and thus lead to activation of JNK (Krishna and Narang, 2008) (Fig. 1). JNK induces activation of transcription factors such as NF- $\kappa$ B p65, ELK-1, c-Fos, c-Jun and STAT1 which play a pivotal role in regulating the immune response to viral infections. In addition, activated JNK induces secretion of proinflammatory cytokines (Das and Muniyappa, 2010) such as TNF $\alpha$ , IFNs, IL1, IL2 and IL6 (Ting et al., 2010). JNK is also associated with regulation of apoptosis via modulation of c-Jun/AP1 and transforming growth factor- $\beta$  (TGF- $\beta$ ) (Xing et al., 2010). Although JNK is important for apoptosis, it also contributes to cell survival. These opposite effects rely on duration and magnitude of the pathway activation as well as concurrent activation of other signaling pathways. Prolonged activation of JNK mediates apoptosis whereas transient activation allows cell survival (Shi et al., 2013). Transient JNK activation is therefore essential for effective virus replication such as HSV-1 (McLean and Bachenheimer, 1999), DENV (Ceballos-Olvera et al., 2010), IAV (Ludwig et al., 2001) and varicella-zoster virus (VZV) (Rahaus et al., 2004).

#### 2.5. p38 MAPK pathway

p38 MAPK pathway modulates macrophage and neutrophil functions including chemotaxis, respiratory burst activity, granular exocytosis, adherence and apoptosis, as well as mediating T cell differentiation and apoptosis, by regulating IFN- $\gamma$  production (O'Sullivan et al., 2009). MKK3 and MKK6 serve as the major MAPKKs responsible for p38 activation (Derijard et al., 1995). MKK3/6 itself are activated by a number of MAPKKs, including MEKK1-3, MLK2/3, ASK1, Tpl2, TAK1, and TAO1/2 (Cuadrado and Nebreda, 2010) (Figs. 1 and 2). After activation, p38 phosphorylates a large number of substrates in both cytoplasm (cPLA2, MNK1/2, MK2/3, HuR, Bax, and Tau) and the nucleus (ATF1/2/6, MEF2, Elk-1, GADD153, Ets1, p53, and MSK1/2) (Cuadrado and Nebreda, 2010) (Fig. 1). These transcription factors regulate expression of RANTES, IL-8 and TNF- $\alpha$ , which are believed to regulate proinflammatory responses following viral infections (Medders and Kaul, 2011).

In conclusion, MAPK pathway positively regulates virus replication in diverse group of viruses (Table 1) except herpesvirus, NDV and polyomavirus where it has negative impact on virus replication cycle. In herpesviruses and polyomaviruses, MAPK/ERK function is performed by MAPK/p38/JNK pathway (Table 1).

### 3. MNK1

MAPK interacting kinase 1 (MNK1) is one of the MAPKs activated kinase encoded by *Mnk1* gene (O'Loughlen et al., 2004). It is expressed in all adult tissues except the brain (Waskiewicz et al., 1997) and is mainly activated by ERK1/2, although p38 can also activate it (Hargett et al., 2005; Roth et al., 2017). The downstream enzymatic activity of MNK1 depends on the upstream regulators (ERK1/2 and p38) which phosphorylate it at specific residues; these phosphorylation sites have been mapped using phospho peptide analysis of MNK1 by mass spectrometry (Cargnello and Roux, 2011). MNK1 has several potential downstream substrates, which include (i) eukaryotic initiation factor 4E (eIF4E) and eukaryotic initiation factor 4 G (eIF4 G) that helps in 5' m7G-dependent

mRNA translation and regulate cell growth and proliferation (Richter and Sonenberg, 2005). (ii) heterogeneous nuclear ribonucleoprotein A 1 (hnRNP A1) that facilitates processing, metabolism and transport of pre-mRNA. Besides, it also helps in production of TNF $\alpha$  and plays an important role in regulating inflammation (Locksley et al., 2001). (iii) p54 together with PSF [polypyrimidine tract binding (PTB)-associated splicing factor] form transcription splicing factor that is implicated in diverse biological functions such as alternative processing and translation of RNAs, as well as regulation of tissue-specific gene expression (Buxade et al., 2008; Valcarcel and Gebauer, 1997). (iv) Cytosolic phospholipase A2 (cPLA2) leads to production of eicosanoids (a second messenger) and plays an important role in immunity and inflammation (Hefner et al., 2000). (v) Sprouty 2 (Spry2) membrane associated protein negatively regulates ERK/MAPK signaling pathway (Cabrita and Christofori, 2008).


### 4. Role of MNK in virus replication

Viruses completely depend on the host translational machinery for synthesis of the viral proteins (Walsh and Mohr, 2004), therefore they have developed strategies to modulate host cell translation for optimizing their replication and spread. Viral mRNAs may have a range of structures at its 5' end. It may be capped with m7G or linked with viral genome-linked proteins (VPg). Uncapped viral mRNAs may carry cis-acting regulatory elements (CREs) such as internal ribosomal entry site (IRES) and protein binding sites. These 5' end structures have been shown to regulate viral mRNA translation (Brown et al., 2014a). MNK1 is involved in regulating both IRES- and cap-dependent viral mRNA translation.

#### 4.1. Role of MNK1 in initiation of cap-dependent viral mRNA translation

Most DNA viruses such as HSV-1, African swine fever virus (ASFV) (Castello et al., 2009), vaccinia virus (Walsh et al., 2008) as well as some RNA viruses such as chikungunya virus (CHIKV) (Walsh et al., 2013), SARS-CoV (Cencic et al., 2011), orthomyxoviruses (Aragon et al., 2000; Yanguz et al., 2012), rhabdoviruses (VSV, rabies virus) (Komarova et al., 2007), reoviruses (Chulu et al., 2010), hantavirus (Mir and Panganiban, 2008), and alphaviruses (Voss et al., 2014) contain a 5' cap in their mRNA. Some of the viruses such as influenza virus steal the 5' cap from the cellular mRNAs (Gu et al., 2015; Plotch et al., 1981) whereas several other viruses synthesize it by using host-encoded capping apparatus, for instance, vaccinia virus and reoviruses (Harwig et al., 2017). Viruses such as flaviviruses (Saeedi and Geiss, 2013) and nidoviruses synthesise the 5' cap by their own capping apparatus (Harwig et al., 2017). Calciviruses and picornaviruses contain VPg at its 5' end (Chaudhry et al., 2006; Goodfellow et al., 2005). These 5' end structures in viral genome mimic 5' m7G cap of the host mRNA, therefore viral RNA is translated like cellular mRNA. These viruses do not completely shutoff host protein synthesis. For initiation of 5' m7G cap-dependent viral mRNA translation, eIF4 F plays a central role in recruiting protein synthesis machinery to fully process capped mRNAs (Fig. 2). eIF4 F consists of a large adaptor protein eIF4 G that binds with eIF4E, a cap-binding protein eIF4 A, an RNA helicase (Svitkin et al., 2005), poly(A)-binding protein (PABA) that links 5' and 3' ends of properly spliced mRNA (Imataka et al., 1998) and eIF3 that assembles mRNA to the 40S ribosomal subunit.

Activated MNK1 may directly bind to eIF4 G in the initiation complex and phosphorylate eIF4E which subsequently binds to 5' cap of mRNA. In resting stage, eIF4E binding protein 1 (4E-BP1) remains associated with eIF4E. This dimer gets dissociated by phosphorylation of 4E-BP1 by metabolic sensor-mammalian target of rapamycin (mTORC1), which may be activated by either MNK1 or by ATP (representing high metabolic energy in the cell) (Waskiewicz et al., 1999). MNK1/eIF4E-mediated viral mRNA translation is known to take place in poxviruses, herpesvirus, alphaviruses, orthomyxoviruses, reoviruses



**Fig. 2. Role of MNK1 in viral mRNA translation.** MAPK pathway is activated by binding of the ligands to their cognate receptors (RTKs/GPCRs). With the help of adaptor proteins (SOS/Grb2), activated receptors convert Ras-GDP to Ras-GTP. Further information in the pathway is conveyed by phosphorylation (activation) of a series of enzymes, namely Raf, MEK1/2, and ERK1/2 which leads to activation of MNK1. Alternatively, MNK1 may be activated by p38. Besides regulating cell proliferation, cell survival and immune response, activated MNK1 also regulates viral mRNA translation. Activated MNK1 leads to dissociation of 4EBP1-eIF4E complex, which results in release (mediated via mTORC1) and subsequently phosphorylation of eIF4E. Activated eIF4E is required for assembly of eIF4 F to initiate cap-dependent viral mRNA translation. Alternatively, MNK1 may modulate SRPK function by directly inhibiting mTORC2-Akt signaling which is prerequisite for cell survival during viral infection. Activated SRPK phosphorylates SRPs that bind with IRES in the viral genome to initiate cap-independent viral mRNA translation. Activated MNK1 also facilitates pre-mRNA and post-translational processing, mediated via hnRNPA1 and PSF respectively.

and coronavirus (Table 1).

#### 4.2. MNK1 in initiation of cap-independent viral mRNA translation

Cap-independent translation is usually mediated by IRES which regulates interaction with eIFs and/or ribosomal proteins, and are able to initiate translation in the middle of a mRNA (Jackson, 2013). The mechanism of association of MNK1 with viral IRES-mediated translation is not completely understood. IRES-containing viruses can misuse activated MNK1 only when the classical function of MNK1 is being usurped, i. e. phosphorylation of eIF4E and its binding with eIF4 G to form eIF4 F complex, to initiation translation (Cuesta et al., 2000) (Fig. 2). Activated MNK1 is misused by the viruses by three distinct mechanisms, which include (i) Dephosphorylation of eIF4E as seen in poliovirus and EMCV (Kleijn et al., 1996). (ii) Cleavage of eIF4 G as seen in poliovirus and pestivirus (Kempf and Barton, 2008; Willcocks et al., 2011) and (iii) Inhibiting 4E-BP1 phosphorylation as seen among adenoviruses (Gingras and Sonenberg, 1997). These events, which are associated with suppressing classical MNK1 function, trigger a switch from cap-dependent to cap-independent translation (Gingras and Sonenberg, 1997), eventually preventing translation of cellular but not viral mRNA.

Four distinct classes of IRES are known: (i) Class I IRES directly cleave eIF4 G (Bordeleau et al., 2006; de Breyne et al., 2009) into two fragments. The larger fragment that contains binding sites for eIF4 A and eIF3 recruits 40s preinitiation complex (PIC) (required for

translation). For example, poliovirus 2A protease cleaves eIF4 G in order to selectively translate viral mRNA (Kempf and Barton, 2008; Willcocks et al., 2011). (ii) class II IRES induce conformational change in eIF4 G but do not result in its cleavage. The host protein shutoff is achieved by inhibiting 4E-BP1 phosphorylation concomitant with recruitment of 40s PIC, for example adenovirus (Gingras and Sonenberg, 1997) and HIV-1 (Borman et al., 2001). (iii) class III IRES utilize prokaryotic-like mode of translation initiation. By interacting with eIF3 and ribosomes, it facilitates positioning of ribosomes at mRNA (Babaylova et al., 2009; Berry et al., 2010; Locker et al., 2007), for example pestivirus, HCV, HIV-1 (Locker et al., 2011) and simian picornavirus (SPV9) (de Breyne et al., 2008). (iv) class IV IRES directly recruit 40S subunits without utilizing any eIFs, for example members of the family *Dicistroviridae* (Jan and Sarnow, 2002; Kamoshita et al., 2009; Wilson et al., 2000). Trans-acting host factors such as SRPKs, hNRNPA1 and PSF (MNK1 downstream substrates) interact with the CREs (positive-stranded RNA viruses and ds RNA viruses) to regulate viral RNA replication, subgenomic mRNA synthesis, translation and encapsidation. (Sullivan and Ahlquist, 1997; Pathak et al., 2012; Wang and Zhang, 1999).

MNK1 regulates IRES-containing mRNA translation by modulating SRPK1/2. However, SRPK1/2 is not a direct substrate for MNK1, its being regulated by MNK1-induced mTORC1-mediated inhibition of mTORC2-AKT signaling (Brown et al., 2014b) (Fig. 2). Ser/Arg (SR)-rich proteins (SRPs) that are prime substrates of SRPK1/2 (Brown et al., 2014b) constitute a large family of nuclear phosphoproteins that found



in both nucleus and cytoplasm (Sanford et al., 2004) and are associated with host-RNA-processing and alternative splicing. Upon direct interaction with viral mRNA, SRPs facilitates splicing of viral pre-mRNA in SINV, HIV-1, CMV (Fukuhara et al., 2006) and HCV (Karakama et al., 2010).

To recruit PIC in class I and class II IRES containing viruses, the SRPs are also believed to interact with the poly (rC) binding protein 2 (PCBP2) (Bedard et al., 2007). PCBP2 may alter IRES conformation to enhance eIF4 G binding, thereby facilitating ribosome recruitment (Blyn et al., 1996; Sweeney et al., 2014). PCBP2/SRP20 interaction is essential for picornavirus mRNA translation (Bedard et al., 2007). However, in IRES class III and class IV, SRPs do not seem to play any role because PIC assembly does not require eIF4 G.

MNK1 substrate hnRNPA1 has been shown to colocalize with viral nucleoprotein in the cytoplasm, particularly in the perinuclear region of virus-infected cells (Shi et al., 2000; Wang and Zhang, 1999) and is believed to regulate not only IRES-dependent translation initiation in enteroviruses (Tolbert et al., 2017), but also non-IRES-dependent translation initiation in SINV (Lin et al., 2009) and HCV (Kim et al., 2007; Li et al., 1997).

PSF, another MNK1 substrate has been shown to directly interact with viral mRNA in order to support posttranscriptional processing as well as translation of viral mRNA (Fig. 2). For example, PSF directly interacts with the cloverleaf CVB3 RNA as well as the IRES elements to regulate viral mRNA translation (Dave et al., 2017). In Hepatitis delta virus (HDV), PSF directly binds to the terminal stem loop domain of the viral RNA to regulate viral mRNA translation (Greco-Stewart et al., 2006). MNK1/SRPK mediated viral mRNA translation occurs in flaviviruses, picornaviruses, retroviruses, polyomaviruses and adenoviruses (Table 1).

#### 4.3. Role of MNK1 in viral genome synthesis

CREs in positive-sense RNA viruses participate in RNA/RNA and RNA/prtein interactions, thereby modulating viral RNA replication/transcription/translation as well as encapsidation (Sullivan and Ahlquist, 1997; Liu et al., 2009; Pathak et al., 2012; Wang and Zhang, 1999). MNK1 substrate hnRNPA1 colocalize with viral RNA in the cytoplasm and regulate genome synthesis of coronavirus mouse hepatitis virus (Shi et al., 2000, 2003). Besides MNK1, mTORC1 (which may be activated by both MNK1 and ATP) has also been shown to support synthesis of the viral genome (Waskiewicz et al., 1999). For example, in a study with HCV, activated mTORC1 was shown to support viral RNA synthesis (Stohr et al., 2016) without any direct involvement of MNK1.

#### 4.4. Role of MNK1 in cell survival during viral infection

mTOR is a metabolic sensor of cells with highly conserved serine/threonine residues (Hall, 2008) and is found in two structurally and functionally distinct multiprotein complexes, mTORC1 and mTORC2. mTORC1 is rapamycin sensitive and contains mTOR, regulatory associated protein of mTOR (RAPTOR) and mammalian lethal with SEC13 protein 8 (mLST8). (Hall, 2008). The high ATP/AMP ratio promotes and activates mTORC1, which leads to the dissociation of eIF4E from 4E-BP, ultimately resulting in enhanced protein synthesis (Brett et al., 2014), to support virus replication (Kuss-Duerkop et al., 2017; McNulty et al., 2013; Stohr et al., 2016). Conversely, antiviral role of mTORC1 has been demonstrated in CHIKV where virus has evolved a mechanism to bypass mTORC1 inhibition via MNK1/eIF4E mediated translation (Joubert et al., 2015) (Fig. 2). Antiviral function of mTORC1 in VSV is believed to be due to mTORC1-dependent type I IFN production, and initiation of autophagy (Alain et al., 2010).

mTORC2 is resistant to rapamycin and contains mTOR, rapamycin-insensitive companion of mammalian target of rapamycin (RICTOR), mammalian stress-activated protein kinase-interaction protein 1 (mSIN1), proline-rich protein 5 (PRR5) and mLST8 (Hall, 2008). Energy

depletion (low ATP levels) leads to adenosine monophosphate-activated protein kinase (AMPK)-mediated accumulation of AMP, which inhibits mTORC1 and activates mTORC2 (Brett et al., 2014). Upregulation of various apoptotic factors and inflammatory mediators during viral infection leads to cellular stress (reduced ATP level), which results in inactivation of mTORC1 and AMPK-mediated activation of mTORC2. Activated mTORC2 then promotes Akt phosphorylation (Vadlakonda et al., 2013) and MNK1 dephosphorylation. The Akt-mTORC2 signaling plays a critical role in survival of cells under stressful conditions such as: (i) activation of PKC $\alpha$  that maintains microtubule integrity (cytoskeletal dynamics) (ii) controlling ion transport and cellular growth via serine/threonine-protein kinase (SGK1) phosphorylation (iii) inhibiting PKR-mediated inflammatory response and upregulation of interferons (iv) interfering caspase protein and p53 mediated apoptosis and (iv) inhibition of MNK1 that shutoff both cap-dependent and cap-independent translation. This global inhibition of cellular protein synthesis aids in immune evasion (Ersahin et al., 2015; Porta et al., 2014). For example, influenza virus (Kuss-Duerkop et al., 2017), vaccinia virus and cowpox virus (Soares et al., 2009) activate mTORC2-Akt signaling which play a critical role in survival of cells at late stage of viral infection. However, in a study, DENV polyprotein was found to suppress only host cell translation at the level of translation initiation whereas viral protein synthesis remained unaffected. DENV infection triggered activation of p38/MNK1 signaling that eventually resulted in phosphorylation of eIF4E, which was sufficient to translate viral genome (Roth et al., 2017).

#### 4.5. MNK1 signaling in immune response to viral infections

Virus has evolved strategies to avoid a complete arrest in synthesis of cellular and viral proteins. Many of them directly modulate PKR activity while others inhibit phosphorylation of eIF2 $\alpha$ . PKR, a member of eIF2 $\alpha$  family of kinases contributes in cellular homeostasis by inhibiting translation of nonessential proteins. It acts as a pattern recognition receptor to sense dsRNA, a replicative intermediate formed in host cells during RNA virus replication. It has three sensor units viz; (i) general control nonrepressed 2 (GCN2) that is activated in response to amino-acid deprivation (ii) PKR-like endoplasmic reticulum kinase (PERK) that is activated in response to overloaded or misfolded proteins in the endoplasmic reticulum and (iii) heme-regulated eIF2 $\alpha$  kinase (HRI) that is activated in response to reduced heme levels as well as heat shock (Dang Do et al., 2009). Activated PKR phosphorylates  $\alpha$  subunit of eIF2- $\alpha$  as well as activating stress pathways- JNK, p38, NF $\kappa$ B (Fernandes, 2016) and protein phosphatase 2 A (PP2 A) (Ahn et al., 2007).

PKR is known to antagonize MNK1 (Castello et al., 2009) and AKT signaling (Pataer et al., 2009). Phosphorylated eIF2 $\alpha$  antagonize MNK1 functions by arresting both cellular and viral mRNA translation (Srivastava et al., 1998). For example, PKR mediated phosphorylation of eIF2 $\alpha$  is shown to inhibit translation of VSV (Connor and Lyles, 2005), rotavirus (Rojas et al., 2010) and alphavirus (Ventoso et al., 2006), besides inducing shutoff of the host cell translational machinery. PKR-mediated activation of NF $\kappa$ B (Dabo and Meurs, 2012), JNK and p38 leads to upregulation of various transcription factors including interferons and cytokines, thereby inducing an antiviral state (Taghavi and Samuel, 2012). PKR-mediated activation of tumor suppressor PP2 A leads to negative regulation of ERK1/2 (Yu et al., 2004) and AKT signaling (Ugi et al., 2004). Inhibition of ERK1/2 impedes MNK1 activation (Waskiewicz et al., 1997) whereas AKT inhibition may negatively affect cell survival (lack of maintenance of microtubule integrity and interference with p53 mediated apoptosis) (Ersahin et al., 2015; Porta et al., 2014), thereby preventing further viral spread and augmenting abortive viral replication. Besides PKR, SPRY2 (MNK1 substrate) also triggers interferon production. In addition, SPRY2 negatively regulates ERK-MNK1 signaling (Cabrita and Christofori, 2008). By upregulating TNF $\alpha$  (Locksley et al., 2001) and eicosanoids (Hefner et al., 2000),



**Table 2**  
Kinase inhibitors as antiviral agents.

Cellular Target	Inhibitor	Virus(es) involved*	References
Abl-family	STI-571 (Gleevec)	Poxvirus, variola virus and monkeypox virus	(Reeves et al., 2005, 2011)
EGFR	Gefitinib (Iressa)	Poxvirus and HCMV	(Herget et al., 2004; Langhammer et al., 2011)
NGFR	AG879	IAV, Sendai virus, HSV-1, MHV, and rotavirus	(Kumar et al., 2011a, b)
PDGFR	Tyrphostin A9 (A9)	IAV, Sendai virus, HSV-1, MHV, and rotavirus	(Kumar et al., 2011a, b)
Src family kinases	TG100572	HSV-1	(Sharma et al., 2011)
Raf (MAP3K)	Vemurafenib	IAV	(Holzberg et al., 2017)
MEK1/2 (MAP2K)	U0126	IAB, IAB, PEDV, Astrovirus, BDV, Coronavirus, JUNV and HSV-1	(Cai et al., 2007; Colao et al., 2017; Kim and Lee, 2015; Ludwig et al., 2004; Moser and Schultz-Cherry, 2008; Planz et al., 2001; Rodriguez et al., 2014)
ERK1/2 (MAPK)	Cl-1040 (PD184352)	IAB	(Haasbach et al., 2017)
	FR180204	DENV and Lentivirus	(Bukong et al., 2010; Sreekanth et al., 2014)
	Ag-126	VEEV	(Voss et al., 2014)
p38 (MAPK)	SB203580	EMCV and HSV-1	(Hirasawa et al., 2003; Walsh and Mohr, 2004)
	SB 202190	Influenza (H5N1, H7N7) virus, EMCV and Avian reovirus	(Borgeling et al., 2014; Chulu et al., 2010; Hirasawa et al., 2003)
JNK (MAPK)	AS601245	IAB	(Nacken et al., 2012)
	SP600125	IAB and HCMV	(Nacken et al., 2012; Zhang et al., 2015)
MNK1	CGP57380	HSV-1, Poxvirus and HCMV	(Walsh et al., 2008; Walsh and Mohr, 2004; Walsh et al., 2005)
eIF4E/eIF4 G	4E2RCat	Coronavirus	(Cencic et al., 2011)
mTORC1	Rapamycin	HCV	(Stohr et al., 2016)
Akt	MK2206	IAB	(Denisova et al., 2014)

\*Abbreviations: IAV: Influenza A virus, DENV: dengue virus, EMCV: Encephalomyocarditis Virus, HCV: Hepatitis C virus, HCMV: Human cytomegalovirus, HSV: Herpes simplex virus, MHV: Mouse hepatitis virus, PEDV: Porcine epidemic diarrhoea virus, JUNV: Junin virus, VEEV: Venezuelan equine encephalitis virus.

MNK1 substrates- hnRNPA1 and cPLA2 also play an important role in immunity and inflammation.

## 5. Genome-wide siRNA screens to identify cellular factors required for virus replication

Gene silencing through RNAi includes sequence-specific targeting of mRNAs by delivering 20–25 nucleotide long dsRNA homologous to gene concerned. RNAi has enabled systematic exploration of host genes involved in virus replication and virulence, thereby allowing identification of novel drug targets (Radoshitzky et al., 2016). Since the last decade, a large number of genome-scale RNAi screens have been conducted, which identified cellular factors required for WNV (Krishnan et al., 2008), DENV (Morchang et al., 2017; Sessions et al., 2009), HIV-1 (Zhou et al., 2008), HCV (Li et al., 2009; Lupberger et al., 2015; Tai et al., 2009), IAV (Han et al., 2018; Karlas et al., 2010), JUNV (Lavanya et al., 2013), coronavirus (Staff, 2015), poxvirus (Kilcher et al., 2014), herpesvirus (Griffiths et al., 2013) and polyomavirus (Zhao and Imperiale, 2017) replication. siRNA screens reveal both proviral and antiviral cellular factors and therefore help in understanding the precise nature of virus-host interactions (Watanabe et al., 2010).

Using reductionist approach, only few of the host factors were known to be implicated in virus replication. However, genome-wide siRNA screens enabled to identify almost every potential host candidate gene that may regulate virus replication. Each virus requires over a thousand host proteins to effectively replicate inside the host. Each siRNA screen identified multiple MAPK components that influence virus replication. Nevertheless, genomewide siRNA screens conducted on Zika Virus (ZIKV), DENV (Savidis et al., 2016), HCV (Lupberger et al., 2015; Tai et al., 2009), Ebola Virus (Kolokoltsov et al., 2009) and SARS-CoV (de Wilde et al., 2015) have identified MNK1 as one of the cellular factor that regulates virus replication. MNK1 has been identified as a proviral factor in all the siRNA screens conducted so far. However, MNK2 has both proviral and antiviral (de Wilde et al., 2015) functions, depending on the virus prototypes involved. More detailed functional analyses of these host genes (identified in genome-wide screens) will provide insights for development of novel antiviral therapeutics.

## 6. Kinase inhibitors as antiviral agents

In 2002, kinome-the protein kinase complement of the human genome, identified 518 protein kinase genes. Protein kinases play a pivotal role in cellular transduction signaling. Their malfunction, hyperactivity or overexpression is usually associated with disease. Kinase is one of the most intensively studied classes of drug targets (Zhang et al., 2009). Understanding the structural basis of kinase is essential for developing selective inhibitors to target most possible member of the kinome (Cao et al., 2010). Over 80 kinase inhibitors have been evaluated to some stage of clinical trial and over 30 distinct kinase inhibitors have been developed to the level of a Phase I clinical trial (Zhang et al., 2009). The vast majority of these inhibitors were investigated for the treatment of cancer (Player, 2009). However, dysregulation of kinase function has been implicated in several other disorders, including metabolic, immunological, neurological and infectious disease. Most of the known kinase inhibitors target at the kinase activation loop in the ATP binding site (Zhang et al., 2009).

Besides genome-wide siRNA screens, screening kinase inhibitor libraries has also enabled identification of kinases required for efficient virus replication (Kumar et al., 2011a). These inhibitor libraries identified both virus-specific and broad-spectrum antiviral agents (Kumar et al., 2011b). While some of the inhibitors were shown to impair single step of virus life cycle (Kumar et al., 2008), others block multiple steps of virus replication cycle (Kumar et al., 2011a). Kinase inhibitors that target MAPK pathway and may have potential to act as antiviral agents are summarized in Table 2. More comprehensive information about the kinase inhibitors could be accessed elsewhere in several excellent reviews (Bogoyevitch et al., 2004; Burkhard and Shapiro, 2010; Gangwal et al., 2013; Genovese, 2009; Janne et al., 2013; Kim and Sim, 2012; Levitzki and Gazit, 1995; Matsuda and Fukumoto, 2011; Michelangeli and East, 2011; Sweeney and Firestein, 2006; Uehling and Harris, 2015).

## 7. Concluding remarks

Classically, antiviral agents are developed by targeting certain viral components. Viral genome is highly unstable and undergoes frequent mutations; under selection pressure of a drug, virus quickly acquires

drug resistance at druggable sites. After the advent of high throughput genome sequencing and genome-wide siRNA screens, thousands of cellular factors that support virus replication have been identified. Since it's not quite easy for the virus to replace the missing cellular functions by mutations, host-directed antiviral therapies usually do not induce generation of drug-resistant virus variants. In this review, we have described numerous cellular factors/cell signaling pathways which are exploited by the viruses for their effective replication. These cellular factors may serve as potential drug targets to develop novel antiviral therapeutics. However, their further validation, in vivo efficacy and clinical trials are essential before actually translating them from research into the clinical settings.

## Acknowledgments

Science and Engineering Research Board (India) supported this work [grant number SB/SO/AS-20/2014]. The funding agency has no role in design, data collection and interpretation, or the decision to submit this work for publication.

## References

- Ahn, J.H., McAvoy, T., Rakhilin, S.V., Nishi, A., Greengard, P., Nairn, A.C., 2007. Protein kinase A activates protein phosphatase 2A by phosphorylation of the B56delta subunit. *Proc. Natl. Acad. Sci. U. S. A.* 104 (8), 2979–2984.
- Alain, T., Lun, X., Martineau, Y., Sean, P., Pulendran, B., Petroulakis, E., Zemp, F.J., Lemay, C.G., Roy, D., Bell, J.C., Thomas, G., Kozma, S.C., Forsyth, P.A., Costa-Mattioli, M., Sonenberg, N., 2010. Vesicular stomatitis virus oncolysis is potentiated by impairing mTORC1-dependent type I IFN production. *Proc. Natl. Acad. Sci. U. S. A.* 107 (4), 1576–1581.
- Alcorn, M.J., Booth, J.L., Coggeshall, K.M., Metcalf, J.P., 2001. Adenovirus type 7 induces interleukin-8 production via activation of extracellular regulated kinase 1/2. *J. Virol.* 75 (14), 6450–6459.
- Alessi, D.R., Saito, Y., Campbell, D.G., Cohen, P., Sithanandam, G., Rapp, U., Ashworth, A., Marshall, C.J., Cowley, S., 1994. Identification of the sites in MAP kinase kinase-1 phosphorylated by p74raf-1. *EMBO J.* 13 (7), 1610–1619.
- Andrade, A.A., Silva, P.N., Pereira, A.C., De Sousa, L.P., Ferreira, P.C., Gazzinelli, R.T., Kroon, E.G., Ropert, C., Bonjardim, C.A., 2004. The vaccinia virus-stimulated mitogen-activated protein kinase (MAPK) pathway is required for virus multiplication. *Biochem. J.* 381 (Pt 2), 437–446.
- Aragon, T., de la Luna, S., Novoa, I., Carrasco, L., Ortin, J., Nieto, A., 2000. Eukaryotic translation initiation factor 4G1 is a cellular target for NS1 protein, a translational activator of influenza virus. *Mol. Cell. Biol.* 20 (17), 6259–6268.
- Babaylova, E., Graifer, D., Malygin, A., Stahl, J., Shatsky, I., Karpova, G., 2009. Positioning of subdomain IIIId and apical loop of domain II of the hepatitis C IRES on the human 40S ribosome. *Nucleic Acids Res.* 37 (4), 1141–1151.
- Bedard, K.M., Daijogo, S., Semler, B.L., 2007. A nucleocytoplasmic SR protein functions in viral IRES-mediated translation initiation. *EMBO J.* 26 (2), 459–467.
- Benn, J., Schneider, R.J., 1994. Hepatitis B virus HBx protein activates Ras-GTP complex formation and establishes a Ras, Raf, MAP kinase signaling cascade. *Proc. Natl. Acad. Sci. U. S. A.* 91 (22), 10350–10354.
- Bennasroune, A., Gardin, A., Aunis, D., Cremel, G., Hubert, P., 2004. Tyrosine kinase receptors as attractive targets of cancer therapy. *Crit. Rev. Oncol. Hematol.* 50 (1), 23–38.
- Bergmann, M., Garcia-Sastre, A., Carnero, E., Pehamberger, H., Wolff, K., Palese, P., Muster, T., 2000. Influenza virus NS1 protein counteracts PKR-mediated inhibition of replication. *J. Virol.* 74 (13), 6203–6206.
- Berry, K.E., Waghray, S., Doudna, J.A., 2010. The HCV IRES pseudoknot positions the initiation codon on the 40S ribosomal subunit. *RNA* 16 (8), 1559–1569.
- Bian, J., Wang, K., Kong, X., Liu, H., Chen, F., Hu, M., Zhang, X., Jiao, X., Ge, B., Wu, Y., Meng, S., 2011. Caspase- and p38-MAPK-dependent induction of apoptosis in A549 lung cancer cells by Newcastle disease virus. *Arch. Virol.* 156 (8), 1335–1344.
- Biou, V., Cherfils, J., 2004. Structural principles for the multispecificity of small GTP-binding proteins. *Biochemistry* 43 (22), 6833–6840.
- Blyn, L.B., Swiderek, K.M., Richards, O., Stahl, D.C., Semler, B.L., Ehrenfeld, E., 1996. Poly(rC) binding protein 2 binds to stem-loop IV of the poliovirus RNA 5' noncoding region: identification by automated liquid chromatography-tandem mass spectrometry. *Proc. Natl. Acad. Sci. U. S. A.* 93 (20), 11115–11120.
- Bogoyevitch, M.A., Boehm, I., Oakley, A., Ketterman, A.J., Barr, R.K., 2004. Targeting the JNK MAPK cascade for inhibition: basic science and therapeutic potential. *Biochim. Biophys. Acta* 1697 (1–2), 89–101.
- Bollag, G., McCormick, F., 1991. Differential regulation of rasGAP and neurofibromatosis gene product activities. *Nature* 351 (6327), 576–579.
- Bondeva, T., Balla, A., Varnai, P., Balla, T., 2002. Structural determinants of Ras-Raf interaction analyzed in live cells. *Mol. Biol. Cell.* 13 (7), 2323–2333.
- Bordeleau, M.E., Mori, A., Oberer, M., Lindqvist, L., Chard, L.S., Higa, T., Belsham, G.J., Wagner, G., Tanaka, J., Pelletier, J., 2006. Functional characterization of IRESes by an inhibitor of the RNA helicase eIF4A. *Nat. Chem. Biol.* 2 (4), 213–220.
- Borgeling, Y., Schmolke, M., Viemann, D., Nordhoff, C., Roth, J., Ludwig, S., 2014. Inhibition of p38 mitogen-activated protein kinase impairs influenza virus-induced primary and secondary host gene responses and protects mice from lethal H5N1 infection. *J. Biol. Chem.* 289 (1), 13–27.
- Borman, A.M., Michel, Y.M., Kean, K.M., 2001. Detailed analysis of the requirements of hepatitis A virus internal ribosome entry segment for the eukaryotic initiation factor complex eIF4F. *J. Virol.* 75 (17), 7864–7871.
- Boulton, T.G., Yancopoulos, G.D., Gregory, J.S., Slaughter, C., Moomaw, C., Hsu, J., Cobb, M.H., 1990. An insulin-stimulated protein kinase similar to yeast kinases involved in cell cycle control. *Science* 249 (4964), 64–67.
- Brett, K.E., Ferraro, Z.M., Yockell-Lelievre, J., Gruslin, A., Adamo, K.B., 2014. Maternal-fetal nutrient transport in pregnancy pathologies: the role of the placenta. *Int. J. Mol. Sci.* 15 (9), 16153–16185.
- Brown, M.C., Bryant, J.D., Dobrikova, E.Y., Shveygert, M., Bradrick, S.S., Chandramohan, V., Bigner, D.D., Gromeier, M., 2014a. Induction of viral, 7-methyl-guanosine cap-independent translation and oncolysis by mitogen-activated protein kinase-interacting kinase-mediated effects on the serine/arginine-rich protein kinase. *J. Virol.* 88 (22), 13135–13148.
- Brown, M.C., Dobrikov, M.I., Gromeier, M., 2014b. Mitogen-activated protein kinase-interacting kinase regulates mTOR/AKT signaling and controls the serine/arginine-rich protein kinase-responsive type 1 internal ribosome entry site-mediated translation and viral oncolysis. *J. Virol.* 88 (22), 13149–13160.
- Brucher, B.L., Jamall, I.S., 2014. Cell-cell communication in the tumor microenvironment, carcinogenesis, and anticancer treatment. *Cell. Physiol. Biochem.* 34 (2), 213–243.
- Bukong, T.N., Hall, W.W., Jacque, J.M., 2010. Lentivirus-associated MAPK/ERK2 phosphorylates EMD and regulates infectivity. *J. Gen. Virol.* 91 (Pt 9), 2381–2392.
- Burkhard, K., Shapiro, P., 2010. Use of inhibitors in the study of MAP kinases. *Methods Mol. Biol.* 661, 107–122.
- Buxade, M., Morrice, N., Krebs, D.L., Proud, C.G., 2008. The PSF/p54nrb complex is a novel Mnk substrate that binds the mRNA for tumor necrosis factor alpha. *J. Biol. Chem.* 283 (1), 57–65.
- Cabrita, M.A., Cristofori, G., 2008. Sprouty proteins, masterminds of receptor tyrosine kinase signaling. *Angiogenesis* 11 (1), 53–62.
- Cagnol, S., Chambard, J.C., 2010. ERK and cell death: mechanisms of ERK-induced cell death—apoptosis, autophagy and senescence. *FEBS J.* 277 (1), 2–21.
- Cai, Y., Liu, Y., Zhang, X., 2007. Suppression of coronavirus replication by inhibition of the MEK signaling pathway. *J. Virol.* 81 (2), 446–456.
- Campbell, S.L., Khosravi-Far, R., Rossman, K.L., Clark, G.J., Der, C.J., 1998. Increasing complexity of Ras signaling. *Oncogene* 17 (11 Reviews), 1395–1413.
- Cao, R., Mi, N., Zhang, H., 2010. 3D-QSAR study of c-Src kinase inhibitors based on docking. *J. Mol. Model.* 16 (2), 361–375.
- Cao, W., Vyboh, K., Routy, B., Chababi-Atallah, M., Lemire, B., Routy, J.P., 2015. Imatinib for highly chemoresistant Kaposi sarcoma in a patient with long-term HIV control: a case report and literature review. *Curr. Oncol.* 22 (5), e395–399.
- Cargnello, M., Roux, P.P., 2011. Activation and function of the MAPKs and their substrates, the MAPK-activated protein kinases. *Microbiol. Mol. Biol. Rev.* 75 (1), 50–83.
- Castello, A., Quintas, A., Sanchez, E.G., Sabina, P., Nogal, M., Carrasco, L., Revilla, Y., 2009. Regulation of host translational machinery by African swine fever virus. *PLoS Pathog.* 5 (8), e1000562.
- Cattaneo, F., Guerra, G., Parisi, M., De Marinis, M., Tafuri, D., Cinelli, M., Ammendola, R., 2014. Cell-surface receptors transactivation mediated by G protein-coupled receptors. *Int. J. Mol. Sci.* 15 (11), 19700–19728.
- Ceballos-Olvera, I., Chavez-Salinas, S., Medina, F., Ludert, J.E., del Angel, R.M., 2010. JNK phosphorylation, induced during dengue virus infection, is important for viral infection and requires the presence of cholesterol. *Virology* 396 (1), 30–36.
- Cegielska, A., Shaffer, S., Derua, R., Goris, J., Virshup, D.M., 1994. Different oligomeric forms of protein phosphatase 2A activate and inhibit simian virus 40 DNA replication. *Mol. Cell. Biol.* 14 (7), 4616–4623.
- Cencic, R., Desforges, M., Hall, D.R., Kozakov, D., Du, Y., Min, J., Dingleline, R., Fu, H., Vajda, S., Talbot, P.J., Pelletier, J., 2011. Blocking eIF4E-eIF4G interaction as a strategy to impair coronavirus replication. *J. Virol.* 85 (13), 6381–6389.
- Chambard, J.C., Lefloch, R., Pouyssegur, J., Lenormand, P., 2007. ERK implication in cell cycle regulation. *Biochim. Biophys. Acta* 1773 (8), 1299–1310.
- Chappell, W.H., Steelman, L.S., Long, J.M., Kempf, R.C., Abrams, S.L., Franklin, R.A., Basecke, J., Stivala, F., Donia, M., Fagone, P., Malaponte, G., Mazzarino, M.C., Nicoletti, F., Libra, M., Maksimovic-Ivanic, D., Mijatovic, S., Montalto, G., Cervello, M., Laidler, P., Milella, M., Tafuri, A., Bonati, A., Evangelisti, C., Cocco, L., Martelli, A.M., McCubrey, J.A., 2011. Ras/Raf/MEK/ERK and PI3K/PTEN/Akt/mTOR inhibitors: rationale and importance to inhibiting these pathways in human health. *Oncotarget* 2 (3), 135–164.
- Chase, A.J., Daijogo, S., Semler, B.L., 2014. Inhibition of poliovirus-induced cleavage of cellular protein PCBP2 reduces the levels of viral RNA replication. *J. Virol.* 88 (6), 3192–3201.
- Chaudhry, Y., Nayak, A., Bordeleau, M.E., Tanaka, J., Pelletier, J., Belsham, G.J., Roberts, L.O., Goodfellow, I.G., 2006. Calciviruses differ in their functional requirements for eIF4F components. *J. Biol. Chem.* 281 (35), 25315–25325.
- Chu, J.J., Yang, P.L., 2007. C-src protein kinase inhibitors block assembly and maturation of dengue virus. *Proc. Natl. Acad. Sci. U. S. A.* 104 (9), 3520–3525.
- Chulu, J.L., Huang, W.R., Wang, L., Shih, W.S., Liu, H.J., 2010. Avian reovirus non-structural protein p17-induced G(2)/M cell cycle arrest and host cellular protein translation shutoff involve activation of p53-dependent pathways. *J. Virol.* 84 (15), 7683–7694.
- Chuluunbaatar, U., Roller, R., Feldman, M.E., Brown, S., Shokat, K.M., Mohr, I., 2010. Constitutive mTORC1 activation by a herpesvirus Akt surrogate stimulates mRNA translation and viral replication. *Genes Dev.* 24 (23), 2627–2639.
- Colao, I., Pennisi, R., Venuti, A., Nygardas, M., Heikkila, O., Hukkanen, V., Sciortino,

- M.T., 2017. The ERK-1 function is required for HSV-1-mediated G1/S progression in HEP-2 cells and contributes to virus growth. *Sci. Rep.* 7 (1), 9176.
- Connor, J.H., Lyles, D.S., 2005. Inhibition of host and viral translation during vesicular stomatitis virus infection. eIF2 is responsible for the inhibition of viral but not host translation. *J. Biol. Chem.* 280 (14), 13512–13519.
- Cseh, B., Doma, E., Baccarini, M., 2014. RAF<sup>W</sup> neighborhood: protein-protein interaction in the Raf/Mek/Erk pathway. *FEBS Lett.* 588 (15), 2398–2406.
- Cuadrado, A., Nebreda, A.R., 2010. Mechanisms and functions of p38 MAPK signalling. *Biochem. J.* 429 (3), 403–417.
- Cuesta, R., Xi, Q., Schneider, R.J., 2000. Adenovirus-specific translation by displacement of kinase Mnk1 from cap-initiation complex eIF4F. *EMBO J.* 19 (13), 3465–3474.
- Cuesta, R., Xi, Q., Schneider, R.J., 2004. Structural basis for competitive inhibition of eIF4G-Mnk1 interaction by the adenovirus 100-kilodalton protein. *J. Virol.* 78 (14), 7707–7716.
- Dabo, S., Meurs, E.F., 2012. dsRNA-dependent protein kinase PKR and its role in stress, signaling and HCV infection. *Viruses* 4 (11), 2598–2635.
- Dai, M., Feng, M., Ye, Y., Wu, X., Liu, D., Liao, M., Cao, W., 2016. Exogenous avian leukosis virus-induced activation of the ERK/AP1 pathway is required for virus replication and correlates with virus-induced tumorigenesis. *Sci. Rep.* 6, 19226.
- Dang Do, A.N., Kimball, S.R., Cavener, D.R., Jefferson, L.S., 2009. eIF2alpha kinases GCN2 and PERK modulate transcription and translation of distinct sets of mRNAs in mouse liver. *Physiol. Genom.* 38 (3), 328–341.
- Das, K.C., Muniyappa, H., 2010. c-Jun-NH2 terminal kinase (JNK)-mediates AP-1 activation by thioredoxin: phosphorylation of cJun, JunB, and Fra-1. *Mol. Cell. Biochem.* 337 (1–2), 53–63.
- Dave, P., George, B., Sharma, D.K., Das, S., 2017. Polypyrimidine tract-binding protein (PTB) and PTB-associated splicing factor in CVB3 infection: an ITAF for an ITAF. *Nucleic Acids Res.* 45 (15), 9068–9084.
- de Brejne, S., Yu, Y., Pestova, T.V., Hellen, C.U., 2008. Factor requirements for translation initiation on the Simian picornavirus internal ribosomal entry site. *RNA* 14 (2), 367–380.
- de Brejne, S., Yu, Y., Unbehaun, A., Pestova, T.V., Hellen, C.U., 2009. Direct functional interaction of initiation factor eIF4G with type 1 internal ribosomal entry sites. *Proc. Natl. Acad. Sci. U. S. A.* 106 (23), 9197–9202.
- de Wilde, A.H., Wansee, K.F., Scholte, F.E., Goeman, J.J., Ten Dijke, P., Snijder, E.J., Kikkert, M., van Hemert, M.J., 2015. A kinome-wide small interfering RNA screen identifies proviral and antiviral host factors in severe acute respiratory syndrome coronavirus replication, including double-stranded RNA-activated protein kinase and early secretory pathway proteins. *J. Virol.* 89 (16), 8318–8333.
- Della Rocca, G.J., van Biesen, T., Daaka, Y., Luttrell, D.K., Luttrell, L.M., Lefkowitz, R.J., 1997. Ras-dependent mitogen-activated protein kinase activation by G protein-coupled receptors. Convergence of Gi- and Gq-mediated pathways on calcium/calmodulin, Pyk2, and Src kinase. *J. Biol. Chem.* 272 (31), 19125–19132.
- Denef, C., 2008. Paracrinicity: the story of 30 years of cellular pituitary crosstalk. *J. Neuroendocrinol.* 20 (1), 1–70.
- Denisova, O.V., Soderholm, S., Virtanen, S., Von Schantz, C., Bychkov, D., Vashchinkina, E., Desloovere, J., Tynell, J., Ikonen, N., Theisen, L.L., Nyman, T.A., Matikainen, S., Kallioniemi, O., Julkunen, I., Muller, C.P., Saelens, X., Verkhusha, V.V., Kainov, D.E., 2014. Akt inhibitor MK2206 prevents influenza pH1N1 virus infection in vitro. *Antimicrob. Agents Chemother.* 58 (7), 3689–3696.
- Derijard, B., Raingeaud, J., Barrett, T., Wu, I.H., Han, J., Ulevitch, R.J., Davis, R.J., 1995. Independent human MAP-kinase signal transduction pathways defined by MEK and MKK isoforms. *Science* 267 (5198), 682–685.
- Ersahin, T., Tuncbag, N., Cetin-Atalay, R., 2015. The PI3K/AKT/mTOR interactive pathway. *Mol. Biosyst.* 11 (7), 1946–1954.
- Fauci, A.S., Challberg, M.D., 2005. Host-based antiviral therapeutic strategies: turning the tables. *J. Clin. Invest.* 115 (2), 231–233.
- Fernandes, J., 2016. Oncogenes: the passport for viral oncolysis through PKR inhibition. *Biomark. Cancer* 8, 101–110.
- Flory, E., Weber, C.K., Chen, P., Hoffmeyer, A., Jassoy, C., Rapp, U.R., 1998. Plasma membrane-targeted Raf kinase activates NF-kappaB and human immunodeficiency virus type 1 replication in T lymphocytes. *J. Virol.* 72 (4), 2788–2794.
- Fujioka, Y., Tsuda, M., Hattori, T., Sasaki, J., Sasaki, T., Miyazaki, T., Ohba, Y., 2011. The Ras-PI3K signaling pathway is involved in clathrin-independent endocytosis and the internalization of influenza viruses. *PLoS One* 6 (1), e16324.
- Fukuhara, T., Hosoya, T., Shimizu, S., Sumi, K., Oshiro, T., Yoshinaka, Y., Suzuki, M., Yamamoto, N., Herzenberg, L.A., Herzenberg, L.A., Hagiwara, M., 2006. Utilization of host SR protein kinases and RNA-splicing machinery during viral replication. *Proc. Natl. Acad. Sci. U. S. A.* 103 (30), 11329–11333.
- Gangwal, R.P., Bhadauriya, A., Damre, M.V., Dhoke, G.V., Sangamwar, A.T., 2013. p38 Mitogen-activated protein kinase inhibitors: a review on pharmacophore mapping and QSAR studies. *Curr. Top. Med. Chem.* 13 (9), 1015–1035.
- Garnett, M.J., Marais, R., 2004. Guilty as charged: B-RAF is a human oncogene. *Cancer Cell* 6 (4), 313–319.
- Gaur, P., Munjal, A., Lal, S.K., 2011. Influenza virus and cell signaling pathways. *Med. Sci. Monit.* 17 (6), RA148–154.
- Genovese, M.C., 2009. Inhibition of p38: has the fat lady sung? *Arthritis Rheum* 60 (2), 317–320.
- Gingras, A.C., Sonenberg, N., 1997. Adenovirus infection inactivates the translational inhibitors 4E-BP1 and 4E-BP2. *Virology* 237 (1), 182–186.
- Gobert Gosse, S., Bourgin, C., Liu, W.Q., Garbay, C., Mouchiroud, G., 2005. M-CSF stimulated differentiation requires persistent MEK activity and MAPK phosphorylation independent of Grb2-Sos association and phosphatidylinositol 3-kinase activity. *Cell. Signal.* 17 (11), 1352–1362.
- Goodfellow, I., Chaudhry, Y., Giodasi, I., Gerondopoulos, A., Natoni, A., Labrie, L., Laliberté, J.F., Roberts, L., 2005. Calicivirus translation initiation requires an interaction between VPg and eIF 4 E. *EMBO Rep.* 6 (10), 968–972.
- Greco-Stewart, V.S., Thibault, C.S., Pelchat, M., 2006. Binding of the polypyrimidine tract-binding protein-associated splicing factor (PSF) to the hepatitis delta virus RNA. *Virology* 356 (1–2), 35–44.
- Gretton, S., Hughes, M., Harris, M., 2010. Hepatitis C virus RNA replication is regulated by Ras-Erk signalling. *J. Gen. Virol.* 91 (Pt 3), 671–680.
- Griffiths, S.J., Koegl, M., Boutell, C., Zenner, H.L., Crump, C.M., Pica, F., Gonzalez, O., Friedel, G.C., Barry, G., Martin, K., Craigon, M.H., Chen, R., Kaza, L.N., Fossum, E., Fazakerley, J.K., Efstathiou, S., Volpi, A., Zimmer, R., Ghazal, P., Haas, J., 2013. A systematic analysis of host factors reveals a Med23-interferon-lambda regulatory axis against herpes simplex virus type 1 replication. *PLoS Pathog* 9 (8), e1003514.
- Gu, W., Gallagher, G.R., Dai, W., Liu, P., Li, R., Trombly, M.I., Gammon, D.B., Mello, C.C., Wang, J.P., Finberg, R.W., 2015. Influenza A virus preferentially snatches noncoding RNA caps. *RNA* 21 (12), 2067–2075.
- Haasbach, E., Muller, C., Ehrhardt, C., Schreiber, A., Pleschka, S., Ludwig, S., Planz, O., 2017. The MEK-inhibitor CI-1040 displays a broad anti-influenza virus activity in vitro and provides a prolonged treatment window compared to standard of care in vivo. *Antivir. Res.* 142, 178–184.
- Hall, M.N., 2008. mTOR-what does it do? *Transpl. Proc.* 40 (10 Suppl), S5–8.
- Hamden, K.E., Whitman, A.G., Ford, P.W., Shelton, J.G., McCubrey, J.A., Akula, S.M., 2005. Raf and VEGF: emerging therapeutic targets in Kaposi's sarcoma-associated herpesvirus infection and angiogenesis in hematopoietic and nonhematopoietic tumors. *Leukemia* 19 (1), 18–26.
- Han, Z., Harty, R.N., 2007. Influence of calcium/calmodulin on budding of Ebola VLPs: implications for the involvement of the Ras/Raf/MEK/ERK pathway. *Virus Genes* 35 (3), 511–520.
- Han, J., Perez, J.T., Chen, C., Li, Y., Benitez, A., Kandasamy, M., Lee, Y., Andrade, J., tenOever, B., Manicassamy, B., 2018. Genome-wide CRISPR/Cas9 screen identifies host factors essential for influenza virus replication. *Cell. Rep.* 23 (2), 596–607.
- Hargett, D., McLean, T., Bachheimer, S.L., 2005. Herpes simplex virus ICP27 activation of stress kinases JNK and p38. *J. Virol.* 79 (13), 8348–8360.
- Harwig, A., Landick, R., Berkhout, B., 2017. The Battle of RNA synthesis: virus versus host. *Viruses* 9 (10).
- Hefner, Y., Borsch-Haubold, A.G., Murakami, M., Wilde, J.I., Pasquet, S., Schieltz, D., Ghomashchi, F., Yates 3rd, J.R., Armstrong, C.G., Paterson, A., Cohen, P., Fukunaga, R., Hunter, T., Kudo, I., Watson, S.P., Gelb, M.H., 2000. Serine 727 phosphorylation and activation of cytosolic phospholipase A2 by MNK1-related protein kinases. *J. Biol. Chem.* 275 (48), 37542–37551.
- Hemonnot, B., Cartier, C., Gay, B., Rebuffat, S., Bardy, M., Devaux, C., Boyer, V., Briant, L., 2004. The host cell MAP kinase ERK-2 regulates viral assembly and release by phosphorylating the p6gag protein of HIV-1. *J. Biol. Chem.* 279 (31), 32426–32434.
- Herget, T., Freitag, M., Morbitzer, M., Kupfer, R., Stamminger, T., Marschall, M., 2004. Novel chemical class of pUL97 protein kinase-specific inhibitors with strong anticytomegaloviral activity. *Antimicrob. Agents Chemother.* 48 (11), 4154–4162.
- Hirasawa, K., Kim, A., Han, H.S., Han, J., Jun, H.S., Yoon, J.W., 2003. Effect of p38 mitogen-activated protein kinase on the replication of encephalomyocarditis virus. *J. Virol.* 77 (10), 5649–5656.
- Hirsch, A.J., Medigeshi, G.R., Meyers, H.L., DeFilippis, V., Fruh, K., Briese, T., Lipkin, W.I., Nelson, J.A., 2005. The Src family kinase c-Yes is required for maturation of West Nile virus particles. *J. Virol.* 79 (18), 11943–11951.
- Holzberg, M., Boergeling, Y., Schrader, T., Ludwig, S., Ehrhardt, C., 2017. Vemurafenib limits influenza A virus propagation by targeting multiple signaling pathways. *Front. Microbiol.* 8, 2426.
- Huber, M., Watson, K.A., Selinka, H.C., Carthy, C.M., Klingel, K., McManus, B.M., Kandolf, R., 1999. Cleavage of RasGAP and phosphorylation of mitogen-activated protein kinase in the course of coxsackievirus B3 replication. *J. Virol.* 73 (5), 3587–3594.
- Imataka, H., Gradi, A., Sonenberg, N., 1998. A newly identified N-terminal amino acid sequence of human eIF4G binds poly(A)-binding protein and functions in poly(A)-dependent translation. *EMBO J.* 17 (24), 7480–7489.
- Jackson, R.J., 2013. The current status of vertebrate cellular mRNA IRESs. *Cold Spring Harb. Perspect. Biol.* 5 (2).
- Jan, E., Sarnow, P., 2002. Factorless ribosome assembly on the internal ribosome entry site of cricket paralysis virus. *J. Mol. Biol.* 324 (5), 889–902.
- Janne, P.A., Shaw, A.T., Pereira, J.R., Jeannin, G., Vansteenkiste, J., Barrios, C., Franke, F.A., Grinstead, L., Zazulina, V., Smith, P., Smith, I., Crino, L., 2013. Selumetinib plus docetaxel for KRAS-mutant advanced non-small-cell lung cancer: a randomised, multicentre, placebo-controlled, phase 2 study. *Lancet Oncol.* 14 (1), 38–47.
- Jham, B.C., Montaner, S., 2010. The Kaposi's sarcoma-associated herpesvirus G protein-coupled receptor: lessons on dysregulated angiogenesis from a viral oncogene. *J. Cell. Biochem.* 110 (1), 1–9.
- Jordan, B.A., Devi, L.A., 1999. G-protein-coupled receptor heterodimerization modulates receptor function. *Nature* 399 (6737), 697–700.
- Joubert, P.E., Stapleford, K., Guivel-Benhassine, F., Vignuzzi, M., Schwartz, O., Albert, M.L., 2015. Inhibition of mTORC1 enhances the translation of chikungunya proteins via the activation of the Mnk/eIF4E pathway. *PLoS Pathog* 11 (8), e1005091.
- Kamoshita, N., Nomoto, A., RajBhandary, U.L., 2009. Translation initiation from the ribosomal A site or the P site, dependent on the conformation of RNA pseudoknot I in dicistronic RNAs. *Mol. Cell.* 35 (2), 181–190.
- Karakama, Y., Sakamoto, N., Itsui, Y., Nakagawa, M., Tasaka-Fujita, M., Nishimura-Sakurai, Y., Kakinuma, S., Oooka, M., Azuma, S., Tsuchiya, K., Onogi, H., Hagiwara, M., Watanabe, M., 2010. Inhibition of hepatitis C virus replication by a specific inhibitor of serine-arginine-rich protein kinase. *Antimicrob. Agents Chemother.* 54 (8), 3179–3186.
- Karlas, A., Machuy, N., Shin, Y., Pleissner, K.P., Artarini, A., Heuer, D., Becker, D., Khalil, H., Ogilvie, L.A., Hess, S., Maurer, A.P., Muller, E., Wolff, T., Rudel, T., Meyer, T.F.,



2010. Genome-wide RNAi screen identifies human host factors crucial for influenza virus replication. *Nature* 463 (7282), 818–822.
- Kempf, B.J., Barton, D.J., 2008. Poliovirus 2A(Pro) increases viral mRNA and polysome stability coordinately in time with cleavage of eIF4G. *J. Virol.* 82 (12), 5847–5859.
- Kilcher, S., Schmidt, F.I., Schneider, C., Kopf, M., Helenius, A., Mercer, J., 2014. siRNA screen of early poxvirus genes identifies the AAA + + ATPase D5 as the virus genome-uncoating factor. *Cell. Host Microbe* 15 (1), 103–112.
- Kim, Y., Lee, C., 2015. Extracellular signal-regulated kinase (ERK) activation is required for porcine epidemic diarrhea virus replication. *Virology* 484, 181–193.
- Kim, D.H., Sim, T., 2012. Novel small molecule Raf kinase inhibitors for targeted cancer therapeutics. *Arch. Pharm. Res.* 35 (4), 605–615.
- Kim, C.S., Seol, S.K., Song, O.K., Park, J.H., Jang, S.K., 2007. An RNA-binding protein, hnRNP A1, and a scaffold protein, septin 6, facilitate hepatitis C virus replication. *J. Virol.* 81 (8), 3852–3865.
- Kleijn, M., Vriens, C.L., Voorma, H.O., Thomas, A.A., 1996. Phosphorylation state of the cap-binding protein eIF4E during viral infection. *Virology* 217 (2), 486–494.
- Kolokoltsov, A.A., Saeed, M.F., Freiberg, A.N., Holbrook, M.R., Davey, R.A., 2009. Identification of novel cellular targets for therapeutic intervention against Ebola virus infection by siRNA screening. *Drug. Dev. Res.* 70 (4), 255–265.
- Komarova, A.V., Real, E., Borman, A.M., Brocard, M., England, P., Tordo, N., Hershey, J.W., Kean, K.M., Jacob, Y., 2007. Rabies virus matrix protein interplay with eIF3, new insights into rabies virus pathogenesis. *Nucleic Acids Res.* 35 (5), 1522–1532.
- Krishna, M., Narang, H., 2008. The complexity of mitogen-activated protein kinases (MAPKs) made simple. *Cell. Mol. Life Sci.* 65 (22), 3525–3544.
- Krishnan, M.N., Ng, A., Sukumaran, B., Gilfoy, F.D., Uchil, P.D., Sultana, H., Brass, A.L., Adametz, R., Tsui, M., Qian, F., Montgomery, R.R., Lev, S., Mason, P.W., Koski, R.A., Elledge, S.J., Xavier, R.J., Agaisse, H., Fikrig, E., 2008. RNA interference screen for human genes associated with West Nile virus infection. *Nature* 455 (7210), 242–245.
- Kuang, E., Fu, B., Liang, Q., Myoung, J., Zhu, F., 2011. Phosphorylation of eukaryotic translation initiation factor 4B (EIF4B) by open reading frame 45/p90 ribosomal S6 kinase (ORF45/RSK) signaling axis facilitates protein translation during Kaposi sarcoma-associated herpesvirus (KSHV) lytic replication. *J. Biol. Chem.* 286 (48), 41171–41182.
- Kujime, K., Hashimoto, S., Gon, Y., Shimizu, K., Horie, T., 2000. p38 mitogen-activated protein kinase and c-jun-NH2-terminal kinase regulate RANTES production by influenza virus-infected human bronchial epithelial cells. *J. Immunol.* 164 (6), 3222–3228.
- Kumar, N., Xin, Z.T., Liang, Y., Ly, H., Liang, Y., 2008. NF-kappaB signaling differentially regulates influenza virus RNA synthesis. *J. Virol.* 82 (20), 9880–9889.
- Kumar, N., Liang, Y., Parslow, T.G., Liang, Y., 2011a. Receptor tyrosine kinase inhibitors block multiple steps of influenza A virus replication. *J. Virol.* 85 (6), 2818–2827.
- Kumar, N., Sharma, N.R., Ly, H., Parslow, T.G., Liang, Y., 2011b. Receptor tyrosine kinase inhibitors that block replication of influenza A and other viruses. *Antimicrob. Agents Chemother.* 55 (12), 5553–5559.
- Kuss-Duerkop, S.K., Wang, J., Mena, I., White, K., Metreveli, G., Sakthivel, R., Mata, M.A., Munoz-Moreno, R., Chen, X., Krammer, F., Diamond, M.S., Chen, Z.J., Garcia-Sastre, A., Fontoura, B.M.A., 2017. Influenza virus differentially activates mTORC1 and mTORC2 signaling to maximize late stage replication. *PLoS Pathog* 13 (9), e1006635.
- Langhammer, S., Koban, R., Yue, C., Ellerbrok, H., 2011. Inhibition of poxvirus spreading by the anti-tumor drug Gefitinib (Iressa). *Antivir. Res.* 89 (1), 64–70.
- Lavanya, M., Cuevas, C.D., Thomas, M., Cherry, S., Ross, S.R., 2013. siRNA screen for genes that affect Junin virus entry uncovers voltage-gated calcium channels as a therapeutic target. *Sci. Transl. Med.* 5 (204), 204ra.
- Le Sommer, C., Barrows, N.J., Bradrick, S.S., Pearson, J.L., Garcia-Blanco, M.A., 2012. G protein-coupled receptor kinase 2 promotes flaviviridae entry and replication. *PLoS Negl. Trop Dis.* 6 (9), e1820.
- Lee, B.J., Koszinowski, U.H., Sarawar, S.R., Adler, H., 2003. A gammaherpesvirus G protein-coupled receptor homologue is required for increased viral replication in response to chemokines and efficient reactivation from latency. *J. Immunol.* 170 (1), 243–251.
- Leite, J.F., Cascio, M., 2001. Structure of ligand-gated ion channels: critical assessment of biochemical data supports novel topology. *Mol. Cell. Neurosci.* 17 (5), 777–792.
- Lemmon, M.A., Schlessinger, J., 2010. Cell signaling by receptor tyrosine kinases. *Cell* 141 (7), 1117–1134.
- Levitzki, A., Gazit, A., 1995. Tyrosine kinase inhibition: an approach to drug development. *Science* 267 (5205), 1782–1788.
- Levy, D.E., Marie, I.J., Durbin, J.E., 2011. Induction and function of type I and III interferon in response to viral infection. *Curr. Opin. Virol.* 1 (6), 476–486.
- Li, H.P., Zhang, X., Duncan, R., Comai, L., Lai, M.M., 1997. Heterogeneous nuclear ribonucleoprotein A1 binds to the transcription-regulatory region of mouse hepatitis virus RNA. *Proc. Natl. Acad. Sci. U. S. A.* 94 (18), 9544–9549.
- Li, Q., Brass, A.L., Ng, A., Hu, Z., Xavier, R.J., Liang, T.J., Elledge, S.J., 2009. A genome-wide genetic screen for host factors required for hepatitis C virus propagation. *Proc. Natl. Acad. Sci. U. S. A.* 106 (38), 16410–16415.
- Li, L., Zhao, G.D., Shi, Z., Qi, L.L., Zhou, L.Y., Fu, Z.X., 2016. The Ras/Raf/MEK/ERK signaling pathway and its role in the occurrence and development of HCC. *Oncol. Lett.* 12 (5), 3045–3050.
- Lim, B.K., Nam, J.H., Gil, C.O., Yun, S.H., Choi, J.H., Kim, D.K., Jeon, E.S., 2005. Cocksackievirus B3 replication is related to activation of the late extracellular signal-regulated kinase (ERK) signal. *Virus Res.* 113 (2), 153–157.
- Lin, J.Y., Shih, S.R., Pan, M., Li, C., Lue, C.F., Stollar, V., Li, M.L., 2009. hnRNP A1 interacts with the 5' untranslated regions of enterovirus 71 and Sindbis virus RNA and is required for viral replication. *J. Virol.* 83 (12), 6106–6114.
- Liu, Y., Wimmer, E., Paul, A.V., 2009. Cis-acting RNA elements in human and animal plus-strand RNA viruses. *Biochim. Biophys. Acta* 1789 (9–10), 495–517.
- Locker, N., Easton, L.E., Lukavsky, P.J., 2007. HCV and CSFV IRES domain II mediate eIF2 release during 80S ribosome assembly. *EMBO J.* 26 (3), 795–805.
- Locker, N., Chamond, N., Sargueil, B., 2011. A conserved structure within the HIV gag open reading frame that controls translation initiation directly recruits the 40S subunit and eIF3. *Nucleic Acids Res.* 39 (6), 2367–2377.
- Locksley, R.M., Killeen, N., Lenardo, M.J., 2001. The TNF and TNF receptor superfamilies: integrating mammalian biology. *Cell* 104 (4), 487–501.
- Ludwig, S., Ehrhardt, C., Neumeier, E.R., Kracht, M., Rapp, U.R., Pleschka, S., 2001. Influenza virus-induced AP-1-dependent gene expression requires activation of the JNK signaling pathway. *J. Biol. Chem.* 276 (24), 10990–10998.
- Ludwig, S., Wolff, T., Ehrhardt, C., Wurzer, W.J., Reinhardt, J., Planz, O., Pleschka, S., 2004. MEK inhibition impairs influenza B virus propagation without emergence of resistant variants. *FEBS Lett.* 561 (1–3), 37–43.
- Ludwig, S., Pleschka, S., Planz, O., Wolff, T., 2006. Ringing the alarm bells: signalling and apoptosis in influenza virus infected cells. *Cell. Microbiol.* 8 (3), 375–386.
- Luo, H., Yanagawa, B., Zhang, J., Luo, Z., Zhang, M., Esfandiari, M., Carthy, C., Wilson, J.E., Yang, D., McManus, B.M., 2002. Cocksackievirus B3 replication is reduced by inhibition of the extracellular signal-regulated kinase (ERK) signaling pathway. *J. Virol.* 76 (7), 3365–3373.
- Lupberger, J., Casanova, C., Fischer, B., Weiss, A., Fofana, I., Fontaine, N., Fujiwara, T., Renaud, M., Kopp, A., Schuster, C., Brino, L., Baumert, T.F., Thoma, C., 2015. PI4K-beta and MKNK1 are regulators of hepatitis C virus IRES-dependent translation. *Sci. Rep.* 5, 13344.
- Mannova, P., Beretta, L., 2005. Activation of the N-Ras-PI3K-Akt-mTOR pathway by hepatitis C virus: control of cell survival and viral replication. *J. Virol.* 79 (14), 8742–8749.
- Margolis, B., Skolnik, E.Y., 1994. Activation of Ras by receptor tyrosine kinases. *J. Am. Soc. Nephrol.* 5 (6), 1288–1299.
- Maroni, P.D., Koul, S., Meacham, R.B., Koul, H.K., 2004. Mitogen activated protein kinase signal transduction pathways in the prostate. *Cell. Commun. Signal.* 2 (1), 5.
- Matsuda, Y., Fukumoto, M., 2011. Sorafenib: complexities of Raf-dependent and Raf-independent signaling are now unveiled. *Med. Mol. Morphol.* 44 (4), 183–189.
- McCubrey, J.A., Steelman, L.S., Chappell, W.H., Abrams, S.L., Wong, E.W., Chang, F., Lehmann, B., Terrian, D.M., Milella, M., Tafuri, A., Stivala, F., Libra, M., Basecke, J., Evangelisti, C., Martelli, A.M., Franklin, R.A., 2007. Roles of the Raf/MEK/ERK pathway in cell growth, malignant transformation and drug resistance. *Biochim. Biophys. Acta* 1773 (8), 1263–1284.
- McLean, T.L., Bachenheimer, S.L., 1999. Activation of cJUN N-terminal kinase by herpes simplex virus type 1 enhances viral replication. *J. Virol.* 73 (10), 8415–8426.
- McNulty, S., Flint, M., Nichol, S.T., Spiropoulou, C.F., 2013. Host mTORC1 signaling regulates andes virus replication. *J. Virol.* 87 (2), 912–922.
- Medders, K.E., Kaul, M., 2011. Mitogen-activated protein kinase p38 in HIV infection and associated brain injury. *J. Neuroimmune Pharmacol.* 6 (2), 202–215.
- Michelangeli, F., East, J.M., 2011. A diversity of SERCA Ca2+ pump inhibitors. *Biochem. Soc. Trans* 39 (3), 789–797.
- Mir, M.A., Panganiban, A.T., 2008. A protein that replaces the entire cellular eIF4E complex. *EMBO J.* 27 (23), 3129–3139.
- Mizutani, T., Fukushi, S., Saijo, M., Kurane, I., Morikawa, S., 2004. Phosphorylation of p38 MAPK and its downstream targets in SARS coronavirus-infected cells. *Biochem Biophys Res. Commun.* 319 (4), 1228–1234.
- Morchang, A., Lee, R.C.H., Yenchitsomanus, P.T., Sreekanth, G.P., Noisakran, S., Chu, J.J.H., Limjindaporn, T., 2017. RNAi screen reveals a role of SPHK2 in dengue virus-mediated apoptosis in hepatic cell lines. *PLoS One* 12 (11), e0188121.
- Moser, L.A., Schultz-Cherry, S., 2008. Suppression of astrovirus replication by an ERK1/2 inhibitor. *J. Virol.* 82 (15), 7475–7482.
- Nacken, W., Ehrhardt, C., Ludwig, S., 2012. Small molecule inhibitors of the c-Jun N-terminal kinase (JNK) possess antiviral activity against highly pathogenic avian and human pandemic influenza A viruses. *Biol. Chem.* 393 (6), 525–534.
- Nakano, H., Shindo, M., Sakon, S., Nishinaka, S., Mihara, M., Yagita, H., Okumura, K., 1998. Differential regulation of IkkappaB kinase alpha and beta by two upstream kinases, NF-kappaB-inducing kinase and mitogen-activated protein kinase/ERK kinase-1. *Proc. Natl. Acad. Sci. U. S. A.* 95 (7), 3537–3542.
- Neves, S.R., Ram, P.T., Iyengar, R., 2002. G protein pathways. *Science* 296 (5573), 1636–1639.
- O'Loughlin, A., Gonzalez, V.M., Pineiro, D., Perez-Morgado, M.I., Salinas, M., Martin, M.E., 2004. Identification and molecular characterization of Mnk1b, a splice variant of human MAP kinase-interacting kinase Mnk1. *Exp. Cell. Res.* 299 (2), 343–355.
- O'Shea, C., Klupsch, K., Choi, S., Bagus, B., Soria, C., Shen, J., McCormick, F., Stokoe, D., 2005. Adenoviral proteins mimic nutrient/growth signals to activate the mTOR pathway for viral replication. *EMBO J.* 24 (6), 1211–1221.
- O'Sullivan, A.W., Wang, J.H., Redmond, H.P., 2009. The role of P38 MAPK and PKC in BLP induced TNF-alpha release, apoptosis, and NFkappaB activation in THP-1 monocyte cells. *J. Surg. Res.* 151 (1), 138–144.
- Pataer, A., Swisher, S.G., Roth, J.A., Logothetis, C.J., Corn, P.G., 2009. Inhibition of RNA-dependent protein kinase (PKR) leads to cancer cell death and increases chemosensitivity. *Cancer Biol. Ther.* 8 (3), 245–252.
- Pathak, K.B., Pogany, J., Xu, K., White, K.A., Nagy, P.D., 2012. Defining the roles of cis-acting RNA elements in tombusvirus replicase assembly in vitro. *J. Virol.* 86 (1), 156–171.
- Pei, R., Zhang, X., Xu, S., Meng, Z., Roggendorf, M., Lu, M., Chen, X., 2012. Regulation of hepatitis C virus replication and gene expression by the MAPK-ERK pathway. *Virol. Sin* 27 (5), 278–285.
- Planz, O., Pleschka, S., Ludwig, S., 2001. MEK-specific inhibitor U0126 blocks spread of Borna disease virus in cultured cells. *J. Virol.* 75 (10), 4871–4877.
- Player, M.R., 2009. Protein kinase inhibitors for the treatment of inflammatory disease. *Curr. Top. Med. Chem.* 9 (7), 598.
- Pleschka, S., Wolff, T., Ehrhardt, C., Hobom, G., Planz, O., Rapp, U.R., Ludwig, S., 2001.



- Influenza virus propagation is impaired by inhibition of the Raf/MEK/ERK signalling cascade. *Nat. Cell. Biol.* 3 (3), 301–305.
- Plotch, S.J., Bouloy, M., Ulmanen, I., Krug, R.M., 1981. A unique cap(m7GpppXm)-dependent influenza virion endonuclease cleaves capped RNAs to generate the primers that initiate viral RNA transcription. *Cell* 23 (3), 847–858.
- Plotnikov, A., Zehorai, E., Procaccia, S., Seger, R., 2011. The MAPK cascades: signaling components, nuclear roles and mechanisms of nuclear translocation. *Biochim. Biophys. Acta* 1813 (9), 1619–1633.
- Porta, C., Paglino, C., Mosca, A., 2014. Targeting PI3K/Akt/mTOR signaling in cancer. *Front. Oncol.* 4, 64.
- Qie, L., Marcellino, D., Herold, B.C., 1999. Herpes simplex virus entry is associated with tyrosine phosphorylation of cellular proteins. *Virology* 256 (2), 220–227.
- Radoshitzky, S.R., Pegoraro, G., Chi, X.O., L, D.N., Chiang, C.Y., Jozwick, L., Clester, J.C., Cooper, C.L., Courier, D., Langan, D.P., Underwood, K., Kuehl, K.A., Sun, M.G., Cai, Y., Yu, S.Q., Burk, R., Zamani, R., Kota, K., Kuhn, J.H., Bavari, S., 2016. siRNA screen identifies trafficking host factors that modulate alphavirus infection. *PLoS Pathog* 12 (3), e1005466.
- Rahaus, M., Desloges, N., Wolff, M.H., 2004. Replication of varicella-zoster virus is influenced by the levels of JNK/SAPK and p38/MAPK activation. *J. Gen. Virol.* 85 (Pt 12), 3529–3540.
- Rajagopalan, S., 2010. Endosomal signaling and a novel pathway defined by the natural killer receptor KIR2DL4 (CD158d). *Traffic* 11 (11), 1381–1390.
- Rajalingam, K., Schreck, R., Rapp, U.R., Albert, S., 2007. Ras oncogenes and their downstream targets. *Biochim. Biophys. Acta* 1773 (8), 1177–1195.
- Raman, M., Chen, W., Cobb, M.H., 2007. Differential regulation and properties of MAPKs. *Oncogene* 26 (22), 3100–3112.
- Ravichandran, V., Jensen, P.N., Major, E.O., 2007. MEK1/2 inhibitors block basal and transforming growth factor 1beta1-stimulated JC virus multiplication. *J. Virol.* 81 (12), 6412–6418.
- Reeves, P.M., Bommarius, B., Lebeis, S., McNulty, S., Christensen, J., Swimm, A., Chahroudi, A., Chavan, R., Feinberg, M.B., Veach, D., Bornmann, W., Sherman, M., Kalman, D., 2005. Disabling poxvirus pathogenesis by inhibition of Abl-family tyrosine kinases. *Nat. Med.* 11 (7), 731–739.
- Reeves, P.M., Smith, S.K., Olson, V.A., Thorne, S.H., Bornmann, W., Damon, I.K., Kalman, D., 2011. Variola and monkeypox viruses utilize conserved mechanisms of virion motility and release that depend on abl and SRC family tyrosine kinases. *J. Virol.* 85 (1), 21–31.
- Richter, J.D., Sonenberg, N., 2005. Regulation of cap-dependent translation by eIF4E inhibitory proteins. *Nature* 433 (7025), 477–480.
- Riolobos, L., Valle, N., Hernando, E., Maroto, B., Kann, M., Almendral, J.M., 2010. Viral oncolysis that targets Raf-1 signaling control of nuclear transport. *J. Virol.* 84 (4), 2090–2099.
- Robbins, D.J., Zhen, E., Owaki, H., Vanderbilt, C.A., Ebert, D., Geppert, T.D., Cobb, M.H., 1993. Regulation and properties of extracellular signal-regulated protein kinases 1 and 2 in vitro. *J. Biol. Chem.* 268 (7), 5097–5106.
- Rodriguez, M.E., Brunetti, J.E., Wachsman, M.B., Scolaro, L.A., Castilla, V., 2014. Raf/MEK/ERK pathway activation is required for Junin virus replication. *J. Gen. Virol.* 95 (Pt 4), 799–805.
- Rojas, M., Arias, C.F., Lopez, S., 2010. Protein kinase R is responsible for the phosphorylation of eIF2alpha in rotavirus infection. *J. Virol.* 84 (20), 10457–10466.
- Rose, B.A., Force, T., Wang, Y., 2010. Mitogen-activated protein kinase signaling in the heart: angels versus demons in a heart-breaking tale. *Physiol. Rev.* 90 (4), 1507–1546.
- Roskoski Jr., R., 2010. RAF protein-serine/threonine kinases: structure and regulation. *Biochem. Biophys. Res. Commun.* 399 (3), 313–317.
- Roskoski Jr., R., 2012. MEK1/2 dual-specificity protein kinases: structure and regulation. *Biochem. Biophys. Res. Commun.* 417 (1), 5–10.
- Roth, H., Magg, V., Uch, F., Mutz, P., Klein, P., Haneke, K., Lohmann, V., Bartenschlager, R., Fackler, O.T., Locker, N., Stoecklin, G., Ruggieri, A., 2017. Flavivirus infection uncouples translation suppression from cellular stress responses. *MBio* 8 (1).
- Roux, P.P., Blenis, J., 2004. ERK and p38 MAPK-activated protein kinases: a family of protein kinases with diverse biological functions. *Microbiol. Mol. Biol. Rev.* 68 (2), 320–344.
- Rushworth, L.K., Hindley, A.D., O'Neill, E., Kolch, W., 2006. Regulation and role of Raf-1/B-Raf heterodimerization. *Mol. Cell. Biol.* 26 (6), 2262–2272.
- Rusnati, M., Urbinati, C., Musulin, B., Ribatti, D., Albini, A., Noonan, D., Marchisone, C., Waltenberger, J., Presta, M., 2001. Activation of endothelial cell mitogen activated protein kinase ERK1/2 by extracellular HIV-1 Tat protein. *Endothelium* 8 (1), 65–74.
- Saeedi, B.J., Geiss, B.J., 2013. Regulation of flavivirus RNA synthesis and capping. *Wiley Interdiscip. Rev. RNA* 4 (6), 723–735.
- Sanford, J.R., Gray, N.K., Beckmann, K., Caceres, J.F., 2004. A novel role for shuttling SR proteins in mRNA translation. *Genes Dev.* 18 (7), 755–768.
- Savidis, G., McDougall, W.M., Meraner, P., Pereira, J.M., Portmann, J.M., Trincucci, G., John, S.P., Aker, A.M., Renzette, N., Robbins, D.R., Guo, Z., Green, S., Kowalik, T.F., Brass, A.L., 2016. Identification of Zika virus and dengue virus dependency factors using functional genomics. *Cell. Rep.* 16 (1), 232–246.
- Schlessinger, J., 2000. Cell signaling by receptor tyrosine kinases. *Cell* 103 (2), 211–225.
- Segaliny, A.I., Tellez-Gabriel, M., Heymann, M.F., Heymann, D., 2015. Receptor tyrosine kinases: characterisation, mechanism of action and therapeutic interests for bone cancers. *J. Bone Oncol.* 4 (1), 1–12.
- Sessions, O.M., Barrows, N.J., Souza-Neto, J.A., Robinson, T.J., Hershey, C.L., Rodgers, M.A., Ramirez, J.L., Dimopoulos, G., Yang, P.L., Pearson, J.L., Garcia-Blanco, M.A., 2009. Discovery of insect and human dengue virus host factors. *Nature* 458 (7241), 1047–1050.
- Sharma, S., Mulik, S., Kumar, N., Suryawanshi, A., Rouse, B.T., 2011. An anti-inflammatory role of VEGFR2/Src kinase inhibitor in herpes simplex virus 1-induced immunopathology. *J. Virol.* 85 (12), 5995–6007.
- Shaul, Y.D., Seger, R., 2007. The MEK/ERK cascade: from signaling specificity to diverse functions. *Biochim. Biophys. Acta* 1773 (8), 1213–1226.
- Shenoy-Scaria, A.M., Kwong, J., Fujita, T., Olszowy, M.W., Shaw, A.S., Lublin, D.M., 1992. Signal transduction through decay-accelerating factor. Interaction of glycosyl-phosphatidylinositol anchor and protein tyrosine kinases p56lck and p59fyn 1. *J. Immunol.* 149 (11), 3535–3541.
- Shi, S.T., Huang, P., Li, H.P., Lai, M.M., 2000. Heterogeneous nuclear ribonucleoprotein A1 regulates RNA synthesis of a cytoplasmic virus. *EMBO J.* 19 (17), 4701–4711.
- Shi, S.T., Yu, G.Y., Lai, M.M., 2003. Multiple type A/B heterogeneous nuclear ribonucleoproteins (hnRNPs) can replace hnRNP A1 in mouse hepatitis virus RNA synthesis. *J. Virol.* 77 (19), 10584–10593.
- Shi, W., Hou, X., Li, X., Peng, H., Shi, M., Jiang, Q., Liu, X., Ji, Y., Yao, Y., He, C., Lei, X., 2013. Differential gene expressions of the MAPK signaling pathway in enterovirus 71-infected rhabdomyosarcoma cells. *Braz. J. Infect. Dis.* 17 (4), 410–417.
- Shmulevitz, M., Marcato, P., Lee, P.W., 2010. Activated Ras signaling significantly enhances reovirus replication and spread. *Cancer Gene Ther.* 17 (1), 69–70.
- Snyder, A., Alsauskas, Z.C., Leventhal, J.S., Rosenstiel, P.E., Gong, P., Chan, J.J., Barley, K., He, J.C., Klotman, M.E., Ross, M.J., Klotman, P.E., 2010. HIV-1 viral protein r induces ERK and caspase-8-dependent apoptosis in renal tubular epithelial cells. *AIDS* 24 (8), 1107–1119.
- Soares, J.A., Leite, F.G., Andrade, L.G., Torres, A.A., De Sousa, L.P., Barcelos, L.S., Teixeira, M.M., Ferreira, P.C., Kroon, E.G., Souto-Padron, T., Bonjardim, C.A., 2009. Activation of the PI3K/Akt pathway early during vaccinia and cowpox virus infections is required for both host survival and viral replication. *J. Virol.* 83 (13), 6883–6899.
- Sreekanth, G.P., Chuncharunee, A., Sirimontaporn, A., Panaampon, J., Srisawat, C., Morchang, A., Malakar, S., Thuwajit, K., Kooptiwut, S., Suttithetmrong, A., Songprakhon, P., Noisakran, S., Yenchitsomanus, P.T., Limjindaporn, T., 2014. Role of ERK1/2 signaling in dengue virus-induced liver injury. *Virus Res.* 188, 15–26.
- Srivastava, S.P., Kumar, K.U., Kaufman, R.J., 1998. Phosphorylation of eukaryotic translation initiation factor 2 mediates apoptosis in response to activation of the double-stranded RNA-dependent protein kinase. *J. Biol. Chem.* 273 (4), 2416–2423.
- Staff, P.P., 2015. Correction: coronavirus cell entry occurs through the endo-/lysosomal pathway in a proteolysis-dependent manner. *PLoS Pathog.* 11 (2), e1004709.
- Steelman, L.S., Pohnert, S.C., Shelton, J.G., Franklin, R.A., Bertrand, F.E., McCubrey, J.A., 2004. JAK/STAT, Raf/MEK/ERK, PI3K/Akt and BCR-ABL in cell cycle progression and leukemogenesis. *Leukemia* 18 (2), 189–218.
- Stohr, S., Costa, R., Sandmann, L., Westhaus, S., Pfaender, S., Anggakusuma, Dazert, E., Meuleman, P., Vondran, F.W., Manns, M.P., Steinmann, E., von Hahn, T., Ciesek, S., 2016. Host cell mTORC1 is required for HCV RNA replication. *Gut* 65 (12), 2017–2028.
- Sturzl, M., Roth, W.K., Brockmeyer, N.H., Zietz, C., Speiser, B., Hofschneider, P.H., 1992. Expression of platelet-derived growth factor and its receptor in AIDS-related Kaposi sarcoma in vivo suggests paracrine and autocrine mechanisms of tumor maintenance. *Proc. Natl. Acad. Sci. U. S. A.* 89 (15), 7046–7050.
- Sullivan, M.L., Ahlquist, P., 1997. cis-Acting Signals in Bromovirus RNA Replication and Gene Expression: Networking With Viral Proteins and Host Factors. pp. 221–230 8.
- Svitkin, Y.V., Herdy, B., Costa-Mattioli, M., Gingras, A.C., Raught, B., Sonenberg, N., 2005. Eukaryotic translation initiation factor 4E availability controls the switch between cap-dependent and internal ribosomal entry site-mediated translation. *Mol. Cell. Biol.* 25 (23), 10556–10565.
- Sweeney, S.E., Firestein, G.S., 2006. Mitogen activated protein kinase inhibitors: where are we now and where are we going? *Ann. Rheum. Dis.* 65 (Suppl 3), iii83–88.
- Sweeney, T.R., Abaeva, I.S., Pestova, T.V., Hellen, C.U., 2014. The mechanism of translation initiation on type 1 picornavirus IRESs. *EMBO J.* 33 (1), 76–92.
- Taghavi, N., Samuel, C.E., 2012. Protein kinase PKR catalytic activity is required for the PKR-dependent activation of mitogen-activated protein kinases and amplification of interferon beta induction following virus infection. *Virology* 427 (2), 208–216.
- Tai, A.W., Benita, Y., Peng, L.F., Kim, S.S., Sakamoto, N., Xavier, R.J., Chung, R.T., 2009. A functional genomic screen identifies cellular cofactors of hepatitis C virus replication. *Cell. Host Microbe* 5 (3), 298–307.
- Tang, H., Delgermaa, L., Huang, F., Oishi, N., Liu, L., He, F., Zhao, L., Murakami, S., 2005. The transcriptional transactivation function of HBx protein is important for its augmentation role in hepatitis B virus replication. *J. Virol.* 79 (9), 5548–5556.
- Thompson, S.R., Sarnow, P., 2003. Enterovirus 71 contains a type I IRES element that functions when eukaryotic initiation factor eIF4G is cleaved. *Virology* 315 (1), 259–266.
- Ting, J.P., Duncan, J.A., Lei, Y., 2010. How the noninflammatory NLRs function in the innate immune system. *Science* 327 (5963), 286–290.
- Tokunaga, K., Kiyokawa, E., Nakaya, M., Otsuka, N., Kojima, A., Kurata, T., Matsuda, M., 1998. Inhibition of human immunodeficiency virus type 1 virion entry by dominant-negative Hck. *J. Virol.* 72 (7), 6257–6259.
- Tolbert, M., Morgan, C.E., Pollum, M., Crespo-Hernandez, C.E., Li, M.L., Brewer, G., Tolbert, B.S., 2017. HnRNP A1 alters the structure of a conserved enterovirus IRES domain to stimulate viral translation. *J. Mol. Biol.* 429 (19), 2841–2858.
- Trahey, M., Wong, G., Halenbeck, R., Rubinfeld, B., Martin, G.A., Ladner, M., Long, C.M., Crosier, W.J., Watt, K., Koths, K., et al., 1988. Molecular cloning of two types of GAP complementary DNA from human placenta. *Science* 242 (4886), 1697–1700.
- Uehling, D.E., Harris, P.A., 2015. Recent progress on MAP kinase pathway inhibitors. *Bioorg. Med. Chem. Lett.* 25 (19), 4047–4056.
- Ugi, S., Imamura, T., Maegawa, H., Egawa, K., Yoshizaki, T., Shi, K., Obata, T., Ebina, Y., Kashiwagi, A., Olefsky, J.M., 2004. Protein phosphatase 2A negatively regulates insulin's metabolic signaling pathway by inhibiting Akt (protein kinase B) activity in 3T3-L1 adipocytes. *Mol. Cell. Biol.* 24 (19), 8778–8789.

- Vadlakonda, L., Dash, A., Pasupuleti, M., Anil Kumar, K., Reddanna, P., 2013. The paradox of Akt-mTOR interactions. *Front. Oncol.* 3, 165.
- Valcarcel, J., Gebauer, F., 1997. Post-transcriptional regulation: the dawn of PTB. *Curr. Biol.* 7 (11), R705–708.
- Ventoso, I., Sanz, M.A., Molina, S., Berlanga, J.J., Carrasco, L., Esteban, M., 2006. Translational resistance of late alphavirus mRNA to eIF2alpha phosphorylation: a strategy to overcome the antiviral effect of protein kinase PKR. *Genes Dev.* 20 (1), 87–100.
- Vert, G., Chory, J., 2011. Crosstalk in cellular signaling: background noise or the real thing? *Dev. Cell.* 21 (6), 985–991.
- Voss, K., Amaya, M., Mueller, C., Roberts, B., Kehn-Hall, K., Bailey, C., Petricoin 3rd, E., Narayanan, A., 2014. Inhibition of host extracellular signal-regulated kinase (ERK) activation decreases new world alphavirus multiplication in infected cells. *Virology* 468–470, 490–503.
- Walsh, D., Mohr, I., 2004. Phosphorylation of eIF4E by Mnk-1 enhances HSV-1 translation and replication in quiescent cells. *Genes Dev.* 18 (6), 660–672.
- Walsh, D., Perez, C., Notary, J., Mohr, I., 2005. Regulation of the translation initiation factor eIF4F by multiple mechanisms in human cytomegalovirus-infected cells. *J. Virol.* 79 (13), 8057–8064.
- Walsh, D., Arias, C., Perez, C., Halladin, D., Escandon, M., Ueda, T., Watanabe-Fukunaga, R., Fukunaga, R., Mohr, I., 2008. Eukaryotic translation initiation factor 4F architectural alterations accompany translation initiation factor redistribution in poxvirus-infected cells. *Mol. Cell. Biol.* 28 (8), 2648–2658.
- Walsh, D., Mathews, M.B., Mohr, I., 2013. Tinkering with translation: protein synthesis in virus-infected cells. *Cold Spring Harb. Perspect. Biol.* 5 (1), a012351.
- Wang, L., Jeng, K.S., Lai, M.M., 2011. Poly(C)-binding protein 2 interacts with sequences required for viral replication in the hepatitis C virus (HCV) 5' untranslated region and directs HCV RNA replication through circularizing the viral genome. *J. Virol.* 85 (16), 7954–7964.
- Wang, Y., Zhang, X., 1999. The nucleocapsid protein of coronavirus mouse hepatitis virus interacts with the cellular heterogeneous nuclear ribonucleoprotein A1 in vitro and in vivo. *Virology* 265 (1), 96–109.
- Waskiewicz, A.J., Flynn, A., Proud, C.G., Cooper, J.A., 1997. Mitogen-activated protein kinases activate the serine/threonine kinases Mnk1 and Mnk2. *EMBO J.* 16 (8), 1909–1920.
- Waskiewicz, A.J., Johnson, J.C., Penn, B., Mahalingam, M., Kimball, S.R., Cooper, J.A., 1999. Phosphorylation of the cap-binding protein eukaryotic translation initiation factor 4E by protein kinase Mnk1 in vivo. *Mol. Cell. Biol.* 19 (3), 1871–1880.
- Watanabe, T., Watanabe, S., Kawaoka, Y., 2010. Cellular networks involved in the influenza virus life cycle. *Cell. Host Microbe* 7 (6), 427–439.
- Weber, F., Wagner, V., Rasmussen, S.B., Hartmann, R., Paludan, S.R., 2006. Double-stranded RNA is produced by positive-strand RNA viruses and DNA viruses but not in detectable amounts by negative-strand RNA viruses. *J. Virol.* 80 (10), 5059–5064.
- Wei, L., Zhu, S., Wang, J., Liu, J., 2012. Activation of the phosphatidylinositol 3-kinase/Akt signaling pathway during porcine circovirus type 2 infection facilitates cell survival and viral replication. *J. Virol.* 86 (24), 13589–13597.
- Wennerberg, K., Rossman, K.L., Der, C.J., 2005. The Ras superfamily at a glance. *J. Cell. Sci.* 118 (Pt 5), 843–846.
- Widmann, C., Gibson, S., Jarpe, M.B., Johnson, G.L., 1999. Mitogen-activated protein kinase: conservation of a three-kinase module from yeast to human. *Physiol. Rev.* 79 (1), 143–180.
- Willcocks, M.M., Locker, N., Gomwalk, Z., Royall, E., Bakhshesh, M., Belsham, G.J., Idamakanti, N., Burroughs, K.D., Reddy, P.S., Hallenbeck, P.L., Roberts, L.O., 2011. Structural features of the Seneca Valley virus internal ribosome entry site (IRES) element: a picornavirus with a pestivirus-like IRES. *J. Virol.* 85 (9), 4452–4461.
- Wilson, J.E., Pestova, T.V., Hellen, C.U., Sarnow, P., 2000. Initiation of protein synthesis from the A site of the ribosome. *Cell* 102 (4), 511–520.
- Wise, L.M., Veikkola, T., Mercer, A.A., Savory, L.J., Fleming, S.B., Caesar, C., Vitali, A., Mäkinen, T., Alitalo, K., Stacker, S.A., 1999. Vascular endothelial growth factor (VEGF)-like protein from orf virus NZ2 binds to VEGFR2 and neuropilin-1. *Proc. Natl. Acad. Sci. U. S. A.* 96 (6), 3071–3076.
- Xie, J., Ajibade, A.O., Ye, F., Kuhne, K., Gao, S.J., 2008. Reactivation of Kaposi's sarcoma-associated herpesvirus from latency requires MEK/ERK, JNK and p38 multiple mitogen-activated protein kinase pathways. *Virology* 371 (1), 139–154.
- Xing, Z., Cardona, C.J., Anunciacion, J., Adams, S., Dao, N., 2010. Roles of the ERK MAPK in the regulation of proinflammatory and apoptotic responses in chicken macrophages infected with H9N2 avian influenza virus. *J. Gen. Virol.* 91 (Pt 2), 343–351.
- Yan, J., Roy, S., Apolloni, A., Lane, A., Hancock, J.F., 1998. Ras isoforms vary in their ability to activate Raf-1 and phosphoinositide 3-kinase. *J. Biol. Chem.* 273 (37), 24052–24056.
- Yan, F., Wang, X.M., Liu, Z.C., Pan, C., Yuan, S.B., Ma, Q.M., 2010. JNK1, JNK2, and JNK3 are involved in P-glycoprotein-mediated multidrug resistance of hepatocellular carcinoma cells. *Hepatobiliary Pancreat. Dis. Int.* 9 (3), 287–295.
- Yanguz, E., Rodriguez, P., Goodfellow, I., Nieto, A., 2012. Influenza virus polymerase confers independence of the cellular cap-binding factor eIF4E for viral mRNA translation. *Virology* 422 (2), 297–307.
- Yin, R., Liu, X., Bi, Y., Xie, G., Zhang, P., Meng, X., Ai, L., Xu, R., Sun, Y., Stoeger, T., Ding, Z., 2015. Expression of Raf kinase inhibitor protein is downregulated in response to Newcastle disease virus infection to promote viral replication. *J. Gen. Virol.* 96 (9), 2579–2586.
- Yu, L.G., Packman, L.C., Weldon, M., Hamlett, J., Rhodes, J.M., 2004. Protein phosphatase 2A, a negative regulator of the ERK signaling pathway, is activated by tyrosine phosphorylation of putative HLA class II-associated protein I (PHAPI)/pp32 in response to the antiproliferative lectin, jacalin. *J. Biol. Chem.* 279 (40), 41377–41383.
- Zebisch, A., Troppmair, J., 2006. Back to the roots: the remarkable RAF oncogene story. *Cell. Mol. Life Sci.* 63 (11), 1314–1330.
- Zhang, J., Yang, P.L., Gray, N.S., 2009. Targeting cancer with small molecule kinase inhibitors. *Nat. Rev. Cancer* 9 (1), 28–39.
- Zhang, H., Niu, X., Qian, Z., Qian, J., Xuan, B., 2015. The c-Jun N-terminal kinase inhibitor SP600125 inhibits human cytomegalovirus replication. *J. Med. Virol.* 87 (12), 2135–2144.
- Zhao, L., Imperiale, M.J., 2017. Identification of Rab18 as an essential host factor for BK polyomavirus infection using a whole-genome RNA interference screen. *mSphere* 2 (4).
- Zhen, Z., Bradel-Tretheway, B., Sumagin, S., Bidlack, J.M., Dewhurst, S., 2005. The human herpesvirus 6 G protein-coupled receptor homolog U51 positively regulates virus replication and enhances cell-cell fusion in vitro. *J. Virol.* 79 (18), 11914–11924.
- Zhou, H., Xu, M., Huang, Q., Gates, A.T., Zhang, X.D., Castle, J.C., Stec, E., Ferrer, M., Strulovici, B., Hazuda, D.J., Espeseth, A.S., 2008. Genome-scale RNAi screen for host factors required for HIV replication. *Cell. Host Microbe* 4 (5), 495–504.