

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/314224808>

# Phytochemical studies on antioxidant activities of two types of Karonda ( *Carissa carandas* ) during storage

Article in *Indian Journal of Horticulture* · December 2016

DOI: 10.5958/0974-0112.2016.00126.2

CITATIONS

0

READS

76

5 authors, including:



**Debashis Dutta**

ICAR-Indian Institute of Farming Systems Research

13 PUBLICATIONS 64 CITATIONS

[SEE PROFILE](#)



**Amit Nath**

29 PUBLICATIONS 359 CITATIONS

[SEE PROFILE](#)



**Dushyant Mishra**

ICAR Indian Institute of Farming Systems Research

9 PUBLICATIONS 16 CITATIONS

[SEE PROFILE](#)



**Nisha Verma**

ICAR-Indian Institute of Farming Systems Research

13 PUBLICATIONS 5 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Value addition of mulberry silk waste/ wool blends to develop handloom [View project](#)



Aonla improvement [View project](#)



## Short communication

# Phytochemical studies on antioxidant activities of two types of *Karonda* (*Carissa carandas*) during storage

Debashis Dutta, Amit Nath\*, Dushyant Mishra, Nisha Verma and Pawan Kumar  
ICAR- Indian Institute of Farming Systems Research, Modipuram 250110, Uttar Pradesh

### ABSTRACT

*Carissa carandas* Linn. is a large dichotomously branched evergreen shrub. It has several medicinal and nutritional properties. An experiment was conducted considering two types, viz., pinkish-green and pinkish-white to compare the phytochemical changes in perspective of antioxidant activity during storage. In fresh fruits, the maximum ascorbic acid content (50.98 mg/100 g) was recorded in pinkish green fruits as compared to pinkish-white (36.86 mg/100 g). Similarly, the scavenging activity of methanol extract of pinkish-green fruits was found significantly higher (89.57%) as compared to pinkish-white fruits (86.27%) in fresh samples. However, the minimum loss in phenol content (11.36%) was recorded in pinkish-white type during storage upto 8 days. Therefore, *karonda* fruits may be considered as an important phyto-chemical source for making different traditional medicines.

**Key words:** Ascorbic acid, antioxidant activity, *Carissa carandas*, flavonoids.

*Carissa carandas* Linn. is commonly known as *karonda* (2n = 22) belongs to Apocynaceae family and found throughout India mainly in the semi-arid regions and have a long history of use in Indian traditional system of medicine. It is a very hard, drought tolerant plant that thrives well on a wide range of soils. The species is native to India and distributed in Sri Lanka, Indonesia, Malaysia, Myanmar and Pakistan. Fruits are generally harvested at immature stage for vegetable purpose, fully ripen fruits are consumed fresh or processed. Traditionally, this fruit is used for making pickles and sometimes candy like products commonly used in ice cream and bakery products as a substitute of cherry. It is a rich source of iron, therefore, in Uttar Pradesh, Uttarakhand and Rajasthan tribal people use for anaemic and pregnant women. It is generally used as live fencing border crops around field boundaries of fruit orchards, protecting crops from wild animals and also provides good returns to the farmers. *Karonda* fruits are the potential source of natural antioxidants. Recent studies have shown that many flavonoids and related polyphenols contribute significantly to the total antioxidant activity of many plant extracts. Though, this fruit is traditionally used as a rich source of minerals, vitamins and phytochemicals but till date no systematic research has been carried out. Based on the above facts, the present study was undertaken on two types of *karonda* (green and white base) to compare the phytochemical changes with reference to antioxidant activity during storage.

The mature fruits of *karonda* were collected from the field of Indian Institute of Farming Systems Research, Modipuram in the month of November, 2013 and 2014. At the time of fruit harvest, physiological parameters such as spade index and leaf area index were also measured with the help of SPAD-502 and canopy analyzer (LAI 2000), respectively. The harvested fruits were washed thoroughly with tap water and dried at room temperature for 2-3 h. The storage studies were conducted with fruits (pinkish-green and pinkish-white, 100 g each) containing 20-25 numbers having average weight of 4-6 g per fruit kept separately in different petri dishes and replicated three times. These fruits were kept in ambient condition ( $18 \pm 2^\circ\text{C}$  &  $40 \pm 5\%$  RH) and analyzed for phytochemical properties, viz., ascorbic acid content, total phenols, flavonoids and antioxidants at two days interval up to eight days in both the years.

Ascorbic acid content was determined by using the 2,6-dichlorophenol indophenol dye method suggested by Freed (4). The amount of ascorbic acid was calculated, and expressed as mg/ 100 g on a fresh weight basis. One gram of both type *karonda* were macerate in a mortar and pestle by using 10 ml methanol three times to extract the phytochemicals. The mixture was centrifuged at 9,000 rpm for 15 min. and the supernatant was decanted into a 50 ml volumetric flask for further chemical analysis. The amount of total phenolic contents in the extracts was determined calorimetrically with the Folin-Ciocalteu reagent, using a slightly modified method of Mansouri *et al.* (9). The results were expressed

\*Corresponding author's E-mail: amitnath2005@gmail.com

as gallic acid equivalents (GAE)/100 g fresh weight. Total flavonoids content of the methanolic extract of *karonda* fruits was determined according to a modified colorimetric method (Zou *et al.*, 13). Briefly, 1.5 ml of fruit extract was taken and 75  $\mu$ l of 5% NaNO<sub>2</sub> solution was added. After 6 min., 150  $\mu$ l of 10% AlCl<sub>3</sub>.6H<sub>2</sub>O was added to the mixture, which was kept at room temperature for 6 min. followed by the addition of 0.5 ml of 1M NaOH and the total volume was made up to 2.5 ml with the addition of deionised water. The resulting solution was mixed well and immediately, the absorbance was measured at 510 nm on a UV-VIS spectrophotometer. For the blank, the extracts were replaced with an equal volume of deionised water. A standard calibration curve was prepared with 0.01, 0.05, 0.1, 0.2, 0.4 and 0.6 mg/ml of quercetin (in deionised water). The total flavonoids content was expressed as the mg equivalents of quercetin (QE) per 100 g fresh weight. The antioxidant activity of the plant extracts and the standard was assessed on the basis of the radical scavenging effect of the stable 1, 1-diphenyl-2-picryl- hydrazyl (DPPH)-free radical activity by modified method (Braca *et al.*, 3). The optical density was recorded and per cent inhibition was calculated (Bors *et al.*, 2). The total antioxidant activity of plant extract was also carried out using FRAP method (Benzie and Strain, 1). Data of 2013 and 2014 for quality parameters were subjected to analysis of variance (ANOVA), mean and standard deviation. Sources of variation were storage time and cultivars. All analysis was performed with a statistical software package SPSS v.11.0 for windows. Values of standard deviation are showed in figures.

At the time of harvest, the physiological parameters such as spade index and leaf area index were recorded. The spade index of *karonda* plants were recorded as 41.57 and 40.12 respectively for pinkish green and pinkish-white, whereas, leaf area index were recorded as 1072.11 and 831.45, respectively for pinkish-green and pinkish-white types.

Ascorbic acid content decreased during the course of storage in both the types of *karonda* (Fig.1). The ascorbic acid content was significantly higher in pinkish-green fruit in comparison of pinkish-white fruits. The ascorbic acid varied from 50.98 to 28.22 mg and 36.86 to 22.15 mg per 100 g for pinkish-green and pinkish-white base *karonda*, respectively during the storage upto 8 days. The ascorbic acid content decreased in pinkish-green and pinkish-white fruits follows the trend of linear equation  $y = -2.6384x + 48.045$  and  $y = -1.88x + 37.864$  with R<sup>2</sup> value 0.92 and 0.96, respectively. A rapid fall in ascorbic acid content (23.32%) was recorded in green fruits as compared to white one upto 2 days during storage (Fig. 1). However, the minimum ascorbic acid

(39.90%) degradation was recorded in white fruits as compared to green fruits upto 8 days during storage. The ascorbic acid content decreased during storage may be due to the enzyme induced biochemical reactions. During storage, oxidizing enzymes like ascorbic acid oxidase, peroxidase, catalase and polyphenol oxidase might help in reducing the ascorbic acid of the fruits (Mapson,10). Whereas, vitamin C or ascorbic acid content in green *karonda* was reported 1.32 mg/g on dry weight basis (Jittawan *et al.*, 8). However, values of ascorbic acid content in *Karanda* were found very close to those values as reported by others (Haque *et al.*, 5).

Further study shows that the total phenols also decreased from 19.73 to 14.48 mg and 15.84 to 14.04 mg GAE per 100 g of green and white *karonda* fruits, respectively during storage. The decrease trend of phenol follow the equation of  $y = -0.7659x + 19.952$  with R<sup>2</sup> value 0.81 and  $y = -0.2405x + 15.858$  with R<sup>2</sup> value 0.97 for green and white fruits, respectively. There was very minimum losses of total phenol content in case of white fruits, whereas 22.76% loss of phenol content was recorded upto 4 days of storage in green fruits. Though, the phenol content of green fruits was higher after harvest as compare to white types but the phenol content in white *karonda* more or less remain constant during 8 days of storage. After losing of 26.61% phenol content in green *karonda* upto 8 days of storage, the phenol content of both the fruits become at par (Fig. 2). It was reported that the levels of total phenolic compounds in the wild fruits varied significantly, from 1.3 to 214 mg gallic acid equivalent (GAE)/g dry weight, where the green based *karonda* contains 1.80 GAE/g dw (Jittawan *et al.*, 8).

It was also found that flavonoid content was not significantly higher in green base than the white base types, but the flavonoids content decreased significantly during the storage of both type of *karonda*




Fig. 1. Changes of Ascorbic acid content in *Carissa carandas* during storage.




Fig. 2. Changes of total phenol content in *Carissa carandas* during storage.




Fig. 3. Changes of total flavonoids content in *Carissa carandas* during storage.

as per the trend of  $y = -0.219x + 5.046$  and  $y = -0.157x + 4.27$ , respectively. The flavonoids content vary from 4.87 to 3.09 mg and 4.21 to 3.01 mg quercetin equivalent mg/g of white and green fruits, respectively. Though, there was a decreasing trend of flavonoids during storage at ambient condition but the magnitude was different. It was found that the flavonoids content decreased by 9% in case of green fruits, whereas the flavonoid content decreased by 15% in case of white *karonda* upto 4 days of storage. After 8 days of storage, the flavonoids content decrease by 36% in green *karonda* as compared to the initial flavonoids content after harvest, whereas it decreased by 28% in white fruits and ultimately the flavonoids content became at par (Fig. 3) at 8 day of storage. Flavonoid content in the methanolic extract of *Carissa carandas* was found to be 2.92 rutin equivalents mg/g of extracts, which is nearly similar to the flavonoid content estimated in both types of *karonda* (Itankar *et al.*, 7). Besides their anti-oxidant activity, flavonoids have demonstrated a wide range of biochemical and pharmacological effects, including anti-inflammatory, anti-viral, anti-allergenic, anti-carcinogenic, anti-aging activity (Hulya, 6), anti-oxidant and anti-allergic effects (Miean and Mohamed, 11).

Different antioxidant compounds may act through different mechanisms and one method alone cannot be utilized to fully evaluate the antioxidant capacity. The antioxidant properties of green and white fruits of *Carissa* was estimated by DPPH and FRAP assays. The scavenging activity of methanol extract of *C. carandas* was significantly higher (89.57%) in case of green fruits compared to white fruits (86.27%). The per cent decrease of scavenging activity follow the trend of  $y = -0.789x + 89.862$  with  $R^2$  value 0.96 and  $y = -0.7605x + 86.6$  with  $R^2 = 0.97$  for green and white fruits, respectively during storage. The

value of percent scavenging activity decrease in similar pattern (Fig. 4) in both types during storage. The ferric reducing ability of plasma (FRAP) assay for antioxidant power was higher (13.93 mg GAE/g) in fresh green fruit extract than white fruits extract (12.91 mg GAE/g). In all the cases the antioxidant properties of green was higher than the white fruited *karonda* but the antioxidant properties were found gradually decreasing during the course of storage as per trend of linear equation  $y = -0.4215x + 13.874$  with  $R^2$  value 0.86 and  $y = -0.5655x + 13.068$  with  $R^2$  value 0.90, respectively. The value of FRAP assay decrease in similar pattern (Fig. 5) in both type of *karonda* fruits during storage. Similar decrease in antioxidant content during storage was reported by (Nath *et al.*, 12) in case of broccoli florets. The decreasing trend of antioxidant properties may be due to the low value of polyphenol content in the sample during storage. There was positive correlation between the polyphenol content and




Fig. 4. Changes of scavenging activity of *Carissa carandas* during storage.




Fig. 5. Changes of ferric reducing ability of plasma (FRAP) in *Carissa carandas* during storage.

antioxidant activity of the fruit extract. Jittawan *et al.* (8) reported the variations in anti-oxidant activity values may be expected if multiple samples of each species had been harvested from different conditions of cultivation, location, ripening stage and season.

Green fruits of *Carissa* are enriched with antioxidant phytochemicals as compared to the white fruits. Though the market value of the white fruits of *Carissa carandas* is higher as compared to the green one, but from the nutritive value the green fruits of *Carissa carandas* is better. Therefore, green fruited *Carissa* may be explored as nutritive fruit for health conscious people since they have high ascorbic acid, phenol and flavonoids in addition to higher antioxidant properties.

## REFERENCES

1. Benzie, I.F. and Strain, J.J. 1999. Ferric reducing/ antioxidant power assay: direct measure of total antioxidant activity of biological fluids and modified version for simultaneous measurement of total antioxidant power and ascorbic acid concentration. *Methods Enzym.* **299**: 15-27.
2. Bors, W., Saran, M. and Elstner, E.F. 1992. Screening for plant anti-oxidants. *In: Modern Methods of Plant Analysis-Plant Toxin Analysis-New Series.* Linskens, H.F., Jackson, J.F. (Eds.), **13**: 277-95.

3. Braca, A., Sortino, C., Politi, M., Morelli, I. and Mendez, J. 2002. Anti-oxidant activity of flavonoids from *Licania licaniaeflora*. *J. Ethnopharm.* **79**: 379-81.
4. Freed, M. 1966. *Method of Vitamin Assay*, New York, Inter Science Pub. Inc.
5. Haque, M.N., Saha, B.K., Karim, M.R. and Bhuiyan, M.N.H. 2009. Evaluation of nutritional and physico-chemical properties of several selected fruits in Bangladesh. *Bangladesh J. Sci. Indus. Res.* **44**: 353-58.
6. Hulya, O.H. 2007. Total antioxidant activities, phenolics, anthocyanins, polyphenols oxidase activities of selected red grape cultivars and their correlations. *Scientia Hort.* **111**: 235-41.
7. Itankar, P.R., Lokhande, S.J., Verma, P.R., Arora, S.K., Sahu, R.A. and Patil, A.T. 2011. Antidiabetic potential of unripe *Carissa carandas* Linn. fruit extract. *J. Ethnopharm.* **135**:430-33
8. Jittawan, K. Sirithon S. and Naret, M. 2011. Phytochemicals, vitamin C and sugar content of Thai wild fruits. *Food Chem.* **612**: 972-81.
9. Mansouri, A., Embarek, G., Kokkalou, E. and Kefalas, P. 2005. Phenolic profile and antioxidant activity of the Algerian ripe date palm fruit (*Phoenix dactylifera*). *Food Chem.* **89**: 411-20.
10. Mapson, C.W. 1970. Vitamins in fruits: Stability of L-ascorbic acid. *In: Biochemistry of Fruits and their Products*, Academic Press, London, pp. 376-87.
11. Miean, H.K. and Mohamed, S. 2001. Flavonoid (myricetin, quercetin, kaempferol, luteolin, and apigenin) content of edible tropical plants. *J. Agri. Food Chem.* **49**: 3106-12.
12. Nath, A., Bagchi, B., Misra, L.K. and Bidyut, D.C. 2011. Changes in post-harvest phytochemical qualities of broccoli florets during ambient and refrigerated storage. *Food Chem.* **127**: 1510-14.
13. Zou, Y., Lu, Y. and Wei, D. 2004. Antioxidant activity of flavonoid-rich extracts of *Hypericum perforatum* L. *in vitro*. *J. Agri. Food Chem.* **52**: 5032-39.

Received : February, 2016; Revised : November, 2016;  
Accepted : November, 2016

## Hort News

The Horticultural Society of India, New Delhi organized 7<sup>th</sup> Indian Horticulture Congress-2016 (an International Meet) on the title "**Doubling Farmers Income through Horticulture**" at Dr B.P. Pal Auditorium, Pusa campus, Indian Agricultural Research Institute, New Delhi from 15-18<sup>th</sup> November, 2016 to commemorate its Platinum Jubilee. It also had its Executive Council meeting, Foundation Day Lecture and Annual General Body Meeting during the congress. Brief highlights of the events are given hereunder;

### 7<sup>th</sup> Indian Horticulture Congress-2016

The Horticultural Society of India, New Delhi in association with ICAR-Indian Agricultural Research Institute, New Delhi organised the 7<sup>th</sup> Indian Horticulture Congress on the theme 'Doubling Farmers' Income through Horticulture' from November 15-18<sup>th</sup>, 2016, at New Delhi. The congress was a grand success. It brought together over 750 researchers, students, extension workers, State Govt. officers, policy makers, corporate & agri-business representatives, progressive farmers, orchardists, entrepreneurs, NGOs, SHG representatives and development practitioners. It also attracted participation from USA, Austria, Nepal, Vietnam etc. During the four day event, deliberations were made to assess the overall strength of technologies developed and advancements made in different sectors in Horticulture with special reference to the main theme of the congress. The focus of entire deliberations made in 21 Technical Sessions was improving the farm incomes, through different interventions.

The Congress was inaugurated on 15<sup>th</sup> Nov, 2016 by the Chief Guest, Sh. S.S. Ahloowalia, Hon'ble Minister for State for Agriculture and Farmers Welfare, GoI, in the august presence of Professor M.S. Swaminathan, Eminent Agricultural Scientist and Chairman of the National Advisory Committee; Dr Gurbachan Singh, Chairman-ASRB, Dr A.K. Singh, Deputy Director General (Agril. Extension & Horticulture Sci.), ICAR; and Dr K.L. Chadha, President HSI and Chairman Organizing Committee, Dr (Mrs.) Ravinder Kaur, Director, IARI, New Delhi, Dr Manmohan Attavar, Chairman, Indo-American Hybrids Seeds, Bengaluru and Dr Pritam Kalia, Organizing Secretary.

The programme was inaugurated by lighting of the ceremonial lamp by dignitaries. Thereafter, Dr K.L. Chadha, President, HSI and Chairman, Organizing Committee welcomed the dignitaries and delegates on the occasion of Platinum Jubilee Celebration of the Horticultural Society of India and the 7<sup>th</sup> Indian Horticultural Congress-2016. During the welcome address, he highlighted the historical events pertaining to the journey of the society and its activities since its

inception in pre-independence era during 1942. He also spoke about the several national and international seminars, conferences, brainstorming sessions, etc. organised in the past and the genesis of holding the Indian Horticulture Congresses. He further informed about the different awards and fellowships conferred to recognise excellence in research and development in different fields of Horticulture. The President HSI, also informed about the 7<sup>th</sup> IHC and the events to be held in 21 Technical Sessions in the form of Lead, and Invited/ Oral presentations and Poster presentations in five sessions on different sub-themes. He also informed that two dedicated sessions, each on Pro-Horticulture Developmental Policy needs; Opportunities for Youth through Horticultural Enterprises and one on Horticulture-led Development in North East Region of the country. Besides, to encourage the young researchers, two Technical Sessions covering 51 oral presentations were also planned.

All the Guests of Honour spoke on the occasion. The Inaugural Address was delivered by the Chief Guest, Sh. S.S. Ahloowalia, Hon'ble Minister for State for Agriculture and Farmers Welfare, Govt. of India, who highlighted the problems of Horticulture production and problems of fruit growers regarding issues concerning marketing, storage, value-addition and trade. He also highlighted the need for tapping potential of horticulture led development in NE regions so that farmers of the region could also benefit from production of high quality produce/ value-added products. Prof M.S. Swaminathan spoke about his association with Horticulture R&D in the country. He empathetically spoke about the role of horticulture in bringing about nutritional and livelihood security in the country. He also emphasised promotion of indigenous fruit and vegetable crops for improving nutrition as well as earnings of the farmers. He applauded the role of HSI in flagging R&D issues by holding conferences and congresses on relevant issues for the benefit of the end users.

Dr Gurbachan Singh, Chairman, ASRB spoke about the role of horticultural crops in greening of grey and marginal areas in the country. He spoke about the climate resilience of horticultural crops and particularly under moisture and abiotic stress situations.

Dr Pritam Kalia, Organizing Secretary and Secretary, HSI presented the vote of thanks.

### HSI Honorary Fellowships and Awards

On this occasion, the Society honoured several scientists, entrepreneurs, and progressive farmers. Honorary Fellowships of HSI were conferred to Sh. Suresh Aggarwal, Chairman, Beej Sheetal Seeds