



Effect of saline water irrigations on physiological, biochemical and yield attributes of dual purpose pearl millet (*Pennisetum glaucum*) varieties

GOVIND MAKARANA¹, ASHWANI KUMAR², R K YADAV³, RAKESH KUMAR⁴, POOJA G. SONI⁵, CHARU LATA⁶ and PARVENDER SHEORAN⁷

ICAR-National Dairy Research Institute, Karnal 132 001

Received: 09 May 2018; Accepted: 31 October 2018

ABSTRACT

A split-plot experiment with four replicates on performance of dual purpose pearl millet (*Pennisetum glaucum* L.) under different saline irrigation water was conducted during *kharif* 2015 at ICAR-CSSRI experimental farm, Nain, Panipat. The experiment was conducted with 4 main-plot treatment consisting of different levels of saline irrigation water [Normal (~0.6 dS/m) and three levels of saline irrigation (EC_{iw} 3, 6 and 9 dS m⁻¹)] and two sub-plot treatments of varieties [ICMV-15111 and AVKB-19]. AVKB 19 variety was found robust and more stable than ICMV 15111 in terms of physiological attributes (relative water content, membrane injury, and chlorophyll content) at all three periodic observations taken at 50 days after sowing (at 1st cut), 30 days and 60 days after 1st cut. Among biochemical attributes, total soluble sugars, proline, epicuticular wax content increased with increasing stress levels where as protein content decreased with increasing level of salinity stress. AVKB 19 variety accumulated significantly higher osmolyte content than ICMV 15111. Irrespective of salinity treatments Na⁺ content (shoot and root) was recorded gradually high with ICMV 15111 than AVKB 19 at all study stages. AVKB 19 variety presented significantly higher shoot and root K⁺ content than ICMV 15111. Data on green fodder yield showed significant differences between two cultivars across different saline water levels. The AVKB-19 variety produced significantly higher mean green fodder yield (27.96 t/ha) as compared to ICMV-15111 (25.51 t/ha). The AVKB-19 resulted into significantly maximum DM yield at the 1st as well as in 2nd cut (5.67 and 5.09 t/ha) over ICMV-15111. AVKB-19 variety produced significantly higher (16.26%) grain yield of 1.93 t/ha as compared to 1.66 t/ha in ICMV-15111. The magnitude of reduction (%) with increasing salinity over good quality irrigation water was observed 10.14, 20.74 and 37.33 with 3.0, 6.0 and 9.0 dS/m salinity water, respectively. Our results concluded that AVKB-19 variety of pearl millet proved superior and may be adapted as a choice for getting higher grain as well as green fodder yield with better physiological and biochemical responses under saline environment. (Exceeding 250 word limit)

Key words: Pearl millet, Physiological attributes Saline water and Yield

Presence of salt affected soils on the earth surface is as old as the history of mankind (Sharma and Choudhari 2012). It has now become a very serious threat for crop production (Ashraf and Wu 1994; Rengasamy 2006) particularly in arid and semi-arid regions, which constitute about one third of the world's land surface. At present about 6.73 million-hectare (Mha) salt affected soils exist in India. Out of total salt affected soils 2.96 Mha are saline and remaining

3.77 Mha are characterized as sodic soils (Sharma and Singh 2015). Most crops are sensitive to salinity caused by high concentrations of salts in the soil. Due to increased population pressure the inter sector competition has been raised, this has resulted into declined availability of fresh water supplies to agriculture particularly in Asian countries. The scarcity of water in association with high salinity is now major problem hindering crop production in these salty lands. Poor quality groundwater extensively occupied (32-84%) in arid and semi-arid parts of India (CGWB 1997) and its indiscriminate use poses a serious threat to sustainability of the natural resources and environment. States like Rajasthan and Haryana do not have sufficient surface water resources for meeting irrigation water requirement and thereby depend on poor quality ground water which is 68% and 55%, respectively in the two states (Lal et al. 2008). The farmers of these states have been over exploiting groundwater for supplementing the limited surface water resources. This over-mining of groundwater is causing decline in water

¹Govind Makarana, Ph D Scholar (e mail: makarana17@gmail.com), ICAR-National Dairy Research Institute, Karnal 132 001, ²Ashwani Kumar, Scientist (e mail: Ashwani.Kumar1@icar.gov.in), ICAR-Central Soil Salinity Research Institute, Karnal-132 001, ³R K Yadav, Principal Scientist (e mail: rk.yadav@icar.gov.in), ⁴Rakesh Kumar, Principal Scientist (e mail: drdudi_rk@rediffmail.com), ⁵P G Soni, Ph D Scholar (e mail: 15ppoojagupta@gmail.com), ⁶Charu Lata, Senior Research Fellow (e mail: charusharmabiotech@gmail.com), ⁷Parvender Sheoran, Principal Scientist (e mail: sheoran76@rediffmail.com)

table at alarming rates in good quality groundwater zones and causing quality deterioration further in these areas. Soil and water salinity cause several physiological disorders in plants, connected with the abnormal concentration of ions in the rhizospheric environment; these can range from a cytotoxic and denaturing effect of the ions themselves (Bernstein 1975) to osmotic stress (Greenway and Munns 1980; Yeo 1983) and alteration of the ion uptake balance (Rains 1972; Flowers and Lauchli 1983). Consequently, it has become imperative to search for suitable crop/genotype alternatives and develop ecologically sustainable and economically sound production systems that can use poor quality water and withstand drought on saline lands.

In this context, Pearl millet (*Pennisetum glaucum* L.) is a promising dual purpose, short duration, quick growing crop with good salinity tolerant characteristics, therefore has an advantage over others cultivated fodder in salt affected areas. Pearl millet has been reported to have high tolerance to salinity and drought thus it can serve as an important crop to ensure good quality fodder for animals in the arid and semi-arid regions of India and elsewhere in the world under similar agro ecologies (Kulkarni et al. 2006; Patel et al. 2008). Several researchers documented that Pearl millet showed minimum yield reduction under saline environment, thus demonstrating its tolerant nature towards salinity. Therefore, present study was conductively undertaken to access the physiological, biochemical and yield responses of pearl millet varieties. By understanding the traits affecting the production potential will help to deliver a suitable alternative genotype that could be helpful to use the poor quality water without compromising the soil health and yield.

MATERIALS AND METHODS

The present study was carried out at ICAR-CSSRI experimental farm, Nain (29°19' N, 76°47' E and 230.5 m above the mean sea level), Panipat, Haryana, India. The climate of the area is semi-arid, with a mean annual rainfall of 678 mm (70-80% of which received during July-September) with the mean annual USWB-open pan evaporation (ET_p) of 1598 mm. The meteorological phenomenon occurred during study period (July-November) are presented in Table 1. The Initial Soil Status of experimental site and salinity build-up in each plot after harvest is presented in Table 2.

The experiment was designed in split-plot arrangements with four replications. The experiment was conducted with

Table 2: Initial Soil Status and salinity build-up in each plot after harvest

Initial Soil Status	Salinity levels of each plots depth wise (cm) after harvest (ECe) (dS m ⁻¹)			
	Sandy loam	Control 1 (0.6)	0-15	5.31
Texture			15-30	5.32
Walkley-Black C (%)	0.3		0-15	6.17
Ece(dS m-1)	6.65	3	15-30	6.18
pH	8.3		0-15	6.83
KMnO4 Oxidizable N (kg/ha)	130.4	6	15-30	6.85
0.5 M NaHCO3 extractable P (kg/ha)	11.6		0-15	7.58
1 N N H 4 O A C extractable K (kg/ha)	248.4	9	15-30	7.61

4 main-plot treatment consisting of levels of irrigation water [normal (~0.6 dS m⁻¹) and three levels of saline irrigations (ECiw 3, 6 and 9 dS m⁻¹)] and two sub-plot treatments of varieties [ICMV-15111 and AVKB-19]. To supplement the total rainfall and maintain of 1.2 ID/CPE ratio 240 mm water was applied through irrigation. To fulfil this purpose check basin method was adopted. Each experimental unit consisted of 4.5 m × 4.5 m plots. The field was deep ploughed by chisel plough to break the hard pan below the plough layer before start of the experiment. The pearl millet cv. AVKB-19 and ICMV-15111 were sown with a seed rate of 12 kg ha⁻¹ during second fortnight of July in 2015 with a row spacing of 30 cm and plant to plant distance at 10 cm. A common dose of nutrients amounting 120 kg N + 60 kg P₂O₅ + 40 kg K₂O were applied in all treatments through urea [CO(NH₂)₂], di-ammonium phosphate((NH₄)₂HPO₄) and sulphate of potash (K₂SO₄). The 1/3rd N and whole P₂O₅ and K₂O was applied as basal, while remaining 2/3rd N was top dressed as urea in two equal splits at 1st cutting and 30 days after 1st cutting. In view of best weed management, two hands weeding at 20 DAS and 30 days after 1st cutting was done to control weeds. The first cut of crop was taken at (50 DAS) at the 8-10 cm above the ground level. Then the crop was left for grain production. The final cut was taken at ~110 DAS for grain purpose. Three observations for physiological parameters were studied, at 50 DAS (1st cut); second observation at 30 days after 1st

Table1: Meteorological observations during study period

Month	Temperature (0C)		Relative humidity (%)		Total Rainfall (mm)	Total Evaporation (mm)
	Mean Maximum	Mean Minimum	MeanMax.	Mean Min.		
20-31 July	32.92	25.4	86.08	69.67	11.4	71.2
1-31 August	32.68	25.1	92.97	73.2	70.2	103.6
1-30 September	33.51	22.8	91.5	59.2	46.8	106.5
1-31 October	32.15	17.77	91.67	46.3	6.4	90.7
1-15 November	27.73	14.19	91.56	47.06	9.2	30.3

cut (80 DAS), and third /final observation at 60 days after 1st cut (110 DAS) from representative (tagged) plants from each plot. *Relative water content* (RWC) was measured following the procedure described by Weatherley (1950). Third fully expanded leaf from the top of the plants sampled was detached from the shoots between 9:00 to 10:00 AM, quickly sealed in humified polythene bags, and transported on ice to the laboratory. Sand was removed with the help of a soft brush. Then the leaves were weighed immediately to take their fresh weight. The leaves were kept in petri dishes full of distilled water for 3 h.

After that the leaves (fully turgid) were weighed again and then kept in oven at 85°C for 72 h till a constant dry weight. These three weights were used to calculate RWC (%) of leaves according to the following formula.

$$\text{RWC (\%)} = \frac{\text{fresh weight} - \text{Dry weight}}{\text{Turgid weight} - \text{Dry weight}} \times 100$$

Membrane injury was estimated according to the method of Dionisio-Sese and Tobita (1998). The leaf discs of 1 cm diameter were kept in 20 ml vials containing 10 ml deionized water at 25°C. After five hours, the electrical conductivity (EC) of the surrounding solution was measured by the Water Analysis Kit (Naina, India Ltd., NDC 732) and designated as ECa. Then the samples were kept in boiling water bath for 50 min to achieve total killing of the tissue. After cooling, the EC of the solution was again measured and designated as ECb. The relative stress injury (RSI)/ Membrane stability index was calculated as follows and expressed in percentage:

$$\text{RSI (\%)} = \frac{\text{ECa}}{\text{ECa} + \text{ECb}} \times 100$$

The *chlorophyll content* was determined using DMSO (Dimethyl sulphoxide) as described by Hiscox and Israelstam (1979) using dimethyl sulfoxide (DMSO). Third leaf from the top of plant was detached and weighed 200 mg of this leaf material was transferred into a test tube containing 5 ml of DMSO. The test tube was then placed into oven at 60°C for about 2 h or more (if required) to facilitate the extraction of pigment. After 2 h and attaining the room temperature, the absorption was recorded at 645 and 665 nm on spectrophotometer. Absorbance at respective wavelengths was also recorded in DMSO and used as blank. Calculations for different pigments were made as per the procedure described by Welburn (1994).

$$\text{Chl 'a' (\mu g/ml)} = 12.19 A_{665} - 3.45 A_{645}$$

$$\text{Chl 'b' (\mu g/ml)} = 21.99 A_{645} - 3.32 A_{665}$$

$$\text{Total chlorophyll} = \text{Chl 'a'} + \text{Chl 'b'}$$

Proline content was estimated using the method of Bates *et al.* (1973). Three hundred mg of leaves was homogenized in 5 ml of 3 per cent sulphosalicylic acid and then centrifuged at 5000 rpm for 15 minutes and supernatant was taken. Two ml of supernatant was taken and 2.0 ml reagent acid ninhydrin + 2.0 ml acetic acid was added. This mixture was then kept in boiling water bath for 1 hr at 100°C and thereafter reaction was terminated

by keeping tubes in ice-bath. Then 4.0 ml of toluene was added. After vigorous shaking, the upper organic phase was taken after attainment of room temperature and absorbance was recorded at 520 nm by using toluene as blank. A standard curve was prepared by using graded concentration of proline in 3% sulphosalicylic acid. The proline content was expressed as $\mu\text{g g}^{-1}$ FW.

Colorimetric method (Ebercon *et al.* 1977) was used for *epicuticular wax* deposition. Plant leaf blade samples, were taken from different treatment wise for wax analysis. The colorimetric method is based on the color change produced due to the reaction of wax with acidic $\text{K}_2\text{Cr}_2\text{O}_7$ reagent. The reagent was prepared by mixing 40 ml deionized water and 20 g powdered potassium bichromate. The resulting slurry was mixed vigorously with 1 litre conc. sulfuric acid and heated (below boiling point) until clear solution was obtained. The individual samples of pearl millet were immersed in 15 ml redistilled chloroform for 15 second. The extract filtered and evaporated on a boiling water bath until smell of chloroform disappeared. After cooling, 12 ml of deionized water was added. Several minutes were allowed for colour development and cooling and then optical density of the sample was read at 590 nm.

Total soluble carbohydrate was determined with the method of Yemm and Willis (1954). The extraction of total soluble carbohydrate was done according to the method of Barnett and Naylor (1966). Fifty mg dry samples of leaves were homogenized separately with 5 ml of 80 per cent (v/v) ethanol. Like this, third extraction was done with 60 per cent (v/v) ethanol. The supernatant from different extractions was pooled and evaporated to dryness in a boiling water bath. Then total volume of extract was made to 10 ml with distilled water. The extract so obtained was used for estimation of total soluble carbohydrate and free amino acids. The residue was used for total soluble protein extraction. Anthrone reagent was prepared by dissolving 0.2 g anthrone in 100 ml concentrated H_2SO_4 . 0.5 ml aliquot of the extract was taken in a test tube and mixed with 4.0 ml of the anthrone reagent. The mixture was heated in boiling water bath for 10 minutes. After cooling, absorbance was recorded at 620 nm. Glucose (10-100 μg) was taken as standard.

Total soluble proteins were determined by Bradford (1976) method using supernatant. Dissolve 100 mg of Coomassie Brilliant Blue G-250 (CBBG-250) in 50 ml of 95% ethanol. Add to this solution 100 ml 85% (w/v) phosphoric acid and make final volume 200 ml with double distilled water. Filter through whatman filter paper No. 1 (if any ppt. occurs), diluted to a final volume of 1 litre and stored at 4°C in amber colour bottle. To 25 μl of the supernatant, 5 ml of the coomassie Brilliant Blue G-250 reagent was added and mixed either by inversion or vortexing. The optical density (O.D.) was measured at 595 nm after 2 minutes and before 1hr against reagent blank. Standard curve was prepared using graded concentration of bovine serum albumin.

For Na^+ and K^+ content, 100 mg of oven dried and well ground plant material was digested with 10 ml of HNO_3

: HClO₄ (3:1) di-acid mixture and readings were taken with flame photometer (PFP7, Jenway, Bibby Scientific, UK) using standard NaCl and KCl. Yields of 1x1 m⁻² was recorded and converted in per hectare. All data recorded were analyzed with the help of analysis of variance (ANOVA) technique using the SAS (Version 9.3, SAS Institute Inc., Cary, NC, USA). The least significant test was used to decipher the main and interaction effects of treatments at 5% level of significance (P<0.05).

RESULTS AND DISCUSSION

Perusal of the physiological and biochemical data indicated significant differences among the varieties and among the treatments with respect to different parameters.

Relative water content (RWC) (%): It is the appropriate measure of plant water status in terms of the physiological consequence of cellular water deficit. Statistical analysis of data indicated that salt stressed pearl millet plants exhibited significant differences for RWC through increasing salinity of irrigation water (Fig 1). The decrease in relative water content (RWC) could be attributed to early symptoms of stress conditions. The maximum relative water content (%) was observed from good quality irrigation water (87.7, 86.135 and 79.32 at 50 DAS, 30 DA 1st cut and 60 DA 1st cut respectively) and the minimum (68.87, 61.65 and 58.04 at 50 DAS, 30 DA 1st cut and 60 DA 1st cut respectively) from irrigation water having 9.0 EC. The minimum significant percentage reduction 10.74, 11.19 and 5.29% was observed at 3.0 dS/m in comparison to good quality water at all periodic observations, whereas the maximum significant was noticed 21.08, 28.42 and 26.82% from irrigation water having 9.0 EC in comparison to good quality water at all sequential periodic observations. Irrespective of water quality, the mean maximum performance with respect to this parameter was found from genotype AVKB 19 at all sequential periodic observations than ICMV 15111 but the difference between genotypes was non-significant at all the three stages. This reduction of RWC in stressed plants might be associated with a decrease in plant vigour (Kumar et al. 2016). High salt concentration in root zone causes osmotic stress which restricts water absorption by the plants and causes cellular dehydration, seems to be primarily responsible for decrease in RWC (Greenway and Munns 1980). These results agree with the findings of Netondo et al. (2004) in sorghum, and Vijayalakshmi et al. (2012) in pearl millet.

Total Chlorophyll content (µg/ml of FW): Chlorophyll concentration has been known as an index for evaluation of source (imparts lush green colour to the fodder, improves radiation absorption and quality), therefore decrease in concentration can be considered as a stomata non-limiting factor under stress conditions. The

observations with respect to chlorophyll content as affected by different irrigation water salinity levels are presented in Figure 2. Mean performance of the genotype AVKB 19 (51.91µg/ml and 39.68 µg/ml at 30 DA 1st cut and 60 DA 1st cut, respectively) were found significantly higher than that ICMV 15111 (45.87 and 37.35 µg/ml) at 30 DA 1st cut and 60 DA 1st cut, respectively), but in case of 1st observation at 50 DAS, there was statistically no difference among both genotypes. The significantly maximum chlorophyll content was found with good quality irrigation water whereas minimum chlorophyll content recorded with EC 9.0 dS/m of irrigation water. Similarly significant variation were noticed by every increasing levels of salinity at 50 DAS and 60 DA 1st cut. The effects of salt stress on chlorophyll degradation, presumably due to increased activity of the enzyme chlorophyllase (Kumar et al. 2016), were characterized as the yellowing of leaves which failed to produce the optimum amounts of photosynthates leading to reduced plant growth and vigour.

Membrane injury index (MI) (%): Electrolyte leakage (an indicator of membrane damage) increased with increase in saline irrigation. Leaves of pearl millet plants irrigated with good quality water exhibited lowest (8.18, 8.56 and




Fig 1 Effect of saline water irrigations on relative water content (%) in dual purpose pearl millet varieties




Fig 2 Effect of saline water irrigations on total chlorophyll concentration (µg/ml) in dual purpose pearl millet varieties

10.42% at 50 DAS, 30DA and 60 DA 1st cut, respectively) membrane injury than all higher level of saline water irrigation (Fig 3).

The maximum MI (24.11, 26.53 and 35.8%) was noticed from irrigation water having 9.0 EC at respective sequential observations. While investigating the genotypic potential for membrane stability, AVKB 19 was found hardier and more stable than ICMV 15111 because it had lower injury index at all three periodic observations taken at 50 DAS, 30 and 60 DA 1st cut. Salt stressed plants exhibit damage of lipid membranes which often results in increased cell permeability and electrolyte leakage from cells (Kumar et al. 2015).

Biochemical attributes: The accumulation of compatible solutes such as soluble sugars, proline (key osmolytes that contribute to osmotic adjustment) etc. which are non-toxic at higher concentrations to cytoplasmic functions allow additional water uptake from the environment and turgor maintenance (Sairam et al. 2002; Misra and Gupta 2005).

The accumulation of soluble carbohydrates in plants has been widely reported as a response to salinity or drought, despite a significant decrease in net CO₂ assimilation rate (Murakezy et al. 2003).

Total soluble sugars (TSS) (mg g⁻¹ DW): The mean minimum TSS was recorded with good quality water at all three periodic observations taken at 50 DAS (7.22 mg g⁻¹ DW), 30 DA 1st cut (10.89 mg g⁻¹ DW) and 60 (13.72 mg g⁻¹ DW) DA 1st cut, whereas the mean maximum TSS was recorded with irrigation water having 9.0 EC at 50 DAS (22.43 mg g⁻¹ DW), 30 DA 1st cut (25.41 mg g⁻¹ DW) and 60 (38.35 mg g⁻¹ DW) DA 1st cut (Table 1). The significant increase in TSS was found at all three sequential periodic observations from 9.0 EC of irrigation water over good quality water. When comparing the varietal difference for TSS observations at 50 DAS and 60 DA 1st cut ICMV 15111 found to contain 13.68 mg g⁻¹ DW and 30.91 mg g⁻¹ DW TSS being significantly higher than AVKB 19 having 10.56 mg g⁻¹ DW and 20.46 mg g⁻¹ DW.

Sugars are the source of energy and carbons needed for adaptive or defensive responses to stresses. The higher sugar content in the extreme salinity levels as an additional mechanism to prevent salt injury (Kumar et al. 2016; 2017). The changes in the carbohydrate are regulated primarily by K⁺ and Cl⁻ ions. Similarly increasing sugar content at higher salinities has been reported in several halophytes (Prado et al. 2000; Ashraf and Harris 2004).

Proline content (µg g⁻¹ FW): is a potent osmoregulator molecule and counteracts the

adverse effects of toxic salt ions in cell vacuoles, contributes to membrane stability and mitigates the effect of NaCl on cell disruption. The mean maximum proline content was recorded from 9.0 EC of irrigation water 3.78, 3.59 and 4.47 at 50 DAS, 30 and 60 DA 1st cut, respectively in comparison to minimum of 0.92, 1.2 and 1.58 at 50 DAS, 30 and 60 DA 1st cut, respectively from good quality irrigation water. The




Fig 3 Effect of saline water irrigations on membrane injury (%) in dual purpose pearl millet varieties




Fig 4 Effect of saline water irrigation on total soluble sugar content (mg g⁻¹ DW) in dual purpose pearl millet varieties




Fig 5 Effect of saline water irrigation on proline content (µg g⁻¹ FW) in dual purpose pearl millet varieties

increase in proline content was found non-significant up to 6.0 EC at 50 DAS and up to 3.0 EC of irrigation water at 30 DA 1st cut stage, whereas the increase in percentage was found significant from every increasing level of salinity at 60 DA 1st cut stage (figure 5). When comparing the varietal potential against irrigation water salinity, variety ICMV 15111 found containing significantly higher proline content than AVKB 19 at 30 DA 1st cut stage, but non-significantly higher at 50 DAS. In case of 60 DA 1st cut stage, variety AVKB 19 was found to contain significantly higher proline content than variety ICMV 15111. Accumulation of these osmolytes can also improve plant tolerance by protecting and stabilizing membranes and enzymes during stress conditions (Ashraf and Harris 2004; Singh et al. 2015; Lata et al. 2017).

Total soluble protein content(TSP)(mg g⁻¹): The proteins that accumulate under stress conditions may provide a storage form of nitrogen that is re-utilized in post-stress recovery and also play a role in osmotic adjustment. The mean maximum total soluble protein content (mg g⁻¹) from good quality water with 15.26, 29.68 and 37.84 at 50 DAS, 30 and 60 DA 1st cut, respectively, whereas mean minimum with 9.0 EC of irrigation water containing 5.99, 11.47 and 24.52 at 50 DAS, 30 and 60 DA 1st cut,

respectively (Table 3). The reduction in TSP was found statistically significant from each increasing salinity level of irrigation water over good quality water irrigation at all three periodic observations. When comparing cultivars responses for total soluble protein content, AVKB 19 found slightly high than ICMV 15111 at all three observations. Several researchers showed in earlier findings that soluble protein contents of leaves decreased in response to salinity (Parida et al. 2002; Wang et al. 2000; Agastian et al. 2000; Kumar et al. 2016; 2017).

Epicuticular wax load (EWL) (mg g⁻¹ DW): It is an effective component of abiotic stresses resistance (avoidance mechanism) in pearl millet. A varying response in EW accumulation was seen among the genotypes (Table 4). When comparing the genotypic potential for epicuticular wax load, ICMV 15111 was seen to accumulate epicuticular wax in significantly very high amounts (2.58 mg g⁻¹ DW, 3.3 mg g⁻¹ DW and 3.76 mg g⁻¹ DW at 50 DAS, 30 and 60 DA 1st cut, respectively) as compared to AVKB 19 with 1.72 mg g⁻¹ DW, 2.91 mg g⁻¹ DW, 2.63 mg g⁻¹ DW at 50 DAS, 30 and 60 DA 1st cut respectively. The mean minimum quantity of EW was found when plants are irrigated with good quality water (0.87 mg g⁻¹ DW, 1.29 mg g⁻¹ DW and 1.68 mg g⁻¹ DW at 50 DAS, 30 and 60 DA 1st cut, respectively, whereas

Table 3: Effect of saline water irrigation on total soluble protein content (mg g⁻¹ DW) in dual purpose pearl millet varieties

Irrigation water quality (dS/m)/Varieties	Total soluble protein content (mg g ⁻¹ DW) at 1st cut (50DAS)			Total soluble protein content (mg g ⁻¹ DW) 30 DA 1st cut			Total soluble protein content (mg g ⁻¹ DW) at grain harvest stage					
	ICMV 15111	AVKB 19	Mean	ICMV 15111	AVKB 19	Mean	ICMV 15111	AVKB 19	Mean			
Control (0.6)	14.43	16.10	15.26	28.38	30.98	29.68	34.94	40.75	37.84			
3	10.32	11.19	10.75	22.40	22.29	22.34	32.44	35.00	33.72			
6	7.71	7.08	7.4	19.77	17.49	18.63	28.55	29.98	29.26			
9	5.36	6.63	5.99	11.56	11.38	11.47	25.57	23.47	24.52			
Mean	9.46	10.25		20.53	20.54		30.37B	32.3A				
Factor	T	V	T×V	V×T	T	V	T×V	V×T	T	V	T×V	V×T
CD (P=0.05)	2.74	NS	NS	NS	2.28	NS	NS	NS	3.94	1.63	3.26	4.57

Table 4 Effect of saline water irrigation on epicuticular wax content (mg g⁻¹ DW) in dual purpose pearl millet varieties.

Irrigation water quality (dS/m)/Varieties	Epicuticular wax content at 1st cut (50DAS)			Epicuticular wax 30 DA 1st cut			Epicuticular wax content at 60 DA 1st cut					
	ICMV 15111	AVKB 19	Mean	ICMV 15111	AVKB 19	Mean	ICMV 15111	AVKB 19	Mean			
Control (0.6)	1.12	0.63	0.87	1.31	1.27	1.29	1.3	2.06	1.68			
3	2.08	1.08	1.58	2.66	2.05	2.35	3.24	1.96	2.6			
6	2.80	1.53	2.17	4.31	3.16	3.73	4.64	2.95	3.8			
9	3.56	3.63	3.99	4.90	5.15	5.03	5.85	4.34	5.1			
Mean	2.58	1.72		3.3	2.91		3.76	2.63				
Factor	T	V	T×V	V×T	T	V	T×V	V×T	T	V	T×V	V×T
CD (P=0.05)	0.75	0.75	NS	NS	0.47	0.36	0.71	0.69	0.47	0.36	0.71	0.69

*T = Treatments (Irrigation at ASM Level), V = Varieties, T × V = Treatments at the same level of Varieties and V × T = Varieties at same level of Treatments, CD-Critical difference, DAS-Days after sowing

the mean maximum amount was obtained ($3.99 \text{ mg g}^{-1} \text{ DW}$, $5.03 \text{ mg g}^{-1} \text{ DW}$ and $5.1 \text{ mg g}^{-1} \text{ DW}$ at 50 DAS, 30 and 60 DA 1st cut respectively). Our findings have shown that with of progress of time under stress pearl millet showed higher deposition of EW because under stress conditions, stomata got closed leading to decreased cuticular permeability of water loss with higher deposition of EW (Blum 1988). Kumar et al. 2016 also reported increased EW load with salt and sodic stress in halophytes.

Ionic relations

Sodium content (% DW): Perusal of data (Figure 6) on Na^+ content of shoots indicated that Na^+ content significantly increased with every increasing level of salinity at all periodic observations, except at 1st with good quality water treatment. The maximum Na^+ concentration was recorded with 9.0 EC of irrigation water (0.21, 0.64 and 0.94) whereas minimum with plot subjected to good quality water (0.06, 0.16 and 0.18) at 1st, 2nd and 3rd observation respectively. Similarly, the minimum root Na^+ was recorded with good quality water treatment containing (0.78, 2.9 and 3.6) and maximum recorded with irrigation water treatment of 9.0 EC (0.78, 2.9 and 3.6) at 1st, 2nd and 3rd observation respectively (Figure 8)..

Potassium content (% DW)

The maximum K^+ content in shoot was recorded with good quality water treatment containing (2.98, 5.16 and 5.25) whereas minimum in irrigation water treatment of 9.0 EC (1.2, 1.79 and 1.5) at 1st, 2nd and 3rd observation respectively (Figure 7). The highest K^+ content in root was recorded from good quality water treatment having (4.13, 6.18 and 7.26), whereas minimum with irrigation water treatment

of 9.0 EC containing (1.66, 3.37 and 3.96) at 1st, 2nd and 3rd observation respectively. A significant reduction was observed with every increasing salinity level of irrigation water. AVKB 19 variety had significantly higher shoot and root K^+ content than ICMV 15111 variety. The higher affinity for K^+ transporters may also act as lower affinity for Na^+ transporters under saline environment, which may reduce K^+ uptake. Our results are supported with the findings of Liu et al. 2000 and Radhouane et al. 2013.

Yield

Green fodder yield: The main effect of irrigation water salinity on green fodder yield of pearl millet has suggested that irrigation with good quality (0.6 dS/m) and low salinity (3.0 dS/m) water produced the maximum green fodder yield of 37.55 and 34.74 t/ha, respectively at 1st cut at 50 DAS (Table 5). It was minimum i.e. 23.97 t/ha (36.17 % reduction from good quality water) with high salinity water (9.0 dS/m) irrigation.

The statistical analysis of results on performance of pearl millet cultivars showed that AVKB-19 produced 32.74 t/ha green fodder yield which was significantly higher as compared to 30.65 t/ha in case of ICMV-15111 on 1st cut taken at 50 DAS. Data on green fodder yield showed significant differences between two cultivars across different saline water levels. The variety AVKB-19 produced significantly higher mean green fodder yield (27.96 t/ha) as compared to ICMV-15111(25.51t/ha). The interaction between water quality and varieties was non-significant for both cuttings. Ghadiri et al. (2005) reported reduced uptake of water by salinity due to the high osmotic potential of the soil solution and high concentrations of specific ions that may cause physiological disorders in the plant tissues




Fig 6: Effect of saline water irrigations on root and shoot Na^+ content (%) in dual purpose pearl millet varieties




Fig 7: Effect of saline water irrigations on root and shoot K⁺ content (%) in dual purpose pearl millet varieties

and reduced yields. The primary effect of high salinity of irrigation water generally renders less water available to plants in spite of the fact that some is still present in the root zone. This is because of the fact that osmotic pressure of the soil solution becomes more negative as the salt concentration increases. Apart from the osmotic effect of salts in the soil solution, excessive concentration and absorption of individual ions such as Na⁺ may prove toxic to the plants and/or may retard the absorption of other essential plant nutrients (Qadir et al. 2008). Yadav et al. (2004); Nadaf et al. (2010), and Zhapayev et al. (2015) also reported similar findings in different crops.

Dry Matter yield: The mean maximum DM yields at

1st (5.89 t/ha) and 2nd cut (5.24 t/ha) were recorded with good quality irrigation water. Application of good quality water resulted into 23.34 and 13.69% higher DM yield at 1st and 2nd cutting, respectively over irrigation water of EC 9.0 dS/m. However, the irrigation water upto EC 3.0 dS/m at 1st cut and upto EC 6.0 dS/m at 2nd cut were found at par with good quality irrigation water (Table 5). This may be due to the fact that combined effect of rainfall and irrigation after 1st cut diluted the salt concentration and helped in leaching the excess salts from the root zone up to irrigation water EC 6.0 dS/m. The AVKB-19 resulted into significantly maximum DM yield at the 1st (5.67 t/ha) as well as 2nd cut (5.09 t/ha) over ICMV-15111. The supply

Table 5: Effect of saline water irrigation on yield of pearl millet varieties

Treatments	Green fodder yield (t/ha)		Dry matter yield (t/ha)		Grain yield (t/ha)
	1 st cut	2 nd cut	1 st cut	2 nd cut	At 110 DAS
<i>Salinity levels (dS/m)</i>					
Control (0.6)	37.55a	31.50a	5.89a	5.25a	2.17a
3	34.74b	28.61b	5.69a	5.13a	1.95a
6	30.52C	25.55c	5.25b	4.88a	1.72b
9	23.97d	21.30d	4.51c	4.53b	1.36b
SEm (±)	0.99	0.94	0.18	0.19	0.12
CD (P=0.05)	2.06	1.95	0.38	0.40	0.24
<i>Varieties</i>					
ICMV-15111	30.65a	25.51a	5.00b	4.79b	4.79b
AVKB-19	32.74b	27.96a	5.67a	5.09a	5.09a
SEm (±)	0.70	0.66	0.13	0.14	0.14
CD (P=0.05)	1.46	1.38	0.27	0.29	0.29

of good quality water significantly enhanced the uptake of macro and micro nutrient by the plants which ultimately lead to higher DM yield due its beneficial effect on the photosynthetic efficiency of plants. Whereas the higher load of salt with increased level of salinity of irrigation water restricted the root growth of plants that in turn reduced the uptake of nutrients leading to leaf chlorosis that reduces the photosynthetic potential of crops which ultimately leads to lower DM yield. The similar results of lower DM yield with increasing salinity levels were also reported by Kumari et al. 2014 and Farissi et al. 2018.

Grain yield: Variety AVKB-19 produced significantly higher (16.26%) grain yield of 1.93 t/ha as compared to 1.66 t/ha in ICMV-15111, and thus has potential to grow in saline environment. Significantly highest (2.17 t/ha) mean grain yield was recorded with good quality water in comparison to 1.95, 1.72 and 1.35 t/ha achieved with irrigation using 3.0, 6.0 and 9.0 dS/m salinity water, respectively (Table 5). The magnitude of reduction (%) with increasing salinity over good quality irrigation water was observed 10.14, 20.74 and 37.33 with 3.0, 6.0 and 9.0 dS/m salinity water, respectively. Decrease in grain yield by increasing salinity of irrigation water may be due to more negative water potential of soil solution causing reduced water and nutrient uptake consequently lower leaf area development in turn reduced net assimilates. Shani and Dudley (2001) also related the yield loss to reduced photosynthesis, high energy and carbohydrate expenses in osmoregulation, and interference with cell functions under saline conditions. Similar results for decreasing grain yields under saline environment were reported in pearl millet by Meena et al. (2012).

Conclusions

Our results suggest that AVKB-19 genotype of pearl millet proved superior and may be adapted as a choice for getting higher grain as well as green fodder yield with better physiological response as compare to the ICMV 15111 under saline environment in north-western region of India and elsewhere under similar agro-climatic conditions.

Acknowledgement

We sincerely acknowledge Director, ICAR-NDRI for financial support, Director, ICAR-CSSRI and In-charge, Agronomy section, ICAR-NDRI, Karnal for providing field staff, lab facilities and assistance in conducting this research.

REFERENCES

- Agastian P, Kingsley S J and Vivekanandan M. 2000. Effect of salinity on photosynthesis and biochemical characteristics in mulberry genotypes. *Photosynthetica* **38**(2): 287–90.
- Ashraf M P J C and Harris P J C. 2004. Potential biochemical indicators of salinity tolerance in plants. *Plant Science* **166**(1): 3–16.
- Ashraf M Y and Wu L. 1994. Breeding for salinity tolerance in plants. *Critical Reviews in Plant Sciences* **13**(1): 17–42.
- Bates L S, Waldren R P and Teare I D. 1973. Rapid determination of free proline for water-stress studies. *Plant and Soil* **39**1: 205–7.
- Bernstein, L. 1975. Effects of salinity and sodicity on plant growth. *Annual Review of Phytopathology* **13**(1): 295–312.
- Blum A. 1988. Plant breeding for stress environments. CRC Press, Boca Raton, Florida, USA. pp. 1–223.
- Bradford M M. 1976. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein dye binding. *Analytical Biochemistry*. **7**: 248–54.
- Central Ground Water Board. 1997. Inland groundwater salinity in India. Central Ground Water Board, Ministry of Water Resources, Government of India, Faridabad.
- Dionisio-Sese M L and Tobita S. 1998. Antioxidant responses of rice seedlings to salinity stress. *Plant Science* **135**1: 1–9.
- Ebercon A, Blum A and Jordan W R. 1977. A rapid colorimetric method for epicuticular wax content of sorghum leaves. *Crop Science*, **17**1: 179–80.
- Farissi M, Mouradi M, Bouizgaren A and Ghoulam C. 2018. Variations in leaf gas exchange, chlorophyll fluorescence and membrane potential of *Medicago sativa* root cortex cells exposed to increased salinity: The role of the antioxidant potential in salt tolerance. *Archives of Biological Sciences* **70**(3): 413–23.
- Flowers T J and Lauchli A. 1983. Sodium versus potassium: substitution and compartmentation. *Encyclopedia of Plant Physiology*. New series.
- Ghadiri H, Dordipour I, Bybordi M and Malakourti M J. 2005. Potential use of Caspian Sea for supplementary irrigation in North Iran. *Agricultural Water Management* **79**: 209–24.
- Greenway H and Munns R. 1980. Mechanisms of salt tolerance in nonhalophytes. *Annual review of Plant Physiology* **31**(1): 149–90.
- Hiscox J T and Israelstam G F. 1979. A method for the extraction of chlorophyll from leaf tissue without maceration. *Canadian Journal of Botany* **57**12: 1332–4.
- Kulkarni V N, Rai K N, Dakheel A J, Ibrahim M, Hebbara M and Vadez V. 2006. Pearl millet germplasm adapted to saline conditions. *International Sorghum and Millets Newsletter* **47**: 103–6.
- Kumar A, Sharma S K, Lata C, Sheokand S and Kulshreshtha N. 2015. Combined effect of boron and salt on polypeptide resolutions in wheat varieties differing in their tolerance. *Indian Journal of Agricultural Sciences*, **85**: 106–12.
- Kumar A, Kumar A, Lata C, and Kumar S. 2016. Eco-physiological responses of *Aeluropus lagopoides* (grass halophyte) and *Suaeda nudiflora* (non-grass halophyte) under individual and interactive sodic and salt stress. *South African Journal of Botany* **105**: 36–44.
- Kumar A, Krishnamurthy S L, Lata C, Kumar P, Devi R, Kulshreshtha N, Yadav R K and Sharma S K. 2016. Effect of dual stress (salinity and drought) on gas exchange attributes and chlorophyll fluorescence characteristics in rice. *Indian Journal of Agricultural Sciences* **86**(6): 19–27.
- Kumar A, Lata C, Krishnamurthy S L, Kumar A, Prasad K R K and Kulshreshtha N. 2017. Physiological and Biochemical Characterization of Rice Varieties under Salt and Drought Stresses. *Journal of Soil Salinity and Water Quality* **9**(2): 167–77.
- Kumari A, Kumar P, Ahmad E, Kumar R, Yadav R K, Datt Chander and Chinchmalatpure A. 2014. Fodder yield and quality of oats fodder (*Avenasativa* L.) as influenced by salinity of irrigation water and applied nitrogen levels. *Indian Journal of Animal Nutrition* **31**: 266–71.
- Lal K, Meena R L, Gupta S K, Saxena C K, Yadav G and Singh

- G. 2008. *Diagnosis and Management of Poor Quality Water and Salt Affected Soils*. Central Soil Salinity Research Institute, Karnal, India. p 311.
- Lata C, Kumar A, Sharma S K, Singh J, Sheokand S, Pooja, Mann A and Rani B. 2017. Tolerance to combined boron and salt stress in wheat varieties: Biochemical and molecular characterization. *Indian Journal of Experimental Biology* **55**: 321–8.
- Liu W, Schachtman D P and Zhang W. 2000. Partial deletion of a loop region in the high affinity K⁺ transporter HKT1 changes ionic permeability leading to increased salt tolerance. *Journal of Biological Chemistry* **275**(36): 27924–32.
- Meena H N, Bhalodia P K, Jat R S and Vekaria L C. 2012. Prospects of using saline water for irrigation in groundnut (*Arachis hypogaea*), pearl millet (*Pennisetum glaucum*) cropping system in saline black soils of Saurashtra. *Indian Journal of Agronomy* **57**(2): 122–6.
- Misra N and Gupta A K. 2005. Effect of salt stress on proline metabolism in two high yielding genotypes of green gram. *Plant Science* **169**: 331–9.
- Murakeozy E P, Nagy Z, Duhaze C, Bouchereau A and Tuba Z. 2003. Seasonal changes in the levels of compatible osmolytes in three halophytic species of inland saline vegetation in Hungary. *Journal of Plant Physiology* **160**(4): 395–401.
- Nadaf S K, Al-Hinai S A, Al-Farsi S M, Al-Lawati A H, Al-Bakri A N, Ahmed M and Hussain, N. 2010. Differential response of salt tolerant pearl millet genotypes to irrigation water salinity. Published in the Monograph on Management of Salt-Affected Soils and Water for Sustainable Agriculture. Mushtaque A, Al-Rawahi S A and Hussain N (Eds). Sultan Qaboos University, Oman. pp 47–60.
- Netondo G W, Onyango J C and Beck E. 2004. Sorghum and salinity: I. Response of growth, water relations, and ion accumulation to NaCl salinity. *Crop Science* **44**(3): 797.
- Parida A, Das A B and Das P. 2002. NaCl stress causes changes in photosynthetic pigments, proteins, and other metabolic components in the leaves of a true mangrove, *Bruguiera parviflora*, in hydroponic cultures. *Journal of Plant Biology* **45**(1): 28–36.
- Patel M R, Sadhu A C, Patel R M, Parmar H P and Kher HR. 2008. Cutting management in different genotypes of forage bajra during summer season. *Research on Crops* **9**(2): 325–7.
- Prado F E, Boem C, Gallardo M and Gonzalez H J A. 2000. Effect of NaCl on germination, growth and soluble sugar content in (*Chenopodium quinoa*) wild seeds. *Botany Academy of Sciences* **41**: 27–34.
- Qadir M, Tubeileh A, Akhtar J, Larbi A, Minhas P S and Khan M A. 2008. Productivity enhancement of salt-affected environments through crop diversification. *Land Degradation and Development* **19**(4): 429–53.
- Radhouane L. 2013. Agronomic and physiological responses of pearl millet ecotype (*Pennisetum glaucum* (L.) R. Br.) to saline irrigation. *Emirates Journal of Food and Agriculture* **25**(2): 109–16
- Rains D W. 1972. Salt transport by plants in relation to salinity. *Annual Review of Plant Physiology* **23**(1): 367–88.
- Rengasamy P. 2006. World salinization with emphasis on Australia. *Journal of Experimental Botany* **57**(5): 1017–23.
- Sairam R K, Veerabhadra R K and Srivastava G C. 2002. Differential responses of wheat genotypes to long term salinity stress in relation to oxidative stress, antioxidant activity and osmolyte concentration. *Plant Science* **163**: 1037–47.
- Shani U and Dudley L M. 2001. Field studies of crop response to water and salt stress. *Soil Science Society of American Journal* **65**: 1522–8.
- Sharma D K and Chaudhari S K. 2012. Agronomic research in salt affected soils of India: An overview. *Indian Journal of Agronomy* **57**(3s): 175–85.
- Sharma DK and Singh A. 2015. Salinity research in India-achievements, challenges and future prospects. *Water Energy International* **58**(6): 35–45.
- Singh A, Sharma P C, Kumar A, Meena M D and Sharma D K. 2015. Salinity induced changes in chlorophyll pigments and ionic relations in bael (*Aegle marmelos* Correa) cultivars. *Journal of Soil Salinity and Water Quality* **71**: 40–4.
- Vijayalakshmi T, Varalaxmi Y, Jainender S, Yadav S K, Vanaja M, Jyothilakshmi N and Maheshwari M. 2012. Physiological and biochemical basis of water-deficit stress tolerance in pearl millet hybrid and parents. *American Journal of Plant Sciences* **3**(12): 1730.
- Wang W, Vinocur B, Altman A. 2003. Plant responses to drought, salinity and extreme temperatures: Towards genetic engineering for stress tolerance. *Planta* **218**: 1–14.
- Weatherley P. 1950. Studies in the water relations of the cotton plant. *New Phytologist* **49**: 81–97.
- Yadav R K, Kumar A, Lal D and Batra L. 2004. Yield responses of winter (rabi) forage crops to irrigation with saline drainage water. *Experimental Agriculture* **40**: 65–75.
- Yemm EW and Willis AJ. 1954. The estimation of carbohydrates in plant extracts by anthrone. *Biochemical Journal* **57**: 508.
- Yeo AR. 1983. Salinity resistance: physiologies and prices. *Physiologia plantarum* **58**(2): 214–22.
- Zhapayev R, Toderich K, Popova V, Tautenov I, Umirzakov S, Bekzhanov S and Karabayev M. 2015. Forage production and nutritional value of Sorghum and Pearl Millet on marginal lands on Priaralie. *Journal of Arid Land Studies* **25**(3): 169–72.