

Initial growth responses of blackgram (*Vigna mungo* L. Hepper) under elevated CO₂ and moisture stress

M. Vanaja, P. Ratnakumar, P. Vagheera, M. Jyothi, P. Raghuram Reddy, N. Jyothi Lakshmi, M. Maheshwari, S.K. Yadav

Central Research Institute for Dryland Agriculture, Santhoshnagar, Hyderabad, India

ABSTRACT

An attempt was made to understand the interactive effect of the elevated CO₂ and moisture stress on germination and initial growth responses of blackgram (*Vigna mungo* L. Hepper). Four open top chambers (OTCs) were used for different conditions: irrigated with ambient CO₂ (365 ppm), irrigated with elevated CO₂ (600 ppm), moisture stress with ambient CO₂ and moisture stress with elevated CO₂. The percentage of germination, germination speed, emergence index, and vigor index were measured on the 5th and 6th day after sowing (DAS). Plants were harvested at different time intervals, i.e. on 7, 14, 21 and 28 DAS; leaf area and dry weights of the seedlings were recorded. It was observed that moisture stress in general reduced the germination in all the conditions and affected seedling growth of blackgram. Germination percentage, emergence index, germination speed and vigor index were increased with elevated CO₂ under both irrigated and moisture stress conditions. Plants grown with elevated CO₂ were taller and attained a greater leaf area along with more biomass than ambient CO₂ levels under irrigated and moisture stress conditions at all time intervals. The increase in the germination, larger leaf area and dry matter of root, shoot and leaf proved that CO₂ enrichment of the atmosphere will be beneficial for the crops for better establishment and greater productivity.

Keywords: elevated CO₂; blackgram; moisture stress; germination; biomass; leaf area

Atmospheric CO₂ concentration, which was about 280 μmol/mol before the industrial revolution, is now 360 μmol/mol and is increasing by 1.8 μmol/mol/year (Houghton et al. 2001) due to man-made activities. Much of the changes in atmospheric CO₂ occurred over the last fifty years (1953 to 2003) and the increase in CO₂ was about 64 ppm (Krull et al. 2005) during this period. This will affect the vegetation, as elevated CO₂ substantially increases photosynthesis and thereby the growth and total biomass, particularly in C₃ plants (Drake et al. 1997). Increased concentrations of elevated CO₂ have already resulted in a global increase of temperatures, and this trend is expected to continue. Changes in temperature are likely to alter precipitation worldwide, decreasing in many areas (Houghton et al. 2001). Together with the higher evapotranspiration from soils, it can result in a prolonged water deficit (Samarakoon and Gifford 1995, Ellsworth 1999). Decreasing water supply causes moisture stress and affects the growth of the plants. Under the present global scenario of CO₂ increase accord-

ing to the Intergovernmental Panel on Climate Change (IPCC 1996), changes in temperature and water availability affect growth and many key metabolic processes in plants. Particularly, there is limited quantitative understanding of the effects of interactions of elevated CO₂ and water deficit (Samarakoon and Gifford 1995).

The crops and weeds responded differently to germination and emergence under elevated CO₂ conditions. Ziska and Bunce (1993) reported that doubling of CO₂ concentration resulted in an increase in the rate and final percentage of germination for alfalfa, and weed species *Amaranthus hybridus* and *Chenopodium album*. Several researchers reported that the response of pulse crops was stronger under elevated CO₂ conditions compared with other cereals and oil seeds. Cowpea showed positive growth responses to elevated CO₂ grown in controlled environmental chambers (Mbikayi et al. 1983). Greengram grown under CO₂ enrichment was taller and attained a greater leaf area along with dry matter than ambient CO₂ grown plants at initial growth

stages (Srivastava et al. 2001). Under elevated CO₂ (600 ppm) and well-watered conditions the response of blackgram was higher than sorghum and sunflower at initial growth stages (Vanaja et al. 2006). Blackgram [*Vigna mungo* (L.) Hepper] is an important source of protein and is cultivated as short duration rainfed crop in semiarid areas of South Asian countries. This paper tried to quantify the response of blackgram to elevated CO₂ and moisture stress in terms of germination and growth at initial stages.

MATERIAL AND METHODS

Plant material and growth conditions. The seeds of blackgram cv. T-9 were sown in four open top chambers (OTCs) of 3 × 3 m, lined with transparent PVC (polyvinylchloride) sheet having 90% transmittance of light. The seeds were sown directly in the soil (alfisol) to study the effect of elevated CO₂ level (600 ± 50 ppm) on germination and initial plant growth up to four weeks. Among the four OTCs, two OTCs were maintained at 600 ± 50 ppm and the elevated levels of CO₂ at crop canopy level was maintained by continuously injecting 100% CO₂ into plenum where it will mix with air from air compressor before entering into the chamber. Other two OTCs were maintained at ambient CO₂ level (365 ppm) without any external CO₂ supply and served as control chambers. The air sample from each chamber was drawn at 3-minute intervals into non-dispersive infrared (NDIR) CO₂ analyzer (California Analytical) and the set ppm of CO₂ concentration (600 ± 50 ppm) was maintained with the help of solenoid valves, rotameters, PCs, Program Logic Control (PLC) and Supervisory Control and Data Acquisition (SCADA) software. Throughout the experimental period continuous measurement of relative humidity and temperature of all the OTCs was possible with the sensors fitted inside the chambers. The temperature of chambers with elevated levels of CO₂ remained nearly the same as in the control chambers. The light intensity in chambers was 80–95% of the outside field. However gentle washing of polythene cover was frequently required to maintain transparency. For each CO₂ level one OTC was maintained stress free and other moisture stressed. Each chamber had four 1 × 1 m plots, each plot with five rows of crop with ten plants in each row. The experimental site was sandy loam in texture, neutral in pH (6.8), low in available nitrogen (225 kg/ha), phosphorus (10 kg/ha) and medium

to high in available potassium (300 kg/ha). The seeds were sown at field capacity and after sowing the moisture stress was imposed with holding the water in two OTCs and in other two the plots were maintained stress free by irrigating at regular intervals. The moisture content of the stressed and control plots were measured gravimetrically and expressed as percentage of moisture. The control plots were maintained at field capacity 16–18% of moisture and stressed plots at 7–9% soil moisture.

Germination studies. Germination measurements were made on the 5th and 6th day from the date of sowing according to ISTA (1976). Standard practices were followed to calculate percentage of germination: germination speed = % germination/days of completion of germination, emergence index = DN/N where DN = emergence (number of seed germination on particular day) and N = days of emergence, vigor index = % of germination × shoot length.

Plant growth. Plants were harvested at weekly intervals with three replications of each treatment, with ten plants for each replication, i.e. thirty plants for each treatment. Roots were carefully washed from the soil and plants were separated into roots, stems and leaves. The measurements were recorded on length of shoot and root for individual plants, i.e. 10 × 3 observations for each parameter, whereas leaf area, stem dry weight, root dry weight and leaf dry weights were recorded for 10 plants/replication, i.e. 2 × 3 observations for each parameter. The root length was recorded on the main root of each plant and root volume of 10 plant roots was measured as ml of water displaced and expressed as ml/10 plants. The leaf area at different time intervals was measured with photo-electronic leaf area meter (LI-3100, LI-COR) and expressed as cm²/10 plants. The dry weights of stem, root and leaf were recorded after thorough drying of the plant material in hot air oven at 65°C and expressed as g/10 plants. All the data were statistically analyzed using a two-way analysis of variance (ANOVA) to test the significance of treatments and their interactions.

RESULTS AND DISCUSSION

The mean values of germination characters of blackgram studied at 5 and 6 DAS under elevated CO₂ (600 ± 50 ppm) and chamber control (365 ppm) for four treatments were presented in Table 1 and growth responses at 28 DAS in Table 2, while the

Table 1. Interactive effect of elevated CO₂ and moisture stress on germination measurements over chamber control of blackgram

Germination measurements	Water status					
	irrigated			moisture stress		
	chamber control (365 ppm)	elevated CO ₂ (600 ppm)	percentage increase	chamber control (365 ppm)	elevated CO ₂ (600 ppm)	percentage increase
Percent of germination (%)	95	100	5.26	80	98	22.0
Emergence index	15.8	16.7	5.69	13.3	16.3	22.5
Vigor index	34.2	41.0	19.8	26.6	35.0	31.5
Germination speed	15.8	16.6	5.06	13.3	15.6	17.2

ANOVA of the above characters at different time intervals in Table 3.

Germination studies in blackgram showed a positive response under elevated CO₂ whereas moisture stress delayed the germination. The percentage increase of germination at 6 DAS under elevated CO₂ over ambient condition was 5.2% and 22%, with irrigated and moisture stress respectively. The emergence index was also increased with elevated CO₂ both under irrigated and stress and the increment was 5.6% and 22% over chamber control,

respectively. The vigor index was increased by 20% and 31% with elevated CO₂ for irrigated and stress conditions, respectively, when compared to the ambient level. The speed of germination was higher with elevated CO₂ levels both under irrigated (5%), and stress conditions (17%) as compared to ambient variants (Table 1).

The growth response of blackgram was positive to elevated CO₂ both under irrigated and moisture stress conditions. The lengths of root and shoot decreased under moisture stress both

Table 2. Interactive effect of elevated CO₂ and moisture stress on growth components over chamber control of blackgram at 28 DAS

Plant measurements	Water status					
	irrigated			moisture stress		
	chamber control (365 ppm)	elevated CO ₂ (600 ppm)	percentage increase	chamber control (365 ppm)	elevated CO ₂ (600 ppm)	percentage increase
Geometry						
Leaf area (cm ²)	913.45	1067.81	16.89	658.43	837.74	27.23
Shoot length (cm)	8.93	10.0	11.98	8.49	9.33	9.89
Root length (cm)	9.78	13.26	35.58	10.04	10.94	8.96
Root volume (ml)	2.0	3.5	75.0	1.5	3.0	100
Dry weights (g)						
Leaf	3.72	5.31	42.74	3.42	4.38	28.07
Stem	1.22	1.66	36.06	0.91	1.31	43.96
Root	0.30	0.42	40.00	0.28	0.36	28.57
Total plant	5.24	7.39	41.03	4.61	6.05	31.24
Root:shoot ratio	0.062	0.060	-3.22	0.062	0.065	4.85

Table 3. ANOVA for various characters at 28 DAS of blackgram crop with elevated CO₂ (600 ppm), and chamber control (365 ppm) under irrigated and moisture stress conditions

Characteristics	Mean sum of squares								
	DF	shoot length	root length	leaf area	total dry weight	stem dry weight	root dry weight	leaf dry weight	root:shoot weight
Replications	2	1.23*	0.230	3704*	0.003	0.005	0.002	0.001	0.00
Conditions	1	11.33**	18.22**	75524**	3.74**	0.153**	0.027**	1.898**	0.00
Treatments	1	3.739**	8.99**	230518**	1.84**	0.152**	0.003	0.829*	0.001*
Time intervals	3	67.73**	81.5**	1678248**	75.7**	3.584**	0.179**	40.860**	0.173**
Conditions × treatments	1	0.263	0.235	314.0	0.16	0.005	0.001	0.094	0.00
Conditions × time intervals	3	0.543	1.234**	20169**	2.05**	0.128**	0.005**	1.042**	0.005**
Treatments × time intervals	3	0.431	0.131	42957**	0.54	0.068**	0.001	0.205	0.00
Conditions × treatments × time intervals	3	0.003	1.839**	87.33	0.07	0.003	0.000	0.070	0.00
Error	30	0.343	0.194	892.91	0.191	0.010	0.001	0.140	0.00

**significance at 1% level, *significance at 5% level

with elevated level of CO₂ and ambient conditions when compared with respective irrigated conditions (Figures 1a, b). The shoot length with elevated CO₂ was higher at different DAS and at 28 DAS it was higher by 12% under irrigated and 9.8% under stress conditions, as compared to the ambient level CO₂. The increment in root length was smaller when compared with the increment in root volume under elevated CO₂. Under irrigated conditions the percentage increase of root length with elevated CO₂ was 35%, whereas for root volume it was 75%. Under moisture stress conditions the response of elevated CO₂ for root length was only 8.96% whereas the root volume doubled (100%) (Figure 1h). This was mainly due to more lateral roots and root hair formation with elevated CO₂ levels. Leaf area was decreased drastically under stress and that was mitigated with elevated CO₂ (Figure 1c) and showed an increase of 17% under irrigated and 27% under stress conditions at 28 DAS.

The response of total dry weight (Figure 1d) to elevated CO₂ was also significant and showed a positive response at different time intervals. Elevated CO₂ increased total dry weights by 41% (irrigated) and 31% (stress) at 28 DAS. The responses of leaf, stem and root dry weight (Figures 1e–g) were also positive and significant with elevated CO₂ both under irrigated and stress conditions.

The percentage increase in dry weight of leaf was 43% and 28%; stem 36% and 44%; root 40% and 28% under irrigated and stress conditions, respectively (Table 2) at 28 DAS. The percentage of root:shoot ratio with elevated CO₂ was negative (–3.2%) under irrigated conditions and positive (4.8%) with moisture stress.

The results of the present investigation clearly showed that with elevated CO₂ level the percentage of germination, speed of germination, emergence index and vigor index were positively increased. The increment of the above-mentioned parameters was higher under moisture stress conditions than under irrigated conditions. The mechanism by which elevated CO₂ affects germination and emergence is unknown. This could be due to the thin seed coat, which turns translucent, and underlined cotyledons can photosynthesize in blackgram as it was reported in *Kalmia* (Heichel and Jaynes 1974). Ziska and Bunce (1993) observed that small seeds have a greater surface to volume ratio and hence a greater diffusing capacity for CO₂. On other hand elevated CO₂ levels may stimulate the internal ethylene production (Esashi et al. 1986); Ziska and Bunce (1993) were speculating that the influence of the elevated CO₂ in stimulating the germination and emergence might be ethylene. To determine the actual effect of elevated CO₂ on germination/emergence and to understand the




Figure 1. Response of blackgram to elevated CO₂ and moisture stress at different days after sowing

underlined mechanisms further a more detailed study is needed.

The percentage of increase in root length was higher under elevated than ambient CO₂ both under irrigated and stress conditions. Increased root length implies the probability of deeper soil

penetration and access to deeper reservoirs of soil moisture. This deeper rooting would be an advantage if climates became drier. Root volume was also increased due to more lateral roots and root hair formation under elevated CO₂ condition. CO₂ enrichment enhances the root growth much

more by increasing its length, volume and weight (Rogers et al. 1992). Increased root biomass due to high CO₂ also indicates greater partitioning of assimilates to the roots. The comparison of partitioning of dry matter allocation to above- and under-ground portions showed that under elevated CO₂ the response of root to shoot ratio was negative under irrigated condition, whereas it was positive under moisture stress condition. The results indicate that proportioning of assimilates was greater to the roots than to the shoots under stress (Table 2). This will provide the plants with a greater support for the growth when grown under high CO₂ and moisture stress environment.

High CO₂ caused a significant increase in growth both in terms of leaf area and dry matter production. The increase in stem and leaf growth with elevated CO₂ caused an overall increase in total biomass in blackgram under irrigated and moisture stress conditions. Similar reports with *Vigna radiata* L. also showed an increase in total biomass under elevated CO₂ conditions (Srivastava et al. 2001); it resulted in a greater leaf growth, which in turn produced greater photosynthetic surface area at the early stages of plant and when combined with higher root growth, it can facilitate better initial establishment and growth under moisture stress environment.

Acknowledgements

This work was carried out in AP-Cess program and authors are grateful to ICAR for the financial support. The support and guidance provided by the Director, CRIDA is highly acknowledged.

REFERENCES

Drake B.G., Gonzalez-Meler M.A., Long S.P. (1997): More efficient plants: A consequence of rising atmospheric CO₂? *Ann. Rev. Plant Physiol. Plant Molec. Biol.*, 48: 609–639.

Ellsworth D.S. (1999): CO₂ enrichment in maturing pine forest: are CO₂ exchange and water stress in the canopy affected? *Plant Cell Environ.*, 22: 461–472.

Esashi Y., Ooshima Y., Michiharu A., Akiko K., Satoh S. (1986): CO₂-enrichment of C₂H₄ production in tissues of imbibed cocklebur seeds. *Aust. J. Plant Physiol.*, 13: 417–429.

Heichel G.H., Jaynes R.A. (1974): Stimulating emergence and growth of kalmia genotype with CO₂. *Hort. Sci.*, 9: 60–62.

Houghton J.T., Ding Y., Griggs D.J., Noguier M., Van der Linden P.J., Xiaosu D. (eds.) (2001): Intergovernmental panel on climate change. In: *Climate change 2001: The scientific basis*. Cambridge Univ. Press.

IPCC (1996): *Climate Change 1995: Summary for policy makers and technical summary of the working group*. In: Houghton J.T., Meria Filho L.G., Callander B.A., Harris N., Kattenberg A., Maskell K. (eds.): *Intergovernmental panel on climate change*. Cambridge Univ. Press, Cambridge, UK.

ISTA (1976): International rules for seed testing. In: *Proc. Int. Seed. Test. Assoc.*, 31: 1–52.

Krull E.S., Skjemstad J.O., Burrows W.H., Bray S.G., Wynn J.G., Bol R., Spouncer L., Harms B. (2005): Recent vegetation changes in central Queensland, Australia: Evidence from δ¹³C and ¹⁴C analyses of soil organic matter. *Geoderma*, 126: 241–259.

Mbikayi N.T., Hileman D.R., Bhattacharya N.C., Ghosh P.P., Biswas P.K. (1983): Effect of CO₂ enrichment on physiology and biomass production in cowpea (*Vigna unguiculata* L.) grown in open top chambers. In: *Proc. Int. Congr. Plant physiology*, New Delhi: 640–645.

Samarakoon A.B., Gifford R.M. (1995): Soil water content under plants at high CO₂ concentration and interaction with the treatment CO₂ effect. A species comparison. *J. Biogeogr.*, 22: 193–202.

Srivastava G.C., Pal M., Das M., Sengupta U.K. (2001): Growth, CO₂ exchange rate and dry matter partitioning in mung bean (*Vigna radiata* L.) grown under elevated CO₂. *Indian J. Exp. Biol.*, 39: 572–577.

Vanaja M., Vagheera P., Ratnakumar P., Jyothi Lakshmi N., Raghuram Reddy P., Yadav S.K., Maheswari M., Venkateswarulu B. (2006): Evaluation of certain rainfed food and oil seed crops for their response to elevated CO₂ at vegetative stage. *Plant Soil Environ.*, 52: 164–170.

Ziska H.L., Bunce A.J. (1993): The influence of elevated CO₂ and temperature on seed germination and emergence from soil. *Field Crops Res.*, 34: 147–157.

Received on April 4, 2006

Corresponding author:

Dr. M. Vanaja, Crop Science Division, Central Research Institute for Dryland Agriculture, Santoshnagar, Hyderabad 500 059 (AP), India
 phone: 91 040 24 530 161, fax: 91 040 24 531 802, e-mail: mvanaja@crida.ernet.in
