



हर कदम, हर डगर
किसानों का हमसफर
भारतीय कृषि अनुसंधान परिषद

AgriSearch with a human touch

**LAND RESOURCE INVENTORY AND SOCIO-ECONOMIC STATUS OF
FARM HOUSEHOLDS FOR WATERSHED PLANNING AND
DEVELOPMENT**

NIRGUDI-1 (4D5C2B1c) MICROWATERSHED

Aland Taluk, Gulbarga District, Karnataka

Karnataka Watershed Development Project – II

SUJALA – III

World Bank funded Project



ICAR – NATIONAL BUREAU OF SOIL SURVEY AND LAND USE PLANNING



ICAR - NBSS & LUP



**WATERSHED DEVELOPMENT DEPARTMENT
GOVT. OF KARNATAKA, BANGALORE**



About ICAR - NBSS&LUP

The ICAR-National Bureau of Soil Survey and Land Use Planning (ICAR-NBSS&LUP), Nagpur, a premier Institute of the Indian Council of Agricultural Research (ICAR), was set up during 1976 with the objective to prepare soil resource maps at national, state and district levels and to provide research inputs in soil resource mapping and its applications, land evaluation, land use planning, land resource management, and database management using GIS for optimising land use on different kinds of soils in the country.

The Bureau has been engaged in carrying out soil resource survey, agro-ecological and soil degradation mapping at the country, state and district levels for qualitative assessment and monitoring the soil health towards viable land use planning. The research activities have resulted in identifying the soil potentials and problems, and the various applications of the soil surveys with the ultimate objective of sustainable agricultural development. The Bureau has the mandate to correlate and classify soils of the country and maintain a National Register of all the established soil series. The Institute is also imparting in-service training to staff of the soil survey agencies in the area of soil survey, land evaluation and soil survey interpretations for land use planning. The Bureau in collaboration with Panjabrao Krishi Vidyapeeth, Akola is running post-graduate teaching and research programme in land resource management, leading to M.Sc. and Ph.D. degrees.

Citation: Rajendra Hegde, Ramesh Kumar, S.C., K.V. Niranjana, S. Srinivas, M.Lalitha, B.A. Dhanorkar, R.S. Reddy and S.K. Singh (2019). "Land Resource Inventory and Socio-Economic Status of Farm Households for Watershed Planning and Development of NIRGUDI-1 (4D5C2B1c) Microwatershed, Aland Taluk, Gulbarga District, Karnataka", ICAR-NBSS&LUP Sujala MWS Publ 10, ICAR – NBSS & LUP, RC, Bangalore. p.83 & 29.

TO OBTAIN COPIES,

Please write to:

Director, ICAR - NBSS & LUP,

Amaravati Road, NAGPUR - 440 033, India

Phone : (0712) 2500386, 2500664, 2500545 (O)

Telefax : 0712-2522534

E-Mail : director@nbsslup.ernet.in

Website URL : nbsslup.in

Or

Head, Regional Centre, ICAR - NBSS&LUP, Hebbal, Bangalore - 560 024

Phone : (080) 23412242, 23510350 (O)

Telefax : 080-23510350

E-Mail : nbssrcb@gmail.com



**LAND RESOURCE INVENTORY AND SOCIO-ECONOMIC
STATUS OF FARM HOUSEHOLDS FOR WATERSHED
PLANNING AND DEVELOPMENT**

NIRGUDI-1 (4D5C2B1c) MICROWATERSHED

Aland Taluk, Gulbarga District, Karnataka

Karnataka Watershed Development Project – II

Sujala-III

World Bank funded Project



**ICAR – NATIONAL BUREAU OF SOIL SURVEY AND LAND USE
PLANNING**



**WATERSHED DEVELOPMENT DEPARTMENT, GOVT. OF
KARNATAKA, BANGALORE**



PREFACE

In Karnataka, as in other Indian States, the livelihoods of rural people are intertwined with farming pursuits. The challenges in agriculture are seriously threatening the livelihood of a large number of farmers as they have been practicing farming in contextual factors beyond their control. Climatic factors are the most important ones and have become much more significant in recent times due to rapid climate changes induced by intensive anthropogenic activities affecting our ecosystem in multiple ways. Climate change has become the reality, it is happening and efforts to evolve and demonstrate climate resilient technologies have become essential. Due to the already over stressed scenario of agrarian sector, the climate change is resulting in manifold increase in the complexities, pushing the rural mass to face more and more unpredictable situations. The rising temperatures and unpredictable rainfall patterns are going to test seriously the informed decisions farmers have to make in order to survive in farming and sustain their livelihood.

It is generally recognized that impacts of climate change shall not be uniform across the globe. It is said that impact of climate change is more severe in South Asia. Based on the analysis of meteorological data, it is predicted that in India, there will be upward trend in mean temperature, downward trend in relative humidity, annual rainfall and number of wet days in a year. Also, in general, phenomena like erratic monsoon, spread of tropical diseases, rise in sea levels, changes in availability of fresh water, frequent floods, droughts, heat waves, storms and hurricanes are predicted. Each one of these adverse situations are already being experienced in various parts of India and also at the global level. Decline in agricultural productivity of small and marginal farmers becoming more vulnerable is already witnessed.

In Karnataka, more than 60 per cent of the population live in rural areas and depend on agriculture and allied activities for their livelihood. Though the state has achieved significant progress in increasing the yield of many crops, there is tremendous pressure on the land resources due to the growing and competing demands of various land uses. This is reflected in the alarming rate of land degradation observed. Already more than 50 per cent of the area is affected by various forms of degradation. If this trend continues, the sustainability of the fragile ecosystem will be badly affected. The adverse effects of change in the climatic factors are putting additional stress on the land resources and the farmers dependent on this.

The natural resources (land, water and vegetation) of the state need adequate and constant care and management, backed by site-specific technological interventions and investments particularly by the government. Detailed database pertaining to the nature of the land resources, their constraints, inherent potentials and suitability for various land based rural enterprises, crops and other uses is a prerequisite for preparing location-specific action plans, which are in tune with the inherent capability of the resources. Any effort to evolve climate resilient technologies has to be based on the baseline scientific database. Then only one can expect effective implementation of climate resilient technologies, monitor the progress, make essential review of the strategy, and finally evaluate the effectiveness of the implemented programs. The information available at present on the land resources of the state are of general nature and useful only for general purpose

planning. Since the need of the hour is to have site-specific information suitable for farm level planning and detailed characterization and delineation of the existing land resources of an area into similar management units is the only option.

ICAR-NBSS&LUP, Regional Centre, Bangalore has taken up a project sponsored by the Karnataka Watershed Development Project-II, (Sujala-III), Government of Karnataka funded by the World Bank under Component -1 Land Resource Inventory. This study was taken up to demonstrate the utility of such a database in reviewing, monitoring and evaluating all the land based watershed development programs on a scientific footing. To meet the requirements of various land use planners at grassroots level, the present study on “Land Resource Inventory and Socio-Economic Status of Farm Households for Watershed Planning and Development of Nirgudi-1 Microwatershed, Aland Taluk, Kalaburgi District, Karnataka” for integrated development was taken up in collaboration with the State Agricultural Universities, IISC, KRSAC, KSNDMC as Consortia partners. The project provides detailed land resource information at cadastral level (1:7920 scale) for all the plots and socio-economic status of farm households covering thirty per cent farmers randomly selected representing landed and landless class of farmers in the microwatershed. The project report with the accompanying maps for the microwatershed will provide required detailed database for evolving effective land use plan, alternative land use options and conservation plans for the planners, administrators, agricultural extension personnel, KVK officials, developmental departments and other land users to manage the land resources in a sustainable manner.

It is hoped that this database will be useful to the planners, administrators and developmental agencies working in the area in not only for formulating location specific developmental schemes but also for their effective monitoring at the village/watershed level.

Nagpur

Date: 10.05.2016

S.K. SINGH

Director, ICAR - NBSS&LUP, Nagpur

Contributors

Dr. RajendraHegde Principal Scientist, Head & Project Leader, Sujala-III Project ICAR-NBSS&LUP, Regional Centre Bangalore	Dr. S.K.Singh Director, ICAR-NBSS&LUP Coordinator, Sujala-III Project Nagpur
Soil Survey, Mapping & Report Preparation	
Dr. B.A. Dhanorkar	Sh. R.S. Reddy
Dr. K.V. Niranjana	Sh. Nagendra, B.R.
	Smt. Chaitra, S.P.
Field Work	
Sh. C.BacheGowda	Sh. Mahesha, D.B.
Sh. Somashekar	Sh. Ashok S. Sindagi
Sh. VenkataGiriyappa	Sh. Veerabhadrapa
Sh. M. Jayaramaiah	Sh. Anand
Sh. Paramesha, K.	Sh. Arun N Kambar
	Sh. Shankarappa, K.
	Sh. Kamalesh K. Avate
	Sh. Sharan Kumar Huppar
	Sh. Yogesh, H.N.
	Sh. Kalaveerachari R. Kammar
GIS Work	
Dr. S.Srinivas	Sh. A.G.Devendra Prasad
Dr. M.Ramesh	Sh. Prakashanaik, M.K.
Sh. D.H.Venkatesh	Sh. Abhijith Sastry, N.S.
Smt.K.Sujatha	Sh. Sudip Kumar Suklabaidya
Smt. K.V.Archana	Sh. Mahamad Ali, M.
Sh. N.Maddileti	Sh. Avinash, K.N.
	Sh. Amar Suputhra, S
	Sh. Anudeep, Y.
	Sh. Deepak, M.J.
	Smt. K.Karunya Lakshmi
	Ms. Seema, K.V.
	Ms. A. Rajab Nisha

Laboratory Analysis	
Dr. K.M.Nair	Dr. H.R. Savitha
Smt. ArtiKoyal	Ms. Steffi Peter
Smt. Parvathy, S.	Ms. Thara, V.R
	Ms. Roopa, G.
	Ms. Swati, H.
	Sh. Shantaveera Swami
	Ms. Shwetha, N.K.
	Smt. Ishrat Haji
	Ms. P. PavanKumari
	Ms. Padmaja
	Ms. Veena, M.
Soil & Water Conservation	
Sh. Sunil P. Maske	
Socio-Economic Analysis	
Dr. S.C. Ramesh Kumar	Sh. M. K. Prakashanaik
	Ms. Sowmya K.B
	Sh.Manjunath M
	Sh.Veerabhadraswamy R
	Sh.Lankesh RS
	Sh.Kalaveerachari R Kammar
	Sh.Pradyumma U
	Sh.Yogesha HN
	Sh.Vijay kumar lamani
	Sh.Arun N Kambar
	Sh.Vinay
	Sh.Basavaraj.Biradar
	Sh.Vinod R
	Sh.Praveenkumar P Achalkar
	Sh.Rajendra D
Watershed Development Department, GoK, Bangalore	
Sh. Rajeev Ranjan IFS Project Director & Commissioner, WDD	Dr. A. Natarajan NRM Consultant, Sujala-III Project
Dr. S.D. Pathak IFS Executive Director & Chief Conservator of Forests, WDD	

PART-A
LAND RESOURCE INVENTORY

Contents

Preface		
Contributors		
Executive Summary		
Chapter 1	Introduction	1
Chapter 2	Geographical Setting	3
2.1	Location and Extent	3
2.2	Geology	4
2.3	Physiography	4
2.4	Drainage	5
2.5	Climate	5
2.6	Natural Vegetation	6
2.7	Land Utilization	7
Chapter 3	Survey Methodology	9
3.1	Base maps	9
3.2	Field Investigation	11
3.3	Laboratory Characterization	12
3.4	Finalization of Soil Map	12
Chapter 4	The Soils	19
4.1	Soils of Basalt Landscape	19
Chapter 5	Interpretation for Land Resource Management	25
5.1	Land Capability Classification	25
5.2	Soil Depth	27
5.3	Surface Soil Texture	28
5.4	Soil Gravelliness	29
5.5	Available Water Capacity	30
5.6	Soil Slope	31
5.7	Soil Erosion	32
Chapter 6	Fertility Status	35
6.1	Soil Reaction (pH)	35
6.2	Electrical Conductivity (EC)	35
6.3	Organic Carbon (OC)	35
6.4	Available Phosphorus	37
6.5	Available Potassium	37
6.6	Available Sulphur	37
6.7	Available Boron	37
6.8	Available Iron	40
6.9	Available Manganese	40
6.10	Available Copper	40
6.11	Available Zinc	40

Chapter 7	Land Suitability for Major Crops	43
7.1	Land suitability for Sorghum	43
7.2	Land suitability for Maize	46
7.3	Land suitability for Red gram	48
7.4	Land suitability for Sunflower	49
7.5	Land suitability for Cotton	51
7.6	Land suitability for Sugarcane	52
7.7	Land suitability for Soybean	53
7.8	Land suitability for Guava	54
7.9	Land suitability for Mango	56
7.10	Land suitability for Sapota	58
7.11	Land suitability for Jackfruit	60
7.12	Land suitability for Jamun	60
7.13	Land Suitability for Musambi	61
7.14	Land Suitability for Lime	62
7.15	Land Suitability for Cashew	64
7.16	Land Suitability for Custard Apple	64
7.17	Land Suitability for Amla	65
7.18	Land Suitability for Tamarind	66
7.19	Land Management Units	67
7.20	Proposed Crop Plan	69
Chapter 8	Soil Health Management	71
Chapter 9	Soil and Water conservation Treatment Plan	75
9.1	Treatment Plan	76
9.2	Recommended Soil and Water Conservation measures	79
9.3	Greening of microwatershed	80
	References	83
	Appendix I	I-X
	Appendix II	XI-XVIII
	Appendix III	XIX-XXIII

LIST OF TABLES

2.1	Mean Monthly Rainfall, PET, 1/2 PET at Aland Taluk, Kalaburgi District	5
2.2	Land Utilization in Aland Taluk	7
3.1	Differentiating Characteristics used for Identifying Soil Series	12
3.2	Soil Legend	17
7.1	Soil-Site Characteristics of Nirgudi-1microwatershed	44
7.2	Crop suitability criteria for Sorghum	45
7.3	Crop suitability criteria for Maize	47
7.4	Crop suitability criteria for Red gram	48
7.5	Crop suitability criteria for Sunflower	50
7.6	Crop suitability criteria for Cotton	51
7.7	Crop suitability criteria for Sugarcane	53
7.8	Crop suitability criteria for Guava	55
7.9	Crop suitability criteria for Mango	57
7.10	Crop suitability criteria for Sapota	59
7.11	Crop suitability criteria for Lime	63
7.12	Proposed Crop Plan for Nirgudi-1Microwatershed	69

LIST OF FIGURES

2.1	Location map of Nirgudi-1 microwatershed	3
2.2	Rock formations in Nirgudi-1 microwatershed	4
2.3	Rainfall distribution in Aland Taluk, Kalaburgi District	6
2.4	Natural Vegetation in Nirgudi-1 microwatershed	6
2.5	Current Land use – Nirgudi-1 microwatershed	8
2.6	Location of Wells- Nirgudi-1 microwatershed	8
3.1	Scanned and Digitized Cadastral map of Nirgudi-1 microwatershed	10
3.2	Satellite image of Nirgudi-1 microwatershed	10
3.3	Cadastral map overlaid on IRS PAN+LISS IV merged imagery of Nirgudi-1 microwatershed	11
3.4	Soil phase or management units of Nirgudi-1 microwatershed	15
5.1	Land Capability Classification of Nirgudi-1 microwatershed	27
5.2	Soil Depth map of Nirgudi-1 microwatershed	28
5.3	Surface Soil Texture map of Nirgudi-1 microwatershed	29
5.4	Soil Gravelliness map of Nirgudi-1 microwatershed	30
5.5	Soil Available Water Capacity map of Nirgudi-1 microwatershed	31
5.6	Soil Slope map of Nirgudi-1 microwatershed	32
5.7	Soil Erosion map of Nirgudi-1 microwatershed	33
6.1	Soil Reaction (pH) map of Nirgudi-1 microwatershed	36
6.2	Electrical Conductivity (EC) map of Nirgudi-1 microwatershed	36
6.3	Soil Organic Carbon (OC) map of Nirgudi-1 microwatershed	38
6.4	Soil Available Phosphorus map of Nirgudi-1 microwatershed	38
6.5	Soil Available Potassium map of Nirgudi-1 microwatershed	39
6.6	Soil Available Sulphur map of Nirgudi-1 microwatershed	39
6.7	Soil Available Boron map of Nirgudi-1 microwatershed	40
6.8	Soil Available Iron map of Nirgudi-1 microwatershed	41
6.9	Soil Available Manganese map of Nirgudi-1 microwatershed	41
6.10	Soil Available Copper map of Nirgudi-1 microwatershed	42
6.11	Soil Available Zinc map of Nirgudi-1 microwatershed	42
7.1	Land Suitability map of Sorghum	46
7.2	Land Suitability map of Maize	47

7.3	Land Suitability map of Red gram	49
7.4	Land Suitability map of Sunflower	50
7.5	Land Suitability map of Cotton	52
7.6	Land Suitability map of Sugarcane	53
7.7	Land Suitability map of Soybean	54
7.8	Land Suitability map of Guava	56
7.9	Land Suitability map of Mango	58
7.10	Land Suitability map of Sapota	59
7.11	Land Suitability map of Jackfruit	60
7.12	Land Suitability map of Jamun	61
7.13	Land Suitability map of Musambi	62
7.14	Land Suitability map of Lime	63
7.15	Land Suitability map of Cashew	64
7.16	Land Suitability map of Custard Apple	65
7.17	Land Suitability map of Amla	66
7.18	Land Suitability map of Tamarind	67
7.19	Land Management Units map of Nirgudi-1 microwatershed	68
9.1	Soil and Water Conservation map of Nirgudi-1 microwatershed	80

EXECUTIVE SUMMARY

The land resource inventory of Nirgudi-1 microwatershed was conducted using village cadastral maps and IRS satellite imagery on 1:7920 scale. The false colour composites of IRS imagery were interpreted for physiography and these physiographic delineations were used as base for mapping soils. The soils were studied in several transects and a soil map was prepared with phases of soil series as mapping units. Random checks were made all over the area outside the transects to confirm and validate the soil map unit boundaries. The soil map shows the geographic distribution and extent, characteristics, classification and use potentials of the soils in the microwatershed.

The present study covers an area of 808 ha in Nirgudi-1 microwatershed in Aland taluk of Kalaburgi district, Karnataka. The climate is semiarid and categorized as drought prone with an average annual rainfall of 786 mm of which about 595 mm is received during south–west monsoon, 116mm during north-east and the remaining 75 mm during the rest of the year. An area of about 98 per cent is covered by soils, two per cent by waterbodies, settlements and others. The salient findings from the land resource inventory are summarized briefly below.

- ❖ The soils belong to 5 soil series and 15 soil phases (management units) and 5 land management units.*
- ❖ The length of crop growing period is about 150 days starting from the 3rd week of June to 1st week of October.*
- ❖ From the master soil map, several interpretative and thematic maps like land capability, soil depth, surface soil texture, soil gravelliness, available water capacity, soil slope and soil erosion were generated.*
- ❖ Soil fertility status maps for macro and micronutrients were generated based on the surface soil samples collected at every 250 m grid interval.*
- ❖ Land suitability for growing major agricultural and horticultural crops were assessed and maps showing degree of suitability along with constraints were generated.*
- ❖ About 94 per cent area is suitable for agriculture and 4 per cent is not suitable for agriculture but well suited for forestry, pasture, agroforestry, silvi-pasture, recreation, installation of wind mills and as habitat for wildlife.*
- ❖ About 20 per cent of the soils are very deep (>150 cm, 40 per cent are moderately shallow to shallow (25-75 cm) and about 38 per cent are very shallow (<25 cm) soils.*
- ❖ About 98 per cent of the area has clayey soils at the surface.*
- ❖ About 39 per cent of the area has non-gravelly soils, 33 per cent gravelly soils (15-35 % gravel) and 26 per cent very gravelly soils (35- 60% gravel).*
- ❖ About 20 per cent of the area has soils that are very high (>200mm/m) in available water capacity, 5 per cent medium (100-150 mm/m) and about 74 per cent low (50-100 mm/m) and very low (<50 mm/m).*
- ❖ About 71 per cent of the area has nearly level (0-1%) to very gently sloping (1-3%) lands and about 27 per cent area is gently (3-5%) to moderately sloping (5-10%) lands.*
- ❖ An area of about 20 per cent has soils that are slightly eroded (e1), 62 per cent moderately eroded (e2) and 15 per cent severely eroded (e3).*
- ❖ An area of about 53 per cent has soils that are moderately alkaline (pH 7.8 to 8.4), one per cent strongly alkaline (pH 8.4 to >9.0), about 34 per cent slightly alkaline (pH 7.3-7.8) and 9 per cent area is neutral (pH 6.5-7.3) in soil reaction.*
- ❖ The Electrical Conductivity (EC) of the soils are dominantly <2 dsm⁻¹ indicating that the soils are non-saline.*
- ❖ About 73 per cent medium (0.5-0.75%), 19 per cent high (>0.75%) and 6 per cent low (<0.5%) in organic carbon.*
- ❖ Major area of 93 per cent has soils that are low (<23 kg/ha) and 5 per cent medium (23-57 kg/ha) in available phosphorus.*

- ❖ About 19 per cent medium (145-337 kg/ha) and 79 per cent high (>337 kg/ha) in available potassium.
- ❖ Available sulphur is low (<10 ppm) in about 23 per cent area, medium (10-20 ppm) in 54 per cent and 21 per cent high (>20 ppm).
- ❖ Available boron is low (<0.5 ppm) in about 84 per cent area and 14 per cent medium (0.5-1.0 ppm).
- ❖ About 5 per cent area has soils that are deficient (<4.5 ppm) in available iron and 93 per cent sufficient (>0.6 ppm).
- ❖ Available manganese and copper are sufficient in all the soils.
- ❖ About 75 per cent area has soils that are deficient (<0.6 ppm) in available zinc and 23 per cent sufficient (>0.6 ppm).
- ❖ The land suitability for 18 major crops grown in the microwatershed were assessed and the areas that are highly suitable (S1) and moderately suitable (S2) are given below. It is however to be noted that a given soil may be suitable for various crops but what specific crop to be grown may be decided by the farmer looking to his capacity to invest on various inputs, marketing infrastructure, price and finally the demand and supply position.

Land suitability for various crops in the microwatershed

Crop	Suitability Area in ha (%)		Crop	Suitability Area in ha (%)	
	Highly suitable (S1)	Moderately suitable (S2)		Highly suitable (S1)	Moderately suitable (S2)
Sorghum	160 (20)	37 (5)	Sapota	-	-
Maize	-	-	Jackfruit	-	-
Red gram	-	197 (24)	Jamun	-	160 (20)
Sunflower	160 (20)	-	Musambi	160 (20)	-
Cotton	160 (20)	37 (5)	Lime	160 (20)	-
Sugarcane	-	-	Cashew	-	-
Soybean	160(20)	37 (5)	Custard apple	160 (20)	37(5)
Guava	-	-	Amla	160 (20)	37 (5)
Mango	-	-	Tamarind	-	1600)

Apart from the individual crop suitability, a proposed crop plan has been prepared for the 5 identified LMUs by considering only the highly and moderately suitable lands for different crops and cropping systems with food, fibre and horticulture crops that helps in maintaining the ecological balance in microwatershed

- ❖ Maintaining soil-health is vital to crop production and conserve soil and land resource base for maintaining ecological balance and to mitigate climate change. For this, several ameliorative measures have been suggested to these problematic soils like saline/alkali, highly eroded, sandy soils etc.,
- ❖ Soil and water conservation treatment plan has been prepared that would help in identifying the sites to be treated and also the type of structures required.
- ❖ As part of the greening programme, several tree species have been suggested to be planted in marginal and submarginal lands, field bunds and also in the hillocks, mounds and ridges.

INTRODUCTION

Soil being a vital natural resource on whose proper use depends the life supporting systems of a country and the socioeconomic development of its people. Soils provide food, fodder, fibre and fuel for meeting the basic human and animal needs. With the ever increasing growth in human and animal population, the demand on soil for more food and fodder production is on the increase. The area available for agriculture is about 51 per cent of the total geographical area and more than 60 per cent of the people are still dependant on agriculture for their livelihood. However, the capacity of a soil to produce is limited and the limits to the production are set by its intrinsic characteristics, agroclimatic setting, and, use and management. There is, therefore, tremendous pressure on land and water resources, which is causing decline in soil-health and stagnation in productivity. The soils have been degrading at an estimated rate of one million hectares per year and ground water levels have been receding at an alarming rate resulting in decline in the ground water resource. Further, land degradation has emerged as a serious problem which has already affected about 38 lakh ha of cultivated area in the State. Soil erosion alone has degraded about 35 lakh ha. Almost all the uncultivated areas are facing various degrees of degradation, particularly soil erosion; salinity and alkalinity has emerged as a major problem (>3.5 lakh ha) in the irrigated areas of the State. Nutrient depletion and declining factor productivity is common in both rainfed and irrigated areas. The degradation is continuing at an alarming rate and there appears to be no systematic effort among the stakeholders to contain this process. In recent times, an aberration of weather due to climate change phenomenon has added another dimension leading to unpredictable situations to be tackled by the farmers.

In this critical juncture, the challenge before us is not only to increase the productivity per unit area which is steadily declining and showing a fatigue syndrome, but also to prevent or at least reduce the severity of degradation. If the situation is not reversed at the earliest, then the sustainability of the already fragile crop production system and the overall ecosystem will be badly affected in the state.

Added to this, every year there is a significant diversion of farm lands and water resources for non-agricultural purposes. Thus, developing strategies to slow down the degradation process or reclaim the soils to normal condition and ensure sustainability of production system are the major issues today. This demands a systematic appraisal of our soil and land resources with respect to their extent, geographic distribution, characteristics, behaviour and use potential, which is very important for developing an effective land use and cropping systems for augmenting agricultural production on a sustainable basis.

The soil and land resource inventories made so far in Karnataka had limited utility because the surveys were of different types, scales and intensities carried out at different times with specific objectives. Hence, there is an urgent need to generate detailed site-specific farm level database on various land resources for all the villages/watersheds in a time

bound manner that would help to protect the valuable soil and land resources and also to stabilize the farm production. Therefore, the land resource inventory required for farm level planning is the one which investigates all the parameters which are critical for productivity viz., soils, site characteristics like slope, erosion, gravelliness and stoniness, climate, water, topography, geology, hydrology, vegetation, crops, land use pattern, animal population, socio-economic conditions, infrastructure, marketing facilities and various schemes and developmental works of the government etc. From the data collected at farm level, the specific problems and potentials of the area can be identified and highlighted, conservation measures required for the area can be planned on a scientific footing, suitability of the area for various uses can be worked out and finally viable and sustainable land use options suitable for each and every land holding can be prescribed.

The Land Resource Inventory is basically done for identifying potential and problem areas, developing sustainable land use plans, estimation of surface run off and water harvesting potential, preparation of soil and water conservation plans, land degradation/desertification etc. The Bureau is presently engaged in developing an LRI methodology using high resolution satellite remote sensing data and Digital Elevation Model (DEM) data to prepare Landscape Ecological Units (LEU) map representing agro-ecosystem as a whole. The LEU is preferred over landform as the base map for LRI. LEU is the assemblage of landform, slope and land use. An attempt has already been made to upscale the soil resource information from 1:250000 and 1:50000 scale to the LEU map in Goa and in some other states. Here, an attempt will be made later to uplink the LRI data generated under Sujala-III Project to the Landscape Ecological Units (LEUs) map. For this, the major physiographic region, *i.e.*, South Deccan Plateau will be taken as an example.

The land resource inventory aims to provide site specific database for Nirgudi-1 microwatershed in Aland Taluk, Kalaburgi District, Karnataka state for the Karnataka Watershed Development Department. The database was generated by using cadastral map of the village as a base along with high resolution IRS LISS IV and Cartosat-1 merged satellite imagery. Later, an attempt will be made to uplink this LRI data generated at 1:7920 scale under Sujala-III Project to the proposed Landscape Ecological Units (LEUs) map.

The study was organized and executed by the ICAR- National Bureau of Soil Survey and Land Use Planning, Regional Centre, Bangalore under Generation of Land Resource Inventory Data Base Component-1 of the Sujala-III Project funded by the World Bank.

GEOGRAPHICAL SETTING

2.1 Location and Extent

The study area of Nirgudi-1 microwatershed (Alur subwatershed) is located in the northeastern part of Karnataka in Aland Taluk, Kalaburgi District, Karnataka State (Fig.2.1). It comprises of part of a Nirgudi and Padsavali villages. It lies between $17^{\circ} 37'$ and $17^{\circ} 39'$ north latitude and between $77^{\circ} 26'$ and $77^{\circ} 27'$ east longitude and covers an area of 808 ha. It is about 15 km from Aland taluk and is surrounded by Chincholi Budruk on west, Chincholi Khurd on southwest, Nagalgaon on southeast, Matki village on east and Maharashtra state on the northern side.




Fig.2.1 Location map of Nirgudi-1 Microwatershed

2.2 Geology

Major rock formation observed in the microwatershed is Basalt (Fig.2.2) or Deccan Trap. The Deccan Traps cover the whole of Bidar, parts of Kalaburgi, Bijapur and Belgaum districts. In all, eight lava flows have been identified in Karnataka horizontally overlying the older formations. The thickness of the individual flows averages about five meters. It is relatively uniform in petrographic character. The most common type is augite basalt. Dominant colour is grayish green and texture ranges from cryptocrystalline to glassy. The rock is often vesicular and scoriaceous filled up with secondary minerals like coloured agate, quartz, calcite and a large variety of zeolites. The Deccan Traps form an excellent building material and also used as road-metal and railway ballast.



Fig. 2.2 Basalt rock formation

2.3 Physiography

Physiographically, the area has been identified as basalt landscape based on geology. The area has been further subdivided into four landforms, viz; mounds/ridges, summits, side slopes and very gently sloping uplands based on slope and its relief features. The elevation ranges from 506 to 546 m. The mounds and ridges are mostly covered by rock outcrops.

2.4 Drainage

The area is drained by several small parallel streams that join Monia nala which further down stream joins Awarja river along its course. Though, it is not a perennial one, during rainy season it carries large quantities of rain water. The microwatershed has only few small tanks which are not capable of storing the water that flows during the rainy season. Due to this, the ground water recharge is very much affected. This is reflected in the failure of many bore wells in the villages. If the available rain water is properly harnessed by constructing new tanks and recharge structures at appropriate places in the village, then the drinking and irrigation needs of the entire area can be easily met. The drainage network is parallel to sub parallel and dendritic.

2.5 Climate

The Kalaburgi district lies in the northern plains of Karnataka and falls under semi arid tract of the state and is categorized as drought prone with average annual rainfall of 743 mm (Table 2.1). Of the total rainfall, maximum of 595 mm is received during the south-west monsoon period from June to September, the north-east monsoon from October to early December contributes about 116 mm, and the remaining 75 mm during the rest of the year. December is the coldest month with mean daily maximum and minimum temperatures being 29.5⁰C and 15⁰ to 10⁰C respectively. During peak summer, temperatures shoot up to 45⁰C. Relative humidity varies from 26 per cent in summer to 62 per cent in winter. Rainfall distribution is shown in Figure 2.3. The average potential evapotranspiration (PET) is 150 mm and varies from a low of 115 mm in December to 232 mm in the month of May. The PET is always higher than precipitation in all the months except August and September. Generally, the length of crop growing period (LGP) is 150 days and starts from 3rd week of June to third week of November.

Table 2.1 Mean Monthly Rainfall, PET, 1/2 PET at Aland Taluk, Kalaburgi District

Sl.No.	Months	Rainfall	PET	1/2 PET
1	January	7.50	126.80	63.40
2	February	3.40	143.90	71.95
3	March	11.30	189.90	94.95
4	April	19.40	209.80	104.90
5	May	32.70	232.20	116.10
6	June	111.00	186.40	93.20
7	July	139.20	152.80	76.40
8	August	172.40	147.60	73.80
9	September	172.30	131.70	65.85
10	October	91.30	145.50	72.75
11	November	19.30	129.80	64.90
12	December	5.80	114.80	57.40
Total		785.6	149.70	




Fig 2.3 Rainfall distribution in Aland Taluk, Kalaburgi District

2.6 Natural Vegetation

The natural vegetation is sparse comprising few tree species, shrubs and herbs. The mounds, ridges and boulders occupy very sizeable area which are under thin to moderately thick forest vegetation. Still, there are some remnants of the past forest cover which can be seen in patches in some ridges and hillocks in the microwatershed (Fig. 2.4).

Apart from the continuing deforestation, the presence of large population of goats, sheep and other cattle in the microwatershed is causing vegetative degradation of whatever little vegetation left in the area. The uncontrolled grazing has left no time for the regeneration of the vegetative cover. This leads to the accelerated rate of erosion on the hill slopes resulting in the formation of deep gullies in the foot slopes that eventually result in the heavy siltation of tanks and reservoirs in the microwatershed.



Fig. 2.4 Natural Vegetation (Scrub) of Nirgudi-1 Microwatershed

2.7 Land Utilization

About 89 per cent area (Table 2.2) in Aland taluk is cultivated at present. An area of about 2 per cent is permanently under pasture, 3 per cent under current fallows and 2 per cent each under nonagricultural land and currently barren. Forests occupy an area of about 2 per cent and the tree cover is in a very poor state. Most of the mounds, ridges and bouldery areas have very poor vegetative cover. Major crops grown in the area are sorghum, maize, cotton, sugarcane, red gram and sapota. While carrying out land resource inventory, the land use/land cover particulars are collected from all the survey numbers and a current land use map of the microwatershed is prepared. The current land use map prepared shows the arable and non-arable lands, other land uses and different types of crops grown in the area. The current land use map of Nirgudi-1 microwatershed is presented in Fig.2.5. Simultaneously, enumeration of wells (bore wells and open wells) in the microwatershed was made and their location in different survey numbers is located on the cadastral map. Map showing the location of wells and other water bodies in the Nirgudi-1 microwatershed is given in Fig.2.6.

Table 2.2 Land Utilization in Aland Taluk

Sl. No.	Agricultural land use	Area (ha)	Per cent
1.	Total geographical area	173417	
2.	Total cultivated area	153806	88.69
3.	Area sown more than once	7910	
4.	Trees and grooves	59	0.034
5.	Forest	2854	1.64
6.	Cultivable wasteland	974	0.56
7.	Permanent Pasture land	3469	2.00
8.	Barren land	3142	1.81
9.	Non- Agriculture land	3465	1.99
10.	Current Fallows	5648	3.25




Fig.2.5 Current Land Use map of Nirgudi-1 Microwatershed




Fig.2.6 Location of Wells map of Nirgudi-1 Microwatershed

SURVEY METHODOLOGY

The purpose of land resource inventory is to delineate similar areas (soil series and phases), which respond or expected to respond similarly to a given level of management. This was achieved in Nirgudi -1 microwatershed by the detailed study of all the soil characteristics (depth, texture, colour, structure, consistence, coarse fragments, porosity, soil reaction, soil horizons etc.) and site (slope of the land, erosion, drainage, occurrence of rock fragments etc.) followed by grouping of similar areas based on soil-site characteristics into homogeneous (management units) units and showing their area extent and geographic distribution on the microwatershed cadastral map. The detailed survey at 1:7920 scale was carried out in 808 ha area. The methodology followed for carrying out land resource inventory was as per the guidelines given in Soil Survey Manual (IARI, 1971; Soil Survey Staff, 2006; Natarajan *et al.*, 2015) which is briefly described below.

3.1 Base Maps

The detailed survey of the land resources occurring in the microwatershed was carried out by using digitized cadastral map as a base. The cadastral map shows field boundaries with their survey numbers, location of tanks, streams and other permanent features of the area (Fig. 3.1). Apart from the cadastral map, remote sensing data products from Cartosat-1 and LISS-IV merged at the scale of 1:7920 were used in conjunction with the cadastral map to identify the rock types, the landscapes, landforms and other surface features. The imagery helped in the identification and delineation of boundaries between hills, uplands and lowlands, water bodies, forest and vegetated areas, roads, habitations and other cultural features of the area (Fig. 3.2). The cadastral map was overlaid on the satellite imagery (Fig.3.3) that helps to identify the parcel boundaries and other permanent features. Apart from cadastral maps and images, toposheets of the area (1:50,000 scale) were used for initial traversing, identification of geology and landforms, drainage features, present land use and also for selection of transects in the microwatershed.




Fig 3.1 Scanned and Digitized Cadastral map of Nirgudi-1 Microwatershed




Fig.3.2 Satellite Image of Nirgudi-1 Microwatershed




Fig.3.3 Cadastral map overlaid on IRS PAN+LISS IV merged imagery of Nirgudi-1 Microwatershed

3.2 Field Investigation

Preliminary traverse of the microwatershed was carried out with the help of cadastral map, imagery and toposheets. While traversing, landforms and physiographic units identified were checked and preliminary soil legend was prepared by studying soils at a few selected places. The field boundaries and survey numbers given on the cadastral sheet were located on the ground by following permanent features like roads, cart tracks, nallas, streams, tanks etc., and wherever changes were noticed, they were incorporated on the microwatershed cadastral map. Then, intensive traversing of each physiographic unit like hills, ridges and uplands was carried out. Based on the variability observed on the surface, transects were selected across the slope covering all the landform units in the microwatershed (Natarajan and Dipak Sarkar, 2010).

In the selected transect, soil profiles were located at closely spaced intervals to take care of any change in the land features like break in slope, erosion, gravel, stones etc. In the selected sites, profiles (vertical cut showing the soil layers from surface to the rock) were opened up to 200 cm or to the depth limited by rock or hard substratum and studied in detail for all their morphological and physical characteristics. The soil and site characteristics were recorded for all profile sites on a standard proforma as per the guidelines given in USDA Soil Survey Manual (Soil Survey Staff, 2012). Apart from the transect study, profiles were also studied at random, almost like in a grid pattern, outside the transect areas.

Based on the soil-site characteristics, the soils were grouped into different soil series (soil series is the most homogeneous unit having similar horizons and properties and behaves similarly for a given level of management). Soil depth, texture, colour, kind of horizon and horizon sequence, amount and nature of gravel present, nature of substratum etc, were used as the major differentiating characteristics for identifying soil series occurring in the area. The differentiating characteristics used for identifying soil series are given in Table 3.1. Based on the above characteristics, 5 soil series were identified in the Nirgudi-1 microwatershed.

Table 3.1 Differentiating Characteristics used for Identifying Soil Series
(Characteristics are of Series Control Section)

SOILS OF BASALT LANDSCAPE							
Sl. No.	Soil Series	Depth (cm)	Colour	Texture	Gravel (%)	Horizon sequence	Calcareousness
1	Margutti (MGT)	<25	10YR3/3,4/3,5/ 7.5YR4/3	c	15-35	Ap- cr	-
2	Novinihala (NHA)	25-50	10YR3/2,3/1,4/2 7.5YR3/4	c	<15	Ap-Bw- cr/R	-
3	Bhimanahalli (BHI)	25-50	10YR3/2,3/3,3/1 7.5YR3/2,4/2	c	15-35	Ap-Bw- cr/R	-
4	Dinsi (DSI)	50-75	10YR3/2,3/3,4/3 3/2	c	<15	Ap-BA- Bss	-
5	Mahagaon (MAN)	>150	10YR3/2,3/1	c	<15	Ap-BA- Bss	-

3.3 Laboratory Characterization

Soil samples were collected from representative master profiles for laboratory characterization by following the methods outlined in the Laboratory Manual (Sarma *et al*, 1987). Surface soil samples collected from farmer's fields (127 samples) for fertility status (major and micronutrients) at 250 m grid interval were analyzed in the laboratory. (Katyal and Rattan, 2003) By linking the soil fertility data to the survey numbers through GIS, soil fertility maps for 11 elements including pH and EC were generated for the microwatershed.

3.4 Finalization of Soil Maps

The area under each soil series was further separated and mapped as soil phases and their boundaries delineated on the cadastral map based on the variations observed in the texture of the surface soil, slope, erosion, presence of gravel, stoniness etc. A soil phase is a subdivision of soil series based mostly on surface features that affect its use and management.

The soil mapping units are shown on the map (Fig.3.4) in the form of symbols. During the survey about 25 profile pits, few minipits and a few auger bores representing different landforms occurring in the microwatershed were studied. All the profile locations

are indicated on the village cadastral map in the form of a triangle. In addition to the profile study, spot observations in the form of minipits, road cuts, terrace cuts etc., were studied to validate the soil boundaries on the soil map. The soil map shows the geographic distribution of 15 mapping units representing 5 soil series occurring in the microwatershed. The soil map unit (soil legend) description is presented in Table 3.2.

The soil phase map (management units) shows the distribution of 15 phases identified and mapped in the microwatershed. Each mapping unit (soil phase) delineated on the map has similar soil and site characteristics. In other words, all the farms or survey numbers included in one phase will have similar management needs and they have to be treated accordingly.

The 15 soil phases identified and mapped in the microwatershed were regrouped into 5 Land Management Units (LMU's) for the purpose of preparing a proposed crop plan for sustained development of the microwatershed. The database (soil phases) generated under LRI was utilized for identifying Land Management Units (LMUs) based on the management needs. One or more than one soil site characteristic having influence on the management have been chosen for identification and delineation of LMUs. For Nirgudi-1 microwatershed, five soil and site characteristics, namely soil depth, soil texture, slope, erosion and gravel content have been considered for defining LMUs. The land management units are expected to behave similarly for a given level of management.




Fig 3.4 Soil phase or management units map of Nirgudi-1 Microwatershed

Table 3.2 Soil Legend

Soil map unit no.	Soil series	Soil Phase	Mapping Unit Description	Area in ha (%)
Soils of Basalt Landscape				
	MGT	Margutti soils are very shallow (<25cm), well drained, have very dark grayish brown to dark brown clayey soils occurring on nearly level to moderately sloping uplands		304.4 (37.6)
1		MGTmB2g1	Clay surface, 1-3 % slope, moderate erosion, gravelly (15-35%)	45.36 (5.61)
2		MGTmB2g2	Clay surface, 1-3 % slope, moderate erosion, very gravelly (35-60%)	52.01 (6.43)
3		MGTmC2g1	Clay surface, 3-5 % slope, moderate erosion, gravelly (15-35%)	57.37 (7.10)
4		MGTmC2g2	Clay surface, 3-5 % slope, moderate erosion, very gravelly (35-60%)	26.17 (3.24)
5		MGTmC3g2	Clay surface, 3-5 % slope, severe erosion, very gravelly (35-60%)	89.60 (11.08)
6		MGTmD3g2	Clay surface, 5-10 % slope, severe erosion, very gravelly (35-60%)	33.87 (4.19)
	NHA	Novinihala soils are shallow (25-50 cm), well drained, have very dark grayish brown to dark brown clayey soils occurring on very gently sloping uplands		276.2 (34.18)
7		NHAmB1g1	Clay surface, slope 1-3%, slight erosion gravelly (15-35%)	4.79 (0.59)
8		NHAmB2	Clay surface, slope 1-3%, moderate erosion	114.38 (14.15)
9		NHAmB2g1	Clay surface, slope 1-3%, moderate erosion, gravelly (15-35%)	133.98 (16.57)
10		NHAmB2g2	Clay surface, slope 1-3%, moderate erosion, gravelly (35-60%)	10.91 (1.35)
11		NHAmC2g1	Clay surface, slope 3-5%, moderate erosion, gravelly (15-35%)	10.64 (1.32)
12		NHAmC2g3	Clay surface, slope 3-5%, moderate erosion, extremely gravelly (60-80%)	1.60 (0.20)
	BHI	Bhimanahalli soils area shallow (25-50 cm), well drained, have very dark gray to brown clay soils occurring on very gently sloping to gently sloping uplands.		14.41 (1.78)
13		BHImB2g1	Clay surface, slope 1-3%, moderate erosion, gravelly (15-35%)	14.41 (1.78)

	DSI	Dinsi soils are moderately shallow (50-75 cm), moderately well drained, have very dark gray to brown clayey soils occurring on very gently sloping uplands		36.73 (4.54)
14		DSImB2	Clay surface, slope 1-3%, moderate erosion	36.73 (4.54)
	MAN	Mahagaon soils are very deep (>150 cm), moderately well drained, have very dark gray to very dark grayish brown cracking clay soils occurring on nearly level to very gently sloping uplands		160.33 (19.83)
15		MANmA1	Clay surface, slope 0-1%, slight erosion	160.33 (19.83)
Miscellaneous Lands				
16	Water body			16.33 (2.02)

THE SOILS

Detailed information pertaining to the nature, extent and distribution of different kinds of soils occurring in Nirgudi-1 microwatershed is provided in this chapter. The microwatershed area has been identified as basalt landscape. In all, 5 soil series were identified in this landscape. Soil formation is the result of the combined effect of environmental and terrain factors that are reflected in soil morphology. In the basalt landscape, it is by parent material and climate. A brief description of each of the 5 soil series identified followed by 15 soil phases (management units) mapped under each series are furnished below. The soils in any one map unit differ from place to place in their depth, texture, slope, gravelliness, erosion or any other site characteristics that affect management. The soil phase map can be used for identifying the suitability of areas for growing specific crops or for other alternative uses and also for deciding the type of conservation structures needed. The detailed information on soil and site-characteristics like soil depth, surface soil texture, slope, erosion, gravelliness, AWC, LCC etc, with respect to each of the soil phase identified is given village/survey number wise for the microwatershed in Appendix-I.

4.1 Soils of Basalt Landscape

In this landscape, 5 soil series are identified and mapped. Of these, Margutti (MGT) soil series occupies maximum area of about 304 ha (38%). The brief description of each series along with the soil phases identified and mapped is given below.

4.1.1 Margutti (MGT) Series: Marguti soils are very shallow (<25cm), well drained, have very dark grayish brown to dark brown clay soils. They have developed from basalt and occur on very gently sloping to moderately sloping uplands.

The total depth of the soil ranges from 10 to 23 cm. The thickness of A horizon ranges from 7 to 18 cm. Its colour is in 10 YR and 7.5 YR hue with value 3 to 4 and chroma 2 to 4. The texture is clay with 15 to 35 per cent gravel. The available water capacity is very low (<50 mm/m).

Six phases were identified:

MGTmB2g1	Clay surface, slope 1-3 %, moderate erosion, gravelly (15-35%)
MGTmB2g2	Clay surface, slope 1-3 %, moderate erosion, very gravelly (35-60 %)
MGTmC2g1	Clay surface, slope 3-5%, moderate erosion, gravelly (15-35%)
MGTmC2g2	Clay surface, slope 3-5%, moderate erosion, very gravelly (35-60%)
MGTmC3g2	Clay surface, slope 3-5%, severe erosion, very gravelly (35-60%)
MGTmD3g2	Clay surface, 5-10 % slope, severe erosion, very gravelly (35-60%)



Landscape and Soil Profile Characteristics of Margutti (MGT) Series

4.1.2 Novanihala (NHA) Series: Novanihala soils are shallow (25-50 cm), well drained, have very dark grayish brown to dark brown clay soils. They have developed from basalt and occur on very gently to gently sloping uplands.

The thickness of the solum ranges from 27 to 48 cm. The thickness of A horizon ranges from 12 to 20 cm. Its colour is in 7.5 YR and 10 YR hue with value 3 to 4 and chroma 2 to 4. The texture varies from sandy clay to clay with 10 to 20 per cent gravel. The thickness of B horizon ranges from 22 to 37 cm. Its colour is in 10 YR and 7.5 YR hue with value 3 to 4 and chroma 2 to 4. Its texture is clay with gravel content of 10-15 per cent. The available water capacity is low (51-100 mm/m).

Six phases were identified:

NHAmB1g1	Clay surface, slope 1-3%, slight erosion, gravelly (15-35%)
NHAmB2	Clay surface, slope 1-3%, moderate erosion
NHAmB2g1	Clay surface, slope 1-3%, moderate erosion, gravelly (15-35%)
NHAmB2g2	Clay surface, slope 1-3%, moderate erosion, very gravelly (35-60%)
NHAmC2g1	Clay surface, slope 3-5%, moderate erosion, gravelly (15-35%)
NHAmC2g3	Clay surface, slope 3-5%, moderate erosion, extremely gravelly (60-80)%



Landscape and Soil Profile Characteristics of Novanihala (NHA) Series

4.1.3 Bhimanahalli (BHI) Series: Bhimanahalli soils are shallow (25-50 cm), well drained, have very dark gray to brown clay soils. They have developed from basalt and occur on very gently sloping uplands.

The thickness of the solum ranges from 29 to 48 cm. The thickness of A horizon ranges from 15 to 20 cm. Its colour is in 7.5 YR and 10 YR hue with value 3 to 4 and chroma 2 to 4. The texture varies from sandy clay to clay with 15 to 25 per cent gravel. The thickness of B horizon ranges from 23 to 33 cm. Its colour is in 10 YR and 7.5 YR hue with value 3 to 4 and chroma 1 to 3. Its texture is clay with gravel content of 15 to 35 per cent. The available water capacity is very low (<50 mm/m).

Only one phases was identified:

BHImB2g1	Clay surface, slope 1-3%, moderate erosion, gravelly (15-35%)
----------	---



Landscape and Soil Profile Characteristics of Bhimanahalli (BHI) Series

4.1.4 Dinsi (DSI) Series: Dinsi soils are moderately shallow (50-75 cm), moderately well drained, have very dark gray to brown clay soils. They have developed from basalt and occur on very gently sloping uplands.

The thickness of the solum ranges from 51 to 71 cm. The thickness of A horizon ranges from 9 to 24 cm. Its colour is in 10 YR hue with value 3 and chroma 1 to 3. The texture is clay with 5 to 10 per cent gravel. The thickness of B horizon ranges from 27 to 62 cm. Its colour is in 10 YR hue with value 3 to 4 and chroma 2 to 4. Its texture is clay with gravel content of less than 15 per cent. The available water capacity is medium (101-150 mm/m).

Only one phase was identified:

DSImB2	Clay surface, slope 1-3%, moderate erosion
--------	--



Landscape and Soil Profile Characteristics of Dinsi (DSI) Series

4.1.5 Mahagaon (MAN) Series: Mahagaon soils are very deep (>150 cm), moderately well drained, have very dark gray to very dark grayish brown cracking clay soils. They have developed from basalt and occur on nearly level lands.

The thickness of the solum ranges from 152 to 195 cm. The thickness of A horizon ranges from 18 to 22 cm. Its colour is in 10 YR and 7.5 YR hue with value 3 to 4 and chroma 1 to 3. The texture is clay with less than 10 per cent gravel. The thickness of B horizon ranges from 130 to 160 cm. Its colour is in 10 YR hue with value 3 to 4 and chroma 1 to 2. Its texture is clay with gravel content of less than 15 per cent. The available water capacity is very high (>200 mm/m).

Only one phase was identified:

MANmA1	Clay surface, 0-1% slope, slight erosion
--------	--



Landscape and Soil Profile Characteristics of Mahagaon (MAN) Series

INTERPRETATION FOR LAND RESOURCE MANAGEMENT

The most important soil and site characteristics that affect the land use and conservation needs of an area are land capability, soil depth, soil texture, coarse fragments, available water capacity, soil slope, soil erosion, soil reaction etc. These are interpreted from the data base generated through land resource inventory and several thematic maps are generated. These would help in identifying the areas suitable for growing crops and, soil and water conservation measures and structures needed thus helping to maintain good soil health for sustained crop production. The various thematic maps generated are described below.

5.1 Land Capability Classification

Land capability classification is an interpretative grouping of soil map units (soil phases) mainly based on inherent soil characteristics, external land features and environmental factors that limit the use of land for agriculture, pasture, forestry, or other uses on a sustained basis (IARI, 1971). The land and soil characteristics used to group the land resources in an area into various land capability classes, subclasses and units are

Land characteristics: Slope, erosion, drainage, rock outcrops.

Climate: Total rainfall and its distribution, and length of crop growing period.

The Land capability classification system is divided into land capability classes, subclasses and units based on the level of information available. Eight land capability classes are recognized. They are

Class I: They are very good lands that have no limitations or very few limitations that restrict their use.

Class II: They are good lands that have minor limitations and require moderate conservation practices.

Class III: They are moderately good lands that have moderate limitations that reduce the choice of crops or that require special conservation practices.

Class IV: They are fairly good lands that have very severe limitations that reduce the choice of crops or that require very careful management.

Class V: Soils in these lands are not likely to erode, but have other limitations like wetness that are impractical to remove and as such not suitable for agriculture, but suitable for pasture or forestry with minor limitations.

Class VI: The lands have severe limitations that make them generally unsuitable for cultivation, but suitable for pasture or forestry with moderate limitations.

Class VII: The lands have very severe limitations that make them unsuitable for cultivation, but suitable for pasture or forestry with major limitations.

Class VIII: Soil and other miscellaneous areas (rock lands) that have very severe limitations that nearly preclude their use for any crop production, but suitable for wildlife, recreation and wind mills.

The land capability subclasses are recognised based on the dominant limitations observed within a given land capability class. The subclasses are designated by adding a lower case letter like 'e', 'w', 's', or 'c' to the class numeral. The subclass "e" indicates that the main hazard is risk of erosion, "w" indicates drainage or wetness as a limitation for plant growth, "s" indicates shallow soil depth, coarse or heavy textures, calcareousness, salinity/alkalinity or gravelliness and "c" indicates limitation due to climate.

The land capability subclasses have been further subdivided into land capability units based on the kinds of limitations present in each subclass. Ten land capability units are used in grouping the soil map units. They are stony or rocky (0), erosion hazard (slope, erosion) (1), coarse texture (sand, loamy sand, sandy loam) (2), fine texture (cracking clay, silty clay) (3), slowly permeable subsoil (4), coarse underlying material (5), salinity/alkali (6), stagnation, overflow, high ground water table (7), soil depth (8) and fertility problems (9). The capability units thus identified have similar soil and land characteristics that respond similarly to a given level of management. The soils of the microwatershed have been classified upto land capability subclass level only.

The 15 soil map units identified in the Nirgudi -1 microwatershed are grouped under 4 land capability classes and 5 land capability subclasses. About 94 per cent area in the microwatershed is suitable for agriculture (Fig. 5.1) and 4 per cent is not suitable for agriculture but well suited for grazing or forestry, recreation and wildlife.

Good cultivable lands (Class II) cover about 24 per cent area and are distributed in the northern, southern, central and southwestern part of the micowatershed with minor problems of soil. Moderately good cultivable lands (Class III) cover maximum area of about 58 per cent and are distributed in all parts of the microwatershed with moderate problems of erosion and soil. The fairly good cultivable lands (class IV) cover only about 11 per cent area. They have severe limitations of erosion and soil and are distributed in the northern, northeastern and central part of the microwatershed.

The class VI lands cover about 4 per cent and are distributed in the southwestern and northwestern part of the microwatershed. They are well suited for pasture, forestry, wild life and recreation.




Fig. 5.1 Land Capability map of Nirgudi-1 Microwatershed

5.2 Soil Depth

Soil depth refers to the depth of the soil occurring above the parent material or hard rock. The depth of the soil determines the effective rooting depth for plants and in accordance with soil texture, mineralogy and gravel content, the capacity of the soil column to hold water and nutrient availability. Soil depth is one of the most important soil characteristic that is used in differentiating soils into different soil series. The soil depth classes used in identifying soils in the field are very shallow (<25 cm), shallow (25-50 cm), moderately shallow (50-75 cm), moderately deep (75-100 cm), deep (100-150 cm) and very deep (>150 cm). They were used to classify the soils into different depth classes and a soil depth map was prepared (Fig. 5.2).

Very deep soils (>150 cm) occur in about 160 ha (20%) and are distributed in the northern, southern and central part of the microwatershed. Moderately shallow (50-75 cm) soils occupy a small area of about 37 ha (5%) and are distributed in the southwestern part of the microwatershed.

Maximum area of about 304 ha (38%) is under very shallow soils (<25 cm) and are distributed in all parts of the microwatershed. Shallow soils (25-50 cm) occupy an area of about 291 ha (36%) in the eastern, southeastern, northeastern and northern part of the microwatershed.

The most productive lands of about 160 ha (20%) with respect to soil rooting depth where all climatically adapted annual and perennial crops can be grown are very deep soils (>150 cm depth) occurring in the northern, southern and central part of the microwatershed.

The most problem lands with an area of about 595 ha (73%) having very shallow (<25 cm) and shallow (25-50 cm) rooting depth occur in all parts of the microwatershed. They are not suitable for growing agricultural crops but well suited for pasture, forestry or other recreational purposes. Occasionally, short duration crops may be grown if rainfall is normal.




Fig. 5.2 Soil Depth map of Nirgudi-1 Microwatershed

5.3 Surface Soil Texture

Texture is an expression to indicate the coarseness or fineness of the soil as determined by the relative proportion of primary particles of sand, silt and clay. It has a direct bearing on the structure, porosity, adhesion and consistence. The surface layer of a soil to a depth of about 25 cm is the layer that is most used by crops and plants. The surface soil textural class provides a guide to understanding soil-water retention and availability, nutrient holding capacity, infiltration, workability, drainage, physical and chemical behaviour, microbial activity and crop suitability.

The entire area of about 792 ha (98%) has soils that are clayey in surface soil texture. The most productive lands (98%) with respect to surface soil texture are the clayey soils (Fig. 5.3) that have high potential for soil-water retention and availability, and nutrient retention

and availability, but have more problems of drainage, infiltration, workability and other physical problems.




Fig. 5.3 Surface Soil Texture map of Nirgudi-1 Microwatershed

5.4 Soil Gravelliness

Gravel is the term used for describing coarse fragments between 2 mm and 7.5 cm diameter and stones for those between 7.5 cm and 25 cm. The presence of gravel and stones in soil reduces the volume of soil responsible for moisture and nutrient storage, drainage, infiltration and runoff and hinders plant growth by impeding root growth and seedling emergence, intercultural operations and farm mechanization.

Maximum area has soils that are nongravelly (<15%) covering about 311 ha (39%) and are distributed in the southeastern, southwestern, western and northern parts the microwatershed (Fig.5.4).

About 267 ha (33%) area in the micro watershed has soils that are gravelly (15-35%) and are distributed in the northeastern, northwestern, southwestern, eastern and western part of the microwatershed followed by soils that are very gravelly (35-60%) covering about 213 ha (26%). They are distributed in the northeastern and northwestern part of the microwatershed. A very small area has soils that are extremely gravelly (60-80%).

The most productive lands with respect to gravelliness are found to be 39 per cent. They are nongravelly with less than 15 per cent gravel and have potential for growing both annual and perennial crops.

The problem soils that are very gravelly (35-60%) are found to cover about 26 per cent area, where only short duration crops can be grown.




Fig. 5.4 Soil Gravelliness map of Nirgudi-1 Microwatershed

5.5 Available Water Capacity

The soil available water capacity (AWC) is estimated based on the ability of the soil column to retain water between the tensions of 0.33 and 15 bar in a depth of 100 cm or the entire solum if the soil is shallower. The AWC of the soils (soil series) as estimated by considering the soil texture, mineralogy, soil depth and gravel content (Sehgal *et al.*, 1990) and accordingly the soil map units were grouped into five AWC classes *viz.* very low (<50 mm/m), low (50-100 mm/m), medium (100-150 mm/m), high (150-200 mm/m) and very high (>200 mm/m) and using these values, an AWC map was prepared (Fig. 5.5).

Major area of about 319 ha (39%) has soils that are very low (<50 mm/m) in available water capacity and are distributed in northwestern, central and northeastern part of the microwatershed. An area of about 276 ha (34%) has soils that are low (51-100 mm/m) in available water capacity and are distributed in the southeastern and eastern part of the microwatershed. Small area in the microwatershed has soils that are medium (101-150 mm/m) in available water capacity. They occur in about 37 ha (5%) and are distributed in the southwestern part of the microwatershed. The soils that are very high (>200 mm/m) in AWC covering about 160 ha (20%) in the microwatershed. They are distributed in the northern part of the microwatershed.

An area of about 160 ha (20%) has soils that have very high potential (>200 mm/m) with regard to available water capacity and are distributed in the central part of the microwatershed. In these areas, if the rainfall is normal and well distributed, all climatically adapted long duration annual and perennial crops can be grown.

About 595 ha (74%) area in the microwatershed has soils that are problematic with regard to available water capacity. Here, only the short or medium duration crops can be grown and the probability of crop failure is very high. These areas are best put to other alternative uses.




Fig. 5.5 Soil Available Water Capacity map of Nirgudi-1 Microwatershed

5.6 Soil Slope

Soil slope refers to the inclination of the surface of the land. It is defined by gradient, shape and length, and is an integral feature of any soil as a natural body. Slope is considered important in soil genesis, land use and land development. The length and gradient of slope influences the rate of runoff, infiltration, erosion and deposition. The soil map units were grouped into four slope classes and a slope map was prepared showing the area extent and geographic distribution of different slope classes in the microwatershed (Fig. 5.6).

Major area of the microwatershed falls under very gently sloping (1-3% slope) slope class. It covers an area of about 413 ha (51%) and is distributed in all parts of the microwatershed. An area of about 185 ha (23 %) in the microwatershed falls under gently sloping (3-5%) slope class and is distributed in the northwestern, northeastern and southern part of the microwatershed. Moderately sloping (5-10% slope) slope class cover a minor area

of about 34 ha (4%) and is distributed in the southeastern part of the microwatershed. Nearly level lands (0-1%) slope class covers about 160 ha (20%) and are distributed in the northern, southern and central parts of the microwatershed.

An area of about 573 ha (71%) in the microwatershed has soils that have high potential in respect of soil slopes. In these areas, all climatically adapted annual and perennial crops can be grown without much soil and water conservation and other land development measures.




Fig. 5.6 Soil Slope map of Nirgudi-1 Microwatershed

5.7 Soil Erosion

Soil erosion refers to the wearing away of the earth's surface by the forces of water, wind and ice involving detachment and transport of soil by raindrop impact. It is used for accelerated soil erosion resulting from disturbance of the natural landscape by burning, excessive grazing and indiscriminate felling of forest trees and tillage, all usually by man. The erosion classes showing an estimate of the current erosion status as judged from field observations in the form of rills, gullies or a carpet of gravel on the surface are recorded. Four erosion classes, viz, slight erosion (e1), moderate erosion (e2), severe erosion (e3) and very severe erosion (e4) are recognized. The soil map units were grouped into different erosion classes and soil erosion map was generated. The area extent and their spatial distribution in the microwatershed is given in Figure 5.7.

Soils that are slightly eroded (e1 class) cover a about area of about 165 ha (20%) and are distributed in the southern, northern and central part of the microwatershed. Soils that are moderately eroded (e2 class) cover maximum area of about 504 ha (62%) in the microwatershed and are distributed in all parts of the microwatershed. Severely eroded (e3 class) soils cover an area of about 123 ha (15%) and are distributed in all parts of the microwatershed.

Top priority is to be given to 123 ha area where they are severely eroded for taking up soil and water conservation and other land development measures followed by moderately eroded lands that cover about 504 ha.




Fig. 5.7 Soil Erosion map of Nirgudi-1 Microwatershed

FERTILITY STATUS

Soil fertility plays an important role in increasing crop yield. The adoption of high yielding varieties that require high amounts of nutrients has resulted in deficiency symptoms in crops and plants due to imbalanced fertilization and poor inherent fertility status. Hence, it is necessary to know the fertility (macro and micro nutrients) status of the soils of the watersheds for assessing the kind and amount of fertilizers required for each of the crop intended to be grown. For this purpose, the surface soil samples collected from the grid points (one soil sample at every 250 m interval) all over the microwatershed through land resource inventory in the year 2014 were analysed for pH, EC, organic carbon, available phosphorus and potassium, and for micronutrients like zinc, copper, iron and manganese, and secondary nutrient sulphur.

Soil fertility data generated has been assessed and individual maps for all the nutrients for the microwatershed have been prepared. The village/survey number wise fertility data for the microwatershed is given in Appendix-II.

6.1 Soil Reaction (pH)

The soil fertility analysis of the Nirgudi-1 microwatershed for soil reaction (pH) showed that about 72 ha (9.0%) area is neutral (pH 6.5-7.3) in reaction and is distributed in the northeastern, northwestern and southeastern part of the microwatershed (Fig.6.1). Slightly alkaline (pH 7.3-7.8) is around 76 ha (34%) area and is distributed in all part of the microwatershed. Maximum area of about 437 ha (53.4%) area is moderately alkaline (pH 7.8-8.4) and is distributed in all parts of the microwatershed. A small area of about 12 ha (1%) is under strongly alkaline (pH 8.4-9.0) and is distributed in the southern part of the microwatershed.

6.2 Electrical Conductivity (EC)

The Electrical Conductivity of the soils of the entire microwatershed area is $<2 \text{ dSm}^{-1}$ (Fig 6.2) and as such the soils in the microwatershed are nonsaline.

6.3 Organic Carbon

The soil organic carbon content of the soils in the microwatershed is medium (0.5-0.75%) in 591 ha (73%) area that are distributed in all parts of the microwatershed (Fig.6.3). High ($>0.75\%$) organic carbon content accounts for 152 ha (19%) area and is distributed in the northeastern, northwestern and central part of the microwatershed. Low ($<0.5\%$) organic carbon content accounts for a small area of 49 ha (6%) in the microwatershed and is distributed in the central and western part of the microwatershed.




Fig.6.1 Soil Reaction (pH) map of Nirgudi-1 Microwatershed




Fig.6.2 Electrical Conductivity (EC) map of Nirgudi-1 Microwatershed

6.4 Available Phosphorus

The soil fertility analysis revealed that available phosphorus is low (<23 kg/ha) in major area of about 754 ha (93%) area and is distributed in all parts of the microwatershed (Fig.6.4). There is an urgent need to increase the dose of phosphorous for all the crops by 25 per cent over the recommended dose to realize better crop performance. About 38 ha (5%) area in the microwatershed is medium (23-57 kg/ha) and is distributed in the southeastern and northwestern part of the microwatershed.

6.5 Available Potassium

Available potassium content is medium (145-337 kg/ha) in 150 ha (19%) area and is distributed in the central, eastern and western part of the microwatershed (Fig.6.5); high available potassium (>337 kg/ ha) content accounts for major area of 636 ha (79%) and is distributed in all parts of the microwatershed. Low available potassium (<145 kg/ha) content accounts for a very minor area of 6 ha and is distributed in the western part of the microwatershed.

6.6 Available Sulphur

Available sulphur content is low (<10 ppm) in 184 ha (23%) area and is distributed in the central, eastern and western part of the microwatershed. Maximum area of about 437 ha (54%) is medium (10-20 ppm) in available sulphur and is distributed in all parts of the microwatershed except southern parts (Fig.6.6). Available sulphur is high (>20 ppm) in an area of 171 ha (21%) and is distributed in the southern part of the microwatershed.

6.7 Available Boron

Available boron content is low (<0.5 ppm) in major area of about 679 ha (84%) and is distributed in all parts of the microwatershed. About 13 ha (14%) has soils that are medium (0.5-1.0 ppm) in available boron (Fig 6.7) and is distributed in the southern, eastern and western part of the microwatershed.




Fig.6.3 Soil Organic Carbon map of Nirgudi-1 Microwatershed




Fig.6.4 Soil available Phosphorus map of Nirgudi-1 Microwatershed




Fig.6.5 Soil available Potassium map of Nirgudi-1 Microwatershed




Fig.6.6 Soil available Sulphur map of Nirgudi-1 Microwatershed




Fig.6.7 Soil available Boron map of Nalgudi-1 Microwatershed

6.8 Available Iron

Available iron content is deficient (<4.5 ppm) in a small area of 38 ha (5%) and is distributed in the southern part of the microwatershed. It is sufficient in 754 ha (93%) area (Fig 6.8) and are distributed in all parts of the microwatershed.

6.9 Available Manganese

Available manganese content is sufficient (>1.0 ppm) in the entire microwatershed area (Fig 6.9).

6.10 Available Copper

Available copper content is sufficient (>0.2 ppm) in the entire microwatershed area (Fig 6.10).

6.11 Available Zinc

Available zinc content is deficient (<0.6 ppm) in major area of about 606 ha (75%) and is distributed in all parts of the microwatershed. It is sufficient (>0.6 ppm) in 186 ha (23%) area (Fig 6.11) and is distributed in the northwestern, southeastern and southwestern part of the microwatershed.




Fig.6.8 Soil available Iron map of Nirgudi-1 Microwatershed




Fig.6.9 Soil available Manganese map of Nirgudi-1 Microwatershed




Fig.6.10 Soil available Copper map of Nirgudi-1 Microwatershed




Fig.6.11 Soil available Zinc map of Nirgudi-1 Microwatershed

LAND SUITABILITY FOR MAJOR CROPS

The soil and land resource units (soil phases) of Nirgudi-1 microwatershed were assessed for their suitability for growing food, fibre, fodder and other horticulture crops by following the procedure as outlined in FAO, 1976 and 1983. Crop requirements were developed for each of the crop from the available research data, and also by referring to Naidu *et. al.* (2006) and Natarajan *et. al.* (2015). The crop requirements were matched with the soil and land characteristics (Table 7.1) to arrive at the crop suitability. In FAO land suitability classification, two orders are recognized. Order S-Suitable and Order N-Not suitable. The orders have classes, subclasses and units. Order-S has three classes, Class S1-Highly Suitable, Class S2-Moderately Suitable and Class S3-Marginally Suitable. Order N has two classes, N1-Currently not Suitable and N2-Permanently not Suitable. There are no subclasses within the class S1 as they will have very minor or no limitations for crop growth. Classes S2 and S3 are divided into subclasses based on the kinds of limitations encountered. The limitations that affect crop production are ‘c’ for erratic rainfall and its distribution and length of growing period (LGP), ‘e’ for erosion hazard, ‘r’ for rooting condition, ‘t’ for lighter or heavy texture, ‘g’ for gravelliness or stoniness, ‘n’ for nutrient availability, ‘l’ for topography, ‘m’ for moisture availability and ‘w’ for drainage. These limitations are indicated as lower case letters to the class symbol. For example, moderately suitable land with the limitations of soil depth and erosion is designated as S2re. For the microwatershed, the soil mapping units were evaluated and classified up to subclass level.

Using the above criteria, the soil map units of the microwatershed were evaluated and land suitability maps for 18 major annual and perennial crops were generated. The detailed information on the kind of suitability of each of the soil phase for the crops assessed are given village/ survey number wise for the microwatershed in Appendix-III.

7.1 Land Suitability for Sorghum (*Sorghum bicolor*)

Sorghum is one of the major crops grown in Karnataka in an area of 11.02 lakh ha in northern Karnataka in Bijapur, Kalaburgi, Raichur, Bidar, Belgaum, Dharwad and Bellary districts. The crop requirements for growing sorghum (Table 7.2) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and land suitability map for growing sorghum was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed is given in Figure. 7.1.

An area of about 160 ha (20%) in the microwatershed has soils that are highly suitable (class S1) for growing sorghum crop. They are distributed mainly in the central, northern and southern part of the microwatershed.

Table 7.1 Soil-Site Characteristics of Nirgudi-1 Microwatershed

Soil Map Units	Climate (P) (mm)	Growing period (Days)	Drainage class	Soil depth (cm)	Soil texture		Gravelliness		AWC (mm/m)	Slope (%)	Erosion	pH	EC	ESP	CEC [Cmol (p ⁺) kg ⁻¹]	BS (%)
					Surface	Sub-surface	Surface (%)	Subsurface (%)								
MGTmB2g1	786	150	WD	<25	c	c	15-35	15-35	<50	1-3	moderate	6.8	0.3	0.2	46	100
MGTmB2g2	786	150	WD	<25	c	c	35-60	15-35	<50	1-3	moderate	6.8	0.3	0.2	46	100
MGTmC2g1	786	150	WD	<25	c	c	15-35	15-35	<50	3-5	moderate	6.8	0.3	0.2	46	100
MGTmC2g2	786	150	WD	<25	c	c	35-60	15-35	<50	3-5	moderate	6.8	0.3	0.2	46	100
MGTmC3g2	786	150	WD	<25	c	c	35-60	15-35	<50	3-5	severe	6.8	0.3	0.2	46	100
MGTmD3g2	786	150	WD	<25	c	c	35-60	15-35	<50	5-10	severe	6.8	0.3	0.2	46	100
NHAmB1g1	786	150	WD	25-50	c	c	15-35	<15	<50	1-3	slight	7.2	0.1	0.3	40	100
NHAmB2	786	150	WD	25-50	c	c	-	<15	<50	1-3	moderate	7.2	0.1	0.3	40	100
NHAmB2g1	786	150	WD	25-50	c	c	15-35	<15	<50	1-3	moderate	7.2	0.1	0.3	40	100
NHAmB2g2	786	150	WD	25-50	c	c	35-60	<15	<50	1-3	moderate	7.2	0.1	0.3	40	100
NHAmC2g1	786	150	WD	25-50	c	c	15-35	<15	<50	3-5	moderate	7.2	0.1	0.3	40	100
NHAmC2g3	786	150	WD	25-50	c	c	60-80	<15	<50	3-5	moderate	7.2	0.1	0.3	40	100
BHImB2g1	786	150	WD	25-50	c	c	15-35	15-35	<50	1-3	moderate	7.0	0.1	0.2	28	100
DSImB2	786	150	MWD	50-75	c	c	-	<15	101-150	1-3	moderate	7.0	0.1	0.3	62	100
MANmA1	786	150	MWD	>150	c	c	-	<15	>200	0-1	slight	8.3	0.2	0.1	58	100

*Symbols and abbreviations are according to Field Guide for LRI under Sujala-III Project, Karnataka

A very small area of about 37 ha (4%) is moderately suitable (class S2) for growing sorghum and are distributed in the southwestern part of the microwatershed. They have moderate limitations of erosion and rooting depth. Marginally suitable lands (class S3) for growing sorghum occupy about 289 ha (36%) and occur in the eastern, northern, southeastern and southwestern part of the microwatershed. They have severe limitations of rooting depth and gravelliness. Major area of about 306 ha (38%) is not suitable for growing sorghum in the microwatershed and occur in the northwestern, central and northeastern part of the microwatershed.

Table 7.2 Crop suitability criteria for Sorghum

Crop requirement		Rating			
Soil –site characteristics	unit	Highly suitable (S1)	Moderately Suitable (S2)	Marginally suitable (S3)	Not suitable (N)
Slope	%	2-3	3-8	8-15	>15
LGP	Days	120-150	120-90	<90	
Soil drainage	class	Well to mod. drained	imperfect	Poorly/excessively	V. poorly
Soil reaction	pH	6.0-8.0	5.5-5.9 8.1-8.5	<5.5 8.6-9.0	>9.0
Surface soil texture	Class	C, cl, sicl, sc	l, sil, sic	Sl, ls	S, fragmental skeletal
Soil depth	Cm	100-75	50-75	30-50	<30
Gravel content	% vol.	5-15	15-30	30-60	>60
Salinity (EC)	dSm ⁻¹	2-4	4-8	8-10	>10
Sodicity (ESP)	%	5-8	8-10	10-15	>15




Fig. 7.1 Land Suitability map of Sorghum

7.2 Land Suitability for Maize (*Zea mays*)

Maize is the most important food crop grown in an area of 13.73 lakh ha in all the district of the state. The crop requirements for growing maize (Table 7.3) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing maize was generated. The area and geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.2.

In Nirgudi-1 microwatershed, there are no lands that are highly (Class S1) or moderately (Class S2) suitable lands for growing maize. The marginally suitable (class S3) lands cover a maximum area of about 486 ha (60%) and occur in all parts of the microwatershed. They have severe limitations of texture and rooting depth. About 306 ha (38%) area is not suitable for growing maize and occur in the all the parts of the microwatershed except eastern part.

Table 7.3 Crop suitability criteria for Maize

Crop requirement		Rating			
Soil –site characteristics	unit	Highly suitable (S1)	Moderately Suitable (S2)	Marginally Suitable (S3)	Not Suitable (N)
Slope	%	<3	3.5	5-8	
LGP	Days	>100	100-80	60-80	
Soil drainage	class	Well drained	Mod. to imperfectly	Poorly/excessively	V.poorly
Soil reaction	pH	5.5-7.5	7.6-8.5	8.6-9.0	
Surface soil texture	Class	l, cl, scl, sil	Sl, sicl, sic	C(s-s), ls	S, fragmental
Soil depth	Cm	>75	50-75	25-50	<25
Gravel content	% vol.	<15	15-35	35-50	>50
Salinity (EC)	dSm ⁻¹	<1.0	1.0-2.0	2.0-4.0	
Sodicity (ESP)	%	<10	10-15	>15	




Fig. 7.2 Land Suitability map of Maize

7.3 Land Suitability for Red gram (*Cajanus cajan*)

Red gram is one of the major pulse crop grown in an area of 8.23 lakh ha mainly in northern Karnataka in Bijapur, Kalaburgi, Raichur, Bidar, Belgaum, Dharwad and Bellary districts. The crop requirements for growing red gram (Table 7.4) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and land suitability map for growing red gram was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed is given in Figure 7.3.

An area of about 197 ha (24%) is moderately suitable (class S2) for red gram and is distributed dominantly in the northern, southern and central part of the microwatershed. They have moderate limitations of texture, rooting depth and erosion.

An area of about 289 ha (36%) is marginally suitable (class S3) for growing red gram and are distributed in all parts of the microwatershed. They have major limitations of rooting depth and erosion. A large area of about 306 ha (38%) is not suitable for growing red gram and occur in the northwestern, central and northeastern part of the microwatershed.

Table 7.4 Crop suitability criteria for Red gram

Crop requirement		Rating			
Soil characteristics	–site unit	Highly suitable (S1)	Moderately Suitable (S2)	Marginally suitable (S3)	Not suitable (N)
Slope	%	<3	3-5	5-10	>10
LGP	Days	>210	180-210	150-180	<150
Soil drainage	class	Well drained	Mod. to well drained	Imperfectly drained	Poorly drained
Soil reaction	pH	6.5-7.5	5.0-6.5 7.6-8.0	8.0-9.0	>9.0
Surface soil texture	Class	l, scl, sil, cl, sl	sicl, c(m)	sic, ls	S, fragmental
Soil depth	Cm	>100	85-100	40-85	<40
Gravel content	% vol.	<20	20-35	35-60	>60
Salinity (EC)	dSm ⁻¹	<1.0	1.0-2.0	>2.0	
Sodicity (ESP)	%	<10	10-15	>15	




Fig. 7.3 Land Suitability map of Red gram

7.4 Land Suitability for Sunflower (*Helianthus annuus*)

Sunflower is the most important oilseed crop grown in an area of 4.1 lakh ha in the State in all the districts. The crop requirements for growing sunflower (Table 7.5) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing sunflower was generated. The area extent and geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.4.

Highly suitable (class S1) lands are found to occur in an area of 160 ha (20%) and are distributed in the northern, southern and central part of the microwatershed. Marginally suitable (class S3) lands are found to occur in a small area of about 37 ha (5%). The soils have limitations of rooting depth. They are dominantly distributed in the southwestern part of the microwatershed. Major area of about 595 ha (74%) is not suitable for growing sunflower and occur in all parts of the microwatershed.

Table 7.5 Crop suitability criteria for Sunflower

Crop requirement		Rating			
Soil characteristics	–site unit	Highly suitable (S1)	Moderately Suitable (S2)	Marginally suitable (S3)	Not suitable (N)
Slope	%	<3	3-5	5-10	>10
LGP	Days	>90	80-90	70-80	<70
Soil drainage	class	Well drained	mod. drained	Well imperfectly drained	Poorly drained
Soil reaction	pH	6.5-8.0	8.1-8.5 5.5-6.4	8.6-9.0; 4.5-5.4	>9.0 <4.5
Surface soil texture	Class	l, cl, sil, sc	Scl, sic, c,	c (>60%), sl	ls, s
Soil depth	Cm	>100	75-100	50-75	<50
Gravel content	% vol.	<15	15-35	35-60	>60
Salinity (EC)	dSm ⁻¹	<1.0	1.0-2.0	>2.0	
Sodicity (ESP)	%	<10	10-15	>15	




Fig. 7.4 Land Suitability map of Sunflower

7.5 Land Suitability for Cotton (*Gossypium hirsutum*)

Cotton is the most important fibre crop grown in the State in about 6.6 lakh ha area in Raichur, Dharwad, Belgaum, Kalaburgi, Bijapur, Bidar, Bellary, Chitradurga and Chamarajnagar districts. The crop requirements for growing cotton (Table 7.6) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing cotton was generated. The area extent and geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.5.

Highly suitable (class S1) lands are found to occur in an area of 160 ha (20%) and are distributed in the northern, southern and central part of the microwatershed. Moderately suitable (class S2) lands are found to occur in a small area of about 37 ha (5%). The soils have moderate limitations of erosion and rooting depth. They are dominantly distributed in the southwestern part of the microwatershed. The marginally suitable (class S3) lands cover about 289 ha (36%) and mainly occur in the northeastern, northwestern and central part of the microwatershed. They have severe limitations of rooting depth and gravelliness. An area of about 306 ha (38%) is not suitable for growing cotton and are distributed in all parts of the microwatershed except eastern part.

Table 7.6 Crop suitability criteria for Cotton

Crop requirement		Rating			
Soil-site characteristics	unit	Highly suitable (S1)	Moderately Suitable (S2)	Marginally suitable (S3)	Not suitable (N)
Slope	%	1-2	2-3	3-5	>5
LGP	Days	180-240	120-180	<120	
Soil drainage	class	Well to moderately well	imperfectly drained	Poor somewhat excessive	Stagnant/excessive
Soil reaction	pH	6.5-7.5	7.6-8.0	8.1-9.0	>9.0 >6.5
Surface soil texture	Class	Sic, c	Sicl, cl	Si, sil, sc, scl, l	Sl, s,ls
Soil depth	Cm	100-150	60-100	30-60	<30
Gravel content	% vol.	<5	5-10	10-15	15-35
CaCO ₃ in root zone	%	<3	3-5	5-10	10-20
Salinity (EC)	dSm ⁻¹	2-4	4.0-8.0	8.0-12	>12
Sodicity (ESP)	%	5-10	10-20	20-30	>30




Fig. 7.5 Land Suitability map of Cotton

7.6 Land Suitability for Sugarcane (*Saccharum officinarum*)

Sugarcane is the most important commercial crop grown in 6.7 lakh ha area in Kalaburgi, Bijapur, Bagalkot, Bidar, Mysore, Chamarajanagar and Mandya districts under irrigated conditions. The crop requirements for growing sugarcane (Table 7.7) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing sugarcane was generated. The area extent and geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.6.

The marginally suitable (class S3) lands cover about 197 ha (25%) area and mainly occur in the northern, central and southwestern part of the microwatershed. They have severe limitations of texture. Major area of about 595 ha (74%) is not suitable for growing sugarcane and occur in all parts of the microwatershed.

Table 7.7 Crop suitability criteria for Sugarcane

Crop requirement		Rating			
Soil-site characteristics	unit	Highly suitable (S1)	Moderately Suitable (S2)	Marginally suitable (S3)	Not suitable (N)
Slope	%	<3	3-5	5-8	>8
Soil drainage	class	Well drained	Mod./imperfectly drained	Poorly drained	V.poor/ excessively drained
Soil reaction	pH	7.0-8.0	6.0-6.9 8.1-9.0	4.0-5.9 9.1-9.5	<4.0/ >9.5
Surface soil texture	Class	l, cl, sil, sicl	C(m/k), sl	C+(ss)	
Soil depth	cm	>100	100-75	75-50	<50
stoniness	%	<15	15-35	35-50	>50
Salinity (EC)	dSm ⁻¹	<2.0	2.0-4.0	4.0-9.0	>9
Sodicity (ESP)	%	<10	10-15	15-25	>25




Fig. 7.6 Land Suitability map of Sugarcane

7.7 Land Suitability for Soybean (*Glycine max*)

Soybean is the most important pulse and oil seed crop grown in about 1.68 lakh ha area in Bijapur, Raichur, Kalaburgi, Dharwad, Belgaum and Bellary districts. The crop

requirements for growing soybean were matched with the soil-site characteristics and a land suitability map for growing soybean was generated. The area extent and geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.7.

Highly suitable (class S1) lands are found to occur in an area of 160 ha (20%) and are distributed in the northern, southern and central part of the microwatershed. Moderately suitable (class S2) lands are found to occur in a small area of about 37 ha (5%). The soils have moderate limitations of erosion and rooting depth. They are dominantly distributed in the western part of the microwatershed. The marginally suitable (class S3) lands cover about 289 ha (36%) area and mainly occur in the northern, northwestern and eastern part of the microwatershed. They have severe limitations of rooting depth and gravelliness. Major area of about 306 ha (38%) is not suitable for growing soybean and occur in all parts of the microwatershed.




Fig. 7.7 Land Suitability map of Soybean

7.8 Land Suitability for Guava (*Psidium guajava*)

Guava is the most important fruit crop grown in the State in Raichur, Dharwad, Belgaum, Kalaburgi, Bijapur, Bidar, Bellary, Chitradurga, Bangalore and Chamarajnarag districts. The crop requirements for growing guava (Table 7.8) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing guava was generated. The area extent and geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.8.

In Nirgudi-1 microwatershed, there are no highly (Class S1) or moderately (Class S2) suitable lands available for growing guava. The marginally suitable (class S3) lands cover an area of about 197 ha (24%) in the microwatershed and mainly occur in the northern, southern, central and southwestern part of the microwatershed. They have severe limitations of texture and rooting depth. Major area of about 595 ha (74%) is not suitable for growing guava and occur in all parts of the microwatershed.

Table 7.8 Crop suitability criteria for Guava

Crop requirement			Rating			
Soil –site characteristics		unit	Highly suitable (S1)	Moderately Suitable (S2)	Marginally suitable (S3)	Not suitable (N)
climate	Temperature in growing season	⁰ C	28-32	33-36 24-27	37-42 20-23	
Soil moisture	Growing period	Days	>150	120-150	90-120	<90
Soil aeration	Soil drainage	class	Well drained	Mod. to imperfectly	poor	Very poor
Nutrient availability	Texture	Class	Scl, l, cl, sil	Sl,sicl,sic.,sc,c	C (<60%)	C (>60%)
	pH	1:2.5	6.0-7.5	7.6-8.0:5.0-5.9	8.1-8.5:4.5-4.9	>8.5:<4.5
	CaCO ₃ in root zone	%	Non calcareous	<10	10-15	>15
Rooting conditions	Soil depth	cm	>100	75-100	50-75	<50
	Gravel content	% vol.	<15	15-35	>35	
Soil toxicity	Salinity	dS/m	<2.0	2.0-4.0	4.0-6.0	
	Sodicity	%	Non sodic	10-15	15-25	>25
Erosion	Slope	%	<3	3-5	5-10	>10




Fig 7.8 Land Suitability map of Guava

7.9 Land Suitability for Mango (*Mangifera indica*)

Mango is the most important fruit crop grown in all the districts of the State. The crop requirements for growing mango (Table 7.9) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing mango was generated. The area and geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.9.

No highly (Class S1) and moderately (Class S2) suitable lands are available for growing mango in the microwatershed. The marginally suitable (class S3) lands cover an area of about 160 ha (20%) and mainly occur in the northern, southern and central part of the microwatershed. They have severe limitations of texture. Major area of about 632 ha (78%) is not suitable for growing mango and occur in all parts of the microwatershed.

Table 7.9 Crop suitability criteria for Mango

Crop requirement			Rating			
soil-site characteristics		unit	Highly suitable (S1)	Moderately Suitable (S2)	Marginally suitable (S3)	Not suitable (N)
climate	Temp in growing season	°C	28-32	24-27 33-35	36-40	20-24
	Min. temp. before flowering	°C	10-15	15-22	>22	
Soil moisture	Growing period	Days	>180	150-180	120-150	<120
Soil aeration	Soil drainage	class	Well drained	Mod. To imperfectly drained	Poor drained	Very poorly drained
	Water table	M	>3	2.50-3.0	2.5-1.5	<1.5
Nutrient availability	Texture	Class	Sc, l, sil, cl	Sl, sc, sic, l, c	C (<60%)	C (>60%),
	pH	1:2.5	5.5-7.5	7.6-8.55.0-5.4	8.6-9.0 4.0-4.9	>9.0 <4.0
	OC	%	High	medium	low	
	CaCO ₃ in root zone	%	Non calcareous	<5	5-10	>10
Rooting conditions	Soil depth	cm	>200	125-200	75-125	<75
	Gravel content	% vol.	Non gravelly	<15	15-35	>35
Soil toxicity	Salinity	dS/m	Non saline	<2.0	2.0-3.0	>3.0
	Sodicity	%	Non sodic	<10	10-15	>15
Erosion	Slope	%	<3	3-5	5-10	




Fig. 7.9 Land Suitability map of Mango

7.10 Land Suitability for Sapota (*Manilkara zapota*)

Sapota is the most important fruit crop grown in almost all the districts of the state. The crop requirements for growing sapota (Table 7.10) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing sapota was generated. The area extent and geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.10.

In Nirgudi-1 microwatershed, there are no lands that are highly (Class S1) or moderately (Class S2) suitable lands for growing sapota. The marginally suitable (class S3) lands cover an area of about 197 ha (24%) and mainly occur in the northern, southern, central and southwestern part of the microwatershed. They have severe limitations of rooting depth and texture. Major area of about 595 ha (74%) is not suitable for growing sapota and occur in all parts of the microwatershed.

Table 7.10 Crop suitability criteria for Sapota

Crop requirement			Rating			
Soil –site characteristics		unit	Highly suitable (S1)	Moderately Suitable (S2)	Marginally suitable (S3)	Not suitable (N)
climate	Temperature in growing season	⁰ C	28-32	33-36 24-27	37-42 20-23	>42 <18
Soil moisture	Growing period	Days	>150	120-150	90-120	<120
Soil aeration	Soil drainage	class	Well drained	Moderately well drained	Imperfectly drained	Poorly drained
Nutrient availability	Texture	Class	Scl, l, cl, sil	Sl, si, cl, sc	C (<60%)	ls, s, C (>60%)
	pH	1:2.5	6.0-7.5	7.6-8.0 5.0-5.9	8.1-9.0 4.5-4.9	>9.0 <4.5
	CaCO ₃ in root zone	%	Non calcareous	<10	10-15	>15
Rooting conditions	Soil depth	cm	>150	75-150	50-75	<50
	Gravel content	% vol.	Non gravelly	<15	15-35	<35
Soil toxicity	Salinity	dS/m	Non saline	Up to 1.0	1.0-2.0	2.0-4.0
	Sodicity	%	Non sodic	10-15	15-25	>25
Erosion	Slope	%	<3	3-5	5-10	>10




Fig. 7.10 Land Suitability map of Sapota

7.11 Land Suitability for Jackfruit (*Artocarpus heterophyllus*)

Jackfruit is the most important fruit crop grown in almost all the districts of the state. The crop requirements for growing jackfruit were matched with the soil-site characteristics and a land suitability map for growing jackfruit was generated. The area extent and geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.11.

No highly (Class S1) and moderately (Class S2) suitable lands are available for growing jackfruit in the microwatershed. The marginally suitable (class S3) lands cover an area of about 197 ha (24%) and mainly occur in the northern, southern, central and southwestern part of the microwatershed. They have severe limitations of texture and rooting depth. Major area of about 595 ha (74%) is not suitable for growing jackfruit and occur in all parts of the microwatershed.




Fig 7.11 Land Suitability map of Jackfruit

7.12 Land Suitability for Jamun (*Syzygium cumini*)

Jamun is the most important fruit crop grown in almost all the districts of the state. The crop requirements for growing jamun were matched with the soil-site characteristics and a land suitability map for growing jamun was generated. The area extent and geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.12.

The moderately suitable (class S2) lands are found to occur in an area of about 160 ha (20%). The soils have moderate limitations of texture. They are dominantly distributed in the northern, southern and central part of the microwatershed.

The marginally suitable (class S3) lands cover about a minor area of 37 ha (5%) and mainly occur in the southwestern part of the microwatershed. They have severe limitations of rooting depth. Major area of about 595 ha (74%) is not suitable for growing jamun and occur in all parts of the microwatershed.




Fig 7.12 Land Suitability map of Jamun

7.13 Land Suitability for Musambi (*Citrus limetta*)

Musambi is the most important fruit crop grown in almost all the districts of the state. The crop requirements for growing musambi were matched with the soil-site characteristics and a land suitability map for growing musambi was generated. The area extent and geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.13.

Highly suitable (class S1) lands are found to occur in an area of about 160 ha (20%) and are distributed in the northern, southern and central part of the microwatershed. The marginally suitable (class S3) lands cover a small area of about 37 ha (5%) in the microwatershed and mainly occur in the southwestern part of the microwatershed. They have

severe limitation of rooting depth. Major area of about 595 ha (74%) is not suitable for growing musambi and occur in all parts of the microwatershed.




Fig 7.13 Land Suitability map of Musambi

7.14 Land Suitability for Lime (*Citrus sp*)

Lime is the most important fruit crop grown in almost all the districts of the state. The crop requirements for growing lime (Table 7.11) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing lime was generated. The area extent and geographic distribution of different suitability subclasses in the microwatershed is given in Figure 7.14.

Highly suitable (class S1) lands are found to occur in an area of about 160 ha (20%) and are distributed in the northern, southern and central part of the microwatershed. The marginally suitable (class S3) lands cover a small area of about 37 ha (5%) in the microwatershed and occur in the southwestern part of the microwatershed. They have severe limitations of rooting depth. Major area of about 595 ha (74%) is not suitable for growing lime and occur in all parts of the microwatershed.

Table 7.11 Crop suitability criteria for Lime

Crop requirement			Rating			
Soil –site characteristics		unit	Highly suitable (S1)	Moderately Suitable (S2)	Marginally suitable (S3)	Not suitable (N)
Climate	Temp in growing season	⁰ C	28-30	31-35 24-27	36-40 20-23	>40 <20
Soil moisture	Growing period	Days	240-265	180-240	150-180	<150
Soil aeration	Soil drainage	class	Well drained	Mod. to imperfectly drained	poorly	Very poorly
Nutrient availability	Texture	Class	Scl, l, sicl, cl, s	Sc, sc, c	C (>70%)	S, ls
	pH	1:2.5	6.0-7.5	5.5-6.4/ 7.6-8.0	4.0-5.4 8.1-8.5	<4.0 >8.5
	CaCO ₃ in root zone	%	Non calcareous	Upto 5	5-10	>10
Rooting condition	Soil depth	cm	>150	100-150	50-100	<50
	Gravel content	% vol.	Non gravelly	15-35	35-55	>55
Soil toxicity	Salinity	dS/m	Non saline	Upto 1.0	1.0-2.5	>2.5
	Sodicity	%	Non sodic	5-10	10-15	>15
Erosion	Slope	%	<3	3-5	5-10	




Fig 7.14 Land Suitability map of Lime

7.15 Land Suitability for Cashew (*Anacardium occidentale*)

Cashew is the most important plantation crop grown in almost all the districts. The crop requirements for growing Cashew were matched with the soil-site characteristics and a land suitability map for growing Cashew was generated. The area and geographic distribution of different suitability subclasses in the microwatershed is given in Figure 7.15. The entire area is not suitable for growing cashew in the microwatershed.




Fig 7.15 Land Suitability map of Cashew

7.16 Land Suitability for Custard Apple (*Annona reticulata*)

Custard apple is the most important fruit crop grown in almost all the districts of the state. The crop requirements for growing custard apple were matched with the soil-site characteristics and a land suitability map for growing custard apple was generated. The area extent and geographic distribution of different suitability subclasses in the microwatershed is given in Figure 7.16.

Highly suitable (class S1) lands are found to occur in an area of 160 ha (20%) and are distributed in the northern, southern and central part of the microwatershed.

Moderately suitable (class S2) lands are found to occur in minor area of about 37 ha (5%). The soils have moderate limitations of erosion and rooting depth. They are dominantly distributed in the southwestern part of the microwatershed. The marginally suitable (class S3) lands cover about 291 ha (36%) area and mainly occur in the northeastern, northwestern and western part of the microwatershed. They have severe limitations of rooting depth and gravelliness. Major area of about 304 ha (38%) is not suitable for growing custard apple and occur in major part of the microwatershed.




Fig 7.16 Land Suitability map of Custard Apple

7.17 Land Suitability for Amla (*Phyllanthus emblica*)

Amla is the most important fruit crop grown in almost all the districts of the state. The crop requirements for growing amla were matched with the soil-site characteristics and a land suitability map for growing amla was generated. The area extent and geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.17.

Highly suitable (class S1) lands are found to occur in an area of 160 ha (20%) and are distributed in the northern and southern part of the microwatershed.

Moderately suitable (class S2) lands are found to occur in a small area of about 37 ha (5%). The soils have moderate limitations of erosion and rooting depth. They are dominantly distributed in the southwestern part of the microwatershed.

The marginally suitable (class S3) lands cover about 291 ha (36%) area and occur in the northeastern, northwestern and western part of the microwatershed. They have severe limitations of rooting depth and gravelliness. Major area of about 304 ha (38%) is not suitable for growing amla and occur in all parts of the microwatershed.




Fig 7.17 Land Suitability map of Amla

7.18 Land Suitability for Tamarind (*Tamarindus indica*)

Tamarind is the most important spice crop raised in all the districts of the state. The crop requirements for growing tamarind were matched with the soil-site characteristics and a land suitability map for growing tamarind was generated. The area extent and geographic distribution of different suitability subclasses in the microwatershed is given in Figure 7.18.

Moderately suitable (class S2) lands are found to occur in an area of about 160 ha (20%). The soils have moderate limitation of rooting depth. They are dominantly distributed in the northern, southern and central part of the microwatershed. Major area of about 632 ha (78%) is not suitable for growing tamarind and occur in all parts of the microwatershed.




Fig 7.18 Land Suitability map of Tamarind

7.19 Land Management Units (LMUs)

The 15 soil map units identified in Nirgudi-1 microwatershed have been regrouped into 5 Land Management Units (LMU's) for the purpose of preparing Proposed Crop Plan. Land Management Units are grouped based on the similarities in respect of the type of soil, the depth of the soil, the surface soil texture, gravel content, AWC, slope, erosion etc. and a Land Management Units map (Fig.7.19) has been prepared. These Land Management Units are expected to behave similarly for a given level of management.

The map units that have been grouped into 5 land management units along with brief description of soil and site characteristics are given below.

LMUs	Soil map units	Soil and site characteristics
1	MGTmC3g2 MGTmD3g2	Very shallow, black soils with slopes of 3-10%, very gravelly (35-60%) and severe erosion
2	MGTmB2g1, MGTmB2g2, MGTmC2g1, MGTmC2g2, NHAmC2g3	Very shallow, black soils with slopes of 1-5%, gravelly to very gravelly and strongly gravelly (15- >60%) and moderate erosion
3	NHAmB1g1, NHAmB2, NHAmB2g1, NHAmB2g2, NHAmC2g1, BHImB2g1	Shallow, black soils with slopes of 1-5 %, gravelly to very gravelly (15-35%) and slight to moderate erosion
4	DSImB2	Moderately shallow, black soils with slopes of 1-3 % and moderate erosion
5	MANmA1	Very deep, black soils with slopes of <1%, gravelly (15-35%) and slight erosion




Fig. 7.19 Land Management Units map of Nirgudi-1 Microwatershed

7.20 Proposed Crop Plan for Nirgudi-1 Microwatershed

After assessing the land suitability for the 18 crops, a proposed crop plan has been prepared for the 5 identified LMUs by considering only the highly (class S1) and moderately (class S2) suitable lands for each of the eighteen crops. The resultant proposed crop plan is presented below in Table 7.12

Table 7.12 Proposed Crop Plan for Nirgudi - 1 Microwatershed

LMUs	Mapping Unit	Survey No	Crops proposed				Suitable Intervention
			Field Crops	Forestry /Grasses	Horticulture Crops (Rainfed Condition)	Horticulture Crops with suitable intervention	
LMU-1	MGTmC3g2 MGTmD3g2	Nirgudi: 150,157,250,260 Padsavli: 23,253,256,257,260	-	Neem, Glyricydia , Silviculture, Agave, Simaroba	-	-	Crescent bunds
LMU-2	MGTmB2g1 MGTmB2g2 MGTmC2g1 MGTmC2g2 NHAmC2g3	Nirgudi: 127,128,129,130, 131,136,151,241,248,249, 251,252, 253,254,257,258, 259,261 Padsavli: 13,251,252,254, 255	Horse gram, Green gram, chick pea	Neem, Glyricydia , Silviculture, Agave, Simaroba	-	-	Crescent bunds
LMU-3	NHAmB1g1 NHAmB2 NHAmB2g1 NHAmB2g2 NHAmC2g1 BHImB2g1	Nirgudi: 132,134,135,137, 138,139,140,141,142,143, 144,145,146,147,148,149, 154,155,156,186,187,188, 189,193,232,255,262,281 Padsavli: 12,14,15,18,24, 25,26,27,28,29,57,264,265	Bajra, Linseed, Green gram, Black gram, Chick pea, Coriander	Subabhul, Neem, Teak	Custard apple, Charoli, Ber, Amla Vegetables: Ladies finger, Brinjal, Cowpea, Flowers: Marigold, Chrysanthemum	Custard apple, Charoli, Ber, Amla Vegetables: Onion, Tomato, Brinjal, Chillies, Bhendi Flowers: Marigold, Chrysanthemum	Drip irrigation, suitable soil and water conservations like cultivation on raised beds with mulches and drip

LMU	Mapping Unit	Survey No	Crops proposed				Recommended Intervention
			Field Crops	Forestry Crop/ Grasses	Horticulture Crops (Rainfed Condition)	Horticulture Crops With suitable intervention	
LMU-4	DSImB2	Padsavli: 258,259,262,263	Sorghum, Cotton, Red Gram, Black gram, Green gram, Soybean, Sesame, Sunflower, Safflower Rabi: Sorghum, Chickpea	Subabul, Neem, Teak	Custard apple, Charoli, Ber, Amla Vegetables: Ladies finger, Brinjal, Cowpea, Flowers: Marigold, Chrysanthemum	Custard apple, Charoli, Ber, Amla, Papaya, Banana, Lime, Citrus Vegetables: Onion, Tomato, Brinjal, Chillies, Bhendi Flowers: Marigold, Chrysanthemum	-do- Graded bunds, Strengthening of field bunds
LMU-5	MANmA1	Nirgudi: 158,159,160,161,162,163,164,165,166,167,168,169,170,171,172,173,174,175,176,177,178,179,180,181,182,183,184,185,233,234,235,236,237,238,239,240,242,243,244,245,246,247 Padsavli: 6,7/1,7/2,7/3,8,9,11,16,17,19,20,21,22,261,266,267,273,274	Sorghum, Cotton, Red Gram Black gram, Green gram, Soybean, Sesame, Sunflower, Safflower, Rabi: Sorghum, Chickpea, Coriander, linseed	-	Vegetables: Ladies finger, Brinjal, Cowpea, Coriander Field crops: Sorghum, Cotton, Red Gram, Sunflower, Safflower, Perennial component: Guava, Tamarind, Sapota, Lime, Mosambi Flowers: Marigold, Chrysanthemum	Banana, Papaya, Lime, Musambi, Guava, Tamrind Vegetables: Onion, Tomato, Brinjal, Chillies, Bhendi Flowers: Marigold, Chrysanthemum	-do- Graded bunds, Strengthening of field bunds

SOIL HEALTH MANAGEMENT

8.1 Soil Health

Soil is fundamental to crop production. Without soil, no food could be produced nor would livestock be fed on a large scale. Because it is finite and fragile, soil is a precious resource that requires special care from its users.

Soil health or the capacity of the soil to function is critical to human survival. Soil health has been defined as: “the capacity of the soil to function as a living system without adverse effect on the ecosystem”. Healthy soils maintain a diverse community of soil organisms that help to form beneficial symbiotic associations with plant roots, recycle essential plant nutrients, improve soil structure with positive repercussions for soil, water and nutrient holding capacity and ultimately improve crop production and also contribute to mitigating climate change by maintaining or increasing its carbon content.

Functional interactions of soil biota with organic and inorganic components, air and water determine a soil’s potential to store and release nutrients and water to plants and to promote and sustain plant growth. Thus, maintaining soil health is vital to crop production and conserve soil resource base for sustaining agriculture.

The most important characteristics of a healthy soil are

- Good soil tilth
- Sufficient soil depth
- Good water storage and good drainage
- Adequate supply, but not excess of nutrients
- Large population of beneficial organisms
- Small proportion of plant pathogens and insect pests
- Low weed pressure
- Free of chemicals and toxins that may harm the crop
- Resistance to degradation
- Resilience when unfavourable conditions occur

Characteristics of Nirgudi-1 Microwatershed

- The soil phases with sizeable area identified in the microwatershed belonged to the soil series of MGT (304 ha), NHA (276 ha), MAN (160 ha), DSI (37 ha) and BHI (14 ha).
- As per land capability classification, nearly 94 per cent area comes under arable land category (Class II, III and IV) and 4 per cent area belongs to nonarable land (class VI) category. The major limitations identified in the arable lands were soil and erosion.
- On the basis of soil reaction, about 432 ha (53%) area is moderately alkaline (pH 7.8-8.4) followed by slightly alkaline (pH 7.3-7.8) soils in 76 ha (9%) and strongly alkaline (pH 8.4-9.0) 12 ha (1.5%). Thus, about 89 per cent of the soils in the microwatershed are alkaline in reaction.

Soil Health Management

The following actions are required to improve the current land husbandry practices that provide a sound basis for the successful adoption of sustainable crop production system.

Alkaline soils

(Slightly alkaline to moderately alkaline soils)

1. Regular addition of organic manure, green manuring, green leaf manuring, crop residue incorporation and mulching needs to be taken up to improve the soil organic matter status.
2. Application of biofertilizers (Azospirillum, Azotobacter, Rhizobium).
3. Application of 25% extra N and P (125 % RDN&P).
4. Application of $ZnSO_4$ – 12.5 kg/ha (once in three years).
5. Application of Boron – 5kg/ha (once in three years).

Neutral soils

1. Regular addition of organic manure, green manuring, green leaf manuring, crop residue incorporation and mulching needs to be taken up to improve the soil organic matter status.
2. Application of biofertilizers, (Azospirillum, Azotobacter, Rhizobium).
3. Application of 100 per cent RDF.
4. Need based micronutrient applications.

Besides the above recommendations, the best transfer of technology options are also to be adopted.

Soil Degradation

Soil erosion is one of the major factor affecting the soil health in the microwatershed. Out of total area of 808 ha in the microwatershed, major area of 627 ha is suffering from either moderate or severe erosion. These areas need immediate soil and water conservation and other land husbandry practices for restoring soil health.

Disseminate information and communicate benefits

Any large scale implementation of soil health management requires that supporting information is made available widely, particularly through channels familiar to farmers and extension workers. Given the very high priority attached to soil health especially by the Central Government on issuing Soil-Health Cards to all the farmers, media outlets like regional, state and national newspapers, Radio and Dooradarshan programs in local languages but also modern information and communication technologies such as cellular phones and the Internet, which can be much more effective in reaching younger farmers.

Inputs for Net Planning and Interventions needed

Net planning in IWMP is focusing on preparation of

1. Soil and Water Conservation Plan for each plot or farm.
2. Productivity enhancement measures/ interventions for existing crops/livestock/other farm enterprises.
3. Diversification of farming mainly with perennial horticultural crops and livestock.
4. Improving livelihood opportunities and income generating activities.

In this connection, how various outputs of Sujala-III are of use in addressing these objectives of Net Planning are briefly presented below.

- ❖ **Soil Depth:** The depth of a soil decides the amount of moisture and nutrients it can hold, what crops can be taken up or not, depending on the rooting depth and the length of growing period available for raising any crop. Deeper the soil, better for a wide variety of crops. If sufficient depth is not available for growing deep rooted crops either choose medium or short duration crops or deeper planting pits need to be opened and additional good quality soil brought from outside has to be filled into the planting pits.
- ❖ **Surface soil texture:** Lighter soil texture in the top soil means, better rain water infiltration, less run-off and soil moisture conservation, less capillary rise and less evaporation losses. Lighter surface textured soils are amenable to good soil tilth and are highly suitable for crops like groundnut, root vegetables (carrot, raddish, potato etc) but not ideal for crops that need stagnant water like lowland paddy. Heavy textured soils are poor in water infiltration and percolation. They are prone for sheet erosion; such soils can be improved by sand mulching. The technology that is developed by the AICRP-Dryland Agriculture, Vijayapura, Karnataka may be adopted.
- ❖ **Gravelliness:** More gravel content is favorable for run-off harvesting but poor in soil moisture storage and nutrient availability. It is a significant parameter that decides the kind of crop to be raised.
- ❖ **Land Capability Classification:** The land capability map shows the areas suitable and not suitable for agriculture and the major constraints in each of the plot/survey number. Hence, one can decide what kind of enterprise is possible in each of these units. In general, erosion and soil are the major constraints in Nirgudi-1 microwatershed.
- ❖ **Organic Carbon:** In about 49 ha (6%) area, the OC content is low (<0.5%), in about 591 ha (73%) area, the OC content is medium (0.5-0.75%) and in about 152 ha (19%) area it is high (>0.75%). The areas that are low and medium in OC needs to be further improved by applying farmyard manure and rotating crops with cereals and legumes or mixed cropping.
- ❖ **Promoting green manuring:** Growing of green manuring crops cost Rs. 1250/ha (green manuring seeds) and about Rs. 2000/ha towards cultivation that totals to Rs. 3250/- per ha. On the other hand, application of organic manure @ 10 tons/ha costs Rs. 5000/ha. The practice needs to be continued for 2-3 years or more. Nitrogen fertilizer needs to be supplemented by 25% in addition to the recommended level in 49 ha area where OC is less

than 0.5-0.75%. For example, for rainfed maize, recommended level is 50 kg N per ha and an additional 12 kg /ha needs to be applied for all the crops grown in these plots.

- ❖ **Available Phosphorus:** In 754 ha (93%) area, the available phosphorus is low and about 38 ha (5%) area it is medium in available phosphorus in the microwatershed. Hence for all the crops, 25% additional P-needs to be applied.
- ❖ **Available Potassium:** Available potassium is medium in 150 ha (19%) area of the microwatershed. Hence, in all these plots, for all crops, an additional 25 % potassium may be applied. It is high in 636 ha (79%) area of the microwatershed.
- ❖ **Available Sulphur:** Available sulphur is a very critical nutrient for oilseed crops. It is low in 184 ha (23%) area of the microwatershed and medium in 437 ha (54%). These areas need to be applied with magnesium sulphate or gypsum or Factamphos (p) fertilizer (13% sulphur) for 2-3 years for the deficiency to be corrected. About 171 ha (21%) area has soils that are high in available sulphur.
- ❖ **Available iron:** It is deficient in a small area of 38 ha (5%) the microwatershed. To manage iron deficiency, iron sulphate @ 25kg /ha needs to be applied for 2-3 years. It is sufficient in the rest of 754 ha (93 %) area in the microwatershed.
- ❖ **Available Zinc:** It is deficient in 606 ha (75%) area of the microwatershed. Application of zinc sulphate @25kg/ha is to be applied. It is sufficient in 186 ha (23%) area in the microwatershed.


Soil alkalinity: The microwatershed has 444 ha area with soils that are moderately to strongly alkaline. These areas need application of gypsum and wherever calcium is in excess, iron pyrites and element sulphur can be recommended. Management practices like treating repeatedly with good quality water to drain out the excess salts and, provision of subsurface drainage and growing of salt tolerant crops like Casuarina, Acasia, Neem, Ber etc., are recommended.

Land Suitability for various crops: Areas that are highly, moderately and marginally suitable for growing various crops are indicated. Along with the suitability, various constraints that are limiting the productivity are also indicated. For example, in case of cotton, gravel content, rooting depth and salinity/alkalinity are the major constraints in various plots. With suitable management interventions, the productivity can be enhanced. In order to increase water holding capacity of light textured soils, growing of green manure crops and application of organic manure is recommended.

SOIL AND WATER CONSERVATION TREATMENT PLAN

For preparing soil and water conservation treatment plan for Nirgudi-1 microwatershed, the land resource inventory database generated under Sujala-III project has been transformed as information through series of interpretative (thematic) maps using soil phase map as a base. The various thematic maps (1:7920 scale) generated were

- Soil depth
- Surface soil texture
- Available water capacity
- Soil slope
- Soil gravelliness
- Land capability
- Present land use and land cover
- Crop suitability maps
- Rainfall map
- Hydrology
- Water Resources
- Socio-economic data
- Contour plan with existing features- Network of waterways, ponthissa boundaries, cut up/ minor terraces etc.
- Cadastral map (1:7920 scale)
- Satellite imagery (1:7920 scale)



Apart from these, Hand Level/ Hydro Marker/ Dumpy Level/ Total Station and Kathedars' List has to be collected.

Steps for Survey and Preparation of Treatment Plan

The boundaries of Land user Groups' and Survey No. boundaries are traced in the field.

- Naming of user groups and farmers
- Identification of arable and non arable lands
- Identification of drainage lines and gullies
- Identification of non treatable areas
- Identification of priority areas in the arable lands
- Treatment plan for arable lands
- Location of water harvesting and recharge structures

9.1 Treatment Plan

The treatment plan recommended for arable lands is briefly described below


9.1.1 Arable Land Treatment

A. BUNDING

Steps for Survey and Preparation of Treatment Plan		USER GROUP-1
Cadastral map (1:7920 scale) is enlarged to a scale of 1:2500 scale		
Existing network of waterways, pothissa boundaries, grass belts, natural drainage lines/ watercourse, cut ups/ terraces are marked on the cadastral map to the scale		
Drainage lines are demarcated into		
Small gullies	(up to 5 ha catchment)	
Medium gullies	(5-15 ha catchment)	
Ravines	(15-25 ha catchment) and	
Halla/Nala	(more than 25ha catchment)	

Measurement of Land Slope

Land slope is estimated or determined by the study and interpretation of contours or by measurement in the field using simple instruments like Hand level or Hydromarker.



Vertical and Horizontal intervals between bunds as recommended by the Watershed Development Department.

Slope percentage	Vertical interval (m)	Corresponding Horizontal Distance (m)
2 - 3%	0.6	24
3 - 4%	0.9	21
4 - 5%	0.9	21
5 - 6%	1.2	21
6 - 7%	1.2	21

Note: (i) The above intervals are maximum.

(ii) Considering the slope class and erosion status (A1....) the intervals have to be decided.

Bund length recording: Considering the contour plan and the existing grass belts/partitions, the bunds are aligned and lengths are measured.

Section of the Bund

Bund section is decided considering the soil texture class and gravelliness class (bgo-loamy sand, <15% gravel). The recommended Sections for different soils are given below.

Recommended Bund Section

Top width (m)	Base width (m)	Height (m)	Side slope (Z:1;H:V)	Cross section (sq m)	Soil Texture	Remarks
0.3	0.9	0.3	01:01	0.18	Sandy loam	Vegetative bund
0.3	1.2	0.3	1.5:1	0.225	Sandy clay	
0.3	1.2	0.5	0.9:1	0.375	Red gravelly soil	
0.3	1.2	0.6	0.75:1	0.45		
0.3	1.5	0.6	01:01	0.54	Red sandy loam	
0.3	2.1	0.6	1.5:1	0.72	Very shallow black soils	
0.45	2	0.75	01:01	0.92		
0.45	2.4	0.75	1.3:1	1.07	Shallow black soils	
0.6	3.1	0.7	1.78:1	1.29	Medium black soils	
0.5	3	0.85	1.47:1	1.49		

Formation of Trench cum Bund

Dimensions of the Borrow Pits/Trenches to be excavated (machinery are decided considering the Bund Section).

Details of Borrow Pit dimensions are given below:

TRENCH CUM BUND

WATER STORAGE AREA

0.45 Sq.m section

IDEAL FOR HORTICULTURE CROPS

'A' FRAME FOR INTERBUND MANAGEMENT

1. ಸಮಪಾತಳಿ ಉಳಿಸುವುದು

2. ಸಮಪಾತಳಿ ಬಿತ್ತನೆ/ನಾಟಿ

Size of Borrow Pits/ Trench recommended for Trench cum Bund (by machinery)

Bund section	Bund length	Earth quantity	Pit				Berm (pit to pit)	Soil depth class
			L(m)	W(m)	D(m)	QUANTITY (m ³)		
m ²	m	m ³	L(m)	W(m)	D(m)	QUANTITY (m ³)	m	
0.375	6	2.25	5.85	0.85	0.45	2.24	0.15	Shallow
0.45	6	2.7	5.4	1.2	0.43	2.79	0.6	Shallow
0.45	6	2.7	5	0.85	0.65	2.76	1	Moderately Shallow
0.54	5.6	3.02	5.5	0.85	0.7	3.27	0.1	Moderately shallow
0.54	5.5	2.97	5	1.2	0.5	3	0.5	Shallow
0.72	6.2	4.46	6	1.2	0.7	5.04	0.2	Moderately shallow
0.72	5.2	3.74	5.1	0.85	0.9	3.9	0.1	Moderately deep

B. Water Ways

- Existing waterways are marked on the cadastral map (1:7920 scale) and their dimensions are recorded.
- Considering the contour plan of the MWS, additional waterways/ modernization of the existing ones can be thought of.
- The design details are given in the Manual.

C. Farm Ponds

Waterways and the catchment area will give an indication on the size of the Farm Pond. Location of the pond can be decided based on the contour plan/ field condition and farmers' need/desire.

D. Diversion Channel

Existing EPT/ CPT are marked on the cadastral map. Looking to the need, these can be modernized or fresh diversion channel can be proposed and runoff from this can be stored in Gokatte/ Recharge ponds.

9.1.2 Non-Arable Land Treatment

Depending on the gravelliness and crops preferred by the farmers, the concerned authorities can decide appropriate treatment plan. The recommended treatments may be Contour Trench, Staggered Trench, Crescent Bund, Boulder Bund or Pebble Bunds are formed in the field.

9.1.3 Treatment of Natural Water Course/ Drainage Lines

- a) The cadastral map has to be updated as regards the network of drainage lines (gullies/ nallas/ hallas) and existing structures are marked to the scale and storage capacity of the existing water bodies are documented.
- b) The drainage line will be demarcated into Upper Reach, Middle Reach and Lower Reach.
- c) Considering the Catchment, Nala bed and bank conditions, suitable structures are decided.
- d) Number of storage structures (Check dam/ Nala bund/ Percolation tank) will be decided considering the commitments and available runoff in water budgeting and quality of water in the wells and site suitability.
- e) Detailed Levelling Survey using Dumpy Level / Total Station has to be carried out to arrive at the site-specific designs as shown in the Manual.
- f) The location of ground water recharge structures are decided by examining the lineaments and fracture zones from geological maps.
- g) Rainfall intensity data of the nearest Rain gauge station is considered for Hydrologic Designs.
- h) Silt load to the Storage/Recharge structures is reduced by providing vegetative, boulder and earthen checks in the natural water course. Location and design details are given in the Manual.

9.2 Recommended Soil and Water Conservation Measures

The appropriate conservation structures best suited for each of the land parcel/ survey number (Appendix-I) are selected based on the slope per cent, severity of erosion, amount of rainfall, land use and soil type. The different kinds of conservation structures recommended are:

1. Graded / Strengthening of bunds
2. Trench cum Bunds (TCB)
3. Trench cum Bunds / Strengthening
4. Crescent Bunds

A map (Fig. 9.1) showing soil and water conservation plan with different kinds of structures recommended has been prepared which shows the spatial distribution and extent of area. An area of about 595 ha (74%) requires crescent bunds or trench cum bunding, 160 ha (20%) area needs graded bunds and about 37 ha (5%) area needs strengthening of bunds.

The conservation plan prepared may be presented to all the stakeholders including farmers and after including their suggestions, the conservation plan for the microwatershed may be finalised in a participatory approach.




Fig. 9.1 Soil and Water Conservation Plan map of Nirgudi-1 Microwatershed

9.3 Greening of Microwatershed

As part of the greening programme in the watersheds, it is envisaged to plant a variety of horticultural and other tree plants that are edible, economical and produce lot of biomass which helps to restore the ecological balance in the watersheds. The lands that are suitable for greening programme are non-arable lands (land capability classes V, VI and VII) and also the lands that are not suitable or marginally suitable and field bunds for growing annual and perennial crops. The method of planting these trees is given below.

It is recommended to open pits during the 1st week of March along the contour and heap the dug out soil on the lower side of the slope in order to harness the flowing water and facilitate weathering of soil in the pit. Exposure of soil in the pit also prevents spread of pests and diseases due to scorching sun rays. The pits should be filled with mixture of soil and organic manure during the second week of April and keep ready with sufficiently tall seedlings produced either in poly bags or in root trainer nurseries so that planting can be done during the 2nd or 3rd week of April depending on the rainfall.

The tree species suitable for the area considering rainfall, temperature and adaptability is listed below; waterlogged areas are recommended to be planted with species like Neral (*Syzgium cumini*) and Bamboo. Dry areas are to be planted with species like Honge, Bevu, Seetaphal etc.

Dry Deciduous Species			Temp (°C)	Rainfall (mm)
1.	Bevu	<i>Azadiracta indica</i>	21-32	400 -1,200
2.	Tapasi	<i>Holoptelia integrifolia</i>	20-30	500 - 1000
3.	Seetaphal	<i>Anona Squamosa</i>	20-40	400 - 1000
4.	Honge	<i>Pongamia pinnata</i>	20 -50	500- 2,500
5.	Kamara	<i>Hardwickia binata</i>	25 -35	400 - 1000
6.	Bage	<i>Albezzia lebbek</i>	20 - 45	500 - 1000
7.	Ficus	<i>Ficus bengalensis</i>	20 - 50	500-2,500
8.	Sisso	<i>Dalbargia Sissoo</i>	20 - 50	500 -2000
9.	Ailanthus	<i>Ailanthus excelsa</i>	20 - 50	500 - 1000
10.	Hale	<i>Wrightia tinctoria</i>	25 - 45	500 - 1000
11.	Uded	<i>Steriospermum chelanoides</i>	25 - 45	500 -2000
12.	Dhupa	<i>Boswellia Serrata</i>	20 - 40	500 - 2000
13.	Nelli	<i>Emblica Officinalis</i>	20 - 50	500 -1500
14.	Honne	<i>Pterocarpus marsupium</i>	20 - 40	500 - 2000
Moist Deciduous Species			Temp (°C)	Rainfall (mm)
15.	Teak	<i>Tectona grandis</i>	20 - 50	500-5000
16.	Nandi	<i>Legarstroemia lanceolata</i>	20 - 40	500 - 4000
17.	Honne	<i>Pterocarpus marsupium</i>	20 - 40	500 - 3000
18.	Mathi	<i>Terminalia alata</i>	20 -50	500 - 2000
19.	Shivane	<i>Gmelina arboria</i>	20 -50	500 -2000
20.	Kindal	<i>T.Paniculata</i>	20 - 40	500 - 1500
21.	Beete	<i>Dalbargia latifolia</i>	20 - 40	500 - 1500
22.	Tare	<i>T. belerica</i>	20 - 40	500 - 2000
23.	Bamboo	<i>Bambusa arundinasia</i>	20 - 40	500 - 2500
24.	Bamboo	<i>Dendrocalamus strictus</i>	20 - 40	500 - 2500
25.	Muthuga	<i>Butea monosperma</i>	20 - 40	400 - 1500
26.	Hippe	<i>Madhuca latifolia</i>	20 - 40	500 - 2000
27.	Sandal	<i>Santalum album</i>	20 - 50	400 - 1000
28.	Nelli	<i>Emblica officinalis</i>	20 - 40	500 - 2000
29.	Nerale	<i>Sizygium cumini</i>	20 - 40	500 - 2000
30.	Dhaman	<i>Grevia tilifolia</i>	20 - 40	500 - 2000
31.	Kaval	<i>Careya arborea</i>	20 - 40	500 - 2000
32.	Harada	<i>Terminalia chebula</i>	20 - 40	500 - 2000

References

1. FAO (1976) Framework for Land Evaluation, Food and Agriculture Organization, Rome.72 pp.
2. FAO (1983) Guidelines for Land Evaluation for Rainfed Agriculture, FAO, Rome, 237 pp.
3. IARI (1971) Soil Survey Manual, All India Soil and Land Use Survey Organization, IARI, New Delhi, 121 pp.
4. Katyal, J.C. and Rattan, R.K. (2003) Secondary and Micronutrients; Research Gap and Future Needs. Fert. News 48 (4); 9-20.
5. Naidu, L.G.K., Ramamurthy, V., Challa, O., Hegde, R. and Krishnan, P. (2006) Manual Soil Site Suitability Criteria for Major Crops, NBSS Publ. No. 129, NBSS & LUP, Nagpur, 118 pp.
6. Natarajan, A. and Dipak Sarkar (2010) Field Guide for Soil Survey, National Bureau of Soil Survey and Land Use Planning (ICAR), Nagpur, India.
7. Natarajan, A., Rajendra Hegde, Raj, J.N. and Shivananda Murthy, H.G. (2015) Implementation Manual for Sujala-III Project, Watershed Development Department, Bengaluru, Karnataka.
8. Sarma, V.A.K., Krishnan, P. and Budihal, S.L. (1987) Laboratory Manual, Tech. Bull. 23, NBSS &LUP, Nagpur.
9. Sehgal, J.L. (1990) Soil Resource Mapping of Different States of India; Why and How?, National Bureau of Soil Survey and Land Use Planning, Nagpur, 49 pp.
10. Shivaprasad, C.R., R.S. Reddy, J. Sehgal and M. Velayuthum (1998) Soils of Karnataka for Optimising Land Use, NBSS Publ. No. 47b, NBSS & LUP, Nagpur, India.
11. Soil Survey Staff (2006) Keys to Soil Taxonomy, Tenth edition, U.S. Department of Agriculture/ NRCS, Washington DC, U.S.A.
12. Soil Survey Staff (2012) Soil Survey Manual, Handbook No. 18, USDA, Washington DC, USA.

Appendix - I

Nirguidi-I Microwatershed

Soil Site and Thematic Information

Village	Sur-vey No.	Total Area (ha)	Soil phase	Land Management Unit	Soil Depth	Surface Soil Texture	Soil Gravel-iness	AWC	Slope	Soil Erosion	CLU Code	WELLS	Land Capability	Conservation Plan
Niragudi	127	13.58	MGTmB2g 2	LMU-2	Very shallow (<25 cm)	Clay	Very gravelly (35-60%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Cultivated Fallow+Redgram (CF+Rg)	Not Available	IIIse	Crescent bund/TCB
Niragudi	128	13.57	MGTmB2g 2	LMU-2	Very shallow (<25 cm)	Clay	Very gravelly (35-60%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Cultivated Fallow+Greengram+Redgram (CF+Gg+Rg)	Not Available	IIIse	Crescent bund/TCB
Niragudi	129	11.57	MGTmB2g 2	LMU-2	Very shallow (<25 cm)	Clay	Very gravelly (35-60%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Cultivated Fallow (CF)	Not Available	IIIse	Crescent bund/TCB
Niragudi	130	9.98	MGTmB2 g1	LMU-2	Very shallow (<25 cm)	Clay	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Cultivated Fallow+Redgram (CF+Rg)	Not Available	IIIse	Crescent bund/TCB
Niragudi	131	7.73	MGTmB2 g1	LMU-2	Very shallow (<25 cm)	Clay	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Grass land+Redgram (G1+Rg)	Not Available	IIIse	Crescent bund/TCB
Niragudi	132	13.49	NHAmB2 g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cultivated Fallow+Scrub land (CF+Sl)	Not Available	IIIse	Crescent bund/TCB
Niragudi	134	11.6	NHAmB2 g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cultivated Fallow+Jowar (CF+Jw)	Not Available	IIIse	Crescent bund/TCB
Niragudi	135	12.09	NHAmB2	LMU-3	Shallow (25-50 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram+Jowar+Redgram (Gg+Jw+Rg)	Not Available	IIIse	Crescent bund/TCB
Niragudi	136	16.14	MGTmB2 g2	LMU-2	Very shallow (<25 cm)	Clay	Very gravelly (35-60%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Cultivated Fallow+Grass land+Redgram+Scrub land (CF+G1+Rg+Sl)	Not Available	IIIse	Crescent bund/TCB
Niragudi	137	3.51	NHAmB2	LMU-3	Shallow (25-50 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cultivated Fallow+Redgram (CF+Rg)	Not Available	IIIse	Crescent bund/TCB
Niragudi	138	6.29	NHAmB2	LMU-3	Shallow (25-50 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cultivated Fallow+Greengram+Redgram (CF+Gg+Rg)	Not Available	IIIse	Crescent bund/TCB
Niragudi	139	6.4	NHAmB2	LMU-3	Shallow (25-50 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Chilli+Greengram+Sugarcane (Ch+Gg+Sc)	Borewell,Openwell	IIIse	Crescent bund/TCB
Niragudi	140	9.51	NHAmB2	LMU-3	Shallow (25-50 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cultivated Fallow+Greengram+Sugarcane (CF+Gg+Sc)	Borewell,Openwell	IIIse	Crescent bund/TCB
Niragudi	141	13.22	NHAmB2 g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cultivated Fallow+Greengram+Redgram (CF+Gg+Rg)	Openwell	IIIse	Crescent bund/TCB

Village	Sur-vey No.	Total Area (ha)	Soil phase	Land Management Unit	Soil Depth	Surface Soil Texture	Soil Gravel-ness	AWC	Slope	Soil Erosion	CLU Code	WELLS	Land Capability	Conservation Plan
Niragudi	142	11.75	NHAmB2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cultivated Fallow+Greengram (CF+Gg)	Openwell	IIIse	Crescent bund/TCB
Niragudi	143	9.36	NHAmB2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram+Redgram (Gg+Rg)	Not Available	IIIse	Crescent bund/TCB
Niragudi	144	10.52	NHAmB2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cultivated Fallow+Jowar+Redgram (CF+Jw+Rg)	Not Available	IIIse	Crescent bund/TCB
Niragudi	145	10.26	NHAmC2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Gently sloping (3-5%)	Moderate	Cultivated Fallow+Redgram+Scrub land (CF+Rg+Sl)	Not Available	IIIse	Crescent bund/TCB
Niragudi	146	2.89	NHAmB2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cultivated Fallow+Sugarcane (CF+Sc)	Borewell	IIIse	Crescent bund/TCB
Niragudi	147	6.96	NHAmB2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cultivated Fallow+Greengram+Redgram (CF+Gg+Rg)	Not Available	IIIse	Crescent bund/TCB
Niragudi	148	11.27	NHAmB2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cultivated Fallow+Greengram+Redgram (CF+Gg+Rg)	Openwell,Openwell	IIIse	Crescent bund/TCB
Niragudi	149	8.79	NHAmB2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cultivated Fallow+Redgram (CF+Rg)	Not Available	IIIse	Crescent bund/TCB
Niragudi	150	13.62	MGTmC3g2	LMU-1	Very shallow (<25 cm)	Clay	Very gravelly (35-60%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Cultivated Fallow+Grass land (CF+Gl)	Not Available	IVse	Crescent bund/TCB
Niragudi	151	1.09	NHAmC2g3	LMU-2	Shallow (25-50 cm)	Clay	Extremely gravelly (60-80%)	Low (51-100 mm/m)	Gently sloping (3-5%)	Moderate	Redgram (Rg)	Not Available	IVse	Crescent bund/TCB
Niragudi	154	4.61	NHAmB2	LMU-3	Shallow (25-50 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Grass land+Redgram (Gl+Rg)	Not Available	IIIse	Crescent bund/TCB
Niragudi	155	3.39	NHAmB2	LMU-3	Shallow (25-50 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cultivated Fallow+Redgram (CF+Rg)	Checkdam	IIIse	Crescent bund/TCB
Niragudi	156	1.81	NHAmB2	LMU-3	Shallow (25-50 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cultivated Fallow (CF)	Not Available	IIIse	Crescent bund/TCB
Niragudi	157	12.72	MGTmC3g2	LMU-1	Very shallow (<25 cm)	Clay	Very gravelly (35-60%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Grass land (Gl)	Farmpond,Checkdam	IVse	Crescent bund/TCB
Niragudi	158	0.68	MANmA1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Cultivated Fallow (CF)	Not Available	IIs	GB/strengthening of Bunds

Village	Sur-vey No.	Total Area (ha)	Soil phase	Land Management Unit	Soil Depth	Surface Soil Texture	Soil Graveliness	AWC	Slope	Soil Erosion	CLU Code	WELLS	Land Capability	Conservation Plan
Niragudi	159	0.84	MANmA1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Cultivated Fallow (CF)	Not Available	IIs	GB/ strengthenin g of Bunds
Niragudi	160	1.13	MANmA1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Redgram+Sugarcane (Rg+Sc)	Not Available	IIs	GB/ strengthenin g of Bunds
Niragudi	161	1.37	MANmA1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Cultivated Fallow (CF)	Openwell,Bor ewell	IIs	GB/ strengthenin g of Bunds
Niragudi	162	0.35	MANmA1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Cultivated Fallow (CF)	Not Available	IIs	GB/ strengthenin g of Bunds
Niragudi	163	0.7	MANmA1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Cultivated Fallow (CF)	Not Available	IIs	GB/ strengthenin g of Bunds
Niragudi	164	0.65	MANmA1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Sugarcane (Sc)	Not Available	IIs	GB/ strengthenin g of Bunds
Niragudi	165	1.23	MANmA1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Cultivated Fallow (CF)	Not Available	IIs	GB/ strengthenin g of Bunds
Niragudi	166	1.28	MANmA1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Cultivated Fallow (CF)	Openwell	IIs	GB/ strengthenin g of Bunds
Niragudi	167	0.75	MANmA1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Cultivated Fallow (CF)	Borewell	IIs	GB/ strengthenin g of Bunds
Niragudi	168	0.74	MANmA1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Cultivated Fallow (CF)	Openwell	IIs	GB/ strengthenin g of Bunds
Niragudi	169	0.59	MANmA1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Cultivated Fallow (CF)	Openwell	IIs	GB/ strengthenin g of Bunds
Niragudi	170	0.93	MANmA1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Cultivated Fallow (CF)	Not Available	IIs	GB/ strengthenin g of Bunds
Niragudi	171	0.4	MANmA1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Cultivated Fallow (CF)	Not Available	IIs	GB/ strengthenin g of Bunds
Niragudi	172	0.47	MANmA1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Cultivated Fallow (CF)	Not Available	IIs	GB/ strengthenin g of Bunds

Village	Sur-vey No.	Total Area (ha)	Soil phase	Land Management Unit	Soil Depth	Surface Soil Texture	Soil Gravel-ness	AWC	Slope	Soil Erosion	CLU Code	WELLS	Land Capability	Conservation Plan
Niragudi	173	0.4	MANmA1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Cultivated Fallow (CF)	Not Available	IIs	GB/ strengthening of Bunds
Niragudi	174	0.29	MANmA1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Cultivated Fallow (CF)	Not Available	IIs	GB/ strengthening of Bunds
Niragudi	175	0.45	MANmA1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Cultivated Fallow (CF)	Not Available	IIs	GB/ strengthening of Bunds
Niragudi	176	0.22	MANmA1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Cultivated Fallow (CF)	Not Available	IIs	GB/ strengthening of Bunds
Niragudi	177	0.55	MANmA1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Cultivated Fallow (CF)	Not Available	IIs	GB/ strengthening of Bunds
Niragudi	178	0.87	MANmA1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Redgram (Rg)	Not Available	IIs	GB/ strengthening of Bunds
Niragudi	179	1.31	MANmA1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Cultivated Fallow (CF)	Not Available	IIs	GB/ strengthening of Bunds
Niragudi	180	0.79	MANmA1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Sugarcane (Sc)	Not Available	IIs	GB/ strengthening of Bunds
Niragudi	181	0.76	MANmA1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Cultivated Fallow (CF)	Not Available	IIs	GB/ strengthening of Bunds
Niragudi	182	0.93	MANmA1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Cultivated Fallow (CF)	Not Available	IIs	GB/ strengthening of Bunds
Niragudi	183	1.23	MANmA1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Cultivated Fallow (CF)	Not Available	IIs	GB/ strengthening of Bunds
Niragudi	184	1.23	MANmA1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Cultivated Fallow (CF)	Not Available	IIs	GB/ strengthening of Bunds
Niragudi	185	1.34	MANmA1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Greengram+Sugarcane (Gg+Sc)	Openwell, Borewell	IIs	GB/ strengthening of Bunds
Niragudi	186	0.52	NHAmB1g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Cultivated Fallow (CF)	Borewell	IIIs	Crescent bund/TCB

Village	Sur-vey No.	Total Area (ha)	Soil phase	Land Management Unit	Soil Depth	Surface Soil Texture	Soil Gravel-ness	AWC	Slope	Soil Erosion	CLU Code	WELLS	Land Capability	Conservation Plan
Niragudi	187	0.63	NHAmB1g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Cultivated Fallow (CF)	Not Available	IIIs	Crescent bund/TCB
Niragudi	188	0.67	NHAmB1g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Cultivated Fallow (CF)	Not Available	IIIs	Crescent bund/TCB
Niragudi	189	0.9	NHAmB1g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Cultivated Fallow (CF)	Not Available	IIIs	Crescent bund/TCB
Niragudi	193	3.65	NHAmB1g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Greengram+Redgram (Gg+Rg)	Not Available	IIIs	Crescent bund/TCB
Niragudi	232	6.12	BHImB2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Cultivated Fallow+Greengram+Redgram (CF+Gg+Rg)	Borewell	IIIse	Crescent bund/TCB
Niragudi	233	2.92	MANmA1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Cultivated Fallow (CF)	Not Available	IIs	GB/ strengthening of Bunds
Niragudi	234	1.16	MANmA1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Cultivated Fallow (CF)	Openwell	IIs	GB/ strengthening of Bunds
Niragudi	235	0.34	MANmA1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Cultivated Fallow (CF)	Not Available	IIs	GB/ strengthening of Bunds
Niragudi	236	1.16	MANmA1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Greengram+Redgram (Gg+Rg)	Not Available	IIs	GB/ strengthening of Bunds
Niragudi	237	0.89	MANmA1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Cultivated Fallow (CF)	Not Available	IIs	GB/ strengthening of Bunds
Niragudi	238	0.74	MANmA1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Cultivated Fallow (CF)	Not Available	IIs	GB/ strengthening of Bunds
Niragudi	239	0.55	MANmA1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Cultivated Fallow+Sugarcane (CF+Sc)	Openwell	IIs	GB/ strengthening of Bunds
Niragudi	240	3.37	MANmA1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Sugarcane (Sc)	Not Available	IIs	GB/ strengthening of Bunds
Niragudi	241	3.74	MGTmC2g2	LMU-2	Very shallow (<25 cm)	Clay	Very gravelly (35-60%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Moderate	Redgram (Rg)	Openwell	IIIse	Crescent bund/TCB
Niragudi	242	1.15	MANmA1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Cultivated Fallow (CF)	Not Available	IIs	GB/ strengthening of Bunds

Village	Sur-vey No.	Total Area (ha)	Soil phase	Land Management Unit	Soil Depth	Surface Soil Texture	Soil Gravel-ness	AWC	Slope	Soil Erosion	CLU Code	WELLS	Land Capability	Conservation Plan
Niragudi	243	0.75	MANmA1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Cultivated Fallow (CF)	Not Available	IIs	GB/strengthening of Bunds
Niragudi	244	0.73	MANmA1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Cultivated Fallow (CF)	Not Available	IIs	GB/strengthening of Bunds
Niragudi	245	1.25	MANmA1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Cultivated Fallow (CF)	Not Available	IIs	GB/strengthening of Bunds
Niragudi	246	10.77	MANmA1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Cultivated Fallow+Redgram (CF+Rg)	Openwell	IIs	GB/strengthening of Bunds
Niragudi	247	4.86	MANmA1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Redgram (Rg)	Borewell,Openwell	IIs	GB/strengthening of Bunds
Niragudi	248	7.87	MGTmC2g1	LMU-2	Very shallow (<25 cm)	Clay	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Moderate	Cultivated Fallow+Redgram (CF+Rg)	Openwell,Openwell	IIIse	Crescent bund/TCB
Niragudi	249	13.82	MGTmC2g1	LMU-2	Very shallow (<25 cm)	Clay	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Moderate	Cultivated Fallow (CF)	Borewell,Borewell	IIIse	Crescent bund/TCB
Niragudi	250	7.29	MGTmD3g2	LMU-1	Very shallow (<25 cm)	Clay	Very gravelly (35-60%)	Very low (<50 mm/m)	Moderately sloping (5-10%)	Severe	Cultivated Fallow+Scrub land (CF+SI)	Not Available	VIse	Crescent bund/TCB
Niragudi	251	6.09	MGTmC2g1	LMU-2	Very shallow (<25 cm)	Clay	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Moderate	Cultivated Fallow+Scrub land (CF+SI)	Not Available	IIIse	Crescent bund/TCB
Niragudi	252	9.36	MGTmC2g1	LMU-2	Very shallow (<25 cm)	Clay	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Moderate	Cultivated Fallow+Scrub land (CF+SI)	Not Available	IIIse	Crescent bund/TCB
Niragudi	253	8.31	MGTmB2g1	LMU-2	Very shallow (<25 cm)	Clay	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Cultivated Fallow (CF)	Not Available	IIIse	Crescent bund/TCB
Niragudi	254	11.66	MGTmC2g2	LMU-2	Very shallow (<25 cm)	Clay	Very gravelly (35-60%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Moderate	Cultivated Fallow+Greengram (CF+Gg)	Not Available	IIIse	Crescent bund/TCB
Niragudi	255	7.54	BHImB2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Cultivated Fallow+Sugarcane+Tomato (CF+Sc+Tm)	Borewell,Borewell,Openwell,Borewell	IIIse	Crescent bund/TCB
Niragudi	257	1.45	MGTmC2g2	LMU-2	Very shallow (<25 cm)	Clay	Very gravelly (35-60%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Moderate	Cultivated Fallow (CF)	Openwell	IIIse	Crescent bund/TCB
Niragudi	258	1.7	MGTmC2g2	LMU-2	Very shallow (<25 cm)	Clay	Very gravelly (35-60%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Moderate	Cultivated Fallow (CF)	Not Available	IIIse	Crescent bund/TCB

Village	Sur-vey No.	Total Area (ha)	Soil phase	Land Management Unit	Soil Depth	Surface Soil Texture	Soil Graveliness	AWC	Slope	Soil Erosion	CLU Code	WELLS	Land Capability	Conservation Plan
Niragudi	259	10.98	MGTmC2g2	LMU-2	Very shallow (<25 cm)	Clay	Very gravelly (35-60%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Moderate	Cultivated Fallow (CF)	Openwell,Farmpound,Farmpound	IIIse	Crescent bund/TCB
Niragudi	260	13.14	MGTmC3g2	LMU-1	Very shallow (<25 cm)	Clay	Very gravelly (35-60%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Cultivated Fallow+Redgram+Sugarcane+Scrub land (CF+Rg+Sc+Sl)	Not Available	IVse	Crescent bund/TCB
Niragudi	261	4.04	MGTmC2g1	LMU-2	Very shallow (<25 cm)	Clay	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Moderate	Cultivated Fallow (CF)	Not Available	IIIse	Crescent bund/TCB
Niragudi	262	8.17	NHAmB2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cultivated Fallow (CF)	Not Available	IIIse	Crescent bund/TCB
Niragudi	281	10.42	NHAmB2g2	LMU-3	Shallow (25-50 cm)	Clay	Very gravelly (35-60%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cultivated Fallow (CF)	Not Available	IIIse	Crescent bund/TCB
Niragudi	STREAM	5.75	Waterbody	Others	Others	Others	Others	Others	Others	Others	Others	Not Available	Others	Others
Padasavali	6	1.24	MANmA1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Onion (On)	Not Available	IIs	GB/strengthening of Bunds
Padasavali	7/1	0.59	MANmA1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Cultivated Fallow (CF)	Not Available	IIs	GB/strengthening of Bunds
Padasavali	7/2	0.86	MANmA1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Cultivated Fallow (CF)	Not Available	IIs	GB/strengthening of Bunds
Padasavali	7/3	0.26	MANmA1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Cultivated Fallow (CF)	Not Available	IIs	GB/strengthening of Bunds
Padasavali	8	4.93	MANmA1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Cultivated Fallow+Greengram+Sugarcane (CF+Gg+Sc)	Openwell,Borewell	IIs	GB/strengthening of Bunds
Padasavali	9	11.21	MANmA1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Greengram+Jowar+Sugarcane (Gg+Jw+Sc)	Borewell,Openwell,Borewell	IIs	GB/strengthening of Bunds
Padasavali	11	7.51	MANmA1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Greengram+Sugarcane (Gg+Sc)	Borewell,Openwell	IIs	GB/strengthening of Bunds
Padasavali	12	8.15	NHAmB2	LMU-3	Shallow (25-50 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram+Sugarcane (Gg+Sc)	Not Available	IIIse	Crescent bund/TCB
Padasavali	13	10.29	MGTmC2g1	LMU-2	Very shallow (<25 cm)	Clay	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Moderate	Greengram+Maize+Redgram+Soyabean (Gg+Mz+Rg+Sb)	Borewell,Openwell	IIIse	Crescent bund/TCB

Village	Sur-vey No.	Total Area (ha)	Soil phase	Land Management Unit	Soil Depth	Surface Soil Texture	Soil Graveliness	AWC	Slope	Soil Erosion	CLU Code	WELLS	Land Capability	Conservation Plan
Padasavali	14	12.34	NHAmB2	LMU-3	Shallow (25-50 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Bajra+Groundnut+Redgram (Bj+Gn+Rg)	Openwell	IIIse	Crescent bund/TCB
Padasavali	15	8.84	NHAmB2	LMU-3	Shallow (25-50 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram+Redgram+Soyabean+Sugarcane (Gg+Rg+Sb+Sc)	Checkdam	IIIse	Crescent bund/TCB
Padasavali	16	7.54	MANmA1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Cultivated Fallow+Sugarcane (CF+Sc)	Not Available	IIs	GB/strengthening of Bunds
Padasavali	17	8.8	MANmA1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Redgram+Soyabean+Sugarcane (Rg+Sb+Sc)	2Borewells, Openwell	IIs	GB/strengthening of Bunds
Padasavali	18	8.88	NHAmB2	LMU-3	Shallow (25-50 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Sunflower (Rg+Sf)	Openwell, Checkdam	IIIse	Crescent bund/TCB
Padasavali	19	11.34	MANmA1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Greengram+Redgram+Sugarcane (Gg+Rg+Sc)	Openwell, Checkdam, Farmpond	IIs	GB/strengthening of Bunds
Padasavali	20	12.62	MANmA1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Redgram+Soyabean+Sugarcane (Rg+Sb+Sc)	Not Available	IIs	GB/strengthening of Bunds
Padasavali	21	7.27	MANmA1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Cultivated Fallow+Redgram (CF+Rg)	Not Available	IIs	GB/strengthening of Bunds
Padasavali	22	8.01	MANmA1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Greengram+Redgram+Sugarcane (Gg+Rg+Sc)	Not Available	IIs	GB/strengthening of Bunds
Padasavali	23	12.05	MGTmC3g2	LMU-1	Very shallow (<25 cm)	Clay	Very gravelly (35-60%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Jawar+Sugarcane (Jw+Sc)	Checkdam, 2Borewells, Openwell	IVse	Crescent bund/TCB
Padasavali	24	11.86	NHAmB2	LMU-3	Shallow (25-50 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cultivated Fallow+Redgram+Scrubland (CF+Rg+Sl)	Not Available	IIIse	Crescent bund/TCB
Padasavali	25	14.4	NHAmB2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut+Jowar+Redgram+Sunflower (Gn+Jw+Rg+Sf)	Not Available	IIIse	Crescent bund/TCB
Padasavali	26	13.25	NHAmB2	LMU-3	Shallow (25-50 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut+Jowar+Redgram+Sunflower (Gn+Jw+Rg+Sf)	Not Available	IIIse	Crescent bund/TCB
Padasavali	27	7.3	NHAmB2	LMU-3	Shallow (25-50 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Bajra+Redgram (Bj+Rg)	Not Available	IIIse	Crescent bund/TCB

Village	Sur-vey No.	Total Area (ha)	Soil phase	Land Management Unit	Soil Depth	Surface Soil Texture	Soil Gravel- liness	AWC	Slope	Soil Erosion	CLU Code	WELLS	Land Capability	Conservation Plan
Padasavali	28	1.58	NHAmB2	LMU-3	Shallow (25-50 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Sugarcane (Sc)	Borewell,Bor ewell	IIIse	Crescent bund/TCB
Padasavali	29	6.74	NHAmB2	LMU-3	Shallow (25-50 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut+Redgram (Gn+Rg)	Not Available	IIIse	Crescent bund/TCB
Padasavali	57	6.69	NHAmB2	LMU-3	Shallow (25-50 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cultivated Fallow+Redgram (CF+Rg)	Not Available	IIIse	Crescent bund/TCB
Padasavali	251	4.92	MGTmB2 g1	LMU-2	Very shallow (<25 cm)	Clay	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Cultivated Fallow+Redgram (CF+Rg)	Borewell,Ope nwell	IIIse	Crescent bund/TCB
Padasavali	252	9.69	MGTmB2 g1	LMU-2	Very shallow (<25 cm)	Clay	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Cultivated Fallow+Redgram (CF+Rg)	Openwell	IIIse	Crescent bund/TCB
Padasavali	253	6.03	MGTmD3 g2	LMU-1	Very shallow (<25 cm)	Clay	Very gravelly (35-60%)	Very low (<50 mm/m)	Moderately sloping (5-10%)	Severe	Cultivated Fallow+Redgram+Scrub land (CF+Rg+Sl)	Not Available	VIse	Crescent bund/TCB
Padasavali	254	10.05	MGTmB2 g1	LMU-2	Very shallow (<25 cm)	Clay	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Cultivated Fallow+Greengram (CF+Gg)	Checkdam,C heckdam	IIIse	Crescent bund/TCB
Padasavali	255	11.04	MGTmB2 g1	LMU-2	Very shallow (<25 cm)	Clay	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Cultivated Fallow+Sugarcane+Tomato (CF+Sc+Tm)	Borewell,Bor ewell	IIIse	Crescent bund/TCB
Padasavali	256	9.51	MGTmD3 g2	LMU-1	Very shallow (<25 cm)	Clay	Very gravelly (35-60%)	Very low (<50 mm/m)	Moderately sloping (5-10%)	Severe	Blackgram+Cultivated Fallow+Jowar+Redgram (Bm+CF+Jw+Rg)	Borewell	VIse	Crescent bund/TCB
Padasavali	257	5.66	MGTmD3 g2	LMU-1	Very shallow (<25 cm)	Clay	Very gravelly (35-60%)	Very low (<50 mm/m)	Moderately sloping (5-10%)	Severe	Cultivated Fallow (CF)	Openwell	VIse	Crescent bund/TCB
Padasavali	258	10.48	DSImB2	LMU-4	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Cultivated Fallow (CF)	Not Available	IIs	TCB/GB/ strengthening of Bunds
Padasavali	259	8.78	DSImB2	LMU-4	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Cultivated Fallow (CF)	Not Available	IIs	TCB/GB/ strengthening of Bunds
Padasavali	260	5.4	MGTmD3 g2	LMU-1	Very shallow (<25 cm)	Clay	Very gravelly (35-60%)	Very low (<50 mm/m)	Moderately sloping (5-10%)	Severe	Cultivated Fallow+Redgram+Sugarcane +Scrub land (CF+Rg+Sc+Sl)	Borewell,Ope nwell	VIse	Crescent bund/TCB
Padasavali	261	13.28	MANmA1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Greengram+Redgram+Scrub land (Gg+Rg+Sl)	Not Available	IIs	GB/ strengthening of Bunds
Padasavali	262	8.17	DSImB2	LMU-4	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Cultivated Fallow (CF)	Not Available	IIs	TCB/GB/ strengthening of Bunds

Village	Sur-vey No.	Total Area (ha)	Soil phase	Land Management Unit	Soil Depth	Surface Soil Texture	Soil Graveliness	AWC	Slope	Soil Erosion	CLU Code	WELLS	Land Capability	Conservation Plan
Padasavali	263	6.6	DSImB2	LMU-4	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram+Soyabean+Sugar cane (Gg+Sb+Sc)	Not Available	IIs	TCB/GB/strengthening of Bunds
Padasavali	264	8.87	NHAmB2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram+Redgram+No Crop (Gg+Rg+NC)	Not Available	IIIse	Crescent bund/TCB
Padasavali	265	7.78	NHAmB2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cultivated Fallow+Redgram (CF+Rg)	Not Available	IIIse	Crescent bund/TCB
Padasavali	266	12.5	MANmA1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Greengram+Jowar+Sugar cane+Scrub land (Gg+Jw+Sc+Sl)	Openwell,Openwell	IIs	GB/strengthening of Bunds
Padasavali	267	7.98	MANmA1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Sugarcane (Sc)	Not Available	IIs	GB/strengthening of Bunds
Padasavali	273	0.71	MANmA1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Greengram+Redgram (Gg+Rg)	Not Available	IIs	GB/strengthening of Bunds
Padasavali	274	0.27	MANmA1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Cultivated Fallow (CF)	Not Available	IIs	GB/strengthening of Bunds
Padasavali	STREAM	1.13	MANmA1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Others	Not Available	IIs	GB/strengthening of Bunds

Appendix – II

Soil Fertility Information

Village	Survey No.	Soil Reaction (pH)	EC	Organic Carbon	Available Phosphorus	Available Potassium	Available Sulphur	Available Boron	Available Iron	Available Manganese	Available Copper	Available Zinc
Niragudi	127	Neutral (pH 6.5-7.3)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	Medium (10-20 ppm)	Low (<0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Niragudi	128	Neutral (pH 6.5-7.3)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	Medium (10-20 ppm)	Low (<0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Niragudi	129	Neutral (pH 6.5-7.3)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	Medium (10-20 ppm)	Low (<0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Niragudi	130	Neutral (pH 6.5-7.3)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	Medium (10-20 ppm)	Low (<0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Sufficient (>0.6 ppm)
Niragudi	131	Slightly alkaline (pH 7.3-7.8)	Non Saline (<2 dsm)	High (>0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	Medium (10-20 ppm)	Low (<0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Sufficient (>0.6 ppm)
Niragudi	132	Moderately alkaline (pH 7.8-8.4)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	Medium (10-20 ppm)	Low (<0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Niragudi	134	Moderately alkaline (pH 7.8-8.4)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	Low (<10 ppm)	Low (<0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Niragudi	135	Moderately alkaline (pH 7.8-8.4)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	Medium (10-20 ppm)	Medium (0.5-1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Niragudi	136	Slightly alkaline (pH 7.3-7.8)	Non Saline (<2 dsm)	High (>0.75 %)	Low (<23 kg/ha)	Medium (145-337 kg/ha)	Low (<10 ppm)	Low (<0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Sufficient (>0.6 ppm)
Niragudi	137	Slightly alkaline (pH 7.3-7.8)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	Medium (10-20 ppm)	Low (<0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Sufficient (>0.6 ppm)
Niragudi	138	Slightly alkaline (pH 7.3-7.8)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	Medium (10-20 ppm)	Low (<0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Niragudi	139	Moderately alkaline (pH 7.8-8.4)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	Medium (10-20 ppm)	Low (<0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Niragudi	140	Moderately alkaline (pH 7.8-8.4)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	Low (<10 ppm)	Low (<0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Niragudi	141	Moderately alkaline (pH 7.8-8.4)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	Medium (145-337 kg/ha)	Medium (10-20 ppm)	Low (<0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Niragudi	142	Slightly alkaline (pH 7.3-7.8)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	Medium (10-20 ppm)	Low (<0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Sufficient (>0.6 ppm)
Niragudi	143	Slightly alkaline (pH 7.3-7.8)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	Medium (145-337 kg/ha)	Medium (10-20 ppm)	Low (<0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)

Village	Survey No.	Soil Reaction (pH)	EC	Organic Carbon	Available Phosphorus	Available Potassium	Available Sulphur	Available Boron	Available Iron	Available Manganese	Available Copper	Available Zinc
Niragudi	260	Moderately alkaline (pH 7.8-8.4)	Non Saline (<2 dsm)	High (>0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	Medium (10-20 ppm)	Low (<0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Niragudi	261	Moderately alkaline (pH 7.8-8.4)	Non Saline (<2 dsm)	High (>0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	Medium (10-20 ppm)	Low (<0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Niragudi	262	Slightly alkaline (pH 7.3-7.8)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Medium (23-57 kg/ha)	High (>337 kg/ha)	Medium (10-20 ppm)	Low (<0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Sufficient (>0.6 ppm)
Niragudi	281	Neutral (pH 6.5-7.3)	Non Saline (<2 dsm)	High (>0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	Medium (10-20 ppm)	Low (<0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Niragudi	STRE AM	Others	Waterbody	Others	Others	Others	Others	Others	Others	Others	Others	Others
Padasavali	6	Moderately alkaline (pH 7.8-8.4)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	High (>20 ppm)	Medium (0.5-1.0 ppm)	Deficient (<4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Padasavali	7/1	Strongly alkaline (pH 8.4-9.0)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	High (>20 ppm)	Medium (0.5-1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Padasavali	7/2	Strongly alkaline (pH 8.4-9.0)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	High (>20 ppm)	Medium (0.5-1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Padasavali	7/3	Strongly alkaline (pH 8.4-9.0)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	High (>20 ppm)	Medium (0.5-1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Padasavali	8	Strongly alkaline (pH 8.4-9.0)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	High (>20 ppm)	Medium (0.5-1.0 ppm)	Deficient (<4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Padasavali	9	Moderately alkaline (pH 7.8-8.4)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	High (>20 ppm)	Low (<0.5 ppm)	Deficient (<4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Padasavali	11	Moderately alkaline (pH 7.8-8.4)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	High (>20 ppm)	Low (<0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Padasavali	12	Slightly alkaline (pH 7.3-7.8)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Medium (23-57 kg/ha)	High (>337 kg/ha)	High (>20 ppm)	Low (<0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Padasavali	13	Moderately alkaline (pH 7.8-8.4)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	High (>20 ppm)	Low (<0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Padasavali	14	Moderately alkaline (pH 7.8-8.4)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	High (>20 ppm)	Low (<0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Padasavali	15	Slightly alkaline (pH 7.3-7.8)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Medium (23-57 kg/ha)	High (>337 kg/ha)	High (>20 ppm)	Low (<0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Padasavali	16	Moderately alkaline (pH 7.8-8.4)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	High (>20 ppm)	Low (<0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Padasavali	17	Moderately alkaline (pH 7.8-8.4)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	High (>20 ppm)	Low (<0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)

Village	Survey No.	Soil Reaction (pH)	EC	Organic Carbon	Available Phosphorus	Available Potassium	Available Sulphur	Available Boron	Available Iron	Available Manganese	Available Copper	Available Zinc
Padasavali	256	Moderately alkaline (pH 7.8-8.4)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	Medium (10-20 ppm)	Low (<0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Padasavali	257	Moderately alkaline (pH 7.8-8.4)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	High (>20 ppm)	Low (<0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Padasavali	258	Moderately alkaline (pH 7.8-8.4)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	Medium (10-20 ppm)	Low (<0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Padasavali	259	Moderately alkaline (pH 7.8-8.4)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	High (>20 ppm)	Low (<0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Sufficient (>0.6 ppm)
Padasavali	260	Moderately alkaline (pH 7.8-8.4)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	High (>20 ppm)	Low (<0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Padasavali	261	Moderately alkaline (pH 7.8-8.4)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	High (>20 ppm)	Medium (0.5-1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Padasavali	262	Moderately alkaline (pH 7.8-8.4)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	High (>20 ppm)	Low (<0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Sufficient (>0.6 ppm)
Padasavali	263	Slightly alkaline (pH 7.3-7.8)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	Medium (10-20 ppm)	Low (<0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Sufficient (>0.6 ppm)
Padasavali	264	Slightly alkaline (pH 7.3-7.8)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	Medium (10-20 ppm)	Low (<0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Sufficient (>0.6 ppm)
Padasavali	265	Moderately alkaline (pH 7.8-8.4)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	Medium (10-20 ppm)	Low (<0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Sufficient (>0.6 ppm)
Padasavali	266	Moderately alkaline (pH 7.8-8.4)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	High (>20 ppm)	Medium (0.5-1.0 ppm)	Deficient (<4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Sufficient (>0.6 ppm)
Padasavali	267	Moderately alkaline (pH 7.8-8.4)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	High (>20 ppm)	Low (<0.5 ppm)	Deficient (<4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Padasavali	273	Strongly alkaline (pH 8.4-9.0)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	High (>20 ppm)	Low (<0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Padasavali	274	Strongly alkaline (pH 8.4-9.0)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	High (>20 ppm)	Medium (0.5-1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Padasavali	STRE AM	Moderately alkaline (pH 7.8-8.4)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	High (>20 ppm)	Low (<0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)

Appendix – III

Soil Fertility Information

Village	Survey No.	Sorghum	Maize	Red-gram	Sun-flower	Cotton	Sugar-cane	Soy-bean	Guava	Mango	Sapota	Jack-fruit	Jamun	Mus-ambi	Lime	Cashew	Custard Apple	Amla	Tamarind
Niragudi	127	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N
Niragudi	128	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N
Niragudi	129	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N
Niragudi	130	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N
Niragudi	131	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N
Niragudi	132	S3r	S3rt	S3r	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Niragudi	134	S3r	S3rt	S3r	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Niragudi	135	S3r	S3rt	S3r	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Niragudi	136	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N
Niragudi	137	S3r	S3rt	S3r	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Niragudi	138	S3r	S3rt	S3r	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Niragudi	139	S3r	S3rt	S3r	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Niragudi	140	S3r	S3rt	S3r	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Niragudi	141	S3r	S3rt	S3r	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Niragudi	142	S3r	S3rt	S3r	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Niragudi	143	S3r	S3rt	S3r	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Niragudi	144	S3r	S3rt	S3r	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Niragudi	145	S3r	S3rt	S3r	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Niragudi	146	S3r	S3rt	S3r	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Niragudi	147	S3r	S3rt	S3r	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Niragudi	148	S3r	S3rt	S3r	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Niragudi	149	S3r	S3rt	S3r	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Niragudi	150	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N
Niragudi	151	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	S3rg	S3rg	N
Niragudi	154	S3r	S3rt	S3r	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Niragudi	155	S3r	S3rt	S3r	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Niragudi	156	S3r	S3rt	S3r	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N

Village	Survey No.	Sorghum	Maize	Red-gram	Sun-flower	Cotton	Sugar-cane	Soy-bean	Guava	Mango	Sapota	Jack-fruit	Jamun	Mus-ambi	Lime	Cas hew	Custard Apple	Amla	Tam-arind
Niragudi	157	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N
Niragudi	158	S1	S3t	S2t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S2r
Niragudi	159	S1	S3t	S2t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S2r
Niragudi	160	S1	S3t	S2t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S2r
Niragudi	161	S1	S3t	S2t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S2r
Niragudi	162	S1	S3t	S2t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S2r
Niragudi	163	S1	S3t	S2t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S2r
Niragudi	164	S1	S3t	S2t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S2r
Niragudi	165	S1	S3t	S2t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S2r
Niragudi	166	S1	S3t	S2t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S2r
Niragudi	167	S1	S3t	S2t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S2r
Niragudi	168	S1	S3t	S2t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S2r
Niragudi	169	S1	S3t	S2t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S2r
Niragudi	170	S1	S3t	S2t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S2r
Niragudi	171	S1	S3t	S2t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S2r
Niragudi	172	S1	S3t	S2t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S2r
Niragudi	173	S1	S3t	S2t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S2r
Niragudi	174	S1	S3t	S2t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S2r
Niragudi	175	S1	S3t	S2t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S2r
Niragudi	176	S1	S3t	S2t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S2r
Niragudi	177	S1	S3t	S2t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S2r
Niragudi	178	S1	S3t	S2t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S2r
Niragudi	179	S1	S3t	S2t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S2r
Niragudi	180	S1	S3t	S2t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S2r
Niragudi	181	S1	S3t	S2t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S2r
Niragudi	182	S1	S3t	S2t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S2r
Niragudi	183	S1	S3t	S2t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S2r
Niragudi	184	S1	S3t	S2t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S2r
Niragudi	185	S1	S3t	S2t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S2r

Village	Survey No.	Sorghum	Maize	Red-gram	Sun-flower	Cotton	Sugar-cane	Soy-bean	Guava	Mango	Sapota	Jack-fruit	Jamun	Mus-ambi	Lime	Cashew	Custard Apple	Amla	Tamarind
Niragudi	186	S3r	S3rt	S3r	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Niragudi	187	S3r	S3rt	S3r	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Niragudi	188	S3r	S3rt	S3r	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Niragudi	189	S3r	S3rt	S3r	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Niragudi	193	S3r	S3rt	S3r	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Niragudi	232	S3r	S3rt	S3r	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Niragudi	233	S1	S3t	S2t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S2r
Niragudi	234	S1	S3t	S2t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S2r
Niragudi	235	S1	S3t	S2t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S2r
Niragudi	236	S1	S3t	S2t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S2r
Niragudi	237	S1	S3t	S2t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S2r
Niragudi	238	S1	S3t	S2t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S2r
Niragudi	239	S1	S3t	S2t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S2r
Niragudi	240	S1	S3t	S2t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S2r
Niragudi	241	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N
Niragudi	242	S1	S3t	S2t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S2r
Niragudi	243	S1	S3t	S2t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S2r
Niragudi	244	S1	S3t	S2t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S2r
Niragudi	245	S1	S3t	S2t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S2r
Niragudi	246	S1	S3t	S2t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S2r
Niragudi	247	S1	S3t	S2t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S2r
Niragudi	248	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N
Niragudi	249	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N
Niragudi	250	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N
Niragudi	251	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N
Niragudi	252	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N
Niragudi	253	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N
Niragudi	254	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N
Niragudi	255	S3r	S3rt	S3r	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N

Village	Survey No.	Sorghum	Maize	Red-gram	Sun-flower	Cotton	Sugar-cane	Soy-bean	Guava	Mango	Sapota	Jack-fruit	Jamun	Mus-ambi	Lime	Cashew	Custard Apple	Amla	Tamarind
Niragudi	257	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N
Niragudi	258	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N
Niragudi	259	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N
Niragudi	260	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N
Niragudi	261	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N
Niragudi	262	S3r	S3rt	S3r	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Niragudi	281	S3rg	S3rt	S3rg	N	S3rg	N	S3rg	N	N	N	N	N	N	N	N	S3rg	S3rg	N
Niragudi	STREAM	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Padasavali	6	S1	S3t	S2t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S2r
Padasavali	7/1	S1	S3t	S2t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S2r
Padasavali	7/2	S1	S3t	S2t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S2r
Padasavali	7/3	S1	S3t	S2t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S2r
Padasavali	8	S1	S3t	S2t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S2r
Padasavali	9	S1	S3t	S2t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S2r
Padasavali	11	S1	S3t	S2t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S2r
Padasavali	12	S3r	S3rt	S3r	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Padasavali	13	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N
Padasavali	14	S3r	S3rt	S3r	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Padasavali	15	S3r	S3rt	S3r	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Padasavali	16	S1	S3t	S2t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S2r
Padasavali	17	S1	S3t	S2t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S2r
Padasavali	18	S3r	S3rt	S3r	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Padasavali	19	S1	S3t	S2t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S2r
Padasavali	20	S1	S3t	S2t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S2r
Padasavali	21	S1	S3t	S2t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S2r
Padasavali	22	S1	S3t	S2t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S2r
Padasavali	23	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N
Padasavali	24	S3r	S3rt	S3r	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N

Village	Survey No.	Sorghum	Maize	Red-gram	Sun-flower	Cotton	Sugar-cane	Soy-bean	Guava	Mango	Sapota	Jack-fruit	Jamun	Mus-ambi	Lime	Cashew	Custard Apple	Amla	Tamarind
Padasavali	25	S3r	S3rt	S3r	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Padasavali	26	S3r	S3rt	S3r	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Padasavali	27	S3r	S3rt	S3r	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Padasavali	28	S3r	S3rt	S3r	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Padasavali	29	S3r	S3rt	S3r	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Padasavali	57	S3r	S3rt	S3r	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Padasavali	251	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N
Padasavali	252	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N
Padasavali	253	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N
Padasavali	254	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N
Padasavali	255	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N
Padasavali	256	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N
Padasavali	257	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N
Padasavali	258	S2re	S3t	S2re	S3r	S2re	S3t	S2re	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2re	S2re	N
Padasavali	259	S2re	S3t	S2re	S3r	S2re	S3t	S2re	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2re	S2re	N
Padasavali	260	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N
Padasavali	261	S1	S3t	S2t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S2r
Padasavali	262	S2re	S3t	S2re	S3r	S2re	S3t	S2re	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2re	S2re	N
Padasavali	263	S2re	S3t	S2re	S3r	S2re	S3t	S2re	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2re	S2re	N
Padasavali	264	S3r	S3rt	S3r	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Padasavali	265	S3r	S3rt	S3r	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Padasavali	266	S1	S3t	S2t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S2r
Padasavali	267	S1	S3t	S2t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S2r
Padasavali	273	S1	S3t	S2t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S2r
Padasavali	274	S1	S3t	S2t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S2r
Padasavali	STREAM	S1	S3t	S2t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S2r

PART-B

SOCIO-ECONOMIC STATUS OF FARM HOUSEHOLDS

CONTENTS

1	Executive summary	1-3
2	Introduction	5
3	Methodology	7-11
4	Results and discussions	13-29

LIST OF TABLES

I. Social status		
1	Human population among sample households	13
2	Basic needs of sample households	15
II. Economic status		
3	Occupational pattern in sample households	15
4	Domestic assets among samples households	16
5	Farm assets among samples households	16
6	Livestock assets among sample households	18
7	Milk produced and Fodder availability of sample households	18
8	Women empowerment of sample households	18
9	Per capita daily consumption of food among the sample households	19
10	Annual average Income from various sources	20
11	Average annual expenditure of sample households	20
12	Distribution of land holding among the sample households	21
13	Land holding among samples households	21
III. Resource use pattern		
14	Number of tree/plants covered in sample farm households	22
15	Present cropping pattern among samples households	22
16	Distribution of soil series in the watershed	23-24
IV. Economic land evaluation		
17	Cropping pattern on major soil series	24
18	Alternative land use options for different size group of farmers (Benefit Cost Ratio)	24
19	Economics Land evaluation and bridging yield gap for different crops	25
20	Estimation of onsite cost of soil erosion	27
21	Ecosystem services of food grain production	27
22	Ecosystem services of fodder production	28
23	Ecosystem services of water supply for crop production	28
24	Farming constraints	29

LIST OF FIGURES

1	Location of study area	8
2	ALPES Framework	9
3	Basic needs of sample households	14
4	Domestic assets among the sample households	16
5	Farm assets among samples households	17
6	Livestock assets among sample households	17
7	Per capita daily consumption of food among the sample households	19
8	Average annual expenditure of sample households	20
9	Present cropping pattern	22
10	Estimation of onsite cost of soil erosion	26
11	Ecosystem services of food grain production	27
12	Ecosystem services of water supply	28

EXECUTIVE SUMMARY

Baseline socioeconomic characterisation is prerequisite to prepare action plan for program implementation and to assess the project performance before making any changes in the watershed development program. The baseline provides appropriate policy direction for enhancing productivity and sustainability in agriculture.

Methodology: *Nirgudi-1 micro-watershed (Alur sub-watershed, Aland taluk, Gulbarga district) is located in between 17°37'–17°39' North latitudes and 77°26'–77°27' East longitudes, covering an area of about 808.47 ha, bounded by ChincholiBudruk, Chincholikurd, Nagalogaon, Matki villages and Maharashtra State with length of growing of period (LGP) 120-150 days. We used soil resource map as basis for sampling farm households to test the hypothesis that soil quality influence crop selection, and conservation investment of farm households. The level of technology adoption and productivity gaps and livelihood patterns were analyses. The cost of soil degradation and ecosystem services were quantified.*

Results: *The socio-economic outputs for Nirgudi-1 micro-watershed (Alur sub-watershed, Aland taluk, Gulbarga district) are presented here.*

Social Indicators;

- ❖ *Male and female ratio is 52.2 to 47.8 per cent to the total sample population.*
- ❖ *Younger age 18 to 50 years group of population is around 73.9 per cent to the total population.*
- ❖ *Literacy population is around 69.6 per cent.*
- ❖ *Social groups belong to general caste is around 80.0 per cent.*
- ❖ *Fire wood is the source of energy for a cooking among all the households.*
- ❖ *About 10 per cent of households have a yashaswini health card.*
- ❖ *About 30.0 percent of the farm households are having MGNREGA card for rural employment.*
- ❖ *Dependence on ration cards for food grains through public distribution system is around 80 per cent.*
- ❖ *Swach bharath program providing closed toilet facilities around 40 per cent of sample households.*
- ❖ *Women participation in decisions making are around 82 per cent of households were found.*

Economic Indicators;

- ❖ *The average land holding is 1.7 ha indicates that majority of farm households are belong to small and medium farmers. The dry land of 48.2 % and irrigated land 51.8 % of total cultivated land area among the sample farmers.*

- ❖ Agriculture is the main occupation among 15.2 per cent and agriculture is the main and agriculture labour is subsidiary occupation for 82.6 per cent of sample households.
- ❖ The average value of domestic assets is around Rs. 21777 per household. Mobile and television are popular media mass communication.
- ❖ The average value of farm assets is around Rs. 166952 per household, about 50.0 per cent of sample farmers having bullock cart.
- ❖ The average value of livestock is around Rs. 30933 per household; about 82 per cent of household are having livestock.
- ❖ The average per capita food consumption is around 758.7 grams (1938.5 kilo calories) against national institute of nutrition (NIN) recommendation at 827 gram. Around 40.5 per cent of sample households are consuming less than the NIN recommendation.
- ❖ The annual average income is around Rs.35394 per household. About 40.0 per cent of farm households are below poverty line.
- ❖ The per capita average monthly expenditure is around Rs.2231.

Environmental Indicators-Ecosystem Services;

- ❖ The value of ecosystem service helps to support investment to decision on soil and water conservation and in promoting sustainable land use.
- ❖ The onsite cost of different soil nutrients lost due to soil erosion is around Rs. 1505 per ha/year. The total cost of annual soil nutrients is around Rs. 1192065 per year for the total area of 808.47 ha.
- ❖ The average value of ecosystem service for food grain production is around Rs. 13470/ha/year. Per hectare food grain production services is maximum in sunflower (Rs. 21495) followed by red gram (Rs. 17447), black gram (Rs.11930) and sorghum (Rs. 3007).
- ❖ The average value of ecosystem service for fodder production is around Rs. 1002/ha/year in sorghum.
- ❖ The data on water requirement for producing one quintal of grain is considered for estimating the total value of water required for crop production. Per hectare value of water used and value of water was maximum in black gram (Rs. 60116) followed by redgram (Rs. 47467), sorghum (Rs. 30628) and sunflower (Rs. 33256).

Economic Land Evaluation;

- ❖ The major cropping pattern is red gram (72.9 %) followed by sunflower (12.0 %) sorghum (10.1 %) and blackgram (5.0 %).
- ❖ In Nirgudi-1 Microwatershed, major soil is Margutti (MGT) series is having very shallow soil depth cover around 37.6 % of area. On this soil farmers are presently black gram (18.5%) and red gram (81.5 %). Soil of granite and granite gneiss landscape of Novinahala (NHA) are also having shallow soil depth cover 34.1 % of area, the crops are redgram (67 %), sorghum (10.1 %) and sunflower (22.9 %).

Bhimanahalli (BHI) soil series having shallow soil depth cover around 1.7 % of areas, crop is red gram. Mahagaon (MAN) soil series having very deep soil depth cover around 19.83 % of area, crops are red gram (60.6 %) and sorghum (39.4 %).

- ❖ *The total cost of cultivation and benefit cost ratio (BCR) in study area for red gram ranges between Rs. 26179/ha in NHA soil (with BCR of 1.63) and Rs. 16343/ha in MGT soil (with BCR of 1.52).*
- ❖ *In sorghum the cost of cultivation range between Rs 19584/ha in MAN soil (with BCR of 1.19) and Rs. 14596/ha in NHA soil (with BCR of 1.30).*
- ❖ *In black gram the cost of cultivation in MGT soil is Rs 22890/ha (with BCR of 1.52) and sunflower the cost of cultivation in NHA soil is Rs 18025/ha (with BCR of 2.19).*
- ❖ *The land management practices reported by the farmers are crop rotation, tillage practices, fertilizer application and use of farm yard manure (FYM). Due to higher wages farmer are following labour saving strategies is not prating soil and water conservation measures. Less ownership of livestock limiting application of FYM.*
- ❖ *It was observed soil quality influences on the type and intensity of land use. More fertilizer applications in deeper soil to maximize returns.*

Suggestions;

- ❖ *Involving farmers in watershed planning helps in strengthening institutional participation.*
- ❖ *The per capita food consumption and monthly income is very low. Diversifying income generation activities from crop and livestock production in order to reduce risk related to drought and market prices.*
- ❖ *Majority of farmers reported that they are not getting timely support/extension services from the concerned development departments.*
- ❖ *By strengthening agricultural extension for providing timely advice improved technology there is scope to increase in net income of farm households.*
- ❖ *By adopting recommended package of practices by following the soil test fertiliser recommendation, there is scope to increase yield in redgram (1.4 to 39.3 %), sorghum (60.4 to 68 %), sunflower (39.3 %) and black gram (10.8 %).*

INTRODUCTION

Watershed Development program aim to restore degraded watersheds in rainfed regions to increase their capacity to capture and store rain water, reduce soil erosion, and improved soil nutrients and carbon contents so they can produce greater agricultural yields and other benefits. As majority of rural poor live in these regions and dependent on natural resources for their livelihood and sustenance, improvements in agricultural yields improve human welfare and simultaneously improve national food security.

Sujala–III watershed development project conceptualised and implemented by the Watershed Development Department of Government of Karnataka with tripartite cost-sharing arrangements. The World Bank through International Development Association provided major portion of plan outlay as a loan to Government of India and in turn loan to Government of Karnataka.

The objectives of Sujala-III is to demonstrate more effective watershed management through greater integration of programs related to rain fed agriculture, innovative and science based approaches and strengthened institutions and capacities. The project is implemented in 11 districts of Bidar, Vijayapura, Gulbarga, Yadgir, Koppal, Gadag, Raichur, Davanagere, Tumkur, Chikkamangalur and Chamarajanagar which have been identified by the Watershed Development Department based on rainfall and socio-economic conditions. The project will be implemented over six years and linked with the centrally financed integrated watershed management programme.

Economic evaluations can better guide in watershed planning and implementation, as well as raise awareness of benefits of ecosystem restoration for food security and poverty alleviation program. The present study aims to characterize socio-economic status of farm households, assess the land and water use status, evaluate the economic viability of land use, prioritize farming constraints and suggest the measures for soil and water conservation for sustainable agriculture.

Objectives of the study

1. To characterize socio-economic status of farm households
2. To evaluate the economic viability of land use and land related constraints
3. To estimate the ecosystem service provided by the watershed and
4. To suggest alternatives for sustainable agriculture production.

METHODOLOGY

Study area

Nirgudi-1 micro-watershed located in north-eastern dry zone of Karnataka (Figure 1): The total geographic area of this zone is about 1.76 M ha covering 8 taluks of Gulbarga district and 3 taluks of Raichur. Net cultivated area in the zone is about 1.31 M ha of which about 0.09 M ha are irrigated. The mean elevation of the zone is 300-450 m MSL. The main soil type is deep to very deep soils with small pockets of shallow to medium black soils. The zone is cropped predominantly during rabi due to insufficient rainfall (465-785 mm). The principal crops of the zone are jowar, bajra, oilseeds, pulses, cotton and sugarcane. It's represented Agro Ecological Sub Region (AESR) 6.2 having LGP 120-150 days.

Nirgudi-1 micro-watershed (Alur sub-watershed, Aland taluk, Gulbarga district) is located in between 17⁰37'–17⁰39' North latitudes and 77⁰26'–77⁰27' East longitudes, covering an area of about 808.47 ha, bounded by ChincholiBudruk, Chincholikhurd, Nagalogaon, Matki villages and Maharashtra State.

Sampling Procedure:

In this study we have followed soil variability as criterion for sampling the farm households. In each micro-watershed the survey numbers and associated soil series are listed. Minimum three farm households for each soil series were taken and summed up to arrive at total sample for analysis.

Sources of data and analysis:

For evaluating the specific objectives of the study, primary data was collected from the sample respondents by personal interview method with the help of pre-tested questionnaire. The data on socio-economic characteristics of respondents such as family size and composition, land holdings, asset position, occupational pattern and education level was collected. The present cropping pattern and the level of input use and yields collected during survey. The data collected from the representative farm households were analysed using Automated Land Potential Evaluation System (Figure 2).

LOCATION MAP OF NIRGUDI-1 MICRO-WATERSHED




Figure 1: Location of study area

Steps followed in socio-economic assessment

- 1 •After the completion of soil profile study link the cadastral number to the soil profile in the micro watershed.
- 2 • Download the names of the farmers who are owning the land for each cadastral number in the Karnataka BHOOMI Website.
- 3 • Compiling the names of the farmers representing for all the soil profiles studied in the micro watershed for socio-economic Survey.
- 4 • Conducting the socioeconomic survey of selected farm households in the micro watershed .
- 5 • Farm households database created using the Automated Land Potential Evaluation System (ALPES) for analysis of socio economic status for each micro watershed .
- 6 • Synthesis of tables and preparation of report for each micro watershed .

KOPPAL-Kavalur Village 1-213

Home Create External Data Database Tools

Main

AUTOMATED LAND POTENTIAL EVALUATION SYSTEM

ALPES

FARM HOUSE HOLD DIGITAL LIBRARY

Form View

KOPPAL-Kavalur Village 1-213

Home Create External Data Database Tools

Main Report

Livelihood Patterns

I. SOCIAL STATUS

- Table 1. Spatial distribution of sample
- Table 3. Total human population in sample HHs
 - Average Family size
 - Age groups
 - Education status
- Table 4. Drinking water facilities
- Table 5. Institutional participation
 - Average age of HHs

Farming Constraints

- Farmers awareness of climate change
 - Reasons
- ICT Use for agril information

II. ECONOMIC STATUS

- Table 6: Migration Details
 - Nature of Job migrates
- Table 7. Domestic assets
- Table 8 Farm assets
- Table 9 Livestock assets
- Table 10 Food consumption in KiloCalories
 - Grams
- Table 11 Annual average income (Rs)
- Table 12 Crop production income
- Table 13 Average annual Expenditure
 - Poverty estimation
 - Priority for crop selection
 - Kind of Government support
 - Socio-economic Indicators

III. RESOURCE USE PATTERN

- Table 14 Land holding sample HHs
- Table 15 Land use among sample HHs
- Table 16 Cropping pattern

IV. ECONOMIC LAND EVALUATION

- Table 17. Cropping pattern on diff soil series
- Table 18. Economics different crops in Soil Series
 - Yield gap analysis of crops

Optimum land use planning

- Input-Output coefficients

Ecosystem

- Food Provision
- Fodder Provision
- Livestock Economics
- Water consumption budgeting

Record: 14 of 1 of 1 No Filter Search

Figure 2: ALPES FRAMEWORK

The sample farmers were post classified in to marginal and small (0.0 to ≤ 2 ha), medium and semi medium (> 2 to ≤ 10 ha) and large (> 10 ha). The steps involved in estimation of soil potential involve estimation of total cost of cultivation, the yield/gross returns and net income per hectare. The cost of inputs such seed, manure and fertilizer, plant protection chemicals, payment towards human and bullock labour and interest on working capita are included under operational costs. In the case of perennial crops, the cost of establishment was estimated by using actual physical requirements and prevailing market prices. Estimation cost included maintenance cost up to bearing period. The value of main product and by product from the crop enterprise at the market rates were the gross returns of the crop. Net returns were worked out by deducting establishment and maintained cost from gross returns.

Operational Cost = cost of seeds, fertilizers, pesticides. Cost of human and bullock labour, cost of machinery, cost of irrigation water + interest on working capital.

Gross returns = Yield (Quintals/hectare)*Price (Rs/Quintal)

Net returns = Gross returns-Operational cost.


Benefit Cost Ratio = Net returns/Total cost.

Economic suitability classes: once each land use –land area combination has been assigned an economic value by the land evaluation, the question arises as to its ‘suitability’, that is, the degree to which it satisfies the land user. The FAO framework defines two suitability orders: ‘S’(suitable if benefit cost ratio (BCR) > 1) and ‘N’(not suitable if (BCR < 1), which are divided into five economic suitability classes: ‘S1’(highly suitable if BCR > 3), ‘S2’(suitable if BCR > 2 and < 3), ‘S3’(Marginally suitable if BCR > 1 and < 2), ‘N1’(Not suitable for economic reasons but physically suitable) and ‘N2’(not suitable for physical reasons). The limit between ‘S3’ and ‘N1’ must be at least at the point of financial feasibility (i.e. net returns, NPV, or IRR > 0 and BCR > 1). The other limits depend on social factors such as farm size, family size, alternative employment or investment possibilities and wealth expectations; these need to be specified for the Soil series.

Economic Valuation of Soil ecosystem services:

The replacement cost approach was followed for estimating the onsite cost of soil erosion, Market price method was followed for estimating the value of food and fodder production. Value transfer methods was followed for estimating the value of water demand by different crops in the micro watershed.

Steps followed in Replacement cost methods for estimation of onsite cost of soil erosion



RESULTS AND DISCUSSIONS

The demographic information shows that the household population dynamics encompasses the socioeconomic status of the farmer. For a rural family, the household size should be optimal to earn a comfortable livelihood through farm and non-farm wage earning. The total number of population in watershed area was 46, out of which 52.2 per cent were males and 47.8 per cent females. Average family size of the households is 4.6. Age is an important factor, which affects the potential employment and mobility status of respondents. The data on age wise distribution of farmers in the sample households indicated that majority of the farmers are coming under the age group of followed by 30 to 50 years (39.1 %), 18 to 30 years (34.8%) and more than 50 years (19.6 %). Hence, in the study area in general, the respondents were of young and middle age, indicating thereby that the households had almost settled with whatever livelihood options they were practicing and sample respondents were young by age who could venture into various options of livelihood sources. Data on literacy indicated that 30.4 per cent of respondents were illiterate and 69.6 per cent literate (Table 1).

Table 1: Human population among sample households in Nirgudi-1 Microwatershed

Particulars	Units	Value
Total human population in sample HHs	Number	46
Male	% to total Population	52.2
Female	% to total Population	47.8
Average family size	Number	4.6
Age group		
0 to 18 years	% to total Population	6.5
18 to 30 years	% to total Population	34.8
30 to 50 years	% to total Population	39.1
>50 years	% to total Population	19.6
Average age	Age in years	36.7
Education Status		
Illiterates	% to total Population	30.4
Literates	% to total Population	69.6
Primary School (<5 class)	% to total Population	15.2
Middle School (6- 8 class)	% to total Population	15.2
High School (9- 10 class)	% to total Population	23.9
Others	% to total Population	15.2

The ethnic groups among the sample farm households found to be 80 per cent belonging to general castes followed by 10 per cent belonging to scheduled tribe (ST) and 10 percent belonging to other backward caste (OBC) (Table 2 and Figure 3). All the sample households are using fire wood as source of fuel for cooking. All the sample farmers are having electricity connection. About 10 per cent are sample households having health cards. About 30 percent of households are having MNREGA job cards for employment generation. About 80 per cent of farm households are having ration cards for

taking food grains from public distribution system. About 40 per cent of farm households are having toilet facilities.




Figure 3: Basic needs of sample households in Nirgudi-1 Microwatershed

The data collected on the source of drinking water in the study area is presented in Table 2. Majority of the sample respondents are having tube well source for water supply for domestic purpose.

Table 2: Basic needs of sample households in Nirgudi-1 Microwatershed

Particulars	Units	Value
Social groups		
ST	% of Households	10.0
OBC	% of Households	10.0
General	% of Households	80.0
Types of fuel use for cooking		
Fire wood	% of Households	100.0
0	% of Households	0.0
Energy supply for home		
Electricity	% of Households	100.0
Number of households having Health card		
Yes	% of Households	10.0
No	% of Households	90.0
MGNREGA Card		
Yes	% of Households	30.0
No	% of Households	70.0
Ration Card		
Yes	% of Households	80.0
No	% of Households	20.0
Households with toilet		
Yes	% of Households	40.0
No	% of Households	60.0
Drinking water facilities		
Tube Well	% of Households	100.00

The occupational pattern (Table 3) among sample households shows that agriculture is the main occupation around 15.2 per cent of farmers followed by agriculture is the main occupation and subsidiary occupations like agricultural labour (82.6 %) and non agricultural labour (2.1 %).

Table 3: Occupational pattern in sample population in Nirgudi-1 Micro-watershed

Occupation		% to total
Main	Subsidiary	
Agriculture	Agriculture	15.2
	Agriculture Labour	82.6
	Non Agriculture Labour	2.1
Family labour availability		Man days/month
Male		52.5
Female		44.0
Total		96.5

The important assets especially with reference to domestic assets were analyzed and are given in Table 4 and Figure 4. The important domestic assets possessed by all categories of farmers are television (100 %) followed by mobile phones (90 %), mixer/grinder (10 %), and motorcycle (10 %). The average value of domestic assets is around Rs. 21777 per household.

Table 4: Domestic assets among the sample households in Nirgudi-1 Micro-watershed

Particulars	% of households	Average value in Rs
Mixer/grinder	10.0	2000
Mobile Phone	90.0	4111
Motorcycle	10.0	70000
Television	100.0	11000
Average Value		21777




Figure 4: Domestic assets among the sample households in Nirgudi-1 Micro-watershed

The most popularly owned farm equipments were sickles, plough, cattle shed; pump sets, chaff cutter, bullock cart, sprayer and thresher. Plough and sickle were commonly present in all the sampled farmers; these were primary implements in agriculture. The per cent of households owned bullock cart (50 %), plough (40 %), sprayer (20 %), tractor (10 %) and weeder (50.0 %). The average value of farm assets is around Rs.166952 per households (Table 5 and Figure 5).

Table 5: Farm assets among samples households in Nirgudi-1 Micro-watershed

Particulars	% of households	Average value in Rs
Bullock cart	50.0	25200
Plough	40.0	6000
Sprayer	20.0	3200
Tractor	10.0	800000
Weeder	50.0	364
Average Value		166952




Figure5: Farm assets among samples households in Nirgudi-1 Micro-watershed




Figure 6: Livestock assets among sample households in Nirgudi-1 Microwatershed

Livestock is an integral component of the conventional farming systems (Table 6 and Figure 6). The highest livestock population is local milching cow were around 27.8 per cent, local dry cow (16.7 %), milching buffalos (16.7 %) and dry buffalos (5.6 %). The average livestock value was Rs 30933 per household.

Table 6: Livestock assets among sample households in Nirgudi-1 Microwatershed

Particulars	% of livestock population	Average value in Rs
Local Dry Cow	16.7	11667
Local Milching Cow	27.8	18000
Dry Buffalos	5.6	60000
Milching Buffalos	16.7	30000
Bullocks	33.3	35000
Average value	30933	

Average milk produced in sample households is 808 litters/annum. Among the farm households the crops are sorghum is the domestic food and fodder for animals. About 1563 kg /ha of average fodder is available per season for the livestock feeding (Table 7).

Table 7: Milk produced and fodder availability of sample households in Nirgudi-1 Micro-watershed

Particulars	
Name of the Livestock	Ltr./Lactation/animal
Local Milching Cow	882
Milching Buffalos	733
Average Milk Produced	808
Fodder produces	
Fodder yield (kg/ha.)	
Sorghum	1563
Average fodder availability	1563
Livestock having households (%)	82
Livestock population (Numbers)	46

A woman participation in decision making in this micro watershed is presented in Table 8. All among women taking decision in her family and agriculture related activities and all the farm households' women earning for her family requirement.

Table 8: Women empowerment of sample households in Nirgudi-1 Micro-watershed % to Grand Total

Particulars	Yes	No
Women participation in local organization activities	00.0	100.0
Women elected as panchayat member	00.0	100.0
Women earning for her family requirement	100.0	00.0
Women taking decision in her family and agriculture related activities	100.0	00.0

The food intake in terms of kilo calorie (kcal) per person per day was calculated and presented in the Table 9 and Figure 7. More quantity of cereals is consumed by sample farmers which accounted for 1743.6.kcal per person. The other important food items consumed was pulses 339.2 kcal followed by cooking oil 302.2 kcal, milk 120.7 kcal, vegetables 54.7 kcal, egg 251.8 kcal and meat 39.7 kcal. In the sampled households,

farmers were consuming less (2851.5 kcal) than NIN- recommended food requirement (2250 kcal).

Table 9: Per capita daily consumption of food among the sample households in Nirgudi-1 Micro-watershed

Particulars	NIN recommendation (gram/ per day/ person)	Present level of consumption (gram/ per day/ person)	Kilo Calories /day/person
Cereals	396	512.8	1743.6
Pulses	43	98.9	339.2
Milk	200	185.6	120.7
Vegetables	143	228.1	54.7
Cooking Oil	31	53.0	302.2
Egg	0.5	167.9	251.8
Meat	14.2	26.2	39.4
Total	827.7	1272.5	2851.5
Threshold of NIN recommendation		827 gram*	2250 Kcal*
% Below NIN		30.0	60.0
% Above NIN		70.0	40.0

Note: * day/person




Figure 7: Per capita daily consumption of food among the sample households in Nirgudi-1 Microwatershed

Annual income of the sample HHs: The average annual household income is around Rs 35394. Major source of income to the farmers in the study area is from Livestock income (Rs 7590) followed by crop production (Rs. 27803). The monthly per capita income is Rs. 1583, which is more than the threshold monthly income of Rs 641 for considering above poverty line. Due to the fact that erratic rainfall and shortage of water, farmers are diverting from crop production activities to enable the household for a

comfortable livelihood. The incomes from the other aforesaid sources are very meagre (Table 10).

Table 10: Annual average income of HHs from various sources in Nirgudi-1 Micro-watershed

Particulars	Income *
Nonfarm income (Rs)	0 (0)
Livestock income (Rs)	7590 (60)
Crop Production (Rs)	27803 (100)
Total Annual Income (Rs)	35394
Average monthly per capita income (Rs)	641
Threshold for Poverty level (Rs 975 per month/person)	
% of households below poverty line	40.0
% of households above poverty line	60.0

* Figure in the parenthesis indicates % of Households




Figure 8: Average annual expenditure of sample HHs in Nirgudi-1 Micro-watershed

Table 11: Average annual expenditure of sample HHs in Nirgudi-1 Microwatershed

Particulars	Value in Rupees	Per cent
Food	48552	39.4
Education	31700	25.7
Clothing	11900	9.7
Social functions	17500	14.2
Health	13500	11.0
Total Expenditure (Rs/year)	123152	100.0
Monthly per capita expenditure (Rs)	2231	

The average annual expenditure of farm households indicated that farmers in the study area spend highest on food (Rs.48552) followed by education, clothing, social function and health. Now a day's education is most important among all of us. In today's competitive world, education is a necessity for man after food, clothing, and shelter. It is the only fundamental way by which a desired change in the society can happen. The average per capita monthly expenditure is around Rs. 2231 and about 40.0 per cent of farm households are below poverty line and 60.0 per of farm households are above poverty line (Table 11 and Figure 8).

Land holding: Total area cultivated by them is 17.3 ha. The average land holding of sample HHs is 1.7 ha. Large number of sample HHs (70 %) belong to small size group with an average holding size of 1.0 ha followed by medium farmers (20 %) with an average holding size of 2.0 ha and a large farmer (20 %) with a average land holding size of 4.1 ha (Table 12).

Table 12: Distribution of land holding among the sample households HHs in Nirgudi-1 Microwatershed

Particulars	Units	Values
Small farmers		
Total Land	ha	7.16
Sample size	Per cent	70
Average land holding	ha	1.0
Medium farmers		
Total Land	ha	2.0
Sample size	Per cent	10
Average land holding	ha	2.0
Large farmers		
Total Land	ha	8.2
Sample size	Per cent	20
Average land holding	ha	4.1
Grand Total		
Total Land	ha	17.3
Sample size	Per cent	100
Average land holding	ha	1.7

Land use: The total land holding in the Nirgudi 1microwatershed is 17.4 ha (Table 13). Of which 48.2 per cent is dry land and 51.8 per cent is irrigated land. The average land holding per household is worked out to be 1.74 ha.

Table 13: Land use among samples households in Nirgudi-1 Microwatershed

Particulars	Per cent	Area in ha
Irrigated land	51.8	9.0
Dry land	48.2	8.4
Fallow Land	0.0	0.0
Total land holding	100.0	17.4
Average land holding	1.74	

In the micro-watershed, the prevalent present land uses under perennial plants are neem trees (73.3 %) followed by banyan tree (6.7 %), mango tree (6.7%), coconut (3.3%) and rosewood (10.0 %) (Table 14).

Table 14: Number of trees/plants covered in sample farm households in Nirgudi-1 Microwatershed

Particulars	Number of Plants/trees	Per cent
Banyan tree(Alada)	2	6.7
Coconut	1	3.3
Mango	2	6.7
Neem trees	22	73.3
Rosewood	3	10.0
Grand Total	30	100.0




Figure 9: Present cropping pattern in Nirgudi-1 Microwatershed

Table 15: Present cropping pattern and cropping intensity in Nirgudi-1 Microwatershed(% to grand total)

Crops	Kharif	Rabi	Grand Total
Black gram	5.0	0.00	5.0
Red gram	72.9	0.0	73.0
Sorghum	4.8	5.3	10.1
Sunflower	12.0	0.0	12.0
Grand Total	94.7	5.3	100.0
Cropping intensity (%)	105		

The land use decisions are usually based on experience of farmers, tradition, expected profit, personal preferences, resources and social requirements. The present dominant crops grown in dry lands in the study area were by red gram (72.9 %) followed by sunflower (12.0 %), sorghum (4.8 %) and black gram (5.0 %) which are taken during kharif and sorghum (5.3 %) during rabi season respectively. The cropping intensity was 105 percent (Table 15 and Figure 9)

Economic land evaluation

The main purpose to characterise the socio-economic systems in the watershed is to identify the existing production constraints and propose the potential/alternate options for agro-technology transfer and for bridging the adoption and yield gap.

In Nirgudi 1 microwatershed, 4 soil series are identified and mapped (Table 16). The distribution of major soil series are Margutti covering an area around 304.38 ha (37.6 %) followed by Novinahala 276.3 ha (34.1%), Mahagaon 160.3 ha (19.8%), Dinsi 36.7 ha (4.5 %) and Bhimanahalli 14.1 ha (1.7%).

Table 16: Distribution of soil series in Nirgudi-1 Micro-watershed

Sl. No	Soil series	Description	Area in ha (%)
1	MGT mB2g 1	Very shallow, black gravelly clay soils developed from weathered basalt on very gently sloping uplands; clay surface on 1-3 % slope, moderately eroded, slightly gravelly, 15-35 per cent gravels.	45.36 (5.61)
	MGT mB2g 2	Very shallow, black gravelly clay soils developed from weathered basalt on very gently sloping uplands; clay surface on 1-3 % slope, moderately eroded, moderately gravelly, 35-60 per cent gravels.	52.01 (6.43)
	MGT mC2g 1	Very shallow, black gravelly clay soils developed from weathered basalt on gently sloping uplands; clay surface on 3-5 % slope, moderately eroded, slightly gravelly, 15-35 per cent gravels.	57.37 (7.10)
	MGT mC2g 2	Very shallow, black gravelly clay soils developed from weathered basalt on gently sloping uplands; clay surface on 3-5 % slope, moderately eroded, moderately gravelly, 35-60 per cent gravels.	26.17 (3.24)
	MGT mC3g 2	Very shallow, black gravelly clay soils developed from weathered basalt on gently sloping uplands; clay surface on 3-5 % slope, severely eroded, moderately gravelly, 35-60 per cent gravels.	89.60 (11.08)
	MGT mD3g 2	Very shallow, black gravelly clay soils developed from weathered basalt on moderately sloping uplands; clay surface on 5-10 % slope, severely eroded, moderately gravelly, 35-60 per cent gravels.	33.87 (4.19)
2	NHA mB1g 1	Shallow, black clayey soils developed from weathered basalt on very gently sloping uplands; clay surface on 1-3% slope, slightly eroded, slightly gravelly, 15-35 per cent gravels.	4.79 (0.59)
	NHA mB2	Shallow, black clayey soils developed from weathered basalt on very gently sloping uplands; clay surface on 1-3% slope, moderately eroded	114.38 (14.15)
	NHA mB2g 1	Shallow, black clayey soils developed from weathered basalt on very gently sloping uplands; clay surface on 1-3% slope, moderately eroded, slightly gravelly, 15-35 per cent gravels.	133.98 (16.57)
	NHA mB2g	Shallow, black clayey soils developed from weathered basalt on very gently sloping uplands; clay surface on 1-3% slope, moderately	10.91 (1.35)

	2	eroded, moderately gravelly, 35-60 per cent gravels.	
	NHA mC2g 1	Shallow, black clayey soils developed from weathered basalt on gently sloping uplands; clay surface on 3-5% slope, moderately eroded, slightly gravelly, 15-35 per cent gravels.	10.64 (1.32)
	NHA mC2g 3	Shallow, black clayey soils developed from weathered basalt on gently sloping uplands; clay surface on 3-5% slope, moderately eroded, highly gravelly, more than 60 per cent gravels.	1.60 (0.20)
	BHI mB2g 1	Shallow, black clay soils developed from weathered basalt on very gently sloping uplands; clay surface on 1-3% slope, moderately eroded, slightly gravelly, 15-35 per cent gravels.	14.41 (1.78)
	DSI mB2	Moderately shallow, black clayey soils developed from weathered basalt on very gently sloping uplands; clay surface on 1-3% slope, moderately eroded.	36.73 (4.54)
3	MAN mA1	Very deep, black clayey soils developed from weathered basalt on nearly level uplands; clay surface on 0-1% slope, slightly eroded	160.33 (19.83)
Water body			16.33 (2.02)

Present cropping pattern on different soil series are given in Table 17. Crops grown on Bhimanahalli soils are red gram. Redgram and sorghum are growing on Mahagaon soils. Black gram and red gram growing on Margutti soils. Red gram, sorghum and sunflower on Novinahala soils can grow.

Table 17: Cropping pattern on major soil series in Nirgudi-1 micro-watershed (Area in per cent)

Soil Series	Soil Depth	Crops	Dry		Grand Total
			Kharif	Rabi	
MGT	Very shallow (<25 cm)	Blackgram	18.5	0.0	18.5
		Redgram	81.5	0.0	81.5
BHI	Shallow (25-50 cm)	Redgram	100	0.0	100
NHA	Shallow (25-50 cm)	Redgram	67.0	0.0	67.0
		Sorghum	0.0	10.1	10.1
		Sunflower	22.9	0.0	22.9
MAN	Very deep (>150 cm)	Redgram	60.6	0.0	60.6
		Sorghum	39.4	0.0	39.4

Land is used for agricultural use for growing cereals, pulse, oilseeds and commercial crops. The soil/ land potential are measures in terms of physical yield and net income. The alternative land use options for each micro-watershed are given below (Table 18).

Table 18: Alternative land use options for different size group of farmers (Benefit Cost Ratio) in Nirgudi-1 Micro-watershed.

Soil Series	Small Farmers	Medium Farmers	Large Farmers
MGT	Black gram(1.52)		Red gram 2.04
NHA	Red gram (1.51) & Sorghum (1.30)	Sunflower (2.19)	Red gram (1.87)
BHI	Red gram (1.73)		
MAN	Red gram (2.14) & Sorghum (1.19)		

The productivity of different crops grown in Nirgudi 1 micro-watershed under potential yield of the crops is given in Table 19.

The data on cost of cultivation and benefit cost ratio (BCR) of different crops is given in Table 19. The total cost of cultivation in study area for red gram ranges between Rs. 26179/ha in NHA soil (with BCR of 1.63) and Rs 16343/ha in MGT soil (with BCR of 2.04), sorghum range between Rs 19584/ha in MAN soil (with of 1.19) and Rs. 14596/ha in NHA soil (with BCR of 1.63), black gram cost of cultivation in MGT soil is Rs. 22890/ha (with BCR of 1.52), sunflower cost of cultivation in NHA soil is Rs.18025/ha (with BCR of 2.19).

Table 19: Economic land evaluation and bridging yield gap for different crops in Nirgudi-1 micro-watershed

Particulars	MGT (<25 cm)		NHA (25-50 cm)			BHI (25-50 cm)	MAN (>150 cm)	
	Black gram	Red gram	Red gram	Sorghum	Sunflower	Red gram	Red gram	Sorghum
Total cost (Rs/ha)	22890	16343	26179	14596	18025	19303	25271	19584
Gross Return (Rs/ha)	34819	33312	41251	18918	39520	33345	54131	23280
Net returns (Rs/ha)	11930	16968	15072	4322	21495	14042	28861	3696
BCR	1.52	2.04	1.63	1.30	2.19	1.73	2.14	1.19
Farmers Practices (FP)								
FYM (t/ha)	2.2	1.5	1.9	1.1	1.0	1.3	1.6	1.3
Nitrogen (kg/ha)	70.5	107.4	69.3	72.7	91.5	94.4	77.9	91.3
Phosphorus (kg/ha)	50.7	68.9	48.5	52.3	57.5	57.5	56.0	86.3
Potash (kg/ha)	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Grain (Qtl/ha)	8.8	7.5	8.6	9.1	10.0	7.5	12.2	11.3
Price of Yield (Rs/Qtl)	4000	4500	4833	2000	4000	4500	4500	2000
Soil test based fertilizer Recommendation (STBR)								
FYM (t/ha)	7.4	7.4	7.4	7.4	6.6	7.4	7.4	7.4
Nitrogen (kg/ha)	13.9	18.5	24.7	81.5	55.2	24.7	24.7	81.5
Phosphorus (kg/ha)	46.3	61.8	61.8	71.0	74.1	61.8	61.8	71.0
Potash (kg/ha)	18.5	24.7	20.6	39.5	37.1	18.5	18.5	29.6
Grain (Qtl/ha)	9.9	12.4	12.4	28.4	16.5	12.4	12.4	28.4
% of Adoption/yield gap (STBR-FP) / (STBR)								
FYM (%)	70.3	79.8	73.7	84.7	84.8	83.1	78.1	83.1
Nitrogen (%)	-407.3	-479.7	-180.7	10.8	-65.9	-282.1	-215.5	-11.9
Phosphorus (%)	-9.4	-11.6	21.5	26.4	22.4	6.9	9.3	-21.5
Potash (%)	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Grain (%)	10.8	39.3	30.4	68.0	39.3	39.3	1.4	60.4
Value of yield and Fertilizer (Rs)								
Additional Cost (Rs/ha)	4707	5023	5925	7994	6622	5881	5771	5965
Additional Benefits (Rs/ha)	4278	21859	18153	38628	25867	21825	786	34310
Net change Income (Rs/ha)	-430	16836	12229	30634	19245	15944	-4985	28345

The data on FYM, Nitrogen, Phosphorus and Potash application by the farmers to different crops and recommended FYM for different crops is given in Table 19. There is a huge gap between FYM application by farmers and recommended FYM in all the crops

across the soils. There is a larger yield gap in crops grown across different soil series. Adequate knowledge about recommended package of practices is the pre-requisite for their use in cultivation of crops. It is a fact that, recommended practices are major contributing factors to yield. Inadequate knowledge about recommended practices leads to their improper adoption. Strengthening of extension services by concerned agency is required to increase adoption of recommended cultivation practices and ultimately reducing the gap. By adopting soil-test fertiliser recommendation, there is scope to increase yield and income to a maximum of Rs 30634 in sorghum and a minimum of Rs 12229 in redgram cultivation.

Economic valuation of Ecosystem Services (ES) was aimed at combining use and non-use values to determine Total Economic Value (TEV) of ES. Ecosystem Services (ES) were valued based on their annual flow or utilization in common monetary units, Rs/year. The valuation of ES was based on market price in 2017 or market cost approaches whichever is applicable and in other cases on value or benefit transfer from previous valuation studies.




Figure 10: Estimation of onsite cost of soil erosion in Nirgudi-1 micro-watershed

The onsite cost of different soil nutrients lost due to soil erosion is given in Table 20 and Figure 10. The average value of soil nutrient loss is around Rs 1505 per ha/year. The total cost of annual soil nutrients is around Rs 1192065 per year for the total area of 808.47 ha.

Table 20: Estimation of onsite cost of soil erosion in Nirgudi-1 Microwatershed

Particulars	Quantity(kg)		Value (Rs)	
	Per ha	Total	Per ha	Total
Organic matter	199.40	157927	1256.24	994942
Phosphorus	0.10	76	4.23	3351
Potash	3.50	2775	70.07	55496
Iron	0.19	149	9.05	7166
Manganese	0.42	335	116.20	92029
Copper	0.07	55	38.93	30829
Zinc	0.01	8	0.40	316
Sulphur	0.24	192	9.71	7690
Boron	0.01	6	0.31	244
Total	203.94	161524	1505.13	1192065




Figure 11: Ecosystem services of food grain production in Nirgudi-1 Micro-watershed

The average value of ecosystem service for food grain production is around Rs. 13470 ha/year (Table 21 and Figure 11). Per hectare food grain production services is maximum in sunflower (Rs.21495) followed by redgram (Rs.17447), black gram (Rs.11930) and sorghum (Rs. 3007).

Table 21: Ecosystem services of food grain production in Nirgudi-1 Micro-watershed

Production items	Crops	Area in ha	Yield (Qtl/ha)	Price (Rs/Qtl)	Gross Returns (Rs/ha)	Cost of Cultivation (Rs/ha)	Net Returns (Rs/ha)
Cereals	Sorghum	1.7	10	2000	20097	17090	3007
Pulses	Black gram	0.9	9	4000	34819	22890	11930
	Red gram	12.8	9	4667	40689	23242	17447
Oil seeds	Sunflower	2	10	4000	39520	18025	21495
Average value		17.4	9.5	3667	33781	20312	13470

The average value of ecosystem service for fodder production is around Rs.1002 /ha/year (Table 22) in sorghum (Rs.1002).

Table 22: Ecosystem services of fodder production in Nirgudi 1 Micro watershed

Production items	Crops	Area in ha	Yield (Qtl/ha)	Price (Rs/Qtl)	Net Returns (Rs/ha)
Cereals	Sorghum	1.7	1.2	850	1002




Figure 12: Ecosystem services of water supply in Nirgudi-1 Micro-watershed

The water demand for production of different crops was worked out in arriving at the ecosystem services of water support to crop growth. The data on water requirement for producing one quintal of grain is considered for estimating the total value of water required for crop production. Per hectare value of water used and value of water was maximum (Table 23 and Figure 12) in black gram (Rs 60116) followed by red gram (Rs 47467), sunflower (Rs 33256) and sorghum (Rs 30628).

Table 23: Ecosystem services of water supply in Nirgudi-1 Microwatershed

Crops	Yield (Qtl/ha)	Virtual water (cubic meter) per ha	Value of Water (Rs/ha)	Water consumption (Cubic meters/Qtl)
Black gram	8.7	6012	60116	691
Red gram	8.7	4747	47467	544
Sorghum	10	3063	30628	305
Sunflower	9.9	3326	33256	337
Average value	37.3	4287	42866	469

The main farming constraints in Nirgudi 1 micro-watershed to be found are less rainfall, non availability fertilizers, high crop pests & diseases, animal pests & diseases, damage of crops by wild animals and non availability of plant protection chemicals.

Majority of farmers depend up on money lender of the sources of loan for purpose of crop production. Farmers to sell the agriculture produce through village market and the farmers getting the agriculture related information on newspaper and television. Farmers reported that they are not getting timely support/extension services from the concerned development department (Table 24).

Table 24: Farming constraints related land resources of sample households in Nirgudi-1 Microwatershed

Sl.No	Particulars	Per cent
1	Less Rainfall	100.0
2	Non availability Fertilizers	11.1
3	High Crop Pests & Diseases	22.2
4	Animal Pests & Diseases	33.3
5	Damage of crops by Wild Animals	100.0
6	Non availability of Plant Protection Chemicals	100.0
7	Source of loan	
	Money Leander	100.0
8	Market for selling	
	Village market	100.0
9	Sources of Agri-Technology information	
	Newspaper	100.0

The findings of the study would be very much useful to the planners and policy makers of the study area to identify the irrationality in the existing production pattern and to suggest appropriate production plans for efficient utilization of their scarce resources resulting in increased net farm incomes and employment. The study also throws light on future potentialities of increasing net farm income and employment under different situations viz., with existing and recommended technology.