



हर कदम, हर डगर  
किसानों का हमसफर  
भारतीय कृषि अनुसंधान परिषद

*AgriSearch with a human touch*

**LAND RESOURCE INVENTORY AND SOCIO-ECONOMIC STATUS OF  
FARM HOUSEHOLDS FOR WATERSHED PLANNING AND  
DEVELOPMENT**

**NIRGUDI WEST (4D5C2B1b) MICROWATERSHED**

**Aland Taluk, Gulbarga District, Karnataka**

**Karnataka Watershed Development Project – II**

**SUJALA – III**

**World Bank funded Project**



**ICAR – NATIONAL BUREAU OF SOIL SURVEY AND LAND USE PLANNING**



ICAR - NBSS & LUP



**WATERSHED DEVELOPMENT DEPARTMENT  
GOVT. OF KARNATAKA, BANGALORE**



## **About ICAR - NBSS&LUP**

The ICAR-National Bureau of Soil Survey and Land Use Planning (ICAR-NBSS&LUP), Nagpur, a premier Institute of the Indian Council of Agricultural Research (ICAR), was set up during 1976 with the objective to prepare soil resource maps at national, state and district levels and to provide research inputs in soil resource mapping and its applications, land evaluation, land use planning, land resource management, and database management using GIS for optimising land use on different kinds of soils in the country.

The Bureau has been engaged in carrying out soil resource survey, agro-ecological and soil degradation mapping at the country, state and district levels for qualitative assessment and monitoring the soil health towards viable land use planning. The research activities have resulted in identifying the soil potentials and problems, and the various applications of the soil surveys with the ultimate objective of sustainable agricultural development. The Bureau has the mandate to correlate and classify soils of the country and maintain a National Register of all the established soil series. The Institute is also imparting in-service training to staff of the soil survey agencies in the area of soil survey, land evaluation and soil survey interpretations for land use planning. The Bureau in collaboration with Panjabrao Krishi Vidyapeeth, Akola is running post-graduate teaching and research programme in land resource management, leading to M.Sc. and Ph.D. degrees.

**Citation:** Rajendra Hegde, Ramesh Kumar, S.C., K.V. Niranjana, S. Srinivas, M.Lalitha, B.A. Dhanorkar, R.S. Reddy and S.K. Singh (2019). "Land Resource Inventory and Socio-Economic Status of Farm Households for Watershed Planning and Development of Nirgudi west (4D5C2B1b) Microwatershed, Aland Taluk, Gulbarga District, Karnataka", ICAR-NBSS&LUP Sujala MWS Publ.12, ICAR – NBSS & LUP, RC, Bangalore. p.83 & 28.

### **TO OBTAIN COPIES,**

**Please write to:**

**Director, ICAR - NBSS & LUP,**

Amaravati Road, NAGPUR - 440 033, India

Phone : (0712) 2500386, 2500664, 2500545 (O)

Telefax : 0712-2522534

E-Mail : director@nbsslup.ernet.in

Website URL : nbsslup.in

Or

**Head, Regional Centre, ICAR - NBSS&LUP, Hebbal, Bangalore - 560 024**

Phone : (080) 23412242, 23510350 (O)

Telefax : 080-23510350

E-Mail : nbssrcb@gmail.com



**LAND RESOURCE INVENTORY AND SOCIO-ECONOMIC  
STATUS OF FARM HOUSEHOLDS FOR WATERSHED  
PLANNING AND DEVELOPMENT**

**NIRGUDI WEST (4D5C2B1b) MICROWATERSHED**

**Aland Taluk, Gulbarga District, Karnataka**

**Karnataka Watershed Development Project – II**

**Sujala-III**

**World Bank funded Project**



**ICAR – NATIONAL BUREAU OF SOIL SURVEY AND LAND USE  
PLANNING**



**WATERSHED DEVELOPMENT DEPARTMENT, GOVT. OF  
KARNATAKA, BANGALORE**





## PREFACE

In Karnataka, as in other Indian States, the livelihoods of rural people are intertwined with farming pursuits. The challenges in agriculture are seriously threatening the livelihood of a large number of farmers as they have been practicing farming in contextual factors beyond their control. Climatic factors are the most important ones and have become much more significant in recent times due to rapid climate changes induced by intensive anthropogenic activities affecting our ecosystem in multiple ways. Climate change has become the reality, it is happening and efforts to evolve and demonstrate climate resilient technologies have become essential. Due to the already over stressed scenario of agrarian sector, the climate change is resulting in manifold increase in the complexities, pushing the rural mass to face more and more unpredictable situations. The rising temperatures and unpredictable rainfall patterns are going to test seriously the informed decisions farmers have to make in order to survive in farming and sustain their livelihood.

It is generally recognized that impacts of climate change shall not be uniform across the globe. It is said that impact of climate change is more severe in South Asia. Based on the analysis of meteorological data, it is predicted that in India, there will be upward trend in mean temperature, downward trend in relative humidity, annual rainfall and number of wet days in a year. Also, in general, phenomena like erratic monsoon, spread of tropical diseases, rise in sea levels, changes in availability of fresh water, frequent floods, droughts, heat waves, storms and hurricanes are predicted. Each one of these adverse situations are already being experienced in various parts of India and also at the global level. Decline in agricultural productivity of small and marginal farmers becoming more vulnerable is already witnessed.

In Karnataka, more than 60 per cent of the population live in rural areas and depend on agriculture and allied activities for their livelihood. Though the state has achieved significant progress in increasing the yield of many crops, there is tremendous pressure on the land resources due to the growing and competing demands of various land uses. This is reflected in the alarming rate of land degradation observed. Already more than 50 per cent of the area is affected by various forms of degradation. If this trend continues, the sustainability of the fragile ecosystem will be badly affected. The adverse effects of change in the climatic factors are putting additional stress on the land resources and the farmers dependent on this.

The natural resources (land, water and vegetation) of the state need adequate and constant care and management, backed by site-specific technological interventions and investments particularly by the government. Detailed database pertaining to the nature of the land resources, their constraints, inherent potentials and suitability for various land based rural enterprises, crops and other uses is a prerequisite for preparing location-specific action plans, which are in tune with the inherent capability of the resources. Any effort to evolve climate resilient technologies has to be based on the baseline scientific database. Then only one can expect effective implementation of climate resilient technologies, monitor the progress, make essential review of the strategy, and finally evaluate the effectiveness of the implemented programs. The information available at

present on the land resources of the state are of general nature and useful only for general purpose planning. Since the need of the hour is to have site-specific information suitable for farm level planning and detailed characterization and delineation of the existing land resources of an area into similar management units is the only option.

ICAR-NBSS&LUP, Regional Centre, Bangalore has taken up a project sponsored by the Karnataka Watershed Development Project-II, (Sujala-III), Government of Karnataka funded by the World Bank under Component -1 Land Resource Inventory. This study was taken up to demonstrate the utility of such a database in reviewing, monitoring and evaluating all the land based watershed development programs on a scientific footing. To meet the requirements of various land use planners at grassroots level, the present study on “Land Resource Inventory and Socio-Economic Status of Farm Households for Watershed Planning and Development of Nirgudi West Microwatershed, Aland Taluk, Kalaburgi District, Karnataka” for integrated development was taken up in collaboration with the State Agricultural Universities, IISC, KSRSAC, KSNDMC as Consortia partners. The project provides detailed land resource information at cadastral level (1:7920 scale) for all the plots and socio-economic status of farm households covering thirty per cent farmers randomly selected representing landed and landless class of farmers in the microwatershed. The project report with the accompanying maps for the microwatershed will provide required detailed database for evolving effective land use plan, alternative land use options and conservation plans for the planners, administrators, agricultural extension personnel, KVK officials, developmental departments and other land users to manage the land resources in a sustainable manner.

It is hoped that this database will be useful to the planners, administrators and developmental agencies working in the area in not only for formulating location specific developmental schemes but also for their effective monitoring at the village/watershed level.

Nagpur

Date:

**S.K. SINGH**

Director, ICAR - NBSS&LUP, Nagpur

## Contributors

<b>Dr. RajendraHegde</b> Principal Scientist, Head & Project Leader, Sujala-III Project ICAR-NBSS&LUP, Regional Centre Bangalore	<b>Dr. S.K.Singh</b> Director, ICAR-NBSS&LUP Coordinator, Sujala-III Project Nagpur
<b>Soil Survey, Mapping &amp; Report Preparation</b>	
Dr. B.A. Dhanorkar	Sh. R.S. Reddy
Dr. K.V. Niranjana	Sh. Nagendra, B.R.
	Smt. Chaitra, S.P.
<b>Field Work</b>	
Sh. C.BacheGowda	Sh. Mahesha, D.B.
Sh. Somashekar	Sh. Ashok S. Sindagi
Sh. VenkataGiriyappa	Sh. Veerabhadrapa
Sh. M. Jayaramaiah	Sh. Anand
Sh. Paramesha, K.	Sh. Arun N Kambar
	Sh. Shankarappa, K.
	Sh. Kamalesh K. Avate
	Sh. Sharan Kumar Huppar
	Sh. Yogesh, H.N.
	Sh. Kalaveerachari R. Kammar
<b>GIS Work</b>	
Dr. S.Srinivas	Sh. A.G.Devendra Prasad
Dr. M.Ramesh	Sh. Prakashanaik, M.K.
Sh. D.H.Venkatesh	Sh. AbhijithSastry, N.S.
Smt.K.Sujatha	Sh. Sudip Kumar Suklabaidya
Smt. K.V.Archana	Sh. Mahamad Ali, M.
Sh. N.Maddileti	Sh. Avinash, K.N.
	Sh. Amar Suputhra, S
	Sh. Anudeep, Y.
	Sh. Deepak, M.J.
	Smt. K.Karunya Lakshmi
	Ms. Seema, K.V.
	Ms. A. Rajab Nisha

<b>Laboratory Analysis</b>	
Dr. K.M.Nair	Dr. H.R. Savitha
Smt. Arti Koyal	Ms. Steffi Peter
Smt. Parvathy, S.	Ms. Thara, V.R
	Ms. Roopa, G.
	Ms. Swati, H.
	Sh. Shantaveera Swami
	Ms. Shwetha, N.K.
	Smt. Ishrat Haji
	Ms. P. PavanKumari
	Ms. Padmaja
	Ms. Veena, M.
<b>Soil &amp; Water Conservation</b>	
Sh. Sunil P. Maske	
<b>Watershed Development Department, GoK, Bangalore</b>	
Sh. Rajeev Ranjan IFS Project Director & Commissioner, WDD	Dr. A. Natarajan NRM Consultant, Sujala-III Project
Dr. S.D. Pathak IFS Executive Director & Chief Conservator of Forests, WDD	



# **PART-A**

## **LAND RESOURCE INVENTORY**



## Contents

Preface		
Contributors		
Executive Summary		
Chapter 1	Introduction	1
Chapter 2	Geographical Setting	3
2.1	Location and Extent	3
2.2	Geology	4
2.3	Physiography	4
2.4	Drainage	4
2.5	Climate	5
2.6	Natural Vegetation	6
2.7	Land Utilization	6
Chapter 3	Survey Methodology	9
3.1	Base maps	9
3.2	Field Investigation	11
3.3	Laboratory Characterization	12
3.4	Finalization of Soil Map	12
Chapter 4	The Soils	19
4.1	Soils of Basalt Landscape	19
Chapter 5	Interpretation for Land Resource Management	25
5.1	Land Capability Classification	25
5.2	Soil Depth	27
5.3	Surface Soil Texture	28
5.4	Soil Gravelliness	29
5.5	Available Water Capacity	30
5.6	Soil Slope	31
5.7	Soil Erosion	32
Chapter 6	Fertility Status	35
6.1	Soil Reaction (pH)	35
6.2	Electrical Conductivity (EC)	35
6.3	Organic Carbon (OC)	35
6.4	Available Phosphorus	37
6.5	Available Potassium	37
6.6	Available Sulphur	37
6.7	Available Boron	37
6.8	Available Iron	40
6.9	Available Manganese	40
6.10	Available Copper	40
6.11	Available Zinc	40

Chapter 7	Land Suitability for Major Crops	43
7.1	Land suitability for Sorghum	43
7.2	Land suitability for Maize	46
7.3	Land suitability for Red gram	47
7.4	Land suitability for Sunflower	49
7.5	Land suitability for Cotton	50
7.6	Land suitability for Sugarcane	52
7.7	Land suitability for Soybean	53
7.8	Land suitability for Guava	54
7.9	Land suitability for Mango	56
7.10	Land suitability for Sapota	57
7.11	Land suitability for Jackfruit	59
7.12	Land suitability for Jamun	60
7.13	Land Suitability for Musambi	61
7.14	Land Suitability for Lime	62
7.15	Land Suitability for Cashew	64
7.16	Land Suitability for Custard Apple	64
7.17	Land Suitability for Amla	65
7.18	Land Suitability for Tamarind	66
7.19	Land Management Units	67
7.20	Proposed Crop Plan	69
Chapter 8	Soil Health Management	71
Chapter 9	Soil and Water conservation Treatment Plan	75
9.1	Treatment Plan	76
9.2	Recommended Soil and Water Conservation measures	79
9.3	Greening of microwatershed	80
	References	83
	Appendix I	I-X
	Appendix II	XI-XVIII
	Appendix III	XIX-XXII

## LIST OF TABLES

2.1	Mean Monthly Rainfall, PET, 1/2 PET at Aland Taluk, Kalburgi District	5
2.2	Land Utilization in Aland Taluk	7
3.1	Differentiating Characteristics used for Identifying Soil Series	12
3.2	Soil Legend	17
7.1	Soil-Site Characteristics of Nirgudi West Microwatershed	44
7.2	Crop suitability criteria for Sorghum	45
7.3	Crop suitability criteria for Maize	46
7.4	Crop suitability criteria for Red gram	48
7.5	Crop suitability criteria for Sunflower	49
7.6	Crop suitability criteria for Cotton	51
7.7	Crop suitability criteria for Sugarcane	52
7.8	Crop suitability criteria for Guava	55
7.9	Crop suitability criteria for Mango	56
7.10	Crop suitability criteria for Sapota	58
7.11	Crop suitability criteria for Lime	63
7.12	Proposed Crop Plan for Nirgudi West Microwatershed	69



## LIST OF FIGURES

2.1	Location map of Nirgudi West Microwatershed	3
2.2	Rock formations in Nirgudi West Microwatershed	4
2.3	Rainfall distribution in Aland Taluk, Kalburgi District	6
2.4	Current Land use map of Nirgudi West Microwatershed	7
2.5	Location of Wells in Nirgudi West Microwatershed	8
3.1	Scanned and Digitized Cadastral map of Nirgudi West Microwatershed	10
3.2	Satellite image of Nirgudi West Microwatershed	10
3.3	Cadastral map overlaid on IRS PAN+LISS IV merged imagery of Nirgudi West Microwatershed	11
3.4	Soil phase or management units of Nirgudi West Microwatershed	15
5.1	Land Capability Classification of Nirgudi West Microwatershed	27
5.2	Soil Depth map of Nirgudi West Microwatershed	28
5.3	Surface Soil Texture map of Nirgudi West Microwatershed	29
5.4	Soil Gravelliness map of Nirgudi West Microwatershed	30
5.5	Soil Available Water Capacity map of Nirgudi West Microwatershed	31
5.6	Soil Slope map of Nirgudi West Microwatershed	32
5.7	Soil Erosion map of Nirgudi West Microwatershed	33
6.1	Soil Reaction (pH) map of Nirgudi West Microwatershed	36
6.2	Electrical Conductivity (EC) map of Nirgudi West Microwatershed	36
6.3	Soil Organic Carbon (OC) map of Nirgudi West Microwatershed	38
6.4	Soil Available Phosphorus map of Nirgudi West Microwatershed	38
6.5	Soil Available Potassium map of Nirgudi West Microwatershed	39
6.6	Soil Available Sulphur map of Nirgudi West Microwatershed	39
6.7	Soil Available Boron map of Nirgudi West Microwatershed	40
6.8	Soil Available Iron map of Nirgudi West Microwatershed	41
6.9	Soil Available Manganese map of Nirgudi West Microwatershed	41
6.10	Soil Available Copper map of Nirgudi West Microwatershed	42
6.11	Soil Available Zinc map of Nirgudi West Microwatershed	42
7.1	Land Suitability map of Sorghum	45
7.2	Land Suitability map of Maize	47
7.3	Land Suitability map of Red gram	48
7.4	Land Suitability map of Sunflower	50
7.5	Land Suitability map of Cotton	51

7.6	Land Suitability map of Sugarcane	53
7.7	Land Suitability map of Soybean	54
7.8	Land Suitability map of Guava	55
7.9	Land Suitability map of Mango	57
7.10	Land Suitability map of Sapota	59
7.11	Land Suitability map of Jackfruit	60
7.12	Land Suitability map of Jamun	61
7.13	Land Suitability map of Musambi	62
7.14	Land Suitability map of Lime	63
7.15	Land Suitability map of Cashew	64
7.16	Land Suitability map of Custard Apple	65
7.17	Land Suitability map of Amla	66
7.18	Land Suitability map of Tamarind	67
7.19	Land Management Units map of Nirgudi West Microwatershed	68
9.1	Soil and Water Conservation map of Nirgudi West Microwatershed	80



## **EXECUTIVE SUMMARY**

*The land resource inventory of Nirgudi West Microwatershed was conducted using village cadastral maps and IRS satellite imagery on 1:7920 scale. The false colour composites of IRS imagery were interpreted for physiography and these physiographic delineations were used as base for mapping soils. The soils were studied in several transects and a soil map was prepared with phases of soil series as mapping units. Random checks were made all over the area outside the transects to confirm and validate the soil map unit boundaries. The soil map shows the geographic distribution and extent, characteristics, classification and use potentials of the soils in the microwatershed.*

*The present study covers an area of 504 ha in Nirgudi West microwatershed in Aland taluk, Kalaburgi district, Karnataka. The climate is semiarid and categorized as drought prone with an average annual rainfall of 786 mm of which about 595 mm is received during south-west monsoon, 116mm during north-east and the remaining 75 mm during the rest of the year. An area of about 92 per cent is covered by soils, 8 per cent by waterbodies, settlements and others. The salient findings from the land resource inventory are summarized briefly below.*

- ❖ *The soils belong to 6 soil series and 18 soil phases (management units) and 5 land management units.*
- ❖ *The length of crop growing period is about 150 days starting from the 3rd week of June to 1st week of October.*
- ❖ *From the master soil map, several interpretative and thematic maps like land capability, soil depth, surface soil texture, soil gravelliness, available water capacity, soil slope and soil erosion were generated.*
- ❖ *Soil fertility status maps for macro and micronutrients were generated based on the surface soil samples collected at every 250 m grid interval.*
- ❖ *Land suitability for growing major agricultural and horticultural crops were assessed and maps showing degree of suitability along with constraints were generated.*
- ❖ *About 80 per cent area is suitable for agriculture and 12 per cent is not suitable for agriculture but well suited for forestry, pasture, agroforestry, silvi-pasture, recreation, installation of wind mills and as habitat for wildlife.*
- ❖ *About 10 per cent of the soils are deep (100-150 cm) to moderately deep (75-100 cm), 60 per cent are moderately shallow to shallow (25-75 cm) and about 23 per cent are very shallow (<25 cm) soils.*
- ❖ *About 92 per cent of the area has clay soils at the surface.*
- ❖ *About 32 per cent of the area has non-gravelly soils (<15% gravel), 37 per cent gravelly soils (15-35% gravel) and 23 per cent very gravelly soils (35- 60% gravel).*
- ❖ *About 8 per cent of the area has soils that are very high (>200mm/m) in available water capacity, 30 per cent medium (100-150 mm/m) and about 55 per cent low (50-100 mm/m) and very low (<50mm/m).*
- ❖ *About 62 per cent of the area has nearly level (0-1%) to very gently sloping (1-3%) lands and about 30 per cent area is gently (3-5%) to moderately sloping (5-10%) lands.*
- ❖ *An area of about 16 per cent has soils that are slightly eroded (e1), 50 per cent moderately eroded (e2) and 26 per cent severely eroded (e3).*
- ❖ *An area of about 41 per cent has soils that are neutral in reaction (pH 6.5 to 7.3), about 28 per cent slightly alkaline (pH 7.3-7.8) and 24 per cent moderately alkaline (pH 7.8 to 8.4).*
- ❖ *The Electrical Conductivity (EC) of the soils are dominantly <2 dSm-1 indicating that the soils are non-saline.*
- ❖ *About 32 per cent medium (0.5-0.75%), 57 per cent high (>0.75%) and 4 per cent low (<0.5%) in organic carbon.*
- ❖ *Major area of 76 per cent has soils that are low (<23 kg/ha), 15 per cent medium (23-57 kg/ha) and only one per cent high (>57 kg/ha) in available phosphorus.*

- ❖ About 25 per cent medium (145-337 kg/ha) and 67 per cent high (>337 kg/ha) in available potassium.
- ❖ Available sulphur is low (<10 ppm) in about one per cent area and medium (10-20 ppm) in 91 per cent.
- ❖ Available boron is low (<0.5 ppm) in about 66 per cent area and 25 per cent medium (0.5-1.0 ppm).
- ❖ About 2 per cent area has soils that are deficient (<4.5 ppm) and 90 per cent sufficient (>0.6 ppm) in available iron.
- ❖ Available manganese and copper are sufficient in all the soils.
- ❖ About 33 per cent area has soils that are deficient (<0.6 ppm) and 60 per cent sufficient (>0.6 ppm) in available zinc.
- ❖ The land suitability for 18 major crops (agricultural and horticultural) grown in the microwatershed were assessed and the areas that are highly suitable (S1) and moderately suitable (S2) are given below. It is however to be noted that a given soil may be suitable for various crops but what specific crop to be grown may be decided by the farmer looking to his capacity to invest on various inputs, marketing infrastructure, price and finally the demand and supply position.

***Land suitability for various crops in the microwatershed***

Crop	Suitability Area in ha (%)		Crop	Suitability Area in ha (%)	
	Highly suitable (S1)	Moderately suitable (S2)		Highly suitable (S1)	Moderately suitable (S2)
Sorghum	24 (5)	165 (33)	Sapota	-	-
Maize	-	140 (28)	Jackfruit	-	-
Red gram	-	189 (37)	Jamun	-	40 (8)
Sunflower	24 (5)	25 (5)	Musambi	24 (5)	25 (5)
Cotton	24 (5)	165 (33)	Lime	24 (5)	25 (5)
Sugarcane	-	140 (28)	Cashew	-	-
Soybean	24(5)	165 (33)	Custard apple	24 (5)	165 (33)
Guava	-	-	Amla	24 (5)	165 (33)
Mango	-	-	Tamarind	24 (5)	16 (4)

Apart from the individual crop suitability, a proposed crop plan has been prepared for the 5 identified LMUs by considering only the highly and moderately suitable lands for different crops and cropping systems with food, fibre and horticulture crops that helps in maintaining the ecological balance in the microwatershed.

- ❖ Maintaining soil-health is vital to crop production and conserve soil and land resource base for maintaining ecological balance and to mitigate climate change. For this, several ameliorative measures have been suggested to these problematic soils like saline/alkali, highly eroded, sandy soils etc.,
- ❖ Soil and water conservation treatment plan has been prepared that would help in identifying the sites to be treated and also the type of structures required.
- ❖ As part of the greening programme, several tree species have been suggested to be planted in marginal and submarginal lands, field bunds and also in the hillocks, mounds and ridges.

## **INTRODUCTION**

Soil being a vital natural resource on whose proper use depends the life supporting systems of a country and the socioeconomic development of its people. Soils provide food, fodder, fibre and fuel for meeting the basic human and animal needs. With the ever increasing growth in human and animal population, the demand on soil for more food and fodder production is on the increase. The area available for agriculture is about 51 per cent of the total geographical area and more than 60 per cent of the people are still dependant on agriculture for their livelihood. However, the capacity of a soil to produce is limited and the limits to the production are set by its intrinsic characteristics, agro-climatic setting, and use and management. There is therefore, tremendous pressure on land and water resources, which is causing decline in soil-health and stagnation in productivity. The soils have been degrading at an estimated rate of one million hectares per year and ground water levels have been receding at an alarming rate resulting in decline in the ground water resource. Further, land degradation has emerged as a serious problem which has already affected about 38 lakh ha of cultivated area in the State. Soil erosion alone has degraded about 35 lakh ha. Almost all the uncultivated areas are facing various degrees of degradation, particularly soil erosion; salinity and alkalinity has emerged as a major problem (>3.5 lakh ha) in the irrigated areas of the State. Nutrient depletion and declining factor productivity is common in both rainfed and irrigated areas. The degradation is continuing at an alarming rate and there appears to be no systematic effort among the stakeholders to contain this process. In recent times, an aberration of weather due to climate change phenomenon has added another dimension leading to unpredictable situations to be tackled by the farmers.

In this critical juncture, the challenge before us is not only to increase the productivity per unit area which is steadily declining and showing a fatigue syndrome, but also to prevent or at least reduce the severity of degradation. If the situation is not reversed at the earliest, then the sustainability of the already fragile crop production system and the overall ecosystem will be badly affected in the state. Added to this, every year there is a significant diversion of farm lands and water resources for non-agricultural purposes. Thus, developing strategies to slow down the degradation process or reclaim the soils to normal condition and ensure sustainability of production system are the major issues today. This demands a systematic appraisal of our soil and land resources with respect to their extent, geographic distribution, characteristics, behaviour and use potential, which is very important for developing an effective land use and cropping systems for augmenting agricultural production on a sustainable basis.

The soil and land resource inventories made so far in Karnataka had limited utility because the surveys were of different types, scales and intensities carried out at different times with specific objectives. Hence, there is an urgent need to generate detailed site-specific farm level database on various land resources for all the villages/watersheds in a time

bound manner that would help to protect the valuable soil and land resources and also to stabilize the farm production. Therefore, the land resource inventory required for farm level planning is the one which investigates all the parameters which are critical for productivity viz., soils, site characteristics (slope, erosion, gravelliness and stoniness) climate, water, topography, geology, hydrology, vegetation, crops, land use pattern, animal population, socio-economic conditions, infrastructure, marketing facilities and various schemes and developmental works of the government etc. From the data collected at farm level, the specific problems and potentials of the area can be identified and highlighted, conservation measures required for the area can be planned on a scientific footing, suitability of the area for various uses can be worked out and finally viable and sustainable land use options suitable for each and every land holding can be prescribed.

The Land Resource Inventory is basically done for identifying potential and problem areas, developing sustainable land use plans, estimation of surface run off and water harvesting potential, preparation of soil and water conservation plans, land degradation/desertification etc. The Bureau is presently engaged in developing an LRI methodology using high resolution satellite remote sensing data and Digital Elevation Model (DEM) data to prepare Landscape Ecological Units (LEU) map representing agro-ecosystem as a whole. The LEU is preferred over landform as the base map for LRI. LEU is the assemblage of landform, slope and land use. An attempt has already been made to upscale the soil resource information from 1:250000 and 1:50000 scale to the LEU map in Goa and in some other states. Here, an attempt will be made later to uplink the LRI data generated under Sujala-III Project to the Landscape Ecological Units (LEUs) map. For this, the major physiographic region, *i.e.*, South Deccan Plateau will be taken as an example.

The land resource inventory aims to provide site specific database for Nirgudi West microwatershed in Aland Taluk, Kalaburgi District, Karnataka state for the Karnataka Watershed Development Department. The database was generated by using cadastral map of the village as a base along with high resolution IRS LISS IV and Cartosat-1 merged satellite imagery. Later, an attempt will be made to uplink this LRI data generated at 1:7920 scale under Sujala-III Project to the proposed Landscape Ecological Units (LEUs) map.

The study was organized and executed by the ICAR- National Bureau of Soil Survey and Land Use Planning, Regional Centre, Bangalore under Generation of Land Resource Inventory Data Base Component-1 of the Sujala-III Project funded by the World Bank.

## GEOGRAPHICAL SETTING

### 2.1 Location and Extent

The study area of Nirgudi West microwatershed (Alur subwatershed) is located in the northeastern part of Karnataka in Aland Taluk, Kalaburgi District, Karnataka State (Fig.2.1). It comprises parts of Nirgudi villages. It lies between  $17^{\circ} 37'$  and  $17^{\circ} 39'$  north latitude and  $77^{\circ} 26'$  and  $77^{\circ} 27'$  east longitude and covers an area of 505 ha. It is about 15 km south of Aland and is surrounded by Chincholi Budruk village in the western part, Padasavli village in the south, Matki village in the east and Maharashtra state in the north and northwestern part of the microwatershed.




Fig.2.1 Location map of Nirgudi West Microwatershed

## 2.2 Geology

Major rock formation observed in the microwatershed is Basalt (Fig.2.2) or Deccan Trap. The Deccan Traps cover the whole of Bidar, parts of Kalaburgi, Bijapur and Belgam districts. In all, eight lava flows have been identified in Karnataka horizontally overlying the older formations. The thickness of the individual flows averages about five meters. It is relatively uniform in petrographic character. The most common type is augite basalt. Dominant colour is grayish green and texture ranges from cryptocrystalline to glassy. The rock is often vesicular and scoriaceous filled up with secondary minerals like coloured agate, quartz, calcite and a large variety of zeolites. The Deccan Traps form an excellent building material and also used as road-metal and railway ballast.



Fig. 2.2 Basalt rock formation

## 2.3 Physiography

Physiographically, the area has been identified as basalt landscape based on geology. The area has been further subdivided into four landforms, viz; mounds/ridges, summits, side slopes and very gently sloping uplands based on slope and its relief features. The elevation ranges from 513 to 569 m. The mounds and ridges are mostly covered by rock outcrops.

## 2.4 Drainage

The area is drained by several small parallel streams that join Monia nala which further downstream joins Awarja river along its course. Though, it is not a perennial one,

during rainy season it carries large quantities of rain water. The microwatershed has only few small tanks which are not capable of storing the water that flows during the rainy season. Due to this, the ground water recharge is very much affected. This is reflected in the failure of many bore wells in the villages. If the available rain water is properly harnessed by constructing new tanks and recharge structures at appropriate places in the village, then the drinking and irrigation needs of the entire area can be easily met. The drainage network is parallel to sub parallel and dendritic.

## 2.5 Climate

The Kalaburgi district lies in the northern plains of Karnataka and falls under semiarid tract of the state and is categorized as drought prone with average annual rainfall of 743 mm (Table 2.1). Of the total rainfall, maximum of 595 mm is received during the south-west monsoon period from June to September, the north-east monsoon from October to early December contributes about 116 mm and the remaining 75 mm during the rest of the year. December is the coldest month with mean daily maximum and minimum temperatures being 29.5<sup>0</sup>C and 15<sup>0</sup> to 10<sup>0</sup>C respectively. During peak summer, temperature shoots up to 45<sup>0</sup>C. Relative humidity varies from 26 per cent in summer to 62 per cent in winter. Rainfall distribution is shown in Figure 2.3. The average Potential Evapotranspiration (PET) is 150 mm and varies from a low of 115 mm in December to 232 mm in the month of May. The PET is always higher than precipitation in all the months except August and September. Generally, the length of crop growing period (LGP) is 150 days and starts from 3<sup>rd</sup> week of June to third week of November.

Table 2.1 Mean Monthly Rainfall, PET, 1/2 PET at Aland Taluk, Kalaburgi District

Sl.No.	Months	Rainfall	PET	1/2 PET
1	January	7.50	126.80	63.40
2	February	3.40	143.90	71.95
3	March	11.30	189.90	94.95
4	April	19.40	209.80	104.90
5	May	32.70	232.20	116.10
6	June	111.00	186.40	93.20
7	July	139.20	152.80	76.40
8	August	172.40	147.60	73.80
9	September	172.30	131.70	65.85
10	October	91.30	145.50	72.75
11	November	19.30	129.80	64.90
12	December	5.80	114.80	57.40
Total		785.6	149.70	




Fig 2.3 Rainfall distribution in Aland Taluk, Kalaburgi District

## 2.6 Natural Vegetation

The natural vegetation is sparse comprising few tree species, shrubs and herbs. The mounds, ridges and boulders occupy very sizeable area which is under thin to moderately thick forest vegetation. Still, there are some remnants of the past forest cover which can be seen in patches in some ridges and hillocks in the microwatershed.

Apart from the continuing deforestation, the presence of large population of goats, sheep and other cattle in the microwatershed is causing vegetative degradation of whatever little vegetation left in the area. The uncontrolled grazing has left no time for the regeneration of the vegetative cover. This leads to the accelerated rate of erosion on the hill slopes resulting in the formation of deep gullies in the foot slopes that eventually result in the heavy siltation of tanks and reservoirs in the microwatershed.

## 2.7 Land Utilization

About 89 per cent area (Table 2.2) in Aland taluk is cultivated at present. An area of about 2 per cent is permanently under pasture, 3 per cent under current fallows and 2 per cent under non agricultural land and currently barren. Forests occupy an area of about 2 per cent and the tree cover is in a very poor state. Most of the mounds, ridges and bouldery areas have very poor vegetative cover. Major crops grown in the area are sorghum, maize, cotton, sugarcane, red gram and sapota. While carrying out land resource inventory, the land use/land cover particulars are collected from all the survey numbers and a current land use map of the microwatershed is prepared. The current land use map prepared shows the arable and non-arable lands, other land uses and different types of crops grown in the area. The current land use map of Nirgudi West microwatershed is presented in Fig 2.4.



Table 2.2 Land Utilization in Aland Taluk

Sl. No.	Agricultural land use	Area ( ha)	Per cent
1.	Total geographical area	173417	
2.	Total cultivated area	153806	88.69
3.	Area sown more than once	7910	
4.	Trees and grooves	59	0.034
5.	Forest	2854	1.64
6.	Cultivable wasteland	974	0.56
7.	Permanent Pasture land	3469	2.00
8.	Barren land	3142	1.81
9.	Non- Agriculture land	3465	1.99
10.	Current Fallows	5648	3.25




Fig.2.4 Current Land Use map of Nirgudi West Microwatershed

Simultaneously, enumeration of wells (bore wells and open wells) in the microwatershed was made and their location in different survey numbers is located on the cadastral map. Map showing the location of wells and other water bodies in the Nirgudi West microwatershed is given in Fig 2.5.




Fig.2.5 Location of Wells in Nirgudi West Microwatershed

## SURVEY METHODOLOGY

The purpose of land resource inventory is to delineate similar areas (soil series and phases), which respond or expected to respond similarly to a given level of management. This was achieved in Nirgudi West microwatershed by the detailed study of all the soil characteristics (depth, texture, colour, structure, consistence, coarse fragments, porosity, soil reaction, soil horizons etc.) and site (slope of the land, erosion, drainage, occurrence of rock fragments etc.) followed by grouping of similar areas based on soil-site characteristics into homogeneous (management units) units and showing their area extent and geographic distribution on the microwatershed cadastral map. The detailed survey at 1:7920 scale was carried out in 505 ha area. The methodology followed for carrying out land resource inventory was as per the guidelines given in Soil Survey Manual (IARI, 1971; Soil Survey Staff, 2006; Natarajan *et al.*, 2015) which is briefly described below.

### 3.1 Base Maps

The detailed survey of the land resources occurring in the microwatershed was carried out by using digitized cadastral map as a base. The cadastral map shows field boundaries with their survey numbers, location of tanks, streams and other permanent features of the area (Fig. 3.1). Apart from the cadastral map, remote sensing data products from Cartosat-1 and LISS-IV merged at the scale of 1:7920 were used in conjunction with the cadastral map to identify the rock types, the landscapes, landforms and other surface features. The imagery helped in the identification and delineation of boundaries between hills, uplands and lowlands, water bodies, forest and vegetated areas, roads, habitations and other cultural features of the area (Fig. 3.2). The cadastral map was overlaid on the satellite imagery (Fig.3.3) that helps to identify the parcel boundaries and other permanent features. Apart from cadastral maps and images, toposheets of the area (1:50,000 scale) were used for initial traversing, identification of geology and landforms, drainage features, present land use and also for selection of transects in the microwatershed.




Fig 3.1 Scanned and Digitized Cadastral map of Nirgudi West Microwatershed




Fig.3.2 Satellite Image of Nirgudi West Microwatershed




Fig.3.3 Cadastral map overlaid on IRS PAN+LISS IV merged imagery of Nirgudi West Microwatershed

### 3.2 Field Investigation

Preliminary traverse of the microwatershed was carried out with the help of cadastral map, imagery and toposheets. While traversing, landforms and physiographic units identified were checked and preliminary soil legend was prepared by studying soils at a few selected places. The field boundaries and survey numbers given on the cadastral sheet were located on the ground by following permanent features like roads, cart tracks, nallas, streams, tanks etc., and wherever changes were noticed, they were incorporated on the microwatershed cadastral map. Then, intensive traversing of each physiographic unit like hills, ridges and uplands was carried out. Based on the variability observed on the surface, transects were selected across the slope covering all the landform units in the microwatershed (Natarajan and Dipak Sarkar, 2010).

In the selected transect, soil profiles were located at closely spaced intervals to take care of any change in the land features like break in slope, erosion, gravel, stones etc. In the selected sites, profiles (vertical cut showing the soil layers from surface to the rock) were opened up to 200 cm or to the depth limited by rock or hard substratum and studied in detail for all their morphological and physical characteristics. The soil and site characteristics were recorded for all profile sites on a standard proforma as per the guidelines given in USDA Soil

Survey Manual (Soil Survey Staff, 2012). Apart from the transect study, profiles were also studied at random, almost like in a grid pattern, outside the transect areas.

Based on the soil-site characteristics, the soils were grouped into different soil series (soil series is the most homogeneous unit having similar horizons and properties and behaves similarly for a given level of management). Soil depth, texture, colour, kind of horizon and horizon sequence, amount and nature of gravel present, nature of substratum etc, were used as the major differentiating characteristics for identifying soil series occurring in the area. The differentiating characteristics used for identifying soil series are given in Table 3.1. Based on the above characteristics, 6 soil series were identified in the Nirgudi West microwatershed.

Table 3.1 Differentiating Characteristics used for Identifying Soil Series  
(Characteristics are of Series Control Section)

SOILS OF BASALT LANDSCAPE							
Sl. no	Soil Series	Depth (cm)	Colour	Text-ure	Gravel (%)	Horizon sequence	Calcar-ousness
1	Margutti (MGT)	<25	10YR3/3,4/3,5/ 7.5YR4/3	c	15-35	Ap- cr	-
2	Novinihala (NHA)	25-50	10YR3/2,3/1,4/2 7.5YR3/4	c	<15	Ap-Bw- cr/R	-
3	Bhimanahalli (BHI)	25-50	10YR3/2,3/3,3/1 7.5YR3/2,4/2	c	15-35	Ap-Bw- cr/R	-
4	Dinsi (DSI)	50-75	10YR3/2,3/3,4/3 3/2	c	<15	Ap-BA- Bss	-
5	Kamalapur (KMP)	75-100	10YR3/2,3/1	c	<15	Ap-Bw- Bss-cr	-
6	Rajanala (RNL)	100-150	10YR3/2,3/1,4/2,4/3	c	<15	Ap-Bw- Bss-cr	-

### 3.3 Laboratory Characterization

Soil samples were collected from representative master profiles for laboratory characterization by following the methods outlined in the Laboratory Manual (Sarma *et al*, 1987). Surface soil samples collected from farmer's fields (79 samples) for fertility status (major and micronutrients) at 250 m grid interval were analyzed in the laboratory (Katyal and Rattan, 2003). By linking the soil fertility data to the survey numbers through GIS, soil fertility maps for all 11 elements including pH and EC were generated for the microwatershed.

### 3.4 Finalization of Soil Maps

The area under each soil series was further separated and mapped as soil phases and their boundaries delineated on the cadastral map based on the variations observed in the

texture of the surface soil, slope, erosion, presence of gravel, stoniness etc. A soil phase is a subdivision of soil series based mostly on surface features that affect its use and management.

The soil mapping units are shown on the map (Fig.3.4) in the form of symbols. During the survey about 26 profile pits, few minipits and a few auger bores representing different landforms occurring in the microwatershed were studied. All the profile locations are indicated on the village cadastral map in the form of a triangle. In addition to the profile study, spot observations in the form of minipits, road cuts, terrace cuts etc., were studied to validate the soil boundaries on the soil map. The soil map shows the geographic distribution of 18 mapping units representing 6 soil series occurring in the microwatershed. The soil map unit (soil legend) description is presented in Table 3.2.

The soil phase map (management units) shows the distribution of 18 phases identified and mapped in the microwatershed. Each mapping unit (soil phase) delineated on the map has similar soil and site characteristics. In other words, all the farms or survey numbers included in one phase will have similar management needs and they have to be treated accordingly.

The 18 soil phases identified and mapped in the microwatershed were regrouped into 5 Land Management Units (LMU's) for the purpose of preparing a proposed crop plan for sustained development of the microwatershed. The database (soil phases) generated under LRI was utilized for identifying Land Management Units (LMUs) based on the management needs. One or more than one soil site characteristic having influence on the management have been chosen for identification and delineation of LMUs. For Nirgudi West microwatershed five soil and site characteristics, namely soil depth, soil texture, slope, erosion and gravel content have been considered for defining LMUs. The land management units are expected to behave similarly for a given level of management.








Fig 3.4 Soil Phase or Management Units map of Nirgudi West Microwatershed



**Table 3.2 Soil Legend**

Soil Map Unit No.	Soil Series	Soil Phase	Mapping Unit Description	Area in ha (%)
<b>Soils of Basalt Landscape</b>				
	MGT	Margutti soils are very shallow (<25cm), well drained, have very dark grayish brown to dark brown clayey soils occurring on nearly level to moderately sloping uplands.		<b>116.5 (23.1)</b>
1		MGTmB2g1	Clay surface, 1-3 % slope, moderate erosion, gravelly (15-35%)	9.69 (1.92)
2		MGTmC3g1	Clay surface, 3-5 % slope, severe erosion, gravelly (15-35%)	9.70 (1.92)
3		MGTmC3g2	Clay surface, 3-5 % slope, severe erosion, gravelly (35-60%)	57.95 (11.48)
4		MGTmD3g2	Clay surface, 5-10 % slope, severe erosion, very gravelly (35-60%)	39.14 (7.76)
	NHA	Novinihala soils are shallow (25-50 cm), well drained, have very dark grayish brown to dark brown clayey soils occurring on very gently sloping uplands.		<b>140.5 (27.8)</b>
5		NHAmB1	Clay surface, slope 1-3%, slight erosion	8.92 (1.77)
6		NHAmB2	Clay surface, slope 1-3%, moderate erosion	63.31 (12.55)
7		NHAmB2g1	Clay surface, slope 1-3%, moderate erosion, gravelly (15-35%)	30.51 (6.05)
8		NHAmC2g1	Clay surface, slope 3-5%, moderate erosion, gravelly (15-35%)	11.18 (2.22)
9		NHAmC3	Clay surface, slope 3-5%, severe erosion	5.17 (1.03)
10		NHAmD3g2	Clay surface, slope 5-10%, severe erosion, gravelly 35-60%	21.42 (4.24)
	BHI	Bhimanahalli soils area shallow (25-50 cm), well drained, have very dark gray to brown clay soils occurring on very gently sloping to gently sloping uplands.		<b>20.30 (4.02)</b>
11		BHImB2g1	Clay surface, slope 1-3%, moderate erosion, gravelly (15-35%)	20.30 (4.02)
	DSI	Dinsi soils are moderately shallow (50-75 cm), moderately well drained, have very dark gray to brown clayey soils occurring on very gently sloping uplands		<b>140.1 (27.8)</b>
12		DSImA1	Clay surface, slope 0-1%, slight erosion	50.20 (9.95)

13		DSImB2g1	Clay surface, slope 1-3%, moderate erosion, gravelly (15-35%)	80.83 (16.02)
14		DSImC2g1	Clay surface, slope 3-5%, moderate erosion, gravelly (15-35%)	9.13 (1.81)
	KMP	Kamalapur soils are moderately deep (75-100 cm), moderately well drained, have very dark gray to very dark grayish brown cracking clay soils occurring on very gently sloping uplands		<b>9.01</b> <b>(1.79)</b>
15		KMPmB2g1	Clay surface, slope 1-3%, moderate erosion, gravelly (15-35%)	9.01 (1.79)
	RNL	Rajanala soils are deep (100-150 cm), moderately well drained, have very dark gray to brown cracking clay soils occurring on very gently sloping uplands		<b>40.0</b> <b>(7.92)</b>
16		RNLmA1	Clay surface, slope 0-1%, slight erosion	18.95 (3.75)
17		RNLmB1g1	Clay surface, slope 1-3%, slight erosion, gravelly (15-35%)	4.62 (0.92)
18		RNLmB2	Clay surface, slope 1-3%, moderate erosion	16.41 (3.25)
<b>Miscellaneous Lands</b>				
19		Habitation		38.14 (7.56)

## THE SOILS

Detailed information pertaining to the nature, extent and distribution of different kinds of soils occurring in Nirgudi West microwatershed is provided in this chapter. The microwatershed area has been identified as basalt landscape. In all, 6 soil series were identified in this landscape. Soil formation is the result of the combined effect of environmental and terrain factors that are reflected in soil morphology. In the basalt landscape, it is by parent material and climate. A brief description of each of the 6 soil series identified followed by 18 soil phases (management units) mapped under each series are furnished below. The soils in any one map unit differ from place to place in their depth, texture, slope, gravelliness, erosion or any other site characteristics that affect management. The soil phase map can be used for identifying the suitability of areas for growing specific crops or for other alternative uses and also for deciding the type of conservation structures needed. The detailed information on soil and site-characteristics like soil depth, surface soil texture, slope, erosion, gravelliness, AWC, LCC etc, with respect to each of the soil phase identified is given village/survey number wise for the microwatershed in Appendix-I.

### 4.1 Soils of Basalt Landscape

In this landscape, 6 soil series are identified and mapped. Of these, Novanihala (NHA) soil series occupies maximum area of about 140 ha (28%) followed by Dinsi (DSI) soils 140 ha (28%) and Margutti soils 117 (23%). The brief description of each series along with the soil phases identified and mapped is given below.

**4.1.1 Margutti (MGT) Series:** Margutti soils are very shallow (<25cm), well drained, have very dark grayish brown to dark brown clay soils. They have developed from basalt and occur on very gently sloping to strongly sloping uplands.

The total depth of the soil ranges from 10 to 23 cm. The thickness of A horizon ranges from 7 to 18 cm. Its colour is in 10 YR and 7.5 YR hue with value 3 to 4 and chroma 2 to 4. The texture is clay with 15 to 35 per cent gravel. The available water capacity is very low (<50 mm/m).

Four phases were identified:

MGTmB2g1	Clay surface, slope 1-3 %, moderate erosion, gravelly (15-35%)
MGTmC3g1	Clay surface, slope 3-5 %, severe erosion, gravelly (15-35 %)
MGTmC3g2	Clay surface, slope 3-5%, severe erosion, very gravelly (35-60%)
MGTmD3g2	Clay surface, 5-10 % slope, severe erosion, very gravelly (35-60%)



Landscape and Soil Profile Characteristics of Margutti (MGT) Series

**4.1.2 Novanihala (NHA) Series:** Novanihala soils are shallow (25-50 cm), well drained, have very dark grayish brown to dark brown clay soils. They have developed from basalt and occur on very gently to moderately sloping uplands.

The thickness of the solum ranges from 27 to 48 cm. The thickness of A horizon ranges from 12 to 20 cm. Its colour is in 7.5 YR and 10 YR hue with value 3 to 4 and chroma 2 to 4. The texture varies from sandy clay to clay with 10 to 20 per cent gravel. The thickness of B horizon ranges from 22 to 37 cm. Its colour is in 10 YR and 7.5 YR hue with value 3 to 4 and chroma 2 to 4. Its texture is clay with gravel content of 10-15 per cent. The available water capacity is low (51-100 mm/m).

Six phases were identified:

NHAmB1	Clay surface, slope 1-3%, slight erosion
NHAmB2	Clay surface, slope 1-3%, moderate erosion
NHAmB2g1	Clay surface, slope 1-3%, moderate erosion, gravelly 15-35%
NHAmC2g1	Clay surface, slope 3-5%, moderate erosion, gravelly 15-35%
NHAmC3	Clay surface, slope 3-5%, severe erosion
NHAmD3g2	Clay surface, slope 5-10%, severe erosion, very gravelly 35-60%



Landscape and Soil Profile Characteristics of Novanihala (NHA) Series

**4.1.3 Bhimanahalli (BHI) Series:** Bhimanahalli soils are shallow (25-50 cm), well drained, have very dark gray to brown clay soils. They have developed from basalt and occur on very gently sloping uplands.

The thickness of the solum ranges from 29 to 48 cm. The thickness of A horizon ranges from 15 to 20 cm. Its colour is in 7.5 YR and 10 YR hue with value 3 to 4 and chroma 2 to 4. The texture varies from sandy clay to clay with 15 to 25 per cent gravel. The thickness of B horizon ranges from 23 to 33 cm. Its colour is in 10 YR and 7.5 YR hue with value 3 to 4 and chroma 1 to 3. Its texture is clay with gravel content of 15 to 35 per cent. The available water capacity is very low (<50 mm/m).

One phase was identified:

BHImB2g1	Clay surface, slope 1-3%, moderate erosion, gravelly 15-35%
----------	-------------------------------------------------------------



Landscape and Soil Profile Characteristics of Bhimanahalli (BHI) Series

**4.1.4 Dinsi (DSI) Series:** Dinsi soils are moderately shallow (50-75 cm), moderately well drained, have very dark gray to brown clay soils. They have developed from basalt and occur on very gently sloping uplands.

The thickness of the solum ranges from 55 to 71 cm. The thickness of A horizon ranges from 9 to 24 cm. Its colour is in 10 YR hue with value 3 and chroma 1 to 3. The texture is clay with 5 to 10 per cent gravel. The thickness of B horizon ranges from 27 to 62 cm. Its colour is in 10 YR hue with value 3 to 4 and chroma 2 to 4. Its texture is clay with gravel content of less than 15 per cent. The available water capacity is medium (101-150 mm/m).

Three phases were identified:

DSImA1	Clay surface, slope 0-1%, slight erosion
DSImB2g1	Clay surface, slope 1-3%, moderate erosion, gravelly 15-35%
DSImC2g1	Clay surface, slope 3-5%, moderate erosion, gravelly 15-35%



Landscape and Soil Profile Characteristics of Dinsi (DSI) Series


**4.1.5 Kamalapur (KMP) Series:** Kamalapur soils are moderately deep (75-100 cm), moderately well drained, have very dark gray to very dark grayish brown cracking clay soils. They have developed from basalt and occur on very gently sloping uplands.

The thickness of the solum ranges from 75 to 95 cm. The thickness of A horizon ranges from 10 to 30 cm. Its colour is in 10 YR and 7.5 YR hue with value 3 to 4 and chroma 1 to 4. The texture is clay with less than 10 per cent gravel. The thickness of B horizon ranges from 45 to 84 cm. Its colour is in 10 YR hue with value 3 to 4 and chroma 1 to 4. Its texture is clay with gravel content of less than 15 per cent. The available water capacity is medium (101-150 mm/m).

One phase was identified:

KMPmB2g1	Clay surface, 1-3% slope, moderate erosion, gravelly 15-35%
----------	-------------------------------------------------------------






Landscape and Soil Profile characteristics of Kamalapur (KMP) Series

**4.1.6 Rajanala (RNL) Series:** Rajanala soils are deep (100-150 cm), moderately well drained, have very dark gray to brown cracking clay soils. They have developed from basalt and occur on very gently to gently sloping uplands.

The thickness of the solum ranges from 125 to 140 cm. The thickness of A horizon ranges from 14 to 23 cm. Its colour is in 10 YR hue with value 3 to 4 and chroma 1 to 2. The texture is clay with less than 10 per cent gravel. The thickness of B horizon ranges from 85 to 130 cm. Its colour is in 10 YR hue with value 3 to 4 and chroma 1 to 3. Its texture is clay with gravel content of less than 15 per cent. The available water capacity is very high (>200 mm/m).

Three phases were identified:

RNLmA1	Clay surface, slope 0-1%, slight erosion
RNLmB1g1	Clay surface, slope 1-3%, slight erosion, gravelly 15-35%
RNLmB2	Clay surface, slope 1-3%, moderate erosion



Landscape and Soil profile Characteristics of Rajanala (RNL) Series

## INTERPRETATION FOR LAND RESOURCE MANAGEMENT

The most important soil and site characteristics that affect the land use and conservation needs of an area are land capability, soil depth, soil texture, coarse fragments, available water capacity, soil slope, soil erosion, soil reaction etc. These are interpreted from the data base generated through land resource inventory and several thematic maps are generated. These would help in identifying the areas suitable for growing crops and, soil and water conservation measures and structures needed thus helping to maintain good soil health for sustained crop production. The various thematic maps generated are described below.

### 5.1 Land Capability Classification

Land capability classification is an interpretative grouping of soil map units (soil phases) mainly based on inherent soil characteristics, external land features and environmental factors that limit the use of land for agriculture, pasture, forestry, or other uses on a sustained basis (IARI, 1971). The land and soil characteristics used to group the land resources in an area into various land capability classes, subclasses and units are

*Soil characteristics:* Soil depth, soil texture, coarse fragments, soil reaction, available water capacity, calcareousness, salinity/alkali *etc.*

*Land characteristics:* Slope, erosion, drainage, rock outcrops.

*Climate:* Total rainfall and its distribution, and length of crop growing period.

The Land capability classification system is divided into land capability classes, subclasses and units based on the level of information available. Eight land capability classes are recognized. They are

*Class I:* They are very good lands that have no limitation or very few limitations that restrict their use.

*Class II:* They are good lands that have minor limitation and require moderate conservation practices.

*Class III:* They are moderately good lands that have moderate limitation that reduce the choice of crops or that require special conservation practices.

*Class IV:* They are fairly good lands have very severe limitations that reduce the choice of crops or that require very careful management.

*Class V:* Soils in these lands are not likely to erode, but have other limitations that are impractical to remove and as such not suitable for agriculture, but suitable for pasture or forestry with minor limitation.

*Class VI:* The lands have severe limitations that make them generally unsuitable for cultivation, but suitable for pasture or forestry with moderate limitation.

*Class VII:* The lands have very severe limitations that make them unsuitable for cultivation, but suitable for pasture or forestry with major limitations.

*Class VIII:* Soil and other miscellaneous areas (rock lands) that have very severe limitations that nearly preclude their use for any crop production, but suitable for wildlife, recreation and wind mills.

The land capability subclasses are recognised based on the dominant limitations observed within a given land capability class. The subclasses are designated by adding a lower case letter like 'e', 'w', 's', or 'c' to the class numeral. The subclass "e" indicates that the main hazard is risk of erosion, "w" indicates drainage or wetness as a limitation for plant growth, "s" indicates shallow soil depth, coarse or heavy textures, calcareousness, salinity/alkalinity or graveliness and "c" indicates limitation due to climate.

The land capability subclasses have been further subdivided into land capability units based on the kinds of limitations present in each subclass. Ten land capability units are used in grouping the soil map units. They are stony or rocky (0), erosion hazard (slope, erosion) (1), coarse texture (sand, loamy sand, sandy loam) (2), fine texture (cracking clay, silty clay) (3), slowly permeable subsoil (4), coarse underlying material (5), salinity/alkali (6), stagnation, overflow, high ground water table (7), soil depth (8) and fertility problems (9). The capability units thus identified have similar soil and land characteristics that respond similarly to a given level of management. The soils of the microwatershed have been classified upto land capability subclass level only.

The 18 soil map units identified in the Nirgudi West microwatershed are grouped under 4 land capability classes and 6 land capability subclasses. About 80 per cent area in the microwatershed is suitable for agriculture (Fig. 5.1) and 12 per cent not suitable for agriculture but well suited for grazing or forestry, recreation and wildlife.

Good cultivable lands (Class II) cover maximum area of about 37 per cent and are distributed in the southeastern, southwestern, northwestern and central part of the microwatershed with problems of soil and erosion. Moderately good cultivable lands (Class III) cover an area of about 29 per cent and are distributed in the southern, western, northeastern and eastern part of the microwatershed with moderate problems of erosion and soil.

The fairly good lands (class IV) cover about 14 per cent of area. They have limitations of erosion and soil and are distributed in the northeastern and central part of the microwatershed.

The non cultivable lands (class VI) cover about 12 per cent and are distributed in the eastern, northern and central part of the microwatershed. They are well suited for pasture, forestry, wild life and recreation.




Fig. 5.1 Land Capability map of Nirgudi West Microwatershed

## 5.2 Soil Depth

Soil depth refers to the depth of the soil occurring above the parent material or hard rock. The depth of the soil determines the effective rooting depth for plants and in accordance with soil texture, mineralogy and gravel content, the capacity of the soil column to hold water and nutrient availability. Soil depth is one of the most important soil characteristic that is used in differentiating soils into different soil series. The soil depth classes used in identifying soils in the field are very shallow soils (<25 cm), shallow (25-50 cm), moderately shallow (50-75 cm), moderately deep (75-100 cm), deep (100-150 cm) and very deep (>150 cm). They were used to classify the soils into different depth classes and a soil depth map was prepared (Fig. 5.2).

Deep soils (100-150 cm) occur in a small area of about 40 ha (8%) and are distributed in the central part of the microwatershed. Moderately deep (75-100 cm) soils occur in a very small area 9 ha (2%). Moderately shallow (50-75 cm) soils occupy about 140 ha (28%) and are distributed in the southwestern, southeastern, central and northwestern part of the microwatershed.

Major area of about 161 ha (32%) is under shallow soils (25-50 cm) and are distributed in the southern, northern, western and northeastern part of the microwatershed. Very shallow soils (<25 cm) occupy an area of about 116 ha (23%) in the northeastern and southwestern part of the microwatershed.

The most productive lands are about 40 ha (8%) with respect to soil rooting depth where all climatically adapted annual and perennial crops can be grown are deep soils (100-150 cm depth) occurring in the central part of the microwatershed.

The most problem lands with a maximum area of about 277 ha (55%) having very shallow (<25 cm) and shallow (25-50 cm) rooting depth occur in all parts of the microwatershed. They are not suitable for growing agricultural crops but well suited for pasture, forestry or other recreational purposes. Occasionally, short duration crops may be grown if rainfall is normal.




Fig. 5.2 Soil Depth map of Nirgudi West Microwatershed

### 5.3 Surface Soil Texture

Texture is an expression to indicate the coarseness or fineness of the soil as determined by the relative proportion of primary particles of sand, silt and clay. It has a direct bearing on the structure, porosity, adhesion and consistence. The surface layer of a soil to a depth of about 25 cm is the layer that is most used by crops and plants. The surface soil textural class provides a guide to understanding soil-water retention and availability, nutrient holding capacity, infiltration, workability, drainage, physical and chemical behaviour, microbial activity and crop suitability.

The entire microwatershed is clayey in surface soil texture (Fig. 5.3). The most productive lands (92%) with respect to surface soil texture are the clayey soils that have high

potential for soil-water retention and availability, and nutrient retention and availability, but have more problems of drainage, infiltration, workability and other physical problems.




Fig. 5.3 Surface Soil Texture map of Nirgudi West Microwatershed

#### 5.4 Soil Gravelliness

Gravel is the term used for describing coarse fragments between 2 mm and 7.5 cm diameter and stones for those between 7.5 cm and 25 cm. The presence of gravel and stones in soil reduces the volume of soil responsible for moisture and nutrient storage, drainage, infiltration and runoff and hinders plant growth by impeding root growth and seedling emergence, intercultural operations and farm mechanization.

Maximum area has soils that are gravelly (15-35%) covering about 185 ha (37%) and are distributed in the southwestern, western and eastern part the microwatershed (Fig.5.4).

About 163 ha (32%) area in the microwatershed has soils that are nongravelly (<15%) and are distributed in the southeastern, central and northeastern part the microwatershed followed by soils that are very gravelly (35-60%) covering about 119 ha (23%) that are distributed in the northern, central and eastern part of the microwatershed.

The most productive lands with respect to gravelliness are found to be 32 per cent. They are nongravelly with less than 15 per cent gravel and have potential for growing both annual and perennial crops.

The problem soils that are very gravelly (35-60%) are found to cover about 23 per cent area, where only short duration crops can be grown.




Fig. 5.4 Soil Gravelliness map of Nirgudi West Microwatershed

### 5.5 Available Water Capacity

The soil available water capacity (AWC) is estimated based on the ability of the soil column to retain water between the tensions of 0.33 and 15 bar in a depth of 100 cm or the entire solum if the soil is shallower. The AWC of the soils (soil series) as estimated by considering the soil texture, mineralogy, soil depth and gravel content (Sehgal *et al.*, 1990) and accordingly the soil map units were grouped into five AWC classes *viz.* very low (<50 mm/m), low (50-100 mm/m), medium (100-150 mm/m), high (150-200 mm/m) and very high (>200 mm/m) and using these values, an AWC map was prepared (Fig. 5.5).

An area of about 137 ha (27%) has soils that are very low (<50 mm/m) in available water capacity and are distributed in the northeastern, central and southern part of the microwatershed. An area of about 141 ha (28%) has soils that are low (51-100 mm/m) in available water capacity and are distributed in the northern, northeastern and western part of the microwatershed. The major area in the microwatershed has soils that are medium (101-150 mm/m) in available water capacity. They occur in an area of about 149 ha (30%) and are distributed in the northwestern, central and southeastern part of the microwatershed and soils that are very high (>200 mm/m) in AWC covering a small area of about 40 ha (8%) in the microwatershed and are distributed in central part of the microwatershed.

An area of about 40 ha (8%) has soils that have very high potential (>200 mm/m) with regard to available water capacity. In these areas, if the rainfall is normal and well distributed,



all climatically adapted long duration annual and perennial crops can be grown. About 278 ha (55%) area in the microwatershed has soils that are problematic with regard to available water capacity. Here, only short duration crops can be grown and the probability of crop failure is very high. These areas are best put to other alternative uses.




Fig. 5.5 Soil Available Water Capacity map of Nirgudi West Microwatershed

## 5.6 Soil Slope

Soil slope refers to the inclination of the surface of the land. It is defined by gradient, shape and length, and is an integral feature of any soil as a natural body. Slope is considered important in soil genesis, land use and land development. The length and gradient of slope influences the rate of runoff, infiltration, erosion and deposition. The soil map units were grouped into four slope classes and a slope map was prepared showing the area extent and geographic distribution of different slope classes in the microwatershed (Fig. 5.6).

Major area of the microwatershed falls under very gently sloping (1-3%) class. It covers an area of about 244 ha (48%) and is distributed in all parts of the microwatershed. An area of about 93 ha (18 %) in the microwatershed falls under gently sloping (3-5%) class and is distributed in the central, southwestern and northeastern part of the microwatershed. Moderately sloping (5-10%) lands cover an area of about 61 ha (12%) and is distributed in the central and northern part of the microwatershed. Nearly level lands (0-1%) slope class covers about 69 ha (14%) and are distributed in the southeastern and central part of the microwatershed.

An area of about 313 ha (62%) in the microwatershed has soils that have high potential in respect of soil slopes. In these areas, all climatically adapted annual and perennial crops can be grown without much soil and water conservation and other land development measures.

An area of about 61 ha (12%) in the microwatershed has soils that are problematic in respect of slopes.




Fig. 5.6 Soil Slope map of Nirgudi West Microwatershed

### 5.7 Soil Erosion

Soil erosion refers to the wearing away of the earth's surface by the forces of water, wind and ice involving detachment and transport of soil by raindrop impact. It is used for accelerated soil erosion resulting from disturbance of the natural landscape by burning, excessive grazing and indiscriminate felling of forest trees and tillage, all usually by man. The erosion classes showing an estimate of the current erosion status as judged from field observations in the form of rills, gullies or a carpet of gravel on the surface are recorded. Four erosion classes, viz, slight erosion (e1), moderate erosion (e2), severe erosion (e3) and very severe erosion (e4) are recognized. The soil map units were grouped into different erosion classes and soil erosion map was generated. The area extent and their spatial distribution in the microwatershed is given in Figure 5.7.

Soils that are slightly eroded (e1 class) covers about 83 ha (16%) and are distributed in the southeastern, northern and central part of the microwatershed. Soils that are moderately

eroded (e2 class) cover a maximum area of about 250 ha (50%) in the microwatershed and are distributed in all parts of the microwatershed. Severely eroded (e3 class) soils cover an area of about 133 ha (26%) and are distributed in the central, southwestern and northeastern part of the microwatershed.

Top priority is to be given to 133 ha area where they are severely eroded for taking up soil and water conservation and other land development measures followed by moderately eroded lands that cover about 250 ha.




Fig. 5.7 Soil Erosion map of Nirgudi West Microwatershed



## FERTILITY STATUS

Soil fertility plays an important role in increasing crop yield. The adoption of high yielding varieties that require high amounts of nutrients has resulted in deficiency symptoms in crops and plants due to imbalanced fertilization and poor inherent fertility status. Hence, it is necessary to know the fertility (macro and micro nutrients) status of the soils of the watersheds for assessing the kind and amount of fertilizers required for each of the crop intended to be grown. For this purpose, the surface soil samples collected from the grid points (one soil sample at every 250 m interval) all over the microwatershed through land resource inventory in the year 2014 were analysed for pH, EC, organic carbon, available phosphorus and potassium and for micronutrients like zinc, boron, copper, iron and manganese and secondary nutrient sulphur.

Soil fertility data generated has been assessed and individual maps for all the nutrients for the microwatershed have been prepared. The village/survey number wise fertility data for the microwatershed is given in Appendix-II.

### 6.1 Soil Reaction (pH)

The soil fertility analysis of the Nirgudi West microwatershed for soil reaction (pH) showed maximum area of about 205 ha (41%) is neutral (pH 6.5-7.3) in reaction and is distributed in the northeastern, northwestern and western part of the microwatershed (Fig.6.1). Slightly alkaline (pH 7.3-7.8) is around 142 ha (28%) area and is distributed in the central, northern, northeastern and eastern part of the microwatershed. Moderately alkaline (pH 7.8-8.4) is around 120 ha (24%) area and is distributed in the central and southern part of the microwatershed.

### 6.2 Electrical Conductivity (EC)

The Electrical Conductivity of the soils of the entire microwatershed area is  $<2 \text{ dSm}^{-1}$  (Fig 6.2) and as such the soils in the microwatershed are nonsaline.

### 6.3 Organic Carbon

The soil organic carbon content of the soils in the microwatershed is medium (0.5-0.75%) in 161 ha (32%) area that are distributed in the southern and northwestern part of the microwatershed (Fig.6.3). High ( $>0.75\%$ ) organic carbon content accounts for maximum area of 286 ha (57%) and is distributed in all parts of the microwatershed. Low ( $<0.5\%$ ) organic carbon content accounts for a very small area of 20 ha (4%) in the microwatershed and is distributed in the southern part of the microwatershed.




Fig.6.1 Soil Reaction (pH) map of Nirgudi West Microwatershed




Fig.6.2 Electrical Conductivity (EC) map of Nirgudi West Microwatershed

#### **6.4 Available Phosphorus**

The soil fertility analysis revealed that available phosphorus is low (<23 kg/ha) in major area of about 383 ha (76%) and is distributed in all parts of the microwatershed (Fig.6.4). There is an urgent need to increase the dose of phosphorous for all the crops by 25 per cent over the recommended dose to realize better crop performance. About 78 ha (15%) area is medium (23-57 kg/ha) and is distributed in the southeastern and northern part of the microwatershed. High available phosphorus (>57 kg/ha) content accounts for very less area of 6 ha (1%) and is distributed in the northwestern part of microwatershed.

#### **6.5 Available Potassium**

Available potassium content is medium (145-337 kg/ha) in 126 ha (25%) area and is distributed in the central, eastern, western and southwestern part of the microwatershed (Fig.6.5) and high available potassium (>337 kg/ ha) content accounts for major area of 341 ha (67%) and is distributed in all parts of the microwatershed.

#### **6.6 Available Sulphur**

Available sulphur content is low (<10 ppm) in a very small area 6 ha (1%). Maximum area of about 459 ha (91%) is medium (10-20 ppm) in available sulphur and is distributed in all parts of the microwatershed (Fig.6.6). Available sulphur is high (>20 ppm) in a very small area of 2 ha in the microwatershed.

#### **6.7 Available Boron**

Available boron content is low (<0.5 ppm) in major area of about 332 ha (66%) and is distributed in all parts of the microwatershed. About 127 ha (25%) has soils that are medium (0.5-1.0 ppm) in available boron (Fig 6.7) and is distributed in the northern, eastern and northwestern part of the microwatershed.




Fig.6.3 Soil Organic Carbon map of Nirgudi West Microwatershed




Fig.6.4 Soil available Phosphorus map of Nirgudi West Microwatershed






Fig.6.5 Soil available Potassium map of Nirgudi West Microwatershed




Fig.6.6 Soil available Sulphur map of Nirgudi West Microwatershed




Fig.6.7 Soil available Boron map of Nirgudi West Microwatershed

## 6.8 Available Iron

Available iron content is deficient (<4.5 ppm) in a small area of 11 ha (2%) and is distributed in the southwestern part of the microwatershed. It is sufficient in major area of 455 ha (90%) and is distributed in all parts of the microwatershed (Fig 6.8).

## 6.9 Available Manganese

Available manganese content is sufficient (>1.0 ppm) in the entire microwatershed area (Fig 6.9).

## 6.10 Available Copper

Available copper content is sufficient (>0.2 ppm) in the entire microwatershed area (Fig 6.10).

## 6.11 Available Zinc

Available zinc content is deficient (<0.6 ppm) in an area of about 165 ha (33%) and is distributed in the southern, central and northwestern part of the microwatershed (Fig 6.11). It is sufficient (>0.6 ppm) in major area of about 302 ha (60%) and is distributed in all parts of the microwatershed except southern part.




Fig.6.8 Soil available Iron map of Nirgudi West Microwatershed




Fig.6.9 Soil available Manganese map of Nirgudi West Microwatershed




Fig.6.10 Soil available Copper map of Nirgudi West Microwatershed




Fig.6.11 Soil available Zinc map of Nirgudi West Microwatershed

## LAND SUITABILITY FOR MAJOR CROPS

The soil and land resource units (soil phases) of Nirgudi West microwatershed were assessed for their suitability for growing food, fodder, fibre and other horticulture crops by following the procedure as outlined in FAO, 1976 and 1983. Crop requirements were developed for each of the crop from the available research data and also by referring to Naidu *et. al.*, (2006) and Natarajan *et. al.*, (2015). The crop requirements were matched with the soil and land characteristics (Table 7.1) to arrive at the crop suitability. In FAO land suitability classification, two orders are recognized. Order S-Suitable and Order N-Not suitable. The orders have classes, subclasses and units. Order-S has three classes, Class S1-Highly Suitable, Class S2-Moderately Suitable and Class S3-Marginally Suitable. Order N has two classes, N1-Currently not Suitable and N2-Permanently not Suitable. There are no subclasses within the class S1 as they will have very minor or no limitations for crop growth. Classes S2 and S3 are divided into subclasses based on the kinds of limitations encountered. The limitations that affect crop production are ‘c’ for erratic rainfall and its distribution and length of growing period (LGP), ‘e’ for erosion hazard, ‘r’ for rooting condition, ‘t’ for lighter or heavy texture, ‘g’ for gravelliness or stoniness, ‘n’ for nutrient availability, ‘l’ for topography, ‘m’ for moisture availability and ‘w’ for drainage. These limitations are indicated as lower case letters to the class symbol. For example, moderately suitable land with the limitations of soil depth and erosion is designated as S2re. For the microwatershed, the soil mapping units were evaluated and classified up to subclass level.

Using the above criteria, the soil map units of the microwatershed were evaluated and land suitability maps for 18 major annual and perennial crops were generated. The detailed information on the kind of suitability of each of the soil phase for the crops assessed are given village/ survey number wise for the microwatershed in Appendix-III.

### 7.1 Land Suitability for Sorghum (*Sorghum bicolor*)

Sorghum is one of the major crops grown in Karnataka in an area of 11.02 lakh ha in northern Karnataka in Bijapur, Kalaburgi, Raichur, Bidar, Belgaum, Dharwad and Bellary districts. The crop requirements for growing sorghum (Table 7.2) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and land suitability map for growing sorghum generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed is given in Figure 7.1.

A very small area of about 24 ha (5%) in the microwatershed has soils that are highly suitable (class S1) for growing sorghum crop. They are distributed mainly in the central part of the microwatershed. An area of about 165 ha (33%) is moderately suitable (class S2) for growing sorghum and are distributed in the southern, central and northwestern part of the microwatershed. They have major limitations of erosion and rooting depth.

**Table 7.1 Soil-Site Characteristics of Nirgudi West Microwatershed**


Soil Map Units	Climate (P) (mm)	Growing period (Days)	Drainage class	Soil depth (cm)	Soil texture		Gravelliness		AWC (mm/m)	Slope (%)	Erosion	pH	EC	ESP	CEC [Cmol (p <sup>+</sup> ) kg <sup>-1</sup> ]	BS (%)
					Surf-ace	Sub-surface	Surface (%)	Subsurfa-ce (%)								
MGTmB2g1	786	150	WD	<25	c	c	15-35	15-35	<50	1-3	Moderate	6.8	0.3	0.2	46	100
MGTmC3g1	786	150	WD	<25	c	c	15-35	15-35	<50	3-5	Severe	6.8	0.3	0.2	46	100
MGTmC3g2	786	150	WD	<25	c	c	35-60	15-35	<50	3-5	Severe	6.8	0.3	0.2	46	100
MGTmD3g2	786	150	WD	<25	c	c	35-60	15-35	<50	5-10	Severe	6.8	0.3	0.2	46	100
BHImB2g1	786	150	WD	25-50	c	c	15-35	15-35	<50	1-3	Moderate	7.0	0.1	0.2	28	100
NHAmB1	786	150	WD	25-50	c	c	-	<15	51-100	1-3	Slight	7.2	0.1	0.3	40	100
NHAmB2	786	150	WD	25-50	c	c	-	<15	51-100	1-3	Moderate	7.2	0.1	0.3	40	100
NHAmB2g1	786	150	WD	25-50	c	c	15-35	<15	51-100	1-3	Moderate	7.2	0.1	0.3	40	100
NHAmC2g1	786	150	WD	25-50	c	c	15-35	<15	51-100	3-5	Moderate	7.2	0.1	0.3	40	100
NHAmC3	786	150	WD	25-50	c	c	-	<15	51-100	3-5	Severe	7.2	0.1	0.3	40	100
NHAmD3g2	786	150	WD	25-50	c	c	35-60	<15	51-100	5-10	Severe	7.2	0.1	0.3	40	100
DSImA1	786	150	MWD	50-75	c	c	-	<15	101-150	0-1	Slight	7.0	0.1	0.3	62	100
DSImB2g1	786	150	MWD	50-75	c	c	15-35	<15	101-150	1-3	Moderate	7.0	0.1	0.3	62	100
DSImC2g1	786	150	MWD	50-75	c	c	15-35	<15	101-150	3-5	Moderate	7.0	0.1	0.3	62	100
KMPmB2g1	786	150	MWD	75-100	c	c	15-35	<15	101-150	1-3	Moderate	6.7	0.2	0.2	43	100
RNLmA1	786	150	MWD	100-150	c	c	-	<15	>200	0-1	Slight	8.4	0.2	0.2	60	100
RNLmB1g1	786	150	MWD	100-150	c	c	15-35	<15	>200	1-3	Slight	8.4	0.2	0.2	60	100
RNLmB2	786	150	MWD	100-150	c	c	-	<15	>200	1-3	Moderate	8.4	0.2	0.2	60	100

\*Symbols and abbreviations are according to Field Guide for LRI under Sujala-III Project, Karnataka

Marginally suitable lands (class S3) for growing sorghum occupy an area of about 139 ha (28%) and occur in the southern, northern and northeastern part of the microwatershed. They have severe limitations of rooting depth and erosion. An area of about 138 ha (27%) is not suitable for growing sorghum in the microwatershed and occur in the central, eastern and northeastern part of the microwatershed.

**Table 7.2 Crop suitability criteria for Sorghum**

Crop requirement		Rating			
Soil –site characteristics	unit	Highly suitable (S1)	Moderately Suitable (S2)	Marginally suitable (S3)	Not suitable (N)
Slope	%	2-3	3-8	8-15	>15
LGP	Days	120-150	120-90	<90	
Soil drainage	class	Well to mod. drained	imperfect	Poorly/ excessively	V. poorly
Soil reaction	pH	6.0-8.0	5.5-5.9 8.1-8.5	<5.5 8.6-9.0	>9.0
Surface soil texture	Class	C, cl, siel, sc	l, sil, sic	Sl, ls	S, fragmental skeletal
Soil depth	cm	100-75	50-75	30-50	<30
Gravel content	% vol.	5-15	15-30	30-60	>60
Salinity (EC)	dSm <sup>-1</sup>	2-4	4-8	8-10	>10
Sodicity (ESP)	%	5-8	8-10	10-15	>15



**Fig. 7.1 Land Suitability map of Sorghum**

## 7.2 Land Suitability for Maize (*Zea mays*)

Maize is the most important food crop grown in an area of 13.73 lakh ha in all the districts of Karnataka. The crop requirements for growing maize (Table 7.3) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing maize was generated. The area and geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.2.

In Nirgudi West watershed, there are no highly (Class S1) suitable lands for growing maize. An area of about 140 ha (28%) is moderately suitable (class S2) for growing maize and are distributed in the southwestern and northwestern part of the microwatershed. They have major limitations of erosion and rooting depth.

The marginally suitable (class S3) lands cover a maximum area of about 188 ha (37%) and occur in all parts of the microwatershed. They have severe limitations of texture, erosion and rooting depth. About 138 ha (27%) area is not suitable for growing maize and occurs in the eastern, southwestern and northeastern part of the microwatershed.

**Table 7.3 Crop suitability criteria for Maize**

Crop requirement		Rating			
Soil –site characteristics	unit	Highly suitable (S1)	Moderately Suitable (S2)	Marginally suitable (S3)	Not suitable (N)
Slope	%	<3	3.5	5-8	
LGP	Days	>100	100-80	60-80	
Soil drainage	class	Well drained	Mod. to imperfectly	Poorly/excessively	V.poorly
Soil reaction	pH	5.5-7.5	7.6-8.5	8.6-9.0	
Surface soil texture	Class	l, cl, scl, sil	Sl, sicl, sic	C(s-s), ls	S,fragmental
Soil depth	Cm	>75	50-75	25-50	<25
Gravel content	% vol.	<15	15-35	35-50	>50
Salinity (EC)	dSm <sup>-1</sup>	<1.0	1.0-2.0	2.0-4.0	
Sodicity (ESP)	%	<10	10-15	>15	






Fig. 7.2 Land Suitability map of Maize


### 7.3 Land Suitability for Red gram (*Cajanus cajan*)

Red gram is one of the major pulse crop grown in an area of 8.23 lakh ha mainly in northern Karnataka in Bijapur, Kalaburgi, Raichur, Bidar, Belgaum, Dharwad and Bellary districts. The crop requirements for growing red gram (Table 7.4) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and land suitability map for growing red gram was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed is given in Figure 7.3.

The major area of about 189 ha (37%) is moderately suitable (class S2) for red gram and is dominantly distributed in the northwestern, southwestern and central part of the microwatershed. They have major limitations of rooting depth, erosion and texture. An area of about 139 ha (28%) is marginally suitable (class S3) for growing red gram and are distributed in the southern, western and northeastern part of the microwatershed. They have major limitations of rooting depth and erosion. An area of about 138 ha (27%) is not suitable for growing red gram and occur in the eastern, northeastern and central part of the microwatershed.

**Table 7.4 Crop suitability criteria for Red gram**

Crop requirement		Rating			
Soil characteristics	unit	Highly suitable (S1)	Moderately Suitable (S2)	Marginally suitable (S3)	Not suitable (N)
Slope	%	<3	3-5	5-10	>10
LGP	Days	>210	180-210	150-180	<150
Soil drainage	class	Well drained	Mod. to well drained	Imperfectly drained	Poorly drained
Soil reaction	pH	6.5-7.5	5.0-6.5 7.6-8.0	8.0-9.0	>9.0
Surface soil texture	Class	l, scl, sil, cl, sl	sicl, c(m)	sic, ls	S, fragmental
Soil depth	Cm	>100	85-100	40-85	<40
Gravel content	% vol.	<20	20-35	35-60	>60
Salinity (EC)	dSm <sup>-1</sup>	<1.0	1.0-2.0	>2.0	
Sodicity (ESP)	%	<10	10-15	>15	



**Fig. 7.3 Land Suitability map of Red gram**

#### 7.4 Land Suitability for Sunflower (*Helianthus annuus*)

Sunflower is the most important oilseed crop grown in an area of 4.1 lakh ha in the State in all the districts. The crop requirements for growing sunflower (Table 7.5) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing sunflower was generated. The area extent and geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.4.

Highly suitable (class S1) lands are found to occur in minor area of 24 ha (5%) and is distributed in the central part of the microwatershed. A small area of about 25 ha (5%) is moderately suitable (class S2) for sunflower and is distributed in the central part of the microwatershed. They have major limitation of erosion.

Marginally suitable (class S3) lands are found to occur in an area of about 140 ha (28%). The soils have severe limitation of rooting depth. They are dominantly distributed in the southwestern, southeastern, and northeastern part of the microwatershed. Major area of about 277 ha (55%) is not suitable for growing sunflower and occur in all parts of the microwatershed.

**Table 7.5 Crop suitability criteria for Sunflower**

Crop requirement		Rating			
Soil –site characteristics	unit	Highly suitable (S1)	Moderately Suitable (S2)	Marginally suitable (S3)	Not suitable (N)
Slope	%	<3	3-5	5-10	>10
LGP	Days	>90	80-90	70-80	<70
Soil drainage	class	Well drained	mod. Well drained	imperfectly drained	Poorly drained
Soil reaction	pH	6.5-8.0	8.1-8.5 5.5-6.4	8.6-9.0; 4.5-5.4	>9.0 <4.5
Surface soil texture	Class	1, cl, sil, sc	Scl, sic, c,	c (>60%), sl	ls, s
Soil depth	Cm	>100	75-100	50-75	<50
Gravel content	% vol.	<15	15-35	35-60	>60
Salinity (EC)	dSm <sup>-1</sup>	<1.0	1.0-2.0	>2.0	
Sodicity (ESP)	%	<10	10-15	>15	




Fig. 7.4 Land Suitability map of Sunflower


### 7.5 Land Suitability for Cotton (*Gossypium hirsutum*)

Cotton is the most important fibre crop grown in the State in about 6.6 lakh ha area in Raichur, Dharwad, Belgaum, Kalaburgi, Bijapur, Bidar, Bellary, Chitradurga and Chamarajnar districts. The crop requirements for growing cotton (Table 7.6) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing cotton was generated. The area extent and geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.5.

Highly suitable (class S1) lands are found to occur in minor area of 24 ha (5%) and distributed in the central part of the microwatershed. Moderately suitable (class S2) lands are found to occur in an area of about 165 ha (33%). The soils have moderate limitations of erosion and rooting depth. They are dominantly distributed in the southwestern, northwestern and central part of the microwatershed. The marginally suitable (class S3) lands cover about 139 ha (28%) area and mainly occur in the northeastern, northern and western part of the microwatershed. They have severe limitations of rooting depth and erosion. An area of about 138 ha (27%) is not suitable for growing cotton and is distributed in the central and northeastern part of the microwatershed.

**Table 7.6 Crop suitability criteria for Cotton**

Crop requirement		Rating			
Soil–site characteristics	unit	Highly suitable (S1)	Moderately Suitable (S2)	Marginally suitable (S3)	Not suitable (N)
Slope	%	1-2	2-3	3-5	>5
LGP	Days	180-240	120-180	<120	
Soil drainage	class	Well to moderately well	imperfectly drained	Poor somewhat excessive	Stagnant/excessive
Soil reaction	pH	6.5-7.5	7.6-8.0	8.1-9.0	>9.0 >6.5
Surface soil texture	Class	Sic, c	Sicl, cl	Si, sil, sc, scl, l	Sl, s,ls
Soil depth	cm	100-150	60-100	30-60	<30
Gravel content	% vol.	<5	5-10	10-15	15-35
CaCO <sub>3</sub> in root zone	%	<3	3-5	5-10	10-20
Salinity (EC)	dSm <sup>-1</sup>	2-4	4.0-8.0	8.0-12	>12
Sodicity (ESP)	%	5-10	10-20	20-30	>30



**Fig. 7.5 Land Suitability map of Cotton**

## 7.6 Land Suitability for Sugarcane (*Saccharum officinarum*)

Sugarcane is the most important commercial crop grown in 6.7 lakh ha area in Kalaburgi, Bijapur, Bagalkot, Bidar, Mysore, Chamarajanagar and Mandya districts under irrigated condition. The crop requirements for growing sugarcane (Table 7.7) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing sugarcane was generated. The area extent and geographical distribution of different suitability subclasses in the microwatershed is given in figure 7.6.

The moderately suitable (class S2) lands cover about 140 ha (28%) area in the microwatershed and mainly occur in the northwestern, central and southeastern part of the microwatershed. They have severe limitations of rooting depth and erosion. The marginally suitable (class S3) lands cover about 49 ha (10%) area and mainly occur in the central part of the microwatershed. They have severe limitations of texture. Major area of about 277 ha (55%) is not suitable for growing sugarcane and occur in all parts of the microwatershed.

**Table 7.7 Crop suitability criteria for Sugarcane**

Crop requirement		Rating			
Soil-site characteristics	unit	Highly suitable (S1)	Moderately Suitable (S2)	Marginally suitable (S3)	Not suitable (N)
Slope	%	<3	3-5	5-8	>8
Soil drainage	class	Well drained	Mod./imperfectly drained	Poorly drained	V.poor/excessively drained
Soil reaction	pH	7.0-8.0	6.0-6.9 8.1-9.0	4.0-5.9 9.1-9.5	<4.0/ >9.5
Surface soil texture	Class	l, cl, sil, sicl	C(m/k), sl	C+(ss)	
Soil depth	cm	>100	100-75	75-50	<50
stoniness	%	<15	15-35	35-50	>50
Salinity (EC)	dSm <sup>-1</sup>	<2.0	2.0-4.0	4.0-9.0	>9
Sodicity (ESP)	%	<10	10-15	15-25	>25




Fig. 7.6 Land Suitability map of Sugarcane

### 7.7 Land Suitability for Soybean (*Glycine max*)

Soybean is the most important pulse and oil seed crop grown in about 1.68 lakh ha area in Bijapur, Raichur, Kalaburgi, Dharwad, Belgaum and Bellary districts. The crop requirements for growing soybean were matched with the soil-site characteristics and a land suitability map for growing soybean was generated. The area extent and geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.7.

Highly suitable (class S1) lands are found to occur in a small area of 24 ha (5%) and are distributed in the central part of the microwatershed. Moderately suitable (class S2) lands are found to occur in an area of about 165 ha (33%). The soils have moderate limitations of erosion and rooting depth. They are dominantly distributed in the central, northwestern and southeastern part of the microwatershed. The marginally suitable (class S3) lands cover about 139 ha (28%) area and mainly occur in the southern, western, northern, and northwestern part of the microwatershed. They have severe limitations of rooting depth and erosion. An area of about 138 ha (37%) is not suitable for growing soybean and occur in the central, southwestern, northeastern and eastern part of the microwatershed.




Fig. 7.7 Land Suitability map of Soybean

## 7.8 Land Suitability for Guava (*Psidium guajava*)


Guava is the most important fruit crop grown in the State in Raichur, Dharwad, Belgaum, Kalaburgi, Bijapur, Bidar, Bellary, Chitradurga and Chamarajnagar districts. The crop requirements for growing guava (Table 7.8) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing guava was generated. The area extent and geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.8.

In Nirgudi West watershed, there are no highly (class S1) and moderately suitable (class S2) lands for growing guava. The marginally suitable (class S3) lands cover an area of about 189 ha (37%) and mainly occur in the central, southern, southwestern, southeastern and northwestern part of the microwatershed. They have severe limitations of texture and rooting depth. Major area of about 277 ha (55%) is not suitable for growing guava and occur in all parts of the microwatershed.



**Table 7.8 Crop suitability criteria for Guava**

Crop requirement			Rating			
Soil –site characteristics		unit	Highly suitable (S1)	Moderately Suitable (S2)	Marginally suitable (S3)	Not suitable (N)
climate	Temperature in growing season	<sup>0</sup> C	28-32	33-36 24-27	37-42 20-23	
Soil moisture	Growing period	Days	>150	120-150	90-120	<90
Soil aeration	Soil drainage	class	Well drained	Mod. to imperfectly	poor	Very poor
Nutrient availability	Texture	Class	Scl, l, cl, sil	Sl,sicl,sic.,sc,c	C (<60%)	C (>60%)
	pH	1:2.5	6.0-7.5	7.6-8.0:5.0-5.9	8.1-8.5:4.5-4.9	>8.5:<4.5
	CaCO <sub>3</sub> in root zone	%	Non calcareous	<10	10-15	>15
Rooting conditions	Soil depth	cm	>100	75-100	50-75	<50
	Gravel content	% vol.	<15	15-35	>35	
Soil toxicity	Salinity	dS/m	<2.0	2.0-4.0	4.0-6.0	
	Sodicity	%	Non sodic	10-15	15-25	>25
Erosion	Slope	%	<3	3-5	5-10	>10



**Fig 7.8 Land Suitability map of Guava**

## 7.9 Land Suitability for Mango (*Mangifera indica*)

Mango is the most important fruit crop grown in all the districts of the State. The crop requirements for growing mango (Table 7.9) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing mango was generated. The area and geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.9.

In Nirgudi West watershed, there are no highly (class S1) and moderately suitable (class S2) lands for growing mango. The marginally suitable (class S3) lands cover an area of about 49 ha (10%) and mainly occur in the central part of the microwatershed. They have severe limitations of texture. Major area of about 417 ha (83%) is not suitable for growing mango and occur in all parts of the microwatershed.

**Table 7.9 Crop suitability criteria for Mango**

Crop requirement			Rating			
soil-site characteristics	unit	Highly suitable (S1)	Moderately Suitable (S2)	Marginally suitable (S3)	Not suitable (N)	
climate	Temp in growing season	<sup>0</sup> C	28-32	24-27 33-35	36-40	20-24
	Min. temp. before flowering	<sup>0</sup> C	10-15	15-22	>22	
Soil moisture	Growing period	Days	>180	150-180	120-150	<120
Soil aeration	Soil drainage	class	Well drained	Mod. To imperfectly drained	Poor drained	Very poorly drained
	Water table	M	>3	2.50-3.0	2.5-1.5	<1.5
Nutrient availability	Texture	Class	Sc, l, sil, cl	Sl, sc, sic, l, c	C (<60%)	C (>60%),
	pH	1:2.5	5.5-7.5	7.6-8.5 5.0-5.4	8.6-9.0 4.0-4.9	>9.0 <4.0
	OC	%	High	medium	low	
	CaCO <sub>3</sub> in root zone	%	Non calcareous	<5	5-10	>10
Rooting conditions	Soil depth	cm	>200	125-200	75-125	<75
	Gravel content	% vol	Non gravelly	<15	15-35	>35
Soil toxicity	Salinity	dS/m	Non saline	<2.0	2.0-3.0	>3.0
	Sodicity	%	Non sodic	<10	10-15	>15
Erosion	Slope	%	<3	3-5	5-10	




Fig. 7.9 Land Suitability map of Mango

### 7.10 Land Suitability for Sapota (*Manilkara zapota*)

Sapota is the most important fruit crop grown in almost all the districts of the state. The crop requirements for growing sapota (Table 7.10) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing sapota was generated. The area extent and geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.10.

No highly (Class S1) and moderately (Class S2) suitable lands are available for growing sapota in the microwatershed. The marginally suitable (class S3) lands cover an area of about 189 ha (37%) and mainly occur in the central, southern, southeastern, western and northwestern part of the microwatershed. They have severe limitations of rooting depth and texture. Major area of about 277 ha (55%) is not suitable for growing sapota and occur in all parts of the microwatershed.

**Table 7.10 Crop suitability criteria for Sapota**

Crop requirement			Rating			
Soil –site characteristics		unit	Highly suitable (S1)	Moderately Suitable (S2)	Marginally suitable (S3)	Not suitable (N)
climate	Temperature in growing season	° C	28-32	33-36 24-27	37-42 20-23	>42 <18
			Soil moisture	Growing period	Days	>150
Soil aeration	Soil drainage	class	Well drained	Moderately well drained	Imperfectly drained	Poorly drained
Nutrient availability	Texture	Class	Scl, l, cl, sil	S1, sicl, sc	C (<60%)	ls, s, C (>60%)
	pH	1:2.5	6.0-7.5	7.6-8.0 5.0-5.9	8.1-9.0 4.5-4.9	>9.0 <4.5
	CaCO <sub>3</sub> in root zone	%	Non calcareous	<10	10-15	>15
Rooting conditions	Soil depth	cm	>150	75-150	50-75	<50
	Gravel content	% vol.	Non gravelly	<15	15-35	<35
Soil toxicity	Salinity	dS/m	Non saline	Up to 1.0	1.0-2.0	2.0-4.0
	Sodicity	%	Non sodic	10-15	15-25	>25
Erosion	Slope	%	<3	3-5	5-10	>10




Fig. 7.10 Land Suitability map of Sapota

### 7.11 Land Suitability for Jackfruit (*Artocarpus heterophyllus*)

Jackfruit is the most important fruit crop grown in southern and western districts of the state. The crop requirements for growing jackfruit were matched with the soil-site characteristics and a land suitability map for growing jackfruit was generated. The area extent and geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.11.

In Nirgudi West microwatershed, there are no highly (Class S1) and moderately (Class S2) suitable lands for growing jackfruit. The marginally suitable (class S3) lands cover an area of about 189 ha (37%) and mainly occur in the southern, southwestern, northwestern and southeastern part of the microwatershed. They have severe limitations of texture and rooting depth. Major area of about 277 ha (55%) is not suitable for growing jackfruit and occur in all parts of the microwatershed.




Fig 7.11 Land Suitability map of Jackfruit

### 7.12 Land Suitability for Jamun (*Syzygium cumini*)

Jamun is the most important fruit crop grown in almost all the districts of the state. The crop requirements for growing jamun were matched with the soil-site characteristics and a land suitability map for growing jamun was generated. The area extent and geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.12.

The moderately suitable (class S2) lands are found to occur in a small area of about 40 ha (8%). The soils have moderate limitations of texture. They are dominantly distributed in central part of the microwatershed. The marginally suitable (class S3) lands cover about an area of 149 ha (30%) and mainly occur in the central, southeastern, southwestern and northwestern part of the microwatershed. They have severe limitations of rooting depth. Major area of about 277 ha (55%) is not suitable for growing jamun and occur in all parts of the microwatershed.




Fig 7.12 Land Suitability map of Jamun

### 7.13 Land Suitability for Musambi (*Citrus limetta*)

Musambi is the most important fruit crop grown in almost all the districts of the state. The crop requirements for growing musambi were matched with the soil-site characteristics and a land suitability map for growing musambi was generated. The area extent and geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.13.

Highly suitable (class S1) lands are found to occur in a small area of about 24 ha (5%) and are distributed in the central part of the microwatershed. The moderately suitable (class S2) lands are found to occur in a small area of about 25 ha (5%). The soils have moderate limitations of erosion and rooting depth. They are dominantly distributed in the central part of the microwatershed. The marginally suitable (class S3) lands cover an area of about 140 ha (28%) and mainly occur in the central, southern, southeastern, southwestern and northwestern part of the microwatershed. They have severe limitations of rooting depth. Major area of about 277 ha (55%) is not suitable for growing musambi and occur in all parts of the microwatershed.




Fig 7.13 Land Suitability map of Musambi

#### 7.14 Land Suitability for Lime (*Citrus sp*)


Lime is the most important fruit crop grown in almost all the districts of the state. The crop requirements for growing lime (Table 7.11) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing lime was generated. The area extent and geographic distribution of different suitability subclasses in the microwatershed is given in Figure 7.14.

Highly suitable (class S1) lands are found to occur in a small area of about 24 ha (5%) and are distributed in the central part of the microwatershed. The moderately suitable (class S2) lands are found to occur in a small area of about 25 ha (5%). The soils have moderate limitations of erosion and rooting depth. They are dominantly distributed in the central part of the microwatershed. The marginally suitable (class S3) lands cover an area of about 140 ha (28%) and mainly occur in the central, southern, southeastern, southwestern and northwestern part of the microwatershed. They have severe limitations of rooting depth. Major area of about 277 ha (55%) is not suitable for growing lime and occur in all parts of the microwatershed.



**Table 7.11 Crop suitability criteria for Lime**

Crop requirement			Rating			
Soil –site characteristics		unit	Highly suitable (S1)	Moderately Suitable (S2)	Marginally suitable (S3)	Not suitable (N)
Climate	Temp in growing season	<sup>0</sup> C	28-30	31-35 24-27	36-40 20-23	>40 <20
Soil moisture	Growing period	Days	240-265	180-240	150-180	<150
Soil aeration	Soil drainage	class	Well drained	Mod. to imperfectly drained	poorly	Very poorly
Nutrient availability	Texture	Class	Scl, l, sicl, cl, s	Sc, sc, c	C (>70%)	S, ls
	pH	1:2.5	6.0-7.5	5.5-6.4/ 7.6-8.0	4.0-5.4 8.1-8.5	<4.0 >8.5
	CaCO <sub>3</sub> in root zone	%	Non-calcareous	Upto 5	5-10	>10
Rooting condition	Soil depth	cm	>150	100-150	50-100	<50
	Gravel content	% vol.	Non gravelly	15-35	35-55	>55
Soil toxicity	Salinity	dS/m	Non saline	Upto 1.0	1.0-2.5	>2.5
	Sodicity	%	Non sodic	5-10	10-15	>15
Erosion	Slope	%	<3	3-5	5-10	



**Fig 7.14 Land Suitability map of Lime**

### 7.15 Land Suitability for Cashew (*Anacardium occidentale*)

Cashew is the most important plantation crop grown mostly in coastal and western part and also in Bidar and Kolar Districts. The crop requirements for growing Cashew were matched with the soil-site characteristics and a land suitability map for growing Cashew was generated. The area and geographic distribution of different suitability subclasses in the microwatershed is given in Figure 7.15.

The entire area is not suitable for growing cashew in the microwatershed.




Fig 7.15 Land Suitability map of Cashew

### 7.16 Land Suitability for Custard Apple (*Annona reticulata*)

Custard apple is the most important fruit crop grown in almost all the districts of the state. The crop requirements for growing custard apple were matched with the soil-site characteristics and a land suitability map for growing custard apple was generated. The area extent and geographic distribution of different suitability subclasses in the microwatershed is given in Figure 7.16.

Highly suitable (class S1) lands are found to occur in a small area of 24 ha (5%) and are distributed in the central part of the microwatershed. Moderately suitable (class S2) lands are found to occur in an area of about 165 ha (33%). The soils have moderate limitations of erosion and rooting depth. They are dominantly distributed in the southern, central, southeastern, southwestern and northwestern part of the microwatershed. The marginally suitable (class S3) lands cover about 139 ha (28%) area and mainly occur in the northern,

northeastern, southern, central and western part of the microwatershed. They have severe limitations of rooting depth and erosion. An area of about 138 ha (27%) is not suitable for growing custard apple and occur in the central, northern and eastern part of the microwatershed.




Fig 7.16 Land Suitability map of Custard Apple

### 7.17 Land Suitability for Amla (*Phyllanthus emblica*)

Amla is the most important fruit crop grown in almost all the districts of the state. The crop requirements for growing amla were matched with the soil-site characteristics and a land suitability map for growing amla was generated. The area extent and geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.17.

Highly suitable (class S1) lands are found to occur in a small area of 24 ha (5%) and are distributed in the central part of the microwatershed. Moderately suitable (class S2) lands are found to occur in an area of about 165 ha (33%). The soils have moderate limitations of erosion and rooting depth. They are dominantly distributed in the southern, central, southeastern, southwestern and northwestern part of the microwatershed. The marginally suitable (class S3) lands cover about 139 ha (28%) area and occur in the northern, northeastern, western and southern part of the microwatershed. They have severe limitations of rooting depth. An area of about 138 ha (27%) is not suitable for growing amla and occur in all parts of the microwatershed.




Fig 7.17 Land Suitability map of Amla

### 7.18 Land Suitability for Tamarind (*Tamarindus indica*)

Tamarind is the most important spice crop raised in almost all the districts of the state. The crop requirements for growing tamarind were matched with the soil-site characteristics and a land suitability map for growing tamarind was generated. The area extent and geographic distribution of different suitability subclasses in the microwatershed is given in Figure 7.18.

Highly suitable (class S1) lands are found to occur in a small area of 24 ha (5%) and are distributed in the central part of the microwatershed. Moderately suitable (class S2) lands are found to occur in a very small area of about 16 ha (3%). The soils have moderate limitations of erosion. They are distributed in the central part of the microwatershed. Marginally suitable (class S3) lands are found to occur in a very small area of about 9 ha (2%). The soils have moderate limitations of rooting depth. They are distributed in the central part of the microwatershed. Major area of about 417 ha (83%) is not suitable for growing tamarind and occurs in all parts of the microwatershed.




Fig 7.18 Land Suitability map of Tamarind

### 7.19 Land Management Units (LMUs)

The 20 soil map units identified in Nirgudi West microwatershed have been regrouped into 5 Land Management Units (LMU's) for the purpose of preparing Proposed Crop Plan. Land Management Units are grouped based on the similarities in respect of the type of soil, the depth of the soil, the surface soil texture, gravel content, AWC, slope, erosion etc. and a Land Management Units map (Fig.7.19) has been prepared. These Land Management Units are expected to behave similarly for a given level of management.

The map units that have been grouped into 5 land management units along with brief description of soil and site characteristics are given below:

LMUs	Soil map units	Soil and site characteristics
1	MGTmC3g1, MGTmC3g2, MGTmD3g2, NHAmD3g2	Very shallow, black soils with slopes of 5-10%, gravelly to very gravelly (15-60%) and severe erosion
2	MGTmB2g1	Very shallow, black soils with slopes of 1-3%, gravelly (15-35%) and moderate erosion
3	NHAmB1, NHAmB2, NHAmB2g1, NHAmC2g1, NHAmC3, BHImB2g1	Shallow, black soils with slopes of 1-5 %, non gravelly to gravelly (<15-35%) and slight to moderate erosion
4	DSImA1, DSImB2g1, DSImC2g1, KMPmB2g1	Moderately shallow, black soils with slopes of <1-5 %, non-gravelly to gravelly (<15-35%) and slight to moderate erosion
5	RNLmA1, RNLmB1g1, RNLmB2	Deep, black soils with slopes of <1-3%, non-gravelly to gravelly (<15-35%) and slight to moderate erosion




Fig. 7.19 Land Management Units map of Nirgudi West Microwatershed

## 7.20 Proposed Crop Plan for Nirgudi West Microwatershed

After assessing the land suitability for the 18 crops, a proposed crop plan has been prepared for the 5 identified LMUs by considering only the highly (class S1) and moderately (class S2) suitable lands for each of the eighteen crops. The resultant proposed crop plan is presented below in Table 7.12.

**Table 7.12 Proposed Crop Plan for Nirgudi West Microwatershed**

LMU	Mapping unit	Survey No	Crops proposed				Suitable Intervention
			Field Crops	Forestry/ Grasses	Horticulture Crops (Rainfed Condition)	Horticulture Crops with suitable Intervention	
LMU-1	MGTmC3g1, MGTmC3g2, MGTmD3g2, NHAmD3g2	<b>Nirgudi:</b> 4,5,6,9,15,16, 279,285,286,287,300, 304,313,316,317	-	Neem, Glyricydia , Silviculture, Agave, Simaroba	-	-	Crescent bunds
LMU-2	MGTmB2g1	<b>Nirgudi:</b> 2,3	Horse gram,	Neem, Glyricydia, Silviculture, Agave, Simaroba	-	-	Crescent bunds
LMU-3	NHAmB1, NHAmB2, NHAmB2g1, NHAmC2g1, NHAmC3, BHImB2g1	<b>Nirgudi:</b> 10,11,12,13, 14,17,18,224,225,231, 232,233,256,257,258, 283,284,291,311,312, 314,315,328	Bajra, Linseed, Green gram, Black gram, Chick pea, Coriander	Subabul, Neem, Teak	Custard apple, Charoli, Ber, Amla <b>Vegetables:</b> Ladies finger, Brinjal, Cowpea, <b>Flowers:</b> Marigold, Chrysanthemum	Custard apple, Charoli, Ber, Amla <b>Vegetables:</b> Onion, Tomato, Brinjal, Chillies, Bhendi <b>Flowers:</b> Marigold, Chrysanthemum	Drip irrigation, suitable soil and water conservations measures like cultivation on raised beds with mulches and drip

LMU-4	DSImA1, DSImB2g1, DSImC2g1, KMPmB2g1	<b>Nirgudi:</b> 206,207,208, 210,211,212,213,214, 215,216,217,218,219, 220,221,222,226,227, 228,229,230,264,265, 267,268,269,270,271, 272,273,274,275,276, 277,278,281,282,288, 289,290,292,293,294, 295,296,297,298,301, 302,303,305,306,307, 308,309,310, 318	Sorghum, Cotton, Red Gram, Black gram, Green gram, Soybean, Sesame, Sunflower, <b>Rabi:</b> Sorghum, Chickpea, Coriander, Safflower	Subabul, Neem, Teak	Custard apple, Charoli, Ber, Amla <b>Vegetables:</b> Ladies finger, Brinjal, Cowpea, <b>Flowers:</b> Marigold, Chrysanthemum	Custard apple, Charoli, Ber, Amla, Papaya, Banana, Lime, Citrus <b>Vegetables:</b> Onion, Tomato, Brinjal, Chillies, Bhendi <b>Flowers:</b> Marigold, Chrysanthemum	-do- Graded bunds, Strengthening of field bunds
LMU-5	RNLmA1, RNLmB1g1, RNLmB2	Nirgudi: 7,8,266,280,299,319, 320,321,322,323,324, 325,326,327	Sorghum, Cotton, Red Gram Black gram, Green gram, Soybean, Sesame, Sunflower, <b>Rabi:</b> Sorghum, Chickpea, Safflower, Coriander, Linseed	-	<b>Vegetables:</b> Ladies finger, Brinjal, Cowpea, Coriander <b>Field crops:</b> Sorghum, Cotton, Red Gram, Sunflower, Safflower, <b>Perennial component:</b> Guava, Sapota, Lime, Musambi, Tamarind <b>Flowers:</b> Marigold, Chrysanthemum	Banana, Papaya, Lime, Musambi, Guava, Tamrind <b>Vegetables:</b> Onion, Tomato, Brinjal, Chillies, Bhendi <b>Flowers:</b> Marigold, Chrysanthemum	-do- Graded bunds, Strengthening of field bunds



## SOIL HEALTH MANAGEMENT

### 8.1 Soil Health

Soil is fundamental to crop production. Without soil, no food could be produced nor would livestock be fed on a large scale. Because it is finite and fragile, soil is a precious resource that requires special care from its users.

Soil health or the capacity of the soil to function is critical to human survival. Soil health has been defined as: “the capacity of the soil to function as a living system without adverse effect on the ecosystem”. Healthy soils maintain a diverse community of soil organisms that help to form beneficial symbiotic associations with plant roots, recycle essential plant nutrients, improve soil structure with positive repercussions for soil, water and nutrient holding capacity and ultimately improve crop production and also contribute to mitigating climate change by maintaining or increasing its carbon content.

Functional interactions of soil biota with organic and inorganic components, air and water determine a soil’s potential to store and release nutrients and water to plants and to promote and sustain plant growth. Thus, maintaining soil health is vital to crop production and conserve soil resource base for sustaining agriculture.

#### **The most important characteristics of a healthy soil are**

- Good soil tilth
- Sufficient soil depth
- Good water storage and good drainage
- Adequate supply, but not excess of nutrients
- Large population of beneficial organisms
- Small proportion of plant pathogens and insect pests
- Low weed pressure
- Free of chemicals and toxins that may harm the crop
- Resistance to degradation
- Resilience when unfavourable conditions occur

#### **Characteristics of Nirgudi West Microwatershed**

- The soil phases with sizeable area identified in the microwatershed belonged to the soil series of NHA (140 ha), DSI (140 ha), MGT (116 ha), RNL (40 ha), BHI (20 ha) and KMP (9 ha).
- As per land capability classification, nearly 86 per cent area falls under arable land category (Class II, III and IV) and 12 per cent area belongs to nonarable land (Class VI) category. The major limitations identified in the arable lands were soil and erosion.
- On the basis of soil reaction, about 205 ha (41%) area is neutral in reaction (pH 6.5-7.3) followed by slightly alkaline (pH 7.3-7.8) 142 ha (28%) and moderately alkaline (pH 8.4-9.0) 120 ha (24%). Thus, about 52 per cent of the soils in the microwatershed are alkaline in reaction.

## **Soil Health Management**

The following actions are required to improve the current land husbandry practices that provide a sound basis for the successful adoption of sustainable crop production system

### **Alkaline soils**

(Slightly alkaline to moderately alkaline soils)

1. Regular addition of organic manure, green manuring, green leaf manuring, crop residue incorporation and mulching needs to be taken up to improve the soil organic matter status.
2. Application of biofertilizers (Azospirillum, Azotobacter, Rhizobium).
3. Application of 25% extra N and P (125 % RDN&P).
4. Application of ZnSO<sub>4</sub> – 12.5 kg/ha (once in three years).
5. Application of Boron – 5kg/ha (once in three years).

### **Neutral soils**

1. Regular addition of organic manure, green manuring, green leaf manuring, crop residue incorporation and mulching needs to be taken up to improve the soil organic matter status.
2. Application of biofertilizers, (Azospirillum, Azotobacter, Rhizobium).
3. Application of 100 per cent RDF.
4. Need based micronutrient applications.

Besides the above recommendations, the best transfer of technology options are also to be adopted.

## **Soil Degradation**

Soil erosion is one of the major factor affecting the soil health in the microwatershed. Out of total area of 466 ha in the microwatershed, major area of 383 ha is suffering from either moderate or severe erosion. These areas need immediate soil and water conservation and other land development, and land husbandry practices for restoring soil health.

## **Disseminate information and communicate benefits**

Any large scale implementation of soil health management requires that supporting information is made available widely, particularly through channels familiar to farmers and extension workers. Given the very high priority attached to soil health especially by the Central Government on issuing Soil-Health Cards to all the farmers, media outlets like regional, state and national newspapers, radio and dooradarshan programs in local languages but also modern information and communication technologies such as cellular phones and the Internet, which can be much more effective in reaching younger farmers.

## Inputs for Net Planning and Interventions needed

Net planning in IWMP is focusing on preparation of

1. Soil and Water Conservation Plan for each plot or farm.
2. Productivity enhancement measures/ interventions for existing crops/livestock/other farm enterprises.
3. Diversification of farming mainly with perennial horticultural crops and livestock.
4. Improving livelihood opportunities and income generating activities.

In this connection, how various outputs of Sujala-III are of use in addressing these objectives of Net Planning are briefly presented below:

- ❖ **Soil Depth:** The depth of a soil decides the amount of moisture and nutrients it can hold, what crops can be taken up or not, depending on the rooting depth and the length of growing period available for raising any crop. Deeper the soil, better for a wide variety of crops. If sufficient depth is not available for growing deep rooted crops either choose medium or short duration crops or deeper planting pits need to be opened and additional good quality soil brought from outside has to be filled into the planting pits.
- ❖ **Surface soil texture:** Lighter soil texture in the top soil means, better rain water infiltration, less run-off and soil moisture conservation, less capillary rise and less evaporation losses. Lighter surface textured soils are amenable to good soil tilth and are highly suitable for crops like groundnut, root vegetables (carrot, radish, potato etc) but not ideal for crops that need stagnant water like lowland paddy. Heavy textured soils are poor in water infiltration and percolation. They are prone for sheet erosion; such soils can be improved by sand mulching. The technology that is developed by the AICRP-Dryland Agriculture, Vijayapura, Karnataka may be adopted.
- ❖ **Gravelliness:** More gravel content is favorable for run-off harvesting but poor in soil moisture storage and nutrient availability. It is a significant parameter that decides the kind of crop to be raised.
- ❖ **Land Capability Classification:** The land capability map shows the areas suitable and not suitable for agriculture and the major constraints in each of the plot/survey number. Hence, one can decide what kind of enterprise is possible in each of these units. In general, erosion and soil are the major constraints in Nirgudi West microwatershed.
- ❖ **Organic Carbon:** In about 20 ha (4%) area, the OC content is low (<0.5%), about 161 ha (32%) area, the OC content is medium (0.5-0.75%) and in about 286 ha (57%) area it is high (>0.75%). The areas that are medium to low in OC needs to be further improved by applying farmyard manure and rotating crops with cereals and legumes or mixed cropping.


- ❖ **Promoting green manuring:** Growing of green manuring crops costs Rs. 1250/ha (green manuring seeds) and about Rs. 2000/ha towards cultivation that totals to Rs. 3250/- per ha. On the other hand, application of organic manure @ 10 tons/ha costs Rs. 5000/ha. The practice needs to be continued for 2-3 years or more. Nitrogen fertilizer needs to be supplemented by 25% in addition to the recommended level in 20 ha area where OC is less than 0.5-0.75%. For example, for rainfed maize, recommended level is 50 kg N per ha and an additional 12 kg /ha needs to be applied for all the crops grown in these plots.
- ❖ **Available Phosphorus:** In 383 ha (76%) area, the available phosphorus is low and about 78 ha (15%) area it is medium in available phosphorus, Hence for all the crops, 25% additional P-needs to be applied.
- ❖ **Available Potassium:** Available potassium is medium in 126 ha (25%) area of the microwatershed. Hence, in all these plots, for all crops, an additional 25 % potassium may be applied. It is high in 341 ha (67%) area of the microwatershed.
- ❖ **Available Sulphur:** Available sulphur is a very critical nutrient for oilseed crops. It is low in 6 ha (1%) area of the microwatershed and medium in 459 ha (91%). These areas need to be applied with magnesium sulphate or gypsum or Factamphos (p) fertilizer (13% sulphur) for 2-3 years for the deficiency to be corrected. Only 2 ha area has soils that are high in available sulphur.
- ❖ **Available iron:** It is deficient in 11 ha (2%) area of the microwatershed. To manage iron deficiency, iron sulphate @ 25kg /ha needs to be applied for 2-3 years. It is sufficient in the rest of 455 ha (90 %) area in the microwatershed.
- ❖ **Available Zinc:** It is deficient in 165 ha (33%) area of the microwatershed. Application of zinc sulphate @25kg/ha is to be applied. It is sufficient in 302 ha (60%) area in the microwatershed.
- ❖ **Soil alkalinity:** The microwatershed has 444 ha area with soils that are moderately to strongly alkaline. These areas need application of gypsum and wherever calcium is in excess, iron pyrites and element sulphur can be recommended. Management practices like treating repeatedly with good quality water to drain out the excess salts and provision of subsurface drainage and growing of salt tolerant crops like Casuarina, Acasia, Neem, Ber et, are recommended.

**Land Suitability for various crops:** Areas that are highly, moderately and marginally suitable for growing various crops are indicated. Along with the suitability, various constraints that are limiting the productivity are also indicated. For example, in case of cotton, gravel content, rooting depth and salinity/alkalinity are the major constraints in various plots. With suitable management interventions, the productivity can be enhanced. In order to increase water holding capacity of light textured soils, growing of green manure crops and application of organic manure is recommended.

## SOIL AND WATER CONSERVATION TREATMENT PLAN

For preparing soil and water conservation treatment plan for Nirgudi West microwatershed, the land resource inventory database generated under Sujala-III project has been transformed as information through series of interpretative (thematic) maps using soil phase map as a base. The various thematic maps (1:7920 scale) generated were

- Soil depth
- Surface soil texture
- Available water capacity
- Soil slope
- Soil gravelliness
- Land capability
- Present land use and land cover
- Crop suitability maps
- Rainfall map
- Hydrology
- Water Resources
- Socio-economic data
- Contour plan with existing features- Network of waterways, pothissa boundaries, cut up/ minor terraces etc.
- Cadastral map (1:7920 scale)
- Satellite imagery (1:7920 scale)



Apart from these, Hand Level/ Hydro Marker/ Dumpy Level/ Total Station and Kathedars' List has to be collected.

### Steps for Survey and Preparation of Treatment Plan

The boundaries of Land user Groups' and Survey No. boundaries are traced in the field.

Naming of user groups and farmers

- Identification of arable and non arable lands
- Identification of drainage lines and gullies
- Identification of non treatable areas
- Identification of priority areas in the arable lands
- Treatment plan for arable lands
- Location of water harvesting and recharge structures

## 9.1 Treatment Plan

The treatment plan recommended for arable lands is briefly described below


### 9.1.1 Arable Land Treatment

#### A. BUNDING

Steps for Survey and Preparation of Treatment Plan		<p><b>USER GROUP-1</b></p> <p><b>CLASSIFICATION OF GULLIES</b></p> <p>ಕೊರಕಲಿನ ವರ್ಗೀಕರಣ</p> <ul style="list-style-type: none"> <li>ಮೇಲ್ಭಾಗ (UPPER REACH) 15 ಹಾ.</li> <li>ಮಧ್ಯಭಾಗ (MIDDLE REACH) 15+10=25 ಹಾ.</li> <li>ಕೆಳಭಾಗ (LOWER REACH) 25 ಹೆಕ್ಟಾರ್ ಗಿಂತ ಅಧಿಕ</li> </ul> <p>POINT OF CONCENTRATION</p>
Cadastral map (1:7920 scale) is enlarged to a scale of 1:2500 scale		
Existing network of waterways, pothissa boundaries, grass belts, natural drainage lines/ watercourse, cut ups/ terraces are marked on the cadastral map to the scale		
Drainage lines are demarcated into		
Small gullies	(up to 5 ha catchment)	
Medium gullies	(5-15 ha catchment)	
Ravines	(15-25 ha catchment) and	
Halla/Nala	(more than 25ha catchment)	

#### Measurement of Land Slope

Land slope is estimated or determined by the study and interpretation of contours or by measurement in the field using simple instruments like Hand level or Hydromarker.



Vertical and Horizontal intervals between bunds as recommended by the Watershed Development Department

Slope percentage	Vertical interval (m)	Corresponding Horizontal Distance (m)
2 - 3%	0.6	24
3 - 4%	0.9	21
4 - 5%	0.9	21
5 - 6%	1.2	21
6 - 7%	1.2	21

**Note:** (i) The above intervals are maximum.

(ii) Considering the slope class and erosion status (A1....) the intervals have to be decided.

**Bund length recording:** Considering the contour plan and the existing grass belts/partitions, the bunds are aligned and lengths are measured.

**Section of the Bund**

Bund section is decided considering the soil texture class and gravelliness class (bg<sub>0</sub>, loamy sand, <15% gravel). The recommended Sections for different soils are given below.


**Recommended Bund Section**

Top width (m)	Base width (m)	Height (m)	Side slope (Z:1;H:V)	Cross section (sq m)	Soil Texture	Remarks
0.3	0.9	0.3	01:01	0.18	Sandy loam	Vegetative bund
0.3	1.2	0.3	1.5:1	0.225	Sandy clay	
0.3	1.2	0.5	0.9:1	0.375	Red gravelly soil	
0.3	1.2	0.6	0.75:1	0.45		
0.3	1.5	0.6	01:01	0.54	Red sandy loam	
0.3	2.1	0.6	1.5:1	0.72	Very shallow black soils	
0.45	2	0.75	01:01	0.92		
0.45	2.4	0.75	1.3:1	1.07	Shallow black soils	
0.6	3.1	0.7	1.78:1	1.29	Medium black soils	
0.5	3	0.85	1.47:1	1.49		

**Formation of Trench cum Bund**

Dimensions of the Borrow Pits/Trenches to be excavated (machinery are decided considering the Bund Section).

Details of Borrow Pit dimensions are given below:




**TRENCH CUM BUND**

WATER STORAGE AREA

0.45 Sq.m section

IDEAL FOR HORTICULTURE CROPS




0.30 m (1')

1.20 m (4')

0.60 m (2')

1.2 m



1. ಸಮಾನಾಕಳಿ ಉಳಿಸುವೆ

2. ಸಮಾನಾಕಳಿ ಬಿತ್ತನೆ/ನಾಟಿ

3. ಸಮಾನಾಕಳಿ ರೇಷುವೆ

**'A' FRAME FOR INTERBUND MANAGEMENT**

### Size of Borrow Pits/ Trench recommended for Trench cum Bund (by machinery)

Bund section	Bund length	Earth quantity	Pit				Berm (pit to pit)	Soil depth class
			L (m)	W (m)	D(m)	QUANTITY (m <sup>3</sup> )		
m <sup>2</sup>	m	m <sup>3</sup>	L (m)	W (m)	D(m)	QUANTITY (m <sup>3</sup> )	m	
0.375	6	2.25	5.85	0.85	0.45	2.24	0.15	Shallow
0.45	6	2.7	5.4	1.2	0.43	2.79	0.6	Shallow
0.45	6	2.7	5	0.85	0.65	2.76	1	Moderately Shallow
0.54	5.6	3.02	5.5	0.85	0.7	3.27	0.1	Moderately shallow
0.54	5.5	2.97	5	1.2	0.5	3	0.5	Shallow
0.72	6.2	4.46	6	1.2	0.7	5.04	0.2	Moderately shallow
0.72	5.2	3.74	5.1	0.85	0.9	3.9	0.1	Moderately deep

#### B. Water Ways

- Existing waterways are marked on the cadastral map (1:7920 scale) and their dimensions are recorded.
- Considering the contour plan of the MWS, additional water ways/ modernization of the existing ones can be thought of.
- The design details are given in the Manual.

#### C. Farm Ponds

Waterways and the catchment area will give an indication on the size of the Farm Pond. Location of the pond can be decided based on the contour plan/ field condition and farmers' need/desire.

#### D. Diversion Channel

Existing EPT/ CPT are marked on the cadastral map. Looking to the need, these can be modernized or fresh diversion channel can be proposed and runoff from this can be stored in Gokatte/ Recharge ponds.

#### 9.1.2 Non-Arable Land Treatment

Depending on the gravelliness and crops preferred by the farmers, the concerned authorities can decide appropriate treatment plan. The recommended treatments may be Contour Trench, Staggered Trench, Crescent Bund, Boulder Bund or Pebble Bunds are formed in the field.



### **9.1.3 Treatment of Natural Water Course/ Drainage Lines**

- a) The cadastral map has to be updated as regards the network of drainage lines (gullies/ nallas/ hallas) and existing structures are marked to the scale and storage capacity of the existing water bodies are documented.
- b) The drainage line will be demarcated into Upper Reach, Middle Reach and Lower Reach.
- c) Considering the Catchment, Nala bed and bank conditions, suitable structures are decided.
- d) Number of storage structures (Check dam/ Nala bund/ Percolation tank) will be decided considering the commitments and available runoff in water budgeting and quality of water in the wells and site suitability.
- e) Detailed Levelling Survey using Dumpy Level / Total Station has to be carried out to arrive at the site-specific designs as shown in the Manual.
- f) The location of ground water recharge structures are decided by examining the lineaments and fracture zones from geological maps.
- g) Rainfall intensity data of the nearest Rain gauge station is considered for Hydrologic Designs.
- h) Silt load to the Storage/Recharge structures is reduced by providing vegetative, boulder and earthen checks in the natural water course. Location and design details are given in the Manual.

### **9.2 Recommended Soil and Water Conservation Measures**

The appropriate conservation structures best suited for each of the land parcel/ survey number (Appendix-I) are selected based on the slope per cent, severity of erosion, amount of rainfall, land use and soil type. The different kinds of conservation structures recommended are:

1. Graded / Strengthening of bunds
2. Trench cum Bunds (TCB)
3. Trench cum Bunds / Strengthening
4. Crescent Bunds

A map (Fig. 9.1) showing soil and water conservation plan with different kinds of structures recommended has been prepared which shows the spatial distribution and extent of area. An area of about 277 ha (55%) requires crescent bund, 149 ha (30%) area requires strengthening of bunds/TCB/GB and about 40 ha (8%) area needs graded bunds/ strengthening of bunds.

The conservation plan prepared may be presented to all the stakeholders including farmers and after including their suggestions, the conservation plan for the microwatershed may be finalised in a participatory approach.




Fig. 9.1 Soil and Water Conservation Plan map of Nirgudi West Microwatershed

### 9.3 Greening of Microwatershed

As part of the greening programme in the watersheds, it is envisaged to plant a variety of horticultural and other tree plants that are edible, economical and produce lot of biomass which helps to restore the ecological balance in the watersheds. The lands that are suitable for greening programme are non-arable lands (land capability classes V, VI and VII) and also the lands that are not suitable or marginally suitable and field bunds for growing annual and perennial crops. The method of planting these trees is given below.

It is recommended to open pits during the 1<sup>st</sup> week of March along the contour and heap the dugout soil on the lower side of the slope in order to harness the flowing water and facilitate weathering of soil in the pit. Exposure of soil in the pit also prevents spread of pests and diseases due to scorching sun rays. The pits should be filled with mixture of soil and organic manure during the second week of April and keep ready with sufficiently tall seedlings produced either in poly bags or in root trainer nurseries so that planting can be done during the 2<sup>nd</sup> or 3<sup>rd</sup> week of April depending on the rainfall.

The tree species suitable for the area considering rainfall, temperature and adaptability is listed below; waterlogged areas are recommended to be planted with species like Neral (*Syzgium cumini*) and Bamboo. Dry areas are to be planted with species like Honge, Bevu, Seetaphal etc.

<b>Dry Deciduous Species</b>			<b>Temp (°C)</b>	<b>Rainfall (mm)</b>
1.	Bevu	<i>Azadiracta indica</i>	21-32	400 -1,200
2.	Tapasi	<i>Holoptelia integrifolia</i>	20-30	500 - 1000
3.	Seetaphal	<i>Anona Squamosa</i>	20-40	400 - 1000
4.	Honge	<i>Pongamia pinnata</i>	20 -50	500- 2,500
5.	Kamara	<i>Hardwickia binata</i>	25 -35	400 - 1000
6.	Bage	<i>Albezzia lebbek</i>	20 - 45	500 - 1000
7.	Ficus	<i>Ficus bengalensis</i>	20 - 50	500-2,500
8.	Sisso	<i>Dalbargia Sissoo</i>	20 - 50	500 -2000
9.	Ailanthus	<i>Ailanthus excelsa</i>	20 - 50	500 - 1000
10.	Hale	<i>Wrightia tinctoria</i>	25 - 45	500 - 1000
11.	Uded	<i>Steriospermum chelanoides</i>	25 - 45	500 -2000
12.	Dhupa	<i>Boswellia Serrata</i>	20 - 40	500 - 2000
13.	Nelli	<i>Emblica Officinalis</i>	20 - 50	500 -1500
14.	Honne	<i>Pterocarpus marsupium</i>	20 - 40	500 - 2000
<b>Moist Deciduous Species</b>			<b>Temp (°C)</b>	<b>Rainfall (mm)</b>
15.	Teak	<i>Tectona grandis</i>	20 - 50	500-5000
16.	Nandi	<i>Legarstroemia lanceolata</i>	20 - 40	500 - 4000
17.	Honne	<i>Pterocarpus marsupium</i>	20 - 40	500 - 3000
18.	Mathi	<i>Terminalia alata</i>	20 -50	500 - 2000
19.	Shivane	<i>Gmelina arboria</i>	20 -50	500 -2000
20.	Kindal	<i>T.Paniculata</i>	20 - 40	500 - 1500
21.	Beete	<i>Dalbargia latifolia</i>	20 - 40	500 - 1500
22.	Tare	<i>T. belerica</i>	20 - 40	500 - 2000
23.	Bamboo	<i>Bambusa arundinasia</i>	20 - 40	500 - 2500
24.	Bamboo	<i>Dendrocalamus strictus</i>	20 - 40	500 - 2500
25.	Muthuga	<i>Butea monosperma</i>	20 - 40	400 - 1500
26.	Hippe	<i>Madhuca latifolia</i>	20 - 40	500 - 2000
27.	Sandal	<i>Santalum album</i>	20 - 50	400 - 1000
28.	Nelli	<i>Emblica officinalis</i>	20 - 40	500 - 2000
29.	Nerale	<i>Sizygium cumini</i>	20 - 40	500 - 2000
30.	Dhaman	<i>Grevia tilifolia</i>	20 - 40	500 - 2000
31.	Kaval	<i>Careya arborea</i>	20 - 40	500 - 2000
32.	Harada	<i>Terminalia chebula</i>	20 - 40	500 - 2000



## References

1. FAO (1976) Framework for Land Evaluation, Food and Agriculture Organization, Rome.72 pp.
2. FAO (1983) Guidelines for Land Evaluation for Rainfed Agriculture, FAO, Rome, 237 pp.
3. IARI (1971) Soil Survey Manual, All India Soil and Land Use Survey Organization, IARI, New Delhi, 121 pp.
4. Katyal, J.C. and Rattan, R.K. (2003) Secondary and Micronutrients; Research Gap and Future Needs. Fert. News 48 (4); 9-20.
5. Naidu, L.G.K., Ramamurthy, V., Challa, O., Hegde, R. and Krishnan, P. (2006) Manual Soil Site Suitability Criteria for Major Crops, NBSS Publ. No. 129, NBSS & LUP, Nagpur, 118 pp.
6. Natarajan, A. and Dipak Sarkar (2010) Field Guide for Soil Survey, National Bureau of Soil Survey and Land Use Planning (ICAR), Nagpur, India.
7. Natarajan, A., Rajendra Hegde, Raj, J.N. and Shivananda Murthy, H.G. (2015) Implementation Manual for Sujala-III Project, Watershed Development Department, Bengaluru, Karnataka.
8. Sarma, V.A.K., Krishnan, P. and Budihal, S.L. (1987) Laboratory Manual, Tech. Bull. 23, NBSS &LUP, Nagpur.
9. Sehgal, J.L. (1990) Soil Resource Mapping of Different States of India; Why and How?, National Bureau of Soil Survey and Land Use Planning, Nagpur, 49 pp.
10. Shivaprasad, C.R., R.S. Reddy, J. Sehgal and M. Velayuthum (1998) Soils of Karnataka for Optimising Land Use, NBSS Publ. No. 47b, NBSS & LUP, Nagpur, India.
11. Soil Survey Staff (2006) Keys to Soil Taxonomy, Tenth edition, U.S. Department of Agriculture/ NRCS, Washington DC, U.S.A.
12. Soil Survey Staff (2012) Soil Survey Manual, Handbook No. 18, USDA, Washington DC, USA.



## Appendix – I

Soil Site and Thematic Information														
Village	Survey No.	Total Area (ha)	Soil phase	Land Management Unit	Soil Depth	Surface Soil Texture	Soil Graveliness	AWC	Slope	Soil Erosion	CLU Code	WELLS	Land Capability	Conservation Plan
Niragudi	1	0.81	Habitation	Others	Others	Others	Others	Others	Others	Others	Grazingland (Gl)	Not Available	Others	Others
Niragudi	2	6.31	MGTmB2g1	LMU-2	Very shallow (<25 cm)	Clay	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar+Redgram (Jw+Rg)	Not Available	IIIse	Crescent bund/TCB
Niragudi	3	0.17	MGTmB2g1	LMU-2	Very shallow (<25 cm)	Clay	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Cultivatedfallow (Cf)	Not Available	IIIse	Crescent bund/TCB
Niragudi	4	8.03	MGTmD3g2	LMU-1	Very shallow (<25 cm)	Clay	Very gravelly (35-60%)	Very low (<50 mm/m)	Moderately sloping (5-10%)	Severe	Redgram+Jowar+ Grazingland (Rg+Jw+Gl)	Not Available	VIse	Crescent bund/TCB
Niragudi	5	6.12	MGTmD3g2	LMU-1	Very shallow (<25 cm)	Clay	Very gravelly (35-60%)	Very low (<50 mm/m)	Moderately sloping (5-10%)	Severe	Jowar + Cultivatedfallow (Jw+Cf)	Not Available	VIse	Crescent bund/TCB
Niragudi	6	9.52	MGTmD3g2	LMU-1	Very shallow (<25 cm)	Clay	Very gravelly (35-60%)	Very low (<50 mm/m)	Moderately sloping (5-10%)	Severe	Redgram+Greengram+Jowar+ Grazingland (Rg+Gg+Jw+Gl)	Not Available	VIse	Crescent bund/TCB
Niragudi	7	1.82	RNLmA1	LMU-5	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Jowar+Sugarcane (Jw+Sc)	Not Available	IIse	GB/strengthening of Bunds
Niragudi	8	4.1	RNLmA1	LMU-5	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Redgram+Sugarcane+Grazingland (Rg+Sc+Gl)	Bore Well, Check Dam	IIse	GB/strengthening of Bunds
Niragudi	9	8.43	MGTmD3g2	LMU-1	Very shallow (<25 cm)	Clay	Very gravelly (35-60%)	Very low (<50 mm/m)	Moderately sloping (5-10%)	Severe	Redgram+Green gram+Jowar+ Grazingland (Rg+Gg+Jw+Gl)	Not Available	VIse	Crescent bund/TCB
Niragudi	10	7.55	NHAmB2	LMU-3	Shallow (25-50 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar+Redgram (Jw+Rg)	Not Available	IIIse	Crescent bund/TCB
Niragudi	11	12.98	NHAmB2	LMU-3	Shallow (25-50 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar+Redgram+ Sugarcane (Jw+Rg+Sc)	Open Well	IIIse	Crescent bund/TCB
Niragudi	12	8.28	NHAmB2	LMU-3	Shallow (25-50 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Jowar+ Grazingland+ Cultivatedfallow (Rg+Jw+Gl+Cf)	Not Available	IIIse	Crescent bund/TCB
Niragudi	13	8.92	NHAmB2	LMU-3	Shallow (25-50 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar+Redgram (Jw+Rg)	Not Available	IIIse	Crescent bund/TCB

Village	Survey No.	Total Area (ha)	Soil phase	Land Management Unit	Soil Depth	Surface Soil Texture	Soil Graveliness	AWC	Slope	Soil Erosion	CLU Code	WELLS	Land Capability	Conservation Plan
Niragudi	14	11.07	NHAmB2	LMU-3	Shallow (25-50 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar+Redgram (Jw+Rg)	Not Available	IIIse	Crescent bund/TCB
Niragudi	15_GR ASS_FI ELD	13.48	MGTmC3g2	LMU-1	Very shallow (<25 cm)	Clay	Very gravelly (35-60%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Grazingland (Gl)	Not Available	IVse	Crescent bund/TCB
Niragudi	16	6.54	MGTmC3g2	LMU-1	Very shallow (<25 cm)	Clay	Very gravelly (35-60%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Redgram+Jowar+ Grazingland (Rg+Jw+Gl)	Not Available	IVse	Crescent bund/TCB
Niragudi	17	10.06	NHAmB2	LMU-3	Shallow (25-50 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar+Redgram (Jw+Rg)	Not Available	IIIse	Crescent bund/TCB
Niragudi	18	10.35	NHAmB2	LMU-3	Shallow (25-50 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar+Redgram (Jw+Rg)	Not Available	IIIse	Crescent bund/TCB
Niragudi	32_GF*	0.83	Habitation	Others	Others	Others	Others	Others	Others	Others	Grazingland (Gl)	Not Available	Others	Others
Niragudi	206	0.57	DSImA1	LMU-4	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Nearly level (0-1%)	Slight	NA	Not Available	IIs	TCB/GB/ strengthening of Bunds
Niragudi	207	0.9	DSImA1	LMU-4	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Nearly level (0-1%)	Slight	NA	Not Available	IIs	TCB/GB/ strengthening of Bunds
Niragudi	208	0.53	DSImA1	LMU-4	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Nearly level (0-1%)	Slight	NA	Not Available	IIs	TCB/GB/ strengthening of Bunds
Niragudi	209	1.82	Habitation	Others	Others	Others	Others	Others	Others	Others	Jowar+Sugarcane (Jw+Sc)	Not Available	Others	Others
Niragudi	210	1.23	DSImA1	LMU-4	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Nearly level (0-1%)	Slight	Cultivatedfallow (Cf)	Not Available	IIs	TCB/GB/ strengthening of Bunds
Niragudi	211	0.83	DSImA1	LMU-4	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Nearly level (0-1%)	Slight	Redgram (Rg)	Not Available	IIs	TCB/GB/ strengthening of Bunds
Niragudi	212	0.92	DSImA1	LMU-4	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Nearly level (0-1%)	Slight	Jowar (Jw)	Not Available	IIs	TCB/GB/ strengthening of Bunds
Niragudi	213	0.77	DSImA1	LMU-4	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Nearly level (0-1%)	Slight	Cultivatedfallow (Cf)	Not Available	IIs	TCB/GB/ strengthening of Bunds



Village	Survey No.	Total Area (ha)	Soil phase	Land Management Unit	Soil Depth	Surface Soil Texture	Soil Graveliness	AWC	Slope	Soil Erosion	CLU Code	WELLS	Land Capability	Conservation Plan
Niragudi	214	0.74	DSImA1	LMU-4	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Nearly level (0-1%)	Slight	Cultivatedfallow (Cf)	Not Available	IIs	TCB/GB/strengthening of Bunds
Niragudi	215	0.94	DSImA1	LMU-4	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Nearly level (0-1%)	Slight	Sugarcane (Sc)	Open Well	IIs	TCB/GB/strengthening of Bunds
Niragudi	216	0.98	DSImA1	LMU-4	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Nearly level (0-1%)	Slight	Sugarcane (Sc)	Not Available	IIs	TCB/GB/strengthening of Bunds
Niragudi	217	1.36	DSImA1	LMU-4	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Nearly level (0-1%)	Slight	Jowar+Redgram (Jw+Rg)	Not Available	IIs	TCB/GB/strengthening of Bunds
Niragudi	218	1.13	DSImA1	LMU-4	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Nearly level (0-1%)	Slight	Jowar+Redgram (Jw+Rg)	Not Available	IIs	TCB/GB/strengthening of Bunds
Niragudi	219	0.89	DSImA1	LMU-4	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Nearly level (0-1%)	Slight	Sugarcane (Sc)	Not Available	IIs	TCB/GB/strengthening of Bunds
Niragudi	220	1	DSImA1	LMU-4	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Nearly level (0-1%)	Slight	Sugarcane (Sc)	Not Available	IIs	TCB/GB/strengthening of Bunds
Niragudi	221	1.07	DSImA1	LMU-4	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Nearly level (0-1%)	Slight	Sugarcane (Sc)	Not Available	IIs	TCB/GB/strengthening of Bunds
Niragudi	222	2.08	DSImA1	LMU-4	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Nearly level (0-1%)	Slight	Redgram+Sugarcane (Rg+Sc)	Open Well	IIs	TCB/GB/strengthening of Bunds
Niragudi	223	0.1	Habitation	Others	Others	Others	Others	Others	Others	Others	NA	Not Available	Others	Others
Niragudi	224	0.22	NHAmC2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Gently sloping (3-5%)	Moderate	Cultivatedfallow (Cf)	Not Available	IIIse	Crescent bund/TCB
Niragudi	225	2.33	NHAmC2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Gently sloping (3-5%)	Moderate	Jowar+Redgram (Jw+Rg)	Not Available	IIIse	Crescent bund/TCB
Niragudi	226	0.92	DSImA1	LMU-4	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Nearly level (0-1%)	Slight	Cultivatedfallow (Cf)	Not Available	IIs	TCB/GB/strengthening of Bunds

Village	Survey No.	Total Area (ha)	Soil phase	Land Management Unit	Soil Depth	Surface Soil Texture	Soil Graveliness	AWC	Slope	Soil Erosion	CLU Code	WELLS	Land Capability	Conservation Plan
Niragudi	227	0.87	DSImA1	LMU-4	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Nearly level (0-1%)	Slight	Jowar (Jw)	Not Available	IIs	TCB/GB/strengthening of Bunds
Niragudi	228	1.11	DSImA1	LMU-4	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Nearly level (0-1%)	Slight	Redgram (Rg)	Open Well	IIs	TCB/GB/strengthening of Bunds
Niragudi	229	1.03	DSImA1	LMU-4	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Nearly level (0-1%)	Slight	Sugarcane (Sc)	Bore Well	IIs	TCB/GB/strengthening of Bunds
Niragudi	230	2.83	DSImA1	LMU-4	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Nearly level (0-1%)	Slight	Sugarcane+Scrubland (Sc+Sl)	Not Available	IIs	TCB/GB/strengthening of Bunds
Niragudi	231	7.7	BHImB2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Sugarcane+Cultivatedfallow (Rg+Sc+Cf)	Open Well	IIIse	Crescent bund/TCB
Niragudi	232	7.35	BHImB2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar+Redgram+Sugarcane(Jw+Rg+Sc)	Open Well	IIIse	Crescent bund/TCB
Niragudi	233	1.09	BHImB2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Scrubland+Jowar (Sl+Jw)	Open Well	IIIse	Crescent bund/TCB
Niragudi	256	3.47	BHImB2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Cultivatedfallow (Cf)	Not Available	IIIse	Crescent bund/TCB
Niragudi	257	2.76	BHImB2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Cultivatedfallow (Cf)	Not Available	IIIse	Crescent bund/TCB
Niragudi	258	2.72	BHImB2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Cultivatedfallow (Cf)	Not Available	IIIse	Crescent bund/TCB
Niragudi	264	5.67	DSImB2g1	LMU-4	Moderately shallow (50-75 cm)	Clay	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIs	TCB/GB/strengthening of Bunds
Niragudi	265	9.57	DSImB2g1	LMU-4	Moderately shallow (50-75 cm)	Clay	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIs	TCB/GB/strengthening of Bunds
Niragudi	266	6.48	RNLmB2	LMU-5	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Cultivatedfallow (Rg+Cf)	Not Available	IIs	GB/strengthening of Bunds

Village	Survey No.	Total Area (ha)	Soil phase	Land Management Unit	Soil Depth	Surface Soil Texture	Soil Graveliness	AWC	Slope	Soil Erosion	CLU Code	WELLS	Land Capability	Conservation Plan
Niragudi	267	0.75	DSImA1	LMU-4	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Nearly level (0-1%)	Slight	Cultivatedfallow (Cf)	Not Available	IIs	TCB/GB/strengthening of Bunds
Niragudi	268	1.08	DSImA1	LMU-4	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Nearly level (0-1%)	Slight	Cultivatedfallow (Cf)	Not Available	IIs	TCB/GB/strengthening of Bunds
Niragudi	269	0.75	DSImA1	LMU-4	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Nearly level (0-1%)	Slight	Cultivatedfallow (Cf)	Not Available	IIs	TCB/GB/strengthening of Bunds
Niragudi	270	0.58	DSImA1	LMU-4	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Nearly level (0-1%)	Slight	Cultivatedfallow (Cf)	Not Available	IIs	TCB/GB/strengthening of Bunds
Niragudi	271	0.59	DSImA1	LMU-4	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Nearly level (0-1%)	Slight	Cultivatedfallow (Cf)	Not Available	IIs	TCB/GB/strengthening of Bunds
Niragudi	272	1.32	DSImA1	LMU-4	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Nearly level (0-1%)	Slight	Jowar + Cultivatedfallow (Jw+Cf)	Not Available	IIs	TCB/GB/strengthening of Bunds
Niragudi	273	0.91	DSImA1	LMU-4	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Nearly level (0-1%)	Slight	Jowar+Redgram (Jw+Rg)	Not Available	IIs	TCB/GB/strengthening of Bunds
Niragudi	274	0.55	DSImA1	LMU-4	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Nearly level (0-1%)	Slight	Cultivatedfallow (Cf)	Not Available	IIs	TCB/GB/strengthening of Bunds
Niragudi	275	0.97	DSImA1	LMU-4	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Nearly level (0-1%)	Slight	Cultivatedfallow (Cf)	Not Available	IIs	TCB/GB/strengthening of Bunds
Niragudi	276	0.82	DSImA1	LMU-4	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Nearly level (0-1%)	Slight	Jowar (Jw)	Not Available	IIs	TCB/GB/strengthening of Bunds
Niragudi	277	0.92	DSImA1	LMU-4	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Nearly level (0-1%)	Slight	Redgram (Rg)	Not Available	IIs	TCB/GB/strengthening of Bunds
Niragudi	278	0.85	DSImA1	LMU-4	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Nearly level (0-1%)	Slight	Redgram (Rg)	Open Well	IIs	TCB/GB/strengthening of Bunds

Village	Survey No.	Total Area (ha)	Soil phase	Land Management Unit	Soil Depth	Surface Soil Texture	Soil Graveliness	AWC	Slope	Soil Erosion	CLU Code	WELLS	Land Capability	Conservation Plan
Niragudi	279	6.89	MGTmC3g2	LMU-1	Very shallow (<25 cm)	Clay	Very gravelly (35-60%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Redgram+Cultivatedfallow+Grazingland (Rg+Cf+G)	Not Available	IVse	Crescent bund/TCB
Niragudi	280	8.06	RNLmB2	LMU-5	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Grazingland (Rg+G)	Not Available	IIse	GB/strengthening of Bunds
Niragudi	281	9.09	DSImB2g1	LMU-4	Moderately shallow (50-75 cm)	Clay	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Sunflower+Redgram+Cultivatedfallow (Sf+Rg+Cf)	Not Available	IIse	TCB/GB/strengthening of Bunds
Niragudi	282	8.41	DSImB2g1	LMU-4	Moderately shallow (50-75 cm)	Clay	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIse	TCB/GB/strengthening of Bunds
Niragudi	283	12.59	NHAmB2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Cultivatedfallow (Rg+Cf)	Not Available	IIIse	Crescent bund/TCB
Niragudi	284	8.44	NHAmB2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar+Grazingland (Jw+G)	Not Available	IIIse	Crescent bund/TCB
Niragudi	285	11.73	MGTmC3g2	LMU-1	Very shallow (<25 cm)	Clay	Very gravelly (35-60%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Redgram+Jowar+Grazingland (Rg+Jw+G)	Check Dam, Check Dam	IVse	Crescent bund/TCB
Niragudi	286	10.74	MGTmC3g2	LMU-1	Very shallow (<25 cm)	Clay	Very gravelly (35-60%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Redgram+Greengram+Jowar+Grazingland (Rg+Gg+Jw+G)	Not Available	IVse	Crescent bund/TCB
Niragudi	287	9.94	MGTmC3g2	LMU-1	Very shallow (<25 cm)	Clay	Very gravelly (35-60%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Redgram+Greengram+Grazingland+Cultivatedfallow (Rg+Gg+G+G+Cf)	Not Available	IVse	Crescent bund/TCB
Niragudi	288	1.21	DSImA1	LMU-4	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Nearly level (0-1%)	Slight	Cultivatedfallow (Cf)	Not Available	IIse	TCB/GB/strengthening of Bunds
Niragudi	289	1.28	DSImA1	LMU-4	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Nearly level (0-1%)	Slight	Redgram (Rg)	Not Available	IIse	TCB/GB/strengthening of Bunds
Niragudi	290	0.72	DSImA1	LMU-4	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Nearly level (0-1%)	Slight	Redgram (Rg)	Not Available	IIse	TCB/GB/strengthening of Bunds

Village	Survey No.	Total Area (ha)	Soil phase	Land Management Unit	Soil Depth	Surface Soil Texture	Soil Graveliness	AWC	Slope	Soil Erosion	CLU Code	WELLS	Land Capability	Conservation Plan
Niragudi	291	1.12	NHAmC2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Gently sloping (3-5%)	Moderate	Cultivatedfallow (Cf)	Bore Well	IIIse	Crescent bund/TCB
Niragudi	292	0.48	DSImA1	LMU-4	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Nearly level (0-1%)	Slight	Redgram (Rg)	Not Available	IIs	TCB/GB/strengthening of Bunds
Niragudi	293	0.24	DSImA1	LMU-4	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Nearly level (0-1%)	Slight	Cultivatedfallow (Cf)	Not Available	IIs	TCB/GB/strengthening of Bunds
Niragudi	294	1.12	DSImA1	LMU-4	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Nearly level (0-1%)	Slight	Jowar+Sugarcan e (Jw+Sc)	Open Well	IIs	TCB/GB/strengthening of Bunds
Niragudi	295	0.56	DSImA1	LMU-4	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Nearly level (0-1%)	Slight	Cultivatedfallow (Cf)	Not Available	IIs	TCB/GB/strengthening of Bunds
Niragudi	296	1.31	DSImA1	LMU-4	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Nearly level (0-1%)	Slight	Cultivatedfallow (Cf)	Not Available	IIs	TCB/GB/strengthening of Bunds
Niragudi	297	1.04	DSImA1	LMU-4	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Nearly level (0-1%)	Slight	Cultivatedfallow (Cf)	Not Available	IIs	TCB/GB/strengthening of Bunds
Niragudi	298	0.85	DSImA1	LMU-4	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Nearly level (0-1%)	Slight	Jowar (Jw)	Not Available	IIs	TCB/GB/strengthening of Bunds
Niragudi	299	0.1	RNLmB2	LMU-5	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Cultivatedfallow (Cf)	Not Available	Iise	GB/strengthening of Bunds
Niragudi	300	4.61	MGTmC3g2	LMU-1	Very shallow (<25 cm)	Clay	Very gravelly (35-60%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Redgram+Green gram+Jowar (Rg+Gg+Jw)	Not Available	IVse	Crescent bund/TCB
Niragudi	301	0.7	DSImA1	LMU-4	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Nearly level (0-1%)	Slight	Redgram (Rg)	Not Available	IIs	TCB/GB/strengthening of Bunds
Niragudi	302	0.81	DSImA1	LMU-4	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Nearly level (0-1%)	Slight	Redgram+Cultivated fallow (Rg+Cf)	Not Available	IIs	TCB/GB/strengthening of Bunds

Village	Survey No.	Total Area (ha)	Soil phase	Land Management Unit	Soil Depth	Surface Soil Texture	Soil Graveliness	AWC	Slope	Soil Erosion	CLU Code	WELLS	Land Capability	Conservation Plan
Niragudi	303	5.98	DSImA1	LMU-4	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Nearly level (0-1%)	Slight	Redgram+Jowar+ Cultivatedfallow(Rg+Jw+Cf)	Open Well	IIs	TCB/GB/ strengthening of Bunds
Niragudi	304	14.13	NHAmD3g2	LMU-1	Shallow (25-50 cm)	Clay	Very gravelly (35-60%)	Low (51-100 mm/m)	Moderately sloping (5-10%)	Severe	Jowar+Sugarcan e+Greengram+C ultivatedfallow+ Grazingland (Jw+Sc+Gg+Cf+ Gl)	Not Available	VIse	Crescent bund/TCB
Niragudi	305	12.71	DSImB2g1	LMU-4	Moderately shallow (50-75 cm)	Clay	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar+Redgram (Jw+Rg)	Not Available	IIse	TCB/GB/ strengthening of Bunds
Niragudi	306	6.94	DSImB2g1	LMU-4	Moderately shallow (50-75 cm)	Clay	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar+Redgram (Jw+Rg)	Not Available	IIse	TCB/GB/ strengthening of Bunds
Niragudi	307	8.62	DSImB2g1	LMU-4	Moderately shallow (50-75 cm)	Clay	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	IIse	TCB/GB/ strengthening of Bunds
Niragudi	308	5.93	DSImB2g1	LMU-4	Moderately shallow (50-75 cm)	Clay	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar+Grazingl and+Cultivated fallow (Jw+Gl+Cf)	Not Available	IIse	TCB/GB/ strengthening of Bunds
Niragudi	309	13.74	DSImB2g1	LMU-4	Moderately shallow (50-75 cm)	Clay	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar+Redgram (Jw+Rg)	Not Available	IIse	TCB/GB/ strengthening of Bunds
Niragudi	310	10.56	DSImB2g1	LMU-4	Moderately shallow (50-75 cm)	Clay	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar+Redgram (Jw+Rg)	Not Available	IIse	TCB/GB/ strengthening of Bunds
Niragudi	311	6.11	NHAmB2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIIse	Crescent bund/TCB
Niragudi	312	8.19	NHAmC3	LMU-3	Shallow (25-50 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Gently sloping (3-5%)	Severe	Redgram+Grazi ng land (Rg+Gl)	Not Available	IVse	Crescent bund/TCB

Village	Survey No.	Total Area (ha)	Soil phase	Land Management Unit	Soil Depth	Surface Soil Texture	Soil Graveliness	AWC	Slope	Soil Erosion	CLU Code	WELLS	Land Capability	Conservation Plan
Niragudi	313	3.73	NHAmD3g2	LMU-1	Shallow (25-50 cm)	Clay	Very gravelly (35-60%)	Low (51-100 mm/m)	Moderately sloping (5-10%)	Severe	Grazingland+Cultivatedfallow (G1+Cf)	Bore Well, Open Well	Vlse	Crescent bund/TCB
Niragudi	314	7.06	NHAmB1	LMU-3	Shallow (25-50 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Redgram+Jowar+ Cultivatedfallow (Rg+Jw+Cf)	Open Well	IIIs	Crescent bund/TCB
Niragudi	315	1.86	NHAmB1	LMU-3	Shallow (25-50 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	NA	Not Available	IIIs	Crescent bund/TCB
Niragudi	316	0.83	MGTmD3g2	LMU-1	Very shallow (<25 cm)	Clay	Very gravelly (35-60%)	Very low (<50 mm/m)	Moderately sloping (5-10%)	Severe	Jowar (Jw)	Not Available	Vlse	Crescent bund/TCB
Niragudi	317	10.64	MGTmD3g2	LMU-1	Very shallow (<25 cm)	Clay	Very gravelly (35-60%)	Very low (<50 mm/m)	Moderately sloping (5-10%)	Severe	Grazingland (G1)	Not Available	Vlse	Crescent bund/TCB
Niragudi	318	9.68	KMPmB2g1	LMU-4	Moderately deep (75-100 cm)	Clay	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Grazingland (G1)	Not Available	Ilse	TCB/GB/strengthening of Bunds
Niragudi	319	1.4	RNLmA1	LMU-5	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Cultivatedfallow (Cf)	Not Available	IIs	GB/strengthening of Bunds
Niragudi	320	5.86	RNLmA1	LMU-5	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Redgram+Cultivatedfallow (Rg+Cf)	Open Well	IIs	GB/strengthening of Bunds
Niragudi	321	0.14	RNLmA1	LMU-5	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	NA	Not Available	IIs	GB/strengthening of Bunds
Niragudi	322	0.75	RNLmA1	LMU-5	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Cultivatedfallow (Cf)	Not Available	IIs	GB/strengthening of Bunds
Niragudi	323	1.2	RNLmA1	LMU-5	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Redgram (Rg)	Not Available	IIs	GB/strengthening of Bunds
Niragudi	324	0.37	RNLmA1	LMU-5	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	NA	Not Available	IIs	GB/strengthening of Bunds
Niragudi	325	1.13	RNLmA1	LMU-5	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Jowar (Jw)	Borewell, Open Well	IIs	GB/strengthening of Bunds

Village	Survey No.	Total Area (ha)	Soil phase	Land Management Unit	Soil Depth	Surface Soil Texture	Soil Graveliness	AWC	Slope	Soil Erosion	CLU Code	WELLS	Land Capability	Conservation Plan
Niragudi	326	0.56	RNLmA1	LMU-5	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	NA	Not Available	IIs	GB/strengthening of Bunds
Niragudi	327	4.81	RNLmA1	LMU-5	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Redgram+Jowar+ Cultivatedfallow(Rg+Jw+Cf)	Not Available	IIs	GB/strengthening of Bunds
Niragudi	328	11.35	NHAmC2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Gently sloping (3-5%)	Moderate	Redgram+Jowar+ Cultivatedfallow(Rg+Jw+Cf)	Open Well	IIIse	Crescent bund/TCB
Niragudi	329	0.07	Habitation	Others	Others	Others	Others	Others	Others	Others	NA	Not Available	Others	Others
Niragudi	330_GF*	0.57	Habitation	Others	Others	Others	Others	Others	Others	Others	Grazingland (GI)	Not Available	Others	Others
Niragudi	SETTLEMENT	8.65	Habitation	Others	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others
Niragudi	STREAM	17.63	Waterbody	Others	Others	Others	Others	Others	Others	Others	Waterbody	Not Available	Others	Others

GF\* - GrassField



## Appendix - II

### Soil Fertility Information

Village	Survey No.	Soil Reaction (pH)	EC	Organic Carbon	Available Phosphorus	Available Potassium	Available Sulphur	Available Boron	Available Iron	Available Manganese	Available Copper	Available Zinc
Niragudi	1	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Niragudi	2	Slightly alkaline (pH 7.3-7.8)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	Medium (145-337 kg/ha)	Medium (10-20 ppm)	Medium (0.5-1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Sufficient (>0.6 ppm)
Niragudi	3	Slightly alkaline (pH 7.3-7.8)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	Medium (145-337 kg/ha)	Medium (10-20 ppm)	Medium (0.5-1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Sufficient (>0.6 ppm)
Niragudi	4	Slightly alkaline (pH 7.3-7.8)	Non Saline (<2 dsm)	High (>0.75 %)	Low (<23 kg/ha)	Medium (145-337 kg/ha)	Medium (10-20 ppm)	Medium (0.5-1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Sufficient (>0.6 ppm)
Niragudi	5	Neutral (pH 6.5-7.3)	Non Saline (<2 dsm)	High (>0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	Medium (10-20 ppm)	Medium (0.5-1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Sufficient (>0.6 ppm)
Niragudi	6	Slightly alkaline (pH 7.3-7.8)	Non Saline (<2 dsm)	High (>0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	Medium (10-20 ppm)	Medium (0.5-1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Sufficient (>0.6 ppm)
Niragudi	7	Moderately alkaline (pH 7.8-8.4)	Non Saline (<2 dsm)	High (>0.75 %)	Medium (23-57 kg/ha)	High (>337 kg/ha)	Medium (10-20 ppm)	Low (<0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Niragudi	8	Slightly alkaline (pH 7.3-7.8)	Non Saline (<2 dsm)	High (>0.75 %)	Medium (23-57 kg/ha)	High (>337 kg/ha)	Medium (10-20 ppm)	Medium (0.5-1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Niragudi	9	Slightly alkaline (pH 7.3-7.8)	Non Saline (<2 dsm)	High (>0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	Medium (10-20 ppm)	Medium (0.5-1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Sufficient (>0.6 ppm)
Niragudi	10	Neutral (pH 6.5-7.3)	Non Saline (<2 dsm)	High (>0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	Medium (10-20 ppm)	Medium (0.5-1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Sufficient (>0.6 ppm)
Niragudi	11	Neutral (pH 6.5-7.3)	Non Saline (<2 dsm)	High (>0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	Medium (10-20 ppm)	Medium (0.5-1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Sufficient (>0.6 ppm)
Niragudi	12	Neutral (pH 6.5-7.3)	Non Saline (<2 dsm)	High (>0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	Medium (10-20 ppm)	Medium (0.5-1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Sufficient (>0.6 ppm)
Niragudi	13	Neutral (pH 6.5-7.3)	Non Saline (<2 dsm)	High (>0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	Medium (10-20 ppm)	Low (<0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Sufficient (>0.6 ppm)
Niragudi	14	Neutral (pH 6.5-7.3)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	Medium (10-20 ppm)	Low (<0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Sufficient (>0.6 ppm)
Niragudi	15_GF*	Neutral (pH 6.5-7.3)	Non Saline (<2 dsm)	High (>0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	Medium (10-20 ppm)	Low (<0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Sufficient (>0.6 ppm)
Niragudi	16	Neutral (pH 6.5-7.3)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	Medium (10-20 ppm)	Low (<0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Sufficient (>0.6 ppm)
Niragudi	17	Slightly alkaline (pH 7.3-7.8)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	Medium (10-20 ppm)	Low (<0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Sufficient (>0.6 ppm)











Village	Survey No.	Soil Reaction (pH)	EC	Organic Carbon	Available Phosphorus	Available Potassium	Available Sulphur	Available Boron	Available Iron	Available Manganese	Available Copper	Available Zinc
Niragudi	322	Moderately alkaline (pH 7.8-8.4)	Non Saline (<2 dsm)	High (>0.75 %)	Medium (23-57 kg/ha)	High (>337 kg/ha)	Medium (10-20 ppm)	Medium (0.5-1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Niragudi	323	Moderately alkaline (pH 7.8-8.4)	Non Saline (<2 dsm)	High (>0.75 %)	Medium (23-57 kg/ha)	High (>337 kg/ha)	Medium (10-20 ppm)	Medium (0.5-1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Niragudi	324	Moderately alkaline (pH 7.8-8.4)	Non Saline (<2 dsm)	High (>0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	Medium (10-20 ppm)	High (>1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Niragudi	325	Moderately alkaline (pH 7.8-8.4)	Non Saline (<2 dsm)	High (>0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	Medium (10-20 ppm)	High (>1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Niragudi	326	Moderately alkaline (pH 7.8-8.4)	Non Saline (<2 dsm)	High (>0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	Medium (10-20 ppm)	Medium (0.5-1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Sufficient (>0.6 ppm)
Niragudi	327	Slightly alkaline (pH 7.3-7.8)	Non Saline (<2 dsm)	High (>0.75 %)	Low (<23 kg/ha)	Medium (145-337 kg/ha)	Medium (10-20 ppm)	High (>1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Sufficient (>0.6 ppm)
Niragudi	328	Slightly alkaline (pH 7.3-7.8)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	Medium (145-337 kg/ha)	Medium (10-20 ppm)	Medium (0.5-1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Niragudi	329	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Niragudi	330_GF*	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Niragudi	SETTLEMENT	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Niragudi	STREAM	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others

GF\* - Grass Field





**Appendix – III**  
**Soil Suitability Information**

Village	Survey No.	Sorghum	Maize	Red gram	Sunflower	Cotton	Sugarcane	Soybean	Guava	Mango	Sapota	Jack fruit	Jamun	Musambi	Lime	Cashew	Custard Apple	Amla	Tamarind
Niragudi	1	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Niragudi	2	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N
Niragudi	3	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N
Niragudi	4	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N
Niragudi	5	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N
Niragudi	6	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N
Niragudi	7	S1	S3t	S3t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S1
Niragudi	8	S1	S3t	S3t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S1
Niragudi	9	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N
Niragudi	10	S3r	S3rt	S3rt	N	S3r	N	S3r	N	N	N	N	N	N	N	N	N	S3r	S3r
Niragudi	11	S3r	S3rt	S3rt	N	S3r	N	S3r	N	N	N	N	N	N	N	N	N	S3r	S3r
Niragudi	12	S3r	S3rt	S3rt	N	S3r	N	S3r	N	N	N	N	N	N	N	N	N	S3r	S3r
Niragudi	13	S3r	S3rt	S3rt	N	S3r	N	S3r	N	N	N	N	N	N	N	N	N	S3r	S3r
Niragudi	14	S3r	S3rt	S3rt	N	S3r	N	S3r	N	N	N	N	N	N	N	N	N	S3r	S3r
Niragudi	15_GF*	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N
Niragudi	16	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N
Niragudi	17	S3r	S3rt	S3rt	N	S3r	N	S3r	N	N	N	N	N	N	N	N	N	S3r	S3r
Niragudi	18	S3r	S3rt	S3rt	N	S3r	N	S3r	N	N	N	N	N	N	N	N	N	S3r	S3r
Niragudi	32_GF*	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Niragudi	206	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2r	S2r	N
Niragudi	207	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2r	S2r	N
Niragudi	208	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2r	S2r	N
Niragudi	209	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Niragudi	210	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2r	S2r	N
Niragudi	211	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2r	S2r	N
Niragudi	212	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2r	S2r	N
Niragudi	213	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2r	S2r	N
Niragudi	214	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2r	S2r	N
Niragudi	215	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2r	S2r	N
Niragudi	216	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2r	S2r	N
Niragudi	217	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2r	S2r	N
Niragudi	218	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2r	S2r	N
Niragudi	219	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2r	S2r	N

Village	Survey No.	Sorghum	Maize	Red gram	Sunflower	Cotton	Sugarcane	Soybean	Guava	Mango	Sapota	Jack fruit	Jamun	Musambi	Lime	Cashew	Custard Apple	Amla	Tamarind
Niragudi	220	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2r	S2r	N
Niragudi	221	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2r	S2r	N
Niragudi	222	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2r	S2r	N
Niragudi	223	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Niragudi	224	S3r	S3rt	S3rt	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Niragudi	225	S3r	S3rt	S3rt	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Niragudi	226	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2r	S2r	N
Niragudi	227	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2r	S2r	N
Niragudi	228	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2r	S2r	N
Niragudi	229	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2r	S2r	N
Niragudi	230	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2r	S2r	N
Niragudi	231	S3r	S3rt	S3rt	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Niragudi	232	S3r	S3rt	S3rt	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Niragudi	233	S3r	S3rt	S3rt	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Niragudi	256	S3r	S3rt	S3rt	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Niragudi	257	S3r	S3rt	S3rt	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Niragudi	258	S3r	S3rt	S3rt	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Niragudi	264	S2re	S2re	S2re	S3r	S2re	S2re	S2re	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2re	S2re	N
Niragudi	265	S2re	S2re	S2re	S3r	S2re	S2re	S2re	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2re	S2re	N
Niragudi	266	S2e	S3t	S3t	S2e	S2e	S3t	S2e	S3t	S3t	S3t	S3t	S2te	S2e	S2e	N	S2e	S2e	S2e
Niragudi	267	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2r	S2r	N
Niragudi	268	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2r	S2r	N
Niragudi	269	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2r	S2r	N
Niragudi	270	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2r	S2r	N
Niragudi	271	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2r	S2r	N
Niragudi	272	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2r	S2r	N
Niragudi	273	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2r	S2r	N
Niragudi	274	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2r	S2r	N
Niragudi	275	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2r	S2r	N
Niragudi	276	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2r	S2r	N
Niragudi	277	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2r	S2r	N
Niragudi	278	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2r	S2r	N
Niragudi	279	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N
Niragudi	280	S2e	S3t	S3t	S2e	S2e	S3t	S2e	S3t	S3t	S3t	S3t	S2te	S2e	S2e	N	S2e	S2e	S2e
Niragudi	281	S2re	S2re	S2re	S3r	S2re	S2re	S2re	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2re	S2re	N

Village	Survey No.	Sorghum	Maize	Red gram	Sunflower	Cotton	Sugarcane	Soybean	Guava	Mango	Sapota	Jackfruit	Jamun	Musambi	Lime	Cashew	Custard Apple	Amla	Tamarind
Niragudi	282	S2re	S2re	S2re	S3r	S2re	S2re	S2re	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2re	S2re	N
Niragudi	283	S3r	S3rt	S3rt	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Niragudi	284	S3r	S3rt	S3rt	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Niragudi	285	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N
Niragudi	286	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N
Niragudi	287	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N
Niragudi	288	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2r	S2r	N
Niragudi	289	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2r	S2r	N
Niragudi	290	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2r	S2r	N
Niragudi	291	S3r	S3rt	S3rt	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Niragudi	292	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2r	S2r	N
Niragudi	293	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2r	S2r	N
Niragudi	294	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2r	S2r	N
Niragudi	295	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2r	S2r	N
Niragudi	296	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2r	S2r	N
Niragudi	297	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2r	S2r	N
Niragudi	298	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2r	S2r	N
Niragudi	299	S2e	S3t	S3t	S2e	S2e	S3t	S2e	S3t	S3t	S3t	S3t	S2te	S2e	S2e	N	S2e	S2e	S2e
Niragudi	300	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N
Niragudi	301	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2r	S2r	N
Niragudi	302	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2r	S2r	N
Niragudi	303	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2r	S2r	N
Niragudi	304	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N
Niragudi	305	S2re	S2re	S2re	S3r	S2re	S2re	S2re	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2re	S2re	N
Niragudi	306	S2re	S2re	S2re	S3r	S2re	S2re	S2re	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2re	S2re	N
Niragudi	307	S2re	S2re	S2re	S3r	S2re	S2re	S2re	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2re	S2re	N
Niragudi	308	S2re	S2re	S2re	S3r	S2re	S2re	S2re	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2re	S2re	N
Niragudi	309	S2re	S2re	S2re	S3r	S2re	S2re	S2re	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2re	S2re	N
Niragudi	310	S2re	S2re	S2re	S3r	S2re	S2re	S2re	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2re	S2re	N
Niragudi	311	S3r	S3rt	S3rt	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Niragudi	312	S3re	S3re	S3re	N	S3re	N	S3re	N	N	N	N	N	N	N	N	S3re	S3r	N
Niragudi	313	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N
Niragudi	314	S3r	S3rt	S3rt	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Niragudi	315	S3r	S3rt	S3rt	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Niragudi	316	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N
Niragudi	317	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N

Village	Survey No.	Sorghum	Maize	Red gram	Sunflower	Cotton	Sugarcane	Soybean	Guava	Mango	Sapota	Jack fruit	Jamun	Musambi	Lime	Cashew	Custard Apple	Amla	Tamarind
Niragudi	318	S2e	S3t	S3t	S2e	S2e	S3t	S2e	S3t	S3t	S3t	S3rt	S3r	S2re	S2re	N	S2e	S2e	S3r
Niragudi	319	S1	S3t	S3t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S1
Niragudi	320	S1	S3t	S3t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S1
Niragudi	321	S1	S3t	S3t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S1
Niragudi	322	S1	S3t	S3t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S1
Niragudi	323	S1	S3t	S3t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S1
Niragudi	324	S1	S3t	S3t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S1
Niragudi	325	S1	S3t	S3t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S1
Niragudi	326	S1	S3t	S3t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S1
Niragudi	327	S1	S3t	S3t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S1
Niragudi	328	S3r	S3rt	S3rt	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Niragudi	329	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Niragudi	330_GF*	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Niragudi	SETTLEMEN T	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Niragudi	STREAM	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others

GF\* - Grass Field

# **PART-B**

**SOCIO-ECONOMIC STATUS OF FARM HOUSEHOLDS**



## CONTENTS

1	Executive summary	1-3
2	Introduction	5
3	Methodology	7-11
4	Results and discussions	13-28





## LIST OF TABLES

<b>I. Social status</b>		
1	Human population among sample households	13
2	Basic needs of sample households	15
3	Migration details among the sample households	15
<b>II. Economic status</b>		
4	Occupational pattern in sample households	16
5	Domestic assets among samples households	17
6	Farm assets among samples households	17
7	Livestock assets among sample households	18
8	Milk produced and Fodder availability of sample households	19
9	Women empowerment of sample households	19
10	Per capita daily consumption of food among the sample households	19
11	Annual average Income from various sources	20
12	Average annual expenditure of sample households	21
13	Distribution of land holding among the sample households	22
14	Land holding among samples households	22
<b>III. Resource use pattern</b>		
15	Number of tree/plants covered in sample farm households	22
16	Present cropping pattern among samples households	23
17	Distribution of soil series in the watershed	23
<b>IV. Economic land evaluation</b>		
18	Cropping pattern on major soil series	24
19	Alternative land use options for different size group of farmers (Benefit Cost Ratio)	24
20	Economics Land evaluation and bridging yield gap for different crops	25
21	Estimation of onsite cost of soil erosion	26
22	Ecosystem services of food grain production	27
23	Ecosystem services of water supply for crop production	27
24	Farming constraints	27

## **LIST OF FIGURES**

1	Location of study area	8
2	ALPES Framework	9
3	Basic needs of sample households	14
4	Domestic assets among the sample households	16
5	Farm assets among samples households	17
6	Livestock assets among sample households	18
7	Per capita daily consumption of food among the sample households	20
8	Average annual expenditure of sample households	21
9	Estimation of onsite cost of soil erosion	26



## EXECUTIVE SUMMARY

*Baseline socioeconomic characterisation is prerequisite to prepare action plan for program implementation and to assess the project performance before making any changes in the watershed development program. The baseline provides appropriate policy direction for enhancing productivity and sustainability in agriculture.*

**Methodology:** *Nirgudi West Microwatershed (Alur sub-watershed, Aland taluk, Gulbarga district) is located in between 17°37'–17°39' North latitudes and 77°26'–77°27' East longitudes, covering an area of about 504.6 ha, bounded by ChincholiBudruk, Matki, Padasavli villages and Maharashtra State with length of growing period (LGP) 120-150 days. We used soil resource map as basis for sampling farm households to test the hypothesis that soil quality influence crop selection, and conservation investment of farm households. The level of technology adoption and productivity gaps and livelihood patterns were analyses. The cost of soil degradation and ecosystem services were quantified.*

**Results:** *The socio-economic outputs for Nirgudi West Microwatershed (Alur sub-watershed, Aland taluk, Gulbarga district) are presented here.*

### **Social Indicators**

- ❖ *Male and female ratio is 53.1 to 46.8 per cent to the total sample population.*
- ❖ *Younger age 18 to 50 years group of population is around 55.3 per cent to the total population.*
- ❖ *Literacy population is around 66.0 per cent.*
- ❖ *Social groups belong to other backward caste (OBC) is around 50.0 per cent.*
- ❖ *Fire wood is the source of energy for a cooking among 90.0 per cent.*
- ❖ *Dependence on ration cards for food grains through public distribution system is around 90 per cent.*
- ❖ *Swach bharath program providing closed toilet facilities among all sample households.*
- ❖ *Rural migration to urban centre for employment is prevalent among 20.0 per cent.*
- ❖ *Women participation in decisions making of agricultural production among the all sample households.*

### **Economic Indicators**

- ❖ *The average land holding is 2.2 ha indicates that majority of farm households are belong to medium and semi-medium farmers. The total cultivated land by dry land condition of the sample farmers.*
- ❖ *Agriculture is the main occupation among 21.3 per cent and agriculture labour is predominant subsidiary occupation for 63.8 per cent of the sample households.*

- ❖ *The average value of domestic assets is around Rs 16948 per household. Mobile and television are popular media mass communication.*
- ❖ *The average value of farm assets is around Rs 120421 per household, about 50 per cent of sample farmers are owning plough.*
- ❖ *The average value of livestock is around Rs 31979 per household; about 40 per cent of household are having livestock.*
- ❖ *The average per capita food consumption is around 897 grams (2149.6 kilo calories) against national institute of nutrition (NIN) recommendation at 827 gram. Around 50 per cent of sample households are consuming less than the NIN recommendation.*
- ❖ *The annual average income is around Rs 120844 per household. About 30.0 per cent of farm households are below poverty line.*
- ❖ *The per capita average monthly expenditure is around Rs 2142.*

#### ***Environmental Indicators-Ecosystem Services***

- ❖ *The value of ecosystem service helps to support investment to decision on soil and water conservation and in promoting sustainable land use.*
- ❖ *The onsite cost of different soil nutrients lost due to soil erosion is around Rs. 3764 per ha/year. The total cost of annual soil nutrients is around Rs. 1754112 per year for the total area of 504.6 ha.*
- ❖ *The average value of ecosystem service for food grain production is around Rs. 10815/ ha/year. Per hectare food grain production services is maximum in red gram (Rs. 20967) followed by soybean (Rs. 662).*
- ❖ *The data on water requirement for producing one quintal of grain is considered for estimating the total value of water required for crop production. The per hectare value of water used and value of water was maximum in redgram (Rs. 58714) and soybean (Rs. 2112).*

#### ***Economic Land Evaluation***

- ❖ *The major cropping pattern is redgram (90.8 %) and soybean (9.2 %).*
- ❖ *In Nirgudi west Microwatershed, major soils are Margutti (MGT) series are having veer shallow soil depth cover around 23.0 % of area. On this soil farmers are presently growing redgram. Novinihala (NHA) are also having shallow soil depth cover around 6.1 % of area, the crops are soybean. Dinsi (DSI) soil series having moderately deep soil depth cover around 27.7 % of areas, crops are red gram. Rajnala (RNL) soil series are having deep soil depth cover around 7.9 % of area. The major crops grown are redgram.*
- ❖ *The total cost of cultivation and benefit cost ratio (BCR) in study area for red gram ranges between Rs.27862/ha in DSI soil (with BCR of 2.19) and Rs.13850/ha in RNL soil (with BCR of 2.53).*

- ❖ *In soybean the cost of cultivation in NHA soil Rs 2290/ha (with BCR of 1.29).*
- ❖ *The land management practices reported by the farmers are crop rotation, tillage practices, fertilizer application and use of farm yard manure (FYM). Due to higher wages farmer are following labour saving strategies is not prating soil and water conservation measures. Less ownership of livestock limiting application of FYM.*
- ❖ *It was observed soil quality influences on the type and intensity of land use. More fertilizer applications in deeper soils to maximize returns.*

### **Suggestions**

- ❖ *Involving farmers in watershed planning helps in strengthening institutional participation.*
- ❖ *The per capita food consumption and monthly income is very low. Diversifying income generation activities from crop and livestock production in order to reduce risk related to drought and market prices.*
- ❖ *Majority of farmers reported that they are not getting timely support/extension services from the concerned development departments.*
- ❖ *By strengthening agricultural extension for providing timely advice improved technology there is scope to increase in net income of farm households.*
- ❖ *By adopting recommended package of practices by following the soil test fertiliser recommendation, there is scope to increase yield in redgram (15.7 to 26.7 %) and soybean (95.0 %).*



## **INTRODUCTION**

Watershed Development program aim to restore degraded watersheds in rainfed regions to increase their capacity to capture and store rain water, reduce soil erosion, and improved soil nutrients and carbon contents so they can produce greater agricultural yields and other benefits. As majority of rural poor live in these regions and dependent on natural resources for their livelihood and sustenance, improvements in agricultural yields improve human welfare and simultaneously improve national food security.

Sujala–III watershed development project conceptualised and implemented by the Watershed Development Department of Government of Karnataka with tripartite cost-sharing arrangements. The World Bank through International Development Association provided major portion of plan outlay as a loan to Government of India and in turn loan to Government of Karnataka.

The objectives of Sujala-III is to demonstrate more effective watershed management through greater integration of programs related to rain fed agriculture, innovative and science based approaches and strengthened institutions and capacities. The project is implemented in 11 districts of Bidar, Vijayapura, Gulbarga, Yadgir, Koppal, Gadag, Raichur, Davanagere, Tumkur, Chikkamangalur and Chamarajanagar which have been identified by the Watershed Development Department based on rainfall and socio-economic conditions. The project will be implemented over six years and linked with the centrally financed integrated watershed management programme.

Economic evaluations can better guide in watershed planning and implementation, as well as raise awareness of benefits of ecosystem restoration for food security and poverty alleviation program. The present study aims to characterize socio-economic status of farm households, assess the land and water use status, evaluate the economic viability of land use, prioritize farming constraints and suggest the measures for soil and water conservation for sustainable agriculture.

### **Objectives of the study**

1. To characterize socio-economic status of farm households
2. To evaluate the economic viability of land use and land related constraints
3. To estimate the ecosystem service provided by the watershed and
4. To suggest alternatives for sustainable agriculture production.





## METHODOLOGY

### *Study area*

Nirgudi West micro-watershed located in North-eastern Dry Zone of Karnataka (Figure 1): The total geographic area of this zone is about 1.76 M ha covering 8 taluks of Gulbarga district and 3 taluks of Raichur. Net cultivated area in the zone is about 1.31 M ha of which about 0.09 M ha are irrigated. The mean elevation of the zone is 300-450 m MSL. The main soil type is deep to very deep soils with small pockets of shallow to medium black soils. The zone is cropped predominantly during rabi due to insufficient rainfall (465-785 mm). The principal crops of the zone are jowar, bajra, oilseeds, pulses, cotton and sugarcane. It's represented Agro Ecological Sub Region (AESR) 6.2 having LGP 120-150 days.

Nirgudi West micro-watershed (Alur sub-watershed, Aland taluk, Gulbarga district) is located in between 17<sup>0</sup>37'–17<sup>0</sup>39' North latitudes and 77<sup>0</sup>26'–77<sup>0</sup>27' East longitudes, covering an area of about 504.6 ha, bounded by Chincholi Budruk, Matki, Padasavli villages and Maharashtra State.

### **Sampling Procedure:**

In this study we have followed soil variability as criterion for sampling the farm households. In each micro-watershed the survey numbers and associated soil series are listed. Minimum three farm households for each soil series were taken and summed up to arrive at total sample for analysis.

### **Sources of data and analysis:**

For evaluating the specific objectives of the study, primary data was collected from the sample respondents by personal interview method with the help of pre-tested questionnaire. The data on socio-economic characteristics of respondents such as family size and composition, land holdings, asset position, occupational pattern and education level was collected. The present cropping pattern and the level of input use and yields collected during survey. The data collected from the representative farm households were analysed using Automated Land Potential Evaluation System (Figure 2).

## LOCATION MAP OF NIRGUDI WEST MICRO-WATERSHED




Figure 1: Location of study area

### Steps followed in socio-economic assessment

- 1 • After the completion of soil profile study link the cadastral number to the soil profile in the micro watershed.
- 2 • Download the names of the farmers who are owning the land for each cadastral number in the Karnataka BHOOMI Website.
- 3 • Compiling the names of the farmers representing for all the soil profiles studied in the micro watershed for socio-economic Survey.
- 4 • Conducting the socioeconomic survey of selected farm households in the micro watershed.
- 5 • Farm households database created using the Automated Land Potential Evaluation System (ALPES) for analysis of socio economic status for each micro watershed.
- 6 • Synthesis of tables and preparation of report for each micro watershed.




Figure 2: ALPES FRAMEWORK

The sample farmers were post classified in to marginal and small (0.0 to  $\leq 2$  ha), medium and semi medium ( $> 2$  to  $\leq 10$  ha) and large ( $> 10$  ha). The steps involved in estimation of soil potential involve estimation of total cost of cultivation, the yield/gross returns and net income per hectare. The cost of inputs such seed, manure and fertilizer, plant protection chemicals, payment towards human and bullock labour and interest on working capita are included under operational costs. In the case of perennial crops, the cost of establishment was estimated by using actual physical requirements and prevailing market prices. Estimation cost included maintenance cost up to bearing period. The value of main product and by product from the crop enterprise at the market rates were the gross returns of the crop. Net returns were worked out by deducting establishment and maintained cost from gross returns.

Operational Cost = cost of seeds, fertilizers, pesticides. Cost of human and bullock labour, cost of machinery, cost of irrigation water + interest on working capital.

Gross returns = Yield (Quintals/hectare)\*Price (Rs/Quintal)

Net returns = Gross returns-Operational cost.


Benefit Cost Ratio = Net returns/Total cost.

Economic suitability classes: once each land use –land area combination has been assigned an economic value by the land evaluation, the question arises as to its ‘suitability’, that is, the degree to which it satisfies the land user. The FAO framework defines two suitability orders: ‘S’(suitable if benefit cost ratio (BCR) $> 1$ ) and ‘N’(not suitable if (BCR $< 1$ ), which are divided into five economic suitability classes: ‘S1’(highly suitable if BCR $> 3$ ), ‘S2’(suitable if BCR $> 2$  and  $< 3$ ), ‘S3’(Marginally suitable if BCR  $> 1$  and  $< 2$ ), ‘N1’(Not suitable for economic reasons but physically suitable) and ‘N2’(not suitable for physical reasons). The limit between ‘S3’ and ‘N1’ must be at least at the point of financial feasibility (i.e. net returns, NPV, or IRR $> 0$  and BCR $> 1$ ). The other limits depend on social factors such as farm size, family size, alternative employment or investment possibilities and wealth expectations; these need to be specified for the Soil series.

### **Economic Valuation of Soil ecosystem services:**

The replacement cost approach was followed for estimating the onsite cost of soil erosion, Market price method was followed for estimating the value of food and fodder production. Value transfer methods was followed for estimating the value of water demand by different crops in the micro watershed.

### Steps followed in Replacement cost methods for estimation of onsite cost of soil erosion





## RESULTS AND DISCUSSIONS

The demographic information shows that the household population dynamics encompasses the socioeconomic status of the farmer. For a rural family, the household size should be optimal to earn a comfortable livelihood through farm and non-farm wage earning. The total number of population in watershed area was 47, out of which 53.1, per cent were males and 46.8, per cent females. Average family size of the households is 4.7. Age is an important factor, which affects the potential employment and mobility status of respondents. The data on age wise distribution of farmers in the sample households indicated that majority of the farmers are coming under the age group of more than 50 years (36.1 %) followed by 18 to 30 years (36.1 %) 30 to 50 years (19.1 %) and 0 to 18 years (8.5 %). Hence, in the study area in general, the respondents were of young and middle age, indicating thereby that the households had almost settled with whatever livelihood options they were practicing and sample respondents were young by age who could venture into various options of livelihood sources. Data on literacy indicated that 34.0 per cent of respondents were illiterate and 66.0 per cent literate (Table 1).

**Table 1: Human population among sample households in Nirgudi west Microwatershed**

Particulars	Units	Value
Total human population in sample HHs	Number	47
Male	% to total Population	53.1
Female	% to total Population	46.8
Average family size	Number	4.7
<b>Age group</b>		
0 to 18 years	% to total Population	8.5
18 to 30 years	% to total Population	36.2s
30 to 50 years	% to total Population	19.1
>50 years	% to total Population	36.1
Average age	Age in years	40.7
<b>Education Status</b>		
Illiterates	% to total Population	34.0
Literates	% to total Population	66.0
Primary School (<5 class)	% to total Population	6.4
Middle School (6- 8 class)	% to total Population	19.2
High School (9- 10 class)	% to total Population	29.8
Others	% to total Population	10.6

The ethnic groups among the sample farm households found to be 50.0 per cent belonging to other backward caste (OBC) followed by 40 per cent belonging to general caste and 10.0 per cent of scheduled caste (SC) (Table 2 and Figure 3). About 90.0 per cent of sample households are using fire wood as source of fuel for cooking. All the sample farmers are having electricity connection. About 90 per cent of farm households



are having ration cards for taking food grains from public distribution system. About 20.0 per cent of farm households are having toilet facilities.




Figure 3: Basic needs of sample households in Nirgudi west Microwatershed

The data collected on the source of drinking water in the study area is presented in Table 2. All the sample respondents are having Tub well source for water supply for domestic purpose.

**Table 2: Basic needs of sample households in Nirgudi West Microwatershed**

Particulars	Units	Value
<b>Social groups</b>		
SC	% of Households	10.0
OBC	% of Households	50.0
General	% of Households	40.0
<b>Types of fuel use for cooking</b>		
Fire wood	% of Households	90.0
Gas	% of Households	10.0
<b>Energy supply for home</b>		
Electricity	% of Households	100
<b>Number of households having Health card</b>		
Yes	% of Households	0.0
No	% of Households	100.0
<b>MGNREGA Card</b>		
Yes	% of Households	0.0
No	% of Households	100.0
<b>Ration Card</b>		
Yes	% of Households	90.0
No	% of Households	10.0
<b>Households with toilet</b>		
Yes	% of Households	20.0
No	% of Households	80.0
<b>Drinking water facilities</b>		
Tube well	% of Households	100.0

The data on migration in Nirgudi west Microwatershed is given in Table 3. It indicated that around 20.0 per cent of samples households were migrated. The average distance travelled for seeking employment is 400 km.

**Table 3: Migration details among the sample households in Nirgudi west Microwatershed**

Particulars	Value
% of households showing migration	20.0
% of persons migrating	6.4
No. of months migrated in a year	15
Average Distance of migration(Km)	400
<b>Nature of job (%)</b>	
Job/wage/work	100.0
Education of the children	0.0

The occupational pattern (Table 4) among sample households shows that agriculture is the main occupation around 21.3 per cent of farmers followed by subsidiary occupations like Agricultural labour (63.8 %). About 10.6 per cent of the households are

private service as main occupation around 2.1 per cent of the population is household Trade and business activity as a main occupation.

**Table 4: Occupational pattern in sample population in Nirgudi West Microwatershed**

Occupation		% to total
Main	Subsidiary	
Agriculture	Agriculture	21.3
	Agriculture Labour	63.8
Govt. service		2.1
Private service		10.6
Trade and business		2.1
Grand Total		100.0
<b>Family labour availability</b>		<b>Man days/month</b>
Male		25.0
Female		20.0
Total		45.0




Figure 4: Domestic assets among the sample households in Nirgudi West Microwatershed

The important assets especially with reference to domestic assets were analyzed and are given in Table 5 and Figure 4. The important domestic assets possessed by all categories of farmers are mobile phones (100 %) followed by television (90.0 %), mixer/grinder (40.0 %), motorcycle (40.0 %) and refrigerator (10.0 %). The average value of domestic assets is around Rs 16948 per households.

**Table 5: Domestic assets among the sample households in Nirgudi West Microwatershed**

Particulars	% of households	Average value in Rs
Mixer/grinder	40.0	2125
Mobile Phone	100	6450
Motor cycle	40.0	56500
Refrigerator	10.0	12000
Television	60.0	7667
Average value	16948	




Figure 5: Farm assets among samples households in Nirgudi West Microwatershed

**Table 6: Farm assets among samples households in Nirgudi West Microwatershed**

Particulars	% of households	Average value in Rs
Bullock cart	50.0	23000
Harvester	10.0	2000
Plough	50.0	6800
Seed cum fertiliser drill	10.0	4500
Sprayer	40.0	6250
Tractor	10.0	800000
Weeder	10.0	400
Average Value	<b>120421</b>	

The most popularly owned farm equipments were sickles, plough, cattle shed; pump sets, chaff cutter, bullock cart, sprayer and thresher. Plough and sickle were

commonly present in all the sampled farmers; these were primary implements in agriculture. The per cent of households owned plough (50.0 %), bullock cart (50.0 %), sprayer (40.0 %), weeder (10.0 %), tractor (10.0 %), seed cum fertiliser drill (10.0 %) and harvester (10.0 %) was found highest among the sample farmers. The average value of farm assets is around Rs 120421 per households (Table 6 and Figure 5).

Livestock is an integral component of the conventional farming systems (Table 7 and Figure 6). The highest livestock population is buffalos were around (40 %) per cent followed by local milching cow (26.6 %), local dry cow (13.3 %) and milching buffalos (20 %). The average livestock value was Rs 31979 per household.

**Table 7: Livestock assets among sample households in Nirgudi West Microwatershed**

Particulars	% of livestock population	Average value in Rs
Local Dry Cow	13.3	15000
Local Milching Cow	26.6	16250
Milching Buffalos	20.0	46667
Bullocks	40.0	50000
Average value	31979	




Figure 6: Livestock assets among sample households in Nirgudi West Microwatershed

Average milk produced in sample households is 1680 liters/ annum of local milching cow and milching buffalos (Table 8). The households having livestock is 78.9 per cent of sample farmers.

**Table 8: Milk produced and fodder availability of sample households in Nirgudi west Microwatershed**

<b>Particulars</b>	
<b>Name of the Livestock</b>	<b>Ltr./Lactation/animal</b>
Local Milching Cow	1175
Milching Buffalos	2353
Average Milk produced	1680
Livestock having households (%)	78.9
Livestock population (Numbers)	25

A woman participation in decision making in this micro-watershed is presented in Table 9. About 10 per cent of women participation in local organisation activates and 100 per cent of women taking decision in her family and agriculture related activities, 100 per cent Women earning for her family requirement.

**Table 9: Women empowerment of sample households in Nirgudi West Microwatershed**

<b>Particulars</b>	<b>% of grand total</b>	
	<b>Yes</b>	<b>No</b>
Women participation in local organization activities	10.0	90.0
Women elected as panchayat member	0.0	100
Women earning for her family requirement	100.0	0.0
Women taking decision in her family and agriculture related activities	100.0	0.0

**Table 10: Per capita daily consumption of food among the sample households in Nirgudi west Microwatershed**

<b>Particulars</b>	<b>NIN recommendation (gram/ per day/ person)</b>	<b>Present level of consumption (gram/ per day/ person)</b>	<b>Kilo Calories /day/person</b>
Cereals	396.0	436.1	1482.9
Pulses	43.0	63.0	216.1
Milk	200.0	131.7	85.6
Vegetables	143.0	133.1	31.9
Cooking Oil	31.0	31.8	181.2
Egg	0.5	83.0	124.5
Meat	14.2	18.1	27.1
<b>Total</b>	<b>827.7</b>	<b>897</b>	<b>2149.6</b>
Threshold of NIN recommendation		827 gram*	2250 Kcal*
% Below NIN		50.0	60.0
% Above NIN		50.0	40.0

Note: \* day/person

The food intake in terms of kilo calorie (kcal) per person per day was calculated and presented in the Table 10 and Figure 7. More quantity of cereals are consumed by sample farmers which accounted for 1482.9 kcal per person. The other important food

items consumed was pulses 216.1 kcal followed by cooking oil 181.2 kcal, milk 85.6 kcal, vegetables 31.9 kcal, egg 124.5 kcal and meat 27.1 kcal. In the sampled households, farmers were consuming less (2149.6 kcal) than NIN- recommended food requirement (2250 kcal).




Figure 7: Per capita daily consumption of food among the sample households in Nirgudi West Microwatershed

**Annual income of the sample HHs:** The average annual household income is around Rs 120844. Major source of income to the farmers in the study area is from livestock (Rs 70395) followed by crop production (Rs. 44549). The income from Non farm income was very low at Rs 5900. The monthly per capita income is Rs.2142, which is less than the threshold monthly income of Rs 975 for considering above poverty line. Due to the fact that erratic rainfall and shortage of water, farmers are diverting from crop production activities to enable the household for a comfortable livelihood. The incomes from the other aforesaid sources are very meagre (Table 11).

**Table 11: Annual average income of HHs from various sources in Nirgudi west Microwatershed**

Particulars	Income *
Nonfarm income (Rs)	5900(20)
Livestock income (Rs)	70395(40)
Crop Production (Rs)	44549(100)
<b>Total Annual Income (Rs)</b>	<b>120844</b>
Average monthly per capita income (Rs)	2142
<b>Threshold for Poverty level (Rs 975 per month/person)</b>	
% of households below poverty line	30.0
% of households above poverty line	70.0

\* Figure in the parenthesis indicates % of Households

The average annual expenditure of farm households indicated that farmers in the study area spend highest on food (Rs. 44803) followed by education, clothing, social function and health. Now a day's education is most important among all of us. In today's competitive world, education is a necessity for man after food, clothing, and shelter. It is the only fundamental way by which a desired change in the society can happen. The average per capita monthly expenditure is around Rs 1991 and about 30.0 per cent of farm households are below poverty line (Table 12 and Figure 8).

**Table 12: Average annual expenditure of sample HHs in Nirgudi West Microwatershed**

Particulars	Value in Rupees	Per cent
Food	44803	40
Education	400	0.4
Clothing	8400	7.5
Social functions	45000	40.1
Health	13700	12.2
Total Expenditure (Rs/year)	112303	100.0
Monthly per capita expenditure (Rs)	1991	




Figure 8: Average annual expenditure of sample HHs in Nirgudi West Microwatershed

**Land holding:** Total area cultivated by them is 22.4 ha. The average land holding of sample HHs is 2.2 ha. Large number of sample HHs (60.0%) belong to medium size group with an average holding size of 2.2 ha followed by small farmers (30.0%) with an



average holding size of 1.1 ha and a large farmer (10.0%) with an average land holding size of 5.4 ha (Table 13)

**Table 13: Distribution of land holding among the sample households in Nirgudi West Microwatershed**

Particulars	Units	Values
<b>Small farmers</b>		
Total land	ha	3.8
Sample size	Per cent	30.0
Average land holding	ha	1.1
<b>Medium farmers</b>		
Total land	ha	13.4
Sample size	Per cent	60.0
Average land holding	ha	2.2
<b>Large farmers</b>		
Total land	ha	5.4
Sample size	Per cent	10.0
Average land holding	ha	5.4
<b>Total sample households</b>		
Total land	ha	22.4
Sample size	Per cent	100
Average land holding	ha	2.2

**Land use:** The total land holding in the Nirgudi West Microwatershed is 22.2 ha is dry land (Table 14). The average land holding per household is worked out to be 2.2 ha.

**Table 14: Land use among samples households in Nirgudi West Microwatershed**

Particulars	Per cent	Area in ha
Irrigated land	0.0	0.0
Dry land	100.0	22.2
Fallow Land	0.0	0.0
Total land holding	100.0	22.2
Average land holding	2.2	

In the Microwatershed, the prevalent present land uses under perennial plants are neem trees (71.4 %), followed by banyan tree (19.1 %), mango (4.8 %) and coconut (4.7 %) (Table 15).

**Table 15: Number of trees/plants covered in sample farm households in Nirgudi west Microwatershed**

Particulars	Number of Plants/trees	Per cent
Banyan tree (Alada)	4	19.1
Neem trees	15	71.4
Mango	1	4.8
Coconut	1	4.7
Grand Total	21	100

The land use decisions are usually based on experience of farmers, tradition, expected profit, personal preferences, resources and social requirements. The present dominant crops grown in dry lands in the study area were by redgram (90.8 %) and

soybean (9.2 %) which are taken during Kharif season respectively (Table 16 and Figure 9).

**Table 16: Present cropping pattern and cropping intensity in Nirgudi west Microwatershed**

Crops	% to Grand Total	
	Kharif	Grand Total
Redgram	90.8	90.8
Soybean	9.2	9.2
<b>Grand Total</b>	100	100

### Economic land evaluation

The main purpose to characterise the socio-economic systems in the watershed is to identify the existing production constraints and propose the potential/alternate options for agro-technology transfer and for bridging the adoption and yield gap.

In Nirgudi west Microwatershed, 6 soil series are identified and mapped (Table 17). The distribution of major soil series are Novinihala covering an area around 140.5 ha (27.9 %) followed by Dinsi 140.1 ha (27.8%), margutti 116.4 ha (23.1%), Bhimanahalli 20.3 ha (4.0 %), Kamalapur 9.0 ha (1.8 %) and Rajnala 40.0 ha (7.9 %).

**Table 17: Distribution of soil series in Nirgudi West Microwatershed**

Sl. No	Soil series	Description	Area in ha (%)
1	MGT	Very shallow, black gravelly clay soils developed from weathered basalt on very gently sloping uplands; clay surface on 1-3 % slope, moderately eroded, slightly gravelly, 15-35 per cent gravels.	116.5 (23.1)
2	BHI	Shallow, black clay soils developed from weathered basalt on very gently sloping uplands; clay surface on 1-3% slope, moderately eroded, slightly gravelly, 15-35 per cent gravels.	20.3 (4.0)
3	NHA	Shallow, black clayey soils developed from weathered basalt on very gently sloping uplands; clay surface on 1-3% slope, slightly eroded.	140.5 (27.9)
4	DSI	Moderately shallow, black clayey soils developed from weathered basalt on nearly level uplands; clay surface on 0-1% slope, slightly eroded.	140.2 (27.8)
5	KMP	Moderately deep, black clayey soils developed from weathered basalt on very gently sloping uplands; clay surface on 1-3% slope, moderately eroded, slightly gravelly, 15-35 per cent gravels.	9.0 (1.8)
6	RNL	Deep, black clayey soils developed from weathered basalt on nearly level uplands; clay surface on 0-1% slope, slightly eroded	40.0 (7.9)
Habitation			38.1 (7.6)

Present cropping pattern on different soil series are given in Table 18. Crops grown on Margutti soils are redgram. Redgram on Kalamundargi soils is grown. Soybean

is grown on Novinihala soils. Redgram on Dinsi soils are grow. Redgram on Rajnala soils are grow.

**Table 18: Cropping pattern on major soil series in Nirgudi west Microwatershed (Area in per cent)**

Soil Series	Soil Depth	Crops	Dry	Grand Total
			Kharif	
MGT	Very shallow (<25 cm)	Redgram	100	100
KGI	Shallow (25-50 cm)	Redgram	100	100
NHA	Shallow (25-50 cm)	Soybean	100	100
DSI	Moderately shallow (50-75 cm)	Redgram	100	100
RNL	Deep (100-150 cm)	Redgram	100	100

Land is used for agricultural use for growing cereals, pulse, oilseeds and commercial crops. The soil/ land potential are measures in terms of physical yield and net income. The alternative land use options for each micro-watershed are given below (Table 19).

**Table19: Alternative land use options for different size group of farmers (Benefit Cost Ratio) in Nirgudi West Microwatershed.**

Soil Series	Small Farmers	Medium Farmers	Large Farmers
MGT	Redgram (1.9)	Redgram(1.7)	
KGI		Redgram (2.1)	
NHA		Soybean (1.2)	
DSI	Redgram (1.4)	Redgram (2.9)	
RNL		Redgram (2.9)	Redgram (2.1)

The productivity of different crops grown in Nirgudi west micro-watershed under potential yield of the crops is given in Table 20.

The data on cost of cultivation and benefit cost ratio (BCR) of different crops is given in Table 20. The total cost of cultivation in study area for red gram ranges between Rs.27862/ha in DSI soil (with BCR of 2.19) and Rs.13850/ha in RNL soil (with BCR of 2.53) and soybean cost of cultivation in NHA soil is Rs 2290/ha (with of BCR 1.29).

The data on FYM, Nitrogen, Phosphorus and Potash application by the farmers to different crops and recommended FYM for different crops is given in Table 20. There is a huge gap between FYM application by farmers and recommended FYM in all the crops across the soils. There is a larger yield gap in crops grown across different soil series. Adequate knowledge about recommended package of practices is the pre-requisite for their use in cultivation of crops. It is a fact that, recommended practices are major contributing factors to yield. Inadequate knowledge about recommended practices leads to their improper adoption. Strengthening of extension services by concerned agency is required to increase adoption of recommended cultivation practices and ultimately reducing the gap. By adopting soil-test fertiliser recommendation, there is scope to

increase yield and income to a maximum of Rs 46217 in soybean and a minimum of Rs 4787 in red gram cultivation.

**Table 20: Economic land evaluation and bridging yield gap for different crops in Nirgudi West Microwatershed**

Particulars	MGT (<25 cm)	KGI (25-50cm)	NHA (25-50 cm)	DSI (5075cm )	RNL (100-150cm)
	Redgram	Redgram	Soybean	Redgram	Redgram
Total cost (Rs/ha)	20812	18787	2290	27862	13850
Gross Return (Rs/ha)	38955	41167	2952	47856	35479
Net returns (Rs/ha)	18143	22380	662	19995	21628
BCR	1.87	2.19	1.29	2.19	2.53
<b>Farmers Practices(FP)</b>					
FYM (t/ha)	2.2	1.7	0.0	3.3	1.6
Nitrogen (kg/ha)	78.1	80.0	0.0	93.3	84.1
Phosphorus (kg/ha)	56.1	57.5	0.0	67.1	58.9
Potash (kg/ha)	0.0	0.0	0.0	0.0	0.0
Grain (Qtl/ha)	9.9	10.4	1.0	12.5	9.1
Price of Yield (Rs/Qtl)	4000	4000	3000	3875	4250
<b>Soil test based fertilizer Recommendation(STBR)</b>					
FYM (t/ha)	7.4	7.4	6.2	7.4	7.4
Nitrogen (kg/ha)	18.5	18.5	20.4	21.6	24.7
Phosphorus (kg/ha)	61.8	61.8	72.6	61.8	55.6
Potash (kg/ha)	21.6	18.5	23.2	23.2	21.6
Grain (Qtl/ha)	12.4	12.4	19.8	12.4	12.4
<b>% of Adoption/yield gap (STBR-FP) / (STBR)</b>					
FYM (%)	70.4	77.5	100.0	55.9	78.9
Nitrogen (%)	-321.6	-331.8	100.0	-331.8	-240.4
Phosphorus (%)	9.1	6.9	100.0	-8.6	-5.9
Potash (%)	100.0	100.0	100.0	100.0	100.0
Grain (%)	20.2	15.7	95.0	-1.2	26.7
<b>Value of yield and Fertilizer (Rs)</b>					
Additional Cost (Rs/ha)	5184	5563	10075	3507	5424
Additional Benefits(Rs/ha)	9971	7733	56292	-581	13996
Net change Income (Rs/ha)	4787	2170	46217	-4088	8572

Economic valuation of Ecosystem Services (ES) was aimed at combining use and non-use values to determine Total Economic Value (TEV) of ES. Ecosystem Services (ES) were valued based on their annual flow or utilization in common monetary units, Rs/year. The valuation of ES was based on market price in 2017 or market cost approaches whichever is applicable, and in other cases on value or benefit transfer from previous valuation studies.

The onsite cost of different soil nutrients lost due to soil erosion is given in Table 21 and Figure 9. The average value of soil nutrient loss is around Rs 3764 per ha/year. The total cost of annual soil nutrients is around Rs 1754112 per year for the total area of 504.6 ha.

**Table 21: Estimation of onsite cost of soil erosion in Nirgudi west micro-watershed**

Particulars	Quantity(kg)		Value (Rs)	
	Per ha	Total	Per ha	Total
Organic matter	524.25	244300	3302.76	1539088
Phosphorous	0.10	48	4.54	2113
Potash	4.01	1869	80.22	37384
Iron	0.43	201	20.74	9665
Manganese	0.95	442	260.55	121414
Cupper	0.15	69	82.40	38397
Zinc	0.02	8	0.67	311
Sulpher	0.30	139	11.93	5560
Boron	0.01	4	0.38	179
Total	530.21	247080	3764.19	1754112




Figure 9: Estimation of onsite cost of soil erosion in Nirgudi West Microwatershed

The average value of ecosystem service for food grain production is around Rs.10815/ ha/year (Table 22 and Figure 11). Per hectare food grain production services is maximum in redgram (Rs. 20967) and soybean (Rs.662).

**Table 22: Ecosystem services of food grain production in Nirgudi West Microwatershed**

Production items	Crops	Area in ha	Yield (Qtl/ha)	Price (Rs/Qtl)	Gross Returns (Rs/ha)	Cost of Cultivation (Rs/ha)	Net Returns (Rs/ha)
Pulses	Redgram	20.06	10.79	4000	43140	22173	20967
Oil seeds	Soybean	20.32	0.98	3000	2952	2290	662
Average value		40.38	5.89	3500	23046	12232	10815

The water demand for production of different crops was worked out in arriving at the ecosystem services of water support to crop growth. The data on water requirement for producing one quintal of grain is considered for estimating the total value of water required for crop production. The per hectare value of water used and value of water was maximum (Table 23 and Figure 12) in redgram (Rs. 58714) and soybean (Rs. 2112).

**Table 23: Ecosystem services of water supply in Nirgudi West Microwatershed**

Crops	Yield (Qtl/ha)	Virtual water (cubic meter) per ha	Value of Water (Rs/ha)	Water consumption (Cubic meters/Qtl)
Redgram	10.8	5871	58714	544
Soybean	1.0	211	2112	215
Average value	5.9	3041	30413	380

**Table 24: Farming constraints related land resources of sample households in Nirgudi west Microwatershed**

Sl. No	Particulars	Per cent
1	Less Rainfall	100.0
2	Lack of good quality seeds	70.0
3	Non availability Fertilizers	40.0
4	High Crop Pests & Diseases	60.0
5	Animal Pests & Diseases	20.0
6	Lack of transportation	50.0
7	Lack of storage	100.0
8	Damage of crops by Wild Animals	100.0
9	Non availability of Plant Protection Chemicals	100.0
10	<b>Source of loan</b>	
	Bank	10.0
	Money Leander	90.0
11	<b>Market for selling</b>	
	Regulated	10.0
	Village market	90.0
12	<b>Sources of Agri-Technology information</b>	
	Mobile	10.0
	Newspaper	80.0
	Television	10.0

The main farming constraints in Nirgudi West Microwatershed to be found are less rainfall, lack of good quality seeds, lack of storage, damage of crops by wild animals

and non availability of plant protection chemicals. Majority of farmers depend up on bank and money lender of the sources of loan for purpose of crop production. Farmers to sell the agriculture produce through regulated and village market and the farmers getting the agriculture related information on newspaper, mobile and television. Farmers reported that they are not getting timely support/extension services from the concerned development department (Table 24).

The findings of the study would be very much useful to the planners and policy makers of the study area to identify the irrationality in the existing production pattern and to suggest appropriate production plans for efficient utilization of their scarce resources resulting in increased net farm incomes and employment. The study also throws light on future potentialities of increasing net farm income and employment under different situations viz., with existing and recommended technology.