



हर कदम, हर डगर
किसानों का हमसफर
भारतीय कृषि अनुसंधान परिषद

AgriSearch with a human touch

**LAND RESOURCE INVENTORY AND SOCIO-ECONOMIC STATUS OF
FARM HOUSEHOLDS FOR WATERSHED PLANNING AND
DEVELOPMENT**

PADASAVLI-1 (4D5C2E1a) MICROWATERSHED

Aland Taluk, Gulbarga District, Karnataka

Karnataka Watershed Development Project – II

SUJALA – III

World Bank funded Project



ICAR – NATIONAL BUREAU OF SOIL SURVEY AND LAND USE PLANNING



ICAR - NBSS & LUP



**WATERSHED DEVELOPMENT DEPARTMENT
GOVT. OF KARNATAKA, BANGALORE**



About ICAR - NBSS&LUP

The ICAR-National Bureau of Soil Survey and Land Use Planning (ICAR-NBSS&LUP), Nagpur, a premier Institute of the Indian Council of Agricultural Research (ICAR), was set up during 1976 with the objective to prepare soil resource maps at national, state and district levels and to provide research inputs in soil resource mapping and its applications, land evaluation, land use planning, land resource management, and database management using GIS for optimising land use on different kinds of soils in the country.

The Bureau has been engaged in carrying out soil resource survey, agro-ecological and soil degradation mapping at the country, state and district levels for qualitative assessment and monitoring the soil health towards viable land use planning. The research activities have resulted in identifying the soil potentials and problems, and the various applications of the soil surveys with the ultimate objective of sustainable agricultural development. The Bureau has the mandate to correlate and classify soils of the country and maintain a National Register of all the established soil series. The Institute is also imparting in-service training to staff of the soil survey agencies in the area of soil survey, land evaluation and soil survey interpretations for land use planning. The Bureau in collaboration with Panjabrao Krishi Vidyapeeth, Akola is running post-graduate teaching and research programme in land resource management, leading to M.Sc. and Ph.D. degrees.

Citation: Rajendra Hegde, Ramesh Kumar, S.C., K.V. Niranjana, S. Srinivas, M.Lalitha, B.A. Dhanorkar, R.S. Reddy and S.K. Singh (2019). "Land Resource Inventory and Socio-Economic Status of Farm Households for Watershed Planning and Development of Padasavli-1(4D5C2E1a) Microwatershed, Aland Taluk, Gulbarga District, Karnataka", ICAR-NBSS&LUP Sujala MWS Publ.13, ICAR – NBSS & LUP, RC, Bangalore. p.85 & 28.

TO OBTAIN COPIES,

Please write to:

Director, ICAR - NBSS & LUP,

Amaravati Road, NAGPUR - 440 033, India

Phone : (0712) 2500386, 2500664, 2500545 (O)

Telefax : 0712-2522534

E-Mail : director@nbsslup.ernet.in

Website URL : nbsslup.in

Or

Head, Regional Centre, ICAR - NBSS&LUP, Hebbal, Bangalore - 560 024

Phone : (080) 23412242, 23510350 (O)

Telefax : 080-23510350

E-Mail : nbssrcb@gmail.com



**LAND RESOURCE INVENTORY AND SOCIO-ECONOMIC
STATUS OF FARM HOUSEHOLDS FOR WATERSHED
PLANNING AND DEVELOPMENT**

PADASAVLI-1 (4D5C2E1a) MICROWATERSHED

Aland Taluk, Gulbarga District, Karnataka

Karnataka Watershed Development Project – II

Sujala-III

World Bank funded Project



**ICAR – NATIONAL BUREAU OF SOIL SURVEY AND LAND USE
PLANNING**



**WATERSHED DEVELOPMENT DEPARTMENT, GOVT. OF
KARNATAKA, BANGALORE**



PREFACE

In Karnataka, as in other Indian States, the livelihoods of rural people are intertwined with farming pursuits. The challenges in agriculture are seriously threatening the livelihood of a large number of farmers as they have been practicing farming in contextual factors beyond their control. Climatic factors are the most important ones and have become much more significant in recent times due to rapid climate changes induced by intensive anthropogenic activities affecting our ecosystem in multiple ways. Climate change has become the reality, it is happening and efforts to evolve and demonstrate climate resilient technologies have become essential. Due to the already over stressed scenario of agrarian sector, the climate change is resulting in manifold increase in the complexities, pushing the rural mass to face more and more unpredictable situations. The rising temperatures and unpredictable rainfall patterns are going to test seriously the informed decisions farmers have to make in order to survive in farming and sustain their livelihood.

It is generally recognized that impacts of climate change shall not be uniform across the globe. It is said that impact of climate change is more severe in South Asia. Based on the analysis of meteorological data, it is predicted that in India, there will be upward trend in mean temperature, downward trend in relative humidity, annual rainfall and number of wet days in a year. Also, in general, phenomena like erratic monsoon, spread of tropical diseases, rise in sea levels, changes in availability of fresh water, frequent floods, droughts, heat waves, storms and hurricanes are predicted. Each one of these adverse situations are already being experienced in various parts of India and also at the global level. Decline in agricultural productivity of small and marginal farmers becoming more vulnerable is already witnessed.

In Karnataka, more than 60 per cent of the population live in rural areas and depend on agriculture and allied activities for their livelihood. Though the state has achieved significant progress in increasing the yield of many crops, there is tremendous pressure on the land resources due to the growing and competing demands of various land uses. This is reflected in the alarming rate of land degradation observed. Already more than 50 per cent of the area is affected by various forms of degradation. If this trend continues, the sustainability of the fragile ecosystem will be badly affected. The adverse effects of change in the climatic factors are putting additional stress on the land resources and the farmers dependent on this.

The natural resources (land, water and vegetation) of the state need adequate and constant care and management, backed by site-specific technological interventions and investments particularly by the government. Detailed database pertaining to the nature of the land resources, their constraints, inherent potentials and suitability for various land based rural enterprises, crops and other uses is a prerequisite for preparing location-specific action plans, which are in tune with the inherent capability of the resources. Any effort to evolve climate resilient technologies has to be based on the baseline scientific database. Then only one can expect effective implementation of climate resilient technologies, monitor the progress, make essential review of the strategy, and finally evaluate the effectiveness of the implemented programs. The information available at present on the land resources of the state are of general nature and useful only for general purpose

planning. Since the need of the hour is to have site-specific information suitable for farm level planning and detailed characterization and delineation of the existing land resources of an area into similar management units is the only option.

ICAR-NBSS&LUP, Regional Centre, Bangalore has taken up a project sponsored by the Karnataka Watershed Development Project-II, (Sujala-III), Government of Karnataka funded by the World Bank under Component -1 Land Resource Inventory. This study was taken up to demonstrate the utility of such a database in reviewing, monitoring and evaluating all the land based watershed development programs on a scientific footing. To meet the requirements of various land use planners at grassroots level, the present study on “Land Resource Inventory and Socio-Economic Status of Farm Households for Watershed Planning and Development of Padasavli-1 Microwatershed, Aland Taluk, Kalaburgi District, Karnataka” for integrated development was taken up in collaboration with the State Agricultural Universities, IISC, KRSRAC, KSNDMC as Consortia partners. The project provides detailed land resource information at cadastral level (1:7920 scale) for all the plots and socio-economic status of farm households covering thirty per cent farmers randomly selected representing landed and landless class of farmers in the microwatershed. The project report with the accompanying maps for the microwatershed will provide required detailed database for evolving effective land use plan, alternative land use options and conservation plans for the planners, administrators, agricultural extension personnel, KVK officials, developmental departments and other land users to manage the land resources in a sustainable manner.

It is hoped that this database will be useful to the planners, administrators and developmental agencies working in the area in not only for formulating location specific developmental schemes but also for their effective monitoring at the village/watershed level.

Nagpur

Date:

S.K. SINGH

Director, ICAR - NBSS&LUP, Nagpur

Contributors

Dr. RajendraHegde Principal Scientist, Head & Project Leader, Sujala-III Project ICAR-NBSS&LUP, Regional Centre Bangalore	Dr. S.K.Singh Director, ICAR-NBSS&LUP Coordinator, Sujala-III Project Nagpur
Soil Survey, Mapping & Report Preparation	
Dr. B.A. Dhanorkar	Sh. R.S. Reddy
Dr. K.V. Niranjana	Sh. Nagendra, B.R.
	Smt. Chaitra, S.P.
Field Work	
Sh. C.BacheGowda	Sh. Mahesha, D.B.
Sh. Somashekar	Sh. Ashok S. Sindagi
Sh. VenkataGiriyappa	Sh. Veerabhadrapa
Sh. M. Jayaramaiah	Sh. Anand
Sh. Paramesha, K.	Sh. Arun N Kambar
	Sh. Shankarappa, K.
	Sh. Kamalesh K. Avate
	Sh. Sharan Kumar Huppar
	Sh. Yogesh, H.N.
	Sh. Kalaveerachari R. Kammar
GIS Work	
Dr. S.Srinivas	Sh. A.G.Devendra Prasad
Dr. M.Ramesh	Sh. Prakashanaik, M.K.
Sh. D.H.Venkatesh	Sh. AbhijithSastry, N.S.
Smt.K.Sujatha	Sh. Sudip Kumar Suklabaidya
Smt. K.V.Archana	Sh. Mahamad Ali, M.
Sh. N.Maddileti	Sh. Avinash, K.N.
	Sh. Amar Suputhra, S
	Sh. Anudeep, Y.
	Sh. Deepak, M.J.
	Smt. K.Karunya Lakshmi
	Ms. Seema, K.V.
	Ms. A. Rajab Nisha

Laboratory Analysis	
Dr. K.M.Nair	Dr. H.R. Savitha
Smt. Arti Koyal	Ms. Steffi Peter
Smt. Parvathy, S.	Ms. Thara, V.R
	Ms. Roopa, G.
	Ms. Swati, H.
	Sh. Shantaveera Swami
	Ms. Shwetha, N.K.
	Smt. Ishrat Haji
	Ms. P. PavanKumari
	Ms. Padmaja
	Ms. Veena, M.
Soil & Water Conservation	
Sh. Sunil P. Maske	
Socio-Economic Analysis	
Dr. S.C. Ramesh Kumar	Sh. M. K. Prakashanaik
	Ms. Sowmya K.B
	Sh.Manjunath M
	Sh.Veerabhadraswamy R
	Sh.Lankesh RS
	Sh.Kalaveerachari R Kammar
	Sh.Pradyumma U
	Sh.Yogesha HN
	Sh.Vijay kumar lamani
	Sh.Arun N Kambar
	Sh.Vinay
	Sh.Basavaraj.Biradar
	Sh.Vinod R
	Sh.Praveenkumar P Achalkar
	Sh.Rajendra D
Watershed Development Department, GoK, Bangalore	
Sh. Rajeev Ranjan IFS Project Director & Commissioner, WDD	Dr. A. Natarajan NRM Consultant, Sujala-III Project
Dr. S.D. Pathak IFS Executive Director & Chief Conservator of Forests, WDD	

PART-A

LAND RESOURCE INVENTORY

Contents

Preface		
Contributors		
Executive Summary		
Chapter 1	Introduction	1
Chapter 2	Geographical Setting	3
2.1	Location and Extent	3
2.2	Geology	4
2.3	Physiography	4
2.4	Drainage	4
2.5	Climate	5
2.6	Natural Vegetation	6
2.7	Land Utilization	7
Chapter 3	Survey Methodology	9
3.1	Base maps	9
3.2	Field Investigation	11
3.3	Laboratory Characterization	12
3.4	Finalization of Soil Map	13
Chapter 4	The Soils	19
4.1	Soils of Basalt Landscape	19
Chapter 5	Interpretation for Land Resource Management	27
5.1	Land Capability Classification	27
5.2	Soil Depth	29
5.3	Surface Soil Texture	30
5.4	Soil Gravelliness	31
5.5	Available Water Capacity	32
5.6	Soil Slope	33
5.7	Soil Erosion	34
Chapter 6	Fertility Status	37
6.1	Soil Reaction (pH)	37
6.2	Electrical Conductivity (EC)	37
6.3	Organic Carbon (OC)	37
6.4	Available Phosphorus	39
6.5	Available Potassium	39
6.6	Available Sulphur	39
6.7	Available Boron	39
6.8	Available Iron	42
6.9	Available Manganese	42
6.10	Available Copper	42
6.11	Available Zinc	42

Chapter 7	Land Suitability for Major Crops	45
7.1	Land suitability for Sorghum	45
7.2	Land suitability for Maize	48
7.3	Land suitability for Red gram	50
7.4	Land suitability for Sunflower	51
7.5	Land suitability for Cotton	53
7.6	Land suitability for Sugarcane	54
7.7	Land suitability for Soybean	56
7.8	Land suitability for Guava	57
7.9	Land suitability for Mango	58
7.10	Land suitability for Sapota	60
7.11	Land suitability for Jackfruit	62
7.12	Land suitability for Jamun	62
7.13	Land Suitability for Musambi	63
7.14	Land Suitability for Lime	64
7.15	Land Suitability for Cashew	66
7.16	Land Suitability for Custard Apple	66
7.17	Land Suitability for Amla	67
7.18	Land Suitability for Tamarind	68
7.19	Land Management Units	69
7.20	Proposed Crop Plan	71
Chapter 8	Soil Health Management	73
Chapter 9	Soil and Water conservation Treatment Plan	77
9.1	Treatment Plan	78
9.2	Recommended Soil and Water Conservation measures	81
9.3	Greening of microwatershed	82
	References	85
	Appendix I	I-VIII
	Appendix II	IX-XIV
	Appendix III	XV-XVII

LIST OF TABLES

2.1	Mean Monthly Rainfall, PET, 1/2 PET at Aland Taluk, Kalaburgi District	5
2.2	Land Utilization in Aland Taluk	7
3.1	Differentiating Characteristics used for Identifying Soil Series	12
3.2	Soil Legend	17
7.1	Soil-Site Characteristics of Padasavli-1 microwatershed	47
7.2	Crop suitability criteria for Sorghum	48
7.3	Crop suitability criteria for Maize	49
7.4	Crop suitability criteria for Red gram	50
7.5	Crop suitability criteria for Sunflower	52
7.6	Crop suitability criteria for Cotton	53
7.7	Crop suitability criteria for Sugarcane	55
7.8	Crop suitability criteria for Guava	57
7.9	Crop suitability criteria for Mango	59
7.10	Crop suitability criteria for Sapota	61
7.11	Crop suitability criteria for Lime	65
7.12	Proposed Crop Plan for Padasavli-1 Microwatershed	71

LIST OF FIGURES

2.1	Location map of Padasavli-1 microwatershed	3
2.2	Rock formations in Padasavli-1 microwatershed	4
2.3	Rainfall distribution in Aland Taluk, Kalaburgi District	6
2.4	Natural Vegetation (Scrub) of Padasavli-1 microwatershed	6
2.5	Current Land use – Padasavli-1 microwatershed	8
2.6	Location of Wells- Padasavli-1 microwatershed	8
3.1	Scanned and Digitized Cadastral map of Padasavli-1 microwatershed	10
3.2	Satellite image of Padasavli-1 microwatershed	10
3.3	Cadastral map overlaid on IRS PAN+LISS IV merged imagery of Padasavli-1 microwatershed	11
3.4	Soil phase or management units of Padasavli-1 microwatershed	15
5.1	Land Capability Classification of Padasavli-1 microwatershed	28
5.2	Soil Depth map of Padasavli-1 microwatershed	30
5.3	Surface Soil Texture map of Padasavli-1 microwatershed	31
5.4	Soil Gravelliness map of Padasavli-1 microwatershed	32
5.5	Soil Available Water Capacity map of Padasavli-1 microwatershed	33
5.6	Soil Slope map of Padasavli-1 microwatershed	34
5.7	Soil Erosion map of Padasavli-1 microwatershed	35
6.1	Soil Reaction (pH) map of Padasavli-1 microwatershed	38
6.2	Electrical Conductivity (EC) map of Padasavli-1 microwatershed	38
6.3	Soil Organic Carbon (OC) map of Padasavli-1 microwatershed	40
6.4	Soil Available Phosphorus map of Padasavli-1 microwatershed	40
6.5	Soil Available Potassium map of Padasavli-1 microwatershed	41
6.6	Soil Available Sulphur map of Padasavli-1 microwatershed	41
6.7	Soil Available Boron map of Padasavli-1 microwatershed	42
6.8	Soil Available Iron map of Padasavli-1 microwatershed	43
6.9	Soil Available Manganese map of Padasavli-1 microwatershed	43
6.10	Soil Available Copper map of Padasavli-1 microwatershed	44
6.11	Soil Available Zinc map of Padasavli-1 microwatershed	44
7.1	Land Suitability map of Sorghum	46
7.2	Land Suitability map of Maize	49
7.3	Land Suitability map of Red gram	51

7.4	Land Suitability map of Sunflower	52
7.5	Land Suitability map of Cotton	54
7.6	Land Suitability map of Sugarcane	55
7.7	Land Suitability map of Soybean	56
7.8	Land Suitability map of Guava	58
7.9	Land Suitability map of Mango	60
7.10	Land Suitability map of Sapota	61
7.11	Land Suitability map of Jackfruit	62
7.12	Land Suitability map of Jamun	63
7.13	Land Suitability map of Musambi	64
7.14	Land Suitability map of Lime	65
7.15	Land Suitability map of Cashew	66
7.16	Land Suitability map of Custard Apple	67
7.17	Land Suitability map of Amla	68
7.18	Land Suitability map of Tamarind	69
7.19	Land Management Units map of Padasavli-1 microwatershed	70
9.1	Soil and Water Conservation map of Padasavli-1 microwatershed	82

EXECUTIVE SUMMARY

The land resource inventory of Padasavli-1 microwatershed was conducted using village cadastral maps and IRS satellite imagery on 1:7920 scale. The false colour composites of IRS imagery were interpreted for physiography and these physiographic delineations were used as base for mapping soils. The soils were studied in several transects and a soil map was prepared with phases of soil series as mapping units. Random checks were made all over the area outside the transects to confirm and validate the soil map unit boundaries. The soil map shows the geographic distribution and extent, characteristics, classification and use potentials of the soils in the microwatershed.

The present study covers an area of 535 ha in Padasavli-1 microwatershed in Aland taluk of Kalaburgi district, Karnataka. The climate is semiarid and categorized as drought prone with an average annual rainfall of 786 mm of which about 595 mm is received during south –west monsoon, 116 mm during north-east and the remaining 75 mm during the rest of the year. An area of about 95 per cent is covered by soils, five per cent by waterbodies, settlements and others. The salient findings from the land resource inventory are summarized briefly below.

- ❖ *The soils belong to 7 soil series and 16 soil phases (management units) and 5 land management units.*
- ❖ *The length of crop growing period is about 150 days starting from the 3rd week of June to 1st week of October.*
- ❖ *From the master soil map, several interpretative and thematic maps like land capability, soil depth, surface soil texture, soil gravelliness, available water capacity, soil slope and soil erosion were generated.*
- ❖ *Soil fertility status maps for macro and micronutrients were generated based on the surface soil samples collected at every 250 m grid interval.*
- ❖ *Land suitability for growing major agricultural and horticultural crops were assessed and maps showing degree of suitability along with constraints were generated.*
- ❖ *About 95 per cent area is suitable for agriculture*
- ❖ *About 11 per cent of the soils are very deep (>150 cm) to moderately deep (75-100 cm), 73 per cent are moderately shallow to shallow (25-75 cm) and about 11 per cent are very shallow (<25 cm) soils.*
- ❖ *About 95 per cent of the area has clayey soils at the surface.*
- ❖ *About 17 per cent of the area has non-gravelly soils, 55 per cent gravelly soils (15-35 % gravel) and 24 per cent has very gravelly (35- 60% gravel) soils.*
- ❖ *About 9 per cent of the area has soils that are very high (>200mm/m) in available water capacity, 18 per cent medium (100-150 mm/m) and about 68 per cent low (50-100 mm/m) and very low (<50mm/m).*
- ❖ *About 87 per cent of the area has nearly level (0-1% slope) to very gently sloping (1-3% slope) lands and about 8 per cent area is gently (3-5% slope) sloping lands.*
- ❖ *An area of about 6 per cent has soils that are slightly eroded (e1), 80 per cent moderately eroded (e2) and 9 per cent severely eroded (e3).*
- ❖ *An area of about 66 per cent has soils that are moderately alkaline to strongly alkaline (pH 7.8 to 9.0), about 20 per cent slightly alkaline (pH 7.3-7.8) and 9 per cent neutral (pH 6.5-7.3)*
- ❖ *The Electrical Conductivity (EC) of the soils are dominantly <2 dsm⁻¹ indicating that the soils are non-saline.*
- ❖ *About 73 per cent medium (0.5-0.75%), 9 per cent high (>0.75%) and 14 per cent low (<0.5%) in organic carbon.*
- ❖ *Major area of 78 per cent has soils that are low (<23 kg/ha), 15 per cent medium (23-57 kg/ha) and 2 per cent high (>57 kg/ha) in available phosphorus.*

- ❖ About 23 per cent medium (145-337 kg/ha) and 72 per cent high (>337 kg/ha) in available potassium.
- ❖ Available sulphur is low (<10 ppm) in about 60 per cent area, medium (10-20 ppm) in 30 per cent and 5 per cent high (>20 ppm).
- ❖ Available boron is low (<0.5 ppm) in about 46 per cent area, 35 per cent medium (0.5-1.0 ppm) and 15 per cent high (>1 ppm) in available boron.
- ❖ About 8 per cent area has soils that are deficient (<4.5 ppm) in available iron and 87 per cent sufficient (>0.6 ppm).
- ❖ Available manganese and copper are sufficient in all the soils.
- ❖ About 69 per cent area has soils that are deficient (<0.6 ppm) in available zinc and 26 per cent sufficient (>0.6 ppm).
- ❖ The land suitability for 18 major crops (agricultural and horticultural) grown in the microwatershed were assessed and the areas that are highly suitable (S1) and moderately suitable (S2) are given below. It is however to be noted that a given soil may be suitable for various crops but what specific crop to be grown may be decided by the farmer looking to his capacity to invest on various inputs, marketing infrastructure, price, and finally the demand and supply position.

Land suitability for various crops in the microwatershed

Crop	Suitability Area in ha (%)		Crop	Suitability Area in ha (%)	
	Highly suitable (S1)	Moderately suitable (S2)		Highly suitable (S1)	Moderately suitable (S2)
Sorghum	33 (6)	131(25)	Sapota	-	-
Maize	-	-	Jackfruit	-	-
Red gram	-	164 (31)	Jamun	-	49 (9)
Sunflower	33(6)	24 (5)	Musambi	33(6)	24 (5)
Cotton	33 (6)	131 (25)	Lime	33(6)	24 (5)
Sugarcane	-	-	Cashew	-	-
Soybean	33(6)	131 (25)	Custard apple	33(6)	131(25)
Guava	-	-	Amla	33(6)	131(25)
Mango	-	-	Tamarind	-	49 9)

Apart from the individual crop suitability, a proposed crop plan has been prepared for the 5 identified LMUs by considering only the highly and moderately suitable lands for different crops and cropping systems with food, fibre and horticulture crops that helps in maintaining the ecological balance in the microwatershed.

- ❖
- ❖ Maintaining soil-health is vital to crop production and conserve soil and land resource base for maintaining ecological balance and to mitigate climate change. For this, several ameliorative measures have been suggested to these problematic soils like saline/alkali, highly eroded, sandy soils etc.,
- ❖ Soil and water conservation treatment plan has been prepared that would help in identifying the sites to be treated and also the type of structures required.
- ❖ As part of the greening programme, several tree species have been suggested to be planted in marginal and submarginal lands and also in the field bunds, hillocks, mounds and ridges.

INTRODUCTION

Soil is a finite natural resource that is central to sustainable agriculture and food security. Over the years, this precious resource is faced with the problems of erosion, salinity, alkalinity, degradation, depletion of nutrients and even decline in availability of land for agriculture. It is a known fact, that it takes thousands of years to form a few centimetres of soil, thus, soil is a precious gift of nature. The area available for agriculture is about 51 per cent of the total geographical area and more than 60 per cent of the people are still dependant on agriculture for their livelihood. However, the capacity of a soil to produce is limited and the limits to the production are set by its intrinsic characteristics, agro-climatic setting, and use and management. There is, therefore, tremendous pressure on land and water resources, which is causing decline in soil-health and stagnation in productivity. As much as 121 m ha of land is reportedly degraded which leads to impaired soil quality. It is imperative that steps are urgently taken to check and reverse land degradation without any further loss of time. The improvements in productivity will have to come from sustainable intensification measures that make the most effective use of land and water resources. Soil erosion alone has degraded about 35 lakh ha. Almost all the uncultivated areas are facing various degrees of degradation, particularly soil erosion; salinity and alkalinity has emerged as a major problem (>3.5 lakh ha) in the irrigated areas of the State. Nutrient depletion and declining factor productivity is common in both rainfed and irrigated areas. The degradation is continuing at an alarming rate and there appears to be no systematic effort among the stakeholders to contain this process. In recent times, an aberration of weather due to climate change phenomenon has added another dimension leading to unpredictable situations to be tackled by the farmers.

In this critical juncture, the challenge before us is not only to increase the productivity per unit area which is steadily declining and showing a fatigue syndrome, but also to prevent or at least reduce the severity of degradation. If the situation is not reversed at the earliest, then the sustainability of the already fragile crop production system and the overall ecosystem will be badly affected in the state. Added to this, every year there is a significant diversion of farm lands and water resources for non-agricultural purposes. Thus, developing strategies to slow down the degradation process or reclaim the soils to normal condition and ensure sustainability of production system are the major issues today. This demands a systematic appraisal of our soil and land resources with respect to their extent, geographic distribution, characteristics, behaviour and uses potential, which is very important for developing an effective land use and cropping systems for augmenting agricultural production on a sustainable basis.

The soil and land resource inventories made so far in Karnataka had limited utility because the surveys were of different types, scales and intensities carried out at different times with specific objectives. Hence, there is an urgent need to generate detailed site-specific farm level database on various land resources for all the villages/watersheds in a time

bound manner that would help to protect the valuable soil and land resources and also to stabilize the farm production. Therefore, the land resource inventory required for farm level planning is the one which investigates all the parameters which are critical for productivity *viz.*, soils, site characteristics like slope, erosion, gravelliness and stoniness, climate, water, topography, geology, hydrology, vegetation, crops, land use pattern, animal population, socio-economic conditions, infrastructure, marketing facilities and various schemes and developmental works of the government etc. From the data collected at farm level, the specific problems and potentials of the area can be identified and highlighted, conservation measures required for the area can be planned on a scientific footing, suitability of the area for various uses can be worked out and finally viable and sustainable land use options suitable for each and every land holding can be prescribed.

The Land Resource Inventory is basically done for identifying potential and problem areas, developing sustainable land use plans, estimation of surface run off and water harvesting potential, preparation of soil and water conservation plans, land degradation/desertification etc. The Bureau is presently engaged in developing an LRI methodology using high resolution satellite remote sensing data and Digital Elevation Model (DEM) data to prepare Landscape Ecological Units (LEU) map representing agro-ecosystem as a whole. The LEU is preferred over landform as the base map for LRI. LEU is the assemblage of landform, slope and land use. An attempt has already been made to upscale the soil resource information from 1:250000 and 1:50000 scale to the LEU map in Goa and other states. Here, an attempt will be made to uplink the LRI data generated under Sujala-III Project to the Landscape Ecological Units (LEUs) map. For this, the major physiographic region, *i.e.*, South Deccan Plateau will be taken as an example.

The land resource inventory aims to provide site specific database for Padasavli-1 microwatershed in Aland Taluk, Kalaburgi District, Karnataka state for the Karnataka Watershed Development Department. The database was generated by using cadastral map of the village as a base along with high resolution IRS LISS IV and Cartosat-1 merged satellite imagery. Later, an attempt will be made to uplink this LRI data generated at 1:7920 scale under Sujala-III Project to the proposed Landscape Ecological Units (LEUs) map.

The study was organized and executed by the ICAR- National Bureau of Soil Survey and Land Use Planning, Regional Centre, Bangalore under Generation of Land Resource Inventory Data Base Component-1 of the Sujala-III Project funded by the World Bank.

GEOGRAPHICAL SETTING

2.1 Location and Extent

The study area of Padasavli-1 microwatershed (Padasavli subwatershed) is located in the northeastern part of Karnataka in Aland Taluk, Kalaburgi District, Karnataka State (Fig.2.1). It comprises parts of Sarasamba and Padasavli villages. It lies between $17^{\circ} 35'$ and $17^{\circ} 36'$ North latitude and $76^{\circ} 25'$ and $76^{\circ} 28'$ east longitude and covers an area of 535 ha. It is about 15 km south of Kalaburgi and is surrounded by Nagalgaon village on the south, Matki village on the northeast, Chincholi khurd village on the north and Hiroli village on the west.

LOCATION MAP OF PADASAVLI-1 MICRO-WATERSHED




Fig.2.1 Location map of Padasavli-1 Microwatershed

2.2 Geology

Major rock formation observed in the microwatershed is Basalt (Fig.2.2) or Deccan Trap. The Deccan Traps cover the whole of Bidar, parts of Kalaburgi, Bijapur and Belgaum districts. In all, eight lava flows have been identified in Karnataka horizontally overlying the older formations. The thickness of the individual flows averages about five metres. It is relatively uniform in petrographic character. The most common type is augite basalt. Dominant colour is grayish green and texture ranges from cryptocrystalline to glassy. The rock is often vesicular and scoriaceous filled up with secondary minerals like coloured agate, quartz, calcite and a large variety of zeolites. The Deccan Traps form an excellent building material and also used as road-metal and railway ballast.



Fig. 2.2 Basalt rock formation

2.3 Physiography

Physiographically, the area has been identified as Basalt landscape based on geology. The area has been further subdivided into four landforms, viz; mounds/ridges, summits, side slopes and very gently sloping uplands based on slope and its relief features. The elevation ranges from 491 to 559 m. The mounds and ridges are mostly covered by rock outcrops.

2.4 Drainage

The area is drained by several small parallel streams that join Monia nala which further downstream joins Awarja River along its course. Though, it is not a perennial one,

during rainy season it carries large quantities of rain water. The microwatershed has only few small tanks which are not capable of storing the water that flows during the rainy season. Due to this, the ground water recharge is very much affected. This is reflected in the failure of many bore wells in the villages. If the available rain water is properly harnessed by constructing new tanks and recharge structures at appropriate places in the villages, then the drinking and irrigation needs of the entire area can be easily met. The drainage network is parallel to sub parallel and dendritic.

2.5 Climate

The Kalaburgi district lies in the northern plains of Karnataka and falls under semiarid tract of the state and is categorized as drought prone with average annual rainfall of 785 mm (Table 2.1). Of the total rainfall, maximum of 595 mm is received during the south-west monsoon period from June to September, the north-east monsoon from October to early December contributes about 116 mm and the remaining 75 mm during the rest of the year. December is the coldest month with mean daily maximum and minimum temperatures being 29.5⁰C and 15⁰ to 10⁰C respectively. During peak summer, temperatures shoot up to 45⁰C. Relative humidity varies from 26 per cent in summer to 62 per cent in winter. Rainfall distribution is shown in Figure 2.3. The average potential evapotranspiration (PET) is 150 mm and varies from a low of 115 mm in December to 232 mm in the month of May. The PET is always higher than precipitation in all the months except August and September. Generally, the length of crop growing period (LGP) is 150 days and starts from 3rd week of June to third week of November.

Table 2.1 Mean Monthly Rainfall, PET, 1/2 PET at Aland Taluk, Kalaburgi District

Sl. No.	Months	Rainfall	PET	1/2 PET
1	January	7.50	126.80	63.40
2	February	3.40	143.90	71.95
3	March	11.30	189.90	94.95
4	April	19.40	209.80	104.90
5	May	32.70	232.20	116.10
6	June	111.00	186.40	93.20
7	July	139.20	152.80	76.40
8	August	172.40	147.60	73.80
9	September	172.30	131.70	65.85
10	October	91.30	145.50	72.75
11	November	19.30	129.80	64.90
12	December	5.80	114.80	57.40
Total		785.6	149.70	




Fig 2.3 Rainfall distribution in Aland Taluk, Kalaburgi District

2.6 Natural Vegetation

The natural vegetation is sparse comprising few tree species, shrubs and herbs. The mounds, ridges and boulders occupy very sizeable area which is under thin to moderately thick forest vegetation. Still, there are some remnants of the past forest cover which can be seen in patches in some ridges and hillocks in the microwatershed (Fig. 2.4).

Apart from the continuing deforestation, the presence of large population of goats, sheep and other cattle in the microwatershed is causing vegetative degradation of whatever little vegetation left in the area. The uncontrolled grazing has left no time for the regeneration of the vegetative cover. This leads to the accelerated rate of erosion on the hill slopes resulting in the formation of deep gullies in the foot slopes that eventually result in the heavy siltation of tanks and reservoirs in the microwatershed.



Fig. 2.4 Natural Vegetation (Scrub) of Padasavli-1 Microwatershed

2.7 Land Utilization

About 89 per cent area (Table 2.2) in Aland taluk is cultivated at present. An area of about 2 per cent is permanently under pasture, 3 per cent under current fallows and 2 per cent each under non agricultural land and currently barren. Forests occupy an area of about 2 per cent and the tree cover is in a very poor state. Most of the mounds, ridges and bouldery areas have very poor vegetative cover. Major crops grown in the area are sorghum, maize, cotton, sugarcane, safflower, groundnut, red gram and sapota. While carrying out land resource inventory, the land use/land cover particulars are collected from all the survey numbers and a current land use map of the microwatershed is prepared. The current land use map prepared shows the arable and non-arable lands, other land uses and different types of crops grown in the area. The current land use map of Padasavli-1 microwatershed is presented in Fig.2.5.

Table 2.2 Land Utilization in Aland Taluk

Sl. No.	Agricultural land use	Area (ha)	Per cent
1.	Total geographical area	173417	
2.	Total cultivated area	153806	88.69
3.	Area sown more than once	7910	
4.	Trees and grooves	59	0.034
5.	Forest	2854	1.64
6.	Cultivable wasteland	974	0.56
7.	Permanent Pasture land	3469	2.00
8.	Barren land	3142	1.81
9.	Non- Agriculture land	3465	1.99
10.	Current Fallows	5648	3.25

Simultaneously, enumeration of wells (bore wells and open wells) in the microwatershed was made and their location in different survey numbers is located on the cadastral map. Map showing the location of wells and other water bodies in the Padasavli-1 microwatershed is given in Figure 2.6.




Fig.2.5 Current Land Use – Padasavli-1 Microwatershed




Fig.2.6 Location of Wells – Padasavli-1 Microwatershed

SURVEY METHODOLOGY

The purpose of land resource inventory is to delineate similar areas (soil series and phases), which respond or expected to respond similarly to a given level of management. This was achieved in Padasavli-1 microwatershed by the detailed study of all the soil characteristics (depth, texture, colour, structure, consistence, coarse fragments, porosity, soil reaction, soil horizons etc.) and site (slope of the land, erosion, drainage, occurrence of rock fragments etc.) followed by grouping of similar areas based on soil-site characteristics into homogeneous (management units) units and showing their area extent and geographic distribution on the microwatershed cadastral map. The detailed survey at 1:7920 scale was carried out in 535 ha area. The methodology followed for carrying out land resource inventory was as per the guidelines given in Soil Survey Manual (IARI, 1971; Soil Survey Staff, 2006; Natarajan *et al.*, 2015) which is briefly described below.

3.1 Base Maps

The detailed survey of the land resources occurring in the microwatershed was carried out by using digitized cadastral map as a base. The cadastral map shows field boundaries with their survey numbers, location of tanks, streams and other permanent features of the area (Fig. 3.1). Apart from the cadastral map, remote sensing data products from Cartosat-1 and LISS-IV merged at the scale of 1:7920 were used in conjunction with the cadastral map to identify the rock types, the landscapes, landforms and other surface features. The imagery helped in the identification and delineation of boundaries between hills, uplands and lowlands, water bodies, forest and vegetated areas, roads, habitations and other cultural features of the area (Fig.3.2). The cadastral map was overlaid on the satellite imagery (Fig.3.3) that helps to identify the parcel boundaries and other permanent features. Apart from cadastral maps and images, toposheets of the area (1:50,000 scale) were used for initial traversing, identification of geology and landforms, drainage features, present land use and also for selection of transects in the microwatershed.




Fig 3.1 Scanned and Digitized Cadastral map of Padasavli-1 Microwatershed




Fig.3.2 Satellite Image of Padasavli-1 Microwatershed




Fig.3.3 Cadastral map overlaid on IRS PAN+LISS IV merged imagery of Padasavli-1 Microwatershed

3.2 Field Investigation

Preliminary traverse of the microwatershed was carried out with the help of cadastral map, imagery and toposheets. While traversing, landforms and physiographic units identified were checked and preliminary soil legend was prepared by studying soils at a few selected places. The field boundaries and survey numbers given on the cadastral sheet were located on the ground by following permanent features like roads, cart tracks, nallas, streams, tanks etc., and wherever changes were noticed, they were incorporated on the microwatershed cadastral map. Then, intensive traversing of each physiographic unit like hills, ridges and uplands was carried out. Based on the variability observed on the surface, transects were selected across the slope covering all the landform units in the microwatershed (Natarajan and Dipak Sarkar, 2010).

In the selected transect, soil profiles were located at closely spaced intervals to take care of any change in the land features like break in slope, erosion, gravel, stones etc. In the selected sites, profiles (vertical cut showing the soil layers from surface to the rock) were opened up to 200 cm or to the depth limited by rock or hard substratum and studied in detail for all their morphological and physical characteristics. The soil and site characteristics were recorded for all profile sites on a standard proforma as per the guidelines given in USDA Soil Survey Manual (Soil Survey Staff, 2012). Apart from the transect study, profiles were also studied at random, almost like in a grid pattern, outside the transect areas.

Based on the soil-site characteristics, the soils were grouped into different soil series (soil series is the most homogeneous unit having similar horizons and properties and behaves similarly for a given level of management). Soil depth, texture, colour, kind of horizon and horizon sequence, amount and nature of gravel present, nature of substratum etc, were used as the major differentiating characteristics for identifying soil series occurring in the area. The differentiating characteristics used for identifying soil series are given in Table 3.1. Based on the above characteristics, 7 soil series were identified in the Padasavli-1 microwatershed.

Table 3.1 Differentiating Characteristics used for Identifying Soil Series
(Characteristics are of Series Control Section)

SOILS OF BASALT LANDSCAPE							
Sl. no	Soil Series	Depth (cm)	Colour	Text-ure	Gravel (%)	Horizon sequence	Calcar-eousness
1	Margutti (MGT)	<25	10YR3/3,4/3,5/4 7.5YR4/3	c	15-35	Ap-R/cr	-
2	Novinihala (NHA)	25-50	10YR3/2,3/1,4/2 7.5YR3/4	c	<15	Ap-Bw-cr/R	-
3	Bhimanahalli (BHI)	25-50	10YR3/2,3/3,3/1 7.5YR3/2,4/2	c	15-35	Ap-Bw-cr/R	-
4	Dinsi (DSI)	50-75	10YR3/2,3/3,4/3 3/2	c	<15	Ap-BA-Bss	-
5	Gutti (GTT)	50-75	10YR3/2, 3/1 7.5YR3/3, 4/3	c	15-35	Ap-Bw-Bss-cr	-
6	Kamalapur (KMP)	75-100	10YR3/2, 3/1	c	<15	Ap-Bw-Bss-cr	-
7	Mahagaon (MAN)	>150	10YR3/2,3/1	c	<15	Ap-BA-Bss	-

3.3 Laboratory Characterization

Soil samples were collected from representative master profiles for laboratory characterization by following the methods outlined in the Laboratory Manual (Sarma *et al*, 1987). Surface soil samples collected from farmer's fields (85 samples) for fertility status (major and micronutrients) at 250 m grid interval were analyzed in the laboratory. (Katyal and Rattan, 2003) By linking the soil fertility data to the survey numbers through GIS, soil fertility maps for all 11 elements including pH and Ec were generated for the micro watershed.

3.4 Finalization of Soil Maps

The area under each soil series was further separated and mapped as soil phases and their boundaries delineated on the cadastral map based on the variations observed in the texture of the surface soil, slope, erosion, presence of gravel, stoniness etc. A soil phase is a subdivision of soil series based mostly on surface features that affect its use and management.

The soil mapping units are shown on the map (Fig.3.4) in the form of symbols. During the survey about 14 profile pits, few minipits and a few auger bores representing different landforms occurring in the microwatershed were studied. All the profile locations are indicated on the village cadastral map in the form of a triangle. In addition to the profile study, spot observations in the form of minipits, road cuts, terrace cuts etc., were studied to validate the soil boundaries on the soil map. The soil map shows the geographic distribution of 16 mapping units representing 7 soil series occurring in the microwatershed. The soil map unit (soil legend) description is presented in Table 3.2.

The soil phase map (management units) shows the distribution of 16 phases identified and mapped in the microwatershed. Each mapping unit (soil phase) delineated on the map has similar soil and site characteristics. In other words, all the farms or survey numbers included in one phase will have similar management needs and they have to be treated accordingly.

The 16 soil phases identified and mapped in the microwatershed were regrouped into 5 Land Management Units (LMU's) for the purpose of preparing a proposed crop plan for sustained development of the microwatershed. The database (soil phases) generated under LRI was utilized for identifying Land Management Units (LMUs) based on the management needs. One or more than one soil site characteristic having influence on the management have been chosen for identification and delineation of LMUs. For Padasavli-1 microwatershed, five soil and site characteristics, namely soil depth, soil texture, slope, erosion and gravel content have been considered for defining LMUs. The land management units are expected to behave similarly for a given level of management.




Fig 3.4 Soil phase or Management Units map of Padasavli-1 Microwatershed

Table 3.2 Soil Legend

Map unit no.	Soil series	Soil Phase	Mapping Unit Description	Area in ha (%)
Soils of Basalt Landscape				
	MGT		Margutti soils are very shallow (<25cm), well drained, have very dark grayish brown to dark brown clayey soils occurring on very gently sloping to gently sloping uplands under cultivation	60.75 (11.37)
1		MGTiC2g1	Sandy clay surface, slope 3-5 %, moderate erosion, gravelly (15-35%)	4.36 (0.82)
2		MGTmB2g1	Clay surface, slope 1-3 %, moderate erosion, gravelly (15-35%)	23.57 (4.41)
3		MGTmB3g1	Clay surface, slope 1-3 %, severe erosion, gravelly (15-35%)	19.26 (3.60)
4		MGTmC3g2	Clay surface, slope 3-5%, severe erosion, very gravelly (35-60%)	13.56 (2.54)
	NHA		Novinihala soils are shallow (25-50 cm), well drained, have very dark grayish brown to dark brown clayey soils occurring on very gently sloping uplands under cultivation	206.5 (38.63)
5		NHAmB2g1	Clay surface, slope 1-3 %, moderate erosion, gravelly (15-35%)	159.24 (29.79)
6		NHAmB2g2	Clay surface, slope 1-3 %, moderate erosion, very gravelly (35-60%)	47.26 (8.84)
	BHI		Bhimanahalli soils are shallow (25-50 cm), well drained, have very dark gray to brown clay soils occurring on very gently sloping to gently sloping uplands under cultivation	72.40 (13.54)
7		BHImB2g2	Clay surface, slope 1-3 %, moderate erosion, very gravelly (35-60 %)	61.59 (11.52)
8		BHImC3g1	Clay surface, slope 3-5%, severe erosion, gravelly (15-35%)	10.81 (2.02)
	DSI		Dinsi soils are moderately shallow (50-75 cm), moderately well drained, have very dark gray to brown clayey soils occurring on very gently to gently sloping uplands	86.83 (16.25)
9		DSImB2	Clay surface, slope 1-3%, moderate erosion	40.56 (7.59)
10		DSImB2g1	Clay surface, slope 1-3%, moderate erosion, gravelly (15-35%)	31.20 (5.84)
11		DSImC2g1	Clay surface, slope 3-5%, moderate erosion, gravelly (15-35%)	10.52 (1.97)

12		DSImC3g2	Clay surface, slope 3-5%, severe erosion, very gravelly (35-60%)	4.55 (0.85)
	GTT	Gutti soils are moderately shallow (50-75 cm), moderately well drained, have very dark gray to brown clayey soils occurring on very gently sloping uplands under cultivation		24.80 (4.64)
13		GTTmB2g1	Clay surface, slope 1-3%, moderate erosion, gravelly (15-35%)	24.80 (4.64)
	KMP	Kamalapur soils are moderately deep (75-100 cm), moderately well drained, have very dark gray to very dark grayish brown cracking clay soils occurring on very gently sloping uplands under cultivation		7.96 (1.49)
14		KMPmB2g1	Clay surface, slope 1-3%, moderate erosion, gravelly (15-35%)	7.96 (1.49)
	MAN	Mahagaon soils are very deep (>150 cm), moderately well drained, have very dark gray to very dark grayish brown cracking clay soils occurring on nearly level to very gently sloping uplands		49.62 (9.28)
15		MANmA1	Clay surface, 0-1% slope, slight erosion	33.27 (6.22)
16		MANmB2	Clay surface, 1-3% slope, moderate erosion	16.35 (3.06)
17	Habitation			25.66 (4.80)

THE SOILS

Detailed information pertaining to the nature, extent and distribution of different kinds of soils occurring in Padasavli-1 microwatershed is provided in this chapter. The microwatershed area has been identified as Basalt landscape. In all, 7 soil series were identified in this landscape. Soil formation is the result of the combined effect of environmental and terrain factors that are reflected in soil morphology. In the Basalt landscape, it is by parent material and climate. A brief description of each of the 7 soil series identified followed by 16 soil phases (management units) mapped under each series are furnished below. The soils in any one map unit differ from place to place in their depth, texture, slope, gravelliness, erosion or any other site characteristics that affect management. The soil phase map can be used for identifying the suitability of areas for growing specific crops or for other alternative uses and also for deciding the type of conservation structures needed. The detailed information on soil and site-characteristics like soil depth, surface soil texture, slope, erosion, gravelliness, AWC, LCC etc, with respect to each of the soil phase identified is given village/survey number wise for the microwatershed in Appendix-I.

4.1 Soils of Basalt Landscape

In this landscape, 7 soil series are identified and mapped. Of these, Novanihala (NHA) soil series occupies maximum area of about 207 ha (39%). The brief description of each series along with the soil phases identified and mapped is given below.

4.1.1 Margutti (MGT) Series: Margutti soils are very shallow (<25cm), well drained, have very dark grayish brown to dark brown clay soils. They have developed from basalt and occur on very gently sloping to moderately sloping uplands.

The total depth of the soil ranges from 10 to 23 cm. The thickness of A horizon ranges from 7 to 18 cm. Its colour is in 10 YR and 7.5 YR hue with value 3 to 4 and chroma 2 to 4. The texture is clay with 15 to 35 per cent gravel. The available water capacity is very low (<50 mm/m).

Four phases were identified:

MGTiC2g1	Sandy clay surface, slope 3-5 %, moderate erosion, gravelly (15-35%)
MGTmB2g1	Clay surface, slope 1-3 %, moderate erosion, gravelly (15-35%)
MGTmB3g1	Clay surface, slope 1-3 %, severe erosion, gravelly (15-35%)
MGTmC3g2	Clay surface, slope 3-5%, severe erosion, very gravelly (35-60%)



Landscape and Soil Profile Characteristics of Margutti (MGT) Series

4.1.2 Novanihala (NHA) Series: Novanihala soils are shallow (25-50 cm), well drained, have very dark grayish brown to dark brown clay soils. They have developed from basalt and occur on very gently sloping uplands.

The thickness of the solum ranges from 27 to 48 cm. The thickness of A horizon ranges from 12 to 20 cm. Its colour is in 7.5 YR and 10 YR hue with value 3 to 4 and chroma 2 to 4. The texture varies from sandy clay to clay with 10 to 20 per cent gravel. The thickness of B horizon ranges from 22 to 37 cm. Its colour is in 10 YR and 7.5 YR hue with value 3 to 4 and chroma 2 to 4. Its texture is clay with gravel content of 10-15 per cent. The available water capacity is low (51-100 mm/m).

Two phases were identified:

NHAmB2g1	Clay surface, slope 1-3 %, moderate erosion, gravelly (15-35%)
NHAmB2g2	Clay surface, slope 1-3 %, moderate erosion, very gravelly (35-60%)



Landscape and Soil Profile Characteristics of Novanihala (NHA) Series

4.1.3 Bhimanahalli (BHI) Series: Bhimanahalli soils are shallow (25-50 cm), well drained, have very dark gray to brown clay soils. They have developed from basalt and occur on very gently sloping to gently sloping uplands.

The thickness of the solum ranges from 29 to 48 cm. The thickness of A horizon ranges from 15 to 20 cm. Its colour is in 7.5 YR and 10 YR hue with value 3 to 4 and chroma 2 to 4. The texture varies from sandy clay to clay with 15 to 25 per cent gravel. The thickness of B horizon ranges from 23 to 33 cm. Its colour is in 10 YR and 7.5 YR hue with value 3 to 4 and chroma 1 to 3. Its texture is clay with gravel content of 15 to 35 per cent. The available water capacity is very low (<50 mm/m).

Two phases were identified:

BHImB2g2	Clay surface, slope 1-3 %, moderate erosion, very gravelly (35-60 %)
BHImC3g1	Clay surface, slope 3-5%, severe erosion, gravelly (15-35%)



Landscape and Soil Profile Characteristics of Bhimanahalli (BHI) Series

4.1.4 Dinsi (DSI) Series: Dinsi soils are moderately shallow (50-75 cm), moderately well drained, have very dark gray to brown clay soils. They have developed from basalt and occur on very gently to gently sloping uplands.

The thickness of the solum ranges from 55 to 71 cm. The thickness of A horizon ranges from 9 to 24 cm. Its colour is in 10 YR hue with value 3 and chroma 1 to 3. The texture is clay with 5 to 10 per cent gravel. The thickness of B horizon ranges from 27 to 62 cm. Its colour is in 10 YR hue with value 3 to 4 and chroma 2 to 4. Its texture is clay with gravel content of less than 15 per cent. The available water capacity is medium (101-150 mm/m).

Four phases were identified:

DSImB2	Clay surface, slope 1-3%, moderate erosion
DSImB2g1	Clay surface, slope 1-3%, moderate erosion, gravelly (15-35%)
DSImC2g1	Clay surface, slope 3-5%, moderate erosion, gravelly (15-35%)
DSImC3g2	Clay surface, slope 3-5%, severe erosion, very gravelly (35-60%)



Landscape and Soil Profile Characteristics of Dinsi (DSI) Series

4.1.5 Gutti (GTT) Series: Gutti soils are moderately shallow (50-75 cm), moderately well drained, have very dark gray to brown clayey soils. They have developed from basalt and occur on very gently sloping uplands.

The thickness of the solum ranges from 24 to 74 cm. The thickness of A horizon ranges from 7 to 23 cm. Its colour is in 10 YR and 7.5 YR hue with value 3 to 4 and chroma 2 to 3. The texture is clay with 10 to 15 per cent gravel. The thickness of B horizon ranges from 28 to 65 cm. Its colour is in 10 YR and 7.5 YR hue with value 3 to 4 and chroma 2 to 3. Its texture is clay with gravel content of 15 to 35 per cent. The available water capacity is low (51-100 mm/m).

Only one phase was identified:

GTTmB2g1	Clay surface, slope 1-3%, moderate erosion, gravelly (15-35%)
----------	---



Landscape and Soil Profile Characteristics of Gutti (GTT) Series

4.1.6 Kamalapur (KMP) Series: Kamalapur soils are moderately deep (75-100 cm), moderately well drained, have very dark gray to very dark grayish brown cracking clay soils. They have developed from basalt and occur on very gently sloping uplands.

The thickness of the solum ranges from 75 to 95 cm. The thickness of A horizon ranges from 10 to 30 cm. Its colour is in 10 YR and 7.5 YR hue with value 3 to 4 and chroma 1 to 4. The texture is clay with less than 10 per cent gravel. The thickness of B horizon ranges from 45 to 84 cm. Its colour is in 10 YR hue with value 3 to 4 and chroma 1 to 4. Its texture is clay with gravel content of less than 15 per cent. The available water capacity is medium (101-150 mm/m).

Only one phase was identified:

KMPmB2g1	Clay surface, slope 1-3%, moderate erosion, gravelly (15-35%)
----------	---



Landscape and Soil Profile Characteristics of Kamalapur (KMP) Series

4.1.7 Mahagaon (MAN) Series: Mahagaon soils are very deep (>150 cm), moderately well drained, have very dark gray to very dark grayish brown cracking clay soils. They have developed from basalt and occur on very gently sloping uplands.

The thickness of the solum ranges from 150 to 195 cm. The thickness of A horizon ranges from 18 to 22 cm. Its colour is in 10 YR and 7.5 YR hue with value 3 to 4 and chroma 1 to 3. The texture is clay with less than 10 per cent gravel. The thickness of B horizon ranges from 130 to 160 cm. Its colour is in 10 YR hue with value 3 to 4 and chroma 1 to 2. Its texture is clay with gravel content of less than 15 per cent. The available water capacity is very high (>200 mm/m).

Two phases were identified:

MANmA1	Clay surface, 0-1% slope, slight erosion
MANmB2	Clay surface, 1-3% slope, moderate erosion



Landscape and Soil Profile Characteristics of Mahagaon (MAN) Series

INTERPRETATION FOR LAND RESOURCE MANAGEMENT

The most important soil and site characteristics that affect the land use and conservation needs of an area are land capability, soil depth, soil texture, coarse fragments, available water capacity, soil slope, soil erosion, soil reaction etc. These are interpreted from the data base generated through land resource inventory and several thematic maps are generated. These would help in identifying the areas suitable for growing crops and, soil and water conservation measures and structures needed thus helping to maintain good soil health for sustained crop production. The various thematic maps generated are described below.

5.1 Land Capability Classification

Land capability classification is an interpretative grouping of soil map units (soil phases) mainly based on inherent soil characteristics, external land features and environmental factors that limit the use of land for agriculture, pasture, forestry, or other uses on a sustained basis (IARI, 1971). The land and soil characteristics used to group the land resources in an area into various land capability classes, subclasses and units are

Soil characteristics: Soil depth, soil texture, coarse fragments, soil reaction, available water capacity, calcareousness, salinity/alkali *etc.*

Land characteristics: Slope, erosion, drainage, rock outcrops.

Climate: Total rainfall and its distribution, and length of crop growing period.

The Land capability classification system is divided into land capability classes, subclasses and units based on the level of information available. Eight land capability classes are recognized. They are

Class I: They are very good lands that have no limitations or very few limitations that restrict their use.

Class II: They are good lands that have minor limitations and require moderate conservation practices.

Class III: They are moderately good lands that have moderate limitations that reduce the choice of crops or that require special conservation practices.

Class IV: They are fairly good lands that have very severe limitations that reduce the choice of crops or that require very careful management.

Class V: Soils in these lands are not likely to erode, but have other limitations like wetness that are impractical to remove and as such not suitable for agriculture, but suitable for pasture or forestry with minor limitations

Class VI: The lands have severe limitations that make them generally unsuitable for cultivation, but suitable for pasture or forestry with moderate limitations.

Class VII: The lands have very severe limitations that make them unsuitable for cultivation, but suitable for pasture or forestry with major limitations.

Class VIII: Soil and other miscellaneous areas (rock lands) that have very severe limitations that nearly preclude their use for any crop production, but suitable for wildlife, recreation and wind mills.

The land capability subclasses are recognised based on the dominant limitations observed within a given land capability class. The subclasses are designated by adding a lower case letter like ‘e’, ‘w’, ‘s’, or ‘c’ to the class numeral. The subclass “e” indicates that the main hazard is risk of erosion, “w” indicates drainage or wetness as a limitation for plant growth, “s” indicates shallow soil depth, coarse or heavy textures, calcareousness, salinity/alkalinity or gravelliness and “c” indicates limitation due to climate.

The land capability subclasses have been further subdivided into land capability units based on the kinds of limitations present in each subclass. Ten land capability units are used in grouping the soil map units. They are stony or rocky (0), erosion hazard (slope, erosion) (1), coarse texture (sand, loamy sand, sandy loam) (2), fine texture (cracking clay, silty clay) (3) slowly permeable subsoil (4), coarse underlying material (5), salinity/alkali (6), stagnation, overflow, high ground water table (7), soil depth (8) and fertility problems (9). The capability units thus identified have similar soil and land characteristics that respond similarly to a given level of management. The soils of the microwatershed have been classified upto land capability subclass level only.

The 16 soil map units identified in the Padasavli-1 microwatershed are grouped under 3 land capability classes and 4 land capability subclasses. About 95 per cent area in the microwatershed is suitable for agriculture (Fig. 5.1) .




Fig. 5.1 Land Capability map of Padasavli-1 Microwatershed

Good cultivable lands (Class II) cover about 31 per cent area and are distributed in the northeastern and southeastern part of the microwatershed with minor problems of soil and erosion. Moderately good cultivable lands (Class III) cover maximum area of about 61 per cent and are distributed in the northwestern and southwestern part of the microwatershed with moderate problems of erosion and soil. The fairly good cultivable lands (class IV) cover a small area of about 4 per cent. They have severe limitations of erosion and soil and are distributed in the northern part of the microwatershed.

5.2 Soil Depth

Soil depth refers to the depth of the soil occurring above the parent material or hard rock. The depth of the soil determines the effective rooting depth for plants and in accordance with soil texture, mineralogy and gravel content, the capacity of the soil column to hold water and nutrient availability. Soil depth is one of the most important soil characteristic that is used in differentiating soils into different soil series. The soil depth classes used in identifying soils in the field are very shallow (<25 cm), shallow (25-50 cm), moderately shallow (50-75 cm), moderately deep (75-100 cm), deep (100-150 cm) and very deep (>150 cm). They were used to classify the soils into different depth classes and a soil depth map was prepared (Fig. 5.2).

Very deep soils (>150 cm) occur in a small area of about 50 ha (9%) and are distributed in the northeastern and southeastern part of the micro watershed. Moderately deep soils (75-100 cm) occur in a very minor area of about 8 ha (1%) area and are distributed in the northeastern part of the microwatershed. Moderately shallow (50-75 cm) soils occupy about 112 ha (21%) and are distributed in the southeastern and northeastern part of the microwatershed.

Area of about 61 ha (11%) is under very shallow (<25 cm) and are distributed in the northern, southern and northeastern part of the microwatershed. Maximum area of about 279 ha (52 %) is under shallow soils (25-50 cm) and are distributed in all parts except in the eastern part of the microwatershed.

The most productive lands 50 ha (9%) with respect to soil rooting depth where all climatically adapted annual and perennial crops can be grown are very deep soils (>150 cm depth) occurring in the southeastern and northeastern part of the microwatershed.

The most problem lands with an area of about 340 ha (64%) having very shallow (<25 cm) and shallow (25-50 cm) occur in all parts of the microwatershed. They are not suitable for growing agricultural crops but well suited for pasture, forestry or other recreational purposes. Occasionally, short duration crops may be grown if rainfall is normal.




Fig. 5.2 Soil Depth map of Padasavli-1 Microwatershed

5.3 Surface Soil Texture

Texture is an expression to indicate the coarseness or fineness of the soil as determined by the relative proportion of primary particles of sand, silt and clay. It has a direct bearing on the structure, porosity, adhesion and consistence. The surface layer of a soil to a depth of about 25 cm is the layer that is most used by crops and plants. The surface soil textural class provides a guide to understanding soil-water retention and availability, nutrient holding capacity, infiltration, workability, drainage, physical and chemical behaviour, microbial activity and crop suitability.

Maximum area of 509 ha (95%) has soils that are clayey and are distributed in all over the microwatershed (Fig. 5.3).

The most productive lands (95%) with respect to surface soil texture are the clayey soils that have high potential for soil-water retention and availability, and nutrient retention and availability, but have problems of drainage, infiltration, workability and other physical problems.




Fig. 5.3 Surface Soil Texture map of Padasavli-1 Microwatershed

5.4 Soil Gravelliness

Gravel is the term used for describing coarse fragments between 2 mm and 7.5 cm diameter and stones for those between 7.5 cm and 25 cm. The presence of gravel and stones in soil reduces the volume of soil responsible for moisture and nutrient storage, drainage, infiltration and runoff and hinders plant growth by impeding root growth and seedling emergence, intercultural operations and farm mechanization.

Maximum area has soils that are gravelly (15-35%) covering about 292 ha (55%) and are distributed in the northwestern, central, southern and northeastern part of the microwatershed (Fig.5.4).

About 90 ha (17%) area in the micro watershed has soils that are nongravelly (<15%) and are distributed in the southeastern and northeastern part of the microwatershed followed by soils that are very gravelly (35-60%) soils covering about 127 ha (24%). They are distributed in the southwestern and northern part of the microwatershed.

The most productive lands with respect to gravelliness are found to be 17 per cent. They are nongravelly with less than 15 per cent gravel and have potential for growing both annual and perennial crops.

The problem soils are those that are very gravelly (35-60%) and gravelly (15-35%) and are found to be 78 per cent where only short duration crops can be grown.




Fig. 5.4 Soil Gravelliness map of Padasavli-1 Microwatershed

5.5 Available Water Capacity

The soil available water capacity (AWC) is estimated based on the ability of the soil column to retain water between the tensions of 0.33 and 15 bar in a depth of 100 cm or the entire solum if the soil is shallower. The AWC of the soils (soil series) as estimated by considering the soil texture, mineralogy, soil depth and gravel content (Sehgal *et al.*, 1990) and accordingly the soil map units were grouped into five AWC classes *viz.* very low (<50 mm/m), low (50-100 mm/m), medium (100-150 mm/m), high (150-200 mm/m) and very high (>200 mm/m) and using these values, an AWC map was prepared (Fig. 5.5).

Area of about 133 ha (25%) has soils that are very low (<50 mm/m) in available water capacity and are distributed in the northern, northwestern and southwestern part of the microwatershed. Major area of about 231 ha (43%) has soils that are low (51-100 mm/m) in available water capacity and are distributed in the southern, northern, central and a small area in southeastern part of the microwatershed. About 95 ha (18%) area in the microwatershed has soils that are medium (101-150 mm/m) in available water capacity and are distributed in the northeastern and southern part of the microwatershed followed by soils that are very high (>200 mm/m) in AWC covering a small area of about 50 ha (9%) in the microwatershed. They are distributed in the northeastern and southeastern part of the microwatershed.

An area of about 145 ha (27%) has soils that have very high potential (>200 mm/m) and medium (101-150 mm/m) with regard to available water capacity. In these areas, if the

rainfall is normal and well distributed, all climatically adapted long duration annual and perennial crops can be grown.

About 133 ha (25%) area in the microwatershed has soils that are problematic with regard to available water capacity. Here, only the short duration crops can be grown and the probability of crop failure is very high. These areas are best put to other alternative uses.




Fig. 5.5 Soil Available Water Capacity map of Padasavli-1 Microwatershed

5.6 Soil Slope

Soil slope refers to the inclination of the surface of the land. It is defined by gradient, shape and length, and is an integral feature of any soil as a natural body. Slope is considered important in soil genesis, land use and land development. The length and gradient of slope influences the rate of runoff, infiltration, erosion and deposition. The soil map units were grouped into four slope classes and a slope map was prepared showing the area extent and geographic distribution of different slope classes in the microwatershed (Fig. 5.6).

Major area of the microwatershed falls under very gently sloping (1-3% slope) slope class. It covers an area of about 432 ha (81%) and is distributed in all parts of the microwatershed. An area of about 44 ha (8 %) in the microwatershed falls under gently sloping (3-5%) slope class and is distributed in the northern, southeastern, northeastern and southern part of the microwatershed.

Nearly level lands (0-1%) slope class cover a small area of about 33 ha (6 %) and are distributed in the eastern part of the microwatershed.

An area of about 465 ha (87%) in the microwatershed has soils that have high potential in respect of soil slopes. In these areas, all climatically adapted annual and perennial crops can be grown without much soil and water conservation and other land development measures.




Fig. 5.6 Soil Slope map of Padasavli-1 Microwatershed

5.7 Soil Erosion

Soil erosion refers to the wearing away of the earth's surface by the forces of water, wind and ice involving detachment and transport of soil by raindrop impact. It is used for accelerated soil erosion resulting from disturbance of the natural landscape by burning, excessive grazing and indiscriminate felling of forest trees and tillage, all usually by man. The erosion classes showing an estimate of the current erosion status as judged from field observations in the form of rills, gullies or a carpet of gravel on the surface are recorded. Four erosion classes, viz, slight erosion (e1), moderate erosion (e2), severe erosion (e3) and very severe erosion (e4) are recognized. The soil map units were grouped into different erosion classes and soil erosion map was generated. The area extent and their spatial distribution in the microwatershed is given in Figure 5.7.

Soils that are slightly eroded (e1 class) cover a very minor area of about 33 ha (6%) and are distributed in the eastern part of the microwatershed. Soils that are moderately eroded (e2 class) cover a maximum area of about 427 ha (80%) and are distributed in all parts of the

microwatershed. Severely eroded (e3 class) soils cover an area of about 48 ha (9%) and are distributed in the northern and a small area in southwestern part of the microwatershed.

Top priority is to be given to 48 ha area where they are severely eroded for taking up soil and water conservation and other land development measures followed by moderately eroded lands that cover about 427 ha.




Fig. 5.7 Soil Erosion map of Padasavli-1 Microwatershed

FERTILITY STATUS

Soil fertility plays an important role in increasing crop yield. The adoption of high yielding varieties that require high amounts of nutrients has resulted in deficiency symptoms in crops and plants due to imbalanced fertilization and poor inherent fertility status. Hence, it is necessary to know the fertility (macro and micro nutrients) status of the soils of the watersheds for assessing the kind and amount of fertilizers required for each of the crop intended to be grown. For this purpose, the surface soil samples collected from the grid points (one soil sample at every 250 m interval) all over the microwatershed through land resource inventory in the year 2014 were analysed for pH, EC, organic carbon, available phosphorus and potassium and for micronutrients like zinc, copper, iron and manganese, and secondary nutrient sulphur.

Soil fertility data generated has been assessed and individual maps for all the nutrients for the microwatershed have been prepared. The village/survey number wise fertility data for the microwatershed is given in Appendix-II.

6.1 Soil Reaction (pH)

The soil fertility analysis of the Padasavli-1 microwatershed for soil reaction (pH) showed that about 107 ha (20%) area is slightly alkaline (pH 7.3-7.8) in reaction and is distributed in the northern, northeastern and southern part of the microwatershed. Maximum area of about 295 ha (55%) is moderately alkaline (pH 7.8-8.4) and is distributed in all parts of the microwatershed. About 60 ha (11%) area is strongly alkaline (pH 8.4-9.0) and is distributed in eastern part of the microwatershed; about 47 ha (9%) area is neutral (pH 6.5-7.3) and is distributed in the northeastern part of the microwatershed.

6.2 Electrical Conductivity (EC)

The Electrical Conductivity of the soils of the entire microwatershed area is $<2 \text{ dSm}^{-1}$ (Fig 6.2) and as such the soils in the microwatershed are nonsaline.

6.3 Organic Carbon

The soil organic carbon content of the soils in the microwatershed is medium (0.5-0.75%) in 390 ha (73%) area that are distributed in all parts of the microwatershed (Fig.6.3). High ($>0.75\%$) organic carbon content accounts for 46 ha (9%) area and is distributed in the southeastern and southwestern part of the microwatershed. Low ($<0.5\%$) organic carbon content accounts for very minor area of 74 ha (14%) in the microwatershed and is distributed in the northern and northeastern part of the microwatershed.




Fig.6.1 Soil Reaction (pH) map of Padasavli-1 Microwatershed




Fig.6.2 Electrical Conductivity (EC) map of Padasavli-1 Microwatershed

6.4 Available Phosphorus

The soil fertility analysis revealed that available phosphorus is low (<23 kg/ha) in major area of about 419 ha (78%) and is distributed in all parts of the microwatershed (Fig.6.4). There is an urgent need to increase the dose of phosphorous for all the crops by 25 per cent over the recommended dose to realize better crop performance. About 78 ha (15%) area is medium (23-57 kg/ha) and is distributed in the northern, southern, eastern and western part of the microwatershed. A minor area of about 12 ha (2%) is high (>57 kg/ha) and is distributed in the southwestern part of the microwatershed.

6.5 Available Potassium

Available potassium content is medium (145-337 kg/ha) in 122 ha (23%) area and is distributed in the central, northern, northeastern and southern part of the microwatershed (Fig.6.5); high available potassium (>337 kg/ ha) content accounts for major area of 387 ha (72%) and is distributed in the northern and southeastern part of the microwatershed.

6.6 Available Sulphur

Available sulphur content is low (<10 ppm) in major area of about 320 ha (60%) and is distributed in the central, southern, southaeastern and eastern part of the microwatershed. Area of about 162 ha (30%) is medium (10-20 ppm) in available sulphur and is distributed in the northern, northeastern and southwestern part of the microwatershed (Fig.6.6). Available sulphur is high (>20 ppm) in small area of about 27 ha (5%) and is distributed in the northeastern part of the microwatershed.

6.7 Available Boron

Available boron content is low (<0.5 ppm) in major area of about 245 ha (46%) and is distributed in the northern, central, northeastern and southwestern part of the microwatershed. About 185 ha (35%) has soils that are medium (0.5-1.0 ppm) in available boron (Fig 6.7) and is distributed in the eastern and small area in western part of the microwatershed. About 79 ha (15%) has soils that are high (>1.0 ppm) in available boron and is distributed in the central and eastern part of the microwatershed.




Fig.6.3 Soil Organic Carbon map of Padasavli-1 Microwatershed




Fig.6.4 Soil available Phosphorus map of Padasavli-1 Microwatershed




Fig.6.5 Soil available Potassium map of Padasavli-1 Microwatershed




Fig.6.6 Soil available Sulphur map of Padasavli-1 Microwatershed




Fig.6.7 Soil available Boron map of Padasavli-1 Microwatershed

6.8 Available Iron

Available iron content is deficient (<4.5 ppm) in a small area of 44 ha (8%) and is distributed in the southeastern part of the microwatershed. It is sufficient in major area of about 465 ha (87%) (Fig 6.8) and are distributed in all parts of the microwatershed.

6.9 Available Manganese

Available manganese content is sufficient (>1.0 ppm) in the entire microwatershed area (Fig 6.9).

6.10 Available Copper

Available copper content is sufficient (>0.2 ppm) in the entire microwatershed area (Fig 6.10).

6.11 Available Zinc

Available zinc content is deficient (<0.6 ppm) in major area of about 368 ha (69%) and is distributed in all parts of the microwatershed (Fig 6.11). It is sufficient (>0.6 ppm) in 141 ha (26%) and is distributed in the northern, southern and eastern part of the microwatershed.




Fig.6.8 Soil available Iron map of Padasavli-1 Microwatershed




Fig.6.9 Soil available Manganese map of Padasavli-1 Microwatershed




Fig.6.10 Soil available Copper map of Padasavli-1 Microwatershed




Fig.6.11 Soil available Zinc map of Padasavli-1 Microwatershed

LAND SUITABILITY FOR MAJOR CROPS

The soil and land resource units (soil phases) of Padasavli-1 microwatershed were assessed for their suitability for growing food, fibre, fodder and other horticulture crops by following the procedure as outlined in FAO, 1976 and 1983. Crop requirements were developed for each of the crop from the available research data and also by referring to Naidu *et. al.* (2006) and Natarajan *et. al.* (2015). The crop requirements were matched with the soil and land characteristics (Table 7.1) to arrive at the crop suitability. In FAO land suitability classification, two orders are recognized. Order S- Suitable and order N- Not suitable. The orders have classes, subclasses and units. Order-S has three classes, Class S1- Highly Suitable, Class S2- Moderately Suitable and Class S3- Marginally Suitable. Order N has two classes, N1- Currently not Suitable and N2- Permanently not Suitable. There are no subclasses within the class S1 as they will have very minor or no limitations for crop growth. Classes S2 and S3 are divided into subclasses based on the kinds of limitations encountered. The limitations that affect crop production are 'c' for erratic rainfall and its distribution and length of growing period (LGP), 'e' for erosion hazard, 'r' for rooting condition, 't' for lighter or heavy texture, 'g' for gravelliness or stoniness, 'n' for nutrient availability, 'l' for topography, 'm' for moisture availability and 'w' for drainage. These limitations are indicated as lower case letters to the class symbol. For example, moderately suitable land with the limitations of soil depth and erosion is designated as S2re. For the microwatershed, the soil mapping units were evaluated and classified up to subclass level.

Using the above criteria, the soil map units of the microwatershed were evaluated and land suitability maps for 18 major annual and perennial crops were generated. The detailed information on the kind of suitability of each of the soil phase for the crops assessed are given village/ survey number wise for the microwatershed in Appendix-III.

7.1 Land Suitability for Sorghum (*Sorghum bicolor*)

Sorghum is one of the major crops grown in Karnataka in an area of 11.02 lakh ha in northern Karnataka in Bijapur, Kalaburgi, Raichur, Bidar, Belgaum, Dharwad and Bellary districts. The crop requirements for growing sorghum (Table 7.2) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and land suitability map for growing sorghum was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed is given in Figure. 7.1.

An area of about 33 ha (6%) in the microwatershed has soils that are highly suitable (class S1) for growing sorghum crop. They are distributed in the eastern part of the microwatershed. About 131 ha (25%) area is moderately suitable (class S2) for growing sorghum and is distributed in the southern and northeastern part the microwatershed. They have minor limitations of erosion and rooting depth. Major area of about 283 ha (53%) is not suitable for growing sorghum and occur in all parts of the microwatershed except eastern

part. They have moderate limitations of rooting depth, erosion and gravelliness. About 61 ha (11%) area is not suitable for sorghum and is distributed in the northern and northeastern part of the microwatwatershed.




Fig. 7.1 Land Suitability map of Sorghum

Table 7.1 Soil-Site Characteristics of Padasavli-1 Microwatershed

Soil Map Units	Climate (P) (mm)	Growing period (Days)	Drainage class	Soil depth (cm)	Soil texture		Gravelliness		AWC (mm/m)	Slope (%)	Erosion	pH	EC	ESP	CEC [Cmol (p+) kg-1]	BS (%)
					Surface	Sub-surface	Surface (%)	Subsurface (%)								
MGTiC2g1	786	150	WD	<25	sc	c	15-35	15-35	<50	3-5	Moderate	6.8	0.3	0.2	46	100
MGTmB2g1	786	150	WD	<25	c	c	15-35	15-35	<50	1-3	Moderate	6.8	0.3	0.2	46	100
MGTmB3g1	786	150	WD	<25	c	c	15-35	15-35	<50	1-3	Severe	6.8	0.3	0.2	46	100
MGTmC3g2	786	150	WD	<25	c	c	35-60	15-35	<50	3-5	Severe	6.8	0.3	0.2	46	100
NHAmB2g1	786	150	WD	25-50	c	c	15-35	<15	51-100	1-3	Moderate	7.2	0.1	0.3	40	100
NHAmB2g2	786	150	WD	25-50	c	c	35-60	<15	51-100	1-3	Moderate	7.2	0.1	0.3	40	100
BHImB2g2	786	150	WD	25-50	c	c	35-60	15-35	<50	1-3	Moderate	7.0	0.1	0.2	28	100
BHImC3g1	786	150	WD	25-50	c	c	15-35	15-35	<50	3-5	Severe	7.0	0.1	0.2	28	100
DSImB2	786	150	MWD	50-75	c	c	-	<15	101-150	3-5	Moderate	7.0	0.1	0.3	62	100
DSImB2g1	786	150	MWD	50-75	c	c	15-35	<15	101-150	1-3	Moderate	7.0	0.1	0.3	62	100
DSImC2g1	786	150	MWD	50-75	c	c	15-35	<15	101-150	3-5	Moderate	7.0	0.1	0.3	62	100
DSImC3g2	786	150	MWD	50-75	c	c	35-60	<15	101-150	3-5	Severe	7.0	0.1	0.3	62	100
GTTmB2g1	786	150	MWD	50-75	c	c	15-35	15-35	51-100	1-3	Moderate	6.5	0.1	0.6	38	91
KMPmB2g1	786	150	MWD	75-100	c	c	15-35	<15	101-150	1-3	Moderate	6.7	0.2	0.2	43	100
MANmA1	786	150	MWD	>150	c	c	-	<15	>200	<1	Slight	8.3	0.2	0.1	58	100
MANmB2	786	150	MWD	>150	c	c	-	<15	>200	1-3	Moderate	8.3	0.2	0.1	58	100

*Symbols and abbreviations are according to Field Guide for LRI under Sujala-III Project, Karnataka

Table 7.2 Crop suitability criteria for Sorghum

Crop requirement		Rating			
Soil –site characteristics	unit	Highly suitable (S1)	Moderately Suitable (S2)	Marginally suitable (S3)	Not suitable (N)
Slope	%	2-3	3-8	8-15	>15
LGP	Days	120-150	120-90	<90	
Soil drainage	class	Well to mod. drained	imperfect	Poorly/excessively	V. poorly
Soil reaction	pH	6.0-8.0	5.5-5.9 8.1-8.5	<5.5 8.6-9.0	>9.0
Surface soil texture	Class	C, cl, sicl, sc	l, sil, sic	Sl, ls	S, fragmental skeletal
Soil depth	Cm	100-75	50-75	30-50	<30
Gravel content	% vol.	5-15	15-30	30-60	>60
Salinity (EC)	dSm ⁻¹	2-4	4-8	8-10	>10
Sodicity (ESP)	%	5-8	8-10	10-15	>15

7.2 Land Suitability for Maize (*Zea mays*)

Maize is the most important food crop grown in an area of 13.73 lakh ha in all the districts of Karnataka. The crop requirements for growing maize (Table 7.3) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing maize was generated. The area and geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.2.

In Padasavli-1 microwatershed, there are no lands that are highly (class S1) or moderately (class S2) suitable lands for growing maize.

The marginally suitable (class S3) lands cover maximum area of about 449 ha (84%) and occur in major part of the microwatershed. They have moderate limitations of texture, erosion, gravelliness and rooting depth. An area of about 61 ha (11%) is not suitable for growing maize and occur in the northern and southwestern part of the microwatershed.

Table 7.3 Crop suitability criteria for Maize

Crop requirement		Rating			
Soil –site characteristics	unit	Highly suitable (S1)	Moderately Suitable (S2)	Marginally suitable (S3)	Not suitable (N)
Slope	%	<3	3.5	5-8	
LGP	Days	>100	100-80	60-80	
Soil drainage	class	Well drained	Mod. to imperfectly	Poorly/ excessively	V.poorly
Soil reaction	pH	5.5-7.5	7.6-8.5	8.6-9.0	
Surface soil texture	Class	l, cl, scl, sil	Sl, sicl, sic	C(s-s), ls	S,fragmental
Soil depth	Cm	>75	50-75	25-50	<25
Gravel content	% vol.	<15	15-35	35-50	>50
Salinity (EC)	dSm ⁻¹	<1.0	1.0-2.0	2.0-4.0	
Sodicity (ESP)	%	<10	10-15	>15	




Fig. 7.2 Land Suitability map of Maize

7.3 Land Suitability for Red gram (*Cajanus cajan*)

Red gram is one of the major pulse crop grown in an area of 8.23 lakh ha mainly in northern Karnataka in Bijapur, Kalaburgi, Raichur, Bidar, Belgaum, Dharwad and Bellary districts. The crop requirements for growing red gram (Table 7.4) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and land suitability map for growing red gram was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed is given in Figure 7.3.

An area of about 164 ha (31%) is moderately suitable (class S2) for red gram and is distributed dominantly in the eastern part of the microwatershed. They have minor limitations of rooting depth, erosion and texture. Marginally suitable lands (class S3) for growing red gram occupy maximum area of about 283 ha (53%) and mainly occur in the northwestern and southwestern part of the microwatershed. They have moderate limitations of rooting depth, erosion and gravelliness. An area of about 61 ha (11%) is not suitable for growing red gram and occur in northern, southern and northeastern part of the microwatershed.

Table 7.4 Crop suitability criteria for Red gram

Crop requirement		Rating			
Soil characteristics	–site unit	Highly suitable (S1)	Moderately Suitable (S2)	Marginally suitable (S3)	Not suitable (N)
Slope	%	<3	3-5	5-10	>10
LGP	Days	>210	180-210	150-180	<150
Soil drainage	class	Well drained	Mod. to well drained	Imperfectly drained	Poorly drained
Soil reaction	pH	6.5-7.5	5.0-6.5 7.6-8.0	8.0-9.0	>9.0
Surface soil texture	Class	l, scl, sil, cl, sl	sicl, c(m)	sic, ls	S, fragmental
Soil depth	Cm	>100	85-100	40-85	<40
Gravel content	% vol.	<20	20-35	35-60	>60
Salinity (EC)	dSm ⁻¹	<1.0	1.0-2.0	>2.0	
Sodicity (ESP)	%	<10	10-15	>15	




Fig. 7.3 Land Suitability map of Red gram

7.4 Land Suitability for Sunflower (*Helianthus annuus*)

Sunflower is the most important oilseed crop grown in an area of 4.1 lakh ha in the State in all the districts. The crop requirements for growing sunflower (Table 7.5) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing sunflower was generated. The area and geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.4.

Highly suitable (class S1) lands are found to occur in a small area of 33 ha (6%) and are distributed in the eastern part of the microwatershed. A very small area of about 24 ha (5%) is moderately suitable (class S2) for red gram and is distributed in the northeastern part of the microwatershed. They have minor limitations of erosion. The marginally suitable (class S3) lands cover about 112 ha (21%) area in the microwatershed and occur in the eastern part of the microwatershed. They have moderate limitations of rooting depth and gravelliness. Major area of about 340 ha (64%) is not suitable for growing sunflower and occur in all parts of the microwatershed except eastern part.

Table 7.5 Crop suitability criteria for Sunflower

Crop requirement		Rating			
Soil –site characteristics	unit	Highly suitable (S1)	Moderately Suitable (S2)	Marginally suitable (S3)	Not suitable (N)
Slope	%	<3	3-5	5-10	>10
LGP	Days	>90	80-90	70-80	<70
Soil drainage	class	Well drained	mod. Well drained	imperfectly drained	Poorly drained
Soil reaction	pH	6.5-8.0	8.1-8.5 5.5-6.4	8.6-9.0; 4.5-5.4	>9.0 <4.5
Surface soil texture	Class	l, cl, sil, sc	Scl, sic, c,	c (>60%), sl	ls, s
Soil depth	Cm	>100	75-100	50-75	<50
Gravel content	% vol.	<15	15-35	35-60	>60
Salinity (EC)	dSm ⁻¹	<1.0	1.0-2.0	>2.0	
Sodicity (ESP)	%	<10	10-15	>15	




Fig. 7.4 Land Suitability map of Sunflower

7.5 Land Suitability for Cotton (*Gossypium hirsutum*)

Cotton is the most important fibre crop grown in the State in about 6.6 lakh ha area in Raichur, Dharwad, Belgaum, Kalaburgi, Bijapur, Bidar, Bellary, Chitradurga and Chamarajnagar districts. The crop requirements for growing cotton (Table 7.6) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing cotton was generated. The area and geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.5.

Highly suitable (class S1) lands are found to occur in a small area of 33 ha (6%) and are distributed in the eastern part of the microwatershed. Moderately suitable (class S2) lands are found to occur in an area of about 131 ha (25%). The soils have minor limitations of erosion and rooting depth. They are dominantly distributed in the eastern part of the microwatershed. The marginally suitable (class S3) lands cover maximum area of about 283 ha (53%) and occur in all parts of the microwatershed except in the eastern part. They have moderate limitations of rooting depth, erosion and gravelliness. About 61 ha (11%) area is not suitable for growing cotton and are distributed in the northern and southern part of the microwatershed.

Table 7.6 Crop suitability criteria for Cotton

Crop requirement		Rating			
Soil-site characteristics	unit	Highly suitable (S1)	Moderately Suitable (S2)	Marginally suitable (S3)	Not suitable (N)
Slope	%	1-2	2-3	3-5	>5
LGP	Days	180-240	120-180	<120	
Soil drainage	class	Well to moderately well	imperfectly drained	Poor somewhat excessive	Stagnant/excessive
Soil reaction	pH	6.5-7.5	7.6-8.0	8.1-9.0	>9.0 >6.5
Surface soil texture	Class	Sic, c	Sicl, cl	Si, sil, sc, scl, l	Sl, s,ls
Soil depth	Cm	100-150	60-100	30-60	<30
Gravel content	% vol.	<5	5-10	10-15	15-35
CaCO ₃ in root zone	%	<3	3-5	5-10	10-20
Salinity (EC)	dSm ⁻¹	2-4	4.0-8.0	8.0-12	>12
Sodicity (ESP)	%	5-10	10-20	20-30	>30




Fig. 7.5 Land Suitability map of Cotton

7.6 Land Suitability for Sugarcane (*Saccharum officinarum*)

Sugarcane is the most important commercial crop grown in 6.7 lakh ha area in Kalaburgi, Bijapur, Bagalkot, Bidar, Mysore, Chamarajanagar and Mandya districts. The crop requirements for growing sugarcane (Table 7.7) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing sugarcane was generated. The area and geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.6.

In Padasavli-1 microwatershed, there are no highly (class S1) and moderately suitable (class S2), lands available for growing sugarcane.

The marginally suitable (class S3) lands cover about 165 ha (31%) area and mainly occur in the eastern part of the microwatershed. They have moderate limitations of texture. Major area of about 344 ha (64%) is not suitable for growing sugarcane and occur in all parts of the microwatershed.

Table 7.7 Crop suitability criteria for Sugarcane

Crop requirement		Rating			
Soil–site characteristics	unit	Highly suitable (S1)	Moderately Suitable (S2)	Marginally suitable (S3)	Not suitable (N)
Slope	%	<3	3-5	5-8	>8
Soil drainage	class	Well drained	Mod./imperfectly drained	Poorly drained	V.poor/excessively drained
Soil reaction	pH	7.0-8.0	6.0-6.9 8.1-9.0	4.0-5.9,9.1-9.5	<4.0/ >9.5
Surface soil texture	Class	l, cl, sil, sicl	C(m/k), sl	C+(ss)	
Soil depth	cm	>100	100-75	75-50	<50
stoniness	%	<15	15-35	35-50	>50
Salinity (EC)	dSm-1	<2.0	2.0-4.0	4.0-9.0	>9
Sodicity (ESP)	%	<10	10-15	15-25	>25




Fig. 7.6 Land Suitability map of Sugarcane

7.7 Land Suitability for Soybean (*Glycine max*)

Soybean is the most important pulse and oilseed crop grown in about 1.68 lakh ha area in the northern districts of the state. The crop requirements for growing soybean were matched with the soil-site characteristics and a land suitability map for growing soybean was generated. The area and geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.7.

Highly suitable (class S1) lands are found to occur in a small area of 33 ha (6%) and are distributed in the eastern part of the microwatershed. Moderately suitable (class S2) lands are found to occur an area of about 131 ha (25%). The soils have minor limitations of erosion and rooting depth. They are dominantly distributed in the eastern part of the microwatershed. The marginally suitable (class S3) lands cover maximum area of about 283 ha (53%) area and mainly occur in the northern, western and southern part of the microwatershed. They have moderate limitations of rooting depth, erosion and gravelliness. About 61 ha (11%) is not suitable for growing soybean and occur in the northern and southern part of the microwatershed.




Fig. 7.7 Land Suitability for Soybean

7.8 Land Suitability for Guava (*Psidium guajava*)

Guava is the most important fruit crop grown in the State in Raichur, Dharwad, Belgaum, Kalaburgi, Bijapur, Bidar, Bellary, Chitradurga and Chamarajnagar districts. The crop requirements for growing guava (Table 7.8) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing guava was generated. The area and geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.8.

In Padasavli-1 microwatershed, there are no highly (class S1) and moderately (class S2) suitable lands available for growing guava. The marginally suitable (class S3) lands cover about 170 ha (32%) area in the microwatershed and mainly occur in the eastern part of the microwatershed. They have moderate limitations of texture, gravelliness and rooting depth.

Major area of about 340 ha (64%) is not suitable for growing guava and occur in all parts of the microwatershed.

Table 7.8 Crop suitability criteria for Guava

Crop requirement			Rating			
Soil –site characteristics		unit	Highly suitable (S1)	Moderately Suitable (S2)	Marginally suitable (S3)	Not suitable (N)
climate	Temperature in growing season	⁰ C	28-32	33-36 24-27	37-42 20-23	
Soil moisture	Growing period	Days	>150	120-150	90-120	<90
Soil aeration	Soil drainage	class	Well drained	Mod. to imperfectly	poor	Very poor
Nutrient availability	Texture	Class	Scl, l, cl, sil	Sl,siel,sic.,sc,c	C (<60%)	C (>60%)
	pH	1:2.5	6.0-7.5	7.6-8.0: 5.0-5.9	8.1-8.5: 4.5-4.9	>8.5: <4.5
	CaCO ₃ in root zone	%	Non calcareous	<10	10-15	>15
Rooting conditions	Soil depth	cm	>100	75-100	50-75	<50
	Gravel content	% vol.	<15	15-35	>35	
Soil toxicity	Salinity	dS/m ⁻¹	<2.0	2.0-4.0	4.0-6.0	
	Sodicity	%	Non sodic	10-15	15-25	>25
Erosion	Slope	%	<3	3-5	5-10	>10




Fig 7.8 Land Suitability for Guava

7.9 Land Suitability for Mango (*Mangifera indica*)

Mango is the most important fruit crop grown in the State in all the districts of the state. The crop requirements for growing Mango (Table 7.9) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing Mango was generated. The area and geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.9.

In Padasavli-1 microwatershed, there are no highly (class S1) and moderately (class S2) suitable lands available for growing mango. The marginally suitable (class S3) lands cover a small area of about 58 ha (11%) and mainly occur in the eastern part of the microwatershed. They have moderate limitations of texture. Major area of about 451 ha (84%) is not suitable for growing mango and occur in all parts of the microwatershed.

Table 7.9 Crop suitability criteria for Mango

Crop requirement			Rating			
soil-site characteristics		unit	Highly suitable (S1)	Moderately Suitable (S2)	Marginally suitable (S3)	Not suitable (N)
Climate	Temp in growing season	⁰ C	28-32	24-27 33-35	36-40	20-24
	Min. temp. before flowering	⁰ C	10-15	15-22	>22	
Soil moisture	Growing period	Days	>180	150-180	120-150	<120
Soil aeration	Soil drainage	class	Well drained	Mod. To imperfectly drained	Poor drained	Very poorly drained
	Water table	M	>3	2.50-3.0	2.5-1.5	<1.5
Nutrient availability	Texture	Class	Sc, l, sil, cl	Sl, sc, sic, l, c	C (<60%)	C (>60%),
	pH	1:2.5	5.5-7.5	7.6-8.55.0-5.4	8.6-9.0 4.0-4.9	>9.0 <4.0
	OC	%	High	medium	low	
	CaCO ₃ in root zone	%	Non calcareous	<5	5-10	>10
Rooting conditions	Soil depth	cm	>200	125-200	75-125	<75
	Gravel content	% vol	Non gravelly	<15	15-35	>35
Soil toxicity	Salinity	dS/m ⁻¹	Non saline	<2.0	2.0-3.0	>3.0
	Sodicity	%	Non sodic	<10	10-15	>15
Erosion	Slope	%	<3	3-5	5-10	




Fig. 7.9 Land Suitability for Mango

7.10 Land Suitability for Sapota (*Manilkara zapota*)

Sapota is the most important fruit crop grown in almost all the districts of the state. The crop requirements for growing sapota (Table 7.10) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing sapota was generated. The area and geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.10.

In Padasavli-1 microwatershed, there are no highly (class S1) and moderately (class S2) suitable lands available for growing sapota. The marginally suitable (class S3) lands cover about 170 ha (32%) area in the microwatershed and mainly occur in the eastern part of the microwatershed. They have moderate limitations of rooting depth, gravelliness and texture. Maximum area of about 340 ha (64%) is not suitable for growing sapota and occur in major parts of the microwatershed.

Table 7.10 Crop suitability criteria for Sapota

Crop requirement			Rating			
Soil –site characteristics		unit	Highly suitable (S1)	Moderately Suitable (S2)	Marginally Suitable (S3)	Not suitable (N)
Climate	Temperature in growing season	⁰ C	28-32	33-36 24-27	37-42 20-23	>42 <18
Soil moisture	Growing period	Days	>150	120-150	90-120	<120
Soil aeration	Soil drainage	class	Well drained	Moderately well drained	Imperfectly drained	Poorly drained
Nutrient availability	Texture	Classes	Scl, l, cl, sil	Sl, sicl, sc	C (<60%)	ls, s, C (>60%)
	pH	1:2.5	6.0-7.5	7.6-8.0 5.0-5.9	8.1-9.0 4.5-4.9	>9.0 <4.5
	CaCO ₃ in root zone	%	Non calcareous	<10	10-15	>15
Rooting conditions	Soil depth	cm	>150	75-150	50-75	<50
	Gravel content	% vol.	Non gravelly	<15	15-35	<35
Soil toxicity	Salinity	dS/m ⁻¹	Non saline	Up to 1.0	1.0-2.0	2.0-4.0
	Sodicity	%	Non sodic	10-15	15-25	>25
Erosion	Slope	%	<3	3-5	5-10	>10




Fig. 7.10 Land Suitability for Sapota

7.11 Land Suitability for Jackfruit (*Artocarpus heterophyllus*)

Jackfruit is the most important fruit crop grown in southern and western districts of the state. The crop requirements for growing jackfruit were matched with the soil-site characteristics and a land suitability map for growing jackfruit was generated. The area and geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.11.

No highly (class S1) and moderately (class S2) suitable lands are available for growing Jackfruit in the microwatershed. The marginally suitable (class S3) lands cover about 170 ha (32%) area in the microwatershed and mainly occur in the eastern part of the microwatershed. They have moderate limitations of rooting depth, gravelliness and texture. Major area of about 340 ha (64%) is not suitable for growing jackfruit and occur in all parts of the microwatershed.




Fig 7.11 Land Suitability for Jackfruit

7.12 Land Suitability for Jamun (*Syzygium cumini*)

Jamun is the most important fruit crop grown in almost all the districts of the state. The crop requirements for growing jamun were matched with the soil-site characteristics and a land suitability map for growing jamun was generated. The area and geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.12.

Moderately suitable (class S2) lands are found to occur in a small area of about 49 ha (9%). The soils have minor limitations of erosion, texture and rooting depth. They are

dominantly distributed in the eastern part of the microwatershed. The marginally suitable (class S3) lands cover about 120 ha (22%) area and occur in the eastern part of the microwatershed. They have moderate limitations of rooting depth and gravelliness. Major area of about 340 ha (64%) is not suitable for growing jamun and occur in all parts of the microwatershed.




Fig 7.12 Land Suitability for Jamun

7.13 Land Suitability for Musambi (*Citrus limetta*)

Musambi is the most important fruit crop grown in almost all the districts of the state. The crop requirements for growing musambi were matched with the soil-site characteristics and a land suitability map for growing musambi was generated. The area and geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.13. Highly suitable (class S1) lands are found to occur in a small area of 33 ha (6%) and are distributed in the eastern part of the microwatershed. Moderately suitable (class S2) lands are found to occur in a small area of about 24 ha (5%). The soils have minor limitations of rooting depth and erosion. They are distributed in the northeastern part of the microwatershed.

The marginally suitable (class S3) lands cover about 112 ha (21%) area and mainly occur in the eastern part of the microwatershed. They have moderate limitations of rooting depth and gravelliness. Major area of about 340 ha (64%) is not suitable for growing musambi and occur in all parts of the microwatershed.




Fig 7.13 Land Suitability for Musambi

7.14 Land Suitability for Lime (*Citrus sp*)

Lime is the most important fruit crop grown in almost all the districts of the state. The crop requirements for growing lime (Table 7.11) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing lime was generated. The area and geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.14.

Highly suitable (class S1) lands are found to occur in a small area of about 33 ha (6%) and are distributed in the eastern part of the microwatershed. Moderately suitable (class S2) lands are found to occur in a very small area of about 24 ha (5%). The soils have minor limitations of rooting depth and erosion. They are dominantly distributed in the northeastern part of the microwatershed. The marginally suitable (class S3) lands cover about 112 ha (21%) area and occur in the eastern part of the microwatershed. They have moderate limitations of rooting depth and gravelliness. Major area of about 340 ha (64%) is not suitable for growing lime and occur in all parts of the microwatershed.

Table 7.11 Crop suitability criteria for Lime

Crop requirement			Rating			
Soil –site characteristics		unit	Highly suitable (S1)	Moderately Suitable (S2)	Marginally suitable (S3)	Not suitable (N)
Climate	Temp in growing season	⁰ C	28-30	31-35 24-27	36-40 20-23	>40 <20
Soil moisture	Growing period	Days	240-265	180-240	150-180	<150
Soil aeration	Soil drainage	class	Well drained	Mod. to imperfectly drained	poorly	Very poorly
Nutrient availability	Texture	Class	Scl, l, sicl, cl, s	Sc, sc, c	C (>70%)	S, ls
	pH	1:2.5	6.0-7.5	5.5-6.4/ 7.6-8.0	4.0-5.4 8.1-8.5	<4.0 >8.5
	CaCO ₃ in root zone	%	Non calcareous	Upto 5	5-10	>10
Rooting condition	Soil depth	cm	>150	100-150	50-100	<50
	Gravel content	% vol.	Non gravelly	15-35	35-55	>55
Soil toxicity	Salinity	dS/m	Non saline	Upto 1.0	1.0-2.5	>2.5
	Sodicity	%	Non sodic	5-10	10-15	>15
Erosion	Slope	%	<3	3-5	5-10	




Fig 7.14 Land Suitability for Lime

7.15 Land Suitability for Cashew (*Anacardium occidentale*)

Cashew is the most important plantation crop grown mostly in coastal and western parts and also in Bidar and Kolar districts of the state. The crop requirements for growing Cashew were matched with the soil-site characteristics and a land suitability map for growing Cashew was generated. The area and geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.15.

The entire area is not suitable for growing cashew in the microwatershed.




Fig 7.15 Land Suitability for Cashew

7.16 Land Suitability for Custard Apple (*Annona reticulata*)

Custard apple is the most important fruit crop grown in almost all the districts of the state. The crop requirements for growing Custard apple were matched with the soil-site characteristics and a land suitability map for growing Custard apple was generated. The area and geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.16.

Highly suitable (class S1) lands are found to occur in a small area of 33 ha (6%) and are distributed in the eastern part of the microwatershed. Moderately suitable (class S2) lands are found to occur in an area of about 131 ha (25%). The soils have minor limitations of erosion and rooting depth. They are dominantly distributed in the eastern part of the microwatershed. The marginally suitable (class S3) lands cover major area of about 284 ha (53%) and occur in all parts of the microwatershed. They have moderate limitations of

rooting depth, erosion and gravelliness. About 61 ha (11%) is not suitable for growing Custard apple and occur in the northern and southern part of the microwatershed.




Fig 7.16 Land Suitability for Custard Apple

7.17 Land Suitability for Amla (*Phyllanthus emblica*)

Amla is the most important fruit crop grown in almost all the districts of the state. The crop requirements for growing amla were matched with the soil-site characteristics and a land suitability map for growing amla was generated. The area and geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.17.

Highly suitable (class S1) lands are found to occur in an area of 33 ha (6%) and are distributed in the eastern part of the microwatershed. Moderately suitable (class S2) lands are found to occur in an area of about 131 ha (25%). The soils have minor limitations of erosion and rooting depth. They are dominantly distributed in the southeastern and eastern part of the microwatershed. The marginally suitable (class S3) lands cover maximum area of about 284 ha (53%) and occur in major part of the microwatershed. They have moderate limitations of rooting depth, erosion and gravelliness. About 61 ha (11%) is not suitable for growing amla and occur in the northern and southern part of the microwatershed.




Fig 7.17 Land Suitability for Amla

7.18 Land Suitability for Tamarind (*Tamarindus indica*)

Tamarind is the most important spice crop grown in all the districts of the state. The crop requirements for growing tamarind were matched with the soil-site characteristics and a land suitability map for growing tamarind was generated. The area and geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.18.

Moderately suitable (class S2) lands are found to occur in a small area of about 49 ha (9%). The soils have minor limitations of rooting depth and erosion. They are distributed in the eastern part of the microwatershed. A very minor area of about 8 ha (1%) is marginally suitable (class S3) for growing tamarind. They have moderate limitations of rooting depth and are distributed in the eastern part of the microwatershed. Major area of about 451 ha (84%) is not suitable for growing tamarind and occur in all parts of the microwatershed.

LMUs	Soil map units	Soil and site characteristics
1	MGTmC3g2	Very shallow, black soils with slopes of 3-5%, very gravelly (35-60%) and severe erosion
2	MGTmC2g1, MGTmB2g1 MGTmB3g1	Very shallow, black soils with slopes of 1-5%, gravelly (15-35%) and moderate to severe erosion
3	NHAmB2g1, NHAmB2g2, BHImB2g2, BHImC3g1, DSImC3g2	Shallow to moderately shallow, black soils with slopes of 1-5 %, gravelly to very gravelly (15-60%) and moderate to severe erosion
4	DSImB2, DSImB2g1, DSImC2g1, GTTmB2g1, KMPmB2g1	Moderately shallow to moderately deep, black soils with slopes of 1-5 %, gravelly (15-35%) and moderate erosion
5	MANmA1, MANmB2	Very deep, black soils with slopes of <1-3% and slight to moderate erosion




Fig. 7.19 Land Management Units map – Padasavli-1 Microwatershed

7.20 Proposed Crop Plan for Padasavli-1 Microwatershed

After assessing the land suitability for the 18 crops, the proposed crop plan has been prepared for the 5 identified LMUs by considering only the highly (class S1) and moderately (class S2) suitable lands for each of the eighteen crops. The resultant proposed crop plan is presented below in Table 7.12

Table 7.12 Proposed Crop Plan for Padasavli-1 Microwatershed

LMUs No	Mapping Units	Survey Number	Field Crops	Forestry/Grasses	Horticulture Crops (Rainfed Condition)	Horticulture Crops With suitable intervention	Suitable Intervention
LMU 1	4 (<25 cm)	Sarasamba: 38,39,44,45,46	-	Neem, Glyricydia , Silviculture, Agave, Simaroba	-	-	Crescent bunds
LMU 2	1,2,3 (<25 cm)	Padasavli: 211,220,275,276/1 Sarasamba: 31,41,42	Horse gram	Neem, Glyricydia , Silviculture, Agave, Simaruba	-	-	Crescent bunds
LMU 3	5,6,7,8,12 (25-75 cm)	Padasavli: 33,34,41,186,187,188, 189,194,201, 203,204, 205,207, 208,209/1, 209/2,210,212,213,214, 215,216,217,218, 219 Sarasamba: 12,13,14,15,16,17, 19,35, 36,37,38,39, 40,43,44, 45, 46,53, 54,55	Bajra, Linseed, Green gram, Black gram, Chick pea, Coriander	Subabul, Neem, Teak	Custard apple, Charoli, Ber, Amla Vegetables: Ladies finger, Brinjal, Cowpea, Flowers: Marigold, Chrysanthemum	Custard apple, Charoli, Ber, Amla Vegetables: Onion, Tomato, Brinjal, Chillies, Bhendi Flowers: Marigold, Chrysanthemum	Drip irrigation, suitable soil and water conservation measures like cultivation on raised beds with mulches and drip

LMU 4	9,10,11,13,14 (50-100 cm)	Padasavli: 10,12,29,31,32,35, 36,54, 195,196, 198/ 1,198/2, 198/3, 199,200,202,206, 276/2,277/1,277/2	Sorghum, Cotton, Red Gram, Black gram, Green gram, Soybean, Sesame, Sunflower, Rabi: Sorghum, Chickpea, Coriander, Safflower	Subabul, Neem, Teak	Custard apple, Charoli, Ber, Amla Vegetables: Ladies finger, Brinjal, Cowpea, Flowers: Marigold, Chrysanthemum	Custard apple, Charoli, Ber, Amla, Papaya, Banana, Lime, Citrus Vegetables: Onion, Tomato, Brinjal, Chillies, Bhendi Flowers: Marigold, Chrysanthemum	-do- Graded bunds, Strengthening of field bunds
LMU 5	15, 16 (>150 cm)	Padasavli: 2/1,2/2,2/3,3,4,5,6,7/2, 7/3,8,37,38/1,38/2,274	Sorghum, Cotton, Red Gram Black gram, Green gram, Soybean, Sesame, Sunflower Rabi: Sorghum, Chickpea , Coriander, Linseed, Safflower	-	Vegetables: Ladies finger, Brinjal, Cowpea, Coriander Field crops: Sorghum, Cotton, Red Gram, Sunflower, Safflower, Perennial component: Guava, Sapota, Lime, Mosambi, Tamarind, Flowers: Marigold, Chrysanthemum	Banana, Papaya, Lime, Musambi, Guava, Tamrind Vegetables: Onion, Tomato, Brinjal, Chillies, Bhendi Flowers: Marigold, Chrysanthemum	-do- Graded bunds, Strengthening of field bunds

SOIL HEALTH MANAGEMENT

8.1 Soil Health

Soil is fundamental to crop production. Without soil, no food could be produced nor would livestock be fed on a large scale. Because it is finite and fragile, soil is a precious resource that requires special care from its users.

Soil health or the capacity of the soil to function is critical to human survival. Soil health has been defined as: “the capacity of the soil to function as a living system without adverse effect on the ecosystem”. Healthy soils maintain a diverse community of soil organisms that help to form beneficial symbiotic associations with plant roots, recycle essential plant nutrients, improve soil structure with positive repercussions for soil, water and nutrient holding capacity and ultimately improve crop production and also contribute to mitigating climate change by maintaining or increasing its carbon content.

Functional interactions of soil biota with organic and inorganic components, air and water determine a soil’s potential to store and release nutrients and water to plants and to promote and sustain plant growth. Thus, maintaining soil health is vital to crop production and conserve soil resource base for sustaining agriculture.

The most important characteristics of a healthy soil are

- Good soil tilth
- Sufficient soil depth
- Good water storage and good drainage
- Adequate supply, but not excess of nutrients
- Large population of beneficial organisms
- Small proportion of plant pathogens and insect pests
- Low weed pressure
- Free of chemicals and toxins that may harm the crop
- Resistance to degradation
- Resilience when unfavourable conditions occur

Characteristics of Padasavli-1 Microwatershed

- The soil phases with sizeable area identified in the microwatershed belonged to the soil series of NHA (207 ha), DSI (87 ha), BHI (72 ha), MGT (61 ha), MAN (50 ha), GTT (25 ha) and KMP (8 ha). As per land capability classification, nearly 95 per cent area falls under arable land category (Class II and IV). The major limitations identified in the arable lands were soil and erosion.
- On the basis of soil reaction, about 402 ha (75%) area is slightly to moderately alkaline (pH 7.3-8.4) followed by strongly alkaline (pH 8.4-9.0) 60 ha (11%). Thus, about 86 per cent of the soils are alkaline in reaction. About 47 ha (9%) area is neutral (pH 6.5-7.3).

Soil Health Management

The following actions are required to improve the current land husbandry practices that provide a sound basis for the successful adoption of sustainable crop production system.

Alkaline soils

(Slightly alkaline to moderately alkaline soils)

1. Regular addition of organic manure, green manuring, green leaf manuring, crop residue incorporation and mulching needs to be taken up to improve the soil organic matter status.
2. Application of biofertilizers (Azospirillum, Azotobacter, Rhizobium).
3. Application of 25% extra N and P (125 % RDN&P).
4. Application of $ZnSO_4$ – 12.5 kg/ha (once in three years).
5. Application of Boron – 5kg/ha (once in three years).

Neutral soils

1. Regular addition of organic manure, green manuring, green leaf manuring, crop residue incorporation and mulching needs to be taken up to improve the soil organic matter status.
2. Application of biofertilizers, (Azospirillum, Azotobacter, Rhizobium).
3. Application of 100 per cent RDF.
4. Need based micronutrient applications.

Besides the above recommendations, the best transfer of technology options are also to be adopted.

Soil Degradation

Soil erosion is one of the major factors affecting the soil health in the micro watershed. Out of total area of 535 ha in the microwatershed, major area of 475 ha is suffering from either moderate or severe erosion. These areas need immediate soil and water conservation and, other land development and land husbandry practices for restoring soil health.

Disseminate information and communicate benefits.

Any large scale implementation of soil health management requires that supporting information is made available widely, particularly through channels familiar to farmers and extension workers. Given the very high priority attached to soil health especially by the Central Government on issuing Soil-Health Cards to all the farmers, media outlets like regional, state and national newspapers, radio and Dooradarshan programs in local languages but also modern information and communication technologies such as cellular phones and the Internet, which can be much more effective in reaching younger farmers.

Inputs for Net Planning and Interventions needed

Net planning in IWMP is focusing on preparation of

1. Soil and Water Conservation Plans for each plot or farm.
2. Productivity enhancement measures/ interventions for existing crops/livestock/other farm enterprises.
3. Diversification of farming mainly with perennial horticultural crops and livestock.
4. Improving livelihood opportunities and income generating activities.

In this connection, how various outputs of Sujala-III are of use in addressing these objectives of Net Planning are briefly presented below.

- ❖ **Soil Depth:** The depth of a soil decides the amount of moisture and nutrients it can hold, what crops can be taken up or not, depending on the rooting depth and the length of growing period available for raising any crop. Deeper the soil, better for a wide variety of crops. If sufficient depth is not available for growing deep rooted crops, either choose medium or short duration crops or deeper planting pits need to be opened and additional good quality soil brought from outside has to be filled into the planting pits.
- ❖ **Surface soil texture:** Lighter soil texture in the top soil means, better rain water infiltration, less run-off and soil moisture conservation, less capillary rise and less evaporation losses. Lighter surface textured soils are amenable to good soil tilth and are highly suitable for crops like groundnut, root vegetables (carrot, radish, potato etc) but not ideal for crops that need stagnant water like lowland paddy. Heavy textured soils are poor in water infiltration and percolation. They are prone for sheet erosion; such soils can be improved by sand mulching. The technology that is developed by the AICRP-Dryland Agriculture, Vijayapura, Karnataka can be adopted.
- ❖ **Gravelliness:** More gravel content is favorable for run-off harvesting but poor in soil moisture storage and nutrient availability. It is a significant parameter that decides the kind of crop to be raised.
- ❖ **Land Capability Classification:** The land capability map shows the areas suitable and not suitable for agriculture and the major constraints in each of the plot/survey number. Hence, one can decide what kind of enterprise is possible in each of these units. In general, erosion and soil are the major constraints in Padasavli-1 microwatershed.
- ❖ **Organic Carbon:** In about 74 ha (14%) area, the OC content is low (<0.5%), in about 390 ha (73%) area, the OC content is medium (0.5-0.75%) and in about 46 ha (6%) area, it is high (>0.75%). The areas that are low and medium in OC needs to be further improved by applying farmyard manure and rotating crops with cereals and legumes or mixed cropping.
- ❖ **Promoting green manuring:** Growing of green manuring crops costs Rs. 1250/ha (green manuring seeds) and about Rs. 2000/ha towards cultivation that totals to Rs. 3250/- per ha. On the other hand, application of organic manure @ 10 tons/ha costs Rs. 5000/ha. The

practice needs to be continued for 2-3 years or more. Nitrogen fertilizer needs to be supplemented by 25% in addition to the recommended level in 65 ha area where OC is less than 0.5-0.75%. For example, for rainfed maize, recommended level is 50 kg N per ha and an additional 12 kg /ha needs to be applied for all the crops grown in these plots.


- ❖ **Available Phosphorus:** In 419 ha (78%) area, the available phosphorus is low and about 78 ha (15%) area it is medium in available phosphorus, Hence for all the crops, 25% additional P-needs to be applied. About 12 ha (2%) area is high in available phosphorus in the microwatershed.
- ❖ **Available Potassium:** Available potassium is medium in 122 ha (23%) area of the microwatershed. Hence, in all these plots, for all crops, an additional 25 % potassium may be applied. It is high in 387 ha (72%) area of the microwatershed.
- ❖ **Available Sulphur:** Available sulphur is a very critical nutrient for oilseed crops. It is low in major area of 320 ha (60%) in the microwatershed and medium in 162 ha (30%). These areas need to be applied with magnesium sulphate or gypsum or Factamphos (p) fertilizer (13% sulphur) for 2-3 years for the deficiency to be corrected. About 27 ha (5 %) is high in available sulphur.
- ❖ **Available iron:** It is deficient in a small area of 44 ha (8%) in the microwatershed. To manage iron deficiency, iron sulphate @ 25kg /ha needs to be applied for 2-3 years. It is sufficient in the rest of 465 ha (87 %) area in the microwatershed.
- ❖ **Available Zinc:** It is deficient in major area of 368 ha (69%) in the microwatershed. Application of zinc sulphate @25kg/ha is to be applied. It is sufficient in 141 ha (26%) area in the microwatershed.
- ❖ **Soil alkalinity:** The microwatershed has 462 ha area with soils that are slightly to strongly alkaline. These areas need application of gypsum and wherever calcium is in excess, iron pyrites and element sulphur can be recommended. Management practices like treating repeatedly with good quality water to drain out the excess salts and provision of subsurface drainage and growing of salt tolerant crops like Casuarina, Acasia, Neem, Ber etc, are recommended.

Land Suitability for various crops: Areas that are highly, moderately and marginally suitable for growing various crops are indicated. Along with the suitability, various constraints that are limiting the productivity are also indicated. For example, in case of cotton, gravel content, rooting depth and salinity/alkalinity are the major constraints in various plots. With suitable management interventions, the productivity can be enhanced. In order to increase water holding capacity of light textured soils, growing of green manure crops and application of organic manure is recommended.

SOIL AND WATER CONSERVATION TREATMENT PLAN

For preparing soil and water conservation treatment plan for Padasavli-1 microwatershed, the land resource inventory database generated under Sujala-III project has been transformed as information through series of interpretative (thematic) maps using soil phase map as a base. The various thematic maps (1:7920 scale) generated were

- Soil depth
- Surface soil texture
- Available water capacity
- Soil slope
- Soil gravelliness
- Land capability
- Present land use and land cover
- Crop suitability maps
- Rainfall map
- Hydrology
- Water Resources
- Socio-economic data
- Contour plan with existing features- Network of waterways, pottissa boundaries, cut up/ minor terraces etc.
- Cadastral map (1:7920 scale)
- Satellite imagery (1:7920 scale)



Apart from these, Hand Level/ Hydro Marker/ Dumpy Level/ Total Station and Kathedars' List has to be collected.

Steps for Survey and Preparation of Treatment Plan

The boundaries of Land user Groups' and Survey No. boundaries are traced in the field.

- Naming of user groups and farmers
- Identification of arable and non arable lands
- Identification of drainage lines and gullies
- Identification of non treatable areas
- Identification of priority areas in the arable lands
- Treatment plan for arable lands
- Location of water harvesting and recharge structures

9.1 Treatment Plan

The treatment plan recommended for arable lands is briefly described below.


9.1.1 Arable Land Treatment

A. BUNDING

Steps for Survey and Preparation of Treatment Plan		USER GROUP-1
Cadastral map (1:7920 scale) is enlarged to a scale of 1:2500 scale		
Existing network of waterways, pothissa boundaries, grass belts, natural drainage lines/ watercourse, cut ups/ terraces are marked on the cadastral map to the scale		
Drainage lines are demarcated into		
Small gullies	(up to 5 ha catchment)	
Medium gullies	(5-15 ha catchment)	
Ravines	(15-25 ha catchment) and	
Halla/Nala	(more than 25ha catchment)	

Measurement of Land Slope

Land slope is estimated or determined by the study and interpretation of contours or by measurement in the field using simple instruments like Hand level or Hydromarker.



Vertical and Horizontal intervals between bunds as recommended by the Watershed Development Department.

Slope percentage	Vertical interval (m)	Corresponding Horizontal Distance (m)
2 - 3%	0.6	24
3 - 4%	0.9	21
4 - 5%	0.9	21
5 - 6%	1.2	21
6 - 7%	1.2	21

Note: (i) The above intervals are maximum.

(ii) Considering the slope class and erosion status (A1....) the intervals have to be decided.

Bund length recording: Considering the contour plan and the existing grass belts/partitions, the bunds are aligned and lengths are measured.

Section of the Bund

Bund section is decided considering the soil texture class and gravelliness class (bg₀, loamy sand, <15% gravel). The recommended Sections for different soils are given below.

Recommended Bund Section



Top width (m)	Base width (m)	Height (m)	Side slope (Z:1;H:V)	Cross section (sq m)	Soil Texture	Remarks
0.3	0.9	0.3	01:01	0.18	Sandy loam	Vegetative bund
0.3	1.2	0.3	1.5:1	0.225	Sandy clay	
0.3	1.2	0.5	0.9:1	0.375	Red gravelly soil	
0.3	1.2	0.6	0.75:1	0.45		
0.3	1.5	0.6	01:01	0.54	Red sandy loam	
0.3	2.1	0.6	1.5:1	0.72	Very shallow black soils	
0.45	2	0.75	01:01	0.92		
0.45	2.4	0.75	1.3:1	1.07	Shallow black soils	
0.6	3.1	0.7	1.78:1	1.29	Medium black soils	
0.5	3	0.85	1.47:1	1.49		

Formation of Trench cum Bund

Dimensions of the Borrow Pits/ Trenches to be excavated (machinery are decided considering the Bund Section).

Details of Borrow Pit dimensions are given below:

TRENCH CUM BUND





0.45 Sq.m section

IDEAL FOR HORTICULTURE CROPS



1. ಸಮಸಾತಲ ಉಳುವೆ
2. ಸಮಸಾತಲ ಬಿತ್ತನೆ/ನಾಟಿ



'A' FRAME FOR INTERBUND MANAGEMENT

Size of Borrow Pits/ Trench recommended for Trench cum Bund (by machinery)

Bund section	Bund length	Earth quantity	Pit				Berm (pit to pit)	Soil depth class
			L(m)	W(m)	D(m)	QUANTITY(m ³)		
m ²	m	m ³					m	
0.375	6	2.25	5.85	0.85	0.45	2.24	0.15	Shallow
0.45	6	2.7	5.4	1.2	0.43	2.79	0.6	Shallow
0.45	6	2.7	5	0.85	0.65	2.76	1	Moderately Shallow
0.54	5.6	3.02	5.5	0.85	0.7	3.27	0.1	Moderately shallow
0.54	5.5	2.97	5	1.2	0.5	3	0.5	Shallow
0.72	6.2	4.46	6	1.2	0.7	5.04	0.2	Moderately shallow
0.72	5.2	3.74	5.1	0.85	0.9	3.9	0.1	Moderately deep

B. Waterways

- Existing waterways are marked on the cadastral map (1:7920 scale) and their dimensions are recorded.
- Considering the contour plan of the MWS, additional waterways/ modernization of the existing ones can be thought of.
- The design details are given in the Manual.

C. Farm Ponds

Waterways and the catchment area will give an indication on the size of the Farm Pond. Location of the pond can be decided based on the contour plan/ field condition and farmers' need/desire.

D. Diversion channel

Existing EPT/ CPT are marked on the cadastral map. Looking to the need, these can be modernized or fresh diversion channel can be proposed and runoff from this can be stored in Gokatte/ Recharge ponds.

9.1.2 Non-Arable Land Treatment

Depending on the gravelliness and crops preferred by the farmers, the concerned authorities can decide appropriate treatment plan. The recommended treatments may be Contour Trench, Staggered Trench, Crescent Bund, Boulder Bund or Pebble Bunds are formed in the field.

9.1.3 Treatment of Natural Water Course/ Drainage Lines

- a) The cadastral map has to be updated as regards the network of drainage lines (gullies/ nalas/ hallas) and existing structures are marked to the scale and storage capacity of the existing water bodies are documented.
- b) The drainage line will be demarcated into Upper Reach, Middle Reach and Lower Reach.
- c) Considering the Catchment, Nala bed and bank conditions, suitable structures are decided.
- d) Number of storage structures (Check dam/ Nala bund/ Percolation tank) will be decided considering the commitments and available runoff in water budgeting and quality of water in the wells and site suitability.
- e) Detailed Levelling Survey using Dumpy Level / Total Station has to be carried out to arrive at the site-specific designs as shown in the Manual.
- f) The location of ground water recharge structures are decided by examining the lineaments and fracture zones from geological maps.
- g) Rainfall intensity data of the nearest Rain gauge station is considered for Hydrologic Designs.
- h) Silt load to the Storage/Recharge structures is reduced by providing vegetative, boulder and earthen checks in the natural water course. Location and design details are given in the Manual.

9.2 Recommended Soil and Water Conservation Measures

The appropriate conservation structures best suited for each of the land parcel/ survey number (Appendix-I) are selected based on the slope per cent, severity of erosion, amount of rainfall, land use and soil type. The different kinds of conservation structures recommended are

1. Graded / Strengthening of bunds
2. Trench cum Bunds (TCB)
3. Trench cum Bunds / Strengthening
4. Crescent Bunds

A map (Fig. 9.1) showing soil and water conservation plan with different kinds of structures recommended has been prepared which shows the spatial distribution and extent of area. An area of about 120 ha (22%) requires trench cum bunding and about 50 ha (9%) area needs graded bunds. The maximum area of about 340 ha (64%) requires crescent bund.

The conservation plan prepared may be presented to all the stakeholders including farmers and after including their suggestions, the conservation plan for the microwatershed may be finalised in a participatory approach.




Fig. 9.1 Soil and Water Conservation Plan map of Padasavli-1 Microwatershed

9.3 Greening of Microwatershed

As part of the greening programme in the watersheds, it is envisaged to plant a variety of horticultural and other tree plants that are edible, economical and produce lot of biomass which helps to restore the ecological balance in the watersheds. The lands that are suitable for greening programme are non-arable lands (land capability classes V, VI and VII) and also the lands that are not suitable or marginally suitable for growing annual and perennial crops. The method of planting these trees is given below.

It is recommended to open pits during the 1st week of March along the contour and heap the dugout soil on the lower side of the slope in order to harness the flowing water and facilitate weathering of soil in the pit. Exposure of soil in the pit also prevents spread of pests and diseases due to scorching sun rays. The pits should be filled with mixture of soil and organic manure during the second week of April and keep ready with sufficiently tall seedlings produced either in poly bags or in root trainer nurseries so that planting can be done during the 2nd or 3rd week of April depending on the rainfall.

The tree species suitable for the area considering rainfall, temperature and adaptability is listed below; waterlogged areas are recommended to be planted with species like Neral (*Sizyium cumini*) and Bamboo. Dry areas are to be planted with species like Honge, Bevu, Seetaphal etc.

Dry Deciduous Species			Temp (°C)	Rainfall (mm)
1.	Bevu	<i>Azadiracta indica</i>	21-32	400 -1,200
2.	Tapasi	<i>Holoptelia integrifolia</i>	20-30	500 - 1000
3.	Seetaphal	<i>Anona Squamosa</i>	20-40	400 - 1000
4.	Honge	<i>Pongamia pinnata</i>	20 -50	500- 2,500
5.	Kamara	<i>Hardwickia binata</i>	25 -35	400 - 1000
6.	Bage	<i>Albezzia lebbek</i>	20 - 45	500 - 1000
7.	Ficus	<i>Ficus bengalensis</i>	20 - 50	500-2,500
8.	Sisso	<i>Dalbargia Sissoo</i>	20 - 50	500 -2000
9.	Ailanthus	<i>Ailanthus excelsa</i>	20 - 50	500 - 1000
10.	Hale	<i>Wrightia tinctoria</i>	25 - 45	500 - 1000
11.	Uded	<i>Steriospermum chelanoides</i>	25 - 45	500 -2000
12.	Dhupa	<i>Boswellia Serrata</i>	20 - 40	500 - 2000
13.	Nelli	<i>Emblica Officinalis</i>	20 - 50	500 -1500
14.	Honne	<i>Pterocarpus marsupium</i>	20 - 40	500 - 2000
Moist Deciduous Species			Temp (°C)	Rainfall (mm)
15.	Teak	<i>Tectona grandis</i>	20 - 50	500-5000
16.	Nandi	<i>Legarstroemia lanceolata</i>	20 - 40	500 - 4000
17.	Honne	<i>Pterocarpus marsupium</i>	20 - 40	500 - 3000
18.	Mathi	<i>Terminalia alata</i>	20 -50	500 - 2000
19.	Shivane	<i>Gmelina arboria</i>	20 -50	500 -2000
20.	Kindal	<i>T.Paniculata</i>	20 - 40	500 - 1500
21.	Beete	<i>Dalbargia latifolia</i>	20 - 40	500 - 1500
22.	Tare	<i>T. belerica</i>	20 - 40	500 - 2000
23.	Bamboo	<i>Bambusa arundinasia</i>	20 - 40	500 - 2500
24.	Bamboo	<i>Dendrocalamus strictus</i>	20 - 40	500 - 2500
25.	Muthuga	<i>Butea monosperma</i>	20 - 40	400 - 1500
26.	Hippe	<i>Madhuca latifolia</i>	20 - 40	500 - 2000
27.	Sandal	<i>Santalum album</i>	20 - 50	400 - 1000
28.	Nelli	<i>Emblica officinalis</i>	20 - 40	500 - 2000
29.	Nerale	<i>Sizygium cumini</i>	20 - 40	500 - 2000
30.	Dhaman	<i>Grevia tilifolia</i>	20 - 40	500 - 2000
31.	Kaval	<i>Careya arborea</i>	20 - 40	500 - 2000
32.	Harada	<i>Terminalia chebula</i>	20 - 40	500 - 2000

References

1. FAO (1976) Framework for Land Evaluation, Food and Agriculture Organization, Rome.72 pp.
2. FAO (1983) Guidelines for Land Evaluation for Rainfed Agriculture, FAO, Rome, 237 pp.
3. IARI (1971) Soil Survey Manual, All India Soil and Land Use Survey Organization, IARI, New Delhi, 121 pp.
4. Katyal, J.C. and Rattan, R.K. (2003) Secondary and Micronutrients; Research Gap and future needs. Fert. News 48 (4); 9-20.
5. Naidu, L.G.K., Ramamurthy, V., Challa, O., Hegde, R. and Krishnan, P. (2006) Manual Soil Site Suitability Criteria for Major Crops, NBSS Publ. No. 129, NBSS &LUP, Nagpur, 118 pp.
6. Natarajan, A. and Dipak Sarkar (2010) Field Guide for Soil Survey, National Bureau of Soil Survey and Land Use Planning (ICAR), Nagpur, India.
7. Natarajan, A., Rajendra Hegde, Raj, J.N. and Shivananda Murthy, H.G. (2015) Implementation Manual for Sujala-III Project, Watershed Development Department, Bengaluru, Karnataka.
8. Sarma, V.A.K., Krishnan, P. and Budihal, S.L. (1987) Laboratory Manual, Tech. Bull. 23, NBSS &LUP, Nagpur.
9. Sehgal, J.L. (1990) Soil Resource Mapping of Different States of India; Why and How?, National Bureau of Soil Survey and Land Use Planning, Nagpur, 49 pp.
10. Shivaprasad, C.R., R.S. Reddy, J. Sehgal and M. Velayuthum (1998) Soils of Karnataka for Optimising Land Use, NBSS Publ. No. 47b, NBSS & LUP, Nagpur, India.
11. Soil Survey Staff (2006) Keys to Soil Taxonomy, Tenth edition, U.S. Department of Agriculture/ NRCS, Washington DC, U.S.A.
12. Soil Survey Staff (2012) Soil Survey Manual, Handbook No. 18, USDA, Washington DC, USA.

Appendix - I

Padasavli -1 Microwatershed														
Soil Site and Thematic Information														
Village	Survey No.	Total Area (ha)	Soil phase	Land Management Unit	Soil Depth	Surface Soil Texture	Soil Graveliness	AWC	Slope	Soil Erosion	CLU Code	WELLS	Land Capability	Conservation Plan
Padasavali	2/1	4.9	MANmA1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Sugarcane (Sc)	2 Bore Wells, Check Dam	IIs	GB/strengthening of Bunds
Padasavali	2/2	2.78	MANmA1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Sugarcane (Sc)	Not Available	IIs	GB/strengthening of Bunds
Padasavali	2/3	1.76	MANmA1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Sugarcane (Sc)	Not Available	IIs	GB/strengthening of Bunds
Padasavali	3	3.49	MANmA1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Sugarcane (Sc)	Bore Well	IIs	GB/strengthening of Bunds
Padasavali	4	5.53	MANmA1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Sugarcane (Sc)	Not Available	IIs	GB/strengthening of Bunds
Padasavali	5	6.25	MANmB2	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Sugarcane+Greengram (Sc+Gg)	Bore Well	Iise	GB/strengthening of Bunds
Padasavali	6	9.66	MANmB2	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Jowar+Greengram+Sugarcane+Groundnut (Rg+Jw+Gg+Sc+Gn)	Farm Pond	Iise	GB/strengthening of Bunds
Padasavali	7/1	0	Water body	Others	Others	Others	Others	Others	Others	Others	NA	Not Available	Others	Others
Padasavali	7/2	0.39	MANmB2	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Cultivated Fallow (CF)	Not Available	Iise	GB/strengthening of Bunds
Padasavali	7/3	1.2	MANmB2	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Cultivated Fallow (CF)	Not Available	Iise	GB/strengthening of Bunds
Padasavali	8	0.1	MANmB2	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	NA	Not Available	Iise	GB/strengthening of Bunds
Padasavali	10	4.24	DSImB2g1	LMU-4	Moderately shallow (50-75 cm)	Clay	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut+Greengram+Sunflower+Jowar (Gn+Gg+Sf+Jw)	Not Available	Iise	TCB/GB/strengthening of Bunds
Padasavali	12	6.35	DSImB2g1	LMU-4	Moderately shallow (50-75 cm)	Clay	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Greengram+Sunflower+Groundnut+Seasamum (Rg+Gg+Sf+Gn+Se)	Not Available	Iise	TCB/GB/strengthening of Bunds
Padasavali	29	3.79	DSImB2g1	LMU-4	Moderately shallow (50-75 cm)	Clay	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Greengram (Rg+Gg)	Not Available	Iise	TCB/GB/strengthening of Bunds

Village	Survey No.	Total Area (ha)	Soil phase	Land Management Unit	Soil Depth	Surface Soil Texture	Soil Graveliness	AWC	Slope	Soil Erosion	CLU Code	WELLS	Land Capability	Conservation Plan
Padasavali	31	6.03	DSImB2g1	LMU-4	Moderately shallow (50-75 cm)	Clay	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Greengram+Groundnut (Rg+Gg+Gn)	Not Available	IIse	TCB/GB/strengthening of Bunds
Padasavali	32	11.53	DSImC2g1	LMU-4	Moderately shallow (50-75 cm)	Clay	Gravelly (15-35%)	Medium (101-150 mm/m)	Gently sloping (3-5%)	Moderate	Redgram+Jowar+Green gram+Sugarcane+Cultivated Fallow (Rg+Jw+Gg+Sc+CF)	Not Available	IIse	TCB/GB/strengthening of Bunds
Padasavali	33	5.32	NHAmB2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Jowar (Rg+Jw)	Check Dam	IIIse	Crescent bund/TCB
Padasavali	34	6.12	NHAmB2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Greengram+Sugarcane (Rg+Gg+Sc)	Bore Well	IIIse	Crescent bund/TCB
Padasavali	35	15.37	GTTmB2g1	LMU-4	Moderately shallow (50-75 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Groundnut+Greengram (Rg+Gn+Gg)	3. Bore Well, Open Well, Farm Pond, Check Dam	IIse	TCB/GB/strengthening of Bunds
Padasavali	36	12.39	GTTmB2g1	LMU-4	Moderately shallow (50-75 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Greengram+Sunflower+Sugarcane (Rg+Gg+Sf+Sc)	Not Available	IIse	TCB/GB/strengthening of Bunds
Padasavali	37	10.23	MANmA1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Sugarcane+Jowar (Sc+Jw)	Open Well, Bore Well	IIs	GB/strengthening of Bunds
Padasavali	38/1	4.06	MANmA1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Sugarcane (Sc)	Not Available	IIs	GB/strengthening of Bunds
Padasavali	38/2	2.11	MANmA1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Sugarcane (Sc)	Not Available	IIs	GB/strengthening of Bunds
Padasavali	41	0.09	NHAmB2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	NA	Not Available	IIIse	Crescent bund/TCB
Padasavali	54	10.23	DSImB2g1	LMU-4	Moderately shallow (50-75 cm)	Clay	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Sugarcane+Redgram+Jowar (Sc+Rg+Jw)	Not Available	IIse	TCB/GB/strengthening of Bunds
Padasavali	186	2.95	NHAmB2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIIse	Crescent bund/TCB
Padasavali	187	7.83	NHAmB2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Jowar+Green gram (Rg+Jw+Gg)	Not Available	IIIse	Crescent bund/TCB
Padasavali	188	10.42	NHAmB2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram (Gg)	Not Available	IIIse	Crescent bund/TCB

Village	Survey No.	Total Area (ha)	Soil phase	Land Management Unit	Soil Depth	Surface Soil Texture	Soil Graveliness	AWC	Slope	Soil Erosion	CLU Code	WELLS	Land Capability	Conservation Plan
Padasavali	189	6.21	NHAmB2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Jowar+Greengram (Rg+Jw+Gg)	Not Available	IIIse	Crescent bund/TCB
Padasavali	194	4.33	NHAmB2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Greengram (Rg+Gg)	Not Available	IIIse	Crescent bund/TCB
Padasavali	195	9.66	DSImB2	LMU-4	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Greengram (Rg+Gg)	Not Available	IIse	TCB/GB/strengthening of Bunds
Padasavali	196	1.8	DSImB2	LMU-4	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Jowar+Greengram (Rg+Jw+Gg)	Not Available	IIse	TCB/GB/strengthening of Bunds
Padasavali	197/1	0.93	Habitation	Others	Others	Others	Others	Others	Others	Others	NA	Not Available	Others	Others
Padasavali	197/2	1.16	Habitation	Others	Others	Others	Others	Others	Others	Others	Redgram (Rg)	Not Available	Others	Others
Padasavali	197/3	0.86	Habitation	Others	Others	Others	Others	Others	Others	Others	Sugarcane (Sc)	Not Available	Others	Others
Padasavali	198/1	1.47	DSImB2	LMU-4	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIse	TCB/GB/strengthening of Bunds
Padasavali	198/2	1.18	DSImB2	LMU-4	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Cultivated Fallow (CF)	Not Available	IIse	TCB/GB/strengthening of Bunds
Padasavali	198/3	1.15	DSImB2	LMU-4	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Sugarcane (Sc)	Not Available	IIse	TCB/GB/strengthening of Bunds
Padasavali	199	3.13	DSImB2	LMU-4	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram+Sugarcane (Gg+Sc)	3.Bore Well, Open Well	IIse	TCB/GB/strengthening of Bunds
Padasavali	200	8.76	DSImB2	LMU-4	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Sugarcane+Greengram+Maize (Rg+Sc+Gg+Mz)	Open Well, Check Dam	IIse	TCB/GB/strengthening of Bunds
Padasavali	201	5.02	NHAmB2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram+Sugarcane (Gg+Sc)	Not Available	IIIse	Crescent bund/TCB
Padasavali	202	7.91	DSImB2	LMU-4	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Sugarcane+Greengram (Sc+Gg)	Not Available	IIse	TCB/GB/strengthening of Bunds
Padasavali	203	5.03	NHAmB2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram+Sugarcane (Gg+Sc)	Not Available	IIIse	Crescent bund/TCB
Padasavali	204	2.06	NHAmB2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Jowar+Green gram (Rg+Jw+Gg)	Not Available	IIIse	Crescent bund/TCB

Village	Survey No.	Total Area (ha)	Soil phase	Land Management Unit	Soil Depth	Surface Soil Texture	Soil Graveliness	AWC	Slope	Soil Erosion	CLU Code	WELLS	Land Capability	Conservation Plan
Padasavali	205	5.3	NHAmB2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Groundnut+Greengram+Sunflower (Rg+Gn+Gg+Sf)	Not Available	IIIse	Crescent bund/TCB
Padasavali	206	7.9	DSImB2	LMU-4	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Tomato+Chilly+Ladyfinger+Brinjal (Tm+Ch+Lf+Br)	4.Bore Well	IIse	TCB/GB/strengthening of Bunds
Padasavali	207	10.15	NHAmB2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Jowar+Greengram+Sunflower (Rg+Jw+Gg+Sf)	Bore Well,Check Dam	IIIse	Crescent bund/TCB
Padasavali	208	10.07	NHAmB2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Jowar+Greengram+Sunflower+Groundnut+Seasamum (Rg+Jw+Gg+Sf+Gn+Se)	Not Available	IIIse	Crescent bund/TCB
Padasavali	209/1	3.67	NHAmB2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Greengram+Sunflower+Jowar (Rg+Gg+Sf+Jw)	Not Available	IIIse	Crescent bund/TCB
Padasavali	209/2	4.7	NHAmB2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Greengram+Sunflower+Jowar (Rg+Gg+Sf+Jw)	Not Available	IIIse	Crescent bund/TCB
Padasavali	210	12.14	NHAmB2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Jowar+Greengram (Rg+Jw+Gg)	Not Available	IIIse	Crescent bund/TCB
Padasavali	211	8.74	MGTmB2g1	LMU-2	Very shallow (<25 cm)	Clay	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Jowar+Greengram (Rg+Jw+Gg)	Not Available	IIIse	Crescent bund/TCB
Padasavali	212	11.44	NHAmB2g2	LMU-3	Shallow (25-50 cm)	Clay	Very gravelly (35-60%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Jowar+Greengram (Rg+Jw+Gg)	Not Available	IIIse	Crescent bund/TCB
Padasavali	213	11.65	NHAmB2g2	LMU-3	Shallow (25-50 cm)	Clay	Very gravelly (35-60%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Sugarcane+Greengram+Jowar+Onion+Brinjal (Sc+Gg+Jw+On+Br)	Open Well, 2. Bore Well	IIIse	Crescent bund/TCB
Padasavali	214	14.96	BHImC3g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Redgram+Jowar+Greengram+Sunflower (Rg+Jw+Gg+Sf)	2. Bore Well	IIIse	Crescent bund/TCB
Padasavali	215	6.71	NHAmB2g2	LMU-3	Shallow (25-50 cm)	Clay	Very gravelly (35-60%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Sugarcane+Scrub land+Groundnut (Sc+Sl+Gn)	Farm Pond	IIIse	Crescent bund/TCB
Padasavali	216	5.57	NHAmB2g2	LMU-3	Shallow (25-50 cm)	Clay	Very gravelly (35-60%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Greengram+Sunflower (Rg+Gg+Sf)	Not Available	IIIse	Crescent bund/TCB

Village	Survey No.	Total Area (ha)	Soil phase	Land Management Unit	Soil Depth	Surface Soil Texture	Soil Graveliness	AWC	Slope	Soil Erosion	CLU Code	WELLS	Land Capability	Conservation Plan
Padasavali	217	3.79	NHAmB2g2	LMU-3	Shallow (25-50 cm)	Clay	Very gravelly (35-60%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Greengram (Rg+Gg)	Not Available	IIIse	Crescent bund/TCB
Padasavali	218	6.77	NHAmB2g2	LMU-3	Shallow (25-50 cm)	Clay	Very gravelly (35-60%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Greengram+Jowar (Rg+Gg+Jw)	Not Available	IIIse	Crescent bund/TCB
Padasavali	219	6.07	NHAmB2g2	LMU-3	Shallow (25-50 cm)	Clay	Very gravelly (35-60%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Greengram+Jowar (Rg+Gg+Jw)	Not Available	IIIse	Crescent bund/TCB
Padasavali	220	2.09	MGTmB3g1	LMU-2	Very shallow (<25 cm)	Clay	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Scrub land+Jowar (Sl+Jw)	Not Available	IVse	Crescent bund/TCB
Padasavali	274	3.19	MANmA1	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Sugarcane (Sc)	Not Available	IIse	GB/strengthening of Bunds
Padasavali	275	12.69	MGTmB3g1	LMU-2	Very shallow (<25 cm)	Clay	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Redgram+Jowar+Greengram+Scrub land (Rg+Jw+Gg+Sl)	Bore Well	IVse	Crescent bund/TCB
Padasavali	276/1	1.87	MGTmB3g1	LMU-2	Very shallow (<25 cm)	Clay	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Jowar+Groundnut (Jw+Gn)	Not Available	IVse	Crescent bund/TCB
Padasavali	276/2	1.65	DSImB2	LMU-4	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Sugarcane+Greengram+Maize (Sc+Gg+Mz)	Bore Well	IIse	TCB/GB/strengthening of Bunds
Padasavali	277/1	3.15	DSImB2	LMU-4	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Groundnut (Rg+Gn)	Bore Well	IIse	TCB/GB/strengthening of Bunds
Padasavali	277/2	2.16	DSImB2	LMU-4	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Sugarcane (Sc)	Not Available	IIse	TCB/GB/strengthening of Bunds
Padasavali	278	4.34	Habitation	Others	Others	Others	Others	Others	Others	Others	Sugarcane+Jowar (Sc+Jw)	Not Available	Others	Others
Padasavali	SETT LEMENT	8.5	Habitation	Others	Others	Others	Others	Others	Others	Others	Others	Not Available	Others	Others
Padasavali	STRE AM	2.1	MANmB2	LMU-5	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Others	Not Available	IIse	GB/strengthening of Bunds
Sarasamba	12	4.49	BHImB2g2	LMU-3	Shallow (25-50 cm)	Clay	Very gravelly (35-60%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Cultivated Fallow (CF)	Not Available	IIIse	Crescent bund/TCB
Sarasamba	13	3.02	BHImB2g2	LMU-3	Shallow (25-50 cm)	Clay	Very gravelly (35-60%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Cultivated Fallow (CF)	Not Available	IIIse	Crescent bund/TCB
Sarasamba	14	2.28	BHImB2g2	LMU-3	Shallow (25-50 cm)	Clay	Very gravelly (35-60%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Cultivated Fallow (CF)	Not Available	IIIse	Crescent bund/TCB

Village	Survey No.	Total Area (ha)	Soil phase	Land Management Unit	Soil Depth	Surface Soil Texture	Soil Graveliness	AWC	Slope	Soil Erosion	CLU Code	WELLS	Land Capability	Conservation Plan
Sarasamba	15	2.72	BHImB2g2	LMU-3	Shallow (25-50 cm)	Clay	Very gravelly (35-60%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Greengram (Rg+Gg)	Not Available	IIIse	Crescent bund/TCB
Sarasamba	16	4.44	BHImB2g2	LMU-3	Shallow (25-50 cm)	Clay	Very gravelly (35-60%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Greengram (Rg+Gg)	Not Available	IIIse	Crescent bund/TCB
Sarasamba	17	0.39	BHImB2g2	LMU-3	Shallow (25-50 cm)	Clay	Very gravelly (35-60%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIIse	Crescent bund/TCB
Sarasamba	19	4.2	BHImB2g2	LMU-3	Shallow (25-50 cm)	Clay	Very gravelly (35-60%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram (Gg)	Not Available	IIIse	Crescent bund/TCB
Sarasamba	31	0.29	MGTmB2g1	LMU-2	Very shallow (<25 cm)	Clay	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	NA	Not Available	IIIse	Crescent bund/TCB
Sarasamba	35	8.75	NHAmB2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Jowar+Greengram+Scrub land (Rg+Jw+Gg+Sl)	Not Available	IIIse	Crescent bund/TCB
Sarasamba	36	6.47	NHAmB2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Sugarcane (Rg+Sc)	Bore Well	IIIse	Crescent bund/TCB
Sarasamba	37	9.49	BHImB2g2	LMU-3	Shallow (25-50 cm)	Clay	Very gravelly (35-60%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Greengram+Sunflower (Rg+Gg+Sf)	Not Available	IIIse	Crescent bund/TCB
Sarasamba	38	6.61	NHAmB2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Groundnut+Greengram+Sugarcane+Onion+Banana (Rg+Gn+Gg+Sc+On+Bn)	Not Available	IIIse	Crescent bund/TCB
Sarasamba	39	10.43	NHAmB2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Greengram+Sunflower+Jowar (Rg+Gg+Sf+Jw)	2.Open Well, 2.Bore Well	IIIse	Crescent bund/TCB
Sarasamba	40	9.06	NHAmB2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Jowar+Greengram+Sugarcane (Rg+Jw+Gg+Sc)	Bore Well	IIIse	Crescent bund/TCB
Sarasamba	41	7.58	MGTmB2g1	LMU-2	Very shallow (<25 cm)	Clay	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Jowar+Greengram+Sunflower (Rg+Jw+Gg+Sf)	Not Available	IIIse	Crescent bund/TCB
Sarasamba	42	5.74	MGTmB2g1	LMU-2	Very shallow (<25 cm)	Clay	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Greengram+Sugarcane+Groundnut (Rg+Gg+Sc+Gn)	Not Available	IIIse	Crescent bund/TCB

Village	Survey No.	Total Area (ha)	Soil phase	Land Management Unit	Soil Depth	Surface Soil Texture	Soil Graveliness	AWC	Slope	Soil Erosion	CLU Code	WELLS	Land Capability	Conservation Plan
Sarasamba	43	9.66	NHAmB2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Greengram+Sunflower+Sugarcane (Rg+Gg+Sf+Sc)	CheckDam, Open Well, Bore Well	IIIse	Crescent bund/TCB
Sarasamba	44	8.61	NHAmB2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Greengram+Jowar (Rg+Gg+Jw)	Not Available	IIIse	Crescent bund/TCB
Sarasamba	45	9.9	BHImB2g2	LMU-3	Shallow (25-50 cm)	Clay	Very gravelly (35-60%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Cultivated Fallow (CF)	Not Available	IIIse	Crescent bund/TCB
Sarasamba	46	8.36	NHAmB2g1	LMU-3	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Greengram+Sunflower (Rg+Gg+Sf)	Not Available	IIIse	Crescent bund/TCB
Sarasamba	53	6.81	BHImB2g2	LMU-3	Shallow (25-50 cm)	Clay	Very gravelly (35-60%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Cultivated Fallow (CF)	Not Available	IIIse	Crescent bund/TCB
Sarasamba	54	7.16	BHImB2g2	LMU-3	Shallow (25-50 cm)	Clay	Very gravelly (35-60%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Cultivated Fallow+Redgram+Greengram (CF+Rg+Gg)	Not Available	IIIse	Crescent bund/TCB
Sarasamba	55	10.54	BHImB2g2	LMU-3	Shallow (25-50 cm)	Clay	Very gravelly (35-60%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Cultivated Fallow (CF)	Not Available	IIIse	Crescent bund/TCB

Appendix - II

Soil Fertility Information

Village	Survey No.	Soil Reaction (pH)	EC	Organic Carbon	Available Phosphorus	Available Potassium	Available Sulphur	Available Boron	Available Iron	Available Manganese	Available Copper	Available Zinc
Padasavali	2/1	Strongly alkaline (pH 8.4-9.0)	Non Saline (<2 dsm)	High (>0.75 %)	Medium (23-57 kg/ha)	High (>337 kg/ha)	Medium (10-20 ppm)	High (>1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Sufficient (>0.6 ppm)
Padasavali	2/2	Strongly alkaline (pH 8.4-9.0)	Non Saline (<2 dsm)	High (>0.75 %)	Medium (23-57 kg/ha)	High (>337 kg/ha)	High (>20 ppm)	High (>1.0 ppm)	Deficient (<4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Sufficient (>0.6 ppm)
Padasavali	2/3	Moderately alkaline (pH 7.8-8.4)	Non Saline (<2 dsm)	High (>0.75 %)	Medium (23-57 kg/ha)	High (>337 kg/ha)	High (>20 ppm)	High (>1.0 ppm)	Deficient (<4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Sufficient (>0.6 ppm)
Padasavali	3	Strongly alkaline (pH 8.4-9.0)	Non Saline (<2 dsm)	High (>0.75 %)	Medium (23-57 kg/ha)	High (>337 kg/ha)	High (>20 ppm)	Medium (0.5-1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Sufficient (>0.6 ppm)
Padasavali	4	Strongly alkaline (pH 8.4-9.0)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Medium (23-57 kg/ha)	High (>337 kg/ha)	Medium (10-20 ppm)	High (>1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Sufficient (>0.6 ppm)
Padasavali	5	Strongly alkaline (pH 8.4-9.0)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	Medium (10-20 ppm)	High (>1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Padasavali	6	Strongly alkaline (pH 8.4-9.0)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	Medium (10-20 ppm)	Medium (0.5-1.0 ppm)	Deficient (<4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Padasavali	7/1	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Padasavali	7/2	Strongly alkaline (pH 8.4-9.0)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	High (>20 ppm)	Medium (0.5-1.0 ppm)	Deficient (<4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Padasavali	7/3	Strongly alkaline (pH 8.4-9.0)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	High (>20 ppm)	Medium (0.5-1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Padasavali	8	Strongly alkaline (pH 8.4-9.0)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	High (>20 ppm)	Medium (0.5-1.0 ppm)	Deficient (<4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Padasavali	10	Moderately alkaline (pH 7.8-8.4)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	Medium (10-20 ppm)	Low (<0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Padasavali	12	Slightly alkaline (pH 7.3-7.8)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	Medium (10-20 ppm)	Low (<0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Padasavali	29	Neutral (pH 6.5-7.3)	Non Saline (<2 dsm)	Low (<0.5 %)	Low (<23 kg/ha)	High (>337 kg/ha)	Medium (10-20 ppm)	Low (<0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Padasavali	31	Neutral (pH 6.5-7.3)	Non Saline (<2 dsm)	Low (<0.5 %)	Low (<23 kg/ha)	High (>337 kg/ha)	Low (<10 ppm)	Low (<0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Padasavali	32	Neutral (pH 6.5-7.3)	Non Saline (<2 dsm)	Low (<0.5 %)	Low (<23 kg/ha)	High (>337 kg/ha)	Low (<10 ppm)	Low (<0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Padasavali	33	Moderately alkaline (pH 7.8-8.4)	Non Saline (<2 dsm)	Low (<0.5 %)	Low (<23 kg/ha)	Medium (145-337 kg/ha)	Low (<10 ppm)	Medium (0.5-1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)

Village	Survey No.	Soil Reaction (pH)	EC	Organic Carbon	Available Phosphorus	Available Potassium	Available Sulphur	Available Boron	Available Iron	Available Manganese	Available Copper	Available Zinc
Padasavali	34	Moderately alkaline (pH 7.8-8.4)	Non Saline (<2 dsm)	Low (<0.5 %)	Low (<23 kg/ha)	High (>337 kg/ha)	Low (<10 ppm)	Low (<0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Padasavali	35	Moderately alkaline (pH 7.8-8.4)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	Low (<10 ppm)	Medium (0.5-1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Padasavali	36	Moderately alkaline (pH 7.8-8.4)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	Low (<10 ppm)	Medium (0.5-1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Padasavali	37	Strongly alkaline (pH 8.4-9.0)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	Medium (10-20 ppm)	Medium (0.5-1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Padasavali	38/1	Strongly alkaline (pH 8.4-9.0)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	Medium (10-20 ppm)	High (>1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Padasavali	38/2	Strongly alkaline (pH 8.4-9.0)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Medium (23-57 kg/ha)	High (>337 kg/ha)	Medium (10-20 ppm)	High (>1.0 ppm)	Deficient (<4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Sufficient (>0.6 ppm)
Padasavali	41	Moderately alkaline (pH 7.8-8.4)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Medium (23-57 kg/ha)	High (>337 kg/ha)	Medium (10-20 ppm)	High (>1.0 ppm)	Deficient (<4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Sufficient (>0.6 ppm)
Padasavali	54	Moderately alkaline (pH 7.8-8.4)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	Medium (10-20 ppm)	Medium (0.5-1.0 ppm)	Deficient (<4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Padasavali	186	Moderately alkaline (pH 7.8-8.4)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	Low (<10 ppm)	Medium (0.5-1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Padasavali	187	Moderately alkaline (pH 7.8-8.4)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	Low (<10 ppm)	Medium (0.5-1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Padasavali	188	Moderately alkaline (pH 7.8-8.4)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	Low (<10 ppm)	Medium (0.5-1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Padasavali	189	Slightly alkaline (pH 7.3-7.8)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	Medium (145-337 kg/ha)	Low (<10 ppm)	Low (<0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Sufficient (>0.6 ppm)
Padasavali	194	Moderately alkaline (pH 7.8-8.4)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	Medium (145-337 kg/ha)	Low (<10 ppm)	Medium (0.5-1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Padasavali	195	Moderately alkaline (pH 7.8-8.4)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	Low (<10 ppm)	Medium (0.5-1.0 ppm)	Deficient (<4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Padasavali	196	Moderately alkaline (pH 7.8-8.4)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	Low (<10 ppm)	Medium (0.5-1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Sufficient (>0.6 ppm)
Padasavali	197/1	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Padasavali	197/2	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Padasavali	197/3	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Padasavali	198/1	Moderately alkaline (pH 7.8-8.4)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	Low (<10 ppm)	Medium (0.5-1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)

Village	Survey No.	Soil Reaction (pH)	EC	Organic Carbon	Available Phosphorus	Available Potassium	Available Sulphur	Available Boron	Available Iron	Available Manganese	Available Copper	Available Zinc
Padasavali	214	Moderately alkaline (pH 7.8-8.4)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	Low (<10 ppm)	High (>1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Padasavali	215	Moderately alkaline (pH 7.8-8.4)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	Medium (145-337 kg/ha)	Medium (10-20 ppm)	Low (<0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Padasavali	216	Slightly alkaline (pH 7.3-7.8)	Non Saline (<2 dsm)	Low (<0.5 %)	Low (<23 kg/ha)	High (>337 kg/ha)	Medium (10-20 ppm)	Low (<0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Sufficient (>0.6 ppm)
Padasavali	217	Slightly alkaline (pH 7.3-7.8)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	Medium (145-337 kg/ha)	Medium (10-20 ppm)	Medium (0.5-1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Sufficient (>0.6 ppm)
Padasavali	218	Slightly alkaline (pH 7.3-7.8)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Medium (23-57 kg/ha)	Medium (145-337 kg/ha)	Medium (10-20 ppm)	Medium (0.5-1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Sufficient (>0.6 ppm)
Padasavali	219	Moderately alkaline (pH 7.8-8.4)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	Medium (10-20 ppm)	Low (<0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Padasavali	220	Moderately alkaline (pH 7.8-8.4)	Non Saline (<2 dsm)	Low (<0.5 %)	Low (<23 kg/ha)	High (>337 kg/ha)	Low (<10 ppm)	Low (<0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Padasavali	274	Strongly alkaline (pH 8.4-9.0)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	High (>20 ppm)	Medium (0.5-1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Sufficient (>0.6 ppm)
Padasavali	275	Moderately alkaline (pH 7.8-8.4)	Non Saline (<2 dsm)	Low (<0.5 %)	Low (<23 kg/ha)	High (>337 kg/ha)	High (>20 ppm)	Medium (0.5-1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Padasavali	276/1	Moderately alkaline (pH 7.8-8.4)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	Medium (10-20 ppm)	High (>1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Padasavali	276/2	Moderately alkaline (pH 7.8-8.4)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	Low (<10 ppm)	High (>1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Padasavali	277/1	Moderately alkaline (pH 7.8-8.4)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	Medium (10-20 ppm)	Medium (0.5-1.0 ppm)	Deficient (<4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Padasavali	277/2	Moderately alkaline (pH 7.8-8.4)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	Low (<10 ppm)	Medium (0.5-1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Padasavali	278	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Padasavali	settlement	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Padasavali	STREAM	Strongly alkaline (pH 8.4-9.0)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	Medium (10-20 ppm)	Medium (0.5-1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Deficient (<0.6 ppm)
Sarasamba	12	Neutral (pH 6.5-7.3)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Medium (23-57 kg/ha)	High (>337 kg/ha)	Medium (10-20 ppm)	Low (<0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Sufficient (>0.6 ppm)
Sarasamba	13	Neutral (pH 6.5-7.3)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	Medium (10-20 ppm)	Low (<0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Sufficient (>0.6 ppm)
Sarasamba	14	Neutral (pH 6.5-7.3)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Low (<23 kg/ha)	High (>337 kg/ha)	Medium (10-20 ppm)	Low (<0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Sufficient (>0.6 ppm)

Village	Survey No.	Soil Reaction (pH)	EC	Organic Carbon	Available Phosphorus	Available Potassium	Available Sulphur	Available Boron	Available Iron	Available Manganese	Available Copper	Available Zinc
Sarasamba	54	Slightly alkaline (pH 7.3-7.8)	Non Saline (<2 dsm)	Medium (0.5-0.75 %)	Medium (23-57 kg/ha)	High (>337 kg/ha)	Medium (10-20 ppm)	Low (<0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Sufficient (>0.6 ppm)
Sarasamba	55	Neutral (pH 6.5-7.3)	Non Saline (<2 dsm)	High (>0.75 %)	High (> 57 kg/ha)	High (>337 kg/ha)	Medium (10-20 ppm)	Low (<0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (>1.0 ppm)	Sufficient (>0.2 ppm)	Sufficient (>0.6 ppm)

Appendix – III
Soil Suitability Information

Village	Survey No.	Sorghum	Maize	Red gram	Sun-flower	Cotton	Sugar-cane	Soy-bean	Guava	Mango	Sapota	Jack-fruit	Jamun	Musa-mbi	Lime	Cashew	Custard Apple	Amla	Tamarind
Padasavali	2/1	S1	S3t	S2t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S2r
Padasavali	2/2	S1	S3t	S2t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S2r
Padasavali	2/3	S1	S3t	S2t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S2r
Padasavali	3	S1	S3t	S2t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S2r
Padasavali	4	S1	S3t	S2t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S2r
Padasavali	5	S2e	S3t	S2te	S2e	S2e	S3t	S2e	S3t	S3t	S3t	S3t	S2te	S2e	S2e	N	S2e	S2e	S2re
Padasavali	6	S2e	S3t	S2te	S2e	S2e	S3t	S2e	S3t	S3t	S3t	S3t	S2te	S2e	S2e	N	S2e	S2e	S2re
Padasavali	7/1	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Padasavali	7/2	S2e	S3t	S2te	S2e	S2e	S3t	S2e	S3t	S3t	S3t	S3t	S2te	S2e	S2e	N	S2e	S2e	S2re
Padasavali	7/3	S2e	S3t	S2te	S2e	S2e	S3t	S2e	S3t	S3t	S3t	S3t	S2te	S2e	S2e	N	S2e	S2e	S2re
Padasavali	8	S2e	S3t	S2te	S2e	S2e	S3t	S2e	S3t	S3t	S3t	S3t	S2te	S2e	S2e	N	S2e	S2e	S2re
Padasavali	10	S2re	S3t	S2re	S3r	S2re	S3t	S2re	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2re	S2re	N
Padasavali	12	S2re	S3t	S2re	S3r	S2re	S3t	S2re	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2re	S2re	N
Padasavali	29	S2re	S3t	S2re	S3r	S2re	S3t	S2re	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2re	S2re	N
Padasavali	31	S2re	S3t	S2re	S3r	S2re	S3t	S2re	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2re	S2re	N
Padasavali	32	S2re	S3t	S2re	S3r	S2re	S3t	S2re	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2re	S2re	N
Padasavali	33	S3r	S3rt	S3r	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Padasavali	34	S3r	S3rt	S3r	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Padasavali	35	S2re	S3t	S2re	S3r	S2re	S3t	S2re	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2re	S2re	N
Padasavali	36	S2re	S3t	S2re	S3r	S2re	S3t	S2re	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2re	S2re	N
Padasavali	37	S1	S3t	S2t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S2r
Padasavali	38/1	S1	S3t	S2t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S2r
Padasavali	38/2	S1	S3t	S2t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S2r
Padasavali	41	S3r	S3rt	S3r	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Padasavali	54	S2re	S3t	S2re	S3r	S2re	S3t	S2re	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2re	S2re	N
Padasavali	186	S3r	S3rt	S3r	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Padasavali	187	S3r	S3rt	S3r	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Padasavali	188	S3r	S3rt	S3r	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Padasavali	189	S3r	S3rt	S3r	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Padasavali	194	S3r	S3rt	S3r	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Padasavali	195	S2re	S3t	S2re	S3r	S2re	S3t	S2re	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2re	S2re	N
Padasavali	196	S2re	S3t	S2re	S3r	S2re	S3t	S2re	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2re	S2re	N
Padasavali	197/1	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Padasavali	197/2	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Padasavali	197/3	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others

Village	Survey No.	Sorghum	Maize	Red gram	Sun-flower	Cotton	Sugar-cane	Soy-bean	Guava	Mango	Sapota	Jack-fruit	Jamun	Musa-mbi	Lime	Cashew	Custard Apple	Amla	Tamarind
Padasavali	198/1	S2re	S3t	S2re	S3r	S2re	S3t	S2re	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2re	S2re	N
Padasavali	198/2	S2re	S3t	S2re	S3r	S2re	S3t	S2re	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2re	S2re	N
Padasavali	198/3	S2re	S3t	S2re	S3r	S2re	S3t	S2re	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2re	S2re	N
Padasavali	199	S2re	S3t	S2re	S3r	S2re	S3t	S2re	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2re	S2re	N
Padasavali	200	S2re	S3t	S2re	S3r	S2re	S3t	S2re	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2re	S2re	N
Padasavali	201	S3r	S3rt	S3r	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Padasavali	202	S2re	S3t	S2re	S3r	S2re	S3t	S2re	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2re	S2re	N
Padasavali	203	S3r	S3rt	S3r	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Padasavali	204	S3r	S3rt	S3r	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Padasavali	205	S3r	S3rt	S3r	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Padasavali	206	S2re	S3t	S2re	S3r	S2re	S3t	S2re	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2re	S2re	N
Padasavali	207	S3r	S3rt	S3r	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Padasavali	208	S3r	S3rt	S3r	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Padasavali	209/1	S3r	S3rt	S3r	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Padasavali	209/2	S3r	S3rt	S3r	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Padasavali	210	S3r	S3rt	S3r	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Padasavali	211	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N
Padasavali	212	S3rg	S3rg	S3rg	N	S3rg	N	S3rg	N	N	N	N	N	N	N	N	S3rg	S3rg	N
Padasavali	213	S3rg	S3rg	S3rg	N	S3rg	N	S3rg	N	N	N	N	N	N	N	N	S3rg	S3rg	N
Padasavali	214	S3re	S3re	S3re	N	S3re	N	S3re	N	N	N	N	N	N	N	N	S3re	S3re	N
Padasavali	215	S3rg	S3rg	S3rg	N	S3rg	N	S3rg	N	N	N	N	N	N	N	N	S3rg	S3rg	N
Padasavali	216	S3rg	S3rg	S3rg	N	S3rg	N	S3rg	N	N	N	N	N	N	N	N	S3rg	S3rg	N
Padasavali	217	S3rg	S3rg	S3rg	N	S3rg	N	S3rg	N	N	N	N	N	N	N	N	S3rg	S3rg	N
Padasavali	218	S3rg	S3rg	S3rg	N	S3rg	N	S3rg	N	N	N	N	N	N	N	N	S3rg	S3rg	N
Padasavali	219	S3rg	S3rg	S3rg	N	S3rg	N	S3rg	N	N	N	N	N	N	N	N	S3rg	S3rg	N
Padasavali	220	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N
Padasavali	274	S1	S3t	S2t	S1	S1	S3t	S1	S3t	S3t	S3t	S3t	S2t	S1	S1	N	S1	S1	S2r
Padasavali	275	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N
Padasavali	276/1	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N
Padasavali	276/2	S2re	S3t	S2re	S3r	S2re	S3t	S2re	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2re	S2re	N
Padasavali	277/1	S2re	S3t	S2re	S3r	S2re	S3t	S2re	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2re	S2re	N
Padasavali	277/2	S2re	S3t	S2re	S3r	S2re	S3t	S2re	S3rt	N	S3rt	S3rt	S3r	S3r	S3r	N	S2re	S2re	N
Padasavali	278	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Padasavali	Settle-ment	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Padasavali	STREAM	S2e	S3t	S2te	S2e	S2e	S3t	S2e	S3t	S3t	S3t	S3t	S2te	S2e	S2e	N	S2e	S2e	S2re

Village	Survey No.	Sorghum	Maize	Red gram	Sun-flower	Cotton	Sugar-cane	Soy-bean	Guava	Mango	Sapota	Jack-fruit	Jamun	Musa-mbi	Lime	Cashew	Custard Apple	Amla	Tamarind
Sarasamba	12	S3rg	S3rg	S3rg	N	S3rg	N	S3rg	N	N	N	N	N	N	N	N	S3rg	S3rg	N
Sarasamba	13	S3rg	S3rg	S3rg	N	S3rg	N	S3rg	N	N	N	N	N	N	N	N	S3rg	S3rg	N
Sarasamba	14	S3rg	S3rg	S3rg	N	S3rg	N	S3rg	N	N	N	N	N	N	N	N	S3rg	S3rg	N
Sarasamba	15	S3rg	S3rg	S3rg	N	S3rg	N	S3rg	N	N	N	N	N	N	N	N	S3rg	S3rg	N
Sarasamba	16	S3rg	S3rg	S3rg	N	S3rg	N	S3rg	N	N	N	N	N	N	N	N	S3rg	S3rg	N
Sarasamba	17	S3rg	S3rg	S3rg	N	S3rg	N	S3rg	N	N	N	N	N	N	N	N	S3rg	S3rg	N
Sarasamba	19	S3rg	S3rg	S3rg	N	S3rg	N	S3rg	N	N	N	N	N	N	N	N	S3rg	S3rg	N
Sarasamba	31	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N
Sarasamba	35	S3r	S3rt	S3r	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Sarasamba	36	S3r	S3rt	S3r	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Sarasamba	37	S3rg	S3rg	S3rg	N	S3rg	N	S3rg	N	N	N	N	N	N	N	N	S3rg	S3rg	N
Sarasamba	38	S3r	S3rt	S3r	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Sarasamba	39	S3r	S3rt	S3r	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Sarasamba	40	S3r	S3rt	S3r	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Sarasamba	41	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N
Sarasamba	42	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N
Sarasamba	43	S3r	S3rt	S3r	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Sarasamba	44	S3r	S3rt	S3r	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Sarasamba	45	S3rg	S3rg	S3rg	N	S3rg	N	S3rg	N	N	N	N	N	N	N	N	S3rg	S3rg	N
Sarasamba	46	S3r	S3rt	S3r	N	S3r	N	S3r	N	N	N	N	N	N	N	N	S3r	S3r	N
Sarasamba	53	S3rg	S3rg	S3rg	N	S3rg	N	S3rg	N	N	N	N	N	N	N	N	S3rg	S3rg	N
Sarasamba	54	S3rg	S3rg	S3rg	N	S3rg	N	S3rg	N	N	N	N	N	N	N	N	S3rg	S3rg	N
Sarasamba	55	S3rg	S3rg	S3rg	N	S3rg	N	S3rg	N	N	N	N	N	N	N	N	S3rg	S3rg	N

PART-B

SOCIO-ECONOMIC STATUS OF FARM HOUSEHOLDS

CONTENTS

1	Executive summary	1-3
2	Introduction	5
3	Methodology	7-11
4	Results and discussions	13-28

LIST OF TABLES

I. Social status		
1	Human population among sample households	13
2	Basic needs of sample households	15
II. Economic status		
3	Occupational pattern in sample households	15
4	Domestic assets among samples households	15
5	Farm assets among samples households	17
6	Livestock assets among sample households	17
7	Milk produced and Fodder availability of sample households	18
8	Women empowerment of sample households	18
9	Per capita daily consumption of food among the sample households	18
10	Annual average Income from various sources	19
11	Average annual expenditure of sample households	20
12	Distribution of land holding among the sample households	21
13	Land holding among samples households	21
III. Resource use pattern		
14	Present cropping pattern among samples households	22
15	Distribution of soil series in the watershed	22-23
IV. Economic land evaluation		
16	Cropping pattern on major soil series	23
17	Alternative land use options for different size group of farmers (Benefit Cost Ratio)	24
18	Economics Land evaluation and bridging yield gap for different crops	24
19	Estimation of onsite cost of soil erosion	25
20	Ecosystem services of food grain production	27
21	Ecosystem services of water supply for crop production	27
22	Farming constraints	28

LIST OF FIGURES

1	Location of study area	8
2	ALPES Framework	9
3	Basic needs of sample households	14
4	Domestic assets among the sample households	16
5	Farm assets among samples households	16
6	Livestock assets among sample households	17
7	Per capita daily consumption of food among the sample households	19
8	Average annual expenditure of sample households	20
9	Present cropping pattern	22
10	Estimation of onsite cost of soil erosion	26
11	Ecosystem services of food grain production	26
12	Ecosystem services of water supply	27

EXECUTIVE SUMMARY

Baseline socioeconomic characterisation is prerequisite to prepare action plan for program implementation and to assess the project performance before making any changes in the watershed development program. The baseline provides appropriate policy direction for enhancing productivity and sustainability in agriculture.

Methodology: *Padasavli-1 micro-watershed (Padasavli sub-watershed, Aland taluk, Gulbarga district) is located in between 17°35'–17°36' North latitudes and 76°25'–76°28' East longitudes, covering an area of about 534.63 ha, bounded by Chincholi Khurd, Hiroli, Nagalogaon and Matki villages with length of growing period (LGP) 120-150 days. We used soil resource map as basis for sampling farm households to test the hypothesis that soil quality influence crop selection, and conservation investment of farm households. The level of technology adoption and productivity gaps and livelihood patterns were analyses. The cost of soil degradation and ecosystem services were quantified.*

Results: *The socio-economic outputs for Padasavli-1 micro-watershed (Padasavli sub-watershed, Aland taluk, Gulbarga district) are presented here.*

Social Indicators

- ❖ *Male and female ratio is 57 to 42.1 Per cent to the total sample population.*
- ❖ *Younger age 18 to 50 years group of population is 60.2 around per cent to the total population.*
- ❖ *Literacy population is around 76.4 per cent.*
- ❖ *Social groups belong to other backward caste (OBC) among all sample households.*
- ❖ *Firewood is the source of energy for a cooking among all sample households.*
- ❖ *About 20 per cent of households have a yashaswini health card.*
- ❖ *Dependence on ration cards for food grains through public distribution system is around 89 per cent of sample households.*
- ❖ *Swach bharrath program providing closed toilet facilities around 30 per cent of sample households*
- ❖ *Women participation in decisions making for agriculture production among all the sample households.*

Economic Indicators

- ❖ *The average land holding is 2.67 ha indicates that majority of farm households are belong to small and medium farmers. The total cultivated area by dry land condition among the sample farmers.*
- ❖ *Agriculture is the main occupation among 44.7 per cent and agriculture is the main and agriculture labour is a subsidiary occupation about 55.3 per cent of sample households.*

- ❖ *The average value of domestic assets is around Rs. 6833. per household. Mobile and television are popular media mass communication.*
- ❖ *The average value of farm assets is around Rs. 3721 per household, about 50 per cent of sample farmers having plough and bullock cart (50 %).*
- ❖ *The average value of livestock is around Rs. 28125 per household; about 78.5 per cent of household are having livestock.*
- ❖ *The average per capita food consumption is around 844 grams (1774 kilo calories) against national institute of nutrition (NIN) recommendation at 827 gram. Around 60 per cent of sample households are consuming less than the NIN recommendation.*
- ❖ *The annual average income is around Rs. 65120 per household. About 70 per cent of farm households are below poverty line.*
- ❖ *The per capita average monthly expenditure is around Rs.1995.*

Environmental Indicators-Ecosystem Services

- ❖ *The value of ecosystem service helps to support investment to decision on soil and water conservation and in promoting sustainable land use.*
- ❖ *The onsite cost of different soil nutrients lost due to soil erosion is around Rs. 1260 per ha/year. The total cost of annual soil nutrients is around Rs. 640239 per year for the total area of 534 ha.*
- ❖ *The average value of ecosystem service for food grain production is around Rs 7231/ha/year. Per hectare food grain production services is maximum in red gram (Rs. 15388) followed by sunflower (Rs. 5371) and greengram (Rs.934).*
- ❖ *The data on water requirement for producing one quintal of grain is considered for estimating the total value of water required for crop production. The per hectare value of water used and value of water was maximum in red gram (Rs. 42840), followed by green gram (Rs. 34116) and sunflower (Rs. 24122).*

Economic Land Evaluation

- ❖ *The major cropping pattern is red gram (83.7 %) followed by green gram (8.7 %) and sunflower (7.6 %).*
- ❖ *In Padasavali-1 Microwatershed, major soil series are Novinihala series are having shallow soil depth covers around 38.7 % of area the major crops are green gram (60 %) and redgram (40 %). Mahagaon soil series having are very deep soils depth covers around 9.22 % of area the crops are redgram. Marugutti soil series are having very shallow depth covers around 9.2 % of area the crops are redgram (44.4 %) and sunflower (55.6 %). Gutti soil series are having moderately shallow depth covers around (4.64) of area the crops are red gram.*
- ❖ *The total cost of cultivation and benefit cost ratio (BCR) in study area for red gram ranges between Rs. 32931/ha in NHA soil (with BCR of 1.13) and Rs. 18116/ha in MAN soil (with BCR of 2.21).*

- ❖ *In sunflower the cost of cultivation in MGT soil is Rs.16128/ha (with BCR of 1.33) and green gram the cost of cultivation in NHA soil is Rs.18826/ha (with BCR of 1.08).*
- ❖ *The land management practices reported by the farmers are crop rotation, tillage practices, fertilizer application and use of farm yard manure (FYM). Due to higher wages farmer are following labour saving strategies is not prating soil and water conservation measures. Less ownership of livestock limiting application of FYM.*
- ❖ *It was observed soil quality influences on the type and intensity of land use. More fertilizer applications in deeper soil to maximize returns.*

Suggestions

- ❖ *Involving farmers in watershed planning helps in strengthening institutional participation.*
- ❖ *The per capita food consumption and monthly income is very low. Diversifying income generation activities from crop and livestock production in order to reduce risk related to drought and market prices.*
- ❖ *Majority of farmers reported that they are not getting timely support/extension services from the concerned development departments.*
- ❖ *By strengthening agricultural extension for providing timely advice improved technology there is scope to increase in net income of farm households.*
- ❖ *By adopting recommended package of practices by following the soil test fertiliser recommendation, there is scope to increase yield in red gram (40.1 to 50.3 %), followed by sunflower (56.0 %) and greengram (42.2 %).*

INTRODUCTION

Watershed Development program aim to restore degraded watersheds in rainfed regions to increase their capacity to capture and store rain water, reduce soil erosion, and improved soil nutrients and carbon contents so they can produce greater agricultural yields and other benefits. As majority of rural poor live in these regions and dependent on natural resources for their livelihood and sustenance, improvements in agricultural yields improve human welfare and simultaneously improve national food security.

Sujala–III watershed development project conceptualised and implemented by the Watershed Development Department of Government of Karnataka with tripartite cost-sharing arrangements. The World Bank through International Development Association provided major portion of plan outlay as a loan to Government of India and in turn loan to Government of Karnataka.

The objectives of Sujala-III is to demonstrate more effective watershed management through greater integration of programs related to rain fed agriculture, innovative and science based approaches and strengthened institutions and capacities. The project is implemented in 11 districts of Bidar, Vijayapura, Gulbarga, Yadgir, Koppal, Gadag, Raichur, Davanagere, Tumkur, Chikkamangalur and Chamarajanagar which have been identified by the Watershed Development Department based on rainfall and socio-economic conditions. The project will be implemented over six years and linked with the centrally financed integrated watershed management programme.

Economic evaluations can better guide in watershed planning and implementation, as well as raise awareness of benefits of ecosystem restoration for food security and poverty alleviation program. The present study aims to characterize socio-economic status of farm households, assess the land and water use status, evaluate the economic viability of land use, prioritize farming constraints and suggest the measures for soil and water conservation for sustainable agriculture.

Objectives of the study

1. To characterize socio-economic status of farm households
2. To evaluate the economic viability of land use and land related constraints
3. To estimate the ecosystem service provided by the watershed and
4. To suggest alternatives for sustainable agriculture production.

METHODOLOGY

Study area

Padasavli-1 micro-watershed is located in southern dry zone of Karnataka (Figure 1) the total geographic area of this zone is about 1.76 M ha covering 8 taluks of Gulbarga district and 3 taluks of Raichur. Net cultivated area in the zone is about 1.31 M ha of which about 0.09 M ha are irrigated. The mean elevation of the zone is 300-450 m MSL. The main soil type is deep to very deep soils with small pockets of shallow to medium black soils. The zone is cropped predominantly during rabi due to insufficient rainfall (465-785 mm). The principal crops of the zone are jowar, bajra, oilseeds, pulses, cotton and sugarcane. It represented Agro Ecological Region (AER)-3 having length of growing period (LPG) 120-150 days.

Padasavli-1 micro-watershed (Padasavli sub-watershed, Aland taluk, Gulbarga district) is located in between 17⁰35'–17⁰36' North latitudes and 76⁰25'–76⁰28' East longitudes, covering an area of about 534.63 ha, bounded by Chincholi Khurd, Hiroli, Nagalogaon villages.

Sampling Procedure

In this study we have followed soil variability as criterion for sampling the farm households. In each micro-watershed the survey numbers and associated soil series are listed. Minimum three farm households for each soil series were taken and summed up to arrive at total sample for analysis.

Sources of data and analysis

For evaluating the specific objectives of the study, primary data was collected from the sample respondents by personal interview method with the help of pre-tested questionnaire. The data on socio-economic characteristics of respondents such as family size and composition, land holdings, asset position, occupational pattern and education level was collected. The present cropping pattern and the level of input use and yields collected during survey. The data collected from the representative farm households were analysed using Automated Land Potential Evaluation System (Figure 2).

LOCATION MAP OF PADASAVLI-1 MICRO-WATERSHED





Figure 1: Location of study area

Steps followed in socio-economic assessment

- 1 • After the completion of soil profile study link the cadastral number to the soil profile in the micro watershed.
- 2 • Download the names of the farmers who are owning the land for each cadastral number in the Karnataka BHOOMI Website.
- 3 • Compiling the names of the farmers representing for all the soil profiles studied in the micro watershed for socio-economic Survey.
- 4 • Conducting the socioeconomic survey of selected farm households in the micro watershed .
- 5 • Farm households database created using the Automated Land Potential Evaluation System (ALPES) for analysis of socio economic status for each micro watershed .
- 6 • Synthesis of tables and preparation of report for each micro watershed .



The sample farmers were post classified in to marginal and small (0.0 to ≤ 2 ha), medium and semi medium (>2 to ≤ 10 ha) and large (>10 ha). The steps involved in estimation of soil potential involve estimation of total cost of cultivation, the yield/gross returns and net income per hectare. The cost of inputs such seed, manure and fertilizer, plant protection chemicals, payment towards human and bullock labour and interest on working capita are included under operational costs. In the case of perennial crops, the cost of establishment was estimated by using actual physical requirements and prevailing market prices. Estimation cost included maintenance cost up to bearing period. The value of main product and by product from the crop enterprise at the market rates were the gross returns of the crop. Net returns were worked out by deducting establishment and maintained cost from gross returns.

Operational Cost = cost of seeds, fertilizers, pesticides. Cost of human and bullock labour, cost of machinery, cost of irrigation water + interest on working capital.

Gross returns = Yield (Quintals/hectare)*Price (Rs/Quintal)

Net returns = Gross returns-Operational cost.


Benefit Cost Ratio = Net returns/Total cost.

Economic suitability classes: once each land use –land area combination has been assigned an economic value by the land evaluation, the question arises as to its ‘suitability’, that is, the degree to which it satisfies the land user. The FAO framework defines two suitability orders: ‘S’(suitable if benefit cost ratio (BCR) >1) and ‘N’(not suitable if (BCR <1), which are divided into five economic suitability classes: ‘S1’(highly suitable if BCR >3), ‘S2’(suitable if BCR >2 and <3), ‘S3’(Marginally suitable if BCR >1 and <2), ‘N1’(Not suitable for economic reasons but physically suitable) and ‘N2’(not suitable for physical reasons). The limit between ‘S3’ and ‘N1’ must be at least at the point of financial feasibility (i.e. net returns, NPV, or IRR >0 and BCR >1). The other limits depend on social factors such as farm size, family size, alternative employment or investment possibilities and wealth expectations; these need to be specified for the soil series.

Economic Valuation of Soil ecosystem services

The replacement cost approach was followed for estimating the onsite cost of soil erosion, Market price method was followed for estimating the value of food and fodder production. Value transfer methods was followed for estimating the value of water demand by different crops in the micro watershed.

Steps followed in Replacement cost methods for estimation of onsite cost of soil erosion



RESULTS AND DISCUSSIONS

The demographic information shows that the household population dynamics encompasses the socioeconomic status of the farmer. For a rural family, the household size should be optimal to earn a comfortable livelihood through farm and non-farm wage earning. The total number of population in watershed area was 38, out of which 57.8 per cent were males and 42.11 per cent females. Average family size of the households is 3.8. Age is an important factor, which affects the potential employment and mobility status of respondents. The data on age wise distribution of farmers in the sample households indicated that majority of the farmers are coming under the age group of 30 to 50 years (42.11 %) followed by 0 to 18 years (23.6 %), 18 to 30 years (18.4 %) and more than 50 years (15.7 %). Hence, in the study area in general, the respondents were of young and middle age, indicating thereby that the households had almost settled with whatever livelihood options they were practicing and sample respondents were young by age who could venture into various options of livelihood sources. Data on literacy indicated that 23.6 per cent of respondents were illiterate and 76.4 per cent literate (Table 1).

Table 1: Human population among sample households in Padasavali-1 Microwatershed

Particulars	Units	Value
Total human population in sample HHs	Number	38.0
Male	% to total Population	57.8
Female	% to total Population	42.11
Average family size	Number	3.8
Age group		
0 to 18 years	% to total Population	23.6
18 to 30 years	% to total Population	18.4
30 to 50 years	% to total Population	42.11
>50 years	% to total Population	15.7
Average age	Age in years	34.1
Education Status		
Illiterates	% to total Population	23.6
Literates	% to total Population	76.4
Primary School (<5 class)	% to total Population	15.7
Middle School (6- 8 class)	% to total Population	21.5
High School (9- 10 class)	% to total Population	23.6
Others	% to total Population	15.7

The ethnic groups among the sample farm households found to be among all belonging to other backward caste (OBC) (Table 2 and Figure 3). All the sample households are using firewood as source of fuel for cooking. All the sample farmers are having electricity connection. About 20 per cent are sample households having health cards. About 89.0 per cent of farm households are having ration cards for taking food

grains from public distribution system. About 30 per cent of farm households are having toilet facilities.




Figure 3: Basic needs of sample households in Padasavali-1 Microwatershed

The data collected on the source of drinking water in the study area is presented in Table 2. All the sample respondents are having Tube well source for water supply for domestic purpose.

Table 2: Basic needs of sample households in Padasavali-1 Microwatershed

Particulars	Units	Value
Social groups		
OBC	% of Households	100
Types of fuel use for cooking		
Firewood	% of Households	100
Energy supply for home		
Electricity	% of Households	100
Number of households having Health card		
Yes	% of Households	20.0
No	% of Households	80.0
MGNREGA Card		
No	% of Households	100
Ration Card		
Yes	% of Households	89.0
No	% of Households	11.0
Households with toilet		
Yes	% of Households	30.0
No	% of Households	70.0
Drinking water facilities		
Tube well	% of Households	100

The occupational pattern (Table 3) among sample households shows that agriculture is the main occupation around 44.7 per cent followed by subsidiary occupation is agriculture labour 55.3 per cent of sample households.

Table 3: Occupational pattern in sample population in Padasavali-1 Microwatershed

Occupation		% to total
Main	Subsidiary	
Agriculture	Agriculture	44.7
	Agriculture Labour	55.3
Grand Total		100.0
Family labour availability		Man days/month
Male		25.0
Female		20.0
Total		45.0

Table 4: Domestic assets among the sample households in Padasavali-1 Microwatershed

Particulars	% of households	Average value in Rs
Mobile phone	90	2666
Television	100	11000
Average value	6833	

The important assets especially with reference to domestic assets were analyzed and are given in Table 4 and Figure 4. The important domestic assets possessed by all categories of farmers are television (100 %) and mobile phone (90 %). The average value of domestic assets is around Rs 6833 per households.




Figure 4: Domestic assets among the sample households in Padasavali-1 Microwatershed




Figure 5: Farm assets among samples households in Padasavali-1 Microwatershed

The most popularly owned farm equipments were sickles, plough, cattle shed; pump sets, chaff cutter, bullock cart, sprayer and thresher. Plough and sickle were commonly present in all the sampled farmers; these were primary implements in agriculture. The per cent of households owned plough (50.0 %), bullock cart (50.0 %), weeder (40 %) and sprayer (10 %). The average value of farm assets is around Rs 3721 per households (Table 5 and Figure 5).

Table 5: Farm assets among samples households in Padasavali-1 Microwatershed

Particulars	% of households	Average value in Rs
Plough	50.0	5000
Bullock cart	50.0	7100
Sprayer	10	2500
Weeder	40.0	287
Average value		3721

Table 6: Livestock assets among sample households in Padasavali-1 Microwatershed

Particulars	% of livestock population	Average value in Rs
Local dry cow	18.1	12500
Local Milching Cow	9.9	15000
Crossbred Dry cow	9.9	15000
Bullocks	63.6	70000
Average value		28125




Figure 6: Livestock assets among sample households in Padasavali-1 micro-watershed

Livestock is an integral component of the conventional farming systems (Table 6 and Figure 6). The highest livestock population is bullocks were around 63.6 per cent, local dry cow (18.1 %), local milching cow (9.9 %) and crossbred dry cow (9.9 %). The average livestock value was Rs. 28125 per household.

Average milk produced in sample households is 900 liters/ annum among the farm households. Livestock having population are 78.5 percent and 18 livestock population numbers in the livestock. About 1250 kg of average fodder is available per season for the livestock feeding (Table 7).

Table 7: Milk produced and fodder availability of sample households in Padasavali-1 Microwatershed

Particulars	
Name of the Livestock	Ltr./Lactation/animal
Local Milching Cow	900
Fodder produces	
Fodder yield (kg/ha.)	
Sorghum	1250
Livestock having households (%)	78.5
Livestock population (Numbers)	18

A woman participation in decision making in this Microwatershed is presented in Table 8. Among all women earning for her family requirement and among all women taking decision in her family and agriculture related activities in these study area.

Table 8: Women empowerment of sample households in Padasavali-1 Microwatershed % to Grand Total

Particulars	Yes	No
Women participation in local organization activities	0.0	100.0
Women elected as panchayat member	0.0	100.0
Women earning for her family requirement	100.0	0.0
Women taking decision in her family and agriculture related activities	100.0	0.0

Table 9: Per capita daily consumption of food among the sample households in Padasavali-1 Microwatershed

Particulars	NIN recommendation (gram/ per day/ person)	Present level of consumption (gram/ per day/ person)	Kilo Calories /day/person
Cereals	396.0	325	1106
Pulses	43.0	46	160
Milk	200.0	143	93
Vegetables	143.0	147	35
Cooking Oil	31.0	25	145
Egg	0.5	136	204
Meat	14.2	18	28
Total	827.7	844	1774
Threshold of NIN recommendation		827 gram*	2250 Kcal*
% Below NIN		60.0	90.0
% Above NIN		40.0	10.0

The food intake in terms of kilo calorie (kcal) per person per day was calculated and presented in the Table 9 and Figure 7. More quantity of cereals is consumed by sample farmers which accounted for 1106 kcal per person. The other important food

items consumed was pulses 160 kcal followed by milk 93 kcal, vegetables 35 kcal, cooking oil 145 kcal, egg 204 kcal and meat 28 kcal. In the sampled households, farmers were consuming less (1774 kcal) than NIN- recommended food requirement (2250 kcal).

Note: * day/person




Figure 7: Per capita daily consumption of food among the sample households in Padasavali-1 Microwatershed

Table 10: Annual average income of HHs from various sources in Padasavali-1 Microwatershed

Particulars	Income *
Nonfarm income	0 (0)
Livestock income (Rs)	19640 (10)
Crop Production (Rs)	45480 (100)
Total Annual Income (Rs)	65120
Average monthly per capita income (Rs)	1428
Threshold for Poverty level (Rs 975 per month/person)	
% of households below poverty line	70.0
% of households above poverty line	30.0

* Figure in the parenthesis indicates % of Households

Annual income of the sample HHs: The average annual household income is around Rs 65120. Major source of income to the farmers in the study area is from crop production (Rs 45480). The monthly per capita income is Rs. 1428, which is above than the threshold monthly income of Rs 975 for considering above poverty line. Due to the

fact that erratic rainfall and shortage of water, farmers are diverting from crop production activities to enable the household for a comfortable livelihood. The incomes from the other aforesaid sources are very meagre (Table 10).

The average annual expenditure of farm households indicated that farmers in the study area spend highest on food (Rs. 35076) followed by education, clothing, social function and health. Now a day's education is most important among all of us. In today's competitive world, education is a necessity for man after food, clothing, and shelter. It is the only fundamental way by which a desired change in the society can happen. The average per capita monthly expenditure is around Rs 1995 and about 70 per cent of farm households are below poverty line (Table 11 and Figure 8).

Table 11: Average annual expenditure of sample HHs in Padasavali-1 Microwatershed

Particulars	Value in Rupees	Per cent
Food	35076	52.5
Education	3600	2.5
Clothing	4400	8.5
Social functions	43000	18.4
Health	4900	18.1
Total Expenditure (Rs/year)	90976	100
Monthly per capita expenditure (Rs)	1995	




Figure 8: Average annual expenditure of sample HHs in Padasavali-1 Microwatershed

Land holding: Total area cultivated by them is 26.6 ha. The average land holding of sample HHs is 2.7 ha. Large number of sample HHs (40%) belong to small size group with an average holding size of 1.1 ha followed by medium farmer (40%) with a average land holding size of 2.9 ha and large farmers (20%) with a average land holding size of 2.7 ha (Table 12)

Table 12: Distribution of land holding among the sample households in Padasavali-1 Microwatershed

Particulars	Units	Values
Small farmers		
Total land	ha	4.48
Sample size	Per cent	40.0
Average land holding	ha	1.12
Medium farmers		
Total land	ha	11.6
Sample size	Per cent	40
Average land holding	ha	2.92
Large farmers		
Total land	ha	10.5
Sample size	Per cent	20
Average land holding	ha	5.27
Total sample households		
Total land	ha	26.6
Sample size	Per cent	100
Average land holding	ha	2.67

Land use: The total land holding in the Padasavali-1 Microwatershed is 26.6 ha (Table 13).of which 26.6 ha is dry land. The average land holding per household is worked out to be 2.67 ha.

Table 13: Land use among samples households in Padasavali-1 Microwatershed

Particulars	Per cent	Area in ha
Dry land	100	26.6
Irrigated land	0.0	0.0
Fallow Land	0.0	0.0
Total land holding	100	26.6
Average land holding	2.67	

In the micro-watershed, the prevalent present land uses under perennial plants are neem tress (100 %).

The land use decisions are usually based on experience of farmers, tradition, expected profit, personal preferences, resources and social requirements.

Table 14: Present cropping pattern and cropping intensity in Padasavali-1 Microwatershed% to Grand Total

Crops	Kharif	Grand Total
Redgram	83.7	83.7
Greengram	8.7	8.7
Sunflower	7.6	7.6
Grand Total	100	100

The present dominant crops grown in dry lands in the study area were by red gram (83.7 %), green gram (8.7 %) and sunflower (7.6 %).which are taken during kharif (Table 14 season (Figure 9).




Figure: 9 Present cropping pattern and cropping intensity in Padasavali-1 Microwatershed

Economic land evaluation

The main purpose to characterise the socio-economic systems in the watershed is to identify the existing production constraints and propose the potential/alternate options for agro-technology transfer and for bridging the adoption and yield gap.

In Padasavali-1 Microwatershed, 7 soil series are identified and mapped (Table 15). The distribution of major soil series are Novinihala covering on area 206 (38.7 %) followed by Dinsi 86.5 ha (14.5 %), Bhimanahalli 72.5 ha (13.5 %), Mahagaon 49.2ha (49.2 %), Margutti 47.3 ha (9.02 %), Gutti 24.8 ha (4.64 %) and Kamalapur 7.96 ha (1.49 %).

Table 15: Distribution of soil series in Padasavali-1 Microwatershed

Sl. No	Soil Series	Description	Area in ha (%)
1	MGT	Very shallow, black gravelly clay soils developed from weathered basalt on gently sloping uplands; sandy clay surface on 3-5 % slope, severely eroded, slightly gravelly, 15-35 per cent gravels.	47.3 (9.02)
2	NHA	Shallow, black clayey soils developed from weathered basalt on very gently sloping uplands; clay surface on 1-3% slope, slightly eroded, slightly gravelly, 15-35 per cent gravels.	206.2 (38.7)
3	BHI	Shallow, black clay soils developed from weathered basalt on very gently sloping uplands; clay surface on 1-3% slope, moderately eroded, moderately gravelly, 35-60 per cent gravels.	72.5 (13.52)
4	DSI	Moderately shallow, black clay soils developed from weathered basalt on very gently sloping uplands; clay surface on 1-3% slope, moderately eroded	86.50 (14.59)
5	GTT	Moderately shallow, black clayey soils developed from weathered basalt on very gently sloping uplands; clay surface on 1-3% slope, moderately eroded , slightly gravelly, 15-35 per cent gravels.	24.80 (4.64)
6	KMP	Moderately deep, black clayey soils developed from weathered basalt on very gently sloping uplands; clay surface on 1-3% slope, moderately eroded , slightly gravelly, 15-35 per cent gravels.	7.96 (1.49)
7	MAN	Very deep, black clayey soils developed from weathered basalt on nearly level uplands; clay surface on 0-1% slope, slightly eroded	49.27 (9.22)

Present cropping pattern on different soil series are given in Table 16. Crops grown on Dhoandothi soils are redgram and sorghum. Redgram on Dargah soils is grown

Table 16: Cropping pattern on major soil series in Padasavali-1 Microwatershed (Area in per cent)

Soil Series	Soil Depth	Crops	Dry	Grand Total
			Kharif	
Margutti	Very shallow (<25 cm)	Redgram	44.4	44.4
		Sunflower	55.9	55.9
Novinihala	Shallow (25-50 cm)	Greengram	60	60
		Redgram	40	40
Gutti	Moderately shallow (50-75 cm)	Redgram	100	100
Mahagaon	Very deep (>150 cm)	Redgram	100	100

Land is used for agricultural use for growing cereals, pulse, oilseeds and commercial crops. The soil/ land potential are measures in terms of physical yield and net income. The alternative land use options for each Microwatershed are given below (Table 17).

Table 17: Alternative land use options for different size group of farmers (Benefit Cost Ratio) in Padasavali-1 Microwatershed.

Soil Series	Small farmers	Medium farmers	Large farmers
GTT		Redgram (1.95)	
MAN	Redgram (1.25)	Redgram (1.58)	Redgram (3.01)
MGT	Redgram (1.49)	Sunflower (1.33)	
NHA	Green gram (1.08) & Redgram (1.13).	Redgram (2.52).	Redgram (2.52).

The productivity of different crops grown in Padasavali-1 Microwatershed under potential yield of the crops is given in Table 18.

Table18: Economic land evaluation and bridging yield gap for different crops in Padasavali-1 Microwatershed

Particulars	MGT (<25 cm)		NHA (25-50 cm)		GTT (50-75cm)	MAN (>150 cm)
	Red gram	Sun flower	Green gram	Red gram	Red gram	Red gram
Total cost (Rs/ha)	20422	16128	18826	32931	19150	18116
Gross Return (Rs/ha)	30344	21499	20254	37050	37455	34873
Net returns (Rs/ha)	9922	5371	1428	4119	18305	16757
BCR	1.49	1.33	1.08	1.13	1.95	2.21
Farmers Practices (FP)						
FYM (t/ha)	1.8	1.5	1.7	2.5	2.0	1.6
Nitrogen (kg/ha)	78.6	88.5	99.2	148.8	111.9	67.6
Phosphorus (kg/ha)	56.5	55.6	57.5	86.3	69.1	60.9
Potash (kg/ha)	0.0	0.0	0.0	0.0	0.0	0.0
Grain (Qtl/ha)	6.1	7.3	5.0	7.5	10.2	7.4
Price of Yield (Rs/Qtl)	5000	3000	4000	5000	3750	4750
Soil test based fertilizer Recommendation (STBR)						
FYM (t/ha)	7.4	6.6	7.4	7.4	7.4	7.4
Nitrogen (kg/ha)	30.9	55.2	18.5	24.7	24.7	21.6
Phosphorus (kg/ha)	61.8	74.1	46.3	61.8	61.8	61.8
Potash (kg/ha)	18.5	37.1	27.8	24.7	18.5	18.5
Grain (Qtl/ha)	12.4	16.5	8.6	12.4	12.4	12.4
% of Adoption/yield gap (STBR-FP) / (STBR)						
FYM (%)	75.1	78.0	77.5	66.3	72.3	78.4
Nitrogen (%)	-154.7	-60.4	-435.3	-502.2	-353.0	-212.8
Phosphorus (%)	8.5	25.0	-24.2	-39.7	-11.9	1.3
Potash (%)	100.0	100.0	100.0	100.0	100.0	100.0
Grain (%)	50.3	56.0	42.2	39.3	17.0	40.1
Value of yield and Fertilizer (Rs)						
Additional Cost (Rs/ha)	5595	6291	4839	2837	4361	5664
Additional Benefits (Rs/ha)	31037	27640	14580	24250	7890	23532
Net change Income (Rs/ha)	25442	21349	9741	21413	3530	17868

The data on cost of cultivation and benefit cost ratio (BCR) of different crops is given in Table 18. The total cost of cultivation in study area for red gram ranges between

Rs. 32931/ha in NHA soil (with BCR of 1.13) and Rs. 18116/ha in MAN soil (with BCR of 2.21), sunflower cost of cultivation in MGT soil is Rs.16128/ha (with BCR of 1.33) and greengram cost of cultivation in NHA soil is Rs.18826/ha (with BCR of 1.08).

The data on FYM, Nitrogen, Phosphorus and Potash application by the farmers to different crops and recommended FYM for different crops is given in Table 18. There is a huge gap between FYM application by farmers and recommended FYM in all the crops across the soils. There is a larger yield gap in crops grown across different soil series. Adequate knowledge about recommended package of practices is the pre-requisite for their use in cultivation of crops. It is a fact that, recommended practices are major contributing factors to yield. Inadequate knowledge about recommended practices leads to their improper adoption. Strengthening of extension services by concerned agency is required to increase adoption of recommended cultivation practices and ultimately reducing the gap. By adopting soil-test fertiliser recommendation, there is scope to increase yield and income to a maximum of Rs 25442 in redgram a minimum of Rs 3530 in redgram cultivation.

Economic valuation of Ecosystem Services (ES) was aimed at combining use and non-use values to determine Total Economic Value (TEV) of ES. Ecosystem Services (ES) were valued based on their annual flow or utilization in common monetary units, Rs/year. The valuation of ES was based on market price in 2017 or market cost approaches whichever is applicable, and in other cases on value or benefit transfer from previous valuation studies.

The onsite cost of different soil nutrients lost due to soil erosion is given in Table 19 and Figure 10. The average value of soil nutrient loss is around Rs 1260 per ha/year. The total cost of annual soil nutrients is around Rs.640239 per year for the total area of 534 ha.

Table 19: Estimation of onsite cost of soil erosion in Padasavali-1 Microwatershed

Particulars	Quantity(kg)		Value (Rs)	
	Per ha	Total	Per ha	Total
Organic matter	167.77	85227	1056.94	536927
Phosphorous	0.10	50	4.36	2216
Potash	3.29	1672	65.82	33434
Iron	0.18	92	8.72	4429
Manganese	0.31	157	84.95	43153
Copper	0.06	29	32.00	16256
Zinc	0.01	4	0.33	169
Sulphurs	0.17	86	6.81	3457
Boron	0.01	5	0.39	198
Total	171.89	87322	1260.31	640239




Figure 10: Estimation of onsite cost of soil erosion in Padasavali-1 Microwatershed

The average value of ecosystem service for food grain production is around Rs 7231/ ha/year (Table 20 and Figure 11). Per hectare a food grain production service is maximum in red gram (Rs 15388) followed by sunflower (Rs 5371) and greengram (Rs.934).




Figure: 11 Ecosystem services of food grain production in Padasavali-1 Microwatershed

The water demand for production of different crops was worked out in arriving at the ecosystem services of water support to crop growth. The data on water requirement for producing one quintal of grain is considered for estimating the total value of water

required for crop production. The per hectare value of water used and value of in redgram (Rs.42840) followed by green gram (Rs. 34116) and sunflower (Rs.24122) (Table 21 and Figure 12)

Table 20: Ecosystem services of food grain production in Padasavali-1 Microwatershed

Production items	Crops	Area in ha	Yield (Qtl/ha)	Price (Rs/Qtl)	Gross Returns (Rs/ha)	Cost of Cultivation (Rs/ha)	Net Returns (Rs/ha)
Pulses	Greengram	1.2	5	4000	19760	18826	934
	Redgram	23.4	8	4563	35903	20514	15388
Oil seeds	Sunflower	2.1	7	3000	21499	16128	5371
Average value		26.7	6.6	3854	25720	18489	7231

Table 21: Ecosystem services of water supply in Padasavali-1 Microwatershed

Crops	Yield (Qtl/ha)	Virtual water (cubic meter) per ha	Value of Water (Rs/ha)	Water consumption (Cubic meters/Qtl)
Greengram	4.9	3412	34116	691
Redgram	7.9	4284	42840	544
Sunflower	7.2	2412	24122	337
Average value	6.6	3369	33692	524




Figure 12: Ecosystem services of water supply in Padasavali-1 Microwatershed

The main farming constraints in Padasavali-1 micro-watershed to be found are less rainfall, non availability fertilizers, high crop pests & diseases, lack of transportations, lack of storage, damage of crops pests & diseases, non availability of plant protection chemicals. Majority of farmers depend up on money lender and bank of the sources of loan for purpose of crop production. Farmers to sell the agriculture produce

through village market and regulated the farmers getting the agriculture related information on newspaper. Farmers reported that they are not getting timely support/extension services from the concerned development department (Table 22).

Table 22: Farming constraints related land resources of sample households in Padasavali-1 Microwatershed

Sl. No	Particulars	Per cent
1	Less Rainfall	90.0
2	Non availability fertilizers	10.0
3	High crop pests & Diseases	80.0
4	Lack of transportations	40.0
5	Lack of storage	20.0
6	Damage of crops pests & Diseases	100.0
7	Non availability of plant protection chemicals	80.0
	Source of loan	
8	Money Leander	90.0
	Bank	10.0
	Market for selling	
9	Village market	90.0
	Regulated	10.0
	Sources of Agri-Technology information	
10	Newspaper	100

The findings of the study would be very much useful to the planners and policy makers of the study area to identify the irrationality in the existing production pattern and to suggest appropriate production plans for efficient utilization of their scarce resources resulting in increased net farm incomes and employment. The study also throws light on future potentialities of increasing net farm income and employment under different situations viz., with existing and recommended technology.