



हर कदम, हर डगर  
किसानों का हमसफर  
भारतीय कृषि अनुसंधान परिषद

*AgriSearch with a human touch*

**LAND RESOURCE INVENTORY AND SOCIO-ECONOMIC STATUS OF  
FARM HOUSEHOLDS FOR WATERSHED PLANNING AND  
DEVELOPMENT**

**RAGHUNATHANAHALLI-2 (4D4A2M3a) MICROWATERSHED**

Alavandi Hobli, Koppal Taluk & District, Karnataka

**Karnataka Watershed Development Project – II**

**SUJALA – III**

**World Bank funded Project**



**THE WORLD BANK**



**ICAR – NATIONAL BUREAU OF SOIL SURVEY AND LAND USE PLANNING**



ICAR - NBSS & LUP



**WATERSHED DEVELOPMENT DEPARTMENT  
GOVT. OF KARNATAKA, BANGALORE**



## **About ICAR - NBSS&LUP**

The ICAR-National Bureau of Soil Survey and Land Use Planning (ICAR-NBSS&LUP), Nagpur, a premier Institute of the Indian Council of Agricultural Research (ICAR), was set up during 1976 with the objective to prepare soil resource maps at national, state and district levels and to provide research inputs in soil resource mapping and its applications, land evaluation, land use planning, land resource management, and database management using GIS for optimising land use on different kinds of soils in the country.

The Bureau has been engaged in carrying out soil resource survey, agro-ecological and soil degradation mapping at the country, state and district levels for qualitative assessment and monitoring the soil health towards viable land use planning. The research activities have resulted in identifying the soil potentials and problems, and the various applications of the soil surveys with the ultimate objective of sustainable agricultural development. The Bureau has the mandate to correlate and classify soils of the country and maintain a National Register of all the established soil series. The Institute is also imparting in-service training to staff of the soil survey agencies in the area of soil survey, land evaluation and soil survey interpretations for land use planning. The Bureau in collaboration with Panjabrao Krishi Vidyapeeth, Akola is running post-graduate teaching and research programme in land resource management, leading to M.Sc. and Ph.D. degrees.

**Citation:** Rajendra Hegde, Ramesh Kumar, S.C., K.V. Niranjana, S. Srinivas, M.Lalitha, B.A. Dhanorkar, R.S. Reddy and S.K. Singh (2019). "Land resource inventory and socio-economic status of farm households for watershed planning and development of Raghunathanahalli-2 (4D4A2M3a) Microwatershed, Koppal Taluk and District, Karnataka", ICAR-NBSS&LUP Sujala MWS Publ.543, ICAR – NBSS & LUP, RC, Bangalore. p.137 &28.

### **TO OBTAIN COPIES,**

**Please write to:**

**Director, ICAR - NBSS & LUP,**

Amaravati Road, NAGPUR - 440 033, India

Phone : (0712) 2500386, 2500664, 2500545 (O)

Telefax : 0712-2522534

E-Mail : [director@nbsslup.ernet.in](mailto:director@nbsslup.ernet.in)

Website URL : [nbsslup.in](http://nbsslup.in)

Or

**Head, Regional Centre, ICAR - NBSS&LUP, Hebbal, Bangalore - 560 024**

Phone : (080) 23412242, 23510350 (O)

Telefax : 080-23510350

E-Mail : [nbssrcb@gmail.com](mailto:nbssrcb@gmail.com)



**LAND RESOURCE INVENTORY AND SOCIO-ECONOMIC  
STATUS OF FARM HOUSEHOLDS FOR WATERSHED  
PLANNING AND DEVELOPMENT  
RAGHUNATHANAHALLI-2 (4D4A2M3a)  
MICROWATERSHED**

**Alavandi Hobli, Koppal Taluk & District, Karnataka**

**Karnataka Watershed Development Project – II**

**Sujala-III**

**World Bank funded Project**



**THE WORLD BANK**



**ICAR – NATIONAL BUREAU OF SOIL SURVEY AND LAND USE  
PLANNING**



ICAR - NBSS & LUP



**WATERSHED DEVELOPMENT DEPARTMENT, GOVT. OF  
KARNATAKA, BANGALORE**







## PREFACE

In Karnataka, as in other Indian States, the livelihoods of rural people are intertwined with farming pursuits. The challenges in agriculture are seriously threatening the livelihood of a large number of farmers as they have been practicing farming in contextual factors beyond their control. Climatic factors are the most important ones and have become much more significant in recent times due to rapid climate changes induced by intensive anthropogenic activities affecting our ecosystem in multiple ways. Climate change has become the reality, it is happening and efforts to evolve and demonstrate climate resilient technologies have become essential. Due to the already over stressed scenario of agrarian sector, the climate change is resulting in manifold increase in the complexities, pushing the rural mass to face more and more unpredictable situations. The rising temperatures and unpredictable rainfall patterns are going to test seriously the informed decisions farmers have to make in order to survive in farming and sustain their livelihood.

It is generally recognized that impacts of climate change shall not be uniform across the globe. It is said that impact of climate change is more severe in South Asia. Based on the analysis of meteorological data, it is predicted that in India, there will be upward trend in mean temperature, downward trend in relative humidity, annual rainfall and number of wet days in a year. Also, in general, phenomena like erratic monsoon, spread of tropical diseases, rise in sea levels, changes in availability of fresh water, frequent floods, droughts, heat waves, storms and hurricanes are predicted. Each one of these adverse situations are already being experienced in various parts of India and also at the global level. Decline in agricultural productivity of small and marginal farmers becoming more vulnerable is already witnessed.

In Karnataka, more than 60 per cent of the population live in rural areas and depend on agriculture and allied activities for their livelihood. Though the state has achieved significant progress in increasing the yield of many crops, there is tremendous pressure on the land resources due to the growing and competing demands of various land uses. This is reflected in the alarming rate of land degradation observed. Already more than 50 per cent of the area is affected by various forms of degradation. If this trend continues, the sustainability of the fragile ecosystem will be badly affected. The adverse effects of change in the climatic factors are putting additional stress on the land resources and the farmers dependent on this.

The natural resources (land, water and vegetation) of the state need adequate and constant care and management, backed by site-specific technological interventions and investments particularly by the government. Detailed database pertaining to the nature of the land resources, their constraints, inherent potentials and suitability for various land

based rural enterprises, crops and other uses is a prerequisite for preparing location-specific action plans, which are in tune with the inherent capability of the resources. Any effort to evolve climate resilient technologies has to be based on the baseline scientific database. Then only one can expect effective implementation of climate resilient technologies, monitor the progress, make essential review of the strategy, and finally evaluate the effectiveness of the implemented programs. The information available at present on the land resources of the state are of general nature and useful only for general purpose planning. Since the need of the hour is to have site-specific information suitable for farm level planning and detailed characterization and delineation of the existing land resources of an area into similar management units is the only option.

ICAR-NBSS&LUP, Regional Centre, Bangalore has taken up a project sponsored by the Karnataka Watershed Development Project-II, (Sujala-III), Government of Karnataka funded by the World Bank under Component -1 Land Resource Inventory. This study was taken up to demonstrate the utility of such a database in reviewing, monitoring and evaluating all the land based watershed development programs on a scientific footing. To meet the requirements of various land use planners at grassroots level, the present study on “Land Resource Inventory and Socio-Economic Status of Farm Households for Watershed Planning and Development of for Raghunathanahalli-2 microwatershed in Koppal Taluk, Koppal District, Karnataka” for integrated development was taken up in collaboration with the State Agricultural Universities, IISC, KSRSAC, KSNDMC as Consortia partners. The project provides detailed land resource information at cadastral level (1:7920 scale) for all the plots and socio-economic status of farm households covering thirty per cent farmers randomly selected representing landed and landless class of farmers in the micro-watershed. The project report with the accompanying maps for the microwatershed will provide required detailed database for evolving effective land use plan, alternative land use options and conservation plans for the planners, administrators, agricultural extension personnel, KVK officials, developmental departments and other land users to manage the land resources in a sustainable manner.

It is hoped that this database will be useful to the planners, administrators and developmental agencies working in the area in not only for formulating location specific developmental schemes but also for their effective monitoring at the village/watershed level.

Nagpur

Date:10-11-2019

**S.K. SINGH**

Director, ICAR - NBSS&LUP,Nagpur

## Contributors

<b>Dr. Rajendra Hegde</b> Principal Scientist, Head & Project Leader, Sujala-III Project ICAR-NBSS&LUP, Regional Centre, Bangalore	<b>Dr. S.K.Singh</b> Director, ICAR-NBSS&LUP Coordinator, Sujala-III Project Nagpur
<b>Soil Survey, Mapping &amp; Report Preparation</b>	
Dr. K.V. Niranjana	Sh. R.S. Reddy
Dr. B.A. Dhanorkar	Smt. Chaitra, S.P.
	Dr. Gopali Bardhan
	Sh. Somashekar T.N
	Ms. Arpitha G.M
	Dr. Mahendra kumar M.B
	Sh. Tirupati Meti
<b>Field Work</b>	
Sh. C. Bache Gowda	Sh. Mayur Patil
Sh. Somashekar	Sh. Arun Kumar, S.
Sh. M. Jayaramaiah	Sh. Sunil Raj
	Sh. Yogesh Kumar, B.
	Sh. Vikas, N.K.
	Sh. Arun Kumar, S.G.
	Sh. Umesh Jادیappa Madolli
	Sh. Praveen Kumar P. Achalkar
	Sh. Veerabhadraswamy
	Sh. Vinay
	Sh. Shankarappa, K.
	Sh. Lankesh, R.S.
	Sh. Appanna B. Hattigoudar
	Sh. Maharudra
<b>GIS Work</b>	
Dr. S.Srinivas	Sh. A.G.Devendra Prasad
Sh. D.H.Venkatesh	Sh. Abhijith Sastry, N.S.
Smt. K.Sujatha	Smt. Shyla, B.
Smt. K.V.Archana	Smt. Swetha ,K.
Sh. N.Maddileti	Ms. Vidya, P.C.
	Sh. Deepak, M.J.
	Smt. K.Karunya Lakshmi
	Ms. Seema, K.V.

<b>Laboratory Analysis</b>	
Dr. M. Lalitha	Sh. Vindhya, N.G.
Smt. Arti Koyal	Ms. P. Pavanakumari, P.
Smt. Parvathy, S.	Ms. Rashmi, N.
	Ms. Leelavathy, K.U.
	Smt. Usha Kiran, G.
<b>Socio-Economic Analysis</b>	
<b>Dr. S.C. Ramesh Kumar</b>	Sh. M.K. Prakashanaik
	Ms. Karuna V. Kulkarni
	Mrs. Sowmya A.N
	Sh. Vinod R
	Sh. Basavaraja
	Sh. Vijay Kumar Lamani
	Ms. Sowmya K.B
	Mrs. Prathibha, D.G
	Sh. Rajendra,D
<b>Soil &amp; Water Conservation</b>	
Sh. Sunil P. Maske	
<b>Watershed Development Department, GoK, Bangalore</b>	
Sh. Rajeev Ranjan IFS Project Director & Commissioner, WDD	Dr. A. Natarajan NRM Consultant, Sujala-III Project
Dr. S.D. Pathak IFS Executive Director & Chief Conservator of Forests, WDD	

# **PART-A**

## **LAND RESOURCE INVENTORY**



## Contents

Preface		
Contributors		
Executive Summary		
Chapter 1	Introduction	1
Chapter 2	Geographical Setting	3
2.1	Location and Extent	3
2.2	Geology	3
2.3	Physiography	4
2.4	Drainage	5
2.5	Climate	5
2.6	Natural Vegetation	6
2.7	Land Utilization	7
Chapter 3	Survey Methodology	11
3.1	Base maps	11
3.2	Image Interpretation for Physiography	11
3.3	Field Investigation	14
3.4	Soil mapping	16
3.5	Laboratory Characterization	16
3.6	Land Management Units	17
Chapter 4	The Soils	21
4.1	Soils of Granite Gneiss Landscape	21
4.2	Soils of Alluvial Landscape	23
Chapter 5	Interpretation for Land Resource Management	37
5.1	Land Capability Classification	37
5.2	Soil Depth	39
5.3	Surface Soil Texture	40
5.4	Soil Gravelliness	41
5.5	Available Water Capacity	42
5.6	Soil Slope	43
5.7	Soil Erosion	44
Chapter 6	Fertility Status	47
6.1	Soil Reaction (pH)	47
6.2	Electrical Conductivity (EC)	47
6.3	Organic Carbon (OC)	47
6.4	Available Phosphorus	47
6.5	Available Potassium	48
6.6	Available Sulphur	48
6.7	Available Boron	51
6.8	Available Iron	51
6.9	Available Manganese	51
6.10	Available Copper	51
6.11	Available Zinc	51
Chapter 7	Land Suitability for Major Crops	55
7.1	Land Suitability for Sorghum	55
7.2	Land Suitability for Maize	56
7.3	Land Suitability for Bajra	57

7.4	Land Suitability for Groundnut	58
7.5	Land Suitability for Sunflower	59
7.6	Land Suitability for Cotton	60
7.7	Land Suitability for Red gram	61
7.8	Land Suitability for Bengal gram	62
7.9	Land Suitability for Chilli	63
7.10	Land Suitability for Tomato	64
7.11	Land Suitability for Brinjal	65
7.12	Land Suitability for Onion	66
7.13	Land Suitability for Bhendi	67
7.14	Land Suitability for Drumstick	68
7.15	Land Suitability for Mulberry	69
7.16	Land Suitability for Mango	70
7.17	Land Suitability for Sapota	71
7.18	Land Suitability for Pomegranate	72
7.19	Land Suitability for Guava	73
7.20	Land Suitability for Jackfruit	74
7.21	Land Suitability for Jamun	75
7.22	Land Suitability for Musambi	76
7.23	Land Suitability for Lime	77
7.24	Land Suitability for Cashew	78
7.25	Land Suitability for Custard apple	79
7.26	Land Suitability for Amla	80
7.27	Land Suitability for Tamarind	81
7.28	Land Suitability for Marigold	82
7.29	Land Suitability for Chrysanthemum	83
7.30	Land Suitability for Jasmine	84
7.31	Land Suitability for Crossandra	85
7.32	Land Management Units (LMUs)	86
7.33	Proposed Crop Plan for Raghunathanahalli-2 Microwatershed	87
Chapter 8	Soil Health Management	123
8.1	Soil health	123
Chapter 9	Soil and Water conservation Treatment Plan	129
9.1	Treatment Plan	129
9.2	Recommended Soil and Water Conservation measures	133
9.3	Greening of microwatershed	134
	References	137
	Appendix I	I-VI
	Appendix II	VII-XII
	Appendix III	XIII-XVII



## LIST OF TABLES

2.1	Mean Monthly Rainfall, PET, ½ PET at Koppal Taluk and District	5
2.2	Land Utilization in Koppal District	7
3.1	Differentiating Characteristics used for Identifying Soil Series	15
3.2	Soil map unit description of Raghunathanahalli-2 microwatershed	17
4.1	Physical and chemical characteristics of soil series identified in Raghunathanahalli-2 microwatershed	29
7.1	Soil-Site Characteristics of Raghunathanahalli-2 microwatershed	87
7.2	Land suitability criteria for Sorghum	88
7.3	Land suitability criteria for Maize	89
7.4	Land suitability criteria for Bajra	90
7.5	Land suitability criteria for Groundnut	91
7.6	Land suitability criteria for Sunflower	92
7.7	Land suitability criteria for Cotton	93
7.8	Land suitability criteria for Red gram	94
7.9	Land suitability criteria for Bengal gram	95
7.10	Land suitability criteria for Chilli	96
7.11	Land suitability criteria for Tomato	97
7.12	Land suitability criteria for Brinjal	98
7.13	Land suitability criteria for Onion	99
7.14	Land suitability criteria for Bhendi	100
7.15	Land suitability criteria for Drumstick	101
7.16	Land suitability criteria for Mulberry	102
7.17	Land suitability criteria for Mango	103
7.18	Land suitability criteria for Sapota	104
7.19	Land suitability criteria for Pomegranate	105
7.20	Land suitability criteria for Guava	106
7.21	Land suitability criteria for Jackfruit	107
7.22	Land suitability criteria for Jamun	108
7.23	Land suitability criteria for Musambi	109

7.24	Land suitability criteria for Lime	110
7.25	Land suitability criteria for Cashew	111
7.26	Land suitability criteria for Custard apple	112
7.27	Land suitability criteria for Amla	113
7.28	Land suitability criteria for Tamarind	114
7.29	Land suitability criteria for Marigold	115
7.30	Land suitability criteria for Chrysanthemum	116
7.31	Land suitability criteria for Jasmine	117
7.32	Land suitability criteria for Crossandra	118
7.33	Proposed Crop Plan for Raghunathanahalli-2 Microwatershed	1213

## LIST OF FIGURES

2.1	Location map of Raghunathanahalli-2 Microwatershed	3
2.2a	Granite and granite gneiss rocks	4
2.2b	Alluvium	4
2.3	Rainfall distribution in Koppal Taluk and District	6
2.4	Natural vegetation of Raghunathanahalli-2 Microwatershed	6
2.5	Different crops and cropping systems in Raghunathanahalli-2 Microwatershed	8
2.6	Current Land use – Raghunathanahalli-2 Microwatershed	9
2.7	Location of Wells -Raghunathanahalli-2 Microwatershed	9
3.1	Scanned and Digitized Cadastral map of Raghunathanahalli-2 Microwatershed	13
3.2	Satellite image of Raghunathanahalli-2 Microwatershed	13
3.3	Cadastral map overlaid on IRS PAN+LISS IV merged imagery of Raghunathanahalli-2 Microwatershed	14
3.4	Location of profiles in a transect	15
3.5	Soil phase or management units-Raghunathanahalli-2 Microwatershed	19
5.1	Land Capability Classification of Raghunathanahalli-2 Microwatershed	39
5.2	Soil Depth map of Raghunathanahalli-2 Microwatershed	40
5.3	Surface Soil Texture map of Raghunathanahalli-2 Microwatershed	41
5.4	Soil Gravelliness map of Raghunathanahalli-2 Microwatershed	42
5.5	Soil Available Water Capacity map of Raghunathanahalli-2 Microwatershed	43
5.6	Soil Slope map of Raghunathanahalli-2 Microwatershed	44
5.7	Soil Erosion map of Raghunathanahalli-2 Microwatershed	45
6.1	Soil Reaction (pH) map of Raghunathanahalli-2 Microwatershed	48
6.2	Electrical Conductivity (EC) map of Raghunathanahalli-2 Microwatershed	49
6.3	Soil Organic Carbon (OC) map of Raghunathanahalli-2 Microwatershed	49
6.4	Soil Available Phosphorus map of Raghunathanahalli-2 Microwatershed	50
6.5	Soil Available Potassium map of Raghunathanahalli-2 Microwatershed	50
6.6	Soil Available Sulphur map of Raghunathanahalli-2 Microwatershed	51
6.7	Soil Available Boron map of Raghunathanahalli-2 Microwatershed	52
6.8	Soil Available Iron map of Raghunathanahalli-2 Microwatershed	52
6.9	Soil Available Manganese map of Raghunathanahalli-2 Microwatershed	53
6.10	Soil Available Copper map of Raghunathanahalli-2 Microwatershed	53

6.11	Soil Available Zinc map of Raghunathanahalli-2 Microwatershed	54
7.1	Land suitability map of Sorghum	56
7.2	Land suitability map of Maize	57
7.3	Land suitability map of Bajra	58
7.4	Land suitability map of Groundnut	59
7.5	Land suitability map of Sunflower	60
7.6	Land suitability map of Cotton	61
7.7	Land suitability map of Redgram	62
7.8	Land suitability map of Bengal gram	63
7.9	Land suitability map of Chilli	64
7.10	Land suitability map of Tomato	65
7.11	Land suitability map of Brinjal	66
7.12	Land suitability map of Onion	67
7.13	Land suitability map of Bhendi	68
7.14	Land suitability map of Drumstick	69
7.15	Land suitability map of Mulberry	70
7.16	Land suitability map of Mango	71
7.17	Land suitability map of Sapota	72
7.18	Land suitability map of Pomegranate	73
7.19	Land suitability map of Guava	74
7.20	Land suitability map of Jackfruit	75
7.21	Land suitability map of Jamun	76
7.22	Land suitability map of Musambi	77
7.23	Land suitability map of Lime	78
7.24	Land suitability map of Cashew	79
7.25	Land suitability map of Custard apple	80
7.26	Land suitability map of Amla	81
7.27	Land suitability map of Tamarind	82
7.28	Land suitability map of Marigold	83
7.29	Land suitability map of Chrysanthemum	84
7.30	Land suitability map of Jasmine	85
7.31	Land suitability map of Crossandra	86
7.32	Land Management Unit map of Raghunathanahalli-2 microwatershed	120
9.1	Soil and water conservation Plan map of Raghunathanahalli-2 Microwatershed	134

## **EXECUTIVE SUMMARY**

*The land resource inventory of Raghunathanahalli-2 microwatershed was conducted using village cadastral maps and IRS satellite imagery on 1:7920 scale. The false colour composites of IRS imagery were interpreted for physiography and these physiographic delineations were used as base for mapping soils. The soils were studied in several transects and a soil map was prepared with phases of soil series as mapping units. Random checks were made all over the area outside the transects to confirm and validate the soil map unit boundaries. The soil map shows the geographic distribution and extent, characteristics, classification, behavior and use potentials of the soils in the microwatershed.*

*The present study covers an area of 441 ha in Koppal taluk and district, Karnataka. The climate is semiarid and categorized as drought - prone with an average annual rainfall of 662 mm, of which about 424 mm is received during south-west monsoon, 161 mm during north-east and the remaining 77 mm during the rest of the year.*

*An area of 99 per cent is covered by soils and 1 per cent is by habitation and settlements. The salient findings from the land resource inventory are summarized briefly below.*

- ❖ The soils belong to 10 soil series and 17 soil phases (management units) and 6 Land Management Units.*
- ❖ The length of crop growing period is <90 days and starts from 2<sup>nd</sup> week of August to 2<sup>nd</sup> week of November.*
- ❖ From the master soil map, several interpretative and thematic maps like land capability, soil depth, surface soil texture, soil gravelliness, available water capacity, soil slope and soil erosion were generated.*
- ❖ Soil fertility status maps for macro and micronutrients were generated based on the surface soil samples collected at every 320 m grid interval.*
- ❖ Land suitability for growing 31 major agricultural and horticultural crops were assessed and maps showing the degree of suitability along with constraints were generated.*
- ❖ An area of about 99 per cent is suitable for agriculture.*
- ❖ An area of about 27 per cent of the soils are very shallow to shallow (<25-50 cm), 64 per cent of the soils are moderately shallow to moderately deep (50-100 cm) and 8 per cent soils are deep to very deep (100->150 cm).*
- ❖ Entire cultivated area of the microwatershed falls under clayey soils at the surface.*
- ❖ An area of about 29 per cent area has non-gravelly (<15% gravel) soils and 70 per cent has gravelly to very gravelly (15-60% gravel) soils.*
- ❖ An area of about 74 per cent area is very low to low (<50-100 mm/m), 17 per cent area is medium (101-150 mm/m) and 8 per cent area is very high (>200 mm/m) in available water capacity.*

- ❖ *An area of about 11 per cent area of the microwatershed has nearly level (0-1% slope) lands and 87 per cent area of the microwatershed has very gently sloping (1-3% slope) lands.*
- ❖ *An area of about 41 per cent area is moderately (e2) eroded and about 58 per cent area is slightly (e1) eroded.*
- ❖ *Entire cultivated area of the microwatershed soils are moderately alkaline to strongly alkaline (pH 7.8-9.0) in soil reaction.*
- ❖ *The Electrical Conductivity (EC) of the soils in the entire cultivated area of the microwatershed is  $<2 \text{ dsm}^{-1}$  indicating that the soils are non-saline.*
- ❖ *Organic carbon is medium (0.5-0.75%) in 77 per cent area and low ( $<0.5\%$ ) in 22 per cent area of the microwatershed.*
- ❖ *An area of about 4 per cent is medium (23-57 kg/ha) and 95 per cent is low ( $<23 \text{ kg/ha}$ ) in available phosphorus.*
- ❖ *Entire cultivated area of the microwatershed is high ( $>337 \text{ kg/ha}$ ) in available potassium.*
- ❖ *Available sulphur is medium (10-20 ppm) in 30 per cent area and high ( $>20 \text{ ppm}$ ) in 69 per cent area of the microwatershed.*
- ❖ *An area of about 80 per cent is low ( $<0.5 \text{ ppm}$ ) and 19 per cent is medium (0.5-1.0 ppm) in available boron content.*
- ❖ *An area of about 67 per cent is sufficient ( $>4.5 \text{ ppm}$ ) and 32 per cent is deficient ( $<4.5 \text{ ppm}$ ) in available iron content.*
- ❖ *Entire cultivated area of the microwatershed is sufficient ( $>1.0 \text{ ppm}$ ) in available manganese content.*
- ❖ *Entire cultivated area of the microwatershed is sufficient ( $>0.2 \text{ ppm}$ ) in available copper content.*
- ❖ *Entire cultivated area of the microwatershed is deficient ( $<0.6 \text{ ppm}$ ) in available zinc content.*
- ❖ *The land suitability for 31 major crops grown in the microwatershed was assessed and the areas that are highly suitable (S1) and moderately suitable (S2) are given below. It is however to be noted that a given soil may be suitable for various crops but what specific crop to be grown may be decided by the farmer looking to his capacity to invest on various inputs, marketing infrastructure, market price and finally the demand and supply position.*

**Land suitability for various crops in the microwatershed**

Crop	Suitability Area in ha (%)		Crop	Suitability Area in ha (%)	
	Highly suitable (S1)	Moderately suitable (S2)		Highly suitable (S1)	Moderately suitable (S2)
Sorghum	21(5)	296(67)	Sapota	-	152(35)
Maize	-	300(68)	Pomegranate	-	260(59)
Bajra	-	317(72)	Guava	-	152(35)
Groundnut	-	152(35)	Jackfruit	-	152(35)
Sunflower	14(3)	246(56)	Jamun	-	34(8)
Cotton	21(5)	297(67)	Musambi	14(3)	246(56)
Red gram	-	210(48)	Lime	14(3)	246(56)
Bengalgram	21(5)	296(67)	Cashew	-	-
Chilli	-	169(38)	Custard apple	21(5)	296(67)
Tomato	-	165(37)	Amla	-	316(72)
Brinjal	-	316(72)	Tamarind	-	34(8)
Onion	-	165(38)	Marigold	-	316(72)
Bhendi	-	316(72)	Chrysanthemum	-	316(72)
Drumstick	-	260(59)	Jasmine	-	208(47)
Mulberry	-	240(55)	Crossandra	-	215(49)
Mango	-	-			

Apart from the individual crop suitability, a proposed crop plan has been prepared for the 6 identified LMUs by considering only the highly and moderately suitable lands for different crops and cropping systems with food, fodder, fibre and other horticulture crops.

- ❖ Maintaining soil-health is vital for crop production and conserve soil and land resource base for maintaining ecological balance and to mitigate climate change. For this, several ameliorative measures have been suggested for these problematic soils like saline/alkali, highly eroded, sandy soils etc.,
- ❖ Soil and water conservation treatment plan has been prepared that would help in identifying the sites to be treated and also the type of structures required.
- ❖ As part of the greening programme, several tree species have been suggested to be planted in marginal and submarginal lands, field bunds and also in the hillocks, mounds and ridges. That would help in supplementing the farm income, provide fodder and fuel, and generate lot of biomass which in turn would help in maintaining the ecological balance and contribute to mitigating the climate change.





## **INTRODUCTION**

Soil is a finite natural resource that is central to sustainable agriculture and food security. Over the years, this precious resource is faced with the problems of erosion, salinity, alkalinity, degradation, depletion of nutrients and even decline in availability of land for agriculture. It is a known fact, that it takes thousands of years to form a few centimetres of soil, thus, soil is a precious gift of nature. The area available for agriculture is about 51 per cent of the total geographical area and more than 60 per cent of the people are still dependant on agriculture for their livelihood. However, the capacity of a soil to produce is limited and the limits to the production are set by its intrinsic characteristics, agro-climatic setting, and use and management. There is, therefore, tremendous pressure on land and water resources, which is causing decline in soil-health and stagnation in productivity. As much as 121 m ha of land is reportedly degraded which leads to impaired soil quality. It is imperative that steps are urgently taken to check and reverse land degradation without any further loss of time. The improvements in productivity will have to come from sustainable intensification measures that make the most effective use of land and water resources. Soil erosion alone has degraded about 35 lakh ha. Almost all the uncultivated areas are facing various degrees of degradation, particularly soil erosion; salinity and alkalinity has emerged as a major problem in more than 3.5 lakh ha in the irrigated areas of the state. Nutrient depletion and declining factor productivity is common in both rainfed and irrigated areas. The degradation is continuing at an alarming rate and there appears to be no systematic effort among the stakeholders to contain this process. In recent times, an aberration of weather due to climate change phenomenon has added another dimension leading to unpredictable situations to be tackled by the farmers. In this critical juncture, the challenge before us is not only to increase the productivity per unit area which is steadily declining and showing a fatigue syndrome, but also to prevent or at least reduce the severity of degradation. If the situation is not reversed at the earliest, then the sustainability of the already fragile crop production system and the overall ecosystem will be badly affected in the state.

Added to this, every year there is a significant diversion of farm lands and water resources for non-agricultural purposes. Thus, developing strategies to slow down the degradation process or reclaim the soils to normal condition and ensure sustainability of production system are the major issues today. This demands a systematic appraisal of our soil and land resources with respect to their extent, geographic distribution, characteristics, behaviour and uses potential, which is very important for developing an effective land use and cropping systems for augmenting agricultural production on a sustainable basis. The soil and land resource inventories made so far in Karnataka had limited utility because the surveys were of different types, scales and intensities carried out at different times with specific objectives. Hence, there is an urgent need to generate

detailed site-specific farm level database on various land resources for all the villages/watersheds in a time bound manner that would help to protect the valuable soil and land resources and also to stabilize the farm production.

Therefore, the land resource inventory required for farm level planning is the one which investigates all the parameters which are critical for productivity *viz.*, soils, site characteristics like slope, erosion, gravelliness and stoniness, climate, water, topography, geology, hydrology, vegetation, crops, land use pattern, animal population, socio-economic conditions, infrastructure, marketing facilities and various schemes and developmental works of the government etc. From the data collected at farm level, the specific problems and potentials of the area can be identified and highlighted, conservation measures required for the area can be planned on a scientific footing, suitability of the area for various uses can be worked out and finally viable and sustainable land use options suitable for each and every land holding can be prescribed.

The Land Resource Inventory is basically done for identifying potential and problem areas, developing sustainable land use plans, estimation of surface run off and water harvesting potential, preparation of soil and water conservation plans, land degradation/desertification etc. The Bureau is presently engaged in developing an LRI methodology using high resolution satellite remote sensing data and Digital Elevation Model (DEM) data to prepare Landscape Ecological Units (LEU) map representing agro-ecosystem as a whole. The LEU is preferred over landform as the base map for LRI. LEU is the assemblage of landform, slope and land use. An attempt was made to upscale the soil resource information from 1:250000 and 1:50000 scale to the LEU map in Goa and other states.

The land resource inventory aims to provide site-specific database for Raghunathanahalli-2 microwatershed in Koppal Taluk and District, Karnataka State for the Karnataka Watershed Development Department. The database was generated by using cadastral map of the village as a base along with high resolution IRS LISS IV and Cartosat-1 merged satellite imagery. Later, an attempt will be made to uplink this LRI data generated at 1:7920 scales under Sujala-III Project to the proposed Landscape Ecological Units (LEUs) map.

The study was organized and executed by the ICAR- National Bureau of Soil Survey and Land Use Planning, Regional Centre, Bangalore under Generation of Land Resource Inventory Data Base Component-1 of the Sujala-III Project funded by the World Bank.

## GEOGRAPHICAL SETTING

### 2.1 Location and Extent

The Raghunathanahalli-2 Microwatershed is located in the central part of northern Karnataka in Koppal Taluk, Koppal District, Karnataka State (Fig.2.1). It comprises parts of Belagatti, Gattareddyhala, Hatti & Raghunathahalli villages. It lies between  $15^{\circ}13'$  –  $15^{\circ}14'$  North latitudes and  $75^{\circ}54'$  –  $75^{\circ}56'$  East longitudes and covers an area of 441 ha. It is about 32 km from Koppal town and is surrounded by Gattareddyhala village on the north and east, Belagatti village on the south, Raghunathahalli village on the east and Hatti village on the southeastern side of the microwatershed.




Fig.2.1 Location map of Raghunathanahalli-2 Microwatershed

### 2.2 Geology

Major rock formation observed in the microwatershed are granite gneiss and alluvium (Figs.2.2 a & b). Granite gneisses are essentially pink to gray and are coarse to

medium grained. They consist primarily of quartz, feldspar, biotite and hornblende. The gray granite gneisses are highly weathered, fractured and fissured upto a depth of about 10 m. Dolerite dykes and quartz veins are common with variable width and found to occur in Bettageri village. The soil thickness of the alluvium generally is limited to less than a meter, except in river valleys where it is very deep extending to tens of meters. Such soils are transported and represent palaeo black soils originally formed at higher elevation, but now occupying river valleys.



Fig.2.2a Granite and granite gneiss rocks



Fig.2.2b Alluvium

### 2.3 Physiography

Physiographically, the area has been identified as Granite gneiss and Alluvial landscapes based on geology. The microwatershed area has been further divided into mounds/ridges, summits, side slopes and very gently sloping uplands and nearly level plains based on slope and its relief features. The elevation ranges from 529-553 m in the gently sloping uplands. The mounds and ridges are mostly covered by rock outcrops.

## 2.4 Drainage

The area is drained by several small seasonal streams that join Hire *halla* and Chenna *halla* along its course. Though, the streams are not perennial, during rainy season they carry large quantities of rain water. The microwatershed has only few small tanks which are not able to store the water flowing during the rainy season. Due to this, the ground water recharge is very much affected in the villages. This is reflected in the failure of many bore wells in the villages. If the available rain water is properly harnessed by constructing tanks and recharge structures at appropriate places in the villages, then the drinking and irrigation needs of the area can be easily met. The drainage network is dendritic to sub parallel.

## 2.5 Climate

The district falls under semiarid tract of the state and is categorized as drought - prone with total annual rainfall of 662 mm (Table 2.1) Of this, a maximum of 424 mm precipitation takes place during south–west monsoon period from June to September, north-east monsoon contributes about 161 mm and prevails from October to early December and the remaining 77 mm received during the rest of the year. The winter season is from December to February. During April and May, the temperatures reach up to 45°C and in December and January, the temperatures will go down to 16°C. Rainfall distribution is shown in Figure 2.3. The average Potential Evapo Transpiration (PET) is 145 mm and varies from a low of 101 mm in December and 193 mm in the months of May. The PET is always higher than precipitation in all the months except in the month of September. Generally, the Length of crop Growing Period (LGP) is <90 days and starts from 2nd week of August to 2nd week of November.

**Table 2.1 Mean Monthly Rainfall, PET, 1/2 PET at Koppal Taluk and District**

Sl. No.	Months	Rainfall	PET	1/2 PET
1	January	1.60	116.70	58.35
2	February	1.50	129.20	64.60
3	March	14.10	169.80	84.90
4	April	18.10	180.60	90.30
5	May	41.60	193.50	96.75
6	June	85.80	167.90	83.95
7	July	72.10	156.20	78.10
8	August	110.50	152.50	76.25
9	September	155.60	138.50	69.25
10	October	116.30	122.30	61.15
11	November	36.00	106.40	53.20
12	December	9.10	101.00	50.50
<b>TOTAL</b>		<b>662.30</b>	<b>144.55</b>	






Fig. 2.3 Rainfall distribution in Koppal Taluk and District

### 2.6 Natural Vegetation

The natural vegetation is sparse comprising few tree species, shrubs and herbs. The mounds, ridges and boulders occupy sizeable areas which are under thin to moderately thick forest vegetation. Still, there are some remnants of the past forest cover which can be seen in patches in some ridges and hillocks in the microwatershed (Fig 2.4).

Apart from the continuing deforestation, the presence of large population of goats, sheep and other cattle in the microwatershed is causing vegetative degradation of whatever little vegetation left in the area. The uncontrolled grazing has left no time for the regeneration of the vegetative cover. This leads to the accelerated rate of erosion on the hill slopes, resulting in the formation of deep gullies in the foot slopes and eventually resulting in the heavy siltation of few tanks and reservoirs in the microwatershed.



Fig 2.4 Natural vegetation of Raghunathanahalli-2 microwatershed

## 2.7 Land Utilization

About 91 per cent area (Table 2.2) in Koppal district is cultivated at present and about 17 per cent of the area is sown more than once. An area of about 3 per cent is currently barren. Forests occupy a small area of about 5 per cent and the tree cover is in a very poor state. Most of the mounds, ridges and bouldery areas have very poor vegetative cover. Major crops grown in the area are sorghum, maize, bajra, cotton, safflower, sunflower, red gram, horse gram, onion, mulberry, pomegranate, sugarcane, Bengalgram, marigold and groundnut (Fig 2.5). While carrying out land resource inventory, the land use/land cover particulars are collected from all the survey numbers and a current land use map of the microwatershed is prepared. The current land use map prepared shows the arable and non-arable lands, other land uses and different types of crops grown in the area. The current land use map of Raghunathanahalli-2 Microwatershed is presented in Fig.2.6. Simultaneously, enumeration of existing wells (bore wells and open wells) and other soil and water conservation structures in the microwatershed is made and their location in different survey numbers is marked on the cadastral map. Map showing the location of wells in Raghunathanahalli-2 Microwatershed is given Fig.2.7.

**Table 2.2 Land Utilization in Koppal District**

Sl. No.	Agricultural land use	Area ( ha)	Per cent
1	Total geographical area	552495	
2	Total cultivated area	500542	90.6
3	Area sown more than once	92696	16.8
4	Trees and groves	210	0.04
5	Cropping intensity	-	118
6	Forest	29451	5.33
7	Cultivable wasteland	2568	0.46
8	Permanent Pasture land	14675	2.66
9	Barren land	16627	3.01
10	Non agricultural land	40591	7.35
11	Current fallow	19660	3.56





Groundnut



Sunflower



Cotton



Red gram



Onion



Marigold

Fig.2.5 Different crops and cropping systems in Raghunathanahalli-2 Microwatershed






Fig.2.6 Current Land Use – Raghunathanahalli-2 Microwatershed




Fig.2.7 Location of wells - Raghunathanahalli-2 Microwatershed



## SURVEY METHODOLOGY

The purpose of land resource inventory is to delineate similar areas (soil series and phases), which respond or expected to respond similarly for a given level of management. This was achieved in Raghunathanahalli-2 Microwatershed by the detailed study of all the soil characteristics (depth, texture, colour, structure, consistence, coarse fragments, porosity, soil reaction, soil horizons etc.) and site characteristics (slope, erosion, drainage, occurrence of rock fragments etc.) followed by grouping of similar areas based on soil-site characteristics into homogeneous (management units) units and showing their extent and geographic distribution on the microwatershed cadastral map. The detailed soil survey at 1:7920 scale was carried out in 441 ha area. The methodology followed for carrying out land resource inventory was as per the guidelines given in Soil Survey Manual (IARI, 1971; Soil Survey Staff, 2006; Natarajan *et al.*, 2015) which is briefly described below.

### 3.1 Base Maps

The detailed survey of the land resources occurring in the microwatershed was carried out by using digitized cadastral map and satellite imagery as base supplied by the KRSRSAC. The cadastral map shows field boundaries with their survey numbers, location of tanks, streams and other permanent features of the area (Fig. 3.1). Apart from the cadastral map, remote sensing data products from Cartosat-1 and LISS IV merged at the scale of 1:7920 were used in conjunction with the cadastral map to identify the geology, landscapes, landforms and other surface features. The imagery helped in the identification and delineation of boundaries between hills, uplands and lowlands, water bodies, forest and vegetated areas, roads, habitations and other cultural features of the area (Fig.3.2).The cadastral map was overlaid on the satellite imagery (Fig.3.3) that helps to identify the parcel boundaries and other permanent features. Apart from cadastral maps and images, toposheets of the area (1:50,000 scale) were used for initial traversing, identification of geology, landscapes and landforms, drainage features, present land use and also for selection of transects in the microwatershed.

### 3.2 Image Interpretation for Physiography

False Colour Composites (FCC) of Cartosat-I and LISS-IV merged satellite data covering the microwatershed area was visually interpreted using image interpretation elements and all the available collateral data with local knowledge. The delineated physiographic boundaries were transferred on to a cadastral map overlaid on satellite imagery. Physiographically, the area has been identified as granite gneiss and alluvial landscapes and is divided into landforms such as uplands, summits and very gently sloping based on slope. They were further subdivided into physiographic/ image

interpretation units based on image characteristics. The image interpretation legend for physiography is given below.

### **Image Interpretation Legend for Physiography**

#### **G- Granite gneiss landscape**

- G1 Hills/ Ridges/ Mounds
  - G11 Summits
  - G12 Side slopes
    - G121 Side slopes with dark grey tones
- G2 Uplands
  - G21 Summits
  - G22 Gently sloping uplands
    - G221 Gently sloping uplands, yellowish green (eroded)
    - G222 Gently sloping uplands, yellowish white (severely eroded)
  - G23 Very gently sloping uplands
    - G231 Very gently sloping uplands, yellowish green
    - G232 Very gently sloping uplands, medium green and pink
    - G233 Very gently sloping uplands, pink and green (scrub land)
    - G234 Very gently sloping uplands, medium greenish grey
    - G235 Very gently sloping uplands, yellowish white (eroded)
    - G236 Very gently sloping uplands, dark green
    - G237 Very gently sloping uplands, medium pink (coconut garden)
    - G238 Very gently sloping uplands, pink and bluish white (eroded)

#### **DSe Alluvial landscape**

##### **DSe 1 Summit**

- DSe 11 Nearly level Summit with dark grey tone
- DSe 12 Nearly level Summit with medium grey tone
- DSe 13 Nearly level Summit with whitish grey tone
- DSe 14 Nearly level Summit with whitish tone (Calcareousness)
- DSe 15 Nearly level Summit with pinkish grey tone
- DSe 16 Nearly level Summit with medium pink tone
- DSe 17 Nearly level Summit with bluish white tone
- DSe 18 Nearly level Summit with greenish grey tone

##### **DSe 2 Very gently sloping**

- DSe 21 Very gently sloping, whitish tone
- DSe 22 Very gently sloping, greyish pink tone
- DSe 23 Very gently sloping, whitish grey tone
- DSe 24 Very gently sloping, medium grey tone
- DSe 25 Very gently sloping, medium pink tone
- DSe 26 Very gently sloping, dark grey tone
- DSe 27 Very gently sloping, bluish grey tone
- DSe 28 Very gently sloping, greenish grey tone
- DSe 29 Very gently sloping, Pinkish grey




Fig 3.1 Scanned and Digitized Cadastral map of Raghunathanahalli-2 Microwatershed




Fig.3.2 Satellite Image of Raghunathanahalli-2 Microwatershed




Fig.3.3 Cadastral map overlaid on IRS PAN+LISS IV merged imagery of Raghunathanahalli-2 Microwatershed

### 3.3 Field Investigation

The field boundaries and survey numbers given on the cadastral sheet were located on the ground by following permanent features like roads, cart tracks, *nallas*, streams, tanks etc., and wherever changes were noticed, they were incorporated on the microwatershed cadastral map. Preliminary traverse of the microwatershed was carried out with the help of cadastral map, imagery and toposheets. While traversing, landforms and physiographic units identified were checked and preliminary soil legend was prepared by studying soils at few selected places. Then, intensive traversing of each physiographic unit like uplands and plains was carried out. Based on the variability observed on the surface, transects (Fig 3.4) were selected across the slope covering all the landform units in the microwatershed (Natarajan and Dipak Sarkar, 2010).






Fig: 3.4. Location of profiles in a transect

In the selected transect, soil profiles (Fig.3.4) were located at closely spaced intervals to take care of any change in the land features like break in slope, erosion, gravel, stones etc. In the selected sites, profiles (vertical cut showing the soil layers from surface to the rock) were opened up to 200 cm or to the depth limited by rock or hard substratum and studied in detail for all their morphological and physical characteristics. The soil and site characteristics were recorded for all profile sites on a standard proforma as per the guidelines given in USDA Soil Survey Manual (Soil Survey Staff, 2012). Apart from the transect study, profiles were also studied at random, almost like in a grid pattern, outside the transect areas to validate the soil map unit boundaries.

Based on the soil characteristics, the soils were grouped into different soil series. Soil series is the most homogeneous unit having similar horizons and properties and behaves similarly for a given level of management. Soil depth, texture, colour, kind of horizon and horizon sequence, amount and nature of gravel present, calcareousness, nature of substratum etc, were used as the major differentiating characteristics for identifying soil series occurring in the area. The differentiating characteristics used for identifying the soil series are given in Table 3.1. Based on the above characteristics, 10 soil series were identified in Raghunathanahalli-2 Microwatershed.

**Table 3.1 Differentiating Characteristics used for identifying Soil Series  
(Characteristics are of Series Control Section)**

Sl. No.	Soil Series	Depth (cm)	Colour (moist)	Texture	Gravel (%)	Horizon sequence	Calcareousness
<b>SOILS OF GRANITE GNEISS LANDSCAPE</b>							
1	Belagatti (BGT)	<25	10 YR3/1, 3/2, 4/2	gc	>35	Ap-Crk	es
2	Bhimanakunte (BMK)	75-100	5YR 3/3, 4/6,	gsc-gc	15-35	Ap-Bt-Ck	e-es

SOILS OF ALLUVIAL LANDSCAPE							
3	Muttal (MTL)	25-50	10YR3/2,3/3,4/2 7.5YR3/2,3/3,6/4	gc	15-35	Ap-Bw-Ck	e-ev
4	Kyasalapura (KSP)	50-75	5YR 3/2, 3/3, 3/4	gscl	15-35	Ap-Bt-Ck	e-es
5	Ravanaki (RNK)	50-75	7.5YR3/2,3/3,5/2,5/3 10YR3/1,3/2,4/1,4/2,5/1,6/1	c	<15	Ap-Bw-Cr	e-ev
6	Dambarahalli (DRL)	75-100	10YR 2/1, 3/1, 4/3	c	<15	Ap-Bss	e-es
7	Narasapura (NSP)	75-100	10 YR 3/1, 3/2, 4/2,	c	<15	Ap-Bw-Cr	e-es
8	Gatareddihal (GRH)	100-150	10YR2/1,3/1 2.5Y 4/3, 5/4	c	<15	Ap-Bss-Bck-Cr	es
9	Alawandi (AWD)	>150	10 YR 2/1, 3/2,	c	<15	Ap-Bss	e-es
10	Murlapur (MLR)	>150	10YR 2/1, 2/2, 3/1, 3/2, 4/1	c	10-20	Ap-Bss	e-es

### 3.4 Soil Mapping

The area under each soil series was further separated into soil phases and their boundaries delineated on the cadastral map based on the variations observed in the texture of the surface soil, slope, erosion, presence of gravel, stoniness etc. A soil phase is a subdivision of soil series based mostly on surface features that affect its use and management. The soil mapping units are shown on the map (Fig.3.5) in the form of symbols. During the survey many soil profile pits, few minipits and a few auger bores representing different landforms occurring in the microwatershed were studied. In addition to the profile study, spot observations in the form of minipits, road cuts, terrace cuts etc., were studied to validate the soil boundaries on the soil map. The soil map shows the geographic distribution and area extent of 17 mapping units representing 10 soil series occurring in the microwatershed. The soil map unit (soil legend) description is presented in Table 3.2. The soil phase map (management units) shows the distribution of 17 phases mapped in the microwatershed. Each mapping unit (soil phase) delineated on the map has similar soil and site characteristics. In other words, all the farms or survey numbers included in one phase will have similar management needs and have to be treated accordingly.

### 3.5 Laboratory Characterization

Soil samples for each series were collected from representative master profiles for laboratory characterization by following the methods outlined in the Laboratory Manual (Sarma *et al*, 1987). Surface soil samples collected in the year 2018 from Raghunathanahalli-2 farmer's fields (43 samples) for fertility status (major and micronutrients) at 320 m grid interval were analyzed in the laboratory (Katyal and Rattan, 2003). By linking the soil fertility data to the survey numbers through GIS, soil fertility maps were generated using Kriging method for the microwatershed.



### 3.6 Land Management Units (LMUs)

The 17 soil phases identified and mapped in the microwatershed were regrouped into 6 Land Management Units (LMU's) for the purpose of preparing a Proposed Crop Plan for sustained development of the microwatershed. The database (soil phases) generated under LRI was utilized for identifying Land Management Units (LMU's) based on the management needs. One or more than one soil site characteristic having influence on the management have been chosen for identification and delineation of LMUs. For Raghunathanahalli-2 Microwatershed, five soil and site characteristics, namely soil depth, soil texture, slope erosion and gravel content have been considered for defining LMUs. The land management units are expected to behave similarly for a given level of management.

**Table 3.2 Soil map unit description of Raghunathanahalli-2 Microwatershed**

Soil map unit No*	Soil Series	Soil Phase Symbol	Mapping Unit Description	Area in ha (%)
<b>SOILS OF GRANITE GNEISS LANDSCAPE</b>				
	BGT		Belagatti soils are very shallow (< 25 cm), well drained, have very dark gray to very dark grayish brown, calcareous, black gravelly clay soils occurring on very gently to gently sloping uplands under cultivation	<b>66 (15.07)</b>
11		BGTmB2g2	Clay surface, slope 1-3%, moderate erosion, very gravelly (35-60%)	66 (15.07)
	BMK		Bhimanakunte soils are moderately deep (75-100 cm), well drained, have dark reddish brown to yellowish red, calcareous, red gravelly sandy clay to clay soils occurring on very gently sloping uplands under cultivation	<b>152 (34.49)</b>
153		BMKiA1	Sandy clay surface, slope 0-1%, slight erosion	34(7.62)
154		BMKiB1g1	Sandy clay surface, slope 1-3%, slight erosion, gravelly (15-35%)	76 (17.35)
155		BMKiB1g2	Sandy clay surface, slope 1-3%, slight erosion, very gravelly (35-60%)	42 (9.52)
<b>SOILS OF ALLUVIAL LANDSCAPE</b>				
	MTL		Muttal soils are shallow (25-50 cm), well drained, have very dark grayish brown to dark brown, calcareous, black gravelly clay soils occurring on nearly level to gently sloping plains under cultivation	<b>53 (11.98)</b>
311		MTLmB2g1	Clay surface, slope 1-3%, moderate erosion, gravelly (15-35%)	18 (4.06)
312		MTLmB2g2	Clay surface, slope 1-3%, moderate erosion, very gravelly (35-60%)	35 (7.92)
	KSP		Kyasalapura soils are moderately shallow (50-75 cm), well drained, have dark reddish brown, calcareous, red gravelly sandy clay loam soils occurring on very gently sloping plains under cultivation	<b>13 (3.0)</b>
323		KSPiB1g2	Sandy clay surface, slope 1-3%, slight erosion,	6

Soil map unit No*	Soil Series	Soil Phase Symbol	Mapping Unit Description	Area in ha (%)
			very gravelly (35-60%)	(1.33)
325		KSPiB2g1	Sandy clay surface, slope 1-3%, moderate erosion, gravelly (15-35%)	7(1.67)
	RNK		Ravanaki soils are moderately shallow (50-75 cm), moderately well drained, have dark brown to very dark grayish brown and dark gray, calcareous, black clay soils occurring on nearly level to very gently sloping plains under cultivation	<b>42.17 (9.61)</b>
333		RNKmB1	Clay surface, slope 1-3%, slight erosion	12(2.83)
334		RNKmB1g1	Clay surface, slope 1-3%, slight erosion, gravelly (15-35%)	30(6.74)
337		RNKmB2g1	Clay surface, slope 1-3%, moderate erosion, gravelly (15-35%)	0.17 (0.04)
	DRL		Dambarahalli soils are moderately deep (75-100 cm), moderately well drained, have dark brown to very dark gray, calcareous, black cracking clay soils occurring on nearly level to very gently sloping plains under cultivation	<b>68 (15.42)</b>
344		DRLmA1	Clay surface, slope 0-1%, slight erosion	17(3.86)
350		DRLmB2	Clay surface, slope 1-3%, moderate erosion	51(11.56)
	NSP		Narasapura soils are moderately deep (75-100 cm), moderately well drained, have dark grayish brown to very dark grayish brown and very dark gray, sodic, calcareous, black cracking clay soils occurring on nearly level to very gently sloping plains under cultivation	<b>6 (1.47)</b>
357		NSPiB1g1	Sandy clay surface, slope 1-3%, slight erosion, gravelly (15-35%)	6 (1.47)
	GRH		Gatareddihal soils are deep (100-150 cm), moderately well drained, have light olive brown to very dark gray, sodic, calcareous, black cracking clay soils occurring on nearly level to very gently sloping plains under cultivation	<b>14 (3.21)</b>
371		GRHmB1	Clay surface, slope 1-3%, slight erosion	14(3.21)
	AWD		Alawandi soils are very deep (>150 cm), moderately well drained, have very dark grayish brown to black, calcareous, black cracking clay soils occurring on nearly level to very gently sloping plains under cultivation	<b>4 (0.87)</b>
425		AWDmB2g1	Clay surface, slope 1-3%, moderate erosion, gravelly (15-35%)	4 (0.87)
	MLR		Murlapur soils are very deep (>150 cm), moderately well drained, have very dark grayish brown to very dark gray, calcareous, black cracking clay soils occurring on nearly level to very gently sloping plains under cultivation	<b>16 (3.74)</b>
417		MLRmB1g2	Clay surface, slope 1-3%, slight erosion, very gravelly (35-60%)	16 (3.74)
1000		Others	Water body	<b>5(1.11)</b>

\*Soil map unit numbers are continuous for the taluk, not for the microwatershed




Fig 3.5 Soil Phase or Management Units- Raghunathanahalli-2 Microwatershed



## THE SOILS

Detailed information pertaining to the nature, extent and distribution of different kinds of soils occurring in Raghunathanahalli-2 Microwatershed is provided in this chapter. The microwatershed area has been identified as granite gneiss and alluvial landscapes based on geology. In all, 10 soil series are identified. Soil formation is the result of the combined effect of environmental and terrain factors that are reflected in soil morphology. The soil formation is dominantly influenced by the parent material, climate, time and relief.

A brief description of each of the 10 soil series identified followed by 17 soil phases (management units) mapped (Fig. 3.5) are furnished below. The physical and chemical characteristics of soil series identified in Raghunathanahalli-2 microwatershed are given in Table 4.1 along with soil classification. The soils in any one map unit differ from place to place in their depth, texture, slope, gravelliness, erosion or any other site characteristic that affect management. The soil phase map can be used for identifying the suitability of areas for growing specific crops or for other alternative uses and also for deciding the type of conservation structures needed. The detailed information on soil and site-characteristics like soil depth, surface soil texture, slope, erosion, gravelliness, AWC, LCC etc, with respect to each of the soil phase identified is given village/survey number wise for the microwatershed in Appendix-I.

### 4.1 Soils of Granite gneiss landscape

In this landscape, 2 soil series are identified and mapped. Of these, Bhimanakunte (BMK) series occupies major area of 152 ha (34%) and Belagatti (BGT) 66 ha (15%). The brief description of each soil series along with the soil phases identified and mapped is given below.

**4.1.1 Belagatti (BGT) Series:** Belagatti soils are very shallow (< 25 cm), well drained, have dark gray to dark grayish brown gravelly clay soils. They have developed from weathered granite gneiss and occur on very gently sloping uplands. The Belagatti series has been classified as a member of the clayey-skeletal, mixed (calc), isohyperthermic family of Lithic Ustorthents.

The thickness of the soil is less than 25 cm. Its colour is in 10 YR hue with value 3 to 4 and chroma 1 to 2. The texture is clay with more than 35 per cent gravel, and the available water capacity is very low (<50 mm/m). Only one soil phase was identified and mapped.





Landscape and soil profile characteristics of Belagatti (BGT) Series

**4.1.2 Bhimanakunte (BMK) Series:** Bhimanakunte soils are moderately deep (75-100 cm), well drained, have very dark reddish brown to yellowish red gravelly, calcareous sandy clay to clay soils. They have developed from weathered granite gneiss and occur on nearly level to very gently sloping uplands under cultivation. The Bhimanakunte series has been classified as a member of the fine, mixed (calc), isohyperthermic family of Typic Haplustalfs.

The thickness of the solum ranges from 75 to 99 cm. The thickness of A-horizon ranges from 15 to 18 cm. Its colour is in 5 YR hue with value 3 to 4 and chroma 2 to 4. The texture is sandy clay loam to clay. The thickness of B-horizon ranges from 57 to 82 cm. Its colour is in 2.5 YR, 5 YR and 7.5 YR hue with value and chroma ranging from 3 to 4. Its texture is sandy clay to clay soil with 15 to 35 per cent gravel. The available water capacity is low (51-100 mm/m). Three soil phases were identified and mapped.



Landscape and soil profile characteristics of Bhimanakunte (BMK) Series

## 4.2 Soils of Alluvial landscape

In this landscape, 8 soil series were identified and mapped. Of these, Dambarahalli (DRL) series occupies major area of 68 ha (15%) followed by Muttal (MTL) 53 ha (12%), Ravanaki (RNK) 42 ha (10%), Murlapur (MLR) 16 ha (4%), Gatareddihal (GRH) 14 ha (3%), Kyasalapura (KSP) 13 ha (3%), Narasapura (NSP) 6 ha (1%) and Alawandi (AWD) 4 ha (1%). The brief description along with the soil phases identified and mapped is given below.

**4.2.1 Muttal (MTL) Series:** Muttal soils are shallow (25-50 cm), well drained, have dark brown to very dark grayish brown, calcareous gravelly clay soils. They have developed from alluvium and occur on nearly level to very gently sloping plains. The Muttal series has been classified as a member of the clayey, mixed (calc), isohyperthermic family of (Paralithic) Haplustepts.

The thickness of the solum ranges from 30 to 50 cm. The thickness of A-horizon ranges from 15 to 18 cm. Its colour is in 7.5 YR and 10 YR hue with value 2 to 3 and chroma 2.5 to 4. The texture varies from sandy clay to clay with 10 to 15 per cent gravel. The thickness of B horizon ranges from 18 to 32 cm. Its colour is in 10 YR and 7.5 YR hue with value 2 to 6 and chroma 2 to 4. Its texture is sandy clay to clay. The available water capacity is low (51-100 mm/m). Two soil phases were identified and mapped.



Landscape and soil profile characteristics of Muttal (MTL) Series

**4.2.2 Kyasalapura (KSP) Series:** Kyasalapura soils are moderately shallow (50-75cm), well drained, have dark reddish brown, gravelly sandy clay loam soils. They are developed from alluvium and occur on very gently sloping plains under cultivation. The Kyasalapura series has been classified as a member of the fine-loamy, mixed (calc), isohyperthermic family of Typic Haplustalfs.



The thickness of the solum ranges from 53 to 75 cm. The thickness of A-horizon ranges from 17 to 23 cm. Its colour is in 2.5YR, 5 YR and 7.5 YR hue with value 3 to 5 and chroma 2 to 4. The texture varies from sandy clay loam to sand clay with 15 to 30 per cent gravel. The thickness of B-horizon varies from 33 to 55 cm. Its colour is in 2.5 YR and 5 YR hue with value 3 and chroma 2 to 4. Texture is sandy clay loam to sandy clay with 15 to 35 per cent gravel. The available water capacity is very low (<50mm/m). Two soil phases were identified and mapped.



Landscape and soil profile characteristics of Kyasalapura (KSP) Series

**4.2.3 Ravanaki (RNK) Series:** Ravanaki soils are moderately shallow (50-75 cm), well drained, have dark brown to very dark grayish brown, calcareous clay soils. They have developed from alluvium and occur on nearly level to very gently sloping plains. The Ravanaki series has been classified as a member of the very-fine, smectitic (calc), isohyperthermic family of Typic Haplustepts.

The thickness of the solum ranges from 50 to 75 cm. The thickness of A-horizon ranges from 15 to 20 cm. Its colour is in 7.5 YR and 10 YR hue with value 2 to 3 and chroma 2.5 to 4. The texture varies from sandy clay to clay with 10 to 15 per cent gravel. The thickness of B horizon ranges from 35 to 60 cm. Its colour is in 10 YR and 7.5 YR hue with value 2 to 6 and chroma 2 to 4. Its texture is sandy clay to clay with gravel content of 10 to 20 per cent. The available water capacity is low (51-100 mm/m). Three soil phases were identified and mapped.





Landscape and Soil Profile Characteristics of Ravanaki (RNK) Series

**4.2.4 Dambarahalli (DRL) Series:** Dambarahalli soils are moderately deep (75-100 cm), moderately well drained, have black and very dark gray to dark brown, calcareous cracking clay soils. They have developed from alluvium and occur on very gently to gently sloping plains under cultivation. The Dambarahalli series has been classified as a member of the very-fine, smectitic (calc), isohyperthermic family of Typic Haplusterts.

The thickness of the solum ranges from 75 to 99 cm. The thickness of A-horizon ranges from 13 to 24 cm. Its colour is in 10 YR hue with value 3 to 4 and chroma 1 to 2. The texture is clay. The thickness of B horizon ranges from 54 to 85 cm. Its colour is in 10 YR hue with value 2 to 4 and chroma 1 to 3. Its texture is clay and is calcareous. The available water capacity is high (151-200 mm/m). Two soil phases were identified and mapped.



Landscape and soil profile characteristics of Dambarahalli (DRL) Series

**4.2.5 Narsapura (NSP) Series:** Narsapura soils are moderately deep (75-100 cm), moderately well drained, have dark grayish brown to very dark grayish brown and very dark gray, sodic, calcareous, black cracking clay soils. They have developed from alluvium and occur on very gently sloping plains. The Narsapura series has been classified as a member of the very-fine, smectitic (calc), isohyperthermic, family of Vertic Haplustepts.

The thickness of the solum is 76 to 98 cm. The thickness of A-horizon ranges from 15 to 19 cm. Its colour is in 10 YR hue with value 3 and chroma 1 to 2. The texture is clay with no gravel. The thickness of B horizon ranges from 57 to 83 cm. Its colour is in 10 YR hue with value 3 to 5 and chroma 1 to 3. Its texture is clay and is calcareous. The available water capacity is medium (101-150 mm/m). Only one soil phase was identified and mapped.



Landscape and soil profile characteristics of Narsapura (NSP) Series

**4.2.6 Gatareddihal (GRH) Series:** Gatareddihal soils are deep (100-150 cm), moderately well drained, have black or dark grey to light olive brown, sodic, calcareous cracking clay soils. They are developed from alluvium and occur on nearly level to very gently sloping plains under cultivation. The Gatareddihal series has been classified as a member of the very-fine, smectitic (calc), isohyperthermic family of Sodic Haplusterts.

The thickness of the solum ranges from 102 to 149 cm. The thickness of A-horizon ranges from 12 to 19 cm. Its colour is in 7.5 YR, 10 YR hue with value 3 to 4 and chroma 1 to 6. The texture is sandy clay loam to clay. The thickness of B-horizon ranges from 86 to 117 cm. Its colour is in 10 YR and 7.5 YR hue with value 3 and chroma 2 to 6. Texture is clay with less than 15 per cent gravel. The available water capacity is very high (>200 mm/m). Only one soil phase was identified and mapped.






Landscape and soil profile characteristics of Gatareddihal (GRH) Series

**4.2.7 Alawandi (AWD) Series:** Alawandi soils are very deep (>150 cm), moderately well drained, have black to very dark grayish brown, calcareous cracking clay soils. They have developed from alluvium and occur on nearly level to very gently sloping plains under cultivation. The Alawandi series has been classified as a member of the fine smectitic (calc), isohyperthermic family of Typic Haplusterts.

The thickness of the solum is more than 150 cm. The thickness of A-horizon ranges from 16 to 26 cm. Its colour is in 10 YR hue with value 2 to 3 and chroma 1 to 2. The texture varies from sandy clay to clay. The thickness of B horizon is more than 150 cm. Its colour is in 10 YR hue with value 2 to 3 and chroma 1 to 3. Its texture is clay and is calcareous. The available water capacity is very high (>200 mm/m). Only one soil phase was identified and mapped.



Landscape and soil Profile Characteristics of Alawandi (AWD) Series

**4.2.8 Murlapur (MLR) series:** Murlapur soils are very deep (>150 cm), moderately well drained, have very dark grayish brown to very dark gray, calcareous black cracking clay soils. They have developed from alluvium and occur on nearly level to very gently sloping plains. The Murlapur series has been classified as a member of the very fine, smectitic (calc), isohyperthermic family of Typic Haplusterts.

The thickness of the solum is >150 cm. The thickness of A-horizon ranges from 20 to 25 cm. Its colour is in 10 YR hue with value 3 and chroma 1. The texture is clay with no gravel. The thickness of B horizon ranges from 150 to 190 cm. Its colour is in 10 YR hue with value 3 to 4 and chroma 1 to 2. Its texture is clay. The available water capacity is very high (>200 mm/m). Only one soil phase was identified and mapped.



Landscape and soil profile characteristics of Murlapur (MLR) Series

**Table: 4.1 Physical and Chemical Characteristics of Soil Series identified in Raghunathanahalli-2 Microwatershed**

**Series Name:** Belagatti (BGT), **Pedon:** A2/RM-5

**Location:** 15°19'10.8"N, 75°57'48.1"E, Kavalura village, Koppal Taluk and District

**Analysis at:** NBSS&LUP, Regional Centre, Bangalore. **Classification:** Clayey-skeletal, mixed (calc), isohyperthermic Lithic Ustorthents

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-23	Ap	36.14	20.34	43.52	10.87	6.93	5.97	8.42	3.94	40	c	29.53	17.97

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO <sub>3</sub>	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP
	Water	CaCl <sub>2</sub>	M KCl				Ca	Mg	K	Na	Total				
							cmol kg <sup>-1</sup>								
0-23	8.4			0.157	0.12	18.24			0.73	0.50		44.84	1.03		1.11

*Contd....*

**Series Name:** Muttal (MTL), **Pedon:** RM-13

**Location:** 15°14'30.8"N, 75°56'50.6"E, Gatareddihalla village, Koppal Taluk and District

**Analysis at:** NBSS&LUP, Regional Centre, Bangalore. **Classification:** Clayey, mixed (calc), isohyperthermic (Paralithic) Haplustepts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-20	Ap	39.05	13.74	47.21	3.05	5.05	8.21	14.63	8.11	15-30	c	29.95	17.94
20-34	Bwk	28.77	19.57	51.66	4.81	4.71	4.92	9.09	5.24	10	c	33.44	21.56

Depth (cm)	pH (1:2.5)			E.C. (1:2.5) dS m <sup>-1</sup>	O.C. %	CaCO <sub>3</sub> %	Exchangeable bases					CEC	CEC/Clay	Base saturation %	ESP %
	Water	CaCl <sub>2</sub>	M KCl				Ca	Mg	K	Na	Total				
							cmol kg <sup>-1</sup>								
0-20	8.27	-	-	0.202	0.79	6.10	-	-	0.62	0.25	-	36.64	0.78	-	0.69
20-34	8.36	-	-	0.177	0.99	23.04	-	-	0.29	0.38	-	39.60	0.77	-	0.96

*Contd....*

**Series Name:** Ravanaki (RNK), **Pedon:** RM-20

**Location:** 15°14'22.7"N, 75°57'45.8"E, Gatareddihalla village, Koppal Taluk and District

**Analysis at:** NBSS&LUP, Regional Centre, Bangalore. **Classification:** Very-fine, smectitic (calc), isohyperthermic Typic Haplustepts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-28	Ap	24.43	17.76	57.81	5.30	3.89	3.78	7.14	4.32	20	c	41.40	29.60
28-55	Bw	18.77	15.59	65.64	2.74	3.73	2.85	4.83	4.61	10	c	46.71	35.18

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO <sub>3</sub>	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP			
	Water	CaCl <sub>2</sub>	M KCl				dS m <sup>-1</sup>	%	%	Ca	Mg					K	Na	Total
										cmol kg <sup>-1</sup>						%	%	
0-28	8.86	-	-	0.483	0.63	15.48	-	-	0.86	6.27	-	37.00	0.64	-	6.78			
28-55	8.61	-	-	1.4	0.23	13.68	-	-	0.68	12.27	-	53.20	0.81	-	9.22			

*Contd....*

**Series Name:** Dombarahalli (DRL), **Pedon:** R-8

**Location:** 15°13'96.2"N, 75°57'48.6" E Ragunathanahalli village, Koppal Taluk and District

**Analysis at:** NBSS&LUP, Regional Centre, Bangalore.

**Classification:** Very-fine, smectitic (calc), isohyperthermic Typic Haplusterts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-15	Ap	28.25	19.48	52.27	4.76	4.44	4.87	8.23	5.95	-	c	39.86	27.20
15-27	BA1	21.55	20.00	58.45	3.76	2.76	3.43	6.30	5.30	-	c	46.35	34.84
27-45	Bss1	14.86	20.89	64.25	2.46	2.23	2.23	3.91	4.02	-	c	57.99	41.06
45-80	Bss2	10.42	19.04	70.54	1.74	1.97	1.27	2.78	2.66	-	c	66.36	36.24

Depth (cm)	pH (1:2.5)			E.C. (1:2.5) dS m <sup>-1</sup>	O.C. %	CaCO <sub>3</sub> %	Exchangeable bases					CEC	CEC/Clay	Base saturation %	ESP %
	Water	CaCl <sub>2</sub>	M KCl				Ca	Mg	K	Na	Total				
0-15	8.78	-	-	0.42	0.32	12.35	-	-	0.59	4.25	-	49.70	0.95	100.00	5.62
15-27	9.03	-	-	0.61	0.30	12.48	-	-	0.30	8.96	-	57.23	0.98	100.00	10.07
27-45	9.10	-	-	0.67	0.34	11.70	-	-	0.25	11.85	-	60.71	0.95	100.00	14.05
45-80	9.18	-	-	0.86	0.32	13.39	-	-	0.27	15.40	-	63.33	0.90	100.00	18.45

Contd....



**Series Name:** Narsapura (NSP), **Pedon:** A2/RM-2

**Location:** 15°19'86.9"N, 75°57'86.1"E, Kavalura village, Koppal Taluk and District

**Analysis at:** NBSS&LUP, Regional Centre, Bangalore. **Classification:** Very-fine, smectitic (calc), isohyperthermic Vertic Haplustepts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-29	Ap	31.32	16.52	52.16	5.51	5.40	5.51	9.83	5.08	10	c	38.86	27.64
29-52	Bw1	13.30	22.08	64.62	2.52	2.41	2.41	3.67	2.29	05	c	49.88	40.05
52-77	BW2	13.22	17.39	69.40	3.56	2.41	1.95	2.76	2.53	05	c	51.33	41.55

Depth (cm)	pH (1:2.5)			E.C. (1:2.5) dS m <sup>-1</sup>	O.C. %	CaCO <sub>3</sub> %	Exchangeable bases					CEC	CEC/Clay	Base saturation %	ESP %
	Water	CaCl <sub>2</sub>	M KCl				Ca	Mg	K	Na	Total				
0-29	9.16	-	-	0.615	0.23	9.36	-	-	0.72	10.98	-	51.09	0.98	-	8.60
29-52	8.69	-	-	2.01	0.5	8.64	-	-	0.55	24.42	-	60.63	0.94	-	16.11
52-77	8.52	-	-	2.68	0.46	7.68	-	-	0.50	25.65	-	60.74	0.88	-	16.90

*Contd....*

**Series Name:** Gatareddihal (GRH) Pedon: R-7

**Location:** 15°14'20.8"N, 76°04'28.4" E Gudlanur village, Koppal Taluk and District

**Analysis at:** NBSS&LUP, Regional Centre, Bangalore.

**Classification:** Very-fine, smectitic (calc), isohyperthermic Sodic Haplusterts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-18	Ap	20.07	19.71	60.23	1.76	3.75	3.64	3.42	7.50	-	c	41.70	29.56
18-51	Bss1	15.11	17.47	67.42	3.16	3.04	2.25	3.38	3.27	-	c	59.43	38.52
51-80	Bss2	13.19	18.74	68.07	1.80	2.93	2.37	3.04	3.04	-	c	60.69	40.91
80-107	Bss3	17.54	19.50	62.96	2.46	4.13	3.24	4.25	3.46	-	c	57.25	37.31
107-131	BC	9.42	17.48	73.10	1.48	1.82	1.36	1.93	2.84	-	c	64.62	43.98

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO <sub>3</sub>	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP			
	Water	CaCl <sub>2</sub>	M KCl				dS m <sup>-1</sup>	%	%	Ca	Mg					K	Na	Total
										cmol kg <sup>-1</sup>						%	%	
0-18	9.08	-	-	0.23	0.33	6.89	-	-	0.70	6.36	-	63.21	1.05	100.00	7.11			
18-51	9.19	-	-	0.61	0.49	9.10	-	-	0.54	14.20	-	66.05	0.98	100.00	15.98			
51-80	9.27	-	-	0.56	0.29	9.36	-	-	0.49	14.75	-	65.63	0.96	100.00	17.07			
80-107	9.28	-	-	0.57	0.39	9.62	-	-	0.44	14.64	-	63.95	1.02	100.00	17.49			
107-131	9.04	-	-	1.08	0.31	8.32	-	-	0.52	16.40	-	68.36	0.94	100.00	17.30			

Contd....

**Series Name:** Alawandi (AWD)

**Pedon:** R-16

**Location :** 15°13'08.2"N, 76°15'27.3" E Neeralagi village, Koppal Taluk and District

**Analysis at:** NBSS&LUP, Regional Centre, Bangalore.

**Classification:** Fine, smectitic (calc), isohyperthermic Typic Haplusterts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-17	Ap	20.88	25.75	53.37	3.31	4.31	4.31	5.19	3.76	-	c	33.11	25.58
17-39	Bss1	25.99	19.79	54.22	5.04	5.48	5.04	5.92	4.50	-	c	33.11	26.23
39-70	Bss2	26.76	17.80	55.44	2.93	5.31	5.53	7.37	5.63	-	c	36.15	28.67
70-111	Bss3	23.83	20.25	55.93	4.15	4.81	4.92	6.01	3.93	-	c	43.60	33.71
111-139	Bss4	21.21	20.40	58.40	2.79	4.80	4.91	5.25	3.46	-	c	46.92	36.28
139-162	Bss5	13.15	20.96	65.90	1.69	2.47	2.36	3.37	3.26	-	c	54.96	41.81

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO <sub>3</sub>	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP
	Water	CaCl <sub>2</sub>	M KCl				Ca	Mg	K	Na	Total				
				dS m <sup>-1</sup>	%	%	cmol kg <sup>-1</sup>					%	%		
0-17	8.10	-	-	0.37	0.52	9.48	-	-	0.40	1.56	-	51.30	0.96	100.00	1.22
17-39	8.60	-	-	0.24	0.52	9.60	-	-	0.14	4.60	-	52.60	0.97	100.00	3.50
39-70	8.89	-	-	0.27	0.52	9.48	-	-	0.16	2.41	-	53.90	0.97	100.00	1.78
70-111	9.10	-	-	0.35	0.54	11.28	-	-	0.15	8.95	-	54.10	0.97	100.00	6.61
111-139	9.15	-	-	0.41	0.58	10.80	-	-	0.15	7.36	-	56.10	0.96	100.00	5.24
139-162	9.16	-	-	0.50	0.50	15.48	-	-	0.19	10.19	-	61.66	0.94	100.00	6.61

Contd....

**Series Name:** Murlapur (MLR), **Pedon:** R-A1/16

**Location:** 15°19'42.9"N, 75°55'84.7"E, Kavalura village, Koppal Taluk and District

**Analysis at:** NBSS&LUP, Regional Centre, Bangalore. **Classification:** Very-fine, smectitic (calc), isohyperthermic Typic Haplusterts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-30	Ap	27.97	13.96	58.07	4.22	4.77	6.66	8.10	4.22	10	c	36.24	25.90
30-53	BA	26.34	17.48	56.17	4.17	5.05	6.04	7.24	3.84	05	c	38.55	28.98
53-83	Bss1	19.35	19.55	61.10	3.13	3.91	4.03	5.48	2.80	05	c	44.48	33.69
83-105	Bss2	16.63	17.47	65.90	2.70	3.93	2.92	3.93	3.15	<5	c	50.55	38.11
105-160	Bss3	14.69	20.34	64.97	0.79	2.26	4.07	4.18	3.39	<5	c	51.54	40.19

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO <sub>3</sub>	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP
	Water	CaCl <sub>2</sub>	M KCl				Ca	Mg	K	Na	Total				
				dS m <sup>-1</sup>	%	%	cmol kg <sup>-1</sup>						%	%	
0-30	9.19	-	-	0.313	0.57	10.08	-	-	0.64	5.67	-	42.08	0.72	-	5.39
30-53	9.22	-	-	0.449	0.24	13.08	-	-	0.35	8.23	-	41.02	0.73	-	8.02
53-83	9.17	-	-	0.377	0.82	16.92	-	-	0.39	14.28	-	51.20	0.84	-	11.16
83-105	9.18	-	-	0.477	0.61	15.48	-	-	0.35	13.19	-	53.11	0.81	-	9.94
105-160	9.01	-	-	1.17	0.24	16.92	-	-	0.43	19.61	-	53.95	0.83	-	14.54

## INTERPRETATION FOR LAND RESOURCE MANAGEMENT

The most important soil and site characteristics that affect the land use and conservation needs of an area are land capability, land irrigability, soil depth, soil texture, coarse fragments, available water capacity, soil slope, soil erosion, soil reaction etc. These are interpreted from the data base generated through land resource inventory and several thematic maps are generated. These would help in identifying the areas suitable for growing crops and, soil and water conservation measures and structures needed thus helping to maintain good soil health for sustained crop production. The various thematic maps generated are described below.

### 5.1 Land Capability Classification

Land capability classification is an interpretative grouping of soil map units (soil phases) mainly based on inherent soil characteristics, external land features and environmental factors that limit the use of land for agriculture, pasture, forestry or other uses on a sustained basis (IARI, 1971). The land and soil characteristics used to group the land resources in an area into various land capability classes, subclasses and units are

*Soil characteristics:* Soil depth, soil texture, coarse fragments, soil reaction, available water capacity, calcareousness, salinity/alkali *etc.*

*Land characteristics:* Slope, erosion, drainage and rock-outcrops.

*Climate:* Total rainfall and its distribution, and length of crop growing period.

The Land Capability Classification system is divided into land capability classes, subclasses and units based on the level of information available. Eight land capability classes are recognized. They are

*Class I:* They are very good lands that have no limitations or very few limitations that restrict their use.

*Class II:* They are good lands that have minor limitations and require moderate conservation practices.

*Class III:* They are moderately good lands that have severe limitations that reduce the choice of crops or that require special conservation practices.

*Class IV:* They are fairly good lands that have very severe limitations that reduce the choice of crops or that require very careful management.

*Class V:* Soils in these lands are not likely to erode, but have other limitations like wetness that are impractical to remove and as such not suitable for agriculture, but suitable for pasture or forestry with minor limitations.

*Class VI:* The lands have severe limitations that make them generally unsuitable for cultivation, but suitable for pasture or forestry with moderate limitations.

*Class VII:* The lands have very severe limitations that make them unsuitable for cultivation, but suitable for pasture or forestry with major limitations.

*Class VIII:* Soil and other miscellaneous areas (rock lands) that have very severe limitations that nearly preclude their use for any crop production, but suitable for wildlife, recreation and installation of wind mills.

The land capability subclasses are recognised based on the dominant limitations observed within a given land capability class. The subclasses are designated by adding a lower case letter like ‘e’, ‘w’, ‘s’, or ‘c’ to the class numeral. The subclass “e” indicates that the main hazard is risk of erosion, “w” indicates drainage or wetness as a limitation for plant growth, “s” indicates shallow soil depth, coarse or heavy textures, calcareousness, salinity/alkali or gravelliness and “c” indicates limitation due to climate.

The land capability subclasses have been further subdivided into land capability units based on the kinds of limitations present in each subclass. Ten land capability units are identified in grouping the soil map units. They are stony or rocky (0), erosion hazard (slope, erosion) (1), coarse texture (sand, loamy sand, sandy loam) (2), fine texture (cracking clay, silty clay) (3), slowly permeable subsoil (4), coarse underlying material (5), salinity/alkali (6), stagnation, overflow, high ground water table (7), soil depth (8) and fertility problems (9). The capability units thus identified have similar soil and land characteristics that respond similarly to a given level of management. The soils of the microwatershed have been classified upto land capability subclass level.

The 17 soil map units identified in the Raghunathanahalli-2 microwatershed are grouped under 3 Land capability classes and 5 land capability subclasses (Fig. 5.1). Entire cultivated area of about 435 ha (99%) is suitable for agriculture. An area of about 5 ha (1%) is under habitation and settlements.

Maximum area of about 312 ha (71%) is good lands (Class II) and distributed in the major part of the microwatershed with minor problems of soil and erosion. An area about 57 ha (13%) is moderately good lands (Class III) and distributed in the central, western, eastern and northeastern part of the microwatershed with moderate limitations of soil and erosion. Fairly good lands (Class IV) cover an area of about 66 ha (15%) and distributed in the northeastern and eastern part of the microwatershed with severe limitations of soil and erosion.






Fig. 5.1 Land Capability map of Raghunathanahalli-2 Microwatershed

## 5.2 Soil Depth

Soil depth refers to the depth of the soil occurring above the parent material or hard rock. The depth of the soil determines the effective rooting depth for plants and in accordance with soil texture, mineralogy and gravel content, the capacity of the soil column to hold water and nutrient availability. Soil depth is one of the most important soil characteristic that is used in differentiating soils into different soil series. The soil depth classes used in identifying soils in the field are very shallow (<25 cm), shallow (25-50 cm), moderately shallow (50-75 cm), moderately deep (75-100 cm), deep (100-150 cm) and very deep (>150 cm). They were used to classify the soils into different depth classes and a soil depth map was generated (Fig. 5.2).

Very shallow (<25 cm) soils cover an area of about 66 ha (15%) and occur in the northeastern and eastern part of the microwatershed. An area of about 53 ha (12%) is under shallow (25-50 cm) soils and distributed in the central, eastern and western part of the microwatershed. Moderately shallow (50-75 cm) soils cover an area of about 56 ha (13%) and occur in the southwestern, eastern and northern part of the microwatershed. Moderately deep (75-100 cm) soils cover a major area of about 226 ha (51%) and distributed in the major part of the microwatershed. An area of about 14 ha (3%) is under deep (100-150 cm) soils and occur in the southern and eastern part of the microwatershed. Very deep (>150 cm) soils occupy an area of about 20 ha (5%) and occur in the southern, southwestern and northeastern part of the microwatershed.

The most productive lands cover about 34 ha (8%) where all climatically adapted long duration crops can be grown. The problem soils cover about 119 ha (27%) area where only short duration crops can be grown and the probability of crop failure is high.




Fig. 5.2 Soil Depth map of Raghunathanahalli-2 Microwatershed

### 5.3 Surface Soil Texture

Texture is an expression to indicate the coarseness or fineness of the soil as determined by the relative proportion of primary particles of sand, silt and clay. It has a direct bearing on the structure, porosity, adhesion and consistence. The surface layer of a soil to a depth of about 25 cm is the layer that is most used by crops and plants. The surface soil textural class provides a guide to understanding soil-water retention and availability, nutrient holding capacity, infiltration, workability, drainage, physical and chemical behaviour, microbial activity and crop suitability. The textural classes used for LRI were used to classify and a surface soil texture map showing sandy, loamy and clayey at the surface was generated. The area extent and their geographical distribution in the microwatershed is shown in Fig.5.3.

An entire cultivated area of about 436 ha (99%) has soils that are clayey (sandy clay and clay) at the surface. Thus, entire cultivated area of the microwatershed falls under clayey soils at the surface.

Entire cultivated area has most productive lands with respect to surface soil texture where they are clayey soils. These soils have high potential for soil-water

retention and availability, and nutrient retention and availability, but have more problems of drainage, infiltration, workability and other physical problems.




Fig. 5.3 Surface Soil Texture map of Raghunathanahalli-2 Microwatershed

#### 5.4 Soil Gravelliness

Gravel is the term used for describing coarse fragments between 2 mm and 7.5 cm diameter and stones for those between 7.5 cm and 25 cm. The presence of gravel and stones in soil reduces the volume of soil responsible for moisture and nutrient storage, drainage, infiltration and runoff, and hinders plant growth by impeding root growth and seedling emergence, intercultural operations and farm mechanization. The gravelliness classes used in LRI were used to classify the soils and using these classes, a gravelliness map was generated. The area extent and their spatial distribution in the microwatershed is shown in Fig.5.4.

An area of about 128 ha (29%) has non gravelly (<15%) soils and occur in the central, southern, eastern, southwestern, western and northern part of the microwatershed. An area of about 142 ha (32%) has gravelly (15-35%) soils and distributed in the eastern, southern, southwestern, northwestern, northern and northeastern part of the microwatershed. Maximum area of about 166 ha (38%) has very gravelly (35-60%) soils and occur in the central, southern, southwestern, western, northern, northeastern and eastern part of the microwatershed.

An area of about 128 ha (29%) are most productive lands with respect to gravelliness. They are non-gravelly with less than 15 per cent gravel and have potential

for growing both annual and perennial crops. The problem lands cover a major area of about 308 ha (70%) that are gravelly to very gravelly where only medium or short duration crops can be grown.




Fig. 5.4 Soil Gravelliness map of Raghunathanahalli-2 Microwatershed

### 5.5 Available Water Capacity

The soil available water capacity (AWC) is estimated based on the ability of the soil column to retain water between the tensions of 0.33 and 15 bar in a depth of 100 cm or the entire solum if the soil is shallower. The AWC of the soils (soil series) as estimated by considering the soil texture, mineralogy, soil depth and gravel content (Sehgal *et al.*, 1990) and accordingly the soil map units were grouped into five AWC classes *viz.* very low (<50 mm/m), low (50-100 mm/m), medium (100-150 mm/m), high (150-200 mm/m) and very high (>200 mm/m) and using these values, an AWC map was generated (Fig. 5.5).

An area of about 80 ha (18%) has soils that are very low (<50 mm/m) in available water capacity and distributed in the northern, northeastern and eastern part of the microwatershed. Low (51-100 mm/m) in available water capacity cover a major area of about 247 ha (56%) and occur in the major part of the microwatershed. An area of about 74 ha (17%) is medium (101-150 mm/m) in available water capacity and occur in the western and southwestern part of the microwatershed. An area of about 34 ha (8%) is very high (>200 mm/m) in available water capacity and occur in the northeastern, southwestern, southern and eastern part of the microwatershed.



An area of about 327 ha (74%) in the microwatershed has soils that are problematic with regard to available water capacity. Here, only short duration crops can be grown and the probability of crop failure is very high. These areas are best put to other alternative uses. An area of about 34 ha (8%) has soils that have very high potential (>200 mm/m) with regard to available water capacity where all climatically adapted long duration crops can be grown successfully.




Fig. 5.5 Soil Available Water Capacity map of Raghunathanahalli-2 Microwatershed

## 5.6 Soil Slope

Soil slope refers to the inclination of the surface of the land. It is defined by gradient, shape and length, and is an integral feature of any soil as a natural body. Slope is considered important in soil genesis, land use and land development. The length and gradient of slope influences the rate of runoff, infiltration, erosion and deposition. The soil map units were grouped into different slope classes and a slope map was generated showing the area extent and their geographic distribution of different slope classes in the microwatershed (Fig. 5.6).

An area of about 51 ha (11%) falls under nearly level (0-1% slope) lands and distributed in the northern and western part of the microwatershed. Maximum area of about 385 ha (87%) falls under very gently sloping (1-3% slope) lands and distributed in the major part of the microwatershed.

Entire cultivated area of the microwatershed has soils that have high potential in respect of soil slopes. In these areas, all climatically adapted annual and perennial crops

can be grown without much soil and water conservation and other land development measures.




Fig. 5.6 Soil Slope map of Raghunathanahalli-2 Microwatershed

## 5.7 Soil Erosion

Soil erosion refers to the wearing away of the earth's surface by the forces of water, wind and ice involving detachment and transport of soil by raindrop impact. It is used for accelerated soil erosion resulting from disturbance of the natural landscape by burning, excessive grazing and indiscriminate felling of forest trees and tillage, all usually by man. The erosion classes showing an estimate of the current erosion status as judged from field observations in the form of rills, gullies or a carpet of gravel on the surface are recorded. Four erosion classes, viz, slight erosion (e1), moderate erosion (e2), severe erosion (e3) and very severe erosion (e4) are recognized. The soil map units were grouped into different erosion classes and a soil erosion map generated. The area extent and their spatial distribution in the microwatershed is given in Figure 5.7.

Soils that are slightly eroded (e1 class) cover a major area of about 254 ha (58%) and distributed in the major part of the microwatershed. Soils that are moderately eroded (e2 class) cover an area of about 182 ha (41%) and distributed in the central, southwestern, northern, northeastern and eastern part of the microwatershed.

An area of about 182 ha (41%) in the microwatershed is problematic because of moderate erosion. For these areas, taking up soil and water conservation and other land development measures are needed.






Fig. 5.7 Soil Erosion map of Raghunathanahalli-2 Microwatershed



## **FERTILITY STATUS**

Soil fertility plays an important role in increasing crop yield. The adoption of high yielding varieties that require high amounts of nutrients has resulted in deficiency symptoms in crops and plants due to imbalanced fertilization and poor inherent fertility status, as these areas are characterised by low rainfall and high temperatures. Hence, it is necessary to know the fertility (macro and micro nutrients) status of the soils of the watersheds for assessing the kind and amount of fertilizers required for each of the crop intended to be grown. For this purpose, the surface soil samples collected from the grid points (one soil sample at every 320 m grid interval) all over the microwatershed through land resource inventory in the year 2018 were analysed for pH, EC, organic carbon, available phosphorus and potassium, and for micronutrients like zinc, boron, copper, iron and manganese, and secondary nutrient sulphur.

Soil fertility data generated has been assessed and individual maps for all the nutrients for the microwatershed have been generated using the Kriging method under GIS. The village/survey number wise fertility data for the microwatershed is given in Appendix-II.

### **6.1 Soil Reaction (pH)**

The soil analysis of the Raghunathanahalli-2 microwatershed for soil reaction (pH) showed that an area of about 58 ha (13%) is under moderately alkaline (pH 7.8-8.4) in soil reaction and distributed in the western, northwestern, northern and northeastern part of the microwatershed. Maximum area of about 378 ha (86%) is under strongly alkaline (pH 8.4-9.0) and occur in the major part of the microwatershed (Fig.6.1). Thus, entire cultivated area of the microwatershed falls under alkaline condition.

### **6.2 Electrical Conductivity (EC)**

The Electrical Conductivity of the soils of the entire microwatershed area is  $<2 \text{ dS m}^{-1}$  (Fig 6.2) and as such the soils are non-saline.

### **6.3 Organic Carbon (OC)**

The soil organic carbon content (an index of available Nitrogen) of the microwatershed is medium (0.5-0.75%) in a major area of about 338 ha (77%) and occur in the major part of the microwatershed. An area of about 98 ha (22%) is low ( $<0.5\%$ ) and distributed in the southwestern, eastern and western part of the microwatershed (Fig.6.3).

### **6.4 Available Phosphorus**

An area of about 18 ha (4%) is medium (23-57 kg/ha) in available phosphorus and distributed in the northern part of the microwatershed. Low ( $<23 \text{ kg/ha}$ ) in available

phosphorus cover a major area of about 418 ha (95%) and distributed in the major part of the microwatershed (Fig 6.4).

### 6.5 Available Potassium

Entire cultivated area of the microwatershed is high (>337 kg/ha) in available potassium (Fig.6.5).

### 6.6 Available Sulphur

An area of about 133 ha (30%) is medium (10-20 ppm) in available sulphur and occur in the western and southwestern part of the microwatershed. Maximum area of about 303 ha (69%) is high (>20 ppm) in available sulphur and distributed in the major part of the microwatershed (Fig.6.6).




Fig.6.1 Soil Reaction (pH) map of Raghunathanahalli-2 Microwatershed




Fig.6.2 Electrical Conductivity (EC) map of Raghunathanahalli-2 Microwatershed




Fig.6.3 Soil Organic Carbon (OC) map of Raghunathanahalli-2 Microwatershed




Fig.6.4 Soil Available Phosphorus map of Raghunathanahalli-2 Microwatershed .




Fig.6.5 Soil Available Potassium map of Raghunathanahalli-2 Microwatershed






Fig.6.6 Soil Available Sulphur map of Raghunathanahalli-2 Microwatershed

### 6.7 Available Boron

Maximum area of about 352 ha (80%) is low (<0.5 ppm) in available boron and distributed in the major part of the microwatershed. An area of about 84 ha (19%) is medium (0.5-1.0 ppm) in available boron and occur in the western and southwestern part of the microwatershed (Fig.6.7).

### 6.8 Available Iron

Available iron content is sufficient (>4.5 ppm) in a major area of about 296 ha (67%) and distributed in the major part of the microwatershed. An area of about 140 ha (32%) is deficient (<4.5 ppm) and distributed in the southwestern and western part of the microwatershed (Fig 6.8).

### 6.9 Available Manganese

Available manganese content is sufficient (>1.0 ppm) in the entire cultivated area of the microwatershed (Fig 6.9).

### 6.10 Available Copper

Available copper content is sufficient (>0.2 ppm) in the entire cultivated area of the microwatershed (Fig 6.10).

### 6.11 Available Zinc

Available zinc content is deficient (<0.6 ppm) in the entire cultivated area of the microwatershed (Fig 6.11).






Fig.6.7 Soil Available Boron map of Raghunathanahalli-2 Microwatershed




Fig.6.8 Soil Available Iron map of Raghunathanahalli-2 Microwatershed




Fig.6.9 Soil Available Manganese map of Raghunathanahalli-2 Microwatershed




Fig.6.10 Soil Available Copper map of Raghunathanahalli-2 Microwatershed




Fig.6.11 Soil Available Zinc map of Raghunathanahalli-2 Microwatershed

## LAND SUITABILITY FOR MAJOR CROPS

The soil and land resource units (soil phases) of Raghunathanahalli-2 Microwatershed were assessed for their suitability for growing food, fodder, fibre and other horticulture crops by following the procedure as outlined in FAO, 1976 and 1983. Crop requirements were developed for each of the crop from the available research data and also by referring to Naidu *et. al.* (2006) and Natarajan *et. al.* (2015). The soil and land characteristics (Table 7.1) were matched with the crop requirements (Tables 7.2 to 7.32) to arrive at the crop suitability. The soil and land characteristics table and crop requirements tables are given at the end of the chapter. In FAO land suitability classification, two orders are recognized. Order S-Suitable and Order N- Not suitable. The orders have Classes, subclasses and units. Order-S has three classes, Class S1-Highly Suitable, Class S2-Moderately Suitable and Class S3-Marginally Suitable. Order N has two classes, N1- Currently not Suitable and N2-Permanently not Suitable. There are no subclasses within the Class S1 as they will have very minor or no limitations for crop growth. Classes S2, S3, N1 and N2 are divided into subclasses based on the kinds of limitations encountered. The limitations that affect crop production are ‘c’ for erratic rainfall and its distribution and length of growing period (LGP), ‘e’ for erosion hazard, ‘r’ for rooting condition, ‘t’ for lighter or heavy texture, ‘g’ for gravelliness or stoniness, ‘n’ for nutrient availability, ‘l’ for topography, ‘m’ for moisture availability, ‘z’ for calcareousness and ‘w’ for drainage. These limitations are indicated as lower case letters to the class symbol. For example, moderately suitable lands with the limitations of soil depth and erosion are designated as S2re. For the microwatershed, the soil mapping units were evaluated and classified up to subclass level.

Using the above criteria, the soil map units of the microwatershed were evaluated and land suitability maps for 31 major agricultural and horticultural crops were generated. The detailed information on the kind of suitability of each of the soil phase for the crops assessed are given village/ survey number wise for the microwatershed in Appendix-III

### 7.1 Land Suitability for Sorghum (*Sorghum bicolor*)

Sorghum is one of the major food crop grown in Karnataka in an area of 10.47 lakh ha in Bijapur, Gulbarga, Raichur, Bidar, Belgaum, Dharwad, Bellary, Chitradurga, Mysore and Chamarajnar districts. The crop requirements for growing sorghum (Table 7.2) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and land a suitability map for growing sorghum was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure. 7.1.

An area of about 21 ha (5%) is highly suitable (Class S1) for growing sorghum and occur in the eastern and southern part of the microwatershed. Maximum area of about 296 ha (67%) is moderately suitable (Class S2) for growing sorghum and distributed in

the major part of the microwatershed. They have minor limitations of rooting depth, gravelliness, calcareousness, nutrient availability and texture. An area of about 53 ha (12%) is marginally suitable (Class S3) for growing sorghum and occur in the central, eastern and western part of the microwatershed with moderate limitations of rooting depth and calcareousness. Currently not suitable (Class N1) lands for growing sorghum cover an area of about 66 ha (15%) and distributed in the northeastern and eastern part of the microwatershed with severe limitations of gravelliness and rooting depth.




Fig. 7.1 Land Suitability map of Sorghum

## 7.2 Land Suitability for Maize (*Zea mays*)

Maize is one of the most important food crop grown in an area of 13.37 lakh ha in almost all the districts of the State. The crop requirements for growing maize (Table 7.3) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing maize was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.2.

Maximum area of about 300 ha (68%) is moderately suitable (Class S2) for growing maize and distributed in the major part of the microwatershed with minor limitations of texture, rooting depth, calcareousness and gravelliness. An area of about 69 ha (16%) is marginally suitable (Class S3) for growing maize and occur in the central, western, southwestern, southern and eastern part of the microwatershed with moderate limitations of texture, calcareousness and gravelliness. Currently not suitable (Class N1) lands for growing maize cover an area of about 66 ha (15%) and distributed in the



northeastern and eastern part of the microwatershed with severe limitations of gravelliness and rooting depth.




Fig. 7.2 Land Suitability map of Maize

### 7.3 Land Suitability for Bajra (*Pennisetum glaucum*)

Bajra is one of the major food crop grown in an area of 2.34 lakh ha in Karnataka in the northern districts. The crop requirements (Table 7.4) for growing bajra were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and land suitability map for growing bajra was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.3.

Maximum area of about 317 ha (72%) is moderately suitable (Class S2) for growing bajra and distributed in the major part of the microwatershed with minor limitations of texture, rooting depth, calcareousness and gravelliness. An area of about 53 ha (12%) is marginally suitable (Class S3) for growing bajra and distributed in the central, western and eastern part of the microwatershed with moderate limitations of rooting depth and calcareousness. Currently not suitable (Class N1) lands for growing bajra cover an area of about 66 ha (15%) and distributed in the northeastern and eastern part of the microwatershed with severe limitations of gravelliness and rooting depth.




Fig. 7.3 Land Suitability map of Bajra

#### 7.4 Land Suitability for Groundnut (*Arachis hypogaea*)

Groundnut is one of the major oilseed crop grown in an area of 6.54 lakh ha in Karnataka in most of the districts either as rainfed or irrigated crop. The crop requirements for growing groundnut (Table 7.5) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing groundnut was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.4.

An area of about 152 ha (35%) is moderately suitable (Class S2) for growing groundnut and distributed in the central, western, northwestern, northern and northeastern part of the microwatershed. They have minor limitation of calcareousness. Maximum area of about 218 ha (49%) is marginally suitable (Class S3) for growing groundnut and distributed in the central, northern, northeastern, eastern, southern, southwestern and western part of the microwatershed with moderate limitations of rooting depth, calcareousness and texture. Currently not suitable (Class N1) lands for growing groundnut cover an area of about 66 ha (15%) and distributed in the northeastern and eastern part of the microwatershed with severe limitations of gravelliness and rooting depth.






Fig. 7.4 Land Suitability map of Groundnut

### 7.5 Land Suitability for Sunflower (*Helianthus annus*)

Sunflower is one of the most important oilseed crop grown in an area of 3.56 lakh ha in the State in all the districts. The crop requirements for growing sunflower (Table 7.6) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing sunflower was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.5.

An area of about 14 ha (3%) is highly suitable (Class S1) for growing sunflower and distributed in the eastern and southern part of the microwatershed. Maximum area of about 246 ha (56%) is moderately suitable (Class S2) for growing sunflower and distributed in the major part of the microwatershed with minor limitations of gravelliness, rooting depth and calcareousness. An area of about 56 ha (13%) is marginally suitable (Class S3) for growing sunflower and occur in the northern, southwestern, southern and eastern part of the microwatershed with moderate limitations of calcareousness and rooting depth. An area of about 119 ha (27%) is currently not suitable (Class N1) for growing sunflower and occur in the central, western, eastern and northeastern part of the microwatershed with severe limitations of rooting depth, calcareousness and gravelliness.




Fig. 7.5 Land Suitability map of Sunflower

## 7.6 Land Suitability for Cotton (*Gossypium hirsutum*)

Cotton is one of the most important fibre crop grown in the state in about 8.75 lakh ha area in Raichur, Dharwad, Belgaum, Kalaburagi, Bijapur, Bidar, Bellary, Chitradurga and Chamarajnagar districts. The crop requirements for growing cotton (Table 7.7) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing cotton was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.6.

An area of about 21 ha (5%) is highly suitable (Class S1) for growing cotton and occur in the southern and eastern part of the microwatershed. Maximum area of about 297 ha (67%) is moderately suitable (Class S2) for growing cotton and distributed in the major part of the microwatershed with minor limitations of gravelliness, calcareousness and rooting depth. An area of about 53 ha (12%) is marginally suitable (Class S3) for growing cotton and occur in the central, western and eastern part of the microwatershed with moderate limitations of rooting depth and calcareousness. Currently not suitable (Class N1) lands for growing cotton cover an area of about 66 ha (15%) and distributed in the eastern and northeastern part of the microwatershed with severe limitations of gravelliness and rooting depth.




Fig. 7.6 Land Suitability map of Cotton

### 7.7 Land Suitability for Red gram (*Cajanus cajana*)

Red gram is one of the major pulse crop grown in an area of 7.28 lakh ha mainly in northern Karnataka in Bijapur, Kalaburagi, Raichur, Bidar, Belgaum, Dharwad and Bellary districts. The crop requirements for growing red gram (Table 7.8) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing red gram was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed is given in Figure 7.7.

Maximum area of about 210 ha (48%) is moderately suitable (Class S2) for growing red gram and occur in the western, northwestern, northern, northeastern, southern and eastern part of the microwatershed. They have minor limitations of texture, calcareousness and gravelliness. An area of about 107 ha (24%) is marginally suitable (Class S3) for growing red gram and distributed in the central, northern, western, southwestern and eastern part of the microwatershed with moderate limitations of rooting depth and calcareousness. An area of about 119 ha (27%) is currently not suitable (Class N1) for growing red gram and occur in the central, western, eastern and northeastern part of the microwatershed with severe limitations of rooting depth, calcareousness and gravelliness.




Fig. 7.7 Land Suitability map of Red gram

## 7.8 Land Suitability for Bengal gram (*Cicer aerativum*)

Bengal gram is one of the most important pulse crop grown in about 9.39 lakh ha area in Bijapur, Raichur, Kalaburagi, Dharwad, Belgaum and Bellary districts. The crop requirements for growing Bengal gram (Table 7.9) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing Bengal gram was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.8.

Highly suitable (Class S1) lands for growing bengal gram occur in an area of 21 ha (5%) and distributed in the southern and eastern part of the microwatershed. Maximum area of about 296 ha (67%) is moderately suitable (Class S2) for growing bengal gram and distributed in the major part of the microwatershed. They have minor limitations of rooting depth, gravelliness, calcareousness and texture. An area of about 53 ha (12%) is marginally suitable (Class S3) for growing bengal gram and occur in the central, western and eastern part of the microwatershed with moderate limitations of rooting depth and calcareousness. Currently not suitable (Class N1) lands for growing bengal gram cover an area of about 66 ha (15%) and distributed in the eastern and northeastern part of the microwatershed with severe limitations of gravelliness and rooting depth.




Fig. 7.8 Land Suitability map of Bengal gram

### 7.9 Land Suitability for Chilli (*Capsicum annum L*)

Chilli is one of the major spice crop grown in an area of 0.42 lakh ha in Karnataka State. The crop requirements for growing chilli (Table 7.10) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing chilli was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.9.

An area of about 169 ha (38%) is moderately suitable (Class S2) for growing chilli and distributed in the western, northwestern, northern and northeastern part of the microwatershed with minor limitations of gravelliness, texture, calcareousness and rooting depth. Major area of about 201 ha (45%) is marginally suitable (Class S3) for growing chilli and occur in the central, western, southwestern, southern and eastern part of the microwatershed with moderate limitations of texture, rooting depth and calcareousness. Currently not suitable (Class N1) lands for growing chilli cover an area of about 66 ha (15%) and distributed in the eastern and northeastern part of the microwatershed with severe limitations of gravelliness and rooting depth.






Fig. 7.9 Land Suitability map of Chilli

### 7.10 Land Suitability for Tomato (*Solanum lycopersicum*)

Tomato is one of the most important vegetable crop grown in an area of 0.65 lakh ha in almost all the districts of the State. The crop requirements (Table 7.11) for growing tomato were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing tomato was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.10.

An area of about 165 ha (37%) is moderately suitable (Class S2) for growing tomato and distributed in the western, northwestern, northern and northeastern part of the microwatershed with minor limitations of gravelliness, calcareousness and rooting depth. Major area of about 205 ha (46%) is marginally suitable (Class S3) for growing tomato and occur in the central, western, southwestern, southern, eastern and northeastern part of the microwatershed with moderate limitations of texture, rooting depth and calcareousness. Currently not suitable (Class N1) lands for growing tomato cover an area of about 66 ha (15%) and distributed in the eastern and northeastern part of the microwatershed with severe limitations of gravelliness and rooting depth.




Fig. 7.10 Land Suitability map of Tomato

### 7.11 Land Suitability for Brinjal (*Solanum melongena*)

Brinjal is one of the most important vegetable crop grown in the state. The crop requirements for growing brinjal (Table 7.12) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing Brinjal was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.11.

Maximum area of about 316 ha (72%) is moderately suitable (Class S2) for brinjal and distributed in the major part of the microwatershed. They have minor limitations of gravelliness, rooting depth, calcareousness and texture. An area about of 53 ha (12%) is marginally suitable (Class S3) and distributed in the central, western and eastern part of the microwatershed with moderate limitation of rooting depth. Currently not suitable (Class N1) lands for growing brinjal cover an area of about 66 ha (15%) and distributed in the eastern and northeastern part of the microwatershed with severe limitation of rooting depth.






Fig 7.11 Land Suitability map of Brinjal

## 7.12 Land Suitability for Onion (*Allium cepa L.*)

Onion is one of the most important vegetable crop grown in the state. The crop requirements for growing onion (Table 7.13) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing onion was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.12.

An area of about 165 ha (38%) is moderately suitable (Class S2) for onion and distributed in the western, northwestern, northern and northeastern part of the microwatershed. They have minor limitations of rooting depth, texture, calcareousness and gravelliness. Major area of about 204 ha (46%) is marginally suitable (Class S3) and distributed in the central, western, southwestern, southern, eastern and northeastern part of the microwatershed with moderate limitations of rooting depth, calcareousness and texture. Currently not suitable (Class N1) lands for growing onion cover an area of about 66 ha (15%) and distributed in the eastern and northeastern part of the microwatershed with severe limitation of rooting depth.




Fig 7.12 Land Suitability map of Onion

### 7.13 Land Suitability for Bhendi (*Abelmoschus esculentus*)

Bhendi is one of the most important vegetable crop grown in the state. The crop requirements for growing bhendi (Table 7.14) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing bhendi was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.13.

Maximum area of about 316 ha (72%) is moderately suitable (Class S2) for bhendi and distributed in the major part of the microwatershed. They have minor limitations of gravelliness, rooting depth, calcareousness and texture. An area about of 53 ha (12%) is marginally suitable (Class S3) and distributed in the central, western and eastern part of the microwatershed with moderate limitation of rooting depth. Currently not suitable (Class N1) lands for growing bhendi cover an area of about 66 ha (15%) and distributed in the eastern and northeastern part of the microwatershed with severe limitation of rooting depth.




Fig 7.13 Land Suitability map of Bhendi

#### 7.14 Land Suitability for Drumstick (*Moringa oleifera*)

Drumstick is one of the most important vegetable crop grown in 2403 ha area in the state. The crop requirements for growing drumstick (Table 7.15) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing drumstick was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.14.

Maximum area of 260 ha (59%) is moderately suitable (Class S2) for growing drumstick and distributed in the major part of the microwatershed with minor limitations of texture, rooting depth and calcareousness. An area of about 56 ha (13%) is marginally suitable (Class S3) for growing drumstick and occur in the southern, southwestern, eastern and northern part of the microwatershed with moderate limitations of rooting depth and calcareousness. An area of about 119 ha (27%) is currently not suitable (Class N1) for growing drumstick and occur in the central, western, eastern and northeastern part of the microwatershed with severe limitations of rooting depth, calcareousness and gravelliness.




Fig. 7.14 Land Suitability map of Drumstick

### 7.15 Land Suitability for Mulberry (*Morus nigra*)

Mulberry is the most important leaf crop grown for rearing silkworms in about 1.66 lakh ha in all the districts of the state. The crop requirements for growing mulberry (Table 7.16) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing mulberry was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.15.

Moderately suitable (Class S2) lands occupy a major area of about 240 ha (55%) and occur in the major part of the microwatershed. They have minor limitations of texture, calcareousness and gravelliness. Marginally suitable (Class S3) lands cover an area of about 76 ha (17%) and occur in the southern, southwestern, eastern, northeastern and northern part of the microwatershed. They have moderate limitations of texture, rooting depth and calcareousness. An area of about 119 ha (27%) is currently not suitable (Class N1) for growing mulberry and occur in the central, western, eastern and northeastern part of the microwatershed with severe limitations of rooting depth, calcareousness and gravelliness.




Fig. 7.15 Land Suitability map of Mulberry

### 7.16 Land Suitability for Mango (*Mangifera indica*)

Mango is one of the most important fruit crop grown in about 1.73 lakh ha in almost all the districts of the State. The crop requirements (Table 7.17) for growing mango were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing mango was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.16.

Marginally suitable (Class S3) lands cover a major area of about 260 ha (59%) and occur in the major part of the microwatershed. They have moderate limitations of texture, rooting depth and calcareousness. An area of about 175 ha (40%) is currently not suitable (Class N1) for growing mango and occur in the central, western, southwestern, eastern, northeastern and northern part of the microwatershed with severe limitations of gravelliness, calcareousness, texture and rooting depth.




Fig. 7.16 Land Suitability map of Mango

### 7.17 Land Suitability for Sapota (*Manilkara zapota*)

Sapota is one of the most important fruit crop grown in an area of about 29373 ha in almost all the districts of the state. The crop requirements (Table 7.18) for growing sapota were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing sapota was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.17.

An area of about 152 ha (35%) is moderately suitable (Class S2) for growing sapota and distributed in the western, northwestern, northern and northeastern part of the microwatershed with minor limitations of calcareousness and rooting depth. Major area of about 164 ha (37%) is marginally (Class S3) suitable for growing sapota and occur in the central, western, southwestern, southern, eastern, northeastern and northern part of the microwatershed with moderate limitations of texture, rooting depth and calcareousness. An area of about 119 ha (27%) is currently not suitable (Class N1) for growing sapota and occur in the central, western, eastern and northeastern part of the microwatershed with severe limitations of rooting depth, calcareousness and gravelliness.






Fig. 7.17 Land Suitability map of Sapota

### 7.18 Land Suitability for Pomegranate (*Punica granatum*)

Pomegranate is one of the commercially grown fruit crop in about 18488 ha in Karnataka mainly in Bijapur, Bagalkot, Koppal, Gadag and Chitradurga districts. The crop requirements for growing pomegranate (Table 7.19) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing pomegranate was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.18.

Maximum area of about 260 ha (59%) is moderately suitable (Class S2) for growing pomegranate and occur in the major part of the microwatershed with minor limitations of texture, rooting depth and calcareousness. Marginally suitable (Class S3) lands cover an area of about 53 ha (13%) and occur in the southern, southwestern, eastern and northern part of the microwatershed. They have moderate limitations of rooting depth and calcareousness. An area of about 119 ha (27%) is currently not suitable (Class N1) for growing pomegranate and occur in the central, western, eastern and northeastern part of the microwatershed with severe limitations of rooting depth, calcareousness and gravelliness.




Fig. 7.18 Land Suitability map of Pomegranate

### 7.19 Land Suitability for Guava (*Psidium guajava*)

Guava is one of the most important fruit crop grown in an area of about 6558 ha in almost all the districts of the state. The crop requirements (Table 7.20) for growing guava were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing guava was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.19.

An area of about 152 ha (35%) is moderately suitable (Class S2) for growing guava and distributed in the western, northwestern, northern and northeastern part of the microwatershed with minor limitations of rooting depth and calcareousness. Maximum area of about 165 ha (37%) is marginally (Class S3) suitable for growing guava and occur in the central, western, southwestern, southern, eastern, northeastern and northern part of the microwatershed with moderate limitations of texture, rooting depth and calcareousness. An area of about 119 ha (27%) is currently not suitable (Class N1) for growing guava and occur in the central, western, eastern and northeastern part of the microwatershed with severe limitations of rooting depth, texture and gravelliness.




Fig. 7.19 Land Suitability map of Guava

## 7.20 Land Suitability for Jackfruit (*Artocarpus heterophyllus*)

Jackfruit is one of the most important fruit crop grown in 5368 ha in all the districts of the state. The crop requirements (Table 7.21) for growing jackfruit were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing jackfruit was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in figure 7.20.

An area of about 152 ha (35%) is moderately suitable (Class S2) for growing jackfruit and distributed in the western, northwestern, northern and northeastern part of the microwatershed with minor limitations of calcareousness and rooting depth. Major area of about 165 ha (37%) is marginally (Class S3) suitable for growing jackfruit and occur in the central, western, southwestern, southern, eastern, northeastern and northern part of the microwatershed with moderate limitations of texture, rooting depth and calcareousness. An area of about 119 ha (27%) is currently not suitable (Class N1) for growing jackfruit and occur in the central, western, eastern and northeastern part of the microwatershed with severe limitations of rooting depth, texture and graveliness.




Fig. 7.20 Land Suitability map of Jackfruit

### 7.21 Land Suitability for Jamun (*Syzygium cumini*)

Jamun is one of the important fruit crop grown in almost all the districts of the state. The crop requirements (Table 7.22) for growing jamun were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing jamun was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.21.

An area of about 34 ha (8%) is moderately suitable (Class S2) for growing jamun and occur in the southwestern, southern, eastern and northeastern part of the microwatershed with minor limitations of texture, rooting depth and calcareousness. Marginally suitable (Class S3) lands cover a major area of about 281 ha (64%) and occur in the major part of the microwatershed. They have moderate limitations of rooting depth, texture and calcareousness. An area of about 119 ha (27%) is currently not suitable (Class N1) for growing jamun and occur in the central, western, eastern and northeastern part of the microwatershed with severe limitations of rooting depth, texture and gravelliness.




Fig. 7.21 Land Suitability map of Jamun

## 7.22 Land Suitability for Musambi (*Citrus limetta*)

Musambi is one of the most important fruit crop grown in an area of 5446 ha in almost all the districts of the state. The crop requirements (Table 7.23) for growing musambi were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing musambi was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.22.

Highly suitable (Class S1) lands for growing musambi cover an area of about 14 ha (3%) and occur in the southern and eastern part of the microwatershed. Maximum area of about 246 ha (56%) is moderately suitable (Class S2) for growing musambi and occur in the major part of the microwatershed with minor limitations of rooting depth, calcareousness and gravelliness. Marginally suitable (Class S3) lands cover an area of about 56 ha (13%) and occur in the southern, southwestern, eastern and northern part of the microwatershed. They have moderate limitations of rooting depth and calcareousness. An area of about 119 ha (27%) is currently not suitable (Class N1) for growing musambi and occur in the central, western, eastern and northeastern part of the microwatershed with severe limitations of rooting depth, calcareousness and gravelliness.




Fig. 7.22 Land Suitability map of Musambi

### 7.23 Land Suitability for Lime (*Citrus sp*)

Lime is one of the most important fruit crop grown in an area of 11752 ha in almost all the districts of the State. The crop requirements (Table 7.24) for growing lime (Table 7.15) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing lime was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.23.

An area of about 14 ha (3%) is highly suitable (Class S1) for growing lime and occur in the southern and eastern part of the microwatershed. Maximum area of about 246 ha (56%) is moderately suitable (Class S2) for growing lime and occur in the major part of the microwatershed with minor limitations of rooting depth, calcareousness and gravelliness. Marginally suitable (Class S3) lands cover an area of about 56 ha (13%) and occur in the southern, southwestern, eastern and northern part of the microwatershed. They have moderate limitations of rooting depth and calcareousness. An area of about 119 ha (27%) is currently not suitable (Class N1) for growing lime and occur in the central, western, eastern and northeastern part of the microwatershed with severe limitations of rooting depth, calcareousness and gravelliness.






Fig. 7.23 Land Suitability map of Lime

#### 7.24 Land Suitability for Cashew (*Anacardium occidentale*)

Cashew is one of the most important nut crop grown in an area of 7052 ha in almost all the districts of the State. The crop requirements (Table 7.25) for growing cashew were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing cashew was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.24.

Currently not suitable (Class N1) lands cover an entire cultivated area of the microwatershed with severe limitations of texture, rooting depth, calcareousness, nutrient availability and gravelliness.




Fig. 7.24 Land Suitability map of Cashew

## 7.25 Land Suitability for Custard Apple (*Annona reticulata*)

Custard apple is one of the most important fruit crop grown in 1426 ha in almost all the districts of the State. The crop requirements (Table 7.26) for growing custard apple were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing custard apple was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.25.

An area of about 21 ha (5%) is highly suitable (Class S1) for growing custard apple and occur in the southern and eastern part of the microwatershed. Major area of about 296 ha (67%) is moderately suitable (Class S2) for growing custard apple and occur in the major part of the microwatershed with minor limitations of rooting depth, calcareousness and gravelliness. Marginally suitable (Class S3) lands cover an area of about 53 ha (12%) for growing custard apple and occur in the central, western and eastern part of the microwatershed. They have moderate limitations of calcareousness and gravelliness. Currently not suitable (Class N1) lands cover an area of about 66 ha (15%) and distributed in the eastern and northeastern part of the microwatershed with severe limitations of rooting depth and gravelliness.




Fig. 7.25 Land Suitability map of Custard Apple

## 7.26 Land Suitability for Amla (*Phyllanthus emblica*)

Amla is one of the most important fruit and medicinal crop grown in an area of 151 ha and distributed in almost all the districts of the state. The crop requirements for (Table 7.27) growing amla were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing amla was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.26.

Major area of about 316 ha (72%) is moderately suitable (Class S2) for growing amla and occur in the major part of the microwatershed with minor limitations of rooting depth, calcareousness, texture and gravelliness. An area of about 53 ha (12%) is marginally suitable (Class S3) for growing amla and occur in the central, western and eastern part of the microwatershed with moderate limitations of calcareousness and texture. Currently not suitable (Class N1) lands cover an area of about 66 ha (15%) and distributed in the eastern and northeastern part of the microwatershed with severe limitations of rooting depth and gravelliness.




Fig. 7.26 Land Suitability map of Amla

### 7.27 Land Suitability for Tamarind (*Tamarindus indica*)

Tamarind is one of the most important spice crop grown in 14897 ha in all the districts of the state. The crop requirements (Table 7.28) for growing tamarind were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing tamarind was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.27.

An area of about 34 ha (8%) is moderately suitable (Class S2) for growing tamarind and occur in the southwestern, southern, eastern and northeastern part of the microwatershed with minor limitations of rooting depth, calcareousness and texture. Marginally suitable (Class S3) lands cover a major area of 226 ha (51%) for growing tamarind and occur in the major part of the microwatershed. They have moderate limitations of rooting depth and calcareousness. An area of about 174 ha (40%) is currently not suitable (Class N1) for growing tamarind and distributed in the central, western, southwestern, eastern, northeastern and northern part of the microwatershed. They have severe limitations of rooting depth, calcareousness and graveliness.




Fig. 7.27 Land Suitability map of Tamarind

## 7.28 Land Suitability for Marigold (*Tagetes erecta*)

Marigold is one of the most important flower crop grown in an area of 9108 ha in almost all the districts of the state. The crop requirements (Table 7.29) for growing marigold were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing marigold was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.28.

Maximum area of about 316 ha (72%) is moderately suitable (Class S2) for growing marigold and distributed in the major part of the microwatershed with minor limitations of texture, rooting depth, calcareousness and gravelliness. An area of about 53 ha (12%) is marginally suitable (Class S3) for growing marigold and occur in the central, western and eastern part of the microwatershed with moderate limitations of rooting depth and calcareousness. Currently not suitable (Class N1) lands cover an area of about 66 ha (15%) and distributed in the eastern and northeastern part of the microwatershed with severe limitations of rooting depth and gravelliness.




Fig. 7.28 Land Suitability map of Marigold

### 7.29 Land Suitability for Chrysanthemum (*Chrysanthemum indicum*)

Chrysanthemum is one of the most important flower crop grown in an area of 4978 ha in almost all the districts of the State. The crop requirements (Table 7.30) for growing chrysanthemum were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing chrysanthemum was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed is given in Figure 7.29.

Maximum area of about 316 ha (72%) is moderately suitable (Class S2) for growing chrysanthemum and distributed in the major part of the microwatershed with minor limitations of texture, rooting depth, calcareousness and gravelliness. An area of about 53 ha (12%) is marginally suitable (Class S3) for growing chrysanthemum and occur in the central, western and eastern part of the microwatershed with moderate limitations of rooting depth and calcareousness. Currently not suitable (Class N1) lands cover an area of about 66 ha (15%) and distributed in the eastern and northeastern part of the microwatershed with severe limitations of rooting depth and gravelliness.






Fig. 7.29 Land Suitability map of Chrysanthemum

### 7.30 Land Suitability for Jasmine (*Jasminum sp.*)

Jasmine is one of the most important flower crop grown in an area of 803 ha in almost all the districts of the State. The crop requirements (Table 7.31) for growing jasmine were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing jasmine was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.30.

Maximum area of about 208 ha (47%) is moderately suitable (Class S2) for growing jasmine and occur in the southern, southwestern, eastern, western, northwestern, northern and northeastern part of the microwatershed. They have minor limitations of gravelliness, calcareousness and rooting depth. An area of about 162 ha (37%) is marginally suitable (Class S3) for growing jasmine and occur in the central, western, southwestern, southern, eastern and northeastern part of the microwatershed. They have moderate limitations of rooting depth, texture and calcareousness. Currently not suitable (Class N1) lands cover an area of about 66 ha (15%) and distributed in the eastern and northeastern part of the microwatershed with severe limitations of rooting depth and gravelliness.




Fig. 7.30 Land Suitability map of Jasmine

### 7. 31 Land Suitability for Crossandra (*Crossandra infundibuliformis.*)

Crossandra is one of the most important flower crop grown in all the districts of the state. The crop requirements (Table 7.32) for growing crossandra were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing crossandra was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.31.

Maximum area of about 215 ha (49%) is moderately suitable (Class S2) for growing crossandra and occur in the central, southwestern, western, northwestern, northern and northeastern part of the microwatershed. They have minor limitations of gravelliness, calcareousness and rooting depth. An area of about 153 ha (35%) is marginally suitable (Class S3) for growing crossandra and occur in the central, western, southwestern, southern, eastern and northeastern part of the microwatershed. They have moderate limitations of rooting depth, texture and calcareousness. Currently not suitable (Class N1) lands cover an area of about 66 ha (15%) and distributed in the eastern and northeastern part of the microwatershed with severe limitations of rooting depth and gravelliness.




Fig. 7.31 Land Suitability map of Crossandra

**Table 7.1 Soil-Site Characteristics of Raghunathanahalli-2 Microwatershed**

Soil Map Units	Climate (P) (mm)	Growing period (Days)	Drainage Class	Soil depth (cm)	Soil texture		Gravelliness		AWC (mm/m)	Slope (%)	Erosion	pH	EC	ESP	CEC [Cmol (p+)kg-1]	BS (%)
					Surf-ace	Sub-surface	Sur- face	Sub- surface								
BGTmB2g2	662	<90	WD	<25	c	gc	35-60	>35	<50	1-3	Moderate	8.4	0.157	1.11	44.84	-
BMKiA1	662	<90	WD	75-100	sc	gsc-gc	<15	15-35	51-100	0-1	Slight	-	-	-	-	-
BMKiB1g1	662	<90	WD	75-100	sc	gsc-gc	15-35	15-35	51-100	1-3	Slight	-	-	-	-	-
BMKiB1g2	662	<90	WD	75-100	sc	gsc-gc	35-60	15-35	51-100	1-3	Slight	-	-	-	-	-
MTLmB2g1	662	<90	WD	25-50	c	gc	15-35	15-35	51-100	1-3	Moderate	8.27	0.202	0.69	36.64	-
MTLmB2g2	662	<90	WD	25-50	c	gc	35-60	15-35	51-100	1-3	Moderate	8.27	0.202	0.69	36.64	-
KSPiB1g2	662	<90	WD	50-75	sc	gscl	35-60	15-35	<50	1-3	Slight	-	-	-	-	-
KSPiB2g1	662	<90	WD	50-75	sc	gscl	15-35	15-35	<50	1-3	Moderate	-	-	-	-	-
RNKmB1	662	<90	MWD	50-75	c	c	<15	<15	51-100	1-3	Slight	8.86	0.483	6.78	37.00	-
RNKmB1g1	662	<90	MWD	50-75	c	c	15-35	<15	51-100	1-3	Slight	8.86	0.483	6.78	37.00	-
RNKmB2g1	662	<90	MWD	50-75	c	c	15-35	<15	51-100	1-3	Moderate	8.86	0.483	6.78	37.00	-
DRLmA1	662	<90	MWD	75-100	c	c	<15	<15	151-200	0-1	Slight	8.78	0.42	5.62	49.70	100
DRLmB2	662	<90	MWD	75-100	c	c	<15	<15	151-200	1-3	Moderate	8.78	0.42	5.62	49.70	100
NSPiB1g1	662	<90	MWD	75-100	sc	c	15-35	<15	101-150	1-3	Slight	9.16	0.615	8.60	51.09	-
GRHmB1	662	<90	MWD	100-150	c	c	<15	<15	>200	1-3	Slight	9.08	0.23	7.11	63.21	100
AWDmB2g1	662	<90	MWD	>150	c	c	15-35	<15	>200	1-3	Moderate	8.10	0.37	1.22	51.30	100
MLRmB1g2	662	<90	MWD	>150	c	c	35-60	10-20	>200	1-3	Slight	9.19	0.313	5.39	42.08	-

\*Symbols and abbreviations are according to Field Guide for LRI under Sujala-III

**Table 7.2 Land suitability criteria for Sorghum**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	26–30	30–34; 24–26	34–40; 20–24	>40; <20
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristics					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, c (red), c (black)	scl, cl	ls, sl	-
	pH	1:2.5	5.5-7.8	5.0-5.5 7.8-9.0	>9.0	-
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	10-15
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

**Table 7.3 Land suitability criteria for Maize**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	30-34	35-38 26-30	38-40 26-20	
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc	c (red), c (black)	ls, sl	-
	pH	1:2.5	5.5-7.8	5.0-5.5 7.8-9.0	>9.0	-
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	-
Erosion hazard	Slope	%	0-3	3-5	5-10	>10



**Table 7.4 Land suitability criteria for Bajra**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-32	33-38 24-27	39-40 20-23	<20
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm	500-750	400-500	200-400	<200
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	Sl, scl, cl,sc,c (red)	C (black)	ls	-
	pH	1:2.5	6.0-7.8	5.0-5.5 7.8-9.0	5.5-6.0 >9.0	
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	15-35	35-60	>60	
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	1-3	3-5	5-10	>10

**Table 7.5 Land suitability criteria for Groundnut**

Land use requirement			Rating			
Soil –site characteristics	Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)	
Climatic regime	Mean temperature in growing season	°C	24–33	22–24; 33–35	20–22; 35–40	<20; >40
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. Well drained	Poorly drained	Very Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl	sl,cl, sc	c (red), c (black), ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<35	35-60	>60	
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.6 Land suitability criteria for Sunflower**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	24–30	30–34; 20–24	34–38; 16–20	>38; <16
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	mod. Well drained	-	Poorly to very drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	cl, sc,c (red), c (black)	scl	ls, sl	-
	pH	1:2.5	6.5-7.8	7.8-8.4 5.5-6.5	8.4-9.0; 5.0-5.5	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.7 Land suitability criteria for Cotton**

Land use requirement		Rating				
Soil –site characteristics	Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)	
Climatic regime	Mean temperature in growing season	°C	22-32	>32	<19	-
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well to moderately well	Poorly drained/Somewhat excessively drained	-	very poorly/excessively drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, c (red,black)	cl	scl	ls, sl
	pH	1:2.5	6.5-7.8	7.8-8.4	5.5-6.5 8.4->9.0	<5.5
	CEC	C mol (p+)Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	50-100	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	<3	3-5	-	>5

**Table 7.8 Land suitability criteria for Red gram**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	30-35(G) 20-25(AV) 15-18 (F&PS) 35-40(M)	25-30(G) 20-25 (AV) 12-15 (F&PS) 30-35(M)	20-25(G) 15-20(AV) 10-12 (F&PS) 25-30(M)	< 20 <15 <10 <25
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. Well drained	Poorly drained	Very Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, c (red)	c (black),sl, scl, cl	ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-9.0	5.0-5.5 >9.0	-
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-50	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<1.0	1.0-2.0	>2.0	
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.9 Land suitability criteria for Bengal gram**

Land use requirement			Rating			
Soil –site characteristics	Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)	
Climatic regime	Mean temperature in growing season	°C	20–25	25–30; 15–20	30–35; 10–15	>35; <10
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. Well drained	Poorly drained	Very Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	C (black)	-	c (red), scl, cl, sc	ls, sl
	pH	1:2.5	6.0-7.8	5.0-6.0 7.8-9.0	>9.0	-
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	-
Erosion hazard	Slope	%	<3	3-5	5-10	>10



**Table 7.10 Land suitability criteria for Chilli**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	25-32	33-35 20-25	35-38 <20	>38
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc	c (black), sl	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.11 Land suitability criteria for Tomato**

Land use requirement		Rating				
Soil –site characteristics	Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)	
Climatic regime	Mean temperature in growing season	°C	25-28	29-32 20-24	15-19 33-36	<15 >36
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl, scl, cl, sc, c (red)	-	ls, c(black)	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.12 Land suitability criteria for Brinjal**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	Well drained	Moderately well drained	Poorly drained	V. Poorly drained
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class				
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl, scl, cl, sc c (red)	-	ls, c (black)	-
	pH	1:2.5	6.0-7.3	7.3-8.4 5.0-6.0	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	>60
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.13 Land suitability criteria for Onion**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	20-30	30-35	35-40	>40
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately /imperfectly	-	Poorly to V poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl,scl,cl,sc,c (red)	-	c (Black),ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<1.0	1.0-2.0	2.0-4.0	<4
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.14 Land suitability criteria for Bhendi**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	25-28	29-32 20-24	15-19 33-36	<15 >36
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Imperfectly drained	Poorly to very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl,sc, c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.15 Land suitability criteria for Drumstick**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, scl, cl, c (red)	sl, c (black)	ls	s
	pH	1:2.5	6.0-7.3	5.0-5.5 7.3-7.8	5.5-6.0 7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<35	35-60	60-80	>80
Soil toxicity	Salinity (EC saturation extract)	dS/m				
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-10	-	>10



**Table 7.16 Land suitability criteria for Mulberry**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	24–28	22–24; 28–32	32–38; 22–18	>38; <18
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V. Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, cl, scl	c (red)	c (black), sl, ls	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.8-8.4	7.3-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	0-35	35-60	60-80	>80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

**Note:** Suitability evaluation only for Mulberry leaf not for Silk worm rearing

**Table 7.17 Land suitability criteria for Mango**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-32	24-27 33-35	36-40	20-24
	Min temp. before flowering	°C	10-15	15-22	>22	-
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration	Days				
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	-	ls, sl, c (black)	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>150	100-150	75-100	<75
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.18 Land suitability criteria for Sapota**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-32	33-36 24-27	37-42 20-23	>42 <18
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	-	Poorly to very drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	sl	ls, c (black)	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.19 Land suitability criteria for Pomegranate**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	30-34	35-38 25-29	39-40 15-24	
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl,cl, sc, c (red)	c (black),sl	ls	-
	pH	1:2.5	5.5-7.8	7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.20 Land suitability criteria for Guava**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-32	33-36 24-27	37-42 20-23	
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	sl	c (black), ls	-
	pH	1:2.5	6.0-7.8	5.0-6.0	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.21 Land suitability criteria for Jackfruit**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. well	Poorly	V. Poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	-	sl, ls, c (black)	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	>60
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10-

**Table 7.22 Land suitability criteria for Jamun**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well	Mod. well	Poorly	V.Poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c(red)	sl, c (black)	ls	-
	pH	1:2.5	6.0-7.8	5.0-6.0	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>150	100-150	50-100	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	>60
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10



**Table 7.23 Land suitability criteria for Musambi**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-30	31-35 24-27	36-40 20-23	>40 <20
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately drained	poorly	Very poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c	sl	ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.24 Land suitability criteria for Lime**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-30	31-35 24-27	36-40 20-23	>40 <20
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately drained	poorly	Very poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c	sl	ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.25 Land suitability criteria for Cashew**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	32 to 34	28 to 32; 34 to 38	24 to 28; 38 to 40	<20; >40
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	-	sl, ls	c (black)
	pH	1:2.5	5.5-6.5	5.0-5.5 6.5-7.3	7.3-7.8	>7.8
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-10	>10	-

**Table 7.26 Land suitability criteria for Custard apple**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	Scl, cl, sc, c (red), c (black)	-	Sl, ls	-
	pH	1:2.5	6.0-7.3	5.5-6.0 7.3-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15-35	35-60	60-80	-
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	>5	-

**Table 7.27 Land suitability criteria for Amla**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod.well drained	Poorly drained	V. Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	c (black)	ls, sl	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15-35	35-60	60-80	-
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

**Table 7.28 Land suitability criteria for Tamarind**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod.well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl,sc, c (red)	sl, c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>150	100-150	75-100	<75
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

**Table 7.29 Land suitability criteria for Marigold**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	18-23	17-15 24-35	35-40 10-14	>40 <10
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl,scl, cl, sc, c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%				
Erosion hazard	Slope	%	<3	3-5	5-10	>10



**Table 7.30 Land suitability criteria for Chrysanthemum**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	18-23	17-15 24-35	35-40 10-14	>40 <10
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl,scl, cl, sc, c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%				
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.31 Land suitability criteria for Jasmine (irrigated)**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	18-23	17-15 24-35	35-40 10-14	-
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	sl	ls, c (black)	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%				
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.32 Land suitability criteria for Crossandra**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	-	Poorly to very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c(red)	sl,	c (black),ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%				
Erosion hazard	Slope	%	<3	3-5	5-10	>10

### 7.32 Land Management Units (LMUs)

The 17 soil map units identified in Raghunathanahalli-2 microwatershed have been grouped into 6 Land Management Units (LMUs) for the purpose of preparing a Proposed Crop Plan. Land Management Units are grouped based on the similarities in respect of the type of soil, the depth of the soil, the surface soil texture, gravel content, AWC, slope, erosion etc. and a Land Management Unit map (Fig.7.32) has been generated. These Land Management Units are expected to behave similarly for a given level of management.

The map units that have been grouped into 6 Land Management Units along with brief description of soil and site characteristics are given below.

LMUs	Mapping unit	Soil and site characteristics
1	425.AWDmB2g1 417.MLRmB1g2 371.GRHmB1 344.DRLmA1 350.DRLmB2 357.NSPiB1g1	Moderately deep to very deep, black calcareous clay soils, 0-3% slope, slight to moderate erosion, non-gravelly to very gravelly (<15-60%).
2	153.BMKiA1 154.BMKiB1g1 155.BMKiB1g2	Moderately deep, red calcareous clay soils, 0-3% slope, slight erosion, non-gravelly to very gravelly (<15-60%).
3	333.RNKmB1 334.RNKmB1g1 337.RNKmB2g1	Moderately shallow, black calcareous clay soils, 1-3% slope, slight to moderate erosion, non-gravelly to gravelly (<15-35%).
4	323.KSPiB1g2 325.KSPiB2g1	Moderately shallow, red calcareous sandy clay to sandy clay loam soils, 1-3% slope, slight to moderate erosion, gravelly to very gravelly (15-60%).
5	311.MTLmB2g1 312.MTLmB2g2	Shallow, black calcareous clay soils, 1-3% slope, moderate erosion, gravelly to very gravelly (15-60%).
6	11.BGTmB2g2	Very shallow, black gravelly clay soils, 1-3% slope, moderate erosion, very gravelly (35-60%).




Fig 7.32 Land Management Units map of Raghunathanahalli-2 microwatershed

### 7.33 Proposed Crop Plan for Raghunathanahalli-2 Microwatershed

After assessing the land suitability for the 31 crops, the proposed crop plan has been prepared for the 6 identified LMUs by considering only the highly (Class S1) and moderately (Class S2) suitable lands for each of the 31 crops. The resultant proposed crop plan is presented in Table 7.33.

**Table 7.33 Proposed Crop Plan for Raghunathanahalli-2 Microwatershed**

LMU	Soil Map Units	Survey Number	Soil and site characteristics	Field Crops	Horticulture Crops	Suitable Interventions
1	425.AWDmB2g1 417.MLRmB1g2 371.GRHmB1 344.DRLmA1 350.DRLmB2 357.NSPiB1g1	<b>Belagatti</b> : 7,8,9 ,10,11,12, 76 <b>Gattareddyhala</b> :119,176 ,177,181,183 <b>Hatti</b> : 12,13,14	Moderately deep to very deep, black calcareous clay soils, 0-3% slope, slight to moderate erosion, non-gravelly to very gravelly (<15-60%).	Maize, Sorghum, Sunflower, Cotton, Bengal gram, Safflower, Linseed, Bajra , Soybean	<b>Fruit crops:</b> Mango, Sapota, Pomegranate, Jamun, Lime, Musambi, Tamarind, Amla, Custard apple <b>Vegetables:</b> Drumstick, Chilli, Coriander, Tomato, Bhendi <b>Flowers:</b> Marigold, Chrysanthemum, Crossandra, Jasmine	Application of FYM, Biofertilizers and micronutrients, drip irrigation, mulching, suitable soil and water conservation practices
2	153.BMKiA1 154.BMKiB1g1 155.BMKiB1g2	<b>Gattareddyhala</b> :121,124 ,125,126,127,128,129,130,149,150,151,152,153,154,155,156,161,163,164,165,166,167,168,169,170,171,172,173,174,175,182,184,185,186,187,188,189,190,191,192, 193,194, 195	Moderately deep, red calcareous clay soils, 0-3% slope, slight erosion, non-gravelly to very gravelly (<15-60%).	Maize, Sorghum, Sunflower, Bajra, Finger millet, Groundnut, Red gram, Cowpea, Field bean, Castor	<b>Fruit crops:</b> Pomegranate, Guava, Sapota, Jackfruit, Tamarind, Lime, Musambi, Amla, Custard apple <b>Vegetable crops:</b> Drumstick, Tomato, Chilli, Brinjal, Onion, Curry leaves <b>Flower crops:</b> Marigold, Chrysanthemum, Jasmine	Drip irrigation, mulching, suitable soil and water conservation practices (Crescent Bunding with Catch Pit etc)
3	333.RNKmB1 334.RNKmB1g1 337.RNKmB2g1	<b>Belagatti</b> : 78,79,80,81,82,83 <b>Hatti</b> : 15,18 <b>Raghunathahalli</b> : 25,26	Moderately shallow, black calcareous clay soils, 1-3% slope, slight to moderate erosion, non-	Sorghum, Bajra, Bengal gram, Linseed, Safflower, Coriander	<b>Fruit crops:</b> Amla, Custard apple <b>Flower crops:</b> Marigold, Jasmine, Chrysanthemum	Application of FYM, Biofertilizers and micronutrients, drip irrigation, mulching, suitable soil and water

LMU	Soil Map Units	Survey Number	Soil and site characteristics	Field Crops	Horticulture Crops	Suitable Interventions
			gravelly to gravelly (<15-35%).			conservation practices
4	323.KSPiB1g2 325.KSPiB2g1	<b>Gattareddyhala:</b> 123,159,160,162,202	Moderately shallow, red calcareous sandy clay to sandy clay loam soils, 1-3% slope, slight to moderate erosion, gravelly to very gravelly (15-60%).	Sorghum, Groundnut, Bajra, Green gram, Black gram, Cowpea, Horse gram, Castor	<b>Fruit crops:</b> Lime, Musambi, Amla, Custard apple, Cashew <b>Flower crops:</b> Marigold, Chrysanthemum, Crossandra, Jasmine	Drip irrigation, Mulching, suitable soil and water conservation practices (Crescent Bunding with Catch Pit etc)
5	311.MTLmB2g1 312.MTLmB2g2	<b>Belagatti :</b> 77 <b>Gattareddyhala:</b> 147,148,178,179,180 <b>Hatti :</b> 16,17	Shallow, black calcareous clay soils, 1-3% slope, moderate erosion, gravelly to very gravelly (15-60%).	Bengal gram	<b>Agri-Silvi-Pasture:</b> Hybrid Napier, <i>Styloxanthes hamata</i> , <i>Styloxanthes scabra</i>	Use of short duration varieties, sowing across the slope
6	11.BGTmB2g2	<b>Gattareddyhala:</b> 131,132,133,134,135,136,137,138,139,140,141,142,143,144,145,146	Very shallow, black gravelly clay soils, 1-3% slope, moderate erosion, very gravelly (35-60%).	-	<b>Agri-Silvi-Pasture:</b> <i>Styloxanthes hamata</i> , <i>Styloxanthes scabra</i>	Suitable soil and water conservation practices



## SOIL HEALTH MANAGEMENT

### 8.1 Soil Health

Soil health is basic to plant health and plant health is basic to human and bovine health. Soil is fundamental to crop production. Without soil, no food could be produced nor would livestock be fed on a large scale. Because it is finite and fragile, soil is a precious resource that requires special care from its users.

Soil health or the capacity of the soil to function is critical to human survival. Soil health has been defined as: “the capacity of the soil to function as a living system without adverse effect on the ecosystem”. Healthy soils maintain a diverse community of soil organisms that help to form beneficial symbiotic associations with plant roots, recycle essential plant nutrients, improve soil structure with positive repercussions for soil, water and nutrient holding capacity and ultimately improve crop production and also contribute to mitigating climate change by maintaining or increasing its carbon content.

Functional interactions of soil biota with organic and inorganic components, air and water determine a soil’s potential to store and release nutrients, and water to plants and to promote and sustain plant growth. Thus, maintaining soil health is vital to crop production and conserve soil resource base for sustaining agriculture.

#### **The most important characteristics of a healthy soil are**

- Good soil tilth
- Sufficient soil depth
- Good water storage and good drainage
- Adequate supply, but not excess of nutrients
- Large population of beneficial organisms
- Small proportion of plant pathogens and insect pests
- Low weed pressure
- Free of chemicals and toxins that may harm the crop
- Resistance to degradation
- Resilience when unfavorable conditions occur

#### **Characteristics of Raghunathanahalli-2 Microwatershed**

- ❖ The soil phases with sizeable area identified in the microwatershed belonged to the soil series of Bhimanakunte (BMK) series occupies major area of 152 ha (34%) followed by Dambarahalli (DRL) 68 ha (15%), Belagatti (BGT) 66 ha (15%), Muttal (MTL) 53 ha (12%), Ravanaki (RNK) 42 ha (10%), Murlapur (MLR) 16 ha (4%), Gatareddihal (GRH) 14 ha (3%), Kyalapura (KSP) 13 ha (3%), Narasapura (NSP) 6 ha (1%) and Alawandi (AWD) 4 ha (1%).
- ❖ As per land capability classification, maximum area of about 312 ha (71%) in the microwatershed falls under good lands (Class II) with minor limitations of soil and

erosion. An area of about 57 ha (13%) is under moderately good lands (Class III) with severe limitations of soil and erosion. Fairly good lands (Class IV) cover an area of about 66 ha (15%) with very severe limitations of soil and erosion.

- ❖ On the basis of soil reaction, an entire cultivated area of the microwatershed falls under moderately alkaline to strongly alkaline (pH 7.8-9.0) in soil reaction.

### **Soil Health Management**

The following actions are required to improve the current land husbandry practices that provide a sound basis for the successful adoption of sustainable crop production system.

#### **Alkaline soils**

Moderately alkaline to strongly alkaline soils cover an entire cultivated area of the microwatershed.

1. Regular addition of organic manure, green manuring, green leaf manuring, crop residue incorporation and mulching needs to be taken up to improve the soil organic matter status.
2. Application of Biofertilizers (Azospirillum, Azotobacter, Rhizobium).
3. Application of 25% extra N and P (125 % RDN&P).
4. Application of ZnSO<sub>4</sub> – 12.5 kg/ha (once in three years).
5. Application of Boron – 5 kg/ha (once in three years).

Besides the above recommendations, the best transfer of technology options are also to be adopted.

#### **Soil Degradation**

Soil erosion is one of the major factor affecting the soil health in the microwatershed. An area of about 182 ha (41%) is suffering from moderate erosion. These areas need immediate soil and water conservation and, other land development and land husbandry practices for restoring soil health.

#### **Dissemination of Information and Communication of Benefits**

Any large scale implementation of soil health management requires that supporting information is made available widely, particularly through channels familiar to farmers and extension workers. Given the very high priority attached to soil health especially by the Central Government on issuing Soil-Health Cards to all the farmers, media outlets like Regional, State and National Newspapers, Radio and Dooradarshan programs in local languages but also modern information and communication technologies such as Cellular phones and the Internet, which can be much more effective in reaching the younger farmers.

## **Inputs for Net Planning (Saturation Plan) and Interventions needed**

Net planning in IWMP is focusing on preparation of

1. Soil and Water Conservation Treatment Plans for each plot or farm.
2. Productivity enhancement measures/ interventions for existing crops/livestock/other farm enterprises.
3. Diversification of farming mainly with perennial horticultural crops and livestock.
4. Improving livelihood opportunities and income generating activities.

In this connection, how various outputs of Sujala-III are of use in addressing these objectives of Net Planning are briefly presented below.

- ❖ **Soil Depth:** The depth of a soil decides the amount of moisture and nutrients it can hold, what crops can be taken up or not, depending on the rooting depth and the length of growing period available for raising any crop. Deeper the soil, better for a wide variety of crops. If sufficient depth is not available for growing deep rooted crops, either choose medium or short duration crops or deeper planting pits need to be opened and additional good quality soil brought from outside has to be filled into the planting pits.
- ❖ **Surface Soil Texture:** Lighter soil texture in the top soil means, better rain water infiltration, less run-off and soil moisture conservation, less capillary rise and less evaporation losses. Lighter surface textured soils are amenable to good soil tilth and are highly suitable for crops like groundnut, root vegetables (carrot, radish, potato etc) but not ideal for crops that need stagnant water like lowland paddy. Heavy textured soils are poor in water infiltration and percolation. They are prone for sheet erosion; such soils can be improved by sand mulching. The technology that is developed by the AICRP-Dryland Agriculture, Vijayapura, Karnataka can be adopted.
- ❖ **Gravelliness:** More gravel content is favorable for run-off harvesting but poor in soil moisture storage and nutrient availability. It is a significant parameter that decides the kind of crop to be raised.
- ❖ **Land Capability Classification:** The land capability map shows the areas suitable and not suitable for agriculture and the major constraints in each of the plot/survey number. Hence, one can decide what kind of enterprise is possible in each of these units. In general, erosion and soil are the major constraints in Raghunathanahalli-2 Microwatershed.
- ❖ **Organic Carbon:** The OC content is medium (0.5-0.75%) in an area of about 338 ha (77%) and low (<0.5%) in 98 ha (22%) area. These areas needs to be further improved by applying farmyard manure and rotating crops with cereals and legumes or mixed cropping.
- ❖ **Promoting Green Manuring:** Growing of green manuring crops costs Rs. 1250/ha (green manuring seeds) and about Rs. 2000/ha towards cultivation that totals to Rs. 3250/- per ha. On the other hand, application of organic manure @ 10 tons/ha costs Rs. 5000/ha. The practice needs to be continued for 2-3 years or more. Nitrogen

fertilizer needs to be supplemented by 25% in addition to the recommended level in 436 ha (99%) area where OC is low and medium (<0.5-0.75%). For example, for rainfed maize, recommended level is 50 kg N per ha and an additional 12 kg /ha needs to be applied for all the crops grown in these plots.

- ❖ **Available Phosphorus:** An area of about 18 ha (4%) is medium (23-57 kg/ha) and an area of about 418 ha (95%) is low (<23 kg/ha) in available phosphorus content. Hence all the plots, where available phosphorus is low and medium, for all the crops, 25% additional P-needs to be applied.
- ❖ **Available Potassium:** Entire cultivated area of the microwatershed is high (>337 kg/ha) in available potassium.
- ❖ **Available Sulphur:** Available sulphur is a very critical nutrient for oilseed crops, Available sulphur content is high (>20 ppm) in an area of about 303 ha (69%) and medium (10-20ppm) in 133 ha (30%) area of the microwatershed. Medium areas need to be applied with magnesium sulphate or gypsum or Factamphos (p) fertilizer (13% of sulphur) for 2-3 years for the deficiency to be corrected.
- ❖ **Available Boron:** An area of about 352 ha (80%) is low (<0.5 ppm) and an area of about 84 ha (19%) is medium (0.5-1.0 ppm) in available boron content. Low and medium (<0.5-1.0 ppm) areas need to be applied with sodium borate @ 10kg/ha as soil application or 0.2% borax as foliar spray to correct the deficiency.
- ❖ **Available Iron:** Available iron content is deficient (<4.5 ppm) in 140 ha (32%) area and sufficient (>4.5 ppm) in 296 ha (67%) area of the microwatershed. For deficient areas, iron sulphate @ 25 kg/ha needs to be applied for 2-3 years to correct the deficiency.
- ❖ **Available Manganese:** Entire cultivated area of the microwatershed is sufficient (>1.0 ppm) in the available manganese content.
- ❖ **Available Copper:** Entire cultivated area of the microwatershed is sufficient (>0.2 ppm) in the available copper content.
- ❖ **Available Zinc:** Entire cultivated area of the microwatershed is deficient (<0.6 ppm) in available zinc content. For deficient areas, application of zinc sulphate @ 25kg/ha is recommended.
- ❖ **Soil Alkalinity:** Entire cultivated area of the microwatershed has soils that are moderately alkaline to strongly alkaline. These areas need application of gypsum and wherever calcium is in excess, iron pyrites and element sulphur can be recommended. Management practices like treating repeatedly with good quality water to drain out the excess salts and provision of subsurface drainage and growing of salt tolerant crops like Casuarina, Acasia, Neem, Ber etc, are recommended.


- ❖ **Land Suitability for various crops:** Areas that are highly, moderately and marginally suitable and not suitable for growing various crops are indicated. Along with the suitability, various constraints that are limiting the productivity are also indicated. For example, in case of cotton, gravel content, rooting depth and salinity/alkalinity are the major constraints in various plots. With suitable management interventions, the productivity can be enhanced. In order to increase water holding capacity of light textured soils, growing of green manure crops and application of organic manure is recommended.



## SOIL AND WATER CONSERVATION TREATMENT PLAN

For preparing soil and water conservation treatment plan for Raghunathanahalli-2 Microwatershed, the land resource inventory database generated under Sujala-III project has been transformed as information through series of interpretative (thematic) maps using soil phase map as a base. The various thematic maps (1:7920 scale) generated were

- Soil depth
  - Surface soil texture
  - Available water capacity
  - Soil slope
  - Soil gravelliness
  - Land capability
  - Present land use and land cover
  - Crop suitability maps
  - Rainfall map
  - Hydrology
  - Water Resources
  - Socio-economic data
  - Contour plan with existing features- network of waterways, pothissa boundaries, cut up/ minor terraces etc.
  - Cadastral map (1:7920 scale)
  - Satellite imagery (1:7920 scale)
- Apart from these, Hand Level/ Hydro Marker/ Dumpy Level/ Total Station and Kathedars' List needs to be collected.



### Steps for Survey and Preparation of Treatment Plan

The boundaries of Land User Groups' and Survey No. boundaries are traced in the field.

- Naming of user groups and farmers
- Identification of arable and non arable lands
- Identification of drainage lines and gullies
- Identification of non treatable areas
- Identification of priority areas in the arable lands
- Treatment plan for arable lands
- Location of water harvesting and recharge structures


### 9.1 Treatment Plan

The treatment plan recommended for arable lands is briefly described below.




### 9.1.1 Arable Land Treatment

#### A. BUNDING

Steps for Survey and Preparation of Treatment Plan		<b>USER GROUP-1</b>  CLASSIFICATION OF GULLIES  ಕೊರಕಲಿನ ವರ್ಗೀಕರಣ  
Cadastral map (1:7920 scale) is enlarged to a scale of 1:2500 scale		
Existing network of waterways, pothissa boundaries, grass belts, natural drainage lines/ watercourse, cut ups/ terraces are marked on the cadastral map to the scale		
Drainage lines are demarcated into		
Small gullies	(up to 5 ha catchment)	
Medium gullies	(5-15 ha catchment)	
Ravines	(15-25 ha catchment) and	
Halla/Nala	(more than 25ha catchment)	

#### Measurement of Land Slope

Land slope is estimated or determined by the study and interpretation of contours or by measurement in the field using simple instruments like Hand Level or Hydro-marker.



Vertical and Horizontal intervals between bunds as recommended by the Watershed Development Department.

Slope percentage	Vertical interval (m)	Corresponding Horizontal Distance (m)
2 - 3%	0.6	24
3 - 4%	0.9	21
4 - 5%	0.9	21
5 - 6%	1.2	21
6 - 7%	1.2	21

**Note:** (i) The above intervals are maximum.

(ii) Considering the slope class and erosion status (A1... A= 0-1% slope, 1= slight erosion) the intervals have to be decided.

**Bund length recording:** Considering the contour plan and the existing grass belts/partitions, the bunds are aligned and lengths are measured.

### Section of the Bund

Bund section is decided considering the soil texture class and gravelliness class (bg0 .....b= loamy sand, g0 = <15% gravel). The recommended sections for different soils are given below.

#### Recommended Bund Section

Top width (m)	Base width (m)	Height (m)	Side slope (Z:1;H:V)	Cross section (sq m)	Soil Texture	Remarks
0.3	0.9	0.3	01:01	0.18	Sandy loam	Vegetative bund
0.3	1.2	0.3	1.5:1	0.225	Sandy clay	
0.3	1.2	0.5	0.9:1	0.375	Red gravelly soils	
0.3	1.2	0.6	0.75:1	0.45		
0.3	1.5	0.6	01:01	0.54	Red sandy loam	
0.3	2.1	0.6	1.5:1	0.72	Very shallow black clayey soils	
0.45	2	0.75	01:01	0.92		
0.45	2.4	0.75	1.3:1	1.07	Shallow black clayey soils	
0.6	3.1	0.7	1.78:1	1.29	Medium black clayey soils	
0.5	3	0.85	1.47:1	1.49		

### Formation of Trench cum Bund

Dimensions of the Borrow Pits/ Trenches to be excavated (machinery are decided considering the Bund Section).

Details of Borrow Pit dimensions are given below

#### TRENCH CUM BUND

IDEAL FOR HORTICULTURE CROPS

#### 'A' FRAME FOR INTERBUND MANAGEMENT

1. ಸಮವಾಹಕಳ ಉಳಿಸುವಿಕೆ
2. ಸಮವಾಹಕಳ ಬಿತ್ತರಣೆ/ನಾಟ

### Size of Borrow Pits/ Trench recommended for Trench cum Bund (by machinery)

Bund section	Bund length	Earth quantity	Pit				Berm (pit to pit)	Soil depth Class
			L(m)	W(m)	D(m)	Quantity (m <sup>3</sup> )		
0.375	6	2.25	5.85	0.85	0.45	2.24	0.15	Shallow
0.45	6	2.7	5.4	1.2	0.43	2.79	0.6	Shallow
0.45	6	2.7	5	0.85	0.65	2.76	1	Moderately Shallow
0.54	5.6	3.02	5.5	0.85	0.7	3.27	0.1	Moderately shallow
0.54	5.5	2.97	5	1.2	0.5	3	0.5	Shallow
0.72	6.2	4.46	6	1.2	0.7	5.04	0.2	Moderately shallow
0.72	5.2	3.74	5.1	0.85	0.9	3.9	0.1	Moderately deep

#### B. Waterways

- Existing waterways are marked on the cadastral map (1:7920 scale) and their dimensions are recorded.
- Considering the contour plan of the MWS, additional waterways/ modernization of the existing ones can be thought of.
- The design details are given in the Manual.

#### C. Farm Ponds

Waterways and the catchment area will give an indication on the size of the Farm Pond. Location of the pond can be decided based on the contour plan/ field condition and farmers' need/desire.

#### D. Diversion Channel

Existing EPT/ CPT are marked on the cadastral map. Looking to the need, these can be modernized or fresh diversion channel can be proposed and runoff from this can be stored in *Gokatte*/ Recharge ponds.

#### 9.1.2 Non-Arable Land Treatment

Depending on the gravelliness and crops preferred by the farmers, the concerned authorities can decide appropriate treatment plan. The recommended treatments may be Contour Trench, Staggered Trench, Crescent Bund, Boulder Bund or Pebble Bund.

### 9.1.3 Treatment of Natural Water Course/ Drainage Lines

- a) The cadastral map has to be updated as regards the network of drainage lines (gullies/ *nalas*/ *hallas*) and existing structures are marked to the scale and storage capacity of the existing water bodies are documented.
- b) The drainage line will be demarcated into Upper Reach, Middle Reach and Lower Reach.
- c) Considering the Catchment, *Nala* bed and bank conditions, suitable structures are decided.
- d) Number of storage structures (Check dam/ *Nala* bund/ Percolation tank) will be decided considering the commitments and available runoff in water budgeting and quality of water in the wells and site suitability.
- e) Detailed Leveling Survey using Dumpy Level / Total Station has to be carried out to arrive at the site-specific designs as shown in the Manual.
- f) The location of ground water recharge structures are decided by examining the lineaments and fracture zones from geological maps.
- g) Rainfall intensity data of the nearest Rain Gauge Station is considered for Hydrologic Designs.
- h) Silt load to the Storage/Recharge Structures is reduced by providing vegetative, boulder and earthen checks in the natural water course. Location and design details are given in the Manual.

### 9.2 Recommended Soil and Water Conservation Measures

The appropriate conservation structures best suited for each of the land parcel/ survey number (Appendix-I) are selected based on the slope per cent, severity of erosion, amount of rainfall, land use and soil type. The different kinds of conservation structures recommended are

1. Graded / Strengthening of Bunds
2. Trench cum Bunds (TCB)
3. Trench cum Bunds / Strengthening
4. Crescent Bunds

A map (Fig. 9.1) showing soil and water conservation plan with different kinds of structures recommended has been prepared which shows the spatial distribution and extent of area. An area of about 132 ha (30%) needs Trench cum Bunding, 253 ha (58%) needs Graded Bunding and 51 ha (11%) needs strengthening of existing bunds.

The conservation plan prepared may be presented to all the stakeholders including farmers and after considering their suggestions, the conservation plan for the microwatershed may be finalised in a participatory approach.




Fig. 9.1 Soil and Water Conservation Plan map of Raghunathanahalli-2 Microwatershed

### 9.3 Greening of Microwatershed

As part of the greening programme in the watersheds, it is envisaged to plant a variety of horticultural and other tree plants that are edible, economical and produce lot of biomass which helps to restore the ecological balance in the watersheds. The lands that are suitable for greening programme are non-arable lands (land capability classes V, VI VII and VIII) and also the lands that are not suitable or marginally suitable for growing annual and perennial crops. The method of planting these trees is given below.

It is recommended to open the pits during the 1st week of March along the contour and heap the dug-out soil on the lower side of the slope in order to harness the flowing water and facilitate weathering of soil in the pit. Exposure of soil in the pit also prevents spread of pests and diseases due to scorching sun rays. The pits should be filled with mixture of soil and organic manure during the second week of April and keep ready with sufficiently tall seedlings produced either in poly bags or in root trainer nurseries so that planting can be done during the 2nd or 3rd week of April depending on the rainfall.

The tree species suitable for the area considering rainfall, temperature and adaptability is listed below; waterlogged areas are recommended to be planted with species like Neral (*Syzgium cumini*) and Bamboo. Dry areas are to be planted with species like Honge, Bevu, Seetaphal etc.

<b>Dry Deciduous Species</b>			<b>Temp (°C)</b>	<b>Rainfall (mm)</b>
1.	Bevu	<i>Azadiracta indica</i>	21–32	400 –1,200
2.	Tapasi	<i>Holoptelia integrifolia</i>	20-30	500 - 1000
3.	Seetaphal	<i>Anona Squamosa</i>	20-40	400 - 1000
4.	Honge	<i>Pongamia pinnata</i>	20 -50	500–2,500
5.	Kamara	<i>Hardwickia binata</i>	25 -35	400 - 1000
6.	Bage	<i>Albezzia lebbek</i>	20 - 45	500 - 1000
7.	Ficus	<i>Ficus bengalensis</i>	20 - 50	500–2,500
8.	Sisso	<i>Dalbargia Sissoo</i>	20 - 50	500 -2000
9.	Ailanthus	<i>Ailanthus excelsa</i>	20 - 50	500 - 1000
10.	Hale	<i>Wrightia tinctoria</i>	25 - 45	500 - 1000
11.	Uded	<i>Steriospermum chelanoides</i>	25 - 45	500 -2000
12.	Dhupa	<i>Boswella Serrata</i>	20 - 40	500 - 2000
13.	Nelli	<i>Emblca Officinalis</i>	20 - 50	500 -1500
14.	Honne	<i>Pterocarpus marsupium</i>	20 - 40	500 - 2000
<b>Moist Deciduous Species</b>			<b>Temp (°C)</b>	<b>Rainfall (mm)</b>
15.	Teak	<i>Tectona grandis</i>	20 - 50	500-5000
16.	Nandi	<i>Legarstroemia lanceolata</i>	20 - 40	500 - 4000
17.	Honne	<i>Pterocarpus marsupium</i>	20 - 40	500 - 3000
18.	Mathi	<i>Terminalia alata</i>	20 -50	500 - 2000
19.	Shivane	<i>Gmelina arboria</i>	20 -50	500 -2000
20.	Kindal	<i>T.Paniculata</i>	20 - 40	500 - 1500
21.	Beete	<i>Dalbargia latifolia</i>	20 - 40	500 - 1500
22.	Tare	<i>T. belerica</i>	20 - 40	500 - 2000
23.	Bamboo	<i>Bambusa arundinasia</i>	20 - 40	500 - 2500
24.	Bamboo	<i>Dendrocalamus strictus</i>	20 – 40	500 – 2500
25.	Muthuga	<i>Butea monosperma</i>	20 - 40	400 - 1500
26.	Hippe	<i>Madhuca latifolia</i>	20 - 40	500 - 2000
27.	Sandal	<i>Santalum album</i>	20 - 50	400 - 1000
28.	Nelli	<i>Emblca officinalis</i>	20 - 40	500 - 2000
29.	Nerale	<i>Sizyzium cumini</i>	20 - 40	500 - 2000
30.	Dhaman	<i>Grevia tilifolia</i>	20 - 40	500 - 2000
31.	Kaval	<i>Careya arborea</i>	20 - 40	500 - 2000
32.	Harada	<i>Terminalia chebula</i>	20 - 40	500 - 2000



## References

1. FAO (1976) Framework for Land Evaluation, Food and Agriculture Organization, Rome.72 pp.
2. FAO (1983) Guidelines for Land Evaluation for Rainfed Agriculture, FAO, Rome, 237 pp.
3. IARI (1971) Soil Survey Manual, All India Soil and Land Use Survey Organization, IARI, New Delhi, 121 pp.
4. Katyal, J.C. and Rattan, R.K. (2003) Secondary and Micronutrients; Research Gap and future needs. Fert. News 48 (4); 9-20.
5. Naidu, L.G.K., Ramamurthy, V., Challa, O., Hegde, R. and Krishnan, P. (2006) Manual Soil Site Suitability Criteria for Major Crops, NBSS Publ. No. 129, NBSS &LUP, Nagpur, 118 pp.
6. Natarajan, A. and Dipak Sarkar (2010) Field Guide for Soil Survey, National Bureau of Soil Survey and Land Use Planning (ICAR), Nagpur, India.
7. Natarajan, A., Rajendra Hegde, Raj, J.N. and Shivananda Murthy, H.G. (2015) Implementation Manual for Sujala-III Project, Watershed Development Department, Bengaluru, Karnataka.
8. Sarma, V.A.K., Krishnan, P. and Budihal, S.L. (1987) Laboratory Manual, Tech. Bull. 23, NBSS &LUP, Nagpur.
9. Sehgal, J.L. (1990) Soil Resource Mapping of Different States of India; Why and How?, National Bureau of Soil Survey and Land Use Planning, Nagpur, 49 pp.
10. Shivaprasad, C.R., R.S. Reddy, J. Sehgal and M. Velayuthum (1998) Soils of Karnataka for Optimizing Land Use, NBSS Publ. No. 47b, NBSS & LUP, Nagpur, India.
11. Soil Survey Staff (2006) Keys to Soil Taxonomy, Tenth edition, U.S. Department of Agriculture/ NRCS, Washington DC, U.S.A.
12. Soil Survey Staff (2012) Soil Survey Manual, Handbook No. 18, USDA, Washington DC, USA.





**Appendix I**  
**Raghunathanahalli-2 (2M3a) Microwatershed**  
**Soil Phase Information**

Village	SY NO	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	WELLS	Land Capability	Conservation Plan
Hatti	12	1.4	GRHmB 1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Fallow land (Fl)	Not Available	IIs	Graded bunding
Hatti	13	1.05	GRHmB 1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Sunflower+Current fallow (Sf+Cf)	Not Available	IIs	Graded bunding
Hatti	14	0.9	GRHmB 1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Maize+Bengalgram (Mz+Bg)	Not Available	IIs	Graded bunding
Hatti	15	0.56	RNKmB 1	LMU-3	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	IIs	Graded bunding
Hatti	16	0.28	MTLmB 2g1	LMU-5	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIIs	Graded bunding
Hatti	17	1.22	MTLmB 2g1	LMU-5	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIIs	Graded bunding
Hatti	18	0.51	RNKmB 1	LMU-3	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	IIs	Graded bunding
NA	NA	94.67	DRLmB 2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIs	Graded bunding
Raghunathahalli	25	0.16	RNKmB 1	LMU-3	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Jowar (Jw)	Not Available	IIs	Graded bunding
Raghunathahalli	26	0.27	RNKmB 2g1	LMU-3	Moderately shallow (50-75 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Sunflower (Sf)	Not Available	IIs	Graded bunding
Belagatti	7	3.76	MLRmB 1g2	LMU-1	Very deep (>150 cm)	Clay	Very gravelly (35-60%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Jowar+Bengalgram (Jw+Bg)	Not Available	IIs	Graded bunding
Belagatti	8	3.05	MLRmB 1g2	LMU-1	Very deep (>150 cm)	Clay	Very gravelly (35-60%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Jowar+Fallow land (Jw+Fl)	1 Borewell	IIs	Graded bunding
Belagatti	9	4.14	NSPiB1 g1	LMU-1	Moderately deep (75-100 cm)	Sandy clay	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Slight	Current fallow+Fallow land (Cf+Fl)	Not Available	IIs	Graded bunding
Belagatti	10	3.78	NSPiB1 g1	LMU-1	Moderately deep (75-100 cm)	Sandy clay	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Slight	Jowar+Bengalgram (Jw+Bg)	Not Available	IIs	Graded bunding
Belagatti	11	2.45	GRHmB 1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Current fallow (Cf)	Not Available	IIs	Graded bunding
Belagatti	12	1.76	GRHmB 1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Fallow land (Fl)	Not Available	IIs	Graded bunding
Belagatti	76	0.8	MLRmB 1g2	LMU-1	Very deep (>150 cm)	Clay	Very gravelly (35-60%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Current fallow (Cf)	Not Available	IIs	Graded bunding
Belagatti	77	4.86	MTLmB 2g2	LMU-5	Shallow (25-50 cm)	Clay	Very gravelly (35-60%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Fallow land (Fl)	Not Available	IIIs	Graded bunding
Belagatti	78	4.84	RNKmB 1	LMU-3	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Fallow land (Fl)	Not Available	IIs	Graded bunding
Belagatti	79	5.76	RNKmB 1	LMU-3	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Currentfallow+Fallow land (Cf+Fl)	Not Available	IIs	Graded bunding

Village	SY NO	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	WELLS	Land Capability	Conservation Plan
Belagatti	80	3.27	RNKmB 1	LMU-3	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Current fallow+F allow land (Cf+Fl)	Not Available	IIs	Graded bunding
Belagatti	81	3.61	RNKmB 1g1	LMU-3	Moderately shallow (50-75 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Fallow land (Fl)	Not Available	IIs	Graded bunding
Belagatti	82	4.06	RNKmB 1g1	LMU-3	Moderately shallow (50-75 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Jowar+Maize (Jw+Mz)	Not Available	IIs	Graded bunding
Belagatti	83	17.68	RNKmB 1g1	LMU-3	Moderately shallow (50-75 cm)	Clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Fallow land+Current fallow (Fl+Cf)	Not Available	IIs	Graded bunding
Gattared dyhala	109	0.04	Waterbody	Others	Others	Others	Others	Others	Others	Others	Fallow land (Fl)	Not Available	Others	Others
Gattared dyhala	110	0.38	Waterbody	Others	Others	Others	Others	Others	Others	Others	Sunflower (Sf)	Not Available	Others	Others
Gattared dyhala	119	4.4	AWDm B2g1	LMU-1	Very deep (>150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar+Bengalgram (Jw+Bg)	Not Available	IIIe	Graded bunding
Gattared dyhala	121	0.16	BMKiB 1g2	LMU-2	Moderately deep (75-100 cm)	Sandy clay	Very gravelly (35-60%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Sunflower (Sf)	Not Available	IIs	Trench cum bunding
Gattared dyhala	123	2.85	KSPiB1 g2	LMU-4	Moderately shallow (50-75 cm)	Sandy clay	Very gravelly (35-60%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Slight	Groundnut+Bajra (Gn+Bj)	Not Available	IIs	Trench cum bunding
Gattared dyhala	124	2.61	BMKiB 1g2	LMU-2	Moderately deep (75-100 cm)	Sandy clay	Very gravelly (35-60%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Current fallow (Cf)	Not Available	IIs	Trench cum bunding
Gattared dyhala	125	4.67	BMKiB 1g2	LMU-2	Moderately deep (75-100 cm)	Sandy clay	Very gravelly (35-60%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Seasamum+Bajra+Maize (Sf+Bj+Mz)	Not Available	IIs	Trench cum bunding
Gattared dyhala	126	4.11	BMKiB 1g2	LMU-2	Moderately deep (75-100 cm)	Sandy clay	Very gravelly (35-60%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Redgram+Sunflower (Rg+Sf)	Not Available	IIs	Trench cum bunding
Gattared dyhala	127	4.76	BMKiB 1g2	LMU-2	Moderately deep (75-100 cm)	Sandy clay	Very gravelly (35-60%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Jowar (Jw)	Not Available	IIs	Trench cum bunding
Gattared dyhala	128	2.7	BMKiB 1g2	LMU-2	Moderately deep (75-100 cm)	Sandy clay	Very gravelly (35-60%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Current fallow (Cf)	Not Available	IIs	Trench cum bunding
Gattared dyhala	129	0.71	BMKiB 1g2	LMU-2	Moderately deep (75-100 cm)	Sandy clay	Very gravelly (35-60%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Sunflower (Sf)	Not Available	IIs	Trench cum bunding
Gattared dyhala	130	2.86	BMKiB 1g2	LMU-2	Moderately deep (75-100 cm)	Sandy clay	Very gravelly (35-60%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Sunflower (Sf)	Not Available	IIs	Trench cum bunding
Gattared dyhala	131	3.01	BGTmB 2g2	LMU-6	Very shallow (<25 cm)	Clay	Very gravelly (35-60%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Fallow land (Fl)	Not Available	IVes	Graded bunding
Gattared dyhala	132	3.63	BGTmB 2g2	LMU-6	Very shallow (<25 cm)	Clay	Very gravelly (35-60%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Fallow land+Current fallow (Fl+Cf)	1 Borewell	IVes	Graded bunding
Gattared dyhala	133	8.54	BGTmB 2g2	LMU-6	Very shallow (<25 cm)	Clay	Very gravelly (35-60%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow+Maize (Cf+Mz)	2 Borewell	IVes	Graded bunding
Gattared dyhala	134	1.81	BGTmB 2g2	LMU-6	Very shallow (<25 cm)	Clay	Very gravelly (35-60%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow (Cf)	Not Available	IVes	Graded bunding
Gattared dyhala	135	4.52	BGTmB 2g2	LMU-6	Very shallow (<25 cm)	Clay	Very gravelly (35-60%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	IVes	Graded bunding
Gattared dyhala	136	0.99	BGTmB 2g2	LMU-6	Very shallow (<25 cm)	Clay	Very gravelly (35-60%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	IVes	Graded bunding

Village	SY NO	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	WELLS	Land Capability	Conservation Plan
Gattared dyhala	137	2.18	BGTmB 2g2	LMU-6	Very shallow (<25 cm)	Clay	Very gravelly (35-60%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Fallow land (Fl)	Not Available	Ives	Graded bunding
Gattared dyhala	138	6.87	BGTmB 2g2	LMU-6	Very shallow (<25 cm)	Clay	Very gravelly (35-60%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow (Cf)	Not Available	Ives	Graded bunding
Gattared dyhala	139	4.26	BGTmB 2g2	LMU-6	Very shallow (<25 cm)	Clay	Very gravelly (35-60%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Fallow land (Fl)	Not Available	Ives	Graded bunding
Gattared dyhala	140	4.57	BGTmB 2g2	LMU-6	Very shallow (<25 cm)	Clay	Very gravelly (35-60%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow (Cf)	Not Available	Ives	Graded bunding
Gattared dyhala	141	4.8	BGTmB 2g2	LMU-6	Very shallow (<25 cm)	Clay	Very gravelly (35-60%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow (Cf)	Not Available	Ives	Graded bunding
Gattared dyhala	142	5.19	BGTmB 2g2	LMU-6	Very shallow (<25 cm)	Clay	Very gravelly (35-60%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Fallow land (Fl)	Not Available	Ives	Graded bunding
Gattared dyhala	143	1.03	BGTmB 2g2	LMU-6	Very shallow (<25 cm)	Clay	Very gravelly (35-60%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow (Cf)	Not Available	Ives	Graded bunding
Gattared dyhala	144	2.36	BGTmB 2g2	LMU-6	Very shallow (<25 cm)	Clay	Very gravelly (35-60%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Fallow land (Fl)	Not Available	Ives	Graded bunding
Gattared dyhala	145	6.35	BGTmB 2g2	LMU-6	Very shallow (<25 cm)	Clay	Very gravelly (35-60%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow+Fallow land (Cf+Fl)	Not Available	Ives	Graded bunding
Gattared dyhala	146	4.86	BGTmB 2g2	LMU-6	Very shallow (<25 cm)	Clay	Very gravelly (35-60%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Fallow land (Fl)	Not Available	Ives	Graded bunding
Gattared dyhala	147	4.88	MTLmB 2g2	LMU-5	Shallow (25-50 cm)	Clay	Very gravelly (35-60%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	IIes	Graded bunding
Gattared dyhala	148	4.92	MTLmB 2g2	LMU-5	Shallow (25-50 cm)	Clay	Very gravelly (35-60%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Bajra (Bj)	Not Available	IIes	Graded bunding
Gattared dyhala	149	7.66	BMKiB 1g2	LMU-2	Moderately deep (75-100 cm)	Sandy clay	Very gravelly (35-60%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Eucalyptus (Eu)	Not Available	IIs	Trench cum bunding
Gattared dyhala	150	3.42	BMKiB 1g1	LMU-2	Moderately deep (75-100 cm)	Sandy clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Maize+Bajra+Groundnut (Mz+Bj+Gn)	Not Available	IIs	Trench cum bunding
Gattared dyhala	151	3.09	BMKiB 1g1	LMU-2	Moderately deep (75-100 cm)	Sandy clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Groundnut+Maize+Bajra (Gn+Mz+Bj)	Not Available	IIs	Trench cum bunding
Gattared dyhala	152	3.8	BMKiB 1g1	LMU-2	Moderately deep (75-100 cm)	Sandy clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Drumstick+Current fallow (Ds+Cf)	Not Available	IIs	Trench cum bunding
Gattared dyhala	153	2.13	BMKiB 1g1	LMU-2	Moderately deep (75-100 cm)	Sandy clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Bajra (Bj)	Not Available	IIs	Trench cum bunding
Gattared dyhala	154	2.86	BMKiB 1g1	LMU-2	Moderately deep (75-100 cm)	Sandy clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Marigold (Mg)	1 Borewell	IIs	Trench cum bunding
Gattared dyhala	155	2.53	BMKiB 1g1	LMU-2	Moderately deep (75-100 cm)	Sandy clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Bajra (Bj)	Not Available	IIs	Trench cum bunding
Gattared dyhala	156	0.73	BMKiB 1g1	LMU-2	Moderately deep (75-100 cm)	Sandy clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Current fallow (Cf)	Not Available	IIs	Trench cum bunding
Gattared dyhala	159	1.23	KSPiB2 g1	LMU-4	Moderately shallow (50-75 cm)	Sandy clay	Gravelly (15-35%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Trench cum bunding
Gattared dyhala	160	1.49	KSPiB2 g1	LMU-4	Moderately shallow (50-75 cm)	Sandy clay	Gravelly (15-35%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIes	Trench cum bunding

Village	SY NO	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Graveliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	WELLS	Land Capability	Conservation Plan
Gattared dyhala	161	1.98	BMKiB 1g2	LMU-2	Moderately deep (75-100 cm)	Sandy clay	Very gravelly (35-60%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Current fallow (Cf)	Not Available	IIs	Trench cum bunding
Gattared dyhala	162	3.64	KSPiB2 g1	LMU-4	Moderately shallow (50-75 cm)	Sandy clay	Gravelly (15-35%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Bajra (Bj)	Not Available	IIs	Trench cum bunding
Gattared dyhala	163	1.47	BMKiA 1	LMU-2	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Nearly level (0-1%)	Slight	Bajra (Bj)	Not Available	IIs	Graded bunding
Gattared dyhala	164	2.88	BMKiA 1	LMU-2	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Nearly level (0-1%)	Slight	Sunflower (Sf)	Not Available	IIs	Graded bunding
Gattared dyhala	165	1.26	BMKiB 1g1	LMU-2	Moderately deep (75-100 cm)	Sandy clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Bajra (Bj)	Not Available	IIs	Trench cum bunding
Gattared dyhala	166	0.65	BMKiB 1g1	LMU-2	Moderately deep (75-100 cm)	Sandy clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Current fallow (Cf)	Not Available	IIs	Trench cum bunding
Gattared dyhala	167	0.53	BMKiB 1g1	LMU-2	Moderately deep (75-100 cm)	Sandy clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Current fallow (Cf)	Not Available	IIs	Trench cum bunding
Gattared dyhala	168	2.01	BMKiA 1	LMU-2	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Nearly level (0-1%)	Slight	Groundnut+Cotton (Gn+Ct)	Not Available	IIs	Graded bunding
Gattared dyhala	169	0.81	BMKiA 1	LMU-2	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Nearly level (0-1%)	Slight	Bajra (Bj)	Not Available	IIs	Graded bunding
Gattared dyhala	170	0.85	BMKiA 1	LMU-2	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Nearly level (0-1%)	Slight	Groundnut (Gn)	Not Available	IIs	Graded bunding
Gattared dyhala	171	1.38	BMKiA 1	LMU-2	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Nearly level (0-1%)	Slight	Maize (Mz)	Not Available	IIs	Graded bunding
Gattared dyhala	172	0.93	BMKiA 1	LMU-2	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Nearly level (0-1%)	Slight	Maize (Mz)	Not Available	IIs	Graded bunding
Gattared dyhala	173	2.28	BMKiA 1	LMU-2	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Nearly level (0-1%)	Slight	Maize (Mz)	Not Available	IIs	Graded bunding
Gattared dyhala	174	6.02	BMKiA 1	LMU-2	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Nearly level (0-1%)	Slight	Maize (Mz)	1 Borewell	IIs	Graded bunding
Gattared dyhala	175	7.64	BMKiA 1	LMU-2	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Nearly level (0-1%)	Slight	Fallow land (Fl)	1 Borewell	IIs	Graded bunding
Gattared dyhala	176	3.07	DRLmB 2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Fallow land (Fl)	Not Available	IIs	Graded bunding
Gattared dyhala	177	2.62	DRLmB 2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow+Sunflower (Cf+Sf)	Not Available	IIs	Graded bunding
Gattared dyhala	178	0.95	MTLmB 2g2	LMU-5	Shallow (25-50 cm)	Clay	Very gravelly (35-60%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Fallow land (Fl)	Not Available	IIIs	Graded bunding
Gattared dyhala	179	7.13	MTLmB 2g2	LMU-5	Shallow (25-50 cm)	Clay	Very gravelly (35-60%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Fallow land (Fl)	Not Available	IIIs	Graded bunding
Gattared dyhala	180	4.58	MTLmB 2g2	LMU-5	Shallow (25-50 cm)	Clay	Very gravelly (35-60%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Fallow land (Fl)	Not Available	IIIs	Graded bunding
Gattared dyhala	181	9.92	DRLmA 1	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Nearly level (0-1%)	Slight	Redgram+Sunflower+Cotton (Rg+Sf+Ct)	Not Available	IIs	Graded bunding
Gattared dyhala	182	1.44	BMKiB 1g1	LMU-2	Moderately deep (75-100 cm)	Sandy clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Redgram (Rg)	Not Available	IIs	Trench cum bunding
Gattared dyhala	183	10.69	DRLmA 1	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Nearly level (0-1%)	Slight	Eucalyptus+Jowar+Sunflower (Eu+Jw+Sf)	Not Available	IIs	Graded bunding

Village	SY NO	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Graveliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	WELLS	Land Capability	Conservation Plan
Gattared dyhala	184	1.39	BMKiB 1g1	LMU-2	Moderately deep (75-100 cm)	Sandy clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Redgram (Rg)	Not Available	IIs	Trench cum bunding
Gattared dyhala	185	2.27	BMKiB 1g1	LMU-2	Moderately deep (75-100 cm)	Sandy clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Redgram (Rg)	Not Available	IIs	Trench cum bunding
Gattared dyhala	186	4.4	BMKiB 1g1	LMU-2	Moderately deep (75-100 cm)	Sandy clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Bajra+Current fallow (Bj+Cf)	Not Available	IIs	Trench cum bunding
Gattared dyhala	187	4.1	BMKiB 1g1	LMU-2	Moderately deep (75-100 cm)	Sandy clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	IIs	Trench cum bunding
Gattared dyhala	188	1.05	BMKiB 1g1	LMU-2	Moderately deep (75-100 cm)	Sandy clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Groundnut (Gn)	Not Available	IIs	Trench cum bunding
Gattared dyhala	189	6.77	BMKiB 1g2	LMU-2	Moderately deep (75-100 cm)	Sandy clay	Very gravelly (35-60%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Current fallow+Maize (Cf+Mz)	1 Borewell	IIs	Trench cum bunding
Gattared dyhala	190	3.55	BMKiB 1g1	LMU-2	Moderately deep (75-100 cm)	Sandy clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Groundnut (Gn)	Not Available	IIs	Trench cum bunding
Gattared dyhala	191	5.36	BMKiB 1g1	LMU-2	Moderately deep (75-100 cm)	Sandy clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Redgram+Onion (Rg+On)	1 Borewell	IIs	Trench cum bunding
Gattared dyhala	192	1.03	BMKiB 1g2	LMU-2	Moderately deep (75-100 cm)	Sandy clay	Very gravelly (35-60%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Fallow land (Fl)	Not Available	IIs	Trench cum bunding
Gattared dyhala	193	1.3	BMKiB 1g1	LMU-2	Moderately deep (75-100 cm)	Sandy clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Fallow land (Fl)	Not Available	IIs	Trench cum bunding
Gattared dyhala	194	1.32	BMKiB 1g1	LMU-2	Moderately deep (75-100 cm)	Sandy clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Onion (On)	Not Available	IIs	Trench cum bunding
Gattared dyhala	195	0.62	BMKiB 1g2	LMU-2	Moderately deep (75-100 cm)	Sandy clay	Very gravelly (35-60%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Redgram+Sunflower+Bajra (Rg+Sf+Bj)	Not Available	IIs	Trench cum bunding
Gattared dyhala	202	0.0002	KSPiB2 g1	LMU-4	Moderately shallow (50-75 cm)	Sandy clay	Gravelly (15-35%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow (Cf)	Not Available	IIs	Trench cum bunding















Village	Survey Number	Soil Reaction	Salinity	Organic Carbon	Available Phosphorus	Available Potassium	Available Sulphur	Available Boron	Available Iron	Available Manganese	Available Copper	Available Zinc
Gattaredd yhala	191	Strongly alkaline (pH 8.4 - 9.0)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Low (< 23 kg/ha)	High (> 337 kg/ha)	High (> 20 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Gattaredd yhala	192	Strongly alkaline (pH 8.4 - 9.0)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Low (< 23 kg/ha)	High (> 337 kg/ha)	High (> 20 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Gattaredd yhala	193	Strongly alkaline (pH 8.4 - 9.0)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Low (< 23 kg/ha)	High (> 337 kg/ha)	High (> 20 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Gattaredd yhala	194	Strongly alkaline (pH 8.4 - 9.0)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Low (< 23 kg/ha)	High (> 337 kg/ha)	High (> 20 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Gattaredd yhala	195	Strongly alkaline (pH 8.4 - 9.0)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Low (< 23 kg/ha)	High (> 337 kg/ha)	High (> 20 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Gattaredd yhala	202	Strongly alkaline (pH 8.4 - 9.0)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	High (> 337 kg/ha)	High (> 20 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)



**Appendix III**  
**Raghunathanahalli-2 (2M3a) Microwatershed**  
**Soil Suitability Information**

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Jasmine	Bhendi	Brinjal	Crossandra	Drumstick	Mulberry	Onion
Hatti	12	S3t	S2t	S3t	S1	S3t	S1	S2r	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S3t	S2t	S2t	S2t	S3t	S2t	S2t	S3t	S2t	S2t	S3t	
Hatti	13	S3t	S2t	S3t	S1	S3t	S1	S2r	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S3t	S2t	S2t	S2t	S2t	S3t	S2t	S2t	S3t	S2t	S2t	S3t
Hatti	14	S3t	S2t	S3t	S1	S3t	S1	S2r	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S3t	S2t	S2t	S2t	S2t	S3t	S2t	S2t	S3t	S2t	S2t	S3t
Hatti	15	N1rz	S2tz	S3rz	S2rz	S3tz	S2rz	N1rz	S3rz	S2rz	S3rz	S3rz	S2rz	S3tz	S2rz	N1tz	S3tz	S3rz	S3tz	S3tz	S3tz	S2rz	S2rz	S3rz	S2tz	S2rz	S2rt	S2rt	S3rz	S3rz	S3rz	S3tz
Hatti	16	N1rt	S3tz	N1rz	S3rz	N1rt	S3rz	N1rz	N1rz	S3rz	N1rz	N1rz	S3tz	N1rt	S3zg	N1rt	N1rt	N1rz	S3tz	S3rz	S3rz	S3rz	S3rz	N1rz	S3rz	S3rz	S3r	S3r	S3rz	N1rz	N1rz	S3rt
Hatti	17	N1rt	S3tz	N1rz	S3rz	N1rt	S3rz	N1rz	N1rz	S3rz	N1rz	N1rz	S3tz	N1rt	S3zg	N1rt	N1rt	N1rz	S3tz	S3rz	S3rz	S3rz	S3rz	N1rz	S3rz	S3rz	S3r	S3r	S3rz	N1rz	N1rz	S3rt
Hatti	18	N1rz	S2tz	S3rz	S2rz	S3tz	S2rz	N1rz	S3rz	S2rz	S3rz	S3rz	S2rz	S3tz	S2rz	N1tz	S3tz	S3rz	S3tz	S3tz	S3tz	S2rz	S2rz	S3rz	S2tz	S2rz	S2rt	S2rt	S3rz	S3rz	S3rz	S3tz
NA	NA	S3rz	S2tz	S3tz	S2nz	S3tz	S2rz	S3rz	S2rz	S2rz	S2rz	S3rz	S2tz	S3tz	S2z	N1tz	S3rz	S2rz	S3tz	S3tz	S3tz	S2tz	S2tz	S2rt	S2tz	S3tz	S2tz	S2tz	S2z	S2rz	S2tz	S3tz
Raghunat hahalli	25	N1rz	S2tz	S3rz	S2rz	S3tz	S2rz	N1rz	S3rz	S2rz	S3rz	S3rz	S2rz	S3tz	S2rz	N1tz	S3tz	S3rz	S3tz	S3tz	S3tz	S2rz	S2rz	S3rz	S2tz	S2rz	S2rt	S2rt	S3rz	S3rz	S3rz	S3tz
Raghunat hahalli	26	N1rz	S2tz	S3rz	S2rz	S3tz	S2rz	N1rz	S3rz	S2rz	S3rz	S3rz	S2rz	S3tz	S2rz	N1tz	S3tz	S3rz	S3tz	S3tz	S3tz	S2rz	S2rz	S3rz	S2tz	S2rz	S2rt	S2rt	S3rz	S3rz	S3rz	S3tz
Belagatti	7	S3tz	S3g	S3tz	S2zg	S3tz	S2zg	S2tz	S2zg	S2gz	S2zg	S2tz	S2tz	S3tz	S2gz	N1tz	S2tz	S2zg	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S3tz	S3tz
Belagatti	8	S3tz	S3g	S3tz	S2zg	S3tz	S2zg	S2tz	S2zg	S2gz	S2zg	S2tz	S2tz	S3tz	S2gz	N1tz	S2tz	S2zg	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S3tz	S3tz
Belagatti	9	S3rt	S2tz	S3rt	S1	S3t	S1	S3r	S2rg	S1	S2rg	S2gt	S2gt	S3t	S1	N1t	S3rt	S2rg	S3t	S3t	S3t	S2tg	S2tg	S2rt	S2t	S3t	S2tz	S2tz	S3t	S2rt	S2gt	S3tz
Belagatti	10	S3rt	S2tz	S3rt	S1	S3t	S1	S3r	S2rg	S1	S2rg	S2gt	S2gt	S3t	S1	N1t	S3rt	S2rg	S3t	S3t	S3t	S2tg	S2tg	S2rt	S2t	S3t	S2tz	S2tz	S3t	S2rt	S2gt	S3tz
Belagatti	11	S3t	S2t	S3t	S1	S3t	S1	S2r	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S3t	S2t	S2t	S2t	S2t	S3t	S2t	S2t	S3t	S2t	S2t	S3t
Belagatti	12	S3t	S2t	S3t	S1	S3t	S1	S2r	S1	S1	S1	S2t	S2t	S3t	S1	N1t	S2rt	S1	S3t	S3t	S3t	S2t	S2t	S2t	S2t	S3t	S2t	S2t	S3t	S2t	S2t	S3t
Belagatti	76	S3tz	S3g	S3tz	S2zg	S3tz	S2zg	S2tz	S2zg	S2gz	S2zg	S2tz	S2tz	S3tz	S2gz	N1tz	S2tz	S2zg	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S3tz	S3tz
Belagatti	77	N1rt	S3tz	N1rz	S3rz	N1rt	S3rz	N1rz	N1rz	S3rz	N1rz	N1rz	S3tz	N1rt	S3zg	N1rt	N1rt	N1rz	S3tz	S3rz	S3rz	S3rz	S3rz	N1rz	S3rz	S3rz	S3r	S3r	S3rz	N1rz	N1rz	S3rt
Belagatti	78	N1rz	S2tz	S3rz	S2rz	S3tz	S2rz	N1rz	S3rz	S2rz	S3rz	S3rz	S2rz	S3tz	S2rz	N1tz	S3tz	S3rz	S3tz	S3tz	S3tz	S2rz	S2rz	S3rz	S2tz	S2rz	S2rt	S2rt	S3rz	S3rz	S3rz	S3tz
Belagatti	79	N1rz	S2tz	S3rz	S2rz	S3tz	S2rz	N1rz	S3rz	S2rz	S3rz	S3rz	S2rz	S3tz	S2rz	N1tz	S3tz	S3rz	S3tz	S3tz	S3tz	S2rz	S2rz	S3rz	S2tz	S2rz	S2rt	S2rt	S3rz	S3rz	S3rz	S3tz
Belagatti	80	N1rz	S2tz	S3rz	S2rz	S3tz	S2rz	N1rz	S3rz	S2rz	S3rz	S3rz	S2rz	S3tz	S2rz	N1tz	S3tz	S3rz	S3tz	S3tz	S3tz	S2rz	S2rz	S3rz	S2tz	S2rz	S2rt	S2rt	S3rz	S3rz	S3rz	S3tz
Belagatti	81	N1rz	S2tz	S3rz	S2rz	S3tz	S2rz	N1rz	S3rz	S2rz	S3rz	S3rz	S2rz	S3tz	S2rz	N1tz	S3tz	S3rz	S3tz	S3tz	S3tz	S2rz	S2rz	S3rz	S2tz	S2rz	S2rt	S2rt	S3rz	S3rz	S3rz	S3tz
Belagatti	82	N1rz	S2tz	S3rz	S2rz	S3tz	S2rz	N1rz	S3rz	S2rz	S3rz	S3rz	S2rz	S3tz	S2rz	N1tz	S3tz	S3rz	S3tz	S3tz	S3tz	S2rz	S2rz	S3rz	S2tz	S2rz	S2rt	S2rt	S3rz	S3rz	S3rz	S3tz

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Jasmine	Bhendi	Brinjal	Crossandra	Drumstick	Mulberry	Onion	
Belagatti	83	N1rz	S2tz	S3rz	S2rz	S3tz	S2rz	N1rz	S3rz	S2rz	S3rz	S3rz	S2rz	S3tz	S2rz	N1tz	S3tz	S3rz	S3tz	S3tz	S3tz	S2rz	S2rz	S3rz	S2tz	S2rz	S2rt	S2rt	S3rz	S3rz	S3rz	S3tz	
Gattared dyhala	109	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs
Gattared dyhala	110	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs
Gattared dyhala	119	S3tz	S2tz	S3tz	S2z	S3tz	S2zg	S2tz	S2z	S2gz	S2z	S2gt	S2tz	S3tz	S2z	N1tz	S2tz	S2z	S3tz	S2tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S3tz	S3tz	
Gattared dyhala	121	S3rz	S2z	S2rz	S2tz	S2rz	S2z	S3rz	S2rz	S2tz	S2rz	S2z	S2z	S2rz	S2z	N1tz	S3rz	S2rz	S2z	S2z	S2z	S2z	S2z	S2rz	S2z	S2z	S2tz	S2z	S2z	S2rz	S2z	S2tz	
Gattared dyhala	123	N1rz	S2rz	S3rz	S2rz	S3rz	S2rz	N1rz	S3rz	S2rz	S3rz	S3rz	S2rz	S3rz	S2rz	N1n	S3rz	S3rz	S3rt	S2rz	S2rz	S2rz	S2rz	S3rz	S2rz	S2rz	S2rg	S2rg	S2rz	S3rz	S3rz	S2rg	
Gattared dyhala	124	S3rz	S2z	S2rz	S2tz	S2rz	S2z	S3rz	S2rz	S2tz	S2rz	S2z	S2z	S2rz	S2z	N1tz	S3rz	S2rz	S2z	S2z	S2z	S2z	S2z	S2rz	S2z	S2z	S2tz	S2z	S2z	S2rz	S2z	S2tz	
Gattared dyhala	125	S3rz	S2z	S2rz	S2tz	S2rz	S2z	S3rz	S2rz	S2tz	S2rz	S2z	S2z	S2rz	S2z	N1tz	S3rz	S2rz	S2z	S2z	S2z	S2z	S2z	S2rz	S2z	S2z	S2tz	S2z	S2z	S2rz	S2z	S2tz	
Gattared dyhala	126	S3rz	S2z	S2rz	S2tz	S2rz	S2z	S3rz	S2rz	S2tz	S2rz	S2z	S2z	S2rz	S2z	N1tz	S3rz	S2rz	S2z	S2z	S2z	S2z	S2z	S2rz	S2z	S2z	S2tz	S2z	S2z	S2rz	S2z	S2tz	
Gattared dyhala	127	S3rz	S2z	S2rz	S2tz	S2rz	S2z	S3rz	S2rz	S2tz	S2rz	S2z	S2z	S2rz	S2z	N1tz	S3rz	S2rz	S2z	S2z	S2z	S2z	S2z	S2rz	S2z	S2z	S2tz	S2z	S2z	S2rz	S2z	S2tz	
Gattared dyhala	128	S3rz	S2z	S2rz	S2tz	S2rz	S2z	S3rz	S2rz	S2tz	S2rz	S2z	S2z	S2rz	S2z	N1tz	S3rz	S2rz	S2z	S2z	S2z	S2z	S2z	S2rz	S2z	S2z	S2tz	S2z	S2z	S2rz	S2z	S2tz	
Gattared dyhala	129	S3rz	S2z	S2rz	S2tz	S2rz	S2z	S3rz	S2rz	S2tz	S2rz	S2z	S2z	S2rz	S2z	N1tz	S3rz	S2rz	S2z	S2z	S2z	S2z	S2z	S2rz	S2z	S2z	S2tz	S2z	S2z	S2rz	S2z	S2tz	
Gattared dyhala	130	S3rz	S2z	S2rz	S2tz	S2rz	S2z	S3rz	S2rz	S2tz	S2rz	S2z	S2z	S2rz	S2z	N1tz	S3rz	S2rz	S2z	S2z	S2z	S2z	S2z	S2rz	S2z	S2z	S2tz	S2z	S2z	S2rz	S2z	S2tz	
Gattared dyhala	131	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1r	N1r	N1rg	N1rg	N1rg	N1r
Gattared dyhala	132	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1r	N1r	N1rg	N1rg	N1rg	N1r
Gattared dyhala	133	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1r	N1r	N1rg	N1rg	N1rg	N1r	
Gattared dyhala	134	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1r	N1r	N1rg	N1rg	N1rg	N1r	
Gattared dyhala	135	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1r	N1r	N1rg	N1rg	N1rg	N1r	
Gattared dyhala	136	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1r	N1r	N1rg	N1rg	N1rg	N1r	
Gattared dyhala	137	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1r	N1r	N1rg	N1rg	N1rg	N1r	
Gattared dyhala	138	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1r	N1r	N1rg	N1rg	N1rg	N1r	

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Jasmine	Bhendi	Brinjal	Crossandra	Drumstick	Mulberry	Onion	
Gattared dyhala	139	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1r	N1r	N1rg	N1rg	N1rg	N1r	
Gattared dyhala	140	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1r	N1r	N1rg	N1rg	N1rg	N1r	
Gattared dyhala	141	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1r	N1r	N1rg	N1rg	N1rg	N1r	
Gattared dyhala	142	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1r	N1r	N1rg	N1rg	N1rg	N1r	
Gattared dyhala	143	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1r	N1r	N1rg	N1rg	N1rg	N1r	
Gattared dyhala	144	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1r	N1r	N1rg	N1rg	N1rg	N1r	
Gattared dyhala	145	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1r	N1r	N1rg	N1rg	N1rg	N1r	
Gattared dyhala	146	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1rg	N1r	N1r	N1rg	N1rg	N1rg	N1r	
Gattared dyhala	147	N1rt	S3tz	N1rz	S3rz	N1rt	S3rz	N1rz	N1rz	S3rz	N1rz	N1rz	S3tz	N1rt	S3zg	N1rt	N1rt	N1rz	S3tz	S3rz	S3rz	S3rz	S3rz	N1rz	S3rz	S3rz	S3r	S3r	S3rz	N1rz	N1rz	S3rt	
Gattared dyhala	148	N1rt	S3tz	N1rz	S3rz	N1rt	S3rz	N1rz	N1rz	S3rz	N1rz	N1rz	S3tz	N1rt	S3zg	N1rt	N1rt	N1rz	S3tz	S3rz	S3rz	S3rz	S3rz	N1rz	S3rz	S3rz	S3r	S3r	S3rz	N1rz	N1rz	S3rt	
Gattared dyhala	149	S3rz	S2z	S2rz	S2tz	S2rz	S2z	S3rz	S2rz	S2tz	S2rz	S2z	S2z	S2rz	S2z	N1tz	S3rz	S2rz	S2z	S2z	S2z	S2z	S2z	S2z	S2rz	S2z	S2z	S2tz	S2z	S2z	S2rz	S2z	S2tz
Gattared dyhala	150	S3rz	S2zg	S2rz	S2tz	S2rz	S2zg	S3rz	S2rz	S2tz	S2rz	S2zg	S2z	S2rz	S2z	N1tz	S3rz	S2rz	S2z	S2zg	S2zg	S2zg	S2zg	S2rz	S2zg	S2zg	S2tz	S2z	S2zg	S2rz	S2z	S2tz	
Gattared dyhala	151	S3rz	S2zg	S2rz	S2tz	S2rz	S2zg	S3rz	S2rz	S2tz	S2rz	S2zg	S2z	S2rz	S2z	N1tz	S3rz	S2rz	S2z	S2zg	S2zg	S2zg	S2zg	S2rz	S2zg	S2zg	S2tz	S2z	S2zg	S2rz	S2z	S2tz	
Gattared dyhala	152	S3rz	S2zg	S2rz	S2tz	S2rz	S2zg	S3rz	S2rz	S2tz	S2rz	S2zg	S2z	S2rz	S2z	N1tz	S3rz	S2rz	S2z	S2zg	S2zg	S2zg	S2zg	S2rz	S2zg	S2zg	S2tz	S2z	S2zg	S2rz	S2z	S2tz	
Gattared dyhala	153	S3rz	S2zg	S2rz	S2tz	S2rz	S2zg	S3rz	S2rz	S2tz	S2rz	S2zg	S2z	S2rz	S2z	N1tz	S3rz	S2rz	S2z	S2zg	S2zg	S2zg	S2zg	S2rz	S2zg	S2zg	S2tz	S2z	S2zg	S2rz	S2z	S2tz	
Gattared dyhala	154	S3rz	S2zg	S2rz	S2tz	S2rz	S2zg	S3rz	S2rz	S2tz	S2rz	S2zg	S2z	S2rz	S2z	N1tz	S3rz	S2rz	S2z	S2zg	S2zg	S2zg	S2zg	S2rz	S2zg	S2zg	S2tz	S2z	S2zg	S2rz	S2z	S2tz	
Gattared dyhala	155	S3rz	S2zg	S2rz	S2tz	S2rz	S2zg	S3rz	S2rz	S2tz	S2rz	S2zg	S2z	S2rz	S2z	N1tz	S3rz	S2rz	S2z	S2zg	S2zg	S2zg	S2zg	S2rz	S2zg	S2zg	S2tz	S2z	S2zg	S2rz	S2z	S2tz	
Gattared dyhala	156	S3rz	S2zg	S2rz	S2tz	S2rz	S2zg	S3rz	S2rz	S2tz	S2rz	S2zg	S2z	S2rz	S2z	N1tz	S3rz	S2rz	S2z	S2zg	S2zg	S2zg	S2zg	S2rz	S2zg	S2zg	S2tz	S2z	S2zg	S2rz	S2z	S2tz	
Gattared dyhala	159	N1rz	S2rz	S3rz	S2rz	S3rz	S2rz	N1rz	S3rz	S2rz	S3rz	S3rz	S2rz	S3rz	S2rz	N1n	S3rz	S3rz	S3rt	S2rz	S2rz	S2rz	S2rz	S3rz	S2rz	S2rz	S2r	S2r	S2rz	S3rz	S3rz	S2r	
Gattared dyhala	160	N1rz	S2rz	S3rz	S2rz	S3rz	S2rz	N1rz	S3rz	S2rz	S3rz	S3rz	S2rz	S3rz	S2rz	N1n	S3rz	S3rz	S3rt	S2rz	S2rz	S2rz	S2rz	S3rz	S2rz	S2rz	S2r	S2r	S2rz	S3rz	S3rz	S2r	
Gattared dyhala	161	S3rz	S2z	S2rz	S2tz	S2rz	S2z	S3rz	S2rz	S2tz	S2rz	S2z	S2z	S2rz	S2z	N1tz	S3rz	S2rz	S2z	S2z	S2z	S2z	S2z	S2rz	S2z	S2z	S2tz	S2z	S2z	S2rz	S2z	S2tz	



Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Jasmine	Bhendi	Brinjal	Crossandra	Drumstick	Mulberry	Onion
Gattared dyhala	162	N1rz	S2rz	S3rz	S2rz	S3rz	S2rz	N1rz	S3rz	S2rz	S3rz	S3rz	S2rz	S3rz	S2rz	N1n	S3rz	S3rz	S3rt	S2rz	S2rz	S2rz	S2rz	S3rz	S2rz	S2rz	S2r	S2r	S2rz	S3rz	S3rz	S2r
Gattared dyhala	163	S3rz	S2z	S2rz	S2tz	S2rz	S2z	S3rz	S2rz	S2tz	S2rz	S2z	S2z	S2rz	S2z	N1tz	S3rz	S2rz	S2z	S2z	S2z	S2z	S2z	S2rz	S2z	S2z	S2tz	S2z	S2z	S2rz	S2z	S2tz
Gattared dyhala	164	S3rz	S2z	S2rz	S2tz	S2rz	S2z	S3rz	S2rz	S2tz	S2rz	S2z	S2z	S2rz	S2z	N1tz	S3rz	S2rz	S2z	S2z	S2z	S2z	S2z	S2rz	S2z	S2z	S2tz	S2z	S2z	S2rz	S2z	S2tz
Gattared dyhala	165	S3rz	S2zg	S2rz	S2tz	S2rz	S2zg	S3rz	S2rz	S2tz	S2rz	S2zg	S2z	S2rz	S2z	N1tz	S3rz	S2rz	S2z	S2zg	S2zg	S2zg	S2zg	S2rz	S2zg	S2zg	S2tz	S2z	S2zg	S2rz	S2z	S2tz
Gattared dyhala	166	S3rz	S2zg	S2rz	S2tz	S2rz	S2zg	S3rz	S2rz	S2tz	S2rz	S2zg	S2z	S2rz	S2z	N1tz	S3rz	S2rz	S2z	S2zg	S2zg	S2zg	S2zg	S2rz	S2zg	S2zg	S2tz	S2z	S2zg	S2rz	S2z	S2tz
Gattared dyhala	167	S3rz	S2zg	S2rz	S2tz	S2rz	S2zg	S3rz	S2rz	S2tz	S2rz	S2zg	S2z	S2rz	S2z	N1tz	S3rz	S2rz	S2z	S2zg	S2zg	S2zg	S2zg	S2rz	S2zg	S2zg	S2tz	S2z	S2zg	S2rz	S2z	S2tz
Gattared dyhala	168	S3rz	S2z	S2rz	S2tz	S2rz	S2z	S3rz	S2rz	S2tz	S2rz	S2z	S2z	S2rz	S2z	N1tz	S3rz	S2rz	S2z	S2z	S2z	S2z	S2z	S2rz	S2z	S2z	S2tz	S2z	S2z	S2rz	S2z	S2tz
Gattared dyhala	169	S3rz	S2z	S2rz	S2tz	S2rz	S2z	S3rz	S2rz	S2tz	S2rz	S2z	S2z	S2rz	S2z	N1tz	S3rz	S2rz	S2z	S2z	S2z	S2z	S2z	S2rz	S2z	S2z	S2tz	S2z	S2z	S2rz	S2z	S2tz
Gattared dyhala	170	S3rz	S2z	S2rz	S2tz	S2rz	S2z	S3rz	S2rz	S2tz	S2rz	S2z	S2z	S2rz	S2z	N1tz	S3rz	S2rz	S2z	S2z	S2z	S2z	S2z	S2rz	S2z	S2z	S2tz	S2z	S2z	S2rz	S2z	S2tz
Gattared dyhala	171	S3rz	S2z	S2rz	S2tz	S2rz	S2z	S3rz	S2rz	S2tz	S2rz	S2z	S2z	S2rz	S2z	N1tz	S3rz	S2rz	S2z	S2z	S2z	S2z	S2z	S2rz	S2z	S2z	S2tz	S2z	S2z	S2rz	S2z	S2tz
Gattared dyhala	172	S3rz	S2z	S2rz	S2tz	S2rz	S2z	S3rz	S2rz	S2tz	S2rz	S2z	S2z	S2rz	S2z	N1tz	S3rz	S2rz	S2z	S2z	S2z	S2z	S2z	S2rz	S2z	S2z	S2tz	S2z	S2z	S2rz	S2z	S2tz
Gattared dyhala	173	S3rz	S2z	S2rz	S2tz	S2rz	S2z	S3rz	S2rz	S2tz	S2rz	S2z	S2z	S2rz	S2z	N1tz	S3rz	S2rz	S2z	S2z	S2z	S2z	S2z	S2rz	S2z	S2z	S2tz	S2z	S2z	S2rz	S2z	S2tz
Gattared dyhala	174	S3rz	S2z	S2rz	S2tz	S2rz	S2z	S3rz	S2rz	S2tz	S2rz	S2z	S2z	S2rz	S2z	N1tz	S3rz	S2rz	S2z	S2z	S2z	S2z	S2z	S2rz	S2z	S2z	S2tz	S2z	S2z	S2rz	S2z	S2tz
Gattared dyhala	175	S3rz	S2z	S2rz	S2tz	S2rz	S2z	S3rz	S2rz	S2tz	S2rz	S2z	S2z	S2rz	S2z	N1tz	S3rz	S2rz	S2z	S2z	S2z	S2z	S2z	S2rz	S2z	S2z	S2tz	S2z	S2z	S2rz	S2z	S2tz
Gattared dyhala	176	S3rz	S2tz	S3tz	S2nz	S3tz	S2rz	S3rz	S2rz	S2rz	S2rz	S3rz	S2tz	S3tz	S2z	N1tz	S3rz	S2rz	S3tz	S3tz	S3tz	S2tz	S2tz	S2rt	S2tz	S3tz	S2tz	S2tz	S2z	S2rz	S2tz	S3tz
Gattared dyhala	177	S3rz	S2tz	S3tz	S2nz	S3tz	S2rz	S3rz	S2rz	S2rz	S2rz	S3rz	S2tz	S3tz	S2z	N1tz	S3rz	S2rz	S3tz	S3tz	S3tz	S2tz	S2tz	S2rt	S2tz	S3tz	S2tz	S2tz	S2z	S2rz	S2tz	S3tz
Gattared dyhala	178	N1rt	S3tz	N1rz	S3rz	N1rt	S3rz	N1rz	N1rz	S3rz	N1rz	N1rz	S3tz	N1rt	S3zg	N1rt	N1rt	N1rz	S3tz	S3rz	S3rz	S3rz	S3rz	N1rz	S3rz	S3rz	S3r	S3r	S3rz	N1rz	N1rz	S3rt
Gattared dyhala	179	N1rt	S3tz	N1rz	S3rz	N1rt	S3rz	N1rz	N1rz	S3rz	N1rz	N1rz	S3tz	N1rt	S3zg	N1rt	N1rt	N1rz	S3tz	S3rz	S3rz	S3rz	S3rz	N1rz	S3rz	S3rz	S3r	S3r	S3rz	N1rz	N1rz	S3rt
Gattared dyhala	180	N1rt	S3tz	N1rz	S3rz	N1rt	S3rz	N1rz	N1rz	S3rz	N1rz	N1rz	S3tz	N1rt	S3zg	N1rt	N1rt	N1rz	S3tz	S3rz	S3rz	S3rz	S3rz	N1rz	S3rz	S3rz	S3r	S3r	S3rz	N1rz	N1rz	S3rt
Gattared dyhala	181	S3rz	S2tz	S3tz	S2nz	S3tz	S2rz	S3rz	S2rz	S2z	S2rz	S2tz	S2tz	S3tz	S2z	N1tz	S3rz	S2rz	S3tz	S3tz	S3tz	S2tz	S2tz	S2rt	S2tz	S3tz	S2tz	S2tz	S3tz	S2rz	S2tz	S3tz
Gattared dyhala	182	S3rz	S2zg	S2rz	S2tz	S2rz	S2zg	S3rz	S2rz	S2tz	S2rz	S2zg	S2z	S2rz	S2z	N1tz	S3rz	S2rz	S2z	S2zg	S2zg	S2zg	S2zg	S2rz	S2zg	S2zg	S2tz	S2z	S2zg	S2rz	S2z	S2tz

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Jasmine	Bhendi	Brinjal	Crossandra	Drumstick	Mulberry	Onion
Gattared dyhala	183	S3rz	S2tz	S3tz	S2nz	S3tz	S2rz	S3rz	S2rz	S2z	S2rz	S2tz	S2tz	S3tz	S2z	N1tz	S3rz	S2rz	S3tz	S3tz	S3tz	S2tz	S2tz	S2rt	S2tz	S3tz	S2tz	S2tz	S3tz	S2rz	S2tz	S3tz
Gattared dyhala	184	S3rz	S2zg	S2rz	S2tz	S2rz	S2zg	S3rz	S2rz	S2tz	S2rz	S2zg	S2z	S2rz	S2z	N1tz	S3rz	S2rz	S2z	S2zg	S2zg	S2zg	S2zg	S2rz	S2zg	S2zg	S2tz	S2z	S2zg	S2rz	S2z	S2tz
Gattared dyhala	185	S3rz	S2zg	S2rz	S2tz	S2rz	S2zg	S3rz	S2rz	S2tz	S2rz	S2zg	S2z	S2rz	S2z	N1tz	S3rz	S2rz	S2z	S2zg	S2zg	S2zg	S2zg	S2rz	S2zg	S2zg	S2tz	S2z	S2zg	S2rz	S2z	S2tz
Gattared dyhala	186	S3rz	S2zg	S2rz	S2tz	S2rz	S2zg	S3rz	S2rz	S2tz	S2rz	S2zg	S2z	S2rz	S2z	N1tz	S3rz	S2rz	S2z	S2zg	S2zg	S2zg	S2zg	S2rz	S2zg	S2zg	S2tz	S2z	S2zg	S2rz	S2z	S2tz
Gattared dyhala	187	S3rz	S2zg	S2rz	S2tz	S2rz	S2zg	S3rz	S2rz	S2tz	S2rz	S2zg	S2z	S2rz	S2z	N1tz	S3rz	S2rz	S2z	S2zg	S2zg	S2zg	S2zg	S2rz	S2zg	S2zg	S2tz	S2z	S2zg	S2rz	S2z	S2tz
Gattared dyhala	188	S3rz	S2zg	S2rz	S2tz	S2rz	S2zg	S3rz	S2rz	S2tz	S2rz	S2zg	S2z	S2rz	S2z	N1tz	S3rz	S2rz	S2z	S2zg	S2zg	S2zg	S2zg	S2rz	S2zg	S2zg	S2tz	S2z	S2zg	S2rz	S2z	S2tz
Gattared dyhala	189	S3rz	S2z	S2rz	S2tz	S2rz	S2z	S3rz	S2rz	S2tz	S2rz	S2z	S2z	S2rz	S2z	N1tz	S3rz	S2rz	S2z	S2z	S2z	S2z	S2z	S2rz	S2z	S2z	S2tz	S2z	S2z	S2rz	S2z	S2tz
Gattared dyhala	190	S3rz	S2zg	S2rz	S2tz	S2rz	S2zg	S3rz	S2rz	S2tz	S2rz	S2zg	S2z	S2rz	S2z	N1tz	S3rz	S2rz	S2z	S2zg	S2zg	S2zg	S2zg	S2rz	S2zg	S2zg	S2tz	S2z	S2zg	S2rz	S2z	S2tz
Gattared dyhala	191	S3rz	S2zg	S2rz	S2tz	S2rz	S2zg	S3rz	S2rz	S2tz	S2rz	S2zg	S2z	S2rz	S2z	N1tz	S3rz	S2rz	S2z	S2zg	S2zg	S2zg	S2zg	S2rz	S2zg	S2zg	S2tz	S2z	S2zg	S2rz	S2z	S2tz
Gattared dyhala	192	S3rz	S2z	S2rz	S2tz	S2rz	S2z	S3rz	S2rz	S2tz	S2rz	S2z	S2z	S2rz	S2z	N1tz	S3rz	S2rz	S2z	S2z	S2z	S2z	S2z	S2rz	S2z	S2z	S2tz	S2z	S2z	S2rz	S2z	S2tz
Gattared dyhala	193	S3rz	S2zg	S2rz	S2tz	S2rz	S2zg	S3rz	S2rz	S2tz	S2rz	S2zg	S2z	S2rz	S2z	N1tz	S3rz	S2rz	S2z	S2zg	S2zg	S2zg	S2zg	S2rz	S2zg	S2zg	S2tz	S2z	S2zg	S2rz	S2z	S2tz
Gattared dyhala	194	S3rz	S2zg	S2rz	S2tz	S2rz	S2zg	S3rz	S2rz	S2tz	S2rz	S2zg	S2z	S2rz	S2z	N1tz	S3rz	S2rz	S2z	S2zg	S2zg	S2zg	S2zg	S2rz	S2zg	S2zg	S2tz	S2z	S2zg	S2rz	S2z	S2tz
Gattared dyhala	195	S3rz	S2z	S2rz	S2tz	S2rz	S2z	S3rz	S2rz	S2tz	S2rz	S2z	S2z	S2rz	S2z	N1tz	S3rz	S2rz	S2z	S2z	S2z	S2z	S2z	S2rz	S2z	S2z	S2tz	S2z	S2z	S2rz	S2z	S2tz
Gattared dyhala	202	N1rz	S2rz	S3rz	S2rz	S3rz	S2rz	N1rz	S3rz	S2rz	S3rz	S3rz	S2rz	S3rz	S2rz	N1n	S3rz	S3rz	S3rt	S2rz	S2rz	S2rz	S2rz	S3rz	S2rz	S2rz	S2r	S2r	S2rz	S3rz	S3rz	S2r



# **PART-B**

**SOCIO-ECONOMIC STATUS OF FARM HOUSEHOLDS**



## CONTENTS

1	Findings of the socio-economic survey	1-2
2	Introduction	3
3	Methodology	5-6
4	Salient features of the survey	7-24
5	Summary	25-28



### LIST OF TABLES

1	Households sampled for socio economic survey	7
2	Population characteristics	7
3	Age wise classification of household members	7
4	Education level of household members	8
5	Occupation of household heads	8
6	Occupation of family members	8
7	Institutional participation of household members	9
8	Type of house owned by households	9
9	Durable assets owned by households	9
10	Average value of durable assets owned by households	10
11	Farm implements owned by households	10
12	Average value of farm implements	10
13	Livestock possession by households	11
14	Average labour availability	11
15	Adequacy of hired labour	11
16	Migration among the households	11
17	Purpose of migration	11
18	Distribution of land (ha)	12
19	Average land value (Rs./ha)	12
20	Status of bore wells	12
21	Source of irrigation	12
22	Depth of water(Avg in meters)	12
23	Irrigated area (ha)	13
24	Cropping pattern	13
25	Cropping intensity	13
26	Possession of bank account and saving	13
27	Borrowing status	13
28.a	Cost of cultivation of Sunflower	14
28.b	Cost of cultivation of Red gram	15
28.c	Cost of cultivation of Sorghum	16
28.d	Cost of cultivation of Groundnut	17
28.e	Cost of cultivation of Maize	18



29	Adequacy of fodder	19
30	Annual gross income	19
31	Average annual expenditure	19
32	Horticultural species grown	19
33	Forest species grown	20
34	Average additional investment capacity	20
35	Source of funds for additional investment	20
36	Marketing of the agricultural produce	21
37	Marketing channels used for sale of agricultural produce	21
38	Mode of transport of agricultural produce	21
39	Incidence of soil and water erosion problems	21
40	Interest shown towards soil testing	22
41	Usage pattern of fuel for domestic use	22
42	Source of drinking water	22
43	Source of light	22
44	Existence of sanitary toilet facility	23
45	Possession of public distribution system (PDS) card	23
46	Participation in NREGA programme	23
47	Adequacy of food items	23
48	Inadequacy of food items	24
49	Farming constraints experienced	24

**FINDINGS OF THE SOCIO-ECONOMIC SURVEY**

- ❖ *The survey was conducted in Raghunathanahalli-2 is located at North latitude 15° 14' 31.598" and 15° 13' 30.436" and East longitude 75° 56' 30.124" and 75° 54' 44.824" covering an area of about 440.93 ha coming under Raghunathahalli, Gattareddyhala and Belagatti Villages of Koppal taluk.*
- ❖ *Socio-economic analysis of Raghunathanahalli-2 micro watersheds of Murlapura sub-watershed, Koppal taluk & District indicated that, out of the total sample of 35 farmers were sampled in Raghunathanahalli-2 micro-watershed among households surveyed 15 (42.86%) were marginal, 12 (34.29%) were small and 8 (22.86 %) were semi medium farmers.*
- ❖ *The population characteristics of households indicated that, there were 96 (56.47%) men and 74 (43.53 %) were women. The average population of marginal farmers was 5.06, small farmers were 4.83 and semi medium farmers were 4.5.*
- ❖ *Majority of the respondents (47.06%) were in the age group of 16-35 years.*
- ❖ *Education level of the sample households indicated that, there were 26.47 per cent illiterates, 68.83 per cent pre university education and 7.65 per cent attained graduation.*
- ❖ *About, 77.14 per cent of household heads practicing agriculture and 22.86 per cent of the household heads were engaged as agricultural labourers.*
- ❖ *Agriculture was the major occupation for 62.94 per cent of the household members.*
- ❖ *In the study area, 97.14 per cent of the households possess katcha house.*
- ❖ *The durable assets owned by the households showed that, 77.14 per cent possess TV, 34.29 per cent possess mixer grinder, 77.14 per cent possess mobile phones and 28.57 per cent possess motor cycles.*
- ❖ *Farm implements owned by the households indicated that, 11.43 per cent of the households possess plough, 8.57 per cent possess bullock cart.*
- ❖ *Regarding livestock possession by the households, 5.71 per cent possess local cow and 5.71 per cent possess buffalo.*
- ❖ *The average labour availability in the study area showed that, own labour men available in the micro watershed was 1.81, women available in the micro watershed was 1.44, hired labour (men) available was 6.58 and hired labour (women) available was 6.31.*
- ❖ *In the study area, about 0.59 per cent of the respondents migrated from the micro watershed in search of jobs.*
- ❖ *Out of the total land holding of the sample respondents 85.19 per cent (44.86 ha) of the area is under dry condition and the remaining 5.56 per cent area is irrigated land.*
- ❖ *There were 2.00 live bore wells among the sampled households.*
- ❖ *Bore well was the major source of irrigation for 5.71 per cent of the households.*

- ❖ *The major crops grown by sample farmers are Sunflower, Redgram, Sorghum, Groundnut and Maize and cropping intensity was recorded as 83.72 per cent.*
- ❖ *Out of the sample households 8.57 percent possessed bank account.*
- ❖ *About 8.57 per cent of the respondents borrowed credit from various sources.*
- ❖ *Per hectare cost of cultivation for Sunflower, Redgram, Sorghum, Groundnut and Maize was Rs.38349.84, 55753.72, 30816.36, 75904.68 and 30591.23 with benefit cost ratio of 1:0.80, 1: 2.10, 1: 0.50, 1: 1.60 and 1:0.70 respectively.*
- ❖ *Further, 42.86 per cent of the households opined that dry fodder was adequate and 8.57 per cent of the households have opined that the green fodder was adequate.*
- ❖ *The average annual gross income of the farmers was Rs. 78168.00 in micro-watershed, of which Rs. 25131.43 comes from agriculture.*
- ❖ *Sampled households have grown 12 horticulture trees and 108 forestry trees together in the fields and back yards.*
- ❖ *Households have an average investment capacity of Rs. 1571.43 for land development and Rs. 142.86 for irrigation facility.*
- ❖ *Source of funds for additional investment is concerned, 25.71 per cent depends on bank loan for land development activities.*
- ❖ *Regarding marketing channels, 100 per cent of the households have sold agricultural produce to the local/village merchants.*
- ❖ *Further, 100 per cent of the households have used tractor for the transport of agriculture commodity.*
- ❖ *Majority of the farmers (60.00%) have experienced soil and water erosion problems in the watershed and 54.29 per cent of the households were interested towards soil testing.*
- ❖ *Fire was the major source of fuel for domestic use for 100 per cent of the households.*
- ❖ *Piped supply was the major source for drinking water for 48.57 per cent of the households.*
- ❖ *Electricity was the major source of light for 100.00 per cent of the households.*
- ❖ *In the study area, 54.29 per cent of the households possess toilet facility.*
- ❖ *Regarding possession of PDS card, 100.00 per cent of the households possessed BPL card.*
- ❖ *Households opined that, the requirement of cereals (100.00%), pulses (60.00%) and oilseeds (22.86%) are adequate for consumption.*
- ❖ *Farming constraints experienced by households in the micro watersheds were lower fertility status of the soil (100.00%) wild animal menace on farm field (51.43%), frequent incidence of pest and diseases (91.43%), inadequacy of irrigation water (2.86%), high cost of fertilizers and plant protection chemicals (91.43%), high rate of interest on credit (40.00%), low price for the agricultural commodities (88.57%), lack of marketing facilities in the area (80.00%), inadequate extension services (5.71%), lack of transport for safe transport of the agricultural produce to the market (74.29%).*

## **INTRODUCTION**

Soil and water are the two precious natural resources which are essential for crop production and existence of life on earth. Rainfed agriculture is under severe stress due to various constraints related to agriculture like uneven and erratic distribution of rainfall, indiscriminate use of fertilizers, chemicals and pesticides, adoption of improper land management practices, soil erosion, decline in soil fertility, decline in ground water resources leading to low crop productivity. The area under rainfed agriculture has to be managed effectively using the best available practices to enhance the production of food, fodder and fuel. This is possible if the land resources are characterized at each parcel of land through detailed land resource inventory using the best available techniques of remote sensing, GPS and GIS. The watershed development programs are aimed at the sustainable distribution of its resources and the process of creating and implementing plans, programs, and projects to sustain and enhance watershed functions that affect the plant, animal and human communities within a watershed boundary.

World Bank funded KWDP II, SUJALA III project was implemented in with Broad objective of demonstrating more effective watershed management through greater integration of programmes related to rain-fed agriculture, innovative and science based approaches and strengthen institutional capacities and If successful, it is expected that the systems and tools could be mainstreamed into the overall IWMP in the State of Karnataka and in time, throughout other IWMP operations in India. With this background the socio-economic survey has been carried out with following specific objectives:

1. To understand the demographic features of the households in the micro-watershed
2. To understand the extent of family labour available and additional employment opportunities available within the village.
3. To know the status of assets of households in the micro-watershed for suggesting possible improvements.
4. To study the cropping pattern, cropped area and productivity levels of different households in micro-watershed.
5. To determine the type and extent of livestock owned by different categories of HHs
6. Availability of fodder and level of livestock management.

### **Scope and importance of survey**

Survey helps in identification of different socio-economic and resource use-patterns of farmers at the Micro watershed. Household survey provides demographic features, labour force, and levels of education; land ownership and asset position (including livestock and other household assets) of surveyed households; and cropping patterns, input intensities, and average crop yields from farmers' fields. It also discusses crop utilization and the degree of commercialization of production in the areas; farmers' access to and utilization of credit from formal and informal sources; and the level of adoption and use of soil, water, and pest management technologies.



## METHODOLOGY

The description of the methods, components selected for the survey and procedures followed in conducting the baseline survey are furnished under the following heads.

### **1. Description of the study area**

Koppal district is an administrative district in the state of Karnataka in India. In the past Koppal was referred to as 'Kopana Nagara'. Koppal, now a district headquarters is ancient Kopana a major holy place of the Jainas. The district occupies an area of 7,190 km<sup>2</sup> and has a population of 1,196,089, which 16.58% were urban as of 2001. The Koppal district was formed after split of Raichur district.

Geographers are very particular about the physiography or relief of a region. It plays a very important role in the spatial analysis of agricultural situation of the study area. The undulating topography with black cotton soil shrips, cut across by numerous nalas or streams is the major characteristic feature of the study region. Three physiographic divisions have made considering the local conditions of landforms and crops grown in the district. On the basis of physiography, Koppal district can be divided into three major divisions. They are (a) Koppal & Yelburga plateau, (b) Maidan division, (c) Tungabhadra valley. The district is part of Krishna basin the main streams draining the area are Maskinala, Ilkal-nadi and Hirenala. These are Ephemeral in nature, these come under Tungabhadra sub-basin. The drainage exhibit dentritic to subdentritic with drainage density varies from 1.4 to 7.0kms/sq.km.

According to the 2011 census Koppal district has a population of 1,391,292, roughly equal to the nation of Swaziland or the US state of Hawaii. This gives it a ranking of 350th in India (out of a total of 640). The district has a population density of 250 inhabitants per square kilometre (650/sq mi). Its population growth rate over the decade 2001-2011 was 16.32%. Koppal has a sex ratio of 983 females for every 1000 males, and a literacy rate of 67.28%.

### **2. Locale of the survey and description of the micro-watershed and**

The study was conducted in Raghunathanahalli-2 micro-watershed (Murlapura sub-watershed, Koppal taluk & District) is located at North latitude 15<sup>o</sup> 14' 31.598" and 15<sup>o</sup> 13' 30.436" and East longitude 75<sup>o</sup> 56' 30.124" and 75<sup>o</sup> 54' 44.824" covering an area of about 440.93 ha bounded by under Raghunathahalli, Gattareddyhala and Belagatti Villages.

### **3. Selection of the respondents for the study**

The micro-watershed is marked with 320 square meters grids. One farmer from every alternate grid in the micro-watershed was selected for the study and interviewed for socio-economic data. Totally 35 households were interviewed for the survey.

#### **4. The parameters considered for socio-economic survey of households**

Two forms of data were collected from the micro-watershed which includes primary data from the farm households and secondary data about the villages under the micro-watershed jurisdiction.

The following parameters were considered for the primary data collection about the socio-economic data of the households, (1) Demographic information, (2) Farm and durable assets owned by households, (3) Livestock possession, (4) Labour availability, (5) Level of migration in the village, Land holding, (7) Cropping pattern, (8) Source of irrigation, (9) Borrowing status, (10) Cost of cultivation of major crops, (11) Economics of subsidiary activities, (12) Fodder availability, (13) Family annual income from different sources, (14) Horticulture and forestry species grown, (15) Additional investment capacity, (16) Marketing practices, (17) Status of soil and water conservation structure, (18) Access to basic needs and (19) Constraints and suggestion.

The following parameters were considered for the secondary data regarding the villages under the micro-watershed jurisdiction, (1) Number of villages in each micro-watershed jurisdiction, (2) Village wise number of households, (3) Geographical area of the villages, (4) Cultivable area including rainfed and irrigated, (5) Number and type of house in each village, (6) Human and livestock population, (7) Facilities in the village such as roads, transport facility for conveyance, drinking water supply, street light and (8) Community based organizations in the villages.

#### **5. Development of interview schedule and data collection**

Taking into the consideration the objectives of the survey, an interview schedule was prepared after thorough consultation with the experts in the field of social sciences. A comprehensive interview schedule covering all the major parameters for measuring the socio-economic situation was developed.

#### **6. Tools used to analyze the data**

The statistical components such as frequency and percentage were used to analyze the data.

#### **Abbreviations used in the report**

LL=Landless

MF=Marginal Farmers

SF=Small farmers

SMF=Semi medium farmers

MDF=Medium farmers

LF=Large Farmers

### FINDINGS OF THE SURVEY

This chapter deals with systematic presentation of results of the survey. Keeping in view the objectives, the salient features of the survey are presented under the following headings.

**Households sampled for socio-economic survey:** The data on households sampled for socio economic survey in Raghunathanahalli-2 Micro watershed is presented in Table 1 and it indicated that 35 farmers were sampled in Raghunathanahalli-2 micro-watershed among households surveyed 15 (42.86%) were marginal, 12 (34.29%) were small and 8 (22.86 %) were semi medium farmers.

**Table 1. Households sampled for socio economic survey in Raghunathanahalli-2 micro-watershed**

Sl.No.	Particulars	LL (0)		MF (15)		SF (12)		SMF (8)		All (35)	
		N	%	N	%	N	%	N	%	N	%
1	Farmers	0	0	15	42.9	12	34.3	8	22.9	35	100

**Population characteristics:** The population characteristics of households sampled for socio-economic survey in Raghunathanahalli-2 Micro watershed is presented in Table 2. The data indicated that, there were 96 (56.47%) men and 74 (43.53%) were women. The average population of marginal farmers was 5.06, small farmers were 4.83 and semi medium farmers were 4.5.

**Table 2. Population characteristics in Raghunathanahalli-2 micro-watershed**

Sl.No.	Particulars	LL (0)		MF (76)		SF (58)		SMF (36)		All (170)	
		N	%	N	%	N	%	N	%	N	%
1	Men	0	0	49	64	28	48	19	52.8	96	56.5
2	Women	0	0	27	36	30	52	17	47.2	74	43.5
Total		0	100	76	100	58	100	36	100	170	100
Average		0.0		5.06		4.83		4.5		4.85	

**Age wise classification of population:** The age wise classification of household members in Raghunathanahalli-2 Micro watershed is presented in Table 3. The indicated that, 23 (13.53%) of population were 0-15 years of age, 80 (47.06%) were 16-35 years of age, 44(25.88%) were 36-60 years of age and 23 (13.53 %) were above 61 years of age.

**Table 3: Age wise classification of members of the household in Raghunathanahalli-2 micro-watershed**

Sl.No.	Particulars	LL (0)		MF (76)		SF (58)		SMF (36)		All (170)	
		N	%	N	%	N	%	N	%	N	%
1	0-15 years of age	0	0	10	13.2	13	22.4	0	0	23	13.53
2	16-35 years of age	0	0	41	54	21	36.2	18	50	80	47.06
3	36-60 years of age	0	0	15	19.7	16	27.6	13	36.11	44	25.88
4	> 61 years	0	0	10	13.2	8	13.8	5	13.89	23	13.53
Total		0	100	76	100	58	100	36	100	170	100



**Education level of household members:** Education level of household members in Raghunathanahalli-2 Micro watershed is presented in Table 4. The results indicated that, there were 26.47 per cent of illiterates, 34.12 per cent of them had primary school education, 4.12 per cent middle school education, 17.65 per cent high school education, 6.47 per cent of them had PUC education, 0.59 per cent of them had Diploma, 7.65 per cent attained graduation and 1.18 them had other education.

**Table 4. Education level of members of the household in Raghunathanahalli-2 micro-watershed**

Sl.No.	Particulars	LL (0)		MF (76)		SF (58)		SMF (36)		All (170)	
		N	%	N	%	N	%	N	%	N	%
1	Illiterate	0	0	26	34.2	15	25.9	4	11.1	45	26.5
2	Primary School	0	0	19	25	22	37.9	17	47.2	58	34.1
3	Middle School	0	0	3	3.95	3	5.17	1	2.78	7	4.12
4	High School	0	0	12	15.8	14	24.1	4	11.1	30	17.7
5	PUC	0	0	8	10.5	2	3.45	1	2.78	11	6.47
6	Diploma	0	0	1	1.32	0	0	0	0	1	0.59
7	ITI	0	0	0	0	1	1.72	1	2.78	2	1.18
8	Degree	0	0	5	6.58	1	1.72	7	19.4	13	7.65
9	Masters	0	0	0	0	0	0	1	2.78	1	0.59
10	Others	0	0	2	2.63	0	0	0	0	2	1.18
Total		0	100	76	100	58	100	36	100	170	100

**Occupation of head of households:** The data regarding the occupation of the household heads in Raghunathanahalli-2 Micro watershed is presented in Table 5. The results indicate that, 77.14 per cent of households heads were practicing agriculture, 22.86 per cent of the household heads were agricultural Labour.

**Table 5: Occupation of heads of households in Raghunathanahalli-2 micro-watershed**

Sl.No.	Particulars	LL (0)		MF (15)		SF (12)		SMF (8)		All (35)	
		N	%	N	%	N	%	N	%	N	%
1	Agriculture	0	0	10	67	9	75	8	100	27	77.14
2	Agricultural Labour	0	0	5	33	3	25	0	0	8	22.86
Total		0	100	15	100	12	100	8	100	35	100

**Table 6: Occupation of members of the household in Raghunathanahalli-2 micro-watershed**

Sl.No.	Particulars	LL (0)		MF (76)		SF (58)		SMF (36)		All (170)	
		N	%	N	%	N	%	N	%	N	%
1	Agriculture	0	0	51	67.1	28	48.28	28	77.78	107	62.9
2	Agricultural Labour	0	0	9	11.8	13	22.41	0	0	22	12.9
3	General Labour	0	0	1	1.32	2	3.45	0	0	3	1.76
4	Private Service	0	0	1	1.32	1	1.72	1	2.78	3	1.76
5	Student	0	0	12	15.8	14	24.14	7	19.44	33	19.4
6	Children	0	0	2	2.63	0	0	0	0	2	1.18
Total		0	100	76	100	58	100	36	100	170	100

**Occupation of the members of the household:** The data regarding the occupation of the household members in Raghunathanahalli-2 Micro watershed is presented in Table 6. The results indicate that, agriculture was the major occupation for 62.94 per cent of the household members, 12.94 per cent were agricultural labour, 1.76 per cent were general labour, 19.41 per cent were working in pursuing education and 1.18 per cent were children.

**Institutional Participation of household members:** The data regarding the institutional participation of the household members in Raghunathanahalli-2 Micro watershed is presented in Table 7. The results show that, out of the total family members in the households 100 per cent of them were not participating in any of the institutions.

**Table 7: Institutional Participation of household member in Raghunathanahalli-2 micro-watershed**

Sl.No.	Particulars	LL (0)		MF (76)		SF (58)		SMF (36)		All (170)	
		N	%	N	%	N	%	N	%	N	%
1	No Participation	0	0	76	100	58	100	36	100	170	100
	Total	0	100	76	100	58	100	36	100	170	100

**Type of house owned:** The data regarding the type of house owned by the households in Raghunathanahalli-2 Micro watershed is presented in Table 8. The results indicate that, 2.86 percent possess thatched house, 97.14 per cent of the households possess katcha house.

**Table 8. Type of house owned by households in Raghunathanahalli-2 micro-watershed**

Sl.No.	Particulars	LL (0)		MF (15)		SF (12)		SMF (8)		All (35)	
		N	%	N	%	N	%	N	%	N	%
1	Thatched	0	0	1	6.7	0	0	0	0	1	2.86
2	Katcha	0	0	14	93	12	100	8	100	34	97.14
	Total	0	100	15	100	12	100	8	100	35	100

**Durable assets owned by the households:** The data regarding the Durable Assets owned by the households in Raghunathanahalli-2 Micro watershed is presented in Table 9. The result shows that, 77.14 per cent possess TV, 34.29 per cent possess mixer grinder, 17.14 per cent possess Bicycle, 28.57 per cent possess motor cycle, 77.14 per cent possess mobile phones.

**Table 9. Durable assets owned by households in Raghunathanahalli-2 micro-watershed**

Sl.No.	Particulars	LL (0)		MF (15)		SF (12)		SMF (8)		All (35)	
		N	%	N	%	N	%	N	%	N	%
1	Television	0	0	14	93	6	50	7	88	27	77.14
2	Mixer/Grinder	0	0	8	53	1	8.33	3	38	12	34.29
3	Bicycle	0	0	2	13	2	16.7	2	25	6	17.14
4	Motor Cycle	0	0	6	40	1	8.33	3	38	10	28.57
5	Mobile Phone	0	0	14	93	8	66.7	5	63	27	77.14
6	Blank	0	0	0	0	1	8.33	1	13	2	5.71

**Average value of durable assets:** The data regarding the average value of durable assets owned by the households in Raghunathanahalli-2 Micro watershed is presented in Table 10. The result shows that, the average value of television was Rs.7685.00, mixer grinder was Rs.1833.00, bicycle was Rs.2166.00, motor cycle was Rs. 41000.00, mobile phone was Rs.2129.00.

**Table 10. Average value of durable assets owned in Raghunathanahalli-2 micro-watershed**

Sl.No.	Particulars	Average Value (Rs.)				
		LL (0)	MF (15)	SF (12)	SMF (8)	All (35)
1	Television	0	7892	7666	7285	7685
2	Mixer/Grinder	0	1850	2000	1733	1833
3	Bicycle	0	2000	1500	3000	2166
4	Motor Cycle	0	45500	42000	31666	41000
5	Mobile Phone	0	2860	1730	1272	2129

**Farm implements owned:** The data regarding the farm implements owned by the households in Raghunathanahalli-2 Micro watershed is presented in Table 11. About 8.57 per cent of the households possess Bullock Cart, 11.43 per cent possess plough and 17.14 per cent possess Weeder.

**Table 11. Farm implements owned in Raghunathanahalli-2 micro-watershed**

Sl.No.	Particulars	LL (0)		MF (15)		SF (12)		SMF (8)		All (35)	
		N	%	N	%	N	%	N	%	N	%
1	Bullock Cart	0	0	2	13.3	1	8.33	0	0	3	8.57
2	Plough	0	0	2	13.3	1	8.33	1	12.5	4	11.43
3	Weeder	0	0	4	26.7	2	16.67	0	0	6	17.14
4	Thresher	0	0	0	0	1	8.33	0	0	1	2.86
5	Blank	0	0	8	53.3	8	66.67	7	87.5	23	65.71

**Average value of farm implements:** The data regarding the average value of farm Implements owned by the households in Raghunathanahalli-2 Micro watershed is presented in Table 12. The results show that the average value of plough was Rs.1200.00, bullock Cart was Rs.13333.00 and weeder was Rs.96.00.

**Table 12. Average value of farm implements in Raghunathanahalli-2 micro-watershed**

Sl.No.	Particulars	Average Value (Rs.)				
		LL (0)	MF (15)	SF (12)	SMF (8)	All (35)
1	Bullock Cart	0	10000	20000	0	13333
2	Plough	0	1200	1200	1200	1200
3	Weeder	0	110	70	0	96
4	Thresher	0	0	250	0	250

**Livestock possession by the households:** The data regarding the Livestock possession by the households in Raghunathanahalli-2 Micro watershed is presented in Table 13. The results indicate that, 8.57 per cent of the households possess bullocks, 5.71 per cent possess local cow, 5.71 per cent possess buffalo, 2.86 per cent were poultry birds.

**Table 13. Livestock possession by households in Raghunathanahalli-2 micro-watershed**

Sl.No.	Particulars	LL (0)		MF (15)		SF (12)		SMF (8)		All (35)	
		N	%	N	%	N	%	N	%	N	%
1	Bullock	0	0	2	13	1	8.33	0	0	3	8.57
2	Local cow	0	0	1	6.7	1	8.33	0	0	2	5.71
3	Buffalo	0	0	2	13	0	0	0	0	2	5.71
4	Poultry birds	0	0	1	6.7	0	0	0	0	1	2.86
5	blank	0	0	10	67	11	91.67	8	100	29	82.86

**Average Labour availability:** The data regarding the average labour availability in Raghunathanahalli-2 Micro watershed is presented in Table 14. The indicated that, own labour men available in the micro watershed was 1.81, women available in the micro watershed was 1.44, hired labour (men) available was 6.58 and hired labour (women) available was 6.31.

**Table 14. Average labour availability in Raghunathanahalli-2 micro-watershed**

Sl.No.	Particulars	LL (0)	MF (15)	SF (12)	SMF (8)	All (35)
		N	N	N	N	N
1	Hired labour Female	0	5.27	8.25	5.44	6.31
2	Own Labour Female	0	1.47	1.75	1	1.44
3	Own labour Male	0	2	1.58	1.78	1.81
4	Hired labour Male	0	4.73	9.25	6.11	6.58

**Adequacy of hired labour:** The data regarding the adequacy of hired labour in Raghunathanahalli-2 Micro watershed is presented in Table 15. The results indicate that, 100 per cent of the household opined that hired labour was adequate.

**Table 15. Adequacy of hired labour in Raghunathanahalli-2 micro-watershed**

Sl.No.	Particulars	LL (0)		MF (15)		SF (12)		SMF (8)		All (35)	
		N	%	N	%	N	%	N	%	N	%
1	Adequate	0	0	15	100	12	100	8	100	35	100

**Migration among the households:** The data regarding the migration (Table 16) indicate that, 0.59 percent of the population was being migrated from the micro watershed.

**Table 16. Migration among the households in Raghunathanahalli-2 micro-watershed**

Sl.No.	Particulars	LL (0)		MF (76)		SF (58)		SMF (36)		All (170)	
		N	%	N	%	N	%	N	%	N	%
1	Migration	0	0.00	0	0.00	0	0.00	1	2.78	1	0.59

**Purpose of migration:** The data regarding the purpose of migration (Table 17) indicate that, 100.00 percent of them went for the purpose of job/wage/work.

**Table 17. Purpose of migration by members of households in Raghunathanahalli-2 micro-watershed**

Sl.No.	Particulars	LL (0)		MF (0)		SF (0)		SMF (1)		All (1)	
		N	%	N	%	N	%	N	%	N	%
1	Job/wage/work	0	0	0	0	0	0	1	100	1	100
Total		0	100	0	100	0	100	1	100	1	100

**Distribution of land (ha):** The data regarding the distribution of land (ha) in Raghunathanahalli-2 Micro watershed is presented in Table 18. The results indicate that, 38.22 ha (85.19%) of dry land and 2.49 ha (5.56 %) of irrigated land.

**Table 18. Distribution of land (ha) in Raghunathanahalli-2 micro-watershed**

Sl.No.	Particulars	LL (0)		MF (15)		SF (12)		SMF (8)		All (35)	
		N	%	N	%	N	%	N	%	N	%
1	Dry	0	0	9.81	91.82	11.24	81.84	17.17	83.97	38.22	85.19
2	Irrigated	0	0	0	0	2.49	18.16	0	0	2.49	5.56
3	Permanent Fallow	0	0	0.87	8.18	0	0	3.28	16.03	4.15	9.26
	Total	0	100	10.7	100	13.73	100	20.45	100	44.86	100

**Average value of land (ha):** The data regarding the average land value (Rs./ha) in Raghunathanahalli-2 Micro watershed is presented in Table 19. The results show that the average value of dry land was Rs.306036.22 and the average value of irrigated land was Rs.481168.82.

**Table 19. Average value of land (ha) in Raghunathanahalli-2 micro-watershed**

Sl.No.	Particulars	LL (0)	MF (15)	SF (12)	SMF (8)	All (35)
		N	N	N	N	N
1	Dry	0	377021.5	284623.7	279490.8	306036.2
2	Irrigated	0	0	481168.8	0	481168.8
3	Permanent Fallow	0	343055.5	0	304938.3	312963

**Status of bore wells:** The data regarding the status of bore wells in Raghunathanahalli-2 Micro watershed is presented in Table 20. The results indicate that, there were 2 functioning bore wells among the sampled households in micro watershed.

**Table 20. Status of bore wells in Raghunathanahalli-2 micro-watershed**

Sl.No.	Particulars	LL (0)	MF (15)	SF (12)	SMF (8)	All (35)
		N	N	N	N	N
1	Functioning	0	0	2	0	2

**Source of irrigation:** The data regarding the source of irrigation in Raghunathanahalli-2 Micro watershed is presented in Table 21. The results show that, bore well for 5.71 per cent of the households.

**Table 21. Source of irrigation in Raghunathanahalli-2 micro-watershed**

Sl.No.	Particulars	LL (0)		MF (15)		SF (12)		SMF (8)		All (35)	
		N	%	N	%	N	%	N	%	N	%
1	Bore Well	0	0	0	0	2	16.67	0	0	2	5.71

**Depth of water (Avg. In meters):** The data regarding the depth of water in Raghunathanahalli-2 Micro watershed is presented in Table 22. The results revealed that, the depth of bore well was 6.10 meter.

**Table 22. Depth of water (Avg. In meters) in Raghunathanahalli-2 micro-watershed**

Sl.No.	Particulars	LL (0)	MF (15)	SF (12)	SMF (8)	All (35)
		N	N	N	N	N
1	Bore Well	0	0	17.78	0	6.1

**Irrigated Area (ha):** The data regarding the irrigated area (ha) in Raghunathanahalli-2 Micro watershed is presented in Table 23. The results indicate that, the availability of irrigation water was used for kharif crops was 1.68 ha.

**Table 23. Irrigated Area (ha) in Raghunathanahalli-2 micro-watershed**

Sl.No.	Particulars	LL (0)	MF (15)	SF (12)	SMF (8)	All (35)
1	Kharif	0	0	1.68	0	1.68

**Cropping pattern:** The data regarding the cropping pattern in Raghunathanahalli-2 Micro watershed is presented in Table 24. The results indicate that, farmers have grown Sunflower (17.72 ha), Maize (8.57 ha), Sorghum (6.04 ha), Bajra (4.78 ha) Bengal gram (2.45 ha) and Groundnut (0.40 ha).

**Table 24. Cropping pattern in Raghunathanahalli-2 micro-watershed**

Sl.No.	Particulars	LL (0)	MF (15)	SF (12)	SMF (8)	All (35)
1	Kharif - Sunflower	0	1.68	6.3	9.74	17.72
2	Kharif - Maize	0	1.7	1.56	5.31	8.57
3	Kharif - Sorghum	0	4.83	1.21	0	6.04
4	Kharif - Bajra	0	1.13	2.43	1.21	4.78
5	Kharif - Bengal gram	0	0	2.05	0.4	2.45
6	Kharif - Groundnut	0	0.4	0	0	0.4
Total		0	9.74	13.56	16.67	39.97

**Cropping intensity:** The data regarding the cropping intensity in Raghunathanahalli-2 Micro watershed is presented in Table 25. The results indicate that, the cropping intensity was 83.72 per cent.

**Table 25. Cropping intensity (%) in Raghunathanahalli-2 micro-watershed**

Sl.No.	Particulars	LL (0)	MF (15)	SF (12)	SMF (8)	All (35)
1	Cropping Intensity	0	91.83	75.17	87.28	83.72

**Possession of bank account and savings:** The data regarding the possession of bank account and saving in Raghunathanahalli-2 micro-watershed is presented in Table 26. The results indicate that, 8.57 cent of the households posses bank account.

**Table 26. Possession of Bank account and savings in Raghunathanahalli-2 micro-watershed**

Sl.No.	Particulars	LL (0)		MF (15)		SF (12)		SMF (8)		All (35)	
		N	%	N	%	N	%	N	%	N	%
1	Account	0	0	2	13.33	0	0	1	12.5	3	8.57

**Borrowing status:** The data regarding the borrowing status in Raghunathanahalli-2 micro-watershed is presented in Table 27. The results indicate that, 8.57 percent of the sample farmers have borrowed credit from different sources.

**Table 27. Borrowing status in Raghunathanahalli-2 micro-watershed**

Sl.No.	Particulars	LL (0)		MF (15)		SF (12)		SMF (8)		All (35)	
		N	%	N	%	N	%	N	%	N	%
1	Credit Availed	0	0	2	13.33	0	0	1	12.5	3	8.57

**Cost of Cultivation of Sunflower:** The data regarding the cost of cultivation (Rs/ha) of Sunflower in Raghunathanahalli-2 micro watershed is presented in Table 28.a. The results indicate that, the total cost of cultivation (Rs/ha) for Sunflower was Rs. 38349.84. The gross income realized by the farmers was Rs. 31333.91. The net income from Sunflower cultivation was Rs.-7015.93, thus the benefit cost ratio was found to be 1:0.80.

**Table 28(a). Cost of Cultivation of Sunflower in Raghunathanahalli-2 micro-watershed**

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
<b>I</b>	<b>Cost A1</b>				
1	Hired Human Labour	Man days	43.96	8097.41	21.11
2	Bullock	Pairs/day	1.48	790.71	2.06
3	Tractor	Hours	3.83	3599.09	9.38
4	Machinery	Hours	0	0	0
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	9.36	5110.6	13.33
6	Seed Inter Crop	Kgs.	0	0	0
7	FYM	Quintal	3.09	2701.87	7.05
8	Fertilizer + micronutrients	Quintal	4.65	4546.2	11.85
9	Pesticides (PPC)	Kgs / liters	2.53	3195.64	8.33
10	Irrigation	Number	0	0	0
13	Depreciation charges		0	106.69	0.28
14	Land revenue and Taxes		0	4.28	0.01
<b>II</b>	<b>Cost B1</b>				
16	Interest on working capital			1866.57	4.87
17	<b>Cost B1 = (Cost A1 + sum of 15 and 16)</b>			30019.05	78.28
<b>III</b>	<b>Cost B2</b>				
18	Rental Value of Land			392.22	1.02
19	<b>Cost B2 = (Cost B1 + Rental value)</b>			30411.28	79.3
<b>IV</b>	<b>Cost C1</b>				
20	Family Human Labour		20.83	4451.81	11.61
21	<b>Cost C1 = (Cost B2 + Family Labour)</b>			34863.09	90.91
<b>V</b>	<b>Cost C2</b>				
22	Risk Premium			0.4	0
23	<b>Cost C2 = (Cost C1 + Risk Premium)</b>			34863.49	90.91
<b>VI</b>	<b>Cost C3</b>				
24	Managerial Cost			3486.35	9.09
25	<b>Cost C3 = (Cost C2 + Managerial Cost)</b>			38349.84	100
<b>VII</b>	<b>Economics of the Crop</b>				
		a) Main Product (q)	10.35	31333.91	
a.	Main Product	b) Main Crop Sales Price (Rs.)		3026.67	
b.	Gross Income (Rs.)			31333.91	
c.	Net Income (Rs.)			-7015.93	
d.	Cost per Quintal (Rs./q.)			3704.36	
e.	Benefit Cost Ratio (BC Ratio)			1:0.8	

**Cost of Cultivation of Redgram:** The data regarding the cost of cultivation (Rs/ha) of Redgram in Raghunathanahalli-2 micro watershed is presented in Table 28.b. The results indicate that, the total cost of cultivation (Rs/ha) for Redgram was Rs. 55753.72. The gross income realized by the farmers was Rs. 117886.36. The net income from Redgram cultivation was Rs.62132.64, thus the benefit cost ratio was found to be 1:2.10.

**Table 28(b). Cost of Cultivation of Redgram in Raghunathanahalli-2 micro-watershed**

Sl. No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
<b>I</b>	<b>Cost A1</b>				
1	Hired Human Labour	Man days	85.33	13697.27	24.57
2	Bullock	Pairs/day	0	0	0
3	Tractor	Hours	6.74	6736.36	12.08
4	Machinery	Hours	0	0	0
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	11.23	1347.27	2.42
6	Seed Inter Crop	Kgs.	0	0	0
7	FYM	Quintal	0	0	0
8	Fertilizer + micronutrients	Quintal	8.98	7185.45	12.89
9	Pesticides (PPC)	Kgs / liters	4.49	6287.27	11.28
10	Irrigation	Number	0	0	0
13	Depreciation charges		0	0.04	0
14	Land revenue and Taxes		0	4.94	0.01
<b>II</b>	<b>Cost B1</b>				
16	Interest on working capital			1778.4	3.19
17	<b>Cost B1 = (Cost A1 + sum of 15 and 16)</b>			37037.02	66.43
<b>III</b>	<b>Cost B2</b>				
18	Rental Value of Land			400	0.72
19	<b>Cost B2 = (Cost B1 + Rental value)</b>			37437.02	67.15
<b>IV</b>	<b>Cost C1</b>				
20	Family Human Labour		58.38	13248.18	23.76
21	<b>Cost C1 = (Cost B2 + Family Labour)</b>			50685.2	90.91
<b>V</b>	<b>Cost C2</b>				
22	Risk Premium			0	0
23	<b>Cost C2 = (Cost C1 + Risk Premium)</b>			50685.2	90.91
<b>VI</b>	<b>Cost C3</b>				
24	Managerial Cost			5068.52	9.09
25	<b>Cost C3 = (Cost C2 + Managerial Cost)</b>			55753.72	100
<b>VII</b>	<b>Economics of the Crop</b>				
a.	Main Product	a) Main Product (q)		33.68	117886.36
		b) Main Crop Sales Price (Rs.)			3500
b.	Gross Income (Rs.)			117886.36	
c.	Net Income (Rs.)			62132.64	
d.	Cost per Quintal (Rs./q.)			1655.31	
e.	Benefit Cost Ratio (BC Ratio)			1:2.1	



**Cost of Cultivation of Sorghum:** The data regarding the cost of cultivation (Rs/ha) of Sorghum in Raghunathanahalli-2 micro watershed is presented in Table 28.c. The results indicate, the total cost of cultivation (Rs/ha) for Sorghum was Rs.30816.36. The gross income realized by the farmers was Rs. 16676.95. The net income from Sorghum cultivation was Rs. -14139.41, thus the benefit cost ratio was found to be 1:0.50.

**Table 28(c). Cost of Cultivation of Sorghum in Raghunathanahalli-2 micro-watershed**

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
<b>I</b>	<b>Cost A1</b>				
1	Hired Human Labour	Man days	44.32	7715.47	25.04
2	Bullock	Pairs/day	1.48	935.72	3.04
3	Tractor	Hours	2.95	2748.01	8.92
4	Machinery	Hours	0.33	261.38	0.85
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	12.81	1775.45	5.76
7	FYM	Quintal	1.72	343.06	1.11
8	Fertilizer + micronutrients	Quintal	4.4	3826.47	12.42
9	Pesticides (PPC)	Kgs / liters	2.78	3891.12	12.63
13	Depreciation charges		0	1.25	0
14	Land revenue and Taxes		0	4.47	0.01
<b>II</b>	<b>Cost B1</b>				
16	Interest on working capital			1180.41	3.83
17	<b>Cost B1 = (Cost A1 + sum of 15 and 16)</b>			22682.8	73.61
<b>III</b>	<b>Cost B2</b>				
18	Rental Value of Land			375	1.22
19	<b>Cost B2 = (Cost B1 + Rental value)</b>			23057.8	74.82
<b>IV</b>	<b>Cost C1</b>				
20	Family Human Labour		22.2	4956.44	16.08
21	<b>Cost C1 = (Cost B2 + Family Labour)</b>			28014.25	90.91
<b>V</b>	<b>Cost C2</b>				
22	Risk Premium			0.63	0
23	<b>Cost C2 = (Cost C1 + Risk Premium)</b>			28014.87	90.91
<b>VI</b>	<b>Cost C3</b>				
24	Managerial Cost			2801.49	9.09
25	<b>Cost C3 = (Cost C2 + Managerial Cost)</b>			30816.36	100
<b>VII</b>	<b>Economics of the Crop</b>				
a.	Main Product	a) Main Product (q)		12.77	16413.81
		b) Main Crop Sales Price (Rs.)			1285.71
	By Product	e) Main Product (q)		1.41	263.14
		f) Main Crop Sales Price (Rs.)			187.14
b.	Gross Income (Rs.)			16676.95	
c.	Net Income (Rs.)			-14139.41	
d.	Cost per Quintal (Rs./q.)			2413.88	
e.	Benefit Cost Ratio (BC Ratio)			1:0.5	

**Cost of Cultivation of Groundnut:** The data regarding the cost of cultivation (Rs/ha) of Groundnut in Raghunathanahalli-2 micro watershed is presented in Table 28.d. The results indicate that, the total cost of cultivation (Rs/ha) for Groundnut was Rs. 75904.68. The gross income realized by the farmers was Rs.120412.50. The net income from Groundnut cultivation was Rs. 44507.82, thus the benefit cost ratio was found to be 1:1.60.

**Table 28(d). Cost of Cultivation of Groundnut in Raghunathanahalli-2 micro-watershed**

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
<b>I</b>	<b>Cost A1</b>				
1	Hired Human Labour	Man days	56.81	10621	13.99
2	Bullock	Pairs/day	1.85	1235	1.63
3	Tractor	Hours	2.47	4940	6.51
4	Machinery	Hours	0	0	0
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	123.5	19760	26.03
6	Seed Inter Crop	Kgs.	0	0	0
7	FYM	Quintal	4.94	5928	7.81
8	Fertilizer + micronutrients	Quintal	4.94	5742.75	7.57
9	Pesticides (PPC)	Kgs / liters	2.47	2964	3.9
10	Irrigation	Number	0	0	0
13	Depreciation charges		0	247.02	0.33
14	Land revenue and Taxes		0	4.94	0.01
<b>II</b>	<b>Cost B1</b>				
16	Interest on working capital			4127.37	5.44
17	<b>Cost B1 = (Cost A1 + sum of 15 and 16)</b>			55570.08	73.21
<b>III</b>	<b>Cost B2</b>				
18	Rental Value of Land			466.67	0.61
19	<b>Cost B2 = (Cost B1 + Rental value)</b>			56036.75	73.83
<b>IV</b>	<b>Cost C1</b>				
20	Family Human Labour		40.76	12967.5	17.08
21	<b>Cost C1 = (Cost B2 + Family Labour)</b>			69004.25	90.91
<b>V</b>	<b>Cost C2</b>				
22	Risk Premium			0	0
23	<b>Cost C2 = (Cost C1 + Risk Premium)</b>			69004.25	90.91
<b>VI</b>	<b>Cost C3</b>				
24	Managerial Cost			6900.43	9.09
25	<b>Cost C3 = (Cost C2 + Managerial Cost)</b>			75904.68	100
<b>VII</b>	<b>Economics of the Crop</b>				
a.	Main Product	a) Main Product (q)	40.14	120412.5	
		b) Main Crop Sales Price (Rs.)		3000	
b.	Gross Income (Rs.)			120412.5	
c.	Net Income (Rs.)			44507.82	
d.	Cost per Quintal (Rs./q.)			1891.12	
e.	Benefit Cost Ratio (BC Ratio)			1:1.6	

**Cost of Cultivation of Maize:** The data regarding the cost of cultivation (Rs/ha) of Maize in Raghunathanahalli-2 micro watershed is presented in Table 28.e. The results indicate that, the total cost of cultivation (Rs/ha) for Maize was Rs.30591.23. The gross income realized by the farmers was Rs. 22221.03. The net income from Maize cultivation was Rs. -8370.20, thus the benefit cost ratio was found to be 1:0.70.

**Table 28(e). Cost of Cultivation of Maize in Raghunathanahalli-2 micro-watershed**

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
<b>I</b>	<b>Cost A1</b>				
1	Hired Human Labour	Man days	31.27	5365.72	17.54
2	Bullock	Pairs/day	2.22	1274	4.16
3	Tractor	Hours	5.27	4744.38	15.51
4	Machinery	Hours	1.27	1012.57	3.31
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	17.26	2281.34	7.46
6	Seed Inter Crop	Kgs.	0	0	0
7	FYM	Quintal	2.59	1505.67	4.92
8	Fertilizer + micronutrients	Quintal	3.7	3646.22	11.92
9	Pesticides (PPC)	Kgs / liters	0.35	347.4	1.14
10	Irrigation	Number	0	0	0
13	Depreciation charges		0	6.36	0.02
14	Land revenue and Taxes		0	3.71	0.01
<b>II</b>	<b>Cost B1</b>				
16	Interest on working capital			933.77	3.05
17	<b>Cost B1 = (Cost A1 + sum of 15 and 16)</b>			21121.13	69.04
<b>III</b>	<b>Cost B2</b>				
18	Rental Value of Land			358.33	1.17
19	<b>Cost B2 = (Cost B1 + Rental value)</b>			21479.46	70.21
<b>IV</b>	<b>Cost C1</b>				
20	Family Human Labour		27.54	6330	20.69
21	<b>Cost C1 = (Cost B2 + Family Labour)</b>			27809.46	90.91
<b>V</b>	<b>Cost C2</b>				
22	Risk Premium			0.75	0
23	<b>Cost C2 = (Cost C1 + Risk Premium)</b>			27810.21	90.91
<b>VI</b>	<b>Cost C3</b>				
24	Managerial Cost			2781.02	9.09
25	<b>Cost C3 = (Cost C2 + Managerial Cost)</b>			30591.23	100
<b>VII</b>	<b>Economics of the Crop</b>				
a.	Main Product	a) Main Product (q)		14.34	21145.76
		b) Main Crop Sales Price (Rs.)			1475
	By Product	e) Main Product (q)		1.19	1075.27
		f) Main Crop Sales Price (Rs.)			900
b.	Gross Income (Rs.)			22221.03	
c.	Net Income (Rs.)			-8370.2	
d.	Cost per Quintal (Rs./q.)			2133.86	
e.	Benefit Cost Ratio (BC Ratio)			1:0.7	

**Adequacy of fodder:** The data regarding the adequacy of fodder in Raghunathanahalli-2 Micro watershed is presented in Table 29. The results indicate that, 42.86 per cent of the households opined that dry fodder was adequate. With respect to green fodder availability, 8.57 percent of them opined it was sufficient.

**Table 29. Adequacy of fodder in Raghunathanahalli-2 micro-watershed**

Sl.No.	Particulars	LL (0)		MF (15)		SF (12)		SMF (8)		All (35)	
		N	%	N	%	N	%	N	%	N	%
1	Adequate-Dry Fodder	0	0	6	40	7	58.33	2	25	15	42.86
2	Adequate-Green Fodder	0	0	1	6.67	1	8.33	1	12.5	3	8.57

**Average annual gross income:** The data regarding the annual gross income in Raghunathanahalli-2 Micro watershed is presented in Table 30. The results indicate that, the farmers have annual gross income of Rs. 78168.00 in micro-watershed, of which Rs. 25131.43 is from agriculture itself.

**Table 30. Average annual gross income in Raghunathanahalli-2 micro-watershed**

Sl.No.	Particulars	LL (0)	MF (15)	SF (12)	SMF (8)	All (35)
		Rs.	Rs.	Rs.	Rs.	Rs.
1	Service/salary	0	49600	20000	81000	46628.6
2	Wage	0	2133.33	9666.67	1500	4571.43
3	Agriculture	0	25600	20912.5	30581.3	25131.4
4	Dairy Farm	0	2133.33	2690	0	1836.57
Income(Rs.)		0	79466.7	53269.2	113081	78168

**Average annual Expenditure:** The data regarding the average annual expenditure in Raghunathanahalli-2 Micro watershed is presented in Table 31. The results indicate that, the farmers have annual gross expenditure of Rs. 374992.86 in micro-watershed, of which Rs. 17314.29 is from agriculture itself.

**Table 31. Average annual Expenditure in Raghunathanahalli-2 micro-watershed**

Sl.No.	Particulars	LL (0)	MF (15)	SF (12)	SMF (8)	All (35)
		Rs.	Rs.	Rs.	Rs.	Rs.
1	Service/salary	0	98750	100000	82500	18857.1
2	Wage	0	8000	6000	2000	857.14
3	Agriculture	0	15428.6	19600	27714.3	17314.3
4	Dairy Farm	0	7000	8000	0	857.14
Total		0	129179	133600	112214	374993

**Horticulture species grown:** The data regarding horticulture species grown in Raghunathanahalli-2 Micro watershed is presented in Table 32. The results indicate that, the total number of horticultural trees grown (both field and backyard) by the sampled households were coconut (12).

**Table 32. Horticulture species grown in Raghunathanahalli-2 micro-watershed**

Sl.No.	Particulars	LL (0)		MF (15)		SF (12)		SMF (8)		All (35)	
		F	B	F	B	F	B	F	B	F	B
1	Coconut	0	0	10	0	2	0	0	0	12	0

\*F= Field B=Back Yard

**Forest species grown:** The data regarding forest species grown in Raghunathanahalli-2 Micro watershed is presented in Table 33. The results indicate that, households have planted 1 teak trees, 92 neem trees, 4 acacia trees, 11 banyan trees together in both field and backyard.

**Table 33. Forest species grown in Raghunathanahalli-2 micro-watershed**

Sl.No.	Particulars	LL (0)		MF (15)		SF (12)		SMF (8)		All (35)	
		F	B	F	B	F	B	F	B	F	B
1	Teak	0	0	0	0	1	0	0	0	1	0
2	Neem	0	0	19	0	20	0	53	0	92	0
3	Acacia	0	0	4	0	0	0	0	0	4	0
4	Banyan	0	0	2	0	3	0	6	0	11	0

\*F= Field B=Back Yard

**Average additional investment capacity:** The data regarding average additional investment capacity in Raghunathanahalli-2 Micro watershed is presented in Table 34. The results indicate that, households have an average investment capacity of Rs. 1571.43 for land development, Rs. 142.86 for creation of irrigation facility, Rs.542.86 for adoption of improved livestock breeds, Rs.57.14 adoption of improved crop production activities.

**Table 34. Average additional investment capacity of households in Raghunathanahalli-2 micro-watershed**

Sl.No.	Particulars	LL (0)	MF (15)	SF (12)	SMF (8)	All (35)
		Rs.	Rs.	Rs.	Rs.	Rs.
1	Land development	0	1066.67	3250	0	1571.43
2	Irrigation facility	0	0	416.67	0	142.86
3	Improved crop production	0	333.33	1166.67	0	542.86
4	Improved livestock management	0	133.33	0	0	57.14

**Source of funds for additional investment:** The data regarding source of funds for additional investment in Raghunathanahalli-2 Micro watershed is presented in Table 35. The results indicate that, the sources of finance raised from bank as a loan and from own sources for land development were 25.71, for irrigation facility was 2.86 and 142.86 per cent.

**Table 35. Source of funds for additional investment in Raghunathanahalli-2 micro-watershed**

Sl.No	Item	Land development		Irrigation facility		Improved crop production		Improved livestock management	
		N	%	N	%	N	%	N	%
1	Loan from bank	9	25.71	1	2.86	4	11.43	1	2.86

**Marketing of agricultural produce:** The data regarding marketing of the agricultural produce in Raghunathanahalli-2 Micro watershed is presented in Table 36. The results indicated that, 100.00 percent of output of Bajra was sold in the market with average price of Rs. 1612.50; 76.92 percent of output of Bengal gram was sold in the market with average price of Rs. 5600.00; 100.00 percent of output of Groundnut was sold in the

market with average price of Rs. 3000.00; 100.00 percent of output of Maize was sold in the market with average price of Rs. 1475.00 and 100.00 percent of output of Red gram was sold in the market with average price of Rs. 3500.00.

**Table 36. Marketing of agricultural produce in Raghunathanahalli-2 micro-watershed**

Sl. No	Crops	Output obtained (q)	Output retained (q)	Output sold (q)	Output sold (%)	Avg. Price obtained (Rs/q)
1	Bajra	46	0	46	100	1613
2	Bengal gram (Kadale)	13	3	10	77	5600
3	Groundnut	45	0	45	100	3000
4	Maize	102	0	102	100	1475
5	Red gram	15	0	15	100	3500
6	Sorghum	68	0	68	100	1286
7	Sunflower	133	0	133	100	3027

**Marketing channels used for sale of agricultural produce:** The data regarding marketing channels used for sale of agricultural produce in Raghunathanahalli-2 Micro watershed is presented in Table 37. The results indicated that, 111.43 cent of the households have sold agricultural produce to the local/village merchants.

**Table 37. Marketing channels used for sale of agricultural produce in Raghunathanahalli-2 micro-watershed**

Sl.No.	Particulars	LL (0)		MF (15)		SF (12)		SMF (8)		All (35)	
		N	%	N	%	N	%	N	%	N	%
1	Local/village Merchant	0	0	15	100	15	125	9	113	39	111.4

**Mode of transport of agricultural produce:** The data regarding mode of transport of agricultural produce in Raghunathanahalli-2 Micro watershed is presented in Table 38. The results indicated that, 97.24 cent of the households have used tractor, 2.86 per cent have used Cart for the transport of agriculture commodity.

**Table 38. Mode of transport of agricultural produce in Raghunathanahalli-2 micro-watershed**

Sl.No.	Particulars	LL (0)		MF (15)		SF (12)		SMF (8)		All (35)	
		N	%	N	%	N	%	N	%	N	%
1	Cart	0	0	0	0	1	8.33	0	0	1	2.86
2	Tractor	0	0	15	100	14	117	9	113	34	97.24

**Table 39. Incidence of soil and water erosion problems in Raghunathanahalli-2 micro-watershed**

Sl. No.	Particulars	LL (0)		MF (15)		SF (12)		SMF (8)		All (35)	
		N	%	N	%	N	%	N	%	N	%
1	Soil and water erosion problems in the farm	0	0	7	47	11	91.7	3	38	21	60

**Incidence of soil and water erosion problems:** The data regarding incidence of incidence of soil and water erosion problems in Raghunathanahalli-2 Micro watershed is

presented in Table 39. The results indicate that, 60.00 per cent of the households have experienced soil and water erosion problems.

**Interest towards soil testing:** The data regarding Interest shown towards soil testing in Raghunathanahalli-2 Micro watershed is presented in Table 40. The results indicated that, 54.29 per cent of the households were interested towards soil testing.

**Table 40. Interest regarding soil testing in Raghunathanahalli-2 micro-watershed**

Sl.No.	Particulars	LL (0)		MF (15)		SF (12)		SMF (8)		All (35)	
		N	%	N	%	N	%	N	%	N	%
1	Interest in soil test	0	0	7	47	9	75	3	38	19	54.29

**Usage pattern of fuel for domestic use:** The data on usage pattern of fuel for domestic use in Raghunathanahalli-2 Micro watershed is presented in Table 41. The results indicated that, firewood was the major source of fuel for domestic use for 100 per cent of the households.

**Table 41. Usage pattern of fuel for domestic use in Raghunathanahalli-2 micro-watershed**

Sl.No.	Particulars	LL (0)		MF (15)		SF (12)		SMF (8)		All (35)	
		N	%	N	%	N	%	N	%	N	%
1	Fire Wood	0	0	15	100	12	100	8	100	35	100

**Source of drinking water:** The data on source of drinking water in Raghunathanahalli-2 Micro watershed is presented in Table 42. The results indicated that, tank supply of water was the major source for drinking water for 2.86 per cent of the households followed by piped waters supply (48.57 %), bore well water (40.00%) and open well water for (5.71%).

**Table 42. Source of drinking water in Raghunathanahalli-2 micro-watershed**

Sl.No.	Particulars	LL (0)		MF (15)		SF (12)		SMF (8)		All (35)	
		N	%	N	%	N	%	N	%	N	%
1	Piped supply	0	0	8	53.3	5	41.67	4	50	17	48.57
2	Bore Well	0	0	5	33.3	6	50	3	37.5	14	40
3	Open well	0	0	0	0	1	8.33	1	12.5	2	5.71
4	Lake/ Tank	0	0	1	6.67	0	0	0	0	1	2.86

**Source of light:** The data on source of light in Raghunathanahalli-2 Micro watershed is presented in Table 43. The results indicated that, electricity was the major source of light for 100.00 per cent of the households.

**Table 43. Source of light in Raghunathanahalli-2 micro-watershed**

Sl.No.	Particulars	LL (0)		MF (15)		SF (12)		SMF (8)		All (35)	
		N	%	N	%	N	%	N	%	N	%
1	Electricity	0	0	15	100	12	100	8	100	35	100

**Existence of sanitary toilet facility:** The data on availability of toilet facility in Raghunathanahalli-2 Micro watershed is presented in Table 44. The results indicated that, 54.29 per cent of the households possess toilets.

**Table 44. Existence of sanitary toilet facility in Raghunathanahalli-2 micro-watershed**

Sl.No.	Particulars	LL (0)		MF (15)		SF (12)		SMF (8)		All (35)	
		N	%	N	%	N	%	N	%	N	%
1	Sanitary toilet facility	0	0	10	67	5	41.67	4	50	19	54.3

**Possession of PDS card:** The data regarding possession of PDS card in Raghunathanahalli-2 Micro watershed is presented in Table 45. The results indicated that, 100.00 per cent of the households possessed BPL card.

**Table 45. Possession of PDS card in Raghunathanahalli-2 micro-watershed**

Sl.No.	Particulars	LL (0)		MF (15)		SF (12)		SMF (8)		All (35)	
		N	%	N	%	N	%	N	%	N	%
1	BPL	0	0	15	100	12	100	8	100	35	100

**Participation in NREGA programme:** The data regarding Participation in NREGA programme in Raghunathanahalli-2 Micro watershed is presented in Table 46. The results indicated that, only 17.14 percent of the households have participated in NREGA programme.

**Table 46. Participation in NREGA programme in Raghunathanahalli-2 micro-watershed**

Sl.No.	Particulars	LL (0)		MF (15)		SF (12)		SMF (8)		All (35)	
		N	%	N	%	N	%	N	%	N	%
1	Participation in NREGA programme	0	100	4	26.7	0	0	2	25	6	17.1

**Adequacy of food items:** The data regarding adequacy of food items in Raghunathanahalli-2 Micro watershed is presented in Table 47. The results indicated that, the extent of adequacy of food items for cereals, pulses, Oilseeds and vegetables were 100.00, 60.00, 22.86, 65.71 per cent respectively, similarly for milk (68.57%), Egg (51.43%), and Meat (54.29%).

**Table 47. Adequacy of food items in Raghunathanahalli-2 micro-watershed**

Sl.No.	Particulars	LL (0)		MF (15)		SF (12)		SMF (8)		All (35)	
		N	%	N	%	N	%	N	%	N	%
1	Cereals	0	0	15	100	12	100	8	100	35	100
2	Pulses	0	0	7	46.7	8	66.67	6	75	21	60
3	Oilseed	0	0	1	6.67	3	25	4	50	8	22.86
4	Vegetables	0	0	11	73.3	7	58.33	5	62.5	23	65.71
5	Milk	0	0	12	80	7	58.33	5	62.5	24	68.57
6	Egg	0	0	9	60	5	41.67	4	50	18	51.43
7	Meat	0	0	10	66.7	5	41.67	4	50	19	54.29

**Inadequacy of food items:** The data regarding in adequacy of food items in Raghunathanahalli-2 Micro watershed is presented in Table 48. The results indicated that, the extent of in adequacy of food items for pulses, Oilseeds and vegetables were 40.00, 74.29, 34.29 and 42.86 per cent respectively, similarly for fruits (100.00%), milk (31.43%), egg (48.57%) and meat (42.86%).



**Table 48. Inadequacy of food items in Raghunathanahalli-2 micro-watershed**

Sl.No.	Particulars	LL (0)		MF (15)		SF (12)		SMF (8)		All (35)	
		N	%	N	%	N	%	N	%	N	%
1	Pulses	0	0	8	53.3	4	33.33	2	25	14	40
2	Oilseed	0	0	13	86.7	9	75	4	50	26	74.29
3	Vegetables	0	0	4	26.7	5	41.67	3	37.5	12	34.29
4	Fruits	0	0	15	100	12	100	8	100	35	100
5	Milk	0	0	3	20	5	41.67	3	37.5	11	31.43
6	Egg	0	0	6	40	7	58.33	4	50	17	48.57
7	Meat	0	0	4	26.7	7	58.33	4	50	15	42.86

**Farming constraints:** The data regarding farming constraints experienced by households in Raghunathanahalli-2 Micro watershed is presented in Table 49. The results indicated that, lower fertility status of the soil was the constraint experienced by (100.00 %) per cent of the households, wild animal menace on farm field (51.43%), frequent incidence of pest and diseases (91.43%), inadequacy of irrigation water (2.86%), high cost of fertilizers and plant protection chemicals (91.43%), high rate of interest on credit (40.00%), low price for the agricultural commodities (88.57 %), lack of marketing facilities in the area (80.00%), inadequate extension services (5.71 %), lack of transport for safe transport of the agricultural produce to the market (74.29%).

**Table 49. Farming constraints experienced in Raghunathanahalli-2 micro-watershed**

SN	Particulars	LL (0)		MF (15)		SF (12)		SMF (8)		All (35)	
		N	%	N	%	N	%	N	%	N	%
1	Lower fertility status of the soil	0	0	15	100	12	100	8	100	35	100
2	Wild animal menace on farm field	0	0	6	40	8	66.67	4	50	18	51.43
3	Frequent incidence of pest and diseases	0	0	13	86.67	11	91.67	8	100	32	91.43
4	Inadequacy of irrigation water	0	0	0	0	1	8.33	0	0	1	2.86
5	High cost of Fertilizers and plant protection chemicals	0	0	13	86.67	11	91.67	8	100	32	91.43
6	High rate of interest on credit	0	0	6	40	5	41.67	3	37.5	14	40
7	Low price for the agricultural commodities	0	0	14	93.33	9	75	8	100	31	88.57
8	Lack of marketing facilities in the area	0	0	13	86.67	8	66.67	7	87.5	28	80
9	Inadequate extension services	0	0	1	6.67	1	8.33	0	0	2	5.71
10	Lack of transport for safe transport of the Agril produce to the market.	0	0	12	80	9	75	5	62.5	26	74.29

**SUMMARY AND IMPLICATIONS**

In order to assess the socio-economic condition of the farmers in the watershed 35 households located in the micro watershed were interviewed for the survey. The study was conducted in Raghunathanahalli-2 micro-watershed (Murlapura sub-watershed, Koppal taluk & District) is located at North latitude 15° 14' 31.598" and 15° 13' 30.436" and East longitude 75° 56' 30.124" and 75° 54' 44.824" covering an area of about 440.93 ha bounded by under Raghunathahalli, Gattareddyhala and Belagatti Villages.

Socio-economic analysis of Raghunathanahalli-2 micro watersheds of Murlapura sub-watershed, Koppal taluk & District indicated that, out of the total sample of 35 farmers were sampled in Raghunathanahalli-2 micro-watershed among households surveyed 15 (42.86%) were marginal, 12 (34.29%) were small and 8 (22.86 %) were semi medium farmers. The population characteristics of households indicated that, there were 96 (56.47%) men and 74 (43.53 %) were women. The average population of marginal farmers was 5.06, small farmers were 4.83 and semi medium farmers were 4.5. Majority of the respondents (47.06%) were in the age group of 16-35 years.

Education level of the sample households indicated that, there were 26.47 per cent illiterates, 68.83 per cent pre university education and 7.65 per cent attained graduation. About, 77.14 per cent of household heads practicing agriculture and 22.86 per cent of the household heads were engaged as agricultural labourers. Agriculture was the major occupation for 62.94 per cent of the household members.

In the study area, 97.14 per cent of the households possess katcha house. The durable assets owned by the households showed that, 77.14 per cent possess TV, 34.29 per cent possess mixer grinder, 77.14 per cent possess mobile phones and 28.57 per cent possess motor cycles. Farm implements owned by the households indicated that, 11.43 per cent of the households possess plough, 8.57 per cent possess bullock cart.

Regarding livestock possession by the households, 5.71 per cent possess local cow and 5.71 per cent possess buffalo. The average labour availability in the study area showed that, own labour men available in the micro watershed was 1.81, women available in the micro watershed was 1.44, hired labour (men) available was 6.58 and hired labour (women) available was 6.31.

In the study area, about 0.59 per cent of the respondents migrated from the micro watershed in search of jobs. Out of the total land holding of the sample respondents 85.19 per cent (44.86 ha) of the area is under dry condition and the remaining 5.56 per cent area is irrigated land. There were 2.00 live bore wells among the sampled households.

Bore well was the major source of irrigation for 5.71 per cent of the households. The major crops grown by sample farmers are Sunflower, Redgram, Sorghum, Groundnut

and Maize and cropping intensity was recorded as 83.72 per cent. Out of the sample households 8.57 percent possessed bank account.

About 8.57 per cent of the respondents borrowed credit from various sources. Per hectare cost of cultivation for Sunflower, Redgram, Sorghum, Groundnut and Maize was Rs.38349.84, 55753.72, 30816.36, 75904.68 and 30591.23 with benefit cost ratio of 1:0.80, 1: 2.10, 1: 0.50, 1: 1.60 and 1:0.70 respectively.

Further, 42.86 per cent of the households opined that dry fodder was adequate and 8.57 per cent of the households have opined that the green fodder was adequate. The average annual gross income of the farmers was Rs. 78168.00 in micro-watershed, of which Rs. 25131.43 comes from agriculture.

Sampled households have grown 12 horticulture trees and 108 forestry trees together in the fields and back yards. Households have an average investment capacity of Rs. 1571.43 for land development and Rs. 142.86 for irrigation facility. Source of funds for additional investment is concerned, 25.71 per cent depends on bank loan for land development activities.

Regarding marketing channels, 100 per cent of the households have sold agricultural produce to the local/village merchants. Further, 100 per cent of the households have used tractor for the transport of agriculture commodity. Majority of the farmers (60.00%) have experienced soil and water erosion problems in the watershed and 54.29 per cent of the households were interested towards soil testing.

Fire was the major source of fuel for domestic use for 100 per cent of the households. Piped supply was the major source for drinking water for 48.57 per cent of the households. Electricity was the major source of light for 100.00 per cent of the households.

In the study area, 54.29 per cent of the households possess toilet facility. Regarding possession of PDS card, 100.00 per cent of the households possessed BPL card. Households opined that, the requirement of cereals (100.00%), pulses (60.00%) and oilseeds (22.86%) are adequate for consumption.

Farming constraints experienced by households in the micro watersheds were lower fertility status of the soil (100.00%) wild animal menace on farm field (51.43%), frequent incidence of pest and diseases (91.43%), inadequacy of irrigation water (2.86%), high cost of fertilizers and plant protection chemicals (91.43%), high rate of interest on credit (40.00%), low price for the agricultural commodities (88.57%), lack of marketing facilities in the area (80.00%), inadequate extension services (5.71%), lack of transport for safe transport of the agricultural produce to the market (74.29%).

### **Implications of the survey**

- ✓ Result indicated that, there were 26.47 per cent were illiterate hence, extension methodologies such as demonstration, street play, drama, video shows will be effective in dissemination of the technologies in the micro watershed.
- ✓ The data indicate that, 97.14 per cent of the households possess katcha house. Hence, the development department while implementing the watershed plan should focus on agriculture to enhance the productivity of major crops in the area to increase the income of the farmers.
- ✓ Results indicated that the local institutional participation of the household members in the micro watershed is minimal hence, activities like membership campaign, awareness creation about the benefits of membership in local institutions and strengths of organized groups must be conveyed.
- ✓ Majority of the households in the watershed have experience in use of mobile phones, and television hence, these mass media can be effectively utilized for transfer of technology as well as for information dissemination.
- ✓ The farm machinery/implement possession in the micro watershed was found to be minimum the reasons may lack of knowledge or lack of financial ability which can be addressed through training on use of different farm implements, providing information on different sources of finance for purchase of farm implements.
- ✓ The possession of livestock such as crossbred cow found is less hence, farmers must be made aware of the benefits of crossbred cow in increased milk production.
- ✓ The possession of livestock such as sheep, goat and poultry was found to be low hence, farmers may be informed the role of subsidiary enterprises in enhancing the income and information on financial support for subsidiary activities.
- ✓ The data indicate that, job/work was the reason for all the migrants hence, farmers may be trained on profitable agriculture or self employment such as animal husbandry, plate making, sheep rearing, goat rearing, rabbit rearing with suitable information on sources of financial support.
- ✓ The results indicate that there was a change in quality of life due to migration hence, the developmental departments should take actions to arrest migration and to improve the quality of the life in rural areas.
- ✓ Households possess 38.22ha (85.19 %) of dry land and 2.49ha (5.56 %) of irrigated land hence, the availability of the dry land agricultural technologies such as short duration crops, high yielding drought resistance crop varieties, drip irrigation technology and subsidy information will be helpful for the farmers to enhance the productivity of land and as well as farmers income.
- ✓ Few of the bore well in micro watershed found non functional hence, farmers may be trained on possibility of bore well rejuvenation.
- ✓ Bore well was major source of irrigation for 5.71 per cent of the households. hence, in order to increase the area under irrigation as well as to increase the water use efficiency farmers may trained on drip irrigation and provide the information on

- subsidy for drip irrigation equipment's along with the information on different agencies which provides the financial assistance for drip irrigation.
- ✓ The cropping intensity in the micro watershed was found to be (83.72 %) hence, care must be taken by the implementing agency to bring uncultivated land into cultivation through suitable measures.
  - ✓ Many of the household members have borrowed loan from cooperative banks which has higher rate of interest hence, farmers may be sensitized on the different sources of credit with lesser interest rate such SHGs etc.
  - ✓ The results indicated the non availability of both green and dry fodder throughout the year hence, fodder development activities can be taken up in the micro watershed.
  - ✓ The average annual gross income of the households Rs.25131.43 from agriculture, Rs.0.00 from business and Rs. 4571.43 from wages and. Agriculture was found to be the major source of income for households hence; the development activities should focus on productivity enhancement, marketing arrangements and agricultural technology dissemination to have a direct impact on the farmers.
  - ✓ The cultivation of forest species is found minimal hence, information and production technology related to agro-forestry and integrated farming system.
  - ✓ The data indicated that, 60.00 per cent of the households have experienced soil and water erosion problems. Hence, those farmers who reported the soil and water erosion problems may be given attention while implementation of the watershed development plan.
  - ✓ The data indicated that, 54.29 per cent of the households have interest in soil testing hence, farmers must be provided with the information on various institutions which are involved in soil testing for the benefit of the farmers.
  - ✓ Except summer ploughing the adoption of other soil and water conservation structures is minimum hence, the farmers in the micro watershed should be sensitized on the use of different conservation structures for soil water conservation.
  - ✓ Cereals and pulses found be adequate for per cent of the households respectively hence, farm households and the farm women must be trained on importance of balanced nutrition and role of vegetable, milk, egg, meat in balanced diet.
  - ✓ Lower fertility status of the soil (100.00%), wild animal menace on farm field (51.43%), frequent incidence of pest and diseases (91.43%), high cost of fertilizers and plant protection chemicals (91.43%), high rate of interest on credit (40.00%), low price for the agricultural commodities (88.57%), lack of marketing facilities in the area (80.00%), inadequate extension services (5.71%), lack of transport for safe transport of the agricultural produce to the market (74.29%) were the major farming constraints experienced hence, these constraints must be addressed immediately for the welfare of the farmers. Awareness to be created among the farmers to approach nearest KVKs/RSKs and other developmental departments for technical and for subsidized inputs and utilize the well established regulated markets, approaching the contract firms, direct markets to avoid the involvement of middlemen.