



हर कदम, हर डगर
किसानों का हमसफर
भारतीय कृषि अनुसंधान परिषद

AgriSearch with a human touch

**LAND RESOURCE INVENTORY AND SOCIO-ECONOMIC STATUS OF
FARM HOUSEHOLDS FOR WATERSHED PLANNING AND
DEVELOPMENT**

HALIGERI-3 (4D4A1T1a) MICROWATERSHED

Koppal Taluk & District, Karnataka

Karnataka Watershed Development Project – II

SUJALA – III

World Bank funded Project



THE WORLD BANK



ICAR – NATIONAL BUREAU OF SOIL SURVEY AND LAND USE PLANNING



ICAR - NBSS & LUP

**WATERSHED DEVELOPMENT DEPARTMENT
GOVT. OF KARNATAKA, BANGALORE**



About ICAR - NBSS&LUP

The ICAR-National Bureau of Soil Survey and Land Use Planning (ICAR-NBSS&LUP), Nagpur, a premier Institute of the Indian Council of Agricultural Research (ICAR), was set up during 1976 with the objective to prepare soil resource maps at national, state and district levels and to provide research inputs in soil resource mapping and its applications, land evaluation, land use planning, land resource management, and database management using GIS for optimising land use on different kinds of soils in the country.

The Bureau has been engaged in carrying out soil resource survey, agro-ecological and soil degradation mapping at the country, state and district levels for qualitative assessment and monitoring the soil health towards viable land use planning. The research activities have resulted in identifying the soil potentials and problems, and the various applications of the soil surveys with the ultimate objective of sustainable agricultural development. The Bureau has the mandate to correlate and classify soils of the country and maintain a National Register of all the established soil series. The Institute is also imparting in-service training to staff of the soil survey agencies in the area of soil survey, land evaluation and soil survey interpretations for land use planning. The Bureau in collaboration with Panjabrao Krishi Vidyapeeth, Akola is running post-graduate teaching and research programme in land resource management, leading to M.Sc. and Ph.D. degrees.

Citation: Rajendra Hegde, Ramesh Kumar, S.C., K.V. Niranjana, S. Srinivas, M.Lalitha, B.A. Dhanorkar, R.S. Reddy, and S.K. Singh (2019). "Land Resource Inventory and socio-economic status of farm households for Watershed Planning and Development of Haligeri-3 (4D4A1T1a) Microwatershed, Koppal Taluk and District, Karnataka", ICAR-NBSS&LUP Sujala MWS Publ. 556, ICAR – NBSS & LUP, RC, Bangalore. P.155 & 29.

TO OBTAIN COPIES,

Please write to:

Director, ICAR - NBSS & LUP,

Amaravati Road, NAGPUR - 440 033, India

Phone : (0712) 2500386, 2500664, 2500545 (O)

Telefax : 0712-2522534

E-Mail : director@nbsslup.ernet.in

Website URL : nbsslup.in

Or

Head, Regional Centre, ICAR - NBSS&LUP, Hebbal, Bangalore - 560 024

Phone : (080) 23412242, 23510350 (O)

Telefax : 080-23510350

E-Mail : nbssrcb@gmail.com



**LAND RESOURCE INVENTORY AND SOCIO-ECONOMIC
STATUS OF FARM HOUSEHOLDS FOR WATERSHED
PLANNING AND DEVELOPMENT**

HALIGERI-3 (4D4A1T1a) MICROWATERSHED

Koppal Taluk & District, Karnataka

Karnataka Watershed Development Project – II

Sujala-III

World Bank funded Project



THE WORLD BANK



**ICAR – NATIONAL BUREAU OF SOIL SURVEY AND LAND
USE PLANNING**



ICAR - NBSS & LUP



**WATERSHED DEVELOPMENT DEPARTMENT, GOVT. OF
KARNATAKA, BANGALORE**



PREFACE

In Karnataka, as in other Indian States, the livelihoods of rural people are intertwined with farming pursuits. The challenges in agriculture are seriously threatening the livelihood of a large number of farmers as they have been practicing farming in contextual factors beyond their control. Climatic factors are the most important ones and have become much more significant in recent times due to rapid climate changes induced by intensive anthropogenic activities affecting our ecosystem in multiple ways. Climate change has become the reality, it is happening and efforts to evolve and demonstrate climate resilient technologies have become essential. Due to the already over stressed scenario of agrarian sector, the climate change is resulting in manifold increase in the complexities, pushing the rural mass to face more and more unpredictable situations. The rising temperatures and unpredictable rainfall patterns are going to test seriously the informed decisions farmers have to make in order to survive in farming and sustain their livelihood.

It is generally recognized that impacts of climate change shall not be uniform across the globe. It is said that impact of climate change is more severe in South Asia. Based on the analysis of meteorological data, it is predicted that in India, there will be upward trend in mean temperature, downward trend in relative humidity, annual rainfall and number of wet days in a year. Also, in general, phenomena like erratic monsoon, spread of tropical diseases, rise in sea levels, changes in availability of fresh water, frequent floods, droughts, heat waves, storms and hurricanes are predicted. Each one of these adverse situations are already being experienced in various parts of India and also at the global level. Decline in agricultural productivity of small and marginal farmers becoming more vulnerable is already witnessed.

In Karnataka, more than 60 per cent of the population live in rural areas and depend on agriculture and allied activities for their livelihood. Though the state has achieved significant progress in increasing the yield of many crops, there is tremendous pressure on the land resources due to the growing and competing demands of various land uses. This is reflected in the alarming rate of land degradation observed. Already more than 50 per cent of the area is affected by various forms of degradation. If this trend continues, the sustainability of the fragile ecosystem will be badly affected. The adverse effects of change in the climatic factors are putting additional stress on the land resources and the farmers dependent on this.

The natural resources (land, water and vegetation) of the state need adequate and constant care and management, backed by site-specific technological interventions and investments particularly by the government. Detailed database pertaining to the nature of

the land resources, their constraints, inherent potentials and suitability for various land based rural enterprises, crops and other uses is a prerequisite for preparing location-specific action plans, which are in tune with the inherent capability of the resources. Any effort to evolve climate resilient technologies has to be based on the baseline scientific database. Then only one can expect effective implementation of climate resilient technologies, monitor the progress, make essential review of the strategy, and finally evaluate the effectiveness of the implemented programs. The information available at present on the land resources of the state are of general nature and useful only for general purpose planning. Since the need of the hour is to have site-specific information suitable for farm level planning and detailed characterization and delineation of the existing land resources of an area into similar management units is the only option.

ICAR-NBSS&LUP, Regional Centre, Bangalore has taken up a project sponsored by the Karnataka Watershed Development Project-II, (Sujala-III), Government of Karnataka funded by the World Bank under Component -1 Land Resource Inventory. This study was taken up to demonstrate the utility of such a database in reviewing, monitoring and evaluating all the land based watershed development programs on a scientific footing. To meet the requirements of various land use planners at grassroots level, the present study on “Land Resource Inventory and Socio-Economic Status of Farm Households for Watershed Planning and Development of for Haligeri-3microwatershed in Koppal Taluk and District, Karnataka” for integrated development was taken up in collaboration with the State Agricultural Universities, IISC, KRSRAC, KSNDMC as Consortia partners. The project provides detailed land resource information at cadastral level (1:7920 scale) for all the plots and socio-economic status of farm households covering thirty per cent farmers randomly selected representing landed and landless class of farmers in the micro-watershed. The project report with the accompanying maps for the microwatershed will provide required detailed database for evolving effective land use plan, alternative land use options and conservation plans for the planners, administrators, agricultural extension personnel, KVK officials, developmental departments and other land users to manage the land resources in a sustainable manner.

It is hoped that this database will be useful to the planners, administrators and developmental agencies working in the area in not only for formulating location specific developmental schemes but also for their effective monitoring at the village/watershed level.

Nagpur

Date:15-11-2019

S.K. SINGH

Director, ICAR - NBSS&LUP, Nagpur

Contributors

| | |
|---|--|
| Dr. Rajendra Hegde Principal Scientist, Head & Project Leader, Sujala-III Project ICAR-NBSS&LUP, Regional Centre, Bangalore | Dr. S.K.Singh Director, ICAR-NBSS&LUP Coordinator, Sujala-III Project Nagpur |
| Soil Survey, Mapping & Report Preparation | |
| Dr. K.V. Niranjana | Sh. R.S. Reddy |
| Dr. B.A. Dhanorkar | Smt. Chaitra, S.P. |
| | Dr. Gopali Bardhan |
| | Mr. Somashekar T.N |
| | Ms. Arpitha G.M |
| | Dr. Mahendra kumar M.B |
| | |
| Field Work | |
| Sh. C. Bache Gowda | Sh. Mayur Patil |
| Sh. Somashekar | Sh. Arun Kumar, S. |
| Sh. M. Jayaramaiah | Sh. Sunil Raj |
| | Sh. Yogesh Kumar, B. |
| | Sh. Vikas, N.K. |
| | Sh. Arun Kumar, S.G. |
| | Sh. Umesh Jادیappa Madolli |
| | Sh. Praveen Kumar P. Achalkar |
| | Sh. Veerabhadraswamy |
| | Sh. Vinay |
| | Sh. Shankarappa, K. |
| | Sh. Lankesh, R.S. |
| | Sh. Appanna B. Hattigoudar |
| | Sh. Maharudra |
| GIS Work | |
| Dr. S.Srinivas | Sh. A.G.Devendra Prasad |
| Sh. D.H.Venkatesh | Sh. Abhijith Sastry, N.S. |
| Smt. K.Sujatha | Smt. Shyla, B. |
| Smt. K.V.Archana | Smt. Swetha ,K. |
| Sh. N.Maddileti | Ms. Vidya, P.C. |
| | Sh. Deepak, M.J. |
| | Smt. K.Karunya Lakshmi |
| | Ms. Seema, K.V. |

| Laboratory Analysis | |
|--|--|
| Dr. M. Lalitha | Sh. Vindhya, N.G. |
| Smt. Arti Koyal | Ms. P. Pavanakumari, P. |
| Smt. Parvathy, S. | Ms. Rashmi, N. |
| | Ms. Leelavathy, K.U. |
| | Smt. Usha Kiran, G. |
| | |
| Socio-Economic Analysis | |
| Dr. S.C. Ramesh Kumar | Sh. M.K. Prakashanaik |
| | Ms. Karuna V. Kulkarni |
| | Mrs. Sowmya A.N |
| | Sh. Vinod R |
| | Sh. Basavaraja |
| | Sh. Vijay Kumar Lamani |
| | Ms. Sowmya K.B |
| | Mrs. Prathibha, D.G |
| | Sh. Rajendra,D |
| | |
| Soil & Water Conservation | |
| Sh. Sunil P. Maske | |
| | |
| Watershed Development Department, GoK, Bangalore | |
| Sh. Prabhash Chandra Ray, IFS Project Director & Commissioner, WDD | Dr. A. Natarajan NRM Consultant, Sujala-III Project |
| Sh. A. Padmaya Naik, Director (In-Charge) Executive Director, KWDP-II, Sujala-III, WDD | |

PART-A
LAND RESOURCE INVENTORY

Contents

| | | |
|-------------------|---|----|
| Preface | | |
| Contributors | | |
| Executive Summary | | |
| Chapter 1 | Introduction | 1 |
| Chapter 2 | Geographical Setting | 3 |
| 2.1 | Location and Extent | 3 |
| 2.2 | Geology | 3 |
| 2.3 | Physiography | 4 |
| 2.4 | Drainage | 5 |
| 2.5 | Climate | 5 |
| 2.6 | Natural Vegetation | 6 |
| 2.7 | Land Utilization | 7 |
| Chapter 3 | Survey Methodology | 11 |
| 3.1 | Base maps | 11 |
| 3.2 | Image Interpretation for Physiography | 11 |
| 3.3 | Field Investigation | 14 |
| 3.4 | Soil mapping | 16 |
| 3.5 | Land Management Units | 17 |
| 3.5 | Laboratory Characterization | 17 |
| Chapter 4 | The Soils | 23 |
| 4.1 | Soils of Granite Gneiss Landscape | 23 |
| 4.2 | Soils of Alluvial Landscape | 31 |
| Chapter 5 | Interpretation for Land Resource Management | 53 |
| 5.1 | Land Capability Classification | 53 |
| 5.2 | Soil Depth | 55 |
| 5.3 | Surface Soil Texture | 56 |
| 5.4 | Soil Gravelliness | 57 |
| 5.5 | Available Water Capacity | 58 |
| 5.6 | Soil Slope | 59 |
| 5.7 | Soil Erosion | 60 |
| Chapter 6 | Fertility Status | 63 |
| 6.1 | Soil Reaction (pH) | 63 |
| 6.2 | Electrical Conductivity (EC) | 63 |
| 6.3 | Organic Carbon (OC) | 63 |
| 6.4 | Available Phosphorus | 63 |
| 6.5 | Available Potassium | 66 |
| 6.6 | Available Sulphur | 66 |
| 6.7 | Available Boron | 66 |
| 6.8 | Available Iron | 66 |
| 6.9 | Available Manganese | 66 |
| 6.10 | Available Copper | 66 |
| 6.11 | Available Zinc | 66 |

| | | |
|-----------|--|-----------|
| Chapter 7 | Land Suitability for Major Crops | 71 |
| 7.1 | Land suitability for Sorghum | 71 |
| 7.2 | Land suitability for Maize | 72 |
| 7.3 | Land suitability for Bajra | 73 |
| 7.4 | Land suitability for Redgram | 74 |
| 7.5 | Land suitability for Bengalgram | 75 |
| 7.6 | Land suitability for Groundnut | 76 |
| 7.7 | Land suitability for Sunflower | 77 |
| 7.8 | Land suitability for Cotton | 78 |
| 7.9 | Land suitability for Chilli | 79 |
| 7.10 | Land suitability for Tomato | 80 |
| 7.11 | Land suitability for Brinjal | 81 |
| 7.12 | Land suitability for Onion | 82 |
| 7.13 | Land suitability for Bhendi | 83 |
| 7.14 | Land suitability for Drumstick | 84 |
| 7.15 | Land suitability for Mulberry | 85 |
| 7.16 | Land suitability for Mango | 86 |
| 7.17 | Land Suitability for Sapota | 87 |
| 7.18 | Land suitability for Pomegranate | 88 |
| 7.19 | Land suitability for Guava | 89 |
| 7.20 | Land Suitability for Jackfruit | 90 |
| 7.21 | Land Suitability for Jamun | 91 |
| 7.22 | Land Suitability for Musambi | 92 |
| 7.23 | Land Suitability for Lime | 93 |
| 7.24 | Land Suitability for Cashew | 94 |
| 7.25 | Land Suitability for Custard apple | 95 |
| 7.26 | Land suitability for Amla | 96 |
| 7.27 | Land suitability for Tamarind | 97 |
| 7.28 | Land suitability for Marigold | 98 |
| 7.29 | Land suitability for Chrysanthemum | 99 |
| 7.30 | Land suitability for Jasmine | 100 |
| 7.31 | Land suitability for Crossandra | 101 |
| 7.32 | Land Management Units | 136 |
| 7.33 | Proposed Crop Plan | 137 |
| Chapter 8 | Soil Health Management | 141 |
| Chapter 9 | Soil and Water conservation Treatment Plan | 147 |
| 9.1 | Treatment Plan | 147 |
| 9.2 | Recommended Soil and Water Conservation measures | 151 |
| 9.3 | Greening of microwatershed | 152 |
| | References | 155 |
| | Appendix I | I-VIII |
| | Appendix II | IX-XVI |
| | Appendix III | XVII-XXIV |

LIST OF TABLES

| | | |
|------|--|-----|
| 2.1 | Mean Monthly Rainfall, PET, 1/2 PET at Koppal Taluk and District | 5 |
| 2.2 | Land Utilization in Koppal District | 7 |
| 3.1 | Differentiating Characteristics used for Identifying Soil Series | 15 |
| 3.2 | Soil map unit description of Haligeri-3 microwatershed | 17 |
| 4.1 | Physical and chemical characteristics of soil series identified in Haligeri-3 microwatershed | 36 |
| 7.1 | Soil-Site Characteristics of Haligeri-3 microwatershed | 103 |
| 7.2 | Land suitability for Sorghum | 105 |
| 7.3 | Land suitability for Maize | 106 |
| 7.4 | Land suitability for Bajra | 107 |
| 7.5 | Land suitability for Redgram | 108 |
| 7.6 | Land suitability for Bengalgram | 109 |
| 7.7 | Land suitability for Groundnut | 110 |
| 7.8 | Land suitability for Sunflower | 111 |
| 7.9 | Land suitability for Cotton | 112 |
| 7.10 | Land suitability for Chilli | 113 |
| 7.11 | Land suitability for Tomato | 114 |
| 7.12 | Land suitability for Brinjal | 115 |
| 7.13 | Land suitability for Onion | 116 |
| 7.14 | Land suitability for Bhendi | 117 |
| 7.15 | Land suitability for Drumstick | 118 |
| 7.16 | Land suitability for Mulberry | 119 |
| 7.17 | Land suitability for Mango | 120 |
| 7.18 | Land Suitability for Sapota | 121 |
| 7.19 | Land suitability for Pomegranate | 122 |
| 7.20 | Land suitability for Guava | 123 |
| 7.21 | Land suitability for Jackfruit | 124 |
| 7.22 | Land suitability for Jamun | 125 |
| 7.23 | Land Suitability for Musambi | 126 |
| 7.24 | Land Suitability for Lime | 127 |
| 7.25 | Land Suitability for Cashew | 128 |
| 7.26 | Land Suitability for Custard apple | 129 |

| | | |
|------|--|-----|
| 7.27 | Land Suitability for Amla | 130 |
| 7.28 | Land Suitability for Tamarind | 131 |
| 7.29 | Land Suitability for Marigold | 132 |
| 7.30 | Land Suitability for Chrysanthemum | 133 |
| 7.31 | Land suitability for Jasmine | 134 |
| 7.32 | Land suitability for Crossandra | 135 |
| 7.33 | Proposed Crop Plan for Haligeri-3 Microwatershed | 138 |

LIST OF FIGURES

| | | |
|------|---|----|
| 2.1 | Location map of Haligeri-3 Microwatershed | 3 |
| 2.2a | Granite and granite gneiss rocks | 4 |
| 2.2b | Alluvial rocks | 4 |
| 2.3 | Rainfall distribution in Koppal Taluk, Koppal District | 6 |
| 2.4 | Natural vegetation of Haligeri-3 microwatershed | 6 |
| 2.5 | Different crops and cropping systems in Haligeri-3 Microwatershed | 7 |
| 2.6 | Current Land use – Haligeri-3 Microwatershed | 9 |
| 2.7 | Location of Wells- Haligeri-3 Microwatershed | 9 |
| 3.1 | Scanned and Digitized Cadastral map of Haligeri-3 Microwatershed | 13 |
| 3.2 | Satellite image of Haligeri-3 Microwatershed | 13 |
| 3.3 | Cadastral map overlaid on IRS PAN+LISS IV merged imagery of Haligeri-3 Microwatershed | 14 |
| 3.4 | Location of profiles in a transect | 15 |
| 3.5 | Soil phase or management units of Haligeri-3 Microwatershed | 21 |
| 5.1 | Land Capability Classification of Haligeri-3 Microwatershed | 54 |
| 5.2 | Soil Depth map of Haligeri-3 Microwatershed | 55 |
| 5.3 | Surface Soil Texture map of Haligeri-3 Microwatershed | 57 |
| 5.4 | Soil Gravelliness map of Haligeri-3 Microwatershed | 58 |
| 5.5 | Soil Available Water Capacity map of Haligeri-3 Microwatershed | 59 |
| 5.6 | Soil Slope map of Haligeri-3 Microwatershed | 60 |
| 5.7 | Soil Erosion map of Haligeri-3 Microwatershed | 61 |
| 6.1 | Soil Reaction (pH) map of Haligeri-3 Microwatershed | 64 |
| 6.2 | Electrical Conductivity (EC) map of Haligeri-3 Microwatershed | 65 |
| 6.3 | Soil Organic Carbon (OC) map of Haligeri-3 Microwatershed | 65 |
| 6.4 | Soil Available Phosphorus map of Haligeri-3 Microwatershed | 67 |
| 6.5 | Soil Available Potassium map of Haligeri-3 Microwatershed | 67 |
| 6.6 | Soil Available Sulphur map of Haligeri-3 Microwatershed | 68 |
| 6.7 | Soil Available Boron map of Haligeri-3 Microwatershed | 68 |
| 6.8 | Soil Available Iron map of Haligeri-3 Microwatershed | 69 |
| 6.9 | Soil Available Manganese map of Haligeri-3 Microwatershed | 69 |
| 6.10 | Soil Available Copper map of Haligeri-3 Microwatershed | 70 |
| 6.11 | Soil Available Zinc map of Haligeri-3 Microwatershed | 70 |

| | | |
|------|--|-----|
| 7.1 | Land suitability map of Sorghum | 72 |
| 7.2 | Land suitability map of Maize | 73 |
| 7.3 | Land suitability map of Bajra | 74 |
| 7.4 | Land suitability map of Redgram | 75 |
| 7.5 | Land suitability map of Bengalgram | 76 |
| 7.6 | Land suitability map of Groundnut | 77 |
| 7.7 | Land suitability map of Sunflower | 78 |
| 7.8 | Land suitability map of Cotton | 79 |
| 7.9 | Land suitability map of Chilli | 80 |
| 7.10 | Land suitability map of Tomato | 81 |
| 7.11 | Land suitability map of Brinjal | 82 |
| 7.12 | Land suitability map of Onion | 83 |
| 7.13 | Land suitability map of Bhendi | 84 |
| 7.14 | Land suitability map of Drumstick | 85 |
| 7.15 | Land suitability map of Mulberry | 86 |
| 7.16 | Land suitability map of Mango | 87 |
| 7.17 | Land Suitability map of Sapota | 88 |
| 7.18 | Land suitability for Pomegranate | 89 |
| 7.19 | Land suitability map of Guava | 90 |
| 7.20 | Land Suitability map of Jackfruit | 91 |
| 7.21 | Land Suitability map of Jamun | 92 |
| 7.22 | Land Suitability map of Musambi | 93 |
| 7.23 | Land Suitability map of Lime | 94 |
| 7.24 | Land Suitability map of Cashew | 95 |
| 7.25 | Land Suitability map of Custard apple | 96 |
| 7.26 | Land suitability map of Amla | 97 |
| 7.27 | Land suitability map of Tamarind | 98 |
| 7.28 | Land suitability map of Marigold | 99 |
| 7.29 | Land suitability map of Chrysanthemum | 100 |
| 7.30 | Land suitability map of Jasmine | 101 |
| 7.31 | Land suitability map of Crossandra | 102 |
| 7.32 | Land Management Units map of Haligeri-3 microwatershed | 137 |
| 9.1 | Soil and water conservation map of Haligeri-3 microwatershed | 152 |

EXECUTIVE SUMMARY

The land resource inventory of Haligeri-3 microwatershed was conducted using village cadastral maps and IRS satellite imagery on 1:7920 scale. The false colour composites of IRS imagery were interpreted for physiography and these physiographic delineations were used as base for mapping soils. The soils were studied in several transects and a soil map was prepared with phases of soil series as mapping units. Random checks were made all over the area outside the transects to confirm and validate the soil map unit boundaries. The soil map shows the geographic distribution and extent, characteristics, classification, behavior and use potentials of the soils in the Microwatershed.

The present study covers an area of 802 ha in Koppal taluk and district, Karnataka. The climate is semiarid and categorized as drought - prone with an average annual rainfall of 662 mm, of which about 424 mm is received during south –west monsoon, 161 mm during north-east and the remaining 77 mm during the rest of the year. An area of about 98 per cent is covered by soil, <1 per cent by railways and <1 per cent by water bodies. The salient findings from the land resource inventory are summarized briefly below.

- ❖ *The soils belong to 18 soil series and 31 soil phases (management units) and 9 land management units.*
- ❖ *The length of crop growing period is <90 days and starts from 2nd week of August to 2nd week of November.*
- ❖ *From the master soil map, several interpretative and thematic maps like land capability, soil depth, surface soil texture, soil gravelliness, available water capacity, soil slope and soil erosion were generated.*
- ❖ *Soil fertility status maps for macro and micronutrients were generated based on the surface soil samples collected at every 320 m grid interval.*
- ❖ *Land suitability for growing 31 major agricultural and horticultural crops were assessed and maps showing the degree of suitability along with constraints were generated.*
- ❖ *Entire area is suitable for agriculture.*
- ❖ *About 6 per cent of the soils are shallow (25-50cm), 24 per cent of the soils are moderately shallow (50-75 cm), 34 per cent moderately deep (75- 100 cm) and 36 per cent is deep to very deep (100->150cm) soils.*
- ❖ *About 16 per cent loamy (sandy loam and sandy clay loam) and 84 per cent has clayey (sandy clay and clay) soils at the surface.*
- ❖ *About 58 per cent of the area has non-gravelly (<15%), 38 per cent gravelly (15-35%) and 3 per cent is very gravelly (35-60%) soils.*
- ❖ *With respect to available water capacity 8 per cent of the area has very low (<50mm/m), 38 per cent of the area has low (51-100 mm/m), 37 per cent medium*

(101-150 mm/m) and 16 per cent area is very high (>200mm/m) in available water capacity.

- ❖ An area of about 5 per cent has nearly level (0-1%) and 94 per cent has very gently sloping (1-3%) lands.
- ❖ An area of about 32 per cent is slightly eroded (e1) and 67 per cent is moderately eroded (e2).
- ❖ An area of about 5 per cent is neutral (pH 6.5 to 7.3), 61 per cent is slightly alkaline (pH 7.3 to 7.8), 31 per cent is strongly alkaline (pH 8.4-9.0) and 2 per cent is very strongly alkaline (pH>9.0).
- ❖ The Electrical Conductivity (EC) of the soils are $<2 \text{ dsm}^{-1}$ indicating that soils are non saline.
- ❖ Organic carbon is low (<0.5%) in 33 per cent, medium (0.5-0.75%) in 43 per cent and 24 per cent is high (>0.75%).
- ❖ Available phosphorus is low (<23 kg/ha) in 31 per cent, medium (<23 kg/ha) in 66 per cent and high (>57 kg/ha) in 2 per cent area of the soils.
- ❖ Available potassium is medium (145-337 kg/ha) in 15 per cent and high (>337 kg/ha) in 84 per cent area of the soils.
- ❖ Available sulphur is low (<10 ppm) in 37 per cent, medium (10-20 ppm) in 58 per cent and high (>20 ppm) in 4 per cent area of the soils.
- ❖ Available boron is low (<0.5 ppm) in 37 per cent, medium (0.5-1.0 ppm) in 56 per cent and high (>1.0 ppm) in 6 per cent area of the microwatershed.
- ❖ Available iron is deficient (<4.5ppm) in the entire area of the microwatershed.
- ❖ Available zinc is deficient (<0.6 ppm) in the entire area of the microwatershed.
- ❖ Available manganese and copper are sufficient in the entire area.
- ❖ The land suitability for 31 major agricultural and horticultural crops grown in the microwatershed was assessed and the areas that are highly suitable (class S1) and moderately suitable (class S2) are given below. It is however to be noted that a given soil may be suitable for various crops but what specific crop to be grown may be decided by the farmer looking to his capacity to invest on various inputs, marketing infrastructure, market price, and finally the demand and supply position.

Land suitability for various crops in the microwatershed

| Crop | Suitability Area in ha (%) | | Crop | Suitability Area in ha (%) | |
|-------------|----------------------------|--------------------------|---------------|----------------------------|--------------------------|
| | Highly suitable (S1) | Moderately suitable (S2) | | Highly suitable (S1) | Moderately suitable (S2) |
| Sorghum | 35(4) | 515(64) | Sapota | 35(4) | 234(29) |
| Maize | 35(4) | 514(64) | Pomegranate | 35(4) | 225(28) |
| Bajra | 35(4) | 560(70) | Guava | 35(4) | 234(29) |
| Redgram | 35(4) | 504(63) | Jackfruit | 35(4) | 234(16) |
| Bengal gram | - | 290(36) | Jamun | - | 161(20) |
| Groundnut | - | 293(37) | Musambi | 35(4) | 524(65) |
| Sunflower | 35(4) | 524(65) | Lime | 35(4) | 524(65) |
| Cotton | - | 549(69) | Cashew | 35(4) | 234(29) |
| Chilli | 35(4) | 515(64) | Custard apple | 35(4) | 535(72) |
| Tomato | 35(4) | 225(28) | Amla | 35(4) | 290(36) |
| Brinjal | 35(4) | 225(28) | Tamarind | - | 161(20) |
| Onion | 35(4) | 221(28) | Marigold | 35(4) | 511(64) |
| Bhendi | 35(4) | 225(28) | Chrysanthemum | 35(4) | 511(64) |
| Drumstick | 35(4) | 238(30) | Jasmine | 35(4) | 221(28) |
| Mulberry | 35(4) | 235(29) | Crossandra | 35(4) | 221(28) |
| Mango | - | 161(20) | - | - | - |

- ❖ *Apart from the individual crop suitability, a proposed crop plan has been prepared for the 9 identified LMUs by considering only the highly and moderately suitable lands for different crops and cropping systems with food, fodder, fibre and other horticulture crops.*
- ❖ *Maintaining soil-health is vital for crop production and conserve soil and land resource base for maintaining ecological balance and to mitigate climate change. For this, several ameliorative measures have been suggested to these problematic soils like saline/alkali, highly eroded, sandy soils etc.,*
- ❖ *Soil and water conservation and drainage line treatment plans have been prepared that would help in identifying the sites to be treated and also the type of structures required.*
- ❖ *As part of the greening programme, several tree species have been suggested to be planted in marginal and submarginal lands, field bunds and also in the hillocks, mounds and ridges. That would help in supplementing the farm income, provide fodder and fuel, and generate lot of biomass which in turn would help in maintaining the ecological balance and contribute to mitigating the climate change.*

INTRODUCTION

Land is a scarce resource and basic unit for any material production. It can support the needs of the growing population, provided they use the land in a rational and judicious manner. But what is happening in many areas of the state is a cause for concern to everyone involved in the management of land resources at the grassroots level. The area available for agriculture is about 51 per cent of the total area and more than 60 per cent of the people are still dependant on agriculture for their livelihood. The limited land area is under severe stress and strain due to increasing population pressure and competing demands of various land uses. Due to this, every year there is significant diversion of farm lands and water resources for non-agricultural purposes. Apart from this, due to lack of interest in farmers for farming, large tracts of cultivable lands are turning into fallows in many areas and this trend is continuing at an alarming rate.

Further, land degradation has emerged as a serious problem which has already affected about 38 lakh ha of cultivated area in the state. Soil erosion alone has degraded about 35 lakh ha. Almost all the uncultivated areas are facing various degrees of degradation, particularly soil erosion. Salinity and alkalinity has emerged as a major problem in more than 3.5 lakh ha in the irrigated areas of the state. Nutrient depletion and declining factor productivity is common in both rainfed and irrigated areas. The degradation is continuing at an alarming rate and there appears to be no systematic effort among the stakeholders to contain this process. In recent times, an aberration of weather due to climate change phenomenon has added another dimension leading to unpredictable situations to be tackled by the farmers.

In this critical juncture, the challenge before us is not only to increase the productivity per unit area which is steadily declining and showing a fatigue syndrome, but also to prevent or at least reduce the severity of degradation. If the situation is not reversed at the earliest, then the sustainability of the already fragile crop production system and the overall ecosystem will be badly affected in the state. The continued neglect and unscientific use of the resources for a long time has led to the situation observed at present in the state. It is a known fact and established beyond doubt by many studies in the past that the cause for all kinds of degradation is the neglect and irrational use of the land resources. Hence, there is urgent need to generate a detailed site-specific farm level database on various land resources for all the villages/watersheds in a time bound manner that would help to protect the valuable soil and land resources and also to stabilize the farm production.

Therefore, the land resource inventory required for farm level planning is the one which investigates not only the surface but also consider the other parameters which are critical for productivity *viz.*, soils, climate, water, minerals and rocks, topography, geology, hydrology, vegetation, crops, land use pattern, animal population, socio-

economic conditions, infrastructure, marketing facilities and various schemes and developmental works of the government etc. From the data collected at farm level, the specific problems and potentials of the area can be identified and highlighted, conservation measures required for the area can be planned on a scientific footing, suitability of the area for various uses can be worked out and finally viable and sustainable land use options suitable for each and every land holding can be prescribed.

The Land Resource Inventory is basically done for identifying potential and problem areas, developing sustainable land use plans, estimation of surface run off and water harvesting potential, preparation of soil and water conservation plans, land degradation/desertification etc. The Bureau is presently engaged in developing an LRI methodology using high resolution satellite remote sensing data and Digital Elevation Model (DEM) data to prepare Landscape Ecological Units (LEU) map representing agro-ecosystem as a whole. The LEU is preferred over landform as the base map for LRI. LEU is the assemblage of landform, slope and land use. An attempt was made to upscale the soil resource information from 1:250000 and 1:50000 scale to the LEU map in Goa and other states.

The land resource inventory aims to provide site-specific database for Haligeri-3 Microwatershed in Koppal Taluk, Koppal District, Karnataka State for the Karnataka Watershed Development Department. The database was generated by using cadastral map of the village as a base along with high resolution IRS LISS IV and Cartosat-1 merged satellite imagery. Later, an attempt will be made to uplink this LRI data generated at 1:7920 scale under Sujala-III Project to the proposed Landscape Ecological Units (LEUs) map.

The study was organized and executed by the ICAR- National Bureau of Soil Survey and Land Use Planning, Regional Centre, Bangalore under Generation of Land Resource Inventory Data Base Component-1 of the Sujala-III Project funded by the World Bank.

GEOGRAPHICAL SETTING

2.1 Location and Extent

The Haligeri-3 micro-watershed is located in the central part of Karnataka in Koppal taluk and district (Fig 2.1). It lies between $15^{\circ} 24'$ and $15^{\circ} 22'$ North latitudes and $76^{\circ} 4'$ and $76^{\circ} 2'$ East longitudes, and covers an area of about 802 ha. It comprises parts of Banapura, Thalabala, Thalakalla, Halageri and veerapura villages. It is about 14 km from Koppal town and is bounded by Halageri on the east, Thalakalla on the south, northeast, veerapura on the northeast and Banapura on the northern side of the microwatershed.




Fig.2.1 Location map of Haligeri-3 Microwatershed

2.2 Geology

Major rock formations observed in the microwatershed are granite gneiss and alluvium (Fig.2.2 a and b). Granite gneisses are essentially pink to gray and are coarse to medium grained. They consist primarily of quartz, feldspar, biotite and hornblende. The

gray granite gneisses are highly weathered, fractured and fissured upto a depth of about 10 m. Dolerite dykes and quartz veins are common with variable width and found to occur in Haligeri-3 village. The thickness of the alluvium generally is limited to less than a meter, except in river valleys where it is very deep extending to tens of meters. Such soils are transported and represent paleo black soils originally formed at higher elevation, but now occupying river valleys.



Fig.2.2 a Granite and granite gneiss rocks



Fig.2.2 b Alluvium

2.3 Physiography

Physiographically, the area has been identified as Granite gneiss and Alluvial landscapes based on geology. The microwatershed area has been further divided into mounds/ridges, summits, side slopes and very gently sloping uplands and nearly level

plains based on slope and its relief features. The elevation ranges from 520 to 544 m in the gently sloping uplands. The mounds and ridges are mostly covered by rock outcrops.

2.4 Drainage

The area is drained by several small seasonal streams that join Hire *halla* and Chenna *halla* along its course. Though, the streams are not perennial, during rainy season they carry large quantities of rain water. The microwatershed has only few small tanks which are not able to store the water flowing during the rainy season. Due to this, the ground water recharge is very much affected in the villages. This is reflected in the failure of many bore wells in the villages. If the available rain water is properly harnessed by constructing tanks and recharge structures at appropriate places in the villages, then the drinking and irrigation needs of the area can be easily met. The drainage network is dendritic to sub parallel.

2.5 Climate

The district falls under semiarid tract of the state and is categorized as drought - prone with total annual rainfall of 662 mm (Table 2.1). Of this, a maximum of 424 mm precipitation is received during south–west monsoon period from June to September, north-east monsoon contributes about 161 mm and prevails from October to early December and the remaining 77 mm is received during the rest of the year. The winter season is from December to February. During April and May, the temperatures reach up to 45°C and in December and January, the temperatures will go down to 16°C. Rainfall distribution is shown in Figure 2.3. The average Potential Evapo Transpiration (PET) is 145 mm and varies from a low of 101 mm in December to 193 mm in the month of May. The PET is always higher than precipitation in all the months except in the month of September. Generally, the Length of crop Growing Period (LGP) is <90 days and starts from 2nd week of August to 2nd week of November.

Table 2.1 Mean Monthly Rainfall, PET, 1/2 PET at Koppal Taluk and District

| Sl. no. | Months | Rainfall | PET | 1/2 PET |
|--------------|-----------|---------------|---------------|---------|
| 1 | January | 1.60 | 116.70 | 58.35 |
| 2 | February | 1.50 | 129.20 | 64.60 |
| 3 | March | 14.10 | 169.80 | 84.90 |
| 4 | April | 18.10 | 180.60 | 90.30 |
| 5 | May | 41.60 | 193.50 | 96.75 |
| 6 | June | 85.80 | 167.90 | 83.95 |
| 7 | July | 72.10 | 156.20 | 78.10 |
| 8 | August | 110.50 | 152.50 | 76.25 |
| 9 | September | 155.60 | 138.50 | 69.25 |
| 10 | October | 116.30 | 122.30 | 61.15 |
| 11 | November | 36.00 | 106.40 | 53.20 |
| 12 | December | 9.10 | 101.00 | 50.50 |
| TOTAL | | 662.30 | 144.55 | |




Fig. 2.3 Rainfall distribution in Koppal Taluk and District

2.6 Natural Vegetation

The natural vegetation is sparse comprising few tree species, shrubs and herbs. The mounds, ridges and boulders occupy sizeable areas which are under thin to moderately thick forest vegetation. Still, there are some remnants of the past forest cover which can be seen in patches in some ridges and hillocks in the microwatershed (Fig 2.4).

Apart from the continuing deforestation, the presence of large population of goats, sheep and other cattle in the microwatershed is causing vegetative degradation of whatever little vegetation left in the area. The uncontrolled grazing has left no time for the regeneration of the vegetative cover. This leads to the accelerated rate of erosion on the hill slopes, resulting in the formation of deep gullies in the foot slopes and eventually resulting in the heavy siltation of few tanks and reservoirs in the microwatershed.



Fig 2.4 Natural vegetation of Haligeri-3 Microwatershed

2.7 Land Utilization

About 91 per cent area (Table 2.2) in Koppal district is cultivated at present and about 17 per cent of the area is sown more than once. An area of about 3 per cent is currently barren. Forests occupy a small area of about 5 per cent and the tree cover is in a very poor state. Most of the mounds, ridges and boulder areas have very poor vegetative cover. Major crops grown in the area are sorghum, maize, bajra, cotton, safflower, sunflower, red gram, horse gram, onion, mulberry, pomegranate, sugarcane, bengalgram and groundnut (Fig 2.5 a and b). While carrying out land resource inventory, the land use/land cover particulars are collected from all the survey numbers and a current land use map of the microwatershed is prepared. The current land use map prepared shows the arable and non-arable lands, other land uses and different types of crops grown in the area. The current land use map of Haligeri-3 Microwatershed is presented in Fig.2.6. Simultaneously, enumeration of existing wells (bore wells) is made and their location in different survey numbers is marked on the cadastral map. Map showing the location of wells in Haligeri-3 Microwatershed is given in Fig 2.7.

Table 2.2 Land Utilization in Koppal District

| Sl. No. | Agricultural land use | Area (ha) | Per cent |
|---------|--------------------------|------------|----------|
| 1 | Total geographical area | 552495 | |
| 2 | Total cultivated area | 500542 | 90.6 |
| 3 | Area sown more than once | 92696 | 16.8 |
| 4 | Trees and groves | 210 | 0.04 |
| 5 | Cropping intensity | - | 118 |
| 6 | Forest | 29451 | 5.33 |
| 7 | Cultivable wasteland | 2568 | 0.46 |
| 8 | Permanent Pasture land | 14675 | 2.66 |
| 9 | Barren land | 16627 | 3.01 |
| 10 | Non agricultural land | 40591 | 7.35 |
| 11 | Current fallow | 19660 | 3.56 |



Fig.2.5 (a) Different crops and cropping systems in Haligeri-3 Microwatershed




Fig.2.5 (b) Different crops and cropping systems in Haligeri-3 Microwatershed




Fig.2.6 Current Land Use – Haligeri-3 Microwatershed




Fig.2.7 Location of wells– Haligeri-3 Microwatershed

SURVEY METHODOLOGY

The purpose of land resource inventory is to delineate similar areas (soil series and phases), which respond or expected to respond similarly for a given level of management. This was achieved in Haligeri-3 microwatershed by the detailed study of all the soil characteristics (depth, texture, colour, structure, consistence, coarse fragments, porosity, soil reaction, soil horizons etc.) and site characteristics (slope, erosion, drainage, occurrence of rock fragments etc.) followed by grouping of similar areas based on soil-site characteristics into homogeneous (management units) units and showing their extent and geographic distribution on the microwatershed cadastral map. The detailed soil survey at 1:7920 scale was carried out in 802 ha area. The methodology followed for carrying out land resource inventory was as per the guidelines given in Soil Survey Manual (IARI, 1971; Soil Survey Staff, 2006; Natarajan *et al.*, 2015) which is briefly described below.

3.1 Base Maps

The detailed survey of the land resources occurring in the microwatershed was carried out by using digitized cadastral map and satellite imagery as base supplied by the KRSRAC. The cadastral map shows field boundaries with their survey numbers, location of tanks, streams and other permanent features of the area (Fig. 3.1). Apart from the cadastral map, remote sensing data products from Cartosat-1 and LISS IV merged at the scale of 1:7920 were used in conjunction with the cadastral map to identify the geology, landscapes, landforms and other surface features. The imagery helped in the identification and delineation of boundaries between hills, uplands and lowlands, water bodies, forest and vegetated areas, roads, habitations and other cultural features of the area (Fig.3.2).The cadastral map was overlaid on the satellite imagery (Fig.3.3) that helps to identify the parcel boundaries and other permanent features. Apart from cadastral maps and images, toposheets of the area (1:50,000 scale) were used for initial traversing, identification of geology, landscapes and landforms, drainage features, present land use and also for selection of transects in the microwatershed.

3.2 Image Interpretation for Physiography

False Colour Composites (FCC) of Cartosat-I and LISS-IV merged satellite data covering the microwatershed area was visually interpreted using image interpretation elements and all the available collateral data with local knowledge. The delineated physiographic boundaries were transferred on to a cadastral map overlaid on satellite imagery. Physiographically, the area has been identified as granite gneiss and alluvial landscapes and is divided into landforms such as ridges, mounds and uplands based on slope. They were further subdivided into physiographic/ image interpretation units based on image characteristics. The image interpretation legend for physiography is given below.

Image Interpretation Legend for Physiography

G- Granite gneiss landscape

- G1 Hills/ Ridges/ Mounds
 - G11 Summits
 - G12 Side slopes
 - G121 Side slopes with dark grey tones
- G2 Uplands
 - G21 Summits
 - G22 Gently sloping uplands
 - G221 Gently sloping uplands, yellowish green (eroded)
 - G222 Gently sloping uplands, yellowish white (severely eroded)
 - G23 Very gently sloping uplands
 - G231 Very gently sloping uplands, yellowish green
 - G232 Very gently sloping uplands, medium green and pink
 - G233 Very gently sloping uplands, pink and green (scrub land)
 - G234 Very gently sloping uplands, medium greenish grey
 - G235 Very gently sloping uplands, yellowish white (eroded)
 - G236 Very gently sloping uplands, dark green
 - G237 Very gently sloping uplands, medium pink (coconut garden)
 - G238 Very gently sloping uplands, pink and bluish white (eroded)

DSe -Alluvial landscape

DSe 1 Summit

- DSe 11 Nearly level Summit with dark grey tone
- DSe 12 Nearly level Summit with medium grey tone
- DSe 13 Nearly level Summit with whitish grey tone
- DSe 14 Nearly level Summit with whitish tone (Calcareousness)
- DSe 15 Nearly level Summit with pinkish grey tone
- DSe 16 Nearly level Summit with medium pink tone
- DSe 17 Nearly level Summit with bluish white tone
- DSe 18 Nearly level Summit with greenish grey tone

DSe 2 Very gently sloping

- DSe 21 Very gently sloping, whitish tone
- DSe 22 Very gently sloping, greyish pink tone
- DSe 23 Very gently sloping, whitish grey tone
- DSe 24 Very gently sloping, medium grey tone
- DSe 25 Very gently sloping, medium pink tone
- DSe 26 Very gently sloping, dark grey tone
- DSe 27 Very gently sloping, bluish grey tone
- DSe 28 Very gently sloping, greenish grey tone
- DSe 29 Very gently sloping, Pinkish grey




Fig 3.1 Scanned and Digitized Cadastral map of Haligeri-3 Microwatershed




Fig.3.2 Satellite Image of Haligeri-3 Microwatershed




Fig.3.3 Cadastral map overlaid on IRS PAN+LISS IV merged imagery of Haligeri-3 Microwatershed

3.3 Field Investigation

The field boundaries and survey numbers given on the cadastral sheet were located on the ground by following permanent features like roads, cart tracks, *nallas*, streams, tanks etc., and wherever changes were noticed, they were incorporated on the microwatershed cadastral map. Preliminary traverse of the microwatershed was carried out with the help of cadastral map, imagery and toposheets. While traversing, landforms and physiographic units identified were checked and preliminary soil legend was prepared by studying soils at few selected places. Then, intensive traversing of each physiographic unit like hills, ridges, uplands and plains was carried out. Based on the variability observed on the surface, transects (Fig 3.4) were selected across the slope covering all the landform units in the microwatershed (Natarajan and Dipak Sarkar, 2010).




Fig: 3.4. Location of profiles in a transect

In the selected transect, soil profiles (Fig.3.4) were located at closely spaced intervals to take care of any change in the land features like break in slope, erosion, gravel, stones etc. In the selected sites, profiles (vertical cut showing the soil layers from surface to the rock) were opened upto 200 cm or to the depth limited by rock or hard substratum and studied in detail for all their morphological and physical characteristics. The soil and site characteristics were recorded for all profile sites on a standard proforma as per the guidelines given in USDA Soil Survey Manual (Soil Survey Staff, 2012). Apart from the transect study, profiles were also studied at random, almost like in a grid pattern, outside the transect areas to validate the soil map unit boundaries.

Based on the soil characteristics, the soils were grouped into different soil series. Soil series is the most homogeneous unit having similar horizons and properties and behaves similarly for a given level of management. Soil depth, texture, colour, kind of horizon and horizon sequence, amount and nature of gravel present, calcareousness, nature of substratum etc, were used as the major differentiating characteristics for identifying soil series occurring in the area. The differentiating characteristics used for identifying the soil series are given in Table 3.1. Based on the above characteristics, 18 soil series were identified in Haligeri-3 microwatershed.

**Table 3.1 Differentiating Characteristics used for identifying Soil Series
(Characteristics are of Series Control Section)**

| Sl. No | Soil Series | Depth (cm) | Colour (moist) | Texture | Gravel (%) | Horizon sequence | Calcareousness |
|--|----------------------------|------------|---------------------------------|---------|------------|------------------|----------------|
| Soils of Granite Gneiss Landscape | | | | | | | |
| 1 | Chikkasavanur (CSR) | 25-50 | 7.5YR3/2,2/3,3/4 | scl | <15 | Ap-Bw-Cr | - |
| 2 | Harve (HRV) | 25-50 | 2.5YR3/4,3/6 5YR3/3,4/4,3/4 | gscl | >35 | Ap-Bt-Cr- | - |
| 3 | Thammadahalli (TDH) | 50-75 | 2.5YR2.5/4,3/6 | sc-c | <15 | Ap-Bt-Cr | - |
| 4 | Lakkur (LKR) | 50-75 | 2.5YR 2.5/3, 2.5/4, 3/4, 3/6 | gsc | 40-60 | Ap-Bt-Bc-Cr | - |

| Sl. No | Soil Series | Depth (cm) | Colour (moist) | Texture | Gravel (%) | Horizon sequence | Calcareousness |
|------------------------------------|---------------------|------------|---|---------|------------|------------------|----------------|
| 5 | Mukhadahalli (MKH) | 50-75 | 5YR3/3,3/4,4/3, 5/4,6/6 2.5YR3/4 | gsc | >35 | Ap-Bt-Cr | - |
| 6 | Bidanagere (BDG) | 75-100 | 5YR3/3,3/4,4/3,5/4 2.5YR3/4 | gc | 35-60 | Ap-Bt-Cr | - |
| 7 | Hooradhahalli (HDH) | 75-100 | 2.5YR2.5/4,3/4, 3/6 | gsc-gc | >35 | Ap-Bt-Cr | - |
| 8 | Bisarahalli (BSR) | 75-100 | 5 YR 3/3, 3/4 | gsc | 15-35 | Ap-Bt-Cr | - |
| 9 | Balapur (BPR) | 100-150 | 2.5YR2.5/4,3/4 | gsc-gc | >35 | Ap-Bt-Cr | - |
| 10 | Nagalapur (NGP) | 100-150 | 5YR2.5/2,3/2, 2.5YR3/6,4/6 | gsc | >35 | Ap-Bt-Cr | - |
| 11 | Giddadapalya (GDP) | 100-150 | 2.5YR3/4, 3/6 | gsc-gc | 35-60 | Ap-Bt-Cr | - |
| Soils of Alluvial Landscape | | | | | | | |
| 12 | Muttal (MTL) | 25-50 | 10YR3/2,3/3,4/2 7.5YR3/2,3/3,6/4 | gc | 15-35 | Ap-Bw-Ck | e-ev |
| 13 | Ravanaki (RNK) | 50-75 | 7.5YR3/2,3/3,5/2,5/3 10YR3/1,3/2,4/1, 4/2, 5/1,6/1 | c | <15 | Ap-Bw-Cr | e-ev |
| 14 | Dambarahalli (DRL) | 75-100 | 10YR 2/1, 3/1, 4/3 | c | <15 | Ap-Bss-Ck | e-es |
| 15 | Narasapura (NSP) | 75-100 | 10 YR 3/1, 3/2, 4/2, | c | - | Ap-Bw-Cr | e-es |
| 16 | Kavalur (KVR) | 100-150 | 10 YR 2/2, 3/1, 3/2, 3/3, 4/4 | c | - | Ap-Bss-Bck-Cr | es-ev |
| 17 | Lakshmgudda (LGD) | 100-150 | 10YR3/1,3/2,4/1,4/2,7.5YR3/1,3/2,5/1, 2.5Y5/2,5/3,6/3 | c | <15 | Ap-Bss-Ck | es |
| 18 | Alawandi (AWD) | >150 | 10 YR 2/1, 3/2, | c | <15 | Ap-Bss | e-es |

3.4 Soil Mapping

The area under each soil series was further separated into soil phases and their boundaries delineated on the cadastral map based on the variations observed in the texture of the surface soil, slope, erosion, presence of gravel, stoniness etc. A soil phase is a subdivision of soil series based mostly on surface features that affect its use and management. The soil mapping units are shown on the map (Fig.3.5) in the form of symbols. During the survey many soil profile pits, few mini pits and a few auger bores representing different landforms occurring in the microwatershed were studied. In addition to the profile study, spot observations in the form of mini pits, road cuts, terrace cuts etc., were studied to validate the soil boundaries on the soil map.

The soil map shows the geographic distribution of 31 mapping units representing 18 soil series occurring in the microwatershed. The soil map unit (soil legend) description is presented in Table 3.2. The soil phase map (management units) shows the distribution of 31 phases mapped in the microwatershed. Each mapping unit (soil phase) delineated on the map has similar soil and site characteristics. In other words, all the farms or survey numbers included in one soil phase will have similar management needs and have to be treated accordingly.

3.5 Land Management Units

The 31 soil phases identified and mapped in the microwatershed were regrouped into nine Land Management Units (LMU's) for the purpose of preparing a Proposed Crop Plan for sustained development of the microwatershed. The database (soil phases) generated under LRI was utilized for identifying Land Management Units (LMU's) based on the management needs. One or more than one soil site characteristic having influence on the management have been chosen for identification and delineation of LMUs. For Haligeri-3 microwatershed, five soil and site characteristics, namely the soil depth, soil texture, slope, erosion and gravel content have been considered for defining LMUs. The land management units are expected to behave similarly for a given level of management.

3.5 Laboratory Characterization

Soil samples for each soil series soil were collected from representative master profiles for laboratory characterization by following the methods outlined in the Laboratory Manual (Sarma *et al*, 1987). Surface soil samples collected in the year 2018 from farmer's fields in Haligeri-3 microwatershed for fertility status (major and micronutrients) at 320 m grid interval were analyzed in the laboratory (Katyal and Rattan, 2003). By linking the soil fertility data to the survey numbers through GIS, soil fertility maps were generated using Kriging method for the microwatershed.

Table 3.2 Soil map unit description of Haligeri-3 Microwatershed

| Soil map unit No* | Soil Series | Soil Phase Symbol | Mapping Unit Description | Area in ha (%) |
|--|-------------|---|---|-------------------------------|
| Soils of Granite gneiss Landscape | | | | |
| | CSR | Chikkasavanur soils are shallow (25-50 cm), well drained, have dark brown to light yellowish brown, sandy clay loam soils occurring on nearly level to very gently sloping uplands under cultivation | | 13 (1.68) |
| 39 | | CSRiB2 | Sandy clay surface, slope 1-3%, moderate erosion | 13 (1.68) |
| | HRV | Harve soils are shallow (25-50 cm), well drained, dark red to dark reddish brown, red gravelly sandy clay loam soils occurring on nearly level to gently sloping uplands under cultivation | | 6 (0.7) |
| 29 | | HRViA1g1 | Sandy clay surface, slope 0-1%, slight erosion, gravelly (15-35%) | 6 (0.7) |
| | TDH | Thammadahalli soils are moderately shallow (50-75cm), well drained, have brown to very dark brown and dark reddish brown sandy clay to clay soils occurring on nearly level to gently sloping uplands | | 10.17 (1.26) |
| 59 | | TDHhB2g2 | Sandy clay loam surface, slope 1-3%, moderate erosion, very gravelly (35-60%) | 0.17 (0.02) |
| 60 | | TDHiB1 | Sandy clay surface, slope 1-3%, slight erosion | 10 (1.24) |

| Soil map unit No* | Soil Series | Soil Phase Symbol | Mapping Unit Description | Area in ha (%) |
|-------------------|-------------|-------------------|---|-----------------------------|
| | LKR | | Lakkur soils are moderately shallow (50-75 cm), well drained, have dark reddish brown to dark red, gravelly sandy clay soils occurring on very gently to moderately sloping uplands under cultivation | 17 (2.15) |
| 43 | | LKRcB2g1 | Sandy loam surface, slope 1-3%, moderate erosion, gravelly (15-35%) | 17 (2.15) |
| | MKH | | Mukhadahalli soils are moderately shallow (50-75 cm), well drained, have dark brown to reddish brown red, gravelly sandy clay soils occurring on gently very gently to gently sloping uplands under cultivation | 28 (3.5) |
| 90 | | MKHiB2g1 | Sandy clay surface, slope 1-3%, moderate erosion, gravelly (15-35%) | 28 (3.5) |
| | BDG | | Bidanagere soils are moderately deep (75-100 cm), well drained, have dark reddish brown, red gravelly clay soils occurring on nearly level to gently sloping uplands under cultivation | 31 (3.9) |
| 192 | | BDGiB1g1 | Sandy clay surface, slope 1-3%, slight erosion, gravelly (15-35%) | 31 (3.9) |
| | HDH | | Hooradhahalli soils are moderately deep (75-100 cm), well drained, have dark red to dark reddish brown, red gravelly sandy clay to clay soils occurring on nearly level to moderately sloping uplands under cultivation | 73 (9.15) |
| 109 | | HDHcB1g1 | Sandy loam surface, slope 1-3%, slight erosion, gravelly (15-35%) | 16 (1.97) |
| 123 | | HDHhB2g1 | Sandy clay loam surface, slope 1-3%, moderate erosion, gravelly (15-35%) | 19 (2.42) |
| 125 | | HDHiB1 | Sandy clay surface, slope 1-3%, slight erosion | 20 (2.49) |
| 128 | | HDHiB2g1 | Sandy clay surface, slope 1-3%, moderate erosion, gravelly (15-35%) | 18 (2.27) |
| | BSR | | Bisarahalli soils are moderately deep (75-100 cm), well drained, have dark reddish brown, red gravelly sandy clay soils occurring on very gently sloping uplands under cultivation | 4 (0.45) |
| 166 | | BSRiB1g2 | Sandy clay surface, slope 1-3%, slight erosion, very gravelly (35-60%) | 4 (0.45) |
| | BPR | | Balapur soils are deep (100-150 cm), well drained, have dark reddish brown to dark red gravelly sandy clay to clay soils occurring on nearly level to gently sloping uplands under cultivation | 90 (11.96) |
| 220 | | BPRcA1 | Sandy loam surface, slope 0-1%, slight erosion | 9 (1.11) |
| 231 | | BPRhB2g1 | Sandy clay loam surface, slope 1-3%, moderate erosion, gravelly (15-35%) | 24 (3.03) |
| 232 | | BPRhB2g2 | Sandy clay loam surface, slope 1-3%, moderate erosion, very gravelly (35-60%) | 20 (2.47) |

| Soil map unit No* | Soil Series | Soil Phase Symbol | Mapping Unit Description | Area in ha (%) |
|------------------------------------|-------------|--|---|-----------------------------|
| 239 | | BPRiB2 | Sandy clay surface, slope 1-3%, moderate erosion | 14 (1.74) |
| 240 | | BPRmB2 | Clay surface, slope 1-3%, moderate erosion | 23 (2.91) |
| | NGP | Nagalapur soils are deep (100-150 cm), well drained, have dark reddish brown to dark red, gravelly sandy clay soils occurring on nearly level to gently sloping uplands under cultivation | | 36 (4.47) |
| 255 | | NGPhA1 | Sandy clay loam surface, slope 0-1%, slight erosion | 18 (2.25) |
| 262 | | NGPiB1 | Sandy clay surface, slope 1-3%, slight erosion | 18 (2.22) |
| | GDP | Giddadapalya soils are deep (100-150 cm), well drained, have dark reddish brown to dark red, gravelly sandy clay to clay soils occurring on very gently sloping uplands under cultivation | | 35 (4.38) |
| 267 | | GDPcB2 | Sandy loam surface, slope 1-3%, moderate erosion | 2 (0.29) |
| 269 | | GDPiB2 | Sandy clay surface, slope 1-3%, moderate erosion | 33 (4.09) |
| Soils of Alluvial Landscape | | | | |
| | MTL | Muttal soils are shallow (25-50 cm), well drained, have very dark grayish brown to dark brown, calcareous black gravelly clay soils occurring on nearly level to gently sloping plains under cultivation | | 25 (3.16) |
| 311 | | MTLmB2g1 | Clay surface, slope 1-3%, moderate erosion, gravelly (15-35%) | 25 (3.16) |
| | RNK | Ravanaki soils are moderately shallow (50-75 cm), moderately well drained, have dark brown to very dark grayish brown and dark gray, calcareous black clay soils occurring on nearly level to very gently sloping plains under cultivation | | 134 (16.7) |
| 336 | | RNKmB2 | Clay surface, slope 1-3%, moderate erosion | 57 (7.11) |
| 337 | | RNKmB2g1 | Clay surface, slope 1-3%, moderate erosion, gravelly (15-35%) | 77 (9.59) |
| | DRL | Dambarahalli soils are moderately deep (75-100 cm), moderately well drained, have dark brown to very dark gray, calcareous black cracking clay soils occurring on nearly level to very gently sloping plains under cultivation | | 163 (20.3) |
| 350 | | DRLmB2 | Clay surface, slope 1-3%, moderate erosion | 118 (14.74) |
| 351 | | DRLmB2g1 | Clay surface, slope 1-3%, moderate erosion, gravelly (15-35%) | 45 (5.56) |
| | NSP | Narasapura soils are moderately deep (75-100 cm), | | 1 |

| Soil map unit No* | Soil Series | Soil Phase Symbol | Mapping Unit Description | Area in ha (%) |
|-------------------|-------------|-------------------|--|------------------------|
| | | | moderately well drained, have dark grayish brown to very dark grayish brown and very dark gray, sodic black calcareous clay soils occurring on nearly level to very gently sloping plains under cultivation | (0.11) |
| 362 | | NSPmB2 | Clay surface, slope 1-3%, moderate erosion | 1 (0.11) |
| | KVR | | Kavalur soils are deep (100-150 cm), moderately well drained, have dark yellowish brown to very dark grayish brown, calcareous cracking black clay soils occurring on nearly level to very gently sloping plains under cultivation | 20 (2.47) |
| 386 | | KVRmA1 | Clay surface, slope 0-1%, slight erosion | 5 (0.59) |
| 388 | | KVRmB1 | Clay surface, slope 1-3%, slight erosion | 15 (1.88) |
| | LGD | | Lakshangudda soils are deep (100-150 cm), well drained, have light olive brown to very dark gray calcareous clayey soils occurring on nearly level to very gently sloping plains under cultivation | 104 (12.93) |
| 393 | | LGDmB1 | Clay surface, slope 1-3%, slight erosion | 104 (12.93) |
| 424 | AWD | | Alawandi soils are very deep (>150 cm), moderately well drained, have very dark grayish brown to black , calcareous black cracking clay soils occurring on nearly level to very gently sloping plains under cultivation | 4 (0.48) |
| | | AWDmB2 | Clay surface, slope 1-3%, moderate erosion | 4 (0.48) |
| 992 | | Railway | | 1 (0.15) |
| 1000 | | Others | Water body | 6 (0.79) |

*Soil map unit numbers are continuous for the taluk, not the microwatersheds




Fig 3.5 Soil Phase or Management Units- Haligeri-3 Microwatershed

THE SOILS

Detailed information pertaining to the nature, extent and distribution of different kinds of soils occurring in Haligeri-3 microwatershed is provided in this chapter. The microwatershed area has been identified as granite gneiss and alluvial landscapes based on geology. In all, 18 soil series were identified. Soil formation is the result of the combined effect of environmental and terrain factors that are reflected in soil morphology. The soil formation is dominantly influenced by the parent material, climate, time and relief.

A brief description of each of the 18 soil series identified followed by 31 soil phases (management units) mapped (Fig. 3.5) are furnished below. The physical and chemical characteristics of soil series identified in Haligeri-3 microwatershed are given in Table 4.1 along with soil classification. The soils in any one map unit differ from place to place in their depth, texture, slope, gravelliness, erosion or any other site characteristic that affect management. The soil phase map can be used for identifying the suitability of areas for growing specific crops or for other alternative uses and also for deciding the type of conservation structures needed. The detailed information on soil and site-characteristics like soil depth, surface soil texture, slope, erosion, gravelliness, AWC, LCC etc, with respect to each of the soil phase identified is given village/survey number wise for the microwatershed in Appendix-I.

4.1 Soils of Granite gneiss Landscape

In this landscape, 11 soil series were identified and mapped. Of these series, Balapur (BPR) series occupies a maximum area of 90 ha (12%) and followed by Hooradhahalli (HDH) and others occupy minor area. The brief description of the soil series along with the soil phases identified and mapped is given below.

4.1.1 Chikkasavanur (CSR) Series: Chikkasavanur soils are shallow (25-50 cm), well drained, have dark brown to light yellowish brown sandy clay loam soils. They have developed from weathered granite gneiss and occur on very gently sloping uplands. The Chikkasavanur series has been classified as a member of the loamy, mixed, isohyperthermic family of (Paralithic) Haplustepts.

The thickness of the solum ranges from 32 to 49 cm. The thickness of A horizon ranges from 12 to 23 cm. Its colour is in 7.5 YR and 10 YR hue with value 2.5 to 4 and chroma 3 to 6. The texture varies from sandy loam to clay with 10 to 20 per cent gravel. The thickness of B horizon ranges from 16 to 32 cm. Its colour is in 7.5 YR and 5 YR hue with value 3 and chroma 2 to 4. Its texture is sandy clay loam with gravel content of < 15 per cent. The available water capacity is low (50-100 mm/m). One soil phase was identified and mapped.



Landscape and soil profile characteristics of Chikkasavanur (CSR) Series

4.1.2 Harve (HRV) Series: Harve soils are shallow (25-50 cm), well drained, have reddish brown to dark red gravelly sandy clay loam soils. They have developed from weathered granite gneiss and occur on very gently to moderately sloping uplands. The Harve series has been classified as a member of the loamy-skeletal, mixed isohyperthermic family of (Paralithic) Rhodustalfs.

The thickness of the solum ranges from 28 to 48 cm. The thickness of A-horizon ranges from 12 to 17 cm. Its colour is in 5YR and 2.5 YR hue with value 3 to 4 and chroma 4 to 6. The texture varies from loamy sand to sandy loam with 20 to 60 per cent gravel. The thickness of B-horizon ranges from 16 to 32 cm. Its colour is in 2.5 YR and 5 YR hue with value 3 to 4 and chroma 4 to 6. Its texture is sandy clay loam with gravel content of more than 35 per cent. The available water capacity is very low (<50mm/m). One soil phase was identified and mapped.



Landscape and soil profile characteristics of Harve (HRV) Series

4.1.3 Thammadahalli (TDH) Series: Thammadahalli soils are moderately shallow (50-75cm), well drained, have brown to very dark brown and dark reddish brown sandy clay to clay soils. They have developed from weathered granite gneiss and occur on nearly level to gently sloping uplands. The Thammadahalli series has been classified as a member of the fine, mixed, isohyperthermic family of Rhodic Paleustalfs.

The thickness of the solum ranges from 54 to 75 cm. The thickness of A horizon ranges from 11 to 19 cm. Its colour is in 7.5 YR, 5YR and 2.5 YR hue with value 2.5 to 4 and chroma 2 to 6. The texture varies from sandy clay loam to clay with 10 to 20 per cent gravel. The thickness of B horizon ranges from 43 to 60 cm. Its colour is in 2.5 YR hue with value 3 and chroma 4 to 6. Its texture is sandy clay to clay. The available water capacity is medium (100-150 mm/m). One soil phase was identified and mapped.



Landscape and soil profile characteristics of Thammadahalli (TDH) Series

4.1.4 Lakkur (LKR) Series: Lakkur soils are moderately shallow (50-75cm), well drained, have reddish brown to dark red gravelly sandy clay soils. They have developed from weathered granite gneiss and occur on nearly level to very gently and gently sloping uplands. The Lakkur series has been classified as a member of the clayey-skeletal, mixed, isohyperthermic family of Typic Rhodustalfs.

The thickness of the solum ranges from 51 to 74 cm. The thickness of A horizon ranges from 12 to 18 cm. Its colour is in 5YR and 2.5 YR hue with value 3 to 4 and chroma 4 to 6. The texture varies from loamy sand to sandy clay loam with 15 to 50 per cent gravel. The thickness of B horizon ranges from 39 to 58 cm. Its colour is in 2.5 YR hue with value 3 to 4 and chroma 4 to 6. Texture is sandy clay with 40 to 60 per cent gravel. The available water capacity is low (50-100 mm/m). One soil phase was identified and mapped.



Landscape and soil profile characteristics of Lakkur (LKR) Series

4.1.5 Mukhadahalli (MKH) Series: Mukhadahalli soils are moderately shallow (50-75 cm), well drained, have dark brown to reddish brown gravelly sandy clay soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands. The Mukhadahalli series has been classified as a member of the clayey-skeletal, mixed, isohyperthermic family of Typic Haplustalfs.

The thickness of the solum ranges from 51 to 72 cm. The thickness of A horizon ranges from 12 to 17 cm. Its colour is in 5 YR and 7.5 YR hue with value 3 to 4 and chroma 2 to 4. The texture varies from loamy sand to sandy loam with 20 to 45 per cent gravel. The thickness of B horizon ranges from 40 to 68 cm. Its colour is in 2.5 YR and 5 YR hue with value and chroma 3 to 6. Texture is sandy clay loam to sandy clay with 35 to 50 per cent gravel. The available water capacity is low (50-100 mm/m). One soil phase was identified and mapped.



Landscape and soil profile characteristics of Mukhadahalli (MKH) Series

4.1.6 Bidanagere (BDG) Series: Bidanagere soils are moderately deep (75-100 cm), well drained, have dark reddish brown gravelly clay soils. They have developed from weathered granite gneiss and occur on very gently sloping uplands under cultivation. The Bidanagere series has been classified as a member of the clayey-skeletal, mixed, isohyperthermic family of Rhodic Paleustalfs.

The thickness of the solum ranges from 78 to 99 cm. The thickness of A-horizon ranges from 12 to 19 cm. Its colour is in 2.5 YR and 5 YR hue with value 2 to 3 and chroma 3 to 4. The texture varies from sandy clay loam to sandy clay with 10 to 20 per cent gravel. The thickness of B-horizon ranges from 68 to 85 cm. Its colour is in 5 YR and 2.5 YR hue with value 3 to 5 and chroma 3 to 4. Its texture is gravelly clay with gravel content of 35-60 per cent. The available water capacity is very low (<50 mm/m). One soil phase was identified and mapped.



Landscape Soil Profile Characteristics of Bidanagere (BDG) Series

4.1.7 Hooradhahalli (HDH) Series: Hooradhahalli soils are moderately deep (75-100 cm), well drained, have red to dark red and reddish brown, gravelly sandy clay to clay soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands. The Hooradhahalli series has been classified as a member of the clayey-skeletal, mixed, isohyperthermic family of Rhodic Paleustalfs.

The thickness of the solum ranges from 76 to 100 cm. The thickness of A horizon ranges from 11 to 19 cm. Its colour is in 5 YR and 2.5 YR hue with value 3 to 4 and chroma 3 to 6. The texture varies from loamy sand to sandy clay with 15 to 50 per cent gravel. The thickness of B horizon varies from 65 to 83 cm. Its colour is in 2.5 YR hue with value 2.5 to 3 and chroma 4 to 6. Texture is sandy clay to clay with 35 to 50 per cent gravel. The available water capacity is low (50-100mm/m). Four soil phases were identified and mapped.



Landscape and soil profile characteristics of Hooradhahalli (HDH) Series

4.1.8 Bisarahalli (BSR) Series: Bisarahalli soils are moderately deep (75-100 cm), well drained, have dark reddish brown, gravelly sandy clay red soils. They have developed from weathered granite gneiss and occur on very gently sloping uplands under cultivation. The Bisarahalli series has been classified as a member of the fine, mixed isohyperthermic family of Typic Paleustalfs.

The thickness of the solum ranges from 75 to 98 cm. The thickness of A horizon ranges from 17 to 25 cm. Its colour is in 5 YR hue with value 3 to 4 and chroma 3 to 6. The texture ranges from sandy clay loam to sandy clay with 15 to 35 per cent gravel. The thickness of B horizon ranges from 61 to 79 cm. Its colour is in 5 YR hue with value 3 and chroma 3 to 4. Its texture is gravelly sandy clay with gravel content of 15-35 per cent. The available water capacity is low (50-100 mm/m). One soil phase was identified and mapped.



Landscape and soil profile characteristics of Bisarahalli (BSR) Series

4.1.9 Balapur (BPR) Series: Balapur soils are deep (100-150 cm), well drained, have dark reddish brown to dark red, gravelly sandy clay to clay soils. These soils are developed from weathered granite gneiss and occur on very gently to gently sloping uplands. The Balapur series has been classified as a member of the clayey-skeletal, mixed, isohyperthermic family of Typic Rhodustalfs.

The thickness of the solum ranges from 102 to 147 cm. The thickness of A horizon ranges from 12 to 17cm. Its colour is in 5 YR and 2.5 YR hue with value and chroma 3 to 4. The texture ranges from loamy sand to sandy clay with 15 to 50 per cent gravel. The thickness of B horizon ranges from 90 to 132 cm. Its colour is in 2.5 YR hue with value 2.5 to 3 and chroma 4 to 6. Texture is sandy clay to clay with 35 to 50 per cent gravel. The available water capacity is medium (100-150 mm/m). Five soil phases were identified and mapped.



Landscape and soil profile characteristics of Balapur (BPR) Series

4.1.10 Nagalapur (NGP) Series: Nagalapur soils are deep (100-150 cm), well drained, have dark reddish brown to dark red gravelly sandy clay soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands. The Nagalapur series has been classified as a member of the clayey-skeletal, mixed, isohyperthermic family of Typic Paleustalfs.

The thickness of the solum ranges from 105 to 145 cm. The thickness of A horizon ranges from 14 to 20 cm. Its colour is in 7.5 YR hue with value and chroma 3 to 4. The texture ranges from sandy loam to sandy clay with 10 to 50 per cent gravel. The thickness of B horizon ranges from 90 to 128 cm. Its colour is in 2.5 YR, 5 YR and 7.5 YR hue with value 3 to 5 and chroma 3 to 6. Texture is sandy clay to clay with 35 to 80 per cent gravel. The available water capacity is low (51-100 mm/m). Two soil phases were identified and mapped.



Landscape Soil Profile Characteristics of Nagalapur (NGP) Series

4.1.11 Giddadapalya (GDP) Series: Giddadapalya soils are deep (100-150 cm), well drained, have dark reddish brown to dark red, gravelly sandy clay to clay soils. They are developed from weathered granite gneiss and occur on very gently sloping uplands under cultivation. The Giddadapalya soil series has been classified as a member of the fine, mixed, isohyperthermic family of Rhodic Paleustalfs.

The thickness of the solum ranges from 106 to 145 cm. The thickness of A-horizon ranges from 12 to 13 cm. Its colour is in 5 YR hue with value and chroma 3 to 4. The texture ranges from sandy loam with 10 to 15 per cent gravel. The thickness of B-horizon ranges from 106 to 123 cm. Its colour is in 2.5 YR hue with value 3 to 4 and chroma 3 to 6. Texture is sandy clay to clay with 35 to 75 per cent gravel. The available water capacity is low (51-100 mm/m). Two soil phases were identified and mapped.



Landscape and soil profile characteristics of Giddadapalya (GDP) Series

4.2 Soils of Alluvial Landscape

In this landscape, seven soil series were identified and mapped. Of these, series Dambarahalli (DRL) series occupies maximum area of 163 ha (20%) and followed by Ravanaki (RNK) 134 ha (17%) and others occupy minor area. The brief description of the soil series along with the soil phases identified and mapped is given below.

4.2.1 Muttal (MTL) Series: Muttal soils are shallow (25-50 cm), well drained, have dark brown to very dark grayish brown, calcareous gravelly clay soils. They have developed from alluvium and occur on nearly level to very gently sloping plains. The Muttal series has been classified as a member of the clayey, mixed (calc), isohyperthermic family of (Paralithic) Haplustepts.

The thickness of the solum ranges from 30 to 50 cm. The thickness of A horizon ranges from 15 to 18 cm. Its colour is in 7.5 YR and 10 YR hue with value 2 to 3 and chroma 2.5 to 4. The texture varies from sandy clay to clay with 10 to 15 per cent gravel. The thickness of B horizon ranges from 18 to 32 cm. Its colour is in 10 YR and 7.5 YR hue with value 2 to 6 and chroma 2 to 4. Its texture is sandy clay to clay. The available water capacity is low (51-100 mm/m). One soil phase was identified and mapped.



Landscape and soil profile characteristics of Muttal (MTL) Series

4.2.2 Ravanaki (RNK) Series: Ravanaki soils are moderately shallow (50-75 cm), well drained, have dark brown to very dark grayish brown, calcareous clay soils. They have developed from alluvium and occur on nearly level to very gently sloping plains. The Ravanaki series has been classified as a member of the very fine, smectitic (calc), isohyperthermic family of Typic Haplustepts.

The thickness of the solum ranges from 50 to 75 cm. The thickness of A horizon ranges from 15 to 20 cm. Its colour is in 7.5 YR and 10 YR hue with value 2 to 3 and

chroma 2.5 to 4. The texture varies from sandy clay to clay with 10 to 15 per cent gravel. The thickness of B horizon ranges from 35 to 60 cm. Its colour is in 10 YR and 7.5 YR hue with value 2 to 6 and chroma 2 to 4. Its texture is sandy clay to clay with gravel content of 10 to 20 per cent. The available water capacity is low (51-100 mm/m). Two soil phases were identified and mapped.



Landscape and Soil Profile Characteristics of Ravanaki (RNK) Series

4.2.3 Dambarahalli (DRL) Series: Dambarahalli soils are moderately deep (75-100 cm), moderately well drained, have black and very dark gray to dark brown, calcareous cracking clay soils. They have developed from alluvium and occur on very gently to gently sloping plains under cultivation. The Dambarahalli series has been classified as a member of the very fine, smectitic (calc), isohyperthermic family of Typic Haplusterts.

The thickness of the solum ranges from 75 to 99 cm. The thickness of A horizon ranges from 13 to 24 cm. Its colour is in 10 YR hue with value 3 to 4 and chroma 1 to 2. The texture is clay. The thickness of B horizon ranges from 54 to 85 cm. Its colour is in 10 YR hue with value 2 to 4 and chroma 1 to 3. Its texture is clay and are calcareous. The available water capacity is high (151-200 mm/m). Two soil phases were identified and mapped.



Landscape and soil profile characteristics of Dambarahalli (DRL) Series

4.2.4 Narsapura (NSP) Series: Narsapura soils are moderately deep (75-100 cm), moderately well drained, have dark grayish brown to very dark grayish brown and very dark gray, black calcareous, sodic cracking clay soils. They have developed from alluvium and occur on very gently sloping plains. The Narsapura series has been classified as a member of the very fine, smectitic (calc), isohyperthermic family of Vertic Haplustepts.

The thickness of the solum is 76 to 98 cm. The thickness of A horizon ranges from 15 to 19 cm. Its colour is in 10 YR hue with value 3 and chroma 1 to 2. The texture is clay with no gravel. The thickness of B horizon ranges from 57 to 83 cm. Its colour is in 10 YR hue with value 3 to 5 and chroma 1 to 3. Its texture is clay and are calcareous. The available water capacity is medium (101-150 mm/m). One soil phase was identified and mapped.



Landscape and soil profile characteristics of Narsapura (NSP) series

4.2.5 Kavalur (KVR) Series: Kavalur soils are deep (100-150 cm), moderately well drained, have dark yellowish brown to very dark brown and very dark gray, calcareous black cracking clay soils. They have developed from alluvium and occur on very gently sloping plains. The Kavalur series has been classified as a member of the fine, smectitic (calc), isohyperthermic family of Typic Haplusterts.

The thickness of the solum is 113 to 143 cm. The thickness of A horizon ranges from 9 to 24 cm. Its colour is in 10 YR hue with value 3 and chroma 1. The texture is clay with no gravel. The thickness of B horizon ranges from 89 to 134 cm. Its colour is in 10 YR hue with value 3 and chroma 1. Its texture is clay. The available water capacity is very high (>200 mm/m). Two soil phases were identified and mapped.



Landscape and soil profile characteristics of Kavalur (KVR) series

4.2.6 Lakshangudda (LGD) Series: Lakshangudda soils are deep (100-150 cm), moderately well drained, have light olive brown to very dark gray calcareous clayey soils. They have developed from alluvium and occur on nearly level plains. The Lakshangudda series has been classified as a member of the fine, smectitic (calc), isohyperthermic family of Typic Haplusterts.


The thickness of the solum ranges from 108 to 149 cm. The thickness of A horizon ranges from 16 to 20 cm. Its colour is in 7.5 YR and 10 YR hue with value and chroma 3 to 4. The texture varies from sandy clay to clay with 5 to 10 per cent gravel. The thickness of B horizon ranges from 90 to 132 cm. Its colour is in 2.5 Y, 10 YR and 7.5 YR hue with value 3 to 6 and chroma 1 to 3. Its texture is clay. The available water capacity is high (150-200 mm/m). One soil phase was identified and mapped.



Landscape and Soil Profile Characteristics of Lakshmangudda (LGD) Series

4.2.7 Alawandi (AWD) Series: Alawandi soils are very deep (>150 cm), moderately well drained, have black to very dark grayish brown, calcareous cracking clay soils. They have developed from alluvium and occur on nearly level to very gently sloping plains under cultivation. The Alawandi series has been classified as a member of the fine, smectitic(calc), isohyperthermic family of Typic Haplusterts .

The thickness of the solum is more than 150 cm. The thickness of A horizon ranges from 16 to 26 cm. Its colour is in 10 YR hue with value 2 to 3 and chroma 1 to 2. The texture varies from sandy clay to clay. The thickness of B horizon is more than 150 cm. Its colour is in 10 YR hue with value 2 to 3 and chroma 1 to 3. Its texture is clay and is calcareous. The available water capacity is very high (>200 mm/m). One soil phase was identified and mapped.



Landscape and Soil Profile Characteristics of Alawandi (AWD) Series

Table: 4.1 Physical and Chemical Characteristics of Soil Series identified in Haligeri-3 microwatershed

Series Name: Harve (HRV), **Pedon:**R-10

Location: 15°25'11.63"N, 76°22'03.65"E Jabbaragudda village, Koppal taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bangalore. **Classification:** Loamy-skeletal, mixed, isohyperthermic (Paralithic) Rhodustalfs

| Depth (cm) | Horizon | Size class and particle diameter (mm) | | | | | | | | Coarse fragments w/w (%) | Texture Class (USDA) | % Moisture | |
|------------|---------|---------------------------------------|-------------------|---------------|-----------------------|------------------|-------------------|-----------------|----------------------|--------------------------|----------------------|------------|--------|
| | | Total | | | Sand | | | | | | | 1/3 Bar | 15 Bar |
| | | Sand (2.0-0.05) | Silt (0.05-0.002) | Clay (<0.002) | Very coarse (2.0-1.0) | Coarse (1.0-0.5) | Medium (0.5-0.25) | Fine (0.25-0.1) | Very fine (0.1-0.05) | | | | |
| 0-15 | Ap | 65.64 | 9.07 | 25.28 | 29.04 | 12.99 | 9.00 | 3.48 | 11.15 | 50 | scl | 12.87 | 4.81 |
| 15-29 | Bt1 | 56.13 | 7.75 | 36.12 | 27.81 | 11.43 | 7.21 | 1.44 | 8.24 | 60 | sc | 15.69 | 6.24 |
| 29-47 | Bt2 | 63.42 | 6.53 | 30.05 | 32.38 | 13.93 | 7.48 | 5.74 | 3.89 | 60 | scl | 15.41 | 9.29 |

| Depth (cm) | pH (1:2.5) | | | E.C. (1:2.5) | O.C. | CaCO ₃ | Exchangeable bases | | | | | CEC | CEC/Clay | Base saturation | ESP |
|------------|------------|-------------------|-------|--------------------|------|-------------------|-----------------------|------|------|------|-------|-------|----------|-----------------|------|
| | Water | CaCl ₂ | M KCl | | | | Ca | Mg | K | Na | Total | | | | |
| | | | | dS m ⁻¹ | % | % | cmol kg ⁻¹ | | | | | | % | % | |
| 0-15 | 6.05 | - | - | 0.21 | 0.93 | - | 8.89 | 1.96 | 0.50 | 0.08 | 11.43 | 11.24 | 0.44 | 100.00 | 0.73 |
| 15-29 | 5.99 | - | - | 0.15 | 0.29 | - | 9.72 | 2.75 | 0.51 | 0.09 | 13.07 | 12.71 | 0.35 | 100.00 | 0.74 |
| 29-47 | 6.07 | - | - | 0.11 | 0.38 | - | 9.35 | 2.47 | 0.49 | 0.06 | 12.36 | 12.71 | 0.42 | 97.29 | 0.44 |

Contd...

Soil Series: Thammadahalli (TDH), **Pedon:** TR₁/1

Location: 15°03'41.7"N, 75°36'65.2"E, (4D4A3G2d), Nilogal village, Shirahatti taluk, Gadag district

Analysis at: NBSS&LUP, Regional Centre, Bengaluru **Classification:** Fine, mixed, isohyperthermic Rhodic Paleustalfs

| Depth (cm) | Horizon | Size class and particle diameter (mm) | | | | | | | | Coarse fragments w/w (%) | Texture Class (USDA) | % Moisture | |
|------------|---------|---------------------------------------|-------------------|---------------|-----------------------|------------------|-------------------|-----------------|----------------------|--------------------------|----------------------|------------|--------|
| | | Total | | | Sand | | | | | | | 1/3 Bar | 15 Bar |
| | | Sand (2.0-0.05) | Silt (0.05-0.002) | Clay (<0.002) | Very coarse (2.0-1.0) | Coarse (1.0-0.5) | Medium (0.5-0.25) | Fine (0.25-0.1) | Very fine (0.1-0.05) | | | | |
| 0-25 | Ap | 85.71 | 7.34 | 6.94 | 14.79 | 13.28 | 16.10 | 24.75 | 16.80 | 20 | ls | - | - |
| 25-65 | Bt | 47.76 | 7.96 | 44.28 | 15.30 | 9.78 | 6.24 | 7.91 | 8.53 | 10 | sc | - | - |

| Depth (cm) | pH (1:2.5) | | | E.C. (1:2.5) dS m ⁻¹ | O.C. % | CaCO ₃ % | Exchangeable bases | | | | | CEC | CEC/Clay | Base saturation % | ESP % |
|------------|------------|-------------------|-------|------------------------------------|-----------|------------------------|--------------------|----|------|------|-------|-------|----------|----------------------|----------|
| | Water | CaCl ₂ | M KCl | | | | Ca | Mg | K | Na | Total | | | | |
| | | | | | | | | | | | | | | | |
| 0-25 | 9.19 | - | - | 0.18 | 0.35 | 1.29 | - | - | 0.08 | 0.52 | 0.60 | 3.57 | 0.51 | 100.00 | 5.82 |
| 25-65 | 8.00 | - | - | 0.17 | 0.35 | 0.58 | - | - | 0.15 | 1.31 | 1.46 | 13.87 | 0.31 | 100.00 | 3.78 |

Contd...

Soil Series: Lakkur (LKR), **Pedon:** RM-8.

Location: 15°04'26.3"N, 75°37'84.1"E, (4D4A3I1f), Belhatti village, Shirahatti taluk, Gadag district

Analysis at: NBSS&LUP, Regional Centre, Bengaluru **Classification:** Clayey-skeletal, mixed, isohyperthermic Typic Rhodustalfs

| Depth (cm) | Horizon | Size class and particle diameter (mm) | | | | | | | | Coarse fragments w/w (%) | Texture Class (USDA) | % Moisture | |
|------------|---------|---------------------------------------|-------------------|---------------|-----------------------|------------------|-------------------|-----------------|----------------------|--------------------------|----------------------|------------|--------|
| | | Total | | | Sand | | | | | | | 1/3 Bar | 15 Bar |
| | | Sand (2.0-0.05) | Silt (0.05-0.002) | Clay (<0.002) | Very coarse (2.0-1.0) | Coarse (1.0-0.5) | Medium (0.5-0.25) | Fine (0.25-0.1) | Very fine (0.1-0.05) | | | | |
| 0-21 | Ap | 74.00 | 8.34 | 17.66 | 9.62 | 11.57 | 15.76 | 23.13 | 13.92 | 20 | sl | - | - |
| 21-35 | Bt | 54.37 | 10.48 | 35.14 | 16.33 | 8.64 | 9.69 | 11.59 | 8.11 | 40 | sc | - | - |
| 35-56 | Bc | 48.37 | 13.46 | 38.17 | 10.96 | 7.69 | 9.17 | 11.28 | 9.27 | 60 | sc | - | - |

| Depth (cm) | pH (1:2.5) | | | E.C. (1:2.5) dS m ⁻¹ | O.C. % | CaCO ₃ % | Exchangeable bases | | | | | CEC | CEC/Clay | Base saturation % | ESP % |
|------------|------------|-------------------|-------|---------------------------------|--------|---------------------|--------------------|----|------|------|-------|-------|----------|-------------------|-------|
| | Water | CaCl ₂ | M KCl | | | | Ca | Mg | K | Na | Total | | | | |
| | | | | | | | | | | | | | | | |
| 0-21 | 8.18 | - | - | 0.30 | 0.56 | 0.94 | - | - | 0.31 | 0.55 | 0.86 | 12.19 | 0.69 | 100.00 | 4.51 |
| 21-35 | 8.17 | - | - | 0.30 | 0.52 | 1.29 | - | - | 0.19 | 0.84 | 1.03 | 22.18 | 0.63 | 100.00 | 3.79 |
| 35-56 | 7.95 | - | - | 0.46 | 0.48 | 1.99 | - | - | 0.24 | 0.58 | 0.82 | 22.94 | 0.60 | 100.00 | 2.53 |

Contd...

Series Name: Mukahadahalli (MKH), **Pedon:** R-11

Location: 15°22'05.4"N, 76°04'10.3"E, Halageri village, Koppal Taluk and District

Analysis at: NBSS&LUP, Regional Centre, Bangalore. **Classification:** Clayey-skeletal, mixed, isohyperthermic Typic Haplustalfs

| Depth (cm) | Horizon | Size class and particle diameter (mm) | | | | | | | | Coarse fragments w/w (%) | Texture Class (USDA) | % Moisture | |
|------------|---------|---------------------------------------|-------------------|---------------|-----------------------|------------------|-------------------|-----------------|----------------------|--------------------------|----------------------|------------|--------|
| | | Total | | | Sand | | | | | | | 1/3 Bar | 15 Bar |
| | | Sand (2.0-0.05) | Silt (0.05-0.002) | Clay (<0.002) | Very coarse (2.0-1.0) | Coarse (1.0-0.5) | Medium (0.5-0.25) | Fine (0.25-0.1) | Very fine (0.1-0.05) | | | | |
| 0-19 | Ap | 65.71 | 8.83 | 25.46 | 9.27 | 9.06 | 14.42 | 21.52 | 11.43 | 70 | scl | 16.54 | 8.60 |
| 19-32 | Bt | 55.89 | 11.13 | 32.98 | 6.47 | 9.18 | 11.89 | 19.19 | 9.18 | 50 | scl | 19.24 | 12.78 |
| 32-58 | Bt | 47.95 | 10.41 | 41.63 | 17.52 | 3.78 | 9.13 | 9.55 | 7.97 | 50 | sc | 24.03 | 16.02 |

| Depth (cm) | pH (1:2.5) | | | E.C. (1:2.5) | O.C. | CaCO ₃ | Exchangeable bases | | | | | CEC | CEC/Clay | Base saturation | ESP | | | |
|------------|------------|-------------------|-------|--------------|------|-------------------|--------------------|------|------|-----------------------|-------|-------|----------|-----------------|------|---|----|-------|
| | Water | CaCl ₂ | M KCl | | | | dS m ⁻¹ | % | % | Ca | Mg | | | | | K | Na | Total |
| | | | | | | | | | | cmol kg ⁻¹ | | | | | | | | |
| 0-19 | 7.38 | - | - | 0.09 | 0.2 | 0.00 | 8.97 | 4.32 | 0.26 | 0.22 | 13.77 | 14.84 | 0.58 | 93 | 1.49 | | | |
| 19-32 | 7.5 | - | - | 0.106 | 0.41 | 0.00 | 15.98 | 3.27 | 0.16 | 0.50 | 19.91 | 20.88 | 0.63 | 95 | 2.38 | | | |
| 32-58 | 7.46 | - | - | 0.173 | 0.49 | 0.00 | 19.71 | 4.53 | 0.23 | 1.32 | 25.79 | 25.76 | 0.62 | 100 | 5.11 | | | |

Contd...

Series: Bidanagere (BDG), **Pedon:** RM-3

Location: 13°22'11"N, 76°38'03"E, (4D3D8G1a), Tharabenahalli village, Chikkanayakanahalli Taluk, Tumakuru District.

Analysis at: NBSS&LUP, Regional Centre, Bengaluru **Classification:** Clayey-skeletal, mixed, isohyperthermic Rhodic Paleustalfs

| Depth (cm) | Horizon | Size class and particle diameter (mm) | | | | | | | | Coarse fragments w/w (%) | Texture Class (USDA) | % Moisture | |
|------------|---------|---------------------------------------|-------------------|---------------|-----------------------|------------------|-------------------|-----------------|----------------------|--------------------------|----------------------|------------|--------|
| | | Total | | | Sand | | | | | | | 1/3 Bar | 15 Bar |
| | | Sand (2.0-0.05) | Silt (0.05-0.002) | Clay (<0.002) | Very coarse (2.0-1.0) | Coarse (1.0-0.5) | Medium (0.5-0.25) | Fine (0.25-0.1) | Very fine (0.1-0.05) | | | | |
| 0-20 | Ap | 81.19 | 11.25 | 7.56 | 12.54 | 15.07 | 17.90 | 21.94 | 13.75 | 50 | ls | - | - |
| 20-35 | Bt1 | 57.45 | 11.45 | 31.10 | 12.76 | 11.02 | 10.92 | 12.45 | 10.31 | 50 | scl | - | - |
| 35-92 | Bt2 | 44.63 | 7.85 | 47.52 | 12.40 | 9.61 | 8.37 | 7.75 | 6.51 | 60 | c | - | - |

| Depth (cm) | pH (1:2.5) | | | E.C. (1:2.5) dS m ⁻¹ | O.C. % | CaCO ₃ % | Exchangeable bases | | | | | CEC | CEC/Clay | Base saturation % | ESP % |
|------------|------------|-------------------|-------|---------------------------------|--------|---------------------|--------------------|------|------|------|-------|------|----------|-------------------|-------|
| | Water | CaCl ₂ | M KCl | | | | Ca | Mg | K | Na | Total | | | | |
| | | | | | | | | | | | | | | | |
| 0-20 | 6.24 | - | - | 0.06 | 0.60 | 0.00 | 1.61 | 0.26 | 0.10 | 0.01 | 1.98 | 3.76 | 0.50 | 52.56 | 0.35 |
| 20-35 | 5.99 | - | - | 0.02 | 0.40 | 0.00 | 4.25 | 0.46 | 0.08 | 0.28 | 5.07 | 8.02 | 0.26 | 63.18 | 3.46 |
| 35-92 | 6.70 | - | - | 0.03 | 0.20 | 0.00 | 5.45 | 0.31 | 0.10 | 0.22 | 6.09 | 9.90 | 0.21 | 61.48 | 2.24 |

Contd...

Soil Series: Hooradhahalli (HDH), **Pedon:** RM-69

Location: 13°24'31"N, 76°33'41"E, (4D3D8G2d), Hesarahalli village, Chikkanayakanahalli taluk, Tumukura district

Analysis at: NBSS&LUP, Regional Centre, Bengaluru **Classification:** Clayey-skeletal, mixed, isohyperthermic Rhodic Paleustalfs

| Depth (cm) | Horizon | Size class and particle diameter (mm) | | | | | | | | Coarse fragments w/w (%) | Texture Class (USDA) | % Moisture | |
|------------|---------|---------------------------------------|-------------------|---------------|-----------------------|------------------|-------------------|-----------------|----------------------|--------------------------|----------------------|------------|--------|
| | | Total | | | Sand | | | | | | | 1/3 Bar | 15 Bar |
| | | Sand (2.0-0.05) | Silt (0.05-0.002) | Clay (<0.002) | Very coarse (2.0-1.0) | Coarse (1.0-0.5) | Medium (0.5-0.25) | Fine (0.25-0.1) | Very fine (0.1-0.05) | | | | |
| 0-18 | Ap | 72.56 | 15.17 | 12.27 | 4.57 | 8.33 | 17.38 | 23.88 | 18.39 | 35 | sl | - | - |
| 18-33 | Bt1 | 56.29 | 10.75 | 32.96 | 7.88 | 10.24 | 13.41 | 14.43 | 10.34 | 55 | scl | - | - |
| 33-58 | Bt2 | 46.66 | 10.79 | 42.55 | 10.79 | 9.87 | 8.43 | 9.04 | 8.53 | 55 | sc | - | - |
| 58-90 | Bt3 | 43.09 | 13.63 | 43.27 | 9.90 | 8.25 | 7.32 | 8.76 | 8.87 | 45 | c | - | - |

| Depth (cm) | pH (1:2.5) | | | E.C. (1:2.5) dS m ⁻¹ | O.C. % | CaCO ₃ % | Exchangeable bases | | | | | CEC | CEC/Clay | Base saturation % | ESP % |
|------------|------------|-------------------|-------|---------------------------------|--------|---------------------|--------------------|------|------|------|-------|-------|----------|-------------------|-------|
| | Water | CaCl ₂ | M KCl | | | | Ca | Mg | K | Na | Total | | | | |
| | | | | | | | | | | | | | | | |
| 0-18 | 6.54 | - | - | 0.07 | 0.60 | 0.00 | 2.68 | 1.38 | 0.44 | 0.42 | 4.91 | 5.84 | 0.48 | 84.07 | 7.11 |
| 18-33 | 5.90 | - | - | 0.07 | 0.52 | 0.00 | 3.99 | 1.27 | 0.09 | 0.37 | 5.71 | 8.61 | 0.26 | 66.32 | 4.29 |
| 33-58 | 6.16 | - | - | 0.07 | 0.44 | 0.00 | 4.92 | 1.67 | 0.08 | 0.55 | 7.22 | 10.00 | 0.24 | 72.23 | 5.50 |
| 58-90 | 6.39 | - | - | 0.06 | 0.40 | 0.00 | 4.30 | 2.02 | 0.08 | 0.46 | 6.87 | 9.21 | 0.21 | 74.61 | 5.05 |

Contd...

Series Name: Bisarahalli (BSR) **Pedon:** R-9

Location: 15°25'21.0"N, 76°11'42.0"E Hatti village, Koppal taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bangalore. **Classification:** Fine, mixed, isohyperthermic Typic Paleustalfs

| Depth (cm) | Horizon | Size class and particle diameter (mm) | | | | | | | | Coarse fragments w/w (%) | Texture Class (USDA) | % Moisture | |
|------------|---------|---------------------------------------|-------------------|---------------|-----------------------|------------------|-------------------|-----------------|----------------------|--------------------------|----------------------|------------|--------|
| | | Total | | | Sand | | | | | | | 1/3 Bar | 15 Bar |
| | | Sand (2.0-0.05) | Silt (0.05-0.002) | Clay (<0.002) | Very coarse (2.0-1.0) | Coarse (1.0-0.5) | Medium (0.5-0.25) | Fine (0.25-0.1) | Very fine (0.1-0.05) | | | | |
| 0-14 | Ap | 70.11 | 9.29 | 20.60 | 22.31 | 15.97 | 11.98 | 9.83 | 10.03 | 20 | scl | 13.22 | 7.81 |
| 14-57 | Bt1 | 47.27 | 7.52 | 45.20 | 27.04 | 8.28 | 4.61 | 2.10 | 5.24 | 25 | sc | 16.39 | 13.31 |
| 57-80 | Bt2 | 41.93 | 8.67 | 49.40 | 21.95 | 6.83 | 4.76 | 4.66 | 3.73 | 30 | c | 21.41 | 15.41 |
| 80-99 | Bt3 | 49.02 | 9.87 | 41.11 | 19.90 | 10.78 | 6.84 | 6.42 | 5.08 | 40 | sc | 21.82 | 14.24 |

| Depth (cm) | pH (1:2.5) | | | E.C. (1:2.5) dS m ⁻¹ | O.C. % | CaCO ₃ % | Exchangeable bases | | | | | CEC | CEC/Clay | Base saturation % | ESP % |
|------------|------------|-------------------|-------|---------------------------------|--------|---------------------|-----------------------|------|------|------|-------|-------|----------|-------------------|-------|
| | Water | CaCl ₂ | M KCl | | | | Ca | Mg | K | Na | Total | | | | |
| | | | | | | | cmol kg ⁻¹ | | | | | | | | |
| 0-14 | 6.59 | - | - | 0.12 | 0.73 | - | 4.47 | 1.77 | 0.06 | 0.53 | 6.82 | 8.80 | 0.43 | 77.55 | 6.00 |
| 14-57 | 7.02 | - | - | 0.04 | 0.48 | - | 5.85 | 2.31 | 0.06 | 0.20 | 8.43 | 14.70 | 0.33 | 57.32 | 1.36 |
| 57-80 | 7.00 | - | - | 0.05 | 0.28 | - | 11.74 | 2.26 | 0.08 | 0.22 | 14.31 | 15.60 | 0.32 | 91.73 | 1.44 |
| 80-99 | 6.90 | - | - | 0.06 | 0.18 | - | 13.70 | 2.16 | 0.08 | 0.14 | 16.08 | 16.50 | 0.40 | 97.44 | 0.83 |

Contd...

Soil Series: Balapur (BPR), **Pedon:** RM-78

Location: 13°26'39"N, 76°35'03"E, (4D3D8G2c), Kasaba, Chikkanayakanahalli taluk, Tumakuru district

Analysis at: NBSS&LUP, Regional Centre, Bengaluru **Classification:** Clayey-skeletal, mixed, isohyperthermic Typic Rhodustalfs

| Depth (cm) | Horizon | Size class and particle diameter (mm) | | | | | | | | Coarse fragments w/w (%) | Texture Class (USDA) | % Moisture | |
|------------|---------|---------------------------------------|-------------------|---------------|-----------------------|------------------|-------------------|-----------------|----------------------|--------------------------|----------------------|------------|--------|
| | | Total | | | Sand | | | | | | | 1/3 Bar | 15 Bar |
| | | Sand (2.0-0.05) | Silt (0.05-0.002) | Clay (<0.002) | Very coarse (2.0-1.0) | Coarse (1.0-0.5) | Medium (0.5-0.25) | Fine (0.25-0.1) | Very fine (0.1-0.05) | | | | |
| 0-12 | Ap | 65.66 | 18.66 | 15.68 | 4.14 | 6.16 | 13.33 | 21.82 | 20.20 | - | sl | - | - |
| 12-34 | Bt1 | 61.91 | 11.52 | 26.57 | 2.36 | 6.78 | 12.53 | 21.36 | 18.89 | - | scl | - | - |
| 34-60 | Bt2 | 51.81 | 11.24 | 36.94 | 4.66 | 5.70 | 12.23 | 15.96 | 13.26 | 30 | sc | - | - |
| 60-84 | Bt3 | 46.61 | 9.02 | 44.37 | 14.70 | 6.88 | 7.51 | 8.97 | 8.55 | 55 | sc | - | - |
| 84-112 | Bt4 | 48.75 | 12.92 | 38.33 | 15.73 | 8.13 | 6.87 | 8.23 | 9.79 | 60 | sc | - | - |
| 112-127 | Bc | 50.98 | 24.74 | 24.28 | 5.25 | 4.63 | 5.15 | 10.92 | 25.03 | 50 | scl | - | - |

| Depth (cm) | pH (1:2.5) | | | E.C. (1:2.5) | O.C. | CaCO ₃ | Exchangeable bases | | | | | CEC | CEC/Clay | Base saturation | ESP |
|------------|------------|-------------------|-------|--------------------|------|-------------------|-----------------------|------|------|------|-------|-------|----------|-----------------|------|
| | Water | CaCl ₂ | M KCl | | | | Ca | Mg | K | Na | Total | | | | |
| | | | | dS m ⁻¹ | % | % | cmol kg ⁻¹ | | | | | | % | % | |
| 0-12 | 6.64 | - | - | 0.03 | 0.56 | 0.00 | 1.90 | 1.32 | 0.21 | 0.03 | 3.46 | 5.45 | 0.35 | 63.48 | 0.51 |
| 12-34 | 6.99 | - | - | 0.02 | 0.48 | 0.00 | 3.66 | 1.90 | 0.07 | 0.08 | 5.70 | 7.82 | 0.29 | 72.93 | 0.96 |
| 34-60 | 7.29 | - | - | 0.02 | 0.40 | 0.00 | 5.13 | 2.08 | 0.11 | 0.20 | 7.52 | 11.19 | 0.30 | 67.18 | 1.75 |
| 60-84 | 7.50 | - | - | 0.02 | 0.32 | 0.00 | 5.83 | 6.36 | 0.13 | 0.23 | 12.55 | 12.38 | 0.28 | 101.43 | 1.83 |
| 84-112 | 7.54 | - | - | 0.02 | 0.24 | 0.00 | 6.02 | 6.59 | 0.11 | 0.25 | 12.96 | 12.77 | 0.33 | 101.49 | 1.97 |
| 112-127 | 7.90 | - | - | 0.02 | 0.20 | 0.00 | 8.04 | 3.62 | 0.07 | 0.32 | 12.04 | 12.47 | 0.51 | 96.56 | 2.55 |

Contd...

Series Name: Nagalapur (NGP) **Pedon :** R-10

Location: 15°26'38.0"N, 76°10'27.0" E Budashettynala village, Koppal taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bangalore. **Classification:** Clayey- skeletal, mixed, isohyperthermic Typic Paleustalfs

| Depth (cm) | Horizon | Size class and particle diameter (mm) | | | | | | | | Coarse fragments w/w (%) | Texture Class (USDA) | % Moisture | |
|------------|---------|---------------------------------------|-------------------|---------------|-----------------------|------------------|-------------------|-----------------|----------------------|--------------------------|----------------------|------------|--------|
| | | Total | | | Sand | | | | | | | 1/3 Bar | 15 Bar |
| | | Sand (2.0-0.05) | Silt (0.05-0.002) | Clay (<0.002) | Very coarse (2.0-1.0) | Coarse (1.0-0.5) | Medium (0.5-0.25) | Fine (0.25-0.1) | Very fine (0.1-0.05) | | | | |
| 0-16 | Ap | 78.43 | 6.36 | 15.21 | 25.23 | 18.82 | 14.04 | 13.22 | 7.12 | 30 | sl | 9.32 | 5.56 |
| 16-38 | Bt1 | 46.97 | 8.53 | 44.51 | 14.33 | 12.34 | 7.43 | 6.80 | 6.07 | 30 | sc | 18.70 | 13.79 |
| 38-58 | Bt2 | 51.92 | 7.48 | 40.60 | 20.98 | 10.07 | 7.37 | 7.48 | 6.02 | 40 | sc | 17.93 | 13.75 |
| 58-81 | Bt3 | 54.05 | 7.18 | 38.77 | 27.07 | 10.58 | 5.91 | 5.81 | 4.67 | 50 | sc | 17.92 | 11.87 |
| 81-104 | Bt4 | 59.03 | 8.93 | 32.04 | 21.88 | 13.11 | 8.88 | 8.05 | 7.12 | 50 | scl | 16.63 | 10.55 |
| 104-126 | BC | 62.35 | 9.26 | 28.40 | 21.19 | 14.51 | 9.88 | 8.13 | 8.64 | 60 | scl | 15.03 | 10.06 |

| Depth (cm) | pH (1:2.5) | | | E.C. (1:2.5) | O.C. | CaCO ₃ | Exchangeable bases | | | | | CEC | CEC/Clay | Base saturation | ESP |
|------------|------------|-------------------|-------|--------------------|------|-------------------|-----------------------|------|------|------|-------|-------|----------|-----------------|------|
| | Water | CaCl ₂ | M KCl | | | | Ca | Mg | K | Na | Total | | | | |
| | | | | dS m ⁻¹ | % | % | cmol kg ⁻¹ | | | | | % | % | | |
| 0-16 | 6.77 | - | - | 0.09 | 0.82 | - | 3.52 | 2.14 | 0.18 | 0.03 | 5.87 | 7.10 | 0.47 | 82.70 | 0.46 |
| 16-38 | 6.89 | - | - | 0.06 | 0.57 | - | 9.35 | 3.85 | 0.10 | 0.21 | 13.50 | 14.70 | 0.33 | 91.87 | 1.40 |
| 38-58 | 6.80 | - | - | 0.06 | 0.52 | - | 8.76 | 3.42 | 0.10 | 0.26 | 12.55 | 14.20 | 0.35 | 88.35 | 1.85 |
| 58-81 | 6.84 | - | - | 0.06 | 0.32 | - | 7.67 | 2.77 | 0.10 | 0.58 | 11.12 | 12.90 | 0.33 | 86.18 | 4.48 |
| 81-104 | 6.86 | - | - | 0.05 | 0.20 | - | 6.97 | 2.07 | 0.09 | 0.95 | 10.07 | 11.90 | 0.37 | 84.59 | 7.95 |
| 104-126 | 6.70 | - | - | 0.07 | 0.10 | - | 5.53 | 1.77 | 0.07 | 0.73 | 8.09 | 9.40 | 0.33 | 86.09 | 7.77 |

Contd...

Series Name: Giddadapalya (GDP), **Pedon:** R-8

Location: 15°25'26"N, 76°10'59"E, Kalakeri village, Koppal taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bangalore. **Classification:** Fine, mixed, isohyperthermic Rhodic Paleustalfs

| Depth (cm) | Horizon | Size class and particle diameter (mm) | | | | | | | | Coarse fragments w/w (%) | Texture Class (USDA) | % Moisture | |
|------------|---------|---------------------------------------|-------------------|---------------|-----------------------|------------------|-------------------|-----------------|----------------------|--------------------------|----------------------|------------|--------|
| | | Total | | | Sand | | | | | | | 1/3 Bar | 15 Bar |
| | | Sand (2.0-0.05) | Silt (0.05-0.002) | Clay (<0.002) | Very coarse (2.0-1.0) | Coarse (1.0-0.5) | Medium (0.5-0.25) | Fine (0.25-0.1) | Very fine (0.1-0.05) | | | | |
| 0-16 | Ap | 74.95 | 9.24 | 15.81 | 18.43 | 18.94 | 13.85 | 14.97 | 8.76 | - | sl | 11.88 | 5.09 |
| 16-43 | Bt1 | 41.69 | 13.89 | 44.42 | 9.84 | 10.90 | 7.41 | 7.62 | 5.93 | - | c | 23.13 | 14.53 |
| 43-61 | Bt2 | 47.67 | 6.13 | 46.19 | 21.14 | 10.15 | 5.29 | 6.45 | 4.65 | - | sc | 21.60 | 11.87 |
| 61-83 | Bt3 | 52.52 | 7.10 | 40.38 | 24.42 | 10.59 | 5.66 | 7.55 | 4.30 | 40 | sc | 19.51 | 11.35 |
| 83-119 | Bt4 | 43.76 | 11.59 | 44.65 | 20.15 | 7.56 | 5.77 | 5.46 | 4.83 | 60 | c | 20.80 | 12.06 |
| 119-139 | Bt5 | 54.93 | 9.84 | 35.23 | 29.70 | 10.49 | 5.50 | 5.92 | 3.32 | 50 | sc | 15.24 | 11.97 |

| Depth (cm) | pH (1:2.5) | | | E.C. (1:2.5) | O.C. | CaCO ₃ | Exchangeable bases | | | | | CEC | CEC/Clay | Base saturation | ESP | | | |
|------------|------------|-------------------|-------|--------------|------|-------------------|--------------------|------|------|-----------------------|-------|------|----------|-----------------|------|---|----|-------|
| | Water | CaCl ₂ | M KCl | | | | dS m ⁻¹ | % | % | Ca | Mg | | | | | K | Na | Total |
| | | | | | | | | | | cmol kg ⁻¹ | | | | | | | | |
| 0-16 | 7.88 | - | - | 0.103 | 0.79 | - | 5.98 | 1.35 | 0.05 | 0.22 | 7.60 | 7.8 | 0.49 | 97 | 2.87 | | | |
| 16-43 | 7.81 | - | - | 0.117 | 0.66 | - | 13.99 | 1.97 | 0.08 | 0.46 | 16.50 | 16.9 | 0.38 | 98 | 2.74 | | | |
| 43-61 | 7.74 | - | - | 0.132 | 0.51 | - | 12.70 | 2.18 | 0.08 | 0.69 | 15.64 | 15.9 | 0.34 | 98 | 4.36 | | | |
| 61-83 | 7.72 | - | - | 0.142 | 0.39 | - | 11.46 | 2.22 | 0.08 | 0.66 | 14.41 | 14.6 | 0.36 | 99 | 4.53 | | | |
| 83-119 | 7.58 | - | - | 0.115 | 0.22 | - | 11.30 | 2.70 | 0.09 | 0.73 | 14.82 | 15.3 | 0.34 | 97 | 4.79 | | | |
| 119-139 | 7.50 | - | - | 0.113 | 0.22 | - | 10.03 | 2.19 | 0.07 | 0.65 | 12.95 | 13.2 | 0.37 | 98 | 4.89 | | | |

Contd...

Series Name: Muttal (MTL), **Pedon:** RM-13

Location: 15°14'30.8"N, 75°56'50.6"E, Gatareddihalla village, Koppal Taluk and District

Analysis at: NBSS&LUP, Regional Centre, Bangalore. **Classification:** Clayey, mixed, (calc) isohyperthermic (Paralithic) Haplustepts

| Depth (cm) | Horizon | Size class and particle diameter (mm) | | | | | | | | Coarse fragments w/w (%) | Texture Class (USDA) | % Moisture | |
|------------|---------|---------------------------------------|-------------------|---------------|-----------------------|------------------|-------------------|-----------------|----------------------|--------------------------|----------------------|------------|--------|
| | | Total | | | Sand | | | | | | | 1/3 Bar | 15 Bar |
| | | Sand (2.0-0.05) | Silt (0.05-0.002) | Clay (<0.002) | Very coarse (2.0-1.0) | Coarse (1.0-0.5) | Medium (0.5-0.25) | Fine (0.25-0.1) | Very fine (0.1-0.05) | | | | |
| 0-20 | Ap | 39.05 | 13.74 | 47.21 | 3.05 | 5.05 | 8.21 | 14.63 | 8.11 | 15-30 | c | 29.95 | 17.94 |
| 20-34 | Bwk | 28.77 | 19.57 | 51.66 | 4.81 | 4.71 | 4.92 | 9.09 | 5.24 | 10 | c | 33.44 | 21.56 |

| Depth (cm) | pH (1:2.5) | | | E.C. (1:2.5) | O.C. | CaCO ₃ | Exchangeable bases | | | | | CEC | CEC/Clay | Base saturation | ESP |
|------------|------------|-------------------|-------|--------------------|------|-------------------|-----------------------|----|------|------|-------|-------|----------|-----------------|------|
| | Water | CaCl ₂ | M KCl | | | | Ca | Mg | K | Na | Total | | | | |
| | | | | dS m ⁻¹ | % | % | cmol kg ⁻¹ | | | | | % | % | | |
| 0-20 | 8.27 | - | - | 0.202 | 0.79 | 6.10 | - | - | 0.62 | 0.25 | - | 36.64 | 0.78 | - | 0.69 |
| 20-34 | 8.36 | - | - | 0.177 | 0.99 | 23.04 | - | - | 0.29 | 0.38 | - | 39.60 | 0.77 | - | 0.96 |

Contd...

Series Name: Ravanaki (RNK), **Pedon:** RM-20

Location: 15°14'22.7"N, 75°57'45.8"E, Gatareddihalla village, Koppal taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bangalore

Classification: Very fine, smectitic(calc), isohyperthermic Typic Haplustepts

| Depth (cm) | Horizon | Size class and particle diameter (mm) | | | | | | | | Coarse fragments w/w (%) | Texture Class (USDA) | % Moisture | |
|------------|---------|---------------------------------------|-------------------|---------------|-----------------------|------------------|-------------------|-----------------|----------------------|--------------------------|----------------------|------------|--------|
| | | Total | | | Sand | | | | | | | 1/3 Bar | 15 Bar |
| | | Sand (2.0-0.05) | Silt (0.05-0.002) | Clay (<0.002) | Very coarse (2.0-1.0) | Coarse (1.0-0.5) | Medium (0.5-0.25) | Fine (0.25-0.1) | Very fine (0.1-0.05) | | | | |
| 0-28 | Ap | 24.43 | 17.76 | 57.81 | 5.30 | 3.89 | 3.78 | 7.14 | 4.32 | 20 | c | 41.40 | 29.60 |
| 28-55 | Bw | 18.77 | 15.59 | 65.64 | 2.74 | 3.73 | 2.85 | 4.83 | 4.61 | 10 | c | 46.71 | 35.18 |

| Depth (cm) | pH (1:2.5) | | | E.C. (1:2.5) dS m ⁻¹ | O.C. % | CaCO ₃ % | Exchangeable bases | | | | | CEC | CEC/Clay | Base saturation % | ESP % |
|------------|------------|-------------------|-------|---------------------------------|--------|---------------------|--------------------|----|------|-------|-------|-------|----------|-------------------|-------|
| | Water | CaCl ₂ | M KCl | | | | Ca | Mg | K | Na | Total | | | | |
| | | | | | | | | | | | | | | | |
| 0-28 | 8.86 | - | - | 0.483 | 0.63 | 15.48 | - | - | 0.86 | 6.27 | - | 37.00 | 0.64 | - | 6.78 |
| 28-55 | 8.61 | - | - | 1.4 | 0.23 | 13.68 | - | - | 0.68 | 12.27 | - | 53.20 | 0.81 | - | 9.22 |

Contd...

Series Name: Dombarahalli (DRL), **Pedon:** R-8

Location: 15°13'96.2"N, 75°57'48.6" E Rangunathanahalli village, Koppal taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bangalore.

Classification: Very fine, smectitic (calc), isohyperthermic Typic Haplusterts

| Depth (cm) | Horizon | Size class and particle diameter (mm) | | | | | | | | Coarse fragments w/w (%) | Texture Class (USDA) | % Moisture | |
|------------|---------|---------------------------------------|-------------------|---------------|-----------------------|------------------|-------------------|-----------------|----------------------|--------------------------|----------------------|------------|--------|
| | | Total | | | Sand | | | | | | | 1/3 Bar | 15 Bar |
| | | Sand (2.0-0.05) | Silt (0.05-0.002) | Clay (<0.002) | Very coarse (2.0-1.0) | Coarse (1.0-0.5) | Medium (0.5-0.25) | Fine (0.25-0.1) | Very fine (0.1-0.05) | | | | |
| 0-15 | Ap | 28.25 | 19.48 | 52.27 | 4.76 | 4.44 | 4.87 | 8.23 | 5.95 | - | c | 39.86 | 27.20 |
| 15-27 | BA1 | 21.55 | 20.00 | 58.45 | 3.76 | 2.76 | 3.43 | 6.30 | 5.30 | - | c | 46.35 | 34.84 |
| 27-45 | Bss1 | 14.86 | 20.89 | 64.25 | 2.46 | 2.23 | 2.23 | 3.91 | 4.02 | - | c | 57.99 | 41.06 |
| 45-80 | Bss2 | 10.42 | 19.04 | 70.54 | 1.74 | 1.97 | 1.27 | 2.78 | 2.66 | - | c | 66.36 | 36.24 |

| Depth (cm) | pH (1:2.5) | | | E.C. (1:2.5) dS m ⁻¹ | O.C. % | CaCO ₃ % | Exchangeable bases | | | | | CEC | CEC/Clay | Base saturation % | ESP % |
|------------|------------|-------------------|-------|---------------------------------|--------|---------------------|--------------------|----|------|-------|-------|-------|----------|-------------------|-------|
| | Water | CaCl ₂ | M KCl | | | | Ca | Mg | K | Na | Total | | | | |
| | | | | | | | | | | | | | | | |
| 0-15 | 8.78 | - | - | 0.42 | 0.32 | 12.35 | - | - | 0.59 | 4.25 | - | 49.70 | 0.95 | 100.00 | 5.62 |
| 15-27 | 9.03 | - | - | 0.61 | 0.30 | 12.48 | - | - | 0.30 | 8.96 | - | 57.23 | 0.98 | 100.00 | 10.07 |
| 27-45 | 9.10 | - | - | 0.67 | 0.34 | 11.70 | - | - | 0.25 | 11.85 | - | 60.71 | 0.95 | 100.00 | 14.05 |
| 45-80 | 9.18 | - | - | 0.86 | 0.32 | 13.39 | - | - | 0.27 | 15.40 | - | 63.33 | 0.90 | 100.00 | 18.45 |

Contd...

Series Name: Narsapura (NSP), **Pedon:** A2/RM-2

Location: 15°19'86.9"N, 75°57'86.1"E, Kavalura village, Koppal taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bangalore

Classification: Very fine, smectitic (calc), isohyperthermic Vertic Haplustepts

| Depth (cm) | Horizon | Size class and particle diameter (mm) | | | | | | | | Coarse fragments w/w (%) | Texture Class (USDA) | % Moisture | |
|------------|---------|---------------------------------------|-------------------|---------------|-----------------------|------------------|-------------------|-----------------|----------------------|--------------------------|----------------------|------------|--------|
| | | Total | | | Sand | | | | | | | 1/3 Bar | 15 Bar |
| | | Sand (2.0-0.05) | Silt (0.05-0.002) | Clay (<0.002) | Very coarse (2.0-1.0) | Coarse (1.0-0.5) | Medium (0.5-0.25) | Fine (0.25-0.1) | Very fine (0.1-0.05) | | | | |
| 0-29 | Ap | 31.32 | 16.52 | 52.16 | 5.51 | 5.40 | 5.51 | 9.83 | 5.08 | 10 | c | 38.86 | 27.64 |
| 29-52 | Bw1 | 13.30 | 22.08 | 64.62 | 2.52 | 2.41 | 2.41 | 3.67 | 2.29 | 05 | c | 49.88 | 40.05 |
| 52-77 | BW2 | 13.22 | 17.39 | 69.40 | 3.56 | 2.41 | 1.95 | 2.76 | 2.53 | 05 | c | 51.33 | 41.55 |

| Depth (cm) | pH (1:2.5) | | | E.C. (1:2.5) | O.C. | CaCO ₃ | Exchangeable bases | | | | | CEC | CEC/Clay | Base saturation | ESP | | | |
|------------|------------|-------------------|-------|--------------|------|-------------------|--------------------|---|------|-----------------------|----|-------|----------|-----------------|-------|---|----|-------|
| | Water | CaCl ₂ | M KCl | | | | dS m ⁻¹ | % | % | Ca | Mg | | | | | K | Na | Total |
| | | | | | | | | | | cmol kg ⁻¹ | | | | | | % | % | |
| 0-29 | 9.16 | - | - | 0.615 | 0.23 | 9.36 | - | - | 0.72 | 10.98 | - | 51.09 | 0.98 | - | 8.60 | | | |
| 29-52 | 8.69 | - | - | 2.01 | 0.5 | 8.64 | - | - | 0.55 | 24.42 | - | 60.63 | 0.94 | - | 16.11 | | | |
| 52-77 | 8.52 | - | - | 2.68 | 0.46 | 7.68 | - | - | 0.50 | 25.65 | - | 60.74 | 0.88 | - | 16.90 | | | |

Contd...

Series Name: Kavalura (KVR), **Pedon:** A2/RM-9

Location: 15°18'86.8"N, 75°56'56.3"E, Kavalura village, Koppal Taluk and District

Analysis at: NBSS&LUP, Regional Centre, Bangalore. **Classification:** Fine, smectitic (calc), isohyperthermic Typic Haplusterts

| Depth (cm) | Horizon | Size class and particle diameter (mm) | | | | | | | | Coarse fragments w/w (%) | Texture Class (USDA) | % Moisture | |
|------------|---------|---------------------------------------|-------------------|---------------|-----------------------|------------------|-------------------|-----------------|----------------------|--------------------------|----------------------|------------|--------|
| | | Total | | | Sand | | | | | | | 1/3 Bar | 15 Bar |
| | | Sand (2.0-0.05) | Silt (0.05-0.002) | Clay (<0.002) | Very coarse (2.0-1.0) | Coarse (1.0-0.5) | Medium (0.5-0.25) | Fine (0.25-0.1) | Very fine (0.1-0.05) | | | | |
| 0-24 | Ap | 36.18 | 17.80 | 46.02 | 7.04 | 7.47 | 6.62 | 9.28 | 5.76 | 10 | c | 28.20 | 18.75 |
| 24-50 | Bss1 | 38.79 | 15.36 | 45.85 | 6.25 | 6.25 | 9.70 | 10.67 | 5.93 | 05 | c | 27.16 | 18.81 |
| 50-85 | Bss2 | 36.80 | 14.66 | 48.54 | 9.63 | 8.23 | 7.03 | 7.58 | 4.33 | <5 | c | 30.16 | 22.17 |
| 85-124 | Bss3 | 22.66 | 17.24 | 60.09 | 4.18 | 3.85 | 5.28 | 5.06 | 4.29 | <5 | c | 40.34 | 31.42 |

| Depth (cm) | pH (1:2.5) | | | E.C. (1:2.5) dS m ⁻¹ | O.C. % | CaCO ₃ % | Exchangeable bases | | | | | CEC | CEC/Clay | Base saturation % | ESP % |
|------------|------------|-------------------|-------|---------------------------------|--------|---------------------|--------------------|----|------|------|-------|-------|----------|-------------------|-------|
| | Water | CaCl ₂ | M KCl | | | | Ca | Mg | K | Na | Total | | | | |
| | | | | | | | | | | | | | | | |
| 0-24 | 8.4 | - | - | 0.265 | 0.2 | 8.04 | - | - | 0.97 | 0.65 | | 43.25 | 0.94 | | 0.60 |
| 24-50 | 9.27 | - | - | 0.23 | 0.37 | 8.04 | - | - | 0.31 | 3.21 | | 41.66 | 0.91 | | 3.08 |
| 50-85 | 9.44 | - | - | 0.297 | 0.41 | 8.64 | - | - | 0.35 | 6.43 | | 43.99 | 0.91 | | 5.85 |
| 85-124 | 9.37 | - | - | 0.46 | 0.41 | 11.40 | - | - | 0.42 | 7.99 | | 51.09 | 0.85 | | 6.26 |

Contd...

Series Name: Lakshmgudda (LGD) **Pedon:** R-2

Location: 15°13'08.2"N, 76°15'27.3" E Raghunathanahalli village, Koppal taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bangalore. **Classification:** Fine, smectitic (calc), isohyperthermic Typic Haplusterts

| Depth (cm) | Horizon | Size class and particle diameter (mm) | | | | | | | | Coarse fragments w/w (%) | Texture Class (USDA) | % Moisture | |
|------------|---------|---------------------------------------|-------------------|---------------|-----------------------|------------------|-------------------|-----------------|----------------------|--------------------------|----------------------|------------|--------|
| | | Total | | | Sand | | | | | | | 1/3 Bar | 15 Bar |
| | | Sand (2.0-0.05) | Silt (0.05-0.002) | Clay (<0.002) | Very coarse (2.0-1.0) | Coarse (1.0-0.5) | Medium (0.5-0.25) | Fine (0.25-0.1) | Very fine (0.1-0.05) | | | | |
| 0-17 | Ap | 50.60 | 14.29 | 35.11 | 4.53 | 7.86 | 12.49 | 5.18 | 20.54 | - | sc | 28.99 | 18.05 |
| 17-40 | Bss1 | 40.22 | 16.89 | 42.89 | 3.03 | 7.03 | 9.95 | 13.84 | 6.38 | - | c | 34.09 | 23.60 |
| 40-65 | Bss2 | 37.58 | 17.32 | 45.10 | 2.94 | 6.86 | 10.24 | 11.55 | 5.99 | - | c | 35.23 | 24.68 |
| 65-92 | Bss3 | 30.69 | 19.33 | 49.97 | 2.09 | 5.06 | 8.03 | 8.25 | 7.26 | - | c | 40.92 | 29.53 |
| 92-124 | Bss4 | 29.82 | 21.09 | 49.09 | 2.99 | 5.76 | 7.65 | 3.33 | 10.09 | - | c | 44.40 | 31.52 |
| 124-145 | Bss5 | 28.77 | 22.78 | 48.44 | 2.63 | 5.36 | 7.44 | 8.86 | 4.49 | - | c | 43.05 | 30.08 |

| Depth (cm) | pH (1:2.5) | | | E.C. (1:2.5) | O.C. | CaCO ₃ | Exchangeable bases | | | | | CEC | CEC/Clay | Base saturation | ESP | | | | | | |
|------------|------------|-------------------|-------|--------------|------|-------------------|--------------------|---|------|------|----|-------|----------|-----------------|------|---|----|-------|-----------------------|---|---|
| | Water | CaCl ₂ | M KCl | | | | dS m ⁻¹ | % | % | Ca | Mg | | | | | K | Na | Total | cmol kg ⁻¹ | % | % |
| | | | | | | | | | | | | | | | | | | | | | |
| 0-17 | 8.03 | - | - | 1.93 | 0.94 | 8.84 | - | - | 0.35 | 5.02 | - | 32.37 | 0.92 | 100.00 | 1.82 | | | | | | |
| 17-40 | 7.68 | - | - | 1.85 | 0.98 | 8.97 | - | - | 0.16 | 4.38 | - | 42.18 | 0.98 | 100.00 | 1.66 | | | | | | |
| 40-65 | 7.61 | - | - | 1.75 | 0.94 | 9.36 | - | - | 0.16 | 3.77 | - | 42.84 | 0.95 | 100.00 | 1.32 | | | | | | |
| 65-92 | 7.82 | - | - | 1.65 | 1.07 | 9.23 | - | - | 0.22 | 5.02 | - | 47.85 | 0.96 | 100.00 | 2.82 | | | | | | |
| 92-124 | 8.46 | - | - | 1.10 | 1.13 | 10.40 | - | - | 0.23 | 6.72 | - | 47.31 | 0.96 | 100.00 | 7.95 | | | | | | |
| 124-145 | 8.66 | - | - | 0.94 | 0.88 | 14.17 | - | - | 0.22 | 6.48 | - | 44.80 | 0.92 | 100.00 | 8.17 | | | | | | |

Contd...

Series Name: Alawandi (AWD)

Pedon: R-16

Location : 15°13'08.2"N, 76°15'27.3" E Neeralagi village, Koppal taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bangalore.

Classification: Fine, smectitic (calc), isohyperthermic Typic Haplusterts

| Depth (cm) | Horizon | Size class and particle diameter (mm) | | | | | | | | Coarse fragments w/w (%) | Texture Class (USDA) | % Moisture | |
|------------|---------|---------------------------------------|-------------------|---------------|-----------------------|------------------|-------------------|-----------------|----------------------|--------------------------|----------------------|------------|--------|
| | | Total | | | Sand | | | | | | | 1/3 Bar | 15 Bar |
| | | Sand (2.0-0.05) | Silt (0.05-0.002) | Clay (<0.002) | Very coarse (2.0-1.0) | Coarse (1.0-0.5) | Medium (0.5-0.25) | Fine (0.25-0.1) | Very fine (0.1-0.05) | | | | |
| 0-17 | Ap | 20.88 | 25.75 | 53.37 | 3.31 | 4.31 | 4.31 | 5.19 | 3.76 | - | c | 33.11 | 25.58 |
| 17-39 | Bss1 | 25.99 | 19.79 | 54.22 | 5.04 | 5.48 | 5.04 | 5.92 | 4.50 | - | c | 33.11 | 26.23 |
| 39-70 | Bss2 | 26.76 | 17.80 | 55.44 | 2.93 | 5.31 | 5.53 | 7.37 | 5.63 | - | c | 36.15 | 28.67 |
| 70-111 | Bss3 | 23.83 | 20.25 | 55.93 | 4.15 | 4.81 | 4.92 | 6.01 | 3.93 | - | c | 43.60 | 33.71 |
| 111-139 | Bss4 | 21.21 | 20.40 | 58.40 | 2.79 | 4.80 | 4.91 | 5.25 | 3.46 | - | c | 46.92 | 36.28 |
| 139-162 | Bss5 | 13.15 | 20.96 | 65.90 | 1.69 | 2.47 | 2.36 | 3.37 | 3.26 | - | c | 54.96 | 41.81 |

| Depth (cm) | pH (1:2.5) | | | E.C. (1:2.5) | O.C. | CaCO ₃ | Exchangeable bases | | | | | CEC | CEC/Clay | Base saturation | ESP |
|------------|------------|-------------------|-------|--------------------|------|-------------------|-----------------------|----|------|-------|-------|-------|----------|-----------------|------|
| | Water | CaCl ₂ | M KCl | | | | Ca | Mg | K | Na | Total | | | | |
| | | | | dS m ⁻¹ | % | % | cmol kg ⁻¹ | | | | | % | % | | |
| 0-17 | 8.10 | - | - | 0.37 | 0.52 | 9.48 | - | - | 0.40 | 1.56 | - | 51.30 | 0.96 | 100.00 | 1.22 |
| 17-39 | 8.60 | - | - | 0.24 | 0.52 | 9.60 | - | - | 0.14 | 4.60 | - | 52.60 | 0.97 | 100.00 | 3.50 |
| 39-70 | 8.89 | - | - | 0.27 | 0.52 | 9.48 | - | - | 0.16 | 2.41 | - | 53.90 | 0.97 | 100.00 | 1.78 |
| 70-111 | 9.10 | - | - | 0.35 | 0.54 | 11.28 | - | - | 0.15 | 8.95 | - | 54.10 | 0.97 | 100.00 | 6.61 |
| 111-139 | 9.15 | - | - | 0.41 | 0.58 | 10.80 | - | - | 0.15 | 7.36 | - | 56.10 | 0.96 | 100.00 | 5.24 |
| 139-162 | 9.16 | - | - | 0.50 | 0.50 | 15.48 | - | - | 0.19 | 10.19 | - | 61.66 | 0.94 | 100.00 | 6.61 |

INTERPRETATION FOR LAND RESOURCE MANAGEMENT

The most important soil and site characteristics that affect the land use and conservation needs of an area are land capability, land irrigability, soil depth, soil texture, coarse fragments, available water capacity, soil slope, soil erosion, soil reaction etc. These are interpreted from the data base generated through land resource inventory and several thematic maps are generated. These would help in identifying the areas suitable for growing crops and, soil and water conservation measures and structures needed thus helping to maintain good soil health for sustained crop production. The various thematic maps generated are described below.

5.1 Land Capability Classification

Land capability classification is an interpretative grouping of soil map units (soil phases) mainly based on inherent soil characteristics, external land features and environmental factors that limit the use of land for agriculture, pasture, forestry, or other uses on a sustained basis (IARI, 1971). The land and soil characteristics used to group the land resources in an area into various land capability classes, subclasses and units are

Soil characteristics: Soil depth, soil texture, coarse fragments, soil reaction, available water capacity, calcareousness, salinity/alkali *etc.*

Land characteristics: Slope, erosion, drainage, rock outcrops.

Climate: Total rainfall and its distribution, and length of crop growing period.

The Land Capability Classification system is divided into land capability classes, subclasses and units based on the level of information available. Eight land capability classes are recognized. They are

Class I: They are very good lands that have no limitations or very few limitations that restrict their use.

Class II: They are good lands that have minor limitations and require moderate conservation practices.

Class III: They are moderately good lands that have severe limitations that reduce the choice of crops or that require special conservation practices.

Class IV: They are fairly good lands that have very severe limitations that reduce the choice of crops or that require very careful management.

Class V: Soils in these lands are not likely to erode, but have other limitations like wetness that are impractical to remove and as such not suitable for agriculture, but suitable for pasture or forestry with minor limitations.

Class VI: The lands have severe limitations that make them generally unsuitable for cultivation, but suitable for pasture or forestry with moderate limitations.

Class VII: The lands have very severe limitations that make them unsuitable for cultivation, but suitable for pasture or forestry with major limitations.

Class VIII: Soil and other miscellaneous areas (rock lands) that have very severe limitations that nearly preclude their use for any crop production, but suitable for wildlife, recreation and installation of wind mills.

The land capability subclasses are recognized based on the dominant limitations observed within a given land capability class. The subclasses are designated by adding a lower case letter like ‘e’, ‘w’, ‘s’, or ‘c’ to the class numeral. The subclass “e” indicates that the main hazard is risk of erosion, “w” indicates drainage or wetness as a limitation for plant growth, “s” indicates shallow soil depth, coarse or heavy textures, calcareousness, salinity/alkalinity or gravelliness and “c” indicates limitation due to climate.

The land capability subclasses have been further subdivided into land capability units based on the kinds of limitations present in each subclass. Ten land capability units are used in grouping the soil map units. They are stony or rocky (0), erosion hazard (slope, erosion) (1), coarse texture (sand, loamy sand, sandy loam) (2), fine texture (cracking clay, silty clay) (3), slowly permeable subsoil (4), coarse underlying material (5), salinity/alkali (6), stagnation, overflow, high ground water table (7), soil depth (8) and fertility problems (9). The capability units thus identified have similar soil and land characteristics that respond similarly to a given level of management. The soils of the microwatershed have been classified upto land capability subclass level.

The 31 soil map units identified in the Haligeri-3 Microwatershed are grouped under three land capability classes and five land capability subclasses (Fig. 5.1).




Fig. 5.1 Land Capability map of Haligeri-3 Microwatershed

Entire cultivated area in the microwatershed is suitable for agriculture. Good lands (Class II) cover a maximum area of about 570 ha (71%) and distributed in the major part of the microwatershed with minor problems of soil and erosion. Moderately good lands (Class III) occupy an area of about 90 ha (11%) and distributed in the eastern, southern and western part of the microwatershed with severe limitations of soil and erosion. An area of about 135 ha (17%) is fairly good lands and distributed in the western, southern and eastern part of the microwatershed with very severe limitations of soil and erosion. An area of about 1 ha (<1%) is covered by railway and 6 ha (<1%) is under water body.

5.2 Soil Depth

Soil depth refers to the depth of the soil occurring above the parent material or hard rock. The depth of the soil determines the effective rooting depth for plants and in accordance with soil texture, mineralogy and gravel content, the capacity of the soil column to hold water and nutrient availability. Soil depth is one of the most important soil characteristic that is used in differentiating soils into different soil series. The soil depth classes used in identifying soils in the field are very shallow (<25 cm), shallow (25-50 cm), moderately shallow (50-75 cm), moderately deep (75-100 cm), deep (100-150 cm) and very deep (>150 cm). They were used to classify the soils into different depth classes and a soil depth map was generated (Fig. 5.2). The area extent and their geographical distribution in the microwatershed is given in Fig. 5.2.




Fig. 5.2 Soil Depth map of Haligeri-3 Microwatershed

Shallow soils (25-50 cm) cover about 44 ha (6%) and distributed in the eastern and southern part of the microwatershed. Moderately shallow (50-75 cm) soils cover an area of about 189 ha (24%) and distributed in the eastern, central and western part of the microwatershed. An area of about 272 ha (34%) is moderately deep soils (75-100 cm) and distributed in the eastern, central and western part of the microwatershed. Deep to very deep (100- >150 cm) soils occupy a maximum area of about 289 ha (36%) and distributed in the major part of the microwatershed.

The most productive lands cover about 289 ha (36%) where all climatically adopted long duration crops be grown. Problem soils cover an area of 44 ha (6%) where occasionally short duration crops can be grown. The probability of crop failure is very high

5.3 Surface Soil Texture

Texture is an expression to indicate the coarseness or fineness of the soil as determined by the relative proportion of primary particles of sand, silt and clay. It has a direct bearing on the structure, porosity, adhesion and consistence. The surface layer of a soil to a depth of about 25 cm is the layer that is most used by crops and plants. The surface soil textural class provides a guide to understanding soil-water retention and availability, nutrient holding capacity, infiltration, workability, drainage, physical and chemical behavior, microbial activity and crop suitability. The textural classes used for LRI were used to classify and a surface soil texture map was generated. The area extent and their geographical distribution in the microwatershed is shown in Fig 5.3.

An area of about 126 ha (16%) is loamy (sandy loam and sandy clay loam) at the surface and distributed in the western, central and southern part of the microwatershed. Clayey (sandy clay and clay) soils cover about 669 ha (83%) and are distributed in the major part of the microwatershed.

The most productive lands with respect to surface soil texture are clayey soils that (83%) have high potential for soil-water retention and availability and nutrient retention and availability, but have more problems of drainage, infiltration, workability and other physical problems. The other productive lands are loamy (16%) soils which also have high potential for soil- water retention and nutrient availability but have no drainage or other physical problems.




Fig. 5.3 Surface Soil Texture map of Haligeri-3 Microwatershed

5.4 Soil Gravelliness

Gravel is the term used for describing coarse fragments between 2 mm and 7.5 cm diameter and stones for those between 7.5 cm and 25 cm. The presence of gravel and stones in soil reduces the volume of soil responsible for moisture and nutrient storage, drainage, infiltration and runoff, and hinders plant growth by impeding root growth and seedling emergence, intercultural operations and farm mechanization. The gravelliness classes used in LRI were used to classify the soils and using these classes a gravelliness map was generated. The area extent and their geographic distribution in the microwatershed is shown in Fig. 5.4.

The soils that are non-gravelly (<15% gravel) cover an area of about 464 ha (58%) and distributed in the major part of the microwatershed. An area of about 307 ha (38%) is covered by gravelly (15-35% gravel) soils and are distributed in the northwestern, western and central part of the microwatershed. Very gravelly soils (35-60%) cover an area of about 24 ha (3%) and distributed in the central part of the microwatershed (Fig. 5.4).

The most productive lands with respect to gravelliness are found to be 58 per cent. They are non-gravelly with less than 15 per cent gravel and have potential for growing both annual and perennial crops. Very gravelly soils (35-60%) cover an area of about 3 per cent where only short duration crops can be grown.




Fig. 5.4 Soil Gravelliness map of Haligeri-3 Microwatershed

5.5 Available Water Capacity

The soil available water capacity (AWC) is estimated based on the ability of the soil column to retain water between the tensions of 0.33 and 15 bar in a depth of 100 cm or the entire solum if the soil is shallower. The AWC of the soils (soil series) as estimated by considering the soil texture, mineralogy, soil depth and gravel content (Sehgal *et al.*, 1990) and accordingly the soil map units were grouped into five AWC classes *viz*, very low (<50 mm/m), low (50-100 mm/m), medium (100-150 mm/m), high (150-200 mm/m) and very high (>200 mm/m) and using these values, an AWC map was generated. The area extent and their geographic distribution of different AWC classes in the microwatershed is shown in Fig. 5.5.

An area of about 64 ha (8%) in the microwatershed has soils that are very low (<50 mm/m) in available water capacity and are distributed in the southern and northeastern part of the microwatershed. Maximum area of about 305 ha (38%) has soils that are low (51 to 100 mm/m) in available water capacity and are distributed in the major part of the microwatershed. An area of about 298 ha (37%) has soils that are medium (101-150 mm/m) in available water capacity and are distributed in the northeastern part of the microwatershed. An area of about 127 ha (16%) very high (>200 mm/min) in available water capacity and distributed in the northern and southern part of the microwatershed.

An area of about 369 ha (46%) in the microwatershed has soils that are problematic with regard to available water capacity. Here, only short duration crops can

be grown and the probability of crop failure is very high. These areas are best put to other alternative uses. An area of about 127 ha (16%) has soils that have high potential (>200 mm/m) with regard to available water capacity where all climatically adapted long duration crops can be grown successfully.




Fig. 5.5 Soil Available Water Capacity map of Haligeri-3 Microwatershed

5.6 Soil Slope

Soil slope refers to the inclination of the surface of the land. It is defined by gradient, shape and length, and is an integral feature of any soil as a natural body. Slope is considered important in soil genesis, land use and land development. The length and gradient of slope influences the rate of runoff, infiltration, erosion and deposition. The soil map units were grouped into two slope classes and a slope map was generated showing the area extent and their geographic distribution of different slope classes in the microwatershed (Fig. 5.6).

Nearly level (0-1%) lands cover an area of about 37 ha (5%) and distributed in the western and southern part of the microwatershed. Very gently sloping (1-3%) lands cover a maximum area of about 757 ha (94%) and distributed in the major part of the microwatershed. In all these areas, all climatically adapted annual and perennial crops can be grown without much soil and water conservation and other land development measures.




Fig. 5.6 Soil Slope map of Haligeri-3 Microwatershed

5.7 Soil Erosion

Soil erosion refers to the wearing away of the earth's surface by the forces of water, wind and ice involving detachment and transport of soil by raindrop impact. It is used for accelerated soil erosion resulting from disturbance of the natural landscape by burning, excessive grazing and indiscriminate felling of forest trees and tillage, all usually by man. The erosion classes showing an estimate of the current erosion status as judged from field observations in the form of rills, gullies or a carpet of gravel on the surface are recorded. Four erosion classes, viz, slight erosion (e1), moderate erosion (e2), severe erosion (e3) and very severe erosion (e4) are recognized. The soil map units were grouped into different erosion classes and a soil erosion map generated. The area extent and their spatial distribution in the microwatershed is given in Figure 5.7.

Slightly eroded lands cover an area of about 255 ha (32 %) and distributed in the southern, central and northeastern part of the microwatershed. Maximum area of about 540 ha (67 %) is moderately eroded (e2 class) and distributed in the major part of the microwatershed. Moderately eroded lands are problematic and need appropriate soil and water conservation and other land development measures.




Fig. 5.7 Soil Erosion map of Haligeri-3 Microwatershed

FERTILITY STATUS

Soil fertility plays an important role in increasing crop yield. The adoption of high yielding varieties that require high amounts of nutrients has resulted in deficiency symptoms in crops and plants due to imbalanced fertilization and poor inherent fertility status, as these areas are characterized by low rainfall and high temperatures. Hence, it is necessary to know the fertility (macro and micro nutrients) status of the soils of the watersheds for assessing the kind and amount of fertilizers required for each of the crop intended to be grown. For this purpose, the surface soil samples collected from the grid points (one soil sample at every 320 m grid interval) all over the microwatershed through land resource inventory in the year 2018 were analyzed for pH, EC, organic carbon, available phosphorus and potassium, and for micronutrients like zinc, boron, copper, iron and manganese, and secondary nutrient sulphur.

Soil fertility data generated has been assessed and individual maps for all the nutrients for the microwatershed have been generated by using the Kriging method under GIS. The village/survey number wise fertility data for the microwatershed is given in Appendix-II.

6.1 Soil Reaction (pH)

The soil analysis of the Haligeri-3 microwatershed for soil reaction (pH) showed that neutral (pH 6.5-7.3) soils cover an area of about 37 ha (5%) and distributed in the western part of the microwatershed. Slightly alkaline soils (pH 7.3-7.8) cover a maximum area of about 492 ha (61%) and distributed in the major part of the microwatershed. Strongly to very strongly alkaline (pH 8.4->9.0) soils cover an area of about 266 ha (33%) and distributed in the eastern and northeastern part of the microwatershed (Fig.6.1). An area of about 37 ha (5%) is neutral and 758 ha (94%) is alkaline in reaction.

6.2 Electrical Conductivity (EC)

The Electrical Conductivity of the soils of the entire microwatershed area is $<2 \text{ dSm}^{-1}$ (Fig 6.2) and as such the soils are non-saline.

6.3 Organic Carbon

An area of about 263 ha (33%) is low ($<0.5\%$) and distributed in the northern and eastern part of the microwatershed. Maximum area of about 343 ha (43%) is medium (0.5-0.75%) and distributed in the major part of the microwatershed. An area of about 189 ha (24%) is high ($>0.75\%$) and distributed in the southern and southwestern part of the microwatershed (Fig.6.3).

6.4 Available Phosphorus

An area of about 251 ha (31%) is low ($<23 \text{ kg/ha}$) and distributed in the northwestern part of the microwatershed. Maximum area of about 531 ha (66%) is

medium (23-57 kg/ha) in available phosphorus and distributed in the major part of the microwatershed. An area of about 13 ha (2%) is high (>57 kg/ha) and distributed in the western and northeastern part of the microwatershed. The areas with high phosphorus content reduce 25 per cent from the recommended dose to avoid the excess application of fertilizer. Apply additional 25% phosphorus in areas where it is low and medium (Fig 6.4).




Fig.6.1 Soil Reaction (pH) map of Haligeri-3 Microwatershed




Fig.6.2 Electrical Conductivity (EC) map of Haligeri-3 Microwatershed




Fig.6.3 Soil Organic Carbon map of Haligeri-3 Microwatershed

6.5 Available Potassium

Available potassium is medium (145-337 kg/ha) in 120 ha (15%) and distributed in the western part of the microwatershed. Maximum area of about 675 ha (84%) is high (>337 kg/ha) and distributed in the major part of the microwatershed. The areas with high potassium content reduce 25 per cent from the recommended dose to avoid the excess application of fertilizer. Apply additional 25% potassium in areas where it is medium (Fig 6.5).

6.6 Available Sulphur

Soil analysis of available sulphur content in Haligeri-3 microwatershed showed that an area of about 300 ha (37%) is low and distributed in the western, eastern and central part of the microwatershed. Maximum area of about 462 ha (58%) is medium (10-20 ppm) in available sulphur content and distributed in the major part of the microwatershed. An area of about 32 ha (4%) is high (>20 ppm) and distributed in the southern part of the microwatershed (Fig.6.6). The areas that are low and medium in available sulphur need to be applied with magnesium sulphate or gypsum or factomphos (p) fertilizer (13% sulphur) for 2-3 years for the deficiency to be corrected.

6.7 Available Boron

An area of about 294 ha (37%) is low (< 0.5ppm) in available boron and distributed in the western, central and eastern part of the microwatershed. A maximum area of about 453 ha (56%) is medium (0.5-1.0 ppm) and distributed in the major part of the microwatershed. An area of about 48 ha (6%) is high(>1.0 ppm) and distributed in the southern part of the microwatershed (Fig.6.7).

6.8 Available Iron

Available iron content in the soils of the Haligeri-3 microwatershed is deficient (<4.5 ppm) in an entire area of the microwatershed (Fig 6.8).

6.9 Available Manganese

Available manganese content is sufficient (>1.0 ppm) in the entire microwatershed area (Fig 6.9).

6.10 Available Copper

Available copper content is sufficient (>0.2 ppm) in the entire microwatershed area (Fig 6.10).

6.11 Available Zinc

Available zinc content is deficient (<0.6 ppm) in an entire part of the microwatershed (Fig 6.11).




Fig.6.4 Soil Available Phosphorus map of Haligeri-3 Microwatershed




Fig.6.5 Soil Available Potassium map of Haligeri-3 Microwatershed




Fig.6.6 Soil Available Sulphur map of Haligeri-3 Microwatershed




Fig.6.7 Soil Available Boron map of Haligeri-3 Microwatershed




Fig.6.8 Soil Available Iron map of Haligeri-3 Microwatershed




Fig.6.9 Soil Available Manganese map of Haligeri-3 Microwatershed




Fig.6.10 Soil Available Copper map of Haligeri-3 Microwatershed




Fig.6.11 Soil Available Zinc map of Haligeri-3 Microwatershed

LAND SUITABILITY FOR MAJOR CROPS

The soil and land resource units (soil phases) of Haligeri-3 Microwatershed were assessed for their suitability for growing food, fodder, fibre and other horticulture crops by following the procedure as outlined in FAO, 1976 and 1983. Crop requirements were developed for each of the crop from the available research data and also by referring to Naidu *et. al.* (2006) and Natarajan *et. al.* (2015). The soil and land characteristics were matched with the crop requirements to arrive at the crop suitability. The soil and land characteristics table (Table 7.1) were matched with the crop requirements (Tables 7.2-7.32) to arrive at the crop suitability and the crop requirement tables are given at the end of the chapter. In FAO land suitability classification, two orders are recognized. Order S- Suitable and Order N- Not suitable. The orders have classes, subclasses and units. Order S has three classes, Class S1- Highly Suitable, Class S2- Moderately Suitable and Class S3- Marginally Suitable. Order N has two Classes, N1- Currently not Suitable and N2- Permanently not Suitable. There are no subclasses within the Class S1 as they will have very minor or no limitations for crop growth. Classes S2, S3 and N1 are divided into subclasses based on the kinds of limitations encountered. The limitations that affect crop production are 'c' for erratic rainfall and its distribution and length of growing period (LGP), 'e' for erosion hazard, 'r' for rooting condition, 't' for lighter or heavy texture, 'g' for gravelliness or stoniness, 'n' for nutrient availability, 'l' for topography, 'm' for moisture availability, 's' for sodium 'z' for calcareousness and 'w' for drainage. These limitations are indicated as lower case letters to the class symbol. For example, moderately suitable lands with the limitations of soil depth and erosion are designated as S2re. For the microwatershed, the soil mapping units were evaluated and classified up to subclass level.

Using the above criteria, the soil map units of the microwatershed were evaluated and land suitability maps for 31 major agricultural and horticultural crops were generated. The detailed information on the kind of suitability of each of the soil phase for the crops assessed are given village/ survey number wise for the microwatershed in Appendix-III.

7.1 Land Suitability for Sorghum (*Sorghum bicolor*)

Sorghum is one of the major food crop grown in Karnataka in an area of 10.47 lakh ha in Bijapur, Gulbarga, Raichur, Bidar, Belgaum, Dharwad, Bellary, Chitradurga, Mysore and Chamarajnagar districts. The crop requirements for growing sorghum (Table 7.2) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing sorghum was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.1.

Highly suitable (Class S1) lands occupy an area of about 35 (4%) for growing sorghum and occur in the western and southern northern part of the microwatershed. A

maximum area of about 515 ha (64%) is moderately suitable (Class S2) for growing sorghum and distributed in the major part of the microwatershed with minor limitations of gravelliness, rooting depth and calcareousness. An area of about 244 ha (30%) is marginally suitable for growing sorghum and distributed in the eastern, western and central part of the microwatershed. They have moderate limitations of gravelliness, rooting depth, nutrient availability and calcareousness.




Fig. 7.1 Land Suitability map of Sorghum

7.2 Land Suitability for Maize (*Zea mays*)

Maize is one of the most important food crop grown in an area of 13.37 lakh ha in almost all the districts of the State. The crop requirements for growing maize (Table 7.3) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing maize was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.2.

Highly suitable (Class S1) lands occupy an area of about 35 (4%) for growing maize and occur in the southern and western part of the microwatershed. Maximum area of about 514 ha (64%) is moderately suitable (Class S2) for growing maize and distributed in the major part of the microwatershed with minor limitations of gravelliness, rooting depth, calcareousness and texture. An area of about 244 ha (30%) is marginally suitable for growing maize and distributed in the eastern, central and western part of the microwatershed. They have moderate limitations of gravelliness, nutrient availability, calcareousness and rooting depth.




Fig. 7.2 Land Suitability map of Maize

7.3 Land Suitability for Bajra (*Pennisetum glaucum*)

Bajra is one of the major food crop grown in an area of 2.34 lakh ha in Karnataka in the northern districts. The crop requirements (Table 7.4) for growing bajra were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing bajra was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.3.

Highly suitable (Class S1) lands occupy an area of about 35 ha (4 %) for growing bajra and occur in the southern and western part of the microwatershed. An area of about 560 ha (70%) is moderately suitable (Class S2) for growing bajra and distributed in the major part of the microwatershed with minor limitations of texture, rooting depth, calcareousness and gravelliness. An area of about 199 ha (25%) is marginally suitable for growing bajra and distributed in the central and eastern part of the microwatershed with moderate limitations of gravelliness, nutrient availability, calcareousness and rooting depth.




Fig. 7.3 Land Suitability map of Bajra

7.4 Land Suitability for Redgram (*Cajanus cajan*)

Redgram is one of the most important pulse crop grown in an area of 7.28 lakh ha in almost all the districts of the State. The crop requirements for growing redgram (Table 7.5) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing redgram was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.4.

Highly suitable (Class S1) lands occupy an area of about 35 ha (4%) for growing redgram and occur in the southern and western part of the microwatershed. Maximum area of about 504 ha (63%) is moderately suitable (Class S2) for growing redgram and distributed in the major part of the microwatershed. They have minor limitations of gravelliness, calcareousness, rooting depth and texture. Marginally suitable lands (Class S3) occupy an area of about 210 ha (26%) and occur in the eastern, central and western part of the microwatershed. They have moderate limitations of rooting depth, nutrient availability and gravelliness. Area currently not suitable (Class N1) cover about 44 ha (6%) and distributed in the part of the microwatershed with severe limitation of rooting depth and calcareousness.




Fig. 7.4 Land Suitability map of Redgram

7.5 Land Suitability for Bengal gram (*Cicer arietinum*)

Bengal gram is one of the major pulse crop grown in an area of 9.39 lakh ha in northern Karnataka in Bijapur, Gulbarga, Raichur, Bidar, Belgaum, Dharwad and Bellary districts. The crop requirements for growing Bengal gram (Table 7.6) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing Bengal gram was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.5.

An area of about 290 ha (36%) is moderately suitable (Class S2) for growing bengalgram and are distributed in the southern, northern and northeastern part of the microwatershed. They have minor limitation of calcareousness. Marginally suitable (Class S3) lands cover a maximum area of about 504 ha (63%) and are distributed in the major part of the microwatershed. They have moderate limitations of nutrient availability, gravelliness, texture, calcareousness and rooting depth.




Fig. 7.5 Land Suitability map of Bengal gram

7.6 Land Suitability for Groundnut (*Arachis hypogaea*)

Groundnut is one of the major oilseed crop grown in an area of 6.54 lakh ha in Karnataka in most of the districts either as rainfed or irrigated crop. The crop requirements for growing groundnut (Table 7.7) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing groundnut was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.6.

An area of about 293 ha (37%) is moderately suitable (Class S2) for growing groundnut and distributed in the southern, western and central part of the microwatershed. They have minor limitations of gravelliness, rooting depth and texture. Maximum area of about 366 ha (46%) is marginally suitable (Class S3) for growing groundnut and are distributed in the major part of the microwatershed with moderate limitations of gravelliness, rooting depth, calcareousness and texture. Area currently not suitable (Class N1) cover about 135 ha (17%) and distributed in the central and eastern part of the microwatershed with severe limitation of nutrient availability.




Fig. 7.6 Land Suitability map of Groundnut

7.7 Land Suitability for Sunflower (*Helianthus annus*)

Sunflower is one of the most important oilseed crop grown in an area of 3.56 lakh ha in the State in all the districts. The crop requirements for growing sunflower (Table 7.8) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing sunflower was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.7.

An area of about 35 ha (4%) is highly suitable (Class S1) for growing sunflower and are distributed in the western and southern part of the microwatershed. Maximum area of about 524 ha (65%) is moderately suitable (Class S2) and are distributed in the major part of the microwatershed. They have minor limitations of gravelliness, rooting depth, texture and calcareousness. Marginally suitable (Class S3) lands occupy an area of about 55 ha (7%) and are distributed in the western part of the microwatershed with moderate limitations of rooting depth and gravelliness. Area currently not suitable (Class N1) cover about 179 ha (22%) and distributed in the western and central part of the microwatershed with severe limitations of nutrient availability, calcareousness and rooting depth.




Fig. 7.7 Land Suitability map of Sunflower

7.8 Land Suitability for Cotton (*Gossypium hirsutum*)

Cotton is one of the most important fibre crop grown in the State in about 8.75 lakh ha area in Raichur, Dharwad, Belgaum, Gulbarga, Bijapur, Bidar, Bellary, Chitradurga and Chamarajnagar districts. The crop requirements for growing cotton (Table 7.9) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing cotton was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.8.

Maximum area of about 549 ha (69%) is moderately suitable (Class S2) and are distributed in the major part of the microwatershed. They have minor limitations of rooting depth, calcareousness, texture and gravelliness. Marginally suitable (Class S3) lands occupy an area of about 244 ha (30%) and are distributed in the central, eastern and western part of the microwatershed with moderate limitations of rooting depth, texture, nutrient availability, calcareousness and gravelliness.




Fig. 7.8 Land Suitability map of Cotton

7.9 Land Suitability for Chilli (*Capsicum annum L*)

Chilli is one of the most important spice crop grown in an area of 0.42 lakh ha in Karnataka State. The crop requirements for growing chilli (Table 7.10) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing chilli was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.9.

An area of about 35 ha (4%) is highly suitable (Class S1) for growing chilli and are distributed in the central and western part of the microwatershed. Maximum area of about 515 ha (64%) is moderately suitable (Class S2) and are distributed in the major part of the microwatershed. They have minor limitations of texture, rooting depth, calcareousness and gravelliness. Marginally suitable (Class S3) lands cover an area of about 109 ha (14%) and distributed in the eastern, southern and western part of the microwatershed. They have moderate limitations of gravelliness, calcareousness and rooting depth. Area currently not suitable (Class N1) cover about 135 ha (17%) and distributed in the western, southern and eastern part of the microwatershed with severe limitation of nutrient availability.




Fig. 7.9 Land Suitability map of Chilli

7.10 Land Suitability for Tomato (*Solanum lycopersicum*)

Tomato is one of the most important vegetable crop grown in an area of 0.65 lakh ha in almost all the districts of the State. The crop requirements (Table 7.11) for growing tomato were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing tomato was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.10.

An area of about 35 ha (4%) is highly suitable (Class S1) for growing tomato and are distributed in the southern and western part of the microwatershed. An area of about 225 ha (28%) is moderately suitable (Class S2) and are distributed in the western, southern and central part of the microwatershed. They have minor limitations of rooting depth and gravelliness. Marginally suitable (Class S3) lands cover an area of about 399 ha (50%) and distributed in the major part of the microwatershed. They have moderate limitations of gravelliness, texture and rooting depth. Area currently not suitable (Class N1) cover about 135 ha (17%) and distributed in the central and eastern part of the microwatershed with severe limitation of nutrient availability.




Fig. 7.10 Land Suitability map of Tomato

7.11 Land Suitability for Brinjal (*Solanum melongena*)

Brinjal is one of the most important vegetable crop grown in the state. The crop requirements for growing brinjal (Table 7.12) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing brinjal was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.11.

An area of about 35 ha (4%) is highly suitable (Class S1) for growing Brinjal and are distributed in the southern and eastern part of the microwatershed. An area of about 225 ha (28%) is moderately suitable (Class S2) for growing Brinjal and distributed in the southern, central and western part of the microwatershed with minor limitations of rooting depth and gravelliness. Marginally suitable (Class S3) lands cover an area of about 399 ha (50%) and occur in the major part of the microwatershed with moderate limitations of gravelliness, rooting depth, texture and calcareousness. Area currently not suitable (Class N1) cover about 135 ha (17%) and distributed in the eastern and central part of the microwatershed with severe limitation of nutrient availability.




Fig 7.11 Land Suitability map of Brinjal

7.12 Land Suitability for Onion (*Allium cepa L.*)

Onion is one of the most important vegetable crop grown in the state. The crop requirements for growing onion (Table 7.13) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing onion was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.12.

An area of about 35 ha (4%) is highly suitable (Class S1) for growing onion and are distributed in the southern and eastern part of the microwatershed. Maximum area of about 221 ha (28%) is moderately suitable (Class S2) for growing Onion and distributed in the southern, central and western part of the microwatershed with minor limitations of gravelliness and rooting depth. Marginally suitable (Class S3) lands cover a maximum area of about 403 ha (50%) and occur in the major part of the microwatershed with moderate limitations of gravelliness, rooting depth, texture and calcareousness. Area currently not suitable (Class N1) cover about 135 ha (17%) and distributed in the eastern and central part of the microwatershed with severe limitation of nutrient availability.




Fig 7.12 Land Suitability map of Onion

7.13 Land Suitability for Bhendi (*Abelmoschus esculentus*)

Bhendi is one of the most important vegetable crop grown in the state. The crop requirements for growing bhendi (Table 7.14) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing bhendi was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.13.

An area of about 35 ha (4%) is highly suitable (Class S1) for growing Bhendi and are distributed in the southern and eastern part of the microwatershed. An area of about 225 ha (28%) is moderately suitable (Class S2) for growing Bhendi and distributed in the southern, central and western part of the microwatershed with minor limitations of rooting depth and gravelliness. Marginally suitable (Class S3) lands cover an area of about 399 ha (50%) and occur in the major part of the microwatershed with moderate limitations of gravelliness, rooting depth, texture and calcareousness. Area currently not suitable (Class N1) cover about 135 ha (17%) and distributed in the eastern and central part of the microwatershed with severe limitation of nutrient availability.




Fig 7.13 Land Suitability map of Bhendi

7.14 Land Suitability for Drumstick (*Moringa oleifera*)

Drumstick is one of the most important vegetable crop grown in 2403 ha area in the state. The crop requirements for growing drumstick (Table 7.15) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing drumstick was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.14.

An area of about 35 ha (4%) is highly suitable (Class S1) for growing drumstick and are distributed in the southern and western part of the microwatershed. Moderately suitable (Class S2) lands cover an area of about 238 ha (30%) and are distributed in the southern, central and western part of the microwatershed. They have minor limitations of rooting depth, texture, calcareousness and gravelliness. Marginally suitable (Class S3) lands cover a maximum area of about 341 ha (43%) and occur in the major part of the microwatershed. They have moderate limitations of gravelliness, calcareousness and rooting depth. Area currently not suitable (Class N1) cover about 179 ha (22%) and distributed in the eastern, central and western part of the microwatershed with severe limitations of nutrient availability, rooting depth and calcareousness.




Fig. 7.14 Land Suitability map of Drumstick

7.15 Land Suitability for Mulberry (*Morus nigra*)

Mulberry is the most important leaf crop grown for rearing silkworms in about 1.66 lakh ha in all the districts of the state. The crop requirements for growing mulberry (Table 7.16) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing mulberry was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.15.

An area of about 35 ha (4%) is highly suitable (Class S1) for growing mulberry and are distributed in the central and western part of the microwatershed. An area of about 235 ha (29%) is moderately suitable (Class S2) for growing mulberry and distributed in the southern, western and central part of the microwatershed. They have minor limitations of gravelliness and rooting depth. Marginally suitable (Class S3) lands cover a maximum area of about 345 ha (43%) and occur in the major part of the microwatershed. They have moderate limitations of rooting depth, texture, calcareousness and gravelliness. Area currently not suitable (Class N1) cover about 179 ha (22%) and distributed in the northern part of the microwatershed with severe limitation of rooting depth, calcareousness and nutrient availability.




Fig. 7.15 Land Suitability map of Mulberry

7.16 Land Suitability for Mango (*Mangifera indica*)

Mango is one of the most important fruit crop grown in about 1.73 lakh ha in almost all the districts of the State. The crop requirements (Table 7.17) for growing mango were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing mango was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.16.

An area of about 161 ha (20%) is moderately suitable (Class S2) for growing mango and distributed in the western, central and southern part of the microwatershed. They have minor limitations of rooting depth and gravelliness. Marginally suitable (Class S3) lands cover an area of about 398 ha (50%) and occur in the major part of the microwatershed. They have moderate limitations of rooting depth and texture. Area currently not suitable (Class N1) for growing mango cover about 234 ha (29%) and distributed in the eastern, central and western part of the microwatershed with severe limitations of rooting depth, nutrient availability and calcareousness.




Fig. 7.16 Land Suitability map of Mango

7.17 Land Suitability for Sapota (*Manilkara zapota*)

Sapota is one of the most important fruit crop grown in an area of about 29373 ha in almost all the districts of the state. The crop requirements (Table 7.18) for growing sapota were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing sapota was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.17.

An area of about 35 ha (4%) is highly suitable (Class S1) for growing sapota and are distributed in the southern part of the microwatershed. An area of about 234 ha (29%) is moderately suitable (Class S2) for growing sapota and distributed in the southern, central and western part of the microwatershed. They have minor limitations of rooting depth and gravelliness. Marginally suitable (Class S3) lands cover a maximum area of about 345 ha (43%) and occur in the major part of the microwatershed. They have moderate limitations of rooting depth, gravelliness and texture. Area currently not suitable (Class N1) for growing sapota cover about 179 ha (22%) and distributed in the eastern and central part of the microwatershed with severe limitations of rooting depth, nutrient availability and calcareousness.




Fig. 7.17 Land Suitability map of Sapota

7.18 Land Suitability for Pomegranate (*Punica granatum*)

Pomegranate is one of the commercially grown fruit crop in about 18488 ha in Karnataka mainly in Bijapur, Bagalkot, Koppal, Gadag and Chitradurga districts. The crop requirements for growing pomegranate (Table 7.19) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing pomegranate was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.18.

An area of about 35 ha (4%) is highly suitable (Class S1) for growing pomegranate and are distributed in the southern and western part of the microwatershed. Moderately suitable (Class S2) lands occupy an area of about 225 ha (28%) and are distributed in the southern, central and western part of the microwatershed. They have minor limitations of rooting depth and gravelliness. Marginally suitable (Class S3) lands for growing pomegranate occupy a maximum area of about 399 ha (50%) and are distributed in the major part of the microwatershed with moderate limitations of rooting depth, texture, calcareousness and gravelliness. Area currently not suitable (Class N1) cover about 135 ha (17%) and distributed in the eastern and central part of the microwatershed with severe limitation of nutrient availability.




Fig. 7.18 Land Suitability map of Pomegranate

7.19 Land Suitability for Guava (*Psidium guajava*)

Guava is one of the most important fruit crop grown in an area of about 6558 ha in almost all the districts of the state. The crop requirements (Table 7.20) for growing guava were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing guava was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.19.

An area of about 35 ha (4%) is highly suitable (Class S1) for growing guava and are distributed in the central and western part of the microwatershed. Moderately suitable (Class S2) lands occupy an area of about 234 ha (29%) and are distributed in the southern, central and western part of the microwatershed. They have minor limitations of rooting depth and gravelliness. Marginally suitable (Class S3) lands for growing guava occupy an area of about 345 ha (43%) and are distributed in the major part of the microwatershed with moderate limitations of gravelliness, texture and calcareousness. Area currently not suitable (Class N1) cover about 179 ha (22%) and distributed in the western and central part of the microwatershed with severe limitations of rooting depth, nutrient availability and calcareousness.




Fig. 7.19 Land Suitability map of Guava

7.20 Land Suitability for Jackfruit (*Artocarpus heterophyllus*)

Jackfruit is one of the most important fruit crop grown in 5368 ha in all the districts of the state. The crop requirements (Table.7.21) for growing jackfruit were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing jackfruit was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in figure 7.20.

An area of about 35 ha (4%) is highly suitable (Class S1) for growing jackfruit and are distributed in the southern and western part of the microwatershed. Moderately suitable (Class S2) lands occupy an area of about 234 ha (29%) and are distributed in the southern, central and western part of the microwatershed. They have minor limitations of rooting depth and gravelliness. Marginally suitable (Class S3) lands for growing jackfruit occupy a maximum area of about 345 ha (43%) and are distributed in the major part of the microwatershed with moderate limitations of gravelliness, texture and rooting depth. Area currently not suitable (Class N1) cover about 179 ha (22%) and distributed in the western and central part of the microwatershed with severe limitations of rooting depth, nutrient availability and calcareousness.




Fig. 7.20 Land Suitability map of Jackfruit

7.21 Land Suitability for Jamun (*Syzygium cumini*)

Jamun is an important fruit crop grown in almost all the districts of the state. The crop requirements (Table 7.22) for growing jamun were matched with the soil-site characteristics and a land suitability map for growing jamun was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.21.

Moderately suitable (Class S2) lands occupy an area of about 161 ha (20%) and distributed in the southern and western part of the microwatershed. They have minor limitations of rooting depth and gravelliness. Marginally suitable (Class S3) lands cover a maximum area of about 453 ha (57%) and are distributed in the major part of the microwatershed with moderate limitations of rooting depth, calcareousness and gravelliness. Area currently not suitable (Class N1) cover about 179 ha (22%) and distributed in the eastern and central part of the microwatershed with severe limitations of rooting depth, nutrient availability and calcareousness.




Fig. 7.21 Land Suitability map of Jamun

7.22 Land Suitability for Musambi (*Citrus limetta*)

Musambi is one of the most important fruit crop grown in an area of 5446 ha in almost all the districts of the state. The crop requirements (Table 7.23) for growing musambi were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing musambi was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.22.

An area of about 35 ha (4%) is highly suitable (Class S1) for growing musambi and are distributed in the southern and western part of the microwatershed. Maximum area of about 524 ha (65%) is moderately suitable (Class S2) and occur in the major part of the microwatershed. They have minor limitations of calcareousness, gravelliness and rooting depth. An area of about 55 ha (7%) is marginally suitable (Class S3) for growing musambi and are distributed in the western part of the microwatershed with moderate limitations of gravelliness and rooting depth. Area currently not suitable (Class N1) cover about 179 ha (22%) and distributed in the southern and eastern part of the microwatershed with severe limitations of rooting depth, nutrient availability and calcareousness.




Fig. 7.22 Land Suitability map of Musambi

7.23 Land Suitability for Lime (*Citrus sp*)

Lime is one of the most important fruit crop grown in an area of 11752 ha in almost all the districts of the State. The crop requirements for growing lime (Table 7.24) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing lime was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.23.

An area of about 35 ha (4%) is highly suitable (Class S1) for growing lime and are distributed in the southern and western part of the microwatershed. Maximum area of about 524 ha (65%) is moderately suitable (Class S2) and occur in the major part of the microwatershed. They have minor limitations of calcareousness, gravelliness and rooting depth. An area of about 55 ha (7%) is marginally suitable (Class S3) for growing lime and are distributed in the western part of the microwatershed with moderate limitations of gravelliness and rooting depth. Area currently not suitable (Class N1) cover about 179 ha (22%) and distributed in the southern and eastern part of the microwatershed with severe limitations of rooting depth, nutrient availability and calcareousness.




Fig. 7.23 Land Suitability map of Lime

7.24 Land Suitability for Cashew (*Anacardium occidentale*)

Cashew is one of the most important nut crop grown in an area of 7052 ha in almost all the districts of the State. The crop requirements for growing cashew (Table 7.25) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing cashew was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.24.

An area of about 35 ha (4%) is highly suitable (Class S1) for growing cashew and are distributed in the western and southern part of the microwatershed. An area of about 234 ha (29%) is moderately suitable (Class S2) and occur in the western, central and southern part of the microwatershed. They have minor limitations of gravelliness and rooting depth. An area of about 55 ha (7%) is marginally suitable (Class S3) for growing cashew and are distributed in the western part of the microwatershed with moderate limitations of gravelliness and rooting depth. Maximum area of about 469 ha (59%) is currently not suitable (Class N1) for growing cashew and distributed in the major part of the microwatershed with severe limitations of rooting depth, nutrient availability, calcareousness and texture.




Fig. 7.24 Land Suitability map of Cashew

7.25 Land Suitability for Custard Apple (*Annona reticulata*)

Custard apple is one of the most important fruit crop grown in 1426 ha in almost all the districts of the State. The crop requirements(Table 7.26) for growing custard apple were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing custard apple was generated .The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.25.

An area of about 35 ha (4%) is highly suitable (Class S1) for growing custard apple and are distributed in the western and southern part of the microwatershed. A maximum area of about 535 ha (72%) is moderately suitable (Class S2) and occur in the major part of the microwatershed. They have minor limitations of gravelliness, rooting depth and calcareousness. An area of about 44 ha (6%) is marginally suitable (Class S3) for growing custard apple and are distributed in the western and southern part of the microwatershed with moderate limitations of gravelliness, calcareousness and rooting depth. An area of about 135 ha (17%) is currently not suitable (Class N1) for growing custard apple and distributed in the western and central part of the microwatershed with severe limitation of nutrient availability.




Fig. 7.25 Land Suitability map of Custard Apple

7.26 Land Suitability for Amla (*Phyllanthus emblica*)

Amla is one of the most important fruit and medicinal crop grown in an area of 151 ha and distributed in almost all the districts of the state. The crop requirements (Table 7.27) for growing amla were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing amla was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.26.

An area of about 35 ha (4%) is highly suitable (Class S1) for growing amla and are distributed in the western and southern part of the microwatershed. An area of about 290 ha (36%) is moderately suitable (Class S2) and occur in the western, central and southern part of the microwatershed. They have minor limitations of gravelliness and rooting depth. A maximum area of about 334 ha (42%) is marginally suitable (Class S3) for growing amla and are distributed in the major part of the microwatershed with moderate limitations of gravelliness, calcareousness and rooting depth. An area of about 135 ha (17%) is currently not suitable (Class N1) for growing amla and distributed in the eastern and central part of the microwatershed with severe limitation of nutrient availability.




Fig. 7.26 Land Suitability map of Amla

7.27 Land Suitability for Tamarind (*Tamarindus indica*)

Tamarind is one of the most important spice crop grown in 14897 ha in all the districts of the state. The crop requirements (Table 7.28) for growing tamarind were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing tamarind was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.27.

An area of about 161 ha (20%) is moderately suitable (Class S2) and occur in the southern and central part of the microwatershed. They have minor limitations of rooting depth and gravelliness. Maximum area of about 398 ha (50%) is marginally suitable (Class S3) for growing tamarind and are distributed in the major part of the microwatershed with moderate limitations of rooting depth and calcareousness. An area of about 234 ha (29%) is currently not suitable (Class N1) for growing tamarind and distributed in the eastern and central part of the microwatershed with severe limitations of rooting depth, nutrient availability and calcareousness.




Fig. 7.27 Land Suitability map of Tamarind

7.28 Land Suitability for Marigold (*Tagetes erecta*)

Marigold is one of the most important flower crop grown in an area of 9108 ha in almost all the districts of the state. The crop requirements (Table 7.29) for growing marigold were matched with the soil-site characteristics and a land suitability map for growing marigold was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.28.

An area of about 35 ha (4%) is highly suitable (Class S1) for growing marigold and are distributed in the southern and western part of the microwatershed. Maximum area of about 511 ha (64%) is moderately suitable (Class S2) and occur in the major part of the microwatershed. They have minor limitations of gravelliness, rooting depth, calcareousness and texture. An area of about 113 ha (14%) is marginally suitable (Class S3) for growing marigold and are distributed in the southern, eastern and western part of the microwatershed with moderate limitations of gravelliness, calcareousness and rooting depth. An area of about 135 ha (17%) is currently not suitable (Class N1) for growing marigold and distributed in the eastern and central part of the microwatershed with severe limitation of nutrient availability.




Fig. 7.28 Land Suitability map of Marigold

7.29 Land Suitability for Chrysanthemum (*Chrysanthemum indicum*)

Chrysanthemum is one of the most important flower crop grown in an area of 4978 ha in almost all the districts of the State. The crop requirements (Table 7.30) for growing chrysanthemum were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing chrysanthemum was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed is given in Figure 7.29.

An area of about 35 ha (4%) is highly suitable (Class S1) for growing chrysanthemum and are distributed in the central and western part of the microwatershed. Maximum area of about 511 ha (64%) is moderately suitable (Class S2) and occur in the major part of the microwatershed. They have minor limitations of gravelliness, rooting depth, calcareousness and texture. An area of about 113 ha (14%) is marginally suitable (Class S3) for growing chrysanthemum and are distributed in the eastern, southern and western part of the microwatershed with moderate limitations of gravelliness, calcareousness and rooting depth. An area of about 135 ha (17%) is currently not suitable (Class N1) for growing chrysanthemum and distributed in the eastern and central part of the microwatershed with severe limitation of nutrient availability.




Fig. 7.29 Land Suitability map of Chrysanthemum

7. 30 Land Suitability for Jasmine (*Jasminum sp.*)

Jasmine is one of the most important flower crop grown in an area of 803 ha in almost all the districts of the State. The crop requirements (Table 7.31) for growing jasmine were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing jasmine was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.30.

An area of about 35 ha (4%) is highly suitable (Class S1) for growing jasmine and are distributed in the southern and eastern part of the microwatershed. An area of about 221 ha (28%) is moderately suitable (Class S2) and occur in the southern, western and central part of the microwatershed. They have minor limitations of gravelliness and rooting depth. Maximum area of about 403 ha (50%) is marginally suitable (Class S3) for growing jasmine and are distributed in the major part of the microwatershed with moderate limitations of gravelliness, texture, calcareousness and rooting depth. An area of about 135 ha (17%) is currently not suitable (Class N1) for growing chrysanthemum and distributed in the eastern and central part of the microwatershed with severe limitation of nutrient availability.




Fig. 7.30 Land Suitability map of Jasmine

7. 31 Land Suitability for Crossandra (*Crossandra infundibuliformis*)

Crossandra is one of the most important flower crop grown in almost all the districts of the State. The crop requirements (Table 7.32) for growing crossandra were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing crossandra was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.31.

An area of about 35 ha (4%) is highly suitable (Class S1) for growing crossandra and are distributed in the western and southern part of the microwatershed. An area of about 221 ha (28%) is moderately suitable (Class S2) and occur in the southern, central and western part of the microwatershed. They have minor limitations of gravelliness and rooting depth. Maximum area of about 403 ha (50%) is marginally suitable (Class S3) for growing crossandra and are distributed in the major part of the microwatershed with moderate limitations of gravelliness, rooting depth, calcareousness and texture. An area of about 135 ha (17%) is currently not suitable (Class N1) for growing crossandra and distributed in the eastern and central part of the microwatershed with severe limitation of nutrient availability.




Fig. 7.31 Land Suitability map of Crossandra

Table 7.1 Soil-Site Characteristics of Haligeri-3 Microwatershed

| Soil Map Units | Climate (P) (mm) | Growing period (Days) | Drainage Class | Soil depth (cm) | Soil texture | | Gravelliness | | AWC (mm/m) | Slope (%) | Erosion | pH | EC (dSm ⁻¹) | ESP | CEC [Cmol (p ⁺)kg ⁻¹] | BS (%) |
|----------------|------------------|-----------------------|----------------|-----------------|--------------|-------------|--------------|-------------|------------|-----------|----------|------|-------------------------|------|---|--------|
| | | | | | Surf-ace | Sub-surface | Surf-ace | Sub-surface | | | | | | | | |
| CSRiB2 | 662 | <90 | WD | 25-50 | sc | scl | - | <15 | 51-100 | 1-3 | moderate | - | - | - | - | - |
| HRViA1g1 | 662 | <90 | WD | 25-50 | sc | gsc | 15-35 | >35 | <50 | 0-1 | slight | 6.05 | 0.21 | 0.73 | 11.24 | 100 |
| TDHhB2g2 | 662 | <90 | WD | 50-75 | scl | sc-c | 35-60 | <15 | 101-150 | 1-3 | moderate | 9.19 | 0.18 | 5.82 | 3.57 | 100 |
| TDHiB1 | 662 | <90 | WD | 50-75 | sc | sc-c | - | <15 | 101-150 | 1-3 | slight | 9.19 | 0.18 | 5.82 | 3.57 | 100 |
| LKRcB2g1 | 662 | <90 | WD | 50-75 | sl | gsc | 15-35 | 40-60 | 51-100 | 1-3 | moderate | 8.18 | 0.30 | 4.51 | 12.19 | 100 |
| MKHiB2g1 | 662 | <90 | WD | 50-75 | sc | gsc | 15-35 | >35 | 51-100 | 1-3 | moderate | 7.38 | 0.09 | 1.49 | 14.84 | 93 |
| BDGiB1g1 | 662 | <90 | WD | 75-100 | sc | gc | 15-35 | 35-60 | <50 | 1-3 | slight | 6.24 | 0.06 | 0.35 | 3.76 | 52.56 |
| HDHcB1g1 | 662 | <90 | WD | 75-100 | sl | gsc-gc | 15-35 | >35 | 51-100 | 1-3 | slight | 6.54 | 0.07 | 7.11 | 5.84 | 84.7 |
| HDHhB2g1 | 662 | <90 | WD | 75-100 | scl | gsc-gc | 15-35 | >35 | 51-100 | 1-3 | moderate | 6.54 | 0.07 | 7.11 | 5.84 | 84.7 |
| HDHiB1 | 662 | <90 | WD | 75-100 | sc | gsc-gc | - | >35 | 51-100 | 1-3 | slight | 6.54 | 0.07 | 7.11 | 5.84 | 84.7 |
| HDHiB2g1 | 662 | <90 | WD | 75-100 | sc | gsc-gc | 15-35 | >35 | 51-100 | 1-3 | moderate | 6.54 | 0.07 | 7.11 | 5.84 | 84.7 |
| BSRiB1g2 | 662 | <90 | WD | 75-100 | sc | gsc | 35-60 | 15-35 | 51-100 | 1-3 | moderate | 6.59 | 0.12 | 6.00 | 8.80 | 77.55 |
| BPRcA1 | 662 | <90 | WD | 101-150 | sl | gsc-gc | - | >35 | 51-100 | 0-1 | slight | 6.64 | 0.03 | 0.51 | 5.45 | 63.48 |
| BPRhB2g1 | 662 | <90 | WD | 101-150 | scl | gsc-gc | 15-35 | >35 | 51-100 | 1-3 | moderate | 6.64 | 0.03 | 0.51 | 5.45 | 63.48 |
| BPRhB2g2 | 662 | <90 | WD | 101-150 | scl | gsc-gc | 35-60 | >35 | 51-100 | 1-3 | moderate | 6.64 | 0.03 | 0.51 | 5.45 | 63.48 |
| BPRiB2 | 662 | <90 | WD | 101-150 | sc | gsc-gc | - | >35 | 51-100 | 1-3 | moderate | 6.64 | 0.03 | 0.51 | 5.45 | 63.48 |
| BPRmB2 | 662 | <90 | WD | 101-150 | c | gsc-gc | - | >35 | 51-100 | 1-3 | moderate | 6.64 | 0.03 | 0.51 | 5.45 | 63.48 |
| NGPhA1 | 662 | <90 | WD | 100-150 | scl | gsc | - | >35 | 51-100 | 0-1 | slight | 6.67 | 0.09 | 0.46 | 7.10 | 82.70 |
| NGPiB1 | 662 | <90 | WD | 100-150 | sc | gsc | - | >35 | 51-100 | 1-3 | slight | 6.67 | 0.09 | 0.46 | 7.10 | 82.70 |
| GDPcB2 | 662 | <90 | WD | 100-150 | sl | gsc-gc | - | 30-60 | 51-100 | 1-3 | moderate | 7.88 | 0.10 | 2.87 | 7.8 | 97 |
| GDPiB2 | 662 | <90 | WD | 100-150 | sc | gsc-gc | - | 30-60 | 51-100 | 1-3 | moderate | 7.88 | 0.10 | 2.87 | 7.8 | 97 |
| MTLmB2g1 | 662 | <90 | WD | 25-50 | c | gc | 15-35 | 15-35 | 51-100 | 1-3 | moderate | 8.27 | 0.20 | 0.69 | 36.64 | - |

| Soil Map Units | Climate (P) (mm) | Growing period (Days) | Drainage Class | Soil depth (cm) | Soil texture | | Gravelliness | | AWC (mm/m) | Slope (%) | Erosion | pH | EC (dSm ⁻¹) | ESP | CEC [Cmol (p ⁺)kg ⁻¹] | BS (%) |
|----------------|------------------|-----------------------|----------------|-----------------|--------------|-------------|--------------|-------------|------------|-----------|----------|------|-------------------------|-------|---|--------|
| | | | | | Surf-ace | Sub-surface | Surf-ace | Sub-surface | | | | | | | | |
| RNKmB2 | 662 | <90 | MWD | 50-75 | c | c | - | <15 | 51-100 | 1-3 | moderate | 8.86 | 0.48 | 16.94 | 37.0 | - |
| RNKmB2g1 | 662 | <90 | MWD | 50-75 | c | c | 15-35 | <15 | 51-100 | 1-3 | moderate | 8.86 | 0.48 | 16.94 | 37.0 | - |
| DRLmB2 | 662 | <90 | MWD | 75-100 | c | c | - | <15 | 151-200 | 1-3 | moderate | 8.78 | 0.42 | 5.62 | 49.70 | 100 |
| DRLmB2g1 | 662 | <90 | MWD | 75-100 | c | c | 15-35 | <15 | 151-200 | 1-3 | moderate | 8.78 | 0.42 | 5.62 | 49.70 | 100 |
| NSPmB2 | 662 | <90 | MWD | 75-100 | c | c | - | - | 101-150 | 1-3 | moderate | 9.16 | 0.61 | 8.60 | 51.09 | - |
| KVRmA1 | 662 | <90 | MWD | 100-150 | c | c | - | - | >200 | 0-1 | slight | 8.4 | 0.26 | 0.60 | 43.25 | - |
| KVRmB1 | 662 | <90 | MWD | 100-150 | c | c | - | - | >200 | 1-3 | slight | 8.4 | 0.26 | 0.60 | 43.25 | - |
| LGDmB1 | 662 | <90 | WD | 100-150 | c | c | - | <15 | 151-200 | 1-3 | slight | 8.03 | 1.85 | 1.66 | 42.18 | 100.00 |
| AWDmB2 | 662 | <90 | MWD | >150 | c | c | - | <15 | >200 | 1-3 | moderate | 8.10 | 0.37 | 1.22 | 51.30 | 100 |

Table 7.2 Land suitability criteria for Sorghum

| Land use requirement | | | Rating | | | |
|------------------------------|---|---------------|---------------------------|--------------------------|--------------------------|-------------------|
| Soil –site characteristics | | Unit | Highly suitable (S1) | Moderately suitable (S2) | Marginally suitable (S3) | Not suitable (N1) |
| Climatic regime 1 | Mean temperature in growing season | °C | 26–30 | 30–34; 24–26 | 34–40; 20–24 | >40; <20 |
| | Mean max. temp. in growing season | °C | | | | |
| | Mean min. temp. in growing season | °C | | | | |
| | Mean RH in growing season | % | | | | |
| | Total rainfall | mm | | | | |
| | Rainfall in growing season | mm | | | | |
| Land quality | Soil-site characteristics | | | | | |
| Moisture availability | Length of growing period for short duration | Days | | | | |
| | Length of growing period for long duration | | | | | |
| | AWC | mm/m | | | | |
| Oxygen availability to roots | Soil drainage | Class | Well drained | Moderately well drained | Poorly drained | V.poorly drained |
| | Water logging in growing season | Days | | | | |
| Nutrient availability | Texture | Class | sc, c (red), c (black) | scl, cl | ls, sl | - |
| | pH | 1:2.5 | 5.5-7.8 | 5.0-5.5 7.8-9.0 | >9.0 | - |
| | CEC | C mol (p+)/Kg | | | | |
| | BS | % | | | | |
| | CaCO ₃ in root zone | % | | <5 | 5-10 | 10-15 |
| | OC | % | | | | |
| Rooting conditions | Effective soil depth | cm | >75 | 50-75 | 25-50 | <25 |
| | Stoniness | % | | | | |
| | Coarse fragments | Vol % | <15 | 15-35 | 35-60 | 60-80 |
| Soil toxicity | Salinity (EC saturation extract) | dS/m | <2 | 2-4 | 4-8 | >8 |
| | Sodicity (ESP) | % | 5-10 | 10-15 | >15 | |
| Erosion hazard | Slope | % | 0-3 | 3-5 | 5-10 | >10 |

Table 7.3 Land suitability criteria for Maize

| Land use requirement | | | Rating | | | |
|------------------------------|---|---------------|----------------------|--------------------------|--------------------------|---------------------|
| Soil –site characteristics | | Unit | Highly suitable (S1) | Moderately suitable (S2) | Marginally suitable (S3) | Not suitable (N1) |
| Climatic regime | Mean temperature in growing season | °C | 30-34 | 35-38 26-30 | 38-40 26-20 | |
| | Mean max. temp. in growing season | °C | | | | |
| | Mean min. tempt. in growing season | °C | | | | |
| | Mean RH in growing season | % | | | | |
| | Total rainfall | mm | | | | |
| | Rainfall in growing season | mm | | | | |
| Land quality | Soil-site characteristic | | | | | |
| Moisture availability | Length of growing period for short duration | Days | | | | |
| | Length of growing period for long duration | | | | | |
| | AWC | mm/m | | | | |
| Oxygen availability to roots | Soil drainage | Class | Well drained | Moderately well drained | Poorly drained | Very poorly drained |
| | Water logging in growing season | Days | | | | |
| Nutrient availability | Texture | Class | scl, cl, sc | c (red), c (black) | ls, sl | - |
| | pH | 1:2.5 | 5.5-7.8 | 5.0-5.5 7.8-9.0 | >9.0 | - |
| | CEC | C mol (p+)/Kg | | | | |
| | BS | % | | | | |
| | CaCO3 in root zone | % | | <5 | 5-10 | >10 |
| | OC | % | | | | |
| Rooting conditions | Effective soil depth | cm | >75 | 50-75 | 25-50 | <25 |
| | Stoniness | % | | | | |
| | Coarse fragments | Vol % | <15 | 15-35 | 35-60 | 60-80 |
| Soil toxicity | Salinity (EC saturation extract) | ds/m | <2 | 2-4 | 4-8 | >8 |
| | Sodicity (ESP) | % | 5-10 | 10-15 | >15 | - |
| Erosion hazard | Slope | % | 0-3 | 3-5 | 5-10 | >10 |

Table 7.4 Land suitability criteria for Bajra

| Land use requirement | | | Rating | | | |
|------------------------------|---|----------------|------------------------|--------------------------|--------------------------|---------------------|
| Soil –site characteristics | | Unit | Highly suitable (S1) | Moderately suitable (S2) | Marginally suitable (S3) | Not suitable (N1) |
| Climatic regime | Mean temperature in growing season | °C | 28-32 | 33-38 24-27 | 39-40 20-23 | <20 |
| | Mean max. temp. in growing season | °C | | | | |
| | Mean min. tempt. in growing season | °C | | | | |
| | Mean RH in growing season | % | | | | |
| | Total rainfall | mm | 500-750 | 400-500 | 200-400 | <200 |
| | Rainfall in growing season | mm | | | | |
| Land quality | Soil-site characteristic | | | | | |
| Moisture availability | Length of growing period for short duration | Days | | | | |
| | Length of growing period for long duration | | | | | |
| | AWC | mm/m | | | | |
| Oxygen availability to roots | Soil drainage | Class | Well drained | Moderately well drained | Poorly drained | Very poorly drained |
| | Water logging in growing season | Days | | | | |
| Nutrient availability | Texture | Class | Sl, scl, cl,sc,c (red) | C (black) | ls | - |
| | pH | 1:2.5 | 6.0-7.8 | 5.0-5.5 7.8-9.0 | 5.5-6.0 >9.0 | |
| | CEC | C mol (p+)/ Kg | | | | |
| | BS | % | | | | |
| | CaCO3 in root zone | % | | <5 | 5-10 | >10 |
| | OC | % | | | | |
| Rooting conditions | Effective soil depth | cm | >75 | 50-75 | 25-50 | <25 |
| | Stoniness | % | | | | |
| | Coarse fragments | Vol % | 15-35 | 35-60 | >60 | |
| Soil toxicity | Salinity (EC saturation extract) | dS/m | <2 | 2-4 | 4-8 | >8 |
| | Sodicity (ESP) | % | 5-10 | 10-15 | >15 | |
| Erosion hazard | Slope | % | 1-3 | 3-5 | 5-10 | >10 |

Table 7.5 Land suitability criteria for Red gram

| Land use requirement | | | Rating | | | |
|------------------------------|---|----------------|---|--|---|---------------------------|
| Soil –site characteristics | | Unit | Highly suitable (S1) | Moderately suitable (S2) | Marginally suitable (S3) | Not suitable (N1) |
| Climatic regime | Mean temperature in growing season | °C | 30-35(G) 20-25(AV) 15-18 (F&PS) 35-40(M) | 25-30(G) 20-25 (AV) 12-15 (F&PS) 30-35(M) | 20-25(G) 15-20(AV) 10-12 (F&PS) 25-30(M) | < 20 <15 <10 <25 |
| | Mean max. temp. in growing season | °C | | | | |
| | Mean min. temp. in growing season | °C | | | | |
| | Mean RH in growing season | % | | | | |
| | Total rainfall | mm | | | | |
| | Rainfall in growing season | mm | | | | |
| Land quality | Soil-site characteristic | | | | | |
| Moisture availability | Length of growing period for short duration | Days | | | | |
| | Length of growing period for long duration | | | | | |
| | AWC | mm/m | | | | |
| Oxygen availability to roots | Soil drainage | Class | Well drained | Mod. Well drained | Poorly drained | Very Poorly drained |
| | Water logging in growing season | Days | | | | |
| Nutrient availability | Texture | Class | sc, c (red) | c (black),sl, scl, cl | ls | - |
| | pH | 1:2.5 | 6.0-7.8 | 5.5-6.0 7.8-9.0 | 5.0-5.5 >9.0 | - |
| | CEC | C mol (p+)/ Kg | | | | |
| | BS | % | | | | |
| | CaCO ₃ in root zone | % | | <5 | 5-10 | >10 |
| | OC | % | | | | |
| Rooting conditions | Effective soil depth | cm | >100 | 75-100 | 50-75 | <50 |
| | Stoniness | % | | | | |
| | Coarse fragments | Vol % | <15 | 15-35 | 35-50 | 60-80 |
| Soil toxicity | Salinity (EC saturation extract) | dS/m | <1.0 | 1.0-2.0 | >2.0 | |
| | Sodicity (ESP) | % | 5-10 | 10-15 | >15 | |
| Erosion hazard | Slope | % | <3 | 3-5 | 5-10 | >10 |

Table 7.6 Land suitability criteria for Bengal gram

| Land use requirement | | | Rating | | | |
|------------------------------|---|----------------------|--------------------------|--------------------------|----------------------|---------------------|
| Soil –site characteristics | Unit | Highly suitable (S1) | Moderately suitable (S2) | Marginally suitable (S3) | Not suitable (N1) | |
| Climatic regime | Mean temperature in growing season | °C | 20–25 | 25–30; 15–20 | 30–35; 10–15 | >35; <10 |
| | Mean max. temp. in growing season | °C | | | | |
| | Mean min. tempt. in growing season | °C | | | | |
| | Mean RH in growing season | % | | | | |
| | Total rainfall | mm | | | | |
| | Rainfall in growing season | mm | | | | |
| Land quality | Soil-site characteristic | | | | | |
| Moisture availability | Length of growing period for short duration | Days | | | | |
| | Length of growing period for long duration | | | | | |
| | AWC | mm/m | | | | |
| Oxygen availability to roots | Soil drainage | Class | Well drained | Mod. Well drained | Poorly drained | Very Poorly drained |
| | Water logging in growing season | Days | | | | |
| Nutrient availability | Texture | Class | C (black) | - | c (red), scl, cl, sc | ls, sl |
| | pH | 1:2.5 | 6.0-7.8 | 5.0-6.0 7.8-9.0 | >9.0 | - |
| | CEC | C mol (p+)/Kg | | | | |
| | BS | % | | | | |
| | CaCO ₃ in root zone | % | | <5 | 5-10 | >10 |
| | OC | % | | | | |
| Rooting conditions | Effective soil depth | cm | >75 | 50-75 | 25-50 | <25 |
| | Stoniness | % | | | | |
| | Coarse fragments | Vol % | <15 | 15-35 | 35-60 | 60-80 |
| Soil toxicity | Salinity (EC saturation extract) | dS/m | <2 | 2-4 | 4-8 | >8 |
| | Sodicity (ESP) | % | 5-10 | 10-15 | >15 | - |
| Erosion hazard | Slope | % | <3 | 3-5 | 5-10 | >10 |

Table 7.7 Land suitability criteria for Groundnut

| Land use requirement | | | Rating | | | |
|------------------------------|---|---------------|----------------------|--------------------------|--------------------------|---------------------|
| Soil –site characteristics | | Unit | Highly suitable (S1) | Moderately suitable (S2) | Marginally suitable (S3) | Not suitable (N1) |
| Climatic regime | Mean temperature in growing season | °C | 24–33 | 22–24; 33–35 | 20–22; 35–40 | <20; >40 |
| | Mean max. temp. in growing season | °C | | | | |
| | Mean min. temp. in growing season | °C | | | | |
| | Mean RH in growing season | % | | | | |
| | Total rainfall | mm | | | | |
| | Rainfall in growing season | mm | | | | |
| Land quality | Soil-site characteristic | | | | | |
| Moisture availability | Length of growing period for short duration | Days | | | | |
| | Length of growing period for long duration | | | | | |
| | AWC | mm/m | | | | |
| Oxygen availability to roots | Soil drainage | Class | Well drained | Mod. Well drained | Poorly drained | Very Poorly drained |
| | Water logging in growing season | Days | | | | |
| Nutrient availability | Texture | Class | scl | sl,cl, sc | c (red), c (black), ls | - |
| | pH | 1:2.5 | 6.0-7.8 | 5.5-6.0 7.8-8.4 | 5.0-5.5 8.4-9.0 | >9.0 |
| | CEC | C mol (p+)/Kg | | | | |
| | BS | % | | | | |
| | CaCO ₃ in root zone | % | | <5 | 5-10 | >10 |
| | OC | % | | | | |
| Rooting conditions | Effective soil depth | cm | >75 | 50-75 | 25-50 | <25 |
| | Stoniness | % | | | | |
| | Coarse fragments | Vol % | <35 | 35-60 | >60 | |
| Soil toxicity | Salinity (EC saturation extract) | dS/m | <2 | 2-4 | 4-8 | >8 |
| | Sodicity (ESP) | % | <5 | 5-10 | 10-15 | >15 |
| Erosion hazard | Slope | % | <3 | 3-5 | 5-10 | >10 |

Table 7.8 Land suitability criteria for Sunflower

| Land use requirement | | | Rating | | | |
|------------------------------|---|---------------|---------------------------|--------------------------|--------------------------|------------------------|
| Soil –site characteristics | | Unit | Highly suitable (S1) | Moderately suitable (S2) | Marginally suitable (S3) | Not suitable (N1) |
| Climatic regime | Mean temperature in growing season | °C | 24–30 | 30–34; 20–24 | 34–38; 16–20 | >38; <16 |
| | Mean max. temp. in growing season | °C | | | | |
| | Mean min. temp. in growing season | °C | | | | |
| | Mean RH in growing season | % | | | | |
| | Total rainfall | mm | | | | |
| | Rainfall in growing season | mm | | | | |
| Land quality | Soil-site characteristic | | | | | |
| Moisture availability | Length of growing period for short duration | Days | | | | |
| | Length of growing period for long duration | | | | | |
| | AWC | mm/m | | | | |
| Oxygen availability to roots | Soil drainage | Class | Well drained | mod. Well drained | - | Poorly to very drained |
| | Water logging in growing season | Days | | | | |
| Nutrient availability | Texture | Class | cl, sc,c (red), c (black) | scl | ls, sl | - |
| | pH | 1:2.5 | 6.5-7.8 | 7.8-8.4 5.5-6.5 | 8.4-9.0; 5.0-5.5 | >9.0 |
| | CEC | C mol (p+)/Kg | | | | |
| | BS | % | | | | |
| | CaCO ₃ in root zone | % | | <5 | 5-10 | >10 |
| | OC | % | | | | |
| Rooting conditions | Effective soil depth | cm | >100 | 75-100 | 50-75 | <50 |
| | Stoniness | % | | | | |
| | Coarse fragments | Vol % | <15 | 15-35 | 35-60 | 60-80 |
| Soil toxicity | Salinity (EC saturation extract) | dS/m | <2 | 2-4 | 4-8 | >8 |
| | Sodicity (ESP) | % | <5 | 5-10 | 10-15 | >15 |
| Erosion hazard | Slope | % | <3 | 3-5 | 5-10 | >10 |

Table 7.9 Land suitability criteria for Cotton

| Land use requirement | | | Rating | | | |
|------------------------------|---|--------------|-------------------------|--|--------------------------|---------------------------------|
| Soil –site characteristics | | Unit | Highly suitable (S1) | Moderately suitable (S2) | Marginally suitable (S3) | Not suitable (N1) |
| Climatic regime | Mean temperature in growing season | °C | 22-32 | >32 | <19 | - |
| | Mean max. temp. in growing season | °C | | | | |
| | Mean min. temp. in growing season | °C | | | | |
| | Mean RH in growing season | % | | | | |
| | Total rainfall | mm | | | | |
| | Rainfall in growing season | mm | | | | |
| Land quality | Soil-site characteristic | | | | | |
| Moisture availability | Length of growing period for short duration | Days | | | | |
| | Length of growing period for long duration | | | | | |
| | AWC | mm/m | | | | |
| Oxygen availability to roots | Soil drainage | Class | Well to moderately well | Poorly drained/Some what excessively drained | - | very poorly/excessively drained |
| | Water logging in growing season | Days | | | | |
| Nutrient availability | Texture | Class | sc, c (red,black) | cl | scl | ls, sl |
| | pH | 1:2.5 | 6.5-7.8 | 7.8-8.4 | 5.5-6.5 8.4->9.0 | <5.5 |
| | CEC | C mol (p+)Kg | | | | |
| | BS | % | | | | |
| | CaCO3 in root zone | % | | <5 | 5-10 | >10 |
| | OC | % | | | | |
| Rooting conditions | Effective soil depth | cm | >100 | 50-100 | 25-50 | <25 |
| | Stoniness | % | | | | |
| | Coarse fragments | Vol % | <15 | 15-35 | 35-60 | 60-80 |
| Soil toxicity | Salinity (EC saturation extract) | dS/m | <2 | 2-4 | 4-8 | >8 |
| | Sodicity (ESP) | % | 5-10 | 10-15 | >15 | |
| Erosion hazard | Slope | % | <3 | 3-5 | - | >5 |

Table 7.10 Land suitability criteria for Chili

| Land use requirement | | | Rating | | | |
|------------------------------|---|----------------|----------------------|--------------------------|--------------------------|---------------------|
| Soil –site characteristics | | Unit | Highly suitable (S1) | Moderately suitable (S2) | Marginally suitable (S3) | Not suitable (N1) |
| Climatic regime | Mean temperature in growing season | °C | 25-32 | 33-35 20-25 | 35-38 <20 | >38 |
| | Mean max. temp. in growing season | °C | | | | |
| | Mean min. tempt. in growing season | °C | | | | |
| | Mean RH in growing season | % | | | | |
| | Total rainfall | mm | | | | |
| | Rainfall in growing season | mm | | | | |
| Land quality | Soil-site characteristic | | | | | |
| Moisture availability | Length of growing period for short duration | Days | | | | |
| | Length of growing period for long duration | | | | | |
| | AWC | mm/m | | | | |
| Oxygen availability to roots | Soil drainage | Class | Well drained | Moderately well drained | Poorly drained | Very poorly drained |
| | Water logging in growing season | Days | | | | |
| Nutrient availability | Texture | Class | scl, cl, sc | c (black), sl | ls | - |
| | pH | 1:2.5 | 6.0-7.3 | 5.0-6.0 7.3-8.4 | 8.4-9.0 | >9.0 |
| | CEC | C mol (p+)/ Kg | | | | |
| | BS | % | | | | |
| | CaCO3 in root zone | % | | <5 | 5-10 | >10 |
| | OC | % | | | | |
| Rooting conditions | Effective soil depth | cm | >75 | 50-75 | 25-50 | <25 |
| | Stoniness | % | | | | |
| | Coarse fragments | Vol % | <15 | 15-35 | 35-60 | 60-80 |
| Soil toxicity | Salinity (EC saturation extract) | dS/m | <2 | 2-4 | 4-8 | >8 |
| | Sodicity (ESP) | % | <5 | 5-10 | 10-15 | >15 |
| Erosion hazard | Slope | % | <3 | 3-5 | 5-10 | >10 |

Table 7.11 Land suitability criteria for Tomato

| Land use requirement | | Rating | | | | |
|------------------------------|---|---------------|--------------------------|--------------------------|--------------------------|-------------------|
| Soil –site characteristics | | Unit | Highly suitable (S1) | Moderately suitable (S2) | Marginally suitable (S3) | Not suitable (N1) |
| Climatic regime | Mean temperature in growing season | °C | 25-28 | 29-32 20-24 | 15-19 33-36 | <15 >36 |
| | Mean max. temp. in growing season | °C | | | | |
| | Mean min. tempt. in growing season | °C | | | | |
| | Mean RH in growing season | % | | | | |
| | Total rainfall | mm | | | | |
| | Rainfall in growing season | mm | | | | |
| Land quality | Soil-site characteristic | | | | | |
| Moisture availability | Length of growing period for short duration | Days | | | | |
| | Length of growing period for long duration | | | | | |
| | AWC | mm/m | | | | |
| Oxygen availability to roots | Soil drainage | Class | Well drained | Moderately well drained | Poorly drained | V.poorly drained |
| | Water logging in growing season | Days | | | | |
| Nutrient availability | Texture | Class | sl, scl, cl, sc, c (red) | - | ls, c(black) | - |
| | pH | 1:2.5 | 6.0-7.3 | 5.0-6.0 7.3-8.4 | 8.4-9.0 | >9.0 |
| | CEC | C mol (p+)/Kg | | | | |
| | BS | % | | | | |
| | CaCO3 in root zone | % | | <5 | 5-10 | >10 |
| | OC | % | | | | |
| Rooting conditions | Effective soil depth | cm | >75 | 50-75 | 25-50 | <25 |
| | Stoniness | % | | | | |
| | Coarse fragments | Vol % | <15 | 15-35 | 35-60 | 60-80 |
| Soil toxicity | Salinity (EC saturation extract) | dS/m | <2.0 | 2-4 | 4-8 | >8.0 |
| | Sodicity (ESP) | % | <5 | 5-10 | 10-15 | >15 |
| Erosion hazard | Slope | % | <3 | 3-5 | 5-10 | >10 |

Table 7.12 Land suitability criteria for Brinjal

| Land use requirement | | | Rating | | | |
|------------------------------|---|---------------|-------------------------|--------------------------|--------------------------|-------------------|
| Soil –site characteristics | | Unit | Highly suitable (S1) | Moderately suitable (S2) | Marginally suitable (S3) | Not suitable (N1) |
| Climatic regime | Mean temperature in growing season | °C | Well drained | Moderately well drained | Poorly drained | V. Poorly drained |
| | Mean max. temp. in growing season | °C | | | | |
| | Mean min. tempt. in growing season | °C | | | | |
| | Mean RH in growing season | % | | | | |
| | Total rainfall | mm | | | | |
| | Rainfall in growing season | mm | | | | |
| Land quality | Soil-site characteristic | | | | | |
| Moisture availability | Length of growing period for short duration | Days | | | | |
| | Length of growing period for long duration | | | | | |
| | AWC | mm/m | | | | |
| Oxygen availability to roots | Soil drainage | Class | | | | |
| | Water logging in growing season | Days | | | | |
| Nutrient availability | Texture | Class | sl, scl, cl, sc c (red) | - | ls, c (black) | - |
| | pH | 1:2.5 | 6.0-7.3 | 7.3-8.4 5.0-6.0 | 8.4-9.0 | >9.0 |
| | CEC | C mol (p+)/Kg | | | | |
| | BS | % | | | | |
| | CaCO ₃ in root zone | % | | <5 | 5-10 | >10 |
| | OC | % | | | | |
| Rooting conditions | Effective soil depth | cm | >75 | 50-75 | 25-50 | <25 |
| | Stoniness | % | | | | |
| | Coarse fragments | Vol % | <15 | 15-35 | 35-60 | >60 |
| Soil toxicity | Salinity (EC saturation extract) | ds/m | <2.0 | 2-4 | 4-8 | >8.0 |
| | Sodicity (ESP) | % | <5 | 5-10 | 10-15 | >15 |
| Erosion hazard | Slope | % | <3 | 3-5 | 5-10 | >10 |

Table 7.13 Land suitability criteria for Onion

| Land use requirement | | Rating | | | | |
|------------------------------|---|---------------|----------------------|--------------------------|--------------------------|----------------------------|
| Soil –site characteristics | | Unit | Highly suitable (S1) | Moderately suitable (S2) | Marginally suitable (S3) | Not suitable (N1) |
| Climatic regime | Mean temperature in growing season | °C | 20-30 | 30-35 | 35-40 | >40 |
| | Mean max. temp. in growing season | °C | | | | |
| | Mean min. tempt. in growing season | °C | | | | |
| | Mean RH in growing season | % | | | | |
| | Total rainfall | mm | | | | |
| | Rainfall in growing season | mm | | | | |
| Land quality | Soil-site characteristic | | | | | |
| Moisture availability | Length of growing period for short duration | Days | | | | |
| | Length of growing period for long duration | | | | | |
| | AWC | mm/m | | | | |
| Oxygen availability to roots | Soil drainage | Class | Well drained | Moderately /imperfectly | - | Poorly to V poorly drained |
| | Water logging in growing season | Days | | | | |
| Nutrient availability | Texture | Class | sl,scl,cl,sc,c (red) | - | c (Black),ls | - |
| | pH | 1:2.5 | 6.0-7.3 | 5.0-6.0 7.3-7.8 | 7.8-8.4 | >8.4 |
| | CEC | C mol (p+)/Kg | | | | |
| | BS | % | | | | |
| | CaCO3 in root zone | % | | <5 | 5-10 | >10 |
| | OC | % | | | | |
| Rooting conditions | Effective soil depth | cm | >75 | 50-75 | 25-50 | <25 |
| | Stoniness | % | | | | |
| | Coarse fragments | Vol % | <15 | 15-35 | 35-60 | 60-80 |
| Soil toxicity | Salinity (EC saturation extract) | ds/m | <1.0 | 1.0-2.0 | 2.0-4.0 | <4 |
| | Sodicity (ESP) | % | <5 | 5-10 | 10-15 | >15 |
| Erosion hazard | Slope | % | <3 | 3-5 | 5-10 | >10 |

Table 7.14 Land suitability criteria for Bhendi

| Land use requirement | | | Rating | | | |
|------------------------------|---|---------------|----------------------|--------------------------|--------------------------|-------------------------------|
| Soil –site characteristics | | Unit | Highly suitable (S1) | Moderately suitable (S2) | Marginally suitable (S3) | Not suitable (N1) |
| Climatic regime | Mean temperature in growing season | °C | 25-28 | 29-32 20-24 | 15-19 33-36 | <15 >36 |
| | Mean max. temp. in growing season | °C | | | | |
| | Mean min. tempt. in growing season | °C | | | | |
| | Mean RH in growing season | % | | | | |
| | Total rainfall | mm | | | | |
| | Rainfall in growing season | mm | | | | |
| Land quality | Soil-site characteristic | | | | | |
| Moisture availability | Length of growing period for short duration | Days | | | | |
| | Length of growing period for long duration | | | | | |
| | AWC | mm/m | | | | |
| Oxygen availability to roots | Soil drainage | Class | Well drained | Moderately well drained | Imperfectly drained | Poorly to very poorly drained |
| | Water logging in growing season | Days | | | | |
| Nutrient availability | Texture | Class | scl, cl,sc, c (red) | c (black) | ls | - |
| | pH | 1:2.5 | 6.0-7.3 | 5.0-6.0 7.3-8.4 | 8.4-9.0 | >9.0 |
| | CEC | C mol (p+)/Kg | | | | |
| | BS | % | | | | |
| | CaCO ₃ in root zone | % | | <5 | 5-10 | >10 |
| | OC | % | | | | |
| Rooting conditions | Effective soil depth | cm | >75 | 50-75 | 25-50 | <25 |
| | Stoniness | % | | | | |
| | Coarse fragments | Vol % | <15 | 15-35 | 35-60 | 60-80 |
| Soil toxicity | Salinity (EC saturation extract) | ds/m | <2.0 | 2-4 | 4-8 | >8.0 |
| | Sodicity (ESP) | % | <5 | 5-10 | 10-15 | >15 |
| Erosion hazard | Slope | % | <3 | 3-5 | 5-10 | >10 |

Table 7.15 Land suitability criteria for Drumstick

| Land use requirement | | | Rating | | | |
|------------------------------|---|---------------|----------------------|--------------------------|--------------------------|-------------------|
| Soil –site characteristics | | Unit | Highly suitable (S1) | Moderately suitable (S2) | Marginally suitable (S3) | Not suitable (N1) |
| Climatic regime | Mean temperature in growing season | °C | | | | |
| | Mean max. temp. in growing season | °C | | | | |
| | Mean min. temp. in growing season | °C | | | | |
| | Mean RH in growing season | % | | | | |
| | Total rainfall | mm | | | | |
| | Rainfall in growing season | mm | | | | |
| Land quality | Soil-site characteristic | | | | | |
| Moisture availability | Length of growing period for short duration | Days | | | | |
| | Length of growing period for long duration | | | | | |
| | AWC | mm/m | | | | |
| Oxygen availability to roots | Soil drainage | Class | Well drained | Moderately well drained | Poorly drained | V.Poorly drained |
| | Water logging in growing season | Days | | | | |
| Nutrient availability | Texture | Class | sc, scl, cl, c (red) | sl, c (black) | ls | s |
| | pH | 1:2.5 | 6.0-7.3 | 5.0-5.5 7.3-7.8 | 5.5-6.0 7.8-8.4 | >8.4 |
| | CEC | C mol (p+)/Kg | | | | |
| | BS | % | | | | |
| | CaCO ₃ in root zone | % | | <5 | 5-10 | >10 |
| | OC | % | | | | |
| Rooting conditions | Effective soil depth | cm | >100 | 75-100 | 50-75 | <50 |
| | Stoniness | % | | | | |
| | Coarse fragments | Vol % | <35 | 35-60 | 60-80 | >80 |
| Soil toxicity | Salinity (EC saturation extract) | dS/m | | | | |
| | Sodicity (ESP) | % | <5 | 5-10 | 10-15 | >15 |
| Erosion hazard | Slope | % | <3 | 3-10 | - | >10 |

Table 7.16 Land suitability criteria for Mulberry

| Land use requirement | | | Rating | | | |
|------------------------------|---|---------------|----------------------|--------------------------|--------------------------|-------------------|
| Soil –site characteristics | | Unit | Highly suitable (S1) | Moderately suitable (S2) | Marginally suitable (S3) | Not suitable (N1) |
| Climatic regime | Mean temperature in growing season | °C | 24–28 | 22–24; 28–32 | 32–38; 22–18 | >38; <18 |
| | Mean max. temp. in growing season | °C | | | | |
| | Mean min. tempt. in growing season | °C | | | | |
| | Mean RH in growing season | % | | | | |
| | Total rainfall | mm | | | | |
| | Rainfall in growing season | mm | | | | |
| Land quality | Soil-site characteristic | | | | | |
| Moisture availability | Length of growing period for short duration | Days | | | | |
| | Length of growing period for long duration | | | | | |
| | AWC | mm/m | | | | |
| Oxygen availability to roots | Soil drainage | Class | Well drained | Moderately well drained | Poorly drained | V. Poorly drained |
| | Water logging in growing season | Days | | | | |
| Nutrient availability | Texture | Class | sc, cl, scl | c (red) | c (black), sl, ls | - |
| | pH | 1:2.5 | 5.5-7.3 | 5.0-5.5 7.8-8.4 | 7.3-8.4 | >8.4 |
| | CEC | C mol (p+)/Kg | | | | |
| | BS | % | | | | |
| | CaCO3 in root zone | % | | <5 | 5-10 | >10 |
| | OC | % | | | | |
| Rooting conditions | Effective soil depth | cm | >100 | 75-100 | 50-75 | <50 |
| | Stoniness | % | | | | |
| | Coarse fragments | Vol % | 0-35 | 35-60 | 60-80 | >80 |
| Soil toxicity | Salinity (EC saturation extract) | dS/m | <2 | 2-4 | 4-8 | >8 |
| | Sodicity (ESP) | % | <5 | 5-10 | 10-15 | >15 |
| Erosion hazard | Slope | % | 0-3 | 3-5 | 5-10 | >10 |

Note: Suitability evaluation only for Mulberry leaf not for Silk worm rearing

Table 7.17 Land suitability criteria for Mango

| Land use requirement | | | Rating | | | |
|------------------------------|---|---------------|----------------------|--------------------------|--------------------------|-------------------|
| Soil –site characteristics | | Unit | Highly suitable (S1) | Moderately suitable (S2) | Marginally suitable (S3) | Not suitable (N1) |
| Climatic regime | Mean temperature in growing season | °C | 28-32 | 24-27 33-35 | 36-40 | 20-24 |
| | Min temp. before flowering | °C | 10-15 | 15-22 | >22 | - |
| | Mean max. temp. in growing season | °C | | | | |
| | Mean min. tempt. in growing season | °C | | | | |
| | Mean RH in growing season | % | | | | |
| | Total rainfall | mm | | | | |
| | Rainfall in growing season | mm | | | | |
| Land quality | Soil-site characteristic | | | | | |
| Moisture availability | Length of growing period for short duration | Days | | | | |
| | Length of growing period for long duration | Days | | | | |
| | AWC | mm/m | | | | |
| Oxygen availability to roots | Soil drainage | Class | Well drained | Moderately well drained | Poorly drained | V.Poorly drained |
| | Water logging in growing season | Days | | | | |
| Nutrient availability | Texture | Class | scl, cl, sc, c (red) | - | ls, sl, c (black) | - |
| | pH | 1:2.5 | 5.5-7.3 | 5.0-5.5 7.3-8.4 | 8.4-9.0 | >9.0 |
| | CEC | C mol (p+)/Kg | | | | |
| | BS | % | | | | |
| | CaCO3 in root zone | % | | <5 | 5-10 | >10 |
| | OC | % | | | | |
| Rooting conditions | Effective soil depth | cm | >150 | 100-150 | 75-100 | <75 |
| | Stoniness | % | | | | |
| | Coarse fragments | Vol % | <15 | 15-35 | 35-60 | 60-80 |
| Soil toxicity | Salinity (EC saturation extract) | dS/m | <2.0 | 2-4 | 4-8 | >8.0 |
| | Sodicity (ESP) | % | <5 | 5-10 | 10-15 | >15 |
| Erosion hazard | Slope | % | <3 | 3-5 | 5-10 | >10 |

Table 7.18 Land suitability criteria for Sapota

| Land use requirement | | | Rating | | | |
|------------------------------|---|---------------|----------------------|--------------------------|--------------------------|------------------------|
| Soil –site characteristics | | Unit | Highly suitable (S1) | Moderately suitable (S2) | Marginally suitable (S3) | Not suitable (N1) |
| Climatic regime | Mean temperature in growing season | °C | 28-32 | 33-36 24-27 | 37-42 20-23 | >42 <18 |
| | Mean max. temp. in growing season | °C | | | | |
| | Mean min. temp. in growing season | °C | | | | |
| | Mean RH in growing season | % | | | | |
| | Total rainfall | mm | | | | |
| | Rainfall in growing season | mm | | | | |
| Land quality | Soil-site characteristic | | | | | |
| Moisture availability | Length of growing period for short duration | Days | | | | |
| | Length of growing period for long duration | | | | | |
| | AWC | mm/m | | | | |
| Oxygen availability to roots | Soil drainage | Class | Well drained | Moderately well drained | - | Poorly to very drained |
| | Water logging in growing season | Days | | | | |
| Nutrient availability | Texture | Class | scl, cl, sc, c (red) | sl | ls, c (black) | - |
| | pH | 1:2.5 | 6.0-7.3 | 5.0-6.0 7.3-8.4 | 8.4-9.0 | >9.0 |
| | CEC | C mol (p+)/Kg | | | | |
| | BS | % | | | | |
| | CaCO3 in root zone | % | | <5 | 5-10 | >10 |
| | OC | % | | | | |
| Rooting conditions | Effective soil depth | cm | >100 | 75-100 | 50-75 | <50 |
| | Stoniness | % | | | | |
| | Coarse fragments | Vol % | <15 | 15-35 | 35-60 | 60-80 |
| Soil toxicity | Salinity (EC saturation extract) | dS/m | <2.0 | 2-4 | 4-8 | >8.0 |
| | Sodicity (ESP) | % | <5 | 5-10 | 10-15 | >15 |
| Erosion hazard | Slope | % | <3 | 3-5 | 5-10 | >10 |

Table 7.19 Land suitability criteria for Pomegranate

| Land use requirement | | | Rating | | | |
|------------------------------|---|----------------|----------------------|--------------------------|--------------------------|-------------------|
| Soil –site characteristics | | Unit | Highly suitable (S1) | Moderately suitable (S2) | Marginally suitable (S3) | Not suitable (N1) |
| Climatic regime | Mean temperature in growing season | °C | 30-34 | 35-38 25-29 | 39-40 15-24 | |
| | Mean max. temp. in growing season | °C | | | | |
| | Mean min. temp. in growing season | °C | | | | |
| | Mean RH in growing season | % | | | | |
| | Total rainfall | mm | | | | |
| | Rainfall in growing season | mm | | | | |
| Land quality | Soil-site characteristic | | | | | |
| Moisture availability | Length of growing period for short duration | Days | | | | |
| | Length of growing period for long duration | | | | | |
| | AWC | mm/m | | | | |
| Oxygen availability to roots | Soil drainage | Class | Well drained | Moderately well drained | Poorly drained | V.Poorly drained |
| | Water logging in growing season | Days | | | | |
| Nutrient availability | Texture | Class | scl,cl, sc, c (red) | c (black),sl | ls | - |
| | pH | 1:2.5 | 5.5-7.8 | 7.8-8.4 | 5.0-5.5 8.4-9.0 | >9.0 |
| | CEC | C mol (p+)/ Kg | | | | |
| | BS | % | | | | |
| | CaCO ₃ in root zone | % | | <5 | 5-10 | >10 |
| | OC | % | | | | |
| Rooting conditions | Effective soil depth | cm | >100 | 75-100 | 50-75 | <50 |
| | Stoniness | % | | | | |
| | Coarse fragments | Vol % | <15 | 15-35 | 35-60 | 60-80 |
| Soil toxicity | Salinity (EC saturation extract) | dS/m | <2.0 | 2-4 | 4-8 | >8.0 |
| | Sodicity (ESP) | % | <5 | 5-10 | 10-15 | >15 |
| Erosion hazard | Slope | % | <3 | 3-5 | 5-10 | >10 |

Table 7.20 Land suitability criteria for Guava

| Land use requirement | | | Rating | | | |
|------------------------------|---|---------------|----------------------|--------------------------|--------------------------|-------------------|
| Soil –site characteristics | | Unit | Highly suitable (S1) | Moderately suitable (S2) | Marginally suitable (S3) | Not suitable (N1) |
| Climatic regime | Mean temperature in growing season | °C | 28-32 | 33-36 24-27 | 37-42 20-23 | |
| | Mean max. temp. in growing season | °C | | | | |
| | Mean min. tempt. in growing season | °C | | | | |
| | Mean RH in growing season | % | | | | |
| | Total rainfall | mm | | | | |
| | Rainfall in growing season | mm | | | | |
| Land quality | Soil-site characteristic | | | | | |
| Moisture availability | Length of growing period for short duration | Days | | | | |
| | Length of growing period for long duration | | | | | |
| | AWC | mm/m | | | | |
| Oxygen availability to roots | Soil drainage | Class | Well drained | Moderately well drained | Poorly drained | V.Poorly drained |
| | Water logging in growing season | Days | | | | |
| Nutrient availability | Texture | Class | scl, cl, sc, c (red) | sl | c (black), ls | - |
| | pH | 1:2.5 | 6.0-7.8 | 5.0-6.0 | 7.8-8.4 | >8.4 |
| | CEC | C mol (p+)/Kg | | | | |
| | BS | % | | | | |
| | CaCO3 in root zone | % | | <5 | 5-10 | >10 |
| | OC | % | | | | |
| Rooting conditions | Effective soil depth | cm | >100 | 75-100 | 50-75 | <50 |
| | Stoniness | % | | | | |
| | Coarse fragments | Vol % | <15 | 15-35 | 35-60 | 60-80 |
| Soil toxicity | Salinity (EC saturation extract) | dS/m | <2.0 | 2-4 | 4-8 | >8.0 |
| | Sodicity (ESP) | % | <5 | 5-10 | 10-15 | >15 |
| Erosion hazard | Slope | % | <3 | 3-5 | 5-10 | >10 |

Table 7.21 Land suitability criteria for Jackfruit

| Land use requirement | | | Rating | | | |
|------------------------------|---|---------------|----------------------|--------------------------|--------------------------|-------------------|
| Soil –site characteristics | | Unit | Highly suitable (S1) | Moderately suitable (S2) | Marginally suitable (S3) | Not suitable (N1) |
| Climatic regime | Mean temperature in growing season | °C | | | | |
| | Mean max. temp. in growing season | °C | | | | |
| | Mean min. temp. in growing season | °C | | | | |
| | Mean RH in growing season | % | | | | |
| | Total rainfall | mm | | | | |
| | Rainfall in growing season | mm | | | | |
| Land quality | Soil-site characteristic | | | | | |
| Moisture availability | Length of growing period for short duration | Days | | | | |
| | Length of growing period for long duration | | | | | |
| | AWC | mm/m | | | | |
| Oxygen availability to roots | Soil drainage | Class | Well drained | Mod. well | Poorly | V. Poorly |
| | Water logging in growing season | Days | | | | |
| Nutrient availability | Texture | Class | scl, cl, sc, c (red) | - | sl, ls, c (black) | - |
| | pH | 1:2.5 | 5.5-7.3 | 5.0-5.5 7.3-7.8 | 7.8-8.4 | >8.4 |
| | CEC | C mol (p+)/Kg | | | | |
| | BS | % | | | | |
| | CaCO3 in root zone | % | | <5 | 5-10 | >10 |
| | OC | % | | | | |
| Rooting conditions | Effective soil depth | cm | >100 | 75-100 | 50-75 | <50 |
| | Stoniness | % | | | | |
| | Coarse fragments | Vol % | <15 | 15-35 | 35-60 | >60 |
| Soil toxicity | Salinity (EC saturation extract) | dS/m | <2.0 | 2-4 | 4-8 | >8.0 |
| | Sodicity (ESP) | % | <5 | 5-10 | 10-15 | >15 |
| Erosion hazard | Slope | % | 0-3 | 3-5 | 5-10 | >10- |

Table 7.22 Land suitability criteria for Jamun

| Land use requirement | | Rating | | | | |
|------------------------------|---|----------------------|--------------------------|--------------------------|-------------------|----------|
| Soil –site characteristics | Unit | Highly suitable (S1) | Moderately suitable (S2) | Marginally suitable (S3) | Not suitable (N1) | |
| Climatic regime | Mean temperature in growing season | °C | | | | |
| | Mean max. temp. in growing season | °C | | | | |
| | Mean min. tempt. in growing season | °C | | | | |
| | Mean RH in growing season | % | | | | |
| | Total rainfall | mm | | | | |
| | Rainfall in growing season | mm | | | | |
| Land quality | Soil-site characteristic | | | | | |
| Moisture availability | Length of growing period for short duration | Days | | | | |
| | Length of growing period for long duration | | | | | |
| | AWC | mm/m | | | | |
| Oxygen availability to roots | Soil drainage | Class | Well | Mod. well | Poorly | V.Poorly |
| | Water logging in growing season | Days | | | | |
| Nutrient availability | Texture | Class | scl, cl, sc, c(red) | sl, c (black) | ls | - |
| | pH | 1:2.5 | 6.0-7.8 | 5.0-6.0 | 7.8-8.4 | >8.4 |
| | CEC | C mol (p+)/Kg | | | | |
| | BS | % | | | | |
| | CaCO ₃ in root zone | % | | <5 | 5-10 | >10 |
| | OC | % | | | | |
| Rooting conditions | Effective soil depth | cm | >150 | 100-150 | 50-100 | <50 |
| | Stoniness | % | | | | |
| | Coarse fragments | Vol % | <15 | 15-35 | 35-60 | >60 |
| Soil toxicity | Salinity (EC saturation extract) | dS/m | <2.0 | 2-4 | 4-8 | >8.0 |
| | Sodicity (ESP) | % | <5 | 5-10 | 10-15 | >15 |
| Erosion hazard | Slope | % | 0-3 | 3-5 | 5-10 | >10 |

Table 7.23 Land suitability criteria for Musambi

| Land use requirement | | | Rating | | | |
|------------------------------|---|---------------|----------------------|--------------------------|--------------------------|-------------------|
| Soil –site characteristics | | Unit | Highly suitable (S1) | Moderately suitable (S2) | Marginally suitable (S3) | Not suitable (N1) |
| Climatic regime | Mean temperature in growing season | °C | 28-30 | 31-35 24-27 | 36-40 20-23 | >40 <20 |
| | Mean max. temp. in growing season | °C | | | | |
| | Mean min. temp. in growing season | °C | | | | |
| | Mean RH in growing season | % | | | | |
| | Total rainfall | mm | | | | |
| | Rainfall in growing season | mm | | | | |
| Land quality | Soil-site characteristic | | | | | |
| Moisture availability | Length of growing period for short duration | Days | | | | |
| | Length of growing period for long duration | | | | | |
| | AWC | mm/m | | | | |
| Oxygen availability to roots | Soil drainage | Class | Well drained | Moderately drained | poorly | Very poorly |
| | Water logging in growing season | Days | | | | |
| Nutrient availability | Texture | Class | scl, cl, sc, c | sl | ls | - |
| | pH | 1:2.5 | 6.0-7.8 | 5.5-6.0 7.8-8.4 | 5.0-5.5 8.4-9.0 | >9.0 |
| | CEC | C mol (p+)/Kg | | | | |
| | BS | % | | | | |
| | CaCO ₃ in root zone | % | | <5 | 5-10 | >10 |
| | OC | % | | | | |
| Rooting conditions | Effective soil depth | cm | >100 | 75-100 | 50-75 | <50 |
| | Stoniness | % | | | | |
| | Coarse fragments | Vol % | <15 | 15-35 | 35-60 | 60-80 |
| Soil toxicity | Salinity (EC saturation extract) | dS/m | <2.0 | 2-4 | 4-8 | >8.0 |
| | Sodicity (ESP) | % | <5 | 5-10 | 10-15 | >15 |
| Erosion hazard | Slope | % | <3 | 3-5 | 5-10 | >10 |

Table 7.24 Land suitability criteria for Lime

| Land use requirement | | | Rating | | | |
|------------------------------|---|---------------|----------------------|--------------------------|--------------------------|-------------------|
| Soil –site characteristics | | Unit | Highly suitable (S1) | Moderately suitable (S2) | Marginally suitable (S3) | Not suitable (N1) |
| Climatic regime | Mean temperature in growing season | °C | 28-30 | 31-35 24-27 | 36-40 20-23 | >40 <20 |
| | Mean max. temp. in growing season | °C | | | | |
| | Mean min. tempt. in growing season | °C | | | | |
| | Mean RH in growing season | % | | | | |
| | Total rainfall | mm | | | | |
| | Rainfall in growing season | mm | | | | |
| Land quality | Soil-site characteristic | | | | | |
| Moisture availability | Length of growing period for short duration | Days | | | | |
| | Length of growing period for long duration | | | | | |
| | AWC | mm/m | | | | |
| Oxygen availability to roots | Soil drainage | Class | Well drained | Moderately drained | poorly | Very poorly |
| | Water logging in growing season | Days | | | | |
| Nutrient availability | Texture | Class | scl, cl, sc, c | sl | ls | - |
| | pH | 1:2.5 | 6.0-7.8 | 5.5-6.0 7.8-8.4 | 5.0-5.5 8.4-9.0 | >9.0 |
| | CEC | C mol (p+)/Kg | | | | |
| | BS | % | | | | |
| | CaCO3 in root zone | % | | <5 | 5-10 | >10 |
| | OC | % | | | | |
| Rooting conditions | Effective soil depth | cm | >100 | 75-100 | 50-75 | <50 |
| | Stoniness | % | | | | |
| | Coarse fragments | Vol % | <15 | 15-35 | 35-60 | 60-80 |
| Soil toxicity | Salinity (EC saturation extract) | dS/m | <2.0 | 2-4 | 4-8 | >8.0 |
| | Sodicity (ESP) | % | <5 | 5-10 | 10-15 | >15 |
| Erosion hazard | Slope | % | <3 | 3-5 | 5-10 | >10 |

Table 7.25 Land suitability criteria for Cashew

| Land use requirement | | | Rating | | | |
|------------------------------|---|----------------|----------------------|--------------------------|--------------------------|---------------------|
| Soil –site characteristics | | Unit | Highly suitable (S1) | Moderately suitable (S2) | Marginally suitable (S3) | Not suitable (N1) |
| Climatic regime | Mean temperature in growing season | °C | 32 to 34 | 28 to 32; 34 to 38 | 24 to 28; 38 to 40 | <20; >40 |
| | Mean max. temp. in growing season | °C | | | | |
| | Mean min. tempt. in growing season | °C | | | | |
| | Mean RH in growing season | % | | | | |
| | Total rainfall | mm | | | | |
| | Rainfall in growing season | mm | | | | |
| Land quality | Soil-site characteristic | | | | | |
| Moisture availability | Length of growing period for short duration | Days | | | | |
| | Length of growing period for long duration | | | | | |
| | AWC | mm/m | | | | |
| Oxygen availability to roots | Soil drainage | Class | Well drained | moderately well drained | Poorly drained | Very poorly drained |
| | Water logging in growing season | Days | | | | |
| Nutrient availability | Texture | Class | scl, cl, sc, c (red) | - | sl, ls | c (black) |
| | pH | 1:2.5 | 5.5-6.5 | 5.0-5.5 6.5-7.3 | 7.3-7.8 | >7.8 |
| | CEC | C mol (p+)/ Kg | | | | |
| | BS | % | | | | |
| | CaCO3 in root zone | % | | <5 | 5-10 | >10 |
| | OC | % | | | | |
| Rooting conditions | Effective soil depth | cm | >100 | 75-100 | 50-75 | <50 |
| | Stoniness | % | | | | |
| | Coarse fragments | Vol % | <15 | 15-35 | 35-60 | 60-80 |
| Soil toxicity | Salinity (EC saturation extract) | dS/m | <2 | 2-4 | 4-8 | >8 |
| | Sodicity (ESP) | % | <5 | 5-10 | 10-15 | >15 |
| Erosion hazard | Slope | % | <3 | 3-10 | >10 | - |

Table 7.26 Land suitability criteria for Custard apple

| Land use requirement | | | Rating | | | |
|------------------------------|---|---------------|---------------------------------|--------------------------|--------------------------|-------------------|
| Soil –site characteristics | | Unit | Highly suitable (S1) | Moderately suitable (S2) | Marginally suitable (S3) | Not suitable (N1) |
| Climatic regime | Mean temperature in growing season | °C | | | | |
| | Mean max. temp. in growing season | °C | | | | |
| | Mean min. tempt. in growing season | °C | | | | |
| | Mean RH in growing season | % | | | | |
| | Total rainfall | mm | | | | |
| | Rainfall in growing season | mm | | | | |
| Land quality | Soil-site characteristic | | | | | |
| Moisture availability | Length of growing period for short duration | Days | | | | |
| | Length of growing period for long duration | | | | | |
| | AWC | mm/m | | | | |
| Oxygen availability to roots | Soil drainage | Class | Well drained | Mod. well drained | Poorly drained | V.Poorly drained |
| | Water logging in growing season | Days | | | | |
| Nutrient availability | Texture | Class | Scl, cl, sc, c (red), c (black) | - | Sl, ls | - |
| | pH | 1:2.5 | 6.0-7.3 | 5.5-6.0 7.3-8.4 | 5.0-5.5 8.4-9.0 | >9.0 |
| | CEC | C mol (p+)/Kg | | | | |
| | BS | % | | | | |
| | CaCO3 in root zone | % | | <5 | 5-10 | >10 |
| | OC | % | | | | |
| Rooting conditions | Effective soil depth | cm | >75 | 50-75 | 25-50 | <25 |
| | Stoniness | % | | | | |
| | Coarse fragments | Vol % | <15-35 | 35-60 | 60-80 | - |
| Soil toxicity | Salinity (EC saturation extract) | dS/m | <2.0 | 2-4 | 4-8 | >8.0 |
| | Sodicity (ESP) | % | <5 | 5-10 | 10-15 | >15 |
| Erosion hazard | Slope | % | 0-3 | 3-5 | >5 | - |

Table 7.27 Land suitability criteria for Amla

| Land use requirement | | | Rating | | | |
|------------------------------|---|---------------|----------------------|--------------------------|--------------------------|-------------------|
| Soil –site characteristics | | Unit | Highly suitable (S1) | Moderately suitable (S2) | Marginally suitable (S3) | Not suitable (N1) |
| Climatic regime | Mean temperature in growing season | °C | | | | |
| | Mean max. temp. in growing season | °C | | | | |
| | Mean min. temp. in growing season | °C | | | | |
| | Mean RH in growing season | % | | | | |
| | Total rainfall | mm | | | | |
| | Rainfall in growing season | mm | | | | |
| Land quality | Soil-site characteristic | | | | | |
| Moisture availability | Length of growing period for short duration | Days | | | | |
| | Length of growing period for long duration | | | | | |
| | AWC | mm/m | | | | |
| Oxygen availability to roots | Soil drainage | Class | Well drained | Mod.well drained | Poorly drained | V. Poorly drained |
| | Water logging in growing season | Days | | | | |
| Nutrient availability | Texture | Class | scl, cl, sc, c (red) | c (black) | ls, sl | - |
| | pH | 1:2.5 | 5.5-7.3 | 5.0-5.5 7.3-7.8 | 7.8-8.4 | >8.4 |
| | CEC | C mol (p+)/Kg | | | | |
| | BS | % | | | | |
| | CaCO3 in root zone | % | | <5 | 5-10 | >10 |
| | OC | % | | | | |
| Rooting conditions | Effective soil depth | cm | >75 | 50-75 | 25-50 | <25 |
| | Stoniness | % | | | | |
| | Coarse fragments | Vol % | <15-35 | 35-60 | 60-80 | - |
| Soil toxicity | Salinity (EC saturation extract) | dS/m | <2.0 | 2-4 | 4-8 | >8.0 |
| | Sodicity (ESP) | % | <5 | 5-10 | 10-15 | >15 |
| Erosion hazard | Slope | % | 0-3 | 3-5 | 5-10 | >10 |

Table 7.28 Land suitability criteria for Tamarind

| Land use requirement | | | Rating | | | |
|------------------------------|---|---------------|----------------------|--------------------------|--------------------------|-------------------|
| Soil –site characteristics | | Unit | Highly suitable (S1) | Moderately suitable (S2) | Marginally suitable (S3) | Not suitable (N1) |
| Climatic regime | Mean temperature in growing season | °C | | | | |
| | Mean max. temp. in growing season | °C | | | | |
| | Mean min. tempt. in growing season | °C | | | | |
| | Mean RH in growing season | % | | | | |
| | Total rainfall | mm | | | | |
| | Rainfall in growing season | mm | | | | |
| Land quality | Soil-site characteristic | | | | | |
| Moisture availability | Length of growing period for short duration | Days | | | | |
| | Length of growing period for long duration | | | | | |
| | AWC | mm/m | | | | |
| Oxygen availability to roots | Soil drainage | Class | Well drained | Mod.well drained | Poorly drained | V.Poorly drained |
| | Water logging in growing season | Days | | | | |
| Nutrient availability | Texture | Class | scl, cl,sc, c (red) | sl, c (black) | ls | - |
| | pH | 1:2.5 | 6.0-7.3 | 5.0-6.0 7.3-7.8 | 7.8-8.4 | >8.4 |
| | CEC | C mol (p+)/Kg | | | | |
| | BS | % | | | | |
| | CaCO3 in root zone | % | | <5 | 5-10 | >10 |
| | OC | % | | | | |
| Rooting conditions | Effective soil depth | cm | >150 | 100-150 | 75-100 | <75 |
| | Stoniness | % | | | | |
| | Coarse fragments | Vol % | <15 | 15-35 | 35-60 | 60-80 |
| Soil toxicity | Salinity (EC saturation extract) | dS/m | <2 | 2-4 | 4-8 | >8 |
| | Sodicity (ESP) | % | <5 | 5-10 | 10-15 | >15 |
| Erosion hazard | Slope | % | 0-3 | 3-5 | 5-10 | >10 |

Table 7.29 Land suitability criteria for Marigold

| Land use requirement | | | Rating | | | |
|------------------------------|---|---------------|-------------------------|--------------------------|--------------------------|-------------------|
| Soil –site characteristics | | Unit | Highly suitable (S1) | Moderately suitable (S2) | Marginally suitable (S3) | Not suitable (N1) |
| Climatic regime | Mean temperature in growing season | °C | 18-23 | 17-15 24-35 | 35-40 10-14 | >40 <10 |
| | Mean max. temp. in growing season | °C | | | | |
| | Mean min. tempt. in growing season | °C | | | | |
| | Mean RH in growing season | % | | | | |
| | Total rainfall | mm | | | | |
| | Rainfall in growing season | mm | | | | |
| Land quality | Soil-site characteristic | | | | | |
| Moisture availability | Length of growing period for short duration | Days | | | | |
| | Length of growing period for long duration | | | | | |
| | AWC | mm/m | | | | |
| Oxygen availability to roots | Soil drainage | Class | Well drained | Moderately well drained | Poorly drained | V.Poorly drained |
| | Water logging in growing season | Days | | | | |
| Nutrient availability | Texture | Class | sl,scl, cl, sc, c (red) | c (black) | ls | - |
| | pH | 1:2.5 | 6.0-7.3 | 5.0-6.0 7.3-8.4 | 8.4-9.0 | >9.0 |
| | CEC | C mol (p+)/Kg | | | | |
| | BS | % | | | | |
| | CaCO ₃ in root zone | % | | <5 | 5-10 | >10 |
| | OC | % | | | | |
| Rooting conditions | Effective soil depth | cm | >75 | 50-75 | 25-50 | <25 |
| | Stoniness | % | | | | |
| | Coarse fragments | Vol % | <15 | 15-35 | 35-60 | 60-80 |
| Soil toxicity | Salinity (EC saturation extract) | dS/m | <2.0 | 2-4 | 4-8 | >8.0 |
| | Sodicity (ESP) | % | | | | |
| Erosion hazard | Slope | % | <3 | 3-5 | 5-10 | >10 |

Table 7.30 Land suitability criteria for Chrysanthemum

| Land use requirement | | | Rating | | | |
|------------------------------|---|---------------|-------------------------|--------------------------|--------------------------|-------------------|
| Soil –site characteristics | | Unit | Highly suitable (S1) | Moderately suitable (S2) | Marginally suitable (S3) | Not suitable (N1) |
| Climatic regime | Mean temperature in growing season | °C | 18-23 | 17-15 24-35 | 35-40 10-14 | >40 <10 |
| | Mean max. temp. in growing season | °C | | | | |
| | Mean min. temp. in growing season | °C | | | | |
| | Mean RH in growing season | % | | | | |
| | Total rainfall | mm | | | | |
| | Rainfall in growing season | mm | | | | |
| Land quality | Soil-site characteristic | | | | | |
| Moisture availability | Length of growing period for short duration | Days | | | | |
| | Length of growing period for long duration | | | | | |
| | AWC | mm/m | | | | |
| Oxygen availability to roots | Soil drainage | Class | Well drained | Moderately well drained | Poorly drained | V.Poorly drained |
| | Water logging in growing season | Days | | | | |
| Nutrient availability | Texture | Class | sl,scl, cl, sc, c (red) | c (black) | ls | - |
| | pH | 1:2.5 | 6.0-7.3 | 5.0-6.0 7.3-8.4 | 8.4-9.0 | >9.0 |
| | CEC | C mol (p+)/Kg | | | | |
| | BS | % | | | | |
| | CaCO3 in root zone | % | | <5 | 5-10 | >10 |
| | OC | % | | | | |
| Rooting conditions | Effective soil depth | cm | >75 | 50-75 | 25-50 | <25 |
| | Stoniness | % | | | | |
| | Coarse fragments | Vol % | <15 | 15-35 | 35-60 | 60-80 |
| Soil toxicity | Salinity (EC saturation extract) | dS/m | <2.0 | 2-4 | 4-8 | >8.0 |
| | Sodicity (ESP) | % | | | | |
| Erosion hazard | Slope | % | <3 | 3-5 | 5-10 | >10 |

Table 7.31 Land suitability criteria for Jasmine (irrigated)

| Land use requirement | | | Rating | | | |
|------------------------------|---|----------------------|--------------------------|--------------------------|-------------------|------------------|
| Soil –site characteristics | Unit | Highly suitable (S1) | Moderately suitable (S2) | Marginally suitable (S3) | Not suitable (N1) | |
| Climatic regime | Mean temperature in growing season | °C | 18-23 | 17-15 24-35 | 35-40 10-14 | - |
| | Mean max. temp. in growing season | °C | | | | |
| | Mean min. temp. in growing season | °C | | | | |
| | Mean RH in growing season | % | | | | |
| | Total rainfall | mm | | | | |
| | Rainfall in growing season | mm | | | | |
| Land quality | Soil-site characteristic | | | | | |
| Moisture availability | Length of growing period for short duration | Days | | | | |
| | Length of growing period for long duration | | | | | |
| | AWC | mm/m | | | | |
| Oxygen availability to roots | Soil drainage | Class | Well drained | Moderately well drained | Poorly drained | V.Poorly drained |
| | Water logging in growing season | Days | | | | |
| Nutrient availability | Texture | Class | scl, cl, sc, c (red) | sl | ls, c (black) | - |
| | pH | 1:2.5 | 6.0-7.3 | 5.0-6.0 7.3-8.4 | 8.4-9.0 | >9.0 |
| | CEC | C mol (p+)/Kg | | | | |
| | BS | % | | | | |
| | CaCO3 in root zone | % | | <5 | 5-10 | >10 |
| | OC | % | | | | |
| Rooting conditions | Effective soil depth | cm | >75 | 50-75 | 25-50 | <25 |
| | Stoniness | % | | | | |
| | Coarse fragments | Vol % | <15 | 15-35 | 35-60 | 60-80 |
| Soil toxicity | Salinity (EC saturation extract) | dS/m | <2.0 | 2-4 | 4-8 | >8.0 |
| | Sodicity (ESP) | % | | | | |
| Erosion hazard | Slope | % | <3 | 3-5 | 5-10 | >10 |

7.32 Land suitability criteria for Crossandra

| Land use requirement | | | Rating | | | |
|------------------------------|---|---------------|----------------------|--------------------------|--------------------------|-------------------------------|
| Soil –site characteristics | | Unit | Highly suitable (S1) | Moderately suitable (S2) | Marginally suitable (S3) | Not suitable (N1) |
| Climatic regime | Mean temperature in growing season | °C | | | | |
| | Mean max. temp. in growing season | °C | | | | |
| | Mean min. tempt. in growing season | °C | | | | |
| | Mean RH in growing season | % | | | | |
| | Total rainfall | mm | | | | |
| | Rainfall in growing season | mm | | | | |
| Land quality | Soil-site characteristic | | | | | |
| Moisture availability | Length of growing period for short duration | Days | | | | |
| | Length of growing period for long duration | | | | | |
| | AWC | mm/m | | | | |
| Oxygen availability to roots | Soil drainage | Class | Well drained | Moderately well drained | - | Poorly to very poorly drained |
| | Water logging in growing season | Days | | | | |
| Nutrient availability | Texture | Class | scl, cl, sc, c(red) | sl, | c (black),ls | - |
| | pH | 1:2.5 | 6.0-7.3 | 5.0-6.0 7.3-8.4 | 8.4-9.0 | >9.0 |
| | CEC | C mol (p+)/Kg | | | | |
| | BS | % | | | | |
| | CaCO3 in root zone | % | | <5 | 5-10 | >10 |
| | OC | % | | | | |
| Rooting conditions | Effective soil depth | cm | >75 | 50-75 | 25-50 | <25 |
| | Stoniness | % | | | | |
| | Coarse fragments | Vol % | <15 | 15-35 | 35-60 | 60-80 |
| Soil toxicity | Salinity (EC saturation extract) | dS/m | <2.0 | 2-4 | 4-8 | >8.0 |
| | Sodicity (ESP) | % | | | | |
| Erosion hazard | Slope | % | <3 | 3-5 | 5-10 | >10 |

7.29 Land Management Units (LMUs)

The 31 soil map units identified in Haligeri-3 Microwatershed have been grouped into nine Land Management Units (LMUs) for the purpose of preparing a Proposed Crop Plan. Land Management Units are grouped based on the similarities in respect of the type of soil, the depth of the soil, the surface soil texture, gravel content, AWC, slope, erosion etc. and a Land Management Units map (Fig.7.32) has been generated. These Land Management Units are expected to behave similarly for a given level of management.

The map units that have been grouped into nine Land Management Units along with brief description of soil and site characteristics are given below.

| LMU | Mapping unit | Soil and site characteristics |
|-----|--|--|
| 1 | AWDmB2, KVRmA1, KVRmB1, LGDmB1, DRLmB2, DRLmB2g1 | Moderately deep to very deep, black calcareousness clay soils with slopes of 0-3%, slight to moderate erosion, gravelly (15-35%) |
| 2 | BPRcA1, BPRhB2g1, BPRhB2g2, BPRiB2, BPRmB2, NGPhA1, NGPiB1, BDGiB1g1, HDHcB1g1, HDHhB2g1, HDHiB1, HDHiB2g1, GDPcB2, GDPiB2 | Moderately deep to very deep, red sandy clay to clay soils with slopes of 0-3%, slight to moderate erosion, gravelly to very gravelly (15-60%) |
| 3 | NSPmB2 | Moderately deep, calcareous sodic clay soils with slopes of 1-3%, moderate erosion |
| 4 | BSRiB1g2 | Moderately deep, red sandy clay soils with slopes of 1-3%, slight erosion, very gravelly (15-60%) |
| 5 | TDHhB2g2, TDHiB1 | Moderately shallow, red sandy clay to clay soils with slopes of 1-3%, slight to moderate erosion, gravelly to very gravelly (35-60%) |
| 6 | RNKmB2, RNKmB2g1 | Moderately shallow, black calcareous clay soils with slopes of 1-3%, moderate erosion, gravelly (15-35%) |
| 7 | LKRcB2g1, MKHiB2g1 | Moderately shallow, red gravelly sandy clay soils with slopes of 1-3%, moderate erosion, gravelly (15-35%) |
| 8 | MTLmB2g1 | Shallow, black calcareous clay soils with slopes of 1-3%, moderate erosion, gravelly (15-35%) |
| 9 | CSRiB2, HRViA1g1 | Shallow, red gravelly sandy clay to sandy clay loam soils with slopes of 0-3%, slight to moderate erosion, gravelly (15-35%) |




Fig 7.32 Land Management Units map of Haligeri-3 microwatershed

7.33 Proposed Crop Plan for Haligeri-3 Microwatershed

After assessing the land suitability for the 31 crops, the proposed crop plan has been prepared for the nine identified LMUs by considering only the highly (Class S1) and moderately (Class S2) suitable lands for each of the 31 crops. The resultant proposed crop plan is presented in Table 7.33.

Table 7.33 Proposed Crop Plan for Haligeri-3 Microwatershed

| LMU | Soil Map Units | Survey Number | Field Crops/ Commercial crops | Horticulture Crops (Rainfed/Irrigated) | Suitable Interventions |
|-----|---|---|--|--|--|
| 1 | 424.AWDmB2 386.KVRmA1 388.KVRmB1 393.LGDmB1 350.DRLmB2 351.DRLmB2g1 (Moderately deep to very deep, black calcareousness clay soils) | Banapura: 36,66,67,74,75,76,95,96,97,98,99,100,101,102,107,108,109,110,111,112,113,115,116,117,118,119,120,121,124,125,126,127,128,129,130,131,132,133,134,135,136,137,138,140,143,144,145, Halageri : 67,68 Thalabala : 187 Thalakalla: 98,99,100,101,103,104,105,106,107,108,109,110,111,124,125,126,127,128,129,130,131,132 Veerapura : 186,187,188 | Maize, Sorghum, Sunflower, Bajra, Cotton, Red gram, Bengal gram, Soybean, Safflower, Linseed | Fruit crops : Pomegranate, Jamun, Lime, Musambi, Tamarind, Amla, Custard apple Vegetables: Drumstick, Chillies, Bhendi, Brinjal, Coriander Flowers: Marigold, Chrysanthemum, | Application of FYM, Biofertilizers and micronutrients, drip irrigation, mulching, suitable soil and water conservation practices |
| 2 | 220.BPRcA1 231.BPRhB2g1 232.BPRhB2g2 239.BPRiB2 240.BPRmB2 255.NGPhA1 262.NGPiB1 192.BDGiB1g1 109.HDHcB1g1 123.HDHhB2g1 125.HDHiB1 128.HDHiB2g1 267.GDPcB2 269.GDPiB2 (Moderately deep to | Banapura: 44,46,54(1),56,57,58,59,60,61,155,156,158,159/(1),159/(2),160,161,162,163,164,165,166,167,168,169,170,171,172,173,174,175,176,177,178,179,180,181,182,183,184,185,186,187,190 Halageri : 54,55,56,63,64,65, 358 Thalakalla : 102,113,115,116 | Maize, Sorghum, Sunflower, Groundnut, Bajra, Cotton, Red gram | Fruit crops : Sapota, Pomegranate, Amla, Cashew, Guava, Custard apple, Jack fruit, Jamun, Lime, Musambi Vegetables: Tomato, Chilli, Drumstick, Onion, Bhendi, Brinjal, Curry leaves Flowers: Marigold, Chrysanthemum, Jasmine, Crossandra | Drip irrigation, mulching, suitable soil and water conservation practices (Crescent Bunding with Catch Pit etc) |

| LMU | Soil Map Units | Survey Number | Field Crops/ Commercial crops | Horticulture Crops (Rainfed/Irrigated) | Suitable Interventions |
|-----|--|--|---|---|---|
| | very deep, red sandy clay to clay soils) | | | | |
| 3 | 362.NSPmB2 (Moderately deep, calcareous sodic clay soils) | Halageri :54,55 | - | Agri-Silvi-Pasture Ber, Aonla, Acacia sp. Dhaincha, Rhodes grass, Para grass ,Bermuda grass | Application of gypsum, iron pyrites and elemental sulphur. Addition of farm yard manures, green manures and providing subsurface drainage |
| 4 | 166.BSRiB1g2 (Moderately deep, red sandy clay soils) | Halageri :356,357,359 | Maize, Sorghum, Groundnut, Sunflower, Bajra, Mulberry, Cotton, Red gram | Fruit crops : Sapota, Pomegranate, Amla, Cashew, Custard apple, Guava, Jackfruit, Lime, Musambi, Vegetables: Tomato, Chillies, Drumstick, Onion, Bhendi, Brinjal, Curry leaves Flowers: Marigold, Chrysanthemum, Jasmine, Crossandra | Drip irrigation, mulching, suitable soil and water conservation practises (Crescent Bunding with Catch Pit etc) |
| 5 | 59.TDHhB2g2 60.TDHiB1 (Moderately shallow, red sandy clay to clay soils) | Banapura: 188,189,191,192,195,196 | Maize, Sorghum, Groundnut, Bajra, Cotton, Horse gram, Castor | Fruit crops : Amla, Custard apple Vegetables: Tomato, Chilli, Onion, Bhendi, Brinjal ,Curry leaves Flowers: Marigold, Chrysanthemum, Jasmine, Crossandra | Drip irrigation, mulching, suitable soil and water conservation practices (Crescent Bunding with Catch Pit etc) |

| LMU | Soil Map Units | Survey Number | Field Crops/ Commercial crops | Horticulture Crops (Rainfed/Irrigated) | Suitable Interventions |
|-----|---|--|--------------------------------------|--|---|
| 6 | 336.RNKmB2 337.RNKmB2g1 (Moderately shallow, black calcareous sodic clay soils) | Banapura: 62,63,64,65,68,69,139,141,142,146,147,148,149,150,151,152,153,154,157 Halageri : 62,66,69 | - | Agri-Silvi-Pasture Ber, Aonla, Acacia sp. Dhaincha, Rhodes grass, Para grass ,Bermuda grass | Application of gypsum, iron pyrites and elemental sulphur. Addition of farm yard manures, green manures and providing subsurface drainage |
| 7 | 43.LKRcB2g1 90.MKHiB2g1 (Moderately shallow, red gravelly sandy clay soils) | Banapura: 37,38,40,41,42,43,45,47, 70,71,72,73 | Bajra, Groundnut, Horse gram, Castor | Fruit crops : Amla, Custard apple Vegetables: Curry leaves Flowers: Marigold, Chrysanthemum | Drip irrigation, mulching, suitable soil and water conservation practices (Crescent Bunding with Catch Pit etc) |
| 8 | 311.MTLmB2g1 (Shallow, black calcareous clay soils) | Halageri: 60,61,70,71,72,73,74,75,76 | - | Agri-Silvi-Pasture: Hybrid Napier, <i>Styloxanthes hamata</i> , <i>Styloxanthes scabra</i> | Use of short duration varieties, sowing across the slope |
| 9 | 39.CSRiB2 29.HRViA1g1 (Shallow, red gravelly sandy clay to sandy clay loam soils) | Thalakalla : 112,114 | - | Agri-Silvi-Pasture: Custard apple, Hybrid Napier, <i>Styloxanthes hamata</i> , Glyricidia, <i>Styloxanthes scabra</i> | Use of short duration varieties, sowing across the slope and split application of nitrogen fertilizers |

SOIL HEALTH MANAGEMENT

8.1 Soil Health

Soil health is basic to plant health and plant health is basic to human and bovine health. Soil is fundamental to crop production. Without soil, no food could be produced nor would livestock be fed on a large scale. Because it is finite and fragile, soil is a precious resource that requires special care from its users.

Soil health or the capacity of the soil to function is critical to human survival. Soil health has been defined as: “the capacity of the soil to function as a living system without adverse effect on the ecosystem”. Healthy soils maintain a diverse community of soil organisms that help to form beneficial symbiotic associations with plant roots, recycle essential plant nutrients, improve soil structure with positive repercussions for soil, water and nutrient holding capacity and ultimately improve crop production and also contribute to mitigating climate change by maintaining or increasing its carbon content.

Functional interactions of soil biota with organic and inorganic components, air and water determine a soil’s potential to store and release nutrients, and water to plants and to promote and sustain plant growth. Thus, maintaining soil health is vital to crop production and conserve soil resource base for sustaining agriculture.

The most important characteristics of a healthy soil are

- Good soil tilth
- Sufficient soil depth
- Good water storage and good drainage
- Adequate supply, but not excess of nutrients
- Large population of beneficial organisms
- Small proportion of plant pathogens and insect pests
- Low weed pressure
- Free of chemicals and toxins that may harm the crop
- Resistance to degradation
- Resilience when unfavourable conditions occur

Characteristics of Haligeri-3 Microwatershed

- ❖ The soil phases with sizeable area identified in the microwatershed belonged to the soil series of DRL (163 ha), RNK (134 ha), LGD(104 ha), BPR(90 ha), HDH(73 ha), NGP(36 ha), GDP(35 ha), BDG(31 ha), MKH(28 ha), MTL(25 ha), KVR(20 ha), LKR (17 ha), CSR(13 ha), TDH(10 ha), HRV(6 ha), BSR (4 ha), AWD(4 ha), NSP(1 ha)
- ❖ As per land capability classification, entire area in the microwatershed falls under arable land category (Class II and III). The major limitations identified in the arable lands were soil and erosion.

- ❖ On the basis of soil reaction, an area of about 37 ha (5%) is neutral (pH 6.5-7.3), 492 ha (61%) is slightly alkaline (pH 7.3-7.8), 252 ha (31%) is strongly alkaline (pH 8.4-9.0), and 14 ha(2%) is very strongly alkaline (pH >9.0) in reaction.

Soil Health Management

The following actions are required to improve the current land husbandry practices that provide a sound basis for the successful adoption of sustainable crop production system.

Alkaline soils

An area of about 758 ha (94%) is under alkaline soils. The following actions are recommended.

1. Regular addition of organic manure, green manuring, green leaf manuring, crop residue incorporation and mulching needs to be taken up to improve the soil organic matter status.
2. Application of biofertilizers (Azospirillum, Azotobacter, Rhizobium).
3. Application of 25% extra N and P (125 % RDN&P).
4. Application of ZnSO₄ – 12.5 kg/ha (once in three years).
5. Application of Boron – 5 kg/ha (once in three years).

Neutral soils

Neutral soils cover about 37 ha (5%) and the following actions are recommended.

6. Regular addition of organic manure, green manuring, green leaf manuring, crop residue incorporation and mulching needs to be taken up to improve the soil organic matter status.
7. Application of biofertilizers, (Azospirillum, Azotobacter, Rhizobium).
8. Application of 100 per cent RDF.
9. Need based micronutrient applications.

Soil Degradation

Soil erosion is one of the major factor affecting the soil health in the microwatershed. An area of about 540 ha (67%) is under moderate erosion. The areas with moderate erosion need immediate soil and water conservation and, other land development and land husbandry practices for restoring soil health.

Dissemination of Information and Communication of Benefits

Any large scale implementation of soil health management requires that supporting information is made available widely, particularly through channels familiar to farmers and extension workers. Given the very high priority attached to soil health especially by the Central Government on issuing Soil-Health Cards to all the farmers, media outlets like Regional, State and National Newspapers, Radio and Dooradarshan programs in local languages but also modern information and communication

technologies such as Cellular phones and the Internet, which can be much more effective in reaching the younger farmers.

Inputs for Net Planning (Saturation Plan) and Interventions needed

Net planning in IWMP is focusing on preparation of

1. Soil and Water Conservation Treatment Plans for each plot or farm.
2. Productivity enhancement measures/ interventions for existing crops/livestock/other farm enterprises.
3. Diversification of farming mainly with perennial horticultural crops and livestock.
4. Improving livelihood opportunities and income generating activities.

In this connection, how various outputs of Sujala-III are of use in addressing these objectives of Net Planning are briefly presented below.

- ❖ **Soil Depth:** The depth of a soil decides the amount of moisture and nutrients it can hold, what crops can be taken up or not, depending on the rooting depth and the length of growing period available for raising any crop. Deeper the soil, better for a wide variety of crops. If sufficient depth is not available for growing deep rooted crops, either choose medium or short duration crops or deeper planting pits need to be opened and additional good quality soil brought from outside has to be filled into the planting pits.
- ❖ **Surface Soil Texture:** Lighter soil texture in the top soil means, better rain water infiltration, less run-off and soil moisture conservation, less capillary rise and less evaporation losses. Lighter surface textured soils are amenable to good soil tilth and are highly suitable for crops like groundnut, root vegetables (carrot, radish, potato etc) but not ideal for crops that need stagnant water like lowland paddy. Heavy textured soils are poor in water infiltration and percolation. They are prone for sheet erosion; such soils can be improved by sand mulching. The technology that is developed by the AICRP-Dryland Agriculture, Vijayapura, Karnataka can be adopted.
- ❖ **Gravelliness:** More gravel content is favorable for run-off harvesting but poor in soil moisture storage and nutrient availability. It is a significant parameter that decides the kind of crop to be raised.
- ❖ **Land Capability Classification:** The land capability map shows the areas suitable and not suitable for agriculture and the major constraints in each of the plot/survey number. Hence, one can decide what kind of enterprise is possible in each of these units. In general, erosion and soil are the major constraints in Haligeri-3 Microwatershed.
- ❖ **Organic Carbon:** An area of about 263 ha (33%) is low (<0.5%), 343 ha (43%) is medium (0.5-0.75%) and 189 ha (24%) is high (>0.75%) in OC. The areas that are low and medium in OC needs to be further improved by applying farmyard manure and rotating crops with cereals and legumes or mixed cropping.

- ❖ **Promoting green manuring:** Growing of green manuring crops costs Rs. 1250/ha (green manuring seeds) and about Rs. 2000/ha towards cultivation that totals to Rs. 3250/- per ha. On the other hand, application of organic manure @ 10 tons/ha costs Rs. 5000/ha. The practice needs to be continued for 2-3 years or more. Nitrogen fertilizer needs to be supplemented by 25% in addition to the recommended level in 606 ha area where OC is less than 0.75 per cent. For example, for rainfed maize, recommended level is 50 kg N per ha and an additional 12 kg /ha needs to be applied for all the crops grown in these plots.
- ❖ **Available Phosphorus:** Available phosphorus is low (<23 kg/ha) in 251 ha (31%), medium (23-57 kg/ha) in 531 ha (66%) and high (>57 kg/ha) in 13 ha (2%) area of the microwatershed. The areas with high phosphorus content reduce 25% from the RDF to avoid the excess application of fertilizer and apply additional 25% phosphorus in areas where it is low and medium.
- ❖ **Available Potassium:** Available potassium is medium (145-337 kg/ha) in 120 ha (15%) and high (>337 kg/ha) in 675 ha (84%) area of the microwatershed. The areas with high potassium content reduce 25% from the RDF to avoid the excess application of fertilizer and apply additional 25% potassium in areas where it is medium.
- ❖ **Available Sulphur:** Available sulphur is a very critical nutrient for oilseed crops. Available sulphur is low (<10 ppm) in 300 ha (37%), medium (10-20 ppm) in 462 ha (58%) and high (>20 ppm) in 32 ha (4%) area of the microwatershed. Areas with low and medium in available sulphur need to be applied with magnesium sulphate or gypsum or Factamphos (p) fertilizer (13% sulphur) for 2-3 years for the deficiency to be corrected.
- ❖ **Available Iron:** It is deficient (<4.5 ppm) in the entire area of the microwatershed. To manage iron deficiency iron sulphate @ 25 kg/ha needs to be applied for 2-3 years.
- ❖ **Available Zinc:** It is deficient (<0.6 ppm) in the entire area of the microwatershed. Application of zinc sulphate @ 25kg/ha is to be followed in areas that are deficient in available zinc.
- ❖ **Available Boron:** Available boron is low in (<0.5ppm) 294 ha (37%), medium (0.5-1.0 ppm) in 453 ha (56%) and high(>1.0 ppm) in 48 ha (6%) area in the microwatershed. The areas with low and medium in boron content need to be applied with sodium borate @ 10kg/ha as soil application or 0.2% borax as foliar spray to correct the deficiency.
- ❖ **Available Manganese:** It is sufficient in the entire area of the microwatershed.
- ❖ **Available Copper:** It is sufficient in the entire area of the microwatershed.
- ❖ **Soil Alkalinity:** An area of about 758 ha (93%) in the microwatershed has soils that are slightly to very strongly alkaline. These areas need application of gypsum and wherever calcium is in excess, iron pyrites and element sulphur can be recommended. Management practices like treating repeatedly with good quality


water to drain out the excess salts and provision of subsurface drainage and growing of salt tolerant crops like Casuarina, Acacia, Neem, Ber etc, are recommended.

- ❖ **Land Suitability for various crops:** Areas that are highly, moderately and marginally suitable and not suitable for growing various crops are indicated. Along with the suitability, various constraints that are limiting the productivity are also indicated. For example, in case of cotton, gravel content, rooting depth and salinity/alkalinity are the major constraints in various plots. With suitable management interventions, the productivity can be enhanced. In order to increase water holding capacity of light textured soils, growing of green manure crops and application of organic manure is recommended.

SOIL AND WATER CONSERVATION TREATMENT PLAN

For preparing soil and water conservation treatment plan for Haligeri-3 Microwatershed, the land resource inventory database generated under Sujala-III project has been transformed as information through series of interpretative (thematic) maps using soil phase map as a base. The various thematic maps (1:7920 scale) generated were

- Soil depth
- Surface soil texture
- Available water capacity
- Soil slope
- Soil gravelliness
- Land capability
- Present land use and land cover
- Crop suitability maps
- Rainfall map
- Hydrology
- Water Resources
- Socio-economic data
- Contour plan with existing features- network of waterways, pottissa boundaries, cut up/ minor terraces etc.
- Cadastral map (1:7920 scale)
- Satellite imagery (1:7920 scale)



Apart from these, Hand Level/ Hydro Marker/ Dumpy Level/ Total Station and Kathedars' List to be collected.

Steps for Survey and Preparation of Treatment Plan

The boundaries of Land User Groups' and Survey No. boundaries are traced in the field.

- Naming of user groups and farmers
- Identification of arable and non arable lands
- Identification of drainage lines and gullies
- Identification of non treatable areas
- Identification of priority areas in the arable lands
- Treatment plan for arable lands
- Location of water harvesting and recharge structures

9.1 Treatment Plan

The treatment plan recommended for arable lands is briefly described below.

9.1.1 Arable Land Treatment

A. BUNDING

| Steps for Survey and Preparation of Treatment Plan | | USER GROUP-1 |
|---|----------------------------|--|
| Cadastral map (1:7920 scale) is enlarged to a scale of 1:2500 scale Existing network of waterways, pothissa boundaries, grass belts, natural drainage lines/ watercourse, cut ups/ terraces are marked on the cadastral map to the scale Drainage lines are demarcated into | | |
| Small gullies | (up to 5 ha catchment) | <div style="text-align: center;"> CLASSIFICATION OF GULLIES </div> <div style="text-align: center; background-color: #90EE90; padding: 5px;"> ಕೊರಕಲಿನ ವರ್ಗೀಕರಣ </div> <div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> <p>UPPER REACH</p> <p>MIDDLE REACH</p> <p>LOWER REACH</p> </div> <div style="width: 50%;"> <ul style="list-style-type: none"> • ಮೇಲ್ಭಾಗ 15 Ha. • ಮಧ್ಯಭಾಗ 15+10=25 ಹೆ. • ಕೆಳಭಾಗ 25 ಹೆಕ್ಕೂ ಹೆಚ್ಚು ಅಧಿಕ </div> </div> <div style="text-align: right; margin-top: 10px;"> POINT OF CONCENTRATION </div> |
| Medium gullies | (5-15 ha catchment) | |
| Ravines | (15-25 ha catchment) and | |
| Halla/Nala | (more than 25ha catchment) | |

Measurement of Land Slope

Land slope is estimated or determined by the study and interpretation of contours or by measurement in the field using simple instruments like Hand Level or Hydromarker.



Vertical and Horizontal intervals between bunds as recommended by the Watershed Development Department.

| Slope percentage | Vertical interval (m) | Corresponding Horizontal Distance (m) |
|------------------|-----------------------|---------------------------------------|
| 2 - 3% | 0.6 | 24 |
| 3 - 4% | 0.9 | 21 |
| 4 - 5% | 0.9 | 21 |
| 5 - 6% | 1.2 | 21 |
| 6 - 7% | 1.2 | 21 |

Note: i) The above intervals are maximum.

(ii) Considering the slope class and erosion status (A1... A= 0-1% slope, 1= slight erosion) the intervals have to be decided.

Bund length recording: Considering the contour plan and the existing grass belts/partitions, the bunds are aligned and lengths are measured.

Section of the Bund

Bund section is decided considering the soil texture class and gravelliness class (bg₀b = loamy sand, g₀ = <15% gravel). The recommended sections for different soils are given below.

Recommended Bund Section


| Top width (m) | Base width (m) | Height (m) | Side slope (Z:1;H:V) | Cross section (sq m) | Soil Texture | Remarks |
|---------------|----------------|------------|----------------------|----------------------|---------------------------------|-----------------|
| 0.3 | 0.9 | 0.3 | 01:01 | 0.18 | Sandy loam | Vegetative bund |
| 0.3 | 1.2 | 0.3 | 1.5:1 | 0.225 | Sandy clay | |
| 0.3 | 1.2 | 0.5 | 0.9:1 | 0.375 | Red gravelly soils | |
| 0.3 | 1.2 | 0.6 | 0.75:1 | 0.45 | | |
| 0.3 | 1.5 | 0.6 | 01:01 | 0.54 | Red sandy loam | |
| 0.3 | 2.1 | 0.6 | 1.5:1 | 0.72 | Very shallow clayey black soils | |
| 0.45 | 2 | 0.75 | 01:01 | 0.92 | | |
| 0.45 | 2.4 | 0.75 | 1.3:1 | 1.07 | Shallow clayey black soils | |
| 0.6 | 3.1 | 0.7 | 1.78:1 | 1.29 | Medium clayey black soils | |
| 0.5 | 3 | 0.85 | 1.47:1 | 1.49 | | |

Formation of Trench cum Bund


Dimensions of the Borrow Pits/ Trenches to be excavated (machinery are decided considering the Bund Section).

Details of Borrow Pit dimensions are given below

TRENCH CUM BUND




WATER STORAGE AREA




0.45 Sq.m section

IDEAL FOR HORTICULTURE CROPS

'A' FRAME FOR INTERBUND MANAGEMENT



1. ಸಮಾನಾಕಳ ಉಳುವುದು
2. ಸಮಾನಾಕಳ ಬಿತ್ತನೆ/ನಾಟಿ



Size of Borrow Pits/ Trench recommended for Trench cum Bund (by machinery)

| Bund section | Bund length | Earth quantity | Pit | | | | Berm (pit to pit) | Soil depth Class |
|--------------|-------------|----------------|------|------|------|----------------------------|-------------------|--------------------|
| | | | L(m) | W(m) | D(m) | Quantity (m ³) | | |
| 0.375 | 6 | 2.25 | 5.85 | 0.85 | 0.45 | 2.24 | 0.15 | Shallow |
| 0.45 | 6 | 2.7 | 5.4 | 1.2 | 0.43 | 2.79 | 0.6 | Shallow |
| 0.45 | 6 | 2.7 | 5 | 0.85 | 0.65 | 2.76 | 1 | Moderately Shallow |
| 0.54 | 5.6 | 3.02 | 5.5 | 0.85 | 0.7 | 3.27 | 0.1 | Moderately shallow |
| 0.54 | 5.5 | 2.97 | 5 | 1.2 | 0.5 | 3 | 0.5 | Shallow |
| 0.72 | 6.2 | 4.46 | 6 | 1.2 | 0.7 | 5.04 | 0.2 | Moderately shallow |
| 0.72 | 5.2 | 3.74 | 5.1 | 0.85 | 0.9 | 3.9 | 0.1 | Moderately deep |

B. Waterways

- a) Existing waterways are marked on the cadastral map (1:7920 scale) and their dimensions are recorded.
- b) Considering the contour plan of the MWS, additional waterways/ modernization of the existing ones can be thought of.
- c) The design details are given in the Manual.

C. Farm Ponds

Waterways and the catchment area will give an indication on the size of the Farm Pond. Location of the pond can be decided based on the contour plan/ field condition and farmers' need/desire.

D. Diversion Channel

Existing EPT/ CPT are marked on the cadastral map. Looking to the need, these can be modernized or fresh diversion channel can be proposed and runoff from this can be stored in *Gokatte/ Recharge Ponds*.

9.1.2 Non-Arable Land Treatment

Depending on the gravelliness and crops preferred by the farmers, the concerned authorities can decide appropriate treatment plan. The recommended treatments may be Contour Trench, Staggered Trench, Crescent Bund, Boulder Bund or Pebble Bund.

9.1.3 Treatment of Natural Water Course/ Drainage Lines

- a) The cadastral map has to be updated as regards the network of drainage lines (gullies/ *nalas*/ *hallas*) and existing structures are marked to the scale and storage capacity of the existing water bodies are documented.
- b) The drainage line will be demarcated into Upper Reach, Middle Reach and Lower Reach.
- c) Considering the Catchment, *Nala* bed and bank conditions, suitable structures are decided.
- d) Number of storage structures (Check dam/ *Nala* bund/ Percolation tank) will be decided considering the commitments and available runoff in water budgeting and quality of water in the wells and site suitability.
- e) Detailed Levelling Survey using Dumpy Level / Total Station has to be carried out to arrive at the site-specific designs as shown in the Manual.
- f) The location of ground water recharge structures are decided by examining the lineaments and fracture zones from geological maps.
- g) Rainfall intensity data of the nearest Rain Gauge Station is considered for Hydrologic Designs.
- h) Silt load to the Storage/Recharge Structures is reduced by providing vegetative, boulder and earthen checks in the natural water course. Location and design details are given in the Manual.

9.2 Recommended Soil and Water Conservation Measures

The appropriate conservation structures best suited for each of the land parcel/ survey number (Appendix-I) are selected based on the slope per cent, severity of erosion, amount of rainfall, land use and soil type. The different kinds of conservation structures recommended are

1. Graded / Strengthening of Bunds
2. Trench cum Bunds (TCB)
3. Trench cum Bunds / Strengthening
4. Crescent Bunds

A map (Fig. 9.1) showing soil and water conservation plan with different kinds of structures recommended has been prepared which shows the spatial distribution and extent of area. A maximum area of about 311 ha (39%) needs trench cum bunding, an area of about 446 ha (56%) needs graded bunding and 37 ha (5 %) requires strengthening of existing bunds/ bunding. The conservation plan prepared may be presented to all the stakeholders including farmers and after considering their suggestions, the conservation plan for the microwatershed may be finalized in a participatory approach.




Fig. 9.1 Soil and Water Conservation Plan map of Haligeri-3 Microwatershed

9.3 Greening of Microwatershed

As part of the greening programme in the watersheds, it is envisaged to plant a variety of horticultural and other tree plants that are edible, economical and produce lot of biomass which helps to restore the ecological balance in the watersheds. The lands that are suitable for greening programme are non-arable lands (land capability classes V, VI VII and VIII) and also the lands that are not suitable or marginally suitable for growing annual and perennial crops. The method of planting these trees is given below.

It is recommended to open the pits during the 1st week of March along the contour and heap the dugout soil on the lower side of the slope in order to harness the flowing water and facilitate weathering of soil in the pit. Exposure of soil in the pit also prevents spread of pests and diseases due to scorching sun rays. The pits should be filled with mixture of soil and organic manure during the second week of April and keep ready with sufficiently tall seedlings produced either in poly bags or in root trainer nurseries so that planting can be done during the 2nd or 3rd week of April depending on the rainfall.

The tree species suitable for the area considering rainfall, temperature and adaptability is listed below; waterlogged areas are recommended to be planted with species like Neral (*Syzgium cumini*) and Bamboo. Dry areas are to be planted with species like Honge, Bevu, Seetaphal etc.

| Dry Deciduous Species | | | Temp (°C) | Rainfall (mm) |
|--------------------------------|-----------|----------------------------------|------------------|----------------------|
| 1. | Bevu | <i>Azadiracta indica</i> | 21–32 | 400 –1,200 |
| 2. | Tapasi | <i>Holoptelia integrifolia</i> | 20-30 | 500 - 1000 |
| 3. | Seetaphal | <i>Anona Squamosa</i> | 20-40 | 400 - 1000 |
| 4. | Honge | <i>Pongamia pinnata</i> | 20 -50 | 500–2,500 |
| 5. | Kamara | <i>Hardwickia binata</i> | 25 -35 | 400 - 1000 |
| 6. | Bage | <i>Albezzia lebbek</i> | 20 - 45 | 500 - 1000 |
| 7. | Ficus | <i>Ficus bengalensis</i> | 20 - 50 | 500–2,500 |
| 8. | Sisso | <i>Dalbargia Sissoo</i> | 20 - 50 | 500 -2000 |
| 9. | Ailanthus | <i>Ailanthus excelsa</i> | 20 - 50 | 500 - 1000 |
| 10. | Hale | <i>Wrightia tinctoria</i> | 25 - 45 | 500 - 1000 |
| 11. | Uded | <i>Steriospermum chelanoides</i> | 25 - 45 | 500 -2000 |
| 12. | Dhupa | <i>Boswella Serrata</i> | 20 - 40 | 500 - 2000 |
| 13. | Nelli | <i>Embllica Officinalis</i> | 20 - 50 | 500 -1500 |
| 14. | Honne | <i>Pterocarpus marsupium</i> | 20 - 40 | 500 - 2000 |
| Moist Deciduous Species | | | Temp (°C) | Rainfall (mm) |
| 15. | Teak | <i>Tectona grandis</i> | 20 - 50 | 500-5000 |
| 16. | Nandi | <i>Legarstroemia lanceolata</i> | 20 - 40 | 500 - 4000 |
| 17. | Honne | <i>Pterocarpus marsupium</i> | 20 - 40 | 500 - 3000 |
| 18. | Mathi | <i>Terminalia alata</i> | 20 -50 | 500 - 2000 |
| 19. | Shivane | <i>Gmelina arborea</i> | 20 -50 | 500 -2000 |
| 20. | Kindal | <i>T.Paniculata</i> | 20 - 40 | 500 - 1500 |
| 21. | Beete | <i>Dalbargia latifolia</i> | 20 - 40 | 500 - 1500 |
| 22. | Tare | <i>T. belerica</i> | 20 - 40 | 500 - 2000 |
| 23. | Bamboo | <i>Bambusa arundinasia</i> | 20 - 40 | 500 - 2500 |
| 24. | Bamboo | <i>Dendrocalamus strictus</i> | 20 – 40 | 500 – 2500 |
| 25. | Muthuga | <i>Butea monosperma</i> | 20 - 40 | 400 - 1500 |
| 26. | Hippe | <i>Madhuca latifolia</i> | 20 - 40 | 500 - 2000 |
| 27. | Sandal | <i>Santalum album</i> | 20 - 50 | 400 - 1000 |
| 28. | Nelli | <i>Embllica officinalis</i> | 20 - 40 | 500 - 2000 |
| 29. | Nerale | <i>Sizygium cumini</i> | 20 - 40 | 500 - 2000 |
| 30. | Dhaman | <i>Grevia tilifolia</i> | 20 - 40 | 500 - 2000 |
| 31. | Kaval | <i>Careya arborea</i> | 20 - 40 | 500 - 2000 |
| 32. | Harada | <i>Terminalia chebula</i> | 20 - 40 | 500 - 2000 |

References

1. FAO (1976) Framework for Land Evaluation, Food and Agriculture Organization, Rome.72 pp.
2. FAO (1983) Guidelines for Land Evaluation for Rainfed Agriculture, FAO, Rome, 237 pp.
3. IARI (1971) Soil Survey Manual, All India Soil and Land Use Survey Organization, IARI, New Delhi, 121 pp.
4. Katyal, J.C. and Rattan, R.K. (2003) Secondary and Micronutrients; Research Gap and future needs. Fert. News 48 (4); 9-20.
5. Naidu, L.G.K., Ramamurthy, V., Challa, O., Hegde, R. and Krishnan, P. (2006) Manual Soil Site Suitability Criteria for Major Crops, NBSS Publ. No. 129, NBSS &LUP, Nagpur, 118 pp.
6. Natarajan, A. and Dipak Sarkar (2010) Field Guide for Soil Survey, National Bureau of Soil Survey and Land Use Planning (ICAR), Nagpur, India.
7. Natarajan, A., Rajendra Hegde, Raj, J.N. and Shivananda Murthy, H.G. (2015) Implementation Manual for Sujala-III Project, Watershed Development Department, Bengaluru, Karnataka.
8. Sarma, V.A.K., Krishnan, P. and Budihal, S.L. (1987) Laboratory Manual, Tech. Bull. 23, NBSS &LUP, Nagpur.
9. Sehgal, J.L. (1990) Soil Resource Mapping of Different States of India; Why and How? National Bureau of Soil Survey and Land Use Planning, Nagpur, 49 pp.
10. Shivaprasad, C.R., R.S. Reddy, J. Sehgal and M. Velayuthum (1998) Soils of Karnataka for Optimizing Land Use, NBSS Publ. No. 47b, NBSS & LUP, Nagpur, India.
11. Soil Survey Staff (2006) Keys to Soil Taxonomy, Tenth edition, U.S. Department of Agriculture/ NRCS, Washington DC, U.S.A.
12. Soil Survey Staff (2012) Soil Survey Manual, Handbook No. 18, USDA, Washington DC, USA.

Appendix I
Haligeri-3 (IT1a) Microwatershed
Soil Phase Information

| Village | Survey No | Area (ha) | Soil Phase | LMU | Soil Depth | Surface Soil Texture | Soil Gravelliness | Available Water Capacity | Slope | Soil Erosion | Current Land Use | WELLS | Land Capability | Conservation Plan |
|----------|-----------|-----------|------------|-------|-------------------------------|----------------------|------------------------|--------------------------|----------------------------|--------------|---|---------------|-----------------|--------------------|
| Halageri | 54 | 6.17 | NGPiB1 | LMU-2 | Deep (100-150 cm) | Sandy clay | Non gravelly (<15%) | Low (51-100 mm/m) | Very gently sloping (1-3%) | Slight | Clusterbean+Maize+Jowar+Current Fallow(Cb+Mz+Jw+Cf) | Not Available | IIs | Trench cum bunding |
| Halageri | 55 | 8.66 | BPRiB2 | LMU-2 | Deep (100-150 cm) | Sandy clay | Non gravelly (<15%) | Low (51-100 mm/m) | Very gently sloping (1-3%) | Moderate | Cluster bean+Sugarcane(Cb+S c) | Not Available | IIs | Trench cum bunding |
| Halageri | 56 | 11.06 | BPRiB2 | LMU-2 | Deep (100-150 cm) | Sandy clay | Non gravelly (<15%) | Low (51-100 mm/m) | Very gently sloping (1-3%) | Moderate | Cotton+Sugarcane (Ct+Sc) | Not Available | IIs | Trench cum bunding |
| Halageri | 60 | 0.01 | MTLmB2g1 | LMU-8 | Shallow (25-50 cm) | Clay | Gravelly (15-35%) | Low (51-100 mm/m) | Very gently sloping (1-3%) | Moderate | Not Available (NA) | Not Available | IIIs | Graded bunding |
| Halageri | 61 | 8.65 | MTLmB2g1 | LMU-8 | Shallow (25-50 cm) | Clay | Gravelly (15-35%) | Low (51-100 mm/m) | Very gently sloping (1-3%) | Moderate | Bengalgram+Current Fallow (Bg+Cf) | 1 Bore well | IIIs | Graded bunding |
| Halageri | 62 | 7.96 | RNKmB2 | LMU-6 | Moderately shallow (50-75 cm) | Clay | Non gravelly (<15%) | Medium (101-150 mm/m) | Very gently sloping (1-3%) | Moderate | Safflower (Sa) | Not Available | IVes | Graded bunding |
| Halageri | 63 | 10.07 | HDHiB1 | LMU-2 | Moderately deep (75-100 cm) | Sandy clay | Non gravelly (<15%) | Low (51-100 mm/m) | Very gently sloping (1-3%) | Slight | Bengalgram+Current Fallow (Bg+Cf) | Not Available | IIs | Trench cum bunding |
| Halageri | 64 | 10 | HDHiB1 | LMU-2 | Moderately deep (75-100 cm) | Sandy clay | Non gravelly (<15%) | Low (51-100 mm/m) | Very gently sloping (1-3%) | Slight | Maize+Current fallow (Mz+Cf) | Not Available | IIs | Trench cum bunding |
| Halageri | 65 | 8.9 | NGPhA1 | LMU-2 | Deep (100-150 cm) | Sandy clay loam | Non gravelly (<15%) | Low (51-100 mm/m) | Nearly level (0-1%) | Slight | Sunflower+Onion (Sf+On) | Not Available | IIs | Graded bunding |
| Halageri | 66 | 6.13 | RNKmB2 | LMU-6 | Moderately shallow (50-75 cm) | Clay | Non gravelly (<15%) | Medium (101-150 mm/m) | Very gently sloping (1-3%) | Moderate | Current Fallow (Cf) | Not Available | IVes | Graded bunding |
| Halageri | 67 | 6.26 | DRLmB2 | LMU-1 | Moderately deep (75-100 cm) | Clay | Non gravelly (<15%) | Medium (101-150 mm/m) | Very gently sloping (1-3%) | Moderate | Bengalgram+Maize (Bg+Mz) | Not Available | IIs | Graded bunding |
| Halageri | 68 | 9.59 | DRLmB2 | LMU-1 | Moderately deep (75-100 cm) | Clay | Non gravelly (<15%) | Medium (101-150 mm/m) | Very gently sloping (1-3%) | Moderate | Bengalgram+Maize (Bg+Mz) | Not Available | IIs | Graded bunding |
| Halageri | 69 | 8.07 | RNKmB2 | LMU-6 | Moderately shallow (50-75 cm) | Clay | Non gravelly (<15%) | Medium (101-150 mm/m) | Very gently sloping (1-3%) | Moderate | Bengalgram+Safflower (Bg+Sa) | Not Available | IVes | Graded bunding |
| Halageri | 70 | 6.45 | MTLmB2g1 | LMU-8 | Shallow (25-50 cm) | Clay | Gravelly (15-35%) | Low (51-100 mm/m) | Very gently sloping (1-3%) | Moderate | Bengalgram+Jowar (Bg+Jw) | Not Available | IIIs | Graded bunding |
| Halageri | 71 | 7.97 | MTLmB2g1 | LMU-8 | Shallow (25-50 cm) | Clay | Gravelly (15-35%) | Low (51-100 mm/m) | Very gently sloping (1-3%) | Moderate | Jowar+Cotton+Bengalgram (Jw+Ct+Bg) | Not Available | IIIs | Graded bunding |
| Halageri | 72 | 0.04 | MTLmB2g1 | LMU-8 | Shallow (25-50 cm) | Clay | Gravelly (15-35%) | Low (51-100 mm/m) | Very gently sloping (1-3%) | Moderate | Not Available (NA) | Not Available | IIIs | Graded bunding |
| Halageri | 73 | 1.02 | MTLmB2g1 | LMU-8 | Shallow (25-50 cm) | Clay | Gravelly (15-35%) | Low (51-100 mm/m) | Very gently sloping (1-3%) | Moderate | Jowar (Jw) | Not Available | IIIs | Graded bunding |
| Halageri | 74 | 3.3 | MTLmB2g1 | LMU-8 | Shallow (25-50 cm) | Clay | Gravelly (15-35%) | Low (51-100 mm/m) | Very gently sloping (1-3%) | Moderate | Bengalgram (Bg) | Not Available | IIIs | Graded bunding |
| Halageri | 75 | 1.95 | MTLmB2g1 | LMU-8 | Shallow (25-50 cm) | Clay | Gravelly (15-35%) | Low (51-100 mm/m) | Very gently sloping (1-3%) | Moderate | Sunflower (Sf) | Not Available | IIIs | Graded bunding |
| Halageri | 76 | 0.56 | MTLmB2g1 | LMU-8 | Shallow (25-50 cm) | Clay | Gravelly (15-35%) | Low (51-100 mm/m) | Very gently sloping (1-3%) | Moderate | Current Fallow (Cf) | Not Available | IIIs | Graded bunding |
| Halageri | 356 | 0.19 | BSRiB1g2 | LMU-4 | Moderately deep (75-100 cm) | Sandy clay | Very gravelly (35-60%) | Low (51-100 mm/m) | Very gently sloping (1-3%) | Slight | Not Available (NA) | Not Available | IIs | Trench cum bunding |

| Village | Survey No | Area (ha) | Soil Phase | LMU | Soil Depth | Surface Soil Texture | Soil Gravelliness | Available Water Capacity | Slope | Soil Erosion | Current Land Use | WELLS | Land Capability | Conservation Plan |
|------------|-----------|-----------|------------|-------|-----------------------------|----------------------|------------------------|--------------------------|----------------------------|--------------|--|---------------|-----------------|--------------------|
| Halageri | 357 | 1.6 | BSRiB1g2 | LMU-4 | Moderately deep (75-100 cm) | Sandy clay | Very gravelly (35-60%) | Low (51-100 mm/m) | Very gently sloping (1-3%) | Slight | Current Fallow (Cf) | Not Available | IIs | Trench cum bunding |
| Halageri | 358 | 4.85 | NGPiB1 | LMU-2 | Deep (100-150 cm) | Sandy clay | Non gravelly (<15%) | Low (51-100 mm/m) | Very gently sloping (1-3%) | Slight | Bajra+Pigeon Pea+Cluster bean (Bj+Pp+Cb) | 1 Bore well | IIs | Trench cum bunding |
| Halageri | 359 | 3.48 | BSRiB1g2 | LMU-4 | Moderately deep (75-100 cm) | Sandy clay | Very gravelly (35-60%) | Low (51-100 mm/m) | Very gently sloping (1-3%) | Slight | Cluster bean (Cb) | Not Available | IIs | Trench cum bunding |
| Thalabala | 187 | 0.23 | DRLmB2g1 | LMU-1 | Moderately deep (75-100 cm) | Clay | Gravelly (15-35%) | Medium (101-150 mm/m) | Very gently sloping (1-3%) | Moderate | Bengalgram (Bg) | Not Available | IIs | Graded bunding |
| Thalakalla | 98 | 2.2 | LGDmB1 | LMU-1 | Deep (100-150 cm) | Clay | Non gravelly (<15%) | Very high (>200 mm/m) | Very gently sloping (1-3%) | Slight | Current Fallow (Cf) | Not Available | IIs | Graded bunding |
| Thalakalla | 99 | 8.58 | LGDmB1 | LMU-1 | Deep (100-150 cm) | Clay | Non gravelly (<15%) | Very high (>200 mm/m) | Very gently sloping (1-3%) | Slight | Current Fallow (Cf) | Not Available | IIs | Graded bunding |
| Thalakalla | 100 | 8.44 | LGDmB1 | LMU-1 | Deep (100-150 cm) | Clay | Non gravelly (<15%) | Very high (>200 mm/m) | Very gently sloping (1-3%) | Slight | CurrentFallow+Bengalgram (Cf+Bg) | Not Available | IIs | Graded bunding |
| Thalakalla | 101 | 2.74 | LGDmB1 | LMU-1 | Deep (100-150 cm) | Clay | Non gravelly (<15%) | Very high (>200 mm/m) | Very gently sloping (1-3%) | Slight | Onion (On) | 1 Bore well | IIs | Graded bunding |
| Thalakalla | 102 | 3.13 | BPRmB2 | LMU-2 | Deep (100-150 cm) | Clay | Non gravelly (<15%) | Low (51-100 mm/m) | Very gently sloping (1-3%) | Moderate | Onion+Maize (On+Mz) | 1 Bore well | IIs | Trench cum bunding |
| Thalakalla | 103 | 3.7 | LGDmB1 | LMU-1 | Deep (100-150 cm) | Clay | Non gravelly (<15%) | Very high (>200 mm/m) | Very gently sloping (1-3%) | Slight | Jowar (Jw) | Not Available | IIs | Graded bunding |
| Thalakalla | 104 | 2.83 | LGDmB1 | LMU-1 | Deep (100-150 cm) | Clay | Non gravelly (<15%) | Very high (>200 mm/m) | Very gently sloping (1-3%) | Slight | Current Fallow (Cf) | Not Available | IIs | Graded bunding |
| Thalakalla | 105 | 10.06 | LGDmB1 | LMU-1 | Deep (100-150 cm) | Clay | Non gravelly (<15%) | Very high (>200 mm/m) | Very gently sloping (1-3%) | Slight | Current Fallow+Jowar (Cf+Jw) | Not Available | IIs | Graded bunding |
| Thalakalla | 106 | 1.2 | LGDmB1 | LMU-1 | Deep (100-150 cm) | Clay | Non gravelly (<15%) | Very high (>200 mm/m) | Very gently sloping (1-3%) | Slight | Not Available (NA) | Not Available | IIs | Graded bunding |
| Thalakalla | 107 | 1.54 | DRLmB2 | LMU-1 | Moderately deep (75-100 cm) | Clay | Non gravelly (<15%) | Medium (101-150 mm/m) | Very gently sloping (1-3%) | Moderate | Not Available (NA) | Not Available | IIs | Graded bunding |
| Thalakalla | 108 | 3.02 | LGDmB1 | LMU-1 | Deep (100-150 cm) | Clay | Non gravelly (<15%) | Very high (>200 mm/m) | Very gently sloping (1-3%) | Slight | Jowar (Jw) | Not Available | IIs | Graded bunding |
| Thalakalla | 109 | 4.14 | LGDmB1 | LMU-1 | Deep (100-150 cm) | Clay | Non gravelly (<15%) | Very high (>200 mm/m) | Very gently sloping (1-3%) | Slight | Current Fallow (Cf) | Not Available | IIs | Graded bunding |
| Thalakalla | 110 | 4.11 | LGDmB1 | LMU-1 | Deep (100-150 cm) | Clay | Non gravelly (<15%) | Very high (>200 mm/m) | Very gently sloping (1-3%) | Slight | Current Fallow (Cf) | Not Available | IIs | Graded bunding |
| Thalakalla | 111 | 2.98 | LGDmB1 | LMU-1 | Deep (100-150 cm) | Clay | Non gravelly (<15%) | Very high (>200 mm/m) | Very gently sloping (1-3%) | Slight | Jowar (Jw) | Not Available | IIs | Graded bunding |
| Thalakalla | 112 | 6.47 | CSRiB2 | LMU-9 | Shallow (25-50 cm) | Sandy clay | Non gravelly (<15%) | Very Low (<50 mm/m) | Very gently sloping (1-3%) | Moderate | Current Fallow+Mango (Cf+Mn) | Not Available | IIIs | Trench cum bunding |
| Thalakalla | 113 | 4.64 | GDPiB2 | LMU-2 | Deep (100-150 cm) | Sandy clay | Non gravelly (<15%) | Low (51-100 mm/m) | Very gently sloping (1-3%) | Moderate | Current Fallow (Cf) | Not Available | IIs | Trench cum bunding |
| Thalakalla | 114 | 8.68 | CSRiB2 | LMU-9 | Shallow (25-50 cm) | Sandy clay | Non gravelly (<15%) | Very Low (<50 mm/m) | Very gently sloping (1-3%) | Moderate | Current Fallow (Cf) | Not Available | IIIs | Trench cum bunding |
| Thalakalla | 115 | 7.05 | NGPiB1 | LMU-2 | Deep (100-150 cm) | Sandy clay | Non gravelly (<15%) | Low (51-100 mm/m) | Very gently sloping (1-3%) | Slight | Current Fallow+Fallow land (Cf+Fl) | Not Available | IIs | Trench cum bunding |
| Thalakalla | 116 | 4.27 | NGPiB1 | LMU-2 | Deep (100-150 cm) | Sandy clay | Non gravelly (<15%) | Low (51-100 mm/m) | Very gently sloping (1-3%) | Slight | Current Fallow (Cf) | Not Available | IIs | Trench cum bunding |
| Thalakalla | 124 | 0.88 | AWDmB2 | LMU-1 | Very deep (>150 cm) | Clay | Non gravelly (<15%) | Very high (>200 mm/m) | Very gently sloping (1-3%) | Moderate | Jowar (Jw) | Not Available | IIs | Graded bunding |

| Village | Survey No | Area (ha) | Soil Phase | LMU | Soil Depth | Surface Soil Texture | Soil Gravelliness | Available Water Capacity | Slope | Soil Erosion | Current Land Use | WELLS | Land Capability | Conservation Plan |
|-------------|-----------|-----------|------------|-------|-------------------------------|----------------------|---------------------|--------------------------|----------------------------|--------------|----------------------------------|---------------|-----------------|--------------------|
| Thalaka lla | 125 | 0.62 | AWDmB 2 | LMU-1 | Very deep (>150 cm) | Clay | Non gravelly (<15%) | Very high (>200 mm/m) | Very gently sloping (1-3%) | Moderate | Bengalgram (Bg) | Not Available | Iles | Graded bunding |
| Thalaka lla | 126 | 0.05 | AWDmB 2 | LMU-1 | Very deep (>150 cm) | Clay | Non gravelly (<15%) | Very high (>200 mm/m) | Very gently sloping (1-3%) | Moderate | Not Available (NA) | Not Available | Iles | Graded bunding |
| Thalaka lla | 127 | 0.00 | AWDmB 2 | LMU-1 | Very deep (>150 cm) | Clay | Non gravelly (<15%) | Very high (>200 mm/m) | Very gently sloping (1-3%) | Moderate | Not Available (NA) | Not Available | Iles | Graded bunding |
| Thalaka lla | 128 | 1.82 | AWDmB 2 | LMU-1 | Very deep (>150 cm) | Clay | Non gravelly (<15%) | Very high (>200 mm/m) | Very gently sloping (1-3%) | Moderate | Bengalgram (Bg) | Not Available | Iles | Graded bunding |
| Thalaka lla | 129 | 0.71 | KVRmA1 | LMU-1 | Deep (100-150 cm) | Clay | Non gravelly (<15%) | Very high (>200 mm/m) | Nearly level (0-1%) | Slight | Bengalgram (Bg) | Not Available | Ils | Graded bunding |
| Thalaka lla | 130 | 2 | KVRmA1 | LMU-1 | Deep (100-150 cm) | Clay | Non gravelly (<15%) | Very high (>200 mm/m) | Nearly level (0-1%) | Slight | Jowar (Jw) | Not Available | Ils | Graded bunding |
| Thalaka lla | 131 | 1.52 | KVRmA1 | LMU-1 | Deep (100-150 cm) | Clay | Non gravelly (<15%) | Very high (>200 mm/m) | Nearly level (0-1%) | Slight | Jowar (Jw) | Not Available | Ils | Graded bunding |
| Thalaka lla | 132 | 0.00 | KVRmA1 | LMU-1 | Deep (100-150 cm) | Clay | Non gravelly (<15%) | Very high (>200 mm/m) | Nearly level (0-1%) | Slight | Not Available (NA) | Not Available | Ils | Graded bunding |
| Veerapura | 186 | 0.76 | DRLmB2 | LMU-1 | Moderately deep (75-100 cm) | Clay | Non gravelly (<15%) | Medium (101-150 mm/m) | Very gently sloping (1-3%) | Moderate | Bengalgram (Bg) | Not Available | Iles | Graded bunding |
| Veerapura | 187 | 2.15 | DRLmB2 | LMU-1 | Moderately deep (75-100 cm) | Clay | Non gravelly (<15%) | Medium (101-150 mm/m) | Very gently sloping (1-3%) | Moderate | Jowar (Jw) | Not Available | Iles | Graded bunding |
| Veerapura | 188 | 3.58 | DRLmB2 | LMU-1 | Moderately deep (75-100 cm) | Clay | Non gravelly (<15%) | Medium (101-150 mm/m) | Very gently sloping (1-3%) | Moderate | Jowar (Jw) | Not Available | Iles | Graded bunding |
| Banapura | 36 | 3.84 | DRLmB2g1 | LMU-1 | Moderately deep (75-100 cm) | Clay | Gravelly (15-35%) | Medium (101-150 mm/m) | Very gently sloping (1-3%) | Moderate | Bengalgram+Maize (Bg+Mz) | Not Available | Iles | Graded bunding |
| Banapura | 37 | 3.69 | MKHiB2g1 | LMU-7 | Moderately shallow (50-75 cm) | Sandy clay | Gravelly (15-35%) | Very Low (<50 mm/m) | Very gently sloping (1-3%) | Moderate | Jowar+Maize (Jw+Mz) | Not Available | IIles | Trench cum bunding |
| Banapura | 38 | 3.91 | MKHiB2g1 | LMU-7 | Moderately shallow (50-75 cm) | Sandy clay | Gravelly (15-35%) | Very Low (<50 mm/m) | Very gently sloping (1-3%) | Moderate | Bengalgram+Jowar (Bg+Jw) | Not Available | IIles | Trench cum bunding |
| Banapura | 40 | 4.45 | MKHiB2g1 | LMU-7 | Moderately shallow (50-75 cm) | Sandy clay | Gravelly (15-35%) | Very Low (<50 mm/m) | Very gently sloping (1-3%) | Moderate | Jowar (Jw) | Not Available | IIles | Trench cum bunding |
| Banapura | 41 | 1.62 | LKRCB2g1 | LMU-7 | Moderately shallow (50-75 cm) | Sandy loam | Gravelly (15-35%) | Very Low (<50 mm/m) | Very gently sloping (1-3%) | Moderate | Bengalgram (Bg) | Not Available | IIles | Trench cum bunding |
| Banapura | 42 | 2.64 | LKRCB2g1 | LMU-7 | Moderately shallow (50-75 cm) | Sandy loam | Gravelly (15-35%) | Very Low (<50 mm/m) | Very gently sloping (1-3%) | Moderate | Current Fallow+Groundnut (Cf+Gn) | Not Available | IIles | Trench cum bunding |
| Banapura | 43 | 6.45 | LKRCB2g1 | LMU-7 | Moderately shallow (50-75 cm) | Sandy loam | Gravelly (15-35%) | Very Low (<50 mm/m) | Very gently sloping (1-3%) | Moderate | Current Fallow+Groundnut (Cf+Gn) | Not Available | IIles | Trench cum bunding |
| Banapura | 44 | 4.75 | HDHhB2g1 | LMU-2 | Moderately deep (75-100 cm) | Sandy clay loam | Gravelly (15-35%) | Low (51-100 mm/m) | Very gently sloping (1-3%) | Moderate | Bajra+Cotton (Bj+Ct) | Not Available | Iles | Trench cum bunding |
| Banapura | 45 | 8.35 | LKRCB2g1 | LMU-7 | Moderately shallow (50-75 cm) | Sandy loam | Gravelly (15-35%) | Very Low (<50 mm/m) | Very gently sloping (1-3%) | Moderate | Fallow land+Mango (Fl+Mn) | 1 Bore well | IIles | Trench cum bunding |
| Banapura | 46 | 0.45 | GDPcB2 | LMU-2 | Deep (100-150 cm) | Sandy loam | Non gravelly (<15%) | Low (51-100 mm/m) | Very gently sloping (1-3%) | Moderate | Horsegram (Hg) | Not Available | Iles | Trench cum bunding |
| Banapura | 47 | 0.85 | LKRCB2g1 | LMU-7 | Moderately shallow (50-75 cm) | Sandy loam | Gravelly (15-35%) | Very Low (<50 mm/m) | Very gently sloping (1-3%) | Moderate | Fallow land (Fl) | Not Available | IIles | Trench cum bunding |
| Banapura | 54(1) | 0.03 | GDPcB2 | LMU-2 | Deep (100-150 cm) | Sandy loam | Non gravelly (<15%) | Low (51-100 mm/m) | Very gently sloping (1-3%) | Moderate | Not Available (NA) | Not Available | Iles | Trench cum bunding |

| Village | Survey No | Area (ha) | Soil Phase | LMU | Soil Depth | Surface Soil Texture | Soil Gravelliness | Available Water Capacity | Slope | Soil Erosion | Current Land Use | WELLS | Land Capability | Conservation Plan |
|----------|-----------|-----------|------------|-------|-------------------------------|----------------------|---------------------|--------------------------|----------------------------|--------------|--|---------------|-----------------|--------------------|
| Banapura | 56 | 0.21 | BPRcA1 | LMU-2 | Deep (100-150 cm) | Sandy loam | Non gravelly (<15%) | Low (51-100 mm/m) | Nearly level (0-1%) | Slight | Not Available (NA) | Not Available | IIs | Graded bunding |
| Banapura | 57 | 0.57 | BPRcA1 | LMU-2 | Deep (100-150 cm) | Sandy loam | Non gravelly (<15%) | Low (51-100 mm/m) | Nearly level (0-1%) | Slight | Not Available (NA) | Not Available | IIs | Graded bunding |
| Banapura | 58 | 5.3 | HDHhB2g1 | LMU-2 | Moderately deep (75-100 cm) | Sandy clay loam | Gravelly (15-35%) | Low (51-100 mm/m) | Very gently sloping (1-3%) | Moderate | Bajra (Bj) | Not Available | IIs | Trench cum bunding |
| Banapura | 59 | 6.36 | HDHhB2g1 | LMU-2 | Moderately deep (75-100 cm) | Sandy clay loam | Gravelly (15-35%) | Low (51-100 mm/m) | Very gently sloping (1-3%) | Moderate | Current Fallow (Cf) | Not Available | IIs | Trench cum bunding |
| Banapura | 60 | 6.97 | HDHcB1g1 | LMU-2 | Moderately deep (75-100 cm) | Sandy loam | Gravelly (15-35%) | Low (51-100 mm/m) | Very gently sloping (1-3%) | Slight | Current Fallow+Pigeon Pea (Cf+Pp) | Not Available | IIs | Trench cum bunding |
| Banapura | 61 | 8.98 | HDHcB1g1 | LMU-2 | Moderately deep (75-100 cm) | Sandy loam | Gravelly (15-35%) | Low (51-100 mm/m) | Very gently sloping (1-3%) | Slight | Current Fallow+Pigeon Pea (Cf+Pp) | Not Available | IIs | Trench cum bunding |
| Banapura | 62 | 1.29 | RNKmB2g1 | LMU-6 | Moderately shallow (50-75 cm) | Clay | Gravelly (15-35%) | Medium (101-150 mm/m) | Very gently sloping (1-3%) | Moderate | Not Available (NA) | Not Available | IVes | Graded bunding |
| Banapura | 63 | 2.33 | RNKmB2g1 | LMU-6 | Moderately shallow (50-75 cm) | Clay | Gravelly (15-35%) | Medium (101-150 mm/m) | Very gently sloping (1-3%) | Moderate | Jowar (Jw) | Not Available | IVes | Graded bunding |
| Banapura | 64 | 2.77 | RNKmB2g1 | LMU-6 | Moderately shallow (50-75 cm) | Clay | Gravelly (15-35%) | Medium (101-150 mm/m) | Very gently sloping (1-3%) | Moderate | Bengalgram (Bg) | 2 Bore well | IVes | Graded bunding |
| Banapura | 65 | 2.44 | RNKmB2g1 | LMU-6 | Moderately shallow (50-75 cm) | Clay | Gravelly (15-35%) | Medium (101-150 mm/m) | Very gently sloping (1-3%) | Moderate | Jowar (Jw) | 1 Bore well | IVes | Graded bunding |
| Banapura | 66 | 6.2 | DRLmB2 | LMU-1 | Moderately deep (75-100 cm) | Clay | Non gravelly (<15%) | Medium (101-150 mm/m) | Very gently sloping (1-3%) | Moderate | Bengalgram+Jowar (Bg+Jw) | Not Available | IIs | Graded bunding |
| Banapura | 67 | 2.04 | DRLmB2 | LMU-1 | Moderately deep (75-100 cm) | Clay | Non gravelly (<15%) | Medium (101-150 mm/m) | Very gently sloping (1-3%) | Moderate | Bengalgram (Bg) | Not Available | IIs | Graded bunding |
| Banapura | 68 | 1.11 | RNKmB2g1 | LMU-6 | Moderately shallow (50-75 cm) | Clay | Gravelly (15-35%) | Medium (101-150 mm/m) | Very gently sloping (1-3%) | Moderate | Current Fallow (Cf) | Not Available | IVes | Graded bunding |
| Banapura | 69 | 9.8 | RNKmB2g1 | LMU-6 | Moderately shallow (50-75 cm) | Clay | Gravelly (15-35%) | Medium (101-150 mm/m) | Very gently sloping (1-3%) | Moderate | Bengalgram+Maize+Current Fallow (Bg+Mz+Cf) | Not Available | IVes | Graded bunding |
| Banapura | 70 | 7.73 | MKHhB2g1 | LMU-7 | Moderately shallow (50-75 cm) | Sandy clay | Gravelly (15-35%) | Very Low (<50 mm/m) | Very gently sloping (1-3%) | Moderate | Redgram+Cotton+Maize (Rg+Ct+Mz) | Not Available | IIIs | Trench cum bunding |
| Banapura | 71 | 7.93 | MKHhB2g1 | LMU-7 | Moderately shallow (50-75 cm) | Sandy clay | Gravelly (15-35%) | Very Low (<50 mm/m) | Very gently sloping (1-3%) | Moderate | Bengalgram+Fallow land (Bg+Fl) | Not Available | IIIs | Trench cum bunding |
| Banapura | 72 | 4.63 | MKHhB2g1 | LMU-7 | Moderately shallow (50-75 cm) | Sandy clay | Gravelly (15-35%) | Very Low (<50 mm/m) | Very gently sloping (1-3%) | Moderate | Bengalgram+Cotton (Bg+Ct) | Not Available | IIIs | Trench cum bunding |
| Banapura | 73 | 5 | MKHhB2g1 | LMU-7 | Moderately shallow (50-75 cm) | Sandy clay | Gravelly (15-35%) | Very Low (<50 mm/m) | Very gently sloping (1-3%) | Moderate | Bengalgram+Jowar (Bg+Jw) | Not Available | IIIs | Trench cum bunding |
| Banapura | 74 | 6.96 | DRLmB2g1 | LMU-1 | Moderately deep (75-100 cm) | Clay | Gravelly (15-35%) | Medium (101-150 mm/m) | Very gently sloping (1-3%) | Moderate | Current Fallow+Jowar (Cf+Jw) | Not Available | IIs | Graded bunding |
| Banapura | 75 | 8.45 | DRLmB2g1 | LMU-1 | Moderately deep (75-100 cm) | Clay | Gravelly (15-35%) | Medium (101-150 mm/m) | Very gently sloping (1-3%) | Moderate | Bengalgram (Bg) | Not Available | IIs | Graded bunding |
| Banapura | 76 | 4.01 | DRLmB2g1 | LMU-1 | Moderately deep (75-100 cm) | Clay | Gravelly (15-35%) | Medium (101-150 mm/m) | Very gently sloping (1-3%) | Moderate | Bengalgram (Bg) | Not Available | IIs | Graded bunding |
| Banapura | 95 | 0.83 | DRLmB2g1 | LMU-1 | Moderately deep (75-100 cm) | Clay | Gravelly (15-35%) | Medium (101-150 mm/m) | Very gently sloping (1-3%) | Moderate | Bengalgram (Bg) | Not Available | IIs | Graded bunding |
| Banapura | 96 | 3.62 | DRLmB2g1 | LMU-1 | Moderately deep (75-100 cm) | Clay | Gravelly (15-35%) | Medium (101-150 mm/m) | Very gently sloping (1-3%) | Moderate | Bengalgram (Bg) | Not Available | IIs | Graded bunding |
| Banapura | 97 | 6.65 | DRLmB2g1 | LMU-1 | Moderately deep (75-100 cm) | Clay | Gravelly (15-35%) | Medium (101-150 mm/m) | Very gently sloping (1-3%) | Moderate | Bengalgram+Jowar (Bg+Jw) | Not Available | IIs | Graded bunding |

| Village | Survey No | Area (ha) | Soil Phase | LMU | Soil Depth | Surface Soil Texture | Soil Graveliness | Available Water Capacity | Slope | Soil Erosion | Current Land Use | WELLS | Land Capability | Conservation Plan |
|----------|-----------|-----------|------------|-------|-----------------------------|----------------------|---------------------|--------------------------|----------------------------|--------------|-----------------------------------|---------------|-----------------|-------------------|
| Banapura | 98 | 4.04 | DRLmB2g1 | LMU-1 | Moderately deep (75-100 cm) | Clay | Gravelly (15-35%) | Medium (101-150 mm/m) | Very gently sloping (1-3%) | Moderate | Bengalgram+Jowar (Bg+Jw) | Not Available | Iles | Graded bunding |
| Banapura | 99 | 8.83 | LGDmB1 | LMU-1 | Deep (100-150 cm) | Clay | Non gravelly (<15%) | Very high (>200 mm/m) | Very gently sloping (1-3%) | Slight | Bengalgram+Chilly (Bg+Ch) | Not Available | Ils | Graded bunding |
| Banapura | 100 | 3.44 | LGDmB1 | LMU-1 | Deep (100-150 cm) | Clay | Non gravelly (<15%) | Very high (>200 mm/m) | Very gently sloping (1-3%) | Slight | Jowar (Jw) | Not Available | Ils | Graded bunding |
| Banapura | 101 | 3.7 | LGDmB1 | LMU-1 | Deep (100-150 cm) | Clay | Non gravelly (<15%) | Very high (>200 mm/m) | Very gently sloping (1-3%) | Slight | Jowar (Jw) | Not Available | Ils | Graded bunding |
| Banapura | 102 | 0.4 | LGDmB1 | LMU-1 | Deep (100-150 cm) | Clay | Non gravelly (<15%) | Very high (>200 mm/m) | Very gently sloping (1-3%) | Slight | Not Available (NA) | Not Available | Ils | Graded bunding |
| Banapura | 107 | 4.56 | LGDmB1 | LMU-1 | Deep (100-150 cm) | Clay | Non gravelly (<15%) | Very high (>200 mm/m) | Very gently sloping (1-3%) | Slight | Current Fallow+Jowar (Cf+Jw) | Not Available | Ils | Graded bunding |
| Banapura | 108 | 8.21 | LGDmB1 | LMU-1 | Deep (100-150 cm) | Clay | Non gravelly (<15%) | Very high (>200 mm/m) | Very gently sloping (1-3%) | Slight | Bengalgram+Jowar (Bg+Jw) | Not Available | Ils | Graded bunding |
| Banapura | 109 | 4.53 | DRLmB2 | LMU-1 | Moderately deep (75-100 cm) | Clay | Non gravelly (<15%) | Medium (101-150 mm/m) | Very gently sloping (1-3%) | Moderate | Bengalgram+Current Fallow (Bg+Cf) | Not Available | Iles | Graded bunding |
| Banapura | 110 | 1.83 | DRLmB2 | LMU-1 | Moderately deep (75-100 cm) | Clay | Non gravelly (<15%) | Medium (101-150 mm/m) | Very gently sloping (1-3%) | Moderate | Bengalgram (Bg) | Not Available | Iles | Graded bunding |
| Banapura | 111 | 3.74 | DRLmB2 | LMU-1 | Moderately deep (75-100 cm) | Clay | Non gravelly (<15%) | Medium (101-150 mm/m) | Very gently sloping (1-3%) | Moderate | Bengalgram+Fallow land (Bg+Fl) | Not Available | Iles | Graded bunding |
| Banapura | 112 | 6.33 | LGDmB1 | LMU-1 | Deep (100-150 cm) | Clay | Non gravelly (<15%) | Very high (>200 mm/m) | Very gently sloping (1-3%) | Slight | Bengalgram+Jowar (Bg+Jw) | Not Available | Ils | Graded bunding |
| Banapura | 113 | 3.78 | LGDmB1 | LMU-1 | Deep (100-150 cm) | Clay | Non gravelly (<15%) | Very high (>200 mm/m) | Very gently sloping (1-3%) | Slight | Jowar (Jw) | Not Available | Ils | Graded bunding |
| Banapura | 115 | 1.71 | LGDmB1 | LMU-1 | Deep (100-150 cm) | Clay | Non gravelly (<15%) | Very high (>200 mm/m) | Very gently sloping (1-3%) | Slight | Bengalgram (Bg) | Not Available | Ils | Graded bunding |
| Banapura | 116 | 0.97 | LGDmB1 | LMU-1 | Deep (100-150 cm) | Clay | Non gravelly (<15%) | Very high (>200 mm/m) | Very gently sloping (1-3%) | Slight | Bengalgram (Bg) | Not Available | Ils | Graded bunding |
| Banapura | 117 | 1.01 | LGDmB1 | LMU-1 | Deep (100-150 cm) | Clay | Non gravelly (<15%) | Very high (>200 mm/m) | Very gently sloping (1-3%) | Slight | Bengalgram (Bg) | Not Available | Ils | Graded bunding |
| Banapura | 118 | 2.01 | LGDmB1 | LMU-1 | Deep (100-150 cm) | Clay | Non gravelly (<15%) | Very high (>200 mm/m) | Very gently sloping (1-3%) | Slight | Bengalgram (Bg) | Not Available | Ils | Graded bunding |
| Banapura | 119 | 1.74 | LGDmB1 | LMU-1 | Deep (100-150 cm) | Clay | Non gravelly (<15%) | Very high (>200 mm/m) | Very gently sloping (1-3%) | Slight | Bengalgram (Bg) | Not Available | Ils | Graded bunding |
| Banapura | 120 | 1.9 | LGDmB1 | LMU-1 | Deep (100-150 cm) | Clay | Non gravelly (<15%) | Very high (>200 mm/m) | Very gently sloping (1-3%) | Slight | Jowar (Jw) | Not Available | Ils | Graded bunding |
| Banapura | 121 | 1.08 | KVRmB1 | LMU-1 | Deep (100-150 cm) | Clay | Non gravelly (<15%) | Very high (>200 mm/m) | Very gently sloping (1-3%) | Slight | Jowar (Jw) | Not Available | Ils | Graded bunding |
| Banapura | 124 | 0.005 | KVRmB1 | LMU-1 | Deep (100-150 cm) | Clay | Non gravelly (<15%) | Very high (>200 mm/m) | Very gently sloping (1-3%) | Slight | Not Available (NA) | Not Available | Ils | Graded bunding |
| Banapura | 125 | 3.08 | KVRmB1 | LMU-1 | Deep (100-150 cm) | Clay | Non gravelly (<15%) | Very high (>200 mm/m) | Very gently sloping (1-3%) | Slight | Bengalgram (Bg) | Not Available | Ils | Graded bunding |
| Banapura | 126 | 6.94 | LGDmB1 | LMU-1 | Deep (100-150 cm) | Clay | Non gravelly (<15%) | Very high (>200 mm/m) | Very gently sloping (1-3%) | Slight | Bengalgram+Jowar (Bg+Jw) | Not Available | Ils | Graded bunding |
| Banapura | 127 | 5.53 | LGDmB1 | LMU-1 | Deep (100-150 cm) | Clay | Non gravelly (<15%) | Very high (>200 mm/m) | Very gently sloping (1-3%) | Slight | Bengalgram+Fallow land (Bg+Fl) | Not Available | Ils | Graded bunding |
| Banapura | 128 | 3.01 | DRLmB2 | LMU-1 | Moderately deep (75-100 cm) | Clay | Non gravelly (<15%) | Medium (101-150 mm/m) | Very gently sloping (1-3%) | Moderate | Bengalgram (Bg) | Not Available | Iles | Graded bunding |

| Village | Survey No | Area (ha) | Soil Phase | LMU | Soil Depth | Surface Soil Texture | Soil Gravelliness | Available Water Capacity | Slope | Soil Erosion | Current Land Use | WELLS | Land Capability | Conservation Plan |
|-----------|-----------|-----------|------------|-------|-------------------------------|----------------------|---------------------|--------------------------|----------------------------|--------------|--|---------------|-----------------|-------------------|
| Banapurra | 129 | 3.35 | DRLmB2 | LMU-1 | Moderately deep (75-100 cm) | Clay | Non gravelly (<15%) | Medium (101-150 mm/m) | Very gently sloping (1-3%) | Moderate | Bengalgram (Bg) | Not Available | Iles | Graded bunding |
| Banapurra | 130 | 5.89 | DRLmB2 | LMU-1 | Moderately deep (75-100 cm) | Clay | Non gravelly (<15%) | Medium (101-150 mm/m) | Very gently sloping (1-3%) | Moderate | Bengalgram (Bg) | Not Available | Iles | Graded bunding |
| Banapurra | 131 | 6.89 | DRLmB2 | LMU-1 | Moderately deep (75-100 cm) | Clay | Non gravelly (<15%) | Medium (101-150 mm/m) | Very gently sloping (1-3%) | Moderate | Jowar+Fallow land (Jw+Fl) | Not Available | Iles | Graded bunding |
| Banapurra | 132 | 6.18 | DRLmB2 | LMU-1 | Moderately deep (75-100 cm) | Clay | Non gravelly (<15%) | Medium (101-150 mm/m) | Very gently sloping (1-3%) | Moderate | Bengalgram+Jowar+Current Fallow (Bg+Jw+Cf) | Not Available | Iles | Graded bunding |
| Banapurra | 133 | 6.12 | DRLmB2 | LMU-1 | Moderately deep (75-100 cm) | Clay | Non gravelly (<15%) | Medium (101-150 mm/m) | Very gently sloping (1-3%) | Moderate | Jowar+Bengalgram+Fallowland (Jw+Bg+Fl) | Not Available | Iles | Graded bunding |
| Banapurra | 134 | 8.59 | DRLmB2 | LMU-1 | Moderately deep (75-100 cm) | Clay | Non gravelly (<15%) | Medium (101-150 mm/m) | Very gently sloping (1-3%) | Moderate | Bengalgram+Jowar (Bg+Jw) | Not Available | Iles | Graded bunding |
| Banapurra | 135 | 6.27 | KVRmB1 | LMU-1 | Deep (100-150 cm) | Clay | Non gravelly (<15%) | Very high (>200 mm/m) | Very gently sloping (1-3%) | Slight | Bengalgram (Bg) | Not Available | Ils | Graded bunding |
| Banapurra | 136 | 4.26 | KVRmB1 | LMU-1 | Deep (100-150 cm) | Clay | Non gravelly (<15%) | Very high (>200 mm/m) | Very gently sloping (1-3%) | Slight | Bengalgram (Bg) | Not Available | Ils | Graded bunding |
| Banapurra | 137 | 7.56 | DRLmB2 | LMU-1 | Moderately deep (75-100 cm) | Clay | Non gravelly (<15%) | Medium (101-150 mm/m) | Very gently sloping (1-3%) | Moderate | Bengalgram (Bg) | Not Available | Iles | Graded bunding |
| Banapurra | 138 | 6.39 | DRLmB2 | LMU-1 | Moderately deep (75-100 cm) | Clay | Non gravelly (<15%) | Medium (101-150 mm/m) | Very gently sloping (1-3%) | Moderate | Bengalgram (Bg) | Not Available | Iles | Graded bunding |
| Banapurra | 139 | 9.63 | RNKmB2g1 | LMU-6 | Moderately shallow (50-75 cm) | Clay | Gravelly (15-35%) | Medium (101-150 mm/m) | Very gently sloping (1-3%) | Moderate | Current Fallow+Jowar+Chilly (Cf+Jw+Ch) | Not Available | IVes | Graded bunding |
| Banapurra | 140 | 6.23 | DRLmB2 | LMU-1 | Moderately deep (75-100 cm) | Clay | Non gravelly (<15%) | Medium (101-150 mm/m) | Very gently sloping (1-3%) | Moderate | Bengalgram+Maize (Bg+Mz) | Not Available | Iles | Graded bunding |
| Banapurra | 141 | 2.21 | RNKmB2g1 | LMU-6 | Moderately shallow (50-75 cm) | Clay | Gravelly (15-35%) | Medium (101-150 mm/m) | Very gently sloping (1-3%) | Moderate | Fallow land (Fl) | Not Available | IVes | Graded bunding |
| Banapurra | 142 | 4.09 | RNKmB2g1 | LMU-6 | Moderately shallow (50-75 cm) | Clay | Gravelly (15-35%) | Medium (101-150 mm/m) | Very gently sloping (1-3%) | Moderate | Fallow land+Sunflower (Fl+Sf) | Not Available | IVes | Graded bunding |
| Banapurra | 143 | 6.93 | DRLmB2 | LMU-1 | Moderately deep (75-100 cm) | Clay | Non gravelly (<15%) | Medium (101-150 mm/m) | Very gently sloping (1-3%) | Moderate | Bengalgram+Current Fallow (Bg+Cf) | Not Available | Iles | Graded bunding |
| Banapurra | 144 | 3.62 | DRLmB2 | LMU-1 | Moderately deep (75-100 cm) | Clay | Non gravelly (<15%) | Medium (101-150 mm/m) | Very gently sloping (1-3%) | Moderate | Bengalgram (Bg) | Not Available | Iles | Graded bunding |
| Banapurra | 145 | 4.08 | DRLmB2 | LMU-1 | Moderately deep (75-100 cm) | Clay | Non gravelly (<15%) | Medium (101-150 mm/m) | Very gently sloping (1-3%) | Moderate | Bengalgram (Bg) | Not Available | Iles | Graded bunding |
| Banapurra | 146 | 7.44 | RNKmB2g1 | LMU-6 | Moderately shallow (50-75 cm) | Clay | Gravelly (15-35%) | Medium (101-150 mm/m) | Very gently sloping (1-3%) | Moderate | Bengalgram (Bg) | Not Available | IVes | Graded bunding |
| Banapurra | 147 | 8.13 | RNKmB2g1 | LMU-6 | Moderately shallow (50-75 cm) | Clay | Gravelly (15-35%) | Medium (101-150 mm/m) | Very gently sloping (1-3%) | Moderate | Current Fallow+Pigeon Pea (Cf+Pp) | 1 Bore well | IVes | Graded bunding |
| Banapurra | 148 | 2.58 | RNKmB2g1 | LMU-6 | Moderately shallow (50-75 cm) | Clay | Non gravelly (<15%) | Medium (101-150 mm/m) | Very gently sloping (1-3%) | Moderate | Sunflower (Sf) | Not Available | IVes | Graded bunding |
| Banapurra | 149 | 4.98 | RNKmB2g1 | LMU-6 | Moderately shallow (50-75 cm) | Clay | Non gravelly (<15%) | Medium (101-150 mm/m) | Very gently sloping (1-3%) | Moderate | Sunflower+Maize (Sf+Mz) | Not Available | IVes | Graded bunding |
| Banapurra | 150 | 2.81 | RNKmB2g1 | LMU-6 | Moderately shallow (50-75 cm) | Clay | Non gravelly (<15%) | Medium (101-150 mm/m) | Very gently sloping (1-3%) | Moderate | Cotton+Chilly (Ct+Ch) | Not Available | IVes | Graded bunding |
| Banapurra | 151 | 4.94 | RNKmB2g1 | LMU-6 | Moderately shallow (50-75 cm) | Clay | Non gravelly (<15%) | Medium (101-150 mm/m) | Very gently sloping (1-3%) | Moderate | Maize (Mz) | Not Available | IVes | Graded bunding |

| Village | Survey No | Area (ha) | Soil Phase | LMU | Soil Depth | Surface Soil Texture | Soil Graveliness | Available Water Capacity | Slope | Soil Erosion | Current Land Use | WELLS | Land Capability | Conservation Plan |
|----------|-----------|-----------|------------|-------|-------------------------------|----------------------|------------------------|--------------------------|----------------------------|--------------|---|---------------|-----------------|--------------------|
| Banapura | 152 | 7.29 | RNKmB2g1 | LMU-6 | Moderately shallow (50-75 cm) | Clay | Gravelly (15-35%) | Medium (101-150 mm/m) | Very gently sloping (1-3%) | Moderate | Bengalgram+Jowar+Maize+Pigeon Pea (Bg+Jw+Mz+Pp) | Not Available | Ives | Graded bunding |
| Banapura | 153 | 2.18 | RNKmB2g1 | LMU-6 | Moderately shallow (50-75 cm) | Clay | Gravelly (15-35%) | Medium (101-150 mm/m) | Very gently sloping (1-3%) | Moderate | Bengalgram (Bg) | Not Available | Ives | Graded bunding |
| Banapura | 154 | 1.98 | RNKmB2g1 | LMU-6 | Moderately shallow (50-75 cm) | Clay | Gravelly (15-35%) | Medium (101-150 mm/m) | Very gently sloping (1-3%) | Moderate | Sunflower (Sf) | Not Available | Ives | Graded bunding |
| Banapura | 155 | 7.42 | BDGiB1g1 | LMU-2 | Moderately deep (75-100 cm) | Sandy clay | Gravelly (15-35%) | Low (51-100 mm/m) | Very gently sloping (1-3%) | Slight | Maize (Mz) | Not Available | IIs | Trench cum bunding |
| Banapura | 156 | 6.96 | BDGiB1g1 | LMU-2 | Moderately deep (75-100 cm) | Sandy clay | Gravelly (15-35%) | Low (51-100 mm/m) | Very gently sloping (1-3%) | Slight | Bengalgram (Bg) | Not Available | IIs | Trench cum bunding |
| Banapura | 157 | 8.45 | RNKmB2 | LMU-6 | Moderately shallow (50-75 cm) | Clay | Non gravelly (<15%) | Medium (101-150 mm/m) | Very gently sloping (1-3%) | Moderate | Current Fallow (Cf) | Not Available | Ives | Graded bunding |
| Banapura | 158 | 6.56 | NGPhA1 | LMU-2 | Deep (100-150 cm) | Sandy clay loam | Non gravelly (<15%) | Low (51-100 mm/m) | Nearly level (0-1%) | Slight | Current Fallow+Sunflower (Cf+Sf) | Not Available | IIs | Graded bunding |
| Banapura | 159/(1) | 1.53 | NGPhA1 | LMU-2 | Deep (100-150 cm) | Sandy clay loam | Non gravelly (<15%) | Low (51-100 mm/m) | Nearly level (0-1%) | Slight | Maize (Mz) | Not Available | IIs | Graded bunding |
| Banapura | 159/(2) | 4.36 | BPRhB2g2 | LMU-2 | Deep (100-150 cm) | Sandy clay loam | Very gravelly (35-60%) | Low (51-100 mm/m) | Very gently sloping (1-3%) | Moderate | Maize (Mz) | Not Available | IIs | Trench cum bunding |
| Banapura | 160 | 4.21 | BDGiB1g1 | LMU-2 | Moderately deep (75-100 cm) | Sandy clay | Gravelly (15-35%) | Low (51-100 mm/m) | Very gently sloping (1-3%) | Slight | Current Fallow (Cf) | Not Available | IIs | Trench cum bunding |
| Banapura | 161 | 6.64 | BDGiB1g1 | LMU-2 | Moderately deep (75-100 cm) | Sandy clay | Gravelly (15-35%) | Low (51-100 mm/m) | Very gently sloping (1-3%) | Slight | Sunflower (Sf) | Not Available | IIs | Trench cum bunding |
| Banapura | 162 | 4.07 | BDGiB1g1 | LMU-2 | Moderately deep (75-100 cm) | Sandy clay | Gravelly (15-35%) | Low (51-100 mm/m) | Very gently sloping (1-3%) | Slight | Maize (Mz) | Not Available | IIs | Trench cum bunding |
| Banapura | 163 | 3.81 | BDGiB1g1 | LMU-2 | Moderately deep (75-100 cm) | Sandy clay | Gravelly (15-35%) | Low (51-100 mm/m) | Very gently sloping (1-3%) | Slight | Maize (Mz) | Not Available | IIs | Trench cum bunding |
| Banapura | 164 | 6.5 | BDGiB1g1 | LMU-2 | Moderately deep (75-100 cm) | Sandy clay | Gravelly (15-35%) | Low (51-100 mm/m) | Very gently sloping (1-3%) | Slight | Maize (Mz) | Not Available | IIs | Trench cum bunding |
| Banapura | 165 | 5.31 | BDGiB1g1 | LMU-2 | Moderately deep (75-100 cm) | Sandy clay | Gravelly (15-35%) | Low (51-100 mm/m) | Very gently sloping (1-3%) | Slight | Current Fallow (Cf) | Not Available | IIs | Trench cum bunding |
| Banapura | 166 | 6.99 | GDPiB2 | LMU-2 | Deep (100-150 cm) | Sandy clay | Non gravelly (<15%) | Low (51-100 mm/m) | Very gently sloping (1-3%) | Moderate | Fallow land+Sunflower (Fl+Sf) | Not Available | IIs | Trench cum bunding |
| Banapura | 167 | 3.81 | BPRmB2 | LMU-2 | Deep (100-150 cm) | Clay | Non gravelly (<15%) | Low (51-100 mm/m) | Very gently sloping (1-3%) | Moderate | Sunflower (Sf) | Not Available | IIs | Trench cum bunding |
| Banapura | 168 | 5.48 | BPRhB2g1 | LMU-2 | Deep (100-150 cm) | Sandy clay loam | Gravelly (15-35%) | Low (51-100 mm/m) | Very gently sloping (1-3%) | Moderate | Current Fallow (Cf) | Not Available | IIs | Trench cum bunding |
| Banapura | 169 | 6.42 | BPRhB2g1 | LMU-2 | Deep (100-150 cm) | Sandy clay loam | Gravelly (15-35%) | Low (51-100 mm/m) | Very gently sloping (1-3%) | Moderate | Fallow land+Bajra (Fl+Bj) | Not Available | IIs | Trench cum bunding |
| Banapura | 170 | 7.08 | GDPiB2 | LMU-2 | Deep (100-150 cm) | Sandy clay | Non gravelly (<15%) | Low (51-100 mm/m) | Very gently sloping (1-3%) | Moderate | Fallow land+Bajra (Fl+Bj) | Not Available | IIs | Trench cum bunding |
| Banapura | 171 | 1.86 | BPRcA1 | LMU-2 | Deep (100-150 cm) | Sandy loam | Non gravelly (<15%) | Low (51-100 mm/m) | Nearly level (0-1%) | Slight | Horsegram (Hg) | Not Available | IIs | Graded bunding |
| Banapura | 172 | 3.08 | BPRcA1 | LMU-2 | Deep (100-150 cm) | Sandy loam | Non gravelly (<15%) | Low (51-100 mm/m) | Nearly level (0-1%) | Slight | Horsegram (Hg) | Not Available | IIs | Graded bunding |
| Banapura | 173 | 1.1 | BPRhB2g1 | LMU-2 | Deep (100-150 cm) | Sandy clay loam | Gravelly (15-35%) | Low (51-100 mm/m) | Very gently sloping (1-3%) | Moderate | Groundnut (Gn) | Not Available | IIs | Trench cum bunding |

| Village | Survey No | Area (ha) | Soil Phase | LMU | Soil Depth | Surface Soil Texture | Soil Graveliness | Available Water Capacity | Slope | Soil Erosion | Current Land Use | WELLS | Land Capability | Conservation Plan |
|----------|-----------|-----------|------------|-------|-------------------------------|----------------------|------------------------|--------------------------|----------------------------|--------------|---|---------------|-----------------|--------------------|
| Banapura | 174 | 6.59 | BPRhB2g1 | LMU-2 | Deep (100-150 cm) | Sandy clay loam | Gravelly (15-35%) | Low (51-100 mm/m) | Very gently sloping (1-3%) | Moderate | Current Fallow (Cf) | Not Available | Iles | Trench cum bunding |
| Banapura | 175 | 2.8 | GDPiB2 | LMU-2 | Deep (100-150 cm) | Sandy clay | Non gravelly (<15%) | Low (51-100 mm/m) | Very gently sloping (1-3%) | Moderate | Not Available (NA) | Not Available | Iles | Trench cum bunding |
| Banapura | 176 | 4.88 | BPRhB2g2 | LMU-2 | Deep (100-150 cm) | Sandy clay loam | Very gravelly (35-60%) | Low (51-100 mm/m) | Very gently sloping (1-3%) | Moderate | Current Fallow (Cf) | Not Available | Iles | Trench cum bunding |
| Banapura | 177 | 1.7 | BPRhB2g2 | LMU-2 | Deep (100-150 cm) | Sandy clay loam | Very gravelly (35-60%) | Low (51-100 mm/m) | Very gently sloping (1-3%) | Moderate | Current Fallow (Cf) | Not Available | Iles | Trench cum bunding |
| Banapura | 178 | 2.04 | BPRhB2g2 | LMU-2 | Deep (100-150 cm) | Sandy clay loam | Very gravelly (35-60%) | Low (51-100 mm/m) | Very gently sloping (1-3%) | Moderate | Current Fallow (Cf) | Not Available | Iles | Trench cum bunding |
| Banapura | 179 | 1.8 | GDPiB2 | LMU-2 | Deep (100-150 cm) | Sandy clay | Non gravelly (<15%) | Low (51-100 mm/m) | Very gently sloping (1-3%) | Moderate | Sunflower (Sf) | 2 Bore well | Iles | Trench cum bunding |
| Banapura | 180 | 4.19 | GDPiB2 | LMU-2 | Deep (100-150 cm) | Sandy clay | Non gravelly (<15%) | Low (51-100 mm/m) | Very gently sloping (1-3%) | Moderate | Maize (Mz) | Not Available | Iles | Trench cum bunding |
| Banapura | 181 | 7.38 | GDPiB2 | LMU-2 | Deep (100-150 cm) | Sandy clay | Non gravelly (<15%) | Low (51-100 mm/m) | Very gently sloping (1-3%) | Moderate | Current Fallow+Maize (Cf+Mz) | Not Available | Iles | Trench cum bunding |
| Banapura | 182 | 3.06 | GDPiB2 | LMU-2 | Deep (100-150 cm) | Sandy clay | Non gravelly (<15%) | Low (51-100 mm/m) | Very gently sloping (1-3%) | Moderate | Cluster bean (Cb) | 1 Bore well | Iles | Trench cum bunding |
| Banapura | 183 | 4.58 | BPRmB2 | LMU-2 | Deep (100-150 cm) | Clay | Non gravelly (<15%) | Low (51-100 mm/m) | Very gently sloping (1-3%) | Moderate | Current Fallow+Maize (Cf+Mz) | 2 Bore well | Iles | Trench cum bunding |
| Banapura | 184 | 5.66 | HDHiB2g1 | LMU-2 | Moderately deep (75-100 cm) | Sandy clay | Gravelly (15-35%) | Low (51-100 mm/m) | Very gently sloping (1-3%) | Moderate | Redgram+Horsegram+Current fallow (Rg+Hg+Cf) | Not Available | Iles | Trench cum bunding |
| Banapura | 185 | 5.92 | BPRmB2 | LMU-2 | Deep (100-150 cm) | Clay | Non gravelly (<15%) | Low (51-100 mm/m) | Very gently sloping (1-3%) | Moderate | Maize+Cluster bean (Mz+Cb) | Not Available | Iles | Trench cum bunding |
| Banapura | 186 | 3.54 | HDHiB2g1 | LMU-2 | Moderately deep (75-100 cm) | Sandy clay | Gravelly (15-35%) | Low (51-100 mm/m) | Very gently sloping (1-3%) | Moderate | Current Fallow+Sapota (Cf+Sp) | Not Available | Iles | Trench cum bunding |
| Banapura | 187 | 7.88 | HDHiB2g1 | LMU-2 | Moderately deep (75-100 cm) | Sandy clay | Gravelly (15-35%) | Low (51-100 mm/m) | Very gently sloping (1-3%) | Moderate | Horsegram+Chilly (Hg+Ch) | Not Available | Iles | Trench cum bunding |
| Banapura | 188 | 5.39 | TDHiB1 | LMU-5 | Moderately shallow (50-75 cm) | Sandy clay | Non gravelly (<15%) | Low (51-100 mm/m) | Very gently sloping (1-3%) | Slight | Current Fallow+Sapota (Cf+Sp) | Not Available | Ils | Trench cum bunding |
| Banapura | 189 | 2.5 | TDHiB1 | LMU-5 | Moderately shallow (50-75 cm) | Sandy clay | Non gravelly (<15%) | Low (51-100 mm/m) | Very gently sloping (1-3%) | Slight | Fallow land (Fl) | Not Available | Ils | Trench cum bunding |
| Banapura | 190 | 4.14 | BPRhB2g1 | LMU-2 | Deep (100-150 cm) | Sandy clay loam | Gravelly (15-35%) | Low (51-100 mm/m) | Very gently sloping (1-3%) | Moderate | Chilly (Ch) | 1 Bore well | Iles | Trench cum bunding |
| Banapura | 191 | 2.13 | TDHiB1 | LMU-5 | Moderately shallow (50-75 cm) | Sandy clay | Non gravelly (<15%) | Low (51-100 mm/m) | Very gently sloping (1-3%) | Slight | Chilly (Ch) | Not Available | Ils | Trench cum bunding |
| Banapura | 192 | 1.52 | TDHiB1 | LMU-5 | Moderately shallow (50-75 cm) | Sandy clay | Non gravelly (<15%) | Low (51-100 mm/m) | Very gently sloping (1-3%) | Slight | Chilly (Ch) | Not Available | Ils | Trench cum bunding |
| Banapura | 195 | 0.09 | TDHiB1 | LMU-5 | Moderately shallow (50-75 cm) | Sandy clay | Non gravelly (<15%) | Low (51-100 mm/m) | Very gently sloping (1-3%) | Slight | Not Available (NA) | Not Available | Ils | Trench cum bunding |
| Banapura | 196 | 0.001 | TDHiB1 | LMU-5 | Moderately shallow (50-75 cm) | Sandy clay | Non gravelly (<15%) | Low (51-100 mm/m) | Very gently sloping (1-3%) | Slight | Not Available (NA) | Not Available | Ils | Trench cum bunding |

Appendix III
Haligeri-3 (1T1a) Microwatershed
Soil Suitability Information

| Village | Survey Number | Mango | Maize | Sapota | Sorghum | Guava | Cotton | Tamarind | Lime | Bengal gram | Sunflower | Red gram | Amla | Jackfruit | Custard-apple | Cashew | Jamun | Musambi | Groundnut | Chilly | Tomato | Marigold | Chrysanthemum | Pomegranate | Bajra | Jasmine | Bhendi | Brinjal | Crossandra | Drumstick | Mulberry | Onion | |
|----------|---------------|-------|-------|--------|---------|-------|--------|----------|------|-------------|-----------|----------|------|-----------|---------------|--------|-------|---------|-----------|--------|--------|----------|---------------|-------------|-------|---------|--------|---------|------------|-----------|----------|-------|------|
| Halageri | 54 | S2rg | S2g | S2g | S2g | S2g | S2gt | S2rg | S2g | S3t | S2g | S2g | S2g | S2g | S2g | S2g | S2rg | S2g | S2gt | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | |
| Halageri | 55 | S2rg | S2g | S2g | S2g | S2g | S2gt | S2rg | S2g | S3t | S2g | S2g | S2g | S2g | S2g | S2g | S2rg | S2g | S2gt | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | |
| Halageri | 56 | S2rg | S2g | S2g | S2g | S2g | S2gt | S2rg | S2g | S3t | S2g | S2g | S2g | S2g | S2g | S2g | S2rg | S2g | S2gt | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | |
| Halageri | 60 | N1rz | S3rz | N1rz | S3rz | N1rz | S3rz | N1rz | N1rz | S3rz | N1rz | N1rz | S3rz | N1rz | S3rz | N1rt | N1rz | N1rz | S3rz | S3rz | S3rz | S3rz | S3rz | N1rz | S3rz | S3rz | S3rz | S3rz | S3rz | S3rz | N1rz | N1rz | S3rz |
| Halageri | 61 | N1rz | S3rz | N1rz | S3rz | N1rz | S3rz | N1rz | N1rz | S3rz | N1rz | N1rz | S3rz | N1rz | S3rz | N1rt | N1rz | N1rz | S3rz | S3rz | S3rz | S3rz | S3rz | N1rz | S3rz | S3rz | S3rz | S3rz | S3rz | S3rz | N1rz | N1rz | S3rz |
| Halageri | 62 | N1rn | S3n | N1n | S3n | N1n | S3n | N1rn | N1n | S3n | N1n | S3rn | N1n | N1n | N1tn | N1n | N1n | N1n | N1n | N1n | N1n | N1n | N1n | S3n | N1n | N1n | N1n | N1n | N1n | N1n | N1n | N1n | N1n |
| Halageri | 63 | S3r | S2g | S2rg | S2g | S2rg | S2rg | S3r | S2rg | S3t | S2rg | S2rg | S2g | S2rg | S2g | S2rg | S3r | S2rg | S2gt | S2g | S2g | S2g | S2g | S2rg | S2g | S2g | S2g | S2g | S2g | S2g | S2rg | S2rg | S2g |
| Halageri | 64 | S3r | S2g | S2rg | S2g | S2rg | S2rg | S3r | S2rg | S3t | S2rg | S2rg | S2g | S2rg | S2g | S2rg | S3r | S2rg | S2gt | S2g | S2g | S2g | S2g | S2rg | S2g | S2g | S2g | S2g | S2g | S2g | S2rg | S2rg | S2g |
| Halageri | 65 | S2rg | S2g | S2g | S2g | S2g | S2gt | S2rg | S2g | S3t | S2g | S2g | S2g | S2g | S2g | S2g | S2rg | S2g | S2gt | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g |
| Halageri | 66 | N1rn | S3n | N1n | S3n | N1n | S3n | N1rn | N1n | S3n | N1n | S3rn | N1n | N1n | N1tn | N1n | N1n | N1n | N1n | N1n | N1n | N1n | N1n | S3n | N1n | N1n | N1n | N1n | N1n | N1n | N1n | N1n | N1n |
| Halageri | 67 | S3rt | S2tz | S3t | S2z | S3tz | S2rz | S3rz | S2rz | S2z | S2rz | S2rt | S3z | S3tz | S2z | N1tz | S3rz | S2rz | S3t | S2tz | S3t | S2tz | S2tz | S2rt | S2tz | S3t | S2tz | S3t | S3t | S3z | S3tz | S3tz | |
| Halageri | 68 | S3rt | S2tz | S3t | S2z | S3tz | S2rz | S3rz | S2rz | S2z | S2rz | S2rt | S3z | S3tz | S2z | N1tz | S3rz | S2rz | S3t | S2tz | S3t | S2tz | S2tz | S2rt | S2tz | S3t | S2tz | S3t | S3t | S3z | S3tz | S3tz | |
| Halageri | 69 | N1rn | S3n | N1n | S3n | N1n | S3n | N1rn | N1n | S3n | N1n | S3rn | N1n | N1n | N1tn | N1n | N1n | N1n | N1n | N1n | N1n | N1n | N1n | S3n | N1n | N1n | N1n | N1n | N1n | N1n | N1n | N1n | N1n |
| Halageri | 70 | N1rz | S3rz | N1rz | S3rz | N1rz | S3rz | N1rz | N1rz | S3rz | N1rz | N1rz | S3rz | N1rz | S3rz | N1rt | N1rz | N1rz | S3rz | S3rz | S3rz | S3rz | S3rz | N1rz | S3rz | S3rz | S3rz | S3rz | S3rz | S3rz | N1rz | N1rz | S3rz |
| Halageri | 71 | N1rz | S3rz | N1rz | S3rz | N1rz | S3rz | N1rz | N1rz | S3rz | N1rz | N1rz | S3rz | N1rz | S3rz | N1rt | N1rz | N1rz | S3rz | S3rz | S3rz | S3rz | S3rz | N1rz | S3rz | S3rz | S3rz | S3rz | S3rz | S3rz | N1rz | N1rz | S3rz |
| Halageri | 72 | N1rz | S3rz | N1rz | S3rz | N1rz | S3rz | N1rz | N1rz | S3rz | N1rz | N1rz | S3rz | N1rz | S3rz | N1rt | N1rz | N1rz | S3rz | S3rz | S3rz | S3rz | S3rz | N1rz | S3rz | S3rz | S3rz | S3rz | S3rz | S3rz | N1rz | N1rz | S3rz |
| Halageri | 73 | N1rz | S3rz | N1rz | S3rz | N1rz | S3rz | N1rz | N1rz | S3rz | N1rz | N1rz | S3rz | N1rz | S3rz | N1rt | N1rz | N1rz | S3rz | S3rz | S3rz | S3rz | S3rz | N1rz | S3rz | S3rz | S3rz | S3rz | S3rz | S3rz | N1rz | N1rz | S3rz |
| Halageri | 74 | N1rz | S3rz | N1rz | S3rz | N1rz | S3rz | N1rz | N1rz | S3rz | N1rz | N1rz | S3rz | N1rz | S3rz | N1rt | N1rz | N1rz | S3rz | S3rz | S3rz | S3rz | S3rz | N1rz | S3rz | S3rz | S3rz | S3rz | S3rz | S3rz | N1rz | N1rz | S3rz |
| Halageri | 75 | N1rz | S3rz | N1rz | S3rz | N1rz | S3rz | N1rz | N1rz | S3rz | N1rz | N1rz | S3rz | N1rz | S3rz | N1rt | N1rz | N1rz | S3rz | S3rz | S3rz | S3rz | S3rz | N1rz | S3rz | S3rz | S3rz | S3rz | S3rz | S3rz | N1rz | N1rz | S3rz |
| Halageri | 76 | N1rz | S3rz | N1rz | S3rz | N1rz | S3rz | N1rz | N1rz | S3rz | N1rz | N1rz | S3rz | N1rz | S3rz | N1rt | N1rz | N1rz | S3rz | S3rz | S3rz | S3rz | S3rz | N1rz | S3rz | S3rz | S3rz | S3rz | S3rz | S3rz | N1rz | N1rz | S3rz |
| Halageri | 356 | S3r | S2g | S2rg | S2g | S2rg | S2rg | S3r | S2rg | S3gt | S2rg | S2rg | S2g | S2rg | S2g | S2rg | S3r | S2rg | S2gt | S2g | S2g | S3g | S3g | S2rg | S2g | S3g | S2g | S2g | S3g | S2rg | S2g | S3g | |
| Halageri | 357 | S3r | S2g | S2rg | S2g | S2rg | S2rg | S3r | S2rg | S3gt | S2rg | S2rg | S2g | S2rg | S2g | S2rg | S3r | S2rg | S2gt | S2g | S2g | S3g | S3g | S2rg | S2g | S3g | S2g | S2g | S3g | S2rg | S2g | S3g | |
| Halageri | 358 | S2rg | S2g | S2g | S2g | S2g | S2gt | S2rg | S2g | S3t | S2g | S2g | S2g | S2g | S2g | S2g | S2rg | S2g | S2gt | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | |

| Village | Survey Number | Mango | Maize | Sapota | Sorghum | Guava | Cotton | Tamarind | Lime | Bengal gram | Sunflower | Red gram | Amla | Jackfruit | Custard-apple | Cashew | Jamun | Musambi | Groundnut | Chilly | Tomato | Marigold | Chrysanthemum | Pomegranate | Bajra | Jasmine | Bhendi | Brinjal | Crossandra | Drumstick | Mulberry | Onion | |
|------------|---------------|-------|-------|--------|---------|-------|--------|----------|------|-------------|-----------|----------|------|-----------|---------------|--------|-------|---------|-----------|--------|--------|----------|---------------|-------------|-------|---------|--------|---------|------------|-----------|----------|-------|-----|
| Halageri | 359 | S3r | S2g | S2rg | S2g | S2rg | S2rg | S3r | S2rg | S3gt | S2rg | S2rg | S2g | S2rg | S2g | S2rg | S3r | S2rg | S2gt | S2g | S2g | S3g | S3g | S2rg | S2g | S3g | S2g | S2g | S3g | S2rg | S2g | S3g | |
| Thalabala | 187 | S3rt | S2tz | S3t | S2z | S3tz | S2rz | S3rz | S2rz | S2z | S2rz | S2rt | S3z | S3tz | S2z | N1tz | S3rz | S2rz | S3t | S2tz | S3t | S2tz | S2tz | S2rt | S2tz | S3t | S2tz | S3t | S3t | S3z | S3tz | S3tz | |
| Thalakalla | 98 | S3t | S2tz | S3t | S2z | S3tz | S2z | S3z | S2z | S2z | S2z | S2tz | S3z | S3tz | S2z | N1tz | S3z | S2z | S3tz | S2tz | S3t | S2tz | S2tz | S2tz | S2tz | S3t | S2tz | S3t | S3t | S3z | S3tz | S3tz | |
| Thalakalla | 99 | S3t | S2tz | S3t | S2z | S3tz | S2z | S3z | S2z | S2z | S2z | S2tz | S3z | S3tz | S2z | N1tz | S3z | S2z | S3tz | S2tz | S3t | S2tz | S2tz | S2tz | S2tz | S3t | S2tz | S3t | S3t | S3z | S3tz | S3tz | |
| Thalakalla | 100 | S3t | S2tz | S3t | S2z | S3tz | S2z | S3z | S2z | S2z | S2z | S2tz | S3z | S3tz | S2z | N1tz | S3z | S2z | S3tz | S2tz | S3t | S2tz | S2tz | S2tz | S2tz | S3t | S2tz | S3t | S3t | S3z | S3tz | S3tz | |
| Thalakalla | 101 | S3t | S2tz | S3t | S2z | S3tz | S2z | S3z | S2z | S2z | S2z | S2tz | S3z | S3tz | S2z | N1tz | S3z | S2z | S3tz | S2tz | S3t | S2tz | S2tz | S2tz | S2tz | S3t | S2tz | S3t | S3t | S3z | S3tz | S3tz | |
| Thalakalla | 102 | S2rg | S2g | S2g | S2g | S2g | S2gt | S2rg | S2g | S3t | S2g | S2g | S2g | S2g | S2g | S2g | S2rg | S2g | S2gt | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | |
| Thalakalla | 103 | S3t | S2tz | S3t | S2z | S3tz | S2z | S3z | S2z | S2z | S2z | S2tz | S3z | S3tz | S2z | N1tz | S3z | S2z | S3tz | S2tz | S3t | S2tz | S2tz | S2tz | S2tz | S3t | S2tz | S3t | S3t | S3z | S3tz | S3tz | |
| Thalakalla | 104 | S3t | S2tz | S3t | S2z | S3tz | S2z | S3z | S2z | S2z | S2z | S2tz | S3z | S3tz | S2z | N1tz | S3z | S2z | S3tz | S2tz | S3t | S2tz | S2tz | S2tz | S2tz | S3t | S2tz | S3t | S3t | S3z | S3tz | S3tz | |
| Thalakalla | 105 | S3t | S2tz | S3t | S2z | S3tz | S2z | S3z | S2z | S2z | S2z | S2tz | S3z | S3tz | S2z | N1tz | S3z | S2z | S3tz | S2tz | S3t | S2tz | S2tz | S2tz | S2tz | S3t | S2tz | S3t | S3t | S3z | S3tz | S3tz | |
| Thalakalla | 106 | S3t | S2tz | S3t | S2z | S3tz | S2z | S3z | S2z | S2z | S2z | S2tz | S3z | S3tz | S2z | N1tz | S3z | S2z | S3tz | S2tz | S3t | S2tz | S2tz | S2tz | S2tz | S3t | S2tz | S3t | S3t | S3z | S3tz | S3tz | |
| Thalakalla | 107 | S3rt | S2tz | S3t | S2z | S3tz | S2rz | S3rz | S2rz | S2z | S2rz | S2rt | S3z | S3tz | S2z | N1tz | S3rz | S2rz | S3t | S2tz | S3t | S2tz | S2tz | S2rt | S2tz | S3t | S2tz | S3t | S3t | S3z | S3tz | S3tz | |
| Thalakalla | 108 | S3t | S2tz | S3t | S2z | S3tz | S2z | S3z | S2z | S2z | S2z | S2tz | S3z | S3tz | S2z | N1tz | S3z | S2z | S3tz | S2tz | S3t | S2tz | S2tz | S2tz | S2tz | S3t | S2tz | S3t | S3t | S3z | S3tz | S3tz | |
| Thalakalla | 109 | S3t | S2tz | S3t | S2z | S3tz | S2z | S3z | S2z | S2z | S2z | S2tz | S3z | S3tz | S2z | N1tz | S3z | S2z | S3tz | S2tz | S3t | S2tz | S2tz | S2tz | S2tz | S3t | S2tz | S3t | S3t | S3z | S3tz | S3tz | |
| Thalakalla | 110 | S3t | S2tz | S3t | S2z | S3tz | S2z | S3z | S2z | S2z | S2z | S2tz | S3z | S3tz | S2z | N1tz | S3z | S2z | S3tz | S2tz | S3t | S2tz | S2tz | S2tz | S2tz | S3t | S2tz | S3t | S3t | S3z | S3tz | S3tz | |
| Thalakalla | 111 | S3t | S2tz | S3t | S2z | S3tz | S2z | S3z | S2z | S2z | S2z | S2tz | S3z | S3tz | S2z | N1tz | S3z | S2z | S3tz | S2tz | S3t | S2tz | S2tz | S2tz | S2tz | S3t | S2tz | S3t | S3t | S3z | S3tz | S3tz | |
| Thalakalla | 112 | N1r | S3r | N1r | S3r | N1r | S3rt | N1r | N1r | S3rt | N1r | N1r | S3r | N1r | S3r | N1r | N1r | N1r | S3r | S3r | S3r | S3r | S3r | N1r | S3r | S3r | S3r | S3r | S3r | S3r | N1r | N1r | S3r |
| Thalakalla | 113 | S2r | S1 | S1 | S1 | S1 | S2t | S2r | S1 | S3t | S1 | S1 | S1 | S1 | S1 | S1 | S2r | S1 | S2t | S1 | S1 | S1 | S1 | S1 | S1 | S1 | S1 | S1 | S1 | S1 | S1 | S1 | S1 |
| Thalakalla | 114 | N1r | S3r | N1r | S3r | N1r | S3rt | N1r | N1r | S3rt | N1r | N1r | S3r | N1r | S3r | N1r | N1r | N1r | S3r | S3r | S3r | S3r | S3r | N1r | S3r | S3r | S3r | S3r | S3r | S3r | N1r | N1r | S3r |
| Thalakalla | 115 | S2rg | S2g | S2g | S2g | S2g | S2gt | S2rg | S2g | S3t | S2g | S2g | S2g | S2g | S2g | S2g | S2rg | S2g | S2gt | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | |
| Thalakalla | 116 | S2rg | S2g | S2g | S2g | S2g | S2gt | S2rg | S2g | S3t | S2g | S2g | S2g | S2g | S2g | S2g | S2rg | S2g | S2gt | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | |
| Thalakalla | 124 | S3t | S2tz | S3t | S2z | S3tz | S2z | S3z | S2z | S2z | S2z | S2tz | S3z | S3tz | S2z | N1tz | S3z | S2z | S3t | S2tz | S3t | S2tz | S2tz | S2tz | S2tz | S3t | S2tz | S3t | S3t | S2tz | S3tz | S3tz | |
| Thalakalla | 125 | S3t | S2tz | S3t | S2z | S3tz | S2z | S3z | S2z | S2z | S2z | S2tz | S3z | S3tz | S2z | N1tz | S3z | S2z | S3t | S2tz | S3t | S2tz | S2tz | S2tz | S2tz | S3t | S2tz | S3t | S3t | S2tz | S3tz | S3tz | |
| Thalakalla | 126 | S3t | S2tz | S3t | S2z | S3tz | S2z | S3z | S2z | S2z | S2z | S2tz | S3z | S3tz | S2z | N1tz | S3z | S2z | S3t | S2tz | S3t | S2tz | S2tz | S2tz | S2tz | S3t | S2tz | S3t | S3t | S2tz | S3tz | S3tz | |
| Thalakalla | 127 | S3t | S2tz | S3t | S2z | S3tz | S2z | S3z | S2z | S2z | S2z | S2tz | S3z | S3tz | S2z | N1tz | S3z | S2z | S3t | S2tz | S3t | S2tz | S2tz | S2tz | S2tz | S3t | S2tz | S3t | S3t | S2tz | S3tz | S3tz | |
| Thalakalla | 128 | S3t | S2tz | S3t | S2z | S3tz | S2z | S3z | S2z | S2z | S2z | S2tz | S3z | S3tz | S2z | N1tz | S3z | S2z | S3t | S2tz | S3t | S2tz | S2tz | S2tz | S2tz | S3t | S2tz | S3t | S3t | S2tz | S3tz | S3tz | |

| Village | Survey Number | Mango | Maize | Sapota | Sorghum | Guava | Cotton | Tamarind | Lime | Bengal gram | Sunflower | Red gram | Amla | Jackfruit | Custard-apple | Cashew | Jamun | Musambi | Groundnut | Chilly | Tomato | Marigold | Chrysanthemum | Pomegranate | Bajra | Jasmine | Bhendi | Brinjal | Crossandra | Drumstick | Mulberry | Onion | |
|------------|---------------|-------|-------|--------|---------|-------|--------|----------|------|-------------|-----------|----------|------|-----------|---------------|--------|-------|---------|-----------|--------|--------|----------|---------------|-------------|-------|---------|--------|---------|------------|-----------|----------|-------|-----|
| Thalakalla | 129 | S3t | S2tz | S3t | S2z | S3tz | S2z | S3z | S2z | S2z | S2z | S2tz | S3z | S3tz | S2z | N1tz | S3z | S2z | S3t | S2tz | S3t | S2tz | S2tz | S2tz | S3t | S2tz | S3t | S3t | S3z | S3tz | S3tz | | |
| Thalakalla | 130 | S3t | S2tz | S3t | S2z | S3tz | S2z | S3z | S2z | S2z | S2z | S2tz | S3z | S3tz | S2z | N1tz | S3z | S2z | S3t | S2tz | S3t | S2tz | S2tz | S2tz | S3t | S2tz | S3t | S3t | S3z | S3tz | S3tz | | |
| Thalakalla | 131 | S3t | S2tz | S3t | S2z | S3tz | S2z | S3z | S2z | S2z | S2z | S2tz | S3z | S3tz | S2z | N1tz | S3z | S2z | S3t | S2tz | S3t | S2tz | S2tz | S2tz | S3t | S2tz | S3t | S3t | S3z | S3tz | S3tz | | |
| Thalakalla | 132 | S3t | S2tz | S3t | S2z | S3tz | S2z | S3z | S2z | S2z | S2z | S2tz | S3z | S3tz | S2z | N1tz | S3z | S2z | S3t | S2tz | S3t | S2tz | S2tz | S2tz | S3t | S2tz | S3t | S3t | S3z | S3tz | S3tz | | |
| Veerapura | 186 | S3rt | S2tz | S3t | S2z | S3tz | S2rz | S3rz | S2rz | S2z | S2rz | S2rt | S3z | S3tz | S2z | N1tz | S3rz | S2rz | S3t | S2tz | S3t | S2tz | S2tz | S2rt | S2tz | S3t | S2tz | S3t | S3t | S3z | S3tz | S3tz | |
| Veerapura | 187 | S3rt | S2tz | S3t | S2z | S3tz | S2rz | S3rz | S2rz | S2z | S2rz | S2rt | S3z | S3tz | S2z | N1tz | S3rz | S2rz | S3t | S2tz | S3t | S2tz | S2tz | S2rt | S2tz | S3t | S2tz | S3t | S3t | S3z | S3tz | S3tz | |
| Veerapura | 188 | S3rt | S2tz | S3t | S2z | S3tz | S2rz | S3rz | S2rz | S2z | S2rz | S2rt | S3z | S3tz | S2z | N1tz | S3rz | S2rz | S3t | S2tz | S3t | S2tz | S2tz | S2rt | S2tz | S3t | S2tz | S3t | S3t | S3z | S3tz | S3tz | |
| Banapura | 36 | S3rt | S2tz | S3t | S2z | S3tz | S2rz | S3rz | S2rz | S2z | S2rz | S2rt | S3z | S3tz | S2z | N1tz | S3rz | S2rz | S3t | S2tz | S3t | S2tz | S2tz | S2rt | S2tz | S3t | S2tz | S3t | S3t | S3z | S3tz | S3tz | |
| Banapura | 37 | N1r | S3g | S3rg | S3g | S3rg | S3gt | N1r | S3rg | S3gt | S3rg | S3rg | S2rg | S3rg | S2rg | S3rg | S3rg | S3rg | S2rg | S3g | S3g | S3g | S3g | S3rg | S2rg | S3g | S3g | S3g | S3g | S3g | S3rg | S3rg | S3g |
| Banapura | 38 | N1r | S3g | S3rg | S3g | S3rg | S3gt | N1r | S3rg | S3gt | S3rg | S3rg | S2rg | S3rg | S2rg | S3rg | S3rg | S3rg | S2rg | S3g | S3g | S3g | S3g | S3rg | S2rg | S3g | S3g | S3g | S3g | S3g | S3rg | S3rg | S3g |
| Banapura | 40 | N1r | S3g | S3rg | S3g | S3rg | S3gt | N1r | S3rg | S3gt | S3rg | S3rg | S2rg | S3rg | S2rg | S3rg | S3rg | S3rg | S2rg | S3g | S3g | S3g | S3g | S3rg | S2rg | S3g | S3g | S3g | S3g | S3g | S3rg | S3rg | S3g |
| Banapura | 41 | N1r | S3g | S3rg | S3g | S3rg | S3gt | N1r | S3rg | S3gt | S3rg | S3rg | S2rg | S3rg | S2rg | S3rg | S3rg | S3rg | S2rg | S3g | S3g | S3g | S3g | S3rg | S2rg | S3g | S3g | S3g | S3g | S3g | S3rg | S3rg | S3g |
| Banapura | 42 | N1r | S3g | S3rg | S3g | S3rg | S3gt | N1r | S3rg | S3gt | S3rg | S3rg | S2rg | S3rg | S2rg | S3rg | S3rg | S3rg | S2rg | S3g | S3g | S3g | S3g | S3rg | S2rg | S3g | S3g | S3g | S3g | S3g | S3rg | S3rg | S3g |
| Banapura | 43 | N1r | S3g | S3rg | S3g | S3rg | S3gt | N1r | S3rg | S3gt | S3rg | S3rg | S2rg | S3rg | S2rg | S3rg | S3rg | S3rg | S2rg | S3g | S3g | S3g | S3g | S3rg | S2rg | S3g | S3g | S3g | S3g | S3g | S3rg | S3rg | S3g |
| Banapura | 44 | S3r | S2g | S2rg | S2g | S2rg | S2rg | S3r | S2rg | S3t | S2rg | S2rg | S2g | S2rg | S2g | S2rg | S3r | S2rg | S2gt | S2g | S2g | S2g | S2g | S2rg | S2g | S2g | S2g | S2g | S2g | S2g | S2rg | S2rg | S2g |
| Banapura | 45 | N1r | S3g | S3rg | S3g | S3rg | S3gt | N1r | S3rg | S3gt | S3rg | S3rg | S2rg | S3rg | S2rg | S3rg | S3rg | S3rg | S2rg | S3g | S3g | S3g | S3g | S3rg | S2rg | S3g | S3g | S3g | S3g | S3g | S3rg | S3rg | S3g |
| Banapura | 46 | S2r | S1 | S1 | S1 | S1 | S2t | S2r | S1 | S3t | S1 | S1 | S1 | S1 | S1 | S1 | S2r | S1 | S2t | S1 | S1 | S1 | S1 | S1 | S1 | S1 | S1 | S1 | S1 | S1 | S1 | S1 | S1 |
| Banapura | 47 | N1r | S3g | S3rg | S3g | S3rg | S3gt | N1r | S3rg | S3gt | S3rg | S3rg | S2rg | S3rg | S2rg | S3rg | S3rg | S3rg | S2rg | S3g | S3g | S3g | S3g | S3rg | S2rg | S3g | S3g | S3g | S3g | S3g | S3rg | S3rg | S3g |
| Banapura | 54(1) | S2r | S1 | S1 | S1 | S1 | S2t | S2r | S1 | S3t | S1 | S1 | S1 | S1 | S1 | S1 | S2r | S1 | S2t | S1 | S1 | S1 | S1 | S1 | S1 | S1 | S1 | S1 | S1 | S1 | S1 | S1 | |
| Banapura | 56 | S2rg | S2g | S2g | S2g | S2g | S2gt | S2rg | S2g | S3t | S2g | S2g | S2g | S2g | S2g | S2g | S2rg | S2g | S2gt | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g |
| Banapura | 57 | S2rg | S2g | S2g | S2g | S2g | S2gt | S2rg | S2g | S3t | S2g | S2g | S2g | S2g | S2g | S2g | S2rg | S2g | S2gt | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g |
| Banapura | 58 | S3r | S2g | S2rg | S2g | S2rg | S2rg | S3r | S2rg | S3t | S2rg | S2rg | S2g | S2rg | S2g | S2rg | S3r | S2rg | S2gt | S2g | S2g | S2g | S2g | S2rg | S2g | S2g | S2g | S2g | S2g | S2g | S2rg | S2rg | S2g |
| Banapura | 59 | S3r | S2g | S2rg | S2g | S2rg | S2rg | S3r | S2rg | S3t | S2rg | S2rg | S2g | S2rg | S2g | S2rg | S3r | S2rg | S2gt | S2g | S2g | S2g | S2g | S2rg | S2g | S2g | S2g | S2g | S2g | S2g | S2rg | S2rg | S2g |
| Banapura | 60 | S3r | S2g | S2rg | S2g | S2rg | S2rg | S3r | S2rg | S3t | S2rg | S2rg | S2g | S2rg | S2g | S2rg | S3r | S2rg | S2gt | S2g | S2g | S2g | S2g | S2rg | S2g | S2g | S2g | S2g | S2g | S2g | S2rg | S2rg | S2g |
| Banapura | 61 | S3r | S2g | S2rg | S2g | S2rg | S2rg | S3r | S2rg | S3t | S2rg | S2rg | S2g | S2rg | S2g | S2rg | S3r | S2rg | S2gt | S2g | S2g | S2g | S2g | S2rg | S2g | S2g | S2g | S2g | S2g | S2g | S2rg | S2rg | S2g |
| Banapura | 62 | N1rn | S3n | N1n | S3n | N1n | S3n | N1rn | N1n | S3n | N1n | S3rn | N1n | N1n | N1n | N1tn | N1n | N1n | N1n | N1n | N1n | N1n | N1n | N1n | S3n | N1n | N1n | N1n | N1n | N1n | N1n | N1n | N1n |

| Village | Survey Number | Mango | Maize | Sapota | Sorghum | Guava | Cotton | Tamarind | Lime | Bengal gram | Sunflower | Red gram | Amla | Jackfruit | Custard-apple | Cashew | Jamun | Musambi | Groundnut | Chilly | Tomato | Marigold | Chrysanthemum | Pomegranate | Bajra | Jasmine | Bhendi | Brinjal | Crossandra | Drumstick | Mulberry | Onion | |
|----------|---------------|-------|-------|--------|---------|-------|--------|----------|------|-------------|-----------|----------|------|-----------|---------------|--------|-------|---------|-----------|--------|--------|----------|---------------|-------------|-------|---------|--------|---------|------------|-----------|----------|-------|-----|
| Banapura | 63 | N1rn | S3n | N1n | S3n | N1n | S3n | N1rn | N1n | S3n | N1n | S3rn | N1n | N1n | N1n | N1tn | N1n | N1n | N1n | N1n | N1n | N1n | N1n | N1n | S3n | N1n | N1n | N1n | N1n | N1n | N1n | N1n | |
| Banapura | 64 | N1rn | S3n | N1n | S3n | N1n | S3n | N1rn | N1n | S3n | N1n | S3rn | N1n | N1n | N1n | N1tn | N1n | N1n | N1n | N1n | N1n | N1n | N1n | N1n | S3n | N1n | N1n | N1n | N1n | N1n | N1n | N1n | |
| Banapura | 65 | N1rn | S3n | N1n | S3n | N1n | S3n | N1rn | N1n | S3n | N1n | S3rn | N1n | N1n | N1n | N1tn | N1n | N1n | N1n | N1n | N1n | N1n | N1n | N1n | S3n | N1n | N1n | N1n | N1n | N1n | N1n | N1n | |
| Banapura | 66 | S3rt | S2tz | S3t | S2z | S3tz | S2rz | S3rz | S2rz | S2z | S2rz | S2rt | S3z | S3tz | S2z | N1tz | S3rz | S2rz | S3t | S2tz | S3t | S2tz | S2tz | S2rt | S2tz | S3t | S2tz | S3t | S3t | S3z | S3tz | S3tz | |
| Banapura | 67 | S3rt | S2tz | S3t | S2z | S3tz | S2rz | S3rz | S2rz | S2z | S2rz | S2rt | S3z | S3tz | S2z | N1tz | S3rz | S2rz | S3t | S2tz | S3t | S2tz | S2tz | S2rt | S2tz | S3t | S2tz | S3t | S3t | S3z | S3tz | S3tz | |
| Banapura | 68 | N1rn | S3n | N1n | S3n | N1n | S3n | N1rn | N1n | S3n | N1n | S3rn | N1n | N1n | N1n | N1tn | N1n | N1n | N1n | N1n | N1n | N1n | N1n | N1n | S3n | N1n | N1n | N1n | N1n | N1n | N1n | N1n | |
| Banapura | 69 | N1rn | S3n | N1n | S3n | N1n | S3n | N1rn | N1n | S3n | N1n | S3rn | N1n | N1n | N1n | N1tn | N1n | N1n | N1n | N1n | N1n | N1n | N1n | N1n | S3n | N1n | N1n | N1n | N1n | N1n | N1n | N1n | |
| Banapura | 70 | N1r | S3g | S3rg | S3g | S3rg | S3gt | N1r | S3rg | S3gt | S3rg | S3rg | S2rg | S3rg | S2rg | S3rg | S3rg | S3rg | S2rg | S3g | S3g | S3g | S3g | S3rg | S2rg | S3g | S3g | S3g | S3g | S3g | S3rg | S3rg | S3g |
| Banapura | 71 | N1r | S3g | S3rg | S3g | S3rg | S3gt | N1r | S3rg | S3gt | S3rg | S3rg | S2rg | S3rg | S2rg | S3rg | S3rg | S3rg | S2rg | S3g | S3g | S3g | S3g | S3rg | S2rg | S3g | S3g | S3g | S3g | S3g | S3rg | S3rg | S3g |
| Banapura | 72 | N1r | S3g | S3rg | S3g | S3rg | S3gt | N1r | S3rg | S3gt | S3rg | S3rg | S2rg | S3rg | S2rg | S3rg | S3rg | S3rg | S2rg | S3g | S3g | S3g | S3g | S3rg | S2rg | S3g | S3g | S3g | S3g | S3g | S3rg | S3rg | S3g |
| Banapura | 73 | N1r | S3g | S3rg | S3g | S3rg | S3gt | N1r | S3rg | S3gt | S3rg | S3rg | S2rg | S3rg | S2rg | S3rg | S3rg | S3rg | S2rg | S3g | S3g | S3g | S3g | S3rg | S2rg | S3g | S3g | S3g | S3g | S3g | S3rg | S3rg | S3g |
| Banapura | 74 | S3rt | S2tz | S3t | S2z | S3tz | S2rz | S3rz | S2rz | S2z | S2rz | S2rt | S3z | S3tz | S2z | N1tz | S3rz | S2rz | S3t | S2tz | S3t | S2tz | S2tz | S2rt | S2tz | S3t | S2tz | S3t | S3t | S3z | S3tz | S3tz | |
| Banapura | 75 | S3rt | S2tz | S3t | S2z | S3tz | S2rz | S3rz | S2rz | S2z | S2rz | S2rt | S3z | S3tz | S2z | N1tz | S3rz | S2rz | S3t | S2tz | S3t | S2tz | S2tz | S2rt | S2tz | S3t | S2tz | S3t | S3t | S3z | S3tz | S3tz | |
| Banapura | 76 | S3rt | S2tz | S3t | S2z | S3tz | S2rz | S3rz | S2rz | S2z | S2rz | S2rt | S3z | S3tz | S2z | N1tz | S3rz | S2rz | S3t | S2tz | S3t | S2tz | S2tz | S2rt | S2tz | S3t | S2tz | S3t | S3t | S3z | S3tz | S3tz | |
| Banapura | 95 | S3rt | S2tz | S3t | S2z | S3tz | S2rz | S3rz | S2rz | S2z | S2rz | S2rt | S3z | S3tz | S2z | N1tz | S3rz | S2rz | S3t | S2tz | S3t | S2tz | S2tz | S2rt | S2tz | S3t | S2tz | S3t | S3t | S3z | S3tz | S3tz | |
| Banapura | 96 | S3rt | S2tz | S3t | S2z | S3tz | S2rz | S3rz | S2rz | S2z | S2rz | S2rt | S3z | S3tz | S2z | N1tz | S3rz | S2rz | S3t | S2tz | S3t | S2tz | S2tz | S2rt | S2tz | S3t | S2tz | S3t | S3t | S3z | S3tz | S3tz | |
| Banapura | 97 | S3rt | S2tz | S3t | S2z | S3tz | S2rz | S3rz | S2rz | S2z | S2rz | S2rt | S3z | S3tz | S2z | N1tz | S3rz | S2rz | S3t | S2tz | S3t | S2tz | S2tz | S2rt | S2tz | S3t | S2tz | S3t | S3t | S3z | S3tz | S3tz | |
| Banapura | 98 | S3rt | S2tz | S3t | S2z | S3tz | S2rz | S3rz | S2rz | S2z | S2rz | S2rt | S3z | S3tz | S2z | N1tz | S3rz | S2rz | S3t | S2tz | S3t | S2tz | S2tz | S2rt | S2tz | S3t | S2tz | S3t | S3t | S3z | S3tz | S3tz | |
| Banapura | 99 | S3t | S2tz | S3t | S2z | S3tz | S2z | S3z | S2z | S2z | S2z | S2tz | S3z | S3tz | S2z | N1tz | S3z | S2z | S3tz | S2tz | S3t | S2tz | S2tz | S2tz | S2tz | S3t | S2tz | S3t | S3t | S3z | S3tz | S3tz | |
| Banapura | 100 | S3t | S2tz | S3t | S2z | S3tz | S2z | S3z | S2z | S2z | S2z | S2tz | S3z | S3tz | S2z | N1tz | S3z | S2z | S3tz | S2tz | S3t | S2tz | S2tz | S2tz | S2tz | S3t | S2tz | S3t | S3t | S3z | S3tz | S3tz | |
| Banapura | 101 | S3t | S2tz | S3t | S2z | S3tz | S2z | S3z | S2z | S2z | S2z | S2tz | S3z | S3tz | S2z | N1tz | S3z | S2z | S3tz | S2tz | S3t | S2tz | S2tz | S2tz | S2tz | S3t | S2tz | S3t | S3t | S3z | S3tz | S3tz | |
| Banapura | 102 | S3t | S2tz | S3t | S2z | S3tz | S2z | S3z | S2z | S2z | S2z | S2tz | S3z | S3tz | S2z | N1tz | S3z | S2z | S3tz | S2tz | S3t | S2tz | S2tz | S2tz | S2tz | S3t | S2tz | S3t | S3t | S3z | S3tz | S3tz | |
| Banapura | 107 | S3t | S2tz | S3t | S2z | S3tz | S2z | S3z | S2z | S2z | S2z | S2tz | S3z | S3tz | S2z | N1tz | S3z | S2z | S3tz | S2tz | S3t | S2tz | S2tz | S2tz | S2tz | S3t | S2tz | S3t | S3t | S3z | S3tz | S3tz | |
| Banapura | 108 | S3t | S2tz | S3t | S2z | S3tz | S2z | S3z | S2z | S2z | S2z | S2tz | S3z | S3tz | S2z | N1tz | S3z | S2z | S3tz | S2tz | S3t | S2tz | S2tz | S2tz | S2tz | S3t | S2tz | S3t | S3t | S3z | S3tz | S3tz | |
| Banapura | 109 | S3rt | S2tz | S3t | S2z | S3tz | S2rz | S3rz | S2rz | S2z | S2rz | S2rt | S3z | S3tz | S2z | N1tz | S3rz | S2rz | S3t | S2tz | S3t | S2tz | S2tz | S2rt | S2tz | S3t | S2tz | S3t | S3t | S3z | S3tz | S3tz | |

| Village | Survey Number | Mango | Maize | Sapota | Sorghum | Guava | Cotton | Tamarind | Lime | Bengal gram | Sunflower | Red gram | Amla | Jackfruit | Custard-apple | Cashew | Jamun | Musambi | Groundnut | Chilly | Tomato | Marigold | Chrysanthemum | Pomegranate | Bajra | Jasmine | Bhendi | Brinjal | Crossandra | Drumstick | Mulberry | Onion |
|----------|---------------|-------|-------|--------|---------|-------|--------|----------|------|-------------|-----------|----------|------|-----------|---------------|--------|-------|---------|-----------|--------|--------|----------|---------------|-------------|-------|---------|--------|---------|------------|-----------|----------|-------|
| Banapura | 110 | S3rt | S2tz | S3t | S2z | S3tz | S2rz | S3rz | S2rz | S2z | S2rz | S2rt | S3z | S3tz | S2z | N1tz | S3rz | S2rz | S3t | S2tz | S3t | S2tz | S2tz | S2rt | S2tz | S3t | S2tz | S3t | S3t | S3z | S3tz | S3tz |
| Banapura | 111 | S3rt | S2tz | S3t | S2z | S3tz | S2rz | S3rz | S2rz | S2z | S2rz | S2rt | S3z | S3tz | S2z | N1tz | S3rz | S2rz | S3t | S2tz | S3t | S2tz | S2tz | S2rt | S2tz | S3t | S2tz | S3t | S3t | S3z | S3tz | S3tz |
| Banapura | 112 | S3t | S2tz | S3t | S2z | S3tz | S2z | S3z | S2z | S2z | S2z | S2tz | S3z | S3tz | S2z | N1tz | S3z | S2z | S3tz | S2tz | S3t | S2tz | S2tz | S2tz | S2tz | S3t | S2tz | S3t | S3t | S3z | S3tz | S3tz |
| Banapura | 113 | S3t | S2tz | S3t | S2z | S3tz | S2z | S3z | S2z | S2z | S2z | S2tz | S3z | S3tz | S2z | N1tz | S3z | S2z | S3tz | S2tz | S3t | S2tz | S2tz | S2tz | S2tz | S3t | S2tz | S3t | S3t | S3z | S3tz | S3tz |
| Banapura | 115 | S3t | S2tz | S3t | S2z | S3tz | S2z | S3z | S2z | S2z | S2z | S2tz | S3z | S3tz | S2z | N1tz | S3z | S2z | S3tz | S2tz | S3t | S2tz | S2tz | S2tz | S2tz | S3t | S2tz | S3t | S3t | S3z | S3tz | S3tz |
| Banapura | 116 | S3t | S2tz | S3t | S2z | S3tz | S2z | S3z | S2z | S2z | S2z | S2tz | S3z | S3tz | S2z | N1tz | S3z | S2z | S3tz | S2tz | S3t | S2tz | S2tz | S2tz | S2tz | S3t | S2tz | S3t | S3t | S3z | S3tz | S3tz |
| Banapura | 117 | S3t | S2tz | S3t | S2z | S3tz | S2z | S3z | S2z | S2z | S2z | S2tz | S3z | S3tz | S2z | N1tz | S3z | S2z | S3tz | S2tz | S3t | S2tz | S2tz | S2tz | S2tz | S3t | S2tz | S3t | S3t | S3z | S3tz | S3tz |
| Banapura | 118 | S3t | S2tz | S3t | S2z | S3tz | S2z | S3z | S2z | S2z | S2z | S2tz | S3z | S3tz | S2z | N1tz | S3z | S2z | S3tz | S2tz | S3t | S2tz | S2tz | S2tz | S2tz | S3t | S2tz | S3t | S3t | S3z | S3tz | S3tz |
| Banapura | 119 | S3t | S2tz | S3t | S2z | S3tz | S2z | S3z | S2z | S2z | S2z | S2tz | S3z | S3tz | S2z | N1tz | S3z | S2z | S3tz | S2tz | S3t | S2tz | S2tz | S2tz | S2tz | S3t | S2tz | S3t | S3t | S3z | S3tz | S3tz |
| Banapura | 120 | S3t | S2tz | S3t | S2z | S3tz | S2z | S3z | S2z | S2z | S2z | S2tz | S3z | S3tz | S2z | N1tz | S3z | S2z | S3tz | S2tz | S3t | S2tz | S2tz | S2tz | S2tz | S3t | S2tz | S3t | S3t | S3z | S3tz | S3tz |
| Banapura | 121 | S3t | S2tz | S3t | S2z | S3tz | S2z | S3z | S2z | S2z | S2z | S2tz | S3z | S3tz | S2z | N1tz | S3z | S2z | S3t | S2tz | S3t | S2tz | S2tz | S2tz | S2tz | S3t | S2tz | S3t | S3t | S3z | S3tz | S3tz |
| Banapura | 124 | S3t | S2tz | S3t | S2z | S3tz | S2z | S3z | S2z | S2z | S2z | S2tz | S3z | S3tz | S2z | N1tz | S3z | S2z | S3t | S2tz | S3t | S2tz | S2tz | S2tz | S2tz | S3t | S2tz | S3t | S3t | S3z | S3tz | S3tz |
| Banapura | 125 | S3t | S2tz | S3t | S2z | S3tz | S2z | S3z | S2z | S2z | S2z | S2tz | S3z | S3tz | S2z | N1tz | S3z | S2z | S3t | S2tz | S3t | S2tz | S2tz | S2tz | S2tz | S3t | S2tz | S3t | S3t | S3z | S3tz | S3tz |
| Banapura | 126 | S3t | S2tz | S3t | S2z | S3tz | S2z | S3z | S2z | S2z | S2z | S2tz | S3z | S3tz | S2z | N1tz | S3z | S2z | S3tz | S2tz | S3t | S2tz | S2tz | S2tz | S2tz | S3t | S2tz | S3t | S3t | S3z | S3tz | S3tz |
| Banapura | 127 | S3t | S2tz | S3t | S2z | S3tz | S2z | S3z | S2z | S2z | S2z | S2tz | S3z | S3tz | S2z | N1tz | S3z | S2z | S3tz | S2tz | S3t | S2tz | S2tz | S2tz | S2tz | S3t | S2tz | S3t | S3t | S3z | S3tz | S3tz |
| Banapura | 128 | S3rt | S2tz | S3t | S2z | S3tz | S2rz | S3rz | S2rz | S2z | S2rz | S2rt | S3z | S3tz | S2z | N1tz | S3rz | S2rz | S3t | S2tz | S3t | S2tz | S2tz | S2rt | S2tz | S3t | S2tz | S3t | S3t | S3z | S3tz | S3tz |
| Banapura | 129 | S3rt | S2tz | S3t | S2z | S3tz | S2rz | S3rz | S2rz | S2z | S2rz | S2rt | S3z | S3tz | S2z | N1tz | S3rz | S2rz | S3t | S2tz | S3t | S2tz | S2tz | S2rt | S2tz | S3t | S2tz | S3t | S3t | S3z | S3tz | S3tz |
| Banapura | 130 | S3rt | S2tz | S3t | S2z | S3tz | S2rz | S3rz | S2rz | S2z | S2rz | S2rt | S3z | S3tz | S2z | N1tz | S3rz | S2rz | S3t | S2tz | S3t | S2tz | S2tz | S2rt | S2tz | S3t | S2tz | S3t | S3t | S3z | S3tz | S3tz |
| Banapura | 131 | S3rt | S2tz | S3t | S2z | S3tz | S2rz | S3rz | S2rz | S2z | S2rz | S2rt | S3z | S3tz | S2z | N1tz | S3rz | S2rz | S3t | S2tz | S3t | S2tz | S2tz | S2rt | S2tz | S3t | S2tz | S3t | S3t | S3z | S3tz | S3tz |
| Banapura | 132 | S3rt | S2tz | S3t | S2z | S3tz | S2rz | S3rz | S2rz | S2z | S2rz | S2rt | S3z | S3tz | S2z | N1tz | S3rz | S2rz | S3t | S2tz | S3t | S2tz | S2tz | S2rt | S2tz | S3t | S2tz | S3t | S3t | S3z | S3tz | S3tz |
| Banapura | 133 | S3rt | S2tz | S3t | S2z | S3tz | S2rz | S3rz | S2rz | S2z | S2rz | S2rt | S3z | S3tz | S2z | N1tz | S3rz | S2rz | S3t | S2tz | S3t | S2tz | S2tz | S2rt | S2tz | S3t | S2tz | S3t | S3t | S3z | S3tz | S3tz |
| Banapura | 134 | S3rt | S2tz | S3t | S2z | S3tz | S2rz | S3rz | S2rz | S2z | S2rz | S2rt | S3z | S3tz | S2z | N1tz | S3rz | S2rz | S3t | S2tz | S3t | S2tz | S2tz | S2rt | S2tz | S3t | S2tz | S3t | S3t | S3z | S3tz | S3tz |
| Banapura | 135 | S3t | S2tz | S3t | S2z | S3tz | S2z | S3z | S2z | S2z | S2z | S2tz | S3z | S3tz | S2z | N1tz | S3z | S2z | S3t | S2tz | S3t | S2tz | S2tz | S2tz | S2tz | S3t | S2tz | S3t | S3t | S3z | S3tz | S3tz |
| Banapura | 136 | S3t | S2tz | S3t | S2z | S3tz | S2z | S3z | S2z | S2z | S2z | S2tz | S3z | S3tz | S2z | N1tz | S3z | S2z | S3t | S2tz | S3t | S2tz | S2tz | S2tz | S2tz | S3t | S2tz | S3t | S3t | S3z | S3tz | S3tz |
| Banapura | 137 | S3rt | S2tz | S3t | S2z | S3tz | S2rz | S3rz | S2rz | S2z | S2rz | S2rt | S3z | S3tz | S2z | N1tz | S3rz | S2rz | S3t | S2tz | S3t | S2tz | S2tz | S2rt | S2tz | S3t | S2tz | S3t | S3t | S3z | S3tz | S3tz |
| Banapura | 138 | S3rt | S2tz | S3t | S2z | S3tz | S2rz | S3rz | S2rz | S2z | S2rz | S2rt | S3z | S3tz | S2z | N1tz | S3rz | S2rz | S3t | S2tz | S3t | S2tz | S2tz | S2rt | S2tz | S3t | S2tz | S3t | S3t | S3z | S3tz | S3tz |

| Village | Survey Number | Mango | Maize | Sapota | Sorghum | Guava | Cotton | Tamarind | Lime | Bengal gram | Sunflower | Red gram | Amla | Jackfruit | Custard-apple | Cashew | Jamun | Musambi | Groundnut | Chilly | Tomato | Marigold | Chrysanthemum | Pomegranate | Bajra | Jasmine | Bhendi | Brinjal | Crossandra | Drumstick | Mulberry | Onion | |
|----------|---------------|-------|-------|--------|---------|-------|--------|----------|------|-------------|-----------|----------|------|-----------|---------------|--------|-------|---------|-----------|--------|--------|----------|---------------|-------------|-------|---------|--------|---------|------------|-----------|----------|-------|-----|
| Banapura | 139 | N1rn | S3n | N1n | S3n | N1n | S3n | N1rn | N1n | S3n | N1n | S3rn | N1n | N1n | N1n | N1tn | N1n | N1n | N1n | N1n | N1n | N1n | N1n | N1n | S3n | N1n | N1n | N1n | N1n | N1n | N1n | N1n | |
| Banapura | 140 | S3rt | S2tz | S3t | S2z | S3tz | S2rz | S3rz | S2rz | S2z | S2rz | S2rt | S3z | S3tz | S2z | N1tz | S3rz | S2rz | S3t | S2tz | S3t | S2tz | S2tz | S2rt | S2tz | S3t | S2tz | S3t | S3t | S3z | S3tz | S3tz | |
| Banapura | 141 | N1rn | S3n | N1n | S3n | N1n | S3n | N1rn | N1n | S3n | N1n | S3rn | N1n | N1n | N1n | N1tn | N1n | N1n | N1n | N1n | N1n | N1n | N1n | N1n | S3n | N1n | N1n | N1n | N1n | N1n | N1n | N1n | |
| Banapura | 142 | N1rn | S3n | N1n | S3n | N1n | S3n | N1rn | N1n | S3n | N1n | S3rn | N1n | N1n | N1n | N1tn | N1n | N1n | N1n | N1n | N1n | N1n | N1n | N1n | S3n | N1n | N1n | N1n | N1n | N1n | N1n | N1n | |
| Banapura | 143 | S3rt | S2tz | S3t | S2z | S3tz | S2rz | S3rz | S2rz | S2z | S2rz | S2rt | S3z | S3tz | S2z | N1tz | S3rz | S2rz | S3t | S2tz | S3t | S2tz | S2tz | S2rt | S2tz | S3t | S2tz | S3t | S3t | S3z | S3tz | S3tz | |
| Banapura | 144 | S3rt | S2tz | S3t | S2z | S3tz | S2rz | S3rz | S2rz | S2z | S2rz | S2rt | S3z | S3tz | S2z | N1tz | S3rz | S2rz | S3t | S2tz | S3t | S2tz | S2tz | S2rt | S2tz | S3t | S2tz | S3t | S3t | S3z | S3tz | S3tz | |
| Banapura | 145 | S3rt | S2tz | S3t | S2z | S3tz | S2rz | S3rz | S2rz | S2z | S2rz | S2rt | S3z | S3tz | S2z | N1tz | S3rz | S2rz | S3t | S2tz | S3t | S2tz | S2tz | S2rt | S2tz | S3t | S2tz | S3t | S3t | S3z | S3tz | S3tz | |
| Banapura | 146 | N1rn | S3n | N1n | S3n | N1n | S3n | N1rn | N1n | S3n | N1n | S3rn | N1n | N1n | N1n | N1tn | N1n | N1n | N1n | N1n | N1n | N1n | N1n | N1n | S3n | N1n | N1n | N1n | N1n | N1n | N1n | N1n | |
| Banapura | 147 | N1rn | S3n | N1n | S3n | N1n | S3n | N1rn | N1n | S3n | N1n | S3rn | N1n | N1n | N1n | N1tn | N1n | N1n | N1n | N1n | N1n | N1n | N1n | N1n | S3n | N1n | N1n | N1n | N1n | N1n | N1n | N1n | |
| Banapura | 148 | N1rn | S3n | N1n | S3n | N1n | S3n | N1rn | N1n | S3n | N1n | S3rn | N1n | N1n | N1n | N1tn | N1n | N1n | N1n | N1n | N1n | N1n | N1n | N1n | S3n | N1n | N1n | N1n | N1n | N1n | N1n | N1n | |
| Banapura | 149 | N1rn | S3n | N1n | S3n | N1n | S3n | N1rn | N1n | S3n | N1n | S3rn | N1n | N1n | N1n | N1tn | N1n | N1n | N1n | N1n | N1n | N1n | N1n | N1n | S3n | N1n | N1n | N1n | N1n | N1n | N1n | N1n | |
| Banapura | 150 | N1rn | S3n | N1n | S3n | N1n | S3n | N1rn | N1n | S3n | N1n | S3rn | N1n | N1n | N1n | N1tn | N1n | N1n | N1n | N1n | N1n | N1n | N1n | N1n | S3n | N1n | N1n | N1n | N1n | N1n | N1n | N1n | |
| Banapura | 151 | N1rn | S3n | N1n | S3n | N1n | S3n | N1rn | N1n | S3n | N1n | S3rn | N1n | N1n | N1n | N1tn | N1n | N1n | N1n | N1n | N1n | N1n | N1n | N1n | S3n | N1n | N1n | N1n | N1n | N1n | N1n | N1n | |
| Banapura | 152 | N1rn | S3n | N1n | S3n | N1n | S3n | N1rn | N1n | S3n | N1n | S3rn | N1n | N1n | N1n | N1tn | N1n | N1n | N1n | N1n | N1n | N1n | N1n | N1n | S3n | N1n | N1n | N1n | N1n | N1n | N1n | N1n | |
| Banapura | 153 | N1rn | S3n | N1n | S3n | N1n | S3n | N1rn | N1n | S3n | N1n | S3rn | N1n | N1n | N1n | N1tn | N1n | N1n | N1n | N1n | N1n | N1n | N1n | N1n | S3n | N1n | N1n | N1n | N1n | N1n | N1n | N1n | |
| Banapura | 154 | N1rn | S3n | N1n | S3n | N1n | S3n | N1rn | N1n | S3n | N1n | S3rn | N1n | N1n | N1n | N1tn | N1n | N1n | N1n | N1n | N1n | N1n | N1n | N1n | S3n | N1n | N1n | N1n | N1n | N1n | N1n | N1n | |
| Banapura | 155 | S3r | S2gt | S2rg | S2g | S2rg | S2rg | S3r | S2rg | S3t | S2rg | S2rg | S2g | S2rg | S2g | S2rg | S3r | S2rg | S3t | S2g | S2g | S2g | S2g | S2rg | S2g | S2g | S2g | S2g | S2g | S2g | S2rg | S2rg | S2g |
| Banapura | 156 | S3r | S2gt | S2rg | S2g | S2rg | S2rg | S3r | S2rg | S3t | S2rg | S2rg | S2g | S2rg | S2g | S2rg | S3r | S2rg | S3t | S2g | S2g | S2g | S2g | S2rg | S2g | S2g | S2g | S2g | S2g | S2g | S2rg | S2rg | S2g |
| Banapura | 157 | N1rn | S3n | N1n | S3n | N1n | S3n | N1rn | N1n | S3n | N1n | S3rn | N1n | N1n | N1n | N1tn | N1n | N1n | N1n | N1n | N1n | N1n | N1n | N1n | S3n | N1n | N1n | N1n | N1n | N1n | N1n | N1n | |
| Banapura | 158 | S2rg | S2g | S2g | S2g | S2g | S2gt | S2rg | S2g | S3t | S2g | S2g | S2g | S2g | S2g | S2g | S2rg | S2g | S2gt | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | |
| Banapura | 159/(1) | S2rg | S2g | S2g | S2g | S2g | S2gt | S2rg | S2g | S3t | S2g | S2g | S2g | S2g | S2g | S2g | S2rg | S2g | S2gt | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | |
| Banapura | 159/(2) | S2rg | S3g | S2g | S3g | S2g | S3g | S2rg | S2g | S3gt | S3g | S3g | S2g | S2g | S2g | S2g | S2rg | S2g | S2g | S3g | S3g | S3g | S3g | S2g | S3g | S3g | S3g | S3g | S3g | S3g | S2g | S3g | |
| Banapura | 160 | S3r | S2gt | S2rg | S2g | S2rg | S2rg | S3r | S2rg | S3t | S2rg | S2rg | S2g | S2rg | S2g | S2rg | S3r | S2rg | S3t | S2g | S2g | S2g | S2g | S2rg | S2g | S2g | S2g | S2g | S2g | S2rg | S2rg | S2g | |

| Village | Survey Number | Mango | Maize | Sapota | Sorghum | Guava | Cotton | Tamarind | Lime | Bengal gram | Sunflower | Red gram | Amla | Jackfruit | Custard-apple | Cashew | Jamun | Musambi | Groundnut | Chilly | Tomato | Marigold | Chrysanthemum | Pomegranate | Bajra | Jasmine | Bhendi | Brinjal | Crossandra | Drumstick | Mulberry | Onion | |
|----------|---------------|-------|-------|--------|---------|-------|--------|----------|------|-------------|-----------|----------|------|-----------|---------------|--------|-------|---------|-----------|--------|--------|----------|---------------|-------------|-------|---------|--------|---------|------------|-----------|----------|-------|-----|
| Banapura | 161 | S3r | S2gt | S2rg | S2g | S2rg | S2rg | S3r | S2rg | S3t | S2rg | S2rg | S2g | S2rg | S2g | S2rg | S3r | S2rg | S3t | S2g | S2g | S2g | S2g | S2rg | S2g | S2g | S2g | S2g | S2g | S2rg | S2rg | S2g | |
| Banapura | 162 | S3r | S2gt | S2rg | S2g | S2rg | S2rg | S3r | S2rg | S3t | S2rg | S2rg | S2g | S2rg | S2g | S2rg | S3r | S2rg | S3t | S2g | S2g | S2g | S2g | S2rg | S2g | S2g | S2g | S2g | S2g | S2rg | S2rg | S2g | |
| Banapura | 163 | S3r | S2gt | S2rg | S2g | S2rg | S2rg | S3r | S2rg | S3t | S2rg | S2rg | S2g | S2rg | S2g | S2rg | S3r | S2rg | S3t | S2g | S2g | S2g | S2g | S2rg | S2g | S2g | S2g | S2g | S2g | S2rg | S2rg | S2g | |
| Banapura | 164 | S3r | S2gt | S2rg | S2g | S2rg | S2rg | S3r | S2rg | S3t | S2rg | S2rg | S2g | S2rg | S2g | S2rg | S3r | S2rg | S3t | S2g | S2g | S2g | S2g | S2rg | S2g | S2g | S2g | S2g | S2g | S2rg | S2rg | S2g | |
| Banapura | 165 | S3r | S2gt | S2rg | S2g | S2rg | S2rg | S3r | S2rg | S3t | S2rg | S2rg | S2g | S2rg | S2g | S2rg | S3r | S2rg | S3t | S2g | S2g | S2g | S2g | S2rg | S2g | S2g | S2g | S2g | S2g | S2rg | S2rg | S2g | |
| Banapura | 166 | S2r | S1 | S1 | S1 | S1 | S2t | S2r | S1 | S3t | S1 | S1 | S1 | S1 | S1 | S1 | S2r | S1 | S2t | S1 | S1 | S1 | S1 | S1 | S1 | S1 | S1 | S1 | S1 | S1 | S1 | S1 | |
| Banapura | 167 | S2rg | S2g | S2g | S2g | S2g | S2gt | S2rg | S2g | S3t | S2g | S2g | S2g | S2g | S2g | S2g | S2rg | S2g | S2gt | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | |
| Banapura | 168 | S2rg | S2g | S2g | S2g | S2g | S2gt | S2rg | S2g | S3t | S2g | S2g | S2g | S2g | S2g | S2g | S2rg | S2g | S2gt | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | |
| Banapura | 169 | S2rg | S2g | S2g | S2g | S2g | S2gt | S2rg | S2g | S3t | S2g | S2g | S2g | S2g | S2g | S2g | S2rg | S2g | S2gt | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | |
| Banapura | 170 | S2r | S1 | S1 | S1 | S1 | S2t | S2r | S1 | S3t | S1 | S1 | S1 | S1 | S1 | S1 | S2r | S1 | S2t | S1 | S1 | S1 | S1 | S1 | S1 | S1 | S1 | S1 | S1 | S1 | S1 | S1 | |
| Banapura | 171 | S2rg | S2g | S2g | S2g | S2g | S2gt | S2rg | S2g | S3t | S2g | S2g | S2g | S2g | S2g | S2g | S2rg | S2g | S2gt | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | |
| Banapura | 172 | S2rg | S2g | S2g | S2g | S2g | S2gt | S2rg | S2g | S3t | S2g | S2g | S2g | S2g | S2g | S2g | S2rg | S2g | S2gt | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | |
| Banapura | 173 | S2rg | S2g | S2g | S2g | S2g | S2gt | S2rg | S2g | S3t | S2g | S2g | S2g | S2g | S2g | S2g | S2rg | S2g | S2gt | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | |
| Banapura | 174 | S2rg | S2g | S2g | S2g | S2g | S2gt | S2rg | S2g | S3t | S2g | S2g | S2g | S2g | S2g | S2g | S2rg | S2g | S2gt | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | |
| Banapura | 175 | S2r | S1 | S1 | S1 | S1 | S2t | S2r | S1 | S3t | S1 | S1 | S1 | S1 | S1 | S1 | S2r | S1 | S2t | S1 | S1 | S1 | S1 | S1 | S1 | S1 | S1 | S1 | S1 | S1 | S1 | S1 | |
| Banapura | 176 | S2rg | S3g | S2g | S3g | S2g | S3g | S2rg | S2g | S3gt | S3g | S3g | S2g | S2g | S2g | S2g | S2rg | S2g | S2g | S3g | S3g | S3g | S3g | S2g | S3g | S3g | S3g | S3g | S3g | S3g | S2g | S2g | S3g |
| Banapura | 177 | S2rg | S3g | S2g | S3g | S2g | S3g | S2rg | S2g | S3gt | S3g | S3g | S2g | S2g | S2g | S2g | S2rg | S2g | S2g | S3g | S3g | S3g | S3g | S2g | S3g | S3g | S3g | S3g | S3g | S3g | S2g | S2g | S3g |
| Banapura | 178 | S2rg | S3g | S2g | S3g | S2g | S3g | S2rg | S2g | S3gt | S3g | S3g | S2g | S2g | S2g | S2g | S2rg | S2g | S2g | S3g | S3g | S3g | S3g | S2g | S3g | S3g | S3g | S3g | S3g | S3g | S2g | S2g | S3g |
| Banapura | 179 | S2r | S1 | S1 | S1 | S1 | S2t | S2r | S1 | S3t | S1 | S1 | S1 | S1 | S1 | S1 | S2r | S1 | S2t | S1 | S1 | S1 | S1 | S1 | S1 | S1 | S1 | S1 | S1 | S1 | S1 | S1 | |
| Banapura | 180 | S2r | S1 | S1 | S1 | S1 | S2t | S2r | S1 | S3t | S1 | S1 | S1 | S1 | S1 | S1 | S2r | S1 | S2t | S1 | S1 | S1 | S1 | S1 | S1 | S1 | S1 | S1 | S1 | S1 | S1 | S1 | |
| Banapura | 181 | S2r | S1 | S1 | S1 | S1 | S2t | S2r | S1 | S3t | S1 | S1 | S1 | S1 | S1 | S1 | S2r | S1 | S2t | S1 | S1 | S1 | S1 | S1 | S1 | S1 | S1 | S1 | S1 | S1 | S1 | S1 | |
| Banapura | 182 | S2r | S1 | S1 | S1 | S1 | S2t | S2r | S1 | S3t | S1 | S1 | S1 | S1 | S1 | S1 | S2r | S1 | S2t | S1 | S1 | S1 | S1 | S1 | S1 | S1 | S1 | S1 | S1 | S1 | S1 | S1 | |
| Banapura | 183 | S2rg | S2g | S2g | S2g | S2g | S2gt | S2rg | S2g | S3t | S2g | S2g | S2g | S2g | S2g | S2g | S2rg | S2g | S2gt | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | |
| Banapura | 184 | S3r | S2g | S2rg | S2g | S2rg | S2rg | S3r | S2rg | S3t | S2rg | S2rg | S2g | S2rg | S2g | S2rg | S3r | S2rg | S2gt | S2g | S2g | S2g | S2g | S2rg | S2g | S2g | S2g | S2g | S2g | S2rg | S2rg | S2g | |
| Banapura | 185 | S2rg | S2g | S2g | S2g | S2g | S2gt | S2rg | S2g | S3t | S2g | S2g | S2g | S2g | S2g | S2g | S2rg | S2g | S2gt | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | |
| Banapura | 186 | S3r | S2g | S2rg | S2g | S2rg | S2rg | S3r | S2rg | S3t | S2rg | S2rg | S2g | S2rg | S2g | S2rg | S3r | S2rg | S2gt | S2g | S2g | S2g | S2g | S2rg | S2g | S2g | S2g | S2g | S2g | S2g | S2rg | S2rg | S2g |

| Village | Survey Number | Mango | Maize | Sapota | Sorghum | Guava | Cotton | Tamarind | Lime | Bengal gram | Sunflower | Red gram | Amla | Jackfruit | Custard-apple | Cashew | Jamun | Musambi | Groundnut | Chilly | Tomato | Marigold | Chrysanthemum | Pomegranate | Bajra | Jasmine | Bhendi | Brinjal | Crossandra | Drumstick | Mulberry | Onion | | | |
|----------|---------------|-------|-------|--------|---------|-------|--------|----------|------|-------------|-----------|----------|------|-----------|---------------|--------|-------|---------|-----------|--------|--------|----------|---------------|-------------|-------|---------|--------|---------|------------|-----------|----------|-------|-----|-----|-----|
| Banapura | 187 | S3r | S2g | S2rg | S2g | S2rg | S2rg | S3r | S2rg | S3t | S2rg | S2rg | S2g | S2rg | S2g | S2rg | S3r | S2rg | S2gt | S2g | S2g | S2g | S2g | S2rg | S2g | S2g | S2g | S2g | S2g | S2rg | S2rg | S2rg | S2g | | |
| Banapura | 188 | N1r | S2r | S3r | S2r | S3r | S2r | N1r | S3r | S3t | S3r | S3r | S2r | S3r | S2r | S3r | S3r | S3r | S2rt | S2r | S2r | S2r | S2r | S3r | S2r | S2r | S2r | S2r | S2r | S2r | S2r | S3r | S3r | S2r | |
| Banapura | 189 | N1r | S2r | S3r | S2r | S3r | S2r | N1r | S3r | S3t | S3r | S3r | S2r | S3r | S2r | S3r | S3r | S3r | S2rt | S2r | S2r | S2r | S2r | S3r | S2r | S2r | S2r | S2r | S2r | S2r | S2r | S2r | S3r | S3r | S2r |
| Banapura | 190 | S2rg | S2g | S2g | S2g | S2g | S2gt | S2rg | S2g | S3t | S2g | S2g | S2g | S2g | S2g | S2g | S2rg | S2g | S2gt | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | S2g | |
| Banapura | 191 | N1r | S2r | S3r | S2r | S3r | S2r | N1r | S3r | S3t | S3r | S3r | S2r | S3r | S2r | S3r | S3r | S3r | S2rt | S2r | S2r | S2r | S2r | S3r | S2r | S2r | S2r | S2r | S2r | S2r | S2r | S2r | S3r | S3r | S2r |
| Banapura | 192 | N1r | S2r | S3r | S2r | S3r | S2r | N1r | S3r | S3t | S3r | S3r | S2r | S3r | S2r | S3r | S3r | S3r | S2rt | S2r | S2r | S2r | S2r | S3r | S2r | S2r | S2r | S2r | S2r | S2r | S2r | S2r | S3r | S3r | S2r |
| Banapura | 195 | N1r | S2r | S3r | S2r | S3r | S2r | N1r | S3r | S3t | S3r | S3r | S2r | S3r | S2r | S3r | S3r | S3r | S2rt | S2r | S2r | S2r | S2r | S3r | S2r | S2r | S2r | S2r | S2r | S2r | S2r | S2r | S3r | S3r | S2r |
| Banapura | 196 | N1r | S2r | S3r | S2r | S3r | S2r | N1r | S3r | S3t | S3r | S3r | S2r | S3r | S2r | S3r | S3r | S3r | S2rt | S2r | S2r | S2r | S2r | S3r | S2r | S2r | S2r | S2r | S2r | S2r | S2r | S2r | S3r | S3r | S2r |

PART-B

SOCIO-ECONOMIC STATUS OF FARM HOUSEHOLDS

CONTENTS

| | | |
|---|---------------------------------------|-------|
| 1 | Findings of the socio-economic survey | 1-2 |
| 2 | Introduction | 3 |
| 3 | Methodology | 4-7 |
| 4 | Salient features of the survey | 8-24 |
| 5 | Summary | 26-29 |

LIST OF TABLES

| | | |
|------|--|----|
| 1 | Households sampled for socio economic survey | 8 |
| 2 | Population characteristics | 8 |
| 3 | Age wise classification of household members | 8 |
| 4 | Education level of household members | 9 |
| 5 | Occupation of household heads | 9 |
| 6 | Occupation of family members | 9 |
| 7 | Institutional participation of household members | 10 |
| 8 | Type of house owned by households | 10 |
| 9 | Durable assets owned by households | 10 |
| 10 | Average value of durable assets owned by households | 10 |
| 11 | Farm implements owned by households | 11 |
| 12 | Average value of farm implements | 11 |
| 13 | Livestock possession by households | 11 |
| 14 | Average labour availability | 11 |
| 15 | Adequacy of hired labour | 12 |
| 16 | Distribution of land (ha) | 12 |
| 17 | Average land value (Rs./ha) | 12 |
| 18 | Cropping pattern | 12 |
| 19 | Cropping intensity | 13 |
| 20 | Possession of bank account and saving | 13 |
| 21 | Borrowing status | 13 |
| 22 | Source of credit | 13 |
| 23 | Avg. credit borrowed | 13 |
| 24 | Purpose of credit borrowed from institutional sources | 14 |
| 25 | Repayment status of household from institutional sources | 14 |
| 26 | Opinion on institutional sources of credit | 14 |
| 27.a | Cost of cultivation of Maize | 15 |
| 27.b | Cost of cultivation of Sorghum | 16 |
| 27.c | Cost of cultivation of Bengal gram | 17 |
| 27.d | Cost of cultivation of Sunflower | 18 |
| 28 | Adequacy of fodder | 19 |

| | | |
|----|--|----|
| 29 | Annual gross income | 19 |
| 30 | Average annual expenditure | 19 |
| 31 | Horticultural species grown | 19 |
| 32 | Forest species grown | 20 |
| 33 | Average additional investment capacity | 20 |
| 34 | Source of funds for additional investment | 20 |
| 35 | Marketing of the agricultural produce | 20 |
| 36 | Marketing channels used for sale of agricultural produce | 21 |
| 37 | Mode of transport of agricultural produce | 21 |
| 38 | Incidence of soil and water erosion problems | 21 |
| 39 | Interest shown towards soil testing | 21 |
| 40 | Usage pattern of fuel for domestic use | 21 |
| 41 | Source of drinking water | 22 |
| 42 | Source of light | 22 |
| 43 | Existence of sanitary toilet facility | 22 |
| 44 | Possession of public distribution system (PDS) card | 22 |
| 45 | Participation in NREGA programme | 22 |
| 46 | Adequacy of food items | 23 |
| 47 | Inadequacy of food items | 23 |
| 48 | Farming constraints experienced | 24 |

FINDINGS OF THE SOCIO-ECONOMIC SURVEY

- ❖ *The survey was conducted in Haligeri-3 is located at North latitude 15^o 24' 35.308'' and 15^o 22' 52.919'' and East longitude 76^o 4' 6.789'' and 76^o 2' 12.128'' covering an area of about 719.28 ha coming under Halageri, Bhanapur, Veerapura and Tadakal villages of Koppal taluk.*
- ❖ *Socio-economic analysis of Haligeri-3 micro watersheds of Haligeri sub-watershed, Koppala taluk & District indicated that, out of the total sample of 35 total respondents, 12 (34.29 %) were marginal, 8 (22.86%) were small, 7 (20.00 %) were Semi medium and 3 (8.57 %) were medium farmers.*
- ❖ *The population characteristics of households indicated that, there were 66 (54.55%) men and 55 (45.45 %) were women.*
- ❖ *Majority of the respondents (36.36%) were in the age group of 16-35 years.*
- ❖ *Education level of the sample households indicated that, there were 36.36 per cent illiterates, 63.64 per cent pre university education and 6.61 per cent attained graduation.*
- ❖ *About, 85.71 per cent of household heads practicing agriculture and 14.29 per cent of the household heads were engaged as agricultural labourers. Agriculture was the major occupation for 24.79 per cent of the household members. In the study area, 42.86 per cent of the households possess katcha house and 2.86 per cent possess pucca house.*
- ❖ *The durable assets owned by the households showed that, 88.57 per cent possess TV, 82.86 per cent possess mobile phones and 5.71 per cent possess motor cycles.*
- ❖ *Farm implements owned by the households indicated that, 5.71 per cent possess tractor, 5.71 per cent possess bullock cart.*
- ❖ *Regarding livestock possession by the households, 2.86 per cent possess buffalo.*
- ❖ *The average labour availability in the study area showed that, own labour men available in the micro watershed was 2.85, women available in the micro watershed was 1.03, hired labour (men) available was 6.17 and hired labour (women) available was 6.37.*
- ❖ *Further, 97.14 per cent of the households opined that hired labour was inadequate during the agricultural season. Out of the total land holding of the sample respondents 90.10 per cent (57.23 ha) of the area is under dry condition.*
- ❖ *The major crops grown by sample farmers are Maize, Sorghum, Bengal gram and Sunflower and cropping intensity was recorded as 99.76 per cent.*
- ❖ *Out of the sample households 80.00 percent possessed bank account and 80.00 per cent of them have savings in the account.*
- ❖ *About 68.57 per cent of the respondents borrowed credit from various sources.*
- ❖ *Among the credit borrowed by households, 37.93 per cent have borrowed loan from commercial banks and 27.59 per cent from co-operative/Grameena bank.*

- ❖ Majority of the respondents (100.00%) have borrowed loan for agriculture purpose.
- ❖ Regarding the opinion on institutional sources of credit, 73.68 per cent of the households opined that credit helped to perform timely agricultural operations.
- ❖ The per hectare cost of cultivation for Maize, Sorghum, Bengal gram and Sunflower was Rs.26600.58, 32265.36, 29342.57 and 25256.57 with benefit cost ratio of 1:2.20, 1: 1.70, 1: 1.63 and 1: 1.10 respectively.
- ❖ Further, 60.00 per cent of the households opined that dry fodder was adequate and 5.71 per cent of the households have opined that the green fodder was adequate.
- ❖ The average annual gross income of the farmers was Rs. 81454.37 in micro-watershed, of which Rs. 74740.06 comes from agriculture.
- ❖ Sampled households have grown 3 horticulture trees and 16 forestry trees together in the fields and back yards.
- ❖ About 34.29 per cent of the households shown interest to cultivate horticultural crops. Households have an average investment capacity of Rs. 485.71 for land development. Source of funds for additional investment is concerned, 2.86 per cent depends on own funds and 5.71 per cent depends on bank loan for land development activities.
- ❖ Regarding marketing channels, 28.57 per cent of the households have sold agricultural produce to the local/village merchants, while, 57.14 per cent have sold in regulated markets. Further, 85.71 per cent of the households have used tractor for the transport of agriculture commodity.
- ❖ Majority of the farmers (85.71%) have experienced soil and water erosion problems in the watershed and 80.00 per cent of the households were interested towards soil testing.
- ❖ Fire was the major source of fuel for domestic use for 77.14 per cent of the households and 11.43 per cent households has LPG connection. Piped supply was the major source for drinking water for 88.57 per cent of the households. Electricity was the major source of light for 100.00 per cent of the households. In the study area, 34.29 per cent of the households possess toilet facility. Regarding possession of PDS card, 100.00 per cent of the households possessed BPL card.
- ❖ Households opined that, the requirement of cereals (88.57%), pulses (74.29%) and oilseeds (14.29%) are adequate for consumption.
- ❖ Farming constraints experienced by households in the micro watersheds were lower fertility status of the soil (77.14%) wild animal menace on farm field (82.86%), frequent incidence of pest and diseases (57.14%), inadequacy of irrigation water (34.29%), high cost of fertilizers and plant protection chemicals (28.57%), high rate of interest on credit (8.57%), low price for the agricultural commodities (11.43%), lack of marketing facilities in the area (14.29%), inadequate extension services (31.43%), lack of transport for safe transport of the agricultural produce to the market(40.00%).

INTRODUCTION

Soil and water are the two precious natural resources which are essential for crop production and existence of life on earth. Rainfed agriculture is under severe stress due to various constraints related to agriculture like uneven and erratic distribution of rainfall, indiscriminate use of fertilizers, chemicals and pesticides, adoption of improper land management practices, soil erosion, decline in soil fertility, decline in ground water resources leading to low crop productivity. The area under rainfed agriculture has to be managed effectively using the best available practices to enhance the production of food, fodder and fuel. This is possible if the land resources are characterized at each parcel of land through detailed land resource inventory using the best available techniques of remote sensing, GPS and GIS. The watershed development programs are aimed at the sustainable distribution of its resources and the process of creating and implementing plans, programs, and projects to sustain and enhance watershed functions that affect the plant, animal and human communities within a watershed boundary.

World Bank funded KWDP II, SUJALA III project was implemented in with Broad objective of demonstrating more effective watershed management through greater integration of programmes related to rain-fed agriculture, innovative and science based approaches and strengthen institutional capacities and If successful, it is expected that the systems and tools could be mainstreamed into the overall IWMP in the State of Karnataka and in time, throughout other IWMP operations in India. With this background the socio-economic survey has been carried out with following specific objectives:

1. To understand the demographic features of the households in the micro-watershed
2. To understand the extent of family labour available and additional employment opportunities available within the village.
3. To know the status of assets of households in the micro-watershed for suggesting possible improvements.
4. To study the cropping pattern, cropped area and productivity levels of different households in micro-watershed.
5. To determine the type and extent of livestock owned by different categories of HHs
6. Availability of fodder and level of livestock management.

Scope and importance of survey

Survey helps in identification of different socio-economic and resource use-patterns of farmers at the Micro watershed. Household survey provides demographic features, labour force, and levels of education; land ownership and asset position (including livestock and other household assets) of surveyed households; and cropping patterns, input intensities, and average crop yields from farmers' fields. It also discusses crop utilization and the degree of commercialization of production in the areas; farmers' access to and utilization of credit from formal and informal sources; and the level of adoption and use of soil, water, and pest management technologies.

METHODOLOGY

The description of the methods, components selected for the survey and procedures followed in conducting the baseline survey are furnished under the following heads.

g heads.

1. Description of the study area

Koppal district is an administrative district in the state of Karnataka in India. In the past Koppal was referred to as 'Kopana Nagara'. Koppal, now a district headquarters is ancient Kopana a major holy place of the Jainas. The district occupies an area of 7,190 km² and has a population of 1,196,089, which 16.58% were urban as of 2001. The Koppal district was formed after split of Raichur district.

Geographers are very particular about the physiography or relief of a region. It plays a very important role in the spatial analysis of agricultural situation of the study area. The undulating topography with black cotton soil shrips, cut across by numerous nalas or streams is the major characteristic feature of the study region. Three physiographic divisions have made considering the local conditions of landforms and crops grown in the district. On the basis of physiography, Koppal district can be divided into three major divisions. They are (a) Koppal & Yelburga plateau, (b) Maidan division, (c) Tungabhadra valley. The district is part of Krishna basin the main streams draining the area are Maskinala, Ilkal-nadi and Hirenala. These are Ephemeral in nature, these come under Tungabhadra sub-basin. The drainage exhibit dentritic to subdentritic with drainage density varies from 1.4 to 7.0kms/sq.km.

According to the 2011 census Koppal district has a population of 1,391,292, roughly equal to the nation of Swaziland or the US state of Hawaii. This gives it a ranking of 350th in India (out of a total of 640). The district has a population density of 250 inhabitants per square kilometre (650/sq mi). Its population growth rate over the decade 2001-2011 was 16.32%.Koppal has a sex ratio of 983 females for every 1000 males, and a literacy rate of 67.28%.

2. Locale of the survey and description of the micro-watershed and

The study was conducted in Haligeri-3 micro-watershed (Haligeri sub-watershed, Koppala taluk & District) is located at North latitude 15⁰ 24' 35.308" and 15⁰ 22' 52.919" and East longitude 76⁰ 4' 6.789" and 76⁰ 2' 12.128" covering an area of about 719.28 ha bounded by under Halageri, Bhanapur, Veerapura and Tadakal Villages.

3. Selection of the respondents for the study

The micro-watershed is marked with 320 square meters grids. One farmer from every alternate grid in the micro-watershed was selected for the study and interviewed for socio-economic data. Totally 35 households were interviewed for the survey.

4. The parameters considered for socio-economic survey of households

Two forms of data were collected from the micro-watershed which includes primary data from the farm households and secondary data about the villages under the micro-watershed jurisdiction.

The following parameters were considered for the primary data collection about the socio-economic data of the households, (1) Demographic information, (2) Farm and durable assets owned by households, (3) Livestock possession, (4) Labour availability, (5) Level of migration in the village, Land holding, (7) Cropping pattern, (8) Source of irrigation, (9) Borrowing status, (10) Cost of cultivation of major crops, (11) Economics of subsidiary activities, (12) Fodder availability, (13) Family annual income from different sources, (14) Horticulture and forestry species grown, (15) Additional investment capacity, (16) Marketing practices, (17) Status of soil and water conservation structure, (18) Access to basic needs and (19) Constraints and suggestion.

The following parameters were considered for the secondary data regarding the villages under the micro-watershed jurisdiction, (1) Number of villages in each micro-watershed jurisdiction, (2) Village wise number of households, (3) Geographical area of the villages, (4) Cultivable area including rainfed and irrigated, (5) Number and type of house in each village, (6) Human and livestock population, (7) Facilities in the village such as roads, transport facility for conveyance, drinking water supply, street light and (8) Community based organizations in the villages.

5. Development of interview schedule and data collection

Taking into the consideration the objectives of the survey, an interview schedule was prepared after thorough consultation with the experts in the field of social sciences. A comprehensive interview schedule covering all the major parameters for measuring the socio-economic situation was developed.

6. Tools used to analyze the data

The statistical components such as frequency and percentage were used to analyze the data.

Abbreviations used in the report

LL=Landless

MF=Marginal Farmers

SF=Small farmers

SMF=Semi medium farmers

MDF=Medium farmers

LF=Large Farmers

FINDINGS OF THE SURVEY

This chapter deals with systematic presentation of results of the survey. Keeping in view the objectives, the salient features of the survey are presented under the following headings.

Households sampled for socio-economic survey: The data on households sampled for socio economic survey in Haligeri-3 Micro watershed is presented in Table 1 and it indicated that 35 farmers were sampled in Haligeri-3 micro-watershed among households surveyed 12 (34.29%) were marginal, 8 (22.86%) were small, 7 (20.00 %) were semi medium and 3 (8.57 %) were medium farmers. 5 landless farmers were also interviewed for the survey.

Table 1. Households sampled for socio economic survey in Haligeri-3 micro-watershed

| Sl.No. | Particulars | LL (5) | | MF (12) | | SF (8) | | SMF (7) | | MDF (3) | | All (35) | |
|--------|-------------|--------|------|---------|------|--------|------|---------|----|---------|------|----------|-----|
| | | N | % | N | % | N | % | N | % | N | % | N | % |
| 1 | Farmers | 5 | 14.3 | 12 | 34.3 | 8 | 22.9 | 7 | 20 | 3 | 8.57 | 35 | 100 |

Population characteristics: The population characteristics of households sampled for socio-economic survey in Haligeri-3 Micro watershed is presented in Table 2. The data indicated that, there were 66 (54.55%) men and 55 (45.45%) were women.

Table 2. Population characteristics in Haligeri-3 micro-watershed

| Sl.No. | Particulars | LL (12) | | MF (41) | | SF (29) | | SMF (27) | | MDF (12) | | All (121) | |
|---------|-------------|---------|------|---------|-----|---------|-----|----------|------|----------|-----|-----------|------|
| | | N | % | N | % | N | % | N | % | N | % | N | % |
| 1 | Men | 7 | 58.3 | 20 | 49 | 14 | 48 | 16 | 59.3 | 9 | 75 | 66 | 54.6 |
| 2 | Women | 5 | 41.7 | 21 | 51 | 15 | 52 | 11 | 40.7 | 3 | 25 | 55 | 45.5 |
| Total | | 12 | 100 | 41 | 100 | 29 | 100 | 27 | 100 | 12 | 100 | 121 | 100 |
| Average | | 2.4 | | 3.4 | | 3.6 | | 3.9 | | 4.0 | | 3.5 | |

Age wise classification of population: The age wise classification of household members in Haligeri-3 Micro watershed is presented in Table 3. The indicated that, 17 (14.05%) of population were 0-15 years of age, 44 (36.36%) were 16-35 years of age, 50(41.32%) were 36-60 years of age and 10 (8.26 %) were above 61 years of age.

Table 3: Age wise classification of members of the household in Haligeri-3 micro-watershed

| Sl.No. | Particulars | LL (12) | | MF (41) | | SF (29) | | SMF (27) | | MDF (12) | | All (121) | |
|--------|--------------------|---------|------|---------|------|---------|------|----------|-------|----------|-----|-----------|-------|
| | | N | % | N | % | N | % | N | % | N | % | N | % |
| 1 | 0-15 years of age | 0 | 0 | 5 | 12.2 | 5 | 17.2 | 4 | 14.81 | 3 | 25 | 17 | 14.05 |
| 2 | 16-35 years of age | 4 | 33.3 | 16 | 39 | 14 | 48.3 | 8 | 29.63 | 2 | 17 | 44 | 36.36 |
| 3 | 36-60 years of age | 8 | 66.7 | 17 | 41.5 | 9 | 31 | 12 | 44.44 | 4 | 33 | 50 | 41.32 |
| 4 | > 61 years | 0 | 0 | 3 | 7.32 | 1 | 3.45 | 3 | 11.11 | 3 | 25 | 10 | 8.26 |
| Total | | 12 | 100 | 41 | 100 | 29 | 100 | 27 | 100 | 12 | 100 | 121 | 100 |

Education level of household members: Education level of household members in Haligeri-3 Micro watershed is presented in Table 4. The results indicated that, there were

36.36 per cent of illiterates, 23.14 per cent of them had primary school education, 4.96 per cent middle school education, 12.40 per cent high school education, 11.57 per cent of them had PUC education, 1.65 per cent of them had Diploma, 0.83 per of them had ITI, 6.61 per cent attained graduation, and 2.48 them had other education.

Table 4. Education level of members of the household in Haligeri-3 micro-watershed

| Sl.No. | Particulars | LL (12) | | MF (41) | | SF (29) | | SMF (27) | | MDF (12) | | All (121) | |
|--------|----------------|---------|------|---------|------|---------|------|----------|------|----------|-------|-----------|------|
| | | N | % | N | % | N | % | N | % | N | % | N | % |
| 1 | Illiterate | 11 | 91.7 | 19 | 46.3 | 11 | 37.9 | 3 | 11.1 | 0 | 0 | 44 | 36.4 |
| 2 | Primary School | 0 | 0 | 9 | 22 | 8 | 27.6 | 6 | 22.2 | 5 | 41.67 | 28 | 23.1 |
| 3 | Middle School | 0 | 0 | 2 | 4.88 | 2 | 6.9 | 2 | 7.41 | 0 | 0 | 6 | 4.96 |
| 4 | High School | 0 | 0 | 2 | 4.88 | 1 | 3.45 | 6 | 22.2 | 6 | 50 | 15 | 12.4 |
| 5 | PUC | 1 | 8.33 | 2 | 4.88 | 5 | 17.2 | 5 | 18.5 | 1 | 8.33 | 14 | 11.6 |
| 6 | Diploma | 0 | 0 | 0 | 0 | 2 | 6.9 | 0 | 0 | 0 | 0 | 2 | 1.65 |
| 7 | ITI | 0 | 0 | 1 | 2.44 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0.83 |
| 8 | Degree | 0 | 0 | 5 | 12.2 | 0 | 0 | 3 | 11.1 | 0 | 0 | 8 | 6.61 |
| 9 | Others | 0 | 0 | 1 | 2.44 | 0 | 0 | 2 | 7.41 | 0 | 0 | 3 | 2.48 |
| Total | | 12 | 100 | 41 | 100 | 29 | 100 | 27 | 100 | 12 | 100 | 121 | 100 |

Occupation of head of households: The data regarding the occupation of the household heads in Haligeri-3 Micro watershed is presented in Table 5. The results indicate that, 85.71 per cent of household's heads were practicing agriculture and 14.29 per cent of the household heads were agricultural Labour.

Table 5: Occupation of heads of households in Haligeri-3 micro-watershed

| Sl.No. | Particulars | LL (5) | | MF (12) | | SF (8) | | SMF (7) | | MDF (3) | | All (35) | |
|--------|---------------------|--------|-----|---------|-----|--------|-----|---------|-----|---------|-----|----------|-------|
| | | N | % | N | % | N | % | N | % | N | % | N | % |
| 1 | Agriculture | 2 | 40 | 11 | 92 | 8 | 100 | 6 | 86 | 3 | 100 | 30 | 85.71 |
| 2 | Agricultural Labour | 3 | 60 | 1 | 8.3 | 0 | 0 | 1 | 14 | 0 | 0 | 5 | 14.29 |
| Total | | 5 | 100 | 12 | 100 | 8 | 100 | 7 | 100 | 3 | 100 | 35 | 100 |

Occupation of the members of the household: The data regarding the occupation of the household members in Haligeri-3 Micro watershed is presented in Table 6. The results indicate that, agriculture was the major occupation for 24.79 per cent of the household members, 56.20 per cent were agricultural labour, 1.65 per cent were working in government sector, 14.88 per cent were working in pursuing education and 2.48 per cent were children's.

Table 6: Occupation of members of the household in Haligeri-3 micro-watershed

| Sl.No. | Particulars | LL (12) | | MF (41) | | SF (29) | | SMF (27) | | MDF (12) | | All (121) | |
|--------|---------------------|---------|------|---------|------|---------|-------|----------|-------|----------|-----|-----------|------|
| | | N | % | N | % | N | % | N | % | N | % | N | % |
| 1 | Agriculture | 2 | 16.7 | 11 | 26.8 | 8 | 27.59 | 6 | 22.22 | 3 | 25 | 30 | 24.8 |
| 2 | Agricultural Labour | 9 | 75 | 22 | 53.7 | 15 | 51.72 | 16 | 59.26 | 6 | 50 | 68 | 56.2 |
| 3 | Private Service | 0 | 0 | 1 | 2.44 | 0 | 0 | 1 | 3.7 | 0 | 0 | 2 | 1.65 |
| 4 | Student | 1 | 8.33 | 6 | 14.6 | 6 | 20.69 | 2 | 7.41 | 3 | 25 | 18 | 14.9 |
| 5 | Children | 0 | 0 | 1 | 2.44 | 0 | 0 | 2 | 7.41 | 0 | 0 | 3 | 2.48 |
| Total | | 12 | 100 | 41 | 100 | 29 | 100 | 27 | 100 | 12 | 100 | 121 | 100 |

Institutional Participation of household members: The data regarding the institutional participation of the household members in Haligeri-3 Micro watershed is presented in Table 7. The results show that, out of the total family members in the households 100 per cent were not participating in any of the institutions.

Table 7: Institutional Participation of household member in Haligeri-3 micro-watershed

| Sl.No. | Particulars | LL (12) | | MF (41) | | SF (29) | | SMF (27) | | MDF (12) | | All (121) | |
|--------|------------------|---------|-----|---------|-----|---------|-----|----------|-----|----------|-----|-----------|-----|
| | | N | % | N | % | N | % | N | % | N | % | N | % |
| 1 | No Participation | 12 | 100 | 41 | 100 | 29 | 100 | 27 | 100 | 12 | 100 | 121 | 100 |
| Total | | 12 | 100 | 41 | 100 | 29 | 100 | 27 | 100 | 12 | 100 | 121 | 100 |

Type of house owned: The data regarding the type of house owned by the households in Haligeri-3 Micro watershed is presented in Table 8. The results indicate that, 54.29 percent possess thatched house, 42.86 per cent of the households possess katcha house and 2.86 per cent possess pucca house.

Table 8. Type of house owned by households in Haligeri-3 micro-watershed

| Sl.No. | Particulars | LL (5) | | MF (12) | | SF (8) | | SMF (7) | | MDF (3) | | All (35) | |
|--------|-------------|--------|-----|---------|-----|--------|-----|---------|------|---------|-----|----------|-------|
| | | N | % | N | % | N | % | N | % | N | % | N | % |
| 1 | Thatched | 3 | 60 | 7 | 58 | 4 | 50 | 3 | 42.9 | 2 | 67 | 19 | 54.29 |
| 2 | Katcha | 2 | 40 | 4 | 33 | 4 | 50 | 4 | 57.1 | 1 | 33 | 15 | 42.86 |
| 3 | Pucca/RCC | 0 | 0 | 1 | 8.3 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 2.86 |
| Total | | 5 | 100 | 12 | 100 | 8 | 100 | 7 | 100 | 3 | 100 | 35 | 100 |

Durable assets owned by the households: The data regarding the Durable Assets owned by the households in Haligeri-3 Micro watershed is presented in Table 9. The result shows that, 88.57 per cent possess TV, 5.71 per cent possess Bicycle and motor cycle, 2.86 per cent possess Landline Phone and 82.86 per cent possess mobile phones.

Table 9. Durable assets owned by households in Haligeri-3 micro-watershed

| Sl.No. | Particulars | LL (5) | | MF (12) | | SF (8) | | SMF (7) | | MDF (3) | | All (35) | |
|--------|----------------|--------|----|---------|-----|--------|------|---------|-----|---------|-----|----------|-------|
| | | N | % | N | % | N | % | N | % | N | % | N | % |
| 1 | Television | 2 | 40 | 12 | 100 | 7 | 87.5 | 7 | 100 | 3 | 100 | 31 | 88.57 |
| 2 | Bicycle | 0 | 0 | 1 | 8.3 | 0 | 0 | 1 | 14 | 0 | 0 | 2 | 5.71 |
| 3 | Motor Cycle | 0 | 0 | 0 | 0 | 2 | 25 | 0 | 0 | 0 | 0 | 2 | 5.71 |
| 4 | Landline Phone | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 14 | 0 | 0 | 1 | 2.86 |
| 5 | Mobile Phone | 2 | 40 | 12 | 100 | 6 | 75 | 6 | 86 | 3 | 100 | 29 | 82.86 |

Table 10. Average value of durable assets owned in Haligeri-3 micro-watershed

Average Value (Rs.)

| Sl.No. | Particulars | LL (5) | MF (12) | SF (8) | SMF (7) | MDF (3) | All (35) |
|--------|----------------|--------|---------|--------|---------|---------|----------|
| 1 | Television | 5000 | 30666 | 4071 | 4571 | 13666 | 15467 |
| 2 | Bicycle | 0 | 30000 | 0 | 2500 | 0 | 16250 |
| 3 | Motor Cycle | 0 | 0 | 6000 | 0 | 0 | 6000 |
| 4 | Landline Phone | 0 | 0 | 0 | 1000 | 0 | 1000 |
| 5 | Mobile Phone | 1250 | 3041 | 2500 | 4083 | 1833 | 2896 |

Average value of durable assets: The data regarding the average value of durable assets owned by the households in Haligeri-3 Micro watershed is presented in Table 10. The result shows that, the average value of television was Rs.15467.00, bicycle was Rs.16250.00, motor cycle was Rs. 6000.00, Landline Phone was Rs. 1000.00 and mobile phone was Rs.2896.00.

Farm implements owned: The data regarding the farm implements owned by the households in Haligeri-3 Micro watershed is presented in Table 11. About 5.71 per cent of the households possess Bullock Cart and tractor, 8.57 per cent possess Weeder, 5.71 per cent possess tractor.

Table 11. Farm implements owned in Haligeri-3 micro-watershed

| Sl.No. | Particulars | LL (5) | | MF (12) | | SF (8) | | SMF (7) | | MDF (3) | | All (35) | |
|--------|--------------|--------|---|---------|------|--------|---|---------|------|---------|------|----------|------|
| | | N | % | N | % | N | % | N | % | N | % | N | % |
| 1 | Bullock Cart | 0 | 0 | 1 | 8.33 | 0 | 0 | 1 | 14.3 | 0 | 0 | 2 | 5.71 |
| 2 | Tractor | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 2 | 66.7 | 2 | 5.71 |
| 8 | Weeder | 0 | 0 | 3 | 25 | 0 | 0 | 0 | 0 | 0 | 0 | 3 | 8.57 |

Average value of farm implements: The data regarding the average value of farm Implements owned by the households in Haligeri-3 Micro watershed is presented in Table 12. The results show that the average value of bullock Cart was Rs.20000.00, weeder was Rs.116.00 and tractor was Rs. 253000.

Table 12. Average value of farm implements in Haligeri-3 micro-watershed

Average Value (Rs.)

| Sl.No. | Particulars | LL (5) | MF (12) | SF (8) | SMF (7) | MDF (3) | All (35) |
|--------|--------------|--------|---------|--------|---------|---------|----------|
| 1 | Bullock Cart | 0 | 25000 | 0 | 15000 | 0 | 20000 |
| 2 | Tractor | 0 | 0 | 0 | 0 | 253000 | 253000 |
| 3 | Weeder | 0 | 116 | 0 | 0 | 0 | 116 |

Livestock possession by the households: The data regarding the Livestock possession by the households in Haligeri-3 Micro watershed is presented in Table 13. The results indicate that, 2.86 per cent of the households possess bullocks, buffalo and crossbred cow.

Table 13. Livestock possession by households in Haligeri-3 micro-watershed

| Sl.No. | Particulars | LL (5) | | MF (12) | | SF (8) | | SMF (7) | | MDF (3) | | All (35) | |
|--------|---------------|--------|-----|---------|-----|--------|-----|---------|----|---------|-----|----------|-------|
| | | N | % | N | % | N | % | N | % | N | % | N | % |
| 1 | Bullock | 0 | 0 | 1 | 8.3 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 2.86 |
| 2 | Crossbred cow | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 14 | 0 | 0 | 1 | 2.86 |
| 3 | Buffalo | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 14 | 0 | 0 | 1 | 2.86 |
| 4 | blank | 5 | 100 | 11 | 92 | 8 | 100 | 5 | 71 | 3 | 100 | 32 | 91.43 |

Table 14. Average labour availability in Haligeri-3 micro-watershed

| Sl.No. | Particulars | LL (5) | MF (12) | SF (8) | SMF (7) | MDF (3) | All (35) |
|--------|---------------------|--------|---------|--------|---------|---------|----------|
| 1 | Hired labour Female | 1.4 | 8.67 | 7 | 6.71 | 3 | 6.37 |
| 2 | Own Labour Female | 1 | 1.08 | 1 | 1 | 1 | 1.03 |
| 3 | Own labour Male | 11 | 0.91 | 1.88 | 2 | 1 | 2.85 |
| 4 | Hired labour Male | 1.4 | 8.08 | 7 | 6.71 | 3 | 6.17 |

Average Labour availability: The data regarding the average labour availability in Haligeri-3 Micro watershed is presented in Table 14. The indicated that, own labour men available in the micro watershed was 2.85, women available in the micro watershed was 1.03, hired labour (men) available was 6.17 and hired labour (women) available was 6.37.

Adequacy of hired labour: The data regarding the adequacy of hired labour in Haligeri-3 Micro watershed is presented in Table 15. The results indicate that, 2.86 per cent of the household opined that hired labour was adequate, 97.14 per cent of the household opined that hired labour was Inadequate.

Table 15. Adequacy of hired labour in Haligeri-3 micro-watershed

| Sl.No. | Particulars | LL (5) | | MF (12) | | SF (8) | | SMF (7) | | MDF (3) | | All (35) | |
|--------|-------------|--------|----|---------|-----|--------|-----|---------|-----|---------|-----|----------|------|
| | | N | % | N | % | N | % | N | % | N | % | N | % |
| 1 | Adequate | 1 | 20 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 2.86 |
| 2 | Inadequate | 4 | 80 | 12 | 100 | 8 | 100 | 7 | 100 | 3 | 100 | 34 | 97.1 |

Distribution of land (ha): The data regarding the distribution of land (ha) in Haligeri-3 Micro watershed is presented in Table 16. The results indicate that, 51.56 ha (90.10%) of dry land and.

Table 16. Distribution of land (ha) in Haligeri-3 micro-watershed

| Sl.No. | Particulars | LL (5) | | MF (12) | | SF (8) | | SMF (7) | | MDF (3) | | All (35) | |
|--------|------------------|--------|-----|---------|-----|--------|-----|---------|-----|---------|------|----------|------|
| | | N | % | N | % | N | % | N | % | N | % | N | % |
| 1 | Dry | 0 | 0 | 9.08 | 100 | 10.92 | 100 | 19.42 | 100 | 12.14 | 68.2 | 51.56 | 90.1 |
| 3 | Permanent Fallow | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 5.67 | 31.8 | 5.67 | 9.9 |
| Total | | 0 | 100 | 9.08 | 100 | 10.92 | 100 | 19.42 | 100 | 17.81 | 100 | 57.23 | 100 |

Average value of land (ha): The data regarding the average land value (Rs./ha) in Haligeri-3 Micro watershed is presented in Table 17. The results show that the average value of dry land was Rs.240389.29 and the average value of permanent fallow land was Rs. 35285.71.

Table 17. Average value of land (ha) in Haligeri-3 micro-watershed

| Sl.No. | Particulars | LL (5) | MF (12) | SF (8) | SMF (7) | MDF (3) | All (35) |
|--------|------------------|--------|----------|----------|----------|----------|----------|
| 1 | Dry | 0 | 539349.4 | 311152.3 | 144143.4 | 107033.3 | 240389.3 |
| 3 | Permanent Fallow | 0 | 0 | 0 | 0 | 35285.71 | 35285.71 |

Cropping pattern: The data regarding the cropping pattern in Haligeri-3 Micro watershed is presented in Table 18. The results indicate that, farmers have grown Bengal gram (18.6 ha), Maize (17.24 ha), Sorghum (13.45 ha) and Sunflower (0.89 ha).

Table 18. Cropping pattern in Haligeri-3 micro-watershed

| Sl.No. | Particulars | LL (5) | MF (12) | SF (8) | SMF (7) | MDF (3) | All (35) |
|--------|----------------------|--------|---------|--------|---------|---------|----------|
| 1 | Kharif - Bengal gram | 0 | 4.32 | 4.17 | 6.07 | 4.05 | 18.6 |
| 2 | Kharif - Maize | 0 | 1.37 | 3.24 | 4.54 | 8.1 | 17.24 |
| 3 | Kharif - Sorghum | 0 | 2.32 | 3.32 | 7.81 | 0 | 13.45 |
| 4 | Kharif - Sunflower | 0 | 0.89 | 0 | 0 | 0 | 0.89 |
| Total | | 0 | 8.89 | 10.72 | 18.42 | 12.15 | 50.18 |

Cropping intensity: The data regarding the cropping intensity in Haligeri-3 Micro watershed is presented in Table 19. The results indicate that, the cropping intensity was 99.76 per cent.

Table 19. Cropping intensity (%) in Haligeri-3 micro-watershed

| Sl.No. | Particulars | LL (5) | MF (12) | SF (8) | SMF (7) | MDF (3) | All (35) |
|--------|--------------------|--------|---------|--------|---------|---------|----------|
| 1 | Cropping Intensity | 0 | 100 | 100 | 99.34 | 100 | 99.76 |

Possession of bank account and savings: The data regarding the possession of bank account and saving in Haligeri-3 micro-watershed is presented in Table 20. The results indicate that, 80.00 cent of the households posses bank account and savings.

Table 20. Possession of Bank account and savings in Haligeri-3 micro-watershed

| Sl.No. | Particulars | LL (5) | | MF (12) | | SF (8) | | SMF (7) | | MDF (3) | | All (35) | |
|--------|-------------|--------|---|---------|-----|--------|------|---------|-------|---------|-----|----------|----|
| | | N | % | N | % | N | % | N | % | N | % | N | % |
| 1 | Account | 0 | 0 | 12 | 100 | 7 | 87.5 | 6 | 85.71 | 3 | 100 | 28 | 80 |
| 2 | Savings | 0 | 0 | 12 | 100 | 7 | 87.5 | 6 | 85.71 | 3 | 100 | 28 | 80 |

Borrowing status: The data regarding the borrowing status in Haligeri-3 micro-watershed is presented in Table 21. The results indicate that, 68.57 percent of the sample farmers have borrowed credit from different sources.

Table 21. Borrowing status in Haligeri-3 micro-watershed

| Sl.No. | Particulars | LL (5) | | MF (12) | | SF (8) | | SMF (7) | | MDF (3) | | All (35) | |
|--------|---------------|--------|---|---------|-----|--------|-----|---------|------|---------|-----|----------|-------|
| | | N | % | N | % | N | % | N | % | N | % | N | % |
| 1 | Credit Aailed | 0 | 0 | 12 | 100 | 8 | 100 | 1 | 14.3 | 3 | 100 | 24 | 68.57 |

Source of credit: The data regarding the source of credit aailed by households in Haligeri-3 micro-watershed is presented in Table 22. The results show that, 37.93 per cent have borrowed loan from commercial banks and 27.59 per cent have borrowed loan from Grameena Bank.

Table 22. Source of credit borrowed by households in Haligeri-3 micro-watershed

| Sl.No. | Particulars | LL (0) | | MF (12) | | SF (8) | | SMF (6) | | MDF (3) | | All (29) | |
|--------|-----------------|--------|---|---------|----|--------|------|---------|----|---------|-------|----------|-------|
| | | N | % | N | % | N | % | N | % | N | % | N | % |
| 1 | Commercial Bank | 0 | 0 | 3 | 25 | 2 | 25 | 5 | 83 | 1 | 33.33 | 11 | 37.93 |
| 2 | Grameena Bank | 0 | 0 | 3 | 25 | 3 | 37.5 | 1 | 17 | 1 | 33.33 | 8 | 27.59 |

Avg. Credit amount: The data regarding the avg. Credit amount in Haligeri-3 micro-watershed is presented in Table 23. The results show that, farmers have borrowed Avg. Credit of Rs.36896.55 from different sources.

Table 23. Avg. Credit amount in Haligeri-3 micro-watershed

| Sl.No. | Particulars | MF (12) | SF (8) | SMF (6) | MDF (3) | All (29) |
|--------|----------------|---------|--------|---------|---------|----------|
| 1 | Average Credit | 17916.7 | 23750 | 52500 | 116667 | 36896.6 |

Purpose of credit borrowed (institutional Source): The data regarding the purpose of credit borrowed - Institutional Credit in Haligeri-3 micro-watershed is presented in Table

24. The results indicate that, 100.00 per cent of the households have borrowed loan for agriculture.

Table 24. Purpose of credit borrowed (institutional Source) by households in Haligeri-3 micro-watershed

| SN | Particulars | MF (6) | | SF (5) | | SMF (6) | | MDF (2) | | All (19) | |
|----|------------------------|--------|-----|--------|-----|---------|-----|---------|-----|----------|-----|
| | | N | % | N | % | N | % | N | % | N | % |
| 1 | Agriculture production | 6 | 100 | 5 | 100 | 6 | 100 | 2 | 100 | 19 | 100 |

Repayment status of household (institutional Source): The data regarding the repayment status of credit borrowed from institutional Source by households in Haligeri-3 micro watershed is presented in Table 25. The results indicate that, 100.00 per cent have unpaid.

Table 25. Repayment status of household (institutional Source) in Haligeri-3 micro-watershed

| Sl.No. | Particulars | MF (6) | | SF (5) | | SMF (6) | | MDF (2) | | All (19) | |
|--------|-------------|--------|-----|--------|-----|---------|-----|---------|-----|----------|-----|
| | | N | % | N | % | N | % | N | % | N | % |
| 1 | Un paid | 6 | 100 | 5 | 100 | 6 | 100 | 2 | 100 | 19 | 100 |

Opinion regarding institutional sources of credit: The data regarding the opinion on institutional sources of credit in Haligeri-3 micro watershed is presented in Table 26. The results indicate that, 73.68 per cent of the households opined that credit helped to perform timely agricultural operations, 15.8 per cent of the households opined that easy accessibility of credit, 5.26 per cent higher rate of interest and Forced to sell the produce at low price to repay loan in time.

Table 26. Opinion regarding institutional sources of credit in Haligeri-3 micro-watershed

| Sl.N | Particulars | MF (6) | | SF (5) | | SMF (6) | | MDF (2) | | All (19) | |
|------|---|--------|------|--------|----|---------|-----|---------|----|----------|------|
| | | N | % | N | % | N | % | N | % | N | % |
| 1 | Helped to perform timely agricultural operations | 3 | 50 | 4 | 80 | 6 | 100 | 1 | 50 | 14 | 73.7 |
| 2 | Easy accessibility of credit | 1 | 16.7 | 1 | 20 | 0 | 0 | 1 | 50 | 3 | 15.8 |
| 3 | Higher rate of interest | 1 | 16.7 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 5.26 |
| 4 | Forced to sell the produce at low price to repay loan in time | 1 | 16.7 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 5.26 |

Cost of Cultivation of Maize: The data regarding the cost of cultivation (Rs/ha) of Maize in Haligeri-3 micro watershed is presented in Table 27.a. The results indicate that, the total cost of cultivation (Rs/ha) for Maize was Rs. 26600.58. The gross income realized by the farmers was Rs. 57689.05. The net income from Maize cultivation was Rs.31088.46, thus the benefit cost ratio was found to be 1:2.20.

Table 27(a). Cost of Cultivation of Maize in Haligeri-3 micro-watershed

| Sl.N | Particulars | Units | Phy Units | Value (Rs.) | % to C3 |
|------------|--|--------------------------------|-----------|-------------|----------|
| I | Cost A1 | | | | |
| 1 | Hired Human Labour | Man days | 21.12 | 5089.32 | 19.13 |
| 2 | Bullock | Pairs/day | 2.25 | 1224.67 | 4.6 |
| 3 | Tractor | Hours | 0.25 | 190.3 | 0.72 |
| 4 | Machinery | Hours | 0 | 0 | 0 |
| 5 | Seed Main Crop (Establishment and Maintenance) | Kgs (Rs.) | 28.08 | 3370 | 12.67 |
| 7 | FYM | Quintal | 0 | 0 | 0 |
| 8 | Fertilizer + micronutrients | Quintal | 6.08 | 4433.19 | 16.67 |
| 9 | Pesticides (PPC) | Kgs /liters | 1.84 | 3005.79 | 11.3 |
| 10 | Irrigation | Number | 0 | 0 | 0 |
| 11 | Repairs | | 0 | 0 | 0 |
| 12 | Msc. Charges (Marketing costs etc) | | 0 | 0 | 0 |
| 13 | Depreciation charges | | 0 | 308.77 | 1.16 |
| 14 | Land revenue and Taxes | | 0 | 0 | 0 |
| II | Cost B1 | | | | |
| 16 | Interest on working capital | | | 1298.28 | 4.88 |
| 17 | Cost B1 = (Cost A1 + sum of 15 and 16) | | | 18920.31 | 71.13 |
| III | Cost B2 | | | | |
| 18 | Rental Value of Land | | | 147.92 | 0.56 |
| 19 | Cost B2 = (Cost B1 + Rental value) | | | 19068.23 | 71.68 |
| IV | Cost C1 | | | | |
| 20 | Family Human Labour | | 19.84 | 5104.12 | 19.19 |
| 21 | Cost C1 = (Cost B2 + Family Labour) | | | 24172.35 | 90.87 |
| V | Cost C2 | | | | |
| 22 | Risk Premium | | | 10 | 0.04 |
| 23 | Cost C2 = (Cost C1 + Risk Premium) | | | 24182.35 | 90.91 |
| VI | Cost C3 | | | | |
| 24 | Managerial Cost | | | 2418.23 | 9.09 |
| 25 | Cost C3 = (Cost C2 + Managerial Cost) | | | 26600.58 | 100 |
| VII | Economics of the Crop | | | | |
| a. | Main Product | a) Main Product (q) | | 35.76 | 57221.35 |
| | | b) Main Crop Sales Price (Rs.) | | | 1600 |
| | By Product | e) Main Product (q) | | 7.48 | 467.69 |
| | | f) Main Crop Sales Price (Rs.) | | | 62.5 |
| b. | Gross Income (Rs.) | | | 57689.05 | |
| c. | Net Income (Rs.) | | | 31088.46 | |
| d. | Cost per Quintal (Rs./q.) | | | 743.79 | |
| e. | Benefit Cost Ratio (BC Ratio) | | | 1:2.2 | |

Cost of Cultivation of Sorghum: The data regarding the cost of cultivation (Rs/ha) of Sorghum in Haligeri-3 micro watershed is presented in Table 27.b. The results indicate that, the total cost of cultivation (Rs/ha) for Sorghum was Rs. 32265.36. The gross income realized by the farmers was Rs. 55031.84. The net income from Sorghum cultivation was Rs.22766.48, thus the benefit cost ratio was found to be 1:1.70.

Table 27(b). Cost of Cultivation of Sorghum in Haligeri-3 micro-watershed

| Sl.No | Particulars | Units | Phy Units | Value(Rs.) | % to C3 |
|----------------------------------|--|--------------------------------|-----------|------------|----------|
| I Cost A1 | | | | | |
| 1 | Hired Human Labour | Man days | 40.17 | 8612.72 | 26.69 |
| 2 | Bullock | Pairs/day | 1.76 | 967.17 | 3 |
| 3 | Tractor | Hours | 2.78 | 2081.98 | 6.45 |
| 4 | Machinery | Hours | 0 | 0 | 0 |
| 5 | Seed Main Crop (Establishment and Maintenance) | Kgs (Rs.) | 14.14 | 1607.16 | 4.98 |
| 7 | FYM | Quintal | 11.78 | 1178.06 | 3.65 |
| 8 | Fertilizer + micronutrients | Quintal | 5.37 | 3737.91 | 11.58 |
| 9 | Pesticides (PPC) | Kgs / liters | 2.62 | 3030.2 | 9.39 |
| 10 | Irrigation | Number | 0 | 0 | 0 |
| 11 | Repairs | | 0 | 0 | 0 |
| 12 | Msc. Charges (Marketing costs etc) | | 0 | 0 | 0 |
| 13 | Depreciation charges | | 0 | 14.59 | 0.05 |
| 14 | Land revenue and Taxes | | 0 | 0 | 0 |
| II Cost B1 | | | | | |
| 16 | Interest on working capital | | | 1147.49 | 3.56 |
| 17 | Cost B1 = (Cost A1 + sum of 15 and 16) | | | 22377.29 | 69.35 |
| III Cost B2 | | | | | |
| 18 | Rental Value of Land | | | 166.67 | 0.52 |
| 19 | Cost B2 = (Cost B1 + Rental value) | | | 22543.95 | 69.87 |
| IV Cost C1 | | | | | |
| 20 | Family Human Labour | | 26.84 | 6779.09 | 21.01 |
| 21 | Cost C1 = (Cost B2 + Family Labour) | | | 29323.04 | 90.88 |
| V Cost C2 | | | | | |
| 22 | Risk Premium | | | 9.1 | 0.03 |
| 23 | Cost C2 = (Cost C1 + Risk Premium) | | | 29332.14 | 90.91 |
| VI Cost C3 | | | | | |
| 24 | Managerial Cost | | | 2933.21 | 9.09 |
| 25 | Cost C3 = (Cost C2 + Managerial Cost) | | | 32265.36 | 100 |
| VII Economics of the Crop | | | | | |
| a. | Main Product | a) Main Product (q) | | 33.86 | 53842.87 |
| | | b) Main Crop Sales Price (Rs.) | | | 1590 |
| a. | By Product | e) Main Product (q) | | 19.82 | 1188.97 |
| | | f) Main Crop Sales Price (Rs.) | | | 60 |
| b. | Gross Income (Rs.) | | | 55031.84 | |
| c. | Net Income (Rs.) | | | 22766.48 | |
| d. | Cost per Quintal (Rs./q.) | | | 952.81 | |
| e. | Benefit Cost Ratio (BC Ratio) | | | 1:1.7 | |

Cost of Cultivation of Bengal gram: The data regarding the cost of cultivation (Rs/ha) of Bengal gram in Haligeri-3 micro watershed is presented in Table 27.c. The results indicate, the total cost of cultivation (Rs/ha) for Bengal gram was Rs.29342.57. The gross income realized by the farmers was Rs. 47720.52. The net income from Bengal gram cultivation was Rs. 18377.95, thus the benefit cost ratio was found to be 1:1.63.

Table 27(c). Cost of Cultivation of Bengal gram in Haligeri-3 micro-watershed

| Sl.No | Particulars | Units | Phy Units | Value(Rs.) | % to C3 |
|------------|--|--------------------------------|-----------|------------|---------|
| I | Cost A1 | | | | |
| 1 | Hired Human Labour | Man days | 24.93 | 5512.36 | 18.79 |
| 2 | Bullock | Pairs/day | 1.16 | 639.4 | 2.18 |
| 3 | Tractor | Hours | 2.67 | 2003.49 | 6.83 |
| 4 | Machinery | Hours | 0.25 | 148.2 | 0.51 |
| 5 | Seed Main Crop (Establishment and Maintenance) | Kgs (Rs.) | 47.89 | 4616.08 | 15.73 |
| 7 | FYM | Quintal | 7.62 | 1524.69 | 5.2 |
| 8 | Fertilizer + micronutrients | Quintal | 3.81 | 2628.7 | 8.96 |
| 9 | Pesticides (PPC) | Kgs / liters | 1.83 | 2900.33 | 9.88 |
| 10 | Irrigation | Number | 0 | 0 | 0 |
| 11 | Repairs | | 0 | 0 | 0 |
| 12 | Msc. Charges (Marketing costs etc) | | 0 | 0 | 0 |
| 13 | Depreciation charges | | 0 | 3.23 | 0.01 |
| 14 | Land revenue and Taxes | | 0 | 0 | 0 |
| II | Cost B1 | | | | |
| 16 | Interest on working capital | | | 1401.58 | 4.78 |
| 17 | Cost B1 = (Cost A1 + sum of 15 and 16) | | | 21378.05 | 72.86 |
| III | Cost B2 | | | | |
| 18 | Rental Value of Land | | | 166.67 | 0.57 |
| 19 | Cost B2 = (Cost B1 + Rental value) | | | 21544.72 | 73.42 |
| IV | Cost C1 | | | | |
| 20 | Family Human Labour | | 20.12 | 5120.35 | 17.45 |
| 21 | Cost C1 = (Cost B2 + Family Labour) | | | 26665.06 | 90.88 |
| V | Cost C2 | | | | |
| 22 | Risk Premium | | | 10 | 0.03 |
| 23 | Cost C2 = (Cost C1 + Risk Premium) | | | 26675.06 | 90.91 |
| VI | Cost C3 | | | | |
| 24 | Managerial Cost | | | 2667.51 | 9.09 |
| 25 | Cost C3 = (Cost C2 + Managerial Cost) | | | 29342.57 | 100 |
| VII | Economics of the Crop | | | | |
| a. | Main Product | a) Main Product (q) | | 15 | 47400 |
| | | b) Main Crop Sales Price (Rs.) | | | 3160 |
| | By Product | e) Main Product (q) | | 6.41 | 320.52 |
| | | f) Main Crop Sales Price (Rs.) | | | 50 |
| b. | Gross Income (Rs.) | | | 47720.52 | |
| c. | Net Income (Rs.) | | | 18377.95 | |
| d. | Cost per Quintal (Rs./q.) | | | 606.36 | |
| e. | Benefit Cost Ratio (BC Ratio) | | | 1:1.6 | |

Cost of Cultivation of Sunflower: The data regarding the cost of cultivation (Rs/ha) of Sunflower in Haligeri-3 micro watershed is presented in Table 27.d. The results indicate that, the total cost of cultivation (Rs/ha) for Sunflower was Rs. 25256.57. The gross income realized by the farmers was Rs.28196.35. The net income from Sunflower cultivation was Rs. 2939.78, thus the benefit cost ratio was found to be 1:1.10.

Table 27(d). Cost of Cultivation of Sunflower in Haligeri-3 micro-watershed

| Sl.No | Particulars | Units | Phy Units | Value(Rs.) | % to C3 |
|------------|--|--------------------------------|-----------|------------|---------|
| I | Cost A1 | | | | |
| 1 | Hired Human Labour | Man days | 36.09 | 8571.69 | 33.94 |
| 2 | Bullock | Pairs/day | 0 | 0 | 0 |
| 3 | Tractor | Hours | 0 | 0 | 0 |
| 4 | Machinery | Hours | 1.13 | 676.71 | 2.68 |
| 5 | Seed Main Crop (Establishment and Maintenance) | Kgs (Rs.) | 11.28 | 2819.63 | 11.16 |
| 6 | Seed Inter Crop | Kgs. | 0 | 0 | 0 |
| 7 | FYM | Quintal | 0 | 0 | 0 |
| 8 | Fertilizer + micronutrients | Quintal | 4.51 | 3157.99 | 12.5 |
| 9 | Pesticides (PPC) | Kgs / liters | 2.26 | 2481.28 | 9.82 |
| 10 | Irrigation | Number | 0 | 0 | 0 |
| 11 | Repairs | | 0 | 0 | 0 |
| 12 | Msc. Charges (Marketing costs etc) | | 0 | 0 | 0 |
| 13 | Depreciation charges | | 0 | 563.93 | 2.23 |
| 14 | Land revenue and Taxes | | 0 | 0 | 0 |
| II | Cost B1 | | | | |
| 16 | Interest on working capital | | | 1016.27 | 4.02 |
| 17 | Cost B1 = (Cost A1 + sum of 15 and 16) | | | 19287.5 | 76.37 |
| III | Cost B2 | | | | |
| 18 | Rental Value of Land | | | 166.67 | 0.66 |
| 19 | Cost B2 = (Cost B1 + Rental value) | | | 19454.17 | 77.03 |
| IV | Cost C1 | | | | |
| 20 | Family Human Labour | | 13.53 | 3496.35 | 13.84 |
| 21 | Cost C1 = (Cost B2 + Family Labour) | | | 22950.51 | 90.87 |
| V | Cost C2 | | | | |
| 22 | Risk Premium | | | 10 | 0.04 |
| 23 | Cost C2 = (Cost C1 + Risk Premium) | | | 22960.51 | 90.91 |
| VI | Cost C3 | | | | |
| 24 | Managerial Cost | | | 2296.05 | 9.09 |
| 25 | Cost C3 = (Cost C2 + Managerial Cost) | | | 25256.57 | 100 |
| VII | Economics of the Crop | | | | |
| a. | Main Product | a) Main Product (q) | 5.64 | 28196.35 | |
| | | b) Main Crop Sales Price (Rs.) | | 5000 | |
| b. | Gross Income (Rs.) | | | 28196.35 | |
| c. | Net Income (Rs.) | | | 2939.78 | |
| d. | Cost per Quintal (Rs./q.) | | | 4478.69 | |
| e. | Benefit Cost Ratio (BC Ratio) | | | 1:1.1 | |

Adequacy of fodder: The data regarding the adequacy of fodder in Haligeri-3 Micro watershed is presented in Table 28. The results indicate that, 60.00 per cent of the households opined that dry fodder was adequate. With respect to green fodder availability, 5.71 percent of them opined it was sufficient and 8.57 percent of them opined it was insufficient.

Table 28. Adequacy of fodder in Haligeri-3 micro-watershed

| Sl.No. | Particulars | LL (5) | | MF (12) | | SF (8) | | SMF (7) | | MDF (3) | | All (35) | |
|--------|-------------------------|--------|---|---------|-------|--------|----|---------|------|---------|-----|----------|------|
| | | N | % | N | % | N | % | N | % | N | % | N | % |
| 1 | Adequate-Dry Fodder | 0 | 0 | 6 | 50 | 6 | 75 | 6 | 85.7 | 3 | 100 | 21 | 60 |
| 2 | Adequate-Green Fodder | 0 | 0 | 1 | 8.33 | 0 | 0 | 1 | 14.3 | 0 | 0 | 2 | 5.71 |
| 3 | Inadequate-Green Fodder | 0 | 0 | 2 | 16.67 | 0 | 0 | 1 | 14.3 | 0 | 0 | 3 | 8.57 |

Average annual gross income: The data regarding the annual gross income in Haligeri-3 Micro watershed is presented in Table 29. The results indicate that, the farmers have annual gross income of Rs. 81454.37 in micro-watershed, of which Rs. 74740.06 is from agriculture itself.

Table 29. Average annual gross income in Haligeri-3 micro-watershed

| Sl.No. | Particulars | LL (5) | MF (12) | SF (8) | SMF (7) | MDF (3) | All (35) |
|-------------|----------------|---------|---------|--------|---------|---------|----------|
| 1 | Service/salary | 0 | 0 | 0 | 0 | 33333.3 | 2857.14 |
| 2 | Wage | 15000.2 | 3333.33 | 2500 | 0 | 0 | 3857.17 |
| 3 | Agriculture | 0.2 | 59550.1 | 59775 | 101871 | 236667 | 74740.1 |
| Income(Rs.) | | 15000.4 | 62883.4 | 62275 | 101871 | 270000 | 81454.4 |

Average annual Expenditure: The data regarding the average annual expenditure in Haligeri-3 Micro watershed is presented in Table 30. The results indicate that, the farmers have annual gross expenditure of Rs. 242309.52 in micro-watershed, of which Rs. 31371.43 is from agriculture itself.

Table 30. Average annual Expenditure in Haligeri-3 micro-watershed

| Sl.No. | Particulars | LL (5) | MF (12) | SF (8) | SMF (7) | MDF (3) | All (35) |
|--------|-------------|--------|---------|--------|---------|---------|----------|
| 1 | Wage | 9500 | 5000 | 10000 | 0 | 0 | 1942.86 |
| 2 | Agriculture | 25000 | 26916.7 | 28750 | 27142.9 | 110000 | 31371.4 |
| Total | | 34500 | 31916.7 | 38750 | 27142.9 | 110000 | 242310 |

Horticulture species grown: The data regarding horticulture species grown in Haligeri-3 Micro watershed is presented in Table 31. The results indicate that, the total number of horticultural trees grown (both field and backyard) by the sampled households were Mango (3).

Table 31. Horticulture species grown in Haligeri-3 micro-watershed

| Sl.No. | Particulars | LL (5) | | MF (12) | | SF (8) | | SMF (7) | | MDF (3) | | All (35) | |
|--------|-------------|--------|---|---------|---|--------|---|---------|---|---------|---|----------|---|
| | | F | B | F | B | F | B | F | B | F | B | F | B |
| 1 | Mango | 0 | 0 | 0 | 0 | 1 | 0 | 1 | 0 | 1 | 0 | 3 | 0 |

*F= Field B=Back Yard

Forest species grown: The data regarding forest species grown in Haligeri-3 Micro watershed is presented in Table 32. The results indicate that, households have planted 6 neem trees, 1 tamarind trees and 9 banyan trees together in both field and backyard.

Table 32. Forest species grown in Haligeri-3 micro-watershed

| Sl.No. | Particulars | LL (5) | | MF (12) | | SF (8) | | SMF (7) | | MDF (3) | | All (35) | |
|--------|-------------|--------|---|---------|---|--------|---|---------|---|---------|---|----------|---|
| | | F | B | F | B | F | B | F | B | F | B | F | B |
| 1 | Neem | 0 | 0 | 0 | 0 | 4 | 0 | 1 | 0 | 1 | 0 | 6 | 0 |
| 2 | Tamarind | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 1 | 0 |
| 3 | Banyan | 0 | 0 | 0 | 0 | 1 | 0 | 8 | 0 | 0 | 0 | 9 | 0 |

*F= Field B=Back Yard

Average additional investment capacity: The data regarding average additional investment capacity in Haligeri-3 Micro watershed is presented in Table 33. The results indicate that, households have an average investment capacity of Rs. 485.71 for land development.

Table 33. Average additional investment capacity of households in Haligeri-3 micro-watershed

| Sl.No. | Particulars | LL (5) | MF (12) | SF (8) | SMF (7) | MDF (3) | All (35) |
|--------|------------------|--------|---------|--------|---------|---------|----------|
| 1 | Land development | 0 | 583.33 | 1250 | 0 | 0 | 485.71 |

Source of funds for additional investment: The data regarding source of funds for additional investment in Haligeri-3 Micro watershed is presented in Table 34. The results indicate that, the sources of finance raised from bank as a loan and from soft loan for land development was 5.71 and 2.86 per cent.

Table 34. Source of funds for additional investment in Haligeri-3 micro-watershed

| Sl.No | Item | Land development | |
|-------|----------------|------------------|------|
| | | N | % |
| 1 | Loan from bank | 2 | 5.71 |
| 2 | Soft loan | 1 | 2.86 |

Marketing of agricultural produce: The data regarding marketing of the agricultural produce in Haligeri-3 Micro watershed is presented in Table 35. The results indicated that, 100.00 percent of output of Bengal gram, Jowar, Sorghum and Sunflower was sold in the market and 98.92 percent of output of Maize was sold in the market.

Table 35. Marketing of agricultural produce in Haligeri-3 micro-watershed

| Sl.No | Crops | Output obtained (q) | Output retained (q) | Output sold (q) | Output sold (%) | Avg. Price obtained (Rs/q) |
|-------|------------|---------------------|---------------------|-----------------|-----------------|----------------------------|
| 1 | Bengalgram | 659 | 0 | 659 | 100 | 3160 |
| 2 | Jowar | 20 | 0 | 20 | 100 | 1200 |
| 3 | Maize | 555 | 6 | 549 | 99 | 1600 |
| 4 | Sorghum | 349 | 0 | 349 | 100 | 1590 |
| 5 | Sunflower | 5 | 0 | 5 | 100 | 5000 |

Marketing channels used for sale of agricultural produce: The data regarding marketing channels used for sale of agricultural produce in Haligeri-3 Micro watershed is

presented in Table 36. The results indicated that, 28.57 cent of the households have sold agricultural produce to the local/village merchants and 57.14 per cent of regulated market.

Table 36. Marketing channels used for sale of agricultural produce in Haligeri-3 micro-watershed

| Sl.No. | Particulars | LL (5) | | MF (12) | | SF (8) | | SMF (7) | | MDF (3) | | All (35) | |
|--------|------------------------|--------|---|---------|----|--------|------|---------|-----|---------|------|----------|-------|
| | | N | % | N | % | N | % | N | % | N | % | N | % |
| 1 | Local/village Merchant | 0 | 0 | 5 | 42 | 3 | 37.5 | 0 | 0 | 2 | 66.7 | 10 | 28.57 |
| 2 | Regulated Market | 0 | 0 | 7 | 58 | 5 | 62.5 | 7 | 100 | 1 | 33.3 | 20 | 57.14 |

Mode of transport of agricultural produce: The data regarding mode of transport of agricultural produce in Haligeri-3 Micro watershed is presented in Table 37. The results indicated that, 85.71 cent of the households have used tractor.

Table 37. Mode of transport of agricultural produce in Haligeri-3 micro-watershed

| Sl.No. | Particulars | LL (5) | | MF (12) | | SF (8) | | SMF (7) | | MDF (3) | | All (35) | |
|--------|-------------|--------|---|---------|-----|--------|-----|---------|-----|---------|-----|----------|-------|
| | | N | % | N | % | N | % | N | % | N | % | N | % |
| 1 | Tractor | 0 | 0 | 12 | 100 | 8 | 100 | 7 | 100 | 3 | 100 | 30 | 85.71 |

Incidence of soil and water erosion problems: The data regarding incidence of incidence of soil and water erosion problems in Haligeri-3 Micro watershed is presented in Table 38. The results indicate that, 85.71 per cent of the households have experienced soil and water erosion problems.

Table 38. Incidence of soil and water erosion problems in Haligeri-3 micro-watershed

| Sl.N | Particulars | LL (5) | | MF (12) | | SF (8) | | SMF (7) | | MDF (3) | | All (35) | |
|------|---|--------|---|---------|-----|--------|-----|---------|-----|---------|-----|----------|-------|
| | | N | % | N | % | N | % | N | % | N | % | N | % |
| 1 | Soil and water erosion problems in the farm | 0 | 0 | 12 | 100 | 8 | 100 | 7 | 100 | 3 | 100 | 30 | 85.71 |

Interest towards soil testing: The data regarding Interest shown towards soil testing in Haligeri-3 Micro watershed is presented in Table 39. The results indicated that, 80.00 per cent of the households were interested towards soil testing.

Table 39. Interest regarding soil testing in Haligeri-3 micro-watershed

| Sl. No. | Particulars | LL (5) | | MF (12) | | SF (8) | | SMF (7) | | MDF (3) | | All (35) | |
|---------|-----------------------|--------|---|---------|----|--------|-----|---------|-----|---------|------|----------|----|
| | | N | % | N | % | N | % | N | % | N | % | N | % |
| 1 | Interest in soil test | 0 | 0 | 11 | 92 | 8 | 100 | 7 | 100 | 2 | 66.7 | 28 | 80 |

Table 40. Usage pattern of fuel for domestic use in Haligeri-3 micro-watershed

| Sl.No. | Particulars | LL (5) | | MF (12) | | SF (8) | | SMF (7) | | MDF (3) | | All (35) | |
|--------|-------------|--------|-----|---------|------|--------|----|---------|------|---------|------|----------|-------|
| | | N | % | N | % | N | % | N | % | N | % | N | % |
| 1 | Fire Wood | 5 | 100 | 11 | 91.7 | 4 | 50 | 5 | 71.4 | 2 | 66.7 | 27 | 77.14 |
| 2 | Biogas | 0 | 0 | 0 | 0 | 2 | 25 | 1 | 14.3 | 1 | 33.3 | 4 | 11.43 |
| 3 | LPG | 0 | 0 | 1 | 8.33 | 2 | 25 | 1 | 14.3 | 0 | 0 | 4 | 11.43 |

Usage pattern of fuel for domestic use: The data on usage pattern of fuel for domestic use in Haligeri-3 Micro watershed is presented in Table 40. The results indicated that,

firewood was the major source of fuel for domestic use for 77.14 per cent of the households followed by LPG and Biogas (11.43 %).

Source of drinking water: The data on source of drinking water in Haligeri-3 Micro watershed is presented in Table 41. The results indicated that, piped supply supply of water was the major source for drinking water for 88.57 per cent of the households followed by bore well water (11.43%).

Table 41. Source of drinking water in Haligeri-3 micro-watershed

| Sl.No. | Particulars | LL (5) | | MF (12) | | SF (8) | | SMF (7) | | MDF (3) | | All (35) | |
|--------|--------------|--------|----|---------|------|--------|-----|---------|------|---------|-----|----------|-------|
| | | N | % | N | % | N | % | N | % | N | % | N | % |
| 1 | Piped supply | 3 | 60 | 11 | 91.7 | 8 | 100 | 6 | 85.7 | 3 | 100 | 31 | 88.57 |
| 2 | Bore Well | 2 | 40 | 1 | 8.33 | 0 | 0 | 1 | 14.3 | 0 | 0 | 4 | 11.43 |

Source of light: The data on source of light in Haligeri-3 Micro watershed is presented in Table 42. The results indicated that, electricity was the major source of light for 100.00 per cent of the households.

Table 42. Source of light in Haligeri-3 micro-watershed

| Sl.No. | Particulars | LL (5) | | MF (12) | | SF (8) | | SMF (7) | | MDF (3) | | All (35) | |
|--------|-------------|--------|-----|---------|-----|--------|-----|---------|-----|---------|-----|----------|-----|
| | | N | % | N | % | N | % | N | % | N | % | N | % |
| 1 | Electricity | 5 | 100 | 12 | 100 | 8 | 100 | 7 | 100 | 3 | 100 | 35 | 100 |

Existence of sanitary toilet facility: The data on availability of toilet facility in Haligeri-3 Micro watershed is presented in Table 43. The results indicated that, 34.29 per cent of the households possess toilets.

Table 43. Existence of sanitary toilet facility in Haligeri-3 micro-watershed

| Sl.No. | Particulars | LL (5) | | MF (12) | | SF (8) | | SMF (7) | | MDF (3) | | All (35) | |
|--------|--------------------------|--------|----|---------|-----|--------|-----|---------|----|---------|------|----------|------|
| | | N | % | N | % | N | % | N | % | N | % | N | % |
| 1 | Sanitary toilet facility | 1 | 20 | 1 | 8.3 | 8 | 100 | 1 | 14 | 1 | 33.3 | 12 | 34.3 |

Possession of PDS card: The data regarding possession of PDS card in Haligeri-3 Micro watershed is presented in Table 44. The results indicated that, 100.00 per cent of the households possessed BPL card.

Table 44. Possession of PDS card in Haligeri-3 micro-watershed

| Sl.No. | Particulars | LL (5) | | MF (12) | | SF (8) | | SMF (7) | | MDF (3) | | All (35) | |
|--------|-------------|--------|-----|---------|-----|--------|-----|---------|-----|---------|-----|----------|-----|
| | | N | % | N | % | N | % | N | % | N | % | N | % |
| 1 | BPL | 5 | 100 | 12 | 100 | 8 | 100 | 7 | 100 | 3 | 100 | 35 | 100 |

Participation in NREGA programme: The data regarding Participation in NREGA programme in Haligeri-3 Micro watershed is presented in Table 45. The results indicated that, only 31.43 percent of the participate have participated in NREGA programme.

Table 45. Participation in NREGA programme in Haligeri-3 micro-watershed

| Sl.No | Particulars | LL (5) | | MF (12) | | SF (8) | | SMF (7) | | MDF (3) | | All (35) | |
|-------|----------------------------------|--------|---|---------|----|--------|------|---------|------|---------|----|----------|------|
| | | N | % | N | % | N | % | N | % | N | % | N | % |
| 1 | Participation in NREGA programme | 0 | 0 | 3 | 25 | 3 | 37.5 | 3 | 42.9 | 2 | 67 | 11 | 31.4 |

Adequacy of food items: The data regarding adequacy of food items in Haligeri-3 Micro watershed is presented in Table 46. The results indicated that, the extent of adequacy of food items for cereals, pulses, Oilseeds and vegetables were 88.57, 74.29, 14.29, 2.86 per cent respectively, similarly for Fruits (11.43%), milk (5.71%), Egg (5.71%), and Meat (2.86%).

Table 46. Adequacy of food items in Haligeri-3 micro-watershed

| Sl.No. | Particulars | LL (5) | | MF (12) | | SF (8) | | SMF (7) | | MDF (3) | | All (35) | |
|--------|-------------|--------|----|---------|------|--------|------|---------|------|---------|-------|----------|-------|
| | | N | % | N | % | N | % | N | % | N | % | N | % |
| 1 | Cereals | 2 | 40 | 13 | 108 | 7 | 87.5 | 7 | 100 | 2 | 66.67 | 31 | 88.57 |
| 2 | Pulses | 2 | 40 | 12 | 100 | 5 | 62.5 | 5 | 71.4 | 2 | 66.67 | 26 | 74.29 |
| 3 | Oilseed | 1 | 20 | 3 | 25 | 0 | 0 | 1 | 14.3 | 0 | 0 | 5 | 14.29 |
| 4 | Vegetables | 0 | 0 | 1 | 8.33 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 2.86 |
| 5 | Fruits | 0 | 0 | 1 | 8.33 | 1 | 12.5 | 1 | 14.3 | 1 | 33.33 | 4 | 11.43 |
| 6 | Milk | 0 | 0 | 2 | 16.7 | 0 | 0 | 0 | 0 | 0 | 0 | 2 | 5.71 |
| 7 | Egg | 0 | 0 | 1 | 8.33 | 0 | 0 | 0 | 0 | 1 | 33.33 | 2 | 5.71 |
| 8 | Meat | 0 | 0 | 1 | 8.33 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 2.86 |

Inadequacy of food items: The data regarding in adequacy of food items in Haligeri-3 Micro watershed is presented in Table 47. The results indicated that, the extent of in adequacy of food items for cereals, pulses, Oilseeds and vegetables were 8.57, 17.14, 80.00 and 74.29 per cent respectively, similarly for fruits (65.71%), milk (54.29%), egg (80.00%) and meat (88.57%).

Table 47. Inadequacy of food items in Haligeri-3 micro-watershed

| Sl.No. | Particulars | LL (5) | | MF (12) | | SF (8) | | SMF (7) | | MDF (3) | | All (35) | |
|--------|-------------|--------|-----|---------|------|--------|------|---------|------|---------|-------|----------|-------|
| | | N | % | N | % | N | % | N | % | N | % | N | % |
| 1 | Cereals | 3 | 60 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 3 | 8.57 |
| 2 | Pulses | 2 | 40 | 0 | 0 | 2 | 25 | 2 | 28.6 | 0 | 0 | 6 | 17.14 |
| 3 | Oilseed | 4 | 80 | 10 | 83.3 | 6 | 75 | 6 | 85.7 | 2 | 66.67 | 28 | 80 |
| 4 | Vegetables | 5 | 100 | 10 | 83.3 | 6 | 75 | 4 | 57.1 | 1 | 33.33 | 26 | 74.29 |
| 5 | Fruits | 5 | 100 | 9 | 75 | 5 | 62.5 | 3 | 42.9 | 1 | 33.33 | 23 | 65.71 |
| 6 | Milk | 4 | 80 | 4 | 33.3 | 5 | 62.5 | 5 | 71.4 | 1 | 33.33 | 19 | 54.29 |
| 7 | Egg | 4 | 80 | 11 | 91.7 | 7 | 87.5 | 4 | 57.1 | 2 | 66.67 | 28 | 80 |
| 8 | Meat | 5 | 100 | 10 | 83.3 | 7 | 87.5 | 7 | 100 | 2 | 66.67 | 31 | 88.57 |

Farming constraints: The data regarding farming constraints experienced by households in Haligeri-3 Micro watershed is presented in Table 48. The results indicated that, lower fertility status of the soil was the constraint experienced by (77.14 %) per cent of the households, wild animal menace on farm field (82.86%), frequent incidence of pest and diseases (57.14%), inadequacy of irrigation water (34.29%), high cost of fertilizers and plant protection chemicals' (28.57%), high rate of interest on credit (8.57%), low price for the agricultural commodities (11.43 %), lack of marketing facilities in the area (14.29%), inadequate extension services (31.43 %), lack of transport for safe transport of the agricultural produce to the market (40.00%), less rainfall (51.43%), source of agri-technology information (Newspaper/Tv/Mobile) (40.00%).

Table 48. Farming constraints experienced in Haligeri-3 micro-watershed

| SN | Particulars | LL (5) | | MF (12) | | SF (8) | | SMF (7) | | MDF (3) | | All (35) | |
|----|--|--------|---|---------|-------|--------|-------|---------|-------|---------|-------|----------|-------|
| | | N | % | N | % | N | % | N | % | N | % | N | % |
| 1 | Lower fertility status of the soil | 0 | 0 | 11 | 91.67 | 7 | 87.5 | 7 | 100 | 2 | 66.67 | 27 | 77.14 |
| 2 | Wild animal menace on farm field | 0 | 0 | 11 | 91.67 | 9 | 112.5 | 6 | 85.71 | 3 | 100 | 29 | 82.86 |
| 3 | Frequent incidence of pest and diseases | 0 | 0 | 7 | 58.33 | 5 | 62.5 | 5 | 71.43 | 3 | 100 | 20 | 57.14 |
| 4 | Inadequacy of irrigation water | 0 | 0 | 2 | 16.67 | 4 | 50 | 3 | 42.86 | 3 | 100 | 12 | 34.29 |
| 5 | High cost of Fertilizers and plant protection chemicals | 0 | 0 | 6 | 50 | 1 | 12.5 | 2 | 28.57 | 1 | 33.33 | 10 | 28.57 |
| 6 | High rate of interest on credit | 0 | 0 | 2 | 16.67 | 0 | 0 | 1 | 14.29 | 0 | 0 | 3 | 8.57 |
| 7 | Low price for the agricultural commodities | 0 | 0 | 2 | 16.67 | 1 | 12.5 | 1 | 14.29 | 0 | 0 | 4 | 11.43 |
| 8 | Lack of marketing facilities in the area | 0 | 0 | 2 | 16.67 | 2 | 25 | 1 | 14.29 | 0 | 0 | 5 | 14.29 |
| 9 | Inadequate extension services | 0 | 0 | 3 | 25 | 4 | 50 | 3 | 42.86 | 1 | 33.33 | 11 | 31.43 |
| 10 | Lack of transport for safe transport of the Agril produce to the market. | 0 | 0 | 7 | 58.33 | 3 | 37.5 | 2 | 28.57 | 2 | 66.67 | 14 | 40 |
| 11 | Less rainfall | 0 | 0 | 7 | 58.33 | 4 | 50 | 5 | 71.43 | 2 | 66.67 | 18 | 51.43 |
| 12 | Source of Agri-technology information | 0 | 0 | 5 | 41.67 | 4 | 50 | 2 | 28.57 | 3 | 100 | 14 | 40 |

SUMMARY AND IMPLICATIONS

In order to assess the socio-economic condition of the farmers in the watershed 35 households located in the micro watershed were interviewed for the survey. The study was conducted in Haligeri-3 micro-watershed (Haligeri sub-watershed, Koppala taluk & District) is located at North latitude 15⁰ 24' 35.308" and 15⁰ 22' 52.919" and East longitude 76⁰ 4' 6.789" and 76⁰ 2' 12.128" covering an area of about 719.28 ha bounded by under Haligeri, Bhanapur, Veerapura and Tadakal Villages.

Socio-economic analysis indicated that, out of the total sample of 35 respondents, 12 (34.29%) were marginal, 8(22.86%) were small and 7 (20.00%) were semi medium, 3 (8.57%) were medium farmers. The population characteristics of households indicated that, there were 66 (54.55%) men and 55 (45.45%) were women. Majority of the respondents (36.36%) were in the age group of 35-60 years. Education level of the sample households indicated that, majority there were 36.36 per cent illiterates and only 6.61 per cent attained graduation. About, 85.71 per cent of household heads practicing agriculture and 14.29 per cent of the household heads were engaged as agricultural labourers. Agriculture was the major occupation for 24.79 per cent of the household members.

In the study area, 42.86 per cent of the households possess katcha house and 2.86 per cent possess pucca house. The durable assets owned by the households showed that, 88.57 per cent possess TV and 82.86 per cent possess mobile phones.

The average labour availability in the study area showed that, own labour men available in the micro watershed was 2.85, women available in the micro watershed was 1.03, hired labour (men) available was 6.17 and hired labour (women) available was 6.37.

Out of the total land holding of the sample respondents (57.23 ha), 90.10 per cent of the area is under dry condition. The major crops grown by sample farmers are Maize, Sorghum, Bengal gram and Sunflower and cropping intensity was recorded as 99.76 per cent.

The sample households possessed 80.00 per cent bank account and 80.00 per cent of them have savings in the account. About 68.57 per cent of the respondents borrowed credit from various sources. Among the credit borrowed by households, 37.93 per cent have borrowed loan from commercial banks and 27.59 per cent from Cooperative bank. Majority of the respondents (100.00 %) have borrowed loan for agriculture purpose. Regarding the opinion on institutional sources of credit, 73.68 per cent of the households opined that credit helped to perform timely agricultural operations.

The per hectare cost of cultivation for Maize, Sorghum, Bengal gram and Sunflower was Rs.26600.58 , 32265.36, 29342.57 and 25256.57 with benefit cost ratio of 1:2.20, 1: 1.70, 1: 1.63 and 1: 1.10 respectively.

Further, 60.00 per cent of the households opined that dry fodder was adequate and 5.71 per cent of the households have opined that the green fodder was adequate.

The average annual gross income of the farmers was Rs. 81454.37 in micro-watershed, of which Rs. 74740.06 comes from agriculture.

The total number of horticultural trees grown (both field and backyard) by the sampled households were Mango (3) and the forest species are grown 6 neem trees, 1 tamarind trees and 9 banyan trees together in both field and backyard.

Households have an average investment capacity of Rs. 485.71 for land development. Source of funds raised from bank as a loan and from soft loan for land development was 5.71 and 2.86 per cent.

Regarding marketing channels, 28.57 per cent of the households have sold agricultural produce to the local/village merchants, while, 57.14 per cent have sold by Agents/Traders. Further, 85.71 per cent of the households have used tractor for the transport of agriculture commodity.

Majority of the farmers (85.71 %) have experienced soil and water erosion problems in the watershed and 80.00 per cent of the households were interested towards soil testing.

Firewood connection was the major source of fuel for domestic use for 77.14 per cent of the households and 11.43 per cent households has LPG. Piped supply was the major source for drinking water for 88.57 per cent of the households. Electricity was the major source of light for 100.00 per cent of the households. In the study area, 34.29 per cent of the households possess toilet facility. Regarding possession of PDS card, 100.00 per cent of the households possessed BPL card. Cereals (88.57%), pulses (74.29%), oilseeds (14.29%) were adequate for consumption.

Farming constraints experienced by households in the micro watersheds were lower fertility status of the soil (77.14%) wild animal menace on farm field (82.86%), frequent incidence of pest and diseases (57.14%), inadequacy of irrigation water (34.29%), high cost of fertilizers and plant protection chemicals (28.57%), high rate of interest on credit (8.57%), low price for the agricultural commodities (11.43%), lack of marketing facilities in the area (14.29%), inadequate extension services (31.43%), lack of transport for safe transport of the agricultural produce to the market (40.00%), Less rainfall (51.43%) and Source of Agri-technology information(Newspaper/TV/Mobile) (40.00%).

Implications of the survey

- ✓ Result indicated that, there were 36.36 per cent were illiterate hence, extension methodologies such as demonstration, street play, drama, video shows will be effective in dissemination of the technologies in the micro watershed.

- ✓ The data indicate that, 42.86 per cent of the households possess katcha house. Hence, the development department while implementing the watershed plan should focus on agriculture to enhance the productivity of major crops in the area to increase the income of the farmers.
- ✓ Results indicated that the local institutional participation of the household members in the micro watershed is minimal hence, activities like membership campaign, awareness creation about the benefits of membership in local institutions and strengths of organized groups must be conveyed.
- ✓ Majority of the households in the watershed have experience in use of mobile phones, and television hence, these mass media can be effectively utilized for transfer of technology as well as for information dissemination.
- ✓ The farm machinery/implement possession in the micro watershed was found to be minimum the reasons may lack of knowledge or lack of financial ability which can be addressed through training on use of different farm implements, providing information on different sources of finance for purchase of farm implements.
- ✓ The possession of livestock such as crossbred cow found is less hence, farmers must be made aware of the benefits of crossbred cow in increased milk production.
- ✓ The possession of livestock such as sheep, goat and poultry was found to be low hence, farmers may be informed the role of subsidiary enterprises in enhancing the income and information on financial support for subsidiary activities.
- ✓ Households possess 51.56ha (90.10 %) of dry land hence, the availability of the dry land agricultural technologies such as short duration crops, high yielding drought resistance crop varieties, drip irrigation technology and subsidy information will be helpful for the farmers to enhance the productivity of land and as well as farmers income.
- ✓ Few of the bore well in micro watershed found non functional hence, farmers may be trained on possibility of bore well rejuvenation.
- ✓ The total number of horticultural trees grown (both field and backyard) by the sampled households were Mango (3) and the forest species are grown 6 neem trees, 1 tamarind trees and 9 banyan trees together in both field and backyard. Hence, production technologies related to these crops can be made available to the farmers for better adoption.
- ✓ The cropping intensity in the micro watershed was found to be (99.76 %) hence, care must be taken by the implementing agency to bring uncultivated land into cultivation through suitable measures.
- ✓ Many of the household members have borrowed loan from cooperative banks which has higher rate of interest hence, farmers may be sensitized on the different sources of credit with lesser interest rate such SHGs etc.
- ✓ The results indicated the non availability of both green and dry fodder throughout the year hence, fodder development activities can be taken up in the micro watershed.

- ✓ The average annual gross income of the households Rs.74740.06 from agriculture, Rs.0.00 from business and Rs. 3857.17 from wages and. Agriculture was found to be the major source of income for households hence; the development activities should focus on productivity enhancement, marketing arrangements and agricultural technology dissemination to have a direct impact on the farmers.
- ✓ The cultivation of forest species is found minimal hence, information and production technology related to agro-forestry and integrated farming system.
- ✓ The data indicated that, 85.71 per cent of the households have experienced soil and water erosion problems. Hence, those farmers who reported the soil and water erosion problems may be given attention while implementation of the watershed development plan.
- ✓ The data indicated that, 80.00 per cent of the households have interest in soil testing hence, farmers must be provided with the information on various institutions which are involved in soil testing for the benefit of the farmers.
- ✓ Except summer ploughing the adoption of other soil and water conservation structures is minimum hence, the farmers in the micro watershed should be sensitized on the use of different conservation structures for soil water conservation.
- ✓ Cereals and pulses found be adequate for per cent of the households respectively hence, farm households and the farm women must be trained on importance of balanced nutrition and role of vegetable, milk, egg, meat in balanced diet.
- ✓ Lower fertility status of the soil (77.14%), wild animal menace on farm field (82.86%), frequent incidence of pest and diseases (57.14%), high cost of fertilizers and plant protection chemicals (28.57%), high rate of interest on credit (8.57%), low price for the agricultural commodities (11.43%), lack of marketing facilities in the area (14.29%), inadequate extension services (31.43%), lack of transport for safe transport of the agricultural produce to the market (40.00%) were the major farming constraints experienced hence, these constraints must be addressed immediately for the welfare of the farmers. Awareness to be created among the farmers to approach nearest KVKs/RSKs and other developmental departments for technical and for subsidized inputs and utilize the well established regulated markets, approaching the contract firms, direct markets to avoid the involvement of middlemen.