



हर कदम, हर डगर
किसानों का हमसफर
भारतीय कृषि अनुसंधान परिषद

AgriSearch with a human touch

**LAND RESOURCE INVENTORY AND SOCIO-ECONOMIC STATUS OF
FARM HOUSEHOLDS FOR WATERSHED PLANNING AND
DEVELOPMENT**

YADGIR-RF (4D5B1G1a) MICROWATERSHED

Yadgir Taluk and District, Karnataka

Karnataka Watershed Development Project – II

SUJALA – III

World Bank funded Project



THE WORLD BANK



ICAR – NATIONAL BUREAU OF SOIL SURVEY AND LAND USE PLANNING



ICAR - NBSS & LUP

**WATERSHED DEVELOPMENT DEPARTMENT
GOVT. OF KARNATAKA, BANGALORE**



About ICAR - NBSS&LUP

The ICAR-National Bureau of Soil Survey and Land Use Planning (ICAR-NBSS&LUP), Nagpur, a premier Institute of the Indian Council of Agricultural Research (ICAR), was set up during 1976 with the objective to prepare soil resource maps at national, state and district levels and to provide research inputs in soil resource mapping and its applications, land evaluation, land use planning, land resource management, and database management using GIS for optimising land use on different kinds of soils in the country.

The Bureau has been engaged in carrying out soil resource survey, agro-ecological and soil degradation mapping at the country, state and district levels for qualitative assessment and monitoring the soil health towards viable land use planning. The research activities have resulted in identifying the soil potentials and problems, and the various applications of the soil surveys with the ultimate objective of sustainable agricultural development. The Bureau has the mandate to correlate and classify soils of the country and maintain a National Register of all the established soil series. The Institute is also imparting in-service training to staff of the soil survey agencies in the area of soil survey, land evaluation and soil survey interpretations for land use planning. The Bureau in collaboration with Panjabrao Krishi Vidyapeeth, Akola is running post-graduate teaching and research programme in land resource management, leading to M.Sc. and Ph.D. degrees.

Citation: Rajendra Hegde, Ramesh Kumar, S.C., B.A. Dhanorkar, S. Srinivas, M. Lalitha, K.V. Niranjana, R.S. Reddy and S.K. Singh (2019). "Land resource inventory and socio-economic status of farm households for watershed planning and development of Yadgir Rf (4D5B1G1a) Microwatershed, Yadgir Taluk and District, Karnataka", ICAR-NBSS&LUP Sujala MWS Publ .489, ICAR – NBSS & LUP, RC, Bangalore. p.116 & 34..

TO OBTAIN COPIES,

Please write to:

Director, ICAR - NBSS & LUP,

Amaravati Road, NAGPUR - 440 033, India

Phone : (0712) 2500386, 2500664, 2500545 (O)

Telefax : 0712-2522534

E-Mail : director@nbsslup.ernet.in

Website URL : nbsslup.in

Or

Head, Regional Centre, ICAR - NBSS&LUP, Hebbal, Bangalore - 560 024

Phone : (080) 23412242, 23510350 (O)

Telefax : 080-23510350

E-Mail : nbssrcb@gmail.com

PREFACE

In Karnataka, as in other Indian States, the livelihoods of rural people are intertwined with farming pursuits. The challenges in agriculture are seriously threatening the livelihood of a large number of farmers as they have been practicing farming in contextual factors beyond their control. Climatic factors are the most important ones and have become much more significant in recent times due to rapid climate changes induced by intensive anthropogenic activities affecting our ecosystem in multiple ways. Climate change has become the reality, it is happening and efforts to evolve and demonstrate climate resilient technologies have become essential. Due to the already over stressed scenario of agrarian sector, the climate change is resulting in manifold increase in the complexities, pushing the rural mass to face more and more unpredictable situations. The rising temperatures and unpredictable rainfall patterns are going to test seriously the informed decisions farmers have to make in order to survive in farming and sustain their livelihood.

It is generally recognized that impacts of climate change shall not be uniform across the globe. It is said that impact of climate change is more severe in South Asia. Based on the analysis of meteorological data, it is predicted that in India, there will be upward trend in mean temperature, downward trend in relative humidity, annual rainfall and number of wet days in a year. Also, in general, phenomena like erratic monsoon, spread of tropical diseases, rise in sea levels, changes in availability of fresh water, frequent floods, droughts, heat waves, storms and hurricanes are predicted. Each one of these adverse situations are already being experienced in various parts of India and also at the global level. Decline in agricultural productivity of small and marginal farmers becoming more vulnerable is already witnessed.

In Karnataka, more than 60 per cent of the population live in rural areas and depend on agriculture and allied activities for their livelihood. Though the state has achieved significant progress in increasing the yield of many crops, there is tremendous pressure on the land resources due to the growing and competing demands of various land uses. This is reflected in the alarming rate of land degradation observed. Already more than 50 per cent of the area is affected by various forms of degradation. If this trend continues, the sustainability of the fragile ecosystem will be badly affected. The adverse effects of change in the climatic factors are putting additional stress on the land resources and the farmers dependent on this.

The natural resources (land, water and vegetation) of the state need adequate and constant care and management, backed by site-specific technological interventions and investments particularly by the government. Detailed database pertaining to the nature of the land resources, their constraints, inherent potentials and suitability for various land

based rural enterprises, crops and other uses is a prerequisite for preparing location-specific action plans, which are in tune with the inherent capability of the resources. Any effort to evolve climate resilient technologies has to be based on the baseline scientific database. Then only one can expect effective implementation of climate resilient technologies, monitor the progress, make essential review of the strategy, and finally evaluate the effectiveness of the implemented programs. The information available at present on the land resources of the state are of general nature and useful only for general purpose planning. Since the need of the hour is to have site-specific information suitable for farm level planning and detailed characterization and delineation of the existing land resources of an area into similar management units is the only option.

ICAR-NBSS&LUP, Regional Centre, Bangalore has taken up a project sponsored by the Karnataka Watershed Development Project-II, (Sujala-III), Government of Karnataka funded by the World Bank under Component-1 Land Resource Inventory. This study was taken up to demonstrate the utility of such a database in reviewing, monitoring and evaluating all the land based watershed development programs on a scientific footing. To meet the requirements of various land use planners at grassroots level, the present study on “Land Resource Inventory and Socio-Economic Status of Farm Households for Watershed Planning and Development of Yadgir Rf Microwatershed, Yadgir Taluk and District, Karnataka” for integrated development was taken up in collaboration with the State Agricultural Universities, IISC, KRSRAC, KSNDMC as Consortia partners. The project provides detailed land resource information at cadastral level (1:7920 scale) for all the plots and socio-economic status of farm households covering thirty per cent farmers randomly selected representing landed and landless class of farmers in the microwatershed. The project report with the accompanying maps for the Microwatershed will provide required detailed database for evolving effective land use plan, alternative land use options and conservation plans for the planners, administrators, agricultural extension personnel, KVK officials, developmental departments and other land users to manage the land resources in a sustainable manner.

It is hoped that this database will be useful to the planners, administrators and developmental agencies working in the area in not only for formulating location specific developmental schemes but also for their effective monitoring at the village/watershed level.

Nagpur

Date: 10.05.2017

S.K. SINGH

Director, ICAR - NBSS&LUP, Nagpur

Contributors

Dr. Rajendra Hegde Principal Scientist, Head & Project Leader, Sujala-III Project ICAR-NBSS&LUP, Regional Centre, Bangalore	Dr. S.K.Singh Director, ICAR-NBSS&LUP Coordinator, Sujala-III Project Nagpur
Soil Survey, Mapping & Report Preparation	
Dr. B.A. Dhanorkar	Sh. R.S. Reddy
Dr. K.V. Niranjana	Sh. Venkata Giriappa
	Mr. Somashekar T N
	Smt. Chaitra, S.P.
	Dr. Gopali bardhan
	Ms. Arpitha
	Dr. Mahendra Kumar, M.B.
Field Work	
Sh. C.BacheGowda	Sh. Mahesh, D.B.
Sh. Somashekar	Sh. Ashok S Sindagi
Sh. M. Jayaramaiah	Sh. Veerabhadrapa B.
Sh. Paramesha, K.	Sh. Kailas
Sh. B. M. Narayana Reddy	Sh. Anand
	Sh. Arun N Kambar.
	Sh Kamalesh Awate
	Sh. Sharaan Kumar Huppar
	Sh. Yogesh H.N.
	Sh. Kalaveerachari R Kammar
GIS Work	
Dr. S.Srinivas	Sh. A.G.Devendra Prasad
Sh. D.H.Venkatesh	Sh. Prakashanaik, M.K.
Smt.K.Sujatha	Sh. Abhijith Sastry, N.S.
Smt. K.V.Archana	Sh. Sudip Kumar Suklabaidya
Sh. N. Maddileti	Sh. Avinash, K.N.
	Sh. Amar Suputhra, S
	Sh. Deepak, M.J.
	Smt. K.Karunya Lakshmi
	Ms. Seema, K.V.
	Ms. A. Rajab Nisha

Laboratory Analysis	
Dr. K.M.Nair	Ms. Steffi Peter
Smt. Arti Koyal	Ms. Thara, V.R
Smt. Parvathy	Ms. Roopa, G.
	Ms. Swati, H.
	Sh. Shantaveera Swami
	Ms. Shwetha, N.K.
	Smt. Ishrat Haji
	Ms. P. Pavan Kumari
	Ms. Padmaja
	Ms. Veena, M.
Socio-Economic Analysis	
Dr. S.C. Ramesh Kumar	Sh. M.K. Prakashanaik
	Ms. Karuna V. Kulkarni
	Mrs. Sowmya A.N
	Sh. Vinod R
	Sh. Basavaraja
	Sh. Vijay Kumar Lamani
	Ms. Sowmya K.B
	Mrs. Prathibha, D.G
	Sh. Rajendra,D
Soil & Water Conservation	
Sh. Sunil P. Maske	
Watershed Development Department, GoK, Bangalore	
Sh. Rajeev Ranjan IFS Project Director & Commissioner, WDD	Dr. A. Natarajan NRM Consultant, Sujala-III Project
Dr. S.D. Pathak IFS Executive Director & Chief Conservator of Forests, WDD	

PART-A

LAND RESOURCE INVENTORY

Contents

Preface		
Contributors		
Executive Summary		
Chapter 1	Introduction	1
Chapter 2	Geographical Setting	3
2.1	Location and Extent	3
2.2	Geology	3
2.3	Physiography	4
2.4	Drainage	4
2.5	Climate	4
2.6	Natural Vegetation	6
2.7	Land Utilization	6
Chapter 3	Survey Methodology	9
3.1	Base maps	9
3.2	Image Interpretation for Physiography	9
3.3	Field Investigation	12
3.4	Soil Mapping	13
3.5	Laboratory Characterization	14
Chapter 4	The Soils	17
4.1	Soils of granite gneiss landscape	17
Chapter 5	Interpretation for Land Resource Management	21
5.1	Land Capability Classification	21
5.2	Soil Depth	23
5.3	Surface Soil Texture	24
5.4	Soil Gravelliness	25
5.5	Available Water Capacity	26
5.6	Soil Slope	27
5.7	Soil Erosion	28
Chapter 6	Fertility Status	31
6.1	Soil Reaction (pH)	31
6.2	Electrical Conductivity (EC)	31
6.3	Organic Carbon (OC)	31
6.4	Available Phosphorus	33
6.5	Available Potassium	33
6.6	Available Sulphur	33
6.7	Available Boron	33
6.8	Available Iron	33
6.9	Available Manganese	34
6.10	Available Copper	34
6.11	Available Zinc	38

Chapter 7	Land Suitability for Major Crops	39
7.1	Land suitability for Sorghum	39
7.2	Land suitability for Maize	40
7.3	Land suitability for Bajra	41
7.4	Land suitability for Groundnut	42
7.5	Land suitability for Sunflower	43
7.6	Land suitability for Redgram	44
7.7	Land suitability for Bengal gram	45
7.8	Land suitability for Cotton	46
7.9	Land suitability for Chilli	47
7.10	Land suitability for Tomato	48
7.11	Land suitability for Brinjal	49
7.12	Land suitability for Onion	50
7.13	Land suitability for Bhendi	51
7.14	Land suitability for Drumstick	52
7.15	Land suitability for Mango	53
7.16	Land suitability for Guava	54
7.17	Land suitability for Sapota	55
7.18	Land Suitability for Pomegranate	56
7.19	Land Suitability for Musambi	57
7.20	Land Suitability for Lime	58
7.21	Land Suitability for Amla	59
7.22	Land Suitability for Cashew	60
7.23	Land Suitability for Jackfruit	61
7.24	Land Suitability for Jamun	62
7.25	Land Suitability for Custard apple	63
7.26	Land Suitability for Tamarind	64
7.27	Land Suitability for Mulberry	65
7.28	Land Suitability for Marigold	66
7.29	Land Suitability for Chrysanthemum	67
7.30	Land Management Units	99
7.31	Proposed Crop Plan	99
Chapter 8	Soil Health Management	102
Chapter 9	Soil and Water conservation Treatment Plan	108
9.1	Treatment Plan	109
9.2	Recommended Soil and Water Conservation measures	112
9.3	Greening of Microwatershed	113
	References	116
	Appendix I	
	Appendix II	
	Appendix III	

LIST OF TABLES

2.1	Mean Monthly Rainfall, PET, 1/2 PET at Yadgir Taluk & District	5
2.2	Land Utilization in Yadgir district	7
3.1	Differentiating Characteristics used for Identifying Soil Series	13
3.2	Soil map unit description of Yadgir Rf Microwatershed	14
4.1	Physical and Chemical Characteristics of Soil Series identified in Yadgir Rf microwatershed	19
7.1	Soil-Site Characteristics of Yadgir Rf Microwatershed	69
7.2	Land suitability criteria for Sorghum	70
7.3	Land suitability criteria for Maize	71
7.4	Land suitability criteria for Bajra	72
7.5	Land suitability criteria for Groundnut	73
7.6	Land suitability criteria for Sunflower	74
7.7	Land suitability criteria for Redgram	75
7.8	Land suitability criteria for Bengal gram	76
7.9	Land suitability criteria for Cotton	77
7.10	Land suitability criteria for Chilli	78
7.11	Land suitability criteria for Tomato	79
7.12	Land suitability criteria for Brinjal	80
7.13	Land suitability criteria for Onion	81
7.14	Land suitability criteria for Bhenidi	82
7.15	Land suitability criteria for Drumstick	83
7.16	Land suitability criteria for Mango	84
7.17	Land suitability criteria for Guava	85
7.18	Land suitability criteria for Sapota	86
7.19	Land suitability criteria for Pomegranate	87
7.20	Land suitability criteria for Musambi	88
7.21	Land suitability criteria for Lime	89
7.22	Land suitability criteria for Amla	90
7.23	Land suitability criteria for Cashew	91
7.24	Land suitability criteria for Jackfruit	92
7.25	Land suitability criteria for Jamun	93
7.26	Land suitability criteria for Custard apple	94
7.27	Land suitability criteria for Tamarind	95

7.28	Land suitability criteria for Mulberry	96
7.29	Land suitability criteria for Marigold	97
7.30	Land suitability criteria for Chrysanthemum	98
7.31	Proposed Crop Plan for Yadgir Rf Microwatershed	100

LIST OF FIGURES

2.1	Location map of Yadgir Rf Microwatershed	3
2.2	Granite and granite gneiss rock formation	4
2.3	Rainfall distribution in Yadgir Taluk & District	5
2.4	Natural vegetation of Yadgir Rf Microwatershed	6
2.5	Current Land use map of Yadgir Rf Microwatershed	7
2.6	Different crops and cropping systems in Yadgir Rf Microwatershed	8
3.1	Scanned and Digitized Cadastral map of Yadgir Rf Microwatershed	10
3.2	Satellite image of Yadgir Rf Microwatershed	11
3.3	Cadastral map overlaid on IRS PAN+LISS IV merged imagery of Yadgir Rf Microwatershed	11
3.4	Location of profiles in a transect	12
3.5	Soil phase or management units of Yadgir Rf Microwatershed	15
5.1	Land Capability Classification map of Yadgir Rf Microwatershed	23
5.2	Soil Depth map of Yadgir Rf Microwatershed	24
5.3	Surface Soil Texture map of Yadgir Rf Microwatershed	25
5.4	Soil Gravelliness map of Yadgir Rf Microwatershed	26
5.5	Soil Available Water Capacity map of Yadgir Rf Microwatershed	27
5.6	Soil Slope map of Yadgir Rf Microwatershed	28
5.7	Soil Erosion map of Yadgir Rf Microwatershed	29
6.1	Soil Reaction (pH) map of Yadgir Rf Microwatershed	32
6.2	Electrical Conductivity (EC) map of Yadgir Rf Microwatershed	32
6.3	Soil Organic Carbon (OC) map of Yadgir Rf Microwatershed	33
6.4	Soil Available Phosphorus map of Yadgir Rf Microwatershed	34
6.5	Soil Available Potassium map of Yadgir Rf Microwatershed	35
6.6	Soil Available Sulphur map of Yadgir Rf Microwatershed	35
6.7	Soil Available Boron map of Yadgir Rf Microwatershed	36
6.8	Soil Available Iron map of Yadgir Rf Microwatershed	36
6.9	Soil Available Manganese map of Yadgir Rf Microwatershed	37
6.10	Soil Available Copper map of Yadgir Rf Microwatershed	37
6.11	Soil Available Zinc map of Yadgir Rf Microwatershed	38
7.1	Land suitability for Sorghum	40
7.2	Land suitability for Maize	41

7.3	Land suitability for Bajra	42
7.4	Land suitability for Groundnut	43
7.5	Land suitability for Sunflower	44
7.6	Land suitability for Redgram	45
7.7	Land suitability for Bengal gram	46
7.8	Land suitability for Cotton	47
7.9	Land suitability for Chilli	48
7.10	Land suitability for Tomato	49
7.11	Land suitability for Brinjal	50
7.12	Land suitability for Onion	51
7.13	Land suitability for Bhendi	52
7.14	Land suitable for Drumstick	53
7.15	Land suitability for Mango	54
7.16	Land suitability for Guava	55
7.17	Land suitability for Sapota	56
7.18	Land suitability for Pomegranate	57
7.19	Land suitability for Musambi	58
7.20	Land suitability for Lime	59
7.21	Land suitability for Amla	60
7.22	Land suitability for Cashew	61
7.23	Land suitability for Jackfruit	62
7.24	Land suitability for Jamun	63
7.25	Land suitability for Custard apple	64
7.26	Land suitability for Tamarind	65
7.27	Land suitability for Mulberry	66
7.28	Land suitability for Marigold	67
7.29	Land suitability for Chrysanthemum	68
7.30	Land use classes map of Yadgir Rf Microwatershed	99
9.1	Soil and water conservation map of Yadgir Rf Microwatershed	113

EXECUTIVE SUMMARY

The land resource inventory of Yadgir Rf Microwatershed was conducted using village cadastral maps and IRS satellite imagery on 1:7920 scale. The false colour composites of IRS imagery were interpreted for physiography and the physiographic delineations were used as base for mapping soils. The soils were studied in several transects and a soil map was prepared with phases of soil series as mapping units. Random checks were made all over the area outside the transects to confirm and validate the soil map unit boundaries. The soil map shows the geographic distribution and extent, characteristics, classification, behavior and use potentials of the soils in the microwatershed.

The present study covers an area of 534 ha in Yadgir taluk & district, Karnataka. The climate is semiarid and categorized as drought-prone with an average annual rainfall of 866 mm, of which about 652 mm is received during south-west monsoon, 138 mm during north-east and the remaining 76 mm during the rest of the year. An area of 336 ha in the microwatershed is covered by soils and 198 ha covered by rock outcrops. The salient findings from the land resource inventory are summarized briefly below.

- ❖ The soils belong to 2 soil series and 2 soil phases (management units) and 2 land management units.*
- ❖ The length of crop growing period is about 120-150 days starting from 1st week of June to 4th week of October.*
- ❖ From the master soil map, several interpretative and thematic maps like land capability, soil depth, surface soil texture, soil gravelliness, available water capacity, soil slope and soil erosion were generated.*
- ❖ Soil fertility status maps for macro and micronutrients were generated based on the surface soil samples collected at every 320 m grid interval.*
- ❖ Land suitability for growing 28 major agricultural and horticultural crops was assessed and maps showing the degree of suitability along with constraints were generated.*
- ❖ Only 336 ha area in the microwatershed is suitable for agriculture and 198 ha area is under rocklands, both massive and boundary with little or no soil.*
- ❖ About 61 per cent area are shallow (25-50 cm) and 2 per cent area of the microwatershed has soils that are moderately shallow (50-75cm).*
- ❖ About 61 per cent area in the microwatershed has sandy soils and 9 per cent area loamy soils at the surface.*
- ❖ Maximum of 61 per cent area in the microwatershed is gravelly (15-35%) and 2 per cent is non gravelly (<15%).*
- ❖ About 2 per cent area low (51-100 mm/m) and 61 per cent area very low (<50 mm/m) in available water capacity in the microwatershed.*
- ❖ Entire area in the microwatershed has very gently sloping (1-3% slope) lands.*

- ❖ *Entire area in the microwatershed has moderately (e2) eroded lands.*
- ❖ *An area of about <1 per cent area is slightly acid (pH 6.0-6.5) in soil reaction, 62 per cent area is neutral (pH 6.5-7.3) and one per cent soils are slightly alkaline (pH 7.3-7.8).*
- ❖ *The Electrical Conductivity (EC) of entire soils of the microwatershed is dominantly <math><2 \text{ dsm}^{-1}</math> indicating that the soils are non-saline.*
- ❖ *About 26 per cent of the soils are medium (0.5-0.75%) in organic carbon and high (>0.75%) in 37 per cent area.*
- ❖ *About 16 per cent area is low (<23 kg/ha) in available phosphorus and 47 per area is medium (23-57 kg/ha).*
- ❖ *Entire area in the microwatershed is medium (145-337 kg/ha) in available potassium.*
- ❖ *Entire area in the microwatershed is low (<10 ppm) in available sulphur.*
- ❖ *Entire area in the microwatershed is low (<0.5 ppm) in available boron.*
- ❖ *Available iron is sufficient (>4.5 ppm) in an area of about 47 per cent and deficient (<4.5 ppm) in 16 per cent area of the microwatershed.*
- ❖ *Available manganese and copper are sufficient in all the soils of the microwatershed.*
- ❖ *Available zinc is deficient (<0.6 ppm) in an area of 62 per cent and sufficient (>0.6 ppm) in one per cent area of the microwatershed.*
- ❖ *The land suitability for 28 major crops grown in the microwatershed were assessed and the areas that are highly suitable (S1) and moderately suitable (S2) are given below. It is however to be noted that a given soil may be suitable for various crops but what specific crop to be grown may be decided by the farmer looking to his capacity to invest on various inputs, marketing infrastructure, market price and finally the demand and supply position.*

Land suitability for various crops in the Microwatershed

Crop	Suitability Area in ha (%)		Crop	Suitability Area in ha (%)	
	Highly suitable (S1)	Moderately suitable (S2)		Highly suitable (S1)	Moderately suitable (S2)
<i>Sorghum</i>	-	9(2)	<i>Guava</i>	-	-
<i>Maize</i>	-	9(2)	<i>Sapota</i>	-	-
<i>Bajra</i>	-	9(2)	<i>Pomegranate</i>	-	-
<i>Groundnut</i>	-	9(2)	<i>Musambi</i>	-	-
<i>Sunflower</i>	-	-	<i>Lime</i>	-	-
<i>Redgram</i>	-	-	<i>Amla</i>	-	9(2)
<i>Bengal gram</i>	-	9(2)	<i>Cashew</i>	-	-
<i>Cotton</i>	-	9(2)	<i>Jackfruit</i>	-	-
<i>Chilli</i>	-	9(2)	<i>Jamun</i>	-	-
<i>Tomato</i>	-	9(2)	<i>Custard apple</i>	-	9(2)
<i>Brinjal</i>	-	9(2)	<i>Tamarind</i>	-	-
<i>Bhendi</i>	-	9(2)	<i>Mulberry</i>	-	-
<i>Drumstick</i>	-	-	<i>Marigold</i>	-	9(2)
<i>Mango</i>	-	-	<i>Chrysanthemum</i>	-	9(2)

- ❖ *Apart from the individual crop suitability, a proposed crop plan has been prepared for the identified LMUs by considering only the highly and moderately suitable lands for different crops and cropping systems with food, fodder, fibre and other horticulture crops.*
- ❖ *Maintaining soil-health is vital for crop production and conserve soil and land resource base for maintaining ecological balance and to mitigate climate change. For this, several ameliorative measures have been suggested for these problematic soils like saline/alkali, highly eroded, sandy soils etc.,*
- ❖ *Soil and water conservation treatment plan has been prepared that would help in identifying the sites to be treated and also the type of structures required.*
- ❖ *As part of the greening programme, several tree species have been suggested to be planted in marginal and submarginal lands, field bunds and also in the hillocks, mounds and ridges. This would help in not only supplementing the farm income but also provide fodder and fuel and generate lot of biomass which would help in maintaining an ecological balance and also contribute to mitigating the climate change.*

INTRODUCTION

Land is a scarce resource and basic unit for any material production. It can support the needs of the growing population, provided they use the land in a rational and judicious manner. But what is happening in many areas of the state is a cause for concern to everyone involved in the management of land resources at the grassroots level. The area available for agriculture is about 51 per cent of the total geographical area and more than 60 per cent of the people are still dependant on agriculture for their livelihood. The limited land area is under severe stress and strain due to increasing population pressure and competing demands of various land uses. Due to this, every year there is significant diversion of farm lands and water resources for non-agricultural purposes. Apart from this, due to lack of interest in farmers for farming, large tracts of cultivable lands are turning into fallows in many areas and this trend is continuing at an alarming rate.

Further, land degradation has emerged as a serious problem which has already affected about 38 lakh ha of cultivated area in the state. Soil erosion alone has degraded about 35 lakh ha. Almost all the uncultivated areas are facing various degrees of degradation, particularly soil erosion. Salinity and alkalinity has emerged as a major problem in more than 3.5 lakh ha in the irrigated areas of the state. Nutrient depletion and declining factor productivity is common in both rainfed and irrigated areas. The degradation is continuing at an alarming rate and there appears to be no systematic effort among the stakeholders to contain this process. In recent times, an aberration of weather due to climate change phenomenon has added another dimension leading to unpredictable situations to be tackled by the farmers.

In this critical juncture, the challenge before us is not only to increase the productivity per unit area which is steadily declining and showing a fatigue syndrome, but also to prevent or at least reduce the severity of degradation. If the situation is not reversed at the earliest, then the sustainability of the already fragile crop production system and the overall ecosystem will be badly affected in the state. The continued neglect and unscientific use of the resources for a long time has led to the situation observed at present in the state. It is a known fact and established beyond doubt by many studies in the past that the cause for all kinds of degradation is the neglect and irrational use of the land resources. Hence, there is an urgent need to generate a detailed site-specific farm level database on various land resources for all the villages/watersheds in a time bound manner that would help to protect the valuable soil and land resources and also to stabilize the farm production.

Therefore, the land resource inventory required for farm level planning is the one which investigates not only the surface but also consider the other parameters which are critical for productivity *viz.*, soils, climate, water, minerals and rocks, topography, geology, hydrology, vegetation, crops, land use pattern, animal population, socio-

economic conditions, infrastructure, marketing facilities and various schemes and developmental works of the government etc. From the data collected at farm level, the specific problems and potentials of the area can be identified and highlighted, conservation measures required for the area can be planned on a scientific footing, suitability of the area for various uses can be worked out and finally viable and sustainable land use options suitable for each and every land holding can be prescribed.

The Land Resource Inventory is basically done for identifying the potential and problem areas, developing sustainable land use plans, estimation of surface run off and water harvesting potential, preparation of soil and water conservation plans, land degradation/desertification etc. The Bureau is presently engaged in developing an LRI methodology using high resolution satellite remote sensing data and Digital Elevation Model (DEM) data to prepare Landscape Ecological Units (LEU) map representing agro-ecosystem as a whole. The LEU is preferred over landform as the base map for LRI. LEU is the assemblage of landform, slope and land use. An attempt has already been made to upscale the soil resource information from 1:250000 and 1:50000 scale to the LEU map in Goa and other states.

The land resource inventory aims to provide site-specific database for Yadgir Rf microwatershed in Yadgir Taluk & District, Karnataka State for the Karnataka Watershed Development Department. The database was generated by using cadastral map of the village as a base along with high resolution IRS LISS IV and Cartosat-1 merged satellite imagery. Later, an attempt will be made to uplink this LRI data generated at 1:7920 scale under Sujala-III Project to the proposed Landscape Ecological Units (LEUs) map.

The study was organized and executed by the ICAR- National Bureau of Soil Survey and Land Use Planning, Regional Centre, Bangalore under Generation of Land Resource Inventory Data Base Component-1 of the Sujala-III Project funded by the World Bank.

GEOGRAPHICAL SETTING

2.1 Location and Extent

The Yadgir Rf microwatershed is located in the northern part of Karnataka in Yadgir Taluk & District, Karnataka State (Fig.2.1). It comprises parts of Ashinala, Arakera. K and Ramasamudra villages. It lies between $16^{\circ}47'$ – $16^{\circ}49'$ North latitudes and $77^{\circ}15'$ – $77^{\circ}16'$ East longitudes, covering an area of about 534.1 ha. It is about 12 km southeast of Yadgir town and is surrounded by Ashinala on the north and northwest, Arakera. K on the east and Ramasamudra village on the southern side.




Fig.2.1 Location map of Yadgir Rf microwatershed

2.2 Geology

Major rock formation observed in the microwatershed is granite gneiss. (Figs.2.2). Granite gneisses are essentially pink to gray and are coarse to medium grained. They consist primarily of quartz, feldspar, biotite and hornblende. The gray granite gneisses are highly weathered, fractured and fissured upto a depth of about 10 m. Dolerite dykes and

quartz veins are common with variable width and found to occur in Yadgir Rf microwatershed.



Fig.2.2 Granite and granite gneiss rocks formation

2.3 Physiography

Physiographically, the area has been identified as granite gneiss landscape based on geology. The area has been further subdivided into five landforms, *viz;* mounds/ridges, summits, side slopes and very gently sloping uplands, plains and valleys based on slope and its relief features. The elevation ranges from 427-474 m above MSL. The mounds and ridges are mostly covered by rock outcrops.

2.4 Drainage

The area is drained by several parallel streams like Bori, Amerja and Kanga which finally join the river Bhima along its course. Though, they are not perennial, during rainy season they carry large quantities of rain water. The microwatershed has only few small tanks which are not capable of storing the water that flows during the rainy season. Due to this, the ground water recharge is very much affected. This is reflected in the failure of many bore wells in the villages. If the available rain water is properly harnessed by constructing new tanks and recharge structures at appropriate places in the villages, then the drinking and irrigation needs of the area can be easily met. The drainage network is parallel to sub parallel and dendritic.

2.5 Climate

The Yadgir district lies in the northern plains of Karnataka and falls under semiarid tract of the state and is categorized as drought- prone with total annual rainfall of 866 mm (Table 2.1). Of the total rainfall, maximum of 652 mm is received during the south–west monsoon period from June to September, the north-east monsoon from October to early December contributes about 138 mm and the remaining 76 mm during

the rest of the year. The summer season starts during the middle of February and continues up to the first week of June. The period from December to the middle of February is the coldest season. December is the coldest month with mean daily maximum and minimum temperatures being 29.5⁰C and 10⁰C respectively. During peak summer, temperature shoots up to 45⁰C. Relative humidity varies from 26% in summer to 62% in winter. Rainfall distribution is shown in Figure 2.3. The average Potential Evapo-Transpiration (PET) is 141 mm and varies from a low of 81 mm in December to 199 mm in the month of May. The PET is always higher than precipitation in all the months except end of June to end of September. Generally, the Length of crop Growing Period (LGP) is 120-150 days and starts from 1st week of June to 4th week of October.

Table 2.1 Mean Monthly Rainfall, PET, 1/2 PET at Yadgir Taluk, Yadgir District

Sl. No.	Months	Rainfall	PET	1/2 PET
1	January	4.30	86.0	43.0
2	February	2.30	125.5	62.7
3	March	15.10	166.0	83.0
4	April	18.50	179.8	89.9
5	May	36.0	198.8	97.9
6	June	118.0	175.1	87.5
7	July	171.80	156.3	78.1
8	August	182.9	150.3	75.1
9	September	179.7	142.0	71.0
10	October	105.3	138.5	69.2
11	November	26.4	97.60	48.6
12	December	6.0	80.90	40.4
Total		866.3		




Fig 2.3 Rainfall distribution in Yadgir Taluk and District

2.6 Natural Vegetation

The natural vegetation is sparse comprising few tree species, shrubs and herbs. The mounds, ridges and boulders occupy very sizeable area which is under thin to moderately thick forest vegetation. Still, there are some remnants of the past forest cover which can be seen in patches in some ridges and hillocks in the microwatershed (Fig 2.4).

Apart from the continuing deforestation, the presence of large population of goats, sheep and other cattle in the microwatershed is causing vegetative degradation of whatever little vegetation left in the area. The uncontrolled grazing has left no time for the regeneration of the vegetative cover. This leads to the accelerated rate of erosion on the hill slopes resulting in the formation of deep gullies in the foot slopes that eventually result in the heavy siltation of tanks and reservoirs in the microwatershed.



Fig 2.4 Natural vegetation of Yadgir Rf microwatershed

2.7 Land Utilization

About 72 per cent area (Table 2.2) in Yadgir district is cultivated at present. An area of about 2 per cent is permanently under pasture, 20 per cent under current fallows and 6 per cent under non-agricultural land and 5 per cent under currently barren. Forests occupy an area of about 7 per cent and the tree cover is in a very poor state. Most of the mounds, ridges and bouldery areas have very poor vegetative cover. Major crops grown in the area are sorghum, maize, cotton, sunflower, groundnut, red gram, marigold and sapota. While carrying out land resource inventory, the land use/land cover particulars are collected from all the survey numbers and a current land use map of the microwatershed is prepared. The current land use map prepared shows the arable and non-arable lands, other land uses and different types of crops grown in the area. The current land use map of Yadgir Rf microwatershed is presented in Fig.2.5. The different crops and cropping systems adopted in the microwatershed is presented in Figures 2.6 a & b.

Table 2.2 Land Utilization in Yadgir District

Sl. No.	Agricultural land use	Area (ha)	Per cent
1	Total geographical area	516088	-
2	Total cultivated area	373617	72.4
3	Area sown more than once	74081	14.3
4	Cropping intensity	-	119.8
5	Trees and grooves	737	0.14
6	Forest	33773	6.54
7	Cultivable wasteland	2385	0.46
8	Permanent Pasture land	11755	2.28
9	Barren land	27954	5.41
10	Non- Agriculture land	29623	5.73
11	Current Fallows	105212	20.4




Fig.2.5 Current Land Use map of Yadgir Rf microwatershed





Fig. 2.6. Different Crops and Cropping Systems in Yadgir Rf microwatershed

SURVEY METHODOLOGY

The purpose of land resource inventory is to delineate similar areas (soil series and phases), which respond or expected to respond similarly to a given level of management. This was achieved in Yadgir Rf microwatershed by the detailed study of all the soil characteristics (depth, texture, colour, structure, consistence, coarse fragments, porosity, soil reaction, soil horizons etc.) and site characteristics (slope of the land, erosion, drainage, occurrence of rock fragments etc.) followed by grouping of similar areas based on soil-site characteristics into homogeneous (management units) units, and showing the area extent and their geographic distribution on the microwatershed cadastral map. The detailed survey at 1:7920 scale was carried out in an area of 601.41 ha. The methodology followed for carrying out land resource inventory was as per the guidelines given in Soil Survey Manual (IARI, 1971; Soil Survey Staff, 2006; Natarajan *et al.*, 2015) which is briefly described below.

3.1 Base Maps

The detailed survey of the land resources occurring in the microwatershed was carried out by using digitized cadastral map and IRS satellite imagery as base supplied by KRSRAC. The cadastral map shows field boundaries with their survey numbers, location of tanks, streams and other permanent features of the area (Fig. 3.1). Apart from the cadastral map, remote sensing data products from Cartosat-1 and LISS IV merged at the scale of 1:7920 were used in conjunction with the cadastral map to identify the landscapes, landforms and other surface features. The imagery helped in the identification and delineation of boundaries between hills, uplands and lowlands, water bodies, forest and vegetated areas, roads, habitations and other cultural features of the area (Fig. 3.2). The cadastral map was overlaid on the satellite imagery (Fig. 3.3) that helps to identify the parcel boundaries and other permanent features. Apart from cadastral maps and images, toposheets of the area (1:50,000 scale) were also used for initial traversing, identification of geology and landforms, drainage features, present land use and also for selection of transects in the microwatershed.

3.2 Image Interpretation for Physiography

False Colour Composites (FCCs) of Cartosat-I and LISS-IV merged satellite data covering microwatershed area was visually interpreted using image interpretation elements and all the available collateral data with local knowledge. The delineated physiographic boundaries were transferred on to a cadastral map overlaid on satellite imagery. Physiographically, the area has been identified as granite and granite gneiss landscape. It was divided into five landforms, *viz.*; ridges and mounds, gently and very gently sloping uplands and lowlands based on slope and image characteristics. They were further subdivided into physiographic/image interpretation units based on image characteristics. The image interpretation legend for physiography is given below.

Image Interpretation Legend for Physiography

G- Granite Gneiss Landscape

G1	Hills/ Ridges/ Mounds
G11	Summits
G12	Side slopes
G121	Side slopes with dark grey tones
G2	Uplands
G21	Summits
G22	Gently sloping uplands
G221	Gently sloping uplands, yellowish green (eroded)
G222	Gently sloping uplands, yellowish white (severely eroded)
G23	Very gently sloping uplands
G231	Very gently sloping uplands, yellowish green
G232	Very gently sloping uplands, medium green and pink
G233	Very gently sloping uplands, pink and green (scrub land)
G234	Very gently sloping uplands, medium greenish grey
G235	Very gently sloping uplands, yellowish white (eroded)
G236	Very gently sloping uplands, dark green
G237	Very gently sloping uplands, medium pink (coconut garden)
G238	Very gently sloping uplands, pink and bluish white (eroded)
G24	Valleys/ lowlands
G241	Valleys, pink tones
G242	Valleys gray mixed with pink tones




Fig 3.1 Scanned and Digitized Cadastral map of Yadgir Rf microwatershed




Fig.3.2 Satellite Image of Yadgir Rf microwatershed




Fig.3.3 Cadastral map overlaid on IRS PAN+LISS IV merged imagery of Yadgir Rf microwatershed

3.3 Field Investigation

The field boundaries and survey numbers given on the cadastral sheet were located on the ground by following permanent features like roads, cart tracks, *nallas*, streams, tanks etc., and wherever changes were noticed, they were incorporated on the microwatershed cadastral map. Preliminary traverse of the microwatershed was carried out with the help of cadastral map, imagery and toposheets. While traversing, landforms and physiographic units identified were checked and preliminary soil legend was prepared by studying soils at few selected places. Then, intensive traversing of each physiographic unit like hills, ridges, uplands and valleys was carried out. Based on the variability observed on the surface, transects (Fig. 3.4) were selected across the slope covering all the landform units in the microwatershed (Natarajan and Dipak Sarkar, 2010).




Fig: 3.4. Location of profiles in a transect

In the selected transect, soil profiles were located (Fig. 3.4) at closely spaced intervals to take care of any change in the land features like break in slope, erosion, gravel, stones etc. In the selected sites, profiles (vertical cut showing the soil layers from surface to the rock) were opened upto 200 cm or to the depth limited by rock or hard substratum and studied in detail for all their morphological and physical characteristics. The soil and site characteristics were recorded for all profile sites on a standard proforma as per the guidelines given in USDA Soil Survey Manual (Soil Survey Staff, 2012). Apart from the transect study, profiles were also studied at random, almost like in a grid pattern, outside the transect areas.

Based on the soil characteristics, the soils were grouped into different soil series. Soil series is the most homogeneous unit having similar horizons and properties and behaves similarly for a given level of management. Soil depth, texture, colour, kind of horizon and horizon sequence, calcareousness, amount and nature of gravel present, nature of substratum etc, were used as the major differentiating characteristics for

identifying soil series occurring in the area. The differentiating characteristics used for identifying the soil series are given in Table 3.1. Based on the above characteristics, 2 soil series were identified in the Yadgir Rf microwatershed.

Table 3.1 Differentiating characteristics used for identifying soil series
(Characteristics are of Series Control Section)

Soils of Granite gneiss Landscape							
Sl. no	Soil Series	Depth (cm)	Colour (moist)	Texture	Gravel (%)	Horizon sequence	Calcareousness
1	DPL (Duppali)	50-75	7.5YR 3/3 5YR 3/4	sc	-	Ap-Bt	-
2	HTK (Hattikuni)	25-50	10YR 4/6, 4/4 7.5YR 4/4, 3/3	sl	10-25	Ap-Ac	-

3.4 Soil Mapping

The area under each soil series was further separated into soil phases and their boundaries delineated on the cadastral map based on the variations observed in the texture of the surface soil, slope, erosion, presence of gravel, stoniness etc. A soil phase is a subdivision of soil series based mostly on surface features that affect its use and management. The soil mapping units are shown on the map (Fig.3.5) in the form of symbols. During the survey many profile pits, few minipits and a few auger bores representing different landforms occurring in the microwatershed were studied. In addition to the profile study, spot observations in the form of minipits, road cuts, terrace cuts etc., were studied to validate the soil boundaries on the soil map.

The soil map shows the geographic distribution of 2 mapping units representing 2 soil series occurring in the microwatershed. The soil map unit (soil legend) description is presented in Table 3.2. The soil phase map (management units) shows the distribution of 2 soil phases mapped in the microwatershed. Each mapping unit (soil phase) delineated on the map has similar soil and site characteristics. In other words, all the farms or survey numbers included in one phase will have similar management needs and have to be treated accordingly.

3.5 Land Management Units

The 2 soil phases identified and mapped in the microwatershed were grouped into 2 Land Management Units (LMU's) for the purpose of preparing a Proposed Crop Plan for sustained development of the microwatershed. The database (soil phases) generated under LRI was utilized for identifying Land Management Units (LMU's) based on the management needs. One or more than one soil site characteristic having influence on the management have been chosen for identification and delineation of LMUs. For Yadgir Rf microwatershed, five soil and site characteristics, namely soil depth, soil texture, slope, erosion and gravel content have been considered for defining LMUs. The Land Management Units are expected to behave similarly for a given level of management.

3.6 Laboratory Characterization

Soil samples were collected from representative master profiles for laboratory characterization by following the methods outlined in the Laboratory Manual (Sarma *et al*, 1987). Surface soil samples collected from farmer's fields (50 samples) for fertility status (major and micronutrients) at 320 m grid interval in the year 2018 were analyzed in the laboratory (Katyal and Rattan, 2003). By linking the soil fertility data to the survey numbers through GIS, soil fertility maps were generated by using Kriging method for the microwatershed.

Table 3.2 Soil map unit description of Yadgir Rf microwatershed

Soil No*	Soil Series	Soil Phase	Mapping Unit Description	Area in ha (%)
Soil of Granite Gneiss Landscape				
	DPL		Duppali soils are moderately shallow (50-75 cm), well drained, have dark brown to dark reddish brown, sandy clay red soils occurring on very gently sloping uplands under cultivation	9 (1.66)
25		DPLcB2	Sandy loam surface, slope 1-3%, moderate erosion	9 (1.66)
	HTK		Hattikuni soils are shallow (25-50 cm), well drained, have dark yellowish brown sandy loam soils occurring on very gently sloping uplands under cultivation	327 (61.29)
161		HTKbB2g1	Loamy sand surface, slope 1-3%, moderate erosion, gravelly (15-35%)	327 (61.29)

* Soil map unit numbers are continuous for the taluk, not for the microwatershed




Fig 3.5 Soil phase or Management Units - Yadgir Rf microwatershed

THE SOILS

Detailed information pertaining to the nature, extent and their distribution of different kinds of soils occurring in Yadgir Rf microwatershed is provided in this chapter. The microwatershed area has been identified as granite gneiss landscape based on geology. In this 2 soil series are identified. Soil formation is the result of the combined effect of environmental and terrain factors that are reflected in soil morphology. In the granite gneiss landscape, it is by parent material, relief and climate.

A brief description of each of the 2 soil series identified followed by 2 soil phases (management units) mapped under each series are furnished below. The physical and chemical characteristics of soil series identified in Yadgir Rf microwatershed are given in Table 4.1 along with soil classification. The soils in any one map unit differ from place to place in their depth, texture, slope, gravelliness, erosion or any other site characteristic that affect management. The soil phase map can be used for identifying the suitability of areas for growing specific crops or for other alternative uses and also for deciding the type of conservation structures needed. The detailed information on soil and site-characteristics like soil depth, surface soil texture, slope, erosion, gravelliness, AWC, LCC etc, with respect to each of the soil phase identified is given village/survey number wise for the microwatershed in Appendix-I.

4.1 Soils of granite gneiss landscape

In this landscape, 2 soil series are identified and mapped. Of these, HTK series occupies maximum area of 327 ha (61%) followed by DPL 9 ha (2%). Brief description of these two series identified and number of soil phases mapped is given below.

4.1.1 Hattikuni (HTK) Series: Hattikuni soils are shallow (25-50 cm), well drained, have dark brown to dark yellowish brown sandy loam soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation. The Hattikuni series has been classified as a member of the mixed, isohyperthermic family of Lithic Ustipsamments.

The thickness of the soil ranges from 36 to 50 cm. The thickness of A horizon ranges from 8 to 12 cm. Its colour is in 10YR and 7.5 YR hue with value 3 to 4 and chroma 4 to 6. The texture varies from loamy sand to sandy loam. The thickness of subsurface horizon ranges from 28 to 42 cm. Its colour is in 10 YR and 7.5 YR hue with value 3 to 4 and chroma 4 to 6. Its texture varies from loamy sand to sand and sandy loam. The available water capacity is very low (<50 mm/m). Only one phase was identified and mapped.



Landscape and Soil Profile characteristics of Hattikuni (HTK) Series

4.1.2 Duppali (DPL) Series: Duppali soils are moderately shallow (50-75 cm), well drained, have dark brown to dark reddish brown, sandy clay soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation. The Duppali series has been classified as a member of the fine, mixed, isohyperthermic family of Typic Haplustalfs.

The thickness of the solum ranges from 51-75 cm. Thickness of A horizon ranges from 8 to 15 cm. Its colour is in hue 10 YR with value 4 and chroma 3 to 6. The texture varies from loamy sand to sandy clay. The thickness of B horizon ranges from 55 to 65 cm. Its colour is in 7.5 YR and 5 YR hue with value 3 to 4 and chroma 2 to 4. The texture is sandy clay. The available water capacity is low (51-100 mm/m). Only one phase was identified and mapped.



Landscape and Soil Profile characteristics of Duppali (DPL) Series

Table 4.1 Physical and Chemical Characteristics of Soil Series Identified in Yadgir Rf Microwatershed

Soil Series: Hattikuni (HTK), Pedon: R-7

Location: 16°50'46.5"N 77°10'16.4"E, Yaddalli village, Hattikuni hobli, Yadgir taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bengaluru **Classification:** Mixed, isohyperthermic, Lithic Ustipsamments

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-12	Ap	90.89	5.62	3.49	8.50	13.46	29.86	29.55	9.51	20	s	7.73	3.16
12-22	A1	89.97	6.53	3.50	7.19	13.48	29.48	29.79	10.03	20	s	8.00	3.05
22-45	A2	87.20	6.43	6.38	11.09	14.42	31.55	7.16	22.98	40	ls	7.67	3.96

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO ₃	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP
	Water	CaCl ₂	M KCl				Ca	Mg	K	Na	Total				
				dS m ⁻¹	%	%	cmol kg ⁻¹					%	%		
0-12	6.81	-	-	0.062	0.07	-	2.35	0.50	0.16	0.01	3.02	3.0	0.86	100	0.38
12.0-22	6.80	-	-	0.050	0.21	-	1.67	0.30	0.09	0.01	2.07	2.4	0.69	86.30	0.45
22-45	6.85	-	-	0.044	0.19	-	1.82	0.42	0.10	0.06	2.40	2.6	0.41	92.41	2.17

Contd...

Soil Series: Duppali (DPL) **Pedon:** R-4

Location: 16°37'45.8"N 77°18'93.2"E, Neelahalli village, Balichakra hobli, Yadgir taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bengaluru **Classification:** Fine, mixed, isohyperthermic Typic Haplustalfs

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-7	Ap	85.28	5.38	9.34	13.40	26.09	19.90	20.51	5.38	-	ls	9.30	4.92
7-39	Bt1	48.50	7.08	44.42	16.85	10.41	10.94	6.97	3.33	-	sc	21.31	16.82
39-65	Bt2	50.95	5.29	43.76	23.57	10.36	8.77	5.50	2.75	-	sc	21.99	17.50

Depth (cm)	pH (1:2.5)			E.C. (1:2.5) dS m ⁻¹	O.C. %	CaCO ₃ %	Exchangeable bases					CEC	CEC/Clay	Base saturation %	ESP %
	Water	CaCl ₂	M KCl				Ca	Mg	K	Na	Total				
0-7	6.92	-	-	0.122	0.92	0.00	4.73	1.61	0.19	0.01	6.54	7.10	0.76	92	0.09
7-39	7.00	-	-	0.060	0.62	0.00	13.57	4.78	0.12	0.40	18.87	19.30	0.43	98	2.06
39-65	6.87	-	-	0.072	0.41	0.00	13.69	4.57	0.19	0.65	19.10	19.90	0.45	96	3.25

INTERPRETATION FOR LAND RESOURCE MANAGEMENT

The most important soil and site characteristics that affect the land use and conservation needs of an area are land capability, soil depth, soil texture, coarse fragments, available water capacity, soil slope, soil erosion, soil reaction etc. These are interpreted from the data base generated through land resource inventory and several thematic maps are generated. These would help in identifying the areas suitable for growing crops and, soil and water conservation measures and structures needed thus helping to maintain good soil health for sustained crop production. The various interpretative and thematic maps generated are described below.

5.1 Land Capability Classification

Land capability classification is an interpretative grouping of soil map units (soil phases) mainly based on inherent soil characteristics, external land features and environmental factors that limit the use of land for agriculture, pasture, forestry, or other uses on a sustained basis (IARI, 1971). The land and soil characteristics used to group the land resources in an area into various land capability classes, subclasses and units are

Soil Characteristics: Depth, texture, gravelliness, calcareousness.

Land characteristics: Slope, erosion, drainage, rock outcrops.

Climate: Total rainfall and its distribution, and length of crop growing period.

The Land capability classification system is divided into land capability classes, subclasses and units based on the level of information available. Eight land capability classes are recognized. They are

Class I: They are very good lands that have no limitations or very few limitations that restrict their use.

Class II: They are good lands that have minor limitations and require moderate conservation practices.

Class III: They are moderately good lands that have moderate limitations that reduce the choice of crops or that require special conservation practices.

Class IV: They are fairly good lands that have very severe limitations that reduce the choice of crops or that require very careful management.

Class V: Soils in these lands are not likely to erode, but have other limitations like wetness that are impractical to remove and as such not suitable for agriculture, but suitable for pasture or forestry with minor limitations.

Class VI: The lands have severe limitations that make them generally unsuitable for cultivation, but suitable for pasture or forestry with moderate limitations.

Class VII: The lands have very severe limitations that make them unsuitable for cultivation, but suitable for pasture or forestry with major limitations.

Class VIII: Soil and other miscellaneous areas (rock lands) that have very severe limitations that nearly preclude their use for any crop production, but suitable for wildlife, recreation and installation of wind mills.

The land capability subclasses are recognised based on the dominant limitations observed within a given land capability class. The subclasses are designated by adding a lower case letter like 'e', 'w', 's', or 'c' to the class numeral. The subclass "e" indicates that the main hazard is risk of erosion, "w" indicates drainage or wetness as a limitation for plant growth, "s" indicates shallow soil depth, coarse or heavy textures, calcareousness, salinity/alkalinity or gravelliness and "c" indicates limitation due to climate.

The land capability subclasses have been further subdivided into land capability units based on the kinds of limitations present in each subclass. Ten land capability units are used in grouping the soil map units. They are stony or rocky (0), erosion hazard (slope, erosion) (1), coarse texture (sand, loamy sand, sandy loam) (2), fine texture (cracking clay, silty clay) (3), slowly permeable subsoil (4), coarse underlying material (5), salinity/alkali (6), stagnation, overflow, high ground water table (7), soil depth (8) and fertility problems (9). The capability units thus identified have similar soil and land characteristics that respond similarly to a given level of management. The soils of the microwatershed have been classified upto land capability subclass level.

The 2 soil map units identified in the Yadgir Rf microwatershed are grouped under 2 land capability classes and 2 subclasses. An area of 336 ha (63%) in the microwatershed is suitable for agriculture. About 198 ha (37%) area is having rock outcrops (Fig. 5.1).

Good cultivable lands (Class II) cover an area of about 2 per cent and are distributed in the southern part of the microwatershed with minor problems of soil and erosion. Moderately good cultivable lands (Class III) cover maximum area of about 61 per cent and are distributed in the major part of the microwatershed with moderate problems of soil and erosion.




Fig. 5.1 Land Capability Classification map of Yadgir Rf microwatershed

5.2 Soil Depth

Soil depth refers to the depth of the soil occurring above the parent material or hard rock. The depth of the soil determines the effective rooting depth for plants and in accordance with soil texture, mineralogy and gravel content, the capacity of the soil column to hold water and nutrient availability. Soil depth is one of the most important soil characteristic that is used in differentiating soils into different soil series. The soil depth classes used in identifying soils in the field are very shallow (<25 cm), shallow (25-50 cm), moderately shallow (50-75 cm), moderately deep (75-100 cm), deep (100-150 cm) and very deep (>150 cm). They were used to classify the soils into different depth classes and a soil depth map was generated. The area extent and their geographical distribution in the microwatershed is given in Fig. 5.2.

Shallow (25-50) soils occupy a maximum area of about 327 ha (61%) and are distributed in the major part of the microwatershed. Moderately shallow (50-75 cm) soils occupy an area of 9 ha (2%) and are distributed in the southern part of the microwatershed.

The problematic soils cover entire area of the microwatershed where the soils are shallow and are suitable for short duration crops




Fig. 5.2 Soil depth map of Yadgir Rf microwatershed

5.3 Surface Soil Texture

Texture is an expression to indicate the coarseness or fineness of the soil as determined by the relative proportion of primary particles of sand, silt and clay. It has a direct bearing on the structure, porosity, adhesion and consistence. The surface layer of a soil to a depth of about 25 cm is the layer that is most used by crops and plants. The surface soil textural class provides a guide to understanding soil-water retention and availability, nutrient holding capacity, infiltration, workability, drainage, physical and chemical behaviour, microbial activity and crop suitability. The textural classes used for LRI were used to classify and a surface soil texture map was generated. The area extent and their geographical distribution in the microwatershed is shown in Figure 5.3.

Maximum area of about 327 ha (61%) of the microwatershed has sandy soils at the surface and are distributed in the major part of the microwatershed. An area of 9 ha (2%) has soils that are loamy and are distributed in the southern part of the microwatershed. Loamy soils have high potential for soil-water retention and availability, and nutrient retention and availability. The sandy soils are productive for root and tuber crops, but these soils have the major limitations of moisture and nutrient retention capacity, hence frequent and shallow irrigation with balanced fertilizer application is to be followed in order to get better crop yields.




Fig. 5.3 Surface soil texture map of Yadgir Rf microwatershed

5.4 Soil Gravelliness

Gravel is the term used for describing coarse fragments between 2 mm and 7.5 cm diameter and stones for those between 7.5 cm and 25 cm. The presence of gravel and stones in soil reduces the volume of soil responsible for moisture and nutrient storage, drainage, infiltration and runoff, and hinders plant growth by impeding root growth and seedling emergence, intercultural operations and farm mechanization. The gravelliness classes used in LRI were used to classify the soils and using these classes a gravelliness map was generated. The area extent and their geographic distribution in the microwatershed is shown in Figure 5.4.

Non gravelly (<15%) soils cover an area of about 9 ha (2%) and are distributed in the southern part of the microwatershed. Maximum area of about 327 ha (61%) is gravelly (15-35%) and are distributed in the major part of the microwatershed.

The problem soils (61%) that are gravelly (15-35%), where only short or medium duration crops can be grown. The most productive soils (2%) that are non gravelly (<15%), where all climatically adapted long duration crops can be grown.




Fig. 5.4 Soil gravelliness map of Yadgir Rf microwatershed

5.5 Available Water Capacity

The soil available water capacity (AWC) is estimated based on the ability of the soil column to retain water between the tensions of 0.33 and 15 bar in a depth of 100 cm or the entire solum if the soil is shallower. The AWC of the soils (soil series) as estimated by considering the soil texture, mineralogy, soil depth and gravel content (Sehgal *et al.*, 1990) and accordingly the soil map units were grouped into five AWC classes *viz.*, very low (<50 mm/m), low (50-100 mm/m), medium (100-150 mm/m), high (150-200 mm/m) and very high (>200 mm/m) and using these values, an AWC map was generated. The area extent and their geographic distribution of different AWC classes in the microwatershed is given in Figure 5.5.

Maximum area of about 327ha (61%) in the microwatershed have soils that are very low (<50 mm/m) in available water capacity and are distributed in the major part of the microwatershed. An area of about 9 ha (2%) is low (51-100 mm/m) in available water capacity and are distributed in the southern part of the microwatershed.

Entire area in the microwatershed has soils that are problematic with regard to available water capacity. Here, only short duration crops can be grown and the probability of crop failure is very high. These areas are best put to other alternative uses.




Fig. 5.5 Soil available water capacity map of Yadgir Rf microwatershed

5.6 Soil Slope

Soil slope refers to the inclination of the surface of the land. It is defined by gradient, shape and length, and is an integral feature of any soil as a natural body. Slope is considered important in soil genesis, land use and land development. The length and gradient of slope influences the rate of runoff, infiltration, erosion and deposition. The soil map units were grouped into two slope classes and a slope map was generated showing the area extent and their geographic distribution in the microwatershed (Fig. 5.6).

Entire area of the microwatershed falls under very gently sloping (1-3% slope) lands and are distributed in all parts of the microwatershed.

Entire area in the microwatershed has soils that have high potential in respect of soil slopes. In these areas, all climatically adapted annual and perennial crops can be grown without much soil and water conservation and other land development measures.




Fig. 5.6 Soil slope map of Yadgir Rf microwatershed

5.7 Soil Erosion

Soil erosion refers to the wearing away of the earth's surface by the forces of water, wind and ice involving detachment and transport of soil by raindrop impact. It is used for accelerated soil erosion resulting from disturbance of the natural landscape by burning, excessive grazing and indiscriminate felling of forest trees and tillage, all usually by man. The erosion classes showing an estimate of the current erosion status as judged from field observations in the form of rills, gullies or a carpet of gravel on the surface are recorded. Four erosion classes, viz, slight erosion (e1), moderate erosion (e2), severe erosion (e3) and very severe erosion (e4) are recognized. The soil map units were grouped into different erosion classes and a soil erosion map generated. The area extent and their spatial distribution in the microwatershed is given in Figure 5.7.

Entire area of the microwatershed falls under moderately eroded (e2 class) lands and are distributed in all parts of the microwatershed.

Entire area in the microwatershed is problematic because of moderate erosion. For these areas, taking up of soil and water conservation and other land development measures are needed.




Fig. 5.7 Soil erosion map of Yadgir Rf microwatershed

FERTILITY STATUS

Soil fertility plays an important role in increasing crop yield. The adoption of high yielding varieties that require high amounts of nutrients has resulted in deficiency symptoms in crops and plants due to imbalanced fertilization and poor inherent fertility status as these areas are characterised by low rainfall and high temperatures. Hence, it is necessary to know the fertility (macro and micro nutrients) status of the soils of the watersheds for assessing the kind and amount of fertilizers required for each of the crop intended to be grown. For this purpose, the surface soil samples collected from the grid points (one soil sample at every 320 m interval) all over the microwatershed through land resource inventory in the year 2018 were analysed for pH, EC, organic carbon, available phosphorus and potassium, and for micronutrients like zinc, boron, copper, iron manganese and secondary nutrient sulphur.

Soil fertility data generated has been assessed and individual maps for all the nutrients for the microwatershed have been generated using Kriging method under GIS. The village/survey number wise fertility data for the microwatershed is given in Appendix-II.

6.1 Soil Reaction (pH)

The soil analysis of the Yadgir Rf microwatershed for soil reaction (pH) showed that an area of about 0.13 ha (<1%) is slightly acid (pH 6.0-6.5) and is distributed in the western part of the microwatershed. Maximum area of 333 ha (62%) is neutral (pH 6.5-7.3) and are distributed in the major part of the microwatershed. An area of about 3 ha (<1%) is slightly alkaline (pH 7.3-7.8) and are distributed in the eastern part of the microwatershed (Fig. 6.1). In all, major area of about 333 ha is neutral and 3 ha is under alkaline.

6.2 Electrical Conductivity (EC)

The electrical conductivity of the soils of the entire microwatershed area is <2 dS m⁻¹ (Fig 6.2) and as such the soils are non-saline.

6.3 Organic Carbon

The soil organic carbon content (an index of available nitrogen) in the soils of the microwatershed is high (>0.75 %) in 196 ha (37%) and are distributed in the eastern, central and northern part of the microwatershed. Medium (0.5-0.75%) in about 140 ha (26%) and are distributed in the central, northern and southern part of the microwatershed (Fig. 6.3).




Fig.6.1 Soil reaction (pH) map of Yadgir Rf microwatershed




Fig.6.2 Electrical conductivity (EC) map of Yadgir Rf microwatershed




Fig.6.3 Soil organic carbon map of Yadgir Rf microwatershed

6.4 Available Phosphorus

Available phosphorus content is low (<23 kg/ha) in an area of about 86 ha (16%) and are distributed in the central, northwestern and southern part of the microwatershed. Maximum area about 250 ha (47%) is in medium (23-57 kg/ha) and are distributed in the major area of the microwatershed (Fig. 6.4).

6.5 Available Potassium

Available potassium content is medium (145-337 kg/ha) in the entire microwatershed area (Fig. 6.5).

6.6 Available Sulphur

Available sulphur content is low (<10 ppm) in the entire microwatershed area (Fig. 6.6).

6.7 Available Boron

Available boron content is low (<0.5 ppm) in the entire microwatershed area (Fig. 6.7).

6.8 Available Iron

Available iron content is deficient (<4.5 ppm) in an area of about 84 ha (16%) and are distributed in the eastern part of the microwatershed. Maximum area of about 252 ha (47%) is under sufficient (>4.5 ppm) and are distributed in the major area of the microwatershed (Fig 6.8).

6.9 Available Manganese

Available manganese content is sufficient (>1.0 ppm) in the entire microwatershed area (Fig 6.9).

6.10 Available Copper

Available copper content is sufficient (>0.2 ppm) in the entire microwatershed area (Fig 6.10).




Fig.6.4 Soil available phosphorus map of Yadgir Rf microwatershed




Fig.6.5 Soil available potassium map of Yadgir Rf microwatershed




Fig.6.6 Soil available sulphur map of Yadgir Rf microwatershed




Fig.6.7 Soil available boron map of Yadgir Rf microwatershed




Fig.6.8 Soil available iron map of Yadgir Rf microwatershed




Fig.6.9 Soil available manganese map of Yadgir Rf microwatershed




Fig.6.10 Soil available copper map of Yadgir Rf microwatershed

6.11 Available Zinc

Available zinc content is deficient in a maximum area of 330 ha (62%) (<0.6 ppm) and are distributed in the major part of the microwatershed. Sufficient in 7 ha (1%) (>0.6 ppm) and is distributed in the central and eastern part of the microwatershed (Fig 6.11).




Fig.6.11 Soil available zinc map of Yadgir Rf microwatershed

LAND SUITABILITY FOR MAJOR CROPS

The soil and land resource units (soil phases) of Yadgir Rf microwatershed were assessed for their suitability for growing food, fodder, fibre and other horticulture crops by following the procedure as outlined in FAO, 1976 and 1983. Crop requirements were developed for each of the crop from the available research data and also by referring to Naidu *et. al.* (2006) and Natarajan *et. al* (2015). The crop requirements were matched with the soil and land characteristics (Table 7.1) to arrive at the crop suitability (Table 7.2 to 7.30) are given at the end of the chapter. In FAO land suitability classification, two orders are recognized. Order S-Suitable and Order N-Not suitable. The orders have classes, subclasses and units. Order-S has three classes, Class S1-Highly Suitable, Class S2-Moderately Suitable and Class S3- Marginally Suitable. Order N has two classes, N1- Currently not Suitable and N2- Permanently not Suitable. There are no subclasses within the Class S1 as they will have very minor or no limitations for crop growth. Classes S2 and S3 are divided into subclasses based on the kinds of limitations encountered. The limitations that affect crop production are ‘c’ for erratic rainfall and its distribution and length of growing period (LGP), ‘e’ for erosion hazard, ‘r’ for rooting condition, ‘t’ for lighter or heavy texture, ‘g’ for gravelliness or stoniness, ‘n’ for nutrient availability, ‘l’ for topography, ‘m’ for moisture availability, ‘w’ for drainage and ‘z’ for calcareousness. These limitations are indicated as lower case letters to the class symbol. For example, moderately suitable lands with the limitations of soil depth and erosion are designated as S2re. For the microwatershed, the soil mapping units were evaluated and classified up to subclass level.

Using the above criteria, the soil map units of the microwatershed were evaluated and land suitability maps for 28 major agricultural and horticultural crops were generated. The detailed information on the kind of suitability of each of the soil phase for the crops assessed are given village/ survey number wise for the microwatershed in Appendix-III.

7.1 Land Suitability for Sorghum (*Sorghum bicolor*)

Sorghum is one of the major crop grown in Karnataka in an area of 10.47 lakh ha in Bijapur, Gulbarga, Raichur, Bidar, Belgaum, Dharwad, Bellary, Chitradurga, Mysore and Tumakuru districts. The crop requirements for growing sorghum (Table 7.2) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing sorghum was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.1.

There are no highly suitable (Class S1) lands available for growing sorghum in the microwatershed. An area of about 9 ha (2%) is moderately suitable (Class S2) for growing sorghum and are distributed in the southern part of the microwatershed. They

have minor limitations of rooting depth and calcareousness. Maximum area of about 327 ha (61%) is marginally suitable (Class S3) for growing sorghum and is distributed in the major part of the microwatershed with moderate limitations of rooting depth and texture.




Fig. 7.1 Land suitability map of Sorghum

7.2 Land Suitability for Maize (*Zea mays*)

Maize is one of the most important food crop grown in an area of 13.37 lakh ha in almost all the districts of the State. The crop requirements for growing maize (Table 7.3) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing maize was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.2.

There are no highly suitable (Class S1) lands available for growing maize in the microwatershed. An area of about 9 ha (2%) is moderately suitable (Class S2) for growing maize and are distributed in the southern part of the microwatershed. They have minor limitations of rooting depth and calcareousness. Maximum area of about 327 ha (61%) is marginally suitable (Class S3) for growing maize and is distributed in the major part of the microwatershed with moderate limitations of rooting depth and texture.




Fig. 7.2 Land suitability map of Maize

7.3 Land Suitability for Bajra (*Pennisetum glaucum*)

Bajra is one of the most important millet crop grown in an area of 2.34 lakh ha in the northern districts of Karnataka state. The crop requirements for growing bajra (Table 7.4) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing bajra was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.3.

There are no highly suitable (Class S1) lands available for growing bajra in the microwatershed. An area of about 9 ha (2%) is moderately suitable (Class S2) for growing bajra and are distributed in the southern part of the microwatershed. They have minor limitations of rooting depth and calcareousness. Maximum area of about 327 ha (61%) is marginally suitable (Class S3) for growing bajra and is distributed in the major part of the microwatershed with moderate limitations of rooting depth and texture.




Fig. 7.3 Land suitability map of Bajra

7.4 Land Suitability for Groundnut (*Arachis hypogaea*)

Groundnut is one of the major oilseed crop grown in an area of 6.54 lakh ha in Karnataka in most of the districts either as rainfed or irrigated crop. The crop requirements for growing groundnut (Table 7.5) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing groundnut was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.4.

There are no highly suitable (Class S1) lands available for growing groundnut in the microwatershed. An area of about 9 ha (2%) is moderately suitable (Class S2) for growing groundnut and are distributed in the southern part of the microwatershed. They have minor limitations of rooting depth and calcareousness. Maximum area of about 327 ha (61%) is marginally suitable (Class S3) for growing groundnut and is distributed in the major part of the microwatershed with moderate limitations of rooting depth and texture.




Fig. 7.4 Land suitability map of Groundnut

7.5 Land Suitability for Sunflower (*Helianthus annuus*)

Sunflower is one of the most important oilseed crop grown in an area of 3.56 lakh ha in the State in all the districts. The crop requirements for growing sunflower (Table 7.6) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing sunflower was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.5.

There are no highly suitable (Class S1) and moderately suitable (Class S2) lands available for growing sunflower in the microwatershed. An area of about 9 ha (2%) is marginally suitable (Class S3) for growing sunflower and are distributed in the southern part of the microwatershed. They have moderate limitations of rooting depth and calcareousness. Maximum area of about 327 ha (61%) is currently not suitable (Class N1) for growing sunflower and is distributed in the major part of the microwatershed with severe limitation of rooting depth.




Fig. 7.5 Land suitability map of Sunflower

7.6 Land suitability for Red gram (*Cajanus Cajan*)

Red gram is one of the most important pulse crop grown in an area of 7.28 lakh ha in almost all the districts of the State. The crop requirements for growing red gram (Table 7.7) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing red gram was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.6.

There are no highly suitable (Class S1) and moderately suitable (Class S2) lands available for growing red gram in the microwatershed. An area of about 9 ha (2%) is marginally suitable (Class S3) for growing red gram and are distributed in the southern part of the microwatershed. They have moderate limitations of rooting depth and calcareousness. Maximum area of about 327 ha (61%) is currently not suitable (Class N1) for growing red gram and is distributed in the major part of the microwatershed with severe limitation of rooting depth.




Fig. 7.6 Land suitability map of Red gram

7.7 Land Suitability for Bengal gram (*Cicer aerativum*)

Bengal gram one of is the most important pulse crop grown in about 9.39 lakh ha area in Bijapur, Raichur, Kalaburgi, Dharwad, Belgaum and Bellary districts. The crop requirements for growing Bengal gram (Table 7.8) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing Bengal gram was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.7.

There are no highly suitable (Class S1) lands available for growing Bengal gram in the microwatershed. An area of about 9 ha (2%) is moderately suitable (Class S2) for growing Bengal gram and are distributed in the southern part of the microwatershed. They have minor limitations of rooting depth and calcareousness. Maximum area of about 327 ha (61%) is currently not suitable (Class N1) for growing Bengal gram and is distributed in the major part of the microwatershed with severe limitation of texture.




Fig. 7.7 Land suitability map of Bengal gram.

7.8 Land Suitability for Cotton (*Gossypium hirsutum*)

Cotton is one of the most important fibre crop grown in the State in about 8.75 lakh ha area in Raichur, Dharwad, Belgaum, Kalaburgi, Bijapur, Bidar, Bellary, Chitradurga and Chamarajnagar districts. The crop requirements for growing cotton (Table 7.9) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing cotton was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.8.

There are no highly suitable (Class S1) lands available for growing cotton in the microwatershed. An area of about 9 ha (2%) is moderately suitable (Class S2) for growing cotton and are distributed in the southern part of the microwatershed. They have minor limitations of rooting depth and calcareousness. Maximum area of about 327 ha (61%) is currently not suitable (Class N1) for growing cotton and is distributed in the major part of the microwatershed with severe limitation of texture.




Fig. 7.8 Land suitability map of Cotton

7.9 Land Suitability for Chilli (*Capsicum annuum*)

Chilli is one of the most important spice crop grown in about 0.42 lakh ha in Karnataka state. The crop requirements for growing chilli (Table 7.10) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing chilli was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.9.

There are no highly suitable (Class S1) lands available for growing chilli in the microwatershed. An area of about 9 ha (2%) is moderately suitable (Class S2) for growing chilli and are distributed in the southern part of the microwatershed. They have minor limitations of rooting depth and calcareousness. Maximum area of about 327 ha (61%) is marginally suitable (Class S3) for growing chilli and is distributed in the major part of the microwatershed with moderate limitations of rooting depth and texture.




Fig 7.9 Land suitability map of Chilli

7.10 Land Suitability for Tomato (*Lycopersicon esculentum*)

Tomato is one of the most important vegetable crop grown in about 0.61 lakh ha covering almost all the district of the state. The crop requirements for growing tomato (Table 7.11) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing tomato was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.10.

There are no highly suitable (Class S1) lands available for growing tomato in the microwatershed. An area of about 9 ha (2%) is moderately suitable (Class S2) for growing tomato and are distributed in the southern part of the microwatershed. They have minor limitations of rooting depth and calcareousness. Maximum area of about 327 ha (61%) is marginally suitable (Class S3) for growing tomato and is distributed in the major part of the microwatershed with moderate limitations of rooting depth and texture.




Fig 7.10 Land suitability map of Tomato

7.11 Land Suitability for Brinjal (*Solanum melongena*)

Brinjal is one of the most important vegetable crop grown in the state. The crop requirements for growing brinjal (Table 7.12) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing brinjal was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.11.

There are no highly suitable (Class S1) lands available for growing brinjal in the microwatershed. An area of about 9 ha (2%) is moderately suitable (Class S2) for growing brinjal and are distributed in the southern part of the microwatershed. They have minor limitation of rooting depth. Maximum area of about 327 ha (61%) is marginally suitable (Class S3) for growing brinjal and is distributed in the major part of the microwatershed with moderate limitations of rooting depth and texture.




Fig 7.11 Land suitability map of Brinjal

7.12 Land Suitability for Onion (*Allium cepa L.*)

Onion is one of the most important vegetable crop grown in the state. The crop requirements for growing onion (Table 7.13) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing onion was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.12.

There are no highly suitable lands (Class S1) available for onion in the microwatershed. Moderately suitable lands (Class S2) cover small area of 9 ha (2%) and are distributed in southern part of the microwatershed with minor limitation of rooting depth. Marginally suitable (Class S3) lands occupy major area of the microwatershed 327 ha (37%) with moderate limitation of rooting depth.




Fig 7.12 Land Suitability map of Onion

7.13 Land Suitability for Bhendi (*Abelmoschus esculentus*)

Bhendi is one of the most important vegetable crop grown in the state. The crop requirements for growing bhendi (Table 7.14) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing bhendi was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.13.

There are no highly suitable (Class S1) lands available for growing bhendi in the microwatershed. An area of about 9 ha (2%) is moderately suitable (Class S2) for growing bhendi and are distributed in the southern part of the microwatershed. They have minor limitation of rooting depth. Maximum area of about 327 ha (61%) is marginally suitable (Class S3) for growing bhendi and is distributed in the major part of the microwatershed with moderate limitation of rooting depth.




Fig 7.13 Land suitability map of Bhendi

7.14 Land Suitability for Drumstick (*Moringa oleifera*)

Drumstick is one of the most important vegetable crop grown in about 2403 ha in the state. The crop requirements for growing drumstick (Table 7.15) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing drumstick was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.14.

There are no highly suitable (Class S1) and moderately suitable (Class S2) lands available for growing drumstick in the microwatershed. An area of about 9 ha (2%) is marginally suitable (Class S3) for growing drumstick and are distributed in the southern part of the microwatershed. They have moderate limitations of rooting depth and calcareousness. Maximum area of about 327 ha (61%) is currently not suitable (Class N1) for growing drumstick and is distributed in the major part of the microwatershed with severe limitation of rooting depth.




Fig 7.14 Land suitability map of Drumstick

7.15 Land suitability for Mango (*Mangifera indica*)

Mango is one of the most important fruit crop grown in an area of 1.73 lakh ha in almost all the districts of the state. The crop requirements (Table 7.16) for growing mango were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing mango was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed is given in Figure 7.15.

Entire area of the microwatershed is currently not suitable (Class N1) for mango with severe limitations of rooting depth and calcareousness.




Fig. 7.15 Land suitability map of Mango

7.16 Land suitability for Guava (*Psidium guajava*)

Guava is one of the most important fruit crop grown in an area of 0.06 lakh ha in almost all the districts of the State. The crop requirements (Table 7.17) for growing guava were matched with the soil-site characteristics (7.1) and a land suitability map for growing guava was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.16.

There are no highly suitable (Class S1) and moderately suitable (Class S2) lands available for growing guava in the microwatershed. An area of about 9 ha (2%) is marginally suitable (Class S3) for growing guava and are distributed in the southern part of the microwatershed. They have moderate limitations of rooting depth and calcareousness. Maximum area of about 327 ha (61%) is currently not suitable (Class N1) for growing guava and is distributed in the major part of the microwatershed with severe limitation of rooting depth.




Fig. 7.16 Land suitability map of Guava

7.17 Land suitability for Sapota (*Manilkara zapota*)

Sapota is one of the most important fruit crop grown in an area of 29373 ha in almost all the districts of the State. The crop requirements (Table 7.18) for growing sapota were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing sapota was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.17.

There are no highly suitable (Class S1) and moderately suitable (Class S2) lands available for growing sapota in the microwatershed. An area of about 9 ha (2%) is marginally suitable (Class S3) for growing sapota and are distributed in the southern part of the microwatershed. They have moderate limitations of rooting depth and calcareousness. Maximum area of about 327 ha (61%) is currently not suitable (Class N1) for growing sapota and is distributed in the major part of the microwatershed with severe limitation of rooting depth.




Fig. 7.17 Land suitability map of Sapota

7.18 Land Suitability for Pomegranate (*Punica granatum*)

Pomegranate is one of the most important fruit crop commercially grown in about 18488 ha in Karnataka, mainly in Bijapur, Bagalkot, Koppal, Gadag and Chitradurga districts. The crop requirements for growing pomegranate (Table 7.19) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing pomegranate was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.18.

There are no highly suitable (Class S1) and moderately suitable (Class S2) lands available for growing pomegranate in the microwatershed. An area of about 9 ha (2%) is marginally suitable (Class S3) for growing pomegranate and are distributed in the southern part of the microwatershed. They have moderate limitations of rooting depth and calcareousness. Maximum area of about 327 ha (61%) is currently not suitable (Class N1) for growing pomegranate and is distributed in the major part of the microwatershed with severe limitation of rooting depth.




Fig 7.18 Land suitability map of Pomegranate

7.19 Land Suitability for Musambi (*Citrus limetta*)

Musambi is one of the important fruit crop grown in an area of 3446 ha in almost all the districts of the State. The crop requirements for growing musambi (Table 7.20) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing musambi was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.19.

There are no highly suitable (Class S1) and moderately suitable (Class S2) lands available for growing musambi in the microwatershed. An area of about 9 ha (2%) is marginally suitable (Class S3) for growing musambi and are distributed in the southern part of the microwatershed. They have moderate limitations of rooting depth and calcareousness. Maximum area of about 327 ha (61%) is currently not suitable (Class N1) for growing musambi and is distributed in the major part of the microwatershed with severe limitation of rooting depth.




Fig. 7.19 Land suitability map of Musambi

7.20 Land Suitability for Lime (*Citrus sp*)

Lime is one of the most important fruit crop grown in an area of 0.11 lakh ha in almost all the districts of the State. The crop requirements for growing lime (Table 7.21) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing lime was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7. 20.

There are no highly suitable (Class S1) and moderately suitable (Class S2) lands available for growing lime in the microwatershed. An area of about 9 ha (2%) is marginally suitable (Class S3) for growing lime and are distributed in the southern part of the microwatershed. They have moderate limitations of rooting depth and calcareousness. Maximum area of about 327 ha (61%) is currently not suitable (Class N1) for growing lime and is distributed in the major part of the microwatershed with severe limitation of rooting depth.




Fig. 7.20 Land suitability map of Lime

7.21 Land Suitability for Amla (*Phyllanthus emblica*)

Amla is one of the medicinal fruit crop grown in almost all the districts of the State. The crop requirements for growing amla (Table 7.22) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing amla was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.21.

There are no highly suitable (Class S1) lands available for growing amla in the microwatershed. An area of about 9 ha (2%) is moderately suitable (Class S2) for growing amla and are distributed in the southern part of the microwatershed. They have minor limitations of rooting depth and calcareousness. Maximum area of about 327 ha (61%) is marginally suitable (Class S3) for growing amla and is distributed in the major part of the microwatershed with moderate limitations of rooting depth and texture.




Fig. 7.21 Land suitability map of Amla

7.22 Land Suitability for Cashew (*Anacardium occidentale*)

Cashew is one of the most important plantation nut crop grown in an area of 0.7 lakh ha in almost all the districts of the State. The crop requirements for growing cashew (Table 7.23) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing cashew was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.22.

There are no highly suitable (Class S1) and moderately suitable (Class S2) lands available for growing cashew in the microwatershed. An area of about 9 ha (2%) is marginally suitable (Class S3) for growing cashew and are distributed in the southern part of the microwatershed. They have moderate limitations of rooting depth and calcareousness. Maximum area of about 327 ha (61%) is currently not suitable (Class N1) for growing cashew and is distributed in the major part of the microwatershed with severe limitations of rooting depth and texture.




Fig. 7.22 Land suitability map of Cashew

7. 23 Land Suitability for Jackfruit (*Artocarpus heterophyllus*)

Jackfruit is one of the most important fruit crop grown in an area of 5368 ha in almost all the districts of the State. The crop requirements for growing jackfruit (Table 7.24) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing jackfruit was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.23.

There are no highly suitable (Class S1) and moderately suitable (Class S2) lands available for growing jackfruit in the microwatershed. An area of about 9 ha (2%) is marginally suitable (Class S3) for growing jackfruit and are distributed in the southern part of the microwatershed. They have moderate limitations of rooting depth and calcareousness. Maximum area of about 327 ha (61%) is currently not suitable (Class N1) for growing jackfruit and is distributed in the major part of the microwatershed with severe limitation of rooting depth.




Fig. 7.23 Land suitability map of Jackfruit

7.24 Land Suitability for Jamun (*Syzygium cumini*)

Jamun is one of the important fruit crop grown in almost all the districts of the State. The crop requirements for growing jamun (Table 25) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing jamun was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.24.

There are no highly suitable (Class S1) and moderately suitable (Class S2) lands available for growing jamun in the microwatershed. An area of about 9 ha (2%) is marginally suitable (Class S3) for growing jamun and are distributed in the southern part of the microwatershed. They have minor limitations of rooting depth and calcareousness. Maximum area of about 327 ha (61%) is currently not suitable (Class N1) for growing jamun and is distributed in the major part of the microwatershed with severe limitation of rooting depth.




Fig. 7.24 Land suitability map of Jamun

7.25 Land Suitability for Custard Apple (*Annona reticulata*)

Custard apple is one of the most important fruit crop grown in almost all the districts of the State. The crop requirements for growing custard apple (Table 7.26) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing custard apple was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.25.

There are no highly suitable (Class S1) lands available for growing custard apple in the microwatershed. An area of about 9 ha (2%) is moderately suitable (Class S2) for growing custard apple and are distributed in the southern part of the microwatershed. They have minor limitations of rooting depth and calcareousness. Maximum area of about 327 ha (61%) is marginally suitable (Class S3) for growing custard apple and is distributed in the major part of the microwatershed with moderate limitations of rooting depth and texture.




Fig. 7.25 Land suitability map of Custard Apple

7.26 Land Suitability for Tamarind (*Tamarindus indica*)

Tamarind is one of the most important spice crop grown in almost all the districts of the state. The crop requirements for growing tamarind (Table 7.27) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing tamarind was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Fig. 7.26.

Entire area of the microwatershed is currently not suitable (Class N1) for tamarind with severe limitations of rooting depth and calcareousness.




Fig. 7.26 Land suitability map of Tamarind

7.27 Land Suitability for Mulberry (*Morus nigra*)

Mulberry is one of the important crop grown for rearing silkworms in about 1.6 lakh ha area in all the districts of the state. The crop requirements for growing mulberry (Table 7.28) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing mulberry was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.27.

There are no highly suitable (Class S1) and moderately suitable (Class S2) lands available for growing mulberry in the microwatershed. An area of about 9 ha (2%) is marginally suitable (Class S3) for growing mulberry and are distributed in the southern part of the microwatershed. They have moderate limitations of rooting depth and calcareousness. Maximum area of about 327 ha (61%) is currently not suitable (Class N1) for growing mulberry and is distributed in the major part of the microwatershed with severe limitation of rooting depth.




Fig 7.27 Land suitability map of Mulberry

7.28 Land suitability for Marigold (*Tagetes sps.*)

Marigold is one of the most important flower crop grown in an area of 9108 ha in almost all the districts of the State. The crop requirements (Table 7.29) for growing marigold were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing marigold was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.28.

There are no highly suitable (Class S1) lands available for growing marigold in the microwatershed. An area of about 9 ha (2%) is moderately suitable (Class S2) for growing marigold and are distributed in the southern part of the microwatershed. They have minor limitations of rooting depth and calcareousness. Maximum area of about 327 ha (61%) is marginally suitable (Class S3) for growing marigold and is distributed in the major part of the microwatershed with moderate limitation of rooting depth.




Fig. 7.28 Land suitability map of Marigold

7.29 Land Suitability for Chrysanthemum (*Dendranthema grandiflora*)

Chrysanthemum is one of the most important flower crop grown in an area of 4978 ha in almost all the districts of the State. The crop requirements (Table 7.30) for growing chrysanthemum were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing chrysanthemum was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.29.

There are no highly suitable (Class S1) lands available for growing chrysanthemum in the microwatershed. An area of about 9 ha (2%) is moderately suitable (Class S2) for growing chrysanthemum and are distributed in the southern part of the microwatershed. They have minor limitations of rooting depth and calcareousness. Maximum area of about 327 ha (61%) is marginally suitable (Class S3) for growing chrysanthemum and is distributed in the major part of the microwatershed with moderate limitations of rooting depth.




Fig. 7.29 Land suitability map of Chrysanthemum

Table 7.1 Soil-Site Characteristics of Warakanahalli Microwatershed

Soil Map Units	Climate (P) (mm)	Growing period (Days)	Drainage Class	Soil depth (cm)	Soil texture		Gravelliness		AWC (mm/m)	Slope (%)	Erosion	pH	EC (dSm ⁻¹)	ESP (%)	CEC [Cmol (p ⁺)kg ⁻¹]	BS (%)
					Sur-face	Sub-surface	Surface (%)	Sub-surface (%)								
DPLcB2	866	150	W	50-75	sc	sc	<15	<15	51-100	1-3	Moderate	6.92	0.122	0.09	7.10	92
HTKbB2g1	866	150	W	25-50	sl	sl	15-35	10-25	<50	1-3	Moderate	6.81	0.062	0.38	3.0	100

*Symbols and abbreviations are according to Field Guide for LRI under Sujala-III Project, Karnataka

Table 7.2 Land suitability criteria for Sorghum

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	26–30	30–34; 24–26	34–40; 20–24	>40; <20
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, c (red), c (black)	scl, cl	ls, sl	-
	pH	1:2.5	5.5-7.8	5.0-5.5 7.8-9.0	>9.0	-
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	10-15
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

Table 7.3 Land suitability criteria for Maize

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	30-34	35-38 26-30	38-40 26-20	
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc	c (red), c (black)	ls, sl	-
	pH	1:2.5	5.5-7.8	5.0-5.5 7.8-9.0	>9.0	-
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	-
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

Table 7.4 Land suitability criteria for Bajra

Land use requirement		Rating				
Soil –site characteristics	Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)	
Climatic regime	Mean temperature in growing season	°C	28-32	33-38 24-27	39-40 20-23	<20
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm	500-750	400-500	200-400	<200
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl, scl, cl,sc,c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.8	5.0-5.5 7.8-9.0	5.5-6.0 >9.0	
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	15-35	35-60	>60	
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	1-3	3-5	5-10	>10

Table 7.5 Land suitability criteria for Groundnut

Land use requirement			Rating			
Soil –site characteristics	Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)	
Climatic regime	Mean temperature in growing season	°C	24–33	22–24; 33–35	20–22; 35–40	<20; >40
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. Well drained	Poorly drained	Very Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl	sl,cl, sc	c (red), c (black), ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<35	35-60	>60	
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.6 Land suitability criteria for Sunflower

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	24–30	30–34; 20–24	34–38; 16–20	>38; <16
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	mod. Well drained	-	Poorly to very drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	cl, sc,c (red), c (black)	scl	ls, sl	-
	pH	1:2.5	6.5-7.8	7.8-8.4 5.5-6.5	8.4-9.0; 5.0-5.5	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.7 Land suitability criteria for Redgram

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	30-35(G) 20-25(AV) 15-18 (F&PS) 35-40(M)	25-30(G) 20-25 (AV) 12-15 (F&PS) 30-35(M)	20-25(G) 15-20(AV) 10-12 (F&PS) 25-30(M)	< 20 <15 <10 <25
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. Well drained	Poorly drained	Very Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, c (red)	c (black),sl, scl, cl	ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-9.0	5.0-5.5 >9.0	-
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-50	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<1.0	1.0-2.0	>2.0	
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.8 Land suitability criteria for Bengal gram

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	20–25	25–30; 15–20	30–35; 10–15	>35; <10
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. Well drained	Poorly drained	Very Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	c(black)	-	c (red), scl, cl, sc	ls, sl
	pH	1:2.5	6.0-7.8	5.0-6.0 7.8-9.0	>9.0	-
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	-
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.9 Land suitability criteria for Cotton

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	22-32	>32	<19	-
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well to moderately well	Poorly drained/Some what excessively drained	-	very poorly/excessively drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, c (red,black)	cl	scl	ls, sl
	pH	1:2.5	6.5-7.8	7.8-8.4	5.5-6.5 8.4->9.0	<5.5
	CEC	C mol (p+)Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	50-100	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	<3	3-5	-	>5

Table 7.10 Land suitability criteria for Chilli

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	25-32	33-35 20-25	35-38 <20	>38
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc	c (black), sl	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.11 Land suitability criteria for Tomato

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	25-28	29-32 20-24	15-19 33-36	<15 >36
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl, scl, cl, sc, c (red)	-	ls, c(black)	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.12 Land suitability criteria for Brinjal

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	Well drained	Moderately well drained	Poorly drained	V. Poorly drained
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class				
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl, scl, cl, sc c (red)	-	ls, c (black)	-
	pH	1:2.5	6.0-7.3	7.3-8.4 5.0-6.0	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	>60
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.13 Land suitability criteria for Onion

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	20-30	30-35	35-40	>40
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately /imperfectly	-	Poorly to V poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl,scl,cl,sc,c (red)	-	c (Black),ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<1.0	1.0-2.0	2.0-4.0	<4
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.14 Land suitability criteria for Bhendi

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	25-28	29-32 20-24	15-19 33-36	<15 >36
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Imperfectly drained	Poorly to very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl,sc, c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.15 Land suitability criteria for Drumstick

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, scl, cl, c (red)	sl, c (black)	ls	s
	pH	1:2.5	6.0-7.3	5.0-5.5 7.3-7.8	5.5-6.0 7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<35	35-60	60-80	>80
Soil toxicity	Salinity (EC saturation extract)	ds/m				
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-10	-	>10

Table 7.16 Land suitability criteria for Mango

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-32	24-27 33-35	36-40	20-24
	Min temp. before flowering	°C	10-15	15-22	>22	-
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration	Days				
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V. Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	-	ls, sl, c (black)	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>150	100-150	75-100	<75
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.17 Land suitability criteria for Guava

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-32	33-36 24-27	37-42 20-23	
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	sl	c (black), ls	-
	pH	1:2.5	6.0-7.8	5.0-6.0	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.18 Land suitability criteria for Sapota

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-32	33-36 24-27	37-42 20-23	>42 <18
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	-	Poorly to very drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	sl	ls, c (black)	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.19 Land suitability criteria for Pomegranate

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	30-34	35-38 25-29	39-40 15-24	
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl,cl, sc, c (red)	c (black),sl	ls	-
	pH	1:2.5	5.5-7.8	7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.20 Land suitability criteria for Musambi

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-30	31-35 24-27	36-40 20-23	>40 <20
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately drained	poorly	Very poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c	sl	ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.21 Land suitability criteria for Lime

Land use requirement		Rating				
Soil –site characteristics	Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)	
Climatic regime	Mean temperature in growing season	°C	28-30	31-35 24-27	36-40 20-23	>40 <20
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately drained	poorly	Very poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c	sl	ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.22 Land suitability criteria for Amla

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. well drained	Poorly drained	V. Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	c (black)	ls, sl	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15-35	35-60	60-80	-
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

Table 7.23 Land suitability criteria for Cashew

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	32 to 34	28 to 32; 34 to 38	24 to 28; 38 to 40	<20; >40
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	-	sl, ls	c (black)
	pH	1:2.5	5.5-6.5	5.0-5.5 6.5-7.3	7.3-7.8	>7.8
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-10	>10	-

Table 7.24 Land suitability criteria for Jackfruit

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. well	Poorly	V. Poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	-	sl, ls, c (black)	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	>60
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10-

Table 7.25 Land suitability criteria for Jamun

Land use requirement		Rating				
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well	Mod. well	Poorly	V.Poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c(red)	sl, c (black)	ls	-
	pH	1:2.5	6.0-7.8	5.0-6.0	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>150	100-150	50-100	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	>60
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

Table 7.26 Land suitability criteria for Custard apple

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	Scl, cl, sc, c (red), c (black)	-	Sl, ls	-
	pH	1:2.5	6.0-7.3	5.5-6.0 7.3-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15-35	35-60	60-80	-
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	>5	-

Table 7.27 Land suitability criteria for Tamarind

Land use requirement		Rating				
Soil –site characteristics	Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)	
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod.well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl,sc, c (red)	sl, c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>150	100-150	75-100	<75
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

Table 7.28 Land suitability criteria for Mulberry

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	24–28	22–24; 28–32	32–38; 22–18	>38; <18
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V. Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, cl, scl	c (red)	c (black), sl, ls	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.8-8.4	7.3-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	0-35	35-60	60-80	>80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

Table 7.29 Land suitability criteria for Marigold

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	18-23	17-15 24-35	35-40 10-14	>40 <10
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl,scl, cl, sc, c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%				
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.30 Land suitability criteria for Chrysanthemum

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	18-23	17-15 24-35	35-40 10-14	>40 <10
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl,scl, cl, sc, c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%				
Erosion hazard	Slope	%	<3	3-5	5-10	>10

7.29 Land Management Units (LMUs)

The 2 soil map units identified in Yadgir Rf microwatershed have been grouped into 2 Land Management Units (LMU's) for the purpose of preparing a Proposed Crop Plan. Land Management Units are grouped based on the similarities in respect of the type of soil, the depth of the soil, the surface soil texture, gravel content, AWC, slope, erosion etc. and a Land Management Units map (Fig. 7.30) has been generated. These Land Management Units are expected to behave similarly for a given level of management.

The 2 map units that have been grouped into 2 Land Management Units along with brief description of soil and site characteristics are given below.

LMU	Soil map units	Soil and site characteristics
1	25.DPLcB2	Moderately shallow (50-75 cm), 1-3 % slopes, non-gravelly (<15 %), moderate erosion
2	161.HTKbB2g1	Shallow (25-50cm), 1-3 % slopes, gravelly (15-35%), moderate erosion




Fig. 7.30 Land Management Units Map-Yadgir Rf microwatershed

7.30 Proposed crop plan for Yadgir Rf microwatershed

After assessing the land suitability for the 28 crops, the Proposed Crop Plan has been prepared for the 2 identified LMUs by considering only the highly (Class S1) and moderately (Class S2) suitable lands for each of the 28 crops. The resultant proposed crop plan is presented below in Table 7.31.

Table 7.31 Proposed crop plan for Yadgir Rf microwatershed

LMU	Mapping Units	Survey Number	Soil Characteristics	Field Crops	Horticulture Crops	Suitable Interventions
LMU 1	25.DPLcB2 (Moderately shallow, red, sandy clay soils)	Ramasamudra : 121	Moderately shallow (50-75 cm), 1-3 % slopes, non-gravelly (<15 %), moderate erosion	Sorghum, Maize, Bajra, Red gram , Finger millet	Fruit crops: Amla, Custard apple Vegetables: Tomato, Chilli Flowers: Marigold Chrysanthemum	Drip irrigation, mulching, suitable soil and water conservation practices (Crescent bunding with catch pit etc)
LMU 2	161.HTKbB2g1 (Shallow, black, sandy loam soils)	Arakera K :72,74,75/1,75/2,76	Shallow (25-50cm), 1-3 % slopes, gravelly (15-35%), moderate erosion	-	Agri-Silvi-Pasture: Hybrid Napier, <i>Styloxanthes hamata</i> , <i>Styloxanthes scabra</i>	Use of short duration varieties, sowing across the slope, drip irrigation and mulching is recommended

SOIL HEALTH MANAGEMENT

8.1 Soil Health

Soil health is basic to plant health and plant health is basic to human and bovine health. Soil is fundamental to crop production. Without soil, no food could be produced nor would livestock be fed on a large scale. Because it is finite and fragile, soil is a precious resource that requires special care from its users.

Soil health or the capacity of the soil to function is critical to human survival. Soil health has been defined as: “the capacity of the soil to function as a living system without adverse effect on the ecosystem”. Healthy soils maintain a diverse community of soil organisms that help to form beneficial symbiotic associations with plant roots, recycle essential plant nutrients, improve soil structure with positive repercussions for soil, water and nutrient holding capacity and ultimately improve crop production and also contribute to mitigating climate change by maintaining or increasing its carbon content.

Functional interactions of soil biota with organic and inorganic components, air and water determine a soil’s potential to store and release nutrients and water to plants and to promote and sustain plant growth. Thus, maintaining soil health is vital to crop production and conserve soil resource base for sustaining agriculture.

The most important characteristics of a healthy soil are

- Good soil tilth
- Sufficient soil depth
- Good water storage and good drainage
- Adequate supply, but not excess of nutrients
- Large population of beneficial organisms
- Small proportion of plant pathogens and insect pests
- Low weed pressure
- Free of chemicals and toxins that may harm the crop
- Resistance to degradation
- Resilience when unfavorable conditions occur

Characteristics of Yadgir Rf microwatershed

- ❖ The soil phases identified in the microwatershed belonged to the soil series of HTK 327 ha (61%) and DPL 9 ha (2%).
- ❖ As per land capability classification entire area of the microwatershed falls under arable land category (Class II & III). The major limitations identified in the arable lands were soil and erosion.
- ❖ On the basis of soil reaction, about 0.13 ha (<1%) is slightly acid (pH 6.0-6.5), 333 ha (62%) neutral (pH 6.5-7.3) and 3 ha (<1%) is slightly alkaline (pH 7.3-7.8). Thus, major area of 333 ha is neutral and 3 ha is under neutral.

❖ Soil Health Management

The following actions are required to improve the current land husbandry practices that provide a sound basis for the successful adoption of sustainable crop production system.

Acid soils

Very minor area of <1 ha is under slightly slightly acid.

1. Growing of crops suitable for particular soil pH.
2. Ameliorating the soils through the application of amendments (liming materials).

Liming materials:

1. CaCO_3 (Calcium Carbonate).
2. Dolomite [$\text{Ca Mg} (\text{CO}_3)_2$]
3. Quick lime (Cao)
4. Slaked lime [$\text{Ca} (\text{OH})_2$]

For normal pH and pH 4.8 (35 t/ha) and pH 6.0-7.0 (4 t/ha) lime is required.

Alkaline soils

Slightly alkaline soils occur in 3 ha area.

1. Regular addition of organic manure, green manuring, green leaf manuring, crop residue incorporation and mulching needs to be taken up to improve the soil organic matter status.
2. Application of biofertilizers (Azospirillum, Azotobacter, Rhizobium).
3. Application of 25% extra N and P (125 % RDN&P).
4. Application of ZnSO_4 – 12.5 kg/ha (once in three years).
5. Application of Boron – 5kg/ha (once in three years).

Neutral soils

Neutral soils cover an area about 333 ha in the microwatershed.

1. Regular addition of organic manure, green manuring, green leaf manuring, crop residue incorporation and mulching needs to be taken up to improve the soil organic matter status.
2. Application of biofertilizers, (Azospirillum, Azotobacter, Rhizobium).
3. Application of 100 per cent RDF.
4. Need based micronutrient applications.

Besides the above recommendations, the best transfer of technology options are also to be adopted.

Soil Degradation

Soil erosion is one of the major factor affecting the soil health in the microwatershed. Out of total 534 ha area in the microwatershed, an area of about 336 ha is suffering from moderate erosion. These areas need immediate soil and water

conservation and other land development and land husbandry practices for restoring soil health.

Dissemination of Information and Communication of Benefits

Any large scale implementation of soil health management requires that supporting information is made available widely, particularly through channels familiar to farmers and extension workers. Given the very high priority attached to soil-health especially by the Central Government on issuing Soil-Health Cards to all the farmers, media outlets like Regional, State and National Newspapers, Radio and Dooradarshan programs in local languages but also modern information and communication technologies such as Cellular phones and the Internet, which can be much more effective in reaching the younger farmers.

Inputs for Net Planning (Saturation Plan) and Interventions needed

Net planning (Saturation Plan) in IWMP is focusing on preparation of

1. Soil and Water Conservation Plan for each plot or farm.
2. Productivity enhancement measures/ interventions for existing crops/livestock/other farm enterprises.
3. Diversification of farming mainly with perennial horticultural crops and livestock.
4. Improving livelihood opportunities and income generating activities.

In this connection, how various outputs of Sujala-III are of use in addressing these objectives of Net Planning (Saturation Plan) are briefly presented below.

- ❖ **Soil Depth:** The depth of a soil decides the amount of moisture and nutrients it can hold, what crops can be taken up or not, depending on the rooting depth and the length of growing period available for raising any crop. Deeper the soil, better for a wide variety of crops. If sufficient depth is not available for growing deep rooted crops, either choose medium or short duration crops or deeper planting pits need to be opened and additional good quality soil brought from outside has to be filled into the planting pits.
- ❖ **Surface Soil Texture:** Lighter soil texture in the top soil means, better rain water infiltration, less run-off and soil moisture conservation, less capillary rise and less evaporation losses. Lighter surface textured soils are amenable to good soil tilth and are highly suitable for crops like groundnut, root vegetables (carrot, raddish, potato etc) but not ideal for crops that need stagnant water like lowland paddy. Heavy textured soils are poor in water infiltration and percolation. They are prone for sheet erosion; such soils can be improved by sand mulching. The technology that is developed by the AICRP-Dry land Agriculture, Vijayapura, Karnataka can be adopted.


- ❖ **Gravelliness:** More gravel content is favorable for run-off harvesting but poor in soil moisture storage and nutrient availability. It is a significant parameter that decides the kind of crop to be raised.
- ❖ **Land Capability Classification:** The land capability map shows the areas suitable and not suitable for agriculture and the major constraints in each of the plot/survey number. Hence, one can decide what kind of enterprise is possible in each of these units. In general, erosion and soil are the major constraints in Yadgir Rf microwatershed.
- ❖ **Organic Carbon:** The OC content (an index of available Nitrogen) is high in (>0.75%) in about 196 ha (37%) and medium (0.5-0.75%) in 140 ha (26%) area. The areas that are medium in OC needs to be further improved by applying farmyard manure and rotating crops with cereals and legumes or mixed cropping.
- ❖ **Promoting Green Manuring:** Growing of green manuring crops costs Rs. 1250/ha (green manuring seeds) and about Rs. 2000/ha towards cultivation that totals to Rs. 3250/- per ha. On the other hand, application of organic manure @ 10 tons/ha costs Rs. 5000/ha. The practice needs to be continued for 2-3 years or more. Nitrogen fertilizer needs to be supplemented by 25% in addition to the recommended level in 140 ha area where OC is medium (<0.5-0.75%). For example, for rainfed maize, recommended level is 50 kg N per ha and an additional 12 kg /ha needs to be applied for all the crops grown in these plots.
- ❖ **Available Phosphorus:** Available phosphorus is medium (23-57 kg/ha) in an area of 250 ha (47%) of the microwatershed and low (<23 kg/ha) in 86 ha (16%). In medium and low areas, for all the crops 25% additional P needs to be applied.
- ❖ **Available Potassium:** Available potassium is medium (145-337 kg/ha) in the entire area of the microwatershed. In medium areas, for all the crops 25% additional potassium needs to be applied.
- ❖ **Available Sulphur:** Available sulphur is a very critical nutrient for oilseed crops. It is low (<10 ppm) in the entire area of the microwatershed. Low areas need to be applied with magnesium sulphate or gypsum or Factamphos (p) fertilizer (13% sulphur) for 2-3 years for the deficiency to be corrected.
- ❖ **Available Boron:** Available boron is low (<0.5 ppm) in the entire area of the microwatershed. For these areas, application of sodium borate @ 10 kg/ha as soil application or 0.2 % borax as foliar spray is recommended.
- ❖ **Available Iron:** Available iron is sufficient (>4.5 ppm) in an area of about 252 ha (47 %) and deficient (<4.5 ppm) in an area of 84 ha (16%). For the deficient areas, iron sulphate @ 25 kg/ha need to be applied for 2-3 years.
- ❖ **Available Manganese:** All the soils in the microwatershed are sufficient (>1.0 ppm) in available manganese.
- ❖ **Available Copper:** All the soils in the microwatershed are sufficient (>0.2 ppm) in available copper.

- ❖ **Available Zinc:** Maximum area of about 330 ha (62%) is deficient (<0.6 ppm) and 7 ha (1%) is sufficient in available zinc content. Application of zinc sulphate @25 kg/ha is recommended for deficient areas.
- ❖ **Soil Alkalinity:** An area of 3 ha (<1%) in the microwatershed has soils that are slightly alkaline. These areas need application of gypsum and wherever calcium is in excess, iron pyrites and element sulphur can be recommended. Management practices like treating repeatedly with good quality water to drain out the excess salts and provision of subsurface drainage and growing of salt tolerant crops like Casuarina, Acacia, Neem, Ber etc, are recommended.
- ❖ **Land Suitability for Various Crops:** Areas that are highly, moderately and marginally suitable for growing various crops are indicated. Along with the suitability, various constraints that are limiting the productivity are also indicated. For example, in case of cotton, rooting depth, texture and calcareousness are the major constraints in various plots. With suitable management interventions, the productivity can be enhanced. In order to increase the water holding capacity of light textured soils, growing of green manure crops and application of organic manure is recommended.

SOIL AND WATER CONSERVATION TREATMENT PLAN

For preparing soil and water conservation treatment plan for Yadgir Rf microwatershed, the land resource inventory database generated under Sujala-III project has been transformed as information through series of interpretative (thematic) maps using soil phase map as a base. The various thematic maps (1:7920 scale) generated were

- Soil depth
- Surface soil texture
- Available water capacity
- Soil slope
- Soil gravelliness
- Land capability
- Present land use and land cover
- Crop suitability
- Rainfall
- Hydrology
- Water Resources
- Socio-economic data
- Contour plan with existing features- network of waterways, pottissa boundaries, cut up/ minor terraces etc.
- Cadastral map (1:7920 scale)
- Satellite imagery (1:7920 scale)



Apart from these, Hand Level/ Hydro Marker/ Dumpy Level/ Total Station and Kathedars' List to be collected.

Steps for Survey and Preparation of Treatment Plan

The boundaries of Land User Groups' and Survey No. boundaries are traced in the field.

- Naming of user groups and farmers
- Identification of arable and non arable lands
- Identification of drainage lines and gullies
- Identification of non treatable areas
- Identification of priority areas in the arable lands
- Treatment plan for arable lands
- Location of water harvesting and recharge structures

9.1 Treatment Plan

The treatment plan recommended for arable lands is briefly described below

9.1.1 Arable Land Treatment

A. BUNDING

Steps for Survey and Preparation of Treatment Plan	
<ul style="list-style-type: none"> • Cadastral map (1:7920 scale) is enlarged to a scale of 1:2500 scale • Existing network of waterways, pottissa boundaries, grass belts, natural drainage lines/ watercourse, cut ups/ terraces are marked on the cadastral map to the scale • Drainage lines are demarcated into 	
Small gullies	(up to 5 ha catchment)
Medium gullies	(5-15 ha catchment)
Ravines	(15-25 ha catchment) and
Halla/Nala	(more than 25ha catchment)

USER GROUP-1

CLASSIFICATION OF GULLIES

ಕೊರಕಲಿನ ವರ್ಗೀಕರಣ

UPPER REACH: 15 Ha.

MIDDLE REACH: 15 + 10 = 25 ಹೆ.

LOWER REACH: 25 ಹೆಕ್ಟಾರ್ ನಿಂತ ಅಧಿಕ

POINT OF CONCENTRATION

Measurement of Land Slope

Land slope is estimated or determined by the study and interpretation of contours or by measurement in the field using simple instruments like Hand Level or Hydromarker.



Vertical and Horizontal intervals between bunds as recommended by the Watershed Development Department.

Slope percentage	Vertical interval (m)	Corresponding Horizontal Distance (m)
2 - 3%	0.6	24
3 - 4%	0.9	21
4 - 5%	0.9	21
5 - 6%	1.2	21
6 - 7%	1.2	21

Note: (i) The above intervals are maximum.

(ii) Considering the slope class and erosion status (A1... A=0-1 % slope, 1= slight erosion) the intervals have to be decided.

Bund length recording: Considering the contour plan and the existing grass belts/partitions, the bunds are aligned and lengths are measured.

Section of the Bund

Bund section is decided considering the soil texture class and gravelliness class (bg₀.. b=loamy sand, g₀ = <15% gravel). The recommended Sections for different soils are given below.

Recommended Bund Section

Top width (m)	Base width (m)	Height (m)	Side slope (Z:1;H:V)	Cross section (sq m)	Soil Texture	Remarks
0.3	0.9	0.3	01:01	0.18	Sandy loam	Vegetative bund
0.3	1.2	0.3	1.5:1	0.225	Sandy clay	
0.3	1.2	0.5	0.9:1	0.375	Red gravelly soils	
0.3	1.2	0.6	0.75:1	0.45		
0.3	1.5	0.6	01:01	0.54	Red sandy loam	
0.3	2.1	0.6	1.5:1	0.72	Very shallow black soils	
0.45	2	0.75	01:01	0.92		
0.45	2.4	0.75	1.3:1	1.07	Shallow black soils	
0.6	3.1	0.7	1.78:1	1.29	Medium black soils	
0.5	3	0.85	1.47:1	1.49		

Formation of Trench cum Bund

Dimensions of the Borrow Pits/Trenches to be excavated (machinery are decided considering the Bund Section).

Details of Borrow Pit dimensions are given below:

TRENCH CUM BUND

WATER STORAGE AREA
0.45 Sq.m section
IDEAL FOR HORTICULTURE CROPS

'A' FRAME FOR INTERBUND MANAGEMENT

1. ಸಮಾನಾತಳ ಉಳಿಸುವೆ
2. ಸಮಾನಾತಳ ಬಿತ್ತನೆ/ನಾಟಿ

Size of Borrow Pits/ Trench recommended for Trench cum Bund (by machinery)

Bund section	Bund length	Earth quantity	Pit				Berm (pit to pit)	Soil depth class
			L(m)	W(m)	D(m)	Quantity (m ³)		
m ²	m	m ³					m	
0.375	6	2.25	5.85	0.85	0.45	2.24	0.15	Shallow
0.45	6	2.7	5.4	1.2	0.43	2.79	0.6	Shallow
0.45	6	2.7	5	0.85	0.65	2.76	1	Moderately Shallow
0.54	5.6	3.02	5.5	0.85	0.7	3.27	0.1	Moderately shallow
0.54	5.5	2.97	5	1.2	0.5	3	0.5	Shallow
0.72	6.2	4.46	6	1.2	0.7	5.04	0.2	Moderately shallow
0.72	5.2	3.74	5.1	0.85	0.9	3.9	0.1	Moderately deep

B. Water Ways

1. Existing waterways are marked on the cadastral map (1:7920 scale) and their dimensions are recorded.
2. Considering the contour plan of the MWS, additional waterways/ modernization of the existing ones can be thought of.
3. The design details are given in the Manual.

C. Farm Ponds

Waterways and the catchment area will give an indication on the size of the Farm Pond. Location of the pond can be decided based on the contour plan/ field condition and farmers' need/desire.

D. Diversion Channel

Existing EPT/ CPT are marked on the cadastral map. Looking to the need, these can be modernized or fresh diversion channel can be proposed and runoff from this can be stored in Gokatte/ Recharge ponds.

9.1.2 Non-Arable Land Treatment

Depending on the gravelliness and crops preferred by the farmers, the concerned authorities can decide appropriate treatment plan. The recommended treatments may be Contour Trench, Staggered Trench, Crescent Bund, Boulder Bund or Pebble Bund.

9.1.3 Treatment of Natural Water Course/ Drainage Lines

- a) The cadastral map has to be updated as regards the network of drainage lines (gullies/ *nalas*/ *hallas*) and existing structures are marked to the scale and storage capacity of the existing water bodies are documented.
- b) The drainage line will be demarcated into Upper Reach, Middle Reach and Lower Reach.
- c) Considering the Catchment, *Nala* bed and bank conditions, suitable structures are decided.
- d) Number of storage structures (Check dam/ *Nala* bund/ Percolation tank) will be decided considering the commitments and available runoff from water budgeting and quality of water in the wells and site suitability.
- e) Detailed Leveling Survey using Dumpy Level / Total Station has to be carried out to arrive at the site-specific designs as shown in the Manual.
- f) The location of ground water recharge structures are decided by examining the lineaments and fracture zones from geological maps.
- g) Rainfall intensity data of the nearest Rain Gauge Station is considered for Hydrologic Designs.
- h) Silt load to the Storage/Recharge Structures is reduced by providing vegetative, boulder and earthen checks in the natural water course. Location and design details are given in the Manual.

9.2 Recommended Soil and Water Conservation Measures

The appropriate conservation structures best suited for each of the land parcel/ survey number (Appendix-I) are selected based on the slope per cent, severity of erosion, amount of rainfall, land use and soil type. The different kinds of conservation structures recommended are:

1. Graded / Strengthening of Bunds
2. Trench cum Bunds (TCB)
3. Trench cum Bunds / Strengthening
4. Crescent Bunds

A map (Fig. 9.1) showing soil and water conservation plan with different kinds of structures recommended has been prepared which shows the spatial distribution and extent of area. An area of about 9 ha (2%) needs trench cum bunding and 327 ha (61%) needs graded bunding.

The conservation plan prepared may be presented to all the stakeholders including farmers and after considering their suggestions, the conservation plan for the microwatershed may be finalised in a participatory approach.




Fig. 9.1 Soil and water conservation plan map of Yadgir Rf microwatershed

9.3 Greening of microwatershed

As part of the greening programme in the watersheds, it is envisaged to plant a variety of horticultural and other tree plants that are edible, economical and produce lot of biomass which helps to restore the ecological balance in the watersheds. The lands that are suitable for greening programme are non-arable lands (land capability classes V, VI VII and VIII) and also the lands that are not suitable or marginally suitable and field bunds for growing annual and perennial crops. The method of planting these trees is given below.

It is recommended to open pits during the 1st week of March along the contour and heap the dugout soil on the lower side of the slope in order to harness the flowing water and facilitate weathering of soil in the pit. Exposure of soil in the pit also prevents spread of pests and diseases due to scorching sun rays. The pits should be filled with mixture of soil and organic manure during the second week of April and keep ready with sufficiently tall seedlings produced either in poly bags or in root trainer nurseries so that planting can be done during the 2nd or 3rd week of April depending on the rainfall.

The tree species suitable for the area considering rainfall, temperature and adaptability is listed below; waterlogged areas are recommended to be planted with species like Nerale (*Sizygium cumini*) and Bamboo. Dry areas are to be planted with species like Honge, Bevu, Seetaphal etc.

Dry deciduous species			Temp (°C)	Rainfall (mm)
1.	Bevu	<i>Azadiracta indica</i>	21–32	400 –1,200
2.	Tapasi	<i>Holoptelia integrifolia</i>	20-30	500 - 1000
3.	Seetaphal	<i>Anona Squamosa</i>	20-40	400 - 1000
4.	Honge	<i>Pongamia pinnata</i>	20 -50	500– 2,500
5.	Kamara	<i>Hardwickia binata</i>	25 -35	400 - 1000
6.	Bage	<i>Albezzia lebbek</i>	20 - 45	500 - 1000
7.	Ficus	<i>Ficus bengalensis</i>	20 - 50	500–2,500
8.	Sisso	<i>Dalbargia Sissoo</i>	20 - 50	500 -2000
9.	Ailanthus	<i>Ailanthus excelsa</i>	20 - 50	500 - 1000
10.	Hale	<i>Wrightia tinctoria</i>	25 - 45	500 - 1000
11.	Uded	<i>Steriospermum chelanoides</i>	25 - 45	500 -2000
12.	Dhupa	<i>Boswellia Serrata</i>	20 - 40	500 - 2000
13.	Nelli	<i>Emblica Officinalis</i>	20 - 50	500 -1500
14.	Honne	<i>Pterocarpus marsupium</i>	20 - 40	500 - 2000
Moist Deciduous Species			Temp (°C)	Rainfall (mm)
15.	Teak	<i>Tectona grandis</i>	20 - 50	500-5000
16.	Nandi	<i>Legarstroemia lanceolata</i>	20 - 40	500 - 4000
17.	Honne	<i>Pterocarpus marsupium</i>	20 - 40	500 - 3000
18.	Mathi	<i>Terminalia alata</i>	20 -50	500 - 2000
19.	Shivane	<i>Gmelina arborea</i>	20 -50	500 -2000
20.	Kindal	<i>T.Paniculata</i>	20 - 40	500 - 1500
21.	Beete	<i>Dalbargia latifolia</i>	20 - 40	500 - 1500
22.	Tare	<i>T. belerica</i>	20 - 40	500 - 2000
23.	Bamboo	<i>Bambusa arundinasia</i>	20 - 40	500 - 2500
24.	Bamboo	<i>Dendrocalamus strictus</i>	20 – 40	500 – 2500
25.	Muthuga	<i>Butea monosperma</i>	20 - 40	400 - 1500
26.	Hippe	<i>Madhuca latifolia</i>	20 - 40	500 - 2000
27.	Sandal	<i>Santalum album</i>	20 - 50	400 - 1000
28.	Nelli	<i>Emblica officinalis</i>	20 - 40	500 - 2000
29.	Nerale	<i>Sizyium cumini</i>	20 - 40	500 - 2000
30.	Dhaman	<i>Grevia tilifolia</i>	20 - 40	500 - 2000
31.	Kaval	<i>Careya arborea</i>	20 - 40	500 - 2000
32.	Harada	<i>Terminalia chebula</i>	20 - 40	500 - 2000

References

1. FAO (1976) Framework for Land Evaluation, Food and Agriculture Organization, Rome.72 pp.
2. FAO (1983) Guidelines for Land Evaluation for Rainfed Agriculture, FAO, Rome, 237 pp.
3. IARI (1971) Soil Survey Manual, All India Soil and Land Use Survey Organization, IARI, New Delhi, 121 pp.
4. Katyal, J.C. and Rattan, R.K. (2003) Secondary and Micronutrients; Research Gap and Future Needs. Fert. News 48 (4); 9-20.
5. Naidu, L.G.K., Ramamurthy, V., Challa, O., Hegde, R. and Krishnan, P. (2006) Manual Soil Site Suitability Criteria for Major Crops, NBSS Publ. No. 129, NBSS & LUP, Nagpur, 118 pp.
6. Natarajan, A. and Dipak Sarkar (2010) Field Guide for Soil Survey, National Bureau of Soil Survey and Land Use Planning (ICAR), Nagpur, India.
7. Natarajan, A., Rajendra Hegde, Raj, J.N. and Shivananda Murthy, H.G. (2015) Implementation Manual for Sujala-III Project, Watershed Development Department, Bengaluru, Karnataka.
8. Sarma, V.A.K., Krishnan, P. and Budihal, S.L. (1987) Laboratory Manual, Tech. Bull. 23, NBSS &LUP, Nagpur.
9. Sehgal, J.L. (1990) Soil Resource Mapping of Different States of India; Why and How?, National Bureau of Soil Survey and Land Use Planning, Nagpur, 49 pp.
10. Shivaprasad, C.R., R.S. Reddy, J. Sehgal and M. Velayuthum (1998) Soils of Karnataka for Optimizing Land Use, NBSS Publ. No. 47b, NBSS & LUP, Nagpur, India.
11. Soil Survey Staff (2006) Keys to Soil Taxonomy, Tenth edition, U.S. Department of Agriculture/ NRCS, Washington DC, U.S.A.
12. Soil Survey Staff (2012) Soil Survey Manual, Handbook No. 18, USDA, Washington DC, USA.

Appendix I
Yadgir Rf_(4D5B1G1a) Microwatershed
Soil Phase Information

Village	Survey Number	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Ashinala	139	23.12	RO	RO	RO	RO	RO	RO	RO	RO	RO (Rc)	Not Available	RO	RO
Ashinala	141	5.21	RO	RO	RO	RO	RO	RO	RO	RO	RO (Rc)	Not Available	RO	RO
Arakera .K	72	0.77	HTKbB2g 1	LMU-2	Shallow (25-50 cm)	Loamy sand	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Arakera .K	74	1.76	HTKbB2g 1	LMU-2	Shallow (25-50 cm)	Loamy sand	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IIes	Graded bunding
Arakera .K	75/1	8.19	HTKbB2g 1	LMU-2	Shallow (25-50 cm)	Loamy sand	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Arakera .K	75/2	1.61	HTKbB2g 1	LMU-2	Shallow (25-50 cm)	Loamy sand	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Arakera .K	76	315.03	HTKbB2g 1	LMU-2	Shallow (25-50 cm)	Loamy sand	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Forest (Fo)	Not Available	IIes	Graded bunding
Ramasamudra	118	146.82	RO	RO	RO	RO	RO	RO	RO	RO	Forest (Fo)	Not Available	RO	RO
Ramasamudra	119	11.99	RO	RO	RO	RO	RO	RO	RO	RO	Groundnut+Redgram (Gn+Rg)	Not Available	RO	RO
Ramasamudra	120	4.86	RO	RO	RO	RO	RO	RO	RO	RO	Redgram (Rg)	Not Available	RO	RO
Ramasamudra	121	6.29	DPLcB2	LMU-1	Moderately shallow (50-75 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Trench cum bunding
Ramasamudra	122	5.19	RO	RO	RO	RO	RO	RO	RO	RO	Redgram (Rg)	Not Available	RO	RO
Ramasamudra	123	2.93	RO	RO	RO	RO	RO	RO	RO	RO	Redgram (Rg)	Not Available	RO	RO
Ramasamudra	124	0.32	RO	RO	RO	RO	RO	RO	RO	RO	Greengram+Redgram (Gg+Rg)	Not Available	RO	RO

Appendix II

Yadgir Rf_(4D5B1G1a) Microwatershed

Soil Fertility Information

Village	Survey Number	Soil Reaction	Salinity	Organic Carbon	Available Phosphorus	Available Potassium	Available Sulphur	Available Boron	Available Iron	Available Manganese	Available Copper	Available Zinc
Ashinala	139	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO
Ashinala	141	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO
Arakera .K	72	Neutral (pH 6.5 - 7.3)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Arakera .K	74	Neutral (pH 6.5 - 7.3)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Arakera .K	75/1	Neutral (pH 6.5 - 7.3)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Arakera .K	75/2	Neutral (pH 6.5 - 7.3)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Arakera .K	76	Neutral (pH 6.5 - 7.3)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Ramasamudra	118	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO
Ramasamudra	119	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO
Ramasamudra	120	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO
Ramasamudra	121	Neutral (pH 6.5 - 7.3)	Non saline (<2 dsm)	High (> 0.75 %)	Low (< 23 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Ramasamudra	122	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO
Ramasamudra	123	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO
Ramasamudra	124	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO

Appendix III
Yadgir Rf_(4D5B1G1a) Microwatershed
Soil Suitability Information

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry	
Ashinala	139	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	
Ashinala	141	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO
Arakera .K	72	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Arakera .K	74	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Arakera .K	75/1	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Arakera .K	75/2	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Arakera .K	76	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Ramasamudra	118	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	
Ramasamudra	119	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	
Ramasamudra	120	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	
Ramasamudra	121	N1rz	S2rz	S3rz	S2rz	S3rz	S2rz	N1rz	S3rz	S2rz	S3rz	S3rz	S2rz	S3rz	S2rz	S3rz	S3rz	S2rz	S2r	S2rz	S2rz	S2rz	S2rz	S3rz	S2rz	S2r	S2r	S3rz	S3rz		
Ramasamudra	122	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	
Ramasamudra	123	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	
Ramasamudra	124	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	

PART-B

SOCIO-ECONOMIC STATUS OF FARM HOUSEHOLDS

CONTENTS

1	Findings of the socio-economic survey	1-3
2	Introduction	5
3	Methodology	7-8
4	Salient features of the survey	9-30
5	Summary	31-34

LIST OF TABLES

1	Households sampled for socio economic survey	9
2	Population characteristics	9
3	Age wise classification of household members	9
4	Education level of household members	10
5	Occupation of household heads	10
6	Occupation of family members	11
7	Institutional participation of household members	11
8	Type of house owned by households	11
9	Durable assets owned by households	12
10	Average value of durable assets owned by households	12
11	Farm implements owned by households	12
12	Average value of farm implements	13
13	Livestock possession by households	13
14	Average labour availability	13
15	Adequacy of hired labour	13
16	Migration among the households	14
17	Average distance and duration of migration	14
18	Purpose of migration by members of households	14
19	Distribution of land (ha)	14
20	Average land value (Rs./ha)	14
21	Status of bore wells	15
22	Source of irrigation	15
23	Depth of water (Avg. In meters)	15
24	Irrigated Area (ha)	15
25	Cropping pattern	16
26	Cropping intensity	16
27	Possession of bank account and saving	16
28	Borrowing status	16
29	Source of credit borrowed by households	16
30	Avg. Credit amount	17
31	Purpose of credit borrowed (institutional Source)	17
32	Purpose of credit borrowed (Private Source)	17
33	Repayment status of household (institutional Source)	17

34	Repayment status of household (Private Source)	18
35	Opinion regarding institutional sources of credit	18
36	Opinion regarding Non- institutional sources of credit	18
37.a	Cost of cultivation of Cotton	19
37.b	Cost of cultivation of Redgram	20
37.c	Cost of cultivation of Maize	21
37.d	Cost of Cultivation of Groundnut	22
37.e	Cost of Cultivation of Jowar	23
38	Adequacy of fodder	24
39	Annual gross income	24
40	Average annual expenditure	24
41	Horticultural species grown	24
42	Interest towards cultivation of horticulture crops	25
43	Forest species grown	25
44	Average additional investment capacity	25
45	Source of funds for additional investment	25
46	Marketing of the agricultural produce	26
47	Marketing channels used for sale of agricultural produce	26
48	Mode of transport of agricultural produce	26
49	Incidence of soil and water erosion problems	26
50	Interest shown towards soil testing	27
51	Usage pattern of fuel for domestic use	27
52	Source of drinking water	27
53	Source of light	27
54	Existence of sanitary toilet facility	27
55	Possession of public distribution system (PDS) card	28
56	Participation in NREGA programme	28
57	Adequacy of food items	28
58	Inadequacy of food items	28
59	Response on market surplus of food items	29
60	Farming constraints experienced	29

FINDINGS OF THE SOCIO-ECONOMIC SURVEY

- ❖ *The survey was conducted in Yadgir Rf is located at North latitude 16⁰ 49' 29.862" and 16⁰ 47' 10.021" and East longitude 77⁰ 15' 52.647" and 77⁰ 14' 56.958" covering an area of about 533.85 ha coming under Ramsamudra, Ashinala and Arakera. K Villages of Yadagiri taluk.*
- ❖ *Socio-economic analysis of Yadgir Rf micro watersheds of Ramasamudram sub-watershed, Yadgir taluk & District indicated that, out of the total sample of 34 total respondents, 11 (32.35 %) were marginal, 17 (50.00%) were small, 1 (2.94 %) were semi medium and 5 landless farmers were also interviewed for the survey.*
- ❖ *The population characteristics of households indicated that, there were 93 (55.03%) men and 76 (44.97 %) were women. The average population of landless was 4, marginal farmers were 6.45, small farmers were 4.29 and semi medium farmers were 5.0.*
- ❖ *Majority of the respondents (38.46%) were in the age group of 16-35 years.*
- ❖ *Education level of the sample households indicated that, there were 44.97 per cent illiterates, 50.89 per cent pre university education and 8.28 per cent attained graduation.*
- ❖ *About, 52.94 per cent of household heads practicing agriculture and 29.41 per cent of the household heads were engaged as agricultural labourers.*
- ❖ *Agriculture was the major occupation for 33.73 per cent of the household members.*
- ❖ *In the study area, 61.76 per cent of the households possess katcha house and 38.24 per cent possess pucca house.*
- ❖ *The durable assets owned by the households showed that, 76.47 per cent possess TV, 64.71 per cent possess mixer grinder, 85.29 per cent possess mobile phones and 44.12 per cent possess motor cycles.*
- ❖ *Farm implements owned by the households indicated that, 20.59 per cent of the households possess plough, 2.94 per cent possess tractor, 2.94 per cent possess bullock cart and 11.76 per cent possess sprayer.*
- ❖ *Regarding livestock possession by the households, 2.94 per cent possess local cow.*
- ❖ *The average labour availability in the study area showed that, own labour men available in the micro watershed was 1.64, women available in the micro watershed was 1.36, hired labour (men) available was 8.86 and hired labour (women) available was 13.96.*
- ❖ *In the study area, about 1.18 per cent of the respondents migrated from the micro watershed in search of jobs with an average distance of 525.00 kms for about 9.00 months.*

- ❖ *Out of the total land holding of the sample respondents 100.00 per cent (33.95 ha) of the area is under dry condition.*
- ❖ *There were 1.00 live bore wells and 1.00 dry bore wells among the sampled households*
- ❖ *Bore well was the major source of irrigation for 2.94 per cent of the households.*
- ❖ *The major crops grown by sample farmers are Cotton, Redgram, Maize, Groundnut and Jowar and cropping intensity was recorded as 100.00 per cent.*
- ❖ *Out of the sample households 94.12 percent possessed bank account and 61.76 per cent of them have savings in the account.*
- ❖ *About 64.71 per cent of the respondents borrowed credit from various sources.*
- ❖ *Among the credit borrowed by households, 18.18 per cent have borrowed loan from commercial banks and 59.09 per cent from co-operative/Grameena bank.*
- ❖ *Majority of the respondents (100.00%) have borrowed loan for agriculture purpose.*
- ❖ *Regarding the opinion on institutional sources of credit, 100.00 per cent of the households opined that credit helped to perform timely agricultural operations.*
- ❖ *The per hectare cost of cultivation for Cotton, Redgram, Maize, Groundnut and Jowar was Rs.34406.70, 54597.19, 15086.15, 70279.00 and 19949.72 with benefit cost ratio of 1:1.40, 1: 0.70, 1: 2.50, 1: 0.60 and 1:1.00 respectively.*
- ❖ *Further, 11.76 per cent of the households opined that dry fodder was adequate.*
- ❖ *The average annual gross income of the farmers was Rs. 73020.59 in micro-watershed, of which Rs. 39491.18 comes from agriculture*
- ❖ *Sampled households have grown 10 horticulture trees and 56 forestry trees together in the fields and back yards.*
- ❖ *About 32.35 per cent of the households shown interest to cultivate horticultural crops.*
- ❖ *Households have an average investment capacity of Rs. 735.29 for land development.*
- ❖ *Source of funds for additional investment is concerned, 32.35 per cent depends on bank loan for land development activities.*
- ❖ *Regarding marketing channels, 85.29 per cent of the households have sold agricultural produce to the local/village merchants.*
- ❖ *Further, 32.35 per cent of the households have used tractor for the transport of agriculture commodity.*
- ❖ *Majority of the farmers (35.29%) have experienced soil and water erosion problems in the watershed and 82.35 per cent of the households were interested towards soil testing.*
- ❖ *Fire was the major source of fuel for domestic use for 55.88 per cent of the households and 44.12 per cent households has LPG connection.*

- ❖ *Piped supply was the major source for drinking water for 94.12 per cent of the households.*
- ❖ *Electricity was the major source of light for 100.00 per cent of the households.*
- ❖ *In the study area, 29.41 per cent of the households possess toilet facility.*
- ❖ *Regarding possession of PDS card, 97.06 per cent of the households possessed BPL card, 2.94 per cent of the household's possessed APL card.*
- ❖ *Households opined that, the requirement of cereals (58.82%), pulses (55.88%) and oilseeds (5.88%) are adequate for consumption.*
- ❖ *Farming constraints experienced by households in the micro watersheds were lower fertility status of the soil (94.12%) wild animal menace on farm field (85.29%), frequent incidence of pest and diseases (50.00%), inadequacy of irrigation water (85.29%), high cost of fertilizers and plant protection chemicals (85.29%), high rate of interest on credit (85.29%), low price for the agricultural commodities (85.29%), lack of marketing facilities in the area (76.47%), inadequate extension services (61.76%), lack of transport for safe transport of the agricultural produce to the market (85.29%), Less rainfall (5.88%) and Source of Agri-technology information (Newspaper/ TV/Mobile) (2.94%).*

INTRODUCTION

Soil and water are the two precious natural resources which are essential for crop production and existence of life on earth. Rainfed agriculture is under severe stress due to various constraints related to agriculture like uneven and erratic distribution of rainfall, indiscriminate use of fertilizers, chemicals and pesticides, adoption of improper land management practices, soil erosion, decline in soil fertility, decline in ground water resources leading to low crop productivity. The area under rainfed agriculture has to be managed effectively using the best available practices to enhance the production of food, fodder and fuel. This is possible if the land resources are characterized at each parcel of land through detailed land resource inventory using the best available techniques of remote sensing, GPS and GIS. The watershed development programs are aimed at the sustainable distribution of its resources and the process of creating and implementing plans, programs, and projects to sustain and enhance watershed functions that affect the plant, animal and human communities within a watershed boundary.

World Bank funded KWDP II, SUJALA III project was implemented in with Broad objective of demonstrating more effective watershed management through greater integration of programmes related to rain-fed agriculture, innovative and science based approaches and strengthen institutional capacities and If successful, it is expected that the systems and tools could be mainstreamed into the overall IWMP in the State of Karnataka and in time, throughout other IWMP operations in India. With this background the socio-economic survey has been carried out with following specific objectives:

1. To understand the demographic features of the households in the micro-watershed
2. To understand the extent of family labour available and additional employment opportunities available within the village.
3. To know the status of assets of households in the micro-watershed for suggesting possible improvements.
4. To study the cropping pattern, cropped area and productivity levels of different households in micro-watershed.
5. To determine the type and extent of livestock owned by different categories of HHs
6. Availability of fodder and level of livestock management.

Scope and importance of survey

Survey helps in identification of different socio-economic and resource use-patterns of farmers at the Micro watershed. Household survey provides demographic features, labour force, and levels of education; land ownership and asset position (including livestock and other household assets) of surveyed households; and cropping patterns, input intensities, and average crop yields from farmers' fields. It also discusses crop utilization and the degree of commercialization of production in the areas; farmers' access to and utilization of credit from formal and informal sources; and the level of adoption and use of soil, water, and pest management technologies.

METHODOLOGY

The description of the methods, components selected for the survey and procedures followed in conducting the baseline survey are furnished under the following heads.

1. Description of the study area

Yadgir District is one of the 30 districts of Karnataka state in southern India. This district was carved out from the erstwhile Gulbarga district as the 30th district of Karnataka on 10 April 2010. Yadgir town is the administrative headquarters of the district. The district comprises of 3 taluks namely, Shahapur, Yadgiri and Shorapur (There are 16 hoblies, 117 Gram Panchayats, 4 Municipalities, 8 Towns/ Urban agglomeration and 487 inhabited & 32 un-inhabited villages The district occupies an area of 5,160.88 km².

Yadgir district is the second smallest district in the state, area wise is very rich in cultural traditions. The vast stretch of fertile black soil of the district is known for bumper red gram and jowar crops. The district is a "Daal bowl" of the state. The district is also known for cluster of cement industries and a distinct stone popularly known as "Malakheda Stone". Two main rivers, Krishna and Bhima, and a few tributaries flow in this region. Krishna and Bhima Rivers drain the district. They constitute the two major river basins of the district. Kagna and Amarja are the two sub - basins of Bhima River, which occur within the geographical area of the district

According to the 2011 census Yadgir district has a population of 1, 172,985, roughly equal to the nation of Timor-Leste or the US state of Rhode Island. This gives it a ranking of 404th in India (out of a total of 640). The district has a population density of 224 inhabitants per square kilometre (580/sq mi). Its population growth rate over the decade 2001-2011 was 22.67%. Yadgir has a sex ratio of 984 females for every 1000 males, and a literacy rate of 52.36%.

2. Locale of the survey and description of the micro-watershed and

The study was conducted in Yadgir Rf micro-watershed (Ramasamudram sub-watershed, Yadgir taluk & District) is located at North latitude 16^o 49' 29.862" and 16^o 47' 10.021" and East longitude 77^o 15' 52.647" and 77^o 14' 56.958" covering an area of about 533.85 ha bounded by under Ramsamudra, Ashinala and Arakera. K Villages.

3. Selection of the respondents for the study

The micro-watershed is marked with 320 square meters grids. One farmer from every alternate grid in the micro-watershed was selected for the study and interviewed for socio-economic data. Totally 34 households were interviewed for the survey.

4. The parameters considered for socio-economic survey of households

Two forms of data were collected from the micro-watershed which includes primary data from the farm households and secondary data about the villages under the micro-watershed jurisdiction.

The following parameters were considered for the primary data collection about the socio-economic data of the households, (1) Demographic information, (2) Farm and durable assets owned by households, (3) Livestock possession, (4) Labour availability, (5) Level of migration in the village, Land holding, (7) Cropping pattern, (8) Source of irrigation, (9) Borrowing status, (10) Cost of cultivation of major crops, (11) Economics of subsidiary activities, (12) Fodder availability, (13) Family annual income from different sources, (14) Horticulture and forestry species grown, (15) Additional investment capacity, (16) Marketing practices, (17) Status of soil and water conservation structure, (18) Access to basic needs and (19) Constraints and suggestion.

The following parameters were considered for the secondary data regarding the villages under the micro-watershed jurisdiction, (1) Number of villages in each micro-watershed jurisdiction, (2) Village wise number of households, (3) Geographical area of the villages, (4) Cultivable area including rainfed and irrigated, (5) Number and type of house in each village, (6) Human and livestock population, (7) Facilities in the village such as roads, transport facility for conveyance, drinking water supply, street light and (8) Community based organizations in the villages.

5. Development of interview schedule and data collection

Taking into the consideration the objectives of the survey, an interview schedule was prepared after thorough consultation with the experts in the field of social sciences. A comprehensive interview schedule covering all the major parameters for measuring the socio-economic situation was developed.

6. Tools used to analyze the data

The statistical components such as frequency and percentage were used to analyze the data.

Abbreviations used in the report

LL=Landless

MF=Marginal Farmers

SF=Small farmers

SMF=Semi medium farmers

MDF=Medium farmers

LF=Large Farmers

FINDINGS OF THE SURVEY

This chapter deals with systematic presentation of results of the survey. Keeping in view the objectives, the salient features of the survey are presented under the following headings.

Households sampled for socio-economic survey: The data on households sampled for socio economic survey in Yadgir Rf Micro watershed is presented in Table 1 and it indicated that 34 farmers were sampled in Yadgir Rf micro-watershed among households surveyed 11 (32.35%) were marginal, 17 (50.00%) were small, 1(2.94 %) were semi medium and 5 landless farmers were also interviewed for the survey.

Table 1. Households sampled for socio economic survey in Yadgir Rf micro-watershed

Sl.No.	Particulars	LL (5)		MF (11)		SF (17)		SMF (1)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Farmers	5	14.7	11	32.4	17	50	1	2.94	34	100

Population characteristics: The population characteristics of households sampled for socio-economic survey in Yadgir Rf Micro watershed is presented in Table 2. The data indicated that, there were 93 (55.03%) men and 76 (44.97%) were women. The average population of landless was 4, marginal farmers were 6.45, small farmers were 4.29 and semi medium farmers were 5.0.

Table 2. Population characteristics in Yadgir Rf micro-watershed

Sl.No.	Particulars	LL (20)		MF (71)		SF (73)		SMF (5)		All (169)	
		N	%	N	%	N	%	N	%	N	%
1	Men	11	55	35	49	43	59	4	80	93	55
2	Women	9	45	36	51	30	41	1	20	76	45
Total		20	100	71	100	73	100	5	100	169	100
Average		4.0		6.45		4.29		5.0		4.97	

Age wise classification of population: The age wise classification of household members in Yadgir Rf Micro watershed is presented in Table 3. The indicated that, 33 (19.53%) of population were 0-15 years of age, 65 (38.46%) were 16-35 years of age, 56(33.14%) were 36-60 years of age and 15 (8.88 %) were above 61 years of age.

Table 3: Age wise classification of members of the household in Yadgir Rf micro-watershed

Sl.No.	Particulars	LL (20)		MF (71)		SF (73)		SMF (5)		All (169)	
		N	%	N	%	N	%	N	%	N	%
1	0-15 years of age	2	10	19	26.8	12	16.4	0	0	33	19.53
2	16-35 years of age	12	60	26	36.6	25	34.3	2	40	65	38.46
3	36-60 years of age	6	30	19	26.8	29	39.7	2	40	56	33.14
4	> 61 years	0	0	7	9.86	7	9.59	1	20	15	8.88
Total		20	100	71	100	73	100	5	100	169	100

Education level of household members: Education level of household members in Yadgir Rf Micro watershed is presented in Table 4. The results indicated that, there were 44.97 per cent of illiterates, 23.08 per cent of them had primary school education, 0.59 per cent middle school education, and 13.02 per cent high school education, 7.10 per cent of them had PUC education, 0.59 per cent of them had Diploma, 0.59 per cent of them had ITI, 8.28 per cent attained graduation, 1.18 per cent attained post graduation and 0.59 them had other education.

Table 4. Education level of members of the household in Yadgir Rf micro-watershed

Sl.No.	Particulars	LL (20)		MF (71)		SF (73)		SMF (5)		All (169)	
		N	%	N	%	N	%	N	%	N	%
1	Illiterate	2	10	36	50.7	35	48	3	60	76	45
2	Primary School	5	25	14	19.7	18	24.7	2	40	39	23.1
3	Middle School	0	0	0	0	1	1.37	0	0	1	0.59
4	High School	1	5	10	14.1	11	15.1	0	0	22	13
5	PUC	4	20	3	4.23	5	6.85	0	0	12	7.1
6	Diploma	1	5	0	0	0	0	0	0	1	0.59
7	ITI	0	0	0	0	1	1.37	0	0	1	0.59
8	Degree	7	35	5	7.04	2	2.74	0	0	14	8.28
9	Masters	0	0	2	2.82	0	0	0	0	2	1.18
10	Others	0	0	1	1.41	0	0	0	0	1	0.59
Total		20	100	71	100	73	100	5	100	169	100

Occupation of head of households: The data regarding the occupation of the household heads in Yadgir Rf Micro watershed is presented in Table 5. The results indicate that, 52.94 per cent of households' heads were practicing agriculture and 29.41 per cent of the household heads were agricultural Labour.

Table 5: Occupation of heads of households in Yadgir Rf micro-watershed

Sl.No.	Particulars	LL (5)		MF (11)		SF (17)		SMF (1)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Agriculture	0	0	1	9.1	16	94.12	1	100	18	52.94
2	Agricultural Labour	0	0	9	82	1	5.88	0	0	10	29.41
3	General Labour	2	40	0	0	0	0	0	0	2	5.88
4	Government Service	3	60	0	0	0	0	0	0	3	8.82
5	Private Service	0	0	1	9.1	0	0	0	0	1	2.94
6	Retired	0	0	0	0	1	5.88	0	0	1	2.94
Total		5	100	11	100	18	100	1	100	35	100

Occupation of the members of the household: The data regarding the occupation of the household members in Yadgir Rf Micro watershed is presented in Table 6. The results indicate that, agriculture was the major occupation for 33.73 per cent of the household members, 22.49 per cent were agricultural labour, 3.55 per cent were general labour, 2.96 per cent were working in government sector, 1.78 per cent were working in private service, 1.78 per cent were involved as housewife, and 0.59 per cent were childrens.

Table 6: Occupation of members of the household in Yadgir Rf micro-watershed

Sl.No.	Particulars	LL (20)		MF (71)		SF (73)		SMF (5)		All (169)	
		N	%	N	%	N	%	N	%	N	%
1	Agriculture	0	0	4	5.63	48	65.75	5	100	57	33.7
2	Agricultural Labour	1	5	33	46.5	4	5.48	0	0	38	22.5
3	General Labour	4	20	0	0	2	2.74	0	0	6	3.55
4	Government Service	4	20	1	1.41	0	0	0	0	5	2.96
5	Private Service	1	5	2	2.82	0	0	0	0	3	1.78
6	Retired	0	0	0	0	1	1.37	0	0	1	0.59
7	Student	3	15	23	32.4	18	24.66	0	0	44	26
8	Others	4	20	7	9.86	0	0	0	0	11	6.51
9	Housewife	3	15	0	0	0	0	0	0	3	1.78
10	Children	0	0	1	1.41	0	0	0	0	1	0.59
Total		20	100	71	100	73	100	5	100	169	100

Institutional Participation of household members: The data regarding the institutional participation of the household members in Yadgir Rf Micro watershed is presented in Table 7. The results show that, out of the total family members in the households 1.18 per cent of them were participating in gram panchayat and 98.8 per cent of them are not participating in any of the institutions.

Table 7: Institutional Participation of household member in Yadgir Rf micro-watershed

Sl.No.	Particulars	LL (20)		MF (71)		SF (73)		SMF (5)		All (169)	
		N	%	N	%	N	%	N	%	N	%
1	Gram Panchayat	0	0	0	0	2	2.74	0	0	2	1.18
2	No Participation	20	100	71	100	71	97.3	5	100	167	98.8
Total		20	100	71	100	73	100	5	100	169	100

Type of house owned: The data regarding the type of house owned by the households in Yadgir Rf Micro watershed is presented in Table 8. The results indicate that, 61.76 per cent of the households possess katcha house and 38.24 per cent possess pucca house.

Table 8. Type of house owned by households in Yadgir Rf micro-watershed

Sl.No.	Particulars	LL (5)		MF (11)		SF (17)		SMF (1)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Katcha	2	40	7	64	12	70.59	0	0	21	61.76
2	Pucca/RCC	3	60	4	36	5	29.41	1	100	13	38.24
Total		5	100	11	100	17	100	1	100	34	100

Durable assets owned by the households: The data regarding the Durable Assets owned by the households in Yadgir Rf Micro watershed is presented in Table 9. The result shows that, 76.47 per cent possess TV, 64.71 per cent possess mixer grinder, 2.94 per cent possess refrigerator, 32.35 per cent possess Bicycle, 44.12 per cent possess motor cycle, 85.29 per cent possess mobile phones and 2.94 per cent possess Computer/Laptop.

Table 9. Durable assets owned by households in Yadgir Rf micro-watershed

Sl.No.	Particulars	LL (5)		MF (11)		SF (17)		SMF (1)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Television	5	100	5	45	15	88.2	1	100	26	76.47
2	Mixer/Grinder	4	80	4	36	13	76.5	1	100	22	64.71
3	Refrigerator	1	20	0	0	0	0	0	0	1	2.94
4	Bicycle	1	20	0	0	9	52.9	1	100	11	32.35
5	Motor Cycle	3	60	3	27	8	47.1	1	100	15	44.12
6	Auto	0	0	0	0	1	5.88	0	0	1	2.94
7	Mobile Phone	5	100	8	73	15	88.2	1	100	29	85.29
8	Computer/Laptop	1	20	0	0	0	0	0	0	1	2.94

Average value of durable assets: The data regarding the average value of durable assets owned by the households in Yadgir Rf Micro watershed is presented in Table 10. The result shows that, the average value of television was Rs.3653.00, mixer grinder was Rs.1163.00, refrigerator was 15000.00, bicycle was Rs.1000.00, motor cycle was Rs. 31375.00, mobile phone was Rs.1441.00 and Computer/Laptop was Rs 20000.00.

Table 10. Average value of durable assets owned in Yadgir Rf micro-watershed

Sl.No.	Particulars	Average Value (Rs.)				
		LL (5)	MF (11)	SF (17)	SMF (1)	All (34)
1	Television	5400	5200	2666	2000	3653
2	Mixer/Grinder	1500	1150	1076	1000	1163
3	Refrigerator	15000	0	0	0	15000
4	Bicycle	1000	0	1000	1000	1000
5	Motor Cycle	35000	46666	24000	30000	31375
6	Auto	0	0	600000	0	600000
7	Mobile Phone	2083	2916	679	333	1441
8	Computer/Laptop	20000	0	0	0	20000

Farm implements owned: The data regarding the farm implements owned by the households in Yadgir Rf Micro watershed is presented in Table 11. About 2.94 per cent of the households possess Bullock Cart, 20.59 per cent possess plough and 11.76 per cent possess Sprayer, 61.76 per cent possess Weeder and 2.94 per cent possess tractor.

Table 11. Farm implements owned in Yadgir Rf micro-watershed

Sl.No.	Particulars	LL (5)		MF (11)		SF (17)		SMF (1)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Bullock Cart	0	0	1	9.09	0	0	0	0	1	2.94
2	Plough	0	0	2	18.2	4	23.53	1	100	7	20.59
3	Tractor	0	0	0	0	1	5.88	0	0	1	2.94
4	Sprayer	0	0	0	0	3	17.65	1	100	4	11.76
5	Weeder	2	40	3	27.3	15	88.24	1	100	21	61.76
6	Harvester	0	0	0	0	1	5.88	0	0	1	2.94
7	Thresher	0	0	0	0	1	5.88	0	0	1	2.94
8	Chaff Cutter	0	0	0	0	3	17.65	1	100	4	11.76

Average value of farm implements: The data regarding the average value of farm Implements owned by the households in Yadgir Rf Micro watershed is presented in Table

12. The results show that the average value of plough was Rs.636.00, bullock Cart was Rs.30000.00, seed/fertilizer drill was Rs.4000.00, sprayer and weeder was Rs.39.00 and tractor was Rs. 300000.

Table 12. Average value of farm implements in Yadgir Rf micro-watershed

Sl.No.	Particulars	Average Value (Rs.)				
		LL (5)	MF (11)	SF (17)	SMF (1)	All (34)
1	Bullock Cart	0	30000	0	0	30000
2	Plough	0	2000	500	500	636
3	Tractor	0	0	300000	0	300000
4	Sprayer	0	0	4000	4000	4000
5	Weeder	33	50	39	25	39
6	Harvester	0	0	55000	0	55000
7	Thresher	0	0	60000	0	60000
8	Chaff Cutter	0	0	3000	2000	2750

Livestock possession by the households: The data regarding the Livestock possession by the households in Yadgir Rf Micro watershed is presented in Table 13. The results indicate that, 20.59 per cent of the households possess bullocks and 2.94 per cent possess local cow.

Table 13. Livestock possession by households in Yadgir Rf micro-watershed

Sl.No.	Particulars	LL (5)		MF (11)		SF (17)		SMF (1)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Bullock	0	0	3	27	3	17.65	1	100	7	20.59
2	Local cow	0	0	0	0	1	5.88	0	0	1	2.94
3	blank	0	0	1	9.1	3	17.65	0	0	4	11.76

Average Labour availability: The data regarding the average labour availability in Yadgir Rf Micro watershed is presented in Table 14. The indicated that, own labour men available in the micro watershed was 1.64, women available in the micro watershed was 1.36, hired labour (men) available was 8.86 and hired labour (women) available was 13.96.

Table 14. Average labour availability in Yadgir Rf micro-watershed

Sl.No.	Particulars	LL (5)	MF (11)	SF (17)	SMF (1)	All (34)
		N	N	N	N	N
1	Hired labour Female	0	9.18	17	18	13.96
2	Own Labour Female	0	1.36	1.38	1	1.36
3	Own labour Male	0	1.27	1.81	3	1.64
4	Hired labour Male	0	5.73	10.81	12	8.86

Table 15. Adequacy of hired labour in Yadgir Rf micro-watershed

Sl.No.	Particulars	LL (5)		MF (11)		SF (17)		SMF (1)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Adequate	0	0	11	100	16	94.1	1	100	28	82.4

Adequacy of hired labour: The data regarding the adequacy of hired labour in Yadgir Rf Micro watershed is presented in Table 15. The results indicate that, 82.35 per cent of the household opined that hired labour was adequate.

Migration among the households: The data regarding the migration (Table 16) indicate that, 1.18 percent of the population was being migrated from the micro watershed.

Table 16. Migration among the households in Yadgir Rf micro-watershed

Sl.No.	Particulars	LL (20)		MF (71)		SF (73)		SMF (5)		All (169)	
		N	%	N	%	N	%	N	%	N	%
1	Migration	0	0.00	1	1.41	1	1.37	0	0.00	2	1.18

Average distance and duration of migration: The data regarding the average distance and duration of migration (Table 17) indicate that, people migrated to a distance of 525 kms on an average for 9 months.

Table 17. Average distance and duration of migration in Yadgir Rf micro-watershed

Sl.No.	Particulars	LL (0)	MF (1)	SF (1)	All (2)
		N	N	N	N
1	Avg. Distance (kms)	0	500	550	525
2	Avg. Duration (months)	0	12	6	9

Purpose of migration: The data regarding the purpose of migration (Table 18) indicate that, 50.00 percent of them went for the purpose of job/wage/work and 50.00 percent for education of the children.

Table 18. Purpose of migration by members of households in Yadgir Rf micro-watershed

Sl.No.	Particulars	LL (0)		MF (1)		SF (1)		All (2)	
		N	%	N	%	N	%	N	%
1	Job/wage/work	0	0	1	100	0	0	1	50
2	Education of the children	0	0	0	0	1	100	1	50
Total		0	100	1	100	1	100	2	100

Distribution of land (ha): The data regarding the distribution of land (ha) in Yadgir Rf Micro watershed is presented in Table 19. The results indicate that, 33.95 ha (100.00%) of dry land.

Table 19. Distribution of land (ha) in Yadgir Rf micro-watershed

Sl.No.	Particulars	LL (5)		MF (11)		SF (17)		SMF (1)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Dry	0	0	6.66	100	25.27	100	2.02	100	33.95	100
Total		0	100	6.66	100	25.27	100	2.02	100	33.95	100

Table 20. Average value of land (ha) in Yadgir Rf micro-watershed

Sl.No.	Particulars	LL (5)	MF (11)	SF (17)	SMF (1)	All (34)
		N	N	N	N	N
1	Dry	0	6.66	25.27	2.02	0

Average value of land (ha): The data regarding the average land value (Rs./ha) in Yadgir Rf Micro watershed is presented in Table 20. The results show that the average value of dry land was Rs.0.00.

Status of bore wells: The data regarding the status of bore wells in Yadgir Rf Micro watershed is presented in Table 21. The results indicate that, there were 1 De-functioning bore wells and 1 functioning bore wells among the sampled households in micro watershed.

Table 21. Status of bore wells in Yadgir Rf micro-watershed

Sl.No.	Particulars	LL (5)	MF (11)	SF (17)	SMF (1)	All (34)
		N	N	N	N	N
1	De-functioning	0	1	0	0	1
2	Functioning	0	1	0	0	1

Source of irrigation: The data regarding the source of irrigation in Yadgir Rf Micro watershed is presented in Table 22. The results indicate that bore well for 2.94 per cent of the households.

Table 22. Source of irrigation in Yadgir Rf micro-watershed

Sl.No.	Particulars	LL (5)		MF (11)		SF (17)		SMF (1)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Bore Well	0	0	1	9.09	0	0	0	0	1	2.94

Depth of water (Avg. In meters): The data regarding the depth of water in Yadgir Rf Micro watershed is presented in Table 23. The results revealed that, the depth of bore well was 2.69 meter.

Table 23. Depth of water (Avg. In meters) in Yadgir Rf micro-watershed

Sl.No.	Particulars	LL (5)	MF (11)	SF (17)	SMF (1)	All (34)
		N	N	N	N	N
1	Bore Well	0	8.31	0	0	2.69

Irrigated Area (ha): The data regarding the irrigated area (ha) in Yadgir Rf Micro watershed is presented in Table 24. The results indicate that, the availability of irrigation water was used for kharif crops was 0.81 ha.

Table 24. Irrigated Area (ha) in Yadgir Rf micro-watershed

Sl.No.	Particulars	LL (5)	MF (11)	SF (17)	SMF (1)	All (34)
1	Kharif	0	0.81	0	0	0.81
	Total	0	0.81	0	0	0.81

Cropping pattern: The data regarding the cropping pattern in Yadgir Rf Micro watershed is presented in Table 25. The results indicate that, farmers have grown Maize (7.75 ha), Red gram (togari) (6.19 ha), Groundnut (6.07 ha), Cotton (4.45 ha) Red gram (togari) (4.18 ha), Jowar (2.02 ha) and Groundnut (1.67 ha).

Table 25. Cropping pattern in Yadgir Rf micro-watershed

Sl.No.	Particulars	LL (5)	MF (11)	SF (17)	SMF (1)	All (34)
1	Rabi - Maize	0	0	5.73	2.02	7.75
2	Rabi - Red gram (togari)	0	0	6.19	0	6.19
3	Rabi - Groundnut	0	0	6.07	0	6.07
4	Rabi - Cotton	0	0	4.45	0	4.45
5	Kharif - Red gram (togari)	0	4.18	0	0	4.18
6	Kharif - Jowar	0	0.81	1.21	0	2.02
7	Kharif - Groundnut	0	1.67	0	0	1.67

Cropping intensity: The data regarding the cropping intensity in Yadgir Rf Micro watershed is presented in Table 26. The results indicate that, the cropping intensity was 100.00 per cent.

Table 26. Cropping intensity (%) in Yadgir Rf micro-watershed

Sl.No.	Particulars	LL (5)	MF (11)	SF (17)	SMF (1)	All (34)
1	Cropping Intensity	0	100	100	100	100

Possession of bank account and savings: The data regarding the possession of bank account and saving in Yadgir Rf micro-watershed is presented in Table 27. The results indicate that, 94.12 cent of the households posses bank account and 61.76 per cent of them have savings.

Table 27. Possession of Bank account and savings in Yadgir Rf micro-watershed

Sl.No.	Particulars	LL (5)		MF (11)		SF (17)		SMF (1)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Account	5	100	9	81.82	17	100	1	100	32	94.12
2	Savings	4	80	0	0	16	94.12	1	100	21	61.76

Borrowing status: The data regarding the borrowing status in Yadgir Rf micro-watershed is presented in Table 28. The results indicate that, 64.71 percent of the sample farmers have borrowed credit from different sources.

Table 28. Borrowing status in Yadgir Rf micro-watershed

Sl.No.	Particulars	LL (5)		MF (11)		SF (17)		SMF (1)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Credit Aailed	4	80	1	9.09	16	94.1	1	100	22	64.71

Table 29. Source of credit borrowed by households in Yadgir Rf micro-watershed

Sl.No.	Particulars	LL (5)		MF (11)		SF (17)		SMF (1)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Commercial Bank	0	0	0	0	4	25	0	0	4	18.18
2	Cooperative Bank	0	0	1	100	1	6.25	0	0	2	9.09
3	Grameena Bank	0	0	2	200	10	62.5	1	100	13	59.09
4	Money Lender	0	0	0	0	5	31.3	0	0	5	22.73

Source of credit: The data regarding the source of credit availed by households in Yadgir Rf micro-watershed is presented in Table 29. The results show that, 18.18 per cent have borrowed loan from commercial banks and 9.09 per cent have borrowed loan from

Cooperative bank and 59.09 per cent have borrowed loan from Grameena Bank and 22.73 per cent have borrowed loan from money lender.

Avg. Credit amount: The data regarding the avg. Credit amount in Yadgir Rf micro-watershed is presented in Table 30. The results show that, farmers have borrowed Avg.Credit of Rs.64772.73 from different sources.

Table 30. Avg. Credit amount in Yadgir Rf micro-watershed

Sl.No.	Particulars	LL (4)	MF (1)	SF (16)	SMF (1)	All (22)
		N	N	N	N	N
1	Average Credit	0	150000	76875	45000	64772.7

Purpose of credit borrowed (institutional Source): The data regarding the purpose of credit borrowed - Institutional Credit in Yadgir Rf micro-watershed is presented in Table 31. The results indicate that, 100.00 per cent of the households have borrowed loan for agriculture.

Table 31. Purpose of credit borrowed (institutional Source) by households in Yadgir Rf micro-watershed

SN	Particulars	LL (0)		MF (2)		SF (15)		SMF (1)		All (18)	
		N	%	N	%	N	%	N	%	N	%
1	Agriculture production	0	0	2	100	15	100	1	100	18	100

Purpose of credit borrowed (Private Source): The data regarding the purpose of credit borrowed – Private Source in Yadgir Rf micro-watershed is presented in Table 32. The results indicate that, 60.00 per cent of the households have borrowed loan for agriculture, construction-house (20.00 %) and Social functions like marriage (20.00 %).

Table 32. Purpose of credit borrowed (Private Source) by households in Yadgir Rf micro-watershed

Sl.No.	Particulars	LL (0)		MF (0)		SF (5)		All (5)	
		N	%	N	%	N	%	N	%
1	Agriculture production	0	0	0	0	3	60	3	60
2	Construction-house, Construction-cattle shed	0	0	0	0	1	20	1	20
3	Social functions like marriage	0	0	0	0	1	20	1	20

Repayment status of household (institutional Source): The data regarding the repayment status of credit borrowed from institutional Source by households in Yadgir Rf micro watershed is presented in Table 33. The results indicate that, 5.00 per cent of the households have partially paid and 95.00 per cent have unpaid.

Table 33. Repayment status of household (institutional Source) in Yadgir Rf micro-watershed

Sl.No.	Particulars	LL (0)		MF (0)		SF (5)		SMF (0)		All (5)	
		N	%	N	%	N	%	N	%	N	%
1	Partially paid	0	0	0	0	1	6.25	0	0	1	5
2	Un paid	0	0	3	100	15	93.8	1	100	19	95

Repayment status of household (Private Source): The data regarding the repayment status of credit borrowed from private sources by households in Yadgir Rf micro

watershed is presented in Table 34. The results indicate that, 100.00 per cent of the households have partially paid.

Table 34. Repayment status of household (Private Source) in Yadgir Rf micro-watershed

Sl.No.	Particulars	LL (0)		MF (0)		SF (5)		All (5)	
		N	%	N	%	N	%	N	%
1	Partially paid	0	0	0	0	5	100	5	100

Opinion regarding institutional sources of credit: The data regarding the opinion on institutional sources of credit in Yadgir Rf micro watershed is presented in Table 35. The results indicate that, 100.00 per cent of the households opined that credit helped to perform timely agricultural operations.

Table 35. Opinion regarding institutional sources of credit in Yadgir Rf micro-watershed

Sl. No.	Particulars	LL (0)		MF (3)		SF (16)		SMF (1)		All (20)	
		N	%	N	%	N	%	N	%	N	%
		1	Helped to perform timely agricultural operations	0	0	3	100	16	100	1	100

Opinion regarding Non- institutional sources of credit: The data regarding the opinion on non-institutional sources of credit in Yadgir Rf micro watershed is presented in Table 36. The results indicate that, 100.00 per cent of the households opined that credit helped to perform timely agricultural operations.

Table 36. Opinion regarding Non- institutional sources of credit in Yadgir Rf micro-watershed

Sl.No.	Particulars	LL (0)		MF (0)		SF (5)		All (5)	
		N	%	N	%	N	%	N	%
1	Loan amount was adequate to fulfil the requirement	0	0	0	0	5	100	5	100

Cost of Cultivation of Cotton: The data regarding the cost of cultivation (Rs/ha) of Cotton in Yadgir Rf micro watershed is presented in Table 37.a. The results indicate that, the total cost of cultivation (Rs/ha) for Cotton was Rs. 34406.70. The gross income realized by the farmers was Rs. 48700.17. The net income from Cotton cultivation was Rs.14293.47, thus the benefit cost ratio was found to be 1:1.40.

Table 37(a). Cost of Cultivation of Cotton in Yadgir Rf micro-watershed

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
I	Cost A1				
1	Hired Human Labour	Man days	25.8	5694.72	16.55
2	Bullock	Pairs/day	1.92	1152.67	3.35
3	Tractor	Hours	3.29	2470	7.18
4	Machinery	Hours	0	0	0
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	0.69	504.29	1.47
6	Seed Inter Crop	Kgs.	0	0	0
7	FYM	Quintal	7.2	10806.25	31.41
8	Fertilizer + micronutrients	Quintal	3.02	1797.61	5.22
9	Pesticides (PPC)	Kgs/liters	1.37	1372.22	3.99
10	Irrigation	Number	0	0	0
11	Repairs		0	0	0
12	Msc. Charges (Marketing costs etc)		0	0	0
13	Depreciation charges		0	46.66	0.14
14	Land revenue and Taxes		0	0	0
II	Cost B1				
16	Interest on working capital			1737.65	5.05
17	Cost B1 = (Cost A1 + sum of 15 and 16)			25582.06	74.35
III	Cost B2				
18	Rental Value of Land			283.33	0.82
19	Cost B2 = (Cost B1 + Rental value)			25865.4	75.18
IV	Cost C1				
20	Family Human Labour		20.93	5413.42	15.73
21	Cost C1 = (Cost B2 + Family Labour)			31278.81	90.91
V	Cost C2				
22	Risk Premium			0	0
23	Cost C2 = (Cost C1 + Risk Premium)			31278.81	90.91
VI	Cost C3				
24	Managerial Cost			3127.88	9.09
25	Cost C3 = (Cost C2 + Managerial Cost)			34406.7	100
VII	Economics of the Crop				
a.	Main Product	a) Main Product (q)		8.65	48700.17
		b) Main Crop Sales Price (Rs.)			5633.33
b.	Gross Income (Rs.)			48700.17	
c.	Net Income (Rs.)			14293.47	
d.	Cost per Quintal (Rs./q.)			3979.95	
e.	Benefit Cost Ratio (BC Ratio)			1:1.4	

Cost of Cultivation of Redgram: The data regarding the cost of cultivation (Rs/ha) of Redgram in Yadgir Rf micro watershed is presented in Table 37.b. The results indicate that, the total cost of cultivation (Rs/ha) for Redgram was Rs. 54597.19. The gross income realized by the farmers was Rs. 36298.38. The net income from Redgram cultivation was Rs.-18298.82, thus the benefit cost ratio was found to be 1:0.70.

Table 37(b). Cost of Cultivation of Redgram in Yadgir Rf micro-watershed

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
I	Cost A1				
1	Hired Human Labour	Man days	44.37	9603.07	17.59
2	Bullock	Pairs/day	2.87	2694.83	4.94
3	Tractor	Hours	3.43	2181.58	4
4	Machinery	Hours	0	0	0
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	18.53	1373.48	2.52
6	Seed Inter Crop	Kgs.	0	0	0
7	FYM	Quintal	5.74	17232.56	31.56
8	Fertilizer + micronutrients	Quintal	5.28	4842.34	8.87
9	Pesticides (PPC)	Kgs / liters	1.11	1172.29	2.15
10	Irrigation	Number	0	0	0
11	Repairs		0	0	0
12	Msc. Charges (Marketing costs etc)		0	0	0
13	Depreciation charges		0	135.82	0.25
14	Land revenue and Taxes		0	0	0
II	Cost B1				
16	Interest on working capital			2954.48	5.41
17	Cost B1 = (Cost A1 + sum of 15 and 16)			42190.45	77.28
III	Cost B2				
18	Rental Value of Land			272.22	0.5
19	Cost B2 = (Cost B1 + Rental value)			42462.67	77.77
IV	Cost C1				
20	Family Human Labour		27.78	7171.14	13.13
21	Cost C1 = (Cost B2 + Family Labour)			49633.81	90.91
V	Cost C2				
22	Risk Premium			0	0
23	Cost C2 = (Cost C1 + Risk Premium)			49633.81	90.91
VI	Cost C3				
24	Managerial Cost			4963.38	9.09
25	Cost C3 = (Cost C2 + Managerial Cost)			54597.19	100
VII	Economics of the Crop				
a.	Main Product	a) Main Product (q)	9.55	36259.78	
		b) Main Crop Sales Price (Rs.)		3795.83	
By Product		e) Main Product (q)	0.51	38.59	
		f) Main Crop Sales Price (Rs.)		75	
b.	Gross Income (Rs.)			36298.38	
c.	Net Income (Rs.)			-18298.82	
d.	Cost per Quintal (Rs./q.)			5715.47	
e.	Benefit Cost Ratio (BC Ratio)			1:0.7	

Cost of Cultivation of Maize: The data regarding the cost of cultivation (Rs/ha) of Maize in Yadgir Rf micro watershed is presented in Table 37.c. The results indicate, the total cost of cultivation (Rs/ha) for Maize was Rs.15086.15. The gross income realized by the farmers was Rs. 37957.22. The net income from Maize cultivation was Rs. 22871.07, thus the benefit cost ratio was found to be 1:2.50.

Table 37(c). Cost of Cultivation of Maize in Yadgir Rf micro-watershed

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
I	Cost A1				
1	Hired Human Labour	Man days	21.25	4612.86	30.58
2	Bullock	Pairs/day	1.18	708.46	4.7
3	Tractor	Hours	2.45	1849.56	12.26
4	Machinery	Hours	0	0	0
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	11.74	862.78	5.72
6	Seed Inter Crop	Kgs.	0	0	0
7	FYM	Quintal	0	0	0
8	Fertilizer + micronutrients	Quintal	1.97	1231.31	8.16
9	Pesticides (PPC)	Kgs/liters	0.78	784.13	5.2
10	Irrigation	Number	0	0	0
11	Repairs		0	0	0
12	Msc. Charges (Marketing costs etc)		0	0	0
13	Depreciation charges		0	26.64	0.18
14	Land revenue and Taxes		0	0	0
II	Cost B1				
16	Interest on working capital			345.39	2.29
17	Cost B1 = (Cost A1 + sum of 15 and 16)			10421.12	69.08
III	Cost B2				
18	Rental Value of Land			283.33	1.88
19	Cost B2 = (Cost B1 + Rental value)			10704.45	70.96
IV	Cost C1				
20	Family Human Labour		11.49	3010.23	19.95
21	Cost C1 = (Cost B2 + Family Labour)			13714.68	90.91
V	Cost C2				
22	Risk Premium			0	0
23	Cost C2 = (Cost C1 + Risk Premium)			13714.68	90.91
VI	Cost C3				
24	Managerial Cost			1371.47	9.09
25	Cost C3 = (Cost C2 + Managerial Cost)			15086.15	100
VII	Economics of the Crop				
a.	Main Product	a) Main Product (q)		14.64	34770.72
		b) Main Crop Sales Price (Rs.)			2375
	By Product	c) Main Product (q)		5.31	3186.5
		d) Main Crop Sales Price (Rs.)			600
b.	Gross Income (Rs.)			37957.22	
c.	Net Income (Rs.)			22871.07	
d.	Cost per Quintal (Rs./q.)			1030.45	
e.	Benefit Cost Ratio (BC Ratio)			1:2.5	

Cost of Cultivation of Groundnut: The data regarding the cost of cultivation (Rs/ha) of Groundnut in Yadgir Rf micro watershed is presented in Table 37.d. The results indicate that, the total cost of cultivation (Rs/ha) for Groundnut was Rs. 70279.00. The gross income realized by the farmers was Rs.39755.68. The net income from Groundnut cultivation was Rs. -30523.32, thus the benefit cost ratio was found to be 1:0.60.

Table 37(d). Cost of Cultivation of Groundnut in Yadgir Rf micro-watershed

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
I	Cost A1				
1	Hired Human Labour	Man days	27.07	5798.88	8.25
2	Bullock	Pairs/day	1.53	1106.05	1.57
3	Tractor	Hours	2.8	1992.09	2.83
4	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	133.59	16895.15	24.04
5	FYM	Quintal	12.35	23465	33.39
6	Fertilizer + micronutrients	Quintal	3.45	3008.99	4.28
7	Pesticides (PPC)	Kgs / liters	1.1	1097.78	1.56
8	Depreciation charges		0	1726.26	2.46
II	Cost B1				
9	Interest on working capital			5336.03	7.59
10	Cost B1 = (Cost A1 + sum of 15 and 16)			60426.22	85.98
III	Cost B2				
11	Rental Value of Land			283.33	0.4
12	Cost B2 = (Cost B1 + Rental value)			60709.55	86.38
IV	Cost C1				
13	Family Human Labour		12.42	3180.44	4.53
14	Cost C1 = (Cost B2 + Family Labour)			63890	90.91
V	Cost C2				
15	Risk Premium			0	0
16	Cost C2 = (Cost C1 + Risk Premium)			63890	90.91
VI	Cost C3				
17	Managerial Cost			6389	9.09
18	Cost C3 = (Cost C2 + Managerial Cost)			70279	100
VII	Economics of the Crop				
a.	Main Product	a) Main Product (q)		8.1	37186.88
		b) Main Crop Sales Price (Rs.)			4590
		c) Intercrop (q)		0.7	0
	By Product	d) Main Product (q)		5.35	2568.8
		e) Main Crop Sales Price (Rs.)			480
b.	Gross Income (Rs.)			39755.68	
c.	Net Income (Rs.)			-30523.32	
d.	Cost per Quintal (Rs./q.)			7988.53	
e.	Benefit Cost Ratio (BC Ratio)			1:0.6	

Cost of Cultivation of Jowar: The data regarding the cost of cultivation (Rs/ha) of Jowar in Yadgir Rf micro watershed is presented in Table 37.e. The results indicate that, the total cost of cultivation (Rs/ha) for Jowar was Rs.19949.72. The gross income realized by the farmers was Rs. 20583.33. The net income from Jowar cultivation was Rs. 633.61, thus the benefit cost ratio was found to be 1:1.00.

Table 37(e). Cost of Cultivation of Jowar in Yadgir Rf micro-watershed

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
I	Cost A1				
1	Hired Human Labour	Man days	26.48	4693	23.52
2	Bullock	Pairs/day	0.69	686.11	3.44
3	Tractor	Hours	2.47	1482	7.43
4	Machinery	Hours	0	0	0
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	13.72	1166.39	5.85
6	Seed Inter Crop	Kgs.	0	0	0
7	FYM	Quintal	0	0	0
8	Fertilizer + micronutrients	Quintal	4.53	3924.56	19.67
9	Pesticides (PPC)	Kgs /liters	0	0	0
10	Irrigation	Number	4.94	0	0
11	Repairs		0	1000	5.01
12	Msc. Charges (Marketing costs etc)		0	0	0
13	Depreciation charges		0	0	0
14	Land revenue and Taxes		0	0	0
II	Cost B1				
16	Interest on working capital			610.91	3.06
17	Cost B1 = (Cost A1 + sum of 15 and 16)			13562.97	67.99
III	Cost B2				
18	Rental Value of Land			188.89	0.95
19	Cost B2 = (Cost B1 + Rental value)			13751.86	68.93
IV	Cost C1				
20	Family Human Labour		17.56	4384.25	21.98
21	Cost C1 = (Cost B2 + Family Labour)			18136.11	90.91
V	Cost C2				
22	Risk Premium			0	0
23	Cost C2 = (Cost C1 + Risk Premium)			18136.11	90.91
VI	Cost C3				
24	Managerial Cost			1813.61	9.09
25	Cost C3 = (Cost C2 + Managerial Cost)			19949.72	100
VII	Economics of the Crop				
a.	Main Product	a) Main Product (q)		10.29	20583.33
		b) Main Crop Sales Price (Rs.)			2000
b.	Gross Income (Rs.)				20583.33
c.	Net Income (Rs.)				633.61
d.	Cost per Quintal (Rs./q.)				1938.43
e.	Benefit Cost Ratio (BC Ratio)				1:1

Adequacy of fodder: The data regarding the adequacy of fodder in Yadgir Rf Micro watershed is presented in Table 38. The results indicate that, 11.76 per cent of the households opined that dry fodder was adequate and 2.94 per cent of them opined dry fodder was inadequate.

Table 38. Adequacy of fodder in Yadgir Rf micro-watershed

Sl.No.	Particulars	LL (5)		MF (11)		SF (17)		SMF (1)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Adequate-Dry Fodder	0	0	3	27.27	0	0	1	100	4	11.76
2	Inadequate-Dry Fodder	0	0	0	0	1	5.88	0	0	1	2.94

Average annual gross income: The data regarding the annual gross income in Yadgir Rf Micro watershed is presented in Table 39. The results indicate that, the farmers have annual gross income of Rs. 73020.59 in micro-watershed, of which Rs. 39491.18 is from agriculture itself.

Table 39. Average annual gross income in Yadgir Rf micro-watershed

Sl.No.	Particulars	LL (5)	MF (11)	SF (17)	SMF (1)	All (34)
		Rs.	Rs.	Rs.	Rs.	Rs.
1	Service/salary	120000	0	0	0	17647.1
2	Wage	37000	0	19411.8	25000	15882.4
Income(Rs.)		157000	14818.2	84747.1	94000	73020.6

Average annual Expenditure: The data regarding the average annual expenditure in Yadgir Rf Micro watershed is presented in Table 40. The results indicate that, the farmers have annual gross expenditure of Rs. 620606.06 in micro-watershed, of which Rs. 24735.29 is from agriculture itself.

Table 40. Average annual Expenditure in Yadgir Rf micro-watershed

Sl.No.	Particulars	LL (5)	MF (11)	SF (17)	SMF (1)	All (34)
		Rs.	Rs.	Rs.	Rs.	Rs.
1	Service/salary	500000	0	0	0	14705.9
2	Wage	29166.7	0	0	0	2573.53
3	Agriculture	0	20272.7	32166.7	39000	24735.3
Total		529167	20272.7	32166.7	39000	620606

Horticulture species grown: The data regarding horticulture species grown in Yadgir Rf Micro watershed is presented in Table 41. The results indicate that, the total number of horticultural trees grown (both field and backyard) by the sampled households were coconut (5), clustered apple (2), Guava (1) and Mango (2).

Table 41. Horticulture species grown in Yadgir Rf micro-watershed

Sl.No.	Particulars	LL (5)		MF (11)		SF (17)		SMF (1)		All (34)	
		F	B	F	B	F	B	F	B	F	B
1	Coconut	0	0	0	0	3	2	0	0	3	2
2	Custard apple	0	0	0	0	2	0	0	0	2	0
3	Guava	0	0	0	0	0	1	0	0	0	1
4	Mango	0	0	0	0	2	0	0	0	2	0

*F= Field B=Back Yard

Interest towards cultivation of horticulture crops: The data regarding Table (42) indicates that, 32.35 percent/ None of the households shown interest to cultivate horticultural crops.

Table 42. Interest towards cultivation of horticulture crops in Yadgir Rf micro-watershed

Sl. No.	Particulars	LL (5)		MF (11)		SF (17)		SMF (1)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Interested towards cultivation of horticulture crops	0	0	0	0	10	59	1	100	11	32.4

Forest species grown: The data regarding forest species grown in Yadgir Rf Micro watershed is presented in Table 43. The results indicate that, households have planted 47 neem trees, 3 tamarind trees, 6 banyan trees together in both field and backyard.

Table 43. Forest species grown in Yadgir Rf micro-watershed

Sl.No.	Particulars	LL (5)		MF (11)		SF (17)		SMF (1)		All (34)	
		F	B	F	B	F	B	F	B	F	B
1	Neem	0	0	7	0	33	3	2	2	42	5
2	Tamarind	0	0	0	0	3	0	0	0	3	0
3	Banyan	0	0	0	0	6	0	0	0	6	0

*F= Field B=Back Yard

Average additional investment capacity: The data regarding average additional investment capacity in Yadgir Rf Micro watershed is presented in Table 44. The results indicate that, households have an average investment capacity of Rs. 735.29 for land development and Rs.676.47 for adoption of improved livestock breeds.

Table 44. Average additional investment capacity of households in Yadgir Rf micro-watershed

Sl.No.	Particulars	LL (5)	MF (11)	SF (17)	SMF (1)	All (34)
		Rs.	Rs.	Rs.	Rs.	Rs.
1	Land development	0	2090.91	117.65	0	735.29
2	Improved crop production	0	1909.09	117.65	0	676.47

Source of funds for additional investment: The data regarding source of funds for additional investment in Yadgir Rf Micro watershed is presented in Table 45. The results indicate that, the sources of finance raised from bank as a loan and from own sources for land development were 32.35.

Table 45. Source of funds for additional investment in Yadgir Rf micro-watershed

Sl. No	Item	Land development		Improved crop production	
		N	%	N	%
1	Own funds	11	32.35	11	32.35

Marketing of agricultural produce: The data regarding marketing of the agricultural produce in Yadgir Rf Micro watershed is presented in Table 46. The results indicated that, 100.00 percent of output of Cotton was sold in the market with average price of Rs. 5633.33; 100.00 percent of output of Groundnut was sold in the market with average

price of Rs. 4590.00; 86.21 percent of output of Jowar was sold in the market with average price of Rs. 2000.00; 100.00 percent of output of Maize was sold in the market with average price of Rs. 2375.00 and 95.40 percent of output of Redgram was sold in the market with average price of Rs. 3795.83.

Table 46. Marketing of agricultural produce in Yadgir Rf micro-watershed

Sl.No	Crops	Output obtained (q)	Output retained (q)	Output sold (q)	Output sold (%)	Avg. Price obtained (Rs/q)
1	Cotton	39	0	39	100	5633
2	Groundnut	58	0	58	100	4590
3	Jowar	29	4	25	86	2000
4	Maize	138	0	138	100	2375
5	Redgram	87	4	83	95	3796

Marketing channels used for sale of agricultural produce: The data regarding marketing channels used for sale of agricultural produce in Yadgir Rf Micro watershed is presented in Table 47. The results indicated that, 85.29 cent of the households have sold agricultural produce to the local/village merchants.

Table 47. Marketing channels used for sale of agricultural produce in Yadgir Rf micro-watershed

Sl.No.	Particulars	LL (5)		MF (11)		SF (17)		SMF (1)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Local/village Merchant	0	0	11	100	17	100	1	100	29	85.29

Mode of transport of agricultural produce: The data regarding mode of transport of agricultural produce in Yadgir Rf Micro watershed is presented in Table 48. The results indicated that, 32.35 cent of the households have used tractor and 41.18 per cent have used Cart.

Table 48. Mode of transport of agricultural produce in Yadgir Rf micro-watershed

Sl.No.	Particulars	LL (5)		MF (11)		SF (17)		SMF (1)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Cart	0	0	0	0	13	76.5	1	100	14	41.18
2	Tractor	0	0	7	64	4	23.5	0	0	11	32.35
3	Truck	0	0	4	36	0	0	0	0	4	11.76

Table 49. Incidence of soil and water erosion problems in Yadgir Rf micro-watershed

Sl.No.	Particulars	LL (5)		MF (11)		SF (17)		SMF (1)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Soil and water erosion problems in the farm	0	0	11	100	1	5.88	0	0	12	35.29

Incidence of soil and water erosion problems: The data regarding incidence of incidence of soil and water erosion problems in Yadgir Rf Micro watershed is presented in Table 49. The results indicate that, 35.29 per cent of the households have experienced soil and water erosion problems.

Interest towards soil testing: The data regarding Interest shown towards soil testing in Yadgir Rf Micro watershed is presented in Table 50. The results indicated that, 82.35 per cent of the households were interested towards soil testing.

Table 50. Interest regarding soil testing in Yadgir Rf micro-watershed

Sl.No.	Particulars	LL (5)		MF (11)		SF (17)		SMF (1)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Interest in soil test	0	0	11	100	16	94.1	1	100	28	82.35

Usage pattern of fuel for domestic use: The data on usage pattern of fuel for domestic use in Yadgir Rf Micro watershed is presented in Table 51. The results indicated that, firewood was the major source of fuel for domestic use for 55.88 per cent of the households followed by LPG (44.12%).

Table 51. Usage pattern of fuel for domestic use in Yadgir Rf micro-watershed

Sl.No.	Particulars	LL (5)		MF (11)		SF (17)		SMF (1)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Fire Wood	1	20	5	45.5	12	70.6	1	100	19	55.88
5	LPG	4	80	6	54.6	5	29.4	0	0	15	44.12

Source of drinking water: The data on source of drinking water in Yadgir Rf Micro watershed is presented in Table 52. The results indicated that, piped supply of water was the major source for drinking water for 94.12 per cent of the households followed by bore well waters supply (2.94 %).

Table 52. Source of drinking water in Yadgir Rf micro-watershed

Sl.No.	Particulars	LL (5)		MF (11)		SF (17)		SMF (1)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Piped supply	4	80	11	100	17	100	0	0	32	94.12
2	Bore Well	0	0	0	0	0	0	1	100	1	2.94

Source of light: The data on source of light in Yadgir Rf Micro watershed is presented in Table 53. The results indicated that, electricity was the major source of light for 100.00 per cent of the households.

Table 53. Source of light in Yadgir Rf micro-watershed

Sl.No.	Particulars	LL (5)		MF (11)		SF (17)		SMF (1)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Electricity	5	100	11	100	17	100	1	100	34	100

Existence of sanitary toilet facility: The data on availability of toilet facility in Yadgir Rf Micro watershed is presented in Table 54. The results indicated that, 29.41 per cent of the households possess toilets.

Table 54. Existence of sanitary toilet facility in Yadgir Rf micro-watershed

Sl.No.	Particulars	LL (5)		MF (11)		SF (17)		SMF (1)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Sanitary toilet facility	1	20	2	18	6	35.29	1	100	10	29.4

Possession of PDS card: The data regarding possession of PDS card in Yadgir Rf Micro watershed is presented in Table 55. The results indicated that, 97.06 per cent of the households possessed BPL card, 2.94 per cent possessed APL card.

Table 55. Possession of PDS card in Yadgir Rf micro-watershed

Sl.No.	Particulars	LL (5)		MF (11)		SF (17)		SMF (1)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	APL	1	20	0	0	0	0	0	0	1	2.94
2	BPL	4	80	11	100	17	100	1	100	33	97.06

Participation in NREGA programme: The data regarding Participation in NREGA programme in Yadgir Rf Micro watershed is presented in Table 56. The results indicated that, only 47.06 percent of the households have participated in NREGA programme.

Table 56. Participation in NREGA programme in Yadgir Rf micro-watershed

Sl.No.	Particulars	LL (5)		MF (11)		SF (17)		SMF (1)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Participation in NREGA programme	0	0	0	0	16	94.1	0	0	16	47.1

Adequacy of food items: The data regarding adequacy of food items in Yadgir Rf Micro watershed is presented in Table 57. The results indicated that, the extent of adequacy of food items for cereals, pulses, Oilseeds and vegetables were 58.82, 55.88, 5.88, 32.35 per cent respectively, similarly for Fruits (61.76%), milk (88.24%), Egg (2.94%), and Meat (5.88%).

Table 57. Adequacy of food items in Yadgir Rf micro-watershed

Sl.No.	Particulars	LL (5)		MF (11)		SF (17)		SMF (1)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Cereals	2	40	1	9.09	16	94.12	1	100	20	58.82
2	Pulses	2	40	0	0	16	94.12	1	100	19	55.88
3	Oilseed	0	0	2	18.2	0	0	0	0	2	5.88
4	Vegetables	1	20	9	81.8	1	5.88	0	0	11	32.35
5	Fruits	2	40	2	18.2	16	94.12	1	100	21	61.76
6	Milk	3	60	9	81.8	17	100	1	100	30	88.24
7	Egg	0	0	0	0	1	5.88	0	0	1	2.94
8	Meat	0	0	1	9.09	1	5.88	0	0	2	5.88

Table 58. Inadequacy of food items in Yadgir Rf micro-watershed

Sl.No.	Particulars	LL (5)		MF (11)		SF (17)		SMF (1)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Cereals	1	20	10	90.9	1	5.88	0	0	12	35.29
2	Pulses	1	20	11	100	1	5.88	0	0	13	38.24
3	Oilseed	3	60	9	81.8	17	100	1	100	30	88.24
4	Vegetables	2	40	2	18.2	16	94.12	1	100	21	61.76
5	Fruits	1	20	9	81.8	1	5.88	0	0	11	32.35
6	Milk	1	20	2	18.2	0	0	0	0	3	8.82
7	Egg	3	60	11	100	16	94.12	1	100	31	91.18
8	Meat	2	40	10	90.9	16	94.12	1	100	29	85.29

Inadequacy of food items: The data regarding in adequacy of food items in Yadgir Rf Micro watershed is presented in Table 58. The results indicated that, the extent of in adequacy of food items for cereals, pulses, Oilseeds and vegetables were 35.29, 38.24, 88.24, 61.76 and 85.29 per cent respectively, similarly for fruits (32.35%), milk (8.82%), egg (91.18%) and meat (85.29%).

Response on market surplus of food items: The data regarding adequacy of food items in Yadgir Rf Micro watershed is presented in Table 59. The results indicated that, the extent of adequacy of food items for cereals, pulses, Oilseeds and vegetables were 5.88, 5.88, 5.88, 5.88 per cent respectively, similarly for fruits (5.88%), milk (5.88%), egg (5.88%) and meat (5.88%).

Table 59. Response on market surplus of food items in Yadgir Rf micro-watershed

Sl.No.	Particulars	LL (5)		MF (11)		SF (17)		SMF (1)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Cereals	2	40	0	0	0	0	0	0	2	5.88
2	Pulses	2	40	0	0	0	0	0	0	2	5.88
3	Oilseed	2	40	0	0	0	0	0	0	2	5.88
4	Vegetables	2	40	0	0	0	0	0	0	2	5.88
5	Fruits	2	40	0	0	0	0	0	0	2	5.88
6	Milk	2	40	0	0	0	0	0	0	2	5.88
7	Egg	2	40	0	0	0	0	0	0	2	5.88
8	Meat	2	40	0	0	0	0	0	0	2	5.88

Table 60. Farming constraints experienced in Yadgir Rf micro-watershed

SN	Particulars	LL (5)		MF (11)		SF (17)		SMF (1)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Lower fertility status of the soil	0	0	11	100	20	117.65	1	100	32	94.12
2	Wild animal menace on farm field	0	0	11	100	17	100	1	100	29	85.29
3	Frequent incidence of pest and diseases	0	0	0	0	16	94.12	1	100	17	50
4	Inadequacy of irrigation water	0	0	11	100	17	100	1	100	29	85.29
5	High cost of Fertilizers and plant protection chemicals	0	0	11	100	17	100	1	100	29	85.29
6	High rate of interest on credit	0	0	11	100	17	100	1	100	29	85.29
7	Low price for the agricultural commodities	0	0	11	100	17	100	1	100	29	85.29
8	Lack of marketing facilities in the area	0	0	8	72.73	17	100	1	100	26	76.47
9	Inadequate extension services	0	0	5	45.45	15	88.24	1	100	21	61.76
10	Lack of transport for safe transport of the Agril produce to the market.	0	0	11	100	17	100	1	100	29	85.29
11	Less rainfall	0	0	1	9.09	1	5.88	0	0	2	5.88
12	Source of Agri-technology information	0	0	0	0	1	5.88	0	0	1	2.94

Farming constraints: The data regarding farming constraints experienced by households in Yadgir Rf Micro watershed is presented in Table 60. The results indicated that, lower fertility status of the soil was the constraint experienced by (94.12 %) per cent of the households, wild animal menace on farm field (85.29%), frequent incidence of pest

and diseases (50.00%), inadequacy of irrigation water (85.29%), high cost of fertilizers and plant protection chemicals (85.29%), high rate of interest on credit (85.29%), low price for the agricultural commodities (85.29 %), lack of marketing facilities in the area (76.47%), inadequate extension services (61.76 %), lack of transport for safe transport of the agricultural produce to the market (85.29%), less rainfall (5.88%), source of agri-technology information (Newspaper/Tv/Mobile) (2.94%).

SUMMARY AND IMPLICATIONS

In order to assess the socio-economic condition of the farmers in the watershed 34 households located in the micro watershed were interviewed for the survey. The study was conducted in Yadgir Rf micro-watershed (Ramasamudram sub-watershed, Yadgir taluk & District) is located at North latitude 16⁰ 49' 29.862" and 16⁰ 47' 10.021" and East longitude 77⁰ 15' 52.647" and 77⁰ 14' 56.958" covering an area of about 533.85 ha bounded by under Ramsamudra, Ashinala and Arakera. K Villages.

Socio-economic analysis of Yadgir Rf micro watersheds of Ramasamudram sub-watershed, Yadgir taluk & District indicated that, out of the total sample of 34 total respondents, 11 (32.35 %) were marginal, 17 (50.00%) were small, 1 (2.94 %) were semi medium and 5 landless farmers were also interviewed for the survey. The population characteristics of households indicated that, there were 93 (55.03%) men and 76 (44.97 %) were women. The average population of landless was 4, marginal farmers were 6.45, small farmers were 4.29 and semi medium farmers were 5.0. Majority of the respondents (38.46%) were in the age group of 16-35 years.

Education level of the sample households indicated that, there were 44.97 per cent illiterates, 50.89 per cent pre university education and 8.28 per cent attained graduation. About, 52.94 per cent of household heads practicing agriculture and 29.41 per cent of the household heads were engaged as agricultural labourers. Agriculture was the major occupation for 33.73 per cent of the household members. In the study area, 61.76 per cent of the households possess katcha house and 38.24 per cent possess pucca house. The durable assets owned by the households showed that, 76.47 per cent possess TV, 64.71 per cent possess mixer grinder, 85.29 per cent possess mobile phones and 44.12 per cent possess motor cycles.

Farm implements owned by the households indicated that, 20.59 per cent of the households possess plough, 2.94 per cent possess tractor, 2.94 per cent possess bullock cart and 11.76 per cent possess sprayer. Regarding livestock possession by the households, 2.94 per cent possess local cow. The average labour availability in the study area showed that, own labour men available in the micro watershed was 1.64, women available in the micro watershed was 1.36, hired labour (men) available was 8.86 and hired labour (women) available was 13.96. In the study area, about 1.18 per cent of the respondents migrated from the micro watershed in search of jobs with an average distance of 525.00 kms for about 9.00 months.

Out of the total land holding of the sample respondents 100.00 per cent (33.95 ha) of the area is under dry condition. There were 1.00 live bore wells and 1.00 dry bore wells among the sampled households. Bore well was the major source of irrigation for 2.94 per cent of the households. The major crops grown by sample farmers are Cotton, Redgram,

Maize, Groundnut and Jowar and cropping intensity was recorded as 100.00 per cent. Out of the sample households 94.12 percent possessed bank account and 61.76 per cent of them have savings in the account. About 64.71 per cent of the respondents borrowed credit from various sources. Among the credit borrowed by households, 18.18 per cent have borrowed loan from commercial banks and 59.09 per cent from co-operative/Grameena bank. Majority of the respondents (100.00%) have borrowed loan for agriculture purpose. Regarding the opinion on institutional sources of credit, 100.00 per cent of the households opined that credit helped to perform timely agricultural operations.

The per hectare cost of cultivation for Cotton, Redgram, Maize, Groundnut and Jowar was Rs.34406.70, 54597.19, 15086.15, 70279.00 and 19949.72 with benefit cost ratio of 1:1.40, 1: 0.70, 1: 2.50, 1: 0.60 and 1:1.00 respectively. Further, 11.76 per cent of the households opined that dry fodder was adequate.

The average annual gross income of the farmers was Rs. 73020.59 in micro-watershed, of which Rs. 39491.18 comes from agriculture. Sampled households have grown 10 horticulture trees and 56 forestry trees together in the fields and back yards. About 32.35 per cent of the households shown interest to cultivate horticultural crops.

Households have an average investment capacity of Rs. 735.29 for land development. Source of funds for additional investment is concerned, 32.35 per cent depends on bank loan for land development activities. Regarding marketing channels, 85.29 per cent of the households have sold agricultural produce to the local/village merchants. Further, 32.35 per cent of the households have used tractor for the transport of agriculture commodity. Majority of the farmers (35.29%) have experienced soil and water erosion problems in the watershed and 82.35 per cent of the households were interested towards soil testing.

Fire was the major source of fuel for domestic use for 55.88 per cent of the households and 44.12 per cent households has LPG connection. Piped supply was the major source for drinking water for 94.12 per cent of the households. Electricity was the major source of light for 100.00 per cent of the households. In the study area, 29.41 per cent of the households possess toilet facility. Regarding possession of PDS card, 97.06 per cent of the households possessed BPL card, 2.94 per cent of the household's possessed APL card.

Households opined that, the requirement of cereals (58.82%), pulses (55.88%) and oilseeds (5.88%) are adequate for consumption. Farming constraints experienced by households in the micro watersheds were lower fertility status of the soil (94.12%) wild animal menace on farm field (85.29%), frequent incidence of pest and diseases (50.00%), inadequacy of irrigation water (85.29%), high cost of fertilizers and plant protection chemicals (85.29%), high rate of interest on credit (85.29%), low price for the agricultural commodities (85.29%), lack of marketing facilities in the area (76.47%), inadequate

extension services (61.76%), lack of transport for safe transport of the agricultural produce to the market (85.29%), Less rainfall (5.88%) and Source of Agri-technology information (Newspaper/ TV/Mobile) (2.94%).

Implications of the survey

- ✓ Result indicated that, there were 44.97 per cent were illiterate hence, extension methodologies such as demonstration, street play, drama, video shows will be effective in dissemination of the technologies in the micro watershed.
- ✓ The data indicate that, 61.76 per cent of the households possess katcha house. Hence, the development department while implementing the watershed plan should focus on agriculture to enhance the productivity of major crops in the area to increase the income of the farmers.
- ✓ Results indicated that the local institutional participation of the household members in the micro watershed is minimal hence, activities like membership campaign, awareness creation about the benefits of membership in local institutions and strengths of organized groups must be conveyed.
- ✓ Majority of the households in the watershed have experience in use of mobile phones, and television hence, these mass media can be effectively utilized for transfer of technology as well as for information dissemination.
- ✓ The farm machinery/implement possession in the micro watershed was found to be minimum the reasons may lack of knowledge or lack of financial ability which can be addressed through training on use of different farm implements, providing information on different sources of finance for purchase of farm implements.
- ✓ The possession of livestock such as crossbred cow found is less hence, farmers must be made aware of the benefits of crossbred cow in increased milk production.
- ✓ The possession of livestock such as sheep, goat and poultry was found to be low hence, farmers may be informed the role of subsidiary enterprises in enhancing the income and information on financial support for subsidiary activities.
- ✓ The data indicate that, job/work was the reason for all the migrants hence, farmers may be trained on profitable agriculture or self employment such as animal husbandry, plate making, sheep rearing, goat rearing, rabbit rearing with suitable information on sources of financial support.
- ✓ The results indicate that there was a change in quality of life due to migration hence, the developmental departments should take actions to arrest migration and to improve the quality of the life in rural areas.
- ✓ Households possess 33.95ha (100.00 %) of dry land and the availability of the dryland agricultural technologies such as short duration crops, high yielding drought resistance crop varieties, drip irrigation technology and subsidy information will be helpful for the farmers to enhance the productivity of land and as well as farmers income.
- ✓ Few of the bore well in micro watershed found non functional hence, farmers may be trained on possibility of bore well rejuvenation.

- ✓ Bore well was major source of irrigation for 2.94 per cent of the households. hence, in order to increase the area under irrigation as well as to increase the water use efficiency farmers may trained on drip irrigation and provide the information on subsidy for drip irrigation equipment's along with the information on different agencies which provides the financial assistance for drip irrigation.
- ✓ The cropping intensity in the micro watershed was found to be (100.00 %) hence, care must be taken by the implementing agency to bring uncultivated land into cultivation through suitable measures.
- ✓ The results indicated the non availability of both green and dry fodder throughout the year hence, fodder development activities can be taken up in the micro watershed.
- ✓ The average annual gross income of the households Rs.39491.18 from agriculture and Rs. 15882.35 from wages and. Agriculture was found to be the major source of income for households hence; the development activities should focus on productivity enhancement, marketing arrangements and agricultural technology dissemination to have a direct impact on the farmers.
- ✓ The cultivation of forest species is found minimal hence, information and production technology related to agro-forestry and integrated farming system.
- ✓ The data indicated that, 35.29 per cent of the households have experienced soil and water erosion problems. Hence, those farmers who reported the soil and water erosion problems may be given attention while implementation of the watershed development plan.
- ✓ The data indicated that, 82.35 per cent of the households have interest in soil testing hence, farmers must be provided with the information on various institutions which are involved in soil testing for the benefit of the farmers.
- ✓ Except summer ploughing the adoption of other soil and water conservation structures is minimum hence, the farmers in the micro watershed should be sensitized on the use of different conservation structures for soil water conservation.
- ✓ Cereals and pulses found be adequate for per cent of the households respectively hence, farm households and the farm women must be trained on importance of balanced nutrition and role of vegetable, milk, egg, meat in balanced diet.
- ✓ Lower fertility status of the soil (94.12%), wild animal menace on farm field (85.29%), frequent incidence of pest and diseases (50.00%), high cost of fertilizers and plant protection chemicals (85.29%), high rate of interest on credit (85.29%), low price for the agricultural commodities (85.29%), lack of marketing facilities in the area (76.47%), inadequate extension services (61.76%), lack of transport for safe transport of the agricultural produce to the market (85.29%) were the major farming constraints experienced hence, these constraints must be addressed immediately for the welfare of the farmers. Awareness to be created among the farmers to approach nearest KVKs/RSKs and other developmental departments for technical and for subsidized inputs and utilize the well established regulated markets, approaching the contract firms, direct markets to avoid the involvement of middlemen.