ICAR-NBSS&LUP Sujala MWS Publ.563

हर कदम, हर डगर
किसानों का हमसफर
भारतीय कृषि अनुसंधान परिषद

Agrisearch with a human touch

LAND RESOURCE INVENTORY AND SOCIO-ECONOMIC STATUS OF FARM HOUSEHOLDS FOR WATERSHED PLANNING AND DEVELOPMENT

LAKSHMAPURA-2 (4D4A1U1c) MICROWATERSHED

Koppal Taluk & District, Karnataka

Karnataka Watershed Development Project – II

SUJALA – III

World Bank funded Project

THE WORLD BANK

ICAR – NATIONAL BUREAU OF SOIL SURVEY AND LAND USE PLANNING

ICAR - NBSS & LUP

WATERSHED DEVELOPMENT DEPARTMENT
GOVT. OF KARNATAKA, BANGALORE

About ICAR - NBSS&LUP

The ICAR-National Bureau of Soil Survey and Land Use Planning (ICAR-NBSS&LUP), Nagpur, a premier Institute of the Indian Council of Agricultural Research (ICAR), was set up during 1976 with the objective to prepare soil resource maps at national, state and district levels and to provide research inputs in soil resource mapping and its applications, land evaluation, land use planning, land resource management, and database management using GIS for optimising land use on different kinds of soils in the country.

The Bureau has been engaged in carrying out soil resource survey, agro-ecological and soil degradation mapping at the country, state and district levels for qualitative assessment and monitoring the soil health towards viable land use planning. The research activities have resulted in identifying the soil potentials and problems, and the various applications of the soil surveys with the ultimate objective of sustainable agricultural development. The Bureau has the mandate to correlate and classify soils of the country and maintain a National Register of all the established soil series. The Institute is also imparting in-service training to staff of the soil survey agencies in the area of soil survey, land evaluation and soil survey interpretations for land use planning. The Bureau in collaboration with Panjabrao Krishi Vidyapeeth, Akola is running post-graduate teaching and research programme in land resource management, leading to M.Sc. and Ph.D. degrees.

Citation: Rajendra Hegde, Ramesh Kumar, S.C., K.V. Niranjana, S. Srinivas, M.Lalitha, B.A. Dhanorkar, R.S. Reddy, and S.K. Singh (2019). “Land Resource Inventory and socio-economic status of farm households for Watershed Planning and Development Laksmapura-2 (4D4A1U1c) Microwatershed, Koppal Taluk and District, Karnataka”, ICAR-NBSS&LUP Sujala MWS Publ. 563, ICAR – NBSS & LUP, RC, Bangalore. p.159 & 28.

TO OBTAIN COPIES,

Please write to:
Director, ICAR - NBSS & LUP,

Amaravati Road, NAGPUR - 440 033, India

Phone : (0712) 2500386, 2500664, 2500545 (O)

Telefax : 0712-2522534

E-Mail : director@nbsslup.ernet.in

Website URL : nbsslup.in

Or

Head, Regional Centre, ICAR - NBSS&LUP, Hebbal, Bangalore - 560 024

Phone : (080) 23412242, 23510350 (O)

Telefax : 080-23510350

E-Mail : nbssrcb@gmail.com

LAND RESOURCE INVENTORY AND SOCIO-ECONOMIC STATUS OF FARM HOUSEHOLDS FOR WATERSHED PLANNING AND DEVELOPMENT

LAKSHMAPURA-2 (4D4A1U1c) MICROWATERSHED

Koppal Taluk & District, Karnataka

Karnataka Watershed Development Project – II

Sujala-III

World Bank funded Project

ICAR – NATIONAL BUREAU OF SOIL SURVEY AND LAND USE PLANNING

WATERSHED DEVELOPMENT DEPARTMENT, GOVT. OF KARNATAKA, BANGALORE

PREFACE

In Karnataka, as in other Indian States, the livelihoods of rural people are intertwined with farming pursuits. Thechallenges in agriculture are seriously threatening the livelihood of a large number of farmers as they have been practicing farming in contextual factors beyond their control. Climatic factors are the most important ones and have become much more significant in recent times due to rapid climate changes induced by intensive anthropogenic activities affecting our ecosystem in multiple ways. Climate change has become the reality, it is happening and efforts to evolve and demonstrate climate resilient technologies have become essential. Due to the already over stressed scenario of agrarian sector, the climate change is resulting in manifold increase in the complexities, pushing the rural mass to face more and more unpredictable situations. The rising temperatures and unpredictable rainfall patterns are going to test seriously the informed decisions farmers have to make in order to survive in farming and sustain their livelihood.

It is generally recognized that impacts of climate change shall not be uniform across the globe. It is said that impact of climate change is more severe in South Asia. Based on the analysis of meteorological data, it is predicted that in India, there will be upward trend in mean temperature, downward trend in relative humidity, annual rainfall and number of wet days in a year. Also, in general, phenomena like erratic monsoon, spread of tropical diseases, rise in sea levels, changes in availability of fresh water, frequent floods, droughts, heat waves, storms and hurricanes are predicted. Each one of these adverse situations are already being experienced in various parts of India and also at the global level. Decline in agricultural productivity of small and marginal farmers becoming more vulnerable is already witnessed.

In Karnataka, more than 60 per cent of the population live in rural areas and depend on agriculture and allied activities for their livelihood. Though the state has achieved significant progress in increasing the yield of many crops, there is tremendous pressure on the land resources due to the growing and competing demands of various land uses. This is reflected in the alarming rate of land degradation observed. Already more than 50 per cent of the area is affected by various forms of degradation. If this trend continues, the sustainability of the fragile ecosystem will be badly affected. The adverse effects of change in the climatic factors are putting additional stress on the land resources and the farmers dependent on this.

The natural resources (land, water and vegetation) of the state need adequate and constant care and management, backed by site-specific technological interventions and investments particularly by the government. Detailed database pertaining to the nature of

the land resources, their constraints, inherent potentials and suitability for various land based rural enterprises, crops and other uses is a prerequisite for preparing location-specific action plans, which are in tune with the inherent capability of the resources. Any effort to evolve climate resilient technologies has to be based on the baseline scientific database. Then only one can expect effective implementation of climate resilient technologies, monitor the progress, make essential review of the strategy, and finally evaluate the effectiveness of the implemented programs. The information available at present on the land resources of the state are of general nature and useful only for general purpose planning. Since the need of the hour is to have site-specific information suitable for farm level planning and detailed characterization and delineation of the existing land resources of an area into similar management units is the only option.

ICAR-NBSS&LUP, Regional Centre, Bangalore has taken up a project sponsored by the Karnataka Watershed Development Project-II, (Sujala-III), Government of Karnataka funded by the World Bank under Component -1 Land Resource Inventry. This study was taken up to demonstrate the utility of such a database in reviewing, monitoring and evaluating all the land based watershed development programs on a scientific footing. To meet the requirements of various land use planners at grassroots level, the present study on "Land Resource Inventory and Socio-Economic Status of Farm Households for Watershed Planning and Development of for Lakshmapura-2 microwatershed in Koppal Taluk and District, Karnataka" for integrated development was taken up in collaboration with the State Agricutural Universities, IISC, KSRSAC, KSNDMC as Consortia partners. The project provides detailed land resource information at cadastral level (1:7920 scale) for all the plots and socio-economic status of farm households covering thirty per cent farmers randomely selected representing landed and landless class of farmers in the micro-watershed. The project report with the accompanying maps for the microwatershed will provide required detailed database for evolving effective land use plan, alternative land use options and conservation plans for the planners, administrators, agricutural extention personnel, KVK officials, developmental departments and other land users to manage the land resources in a sustainable manner.

It is hoped that this database will be useful to the planners, administrators and developmental agencies working in the area in not only for formulating location specific developmental schemes but also for their effective monitoring at the village/watershed level.

Nagpur
Date:18-11-2019

S.K. SINGH
Director, ICAR - NBSS&LUP, Nagpur

Contributors

Dr. Rajendra Hegde Principal Scientist, Head & Project Leader, Sujala-III Project ICAR-NBSS&LUP, Regional Centre, Bangalore	**Dr. S.K.Singh** Director, ICAR-NBSS&LUP Coordinator, Sujala-III Project Nagpur
Soil Survey, Mapping & Report Preparation	
Dr. K.V. Niranjana	Sh. R.S. Reddy
Dr. B.A. Dhanorkar	Smt. Chaitra, S.P.
	Dr. Gopali Bardhan
	Mr. Somashekar T.N
	Ms. Arpitha G.M
	Dr. Mahendra kumar M.B
Field Work	
Sh. C. Bache Gowda	Sh. Mayur Patil
Sh. Somashekar	Sh. Arun Kumar, S.
Sh. M. Jayaramaiah	Sh. Sunil Raj
	Sh. Yogesh Kumar, B.
	Sh. Vikas, N.K.
	Sh. Arun Kumar, S.G.
	Sh. Umesh Jadiyappa Madolli
	Sh. Praveen Kumar P. Achalkar
	Sh. Veerabhadraswamy
	Sh. Vinay
	Sh. Shankarappa, K.
	Sh. Lankesh, R.S.
	Sh. Appanna B. Hattigoudar
	Sh. Maharudra
GIS Work	
Dr. S.Srinivas	Sh. A.G.Devendra Prasad
Sh. D.H.Venkatesh	Sh. Abhijith Sastry, N.S.
Smt. K.Sujatha	Smt. Shyla, B.
Smt. K.V.Archana	Smt. Swetha ,K.
Sh. N.Maddileti	Ms. Vidya, P.C.
	Sh. Deepak, M.J.
	Smt. K.Karunya Lakshmi
	Ms. Seema, K.V.

Laboratory Analysis	
Dr. M. Lalitha	Sh. Vindhya, N.G.
Smt. Arti Koyal	Ms. P. Pavanakumari, P.
Smt. Parvathy, S.	Ms. Rashmi, N.
	Ms. Leelavathy, K.U.
	Smt. Usha Kiran, G.
Socio-Economic Analysis	
Dr. S.C. Ramesh Kumar	Sh. M.K. Prakashanaik
	Ms. Karuna V. Kulkarni
	Mrs. Sowmya A.N
	Sh. Vinod R
	Sh. Basavaraja
	Sh. Vijay Kumar Lamani
	Ms. Sowmya K.B
	Mrs. Prathibha, D.G
	Sh. Rajendra,D
Soil & Water Conservation	
Sh. Sunil P. Maske	
Watershed Development Department, GoK, Bangalore	
Sh. Prabhash Chandra Ray, IFS Project Director & Commissioner, WDD	Dr. A. Natarajan NRM Consultant, Sujala-III Project
Sh. A. Padmaya Naik, Director (In-Charge) Executive Director, KWDP-II, Sujala-III, WDD	

PART-A

LAND RESOURCE INVENTORY

Contents

Preface		
Contributors		
Executive Summary		
Chapter 1	Introduction	1
Chapter 2	Geographical Setting	3
2.1	Location and Extent	3
2.2	Geology	3
2.3	Physiography	4
2.4	Drainage	5
2.5	Climate	5
2.6	Natural Vegetation	6
2.7	Land Utilization	7
Chapter 3	Survey Methodology	11
3.1	Base maps	11
3.2	Image Interpretation for Physiography	11
3.3	Field Investigation	14
3.4	Soil mapping	16
3.5	Land Management Units	17
3.6	Laboratory Characterization	17
Chapter 4	The Soils	25
4.1	Soils of Granite Gneiss Landscape	25
4.2	Soils of Alluvial Landscape	35
Chapter 5	Interpretation for Land Resource Management	59
5.1	Land Capability Classification	59
5.2	Soil Depth	61
5.3	Surface Soil Texture	62
5.4	Soil Gravelliness	63
5.5	Available Water Capacity	64
5.6	Soil Slope	65
5.7	Soil Erosion	66
Chapter 6	Fertility Status	69
6.1	Soil Reaction (pH)	69
6.2	Electrical Conductivity (EC)	69
6.3	Organic Carbon (OC)	69
6.4	Available Phosphorus	71
6.5	Available Potassium	71
6.6	Available Sulphur	71
6.7	Available Boron	72
6.8	Available Iron	72
6.9	Available Manganese	75
6.10	Available Copper	75
6.11	Available Zinc	75

Chapter 7	Land Suitability for Major Crops	77
7.1	Land suitability for Sorghum	77
7.2	Land suitability for Maize	78
7.3	Land suitability for Bajra	79
7.4	Land suitability for Redgram	80
7.5	Land suitabilitry for Bengalgram	81
7.6	Land suitability for Groundnut	82
7.7	Land suitability for Sunflower	83
7.8	Land suitability for Cotton	84
7.9	Land suitability for Chilli	85
7.10	Land suitability for Tomato	86
7.11	Land suitability for Brinjal	87
7.12	Land suitability for Onion	88
7.13	Land suitability for Bhendi	89
7.14	Land suitability for Drumstick	90
7.15	Land suitability for Mulberry	91
7.16	Land suitability for Mango	92
7.17	Land Suitability for Sapota	93
7.18	Land suitability for Pomegranate	94
7.19	Land suitability for Guava	95
7.20	Land Suitability for Jackfruit	96
7.21	Land Suitability for Jamun	97
7.22	Land Suitability for Musambi	98
7.23	Land Suitability for Lime	99
7.24	Land Suitability for Cashew	100
7.25	Land Suitability for Custard apple	101
7.26	Land suitability for Amla	102
7.27	Land suitability for Tamarind	103
7.28	Land suitability for Marigold	104
7.29	Land suitability for Chrysanthemum	105
7.30	Land suitability for Jasmine	106
7.31	Land suitability for Crossandra	107
7.32	Land Management Units	142
7.33	Proposed Crop Plan	143
Chapter 8	Soil Health Management	147
Chapter 9	Soil and Water conservation Treatment Plan	151
9.1	Treatment Plan	151
9.2	Recommended Soil and Water Conservation measures	155
9.3	Greening of microwatershed	156
	References	159
	Appendix I	I-VI
	Appendix II	VII-XII
	Appendix III	XIII-XIX

LIST OF TABLES

2.1	Mean Monthly Rainfall, PET, 1/2 PET at Koppal Taluk and District	5
2.2	Land Utilization in Koppal District	7
3.1	Differentiating Characteristics used for Identifying Soil Series	15
3.2	Soil map unit description of Lakshmapura-2 microwatershed	17
4.1	Physical and chemical characteristics of soil series identified in Lakshmapura-2 microwatershed	40
7.1	Soil-Site Characteristics of Lakshmapura-2 microwatershed	109
7.2	Land suitability for Sorghum	111
7.3	Land suitability for Maize	112
7.4	Land suitability for Bajra	113
7.5	Land suitability for Redgram	114
7.6	Land suitability for Bengalgram	115
7.7	Land suitability for Groundnut	116
7.8	Land suitability for Sunflower	117
7.9	Land suitability for Cotton	118
7.10	Land suitability for Chilli	119
7.11	Land suitability for Tomato	120
7.12	Land suitability for Brinjal	121
7.13	Land suitability for Onion	122
7.14	Land suitability for Bhendi	123
7.15	Land suitability for Drumstick	124
7.16	Land suitability for Mulberry	125
7.17	Land suitability for Mango	126
7.18	Land Suitability for Sapota	127
7.19	Land suitability for Pomegranate	128
7.20	Land suitability for Guava	129
7.21	Land suitability for Jackfruit	130
7.22	Land suitability for Jamun	131
7.23	Land Suitability for Musambi	132
7.24	Land Suitability for Lime	133

7.25	Land Suitability for Cashew	134
7.26	Land Suitability for Custard apple	135
7.27	Land Suitability for Amla	136
7.28	Land Suitability for Tamarind	137
7.29	Land Suitability for Marigold	138
7.30	Land Suitability for Chrysanthemum	139
7.31	Land suitability for Jasmine	140
7.32	Land suitability for Crossandra	141
7.33	Proposed Crop Plan for Lakshmapura-2 Microwatershed	144

LIST OF FIGURES

2.1	Location map of Lakshmapura-2 Microwatershed	3
2.2a	Granite and granite gneiss rocks	4
2.2b	Alluvial rocks	4
2.3	Rainfall distribution in Koppal Taluk, Koppal District	6
2.4	Natural vegetation of Lakshmapura-2 microwatershed	6
2.5	Different crops and cropping systems in Lakshmapura-2 Microwatershed	7
2.6	Current Land use – Lakshmapura-2 Microwatershed	9
2.7	Location of Wells- Lakshmapura-2 Microwatershed	9
3.1	Scanned and Digitized Cadastral map of Lakshmapura-2 Microwatershed	13
3.2	Satellite image of Lakshmapura-2 Microwatershed	13
3.3	Cadastral map overlaid on IRS PAN+LISS IV merged imagery of Lakshmapura-2 Microwatershed	14
3.4	Location of profiles in a transect	15
3.5	Soil phase or management units of Lakshmapura-2 Microwatershed	23
5.1	Land Capability Classification of Lakshmapura-2 Microwatershed	60
5.2	Soil Depth map of Lakshmapura-2 Microwatershed	61
5.3	Surface Soil Texture map of Lakshmapura-2 Microwatershed	63
5.4	Soil Gravelliness map of Lakshmapura-2 Microwatershed	64
5.5	Soil Available Water Capacity map of Lakshmapura-2 Microwatershed	65
5.6	Soil Slope map of Lakshmapura-2 Microwatershed	66
5.7	Soil Erosion map of Lakshmapura-2 Microwatershed	67
6.1	Soil Reaction (pH) map of Lakshmapura-2 Microwatershed	70
6.2	Electrical Conductivity (EC) map of Lakshmapura-2 Microwatershed	70
6.3	Soil Organic Carbon (OC) map of Lakshmapura-2 Microwatershed	71
6.4	Soil Available Phosphorus map of Lakshmapura-2 Microwatershed	72
6.5	Soil Available Potassium map of Lakshmapura-2 Microwatershed	73
6.6	Soil Available Sulphur map of Lakshmapura-2 Microwatershed	73
6.7	Soil Available Boron map of Lakshmapura-2 Microwatershed	74
6.8	Soil Available Iron map of Lakshmapura-2 Microwatershed	74
6.9	Soil Available Manganese map of Lakshmapura-2 Microwatershed	75

6.10	Soil Available Copper map of Lakshmapura-2 Microwatershed	76
6.11	Soil Available Zinc map of Lakshmapura-2 Microwatershed	76
7.1	Land suitability map of Sorghum	78
7.2	Land suitability map of Maize	79
7.3	Land suitability map of Bajra	80
7.4	Land suitability map of Redgram	81
7.5	Land suitability map of Bengalgram	82
7.6	Land suitability map of Groundnut	83
7.7	Land suitability map of Sunflower	84
7.8	Land suitability map of Cotton	85
7.9	Land suitability map of Chilli	86
7.10	Land suitability map of Tomato	87
7.11	Land suitability map of Brinjal	88
7.12	Land suitability map of Onion	89
7.13	Land suitability map of Bhendi	90
7.14	Land suitability map of Drumstick	91
7.15	Land suitability map of Mulberry	92
7.16	Land suitability map of Mango	93
7.17	Land Suitability map of Sapota	94
7.18	Land suitability for Pomegranate	95
7.19	Land suitability map of Guava	96
7.20	Land Suitability map of Jackfruit	97
7.21	Land Suitability map of Jamun	98
7.22	Land Suitability map of Musambi	99
7.23	Land Suitability map of Lime	100
7.24	Land Suitability map of Cashew	101
7.25	Land Suitability map of Custard apple	102
7.26	Land suitability map of Amla	103
7.27	Land suitability map of Tamarind	104
7.28	Land suitability map of Marigold	105
7.29	Land suitability map of Chrysanthemum	106
7.30	Land suitability map of Jasmine	107
7.31	Land suitability map of Crossandra	108
7.32	Land Management Units map of Lakshmapura-2 microwatershed	143
9.1	Soil and water conservation map of Lakshmapura-2 microwatershed	156

EXECUTIVE SUMMARY

The land resource inventory of Lakshmapura-2 microwatershed was conducted using village cadastral maps and IRS satellite imagery on 1:7920 scale. The false colour composites of IRS imagery were interpreted for physiography and these physiographic delineations were used as base for mapping soils. The soils were studied in several transects and a soil map was prepared with phases of soil series as mapping units. Random checks were made all over the area outside the transects to confirm and validate the soil map unit boundaries. The soil map shows the geographic distribution and extent, characteristics, classification, behavior and use potentials of the soils in the Microwatershed.

The present study covers an area of 460 ha in Koppal taluk and district, Karnataka. The climate is semiarid and categorized as drought - prone with an average annual rainfall of 662 mm, of which about 424 mm is received during south –west monsoon, 161 mm during north-east and the remaining 77 mm during the rest of the year. An area of about 98 per cent is covered by soil, <1 per cent by railways and 2 per cent by water bodies settlements and others. The salient findings from the land resource inventory are summarized briefly below.

- *The soils belong to 21 soil series and 33 soil phases (management units) and 9 land management units.*
- *The length of crop growing period is <90 days and starts from 2nd week of August to 2nd week of November.*
- *From the master soil map, several interpretative and thematic maps like land capability, soil depth, surface soil texture, soil gravelliness, available water capacity, soil slope and soil erosion were generated.*
- *Soil fertility status maps for macro and micronutrients were generated based on the surface soil samples collected at every 320 m grid interval.*
- *Land suitability for growing 31 major agricultural and horticultural crops were assessed and maps showing the degree of suitability along with constraints were generated.*
- *Entire area is suitable for agriculture.*
- *About 15 per cent of the soils are shallow (25-50cm), 11 per cent of the soils are moderately shallow (50-75 cm), 35 per cent moderately deep (75- 100 cm) and 37 per cent is deep to very deep (100->150cm) soils.*
- *About 14 per cent loamy (sandy clay loam) and 85 per cent has clayey (sandy clay and clay) soils at the surface.*
- *About 46 per cent of the area has non-gravelly (<15%), 44 per cent gravelly (15-35%) and 9 per cent is very gravelly (35-60%) soils.*
- *With respect to available water capacity 16 per cent of the area has very low (<50mm/m), 34 per cent of the area has low (51-100 mm/m), 22 per cent medium*

(101-150 mm/m) and 27 per cent area is high to very high (151->200mm/m) in available water capacity.

- *An area of about 2 per cent has nearly level (0-1%) and 96 per cent has very gently sloping (1-3%) lands.*
- *An area of about 19 per cent is slightly eroded (e1) and 79 per cent is moderately eroded (e2).*
- *An area of about 87 per cent is slightly alkaline (pH 7.3 to 7.8) and 12 per cent is strongly alkaline (pH 8.4-9.0).*
- *The Electrical Conductivity (EC) of the soils are <2 dsm^{-1} indicating that soils are non saline.*
- *Organic carbon is low (<0.5%) in 6 per cent, medium (0.5-0.75%) in 51 per cent and 42 per cent is high (>0.75%).*
- *Available phosphorus is low (<23 kg/ha) in 22 per cent, medium (<23 kg/ha) in 68 per cent and high (>57 kg/ha) in 8 per cent area of the soils.*
- *Available potassium is medium (145-337 kg/ha) in 49 per cent and high (>337 kg/ha) in 50 per cent area of the soils.*
- *Available sulphur is low (<10 ppm) in 78 per cent and medium (10-20 ppm) in 20 per cent area of the soils.*
- *Available boron is low (<0.5 ppm) in 19 per cent and medium (0.5-1.0 ppm) in 80 per cent area of the microwatershed.*
- *Available iron is deficient (<4.5ppm) in 83 per cent and sufficient (>4.5 ppm) in 16 per cent area of the microwatershed.*
- *Available zinc is deficient (<0.6 ppm) in the entire area of the microwatershed.*
- *Available manganese and copper are sufficient in the entire area.*
- *The land suitability for 31 major agricultural and horticultural crops grown in the microwatershed was assessed and the areas that are highly suitable (class S1) and moderately suitable (class S2) are given below. It is however to be noted that a given soil may be suitable for various crops but what specific crop to be grown may be decided by the farmer looking to his capacity to invest on various inputs, marketing infrastructure, market price, and finally the demand and supply position.*

Land suitability for various crops in the microwatershed

Crop	Suitability Area in ha (%)		Crop	Suitability Area in ha (%)	
	Highly suitable (S1)	Moderately suitable (S2)		Highly suitable (S1)	Moderately suitable (S2)
Sorghum	*18(4)*	*313(68)*	*Sapota*	*46(10)*	*127(28)*
Maize	-	*331(72)*	*Pomegranate*	*46(10)*	*263(57)*
Bajra	*47(10)*	*295(64)*	*Guava*	*46(10)*	*127(28)*
Redgram	*18(4)*	*283(62)*	*Jackfruit*	*46(10)*	*127(28)*
Bengal gram	-	*136(30)*	*Jamun*	*18(4)*	*47(10)*
Groundnut	-	*188(41)*	*Musambi*	*46(10)*	*263(57)*
Sunflower	*18(4)*	*283(62)*	*Lime*	*46(10)*	*263(57)*
Cotton	-	*330(72)*	*Cashew*	*46(10)*	*127(28)*
Chilli	*18(4)*	*312(68)*	*Custard apple*	*47(10)*	*295(64)*
Tomato	*18(4)*	*176(38)*	*Amla*	*47(10)*	*159(34)*
Brinjal	*18(4)*	*176(38)*	*Tamarind*	*18(4)*	*47(10)*
Onion	-	*194(42)*	*Marigold*	*18(4)*	*312(68)*
Bhendi	*18(4)*	*312(68)*	*Chrysanthemum*	*18(4)*	*312 (68)*
Drumstick	*46(10)*	*138(30)*	*Jasmine*	*18(4)*	*176(38)*
Mulberry	*33(7)*	*140(30)*	*Crossandra*	*18(4)*	*176(38)*
Mango	*18(4)*	*47(10)*	-	-	-

- *Apart from the individual crop suitability, a proposed crop plan has been prepared for the 9 identified LMUs by considering only the highly and moderately suitable lands for different crops and cropping systems with food, fodder, fibre and other horticulture crops.*
- *Maintaining soil-health is vital for crop production and conserve soil and land resource base for maintaining ecological balance and to mitigate climate change. For this, several ameliorative measures have been suggested to these problematic soils like saline/alkali, highly eroded, sandy soils etc.,*
- *Soil and water conservation and drainage line treatment plans have been prepared that would help in identifying the sites to be treated and also the type of structures required.*
- *As part of the greening programme, several tree species have been suggested to be planted in marginal and submarginal lands, field bunds and also in the hillocks, mounds and ridges. That would help in supplementing the farm income, provide fodder and fuel, and generate lot of biomass which in turn would help in maintaining the ecological balance and contribute to mitigating the climate change.*

Chapter 1

INTRODUCTION

Land is a scarce resource and basic unit for any material production. It can support the needs of the growing population, provided they use the land in a rational and judicious manner. But what is happening in many areas of the state is a cause for concern to everyone involved in the management of land resources at the grassroots level. The area available for agriculture is about 51 per cent of the total area and more than 60 per cent of the people are still dependant on agriculture for their livelihood. The limited land area is under severe stress and strain due to increasing population pressure and competing demands of various land uses. Due to this, every year there is significant diversion of farm lands and water resources for non-agricultural purposes. Apart from this, due to lack of interest in farmers for farming, large tracts of cultivable lands are turning into fallows in many areas and this trend is continuing at an alarming rate.

Further, land degradation has emerged as a serious problem which has already affected about 38 lakh ha of cultivated area in the state. Soil erosion alone has degraded about 35 lakh ha. Almost all the uncultivated areas are facing various degrees of degradation, particularly soil erosion. Salinity and alkalinity has emerged as a major problem in more than 3.5 lakh ha in the irrigated areas of the state. Nutrient depletion and declining factor productivity is common in both rainfed and irrigated areas. The degradation is continuing at an alarming rate and there appears to be no systematic effort among the stakeholders to contain this process. In recent times, an aberration of weather due to climate change phenomenon has added another dimension leading to unpredictable situations to be tackled by the farmers.

In this critical juncture, the challenge before us is not only to increase the productivity per unit area which is steadily declining and showing a fatigue syndrome, but also to prevent or at least reduce the severity of degradation. If the situation is not reversed at the earliest, then the sustainability of the already fragile crop production system and the overall ecosystem will be badly affected in the state. The continued neglect and unscientific use of the resources for a long time has led to the situation observed at present in the state. It is a known fact and established beyond doubt by many studies in the past that the cause for all kinds of degradation is the neglect and irrational use of the land resources. Hence, there is urgent need to generate a detailed site-specific farm level database on various land resources for all the villages/watersheds in a time bound manner that would help to protect the valuable soil and land resources and also to stabilize the farm production.

Therefore, the land resource inventory required for farm level planning is the one which investigates not only the surface but also consider the other parameters which are critical for productivity *viz.*, soils, climate, water, minerals and rocks, topography, geology, hydrology, vegetation, crops, land use pattern, animal population, socio-

economic conditions, infrastructure, marketing facilities and various schemes and developmental works of the government etc. From the data collected at farm level, the specific problems and potentials of the area can be identified and highlighted, conservation measures required for the area can be planned on a scientific footing, suitability of the area for various uses can be worked out and finally viable and sustainable land use options suitable for each and every land holding can be prescribed.

The Land Resource Inventory is basically done for identifying potential and problem areas, developing sustainable land use plans, estimation of surface run off and water harvesting potential, preparation of soil and water conservation plans, land degradation/desertification etc. The Bureau is presently engaged in developing an LRI methodology using high resolution satellite remote sensing data and Digital Elevation Model (DEM) data to prepare Landscape Ecological Units (LEU) map representing agro-ecosystem as a whole. The LEU is preferred over landform as the base map for LRI. LEU is the assemblage of landform, slope and land use. An attempt was made to upscale the soil resource information from 1:250000 and 1:50000 scale to the LEU map in Goa and other states.

The land resource inventory aims to provide site-specific database for Lakshmapura-2 Microwatershed in Koppal Taluk, Koppal District, Karnataka State for the Karnataka Watershed Development Department. The database was generated by using cadastral map of the village as a base along with high resolution IRS LISS IV and Cartosat-1 merged satellite imagery. Later, an attempt will be made to uplink this LRI data generated at 1:7920 scale under Sujala-III Project to the proposed Landscape Ecological Units (LEUs) map.

The study was organized and executed by the ICAR- National Bureau of Soil Survey and Land Use Planning, Regional Centre, Bangalore under Generation of Land Resource Inventory Data Base Component-1 of the Sujala-III Project funded by the World Bank.

Chapter 2

GEOGRAPHICAL SETTING

2.1 Location and Extent

The Lakshmapura-2 micro-watershed is located in the central part of Karnataka in Koppal taluk and district (Fig 2.1). It lies between 15^0 35' and 15^0 38' North latitudes and 76^0 03' and 76^0 06' East longitudes, and covers an area of about 460 ha. It comprises parts of Lakamapura, Thalakalla and Halageri villages. It is about 19 km from Koppal town and is bounded by Halageri on the east and Thalakalla on the north and southwest and Lakamapura on the south and western side of the microwatershed.



Fig.2.1 Location map of Lakshmapura-2 Microwatershed

2.2 Geology

Major rock formations observed in the microwatershed are granite gneiss and alluvium (Fig.2.2 a and b). Granite gneisses are essentially pink to gray and are coarse to medium grained. They consist primarily of quartz, feldspar, biotite and hornblende. The

gray granite gneisses are highly weathered, fractured and fissured upto a depth of about 10 m. Dolerite dykes and quartz veins are common with variable width and found to occur in Lakshmapura-2 village. The thickness of the alluvium generally is limited to less than a meter, except in river valleys where it is very deep extending to tens of meters. Such soils are transported and represent paleo black soils originally formed at higher elevation, but now occupying river valleys.

Fig.2.2 a Granite and granite gneiss rocks

Fig.2.2 b Alluvium

2.3 Physiography

Physiographically, the area has been identified as Granite gneiss and Alluvial landscapes based on geology. The microwatershed area has been further divided into mounds/ridges, summits, side slopes and very gently sloping uplands and nearly level

plains based on slope and its relief features. The elevation ranges from 520 to 544 m in the gently sloping uplands. The mounds and ridges are mostly covered by rock outcrops.

2.4 Drainage

The area is drained by several small seasonal streams that join Hire *halla* and Chenna *halla* along its course. Though, the streams are not perennial, during rainy season they carry large quantities of rain water. The microwatershed has only few small tanks which are not able to store the water flowing during the rainy season. Due to this, the ground water recharge is very much affected in the villages. This is reflected in the failure of many bore wells in the villages. If the available rain water is properly harnessed by constructing tanks and recharge structures at appropriate places in the villages, then the drinking and irrigation needs of the area can be easily met. The drainage network is dendritic to sub parallel.

2.5 Climate

The district falls under semiarid tract of the state and is categorized as drought - prone with total annual rainfall of 662 mm (Table 2.1). Of this, a maximum of 424 mm precipitation is received during south–west monsoon period from June to September, north-east monsoon contributes about 161 mm and prevails from October to early December and the remaining 77 mm is received during the rest of the year. The winter season is from December to February. During April and May, the temperatures reach up to 45°C and in December and January, the temperatures will go down to 16°C. Rainfall distribution is shown in Figure 2.3. The average Potential Evapo Transpiration (PET) is 145 mm and varies from a low of 101 mm in December to 193 mm in the month of May. The PET is always higher than precipitation in all the months except in the month of September. Generally, the Length of crop Growing Period (LGP) is <90 days and starts from 2nd week of August to 2nd week of November.

Table 2.1 Mean Monthly Rainfall, PET, 1/2 PET at Koppal Taluk and District

Sl. no.	Months	Rainfall	PET	1/2 PET
1	January	1.60	116.70	58.35
2	February	1.50	129.20	64.60
3	March	14.10	169.80	84.90
4	April	18.10	180.60	90.30
5	May	41.60	193.50	96.75
6	June	85.80	167.90	83.95
7	July	72.10	156.20	78.10
8	August	110.50	152.50	76.25
9	September	155.60	138.50	69.25
10	October	116.30	122.30	61.15
11	November	36.00	106.40	53.20
12	December	9.10	101.00	50.50
TOTAL		**662.30**	**144.55**	



Fig. 2.3 Rainfall distribution in Koppal Taluk and District

2.6 Natural Vegetation

The natural vegetation is sparse comprising few tree species, shrubs and herbs. The mounds, ridges and boulders occupy sizeable areas which are under thin to moderately thick forest vegetation. Still, there are some remnants of the past forest cover which can be seen in patches in some ridges and hillocks in the microwatershed (Fig 2.4).

Apart from the continuing deforestation, the presence of large population of goats, sheep and other cattle in the microwatershed is causing vegetative degradation of whatever little vegetation left in the area. The uncontrolled grazing has left no time for the regeneration of the vegetative cover. This leads to the accelerated rate of erosion on the hill slopes, resulting in the formation of deep gullies in the foot slopes and eventually resulting in the heavy siltation of few tanks and reservoirs in the microwatershed.

Fig 2.4 Natural vegetation of Lakshmapura-2 Microwatershed

2.7 Land Utilization

About 91 per cent area (Table 2.2) in Koppal district is cultivated at present and about 17 per cent of the area is sown more than once. An area of about 3 per cent is currently barren. Forests occupy a small area of about 5 per cent and the tree cover is in a very poor state. Most of the mounds, ridges and boulder areas have very poor vegetative cover. Major crops grown in the area are sorghum, maize, bajra, cotton, safflower, sunflower, red gram, horse gram, onion, mulberry, pomegranate, sugarcane, bengalgram and groundnut (Fig 2.5 a and b). While carrying out land resource inventory, the land use/land cover particulars are collected from all the survey numbers and a current land use map of the microwatershed is prepared. The current land use map prepared shows the arable and non-arable lands, other land uses and different types of crops grown in the area. The current land use map of Lakshmapura-2 Microwatershed is presented in Fig.2.6. Simultaneously, enumeration of existing wells (bore wells) is made and their location in different survey numbers is marked on the cadastral map. Map showing the location of wells in Lakshmapura-2 Microwatershed is given in Fig 2.7.

Table 2.2 Land Utilization in Koppal District

Sl. No.	Agricultural land use	Area (ha)	Per cent
1	Total geographical area	552495	
2	Total cultivated area	500542	90.6
3	Area sown more than once	92696	16.8
4	Trees and groves	210	0.04
5	Cropping intensity	-	118
6	Forest	29451	5.33
7	Cultivable wasteland	2568	0.46
8	Permanent Pasture land	14675	2.66
9	Barren land	16627	3.01
10	Non agricultural land	40591	7.35
11	Current fallow	19660	3.56

Fig.2.5 (a) Different crops and cropping systems in Lakshmapura-2 Microwatershed

Fig.2.5 (b) Different crops and cropping systems in Lakshmapura-2 Microwatershed



Fig.2.6 Current Land Use – Lakshmapura-2 Microwatershed



Fig.2.7 Location of wells– Lakshmapura-2 Microwatershed

Chapter 3

SURVEY METHODOLOGY

The purpose of land resource inventory is to delineate similar areas (soil series and phases), which respond or expected to respond similarly for a given level of management. This was achieved in Lakshmapura-2 microwatershed by the detailed study of all the soil characteristics (depth, texture, colour, structure, consistence, coarse fragments, porosity, soil reaction, soil horizons etc.) and site characteristics(slope, erosion, drainage, occurrence of rock fragments etc.) followed by grouping of similar areas based on soil-site characteristics into homogeneous (management units) units and showing their extent and geographic distribution on the microwatershed cadastral map. The detailed soil survey at 1:7920 scale was carried out in 460 ha area. The methodology followed for carrying out land resource inventory was as per the guidelines given in Soil Survey Manual (IARI, 1971; Soil Survey Staff, 2006; Natarajan *et al.,* 2015) which is briefly described below.

3.1 Base Maps

The detailed survey of the land resources occurring in the microwatershed was carried out by using digitized cadastral map and satellite imagery as base supplied by the KSRSAC. The cadastral map shows field boundaries with their survey numbers, location of tanks, streams and other permanent features of the area (Fig. 3.1). Apart from the cadastral map, remote sensing data products from Cartosat-1 and LISS IV merged at the scale of 1:7920 were used in conjunction with the cadastral map to identify the geology, landscapes, landforms and other surface features. The imagery helped in the identification and delineation of boundaries between hills, uplands and lowlands, water bodies, forest and vegetated areas, roads, habitations and other cultural features of the area (Fig.3.2).The cadastral map was overlaid on the satellite imagery (Fig.3.3) that helps to identify the parcel boundaries and other permanent features. Apart from cadastral maps and images, toposheets of the area (1:50,000 scale) were used for initial traversing, identification of geology, landscapes and landforms, drainage features, present land use and also for selection of transects in the microwatershed.

3.2 Image Interpretation for Physiography

False Colour Composites (FCC) of Cartosat-I and LISS-IV merged satellite data covering the microwatershed area was visually interpreted using image interpretation elements and all the available collateral data with local knowledge. The delineated physiographic boundaries were transferred on to a cadastral map overlaid on satellite imagery. Physiographically, the area has been identified as granite gneiss and alluvial landscapes and is divided into landforms such as ridges, mounds and uplands based on slope. They were further subdivided into physiographic/ image interpretation units based on image characteristics. The image interpretation legend for physiography is given below.

Image Interpretation Legend for Physiography

G- Granite gneiss landscape

G1 Hills/ Ridges/ Mounds

G11 Summits

G12 Side slopes

G121 Side slopes with dark grey tones

G2 Uplands

G21 Summits

G22 Gently sloping uplands

G221 Gently sloping uplands, yellowish green (eroded)

G222 Gently sloping uplands, yellowish white (severely eroded)

G23 Very gently sloping uplands

G231 Very gently sloping uplands, yellowish green

G232 Very gently sloping uplands, medium green and pink

G233 Very gently sloping uplands, pink and green (scrub land)

G234 Very gently sloping uplands, medium greenish grey

G235 Very gently sloping uplands, yellowish white (eroded)

G236 Very gently sloping uplands, dark green

G237 Very gently sloping uplands, medium pink (coconut garden)

G238 Very gently sloping uplands, pink and bluish white (eroded)

DSe -Alluvial landscape

DSe 1 Summit

DSe 11 Nearly level Summit with dark grey tone

DSe 12 Nearly level Summit with medium grey tone

DSe 13 Nearly level Summit with whitish grey tone

DSe 14 Nearly level Summit with whitish tone (Calcareousness)

DSe 15 Nearly level Summit with pinkish grey tone

DSe 16 Nearly level Summit with medium pink tone

DSe 17 Nearly level Summit with bluish white tone

DSe 18 Nearly level Summit with greenish grey tone

DSe 2 Very genetly sloping

DSe 21 Very gently sloping, whitish tone

DSe 22 Very gently sloping, greyish pink tone

DSe 23 Very gently sloping, whitish grey tone

DSe 24 Very gently sloping, medium grey tone

DSe 25 Very gently sloping, medium pink tone

DSe 26 Very gently sloping, dark grey tone

DSe 27 Very gently sloping, bluish grey tone

DSe 28 Very gently sloping, greenish grey tone

DSe 29 Very gently sloping, Pinkish grey



Fig 3.1 Scanned and Digitized Cadastral map of Lakshmapura-2 Microwatershed



Fig.3.2 Satellite Image of Lakshmapura-2 Microwatershed



Fig.3.3 Cadastral map overlaid on IRS PAN+LISS IV merged imagery of Lakshmapura-2 Microwatershed

3.3 Field Investigation

The field boundaries and survey numbers given on the cadastral sheet were located on the ground by following permanent features like roads, cart tracks, *nallas*, streams, tanks etc., and wherever changes were noticed, they were incorporated on the microwatershed cadastral map. Preliminary traverse of the microwatershed was carried out with the help of cadastral map, imagery and toposheets. While traversing, landforms and physiographic units identified were checked and preliminary soil legend was prepared by studying soils at few selected places. Then, intensive traversing of each physiographic unit like hills, ridges, uplands and plains was carried out. Based on the variability observed on the surface, transects (Fig 3.4) were selected across the slope covering all the landform units in the microwatershed (Natarajan and Dipak Sarkar, 2010).



Fig: 3.4. Location of profiles in a transect

In the selected transect, soil profiles (Fig.3.4) were located at closely spaced intervals to take care of any change in the land features like break in slope, erosion, gravel, stones etc. In the selected sites, profiles (vertical cut showing the soil layers from surface to the rock) were opened upto 200 cm or to the depth limited by rock or hard substratum and studied in detail for all their morphological and physical characteristics. The soil and site characteristics were recorded for all profile sites on a standard proforma as per the guidelines given in USDA Soil Survey Manual (Soil Survey Staff, 2012). Apart from the transect study, profiles were also studied at random, almost like in a grid pattern, outside the transect areas to validate the soil map unit boundaries.

Based on the soil characteristics, the soils were grouped into different soil series. Soil series is the most homogeneous unit having similar horizons and properties and behaves similarly for a given level of management. Soil depth, texture, colour, kind of horizon and horizon sequence, amount and nature of gravel present, calcareousness, nature of substratum etc, were used as the major differentiating characteristics for identifying soil series occurring in the area. The differentiating characteristics used for identifying the soil series are given in Table 3.1. Based on the above characteristics, 21 soil series were identified in Lakshmapura-2 microwatershed.

Table 3.1 Differentiating Characteristics used for identifying Soil Series (Characteristics are of Series Control Section)

Sl. No	Soil Series	Depth (cm)	Colour (moist)	Texture	Gravel (%)	Horizon sequence	Calcareousness
Soils of Granite Gneiss Landscape							
1	**Abbigeri (ABR)**	25-50	2.5YR 3/3, 3/4	gsc	>35	Ap-Bt-Cr	-
2	**Harve (HRV)**	25-50	2.5YR3/4,3/6 5YR3/3,4/4,3/4	gscl	>35	Ap-Bt-Cr-	-
3	**Kanchanahalli (KNH)**	25-50	2.5YR3/4,3/6	sc	<15	Ap-Bt-Cr	-

Sl. No	Soil Series	Depth (cm)	Colour (moist)	Texture	Gravel (%)	Horizon sequence	Calcareo-usness
4	**Hatti (HTI)**	50-75	5 YR 3/3, 3/4,	gsc	15-35	Ap-Bt-Cr	-
5	**Thammadahalli (TDH)**	50-75	2.5YR2.5/4,3/6	sc-c	<15	Ap-Bt-Cr	-
6	**Lakkur (LKR)**	50-75	2.5YR 2.5/3, 2.5/4, 3/4, 3/6	gsc	40-60	Ap-Bt-Bc-Cr	-
7	**Mukhadahalli (MKH)**	50-75	5YR3/3,3/4,4/3, 5/4,6/6 2.5YR3/4	gsc	>35	Ap-Bt-Cr	-
8	**Bisarahalli (BSR)**	75-100	5 YR 3/3, 3/4	gsc	15-35	Ap-Bt-Cr	-
9	**Hooradhahalli (HDH)**	75-100	2.5YR2.5/4,3/4, 3/6	gsc-gc	>35	Ap-Bt-Cr	-
10	**Balapur (BPR)**	100-150	2.5YR2.5/4,3/4	gsc-gc	>35	Ap-Bt-Cr	-
11	**Nagalapur (NGP)**	100-150	5YR2.5/2,3/2, 2.5YR3/6,4/6	gsc	>35	Ap-Bt-Cr	-
12	**Mornal (MNL)**	100-150	5YR 3/4, 2.5 YR 3/4, 4/6	gsc	15-35	Ap-Bt-Cr	-
13	**Hallikere (HLK)**	>150	5YR3/3,3/4 7.5YR3/3,3/4	c	<15	Ap-Bt	-
14	**Ranatur (RTR)**	>150	2.5YR2.5/3,2.5/4, 3/3,4/6	c	-	Ap-Bt	-
Soils of Alluvial Landscape							
15	**Ravanaki (RNK)**	50-75	7.5YR3/2,3/3,5/2,5/3 10YR3/1,3/2,4/1, 4/2, 5/1,6/1	c	<15	Ap-Bw-Cr	e-ev
16	**Dambarahalli (DRL)**	75-100	10YR 2/1, 3/1, 4/3	c	<15	Ap-Bw-Ck	e-es
17	**Gatareddihal (GRH)**	100-150	10YR 2/1, 3/1, 2.5Y 4/3, 5/4	c	<15	Ap-Bss-BC-C	es
18	**Kavalur (KVR)**	100-150	10 YR 2/2, 3/1, 3/2, 3/3, 4/4	c	-	Ap-Bss-Bck-Cr	es-ev
19	**Lakshmangudda (LGD)**	100-150	10YR3/1,3/2,4/1,4/2, 7.5YR3/1,3/2,5/1, 2.5Y5/2,5/3,6/3	c	<15	Ap-Bss-Ck	es
20	**Alawandi (AWD)**	>150	10 YR 2/1, 3/2,	c	<15	Ap-Bss	e-es
21	**Bardur (BDR)**	>150	10YR 2/1, 3/1, 3/2,	c	<15	Ap-Bss	es

3.4 Soil Mapping

The area under each soil series was further separated into soil phases and their boundaries delineated on the cadastral map based on the variations observed in the texture of the surface soil, slope, erosion, presence of gravel, stoniness etc. A soil phase is a subdivision of soil series based mostly on surface features that affect its use and management. The soil mapping units are shown on the map (Fig.3.5) in the form of

symbols. During the survey many soil profile pits, few mini pits and a few auger bores representing different landforms occurring in the microwatershed were studied. In addition to the profile study, spot observations in the form of mini pits, road cuts, terrace cuts etc., were studied to validate the soil boundaries on the soil map.

The soil map shows the geographic distribution of 33 mapping units representing 21 soil series occurring in the microwatershed. The soil map unit (soil legend) description is presented in Table 3.2. The soil phase map (management units) shows the distribution of 33 phases mapped in the microwatershed. Each mapping unit (soil phase) delineated on the map has similar soil and site characteristics. In other words, all the farms or survey numbers included in one soil phase will have similar management needs and have to be treated accordingly.

3.5 Land Management Units

The 33 soil phases identified and mapped in the microwatershed were regrouped into nine Land Management Units (LMU's) for the purpose of preparing a Proposed Crop Plan for sustained development of the microwatershed. The database (soil phases) generated under LRI was utilized for identifying Land Management Units (LMU's) based on the management needs. One or more than one soil site characteristic having influence on the management have been chosen for identification and delineation of LMUs. For Lakshmapura-2 microwatershed, five soil and site characteristics, namely the soil depth, soil texture, slope, erosion and gravel content have been considered for defining LMUs. The land management units are expected to behave similarly for a given level of management.

3.6 Laboratory Characterization

Soil samples for each soil series soil were collected from representative master profiles for laboratory characterization by following the methods outlined in the Laboratory Manual (Sarma *et al*, 1987). Surface soil samples collected in the year 2018 from farmer's fields in Lakshmapura-2 microwatershed for fertility status (major and micronutrients) at 320 m grid interval were analyzed in the laboratory (Katyal and Rattan, 2003). By linking the soil fertility data to the survey numbers through GIS, soil fertility maps were generated using Kriging method for the microwatershed.

Table 3.2 Soil map unit description of Lakshmapura-2 Microwatershed

Soil map unit No*	Soil Series	Soil Phase Symbol	Mapping Unit Description	Area in ha (%)
Soils of Granite gneiss Landscape				
	ABR	Abbigere soils are shallow (25-50 cm), well drained, have dark reddish brown, red gravelly sandy clay soils occurring on very gently sloping uplands under cultivation.		**15 (3.24)**
472		ABRiB2g2	Sandy clay surface, slope 1-3%, moderate erosion, very gravelly (35-60%)	15 (3.24)

Soil map unit No*	Soil Series	Soil Phase Symbol	Mapping Unit Description	Area in ha (%)
	HRV	Harve soils are shallow (25-50 cm), well drained, dark red to dark red dish brown, red gravelly sandy clay loam soils occurring on nearly level to gently sloping uplands under cultivation		**41 (9.1)**
27		HRVhB2g2	Sandy clay loam surface, slope 1-3%, moderate erosion, very gravelly (35-60%)	17 (3.74)
29		HRViA1g1	Sandy clay surface, slope 0-1%, slight erosion, gravelly (15-35%)	4 (0.96)
31		HRViB2g1	Sandy clay surface, slope 1-3%, moderate erosion, gravelly (15-35%)	20 (4.4)
	KNH	Kanchanahalli soils are shallow (25-50 cm), well drained, have dark reddish brown to dark red sandy clay soils occurring on very gently sloping uplands under cultivation		**13 (2.73)**
467		KNHiB1g1	Sandy clay surface, slope 1-3%, slight erosion, gravelly (15-35%)	13 (2.73)
	HTI	Hatti soils are moderately shallow (50-75 cm), well drained, have dark reddish brown, red gravelly sandy clay soils occurring on nearly level to very gently sloping uplands under cultivation		**16 (3.56)**
96		HTIiA1g1	Sandy clay surface, slope 0-1%, slight erosion, gravelly (15-35%)	4 (0.87)
100		HTIiB2	Sandy clay surface, slope 1-3%, moderate erosion	0.45 (0.1)
101		HTIiB2g1	Sandy clay surface, slope 1-3%, moderate erosion, gravelly (15-35%)	12 (2.59)
	TDH	Thammadahalli soils are moderately shallow (50-75cm), well drained, have brown to very dark brown and dark reddish brown, sandy clay to clay soils occurring on nearly level to gently sloping uplands		**13 (2.92)**
60		TDHiB1	Sandy clay surface, slope 1-3%, slight erosion	13 (2.92)
	LKR	Lakkur soils are moderately shallow (50-75 cm), well drained, have dark reddish brown to dark red, red gravelly sandy clay soils occurring on very gently to moderately sloping uplands under cultivation		**2 (0.35)**
54		LKRiB2g1	Sandy clay surface, slope 1-3%, moderate erosion, gravelly (15-35%)	2 (0.35)
	MKH	Mukhadahalli soils are moderately shallow (50-75 cm), well drained, have dark brown to reddish brown, red gravelly sandy clay soils occurring on gently very gently to gently sloping uplands under cultivation		**2 (0.34)**
82		MKHhB1g1	Sandy clay loam surface, slope 1-3%,slight erosion, gravelly (15-35%)	2 (0.34)
	BSR	Bisarahalli soils are moderately deep (75-100 cm), well drained, have dark reddish brown, red gravelly sandy clay soils occurring on very gently sloping uplands under cultivation		**1 (0.27)**
165		BSRiB1g1	Sandy clay surface, slope 1-3%, slight erosion	1 (0.27)

Soil map unit No*	Soil Series	Soil Phase Symbol	Mapping Unit Description	Area in ha (%)
			gravelly (15-35%)	
	HDH	Hooradhahalli soils are moderately deep (75-100 cm), well drained, have dark red to dark reddish brown, red gravelly sandy clay to clay soils occurring on nearly level to moderately sloping uplands under cultivation		**105 (23.19)**
121		HDHhB1g2	Sandy clay loam surface, slope 1-3%, slight erosion, very gravelly (35-60%)	4 (0.95)
123		HDHhB2g1	Sandy clay loam surface, slope 1-3%, moderate erosion, gravelly (15-35%)	24 (5.3)
127		HDHiB2	Sandy clay surface, slope 1-3%, moderate erosion	54 (11.84)
128		HDHiB2g1	Sandy clay surface, slope 1-3%, moderate erosion, gravelly (15-35%)	23 (5.1)
	BPR	Balapur soils are deep (100-150 cm), well drained, have dark reddish brown to dark red, gravelly sandy clay to clay soils occurring on nearly level to gently sloping uplands under cultivation		**15 (3.34)**
231		BPRhB2g1	Sandy clay loam surface, slope 1-3%, moderate erosion, gravelly (15-35%)	15 (3.34)
	NGP	Nagalapur soils are deep (100-150 cm), well drained, have dark reddish brown to dark red, gravelly sandy clay soils occurring on nearly level to gently sloping uplands under cultivation		**4 (0.78)**
266		NGPiB2g2	Sandy clay surface, slope 1-3%, moderate erosion, very gravelly (35-60%)	4 (0.78)
	MNL	Mornal soils are deep (100-150 cm), well drained, have dark reddish brown to red, gravelly sandy clay soils occurring on very gently sloping uplands under cultivation		**28 (6.04)**
208		MNLiB2	Sandy clay surface, slope 1-3%, moderate erosion	10 (2.22)
209		MNLiB2g1	Sandy clay surface, slope 1-3%, moderate erosion, gravelly (15-35%)	18 (3.82)
	HLK	Hallikere soils are very deep (>150 cm), well drained, have dark brown to dark reddish brown clayey soils occurring on nearly level to very gently sloping uplands under cultivation		**13 (2.86)**
274		HLKiB2	Sandy clay surface, slope 1-3%, moderate erosion	13 (2.86)
	RTR	Ranatur soils are very deep (>150 cm), well drained, have dark reddish brown to dark red, clay soils occurring on nearly level to very gently sloping uplands under cultivation		**5 (1.16)**
288		RTRiB2	Sandy clay surface, slope 1-3%, moderate erosion	5 (1.16)
Soils of Alluvial Landscape				
	RNK	Ravanaki soils are moderately shallow (50-75 cm), moderately well drained, have dark brown to very dark grayish brown and dark gray, calcareous, black clay soils occurring on nearly level to very gently sloping plains under cultivation		**17 (3.61)**

Soil map unit No*	Soil Series	Soil Phase Symbol	Mapping Unit Description	Area in ha (%)
337		RNKmB2g1	Clay surface, slope 1-3%, moderate erosion, gravelly (15-35%)	17 (3.61)
	DRL	Dambarahalli soils are moderately deep (75-100 cm), moderately well drained, have dark brown to very dark gray, calcareous black cracking clay soils occurring on nearly level to very gently sloping plains under cultivation		**55 (12.01)**
350		DRLmB2	Clay surface, slope 1-3%, moderate erosion	34 (7.44)
351		DRLmB2g1	Clay surface, slope 1-3%, moderate erosion, gravelly (15-35%)	21 (4.57)
	GRH	Gatareddihal soils are deep (100-150 cm), moderately well drained, have light olive brown to very dark gray, sodic black calcareous cracking clay soils occurring on nearly level to very gently sloping plains under cultivation		**24 (5.19)**
372		GRHmB1g1	Clay surface, slope 1-3%, slight erosion, gravelly (15-35%)	12 (2.57)
373		GRHmB2	Clay surface, slope 1-3%,moderate erosion	12 (2.62)
	KVR	Kavalur soils are deep (100-150 cm), moderately well drained, have dark yellowish brown to very dark grayish brown, calcareous black cracking clay soils occurring on nearly level to very gently sloping plains under cultivation		**22 (4.77)**
386		KVRmA1	Clay surface, slope 0-1%,slight erosion	1 (0.21)
388		KVRmB1	Clay surface, slope 1-3%,slight erosion	8 (1.66)
390		KVRmB2g1	Clay surface, slope 1-3%, moderate erosion, gravelly (15-35%)	13 (2.9)
	LGD	Lakshmangudda soils are deep (100-150 cm), moderately well drained, have light olive brown to very dark gray calcareous black cracking clay soils occurring on nearly level to very gently sloping plains under cultivation		**26 (5.55)**
393		LGDmB1	Clay surface, slope 1-3%,slight erosion	26 (5.55)
	AWD	Alawandi soils are very deep (>150 cm), moderately well drained, have very dark grayish brown to black, calcareous black cracking clay soils occurring on nearly level to very gently sloping plains under cultivation		**11 (2.41)**
424		AWDmB2	Clay surface, slope 1-3%, moderate erosion	11 (2.41)
	BDR	Bardur soils are very deep (>150 cm), moderately well drained, have very dark grayish brown to very dark gray, black calcareous cracking clay soils occurring on nearly level to very gently sloping plains under cultivation		**23 (4.92)**
433		BDRmB2	Clay surface, slope 1-3%, moderate erosion	23 (4.92)
992		Railway		0.30

Soil map unit No*	Soil Series	Soil Phase Symbol	Mapping Unit Description	Area in ha (%)
				(0.06)
1000		Others	Habitation & Waterbody	7 (1.6)

*Soil map unit numbers are continuous for the taluk, not the microwatersheds



Fig 3.5 Soil Phase or Management Units- Lakshmapura-2 Microwatershed

Chapter 4

THE SOILS

Detailed information pertaining to the nature, extent and distribution of different kinds of soils occurring in Lakshmapura-2 microwatershed is provided in this chapter. The microwatershed area has been identified as granite gneiss and alluvial landscapes based on geology. In all, 21 soil series were identified. Soil formation is the result of the combined effect of environmental and terrain factors that are reflected in soil morphology. The soil formation is dominantly influenced by the parent material, climate, time and relief.

A brief description of each of the 21 soil series identified followed by 33 soil phases (management units) mapped (Fig. 3.5) are furnished below. The physical and chemical characteristics of soil series identified in Lakshmapura-2 microwatershed are given in Table 4.1 along with soil classification. The soils in any one map unit differ from place to place in their depth, texture, slope, gravelliness, erosion or any other site characteristic that affect management. The soil phase map can be used for identifying the suitability of areas for growing specific crops or for other alternative uses and also for deciding the type of conservation structures needed. The detailed information on soil and site-characteristics like soil depth, surface soil texture, slope, erosion, gravelliness, AWC, LCC etc, with respect to each of the soil phase identified is given village/survey number wise for the microwatershed in Appendix-I.

4.1 Soils of Granite gneiss Landscape

In this landscape, 14 soil series were identified and mapped. Of these series, Hooradhahalli (HDH) series occupies a maximum area of 105 ha (23%) and others occupy minor area. The brief description of the soil series along with the soil phases identified and mapped is given below.

4.1.1 Abbigere (ABR) Series: Abbigere soils are shallow (25-50 cm), well drained, have dark reddish brown, to red gravelly sandy clay soils. They have developed from weathered granite gneiss and occur on very gently sloping uplands under cultivation. The Abbigere soil series has been classified as a member of the clayey-skeletal, mixed, isohyperthermic family of (Paralithic) Rhodustalfs.

The thickness of the solum ranges from 28 to 48 cm. The thickness of A-horizon ranges from 12 to 17 cm. Its colour is in 5YR and 2.5 YR hue with value 3 to 4 and chroma 3 to 4. The texture is sandy clay with 20 to 35 per cent gravel. The thickness of B-horizon ranges from 16 to 32 cm. Its colour is in 2.5 YR and 5 YR hue with value 2.5 to 4 and chroma 2 to 3. Its texture is sandy clay with gravel content of more than 35 per cent. The available water capacity is very low (<50 mm/m).One soil phase was identified and mapped.

Landscape Soil Profile Characteristics of Abbigere (ABR) Series

4.1.2 Harve (HRV) Series: Harve soils are shallow (25-50 cm), well drained, have reddish brown to dark red gravelly sandy clay loam soils. They have developed from weathered granite gneiss and occur on very gently to moderately sloping uplands. The Harve series has been classified as a member of the loamy-skeletal, mixed, isohyperthermic family of (Paralithic) Rhodustalfs.

The thickness of the solum ranges from 28 to 48 cm. The thickness of A-horizon ranges from 12 to 17 cm. Its colour is in 5YR and 2.5 YR hue with value 3 to 4 and chroma 4 to 6. The texture varies from loamy sand to sandy loam with 20 to 60 per cent gravel. The thickness of B-horizon ranges from 16 to 32 cm. Its colour is in 2.5 YR and 5 YR hue with value 3 to 4 and chroma 4 to 6. Its texture is sandy clay loam with gravel content of more than 35 per cent. The available water capacity is very low (<50mm/m). Three soil phases were identified and mapped.

Landscape and soil profile characteristics of Harve (HRV) Series

4.1.3 Kanchanahalli (KNH) Series: Kanchanahalli soils are shallow (25 -50 cm), well drained, have dark reddish brown, sandy clay soils. They have developed from weathered granite gneiss and occur on very gently sloping uplands. The Kanchanahalli series has been classified as a member of the clayey, mixed, isohyperthermic family of (Paralithic) Rhodustalfs.

The thickness of the solum ranges from 28 to 50 cm. The thickness of A horizon ranges from 12 to 18 cm. Its colour is in 5YR and 2.5 YR hue with value 3 and chroma 4 to 6. The texture varies from sandy clay loam to sandy clay with 10 to 15 per cent gravel. The thickness of B horizon ranges from 16 to 38 cm. Its colour is in 2.5 YR hue with value 3 to 4 and chroma 4 to 6. Its texture is sandy clay with gravel content of < 15 per cent. The available water capacity is low (50-100 mm/m). One soil phase was identified and mapped.

Landscape and soil profile characteristics of Kanchanahalli (KNH) Series

4.1.4 Hatti (HTI) Series: Hatti soils are moderately shallow (50-75cm), well drained, have dark reddish brown, gravelly sandy clay soils. They are developed from weathered granite gneiss and occur on very gently sloping uplands under cultivation. The Hatti series has been classified as a member of the fine, mixed, isohyperthermic family of Typic Paleustalfs.

The thickness of the solum ranges from 57 to 74 cm. The thickness of A horizon ranges from 16 to 20 cm. Its colour is in 5 YR hue with value and chroma 3 to 4. The texture varies from sandy loam to sandy clay loam and sandy clay with 15 to 60 per cent gravel. The thickness of B horizon ranges from 45 to 56 cm. Its colour is in 5 YR hue with value 3 and chroma 3 to 4. Texture is sandy clay with 15 to 35 per cent gravel. The available water capacity is low (50-100 mm/m).Three soil phases were identified and mapped.

Landscape and soil profile characteristics of Hatti (HTI) Series

4.1.5 Thammadahalli (TDH) Series: Thammadahalli soils are moderately shallow (50-75cm), well drained, have brown to very dark brown and dark reddish brown sandy clay to clay soils. They have developed from weathered granite gneiss and occur on nearly level to gently sloping uplands. The Thammadahalli series has been classified as a member of the fine, mixed, isohyperthermic family of Rhodic Paleustalfs.

The thickness of the solum ranges from 54 to 75 cm. The thickness of A horizon ranges from 11 to 19 cm. Its colour is in 7.5 YR, 5YR and 2.5 YR hue with value 2.5 to 4 and chroma 2 to 6. The texture varies from sandy clay loam to clay with 10 to 20 per cent gravel. The thickness of B horizon ranges from 43 to 60 cm. Its colour is in 2.5 YR hue with value 3 and chroma 4 to 6. Its texture is sandy clay to clay. The available water capacity is medium (100-150 mm/m). One soil phase was identified and mapped.

Landscape and soil profile characteristics of Thammadahalli (TDH) Series

4.1.6 Lakkur (LKR) Series: Lakkur soils are moderately shallow (50-75cm), well drained, have reddish brown to dark red gravelly sandy clay soils. They have developed from weathered granite gneiss and occur on nearly level to very gently and gently sloping uplands. The Lakkur series has been classified as a member of the clayey-skeletal, mixed, isohyperthermic family of Typic Rhodustalfs.

The thickness of the solum ranges from 51 to 74 cm. The thickness of A horizon ranges from 12 to 18 cm. Its colour is in 5YR and 2.5 YR hue with value 3 to 4 and chroma 4 to 6. The texture varies from loamy sand to sandy clay loam with 15 to 50 per cent gravel. The thickness of B horizon ranges from 39 to 58 cm. Its colour is in 2.5 YR hue with value 3 to 4 and chroma 4 to 6. Texture is sandy clay with 40 to 60 per cent gravel. The available water capacity is low (50-100 mm/m). One soil phase was identified and mapped.

Landscape and soil profile characteristics of Lakkur (LKR) Series

4.1.7 Mukhadahalli (MKH) Series: Mukhadahalli soils are moderately shallow (50-75 cm), well drained, have dark brown to reddish brown, gravelly sandy clay soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands. The Mukhadahalli series has been classified as a member of the clayey-skeletal, mixed, isohyperthermic family of Typic Haplustalfs.

The thickness of the solum ranges from 51 to 72 cm. The thickness of A horizon ranges from 12 to 17 cm. Its colour is in 5 YR and 7.5 YR hue with value 3 to 4 and chroma 2 to 4. The texture varies from loamy sand to sandy loam with 20 to 45 per cent gravel. The thickness of B horizon ranges from 40 to 68 cm. Its colour is in 2.5 YR and 5 YR hue with value and chroma 3 to 6. Texture is sandy clay loam to sandy clay with 35 to 50 per cent gravel. The available water capacity is low (50-100 mm/m). One soil phase was identified and mapped.

Landscape and soil profile characteristics of Mukhadahalli (MKH) Series

4.1.8 Bisarahalli (BSR) Series: Bisarahalli soils are moderately deep (75-100 cm), well drained, have dark reddish brown, gravelly sandy clay red soils. They have developed from weathered granite gneiss and occur on very gently sloping uplands under cultivation. The Bisarahalli series has been classified as a member of the fine, mixed isohyperthermic family of Typic Paleustalfs.

The thickness of the solum ranges from 75 to 98 cm. The thickness of A horizon ranges from 17 to 25 cm. Its colour is in 5 YR hue with value 3 to 4 and chroma 3 to 6. The texture ranges from sandy clay loam to sandy clay with 15 to 35 per cent gravel. The thickness of B horizon ranges from 61 to 79 cm. Its colour is in 5 YR hue with value 3 and chroma 3 to 4. Its texture is gravelly sandy clay with gravel content of 15-35 per cent. The available water capacity is low (50-100 mm/m). One soil phase was identified and mapped.

Landscape and soil profile characteristics of Bisarahalli (BSR) Series

4.1.9 Hooradhahalli (HDH) Series: Hooradhahalli soils are moderately deep (75-100 cm), well drained, have red to dark red and reddish brown, gravelly sandy clay to clay soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands. The Hooradhahalli series has been classified as a member of the clayey-skeletal, mixed, isohyperthermic family of Rhodic Paleustalfs.

The thickness of the solum ranges from 76 to 100 cm. The thickness of A horizon ranges from 11 to 19 cm. Its colour is in 5 YR and 2.5 YR hue with value 3 to 4 and chroma 3 to 6. The texture varies from loamy sand to sandy clay with 15 to 50 per cent gravel. The thickness of B horizon varies from 65 to 83 cm. Its colour is in 2.5 YR hue with value 2.5 to 3 and chroma 4 to 6. Texture is sandy clay to clay with 35 to 50 per cent gravel. The available water capacity is low (50-100mm/m). Four soil phases were identified and mapped.

Landscape and soil profile characteristics of Hooradhahalli (HDH) Series

4.1.10 Balapur (BPR) Series: Balapur soils are deep (100-150 cm), well drained, have dark reddish brown to dark red, gravelly sandy clay to clay soils. These soils are developed from weathered granite gneiss and occur on very gently to gently sloping uplands. The Balapur series has been classified as a member of the clayey-skeletal, mixed, isohyperthermic family of Typic Rhodustalfs.

The thickness of the solum ranges from 102 to 147 cm. The thickness of A horizon ranges from 12 to 17cm. Its colour is in 5 YR and 2.5 YR hue with value and chroma 3 to 4. The texture ranges from loamy sand to sandy clay with 15 to 50 per cent gravel. The thickness of B horizon ranges from 90 to 132 cm. Its colour is in 2.5 YR hue with value 2.5 to 3 and chroma 4 to 6. Texture is sandy clay to clay with 35 to 50 per cent gravel. The available water capacity is medium (100-150 mm/m). One soil phase was identified and mapped.

Landscape and soil profile characteristics of Balapur (BPR) Series

4.1.11 Nagalapur (NGP) Series: Nagalapur soils are deep (100-150 cm), well drained, have dark reddish brown to dark red gravelly sandy clay soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands. The Nagalapur series has been classified as a member of the clayey-skeletal, mixed, isohyperthermic family of Typic Paleustalfs.

The thickness of the solum ranges from 105 to 145 cm. The thickness of A-horizon ranges from 14 to 20 cm. Its colour is in 7.5 YR hue with value and chroma 3 to 4. The texture ranges from sandy loam to sandy clay with 10 to 50 per cent gravel. The thickness of B horizon ranges from 90 to 128 cm. Its colour is in 2.5 YR, 5 YR and 7.5 YR hue with value 3 to 5 and chroma 3 to 6. Texture is sandy clay to clay with 35 to 80 per cent gravel. The available water capacity is low (51-100 mm/m). One soil phase was identified and mapped.

Landscape Soil Profile Characteristics of Nagalapur (NGP) Series

4.1.12 Mornal (**MNL) Series:** Mornal soils are deep (100-150 cm), well drained have dark reddish brown to dark red gravelly sandy clay soils. They are developed from weathered granite gneiss and occur on very gently sloping uplands under cultivation. The Mornal soil series has been classified as a member of the fine, mixed, isohyperthermic family of Rhodic Paleustalfs.

The thickness of the solum ranges from 112 to 149 cm. The thickness of A-horizon ranges from 15 to 25 cm. Its colour is in 5 YR, 10 YR hue with value 3 to 4 and chroma 2 to 4. The texture is sandy clay loam, sandy clay and clay with 15 to 30 per cent gravel. The thickness of B-horizon ranges from 103 to 131 cm. Its colour is in 2.5 YR and 5 YR hue with value 2.5 to 4 and chroma 3 to 6. Texture is sandy clay loam to sandy clay with 15 to 35 per cent gravel. The available water capacity is medium (101-150 mm/m).Two soil phases were identified and mapped.

Landscape and soil profile characteristics of Mornal (MNL) Series

4.1.13 Hallikere (HLK) Series: Hallikere soils are very deep (>150 cm), well drained, have dark brown and dark reddish brown clayey soils. They have developed from weathered granite gneiss and occur on nearly level to very gently sloping uplands. The Hallikere series has been classified as a member of the fine, mixed, isohyperthermic family of Typic Paleaustalfs.

The thickness of the solum is more than 150 cm. The thickness of A horizon ranges from 11 to 14 cm. Its colour is in 7.5 YR and 5 YR hue with value 3 to 4 and chroma 3 to 4. The texture varies from sandy loam to sandy clay loam. The thickness of B horizon is more than 150 cm. Its colour is in 7.5 YR and 5 YR hue with value and chroma 3 to 4. Its texture is clay. The available water capacity is high (150-200 mm/m). One soil phase was identified and mapped.

Landscape Soil Profile Characteristics of Hallikere (HLK) Series

4.1.14 Ranatur (RTR) Series: Ranatur soils are very deep (> 150 cm), well drained, have dark reddish brown to dark red, clayey soils. They have developed from weathered granite gneiss and occur on very gently sloping uplands. The Ranatur series has been classified as a member of the fine, mixed, isohyperthermic family of Rhodic Paleustalfs.

The thickness of the solum is more than 150 cm. The thickness of A horizon ranges from 8 to 14 cm. Its colour is in 5 YR and 2.5 YR hue with value 2.5 to 4 and chroma 3 to 6. The texture varies from sandy loam to sand clay. The thickness of B horizon is more than 150 cm. Its colour is in 2.5 YR hue with value 2.5 to 3 and chroma 3 to 6. Its texture is clay. The available water capacity is high (150-200 mm/m). One soil phase was identified and mapped.

Landscape and soil profile characteristics of Ranatur (RTR) Series

4.2 Soils of Alluvial Landscape

In this landscape, seven soil series were identified and mapped. Of these, series Dambarahalli (DRL) series occupies maximum area of 55 ha (12%) and others occupy minor area. The brief description of the soil series along with the soil phases identified and mapped is given below.

4.2.1 Ravanaki (RNK) Series: Ravanaki soils are moderately shallow (50-75 cm), moderately well drained, have dark brown to very dark grayish brown, calcareous clay soils. They have developed from alluvium and occur on nearly level to very gently sloping plains. The Ravanaki series has been classified as a member of the very fine, smectitic (calc), isohyperthermic family of Typic Haplustepts.

The thickness of the solum ranges from 50 to 75 cm. The thickness of A horizon ranges from 15 to 20 cm. Its colour is in 7.5 YR and 10 YR hue with value 2 to 3 and chroma 2.5 to 4. The texture varies from sandy clay to clay with 10 to 15 per cent gravel. The thickness of B horizon ranges from 35 to 60 cm. Its colour is in 10 YR and 7.5 YR hue with value 2 to 6 and chroma 2 to 4. Its texture is sandy clay to clay with gravel content of 10 to 20 per cent. The available water capacity is low (51-100 mm/m).One soil phase was identified and mapped.

Landscape and Soil Profile Characteristics of Ravanaki (RNK) Series

4.2.2 Dambarahalli (DRL) Series: Dambarahalli soils are moderately deep (75-100 cm), moderately well drained, have black and very dark gray to dark brown, calcareous cracking clay soils. They have developed from alluvium and occur on very gently to gently sloping plains under cultivation. The Dombarahalli series has been classified as a member of the very fine, smectitic (calc), isohyperthermic family of Typic Haplusterts.

The thickness of the solum ranges from 75 to 99 cm. The thickness of A horizon ranges from 13 to 24 cm. Its colour is in 10 YR hue with value 3 to 4 and chroma 1 to 2.

The texture is clay. The thickness of B horizon ranges from 54 to 85 cm. Its colour is in 10 YR hue with value 2 to 4 and chroma 1 to 3. Its texture is clay and are calcareous. The available water capacity is high (151-200 mm/m). Two soil phases were identified and mapped.

Landscape and soil profile characteristics of Dambarahalli (DRL) Series

4.2.3 Gatareddihal (GRH) Series: Gatareddihal soils are deep (100-150 cm), moderately well drained, have black or dark grey to light olive brown, sodic, calcareous cracking clay soils. They are developed from alluvium and occur on nearly level to very gently sloping plains under cultivation. The Gatareddihal series has been classified as a member of the very fine, smectitic (calc), isohyperthermic family of Sodic Haplusterts.

The thickness of the solum ranges from 102 to 149 cm. The thickness of A-horizon ranges from 12 to 19 cm. Its colour is in 7.5 YR, 10 YR hue with value 3 to 4 and chroma 1 to 6. The texture is sandy clay loam to clay. The thickness of B-horizon ranges from 86 to 117 cm. Its colour is in 10 YR and 7.5 YR hue with value 3 and chroma 2 to 6. Texture is clay with less than 15 per cent gravel. The available water capacity is very high (>200 mm/m). Two soil phases were identified and mapped.

Landscape and soil profile characteristics of Gatareddihal (GRH) Series

4.2.4 Kavalur (KVR) Series: Kavalur soils are deep (100-150 cm), moderately well drained, have dark yellowish brown to very dark brown and very dark gray, calcareous black cracking clay soils They have developed from alluvium and occur on very gently sloping plains. The Kavalur series has been classified as a member of the fine, smectitic (calc), isohyperthermic family of Typic Haplusterts.

The thickness of the solum is 113 to 143 cm. The thickness of A horizon ranges from 9 to 24 cm. Its colour is in 10 YR hue with value 3 and chroma 1. The texture is clay with no gravel. The thickness of B horizon ranges from 89 to 134 cm. Its colour is in 10 YR hue with value 3 and chroma 1. Its texture is clay. The available water capacity is very high (>200 mm/m). Three soil phases were identified and mapped.

Landscape and soil profile characteristics of Kavalur (KVR) series

4.2.5 Lakshmangudda (LGD) Series: Lakshmangudda soils are deep (100-150 cm), moderately well drained, have light olive brown to very dark gray calcareous cracking clayey soils. They have developed from alluvium and occur on nearly level plains. The Lakshmangudda series has been classified as a member of the fine, smectitic (calc), isohyperthermic family of Typic Haplusterts.

The thickness of the solum ranges from 108 to 149 cm. The thickness of A horizon ranges from 16 to 20 cm. Its colour is in 7.5 YR and 10 YR hue with value and chroma 3 to 4. The texture varies from sandy clay to clay with 5 to 10 per cent gravel. The thickness of B horizon ranges from 90 to 132 cm. Its colour is in 2.5 Y, 10 YR and 7.5 YR hue with value 3 to 6 and chroma 1 to 3. Its texture is clay. The available water capacity is high (150-200 mm/m).One soil phase was identified and mapped.

Landscape and Soil Profile Characteristics of Lakshmangudda (LGD) Series

4.2.6 Alawandi (AWD) Series: Alawandi soils are very deep (>150 cm), moderately well drained, have black to very dark grayish brown, calcareous cracking clay soils. They have developed from alluvium and occur on nearly level to very gently sloping plains under cultivation. The Alawandi series has been classified as a member of the fine, smectitic (calc), isohyperthermic family of Typic Haplusterts .

The thickness of the solum is more than 150 cm. The thickness of A horizon ranges from 16 to 26 cm. Its colour is in 10 YR hue with value 2 to 3 and chroma 1 to 2. The texture varies from sandy clay to clay. The thickness of B horizon is more than 150 cm. Its colour is in 10 YR hue with value 2 to 3 and chroma 1 to 3. Its texture is clay and is calcareous. The available water capacity is very high (>200 mm/m). One soil phase was identified and mapped.



Landscape and Soil Profile Characteristics of Alawandi (AWD) Series

4.2.7 Bardur (BDR) Series: Bardur soils are very deep (>150 cm), moderately well drained, have very dark grayish brown to very dark gray, black calcareous cracking clay soils. They are developed from alluvium and occur on nearly level to very gently sloping plains under cultivation. The Bardur series has been classified as a member of the very fine, smectitic (calc), isohyperthermic family of Typic Haplusterts.

The thickness of the solum is more than 150 cm. The thickness of A horizon ranges from 15 to 19 cm. Its colour is in 10 YR hue with value 2 and chroma 1 with clay texture. The thickness of B horizon ranges from 146 to 180 cm. Its colour is in 10 YR hue with value 2 to 3 and chroma 1 to 2. Its texture is clay and is calcareous with less than 15 per cent gravel. The available water capacity is very high (>200 mm/m). One soil phase was identified and mapped.

Landscape and soil profile characteristics of Bardur (BDR) Series

Table: 4.1 Physical and Chemical Characteristics of Soil Series identified in Lakshmapura-2 microwatershed

Series Name: Abbigeri (ABR) **Pedon:** R-11
Location: $15^0$26'14.0"N, $76^0$16'39.0"E Abbigeri village, Koppal taluk and district
Analysis at: NBSS&LUP, Regional Centre, Bangalore. **Classification:** Clayey- skeletal, mixed, isohyperthermic (Paralithic) Rhodustalfs

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand								
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)			1/3 Bar	15 Bar
0-10	Ap	81.18	8.29	10.53	24.31	11.90	19.33	16.07	9.56	20	ls	7.13	3.91
10-25	Bt1	54.32	7.39	38.29	26.64	11.34	5.83	6.24	4.27	40	sc	14.71	11.30
25-40	Bt2	53.84	7.99	38.17	22.10	14.32	6.43	6.85	4.15	50	sc	16.45	12.00

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	$CaCO_3$	Exchangeable bases					CEC	CEC/ Clay	Base saturation	ESP
							Ca	Mg	K	Na	Total				
	Water	$CaCl_2$	M KCl	dS m^{-1}	%	%	cmol kg^{-1}							%	%
0-10	6.13	-	-	0.02	0.81	-	1.56	0.50	0.04	0.01	2.12	3.60	0.34	58.76	0.36
10.-25	6.32	-	-	0.03	0.79	-	5.63	2.41	0.12	0.01	8.17	10.60	0.28	77.07	0.10
25-40	6.27	-	-	0.03	0.64	-	5.41	2.24	0.08	0.01	7.74	12.40	0.32	62.44	0.09

Contd…

Series Name: Harve (HRV), **Pedon:** R-10
Location: $15^0$25'11.63"N, $76^0$22'03.65"E Jabbaragudda village, Koppal taluk and district
Analysis at: NBSS&LUP, Regional Centre, Bangalore. **Classification:** Loamy-skeletal, mixed, isohyperthermic (Paralithic) Rhodustalfs

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand								
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)			1/3 Bar	15 Bar
0-15	Ap	65.64	9.07	25.28	29.04	12.99	9.00	3.48	11.15	50	scl	12.87	4.81
15-29	Bt1	56.13	7.75	36.12	27.81	11.43	7.21	1.44	8.24	60	sc	15.69	6.24
29-47	Bt2	63.42	6.53	30.05	32.38	13.93	7.48	5.74	3.89	60	scl	15.41	9.29

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	$CaCO_3$	Exchangeable bases					CEC	CEC/ Clay	Base saturation	ESP
							Ca	Mg	K	Na	Total				
	Water	$CaCl_2$	M KCl	dS m^{-1}	%	%	cmol kg^{-1}							%	%
0-15	6.05	-	-	0.21	0.93	-	8.89	1.96	0.50	0.08	11.43	11.24	0.44	100.00	0.73
15-29	5.99	-	-	0.15	0.29	-	9.72	2.75	0.51	0.09	13.07	12.71	0.35	100.00	0.74
29-47	6.07	-	-	0.11	0.38	-	9.35	2.47	0.49	0.06	12.36	12.71	0.42	97.29	0.44

Contd…

Series Name: Hatti (HTI), **Pedon:** R-20
Location: $15^0$21'45"N, $76^0$03'06" E Lakshmapura village Koppal taluk and district
Analysis at: NBSS&LUP, Regional Centre, Bangalore. **Classification:** Fine, mixed, isohyperthermic Typic Paleustalfs

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand								
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)			1/3 Bar	15 Bar
0-16	Ap	65.33	12.19	22.48	13.79	11.32	13.37	18.31	8.54	15-20	scl	16.83	5.49
16-41	Bt1	41.54	14.04	44.42	6.47	6.26	9.50	13.36	5.95	15-20	c	27.26	16.64
41-64	Bt2	48.71	8.48	42.81	26.06	7.55	5.38	6.31	3.41	55-60	sc	27.22	12.63

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	$CaCO_3$	Exchangeable bases					CEC	CEC/ Clay	Base saturation	ESP
							Ca	Mg	K	Na	Total				
	Water	$CaCl_2$	M KCl	dS m^{-1}	%	%	cmol kg^{-1}							%	%
0-16	7.11			0.109	0.92		21.06	8.23	0.39	0.06	29.74	20.19	0.90	147	0.30
16-41	7.54			0.220	0.92		21.93	8.47	0.23	0.27	30.90	31.31	0.70	99	0.85
41-64	7.82			0.168	0.55		19.43	7.09	0.31	0.47	27.30	26.57	0.62	103	1.77

Contd…

Soil Series: Thammadahalli (TDH), **Pedon:** $TR_1/1$
Location: $15^0$03'41.7"N, $75^0$36'65.2"E, (4D4A3G2d), Nilogal village, Shirahatti taluk, Gadag district
Analysis at: NBSS&LUP, Regional Centre, Bengaluru **Classification:** Fine, mixed, isohyperthermic Rhodic Paleustalfs

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand								
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)			1/3 Bar	15 Bar
0-25	Ap	85.71	7.34	6.94	14.79	13.28	16.10	24.75	16.80	20	ls	-	-
25-65	Bt	47.76	7.96	44.28	15.30	9.78	6.24	7.91	8.53	10	sc	-	-

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	$CaCO_3$	Exchangeable bases					CEC	CEC/ Clay	Base saturation	ESP
							Ca	Mg	K	Na	Total				
	Water	$CaCl_2$	M KCl	dS m^{-1}	%	%	cmol kg^{-1}							%	%
0-25	9.19	-	-	0.18	0.35	1.29	-	-	0.08	0.52	0.60	3.57	0.51	100.00	5.82
25-65	8.00	-	-	0.17	0.35	0.58	-	-	0.15	1.31	1.46	13.87	0.31	100.00	3.78

Contd…

Soil Series: Lakkur (LKR), **Pedon:** RM-8.
Location: $15^0$04'26.3"N, $75^0$37'84.1"E, (4D4A3I1f), Belhatti village, Shirahatti taluk, Gadag distrtict
Analysis at: NBSS&LUP, Regional Centre, Bengaluru **Classification:** Clayey-skeletal, mixed, isohyperthermic Typic Rhodustalfs

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand								
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)			1/3 Bar	15 Bar
0-21	Ap	74.00	8.34	17.66	9.62	11.57	15.76	23.13	13.92	20	sl	-	-
21-35	Bt	54.37	10.48	35.14	16.33	8.64	9.69	11.59	8.11	40	sc	-	-
35-56	Bc	48.37	13.46	38.17	10.96	7.69	9.17	11.28	9.27	60	sc	-	-

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	$CaCO_3$	Exchangeable bases					CEC	CEC/ Clay	Base saturation	ESP
							Ca	Mg	K	Na	Total				
	Water	$CaCl_2$	M KCl	dS m^{-1}	%	%	cmol kg^{-1}							%	%
0-21	8.18	-	-	0.30	0.56	0.94	-	-	0.31	0.55	0.86	12.19	0.69	100.00	4.51
21-35	8.17	-	-	0.30	0.52	1.29	-	-	0.19	0.84	1.03	22.18	0.63	100.00	3.79
35-56	7.95	-	-	0.46	0.48	1.99	-	-	0.24	0.58	0.82	22.94	0.60	100.00	2.53

Contd…

Series Name: Mukahadahalli (MKH), **Pedon:** R-11
Location: $15^0$22'05.4"N, $76^0$04'10.3"E, Halageri village, Koppal taluk and district
Analysis at: NBSS&LUP, Regional Centre, Bangalore. **Classification:** Clayey-skeletal, mixed, isohyperthermic Typic Haplustalfs

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand								
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)			1/3 Bar	15 Bar
0-19	Ap	65.71	8.83	25.46	9.27	9.06	14.42	21.52	11.43	70	scl	16.54	8.60
19-32	Bt	55.89	11.13	32.98	6.47	9.18	11.89	19.19	9.18	50	scl	19.24	12.78
32-58	Bt	47.95	10.41	41.63	17.52	3.78	9.13	9.55	7.97	50	sc	24.03	16.02

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	$CaCO_3$	Exchangeable bases					CEC	CEC/ Clay	Base saturation	ESP
							Ca	Mg	K	Na	Total				
	Water	$CaCl_2$	M KCl	dS m^{-1}	%	%	cmol kg^{-1}							%	%
0-19	7.38	-	-	0.09	0.2	0.00	8.97	4.32	0.26	0.22	13.77	14.84	0.58	93	1.49
19-32	7.5	-	-	0.106	0.41	0.00	15.98	3.27	0.16	0.50	19.91	20.88	0.63	95	2.38
32-58	7.46	-	-	0.173	0.49	0.00	19.71	4.53	0.23	1.32	25.79	25.76	0.62	100	5.11

Contd…

Series Name: Bisarahalli (BSR), **Pedon:** R-9
Location: $15^0$25'21.0"N, $76^0$11'42.0"E Hatti village, Koppal taluk and district
Analysis at: NBSS&LUP, Regional Centre, Bangalore**.** **Classification:** Fine, mixed, isohyperthermic Typic Paleustalfs

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand								
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)			1/3 Bar	15 Bar
0-14	Ap	70.11	9.29	20.60	22.31	15.97	11.98	9.83	10.03	20	scl	13.22	7.81
14-57	Bt1	47.27	7.52	45.20	27.04	8.28	4.61	2.10	5.24	25	sc	16.39	13.31
57-80	Bt2	41.93	8.67	49.40	21.95	6.83	4.76	4.66	3.73	30	c	21.41	15.41
80-99	Bt3	49.02	9.87	41.11	19.90	10.78	6.84	6.42	5.08	40	sc	21.82	14.24

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	$CaCO_3$	Exchangeable bases					CEC	CEC/ Clay	Base saturation	ESP
							Ca	Mg	K	Na	Total				
	Water	$CaCl_2$	M KCl	dS m^{-1}	%	%	cmol kg^{-1}							%	%
0-14	6.59	-	-	0.12	0.73	-	4.47	1.77	0.06	0.53	6.82	8.80	0.43	77.55	6.00
14-57	7.02	-	-	0.04	0.48	-	5.85	2.31	0.06	0.20	8.43	14.70	0.33	57.32	1.36
57-80	7.00	-	-	0.05	0.28	-	11.74	2.26	0.08	0.22	14.31	15.60	0.32	91.73	1.44
80-99	6.90	-	-	0.06	0.18	-	13.70	2.16	0.08	0.14	16.08	16.50	0.40	97.44	0.83

Contd…

Soil Series: Hooradhahalli (HDH), **Pedon:** RM-69
Location: $13^0$24'31"N, $76^0$33'41"E, (4D3D8G2d), Hesarahalli village, Chikkanayakanahalli taluk, Tumukura district
Analysis at: NBSS&LUP, Regional Centre, Bengaluru **Classification:** Clayey-skeletal, mixed, isohyperthermic Rhodic Paleustalfs

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand								
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)			1/3 Bar	15 Bar
0-18	Ap	72.56	15.17	12.27	4.57	8.33	17.38	23.88	18.39	35	sl	-	-
18-33	Bt1	56.29	10.75	32.96	7.88	10.24	13.41	14.43	10.34	55	scl	-	-
33-58	Bt2	46.66	10.79	42.55	10.79	9.87	8.43	9.04	8.53	55	sc	-	-
58-90	Bt3	43.09	13.63	43.27	9.90	8.25	7.32	8.76	8.87	45	c	-	-

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	$CaCO_3$	Exchangeable bases					CEC	CEC/ Clay	Base saturation	ESP
							Ca	Mg	K	Na	Total				
	Water	$CaCl_2$	M KCl	dS m^{-1}	%	%	cmol kg^{-1}							%	%
0-18	6.54	-	-	0.07	0.60	0.00	2.68	1.38	0.44	0.42	4.91	5.84	0.48	84.07	7.11
18-33	5.90	-	-	0.07	0.52	0.00	3.99	1.27	0.09	0.37	5.71	8.61	0.26	66.32	4.29
33-58	6.16	-	-	0.07	0.44	0.00	4.92	1.67	0.08	0.55	7.22	10.00	0.24	72.23	5.50
58-90	6.39	-	-	0.06	0.40	0.00	4.30	2.02	0.08	0.46	6.87	9.21	0.21	74.61	5.05

Contd…

Soil Series: Balapur (BPR), **Pedon**: RM-78
Location: $13^0$26'39"N, $76^0$35'03"E, (4D3D8G2c), Kasaba, Chikkanayakanahalli taluk, Tumakuru district
Analysis at: NBSS&LUP, Regional Centre, Bengaluru **Classification:** Clayey-skeletal, mixed, isohyperthermic Typic Rhodustalfs

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand								
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)			1/3 Bar	15 Bar
0-12	Ap	65.66	18.66	15.68	4.14	6.16	13.33	21.82	20.20	-	sl	-	-
12-34	Bt1	61.91	11.52	26.57	2.36	6.78	12.53	21.36	18.89	-	scl	-	-
34-60	Bt2	51.81	11.24	36.94	4.66	5.70	12.23	15.96	13.26	30	sc	-	-
60-84	Bt3	46.61	9.02	44.37	14.70	6.88	7.51	8.97	8.55	55	sc	-	-
84-112	Bt4	48.75	12.92	38.33	15.73	8.13	6.87	8.23	9.79	60	sc	-	-
112-127	Bc	50.98	24.74	24.28	5.25	4.63	5.15	10.92	25.03	50	scl	-	-

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	$CaCO_3$	Exchangeable bases					CEC	CEC/ Clay	Base saturation	ESP
							Ca	Mg	K	Na	Total				
	Water	$CaCl_2$	M KCl	dS m^{-1}	%	%	cmol kg^{-1}							%	%
0-12	6.64	-	-	0.03	0.56	0.00	1.90	1.32	0.21	0.03	3.46	5.45	0.35	63.48	0.51
12-34	6.99	-	-	0.02	0.48	0.00	3.66	1.90	0.07	0.08	5.70	7.82	0.29	72.93	0.96
34-60	7.29	-	-	0.02	0.40	0.00	5.13	2.08	0.11	0.20	7.52	11.19	0.30	67.18	1.75
60-84	7.50	-	-	0.02	0.32	0.00	5.83	6.36	0.13	0.23	12.55	12.38	0.28	101.43	1.83
84-112	7.54	-	-	0.02	0.24	0.00	6.02	6.59	0.11	0.25	12.96	12.77	0.33	101.49	1.97
112-127	7.90	-	-	0.02	0.20	0.00	8.04	3.62	0.07	0.32	12.04	12.47	0.51	96.56	2.55

Contd…

Series Name: Nagalapur (NGP) **Pedon :** R-10
Location: $15^0$26'38.0"N, $76^0$10'27.0" E Budashettynala village, Koppal taluk and district
Analysis at: NBSS&LUP, Regional Centre, Bangalore. **Classification:** Clayey- skeletal, mixed, isohyperthermic Typic Paleustalfs

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand								
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)			1/3 Bar	15 Bar
0-16	Ap	78.43	6.36	15.21	25.23	18.82	14.04	13.22	7.12	30	sl	9.32	5.56
16-38	Bt1	46.97	8.53	44.51	14.33	12.34	7.43	6.80	6.07	30	sc	18.70	13.79
38-58	Bt2	51.92	7.48	40.60	20.98	10.07	7.37	7.48	6.02	40	sc	17.93	13.75
58-81	Bt3	54.05	7.18	38.77	27.07	10.58	5.91	5.81	4.67	50	sc	17.92	11.87
81-104	Bt4	59.03	8.93	32.04	21.88	13.11	8.88	8.05	7.12	50	scl	16.63	10.55
104-126	BC	62.35	9.26	28.40	21.19	14.51	9.88	8.13	8.64	60	scl	15.03	10.06

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	$CaCO_3$	Exchangeable bases					CEC	CEC/ Clay	Base saturation	ESP
							Ca	Mg	K	Na	Total				
	Water	$CaCl_2$	M KCl	dS m^{-1}	%	%	cmol kg^{-1}							%	%
0-16	6.77	-	-	0.09	0.82	-	3.52	2.14	0.18	0.03	5.87	7.10	0.47	82.70	0.46
16-38	6.89	-	-	0.06	0.57	-	9.35	3.85	0.10	0.21	13.50	14.70	0.33	91.87	1.40
38-58	6.80	-	-	0.06	0.52	-	8.76	3.42	0.10	0.26	12.55	14.20	0.35	88.35	1.85
58-81	6.84	-	-	0.06	0.32	-	7.67	2.77	0.10	0.58	11.12	12.90	0.33	86.18	4.48
81-104	6.86	-	-	0.05	0.20	-	6.97	2.07	0.09	0.95	10.07	11.90	0.37	84.59	7.95
104-126	6.70	-	-	0.07	0.10	-	5.53	1.77	0.07	0.73	8.09	9.40	0.33	86.09	7.77

Contd…

Series Name: Mornal (MNL), **Pedon:** R-12
Location: $15^0$22'75"N, $76^0$05'16.1" Halageri village, Koppal Taluk and District
Analysis at: NBSS&LUP, Regional Centre, Bangalore. **Classification:** Fine mixed, isohyperthermic Rhodic Paleustalfs

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand								
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)			1/3 Bar	15 Bar
0-17	Ap	81.48	5.14	13.39	14.07	12.15	17.00	27.53	10.73	70	sl	9.64	4.93
17-31	Bt1	51.43	10.24	38.33	6.67	7.72	9.52	19.26	8.25	30	sc	23.97	11.70
31-56	Bt2	45.62	8.77	45.62	17.85	7.31	8.14	8.87	3.44	30	sc	25.94	12.45
56-104	Bt3	53.10	10.62	36.28	21.87	10.30	8.10	7.99	4.84	<30	sc	20.95	10.16
104-126	Bc	54.21	12.88	32.91	12.28	8.84	15.92	10.20	6.97	<30	scl	19.96	10.21

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	$CaCO_3$	Exchangeable bases					CEC	CEC/ Clay	Base saturation	ESP
							Ca	Mg	K	Na	Total				
	Water	$CaCl_2$	M KCl	dS m^{-1}	%	%	cmol kg^{-1}							%	%
0-17	7.89	-	-	0.137	0.33	0.00	4.92	3.35	0.35	0.45	9.07	9.01	0.67	100	5.04
17-31	8.19	-	-	0.31	0.45	0.00	7.24	5.16	0.16	0.15	12.70	13.57	0.35	94	1.12
31-56	8.2	-	-	0.414	0.53	0.00	6.49	5.32	0.11	0.13	12.05	18.55	0.41	65	0.71
56-104	8.64	-	-	0.422	0.37	0.00	6.21	4.64	0.16	0.14	11.15	15.16	0.42	74	0.95
104-126	8.71	-	-	0.436	0.2	0.00	7.06	6.31	0.09	0.33	13.79	14.52	0.44	95	2.31

Contd…

Soil Series: Ranatur (RTR), **Pedon:** TR7-3
Location: $15^0$07'58.3"N, $75^0$38'30.6"E, (4D4A3G2d), Devihal-4 microwatershed, Shirahatti taluk, Gadag district
Analysis at: NBSS&LUP, Regional Centre, Bengaluru **Classification:** Fine, mixed, isohyperthermic Rhodic Paleustalfs

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand								
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)			1/3 Bar	15 Bar
0-10	Ap	80.08	8.23	11.69	7.22	16.46	17.68	21.95	16.77	<5	sl	-	-
10-34	Bt1	44.96	12.64	42.39	3.84	11.42	10.07	11.32	8.31	<5	c	-	-
34-71	Bt2	43.35	13.02	43.63	5.20	10.40	9.77	9.77	8.21	<5	c	-	-
71-100	Bt3	47.00	10.23	42.77	10.43	12.71	9.09	7.54	7.23	<5	sc	-	-
100-138	Bt4	45.04	12.78	42.17	8.37	10.33	9.30	9.19	7.85	<5	sc	-	-
138-170	Bt5	44.63	13.79	41.58	9.19	8.99	8.26	9.40	8.78	<5	c	-	-

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	$CaCO_3$	Exchangeable bases					CEC	CEC/ Clay	Base saturation	ESP
							Ca	Mg	K	Na	Total				
	Water	$CaCl_2$	M KCl	dS m^{-1}	%	%	cmol kg^{-1}							%	%
0-10	6.47	-	-	0.03	0.49	0.00	5.61	1.33	0.13	0.01	7.07	7.07	0.60	100.00	0.41
10-34	6.46	-	-	0.03	0.57	0.00	11.69	3.19	0.14	0.01	15.03	16.87	0.40	89.00	0.06
34-71	7.23	-	-	0.03	0.53	1.20	-	-	0.16	0.01	-	17.33	0.40	100.00	0.06
71-100	7.60	-	-	0.03	0.3	0.30	-	-	0.17	0.04	-	17.21	0.40	100.00	0.23
100-138	7.88	-	-	0.03	0.6	0.42	-	-	0.17	0.15	-	16.30	0.39	100.00	0.92
138-170	8.12	-	-	0.08	0.64	0.60	-	-	0.14	0.06	-	16.87	0.41	100.00	0.36

Contd…

Series Name: Ravanaki (RNK), **Pedon:** RM-20
Location: $15^0$14'22.7"N, $75^0$57'45.8"E, Gatareddihalla village, Koppal taluk and district
Analysis at: NBSS&LUP, Regional Centre, Bangalore **Classification:** Very fine, smectitic (calc), isohyperthermic Typic Haplustepts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand								
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)			1/3 Bar	15 Bar
0-28	Ap	24.43	17.76	57.81	5.30	3.89	3.78	7.14	4.32	20	c	41.40	29.60
28-55	Bw	18.77	15.59	65.64	2.74	3.73	2.85	4.83	4.61	10	c	46.71	35.18

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	$CaCO_3$	Exchangeable bases					CEC	CEC/ Clay	Base saturation	ESP
							Ca	Mg	K	Na	Total				
	Water	$CaCl_2$	M KCl	dS m^{-1}	%	%	cmol kg^{-1}							%	%
0-28	8.86	-	-	0.483	0.63	15.48	-	-	0.86	6.27	-	37.00	0.64	-	6.78
28-55	8.61	-	-	1.4	0.23	13.68	-	-	0.68	12.27	-	53.20	0.81	-	9.22

Contd…

Series Name: Dombarahalli (DRL), **Pedon:** R-8
Location: $15^0$13'96.2"N, $75^0$57'48.6" E Ragunathanahalli village, Koppal taluk and district
Analysis at: NBSS&LUP, Regional Centre, Bangalore. **Classification:** Very fine, smectitic (calc), isohyperthermic Typic Haplusterts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand								
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)			1/3 Bar	15 Bar
0-15	Ap	28.25	19.48	52.27	4.76	4.44	4.87	8.23	5.95	-	c	39.86	27.20
15-27	BA1	21.55	20.00	58.45	3.76	2.76	3.43	6.30	5.30	-	c	46.35	34.84
27-45	Bss1	14.86	20.89	64.25	2.46	2.23	2.23	3.91	4.02	-	c	57.99	41.06
45-80	Bss2	10.42	19.04	70.54	1.74	1.97	1.27	2.78	2.66	-	c	66.36	36.24

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	$CaCO_3$	Exchangeable bases					CEC	CEC/ Clay	Base saturation	ESP
							Ca	Mg	K	Na	Total				
	Water	$CaCl_2$	M KCl	dS m^{-1}	%	%	cmol kg^{-1}							%	%
0-15	8.78	-	-	0.42	0.32	12.35	-	-	0.59	4.25	-	49.70	0.95	100.00	5.62
15-27	9.03	-	-	0.61	0.30	12.48	-	-	0.30	8.96	-	57.23	0.98	100.00	10.07
27-45	9.10	-	-	0.67	0.34	11.70	-	-	0.25	11.85	-	60.71	0.95	100.00	14.05
45-80	9.18	-	-	0.86	0.32	13.39	-	-	0.27	15.40	-	63.33	0.90	100.00	18.45

Contd…

Series Name: Gatareddihal (GRH) Pedon**:** R-7
Location: $15^0$14'20.8"N, $76^0$04'28.4" E Gudlanur village, Koppal taluk and district
Analysis at: NBSS&LUP, Regional Centre, Bangalore. **Classification:** Very fine, smectitic (calc), isohyperthermic Sodic Haplusterts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand								
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)			1/3 Bar	15 Bar
0-18	Ap	20.07	19.71	60.23	1.76	3.75	3.64	3.42	7.50	-	c	41.70	29.56
18-51	Bss1	15.11	17.47	67.42	3.16	3.04	2.25	3.38	3.27	-	c	59.43	38.52
51-80	Bss2	13.19	18.74	68.07	1.80	2.93	2.37	3.04	3.04	-	c	60.69	40.91
80-107	Bss3	17.54	19.50	62.96	2.46	4.13	3.24	4.25	3.46	-	c	57.25	37.31
107-131	BC	9.42	17.48	73.10	1.48	1.82	1.36	1.93	2.84	-	c	64.62	43.98

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	$CaCO_3$	Exchangeable bases					CEC	CEC/ Clay	Base saturation	ESP
							Ca	Mg	K	Na	Total				
	Water	$CaCl_2$	M KCl	dS m^{-1}	%	%	cmol kg^{-1}							%	%
0-18	9.08	-	-	0.23	0.33	6.89	-	-	0.70	6.36	-	63.21	1.05	100.00	7.11
18-51	9.19	-	-	0.61	0.49	9.10	-	-	0.54	14.20	-	66.05	0.98	100.00	15.98
51-80	9.27	-	-	0.56	0.29	9.36	-	-	0.49	14.75	-	65.63	0.96	100.00	17.07
80-107	9.28	-	-	0.57	0.39	9.62	-	-	0.44	14.64	-	63.95	1.02	100.00	17.49
107-131	9.04	-	-	1.08	0.31	8.32	-	-	0.52	16.40	-	68.36	0.94	100.00	17.30

Contd…

Series Name: Kavalura (KVR), **Pedon:** A2/RM-9
Location: $15^0$18'86.8"N, $75^0$56'56.3"E, Kavalura village, Koppal taluk and district
Analysis at: NBSS&LUP, Regional Centre, Bangalore. **Classification:** Fine, smectitic (calc), isohyperthermic Typic Haplusterts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand								
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)			1/3 Bar	15 Bar
0-24	Ap	36.18	17.80	46.02	7.04	7.47	6.62	9.28	5.76	10	c	28.20	18.75
24-50	Bss1	38.79	15.36	45.85	6.25	6.25	9.70	10.67	5.93	05	c	27.16	18.81
50-85	Bss2	36.80	14.66	48.54	9.63	8.23	7.03	7.58	4.33	<5	c	30.16	22.17
85-124	Bss3	22.66	17.24	60.09	4.18	3.85	5.28	5.06	4.29	<5	c	40.34	31.42

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	$CaCO_3$	Exchangeable bases					CEC	CEC/ Clay	Base saturation	ESP
							Ca	Mg	K	Na	Total				
	Water	$CaCl_2$	M KCl	dS m^{-1}	%	%	cmol kg^{-1}							%	%
0-24	8.4	-	-	0.265	0.2	8.04	-	-	0.97	0.65		43.25	0.94		0.60
24-50	9.27	-	-	0.23	0.37	8.04	-	-	0.31	3.21		41.66	0.91		3.08
50-85	9.44	-	-	0.297	0.41	8.64	-	-	0.35	6.43		43.99	0.91		5.85
85-124	9.37	-	-	0.46	0.41	11.40	-	-	0.42	7.99		51.09	0.85		6.26

Contd…

Series Name: Lakshmangudda (LGD) **Pedon:** R-2
Location: $15^0$13'08.2"N, $76^0$15'27.3" E Raghunathanahalli village, Koppal taluk and district
Analysis at: NBSS&LUP, Regional Centre, Bangalore. **Classification:** Fine, smectitic (calc), isohyperthermic Typic Haplusterts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand								
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)			1/3 Bar	15 Bar
0-17	Ap	50.60	14.29	35.11	4.53	7.86	12.49	5.18	20.54	-	sc	28.99	18.05
17-40	Bss1	40.22	16.89	42.89	3.03	7.03	9.95	13.84	6.38	-	c	34.09	23.60
40-65	Bss2	37.58	17.32	45.10	2.94	6.86	10.24	11.55	5.99	-	c	35.23	24.68
65-92	Bss3	30.69	19.33	49.97	2.09	5.06	8.03	8.25	7.26	-	c	40.92	29.53
92-124	Bss4	29.82	21.09	49.09	2.99	5.76	7.65	3.33	10.09	-	c	44.40	31.52
124-145	Bss5	28.77	22.78	48.44	2.63	5.36	7.44	8.86	4.49	-	c	43.05	30.08

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	$CaCO_3$	Exchangeable bases					CEC	CEC/ Clay	Base saturation	ESP
							Ca	Mg	K	Na	Total				
	Water	$CaCl_2$	M KCl	dS m^{-1}	%	%	cmol kg^{-1}							%	%
0-17	8.03	-	-	1.93	0.94	8.84	-	-	0.35	5.02	-	32.37	0.92	100.00	1.82
17-40	7.68	-	-	1.85	0.98	8.97	-	-	0.16	4.38	-	42.18	0.98	100.00	1.66
40-65	7.61	-	-	1.75	0.94	9.36	-	-	0.16	3.77	-	42.84	0.95	100.00	1.32
65-92	7.82	-	-	1.65	1.07	9.23	-	-	0.22	5.02	-	47.85	0.96	100.00	2.82
92-124	8.46	-	-	1.10	1.13	10.40	-	-	0.23	6.72	-	47.31	0.96	100.00	7.95
124-145	8.66	-	-	0.94	0.88	14.17	-	-	0.22	6.48	-	44.80	0.92	100.00	8.17

Contd…

Series Name: Alawandi (AWD), **Pedon:** R-16
Location: : $15^0$13'08.2"N, $76^0$15'27.3" E Neeralagi village, Koppal taluk and district
Analysis at: NBSS&LUP, Regional Centre, Bangalore. **Classification:** Fine, smectitic (calc), isohyperthermic Typic Haplusterts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand								
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)			1/3 Bar	15 Bar
0-17	Ap	20.88	25.75	53.37	3.31	4.31	4.31	5.19	3.76	-	c	33.11	25.58
17-39	Bss1	25.99	19.79	54.22	5.04	5.48	5.04	5.92	4.50	-	c	33.11	26.23
39-70	Bss2	26.76	17.80	55.44	2.93	5.31	5.53	7.37	5.63	-	c	36.15	28.67
70-111	Bss3	23.83	20.25	55.93	4.15	4.81	4.92	6.01	3.93	-	c	43.60	33.71
111-139	Bss4	21.21	20.40	58.40	2.79	4.80	4.91	5.25	3.46	-	c	46.92	36.28
139-162	Bss5	13.15	20.96	65.90	1.69	2.47	2.36	3.37	3.26	-	c	54.96	41.81

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	$CaCO_3$	Exchangeable bases					CEC	CEC/ Clay	Base saturation	ESP
							Ca	Mg	K	Na	Total				
	Water	$CaCl_2$	M KCl	dS m^{-1}	%	%	cmol kg^{-1}							%	%
0-17	8.10	-	-	0.37	0.52	9.48	-	-	0.40	1.56	-	51.30	0.96	100.00	1.22
17-39	8.60	-	-	0.24	0.52	9.60	-	-	0.14	4.60	-	52.60	0.97	100.00	3.50
39-70	8.89	-	-	0.27	0.52	9.48	-	-	0.16	2.41	-	53.90	0.97	100.00	1.78
70-111	9.10	-	-	0.35	0.54	11.28	-	-	0.15	8.95	-	54.10	0.97	100.00	6.61
111-139	9.15	-	-	0.41	0.58	10.80	-	-	0.15	7.36	-	56.10	0.96	100.00	5.24
139-162	9.16	-	-	0.50	0.50	15.48	-	-	0.19	10.19	-	61.66	0.94	100.00	6.61

Contd…

Series Name: Bardur (BDR), **Pedon:** R-4

Location: $15^0$14'31.7"N, $76^0$01'19.1"E, Moranali village, Koppal taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bangalore. **Classification:** Very fine, smectitic (calc), isohyperthermic Typic Haplusterts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand								
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)			1/3 Bar	15 Bar
0-25	Ap	21.78	22.78	55.44	2.17	3.68	4.44	6.61	4.88	-	c	36.78	26.95
25-53	BA	18.62	18.56	62.82	2.23	4.24	3.46	5.24	3.46	-	c	41.25	29.87
53-90	Bss1	15.87	18.60	65.53	2.23	1.34	4.25	3.91	4.13	-	c	44.73	33.64
90-126	Bss2	13.66	20.02	66.32	1.68	2.80	2.35	3.70	3.14	-	c	49.24	38.37
126-152	Bss3	11.64	20.79	67.57	1.69	1.81	1.81	3.50	2.82	-	c	53.50	41.90
152-210	Bss4	11.38	22.78	65.42	2.16	2.16	1.93	3.07	2.05	-	c	51.53	39.64

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	$CaCO_3$	Exchangeable bases					CEC	CEC/ Clay	Base saturation	ESP
							Ca	Mg	K	Na	Total				
	Water	$CaCl_2$	M KCl	dS m^{-1}	%	%	cmol kg^{-1}							%	%
0-25	8.73	-	22.78	0.203	0.24	5.76	-	-	0.65	4.43	-	40.56	0.73	-	4.37
25-53	9.17	-	18.56	0.295	0.45	4.92	-	-	0.32	10.47	-	74.70	1.19	-	5.61
53-90	9.27	-	18.60	0.388	0.66	6.00	-	-	0.24	10.49	-	76.20	1.16	-	5.51
90-126	9.22	-	20.02	0.608	0.57	5.88	-	-	0.21	15.93	-	77.20	1.16	-	8.25
126-152	9.21	-	20.79	0.936	0.33	6.60	-	-	0.37	20.88	-	80.90	1.20	-	10.32
152-210	9.03	-	23.21	1.47	0.33	8.16	-	-	0.24	15.34	-	73.10	1.12	-	8.39

Chapter 5

INTERPRETATION FOR LAND RESOURCE MANAGEMENT

The most important soil and site characteristics that affect the land use and conservation needs of an area are land capability, land irrigability, soil depth, soil texture, coarse fragments, available water capacity, soil slope, soil erosion, soil reaction etc. These are interpreted from the data base generated through land resource inventory and several thematic maps are generated. These would help in identifying the areas suitable for growing crops and, soil and water conservation measures and structures needed thus helping to maintain good soil health for sustained crop production. The various thematic maps generated are described below.

5.1 Land Capability Classification

Land capability classification is an interpretative grouping of soil map units (soil phases) mainly based on inherent soil characteristics, external land features and environmental factors that limit the use of land for agriculture, pasture, forestry, or other uses on a sustained basis (IARI, 1971). The land and soil characteristics used to group the land resources in an area into various land capability classes, subclasses and units are

Soil characteristics: Soil depth, soil texture, coarse fragments, soil reaction, available water capacity, calcareousness, salinity/alkali *etc*.

Land characteristics: Slope, erosion, drainage, rock outcrops.

Climate: Total rainfall and its distribution, and length of crop growing period.

The Land Capability Classification system is divided into land capability classes, subclasses and units based on the level of information available. Eight land capability classes are recognized. They are

Class I: They are very good lands that have no limitations or very few limitations that restrict their use.

Class II: They are good lands that have minor limitations and require moderate conservation practices.

Class III: They are moderately good lands that have severe limitations that reduce the choice of crops or that require special conservation practices.

Class IV: They are fairly good lands that have very severe limitations that reduce the choice of crops or that require very careful management.

Class V: Soils in these lands are not likely to erode, but have other limitations like wetness that are impractical to remove and as such not suitable for agriculture, but suitable for pasture or forestry with minor limitations.

Class VI: The lands have severe limitations that make them generally unsuitable for cultivation, but suitable for pasture or forestry with moderate limitations.

Class VII: The lands have very severe limitations that make them unsuitable for cultivation, but suitable for pasture or forestry with major limitations.

Class VIII: Soil and other miscellaneous areas (rock lands) that have very severe limitations that nearly preclude their use for any crop production, but suitable for wildlife, recreation and installation of wind mills.

The land capability subclasses are recognized based on the dominant limitations observed within a given land capability class. The subclasses are designated by adding a lower case letter like 'e', 'w', 's', or 'c' to the class numeral. The subclass "e" indicates that the main hazard is risk of erosion, "w" indicates drainage or wetness as a limitation for plant growth, "s" indicates shallow soil depth, coarse or heavy textures, calcareousness, salinity/alkalinity or gravelliness and "c" indicates limitation due to climate.

The land capability subclasses have been further subdivided into land capability units based on the kinds of limitations present in each subclass. Ten land capability units are used in grouping the soil map units. They are stony or rocky (0), erosion hazard (slope, erosion) (1), coarse texture (sand, loamy sand, sandy loam) (2), fine texture (cracking clay, silty clay) (3), slowly permeable subsoil (4), coarse underlying material (5), salinity/alkali (6), stagnation, overflow, high ground water table (7), soil depth (8) and fertility problems (9). The capability units thus identified have similar soil and land characteristics that respond similarly to a given level of management. The soils of the microwatershed have been classified upto land capability subclass level.

The 33 soil map units identified in the Lakshmapura-2 Microwatershed are grouped under three land capability classes and six land capability subclasses (Fig. 5.1).



Fig. 5.1 Land Capability map of Lakshmapura-2 Microwatershed

Entire cultivated area in the microwatershed is suitable for agriculture. Good lands (Class II) cover a maximum area of about 339 ha (74%) and distributed in the major part of the microwatershed with minor problems of soil and erosion. Moderately good lands (Class III) occupy an area of about 73 ha (16%) and distributed in the southern, central and northeastern part of the microwatershed with severe limitations of soil and erosion. An area of about 41 ha (9%) is fairly good lands and distributed in the southern part of the microwatershed with very severe limitations of soil and erosion. An area of about <1 ha (<1%) is covered by railway and 7 ha (2%) is under habitation and water body.

5.2 Soil Depth

Soil depth refers to the depth of the soil occurring above the parent material or hard rock. The depth of the soil determines the effective rooting depth for plants and in accordance with soil texture, mineralogy and gravel content, the capacity of the soil column to hold water and nutrient availability. Soil depth is one of the most important soil characteristic that is used in differentiating soils into different soil series. The soil depth classes used in identifying soils in the field are very shallow (<25 cm), shallow (25-50 cm), moderately shallow (50-75 cm), moderately deep (75-100 cm), deep (100-150 cm) and very deep (>150 cm). They were used to classify the soils into different depth classes and a soil depth map was generated (Fig. 5.2). The area extent and their geographical distribution in the microwatershed is given in Fig. 5.2.



Fig. 5.2 Soil Depth map of Lakshmapura-2 Microwatershed

Shallow soils (25-50 cm) cover about 69 ha (15%) and distributed in the eastern, central and southern part of the microwatershed. Moderately shallow (50-75 cm) soils cover an area of about 50 ha (11%) and distributed in the southern and eastern part of the microwatershed. Moderately deep (75-100 cm) soils cover an area of about 163 ha(35%) and distributed in the southern, central and northern part of the microwatershed. Deep to very deep (100- >150 cm) soils occupy an area of about 170 ha (37%) and distributed in the western, southern and northeastern part of the microwatershed.

The most productive lands cover about 170 ha (37%) where all climatically adopted long duration crops be grown. Problem soils cover an area of 69 ha (15%) where occasionally short duration crops can be grown. The probability of crop failure is very high

5.3 Surface Soil Texture

Texture is an expression to indicate the coarseness or fineness of the soil as determined by the relative proportion of primary particles of sand, silt and clay. It has a direct bearing on the structure, porosity, adhesion and consistence. The surface layer of a soil to a depth of about 25 cm is the layer that is most used by crops and plants. The surface soil textural class provides a guide to understanding soil-water retention and availability, nutrient holding capacity, infiltration, workability, drainage, physical and chemical behavior, microbial activity and crop suitability. The textural classes used for LRI were used to classify and a surface soil texture map was generated. The area extent and their geographical distribution in the microwatershed is shown in Fig 5.3.

An area of about 63 ha (14%) is loamy (sandy clay loam) at the surface and distributed in the eastern, central and southern part of the microwatershed. Clayey (sandy clay and clay) soils cover about 389 ha (85%) and are distributed in the major part of the microwatershed.

The most productive lands with respect to surface soil texture are clayey soils that (85%) have high potential for soil-water retention and availability and nutrient retention and availability, but have more problems of drainage, infiltration, workability and other physical problems. The other productive lands are loamy (14%) soils which also have high potential for soil- water retention and nutrient availability but have no drainage or other physical problems.



Fig. 5.3 Surface Soil Texture map of Lakshmapura-2 Microwatershed

5.4 Soil Gravelliness

Gravel is the term used for describing coarse fragments between 2 mm and 7.5 cm diameter and stones for those between 7.5 cm and 25 cm. The presence of gravel and stones in soil reduces the volume of soil responsible for moisture and nutrient storage, drainage, infiltration and runoff, and hinders plant growth by impeding root growth and seedling emergence, intercultural operations and farm mechanization. The gravelliness classes used in LRI were used to classify the soils and using these classes a gravelliness map was generated. The area extent and their geographic distribution in the microwatershed is shown in Fig. 5.4.

The soils that are non-gravelly (<15% gravel) cover an area of about 211 ha (46%) and distributed in the major part of the microwatershed. An area of about 201 ha (44%) is covered by gravelly (15-35% gravel) soils and are distributed in the southern, eastern and central part of the microwatershed. Very gravelly soils (35-60%) cover an area of about 40 ha (9%) and distributed in the western and central part of the microwatershed (Fig. 5.4).

The most productive lands with respect to gravelliness are found to be 46 per cent. They are non-gravelly with less than 15 per cent gravel and have potential for growing both annual and perennial crops. Very gravelly soils (35-60%) cover an area of about 9 per cent where only short duration crops can be grown.



Fig. 5.4 Soil Gravelliness map of Lakshmapura-2 Microwatershed

5.5 Available Water Capacity

The soil available water capacity (AWC) is estimated based on the ability of the soil column to retain water between the tensions of 0.33 and 15 bar in a depth of 100 cm or the entire solum if the soil is shallower. The AWC of the soils (soil series) as estimated by considering the soil texture, mineralogy, soil depth and gravel content (Sehgal *et al.*, 1990) and accordingly the soil map units were grouped into five AWC classes *viz*, very low (<50 mm/m), low (50-100 mm/m), medium (100-150 mm/m), high (150-200 mm/m) and very high (>200 mm/m) and using these values, an AWC map was generated. The area extent and their geographic distribution of different AWC classes in the microwatershed is shown in Fig. 5.5.

An area of about 72 ha (16%) in the microwatershed has soils that are very low (<50 mm/m) in available water capacity and are distributed in the southern, western and central part of the microwatershed. Maximum area of about 157 ha (34%) has soils that are low (51 to 100 mm/m) in available water capacity and are distributed in the eastern, southern and northwestern part of the microwatershed. An area of about 100 ha (22%) has soils that are medium (101-150 mm/m) in available water capacity and are distributed in the southern, central and northern part of the microwatershed. An area of about 123 ha (27%) is high to very high (151->200 mm/min) in available water capacity and distributed in the southern, central and northern part of the microwatershed.

An area of about 229 ha (50%) in the microwatershed has soils that are problematic with regard to available water capacity. Here, only short duration crops can

be grown and the probability of crop failure is very high. These areas are best put to other alternative uses. An area of about 123 ha (27%) has soils that have high potential (>200 mm/m) with regard to available water capacity where all climatically adapted long duration crops can be grown successfully.



Fig. 5.5 Soil Available Water Capacity map of Lakshmapura-2 Microwatershed

5.6 Soil Slope

Soil slope refers to the inclination of the surface of the land. It is defined by gradient, shape and length, and is an integral feature of any soil as a natural body. Slope is considered important in soil genesis, land use and land development. The length and gradient of slope influences the rate of runoff, infiltration, erosion and deposition. The soil map units were grouped into two slope classes and a slope map was generated showing the area extent and their geographic distribution of different slope classes in the microwatershed (Fig. 5.6).

Nearly level (0-1%) lands cover an area of about 9 ha (2%) and distributed in the eastern part of the microwatershed. Very gently sloping (1-3%) lands cover a maximum area of about 443 ha (96%) and distributed in the major part of the microwatershed. In all these areas, all climatically adapted annual and perennial crops can be grown without much soil and water conservation and other land development measures.



Fig. 5.6 Soil Slope map of Lakshmapura-2 Microwatershed

5.7 Soil Erosion

Soil erosion refers to the wearing away of the earth's surface by the forces of water, wind and ice involving detachment and transport of soil by raindrop impact. It is used for accelerated soil erosion resulting from disturbance of the natural landscape by burning, excessive grazing and indiscriminate felling of forest trees and tillage, all usually by man. The erosion classes showing an estimate of the current erosion status as judged from field observations in the form of rills, gullies or a carpet of gravel on the surface are recorded. Four erosion classes, viz, slight erosion (e1), moderate erosion (e2), severe erosion (e3) and very severe erosion (e4) are recognized. The soil map units were grouped into different erosion classes and a soil erosion map generated. The area extent and their spatial distribution in the microwatershed is given in Figure 5.7.

Slightly eroded lands cover an area of about 87 ha (19 %) and distributed in the southern, central and eastern part of the microwatershed. Maximum area of about 365 ha (79%) is moderately eroded (e2 class) and distributed in the major part of the microwatershed. Moderately eroded lands are problematic and need appropriate soil and water conservation and other land development measures.



Fig. 5.7 Soil Erosion map of Lakshmapura-2 Microwatershed

Chapter 6

FERTILITY STATUS

Soil fertility plays an important role in increasing crop yield. The adoption of high yielding varieties that require high amounts of nutrients has resulted in deficiency symptoms in crops and plants due to imbalanced fertilization and poor inherent fertility status, as these areas are characterized by low rainfall and high temperatures. Hence, it is necessary to know the fertility (macro and micro nutrients) status of the soils of the watersheds for assessing the kind and amount of fertilizers required for each of the crop intended to be grown. For this purpose, the surface soil samples collected from the grid points (one soil sample at every 320 m grid interval) all over the microwatershed through land resource inventory in the year 2018 were analyzed for pH, EC, organic carbon, available phosphorus and potassium, and for micronutrients like zinc, boron, copper, iron and manganese, and secondary nutrient sulphur.

Soil fertility data generated has been assessed and individual maps for all the nutrients for the microwatershed have been generated by using the Kriging method under GIS. The village/survey number wise fertility data for the microwatershed is given in Appendix-II.

6.1 Soil Reaction (pH)

The soil analysis of the Lakshmapura-2 microwatershed for soil reaction (pH) showed that slightly alkaline soils (pH 7.3-7.8) cover a maximum area of about 398 ha (87%) and distributed in the major part of the microwatershed. Strongly alkaline (pH 8.4->9.0) soils cover an area of about 54 ha (12%) and distributed in the western part of the microwatershed (Fig.6.1). Entire area in the microwatershed is alkaline in reaction.

6.2 Electrical Conductivity (EC)

The Electrical Conductivity of the soils of the entire microwatershed area is <2 dSm^{-1} (Fig 6.2) and as such the soils are non-saline.

6.3 Organic Carbon

An area of about 26 ha (6%) is low (<0.5%) and distributed in the western part of the microwatershed. Maximum area of about 233 ha (51%) is medium (0.5-0.75%) and distributed in the major part of the microwatershed. An area of about 193 ha (42%) is high (>0.75%) and distributed in the central, eastern, western and northern part of the microwatershed (Fig.6.3).



Fig.6.1 Soil Reaction (pH) map of Lakshmapura-2 Microwatershed



Fig.6.2 Electrical Conductivity (EC) map of Lakshmapura-2 Microwatershed



Fig.6.3 Soil Organic Carbon map of Lakshmapura-2 Microwatershed

6.4 Available Phosphorus

An area of about 102 ha (22%) is low (<23 kg/ha) and distributed in the southwestern and southern part of the microwatershed. Maximum area of about 314 ha (68%) is medium (23-57 kg/ha) in available phosphorus and distributed in the major part of the microwatershed. An area of about 37 ha (8%) is high (>57 kg/ha) and distributed in the western part of the microwatershed. The areas with high phosphorus content reduce 25 per cent from the recommended dose to avoid the excess application of fertilizer. Apply additional 25% phosphorus in areas where it is low and medium (Fig 6.4).

6.5 Available Potassium

Available potassium is medium (145-337 kg/ha) in 224 ha (49%) and distributed in the eastern, southern and central part of the microwatershed. Maximum area of about 228 ha (50%) is high (>337 kg/ha) and distributed in the major part of the microwatershed. The areas with high potassium content reduce 25 per cent from the recommended dose to avoid the excess application of fertilizer. Apply additional 25% potassium in areas where it is medium (Fig 6.5).

6.6 Available Sulphur

Soil analysis of available sulphur content in Lakshmapura-2 microwatershed showed that an area of about 359 ha (78%) is low and distributed in the major part of the microwatershed. An area of about 93 ha (20%) is medium (10-20 ppm) in available

sulphur content and distributed in the eastern and northern part of the microwatershed (Fig.6.6). The areas that are low and medium in available sulphur need to be applied with magnesium sulphate or gypsum or factomphos (p) fertilizer (13% sulphur) for 2-3 years for the deficiency to be corrected.

6.7 Available Boron

An area of about 85 ha (19%) is low (< 0.5ppm) in available boron and distributed in the western and northern part of the microwatershed. A maximum area of about 367 ha (80%) is medium (0.5-1.0 ppm) and distributed in the major part of the microwatershed (Fig.6.7).

6.8 Available Iron

Available iron content in the soils of the Lakshmapura-2 microwatershed is deficient (<4.5 ppm) in 380 ha (83%) and distributed in the major part of the microwatershed. An area of about 72 ha (16%) is sufficient (>4.5 ppm) and distributed in the eastern part of the microwatershed (Fig 6.8).



Fig.6.4 Soil Available Phosphorus map of Lakshmapura-2Microwatershed



Fig.6.5 Soil Available Potassium map of Lakshmapura-2 Microwatershed



Fig.6.6 Soil Available Sulphur map of Lakshmapura-2Microwatershed



Fig.6.7 Soil Available Boron map of Lakshmapura-2 Microwatershed



Fig.6.8 Soil Available Iron map of Lakshmapura-2 Microwatershed

6.9 Available Manganese

Available manganese content is sufficient (>1.0 ppm) in the entire microwatershed area (Fig 6.9).

6.10 Available Copper

Available copper content is sufficient (>0.2 ppm) in the entire microwatershed area (Fig 6.10).

6.11 Available Zinc

Available zinc content is deficient (<0.6 ppm) in an entire part of the microwatershed (Fig 6.11).



Fig.6.9 Soil Available Manganese map of Lakshmapura-2Microwatershed



Fig.6.10 Soil Available Copper map of Lakshmapura-2 Microwatershed



Fig.6.11 Soil Available Zinc map of Lakshmapura-2 Microwatershed

Chapter 7

LAND SUITABILITY FOR MAJOR CROPS

The soil and land resource units (soil phases) of Lakshmapura-2 Microwatershed were assessed for their suitability for growing food, fodder, fibre and other horticulture crops by following the procedure as outlined in FAO, 1976 and 1983. Crop requirements were developed for each of the crop from the available research data and also by referring to Naidu *et. al.* (2006) and Natarajan *et. al* (2015). The soil and land characteristics were matched with the crop requirements to arrive at the crop suitability. The soil and land characteristics table (Table 7.1) were matched with the crop requirements (Tables 7.2-7.32) to arrive at the crop suitability and the crop requirement tables are given at the end of the chapter. In FAO land suitability classification, two orders are recognized. Order S-Suitable and Order N- Not suitable. The orders have classes, subclasses and units. Order-S has three classes, Class S1- Highly Suitable, Class S2- Moderately Suitable and Class S3- Marginally Suitable. Order N has two Classes, N1- Currently not Suitable and N2-Permanently not Suitable. There are no subclasses within the Class S1 as they will have very minor or no limitations for crop growth. Classes S2, S3 and N1 are divided into subclasses based on the kinds of limitations encountered. The limitations that affect crop production are 'c' for erratic rainfall and its distribution and length of growing period (LGP), 'e' for erosion hazard, 'r' for rooting condition, 't' for lighter or heavy texture, 'g' for gravelliness or stoniness, 'n' for nutrient availability, 'l' for topography, 'm' for moisture availability, 's' for sodium 'z' for calcareousness and 'w' for drainage. These limitations are indicated as lower case letters to the class symbol. For example, moderately suitable lands with the limitations of soil depth and erosion are designated as S2re. For the microwatershed, the soil mapping units were evaluated and classified up to subclass level.

Using the above criteria, the soil map units of the microwatershed were evaluated and land suitability maps for 31 major agricultural and horticultural crops were generated. The detailed information on the kind of suitability of each of the soil phase for the crops assessed are given village/ survey number wise for the microwatershed in Appendix-III.

7.1 Land Suitability for Sorghum (*Sorghum bicolor*)

Sorghum is one of the major food crop grown in Karnataka in an area of 10.47 lakh ha in Bijapur, Gulbarga, Raichur, Bidar, Belgaum, Dharwad, Bellary, Chitradurga, Mysore and Chamarajnagar districts. The crop requirements for growing sorghum (Table 7.2) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing sorghum was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.1.

Highly suitable (Class S1) lands occupy an area of about 18 (4%) for growing sorghum and occur in the western and southern part of the microwatershed. A maximum

area of about 313 ha (68%) is moderately suitable (Class S2) for growing sorghum and distributed in the major part of the microwatershed with minor limitations of gravelliness, rooting depth and calcareousness. An area of about 121 ha (30%) is marginally suitable for growing sorghum and distributed in the southern, central and northeastern part of the microwatershed. They have moderate limitations of gravelliness, rooting depth and nutrient availability.



Fig. 7.1 Land Suitability map of Sorghum

7.2 Land Suitability for Maize (*Zea mays*)

Maize is one of the most important food crop grown in an area of 13.37 lakh ha in almost all the districts of the State. The crop requirements for growing maize (Table 7.3) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing maize was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.2.

Maximum area of about 331 ha (72%) is moderately suitable (Class S2) for growing maize and distributed in the major part of the microwatershed with minor limitations of gravelliness, rooting depth, calcareousness and texture. An area of about 121 ha (26%) is marginally suitable for growing maize and distributed in the northern, central and southwestern part of the microwatershed. They have moderate limitations of gravelliness, nutrient availability and rooting depth.



Fig. 7.2 Land Suitability map of Maize

7.3 Land Suitability for Bajra (*Pennisetum glaucum)*

Bajra is one of the major food crop grown in an area of 2.34 lakh ha in Karnataka in the northern districts. The crop requirements (Table 7.4) for growing bajra were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing bajra was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.3.

Highly suitable (Class S1) lands occupy an area of about 47 ha (10 %) for growing bajra and occur in the central and western part of the microwatershed. An area of about 295 ha (64%) is moderately suitable (Class S2) for growing bajra and distributed in the major part of the microwatershed with minor limitations of texture, rooting depth, calcareousness and gravelliness. An area of about 110 ha (24%) is marginally suitable for growing bajra and distributed in the southern, central and northeastern part of the microwatershed with moderate limitations of gravelliness, nutrient availability and rooting depth.



Fig. 7.3 Land Suitability map of Bajra

7.4 Land Suitability for Redgram (*Cajanus cajan*)

Redgram is one of the most important pulse crop grown in an area of 7.28 lakh ha in almost all the districts of the State. The crop requirements for growing redgram (Table 7.5) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing redgram was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.4.

Highly suitable (Class S1) lands occupy an area of about 18 ha (4%) for growing redgram and occur in the central and western part of the microwatershed. Maximum area of about 283 ha (62%) is moderately suitable (Class S2) for growing redgram and distributed in the major part of the microwatershed. They have minor limitations of gravelliness, calcareousness, rooting depth and texture. Marginally suitable lands (Class S3) occupy an area of about 82 ha (18%) and occur in the southern and western part of the microwatershed. They have moderate limitations of rooting depth, nutrient availability and gravelliness. Area currently not suitable (Class N1) cover about 69 ha (15%) and distributed in the southern, central and northern part of the microwatershed with severe limitation of rooting depth.



Fig. 7.4 Land Suitability map of Redgram

7.5 Land Suitability for Bengal gram (*Cicer arietinum*)

Bengal gram is one of the major pulse crop grown in an area of 9.39 lakh ha in northern Karnataka in Bijapur, Gulbarga, Raichur, Bidar, Belgaum, Dharwad and Bell ary districts. The crop requirements for growing Bengal gram (Table 7.6) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing Bengal gram was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.5.

An area of about 136 ha (30%) is moderately suitable (Class S2) for growing bengalgram and are distributed in the southern and central part of the microwatershed. They have minor limitation of calcareousness. Marginally suitable (Class S3) lands cover an area of about 316 ha (69%) and are distributed in the major part of the microwatershed. They have moderate limitations of nutrient availability, gravelliness, texture and rooting depth.



Fig. 7.5 Land Suitability map of Bengal gram

7.6 Land Suitability for Groundnut (*Arachis hypogaea*)

Groundnut is one of the major oilseed crop grown in an area of 6.54 lakh ha in Karnataka in most of the districts either as rainfed or irrigated crop. The crop requirements for growing groundnut (Table 7.7) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing groundnut was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.6.

An area of about 188 ha (41%) is moderately suitable (Class S2) for growing groundnut and distributed in the southern, eastern and northern part of the microwatershed. They have minor limitations of gravelliness, rooting depth and texture. Maximum area of about 225 ha (49%) is marginally suitable (Class S3) for growing groundnut and are distributed in the major part of the microwatershed with moderate limitations of gravelliness, rooting depth, calcareousness and texture. Area currently not suitable (Class N1) cover about 40 ha (9%) and distributed in the southern part of the microwatershed with severe limitation of nutrient availability.



Fig. 7.6 Land Suitability map of Groundnut

7.7 Land Suitability for Sunflower (*Helianthus annus*)

Sunflower is one of the most important oilseed crop grown in an area of 3.56 lakh ha in the State in all the districts. The crop requirements for growing sunflower (Table 7.8) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing sunflower was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.7.

An area of about 18 ha (4%) is highly suitable (Class S1) for growing sunflower and are distributed in the western and central part of the microwatershed. Maximum area of about 283 ha (62%) is moderately suitable (Class S2) and are distributed in the major part of the microwatershed. They have minor limitations of gravelliness, rooting depth and calcareousness. Marginally suitable (Class S3) lands occupy an area of about 41 ha (9%) and are distributed in the western, southern and eastern part of the microwatershed with moderate limitations of rooting depth and gravelliness. Area currently not suitable (Class N1) cover about 109 ha (24%) and distributed in the southern central and northern part of the microwatershed with severe limitations of nutrient availability and rooting depth.



Fig. 7.7 Land Suitability map of Sunflower

7.8 Land Suitability for Cotton (*Gossypium hirsutum*)

Cotton is one of the most important fibre crop grown in the State in about 8.75 lakh ha area in Raichur, Dharwad, Belgaum, Gulbarga, Bijapur, Bidar, Bellary, Chitradurga and Chamarajnagar districts. The crop requirements for growing cotton (Table 7.9) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing cotton was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.8.

Maximum area of about 330 ha (72%) is moderately suitable (Class S2) and are distributed in the major part of the microwatershed. They have minor limitations of rooting depth, calcareousness, texture and gravelliness. Marginally suitable (Class S3) lands occupy an area of about 122 ha (26%) and are distributed in the central, southern and northeastern part of the microwatershed with moderate limitations of rooting depth, texture, nutrient availability and gravelliness.



Fig. 7.8 Land Suitability map of Cotton

7.9 Land Suitability for Chilli (*Capsicum annuum L*)

Chilli is one of the most important spice crop grown in an area of 0.42 lakh ha in Karnataka State. The crop requirements for growing chilli (Table 7.10) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing chilli was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.9.

An area of about 18 ha (4%) is highly suitable (Class S1) for growing chilli and are distributed in the western part of the microwatershed. Maximum area of about 312 ha (68%) is moderately suitable (Class S2) and are distributed in the major part of the microwatershed. They have minor limitations of texture, rooting depth, calcareousness and gravelliness. Marginally suitable (Class S3) lands cover an area of about 81 ha (17%) and distributed in the eastern, southern and northeastern part of the microwatershed. They have moderate limitations of gravelliness and rooting depth. Area currently not suitable (Class N1) cover about 40 ha (9%) and distributed in the southern part of the microwatershed with severe limitation of nutrient availability.



Fig. 7.9 Land Suitability map of Chilli

7.10 Land Suitability for Tomato (*Solanum lycopersicum*)

Tomato is one of the most important vegetable crop grown in an area of 0.65 lakh ha in almost all the districts of the State. The crop requirements (Table 7.11) for growing tomato were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing tomato was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.10.

An area of about 18 ha (4%) is highly suitable (Class S1) for growing tomato and are distributed in the central and western part of the microwatershed. An area of about 176 ha (38%) is moderately suitable (Class S2) and are distributed in the eastern, southern and northern part of the microwatershed. They have minor limitations of rooting depth and gravelliness. Marginally suitable (Class S3) lands cover an area of about 217 ha (47%) and distributed in the major part of the microwatershed. They have moderate limitations of gravelliness, texture and rooting depth. Area currently not suitable (Class N1) cover about 40 ha (9%) and distributed in the southern part of the microwatershed with severe limitation of nutrient availability.



Fig. 7.10 Land Suitability map of Tomato

7.11 Land Suitability for Brinjal (*Solanum melongena*)

Brinjal is one of the most important vegetable crop grown in the state. The crop requirements for growing brinjal (Table 7.12) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing brinjal was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.11.

An area of about 18 ha (4%) is highly suitable (Class S1) for growing Brinjal and are distributed in the western and central part of the microwatershed. An area of about 176 ha (38%) is moderately suitable (Class S2) for growing Brinjal and distributed in the southern, eastern and northern part of the microwatershed with minor limitations of rooting depth and gravelliness. Marginally suitable (Class S3) lands cover an area of about 217 ha (47%) and occur in the major part of the microwatershed with moderate limitations of gravelliness, rooting depth and texture. Area currently not suitable (Class N1) cover about 40 ha (9%) and distributed in the southern part of the microwatershed with severe limitation of nutrient availability.



Suitability subclass	Area in ha (%)	Suitability subclass	Area in ha (%)
S1	18 (4.02)	S3t	136 (29.67)
S2g	147 (31.88)	S3rg	57 (12.33)
S2r	13 (2.92)	N1n	40 (8.8)
S2rg	16 (3.56)	Railway	0.3 (0.06)
S3g	11 (2.42)	Others*	7 (1.6)
S3r	13 (2.73)		

Fig 7.11 Land Suitability map of Brinjal

7.12 Land Suitability for Onion (*Allium cepa L.*,)

Onion is one of the most important vegetable crop grown in the state. The crop requirements for growing onion (Table 7.13) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing onion was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.12.

An area of about 194 ha (42%) is moderately suitable (Class S2) for growing Onion and distributed in the southern, eastern and western part of the microwatershed with minor limitations of gravelliness, texture and rooting depth. Marginally suitable (Class S3) lands cover a maximum area of about 217 ha (47%) and occur in the major part of the microwatershed with moderate limitations of gravelliness, rooting depth, texture and calcareousness. Area currently not suitable (Class N1) cover about 40 ha (9%) and distributed in the southern part of the microwatershed with severe limitation of nutrient availability.



Fig 7.12 Land Suitability map of Onion

7.13 Land Suitability for Bhendi (*Abelmoschus esculentus*)

Bhendi is one of the most important vegetable crop grown in the state. The crop requirements for growing bhendi (Table 7.14) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing bhendi was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.13.

An area of about 18 ha (4%) is highly suitable (Class S1) for growing Bhendi and are distributed in the western and central part of the microwatershed. Maximum area of about 312 ha (68%) is moderately suitable (Class S2) for growing Bhendi and distributed in the major part of the microwatershed with minor limitations of rooting depth, gravelliness, texture and calcareousnesss. Marginally suitable (Class S3) lands cover an area of about 81 ha (17%) and occur in the southern, central and northeastern part of the microwatershed with moderate limitations of gravelliness and rooting depth. Area currently not suitable (Class N1) cover about 40 ha (9%) and distributed in the southern part of the microwatershed with severe limitation of nutrient availability.



Fig 7.13 Land Suitability map of Bhendi

7.14 Land Suitability for Drumstick (*Moringa oleifera)*

Drumstick is one of the most important vegetable crop grown in 2403 ha area in the state. The crop requirements for growing drumstick (Table 7.15) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing drumstick was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.14.

An area of about 46 ha (10%) is highly suitable (Class S1) for growing drumstick and are distributed in the central and western part of the microwatershed. Moderately suitable (Class S2) lands cover an area of about 138 ha (30%) and are distributed in the southern, eastern and northern part of the microwatershed. They have minor limitations of rooting depth, texture, calcareousness and gravelliness. Marginally suitable (Class S3) lands cover a maximum area of about 158 ha (34%) and occur in the major part of the microwatershed. They have moderate limitations of gravelliness, calcareousness and rooting depth. Area currently not suitable (Class N1) cover about 109 ha (24%) and distributed in the northeastern, central and southern part of the microwatershed with severe limitations of nutrient availability and rooting depth.



Fig. 7.14 Land Suitability map of Drumstick

7.15 Land Suitability for Mulberry (*Morus nigra*)

Mulberry is the most important leaf crop grown for rearing silkworms in about 1.66 lakh ha in all the districts of the state. The crop requirements for growing mulberry (Table 7.16) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing mulberry was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.15.

An area of about 33 ha (7%) is highly suitable (Class S1) for growing mulberry and are distributed in the central part of the microwatershed. An area of about 140 ha (30%) is moderately suitable (Class S2) for growing mulberry and distributed in the southern, eastern and northern part of the microwatershed. They have minor limitations of gravelliness, texture and rooting depth. Marginally suitable (Class S3) lands cover an area of about 169 ha (37%) and occur in the major part of the microwatershed. They have moderate limitations of rooting depth, texture, calcareousness and gravelliness. Area currently not suitable (Class N1) cover about 109 ha (24%) and distributed in the southern, central and northeastern part of the microwatershed with severe limitations of rooting depth and nutrient availability.



Fig. 7.15 Land Suitability map of Mulberry

7.16 Land Suitability for Mango (*Mangifera indica*)

Mango is one of the most important fruit crop grown in about 1.73 lakh ha in almost all the districts of the State. The crop requirements (Table 7.17) for growing mango were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing mango was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.16.

An area of about 18 ha (4%) is highly suitable (Class S1) for growing mango and are distributed in the western part of the microwatershed. An area of about 47 ha (10%) is moderately suitable (Class S2) for growing mango and distributed in the northeastern part of the microwatershed. They have minor limitations of rooting depth and gravelliness. Marginally suitable (Class S3) lands cover an area of about 244 ha (53%) and occur in the major part of the microwatershed. They have moderate limitations of rooting depth, gravelliness and texture. Area currently not suitable (Class N1) for growing mango cover about 143 ha (31%) and distributed in the central, southern and northeastern part of the microwatershed with severe limitations of rooting depth and nutrient availability.



Fig. 7.16 Land Suitability map of Mango

7.17 Land Suitability for Sapota (*Manilkara zapota*)

Sapota is one of the most important fruit crop grown in an area of about 29373 ha in almost all the districts of the state. The crop requirements (Table 7.18) for growing sapota were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing sapota was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.17.

An area of about 46 ha (10%) is highly suitable (Class S1) for growing sapota and are distributed in the western and central part of the microwatershed. An area of about 127 ha (28%) is moderately suitable (Class S2) for growing sapota and distributed in the southern, eastern and northern part of the microwatershed. They have minor limitations of rooting depth and gravelliness. Marginally suitable (Class S3) lands cover a maximum area of about 169 ha (37%) and occur in the major part of the microwatershed. They have moderate limitations of rooting depth, gravelliness and texture. Area currently not suitable (Class N1) for growing sapota cover about 109 ha (24%) and distributed in the northeastern, southern and central part of the microwatershed with severe limitations of rooting depth and nutrient availability.



Fig. 7.17 Land Suitability map of Sapota

7.18 Land Suitability for Pomegranate (*Punica granatum*)

Pomegranate is one of the commercially grown fruit crop in about 18488 ha in Karnataka mainly in Bijapur, Bagalkot, Koppal, Gadag and Chitradurga districts. The crop requirements for growing pomegranate (Table 7.19) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing pomegranate was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.18.

An area of about 46 ha (10%) is highly suitable (Class S1) for growing pomegranate and are distributed in the central and western part of the microwatershed. Moderately suitable (Class S2) lands occupy a maximum area of about 263 ha (57%) and are distributed in the major part of the microwatershed. They have minor limitations of rooting depth, texture, calcareousness and gravelliness. Marginally suitable (Class S3) lands for growing pomegranate occupy an area of about 33 ha (7%) and are distributed in the southern and eastern part of the microwatershed with moderate limitations of rooting depth and gravelliness. Area currently not suitable (Class N1) cover about 109 ha (24%) and distributed in the southern and eastern part of the microwatershed with severe limitations of nutrient availability and rooting depth.



Fig. 7.18 Land Suitability map of Pomegranate

7.19 Land Suitability for Guava (*Psidium guajava*)

Guava is one of the most important fruit crop grown in an area of about 6558 ha in almost all the districts of the state. The crop requirements (Table 7.20) for growing guava were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing guava was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.19.

An area of about 46 ha (10%) is highly suitable (Class S1) for growing guava and are distributed in the central and western part of the microwatershed. Moderately suitable (Class S2) lands occupy an area of about 127 ha (28%) and are distributed in the southern, eastern and northwestern part of the microwatershed. They have minor limitations of rooting depth and gravelliness. Marginally suitable (Class S3) lands for growing guava occupy an area of about 169 ha (37%) and are distributed in the major part of the microwatershed with moderate limitations of gravelliness, texture, rooting depth and calcareousness. Area currently not suitable (Class N1) cover about 109 ha (24%) and distributed in the southern, northern and central part of the microwatershed with severe limitations of rooting depth and nutrient availability.



Fig. 7.19 Land Suitability map of Guava

7.20 Land Suitability for Jackfruit (*Artocarpus heterophyllus*)

Jackfruit is one of the most important fruit crop grown in 5368 ha in all the districts of the state. The crop requirements (Table.7.21) for growing jackfruit were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing jackfruit was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in figure 7.20.

An area of about 46 ha (10%) is highly suitable (Class S1) for growing jackfruit and are distributed in the central and western part of the microwatershed. Moderately suitable (Class S2) lands occupy an area of about 127 ha (28%) and are distributed in the southern, central and western part of the microwatershed. They have minor limitations of rooting depth and gravelliness. Marginally suitable (Class S3) lands for growing jackfruit occupy a maximum area of about 169 ha (37%) and are distributed in the major part of the microwatershed with moderate limitations of gravelliness, texture, calcareousness and rooting depth. Area currently not suitable (Class N1) cover about 109 ha (24%) and distributed in the southern, northern and central part of the microwatershed with severe limitations of rooting depth and nutrient availability.



Fig. 7.20 Land Suitability map of Jackfruit

7.21 Land Suitability for Jamun (*Syzygium cumini*)

Jamun is an important fruit crop grown in almost all the districts of the state. The crop requirements (Table 7.22) for growing jamun were matched with the soil-site characteristics and a land suitability map for growing jamun was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.21.

An area of about 18 ha (4%) is highly suitable (Class S1) for growing jackfruit and are distributed in the central and western part of the microwatershed. Moderately suitable (Class S2) lands occupy an area of about 47 ha (10%) and distributed in the northeastern part of the microwatershed. They have minor limitations of rooting depth and gravelliness. Marginally suitable (Class S3) lands cover a maximum area of about 277 ha (60%) and are distributed in the major part of the microwatershed with moderate limitations of rooting depth, calcareousness and gravelliness. Area currently not suitable (Class N1) cover about 109 ha (24%) and distributed in the southern, central and northeastern part of the microwatershed with severe limitations of rooting depth and nutrient availability.



Fig. 7.21 Land Suitability map of Jamun

7.22 Land Suitability for Musambi (*Citrus limetta*)

Musambi is one of the most important fruit crop grown in an area of 5446 ha in almost all the districts of the state. The crop requirements (Table 7.23) for growing musambi were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing musambi was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.22.

An area of about 46 ha (10%) is highly suitable (Class S1) for growing musambi and are distributed in the central and western part of the microwatershed. Maximum area of about 263 ha (57%) is moderately suitable (Class S2) and occur in the major part of the microwatershed. They have minor limitations of calcareousness, gravelliness and rooting depth. An area of about 33 ha (7%) is marginally suitable (Class S3) for growing musambi and are distributed in the southern and eastern part of the microwatershed with moderate limitations of gravelliness and rooting depth. Area currently not suitable (Class N1) cover about 109 ha (24%) and distributed in the southern, central and northeastern part of the microwatershed with severe limitations of rooting depth and nutrient availability.



Fig. 7.22 Land Suitability map of Musambi

7.23 Land Suitability for Lime (*Citrus sp*)

Lime is one of the most important fruit crop grown in an area of 11752 ha in almost all the districts of the State. The crop requirements for growing lime (Table 7.24) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing lime was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.23.

An area of about 46 ha (10%) is highly suitable (Class S1) for growing lime and are distributed in the central and western part of the microwatershed. Maximum area of about 263 ha (57%) is moderately suitable (Class S2) and occur in the major part of the microwatershed. They have minor limitations of calcareousness, gravelliness and rooting depth. An area of about 33 ha (7%) is marginally suitable (Class S3) for growing lime and are distributed in the southern and eastern part of the microwatershed with moderate limitations of gravelliness and rooting depth. Area currently not suitable (Class N1) cover about 109 ha (24%) and distributed in the southern, central and northeastern part of the microwatershed with severe limitations of rooting depth and nutrient availability.



Fig. 7.23 Land Suitability map of Lime

7.24 Land Suitability for Cashew (*Anacardium occidentale)*

Cashew is one of the most important nut crop grown in an area of 7052 ha in almost all the districts of the State. The crop requirements for growing cashew (Table 7.25) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing cashew was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.24.

An area of about 46 ha (10%) is highly suitable (Class S1) for growing cashew and are distributed in the western and central part of the microwatershed. An area of about 127 ha (28%) is moderately suitable (Class S2) and occur in the eastern, northern and southern part of the microwatershed. They have minor limitations of gravelliness and rooting depth. An area of about 33 ha (7%) is marginally suitable (Class S3) for growing cashew and are distributed in the southern and western part of the microwatershed with moderate limitations of gravelliness and rooting depth. Maximum area of about 245 ha (54%) is currently not suitable (Class N1) for growing cashew and distributed in the major part of the microwatershed with severe limitations of rooting depth, nutrient availability, calcareousness and texture.



Fig. 7.24 Land Suitability map of Cashew

7.25 Land Suitability for Custard Apple (*Annona reticulata*)

Custard apple is one of the most important fruit crop grown in 1426 ha in almost all the districts of the State. The crop requirements(Table 7.26) for growing custard apple were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing custard apple was generated .The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.25.

An area of about 47 ha (10%) is highly suitable (Class S1) for growing custard apple and are distributed in the western and central part of the microwatershed. A maximum area of about 295 ha (64%) is moderately suitable (Class S2) and occur in the major part of the microwatershed. They have minor limitations of gravelliness, rooting depth and calcareousness. An area of about 70 ha (15%) is marginally suitable (Class S3) for growing custard apple and are distributed in the central, northern and southern part of the microwatershed with moderate limitations of gravelliness and rooting depth. An area of about 40 ha (9%) is currently not suitable (Class N1) for growing custard apple and distributed in the southern part of the microwatershed with severe limitation of nutrient availability.



Fig. 7.25 Land Suitability map of Custard Apple

7.26 Land Suitability for Amla (*Phyllanthus emblica*)

Amla is one of the most important fruit and medicinal crop grown in an area of 151 ha and distributed in almost all the districts of the state. The crop requirements (Table 7.27) for growing amla were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing amla was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.26.

An area of about 47 ha (10%) is highly suitable (Class S1) for growing amla and are distributed in the western and central part of the microwatershed. An area of about 159 ha (34%) is moderately suitable (Class S2) and occur in the southern, eastern and southern part of the microwatershed. They have minor limitations of gravelliness and rooting depth. A maximum area of about 206 ha (45%) is marginally suitable (Class S3) for growing amla and are distributed in the southern, central and northern part of the microwatershed with moderate limitations of gravelliness, calcareousness and rooting depth. An area of about 40 ha (9%) is currently not suitable (Class N1) for growing amla and distributed in the southern part of the microwatershed with severe limitation of nutrient availability.



Fig. 7.26 Land Suitability map of Amla

7.27 Land Suitability for Tamarind (*Tamarindus indica*)

Tamarind is one of the most important spice crop grown in 14897 ha in all the districts of the state. The crop requirements (Table 7.28) for growing tamarind were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing tamarind was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.27.

An area of about 18 ha (4%) is highly suitable (Class S1) for growing tamarind and are distributed in the central and western part of the microwatershed. An area of about 47 ha (10%) is moderately suitable (Class S2) and occur in the southern and eastern part of the microwatershed. They have minor limitations of rooting depth and gravelliness. Maximum area of about 244 ha (53%) is marginally suitable (Class S3) for growing tamarind and are distributed in the major part of the microwatershed with moderate limitations of rooting depth, gravelliness and calcareousness. An area of about 143 ha (31%) is currently not suitable (Class N1) for growing tamarind and distributed in the southern, eastern and central part of the microwatershed with severe limitations of rooting depth and nutrient availability.



Fig. 7.27 Land Suitability map of Tamarind

7.28 Land Suitability for Marigold (*Tagetes erecta*)

Marigold is one of the most important flower crop grown in an area of 9108 ha in almost all the districts of the state. The crop requirements (Table 7.29) for growing marigold were matched with the soil-site characteristics and a land suitability map for growing marigold was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.28.

An area of about 18 ha (4%) is highly suitable (Class S1) for growing marigold and are distributed in the central and western part of the microwatershed. Maximum area of about 312 ha (68%) is moderately suitable (Class S2) and occur in the major part of the microwatershed. They have minor limitations of gravelliness, rooting depth, calcareousness and texture. An area of about 81 ha (17%) is marginally suitable (Class S3) for growing marigold and are distributed in the southern, central and northern part of the microwatershed with moderate limitations of gravelliness and rooting depth. An area of about 40 ha (9%) is currently not suitable (Class N1) for growing marigold and distributed in the southern part of the microwatershed with severe limitation of nutrient availability.



Fig. 7.28 Land Suitability map of Marigold

7.29 Land Suitability for Chrysanthemum (*Chrysanthemum indicum*)

Chrysanthemum is one of the most important flower crop grown in an area of 4978 ha in almost all the districts of the State. The crop requirements (Table 7.30) for growing chrysanthemum were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing chrysanthemum was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed is given in Figure 7.29.

An area of about 18 ha (4%) is highly suitable (Class S1) for growing chrysanthemum and are distributed in the central and western part of the microwatershed. Maximum area of about 312 ha (68%) is moderately suitable (Class S2) and occur in the major part of the microwatershed. They have minor limitations of gravelliness, rooting depth, calcareousness and texture. An area of about 81 ha (17%) is marginally suitable (Class S3) for growing chrysanthemum and are distributed in the southern, central and northern part of the microwatershed with moderate limitations of gravelliness and rooting depth. An area of about 40 ha (9%) is currently not suitable (Class N1) for growing chrysanthemum and distributed in the southern part of the microwatershed with severe limitation of nutrient availability.



Fig. 7.29 Land Suitability map of Chrysanthemum

7. 30 Land Suitability for Jasmine *(Jasminum sp.)*

Jasmine is one of the most important flower crop grown in an area of 803 ha in almost all the districts of the State. The crop requirements (Table 7.31) for growing jasmine were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing jasmine was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.30.

An area of about 18 ha (4%) is highly suitable (Class S1) for growing jasmine and are distributed in the western part of the microwatershed. An area of about 176 ha (38%) is moderately suitable (Class S2) and occur in the southern, western and central part of the microwatershed. They have minor limitations of gravelliness and rooting depth. Maximum area of about 217 ha (47%) is marginally suitable (Class S3) for growing jasmine and are distributed in the major part of the microwatershed with moderate limitations of gravelliness, texture and rooting depth. An area of about 40 ha (9%) is currently not suitable (Class N1) for growing jasmine and distributed in the southern part of the microwatershed with severe limitation of nutrient availability.



Fig. 7.30 Land Suitability map of Jasmine

7. 31 Land Suitability for Crossandra *(Crossandra infundibuliformis)*

Crossandra is one of the most important flower crop grown in almost all the districts of the State. The crop requirements (Table 7.32) for growing crossandra were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing crossandra was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.31.

An area of about 18 ha (4%) is highly suitable (Class S1) for growing crossandra and are distributed in the western part of the microwatershed. An area of about 176 ha (38%) is moderately suitable (Class S2) and occur in the major part of the microwatershed. They have minor limitations of gravelliness and rooting depth. An area of about 217 ha (47%) is marginally suitable (Class S3) for growing crossandra and are distributed in the major part of the microwatershed with moderate limitations of gravelliness, rooting depth and texture. An area of about 40 ha (9%) is currently not suitable (Class N1) for growing crossandra and distributed in the southern part of the microwatershed with severe limitation of nutrient availability.



Fig. 7.31 Land Suitability map of Crossandra

Table 7.1 Soil-Site Characteristics of Lakshmapura-2 Microwatershed

Soil Map Units	Climate (P) (mm)	Growing period (Days)	Drainage Class	Soil depth (cm)	Soil texture		Gravelliness		AWC (mm/m)	Slope (%)	Erosion	pH	EC (dSm^{-1})	ESP	CEC [Cmol $(p^{+})kg^{-1}$]	BS (%)
					Surf-ace	Sub-surface	Surface	Sub-surface								
ABRiB2g2	662	<90	WD	25-50	sc	sc	35-60	>35	<50	1-3	moderate	6.13	0.02	0.36	3.60	58.76
HRVhB2g2	662	<90	WD	25-50	scl	gscl	35-60	>35	<50	1-3	moderate	6.05	0.21	0.73	11.24	100
HRViA1g1	662	<90	WD	25-50	sc	gscl	15-35	>35	<50	0-1	slight	6.05	0.21	0.73	11.24	100
HRViB2g1	662	<90	WD	25-50	sc	gscl	15-35	>35	<50	1-3	moderate	6.05	0.21	0.73	11.24	100
KNHiB1g1	662	<90	WD	25-50	sc	sc	15-35	<15	51-100	1-3	slight	-	-	-	-	-
HTIiA1g1	662	<90	WD	50-75	sc	gsc	15-35	15-35	51-100		slight	7.11	0.10	0.30	0.90	147
HTIiB2	662	<90	WD	50-75	sc	gsc	-	15-35	51-100	1-3	moderate	7.11	0.10	0.30	0.90	147
HTIiB2g1	662	<90	WD	50-75	sc	gsc	15-35	15-35	51-100	1-3	moderate	7.11	0.10	0.30	0.90	147
TDHiB1	662	<90	WD	50-75	sc	sc-c	-	<15	101-150	1-3	slight	9.19	0.18	5.82	3.57	100
LKRiB2g1	662	<90	WD	50-75	sc	gsc	15-35	40-60	51-100	1-3	moderate	8.18	0.30	4.51	12.19	100
MKHhB1g1	662	<90	WD	50-75	scl	gsc	15-35	>35	51-100	1-3	slight	7.38	0.09	1.49	14.84	93
BSRiB1g1	662	<90	WD	75-10	sc	gsc	15-35	15-35	51-100	1-3	slight	6.59	0.12	6.00	8.80	77.55
HDHhB1g2	662	<90	WD	75-100	scl	gsc-gc	35-60	>35	51-100	1-3	slight	6.54	0.07	7.11	5.84	84.7
HDHhB2g1	662	<90	WD	75-100	scl	gsc-gc	15-35	>35	51-100	1-3	moderate	6.54	0.07	7.11	5.84	84.7
HDHiB2	662	<90	WD	75-100	sc	gsc-gc	-	>35	51-100	1-3	moderate	6.54	0.07	7.11	5.84	84.7
HDHiB2g1	662	<90	WD	75-100	sc	gsc-gc	15-35	>35	51-100	1-3	moderate	6.54	0.07	7.11	5.84	84.7
BPRhB2g1	662	<90	WD	100-150	scl	gsc-gc	15-35	>35	51-100	1-3	moderate	6.64	0.03	0.51	5.45	63.48
NGPiB2g2	662	<90	WD	100-150	sc	gsc	35-60	>35	51-100	1-3	moderate	6.67	0.09	0.46	7.10	82.70
MNLiB2	662	<90	WD	100-150	sc	gsc	-	15-35	101-150	1-3	moderate	7.89	0.13	5.04	9.01	100
MNLiB2g1	662	<90	WD	100-150	sc	gsc	15-35	15-35	101-150	1-3	moderate	7.89	0.13	5.04	9.01	100
HLKiB2	662	<90	WD	>150	sc	c	-	<15	151-200	1-3	moderate	-	-	-	-	-

Soil Map Units	Climate (P) (mm)	Growing period (Days)	Drainage Class	Soil depth (cm)	Soil texture		Gravelliness		AWC (mm/m)	Slope (%)	Erosion	pH	EC (dSm^{-1})	ESP	CEC [$Cmol(p^{+})kg^{-1}$]	BS (%)
					Surf-ace	Sub-surface	Surface	Sub-surface								
RTRiB2	662	<90	WD	>150	sc	c	-	-	151-200	1-3	moderate	5.08	0.03	2.06	9.21	50.50
RNKmB2g1	662	<90	MWD	50-75	c	c	15-35	<15	51-100	1-3	moderate	8.86	0.48	16.9	37.0	-
DRLmB2	662	<90	MWD	75-100	c	c	-	<15	151-200	1-3	moderate	8.78	0.42	5.62	49.70	100
DRLmB2g1	662	<90	MWD	75-100	c	c	15-35	<15	151-200	1-3	moderate	8.78	0.42	5.62	49.70	100
GRHmB1g1	662	<90	MWD	100-150	c	c	15-35	<15	>200	1-3	slight	9.08	0.23	7.11	63.21	100
GRHmB2	662	<90	MWD	100-150	c	c	-	<15	>200	1-3	moderate	9.08	0.23	7.11	63.21	100
KVRmA1	662	<90	MWD	100-150	c	c	-	-	>200	0-1	slight	8.4	0.26	0.60	43.25	-
KVRmB1	662	<90	MWD	100-150	c	c	-	-	>200	1-3	slight	8.4	0.26	0.60	43.25	-
KVRmB2g1	662	<90	MWD	100-150	c	c	15-35	-	>200	1-3	moderate	8.4	0.26	0.60	43.25	-
LGDmB1	662	<90	WD	100-150	c	c	-	<15	151-200	1-3	slight	8.03	1.85	1.66	42.18	100
AWDmB2	662	<90	MWD	>150	c	c	-	<15	>200	1-3	moderate	8.10	0.37	1.22	51.30	100
BDRmB2	662	<90	MWD	>150	c	c	-	<15	>200	1-3	moderate	8.73	0.20	4.37	40.56	-

Table 7.2 Land suitability criteria for Sorghum

Land use requirement			Rating			
Soil –site characteristics		**Unit**	**Highly suitable (S1)**	**Moderately suitable (S2)**	**Marginally suitable (S3)**	**Not suitable (N1)**
Climatic regime1	Mean temperature in growing season	°C	26–30	30–34; 24–26	34–40; 20–24	>40; <20
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristics					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, c (red), c (black)	scl, cl	ls, sl	-
	pH	1:2.5	5.5-7.8	5.0-5.5 7.8-9.0	>9.0	-
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	10-15
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

Table 7.3 Land suitability criteria for Maize

Land use requirement			Rating			
Soil –site characteristics		**Unit**	**Highly suitable (S1)**	**Moderately suitable (S2)**	**Marginally suitable (S3)**	**Not suitable (N1)**
Climatic regime	Mean temperature in growing season	°C	30-34	35-38 26-30	38-40 26-20	
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc	c (red), c (black)	ls, sl	-
	pH	1:2.5	5.5-7.8	5.0-5.5 7.8-9.0	>9.0	-
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	-
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

Table 7.4 Land suitability criteria for Bajra

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-32	33-38 24-27	39-40 20-23	<20
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm	500-750	400-500	200-400	<200
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	Sl, scl, cl,sc,c (red)	C (black)	ls	-
	pH	1:2.5	6.0-7.8	5.0-5.5 7.8-9.0	5.5-6.0 >9.0	
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	15-35	35-60	>60	
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	1-3	3-5	5-10	>10

Table 7.5 Land suitability criteria for Red gram

Land use requirement			Rating			
Soil –site characteristics		**Unit**	**Highly suitable (S1)**	**Moderately suitable (S2)**	**Marginally suitable (S3)**	**Not suitable (N1)**
Climatic regime	Mean temperature in growing season	°C	30-35(G) 20-25(AV) 15-18 (F&PS) 35-40(M)	25-30(G) 20-25 (AV) 12-15 (F&PS) 30-35(M)	20-25(G) 15-20(AV) 10-12 (F&PS) 25-30(M)	< 20 <15 <10 <25
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. Well drained	Poorly drained	Very Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, c (red)	c (black),sl, scl, cl	ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-9.0	5.0-5.5 >9.0	-
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-50	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<1.0	1.0-2.0	>2.0	
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.6 Land suitability criteria for Bengal gram

Land use requirement			Rating			
Soil –site characteristics		**Unit**	**Highly suitable (S1)**	**Moderately suitable (S2)**	**Marginally suitable (S3)**	**Not suitable (N1)**
Climatic regime	Mean temperature in growing season	°C	20–25	25–30; 15–20	30–35; 10–15	>35; <10
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. Well drained	Poorly drained	Very Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	C (black)	-	c (red), scl, cl, sc	ls, sl
	pH	1:2.5	6.0-7.8	5.0-6.0 7.8-9.0	>9.0	-
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	-
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.7 Land suitability criteria for Groundnut

Land use requirement			Rating			
Soil –site characteristics		**Unit**	**Highly suitable (S1)**	**Moderately suitable (S2)**	**Marginally suitable (S3)**	**Not suitable (N1)**
Climatic regime	Mean temperature in growing season	°C	24–33	22–24; 33–35	20–22; 35–40	<20; >40
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. Well drained	Poorly drained	Very Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl	sl,cl, sc	c (red), c (black), ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<35	35-60	>60	
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.8 Land suitability criteria for Sunflower

Land use requirement			Rating			
Soil –site characteristics		**Unit**	**Highly suitable (S1)**	**Moderately suitable (S2)**	**Marginally suitable (S3)**	**Not suitable (N1)**
Climatic regime	Mean temperature in growing season	°C	24–30	30–34; 20–24	34–38; 16–20	>38; <16
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	mod. Well drained	-	Poorly to very drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	cl, sc,c (red), c (black)	scl	ls, sl	-
	pH	1:2.5	6.5-7.8	7.8-8.4 5.5-6.5	8.4-9.0; 5.0-5.5	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.9 Land suitability criteria for Cotton

Land use requirement			Rating			
Soil –site characteristics		**Unit**	**Highly suitable (S1)**	**Moderately suitable (S2)**	**Marginally suitable (S3)**	**Not suitable (N1)**
Climatic regime	Mean temperature in growing season	°C	22-32	>32	<19	-
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well to moderately well	Poorly drained/Some what excessively drained	-	very poorly/excessively drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, c (red,black)	cl	scl	ls, sl
	pH	1:2.5	6.5-7.8	7.8-8.4	5.5-6.5 8.4->9.0	<5.5
	CEC	C mol (p+)Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	50-100	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	<3	3-5	-	>5

Table 7.10 Land suitability criteria for Chilli

Land use requirement			Rating			
Soil –site characteristics		**Unit**	**Highly suitable (S1)**	**Moderately suitable (S2)**	**Marginally suitable (S3)**	**Not suitable (N1)**
Climatic regime	Mean temperature in growing season	°C	25-32	33-35 20-25	35-38 <20	>38
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc	c (black), sl	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.11 Land suitability criteria for Tomato

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	25-28	29-32 20-24	15-19 33-36	<15 >36
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl, scl, cl, sc, c (red)	-	ls, c(black)	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.12 Land suitability criteria for Brinjal

Land use requirement			Rating			
Soil –site characteristics		**Unit**	**Highly suitable (S1)**	**Moderately suitable (S2)**	**Marginally suitable (S3)**	**Not suitable (N1)**
Climatic regime	Mean temperature in growing season	°C	Well drained	Moderately well drained	Poorly drained	V. Poorly drained
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class				
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl, scl, cl, sc c (red)	-	ls, c (black)	-
	pH	1:2.5	6.0-7.3	7.3-8.4 5.0-6.0	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	>60
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.13 Land suitability criteria for Onion

Land use requirement			Rating			
Soil –site characteristics		**Unit**	**Highly suitable (S1)**	**Moderately suitable (S2)**	**Marginally suitable (S3)**	**Not suitable (N1)**
Climatic regime	Mean temperature in growing season	°C	20-30	30-35	35-40	>40
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately /imperfectly	-	Poorly to V poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl,scl,cl,sc,c (red)	-	c (Black),ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<1.0	1.0-2.0	2.0-4.0	<4
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.14 Land suitability criteria for Bhendi

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	25-28	29-32 20-24	15-19 33-36	<15 >36
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Imperfectly drained	Poorly to very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl,sc, c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.15 Land suitability criteria for Drumstick

Land use requirement			Rating			
Soil –site characteristics		**Unit**	**Highly suitable (S1)**	**Moderately suitable (S2)**	**Marginally suitable (S3)**	**Not suitable (N1)**
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, scl, cl, c (red)	sl, c (black)	ls	s
	pH	1:2.5	6.0-7.3	5.0-5.5 7.3-7.8	5.5-6.0 7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<35	35-60	60-80	>80
Soil toxicity	Salinity (EC saturation extract)	dS/m				
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-10	-	>10

Table 7.16 Land suitability criteria for Mulberry

Land use requirement			Rating			
Soil –site characteristics		**Unit**	**Highly suitable (S1)**	**Moderately suitable (S2)**	**Marginally suitable (S3)**	**Not suitable (N1)**
Climatic regime	Mean temperature in growing season	°C	24–28	22–24; 28–32	32–38; 22–18	>38; <18
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V. Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, cl, scl	c (red)	c (black), sl, ls	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.8-8.4	7.3-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	0-35	35-60	60-80	>80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

Note: Suitability evaluation only for Mulberry leaf not for Silk worm rearing

Table 7.17 Land suitability criteria for Mango

Land use requirement			Rating			
Soil –site characteristics		**Unit**	**Highly suitable (S1)**	**Moderately suitable (S2)**	**Marginally suitable (S3)**	**Not suitable (N1)**
Climatic regime	Mean temperature in growing season	°C	28-32	24-27 33-35	36-40	20-24
	Min temp. before flowering	^{0}C	10-15	15-22	>22	-
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration	Days				
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	-	ls, sl, c (black)	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>150	100-150	75-100	<75
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.18 Land suitability criteria for Sapota

Land use requirement			Rating			
Soil –site characteristics		**Unit**	**Highly suitable (S1)**	**Moderately suitable (S2)**	**Marginally suitable (S3)**	**Not suitable (N1)**
Climatic regime	Mean temperature in growing season	°C	28-32	33-36 24-27	37-42 20-23	>42 <18
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	-	Poorly to very drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	sl	ls, c (black)	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.19 Land suitability criteria for Pomegranate

Land use requirement			Rating			
Soil –site characteristics		**Unit**	**Highly suitable (S1)**	**Moderately suitable (S2)**	**Marginally suitable (S3)**	**Not suitable (N1)**
Climatic regime	Mean temperature in growing season	°C	30-34	35-38 25-29	39-40 15-24	
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl,cl, sc, c (red)	c (black),sl	ls	-
	pH	1:2.5	5.5-7.8	7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.20 Land suitability criteria for Guava

Land use requirement			Rating			
Soil –site characteristics		**Unit**	**Highly suitable (S1)**	**Moderately suitable (S2)**	**Marginally suitable (S3)**	**Not suitable (N1)**
Climatic regime	Mean temperature in growing season	°C	28-32	33-36 24-27	37-42 20-23	
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	sl	c (black), ls	-
	pH	1:2.5	6.0-7.8	5.0-6.0	7.8-8.4	>8.4
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.21 Land suitability criteria for Jackfruit

Land use requirement			Rating			
Soil –site characteristics		**Unit**	**Highly suitable (S1)**	**Moderately suitable (S2)**	**Marginally suitable (S3)**	**Not suitable (N1)**
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. well	Poorly	V. Poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	-	sl, ls, c (black)	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	>60
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10-

Table 7.22 Land suitability criteria for Jamun

Land use requirement			Rating			
Soil –site characteristics		**Unit**	**Highly suitable (S1)**	**Moderately suitable (S2)**	**Marginally suitable (S3)**	**Not suitable (N1)**
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well	Mod. well	Poorly	V.Poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c(red)	sl, c (black)	ls	-
	pH	1:2.5	6.0-7.8	5.0-6.0	7.8-8.4	>8.4
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>150	100-150	50-100	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	>60
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

Table 7.23 Land suitability criteria for Musambi

Land use requirement			Rating			
Soil –site characteristics		**Unit**	**Highly suitable (S1)**	**Moderately suitable (S2)**	**Marginally suitable (S3)**	**Not suitable (N1)**
Climatic regime	Mean temperature in growing season	°C	28-30	31-35 24-27	36-40 20-23	>40 <20
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately drained	poorly	Very poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c	sl	ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.24 Land suitability criteria for Lime

Land use requirement			Rating			
Soil –site characteristics		**Unit**	**Highly suitable (S1)**	**Moderately suitable (S2)**	**Marginally suitable (S3)**	**Not suitable (N1)**
Climatic regime	Mean temperature in growing season	°C	28-30	31-35 24-27	36-40 20-23	>40 <20
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately drained	poorly	Very poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c	sl	ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.25 Land suitability criteria for Cashew

Land use requirement			Rating			
Soil –site characteristics		**Unit**	**Highly suitable (S1)**	**Moderately suitable (S2)**	**Marginally suitable (S3)**	**Not suitable (N1)**
Climatic regime	Mean temperature in growing season	°C	32 to 34	28 to 32; 34 to 38	24 to 28; 38 to 40	<20; >40
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	-	sl, ls	c (black)
	pH	1:2.5	5.5-6.5	5.0-5.5 6.5-7.3	7.3-7.8	>7.8
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-10	>10	-

Table 7.26 Land suitability criteria for Custard apple

Land use requirement			Rating			
Soil –site characteristics		**Unit**	**Highly suitable (S1)**	**Moderately suitable (S2)**	**Marginally suitable (S3)**	**Not suitable (N1)**
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	Scl, cl, sc, c (red), c (black)	-	Sl, ls	-
	pH	1:2.5	6.0-7.3	5.5-6.0 7.3-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15-35	35-60	60-80	-
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	>5	-

Table 7.27 Land suitability criteria for Amla

Land use requirement			Rating			
Soil –site characteristics		**Unit**	**Highly suitable (S1)**	**Moderately suitable (S2)**	**Marginally suitable (S3)**	**Not suitable (N1)**
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod.well drained	Poorly drained	V. Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	c (black)	ls, sl	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15-35	35-60	60-80	-
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

Table 7.28 Land suitability criteria for Tamarind

Land use requirement			Rating			
Soil –site characteristics		**Unit**	**Highly suitable (S1)**	**Moderately suitable (S2)**	**Marginally suitable (S3)**	**Not suitable (N1)**
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod.well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl,sc, c (red)	sl, c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>150	100-150	75-100	<75
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

Table 7.29 Land suitability criteria for Marigold

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	18-23	17-15 24-35	35-40 10-14	>40 <10
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl,scl, cl, sc, c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%				
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.30 Land suitability criteria for Chrysanthemum

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	18-23	17-15 24-35	35-40 10-14	>40 <10
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl,scl, cl, sc, c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%				
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.31 Land suitability criteria for Jasmine (irrigated)

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	18-23	17-15 24-35	35-40 10-14	-
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	sl	ls, c (black)	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%				
Erosion hazard	Slope	%	<3	3-5	5-10	>10

7.32 Land suitability criteria for Crossandra

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	-	Poorly to very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c(red)	sl,	c (black),ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%				
Erosion hazard	Slope	%	<3	3-5	5-10	>10

7.32 Land Management Units (LMUs)

The 33 soil map units identified in Lakshmapura-2 Microwatershed have been grouped into nine Land Management Units (LMUs) for the purpose of preparing a Proposed Crop Plan. Land Management Units are grouped based on the similarities in respect of the type of soil, the depth of the soil, the surface soil texture, gravel content, AWC, slope, erosion etc. and a Land Management Units map (Fig.7.32) has been generated. These Land Management Units are expected to behave similarly for a given level of management.

The map units that have been grouped into nine Land Management Units along with brief description of soil and site characteristics are given below.

LMU	Mapping unit	Soil and site characteristics
1	AWDmB2 BDRmB2 KVRmA1 KVRmB1 KVRmB2g1 LGDmB1 DRLmB2 DRLmB2g1	Moderately deep to very deep, black calcareous clay soils with slopes of 0-3%, slight to moderate erosion, gravelly (15-35%)
2	GRHmB1g1, GRHmB2	Deep, sodic clay soils with slopes of 1-3%, slight to moderate erosion, gravelly (15-35%)
3	MNLiB2, MNLiB2g1, BSRiB1g1	Moderately deep to deep, sandy clay soils with slopes of 1-3%, slight to moderate erosion, gravelly (15-35%)
4	HLKiB2, RTRiB2	Very deep, red clay soils with slopes of 1-3%, moderate erosion
5	BPRhB2g1, NGPiB2g2, HDHhB1g2, HDHhB2g1, HDHiB2, HDHiB2g1	Moderately deep to deep, red sandy clay soils with slopes of 1-3%, slight to moderate erosion, gravelly to very gravelly (15-60%)
6	HTIiA1g1, HTIiB2, HTIiB2g1, TDHiB1	Moderately shallow, red sandy clay to clay soils with slopes of 0-3%, slight to moderate erosion, gravelly (15-35%)
7	RNKmB2g1	Moderately shallow, black calcareous clay soils with slopes of 1-3%, moderate erosion, gravelly (15-35%)
8	LKRiB2g1, MKHhB1g1	Moderately shallow, red sandy clay soils with slopes of 1-3%, slight to moderate erosion, gravelly (15-35%)
9	ABRiB2g2, HRVhB2g2, HRViA1g1, HRViB2g1, KNHiB1g1	Very shallow, red gravelly sandy clay soils with slopes of 0-3%, slight to moderate erosion, gravelly to very gravelly (15-60%)



Fig 7.32 Land Management Units map of Lakshmapura-2 microwatershed

7.33 Proposed Crop Plan for Lakshmapura-2 Microwatershed

After assessing the land suitability for the 31 crops, the proposed crop plan has been prepared for the nine identified LMUs by considering only the highly (Class S1) and moderately (Class S2) suitable lands for each of the 31 crops. The resultant proposed crop plan is presented in Table 7.33.

Table 7.33 Proposed Crop Plan for Lakshmapura-2 Microwatershed

LMU	Soil Map Units	Survey Number	Field Crops/ Commercial crops	Horticulture Crops (Rainfed/Irrigated)	Suitable Interventions
1	424.AWDmB2 433.BDRmB2 386.KVRmA1 388.KVRmB1 390.KVRmB2g1 393.LGDmB1 350.DRLmB2 351.DRLmB2g1 (Moderately deep to very deep, black calcareous clay soils)	**Halageri :**322,323 **Lakamapura:**13,14,15, 23,24,25,26,27,34,44,52 ,53,54,55,56,131,132,13 3,134,135,141,142,143, 144,145,146, 150 **Thalakalla:**124,125,12 6,127,128,129,130,131, 143,144,150,151, 234,235,236	Maize, Sorghum, Sunflower, Bajra, Cotton, Red gram, Bengal gram, Soybean, Safflower, Linseed	**Fruit crops** : Pomegranate, Jamun, Lime, Musambi, Tamarind, Amla, Custard apple **Vegetables:** Drumstick, Chillies, Bhendi, Brinjal, Coriander **Flowers:** Marigold, Chrysanthemum,	Application of FYM, Biofertilizers and micronutrients, drip irrigation, mulching, suitable soil and water conservation practices
2	372.GRHmB1g1 373.GRHmB2 (Deep, sodic clay soils)	**Lakamapura:**61,64,65, 66,78,136,137, 139,140	-	**Agri-Silvi-Pasture** Ber, Aonla, Acacia sp. Dhaincha, Rhodes grass, Para grass ,Bermuda grass	Application of gypsum, iron pyrites and elemental sulphur. Addition of farm yard manures, green manures and providing subsurface drainage
3	208.MNLiB2 209.MNLiB2g1 165.BSRiB1g1 (Moderately deep to deep, sandy clay soils)	**Halageri :**332,349 **Lakamapura :**32 **Thalakalla:**155,157,15 8,159,164,166, 167	Maize, Sorghum, Groundnut, Sunflower, Bajra, Mulberry, Cotton, Red gram	**Fruit crops** : Sapota, Pomegranate, Amla, Cashew, Custard apple, Guava, Jackfruit, Lime, Musambi, **Vegetables:** Tomato, Chillies, Drumstick, Onion, Bhendi, Brinjal, Curry leaves **Flowers:** Marigold, Chrysanthemum**,** Jasmine, Crossandra	Drip irrigation, mulching, suitable soil and water conservation practices (Crescent Bunding with Catch Pit etc)
4	274.HLKiB2	**Lakamapura**	Maize, Sorghum,	**Fruit crops** : Mango, Sapota, Guava,	Drip irrigation,

LMU	Soil Map Units	Survey Number	Field Crops/ Commercial crops	Horticulture Crops (Rainfed/Irrigated)	Suitable Interventions
	288.RTRiB2 (Very deep, red clay soils)	9,10,11,12,16,30	Groundnut, Sunflower, Bajra, Mulberry, Cotton, Red gram, Horse gram, Field bean	Tamarind, Pomegranate, Lime, Musambi Cashew, Jackfruit, Jamun Custard apple, Amla **Vegetables:** Tomato, Chillies, Drumstick, Onion, Bhendi, Brinjal, Curry leaves **Flowers:** Marigold, Chrysanthemum, Jasmine, Crossandra	mulching, suitable soil and water conservation practices (Crescent Bunding with Catch Pit etc)
5	231.BPRhB2g1 266.NGPiB2g2 121.HDHhB1g2 123.HDHhB2g1 127.HDHiB2 128.HDHiB2g1 (Moderately deep to deep, red sandy clay soils)	**Halageri:**318,321,324,325,330,331 **Lakamapura:**8,18,19,20,28,33,36,37,38,39,40, 43,45,46,47,48,147, 149 **Thalakalla:**142,145,146,147,148,149,154,156, 168,169	Maize, Sorghum, Sunflower, Groundnut, Bajra, Cotton, Red gram	**Fruit crops :** Sapota, Pomegranate, Amla, Cashew, Guava, Custard apple, Jack fruit, Jamun, Lime, Musambi **Vegetables:** Tomato, Chilli, Drumstick, Onion, Bhendi, Brinjal, Curry leaves **Flowers:** Marigold, Chrysanthemum, Jasmine, Crossandra	Drip irrigation, mulching, suitable soil and water conservation practices (Crescent Bunding with Catch Pit etc)
6	96.HTIiA1g1 100.HTIiB2 101.HTIiB2g1 60.TDHiB1 (Moderately shallow, red sandy clay to clay soils)	**Halageri:**326,329,333 **Lakamapura :**35,49,50,63	Maize, Sorghum, Groundnut, Bajra, Cotton, Horse gram, Castor	**Fruit crops :** Amla, Custard apple **Vegetables:** Tomato, Chilli, Onion, Bhendi, Brinjal, Curry leaves **Flowers:** Marigold, Chrysanthemum, Jasmine, Crossandra	Drip irrigation, mulching, suitable soil and water conservation practises (Crescent Bunding with Catch Pit etc)
7	337.RNKmB2g1 (Moderately shallow, black calcareous clay soils)	**Lakamapura :**57,58,59	-	**Agri-Silvi-Pasture** Ber, Aonla, Acacia sp. Dhaincha, Rhodes grass, Para grass ,Bermuda grass	Application of gypsum, iron pyrites and elemental sulphur. Addition of farm yard manures, green manures and providing

LMU	Soil Map Units	Survey Number	Field Crops/ Commercial crops	Horticulture Crops (Rainfed/Irrigated)	Suitable Interventions
					subsurface drainage
8	54.LKRiB2g1 82.MKHhB1g1 (Moderately shallow, red sandy clay soils	**Halageri :**317 **Lakamapura :**42,67	Bajra, Groundnut, Horse gram, Castor	**Fruit crops :** Amla, Custard apple **Vegetables:** Curry leaves **Flowers:** Marigold, Chrysanthemum	Drip irrigation, mulching, suitable soil and water conservation practices (Crescent Bunding with Catch Pit etc)
9	472.ABRiB2g2 27.HRVhB2g2 29.HRViA1g1 31.HRViB2g1 467.KNHiB1g1 (Very shallow, red gravelly sandy clay soils)	**Halageri :**350 **Lakamapura :**21,22,29,31,51,60,62 **Thalakalla:**120,121,122,123,152,153,160,161,162,165	-	**Agri-Silvi-Pasture:** Custard apple, Hybrid Napier, *Styloxanthes hamata, Glyricidia, Styloxanthes scabra*	Use of short duration varieties, sowing across the slope and split application of nitrogen fertilizers

Chapter 8

SOIL HEALTH MANAGEMENT

8.1 Soil Health

Soil health is basic to plant health and plant health is basic to human and bovine health. Soil is fundamental to crop production. Without soil, no food could be produced nor would livestock be fed on a large scale. Because it is finite and fragile, soil is a precious resource that requires special care from its users.

Soil health or the capacity of the soil to function is critical to human survival. Soil health has been defined as: "the capacity of the soil to function as a living system without adverse effect on the ecosystem". Healthy soils maintain a diverse community of soil organisms that help to form beneficial symbiotic associations with plant roots, recycle essential plant nutrients, improve soil structure with positive repercussions for soil, water and nutrient holding capacity and ultimately improve crop production and also contribute to mitigating climate change by maintaining or increasing its carbon content.

Functional interactions of soil biota with organic and inorganic components, air and water determine a soil's potential to store and release nutrients, and water to plants and to promote and sustain plant growth. Thus, maintaining soil health is vital to crop production and conserve soil resource base for sustaining agriculture.

The most important characteristics of a healthy soil are

- Good soil tilth
- Sufficient soil depth
- Good water storage and good drainage
- Adequate supply, but not excess of nutrients
- Large population of beneficial organisms
- Small proportion of plant pathogens and insect pests
- Low weed pressure
- Free of chemicals and toxins that may harm the crop
- Resistance to degradation
- Resilience when unfavourable conditions occur

Characteristics of Lakshmapura-2 Microwatershed

- The soil phases with sizeable area identified in the microwatershed belonged to the soil series of HDH (105 ha), DRL (55 ha), HRV(41 ha), MNL(28ha), LGD(26 ha), GRH(24 ha), BDR(23 ha), KVR(22 ha), RNK(17 ha), HTI (16 ha), BPR(15 ha), ABR(15 ha), HLK(13 ha), KNH(13 ha), TDH(13 ha), AWD(11 ha), RTR(5 ha), NGP (4 ha), LKR (2 ha), MKH (2 ha) and BSR(1 ha)
- As per land capability classification, entire area in the microwatershed falls under arable land category (Class II, III and IV). The major limitations identified in the arable lands were soil and erosion.

- On the basis of soil reaction, an area of about 398 ha (87%) is slightly alkaline (pH 7.3-7.8) and 54 ha (12%) is strongly alkaline (pH 8.4-9.0) in reaction.

Soil Health Management

The following actions are required to improve the current land husbandry practices that provide a sound basis for the successful adoption of sustainable crop production system.

Alkaline soils

Entire area is under alkaline soils. The following actions are recommended.

1. Regular addition of organic manure, green manuring, green leaf manuring, crop residue incorporation and mulching needs to be taken up to improve the soil organic matter status.
2. Application of biofertilizers (Azospirullum, Azatobacter, Rhizobium).
3. Application of 25% extra N and P (125 % RDN&P).
4. Application of $ZnSO_4$ – 12.5 kg/ha (once in three years).
5. Application of Boron – 5 kg/ha (once in three years).

Soil Degradation

Soil erosion is one of the major factor affecting the soil health in the microwatershed. An area of about 365 ha (79%) is under moderate erosion. The areas with moderate erosion need immediate soil and water conservation and, other land development and land husbandry practices for restoring soil health.

Dissemination of Information and Communication of Benefits

Any large scale implementation of soil health management requires that supporting information is made available widely, particularly through channels familiar to farmers and extension workers. Given the very high priority attached to soil health especially by the Central Government on issuing Soil-Health Cards to all the farmers, media outlets like Regional, State and National Newspapers, Radio and Dooradarshan programs in local languages but also modern information and communication technologies such as Cellular phones and the Internet, which can be much more effective in reaching the younger farmers.

Inputs for Net Planning (Saturation Plan) and Interventions needed

Net planning in IWMP is focusing on preparation of

1. Soil and Water Conservation Treatment Plans for each plot or farm.
2. Productivity enhancement measures/ interventions for existing crops/livestock/other farm enterprises.
3. Diversification of farming mainly with perennial horticultural crops and livestock.
4. Improving livelihood opportunities and income generating activities.

In this connection, how various outputs of Sujala-III are of use in addressing these objectives of Net Planning are briefly presented below.

- **Soil Depth:** The depth of a soil decides the amount of moisture and nutrients it can hold, what crops can be taken up or not, depending on the rooting depth and the length of growing period available for raising any crop. Deeper the soil, better for a wide variety of crops. If sufficient depth is not available for growing deep rooted crops, either choose medium or short duration crops or deeper planting pits need to be opened and additional good quality soil brought from outside has to be filled into the planting pits.
- **Surface Soil Texture:** Lighter soil texture in the top soil means, better rain water infiltration, less run-off and soil moisture conservation, less capillary rise and less evaporation losses. Lighter surface textured soils are amenable to good soil tilth and are highly suitable for crops like groundnut, root vegetables (carrot, raddish, potato etc) but not ideal for crops that need stagnant water like lowland paddy. Heavy textured soils are poor in water infiltration and percolation. They are prone for sheet erosion; such soils can be improved by sand mulching. The technology that is developed by the AICRP-Dryland Agriculture, Vijayapura, Karnataka can be adopted.
- **Gravelliness:** More gravel content is favorable for run-off harvesting but poor in soil moisture storage and nutrient availability. It is a significant parameter that decides the kind of crop to be raised.
- **Land Capability Classification:** The land capability map shows the areas suitable and not suitable for agriculture and the major constraints in each of the plot/survey number. Hence, one can decide what kind of enterprise is possible in each of these units. In general, erosion and soil are the major constraints in Lakshmapura-2 Microwatershed.
- **Organic Carbon:** An area of about 26 ha (6%) is low (<0.5%), 233 ha (51%) is medium (0.5-0.75%) and 193 ha (42%) is high (>0.75%) in OC. The areas that are low and medium in OC needs to be further improved by applying farmyard manure and rotating crops with cereals and legumes or mixed cropping.
- **Promoting green manuring:** Growing of green manuring crops costs Rs. 1250/ha (green manuring seeds) and about Rs. 2000/ha towards cultivation that totals to Rs. 3250/- per ha. On the other hand, application of organic manure @ 10 tons/ha costs Rs. 5000/ha. The practice needs to be continued for 2-3 years or more. Nitrogen fertilizer needs to be supplemented by 25% in addition to the recommended level in 259 ha area where OC is less than 0.75 per cent. For example, for rainfed maize, recommended level is 50 kg N per ha and an additional 12 kg /ha needs to be applied for all the crops grown in these plots.
- **Available Phosphorus**: Available phosphorus is low (<23 kg/ha) in 102 ha (22%), medium (23-57 kg/ha) in 314 ha (68%) and high (>57 kg/ha) in 37 ha (8%) area of the microwatershed. The areas with high phosphorus content reduce 25% from the

RDF to avoid the excess application of fertilizer and apply additional 25% phosphorus in areas where it is low and medium.

- **Available Potassium:** Available potassium is medium (145-337 kg/ha) in 224 ha (49%) and high (>337 kg/ha) in 228 ha (50%) area of the microwatershed. The areas with high potassium content reduce 25% from the RDF to avoid the excess application of fertilizer and apply additional 25% potassium in areas where it is medium.
- **Available Sulphur:** Available sulphur is a very critical nutrient for oilseed crops. Available sulphur is low (<10 ppm) in 359 ha (78%) and medium (10-20 ppm) in 93 ha (20%) area of the microwatershed. Areas with low and medium in available sulphur need to be applied with magnesium sulphate or gypsum or Factamphos (p) fertitilizer (13% sulphur) for 2-3 years for the deficiency to be corrected.
- **Available Iron:** It is deficient (<4.5 ppm) in 380 ha (83%) and sufficient (>4.5 ppm) in 72 ha (16%) area of the microwatershed. To manage iron deficiency iron sulphate @ 25 kg/ha needs to be applied for 2-3 years.
- **Available Zinc:** It is deficient (<0.6 ppm) in the entire area of the microwatershed. Application of zinc sulphate @ 25kg/ha is to be followed in areas that are deficient in available zinc.
- **Available Boron:** Available boron is low in (<0.5ppm) 85 ha (19%) and medium (0.5-1.0 ppm) in 367 ha (80%) area in the microwatershed. The areas with low and medium in boron content need to be applied with sodium borate @ 10kg/ha as soil application or 0.2% borax as foliar spray to correct the deficiency.
- **Available Manganese**: It is sufficient in the entire area of the microwatershed**.**
- **Available Copper:** It is sufficient in the entire area of the microwatershed**.**
- **Soil Alkalinity:** Entire area in the microwatershed has soils that are slightly to strongly alkaline. These areas need application of gypsum and wherever calcium is in excess, iron pyrites and element sulphur can be recommended. Management practices like treating repeatedly with good quality water to drain out the excess salts and provision of subsurface drainage and growing of salt tolerant crops like Casuarina, Acasia, Neem, Ber etc, are recommended.
- **Land Suitability for various crops:** Areas that are highly, moderately and marginally suitable and not suitable for growing various crops are indicated. Along with the suitability, various constraints that are limiting the productivity are also indicated. For example, in case of cotton, gravel content, rooting depth and salinity/alkalinity are the major constraints in various plots. With suitable management interventions, the productivity can be enhanced. In order to increase water holding capacity of light textured soils, growing of green manure crops and application of organic manure is recommended.

Chapter 9

SOIL AND WATER CONSERVATION TREATMENT PLAN

For preparing soil and water conservation treatment plan for Lakshmapura-2 Microwatershed, the land resource inventory database generated under Sujala-III project has been transformed as information through series of interpretative (thematic) maps using soil phase map as a base. The various thematic maps (1:7920 scale) generated were

- Soil depth
- Surface soil texture
- Available water capacity
- Soil slope
- Soil gravelliness
- Land capability
- Present land use and land cover
- Crop suitability maps
- Rainfall map
- Hydrology
- Water Resources
- Socio-economic data
- Contour plan with existing features- network of waterways, pothissa boundaries, cut up/ minor terraces etc.
- Cadastral map (1:7920 scale)
- Satellite imagery (1:7920 scale)



Apart from these, Hand Level/ Hydro Marker/ Dumpy Level/ Total Station and Kathedars' List to be collected.

Steps for Survey and Preparation of Treatment Plan

The boundaries of Land User Groups' and Survey No. boundaries are traced in the field.

- Naming of user groups and farmers
- Identification of arable and non arable lands
- Identification of drainage lines and gullies
- Identification of non treatable areas
- Identification of priority areas in the arable lands
- Treatment plan for arable lands
- Location of water harvesting and recharge structures

9.1 Treatment Plan

The treatment plan recommended for arable lands is briefly described below.

9.1.1 Arable Land Treatment

A. BUNDING

Steps for Survey and Preparation of Treatment Plan		USER GROUP-1
Cadastral map (1:7920 scale) is enlarged to a scale of 1:2500 scale Existing network of waterways, pothissa boundaries, grass belts, natural drainage lines/ watercourse, cut ups/ terraces are marked on the cadastral map to the scale Drainage lines are demarcated into		
Small gullies	(up to 5 ha catchment)	
Medium gullies	(5-15 ha catchment)	
Ravines	(15-25 ha catchment) and	
Halla/Nala	(more than 25ha catchment)	

Measurement of Land Slope

Land slope is estimated or determined by the study and interpretation of contours or by measurement in the field using simple instruments like Hand Level or Hydromarker.





FALL: 1.5 - 0.6 = 0.9 m.

Vertical and Horizontal intervals between bunds as recommended by the Watershed Development Department.

Slope percentage	Vertical interval (m)	Corresponding Horizontal Distance (m)
2 - 3%	0.6	24
3 - 4%	0.9	21
4 - 5%	0.9	21
5 - 6%	1.2	21
6 - 7%	1.2	21

Note: i) The above intervals are maximum.

(ii) Considering the slope class and erosion status (A1… A= 0-1% slope, 1= slight erosion) the intervals have to be decided.

Bund length recording: Considering the contour plan and the existing grass belts/partitions, the bunds are aligned and lengths are measured.

Section of the Bund

Bund section is decided considering the soil texture class and gravelliness class (bg_0 .....b = loamy sand, g_0 = <15% gravel). The recommended sections for different soils are given below.

Recommended Bund Section

Top width (m)	Base width (m)	Height (m)	Side slope (Z:1;H :V)	Cross section (sq m)	Soil Texture	Remarks
0.3	0.9	0.3	01:01	0.18	Sandy loam	Vegetative bund
0.3	1.2	0.3	1.5:1	0.225	Sandy clay	
0.3	1.2	0.5	0.9:1	0.375	Red gravelly soils	
0.3	1.2	0.6	0.75:1	0.45		
0.3	1.5	0.6	01:01	0.54	Red sandy loam	
0.3	2.1	0.6	1.5:1	0.72	Very shallow clayey black soils	
0.45	2	0.75	01:01	0.92		
0.45	2.4	0.75	1.3:1	1.07	Shallow clayey black soils	
0.6	3.1	0.7	1.78:1	1.29	Medium clayey black soils	
0.5	3	0.85	1.47:1	1.49		

Formation of Trench cum Bund

Dimensions of the Borrow Pits/ Trenches to be excavated (machinery are decided considering the Bund Section).

Details of Borrow Pit dimensions are given below



Size of Borrow Pits/ Trench recommended for Trench cum Bund (by machinery)

Bund section	Bund length	Earth quantity	Pit				Berm (pit to pit)	Soil depth Class
m^2	m	m^3	L(m)	W(m)	D(m)	Quantity (m^3)	m	
0.375	6	2.25	5.85	0.85	0.45	2.24	0.15	Shallow
0.45	6	2.7	5.4	1.2	0.43	2.79	0.6	Shallow
0.45	6	2.7	5	0.85	0.65	2.76	1	Moderately Shallow
0.54	5.6	3.02	5.5	0.85	0.7	3.27	0.1	Moderately shallow
0.54	5.5	2.97	5	1.2	0.5	3	0.5	Shallow
0.72	6.2	4.46	6	1.2	0.7	5.04	0.2	Moderately shallow
0.72	5.2	3.74	5.1	0.85	0.9	3.9	0.1	Moderately deep

B. Waterways

a) Existing waterways are marked on the cadastral map (1:7920 scale) and their dimensions are recorded.

b) Considering the contour plan of the MWS, additional waterways/ modernization of the existing ones can be thought of.

c) The design details are given in the Manual.

C. Farm Ponds

Waterways and the catchment area will give an indication on the size of the Farm Pond. Location of the pond can be decided based on the contour plan/ field condition and farmers' need/desire.

D. Diversion Channel

Existing EPT/ CPT are marked on the cadastral map. Looking to the need, these can be modernized or fresh diversion channel can be proposed and runoff from this can be stored in *Gokatte*/ Recharge Ponds.

9.1.2 Non-Arable Land Treatment

Depending on the gravelliness and crops preferred by the farmers, the concerned authorities can decide appropriate treatment plan. The recommended treatments may be Contour Trench, Staggered Trench, Crescent Bund, Boulder Bund or Pebble Bund.

9.1.3 Treatment of Natural Water Course/ Drainage Lines

a) The cadastral map has to be updated as regards the network of drainge lines (gullies/ *nalas/ hallas*) and existing structures are marked to the scale and storage capacity of the existing water bodies are documented.
b) The drainage line will be demarcated into Upper Reach, Middle Reach and Lower Reach.
c) Considering the Catchment, *Nala* bed and bank conditions, suitable structures are decided.
d) Number of storage structures (Check dam/ *Nala* bund/ Percolation tank) will be decided considering the commitments and available runoff in water budgeting and quality of water in the wells and site suitability.
e) Detailed Levelling Survey using Dumpy Level / Total Station has to be carried out to arrive at the site-specific designs as shown in the Manual.
f) The location of ground water recharge structures are decided by examining the lineaments and fracture zones from geological maps.
g) Rainfall intensity data of the nearest Rain Gauge Station is considered for Hydrologic Designs.
h) Silt load to the Storage/Recharge Structures is reduced by providing vegetative, boulder and earthern checks in the natural water course. Location and design details are given in the Manual.

9.2 Recommended Soil and Water Conservation Measures

The appropriate conservation structures best suited for each of the land parcel/ survey number (Appendix-I) are selected based on the slope per cent, severity of erosion, amount of rainfall, land use and soil type. The different kinds of conservation structures recommended are

1. Graded / Strengthening of Bunds
2. Trench cum Bunds (TCB)
3. Trench cum Bunds / Strengthening
4. Crescent Bunds

A map (Fig. 9.1) showing soil and water conservation plan with different kinds of structures recommended has been prepared which shows the spatial distribution and extent of area. A maximum area of about 267 ha (58%) needs trench cum bunding, an area of about 176 ha (38%) needs graded bunding and 9 ha (2 %) requires strengthening of existing bunds/ bunding. The conservation plan prepared may be presented to all the stakeholders including farmers and after considering their suggestions, the conservation plan for the microwatershed may be finalized in a participatory approach.



Fig. 9.1 Soil and Water Conservation Plan map of Lakshmapura-2 Microwatershed

9.3 Greening of Microwatershed

As part of the greening programme in the watersheds, it is envisaged to plant a variety of horticultural and other tree plants that are edible, economical and produce lot of biomass which helps to restore the ecological balance in the watersheds. The lands that are suitable for greening programme are non-arable lands (land capability classes V, VI VII and VIII) and also the lands that are not suitable or marginally suitable for growing annual and perennial crops. The method of planting these trees is given below.

It is recommended to open the pits during the 1st week of March along the contour and heap the dugout soil on the lower side of the slope in order to harness the flowing water and facilitate weathering of soil in the pit. Exposure of soil in the pit also prevents spread of pests and diseases due to scorching sun rays. The pits should be filled with mixture of soil and organic manure during the second week of April and keep ready with sufficiently tall seedlings produced either in poly bags or in root trainer nurseries so that planting can be done during the 2nd or 3rd week of April depending on the rainfall.

The tree species suitable for the area considering rainfall, temperature and adaptability is listed below; waterlogged areas are recommended to be planted with species like Neral (*Sizyzium cumini*) and Bamboo. Dry areas are to be planted with species like Honge, Bevu, Seetaphal *etc*.

Dry Deciduous Species			Temp (°C)	Rainfall (mm)
1.	Bevu	*Azadiracta indica*	21–32	400 –1,200
2.	Tapasi	*Holoptelia integrifolia*	20-30	500 - 1000
3.	Seetaphal	*Anona Squamosa*	20-40	400 - 1000
4.	Honge	*Pongamia pinnata*	20 -50	500–2,500
5.	Kamara	*Hardwikia binata*	25 -35	400 - 1000
6.	Bage	*Albezzia lebbek*	20 - 45	500 - 1000
7.	Ficus	*Ficus bengalensis*	20 - 50	500–2,500
8.	Sisso	*Dalbargia Sissoo*	20 - 50	500 -2000
9.	Ailanthus	*Ailanthus excelsa*	20 - 50	500 - 1000
10.	Hale	*Wrightia tinctoria*	25 - 45	500 - 1000
11.	Uded	*Steriospermum chelanoides*	25 - 45	500 -2000
12.	Dhupa	*Boswella Serrata*	20 - 40	500 - 2000
13.	Nelli	*Emblica Officinalis*	20 - 50	500 -1500
14.	Honne	*Pterocarpus marsupium*	20 - 40	500 - 2000
Moist Deciduous Species			**Temp (°C)**	**Rainfall (mm)**
15.	Teak	*Tectona grandis*	20 - 50	500-5000
16.	Nandi	*Legarstroemia lanceolata*	20 - 40	500 - 4000
17.	Honne	*Pterocarpus marsupium*	20 - 40	500 - 3000
18.	Mathi	*Terminalia alata*	20 -50	500 - 2000
19.	Shivane	*Gmelina arboria*	20 -50	500 -2000
20.	Kindal	*T.Paniculata*	20 - 40	500 - 1500
21.	Beete	*Dalbargia latifolia*	20 - 40	500 - 1500
22.	Tare	*T. belerica*	20 - 40	500 - 2000
23.	Bamboo	*Bambusa arundinasia*	20 - 40	500 - 2500
24.	Bamboo	*Dendrocalamus strictus*	20 – 40	500 – 2500
25.	Muthuga	*Butea monosperma*	20 - 40	400 - 1500
26.	Hippe	*Madhuca latifolia*	20 - 40	500 - 2000
27.	Sandal	*Santalum album*	20 - 50	400 - 1000
28.	Nelli	*Emblica officinalis*	20 - 40	500 - 2000
29.	Nerale	*Sizyzium cumini*	20 - 40	500 - 2000
30.	Dhaman	*Grevia tilifolia*	20 - 40	500 - 2000
31.	Kaval	*Careya arborea*	20 - 40	500 - 2000
32.	Harada	*Terminalia chebula*	20 - 40	500 - 2000

References

1. FAO (1976) Framework for Land Evaluation, Food and Agriculture Organization, Rome.72 pp.
2. FAO (1983) Guidelines for Land Evaluation for Rainfed Agriculture, FAO, Rome, 237 pp.
3. IARI (1971) Soil Survey Manual, All India Soil and Land Use Survey Organization, IARI, New Delhi, 121 pp.
4. Katyal, J.C. and Rattan, R.K. (2003) Secondary and Micronutrients; Research Gap and future needs. Fert. News 48 (4); 9-20.
5. Naidu, L.G.K., Ramamurthy, V., Challa, O., Hegde, R. and Krishnan, P. (2006) Manual Soil Site Suitability Criteria for Major Crops, NBSS Publ. No. 129, NBSS &LUP, Nagpur, 118 pp.
6. Natarajan, A. and Dipak Sarkar (2010) Field Guide for Soil Survey, National Bureau of Soil Survey and Land Use Planning (ICAR), Nagpur, India.
7. Natarajan, A., Rajendra Hegde, Raj, J.N. and Shivananda Murthy, H.G. (2015) Implementation Manual for Sujala-III Project, Watershed Development Department, Bengaluru, Karnataka.
8. Sarma, V.A.K., Krishnan, P. and Budihal, S.L. (1987) Laboratory Manual, Tech. Bull. 23, NBSS &LUP, Nagpur.
9. Sehgal, J.L. (1990) Soil Resource Mapping of Different States of India; Why and How? National Bureau of Soil Survey and Land Use Planning, Nagpur, 49 pp.
10. Shivaprasad, C.R., R.S. Reddy, J. Sehgal and M. Velayuthum (1998) Soils of Karnataka for Optimizing Land Use, NBSS Publ. No. 47b, NBSS & LUP, Nagpur, India.
11. Soil Survey Staff (2006) Keys to Soil Taxonomy, Tenth edition, U.S. Department of Agriculture/ NRCS, Washington DC, U.S.A.
12. Soil Survey Staff (2012) Soil Survey Manual, Handbook No. 18, USDA, Washington DC, USA.

Appendix I

Lakshmapura-2 (1U1c) Microwatershed

Soil Phase Information

Village	Survey Number	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	WELLS	Land Capability	Conservation Plan
Halageri	317	2.87	MKHhB1g1	LMU-8	Moderately shallow (50-75 cm)	Sandy clay loam	Gravelly (15-35%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Slight	Eucalyptus (Eu)	Not Available	IIIs	Trench cum bunding
Halageri	318	2.08	HDHiB2g1	LMU-5	Moderately deep (75-100 cm)	Sandy clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow (Cf)	Not Available	IIes	Trench cum bunding
Halageri	321	2.46	HDHiB2	LMU-5	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut (Gn)	Not Available	IIes	Trench cum bunding
Halageri	322	8.44	LGDmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Jowar+Groundnut (Jw+Gn)	Not Available	IIs	Graded bunding
Halageri	323	5.48	LGDmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Fallow land (Fl)	Not Available	IIs	Graded bunding
Halageri	324	4.8	HDHhB2g1	LMU-5	Moderately deep (75-100 cm)	Sandy clay loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow (Cf)	1 Borewell	IIes	Trench cum bunding
Halageri	325	6.95	HDHhB2g1	LMU-5	Moderately deep (75-100 cm)	Sandy clay loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow (Cf)	Not Available	IIes	Trench cum bunding
Halageri	326	0.05	HTIiA1g1	LMU-6	Moderately shallow (50-75 cm)	Sandy clay	Gravelly (15-35%)	Low (51-100 mm/m)	Nearly level (0-1%)	Slight	Not Available (NA)	Not Available	IIs	Graded bunding
Halageri	329	2.93	HTIiA1g1	LMU-6	Moderately shallow (50-75 cm)	Sandy clay	Gravelly (15-35%)	Low (51-100 mm/m)	Nearly level (0-1%)	Slight	Current fallow (Cf)	Not Available	IIs	Graded bunding
Halageri	330	4.25	BPRhB2g1	LMU-5	Deep (100-150 cm)	Sandy clay loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut (Gn)	1 Borewell	IIes	Trench cum bunding
Halageri	331	3.44	BPRhB2g1	LMU-5	Deep (100-150 cm)	Sandy clay loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow (Cf)	Not Available	IIes	Trench cum bunding
Halageri	332	0.29	BSRiB1g1	LMU-3	Moderately deep (75-100 cm)	Sandy clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Current fallow (Cf)	Not Available	IIs	Trench cum bunding
Halageri	333	0.93	HTIiA1g1	LMU-6	Moderately shallow (50-75 cm)	Sandy clay	Gravelly (15-35%)	Low (51-100 mm/m)	Nearly level (0-1%)	Slight	Maize (Mz)	Not Available	IIs	Graded bunding
Halageri	349	0.77	BSRiB1g1	LMU-3	Moderately deep (75-100 cm)	Sandy clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Current fallow (Cf)	Not Available	IIs	Trench cum bunding
Halageri	350	4.49	HRViA1g1	LMU-9	Shallow (25-50 cm)	Sandy clay	Gravelly (15-35%)	Very Low (<50 mm/m)	Nearly level (0-1%)	Slight	Jowar (Jw)	Not Available	IIIs	Graded bunding
Halageri	351	0.005	Railway	Railway	Railway	Railway	Railway	Railway	Railway	Railway	Not Available (NA)	Not Available	Railway	Railway
Lakamapura	8	0.15	HDHhB1g2	LMU-5	Moderately deep (75-100 cm)	Sandy clay loam	Very gravelly (35-60%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Not Available (NA)	Not Available	IIs	Trench cum bunding
Lakamapura	9	2.07	HLKiB2	LMU-4	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	High (151-200 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow (Cf)	Not Available	IIes	Trench cum bunding
Lakamapura	10	2.45	HLKiB2	LMU-4	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	High (151-200 mm/m)	Very gently sloping (1-3%)	Moderate	Cluster bean (Cb)	3 Borewell	IIes	Trench cum bunding
Lakamapura	11	0.6	HLKiB2	LMU-4	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	High (151-200 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow (Cf)	Not Available	IIes	Trench cum bunding
Lakamapura	12	4.75	HLKiB2	LMU-4	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	High (151-200 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow (Cf)	Not Available	IIes	Trench cum bunding
Lakamapura	13	3.16	DRLmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	IIes	Graded bunding

Village	Survey Number	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	WELLS	Land Capability	Conservation Plan
Lakamapura	14	1.32	DRLmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow (Cf)	Not Available	IIes	Graded bunding
Lakamapura	15	1.69	DRLmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow (Cf)	Not Available	IIes	Graded bunding
Lakamapura	16	2.39	HLKiB2	LMU-4	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	High (151-200 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow (Cf)	Not Available	IIes	Trench cum bunding
Lakamapura	18	6.48	HDHhB2g1	LMU-5	Moderately deep (75-100 cm)	Sandy clay loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow (Cf)	Not Available	IIes	Trench cum bunding
Lakamapura	19	5.55	HDHhB2g1	LMU-5	Moderately deep (75-100 cm)	Sandy clay loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow (Cf)	Not Available	IIes	Trench cum bunding
Lakamapura	20	5.82	HDHhB2g1	LMU-5	Moderately deep (75-100 cm)	Sandy clay loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIes	Trench cum bunding
Lakamapura	21	2.71	ABRiB2g2	LMU-9	Shallow (25-50 cm)	Sandy clay	Very gravelly (35-60%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut (Gn)	Not Available	IIIes	Trench cum bunding
Lakamapura	22	4.56	ABRiB2g2	LMU-9	Shallow (25-50 cm)	Sandy clay	Very gravelly (35-60%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow (Cf)	Not Available	IIIes	Trench cum bunding
Lakamapura	23	6.06	DRLmB2g1	LMU-1	Moderately deep (75-100 cm)	Clay	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow+Banana+Papaya (Cf+Ba+Ppy)	1 Borewell	IIes	Graded bunding
Lakamapura	24	2.68	DRLmB2g1	LMU-1	Moderately deep (75-100 cm)	Clay	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIes	Graded bunding
Lakamapura	25	1.49	LGDmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Not Available (NA)	Not Available	IIs	Graded bunding
Lakamapura	26	3.08	DRLmB2g1	LMU-1	Moderately deep (75-100 cm)	Clay	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow (Cf)	Not Available	IIes	Graded bunding
Lakamapura	27	6.29	LGDmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Current fallow+Jowar (Cf+Jw)	1 Borewell	IIs	Graded bunding
Lakamapura	28	2.6	HDHhB2g1	LMU-5	Moderately deep (75-100 cm)	Sandy clay loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Chilli (Ch)	Not Available	IIes	Trench cum bunding
Lakamapura	29	4.02	KNHiB1g1	LMU-9	Shallow (25-50 cm)	Sandy clay	Gravelly (15-35%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Slight	Current fallow (Cf)	1 Borewell	IIIs	Trench cum bunding
Lakamapura	30	7.12	RTRiB2	LMU-4	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	High (151-200 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow (Cf)	Not Available	IIes	Trench cum bunding
Lakamapura	31	6.58	KNHiB1g1	LMU-9	Shallow (25-50 cm)	Sandy clay	Gravelly (15-35%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Slight	Current fallow+Horsegram (Cf+Hg)	1 Borewell	IIIs	Trench cum bunding
Lakamapura	32	4.98	MNLiB2g1	LMU-3	Deep (100-150 cm)	Sandy clay	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow (Cf)	1 Borewell	IIes	Trench cum bunding
Lakamapura	33	5.36	BPRhB2g1	LMU-5	Deep (100-150 cm)	Sandy clay loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow (Cf)	Not Available	IIes	Trench cum bunding
Lakamapura	34	7.49	LGDmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Bengalgram (Bg)	Not Available	IIs	Graded bunding
Lakamapura	35	6.5	TDHiB1	LMU-6	Moderately shallow (50-75 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	IIs	Trench cum bunding
Lakamapura	36	8.79	HDHiB2	LMU-5	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIes	Trench cum bunding
Lakamapura	37	4.66	HDHiB2	LMU-5	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow (Cf)	Not Available	IIes	Trench cum bunding

Village	Survey Number	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	WELLS	Land Capability	Conservation Plan
Lakamapura	38	2.01	HDHiB2	LMU-5	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow (Cf)	Not Available	IIes	Trench cum bunding
Lakamapura	39	1.12	NGPiB2g2	LMU-5	Deep (100-150 cm)	Sandy clay	Very gravelly (35-60%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow (Cf)	Not Available	IIes	Trench cum bunding
Lakamapura	40	0.69	NGPiB2g2	LMU-5	Deep (100-150 cm)	Sandy clay	Very gravelly (35-60%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow (Cf)	Not Available	IIes	Trench cum bunding
Lakamapura	42	0.27	LKRiB2g1	LMU-8	Moderately shallow (50-75 cm)	Sandy clay	Gravelly (15-35%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow (Cf)	Not Available	IIIes	Trench cum bunding
Lakamapura	43	5.44	HDHiB2g1	LMU-5	Moderately deep (75-100 cm)	Sandy clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Bengalgram+Sunflower (Bg+Sf)	Not Available	IIes	Trench cum bunding
Lakamapura	44	5.62	LGDmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Current fallow (Cf)	Not Available	IIs	Graded bunding
Lakamapura	45	0.75	HDHiB2	LMU-5	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	IIes	Trench cum bunding
Lakamapura	46	1.44	HDHiB2	LMU-5	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	IIes	Trench cum bunding
Lakamapura	47	1.99	HDHiB2	LMU-5	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow (Cf)	Not Available	IIes	Trench cum bunding
Lakamapura	48	3.52	HDHiB2g1	LMU-5	Moderately deep (75-100 cm)	Sandy clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Bengalgram+Sunflower (Bg+Sf)	Not Available	IIes	Trench cum bunding
Lakamapura	49	6.49	HTIiB2g1	LMU-6	Moderately shallow (50-75 cm)	Sandy clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow+Horsegram (Cf+Hg)	Not Available	IIes	Trench cum bunding
Lakamapura	50	4.81	HTIiB2g1	LMU-6	Moderately shallow (50-75 cm)	Sandy clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow (Cf)	Not Available	IIes	Trench cum bunding
Lakamapura	51	4.8	HRVhB2g2	LMU-9	Shallow (25-50 cm)	Sandy clay loam	Very gravelly (35-60%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Drumstick+Sunflower (Ds+Sf)	Not Available	IIIes	Trench cum bunding
Lakamapura	52	4.14	DRLmB2g1	LMU-1	Moderately deep (75-100 cm)	Clay	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow (Cf)	3 Borewell	IIes	Graded bunding
Lakamapura	53	0.61	DRLmB2g1	LMU-1	Moderately deep (75-100 cm)	Clay	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow (Cf)	1 Borewell	IIes	Graded bunding
Lakamapura	54	6.56	DRLmB2g1	LMU-1	Moderately deep (75-100 cm)	Clay	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar+Fallow land (Jw+Fl)	Not Available	IIes	Graded bunding
Lakamapura	55	1.96	DRLmB2g1	LMU-1	Moderately deep (75-100 cm)	Clay	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	1 Borewell	IIes	Graded bunding
Lakamapura	56	3.47	KVRmB2g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow (Cf)	Not Available	IIes	Graded bunding
Lakamapura	57	6.35	RNKmB2g1	LMU-7	Moderately shallow (50-75 cm)	Clay	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Bengalgram (Bg)	Not Available	IVes	Graded bunding
Lakamapura	58	6.95	RNKmB2g1	LMU-7	Moderately shallow (50-75 cm)	Clay	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Bengalgram+Current fallow (Cf)	1 Borewell	IVes	Graded bunding
Lakamapura	59	7.49	RNKmB2g1	LMU-7	Moderately shallow (50-75 cm)	Clay	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Bengalgram (Bg)	1 Borewell	IVes	Graded bunding
Lakamapura	60	2.96	HRVhB2g2	LMU-9	Shallow (25-50 cm)	Sandy clay loam	Very gravelly (35-60%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow (Cf)	Not Available	IIIes	Trench cum bunding
Lakamapura	61	2.99	GRHmB1g1	LMU-2	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Redgram (Rg)	Not Available	IVs	Graded bunding
Lakamapura	62	3.06	HRVhB2g2	LMU-9	Shallow (25-50 cm)	Sandy clay loam	Very gravelly (35-60%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	IIIes	Trench cum bunding

Village	Survey Number	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	WELLS	Land Capability	Conservation Plan
Lakamapura	63	3.42	HTIiB2g1	LMU-6	Moderately shallow (50-75 cm)	Sandy clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow (Cf)	Not Available	IIes	Trench cum bunding
Lakamapura	64	1.86	GRHmB1g1	LMU-2	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Current fallow (Cf)	Not Available	IVs	Graded bunding
Lakamapura	65	1.5	GRHmB1g1	LMU-2	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Jowar (Jw)	Not Available	IVs	Graded bunding
Lakamapura	66	2.26	GRHmB1g1	LMU-2	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Current fallow (Cf)	1 Borewell	IVs	Graded bunding
Lakamapura	67	1.26	LKRiB2g1	LMU-8	Moderately shallow (50-75 cm)	Sandy clay	Gravelly (15-35%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow (Cf)	Not Available	IIIes	Trench cum bunding
Lakamapura	78	2.91	GRHmB2	LMU-2	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow (Cf)	Not Available	IVes	Graded bunding
Lakamapura	131	2.58	BDRmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Bengalgram (Bg)	Not Available	IIes	Graded bunding
Lakamapura	132	3.32	BDRmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Bengalgram (Bg)	Not Available	IIes	Graded bunding
Lakamapura	133	6.37	BDRmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Bengalgram (Bg)	Not Available	IIes	Graded bunding
Lakamapura	134	6.59	DRLmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Bengalgram (Bg)	Not Available	IIes	Graded bunding
Lakamapura	135	6.89	DRLmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar+Bengalgram (Jw+Bg)	Not Available	IIes	Graded bunding
Lakamapura	136	1.42	GRHmB2	LMU-2	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Bengalgram (Bg)	Not Available	IVes	Graded bunding
Lakamapura	137	1.42	GRHmB2	LMU-2	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Bengalgram (Bg)	Not Available	IVes	Graded bunding
Lakamapura	139	0.48	GRHmB2	LMU-2	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow (Cf)	Not Available	IVes	Graded bunding
Lakamapura	140	2.62	GRHmB2	LMU-2	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow (Cf)	Not Available	IVes	Graded bunding
Lakamapura	141	2.26	DRLmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow (Cf)	Not Available	IIes	Graded bunding
Lakamapura	142	4.63	DRLmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Bengalgram (Bg)	Not Available	IIes	Graded bunding
Lakamapura	143	6.04	KVRmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Bengalgram (Bg)	Not Available	IIs	Graded bunding
Lakamapura	144	6.09	KVRmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Bengalgram (Bg)	Not Available	IIs	Graded bunding
Lakamapura	145	5.75	KVRmB2g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Bengalgram (Bg)	Not Available	IIes	Graded bunding
Lakamapura	146	4.13	KVRmB2g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Cluster bean (Cb)	Not Available	IIes	Graded bunding
Lakamapura	147	1.31	HDHhB1g2	LMU-5	Moderately deep (75-100 cm)	Sandy clay loam	Very gravelly (35-60%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Bengalgram (Bg)	Not Available	IIs	Trench cum bunding
Lakamapura	149	0.14	HDHhB1g2	LMU-5	Moderately deep (75-100 cm)	Sandy clay loam	Very gravelly (35-60%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Bengalgram (Bg)	Not Available	IIs	Trench cum bunding
Lakamapura	150	1.58	BDRmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Bengalgram (Bg)	Not Available	IIes	Graded bunding

Village	Survey Number	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	WELLS	Land Capability	Conservation Plan
Thalakalla	119	0.02	Railway	Railway	Railway	Railway	Railway	Railway	Railway	Railway	Not Available (NA)	Not Available	Railway	Railway
Thalakalla	120	1.38	HRViA1g1	LMU-9	Shallow (25-50 cm)	Sandy clay	Gravelly (15-35%)	Very Low (<50 mm/m)	Nearly level (0-1%)	Slight	Current fallow (Cf)	Not Available	IIIs	Graded bunding
Thalakalla	121	0.26	HRViB2g1	LMU-9	Shallow (25-50 cm)	Sandy clay	Gravelly (15-35%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	IIIes	Trench cum bunding
Thalakalla	122	0.21	HRViB2g1	LMU-9	Shallow (25-50 cm)	Sandy clay	Gravelly (15-35%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	IIIes	Trench cum bunding
Thalakalla	123	0.28	HRViB2g1	LMU-9	Shallow (25-50 cm)	Sandy clay	Gravelly (15-35%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Fallow land (Fl)	Not Available	IIIes	Trench cum bunding
Thalakalla	124	0.98	DRLmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	IIes	Graded bunding
Thalakalla	125	0.48	AWDmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow (Cf)	Not Available	IIes	Graded bunding
Thalakalla	126	0.52	AWDmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	IIes	Graded bunding
Thalakalla	127	1	AWDmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	IIes	Graded bunding
Thalakalla	128	0.03	AWDmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIes	Graded bunding
Thalakalla	129	0.66	AWDmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	IIes	Graded bunding
Thalakalla	130	1.9	AWDmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow (Cf)	Not Available	IIes	Graded bunding
Thalakalla	131	0.87	AWDmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Bengalgram (Bg)	Not Available	IIes	Graded bunding
Thalakalla	142	0.91	HDHiB2	LMU-5	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow (Cf)	Not Available	IIes	Trench cum bunding
Thalakalla	143	0.12	AWDmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIes	Graded bunding
Thalakalla	144	3.26	AWDmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Fallow land (Fl)	Not Available	IIes	Graded bunding
Thalakalla	145	2.21	HDHiB2	LMU-5	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Bengalgram (Bg)	Not Available	IIes	Trench cum bunding
Thalakalla	146	4.23	HDHiB2	LMU-5	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow (Cf)	Not Available	IIes	Trench cum bunding
Thalakalla	147	6.93	HDHiB2	LMU-5	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow (Cf)	Not Available	IIes	Trench cum bunding
Thalakalla	148	3.45	HDHiB2	LMU-5	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	IIes	Trench cum bunding
Thalakalla	149	2.31	HDHiB2	LMU-5	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow (Cf)	Not Available	IIes	Trench cum bunding
Thalakalla	150	0.84	DRLmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	IIes	Graded bunding
Thalakalla	151	6.26	DRLmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow+Jowar (Cf+Jw)	Not Available	IIes	Graded bunding
Thalakalla	152	0.13	HRViB2g1	LMU-9	Shallow (25-50 cm)	Sandy clay	Gravelly (15-35%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIIes	Trench cum bunding

Village	Survey Number	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	WELLS	Land Capability	Conservation Plan
Thalakalla	153	4.01	HRViB2g1	LMU-9	Shallow (25-50 cm)	Sandy clay	Gravelly (15-35%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow (Cf)	Not Available	IIIes	Trench cum bunding
Thalakalla	154	5.83	HDHiB2	LMU-5	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow (Cf)	Not Available	IIes	Trench cum bunding
Thalakalla	155	4.41	MNLiB2	LMU-3	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIes	Trench cum bunding
Thalakalla	156	2.31	HDHiB2	LMU-5	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow (Cf)	Not Available	IIes	Trench cum bunding
Thalakalla	157	2.57	MNLiB2g1	LMU-3	Deep (100-150 cm)	Sandy clay	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow (Cf)	Not Available	IIes	Trench cum bunding
Thalakalla	158	2.27	MNLiB2g1	LMU-3	Deep (100-150 cm)	Sandy clay	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	IIes	Trench cum bunding
Thalakalla	159	2.18	MNLiB2g1	LMU-3	Deep (100-150 cm)	Sandy clay	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	3 Borewell	IIes	Trench cum bunding
Thalakalla	160	1.86	HRViB2g1	LMU-9	Shallow (25-50 cm)	Sandy clay	Gravelly (15-35%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow (Cf)	Not Available	IIIes	Trench cum bunding
Thalakalla	161	0.03	HRViA1g1	LMU-9	Shallow (25-50 cm)	Sandy clay	Gravelly (15-35%)	Very Low (<50 mm/m)	Nearly level (0-1%)	Slight	Not Available (NA)	Not Available	IIIs	Graded bunding
Thalakalla	162	6	HRViB2g1	LMU-9	Shallow (25-50 cm)	Sandy clay	Gravelly (15-35%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Sunflower (Sf)	Not Available	IIIes	Trench cum bunding
Thalakalla	163	3.47	BPRhB2g1	LMU-5	Deep (100-150 cm)	Sandy clay loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Sunflower (Sf)	Not Available	IIes	Trench cum bunding
Thalakalla	164	3.39	MNLiB2g1	LMU-3	Deep (100-150 cm)	Sandy clay	Gravelly (15-35%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow (Cf)	Not Available	IIes	Trench cum bunding
Thalakalla	165	4.7	KNHiB1g1	LMU-9	Shallow (25-50 cm)	Sandy clay	Gravelly (15-35%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Slight	Current fallow (Cf)	Not Available	IIIs	Trench cum bunding
Thalakalla	166	3.09	MNLiB2	LMU-3	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow (Cf)	Not Available	IIes	Trench cum bunding
Thalakalla	167	2.73	MNLiB2	LMU-3	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow (Cf)	Not Available	IIes	Trench cum bunding
Thalakalla	168	4.7	HDHiB2g1	LMU-5	Moderately deep (75-100 cm)	Sandy clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow (Cf)	Not Available	IIes	Trench cum bunding
Thalakalla	169	1.72	HDHiB2g1	LMU-5	Moderately deep (75-100 cm)	Sandy clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow (Cf)	Not Available	IIes	Trench cum bunding
Thalakalla	234	2.31	BDRmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow (Cf)	Not Available	IIes	Graded bunding
Thalakalla	235	2.2	BDRmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIes	Graded bunding
Thalakalla	236	0.97	BDRmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow (Cf)	Not Available	IIes	Graded bunding

Appendix II

Lakshmapura-2 (1U1c) Microwatershed

Soil Fertility Information

Village	Survey Number	Soil Reaction	Salinity	Organic Carbon	Available Phosphorus	Available Potassium	Available Sulphur	Available Boron	Available Iron	Available Manganese	Available Copper	Available Zinc
Halageri	317	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	Medium (0.5 – 0.75 %)	Medium (23 – 57 kg/ha)	Medium (145 – 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Halageri	318	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	Medium (0.5 – 0.75 %)	Low (< 23 kg/ha)	Medium (145 – 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Halageri	321	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	High (> 0.75 %)	Low (< 23 kg/ha)	Medium (145 – 337 kg/ha)	Medium (10 – 20 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Halageri	322	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	Medium (0.5 – 0.75 %)	Medium (23 – 57 kg/ha)	Medium (145 – 337 kg/ha)	Medium (10 – 20 ppm)	Medium (0.5 – 1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Halageri	323	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	Medium (0.5 – 0.75 %)	Medium (23 – 57 kg/ha)	Medium (145 – 337 kg/ha)	Medium (10 – 20 ppm)	Medium (0.5 – 1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Halageri	324	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	Medium (0.5 – 0.75 %)	Medium (23 – 57 kg/ha)	Medium (145 – 337 kg/ha)	Medium (10 – 20 ppm)	Medium (0.5 – 1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Halageri	325	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	Medium (0.5 – 0.75 %)	Medium (23 – 57 kg/ha)	Medium (145 – 337 kg/ha)	Medium (10 – 20 ppm)	Medium (0.5 – 1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Halageri	326	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	Medium (0.5 – 0.75 %)	Medium (23 – 57 kg/ha)	Medium (145 – 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Halageri	329	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	Medium (0.5 – 0.75 %)	Medium (23 – 57 kg/ha)	Medium (145 – 337 kg/ha)	Medium (10 – 20 ppm)	Medium (0.5 – 1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Halageri	330	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	Medium (0.5 – 0.75 %)	Medium (23 – 57 kg/ha)	Medium (145 – 337 kg/ha)	Medium (10 – 20 ppm)	Medium (0.5 – 1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Halageri	331	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	Medium (0.5 – 0.75 %)	Medium (23 – 57 kg/ha)	Medium (145 – 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Halageri	332	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	Medium (0.5 – 0.75 %)	Medium (23 – 57 kg/ha)	Medium (145 – 337 kg/ha)	Medium (10 – 20 ppm)	Medium (0.5 – 1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Halageri	333	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	Medium (0.5 – 0.75 %)	Medium (23 – 57 kg/ha)	Medium (145 – 337 kg/ha)	Medium (10 – 20 ppm)	Medium (0.5 – 1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Halageri	349	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	Medium (0.5 – 0.75 %)	Medium (23 – 57 kg/ha)	Medium (145 – 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Halageri	350	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	Medium (0.5 – 0.75 %)	Medium (23 – 57 kg/ha)	Medium (145 – 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Halageri	351	Railway	Railway	Railway	Railway	Railway	Railway	Railway	Railway	Railway	Railway	Railway
Lakamapura	8	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	High (> 0.75 %)	High (> 57 kg/ha)	High (> 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	9	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	High (> 0.75 %)	High (> 57 kg/ha)	High (> 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	10	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	High (> 0.75 %)	High (> 57 kg/ha)	High (> 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	11	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	High (> 0.75 %)	High (> 57 kg/ha)	High (> 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	12	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	High (> 0.75 %)	High (> 57 kg/ha)	High (> 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	13	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	High (> 0.75 %)	High (> 57 kg/ha)	High (> 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)

Village	Survey Number	Soil Reaction	Salinity	Organic Carbon	Available Phosphorus	Available Potassium	Available Sulphur	Available Boron	Available Iron	Available Manganese	Available Copper	Available Zinc
Lakamapura	14	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	High (> 0.75 %)	High (> 57 kg/ha)	High (> 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	15	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	High (> 0.75 %)	High (> 57 kg/ha)	High (> 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	16	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	High (> 0.75 %)	High (> 57 kg/ha)	High (> 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	18	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	High (> 0.75 %)	High (> 57 kg/ha)	High (> 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	19	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 – 57 kg/ha)	High (> 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	20	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 – 57 kg/ha)	High (> 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	21	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 – 57 kg/ha)	High (> 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	22	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	High (> 0.75 %)	High (> 57 kg/ha)	High (> 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	23	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 – 57 kg/ha)	High (> 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	24	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 – 57 kg/ha)	High (> 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	25	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 – 57 kg/ha)	Medium (145 – 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	26	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 – 57 kg/ha)	Medium (145 – 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	27	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	Medium (0.5 – 0.75 %)	Medium (23 – 57 kg/ha)	Medium (145 – 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	28	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	Medium (0.5 – 0.75 %)	Medium (23 – 57 kg/ha)	Medium (145 – 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	29	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	Medium (0.5 – 0.75 %)	Medium (23 – 57 kg/ha)	Medium (145 – 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	30	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	Medium (0.5 – 0.75 %)	Medium (23 – 57 kg/ha)	Medium (145 – 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	31	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	Medium (0.5 – 0.75 %)	Medium (23 – 57 kg/ha)	Medium (145 – 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	32	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	Medium (0.5 – 0.75 %)	Medium (23 – 57 kg/ha)	Medium (145 – 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	33	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	Medium (0.5 – 0.75 %)	Medium (23 – 57 kg/ha)	Medium (145 – 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	34	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	Medium (0.5 – 0.75 %)	Medium (23 – 57 kg/ha)	Medium (145 – 337 kg/ha)	Medium (10 – 20 ppm)	Medium (0.5 – 1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	35	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	Medium (0.5 – 0.75 %)	Medium (23 – 57 kg/ha)	Medium (145 – 337 kg/ha)	Medium (10 – 20 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	36	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 – 57 kg/ha)	Medium (145 – 337 kg/ha)	Medium (10 – 20 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	37	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 – 57 kg/ha)	Medium (145 – 337 kg/ha)	Medium (10 – 20 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	38	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 – 57 kg/ha)	Medium (145 – 337 kg/ha)	Medium (10 – 20 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)

Village	Survey Number	Soil Reaction	Salinity	Organic Carbon	Available Phosphorus	Available Potassium	Available Sulphur	Available Boron	Available Iron	Available Manganese	Available Copper	Available Zinc
Lakamapura	39	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	High (> 0.75 %)	Low (< 23 kg/ha)	Medium (145 – 337 kg/ha)	Medium (10 – 20 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	40	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	High (> 0.75 %)	Low (< 23 kg/ha)	Medium (145 – 337 kg/ha)	Medium (10 – 20 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	42	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	High (> 0.75 %)	Low (< 23 kg/ha)	Medium (145 – 337 kg/ha)	Medium (10 – 20 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	43	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 – 57 kg/ha)	Medium (145 – 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	44	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 – 57 kg/ha)	Medium (145 – 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	45	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 – 57 kg/ha)	Medium (145 – 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	46	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 – 57 kg/ha)	Medium (145 – 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	47	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 – 57 kg/ha)	Medium (145 – 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	48	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 – 57 kg/ha)	Medium (145 – 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	49	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 – 57 kg/ha)	Medium (145 – 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	50	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 – 57 kg/ha)	High (> 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	51	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 – 57 kg/ha)	High (> 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	52	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 – 57 kg/ha)	High (> 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	53	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	High (> 0.75 %)	High (> 57 kg/ha)	High (> 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	54	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	High (> 0.75 %)	High (> 57 kg/ha)	High (> 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	55	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 – 57 kg/ha)	High (> 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	56	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 – 57 kg/ha)	High (> 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	57	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	Medium (0.5 – 0.75 %)	Low (< 23 kg/ha)	High (> 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	58	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	Medium (0.5 – 0.75 %)	Medium (23 – 57 kg/ha)	High (> 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	59	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	Medium (0.5 – 0.75 %)	Medium (23 – 57 kg/ha)	High (> 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	60	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 – 57 kg/ha)	High (> 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	61	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	Medium (0.5 – 0.75 %)	Low (< 23 kg/ha)	High (> 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	62	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	High (> 0.75 %)	Low (< 23 kg/ha)	High (> 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	63	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 – 57 kg/ha)	High (> 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)

Village	Survey Number	Soil Reaction	Salinity	Organic Carbon	Available Phosphorus	Available Potassium	Available Sulphur	Available Boron	Available Iron	Available Manganese	Available Copper	Available Zinc
Lakamapura	64	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 – 57 kg/ha)	Medium (145 – 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	65	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 – 57 kg/ha)	Medium (145 – 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	66	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 – 57 kg/ha)	Medium (145 – 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	67	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	High (> 0.75 %)	Low (< 23 kg/ha)	Medium (145 – 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	78	Strongly alkaline (pH 8.4 – 9.0)	Non saline (<2 dsm)	Medium (0.5 – 0.75 %)	Low (< 23 kg/ha)	High (> 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	131	Strongly alkaline (pH 8.4 – 9.0)	Non saline (<2 dsm)	Low (< 0.5 %)	Low (< 23 kg/ha)	High (> 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	132	Strongly alkaline (pH 8.4 – 9.0)	Non saline (<2 dsm)	Low (< 0.5 %)	Low (< 23 kg/ha)	High (> 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	133	Strongly alkaline (pH 8.4 – 9.0)	Non saline (<2 dsm)	Low (< 0.5 %)	Low (< 23 kg/ha)	High (> 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	134	Strongly alkaline (pH 8.4 – 9.0)	Non saline (<2 dsm)	Low (< 0.5 %)	Low (< 23 kg/ha)	High (> 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	135	Strongly alkaline (pH 8.4 – 9.0)	Non saline (<2 dsm)	Medium (0.5 – 0.75 %)	Low (< 23 kg/ha)	High (> 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	136	Strongly alkaline (pH 8.4 – 9.0)	Non saline (<2 dsm)	Low (< 0.5 %)	Low (< 23 kg/ha)	High (> 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	137	Strongly alkaline (pH 8.4 – 9.0)	Non saline (<2 dsm)	Low (< 0.5 %)	Low (< 23 kg/ha)	High (> 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	139	Strongly alkaline (pH 8.4 – 9.0)	Non saline (<2 dsm)	Medium (0.5 – 0.75 %)	Low (< 23 kg/ha)	High (> 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	140	Strongly alkaline (pH 8.4 – 9.0)	Non saline (<2 dsm)	Medium (0.5 – 0.75 %)	Low (< 23 kg/ha)	High (> 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	141	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	Medium (0.5 – 0.75 %)	Low (< 23 kg/ha)	High (> 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	142	Strongly alkaline (pH 8.4 – 9.0)	Non saline (<2 dsm)	Medium (0.5 – 0.75 %)	Low (< 23 kg/ha)	High (> 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	143	Strongly alkaline (pH 8.4 – 9.0)	Non saline (<2 dsm)	Medium (0.5 – 0.75 %)	Low (< 23 kg/ha)	High (> 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	144	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	Medium (0.5 – 0.75 %)	Low (< 23 kg/ha)	High (> 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	145	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	Medium (0.5 – 0.75 %)	Low (< 23 kg/ha)	High (> 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	146	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	Medium (0.5 – 0.75 %)	Low (< 23 kg/ha)	High (> 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	147	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	Medium (0.5 – 0.75 %)	Low (< 23 kg/ha)	High (> 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	149	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	Medium (0.5 – 0.75 %)	Low (< 23 kg/ha)	High (> 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Lakamapura	150	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	Medium (0.5 – 0.75 %)	Low (< 23 kg/ha)	High (> 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Thalakalla	119	Railway	Railway	Railway	Railway	Railway	Railway	Railway	Railway	Railway	Railway	Railway

Village	Survey Number	Soil Reaction	Salinity	Organic Carbon	Available Phosphorus	Available Potassium	Available Sulphur	Available Boron	Available Iron	Available Manganese	Available Copper	Available Zinc
Thalakalla	120	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 – 57 kg/ha)	Medium (145 – 337 kg/ha)	Medium (10 – 20 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Thalakalla	121	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 – 57 kg/ha)	High (> 337 kg/ha)	Medium (10 – 20 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Thalakalla	122	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 – 57 kg/ha)	High (> 337 kg/ha)	Medium (10 – 20 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Thalakalla	123	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 – 57 kg/ha)	High (> 337 kg/ha)	Medium (10 – 20 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Thalakalla	124	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 – 57 kg/ha)	High (> 337 kg/ha)	Medium (10 – 20 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Thalakalla	125	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 – 57 kg/ha)	High (> 337 kg/ha)	Medium (10 – 20 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Thalakalla	126	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 – 57 kg/ha)	High (> 337 kg/ha)	Medium (10 – 20 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Thalakalla	127	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 – 57 kg/ha)	High (> 337 kg/ha)	Medium (10 – 20 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Thalakalla	128	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 – 57 kg/ha)	High (> 337 kg/ha)	Medium (10 – 20 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Thalakalla	129	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	Medium (0.5 – 0.75 %)	Medium (23 – 57 kg/ha)	High (> 337 kg/ha)	Medium (10 – 20 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Thalakalla	130	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	Medium (0.5 – 0.75 %)	Medium (23 – 57 kg/ha)	High (> 337 kg/ha)	Medium (10 – 20 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Thalakalla	131	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	Medium (0.5 – 0.75 %)	Medium (23 – 57 kg/ha)	Medium (145 – 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Thalakalla	142	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	Medium (0.5 – 0.75 %)	Medium (23 – 57 kg/ha)	Medium (145 – 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Thalakalla	143	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	Medium (0.5 – 0.75 %)	Medium (23 – 57 kg/ha)	Medium (145 – 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Thalakalla	144	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	Medium (0.5 – 0.75 %)	Medium (23 – 57 kg/ha)	High (> 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Thalakalla	145	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	Medium (0.5 – 0.75 %)	Medium (23 – 57 kg/ha)	High (> 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Thalakalla	146	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	Medium (0.5 – 0.75 %)	Medium (23 – 57 kg/ha)	Medium (145 – 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Thalakalla	147	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	Medium (0.5 – 0.75 %)	Medium (23 – 57 kg/ha)	Medium (145 – 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Thalakalla	148	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	Medium (0.5 – 0.75 %)	Medium (23 – 57 kg/ha)	High (> 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Thalakalla	149	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 – 57 kg/ha)	High (> 337 kg/ha)	Medium (10 – 20 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Thalakalla	150	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 – 57 kg/ha)	High (> 337 kg/ha)	Medium (10 – 20 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Thalakalla	151	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 – 57 kg/ha)	High (> 337 kg/ha)	Medium (10 – 20 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Thalakalla	152	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 – 57 kg/ha)	High (> 337 kg/ha)	Medium (10 – 20 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Thalakalla	153	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 – 57 kg/ha)	High (> 337 kg/ha)	Medium (10 – 20 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)

Village	Survey Number	Soil Reaction	Salinity	Organic Carbon	Available Phosphorus	Available Potassium	Available Sulphur	Available Boron	Available Iron	Available Manganese	Available Copper	Available Zinc
Thalakalla	154	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 – 57 kg/ha)	Medium (145 – 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Thalakalla	155	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	Medium (0.5 – 0.75 %)	Medium (23 – 57 kg/ha)	Medium (145 – 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Thalakalla	156	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	Medium (0.5 – 0.75 %)	Medium (23 – 57 kg/ha)	Medium (145 – 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Thalakalla	157	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	Medium (0.5 – 0.75 %)	Medium (23 – 57 kg/ha)	Medium (145 – 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Thalakalla	158	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	Medium (0.5 – 0.75 %)	Medium (23 – 57 kg/ha)	Medium (145 – 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Thalakalla	159	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	Medium (0.5 – 0.75 %)	Medium (23 – 57 kg/ha)	Medium (145 – 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Thalakalla	160	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 – 57 kg/ha)	Medium (145 – 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Thalakalla	161	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	Medium (0.5 – 0.75 %)	Medium (23 – 57 kg/ha)	Medium (145 – 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Thalakalla	162	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	Medium (0.5 – 0.75 %)	Medium (23 – 57 kg/ha)	Medium (145 – 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Thalakalla	163	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	Medium (0.5 – 0.75 %)	Medium (23 – 57 kg/ha)	Medium (145 – 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Thalakalla	164	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	Medium (0.5 – 0.75 %)	Medium (23 – 57 kg/ha)	Medium (145 – 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Thalakalla	165	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	Medium (0.5 – 0.75 %)	Medium (23 – 57 kg/ha)	Medium (145 – 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Thalakalla	166	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	Medium (0.5 – 0.75 %)	Medium (23 – 57 kg/ha)	Medium (145 – 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Thalakalla	167	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	Medium (0.5 – 0.75 %)	Medium (23 – 57 kg/ha)	Medium (145 – 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Thalakalla	168	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	Medium (0.5 – 0.75 %)	Medium (23 – 57 kg/ha)	High (> 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Thalakalla	169	Slightly alkaline (pH 7.3 – 7.8)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 – 57 kg/ha)	High (> 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Thalakalla	234	Strongly alkaline (pH 8.4 – 9.0)	Non saline (<2 dsm)	Low (< 0.5 %)	Low (< 23 kg/ha)	High (> 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Thalakalla	235	Strongly alkaline (pH 8.4 – 9.0)	Non saline (<2 dsm)	Low (< 0.5 %)	Low (< 23 kg/ha)	High (> 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Thalakalla	236	Strongly alkaline (pH 8.4 – 9.0)	Non saline (<2 dsm)	Low (< 0.5 %)	Low (< 23 kg/ha)	High (> 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)

Appendix III

Lakshmapura-2 (1U1c) Microwatershed

Soil Suitability Information

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Jasmine	Bhendi	Brinjal	Crossandra	Drumstick	Mulberry	Onion
Halageri	317	N1r	S3g	S3rg	S3g	S3rg	S3gt	N1r	S3rg	S3gt	S3rg	S3rg	S2rg	S3rg	S2rg	S3rg	S3rg	S3rg	S2rg	S3g	S3g	S3g	S3g	S3rg	S2rg	S3g	S3g	S3g	S3g	S3rg	S3rg	S3g
Halageri	318	S3r	S2g	S2rg	S2g	S2rg	S2rg	S3r	S2rg	S3t	S2rg	S2rg	S2g	S2rg	S2g	S2rg	S3r	S2rg	S2gt	S2g	S2g	S2g	S2g	S2rg	S2g	S2g	S2g	S2g	S2g	S2rg	S2rg	S2g
Halageri	321	S3r	S2g	S2rg	S2g	S2rg	S2rg	S3r	S2rg	S3t	S2rg	S2rg	S2g	S2rg	S2g	S2rg	S3r	S2rg	S2gt	S2g	S2g	S2g	S2g	S2rg	S2g	S2g	S2g	S2g	S2g	S2rg	S2rg	S2g
Halageri	322	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1tz	S3z	S2z	S3tz	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3t	S3t	S3z	S3tz	S3tz
Halageri	323	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1tz	S3z	S2z	S3tz	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3t	S3t	S3z	S3tz	S3tz
Halageri	324	S3r	S2g	S2rg	S2g	S2rg	S2rg	S3r	S2rg	S3t	S2rg	S2rg	S2g	S2rg	S2g	S2rg	S3r	S2rg	S2gt	S2g	S2g	S2g	S2g	S2rg	S2g	S2g	S2g	S2g	S2g	S2rg	S2rg	S2g
Halageri	325	S3r	S2g	S2rg	S2g	S2rg	S2rg	S3r	S2rg	S3t	S2rg	S2rg	S2g	S2rg	S2g	S2rg	S3r	S2rg	S2gt	S2g	S2g	S2g	S2g	S2rg	S2g	S2g	S2g	S2g	S2g	S2rg	S2rg	S2g
Halageri	326	N1r	S2rg	S3r	S2rg	S3r	S2rg	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	S3r	S3r	S3r	S2rt	S2rg	S2rg	S2rg	S2rg	S3r	S2r	S2rg	S2rg	S2rg	S2rg	S3r	S3r	S2rg
Halageri	329	N1r	S2rg	S3r	S2rg	S3r	S2rg	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	S3r	S3r	S3r	S2rt	S2rg	S2rg	S2rg	S2rg	S3r	S2r	S2rg	S2rg	S2rg	S2rg	S3r	S3r	S2rg
Halageri	330	S2rg	S2g	S2g	S2g	S2g	S2gt	S2rg	S2g	S3t	S2g	S2g	S2g	S2g	S2g	S2g	S2rg	S2g	S2gt	S2g	S2g	S2g	S2g	S2g	S2g	S2g	S2g	S2g	S2g	S2g	S2g	S2g
Halageri	331	S2rg	S2g	S2g	S2g	S2g	S2gt	S2rg	S2g	S3t	S2g	S2g	S2g	S2g	S2g	S2g	S2rg	S2g	S2gt	S2g	S2g	S2g	S2g	S2g	S2g	S2g	S2g	S2g	S2g	S2g	S2g	S2g
Halageri	332	S3r	S2g	S2rg	S2g	S2rg	S2rg	S3r	S2rg	S3t	S2rg	S2rg	S1	S2rg	S1	S2rg	S3r	S2rg	S2t	S2g	S2g	S2g	S2g	S2rg	S1	S2g	S2g	S2g	S2g	S2r	S2r	S2g
Halageri	333	N1r	S2rg	S3r	S2rg	S3r	S2rg	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	S3r	S3r	S3r	S2rt	S2rg	S2rg	S2rg	S2rg	S3r	S2r	S2rg	S2rg	S2rg	S2rg	S3r	S3r	S2rg
Halageri	349	S3r	S2g	S2rg	S2g	S2rg	S2rg	S3r	S2rg	S3t	S2rg	S2rg	S1	S2rg	S1	S2rg	S3r	S2rg	S2t	S2g	S2g	S2g	S2g	S2rg	S1	S2g	S2g	S2g	S2g	S2r	S2r	S2g
Halageri	350	N1r	S3rg	N1r	S3rg	N1r	S3rt	N1r	N1r	S3rt	N1r	N1r	S3rg	N1r	S3rg	N1r	N1r	N1r	S3rg	S3rg	S3rg	S3rg	S3rg	N1r	S3rg	S3rg	S3rg	S3rg	S3rg	N1r	N1r	S3rg
Halageri	351	Rail way	Rail way	Rail way	Rail way	Rail way	Rail way	Rail way	Rail way	Rail way	Rail way	Rail way	Rail way	Rail way	Rail way	Rail way	Rail way	Rail way	Rail way	Rail way	Rail way	Rail way	Rail way	Rail way	Rail way	Rail way	Rail way	Rail way	Rail way	Rail way	Rail way	Rail way
Lakama pura	8	S3rg	S3g	S2rg	S3g	S2rg	S3g	S3rg	S2rg	S3gt	S3g	S3g	S2g	S2rg	S2g	S2rg	S3rg	S2rg	S2gt	S3g	S3g	S3g	S3g	S2rg	S2g	S3g	S3g	S3g	S3g	S2rg	S2rg	S3g
Lakama pura	9	S1	S2t	S1	S1	S1	S2t	S1	S1	S3t	S1	S1	S1	S1	S1	S1	S1	S1	S3t	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S2t	S2t
Lakama pura	10	S1	S2t	S1	S1	S1	S2t	S1	S1	S3t	S1	S1	S1	S1	S1	S1	S1	S1	S3t	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S2t	S2t
Lakama pura	11	S1	S2t	S1	S1	S1	S2t	S1	S1	S3t	S1	S1	S1	S1	S1	S1	S1	S1	S3t	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S2t	S2t
Lakama pura	12	S1	S2t	S1	S1	S1	S2t	S1	S1	S3t	S1	S1	S1	S1	S1	S1	S1	S1	S3t	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S2t	S2t
Lakama pura	13	S3rt	S2tz	S3t	S2z	S3tz	S2rz	S3rz	S2rz	S2z	S2rz	S2rt	S3z	S3tz	S2z	N1tz	S3rz	S2rz	S3t	S2tz	S3t	S2tz	S2tz	S2rt	S2tz	S3t	S2tz	S3t	S3t	S3z	S3tz	S3tz

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Jasmine	Bhendi	Brinjal	Crossandra	Drumstick	Mulberry	Onion
Lakamapura	14	S3rt	S2tz	S3t	S2z	S3tz	S2rz	S3rz	S2rz	S2z	S2rz	S2rt	S3z	S3tz	S2z	N1tz	S3rz	S2rz	S3t	S2tz	S3t	S2tz	S2tz	S2rt	S2tz	S3t	S2tz	S3t	S3t	S3z	S3tz	S3tz
Lakamapura	15	S3rt	S2tz	S3t	S2z	S3tz	S2rz	S3rz	S2rz	S2z	S2rz	S2rt	S3z	S3tz	S2z	N1tz	S3rz	S2rz	S3t	S2tz	S3t	S2tz	S2tz	S2rt	S2tz	S3t	S2tz	S3t	S3t	S3z	S3tz	S3tz
Lakamapura	16	S1	S2t	S1	S1	S1	S2t	S1	S1	S3t	S1	S1	S1	S1	S1	S1	S1	S1	S3t	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S2t	S2t
Lakamapura	18	S3r	S2g	S2rg	S2g	S2rg	S2rg	S3r	S2rg	S3t	S2rg	S2rg	S2g	S2rg	S2g	S2rg	S3r	S2rg	S2gt	S2g	S2g	S2g	S2g	S2rg	S2g	S2g	S2g	S2g	S2g	S2rg	S2rg	S2g
Lakamapura	19	S3r	S2g	S2rg	S2g	S2rg	S2rg	S3r	S2rg	S3t	S2rg	S2rg	S2g	S2rg	S2g	S2rg	S3r	S2rg	S2gt	S2g	S2g	S2g	S2g	S2rg	S2g	S2g	S2g	S2g	S2g	S2rg	S2rg	S2g
Lakamapura	20	S3r	S2g	S2rg	S2g	S2rg	S2rg	S3r	S2rg	S3t	S2rg	S2rg	S2g	S2rg	S2g	S2rg	S3r	S2rg	S2gt	S2g	S2g	S2g	S2g	S2rg	S2g	S2g	S2g	S2g	S2g	S2rg	S2rg	S2g
Lakamapura	21	N1r	S3rg	N1r	S3rg	N1r	S3rg	N1r	N1r	S3rt	N1r	N1r	S3rg	N1r	S3rg	N1r	N1r	N1r	S3rg	S3rg	S3rg	S3rg	S3rg	N1r	S3rg	S3rg	S3rg	S3rg	S3rg	N1r	N1r	S3rg
Lakamapura	22	N1r	S3rg	N1r	S3rg	N1r	S3rg	N1r	N1r	S3rt	N1r	N1r	S3rg	N1r	S3rg	N1r	N1r	N1r	S3rg	S3rg	S3rg	S3rg	S3rg	N1r	S3rg	S3rg	S3rg	S3rg	S3rg	N1r	N1r	S3rg
Lakamapura	23	S3rt	S2tz	S3t	S2z	S3tz	S2rz	S3rz	S2rz	S2z	S2rz	S2rt	S3z	S3tz	S2z	N1tz	S3rz	S2rz	S3t	S2tz	S3t	S2tz	S2tz	S2rt	S2tz	S3t	S2tz	S3t	S3t	S3z	S3tz	S3tz
Lakamapura	24	S3rt	S2tz	S3t	S2z	S3tz	S2rz	S3rz	S2rz	S2z	S2rz	S2rt	S3z	S3tz	S2z	N1tz	S3rz	S2rz	S3t	S2tz	S3t	S2tz	S2tz	S2rt	S2tz	S3t	S2tz	S3t	S3t	S3z	S3tz	S3tz
Lakamapura	25	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1tz	S3z	S2z	S3tz	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3t	S3t	S3z	S3tz	S3tz
Lakamapura	26	S3rt	S2tz	S3t	S2z	S3tz	S2rz	S3rz	S2rz	S2z	S2rz	S2rt	S3z	S3tz	S2z	N1tz	S3rz	S2rz	S3t	S2tz	S3t	S2tz	S2tz	S2rt	S2tz	S3t	S2tz	S3t	S3t	S3z	S3tz	S3tz
Lakamapura	27	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1tz	S3z	S2z	S3tz	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3t	S3t	S3z	S3tz	S3tz
Lakamapura	28	S3r	S2g	S2rg	S2g	S2rg	S2rg	S3r	S2rg	S3t	S2rg	S2rg	S2g	S2rg	S2g	S2rg	S3r	S2rg	S2gt	S2g	S2g	S2g	S2g	S2rg	S2g	S2g	S2g	S2g	S2g	S2rg	S2rg	S2g
Lakamapura	29	N1r	S3r	N1r	S3r	N1r	S3r	N1r	N1r	S3r	N1r	N1r	S3r	N1r	S3r	N1r	N1r	N1r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	S3r	S3r	N1r	N1r	S3r
Lakamapura	30	S1	S2t	S1	S1	S1	S2t	S1	S1	S3t	S1	S1	S1	S1	S1	S1	S1	S1	S3t	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S2t
Lakamapura	31	N1r	S3r	N1r	S3r	N1r	S3r	N1r	N1r	S3r	N1r	N1r	S3r	N1r	S3r	N1r	N1r	N1r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	S3r	S3r	N1r	N1r	S3r
Lakamapura	32	S2r	S2g	S1	S2g	S1	S2g	S2r	S1	S3t	S2g	S2g	S1	S1	S1	S1	S2r	S1	S2t	S2g	S2g	S2g	S2g	S1	S1	S2g	S2g	S2g	S2g	S1	S1	S2g
Lakamapura	33	S2rg	S2g	S2g	S2g	S2g	S2gt	S2rg	S2g	S3t	S2g	S2g	S2g	S2g	S2g	S2g	S2rg	S2g	S2gt	S2g	S2g	S2g	S2g	S2g	S2g	S2g	S2g	S2g	S2g	S2g	S2g	S2g
Lakamapura	34	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1tz	S3z	S2z	S3tz	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3t	S3t	S3z	S3tz	S3tz
Lakamapura	35	N1r	S2r	S3r	S2r	S3r	S2r	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	S3r	S3r	S3r	S2rt	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S2r	S2r	S3r	S3r	S2r

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Jasmine	Bhendi	Brinjal	Crossandra	Drumstick	Mulberry	Onion
Lakamapura	36	S3r	S2g	S2rg	S2g	S2rg	S2rg	S3r	S2rg	S3t	S2rg	S2rg	S2g	S2rg	S2g	S2rg	S3r	S2rg	S2gt	S2g	S2g	S2g	S2g	S2rg	S2g	S2g	S2g	S2g	S2g	S2rg	S2rg	S2g
Lakamapura	37	S3r	S2g	S2rg	S2g	S2rg	S2rg	S3r	S2rg	S3t	S2rg	S2rg	S2g	S2rg	S2g	S2rg	S3r	S2rg	S2gt	S2g	S2g	S2g	S2g	S2rg	S2g	S2g	S2g	S2g	S2g	S2rg	S2rg	S2g
Lakamapura	38	S3r	S2g	S2rg	S2g	S2rg	S2rg	S3r	S2rg	S3t	S2rg	S2rg	S2g	S2rg	S2g	S2rg	S3r	S2rg	S2gt	S2g	S2g	S2g	S2g	S2rg	S2g	S2g	S2g	S2g	S2g	S2rg	S2rg	S2g
Lakamapura	39	S2rg	S3g	S2g	S3g	S2g	S3g	S2rg	S2g	S3gt	S3g	S3g	S2g	S2g	S2g	S2g	S2rg	S2g	S2gt	S3g	S3g	S3g	S3g	S2g	S2g	S3g	S3g	S3g	S3g	S2g	S2g	S3g
Lakamapura	40	S2rg	S3g	S2g	S3g	S2g	S3g	S2rg	S2g	S3gt	S3g	S3g	S2g	S2g	S2g	S2g	S2rg	S2g	S2gt	S3g	S3g	S3g	S3g	S2g	S2g	S3g	S3g	S3g	S3g	S2g	S2g	S3g
Lakamapura	42	N1r	S3g	S3rg	S3g	S3rg	S3g	N1r	S3rg	S3gt	S3rg	S3rg	S2rg	S3rg	S2rg	S3rg	S3rg	S3rg	S2rg	S3g	S3g	S3g	S3g	S3rg	S2rg	S3g	S3g	S3g	S3g	S3rg	S3rg	S3g
Lakamapura	43	S3r	S2g	S2rg	S2g	S2rg	S2rg	S3r	S2rg	S3t	S2rg	S2rg	S2g	S2rg	S2g	S2rg	S3r	S2rg	S2gt	S2g	S2g	S2g	S2g	S2rg	S2g	S2g	S2g	S2g	S2g	S2rg	S2rg	S2g
Lakamapura	44	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1tz	S3z	S2z	S3tz	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3t	S3t	S3z	S3tz	S3tz
Lakamapura	45	S3r	S2g	S2rg	S2g	S2rg	S2rg	S3r	S2rg	S3t	S2rg	S2rg	S2g	S2rg	S2g	S2rg	S3r	S2rg	S2gt	S2g	S2g	S2g	S2g	S2rg	S2g	S2g	S2g	S2g	S2g	S2rg	S2rg	S2g
Lakamapura	46	S3r	S2g	S2rg	S2g	S2rg	S2rg	S3r	S2rg	S3t	S2rg	S2rg	S2g	S2rg	S2g	S2rg	S3r	S2rg	S2gt	S2g	S2g	S2g	S2g	S2rg	S2g	S2g	S2g	S2g	S2g	S2rg	S2rg	S2g
Lakamapura	47	S3r	S2g	S2rg	S2g	S2rg	S2rg	S3r	S2rg	S3t	S2rg	S2rg	S2g	S2rg	S2g	S2rg	S3r	S2rg	S2gt	S2g	S2g	S2g	S2g	S2rg	S2g	S2g	S2g	S2g	S2g	S2rg	S2rg	S2g
Lakamapura	48	S3r	S2g	S2rg	S2g	S2rg	S2rg	S3r	S2rg	S3t	S2rg	S2rg	S2g	S2rg	S2g	S2rg	S3r	S2rg	S2gt	S2g	S2g	S2g	S2g	S2rg	S2g	S2g	S2g	S2g	S2g	S2rg	S2rg	S2g
Lakamapura	49	N1r	S2rg	S3r	S2rg	S3r	S2rg	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	S3r	S3r	S3r	S2rt	S2rg	S2rg	S2rg	S2rg	S3r	S2r	S2rg	S2rg	S2rg	S2rg	S3r	S3r	S2rg
Lakamapura	50	N1r	S2rg	S3r	S2rg	S3r	S2rg	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	S3r	S3r	S3r	S2rt	S2rg	S2rg	S2rg	S2rg	S3r	S2r	S2rg	S2rg	S2rg	S2rg	S3r	S3r	S2rg
Lakamapura	51	N1r	S3rg	N1r	S3rg	N1r	S3rt	N1r	N1r	S3rt	N1r	N1r	S3rg	N1r	S3rg	N1r	N1r	N1r	S3rg	S3rg	S3rg	S3rg	S3rg	N1r	S3rg	S3rg	S3rg	S3rg	S3rg	N1r	N1r	S3rg
Lakamapura	52	S3rt	S2tz	S3t	S2z	S3tz	S2rz	S3rz	S2rz	S2z	S2rz	S2rt	S3z	S3tz	S2z	N1tz	S3rz	S2rz	S3t	S2tz	S3t	S2tz	S2tz	S2rt	S2tz	S3t	S2tz	S3t	S3t	S3z	S3tz	S3tz
Lakamapura	53	S3rt	S2tz	S3t	S2z	S3tz	S2rz	S3rz	S2rz	S2z	S2rz	S2rt	S3z	S3tz	S2z	N1tz	S3rz	S2rz	S3t	S2tz	S3t	S2tz	S2tz	S2rt	S2tz	S3t	S2tz	S3t	S3t	S3z	S3tz	S3tz
Lakamapura	54	S3rt	S2tz	S3t	S2z	S3tz	S2rz	S3rz	S2rz	S2z	S2rz	S2rt	S3z	S3tz	S2z	N1tz	S3rz	S2rz	S3t	S2tz	S3t	S2tz	S2tz	S2rt	S2tz	S3t	S2tz	S3t	S3t	S3z	S3tz	S3tz
Lakamapura	55	S3rt	S2tz	S3t	S2z	S3tz	S2rz	S3rz	S2rz	S2z	S2rz	S2rt	S3z	S3tz	S2z	N1tz	S3rz	S2rz	S3t	S2tz	S3t	S2tz	S2tz	S2rt	S2tz	S3t	S2tz	S3t	S3t	S3z	S3tz	S3tz
Lakamapura	56	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1tz	S3z	S2z	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3t	S3t	S3z	S3tz	S3tz
Lakamapura	57	N1rn	S3n	N1n	S3n	N1n	S3n	N1rn	N1n	S3n	N1n	S3rn	N1n	N1n	N1n	N1tn	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n	N1n	N1n	N1n

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Jasmine	Bhendi	Brinjal	Crossandra	Drumstick	Mulberry	Onion
Lakamapura	58	N1rn	S3n	N1n	S3n	N1n	S3n	N1rn	N1n	S3n	N1n	S3rn	N1n	N1n	N1n	N1tn	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n	N1n	N1n	N1n
Lakamapura	59	N1rn	S3n	N1n	S3n	N1n	S3n	N1rn	N1n	S3n	N1n	S3rn	N1n	N1n	N1n	N1tn	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n	N1n	N1n	N1n
Lakamapura	60	N1r	S3rg	N1r	S3rg	N1r	S3rt	N1r	N1r	S3rt	N1r	N1r	S3rg	N1r	S3rg	N1r	N1r	N1r	S3rg	S3rg	S3rg	S3rg	S3rg	N1r	S3rg	S3rg	S3rg	S3rg	S3rg	N1r	N1r	S3rg
Lakamapura	61	N1n	S3n	N1n	S3n	N1n	S3n	N1n	N1n	S3n	N1n	S3n	N1n	N1n	N1n	N1tn	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n	N1n	N1n	N1n
Lakamapura	62	N1r	S3rg	N1r	S3rg	N1r	S3rt	N1r	N1r	S3rt	N1r	N1r	S3rg	N1r	S3rg	N1r	N1r	N1r	S3rg	S3rg	S3rg	S3rg	S3rg	N1r	S3rg	S3rg	S3rg	S3rg	S3rg	N1r	N1r	S3rg
Lakamapura	63	N1r	S2rg	S3r	S2rg	S3r	S2rg	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	S3r	S3r	S3r	S2rt	S2rg	S2rg	S2rg	S2rg	S3r	S2r	S2rg	S2rg	S2rg	S2rg	S3r	S3r	S2rg
Lakamapura	64	N1n	S3n	N1n	S3n	N1n	S3n	N1n	N1n	S3n	N1n	S3n	N1n	N1n	N1n	N1tn	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n	N1n	N1n	N1n
Lakamapura	65	N1n	S3n	N1n	S3n	N1n	S3n	N1n	N1n	S3n	N1n	S3n	N1n	N1n	N1n	N1tn	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n	N1n	N1n	N1n
Lakamapura	66	N1n	S3n	N1n	S3n	N1n	S3n	N1n	N1n	S3n	N1n	S3n	N1n	N1n	N1n	N1tn	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n	N1n	N1n	N1n
Lakamapura	67	N1r	S3g	S3rg	S3g	S3rg	S3g	N1r	S3rg	S3gt	S3rg	S3rg	S2rg	S3rg	S2rg	S3rg	S3rg	S3rg	S2rg	S3g	S3g	S3g	S3g	S3rg	S2rg	S3g	S3g	S3g	S3g	S3rg	S3rg	S3g
Lakamapura	78	N1n	S3n	N1n	S3n	N1n	S3n	N1n	N1n	S3n	N1n	S3n	N1n	N1n	N1n	N1tn	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n	N1n	N1n	N1n
Lakamapura	131	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1tz	S3z	S2z	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3t	S3t	S3z	S3tz	S3tz
Lakamapura	132	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1tz	S3z	S2z	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3t	S3t	S3z	S3tz	S3tz
Lakamapura	133	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1tz	S3z	S2z	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3t	S3t	S3z	S3tz	S3tz
Lakamapura	134	S3rt	S2tz	S3t	S2z	S3tz	S2rz	S3rz	S2rz	S2z	S2rz	S2rt	S3z	S3tz	S2z	N1tz	S3rz	S2rz	S3t	S2tz	S3t	S2tz	S2tz	S2rt	S2tz	S3t	S2tz	S3t	S3t	S3z	S3tz	S3tz
Lakamapura	135	S3rt	S2tz	S3t	S2z	S3tz	S2rz	S3rz	S2rz	S2z	S2rz	S2rt	S3z	S3tz	S2z	N1tz	S3rz	S2rz	S3t	S2tz	S3t	S2tz	S2tz	S2rt	S2tz	S3t	S2tz	S3t	S3t	S3z	S3tz	S3tz
Lakamapura	136	N1n	S3n	N1n	S3n	N1n	S3n	N1n	N1n	S3n	N1n	S3n	N1n	N1n	N1n	N1tn	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n	N1n	N1n	N1n
Lakamapura	137	N1n	S3n	N1n	S3n	N1n	S3n	N1n	N1n	S3n	N1n	S3n	N1n	N1n	N1n	N1tn	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n	N1n	N1n	N1n
Lakamapura	139	N1n	S3n	N1n	S3n	N1n	S3n	N1n	N1n	S3n	N1n	S3n	N1n	N1n	N1n	N1tn	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n	N1n	N1n	N1n
Lakamapura	140	N1n	S3n	N1n	S3n	N1n	S3n	N1n	N1n	S3n	N1n	S3n	N1n	N1n	N1n	N1tn	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n	N1n	N1n	N1n
Lakamapura	141	S3rt	S2tz	S3t	S2z	S3tz	S2rz	S3rz	S2rz	S2z	S2rz	S2rt	S3z	S3tz	S2z	N1tz	S3rz	S2rz	S3t	S2tz	S3t	S2tz	S2tz	S2rt	S2tz	S3t	S2tz	S3t	S3t	S3z	S3tz	S3tz

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Jasmine	Bhendi	Brinjal	Crossandra	Drumstick	Mulberry	Onion
Lakamapura	142	S3rt	S2tz	S3t	S2z	S3tz	S2rz	S3rz	S2rz	S2z	S2rz	S2rt	S3z	S3tz	S2z	N1tz	S3rz	S2rz	S3t	S2tz	S3t	S2tz	S2tz	S2rt	S2tz	S3t	S2tz	S3t	S3t	S3z	S3tz	S3tz
Lakamapura	143	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1tz	S3z	S2z	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3t	S3t	S3z	S3tz	S3tz
Lakamapura	144	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1tz	S3z	S2z	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3t	S3t	S3z	S3tz	S3tz
Lakamapura	145	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1tz	S3z	S2z	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3t	S3t	S3z	S3tz	S3tz
Lakamapura	146	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1tz	S3z	S2z	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3t	S3t	S3z	S3tz	S3tz
Lakamapura	147	S3rg	S3g	S2rg	S3g	S2rg	S3g	S3rg	S2rg	S3gt	S3g	S3g	S2g	S2rg	S2g	S2rg	S3rg	S2rg	S2gt	S3g	S3g	S3g	S3g	S2rg	S2g	S3g	S3g	S3g	S3g	S2rg	S2rg	S3g
Lakamapura	149	S3rg	S3g	S2rg	S3g	S2rg	S3g	S3rg	S2rg	S3gt	S3g	S3g	S2g	S2rg	S2g	S2rg	S3rg	S2rg	S2gt	S3g	S3g	S3g	S3g	S2rg	S2g	S3g	S3g	S3g	S3g	S2rg	S2rg	S3g
Lakamapura	150	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1tz	S3z	S2z	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3t	S3t	S3z	S3tz	S3tz
Thalakalla	119	Rail way	Rail way	Rail way	Rail way	Rail way	Rail way	Rail way	Rail way	Rail way	Rail way	Rail way	Rail way	Rail way	Rail way	Rail way	Rail way	Rail way	Rail way	Rail way	Rail way	Rail way	Rail way	Rail way	Rail way	Rail way	Rail way	Rail way	Rail way	Rail way	Rail way	Rail way
Thalakalla	120	N1r	S3rg	N1r	S3rg	N1r	S3rt	N1r	N1r	S3rt	N1r	N1r	S3rg	N1r	S3rg	N1r	N1r	N1r	S3rg	S3rg	S3rg	S3rg	S3rg	N1r	S3rg	S3rg	S3rg	S3rg	S3rg	N1r	N1r	S3rg
Thalakalla	121	N1r	S3rg	N1r	S3rg	N1r	S3rt	N1r	N1r	S3rt	N1r	N1r	S3rg	N1r	S3rg	N1r	N1r	N1r	S3rg	S3rg	S3rg	S3rg	S3rg	N1r	S3rg	S3rg	S3rg	S3rg	S3rg	N1r	N1r	S3rg
Thalakalla	122	N1r	S3rg	N1r	S3rg	N1r	S3rt	N1r	N1r	S3rt	N1r	N1r	S3rg	N1r	S3rg	N1r	N1r	N1r	S3rg	S3rg	S3rg	S3rg	S3rg	N1r	S3rg	S3rg	S3rg	S3rg	S3rg	N1r	N1r	S3rg
Thalakalla	123	N1r	S3rg	N1r	S3rg	N1r	S3rt	N1r	N1r	S3rt	N1r	N1r	S3rg	N1r	S3rg	N1r	N1r	N1r	S3rg	S3rg	S3rg	S3rg	S3rg	N1r	S3rg	S3rg	S3rg	S3rg	S3rg	N1r	N1r	S3rg
Thalakalla	124	S3rt	S2tz	S3t	S2z	S3tz	S2rz	S3rz	S2rz	S2z	S2rz	S2rt	S3z	S3tz	S2z	N1tz	S3rz	S2rz	S3t	S2tz	S3t	S2tz	S2tz	S2rt	S2tz	S3t	S2tz	S3t	S3t	S3z	S3tz	S3tz
Thalakalla	125	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1tz	S3z	S2z	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3t	S3t	S2tz	S3tz	S3tz
Thalakalla	126	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1tz	S3z	S2z	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3t	S3t	S2tz	S3tz	S3tz
Thalakalla	127	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1tz	S3z	S2z	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3t	S3t	S2tz	S3tz	S3tz
Thalakalla	128	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1tz	S3z	S2z	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3t	S3t	S2tz	S3tz	S3tz
Thalakalla	129	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1tz	S3z	S2z	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3t	S3t	S2tz	S3tz	S3tz
Thalakalla	130	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1tz	S3z	S2z	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3t	S3t	S2tz	S3tz	S3tz
Thalakalla	131	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1tz	S3z	S2z	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3t	S3t	S2tz	S3tz	S3tz

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Jasmine	Bhendi	Brinjal	Crossandra	Drumstick	Mulberry	Onion
Thalakalla	142	S3r	S2g	S2rg	S2g	S2rg	S2rg	S3r	S2rg	S3t	S2rg	S2rg	S2g	S2rg	S2g	S2rg	S3r	S2rg	S2gt	S2g	S2g	S2g	S2g	S2rg	S2g	S2g	S2g	S2g	S2g	S2rg	S2rg	S2g
Thalakalla	143	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1tz	S3z	S2z	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3t	S3t	S2tz	S3tz	S3tz
Thalakalla	144	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1tz	S3z	S2z	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3t	S3t	S2tz	S3tz	S3tz
Thalakalla	145	S3r	S2g	S2rg	S2g	S2rg	S2rg	S3r	S2rg	S3t	S2rg	S2rg	S2g	S2rg	S2g	S2rg	S3r	S2rg	S2gt	S2g	S2g	S2g	S2g	S2rg	S2g	S2g	S2g	S2g	S2g	S2rg	S2rg	S2g
Thalakalla	146	S3r	S2g	S2rg	S2g	S2rg	S2rg	S3r	S2rg	S3t	S2rg	S2rg	S2g	S2rg	S2g	S2rg	S3r	S2rg	S2gt	S2g	S2g	S2g	S2g	S2rg	S2g	S2g	S2g	S2g	S2g	S2rg	S2rg	S2g
Thalakalla	147	S3r	S2g	S2rg	S2g	S2rg	S2rg	S3r	S2rg	S3t	S2rg	S2rg	S2g	S2rg	S2g	S2rg	S3r	S2rg	S2gt	S2g	S2g	S2g	S2g	S2rg	S2g	S2g	S2g	S2g	S2g	S2rg	S2rg	S2g
Thalakalla	148	S3r	S2g	S2rg	S2g	S2rg	S2rg	S3r	S2rg	S3t	S2rg	S2rg	S2g	S2rg	S2g	S2rg	S3r	S2rg	S2gt	S2g	S2g	S2g	S2g	S2rg	S2g	S2g	S2g	S2g	S2g	S2rg	S2rg	S2g
Thalakalla	149	S3r	S2g	S2rg	S2g	S2rg	S2rg	S3r	S2rg	S3t	S2rg	S2rg	S2g	S2rg	S2g	S2rg	S3r	S2rg	S2gt	S2g	S2g	S2g	S2g	S2rg	S2g	S2g	S2g	S2g	S2g	S2rg	S2rg	S2g
Thalakalla	150	S3rt	S2tz	S3t	S2z	S3tz	S2rz	S3rz	S2rz	S2z	S2rz	S2rt	S3z	S3tz	S2z	N1tz	S3rz	S2rz	S3t	S2tz	S3t	S2tz	S2tz	S2rt	S2tz	S3t	S2tz	S3t	S3t	S3z	S3tz	S3tz
Thalakalla	151	S3rt	S2tz	S3t	S2z	S3tz	S2rz	S3rz	S2rz	S2z	S2rz	S2rt	S3z	S3tz	S2z	N1tz	S3rz	S2rz	S3t	S2tz	S3t	S2tz	S2tz	S2rt	S2tz	S3t	S2tz	S3t	S3t	S3z	S3tz	S3tz
Thalakalla	152	N1r	S3rg	N1r	S3rg	N1r	S3rt	N1r	N1r	S3rt	N1r	N1r	S3rg	N1r	S3rg	N1r	N1r	N1r	S3rg	S3rg	S3rg	S3rg	S3rg	N1r	S3rg	S3rg	S3rg	S3rg	S3rg	N1r	N1r	S3rg
Thalakalla	153	N1r	S3rg	N1r	S3rg	N1r	S3rt	N1r	N1r	S3rt	N1r	N1r	S3rg	N1r	S3rg	N1r	N1r	N1r	S3rg	S3rg	S3rg	S3rg	S3rg	N1r	S3rg	S3rg	S3rg	S3rg	S3rg	N1r	N1r	S3rg
Thalakalla	154	S3r	S2g	S2rg	S2g	S2rg	S2rg	S3r	S2rg	S3t	S2rg	S2rg	S2g	S2rg	S2g	S2rg	S3r	S2rg	S2gt	S2g	S2g	S2g	S2g	S2rg	S2g	S2g	S2g	S2g	S2g	S2rg	S2rg	S2g
Thalakalla	155	S2r	S2g	S1	S2g	S1	S2g	S2r	S1	S3t	S2g	S2g	S1	S1	S1	S1	S2r	S1	S2t	S2g	S2g	S2g	S2g	S1	S1	S2g	S2g	S2g	S2g	S1	S1	S2g
Thalakalla	156	S3r	S2g	S2rg	S2g	S2rg	S2rg	S3r	S2rg	S3t	S2rg	S2rg	S2g	S2rg	S2g	S2rg	S3r	S2rg	S2gt	S2g	S2g	S2g	S2g	S2rg	S2g	S2g	S2g	S2g	S2g	S2rg	S2rg	S2g
Thalakalla	157	S2r	S2g	S1	S2g	S1	S2g	S2r	S1	S3t	S2g	S2g	S1	S1	S1	S1	S2r	S1	S2t	S2g	S2g	S2g	S2g	S1	S1	S2g	S2g	S2g	S2g	S1	S1	S2g
Thalakalla	158	S2r	S2g	S1	S2g	S1	S2g	S2r	S1	S3t	S2g	S2g	S1	S1	S1	S1	S2r	S1	S2t	S2g	S2g	S2g	S2g	S1	S1	S2g	S2g	S2g	S2g	S1	S1	S2g
Thalakalla	159	S2r	S2g	S1	S2g	S1	S2g	S2r	S1	S3t	S2g	S2g	S1	S1	S1	S1	S2r	S1	S2t	S2g	S2g	S2g	S2g	S1	S1	S2g	S2g	S2g	S2g	S1	S1	S2g
Thalakalla	160	N1r	S3rg	N1r	S3rg	N1r	S3rt	N1r	N1r	S3rt	N1r	N1r	S3rg	N1r	S3rg	N1r	N1r	N1r	S3rg	S3rg	S3rg	S3rg	S3rg	N1r	S3rg	S3rg	S3rg	S3rg	S3rg	N1r	N1r	S3rg
Thalakalla	161	N1r	S3rg	N1r	S3rg	N1r	S3rt	N1r	N1r	S3rt	N1r	N1r	S3rg	N1r	S3rg	N1r	N1r	N1r	S3rg	S3rg	S3rg	S3rg	S3rg	N1r	S3rg	S3rg	S3rg	S3rg	S3rg	N1r	N1r	S3rg
Thalakalla	162	N1r	S3rg	N1r	S3rg	N1r	S3rt	N1r	N1r	S3rt	N1r	N1r	S3rg	N1r	S3rg	N1r	N1r	N1r	S3rg	S3rg	S3rg	S3rg	S3rg	N1r	S3rg	S3rg	S3rg	S3rg	S3rg	N1r	N1r	S3rg

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Jasmine	Bhendi	Brinjal	Crossandra	Drumstick	Mulberry	Onion
Thalakalla	163	S2rg	S2g	S2g	S2g	S2g	S2gt	S2rg	S2g	S3t	S2g	S2g	S2g	S2g	S2g	S2g	S2rg	S2g	S2gt	S2g	S2g	S2g	S2g	S2g	S2g	S2g	S2g	S2g	S2g	S2g	S2g	S2g
Thalakalla	164	S2r	S2g	S1	S2g	S1	S2g	S2r	S1	S3t	S2g	S2g	S1	S1	S1	S1	S2r	S1	S2t	S2g	S2g	S2g	S2g	S1	S1	S2g	S2g	S2g	S2g	S1	S1	S2g
Thalakalla	165	N1r	S3r	N1r	S3r	N1r	S3r	N1r	N1r	S3r	N1r	N1r	S3r	N1r	S3r	N1r	N1r	N1r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	S3r	S3r	N1r	N1r	S3r
Thalakalla	166	S2r	S2g	S1	S2g	S1	S2g	S2r	S1	S3t	S2g	S2g	S1	S1	S1	S1	S2r	S1	S2t	S2g	S2g	S2g	S2g	S1	S1	S2g	S2g	S2g	S2g	S1	S1	S2g
Thalakalla	167	S2r	S2g	S1	S2g	S1	S2g	S2r	S1	S3t	S2g	S2g	S1	S1	S1	S1	S2r	S1	S2t	S2g	S2g	S2g	S2g	S1	S1	S2g	S2g	S2g	S2g	S1	S1	S2g
Thalakalla	168	S3r	S2g	S2rg	S2g	S2rg	S2rg	S3r	S2rg	S3t	S2rg	S2rg	S2g	S2rg	S2g	S2rg	S3r	S2rg	S2gt	S2g	S2g	S2g	S2g	S2rg	S2g	S2g	S2g	S2g	S2g	S2rg	S2rg	S2g
Thalakalla	169	S3r	S2g	S2rg	S2g	S2rg	S2rg	S3r	S2rg	S3t	S2rg	S2rg	S2g	S2rg	S2g	S2rg	S3r	S2rg	S2gt	S2g	S2g	S2g	S2g	S2rg	S2g	S2g	S2g	S2g	S2g	S2rg	S2rg	S2g
Thalakalla	234	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1tz	S3z	S2z	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3t	S3t	S3z	S3tz	S3tz
Thalakalla	235	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1tz	S3z	S2z	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3t	S3t	S3z	S3tz	S3tz
Thalakalla	236	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1tz	S3z	S2z	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3t	S3t	S3z	S3tz	S3tz

PART-B

SOCIO-ECONOMIC STATUS OF FARM HOUSEHOLDS

CONTENTS

1	Findings of the socio-economic survey	1-2
2	Introduction	3
3	Methodology	5-6
4	Salient features of the survey	7-24
5	Summary	25-28

LIST OF TABLES

1	Households sampled for socio economic survey	7
2	Population characteristics	7
3	Age wise classification of household members	7
4	Education level of household members	8
5	Occupation of household heads	8
6	Occupation of family members	8
7	Institutional participation of household members	9
8	Type of house owned by households	9
9	Durable assets owned by households	9
10	Average value of durable assets owned by households	10
11	Farm implements owned by households	10
12	Average value of farm implements	10
13	Livestock possession by households	10
14	Average labour availability	11
15	Adequacy of hired labour	11
16	Distribution of land (ha)	11
17	Average land value (Rs./ha)	11
18	Status of bore wells	12
19	Source of irrigation	12
20	Depth of water(Avg in meters)	12
21	Irrigated area (ha)	12
22	Cropping pattern	12
23	Cropping intensity	13
24	Possession of bank account and saving	13
25	Borrowing status	13
26.a	Cost of cultivation of Maize	14
26.b	Cost of cultivation of Sunflower	15
26.c	Cost of cultivation of Bengal gram	16
26.d	Cost of cultivation of Groundnut	17
26.e	Cost of cultivation of Sorghum	18

27	Adequacy of fodder	19
28	Annual gross income	19
29	Average annual expenditure	19
30	Horticultural species grown	19
31	Forest species grown	20
32	Average additional investment capacity	20
33	Source of funds for additional investment	20
34	Marketing of the agricultural produce	21
35	Marketing channels used for sale of agricultural produce	21
36	Mode of transport of agricultural produce	21
37	Incidence of soil and water erosion problems	21
38	Interest shown towards soil testing	22
39	Usage pattern of fuel for domestic use	22
40	Source of drinking water	22
41	Source of light	22
42	Existence of sanitary toilet facility	22
43	Possession of public distribution system (PDS) card	23
44	Participation in NREGA programme	23
45	Adequacy of food items	23
46	Inadequacy of food items	23
47	Farming constraints experienced	24

Chapter 1

FINDINGS OF THE SOCIO-ECONOMIC SURVEY

- *The survey was conducted in Lakshmapura-2 is located at North latitude 15^0 22' 54.598" and 15^0 21' 33.894" and East longitude 76^0 3' 55.966" and 76^0 1' 49.0989" covering an area of about 493.28 ha coming under Lakshmapura, Vadagenahalu, Tadakal and Haligeri Villages of Koppal taluk.*
- *Socio-economic analysis of Lakshmapura-2 micro watersheds of Bhanapur sub-watershed, Koppala taluk & District indicated that, out of the total sample of 35 farmers were sampled in Lakshmapura-2 micro-watershed among households surveyed 18 (51.43%) were marginal, 6 (17.14%) were small, 5 (14.29 %) were semi medium and 1 (2.86 %) were medium farmers. 5 landless farmers were also interviewed for the survey.*
- *The population characteristics of households indicated that, there were 75 (60.48%) men and 49 (39.52 %) were women. The average population of landless was 2.8, marginal farmers were 3.8, small farmers were 3.7, semi medium farmers were 3.2 and medium farmers were 4.*
- *Majority of the respondents (37.10%) were in the age group of 16-35 years.*
- *Education level of the sample households indicated that, there were 29.84 per cent illiterates, 70.96 per cent pre university education and 4.03 per cent attained graduation.*
- *About, 82.86 per cent of household heads practicing agriculture and 14.29 per cent of the household heads were engaged as agricultural labourers.*
- *Agriculture was the major occupation for 64.52 per cent of the household members.*
- *In the study area, 97.14 per cent of the households possess katcha house.*
- *The durable assets owned by the households showed that, 100.00 per cent possess TV, 100.00 per cent possess mobile phones and 22.86 per cent possess motor cycles.*
- *Farm implements owned by the households indicated that, 2.86 per cent of the households possess plough, 8.57 per cent possess bullock cart.*
- *Regarding livestock possession by the households, 2.86 per cent possess local cow and 5.71 per cent possess buffalo.*
- *The average labour availability in the study area showed that, own labour men available in the micro watershed was 1.53, women available in the micro watershed was 1.17, hired labour (men) available was 10.63 and hired labour (women) available was 10.73.*
- *Out of the total land holding of the sample respondents 87.34 per cent (32.07 ha) of the area is under dry condition and the remaining 12.66 per cent area is irrigated land.*
- *There were 4.00 live bore wells and 3.00 dry bore wells among the sampled households.*
- *Bore well was the major source of irrigation for 11.43 per cent of the households.*

- *The major crops grown by sample farmers are Maize, Sunflower, Bengal gram, Groundnut and Sorghum and cropping intensity was recorded as 81.77 per cent.*
- *Out of the sample households 31.43 percent possessed bank account and 0.00 per cent of them have savings in the account.*
- *About 31.43 per cent of the respondents borrowed credit from various sources.*
- *The per hectare cost of cultivation for Maize, Sunflower, Bengal gram, Groundnut and Sorghum was Rs.53974.39 , 72039.01, 16550.91, 43647.47 and 25637.44 with benefit cost ratio of 1:1.10, 1: 2.60, 1: 4.50, 1: 2.90 and 1:1.70 respectively.*
- *Further, 17.14 per cent of the households opined that dry fodder was adequate.*
- *The average annual gross income of the farmers was Rs. 53057.14 in micro-watershed, of which Rs. 50914.29 comes from agriculture.*
- *Sampled households have grown 18 horticulture trees and 33 forestry trees together in the fields and back yards.*
- *Households have an average investment capacity of Rs. 3485.71 for land development.*
- *Source of funds for additional investment is concerned, 2.86 per cent depends on own funds and 48.57 per cent depends on bank loan for land development activities.*
- *Regarding marketing channels, 85.71 per cent of the households have sold agricultural produce to the local/village merchants, while, 5.71 per cent have sold in regulated markets.*
- *Further, 91.43 per cent of the households have used tractor for the transport of agriculture commodity. Majority of the farmers (82.86%) have experienced soil and water erosion problems in the watershed and 77.14 per cent of the households were interested towards soil testing.*
- *Fire was the major source of fuel for domestic use for 91.43 per cent of the households and 2.86 per cent households has LPG connection.*
- *Piped supply was the major source for drinking water for 68.57 per cent of the households.*
- *Electricity was the major source of light for 97.14 per cent of the households.*
- *In the study area, 42.86 per cent of the households possess toilet facility.*
- *Regarding possession of PDS card, 97.14 per cent of the households possessed BPL card.*
- *Households opined that, the requirement of cereals (97.14%), pulses (97.14%) and oilseeds (48.57%) are adequate for consumption.*
- *Farming constraints experienced by households in the micro watersheds were lower fertility status of the soil (88.57%) wild animal menace on farm field (80.00%), frequent incidence of pest and diseases (71.43%), inadequacy of irrigation water (8.57%), high cost of fertilizers and plant protection chemicals (74.29%), high rate of interest on credit (62.86%), low price for the agricultural commodities (57.14%), lack of marketing facilities in the area (57.14%), inadequate extension services (5.71%), lack of transport for safe transport of the agricultural produce to the market (37.14%), Less rainfall (2.86%).*

Chapter 2

INTRODUCTION

Soil and water are the two precious natural resources which are essential for crop production and existence of life on earth. Rainfed agriculture is under severe stress due to various constraints related to agriculture like uneven and erratic distribution of rainfall, indiscriminate use of fertilizers, chemicals and pesticides, adoption of improper land management practices, soil erosion, decline in soil fertility, decline in ground water resources leading to low crop productivity. The area under rainfed agriculture has to be managed effectively using the best available practices to enhance the production of food, fodder and fuel. This is possible if the land resources are characterized at each parcel of land through detailed land resource inventory using the best available techniques of remote sensing, GPS and GIS. The watershed development programs are aimed at the sustainable distribution of its resources and the process of creating and implementing plans, programs, and projects to sustain and enhance watershed functions that affect the plant, animal and human communities within a watershed boundary.

World Bank funded KWDP II, SUJALA III project was implemented in with Broad objective of demonstrating more effective watershed management through greater integration of programmes related to rain-fed agriculture, innovative and science based approaches and strengthen institutional capacities and If successful, it is expected that the systems and tools could be mainstreamed into the overall IWMP in the State of Karnataka and in time, throughout other IWMP operations in India. With this background the socio-economic survey has been carried out with following specific objectives:

1. To understand the demographic features of the households in the micro-watershed
2. To understand the extent of family labour available and additional employment opportunities available within the village.
3. To know the status of assets of households in the micro-watershed for suggesting possible improvements.
4. To study the cropping pattern, cropped area and productivity levels of different households in micro-watershed.
5. To determine the type and extent of livestock owned by different categories of HHs
6. Availability of fodder and level of livestock management.

Scope and importance of survey

Survey helps in identification of different socio-economic and resource use-patterns of farmers at the Micro watershed. Household survey provides demographic features, labour force, and levels of education; land ownership and asset position (including livestock and other household assets) of surveyed households; and cropping patterns, input intensities, and average crop yields from farmers' fields. It also discusses crop utilization and the degree of commercialization of production in the areas; farmers' access to and utilization of credit from formal and informal sources; and the level of adoption and use of soil, water,and pest management technologies.

Chapter 3

METHODOLOGY

The description of the methods, components selected for the survey and procedures followed in conducting the baseline survey are furnished under the following heads.

1. Description of the study area

Koppal district is an administrative district in the state of Karnataka in India. In the past Koppal was referred to as 'Kopana Nagara'. Koppal, now a district headquarters is ancient Kopana a major holy place of the Jainas. The district occupies an area of 7,190 km² and has a population of 1,196,089, which 16.58% were urban as of 2001. The Koppal district was formed after split of Raichur district.

Geographers are very particular about the physiography or relief of a region. It plays a very important role in the spatial analysis of agricultural situation of the study area. The undulating topography with black cotton soil shrips, cut across by numerous nalas or streams is the major characteristic feature of the study region. Three physiographic divisions have made considering the local conditions of landforms and crops grown in the district. On the basis of physiography, Koppal district can be divided into three major divisions. They are (a) Koppal & Yelburga plateau, (b) Maidan division, (c) Tungabhadra valley. The district is part of Krishna basin the main streams draining the area are Maskinala, Ilkal-nadi and Hirenala. These are Ephemaral in nature, these come under Tungabhadra sub-basin. The drainage exhibit dentritic to subdentric with drainage density varies from 1.4 to7.0kms/sq.km.

According to the 2011 census Koppal district has a population of 1,391,292, roughly equal to the nation of Swaziland or the US state of Hawaii. This gives it a ranking of 350th in India (out of a total of 640). The district has a population density of 250 inhabitants per square kilometre (650/sq mi). Its population growth rate over the decade 2001-2011 was 16.32%.Koppal has a sex ratio of 983 females for every 1000 males, and a literacy rate of 67.28%.

2. Locale of the survey and description of the micro-watershed and

The study was conducted in Lakshmapura-2 micro-watershed (Bhanapur sub-watershed, Koppala taluk & District) is located at North latitude 15^0 22’ 54.598” and 15^0 21’ 33.894” and East longitude 76^0 3’ 55.966" and 76^0 1’ 49.098”covering an area of about 493.28 ha bounded by under Lakshmapura, Vadagenahalu, Tadakal and Haligeri Villages.

3. Selection of the respondents for the study

The micro-watershed is marked with 320 square meters grids. One farmer from every alternate grid in the micro-watershed was selected for the study and interviewed for socio-economic data. Totally 35 households were interviewed for the survey.

4. The parameters considered for socio-economic survey of households

Two forms of data were collected from the micro-watershed which includes primary data from the farm households and secondary data about the villages under the micro-watershed jurisdiction.

The following parameters were considered for the primary data collection about the socio-economic data of the households, (1) Demographic information, (2) Farm and durable assets owned by households, (3) Livestock possession, (4) Labour availability, (5) Level of migration in the village, Land holding, (7) Cropping pattern, (8) Source of irrigation, (9) Borrowing status, (10) Cost of cultivation of major crops, (11) Economics of subsidiary activities, (12) Fodder availability, (13) Family annual income from different sources, (14) Horticulture and forestry species grown, (15) Additional investment capacity, (16) Marketing practices, (17) Status of soil and water conservation structure, (18) Access to basic needs and (19) Constraints and suggestion.

The following parameters were considered for the secondary data regarding the villages under the micro-watershed jurisdiction, (1) Number of villages in each micro-watershed jurisdiction, (2) Village wise number of households, (3) Geographical area of the villages, (4) Cultivable are a including rainfed and irrigated, (5) Number and type of house in each village, (6) Human and livestock population, (7) Facilities in the village such as roads, transport facility for conveyance, drinking water supply, street light and (8) Community based organizations in the villages.

5. Development of interview schedule and data collection

Taking into the consideration the objectives of the survey, an interview schedule was prepared after thorough consultation with the experts in the field of social sciences. A comprehensive interview schedule covering all the major parameters for measuring the socio-economic situation was developed.

6. Tools used to analyze the data

The statistical components such as frequency and percentage were used to analyze the data.

Abbreviations used in the report

LL=Landless

MF=Marginal Farmers

SF=Small farmers

SMF=Semi medium farmers

MDF=Medium farmers

LF=Large Farmers

Chapter 4

FINDINGS OF THE SURVEY

This chapter deals with systematic presentation of results of the survey. Keeping in view the objectives, the salient features of the survey are presented under the following headings.

Households sampled for socio-economic survey: The data on households sampled for socio economic survey in Lakshmapura-2 Micro watershed is presented in Table 1 and it indicated that 35 farmers were sampled in Lakshmapura-2 micro-watershed among households surveyed 18 (51.43%) were marginal, 6 (17.14%) were small, 5 (14.29 %) were semi medium and 1 (2.86 %) were medium farmers. 5 landless farmers were also interviewed for the survey.

Table 1. Households sampled for socio economic survey in Lakshmapura-2 micro-watershed

Sl.No.	Particulars	LL (5)		MF (18)		SF (6)		SMF (5)		MDF (1)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Farmers	5	14.3	18	51.4	6	17.1	5	14.3	1	2.86	35	100

Population characteristics: The population characteristics of households sampled for socio-economic survey in Lakshmapura-2 Micro watershed is presented in Table 2. The data indicated that, there were 75 (60.48%) men and 49 (39.52%) were women. The average population of landless was 2.8, marginal farmers were 3.8, small farmers were 3.7, semi medium farmers were 3.2 and medium farmers were 4.

Table 2. Population characteristics in Lakshmapura-2 micro-watershed

Sl.No.	Particulars	LL (14)		MF (68)		SF (22)		SMF (16)		MDF (4)		All (124)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Men	9	64.3	39	57	13	59	12	75	2	50	75	60.5
2	Women	5	35.7	29	43	9	41	4	25	2	50	49	39.5
Total		14	100	68	100	22	100	16	100	4	100	124	100
Average		2.8		3.8		3.7		3.2		4.0		3.5	

Age wise classification of population: The age wise classification of household members in Lakshmapura-2 Micro watershed is presented in Table 3. The indicated that, 26 (20.97%) of population were 0-15 years of age, 46 (37.10%) were 16-35 years of age, 31(25.00%) were 36-60 years of age and 21 (16.94 %) were above 61 years of age.

Table 3: Age wise classification of members of the household in Lakshmapura-2 micro-watershed

Sl.No.	Particulars	LL (14)		MF (68)		SF (22)		SMF (16)		MDF (4)		All (124)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	0-15 years of age	0	0	21	30.9	3	13.6	0	0	2	50	26	20.97
2	16-35 years of age	6	42.9	21	30.9	11	50	7	43.75	1	25	46	37.1
3	36-60 years of age	7	50	13	19.1	6	27.3	4	25	1	25	31	25
4	> 61 years	1	7.14	13	19.1	2	9.09	5	31.25	0	0	21	16.94
Total		14	100	68	100	22	100	16	100	4	100	124	100

Education level of household members: Education level of household members in Lakshmapura-2 Micro watershed is presented in Table 4. The results indicated that, there were 29.84 per cent of illiterates, 22.58 per cent of them had primary school education, 8.06 per cent middle school education, 22.58 per cent high school education, 8.87 per cent of them had PUC education, 4.03 per cent attained graduation and 3.23 them had other education.

Table 4. Education level of members of the household in Lakshmapura-2 micro-watershed

Sl.No.	Particulars	LL (14)		MF (68)		SF (22)		SMF (16)		MDF (4)		All (124)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Illiterate	4	28.6	19	27.9	4	18.2	10	62.5	0	0	37	29.8
2	Primary School	3	21.4	16	23.5	6	27.3	1	6.25	2	50	28	22.6
3	Middle School	0	0	9	13.2	1	4.55	0	0	0	0	10	8.06
4	High School	4	28.6	15	22.1	4	18.2	3	18.8	2	50	28	22.6
5	PUC	1	7.14	4	5.88	4	18.2	2	12.5	0	0	11	8.87
6	Degree	2	14.3	0	0	3	13.6	0	0	0	0	5	4.03
7	Masters	0	0	1	1.47	0	0	0	0	0	0	1	0.81
8	Others	0	0	4	5.88	0	0	0	0	0	0	4	3.23
Total		14	100	68	100	22	100	16	100	4	100	124	100

Occupation of head of households: The data regarding the occupation of the household heads in Lakshmapura-2 Micro watershed is presented in Table 5. The results indicate that, 82.86 per cent of households heads were practicing agriculture and 14.29 per cent of the household heads were agricultural Labour.

Table 5: Occupation of heads of households in Lakshmapura-2 micro-watershed

Sl.No.	Particulars	LL (5)		MF (18)		SF (6)		SMF (5)		MDF (1)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Agriculture	1	20	18	100	5	83.33	4	80	1	100	29	82.86
2	Agricultural Labour	3	60	0	0	1	16.67	1	20	0	0	5	14.29
3	General Labour	1	20	0	0	0	0	0	0	0	0	1	2.86
Total		5	100	18	100	6	100	5	100	1	100	35	100

Table 6: Occupation of members of the household in Lakshmapura-2 micro-watershed

Sl.No.	Particulars	LL (14)		MF (68)		SF (22)		SMF (16)		MDF (4)		All (124)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Agriculture	5	35.7	44	64.7	14	63.64	15	93.75	2	50	80	64.5
2	Agricultural Labour	7	50	0	0	2	9.09	1	6.25	0	0	10	8.06
3	General Labour	1	7.14	1	1.47	0	0	0	0	0	0	2	1.61
4	Private Service	0	0	0	0	1	4.55	0	0	0	0	1	0.81
5	Student	1	7.14	19	27.9	5	22.73	0	0	2	50	27	21.8
6	Children	0	0	4	5.88	0	0	0	0	0	0	4	3.23
Total		14	100	68	100	22	100	16	100	4	100	124	100

Occupation of the members of the household: The data regarding the occupation of the household members in Lakshmapura-2 Micro watershed is presented in Table 6. The results indicate that, agriculture was the major occupation for 64.52 per cent of the household members, 8.06 per cent were agricultural labour, 1.61 per cent were general labour, 21.77 per cent were working in pursuing education and 3.23 per cent were children.

Institutional Participation of household members: The data regarding the institutional participation of the household members in Lakshmapura-2 Micro watershed is presented in Table 7. The results show that, out of the total family members in the households 100 per cent of them were participating in taluk panchayat and rest were not participating in any of the institutions.

Table 7: Institutional Participation of household member in Lakshmapura-2 micro-watershed

Sl.No.	Particulars	LL (14)		MF (68)		SF (22)		SMF (16)		MDF (4)		All (124)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	No Participation	14	100	68	100	22	100	16	100	4	100	124	100
Total		14	100	68	100	22	100	16	100	4	100	124	100

Type of house owned: The data regarding the type of house owned by the households in Lakshmapura-2 Micro watershed is presented in Table 8. The results indicate that, 2.86 percent possess thatched house and 97.14 per cent of the households possess katcha house.

Table 8. Type of house owned by households in Lakshmapura-2 micro-watershed

Sl.No.	Particulars	LL (5)		MF (18)		SF (6)		SMF (5)		MDF (1)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Thatched	0	0	1	5.6	0	0	0	0	0	0	1	2.86
2	Katcha	5	100	17	94	6	100	5	100	1	100	34	97.14
Total		5	100	18	100	6	100	5	100	1	100	35	100

Durable assets owned by the households: The data regarding the Durable Assets owned by the households in Lakshmapura-2 Micro watershed is presented in Table 9. The result shows that, 100.00 per cent possess TV, 22.86 per cent possess motor cycle and 100.00 per cent possess mobile phones.

Table 9. Durable assets owned by households in Lakshmapura-2 micro-watershed

Sl.No.	Particulars	LL (5)		MF (18)		SF (6)		SMF (5)		MDF (1)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Television	5	100	18	100	6	100	5	100	1	100	35	100
2	Motor Cycle	0	0	4	22	3	50	0	0	1	100	8	22.86
3	Mobile Phone	5	100	18	100	6	100	5	100	1	100	35	100

Average value of durable assets: The data regarding the average value of durable assets owned by the households in Lakshmapura-2 Micro watershed is presented in Table 10.

The result shows that, the average value of television was Rs.5457.00, motor cycle was Rs. 38125.00 and mobile phone was Rs.2985.00.

Table 10. Average value of durable assets owned in Lakshmapura-2 micro-watershed **Average Value (Rs.)**

Sl.No.	Particulars	LL (5)	MF (18)	SF (6)	SMF (5)	MDF (1)	All (35)
1	Television	4200	5500	6000	5600	7000	5457
2	Motor Cycle	0	39500	38333	0	32000	38125
3	Mobile Phone	1300	3277	4416	1300	6000	2985

Farm implements owned: The data regarding the farm implements owned by the households in Lakshmapura-2 Micro watershed is presented in Table 11. About 8.57 per cent of the households possess Bullock Cart, 2.86 per cent possess plough and 20.00 per cent possess Weeder.

Table 11. Farm implements owned in Lakshmapura-2 micro-watershed

Sl.No.	Particulars	LL (5)		MF (18)		SF (6)		SMF (5)		MDF (1)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Bullock Cart	0	0	1	5.56	0	0	2	40	0	0	3	8.57
2	Plough	0	0	0	0	0	0	1	20	0	0	1	2.86
3	Weeder	0	0	4	22.2	0	0	3	60	0	0	7	20
4	Blank	5	100	14	77.8	6	100	1	20	1	100	27	77.14

Average value of farm implements: The data regarding the average value of farm Implements owned by the households in Lakshmapura-2 Micro watershed is presented in Table 12. The results show that the average value of plough was Rs.1000.00, bullock Cart was Rs.20000.00 and weeder was Rs.85.00.

Table 12. Average value of farm implements in Lakshmapura-2 micro-watershed

Average Value (Rs.)

Sl.No.	Particulars	LL (5)	MF (18)	SF (6)	SMF (5)	MDF (1)	All (35)
1	Bullock Cart	0	20000	0	20000	0	20000
2	Plough	0	0	0	1000	0	1000
3	Weeder	0	90	0	78	0	85

Livestock possession by the households: The data regarding the Livestock possession by the households in Lakshmapura-2 Micro watershed is presented in Table 13. The results indicate that, 14.29 per cent of the households possess bullocks, 2.86 per cent possess local cow and 5.71 per cent possess buffalo.

Table 13. Livestock possession by households in Lakshmapura-2 micro-watershed

Sl.No.	Particulars	LL (5)		MF (18)		SF (6)		SMF (5)		MDF (1)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Bullock	0	0	3	17	0	0	2	40	0	0	5	14.29
2	Local cow	0	0	0	0	1	16.67	0	0	0	0	1	2.86
3	Buffalo	0	0	1	5.6	0	0	1	20	0	0	2	5.71
4	blank	5	100	15	83	5	83.33	3	60	1	100	29	82.86

Average Labour availability: The data regarding the average labour availability in Lakshmapura-2 Micro watershed is presented in Table 14. The indicated that, own labour men available in the micro watershed was 1.53, women available in the micro watershed was 1.17, hired labour (men) available was 10.63 and hired labour (women) available was 10.73.

Table 14. Average labour availability in Lakshmapura-2 micro-watershed

Sl.No.	Particulars	LL (5)	MF (18)	SF (6)	SMF (5)	MDF (1)	All (35)
		N	N	N	N	N	N
1	Hired labour Female	0	7.89	9.17	23	10	10.73
2	Own Labour Female	0	1.06	1	1.8	1	1.17
3	Own labour Male	0	1.39	1.5	2.2	1	1.53
4	Hired labour Male	0	7.72	9.17	23	10	10.63

Adequacy of hired labour: The data regarding the adequacy of hired labour in Lakshmapura-2 Micro watershed is presented in Table 15. The results indicate that, 85.71 per cent of the household opined that hired labour was adequate.

Table 15. Adequacy of hired labour in Lakshmapura-2 micro-watershed

Sl.No.	Particulars	LL (5)		MF (18)		SF (6)		SMF (5)		MDF (1)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Adequate	0	0	18	100	6	100	5	100	1	100	30	85.7

Distribution of land (ha): The data regarding the distribution of land (ha) in Lakshmapura-2 Micro watershed is presented in Table 16. The results indicate that, 28.01 ha (87.34%) of dry land and 4.06 ha (12.66 %) of irrigated land.

Table 16. Distribution of land (ha) in Lakshmapura-2 micro-watershed

Sl.No.	Particulars	LL (5)		MF (18)		SF (6)		SMF (5)		MDF (1)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Dry	0	0	8.24	86.53	6.62	87.39	8.3	82	4.86	100	28.01	87.34
2	Irrigated	0	0	1.28	13.47	0.96	12.61	1.82	18	0	0	4.06	12.66
Total		0	100	9.52	100	7.57	100	10.12	100	4.86	100	32.07	100

Average value of land (ha): The data regarding the average land value (Rs./ha) in Lakshmapura-2 Micro watershed is presented in Table 17. The results show that the average value of dry land was Rs.670943.51 and the average value of irrigated land was Rs.566400.80.

Table 17. Average value of land (ha) in Lakshmapura-2 micro-watershed

Sl.No.	Particulars	LL (5)	MF (18)	SF (6)	SMF (5)	MDF (1)	All (35)
		N	N	N	N	N	N
1	Dry	0	1504322	317247.7	481951.2	61750	670943.5
2	Irrigated	0	1168770	418644.1	219555.6	0	566400.8

Status of bore wells: The data regarding the status of bore wells in Lakshmapura-2 Micro watershed is presented in Table 18. The results indicate that, there were 3 De-

functioning bore wells and 4 functioning bore wells among the sampled households in micro watershed.

Table 18. Status of bore wells in Lakshmapura-2 micro-watershed

Sl.No.	Particulars	LL (5)	MF (18)	SF (6)	SMF (5)	MDF (1)	All (35)
		N	N	N	N	N	N
1	De-functioning	0	0	2	1	0	3
2	Functioning	0	2	1	1	0	4

Source of irrigation: The data regarding the source of irrigation in Lakshmapura-2 Micro watershed is presented in Table 19. The results show that, bore well for 11.43 per cent of the households.

Table 19. Source of irrigation in Lakshmapura-2 micro-watershed

Sl.No.	Particulars	LL (5)		MF (18)		SF (6)		SMF (5)		MDF (1)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Bore Well	0	0	2	11.1	1	16.67	1	20	0	0	4	11.43

Depth of water (Avg. In meters): The data regarding the depth of water in Lakshmapura-2 Micro watershed is presented in Table 20. The results revealed that, the depth of bore well was 9.14 meter.

Table 20. Depth of water (Avg. In meters) in Lakshmapura-2 micro-watershed

Sl.No.	Particulars	LL (5)	MF (18)	SF (6)	SMF (5)	MDF (1)	All (35)
		N	N	N	N	N	N
1	Bore Well	0	7.62	20.32	12.19	0	9.14

Irrigated Area (ha): The data regarding the irrigated area (ha) in Lakshmapura-2 Micro watershed is presented in Table 21. The results indicate that, the availability of irrigation water was used for kharif crops was 3.62 ha.

Table 21. Irrigated Area (ha) in Lakshmapura-2 micro-watershed

Sl.No.	Particulars	LL (5)	MF (18)	SF (6)	SMF (5)	MDF (1)	All (35)
1	Kharif	0	0.84	0.96	1.82	0	3.62
Total		0	0.84	0.96	1.82	0	3.62

Table 22. Cropping pattern in Lakshmapura-2 micro-watershed

Sl.No.	Particulars	LL (5)	MF (18)	SF (6)	SMF (5)	MDF (1)	All (35)
1	Kharif - Maize	0	2.62	2.49	4.92	0	10.03
2	Kharif - Groundnut	0	3.52	2.96	2.02	0	8.5
3	Kharif - Sorghum	0	2.61	0.85	2.02	2.43	7.91
4	Kharif - Bengal gram	0	0	0	0	2.43	2.43
5	Kharif - Sunflower	0	0.51	0.96	0.81	0	2.27
Total		0	9.26	7.25	9.78	4.86	31.14

Cropping pattern: The data regarding the cropping pattern in Lakshmapura-2 Micro watershed is presented in Table 22. The results indicate that, farmers have grown Maize

(10.03 ha), Groundnut (8.50 ha), Sorghum (7.91 ha), Bengal gram (2.43 ha) and Sunflower (2.27 ha).

Cropping intensity: The data regarding the cropping intensity in Lakshmapura-2 Micro watershed is presented in Table 23. The results indicate that, the cropping intensity was 81.77 per cent.

Table 23. Cropping intensity (%) in Lakshmapura-2 micro-watershed

Sl.No.	Particulars	LL (5)	MF (18)	SF (6)	SMF (5)	MDF (1)	All (35)
1	Cropping Intensity	0	98.7	100	83.28	50	81.77

Possession of bank account and savings: The data regarding the possession of bank account and saving in Lakshmapura-2 micro-watershed is presented in Table 24. The results indicate that, 31.43 cent of the households posses bank account.

Table 24. Possession of Bank account and savings in Lakshmapura-2 micro-watershed

Sl.No.	Particulars	LL (5)		MF (18)		SF (6)		SMF (5)		MDF (1)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Account	0	0	5	27.78	3	50	2	40	1	100	11	31.43

Borrowing status: The data regarding the borrowing status in Lakshmapura-2 micro-watershed is presented in Table 25. The results indicate that, 31.43 percent of the sample farmers have borrowed credit from different sources.

Table 25. Borrowing status in Lakshmapura-2 micro-watershed

Sl.No.	Particulars	LL (5)		MF (18)		SF (6)		SMF (5)		MDF (1)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Credit Availed	0	0	5	27.78	3	50	2	40	1	100	11	31.43

Cost of Cultivation of Maize: The data regarding the cost of cultivation (Rs/ha) of Maize in Lakshmapura-2 micro watershed is presented in Table 26.a. The results indicate that, the total cost of cultivation (Rs/ha) for Maize was Rs. 53974.39. The gross income realized by the farmers was Rs. 62151.66. The net income from Maize cultivation was Rs.8177.27, thus the benefit cost ratio was found to be 1:1.10.

Table 26(a). Cost of Cultivation of Maize in Lakshmapura-2 micro-watershed

Sl.No	Particulars		Units	Phy Units	Value(Rs.)	% to C3
I	**Cost A1**					
1	Hired Human Labour		Man days	88.96	15096.41	27.97
2	Bullock		Pairs/day	3.77	1882.63	3.49
3	Tractor		Hours	6.09	6085.1	11.27
4	Seed Main Crop (Establishment and Maintenance)		Kgs (Rs.)	26.74	3208.91	5.95
5	FYM		Quintal	2.06	2470	4.58
6	Fertilizer + micronutrients		Quintal	5.87	4671.74	8.66
7	Pesticides (PPC)		Kgs / liters	3.45	4142.63	7.68
8	Irrigation		Number	11.49	0	0
9	Depreciation charges			0	101.12	0.19
10	Land revenue and Taxes			0	4.94	0.01
II	**Cost B1**					
11	Interest on working capital				1739.19	3.22
12	**Cost B1 = (Cost A1 + sum of 15 and 16)**				39402.68	73
III	**Cost B2**					
13	Rental Value of Land				388.89	0.72
14	**Cost B2 = (Cost B1 + Rental value)**				39791.57	73.72
IV	**Cost C1**					
15	Family Human Labour			44.67	9276.06	17.19
16	**Cost C1 = (Cost B2 + Family Labour)**				49067.63	90.91
V	**Cost C2**					
17	Risk Premium				0	0
18	**Cost C2 = (Cost C1 + Risk Premium)**				49067.63	90.91
VI	**Cost C3**					
19	Managerial Cost				4906.76	9.09
20	**Cost C3 = (Cost C2 + Managerial Cost)**				53974.39	100
VII	**Economics of the Crop**					
a.	Main Product	a) Main Product (q)		51.79	62151.66	
		b) Main Crop Sales Price (Rs.)			1200	
b.	Gross Income (Rs.)				62151.66	
c.	Net Income (Rs.)				8177.27	
d.	Cost per Quintal (Rs./q.)				1042.12	
e.	Benefit Cost Ratio (BC Ratio)				1:1.1	

Cost of Cultivation of Sunflower: The data regarding the cost of cultivation (Rs/ha) of Sunflower in Lakshmapura-2 micro watershed is presented in Table 26.b. The results indicate that, the total cost of cultivation (Rs/ha) for Sunflower was Rs. 72039.01. The gross income realized by the farmers was Rs. 189787.06. The net income from Sunflower cultivation was Rs.117748.05, thus the benefit cost ratio was found to be 1:2.60.

Table 26(b). Cost of Cultivation of Sunflower in Lakshmapura-2 micro-watershed

Sl.No	Particulars		Units	Phy Units	Value(Rs.)	% to C3
I	**Cost A1**					
1	Hired Human Labour		Man days	109.33	18008.24	25
2	Bullock		Pairs/day	3.35	1674.58	2.32
3	Tractor		Hours	6.87	6865.76	9.53
4	Machinery		Hours	1.31	1308.26	1.82
5	Seed Main Crop (Establishment and Maintenance)		Kgs (Rs.)	10.41	4683.84	6.5
6	FYM		Quintal	6.18	7410	10.29
7	Fertilizer + micronutrients		Quintal	4.8	2969.23	4.12
8	Pesticides (PPC)		Kgs / liters	4.53	5434	7.54
9	Depreciation charges			0	1.91	0
10	Land revenue and Taxes			0	4.94	0.01
II	**Cost B1**					
11	Interest on working capital				2459.65	3.41
12	**Cost B1 = (Cost A1 + sum of 15 and 16)**				50820.42	70.55
III	**Cost B2**					
13	Rental Value of Land				708.33	0.98
14	**Cost B2 = (Cost B1 + Rental value)**				51528.75	71.53
IV	**Cost C1**					
15	Family Human Labour			64.27	13961.26	19.38
16	**Cost C1 = (Cost B2 + Family Labour)**				65490.01	90.91
V	**Cost C2**					
17	Risk Premium				0	0
18	**Cost C2 = (Cost C1 + Risk Premium)**				65490.01	90.91
VI	**Cost C3**					
19	Managerial Cost				6549	9.09
20	**Cost C3 = (Cost C2 + Managerial Cost)**				72039.01	100
VII	**Economics of the Crop**					
a.	Main Product	a) Main Product (q)		63.26	189787.06	
		b) Main Crop Sales Price (Rs.)			3000	
b.	Gross Income (Rs.)				189787.06	
c.	Net Income (Rs.)				117748.05	
d.	Cost per Quintal (Rs./q.)				1138.73	
e.	Benefit Cost Ratio (BC Ratio)				1:2.6	

Cost of Cultivation of Bengal gram: The data regarding the cost of cultivation (Rs/ha) of Bengal gram in Lakshmapura-2 micro watershed is presented in Table 26.c. The results indicate, the total cost of cultivation (Rs/ha) for Bengal gram was Rs.16550.91. The gross income realized by the farmers was Rs. 74100.00. The net income from Bengal gram cultivation was Rs. 57549.09, thus the benefit cost ratio was found to be 1:4.50.

Table 26(c). Cost of Cultivation of Bengal gram in Lakshmapura-2 micro-watershed

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
I	**Cost A1**				
1	Hired Human Labour	Man days	18.11	2964	17.91
2	Bullock	Pairs/day	0.41	205.83	1.24
3	Tractor	Hours	4.94	4940	29.85
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	49.4	3705	22.39
6	Seed Inter Crop	Kgs.	0	0	0
7	FYM	Quintal	0	0	0
8	Fertilizer + micronutrients	Quintal	0.82	658.67	3.98
9	Pesticides (PPC)	Kgs / liters	0.41	494	2.98
10	Irrigation	Number	0	0	0
11	Repairs		0	0	0
12	Msc. Charges (Marketing costs etc)		0	0	0
13	Depreciation charges		0	0.01	0
14	Land revenue and Taxes		0	4.94	0.03
II	**Cost B1**				
16	Interest on working capital			582.92	3.52
17	**Cost B1 = (Cost A1 + sum of 15 and 16)**			13555.37	81.9
III	**Cost B2**				
18	Rental Value of Land			400	2.42
19	**Cost B2 = (Cost B1 + Rental value)**			13955.37	84.32
IV	**Cost C1**				
20	Family Human Labour		5.35	1090.92	6.59
21	**Cost C1 = (Cost B2 + Family Labour)**			15046.28	90.91
V	**Cost C2**				
22	Risk Premium			0	0
23	**Cost C2 = (Cost C1 + Risk Premium)**			15046.28	90.91
VI	**Cost C3**				
24	Managerial Cost			1504.63	9.09
25	**Cost C3 = (Cost C2 + Managerial Cost)**			16550.91	100
VII	**Economics of the Crop**				
a.	Main Product	a) Main Product (q)	37.05	74100	
		b) Main Crop Sales Price (Rs.)		2000	
b.	Gross Income (Rs.)			74100	
c.	Net Income (Rs.)			57549.09	
d.	Cost per Quintal (Rs./q.)			446.72	
e.	Benefit Cost Ratio (BC Ratio)			1:4.5	

Cost of Cultivation of Groundnut: The data regarding the cost of cultivation (Rs/ha) of Groundnut in Lakshmapura-2 micro watershed is presented in Table 26.d. The results indicate that, the total cost of cultivation (Rs/ha) for Groundnut was Rs. 43647.47. The gross income realized by the farmers was Rs.125266.01. The net income from Groundnut cultivation was Rs. 81618.54, thus the benefit cost ratio was found to be 1:2.90.

Table 26(d). Cost of Cultivation of Groundnut in Lakshmapura-2 micro-watershed

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
I	**Cost A1**				
1	Hired Human Labour	Man days	30.41	5569.26	12.76
2	Bullock	Pairs/day	3.82	1911.13	4.38
3	Tractor	Hours	2.64	2635.9	6.04
4	Machinery	Hours	0.08	77.19	0.18
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	116.96	14727.2	33.74
6	Seed Inter Crop	Kgs.	0	0	0
7	FYM	Quintal	2.35	2815.39	6.45
8	Fertilizer + micronutrients	Quintal	2.11	1876.61	4.3
9	Pesticides (PPC)	Kgs / liters	1.11	1337.05	3.06
10	Irrigation	Number	0	0	0
11	Repairs		0	0	0
12	Msc. Charges (Marketing costs etc)		0	0	0
13	Depreciation charges		0	24.85	0.06
14	Land revenue and Taxes		0	4.94	0.01
II	**Cost B1**				
16	Interest on working capital			2490.75	5.71
17	**Cost B1 = (Cost A1 + sum of 15 and 16)**			33470.27	76.68
III	**Cost B2**				
18	Rental Value of Land			616.71	1.41
19	**Cost B2 = (Cost B1 + Rental value)**			34086.98	78.1
IV	**Cost C1**				
20	Family Human Labour		26.37	5592.54	12.81
21	**Cost C1 = (Cost B2 + Family Labour)**			39679.52	90.91
V	**Cost C2**				
22	Risk Premium			0	0
23	**Cost C2 = (Cost C1 + Risk Premium)**			39679.52	90.91
VI	**Cost C3**				
24	Managerial Cost			3967.95	9.09
25	**Cost C3 = (Cost C2 + Managerial Cost)**			43647.47	100
VII	**Economics of the Crop**				
a.	Main Product	a) Main Product (q)	34.09	125266.01	
		b) Main Crop Sales Price (Rs.)		3675	
b.	Gross Income (Rs.)			125266.01	
c.	Net Income (Rs.)			81618.54	
d.	Cost per Quintal (Rs./q.)			1280.51	
e.	Benefit Cost Ratio (BC Ratio)			1:2.9	

Cost of Cultivation of Sorghum: The data regarding the cost of cultivation (Rs/ha) of Sorghum in Lakshmapura-2 micro watershed is presented in Table 26.e. The results indicate that, the total cost of cultivation (Rs/ha) for Sorghum was Rs.25637.44. The gross income realized by the farmers was Rs. 43336.42. The net income from Sorghum cultivation was Rs. 17698.97, thus the benefit cost ratio was found to be 1:1.70.

Table 26(e). Cost of Cultivation of Sorghum in Lakshmapura-2 micro-watershed

Sl.No	Particulars		Units	Phy Units	Value(Rs.)	% to C3
I	**Cost A1**					
1	Hired Human Labour		Man days	31.35	5545.57	21.63
2	Bullock		Pairs/day	2.33	1251.5	4.88
3	Tractor		Hours	2.97	2973.79	11.6
4	Machinery		Hours	0	0	0
5	Seed Main Crop (Establishment and Maintenance)		Kgs (Rs.)	12.82	1277.69	4.98
6	Seed Inter Crop		Kgs.	0	0	0
7	FYM		Quintal	1.17	1403.29	5.47
8	Fertilizer + micronutrients		Quintal	1.99	1575.55	6.15
9	Pesticides (PPC)		Kgs / liters	1.86	2233.64	8.71
10	Irrigation		Number	0	0	0
11	Repairs			0	0	0
12	Msc. Charges (Marketing costs etc)			0	0	0
13	Depreciation charges			0	2.03	0.01
14	Land revenue and Taxes			0	4.82	0.02
II	**Cost B1**					
16	Interest on working capital				778.82	3.04
17	**Cost B1 = (Cost A1 + sum of 15 and 16)**				17046.71	66.49
III	**Cost B2**					
18	Rental Value of Land				485.71	1.89
19	**Cost B2 = (Cost B1 + Rental value)**				17532.43	68.39
IV	**Cost C1**					
20	Family Human Labour			27.96	5774.34	22.52
21	**Cost C1 = (Cost B2 + Family Labour)**				23306.77	90.91
V	**Cost C2**					
22	Risk Premium				0	0
23	**Cost C2 = (Cost C1 + Risk Premium)**				23306.77	90.91
VI	**Cost C3**					
24	Managerial Cost				2330.68	9.09
25	**Cost C3 = (Cost C2 + Managerial Cost)**				25637.44	100
VII	**Economics of the Crop**					
a.	Main Product	a) Main Product (q)		30.64	43336.42	
		b) Main Crop Sales Price (Rs.)			1414.29	
b.	Gross Income (Rs.)				43336.42	
c.	Net Income (Rs.)				17698.97	
d.	Cost per Quintal (Rs./q.)				836.68	
e.	Benefit Cost Ratio (BC Ratio)				1:1.7	

Adequacy of fodder: The data regarding the adequacy of fodder in Lakshmapura-2 Micro watershed is presented in Table 27. The results indicate that, 17.14 per cent of the households opined that dry fodder was adequate.

Table 27. Adequacy of fodder in Lakshmapura-2 micro-watershed

Sl.No.	Particulars	LL (5)		MF (18)		SF (6)		SMF (5)		MDF (1)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Adequate-Dry Fodder	0	0	4	22.22	2	33.33	0	0	0	0	6	17.14

Average annual gross income: The data regarding the annual gross income in Lakshmapura-2 Micro watershed is presented in Table 28. The results indicate that, the farmers have annual gross income of Rs. 53057.14 in micro-watershed, of which Rs. 50914.29 is from agriculture itself.

Table 28. Average annual gross income in Lakshmapura-2 micro-watershed

Sl.No.	Particulars	LL (5)	MF (18)	SF (6)	SMF (5)	MDF (1)	All (35)
		Rs.	Rs.	Rs.	Rs.	Rs.	Rs.
1	Wage	0	1666.67	0	9000	0	2142.86
2	Agriculture	0	46111.1	73000	84800	90000	50914.3
	Income(Rs.)	0	47777.8	73000	93800	90000	53057.1

Average annual Expenditure: The data regarding the average annual expenditure in Lakshmapura-2 Micro watershed is presented in Table 29. The results indicate that, the farmers have annual gross expenditure of Rs. 91833.33 in micro-watershed, of which Rs. 16114.29 is from agriculture itself.

Table 29. Average annual Expenditure in Lakshmapura-2 micro-watershed

Sl.No.	Particulars	LL (5)	MF (18)	SF (6)	SMF (5)	MDF (1)	All (35)
		Rs.	Rs.	Rs.	Rs.	Rs.	Rs.
1	Wage	0	5000	0	6666.67	0	1000
2	Agriculture	0	17166.7	20000	23000	20000	16114.3
Total		0	22166.7	20000	29666.7	20000	91833.3

Horticulture species grown: The data regarding horticulture species grown in Lakshmapura-2 Micro watershed is presented in Table 30. The results indicate that, the total number of horticultural trees grown (both field and backyard) by the sampled households was Mango (18).

Table 30. Horticulture species grown in Lakshmapura-2 micro-watershed

Sl.No.	Particulars	LL (5)		MF (18)		SF (6)		SMF (5)		MDF (1)		All (35)	
		F	B	F	B	F	B	F	B	F	B	F	B
1	Mango	0	0	0	0	3	0	15	0	0	0	18	0

***F= Field B=Back Yard**

Forest species grown: The data regarding forest species grown in Lakshmapura-2 Micro watershed is presented in Table 31. The results indicate that, households have planted 27 neem trees, 6 banyan trees together in both field and backyard.

Table 31. Forest species grown in Lakshmapura-2 micro-watershed

Sl.No.	Particulars	LL (5)		MF (18)		SF (6)		SMF (5)		MDF (1)		All (35)	
		F	B	F	B	F	B	F	B	F	B	F	B
1	Neem	0	0	14	0	12	0	1	0	0	0	27	0
2	Banyan	0	0	4	0	2	0	0	0	0	0	6	0

***F= Field B=Back Yard**

Average additional investment capacity: The data regarding average additional investment capacity in Lakshmapura-2 Micro watershed is presented in Table 32. The results indicate that, households have an average investment capacity of Rs. 3485.71 for land development, Rs.1857.14 for adoption of improved livestock breeds and Rs. 114.29 for adoption of subsidiaries enterprises.

Table 32. Average additional investment capacity of households in Lakshmapura-2 micro-watershed

Sl.No.	Particulars	LL (5)	MF (18)	SF (6)	SMF (5)	MDF (1)	All (35)
		Rs.	Rs.	Rs.	Rs.	Rs.	Rs.
1	Land development	0	3500	3833.33	5800	7000	3485.71
2	Improved crop production	0	1388.89	3000	3400	5000	1857.14
3	Subsidiary enterprises	0	0	0	800	0	114.29

Source of funds for additional investment: The data regarding source of funds for additional investment in Lakshmapura-2 Micro watershed is presented in Table 33. The results indicate that, the sources of finance raised from bank as a loan and from own a source for land development was 48.57 and 2.86 per cent.

Table 33. Source of funds for additional investment in Lakshmapura-2 micro-watershed

Sl.No	Item	Land development		Improved crop production		Subsidiary enterprises	
		N	%	N	%	N	%
1	Loan from bank	17	48.57	14	40	1	2.86
2	Soft loan	1	2.86	0	0	0	0

Marketing of agricultural produce: The data regarding marketing of the agricultural produce in Lakshmapura-2 Micro watershed is presented in Table 34. The results indicated that, 100.00 per cent of output of Bengal gram was sold in the market with average price of Rs. 2000.00; 100.00 per cent of output of Groundnut was sold in the market with average price of Rs. 3675.00; 72.05 per cent of output of Maize was sold in the market with average price of Rs. 1200.00; 100.00 per cent of output of Sorghum was sold in the market with average price of Rs. 1414.29 and 100.00 per cent of output of Sunflower was sold in the market with average price of Rs. 3000.00.

Table 34. Marketing of agricultural produce in Lakshmapura-2 micro-watershed

Sl.No	Crops	Output obtained (q)	Output retained (q)	Output sold (q)	Output sold (%)	Avg. Price obtained (Rs/q)
1	Bengal gram	90	0	90	100	2000
2	Groundnut	285	0	285	100	3675
3	Maize	415	116	299	72	1200
4	Sorghum	248	0	248	100	1414
5	Sunflower	98	0	98	100	3000

Marketing channels used for sale of agricultural produce: The data regarding marketing channels used for sale of agricultural produce in Lakshmapura-2 Micro watershed is presented in Table 35. The results indicated that, 85.71 cent of the households have sold agricultural produce to the local/village merchants and 5.71 per cent of regulated market.

Table 35. Marketing channels used for sale of agricultural produce in Lakshmapura-2 micro-watershed

Sl. No.	Particulars	LL (5)		MF (18)		SF (6)		SMF (5)		MDF (1)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Local/village Merchant	0	0	18	100	4	66.7	6	120	2	200	30	85.71
2	Regulated Market	0	0	0	0	2	33.3	0	0	0	0	2	5.71

Mode of transport of agricultural produce: The data regarding mode of transport of agricultural produce in Lakshmapura-2 Micro watershed is presented in Table 36. The results indicated that, 91.43 cent of the households have used tractor for the transport of agriculture commodity.

Table 36. Mode of transport of agricultural produce in Lakshmapura-2 micro-watershed

Sl.No.	Particulars	LL (5)		MF (18)		SF (6)		SMF (5)		MDF (1)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Tractor	0	0	18	100	6	100	6	120	2	200	32	91.43

Incidence of soil and water erosion problems: The data regarding incidence of incidence of soil and water erosion problems in Lakshmapura-2 Micro watershed is presented in Table 37. The results indicate that, 82.86 per cent of the households have experienced soil and water erosion problems.

Table 37. Incidence of soil and water erosion problems in Lakshmapura-2 micro-watershed

Sl. No.	Particulars	LL (5)		MF (18)		SF (6)		SMF (5)		MDF (1)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Soil and water erosion problems in the farm	0	0	17	94	6	100	5	100	1	100	29	82.86

Interest towards soil testing: The data regarding Interest shown towards soil testing in Lakshmapura-2 Micro watershed is presented in Table 38. The results indicated that, 77.14 per cent of the households were interested towards soil testing.

Table 38. Interest regarding soil testing in Lakshmapura-2 micro-watershed

Sl.No.	Particulars	LL (5)		MF (18)		SF (6)		SMF (5)		MDF (1)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Interest in soil test	0	0	15	83	6	100	5	100	1	100	27	77.14

Usage pattern of fuel for domestic use: The data on usage pattern of fuel for domestic use in Lakshmapura-2 Micro watershed is presented in Table 39. The results indicated that, firewood was the major source of fuel for domestic use for 91.43 per cent of the households followed by LPG (2.86%) and Biogas (5.71 %).

Table 39. Usage pattern of fuel for domestic use in Lakshmapura-2 micro-watershed

Sl.No.	Particulars	LL (5)		MF (18)		SF (6)		SMF (5)		MDF (1)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Fire Wood	4	80	17	94.4	5	83.3	5	100	1	100	32	91.43
2	Biogas	0	0	1	5.56	1	16.7	0	0	0	0	2	5.71
3	LPG	1	20	0	0	0	0	0	0	0	0	1	2.86

Source of drinking water: The data on source of drinking water in Lakshmapura-2 Micro watershed is presented in Table 40. The results indicated that, piped waters supply was the major source for drinking water for 68.57 per cent of the households.

Table 40. Source of drinking water in Lakshmapura-2 micro-watershed

Sl. No.	Particulars	LL (5)		MF (18)		SF (6)		SMF (5)		MDF (1)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Piped supply	1	20	13	72.2	5	83.33	5	100	0	0	24	68.57
2	Bore Well	4	80	3	16.7	1	16.67	0	0	1	100	9	25.71

Source of light: The data on source of light in Lakshmapura-2 Micro watershed is presented in Table 41. The results indicated that, electricity was the major source of light for 97.14 per cent of the households.

Table 41. Source of light in Lakshmapura-2 micro-watershed

Sl.No.	Particulars	LL (5)		MF (18)		SF (6)		SMF (5)		MDF (1)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Electricity	5	100	17	94	6	100	5	100	1	100	34	97.1

Existence of sanitary toilet facility: The data on availability of toilet facility in Lakshmapura-2 Micro watershed is presented in Table 42. The results indicated that, 42.86 per cent of the households possess toilets.

Table 42. Existence of sanitary toilet facility in Lakshmapura-2 micro-watershed

Sl.No.	Particulars	LL (5)		MF (18)		SF (6)		SMF (5)		MDF (1)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Sanitary toilet facility	1	20	2	11	6	100	5	100	1	100	15	42.9

Possession of PDS card: The data regarding possession of PDS card in Lakshmapura-2 Micro watershed is presented in Table 43. The results indicated that, 97.14per cent of the households possessed BPL card.

Table 43. Possession of PDS card in Lakshmapura-2 micro-watershed

Sl.No.	Particulars	LL (5)		MF (18)		SF (6)		SMF (5)		MDF (1)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	BPL	5	100	17	94.4	6	100	5	100	1	100	34	97.14

Participation in NREGA programme: The data regarding Participation in NREGA programme in Lakshmapura-2 Micro watershed is presented in Table 44. The results indicated that, only 42.86 percent of the households have participated in NREGA programme.

Table 44. Participation in NREGA programme in Lakshmapura-2 micro-watershed

Sl. No.	Particulars	LL (5)		MF (18)		SF (6)		SMF (5)		MDF (1)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Participation in NREGA programme	0	0	11	61.1	4	66.7	0	0	0	0	15	42.9

Adequacy of food items: The data regarding adequacy of food items in Lakshmapura-2 Micro watershed is presented in Table 45. The results indicated that, the extent of adequacy of food items for cereals, pulses, Oilseeds and vegetables were 97.14, 97.14, 48.57, 17.14 per cent respectively, similarly for milk (22.86%).

Table 45. Adequacy of food items in Lakshmapura-2 micro-watershed

Sl. No.	Particulars	LL (5)		MF (18)		SF (6)		SMF (5)		MDF (1)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Cereals	5	100	17	94.4	6	100	5	100	1	100	34	97.14
2	Pulses	5	100	17	94.4	6	100	5	100	1	100	34	97.14
3	Oilseed	3	60	9	50	1	16.67	4	80	0	0	17	48.57
4	Vegetables	1	20	4	22.2	0	0	1	20	0	0	6	17.14
5	Milk	0	0	4	22.2	1	16.67	2	40	1	100	8	22.86

Table 46. Inadequacy of food items in Lakshmapura-2 micro-watershed

Sl.No.	Particulars	LL (5)		MF (18)		SF (6)		SMF (5)		MDF (1)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Oilseed	2	40	8	44.4	5	83.33	1	20	1	100	17	48.57
2	Vegetables	4	80	12	66.7	6	100	3	60	1	100	26	74.29
3	Fruits	5	100	16	88.9	6	100	5	100	1	100	33	94.29
4	Milk	5	100	12	66.7	5	83.33	3	60	0	0	25	71.43
5	Egg	5	100	16	88.9	6	100	4	80	1	100	32	91.43
6	Meat	5	100	16	88.9	6	100	4	80	1	100	32	91.43

Inadequacy of food items: The data regarding in adequacy of food items in Lakshmapura-2 Micro watershed is presented in Table 46. The results indicated that, the

extent of in adequacy of food items for Oilseeds and vegetables were 48.57, 74.29 per cent respectively, similarly for fruits (94.29%), milk (71.43%), egg (91.43%) and meat (91.43%).

Farming constraints: The data regarding farming constraints experienced by households in Lakshmapura-2 Micro watershed is presented in Table 47. The results indicated that, lower fertility status of the soil was the constraint experienced by (88.57 %) per cent of the households, wild animal menace on farm field (80.00%), frequent incidence of pest and diseases (71.43%), inadequacy of irrigation water (8.57%), high cost of fertilizers and plant protection chemicals (74.29%), high rate of interest on credit (62.86%), low price for the agricultural commodities (57.14 %), lack of marketing facilities in the area (57.14%), inadequate extension services (5.71 %), lack of transport for safe transport of the agricultural produce to the market (37.14%) and less rainfall (2.86%).

Table 47. Farming constraints experienced in Lakshmapura-2 micro-watershed

SN	Particulars	**LL** (5)		**MF** (18)		**SF** (6)		**SMF** (5)		**MDF** (1)		**All (35)**	
		N	**%**	**N**	**%**	**N**	**%**	**N**	**%**	**N**	**%**	**N**	**%**
1	Lower fertility status of the soil	0	0	19	105.56	6	100	5	100	1	100	31	88.57
2	Wild animal menace on farm field	0	0	16	88.89	6	100	5	100	1	100	28	80
3	Frequent incidence of pest and diseases	0	0	14	77.78	6	100	5	100	0	0	25	71.43
4	Inadequacy of irrigation water	0	0	2	11.11	1	16.67	0	0	0	0	3	8.57
5	High cost of Fertilizers and plant protection chemicals	0	0	15	83.33	6	100	4	80	1	100	26	74.29
6	High rate of interest on credit	0	0	14	77.78	5	83.33	3	60	0	0	22	62.86
7	Low price for the agricultural commodities	0	0	11	61.11	4	66.67	4	80	1	100	20	57.14
8	Lack of marketing facilities in the area	0	0	11	61.11	5	83.33	3	60	1	100	20	57.14
9	Inadequate extension services	0	0	1	5.56	0	0	1	20	0	0	2	5.71
10	Lack of transport for safe transport of the Agril produce to the market.	0	0	9	50	3	50	1	20	0	0	13	37.14
11	Less rainfall	0	0	1	5.56	0	0	0	0	0	0	1	2.86

Chapter 5

SUMMARY AND IMPLICATIONS

In order to assess the socio-economic condition of the farmers in the watershed 35 households located in the micro watershed were interviewed for the survey. The study was conducted in Lakshmapura-2 micro-watershed (Bhanapur sub-watershed, Koppala taluk & District) is located at North latitude 15^0 22' 54.598" and 15^0 21' 33.894" and East longitude 76^0 3' 55.966" and 76^0 1' 49.0989"covering an area of about 493.28 ha bounded by under Lakshmapura, Vadagenahalu, Tadakal and Haligeri Villages.

Socio-economic analysis of Lakshmapura-2 micro watersheds of Bhanapur sub-watershed, Koppala taluk & District indicated that, out of the total sample of 35 farmers were sampled in Lakshmapura-2 micro-watershed among households surveyed 18 (51.43%) were marginal, 6 (17.14%) were small, 5 (14.29 %) were semi medium and 1 (2.86 %) were medium farmers. 5 landless farmers were also interviewed for the survey. The population characteristics of households indicated that, there were 75 (60.48%) men and 49 (39.52 %) were women. The average population of landless was 2.8, marginal farmers were 3.8, small farmers were 3.7, semi medium farmers were 3.2 and medium farmers were 4. Majority of the respondents (37.10%) were in the age group of 16-35 years.

Education level of the sample households indicated that, there were 29.84 per cent illiterates, 70.96 per cent pre university education and 4.03 per cent attained graduation. About, 82.86 per cent of household heads practicing agriculture and 14.29 per cent of the household heads were engaged as agricultural labourers. Agriculture was the major occupation for 64.52 per cent of the household members. In the study area, 97.14 per cent of the households possess katcha house.

The durable assets owned by the households showed that, 100.00 per cent possess TV, 100.00 per cent possess mobile phones and 22.86 per cent possess motor cycles. Farm implements owned by the households indicated that, 2.86 per cent of the households possess plough, 8.57 per cent possess bullock cart. Regarding livestock possession by the households, 2.86 per cent possess local cow and 5.71 per cent possess buffalo. The average labour availability in the study area showed that**,** own labour men available in the micro watershed was 1.53, women available in the micro watershed was 1.17, hired labour (men) available was 10.63 and hired labour (women) available was 10.73.

Out of the total land holding of the sample respondents 87.34 per cent (32.07 ha) of the area is under dry condition and the remaining 12.66 per cent area is irrigated land. There were 4.00 live bore wells and 3.00 dry bore wells among the sampled households. Bore well was the major source of irrigation for 11.43 per cent of the households. The major crops grown by sample farmers are Maize, Sunflower, Bengal gram, Groundnut and Sorghum and cropping intensity was recorded as 81.77 per cent.

Out of the sample households 31.43 percent possessed bank account and 0.00 per cent of them have savings in the account. About 31.43 per cent of the respondents borrowed credit from various sources. The per hectare cost of cultivation for Maize, Sunflower, Bengal gram, Groundnut and Sorghum was Rs.53974.39 , 72039.01, 16550.91, 43647.47 and 25637.44 with benefit cost ratio of 1:1.10, 1: 2.60, 1: 4.50, 1: 2.90 and 1:1.70 respectively. Further, 17.14 per cent of the households opined that dry fodder was adequate.

The average annual gross income of the farmers was Rs. 53057.14 in micro-watershed, of which Rs. 50914.29 comes from agriculture. Sampled households have grown 18 horticulture trees and 33 forestry trees together in the fields and back yards. Households have an average investment capacity of Rs. 3485.71 for land development.

Source of funds for additional investment is concerned, 2.86 per cent depends on own funds and 48.57 per cent depends on bank loan for land development activities. Regarding marketing channels, 85.71 per cent of the households have sold agricultural produce to the local/village merchants, while, 5.71 per cent have sold in regulated markets. Further, 91.43 per cent of the households have used tractor for the transport of agriculture commodity.

Majority of the farmers (82.86%) have experienced soil and water erosion problems in the watershed and 77.14 per cent of the households were interested towards soil testing. Fire was the major source of fuel for domestic use for 91.43 per cent of the households and 2.86 per cent households has LPG connection. Piped supply was the major source for drinking water for 68.57 per cent of the households. Electricity was the major source of light for 97.14 per cent of the households. In the study area, 42.86 per cent of the households possess toilet facility.

Regarding possession of PDS card, 97.14 per cent of the households possessed BPL card. Households opined that, the requirement of cereals (97.14%), pulses (97.14%) and oilseeds (48.57%) are adequate for consumption. Farming constraints experienced by households in the micro watersheds were lower fertility status of the soil (88.57%) wild animal menace on farm field (80.00%), frequent incidence of pest and diseases (71.43%), inadequacy of irrigation water (8.57%), high cost of fertilizers and plant protection chemicals (74.29%), high rate of interest on credit (62.86%), low price for the agricultural commodities (57.14%), lack of marketing facilities in the area (57.14%), inadequate extension services (5.71%), lack of transport for safe transport of the agricultural produce to the market (37.14%), Less rainfall (2.86%).

Implications of the survey

- ✓ Result indicated that, there were 29.84 per cent were illiterate hence, extension methodologies such as demonstration, street play, drama, video shows will be effective in dissemination of the technologies in the micro watershed.

- ✓ The data indicate that, 97.14 per cent of the households possess katcha house. Hence, the development department while implementing the watershed plan should focus on agriculture to enhance the productivity of major crops in the area to increase the income of the farmers.
- ✓ Results indicated that the local institutional participation of the household members in the micro watershed is minimal hence, activities like membership campaign, awareness creation about the benefits of membership in local institutions and strengths of organized groups must be conveyed.
- ✓ Majority of the households in the watershed have experience in use of mobile phones, and television hence, these mass media can be effectively utilized for transfer of technology as well as for information dissemination.
- ✓ The farm machinery/implement possession in the micro watershed was found to be minimum the reasons may lack of knowledge or lack of financial ability which can be addressed through training on use of different farm implements, providing information on different sources of finance for purchase of farm implements.
- ✓ The possession of livestock such as crossbred cow found is less hence, farmers must be made aware of the benefits of crossbred cow in increased milk production.
- ✓ The possession of livestock such as sheep, goat and poultry was found to be low hence, farmers may be informed the role of subsidiary enterprises in enhancing the income and information on financial support for subsidiary activities.
- ✓ The data indicate that, job/work was the reason for all the migrants hence, farmers may be trained on profitable agriculture or self employment such has animal husbandry, plate making, sheep rearing, goat rearing, rabbit rearing with suitable information on sources of financial support.
- ✓ The results indicate that there was a change in quality of life due to migration hence, the developmental departments should take actions to arrest migration and to improve the quality of the life in rural areas.
- ✓ Households possess 28.01ha (87.34 %) of dry land and 4.06ha (12.66 %) of irrigated land hence, the availability of the dryland agricultural technologies such as short duration crops, high yielding drought resistance crop varieties, drip irrigation technology and subsidy information will be helpful for the farmers to enhance the productivity of land and as well as farmers income.
- ✓ Few of the bore well in micro watershed found non functional hence, farmers may be trained on possibility of bore well rejuvenation.
- ✓ Bore well was major source of irrigation for 11.43 per cent of the households. hence, in order to increase the area under irrigation as well as to increase the water use efficiency farmers may trained on drip irrigation and provide the information on subsidy for drip irrigation equipment's along with the information on different agencies which provides the financial assistance for drip irrigation.

- ✓ The cropping intensity in the micro watershed was found to be (81.77 %) hence, care must be taken by the implementing agency to bring uncultivated land into cultivation through suitable measures.
- ✓ Many of the household members have borrowed loan from cooperative banks which has higher rate of interest hence, farmers may be sensitized on the different sources of credit with lesser interest rate such SHGs etc.
- ✓ The results indicated the non availability of both green and dry fodder throughout the year hence, fodder development activities can be taken up in the micro watershed.
- ✓ The average annual gross income of the households Rs.50914.29 from agriculture and Rs. 2142.86 from wages. Agriculture was found to be the major source of income for households hence; the development activities should focus on productivity enhancement, marketing arrangements and agricultural technology dissemination to have a direct impact on the farmers.
- ✓ The cultivation of forest species is found minimal hence, information and production technology related to agro-forestry and integrated farming system.
- ✓ The data indicated that, 82.86 per cent of the households have experienced soil and water erosion problems. Hence, those farmers who reported the soil and water erosion problems may be given attention while implementation of the watershed development plan.
- ✓ The data indicated that, 77.14 per cent of the households have interest in soil testing hence, farmers must be provided with the information on various institutions which are involved in soil testing for the benefit of the farmers.
- ✓ Except summer ploughing the adoption of other soil and water conservation structures is minimum hence, the farmers in the micro watershed should be sensitized on the use of different conservation structures for soil water conservation.
- ✓ Cereals and pulses found be adequate for per cent of the households respectively hence, farm households and the farm women must be trained on importance of balanced nutrition and role of vegetable, milk, egg, meat in balanced diet.
- ✓ Lower fertility status of the soil (88.57%), wild animal menace on farm field (80.00%), frequent incidence of pest and diseases (71.43%), high cost of fertilizers and plant protection chemicals (74.29%), high rate of interest on credit (62.86%), low price for the agricultural commodities (57.14%), lack of marketing facilities in the area (57.14%), inadequate extension services (5.71%), lack of transport for safe transport of the agricultural produce to the market (37.14%) were the major farming constraints experienced hence, these constraints must be addressed immediately for the welfare of the farmers. Awareness to be created among the farmers to approach nearest KVKs/RSKs and other developmental departments for technical and for subsidized inputs and utilize the well established regulated markets, approaching the contract firms, direct markets to avoid the involvement of middlemen.