



हर कदम, हर डगर
किसानों का हमसफर
भारतीय कृषि अनुसंधान परिषद

AgriSearch with a human touch

**LAND RESOURCE INVENTORY AND SOCIO-ECONOMIC STATUS OF
FARM HOUSEHOLDS FOR WATERSHED PLANNING AND
DEVELOPMENT**

VOJENAHALLI-2 (4D4A1X1d) MICROWATERSHED

Koppal Taluk and District, Karnataka

Karnataka Watershed Development Project – II

SUJALA – III

World Bank funded Project



THE WORLD BANK



ICAR – NATIONAL BUREAU OF SOIL SURVEY AND LAND USE PLANNING



ICAR - NBSS & LUP

**WATERSHED DEVELOPMENT DEPARTMENT
GOVT. OF KARNATAKA, BANGALORE**



About ICAR - NBSS&LUP

The ICAR-National Bureau of Soil Survey and Land Use Planning (ICAR-NBSS&LUP), Nagpur, a premier Institute of the Indian Council of Agricultural Research (ICAR), was set up during 1976 with the objective to prepare soil resource maps at national, state and district levels and to provide research inputs in soil resource mapping and its applications, land evaluation, land use planning, land resource management, and database management using GIS for optimising land use on different kinds of soils in the country.

The Bureau has been engaged in carrying out soil resource survey, agro-ecological and soil degradation mapping at the country, state and district levels for qualitative assessment and monitoring the soil health towards viable land use planning. The research activities have resulted in identifying the soil potentials and problems, and the various applications of the soil surveys with the ultimate objective of sustainable agricultural development. The Bureau has the mandate to correlate and classify soils of the country and maintain a National Register of all the established soil series. The Institute is also imparting in-service training to staff of the soil survey agencies in the area of soil survey, land evaluation and soil survey interpretations for land use planning. The Bureau in collaboration with Panjabrao Krishi Vidyapeeth, Akola is running post-graduate teaching and research programme in land resource management, leading to M.Sc. and Ph.D. degrees.

Citation: Rajendra Hegde, Ramesh Kumar, S.C., K.V. Niranjana, S. Srinivas, M.Lalitha, B.A. Dhanorkar, R.S. Reddy and S.K. Singh (2019). "Land resource inventory and socio-economic status of farm households for watershed planning and development of Vojenahalli-2 (4D4A1X1d) Microwatershed, Koppal Taluk and District, Karnataka", Sujala MWS Publ .437, ICAR – NBSS & LUP, RC, Bangalore. p.133 & 37..

TO OBTAIN COPIES,

Please write to:

Director, ICAR - NBSS & LUP,

Amaravati Road, NAGPUR - 440 033, India

Phone : (0712) 2500386, 2500664, 2500545 (O)

Telefax : 0712-2522534

E-Mail : director@nbsslup.ernet.in

Website URL : nbsslup.in

Or

Head, Regional Centre, ICAR - NBSS&LUP, Hebbal, Bangalore - 560 024

Phone : (080) 23412242, 23510350 (O)

Telefax : 080-23510350

E-Mail : nbssrcb@gmail.com



**LAND RESOURCE INVENTORY AND SOCIO-ECONOMIC
STATUS OF FARM HOUSEHOLDS FOR WATERSHED
PLANNING AND DEVELOPMENT**

VOJENAHALLI-2 (4D4A1X1d) MICROWATERSHED

Koppal Taluk and District, Karnataka

Karnataka Watershed Development Project – II

Sujala-III

World Bank funded Project



**ICAR – NATIONAL BUREAU OF SOIL SURVEY AND LAND
USE PLANNING**



ICAR - NBSS & LUP



**WATERSHED DEVELOPMENT DEPARTMENT, GOVT. OF
KARNATAKA, BANGALORE**



PREFACE

In Karnataka, as in other Indian States, the livelihoods of rural people are intertwined with farming pursuits. The challenges in agriculture are seriously threatening the livelihood of a large number of farmers as they have been practicing farming in contextual factors beyond their control. Climatic factors are the most important ones and have become much more significant in recent times due to rapid climate changes induced by intensive anthropogenic activities affecting our ecosystem in multiple ways. Climate change has become the reality, it is happening and efforts to evolve and demonstrate climate resilient technologies have become essential. Due to the already over stressed scenario of agrarian sector, the climate change is resulting in manifold increase in the complexities, pushing the rural mass to face more and more unpredictable situations. The rising temperatures and unpredictable rainfall patterns are going to test seriously the informed decisions farmers have to make in order to survive in farming and sustain their livelihood.

It is generally recognized that impacts of climate change shall not be uniform across the globe. It is said that impact of climate change is more severe in South Asia. Based on the analysis of meteorological data, it is predicted that in India, there will be upward trend in mean temperature, downward trend in relative humidity, annual rainfall and number of wet days in a year. Also, in general, phenomena like erratic monsoon, spread of tropical diseases, rise in sea levels, changes in availability of fresh water, frequent floods, droughts, heat waves, storms and hurricanes are predicted. Each one of these adverse situations are already being experienced in various parts of India and also at the global level. Decline in agricultural productivity of small and marginal farmers becoming more vulnerable is already witnessed.

In Karnataka, more than 60 per cent of the population live in rural areas and depend on agriculture and allied activities for their livelihood. Though the state has achieved significant progress in increasing the yield of many crops, there is tremendous pressure on the land resources due to the growing and competing demands of various land uses. This is reflected in the alarming rate of land degradation observed. Already more than 50 per cent of the area is affected by various forms of degradation. If this trend continues, the sustainability of the fragile ecosystem will be badly affected. The adverse effects of change in the climatic factors are putting additional stress on the land resources and the farmers dependent on this.

The natural resources (land, water and vegetation) of the state need adequate and constant care and management, backed by site-specific technological interventions and investments particularly by the government. Detailed database pertaining to the nature of

the land resources, their constraints, inherent potentials and suitability for various land based rural enterprises, crops and other uses is a prerequisite for preparing location-specific action plans, which are in tune with the inherent capability of the resources. Any effort to evolve climate resilient technologies has to be based on the baseline scientific database. Then only one can expect effective implementation of climate resilient technologies, monitor the progress, make essential review of the strategy, and finally evaluate the effectiveness of the implemented programs. The information available at present on the land resources of the state are of general nature and useful only for general purpose planning. Since the need of the hour is to have site-specific information suitable for farm level planning and detailed characterization and delineation of the existing land resources of an area into similar management units is the only option.

ICAR-NBSS&LUP, Regional Centre, Bangalore has taken up a project sponsored by the Karnataka Watershed Development Project-II, (Sujala-III), Government of Karnataka funded by the World Bank under Component -1 Land Resource Inventory. This study was taken up to demonstrate the utility of such a database in reviewing, monitoring and evaluating all the land based watershed development programs on a scientific footing. To meet the requirements of various land use planners at grassroots level, the present study on “Land Resource Inventory and Socio-Economic Status of Farm Households for Watershed Planning and Development of for Vojenahalli-2 microwatershed in Koppal Taluk, and District, Karnataka” for integrated development was taken up in collaboration with the State Agricultural Universities, IISC, KRSRAC, KSNDMC as Consortia partners. The project provides detailed land resource information at cadastral level (1:7920 scale) for all the plots and socio-economic status of farm households covering thirty per cent farmers randomly selected representing landed and landless class of farmers in the micro-watershed. The project report with the accompanying maps for the microwatershed will provide required detailed database for evolving effective land use plan, alternative land use options and conservation plans for the planners, administrators, agricultural extension personnel, KVK officials, developmental departments and other land users to manage the land resources in a sustainable manner.

It is hoped that this database will be useful to the planners, administrators and developmental agencies working in the area in not only for formulating location specific developmental schemes but also for their effective monitoring at the village/watershed level.

Nagpur

Date: 22-10-2019

S.K. SINGH

Director, ICAR - NBSS&LUP Nagpur

Contributors

Dr. Rajendra Hegde Principal Scientist, Head & Project Leader, Sujala-III Project ICAR-NBSS&LUP, Regional Centre, Bangalore	Dr. S.K.Singh Director, ICAR-NBSS&LUP Coordinator, Sujala-III Project Nagpur
Soil Survey, Mapping & Report Preparation	
Dr. K.V. Niranjana	Sh. R.S. Reddy
Dr. B.A. Dhanorkar	Smt. Chaitra, S.P.
	Dr. Gopali Bardhan
	Mr. Somashekar T.N
	Ms. Arpitha G.M
	Dr. Mahendra kumar M.B
Field Work	
Sh. C. Bache Gowda	Sh. Mayur Patil
Sh. Somashekar	Sh. Arun Kumar, S.
Sh. M. Jayaramaiah	Sh. Sunil Raj
	Sh. Yogesh Kumar, B.
	Sh. Vikas, N.K.
	Sh. Arun Kumar, S.G.
	Sh. Umesh Jadiyahappa Madolli
	Sh. Praveen Kumar P. Achalkar
	Sh. Veerabhadraswamy
	Sh. Vinay
	Sh. Shankarappa, K.
	Sh. Lankesh, R.S.
	Sh. Appanna B. Hattigoudar
	Sh. Maharudra
GIS Work	
Dr. S.Srinivas	Sh. A.G.Devendra Prasad
Sh. D.H.Venkatesh	Sh. Abhijith Sastry, N.S.
Smt. K.Sujatha	Smt. Shyla, B.
Smt. K.V.Archana	Smt. Swetha ,K.
Sh. N.Maddileti	Ms. Vidya, P.C.
	Sh. Deepak, M.J.
	Smt. K.Karunya Lakshmi
	Ms. Seema, K.V.

Laboratory Analysis	
Dr. M. Lalitha	Sh. Vindhya, N.G.
Smt. Arti Koyal	Ms. P. Pavanakumari, P.
Smt. Parvathy, S.	Ms. Rashmi, N.
	Ms. Leelavathy, K.U.
	Smt. Usha Kiran, G.
Socio-Economic Analysis	
Dr. S.C. Ramesh Kumar	Sh. M.K. Prakashanaik
	Ms. Shraddha Hegde
	Mrs. Sowmya A N
	Sh. Vijay Kumar
	Sh. Basavaraja
	Sh. Vinod R
	Ms. Sowmya K.B
	Mrs. Prathibha, D.G
	Sh. Rajendra,D
Soil & Water Conservation	
Sh. Sunil P. Maske	
Watershed Development Department, GoK, Bangalore	
Sh. Rajeev Ranjan IFS Project Director & Commissioner, WDD	Dr. A. Natarajan NRM Consultant, Sujala-III Project
Dr. S.D. Pathak IFS Executive Director & Chief Conservator of Forests, WDD	

PART-A

LAND RESOURCE INVENTORY

Contents

Preface		
Contributors		
Executive Summary		
Chapter 1	Introduction	1
Chapter 2	Geographical Setting	3
2.1	Location and Extent	3
2.2	Geology	3
2.3	Physiography	4
2.4	Drainage	5
2.5	Climate	5
2.6	Natural Vegetation	6
2.7	Land Utilization	7
Chapter 3	Survey Methodology	11
3.1	Base maps	11
3.2	Image Interpretation for Physiography	11
3.3	Field Investigation	14
3.4	Soil mapping	16
3.5	Land Management Units	16
3.6	Laboratory Characterization	17
Chapter 4	The Soils	21
4.1	Soils of Granite Gneiss Landscape	21
4.2	Soils of Alluvial Landscape	23
Chapter 5	Interpretation for Land Resource Management	35
5.1	Land Capability Classification	35
5.2	Soil Depth	37
5.3	Surface Soil Texture	38
5.4	Soil Gravelliness	39
5.5	Available Water Capacity	40
5.6	Soil Slope	41
5.7	Soil Erosion	42
Chapter 6	Fertility Status	45
6.1	Soil Reaction (pH)	45
6.2	Electrical Conductivity (EC)	45
6.3	Organic Carbon (OC)	45
6.4	Available Phosphorus	45
6.5	Available Potassium	47
6.6	Available Sulphur	47
6.7	Available Boron	47
6.8	Available Iron	48
6.9	Available Manganese	48
6.10	Available Copper	48
6.11	Available Zinc	52

Chapter 7	Land Suitability for Major Crops	53
7.1	Land suitability for Sorghum	53
7.2	Land suitability for Maize	54
7.3	Land suitability for Bajra	55
7.4	Land suitability for Redgram	56
7.5	Land suitability for Bengalgram	57
7.6	Land suitability for Groundnut	58
7.7	Land suitability for Sunflower	59
7.8	Land suitability for Cotton	60
7.9	Land suitability for Chilli	61
7.10	Land suitability for Tomato	62
7.11	Land suitability for Brinjal	63
7.12	Land suitability for Onion	64
7.13	Land suitability for Bhendi	65
7.14	Land suitability for Drumstick	66
7.15	Land suitability for Mulberry	67
7.16	Land suitability for Mango	68
7.17	Land Suitability for Sapota	69
7.18	Land suitability for Pomegranate	70
7.19	Land suitability for Guava	71
7.20	Land Suitability for Jackfruit	72
7.21	Land Suitability for Jamun	73
7.22	Land Suitability for Musambi	74
7.23	Land Suitability for Lime	75
7.24	Land Suitability for Cashew	76
7.25	Land Suitability for Custard apple	77
7.26	Land suitability for Amla	78
7.27	Land suitability for Tamarind	79
7.28	Land suitability for Marigold	80
7.29	Land suitability for Chrysanthemum	81
7.30	Land suitability for Jasmine	82
7.31	Land suitability for Crossandra	83
7.32	Land Management Units	117
7.33	Proposed Crop Plan	118
Chapter 8	Soil Health Management	121
Chapter 9	Soil and Water conservation Treatment Plan	125
9.1	Treatment Plan	125
9.2	Recommended Soil and Water Conservation measures	129
9.3	Greening of microwatershed	130
	References	133
	Appendix I	I-VIII
	Appendix II	IX-XIV
	Appendix III	XV-XX

LIST OF TABLES

2.1	Mean Monthly Rainfall, PET, 1/2 PET at Koppal Taluk and District	5
2.2	Land Utilization in Koppal District	7
3.1	Differentiating Characteristics used for Identifying Soil Series	15
3.2	Soil map unit description of Vojenahalli-2 microwatershed	17
4.1	Physical and chemical characteristics of soil series identified in Vojenahalli-2 microwatershed	27
7.1	Soil-Site Characteristics of Vojenahalli-2 microwatershed	85
7.2	Land suitability for Sorghum	86
7.3	Land suitability for Maize	87
7.4	Land suitability for Bajra	88
7.5	Land suitability for Redgram	89
7.6	Land suitability for Bengalgram	90
7.7	Land suitability for Groundnut	91
7.8	Land suitability for Sunflower	92
7.9	Land suitability for Cotton	93
7.10	Land suitability for Chilli	94
7.11	Land suitability for Tomato	95
7.12	Land suitability for Brinjal	96
7.13	Land suitability for Onion	97
7.14	Land suitability for Bhendi	98
7.15	Land suitability for Drumstick	99
7.16	Land suitability for Mulberry	100
7.17	Land suitability for Mango	101
7.18	Land Suitability for Sapota	102
7.19	Land suitability for Pomegranate	103
7.20	Land suitability for Guava	104
7.21	Land suitability for Jackfruit	105
7.22	Land suitability for Jamun	106
7.23	Land Suitability for Musambi	107
7.24	Land Suitability for Lime	108

7.25	Land Suitability for Cashew	109
7.26	Land Suitability for Custard apple	110
7.27	Land Suitability for Amla	111
7.28	Land Suitability for Tamarind	112
7.29	Land Suitability for Marigold	113
7.30	Land Suitability for Chrysanthemum	114
7.31	Land suitability for Jasmine	115
7.32	Land suitability for Crossandra	116
7.33	Proposed Crop Plan for Vojenahalli-2 Microwatershed	119

LIST OF FIGURES

2.1	Location map of Vojenahalli-2 Microwatershed	3
2.2a	Granite and granite gneiss rocks	4
2.2b	Alluvial rocks	4
2.3	Rainfall distribution in Koppal Taluk, Koppal District	6
2.4	Natural vegetation of Vojenahalli-2 microwatershed	6
2.5	Different crops and cropping systems in Vojenahalli-2 Microwatershed	7
2.6	Current Land use – Vojenahalli-2 Microwatershed	9
2.7	Location of Wells- Vojenahalli-2 Microwatershed	9
3.1	Scanned and Digitized Cadastral map of Vojenahalli-2 Microwatershed	13
3.2	Satellite image of Vojenahalli-2 Microwatershed	13
3.3	Cadastral map overlaid on IRS PAN+LISS IV merged imagery of Vojenahalli-2 Microwatershed	14
3.4	Location of profiles in a transect	15
3.5	Soil phase or management units of Vojenahalli-2 Microwatershed	19
5.1	Land Capability Classification of Vojenahalli-2 Microwatershed	36
5.2	Soil Depth map of Vojenahalli-2 Microwatershed	37
5.3	Surface Soil Texture map of Vojenahalli-2 Microwatershed	39
5.4	Soil Gravelliness map of Vojenahalli-2 Microwatershed	40
5.5	Soil Available Water Capacity map of Vojenahalli-2 Microwatershed	41
5.6	Soil Slope map of Vojenahalli-2 Microwatershed	42
5.7	Soil Erosion map of Vojenahalli-2 Microwatershed	43
6.1	Soil Reaction (pH) map of Vojenahalli-2 Microwatershed	46
6.2	Electrical Conductivity (EC) map of Vojenahalli-2 Microwatershed	46
6.3	Soil Organic Carbon (OC) map of Vojenahalli-2 Microwatershed	47
6.4	Soil Available Phosphorus map of Vojenahalli-2 Microwatershed	48
6.5	Soil Available Potassium map of Vojenahalli-2 Microwatershed	49
6.6	Soil Available Sulphur map of Vojenahalli-2 Microwatershed	49
6.7	Soil Available Boron map of Vojenahalli-2 Microwatershed	50
6.8	Soil Available Iron map of Vojenahalli-2 Microwatershed	50
6.9	Soil Available Manganese map of Vojenahalli-2 Microwatershed	51
6.10	Soil Available Copper map of Vojenahalli-2 Microwatershed	51

6.11	Soil Available Zinc map of Vojenahalli-2 Microwatershed	52
7.1	Land suitability map of Sorghum	54
7.2	Land suitability map of Maize	55
7.3	Land suitability map of Bajra	56
7.4	Land suitability map of Redgram	57
7.5	Land suitability map of Bengalgram	58
7.6	Land suitability map of Groundnut	59
7.7	Land suitability map of Sunflower	60
7.8	Land suitability map of Cotton	61
7.9	Land suitability map of Chilli	62
7.10	Land suitability map of Tomato	63
7.11	Land suitability map of Brinjal	64
7.12	Land suitability map of Onion	65
7.13	Land suitability map of Bhendi	66
7.14	Land suitability map of Drumstick	67
7.15	Land suitability map of Mulberry	68
7.16	Land suitability map of Mango	69
7.17	Land Suitability map of Sapota	70
7.18	Land suitability for Pomegranate	71
7.19	Land suitability map of Guava	72
7.20	Land Suitability map of Jackfruit	73
7.21	Land Suitability map of Jamun	74
7.22	Land Suitability map of Musambi	75
7.23	Land Suitability map of Lime	76
7.24	Land Suitability map of Cashew	77
7.25	Land Suitability map of Custard apple	78
7.26	Land suitability map of Amla	79
7.27	Land suitability map of Tamarind	80
7.28	Land suitability map of Marigold	81
7.29	Land suitability map of Chrysanthemum	82
7.30	Land suitability map of Jasmine	83
7.31	Land suitability map of Crossandra	84
7.32	Land Management Units map of Vojenahalli-2 microwatershed	117
9.1	Soil and water conservation map of Vojenahalli-2 microwatershed	130

EXECUTIVE SUMMARY

The land resource inventory of Vojenahalli-2 microwatershed was conducted using village cadastral maps and IRS satellite imagery on 1:7920 scale. The false colour composites of IRS imagery were interpreted for physiography and these physiographic delineations were used as base for mapping soils. The soils were studied in several transects and a soil map was prepared with phases of soil series as mapping units. Random checks were made all over the area outside the transects to confirm and validate the soil map unit boundaries. The soil map shows the geographic distribution and extent, characteristics, classification, behavior and use potentials of the soils in the Microwatershed.

The present study covers an area of 576 ha in Koppal taluk and district, Karnataka. The climate is semiarid and categorized as drought - prone with an average annual rainfall of 662 mm, of which about 424 mm is received during south –west monsoon, 161 mm during north-east and the remaining 77 mm during the rest of the year. An area of about 69 per cent is covered by soil, 31 per cent by water bodies, settlements and <1 per cent by Mining/Industrial. The salient findings from the land resource inventory are summarized briefly below.

- ❖ *The soils belong to 7 soil series and 12 soil phases (management units) and 3 land use classes.*
- ❖ *The length of crop growing period is <90 days and starts from 2nd week of August to 2nd week of November.*
- ❖ *From the master soil map, several interpretative and thematic maps like land capability, soil depth, surface soil texture, soil gravelliness, available water capacity, soil slope and soil erosion were generated.*
- ❖ *Soil fertility status maps for macro and micronutrients were generated based on the surface soil samples collected at every 320 m grid interval.*
- ❖ *Land suitability for growing 28 major agricultural and horticultural crops were assessed and maps showing the degree of suitability along with constraints were generated.*
- ❖ *Entire area is suitable for agriculture.*
- ❖ *About 6 per cent of the soils are moderately shallow (50-75 cm), 21 per cent moderately deep (75- 100 cm), 42 per cent is deep to very deep (100->150cm) soils.*
- ❖ *About 3 per cent loamy (sandy clay loam) and 65 per cent has clayey (sandy clay and clay) soils at the surface.*
- ❖ *About 62 per cent of the area has non-gravelly (<15%) soils and 7 per cent has gravelly soils (15-35 % gravel) soils.*
- ❖ *With respect to available water capacity 4 per cent of the area has very low (<50mm/m), 11 per cent of the area has low (51-100 mm/m), 17 per cent medium*

(101-150 mm/m) and 37 per cent area is high to very high (151->200mm/m) in available water capacity.

- ❖ An area of about <1 per cent has nearly level (0-1%) and 68 per cent has very gently sloping (1-3%) lands.
- ❖ An area of about 34 per cent is slightly eroded (e1) and 34 per cent is moderately eroded (e2).
- ❖ An area of about 1 per cent slightly alkaline (pH 7.3 to 7.8), 15 per cent moderately alkaline (pH 7.8 to 8.4), 50 per cent strongly alkaline (pH 8.4 to 9.0) and 3 per cent very strongly alkaline (pH >9.0).
- ❖ The Electrical Conductivity (EC) of the soils are <2 dsm⁻¹ indicating that soils are non saline.
- ❖ Organic carbon is low (<0.5%) in 18 per cent and medium (0.5-0.75%) in 51 per cent area of the microwatershed.
- ❖ Available phosphorus is medium (<23 kg/ha) in the entire area of the soils.
- ❖ Available potassium is medium (145-337 kg/ha) in 61 per cent and high (>337 kg/ha) in 8 per cent area of the soils.
- ❖ Available sulphur is low (<10 ppm) in 35 per cent, medium (10-20 ppm) in 22 per cent and high (>20 ppm) in 12 per cent area of the soils.
- ❖ Available boron is low (<0.5 ppm) in 10 per cent, medium (0.5-1.0 ppm) in 57 per cent and high (>1.0 ppm) in 1 per cent area of the microwatershed.
- ❖ Available iron is deficient (<4.5ppm) in 69 per cent and sufficient (>4.5 ppm) in < 1 per cent of the area.
- ❖ Available zinc is deficient (<0.6 ppm) in 35 per cent and sufficient (>0.6 ppm) in 34 per cent of the microwatershed.
- ❖ Available manganese and copper are sufficient in the entire area.
- ❖ The land suitability for 31 major agricultural and horticultural crops grown in the microwatershed was assessed and the areas that are highly suitable (class S1) and moderately suitable (class S2) are given below. It is however to be noted that a given soil may be suitable for various crops but what specific crop to be grown may be decided by the farmer looking to his capacity to invest on various inputs, marketing infrastructure, market price, and finally the demand and supply position.

Land suitability for various crops in the microwatershed

Crop	Suitability Area in ha (%)		Crop	Suitability Area in ha (%)	
	Highly suitable (S1)	Moderately suitable (S2)		Highly suitable (S1)	Moderately suitable (S2)
Sorghum	28(5)	346 (60)	Sapota	28(5)	21(4)
Maize	28(5)	347 (60)	Pomegranate	28(5)	331(58)
Bajra	28(5)	368(64)	Guava	-	49(9)
Redgram	28(5)	212(37)	Jackfruit	28(5)	21(4)
Bengal gram	-	375(65)	Jamun	28(5)	233(41)
Groundnut	-	49(9)	Musambi	28(5)	331(58)
Sunflower	28 (5)	310 (54)	Lime	28(5)	331(58)
Cotton	28(5)	347(60)	Cashew	-	49(9)
Chilli	28(5)	-	Custard apple	28(5)	368(64)
Tomato	28(5)	-	Amla	28(5)	368(64)
Brinjal	-	375(65)	Tamarind	28(5)	212(37)
Onion	-	28(5)	Marigold	28(5)	347(60)
Bhendi	-	375(65)	Chrysanthemum	28(5)	347 (60)
Drumstick	28(5)	310(54)	Jasmine	28(5)	36(6)
Mulberry	28(5)	281(49)	Crossandra	28(5)	98(17)
Mango	28(5)	98(17)	-	-	-

Apart from the individual crop suitability, a proposed crop plan has been prepared for the 3 identified LMUs by considering only the highly and moderately suitable lands for different crops and cropping systems with food, fodder, fibre and other horticulture crops.

- ❖ *Maintaining soil-health is vital for crop production and conserve soil and land resource base for maintaining ecological balance and to mitigate climate change. For this, several ameliorative measures have been suggested to these problematic soils like saline/alkali, highly eroded, sandy soils etc.,*
- ❖ *Soil and water conservation and drainage line treatment plans have been prepared that would help in identifying the sites to be treated and also the type of structures required.*
- ❖ *As part of the greening programme, several tree species have been suggested to be planted in marginal and submarginal lands, field bunds and also in the hillocks, mounds and ridges. That would help in supplementing the farm income, provide fodder and fuel, and generate lot of biomass which in turn would help in maintaining the ecological balance and contribute to mitigating the climate change.*

INTRODUCTION

Land is a scarce resource and basic unit for any material production. It can support the needs of the growing population, provided they use the land in a rational and judicious manner. But what is happening in many areas of the state is a cause for concern to everyone involved in the management of land resources at the grassroots level. The area available for agriculture is about 51 per cent of the total area and more than 60 per cent of the people are still dependant on agriculture for their livelihood. The limited land area is under severe stress and strain due to increasing population pressure and competing demands of various land uses. Due to this, every year there is significant diversion of farm lands and water resources for non-agricultural purposes. Apart from this, due to lack of interest in farmers for farming, large tracts of cultivable lands are turning into fallows in many areas and this trend is continuing at an alarming rate.

Further, land degradation has emerged as a serious problem which has already affected about 38 lakh ha of cultivated area in the state. Soil erosion alone has degraded about 35 lakh ha. Almost all the uncultivated areas are facing various degrees of degradation, particularly soil erosion. Salinity and alkalinity has emerged as a major problem in more than 3.5 lakh ha in the irrigated areas of the state. Nutrient depletion and declining factor productivity is common in both rainfed and irrigated areas. The degradation is continuing at an alarming rate and there appears to be no systematic effort among the stakeholders to contain this process. In recent times, an aberration of weather due to climate change phenomenon has added another dimension leading to unpredictable situations to be tackled by the farmers.

In this critical juncture, the challenge before us is not only to increase the productivity per unit area which is steadily declining and showing a fatigue syndrome, but also to prevent or at least reduce the severity of degradation. If the situation is not reversed at the earliest, then the sustainability of the already fragile crop production system and the overall ecosystem will be badly affected in the state. The continued neglect and unscientific use of the resources for a long time has led to the situation observed at present in the state. It is a known fact and established beyond doubt by many studies in the past that the cause for all kinds of degradation is the neglect and irrational use of the land resources. Hence, there is urgent need to generate a detailed site-specific farm level database on various land resources for all the villages/watersheds in a time bound manner that would help to protect the valuable soil and land resources and also to stabilize the farm production.

Therefore, the land resource inventory required for farm level planning is the one which investigates not only the surface but also consider the other parameters which are critical for productivity *viz.*, soils, climate, water, minerals and rocks, topography, geology, hydrology, vegetation, crops, land use pattern, animal population, socio-

economic conditions, infrastructure, marketing facilities and various schemes and developmental works of the government etc. From the data collected at farm level, the specific problems and potentials of the area can be identified and highlighted, conservation measures required for the area can be planned on a scientific footing, suitability of the area for various uses can be worked out and finally viable and sustainable land use options suitable for each and every land holding can be prescribed.

The Land Resource Inventory is basically done for identifying potential and problem areas, developing sustainable land use plans, estimation of surface run off and water harvesting potential, preparation of soil and water conservation plans, land degradation/desertification etc. The Bureau is presently engaged in developing an LRI methodology using high resolution satellite remote sensing data and Digital Elevation Model (DEM) data to prepare Landscape Ecological Units (LEU) map representing agro-ecosystem as a whole. The LEU is preferred over landform as the base map for LRI. LEU is the assemblage of landform, slope and land use. An attempt was made to upscale the soil resource information from 1:250000 and 1:50000 scale to the LEU map in Goa and other states.

The land resource inventory aims to provide site-specific database for Vojenahalli-2 Microwatershed in Koppal Taluk, Koppal District, Karnataka State for the Karnataka Watershed Development Department. The database was generated by using cadastral map of the village as a base along with high resolution IRS LISS IV and Cartosat-1 merged satellite imagery. Later, an attempt will be made to uplink this LRI data generated at 1:7920 scale under Sujala-III Project to the proposed Landscape Ecological Units (LEUs) map.

The study was organized and executed by the ICAR- National Bureau of Soil Survey and Land Use Planning, Regional Centre, Bangalore under Generation of Land Resource Inventory Data Base Component-1 of the Sujala-III Project funded by the World Bank.

GEOGRAPHICAL SETTING

2.1 Location and Extent

The Vojenahalli-2 micro-watershed is located in the central part of Karnataka in Koppal taluk and district (Fig 2.1). It lies between 15⁰21' and 15⁰22' North latitudes and 76⁰8' and 76⁰ 10' East longitudes and covers an area of about 576 ha. It comprises parts of Yathnatti, Koppal, Dadhegalla and Ojanahalli villages. It is about 4 km from Koppal town and is bounded by Ojanahalli on the north and east, Yathnatti on the west, Koppal on the south and Daddegalla on the southwestern side of the microwatershed.




Fig.2.1 Location map of Vojenahalli-2 Microwatershed

2.2 Geology

Major rock formations observed in the microwatershed are granite gneiss and alluvium (Fig.2.2 a and b). Granite gneisses are essentially pink to gray and are coarse to medium grained. They consist primarily of quartz, feldspar, biotite and hornblende. The

gray granite gneisses are highly weathered, fractured and fissured upto a depth of about 10 m. Dolerite dykes and quartz veins are common with variable width and found to occur in Vojenahalli-2 village. The thickness of the alluvium generally is limited to less than a meter, except in river valleys where it is very deep extending to tens of meters. Such soils are transported and represent paleo black soils originally formed at higher elevation, but now occupying river valleys.



Fig.2.2 a Granite and granite gneiss rocks



Fig.2.2 b Alluvium

2.3 Physiography

Physiographically, the area has been identified as Granite gneiss and Alluvial landscapes based on geology. The microwatershed area has been further divided into mounds/ridges, summits, side slopes and very gently sloping uplands and nearly level

plains based on slope and its relief features. The elevation ranges from 519 to 548 m in the gently sloping uplands. The mounds and ridges are mostly covered by rock outcrops.

2.4 Drainage

The area is drained by several small seasonal streams that join Hire *halla* and Chenna *halla* along its course. Though, the streams are not perennial, during rainy season they carry large quantities of rain water. The microwatershed has only few small tanks which are not able to store the water flowing during the rainy season. Due to this, the ground water recharge is very much affected in the villages. This is reflected in the failure of many bore wells in the villages. If the available rain water is properly harnessed by constructing tanks and recharge structures at appropriate places in the villages, then the drinking and irrigation needs of the area can be easily met. The drainage network is dendritic to sub parallel.

2.5 Climate

The district falls under semiarid tract of the state and is categorized as drought - prone with total annual rainfall of 662 mm (Table 2.1). Of this, a maximum of 424 mm precipitation is received during south–west monsoon period from June to September, north-east monsoon contributes about 161 mm and prevails from October to early December and the remaining 77 mm is received during the rest of the year. The winter season is from December to February. During April and May, the temperatures reach up to 45°C and in December and January, the temperatures will go down to 16°C. Rainfall distribution is shown in Figure 2.3. The average Potential Evapo Transpiration (PET) is 145 mm and varies from a low of 101 mm in December to 193 mm in the month of May. The PET is always higher than precipitation in all the months except in the month of September. Generally, the Length of crop Growing Period (LGP) is <90 days and starts from 2nd week of August to 2nd week of November.

Table 2.1 Mean Monthly Rainfall, PET, 1/2 PET at Koppal Taluk and District

Sl. no.	Months	Rainfall	PET	1/2 PET
1	January	1.60	116.70	58.35
2	February	1.50	129.20	64.60
3	March	14.10	169.80	84.90
4	April	18.10	180.60	90.30
5	May	41.60	193.50	96.75
6	June	85.80	167.90	83.95
7	July	72.10	156.20	78.10
8	August	110.50	152.50	76.25
9	September	155.60	138.50	69.25
10	October	116.30	122.30	61.15
11	November	36.00	106.40	53.20
12	December	9.10	101.00	50.50
TOTAL		662.30	144.55	




Fig. 2.3 Rainfall distribution in Koppal Taluk and District

2.6 Natural Vegetation

The natural vegetation is sparse comprising few tree species, shrubs and herbs. The mounds, ridges and boulders occupy sizeable areas which are under thin to moderately thick forest vegetation. Still, there are some remnants of the past forest cover which can be seen in patches in some ridges and hillocks in the microwatershed (Fig 2.4).

Apart from the continuing deforestation, the presence of large population of goats, sheep and other cattle in the microwatershed is causing vegetative degradation of whatever little vegetation left in the area. The uncontrolled grazing has left no time for the regeneration of the vegetative cover. This leads to the accelerated rate of erosion on the hill slopes, resulting in the formation of deep gullies in the foot slopes and eventually resulting in the heavy siltation of few tanks and reservoirs in the microwatershed.



Fig 2.4 Natural vegetation of Vojenahalli-2 Microwatershed

2.7 Land Utilization

About 91 per cent area (Table 2.2) in Koppal district is cultivated at present and about 17 per cent of the area is sown more than once. An area of about 3 per cent is currently barren. Forests occupy a small area of about 5 per cent and the tree cover is in a very poor state. Most of the mounds, ridges and boulder areas have very poor vegetative cover. Major crops grown in the area are sorghum, maize, bajra, cotton, safflower, sunflower, red gram, horse gram, onion, mulberry, pomegranate, sugarcane, bengalgram and groundnut (Fig 2.5 a and b). While carrying out land resource inventory, the land use/land cover particulars are collected from all the survey numbers and a current land use map of the microwatershed is prepared. The current land use map prepared shows the arable and non-arable lands, other land uses and different types of crops grown in the area. The current land use map of Vojenahalli-2 Microwatershed is presented in Fig.2.6. Simultaneously, enumeration of existing wells (bore wells) is made and their location in different survey numbers is marked on the cadastral map. Map showing the location of wells in Vojenahalli-2 Microwatershed is given in Fig 2.7.

Table 2.2 Land Utilization in Koppal District

Sl. No.	Agricultural land use	Area (ha)	Per cent
1	Total geographical area	552495	
2	Total cultivated area	500542	90.6
3	Area sown more than once	92696	16.8
4	Trees and groves	210	0.04
5	Cropping intensity	-	118
6	Forest	29451	5.33
7	Cultivable wasteland	2568	0.46
8	Permanent Pasture land	14675	2.66
9	Barren land	16627	3.01
10	Non agricultural land	40591	7.35
11	Current fallow	19660	3.56



Fig.2.5 (a) Different crops and cropping systems in Vojenahalli-2 Microwatershed




Fig.2.5 (b) Different crops and cropping systems in Vojenahalli-2 Microwatershed




Fig.2.6 Current Land Use – Vojenahalli-2 Microwatershed




Fig.2.7 Location of wells– Vojenahalli-2 Microwatershed

SURVEY METHODOLOGY

The purpose of land resource inventory is to delineate similar areas (soil series and phases), which respond or expected to respond similarly for a given level of management. This was achieved in Vojenahalli-2 microwatershed by the detailed study of all the soil characteristics (depth, texture, colour, structure, consistence, coarse fragments, porosity, soil reaction, soil horizons etc.) and site characteristics (slope, erosion, drainage, occurrence of rock fragments etc.) followed by grouping of similar areas based on soil-site characteristics into homogeneous (management units) units and showing their extent and geographic distribution on the microwatershed cadastral map. The detailed soil survey at 1:7920 scale was carried out in 576 ha area. The methodology followed for carrying out land resource inventory was as per the guidelines given in Soil Survey Manual (IARI, 1971; Soil Survey Staff, 2006; Natarajan *et al.*, 2015) which is briefly described below.

3.1 Base Maps

The detailed survey of the land resources occurring in the microwatershed was carried out by using digitized cadastral map and satellite imagery as base supplied by the KRSRSAC. The cadastral map shows field boundaries with their survey numbers, location of tanks, streams and other permanent features of the area (Fig. 3.1). Apart from the cadastral map, remote sensing data products from Cartosat-1 and LISS IV merged at the scale of 1:7920 were used in conjunction with the cadastral map to identify the geology, landscapes, landforms and other surface features. The imagery helped in the identification and delineation of boundaries between hills, uplands and lowlands, water bodies, forest and vegetated areas, roads, habitations and other cultural features of the area (Fig.3.2). The cadastral map was overlaid on the satellite imagery (Fig.3.3) that helps to identify the parcel boundaries and other permanent features. Apart from cadastral maps and images, toposheets of the area (1:50,000 scale) were used for initial traversing, identification of geology, landscapes and landforms, drainage features, present land use and also for selection of transects in the microwatershed.

3.2 Image Interpretation for Physiography

False Colour Composites (FCC) of Cartosat-I and LISS-IV merged satellite data covering the microwatershed area was visually interpreted using image interpretation elements and all the available collateral data with local knowledge. The delineated physiographic boundaries were transferred on to a cadastral map overlaid on satellite imagery. Physiographically, the area has been identified as granite gneiss and alluvial landscapes and is divided into landforms such as ridges, mounds and uplands based on slope. They were further subdivided into physiographic/ image interpretation units based on image characteristics. The image interpretation legend for physiography is given below.

Image Interpretation Legend for Physiography

G- Granite gneiss landscape

- G1 Hills/ Ridges/ Mounds
 - G11 Summits
 - G12 Side slopes
 - G121 Side slopes with dark grey tones
- G2 Uplands
 - G21 Summits
 - G22 Gently sloping uplands
 - G221 Gently sloping uplands, yellowish green (eroded)
 - G222 Gently sloping uplands, yellowish white (severely eroded)
 - G23 Very gently sloping uplands
 - G231 Very gently sloping uplands, yellowish green
 - G232 Very gently sloping uplands, medium green and pink
 - G233 Very gently sloping uplands, pink and green (scrub land)
 - G234 Very gently sloping uplands, medium greenish grey
 - G235 Very gently sloping uplands, yellowish white (eroded)
 - G236 Very gently sloping uplands, dark green
 - G237 Very gently sloping uplands, medium pink (coconut garden)
 - G238 Very gently sloping uplands, pink and bluish white (eroded)

DSe -Alluvial landscape

DSe 1 Summit

- DSe 11 Nearly level Summit with dark grey tone
- DSe 12 Nearly level Summit with medium grey tone
- DSe 13 Nearly level Summit with whitish grey tone
- DSe 14 Nearly level Summit with whitish tone (Calcareousness)
- DSe 15 Nearly level Summit with pinkish grey tone
- DSe 16 Nearly level Summit with medium pink tone
- DSe 17 Nearly level Summit with bluish white tone
- DSe 18 Nearly level Summit with greenish grey tone

DSe 2 Very genetly sloping

- DSe 21 Very gently sloping, whitish tone
- DSe 22 Very gently sloping, greyish pink tone
- DSe 23 Very gently sloping, whitish grey tone
- DSe 24 Very gently sloping, medium grey tone
- DSe 25 Very gently sloping, medium pink tone
- DSe 26 Very gently sloping, dark grey tone
- DSe 27 Very gently sloping, bluish grey tone
- DSe 28 Very gently sloping, greenish grey tone
- DSe 29 Very gently sloping, Pinkish grey




Fig 3.1 Scanned and Digitized Cadastral map of Vojenahalli-2 Microwatershed




Fig.3.2 Satellite Image of Vojenahalli-2 Microwatershed




Fig.3.3 Cadastral map overlaid on IRS PAN+LISS IV merged imagery of Vojenahalli-2 Microwatershed

3.3 Field Investigation

The field boundaries and survey numbers given on the cadastral sheet were located on the ground by following permanent features like roads, cart tracks, *nallas*, streams, tanks etc., and wherever changes were noticed, they were incorporated on the microwatershed cadastral map. Preliminary traverse of the microwatershed was carried out with the help of cadastral map, imagery and toposheets. While traversing, landforms and physiographic units identified were checked and preliminary soil legend was prepared by studying soils at few selected places. Then, intensive traversing of each physiographic unit like hills, ridges, uplands and plains was carried out. Based on the variability observed on the surface, transects (Fig 3.4) were selected across the slope covering all the landform units in the microwatershed (Natarajan and Dipak Sarkar, 2010).




Fig: 3.4. Location of profiles in a transect

In the selected transect, soil profiles (Fig.3.4) were located at closely spaced intervals to take care of any change in the land features like break in slope, erosion, gravel, stones etc. In the selected sites, profiles (vertical cut showing the soil layers from surface to the rock) were opened upto 200 cm or to the depth limited by rock or hard substratum and studied in detail for all their morphological and physical characteristics. The soil and site characteristics were recorded for all profile sites on a standard proforma as per the guidelines given in USDA Soil Survey Manual (Soil Survey Staff, 2012). Apart from the transect study, profiles were also studied at random, almost like in a grid pattern, outside the transect areas to validate the soil map unit boundaries.

Based on the soil characteristics, the soils were grouped into different soil series. Soil series is the most homogeneous unit having similar horizons and properties and behaves similarly for a given level of management. Soil depth, texture, colour, kind of horizon and horizon sequence, amount and nature of gravel present, calcareousness, nature of substratum etc, were used as the major differentiating characteristics for identifying soil series occurring in the area. The differentiating characteristics used for identifying the soil series are given in Table 3.1. Based on the above characteristics, 7 soil series were identified in Vojenahalli-2 microwatershed.

**Table 3.1 Differentiating Characteristics used for identifying Soil Series
(Characteristics are of Series Control Section)**

Soils of Granite Gneiss Landscape							
Sl. No	Soil Series	Depth (cm)	Colour (moist)	Texture	Gravel (%)	Horizon sequence	Calcareousness
1	Hooradhahalli (HDH)	75-100	2.5YR2.5/4,3/4, 3/6	gsc-gc	>35	Ap-Bt-Cr	
2	Giddadapalya (GDP)	100-150	2.5YR3/4, 3/6	gsc-gc	30-60 after 60 cm	Ap-Bt-Cr	-
Soils of Alluvial Landscape							

3	Ravanaki (RNK)	50-75	7.5YR3/2,3/3,5/2,5/3 10YR3/1,3/2,4/1, 4/2, 5/1,6/1	c	<15	Ap-Bw-Cr	e-ev
4	Dambarahalli (DRL)	75-100	10YR 2/1, 3/1, 4/3	c	<15	Ap-Bw-Ck	e-es
5	Kavalur (KVR)	100-150	10 YR 2/2, 3/1, 3/2, 3/3, 4/4	c	-	Ap-Bss-Bck-Cr	es-ev
6	Budagumpa (BGP)	>150	7.5YR3/2,5/1 10YR4/1,4/4	c	<15	Ap-Bw	es
7	Alawandi (AWD)	>150	10 YR 2/1, 3/2,	c	<15	Ap-Bss	e-es

3.4 Soil Mapping

The area under each soil series was further separated into soil phases and their boundaries delineated on the cadastral map based on the variations observed in the texture of the surface soil, slope, erosion, presence of gravel, stoniness etc. A soil phase is a subdivision of soil series based mostly on surface features that affect its use and management. The soil mapping units are shown on the map (Fig.3.5) in the form of symbols. During the survey many soil profile pits, few mini pits and a few auger bores representing different landforms occurring in the microwatershed were studied. In addition to the profile study, spot observations in the form of mini pits, road cuts, terrace cuts etc., were studied to validate the soil boundaries on the soil map.

The soil map shows the geographic distribution of 12 mapping units representing 7 soil series occurring in the microwatershed. The soil map unit (soil legend) description is presented in Table 3.2. The soil phase map (management units) shows the distribution of 12 phases mapped in the microwatershed. Each mapping unit (soil phase) delineated on the map has similar soil and site characteristics. In other words, all the farms or survey numbers included in one soil phase will have similar management needs and have to be treated accordingly.

3.5 Land Management Units

The 12 soil phases identified and mapped in the microwatershed were regrouped into three Land Management Units (LMU's) for the purpose of preparing a Proposed Crop Plan for sustained development of the microwatershed. The database (soil phases) generated under LRI was utilized for identifying Land Management Units (LMU's) based on the management needs. One or more than one soil site characteristic having influence on the management have been chosen for identification and delineation of LMUs. For Vojenahalli-2 microwatershed, five soil and site characteristics, namely the soil depth, soil texture, slope, erosion and gravel content have been considered for defining LMUs. The land management units are expected to behave similarly for a given level of management.

3.5 Laboratory Characterization

Soil samples for each soil series soil were collected from representative master profiles for laboratory characterization by following the methods outlined in the Laboratory Manual (Sarma *et al*, 1987). Surface soil samples collected in the year 2018 from farmer's fields in Vojenahalli-2 microwatershed (53 samples) for fertility status (major and micronutrients) at 320 m grid interval were analyzed in the laboratory (Katyal and Rattan, 2003). By linking the soil fertility data to the survey numbers through GIS, soil fertility maps were generated using Kriging method for the microwatershed.

Table 3.2 Soil map unit description of Vojenahalli-2 Microwatershed

Soil map unit No*	Soil Series	Soil Phase Symbol	Mapping Unit Description	Area in ha (%)
Soils of Granite and Granite gneiss Landscape				
	HDH		Hooradhahalli soils are moderately deep (75-100 cm), well drained, dark red to dark reddish brown, red gravelly sandy clay to clay soils occurring on nearly level to moderately sloping uplands under cultivation	21(3.65)
120		HDHhB1g1	Sandy clay loam surface, slope 1-3%, slight erosion, gravelly (15-35%)	20 (3.47)
125		HDHiB1	Sandy clay surface, slope 1-3%, slight erosion	1 (0.18)
	GDP		Giddadapalya soils are deep (100-150 cm), well drained, have dark reddish brown to dark red gravelly sandy clay to clay soils occurring on very gently sloping uplands under cultivation	28 (4.86)
269		GDPiB2	Sandy clay surface, slope 1-3%, moderate erosion	28 (4.86)
Soils of Alluvial Landscape				
	RNK		Ravanaki soils are moderately shallow (50-75 cm), moderately well drained, have dark brown to very dark grayish brown and dark gray, calcareous clay black soils occurring on nearly level to very gently sloping plains under cultivation	36.12(6.27)
333		RNKmB1	Clay surface, slope 1-3%, slight erosion	36 (6.25)
337		RNKmB2g1	Clay surface, slope 1-3%, moderate erosion, gravelly (15-35%)	0.12(0.02)
	DRL		Dambarahalli soils are moderately deep (75-100 cm), moderately well drained, have dark brown to very dark gray, calcareous black cracking clay soils occurring on nearly level to very gently sloping plains under cultivation	98 (17.09)
350		DRLmB2	Clay surface, slope 1-3%, moderate erosion	98 (17.09)
	KVR		Kavalur soils are deep (100-150 cm), moderately well drained, have dark yellowish brown to very dark grayish brown, calcareous cracking black clay soils occurring on nearly level to very gently sloping plains under cultivation	98.15(17.11)
385		KVRiB2g1	Sandy clay surface, slope 1-3%, moderate erosion, gravelly (15-35%)	0.15 (0.03)

Soil map unit No*	Soil Series	Soil Phase Symbol	Mapping Unit Description	Area in ha (%)
388		KVRmB1	Clay surface , slope 1-3%, slight erosion	78 (13.62)
390		KVRmB2g1	Clay surface , slope 1-3%, moderate erosion, gravelly (15-35%)	20 (3.46)
	BGP	Budagumpa soils are very deep (>150 cm), moderately well drained, have dark yellowish brown to dark brown and dark gray, sodic calcareous black clay soils occurring on nearly level to very gently sloping plains under cultivation		63(10.92)
395		BGPmA1	Clay surface , slope 0-1%, slight erosion	3 (0.46)
396		BGPmB1	Clay surface, slope 1-3%, slight erosion	60 (10.46)
	AWD	Alawandi soils are very deep (>150 cm), moderately well drained, have very dark grayish brown to black , calcareous black cracking clay soils occurring on nearly level to very gently sloping plains under cultivation		51 (8.85)
424		AWDmB2	Clay surface, slope 1-3%, moderate erosion	51 (8.85)
994		Mining/ Industrial	Mining and Industrial area	1 (0.19)
1000		Others	Habitation & waterbody	179(31.25)

*Soil map unit numbers are continuous for the taluk, not the microwatersheds




Fig 3.5 Soil Phase or Management Units- Vojenahalli-2 Microwatershed

THE SOILS

Detailed information pertaining to the nature, extent and distribution of different kinds of soils occurring in Vojenahalli-2 microwatershed is provided in this chapter. The microwatershed area has been identified as granite gneiss and alluvial landscapes based on geology. In all, 7 soil series were identified. Soil formation is the result of the combined effect of environmental and terrain factors that are reflected in soil morphology. The soil formation is dominantly influenced by the parent material, climate, time and relief.

A brief description of each of the 7 soil series identified followed by 12 soil phases (management units) mapped (Fig. 3.5) are furnished below. The physical and chemical characteristics of soil series identified in Vojenahalli-2 microwatershed are given in Table 4.1 along with soil classification. The soils in any one map unit differ from place to place in their depth, texture, slope, gravelliness, erosion or any other site characteristic that affect management. The soil phase map can be used for identifying the suitability of areas for growing specific crops or for other alternative uses and also for deciding the type of conservation structures needed. The detailed information on soil and site-characteristics like soil depth, surface soil texture, slope, erosion, gravelliness, AWC, LCC etc, with respect to each of the soil phase identified is given village/survey number wise for the microwatershed in Appendix-I.

4.1 Soils of Granite gneiss Landscape

In this landscape, 2 soil series were identified and mapped. The brief description of the soil series along with the soil phases identified and mapped is given below.

4.1.1 Hooradhahalli (HDH) Series: Hooradhahalli soils are moderately deep (75-100 cm), well drained, have red to dark red and reddish brown gravelly sandy clay to clay soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands. The Hooradhahalli series has been classified as a member of the clayey-skeletal, mixed, isohyperthermic family of Rhodic Paleustalfs.

The thickness of the solum ranges from 76 to 100 cm. The thickness of A horizon ranges from 11 to 19 cm. Its colour is in 5 YR and 2.5 YR hue with value 3 to 4 and chroma 3 to 6. The texture varies from loamy sand to sandy clay with 15 to 50 per cent gravel. The thickness of B horizon varies from 65 to 83 cm. Its colour is in 2.5 YR hue with value 2.5 to 3 and chroma 4 to 6. Texture is sandy clay to clay with 35 to 50 per cent gravel. The available water capacity is low (50-100mm/m). Two soil phases were identified and mapped.



Landscape and soil profile characteristics of Hooradhahalli (HDH) Series

4.1.2 Giddadapalya (GDP) Series: Giddadapalya soils are deep (100-150 cm), well drained, have dark reddish brown to dark red, gravelly sandy clay to clay soils. They are developed from weathered granite gneiss and occur on very gently sloping uplands under cultivation. The Giddadapalya soil series has been classified as a member of the fine, mixed, isohyperthermic family of Rhodic Paleustalfs.

The thickness of the solum ranges from 106 to 145 cm. The thickness of A-horizon ranges from 12 to 13 cm. Its colour is in 5 YR hue with value and chroma 3 to 4. The texture ranges from sandy loam with 10 to 15 per cent gravel. The thickness of B-horizon ranges from 106 to 123 cm. Its colour is in 2.5 YR hue with value 3 to 4 and chroma 3 to 6. Texture is sandy clay to clay with 35 to 75 per cent gravel. The available water capacity is low (51-100 mm/m). One soil phase was identified and mapped.



Landscape and soil profile characteristics of Giddadapalya (GDP) Series

4.2 Soils of Alluvial Landscape

In this landscape, five soil series were identified and mapped. Of these series, Dambarahalli (DRL) and Kavalur (KVR) series occupies maximum area of 98 ha (17 %) followed by Budagumpa (BGP) 106 ha (20 %) and others occupy minor area. The brief description of the soil series along with the soil phases identified and mapped is given below.

4.2.1 Ravanaki (RNK) Series: Ravanaki soils are moderately shallow (50-75 cm), well drained, have dark brown to very dark grayish brown, calcareous clay soils. They have developed from alluvium and occur on nearly level to very gently sloping plains. The Ravanaki series has been classified as a member of the very fine, smectitic, (calc) isohyperthermic family of Typic Haplustepts.

The thickness of the solum ranges from 50 to 75 cm. The thickness of A horizon ranges from 15 to 20 cm. Its colour is in 7.5 YR and 10 YR hue with value 2 to 3 and chroma 2.5 to 4. The texture varies from sandy clay to clay with 10 to 15 per cent gravel. The thickness of B horizon ranges from 35 to 60 cm. Its colour is in 10 YR and 7.5 YR hue with value 2 to 6 and chroma 2 to 4. Its texture is sandy clay to clay with gravel content of 10 to 20 per cent. The available water capacity is low (51-100 mm/m). Two soil phases were identified and mapped.



Landscape and Soil Profile Characteristics of Ravanaki (RNK) Series

4.2.2 Dambarahalli (DRL) Series: Dambarahalli soils are moderately deep (75-100 cm), moderately well drained, have black and very dark gray to dark brown, calcareous cracking clay soils. They have developed from alluvium and occur on very gently to gently sloping plains under cultivation. The Dombarahalli series has been classified as a member of the very fine, smectitic (calc), isohyperthermic family of Typic Haplusterts.

The thickness of the solum ranges from 75 to 99 cm. The thickness of A horizon ranges from 13 to 24 cm. Its colour is in 10 YR hue with value 3 to 4 and chroma 1 to 2. The texture is clay. The thickness of B horizon ranges from 54 to 85 cm. Its colour is in 10 YR hue with value 2 to 4 and chroma 1 to 3. Its texture is clay and are calcareous. The available water capacity is high (151-200 mm/m). One soil phase was identified and mapped.



Landscape and soil profile characteristics of Dambarahalli (DRL) Series

4.2.3 Kavalur (KVR): Series Kavalur soils are deep (100-150 cm), moderately well drained, have dark yellowish brown to very dark brown and very dark gray, calcareous black cracking clay soils. They have developed from alluvium and occur on very gently sloping plains. The Kavalur series has been classified as a member of the fine, smectitic (calc), isohyperthermic family of Typic Haplusterts.

The thickness of the solum is 113 to 143 cm. The thickness of A horizon ranges from 9 to 24 cm. Its colour is in 10 YR hue with value 3 and chroma 1. The texture is clay with no gravel. The thickness of B horizon ranges from 89 to 134 cm. Its colour is in 10 YR hue with value 3 and chroma 1. Its texture is clay. The available water capacity is very high (>200 mm/m). Three soil phases were identified and mapped.



Landscape and soil profile characteristics of Kavalur (KVR) series

4.2.4 Budagumpa (BGP) Series: Budagumpa soils are very deep (>150 cm), well drained, black calcareous sodic clay soils. They have developed from alluvium and occur on very gently sloping plains under cultivation. The Budagumpa series has been classified as a member of the fine, mixed, (calc) isohyperthermic family of Typic Haplustepts.


The thickness of the solum is more than 150 cm. The thickness of A horizon ranges from 16 to 26 cm. Its colour is in 7.5 YR and 10 YR hue with value 2 to 3 and chroma 2 to 4. The texture varies from sandy clay to clay with 5 to 10 per cent gravel. The thickness of B horizon ranges from 130 to 160 cm. Its colour is in 10 YR and 7.5 YR hue with value 3 to 5 and chroma 1 to 4. Its texture is clay with gravel content of <15 per cent. These soils are calcareous that increase with depth. The available water capacity is very high (>200 mm/m). Two soil phases were identified and mapped.



Landscape and Soil Profile Characteristics of Budagumpa (BGP) Series

4.2.5 Alawandi (AWD) Series: Alawandi soils are very deep (>150 cm), moderately well drained, have black to very dark grayish brown, calcareous cracking clay soils. They have developed from alluvium and occur on nearly level to very gently sloping plains under cultivation. The Alawandi series has been classified as a member of the fine smectitic (calc), isohyperthermic family of Typic Haplusterts .

The thickness of the solum is more than 150 cm. The thickness of A horizon ranges from 16 to 26 cm. Its colour is in 10 YR hue with value 2 to 3 and chroma 1 to 2. The texture varies from sandy clay to clay. The thickness of B horizon is more than 150 cm. Its colour is in 10 YR hue with value 2 to 3 and chroma 1 to 3. Its texture is clay and is calcareous. The available water capacity is very high (>200 mm/m). One soil phase was identified and mapped.



Landscape and Soil Profile Characteristics of Alawandi (AWD) Series

Table: 4.1 Physical and Chemical Characteristics of Soil Series identified in Vojenahalli-2 microwatershed

Soil Series: Hooradhahalli (HDH), **Pedon:** RM-69

Location: 13°24'31"N, 76°33'41"E, (4D3D8G2d), Hesarahalli village, Chikkanayakanahalli taluk, Tumukura district

Analysis at: NBSS&LUP, Regional Centre, Bengaluru **Classification:** Clayey-skeletal, mixed, isohyperthermic Rhodic Paleustalfs

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-18	Ap	72.56	15.17	12.27	4.57	8.33	17.38	23.88	18.39	35	sl	-	-
18-33	Bt1	56.29	10.75	32.96	7.88	10.24	13.41	14.43	10.34	55	scl	-	-
33-58	Bt2	46.66	10.79	42.55	10.79	9.87	8.43	9.04	8.53	55	sc	-	-
58-90	Bt3	43.09	13.63	43.27	9.90	8.25	7.32	8.76	8.87	45	c	-	-

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO ₃	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP			
	Water	CaCl ₂	M KCl				dS m ⁻¹	%	%	Ca	Mg					K	Na	Total
										cmol kg ⁻¹								
0-18	6.54	-	-	0.07	0.60	0.00	2.68	1.38	0.44	0.42	4.91	5.84	0.48	84.07	7.11			
18-33	5.90	-	-	0.07	0.52	0.00	3.99	1.27	0.09	0.37	5.71	8.61	0.26	66.32	4.29			
33-58	6.16	-	-	0.07	0.44	0.00	4.92	1.67	0.08	0.55	7.22	10.00	0.24	72.23	5.50			
58-90	6.39	-	-	0.06	0.40	0.00	4.30	2.02	0.08	0.46	6.87	9.21	0.21	74.61	5.05			

Contd...

Series Name: Giddadapalya (GDP), **Pedon:** R-8

Location: 15°25'26"N, 76°10'59"E, Kalakeri village, Koppal taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bangalore. Classification: Fine, mixed, isohyperthermic Rhodic Paleustalfs

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-16	Ap	74.95	9.24	15.81	18.43	18.94	13.85	14.97	8.76	-	sl	11.88	5.09
16-43	Bt1	41.69	13.89	44.42	9.84	10.90	7.41	7.62	5.93	-	c	23.13	14.53
43-61	Bt2	47.67	6.13	46.19	21.14	10.15	5.29	6.45	4.65	-	sc	21.60	11.87
61-83	Bt3	52.52	7.10	40.38	24.42	10.59	5.66	7.55	4.30	40	sc	19.51	11.35
83-119	Bt4	43.76	11.59	44.65	20.15	7.56	5.77	5.46	4.83	60	c	20.80	12.06
119-139	Bt5	54.93	9.84	35.23	29.70	10.49	5.50	5.92	3.32	50	sc	15.24	11.97

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO ₃	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP			
	Water	CaCl ₂	M KCl				dS m ⁻¹	%	%	Ca	Mg					K	Na	Total
										cmol kg ⁻¹								
0-16	7.88	-	-	0.103	0.79	-	5.98	1.35	0.05	0.22	7.60	7.8	0.49	97	2.87			
16-43	7.81	-	-	0.117	0.66	-	13.99	1.97	0.08	0.46	16.50	16.9	0.38	98	2.74			
43-61	7.74	-	-	0.132	0.51	-	12.70	2.18	0.08	0.69	15.64	15.9	0.34	98	4.36			
61-83	7.72	-	-	0.142	0.39	-	11.46	2.22	0.08	0.66	14.41	14.6	0.36	99	4.53			
83-119	7.58	-	-	0.115	0.22	-	11.30	2.70	0.09	0.73	14.82	15.3	0.34	97	4.79			
119-139	7.50	-	-	0.113	0.22	-	10.03	2.19	0.07	0.65	12.95	13.2	0.37	98	4.89			

Contd...

Series Name: Ravanaki (RNK), **Pedon:** RM-20

Location: 15°14'22.7"N, 75°57'45.8"E, Gatareddihalla village, Koppal taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bangalore. **Classification:** Very fine, smectitic (calc), isohyperthermic Typic Haplustepts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-28	Ap	24.43	17.76	57.81	5.30	3.89	3.78	7.14	4.32	20	c	41.40	29.60
28-55	Bw	18.77	15.59	65.64	2.74	3.73	2.85	4.83	4.61	10	c	46.71	35.18

Depth (cm)	pH (1:2.5)			E.C. (1:2.5) dS m ⁻¹	O.C. %	CaCO ₃ %	Exchangeable bases					CEC	CEC/Clay	Base saturation %	ESP %
	Water	CaCl ₂	M KCl				Ca	Mg	K	Na	Total				
0-28	8.86	-	-	0.483	0.63	15.48	-	-	0.86	6.27	-	37.00	0.64	-	6.78
28-55	8.61	-	-	1.4	0.23	13.68	-	-	0.68	12.27	-	53.20	0.81	-	9.22

Contd...

Series Name: Dombarahalli (DRL)

Pedon: R-8

Location: 15°13'96.2"N, 75°57'48.6" E Rangunathanahalli village, Koppal taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bangalore.

Classification: Very fine, smectitic (calc), isohyperthermic Typic Haplusterts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-15	Ap	28.25	19.48	52.27	4.76	4.44	4.87	8.23	5.95	-	c	39.86	27.20
15-27	BA1	21.55	20.00	58.45	3.76	2.76	3.43	6.30	5.30	-	c	46.35	34.84
27-45	Bss1	14.86	20.89	64.25	2.46	2.23	2.23	3.91	4.02	-	c	57.99	41.06
45-80	Bss2	10.42	19.04	70.54	1.74	1.97	1.27	2.78	2.66	-	c	66.36	36.24

Depth (cm)	pH (1:2.5)			E.C. (1:2.5) dS m ⁻¹	O.C. %	CaCO ₃ %	Exchangeable bases					CEC	CEC/Clay	Base saturation %	ESP %
	Water	CaCl ₂	M KCl				Ca	Mg	K	Na	Total				
0-15	8.78	-	-	0.42	0.32	12.35	-	-	0.59	4.25	-	49.70	0.95	100.00	5.62
15-27	9.03	-	-	0.61	0.30	12.48	-	-	0.30	8.96	-	57.23	0.98	100.00	10.07
27-45	9.10	-	-	0.67	0.34	11.70	-	-	0.25	11.85	-	60.71	0.95	100.00	14.05
45-80	9.18	-	-	0.86	0.32	13.39	-	-	0.27	15.40	-	63.33	0.90	100.00	18.45

Contd...

Series Name: Kavalura (KVR), **Pedon:** A2/RM-9

Location: 15°18'86.8"N, 75°56'56.3"E, Kavalura village, Koppal taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bangalore. **Classification:** Fine, smectitic (calc), isohyperthermic Typic Haplusterts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-24	Ap	36.18	17.80	46.02	7.04	7.47	6.62	9.28	5.76	10	c	28.20	18.75
24-50	Bss1	38.79	15.36	45.85	6.25	6.25	9.70	10.67	5.93	05	c	27.16	18.81
50-85	Bss2	36.80	14.66	48.54	9.63	8.23	7.03	7.58	4.33	<5	c	30.16	22.17
85-124	Bss3	22.66	17.24	60.09	4.18	3.85	5.28	5.06	4.29	<5	c	40.34	31.42

Depth (cm)	pH (1:2.5)			E.C. (1:2.5) dS m ⁻¹	O.C. %	CaCO ₃ %	Exchangeable bases					CEC	CEC/Clay	Base saturation %	ESP %
	Water	CaCl ₂	M KCl				Ca	Mg	K	Na	Total				
0-24	8.4	-	-	0.265	0.2	8.04	-	-	0.97	0.65		43.25	0.94		0.60
24-50	9.27	-	-	0.23	0.37	8.04	-	-	0.31	3.21		41.66	0.91		3.08
50-85	9.44	-	-	0.297	0.41	8.64	-	-	0.35	6.43		43.99	0.91		5.85
85-124	9.37	-	-	0.46	0.41	11.40	-	-	0.42	7.99		51.09	0.85		6.26

Contd...

Series Name: Budagumpa (BGP), **Pedon:** R-21

Location: 15°23'45"N, 76°08'52"E Neregalla village, Koppal taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bangalore. **Classification:** Fine, mixed (calc), isohyperthermic Typic Haplustepts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-16	Ap	58.30	18.10	23.60	6.34	11.75	11.66	17.44	11.10	-	scl	18.24	10.29
16-38	Bw1	44.26	18.39	37.36	4.71	9.79	9.32	12.24	8.19	-	cl	32.99	18.12
38-68	Bw2	37.84	24.91	37.25	3.66	7.51	8.45	10.89	7.32	-	cl	39.50	22.32
68-83	Bw3	19.17	19.89	60.93	0.87	3.47	3.85	6.07	4.91	-	c	47.27	28.52
83-107	Bw4	14.76	23.22	62.02	0.63	2.41	3.25	4.61	3.87	-	c	46.10	29.36
107-131	Bw5	11.86	17.75	70.39	0.85	2.73	2.45	3.20	2.64	-	c	50.52	28.09
131-160	Bw6	14.48	18.21	67.31	2.23	2.50	2.59	3.84	3.31	-	c	59.14	28.35

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO ₃	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP			
	Water	CaCl ₂	M KCl				dS m ⁻¹	%	%	Ca	Mg					K	Na	Total
0-16	9.20	-	-	0.27	0.51	6.24	-	-	0.42	3.11	-	19.60	0.83	100.00	3.84			
16-38	9.29	-	-	0.88	0.35	5.98	-	-	0.17	9.36	-	28.40	0.76	100.00	15.38			
38-68	8.95	-	-	2.37	0.31	4.81	-	-	0.31	24.10	-	34.90	0.94	100.00	42.65			
68-83	8.65	-	-	4.28	0.33	4.42	-	-	0.39	27.95	-	45.10	0.74	100.00	25.94			
83-107	8.10	-	-	9.50	0.30	3.38	-	-	0.44	31.29	-	44.10	0.71	100.00	12.82			
107-131	8.16	-	-	9.32	0.22	2.73	-	-	0.63	37.86	-	47.20	0.67	100.00	20.37			
131-160	8.49	-	-	5.29	0.19	3.51	-	-	0.60	34.82	-	43.70	0.65	100.00	48.66			

Contd...

Series Name: Alawandi (AWD), **Pedon:** R-16

Location : 15°13'08.2"N, 76°15'27.3" E Neeralagi village, Koppal taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bangalore. **Classification:** Fine, smectitic (calc), isohyperthermic Typic Haplusterts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-17	Ap	20.88	25.75	53.37	3.31	4.31	4.31	5.19	3.76	-	c	33.11	25.58
17-39	Bss1	25.99	19.79	54.22	5.04	5.48	5.04	5.92	4.50	-	c	33.11	26.23
39-70	Bss2	26.76	17.80	55.44	2.93	5.31	5.53	7.37	5.63	-	c	36.15	28.67
70-111	Bss3	23.83	20.25	55.93	4.15	4.81	4.92	6.01	3.93	-	c	43.60	33.71
111-139	Bss4	21.21	20.40	58.40	2.79	4.80	4.91	5.25	3.46	-	c	46.92	36.28
139-162	Bss5	13.15	20.96	65.90	1.69	2.47	2.36	3.37	3.26	-	c	54.96	41.81

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO ₃	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP			
	Water	CaCl ₂	M KCl				dS m ⁻¹	%	%	Ca	Mg					K	Na	Total
										cmol kg ⁻¹								
0-17	8.10	-	-	0.37	0.52	9.48	-	-	0.40	1.56	-	51.30	0.96	100.00	1.22			
17-39	8.60	-	-	0.24	0.52	9.60	-	-	0.14	4.60	-	52.60	0.97	100.00	3.50			
39-70	8.89	-	-	0.27	0.52	9.48	-	-	0.16	2.41	-	53.90	0.97	100.00	1.78			
70-111	9.10	-	-	0.35	0.54	11.28	-	-	0.15	8.95	-	54.10	0.97	100.00	6.61			
111-139	9.15	-	-	0.41	0.58	10.80	-	-	0.15	7.36	-	56.10	0.96	100.00	5.24			
139-162	9.16	-	-	0.50	0.50	15.48	-	-	0.19	10.19	-	61.66	0.94	100.00	6.61			

INTERPRETATION FOR LAND RESOURCE MANAGEMENT

The most important soil and site characteristics that affect the land use and conservation needs of an area are land capability, land irrigability, soil depth, soil texture, coarse fragments, available water capacity, soil slope, soil erosion, soil reaction etc. These are interpreted from the data base generated through land resource inventory and several thematic maps are generated. These would help in identifying the areas suitable for growing crops and, soil and water conservation measures and structures needed thus helping to maintain good soil health for sustained crop production. The various thematic maps generated are described below.

5.1 Land Capability Classification

Land capability classification is an interpretative grouping of soil map units (soil phases) mainly based on inherent soil characteristics, external land features and environmental factors that limit the use of land for agriculture, pasture, forestry, or other uses on a sustained basis (IARI, 1971). The land and soil characteristics used to group the land resources in an area into various land capability classes, subclasses and units are

Soil characteristics: Soil depth, soil texture, coarse fragments, soil reaction, available water capacity, calcareousness, salinity/alkali *etc.*

Land characteristics: Slope, erosion, drainage, rock outcrops.

Climate: Total rainfall and its distribution, and length of crop growing period.

The Land Capability Classification system is divided into land capability classes, subclasses and units based on the level of information available. Eight land capability classes are recognized. They are

Class I: They are very good lands that have no limitations or very few limitations that restrict their use.

Class II: They are good lands that have minor limitations and require moderate conservation practices.

Class III: They are moderately good lands that have severe limitations that reduce the choice of crops or that require special conservation practices.

Class IV: They are fairly good lands that have very severe limitations that reduce the choice of crops or that require very careful management.

Class V: Soils in these lands are not likely to erode, but have other limitations like wetness that are impractical to remove and as such not suitable for agriculture, but suitable for pasture or forestry with minor limitations.

Class VI: The lands have severe limitations that make them generally unsuitable for cultivation, but suitable for pasture or forestry with moderate limitations.

Class VII: The lands have very severe limitations that make them unsuitable for cultivation, but suitable for pasture or forestry with major limitations.

Class VIII: Soil and other miscellaneous areas (rock lands) that have very severe limitations that nearly preclude their use for any crop production, but suitable for wildlife, recreation and installation of wind mills.

The land capability subclasses are recognized based on the dominant limitations observed within a given land capability class. The subclasses are designated by adding a lower case letter like ‘e’, ‘w’, ‘s’, or ‘c’ to the class numeral. The subclass “e” indicates that the main hazard is risk of erosion, “w” indicates drainage or wetness as a limitation for plant growth, “s” indicates shallow soil depth, coarse or heavy textures, calcareousness, salinity/alkalinity or gravelliness and “c” indicates limitation due to climate.

The land capability subclasses have been further subdivided into land capability units based on the kinds of limitations present in each subclass. Ten land capability units are used in grouping the soil map units. They are stony or rocky (0), erosion hazard (slope, erosion) (1), coarse texture (sand, loamy sand, sandy loam) (2), fine texture (cracking clay, silty clay) (3), slowly permeable subsoil (4), coarse underlying material (5), salinity/alkali (6), stagnation, overflow, high ground water table (7), soil depth (8) and fertility problems (9). The capability units thus identified have similar soil and land characteristics that respond similarly to a given level of management. The soils of the microwatershed have been classified upto land capability subclass level.

The 12 soil map units identified in the Vojanahalli-2 Microwatershed are grouped under two land capability classes and four land capability subclasses (Fig. 5.1).




Fig. 5.1 Land Capability map of Vojanahalli-2 Microwatershed

Entire cultivated area in the microwatershed is suitable for agriculture. Good lands (Class II) cover an area of about 297 ha (52%) and distributed in the major part of the microwatershed with minor problems of soil and erosion. Moderately good lands (Class III) occupy an area of about 99 ha (17%) and distributed in the southeastern and northwestern part of the microwatershed with severe limitations of soil and erosion. An area of about 179 ha (31%) is covered by habitation and water body and 1 ha (<1%) by mining/Industrial area.

5.2 Soil Depth

Soil depth refers to the depth of the soil occurring above the parent material or hard rock. The depth of the soil determines the effective rooting depth for plants and in accordance with soil texture, mineralogy and gravel content, the capacity of the soil column to hold water and nutrient availability. Soil depth is one of the most important soil characteristic that is used in differentiating soils into different soil series. The soil depth classes used in identifying soils in the field are very shallow (<25 cm), shallow (25-50 cm), moderately shallow (50-75 cm), moderately deep (75-100 cm), deep (100-150 cm) and very deep (>150 cm). They were used to classify the soils into different depth classes and a soil depth map was generated (Fig. 5.2). The area extent and their geographical distribution in the microwatershed is given in Fig. 5.2.




Fig. 5.2 Soil Depth map of Vojenahalli-2 Microwatershed

Moderately shallow (50-75 cm) soils cover an area of about 36 ha (6%) and distributed in the northern part of the microwatershed. An area of about 119 ha (21%) is

moderately deep soils (75-100 cm) and distributed in the northern and central part of the microwatershed. Deep to very deep (100- >150 cm) soils occupy a maximum area of about 240 ha (42%) and distributed in the major part of the microwatershed.

The most productive lands cover about 240 ha (42%) where all climatically adopted long duration crops be grown.

5.3 Surface Soil Texture

Texture is an expression to indicate the coarseness or fineness of the soil as determined by the relative proportion of primary particles of sand, silt and clay. It has a direct bearing on the structure, porosity, adhesion and consistence. The surface layer of a soil to a depth of about 25 cm is the layer that is most used by crops and plants. The surface soil textural class provides a guide to understanding soil-water retention and availability, nutrient holding capacity, infiltration, workability, drainage, physical and chemical behavior, microbial activity and crop suitability. The textural classes used for LRI were used to classify and a surface soil texture map was generated. The area extent and their geographical distribution in the microwatershed is shown in Fig 5.3.

An area of about 20 ha (3%) is loamy (sandy clay loam) at the surface and distributed in the central part of the microwatershed. Clayey (sandy clay and clay) soils cover about 376 ha (65%) and are distributed in the major part of the microwatershed.

The most productive lands with respect to surface soil texture are clayey soils that (65%) have high potential for soil-water retention and availability and nutrient retention and availability, but have more problems of drainage, infiltration, workability and other physical problems. The other productive lands are loamy (3%) soils which also have high potential for soil- water retention and nutrient availability but have no drainage or other physical problems.




Fig. 5.3 Surface Soil Texture map of Vojenahalli-2 Microwatershed

5.4 Soil Gravelliness

Gravel is the term used for describing coarse fragments between 2 mm and 7.5 cm diameter and stones for those between 7.5 cm and 25 cm. The presence of gravel and stones in soil reduces the volume of soil responsible for moisture and nutrient storage, drainage, infiltration and runoff, and hinders plant growth by impeding root growth and seedling emergence, intercultural operations and farm mechanization. The gravelliness classes used in LRI were used to classify the soils and using these classes a gravelliness map was generated. The area extent and their geographic distribution in the microwatershed is shown in Fig. 5.4.

The soils that are non-gravelly (<15% gravel) cover a maximum area of about 356 ha (62 %) and distributed in the major part of the microwatershed. An area of about 40 ha (7 %) is covered by gravelly (15-35% gravel) soils and are distributed in the western and central part of the microwatershed (Fig. 5.4).

The most productive lands with respect to gravelliness are found to be 62 per cent. They are non-gravelly with less than 15 per cent gravel and have potential for growing both annual and perennial crops.




Fig. 5.4 Soil Gravelliness map of Vojenahalli-2 Microwatershed

5.5 Available Water Capacity

The soil available water capacity (AWC) is estimated based on the ability of the soil column to retain water between the tensions of 0.33 and 15 bar in a depth of 100 cm or the entire solum if the soil is shallower. The AWC of the soils (soil series) as estimated by considering the soil texture, mineralogy, soil depth and gravel content (Sehgal *et al.*, 1990) and accordingly the soil map units were grouped into five AWC classes *viz.*, very low (<50 mm/m), low (50-100 mm/m), medium (100-150 mm/m), high (150-200 mm/m) and very high (>200 mm/m) and using these values, an AWC map was generated. The area extent and their geographic distribution of different AWC classes in the microwatershed is shown in Fig. 5.5.

An area of about 21 ha (4%) in the microwatershed has soils that are very low (<50 mm/m) in available water capacity and are distributed in the central part of the microwatershed. An area of about 64 ha (11%) has soils that are low (51 to 100 mm/m) in available water capacity and are distributed in the northern and southeastern part of the microwatershed. An area of about 98 ha (17%) has soils that are medium (101-150 mm/m) in available water capacity and are distributed in the central part of the microwatershed. An area of about 212 ha (37%) is high to very high (151->200 mm/min) in available water capacity and distributed in the major part of the microwatershed.

An area of about 85 ha (15%) in the microwatershed has soils that are problematic with regard to available water capacity. Here, only short duration crops can be grown and the probability of crop failure is very high. These areas are best put to other alternative

uses. An area of about 212 ha (37%) has soils that have high potential (>200 mm/m) with regard to available water capacity where all climatically adapted long duration crops can be grown successfully.




Fig. 5.5 Soil Available Water Capacity map of Vojenahalli-2 Microwatershed

5.6 Soil Slope

Soil slope refers to the inclination of the surface of the land. It is defined by gradient, shape and length, and is an integral feature of any soil as a natural body. Slope is considered important in soil genesis, land use and land development. The length and gradient of slope influences the rate of runoff, infiltration, erosion and deposition. The soil map units were grouped into two slope classes and a slope map was generated showing the area extent and their geographic distribution of different slope classes in the microwatershed (Fig. 5.6).

Nearly level (0-1%) lands cover an area of about 3 ha (<1%) and distributed in the southwestern part of the microwatershed. Very gently sloping (1-3%) lands cover a maximum area of about 393 ha (68%) and distributed in the major part of the microwatershed. In all these areas, all climatically adapted annual and perennial crops can be grown without much soil and water conservation and other land development measures.




Fig. 5.6 Soil Slope map of Vojenahalli-2 Microwatershed

5.7 Soil Erosion

Soil erosion refers to the wearing away of the earth's surface by the forces of water, wind and ice involving detachment and transport of soil by raindrop impact. It is used for accelerated soil erosion resulting from disturbance of the natural landscape by burning, excessive grazing and indiscriminate felling of forest trees and tillage, all usually by man. The erosion classes showing an estimate of the current erosion status as judged from field observations in the form of rills, gullies or a carpet of gravel on the surface are recorded. Four erosion classes, viz, slight erosion (e1), moderate erosion (e2), severe erosion (e3) and very severe erosion (e4) are recognized. The soil map units were grouped into different erosion classes and a soil erosion map generated. The area extent and their spatial distribution in the microwatershed is given in Figure 5.7.

Slightly eroded lands cover an area of about 198 ha (34 %) and distributed in the eastern, western and northern part of the microwatershed. An area of about 198 ha (34 %) is moderately eroded (e2 class) and distributed in the northern and central part of the microwatershed. Moderately eroded lands are problematic and need appropriate soil and water conservation and other land development measures.

FERTILITY STATUS

Soil fertility plays an important role in increasing crop yield. The adoption of high yielding varieties that require high amounts of nutrients has resulted in deficiency symptoms in crops and plants due to imbalanced fertilization and poor inherent fertility status, as these areas are characterized by low rainfall and high temperatures. Hence, it is necessary to know the fertility (macro and micro nutrients) status of the soils of the watersheds for assessing the kind and amount of fertilizers required for each of the crop intended to be grown. For this purpose, the surface soil samples collected from the grid points (one soil sample at every 320 m grid interval) all over the microwatershed through land resource inventory in the year 2018 were analyzed for pH, EC, organic carbon, available phosphorus and potassium, and for micronutrients like zinc, boron, copper, iron and manganese, and secondary nutrient sulphur.

Soil fertility data generated has been assessed and individual maps for all the nutrients for the microwatershed have been generated by using the Kriging method under GIS. The village/survey number wise fertility data for the microwatershed is given in Appendix-II.

6.1 Soil Reaction (pH)

The soil analysis of the Vojenahalli-2 microwatershed for soil reaction (pH) showed that slightly to moderately alkaline (pH 7.3-8.4) soils cover an area of about 94 ha (16%) and distributed in the southeastern and southwestern part of the microwatershed. Maximum area of about 302 ha (52%) is strongly to very strongly alkaline (pH 8.4->9.0) and is distributed in the major part of the microwatershed. (Fig.6.1). Thus, entire area in the microwatershed is alkaline in reaction.

6.2 Electrical Conductivity (EC)

The Electrical Conductivity of the soils of the entire microwatershed area is $<2 \text{ dSm}^{-1}$ (Fig 6.2) and as such the soils are non-saline.

6.3 Organic Carbon

An area of about 105 ha (18%) is low ($<0.5\%$) in OC and distributed in the northern and eastern part of the microwatershed. Maximum area of about 291 ha (51%) is medium (0.5-0.75%) and distributed in the major part of the microwatershed (Fig.6.3).

6.4 Available Phosphorus

Entire area in the microwatershed is medium (23-57 kg/ha) in available phosphorus. Apply additional 25% phosphorus in areas where it is medium (Fig 6.4).




Fig.6.1 Soil Reaction (pH) map of Vojenahalli-2 Microwatershed




Fig.6.2 Electrical Conductivity (EC) map of Vojenahalli-2 Microwatershed




Fig.6.3 Soil Organic Carbon map of Vojenahalli-2 Microwatershed

6.5 Available Potassium

Available potassium is medium (145-337 kg/ha) in 351 ha (61%) and distributed in the major part of the microwatershed. An area of about 45 ha (8%) is high (>337 kg/ha) and distributed in the southwestern and eastern part of the microwatershed. The areas with high potassium content reduce 25 per cent from the recommended dose to avoid the excess application of fertilizer. Apply additional 25% potassium in areas where it is medium (Fig 6.5).

6.6 Available Sulphur

Soil analysis of available sulphur content in Vojenahalli-2 microwatershed showed that an area of about 200 ha (35%) is low and distributed in the major part of the microwatershed. An area of about 127 ha (22%) is medium (10-20 ppm) in available sulphur content and distributed in the northern and eastern part of the microwatershed. An area of about 69 ha (12%) is high (>20 ppm) and distributed in the northwestern and western part of the microwatershed (Fig.6.6). The areas that are low and medium in available sulphur need to be applied with magnesium sulphate or gypsum or factomphos (p) fertilizer (13% sulphur) for 2-3 years for the deficiency to be corrected.

6.7 Available Boron

An area of about 58 ha (10%) is low (<0.5ppm) in available boron and distributed in the eastern part of the microwatershed. Maximum area of about 331 ha (57%) is

medium (0.5-1.0 ppm) and distributed in the major part of the microwatershed (Fig.6.7). An area of about 8 ha (1%) is high (>1.0 ppm) and distributed in the northwestern part of the microwatershed.

6.8 Available Iron

Available iron content in the soils of the Vojenahalli-2 microwatershed is deficient (<4.5 ppm) in a maximum area of about 396 ha (69%) and distributed in the major part of the microwatershed. An area of about < 1 ha (<1 %) showed sufficiency (>4.5 ppm) with respect to iron content and distributed in the eastern part of the microwatershed (Fig 6.8).

6.9 Available Manganese

Available manganese content is sufficient (>1.0 ppm) in the entire microwatershed area (Fig 6.9).

6.10 Available Copper

Available copper content is sufficient (>0.2 ppm) in the entire microwatershed area (Fig 6.10).




Fig.6.4 Soil Available Phosphorus map of Vojenahalli-2 Microwatershed




Fig.6.5 Soil Available Potassium map of Vojenahalli-2 Microwatershed




Fig.6.6 Soil Available Sulphur map of Vojenahalli-2 Microwatershed




Fig.6.7 Soil Available Boron map of Vojenahalli-2 Microwatershed




Fig.6.8 Soil Available Iron map of Vojenahalli-2 Microwatershed




Fig.6.9 Soil Available Manganese map of Vojenahalli-2 Microwatershed




Fig.6.10 Soil Available Copper map of Vojenahalli-2 Microwatershed

6.11 Available Zinc

Available zinc content is deficient (<0.6 ppm) in an area of about 199 ha (35 %) and distributed in the northern and central part of the microwatershed. An area of about 197 ha (34%) is sufficient (>0.6 ppm) and distributed in the eastern and central part of the microwatershed (Fig 6.11).




Fig.6.11 Soil Available Zinc map of Vojenahalli-2 Microwatershed

LAND SUITABILITY FOR MAJOR CROPS

The soil and land resource units (soil phases) of Vojenahalli-2 Microwatershed were assessed for their suitability for growing food, fodder, fibre and other horticulture crops by following the procedure as outlined in FAO, 1976 and 1983. Crop requirements were developed for each of the crop from the available research data and also by referring to Naidu *et. al.* (2006) and Natarajan *et. al.* (2015). The soil and land characteristics were matched with the crop requirements to arrive at the crop suitability. The soil and land characteristics table (Table 7.1) were matched with the crop requirements (Tables 7.2-7.32) to arrive at the crop suitability and the crop requirement tables are given at the end of the chapter. In FAO land suitability classification, two orders are recognized. Order S- Suitable and Order N- Not suitable. The orders have classes, subclasses and units. Order S has three classes, Class S1- Highly Suitable, Class S2- Moderately Suitable and Class S3- Marginally Suitable. Order N has two Classes, N1- Currently not Suitable and N2- Permanently not Suitable. There are no subclasses within the Class S1 as they will have very minor or no limitations for crop growth. Classes S2, S3 and N1 are divided into subclasses based on the kinds of limitations encountered. The limitations that affect crop production are 'c' for erratic rainfall and its distribution and length of growing period (LGP), 'e' for erosion hazard, 'r' for rooting condition, 't' for lighter or heavy texture, 'g' for gravelliness or stoniness, 'n' for nutrient availability, 'l' for topography, 'm' for moisture availability, 's' for sodium 'z' for calcareousness and 'w' for drainage. These limitations are indicated as lower case letters to the class symbol. For example, moderately suitable lands with the limitations of soil depth and erosion are designated as S2re. For the microwatershed, the soil mapping units were evaluated and classified up to subclass level.

Using the above criteria, the soil map units of the microwatershed were evaluated and land suitability maps for 31 major agricultural and horticultural crops were generated. The detailed information on the kind of suitability of each of the soil phase for the crops assessed are given village/ survey number wise for the microwatershed in Appendix-III.

7.1 Land Suitability for Sorghum (*Sorghum bicolor*)

Sorghum is one of the major food crop grown in Karnataka in an area of 10.47 lakh ha in Bijapur, Gulbarga, Raichur, Bidar, Belgaum, Dharwad, Bellary, Chitradurga, Mysore and Chamarajnagar districts. The crop requirements for growing sorghum (Table 7.2) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing sorghum was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.1.

Highly suitable (Class S1) lands occupy an area of about 28 (5%) for growing sorghum and occur in the southern part of the microwatershed. Maximum area of about

346 ha (60%) is moderately suitable (Class S2) for growing sorghum and distributed in the major part of the microwatershed with minor limitations of gravelliness, rooting depth, nutrient availability and calcareousness. An area of about 21 ha (4%) is marginally suitable for growing sorghum and distributed in the central part of the microwatershed. They have moderate limitation of gravelliness.




Fig. 7.1 Land Suitability map of Sorghum

7.2 Land Suitability for Maize (*Zea mays*)

Maize is one of the most important food crop grown in an area of 13.37 lakh ha in almost all the districts of the State. The crop requirements for growing maize (Table 7.3) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing maize was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.2.

An area of about 28 ha (5 %) is highly suitable (Class S1) and distributed in the southern part of the microwatershed. Moderately suitable (Class S2) lands cover an area of about 347 ha(60%) and distributed in the major part of the microwatershed with minor limitations of texture and calcareousness. Marginally suitable (Class S3) lands cover an area of about 21 ha (4%) and occur in the central part of the microwatershed. They have moderate limitation of gravelliness.




Fig. 7.2 Land Suitability map of Maize

7.3 Land Suitability for Bajra (*Pennisetum glaucum*)

Bajra is one of the major food crop grown in an area of 2.34 lakh ha in Karnataka in the northern districts. The crop requirements (Table 7.4) for growing bajra were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing bajra was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.3.

Highly suitable (Class S1) lands occupy an area of about 28 ha (5 %) for growing bajra and occur in the southern part of the microwatershed. Maximum area of about 368 ha (64%) is moderately suitable (Class S2) for growing bajra and distributed in the major part of the microwatershed with minor limitations of texture, calcareousness and gravelliness.




Fig. 7.3 Land Suitability map of Bajra

7.4 Land Suitability for Redgram (*Cajanus cajan*)

Redgram is one of the most important pulse crop grown in an area of 7.28 lakh ha in almost all the districts of the State. The crop requirements for growing redgram (Table 7.5) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing redgram was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.4.

Highly suitable (Class S1) lands occupy an area of about 28 ha (5 %) for growing redgram and occur in the southern part of the microwatershed. An area of about 212 ha (37%) is moderately suitable (Class S2) for growing redgram and distributed in the major part of the microwatershed. They have minor limitations of gravelliness, calcareousness and texture. Marginally suitable lands (Class S3) occupy an area of about 156 ha (27%) and occur in the northern and central part of the microwatershed. They have moderate limitations of rooting depth, gravelliness and calcareousness.




Fig. 7.4 Land Suitability map of Redgram

7.5 Land Suitability for Bengal gram (*Cicer arietinum*)

Bengal gram is one of the major pulse crop grown in an area of 9.39 lakh ha in northern Karnataka in Bijapur, Gulbarga, Raichur, Bidar, Belgaum, Dharwad and Bellary districts. The crop requirements for growing Bengal gram (Table 7.6) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing Bengal gram was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.5.

Maximum area of about 375 ha (65%) is moderately suitable (Class S2) for growing bengalgram and are distributed in the major part of the microwatershed. They have minor limitations of texture, calcareousness, rooting depth and gravelliness. Marginally suitable (Class S3) lands cover an area of about 21 ha (4%) and are distributed in the southern part of the microwatershed. They have moderate limitation of gravelliness.




Fig. 7.5 Land Suitability map of Bengal gram

7.6 Land Suitability for Groundnut (*Arachis hypogaea*)

Groundnut is one of the major oilseed crop grown in an area of 6.54 lakh ha in Karnataka in most of the districts either as rainfed or irrigated crop. The crop requirements for growing groundnut (Table 7.7) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing groundnut was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.6.

An area of about 49 ha (9%) is moderately suitable (Class S2) for growing groundnut and distributed in the southern part of the microwatershed. They have minor limitations of gravelliness and texture. Maximum area of about 347 ha (60%) is marginally suitable (Class S3) for growing groundnut and are distributed in the major part of the microwatershed with moderate limitations of texture and calcareousness.




Fig. 7.6 Land Suitability map of Groundnut

7.7 Land Suitability for Sunflower (*Helianthus annuus*)

Sunflower is one of the most important oilseed crop grown in an area of 3.56 lakh ha in the State in all the districts. The crop requirements for growing sunflower (Table 7.8) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing sunflower was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.7.

An area of about 28 ha (5%) is highly suitable (Class S1) for growing sunflower and are distributed in the southern part of the microwatershed. Maximum area of about 310 ha (54%) is moderately suitable (Class S2) and are distributed in the major part of the microwatershed. They have minor limitations of rooting depth and calcareousness. Marginally suitable (Class S3) lands occupy an area of about 57 ha (10%) and are distributed in the northern and central part of the microwatershed with moderate limitations of rooting depth, calcareousness and gravelliness.




Fig. 7.7 Land Suitability map of Sunflower

7.8 Land Suitability for Cotton (*Gossypium hirsutum*)

Cotton is one of the most important fibre crop grown in the State in about 8.75 lakh ha area in Raichur, Dharwad, Belgaum, Gulbarga, Bijapur, Bidar, Bellary, Chitradurga and Chamarajnagar districts. The crop requirements for growing cotton (Table 7.9) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing cotton was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.8.

An area of about 28 ha (5%) is highly suitable (Class S1) for growing cotton and are distributed in the central part of the microwatershed. Maximum area of about 347 ha (60%) is moderately suitable (Class S2) and are distributed in the major part of the microwatershed. They have minor limitations of rooting depth, calcareousness and gravelliness. Marginally suitable (Class S3) lands occupy an area of about 21 ha (4%) and are distributed in the central part of the microwatershed with moderate limitations of rooting depth and gravelliness.




Fig. 7.8 Land Suitability map of Cotton

7.9 Land Suitability for Chilli (*Capsicum annum L*)

Chilli is one of the most important spice crop grown in an area of 0.42 lakh ha in Karnataka State. The crop requirements for growing chilli (Table 7.10) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing chilli was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.9.

An area of about 28 ha (5%) is highly suitable (Class S1) for growing chilli and are distributed in the southern part of the microwatershed. Marginally suitable (Class S3) lands cover a maximum area of about 368 ha (64%) and distributed in the major part of the microwatershed. They have moderate limitations of gravelliness, texture and calcareousness.




Fig. 7.9 Land Suitability map of Chili

7.10 Land Suitability for Tomato (*Solanum lycopersicum*)

Tomato is one of the most important vegetable crop grown in an area of 0.65 lakh ha in almost all the districts of the State. The crop requirements (Table 7.11) for growing tomato were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing tomato was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.10.

An area of about 28 ha (5%) is highly suitable (Class S1) for growing tomato and are distributed in the southern part of the microwatershed. Marginally suitable (Class S3) lands cover a maximum area of about 368 ha (64%) and distributed in the major part of the microwatershed. They have moderate limitations of gravelliness, texture and calcareousness.




Fig. 7.10 Land Suitability map of Tomato

7.11 Land Suitability for Brinjal (*Solanum melongena*)

Brinjal is one of the most important vegetable crop grown in the state. The crop requirements for growing brinjal (Table 7.12) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing brinjal was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.11.

Maximum area of about 375 ha (65%) is moderately suitable (Class S2) for growing Brinjal and distributed in the major part of the microwatershed with minor limitations of texture, rooting depth, calcareousness and gravelliness. Marginally suitable (Class S3) lands cover an area of about 21 ha (4%) and occur in the central part of the microwatershed with moderate limitation of gravelliness.




Fig 7.11 Land Suitability map of Brinjal

7.12 Land Suitability for Onion (*Allium cepa L.*)

Onion is one of the most important vegetable crop grown in the state. The crop requirements for growing onion (Table 7.13) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing onion was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.12.

An area of about 28 ha (5%) is moderately suitable (Class S2) for growing Onion and distributed in the southern part of the microwatershed with minor limitations of texture and gravelliness. Marginally suitable (Class S3) lands cover a maximum area of about 368 ha (64%) and occur in the major part of the microwatershed with moderate limitations of texture and calcareousness.




Fig 7.12 Land Suitability map of Onion

7.13 Land Suitability for Bhendi (*Abelmoschus esculentus*)

Bhendi is one of the most important vegetable crop grown in the state. The crop requirements for growing bhendi (Table 7.14) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing bhendi was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.13.

Moderately suitable (Class S1) lands occupy a maximum area of about 375 ha (65%) for growing Bhendi and occur in the major part of the microwatershed with minor limitations of rooting depth, texture, gravelliness and calcareousness. An area of about 21 ha (4%) is Marginally suitable (Class S3) for growing Bhendi and distributed in the central part of the microwatershed with moderate limitation of gravelliness.




Fig 7.13 Land Suitability map of Bhendi

7.14 Land Suitability for Drumstick (*Moringa oleifera*)

Drumstick is one of the most important vegetable crop grown in 2403 ha area in the state. The crop requirements for growing drumstick (Table 7.15) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing drumstick was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.14.

An area of about 28 ha (5%) is highly suitable (Class S1) for growing drumstick and are distributed in the southern part of the microwatershed. Moderately suitable (Class S2) lands cover a maximum area of about 310 ha (54%) and are distributed in the major part of the microwatershed. They have minor limitations of rooting depth, calcareousness and texture. Marginally suitable (Class S3) lands cover an area of about 57 ha (10%) and occur in the northern and central part of the microwatershed. They have moderate limitations of gravelliness, calcareousness and rooting depth.




Fig. 7.14 Land Suitability map of Drumstick

7.15 Land Suitability for Mulberry (*Morus nigra*)

Mulberry is the most important leaf crop grown for rearing silkworms in about 1.66 lakh ha in all the districts of the state. The crop requirements for growing mulberry (Table 7.16) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing mulberry was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.15.

An area of about 28 ha (5%) is highly suitable (Class S1) for growing mulberry and are distributed in the southern part of the microwatershed. Maximum area of about 281 ha (49%) is moderately suitable (Class S2) for growing mulberry and distributed in the major part of the microwatershed. They have minor limitations of texture, gravelliness and calcareousness. Marginally suitable (Class S3) lands cover an area of about 87 ha (15%) and occur in the northern part of the microwatershed. They have moderate limitations of rooting depth, texture and calcareousness.




Fig. 7.15 Land Suitability map of Mulberry

7.16 Land Suitability for Mango (*Mangifera indica*)

Mango is one of the most important fruit crop grown in about 1.73 lakh ha in almost all the districts of the State. The crop requirements (Table 7.17) for growing mango were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing mango was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.16.

An area of about 28 ha (5%) is highly suitable (Class S1) for growing mango and are distributed in the southern part of the microwatershed. An area of about 98 ha (17%) is moderately suitable (Class S2) for growing mulberry and distributed in the eastern and northwestern part of the microwatershed. They have minor limitations of calcareousness and rooting depth. Marginally suitable (Class S3) lands cover an area of about 233 ha (41%) and occur in the major part of the microwatershed. They have moderate limitations of rooting depth, gravelliness, texture and calcareousness. Area currently not suitable (Class N1) for growing mango cover about 36 ha (6%) and distributed in the northern part of the microwatershed with severe limitations of rooting depth and calcareousness.




Fig. 7.16 Land Suitability map of Mango

7.17 Land Suitability for Sapota (*Manilkara zapota*)

Sapota is one of the most important fruit crop grown in an area of about 29373 ha in almost all the districts of the state. The crop requirements (Table 7.18) for growing sapota were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing sapota was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.17.

An area of about 28 ha (5%) is highly suitable (Class S1) for growing sapota and are distributed in the southern part of the microwatershed. Moderately suitable (S2) lands cover an area of about 21 ha (4%) and are distributed in the central part of the microwatershed. They have minor limitations of rooting depth and graveliness. Marginally suitable (Class S3) lands cover a maximum area of about 347 ha 60 %) and occur in the major part of the microwatershed. They have moderate limitations of rooting depth, calcareousness and texture.




Fig. 7.17 Land Suitability map of Sapota

7.18 Land Suitability for Pomegranate (*Punica granatum*)

Pomegranate is one of the commercially grown fruit crop in about 18488 ha in Karnataka mainly in Bijapur, Bagalkot, Koppal, Gadag and Chitradurga districts. The crop requirements for growing pomegranate (Table 7.19) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing pomegranate was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.18.

An area of about 28 ha (5%) is highly suitable (Class S1) for growing pomegranate and are distributed in the southern part of the microwatershed. Moderately suitable (Class S2) lands occupy an area of about 331 ha (58%) and are distributed in the major part of the microwatershed. They have minor limitations of rooting depth, gravelliness, calcareousness and texture. Marginally suitable (Class S3) lands for growing pomegranate occupy an area of about 36 ha (6%) and are distributed in the northern part of the microwatershed with moderate limitations of calcareousness and rooting depth.




Fig. 7.18 Land Suitability map of Pomegranate

7.19 Land Suitability for Guava (*Psidium guajava*)

Guava is one of the most important fruit crop grown in an area of about 6558 ha in almost all the districts of the state. The crop requirements (Table 7.20) for growing guava were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing guava was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.19.

Moderately suitable (Class S2) lands occupy an area of about 49 ha (9%) and are distributed in the southern part of the microwatershed. They have minor limitations of rooting depth, gravelliness and texture. Marginally suitable (Class S3) lands for growing guava occupy a maximum area of about 347 ha (60%) and are distributed in the major part of the microwatershed with moderate limitations of calcareousness and texture.




Fig. 7.19 Land Suitability map of Guava

7.20 Land Suitability for Jackfruit (*Artocarpus heterophyllus*)

Jackfruit is one of the most important fruit crop grown in 5368 ha in all the districts of the state. The crop requirements (Table.7.21) for growing jackfruit were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing jackfruit was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in figure 7.20.

An area of about 28 ha (5%) is highly suitable (Class S1) for growing jackfruit and are distributed in the southern part of the microwatershed. Moderately suitable (Class S2) lands cover an area of about 21 ha (4%) and are distributed in the southern and central part of the microwatershed. They have minor limitations of rooting depth and gravelliness. Marginally suitable (Class S3) lands cover a maximum area of about 347 ha (60%) and occur in the major part of the microwatershed. They have moderate limitations of calcareousness and texture.




Fig. 7.20 Land Suitability map of Jackfruit

7.21 Land Suitability for Jamun (*Syzygium cumini*)

Jamun is an important fruit crop grown in almost all the districts of the state. The crop requirements (Table 7.22) for growing jamun were matched with the soil-site characteristics and a land suitability map for growing jamun was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.21.

An area of about 28 ha (5%) is highly suitable (Class S1) for growing jamun and are distributed in the southern part of the microwatershed. Moderately suitable (Class S2) lands occupy an area of about 233 ha (41%) and distributed in the major part of the microwatershed. They have minor limitations of rooting depth, texture, calcareousness and gravelliness. Marginally suitable (Class S3) lands cover a maximum area of about 134 ha (23%) and are distributed in the northern and central part of the microwatershed with moderate limitations of rooting depth, calcareousness and texture.




Fig. 7.21 Land Suitability map of Jamun

7.22 Land Suitability for Musambi (*Citrus limetta*)

Musambi is one of the most important fruit crop grown in an area of 5446 ha in almost all the districts of the state. The crop requirements (Table 7.23) for growing musambi were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing musambi was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.22.

An area of about 28 ha (5%) is highly suitable (Class S1) for growing musambi and are distributed in the southern part of the microwatershed. An area of about 331 ha (58%) is moderately suitable (Class S2) and occur in the major part of the microwatershed. They have minor limitations of calcareousness, gravelliness and rooting depth. An area of about 36 ha (6%) is marginally suitable (Class S3) for growing musambi and are distributed in the northern part of the microwatershed with moderate limitations of calcareousness and rooting depth.




Fig. 7.22 Land Suitability map of Musambi

7.23 Land Suitability for Lime (*Citrus sp*)

Lime is one of the most important fruit crop grown in an area of 11752 ha in almost all the districts of the State. The crop requirements for growing lime (Table 7.24) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing lime was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.23.

An area of about 28 ha (5%) is highly suitable (Class S1) for growing lime and are distributed in the southern part of the microwatershed. An area of about 331 ha (58%) is moderately suitable (Class S2) and occur in the major part of the microwatershed. They have minor limitations of calcareousness, gravelliness and rooting depth. An area of about 36 ha (6%) is marginally suitable (Class S3) for growing lime and are distributed in the northern part of the microwatershed with moderate limitations of calcareousness and rooting depth.




Fig. 7.23 Land Suitability map of Lime

7.24 Land Suitability for Cashew (*Anacardium occidentale*)

Cashew is one of the most important nut crop grown in an area of 7052 ha in almost all the districts of the State. The crop requirements for growing cashew (Table 7.25) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing cashew was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.24.

An area of about 49 ha (9%) is moderately suitable (Class S2) and occur in the southern part of the microwatershed. They have minor limitations of gravelliness, texture and rooting depth. Maximum area of about 347 ha (60%) is currently not suitable (Class N1) for growing cashew and distributed in the major part of the microwatershed with severe limitations of texture and calcareousness.




Fig. 7.24 Land Suitability map of Cashew

7.25 Land Suitability for Custard Apple (*Annona reticulata*)

Custard apple is one of the most important fruit crop grown in 1426 ha in almost all the districts of the State. The crop requirements(Table 7.26) for growing custard apple were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing custard apple was generated .The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.25.

An area of about 28 ha (5%) is highly suitable (Class S1) for growing custard apple and are distributed in the southern part of the microwatershed. Moderately suitable (Class S2) lands cover a maximum area of about 368 ha (64%) and occur in the major part of the microwatershed. They have minor limitations of rooting depth, calcareousness and gravelliness.




Fig. 7.25 Land Suitability map of Custard Apple

7.26 Land Suitability for Amla (*Phyllanthus emblica*)

Amla is one of the most important fruit and medicinal crop grown in an area of 151 ha and distributed in almost all the districts of the state. The crop requirements (Table 7.27) for growing amla were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing amla was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.26.

An area of about 28 ha (5%) is highly suitable (Class S1) for growing amla and are distributed in the southern part of the microwatershed. Moderately suitable (Class S2) lands cover a maximum area of about 368 ha (64%) and occur in the major part of the microwatershed. They have minor limitations of rooting depth, graveliness, calcareousness and texture.




Fig. 7.26 Land Suitability map of Amla

7.27 Land Suitability for Tamarind (*Tamarindus indica*)

Tamarind is one of the most important spice crop grown in 14897 ha in all the districts of the state. The crop requirements (Table 7.28) for growing tamarind were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing tamarind was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.27.

An area of about 28 ha (5%) is highly suitable (Class S1) for growing tamarind and are distributed in the southern part of the microwatershed. An area of about 212 ha (37%) is moderately suitable (Class S2) and occur in the eastern and northern part of the microwatershed. They have minor limitations of rooting depth, texture and calcareousness. An area of about 119 ha (21%) is marginally suitable (Class S3) for growing tamarind and are distributed in the central part of the microwatershed with moderate limitations of rooting depth, calcareousness and gravelliness. An area of about 36 ha (6%) is currently not suitable (Class N1) for growing tamarind and distributed in the northern part of the microwatershed with severe limitations of rooting depth and calcareousness.




Fig. 7.27 Land Suitability map of Tamarind

7.28 Land Suitability for Marigold (*Tagetes erecta*)

Marigold is one of the most important flower crop grown in an area of 9108 ha in almost all the districts of the state. The crop requirements (Table 7.29) for growing marigold were matched with the soil-site characteristics and a land suitability map for growing marigold was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.28.

An area of about 28 ha (5%) is highly suitable (Class S1) for growing marigold and are distributed in the southern part of the microwatershed. Maximum area of about 347 ha (60%) is moderately suitable (Class S2) and occur in the major part of the microwatershed. They have minor limitations of calcareousness, rooting depth and texture. An area of about 21 ha (4%) is marginally suitable (Class S3) for growing marigold and are distributed in the central part of the microwatershed with moderate limitation of gravelliness.




Fig. 7.28 Land Suitability map of Marigold

7.29 Land Suitability for Chrysanthemum (*Chrysanthemum indicum*)

Chrysanthemum is one of the most important flower crop grown in an area of 4978 ha in almost all the districts of the State. The crop requirements (Table 7.30) for growing chrysanthemum were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing chrysanthemum was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed is given in Figure 7.29.

An area of about 28 ha (5%) is highly suitable (Class S1) for growing chrysanthemum and are distributed in the central part of the microwatershed. Maximum area of about 347 ha (60%) is moderately suitable (Class S2) and occur in the major part of the microwatershed. They have minor limitations of rooting depth, calcareousness and texture. An area of about 21 ha (4%) is marginally suitable (Class S3) for growing chrysanthemum and are distributed in the central part of the microwatershed with moderate limitation of gravelliness.




Fig. 7.29 Land Suitability map of Chrysanthemum

7. 30 Land Suitability for Jasmine (*Jasminum sp.*)

Jasmine is one of the most important flower crop grown in an area of 803 ha in almost all the districts of the State. The crop requirements (Table 7.31) for growing jasmine were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing jasmine was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.30.

An area of about 28 ha (5%) is highly suitable (Class S1) for growing jasmine and are distributed in the southern part of the microwatershed. An area of about 36 ha (6%) is moderately suitable (Class S2) and occur in the northern part of the microwatershed. They have minor limitation of calcareousness and rooting depth. Maximum area of about 332 ha (58%) is marginally suitable (Class S3) for growing jasmine and are distributed in the major part of the microwatershed with moderate limitations of gravelliness, texture and calcareousness.




Fig. 7.30 Land Suitability map of Jasmine

7. 31 Land Suitability for Crossandra (*Crossandra infundibuliformis*)

Crossandra is one of the most important flower crop grown in almost all the districts of the State. The crop requirements (Table 7.32) for growing crossandra were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing crossandra was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.31.

An area of about 28 ha (5%) is highly suitable (Class S1) for growing crossandra and are distributed in the southern part of the microwatershed. An area of about 98 ha (17%) is moderately suitable (Class S2) and occur in the central part of the microwatershed. They have minor limitation of calcareousness. Maximum area of about 269 ha (47%) is marginally suitable (Class S3) for growing crossandra and are distributed in the major part of the microwatershed with moderate limitations of gravelliness, rooting depth, calcareousness and texture.




Fig. 7.31 Land Suitability map of Crossandra

Table 7.1 Soil-Site Characteristics of Vojnahalli-2 Microwatershed

Soil Map Units	Climate (P) (mm)	Growing period (Days)	Drainage Class	Soil depth (cm)	Soil texture		Gravelliness		AWC (mm/m)	Slope (%)	Erosion	pH	EC (dSm ⁻¹)	ESP	CEC [Cmol (p ⁺)kg ⁻¹]	BS (%)
					Surf-ace	Sub-surface	Sur-face	Sub-surface								
HDHhB1g1	662	<90	WD	75-100	scl	gsc-gc	15-35	>35	51-100	1-3	slight	6.54	0.07	7.11	5.84	84.7
HDHiB1	662	<90	WD	75-100	sc	gsc-gc	-	>35	51-100	1-3	slight	6.54	0.07	7.11	5.84	84.7
GDPiB2	662	<90	WD	100-150	sc	gsc-gc	-	30-60	51-100	1-3	moderate	7.88	0.10	2.87	7.8	97
RNKmB1	662	<90	MWD	50-75	c	c	-	<15	101-150	1-3	slight	8.86	0.48	16.94	37.0	-
RNKmB2g1	662	<90	MWD	50-75	c	c	15-35	<15	101-150	1-3	moderate	8.86	0.48	16.94	37.0	-
DRLmB2	662	<90	MWD	75-100	c	c	-	<15	151-200	1-3	moderate	8.78	0.42	5.62	49.70	100
KVRiB2g1	662	<90	MWD	100-150	sc	c	-	-	>200	1-3	moderate	8.4	0.26	0.60	43.25	-
KVRmB1	662	<90	MWD	100-150	c	c	-	-	>200	1-3	slight	8.4	0.26	0.60	43.25	-
KVRmB2g1	662	<90	MWD	100-150	c	c	15-35	-	>200	1-3	moderate	8.4	0.26	0.60	43.25	-
BGPmA1	662	<90	MWD	>150	c	c	-	<15	>200	0-1	slight	9.20	0.27	3.84	19.60	100
BGPmB1	662	<90	MWD	>150	c	c	-	<15	>200	1-3	slight	9.20	0.27	3.84	19.60	100
AWDmB2	662	<90	MWD	>150	c	c	-	<15	>200	1-3	moderate	8.10	0.37	1.22	51.30	100

Table 7.2 Land suitability criteria for Sorghum

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime 1	Mean temperature in growing season	°C	26–30	30–34; 24–26	34–40; 20–24	>40; <20
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristics					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V. poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, c (red), c (black)	scl, cl	ls, sl	-
	pH	1:2.5	5.5-7.8	5.0-5.5 7.8-9.0	>9.0	-
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	10-15
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

Table 7.3 Land suitability criteria for Maize

Land use requirement		Rating				
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	30-34	35-38 26-30	38-40 26-20	
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc	c (red), c (black)	ls, sl	-
	pH	1:2.5	5.5-7.8	5.0-5.5 7.8-9.0	>9.0	-
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	-
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

Table 7.4 Land suitability criteria for Bajra

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-32	33-38 24-27	39-40 20-23	<20
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm	500-750	400-500	200-400	<200
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	Sl, scl, cl,sc,c (red)	C (black)	ls	-
	pH	1:2.5	6.0-7.8	5.0-5.5 7.8-9.0	5.5-6.0 >9.0	
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	15-35	35-60	>60	
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	1-3	3-5	5-10	>10

Table 7.5 Land suitability criteria for Red gram

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	30-35(G) 20-25(AV) 15-18 (F&PS) 35-40(M)	25-30(G) 20-25 (AV) 12-15 (F&PS) 30-35(M)	20-25(G) 15-20(AV) 10-12 (F&PS) 25-30(M)	< 20 <15 <10 <25
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. Well drained	Poorly drained	Very Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, c (red)	c (black),sl, scl, cl	ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-9.0	5.0-5.5 >9.0	-
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-50	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<1.0	1.0-2.0	>2.0	
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.6 Land suitability criteria for Bengal gram

Land use requirement			Rating			
Soil –site characteristics	Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)	
Climatic regime	Mean temperature in growing season	°C	20–25	25–30; 15–20	30–35; 10–15	>35; <10
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. Well drained	Poorly drained	Very Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	C (black)	-	c (red), scl, cl, sc	ls, sl
	pH	1:2.5	6.0-7.8	5.0-6.0 7.8-9.0	>9.0	-
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	-
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.7 Land suitability criteria for Groundnut

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	24–33	22–24; 33–35	20–22; 35–40	<20; >40
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. Well drained	Poorly drained	Very Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl	sl,cl, sc	c (red), c (black), ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<35	35-60	>60	
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.8 Land suitability criteria for Sunflower

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	24–30	30–34; 20–24	34–38; 16–20	>38; <16
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	mod. Well drained	-	Poorly to very drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	cl, sc,c (red), c (black)	scl	ls, sl	-
	pH	1:2.5	6.5-7.8	7.8-8.4 5.5-6.5	8.4-9.0; 5.0-5.5	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.9 Land suitability criteria for Cotton

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	22-32	>32	<19	-
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well to moderately well	Poorly drained/Some what excessively drained	-	very poorly/excessively drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, c (red,black)	cl	scl	ls, sl
	pH	1:2.5	6.5-7.8	7.8-8.4	5.5-6.5 8.4->9.0	<5.5
	CEC	C mol (p+)Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	50-100	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	<3	3-5	-	>5

Table 7.10 Land suitability criteria for Chili

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	25-32	33-35 20-25	35-38 <20	>38
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc	c (black), sl	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.11 Land suitability criteria for Tomato

Land use requirement		Rating				
Soil –site characteristics	Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)	
Climatic regime	Mean temperature in growing season	°C	25-28	29-32 20-24	15-19 33-36	<15 >36
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl, scl, cl, sc, c (red)	-	ls, c(black)	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.12 Land suitability criteria for Brinjal

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	Well drained	Moderately well drained	Poorly drained	V. Poorly drained
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class				
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl, scl, cl, sc c (red)	-	ls, c (black)	-
	pH	1:2.5	6.0-7.3	7.3-8.4 5.0-6.0	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	>60
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.13 Land suitability criteria for Onion

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	20-30	30-35	35-40	>40
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately /imperfectly	-	Poorly to V poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl,scl,cl,sc,c (red)	-	c (Black),ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<1.0	1.0-2.0	2.0-4.0	<4
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.14 Land suitability criteria for Bhendi

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	25-28	29-32 20-24	15-19 33-36	<15 >36
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Imperfectly drained	Poorly to very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl,sc, c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.15 Land suitability criteria for Drumstick

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, scl, cl, c (red)	sl, c (black)	ls	s
	pH	1:2.5	6.0-7.3	5.0-5.5 7.3-7.8	5.5-6.0 7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<35	35-60	60-80	>80
Soil toxicity	Salinity (EC saturation extract)	dS/m				
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-10	-	>10

Table 7.16 Land suitability criteria for Mulberry

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	24–28	22–24; 28–32	32–38; 22–18	>38; <18
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V. Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, cl, scl	c (red)	c (black), sl, ls	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.8-8.4	7.3-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	0-35	35-60	60-80	>80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

Note: Suitability evaluation only for Mulberry leaf not for Silk worm rearing

Table 7.17 Land suitability criteria for Mango

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-32	24-27 33-35	36-40	20-24
	Min temp. before flowering	°C	10-15	15-22	>22	-
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration	Days				
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	-	ls, sl, c (black)	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>150	100-150	75-100	<75
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.18 Land suitability criteria for Sapota

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-32	33-36 24-27	37-42 20-23	>42 <18
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	-	Poorly to very drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	sl	ls, c (black)	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.19 Land suitability criteria for Pomegranate

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	30-34	35-38 25-29	39-40 15-24	
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl,cl, sc, c (red)	c (black),sl	ls	-
	pH	1:2.5	5.5-7.8	7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.20 Land suitability criteria for Guava

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-32	33-36 24-27	37-42 20-23	
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	sl	c (black), ls	-
	pH	1:2.5	6.0-7.8	5.0-6.0	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.21 Land suitability criteria for Jackfruit

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. well	Poorly	V. Poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	-	sl, ls, c (black)	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	>60
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10-

Table 7.22 Land suitability criteria for Jamun

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well	Mod. well	Poorly	V.Poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c(red)	sl, c (black)	ls	-
	pH	1:2.5	6.0-7.8	5.0-6.0	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>150	100-150	50-100	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	>60
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

Table 7.23 Land suitability criteria for Musambi

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-30	31-35 24-27	36-40 20-23	>40 <20
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately drained	poorly	Very poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c	sl	ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.24 Land suitability criteria for Lime

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-30	31-35 24-27	36-40 20-23	>40 <20
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately drained	poorly	Very poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c	sl	ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.25 Land suitability criteria for Cashew

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	32 to 34	28 to 32; 34 to 38	24 to 28; 38 to 40	<20; >40
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	-	sl, ls	c (black)
	pH	1:2.5	5.5-6.5	5.0-5.5 6.5-7.3	7.3-7.8	>7.8
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-10	>10	-

Table 7.26 Land suitability criteria for Custard apple

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	Scl, cl, sc, c (red), c (black)	-	Sl, ls	-
	pH	1:2.5	6.0-7.3	5.5-6.0 7.3-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15-35	35-60	60-80	-
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	>5	-

Table 7.27 Land suitability criteria for Amla

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod.well drained	Poorly drained	V. Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	c (black)	ls, sl	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15-35	35-60	60-80	-
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

Table 7.28 Land suitability criteria for Tamarind

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod.well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl,sc, c (red)	sl, c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>150	100-150	75-100	<75
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

Table 7.29 Land suitability criteria for Marigold

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	18-23	17-15 24-35	35-40 10-14	>40 <10
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl,scl, cl, sc, c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%				
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.30 Land suitability criteria for Chrysanthemum

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	18-23	17-15 24-35	35-40 10-14	>40 <10
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl,scl, cl, sc, c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%				
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.31 Land suitability criteria for Jasmine (irrigated)

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	18-23	17-15 24-35	35-40 10-14	-
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	sl	ls, c (black)	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%				
Erosion hazard	Slope	%	<3	3-5	5-10	>10

7.32 Land suitability criteria for Crossandra

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	-	Poorly to very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c(red)	sl,	c (black),ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%				
Erosion hazard	Slope	%	<3	3-5	5-10	>10

7.29 Land Management Units (LMUs)

The 12 soil map units identified in Vojenahalli-2 Microwatershed have been grouped into three Land Management Units (LMUs) for the purpose of preparing a Proposed Crop Plan. Land Management Units are grouped based on the similarities in respect of the type of soil, the depth of the soil, the surface soil texture, gravel content, AWC, slope, erosion etc. and a Land Management Units map (Fig.7.32) has been generated. These Land Management Units are expected to behave similarly for a given level of management.

The map units that have been grouped into three Land Management Units along with brief description of soil and site characteristics are given below.

LMU	Mapping unit	Soil and site characteristics
1	AWDmB2, BGPmA1, BGPmB1, KVRiB2g1, KVRmB1, KVRmB2g1, DRLmB2	Moderately deep to very deep, black calcareous clay soils with slopes of 0-3%, slight to moderate erosion, gravelly (15-35%)
2	GDPiB2, HDHhB1g1, HDHiB1	Moderately deep to deep, red gravelly sandy clay to clay soils with slopes of 1-3%, slight to moderate erosion, gravelly (15-35%)
3	RNKmB1, RNKmB2g1	Moderately shallow, black calcareous clay soils with slopes of 1-3%, slight to moderate erosion, gravelly (15-35%)




Fig 7.32 Land Management Units map of Vojenahalli-2 microwatershed

7.30 Proposed Crop Plan for Vojenahalli-2 Microwatershed

After assessing the land suitability for the 31 crops, the proposed crop plan has been prepared for the three identified LMUs by considering only the highly (Class S1) and moderately (Class S2) suitable lands for each of the 28 crops. The resultant proposed crop plan is presented in Table 7.30.

Table 7.30 Proposed Crop Plan for Vojenahalli-2 Microwatershed

LMU	Soil Map Units	Survey Number	Field Crops/ Commercial crops	Horticulture Crops (Rainfed/Irrigated)	Suitable Interventions
1	424.AWDmB2 395.BGPmA1 396.BGPmB1 385.KVRiB2g1 388.KVRmB1 390.KVRmB2g1 350.DRLmB2 (Moderately deep to very deep, black calcareous clay soils)	Ojanahalli: 39,40,48,50,52,53,54,55,56,57,58,59,60,61,62,63,64,65,66,67,68,69,70,73,74,77,78,79,80,81,82,83,84,85,86,87,88,89,90,91,104,115,116,117,118,119,120,121,122,123,124,126,127,146 Koppal: 25,26,27,28,29,30,106,107,686,701,702,703,704,709,710,711,713, 714,715,716	Maize, Sorghum, Sunflower, Cotton, Bengal gram, Safflower, Linseed, Bajra , Soybean	Fruit crops: Sapota, Pomegranate, Jamun, Lime, Musambi, Tamarind, Amla, Custard apple Vegetables: Drumstick, Chilli, Coriander, Tomato, Bhendi Flowers: Marigold, Chrysanthemum, Crossandra, Jasmine	Application of FYM, Biofertilizers and micronutrients, drip irrigation, mulching, suitable soil and water conservation practices
2	269.GDPiB2 120.HDHhB1g1 125.HDHiB1 (Moderately deep to deep, red gravelly sandy clay to clay soils)	Ojanahalli: 51 Koppal: 37,42,43,44,45,46,51,52,53,54,55,57,61	Groundnut, Bajra, Horse gram, Castor, Mulberry	Fruit crops: Musambi, Lime, Jamun, Jackfruit Amla, Custard apple, Tamarind Vegetable crops: Drumstick, Curry leaves	Drip irrigation, mulching, suitable soil and water conservation practices (Crescent Bunding with Catch Pit etc)
3	333.RNKmB1 337.RNKmB2g1 (Moderately shallow, black calcareous clay soils)	Ojanahalli: 37,92,93,94,95,96,97,102,103,114	Sorghum, Bajra, Bengal gram, linseed, Safflower, Coriander	Fruit crops: Amla, Custard apple Flower crops: Marigold, Jasmine Chrysanthemum	Application of FYM, Biofertilizers and micronutrients, drip irrigation, mulching, suitable soil and water conservation practices

SOIL HEALTH MANAGEMENT

8.1 Soil Health

Soil health is basic to plant health and plant health is basic to human and bovine health. Soil is fundamental to crop production. Without soil, no food could be produced nor would livestock be fed on a large scale. Because it is finite and fragile, soil is a precious resource that requires special care from its users.

Soil health or the capacity of the soil to function is critical to human survival. Soil health has been defined as: “the capacity of the soil to function as a living system without adverse effect on the ecosystem”. Healthy soils maintain a diverse community of soil organisms that help to form beneficial symbiotic associations with plant roots, recycle essential plant nutrients, improve soil structure with positive repercussions for soil, water and nutrient holding capacity and ultimately improve crop production and also contribute to mitigating climate change by maintaining or increasing its carbon content.

Functional interactions of soil biota with organic and inorganic components, air and water determine a soil’s potential to store and release nutrients, and water to plants and to promote and sustain plant growth. Thus, maintaining soil health is vital to crop production and conserve soil resource base for sustaining agriculture.

The most important characteristics of a healthy soil are

- Good soil tilth
- Sufficient soil depth
- Good water storage and good drainage
- Adequate supply, but not excess of nutrients
- Large population of beneficial organisms
- Small proportion of plant pathogens and insect pests
- Low weed pressure
- Free of chemicals and toxins that may harm the crop
- Resistance to degradation
- Resilience when unfavourable conditions occur

Characteristics of Vojenahalli-2 Microwatershed

- ❖ The soil phases with sizeable area identified in the microwatershed belonged to the soil series of KVR (98 ha), DRL (98 ha), BGP (63 ha), AWD (51 ha), RNK (36 ha), GDP (28 ha) and HDH (21 ha).
- ❖ As per land capability classification, entire area in the microwatershed falls under arable land category (Class II and III). The major limitations identified in the arable lands were soil and erosion.

- ❖ On the basis of soil reaction, an area of about 8 ha (1%) is slightly alkaline (pH 7.3-7.8), 86 (15%) is moderately alkaline, 287 ha (50%) is strongly alkaline (pH 8.4-9.0) and 15 ha (3%) is very strongly alkaline (pH >9.0) in reaction.

Soil Health Management

The following actions are required to improve the current land husbandry practices that provide a sound basis for the successful adoption of sustainable crop production system.

Alkaline soils

An area of about 396 ha (69%) is under alkaline soils. The following actions are recommended.

1. Regular addition of organic manure, green manuring, green leaf manuring, crop residue incorporation and mulching needs to be taken up to improve the soil organic matter status.
2. Application of biofertilizers (Azospirillum, Azotobacter, Rhizobium).
3. Application of 25% extra N and P (125 % RDN&P).
4. Application of ZnSO₄ – 12.5 kg/ha (once in three years).
5. Application of Boron – 5 kg/ha (once in three years).

Soil Degradation

Soil erosion is one of the major factor affecting the soil health in the microwatershed. An area of about 198 ha (34%) is under moderate erosion. The areas with moderate erosion need immediate soil and water conservation and, other land development and land husbandry practices for restoring soil health.

Dissemination of Information and Communication of Benefits

Any large scale implementation of soil health management requires that supporting information is made available widely, particularly through channels familiar to farmers and extension workers. Given the very high priority attached to soil health especially by the Central Government on issuing Soil-Health Cards to all the farmers, media outlets like Regional, State and National Newspapers, Radio and Dooradarshan programs in local languages but also modern information and communication technologies such as Cellular phones and the Internet, which can be much more effective in reaching the younger farmers.

Inputs for Net Planning (Saturation Plan) and Interventions needed

Net planning in IWMP is focusing on preparation of

1. Soil and Water Conservation Treatment Plans for each plot or farm.
2. Productivity enhancement measures/ interventions for existing crops/livestock/other farm enterprises.
3. Diversification of farming mainly with perennial horticultural crops and livestock.

4. Improving livelihood opportunities and income generating activities.

In this connection, how various outputs of Sujala-III are of use in addressing these objectives of Net Planning are briefly presented below.

- ❖ **Soil Depth:** The depth of a soil decides the amount of moisture and nutrients it can hold, what crops can be taken up or not, depending on the rooting depth and the length of growing period available for raising any crop. Deeper the soil, better for a wide variety of crops. If sufficient depth is not available for growing deep rooted crops, either choose medium or short duration crops or deeper planting pits need to be opened and additional good quality soil brought from outside has to be filled into the planting pits.
- ❖ **Surface Soil Texture:** Lighter soil texture in the top soil means, better rain water infiltration, less run-off and soil moisture conservation, less capillary rise and less evaporation losses. Lighter surface textured soils are amenable to good soil tilth and are highly suitable for crops like groundnut, root vegetables (carrot, radish, potato etc) but not ideal for crops that need stagnant water like lowland paddy. Heavy textured soils are poor in water infiltration and percolation. They are prone for sheet erosion; such soils can be improved by sand mulching. The technology that is developed by the AICRP-Dryland Agriculture, Vijayapura, Karnataka can be adopted.
- ❖ **Gravelliness:** More gravel content is favorable for run-off harvesting but poor in soil moisture storage and nutrient availability. It is a significant parameter that decides the kind of crop to be raised.
- ❖ **Land Capability Classification:** The land capability map shows the areas suitable and not suitable for agriculture and the major constraints in each of the plot/survey number. Hence, one can decide what kind of enterprise is possible in each of these units. In general, erosion and soil are the major constraints in Vojenahalli-2 Microwatershed.
- ❖ **Organic Carbon:** An area of about 105 ha (18%) is low (<0.5%) and 291 ha (51%) is medium (0.5-0.75%) in OC. The areas that are low and medium in OC needs to be further improved by applying farmyard manure and rotating crops with cereals and legumes or mixed cropping.
- ❖ **Promoting green manuring:** Growing of green manuring crops costs Rs. 1250/ha (green manuring seeds) and about Rs. 2000/ha towards cultivation that totals to Rs. 3250/- per ha. On the other hand, application of organic manure @ 10 tons/ha costs Rs. 5000/ha. The practice needs to be continued for 2-3 years or more. Nitrogen fertilizer needs to be supplemented by 25% in addition to the recommended level in 396 ha area where OC is less than 0.75 per cent. For example, for rainfed maize, recommended level is 50 kg N per ha and an additional 12 kg /ha needs to be applied for all the crops grown in these plots.
- ❖ **Available Phosphorus:** Available phosphorus is medium in (23-57 kg/ha) in the entire area. The areas with high phosphorus content reduce 25% from the RDF to


avoid the excess application of fertilizer and apply additional 25% phosphorus in areas where it is low and medium.

- ❖ **Available Potassium:** Available potassium is medium (145-337 kg/ha) in 351 ha (61%) and high (>337 kg/ha) in 45 ha (8%) area of the microwatershed. The areas with high potassium content reduce 25% from the RDF to avoid the excess application of fertilizer and apply additional 25% potassium in areas where it is medium.
- ❖ **Available Sulphur:** Available sulphur is a very critical nutrient for oilseed crops. Available sulphur is low (<10 ppm) in 200 ha (35%), medium in 127 ha (22%) and high (>20 ppm) in 69 ha (12%) area of the microwatershed. Areas with low and medium in available sulphur need to be applied with magnesium sulphate or gypsum or Factamphos (p) fertilizer (13% sulphur) for 2-3 years for the deficiency to be corrected.
- ❖ **Available Iron:** It is deficient (<4.5 ppm) in 396 ha (69 %) and sufficient (>4.5 ppm) in <1 ha (<1 %) area of the microwatershed. To manage iron deficiency iron sulphate @ 25 kg/ha needs to be applied for 2-3 years.
- ❖ **Available Zinc:** It is deficient (<0.6 ppm) in the 199 ha (35%) and sufficient (>0.6 ppm) in 197 ha (34 %) area of the microwatershed. Application of zinc sulphate @ 25kg/ha is to be followed in areas that are deficient in available zinc.
- ❖ **Available Boron:** Available boron is low in (<0.5ppm) 58 ha (10%), medium (0.5-1.0 ppm) in 331 ha (57%) and high (>1.0 ppm) in 8 ha (1%) area in the microwatershed. The areas with low and medium in boron content need to be applied with sodium borate @ 10kg/ha as soil application or 0.2% borax as foliar spray to correct the deficiency.
- ❖ **Available Manganese:** It is sufficient in the entire area of the microwatershed.
- ❖ **Available Copper:** It is sufficient in the entire area of the microwatershed.
- ❖ **Soil Alkalinity:** An area of about 396 ha (69%) in the microwatershed has soils that are slightly to very strongly alkaline. These areas need application of gypsum and wherever calcium is in excess, iron pyrites and element sulphur can be recommended. Management practices like treating repeatedly with good quality water to drain out the excess salts and provision of subsurface drainage and growing of salt tolerant crops like Casuarina, Acasia, Neem, Ber etc, are recommended.
- ❖ **Land Suitability for various crops:** Areas that are highly, moderately and marginally suitable and not suitable for growing various crops are indicated. Along with the suitability, various constraints that are limiting the productivity are also indicated. For example, in case of cotton, gravel content, rooting depth and salinity/alkalinity are the major constraints in various plots. With suitable management interventions, the productivity can be enhanced. In order to increase water holding capacity of light textured soils, growing of green manure crops and application of organic manure is recommended.

SOIL AND WATER CONSERVATION TREATMENT PLAN

For preparing soil and water conservation treatment plan for Vojenahalli-2 Microwatershed, the land resource inventory database generated under Sujala-III project has been transformed as information through series of interpretative (thematic) maps using soil phase map as a base. The various thematic maps (1:7920 scale) generated were

- Soil depth
- Surface soil texture
- Available water capacity
- Soil slope
- Soil gravelliness
- Land capability
- Present land use and land cover
- Crop suitability maps
- Rainfall map
- Hydrology
- Water Resources
- Socio-economic data
- Contour plan with existing features- network of waterways, pottissa boundaries, cut up/ minor terraces etc.
- Cadastral map (1:7920 scale)
- Satellite imagery (1:7920 scale)



Apart from these, Hand Level/ Hydro Marker/ Dumpy Level/ Total Station and Kathedars' List to be collected.

Steps for Survey and Preparation of Treatment Plan

The boundaries of Land User Groups' and Survey No. boundaries are traced in the field.


- Naming of user groups and farmers
- Identification of arable and non arable lands
- Identification of drainage lines and gullies
- Identification of non treatable areas
- Identification of priority areas in the arable lands
- Treatment plan for arable lands
- Location of water harvesting and recharge structures

9.1 Treatment Plan

The treatment plan recommended for arable lands is briefly described below.


9.1.1 Arable Land Treatment

A. BUNDING

Steps for Survey and Preparation of Treatment Plan		USER GROUP-1 CLASSIFICATION OF GULLIES 
Cadastral map (1:7920 scale) is enlarged to a scale of 1:2500 scale		
Existing network of waterways, pottissa boundaries, grass belts, natural drainage lines/ watercourse, cut ups/ terraces are marked on the cadastral map to the scale		
Drainage lines are demarcated into		
Small gullies	(up to 5 ha catchment)	
Medium gullies	(5-15 ha catchment)	
Ravines	(15-25 ha catchment) and	
<i>Halla/Nala</i>	(more than 25ha catchment)	

Measurement of Land Slope

Land slope is estimated or determined by the study and interpretation of contours or by measurement in the field using simple instruments like Hand Level or Hydromarker.



Vertical and Horizontal intervals between bunds as recommended by the Watershed Development Department.

Slope percentage	Vertical interval (m)	Corresponding Horizontal Distance (m)
2 - 3%	0.6	24
3 - 4%	0.9	21
4 - 5%	0.9	21
5 - 6%	1.2	21
6 - 7%	1.2	21

Note: i) The above intervals are maximum.

(ii) Considering the slope class and erosion status (A1... A= 0-1% slope, 1= slight erosion) the intervals have to be decided.

Bund length recording: Considering the contour plan and the existing grass belts/partitions, the bunds are aligned and lengths are measured.

Section of the Bund

Bund section is decided considering the soil texture class and gravelliness class (bg₀b = loamy sand, g₀ = <15% gravel). The recommended sections for different soils are given below.

Recommended Bund Section



Top width (m)	Base width (m)	Height (m)	Side slope (Z:1;H:V)	Cross section (sq m)	Soil Texture	Remarks
0.3	0.9	0.3	01:01	0.18	Sandy loam	Vegetative bund
0.3	1.2	0.3	1.5:1	0.225	Sandy clay	
0.3	1.2	0.5	0.9:1	0.375	Red gravelly soils	
0.3	1.2	0.6	0.75:1	0.45		
0.3	1.5	0.6	01:01	0.54	Red sandy loam	
0.3	2.1	0.6	1.5:1	0.72	Very shallow clayey black soils	
0.45	2	0.75	01:01	0.92		
0.45	2.4	0.75	1.3:1	1.07	Shallow clayey black soils	
0.6	3.1	0.7	1.78:1	1.29	Medium clayey black soils	
0.5	3	0.85	1.47:1	1.49		

Formation of Trench cum Bund

Dimensions of the Borrow Pits/ Trenches to be excavated (machinery are decided considering the Bund Section).

Details of Borrow Pit dimensions are given below

TRENCH CUM BUND





WATER STORAGE AREA

0.45 Sq.m section

IDEAL FOR HORTICULTURE CROPS

'A' FRAME FOR INTERBUND MANAGEMENT



1. ಸಮಸಾತಳ ಉಳುವು
2. ಸಮಸಾತಳ ಬಿತ್ತನೆ/ನಾಟಿ

Size of Borrow Pits/ Trench recommended for Trench cum Bund (by machinery)

Bund section	Bund length	Earth quantity	Pit				Berm (pit to pit)	Soil depth Class
			L(m)	W(m)	D(m)	Quantity (m ³)		
0.375	6	2.25	5.85	0.85	0.45	2.24	0.15	Shallow
0.45	6	2.7	5.4	1.2	0.43	2.79	0.6	Shallow
0.45	6	2.7	5	0.85	0.65	2.76	1	Moderately Shallow
0.54	5.6	3.02	5.5	0.85	0.7	3.27	0.1	Moderately shallow
0.54	5.5	2.97	5	1.2	0.5	3	0.5	Shallow
0.72	6.2	4.46	6	1.2	0.7	5.04	0.2	Moderately shallow
0.72	5.2	3.74	5.1	0.85	0.9	3.9	0.1	Moderately deep

B. Waterways

- a) Existing waterways are marked on the cadastral map (1:7920 scale) and their dimensions are recorded.
- b) Considering the contour plan of the MWS, additional waterways/ modernization of the existing ones can be thought of.
- c) The design details are given in the Manual.

C. Farm Ponds

Waterways and the catchment area will give an indication on the size of the Farm Pond. Location of the pond can be decided based on the contour plan/ field condition and farmers' need/desire.

D. Diversion Channel

Existing EPT/ CPT are marked on the cadastral map. Looking to the need, these can be modernized or fresh diversion channel can be proposed and runoff from this can be stored in *Gokatte/ Recharge Ponds*.

9.1.2 Non-Arable Land Treatment

Depending on the gravelliness and crops preferred by the farmers, the concerned authorities can decide appropriate treatment plan. The recommended treatments may be Contour Trench, Staggered Trench, Crescent Bund, Boulder Bund or Pebble Bund.

9.1.3 Treatment of Natural Water Course/ Drainage Lines

- a) The cadastral map has to be updated as regards the network of drainage lines (gullies/ *nalas/ hallas*) and existing structures are marked to the scale and storage capacity of the existing water bodies are documented.
- b) The drainage line will be demarcated into Upper Reach, Middle Reach and Lower Reach.
- c) Considering the Catchment, *Nala* bed and bank conditions, suitable structures are decided.
- d) Number of storage structures (Check dam/ *Nala* bund/ Percolation tank) will be decided considering the commitments and available runoff in water budgeting and quality of water in the wells and site suitability.
- e) Detailed Levelling Survey using Dumpy Level / Total Station has to be carried out to arrive at the site-specific designs as shown in the Manual.
- f) The location of ground water recharge structures are decided by examining the lineaments and fracture zones from geological maps.
- g) Rainfall intensity data of the nearest Rain Gauge Station is considered for Hydrologic Designs.
- h) Silt load to the Storage/Recharge Structures is reduced by providing vegetative, boulder and earthen checks in the natural water course. Location and design details are given in the Manual.

9.2 Recommended Soil and Water Conservation Measures

The appropriate conservation structures best suited for each of the land parcel/ survey number (Appendix-I) are selected based on the slope per cent, severity of erosion, amount of rainfall, land use and soil type. The different kinds of conservation structures recommended are

1. Graded / Strengthening of Bunds
2. Trench cum Bunds (TCB)
3. Trench cum Bunds / Strengthening
4. Crescent Bunds

A map (Fig. 9.1) showing soil and water conservation plan with different kinds of structures recommended has been prepared which shows the spatial distribution and extent of area. A maximum area of about 344 ha (60 %) needs graded bunding, an area of about 49 ha (9 %) needs trench cum bunding and 3 ha (<1 %) requires strengthening of existing bunds/ bunding. The conservation plan prepared may be presented to all the stakeholders including farmers and after considering their suggestions, the conservation plan for the microwatershed may be finalized in a participatory approach.




Fig. 9.1 Soil and Water Conservation Plan map of Vojenahalli-2 Microwatershed

9.3 Greening of Microwatershed

As part of the greening programme in the watersheds, it is envisaged to plant a variety of horticultural and other tree plants that are edible, economical and produce lot of biomass which helps to restore the ecological balance in the watersheds. The lands that are suitable for greening programme are non-arable lands (land capability classes V, VI VII and VIII) and also the lands that are not suitable or marginally suitable for growing annual and perennial crops. The method of planting these trees is given below.

It is recommended to open the pits during the 1st week of March along the contour and heap the dugout soil on the lower side of the slope in order to harness the flowing water and facilitate weathering of soil in the pit. Exposure of soil in the pit also prevents spread of pests and diseases due to scorching sun rays. The pits should be filled with mixture of soil and organic manure during the second week of April and keep ready with sufficiently tall seedlings produced either in poly bags or in root trainer nurseries so that planting can be done during the 2nd or 3rd week of April depending on the rainfall.

The tree species suitable for the area considering rainfall, temperature and adaptability is listed below; waterlogged areas are recommended to be planted with species like Neral (*Syzizium cumini*) and Bamboo. Dry areas are to be planted with species like Honge, Bevu, Seetaphal etc.

Dry Deciduous Species			Temp (°C)	Rainfall (mm)
1.	Bevu	<i>Azadiracta indica</i>	21–32	400 –1,200
2.	Tapasi	<i>Holoptelia integrifolia</i>	20-30	500 - 1000
3.	Seetaphal	<i>Anona Squamosa</i>	20-40	400 - 1000
4.	Honge	<i>Pongamia pinnata</i>	20 -50	500–2,500
5.	Kamara	<i>Hardwickia binata</i>	25 -35	400 - 1000
6.	Bage	<i>Albezzia lebbek</i>	20 - 45	500 - 1000
7.	Ficus	<i>Ficus bengalensis</i>	20 - 50	500–2,500
8.	Sisso	<i>Dalbargia Sissoo</i>	20 - 50	500 -2000
9.	Ailanthus	<i>Ailanthus excelsa</i>	20 - 50	500 - 1000
10.	Hale	<i>Wrightia tinctoria</i>	25 - 45	500 - 1000
11.	Uded	<i>Steriospermum chelanoides</i>	25 - 45	500 -2000
12.	Dhupa	<i>Boswella Serrata</i>	20 - 40	500 - 2000
13.	Nelli	<i>Emblia Officinalis</i>	20 - 50	500 -1500
14.	Honne	<i>Pterocarpus marsupium</i>	20 - 40	500 - 2000
Moist Deciduous Species			Temp (°C)	Rainfall (mm)
15.	Teak	<i>Tectona grandis</i>	20 - 50	500-5000
16.	Nandi	<i>Legarstroemia lanceolata</i>	20 - 40	500 - 4000
17.	Honne	<i>Pterocarpus marsupium</i>	20 - 40	500 - 3000
18.	Mathi	<i>Terminalia alata</i>	20 -50	500 - 2000
19.	Shivane	<i>Gmelina arborea</i>	20 -50	500 -2000
20.	Kindal	<i>T.Paniculata</i>	20 - 40	500 - 1500
21.	Beete	<i>Dalbargia latifolia</i>	20 - 40	500 - 1500
22.	Tare	<i>T. belerica</i>	20 - 40	500 - 2000
23.	Bamboo	<i>Bambusa arundinasia</i>	20 - 40	500 - 2500
24.	Bamboo	<i>Dendrocalamus strictus</i>	20 – 40	500 – 2500
25.	Muthuga	<i>Butea monosperma</i>	20 - 40	400 - 1500
26.	Hippe	<i>Madhuca latifolia</i>	20 - 40	500 - 2000
27.	Sandal	<i>Santalum album</i>	20 - 50	400 - 1000
28.	Nelli	<i>Emblia officinalis</i>	20 - 40	500 - 2000
29.	Nerale	<i>Sizyium cumini</i>	20 - 40	500 - 2000
30.	Dhaman	<i>Grevia tilifolia</i>	20 - 40	500 - 2000
31.	Kaval	<i>Careya arborea</i>	20 - 40	500 - 2000
32.	Harada	<i>Terminalia chebula</i>	20 - 40	500 - 2000

References

1. FAO (1976) Framework for Land Evaluation, Food and Agriculture Organization, Rome. 72 pp.
2. FAO (1983) Guidelines for Land Evaluation for Rainfed Agriculture, FAO, Rome, 237 pp.
3. IARI (1971) Soil Survey Manual, All India Soil and Land Use Survey Organization, IARI, New Delhi, 121 pp.
4. Katyal, J.C. and Rattan, R.K. (2003) Secondary and Micronutrients; Research Gap and future needs. Fert. News 48 (4); 9-20.
5. Naidu, L.G.K., Ramamurthy, V., Challa, O., Hegde, R. and Krishnan, P. (2006) Manual Soil Site Suitability Criteria for Major Crops, NBSS Publ. No. 129, NBSS &LUP, Nagpur, 118 pp.
6. Natarajan, A. and Dipak Sarkar (2010) Field Guide for Soil Survey, National Bureau of Soil Survey and Land Use Planning (ICAR), Nagpur, India.
7. Natarajan, A., Rajendra Hegde, Raj, J.N. and Shivananda Murthy, H.G. (2015) Implementation Manual for Sujala-III Project, Watershed Development Department, Bengaluru, Karnataka.
8. Sarma, V.A.K., Krishnan, P. and Budihal, S.L. (1987) Laboratory Manual, Tech. Bull. 23, NBSS &LUP, Nagpur.
9. Sehgal, J.L. (1990) Soil Resource Mapping of Different States of India; Why and How? National Bureau of Soil Survey and Land Use Planning, Nagpur, 49 pp.
10. Shivaprasad, C.R., R.S. Reddy, J. Sehgal and M. Velayuthum (1998) Soils of Karnataka for Optimizing Land Use, NBSS Publ. No. 47b, NBSS & LUP, Nagpur, India.
11. Soil Survey Staff (2006) Keys to Soil Taxonomy, Tenth edition, U.S. Department of Agriculture/ NRCS, Washington DC, U.S.A.
12. Soil Survey Staff (2012) Soil Survey Manual, Handbook No. 18, USDA, Washington DC, USA.

Appendix I
Vojenahlli-2 (1X2d) Microwatershed
Soil Phase Information

Village	Survey NO	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Dadhegalla	RIVER	0.11	Waterbody	Others	Others	Others	Others	Others	Others	Others	Waterbody	Not Available	Others	Others
Koppal	17	1.09	Habitation	Others	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others
Koppal	18	0.001	Habitation	Others	Others	Others	Others	Others	Others	Others	Not Available (NA)	Not Available	Others	Others
Koppal	19	1.82	Habitation	Others	Others	Others	Others	Others	Others	Others	Not Available (NA)	Not Available	Others	Others
Koppal	20	0.47	Habitation	Others	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others
Koppal	21	4.74	Habitation	Others	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others
Koppal	22	5.12	Habitation	Others	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others
Koppal	23	7.35	Habitation	Others	Others	Others	Others	Others	Others	Others	Fallow land (Fl)	Not Available	Others	Others
Koppal	24	3.45	Habitation	Others	Others	Others	Others	Others	Others	Others	Fallow land (Fl)	Not Available	Others	Others
Koppal	25	10.05	DRLmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Habitation+Maize (Hb+Mz)	Not Available	Iles	Graded bunding
Koppal	26	4.14	DRLmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	Iles	Graded bunding
Koppal	27	6.98	DRLmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	Iles	Graded bunding
Koppal	28	5.9	AWDmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Habitation	Not Available	IIIe	Graded bunding
Koppal	29	5.84	DRLmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Fallow land+Drumstick (Fl+Ds)	Not Available	Iles	Graded bunding
Koppal	30	5.7	DRLmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Habitation+Maize (Hb+Mz)	Not Available	Iles	Graded bunding
Koppal	31	7.7	Habitation	Others	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others
Koppal	32	2.64	Habitation	Others	Others	Others	Others	Others	Others	Others	Not Available (NA)	Not Available	Others	Others
Koppal	33	2.33	Habitation	Others	Others	Others	Others	Others	Others	Others	Not Available (NA)	Not Available	Others	Others
Koppal	34	2.18	Habitation	Others	Others	Others	Others	Others	Others	Others	Not Available (NA)	Not Available	Others	Others
Koppal	35	4.23	Habitation	Others	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others
Koppal	36	4.75	Habitation	Others	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others
Koppal	37	4.11	HDHhB1g1	LMU-2	Moderately deep (75-100 cm)	Sandy clay loam	Gravelly (15-35%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Slight	Fallow land (Fl)	Not Available	Iis	Trench cum bunding

Village	Survey NO	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Koppal	38	3.27	Habitation	Others	Others	Others	Others	Others	Others	Others	Fallow land (Fl)	Not Available	Others	Others
Koppal	39	5.69	Habitation	Others	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others
Koppal	40	4.04	Habitation	Others	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others
Koppal	41	4.17	Habitation	Others	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others
Koppal	42	10.5	HDHhB1g 1	LMU-2	Moderately deep (75-100 cm)	Sandy clay loam	Gravelly (15-35%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Slight	Cotton+Maize (Ct+Mz)	Not Available	Iis	Trench cum bunding
Koppal	43	3.3	HDHhB1g 1	LMU-2	Moderately deep (75-100 cm)	Sandy clay loam	Gravelly (15-35%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Slight	Not Available (NA)	Not Available	Iis	Trench cum bunding
Koppal	44	0.14	HDHhB1g 1	LMU-2	Moderately deep (75-100 cm)	Sandy clay loam	Gravelly (15-35%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Slight	Not Available (NA)	Not Available	Iis	Trench cum bunding
Koppal	45	1.43	HDHhB1g 1	LMU-2	Moderately deep (75-100 cm)	Sandy clay loam	Gravelly (15-35%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Slight	Sugarcane (Sc)	Not Available	Iis	Trench cum bunding
Koppal	46	8.25	GDPiB2	LMU-2	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Drumstick+Sugar cane (Ds+Sc)	Not Available	IIles	Trench cum bunding
Koppal	47	0.44	Habitation	Others	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others
Koppal	48	1.34	Habitation	Others	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others
Koppal	49	3.7	Habitation	Others	Others	Others	Others	Others	Others	Others	Habitation	1 Borewell	Others	Others
Koppal	50	4.27	Habitation	Others	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others
Koppal	51	2.21	GDPiB2	LMU-2	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIles	Trench cum bunding
Koppal	52	1.94	GDPiB2	LMU-2	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	1 Borewell	IIles	Trench cum bunding
Koppal	53	2.49	GDPiB2	LMU-2	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Habitation	Not Available	IIles	Trench cum bunding
Koppal	54	9.9	GDPiB2	LMU-2	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIles	Trench cum bunding
Koppal	55	0.07	GDPiB2	LMU-2	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIles	Trench cum bunding
Koppal	57	1.75	GDPiB2	LMU-2	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIles	Trench cum bunding
Koppal	58	2.69	Habitation	Others	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others
Koppal	59	2.8	Habitation	Others	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others
Koppal	60	2.81	Habitation	Others	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others
Koppal	61	0.42	GDPiB2	LMU-2	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Maize+Redgram (Mz+Rg)	Not Available	IIles	Trench cum bunding
Koppal	65	0.001	Habitation	Others	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others
Koppal	106	1.84	BGPmB1	LMU-1	Very deep (>150)	Clay	Non gravelly	High (151-200)	Very gently	Slight	Maize (Mz)	Not	Iis	Graded

Village	Survey NO	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Graveliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
					cm)		(<15%)	mm/m)	sloping (1-3%)			Available		bunding
Koppal	107	1.32	BGPmB1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	High (151-200 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	IIs	Graded bunding
Koppal	686	0.001	BGPmA1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	High (151-200 mm/m)	Nearly level (0-1%)	Slight	Habitation	Not Available	IIs	Graded bunding
Koppal	693	0.33	Habitation	Others	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others
Koppal	694	2.39	Habitation	Others	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others
Koppal	695	1.28	Habitation	Others	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others
Koppal	696	13.68	Habitation	Others	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others
Koppal	699	0.07	Habitation	Others	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others
Koppal	700	1	Habitation	Others	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others
Koppal	701	10.08	BGPmB1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	High (151-200 mm/m)	Very gently sloping (1-3%)	Slight	Habitation	1 Borewell	IIs	Graded bunding
Koppal	702	0.68	BGPmB1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	High (151-200 mm/m)	Very gently sloping (1-3%)	Slight	Not Available (NA)	Not Available	IIs	Graded bunding
Koppal	703	9.92	BGPmB1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	High (151-200 mm/m)	Very gently sloping (1-3%)	Slight	Maize+Paddy (Mz+Pd)	Not Available	IIs	Graded bunding
Koppal	704	4.06	BGPmB1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	High (151-200 mm/m)	Very gently sloping (1-3%)	Slight	Habitation	Not Available	IIs	Graded bunding
Koppal	705	9.31	Habitation	Others	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others
Koppal	706	5.4	Habitation	Others	Others	Others	Others	Others	Others	Others	Habitation	1 Borewell	Others	Others
Koppal	707	10.15	Habitation	Others	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others
Koppal	708	2.45	Habitation	Others	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others
Koppal	709	5.7	KVRmB2g 1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Habitation+Maize (Hb+Mz)	1 Borewell	IIIs	Graded bunding
Koppal	710	3.25	KVRmB2g 1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIIs	Graded bunding
Koppal	711	8.22	KVRmB2g 1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIIs	Graded bunding
Koppal	712	5.43	Habitation	Others	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others
Koppal	713	0.19	KVRmB2g 1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Habitation	Not Available	IIIs	Graded bunding
Koppal	714	7.04	AWDmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIIe	Graded bunding
Koppal	715	4.74	AWDmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIIe	Graded bunding
Koppal	716	3.69	AWDmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIIe	Graded bunding

Village	Survey NO	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Koppal	717	3.52	Habitation	Others	Others	Others	Others	Others	Others	Others	Maize (Mz)	Not Available	Others	Others
Koppal	718	4.15	Habitation	Others	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others
Koppal	719	4.5	Habitation	Others	Others	Others	Others	Others	Others	Others	Fallow land (Fl)	Not Available	Others	Others
Koppal	720	3.77	Habitation	Others	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others
Koppal	721	0.08	Habitation	Others	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others
Koppal	722	0.16	Habitation	Others	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others
Koppal	723	0.7	Habitation	Others	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others
Koppal	725	0.15	Habitation	Others	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others
Ojanahalli	37	1.16	RNKmB1	LMU-3	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	1 Borewell	IIs	Graded bunding
Ojanahalli	39	0.63	KVRmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	IIs	Graded bunding
Ojanahalli	40	1.59	KVRmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	IIs	Graded bunding
Ojanahalli	48	1.27	BGPmB1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	High (151-200 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	IIs	Graded bunding
Ojanahalli	50	1.32	KVRmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	IIs	Graded bunding
Ojanahalli	51	0.001	HDHiB1	LMU-2	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	IIs	Trench cum bunding
Ojanahalli	52	3.11	KVRmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	2 Borewell	IIs	Graded bunding
Ojanahalli	53	7.26	KVRmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	1 Borewell	IIs	Graded bunding
Ojanahalli	54	3.81	BGPmB1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	High (151-200 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	IIs	Graded bunding
Ojanahalli	55	3.35	BGPmB1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	High (151-200 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	IIs	Graded bunding
Ojanahalli	56	2.53	BGPmB1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	High (151-200 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	IIs	Graded bunding
Ojanahalli	57	1.6	BGPmB1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	High (151-200 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	IIs	Graded bunding
Ojanahalli	58	5.76	KVRmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	IIs	Graded bunding
Ojanahalli	59	1.05	BGPmB1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	High (151-200 mm/m)	Very gently sloping (1-3%)	Slight	Habitation	Not Available	IIs	Graded bunding
Ojanahalli	60	8.2	KVRmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	IIs	Graded bunding
Ojanahalli	61	0.5	KVRmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	IIs	Graded bunding

Village	Survey NO	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Graveliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Ojanahalli	62	4.94	KVRmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	IIs	Graded bunding
Ojanahalli	63	0.53	KVRmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	IIs	Graded bunding
Ojanahalli	64	3.66	KVRmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	IIs	Graded bunding
Ojanahalli	65	2.09	KVRmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	IIs	Graded bunding
Ojanahalli	66	3.77	KVRmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	IIs	Graded bunding
Ojanahalli	67	3.57	KVRmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	IIs	Graded bunding
Ojanahalli	68	4.44	BGPmB1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	High (151-200 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	IIs	Graded bunding
Ojanahalli	69	2.01	BGPmB1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	High (151-200 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	IIs	Graded bunding
Ojanahalli	70	4.24	BGPmB1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	High (151-200 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	IIs	Graded bunding
Ojanahalli	73	1.73	BGPmB1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	High (151-200 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	IIs	Graded bunding
Ojanahalli	74	1.59	BGPmB1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	High (151-200 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	IIs	Graded bunding
Ojanahalli	77	4.06	BGPmB1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	High (151-200 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	IIs	Graded bunding
Ojanahalli	78	6.88	KVRmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	IIs	Graded bunding
Ojanahalli	79	6.5	KVRmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	IIs	Graded bunding
Ojanahalli	80	6.87	KVRmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	1 Borewell	IIs	Graded bunding
Ojanahalli	81	8.69	KVRmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	IIs	Graded bunding
Ojanahalli	82	9.75	KVRmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	IIs	Graded bunding
Ojanahalli	83	0.93	KVRmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	1 Borewell	IIs	Graded bunding
Ojanahalli	84	13.2	DRLmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow (Cf)	1 Borewell	IIs	Graded bunding
Ojanahalli	85	1.05	DRLmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIs	Graded bunding
Ojanahalli	86	0.15	DRLmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow (Cf)	Not Available	IIs	Graded bunding
Ojanahalli	87	4.65	DRLmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	1 Borewell	IIs	Graded bunding
Ojanahalli	88	8.41	DRLmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIs	Graded bunding
Ojanahalli	89	8.19	DRLmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIs	Graded bunding

Village	Survey NO	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Graveliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Ojanahalli	90	2.86	DRLmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	Iles	Graded bunding
Ojanahalli	91	4.19	DRLmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	Iles	Graded bunding
Ojanahalli	92	7.77	RNKmB1	LMU-3	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	IIs	Graded bunding
Ojanahalli	93	2.32	RNKmB1	LMU-3	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	IIs	Graded bunding
Ojanahalli	94	2.79	RNKmB1	LMU-3	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	IIs	Graded bunding
Ojanahalli	95	5.37	RNKmB1	LMU-3	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	IIs	Graded bunding
Ojanahalli	96	0.56	RNKmB1	LMU-3	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	IIs	Graded bunding
Ojanahalli	97	0.13	RNKmB1	LMU-3	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	IIs	Graded bunding
Ojanahalli	102	5.68	RNKmB1	LMU-3	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	IIs	Graded bunding
Ojanahalli	103	4.68	RNKmB1	LMU-3	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	IIs	Graded bunding
Ojanahalli	104	6.5	DRLmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	Iles	Graded bunding
Ojanahalli	114	0.001	RNKmB1	LMU-3	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Slight	Maize (Mz)	Not Available	IIs	Graded bunding
Ojanahalli	115	6.65	DRLmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	Iles	Graded bunding
Ojanahalli	116	0.42	AWDmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIIe	Graded bunding
Ojanahalli	117	1.61	AWDmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIIe	Graded bunding
Ojanahalli	118	3.05	AWDmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIIe	Graded bunding
Ojanahalli	119	3.23	AWDmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIIe	Graded bunding
Ojanahalli	120	3.13	AWDmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIIe	Graded bunding
Ojanahalli	121	8.42	DRLmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	Iles	Graded bunding
Ojanahalli	122	7.91	AWDmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Coconut (Cn)	1 Borewell	IIIe	Graded bunding
Ojanahalli	123	7.44	AWDmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIIe	Graded bunding
Ojanahalli	124	0.01	AWDmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Coconut (Cn)	Not Available	IIIe	Graded bunding
Ojanahalli	126	2.41	AWDmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIIe	Graded bunding
Ojanahalli	127	1.74	AWDmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIIe	Graded bunding

Village	Survey NO	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Ojanahalli	146	0.01	KVRiB2g1	LMU-1	Deep (100-150 cm)	Sandy clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIIes	Graded bunding
Yathnatti	RIVER	4.78	Waterbody	Others	Others	Others	Others	Others	Others	Others	Waterbody	Not Available	Others	Others

Appendix II
Vojenahlli-2 (1X2d) Microwatershed
Soil Fertility Information

Village	Survey No	Soil Reaction	Salinity	Organic Carbon	Available Phosphorus	Available Potassium	Available Sulphur	Available Boron	Available Iron	Available Manganese	Available Copper	Available Zinc
Dadhegalla	RIVER	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Koppal	17	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Koppal	18	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Koppal	19	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Koppal	20	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Koppal	21	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Koppal	22	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Koppal	23	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Koppal	24	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Koppal	25	Moderately alkaline (pH 7.8 - 8.4)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	Medium (0.5 - 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Koppal	26	Moderately alkaline (pH 7.8 - 8.4)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	Medium (0.5 - 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Koppal	27	Moderately alkaline (pH 7.8 - 8.4)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	Medium (0.5 - 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Koppal	28	Strongly alkaline (pH 8.4 - 9.0)	Non saline (<2 dsm)	Low (< 0.5 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	Medium (0.5 - 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Koppal	29	Strongly alkaline (pH 8.4 - 9.0)	Non saline (<2 dsm)	Low (< 0.5 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	Medium (0.5 - 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Koppal	30	Strongly alkaline (pH 8.4 - 9.0)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	Medium (0.5 - 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Koppal	31	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Koppal	32	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Koppal	33	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Koppal	34	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Koppal	35	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Koppal	36	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Koppal	37	Strongly alkaline (pH 8.4 - 9.0)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	Medium (0.5 - 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Koppal	38	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Koppal	39	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Koppal	40	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Koppal	41	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others

Village	Survey No	Soil Reaction	Salinity	Organic Carbon	Available Phosphorus	Available Potassium	Available Sulphur	Available Boron	Available Iron	Available Manganese	Available Copper	Available Zinc
Koppal	699	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Koppal	700	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Koppal	701	Moderately alkaline (pH 7.8 - 8.4)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	High (> 337 kg/ha)	High (> 20 ppm)	Medium (0.5 - 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Koppal	702	Moderately alkaline (pH 7.8 - 8.4)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	High (> 337 kg/ha)	High (> 20 ppm)	Medium (0.5 - 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Koppal	703	Moderately alkaline (pH 7.8 - 8.4)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	High (> 20 ppm)	Medium (0.5 - 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Koppal	704	Strongly alkaline (pH 8.4 - 9.0)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	Medium (10 - 20 ppm)	Medium (0.5 - 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Koppal	705	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Koppal	706	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Koppal	707	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Koppal	708	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Koppal	709	Moderately alkaline (pH 7.8 - 8.4)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	High (> 20 ppm)	Medium (0.5 - 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Koppal	710	Moderately alkaline (pH 7.8 - 8.4)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	High (> 20 ppm)	Medium (0.5 - 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Koppal	711	Strongly alkaline (pH 8.4 - 9.0)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	High (> 20 ppm)	Medium (0.5 - 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Koppal	712	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Koppal	713	Strongly alkaline (pH 8.4 - 9.0)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	High (> 20 ppm)	High (> 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Koppal	714	Strongly alkaline (pH 8.4 - 9.0)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	High (> 20 ppm)	High (> 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Koppal	715	Strongly alkaline (pH 8.4 - 9.0)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	High (> 20 ppm)	Medium (0.5 - 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Koppal	716	Strongly alkaline (pH 8.4 - 9.0)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	High (> 20 ppm)	Medium (0.5 - 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Koppal	717	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Koppal	718	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Koppal	719	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Koppal	720	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Koppal	721	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Koppal	722	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Koppal	723	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Koppal	725	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Ojanahalli	37	Moderately alkaline (pH 7.8 - 8.4)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	Medium (10 - 20 ppm)	Low (< 0.5 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)

Appendix III
Vojenahlli-2 (1X2d) Microwatershed
Soil Suitability Information

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Jasmine	Bhendi	Brinjal	Crossandra	Drumstick	Mulberry	Onion	
Dadhegalla	RIVER	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	
Koppal	17	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	
Koppal	18	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	
Koppal	19	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	
Koppal	20	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	
Koppal	21	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	
Koppal	22	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	
Koppal	23	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	
Koppal	24	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	
Koppal	25	S3rz	S2tz	S3tz	S2nz	S3tz	S2rz	S3rz	S2rz	S2rz	S2rz	S3rz	S2tz	S3tz	S2z	N1tz	S3rz	S2rz	S3tz	S3tz	S3tz	S2tz	S2tz	S2rt	S2tz	S3tz	S2tz	S2tz	S2z	S2rz	S2tz	S3tz	
Koppal	26	S3rz	S2tz	S3tz	S2nz	S3tz	S2rz	S3rz	S2rz	S2rz	S2rz	S3rz	S2tz	S3tz	S2z	N1tz	S3rz	S2rz	S3tz	S3tz	S3tz	S2tz	S2tz	S2rt	S2tz	S3tz	S2tz	S2tz	S2z	S2rz	S2tz	S3tz	
Koppal	27	S3rz	S2tz	S3tz	S2nz	S3tz	S2rz	S3rz	S2rz	S2rz	S2rz	S3rz	S2tz	S3tz	S2z	N1tz	S3rz	S2rz	S3tz	S3tz	S3tz	S2tz	S2tz	S2rt	S2tz	S3tz	S2tz	S2tz	S2z	S2rz	S2tz	S3tz	
Koppal	28	S3tz	S2tz	S3tz	S2z	S3tz	S2z	S2tz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2tz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S3tz	S3tz	
Koppal	29	S3rz	S2tz	S3tz	S2nz	S3tz	S2rz	S3rz	S2rz	S2rz	S2rz	S3rz	S2tz	S3tz	S2z	N1tz	S3rz	S2rz	S3tz	S3tz	S3tz	S2tz	S2tz	S2rt	S2tz	S3tz	S2tz	S2tz	S2z	S2rz	S2tz	S3tz	
Koppal	30	S3rz	S2tz	S3tz	S2nz	S3tz	S2rz	S3rz	S2rz	S2rz	S2rz	S3rz	S2tz	S3tz	S2z	N1tz	S3rz	S2rz	S3tz	S3tz	S3tz	S2tz	S2tz	S2rt	S2tz	S3tz	S2tz	S2tz	S2z	S2rz	S2tz	S3tz	
Koppal	31	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	
Koppal	32	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	
Koppal	33	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	
Koppal	34	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	
Koppal	35	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	
Koppal	36	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	
Koppal	37	S3rg	S3g	S2rg	S3g	S2rg	S3rg	S3rg	S2rg	S3g	S3rg	S3g	S2rg	S2rg	S2rg	S2rg	S2rg	S2rg	S2g	S3g	S3g	S3g	S3g	S2rg	S2g	S3g	S3g	S3g	S3g	S3g	S3g	S2g	S3g
Koppal	38	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	
Koppal	39	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Jasmine	Bhendi	Brinjal	Crossandra	Drumstick	Mulberry	Onion	
Koppal	723	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Koppal	725	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Ojanahalli	37	N1rz	S2tz	S3rz	S2rz	S3tz	S2rz	N1rz	S3rz	S2rz	S3rz	S3rz	S2rz	S3tz	S2rz	N1tz	S3tz	S3rz	S3tz	S3tz	S3tz	S2rz	S2rz	S3rz	S2tz	S2rz	S2rt	S2rt	S3rz	S3rz	S3rz	S3tz	
Ojanahalli	39	S2rz	S2tz	S3tz	S2z	S3tz	S2z	S2rz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2rz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	
Ojanahalli	40	S2rz	S2tz	S3tz	S2z	S3tz	S2z	S2rz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2rz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	
Ojanahalli	48	S3tz	S2tz	S3tz	S2z	S3tz	S2z	S2tz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2tz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	
Ojanahalli	50	S2rz	S2tz	S3tz	S2z	S3tz	S2z	S2rz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2rz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	
Ojanahalli	51	S3rg	S3g	S2rg	S3g	S2rg	S3rg	S3rg	S2rg	S3g	S3rg	S3g	S2rg	S2rg	S2rg	S2rg	S2rg	S2rg	S2g	S3g	S3g	S3g	S3g	S2rg	S2g	S3g	S3g	S3g	S3g	S3g	S3g	S2g	S3g
Ojanahalli	52	S2rz	S2tz	S3tz	S2z	S3tz	S2z	S2rz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2rz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	
Ojanahalli	53	S2rz	S2tz	S3tz	S2z	S3tz	S2z	S2rz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2rz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	
Ojanahalli	54	S3tz	S2tz	S3tz	S2z	S3tz	S2z	S2tz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2tz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	
Ojanahalli	55	S3tz	S2tz	S3tz	S2z	S3tz	S2z	S2tz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2tz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	
Ojanahalli	56	S3tz	S2tz	S3tz	S2z	S3tz	S2z	S2tz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2tz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	
Ojanahalli	57	S3tz	S2tz	S3tz	S2z	S3tz	S2z	S2tz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2tz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	
Ojanahalli	58	S2rz	S2tz	S3tz	S2z	S3tz	S2z	S2rz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2rz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	
Ojanahalli	59	S3tz	S2tz	S3tz	S2z	S3tz	S2z	S2tz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2tz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	
Ojanahalli	60	S2rz	S2tz	S3tz	S2z	S3tz	S2z	S2rz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2rz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	
Ojanahalli	61	S2rz	S2tz	S3tz	S2z	S3tz	S2z	S2rz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2rz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	
Ojanahalli	62	S2rz	S2tz	S3tz	S2z	S3tz	S2z	S2rz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2rz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	
Ojanahalli	63	S2rz	S2tz	S3tz	S2z	S3tz	S2z	S2rz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2rz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	
Ojanahalli	64	S2rz	S2tz	S3tz	S2z	S3tz	S2z	S2rz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2rz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	
Ojanahalli	65	S2rz	S2tz	S3tz	S2z	S3tz	S2z	S2rz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2rz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	
Ojanahalli	66	S2rz	S2tz	S3tz	S2z	S3tz	S2z	S2rz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2rz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	
Ojanahalli	67	S2rz	S2tz	S3tz	S2z	S3tz	S2z	S2rz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2rz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	
Ojanahalli	68	S3tz	S2tz	S3tz	S2z	S3tz	S2z	S2tz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2tz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	
Ojanahalli	69	S3tz	S2tz	S3tz	S2z	S3tz	S2z	S2tz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2tz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	
Ojanahalli	70	S3tz	S2tz	S3tz	S2z	S3tz	S2z	S2tz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2tz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Jasmine	Bhendi	Brinjal	Crossandra	Drumstick	Mulberry	Onion
Ojanahalli	115	S3rz	S2tz	S3tz	S2nz	S3tz	S2rz	S3rz	S2rz	S2rz	S2rz	S3rz	S2tz	S3tz	S2z	N1tz	S3rz	S2rz	S3tz	S3tz	S3tz	S2tz	S2tz	S2rt	S2tz	S3tz	S2tz	S2tz	S2z	S2rz	S2tz	S3tz
Ojanahalli	116	S3tz	S2tz	S3tz	S2z	S3tz	S2z	S2tz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2tz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S3tz	S3tz
Ojanahalli	117	S3tz	S2tz	S3tz	S2z	S3tz	S2z	S2tz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2tz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S3tz	S3tz
Ojanahalli	118	S3tz	S2tz	S3tz	S2z	S3tz	S2z	S2tz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2tz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S3tz	S3tz
Ojanahalli	119	S3tz	S2tz	S3tz	S2z	S3tz	S2z	S2tz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2tz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S3tz	S3tz
Ojanahalli	120	S3tz	S2tz	S3tz	S2z	S3tz	S2z	S2tz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2tz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S3tz	S3tz
Ojanahalli	121	S3rz	S2tz	S3tz	S2nz	S3tz	S2rz	S3rz	S2rz	S2rz	S2rz	S3rz	S2tz	S3tz	S2z	N1tz	S3rz	S2rz	S3tz	S3tz	S3tz	S2tz	S2tz	S2rt	S2tz	S3tz	S2tz	S2tz	S2z	S2rz	S2tz	S3tz
Ojanahalli	122	S3tz	S2tz	S3tz	S2z	S3tz	S2z	S2tz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2tz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S3tz	S3tz
Ojanahalli	123	S3tz	S2tz	S3tz	S2z	S3tz	S2z	S2tz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2tz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S3tz	S3tz
Ojanahalli	124	S3tz	S2tz	S3tz	S2z	S3tz	S2z	S2tz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2tz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S3tz	S3tz
Ojanahalli	126	S3tz	S2tz	S3tz	S2z	S3tz	S2z	S2tz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2tz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S3tz	S3tz
Ojanahalli	127	S3tz	S2tz	S3tz	S2z	S3tz	S2z	S2tz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2tz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S3tz	S3tz
Ojanahalli	146	S2rz	S2tz	S3tz	S2z	S3tz	S2z	S2rz	S2z	S2gz	S2z	S2gt	S2tz	S3tz	S2z	N1tz	S2rz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz
Yathnatti	RIVER	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others

PART-B

SOCIO-ECONOMIC STATUS OF FARM HOUSEHOLDS

CONTENTS

1.	Salient findings of the survey	1-5
2.	Introduction	7
3	Methodology	9
4	Salient features of the survey	11-31
5	Summary	33-37

LIST OF TABLES

1	Households sampled for socio economic survey	11
2	Population characteristics	11
3	Age wise classification of household members	11
4	Education level of household members	12
5	Occupation of household heads	12
6	Occupation of family members	12
7	Institutional participation of household members	13
8	Type of house owned by households	13
9	Durable assets owned by households	13
10	Average value of durable assets owned by households	13
11	Farm implements owned by households	14
12	Average value of farm implements	14
13	Livestock possession by households	15
14	Average labour availability	15
15	Adequacy of hired labour	15
16	Distribution of land (ha)	16
17	Average land value (Rs./ha)	16
18	Status of bore wells	16
19	Source of irrigation	16
20	Depth of water	16
21	Irrigated area (ha)	16
22	Cropping pattern	17
23	Cropping intensity	17
24	Possession of bank account and saving	17
25	Borrowing status	17
26	Cost of cultivation of Bajra	18
27	Cost of cultivation of Groundnut	19
28	Cost of cultivation of Maize	20
29	Cost of cultivation of Navane	21
30	Cost of cultivation of Onion	22
31	Cost of cultivation of Red gram	23

32	Cost of cultivation of Sunflower	24
33	Cost of cultivation of Watermelon	25
34	Adequacy of fodder	26
35	Annual gross income	26
36	Average annual expenditure	26
37	Horticulture species grown	27
38	Forest species grown	27
39	Average additional investment capacity	27
40	Marketing of the agricultural produce	27
41	Marketing channels used for sale of agricultural produce	28
42	Mode of transport of agricultural produce	28
43	Incidence of soil and water erosion problems	28
44	Interest towards soil testing	28
45	Usage pattern of fuel for domestic use	29
46	Source of drinking water	29
47	Source of light	29
48	Existence of sanitary toilet facility	29
49	Possession of public distribution system(PDS) card	29
50	Participation in NREGA programme	30
51	Adequacy of food items	30
52	Response on inadequacy of food items	30
53	Response on market surplus of food items	31
54	Farming constraints experienced	31

SALIENT FINDINGS OF THE SURVEY

- ❖ *The data indicated that there were 106 (54.64%) men and 88 (45.36%) women among the sampled households.*
- ❖ *The average family size of landless farmers' was 4, marginal farmers' was 5, small farmers' was 6.6, semi medium farmers' was 4.7 and medium farmers' was 5.6.*
- ❖ *The data indicated that, 43 (22.16%) people were in 0-15 years of age, 82 (42.27%) were in 16-35 years of age, 55 (28.35%) were in 36-60 years of age and 14 (7.22%) were above 61 years of age.*
- ❖ *The results indicated that Vojenahalli-2 had 27.32 per cent illiterates, 29.38 per cent of them had primary school education, 7.73 per cent of them had middle school education, 21.13 per cent of them had high school education, 9.28 per cent of them had PUC education and 2.58 per cent of them had degree.*
- ❖ *The results indicate that, 68.57 per cent of the household heads were practicing agriculture, 28.57 per cent of the household heads were agricultural labourers and 2.86 per cent of the household heads were trade and business and housewives.*
- ❖ *The results indicate that agriculture was the major occupation for 25.77 per cent of the household members, 39.18 per cent were agricultural labourers, 0.52 per cent were private service and trade and business, 24.74 per cent were student, 5.67 per cent were housewives and 2.58 per cent were children.*
- ❖ *The results show that, 100 per cent of the population in the micro watershed has not participated in any local institutions.*
- ❖ *The results indicate that 77.14 per cent of the households possess katcha house, 17.14 per cent of them possess pucca/RCC house and 5.71 per cent of them possess semi pucca house.*
- ❖ *The results show that 85.71 per cent of the households possess TV, 77.14 per cent of them possess mixer/grinder, 25.71 per cent of them possess bicycle, 37.14 per cent of the households possess motor cycle and 97.14 per cent of the households possess mobile phones.*
- ❖ *The results show that the average value of television was Rs. 7,066, mixer grinder was Rs. 2,344, bicycle was Rs. 2,711, motor cycle was Rs. 41,923 and mobile phone was Rs. 2,220.*
- ❖ *About 20 per cent of the households possess bullock cart, 45.71 per cent of them possess plough, 5.71 per cent of them possess seed/fertilizer drill and tractor, 2.86 per cent of them possess irrigation pump, power tiller and harvester, 20 per cent of them possess sprayer and 57.14 per cent of them possess weeder.*
- ❖ *The results show that the average value of bullock cart was Rs. 13,625, plough was Rs. 3,381, seed/fertilizer drill was Rs.35,000, irrigation pump was Rs.75,000, power tiller was Rs. 50,000, tractor was Rs.500,000, sprayer was Rs.3,285, weeder was Rs. 70 and harvester was Rs.80.*

- ❖ *The results indicate that, 37.14 per cent of the households possess bullocks and local cow, 2.86 per cent of the households possess crossbreed cow and 8.57 per cent of the households possess local cow.*
- ❖ *The results indicate that, average own labour men available in the micro watershed was 1.56, average own labour (women) available was 1.41, average hired labour (men) available was 12 and average hired labour (women) available was 12.18.*
- ❖ *The results indicate that 45.71 per cent of the households opined that the hired labour was adequate and 48.57 per cent of the households opined that hired labour was inadequate.*
- ❖ *The results indicate that, households of the Vojenahalli-2 micro-watershed possess 38.66 ha (76.48%) of dry land and 11.89 ha (23.52%) of irrigated land. Marginal farmers possess 5.33 ha (92.95%) of dry land and 0.40 ha (7.05%) of irrigated land. Small farmers possess 16.51 ha (96.8%) of dry land and 0.55 ha (3.20%) irrigated land. Semi medium farmers possess 10.34 ha (65.16%) of dry land and 5.53 ha (34.84%) of irrigated land. Medium farmers possess 6.48 ha (54.50%) of dry land and 5.41 ha (45.50 %) of irrigated land.*
- ❖ *The results indicate that, the average value of dry land was Rs. 307,715.66 and the average value of irrigated land was Rs. 420,497.11. In case of marginal famers, the average land value was Rs. 468,512.90 for dry land the average land value was Rs. 2,470,000. In case of small famers, the average land value was Rs. 272,493.26 for dry land, Rs.182,962.96 for irrigated land. In case of semi medium famers, the average land value was Rs. 280,352.26 for dry land and Rs. 506,295.75 for irrigated land. In case of medium farmers, the average land value was Rs. 308,750 and the average value was Rs. 203,368.27 for irrigated land.*
- ❖ *The results indicate that, there were 1 de-functioning and 7 functioning bore wells in the micro watershed.*
- ❖ *The results indicate that, bore well was the major irrigation source in the micro water shed for 20 per cent of the farmers.*
- ❖ *The results indicate that, the depth of bore well was found to be 23.33 meters.*
- ❖ *The results indicate that marginal and small, semi medium and medium farmers had an irrigated area of 0.40 ha, 5.53 ha and 2.11 ha respectively.*
- ❖ *The results indicate that, farmers have grown bajra (3.47 ha), groundnut (1.21 ha), maize (37.21 ha), navane (2.02 ha), onion (0.81 ha). Red gram (1.69 ha), sunflower (4.57 ha) and watermelon (0.40 ha). Marginal farmers have grown bajra, and maize. Small farmers have grown bajra, maize, navane, red gram and sunflower. Semi medium farmers have grown maize, onion, red gram and watermelon. Medium farmers have grown groundnut, maize and sunflower.*
- ❖ *The results indicate that, the cropping intensity in Vojenahalli-2 micro-watershed was found to be 78.89 per cent.*

- ❖ *The results indicate that, 42.86 per cent of the households have bank account and savings.*
- ❖ *The results indicate that, 42.86 per cent of the households have availed credit from different sources.*
- ❖ *The results indicate that, the total cost of cultivation for bajra was Rs. 42621.58. The gross income realized by the farmers was Rs. 19878.02. The net income from bajra cultivation was Rs. -22743.56. Thus the benefit cost ratio was found to be 1:0.47.*
- ❖ *The total cost of cultivation for groundnut was Rs. 30409.08. The gross income realized by the farmers was Rs. 55986.67. The net income from groundnut cultivation was Rs. 25577.58. Thus the benefit cost ratio was found to be 1:1.84.*
- ❖ *The total cost of cultivation for maize was Rs. 36307.28. The gross income realized by the farmers was Rs. 36634.06. The net income from maize cultivation was Rs. 326.78. Thus the benefit cost ratio was found to be 1:1.01.*
- ❖ *The total cost of cultivation for maize was Rs. 36307.28. The gross income realized by the farmers was Rs. 36634.06. The net income from maize cultivation was Rs. 326.78. Thus the benefit cost ratio was found to be 1:1.01.*
- ❖ *The total cost of cultivation for onion was Rs. 30525.33. The gross income realized by the farmers was Rs. 37050. The net income from onion cultivation was Rs. 6524.67. Thus the benefit cost ratio was found to be 1:1.21.*
- ❖ *The total cost of cultivation for red gram was Rs. 35775.70. The gross income realized by the farmers was Rs. 20601.84. The net income from red gram cultivation was Rs. -15173.86. Thus the benefit cost ratio was found to be 1:0.58.*
- ❖ *The total cost of cultivation for sunflower was Rs. 32563.90. The gross income realized by the farmers was Rs. 30187.33. The net income from sunflower cultivation was Rs. -2376.57. Thus the benefit cost ratio was found to be 1:0.93.*
- ❖ *The total cost of cultivation for watermelon was Rs. 228916.72. The gross income realized by the farmers was Rs. 185250.00. The net income from watermelon cultivation was Rs. -43666.72. Thus the benefit cost ratio was found to be 1:0.81.*
- ❖ *The results indicate that, 17.14 per cent of the households opined that dry fodder was adequate, 14.71 per cent of the households opined that green fodder was adequate, 31.43 per cent opined that dry fodder was inadequate and 2.86 per cent opined that green fodder was inadequate.*
- ❖ *The results indicate that the annual gross income was Rs. 53,000 for landless farmers, for marginal farmers it was Rs. 38,537.50, for small farmers it was Rs. 62,961.54, for semi medium farmers it was Rs. 80,250 and for medium farmers it was Rs. 120,333.33.*
- ❖ *The results indicate that the average annual expenditure is Rs. 6,160.64. For landless households it was Rs. 11,555.56, for marginal farmers it was Rs 4,700, for*

small farmers it was Rs. 4,043.39, for semi medium farmers it was Rs. 4,640.63 and for medium farmers it was Rs. 17,888.89.

- ❖ *The results indicate that, sampled households have grown 2 coconut trees in their field. The results indicate that, households have planted 41 neem, 2 tamarind and 4 banyan trees in their field. Also, the households have planted 5 neem, trees in their backyard.*
- ❖ *The results indicated that, households have an average investment capacity of Rs. 3,200 for land development, and Rs. 1,000 for improved crop production.*
- ❖ *The results indicated that, bajra, groundnut, navane, onion, red gram, sunflower and watermelon were sold to the extent of 100 per cent and maize was sold to the extent of 95.58 per cent.*
- ❖ *The results indicated that, about 48.57 per cent of the farmers sold their produce to local/village merchants, 62.86 per cent of the farmers sold their produce to regulated market and 5.71 per cent of the farmers sold their produce to cooperative marketing society.*
- ❖ *The results indicated that, 114.29 per cent of them used tractor as a mode of transportation.*
- ❖ *The results indicated that, 60 per cent of the households have experienced soil and water erosion problems in the farm.*
- ❖ *The results indicated that, 74.29 per cent have shown interest in soil test.*
- ❖ *The results indicated that, 100 per cent of the households used firewood as a source of fuel.*
- ❖ *The results indicated that, piped supply was the major source of drinking water for 51.43 per cent of the households, bore well was the source of drinking water for 45.71 per cent of the households and lake/tank was the source of drinking water for 2.86 per cent of the households in micro watershed.*
- ❖ *Electricity was the major source of light for 100 per cent of the households in micro watershed.*
- ❖ *The results indicated that, 31.43 per cent of the households possess sanitary toilet facility.*
- ❖ *The results indicated that, 2.86 per cent of the sampled households possessed BPL card and 97.14 per cent of the sampled households possessed BPL card.*
- ❖ *The results indicated that, 42.86 per cent of the households participated in NREGA programme.*
- ❖ *The results indicated that, cereals were adequate for 94.29 per cent of the households, pulses were adequate for 57.14 per cent, oilseeds and fruits were adequate for 11.43 per cent, vegetables were adequate for 8.57, milk was adequate for 51.43 per cent, eggs were adequate for 48.57 per cent and meat were adequate for 45.71 per cent.*

- ❖ *The results indicated that, cereals were in adequate for 5.71 per cent of the households, pulses were inadequate for 42.86 per cent, oilseeds and fruits were inadequate for 85.71 per cent, vegetables were inadequate for 88.57 per cent, milk was inadequate for 42.86 per cent, eggs were inadequate for 40 per cent and meat was inadequate for 2.86 per cent of the households.*
- ❖ *The results indicated that, oilseeds and vegetables were market surplus for 2.86 per cent of the households.*
- ❖ *The results indicated that, lower fertility status of the soil, high rate of interest on credit and lack of transport for safe transport of the agricultural produce to the market was the constraint experienced by 45.71 per cent of the households, wild animal menace on farm field and less rainfall (57.14%), frequent incidence of pest and diseases (68.57%), inadequacy of irrigation water (54.29%), high cost of fertilizers and plant protection chemicals (51.43%), low price for the agricultural commodities and lack of marketing facilities in the area (20%), inadequate extension services (8.57%), and Source of Agri-technology information (42.86%).*

INTRODUCTION

Soil and water are the two precious natural resources which are essential for crop production and existence of life on earth. Rainfed agriculture is under severe stress due to various constraints related to agriculture like uneven and erratic distribution of rainfall, indiscriminate use of fertilizers, chemicals and pesticides, adoption of improper land management practices, soil erosion, decline in soil fertility, decline in ground water resources leading to low crop productivity. The area under rainfed agriculture has to be managed effectively using the best available practices to enhance the production of food, fodder and fuel. This is possible if the land resources are characterized at each parcel of land through detailed land resource inventory using the best available techniques of remote sensing, GPS and GIS. The watershed development programs are aimed at the sustainable distribution of its resources and the process of creating and implementing plans, programs, and projects to sustain and enhance watershed functions that affect the plant, animal and human communities within a watershed boundary.

World Bank funded KWDP II, SUJALA III project was implemented in with Broad objective of demonstrating more effective watershed management through greater integration of programmes related to rain-fed agriculture, innovative and science based approaches and strengthen institutional capacities and If successful, it is expected that the systems and tools could be mainstreamed into the overall IWMP in the State of Karnataka and in time, throughout other IWMP operations in India. With this background the socio-economic survey has been carried out with following specific objectives:

1. To understand the demographic features of the households in the micro-watershed
2. To understand the extent of family labour available and additional employment opportunities available within the village.
3. To know the status of assets of households in the micro-watershed for suggesting possible improvements.
4. To study the cropping pattern, cropped area and productivity levels of different households in micro-watershed.
5. To determine the type and extent of livestock owned by different categories of HHs
6. Availability of fodder and level of livestock management.

Scope and importance of survey

Survey helps in identification of different socio-economic and resource use-patterns of farmers at the Micro watershed. Household survey provides demographic features, labour force, and levels of education; land ownership and asset position (including livestock and other household assets) of surveyed households; and cropping patterns, input intensities, and average crop yields from farmers' fields. It also discusses crop utilization and the degree of commercialization of production in the areas; farmers' access to and utilization of credit from formal and informal sources; and the level of adoption and use of soil, water, and pest management technologies.

METHODOLOGY

The description of the methods, components selected for the survey and procedures followed in conducting the baseline survey are furnished under the following heads.

Description of the study area

Koppal district is an administrative district in the state of Karnataka in India. In the past Koppal was referred to as 'Kopana Nagara'. Koppal, now a district headquarters is ancient Kopana a major holy place of the Jainas. The district occupies an area of 7,190 km² and has a population of 1,196,089, which 16.58% were urban as of 2001. The Koppal district was formed after split of Raichur district.

Geographers are very particular about the physiography or relief of a region. It plays a very important role in the spatial analysis of agricultural situation of the study area. The undulating topography with black cotton soil shrips, cut across by numerous nalas or streams is the major characteristic feature of the study region. Three physiographic divisions have made considering the local conditions of landforms and crops grown in the district. On the basis of physiography, Koppal district can be divided into three major divisions. They are (a) Koppal & Yelburga plateau, (b) Maidan division, (c) Tungabhadra valley. The district is part of Krishna basin the main streams draining the area are Maskinala, Ilkal-nadi and Hirenala. These are Ephemeral in nature, these come under Tungabhadra sub-basin. The drainage exhibit dentritic to subdentritic with drainage density varies from 1.4 to 7.0kms/sq.km.

According to the 2011 census Koppal district has a population of 1,391,292, roughly equal to the nation of Swaziland or the US state of Hawaii. This gives it a ranking of 350th in India (out of a total of 640). The district has a population density of 250 inhabitants per square kilometre (650/sq mi). Its population growth rate over the decade 2001-2011 was 16.32%. Koppal has a sex ratio of 983 females for every 1000 males, and a literacy rate of 67.28%.

Description of the micro watershed

Vojenahalli-2 micro-watershed in Bhagyanagara sub-watershed (Koppal taluk and district) is located in between 15^o22'57.185'' to 15^o 21'24.377'' North latitudes and 76^o 10'36.16'' to 76^o8'1.309'' East longitudes, covering an area of about 576.13 ha, bounded by Ojanahalli and Koppla villages.

Methodology followed in assessing socio-economic status of households

In order to assess the socio-economic condition of the farmers in the watershed a comprehensive questionnaire was prepared. Major components such as demographic conditions, migration details, food consumption and family expenditure pattern, material possession, land holding, land use management, cropping pattern, cost of cultivation of crops, livestock management. The statistical components such as frequency and percentage were used to analyze the data. About 35 households located in the micro-watershed were interviewed for the survey.

SALIENT FEATURES OF THE SURVEY

Households sampled for socio-economic survey: The data on households sampled for socio economic survey in Vojenahalli-2 micro-watershed is presented in Table 1 and it indicated that 35 farmers were sampled in Vojenahalli-2 micro-watershed among them 3 (8.57%) were landless and medium farmers, 8 (22.86%) were marginal farmers and semi medium farmers and 13 (37.14%) were small farmers.

Table 1: Households sampled for socio economic survey in Vojenahalli-2 micro-watershed

Sl.No.	Particulars	LL (3)		MF (8)		SF (13)		SMF (8)		MDF (3)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Farmers	3	8.57	8	22.86	13	37.14	8	22.86	3	8.57	35	100.00

Population characteristics: The population characteristics of households sampled for socio-economic survey in Vojenahalli-2 micro-watershed is presented in Table 2. The data indicated that there were 106 (54.64%) men and 88 (45.36%) women among the sampled households. The average family size of landless farmers' was 4, marginal farmers' was 5, small farmers' was 6.6, semi medium farmers' was 4.7 and medium farmers' was 5.6.

Table 2: Population characteristics of Vojenahalli-2 micro-watershed

Sl.No.	Particulars	LL (12)		MF (40)		SF (87)		SMF (38)		MDF (17)		All (194)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Men	5	41.67	22	55.00	46	52.87	20	52.63	13	76.47	106	54.64
2	Women	7	58.33	18	45.00	41	47.13	18	47.37	4	23.53	88	45.36
	Total	12	100.00	40	100.00	87	100.00	38	100.00	17	100.00	194	100.00
	Average	4		5		6.6		4.7		5.6		5.5	

Age wise classification of population: The age wise classification of household members in Vojenahalli-2 micro-watershed is presented in Table 3. The data indicated that, 43 (22.16%) people were in 0-15 years of age, 82 (42.27%) were in 16-35 years of age, 55 (28.35%) were in 36-60 years of age and 14 (7.22%) were above 61 years of age.

Table 3: Age wise classification of household members in Vojenahalli-2 micro-watershed

Sl. No.	Particulars	LL (12)		MF (40)		SF (87)		SMF (38)		MDF(17)		All (194)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	0-15 years of age	1	8.33	7	17.50	19	21.84	11	28.95	5	29.41	43	22.16
2	16-35 years of age	6	50	19	47.50	39	44.83	11	28.95	7	41.18	82	42.27
3	36-60 years of age	5	41.67	13	32.50	22	25.29	12	31.58	3	17.65	55	28.35
4	> 61 years	0	0	1	2.50	7	8.05	4	10.53	2	11.76	14	7.22
	Total	12	100	40	100	87	100	38	100	17	100	194	100

Education level of household members: Education level of household members in Vojenahalli-2 micro-watershed is presented in Table 4. The results indicated that Vojenahalli-2 had 27.32 per cent illiterates, 29.38 per cent of them had primary school education, 7.73 per cent of them had middle school education, 21.13 per cent of them had

high school education, 9.28 per cent of them had PUC education and 2.58 per cent of them had degree.

Table 4. Education level of household members in Vojenahalli-2 micro-watershed

Sl.No.	Particulars	LL (12)		MF (40)		SF (87)		SMF (38)		MDF (17)		All (194)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Illiterate	6	50.00	8	20.00	23	26.44	10	26.32	6	35.29	53	27.32
2	Primary School	0	0.00	11	27.50	27	31.03	14	36.84	5	29.41	57	29.38
3	Middle School	1	8.33	3	7.50	8	9.20	1	2.63	2	11.76	15	7.73
4	High School	5	41.67	11	27.50	15	17.24	9	23.68	1	5.88	41	21.13
5	PUC	0	0.00	5	12.50	11	12.64	2	5.26	0	0.00	18	9.28
6	Degree	0	0.00	1	2.50	1	1.15	2	5.26	1	5.88	5	2.58
7	Others	0	0.00	1	2.50	2	2.30	0	0.00	2	11.76	5	2.58
Total		12	100.00	40	100.00	87	100.00	38	100.00	17	100.00	194	100.00

Occupation of household heads: The data regarding the occupation of the household heads in Vojenahalli-2 micro-watershed is presented in Table 5. The results indicate that, 68.57 per cent of the household heads were practicing agriculture, 28.57 per cent of the household heads were agricultural labourers and 2.86 per cent of the household heads were trade and business and housewives.

Table 5: Occupation of household heads in Vojenahalli-2 micro-watershed

Sl.No.	Particulars	LL (3)		MF (8)		SF (13)		SMF (8)		MDF (3)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Agriculture	0	0.00	5	62.50	10	76.92	7	87.50	2	66.67	24	68.57
2	Agricultural Labour	3	100.00	3	37.50	3	23.08	0	0.00	1	33.33	10	28.57
3	Trade & Business	0	0.00	0	0.00	0	0.00	1	12.50	0	0.00	1	2.86
4	Housewife	0	0.00	0	0.00	1	7.69	0	0.00	0	0.00	1	2.86
Total		3	100.00	8	100.00	14	100.00	8	100.00	3	100.00	36	100.00

Occupation of the household members: The data regarding the occupation of the household members in Vojenahalli-2 micro-watershed is presented in Table 6. The results indicate that agriculture was the major occupation for 25.77 per cent of the household members, 39.18 per cent were agricultural labourers, 0.52 per cent were private service and trade and business, 24.74 per cent were student, 5.67 per cent were housewives and 2.58 per cent were children.

Table 6: Occupation of family members in Vojenahalli-2 micro-watershed

Sl.No.	Particulars	LL (12)		MF (40)		SF (87)		SMF (38)		MDF (17)		All (194)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Agriculture	0	0	12	30	20	22.99	14	36.84	4	23.53	50	25.77
2	Agricultural Labour	11	91.67	17	42.50	35	40.23	6	15.79	7	41.18	76	39.18
3	Private Service	0	0	0	0	1	1.15	0	0	0	0	1	0.52
4	Trade & Business	0	0	0	0	0	0	1	2.63	0	0	1	0.52
5	Student	1	8.33	8	20	24	27.59	12	31.58	3	17.65	48	24.74
6	Others	0	0	0	0	0	0	2	5.26	0	0	2	1.03
7	Housewife	0	0	2	5	5	5.75	3	7.89	1	5.88	11	5.67
8	Children	0	0	1	2.50	2	2.30	0	0	2	11.76	5	2.58
Total		12	100	40	100	87	100	38	100	17	100	194	100

Institutional participation of the household members: The data regarding the institutional participation of the household members in Vojenahalli-2 micro-watershed is presented in Table 7. The results show that, 100 per cent of the population in the micro watershed has not participated in any local institutions.

Table 7. Institutional Participation of household members in Vojenahalli-2 micro-watershed

Sl.No.	Particulars	LL (12)		MF (40)		SF (87)		SMF (38)		MDF (17)		All (194)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	No Participation	12	100.00	40	100.00	87	100.00	38	100.00	17	100.00	194	100.00
	Total	12	100.00	40	100.00	87	100.00	38	100.00	17	100.00	194	100.00

Type of house owned: The data regarding the type of house owned by the households in Vojenahalli-2 micro-watershed is presented in Table 8. The results indicate that 77.14 per cent of the households possess katcha house, 17.14 per cent of them possess pucca/RCC house and 5.71 per cent of them possess semi pucca house.

Table 8. Type of house owned by households in Vojenahalli-2 micro-watershed

Sl.No.	Particulars	LL (3)		MF (8)		SF (13)		SMF (8)		MDF (3)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Katcha	3	100.00	7	87.50	11	84.62	3	37.50	3	100.00	27	77.14
2	Pucca/RCC	0	0.00	0	0.00	2	15.38	4	50.00	0	0.00	6	17.14
3	Semi pucca	0	0.00	1	12.50	0	0.00	1	12.50	0	0.00	2	5.71
	Total	3	100.00	8	100.00	13	100.00	8	100.00	3	100.00	35	100.00

Durable Assets owned by the households: The data regarding the Durable Assets owned by the households in Vojenahalli-2 micro-watershed is presented in Table 9. The results show that 85.71 per cent of the households possess TV, 77.14 per cent of them possess mixer/grinder, 25.71 per cent of them possess bicycle, 37.14 per cent of the households possess motor cycle and 97.14 per cent of the households possess mobile phones.

Table 9. Durable Assets owned by households in Vojenahalli-2 micro-watershed

Sl.No.	Particulars	LL (3)		MF (8)		SF (13)		SMF (8)		MDF (3)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Television	2	66.67	7	87.50	12	92.31	6	75.00	3	100.00	30	85.71
2	Mixer/Grinder	1	33.33	6	75.00	11	84.62	7	87.50	2	66.67	27	77.14
3	Bicycle	0	0.00	2	25.00	6	46.15	1	12.50	0	0.00	9	25.71
4	Motor Cycle	0	0.00	1	12.50	6	46.15	5	62.50	1	33.33	13	37.14
5	Mobile Phone	3	100.00	7	87.50	13	100.00	8	100.00	3	100.00	34	97.14

Table 10. Average value of durable assets owned by households in Vojenahalli-2 micro-watershed

Sl.No.	Particulars	Average value (Rs.)					
		LL (3)	MF (8)	SF (13)	SMF (8)	MDF (3)	All (35)
1	Television	9,000.00	7,142.00	6,250.00	7,666.00	7,666.00	7,066.00
2	Mixer/Grinder	2,000.00	2,333.00	2,681.00	1,971.00	2,000.00	2,344.00
3	Bicycle	0.00	1,700.00	3,000.00	3,000.00	0.00	2,711.00
4	Motor Cycle	0.00	35,000.00	38,333.00	49,000.00	35,000.00	41,923.00
5	Mobile Phone	2,000.00	2,555.00	1,818.00	2,416.00	3,250.00	2,220.00

Average value of durable assets: The data regarding the average value of durable assets owned by the households in Vojenahalli-2 micro-watershed is presented in Table 10. The results show that the average value of television was Rs. 7,066, mixer grinder was Rs. 2,344, bicycle was Rs. 2,711, motor cycle was Rs. 41,923 and mobile phone was Rs. 2,220.

Farm Implements owned: The data regarding the farm implements owned by the households in Vojenahalli-2 micro-watershed is presented in Table 11. About 20 per cent of the households possess bullock cart, 45.71 per cent of them possess plough, 5.71 per cent of them possess seed/fertilizer drill and tractor, 2.86 per cent of them possess irrigation pump, power tiller and harvester, 20 per cent of them possess sprayer and 57.14 per cent of them possess weeder.

Table 11. Farm Implements owned by households in Vojenahalli-2 micro-watershed

Sl.No.	Particulars	LL (3)		MF (8)		SF (13)		SMF (8)		MDF (3)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Bullock Cart	0	0.00	3	37.50	2	15.38	2	25.00	0	0.00	7	20.00
2	Plough	1	33.33	4	50.00	6	46.15	5	62.50	0	0.00	16	45.71
3	Seed/Fertilizer Drill	0	0.00	0	0.00	0	0.00	2	25.00	0	0.00	2	5.71
4	Irrigation Pump	0	0.00	0	0.00	0	0.00	1	12.50	0	0.00	1	2.86
5	Power Tiller	0	0.00	0	0.00	1	7.69	0	0.00	0	0.00	1	2.86
6	Tractor	0	0.00	0	0.00	1	7.69	1	12.50	0	0.00	2	5.71
7	Sprayer	0	0.00	1	12.50	4	30.77	2	25.00	0	0.00	7	20.00
8	Weeder	0	0.00	5	62.50	6	46.15	7	87.50	2	66.67	20	57.14
9	Harvester	0	0.00	0	0.00	0	0.00	1	12.50	0	0.00	1	2.86
10	Blank	2	66.67	3	37.50	7	53.85	1	12.50	1	33.33	14	40.00

Average value of farm implements: The data regarding the average value of farm Implements owned by the households in Vojenahalli-2 micro-watershed is presented in Table 12. The results show that the average value of bullock cart was Rs. 13,625, plough was Rs. 3,381, seed/fertilizer drill was Rs.35,000, irrigation pump was Rs.75,000, power tiller was Rs. 50,000, tractor was Rs.500,000, sprayer was Rs.3,285, weeder was Rs. 70 and harvester was Rs.80.

Table 12. Average value of farm implements owned by households in Vojenahalli-2 micro-watershed

Sl.No.	Particulars	Average Value (Rs.)					
		LL (3)	MF (8)	SF (13)	SMF (8)	MDF (3)	All (35)
1	Bullock Cart	0.00	11,750.00	11,000.00	20,000.00	0.00	13,625.00
2	Plough	1,500.00	2,250.00	4,833.00	2,920.00	0.00	3,381.00
3	Seed/Fertilizer Drill	0.00	0.00	0.00	35,000.00	0.00	35,000.00
4	Irrigation Pump	0.00	0.00	0.00	75,000.00	0.00	75,000.00
5	Power Tiller	0.00	0.00	50,000.00	0.00	0.00	50,000.00
6	Tractor	0.00	0.00	500,000.00	500,000.00	0.00	500,000.00
7	Sprayer	0.00	2,000.00	3,500.00	3,500.00	0.00	3,285.00
8	Weeder	0.00	92.00	62.00	61.00	83.00	70.00
9	Harvester	0.00	0.00	0.00	80.00	0.00	80.00
10	Blank	1.00	1.00	1.00	1.00	1.00	1.00

Livestock possession by the households: The data regarding the Livestock possession by the households in Vojenahalli-2 micro-watershed is presented in Table 13. The results indicate that, 37.14 per cent of the households possess bullocks and local cow, 2.86 per cent of the households possess crossbred cow and 8.57 per cent of the households possess local cow.

Table 13. Livestock possession by households in Vojenahalli-2 micro-watershed

Sl.No.	Particulars	LL (3)		MF (8)		SF (13)		SMF (8)		MDF (3)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Bullock	0	0.00	4	50.00	4	30.77	3	37.50	2	66.67	13	37.14
2	Local cow	0	0.00	3	37.50	5	38.46	4	50.00	1	33.33	13	37.14
3	Crossbred cow	0	0.00	0	0.00	1	7.69	0	0.00	0	0.00	1	2.86
4	Sheep	0	0.00	1	12.50	1	7.69	0	0.00	1	33.33	3	8.57
5	blank	3	100.00	4	50.00	5	38.46	4	50.00	0	0.00	16	45.71

Average Labour availability: The data regarding the average labour availability in Vojenahalli-2 micro-watershed is presented in Table 14. The results indicate that, average own labour men available in the micro watershed was 1.56, average own labour (women) available was 1.41, average hired labour (men) available was 12 and average hired labour (women) available was 12.18.

Table 14. Average Labour availability in Vojenahalli-2 micro-watershed

Sl.No.	Particulars	LL (3)	MF (8)	SF (13)	SMF (8)	MDF (3)	All (35)
1	Hired labour Female	0.00	15.50	7.77	15.44	12.67	12.18
2	Own Labour Female	0.00	1.63	1.54	1.00	1.33	1.41
3	Own labour Male	0.00	1.63	1.69	1.13	2.00	1.56
4	Hired labour Male	0.00	15.38	7.46	15.56	12.00	12.00

Adequacy of Hired Labour: The data regarding the adequacy of hired labour in Vojenahalli-2 micro-watershed is presented in Table 15. The results indicate that 45.71 per cent of the households opined that the hired labour was adequate and 48.57 per cent of the households opined that hired labour was inadequate.

Table 15. Adequacy of Hired Labour in Vojenahalli-2 micro-watershed

Sl.No.	Particulars	LL (3)		MF (8)		SF (13)		SMF (8)		MDF (3)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Adequate	0	0.00	5	62.50	3	23.08	6	75.00	2	66.67	16	45.71
2	Inadequate	0	0.00	3	37.50	10	76.92	3	37.50	1	33.33	17	48.57

Distribution of land (ha): The data regarding the distribution of land (ha) in Vojenahalli-2 micro-watershed is presented in Table 16. The results indicate that, households of the Vojenahalli-2 micro-watershed possess 38.66 ha (76.48%) of dry land and 11.89 ha (23.52%) of irrigated land. Marginal farmers possess 5.33 ha (92.95%) of dry land and 0.40 ha (7.05%) of irrigated land. Small farmers possess 16.51 ha (96.8%) of dry land and 0.55 ha (3.20%) irrigated land. Semi medium farmers possess 10.34 ha (65.16%) of dry land and 5.53 ha (34.84%) of irrigated land. Medium farmers possess 6.48 ha (54.50%) of dry land and 5.41 ha (45.50 %) of irrigated land.

Table 16. Distribution of land (Ha) in Vojenahalli-2 micro-watershed

Sl.No.	Particulars	MF (8)		SF (13)		SMF (8)		MDF (3)		All (35)	
		ha	%	ha	%	ha	%	ha	%	ha	%
1	Dry	5.33	92.95	16.51	96.80	10.34	65.16	6.48	54.50	38.66	76.48
2	Irrigated	0.40	7.05	0.55	3.20	5.53	34.84	5.41	45.50	11.89	23.52
	Total	5.74	100.00	17.05	100.00	15.87	100.00	11.88	100.00	50.54	100.00

Average land value (Rs./ha): The data regarding the average land value (Rs./ha) in Vojenahalli-2 micro-watershed is presented in Table 17. The results indicate that, the average value of dry land was Rs. 307,715.66 and the average value of irrigated land was Rs. 420,497.11.

Table 17. Average land value (Rs./ha) in Vojenahalli-2 micro-watershed

Sl.No.	Particulars	LL (3)	MF (8)	SF (13)	SMF (8)	MDF (3)	All (35)
1	Dry	0	468,512.90	272,493.26	280,352.26	308,750	307,715.66
2	Irrigated	0	2,470,000	182,962.96	506,295.75	203,368.27	420,497.11

Status of bore wells: The data regarding the status of bore wells in Vojenahalli-2 micro-watershed is presented in Table 18. The results indicate that, there were 1 de-functioning and 7 functioning bore wells in the micro watershed.

Table 18. Status of bore wells in Vojenahalli-2 micro-watershed

Sl.No.	Particulars	LL (3)	MF (8)	SF (13)	SMF (8)	MDF (3)	LF (0)	All (35)
1	De-functioning	0	0	0	1	0	0	1
2	Functioning	0	1	1	4	1	0	7

Source of irrigation: The data regarding the source of irrigation in Vojenahalli-2 micro-watershed is presented in Table 19. The results indicate that, bore well was the major irrigation source in the micro water shed for 20 per cent of the farmers.

Table 19. Source of irrigation in Vojenahalli-2 micro-watershed

Sl.No.	Particulars	LL (3)		MF (8)		SF (13)		SMF (8)		MDF (3)		LF (0)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Bore Well	0	0.00	1	12.50	1	7.69	4	50.00	1	33.33	0	0.00	7	20.00

Depth of water (Avg in meters): The data regarding the depth of water in Vojenahalli-2 micro-watershed is presented in Table 20. The results indicate that, the depth of bore well was found to be 23.23 meters.

Table 20. Depth of water (Avg in meters) in Vojenahalli-2 micro-watershed

Sl.No.	Particulars	LL (3)	MF (8)	SF (13)	SMF (8)	MDF (3)	LF (0)	All (35)
1	Bore Well	0.00	13.34	10.74	57.53	35.56	0.00	23.23

Table 21. Irrigated Area (ha) in Vojenahalli-2 micro-watershed

Sl.No.	Particulars	LL (3)	MF (8)	SF (13)	SMF (8)	MDF (3)	All (35)
1	Kharif	0.00	0.40	0.40	5.53	2.11	8.45
	Total	0.00	0.40	0.40	5.53	2.11	8.45

Irrigated Area (ha): The data regarding the irrigated area (ha) in Vojenahalli-2 micro-watershed is presented in Table 21. The results indicate that marginal and small, semi

medium and medium farmers had an irrigated area of 0.40 ha, 5.53 ha and 2.11 ha respectively.

Cropping pattern: The data regarding the cropping pattern in Vojenahalli-2 micro-watershed is presented in Table 22. The results indicate that, farmers have grown bajra (3.47 ha), groundnut (1.21 ha), maize (37.21 ha), navane (2.02 ha), onion (0.81 ha), Red gram (1.69 ha), sunflower (4.57 ha) and watermelon (0.40 ha). Marginal farmers have grown bajra, and maize. Small farmers have grown bajra, maize, navane, red gram and sunflower. Semi medium farmers have grown maize, onion, red gram and watermelon. Medium farmers have grown groundnut, maize and sunflower.

Table 22. Cropping pattern in Vojenahalli-2 micro-watershed (Area in ha)

Sl.No.	Particulars	LL (3)	MF (8)	SF (13)	SMF (8)	MDF (3)	All (35)
1	Kharif - Bajra	0.00	1.73	1.74	0.00	0.00	3.47
2	Kharif - Groundnut	0.00	0.00	0.00	0.00	1.21	1.21
3	Kharif - Maize	0.00	3.48	11.01	13.67	9.05	37.21
4	Kharif - Navane	0.00	0.00	2.02	0.00	0.00	2.02
5	Kharif - Onion	0.00	0.00	0.00	0.81	0.00	0.81
6	Kharif - Red gram	0.00	0.00	0.81	0.88	0.00	1.69
7	Kharif - Sunflower	0.00	0.00	2.96	0.00	1.62	4.57
8	Kharif - Watermelon	0.00	0.00	0.00	0.40	0.00	0.40
Total		0.00	5.74	18.54	15.77	11.89	51.93

Cropping intensity: The data regarding the cropping intensity in Vojenahalli-2 micro-watershed is presented in Table 23. The results indicate that, the cropping intensity in Vojenahalli-2 micro-watershed was found to be 78.89 per cent.

Table 23. Cropping intensity (%) in Vojenahalli-2 micro-watershed

Sl.No.	Particulars	LL (3)	MF (8)	SF (13)	SMF (8)	MDF (3)	All (35)
1	Cropping intensity	0.00	100.00	66.92	91.67	78.27	78.89

Possession of Bank account and savings: The data regarding the possession of bank account and saving in Vojenahalli-2 micro-watershed is presented in Table 24. The results indicate that, 42.86 per cent of the households have bank account and savings.

Table 24. Possession of Bank account and savings in Vojenahalli-2 micro-watershed

Sl.No.	Particulars	LL (3)		MF (8)		SF (13)		SMF (8)		MDF (3)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Account	0	0.00	3	37.50	9	69.23	2	25.00	1	33.33	15	42.86
2	Savings	0	0.00	3	37.50	9	69.23	2	25.00	1	33.33	15	42.86

Borrowing status: The data regarding the borrowing status in Vojenahalli-2 micro-watershed is presented in Table 25. The results indicate that, 42.86 per cent of the households have availed credit from different sources.

Table 25. Borrowing status in Vojenahalli-2 micro-watershed

Sl.No.	Particulars	LL (3)		MF (8)		SF (13)		SMF (8)		MDF (3)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Credit Availed	0	0.00	3	37.50	9	69.23	2	25.00	1	33.33	15	42.86

Cost of cultivation of Bajra: The data regarding the cost of cultivation of bajra in Vojenahalli-2 micro-watershed is presented in Table 26. The results indicate that, the total cost of cultivation for bajra was Rs. 42621.58. The gross income realized by the farmers was Rs. 19878.02. The net income from bajra cultivation was Rs. -22743.56. Thus the benefit cost ratio was found to be 1:0.47.

Table 26. Cost of Cultivation of bajra in Vojenahalli-2 micro-watershed

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
I	Cost A1				
1	Hired Human Labour	Man days	54.74	12064.46	28.31
2	Bullock	Pairs/day	4.36	2395.77	5.62
3	Tractor	Hours	2.73	2047.48	4.80
4	Machinery	Hours	2.33	1400.70	3.29
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	7.26	831.69	1.95
6	Seed Inter Crop	Kgs.	0.00	0.00	0.00
7	FYM	Quintal	18.40	3679.26	8.63
8	Fertilizer + micronutrients	Quintal	5.33	7628.97	17.90
9	Pesticides (PPC)	Kgs / liters	1.41	1854.04	4.35
10	Irrigation	Number	0.00	0.00	0.00
11	Repairs		0.00	0.00	0.00
12	Msc. Charges (Marketing costs etc)		0.00	0.00	0.00
13	Depreciation charges		0.00	300.39	0.70
14	Land revenue and Taxes		0.00	0.00	0.00
II	Cost B1				
16	Interest on working capital			1680.47	3.94
17	Cost B1 = (Cost A1 + sum of 15 and 16)			33883.23	79.50
III	Cost B2				
18	Rental Value of Land			166.67	0.39
19	Cost B2 = (Cost B1 + Rental value)			34049.90	79.89
IV	Cost C1				
20	Family Human Labour		17.28	4686.99	11.00
21	Cost C1 = (Cost B2 + Family Labour)			38736.89	90.89
V	Cost C2				
22	Risk Premium			10.00	0.02
23	Cost C2 = (Cost C1 + Risk Premium)			38746.89	90.91
VI	Cost C3				
24	Managerial Cost			3874.69	9.09
25	Cost C3 = (Cost C2 + Managerial Cost)			42621.58	100.00
VII	Economics of the Crop				
a.	Main Product	a) Main Product (q)	11.77	17651.32	
		b) Main Crop Sales Price (Rs.)		1500.00	
	By Product	e) Main Product (q)	5.94	2226.70	
		f) Main Crop Sales Price (Rs.)		375.00	
b.	Gross Income (Rs.)			19878.02	
c.	Net Income (Rs.)			-22743.56	
d.	Cost per Quintal (Rs./q.)			3621.96	
e.	Benefit Cost Ratio (BC Ratio)			1:0.47	

Cost of Cultivation of Groundnut: The data regarding the cost of cultivation of groundnut in Vojenahalli-2 micro-watershed is presented in Table 27. The results indicate that, the total cost of cultivation for groundnut was Rs. 30409.08. The gross income realized by the farmers was Rs. 55986.67. The net income from groundnut cultivation was Rs. 25577.58. Thus the benefit cost ratio was found to be 1:1.84.

Table 27. Cost of Cultivation of Groundnut in Vojenahalli-2 micro-watershed

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
I	Cost A1				
1	Hired Human Labour	Man days	40.34	8480.33	27.89
2	Bullock	Pairs/day	1.65	905.67	2.98
3	Tractor	Hours	2.47	1852.50	6.09
4	Machinery	Hours	1.65	988.00	3.25
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	24.70	2223.00	7.31
6	Seed Inter Crop	Kgs.	0.00	0.00	0.00
7	FYM	Quintal	8.23	1646.67	5.42
8	Fertilizer + micronutrients	Quintal	4.12	3787.33	12.45
9	Pesticides (PPC)	Kgs / liters	0.82	576.33	1.90
10	Irrigation	Number	4.12	0.00	0.00
11	Repairs		0.00	0.00	0.00
12	Msc. Charges (Marketing costs etc)		0.00	0.00	0.00
13	Depreciation charges		0.00	6.59	0.02
14	Land revenue and Taxes		0.00	0.00	0.00
II	Cost B1				
16	Interest on working capital			989.20	3.25
17	Cost B1 = (Cost A1 + sum of 15 and 16)			21455.62	70.56
III	Cost B2				
18	Rental Value of Land			333.33	1.10
19	Cost B2 = (Cost B1 + Rental value)			21788.95	71.65
IV	Cost C1				
20	Family Human Labour		22.23	5845.67	19.22
21	Cost C1 = (Cost B2 + Family Labour)			27634.62	90.88
V	Cost C2				
22	Risk Premium			10.00	0.03
23	Cost C2 = (Cost C1 + Risk Premium)			27644.62	90.91
VI	Cost C3				
24	Managerial Cost			2764.46	9.09
25	Cost C3 = (Cost C2 + Managerial Cost)			30409.08	100.00
VII	Economics of the Crop				
a.	Main Product	a) Main Product (q)	16.47	49400.00	
		b) Main Crop Sales Price (Rs.)		3000.00	
	By Product	e) Main Product (q)	8.23	6586.67	
		f) Main Crop Sales Price (Rs.)		800.00	
b.	Gross Income (Rs.)			55986.67	
c.	Net Income (Rs.)			25577.58	
d.	Cost per Quintal (Rs./q.)			1846.71	
e.	Benefit Cost Ratio (BC Ratio)			1:1.84	

Cost of Cultivation of Maize: The data regarding the cost of cultivation of maize in Vojenahalli-2 micro-watershed is presented in Table 28. The results indicate that, the total cost of cultivation for maize was Rs. 36307.28. The gross income realized by the farmers was Rs. 36634.06. The net income from maize cultivation was Rs. 326.78. Thus the benefit cost ratio was found to be 1:1.01.

Table 28. Cost of Cultivation of Maize in Vojenahalli-2 micro-watershed

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
I	Cost A1				
1	Hired Human Labour	Man days	34.98	8141.71	22.42
2	Bullock	Pairs/day	2.69	1570.46	4.33
3	Tractor	Hours	1.91	1399.97	3.86
4	Machinery	Hours	0.71	488.22	1.34
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	17.02	2290.92	6.31
6	Seed Inter Crop	Kgs.	0.00	0.00	0.00
7	FYM	Quintal	7.76	4295.52	11.83
8	Fertilizer + micronutrients	Quintal	5.89	5525.10	15.22
9	Pesticides (PPC)	Kgs / liters	1.19	1139.61	3.14
10	Irrigation	Number	8.65	0.00	0.00
11	Repairs		0.00	0.00	0.00
12	Msc. Charges (Marketing costs etc)		0.00	0.00	0.00
13	Depreciation charges		0.00	983.03	2.71
14	Land revenue and Taxes		0.00	3.35	0.01
II	Cost B1				
16	Interest on working capital			1590.64	4.38
17	Cost B1 = (Cost A1 + sum of 15 and 16)			27428.54	75.55
III	Cost B2				
18	Rental Value of Land			337.53	0.93
19	Cost B2 = (Cost B1 + Rental value)			27766.07	76.48
IV	Cost C1				
20	Family Human Labour		20.37	5236.36	14.42
21	Cost C1 = (Cost B2 + Family Labour)			33002.43	90.90
V	Cost C2				
22	Risk Premium			4.19	0.01
23	Cost C2 = (Cost C1 + Risk Premium)			33006.62	90.91
VI	Cost C3				
24	Managerial Cost			3300.66	9.09
25	Cost C3 = (Cost C2 + Managerial Cost)			36307.28	100.00
VII	Economics of the Crop				
a.	Main Product	a) Main Product (q)		21.76	33076.09
		b) Main Crop Sales Price (Rs.)			1520.37
	By Product	e) Main Product (q)		6.28	3557.97
		f) Main Crop Sales Price (Rs.)			566.67
b.	Gross Income (Rs.)			36634.06	
c.	Net Income (Rs.)			326.78	
d.	Cost per Quintal (Rs./q.)			1668.89	
e.	Benefit Cost Ratio (BC Ratio)			1:1.01	

Cost of Cultivation of Navane: The data regarding the cost of cultivation of navane in Vojenahalli-2 micro-watershed is presented in Table 29. The results indicate that, the total cost of cultivation for navane was Rs. 74337.15. The gross income realized by the farmers was Rs. 45448.00. The net income from navane cultivation was Rs. -28889.15. Thus the benefit cost ratio was found to be 1:0.61.

Table 29. Cost of Cultivation of Navane in Vojenahalli-2 micro-watershed

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
I	Cost A1				
1	Hired Human Labour	Man days	71.63	15314.00	20.60
2	Bullock	Pairs/day	12.35	6792.50	9.14
3	Tractor	Hours	2.47	1852.50	2.49
4	Machinery	Hours	2.47	1482.00	1.99
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	12.35	988.00	1.33
6	Seed Inter Crop	Kgs.	0.00	0.00	0.00
7	FYM	Quintal	24.70	4940.00	6.65
8	Fertilizer + micronutrients	Quintal	9.88	13832.00	18.61
9	Pesticides (PPC)	Kgs /liters	2.47	3705.00	4.98
10	Irrigation	Number	0.00	0.00	0.00
11	Repairs		0.00	0.00	0.00
12	Msc. Charges (Marketing costs etc)		0.00	0.00	0.00
13	Depreciation charges		0.00	1106.56	1.49
14	Land revenue and Taxes		0.00	0.00	0.00
II	Cost B1				
16	Interest on working capital			2817.00	3.79
17	Cost B1 = (Cost A1 + sum of 15 and 16)			52829.56	71.07
III	Cost B2				
18	Rental Value of Land			166.67	0.22
19	Cost B2 = (Cost B1 + Rental value)			52996.23	71.29
IV	Cost C1				
20	Family Human Labour		56.81	14573.00	19.60
21	Cost C1 = (Cost B2 + Family Labour)			67569.23	90.90
V	Cost C2				
22	Risk Premium			10.00	0.01
23	Cost C2 = (Cost C1 + Risk Premium)			67579.23	90.91
VI	Cost C3				
24	Managerial Cost			6757.92	9.09
25	Cost C3 = (Cost C2 + Managerial Cost)			74337.15	100.00
VII	Economics of the Crop				
a.	Main Product	a) Main Product (q)		19.76	45448.00
		b) Main Crop Sales Price (Rs.)			2300.00
b.	Gross Income (Rs.)			45448.00	
c.	Net Income (Rs.)			-28889.15	
d.	Cost per Quintal (Rs./q.)			3762.00	
e.	Benefit Cost Ratio (BC Ratio)			1:0.61	

Cost of Cultivation of Onion: The data regarding the cost of cultivation of onion in Vojenahalli-2 micro-watershed is presented in Table 30. The results indicate that, the total cost of cultivation for onion was Rs. 30525.33. The gross income realized by the farmers was Rs. 37050. The net income from onion cultivation was Rs. 6524.67. Thus the benefit cost ratio was found to be 1:1.21.

Table 30. Cost of Cultivation of Onion in Vojenahalli-2 micro-watershed

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
I	Cost A1				
1	Hired Human Labour	Man days	29.64	4940.00	16.18
2	Bullock	Pairs/day	9.88	5928.00	19.42
3	Tractor	Hours	1.24	988.00	3.24
4	Machinery	Hours	1.24	988.00	3.24
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	3.71	852.15	2.79
6	Seed Inter Crop	Kgs.	0.00	0.00	0.00
7	FYM	Quintal	2.47	494.00	1.62
8	Fertilizer + micronutrients	Quintal	4.94	4544.80	14.89
9	Pesticides (PPC)	Kgs /liters	1.24	1235.00	4.05
10	Irrigation	Number	0.00	0.00	0.00
11	Repairs		0.00	0.00	0.00
12	Msc. Charges (Marketing costs etc)		0.00	0.00	0.00
13	Depreciation charges		0.00	41.99	0.14
14	Land revenue and Taxes		0.00	3.29	0.01
II	Cost B1				
16	Interest on working capital			855.23	2.80
17	Cost B1 = (Cost A1 + sum of 15 and 16)			20870.47	68.37
III	Cost B2				
18	Rental Value of Land			333.33	1.09
19	Cost B2 = (Cost B1 + Rental value)			21203.80	69.46
IV	Cost C1				
20	Family Human Labour		29.64	6545.50	21.44
21	Cost C1 = (Cost B2 + Family Labour)			27749.30	90.91
V	Cost C2				
22	Risk Premium			1.00	0.00
23	Cost C2 = (Cost C1 + Risk Premium)			27750.30	90.91
VI	Cost C3				
24	Managerial Cost			2775.03	9.09
25	Cost C3 = (Cost C2 + Managerial Cost)			30525.33	100.00
VII	Economics of the Crop				
a.	Main Product	a) Main Product (q)		18.53	37050.00
		b) Main Crop Sales Price (Rs.)			2000.00
b.	Gross Income (Rs.)			37050.00	
c.	Net Income (Rs.)			6524.67	
d.	Cost per Quintal (Rs./q.)			1647.79	
e.	Benefit Cost Ratio (BC Ratio)			1:1.21	

Cost of Cultivation of Red gram: The data regarding the cost of cultivation of red gram in Vojenahalli-2 micro-watershed is presented in Table 31. The results indicate that, the total cost of cultivation for red gram was Rs. 35775.70. The gross income realized by the farmers was Rs. 20601.84. The net income from red gram cultivation was Rs. -15173.86. Thus the benefit cost ratio was found to be 1:0.58.

Table 31. Cost of Cultivation of Red gram in Vojenahalli-2 micro-watershed

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
I	Cost A1				
1	Hired Human Labour	Man days	47.62	10423.85	29.14
2	Bullock	Pairs/day	2.27	1246.33	3.48
3	Tractor	Hours	1.80	1351.14	3.78
4	Machinery	Hours	0.62	370.50	1.04
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	5.92	535.36	1.50
6	Seed Inter Crop	Kgs.	0.00	0.00	0.00
7	FYM	Quintal	11.84	2368.03	6.62
8	Fertilizer + micronutrients	Quintal	4.74	6630.48	18.53
9	Pesticides (PPC)	Kgs / liters	1.18	1776.02	4.96
10	Irrigation	Number	0.00	0.00	0.00
11	Repairs		0.00	0.00	0.00
12	Msc. Charges (Marketing costs etc)		0.00	0.00	0.00
13	Depreciation charges		0.00	242.48	0.68
14	Land revenue and Taxes		0.00	0.00	0.00
II	Cost B1				
16	Interest on working capital			1358.39	3.80
17	Cost B1 = (Cost A1 + sum of 15 and 16)			26302.57	73.52
III	Cost B2				
18	Rental Value of Land			166.67	0.47
19	Cost B2 = (Cost B1 + Rental value)			26469.23	73.99
IV	Cost C1				
20	Family Human Labour		22.86	6044.14	16.89
21	Cost C1 = (Cost B2 + Family Labour)			32513.37	90.88
V	Cost C2				
22	Risk Premium			10.00	0.03
23	Cost C2 = (Cost C1 + Risk Premium)			32523.37	90.91
VI	Cost C3				
24	Managerial Cost			3252.34	9.09
25	Cost C3 = (Cost C2 + Managerial Cost)			35775.70	100.00
VII	Economics of the Crop				
a.	Main Product	a) Main Product (q)		7.10	20601.84
		b) Main Crop Sales Price (Rs.)			2900.00
b.	Gross Income (Rs.)			20601.84	
c.	Net Income (Rs.)			-15173.86	
d.	Cost per Quintal (Rs./q.)			5035.94	
e.	Benefit Cost Ratio (BC Ratio)			1:0.58	

Cost of Cultivation of Sunflower: The data regarding the cost of cultivation of sunflower in Vojenahalli-2 micro-watershed is presented in Table 32. The results indicate that, the total cost of cultivation for sunflower was Rs. 32563.90. The gross income realized by the farmers was Rs. 30187.33. The net income from sunflower cultivation was Rs. -2376.57. Thus the benefit cost ratio was found to be 1:0.93.

Table 32. Cost of Cultivation of Sunflower in Vojenahalli-2 micro-watershed

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
I	Cost A1				
1	Hired Human Labour	Man days	40.90	8639.39	26.53
2	Bullock	Pairs/day	0.80	457.98	1.41
3	Tractor	Hours	2.41	1841.27	5.65
4	Machinery	Hours	0.68	409.80	1.26
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	10.27	2413.86	7.41
6	Seed Inter Crop	Kgs.	0.00	0.00	0.00
7	FYM	Quintal	10.73	2145.66	6.59
8	Fertilizer + micronutrients	Quintal	4.98	6482.91	19.91
9	Pesticides (PPC)	Kgs / liters	0.96	1129.74	3.47
10	Irrigation	Number	0.00	0.00	0.00
11	Repairs		0.00	0.00	0.00
12	Msc. Charges (Marketing costs etc)		0.00	0.00	0.00
13	Depreciation charges		0.00	70.71	0.22
14	Land revenue and Taxes		0.00	0.82	0.00
II	Cost B1				
16	Interest on working capital			1461.59	4.49
17	Cost B1 = (Cost A1 + sum of 15 and 16)			25053.73	76.94
III	Cost B2				
18	Rental Value of Land			208.33	0.64
19	Cost B2 = (Cost B1 + Rental value)			25262.07	77.58
IV	Cost C1				
20	Family Human Labour		16.78	4333.73	13.31
21	Cost C1 = (Cost B2 + Family Labour)			29595.79	90.89
V	Cost C2				
22	Risk Premium			7.75	0.02
23	Cost C2 = (Cost C1 + Risk Premium)			29603.54	90.91
VI	Cost C3				
24	Managerial Cost			2960.35	9.09
25	Cost C3 = (Cost C2 + Managerial Cost)			32563.90	100.00
VII	Economics of the Crop				
a.	Main Product	a) Main Product (q)		10.06	30187.33
		b) Main Crop Sales Price (Rs.)			3000.00
b.	Gross Income (Rs.)				30187.33
c.	Net Income (Rs.)				-2376.57
d.	Cost per Quintal (Rs./q.)				3236.18
e.	Benefit Cost Ratio (BC Ratio)				1:0.93

Cost of Cultivation of Watermelon: The data regarding the cost of cultivation of watermelon in Vojenahalli-2 micro-watershed is presented in Table 33. The results indicate that, the total cost of cultivation for watermelon was Rs. 228916.72. The gross income realized by the farmers was Rs. 185250.00. The net income from watermelon cultivation was Rs. -43666.72. Thus the benefit cost ratio was found to be 1:0.81.

Table 33. Cost of Cultivation of Watermelon in Vojenahalli-2 micro-watershed

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
I	Cost A1				
1	Hired Human Labour	Man days	98.80	22724.00	9.93
2	Bullock	Pairs/day	0.00	0.00	0.00
3	Tractor	Hours	4.94	3705.00	1.62
4	Machinery	Hours	2.47	1482.00	0.65
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	4.94	14820.00	6.47
6	Seed Inter Crop	Kgs.	0.00	0.00	0.00
7	FYM	Quintal	49.40	98800.00	43.16
8	Fertilizer + micronutrients	Quintal	9.88	13832.00	6.04
9	Pesticides (PPC)	Kgs / liters	2.47	1976.00	0.86
10	Irrigation	Number	0.00	0.00	0.00
11	Repairs		0.00	0.00	0.00
12	Msc. Charges (Marketing costs etc)		0.00	0.00	0.00
13	Depreciation charges		0.00	25011.22	10.93
14	Land revenue and Taxes		0.00	0.00	0.00
II	Cost B1				
16	Interest on working capital			15532.56	6.79
17	Cost B1 = (Cost A1 + sum of 15 and 16)			197882.78	86.44
III	Cost B2				
18	Rental Value of Land			333.33	0.15
19	Cost B2 = (Cost B1 + Rental value)			198216.11	86.59
IV	Cost C1				
20	Family Human Labour		37.05	9880.00	4.32
21	Cost C1 = (Cost B2 + Family Labour)			208096.11	90.90
V	Cost C2				
22	Risk Premium			10.00	0.00
23	Cost C2 = (Cost C1 + Risk Premium)			208106.11	90.91
VI	Cost C3				
24	Managerial Cost			20810.61	9.09
25	Cost C3 = (Cost C2 + Managerial Cost)			228916.72	100.00
VII	Economics of the Crop				
a.	Main Product	a) Main Product (q)		370.50	185250.00
		b) Main Crop Sales Price (Rs.)			500.00
b.	Gross Income (Rs.)			185250.00	
c.	Net Income (Rs.)			-43666.72	
d.	Cost per Quintal (Rs./q.)			617.86	
e.	Benefit Cost Ratio (BC Ratio)			1:0.81	

Adequacy of fodder: The data regarding the adequacy of fodder in Vojenahalli-2 micro-watershed is presented in Table 34. The results indicate that, 17.14 per cent of the households opined that dry fodder was adequate, 14.71 per cent of the households opined that green fodder was adequate, 31.43 per cent opined that dry fodder was inadequate and 2.86 per cent opined that green fodder was inadequate.

Table 34. Adequacy of fodder in Vojenahalli-2 micro-watershed

Sl.No.	Particulars	LL (3)		MF (8)		SF (13)		SMF (8)		MDF (3)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Adequate-Dry Fodder	0	0.00	1	12.50	2	15.38	2	25.00	1	33.33	6	17.14
2	Inadequate-Dry Fodder	0	0.00	3	37.50	6	46.15	1	12.50	1	33.33	11	31.43
3	Adequate-Green Fodder	0	0.00	4	50.00	7	53.85	3	37.50	2	66.67	16	45.71
4	Inadequate-Green Fodder	0	0.00	0	0.00	1	7.69	0	0.00	0	0.00	1	2.86

Annual gross income: The data regarding the annual gross income in Vojenahalli-2 micro-watershed is presented in Table 35. The results indicate that the annual gross income was Rs. 53,000 for landless farmers, for marginal farmers it was Rs. 38,537.50, for small farmers it was Rs. 62,961.54, for semi medium farmers it was Rs. 80,250 and for medium farmers it was Rs. 120,333.33.

Table 35. Annual gross income in Vojenahalli-2 micro-watershed

(Avg value in Rs.)

Sl.No.	Particulars	LL (3)	MF (8)	SF (13)	SMF (8)	MDF (3)	All (35)
1	Service/salary	0.00	7,500.00	8,076.92	16,250.00	20,000.00	10,142.86
2	Business	0.00	0.00	4,615.38	0.00	0.00	1,714.29
3	Wage	53,000.00	11,875.00	12,692.31	1,875.00	8,333.33	13,114.29
4	Agriculture	0.00	17,912.50	35,500.00	62,125.00	92,000.00	39,365.71
5	Dairy Farm	0.00	625.00	923.08	0.00	0.00	485.71
6	Goat Farming	0.00	625.00	1,153.85	0.00	0.00	571.43
	Income(Rs.)	53,000.00	38,537.50	62,961.54	80,250.00	120,333.33	65,394.29

Average annual expenditure: The data regarding the average annual expenditure in Vojenahalli-2 micro-watershed is presented in Table 36. The results indicate that the average annual expenditure is Rs. 6,160.64. For landless households it was Rs. 11,555.56, for marginal farmers it was Rs 4,700, for small farmers it was Rs. 4,043.39, for semi medium farmers it was Rs. 4,640.63 and for medium farmers it was Rs. 17,888.89.

Table 36. Average annual expenditure in Vojenahalli-2 micro-watershed

(Avg value in Rs.)

Sl.No.	Particulars	LL (3)	MF (8)	SF (13)	SMF (8)	MDF (3)	All (35)
1	Service/salary	0.00	20,000.00	10,666.67	4,000.00	0.00	1,714.29
2	Business	0.00	0.00	15,000.00	0.00	0.00	428.57
3	Wage	34,666.67	8,400.00	6,333.33	5,000.00	7,000.00	6,342.86
4	Agriculture	0.00	8,000.00	14,230.77	28,125.00	46,666.67	17,542.86
5	Dairy Farm	0.00	1,000.00	1,333.33	0.00	0.00	171.43
6	Goat Farming	0.00	200.00	5,000.00	0.00	0.00	148.57
	Total	34,666.67	37,600.00	52,564.10	37,125.00	53,666.67	215,622.44
	Average	11,555.56	4,700.00	4,043.39	4,640.63	17,888.89	6,160.64

Horticulture species grown: The data regarding horticulture species grown in Vojenahalli-2 micro-watershed is presented in Table 37. The results indicate that, sampled households have grown 2 coconut trees in their field.

Table 37. Horticulture species grown in Vojenahalli-2 micro-watershed

Sl.No.	Particulars	LL (3)		MF (8)		SF (13)		SMF (8)		MDF (3)		LF (0)		All (35)	
		F	B	F	B	F	B	F	B	F	B	F	B	F	B
1	Coconut	0	0	0	0	1	0	0	0	1	0	0	0	2	0

*F= Field B=Back Yard

Forest species grown: The data regarding forest species grown in Vojenahalli-2 micro-watershed is presented in Table 38. The results indicate that, households have planted 41 neem, 2 tamarind and 4 banyan trees in their field. Also, the households have planted 5 neem, trees in their backyard.

Table 38: Forest species grown in Vojenahalli-2 micro-watershed

Sl.No.	Particulars	LL (3)		MF (8)		SF (13)		SMF (8)		MDF (3)		LF (0)		All (35)	
		F	B	F	B	F	B	F	B	F	B	F	B	F	B
1	Neem	0	0	7	2	15	3	15	0	4	0	0	0	41	5
2	Tamarind	0	0	0	0	2	0	0	0	0	0	0	0	2	0
3	Banyan	0	0	0	0	3	0	0	0	1	0	0	0	4	0

*F= Field B=Back Yard

Average Additional investment capacity: The data regarding average additional investment capacity in Vojenahalli-2 micro-watershed is presented in Table 39. The results indicated that, households have an average investment capacity of Rs. 3,200 for land development, and Rs. 1,000 for improved crop production.

Table 39: Source of funds for additional investment capacity in Vojenahalli-2 micro-watershed

Sl.No.	Particulars	LL (3)	MF (8)	SF (13)	SMF (8)	MDF (3)	All (35)
1	Land development	0.00	2,125.00	769.23	6,875.00	10,000.00	3,200.00
2	Improved crop production	0.00	500.00	307.69	2,125.00	3,333.33	1,000.00

Table 40. Marketing of the agricultural produce in Vojenahalli-2 micro-watershed

Sl.No	Crops	Output obtained (q)	Output retained (q)	Output sold (q)	Output sold (%)	Avg. Price obtained (Rs/q)
1	Bajra	41.0	0.0	41.0	100.0	1500.0
2	Groundnut	20.0	0.0	20.0	100.0	3000.0
3	Maize	769.0	34.0	735.0	95.58	1520.37
4	Navane	8.0	0.0	8.0	100.0	2300.0
5	Onion	15.0	0.0	15.0	100.0	2000.0
6	Redgram	12.0	0.0	12.0	100.0	2900.0
7	Sunflower	46.0	0.0	46.0	100.0	3000.0
8	Water Melon	150.0	0.0	150.0	100.0	500.0

Marketing of the agricultural produce: The data regarding marketing of the agricultural produce in Vojenahalli-2 micro-watershed is presented in Table 40. The

results indicated that, bajra, groundnut, navane, onion, red gram, sunflower and watermelon were sold to the extent of 100 per cent and maize was sold to the extent of 95.58 per cent.

Marketing Channels used for sale of agricultural produce: The data regarding marketing channels used for sale of agricultural produce in Vojenahalli-2 micro-watershed is presented in Table 41. The results indicated that, about 48.57 per cent of the farmers sold their produce to local/village merchants, 62.86 per cent of the farmers sold their produce to regulated market and 5.71 per cent of the farmers sold their produce to cooperative marketing society.

Table 41. Marketing Channels used for sale of agricultural produce in Vojenahalli-2 micro-watershed

Sl. No.	Particulars	LL (3)		MF (8)		SF (13)		SMF (8)		MDF (3)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Local/village Merchant	0	0.00	5	62.50	4	30.77	7	87.50	1	33.33	17	48.57
2	Regulated Market	0	0.00	3	37.50	13	100.00	3	37.50	3	100.00	22	62.86
3	Cooperative marketing Society	0	0.00	0	0.00	0	0.00	1	12.50	1	33.33	2	5.71

Mode of transport of agricultural produce: The data regarding mode of transport of agricultural produce in Vojenahalli-2 micro-watershed is presented in Table 42. The results indicated that, 114.29 per cent of them used tractor as a mode of transportation.

Table 42. Mode of transport of agricultural produce in Vojenahalli-2 micro-watershed

Sl.No.	Particulars	LL (3)		MF (8)		SF (13)		SMF (8)		MDF (3)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Tractor	0	0.00	8	100.00	16	123.08	11	137.50	5	166.67	40	114.29

Incidence of soil and water erosion problems: The data regarding incidence of soil and water erosion problems in Vojenahalli-2 micro-watershed is presented in Table 43. The results indicated that, 60 per cent of the households have experienced soil and water erosion problems in the farm.

Table 43. Incidence of soil and water erosion problems in Vojenahalli-2 micro-watershed

Sl. No.	Particulars	LL (3)		MF (8)		SF (13)		SMF(8)		MDF (3)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Soil and water erosion problems in the farm	0	0.00	5	62.50	10	76.92	4	50.00	2	66.67	21	60.00

Interest shown towards soil testing: The data regarding Interest shown towards soil testing in Vojenahalli-2 micro-watershed is presented in Table 44. The results indicated that, 74.29 per cent have shown interest in soil test.

Table 44. Interest shown towards soil testing in Vojenahalli-2 micro-watershed

Sl.No.	Particulars	LL (3)		MF (8)		SF (13)		SMF (8)		MDF (3)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Interest in soil test	0	0.00	7	87.50	10	76.92	6	75.00	3	100.00	26	74.29

Usage pattern of fuel for domestic use: The data regarding usage pattern of fuel for domestic use in Vojenahalli-2 micro-watershed is presented in Table 45. The results indicated that, 100 per cent of the households used firewood as a source of fuel.

Table 45. Usage pattern of fuel for domestic use in Vojenahalli-2 micro-watershed

Sl.No.	Particulars	LL (3)		MF (8)		SF (13)		SMF (8)		MDF (3)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Fire Wood	3	100.00	8	100.00	13	100.00	8	100.00	3	100.00	35	100.00

Source of drinking water: The data regarding source of drinking water in Vojenahalli-2 micro-watershed is presented in Table 46. The results indicated that, piped supply was the major source of drinking water for 51.43 per cent of the households, bore well was the source of drinking water for 45.71 per cent of the households and lake/tank was the source of drinking water for 2.86 per cent of the households in micro watershed.

Table 46. Source of drinking water in Vojenahalli-2 micro-watershed

Sl.No.	Particulars	LL (3)		MF (8)		SF (13)		SMF (8)		MDF (3)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Piped supply	3	100.00	5	62.50	3	23.08	6	75.00	1	33.33	18	51.43
2	Bore Well	0	0.00	3	37.50	10	76.92	2	25.00	1	33.33	16	45.71
3	Lake/ Tank	0	0.00	0	0.00	0	0.00	0	0.00	1	33.33	1	2.86

Source of light: The data regarding source of light in Vojenahalli-2 micro-watershed is presented in Table 47. The results indicated that, Electricity was the major source of light for 100 per cent of the households in micro watershed.

Table 47. Source of light in Vojenahalli-2 micro-watershed

Sl.No.	Particulars	LL (3)		MF (8)		SF (13)		SMF (8)		MDF (3)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Electricity	3	100.00	8	100.00	13	100.00	8	100.00	3	100.00	35	100.00

Existence of Sanitary toilet facility: The data regarding existence of sanitary toilet facility in Vojenahalli-2 micro-watershed is presented in Table 48. The results indicated that, 31.43 per cent of the households possess sanitary toilet facility.

Table 48. Existence of Sanitary toilet facility in Vojenahalli-2 micro-watershed

Sl.No.	Particulars	LL (3)		MF (8)		SF (13)		SMF (8)		MDF (3)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Sanitary toilet facility	1	33.33	2	25.00	5	38.46	2	25.00	1	33.33	11	31.43

Possession of PDS card: The data regarding possession of PDS card in Vojenahalli-2 micro-watershed is presented in Table 49. The results indicated that, 2.86 per cent of the sampled households possessed BPL card and 97.14 per cent of the sampled households possessed APL card.

Table 49. Possession of PDS card in Vojenahalli-2 micro-watershed

Sl.No.	Particulars	LL (3)		MF (8)		SF (13)		SMF (8)		MDF (3)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	APL	0	0.00	0	0.00	1	7.69	0	0.00	0	0.00	1	2.86
2	BPL	3	100.00	8	100.00	12	92.31	8	100.00	3	100.00	34	97.14

Participation in NREGA program: The data regarding participation in NREGA programme in Vojenahalli-2 micro-watershed is presented in Table 50. The results indicated that, 42.86 per cent of the households participated in NREGA programme.

Table 50. Participation in NREGA programme in Vojenahalli-2 micro-watershed

Sl.No.	Particulars	LL (3)		MF (8)		SF (13)		SMF (8)		MDF (3)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Participation in NREGA programme	3	100.00	2	25.00	5	38.46	3	37.50	2	66.67	15	42.86

Adequacy of food items: The data regarding adequacy of food items in Vojenahalli-2 micro-watershed is presented in Table 51. The results indicated that, cereals were adequate for 94.29 per cent of the households, pulses were adequate for 57.14 per cent, oilseeds and fruits were adequate for 11.43 per cent, vegetables were adequate for 8.57, milk was adequate for 51.43 per cent, eggs were adequate for 48.57 per cent and meat were adequate for 45.71 per cent.

Table 51. Adequacy of food items in Vojenahalli-2 micro-watershed

Sl.No.	Particulars	LL (3)		MF (8)		SF (13)		SMF (8)		MDF (3)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Cereals	3	100.00	8	100.00	11	84.62	8	100.00	3	100.00	33	94.29
2	Pulses	3	100.00	4	50.00	5	38.46	6	75.00	2	66.67	20	57.14
3	Oilseed	0	0.00	3	37.50	1	7.69	0	0.00	0	0.00	4	11.43
4	Vegetables	0	0.00	1	12.50	0	0.00	2	25.00	0	0.00	3	8.57
5	Fruits	0	0.00	0	0.00	4	30.77	0	0.00	0	0.00	4	11.43
6	Milk	3	100.00	5	62.50	5	38.46	3	37.50	2	66.67	18	51.43
7	Egg	3	100.00	4	50.00	4	30.77	4	50.00	2	66.67	17	48.57
8	Meat	3	100.00	4	50.00	3	23.08	4	50.00	2	66.67	16	45.71

Response on Inadequacy of food items: The data regarding inadequacy of food items in Vojenahalli-2 micro-watershed is presented in Table 52. The results indicated that, cereals were in adequate for 5.71 per cent of the households, pulses were inadequate for 42.86 per cent, oilseeds and fruits were inadequate for 85.71 per cent, vegetables were inadequate for 88.57 per cent, milk was inadequate for 42.86 per cent, eggs were inadequate for 40 per cent and meat was inadequate for 2.86 per cent of the households.

Table 52. Response on Inadequacy of food items in Vojenahalli-2 micro-watershed

Sl.No.	Particulars	LL (3)		MF (8)		SF (13)		SMF (8)		MDF (3)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Cereals	0	0.00	0	0.00	2	15.38	0	0.00	0	0.00	2	5.71
2	Pulses	0	0.00	4	50.00	8	61.54	2	25.00	1	33.33	15	42.86
3	Oilseed	3	100.00	5	62.50	12	92.31	7	87.50	3	100.00	30	85.71
4	Vegetables	3	100.00	6	75.00	13	100.00	6	75.00	3	100.00	31	88.57
5	Fruits	3	100.00	8	100.00	8	61.54	8	100.00	3	100.00	30	85.71
6	Milk	0	0.00	3	37.50	6	46.15	5	62.50	1	33.33	15	42.86
7	Egg	0	0.00	3	37.50	7	53.85	3	37.50	1	33.33	14	40.00
8	Meat	0	0.00	0	0.00	0	0.00	1	12.50	0	0.00	1	2.86

Response on Market surplus of food items: The data regarding market surplus of food items in Vojenahalli-2 micro-watershed is presented in Table 53. The results indicated that, oilseeds and vegetables were market surplus for 2.86 per cent of the households.

Table 53. Response on Market surplus of food items in Vojenahalli-2 micro-watershed

Sl.No.	Particulars	LL (3)		MF (8)		SF (13)		SMF (8)		MDF (3)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Oilseed	0	0.00	0	0.00	0	0.00	1	12.50	0	0.00	1	2.86
2	Vegetables	0	0.00	1	12.50	0	0.00	0	0.00	0	0.00	1	2.86

Farming constraints: The data regarding farming constraints experienced by households in Vojenahalli-2 micro-watershed is presented in Table 54. The results indicated that, lower fertility status of the soil, high rate of interest on credit and lack of transport for safe transport of the agricultural produce to the market was the constraint experienced by 45.71 per cent of the households, wild animal menace on farm field and less rainfall (57.14%), frequent incidence of pest and diseases (68.57%), inadequacy of irrigation water (54.29%), high cost of fertilizers and plant protection chemicals (51.43%), low price for the agricultural commodities and lack of marketing facilities in the area (20%), inadequate extension services (8.57%), and Source of Agri-technology information (42.86%).

Table 54. Farming constraints Experienced in Vojenahalli-2 micro-watershed

Sl. No.	Particulars	MF (8)		SF (13)		SMF(8)		MDF (3)		All (35)	
		N	%	N	%	N	%	N	%	N	%
1	Lower fertility status of the soil	3	37.50	8	61.54	3	37.50	2	66.67	16	45.71
2	Wild animal menace on farm field	7	87.50	5	38.46	6	75	2	66.67	20	57.14
3	Frequent incidence of pest and diseases	5	62.50	9	69.23	7	87.50	3	100	24	68.57
4	Inadequacy of irrigation water	5	62.50	6	46.15	6	75	2	66.67	19	54.29
5	High cost of Fertilizers and plant protection chemicals	5	62.50	6	46.15	4	50	3	100	18	51.43
6	High rate of interest on credit	3	37.50	7	53.85	4	50	2	66.67	16	45.71
7	Low price for the agricultural commodities	2	25	3	23.08	2	25	0	0	7	20
8	Lack of marketing facilities in the area	2	25	4	30.77	0	0	1	33.33	7	20
9	Inadequate extension services	1	12.50	1	7.69	1	12.50	0	0	3	8.57
10	Lack of transport for safe transport of the Agril produce to the market.	4	50	9	69.23	2	25	1	33.33	16	45.71
11	Less rainfall	5	62.50	10	76.92	4	50	1	33.33	20	57.14
12	Source of Agri-technology information(Newspaper/TV/Mobile)	2	25	8	61.54	4	50	1	33.33	15	42.86

SUMMARY

In order to assess the socio-economic condition of the farmers in the watershed a comprehensive questionnaire was prepared. Major components such as demographic conditions, migration details, food consumption and family expenditure pattern, material possession, land holding, land use management, cropping pattern, cost of cultivation of crops, livestock management. The statistical components such as frequency and percentage were used to analyze the data. About 35 households located in the micro watershed were interviewed for the survey.

The data indicated that there were 106 (54.64%) men and 88 (45.36%) women among the sampled households. The average family size of landless farmers' was 4, marginal farmers' was 5, small farmers' was 6.6, semi medium farmers' was 4.7 and medium farmers' was 5.6. The data indicated that, 43 (22.16%) people were in 0-15 years of age, 82 (42.27%) were in 16-35 years of age, 55 (28.35%) were in 36-60 years of age and 14 (7.22%) were above 61 years of age.

The results indicated that Vojenahalli-2 had 27.32 per cent illiterates, 29.38 per cent of them had primary school education, 7.73 per cent of them had middle school education, 21.13 per cent of them had high school education, 9.28 per cent of them had PUC education and 2.58 per cent of them had degree.

The results indicate that, 68.57 per cent of the household heads were practicing agriculture, 28.57 per cent of the household heads were agricultural labourers and 2.86 per cent of the household heads were trade and business and housewives. The results indicate that agriculture was the major occupation for 25.77 per cent of the household members, 39.18 per cent were agricultural labourers, 0.52 per cent were private service and trade and business, 24.74 per cent were student, 5.67 per cent were housewives and 2.58 per cent were children.

The results show that, 100 per cent of the population in the micro watershed has not participated in any local institutions. The results indicate that 77.14 per cent of the households possess katcha house, 17.14 per cent of them possess pucca/RCC house and 5.71 per cent of them possess semi pacca house.

The results show that 85.71 per cent of the households possess TV, 77.14 per cent of them possess mixer/grinder, 25.71 per cent of them possess bicycle, 37.14 per cent of the households possess motor cycle and 97.14 per cent of the households possess mobile phones. The results show that the average value of television was Rs. 7,066, mixer grinder was Rs. 2,344, bicycle was Rs. 2,711, motor cycle was Rs. 41,923 and mobile phone was Rs. 2,220.

About 20 per cent of the households possess bullock cart, 45.71 per cent of them possess plough, 5.71 per cent of them possess seed/fertilizer drill and tractor, 2.86 per

cent of them possess irrigation pump, power tiller and harvester, 20 per cent of them possess sprayer and 57.14 per cent of them possess weeder. The results show that the average value of bullock cart was Rs. 13,625, plough was Rs. 3,381, seed/fertilizer drill was Rs.35,000, irrigation pump was Rs.75,000, power tiller was Rs. 50,000, tractor was Rs.500,000, sprayer was Rs.3,285, weeder was Rs. 70 and harvester was Rs.80.

The results indicate that, 37.14 per cent of the households possess bullocks and local cow, 2.86 per cent of the households possess crossbreed cow and 8.57 per cent of the households possess local cow .

The results indicate that, average own labour men available in the micro watershed was 1.56, average own labour (women) available was 1.41, average hired labour (men) available was 12 and average hired labour (women) available was 12.18. The results indicate that 45.71 per cent of the households opined that the hired labour was adequate and 48.57 per cent of the households opined that hired labour was inadequate.

The results indicate that, households of the Vojenahalli-2 micro-watershed possess 38.66 ha (76.48%) of dry land and 11.89 ha (23.52%) of irrigated land. Marginal farmers possess 5.33 ha (92.95%) of dry land and 0.40 ha (7.05%) of irrigated land. Small farmers possess 16.51 ha (96.8%) of dry land and 0.55 ha (3.20%) irrigated land. Semi medium farmers possess 10.34 ha (65.16%) of dry land and 5.53 ha (34.84%) of irrigated land. Medium farmers possess 6.48 ha (54.50%)of dry land and 5.41 ha (45.50 %) of irrigated land.

The results indicate that, the average value of dry land was Rs. 307,715.66 and the average value of irrigated land was Rs. 420,497.11. In case of marginal famers, the average land value was Rs. 468,512.90 for dry land the average land value was Rs. 2,470,000. In case of small famers, the average land value was Rs. 272,493.26 for dry land, Rs.182,962.96 for irrigated land. In case of semi medium famers, the average land value was Rs. 280,352.26 for dry land and Rs. 506,295.75 for irrigated land. In case of medium farmers, the average land value was Rs. 308,750 and the average value was Rs. 203,368.27 for irrigated land.

The results indicate that, there were 1 de-functioning and 7 functioning bore wells in the micro watershed. The results indicate that, bore well was the major irrigation source in the micro water shed for 20 per cent of the farmers. The results indicate that, the depth of bore well was found to be 23.33 meters.

The results indicate that marginal and small, semi medium and medium farmers had an irrigated area of 0.40 ha, 5.53 ha and 2.11 ha respectively. The results indicate that, farmers have grown bajra (3.47 ha), groundnut (1.21 ha), maize (37.21 ha), navane (2.02 ha), onion (0.81 ha). Red gram (1.69 ha), sunflower (4.57 ha) and watermelon (0.40 ha). Marginal farmers have grown bajra, and maize. Small farmers have grown bajra, maize, navane, red gram and sunflower. Semi medium farmers have grown maize, onion,

red gram and watermelon. Medium farmers have grown groundnut, maize and sunflower. The results indicate that, the cropping intensity in Vojenahalli-2 micro-watershed was found to be 78.89 per cent.

The results indicate that, 42.86 per cent of the households have bank account and savings. The results indicate that, 42.86 per cent of the households have availed credit from different sources.

The results indicate that, the total cost of cultivation for bajra was Rs. 42621.58. The gross income realized by the farmers was Rs. 19878.02. The net income from bajra cultivation was Rs. -22743.56. Thus the benefit cost ratio was found to be 1:0.47. The total cost of cultivation for groundnut was Rs. 30409.08. The gross income realized by the farmers was Rs. 55986.67. The net income from groundnut cultivation was Rs. 25577.58. Thus the benefit cost ratio was found to be 1:1.84. The total cost of cultivation for maize was Rs. 36307.28. The gross income realized by the farmers was Rs. 36634.06. The net income from maize cultivation was Rs. 326.78. Thus the benefit cost ratio was found to be 1:1.01. The total cost of cultivation for onion was Rs. 30525.33. The gross income realized by the farmers was Rs. 37050. The net income from onion cultivation was Rs. 6524.67. Thus the benefit cost ratio was found to be 1:1.21. The total cost of cultivation for red gram was Rs. 35775.70. The gross income realized by the farmers was Rs. 20601.84. The net income from red gram cultivation was Rs. -15173.86. Thus the benefit cost ratio was found to be 1:0.58. The total cost of cultivation for sunflower was Rs. 32563.90. The gross income realized by the farmers was Rs. 30187.33. The net income from sunflower cultivation was Rs. -2376.57. Thus the benefit cost ratio was found to be 1:0.93. The total cost of cultivation for watermelon was Rs. 228916.72. The gross income realized by the farmers was Rs. 185250.00. The net income from watermelon cultivation was Rs. -43666.72. Thus the benefit cost ratio was found to be 1:0.81.

The results indicate that, 17.14 per cent of the households opined that dry fodder was adequate, 14.71 per cent of the households opined that green fodder was adequate, 31.43 per cent opined that dry fodder was inadequate and 2.86 per cent opined that green fodder was inadequate.

The results indicate that the annual gross income was Rs. 53,000 for landless farmers, for marginal farmers it was Rs. 38,537.50, for small farmers it was Rs. 62,961.54, for semi medium farmers it was Rs. 80,250 and for medium farmers it was Rs. 120,333.33. The results indicate that the average annual expenditure is Rs. 6,160.64. For landless households it was Rs. 11,555.56, for marginal farmers it was Rs 4,700, for small farmers it was Rs. 4,043.39, for semi medium farmers it was Rs. 4,640.63 and for medium farmers it was Rs. 17,888.89.

The results indicate that, sampled households have grown 2 coconut trees in their field. The results indicate that, households have planted 41 neem, 2 tamarind and 4 banyan trees in their field. Also, the households have planted 5 neem, trees in their backyard.

The results indicated that, households have an average investment capacity of Rs. 3.200 for land development, and Rs. 1,000 for improved crop production.

The results indicated that, bajra, groundnut, navane, onion, red gram, sunflower and watermelon were sold to the extent of 100 per cent and maize was sold to the extent of 95.58 per cent.

The results indicated that, about 48.57 per cent of the farmers sold their produce to local/village merchants, 62.86 per cent of the farmers sold their produce to regulated market and 5.71 per cent of the farmers sold their produce to cooperative marketing society. The results indicated that, 114.29 per cent of them used tractor as a mode of transportation.

The results indicated that, 60 per cent of the households have experienced soil and water erosion problems in the farm. The results indicated that, 74.29 per cent have shown interest in soil test.

The results indicated that, 100 per cent of the households used firewood as a source of fuel. The results indicated that, piped supply was the major source of drinking water for 51.43 per cent of the households, bore well was the source of drinking water for 45.71 per cent of the households and lake/tank was the source of drinking water for 2.86 per cent of the households in micro watershed.

Electricity was the major source of light for 100 per cent of the households in micro watershed. The results indicated that, 31.43 per cent of the households possess sanitary toilet facility. The results indicated that, 2.86 per cent of the sampled households possessed BPL card and 97.14 per cent of the sampled households possessed BPL card. The results indicated that, 42.86 per cent of the households participated in NREGA programme.

The results indicated that, cereals were adequate for 94.29 per cent of the households, pulses were adequate for 57.14 per cent, oilseeds and fruits were adequate for 11.43 per cent, vegetables were adequate for 8.57, milk was adequate for 51.43 per cent, eggs were adequate for 48.57 per cent and meat were adequate for 45.71 per cent.

The results indicated that, cereals were in adequate for 5.71 per cent of the households, pulses were inadequate for 42.86 per cent, oilseeds and fruits were inadequate for 85.71 per cent, vegetables were inadequate for 88.57 per cent, milk was inadequate for 42.86 per cent, eggs were inadequate for 40 per cent and meat was inadequate for 2.86 per cent of the households.

The results indicated that, oilseeds and vegetables were market surplus for 2.86 per cent of the households.

The results indicated that, lower fertility status of the soil, high rate of interest on credit and lack of transport for safe transport of the agricultural produce to the market was the constraint experienced by 45.71 per cent of the households, wild animal menace on farm field and less rainfall (57.14%), frequent incidence of pest and diseases (68.57%), inadequacy of irrigation water (54.29%), high cost of fertilizers and plant protection chemicals (51.43%), low price for the agricultural commodities and lack of marketing facilities in the area (20%), inadequate extension services (8.57%), and Source of Agri-technology information (42.86%).