



हर कदम, हर डगर
किसानों का हमसफर
भारतीय कृषि अनुसंधान परिषद

Agrisearch with a human touch

**LAND RESOURCE INVENTORY AND SOCIO-ECONOMIC STATUS OF
FARM HOUSEHOLDS FOR WATERSHED PLANNING AND
DEVELOPMENT**

BHIMANAHALLI-2 (4D5B1A2b) MICROWATERSHED

Yadgir Taluk & District, Karnataka

Karnataka Watershed Development Project – II

SUJALA – III

World Bank funded Project



THE WORLD BANK



ICAR – NATIONAL BUREAU OF SOIL SURVEY AND LAND USE PLANNING



ICAR - NBSS & LUP

**WATERSHED DEVELOPMENT DEPARTMENT
GOVT. OF KARNATAKA, BANGALORE**



About ICAR - NBSS&LUP

The ICAR-National Bureau of Soil Survey and Land Use Planning (ICAR-NBSS&LUP), Nagpur, a premier Institute of the Indian Council of Agricultural Research (ICAR), was set up during 1976 with the objective to prepare soil resource maps at national, state and district levels and to provide research inputs in soil resource mapping and its applications, land evaluation, land use planning, land resource management, and database management using GIS for optimising land use on different kinds of soils in the country.

The Bureau has been engaged in carrying out soil resource survey, agro-ecological and soil degradation mapping at the country, state and district levels for qualitative assessment and monitoring the soil health towards viable land use planning. The research activities have resulted in identifying the soil potentials and problems, and the various applications of the soil surveys with the ultimate objective of sustainable agricultural development. The Bureau has the mandate to correlate and classify soils of the country and maintain a National Register of all the established soil series. The Institute is also imparting in-service training to staff of the soil survey agencies in the area of soil survey, land evaluation and soil survey interpretations for land use planning. The Bureau in collaboration with Panjabrao Krishi Vidyapeeth, Akola is running post-graduate teaching and research programme in land resource management, leading to M.Sc. and Ph.D. degrees.

Citation: Rajendra Hegde, Ramesh Kumar, S.C., B.A. Dhanorkar, S. Srinivas, M. Lalitha, K.V. Niranjana, R.S. Reddy and S.K. Singh (2019). Land resource inventory and socio-economic status of farm households for watershed planning and development of Bhimanahalli-2 (4D5B1A2b) Microwatershed, Yadgir Taluk and District, Karnataka", ICAR-NBSS&LUP Sujala MWS Publ .521, ICAR – NBSS & LUP, RC, Bangalore. p.137 & 28.

TO OBTAIN COPIES,

Please write to:

Director, ICAR - NBSS & LUP,

Amaravati Road, NAGPUR - 440 033, India

Phone : (0712) 2500386, 2500664, 2500545 (O)

Telefax : 0712-2522534

E-Mail : director@nbsslup.ernet.in

Website URL : nbsslup.in

Or

Head, Regional Centre, ICAR - NBSS&LUP, Hebbal, Bangalore - 560 024

Phone : (080) 23412242, 23510350 (O)

Telefax : 080-23510350

E-Mail : nbssrcb@gmail.com



**LAND RESOURCE INVENTORY AND SOCIO-ECONOMIC
STATUS OF FARM HOUSEHOLDS FOR WATERSHED
PLANNING AND DEVELOPMENT**

BHIMANAHALLI-2 (4D5B1A2b) MICROWATERSHED

Yadgir Taluk & District, Karnataka

Karnataka Watershed Development Project – II

Sujala-III

World Bank funded Project



THE WORLD BANK



**ICAR – NATIONAL BUREAU OF SOIL SURVEY AND LAND USE
PLANNING**



ICAR - NBSS & LUP



**WATERSHED DEVELOPMENT DEPARTMENT, GOVT. OF
KARNATAKA, BANGALORE**



PREFACE

In Karnataka, as in other Indian States, the livelihoods of rural people are intertwined with farming pursuits. The challenges in agriculture are seriously threatening the livelihood of a large number of farmers as they have been practicing farming in contextual factors beyond their control. Climatic factors are the most important ones and have become much more significant in recent times due to rapid climate changes induced by intensive anthropogenic activities affecting our ecosystem in multiple ways. Climate change has become the reality, it is happening and efforts to evolve and demonstrate climate resilient technologies have become essential. Due to the already over stressed scenario of agrarian sector, the climate change is resulting in manifold increase in the complexities, pushing the rural mass to face more and more unpredictable situations. The rising temperatures and unpredictable rainfall patterns are going to test seriously the informed decisions farmers have to make in order to survive in farming and sustain their livelihood.

It is generally recognized that impacts of climate change shall not be uniform across the globe. It is said that impact of climate change is more severe in South Asia. Based on the analysis of meteorological data, it is predicted that in India, there will be upward trend in mean temperature, downward trend in relative humidity, annual rainfall and number of wet days in a year. Also, in general, phenomena like erratic monsoon, spread of tropical diseases, rise in sea levels, changes in availability of fresh water, frequent floods, droughts, heat waves, storms and hurricanes are predicted. Each one of these adverse situations are already being experienced in various parts of India and also at the global level. Decline in agricultural productivity of small and marginal farmers becoming more vulnerable is already witnessed.

In Karnataka, more than 60 per cent of the population live in rural areas and depend on agriculture and allied activities for their livelihood. Though the state has achieved significant progress in increasing the yield of many crops, there is tremendous pressure on the land resources due to the growing and competing demands of various land uses. This is reflected in the alarming rate of land degradation observed. Already more than 50 per cent of the area is affected by various forms of degradation. If this trend continues, the sustainability of the fragile ecosystem will be badly affected. The adverse effects of change in the climatic factors are putting additional stress on the land resources and the farmers dependent on this.

The natural resources (land, water and vegetation) of the state need adequate and constant care and management, backed by site-specific technological interventions and investments particularly by the government. Detailed database pertaining to the nature of the land resources, their constraints, inherent potentials and suitability for various land

based rural enterprises, crops and other uses is a prerequisite for preparing location-specific action plans, which are in tune with the inherent capability of the resources. Any effort to evolve climate resilient technologies has to be based on the baseline scientific database. Then only one can expect effective implementation of climate resilient technologies, monitor the progress, make essential review of the strategy, and finally evaluate the effectiveness of the implemented programs. The information available at present on the land resources of the state are of general nature and useful only for general purpose planning. Since the need of the hour is to have site-specific information suitable for farm level planning and detailed characterization and delineation of the existing land resources of an area into similar management units is the only option.

ICAR-NBSS&LUP, Regional Centre, Bangalore has taken up a project sponsored by the Karnataka Watershed Development Project-II, (Sujala-III), Government of Karnataka funded by the World Bank under Component-1 Land Resource Inventory. This study was taken up to demonstrate the utility of such a database in reviewing, monitoring and evaluating all the land based watershed development programs on a scientific footing. To meet the requirements of various land use planners at grassroots level, the present study on Land Resource Inventory and Socio-Economic Status of Farm Households for Watershed Planning and Development of Bhimanahalli-2 microwatershed in Yadgir Taluk and District, Karnataka” for integrated development was taken up in collaboration with the State Agricultural Universities, IISC, KRSRAC, KSNDMC as Consortia partners. The project provides detailed land resource information at cadastral level (1:7920 scale) for all the plots and socio-economic status of farm households covering thirty per cent farmers randomly selected representing landed and landless class of farmers in the micro-watershed. The project report with the accompanying maps for the microwatershed will provide required detailed database for evolving effective land use plan, alternative land use options and conservation plans for the planners, administrators, agricultural extension personnel, KVK officials, developmental departments and other land users to manage the land resources in a sustainable manner.

It is hoped that this database will be useful to the planners, administrators and developmental agencies working in the area in not only for formulating location specific developmental schemes but also for their effective monitoring at the village/watershed level.

Nagpur

Date: 05-11-2019

S.K. SINGH

Director, ICAR - NBSS&LUP, Nagpur

Contributors

Dr. Rajendra Hegde Principal Scientist, Head & Project Leader, Sujala-III Project ICAR-NBSS&LUP, Regional Centre, Bangalore	Dr. S.K.Singh Director, ICAR-NBSS&LUP Coordinator, Sujala-III Project Nagpur
Soil Survey, Mapping & Report Preparation	
Dr. B.A. Dhanorkar	Sh. R.S. Reddy
Dr. K.V. Niranjana	Sh. Venkata Giriyappa
	Sh. Somashekar T N
	Smt. Chaitra, S.P.
	Dr. Gayathri. B.
	Dr. Gopali bardhan
	Dr. H.R. Savitha
	Sh. Nagendra, B.R
Field Work	
Sh. C.BacheGowda	Sh. Mahesh, D.B.
Sh. Somashekar	Sh. Ashok S Sindagi
Sh. M. Jayaramaiah	Sh. Veerabhadrapa B.
Sh. Paramesha, K.	Sh. Shankarappa
Sh. B. M. Narayana Reddy	Sh. Anand
	Sh. Arun N Kambar.
	Sh Kamalesh Awate
	Sh. Sharaan Kumar Huppar
	Sh. Yogesh H.N.
	Sh. Kalaveerachari R Kammar
GIS Work	
Dr. S.Srinivas	Sh. A.G.Devendra Prasad
Sh. D.H.Venkatesh	Sh. Prakashanaik, M.K.
Smt.K.Sujatha	Sh. Abhijith Sastry, N.S.
Smt. K.V.Archana	Sh. Sudip Kumar Suklabaidya
Sh. N. Maddileti	Sh. Avinash, K.N.
	Sh. Amar Suputhra, S
	Sh. Deepak, M.J.
	Smt. K.Karunya Lakshmi
	Ms. Seema, K. V.
	Ms. A. Rajab Nisha

Laboratory Analysis	
Dr. K.M.Nair	Ms. Steffi Peter
Smt. Arti Koyal	Ms. Thara, V.R
Smt. Parvathy	Ms. Roopa, G.
	Ms. Swati, H.
	Sh. Shantaveera Swami
	Ms. Shwetha, N.K.
	Smt. Ishrat Haji
	Ms. P. Pavan Kumari
	Ms. Padmaja
	Ms. Veena, M.
Socio-Economic Analysis	
Dr. S.C. Ramesh Kumar	Sh. M.K. Prakashanaik
	Ms. Karuna V. Kulkarni
	Mrs. Sowmya A.N
	Sh. Vinod R
	Sh. Basavaraja
	Sh. Vijay Kumar Lamani
	Ms. Sowmya K.B
	Mrs. Prathibha, D.G
	Sh. Rajendra,D
Soil & Water Conservation	
Sh. Sunil P. Maske	
Watershed Development Department, GoK, Bangalore	
Sh. Rajeev Ranjan IFS Project Director & Commissioner, WDD	Dr. A. Natarajan NRM Consultant, Sujala-III Project
Dr. S.D. Pathak IFS Executive Director & Chief Conservator of Forests, WDD	

PART-A

LAND RESOURCE INVENTORY

Contents

Preface		
Contributors		
Executive Summary		
Chapter 1	Introduction	1
Chapter 2	Geographical Setting	3
2.1	Location and Extent	3
2.2	Geology	3
2.3	Physiography	5
2.4	Drainage	5
2.5	Climate	5
2.6	Natural Vegetation	6
2.7	Land Utilization	7
Chapter 3	Survey Methodology	11
3.1	Base maps	11
3.2	Image Interpretation for Physiography	11
3.3	Field Investigation	14
3.4	Soil Mapping	16
3.5	Land Management Units	17
3.6	Laboratory Characterization	17
Chapter 4	The Soils	23
4.1	Soils of granite gneiss landscape	23
4.2	Soils of alluvial landscape	30
Chapter 5	Interpretation for Land Resource Management	43
5.1	Land Capability Classification	43
5.2	Soil Depth	45
5.3	Surface Soil Texture	46
5.4	Soil Gravelliness	47
5.5	Available Water Capacity	48
5.6	Soil Slope	49
5.7	Soil Erosion	50
Chapter 6	Fertility Status	53
6.1	Soil Reaction (pH)	53
6.2	Electrical Conductivity (EC)	53
6.3	Organic Carbon (OC)	53
6.4	Available Phosphorus	55
6.5	Available Potassium	55
6.6	Available Sulphur	55
6.7	Available Boron	55
6.8	Available Iron	55
6.9	Available Manganese	58
6.10	Available Copper	58

6.11	Available Zinc	58
Chapter 7	Land Suitability for Major Crops	61
7.1	Land suitability for Sorghum	61
7.2	Land suitability for Maize	62
7.3	Land suitability for Bajra	63
7.4	Land suitability for Groundnut	64
7.5	Land suitability for Sunflower	65
7.6	Land suitability for Redgram	66
7.7	Land suitability for Bengal gram	67
7.8	Land suitability for Cotton	68
7.9	Land suitability for Chilli	69
7.10	Land suitability for Tomato	70
7.11	Land suitability for Brinjal	71
7.12	Land suitability for Onion	72
7.13	Land suitability for Bhendi	73
7.14	Land suitability for Drumstick	74
7.15	Land suitability for Mango	75
7.16	Land suitability for Guava	76
7.17	Land suitability for Sapota	77
7.18	Land Suitability for Pomegranate	78
7.19	Land Suitability for Musambi	79
7.20	Land Suitability for Lime	80
7.21	Land Suitability for Amla	81
7.22	Land Suitability for Cashew	82
7.23	Land Suitability for Jackfruit	83
7.24	Land Suitability for Jamun	84
7.25	Land Suitability for Custard apple	85
7.26	Land Suitability for Tamarind	86
7.27	Land Suitability for Mulberry	87
7.28	Land Suitability for Marigold	88
7.29	Land Suitability for Chrysanthemum	89
7.30	Land Management Units (LMUs)	121
7.31	Proposed Crop Plan for Bhimanahalli-2 Microwatershed	122
Chapter 8	Soil Health Management	125
Chapter 9	Soil and Water conservation Treatment Plan	129
9.1	Treatment Plan	129
9.2	Recommended Soil and Water Conservation measures	133
9.3	Greening of Microwatershed	134
	References	137
	Appendix I	I-VIII
	Appendix II	IX-XVII
	Appendix III	XVIII-XXIV

LIST OF TABLES

2.1	Mean Monthly Rainfall, PET, 1/2 PET at Yadgir Taluk & District	6
2.2	Land Utilization in Yadgir district	7
3.1	Differentiating Characteristics used for Identifying Soil Series	16
3.2	Soil map unit description of Bhimanahalli-2 Microwatershed	17
4.1	Physical and Chemical Characteristics of Soil Series identified in Bhimanahalli-2 microwatershed	32
7.1	Soil-Site Characteristics of Bhimanahalli-2 Microwatershed	91
7.2	Crop suitability for Sorghum	92
7.3	Crop suitability for Maize	93
7.4	Crop suitability for Bajra	94
7.5	Crop suitability for Groundnut	95
7.6	Crop suitability for Sunflower	96
7.7	Crop suitability for Redgram	97
7.8	Crop suitability for Bengal gram	98
7.9	Crop suitability for Cotton	99
7.10	Crop suitability for Chilli	100
7.11	Crop suitability for Tomato	101
7.12	Crop suitability for Brinjal	102
7.13	Crop suitability for Onion	103
7.14	Crop suitability for Bhendi	104
7.15	Crop suitability for Drumstick	105
7.16	Crop suitability for Mango	106
7.17	Crop suitability for Guava	107
7.18	Crop suitability for Sapota	108
7.19	Crop suitability for Pomegranate	109
7.20	Crop suitability for Musambi	110
7.21	Crop suitability for Lime	111
7.22	Crop suitability for Amla	112
7.23	Crop suitability for Cashew	113
7.24	Crop suitability for Jackfruit	114
7.25	Crop suitability for Jamun	115
7.26	Crop suitability for Custard apple	116

7.27	Crop suitability for Tamarind	117
7.28	Crop suitability for Mulberry	118
7.29	Crop suitability for Marigold	119
7.30	Crop suitability for Chrysanthemum	120
7.31	Proposed Crop Plan for Bhimanahalli-2 Microwatershed	123

LIST OF FIGURES

2.1	Location map of Bhimanahalli-2 Microwatershed	3
2.2	Granite and granite gneiss rock formation	4
2.3	Rainfall distribution in Yadgir Taluk & District	6
2.4	Natural vegetation of Bhimanahalli-2 Microwatershed	7
2.5	Current Land use map of Bhimanahalli-2 Microwatershed	8
2.76a	Different crops and cropping systems in Bhimanahalli-2 Microwatershed	8
2.6 b	Different crops and cropping systems in Bhimanahalli-2 Microwatershed	9
3.1	Scanned and Digitized Cadastral map of Bhimanahalli-2 Microwatershed	13
3.2	Satellite image of Bhimanahalli-2 Microwatershed	13
3.3	Cadastral map overlaid on IRS PAN+LISS IV merged imagery of Bhimanahalli-2 Microwatershed	14
3.4	Location of profiles in a transect	15
3.5	Soil phase or management units of Bhimanahalli-2 Microwatershed	21
5.1	Land Capability Classification map of Bhimanahalli-2 Microwatershed	45
5.2	Soil Depth map of Bhimanahalli-2 Microwatershed	46
5.3	Surface Soil Texture map of Bhimanahalli-2 Microwatershed	47
5.4	Soil Gravelliness map of Bhimanahalli-2 Microwatershed	48
5.5	Soil Available Water Capacity map Bhimanahalli-2 Microwatershed	49
5.6	Soil Slope map of Bhimanahalli-2 Microwatershed	50
5.7	Soil Erosion map of Bhimanahalli-2 Microwatershed	51
6.1	Soil Reaction (pH) map of Bhimanahalli-2 Microwatershed	54
6.2	Electrical Conductivity (EC) map of Bhimanahalli-2 Microwatershed	54
6.3	Soil Organic Carbon (OC) map of Bhimanahalli-2 Microwatershed	55
6.4	Soil Available Phosphorus map of Bhimanahalli-2 Microwatershed	56
6.5	Soil Available Potassium map of Bhimanahalli-2 Microwatershed	56
6.6	Soil Available Sulphur map of Bhimanahalli-2 Microwatershed	57
6.7	Soil Available Boron map of Bhimanahalli-2 Microwatershed	57
6.8	Soil Available Iron map of Bhimanahalli-2 Microwatershed	58
6.9	Soil Available Manganese map of Bhimanahalli-2 Microwatershed	59
6.10	Soil Available Copper map of Bhimanahalli-2 Microwatershed	59

6.11	Soil Available Zinc map of Bhimanahalli-2 Microwatershed	60
7.1	Land suitability for Sorghum	62
7.2	Land suitability for Maize	63
7.3	Land suitability for Bajra	64
7.4	Land suitability for Groundnut	65
7.5	Land suitability for Sunflower	66
7.6	Land suitability for Redgram	67
7.7	Land suitability for Bengal gram	68
7.8	Land suitability for Cotton	69
7.9	Land suitability for Chilli	70
7.10	Land suitability for Tomato	71
7.11	Land suitable for Brinjal	72
7.12	Land suitable for Onion	73
7.13	Land suitable for Bhendi	74
7.14	Land suitable for Drumstick	75
7.15	Land suitability for Mango	76
7.16	Land suitability for Guava	77
7.17	Land suitability for Sapota	78
7.18	Land suitability for Pomegranate	79
7.19	Land suitability for Musambi	80
7.20	Land suitability for Lime	81
7.21	Land suitability for Amla	82
7.22	Land suitability for Cashew	83
7.23	Land suitability for Jackfruit	84
7.24	Land suitability for Jamun	85
7.25	Land suitability for Custard apple	86
7.26	Land suitability for Tamarind	87
7.27	Land suitability for Mulberry	88
7.28	Land suitability for Marigold	89
7.29	Land suitability for Chrysanthemum	90
7.30	Land management units map of Bhimanahalli-2 Microwatershed	121
9.1	Soil and water conservation plan map of Bhimanahalli-2 Microwatershed	134

EXECUTIVE SUMMARY

The land resource inventory of Bhimanahalli-2 Microwatershed was conducted using village cadastral maps and IRS satellite imagery on 1:7920 scale. The false colour composites of IRS imagery were interpreted for physiography and the physiographic delineations were used as base for mapping soils. The soils were studied in several transects and a soil map was prepared with phases of soil series as mapping units. Random checks were made all over the area outside the transects to confirm and validate the soil map unit boundaries. The soil map shows the geographic distribution and extent, characteristics, classification, behavior and use potentials of the soils in the microwatershed.

The present study covers an area of 573 ha in Yadgir taluk & district, Karnataka. The climate is semiarid and categorized as drought-prone with an average annual rainfall of 866 mm, of which about 652 mm is received during south-west monsoon, 138 mm during north-east and the remaining 76 mm during the rest of the year. An area of 442 ha in the microwatershed is covered by soils, about 12 ha by quarry, an area of 43 ha by rock outcrops and about 77 ha cover by forest. The salient findings from the land resource inventory are summarized briefly below.

- ❖ The soils belong to 11 soil series and 13 soil phases (management units) and 8 land management units.*
- ❖ The length of crop growing period is about 120-150 days starting from 1st week of June to 4th week of October.*
- ❖ From the master soil map, several interpretative and thematic maps like land capability, soil depth, surface soil texture, soil gravelliness, available water capacity, soil slope and soil erosion were generated.*
- ❖ Soil fertility status maps for macro and micronutrients were generated based on the surface soil samples collected at every 320 m grid interval.*
- ❖ Land suitability for growing 29 major agricultural and horticultural crops was assessed and maps showing the degree of suitability along with constraints were generated.*
- ❖ An area of about 77 per cent is suitable for agriculture in the microwatershed.*
- ❖ About 20 per cent area of the microwatershed has soils that are deep to very deep (100- >150 cm), whereas 29 per cent soils are moderately shallow (50-75 cm), 29 per cent soils are very shallow to shallow (<25-50 cm) in the microwatershed.*
- ❖ About 7 percent soils are loamy and 70 per cent is clayey soils at the surface.*
- ❖ Entire cultivated area is non gravelly (<15%) in the microwatershed.*
- ❖ About 20 per cent area of the microwatershed is very high (>200 mm/m) in available water capacity and 58 per cent soils are low (51-100 mm/m) and very low (<50mm/m) in available water capacity.*
- ❖ An area of about 77 per cent is very gently sloping (1-3% slope) lands and about <1 per cent is nearly level (0-1% slope) lands.*
- ❖ An area of about 69 per cent is moderately (e2) eroded and about 8 per cent is slightly (e1) eroded in the microwatershed.*

- ❖ *An area of about 38 per cent is neutral (pH 6.5-7.3) about 30 per cent is slightly alkaline (pH 7.3-7.8) and about 8 per cent is moderately alkaline (pH 7.8-8.4) in reaction in the microwatershed.*
- ❖ *The Electrical Conductivity (EC) of the soils in the entire cultivated area of the microwatershed is $<2 \text{ ds}^{\text{m}^{-1}}$ indicating that the soils are non-saline.*
- ❖ *An area of 75 per cent is high ($>0.75\%$) and 2 per cent is medium (0.50-0.75%) in organic carbon content.*
- ❖ *An area of 14 per cent is medium (23-57 kg/ha) and 63 percent soils are high ($>57 \text{ kg/ha}$) in available phosphorus.*
- ❖ *Maximum area of about 77 percent is high ($>337\text{kg/ha}$) and about <1 percent is medium (145-337kg/ha) in available potassium.*
- ❖ *Available sulphur content is low ($<10 \text{ ppm}$) in the entire cultivated area of the microwatershed.*
- ❖ *Available boron content is low ($<0.5 \text{ ppm}$) in the entire cultivated area of the microwatershed.*
- ❖ *Available iron content is sufficient ($>4.5 \text{ ppm}$) in an area of 76 per cent and about deficient ($<4.5 \text{ ppm}$) in about <1 per cent in the microwatershed.*
- ❖ *Available manganese and copper are sufficient in all the soils of the microwatershed.*
- ❖ *Available zinc content is deficient ($<0.6 \text{ ppm}$) in the entire cultivated area of the microwatershed.*
- ❖ *The land suitability for 29 major crops grown in the microwatershed were assessed and the areas that are highly suitable (S1) and moderately suitable (S2) are given below. It is however to be noted that a given soil may be suitable for various crops but what specific crop to be grown may be decided by the farmer looking to his capacity to invest on various inputs, marketing infrastructure, market price and finally the demand and supply position.*

Land suitability for various crops in the Microwatershed

Crop	Suitability Area in ha (%)		Crop	Suitability Area in ha (%)	
	Highly suitable (S1)	Moderately suitable (S2)		Highly suitable (S1)	Moderately suitable (S2)
<i>Sorghum</i>	-	278(49)	<i>Guava</i>	-	-
<i>Maize</i>	-	278(49)	<i>Sapota</i>	-	-
<i>Bajra</i>	-	275(48)	<i>Pomegranate</i>	-	57(10)
<i>Groundnut</i>	-	163(29)	<i>Musambi</i>	-	57(10)
<i>Sunflower</i>	-	57(10)	<i>Lime</i>	-	57(10)
<i>Redgram</i>	-	111(19)	<i>Amla</i>	-	164(29)
<i>Bengal gram</i>	-	57(10)	<i>Cashew</i>	-	-
<i>Cotton</i>	-	199(35)	<i>Jackfruit</i>	-	-
<i>Chilli</i>	-	221(38)	<i>Jamun</i>	-	-
<i>Tomato</i>	-	164(29)	<i>Custard apple</i>	-	221(38)
<i>Brinjal</i>	-	164(29)	<i>Tamarind</i>	-	-
<i>Onion</i>	-	164(29)	<i>Mulberry</i>	-	-
<i>Bhendi</i>	-	221(38)	<i>Marigold</i>	-	221(38)
<i>Drumstick</i>	-	-	<i>Chrysanthemum</i>	-	221(38)
<i>Mango</i>	-	-			

- ❖ *Apart from the individual crop suitability, a proposed crop plan has been prepared for the identified LMUs by considering only the highly and moderately suitable lands for different crops and cropping systems with food, fodder, fiber and horticulture crops.*
- ❖ *Maintaining soil-health is vital to crop production and conserve soil and land resource base for maintaining ecological balance and to mitigate climate change. For this, several ameliorative measures have been suggested to these problematic soils like saline/alkali, highly eroded, sandy soils etc.*
- ❖ *Soil and water conservation treatment plan has been prepared that would help in identifying the sites to be treated and also the type of structures required.*
- ❖ *As part of the greening programme, several tree species have been suggested to be planted in marginal and submarginal lands, field bunds and also in the hillocks, mounds and ridges. This would help in not only supplementing the farm income but also provide fodder and fuel to generate lot of biomass which would help in maintaining an ecological balance and also contribute to mitigating the climate change.*

INTRODUCTION

Land is a scarce resource and basic unit for any material production. It can support the needs of the growing population, provided they use the land in a rational and judicious manner. But what is happening in many areas of the state is a cause for concern to everyone involved in the management of land resources at the grassroots level. The area available for agriculture is about 51 per cent of the total geographical area and more than 60 per cent of the people are still dependant on agriculture for their livelihood. The limited land area is under severe stress and strain due to increasing population pressure and competing demands of various land uses. Due to this, every year there is significant diversion of farm lands and water resources for non-agricultural purposes. Apart from this, due to lack of interest in farmers for farming, large tracts of cultivable lands are turning into fallows in many areas and this trend is continuing at an alarming rate.

Further, land degradation has emerged as a serious problem which has already affected about 38 lakh ha of cultivated area in the state. Soil erosion alone has degraded about 35 lakh ha. Almost all the uncultivated areas are facing various degrees of degradation, particularly soil erosion. Salinity and alkalinity has emerged as a major problem in more than 3.5 lakh ha in the irrigated areas of the state. Nutrient depletion and declining factor productivity is common in both rainfed and irrigated areas. The degradation is continuing at an alarming rate and there appears to be no systematic effort among the stakeholders to contain this process. In recent times, an aberration of weather due to climate change phenomenon has added another dimension leading to unpredictable situations to be tackled by the farmers.

In this critical juncture, the challenge before us is not only to increase the productivity per unit area which is steadily declining and showing a fatigue syndrome, but also to prevent or at least reduce the severity of degradation. If the situation is not reversed at the earliest, then the sustainability of the already fragile crop production system and the overall ecosystem will be badly affected in the state. The continued neglect and unscientific use of the resources for a long time has led to the situation observed at present in the state. It is a known fact and established beyond doubt by many studies in the past that the cause for all kinds of degradation is the neglect and irrational use of the land resources. Hence, there is an urgent need to generate a detailed site-specific farm level database on various land resources for all the villages/watersheds in a time bound manner that would help to protect the valuable soil and land resources and also to stabilize the farm production.

Therefore, the land resource inventory required for farm level planning is the one which investigates not only the surface but also consider the other parameters which are critical for productivity *viz.*, soils, climate, water, minerals and rocks, topography, geology, hydrology, vegetation, crops, land use pattern, animal population, socio-

economic conditions, infrastructure, marketing facilities and various schemes and developmental works of the government etc. From the data collected at farm level, the specific problems and potentials of the area can be identified and highlighted, conservation measures required for the area can be planned on a scientific footing, suitability of the area for various uses can be worked out and finally viable and sustainable land use options suitable for each and every land holding can be prescribed.

The Land Resource Inventory is basically done for identifying the potential and problem areas, developing sustainable land use plans, estimation of surface run off and water harvesting potential, preparation of soil and water conservation plans, land degradation/desertification etc. The Bureau is presently engaged in developing an LRI methodology using high resolution satellite remote sensing data and Digital Elevation Model (DEM) data to prepare Landscape Ecological Units (LEU) map representing agro-ecosystem as a whole. The LEU is preferred over landform as the base map for LRI. LEU is the assemblage of landform, slope and land use. An attempt has already been made to upscale the soil resource information from 1:250000 and 1:50000 scale to the LEU map in Goa and other states.

The land resource inventory aims to provide site specific database for Bhimanahalli-2 microwatershed in Yadgir Taluk & District, Karnataka State for the Karnataka Watershed Development Department. The database was generated by using cadastral map of the village as a base along with high resolution IRS LISS IV and Cartosat-1 merged satellite imagery. Later, an attempt will be made to uplink this LRI data generated at 1:7920 scale under Sujala-III Project to the proposed Landscape Ecological Units (LEUs) map.

The study was organized and executed by the ICAR- National Bureau of Soil Survey and Land Use Planning, Regional Centre, Bangalore under Generation of Land Resource Inventory Data Base Component-1 of the Sujala-III Project funded by the World Bank.

GEOGRAPHICAL SETTING

2.1 Location and Extent

The Bhimanahalli-2 microwatershed is located in the northern part of Karnataka in Yadgir Taluk & District, Karnataka State (Fig.2.1). It comprises parts of Bheemanahalli, Handaraki and Ramathirtha villages. It lies between $16^{\circ} 56'$ and $16^{\circ} 58'$ North latitudes and $77^{\circ} 11'$ and $77^{\circ} 13'$ East longitudes, covering an area of about 573 ha. It is on northern side of Yadgir town and is surrounded by Bheemanahalli on the west and southwest, Handaraki on the northwest, east and southeast, Ramathirtha on the northwest and Motahalli on southern side of the microwatershed.




Fig.2.1 Location map of Bhimanahalli-2 Microwatershed

2.2 Geology

Major rock formations observed in the microwatershed are granite gneiss and alluvium (Figs.2.2a and b). Granite gneisses are essentially pink to gray and are coarse to

medium grained. They consist primarily of quartz, feldspar, biotite and hornblende. The gray granite gneisses are highly weathered, fractured and fissured upto a depth of about 10 m. Dolerite dykes and quartz veins are common with variable width and found to occur in Bhimanahalli-2 microwatershed. The most widespread and characteristic development of alluvium in the watershed region lying between the rivers Krishna and Bhima is a wide belt, the underlying formation is gneiss and alluvial soils occur over gneiss, limestone and shale. The thickness of the alluvium generally is limited to less than a meter, except in river valleys where it is very deep extending to tens of meters. Such soils are transported and represent palaeo black soils originally formed at higher elevation, but now occupying river valleys.



Fig.2.2a Granite and granite gneiss rocks



Fig. 2.2b Alluvium

2.3 Physiography

Physiographically, the area has been identified as granite gneiss and alluvium based landscapes on geology. The area has been further subdivided into five landforms, viz; mounds/ridges, summits, side slopes and very gently sloping uplands, plains and valleys based on slope and its relief features. The elevation ranges from 490-518 m above MSL. The mounds and ridges are mostly covered by rock outcrops.

2.4 Drainage

The area is drained by several parallel streams like Bori, Amerja and Kanga which finally join the river Bhima along its course. Though, they are not perennial, during rainy season they carry large quantities of rain water. The microwatershed has only few small tanks which are not capable of storing the water that flows during the rainy season. Due to this, the ground water recharge is very much affected. This is reflected in the failure of many bore wells in the villages. If the available rain water is properly harnessed by constructing new tanks and recharge structures at appropriate places in the villages, then the drinking and irrigation needs of the area can be easily met. The drainage network is parallel to sub parallel and dendritic.

2.5 Climate

The Yadgir district lies in the northern plains of Karnataka and falls under semiarid tract of the state and is categorized as drought- prone with total annual rainfall of 866 mm (Table 2.1). Of the total rainfall, maximum of 652 mm is received during the south-west monsoon period from June to September; the north-east monsoon from October to early December contributes about 138 mm and the remaining 76 mm during the rest of the year. The summer season starts during the middle of February and continues up to the first week of June. The period from December to the middle of February is the coldest season. December is the coldest month with mean daily maximum and minimum temperatures being 29.5⁰C and 10⁰C respectively. During peak summer, temperature shoots up to 45⁰C. Relative humidity varies from 26% in summer to 62% in winter. Rainfall distribution is shown in Figure 2.3. The average Potential Evapo-Transpiration (PET) is 141 mm and varies from a low of 81 mm in December to 199 mm in the month of May. The PET is always higher than precipitation in all the months except end of June to end of September. Generally, the Length of crop Growing Period (LGP) is 120-150 days and starts from 1st week of June to 4th week of October.

Table 2.1 Mean Monthly Rainfall, PET, 1/2 PET at Yadgir Taluk, Yadgir District

Sl. No.	Months	Rainfall	PET	1/2 PET
1	January	4.30	86.0	43.0
2	February	2.30	125.5	62.7
3	March	15.10	166.0	83.0
4	April	18.50	179.8	89.9
5	May	36.0	198.8	97.9
6	June	118.0	175.1	87.5
7	July	171.80	156.3	78.1
8	August	182.9	150.3	75.1
9	September	179.7	142.0	71.0
10	October	105.3	138.5	69.2
11	November	26.4	97.60	48.6
12	December	6.0	80.90	40.4
Total		866.3		




Fig 2.3 Rainfall distribution in Yadgir Taluk, Yadgir District

2.6 Natural Vegetation

The natural vegetation is sparse comprising few tree species, shrubs and herbs. The mounds, ridges and boulders occupy very sizeable area which is under thin to moderately thick forest vegetation. Still, there are some remnants of the past forest cover which can be seen in patches in some ridges and hillocks in the microwatershed (Fig 2.4).

Apart from the continuing deforestation, the presence of large population of goats, sheep and other cattle in the microwatershed is causing vegetative degradation of whatever little vegetation left in the area. The uncontrolled grazing has left no time for the regeneration of the vegetative cover. This leads to the accelerated rate of erosion on the hill slopes resulting in the formation of deep gullies in the foot slopes that eventually result in the heavy siltation of tanks and reservoirs in the microwatershed.



Fig 2.4 Natural vegetation of Bhimanahalli-2 Microwatershed

2.7 Land Utilization

About 72 per cent area (Table 2.2) in Yadgir district is cultivated at present. An area of about 2 per cent is permanently under pasture, 20 per cent under current fallows and 6 per cent under non-agricultural land and 5 per cent under currently barren. Forests occupy an area of about 7 per cent and the tree cover is in a very poor state. Most of the mounds, ridges and bouldery areas have very poor vegetative cover. Major crops grown in the area are sorghum, maize, cotton, sunflower, groundnut, red gram, mango, pomegranate, marigold and sapota. While carrying out land resource inventory, the land use/land cover particulars are collected from all the survey numbers and a current land use map of the microwatershed is prepared. The current land use map prepared shows the arable and non-arable lands, other land uses and different types of crops grown in the area. The current land use map of Bhimanahalli-2 microwatershed is presented in Fig.2.5. The different crops and cropping systems adopted in the microwatershed are presented in Figures 2.6 a & b.

Table 2.2 Land Utilization in Yadgir District

Sl. No.	Agricultural land use	Area (ha)	Per cent
1	Total geographical area	516088	-
2	Total cultivated area	373617	72.4
3	Area sown more than once	74081	14.3
4	Cropping intensity	-	119.8
5	Trees and grooves	737	0.14
6	Forest	33773	6.54
7	Cultivable wasteland	2385	0.46
8	Permanent Pasture land	11755	2.28
9	Barren land	27954	5.41
10	Non- Agriculture land	29623	5.73
11	Current Fallows	105212	20.4




Fig.2.5 Current Land Use map of Bhimanahalli-2 Microwatershed



Fig. 2.6 a. Different Crops and Cropping Systems in Bhimanahalli-2 Microwatershed




Fig. 2.6 b. Different Crops and Cropping Systems in Bhimanahalli-2 Microwatershed

SURVEY METHODOLOGY

The purpose of land resource inventory is to delineate similar areas (soil series and phases), which respond or expected to respond similarly to a given level of management. This was achieved in Bhimanahalli-2 microwatershed by the detailed study of all the soil characteristics (depth, texture, colour, structure, consistence, coarse fragments, porosity, soil reaction, soil horizons etc.) and site characteristics (slope of the land, erosion, drainage, occurrence of rock fragments etc.) followed by grouping of similar areas based on soil-site characteristics into homogeneous (management units) units, and showing the area extent and their geographic distribution on the microwatershed cadastral map. The detailed survey at 1:7920 scale was carried out in an area of 573 ha. The methodology followed for carrying out land resource inventory was as per the guidelines given in Soil Survey Manual (IARI, 1971; Soil Survey Staff, 2006; Natarajan *et al.*, 2015) which is briefly described below.

3.1 Base Maps

The detailed survey of the land resources occurring in the microwatershed was carried out by using digitized cadastral and IRS satellite imagery map as base supplied by KSRSAC. The cadastral map shows field boundaries with their survey numbers, location of tanks, streams and other permanent features of the area (Fig. 3.1). Apart from the cadastral map, remote sensing data products from Cartosat-1 and LISS IV merged at the scale of 1:7920 were used in conjunction with the cadastral map to identify the landscapes, landforms and other surface features. The imagery helped in the identification and delineation of boundaries between hills, uplands and lowlands, water bodies, forest and vegetated areas, roads, habitations and other cultural features of the area (Fig. 3.2). The cadastral map was overlaid on the satellite imagery (Fig. 3.3) that helps to identify the parcel boundaries and other permanent features. Apart from cadastral maps and images, toposheets of the area (1:50,000 scale) were also used for initial traversing, identification of geology and landforms, drainage features, present land use and also for selection of transects in the microwatershed.

3.2 Image Interpretation for Physiography

False Colour Composites (FCCs) of Cartosat-I and LISS-IV merged satellite data covering microwatershed area was visually interpreted using image interpretation elements and all the available collateral data with local knowledge. The delineated physiographic boundaries were transferred on to a cadastral map overlaid on satellite imagery. Physiographically, the area has been identified as granite gneiss and alluvial landscapes. It was divided into five landforms, *viz*; ridges and mounds, gently and very gently sloping uplands and lowlands based on slope and image characteristics. They were further subdivided into physiographic/image interpretation units based on image characteristics. The image interpretation legend for physiography is given below.

Image Interpretation Legend for Physiography

G- Granite Gneiss Landscape

G1	Hills/ Ridges/ Mounds
G11	Summits
G12	Side slopes
	G121 Side slopes with dark grey tones
G2	Uplands
G21	Summits
G22	Gently sloping uplands
	G221 Gently sloping uplands, yellowish green (eroded)
	G222 Gently sloping uplands, yellowish white (severely eroded)
G23	Very gently sloping uplands
	G231 Very gently sloping uplands, yellowish green
	G232 Very gently sloping uplands, medium green and pink
	G233 Very gently sloping uplands, pink and green (scrub land)
	G234 Very gently sloping uplands, medium greenish grey
	G235 Very gently sloping uplands, yellowish white (eroded)
	G236 Very gently sloping uplands, dark green
	G237 Very gently sloping uplands, medium pink (coconut garden)
	G238 Very gently sloping uplands, pink and bluish white (eroded)
G3	Valleys/ lowlands
G31	Valleys, pink tones
G32	Valleys gray mixed with pink tones

DSe – Alluvial Landscape

DSe 1 – Summit

DSe 11 –

DSe 12 –

DSe 2 – Very gently sloping

DSe 21 – Very gently sloping, dark gray tone

DSe 22 – Very gently sloping, medium gray tone

DSe 23 – Very gently sloping, yellowish grey tone

DSe 24 – Very gently sloping, whitish grey tone

DSe 25 – Very gently sloping, whitish/ eroded/ calcareous tone

DSe 26- Very gently sloping, medium pink

DSe 3 – Valley/ Lowland

DSe 31 – Whitish gray/Calcareous

DSe 32 – Gray with pink patches

DSe 33 – Medium gray tone

DSe 34 – Lightish gray tone

DSe 35 – Dark gray tone




Fig 3.1 Scanned and Digitized Cadastral map of Bhimanahalli-2 Microwatershed




Fig.3.2 Satellite Image of Bhimanahalli-2 Microwatershed




Fig.3.3 Cadastral map overlaid on IRS PAN+LISS IV merged imagery of Bhimanahalli-2 Microwatershed

3.3 Field Investigation

The field boundaries and survey numbers given on the cadastral sheet were located on the ground by following permanent features like roads, cart tracks, *nallas*, streams, tanks etc., and wherever changes were noticed, they were incorporated on the microwatershed cadastral map. Preliminary traverse of the microwatershed was carried out with the help of cadastral map, imagery and toposheets. While traversing, landforms and physiographic units identified were checked and preliminary soil legend was prepared by studying soils at few selected places. Then, intensive traversing of each physiographic unit like hills, ridges, uplands and valleys was carried out. Based on the variability observed on the surface, transects (Fig. 3.4) were selected across the slope covering all the landform units in the microwatershed (Natarajan and Dipak Sarkar, 2010).




Fig: 3.4. Location of profiles in a transect

In the selected transect, soil profiles were located (Fig. 3.4) at closely spaced intervals to take care of any change in the land features like break in slope, erosion, gravel, stones etc. In the selected sites, profiles (vertical cut showing the soil layers from surface to the rock) were opened upto 200 cm or to the depth limited by rock or hard substratum and studied in detail for all their morphological and physical characteristics. The soil and site characteristics were recorded for all profile sites on a standard proforma as per the guidelines given in USDA Soil Survey Manual (Soil Survey Staff, 2012). Apart from the transect study, profiles were also studied at random, almost like in a grid pattern, outside the transect areas.

Based on the soil characteristics, the soils were grouped into different soil series. Soil series is the most homogeneous unit having similar horizons and properties and behaves similarly for a given level of management. Soil depth, texture, colour, kind of horizon and horizon sequence, calcareousness, amount and nature of gravel present, nature of substratum etc, were used as the major differentiating characteristics for identifying soil series occurring in the area. The differentiating characteristics used for identifying the soil series are given in Table 3.1. Based on the above characteristics, 11 soil series were identified in the Bhimanahalli-2 microwatershed.

Table 3.1 Differentiating Characteristics used for identifying Soil Series
(Characteristics are of Series Control Section)

Soils of Granite gneiss Landscape							
Sl. no	Soil Series	Depth (cm)	Colour (moist)	Texture	Gravel (%)	Horizon sequence	Calcareousness
1	BDP (Baddeppalli)	<25	7.5YR 3/2,3/4 5YR 3/4	scl	<15	Ap-AC	es
2	HTK (Hattikuni)	25-50	10YR 4/6, 4/4 7.5YR 4/4, 3/3	sl	10-25	Ap-AC	-
3	BDL (Badiyala)	25-50	7.5YR2.5/3,2.5/ 3/3,10YR 3/4,4/3	sl	<15	Ap-Bw	e
4	DSB (Dastharabad)	25-50	7.5YR 3/3	g c	35-60	Ap-Bt-Cr	-
5	VNK (Vanakanahalli)	25-50	2.5YR 3/4	sc	<15	Ap-Bt-Cr	-
6	JNK (Jinkera)	50-75	10YR 3/1,3/2 7.5YR 3/4	scl	<15	Ap-Bw	e
7	NGP (Naglapur)	100-150	10YR 3/2,3/1,2/1	c	<15	Ap-Bss	es
8	ANR (Anur)	100-150	10YR 4/3,4/1	c	<15	Ap-Bw	es
9	MDG (Mundargi)	100-150	10YR 4/4,3/3 7.5YR 4/4	scl	<15	Ap-Bw	-
10	BMN (Bhimanahalli)	>150	10YR 3/1	c	<15	Ap-Bss	es
Soils of Alluvial Landscape							
11	BLD (Balched)	50-75	10 YR 3/2,2/1	cl	<15	Ap-Bw	e

3.4 Soil Mapping

The area under each soil series was further separated into soil phases and their boundaries delineated on the cadastral map based on the variations observed in the texture of the surface soil, slope, erosion, presence of gravel, stoniness etc. A soil phase is a subdivision of soil series based mostly on surface features that affect its use and management. The soil mapping units are shown on the map (Fig.3.5) in the form of symbols. During the survey many profile pits, few minipits and a few auger bores representing different landforms occurring in the microwatershed were studied. In addition to the profile study, spot observations in the form of minipits, road cuts, terrace cuts etc., were studied to validate the soil boundaries on the soil map.

The soil map shows the geographic distribution of 13 mapping units representing 11 soil series occurring in the microwatershed. The soil map unit (soil legend) description is presented in Table 3.2. The soil phase map (management units) shows the distribution of 13 soil phases mapped in the microwatershed. Each mapping unit (soil phase) delineated on the map has similar soil and site characteristics. In other words, all the farms or survey numbers included in one phase will have similar management needs and have to be treated accordingly.

3.5 Land Management Units

The 13 soil phases identified and mapped in the microwatershed were grouped into 8 Land Management Units (LMU's) for the purpose of preparing a Proposed Crop Plan for sustained development of the microwatershed. The database (soil phases) generated under LRI was utilized for identifying Land Management Units (LMU's) based on the management needs. One or more than one soil site characteristic having influence on the management have been chosen for identification and delineation of LMUs. For Bhimanahalli-2 microwatershed, five soil and site characteristics, namely soil depth, soil texture, slope, erosion and gravel content have been considered for defining LMUs. The land use classes are expected to behave similarly for a given level of management.

3.6 Laboratory Characterization

Soil samples were collected from representative master profiles for laboratory characterization by following the methods outlined in the Laboratory Manual (Sarma *et al.*, 1987). Surface soil samples collected from farmer's fields for fertility status (major and micronutrients) at 320 m grid interval in the year 2018 were analyzed in the laboratory (Katyala and Rattan, 2003). By linking the soil fertility data to the survey numbers through GIS, soil fertility maps were generated by using Kriging method for the microwatershed.

Table 3.2 Soil map unit description of Bhimanahalli-2 Microwatershed

*Soil map unit No.	Soil Series	Soil Phase	Mapping Unit Description	Area in ha (%)
Soils of Granite and Granite Gneiss Landscape				
	BDP	Baddeppalli soils are very shallow (<25 cm), well drained, have dark brown to dark reddish brown, calcareous sandy clay loam soils occurring on very gently sloping uplands under cultivation		68 (11.9)
1		BDPiB2	Sandy clay surface, slope 1-3%, moderate erosion	68 (11.9)
	HTK	Hattikuni soils are shallow (25-50 cm), well drained, have dark yellowish brown sandy loam soils occurring on very gently sloping uplands under cultivation		7 (1.17)
165		HTKcB2	Sandy loam surface, slope 1-3%, moderate erosion	7 (1.17)
	BDL	Badiyala soils are shallow (25-50 cm), well drained, have dark brown to very dark brown and dark yellowish brown, slightly calcareous sandy loam soils occurring on very gently to gently sloping uplands under cultivation		32 (5.55)
5		BDLiB2	Sandy clay surface, slope 1-3%, moderate erosion	32 (5.55)
	DSB	Dastharabad soils are shallow (25-50 cm), well		45

*Soil map unit No.	Soil Series	Soil Phase	Mapping Unit Description	Area in ha (%)
			drained, have dark brown to very dark brown, gravelly clay soils occurring on very gently to gently sloping uplands under cultivation	(7.84)
108		DSBiB2	Sandy clay surface, slope 1-3%, moderate erosion	45 (7.84)
	VNK		Vanakanahalli soils are shallow (25-50 cm), well drained, have dark reddish brown, sandy clay red soils occurring on very gently to moderately sloping uplands under cultivation	13 (2.19)
10		VNKiB2	Sandy clay surface, slope 1-3%, moderate erosion	13 (2.19)
	JNK		Jinkera soils are moderately shallow (50-75 cm), well drained, have dark brown to very dark grayish brown, slightly calcareous sandy clay loam soils occurring on very gently sloping uplands under cultivation	21 (3.73)
110		JNKhB2	Sandy clay loam surface, slope 1-3%, moderate erosion	21 (3.73)
	NGP		Nagalapur soils are deep (100-150 cm), moderately well drained, have very dark gray to very dark grayish brown, black calcareous cracking clay soils occurring on very gently sloping uplands under cultivation	19 (3.25)
49		NGPmB2	Clay surface, slope 1-3%, moderate erosion	19 (3.25)
	ANR		Anur soils are deep (100-150 cm), moderately well drained, have dark gray to brown, calcareous sodic cracking clay soils occurring on very gently sloping uplands under cultivation	3 (0.48)
55		ANRiB2	Sandy clay surface, slope 1-3%, moderate erosion	3 (0.48)
	MDG		Mundargi soils are deep (100-150 cm), well drained, have brown to dark yellowish brown, sandy clay loam soils occurring on very gently sloping uplands under cultivation	54 (9.49)
148		MDGhB2	Sandy clay loam surface, slope 1-3%, moderate erosion	10 (1.83)
170		MDGmB1	Clay surface, slope 1-3%, slight erosion	44 (7.66)
	BMN		Bhimanahalli soils are very deep (>150 cm), moderately well drained, have very dark gray, calcareous cracking clay black soils occurring on very gently sloping uplands under cultivation	38 (6.64)
159		BMNmA1	Clay surface, slope 0-1%, slight erosion	1 (0.21)
62		BMNmB2	Clay surface, slope 1-3%, moderate erosion	37 (6.43)

*Soil map unit No.	Soil Series	Soil Phase	Mapping Unit Description	Area in ha (%)
Soils of Alluvial Landscape				
	BLD		Balched soils are moderately shallow (50-75 cm), moderately well drained, have black to very dark grayish brown, slightly calcareous clay loam soils, occurring on very gently to gently sloping plains under cultivation	142 (24.85)
76		BLDmB2	Clay surface, slope 1-3%, moderate erosion	142 (24.85)
900		Forest		77 (13.38)
993		Quarry		12 (2.03)
999		Rock outcrops	Rock lands, both massive and bouldery with little or no soil	43 (7.5)




Fig 3.5 Soil Phase or Management Units - Bhimanahalli-2 Microwatershed

THE SOILS

Detailed information pertaining to the nature, extent and their distribution of different kinds of soils occurring in Bhimanahalli-2 microwatershed is provided in this chapter. The microwatershed area has been identified as granite gneiss and alluvial landscapes based on geology. In all, 11 soil series are identified. Soil formation is the result of the combined effect of environmental and terrain factors that are reflected in soil morphology. In the granite gneiss landscape, it is by parent material, relief and climate.

A brief description of each of the 11 soil series identified followed by 13 soil phases (management units) mapped under each series are furnished below. The physical and chemical characteristics of soil series identified in Bhimanahalli-2 microwatershed are given in Table 4.1 along with soil classification. The soils in any one map unit differ from place to place in their depth, texture, slope, gravelliness, erosion or any other site characteristic that affect management. The soil phase map can be used for identifying the suitability of areas for growing specific crops or for other alternative uses and also for deciding the type of conservation structures needed. The detailed information on soil and site-characteristics like soil depth, surface soil texture, slope, erosion, gravelliness, AWC, LCC etc, with respect to each of the soil phase identified is given village/survey number wise for the microwatershed in Appendix-I.

4.1 Soils of granite gneiss landscape

In this landscape, 10 soil series are identified and mapped. BDP series occupies maximum area of 68 ha (12%) followed by MDG 54 ha (10%), DSB 45 ha (8%), BMN 38 ha (7%), BDL 32 ha (6%), JNK 21 ha (4%), NGP 19 ha (3%), VNK 13 ha (2%), HTK 7 ha (1%) and ANR 3 ha (<1%). Brief description of each series identified and number of soil phases mapped is given below.

4.1.1 Baddeppalli (BDP) Series: Baddeppalli soils are very shallow (<25cm), well drained, have dark brown to dark reddish brown, calcareous sandy clay loam soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation. The Baddeppalli series has been classified as a member of the loamy, mixed (calcareous), isohyperthermic family of Lithic Ustorthents.

The thickness of the soil is less than 25 cm. Its colour is in 7.5 YR and 5 YR hue with value 3 and chroma 2 to 4. The texture varies from sandy clay loam to sandy clay and is calcareous. The available water capacity is very low (<50 mm/m). One soil phase was identified and mapped.



Landscape and Soil Profile characteristics of Baddeppalli (BDP) Series

4.1.2 Hattikuni (HTK) Series: Hattikuni soils are shallow (25-50 cm), well drained, have dark brown to dark yellowish brown sandy loam soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation. The Hattikuni series has been classified as a member of the mixed, isohyperthermic family of Lithic Ustipsamments.

The thickness of the soil ranges from 36 to 50 cm. The thickness of A horizon ranges from 8 to 12 cm. Its colour is in 10YR and 7.5 YR hue with value 3 to 4 and chroma 4 to 6. The texture varies from loamy sand to sandy loam. The thickness of subsurface horizon ranges from 28 to 42 cm. Its colour is in 10 YR and 7.5 YR hue with value 3 to 4 and chroma 4 to 6. Its texture varies from loamy sand to sand and sandy loam. The available water capacity is very low (<50 mm/m). One soil phase was identified and mapped.



Landscape and Soil Profile characteristics of Hattikuni (HTK) Series

4.1.3 Badiyala (BDL) Series: Badiyala soils are shallow (25-50 cm), well drained, have very dark brown to dark yellow brown and dark brown, slightly calcareous sandy loam soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation. The Badiyala series has been classified as a member of the coarse-loamy, mixed, isohyperthermic family of Fluventic Haplustepts

The thickness of the solum ranges from 28 to 50 cm. The thickness of A horizon ranges from 4 to 12 cm. Its colour is in 10YR hue with value 3 to 4 and chroma 3 to 4. The texture is loamy sand, sandy clay loam and sandy clay. The thickness of B horizon ranges from 27 to 45 cm. Its colour is in 10 YR and 7.5 YR hue with value 2 to 4 and chroma 3 to 4. Its texture is sandy loam to sandy clay loam and is slightly calcareous. The available water capacity is very low (<50mm/m). One soil phase was identified and mapped.



Landscape and Soil Profile characteristics of Badiyala (BDL) Series

4.1.4 Dastharabad (DSB) Series: Dastharabad soils are shallow (25-50 cm), well drained, have dark brown, gravelly clay soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation. The Dastharabad series has been classified as a member of the clayey-skeletal, mixed, isohyperthermic family of (Paralithic) Haplustalfs.

The thickness of the solum ranges from 28 to 50 cm. The thickness of A horizon ranges from 9 to 14 cm. Its colour is in 10 YR and 7.5 YR hue with value and chroma of 3 to 4. The texture varies from sandy loam to sandy clay. The thickness of B horizon ranges from 28 to 40 cm. Its colour is in 7.5 YR hue with value 3 and chroma 3 to 4. The texture is sandy clay to clay with 35-60 per cent gravel. The available water capacity is very low (<50 mm/m). One soil phase was identified and mapped.



Landscape and Soil Profile characteristics of Dastharabad (DSB) Series

4.1.5 Vanakanahalli (VNK) Series: Vanakanahalli soils are shallow (25-50 cm), well drained, have dark reddish brown sandy clay red soils. They have developed from weathered granite gneiss and occur on very gently sloping uplands under cultivation. The Vanakanahalli series has been classified as a member of the clayey, mixed isohyperthermic family of (Paralthic) Haplustalfs.

The thickness of the solum ranges from 25 to 49 cm. The thickness of A horizon ranges from 7 to 16 cm. Its colour is in 2.5 YR and 5 YR with value 3 and chroma 2 to 4. The texture is sandy loam to sandy clay loam and sandy clay. The thickness of B horizon ranges from 20 to 40 cm. Its colour is in 2.5 YR and 5 YR hue with value 3 to 4 and chroma 3 to 4. Its texture is sandy clay. The available water capacity is very low (<50 mm/m). One soil phase was identified and mapped.



Landscape and Soil Profile characteristics of Vanakanahalli (VNK) Series

4.1.6 Jinkera (JNK) Series: Jinkera soils are moderately shallow (50-75 cm), well drained, have very dark gray to very dark grayish brown and dark brown, slightly calcareous sandy clay loam soils. They are developed from weathered granite gneiss and occur on very gently sloping uplands under cultivation. The Jinkera series has been classified as a member of the fine-loamy, mixed, isohyperthermic family of Typic Haplustepts.

The thickness of the solum ranges from 51-75 cm. Thickness of A horizon ranges from 6 to 11 cm. Its colour is in hue 10 YR and 7.5 YR with value and chroma of 3 to 4. The texture varies from sandy loam to sandy clay. The thickness of B horizon ranges from 53 to 66 cm. Its colour is in 10 YR and 7.5 YR hue with value and chroma of 2 to 4. The texture varies from sandy clay loam to sandy clay and is slightly calcareous. The available water capacity is low (51-100 mm/m). One soil phase was identified and mapped.



Landscape and Soil Profile characteristics of Jinkera (JNK) Series

4.1.7 Naglapur (NGP) Series: Naglapur soils are deep (100-150 cm), moderately well drained, have black to very dark grayish brown, calcareous cracking clay soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation. The Naglapur series has been classified as a member of the very fine, smectitic (calcareous), isohyperthermic family of Typic Haplusterts.

The thickness of the solum ranges from 110 to 150 cm. The thickness of A horizon ranges from 6 to 25 cm. Its colour is in 10 YR hue with value 3 to 5 and chroma 1 to 3. The texture varies from sandy loam to sandy clay and clay. The thickness of B horizon ranges from 110 to 141 cm. Its colour is in 10 YR hue with value 2 to 3 and chroma 1 to 2. Texture is clay and is calcareous. The available water capacity is very high (>200 mm/m). One soil phase was identified and mapped.



Landscape and Soil Profile characteristics of Naglapur (NGP) Series

4.1.8 Anur (ANR) Series: Anur soils are deep (100-150 cm), moderately well drained, have dark gray to dark brown, calcareous sodic clay soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation. The Anur series has been classified as a member of the fine, mixed (calcareous), isohyperthermic family of Typic Haplustepts.

The thickness of the solum ranges from 102 to 148 cm. The thickness of A-horizon ranges from 9 to 17 cm. Its colour is in 10 YR hue with value 3 to 4 and chroma 2 to 4. The texture ranges from loamy sand to sandy clay loam and sandy clay and are calcareous. The thickness of B horizon ranges from 102 to 135 cm. Its colour is in 10 YR hue with value 3 to 5 and chroma 1 to 6. Texture is sandy clay loam to sandy clay and clay and is calcareous sodic soils. The available water capacity is very high (>200 mm/m). One soil phase was identified and mapped.



Landscape and Soil Profile characteristics of Anur (ANR) Series

4.1.9 Mundargi (MDG) Series: Mundargi soils are deep (100-150 cm), well drained, have dark brown to dark yellowish brown, sandy clay loam soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation. The Mundargi series has been classified as a member of the fine-loamy, mixed, isohyperthermic family of Fluventic Haplustepts.

The thickness of the solum ranges from 100 to 149 cm. The thickness of A horizon ranges from 8 to 20 cm. Its colour is in 10 YR hue with value 3 and chroma 1 to 4. The texture ranges from sandy loam to sandy clay loam and sandy clay. The thickness of B horizon ranges from 105 to 140 cm. Its colour is in 10 YR hue with value 3 to 4 and chroma 2 to 4. The texture varies from sandy loam to sandy clay loam and sandy clay. The available water capacity is very high (>200 mm/m). Two soil phases were identified and mapped.



Landscape and Soil Profile characteristics of Mundargi (MDG) Series

4.1.10 Bhimanahalli (BMN) Series: Bhimanahalli soils are very deep (>150 cm), moderately well drained, have very dark gray calcareous cracking clay soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation. The Bhimanahalli series has been classified as a member of the fine, smectitic (calcareous), isohyperthermic family of Typic Haplusterts.

The thickness of the solum is more than 150 cm. The thickness of A horizon ranges from 6 to 13 cm. Its colour is in 10 YR hue with value 3 to 4 and chroma 1 to 2 with clay texture. The thickness of B horizon ranges from 163 to 176 cm. Its colour is in 10 YR hue with value 2 to 4 and chroma 1. Its texture is clay and is calcareous. The available water capacity is very high (>200 mm/m). Two soil phases were identified and mapped.



Landscape and Soil Profile characteristics of Bhimanahalli (BMN) Series

4.2 Soils of alluvial landscape

In this landscape, only one soil series is identified and mapped. BLD series occupies an area of 142 ha (25%). Brief description of this series identified and number of soil phases mapped is given below.

4.2.1 Balched (BLD) Series: Balched soils are moderately shallow (50-75 cm), moderately well drained, have black to very dark grayish brown, slightly calcareous clay loam soils. They are developed from alluvium and occur on very gently to gently sloping plains under cultivation. The Balched series has been classified as a member of the fine, mixed, isohyperthermic family of Typic Haplustepts.

The thickness of the solum ranges from 50-75 cm. Thickness of A horizon ranges from 5 to 10 cm. Its colour is in hue 10 YR and 7.5 YR with value 3 to 4 and chroma 1 to 3. The texture varies from sandy clay to clay. The thickness of B horizon ranges from 41 to 69 cm. Its colour is in 10 YR hue with value 2 to 4 and chroma 1 to 2. The texture is clay loam and is slightly calcareous. The available water capacity is medium (101-150 mm/m). One soil phase was identified and mapped.



Landscape and Soil Profile characteristics of Balched (BLD) Series

Table: 4.1 Physical and Chemical Characteristics of Soil Series identified in Bhimanahalli-2 microwatershed

Soil Series: Baddeppalli (BDP) **Pedon:** R-11

Location: 16°43'84.4"N 77°14'06.4"E, Halagera village, Yadgir hobli, Yadgir taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bengaluru **Classification:** Loamy, mixed (calcareous), isohyperthermic Lithic Ustorthents

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-16	Ap	58.67	17.02	24.31	19.03	13.74	9.62	10.57	5.71	<15	scl	16.19	8.18

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO ₃	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP
	Water	CaCl ₂	M KCl				Ca	Mg	K	Na	Total				
				dS m ⁻¹	%	%	cmol kg ⁻¹						%	%	
0-16	8.58	-	-	0.262	1.60	7.67	-	-	0.24	0.06	-	18.10	0.74	100	0.35

Contd...

Soil Series: Hattikuni (HTK), Pedon: R-7

Location: 16°50'46.5"N 77°10'16.4"E, Yaddalli village, Hattikuni hobli, Yadgir taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bengaluru **Classification:** Mixed, isohyperthermic Lithic Ustipsamments

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-12	Ap	90.89	5.62	3.49	8.50	13.46	29.86	29.55	9.51	20	s	7.73	3.16
12-22	A1	89.97	6.53	3.50	7.19	13.48	29.48	29.79	10.03	20	s	8.00	3.05
22-45	A2	87.20	6.43	6.38	11.09	14.42	31.55	7.16	22.98	40	ls	7.67	3.96

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO ₃	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP			
	Water	CaCl ₂	M KCl				dS m ⁻¹	%	%	Ca	Mg					K	Na	Total
										cmol kg ⁻¹								
0-12	6.81	-	-	0.062	0.07	-	2.35	0.50	0.16	0.01	3.02	3.0	0.86	100	0.38			
12.0-22	6.80	-	-	0.050	0.21	-	1.67	0.30	0.09	0.01	2.07	2.4	0.69	86.30	0.45			
22-45	6.85	-	-	0.044	0.19	-	1.82	0.42	0.10	0.06	2.40	2.6	0.41	92.41	2.17			

Contd...

Soil Series: Badiyala (BDL) **Pedon:** R-5

Location: 16°37'10.0"N 77°20'21.5", Gudalagunta village, Balichakra hobli, Yadgir taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bengaluru **Classification:** Coarse-loamy, mixed, isohyperthermic Fluventic Haplustepts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-12	Ap	87.13	7.04	5.83	10.03	24.32	23.61	23.51	5.67	<15	ls	6.27	2.44
12-28	Bw	64.63	13.30	22.07	6.74	13.07	22.30	17.01	5.50	<15	scl	16.34	7.83
28-50	BC	73.11	12.02	14.87	3.93	16.03	26.89	18.41	7.86	<15	sl	12.94	5.47

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO ₃	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP			
	Water	CaCl ₂	M KCl				dS m ⁻¹	%	%	Ca	Mg					K	Na	Total
										cmol kg ⁻¹								
0-12	6.20	-	-	0.074	1.00	0.00	2.80	0.98	0.14	0.01	3.92	4.20	0.72	93	0.20			
12-28	9.04	-	-	0.253	0.80	3.20	-	-	0.16	0.69	-	16.90	0.77	100	4.09			
28-50	9.41	-	-	0.364	1.10	3.60	-	-	0.16	1.39	-	11.10	0.75	100	12.52			

Contd...

Soil Series: Dastharabad (DSB) **Pedon:** R-17

Location: 16°31' 98.6"N 77°22'93.0"E, Duppalli village, Sydhapura hobli, Yadgir taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bengaluru **Classification:** Clayey-skeletal, mixed, isohyperthermic (Paralithic) Haplustalfs

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-6	Ap	90.51	4.84	4.64	7.06	8.07	37.24	26.03	12.11	35	s	5.32	1.59
6-17	Bt1	49.11	8.08	42.81	10.67	15.44	10.00	8.44	4.56	20	sc	20.68	13.16
17-43	Bt2	39.54	2.84	57.63	12.89	9.14	7.71	6.83	2.97	50	c	26.69	18.50

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO ₃	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP			
	Water	CaCl ₂	M KCl				dS m ⁻¹	%	%	Ca	Mg					K	Na	Total
										cmol kg ⁻¹								
0-6	5.93	-	-	0.04	0.67	0.00	2.00	0.54	0.07	0.01	2.61	3.60	0.78	73	0.14			
6-17	7.31	-	-	0.110	0.91	0.91	11.19	3.37	0.12	0.49	15.00	15.20	0.36	100	3.22			
17-43	6.64	-	-	0.048	0.76	0.00	18.81	5.57	0.23	0.09	24.70	24.90	0.43	99	0.38			

Contd...

Soil Series: Vanakanahalli (VNK) **Pedon:** R-15

Location: 16°43'49.5"N 77°17'17.9"E, Yaleri village, Balichakra hobli, Yadgiri taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bengaluru **Classification:** Clayey, mixed isohyperthermic Paralthic Haplustalfs

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-18	Ap	82.61	8.09	9.30	6.77	8.59	21.13	34.58	11.53	-	ls	8.85	3.53
18-61	Bt	54.51	8.73	36.77	4.93	6.18	14.15	20.75	8.49	-	sc	18.88	11.63

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO ₃	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP			
	Water	CaCl ₂	M KCl				dS m ⁻¹	%	%	Ca	Mg					K	Na	Total
0-18	5.37	-	-	0.11	0.60	0.00	2.96	1.45	0.13	0.14	4.68	6.27	0.67	75	2.22			
18-61	4.71	-	-	0.05	0.81	0.00	5.56	2.24	0.10	0.05	7.95	13.31	0.36	60	0.38			

Contd...

Soil Series: Jinkera (JNK) **Pedon:** R-1

Location: 16°45'13.5"N 77°10'59.8"E, Varkanahalli village, Yadgir hobli, Yadgir taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bengaluru **Classification:** Fine-loamy, mixed, isohyperthermic Typic Haplustepts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-15	Ap	66.84	13.62	19.54	12.15	21.22	11.23	12.56	9.68	10	sl	14.42	7.70
15-38	Bw1	59.08	12.11	28.81	12.53	12.42	17.85	8.77	7.52	20	scl	18.21	12.23
38-52	Bw2	68.21	11.68	20.11	17.90	21.81	10.60	10.80	7.10	10	scl	14.54	8.96

Depth (cm)	pH (1:2.5)			E.C. (1:2.5) dS m ⁻¹	O.C. %	CaCO ₃ %	Exchangeable bases					CEC	CEC/Clay	Base saturation %	ESP %
	Water	CaCl ₂	M KCl				Ca	Mg	K	Na	Total				
0-15	8.42	-	-	0.148	0.70	0.65	-	-	0.15	0.03	-	14.50	0.74	100	0.18
15-38	8.38	-	-	0.226	0.31	2.21	-	-	0.09	0.23	-	21.70	0.75	100	1.05
38-52	8.40	-	-	0.195	0.25	1.17	-	-	0.07	0.19	-	15.90	0.79	100	1.23

Contd...

Soil Series: Naglapur (NGP) **Pedon:** R-8

Location: 16°52'84.1"N 77°22'99.4"E, Gurumitkal village, Gurumitkal hobli, Yadgir taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bengaluru, **Classification:** Very fine, smectitic (calcareous), isohyperthermic Typic Haplusterts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-10	Ap	7.53	19.88	72.59	1.00	0.78	0.89	2.10	2.77	-	c	44.31	32.79
10-35	Bss1	6.55	18.76	74.68	0.80	0.92	0.80	1.72	2.30	-	c	43.09	31.62
35-60	Bss2	6.58	21.05	72.37	0.69	0.46	1.04	1.50	2.89	-	c	46.52	32.52
60-102	Bss3	7.48	19.74	72.78	1.61	1.38	0.69	1.61	2.19	-	c	51.12	35.62

Depth (cm)	pH (1:2.5)			E.C. (1:2.5) dS m ⁻¹	O.C. %	CaCO ₃ %	Exchangeable bases					CEC	CEC/Clay	Base saturation %	ESP %
	Water	CaCl ₂	M KCl				Ca	Mg	K	Na	Total				
0-10	7.42	-	-	0.24	0.84	1.30	-	-	0.84	0.15	-	67.10	0.92	100	0.22
10-35	8.52	-	-	0.291	0.64	2.86	-	-	0.17	0.29	-	65.20	0.87	100	0.45
35-60	7.89	-	-	0.134	0.62	4.55	-	-	0.15	0.20	-	65.00	0.90	100	0.30
60-102	8.68	-	-	0.213	0.54	8.32	-	-	0.17	0.15	-	64.10	0.88	100	0.24

Contd...

Soil Series: Anur (ANR) **Pedon:** R-15

Location: 16°32'45.0"N 77°23'57.4"E, Duppalli village, Sydhapura hobli, Yadgir taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bengaluru **Classification:** Fine, mixed (calcareous), isohyperthermic Typic Haplustepts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-18	Ap	64.60	13.44	21.96	7.33	10.42	18.68	20.12	8.05	<15	scl	16.59	7.96
18-49	Bw1	56.66	12.19	31.15	4.73	9.80	18.66	17.02	6.45	-	scl	33.38	13.51
49-95	Bw2	39.94	17.81	42.25	3.09	3.30	15.44	10.65	7.45	<15	c	44.68	25.23
95-123	Bw3	30.65	17.58	51.77	1.50	5.57	10.18	9.65	3.75	<15	c	54.94	32.07

Depth (cm)	pH (1:2.5)			E.C. (1:2.5) dS m ⁻¹	O.C. %	CaCO ₃ %	Exchangeable bases					CEC	CEC/Clay	Base saturation %	ESP %
	Water	CaCl ₂	M KCl				Ca	Mg	K	Na	Total				
0-18	10.17	-	-	0.365	0.48	6.11	-	-	0.25	3.52	-	19.90	0.91	100	7.08
18-49	10.32	-	-	1.38	0.30	6.76	-	-	0.21	16.03	-	24.60	0.79	100	26.07
49-95	10.08	-	-	2.55	0.17	6.11	-	-	0.33	21.49	-	32.60	0.77	100	26.36
95-123	9.92	-	-	2.56	0.12	7.93	-	-	0.51	26.03	-	36.00	0.70	100	28.92

Contd...

Soil Series: Mundargi (MDG) **Pedon:** R-2

Location: 16°46'82.4"N 77°04'85.2"E, Thumakura village, Yadgir hobli, Yadgir taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bengaluru **Classification:** Fine-loamy, mixed, isohyperthermic Fluventic Haplustepts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-9	Ap	81.23	12.97	5.80	4.84	10.19	14.83	37.94	13.42	<15	ls	11.75	3.31
9-20	A2	76.82	16.19	6.98	4.96	10.12	20.75	27.53	13.46	-	ls	14.52	3.99
20-46	Bw1	42.43	17.43	40.15	2.26	5.59	11.49	14.93	8.16	-	c	34.90	21.14
46-90	Bw2	54.51	16.56	28.93	4.72	5.03	19.92	16.67	8.18	-	scl	36.73	18.88
90-110	Bw3	53.69	11.00	35.30	9.57	9.89	16.23	13.01	4.99	-	sc	38.72	20.53

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO ₃	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP						
	Water	CaCl ₂	M KCl				dS m ⁻¹	%	%	Ca	Mg					K	Na	Total	cmol kg ⁻¹	%	%
0-9	8.2	-	-	0.399	0.44	0.78	-	-	0.16	0.38	-	4.90	0.84	100	3.08						
9-20	8.44	-	-	0.075	0.29	1.82	-	-	0.05	0.35	-	4.90	0.70	100	2.88						
20-46	9.39	-	-	0.451	0.32	2.73	-	-	0.12	5.22	-	20.77	0.52	100	10.06						
46-90	9.75	-	-	0.616	0.24	3.25	-	-	0.12	5.72	-	16.56	0.57	100	13.82						
90-110	9.72	-	-	0.725	0.24	3.64	-	-	0.14	6.84	-	19.76	0.56	100	13.836						

Contd...

Soil Series: Bhimanahalli (BMN) **Pedon:** R-3

Location: 16°31'82.4"N 77°12'70.8"E, Bheemanahalli village, Sydhapura hobli, Yadgir taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bengaluru **Classification:** Fine, smectitic (calcareous), isohyperthermic Typic Haplusterts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-8	Ap	20.34	19.94	59.72	2.68	5.03	3.75	5.25	3.64	-	c	50.19	33.49
8-40	Bss1	19.61	22.76	57.62	1.94	2.59	5.28	4.96	4.85	-	c	43.22	29.05
40-70	Bss2	21.25	17.65	61.10	3.02	5.26	3.91	5.48	3.58	-	c	44.30	30.25
70-120	Bss3	19.08	22.29	58.63	1.75	5.04	3.84	5.15	3.29	-	c	43.26	30.31
120-170	Bss4	11.11	20.44	68.45	2.04	1.93	1.70	2.83	2.61	-	c	51.33	33.51

Depth (cm)	pH (1:2.5)			E.C. (1:2.5) dS m ⁻¹	O.C. %	CaCO ₃ %	Exchangeable bases					CEC	CEC/Clay	Base saturation %	ESP %
	Water	CaCl ₂	M KCl				Ca	Mg	K	Na	Total				
0-8	8.2	-	-	0.284	0.72	4.94	-	-	1.20	0.34	-	52.70	0.88	100	0.65
8-40	8.44	-	-	0.139	0.40	7.28	-	-	0.30	0.48	-	52.06	0.90	100	0.93
40-70	8.32	-	-	0.202	0.40	6.37	-	-	0.18	0.40	-	52.52	0.86	100	0.77
70-120	9.3	-	-	0.282	0.36	6.89	-	-	0.27	0.38	-	50.97	0.87	100	0.75
120-170	8.47	-	-	0.305	0.37	8.19	-	-	0.28	0.91	-	58.19	0.85	100	1.57

Contd...

Soil Series: Balched (BLD) **Pedon:** R-40

Location: 16⁰44'19.4"N 77⁰19'40.9"E Yaleri village, Balichakra hobli, Yadgir taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bengaluru **Classification:** Fine, mixed, isohyperthermic Typic Haplustepts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-7	Ap	38.19	26.03	35.79	2.32	6.22	9.60	14.87	5.17	15	cl	22.13	11.07
7-28	Bw1	37.87	23.59	38.54	3.30	6.06	9.15	12.77	6.60	-	cl	23.75	14.43
28-54	Bw2	35.71	28.94	35.36	4.10	2.16	10.46	11.76	7.23	-	cl	25.47	16.56

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO ₃	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP			
	Water	CaCl ₂	M KCl				dS m ⁻¹	%	%	Ca	Mg					K	Na	Total
										cmol kg ⁻¹								
0-7	8.19	-	-	0.22	0.54	2.32	27.16	6.43	0.38	0.31	34.28	38.20	1.07	90	0.80			
7-28	8.56	-	-	0.14	0.42	3.18	29.26	6.83	0.14	0.51	36.75	39.91	1.04	92	1.27			
28-54	8.70	-	-	0.16	0.38	3.92	29.79	7.14	0.08	0.91	37.92	42.91	1.21	88	2.13			

INTERPRETATION FOR LAND RESOURCE MANAGEMENT

The most important soil and site characteristics that affect the land use and conservation needs of an area are land capability, soil depth, soil texture, coarse fragments, available water capacity, soil slope, soil erosion, soil reaction etc. These are interpreted from the data base generated through land resource inventory and several thematic maps are generated. These would help in identifying the areas suitable for growing crops and, soil and water conservation measures and structures needed thus helping to maintain good soil health for sustained crop production. The various interpretative and thematic maps generated are described below.

5.1 Land Capability Classification

Land capability classification is an interpretative grouping of soil map units (soil phases) mainly based on inherent soil characteristics, external land features and environmental factors that limit the use of land for agriculture, pasture, forestry, or other uses on a sustained basis (IARI, 1971). The land and soil characteristics used to group the land resources in an area into various land capability classes, subclasses and units are

Soil Characteristics: Depth, texture, gravelliness, calcareousness.

Land characteristics: Slope, erosion, drainage, rock outcrops.

Climate: Total rainfall and its distribution, and length of crop growing period.

The Land capability classification system is divided into land capability classes, subclasses and units based on the level of information available. Eight land capability classes are recognized. They are

Class I: They are very good lands that have no limitations or very few limitations that restrict their use.

Class II: They are good lands that have minor limitations and require moderate conservation practices.

Class III: They are moderately good lands that have moderate limitations that reduce the choice of crops or that require special conservation practices.

Class IV: They are fairly good lands that have very severe limitations that reduce the choice of crops or that require very careful management.

Class V: Soils in these lands are not likely to erode, but have other limitations like wetness that are impractical to remove and as such not suitable for agriculture, but suitable for pasture or forestry with minor limitations.

Class VI: The lands have severe limitations that make them generally unsuitable for cultivation, but suitable for pasture or forestry with moderate limitations.

Class VII: The lands have very severe limitations that make them unsuitable for cultivation, but suitable for pasture or forestry with major limitations.

Class VIII: Soil and other miscellaneous areas (rock lands) that have very severe limitations that nearly preclude their use for any crop production, but suitable for wildlife, recreation and installation of wind mills.

The land capability subclasses are recognised based on the dominant limitations observed within a given land capability class. The subclasses are designated by adding a lower case letter like 'e', 'w', 's', or 'c' to the class numeral. The subclass "e" indicates that the main hazard is risk of erosion, "w" indicates drainage or wetness as a limitation for plant growth, "s" indicates shallow soil depth, coarse or heavy textures, calcareousness, salinity/alkalinity or gravelliness and "c" indicates limitation due to climate.

The land capability subclasses have been further subdivided into land capability units based on the kinds of limitations present in each subclass. Ten land capability units are used in grouping the soil map units. They are stony or rocky (0), erosion hazard (slope, erosion) (1), coarse texture (sand, loamy sand, sandy loam) (2), fine texture (cracking clay, silty clay) (3), slowly permeable subsoil (4), coarse underlying material (5), salinity/alkali (6), stagnation, overflow, high ground water table (7), soil depth (8) and fertility problems (9). The capability units thus identified have similar soil and land characteristics that respond similarly to a given level of management. The soils of the microwatershed have been classified upto land capability subclass level.

The 13 soil map units identified in the Bhimanahalli-2 microwatershed are grouped under 3 land capability classes and 5 subclasses. An area about 442 ha (77%) in the microwatershed is suitable for agriculture, about 43 ha (8%) covered by rock outcrops, about 12 ha (2%) covered by quarry and about 77 ha (13%) covered by forest in the microwatershed. (Fig. 5.1).




Fig. 5.1 Land Capability map of Bhimanahalli-2 Microwatershed

Good lands (Class II) cover an area of 275 ha (48%) and are distributed in the major part of the microwatershed. They have minor limitations of soil and erosion. Moderately good lands (Class III) cover an area of 96 ha (17%) and are distributed in the central, northwestern and southern part of the microwatershed. They have minor limitations of soil and erosion. Fairly good lands (Class IV) cover an area of about 71 ha (12%) and are distributed in the northwestern and southern part of the microwatershed. They have very severe limitations of soil and erosion.

5.2 Soil Depth

Soil depth refers to the depth of the soil occurring above the parent material or hard rock. The depth of the soil determines the effective rooting depth for plants and in accordance with soil texture, mineralogy and gravel content, the capacity of the soil column to hold water and nutrient availability. Soil depth is one of the most important soil characteristic that is used in differentiating soils into different soil series. The soil depth classes used in identifying soils in the field are very shallow (<25 cm), shallow (25-50 cm), moderately shallow (50-75 cm), moderately deep (75-100 cm), deep (100-150 cm) and very deep (>150 cm). They were used to classify the soils into different depth classes and a soil depth map was generated. The area extent and their geographical distribution in the microwatershed is given in Fig. 5.2.




Fig. 5.2 Soil Depth map of Bhimanahalli-2 Microwatershed

Shallow (25-50 cm) and very shallow (<25 cm) soils cover an area of 164 ha (29%) and are distributed in the major part of the microwatershed. Moderately shallow (50-75 cm) soils cover an area of 164 ha (29%) and are distributed in the central, southern, southeastern, western, northern and northeastern part of the microwatershed. Deep (100-150 cm) soils cover an area of 76 ha (13%) and are distributed in the eastern, southern, southeastern and northwestern part of the microwatershed. Very deep (>150 cm) soils cover an area of 38 ha (7%) and are distributed in the western and northeastern part of the microwatershed.

The most productive lands 114 ha (20%) with respect to soil rooting depth where all climatically adapted annual and perennial crops can be grown are deep to very deep (100 - > 150 cm) soils. Problem soil covering an area of 164 ha (29%) are shallow and very shallow soils, where only short duration crops can be grown occasionally and the probability of crop failure is very high.

5.3 Surface Soil Texture

Texture is an expression to indicate the coarseness or fineness of the soil as determined by the relative proportion of primary particles of sand, silt and clay. It has a direct bearing on the structure, porosity, adhesion and consistence. The surface layer of a soil to a depth of about 25 cm is the layer that is most used by crops and plants. The surface soil textural class provides a guide to understanding soil-water retention and availability, nutrient holding capacity, infiltration, workability, drainage, physical and

chemical behaviour, microbial activity and crop suitability. The textural classes used for LRI were used to classify and a surface soil texture map was generated. The area extent and their geographical distribution in the microwatershed is shown in Figure 5.3.




Fig. 5.3 Surface Soil Texture map of Bhimanahalli-2 Microwatershed

An area of 39 ha (7%) has soils that are loamy at the surface and occur in the southern and southeastern part of the microwatershed. An area of 393 ha (70%) has soils that are clayey at the surface and occur in the major part of the microwatershed.

Major area of 432 ha (77%) the microwatershed is most productive with respect to surface soil texture. The clayey soils (70%) have high potential for soil-water retention and availability, and nutrient retention and availability, but have more problems of drainage, infiltration, workability and other physical problems. The other productive lands are loamy soils (7%) which also have high potential for soil-water retention and nutrient availability but have no drainage or other physical problems.

5.4 Soil Gravelliness

Gravel is the term used for describing coarse fragments between 2 mm and 7.5 cm diameter and stones for those between 7.5 cm and 25 cm. The presence of gravel and stones in soil reduces the volume of soil responsible for moisture and nutrient storage, drainage, infiltration and runoff, and hinders plant growth by impeding root growth and seedling emergence, intercultural operations and farm mechanization. The gravelliness classes used in LRI were used to classify the soils and using these classes a gravelliness

map was generated. The area extent and their geographic distribution in the microwatershed is shown in Figure 5.4.




Fig. 5.4 Soil Gravelliness map of Bhimanahalli-2 Microwatershed

Entire cultivated area of about 442 ha (77%) is non gravelly (<15%), in the microwatershed.

The most productive soils (77%) that are non gravelly (<15%), where all climatically adapted long duration crops can be grown.

5.5 Available Water Capacity

The soil available water capacity (AWC) is estimated based on the ability of the soil column to retain water between the tensions of 0.33 and 15 bar in a depth of 100 cm or the entire solum if the soil is shallower. The AWC of the soils (soil series) as estimated by considering the soil texture, mineralogy, soil depth and gravel content (Sehgal *et al.*, 1990) and accordingly the soil map units were grouped into five AWC classes *viz*, very low (<50 mm/m), low (50-100 mm/m), medium (100-150 mm/m), high (150-200 mm/m) and very high (>200 mm/m) and using these values, an AWC map was generated. The area extent and their geographic distribution of different AWC classes in the microwatershed is given in Figure 5.5.




Fig. 5.5 Soil Available Water Capacity map of Bhimanahalli-2 Microwatershed

An area of about 164 ha (29%) and 164 ha (29%) each are low (51-100 mm/m) and very low (<50 mm) in available water capacity in the microwatershed and are distributed in the major part of the microwatershed and about 114 ha (20%) is very high (>200 mm/m) in available water capacity and are distributed in the all parts of the microwatershed.

An area of 328 ha (57%) in the microwatershed is problematic with regard to available water capacity. Here, only short duration crops can be grown and the probability of crop failure is very high. These areas are best put to other alternative uses. An area of 114 ha (20%) is potential, where all climatically adapted long duration crops can be grown.

5.6 Soil Slope

Soil slope refers to the inclination of the surface of the land. It is defined by gradient, shape and length, and is an integral feature of any soil as a natural body. Slope is considered important in soil genesis, land use and land development. The length and gradient of slope influences the rate of runoff, infiltration, erosion and deposition. The soil map units were grouped into two slope classes and a slope map was generated showing the area extent and their geographic distribution in the microwatershed (Fig. 5.6).

Maximum area of about 440 ha (77%) falls under very gently sloping (1-3% slope) lands and are distributed in the major part of the microwatershed. An area of about

1 ha (<1%) falls under nearly level (0-1% slope) lands and are distributed in the northeastern part of the microwatershed.




Fig. 5.6 Soil Slope map of Bhimanahalli-2 Microwatershed

Entire cultivated area in the microwatershed is highly potential in respect of soil slopes. In these areas, all climatically adapted annual and perennial crops can be grown without much soil and water conservation and other land development measures.

5.7 Soil Erosion

Soil erosion refers to the wearing away of the earth's surface by the forces of water, wind and ice involving detachment and transport of soil by raindrop impact. It is used for accelerated soil erosion resulting from disturbance of the natural landscape by burning, excessive grazing and indiscriminate felling of forest trees and tillage, all usually by man. The erosion classes showing an estimate of the current erosion status as judged from field observations in the form of rills, gullies or a carpet of gravel on the surface are recorded. Four erosion classes, viz, slight erosion (e1), moderate erosion (e2), severe erosion (e3) and very severe erosion (e4) are recognized. The soil map units were grouped into different erosion classes and a soil erosion map generated. The area extent and their spatial distribution in the microwatershed is given in Figure 5.7.

Soils that are slightly eroded (e1 class) cover an area of 45 ha (8%) and are distributed in the eastern and northeastern part of the microwatershed. Soils that are moderately eroded (e2 class) cover a maximum area of 396 ha (69%) and are distributed

in the major part of the microwatershed. Moderately eroded soils are problematic and require proper soil and water conservation.




Fig. 5.7 Soil Erosion map of Bhimanahalli-2 Microwatershed

FERTILITY STATUS

Soil fertility plays an important role in increasing crop yield. The adoption of high yielding varieties that require high amounts of nutrients has resulted in deficiency symptoms in crops and plants due to imbalanced fertilization and poor inherent fertility status as these areas are characterised by low rainfall and high temperatures. Hence, it is necessary to know the fertility (macro and micro nutrients) status of the soils of the watersheds for assessing the kind and amount of fertilizers required for each of the crop intended to be grown. For this purpose, the surface soil samples collected from the grid points (one soil sample at every 320 m interval) all over the microwatershed through land resource inventory in the year 2018 were analysed for pH, EC, organic carbon, available phosphorus and potassium, and for micronutrients like zinc, boron, copper, iron and manganese, and secondary nutrient sulphur.

Soil fertility data generated has been assessed and individual maps for all the nutrients for the microwatershed have been generated using Kriging method under GIS. The village/survey number wise fertility data for the microwatershed is given in Appendix-II.

6.1 Soil Reaction (pH)

The soil analysis of the Bhimanahalli-2 microwatershed for soil reaction (pH) showed that an area of about 220 ha (38%) is neutral (pH 6.5-7.3) and are distributed in the major part of the microwatershed. About 174 ha (30%) is slightly alkaline (pH 7.3-7.8) and are distributed in the all parts of the microwatershed and about 47 ha (8%) is moderately alkaline (pH 7.8-8.4) and are distributed in the southern, western and eastern part of the microwatershed (fig.6.1).

6.2 Electrical Conductivity (EC)

The Electrical Conductivity of the soils in the microwatershed area is $<2 \text{ dS m}^{-1}$ (Fig 6.2) and as such the soils are non-saline.

6.3 Organic Carbon

The soil organic carbon content (an index of available Nitrogen) in the soils of the microwatershed is medium (0.5-0.75%) in about 10 ha (2%) and are distributed in the southeastern and western part of the microwatershed and 432 ha (75%) is high ($>0.75\%$) in organic carbon and are distributed in the major part of the microwatershed (Fig. 6.3).




Fig.6.1 Soil Reaction (pH) map of Bhimanahalli-2 Microwatershed




Fig.6.2 Electrical Conductivity (EC) map of Bhimanahalli-2 Microwatershed




Fig.6.3 Soil Organic Carbon map of Bhimanahalli-2 Microwatershed

6.4 Available Phosphorus

Available phosphorus content is high (>57 kg/ha) covering an area of about 363 ha (63%) and occur in the major part of the microwatershed. Medium (23-57 kg/ha) in an area of about 79 ha (14%) and occur in the western, eastern and northeastern part of the microwatershed (Fig. 6.4).

6.5 Available Potassium

Available potassium content is high (>337 kg/ha) in an area of 439 ha (77%) and occur in the major part of the microwatershed and medium (145-337 kg/ha) in an area of 3 ha (<1%) and occur in the southern part of the microwatershed (Fig.6.5).

6.6 Available Sulphur

Available sulphur content is low (<10 ppm) in the entire cultivated area of the microwatershed (Fig. 6.6).

6.7 Available Boron

Available boron content is low (<0.5 ppm) in the entire cultivated area of the microwatershed (Fig. 6.7).

6.8 Available Iron

Available iron content is sufficient (>4.5 ppm) in an area of 436 ha (76%) and occur in the major part of the microwatershed and deficient (<4.5 ppm) in an area of 5 ha (<1%) and occur in the northeastern part of the microwatershed (Fig 6.8).




Fig.6.4 Soil Available Phosphorus map of Bhimanahalli-2 Microwatershed




Fig.6.5 Soil Available Potassium map of Bhimanahalli-2 Microwatershed




Fig.6.6 Soil Available Sulphur map of Bhimanahalli-2 Microwatershed




Fig.6.7 Soil Available Boron map of Bhimanahalli-2 Microwatershed




Fig.6.8 Soil Available Iron map of Bhimanahalli-2 Microwatershed

6.9 Available Manganese

Available manganese content is sufficient (>1.0 ppm) in the entire cultivated area of the microwatershed (Fig 6.9).

6.10 Available Copper

Available copper content is sufficient (>0.2 ppm) in the entire cultivated area of the microwatershed area (Fig 6.10).

6.11 Available Zinc

Available zinc content is deficient (<0.6 ppm) in the entire cultivated area of the microwatershed (Fig 6.11).




Fig.6.9 Soil Available Manganese map of Bhimanahalli-2 Microwatershed




Fig.6.10 Soil Available Copper map of Bhimanahalli-2 Microwatershed




Fig.6.11 Soil Available Zinc map of Bhimanahalli-2 Microwatershed

LAND SUITABILITY FOR MAJOR CROPS

The soil and land resource units (soil phases) of Bhimanahalli-2 microwatershed were assessed for their suitability for growing food, fodder, fibre and other horticulture crops by following the procedure as outlined in FAO, 1976 and 1983. Crop requirements were developed for each of the crop from the available research data and also by referring to Naidu *et. al.* (2006) and Natarajan *et. al* (2015). The soil and land characteristics were matched with the crop requirement to arrive at the crop suitability. The soil and land characteristics (Table 7.1) and crop requirement (Table 7.2 to 7.30) are given at the end of the chapter. In FAO land suitability classification, two orders are recognized. Order S-Suitable and Order N-Not suitable. The orders have classes, subclasses and units. Order S has three classes, Class S1-Highly Suitable, Class S2-Moderately Suitable and Class S3- Marginally Suitable. Order N has two classes, N1-Currently not Suitable and N2- Permanently not Suitable. There are no subclasses within the Class S1 as they will have very minor or no limitations for crop growth. Classes S2, S3 and N1 are divided into subclasses based on the kinds of limitations encountered. The limitations that affect crop production are ‘c’ for erratic rainfall and its distribution and length of growing period (LGP), ‘e’ for erosion hazard, ‘r’ for rooting condition, ‘t’ for lighter or heavy texture, ‘g’ for gravelliness or stoniness, ‘n’ for nutrient availability, ‘l’ for topography, ‘m’ for moisture availability, ‘w’ for drainage and ‘z’ for calcareousness. These limitations are indicated as lower case letters to the Class symbol. For example, moderately suitable lands with the limitations of soil depth and erosion are designated as S2re. For the microwatershed, the soil mapping units were evaluated and classified up to subclass level.

Using the above criteria, the soil map units of the microwatershed were evaluated and land suitability maps for 29 major annual and perennial crops were generated. The detailed information on the kind of suitability of each of the soil phase for the crops assessed are given village/ survey number wise for the microwatershed in Appendix-IV.

7.1 Land Suitability for Sorghum (*Sorghum bicolor*)

Sorghum is one of the major food crop grown in Karnataka in an area of 10.47 lakh ha in Bijapur, Gulbarga, Raichur, Bidar, Belgaum, Dharwad, Bellary, Chitradurga, Mysore and Tumakuru districts. The crop requirements for growing sorghum (Table 7.2) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing sorghum was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.1.

An area of about 278 ha (49%) is moderately suitable (Class S2) for growing sorghum and are distributed in the major part of the microwatershed. They have minor limitations of rooting depth, texture, nutrient availability and calcareousness. About 99 ha

(17%) is marginally suitable (Class S3) for growing sorghum and are distributed in the central, southern and northwestern part of the microwatershed with moderate limitations of texture, nutrient availability, gravelliness and rooting depth. About 68 ha (12%) is currently not suitable (Class N1) for growing sorghum and is distributed in the northern and northwestern part of the microwatershed with severe limitation of rooting depth.




Fig. 7.1 Land Suitability map of Sorghum

7.2 Land Suitability for Maize (*Zea mays*)

Maize is one of the most important food crop grown in an area of 13.37 lakh ha in almost all the districts of the State. The crop requirements for growing maize (Table 7.3) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing maize was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.2.

An area of about 278 ha (49%) is moderately suitable (Class S2) for growing maize and are distributed in the major part of the microwatershed. They have minor limitations of rooting depth, texture, nutrient availability and calcareousness. About 99 ha (17%) is marginally suitable (Class S3) for growing maize and are distributed in the central, southern and northwestern part of the microwatershed with moderate limitations of texture, nutrient availability, gravelliness and rooting depth. About 68 ha (12%) is currently not suitable (Class N1) for growing maize and is distributed in the northern and northwestern part of the microwatershed with severe limitation of rooting depth.




Fig. 7.2 Land Suitability map of Maize

7.3 Land Suitability for Bajra (*Pennisetum glaucum*)

Bajra is one of the most important millet crop grown in an area of 2.34 lakh ha in the northern districts of Karnataka state. The crop requirements for growing bajra (Table 7.4) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing bajra was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.3.

An area of about 275 ha (48%) is moderately suitable (Class S2) for growing bajra and are distributed in the major part of the microwatershed. They have minor limitations of nutrient availability, rooting depth, texture and calcareousness. About 99 ha (17%) is marginally suitable (Class S3) for growing bajra and are distributed in the central, southern and northwestern part of the microwatershed with moderate limitations of texture, nutrient availability and rooting depth. About 68 ha (12%) is currently not suitable (Class N1) for growing bajra and is distributed in the northern and northwestern part of the microwatershed with severe limitation of rooting depth.




Fig. 7.3 Land Suitability map of Bajra

7.4 Land Suitability for Groundnut (*Arachis hypogaea*)

Groundnut is one of the major oilseed crop grown in an area of 6.54 lakh ha in Karnataka in most of the districts either as rainfed or irrigated crop. The crop requirements for growing groundnut (Table 7.5) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing groundnut was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.4.

An area of about 163 ha (29%) is moderately suitable (Class S2) for growing groundnut and are distributed in the central, southern, western, northeastern and southeastern part of the microwatershed. They have minor limitations of rooting depth and texture. An area of about 207 ha (36%) is marginally suitable (Class S3) for growing groundnut and are distributed in the major part of the microwatershed. They have moderate limitations of calcareousness, rooting depth, nutrient availability and texture. About 71 ha (12%) is currently not suitable (Class N1) for growing groundnut and is distributed in the northwestern and southern part of the microwatershed with severe limitation of rooting depth.




Fig. 7.4 Land Suitability map of Groundnut

7.5 Land Suitability for Sunflower (*Helianthus annus*)

Sunflower is one of the most important oilseed crop grown in an area of 3.56 lakh ha in the State in all the districts. The crop requirements for growing sunflower (Table 7.6) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing sunflower was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.5.

An area of about 57 ha (10%) is moderately suitable (Class S2) for sunflower and are distributed in the western, northwestern and northeastern part of the microwatershed. They have minor limitation of calcareousness. An area of about 218 ha (38%) is marginally suitable (Class S3) for growing sunflower and are distributed in the major part of the microwatershed. They have moderate limitations of nutrient availability and rooting depth. About 167 ha (29%) is currently not suitable (Class N1) for growing sunflower and is distributed in the central, southern, northern and northwestern part of the microwatershed with severe limitations of rooting depth and nutrient availability.




Fig. 7.5 Land Suitability map of Sunflower

7.6 Land Suitability for Red gram (*Cajanus Cajan*)

Redgram is one of the most important pulse crop grown in an area of 7.28 lakh ha in almost all the districts of the State. The crop requirements for growing red gram (Table 7.7) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing redgram was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.6.

An area of about 111 ha (19%) is moderately suitable (Class S2) for growing redgram and are distributed in the eastern, southern, western and northwestern part of the microwatershed. They have minor limitations of nutrient availability, texture and calcareousness. An area of about 167 ha (29%) is marginally suitable (Class S3) for growing redgram and are distributed in the major part of the microwatershed. They have moderate limitations of rooting depth and nutrient availability. About 164 ha (29%) is currently not suitable (Class N1) for growing redgram and is distributed in the central, southern, northern and northwestern part of the microwatershed with severe limitation of rooting depth.




Fig. 7.6 Land Suitability map of Redgram

7.7 Land Suitability for Bengal gram (*Cicer aerativum*)

Bengal gram is one of the most important pulse crop grown in about 9.39 lakh ha area in Bijapur, Raichur, Kalaburgi, Dharwad, Belgaum and Bellary districts. The crop requirements for growing Bengal gram (Table 7.8) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing Bengal gram was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.7.

An area of about 57 ha (10%) is moderately suitable (Class S2) for bengalgram and are distributed in the western, northwestern and northeastern part of the microwatershed. They have minor limitation of calcareousness. An area of about 278 ha (49%) is marginally suitable (Class S3) for growing bengalgram and are distributed in the major part of the microwatershed. They have moderate limitations of rooting depth, nutrient availability and texture. About 106 ha (19%) is currently not suitable (Class N1) for growing bengalgram and is distributed in the southern, northwestern and northern part of the microwatershed with severe limitations of rooting depth and texture.




Fig. 7.7 Land Suitability map of Bengal gram.

7.8 Land Suitability for Cotton (*Gossypium hirsutum*)

Cotton is one of the most important fibre crop grown in the State in about 8.75 lakh ha area in Raichur, Dharwad, Belgaum, Kalaburgi, Bijapur, Bidar, Bellary, Chitradurga and Chamarajnagar districts. The crop requirements for growing cotton (Table 7.9) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing cotton was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.8.

An area of about 199 ha (35%) is moderately suitable (Class S2) for cotton and are distributed in the major part of the microwatershed. They have minor limitations of rooting depth, texture and calcareousness. Marginally suitable lands (Class S3) for growing cotton occupy an area of about 136 ha (24%) and occur in the eastern, central, northeastern, southern and northwestern part of the microwatershed. They have moderate limitations of rooting depth, texture, gravelliness and nutrient availability. About 106 ha (19%) is currently not suitable (Class N1) and is distributed in the southern, northwestern and northern part of the microwatershed with severe limitations of rooting depth and texture.




Fig. 7.8 Land Suitability map of Cotton

7.9 Land Suitability for Chilli (*Capsicum annuum*)

Chilli is one of the most important spice crop grown in about 0.42 lakh ha in Karnataka State. The crop requirements for growing chilli (Table 7.10) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing chilli was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.9.

An area of about 221 ha (38%) is moderately suitable (Class S2) for growing chilli and are distributed in the major part of the microwatershed. They have minor limitations of rooting depth, texture and calcareousness. About 150 ha (26%) is marginally suitable (Class S3) for growing chilli and are distributed in the central, southern, eastern and northwestern part of the microwatershed with moderate limitations of rooting depth, nutrient availability and gravelliness. About 71 ha (12%) is currently not suitable (Class N1) for growing chilli and is distributed in the northern and northwestern part of the microwatershed with severe limitation of rooting depth.




Fig 7.9 Land Suitability map of Chilli

7.10 Land Suitability for Tomato (*Lycopersicon esculentum*)

Tomato is one of the most important vegetable crop grown in about 0.61 lakh ha covering almost all the district of the state. The crop requirements for growing tomato (Table 7.11) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing tomato was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.10.

An area of about 164 ha (29%) is moderately suitable (Class S2) for growing tomato and are distributed in the central, southern, western, northeastern and southeastern part of the microwatershed. They have minor limitations of rooting depth and texture. An area of about 207 ha (36%) is marginally suitable (Class S3) for growing tomato and are distributed in the major part of the microwatershed. They have moderate limitations of gravelliness, rooting depth, nutrient availability and texture. About 71 ha (12%) is currently not suitable (Class N1) for growing tomato and is distributed in the northwestern and southern part of the microwatershed with severe limitations of rooting depth and nutrient availability.




Fig 7.10 Land Suitability map of Tomato

7.11 Land Suitability for Brinjal (*Solanum melongena*)

Brinjal is one of the most important vegetable crop grown in the state. The crop requirements for growing brinjal (Table 7.12) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing Brinjal was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.11.

An area of about 164 ha (29%) is moderately suitable (Class S2) for growing brinjal and are distributed in the central, southern, western, northeastern and southeastern part of the microwatershed. They have minor limitation of rooting depth. An area of about 207 ha (36%) is marginally suitable (Class S3) for growing brinjal and are distributed in the major part of the microwatershed. They have moderate limitations of gravelliness, rooting depth, nutrient availability and texture. About 71 ha (12%) is currently not suitable (Class N1) for growing brinjal and is distributed in the northwestern, northern and southern part of the microwatershed with severe limitations of rooting depth and nutrient availability.




Fig 7.11 Land Suitability map of Brinjal

7.12 Land Suitability for Onion (*Allium cepa L.*)

Onion is one of the most important vegetable crop grown in the state. The crop requirements for growing onion (Table 7.13) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing onion was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.12.

An area of about 164 ha (29%) is moderately suitable (Class S2) for growing onion and are distributed in the central, southern, western, northeastern and southeastern part of the microwatershed. They have minor limitation of rooting depth. An area of about 207 ha (36%) is marginally suitable (Class S3) for growing onion and are distributed in the major part of the microwatershed. They have moderate limitations of gravelliness, rooting depth, nutrient availability and texture. About 71 ha (12%) is currently not suitable (Class N1) for growing onion and is distributed in the northwestern, northern and southern part of the microwatershed with severe limitations of rooting depth and nutrient availability.




Fig 7.12 Land Suitability map of Onion

7.13 Land Suitability for Bhendi (*Abelmoschus esculentus*)

Bhendi is one of the most important vegetable crop grown in the state. The crop requirements for growing bhendi (Table 7.14) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing bhendi was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.13.

An area of about 221 ha (38%) is moderately suitable (Class S2) for growing bhendi and are distributed in the major part of the microwatershed. They have minor limitations of rooting depth, texture and calcareousness. About 150 ha (26%) is marginally suitable (Class S3) for growing bhendi and are distributed in the central, southern, eastern and northwestern part of the microwatershed with moderate limitations of rooting depth, nutrient availability and gravelliness. About 71 ha (12%) is currently not suitable (Class N1) for growing bhendi and is distributed in the northern and northwestern part of the microwatershed with severe limitations of rooting depth and nutrient availability.




Fig 7.13 Land Suitability map of Bhenidi

7.14 Land Suitability for Drumstick (*Moringa oleifera*)

Drumstick is one of the most important vegetable crop grown in about 2403 ha in the state. The crop requirements for growing drumstick (Table 7.15) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing drumstick was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.14.

An area of about 221 ha (38%) is marginally suitable (Class S3) for growing drumstick and are distributed in the all parts of the microwatershed. They have moderate limitations of rooting depth and calcareousness. About 221 ha (39%) is currently not suitable (Class N1) for growing drumstick and is distributed in the major part of the microwatershed with severe limitations of rooting depth and nutrient availability.




Fig 7.14 Land Suitability map of Drumstick

7.15 Land Suitability for Mango (*Mangifera indica*)

Mango is one of the most important fruit crop grown in an area of 1.73 lakh ha in almost all the districts of the State. The crop requirements (Table 7.16) for growing mango were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing mango was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed is given in Figure 7.15.

An area of about 111 ha (19%) is marginally suitable (Class S3) for mango and are distributed in the western, northwestern, southern, southeastern and northeastern part of the microwatershed. They have moderate limitations of texture and nutrient availability. Currently not suitable (Class N1) lands for growing mango occupy an area about 331 ha (59%) and occur in the major part of the microwatershed. They have severe limitations of rooting depth and nutrient availability.




Fig. 7.15 Land Suitability map of Mango

7.16 Land Suitability for Guava (*Psidium guajava*)

Guava is one of the most important fruit crop grown in an area of 0.06 lakh ha in almost all the districts of the State. The crop requirements (Table 7.17) for growing guava were matched with the soil-site characteristics (7.1) and a land suitability map for growing guava was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.16.

An area of about 221 ha (38%) is marginally suitable (Class S3) for growing guava and are distributed in the all parts of the microwatershed. They have moderate limitations of rooting depth, texture and calcareousness. About 221 ha (39%) is currently not suitable (Class N1) for growing guava and is distributed in the major part of the microwatershed with severe limitations of rooting depth and nutrient availability.




Fig. 7.16 Land Suitability map of Guava

7.17 Land Suitability for Sapota (*Manilkara zapota*)

Sapota is one of the most important fruit crop grown in an area of 29373 ha in almost all the districts of the State. The crop requirements (Table 7.18) for growing sapota were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing sapota was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.17.

An area of about 277 ha (48%) is marginally suitable (Class S3) for growing sapota and are distributed in the major part of the microwatershed. They have moderate limitations of rooting depth, nutrient availability and texture. About 167 ha (29%) is currently not suitable (Class N1) for growing sapota and is distributed in the central, southern, northern and northwestern part of the microwatershed with severe limitations of rooting depth and nutrient availability.




Fig. 7.17 Land Suitability map of Sapota

7.18 Land Suitability for Pomegranate (*Punica granatum*)

Pomegranate is one of the most important fruit crop commercially grown in about 18488 ha in Karnataka, mainly in Bijapur, Bagalkot, Koppal, Gadag and Chitradurga districts. The crop requirements for growing pomegranate (Table 7.19) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing pomegranate was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.18.

An area of about 57 ha (10%) is moderately suitable (Class S2) for pomegranate and are distributed in the western, northwestern and northeastern part of the microwatershed. They have minor limitations of texture and calcareousness. An area of about 218 ha (38%) is marginally suitable (Class S3) for growing pomegranate and are distributed in the major part of the microwatershed. They have moderate limitations of nutrient availability and rooting depth. About 167 ha (29%) is currently not suitable (Class N1) for growing pomegranate and is distributed in the central, southern, northern and northwestern part of the microwatershed with severe limitations of rooting depth and nutrient availability.




Fig 7.18 Land Suitability map of Pomegranate

7.19 Land Suitability for Musambi (*Citrus limetta*)

Musambi is one of the important fruit crop grown in an area of 3446 ha in almost all the districts of the State. The crop requirements for growing musambi (Table 7.20) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing musambi was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.19.

An area of about 57 ha (10%) is moderately suitable (Class S2) for musambi and are distributed in the western, northwestern and northeastern part of the microwatershed. They have minor limitation of calcareousness. An area of about 218 ha (38%) is marginally suitable (Class S3) for growing musambi and are distributed in the major part of the microwatershed. They have moderate limitations of nutrient availability and rooting depth. About 167 ha (29%) is currently not suitable (Class N1) for growing musambi and is distributed in the central, southern, northern and northwestern part of the microwatershed with severe limitations of rooting depth and nutrient availability.




Fig. 7.19 Land Suitability map of Musambi

7.20 Land Suitability for Lime (*Citrus sp*)

Lime is one of the most important fruit crop grown in an area of 0.11 lakh ha in almost all the districts of the State. The crop requirements for growing lime (Table 7.21) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing lime was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7. 20.

An area of about 57 ha (10%) is moderately suitable (Class S2) for lime and are distributed in the western, northwestern and northeastern part of the microwatershed. They have minor limitation of calcareousness. An area of about 218 ha (38%) is marginally suitable (Class S3) for growing lime and are distributed in the major part of the microwatershed. They have moderate limitations of nutrient availability and rooting depth. About 167 ha (29%) is currently not suitable (Class N1) for growing lime and is distributed in the central, southern, northern and northwestern part of the microwatershed with severe limitations of rooting depth and nutrient availability.




Fig. 7.20 Land Suitability map of Lime

7.21 Land Suitability for Amla (*Phyllanthus emblica*)

Amla is one of the medicinal fruit crop grown in almost all the districts of the State. The crop requirements for growing amla (Table 7.22) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing amla was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.21.

An area of about 164 ha (29%) is moderately suitable (Class S2) for growing amla and are distributed in the major part of the microwatershed. They have minor limitation of rooting depth. An area of about 152 ha (27%) is marginally suitable (Class S3) for growing amla and are distributed in the central, southern, western, northwestern and northeastern part of the microwatershed. They have moderate limitations of calcareousness, rooting depth and texture. About 125 ha (22%) is currently not suitable (Class N1) for growing amla and is distributed in the southern, eastern, northeastern and southeastern part of the microwatershed with severe limitations of rooting depth and nutrient availability.




Fig. 7.21 Land Suitability map of Amla

7.22 Land Suitability for Cashew (*Anacardium occidentale*)

Cashew is one of the most important plantation nut crop grown in an area of 0.7 lakh ha in almost all the districts of the State. The crop requirements for growing cashew (Table 7.23) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing cashew was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.22.

Marginally (Class S3) suitable lands for growing cashew occur in an area of 142 ha (25%) and are distributed in the southern, central, western and northeastern part of the microwatershed. They have moderate limitation of rooting depth. Currently not suitable (Class N1) lands for growing cashew occur in a maximum area of 300 ha (52%) and are distributed in the major part of the microwatershed with severe limitations of texture, rooting depth and nutrient availability.




Fig. 7.22 Land Suitability map of Cashew

7. 23 Land Suitability for Jackfruit (*Artocarpus heterophyllus*)

Jackfruit is one of the most important fruit crop grown in an area of 5368 ha in almost all the districts of the State. The crop requirements for growing jackfruit (Table 7.24) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing jackfruit was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.23.

An area of about 221 ha (38%) is marginally suitable (Class S3) for growing jackfruit and are distributed in the all parts of the microwatershed. They have moderate limitations of rooting depth, texture and calcareousness. About 221 ha (39%) is currently not suitable (Class N1) for growing jackfruit and is distributed in the major part of the microwatershed with severe limitations of rooting depth and nutrient availability.




Fig. 7.23 Land Suitability map of Jackfruit

7.24 Land Suitability for Jamun (*Syzygium cumini*)

Jamun is an important fruit crop grown in almost all the districts of the State. The crop requirements for growing jamun (Table 25) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing jamun was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.24.

An area of about 221 ha (38%) is marginally suitable (Class S3) for growing jamun and are distributed in the all parts of the microwatershed. They have moderate limitations of rooting depth and calcareousness. About 221 ha (39%) is currently not suitable (Class N1) for growing jamun and is distributed in the major part of the microwatershed with severe limitations of rooting depth and nutrient availability.




Fig. 7.24 Land Suitability map of Jamun

7.25 Land Suitability for Custard Apple (*Annona reticulata*)

Custard apple is one of the most important fruit crop grown in almost all the districts of the State. The crop requirements for growing custard apple (Table 7.26) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing custard apple was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.25.

An area of about 221 ha (38%) is moderately suitable (Class S2) for growing custard apple and are distributed in the major part of the microwatershed. They have minor limitations of rooting depth and calcareousness. About 149 ha (26%) is marginally suitable (Class S3) for growing custard apple and are distributed in the eastern, central, southern, southeastern and northwestern part of the microwatershed with moderate limitations of nutrient availability, texture and rooting depth. About 71 ha (12%) is currently not suitable (Class N1) and is distributed in the northern and northwestern part of the microwatershed with severe limitations of rooting depth and nutrient availability.




Fig. 7.25 Land Suitability map of Custard Apple

7.26 Land Suitability for Tamarind (*Tamarindus indica*)

Tamarind is one of the most important spice crop grown in almost all the districts of the state. The crop requirements for growing tamarind (Table 7.27) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing tamarind was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Fig. 7.26.

An area of about 57 ha (10%) is marginally suitable (Class S3) for tamarind and are distributed in the western and northwestern part of the microwatershed. They have moderate limitation of calcareousness. Currently not suitable (Class N1) lands for growing tamarind occupy an area about 385 ha (67%) and occur in the major part of the microwatershed. They have severe limitations of nutrient availability and rooting depth.




Fig. 7.26 Land Suitability map of Tamarind

7.27 Land Suitability for Mulberry (*Morus nigra*)

Mulberry is the important leaf crop grown for rearing of silkworms in about 1.6 lakh ha area in all the districts of the state. The crop requirements for growing mulberry (Table 7.28) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing mulberry was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.27.

An area of about 221 ha (38%) is marginally suitable (Class S3) for growing mulberry and are distributed in the all parts of the microwatershed. They have moderate limitations of rooting depth, texture and calcareousness. About 221 ha (39%) is currently not suitable (Class N1) for growing jamun and is distributed in the major part of the microwatershed with severe limitations of rooting depth and nutrient availability.




Fig 7.27 Land Suitability map of Mulberry

7.28 Land Suitability for Marigold (*Tagetes sps.*)

Marigold is one of the most important flower crop grown in an area of 9108 ha in almost all the districts of the State. The crop requirements (Table 7.29) for growing marigold were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing marigold was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.28.

An area of about 221 ha (38%) is moderately suitable (Class S2) for growing marigold and are distributed in the major part of the microwatershed. They have minor limitations of rooting depth, texture and calcareousness. About 150 ha (26%) is marginally suitable (Class S3) for growing marigold and are distributed in the central, southern, eastern and northwestern part of the microwatershed with moderate limitations of rooting depth, nutrient availability and gravelliness. About 71 ha (12%) is currently not suitable (Class N1) for growing marigold and is distributed in the northern and northwestern part of the microwatershed with severe limitations of rooting depth and nutrient availability.




Fig. 7.28 Land Suitability map of Marigold

7.29 Land Suitability for Chrysanthemum (*Dendranthema grandiflora*)

Chrysanthemum is one of the most important flower crop grown in an area of 4978 ha in almost all the districts of the State. The crop requirements (Table 7.30) for growing chrysanthemum were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing chrysanthemum was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.29.

An area of about 221 ha (38%) is moderately suitable (Class S2) for growing chrysanthemum and are distributed in the major part of the microwatershed. They have minor limitations of rooting depth, texture and calcareousness. About 150 ha (26%) is marginally suitable (Class S3) for growing chrysanthemum and are distributed in the central, southern, eastern and northwestern part of the microwatershed with moderate limitations of rooting depth, nutrient availability and gravelliness. About 71 ha (12%) is currently not suitable (Class N1) for growing chrysanthemum and is distributed in the northern and northwestern part of the microwatershed with severe limitations of rooting depth and nutrient availability.




Fig. 7.29 Land Suitability map of Chrysanthemum

Table 7.1 Soil-Site Characteristics of Bhimanahalli-2 Microwatershed

Soil Map Units	Climate (P) (mm)	Growing period (Days)	Drainage Class	Soil depth (cm)	Soil texture		Gravelliness		AWC (mm/m)	Slope (%)	Erosion	pH	EC (dSm ⁻¹)	ESP (%)	CEC [Cmol (p ⁺)kg ⁻¹]	BS (%)
					Sur-face	Sub-surface	Surface (%)	Sub-surface (%)								
BDPiB2	866	150	WD	<25	sc	scl	<15	<15	<50	1-3	moderate	8.58	0.262	0.35	18.10	100
HTKcB2	866	150	WD	25-50	sl	sl	<15	10-25	<50	1-3	moderate	6.81	0.062	0.38	3.0	100
BDLiB2	866	150	WD	25-50	sc	sl	<15	<15	<50	1-3	moderate	6.20	0.074	0.20	4.20	93
DSBiB2	866	150	WD	25-50	sc	g c	<15	35-60	<50	1-3	moderate	5.93	0.04	0.14	3.60	73
VNKib2	866	150	WD	25-50	sc	sc	<15	<15	<50	1-3	moderate	5.37	0.11	2.22	6.27	75
JNKhB2	866	150	W	50-75	scl	scl	<15	<15	51-100	1-3	moderate	8.42	0.148	0.18	14.50	100
NGPmB2	866	150	MWD	100-150	c	c	<15	<15	>200	1-3	moderate	7.42	0.24	0.22	67.10	100
ANRiB2	866	150	MWD	100-150	sc	c	<15	<15	>200	1-3	moderate	10.17	0.365	7.08	19.90	100
MDGhB2	866	150	WD	100-150	scl	scl	<15	<15	>200	1-3	moderate	8.20	0.399	3.08	4.90	100
MDGmB1	866	150	WD	100-150	c	scl	<15	<15	>200	1-3	moderate	8.20	0.399	3.08	4.90	100
BMNmA1	866	150	MWD	>150	c	c	<15	<15	>200	0-1	slight	8.20	0.284	0.65	52.70	100
BMNmB2	866	150	MWD	>150	c	c	<15	<15	>200	1-3	moderate	8.20	0.284	0.65	52.70	100
BLDmB2	866	150	Mw	50-75	c	cl	<15	<15	101-150	1-3	moderate	8.19	0.22	0.80	38.20	90

*Symbols and abbreviations are according to Field Guide for LRI under Sujala-III Project, Karnataka

Table 7.2 Land suitability criteria for Sorghum

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	26–30	30–34; 24–26	34–40; 20–24	>40; <20
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, c (red), c (black)	scl, cl	ls, sl	-
	pH	1:2.5	5.5-7.8	5.0-5.5 7.8-9.0	>9.0	-
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	10-15
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

Table 7.3 Land suitability criteria for Maize

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	30-34	35-38 26-30	38-40 26-20	
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	Mm				
	Rainfall in growing season	Mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc	c (red), c (black)	ls, sl	-
	pH	1:2.5	5.5-7.8	5.0-5.5 7.8-9.0	>9.0	-
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	Cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	-
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

Table 7.4 Land suitability criteria for Bajra

Land use requirement		Rating				
Soil –site characteristics	Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)	
Climatic regime	Mean temperature in growing season	°C	28-32	33-38 24-27	39-40 20-23	<20
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm	500-750	400-500	200-400	<200
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl, scl, cl,sc,c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.8	5.0-5.5 7.8-9.0	5.5-6.0 >9.0	
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	15-35	35-60	>60	
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	1-3	3-5	5-10	>10

Table 7.5 Land suitability criteria for Groundnut

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	24–33	22–24; 33–35	20–22; 35–40	<20; >40
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	Mm				
	Rainfall in growing season	Mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. Well drained	Poorly drained	Very Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl	sl,cl, sc	c (red), c (black), ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	Cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<35	35-60	>60	
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.6 Land suitability criteria for Sunflower

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	24–30	30–34; 20–24	34–38; 16–20	>38; <16
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	mod. Well drained	-	Poorly to very drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	cl, sc,c (red), c (black)	scl	ls, sl	-
	pH	1:2.5	6.5-7.8	7.8-8.4 5.5-6.5	8.4-9.0; 5.0-5.5	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.7 Land suitability criteria for Redgram

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	30-35(G) 20-25(AV) 15-18 (F&PS) 35-40(M)	25-30(G) 20-25 (AV) 12-15 (F&PS) 30-35(M)	20-25(G) 15-20(AV) 10-12 (F&PS) 25-30(M)	< 20 <15 <10 <25
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	Mm				
	Rainfall in growing season	Mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. Well drained	Poorly drained	Very Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, c (red)	c (black),sl, scl, cl	ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-9.0	5.0-5.5 >9.0	-
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	Cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-50	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<1.0	1.0-2.0	>2.0	
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.8 Land suitability criteria for Bengal gram

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	20–25	25–30; 15–20	30–35; 10–15	>35; <10
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. Well drained	Poorly drained	Very Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	c(black)	-	c (red), scl, cl, sc	ls, sl
	pH	1:2.5	6.0-7.8	5.0-6.0 7.8-9.0	>9.0	-
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	-
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.9 Land suitability criteria for Cotton

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	22-32	>32	<19	-
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well to moderately well	Poorly drained/Some what excessively drained	-	very poorly/excessively drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, c (red,black)	cl	scl	ls, sl
	pH	1:2.5	6.5-7.8	7.8-8.4	5.5-6.5 8.4->9.0	<5.5
	CEC	C mol (p+)Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	50-100	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	<3	3-5	-	>5

Table 7.10 Land suitability criteria for Chilli

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	25-32	33-35 20-25	35-38 <20	>38
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc	c (black), sl	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.11 Land suitability criteria for Tomato

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	25-28	29-32 20-24	15-19 33-36	<15 >36
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl, scl, cl, sc, c (red)	-	ls, c(black)	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.12 Land suitability criteria for Brinjal

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	Well drained	Moderately well drained	Poorly drained	V. Poorly drained
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class				
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl, scl, cl, sc c (red)	-	ls, c (black)	-
	pH	1:2.5	6.0-7.3	7.3-8.4 5.0-6.0	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	>60
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.13 Land suitability criteria for Onion

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	20-30	30-35	35-40	>40
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately /imperfectly	-	Poorly to V poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl,scl,cl,sc,c (red)	-	c (Black),ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<1.0	1.0-2.0	2.0-4.0	<4
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.14 Land suitability criteria for Bhendi

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	25-28	29-32 20-24	15-19 33-36	<15 >36
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Imperfectly drained	Poorly to very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl,sc, c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.15 Land suitability criteria for Drumstick

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, scl, cl, c (red)	sl, c (black)	ls	S
	pH	1:2.5	6.0-7.3	5.0-5.5 7.3-7.8	5.5-6.0 7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<35	35-60	60-80	>80
Soil toxicity	Salinity (EC saturation extract)	ds/m				
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-10	-	>10

Table 7.16 Land suitability criteria for Mango

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-32	24-27 33-35	36-40	20-24
	Min temp. before flowering	°C	10-15	15-22	>22	-
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration	Days				
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V. Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	-	ls, sl, c (black)	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>150	100-150	75-100	<75
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.17 Land suitability criteria for Guava

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-32	33-36 24-27	37-42 20-23	
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	sl	c (black), ls	-
	pH	1:2.5	6.0-7.8	5.0-6.0	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.18 Land suitability criteria for Sapota

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-32	33-36 24-27	37-42 20-23	>42 <18
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	-	Poorly to very drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	sl	ls, c (black)	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.19 Land suitability criteria for Pomegranate

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	30-34	35-38 25-29	39-40 15-24	
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl,cl, sc, c (red)	c (black),sl	ls	-
	pH	1:2.5	5.5-7.8	7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.20 Land suitability criteria for Musambi

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-30	31-35 24-27	36-40 20-23	>40 <20
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately drained	poorly	Very poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c	sl	ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.21 Land suitability criteria for Lime

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-30	31-35 24-27	36-40 20-23	>40 <20
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately drained	poorly	Very poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c	sl	ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.22 Land suitability criteria for Amla

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. well drained	Poorly drained	V. Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	c (black)	ls, sl	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15-35	35-60	60-80	-
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

Table 7.23 Land suitability criteria for Cashew

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	32 to 34	28 to 32; 34 to 38	24 to 28; 38 to 40	<20; >40
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	-	sl, ls	c (black)
	pH	1:2.5	5.5-6.5	5.0-5.5 6.5-7.3	7.3-7.8	>7.8
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-10	>10	-

Table 7.24 Land suitability criteria for Jackfruit

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. well	Poorly	V. Poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	-	sl, ls, c (black)	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	>60
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10-

Table 7.25 Land suitability criteria for Jamun

Land use requirement		Rating				
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well	Mod. well	Poorly	V.Poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c(red)	sl, c (black)	ls	-
	pH	1:2.5	6.0-7.8	5.0-6.0	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>150	100-150	50-100	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	>60
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

Table 7.26 Land suitability criteria for Custard apple

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	Scl, cl, sc, c (red), c (black)	-	Sl, ls	-
	pH	1:2.5	6.0-7.3	5.5-6.0 7.3-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15-35	35-60	60-80	-
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	>5	-

Table 7.27 Land suitability criteria for Tamarind

Land use requirement		Rating				
Soil –site characteristics	Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)	
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod.well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl,sc, c (red)	sl, c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>150	100-150	75-100	<75
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

Table 7.28 Land suitability criteria for Mulberry

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	24–28	22–24; 28–32	32–38; 22–18	>38; <18
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V. Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, cl, scl	c (red)	c (black), sl, ls	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.8-8.4	7.3-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	0-35	35-60	60-80	>80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

Table 7.29 Land suitability criteria for Marigold

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	18-23	17-15 24-35	35-40 10-14	>40 <10
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl,scl, cl, sc, c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%				
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.30 Land suitability criteria for Chrysanthemum

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	18-23	17-15 24-35	35-40 10-14	>40 <10
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl,scl, cl, sc, c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%				
Erosion hazard	Slope	%	<3	3-5	5-10	>10

7.30 Land Management Units (LMUs)

The 13 soil map units identified in Bhimanahalli-2 microwatershed have been grouped into 8 Land Management Units (LMUs) for the purpose of preparing a Proposed Crop Plan. Land Management Units are grouped based on the similarities in respect of the type of soil, the depth of the soil, the surface soil texture, gravel content, AWC, slope, erosion etc. and a Land Management Units map (Fig. 7.30) has been generated. These Land Management Units are expected to behave similarly for a given level of management.




Fig. 7.30 Land Management Units Map Bhimanahalli-2 Microwatershed

The map units that have been grouped into 8 Land Management Units along with brief description of soil and site characteristics are given below.

LMU	Soil map units	Soil and site characteristics
1	159.BMNmA1 62.BMNmB2	Very deep, black calcareous clay soils (>150cm), 1-3 % slopes non-gravelly (<15%), slight to moderate erosion.
2	55.ANRiB2	Deep, sodic clay soils (100-150cm), 1-3 % slopes, non-gravelly (<15%), moderate erosion.
3	148.MDGhB2 170.MDGmB1	Deep, sandy clay loam and strongly alkaline soils (100-150 cm), 1- 3% slopes, non- gravelly (<15%), moderate erosion.
4	49.NGPmB2	Deep, black calcareous clay soils (100-150 cm), 1-3 % slopes, non-gravelly (<15%), moderate erosion.
5	76.BLDmB2	Moderately shallow, clay loam soils (50-75 cm), 1- 3% slopes, non- gravelly (<15 %), moderate erosion.
6	110.JNKhB2	Moderately shallow, sandy clay loam soils (50-75 cm), 1- 3% slopes, non- gravelly (<15%), moderate erosion.
7	5.BDLiB2 108.DSBiB2 165.HTKcB2 10.VNKiB2	Shallow sandy clay loam to sandy loam soils (25-50 cm), 1- 3% slopes, non- gravelly (<15 %), moderate erosion.
8	1.BDPiB2	Very shallow, sandy clay loam soils (<25 cm), 1- 3% slopes, non- gravelly (<15%), moderate erosion.

7.31 Proposed Crop Plan for Bhimanahalli-2 Microwatershed

After assessing the land suitability for the 29 crops, the Proposed Crop Plan has been prepared for the 8 identified LMUs by considering only the highly (Class S1) and moderately (Class S2) suitable lands for each of the 29 crops. The resultant proposed crop plan is presented below in Table 7.31.

Table 7.31 Proposed Crop Plan for Bhimanahalli-2 Microwatershed

LMU	Soil Map Units	Survey Number	Field Crops/ Commercial crops	Horticulture Crops (Rainfed/Irrigated)	Suitable Interventions
1	159.BMNmA1 62.BMNmB2	Bheemanahalli: 2(2),3,10,11, 26,27,28,29(1),29(2),29(3),29 (4),60,61,76,77,78,79,281,28 2,283,287(1) Handaraki : 421,422	Maize, Sorghum, Sunflower, Cotton, Red gram, Bengal gram, Bajra	Fruit crops: Lime, Musambi, Custard apple, Pomegranate Vegetables: Chilli, Bhendi Flowers: Marigold, Chrysanthemum	Application of FYM, Bio- fertilizers and micronutrients, drip irrigation, mulching, suitable soil and water conservation practices
2	55.ANRiB2	Bheemanahalli : 70	-	Agri-Silvi-Pasture Ber, Aonla, Acacia sp. Dhaincha, Rhodes grass, Para grass ,Bermuda grass	Application of gypsum, iron pyrites and elemental sulphur. Addition of farm yard manure, green manures and providing subsurface drainage
3	148.MDGhB2 170.MDGmB1	Bheemanahalli: 71,72(1),72(2),75,92 Handaraki: 253,254,255,256, 257, 258,259,260,261,343, 344,345,346	Sorghum, Maize, Bajra	Agri-Silvi-Pasture Ber, Aonla, Acacia sp. Dhaincha, Rhodes grass, Para grass ,Bermuda grass	Application of FYM, Bio- fertilizers and micronutrients, drip irrigation, mulching, suitable soil and water conservation practices
4	49.NGPmB2	Ramathirtha: 128,134,180,18 4,185	Maize, Sorghum, Sunflower, Cotton, Red gram, Bengalgram, Bajra	Fruit crops: Lime, Musambi, Custard apple, Pomegranate Vegetables: Chilli, Bhendi Flowers: Marigold, Chrysanthemum	Application of FYM, Bio- fertilizers and micronutrients, drip irrigation, mulching, suitable soil and water conservation practices
5	76.BLDmB2	Bheemanahalli: 4,5,6,7,19,20 ,21,22,23(1),24(1),24(2),25(1) ,25(2),30(3),34,35,38,40,41(1),41(2),42,44,45,51,52,53,55	Maize, Sorghum, Groundnut, Cotton, Bajra	Fruit crops: Amla, Custard apple Vegetables: Tomato, Chilli, Onion, Bhendi	Application of FYM, Biofertilizers and micronutrients, drip irrigation, mulching, suitable

LMU	Soil Map Units	Survey Number	Field Crops/ Commercial crops	Horticulture Crops (Rainfed/Irrigated)	Suitable Interventions
		,56,63, 64, 66(1),66(2), 66(3),66(4) Handaraki: 349,350,351,352, 360,419, 420		Flowers: Marigold, Chrysanthemum	soil and water conservation practices
6	110.JNKbB2	Handaraki: 247,248,249(10), 249/4	Maize, Sorghum Groundnut, Bajra	Fruit crops: Amla, Custard apple Vegetables: Tomato, Chilli, Brinjal, Bhendi, Onion Flowers: Marigold, Chrysanthemum	Application of FYM, Biofertilizers and micronutrients, drip irrigation, mulching, suitable soil and water conservation practices
7	5.BDLiB2 108.DSBiB2 165.HTKcB2 10.VNKiB2	Bheemanahalli: 12,14,23(2),3- 0(1),30(2),31,32,33,36,47,48, 54,57,58,59,62,65,67,68,69(1 ,69(2),73,74 275,276,277, 278,279	-	Agri-Silvi-Pasture: Custard apple, Hybrid Napier, <i>Styloxanthes hamata</i> , <i>Styloxanthes scabra</i>	Use of short duration varieties, sowing across the slope, drip irrigation is recommended
8	1.BDPiB2	Bheemanahalli : ,8,9(1),9(2) ,13,15,16,18,37 Handaraki : 370,371 Ramathirtha: 126,127,135,13 6,178,181, 182,183,186,187	-	Hybrid Napier, <i>Styloxanthes hamata</i> , <i>Styloxanthes scabra</i>	Use of short duration varieties, sowing across the slope, drip irrigation is recommended

SOIL HEALTH MANAGEMENT

8.1 Soil Health

Soil health is basic to plant health and plant health is basic to human and bovine health. Soil is fundamental to crop production. Without soil, no food could be produced nor would livestock be fed on a large scale. Because it is finite and fragile, soil is a precious resource that requires special care from its users.

Soil health or the capacity of the soil to function is critical to human survival. Soil health has been defined as: “the capacity of the soil to function as a living system without adverse effect on the ecosystem”. Healthy soils maintain a diverse community of soil organisms that help to form beneficial symbiotic associations with plant roots, recycle essential plant nutrients, improve soil structure with positive repercussions for soil, water and nutrient holding capacity and ultimately improve crop production and also contribute to mitigating climate change by maintaining or increasing its carbon content.

Functional interactions of soil biota with organic and inorganic components, air and water determine a soil’s potential to store and release nutrients and water to plants and to promote and sustain plant growth. Thus, maintaining soil health is vital to crop production and conserve soil resource base for sustaining agriculture.

The most important characteristics of a healthy soil are

- Good soil tilth
- Sufficient soil depth
- Good water storage and good drainage
- Adequate supply, but not excess of nutrients
- Large population of beneficial organisms
- Small proportion of plant pathogens and insect pests
- Low weed pressure
- Free of chemicals and toxins that may harm the crop
- Resistance to degradation
- Resilience when unfavorable conditions occur

Characteristics of Bhimanahalli-2 Microwatershed

- ❖ The soil phases identified in the microwatershed belonged to different soil series, BLD series occupies maximum area of 142 ha (25%), BDP 68 ha (12%) followed by MDG 54 ha (10%), DSB 45 ha (8%), BMN 38 ha (7%), BDL 32 ha (6%), JNK 21 ha (4%), NGP 19 ha (3%), VNK 13 ha (2%), HTK 7 ha (1%) and ANR 3 ha (<1%).
- ❖ As per land capability classification an area of 442 ha in the microwatershed falls under arable land category (Class II, III & IV). The major limitations identified in the arable lands were soil and erosion.

- ❖ On the basis of soil reaction an area of about 220 ha (38%) is neutral (pH 6.5-7.3), about 174 ha (30%) is slightly alkaline (pH 7.3-7.8) and a about 47 ha (8%) is moderately alkaline (pH 7.8-8.4) soil reaction in the microwatershed.

❖ **Soil Health Management**

The following actions are required to improve the current land husbandry practices that provide a sound basis for the successful adoption of sustainable crop production system.

Alkaline soils

Slightly alkaline soils cover an area of about 221 ha in the microwatershed

1. Regular addition of organic manure, green manuring, green leaf manuring, crop residue incorporation and mulching needs to be taken up to improve the soil organic matter status.
2. Application of Biofertilizers (Azospirillum, Azotobacter, Rhizobium).
3. Application of 25% extra N and P (125 % RDN&P).
4. Application of ZnSO₄ – 12.5 kg/ha (once in three years).
5. Application of Boron – 5kg/ha (once in three years).

Neutral soils

Entire cultivated area of about 220 ha is under neutral soils.

1. Regular addition of organic manure, green manuring, green leaf manuring, crop residue incorporation and mulching needs to be taken up to improve the soil organic matter status.
2. Application of Biofertilizers, (Azospirillum, Azotobacter, Rhizobium).
3. Application of 100 per cent RDF.
4. Need based micronutrient applications.

Besides the above recommendations, the best transfer of technology options are also to be adopted.

Soil Degradation

Soil erosion is one of the major factor affecting the soil health in the microwatershed. Out of total 573 ha area in the microwatershed, an area of about 45 ha (8%) is slight erosion and 396 ha (69%) is moderate erosion. These areas need immediate soil and water conservation and, other land development and land husbandry practices for restoring soil health.

Dissemination of Information and Communication of Benefits

Any large scale implementation of soil health management requires that supporting information is made available widely, particularly through channels familiar to farmers and extension workers. Given the very high priority attached to soil-health especially by the Central Government on issuing Soil-Health Cards to all the farmers, media outlets like Regional, State and National Newspapers, Radio and Dooradarshan

programs in local languages but also modern information and communication technologies such as Cellular phones and the Internet, which can be much more effective in reaching the younger farmers.

Inputs for Net Planning (Saturation Plan) and Interventions needed

Net planning (Saturation Plan) in IWMP is focusing on preparation of

1. Soil and Water Conservation Plan for each plot or farm.
2. Productivity enhancement measures/ interventions for existing crops/livestock/other farm enterprises.
3. Diversification of farming mainly with perennial horticultural crops and livestock.
4. Improving livelihood opportunities and income generating activities.

In this connection, how various outputs of Sujala-III are of use in addressing these objectives of Net Planning (Saturation Plan) are briefly presented below.


- ❖ **Soil Depth:** The depth of a soil decides the amount of moisture and nutrients it can hold, what crops can be taken up or not, depending on the rooting depth and the length of growing period available for raising any crop. Deeper the soil, better for a wide variety of crops. If sufficient depth is not available for growing deep rooted crops, either choose medium or short duration crops or deeper planting pits need to be opened and additional good quality soil brought from outside has to be filled into the planting pits.
- ❖ **Surface Soil Texture:** Lighter soil texture in the top soil means, better rain water infiltration, less run-off and soil moisture conservation, less capillary rise and less evaporation losses. Lighter surface textured soils are amenable to good soil tilth and are highly suitable for crops like groundnut, root vegetables (carrot, radish, potato etc) but not ideal for crops that need stagnant water like lowland paddy. Heavy textured soils are poor in water infiltration and percolation. They are prone for sheet erosion; such soils can be improved by sand mulching. The technology that is developed by the AICRP-Dryland Agriculture, Vijayapura, Karnataka can be adopted.
- ❖ **Gravelliness:** More gravel content is favorable for run-off harvesting but poor in soil moisture storage and nutrient availability. It is a significant parameter that decides the kind of crop to be raised.
- ❖ **Land Capability Classification:** The land capability map shows the areas suitable and not suitable for agriculture and the major constraints in each of the plot/survey number. Hence, one can decide what kind of enterprise is possible in each of these units. In general, erosion and soil are the major constraints in Bhimanahalli-2 microwatershed.
- ❖ **Organic Carbon:** The OC content (an index of available Nitrogen) is high (>0.75%) in an area of 432 ha (75%) and medium (0.5-0.75%) in about 10 ha (2%). The areas that are medium in OC needs to be further improved by applying farmyard manure and rotating crops with cereals and legumes or mixed cropping.

- ❖ **Promoting Green Manuring:** Growing of green manuring crops costs Rs. 1250/ha (green manuring seeds) and about Rs. 2000/ha towards cultivation that totals to Rs. 3250/- per ha. On the other hand, application of organic manure @ 10 tons/ha costs Rs. 5000/ha. The practice needs to be continued for 2-3 years or more. Nitrogen fertilizer needs to be supplemented by 25% in addition to the recommended level in 10 ha area where OC is medium (0.5-0.75%). For example, a rainfed maize, recommended level is 50 kg N per ha and an additional 12 kg /ha needs to be applied for all the crops grown in these plots.
- ❖ **Available Phosphorus:** Available Phosphorus is high (>57 kg/ha) covering an area of 363 ha (63%), medium (23-57 kg/ha) covering an area of 79 ha (14%) in the microwatershed. For all the crops 25% additional P needs to be applied where available P is medium.
- ❖ **Available Potassium:** Available potassium is high (>337 kg/ha) covering an area of 439 ha (77%) and medium (145-337 kg/ha) covering an area of 3 ha (<1%) in the microwatershed. All the plots, where available potassium is medium, additional 25% potassium may be applied.
- ❖ **Available Sulphur:** Available sulphur is a very critical nutrient for oilseed crops. It is low (<10 ppm) in the entire cultivated area of the microwatershed. Low areas need to be applied with magnesium sulphate or gypsum or Factamphos (p) fertilizer (13% sulphur) for 2-3 years for the deficiency to be corrected.
- ❖ **Available Boron:** Available boron content is low (<0.5 ppm) in the entire cultivated area of the microwatershed. For these low areas, application of sodium borate @ 10 kg/ha as soil application or 0.2 % borax as foliar spray is recommended.
- ❖ **Available Iron:** Available iron content is sufficient (>4.5 ppm) in an area of 436 ha (76%) and about deficient (<4.5 ppm) in about 5 ha (<1%) in the microwatershed. Deficient areas need to be applied with iron sulphate@25 kg/ha for 2-3 years.
- ❖ **Available Manganese:** Entire cultivated area in the microwatershed is sufficient in the available manganese content.
- ❖ **Available Copper:** Entire cultivated area in the microwatershed is sufficient in available copper content.
- ❖ **Available Zinc:** Available zinc content is deficient (<0.6 ppm) in the entire cultivated area of microwatershed. Application of zinc sulphate @25 kg/ha is recommended for the deficient areas.
- ❖ **Land Suitability for various crops:** Areas that are highly, moderately and marginally suitable for growing various crops are indicated. Along with the suitability, various constraints that are limiting the productivity are also indicated. For example, in case of cotton, gravel content, rooting depth and salinity/alkalinity are the major constraints in various plots. With suitable management interventions, the productivity can be enhanced. In order to increase the water holding capacity of light textured soils, growing of green manure crops and application of organic manure is recommended.

SOIL AND WATER CONSERVATION TREATMENT PLAN

For preparing soil and water conservation treatment plan for Bhimanahalli-2 microwatershed, the land resource inventory database generated under Sujala-III project has been transformed as information through series of interpretative (thematic) maps using soil phase map as a base. The various thematic maps (1:7920 scale) generated were

- Soil depth
- Surface soil texture
- Available water capacity
- Soil slope
- Soil gravelliness
- Land capability
- Present land use and land cover
- Crop suitability
- Rainfall
- Hydrology
- Water Resources
- Socio-economic data
- Contour plan with existing features- network of waterways, pottissa boundaries, cut up/ minor terraces etc.
- Cadastral map (1:7920 scale)
- Satellite imagery (1:7920 scale)



Apart from these, Hand Level/ Hydro Marker/ Dumpy Level/ Total Station and Kathedars' List to be collected.

Steps for Survey and Preparation of Treatment Plan

The boundaries of Land User Groups' and Survey No. boundaries are traced in the field.

- Naming of user groups and farmers
- Identification of arable and non arable lands
- Identification of drainage lines and gullies
- Identification of non treatable areas
- Identification of priority areas in the arable lands
- Treatment plan for arable lands
- Location of water harvesting and recharge structures

9.1 Treatment Plan

The treatment plan recommended for arable lands is briefly described below

9.1.1 Arable Land Treatment


A. BUNDING

Steps for Survey and Preparation of Treatment Plan	
<ul style="list-style-type: none"> • Cadastral map (1:7920 scale) is enlarged to a scale of 1:2500 scale • Existing network of waterways, pothissa boundaries, grass belts, natural drainage lines/ watercourse, cut ups/ terraces are marked on the cadastral map to the scale • Drainage lines are demarcated into 	
Small gullies	(up to 5 ha catchment)
Medium gullies	(5-15 ha catchment)
Ravines	(15-25 ha catchment) and
Halla/Nala	(more than 25ha catchment)

USER GROUP-1

CLASSIFICATION OF GULLIES

ಕೊರಕಲಿನ ವರ್ಗೀಕರಣ



Measurement of Land Slope

Land slope is estimated or determined by the study and interpretation of contours or by measurement in the field using simple instruments like Hand Level or Hydromarker.



Vertical and Horizontal intervals between bunds as recommended by the Watershed Development Department.

Slope percentage	Vertical interval (m)	Corresponding Horizontal Distance (m)
2 - 3%	0.6	24
3 - 4%	0.9	21
4 - 5%	0.9	21
5 - 6%	1.2	21
6 - 7%	1.2	21

Note: (i) The above intervals are maximum.

(ii) Considering the slope class and erosion status (A1... A=0-1 % slope, 1= slight erosion) the intervals have to be decided.

Bund length recording: Considering the contour plan and the existing grass belts/partitions, the bunds are aligned and lengths are measured.

Section of the Bund

Bund section is decided considering the soil texture class and gravelliness class (bg₀... b=loamy sand, g₀ = <15% gravel). The recommended Sections for different soils are given below.

Recommended Bund Section



Top width (m)	Base width (m)	Height (m)	Side slope (Z:1;H:V)	Cross section (sq m)	Soil Texture	Remarks
0.3	0.9	0.3	01:01	0.18	Sandy loam	Vegetative bund
0.3	1.2	0.3	1.5:1	0.225	Sandy clay	
0.3	1.2	0.5	0.9:1	0.375	Red gravelly soils	
0.3	1.2	0.6	0.75:1	0.45		
0.3	1.5	0.6	01:01	0.54	Red sandy loam	
0.3	2.1	0.6	1.5:1	0.72	Very shallow black soils	
0.45	2	0.75	01:01	0.92		
0.45	2.4	0.75	1.3:1	1.07	Shallow black soils	
0.6	3.1	0.7	1.78:1	1.29	Medium black soils	
0.5	3	0.85	1.47:1	1.49		

Formation of Trench cum Bund

Dimensions of the Borrow Pits/Trenches to be excavated (machinery are decided considering the Bund Section).

Details of Borrow Pit dimensions are given below:


TRENCH CUM BUND

0.45 Sq.m section


IDEAL FOR HORTICULTURE CROPS

'A' FRAME FOR INTERBUND MANAGEMENT



1. ಸಮವಾತಳ ಉಳುಮೆ

2. ಸಮವಾತಳ ಬಿತ್ತನೆ/ನಾಟಿ



Size of Borrow Pits/ Trench recommended for Trench cum Bund (by machinery)

Bund section	Bund length	Earth quantity	Pit				Berm (pit to pit)	Soil depth class
			L(m)	W(m)	D(m)	Quantity (m ³)		
m ²	M	m ³					m	
0.375	6	2.25	5.85	0.85	0.45	2.24	0.15	Shallow
0.45	6	2.7	5.4	1.2	0.43	2.79	0.6	Shallow
0.45	6	2.7	5	0.85	0.65	2.76	1	Moderately Shallow
0.54	5.6	3.02	5.5	0.85	0.7	3.27	0.1	Moderately shallow
0.54	5.5	2.97	5	1.2	0.5	3	0.5	Shallow
0.72	6.2	4.46	6	1.2	0.7	5.04	0.2	Moderately shallow
0.72	5.2	3.74	5.1	0.85	0.9	3.9	0.1	Moderately deep

B. Water Ways

- Existing waterways are marked on the cadastral map (1:7920 scale) and their dimensions are recorded.
- Considering the contour plan of the MWS, additional waterways/ modernization of the existing ones can be thought of.
- The design details are given in the Manual.

C. Farm Ponds

Waterways and the catchment area will give an indication on the size of the Farm Pond. Location of the pond can be decided based on the contour plan/ field condition and farmers' need/desire.

D. Diversion Channel

Existing EPT/ CPT are marked on the cadastral map. Looking to the need, these can be modernized or fresh diversion channel can be proposed and runoff from this can be stored in Gokatte/ Recharge ponds.

9.1.2 Non-Arable Land Treatment

Depending on the gravelliness and crops preferred by the farmers, the concerned authorities can decide appropriate treatment plan. The recommended treatments may be Contour Trench, Staggered Trench, Crescent Bund, Boulder Bund or Pebble Bund.

9.1.3 Treatment of Natural Water Course/ Drainage Lines

- a) The cadastral map has to be updated as regards the network of drainage lines (gullies/ *nalas/ hallas*) and existing structures are marked to the scale and storage capacity of the existing water bodies are documented.
- b) The drainage line will be demarcated into Upper Reach, Middle Reach and Lower Reach.
- c) Considering the Catchment, *Nala* bed and bank conditions, suitable structures are decided.
- d) Number of storage structures (Check dam/ *Nala* bund/ Percolation tank) will be decided considering the commitments and available runoff from water budgeting and quality of water in the wells and site suitability.
- e) Detailed Leveling Survey using Dumpy Level / Total Station has to be carried out to arrive at the site-specific designs as shown in the Manual.
- f) The location of ground water recharge structures are decided by examining the lineaments and fracture zones from geological maps.
- g) Rainfall intensity data of the nearest Rain Gauge Station is considered for Hydrologic Designs.
- h) Silt load to the Storage/Recharge structures is reduced by providing vegetative, boulder and earthen checks in the natural water course. Location and design details are given in the Manual.

9.2 Recommended Soil and Water Conservation Measures

The appropriate conservation structures best suited for each of the land parcel/ survey number (Appendix-I) are selected based on the slope per cent, severity of erosion, amount of rainfall, land use and soil type. The different kinds of conservation structures recommended are:

1. Graded / Strengthening of Bunds
2. Trench cum Bunds (TCB)
3. Trench cum Bunds / Strengthening
4. Crescent Bunds

A map (Fig. 9.1) showing soil and water conservation plan with different kinds of structures recommended has been prepared which shows the spatial distribution and extent of area. An area of about 126 ha (22%) needs trench cum bunding, about 315 ha (55%) needs Graded bunding and about 1 ha (<1%) requires strengthening of existing bunds.

The conservation plan prepared may be presented to all the stakeholders including farmers and after considering their suggestions, the conservation plan for the microwatershed may be finalised in a participatory approach.




Fig. 9.1 Soil and Water Conservation Plan map of Bhimanahalli-2 Microwatershed

9.3 Greening of Microwatershed

As part of the greening programme in the watersheds, it is envisaged to plant a variety of horticultural and other tree plants that are edible, economical and produce lot of biomass which helps to restore the ecological balance in the watersheds. The lands that are suitable for greening programme are non-arable lands (land capability classes V, VI VII and VIII) and also the lands that are not suitable or marginally suitable and field bunds for growing annual and perennial crops. The method of planting these trees is given below.

It is recommended to open pits during the 1st week of March along the contour and heap the dug out soil on the lower side of the slope in order to harness the flowing water and facilitate weathering of soil in the pit. Exposure of soil in the pit also prevents spread of pests and diseases due to scorching sun rays. The pits should be filled with mixture of soil and organic manure during the second week of April and keep ready with sufficiently tall seedlings produced either in poly bags or in root trainer nurseries so that planting can be done during the 2nd or 3rd week of April depending on the rainfall.

The tree species suitable for the area considering rainfall, temperature and adaptability is listed below; waterlogged areas are recommended to be planted with species like Nerale (*Syzgium cumini*) and Bamboo. Dry areas are to be planted with species like Honge, Bevu, Seetaphal etc.

Dry Deciduous Species			Temp (°C)	Rainfall (mm)
1.	Bevu	<i>Azadiracta indica</i>	21–32	400 –1,200
2.	Tapasi	<i>Holoptelia integrifolia</i>	20-30	500 - 1000
3.	Seetaphal	<i>Anona Squamosa</i>	20-40	400 - 1000
4.	Honge	<i>Pongamia pinnata</i>	20 -50	500– 2,500
5.	Kamara	<i>Hardwickia binata</i>	25 -35	400 - 1000
6.	Bage	<i>Albezzia lebbek</i>	20 - 45	500 - 1000
7.	Ficus	<i>Ficus bengalensis</i>	20 - 50	500–2,500
8.	Sisso	<i>Dalbargia Sissoo</i>	20 - 50	500 -2000
9.	Ailanthus	<i>Ailanthus excelsa</i>	20 - 50	500 - 1000
10.	Hale	<i>Wrightia tinctoria</i>	25 - 45	500 - 1000
11.	Uded	<i>Steriospermum chelanoides</i>	25 - 45	500 -2000
12.	Dhupa	<i>Boswella Serrata</i>	20 - 40	500 - 2000
13.	Nelli	<i>Embllica Officinalis</i>	20 - 50	500 -1500
14.	Honne	<i>Pterocarpus marsupium</i>	20 - 40	500 - 2000
Moist Deciduous Species			Temp (°C)	Rainfall (mm)
15.	Teak	<i>Tectona grandis</i>	20 - 50	500-5000
16.	Nandi	<i>Legarstroemia lanceolata</i>	20 - 40	500 - 4000
17.	Honne	<i>Pterocarpus marsupium</i>	20 - 40	500 - 3000
18.	Mathi	<i>Terminalia alata</i>	20 -50	500 - 2000
19.	Shivane	<i>Gmelina arborea</i>	20 -50	500 -2000
20.	Kindal	<i>T.Paniculata</i>	20 - 40	500 - 1500
21.	Beete	<i>Dalbargia latifolia</i>	20 - 40	500 - 1500
22.	Tare	<i>T. belerica</i>	20 - 40	500 - 2000
23.	Bamboo	<i>Bambusa arundinasia</i>	20 - 40	500 - 2500
24.	Bamboo	<i>Dendrocalamus strictus</i>	20 – 40	500 – 2500
25.	Muthuga	<i>Butea monosperma</i>	20 - 40	400 - 1500
26.	Hippe	<i>Madhuca latifolia</i>	20 - 40	500 - 2000
27.	Sandal	<i>Santalum album</i>	20 - 50	400 - 1000
28.	Nelli	<i>Embllica officinalis</i>	20 - 40	500 - 2000
29.	Nerale	<i>Sizygium cumini</i>	20 - 40	500 - 2000
30.	Dhaman	<i>Grevia tilifolia</i>	20 - 40	500 - 2000
31.	Kaval	<i>Careya arborea</i>	20 - 40	500 - 2000
32.	Harada	<i>Terminalia chebula</i>	20 - 40	500 - 2000

References

1. FAO (1976) Framework for Land Evaluation, Food and Agriculture Organization, Rome. 72 pp.
2. FAO (1983) Guidelines for Land Evaluation for Rainfed Agriculture, FAO, Rome, 237 pp.
3. IARI (1971) Soil Survey Manual, All India Soil and Land Use Survey Organization, IARI, New Delhi, 121 pp.
4. Katyal, J.C. and Rattan, R.K. (2003) Secondary and Micronutrients; Research Gap and Future Needs. Fert. News 48 (4); 9-20.
5. Naidu, L.G.K., Ramamurthy, V., Challa, O., Hegde, R. and Krishnan, P. (2006) Manual Soil Site Suitability Criteria for Major Crops, NBSS Publ. No. 129, NBSS & LUP, Nagpur, 118 pp.
6. Natarajan, A. and Dipak Sarkar (2010) Field Guide for Soil Survey, National Bureau of Soil Survey and Land Use Planning (ICAR), Nagpur, India.
7. Natarajan, A., Rajendra Hegde, Raj, J.N. and Shivananda Murthy, H.G. (2015) Implementation Manual for Sujala-III Project, Watershed Development Department, Bengaluru, Karnataka.
8. Sarma, V.A.K., Krishnan, P. and Budihal, S.L. (1987) Laboratory Manual, Tech. Bull. 23, NBSS & LUP, Nagpur.
9. Sehgal, J.L. (1990) Soil Resource Mapping of Different States of India; Why and How?, National Bureau of Soil Survey and Land Use Planning, Nagpur, 49 pp.
10. Shivaprasad, C.R., R.S. Reddy, J. Sehgal and M. Velayuthum (1998) Soils of Karnataka for Optimizing Land Use, NBSS Publ. No. 47b, NBSS & LUP, Nagpur, India.
11. Soil Survey Staff (2006) Keys to Soil Taxonomy, Tenth edition, U.S. Department of Agriculture/ NRCS, Washington DC, U.S.A.
12. Soil Survey Staff (2012) Soil Survey Manual, Handbook No. 18, USDA, Washington DC, USA.

Appendix I
Bhimanahalli-2 (1A2b) Microwatershed
Soil Phase Information

Village	Survey Number	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Bheema nahalli	2(2)	0.04	BMNmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Ies	Graded bunding
Bheema nahalli	23(1)	2.11	BLDmB2	LMU-5	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Ies	Graded bunding
Bheema nahalli	23(2)	1.82	DSBiB2	LMU-7	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	IIes	Trench cum bunding
Bheema nahalli	24(1)	1.82	BLDmB2	LMU-5	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	Ies	Graded bunding
Bheema nahalli	24(2)	2.34	BLDmB2	LMU-5	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	Ies	Graded bunding
Bheema nahalli	25(1)	3.98	BLDmB2	LMU-5	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Ies	Graded bunding
Bheema nahalli	25(2)	0.23	BLDmB2	LMU-5	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Ies	Graded bunding
Bheema nahalli	287(1)	0.22	BMNmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Ies	Graded bunding
Bheema nahalli	29(1)	1	BMNmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Ies	Graded bunding
Bheema nahalli	29(2)	0.49	BMNmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	Ies	Graded bunding
Bheema nahalli	29(3)	1.05	BMNmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Ies	Graded bunding
Bheema nahalli	29(4)	1.01	BMNmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Ies	Graded bunding
Bheema nahalli	30(1)	1.91	DSBiB2	LMU-7	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	IIes	Trench cum bunding
Bheema nahalli	30(2)	0.79	DSBiB2	LMU-7	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	IIes	Trench cum bunding
Bheema nahalli	30(3)	0.33	BLDmB2	LMU-5	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	Ies	Graded bunding
Bheema nahalli	41(1)	1.8	BLDmB2	LMU-5	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Ies	Graded bunding
Bheema nahalli	41(2)	1.71	BLDmB2	LMU-5	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Ies	Graded bunding
Bheema nahalli	50(1)	2.38	RO	RO	RO	RO	RO	RO	RO	RO	Scrub land (Sl)	Not Available	RO	RO
Bheema nahalli	50(2)	3.21	RO	RO	RO	RO	RO	RO	RO	RO	Jowar (Jw)	Not Available	RO	RO
Bheema nahalli	66(1)	2.43	BLDmB2	LMU-5	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Ies	Graded bunding
Bheema nahalli	66(2)	0.38	BLDmB2	LMU-5	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Ies	Graded bunding
Bheema nahalli	66(3)	0.37	BLDmB2	LMU-5	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Ies	Graded bunding
Bheema nahalli	66(4)	0.53	BLDmB2	LMU-5	Moderately	Clay	Non gravelly	Low (51-100	Very gently	Moderate	Redgram (Rg)	Not Available	Ies	Graded

Village	Survey Number	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
nahalli					shallow (50-75 cm)		(<15%)	mm/m)	sloping (1-3%)	te		Available		bunding
Bheema nahalli	69(1)	0.73	BDLiB2	LMU-7	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	IIes	Graded bunding
Bheema nahalli	69(2)	2.54	BDLiB2	LMU-7	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Bheema nahalli	72(1)	0.8	MDGhB2	LMU-3	Deep (100-150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Ies	Graded bunding
Bheema nahalli	72(2)	1.63	MDGhB2	LMU-3	Deep (100-150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Ies	Graded bunding
Bheema nahalli	9(1)	2.2	BDPiB2	LMU-8	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	IVs	Trench cum bunding
Bheema nahalli	9(2)	1.63	BDPiB2	LMU-8	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IVs	Trench cum bunding
Bheema nahalli	3	0.51	BMNmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Ies	Graded bunding
Bheema nahalli	4	4.34	BLDmB2	LMU-5	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Ies	Graded bunding
Bheema nahalli	5	5.66	BLDmB2	LMU-5	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Ies	Graded bunding
Bheema nahalli	6	5.16	BLDmB2	LMU-5	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Ies	Graded bunding
Bheema nahalli	7	6.56	BLDmB2	LMU-5	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Ies	Graded bunding
Bheema nahalli	8	5.59	BDPiB2	LMU-8	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IVs	Trench cum bunding
Bheema nahalli	10	3.49	BMNmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Ies	Graded bunding
Bheema nahalli	11	4.31	BMNmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Ies	Graded bunding
Bheema nahalli	12	1.05	VNKiB2	LMU-7	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Trench cum bunding
Bheema nahalli	13	4.91	BDPiB2	LMU-8	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IVs	Trench cum bunding
Bheema nahalli	14	1	VNKiB2	LMU-7	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Trench cum bunding
Bheema nahalli	15	2.84	BDPiB2	LMU-8	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IVs	Trench cum bunding
Bheema nahalli	16	3.36	BDPiB2	LMU-8	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IVs	Trench cum bunding
Bheema nahalli	17	9.41	RO	RO	RO	RO	RO	RO	RO	RO	Redgram+Fallow land+Scrub land (Rg+Fl+Sl)	Not Available	RO	RO
Bheema nahalli	18	4.25	BDPiB2	LMU-8	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IVs	Trench cum bunding
Bheema nahalli	19	5.76	BLDmB2	LMU-5	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Ies	Graded bunding
Bheema nahalli	20	3.64	BLDmB2	LMU-5	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Scrub land (Rg+Sl)	Not Available	Ies	Graded bunding

Village	Survey Number	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Bheema nahalli	21	5.47	BLDmB2	LMU-5	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Ies	Graded bunding
Bheema nahalli	22	2.33	BLDmB2	LMU-5	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Ies	Graded bunding
Bheema nahalli	26	2	BMNmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Ies	Graded bunding
Bheema nahalli	27	2.34	BMNmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Ies	Graded bunding
Bheema nahalli	28	0.03	BMNmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	Ies	Graded bunding
Bheema nahalli	31	4.05	DSBiB2	LMU-7	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	IIes	Trench cum bunding
Bheema nahalli	32	0.95	DSBiB2	LMU-7	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	IIes	Trench cum bunding
Bheema nahalli	33	2.22	DSBiB2	LMU-7	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	IIes	Trench cum bunding
Bheema nahalli	34	5.89	BLDmB2	LMU-5	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Ies	Graded bunding
Bheema nahalli	35	3.97	BLDmB2	LMU-5	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Scrub land (SI)	Not Available	Ies	Graded bunding
Bheema nahalli	36	9.47	DSBiB2	LMU-7	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Trench cum bunding
Bheema nahalli	37	2.52	BDPiB2	LMU-8	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IVs	Trench cum bunding
Bheema nahalli	38	4.03	BLDmB2	LMU-5	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Cotton (Rg+Ct)	Not Available	Ies	Graded bunding
Bheema nahalli	40	0.04	BLDmB2	LMU-5	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Ies	Graded bunding
Bheema nahalli	42	3.21	BLDmB2	LMU-5	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Ies	Graded bunding
Bheema nahalli	43	10.08	RO	RO	RO	RO	RO	RO	RO	RO	Scrub land (SI)	Not Available	RO	RO
Bheema nahalli	44	3.02	BLDmB2	LMU-5	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Ies	Graded bunding
Bheema nahalli	45	2.48	BLDmB2	LMU-5	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	Ies	Graded bunding
Bheema nahalli	46	3.79	RO	RO	RO	RO	RO	RO	RO	RO	Jowar+Scrub land (Jw+SI)	Not Available	RO	RO
Bheema nahalli	47	6.09	DSBiB2	LMU-7	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	IIes	Trench cum bunding
Bheema nahalli	48	8.77	DSBiB2	LMU-7	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Cotton (Rg+Ct)	Not Available	IIes	Trench cum bunding
Bheema nahalli	49	2.94	RO	RO	RO	RO	RO	RO	RO	RO	Scrub land (SI)	Not Available	RO	RO
Bheema nahalli	51	4.06	BLDmB2	LMU-5	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	Ies	Graded bunding
Bheema nahalli	52	5.54	BLDmB2	LMU-5	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	Ies	Graded bunding

Village	Survey Number	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Bheema nahalli	53	2.96	BLDmB2	LMU-5	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iles	Graded bunding
Bheema nahalli	54	6.82	DSBiB2	LMU-7	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIles	Trench cum bunding
Bheema nahalli	55	3.91	BLDmB2	LMU-5	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Bheema nahalli	56	6.07	BLDmB2	LMU-5	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Bheema nahalli	57	4.59	DSBiB2	LMU-7	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	IIles	Trench cum bunding
Bheema nahalli	58	6.58	DSBiB2	LMU-7	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	IIles	Trench cum bunding
Bheema nahalli	59	1.76	DSBiB2	LMU-7	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	IIles	Trench cum bunding
Bheema nahalli	60	6.5	BMNmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	Iles	Graded bunding
Bheema nahalli	61	4.39	BMNmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	Iles	Graded bunding
Bheema nahalli	62	3.78	BDLiB2	LMU-7	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIles	Graded bunding
Bheema nahalli	63	9.11	BLDmB2	LMU-5	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Bheema nahalli	64	7.93	BLDmB2	LMU-5	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Bheema nahalli	65	5.18	BDLiB2	LMU-7	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIles	Graded bunding
Bheema nahalli	67	0.37	BDLiB2	LMU-7	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIles	Graded bunding
Bheema nahalli	68	6.02	BDLiB2	LMU-7	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	IIles	Graded bunding
Bheema nahalli	70	2.2	ANRiB2	LMU-2	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IVes	Graded bunding
Bheema nahalli	71	1.93	MDGhB2	LMU-3	Deep (100-150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Bheema nahalli	73	4.01	BDLiB2	LMU-7	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Jowar (Rg+Jw)	Not Available	IIles	Graded bunding
Bheema nahalli	74	0.12	BDLiB2	LMU-7	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Jowar (Rg+Jw)	Not Available	IIles	Graded bunding
Bheema nahalli	75	1.47	MDGhB2	LMU-3	Deep (100-150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Bheema nahalli	76	1.01	BMNmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	Iles	Graded bunding
Bheema nahalli	77	0.3	BMNmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	Iles	Graded bunding
Bheema nahalli	78	0.77	BMNmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	Iles	Graded bunding
Bheema nahalli	79	1.23	BMNmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	Iles	Graded bunding

Village	Survey Number	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Bheema nahalli	92	0.77	MDGhB2	LMU-3	Deep (100-150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Bheema nahalli	275	2.37	VNKiB2	LMU-7	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Trench cum bunding
Bheema nahalli	276	1.55	VNKiB2	LMU-7	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Trench cum bunding
Bheema nahalli	277	1.6	VNKiB2	LMU-7	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Trench cum bunding
Bheema nahalli	278	3.28	VNKiB2	LMU-7	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Trench cum bunding
Bheema nahalli	279	0.31	VNKiB2	LMU-7	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Trench cum bunding
Bheema nahalli	281	1.25	BMNmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Bheema nahalli	282	2.2	BMNmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Handara ki	249(10)	0.86	JNKhB2	LMU-6	Moderately shallow (50-75 cm)	Sandy clay loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Scrub land (Rg+Sl)	Not Available	Iles	Graded bunding
Handara ki	247	0.11	JNKhB2	LMU-6	Moderately shallow (50-75 cm)	Sandy clay loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Scrub land (Rg+Sl)	Not Available	Iles	Graded bunding
Handara ki	248	2.05	JNKhB2	LMU-6	Moderately shallow (50-75 cm)	Sandy clay loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Handara ki	249/1	68.15	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Scrub land (Sl)	Not Available	Forest	Forest
Handara ki	249/4	4.74	JNKhB2	LMU-6	Moderately shallow (50-75 cm)	Sandy clay loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Scrub land (Rg+Sl)	Not Available	Iles	Graded bunding
Handara ki	249/5	4.84	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Redgram (Rg)	Not Available	Forest	Forest
Handara ki	249/7	2.28	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Scrub land (Sl)	Not Available	Forest	Forest
Handara ki	249/8	1.35	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Scrub land (Sl)	Not Available	Forest	Forest
Handara ki	249/9	0.67	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Scrub land (Sl)	Not Available	Forest	Forest
Handara ki	250	4.49	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Redgram (Rg)	Not Available	Forest	Forest
Handara ki	251	7.2	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Scrub land (Sl)	Not Available	Forest	Forest
Handara ki	252	6.1	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Redgram+Jowar (Rg+Jw)	Not Available	Forest	Forest
Handara ki	253	2.8	MDGmB1	LMU-3	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Redgram (Rg)	Not Available	IIs	Graded bunding
Handara ki	254	3.82	MDGmB1	LMU-3	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Redgram (Rg)	Not Available	IIs	Graded bunding
Handara ki	255	5.03	MDGmB1	LMU-3	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Redgram (Rg)	Not Available	IIs	Graded bunding
Handara ki	256	7.5	MDGmB1	LMU-3	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Redgram (Rg)	Not Available	IIs	Graded bunding

Village	Survey Number	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Handara ki	257	1.62	MDGmB1	LMU-3	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Redgram (Rg)	Not Available	IIs	Graded bunding
Handara ki	258	3.83	MDGmB1	LMU-3	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Redgram (Rg)	Not Available	IIs	Graded bunding
Handara ki	259	6.56	MDGmB1	LMU-3	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Redgram+Jowar (Rg+Jw)	Not Available	IIs	Graded bunding
Handara ki	260	0.52	MDGmB1	LMU-3	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Redgram+Jowar (Rg+Jw)	Not Available	IIs	Graded bunding
Handara ki	261	0.72	MDGmB1	LMU-3	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Redgram (Rg)	Not Available	IIs	Graded bunding
Handara ki	343	3.52	MDGmB1	LMU-3	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Redgram+Cotton+Jowar (Rg+Ct+Jw)	Not Available	IIs	Graded bunding
Handara ki	344	8.77	MDGmB1	LMU-3	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Redgram (Rg)	Not Available	IIs	Graded bunding
Handara ki	345	1.08	MDGmB1	LMU-3	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Jowar (Jw)	Not Available	IIs	Graded bunding
Handara ki	346	1.24	MDGmB1	LMU-3	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Jowar (Jw)	Not Available	IIs	Graded bunding
Handara ki	347	7.65	RO	RO	RO	RO	RO	RO	RO	RO	Quarry	Not Available	RO	RO
Handara ki	348	7.14	Quarry	Quarry	Quarry	Quarry	Quarry	Quarry	Quarry	Quarry	Quarry	Not Available	Quarry	Quarry
Handara ki	349	3.86	BLDmB2	LMU-5	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	IIs	Graded bunding
Handara ki	350	2.06	BLDmB2	LMU-5	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	IIs	Graded bunding
Handara ki	351	4.84	BLDmB2	LMU-5	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	IIs	Graded bunding
Handara ki	352	3.57	BLDmB2	LMU-5	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	RO	Not Available	IIs	Graded bunding
Handara ki	353	3.05	RO	RO	RO	RO	RO	RO	RO	RO	RO	Not Available	RO	RO
Handara ki	354	1.81	RO	RO	RO	RO	RO	RO	RO	RO	RO	Not Available	RO	RO
Handara ki	360	0.23	BLDmB2	LMU-5	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	IIs	Graded bunding
Handara ki	370	3.38	BDPiB2	LMU-8	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IVs	Trench cum bunding
Handara ki	371	0.16	BDPiB2	LMU-8	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IVs	Trench cum bunding
Handara ki	419	1.66	BLDmB2	LMU-5	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	IIs	Graded bunding
Handara ki	420	0.04	BLDmB2	LMU-5	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	IIs	Graded bunding
Handara ki	421	0.64	BMNmA1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Cotton (Ct)	Not Available	IIs	Graded bunding
Handara ki	422	0.06	BMNmA1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Cotton (Ct)	Not Available	IIs	Graded bunding

Village	Survey Number	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Ramathirtha	126	2.34	BDPiB2	LMU-8	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IVs	Trench cum bunding
Ramathirtha	127	3.71	BDPiB2	LMU-8	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IVs	Trench cum bunding
Ramathirtha	128	3.65	NGPmB2	LMU-4	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	IIs	Graded bunding
Ramathirtha	134	2.92	NGPmB2	LMU-4	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIs	Graded bunding
Ramathirtha	135	3.14	BDPiB2	LMU-8	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	IVs	Trench cum bunding
Ramathirtha	136	0.02	BDPiB2	LMU-8	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	IVs	Trench cum bunding
Ramathirtha	178	1.13	BDPiB2	LMU-8	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IVs	Trench cum bunding
Ramathirtha	180	1.83	NGPmB2	LMU-4	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIs	Graded bunding
Ramathirtha	181	7.25	BDPiB2	LMU-8	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IVs	Trench cum bunding
Ramathirtha	182	8.01	BDPiB2	LMU-8	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IVs	Trench cum bunding
Ramathirtha	183	4.44	BDPiB2	LMU-8	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IVs	Trench cum bunding
Ramathirtha	184	2.11	NGPmB2	LMU-4	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIs	Graded bunding
Ramathirtha	185	3.53	NGPmB2	LMU-4	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIs	Graded bunding
Ramathirtha	186	1.97	BDPiB2	LMU-8	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IVs	Trench cum bunding
Ramathirtha	187	2.55	BDPiB2	LMU-8	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IVs	Trench cum bunding

Appendix III
Bhimanahalli-2 (1A2b) Microwatershed
Soil Suitability Information

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry
Bheemanahalli	2(2)	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz
Bheemanahalli	23(1)	N1r	S2r	S3r	S2rt	S3r	S2rt	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	S3r	S3r	S3r	S2rt	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r
Bheemanahalli	23(2)	N1r	S3rg	N1r	S3rg	N1r	S3rg	N1r	N1r	S3rt	N1r	N1r	S3r	N1r	S3r	N1r	N1r	N1r	S3r	S3rg	S3rg	S3rg	S3rg	S3rg	N1r	S3r	S3rg	S3rg	N1r	N1r
Bheemanahalli	24(1)	N1r	S2r	S3r	S2rt	S3r	S2rt	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	S3r	S3r	S3r	S2rt	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r
Bheemanahalli	24(2)	N1r	S2r	S3r	S2rt	S3r	S2rt	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	S3r	S3r	S3r	S2rt	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r
Bheemanahalli	25(1)	N1r	S2r	S3r	S2rt	S3r	S2rt	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	S3r	S3r	S3r	S2rt	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r
Bheemanahalli	25(2)	N1r	S2r	S3r	S2rt	S3r	S2rt	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	S3r	S3r	S3r	S2rt	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r
Bheemanahalli	287(1)	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz
Bheemanahalli	29(1)	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz
Bheemanahalli	29(2)	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz
Bheemanahalli	29(3)	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz
Bheemanahalli	29(4)	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz
Bheemanahalli	30(1)	N1r	S3rg	N1r	S3rg	N1r	S3rg	N1r	N1r	S3rt	N1r	N1r	S3r	N1r	S3r	N1r	N1r	N1r	S3r	S3rg	S3rg	S3rg	S3rg	S3rg	N1r	S3r	S3rg	S3rg	N1r	N1r
Bheemanahalli	30(2)	N1r	S3rg	N1r	S3rg	N1r	S3rg	N1r	N1r	S3rt	N1r	N1r	S3r	N1r	S3r	N1r	N1r	N1r	S3r	S3rg	S3rg	S3rg	S3rg	S3rg	N1r	S3r	S3rg	S3rg	N1r	N1r
Bheemanahalli	30(3)	N1r	S2r	S3r	S2rt	S3r	S2rt	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	S3r	S3r	S3r	S2rt	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r
Bheemanahalli	41(1)	N1r	S2r	S3r	S2rt	S3r	S2rt	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	S3r	S3r	S3r	S2rt	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r
Bheemanahalli	41(2)	N1r	S2r	S3r	S2rt	S3r	S2rt	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	S3r	S3r	S3r	S2rt	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r
Bheemanahalli	50(1)	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO
Bheemanahalli	50(2)	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brimjal	Bhendi	Drumstick	Mulberry	
Bheeman ahalli	66(1)	N1r	S2r	S3r	S2rt	S3r	S2rt	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	S3r	S3r	S3r	S2rt	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Bheeman ahalli	66(2)	N1r	S2r	S3r	S2rt	S3r	S2rt	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	S3r	S3r	S3r	S2rt	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Bheeman ahalli	66(3)	N1r	S2r	S3r	S2rt	S3r	S2rt	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	S3r	S3r	S3r	S2rt	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Bheeman ahalli	66(4)	N1r	S2r	S3r	S2rt	S3r	S2rt	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	S3r	S3r	S3r	S2rt	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Bheeman ahalli	69(1)	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Bheeman ahalli	69(2)	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Bheeman ahalli	72(1)	S3n	S2n	S3n	S2tn	N1n	S3tn	N1n	S3n	S3t	S3n	S2tn	N1n	N1n	S3n	N1n	N1n	S3n	S3n	N1n	S3n	S3n	S3n	S3n	S3n	S2n	S3n	S3n	N1n	N1n	
Bheeman ahalli	72(2)	S3n	S2n	S3n	S2tn	N1n	S3tn	N1n	S3n	S3t	S3n	S2tn	N1n	N1n	S3n	N1n	N1n	S3n	S3n	N1n	S3n	S3n	S3n	S3n	S3n	S2n	S3n	S3n	N1n	N1n	
Bheeman ahalli	9(1)	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Bheeman ahalli	9(2)	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Bheeman ahalli	3	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz	
Bheeman ahalli	4	N1r	S2r	S3r	S2rt	S3r	S2rt	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	S3r	S3r	S3r	S2rt	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Bheeman ahalli	5	N1r	S2r	S3r	S2rt	S3r	S2rt	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	S3r	S3r	S3r	S2rt	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Bheeman ahalli	6	N1r	S2r	S3r	S2rt	S3r	S2rt	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	S3r	S3r	S3r	S2rt	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Bheeman ahalli	7	N1r	S2r	S3r	S2rt	S3r	S2rt	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	S3r	S3r	S3r	S2rt	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Bheeman ahalli	8	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Bheeman ahalli	10	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz	
Bheeman ahalli	11	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz	
Bheeman ahalli	12	N1r	S3r	N1r	S3r	N1r	S3r	N1r	N1r	S3rt	N1r	N1r	S3r	N1r	S3r	N1r	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Bheeman ahalli	13	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Bheeman ahalli	14	N1r	S3r	N1r	S3r	N1r	S3r	N1r	N1r	S3rt	N1r	N1r	S3r	N1r	S3r	N1r	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brimjal	Bhendi	Drumstick	Mulberry	
Bheeman ahalli	15	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	
Bheeman ahalli	16	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Bheeman ahalli	17	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO
Bheeman ahalli	18	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Bheeman ahalli	19	N1r	S2r	S3r	S2rt	S3r	S2rt	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	S3r	S3r	S3r	S2rt	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Bheeman ahalli	20	N1r	S2r	S3r	S2rt	S3r	S2rt	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	S3r	S3r	S3r	S2rt	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Bheeman ahalli	21	N1r	S2r	S3r	S2rt	S3r	S2rt	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	S3r	S3r	S3r	S2rt	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Bheeman ahalli	22	N1r	S2r	S3r	S2rt	S3r	S2rt	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	S3r	S3r	S3r	S2rt	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Bheeman ahalli	26	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz	
Bheeman ahalli	27	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz	
Bheeman ahalli	28	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz	
Bheeman ahalli	31	N1r	S3rg	N1r	S3rg	N1r	S3rg	N1r	N1r	S3rt	N1r	N1r	S3r	N1r	S3r	N1r	N1r	N1r	S3r	S3rg	S3rg	S3rg	S3rg	S3rg	N1r	S3r	S3rg	S3rg	N1r	N1r	
Bheeman ahalli	32	N1r	S3rg	N1r	S3rg	N1r	S3rg	N1r	N1r	S3rt	N1r	N1r	S3r	N1r	S3r	N1r	N1r	N1r	S3r	S3rg	S3rg	S3rg	S3rg	S3rg	N1r	S3r	S3rg	S3rg	N1r	N1r	
Bheeman ahalli	33	N1r	S3rg	N1r	S3rg	N1r	S3rg	N1r	N1r	S3rt	N1r	N1r	S3r	N1r	S3r	N1r	N1r	N1r	S3r	S3rg	S3rg	S3rg	S3rg	S3rg	N1r	S3r	S3rg	S3rg	N1r	N1r	
Bheeman ahalli	34	N1r	S2r	S3r	S2rt	S3r	S2rt	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	S3r	S3r	S3r	S2rt	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Bheeman ahalli	35	N1r	S2r	S3r	S2rt	S3r	S2rt	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	S3r	S3r	S3r	S2rt	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Bheeman ahalli	36	N1r	S3rg	N1r	S3rg	N1r	S3rg	N1r	N1r	S3rt	N1r	N1r	S3r	N1r	S3r	N1r	N1r	N1r	S3r	S3rg	S3rg	S3rg	S3rg	S3rg	N1r	S3r	S3rg	S3rg	N1r	N1r	
Bheeman ahalli	37	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Bheeman ahalli	38	N1r	S2r	S3r	S2rt	S3r	S2rt	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	S3r	S3r	S3r	S2rt	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Bheeman ahalli	40	N1r	S2r	S3r	S2rt	S3r	S2rt	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	S3r	S3r	S3r	S2rt	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Bheeman ahalli	42	N1r	S2r	S3r	S2rt	S3r	S2rt	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	S3r	S3r	S3r	S2rt	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brimjal	Bhendi	Drumstick	Mulberry	
Bheeman ahalli	43	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	
Bheeman ahalli	44	N1r	S2r	S3r	S2rt	S3r	S2rt	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	S3r	S3r	S3r	S2rt	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Bheeman ahalli	45	N1r	S2r	S3r	S2rt	S3r	S2rt	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	S3r	S3r	S3r	S2rt	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Bheeman ahalli	46	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO
Bheeman ahalli	47	N1r	S3rg	N1r	S3rg	N1r	S3rg	N1r	N1r	S3rt	N1r	N1r	S3r	N1r	S3r	N1r	N1r	N1r	S3r	S3rg	S3rg	S3rg	S3rg	S3rg	N1r	S3r	S3rg	S3rg	N1r	N1r	
Bheeman ahalli	48	N1r	S3rg	N1r	S3rg	N1r	S3rg	N1r	N1r	S3rt	N1r	N1r	S3r	N1r	S3r	N1r	N1r	N1r	S3r	S3rg	S3rg	S3rg	S3rg	S3rg	N1r	S3r	S3rg	S3rg	N1r	N1r	
Bheeman ahalli	49	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO	RO
Bheeman ahalli	51	N1r	S2r	S3r	S2rt	S3r	S2rt	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	S3r	S3r	S3r	S2rt	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Bheeman ahalli	52	N1r	S2r	S3r	S2rt	S3r	S2rt	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	S3r	S3r	S3r	S2rt	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Bheeman ahalli	53	N1r	S2r	S3r	S2rt	S3r	S2rt	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	S3r	S3r	S3r	S2rt	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Bheeman ahalli	54	N1r	S3rg	N1r	S3rg	N1r	S3rg	N1r	N1r	S3rt	N1r	N1r	S3r	N1r	S3r	N1r	N1r	N1r	S3r	S3rg	S3rg	S3rg	S3rg	S3rg	N1r	S3r	S3rg	S3rg	N1r	N1r	
Bheeman ahalli	55	N1r	S2r	S3r	S2rt	S3r	S2rt	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	S3r	S3r	S3r	S2rt	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Bheeman ahalli	56	N1r	S2r	S3r	S2rt	S3r	S2rt	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	S3r	S3r	S3r	S2rt	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Bheeman ahalli	57	N1r	S3rg	N1r	S3rg	N1r	S3rg	N1r	N1r	S3rt	N1r	N1r	S3r	N1r	S3r	N1r	N1r	N1r	S3r	S3rg	S3rg	S3rg	S3rg	S3rg	N1r	S3r	S3rg	S3rg	N1r	N1r	
Bheeman ahalli	58	N1r	S3rg	N1r	S3rg	N1r	S3rg	N1r	N1r	S3rt	N1r	N1r	S3r	N1r	S3r	N1r	N1r	N1r	S3r	S3rg	S3rg	S3rg	S3rg	S3rg	N1r	S3r	S3rg	S3rg	N1r	N1r	
Bheeman ahalli	59	N1r	S3rg	N1r	S3rg	N1r	S3rg	N1r	N1r	S3rt	N1r	N1r	S3r	N1r	S3r	N1r	N1r	N1r	S3r	S3rg	S3rg	S3rg	S3rg	S3rg	N1r	S3r	S3rg	S3rg	N1r	N1r	
Bheeman ahalli	60	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz	
Bheeman ahalli	61	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz	
Bheeman ahalli	62	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Bheeman ahalli	63	N1r	S2r	S3r	S2rt	S3r	S2rt	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	S3r	S3r	S3r	S2rt	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Bheeman ahalli	64	N1r	S2r	S3r	S2rt	S3r	S2rt	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	S3r	S3r	S3r	S2rt	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brimjal	Bhendi	Drumstick	Mulberry
Bheeman ahalli	65	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r
Bheeman ahalli	67	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r
Bheeman ahalli	68	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r
Bheeman ahalli	70	N1n	S3n	N1n	S3n	N1n	S3n	N1n	N1n	S3n	N1n	S3n	N1n	N1n	N1n	N1tn	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n
Bheeman ahalli	71	S3n	S2n	S3n	S2tn	N1n	S3tn	N1n	S3n	S3t	S3n	S2tn	N1n	N1n	S3n	N1n	N1n	S3n	S3n	N1n	S3n	S3n	S3n	S3n	S3n	S2n	S3n	S3n	N1n	N1n
Bheeman ahalli	73	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r
Bheeman ahalli	74	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r
Bheeman ahalli	75	S3n	S2n	S3n	S2tn	N1n	S3tn	N1n	S3n	S3t	S3n	S2tn	N1n	N1n	S3n	N1n	N1n	S3n	S3n	N1n	S3n	S3n	S3n	S3n	S3n	S2n	S3n	S3n	N1n	N1n
Bheeman ahalli	76	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz
Bheeman ahalli	77	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz
Bheeman ahalli	78	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz
Bheeman ahalli	79	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz
Bheeman ahalli	92	S3n	S2n	S3n	S2tn	N1n	S3tn	N1n	S3n	S3t	S3n	S2tn	N1n	N1n	S3n	N1n	N1n	S3n	S3n	N1n	S3n	S3n	S3n	S3n	S3n	S2n	S3n	S3n	N1n	N1n
Bheeman ahalli	275	N1r	S3r	N1r	S3r	N1r	S3r	N1r	N1r	S3rt	N1r	N1r	S3r	N1r	S3r	N1r	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r
Bheeman ahalli	276	N1r	S3r	N1r	S3r	N1r	S3r	N1r	N1r	S3rt	N1r	N1r	S3r	N1r	S3r	N1r	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r
Bheeman ahalli	277	N1r	S3r	N1r	S3r	N1r	S3r	N1r	N1r	S3rt	N1r	N1r	S3r	N1r	S3r	N1r	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r
Bheeman ahalli	278	N1r	S3r	N1r	S3r	N1r	S3r	N1r	N1r	S3rt	N1r	N1r	S3r	N1r	S3r	N1r	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r
Bheeman ahalli	279	N1r	S3r	N1r	S3r	N1r	S3r	N1r	N1r	S3rt	N1r	N1r	S3r	N1r	S3r	N1r	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r
Bheeman ahalli	281	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz
Bheeman ahalli	282	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz
Handara ki	249(10)	N1r	S2r	S3r	S2rt	S3r	S3t	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	N1n	S3r	S3r	S2r	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brimjal	Bhendi	Drumstick	Mulberry	
Handara ki	247	N1r	S2r	S3r	S2rt	S3r	S3t	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	N1n	S3r	S3r	S2r	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Handara ki	248	N1r	S2r	S3r	S2rt	S3r	S3t	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	N1n	S3r	S3r	S2r	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Handara ki	249/1	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	
Handara ki	249/4	N1r	S2r	S3r	S2rt	S3r	S3t	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	N1n	S3r	S3r	S2r	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Handara ki	249/5	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	
Handara ki	249/7	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st
Handara ki	249/8	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st
Handara ki	249/9	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st
Handara ki	250	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st
Handara ki	251	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st
Handara ki	252	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st
Handara ki	253	S3n	S2n	S3n	S2tn	N1n	S3tn	N1n	S3n	S3t	S3n	S2tn	N1n	N1n	S3n	N1n	N1n	S3n	S3n	N1n	S3n	S3n	S3n	S3n	S3n	S3n	S2n	S3n	S3n	N1n	N1n
Handara ki	254	S3n	S2n	S3n	S2tn	N1n	S3tn	N1n	S3n	S3t	S3n	S2tn	N1n	N1n	S3n	N1n	N1n	S3n	S3n	N1n	S3n	S3n	S3n	S3n	S3n	S3n	S2n	S3n	S3n	N1n	N1n
Handara ki	255	S3n	S2n	S3n	S2tn	N1n	S3tn	N1n	S3n	S3t	S3n	S2tn	N1n	N1n	S3n	N1n	N1n	S3n	S3n	N1n	S3n	S3n	S3n	S3n	S3n	S3n	S2n	S3n	S3n	N1n	N1n
Handara ki	256	S3n	S2n	S3n	S2tn	N1n	S3tn	N1n	S3n	S3t	S3n	S2tn	N1n	N1n	S3n	N1n	N1n	S3n	S3n	N1n	S3n	S3n	S3n	S3n	S3n	S3n	S2n	S3n	S3n	N1n	N1n
Handara ki	257	S3n	S2n	S3n	S2tn	N1n	S3tn	N1n	S3n	S3t	S3n	S2tn	N1n	N1n	S3n	N1n	N1n	S3n	S3n	N1n	S3n	S3n	S3n	S3n	S3n	S3n	S2n	S3n	S3n	N1n	N1n
Handara ki	258	S3n	S2n	S3n	S2tn	N1n	S3tn	N1n	S3n	S3t	S3n	S2tn	N1n	N1n	S3n	N1n	N1n	S3n	S3n	N1n	S3n	S3n	S3n	S3n	S3n	S3n	S2n	S3n	S3n	N1n	N1n
Handara ki	259	S3n	S2n	S3n	S2tn	N1n	S3tn	N1n	S3n	S3t	S3n	S2tn	N1n	N1n	S3n	N1n	N1n	S3n	S3n	N1n	S3n	S3n	S3n	S3n	S3n	S3n	S2n	S3n	S3n	N1n	N1n
Handara ki	260	S3n	S2n	S3n	S2tn	N1n	S3tn	N1n	S3n	S3t	S3n	S2tn	N1n	N1n	S3n	N1n	N1n	S3n	S3n	N1n	S3n	S3n	S3n	S3n	S3n	S3n	S2n	S3n	S3n	N1n	N1n
Handara ki	261	S3n	S2n	S3n	S2tn	N1n	S3tn	N1n	S3n	S3t	S3n	S2tn	N1n	N1n	S3n	N1n	N1n	S3n	S3n	N1n	S3n	S3n	S3n	S3n	S3n	S3n	S2n	S3n	S3n	N1n	N1n
Handara ki	343	S3n	S2n	S3n	S2tn	N1n	S3tn	N1n	S3n	S3t	S3n	S2tn	N1n	N1n	S3n	N1n	N1n	S3n	S3n	N1n	S3n	S3n	S3n	S3n	S3n	S3n	S2n	S3n	S3n	N1n	N1n

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brimjal	Bhendi	Drumstick	Mulberry	
Handaraki	344	S3n	S2n	S3n	S2tn	N1n	S3tn	N1n	S3n	S3t	S3n	S2tn	N1n	N1n	S3n	N1n	N1n	S3n	S3n	N1n	S3n	S3n	S3n	S3n	S3n	S2n	S3n	S3n	N1n	N1n	
Handaraki	345	S3n	S2n	S3n	S2tn	N1n	S3tn	N1n	S3n	S3t	S3n	S2tn	N1n	N1n	S3n	N1n	N1n	S3n	S3n	N1n	S3n	S3n	S3n	S3n	S3n	S2n	S3n	S3n	N1n	N1n	
Handaraki	346	S3n	S2n	S3n	S2tn	N1n	S3tn	N1n	S3n	S3t	S3n	S2tn	N1n	N1n	S3n	N1n	N1n	S3n	S3n	N1n	S3n	S3n	S3n	S3n	S3n	S2n	S3n	S3n	N1n	N1n	
Handaraki	347	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0
Handaraki	348	Quar	Quar	Quar	Quar	Quar	Quar	Quar	Quar	Quar	Quar	Quar	Quar	Quar	Quar	Quar	Quar	Quar	Quar	Quar	Quar	Quar	Quar	Quar	Quar	Quar	Quar	Quar	Quar	Quar	
Handaraki	349	N1r	S2r	S3r	S2rt	S3r	S2rt	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	S3r	S3r	S3r	S2rt	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Handaraki	350	N1r	S2r	S3r	S2rt	S3r	S2rt	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	S3r	S3r	S3r	S2rt	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Handaraki	351	N1r	S2r	S3r	S2rt	S3r	S2rt	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	S3r	S3r	S3r	S2rt	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Handaraki	352	N1r	S2r	S3r	S2rt	S3r	S2rt	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	S3r	S3r	S3r	S2rt	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Handaraki	353	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0
Handaraki	354	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0	R0
Handaraki	360	N1r	S2r	S3r	S2rt	S3r	S2rt	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	S3r	S3r	S3r	S2rt	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Handaraki	370	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Handaraki	371	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Handaraki	419	N1r	S2r	S3r	S2rt	S3r	S2rt	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	S3r	S3r	S3r	S2rt	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Handaraki	420	N1r	S2r	S3r	S2rt	S3r	S2rt	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	S3r	S3r	S3r	S2rt	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Handaraki	421	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz	
Handaraki	422	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz	
Ramathirtha	126	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Ramathirtha	127	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Ramathirtha	128	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3t	S3tz	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz	

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brimjal	Bhendi	Drumstick	Mulberry	
Ramathir tha	134	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3t	S3tz	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz	
Ramathir tha	135	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Ramathir tha	136	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Ramathir tha	178	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Ramathir tha	180	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3t	S3tz	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz	
Ramathir tha	181	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Ramathir tha	182	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Ramathir tha	183	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Ramathir tha	184	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3t	S3tz	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz	
Ramathir tha	185	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3t	S3tz	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz	
Ramathir tha	186	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Ramathir tha	187	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r

RO-Rock outcrops

PART-B

SOCIO-ECONOMIC STATUS OF FARM HOUSEHOLDS

CONTENTS

1.	Findings of the socio-economic survey	1-2
2.	Introduction	3
3	Methodology	5-6
4	Salient features of the survey	7-23
5	Summary	25-28

LIST OF TABLES

1	Households sampled for socio economic survey	7
2	Population characteristics	7
3	Age wise classification of household members	7
4	Education level of household members	8
5	Occupation of household heads	8
6	Occupation of family members	9
7	Institutional participation of household members	9
8	Type of house owned by households	9
9	Durable assets owned by households	10
10	Average value of durable assets owned by households	10
11	Farm implements owned by households	10
12	Average value of farm implements	11
13	Livestock possession by households	11
14	Average labour availability	11
15	Adequacy of hired labour	11
16	Distribution of land (ha)	12
17	Average land value (Rs./ha)	12
18	Cropping pattern	12
19	Cropping intensity	12
20	Possession of bank account and saving	13
21	Borrowing status	13
22	Source of credit	13
23	Avg. credit borrowed	13
24	Purpose of credit borrowed from institutional sources	14
25	Repayment status of household from institutional sources	14
26	Opinion on institutional sources of credit	14
27.,a	Cost of cultivation of Red gram	15
27.b	Cost of cultivation of Maize	16
27.c	Cost of cultivation of Cotton	17
28	Adequacy of fodder	18

29	Annual gross income	18
30	Average annual expenditure	18
31	Horticultural species grown	18
32	Forest species grown	19
33	Average additional investment capacity	19
34	Source of funds for additional investment	19
35	Marketing of the agricultural produce	19
36	Marketing channels used for sale of agricultural produce	20
37	Mode of transport of agricultural produce	20
38	Incidence of soil and water erosion problems	20
39	Interest shown towards soil testing	20
40	Usage pattern of fuel for domestic use	21
41	Source of drinking water	21
42	Source of light	21
43	Existence of sanitary toilet facility	21
44	Possession of public distribution system (PDS) card	22
45	Participation in NREGA programme	22
46	Adequacy of food items	22
47	Inadequacy of food items	22
48	Farming constraints experienced	28

FINDINGS OF THE SOCIO-ECONOMIC SURVEY

- ❖ *The survey was conducted in Bhimanahalli-2 is located at North latitude 16° 58' 39.868" and 16° 56' 44.222" and East longitude 77° 13' 58.007" and 77° 11' 56.808" covering an area of about 561.68 ha coming under Bhimanahalli and Handarki villages of Yadagiri taluk.*
- ❖ *Socio-economic analysis of Bhimanahalli-2 micro watersheds of Motanahalli sub-watershed, Chitapura taluk & Kalaburagi District indicated that, out of the total sample of 36 total respondents, 8 (22.22 %) were marginal, 13 (36.11%) were small and 13 (36.11 %) were Semi medium*
- ❖ *The population characteristics of households indicated that, there were 81 (59.12%) men and 56 (40.88 %) were women.*
- ❖ *Majority of the respondents (43.07%) were in the age group of 16-35 years.*
- ❖ *Education level of the sample households indicated that, there were 30.66 per cent illiterates, 63.51 per cent pre university education and 6.57 per cent attained graduation.*
- ❖ *About, 72.22 per cent of household heads practicing agriculture and 16.67 per cent of the household heads were engaged as agricultural labourers.*
- ❖ *Agriculture was the major occupation for 48.91 per cent of the household members.*
- ❖ *In the study area, 94.44 per cent of the households possess katcha house and 5.56 per cent possess pucca house.*
- ❖ *The durable assets owned by the households showed that, 83.33 per cent possess TV, 66.67 per cent possess mixer grinder, 108.33 per cent possess mobile phones and 33.33 per cent possess motor cycles.*
- ❖ *Farm implements owned by the households indicated that, 22.22 per cent of the households possess plough, 5.56 per cent possess bullock cart and 8.33 per cent possess sprayer.*
- ❖ *The average labour availability in the study area showed that, own labour men available in the micro watershed was 1.56, women available in the micro watershed was 1.29, hired labour (men) available was 8 and hired labour (women) available was 13.24.*
- ❖ *Out of the total land holding of the sample respondents 100.00 per cent (59.06 ha) of the area is under dry condition.*
- ❖ *The major crops grown by sample farmers are Red gram, Maize and Cotton and cropping intensity was recorded as 99.97 per cent.*
- ❖ *Out of the sample households 97.22 percent possessed bank account and 63.89 per cent of them have savings in the account.*
- ❖ *About 44.44 per cent of the respondents borrowed credit from various sources.*
- ❖ *Among the credit borrowed by households, 20.00 per cent have borrowed loan from commercial banks and 68.00 per cent from co-operative/Grameena bank.*
- ❖ *Majority of the respondents (100.00%) have borrowed loan for agriculture purpose.*

- ❖ *Regarding the opinion on institutional sources of credit, 96.15 per cent of the households opined that credit helped to perform timely agricultural operations,*
- ❖ *The per hectare cost of cultivation for Red gram, Maize, Cotton, was Rs.31594.21 , 20566.68 and 20292.15 with benefit cost ratio of 1:1.30, 1: 3.50 and 1: 1.50 respectively.*
- ❖ *The average annual gross income of the farmers was Rs. 111534.72 in micro-watershed, of which Rs. 65118.06 comes from agriculture.*
- ❖ *Sampled households have grown 5 horticulture trees and 118 forestry trees together in the fields and back yards.*
- ❖ *About 75.00 per cent of the households shown interest to cultivate horticultural crops.*
- ❖ *Households have an average investment capacity of Rs. 1083.33 for land development.*
- ❖ *Source of funds for additional investment is concerned, 27.78 per cent depends on bank loan for land development activities.*
- ❖ *Regarding marketing channels, 94.44 per cent of the households have sold agricultural produce to the local/village merchants, while, 8.33 per cent have sold in regulated markets.*
- ❖ *Further, 13.89 per cent of the households have used tractor for the transport of agriculture commodity.*
- ❖ *Majority of the farmers (30.56%) have experienced soil and water erosion problems in the watershed and 105.56 per cent of the households were interested towards soil testing.*
- ❖ *Fire was the major source of fuel for domestic use for 75.00 per cent of the households and 63.89 per cent households has LPG connection.*
- ❖ *Piped supply was the major source for drinking water for 100.00 per cent of the households.*
- ❖ *Electricity was the major source of light for 100.00 per cent of the households.*
- ❖ *In the study area, 69.44 per cent of the households possess toilet facility.*
- ❖ *Regarding possession of PDS card, 94.44 per cent of the households possessed BPL card, 2.78 per cent of the household's possessed APL card and 2.78 per cent of the household's were not having ration cards.*
- ❖ *Households opined that, the requirement of cereals (100.00%), pulses (10.00%) and oilseeds (52.78%) are adequate for consumption.*
- ❖ *Farming constraints experienced by households in the micro watersheds were lower fertility status of the soil (100.00%) wild animal menace on farm field (94.44%), frequent incidence of pest and diseases (91.67%), inadequacy of irrigation water (100.00%), high cost of fertilizers and plant protection chemicals (100.00%), high rate of interest on credit (100.00%), low price for the agricultural commodities (100.00%), lack of marketing facilities in the area (101.00%), inadequate extension services (97.22%), lack of transport for safe transport of the agricultural produce to the market (94.44%),*

INTRODUCTION

Soil and water are the two precious natural resources which are essential for crop production and existence of life on earth. Rainfed agriculture is under severe stress due to various constraints related to agriculture like uneven and erratic distribution of rainfall, indiscriminate use of fertilizers, chemicals and pesticides, adoption of improper land management practices, soil erosion, decline in soil fertility, decline in ground water resources leading to low crop productivity. The area under rainfed agriculture has to be managed effectively using the best available practices to enhance the production of food, fodder and fuel. This is possible if the land resources are characterized at each parcel of land through detailed land resource inventory using the best available techniques of remote sensing, GPS and GIS. The watershed development programs are aimed at the sustainable distribution of its resources and the process of creating and implementing plans, programs, and projects to sustain and enhance watershed functions that affect the plant, animal and human communities within a watershed boundary.

World Bank funded KWDP II, SUJALA III project was implemented in with Broad objective of demonstrating more effective watershed management through greater integration of programmes related to rain-fed agriculture, innovative and science based approaches and strengthen institutional capacities and If successful, it is expected that the systems and tools could be mainstreamed into the overall IWMP in the State of Karnataka and in time, throughout other IWMP operations in India. With this background the socio-economic survey has been carried out with following specific objectives:

1. To understand the demographic features of the households in the micro-watershed
2. To understand the extent of family labour available and additional employment opportunities available within the village.
3. To know the status of assets of households in the micro-watershed for suggesting possible improvements.
4. To study the cropping pattern, cropped area and productivity levels of different households in micro-watershed.
5. To determine the type and extent of livestock owned by different categories of HHs
6. Availability of fodder and level of livestock management.

Scope and importance of survey

Survey helps in identification of different socio-economic and resource use-patterns of farmers at the Micro watershed. Household survey provides demographic features, labor force, and levels of education; land ownership and asset position (including livestock and other household assets) of surveyed households; and cropping patterns, input intensities, and average crop yields from farmers' fields. It also discusses crop utilization and the degree of commercialization of production in the areas; farmers' access to and utilization of credit from formal and informal sources; and the level of adoption and use of soil, water, and pest management technologies.

METHODOLOGY

The description of the methods, components selected for the survey and procedures followed in conducting the baseline survey are furnished under the following heads.

1. Description of the study area

Kalaburagi district is one of the three districts that were transferred from Hyderabad State to Karnataka state at the time of re-organization of the state in 1956. The district is one among the 30 districts of Karnataka State. It is located in the Northern part of the state and lies between 76°.04' and 77°.42 east longitude, and 17°.12' and 17°.46' north latitude, covering an area of 10,951 km². It is bounded on the west by Bijapur district of Karnataka and Sholapur district of Maharashtra, on the west by RangaReddy and Medak district of Telegana State, on the north by Bidar district and Osmanabad district of Maharashtra and on the south by Yadgir district of Karnataka. Kalaburagi is famous for toordal Pigeon pea and the limestone deposits are more in Kalaburagi District.

The District was under the rule of Nijam s of Hyderabad before independence. The district has a rich background of knowledge and culture. The existence of university at Nagai in Chitapur, Vignaneeshwaras Mitakshara, Nrupatungas Kavirajmarg and the religious and social revolution led by Shivsharanas and the Sufi saint Banda Nawaz are all evidence of it. However, due to erratic rainfall and continuous occurrence of droughts in the 19th century the life of the people was never smooth and secure. Further during the Nizams period, the district could not develop due to the negligence and inefficient administration.

Kalaburagi is situated in Deccan Plateau located at 17.33°N 76.83°E and the general elevation ranges from 300 to 750 meters above mean sea level. Two main rivers, Krishna and Bhima, flow in the district. Black soil is predominant soil type in the district. The district has a large number of tanks which, in addition to the rivers, irrigate the land. The Upper Krishna Project is major irrigation venture in the district. Bajra, toor, sugarcane, groundnut, sunflower, sesame, castor bean, black gram, jowar, wheat, cotton, ragi, Bengal gram, and linseed are grown in this district.

According to the 2011 census Kalaburagi district has a population of 2,564,892. The district has a population density of 233 inhabitants per square kilometre (600/sq mi). Kalaburagi has a sex ratio of 962 females for every 1000 males, and a literacy rate of 65.65%.

2. Locale of the survey and description of the micro-watershed and

The study was conducted in Bhimanahalli-2 micro-watershed (Motanahalli sub-watershed, Chitapura taluk & Kalaburagi District) is located North latitude 16⁰ 58' 39.868" and 16⁰ 56' 44.222" and East longitude 77⁰ 13' 58.007" and 77⁰ 11' 56.808" covering an area of about 561.68 ha bounded by under Bhimanahalli and Handarki Villages.

3. Selection of the respondents for the study

The micro-watershed is marked with 320 square meters grids. One farmer from every alternate grid in the micro-watershed was selected for the study and interviewed for socio-economic data. Totally 36 households were interviewed for the survey.

4. The parameters considered for socio-economic survey of households

Two forms of data were collected from the micro-watershed which includes primary data from the farm households and secondary data about the villages under the micro-watershed jurisdiction.

The following parameters were considered for the primary data collection about the socio-economic data of the households, (1) Demographic information, (2) Farm and durable assets owned by households, (3) Livestock possession, (4) Labour availability, (5) Level of migration in the village, Land holding, (7) Cropping pattern, (8) Source of irrigation, (9) Borrowing status, (10) Cost of cultivation of major crops, (11) Economics of subsidiary activities, (12) Fodder availability, (13) Family annual income from different sources, (14) Horticulture and forestry species grown, (15) Additional investment capacity, (16) Marketing practices, (17) Status of soil and water conservation structure, (18) Access to basic needs and (19) Constraints and suggestion.

The following parameters were considered for the secondary data regarding the villages under the micro-watershed jurisdiction, (1) Number of villages in each micro-watershed jurisdiction, (2) Village wise number of households, (3) Geographical area of the villages, (4) Cultivable area including rainfed and irrigated, (5) Number and type of house in each village, (6) Human and livestock population, (7) Facilities in the village such as roads, transport facility for conveyance, drinking water supply, street light and (8) Community based organizations in the villages.

5. Development of interview schedule and data collection

Taking into the consideration the objectives of the survey, an interview schedule was prepared after thorough consultation with the experts in the field of social sciences. A comprehensive interview schedule covering all the major parameters for measuring the socio-economic situation was developed.

6. Tools used to analyze the data

The statistical components such as frequency and percentage were used to analyze the data.

Abbreviations used in the report

LL=Landless

MF=Marginal Farmers

SF=Small farmers

SMF=Semi medium farmers

MDF=Medium farmers

LF=Large Farmers

FINDINGS OF THE SURVEY

This chapter deals with systematic presentation of results of the survey. Keeping in view the objectives, the salient features of the survey are presented under the following headings.

Households sampled for socio-economic survey: The data on households sampled for socio economic survey in Bhimanahalli-2 Micro watershed is presented in Table 1 and it indicated that 36 farmers were sampled in Bhimanahalli-2 micro-watershed among households surveyed 8 (22.22%) were marginal, 13 (36.11%) were small and 13 (36.11 %) were semi medium. 2 landless farmers were also interviewed for the survey.

Table 1. Households sampled for socio economic survey in Bhimanahalli-2 micro-watershed

Sl.No.	Particulars	LL (2)		MF (8)		SF (13)		SMF (13)		All (36)	
		N	%	N	%	N	%	N	%	N	%
1	Farmers	2	5.56	8	22.2	13	36.1	13	36.1	36	100

Population characteristics: The population characteristics of households sampled for socio-economic survey in Bhimanahalli-2 Micro watershed is presented in Table 2. The data indicated that, there were 81 (59.12%) men and 56 (40.88%) were women.

Table 2. Population characteristics in Bhimanahalli-2 micro-watershed

Sl.No.	Particulars	LL (9)		MF (33)		SF (52)		SMF (43)		All (137)	
		N	%	N	%	N	%	N	%	N	%
1	Men	5	55.6	21	64	33	63	22	51.2	81	59.1
2	Women	4	44.4	12	36	19	37	21	48.8	56	40.9
Total		9	100	33	100	52	100	43	100	137	100
Average		4.5		4.1		4.0		3.3		3.8	

Age wise classification of population: The age wise classification of household members in Bhimanahalli-2 Micro watershed is presented in Table 3. The indicated that, 21 (15.33%) of population were 0-15 years of age, 59 (43.07%) were 16-35 years of age, 46(33.58%) were 36-60 years of age and 11 (8.03 %) were above 61 years of age.

Table 3: Age wise classification of members of the household in Bhimanahalli-2 micro-watershed

Sl.No.	Particulars	LL (9)		MF (33)		SF (52)		SMF (43)		All (137)	
		N	%	N	%	N	%	N	%	N	%
1	0-15 years of age	2	22.2	5	15.2	9	17.3	5	11.63	21	15.33
2	16-35 years of age	5	55.6	14	42.4	23	44.2	17	39.53	59	43.07
3	36-60 years of age	2	22.2	12	36.4	16	30.8	16	37.21	46	33.58
4	> 61 years	0	0	2	6.06	4	7.69	5	11.63	11	8.03
Total		9	100	33	100	52	100	43	100	137	100

Education level of household members: Education level of household members in Bhimanahalli-2 Micro watershed is presented in Table 4. The results indicated that, there were 30.66 per cent of illiterates, 29.20 per cent of them had primary school education, 3.65 per cent middle school education and masters education, 13.14 per cent high school education, 8.76 per cent of them had PUC education, 1.46 per cent of them had Diploma, 6.57 per cent attained graduation, and 2.92 them had other education.

Table 4. Education level of members of the household in Bhimanahalli-2 micro-watershed

Sl.No.	Particulars	LL (9)		MF (33)		SF (52)		SMF (43)		All (137)	
		N	%	N	%	N	%	N	%	N	%
1	Illiterate	4	44.4	13	39.4	11	21.2	14	32.6	42	30.7
2	Primary School	3	33.3	6	18.2	20	38.5	11	25.6	40	29.2
3	Middle School	1	11.1	1	3.03	1	1.92	2	4.65	5	3.65
4	High School	1	11.1	4	12.1	9	17.3	4	9.3	18	13.1
5	PUC	0	0	5	15.2	6	11.5	1	2.33	12	8.76
6	Diploma	0	0	0	0	0	0	2	4.65	2	1.46
7	Degree	0	0	1	3.03	3	5.77	5	11.6	9	6.57
8	Masters	0	0	0	0	1	1.92	4	9.3	5	3.65
9	Others	0	0	3	9.09	1	1.92	0	0	4	2.92
Total		9	100	33	100	52	100	43	100	137	100

Occupation of head of households: The data regarding the occupation of the household heads in Bhimanahalli-2 Micro watershed is presented in Table 5. The results indicate that, 72.22 per cent of household's heads were practicing agriculture, 16.67 per cent of the household heads were agricultural Labour, and 5.56 per cent of the household's heads were private services and student (2.78%).

Table 5: Occupation of heads of households in Bhimanahalli-2 micro-watershed

Sl.No.	Particulars	LL (2)		MF (8)		SF (13)		SMF (13)		All (36)	
		N	%	N	%	N	%	N	%	N	%
1	Agriculture	0	0	7	88	7	53.85	12	92	26	72.22
2	Agricultural Labour	2	100	0	0	4	30.77	0	0	6	16.67
3	Private Service	0	0	0	0	0	0	2	15	2	5.56
4	Student	0	0	0	0	1	7.69	0	0	1	2.78
Total		2	100	8	100	13	100	14	100	37	100

Occupation of the members of the household: The data regarding the occupation of the household members in Bhimanahalli-2 Micro watershed is presented in Table 6. The results indicate that, agriculture was the major occupation for 48.91 per cent of the household members, 10.22 per cent were agricultural labour, 5.84 per cent were general labour, 10.2 per cent were working in private services, 17.52 per cent were working in pursuing education, 2.92 per cent were involved as housewife and children's.

Table 6: Occupation of members of the household in Bhimanahalli-2 micro-watershed

Sl.No.	Particulars	LL (9)		MF (33)		SF (52)		SMF (43)		All (137)	
		N	%	N	%	N	%	N	%	N	%
1	Agriculture	0	0	16	48.5	25	48.08	26	60.47	67	48.9
2	Agricultural Labour	4	44.4	2	6.06	8	15.38	0	0	14	10.2
3	General Labour	2	22.2	6	18.2	0	0	0	0	8	5.84
4	Private Service	0	0	0	0	6	11.54	8	18.6	14	10.2
5	Student	3	33.3	5	15.2	10	19.23	6	13.95	24	17.5
6	Others	0	0	1	3.03	1	1.92	0	0	2	1.46
7	Housewife	0	0	0	0	1	1.92	3	6.98	4	2.92
8	Children	0	0	3	9.09	1	1.92	0	0	4	2.92
9	Retired	0	0	0	0	0	0	0	0	0	0
Total		9	100	33	100	52	100	43	100	137	100

Institutional Participation of household members: The data regarding the institutional participation of the household members in Bhimanahalli-2 Micro watershed is presented in Table 7. The results show that, out of the total family members in the households 0.73 per cent of them are participating in diary cooperative, 6.57 per cent of them were participating in self help group and 92.7 per cent were not participating in any of the institutions.

Table 7: Institutional Participation of household member in Bhimanahalli-2 micro-watershed

Sl.No.	Particulars	LL (9)		MF (33)		SF (52)		SMF (43)		All (137)	
		N	%	N	%	N	%	N	%	N	%
1	Self Help Group	1	11	2	6.06	3	5.77	3	6.98	9	6.57
2	Dairy Cooperative	0	0	0	0	0	0	1	2.33	1	0.73
3	No Participation	8	89	31	93.9	49	94.2	39	90.7	127	92.7
Total		9	100	33	100	52	100	43	100	137	100

Type of house owned: The data regarding the type of house owned by the households in Bhimanahalli-2 Micro watershed is presented in Table 8. The results indicate that, 16.67 percent possess thatched house, 94.44 per cent of the households possess katcha house and 5.56 per cent possess pucca house.

Table 8. Type of house owned by households in Bhimanahalli-2 micro-watershed

Sl.No.	Particulars	LL (2)		MF (8)		SF (13)		SMF (13)		All (36)	
		N	%	N	%	N	%	N	%	N	%
1	Thatched	0	0	2	25	4	30.77	0	0	6	16.67
2	Katcha	2	100	10	125	9	69.23	13	100	34	94.44
3	Pucca/RCC	0	0	0	0	0	0	2	15.4	2	5.56
Total		2	100	12	100	13	100	15	100	42	100

Durable assets owned by the households: The data regarding the Durable Assets owned by the households in Bhimanahalli-2 Micro watershed is presented in Table 9. The results

shows that, 83.33 per cent possess TV, 66.67 per cent possess mixer grinder, 63.89 per cent possess Bicycle, 33.33 per cent possess motor cycle and 108.33 per cent possess mobile phones.

Table 9. Durable assets owned by households in Bhimanahalli-2 micro-watershed

Sl.No.	Particulars	LL (2)		MF (8)		SF (13)		SMF (13)		All (36)	
		N	%	N	%	N	%	N	%	N	%
1	Television	2	100	6	75	8	61.5	14	108	30	83.33
2	Mixer/Grinder	1	50	6	75	7	53.9	10	77	24	66.67
3	Bicycle	1	50	6	75	7	53.9	9	69	23	63.89
4	Motor Cycle	0	0	1	13	4	30.8	7	54	12	33.33
5	Mobile Phone	2	100	11	138	12	92.3	14	108	39	108.33

Average value of durable assets: The data regarding the average value of durable assets owned by the households in Bhimanahalli-2 Micro watershed is presented in Table 10. The result shows that, the average value of television was Rs.2306.00, mixer grinder was Rs.1000.00, bicycle was Rs.1000.00, motor cycle was Rs. 31666.00 and mobile phone was Rs.927.00.

Table 10. Average value of durable assets owned in Bhimanahalli-2 micro-watershed
Average Value (Rs.)

Sl.No.	Particulars	LL (2)	MF (8)	SF (13)	SMF (13)	All (36)
1	Television	1100	2000	2375	2571	2306
2	Mixer/Grinder	1000	1000	1000	1000	1000
3	Bicycle	1000	1000	1000	1000	1000
4	Motor Cycle	0	30000	30000	32857	31666
5	Mobile Phone	1000	1083	981	769	927

Farm implements owned: The data regarding the farm implements owned by the households in Bhimanahalli-2 Micro watershed is presented in Table 11. About 5.56 per cent of the households possess Bullock Cart, 22.22 per cent possess plough, 8.33 per cent possess Sprayer, 86.11 per cent possess Weeder and 16.67 per cent possess chaff cutter.

Table 11. Farm implements owned in Bhimanahalli-2 micro-watershed

Sl.No.	Particulars	LL (2)		MF (8)		SF (13)		SMF (13)		All (36)	
		N	%	N	%	N	%	N	%	N	%
1	Bullock Cart	0	0	1	12.5	1	7.69	0	0	2	5.56
2	Plough	0	0	3	37.5	3	23.08	2	15.4	8	22.22
3	Sprayer	0	0	1	12.5	1	7.69	1	7.69	3	8.33
4	Weeder	2	100	6	75	7	53.85	16	123	31	86.11
5	Chaff Cutter	0	0	1	12.5	3	23.08	2	15.4	6	16.67
6	Blank	0	0	1	12.5	6	46.15	2	15.4	9	25

Average value of farm implements: The data regarding the average value of farm Implements owned by the households in Bhimanahalli-2 Micro watershed is presented in

Table 12. The results show that the average value of plough was Rs.568.00, bullock Cart was Rs.20000.00, sprayer was Rs.4000.00, weeder was Rs.26.00 and chaff cutter was Rs.3233.

Table 12. Average value of farm implements in Bhimanahalli-2 micro-watershed

Sl.No.	Particulars	Average Value (Rs.)				
		LL (2)	MF (8)	SF (13)	SMF (13)	All (36)
1	Bullock Cart	0	20000	20000	0	20000
2	Plough	0	722	500	500	568
3	Sprayer	0	4000	4000	4000	4000
4	Weeder	25	26	25	27	26
5	Chaff Cutter	0	4000	2466	4000	3233

Livestock possession by the households: The data regarding the Livestock possession by the households in Bhimanahalli-2 Micro watershed is presented in Table 13. The results indicate that, 25.00 per cent of the households possess bullocks, 13.89 per cent possess local cow, 2.78 per cent possess crossbred cow and sheep.

Table 13. Livestock possession by households in Bhimanahalli-2 micro-watershed

Sl.No.	Particulars	LL (2)		MF (8)		SF (13)		SMF (13)		All (36)	
		N	%	N	%	N	%	N	%	N	%
1	Bullock	0	0	3	38	4	30.77	2	15	9	25
2	Local cow	0	0	1	13	3	23.08	1	7.7	5	13.89
3	Crossbred cow	0	0	0	0	0	0	1	7.7	1	2.78
4	Sheep	0	0	0	0	1	7.69	0	0	1	2.78
5	blank	2	100	5	63	8	61.54	13	100	28	77.78

Average Labour availability: The data regarding the average labour availability in Bhimanahalli-2 Micro watershed is presented in Table 14. The indicated that, own labour men available in the micro watershed was 1.56, women available in the micro watershed was 1.29, hired labour (men) available was 8 and hired labour (women) available was 13.24.

Table 14. Average labour availability in Bhimanahalli-2 micro-watershed

Sl.No.	Particulars	LL (2)	MF (8)	SF (13)	SMF (13)	All (36)
1	Hired labour Female	0	8.25	10.23	19.31	13.24
2	Own Labour Female	0	1.38	1.23	1.31	1.29
3	Own labour Male	0	1.75	1.69	1.31	1.56
4	Hired labour Male	0	4	5.54	12.92	8

Table 15. Adequacy of hired labour in Bhimanahalli-2 micro-watershed

Sl.No.	Particulars	LL (2)		MF (8)		SF (13)		SMF (13)		All (36)	
		N	%	N	%	N	%	N	%	N	%
1	Adequate	0	0	8	100	14	108	13	100	35	97.2

Adequacy of hired labour: The data regarding the adequacy of hired labour in Bhimanahalli-2 Micro watershed is presented in Table 15. The results indicate that, 97.22 per cent of the household opined that hired labour was adequate.

Distribution of land (ha): The data regarding the distribution of land (ha) in Bhimanahalli-2 Micro watershed is presented in Table 16. The results indicate that, 59.06 ha (100.00%) of dry land.

Table 16. Distribution of land (ha) in Bhimanahalli-2 micro-watershed

Sl.No.	Particulars	LL (2)		MF (8)		SF (13)		SMF (13)		All (36)	
		N	%	N	%	N	%	N	%	N	%
1	Dry	0	0	5.68	100	17.05	100	36.34	100	59.06	100
	Total	0	100	5.68	100	17.05	100	36.34	100	59.06	100

Average value of land (ha): The data regarding the average land value (Rs./ha) in Bhimanahalli-2 Micro watershed is presented in Table 17. The results show that the average value of dry land was Rs.255563.93.

Table 17. Average value of land (ha) in Bhimanahalli-2 micro-watershed

Sl.No.	Particulars	LL (2)	MF (8)	SF (13)	SMF (13)	All (36)
1	Dry	0	440128.3	340123.5	187058.7	255563.9

Cropping pattern: The data regarding the cropping pattern in Bhimanahalli-2 Micro watershed is presented in Table 18. The results indicate that, farmers have grown red gram (37.09 ha), Maize (16.87 ha), kharif red gram (14.30 ha), cotton (3.24 ha) and rabi cotton (2.43 ha).

Table 18. Cropping pattern in Bhimanahalli-2 micro-watershed

Sl.No.	Particulars	LL (2)	MF (8)	SF (13)	SMF (13)	All (36)
1	Rabi - Red gram	0	5.12	6.17	25.79	37.09
2	Rabi - Maize	0	0	2.5	14.37	16.87
3	Kharif - Red gram	0	0.56	8.36	5.38	14.3
4	Kharif - Cotton	0	0	0	3.24	3.24
5	Rabi - Cotton	0	0	0	2.43	2.43
	Total	0	5.68	17.03	51.21	73.93

Cropping intensity: The data regarding the cropping intensity in Bhimanahalli-2 Micro watershed is presented in Table 19. The results indicate that, the cropping intensity was 99.97 per cent.

Table 19. Cropping intensity (%) in Bhimanahalli-2 micro-watershed

Sl.No.	Particulars	LL (2)	MF (8)	SF (13)	SMF (13)	All (36)
1	Cropping Intensity	0	100	99.88	100	99.97

Possession of bank account and savings: The data regarding the possession of bank account and saving in Bhimanahalli-2 micro-watershed is presented in Table 20. The results indicate that, 97.22 cent of the households posses bank account and 63.89 per cent of them have savings.

Table 20. Possession of Bank account and savings in Bhimanahalli-2 micro-watershed

Sl.No.	Particulars	LL (2)		MF (8)		SF (13)		SMF (13)		All (36)	
		N	%	N	%	N	%	N	%	N	%
1	Account	2	100	8	100	13	100	12	92.31	35	97.22
2	Savings	0	0	6	75	7	53.85	10	76.92	23	63.89

Borrowing status: The data regarding the borrowing status in Bhimanahalli-2 micro-watershed is presented in Table 21. The results indicate that, 44.44 percent of the sample farmers have borrowed credit from different sources.

Table 21. Borrowing status in Bhimanahalli-2 micro-watershed

Sl.No.	Particulars	LL (2)		MF (8)		SF (13)		SMF (13)		All (36)	
		N	%	N	%	N	%	N	%	N	%
1	Credit Aailed	2	100	6	75	7	53.9	1	7.69	16	44.44

Source of credit: The data regarding the source of credit availed by households in Bhimanahalli-2 micro-watershed is presented in Table 22. The results show that, 20.00 per cent have borrowed loan from commercial banks, 12.00 per cent have borrowed loan from Cooperative bank and 68.00 per cent have borrowed loan from Grameena Bank.

Table 22. Source of credit borrowed by households in Bhimanahalli-2 micro-watershed

Sl.No.	Particulars	LL (2)		MF (6)		SF (7)		SMF (10)		All (25)	
		N	%	N	%	N	%	N	%	N	%
1	Commercial Bank	0	0	1	16.7	1	14.3	3	30	5	20
2	Cooperative Bank	0	0	1	16.7	1	14.3	1	10	3	12
3	Grameena Bank	0	0	3	50	6	85.7	8	80	17	68

Avg. Credit amount: The data regarding the avg. Credit amount in Bhimanahalli-2 micro-watershed is presented in Table 23. The results show that, farmers have borrowed Avg. Credit of Rs.130200.00 from different sources.

Table 23. Avg. Credit amount in Bhimanahalli-2 micro-watershed

Sl.No.	Particulars	LL (2)	MF (6)	SF (7)	SMF (10)	All (25)
1	Average Credit	0	17500	57142.9	275000	130200

Purpose of credit borrowed (institutional Source): The data regarding the purpose of credit borrowed - Institutional Credit in Bhimanahalli-2 micro-watershed is presented in Table 24. The results indicate that, 100.00 per cent of the households have borrowed loan for agriculture.

Table 24. Purpose of credit borrowed (institutional Source) by households in Bhimanahalli-2 micro-watershed

SN	Particulars	LL (0)		MF (5)		SF (8)		SMF (13)		All (26)	
		N	%	N	%	N	%	N	%	N	%
1	Agriculture production	0	0	5	100	8	100	13	100	26	100

Repayment status of household (institutional Source): The data regarding the repayment status of credit borrowed from institutional Source by households in Bhimanahalli-2 micro watershed is presented in Table 25. The results indicate that, 96.15 per cent have unpaid and 3.85 percent have fully paid.

Table 25. Repayment status of household (institutional Source) in Bhimanahalli-2 micro-watershed

Sl.No.	Particulars	LL (0)		MF (5)		SF (8)		SMF (13)		All (26)	
		N	%	N	%	N	%	N	%	N	%
1	Un paid	0	0	5	100	7	87.5	13	100	25	96.15
2	Fully paid	0	0	0	0	1	12.5	0	0	1	3.85

Opinion regarding institutional sources of credit: The data regarding the opinion on institutional sources of credit in Bhimanahalli-2 micro watershed is presented in Table 26. The results indicate that, 96.15 per cent of the households opined that credit helped to perform timely agricultural operations.

Table 26. Opinion regarding institutional sources of credit in Bhimanahalli-2 micro-watershed

Sl.No	Particulars	MF (5)		SF (8)		SMF (13)		All (26)	
		N	%	N	%	N	%	N	%
1	Helped to perform timely agricultural operations	5	100	8	100	12	92	25	96.2

Cost of Cultivation of Red gram: The data regarding the cost of cultivation (Rs/ha) of Red gram in Bhimanahalli-2 micro watershed is presented in Table 27.a. The results indicate that, the total cost of cultivation (Rs/ha) for Red gram was Rs. 31594.21. The gross income realized by the farmers was Rs. 41712.47. The net income from Red gram cultivation was Rs.10118.26, thus the benefit cost ratio was found to be 1:1.30.

Table 27(a). Cost of Cultivation of Red gram in Bhimanahalli-2 micro-watershed

Sl. No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
I	Cost A1				
1	Hired Human Labour	Man days	35.55	7399.84	23.42
2	Bullock	Pairs/day	1.43	1016.44	3.22
3	Tractor	Hours	4.02	3432.24	10.86
4	Machinery	Hours	0	0	0
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	14.28	1394.27	4.41
7	FYM	Quintal	2.53	3708.91	11.74
8	Fertilizer + micronutrients	Quintal	4.1	3088.7	9.78
9	Pesticides (PPC)	Kgs/liters	1.74	1324.25	4.19
10	Irrigation	Number	0	0	0
11	Repairs		0	0	0
12	Msc. Charges (Marketing costs etc)		0	0	0
13	Depreciation charges		0	80.76	0.26
14	Land revenue and Taxes		0	0	0
II	Cost B1				
16	Interest on working capital			1141.93	3.61
17	Cost B1 = (Cost A1 + sum of 15 and 16)			22587.34	71.49
III	Cost B2				
18	Rental Value of Land			306	0.97
19	Cost B2 = (Cost B1 + Rental value)			22893.34	72.46
IV	Cost C1				
20	Family Human Labour		23.12	5828.67	18.45
21	Cost C1 = (Cost B2 + Family Labour)			28722.01	90.91
V	Cost C2				
22	Risk Premium			0	0
23	Cost C2 = (Cost C1 + Risk Premium)			28722.01	90.91
VI	Cost C3				
24	Managerial Cost			2872.2	9.09
25	Cost C3 = (Cost C2 + Managerial Cost)			31594.21	100
VII	Economics of the Crop				
a.	Main Product	a) Main Product (q)		7.74	41712.47
		b) Main Crop Sales Price (Rs.)			5388
b.	Gross Income (Rs.)			41712.47	
c.	Net Income (Rs.)			10118.26	
d.	Cost per Quintal (Rs./q.)			4081.02	
e.	Benefit Cost Ratio (BC Ratio)			1:1.3	

Cost of Cultivation of Maize: The data regarding the cost of cultivation (Rs/ha) of Maize in Bhimanahalli-2 micro watershed is presented in Table 27.b. The results indicate that, the total cost of cultivation (Rs/ha) for Maize was Rs. 20566.68. The gross income realized by the farmers was Rs. 71167.81. The net income from Maize cultivation was Rs.50601.13, thus the benefit cost ratio was found to be 1:3.50.

Table 27(b). Cost of Cultivation of Maize in Bhimanahalli-2 micro-watershed

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
I	Cost A1				
1	Hired Human Labour	Man days	25.21	5403.79	26.27
2	Bullock	Pairs/day	0.92	551.15	2.68
3	Tractor	Hours	2.99	2243.53	10.91
4	Machinery	Hours	0	0	0
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	22.45	1471.14	7.15
7	FYM	Quintal	1.01	2475.83	12.04
8	Fertilizer + micronutrients	Quintal	3.43	2735.89	13.3
9	Pesticides (PPC)	Kgs / liters	0	0	0
10	Irrigation	Number	0	0	0
11	Repairs		0	0	0
12	Msc. Charges (Marketing costs etc)		0	0	0
13	Depreciation charges		0	16.85	0.08
14	Land revenue and Taxes		0	0	0
II	Cost B1				
16	Interest on working capital			801.94	3.9
17	Cost B1 = (Cost A1 + sum of 15 and 16)			15700.13	76.34
III	Cost B2				
18	Rental Value of Land			283.33	1.38
19	Cost B2 = (Cost B1 + Rental value)			15983.46	77.72
IV	Cost C1				
20	Family Human Labour		10.67	2713.51	13.19
21	Cost C1 = (Cost B2 + Family Labour)			18696.98	90.91
V	Cost C2				
22	Risk Premium			0	0
23	Cost C2 = (Cost C1 + Risk Premium)			18696.98	90.91
VI	Cost C3				
24	Managerial Cost			1869.7	9.09
25	Cost C3 = (Cost C2 + Managerial Cost)			20566.68	100
VII	Economics of the Crop				
a.	Main Product	a) Main Product (q)		19.64	70160.51
		b) Main Crop Sales Price (Rs.)			3571.43
	By Product	e) Main Product (q)		1.18	1007.3
		f) Main Crop Sales Price (Rs.)			857.14
b.	Gross Income (Rs.)			71167.81	
c.	Net Income (Rs.)			50601.13	
d.	Cost per Quintal (Rs./q.)			1046.92	
e.	Benefit Cost Ratio (BC Ratio)			1:3.5	

Cost of Cultivation of Cotton: The data regarding the cost of cultivation (Rs/ha) of Cotton in Bhimanahalli-2 micro watershed is presented in Table 27.c. The results indicate, the total cost of cultivation (Rs/ha) for Cotton was Rs.20292.15. The gross income realized by the farmers was Rs. 31518.23. The net income from Cotton cultivation was Rs. 11226.08, thus the benefit cost ratio was found to be 1:1.50.

Table 27(c). Cost of Cultivation of Cotton in Bhimanahalli-2 micro-watershed

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
I	Cost A1				
1	Hired Human Labour	Man days	25.52	5035.2	24.81
2	Bullock	Pairs/day	0.67	586.63	2.89
3	Tractor	Hours	1.7	1389.38	6.85
4	Machinery	Hours	0	0	0
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	4.12	3334.5	16.43
6	Seed Inter Crop	Kgs.	0	0	0
7	FYM	Quintal	0.62	1543.75	7.61
8	Fertilizer + micronutrients	Quintal	2.98	2737.58	13.49
9	Pesticides (PPC)	Kgs / liters	1.44	1358.5	6.69
10	Irrigation	Number	0	0	0
11	Repairs		0	0	0
12	Msc. Charges (Marketing costs etc)		0	0	0
13	Depreciation charges		0	0.41	0
14	Land revenue and Taxes		0	0	0
II	Cost B1				
16	Interest on working capital			1076.92	5.31
17	Cost B1 = (Cost A1 + sum of 15 and 16)			17062.87	84.09
III	Cost B2				
18	Rental Value of Land			283.33	1.4
19	Cost B2 = (Cost B1 + Rental value)			17346.2	85.48
IV	Cost C1				
20	Family Human Labour		4.48	1101.21	5.43
21	Cost C1 = (Cost B2 + Family Labour)			18447.41	90.91
V	Cost C2				
22	Risk Premium			0	0
23	Cost C2 = (Cost C1 + Risk Premium)			18447.41	90.91
VI	Cost C3				
24	Managerial Cost			1844.74	9.09
25	Cost C3 = (Cost C2 + Managerial Cost)			20292.15	100
VII	Economics of the Crop				
a.	Main Product	a) Main Product (q)		6.43	31518.23
		b) Main Crop Sales Price (Rs.)			4900
b.	Gross Income (Rs.)				31518.23
c.	Net Income (Rs.)				11226.08
d.	Cost per Quintal (Rs./q.)				3154.73
e.	Benefit Cost Ratio (BC Ratio)				1:1.5

Adequacy of fodder: The data regarding the adequacy of fodder in Bhimanahalli-2 Micro watershed is presented in Table 28. The results indicate that, 5.56 per cent of them opined dry fodder was inadequate.

Table 28. Adequacy of fodder in Bhimanahalli-2 micro-watershed

Sl.No.	Particulars	LL (2)		MF (8)		SF (13)		SMF (13)		All (36)	
		N	%	N	%	N	%	N	%	N	%
1	Inadequate-Dry Fodder	0	0	0	0	1	7.69	1	7.69	2	5.56

Average annual gross income: The data regarding the annual gross income in Bhimanahalli-2 Micro watershed is presented in Table 29. The results indicate that, the farmers have annual gross income of Rs. 111534.72 in micro-watershed, of which Rs. 65118.06 is from agriculture itself.

Table 29. Average annual gross income in Bhimanahalli-2 micro-watershed

Sl.No.	Particulars	LL (2)	MF (8)	SF (13)	SMF (13)	All (36)
1	Service/salary	0	0	0	23076.9	8333.33
2	Wage	0	72875	36000	22307.7	37250
3	Agriculture	85000	53212.5	39930.8	94573.1	65118.1
4	Dairy Farm	0	0	0	2307.69	833.33
	Income(Rs.)	85000	126088	75930.8	142265	111535

Average annual Expenditure: The data regarding the average annual expenditure in Bhimanahalli-2 Micro watershed is presented in Table 30. The results indicate that, the farmers have annual gross expenditure of Rs. 130698.72 in micro-watershed, of which Rs. 38055.56 is from agriculture itself.

Table 30. Average annual Expenditure in Bhimanahalli-2 micro-watershed

Sl.No.	Particulars	LL (2)	MF (8)	SF (13)	SMF (13)	All (36)
1	Agriculture	28000	21083.3	23461.5	58153.9	38055.6
	Total	28000	21083.3	23461.5	58153.9	130699

Horticulture species grown: The data regarding horticulture species grown in Bhimanahalli-2 Micro watershed is presented in Table 31. The results indicate that, the total number of horticultural trees grown (both field and backyard) by the sampled households were coconut (5).

Table 31. Horticulture species grown in Bhimanahalli-2 micro-watershed

Sl.No.	Particulars	LL (2)		MF (8)		SF (13)		SMF (13)		MDF (0)		LF (0)		All (36)	
		F	B	F	B	F	B	F	B	F	B	F	B	F	B
1	Coconut	0	0	0	0	0	0	1	4	0	0	0	0	1	4

*F= Field B=Back Yard

Forest species grown: The data regarding forest species grown in Bhimanahalli-2 Micro watershed is presented in Table 32. The results indicate that, households have planted 112 neem trees, 4 tamarind trees and 2 banyan trees together in both field and backyard.

Table 32. Forest species grown in Bhimanahalli-2 micro-watershed

Sl.No.	Particulars	LL (2)		MF (8)		SF (13)		SMF (13)		All (36)	
		F	B	F	B	F	B	F	B	F	B
1	Neem	0	0	4	2	4	0	94	8	102	10
2	Tamarind	0	0	0	0	0	0	4	0	4	0
3	Banyan	0	0	0	0	0	0	2	0	2	0

*F= Field B=Back Yard

Average additional investment capacity: The data regarding average additional investment capacity in Bhimanahalli-2 Micro watershed is presented in Table 33. The results indicate that, households have an average investment capacity of Rs. 1083.33 for land development, Rs.2000.00 for adoption of improved crop production and Rs.83.33 for adoption of improved livestock management.

Table 33. Average additional investment capacity of households in Bhimanahalli-2 micro-watershed

Sl.No.	Particulars	LL (2)	MF (8)	SF (13)	SMF (13)	All (36)
1	Land development	0	750	1923.08	615.38	1083.33
2	Improved crop production	0	1250	3538.46	1230.77	2000
3	Improved livestock management	0	0	0	230.77	83.33

Source of funds for additional investment: The data regarding source of funds for additional investment in Bhimanahalli-2 Micro watershed is presented in Table 34. The results indicate that, the sources of finance raised from own sources for land development and improved crop production was 27.78 per cent and for improved livestock adoption was 2.78 per cent.

Table 34. Source of funds for additional investment in Bhimanahalli-2 micro-watershed

Sl.No	Item	Land development		Improved crop production		Improved livestock management	
		N	%	N	%	N	%
1	Own funds	10	27.78	10	27.78	1	2.78

Table 35. Marketing of agricultural produce in Bhimanahalli-2 micro-watershed

Sl.No	Crops	Output obtained (q)	Output retained (q)	Output sold (q)	Output sold (%)	Avg. Price obtained (Rs/q)
1	Cotton	35	0	35	100	4900
2	Maize	262	36	226	86	3125
3	Redgram	242	52	190	79	4989

Marketing of agricultural produce: The data regarding marketing of the agricultural produce in Bhimanahalli-2 Micro watershed is presented in Table 35. The results indicated

that, 100.00 percent of output of cotton was sold in the market; 86.26 percent of output of maize was sold in the market and 78.51 percent of output of red gram was sold in the market.

Marketing channels used for sale of agricultural produce: The data regarding marketing channels used for sale of agricultural produce in Bhimanahalli-2 Micro watershed is presented in Table 36. The results indicated that, 94.44 cent of the households have sold agricultural produce to the local/village merchants and 8.33 per cent of regulated market.

Table 36. Marketing channels used for sale of agricultural produce in Bhimanahalli-2 micro-watershed

Sl.No.	Particulars	LL (2)		MF (8)		SF (13)		SMF (13)		All (36)	
		N	%	N	%	N	%	N	%	N	%
1	Local/village Merchant	0	0	7	88	11	84.6	16	123	34	94.44
2	Regulated Market	0	0	1	13	2	15.4	0	0	3	8.33

Mode of transport of agricultural produce: The data regarding mode of transport of agricultural produce in Bhimanahalli-2 Micro watershed is presented in Table 37. The results indicated that, 13.89 cent of the households have used tractor and 88.89 per cent have used Cart.

Table 37. Mode of transport of agricultural produce in Bhimanahalli-2 micro-watershed

Sl.No.	Particulars	LL (2)		MF (8)		SF (13)		SMF (13)		All (36)	
		N	%	N	%	N	%	N	%	N	%
1	Cart	0	0	8	100	13	100	11	84.6	32	88.89
2	Tractor	0	0	0	0	0	0	5	38.5	5	13.89

Incidence of soil and water erosion problems: The data regarding incidence of incidence of soil and water erosion problems in Bhimanahalli-2 Micro watershed is presented in Table 38. The results indicate that, 30.56 per cent of the households have experienced soil and water erosion problems.

Table 38. Incidence of soil and water erosion problems in Bhimanahalli-2 micro-watershed

Sl.No	Particulars	LL (2)		MF (8)		SF (13)		SMF (13)		All (36)	
		N	%	N	%	N	%	N	%	N	%
1	Soil and water erosion problems in the farm	0	0	2	25	6	46.2	3	23	11	30.56

Table 39. Interest regarding soil testing in Bhimanahalli-2 micro-watershed

Sl.No.	Particulars	LL (2)		MF (8)		SF (13)		SMF (13)		All (36)	
		N	%	N	%	N	%	N	%	N	%
1	Interest in soil test	0	0	8	100	15	115	15	115	38	105.6

Interest towards soil testing: The data regarding Interest shown towards soil testing in Bhimanahalli-2 Micro watershed is presented in Table 39. The results indicated that, 105.56 per cent of the households were interested towards soil testing.

Usage pattern of fuel for domestic use: The data on usage pattern of fuel for domestic use in Bhimanahalli-2 Micro watershed is presented in Table 40. The results indicated that, firewood was the major source of fuel for domestic use for 75.00 per cent of the households followed by LPG (63.89%).

Table 40. Usage pattern of fuel for domestic use in Bhimanahalli-2 micro-watershed

Sl.No.	Particulars	LL (2)		MF (8)		SF (13)		SMF (13)		All (36)	
		N	%	N	%	N	%	N	%	N	%
1	Fire Wood	2	100	7	87.5	8	61.5	10	76.9	27	75
2	LPG	0	0	10	125	5	38.5	8	61.5	23	63.89

Source of drinking water: The data on source of drinking water in Bhimanahalli-2 Micro watershed is presented in Table 41. The results indicated that, piped supply of water was the major source for drinking water for 100 per cent of the households.

Table 41. Source of drinking water in Bhimanahalli-2 micro-watershed

Sl.No.	Particulars	LL (2)		MF (8)		SF (13)		SMF (13)		All (36)	
		N	%	N	%	N	%	N	%	N	%
1	Piped supply	2	100	8	100	13	100	13	100	36	100

Source of light: The data on source of light in Bhimanahalli-2 Micro watershed is presented in Table 42. The results indicated that, electricity was the major source of light for 100.00 per cent of the households

Table 42. Source of light in Bhimanahalli-2 micro-watershed

Sl.No.	Particulars	LL (2)		MF (8)		SF (13)		SMF (13)		All (36)	
		N	%	N	%	N	%	N	%	N	%
1	Electricity	2	100	8	100	13	100	13	100	36	100

Existence of sanitary toilet facility: The data on availability of toilet facility in Bhimanahalli-2 Micro watershed is presented in Table 43. The results indicated that, 69.44 per cent of the households possess toilets.

Table 43. Existence of sanitary toilet facility in Bhimanahalli-2 micro-watershed

Sl.No.	Particulars	LL (2)		MF (8)		SF (13)		SMF (13)		All (36)	
		N	%	N	%	N	%	N	%	N	%
1	Sanitary toilet facility	1	50	9	113	10	76.92	5	38	25	69.4

Possession of PDS card: The data regarding possession of PDS card in Bhimanahalli-2 Micro watershed is presented in Table 44. The results indicated that, 94.44 per cent of the

households possessed BPL card, 2.78 per cent possessed APL card and 2.78 per cent do not possess PDS card.

Table 44. Possession of PDS card in Bhimanahalli-2 micro-watershed

Sl.No.	Particulars	LL (2)		MF (8)		SF (13)		SMF (13)		All (36)	
		N	%	N	%	N	%	N	%	N	%
1	APL	0	0	0	0	0	0	1	7.7	1	2.78
2	BPL	2	100	8	100	12	92.31	12	92	34	94.44
3	Not Possessed	0	0	0	0	1	7.69	0	0	1	2.78

Participation in NREGA programme: The data regarding Participation in NREGA programme in Bhimanahalli-2 Micro watershed is presented in Table 45. The results indicated that, only 55.56 percent of the participate have participated in NREGA programme.

Table 45. Participation in NREGA programme in Bhimanahalli-2 micro-watershed

Sl.No.	Particulars	LL (2)		MF (8)		SF (13)		SMF (13)		All (36)	
		N	%	N	%	N	%	N	%	N	%
1	Participation in NREGA programme	0	0	1	12.5	9	69.2	10	76.9	20	55.6

Adequacy of food items: The data regarding adequacy of food items in Bhimanahalli-2 Micro watershed is presented in Table 46. The results indicated that, the extent of adequacy of food items for cereals, pulses, Oilseeds and vegetables were 100.00, 100.00, 52.78, 30.56 per cent respectively, similarly for Fruits (33.33%) and milk (38.89%).

Table 46. Adequacy of food items in Bhimanahalli-2 micro-watershed

Sl.No.	Particulars	LL (2)		MF (8)		SF (13)		SMF (13)		All (36)	
		N	%	N	%	N	%	N	%	N	%
1	Cereals	2	100	8	100	13	100	13	100	36	100
2	Pulses	2	100	8	100	13	100	13	100	36	100
3	Oilseed	0	0	3	37.5	8	61.54	8	61.5	19	52.78
4	Vegetables	0	0	2	25	6	46.15	3	23.1	11	30.56
5	Fruits	1	50	4	50	3	23.08	4	30.8	12	33.33
6	Milk	1	50	6	75	3	23.08	4	30.8	14	38.89

Table 47. Inadequacy of food items in Bhimanahalli-2 micro-watershed

Sl.No.	Particulars	LL (2)		MF (8)		SF (13)		SMF (13)		All (36)	
		N	%	N	%	N	%	N	%	N	%
1	Oilseed	2	100	5	62.5	5	38.46	5	38.5	17	47.22
2	Vegetables	2	100	6	75	7	53.85	10	76.9	25	69.44
3	Fruits	1	50	4	50	10	76.92	10	76.9	25	69.44
4	Milk	1	50	4	50	11	84.62	13	100	29	80.56
5	Egg	2	100	7	87.5	13	100	13	100	36	100
8	Meat	2	100	7	87.5	13	100	13	100	36	100

Inadequacy of food items: The data regarding in adequacy of food items in Bhimanahalli-2 Micro watershed is presented in Table 47. The results indicated that, the extent of in

adequacy of food items for Oilseeds and vegetables were 47.22 and 69.44 per cent respectively, similarly for fruits (69.44%), milk (80.56%), egg (100.00%) and meat (100.00%).

Farming constraints: The data regarding farming constraints experienced by households in Bhimanahalli-2 Micro watershed is presented in Table 48. The results indicated that, lower fertility status of the soil was the constraint experienced by (100.00 %) per cent of the households, wild animal menace on farm field (94.44%), frequent incidence of pest and diseases (91.67%), inadequacy of irrigation water (100.00%), high cost of fertilizers and plant protection chemicals (100.00%), high rate of interest on credit (100.00%), low price for the agricultural commodities (100.00 %), lack of marketing facilities in the area (100%), inadequate extension services (97.22 %), lack of transport for safe transport of the agricultural produce to the market (94.44%).

Table 48. Farming constraints experienced in Bhimanahalli-2 micro-watershed

SN	Particulars	LL (2)		MF (8)		SF (13)		SMF (13)		All (36)	
		N	%	N	%	N	%	N	%	N	%
1	Lower fertility status of the soil	0	0	8	100	16	123.08	15	115.38	36	100
2	Wild animal menace on farm field	0	0	8	100	13	100	13	100	34	94.44
3	Frequent incidence of pest and diseases	0	0	8	100	12	92.31	13	100	33	91.67
4	Inadequacy of irrigation water	0	0	8	100	13	100	13	100	36	100
5	High cost of Fertilizers and plant protection chemicals	0	0	8	100	13	100	13	100	36	100
6	High rate of interest on credit	0	0	8	100	13	100	13	100	36	100
7	Low price for the agricultural commodities	0	0	8	100	13	100	13	100	36	100
8	Lack of marketing facilities in the area	0	0	8	100	13	100	13	100	36	100
9	Inadequate extension services	0	0	8	100	14	107.69	13	100	35	97.22
10	Lack of transport for safe transport of the Agril produce to the market.	0	0	8	100	13	100	13	100	34	94.44

SUMMARY AND IMPLICATIONS

In order to assess the socio-economic condition of the farmers in the watershed 36 households located in the micro watershed were interviewed for the survey. The study was conducted in Bhimanahalli-2 micro-watershed (Motanahalli sub-watershed, Chitapura taluk & Kalaburagi District) is located at North latitude $16^{\circ} 58' 39.868''$ and $16^{\circ} 56' 44.222''$ and East longitude $77^{\circ} 13' 58.007''$ and $77^{\circ} 11' 56.808''$ covering an area of about 561.68 ha bounded by under Bhimanahalli and Handarki Villages.

Socio-economic analysis indicated that, out of the total sample of 36 respondents, 8 (22.22%) were marginal, 13(36.11%) were small and 13 (36.11%) were semi medium farmers. The population characteristics of households indicated that, there were 81 (59.12%) men and 56 (40.88%) were women. Majority of the respondents (43.07%) were in the age group of 35-60 years. Education level of the sample households indicated that, majority there were 30.66 per cent illiterates and only 6.57 per cent attained graduation. About, 72.22 per cent of household heads practicing agriculture and 16.67 per cent of the household heads were engaged as agricultural labourers. Agriculture was the major occupation for 48.91 per cent of the household members.

In the study area, 94.44 per cent of the households possess katcha house and 5.56 per cent possess pucca house. The durable assets owned by the households showed that, 83.33 per cent possess TV, 66.67 per cent possess mixer grinder and 108.33 per cent possess mobile phones. Farm implements owned by the households indicated that, 22.22 per cent of the households possess plough and only 8.33 per cent sprayer. Regarding livestock possession by the households, 13.89 per cent possess local cow.

The average labour availability in the study area showed that, own labour men available in the micro watershed was 1.56, women available in the micro watershed was 1.29, hired labour (men) available was 8 and hired labour (women) available was 13.24.

Out of the total land holding of the sample respondents (59.06 ha), 100.00 per cent of the area is under dry condition. The major crops grown by sample farmers are Red gram, Maize and Cotton and cropping intensity was recorded as 99.97 per cent.

The sample households possessed 97.22 per cent bank account and 63.89 per cent of them have savings in the account. About 44.44 per cent of the respondents borrowed credit from various sources. Among the credit borrowed by households, 20.00 per cent have borrowed loan from commercial banks and 68.00 per cent from Cooperative bank. Majority of the respondents (100.00 %) have borrowed loan for agriculture purpose. Regarding the opinion on institutional sources of credit, 96.15 per cent of the households opined that credit helped to perform timely agricultural operations.

The per hectare cost of cultivation for Red gram, Maize and Cotton was Rs.31594.21 , 20566.68 and 20292.15 with benefit cost ratio of 1:1.30, 1: 3.50 and 1: 1.50, respectively.

The average annual gross income of the farmers was Rs. 111534.72 in micro-watershed, of which Rs. 65118.06 comes from agriculture.

The total number of horticultural trees grown (both field and backyard) by the sampled households were coconut (5) and forest species are grown 112 neem trees, 4 tamarind trees and 2 banyan trees together in both field and backyard.

Households have an average investment capacity of Rs. 1083.33 for land development, Rs.2000.00 for adoption of improved crop production and Rs.83.33 for adoption of improved livestock management. Source of funds from own sources for land development and improved crop production was 27.78 per cent and for improved livestock adoption was 2.78 per cent.

Regarding marketing channels, 94.44 per cent of the households have sold agricultural produce to the local/village merchants, while, 8.33 per cent have sold by Agents/Traders. Further, 13.89 per cent of the households have used tractor for the transport of agriculture commodity.

Majority of the farmers (30.56 %) have experienced soil and water erosion problems in the watershed and 105.56 per cent of the households were interested towards soil testing.

Firewood connection was the major source of fuel for domestic use for 75.00 per cent of the households and 63.89 per cent households has LPG. Piped supply was the major source for drinking water for 100.00 per cent of the households. Electricity was the major source of light for 100.00 per cent of the households. In the study area, 69.44 per cent of the households possess toilet facility. Regarding possession of PDS card, 94.44 per cent of the households possessed BPL card and 2.78 per cent do not possess PDS card. Cereals (100.00%), pulses (10.00%), oilseeds (52.78%) were adequate for consumption.

Farming constraints experienced by households in the micro watersheds were lower fertility status of the soil (100.00%) wild animal menace on farm field (94.44%), frequent incidence of pest and diseases (91.67%), inadequacy of irrigation water (100.00%), high cost of fertilizers and plant protection chemicals (100.00%), high rate of interest on credit (100.00%), low price for the agricultural commodities (100.00%), lack of marketing facilities in the area (101.00%), inadequate extension services (97.22%), lack of transport for safe transport of the agricultural produce to the market (94.44%),.

Implications of the survey

- ✓ Result indicated that, there were 30.66 per cent were illiterate hence, extension methodologies such as demonstration, street play, drama, video shows will be effective in dissemination of the technologies in the micro watershed.

- ✓ The data indicate that, 94.44 per cent of the households possess katcha house. Hence, the development department while implementing the watershed plan should focus on agriculture to enhance the productivity of major crops in the area to increase the income of the farmers.
- ✓ Results indicated that the local institutional participation of the household members in the micro watershed is minimal hence, activities like membership campaign, awareness creation about the benefits of membership in local institutions and strengths of organized groups must be conveyed.
- ✓ Majority of the households in the watershed have experience in use of mobile phones, and television hence, these mass media can be effectively utilized for transfer of technology as well as for information dissemination.
- ✓ The farm machinery/implement possession in the micro watershed was found to be minimum the reasons may lack of knowledge or lack of financial ability which can be addressed through training on use of different farm implements, providing information on different sources of finance for purchase of farm implements.
- ✓ The possession of livestock such as crossbred cow found is less hence, farmers must be made aware of the benefits of crossbred cow in increased milk production.
- ✓ The possession of livestock such as sheep, goat and poultry was found to be low hence, farmers may be informed the role of subsidiary enterprises in enhancing the income and information on financial support for subsidiary activities.
- ✓ The data indicate that, job/work was the reason for all the migrants hence, farmers may be trained on profitable agriculture or self employment such as animal husbandry, plate making, sheep rearing, goat rearing, rabbit rearing with suitable information on sources of financial support.
- ✓ The results indicate that there was a change in quality of life due to migration hence, the developmental departments should take actions to arrest migration and to improve the quality of the life in rural areas.
- ✓ Households possess 59.06ha (100.00 %) of dry land hence, the availability of the dryland agricultural technologies such as short duration crops, high yielding drought resistance crop varieties, drip irrigation technology and subsidy information will be helpful for the farmers to enhance the productivity of land and as well as farmers income.
- ✓ Few of the bore well in micro watershed found non functional hence, farmers may be trained on possibility of bore well rejuvenation.
- ✓ The total number of horticultural trees grown (both field and backyard) by the sampled households were coconut (5) and forest species are grown 112 neem trees, 4 tamarind trees and 2 banyan trees together in both field and backyard. Hence, production technologies related to these crops can be made available to the farmers for better adoption.

- ✓ The cropping intensity in the micro watershed was found to be (99.97 %) hence, care must be taken by the implementing agency to bring uncultivated land into cultivation through suitable measures.
- ✓ Many of the household members have borrowed loan from cooperative banks which has higher rate of interest hence, farmers may be sensitized on the different sources of credit with lesser interest rate such SHGs etc.
- ✓ The results indicated the non availability of both green and dry fodder throughout the year hence, fodder development activities can be taken up in the micro watershed.
- ✓ The average annual gross income of the households Rs.65118.06 from agriculture, and Rs. 37250.00 from wages and. Agriculture was found to be the major source of income for households hence; the development activities should focus on productivity enhancement, marketing arrangements and agricultural technology dissemination to have a direct impact on the farmers.
- ✓ The cultivation of forest species is found minimal hence, information and production technology related to agro-forestry and integrated farming system.
- ✓ The data indicated that, 30.56 per cent of the households have experienced soil and water erosion problems. Hence, those farmers who reported the soil and water erosion problems may be given attention while implementation of the watershed development plan.
- ✓ The data indicated that, 105.56 per cent of the households have interest in soil testing hence, farmers must be provided with the information on various institutions which are involved in soil testing for the benefit of the farmers.
- ✓ Except summer ploughing the adoption of other soil and water conservation structures is minimum hence, the farmers in the micro watershed should be sensitized on the use of different conservation structures for soil water conservation.
- ✓ Cereals and pulses found be adequate for per cent of the households respectively hence, farm households and the farm women must be trained on importance of balanced nutrition and role of vegetable, milk, egg, meat in balanced diet.
- ✓ Lower fertility status of the soil (100.00%), wild animal menace on farm field (94.44%), frequent incidence of pest and diseases (91.67%), high cost of fertilizers and plant protection chemicals (100.00%), high rate of interest on credit (100.00%), low price for the agricultural commodities (100.00%), lack of marketing facilities in the area (101.00%), inadequate extension services (97.22%), lack of transport for safe transport of the agricultural produce to the market (94.44%) were the major farming constraints experienced hence, these constraints must be addressed immediately for the welfare of the farmers. Awareness to be created among the farmers to approach nearest KVKs/RSKs and other developmental departments for technical and for subsidized inputs and utilize the well established regulated markets, approaching the contract firms, direct markets to avoid the involvement of middlemen.