



हर कदम, हर डगर  
किसानों का हमसफर  
भारतीय कृषि अनुसंधान परिषद

*AgriSearch with a human touch*

**LAND RESOURCE INVENTORY AND SOCIO-ECONOMIC STATUS OF  
FARM HOUSEHOLDS FOR WATERSHED PLANNING AND  
DEVELOPMENT**

**KOTAGEI-2 (4D5B1B2d) MICROWATERSHED**

**Yadgir Taluk and District, Karnataka**

**Karnataka Watershed Development Project – II**

**SUJALA – III**

**World Bank funded Project**



**THE WORLD BANK**



**ICAR – NATIONAL BUREAU OF SOIL SURVEY AND LAND USE PLANNING**



ICAR - NBSS & LUP

**WATERSHED DEVELOPMENT DEPARTMENT  
GOVT. OF KARNATAKA, BANGALORE**



## **About ICAR - NBSS&LUP**

The ICAR-National Bureau of Soil Survey and Land Use Planning (ICAR-NBSS&LUP), Nagpur, a premier Institute of the Indian Council of Agricultural Research (ICAR), was set up during 1976 with the objective to prepare soil resource maps at national, state and district levels and to provide research inputs in soil resource mapping and its applications, land evaluation, land use planning, land resource management, and database management using GIS for optimising land use on different kinds of soils in the country.

The Bureau has been engaged in carrying out soil resource survey, agro-ecological and soil degradation mapping at the country, state and district levels for qualitative assessment and monitoring the soil health towards viable land use planning. The research activities have resulted in identifying the soil potentials and problems, and the various applications of the soil surveys with the ultimate objective of sustainable agricultural development. The Bureau has the mandate to correlate and classify soils of the country and maintain a National Register of all the established soil series. The Institute is also imparting in-service training to staff of the soil survey agencies in the area of soil survey, land evaluation and soil survey interpretations for land use planning. The Bureau in collaboration with Panjabrao Krishi Vidyapeeth, Akola is running post-graduate teaching and research programme in land resource management, leading to M.Sc. and Ph.D. degrees.

**Citation:** Rajendra Hegde, Ramesh Kumar, S.C., B.A. Dhanorkar, S. Srinivas, M. Lalitha, K.V. Niranjana, R.S. Reddy and S.K. Singh (2019).“Land resource inventory and socio-economic status of farm households for watershed planning and development of Kotagei-2 (4D5B1B2d) Microwatershed, Yadgir Taluk and District, Karnataka”, ICAR-NBSS &LUP Sujala MWS Publ.504, ICAR – NBSS & LUP, RC, Bangalore. p.129 & 37.

### **TO OBTAIN COPIES,**

**Please write to:**

**Director, ICAR - NBSS & LUP,**

Amaravati Road, NAGPUR - 440 033, India

Phone : (0712) 2500386, 2500664, 2500545 (O)

Telefax : 0712-2522534

E-Mail : [director@nbsslup.ernet.in](mailto:director@nbsslup.ernet.in)

Website URL : [nbsslup.in](http://nbsslup.in)

Or

**Head, Regional Centre, ICAR - NBSS&LUP, Hebbal, Bangalore - 560 024**

Phone : (080) 23412242, 23510350 (O)

Telefax : 080-23510350

E-Mail : [nbssrcb@gmail.com](mailto:nbssrcb@gmail.com)



**LAND RESOURCE INVENTORY AND SOCIO-ECONOMIC  
STATUS OF FARM HOUSEHOLDS FOR WATERSHED  
PLANNING AND DEVELOPMENT  
KOTAGEI-2 (4D5B1B2d) MICROWATERSHED  
Yadgir Taluk and District, Karnataka**

**Karnataka Watershed Development Project – II**

**Sujala-III**

**World Bank funded Project**



**ICAR – NATIONAL BUREAU OF SOIL SURVEY AND LAND  
USE PLANNING**



**WATERSHED DEVELOPMENT DEPARTMENT, GOVT. OF  
KARNATAKA, BANGALORE**







## PREFACE

In Karnataka, as in other Indian States, the livelihoods of rural people are intertwined with farming pursuits. The challenges in agriculture are seriously threatening the livelihood of a large number of farmers as they have been practicing farming in contextual factors beyond their control. Climatic factors are the most important ones and have become much more significant in recent times due to rapid climate changes induced by intensive anthropogenic activities affecting our ecosystem in multiple ways. Climate change has become the reality, it is happening and efforts to evolve and demonstrate climate resilient technologies have become essential. Due to the already over stressed scenario of agrarian sector, the climate change is resulting in manifold increase in the complexities, pushing the rural mass to face more and more unpredictable situations. The rising temperatures and unpredictable rainfall patterns are going to test seriously the informed decisions farmers have to make in order to survive in farming and sustain their livelihood.

It is generally recognized that impacts of climate change shall not be uniform across the globe. It is said that impact of climate change is more severe in South Asia. Based on the analysis of meteorological data, it is predicted that in India, there will be upward trend in mean temperature, downward trend in relative humidity, annual rainfall and number of wet days in a year. Also, in general, phenomena like erratic monsoon, spread of tropical diseases, rise in sea levels, changes in availability of fresh water, frequent floods, droughts, heat waves, storms and hurricanes are predicted. Each one of these adverse situations are already being experienced in various parts of India and also at the global level. Decline in agricultural productivity of small and marginal farmers becoming more vulnerable is already witnessed.

In Karnataka, more than 60 per cent of the population live in rural areas and depend on agriculture and allied activities for their livelihood. Though the state has achieved significant progress in increasing the yield of many crops, there is tremendous pressure on the land resources due to the growing and competing demands of various land uses. This is reflected in the alarming rate of land degradation observed. Already more than 50 per cent of the area is affected by various forms of degradation. If this trend continues, the sustainability of the fragile ecosystem will be badly affected. The adverse effects of change in the climatic factors are putting additional stress on the land resources and the farmers dependent on this.

The natural resources (land, water and vegetation) of the state need adequate and constant care and management, backed by site-specific technological interventions and investments particularly by the government. Detailed database pertaining to the nature of the land resources, their constraints, inherent potentials and suitability for various land

based rural enterprises, crops and other uses is a prerequisite for preparing location-specific action plans, which are in tune with the inherent capability of the resources. Any effort to evolve climate resilient technologies has to be based on the baseline scientific database. Then only one can expect effective implementation of climate resilient technologies, monitor the progress, make essential review of the strategy, and finally evaluate the effectiveness of the implemented programs. The information available at present on the land resources of the state are of general nature and useful only for general purpose planning. Since the need of the hour is to have site-specific information suitable for farm level planning and detailed characterization and delineation of the existing land resources of an area into similar management units is the only option.

ICAR-NBSS&LUP, Regional Centre, Bangalore has taken up a project sponsored by the Karnataka Watershed Development Project-II, (Sujala-III), Government of Karnataka funded by the World Bank under Component-1 Land Resource Inventory. This study was taken up to demonstrate the utility of such a database in reviewing, monitoring and evaluating all the land based watershed development programs on a scientific footing. To meet the requirements of various land use planners at grassroots level, the present study on Land Resource Inventory and Socio-Economic Status of Farm Households for Watershed Planning and Development of for Kotageri-2 microwatershed in Yadgir Taluk and District, Karnataka” for integrated development was taken up in collaboration with the State Agricultural Universities, IISC, KRSRAC, KSNDMC as Consortia partners. The project provides detailed land resource information at cadastral level (1:7920 scale) for all the plots and socio-economic status of farm households covering thirty per cent farmers randomly selected representing landed and landless class of farmers in the microwatershed. The project report with the accompanying maps for the microwatershed will provide required detailed database for evolving effective land use plan, alternative land use options and conservation plans for the planners, administrators, agricultural extension personnel, KVK officials, developmental departments and other land users to manage the land resources in a sustainable manner.

It is hoped that this database will be useful to the planners, administrators and developmental agencies working in the area in not only for formulating location specific developmental schemes but also for their effective monitoring at the village/watershed level.

Nagpur

Date: 01-11-2019

**S.K. SINGH**

Director, ICAR - NBSS&LUP, Nagpur

## Contributors

<b>Dr. Rajendra Hegde</b> Principal Scientist, Head & Project Leader, Sujala-III Project ICAR-NBSS&LUP, Regional Centre, Bangalore	<b>Dr. S.K.Singh</b> Director, ICAR-NBSS&LUP Coordinator, Sujala-III Project Nagpur
<b>Soil Survey, Mapping &amp; Report Preparation</b>	
Dr. B.A. Dhanorkar	Sh. R.S. Reddy
Dr. K.V. Niranjana	Smt. Chaitra, S.P.
	Dr. Gopali Bardhan
	Dr. Mahendra Kumar M.B
	Mr. Somashekar T.N
	Ms. Arpitha G.M
<b>Field Work</b>	
Sh. C.BacheGowda	Sh. Mahesh, D.B.
Sh. Somashekar	Sh. Ashok S Sindagi
Sh. M. Jayaramaiah	Sh. Veerabhadrapa B.
Sh. Paramesha, K.	Sh. Shankarappa
Sh. B. M. Narayana Reddy	Sh. Anand
	Sh. Arun N Kambar.
	Sh Kamalesh Awate
	Sh. Sharaan Kumar Huppar
	Sh. Yogesh H.N.
	Sh. Kalaveerachari R Kammar
<b>GIS Work</b>	
Dr. S.Srinivas	Sh. A.G.Devendra Prasad
Sh. D.H.Venkatesh	Sh. Prakashanaik, M.K.
Smt.K.Sujatha	Sh. Abhijith Sastry, N.S.
Smt. K.V.Archana	Sh. Sudip Kumar Suklabaidya
Sh. N. Maddileti	Sh. Avinash, K.N.
	Sh. Amar Suputhra, S
	Sh. Deepak, M.J.
	Smt. K.Karunya Lakshmi
	Ms. Seema, K.V.
	Ms. A. Rajab Nisha

<b>Laboratory Analysis</b>	
Dr. M. Lalitha	Sh. Vindhya, N.G.
Smt. Arti Koyal	Ms. P. Pavanakumari, P.
Smt. Parvathy, S.	Ms. Rashmi, N.
	Ms. Leelavathy, K.U.
	Smt. Usha Kiran, G.
<b>Socio-Economic Analysis</b>	
Dr. S.C. Ramesh Kumar	Sh. M.K. Prakashanaik
	Ms. Karuna V. Kulkarni
	Mrs. Sowmya A.N
	Sh. Vinod R
	Sh. Basavaraja
	Sh. Vijay Kumar Lamani
	Ms. Sowmya K.B
	Mrs. Prathibha, D.G
	Sh. Rajendra,D
<b>Soil &amp; Water Conservation</b>	
Sh. Sunil P. Maske	
<b>Watershed Development Department, GoK, Bangalore</b>	
Sh. Prabhash Chandra Ray, IFS Project Director & Commissioner, WDD	Dr. A. Natarajan NRM Consultant, Sujala-III Project
Sh. A. Padmaya Naik, Director (In-Charge) Executive Director, KWDP-II, Sujala-III, WDD	

# **PART-A**

## **LAND RESOURCE INVENTORY**



## Contents

Preface		
Contributors		
Executive Summary		
Chapter 1	Introduction	1
Chapter 2	Geographical Setting	3
2.1	Location and Extent	3
2.2	Geology	3
2.3	Physiography	4
2.4	Drainage	4
2.5	Climate	4
2.6	Natural Vegetation	6
2.7	Land Utilization	6
Chapter 3	Survey Methodology	11
3.1	Base maps	11
3.2	Image Interpretation for Physiography	11
3.3	Field Investigation	14
3.4	Soil Mapping	16
3.5	Land Management Units	16
3.6	Laboratory Characterization	17
Chapter 4	The Soils	21
4.1	Soils of granite gneiss landscape	21
Chapter 5	Interpretation for Land Resource Management	35
5.1	Land Capability Classification	35
5.2	Soil Depth	37
5.3	Surface Soil Texture	38
5.4	Soil Gravelliness	39
5.5	Available Water Capacity	40
5.6	Soil Slope	41
5.7	Soil Erosion	42
Chapter 6	Fertility Status	45
6.1	Soil Reaction (pH)	45
6.2	Electrical Conductivity (EC)	45
6.3	Organic Carbon (OC)	45
6.4	Available Phosphorus	47
6.5	Available Potassium	47
6.6	Available Sulphur	47
6.7	Available Boron	48
6.8	Available Iron	48
6.9	Available Manganese	48
6.10	Available Copper	48
6.11	Available Zinc	52

Chapter 7	Land Suitability for Major Crops	53
7.1	Land suitability for Sorghum	53
7.2	Land suitability for Maize	54
7.3	Land suitability for Bajra	55
7.4	Land suitability for Groundnut	56
7.5	Land suitability for Sunflower	57
7.6	Land suitability for Redgram	58
7.7	Land suitability for Bengal gram	59
7.8	Land suitability for Cotton	60
7.9	Land suitability for Chilli	61
7.10	Land suitability for Tomato	62
7.11	Land suitability for Brinjal	63
7.12	Land suitability for Onion	64
7.13	Land suitability for Bhendi	65
7.14	Land suitability for Drumstick	66
7.15	Land suitability for Mango	67
7.16	Land suitability for Guava	68
7.17	Land suitability for Sapota	69
7.18	Land Suitability for Pomegranate	70
7.19	Land Suitability for Musambi	71
7.20	Land Suitability for Lime	72
7.21	Land Suitability for Amla	73
7.22	Land Suitability for Cashew	74
7.23	Land Suitability for Jackfruit	75
7.24	Land Suitability for Jamun	76
7.25	Land Suitability for Custard apple	77
7.26	Land Suitability for Tamarind	78
7.27	Land Suitability for Mulberry	79
7.28	Land Suitability for Marigold	80
7.29	Land Suitability for Chrysanthemum	81
7.30	Land Management Units (LMUs)	113
7.31	Proposed Crop Plan for Kotageri-2 Microwatershed	114
Chapter 8	Soil Health Management	117
Chapter 9	Soil and Water conservation Treatment Plan	121
9.1	Treatment Plan	121
9.2	Recommended Soil and Water Conservation measures	125
9.3	Greening of Microwatershed	126
	References	129
	Appendix I	I-XII
	Appendix II	XIII-XXIV
	Appendix III	XXV-XXXIV



## LIST OF TABLES

2.1	Mean Monthly Rainfall, PET, 1/2 PET at Yadgir Taluk & District	5
2.2	Land Utilization in Yadgir district	7
3.1	Differentiating Characteristics used for Identifying Soil Series	16
3.2	Soil map unit description of Kotageri-2 Microwatershed	17
4.1	Physical and Chemical Characteristics of Soil Series identified in Kotageri-2 microwatershed	27
7.1	Soil-Site Characteristics of Kotageri-2 Microwatershed	83
7.2	Crop suitability for Sorghum	84
7.3	Crop suitability for Maize	85
7.4	Crop suitability for Bajra	86
7.5	Crop suitability for Groundnut	87
7.6	Crop suitability for Sunflower	88
7.7	Crop suitability for Redgram	89
7.8	Crop suitability for Bengal gram	90
7.9	Crop suitability for Cotton	91
7.10	Crop suitability for Chilli	92
7.11	Crop suitability for Tomato	93
7.12	Crop suitability for Brinjal	94
7.13	Crop suitability for Onion	95
7.14	Crop suitability for Bhendi	96
7.15	Crop suitability for Drumstick	97
7.16	Crop suitability for Mango	98
7.17	Crop suitability for Guava	99
7.18	Crop suitability for Sapota	100
7.19	Crop suitability for Pomegranate	101
7.20	Crop suitability for Musambi	102
7.21	Crop suitability for Lime	103
7.22	Crop suitability for Amla	104
7.23	Crop suitability for Cashew	105
7.24	Crop suitability for Jackfruit	106
7.25	Crop suitability for Jamun	107
7.26	Crop suitability for Custard apple	108

7.27	Crop suitability for Tamarind	109
7.28	Crop suitability for Mulberry	110
7.29	Crop suitability for Marigold	111
7.30	Crop suitability for Chrysanthemum	112
7.31	Proposed Crop Plan for Kotageri-2 Microwatershed	115

## LIST OF FIGURES

2.1	Location map of Kotageri-2 Microwatershed	3
2.2	Granite and granite gneiss rock formation	4
2.3	Rainfall distribution in Yadgir Taluk & District	5
2.4	Natural vegetation of Kotageri-2 Microwatershed	6
2.5	Current Land use map of Kotageri-2 Microwatershed	7
2.6 a&b	Different crops and cropping systems in Kotageri-2 Microwatershed	8
3.1	Scanned and Digitized Cadastral map of Kotageri-2 Microwatershed	13
3.2	Satellite image of Kotageri-2 Microwatershed	13
3.3	Cadastral map overlaid on IRS PAN+LISS IV merged imagery of Kotageri-2 Microwatershed	14
3.4	Location of profiles in a transect	15
3.5	Soil phase or management units of Kotageri-2 Microwatershed	19
5.1	Land Capability Classification map of Kotageri-2 Microwatershed	37
5.2	Soil Depth map of Kotageri-2 Microwatershed	38
5.3	Surface Soil Texture map of Kotageri-2 Microwatershed	39
5.4	Soil Gravelliness map of Kotageri-2 Microwatershed	40
5.5	Soil Available Water Capacity map Kotageri-2 Microwatershed	41
5.6	Soil Slope map of Kotageri-2 Microwatershed	42
5.7	Soil Erosion map of Kotageri-2 Microwatershed	43
6.1	Soil Reaction (pH) map of Kotageri-2 Microwatershed	46
6.2	Electrical Conductivity (EC) map of Kotageri-2 Microwatershed	46
6.3	Soil Organic Carbon (OC) map of Kotageri-2 Microwatershed	47
6.4	Soil Available Phosphorus map of Kotageri-2 Microwatershed	48
6.5	Soil Available Potassium map of Kotageri-2 Microwatershed	49
6.6	Soil Available Sulphur map of Kotageri-2 Microwatershed	49
6.7	Soil Available Boron map of Kotageri-2 Microwatershed	50
6.8	Soil Available Iron map of Kotageri-2 Microwatershed	50
6.9	Soil Available Manganese map of Kotageri-2 Microwatershed	51
6.10	Soil Available Copper map of Kotageri-2 Microwatershed	51
6.11	Soil Available Zinc map of Kotageri-2 Microwatershed	52
7.1	Land suitability for Sorghum	54
7.2	Land suitability for Maize	55
7.3	Land suitability for Bajra	56
7.4	Land suitability for Groundnut	57
7.5	Land suitability for Sunflower	58

7.6	Land suitability for Redgram	59
7.7	Land suitability for Bengal gram	60
7.8	Land suitability for Cotton	61
7.9	Land suitability for Chilli	62
7.10	Land suitability for Tomato	63
7.11	Land suitable for Brinjal	64
7.12	Land suitable for Onion	65
7.13	Land suitable for Bhendi	66
7.14	Land suitable for Drumstick	67
7.15	Land suitability for Mango	68
7.16	Land suitability for Guava	69
7.17	Land suitability for Sapota	70
7.18	Land suitability for Pomegranate	71
7.19	Land suitability for Musambi	72
7.20	Land suitability for Lime	73
7.21	Land suitability for Amla	74
7.22	Land suitability for Cashew	75
7.23	Land suitability for Jackfruit	76
7.24	Land suitability for Jamun	77
7.25	Land suitability for Custard apple	78
7.26	Land suitability for Tamarind	79
7.27	Land suitability for Mulberry	80
7.28	Land suitability for Marigold	81
7.29	Land suitability for Chrysanthemum	82
7.30	Land management units map of Kotageri-2 Microwatershed	113
9.1	Soil and water conservation plan map of Kotageri-2 Microwatershed	126

## **EXECUTIVE SUMMARY**

*The land resource inventory of Kotageri-2 Microwatershed was conducted using village cadastral maps and IRS satellite imagery on 1:7920 scale. The false colour composites of IRS imagery were interpreted for physiography and the physiographic delineations were used as base for mapping soils. The soils were studied in several transects and a soil map was prepared with phases of soil series as mapping units. Random checks were made all over the area outside the transects to confirm and validate the soil map unit boundaries. The soil map shows the geographic distribution and extent, characteristics, classification, behavior and use potentials of the soils in the microwatershed.*

*The present study covers an area of 302 ha in Yadgir taluk & district, Karnataka. The climate is semiarid and categorized as drought-prone with an average annual rainfall of 866 mm, of which about 652 mm is received during south-west monsoon, 138 mm during north-east and the remaining 76 mm during the rest of the year. An area of 265 ha in the microwatershed is covered by soils and about 37 ha by others (Habitation and water body). The salient findings from the land resource inventory are summarized briefly below.*

- ❖ The soils belong to 8 soil series and 11 soil phases (management units) and 5 land management units.*
- ❖ The length of crop growing period is about 120-150 days starting from 1<sup>st</sup> week of June to 4<sup>th</sup> week of October.*
- ❖ From the master soil map, several interpretative and thematic maps like land capability, soil depth, surface soil texture, soil gravelliness, available water capacity, soil slope and soil erosion were generated.*
- ❖ Soil fertility status maps for macro and micronutrients were generated based on the surface soil samples collected at every 320 m grid interval.*
- ❖ Land suitability for growing 29 major agricultural and horticultural crops was assessed and maps showing the degree of suitability along with constraints were generated.*
- ❖ An area of about 88 per cent is suitable for agriculture in the microwatershed.*
- ❖ About 5 per cent soils are moderately deep (75-100), whereas 6 per cent soils are moderately shallow (50-75 cm) and 76 per cent soils are shallow (25-50 cm) and very shallow (<25cm) in the microwatershed.*
- ❖ About 4 percent soils are sandy, 32 percent soils are loamy and 52 per cent is clayey soils at the surface.*
- ❖ An area of about 79 per cent is non gravelly (<15%) soils, about 8 per cent soils are gravelly (15-35%) in the microwatershed.*
- ❖ About <1 per cent soils are medium (101-150 mm/m) and 88 per cent soils are low (51-100 mm/m) and very low (<50mm/m) in available water capacity.*

- ❖ *An area of about 84 per cent is very gently sloping (1-3% slope) lands and about 4 per cent is gently sloping (3-5% slope) lands.*
- ❖ *An area of about 40 per cent is moderately (e2) eroded and about 48 per cent is severely (e3) eroded in the microwatershed.*
- ❖ *Entire cultivated area in the microwatershed is neutral (pH 6.5-7.3), in soil reaction.*
- ❖ *The Electrical Conductivity (EC) of the soils in the entire cultivated area of the microwatershed is  $<2 \text{ ds}^{\text{m}^{-1}}$  indicating that the soils are non-saline.*
- ❖ *An area of  $<1$  per cent is high ( $>0.75\%$ ), 19 percent is medium (0.50-0.75%) and 69 percent is low ( $<0.50\%$ ) in organic carbon content.*
- ❖ *An area of about 56 percent is medium (23-57 kg/ha) and 32 percent soils are high ( $>57$  kg/ha) in available phosphorus.*
- ❖ *An area of about 6 percent is low ( $<145$  kg/ha) and about 81 percent is medium (145-337kg/ha) in available potassium.*
- ❖ *Available sulphur is medium (10-20 ppm) in an area of about 62 percent and low ( $<10$  ppm) in about 26 per cent of microwatershed.*
- ❖ *Available boron content is medium (0.5-1.0 ppm) in an area of 15 per cent and about 73 per cent is low ( $<0.5$  ppm) in available boron in the microwatershed.*
- ❖ *Available iron content is sufficient ( $>4.5$ ppm) in the entire cultivated area of the microwatershed.*
- ❖ *Available manganese and copper are sufficient in all the soils of the microwatershed.*
- ❖ *Available zinc content is deficient ( $<0.6$  ppm) covering a maximum area of 72 per cent and sufficient ( $>0.6$  ppm) about 15 per cent of the microwatershed.*
- ❖ *The land suitability for 29 major crops grown in the microwatershed were assessed and the areas that are highly suitable (S1) and moderately suitable (S2) are given below. It is however to be noted that a given soil may be suitable for various crops but what specific crop to be grown may be decided by the farmer looking to his capacity to invest on various inputs, marketing infrastructure, market price and finally the demand and supply position.*

**Land suitability for various crops in the Microwatershed**

<b>Crop</b>	<b>Suitability Area in ha (%)</b>		<b>Crop</b>	<b>Suitability Area in ha (%)</b>	
	<b>Highly suitable (S1)</b>	<b>Moderately suitable (S2)</b>		<b>Highly suitable (S1)</b>	<b>Moderately suitable (S2)</b>
<i>Sorghum</i>	<1(<1)	33(11)	<i>Guava</i>	-	15(5)
<i>Maize</i>	15(5)	18(6)	<i>Sapota</i>	-	15(5)
<i>Bajra</i>	15(5)	18(6)	<i>Pomegranate</i>	-	15(5)
<i>Groundnut</i>	15(5)	18(6)	<i>Musambi</i>	-	15(5)
<i>Sunflower</i>	-	15(5)	<i>Lime</i>	-	15(5)
<i>Redgram</i>	-	15(5)	<i>Amla</i>	15(5)	18(6)
<i>Bengal gram</i>	-	-	<i>Cashew</i>	-	<1(<1)
<i>Cotton</i>	-	<1(<1)	<i>Jackfruit</i>	-	15(5)
<i>Chilli</i>	15(5)	18(6)	<i>Jamun</i>	-	-
<i>Tomato</i>	15(5)	18(6)	<i>Custard apple</i>	15(5)	18(6)
<i>Brinjal</i>	15(5)	18(6)	<i>Tamarind</i>	-	-
<i>Onion</i>	15(5)	18(6)	<i>Mulberry</i>	-	15(5)
<i>Bhendi</i>	15(5)	18(6)	<i>Marigold</i>	15(5)	18(6)
<i>Drumstick</i>	-	15(5)	<i>Chrysanthemum</i>	15(5)	18(6)
<i>Mango</i>	-	-			

- ❖ *Apart from the individual crop suitability, a proposed crop plan has been prepared for the identified LMUs by considering only the highly and moderately suitable lands for different crops and cropping systems with food, fodder, fiber and horticulture crops.*
- ❖ *Maintaining soil-health is vital to crop production and conserve soil and land resource base for maintaining ecological balance and to mitigate climate change. For this, several ameliorative measures have been suggested to these problematic soils like saline/alkali, highly eroded, sandy soils etc.*
- ❖ *Soil and water conservation treatment plan has been prepared that would help in identifying the sites to be treated and also the type of structures required.*
- ❖ *As part of the greening programme, several tree species have been suggested to be planted in marginal and submarginal lands, field bunds and also in the hillocks, mounds and ridges. This would help in not only supplementing the farm income but also provide fodder and fuel to generate lot of biomass which would help in maintaining an ecological balance and also contribute to mitigating the climate change.*





## **INTRODUCTION**

Land is a scarce resource and basic unit for any material production. It can support the needs of the growing population, provided they use the land in a rational and judicious manner. But what is happening in many areas of the state is a cause for concern to everyone involved in the management of land resources at the grassroots level. The area available for agriculture is about 51 per cent of the total geographical area and more than 60 per cent of the people are still dependant on agriculture for their livelihood. The limited land area is under severe stress and strain due to increasing population pressure and competing demands of various land uses. Due to this, every year there is significant diversion of farm lands and water resources for non-agricultural purposes. Apart from this, due to lack of interest in farmers for farming, large tracts of cultivable lands are turning into fallows in many areas and this trend is continuing at an alarming rate.

Further, land degradation has emerged as a serious problem which has already affected about 38 lakh ha of cultivated area in the state. Soil erosion alone has degraded about 35 lakh ha. Almost all the uncultivated areas are facing various degrees of degradation, particularly soil erosion. Salinity and alkalinity has emerged as a major problem in more than 3.5 lakh ha in the irrigated areas of the state. Nutrient depletion and declining factor productivity is common in both rainfed and irrigated areas. The degradation is continuing at an alarming rate and there appears to be no systematic effort among the stakeholders to contain this process. In recent times, an aberration of weather due to climate change phenomenon has added another dimension leading to unpredictable situations to be tackled by the farmers.

In this critical juncture, the challenge before us is not only to increase the productivity per unit area which is steadily declining and showing a fatigue syndrome, but also to prevent or at least reduce the severity of degradation. If the situation is not reversed at the earliest, then the sustainability of the already fragile crop production system and the overall ecosystem will be badly affected in the state. The continued neglect and unscientific use of the resources for a long time has led to the situation observed at present in the state. It is a known fact and established beyond doubt by many studies in the past that the cause for all kinds of degradation is the neglect and irrational use of the land resources. Hence, there is an urgent need to generate a detailed site-specific farm level database on various land resources for all the villages/watersheds in a time bound manner that would help to protect the valuable soil and land resources and also to stabilize the farm production.

Therefore, the land resource inventory required for farm level planning is the one which investigates not only the surface but also consider the other parameters which are critical for productivity *viz.*, soils, climate, water, minerals and rocks, topography, geology, hydrology, vegetation, crops, land use pattern, animal population, socio-

economic conditions, infrastructure, marketing facilities and various schemes and developmental works of the government etc. From the data collected at farm level, the specific problems and potentials of the area can be identified and highlighted, conservation measures required for the area can be planned on a scientific footing, suitability of the area for various uses can be worked out and finally viable and sustainable land use options suitable for each and every land holding can be prescribed.

The Land Resource Inventory is basically done for identifying the potential and problem areas, developing sustainable land use plans, estimation of surface run off and water harvesting potential, preparation of soil and water conservation plans, land degradation/desertification etc. The Bureau is presently engaged in developing an LRI methodology using high resolution satellite remote sensing data and Digital Elevation Model (DEM) data to prepare Landscape Ecological Units (LEU) map representing agro-ecosystem as a whole. The LEU is preferred over landform as the base map for LRI. LEU is the assemblage of landform, slope and land use. An attempt has already been made to upscale the soil resource information from 1:250000 and 1:50000 scale to the LEU map in Goa and other states.

The land resource inventory aims to provide site specific database for Kotageri-2 microwatershed in Yadgir Taluk & District, Karnataka State for the Karnataka Watershed Development Department. The database was generated by using cadastral map of the village as a base along with high resolution IRS LISS IV and Cartosat-1 merged satellite imagery. Later, an attempt will be made to uplink this LRI data generated at 1:7920 scale under Sujala-III Project to the proposed Landscape Ecological Units (LEUs) map.

The study was organized and executed by the ICAR- National Bureau of Soil Survey and Land Use Planning, Regional Centre, Bangalore under Generation of Land Resource Inventory Data Base Component-1 of the Sujala-III Project funded by the World Bank.

## GEOGRAPHICAL SETTING

### 2.1 Location and Extent

The Kotageri-2 microwatershed is located in the northern part of Karnataka in Yadgir Taluk & District, Karnataka State (Fig.2.1). It comprises parts of Kootagera, Fatthepura and Gajarakota villages. It lies between  $16^{\circ} 52'$  and  $16^{\circ} 54'$  North latitudes and  $77^{\circ} 13'$  and  $77^{\circ} 15'$  East longitudes, covering an area of about 302 ha. It is on the north side of Yadgir town and is surrounded by Kootagera on the north, northeast and northwest and Gajarakota on southeast, northeast and east side and Fatthepura on southern side of the microwatershed.




Fig.2.1 Location map of Kotageri-2 Microwatershed

### 2.2 Geology

Major rock formations observed in the microwatershed are granite gneiss (Figs.2.2a). Granite gneisses are essentially pink to gray and are coarse to medium grained. They

consist primarily of quartz, feldspar, biotite and hornblende. The gray granite gneisses are highly weathered, fractured and fissured upto a depth of about 10 m. Dolerite dykes and quartz veins are common with variable width and found to occur in Kotageri-2 microwatershed.



Fig.2.2a Granite and granite gneiss rocks

### **2.3 Physiography**

Physiographically, the area has been identified as granite gneiss based on geology. The area has been further subdivided into five landforms, *viz*; mounds/ridges, summits, side slopes and very gently sloping uplands, plains and valleys based on slope and its relief features. The elevation ranges from 524-554 m above MSL. The mounds and ridges are mostly covered by rock outcrops.

### **2.4 Drainage**

The area is drained by several parallel streams like Bori, Amerja and Kanga which finally join the river Bhima along its course. Though, they are not perennial, during rainy season they carry large quantities of rain water. The microwatershed has only few small tanks which are not capable of storing the water that flows during the rainy season. Due to this, the ground water recharge is very much affected. This is reflected in the failure of many bore wells in the villages. If the available rain water is properly harnessed by constructing new tanks and recharge structures at appropriate places in the villages, then the drinking and irrigation needs of the area can be easily met. The drainage network is parallel to sub parallel and dendritic.


### **2.5 Climate**

The Yadgir district lies in the northern plains of Karnataka and falls under semiarid tract of the state and is categorized as drought- prone with total annual rainfall of 866 mm (Table 2.1). Of the total rainfall, maximum of 652 mm is received during the south–west monsoon period from June to September; the north-east monsoon from

October to early December contributes about 138 mm and the remaining 76 mm during the rest of the year. The summer season starts during the middle of February and continues up to the first week of June. The period from December to the middle of February is the coldest season. December is the coldest month with mean daily maximum and minimum temperatures being 29.5<sup>0</sup>C and 10<sup>0</sup>C respectively. During peak summer, temperature shoots up to 45<sup>0</sup>C. Relative humidity varies from 26% in summer to 62% in winter. Rainfall distribution is shown in Figure 2.3. The average Potential Evapo-Transpiration (PET) is 141 mm and varies from a low of 81 mm in December to 199 mm in the month of May. The PET is always higher than precipitation in all the months except end of June to end of September. Generally, the Length of crop Growing Period (LGP) is 120-150 days and starts from 1<sup>st</sup> week of June to 4<sup>th</sup> week of October.

**Table 2.1 Mean Monthly Rainfall, PET, 1/2 PET at Yadgir Taluk, Yadgir District**

Sl. No.	Months	Rainfall	PET	1/2 PET
1	January	4.30	86.0	43.0
2	February	2.30	125.5	62.7
3	March	15.10	166.0	83.0
4	April	18.50	179.8	89.9
5	May	36.0	198.8	97.9
6	June	118.0	175.1	87.5
7	July	171.80	156.3	78.1
8	August	182.9	150.3	75.1
9	September	179.7	142.0	71.0
10	October	105.3	138.5	69.2
11	November	26.4	97.60	48.6
12	December	6.0	80.90	40.4
<b>Total</b>		<b>866.3</b>		



**Fig 2.3 Rainfall distribution in Yadgir Taluk, Yadgir District**



## 2.6 Natural Vegetation

The natural vegetation is sparse comprising few tree species, shrubs and herbs. The mounds, ridges and boulders occupy very sizeable area which is under thin to moderately thick forest vegetation. Still, there are some remnants of the past forest cover which can be seen in patches in some ridges and hillocks in the microwatershed (Fig 2.4).

Apart from the continuing deforestation, the presence of large population of goats, sheep and other cattle in the microwatershed is causing vegetative degradation of whatever little vegetation left in the area. The uncontrolled grazing has left no time for the regeneration of the vegetative cover. This leads to the accelerated rate of erosion on the hill slopes resulting in the formation of deep gullies in the foot slopes that eventually result in the heavy siltation of tanks and reservoirs in the microwatershed.




Fig 2.4 Natural vegetation of Kotageri-2 Microwatershed

## 2.7 Land Utilization

About 72 per cent area (Table 2.2) in Yadgir district is cultivated at present. An area of about 2 per cent is permanently under pasture, 20 per cent under current fallows and 6 per cent under non-agricultural land and 5 per cent under currently barren. Forests occupy an area of about 7 per cent and the tree cover is in a very poor state. Most of the mounds, ridges and bouldery areas have very poor vegetative cover. Major crops grown in the area are sorghum, maize, cotton, sunflower, groundnut, red gram, mango, pomegranate, marigold and sapota. While carrying out land resource inventory, the land use/land cover particulars are collected from all the survey numbers and a current land use map of the microwatershed is prepared. The current land use map prepared shows the arable and non-arable lands, other land uses and different types of crops grown in the area. The current land use map of Kotageri-2 microwatershed is presented in Fig.2.5. The different crops and cropping systems adopted in the microwatershed are presented in Figures 2.6 a & b.

**Table 2.2 Land Utilization in Yadgir District**

Sl. No.	Agricultural land use	Area ( ha)	Per cent
1	Total geographical area	516088	-
2	Total cultivated area	373617	72.4
3	Area sown more than once	74081	14.3
4	Cropping intensity	-	119.8
5	Trees and grooves	737	0.14
6	Forest	33773	6.54
7	Cultivable wasteland	2385	0.46
8	Permanent Pasture land	11755	2.28
9	Barren land	27954	5.41
10	Non- Agriculture land	29623	5.73
11	Current Fallows	105212	20.4



**Fig.2.5 Current Land Use map of Kotageri-2 Microwatershed**







Fig. 2.6 a. Different Crops and Cropping Systems in Kotageri-2 Microwatershed





Mango



Pomegranate



Cotton



Sorghum

Fig. 2.6 b. Different Crops and Cropping Systems in Kotageri-2 Microwatershed



## SURVEY METHODOLOGY

The purpose of land resource inventory is to delineate similar areas (soil series and phases), which respond or expected to respond similarly to a given level of management. This was achieved in Kotageri-2 microwatershed by the detailed study of all the soil characteristics (depth, texture, colour, structure, consistence, coarse fragments, porosity, soil reaction, soil horizons etc.) and site characteristics (slope of the land, erosion, drainage, occurrence of rock fragments etc.) followed by grouping of similar areas based on soil-site characteristics into homogeneous (management units) units, and showing the area extent and their geographic distribution on the microwatershed cadastral map. The detailed survey at 1:7920 scale was carried out in an area of 301 ha. The methodology followed for carrying out land resource inventory was as per the guidelines given in Soil Survey Manual (IARI, 1971; Soil Survey Staff, 2006; Natarajan *et al.*, 2015) which is briefly described below.

### 3.1 Base Maps

The detailed survey of the land resources occurring in the microwatershed was carried out by using digitized cadastral and IRS satellite imagery map as base supplied by KSRSAC. The cadastral map shows field boundaries with their survey numbers, location of tanks, streams and other permanent features of the area (Fig. 3.1). Apart from the cadastral map, remote sensing data products from Cartosat-1 and LISS IV merged at the scale of 1:7920 were used in conjunction with the cadastral map to identify the landscapes, landforms and other surface features. The imagery helped in the identification and delineation of boundaries between hills, uplands and lowlands, water bodies, forest and vegetated areas, roads, habitations and other cultural features of the area (Fig. 3.2). The cadastral map was overlaid on the satellite imagery (Fig. 3.3) that helps to identify the parcel boundaries and other permanent features. Apart from cadastral maps and images, toposheets of the area (1:50,000 scale) were also used for initial traversing, identification of geology and landforms, drainage features, present land use and also for selection of transects in the microwatershed.

### 3.2 Image Interpretation for Physiography

False Colour Composites (FCCs) of Cartosat-I and LISS-IV merged satellite data covering microwatershed area was visually interpreted using image interpretation elements and all the available collateral data with local knowledge. The delineated physiographic boundaries were transferred on to a cadastral map overlaid on satellite imagery. Physiographically, the area has been identified as granite gneiss landscape. It was divided into five landforms, *viz.*; ridges and mounds, gently and very gently sloping uplands and lowlands based on slope and image characteristics. They were further subdivided into physiographic/image interpretation units based on image characteristics. The image interpretation legend for physiography is given below.

## Image Interpretation Legend for Physiography

### G- Granite Gneiss Landscape

G1	Hills/ Ridges/ Mounds
G11	Summits
G12	Side slopes
G121	Side slopes with dark grey tones
G2	Uplands
G21	Summits
G22	Gently sloping uplands
G221	Gently sloping uplands, yellowish green (eroded)
G222	Gently sloping uplands, yellowish white (severely eroded)
G23	Very gently sloping uplands
G231	Very gently sloping uplands, yellowish green
G232	Very gently sloping uplands, medium green and pink
G233	Very gently sloping uplands, pink and green (scrub land)
G234	Very gently sloping uplands, medium greenish grey
G235	Very gently sloping uplands, yellowish white (eroded)
G236	Very gently sloping uplands, dark green
G237	Very gently sloping uplands, medium pink (coconut garden)
G238	Very gently sloping uplands, pink and bluish white (eroded)
G24	Valleys/ lowlands
G241	Valleys, pink tones
G242	Valleys gray mixed with pink tones






Fig 3.1 Scanned and Digitized Cadastral map of Kotageri-2 Microwatershed




Fig.3.2 Satellite Image of Kotageri-2 Microwatershed




Fig.3.3 Cadastral map overlaid on IRS PAN+LISS IV merged imagery of Kotageri-2 Microwatershed

### 3.3 Field Investigation

The field boundaries and survey numbers given on the cadastral sheet were located on the ground by following permanent features like roads, cart tracks, *nallas*, streams, tanks etc., and wherever changes were noticed, they were incorporated on the microwatershed cadastral map. Preliminary traverse of the microwatershed was carried out with the help of cadastral map, imagery and toposheets. While traversing, landforms and physiographic units identified were checked and preliminary soil legend was prepared by studying soils at few selected places. Then, intensive traversing of each physiographic unit like hills, ridges, uplands and valleys was carried out. Based on the variability observed on the surface, transects (Fig. 3.4) were selected across the slope covering all the landform units in the microwatershed (Natarajan and Dipak Sarkar, 2010).




Fig: 3.4. Location of profiles in a transect

In the selected transect, soil profiles were located (Fig. 3.4) at closely spaced intervals to take care of any change in the land features like break in slope, erosion, gravel, stones etc. In the selected sites, profiles (vertical cut showing the soil layers from surface to the rock) were opened upto 200 cm or to the depth limited by rock or hard substratum and studied in detail for all their morphological and physical characteristics. The soil and site characteristics were recorded for all profile sites on a standard proforma as per the guidelines given in USDA Soil Survey Manual (Soil Survey Staff, 2012). Apart from the transect study, profiles were also studied at random, almost like in a grid pattern, outside the transect areas.

Based on the soil characteristics, the soils were grouped into different soil series. Soil series is the most homogeneous unit having similar horizons and properties and behaves similarly for a given level of management. Soil depth, texture, colour, kind of horizon and horizon sequence, calcareousness, amount and nature of gravel present, nature of substratum etc, were used as the major differentiating characteristics for identifying soil series occurring in the area. The differentiating characteristics used for identifying the soil series are given in Table 3.1. Based on the above characteristics, 8 soil series were identified in the Kotageri-2 microwatershed.

**Table 3.1 Differentiating Characteristics used for identifying Soil Series**  
(Characteristics are of Series Control Section)

<b>Soils of Granite gneiss Landscape</b>							
<b>Sl. no</b>	<b>Soil Series</b>	<b>Depth (cm)</b>	<b>Colour (moist)</b>	<b>Texture</b>	<b>Gravel (%)</b>	<b>Horizon sequence</b>	<b>Calcareousness</b>
1	BDP (Baddeppalli)	<25	7.5YR 3/2,3/4 5YR 3/4	scl	<15	Ap-AC	es
2	KKR (Kakalawar)	<25	7.5YR 4/3 10YR 6/3	sl	10-15	Ap-AC	-
3	BDL (Badiyala)	25-50	7.5YR2.5/3,2.5/ 3/3,10YR 3/4,4/3	sl	<15	Ap-Bw	e
4	DSB (Dastharabad)	25-50	7.5YR 3/3	g c	35-60	Ap-Bt-Cr	-
5	JNK (Jinkera)	50-75	10YR 3/1,3/2 7.5YR 3/4	scl	<15	Ap-Bw	e
6	GWD (Gowdagera)	75-100	10YR 3/1,3/2,4/2	scl	<15	Ap-Bw	es
7	SHT (Shettalli)	75-100	10YR 3/1	gsc	15-35	Ap-Bw	e
8	PGP (Poglapur)	75-100	5YR 4/6,3/3 2.5 YR3/4, 3/6	sc	<15	Ap-Bt	-

### 3.4 Soil Mapping

The area under each soil series was further separated into soil phases and their boundaries delineated on the cadastral map based on the variations observed in the texture of the surface soil, slope, erosion, presence of gravel, stoniness etc. A soil phase is a subdivision of soil series based mostly on surface features that affect its use and management. The soil mapping units are shown on the map (Fig.3.5) in the form of symbols. During the survey many profile pits, few minipits and a few auger bores representing different landforms occurring in the microwatershed were studied. In addition to the profile study, spot observations in the form of minipits, road cuts, terrace cuts etc., were studied to validate the soil boundaries on the soil map.

The soil map shows the geographic distribution of 11 mapping units representing 8 soil series occurring in the microwatershed. The soil map unit (soil legend) description is presented in Table 3.2. The soil phase map (management units) shows the distribution of 11 soil phases mapped in the microwatershed. Each mapping unit (soil phase) delineated on the map has similar soil and site characteristics. In other words, all the farms or survey numbers included in one phase will have similar management needs and have to be treated accordingly.

### 3.5 Land Management Units

The 11 soil phases identified and mapped in the microwatershed were grouped into 5 Land Management Units (LMU's) for the purpose of preparing a Proposed Crop Plan for sustained development of the microwatershed. The database (soil phases)



generated under LRI was utilized for identifying Land Management Units (LMU's) based on the management needs. One or more than one soil site characteristic having influence on the management have been chosen for identification and delineation of LMUs. For Kotageri-2 microwatershed, five soil and site characteristics, namely soil depth, soil texture, slope, erosion and gravel content have been considered for defining LMUs. The land use classes are expected to behave similarly for a given level of management.

### 3.6 Laboratory Characterization

Soil samples were collected from representative master profiles for laboratory characterization by following the methods outlined in the Laboratory Manual (Sarma *et al.*, 1987). Surface soil samples collected from farmer's fields (28 samples) for fertility status (major and micronutrients) at 320 m grid interval in the year 2019 were analyzed in the laboratory (Katyal and Rattan, 2003). By linking the soil fertility data to the survey numbers through GIS, soil fertility maps were generated by using Kriging method for the microwatershed.

**Table 3.2 Soil map unit description of Kotageri-2 Microwatershed**

*Soil map unit No.	Soil Series	Soil Phase	Mapping Unit Description	Area in ha (%)
	BDP		Baddeppalli soils are very shallow (<25 cm), well drained, have dark brown to dark reddish brown, calcareous sandy clay loam soils occurring on very gently sloping uplands under cultivation	<b>132 (43.62)</b>
119		BDPiB3	Sandy clay surface, slope 1-3%, severe erosion	132 (43.62)
	KKR		Kakalawar soils are very shallow (<25 cm), well drained, have dark brown sandy loam soils occurring on very gently sloping uplands under cultivation	<b>17 (5.77)</b>
175		KKRcB2	Sandy loam surface, slope 1-3%, moderate erosion	17 (5.77)
	BDL		Badiyala soils are shallow (25-50 cm), well drained, have dark brown to very dark brown and dark yellowish brown, slightly calcareous sandy loam soils occurring on very gently to gently sloping uplands under cultivation	<b>67 (22.28)</b>
3		BDLbC3	Loamy sand surface, slope 3-5%, severe erosion	12 (4.05)
4		BDLhB2	Sandy clay loam surface, slope 1-3%, moderate erosion	23 (7.72)
162		BDLhB2g1	Sandy clay loam surface, slope 1-3%, moderate erosion, gravelly (15-35%)	26 (8.47)
5		BDLiB2	Sandy clay surface, slope 1-3%, moderate erosion	6 (2.04)
	DSB		Dastharabad soils are shallow (25-50 cm), well drained, have dark brown to very dark brown, gravelly	<b>15 (4.92)</b>

*Soil map unit No.	Soil Series	Soil Phase	Mapping Unit Description	Area in ha (%)
			clay soils occurring on very gently to gently sloping uplands under cultivation	
121		DSBcB2	Sandy loam surface, slope 1-3%, moderate erosion	15 (4.92)
	JNK		Jinkera soils are moderately shallow (50-75 cm), well drained, have dark brown to very dark grayish brown, slightly calcareous sandy clay loam soils occurring on very gently sloping uplands under cultivation	<b>18</b> <b>(5.87)</b>
22		JNKiB2	Sandy clay surface, slope 1-3%, moderate erosion	18 (5.87)
	GWD		Gowdagera soils are moderately deep (75-100 cm), moderately well drained, have dark grayish brown to very dark grayish brown, calcareous, sodic sandy clay loam soils occurring on very gently sloping uplands under cultivation	<b>1</b> <b>(0.17)</b>
127		GWDmB2	Clay surface, slope 1-3%, moderate erosion	1 (0.17)
	SHT		Shettalli soils are moderately deep (75-100 cm), well drained, have very dark gray, gravelly sandy clay soils occurring on very gently sloping uplands under cultivation	<b>15</b> <b>(4.98)</b>
36		SHTbB2	Sandy clay loam surface, slope 1-3%, moderate erosion	15 (4.98)
	PGP		Poglapur soils are moderately deep (75-100 cm), well drained, have brown to dark reddish brown and yellowish red, sandy clay soils occurring on very gently sloping uplands under cultivation	<b>0.40</b> <b>(0.13)</b>
40		PGPcB2	Sandy loam surface, slope 1-3%, moderate erosion	0.40 (0.13)
1000		Others	Habitation and Water body	37 (12.26)




Fig 3.5 Soil Phase or Management Units - Kotageri-2 Microwatershed



## THE SOILS

Detailed information pertaining to the nature, extent and their distribution of different kinds of soils occurring in Kotageri-2 microwatershed is provided in this chapter. The microwatershed area has been identified as granite gneiss landscape based on geology. In all, 8 soil series are identified. Soil formation is the result of the combined effect of environmental and terrain factors that are reflected in soil morphology. In the granite gneiss landscape, it is by parent material, relief and climate.

A brief description of each of the 8 soil series identified followed by 11 soil phases (management units) mapped under each series are furnished below. The physical and chemical characteristics of soil series identified in Kotageri-2 microwatershed are given in Table 4.1 along with soil classification. The soils in any one map unit differ from place to place in their depth, texture, slope, gravelliness, erosion or any other site characteristic that affect management. The soil phase map can be used for identifying the suitability of areas for growing specific crops or for other alternative uses and also for deciding the type of conservation structures needed. The detailed information on soil and site-characteristics like soil depth, surface soil texture, slope, erosion, gravelliness, AWC, LCC etc, with respect to each of the soil phase identified is given village/survey number wise for the microwatershed in Appendix-I.

### 4.1 Soils of granite gneiss landscape

In this landscape, 8 soil series are identified and mapped. BDP series occupies maximum area of 132 ha (44%) followed by BDL 67 ha (22%), JNK 18 ha (6%), KKR 17 ha (6%), SHT 15 ha (5%), DSB 15 ha (5%), GWD 1 ha (<1%) and PGP 5 ha (<1%). Brief description of each series identified and number of soil phases mapped is given below.

**4.1.1 Baddeppalli (BDP) Series:** Baddeppalli soils are very shallow (<25cm), well drained, have dark brown to dark reddish brown, calcareous sandy clay loam soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation. The Baddeppalli series has been classified as a member of the loamy, mixed (calcareous), isohyperthermic family of Lithic Ustorthents.

The thickness of the soil is less than 25 cm. Its colour is in 7.5 YR and 5 YR hue with value 3 and chroma 2 to 4. The texture varies from sandy clay loam to sandy clay and is calcareous. The available water capacity is very low (<50 mm/m). One phase was identified and mapped.



Landscape and Soil Profile characteristics of Baddeppalli (BDP) Series

**4.1.2 Kakalawar (KKR) Series:** Kakalawar soils are very shallow (<25cm), well drained, have dark brown to light brown, sandy loam soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation. The Kakalawar series has been classified as a member of the mixed, isohyperthermic family of Lithic Ustipsamments.

The thickness of the soil is less than 25 cm. Its colour is in 10 YR and 7.5 YR hue with value 4 to 6 and chroma 3 to 4. The texture varies from loamy sand to sand. The available water capacity is very low (<50 mm/m). One soil phase was identified and mapped.



Landscape and Soil Profile characteristics of Kakalawar (KKR) Series



**4.1.3 Badiyala (BDL) Series:** Badiyala soils are shallow (25-50 cm), well drained, have very dark brown to dark yellow brown and dark brown, slightly calcareous sandy loam soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation. The Badiyala series has been classified as a member of the coarse-loamy, mixed, isohyperthermic family of Fluventic Haplustepts.

The thickness of the solum ranges from 28 to 50 cm. The thickness of A horizon ranges from 4 to 12 cm. Its colour is in 10YR hue with value 3 to 4 and chroma 3 to 4. The texture is loamy sand, sandy clay loam and sandy clay. The thickness of B horizon ranges from 27 to 45 cm. Its colour is in 10 YR and 7.5 YR hue with value 2 to 4 and chroma 3 to 4. Its texture is sandy loam to sandy clay loam and is slightly calcareous. The available water capacity is very low (<50mm/m). Four phases were identified and mapped.



Landscape and Soil Profile characteristics of Badiyala (BDL) Series

**4.1.4 Dastharabad (DSB) Series:** Dastharabad soils are shallow (25-50 cm), well drained, have dark brown, gravelly clay soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation. The Dastharabad series has been classified as a member of the clayey-skeletal, mixed, isohyperthermic family of (Paralithic) Haplustalfs.

The thickness of the solum ranges from 28 to 50 cm. The thickness of A horizon ranges from 9 to 14 cm. Its colour is in 10 YR and 7.5 YR hue with value and chroma of 3 to 4. The texture varies from sandy loam to sandy clay. The thickness of B horizon ranges from 28 to 40 cm. Its colour is in 7.5 YR hue with value 3 and chroma 3 to 4. The texture is sandy clay to clay with 35-60 per cent gravel. The available water capacity is very low (<50 mm/m). One phase was identified and mapped.



Landscape and Soil Profile characteristics of Dastharabad (DSB) Series

**4.1.5 Jinkera (JNK) Series:** Jinkera soils are moderately shallow (50-75 cm), well drained, have very dark gray to very dark grayish brown and dark brown, slightly calcareous sandy clay loam soils. They are developed from weathered granite gneiss and occur on very gently sloping uplands under cultivation. The Jinkera series has been classified as a member of the fine-loamy, mixed, isohyperthermic family of Typic Haplustepts.

The thickness of the solum ranges from 51-75 cm. Thickness of A horizon ranges from 6 to 11 cm. Its colour is in hue 10 YR and 7.5 YR with value and chroma of 3 to 4. The texture varies from sandy loam to sandy clay. The thickness of B horizon ranges from 53 to 66 cm. Its colour is in 10 YR and 7.5 YR hue with value and chroma of 2 to 4. The texture varies from sandy clay loam to sandy clay and is slightly calcareous. The available water capacity is low (51-100 mm/m). One phase was identified and mapped.



Landscape and Soil Profile characteristics of Jinkera (JNK) Series



**4.1.6 Gowdagera (GWD) Series:** Gowdagera soils are moderately deep (75-100 cm), moderately well drained, have very dark gray to dark grayish brown, calcareous sodic sandy clay loam soils. They are developed from weathered granite gneiss and occur on very gently sloping uplands under cultivation. The Gowdagera series has been classified as a member of the fine-loamy, mixed (calcareous), isohyperthermic family of Typic Haplustepts.

The thickness of the solum ranges from 76 to 100 cm. The thickness of A horizon ranges from 8 to 16 cm. Its colour is in hue 10 YR with value 3 to 4 and chroma 2 to 4. Its texture varies from sandy loam to sandy clay loam. The thickness of B horizon ranges from 61 to 91 cm. Its colour is in hue 10 YR with value 2 to 4 and chroma 1 to 4. Its texture is sandy clay loam to sandy clay and is calcareous sodic soils. The available water capacity is medium (101-150 mm/m). One phase was identified and mapped.



Landscape and Soil Profile characteristics of Gowdagera (GWD) Series

**4.1.7 Shettalli (SHT) Series:** Shettalli soils are moderately deep (75-100 cm), well drained, have very dark gray, slightly calcareous gravelly sandy clay soils. They are developed from weathered granite gneiss and occur on very gently sloping uplands under cultivation. The Shettalli series has been classified as a member of the fine, mixed, isohyperthermic family of Typic Haplustepts.

The thickness of the solum ranges from 78 to 100 cm. The thickness of A horizon ranges from 7 to 12 cm. Its colour is in hue 7.5 YR with value and chroma of 3 to 4. Its texture varies from sandy loam to sandy clay with 20 per cent gravel. The thickness of B horizon ranges from 68 to 92 cm. Its colour is in hue 7.5 YR with value 2 to 4 and chroma 1 to 3. Its texture is sandy clay with 15-35 per cent gravel and is slightly calcareous. The available water capacity is low (51-100 mm/m). One phase was identified and mapped.



Landscape and Soil Profile characteristics of Shettalli (SHT) Series

**4.1.8 Poglapur (PGP) Series:** Poglapur soils are moderately deep (75-100 cm), well drained, have dark brown to dark reddish brown and yellowish red sandy clay red soils. They have developed from granite gneiss and occur on very gently sloping uplands under cultivation. The Poglapur series has been classified as a member of the fine, mixed, isohyperthermic family of Rhodic Paleustalfs.

The thickness of the solum ranges from 78 to 100 cm. The thickness of A horizon ranges from 8 to 17 cm. Its colour is in 7.5 YR hue with value 3 and chroma 3 to 4. Its texture varies from loamy sand to sandy clay loam and sandy clay. The thickness of B horizon ranges from 65 to 92 cm. Its colour is in 2.5 YR and 5 YR hue with value 2 to 4 and chroma 2 to 6. Its texture is sandy clay and clay. The available water capacity is medium (101-150 mm/m). One phase was identified and mapped.



Landscape and Soil Profile characteristics of Poglapur (PGP) Series

**Table: 4.1 Physical and Chemical Characteristics of Soil Series identified in Kotageri-2 microwatershed**

**Soil Series:** Baddeppalli (BDP) **Pedon:** R-11

**Location:** 16°43'84.4"N 77°14'06.4"E, Halagera village, Yadgir hobli, Yadgir taluk and district

**Analysis at:** NBSS&LUP, Regional Centre, Bengaluru **Classification:** Loamy, mixed (calcareous), isohyperthermic Lithic Ustorthents

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-16	Ap	58.67	17.02	24.31	19.03	13.74	9.62	10.57	5.71	<15	scl	16.19	8.18

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO <sub>3</sub>	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP						
	Water	CaCl <sub>2</sub>	M KCl				dS m <sup>-1</sup>	%	%	Ca	Mg					K	Na	Total	cmol kg <sup>-1</sup>	%	%
0-16	8.58	-	-	0.262	1.60	7.67	-	-	0.24	0.06	-	18.10	0.74	100	0.35						

*Contd...*

**Soil Series:** Kakalawar (KKR), Pedon: R-7

**Location:** 16°50'25.9"N 77°15'97.1"E, Yampada village, Gurumitkal hobli, Yadgir taluk and district

**Analysis at:** NBSS&LUP, Regional Centre, Bengaluru **Classification:** Mixed, isohyperthermic Lithic Ustipsamments

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-22	Ap	83.81	10.37	5.82	17.31	20.65	17.91	5.67	22.27	10-20	ls	9.77	4.65

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO <sub>3</sub>	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP			
	Water	CaCl <sub>2</sub>	M KCl				dS m <sup>-1</sup>	%	%	Ca	Mg					K	Na	Total
										cmol kg <sup>-1</sup>						%	%	
0-22	5.85	-	-	0.027	0.19	-	0.72	0.21	0.62	0.03	1.58	2.6	0.45	60.90	1.17			

*Contd...*



**Soil Series:** Badiyala (BDL) **Pedon:** R-5

**Location:** 16°37'10.0"N 77°20'21.5", Gudalagunta village, Balichakra hobli, Yadgir taluk and district

**Analysis at:** NBSS&LUP, Regional Centre, Bengaluru **Classification:** Coarse-loamy, mixed, isohyperthermic Fluventic Haplustepts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-12	Ap	87.13	7.04	5.83	10.03	24.32	23.61	23.51	5.67	<15	ls	6.27	2.44
12-28	Bw	64.63	13.30	22.07	6.74	13.07	22.30	17.01	5.50	<15	scl	16.34	7.83
28-50	BC	73.11	12.02	14.87	3.93	16.03	26.89	18.41	7.86	<15	sl	12.94	5.47

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO <sub>3</sub>	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP			
	Water	CaCl <sub>2</sub>	M KCl				dS m <sup>-1</sup>	%	%	Ca	Mg					K	Na	Total
										cmol kg <sup>-1</sup>								
0-12	6.20	-	-	0.074	1.00	0.00	2.80	0.98	0.14	0.01	3.92	4.20	0.72	93	0.20			
12-28	9.04	-	-	0.253	0.80	3.20	-	-	0.16	0.69	-	16.90	0.77	100	4.09			
28-50	9.41	-	-	0.364	1.10	3.60	-	-	0.16	1.39	-	11.10	0.75	100	12.52			

*Contd...*

**Soil Series:** Dastharabad (DSB) **Pedon:** R-17

**Location:** 16°31' 98.6"N 77°22'93.0"E, Duppalli village, Sydhapura hobli, Yadgir taluk and district

**Analysis at:** NBSS&LUP, Regional Centre, Bengaluru **Classification:** Clayey-skeletal, mixed, isohyperthermic (Paralithic) Haplustalfs

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-6	Ap	90.51	4.84	4.64	7.06	8.07	37.24	26.03	12.11	35	s	5.32	1.59
6-17	Bt1	49.11	8.08	42.81	10.67	15.44	10.00	8.44	4.56	20	sc	20.68	13.16
17-43	Bt2	39.54	2.84	57.63	12.89	9.14	7.71	6.83	2.97	50	c	26.69	18.50

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO <sub>3</sub>	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP			
	Water	CaCl <sub>2</sub>	M KCl				dS m <sup>-1</sup>	%	%	Ca	Mg					K	Na	Total
										cmol kg <sup>-1</sup>						%	%	
0-6	5.93	-	-	0.04	0.67	0.00	2.00	0.54	0.07	0.01	2.61	3.60	0.78	73	0.14			
6-17	7.31	-	-	0.110	0.91	0.91	11.19	3.37	0.12	0.49	15.00	15.20	0.36	100	3.22			
17-43	6.64	-	-	0.048	0.76	0.00	18.81	5.57	0.23	0.09	24.70	24.90	0.43	99	0.38			

*Contd...*

**Soil Series:** Jinkera (JNK) **Pedon:** R-1

**Location:** 16°45'13.5"N 77°10'59.8"E, Varkanahalli village, Yadgir hobli, Yadgir taluk and district

**Analysis at:** NBSS&LUP, Regional Centre, Bengaluru **Classification:** Fine-loamy, mixed, isohyperthermic Typic Haplustepts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-15	Ap	66.84	13.62	19.54	12.15	21.22	11.23	12.56	9.68	10	sl	14.42	7.70
15-38	Bw1	59.08	12.11	28.81	12.53	12.42	17.85	8.77	7.52	20	scl	18.21	12.23
38-52	Bw2	68.21	11.68	20.11	17.90	21.81	10.60	10.80	7.10	10	scl	14.54	8.96

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO <sub>3</sub>	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP
	Water	CaCl <sub>2</sub>	M KCl				Ca	Mg	K	Na	Total				
				dS m <sup>-1</sup>	%	%	cmol kg <sup>-1</sup>							%	%
0-15	8.42	-	-	0.148	0.70	0.65	-	-	0.15	0.03	-	14.50	0.74	100	0.18
15-38	8.38	-	-	0.226	0.31	2.21	-	-	0.09	0.23	-	21.70	0.75	100	1.05
38-52	8.40	-	-	0.195	0.25	1.17	-	-	0.07	0.19	-	15.90	0.79	100	1.23

*Contd...*

**Soil Series:** Gowdagera (GWD) **Pedon:** R-13

**Location:** 16°38'24.4"N 77°21'24.0"E, Madhawara village, Balichakara hobli, Yadgir taluk and district

**Analysis at:** NBSS&LUP, Regional Centre, Bengaluru **Classification:** Fine-loamy, mixed (calcareous), isohyperthermic Typic Haplustepts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-18	Ap	79.61	13.94	6.45	14.17	17.53	23.65	17.02	7.24	-	ls	11.36	3.86
18-42	BW1	69.09	10.58	21.06	10.54	16.58	22.01	14.43	5.53	-	scl	31.62	12.30
42-81	Bw2	51.37	13.51	35.60	7.59	10.55	16.24	11.60	5.38	-	sc	67.57	26.89

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO <sub>3</sub>	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP						
	Water	CaCl <sub>2</sub>	M KCl				dS m <sup>-1</sup>	%	%	Ca	Mg					K	Na	Total	cmol kg <sup>-1</sup>	%	%
0-18	9.89	-	-	0.74	0.66	1.20	-	-	0.18	3.63	-	8.35	1.29	100	17.40						
18-42	10.82	-	-	1.60	0.27	5.76	-	-	0.19	19.23	-	15.84	0.75	100	40.17						
42-81	10.83	-	-	2.30	0.27	7.80	-	-	0.40	26.71	-	26.54	0.75	100	40.27						

*Contd...*



**Soil Series:** Shettalli (SHT) **Pedon:** R-14

**Location:** 16°47'21.1"N 77°04'91.1"E, Thumakura village, Yadgir hobli, Yadgir taluk and district

**Analysis at:** NBSS&LUP, Regional Centre, Bengaluru **Classification:** Fine, mixed, isohyperthermic Typic Haplustepts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-14	Ap	74.39	10.89	14.73	5.64	8.30	21.00	28.89	10.55	50	sl	12.58	4.51
14-35	Bw1	54.37	14.73	30.90	3.58	5.90	15.38	21.71	7.80	25	scl	20.37	10.92
35-63	Bw2	41.16	20.63	38.21	1.71	1.71	10.61	13.61	13.50	30	cl	24.34	15.03
63-83	Bw3	36.96	21.52	41.51	4.31	5.28	8.94	12.39	6.03	35	c	24.76	16.17

Depth (cm)	pH (1:2.5)			E.C. (1:2.5) dS m <sup>-1</sup>	O.C. %	CaCO <sub>3</sub> %	Exchangeable bases					CEC	CEC/Clay	Base saturation %	ESP %
	Water	CaCl <sub>2</sub>	M KCl				Ca	Mg	K	Na	Total				
0-14	7.26	-	-	0.199	0.91	0.13	-	-	0.28	0.09	-	10.60	0.72	100	0.86
14-35	7.05	-	-	0.051	0.80	1.17	-	-	0.12	0.09	-	18.20	0.59	100	0.48
35-63	7.67	-	-	0.238	0.70	2.86	-	-	0.14	0.16	-	24.40	0.64	100	0.64
63-83	8.67	-	-	0.142	0.20	12.48	-	-	0.13	0.23	-	27.40	0.66	100	0.84

Contd...

**Soil Series:** Poglapur (PGP) **Pedon:** R-6

**Location:** 16°34'45.2"N 77°10'96.4"E, Anura B village, Sydhapura hobli, Yadgir taluk and district

**Analysis at:** NBSS&LUP, Regional Centre, Bengaluru **Classification:** Fine, mixed, isohyperthermic Rhodic Paleustalfs

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-15	Ap	91.81	4.70	3.49	17.80	30.23	15.57	20.93	7.28	-	s	4.94	2.29
15-50	Bt1	46.83	4.99	48.17	11.92	16.22	8.59	6.77	3.33	10	sc	24.59	17.37
50-90	Bt2	45.81	4.73	49.46	17.10	14.09	6.45	5.16	3.01	15	sc	24.44	16.57
90-125	Bt3	58.92	5.86	35.22	28.51	10.45	10.98	5.49	3.48	15	sc	21.73	10.30

Depth (cm)	pH (1:2.5)			E.C. (1:2.5) dS m <sup>-1</sup>	O.C. %	CaCO <sub>3</sub> %	Exchangeable bases					CEC	CEC/Clay	Base saturation %	ESP %
	Water	CaCl <sub>2</sub>	M KCl				Ca	Mg	K	Na	Total				
0-15	6.83	-	-	0.210	0.76	0.00	1.79	0.88	0.41	0.09	3.16	3.15	0.90	100	2.83
15-50	6.20	-	-	0.105	0.48	0.00	12.27	4.45	0.30	0.39	17.40	17.54	0.36	99	2.22
50-90	6.23	-	-	0.080	0.40	0.00	11.51	3.92	0.28	0.37	16.09	17.33	0.35	93	2.16
90-125	6.49	-	-	0.068	0.20	0.00	11.19	3.62	0.27	0.40	15.49	17.43	0.49	89	2.29

## INTERPRETATION FOR LAND RESOURCE MANAGEMENT

The most important soil and site characteristics that affect the land use and conservation needs of an area are land capability, soil depth, soil texture, coarse fragments, available water capacity, soil slope, soil erosion, soil reaction etc. These are interpreted from the data base generated through land resource inventory and several thematic maps are generated. These would help in identifying the areas suitable for growing crops and, soil and water conservation measures and structures needed thus helping to maintain good soil health for sustained crop production. The various interpretative and thematic maps generated are described below.

### 5.1 Land Capability Classification

Land capability classification is an interpretative grouping of soil map units (soil phases) mainly based on inherent soil characteristics, external land features and environmental factors that limit the use of land for agriculture, pasture, forestry, or other uses on a sustained basis (IARI, 1971). The land and soil characteristics used to group the land resources in an area into various land capability classes, subclasses and units are

*Soil Characteristics:* Depth, texture, gravelliness, calcareousness.

*Land characteristics:* Slope, erosion, drainage, rock outcrops.

*Climate:* Total rainfall and its distribution, and length of crop growing period.

The Land capability classification system is divided into land capability classes, subclasses and units based on the level of information available. Eight land capability classes are recognized. They are

*Class I:* They are very good lands that have no limitations or very few limitations that restrict their use.

*Class II:* They are good lands that have minor limitations and require moderate conservation practices.

*Class III:* They are moderately good lands that have moderate limitations that reduce the choice of crops or that require special conservation practices.

*Class IV:* They are fairly good lands that have very severe limitations that reduce the choice of crops or that require very careful management.

*Class V:* Soils in these lands are not likely to erode, but have other limitations like wetness that are impractical to remove and as such not suitable for agriculture, but suitable for pasture or forestry with minor limitations.

*Class VI:* The lands have severe limitations that make them generally unsuitable for cultivation, but suitable for pasture or forestry with moderate limitations.

*Class VII:* The lands have very severe limitations that make them unsuitable for cultivation, but suitable for pasture or forestry with major limitations.

*Class VIII:* Soil and other miscellaneous areas (rock lands) that have very severe limitations that nearly preclude their use for any crop production, but suitable for wildlife, recreation and installation of wind mills.

The land capability subclasses are recognised based on the dominant limitations observed within a given land capability class. The subclasses are designated by adding a lower case letter like 'e', 'w', 's', or 'c' to the class numeral. The subclass "e" indicates that the main hazard is risk of erosion, "w" indicates drainage or wetness as a limitation for plant growth, "s" indicates shallow soil depth, coarse or heavy textures, calcareousness, salinity/alkalinity or gravelliness and "c" indicates limitation due to climate.

The land capability subclasses have been further subdivided into land capability units based on the kinds of limitations present in each subclass. Ten land capability units are used in grouping the soil map units. They are stony or rocky (0), erosion hazard (slope, erosion) (1), coarse texture (sand, loamy sand, sandy loam) (2), fine texture (cracking clay, silty clay) (3), slowly permeable subsoil (4), coarse underlying material (5), salinity/alkali (6), stagnation, overflow, high ground water table (7), soil depth (8) and fertility problems (9). The capability units thus identified have similar soil and land characteristics that respond similarly to a given level of management. The soils of the microwatershed have been classified upto land capability subclass level.

The 11 soil map units identified in the Kotageri-2 microwatershed are grouped under 3 land capability classes and 3 subclasses. An area about 265 ha (88%) in the microwatershed is suitable for agriculture and others cover an area of 37 ha (12%) in the microwatershed. (Fig. 5.1).




Fig. 5.1 Land Capability map of Kotageri-2 Microwatershed

Good lands (Class II) cover an area of 33 ha (11%) and are distributed in the northern and southeastern part of the microwatershed. They have minor limitations of soil and erosion. Moderately good lands (Class III) cover an area of about 70 ha (23%) and are distributed in the northern, central and southeastern part of the microwatershed. They have moderate limitations of soil and erosion. Fairly good lands (Class IV) cover an area of about 162 ha (54%) and are distributed in the major part of the microwatershed. They have moderate limitations of soil and erosion.

## 5.2 Soil Depth

Soil depth refers to the depth of the soil occurring above the parent material or hard rock. The depth of the soil determines the effective rooting depth for plants and in accordance with soil texture, mineralogy and gravel content, the capacity of the soil column to hold water and nutrient availability. Soil depth is one of the most important soil characteristic that is used in differentiating soils into different soil series. The soil depth classes used in identifying soils in the field are very shallow (<25 cm), shallow (25-50 cm), moderately shallow (50-75 cm), moderately deep (75-100 cm), deep (100-150 cm) and very deep (>150 cm). They were used to classify the soils into different depth classes and a soil depth map was generated. The area extent and their geographical distribution in the microwatershed is given in Fig. 5.2.




Fig. 5.2 Soil Depth map of Kotageri-2 Microwatershed

Very shallow (<25cm) soils cover an area of 149 ha (49%) and are distributed in the major part of the microwatershed. Shallow (25-50 cm) soils cover an area of 82 ha (27%) and are distributed in the central, southeastern and northwestern part of the microwatershed. Moderately shallow (50-75 cm) soils cover an area of 18 ha (6%) and are distributed in the southeastern part of the microwatershed. Moderately deep (75-100 cm) soils cover an area of 16 ha (5%) and are distributed in the northern part of the microwatershed.

Problem soils cover about 231 ha (76%) area where short duration crops can be grown and probability of crop failure is high.

### 5.3 Surface Soil Texture

Texture is an expression to indicate the coarseness or fineness of the soil as determined by the relative proportion of primary particles of sand, silt and clay. It has a direct bearing on the structure, porosity, adhesion and consistence. The surface layer of a soil to a depth of about 25 cm is the layer that is most used by crops and plants. The surface soil textural class provides a guide to understanding soil-water retention and availability, nutrient holding capacity, infiltration, workability, drainage, physical and chemical behaviour, microbial activity and crop suitability. The textural classes used for LRI were used to classify and a surface soil texture map was generated. The area extent and their geographical distribution in the microwatershed is shown in Figure 5.3.




Fig. 5.3 Surface Soil Texture map of Kotageri-2 Microwatershed

An area of about 12 ha (4%) is sandy and is distributed in the northwestern part of the microwatershed. An area of 97 ha (32%) has soils that are loamy at the surface and occur in the central, southeastern, eastern and northern part of the microwatershed. An area of 156 ha (52%) has soils that are clayey at the surface and occur in the major part of the microwatershed.

Major area of 253 ha (84%) the microwatershed is most productive with respect to surface soil texture. The clayey soils (52%) have high potential for soil-water retention and availability, and nutrient retention and availability, but have more problems of drainage, infiltration, workability and other physical problems. The other productive lands are loamy soils (32%) which also have high potential for soil-water retention and nutrient availability but have no drainage or other physical problems. The sandy soils (4%) are problematic but productive for root and tuber crops, but these soils have the major limitation of moisture and nutrient retention capacity, hence frequent and shallow irrigation with balanced fertilizer application is to be followed in order to get better crop yields.

#### 5.4 Soil Gravelliness

Gravel is the term used for describing coarse fragments between 2 mm and 7.5 cm diameter and stones for those between 7.5 cm and 25 cm. The presence of gravel and stones in soil reduces the volume of soil responsible for moisture and nutrient storage, drainage, infiltration and runoff, and hinders plant growth by impeding root growth and seedling emergence, intercultural operations and farm mechanization. The gravelliness



classes used in LRI were used to classify the soils and using these classes a gravelliness map was generated. The area extent and their geographic distribution in the microwatershed is shown in Figure 5.4.




Fig. 5.4 Soil Gravelliness map of Kotageri-2 Microwatershed

An area of about 239 ha (79%) is non gravelly (<15%) soils and are distributed in the major part of the microwatershed. About 26 ha (8%) is gravelly (15-35%) soils and are distributed in the northwestern part of the microwatershed.

The most productive soils (79%) that are non gravelly (<15%), where all climatically adapted long duration crops can be grown.

### 5.5 Available Water Capacity

The soil available water capacity (AWC) is estimated based on the ability of the soil column to retain water between the tensions of 0.33 and 15 bar in a depth of 100 cm or the entire solum if the soil is shallower. The AWC of the soils (soil series) as estimated by considering the soil texture, mineralogy, soil depth and gravel content (Sehgal *et al.*, 1990) and accordingly the soil map units were grouped into five AWC classes *viz*, very low (<50 mm/m), low (50-100 mm/m), medium (100-150 mm/m), high (150-200 mm/m) and very high (>200 mm/m) and using these values, an AWC map was generated. The area extent and their geographic distribution of different AWC classes in the microwatershed is given in Figure 5.5.




Fig. 5.5 Soil Available Water Capacity map of Kotageri-2 Microwatershed

An area of about 33 ha (11%) and 231 ha (77%) in the microwatershed has soils that are low (51-100 mm/m) and very low (<50mm/m) in available water capacity and are distributed in the major part of the microwatershed and about 1 ha (<1%) is medium (51-100 mm/m) in available water capacity and are distributed in the northern part of the microwatershed.

An area of 264 ha (88%) in the microwatershed has problematic with regard to available water capacity. Here, only short duration crops can be grown and the probability of crop failure is very high. These areas are best put to other alternative uses.

## 5.6 Soil Slope

Soil slope refers to the inclination of the surface of the land. It is defined by gradient, shape and length, and is an integral feature of any soil as a natural body. Slope is considered important in soil genesis, land use and land development. The length and gradient of slope influences the rate of runoff, infiltration, erosion and deposition. The soil map units were grouped into two slope classes and a slope map was generated showing the area extent and their geographic distribution in the microwatershed (Fig. 5.6).

An area of 253 ha (84%) falls under very gently sloping (1-3% slope) lands and are distributed in the major part of the microwatershed and about 12 ha (4%) falls under gently sloping (3-5% slope) lands and are distributed in the southwestern part of the microwatershed.




Fig. 5.6 Soil Slope map of Kotageri-2 Microwatershed

An area of 253 ha (84%) in the microwatershed has high potential in respect of soil slopes. In these areas, all climatically adapted annual and perennial crops can be grown without much soil and water conservation and other land development measures.

### 5.7 Soil Erosion

Soil erosion refers to the wearing away of the earth's surface by the forces of water, wind and ice involving detachment and transport of soil by raindrop impact. It is used for accelerated soil erosion resulting from disturbance of the natural landscape by burning, excessive grazing and indiscriminate felling of forest trees and tillage, all usually by man. The erosion classes showing an estimate of the current erosion status as judged from field observations in the form of rills, gullies or a carpet of gravel on the surface are recorded. Four erosion classes, viz, slight erosion (e1), moderate erosion (e2), severe erosion (e3) and very severe erosion (e4) are recognized. The soil map units were grouped into different erosion classes and a soil erosion map generated. The area extent and their spatial distribution in the microwatershed is given in Figure 5.7.

An area of 121 ha (40%) is moderately eroded (e2 class) lands and are distributed in the central, southeastern, northern and northwestern part of the microwatershed in the microwatershed and about 144 ha (48%) is severely eroded (e3 class) lands and are distributed in the major part of the microwatershed.

Entire cultivated area in the microwatershed has problematic because of moderate and severe erosion. For these areas, taking up soil and water conservation and other land development measures are needed.




Fig. 5.7 Soil Erosion map of Kotageri-2 Microwatershed



## **FERTILITY STATUS**

Soil fertility plays an important role in increasing crop yield. The adoption of high yielding varieties that require high amounts of nutrients has resulted in deficiency symptoms in crops and plants due to imbalanced fertilization and poor inherent fertility status as these areas are characterised by low rainfall and high temperatures. Hence, it is necessary to know the fertility (macro and micro nutrients) status of the soils of the watersheds for assessing the kind and amount of fertilizers required for each of the crop intended to be grown. For this purpose, the surface soil samples collected from the grid points (one soil sample at every 320 m interval) all over the microwatershed through land resource inventory in the year 2019 were analysed for pH, EC, organic carbon, available phosphorus and potassium, and for micronutrients like zinc, boron, copper, iron and manganese, and secondary nutrient sulphur.

Soil fertility data generated has been assessed and individual maps for all the nutrients for the microwatershed have been generated using Kriging method under GIS. The village/survey number wise fertility data for the microwatershed is given in Appendix-II.

### **6.1 Soil Reaction (pH)**

The soil analysis of the Kotageri-2 microwatershed for soil reaction (pH) showed that an area of 252 ha (83%) in the microwatershed is neutral (pH 6.5-7.3) and small area of 13 ha (4%) is slightly alkaline (pH 7.3-7.8) in soil reaction (fig.6.1).

### **6.2 Electrical Conductivity (EC)**

The Electrical Conductivity of the soils in the microwatershed area is  $<2 \text{ dS m}^{-1}$  (Fig 6.2) and as such the soils are non-saline.

### **6.3 Organic Carbon**

The soil organic carbon content (an index of available Nitrogen) in the soils of the microwatershed is low ( $<0.5\%$ ) in organic carbon about 207 ha (69%) and are distributed in the major part of the microwatershed. Medium (0.5-0.75%) in organic carbon in about 58 ha (19%) and are distributed in the southeastern, northern and northwestern part of the microwatershed and  $<1 \text{ ha}$  ( $<1\%$ ) is high ( $>0.75\%$ ) in organic carbon and are distributed in the minor part of the microwatershed (Fig. 6.3).






Fig.6.1 Soil Reaction (pH) map of Kotageri-2 Microwatershed




Fig.6.2 Electrical Conductivity (EC) map of Kotageri-2 Microwatershed






Fig.6.3 Soil Organic Carbon map of Kotageri-2 Microwatershed

#### 6.4 Available Phosphorus

Available phosphorus content is medium (23-57 kg/ha) covering an area of about 168 ha (56%) and occur in the major part of the microwatershed and high (>57 kg/ha) covering an area of about 97 ha (32%) and occur in the central, eastern and northern part of the microwatershed (Fig. 6.4).

#### 6.5 Available Potassium

Available potassium content is low (<145 kg/ha) covering an area of about 19 ha (6%) and occur in the southeastern and southwestern part of the microwatershed and medium (145-337 kg/ha) covering a maximum area of about 245 ha (81%) and occur in the major part of the microwatershed (Fig.6.5).

#### 6.6 Available Sulphur

Available sulphur is medium (10-20 ppm) in an area of about 186 ha (62%) and occur in the major part of the microwatershed and low (<10 ppm) covering an area of about 79 ha (26%) and occur in the northern, eastern and southeastern part of the microwatershed (Fig. 6.6).

### 6.7 Available Boron

Available boron content is medium (0.5-1.0 ppm) in an area of 45 ha (15%) and are distributed in the northern, southeastern, southwestern part of the microwatershed. An area of about 219 ha (73%) is low (<0.5 ppm) in available boron and are distributed in the major part of the microwatershed (Fig. 6.7).

### 6.8 Available Iron

Available iron content is sufficient (>4.5 ppm) in the entire cultivated area of the microwatershed (Fig 6.8).

### 6.9 Available Manganese

Available manganese content is sufficient (>1.0 ppm) in the entire cultivated area of the microwatershed (Fig 6.9).

### 6.10 Available Copper

Available copper content is sufficient (>0.2 ppm) in the entire cultivated area of the microwatershed area (Fig 6.10).




Fig.6.4 Soil Available Phosphorus map of Kotageri-2 Microwatershed




Fig.6.5 Soil Available Potassium map of Kotageri-2 Microwatershed




Fig.6.6 Soil Available Sulphur map of Kotageri-2 Microwatershed




Fig.6.7 Soil Available Boron map of Kotageri-2 Microwatershed




Fig.6.8 Soil Available Iron map of Kotageri-2 Microwatershed




Fig.6.9 Soil Available Manganese map of Kotageri-2 Microwatershed




Fig.6.10 Soil Available Copper map of Kotageri-2 Microwatershed



## 6.11 Available Zinc

Available zinc content is deficient (<0.6 ppm) covering a maximum area of 218 ha (72%) and are distributed in the major part of the microwatershed and sufficient (>0.6 ppm) an area of 47 ha (15%) and are distributed in the northern, central and southeastern part of the microwatershed (Fig 6.11).




Fig.6.11 Soil Available Zinc map of Kotageri-2 Microwatershed

## LAND SUITABILITY FOR MAJOR CROPS

The soil and land resource units (soil phases) of Kotageri-2 microwatershed were assessed for their suitability for growing food, fodder, fibre and other horticulture crops by following the procedure as outlined in FAO, 1976 and 1983. Crop requirements were developed for each of the crop from the available research data and also by referring to Naidu *et. al.* (2006) and Natarajan *et. al* (2015). The soil and land characteristics were matched with the crop requirement to arrive at the crop suitability. The soil and land characteristics (Table 7.1) and crop requirement (Table 7.2 to 7.30) are given at the end of the chapter. In FAO land suitability classification, two orders are recognized. Order S-Suitable and Order N-Not suitable. The orders have classes, subclasses and units. Order-S has three classes, Class S1-Highly Suitable, Class S2-Moderately Suitable and Class S3- Marginally Suitable. Order N has two classes, N1-Currently not Suitable and N2-Permanently not Suitable. There are no subclasses within the Class S1 as they will have very minor or no limitations for crop growth. Classes S2, S3 and N1 are divided into subclasses based on the kinds of limitations encountered. The limitations that affect crop production are ‘c’ for erratic rainfall and its distribution and length of growing period (LGP), ‘e’ for erosion hazard, ‘r’ for rooting condition, ‘t’ for lighter or heavy texture, ‘g’ for gravelliness or stoniness, ‘n’ for nutrient availability, ‘l’ for topography, ‘m’ for moisture availability, ‘w’ for drainage and ‘z’ for calcareousness. These limitations are indicated as lower case letters to the Class symbol. For example, moderately suitable lands with the limitations of soil depth and erosion are designated as S2re. For the microwatershed, the soil mapping units were evaluated and classified up to subclass level.

Using the above criteria, the soil map units of the microwatershed were evaluated and land suitability maps for 29 major annual and perennial crops were generated. The detailed information on the kind of suitability of each of the soil phase for the crops assessed are given village/ survey number wise for the microwatershed in Appendix-IV.

### 7.1 Land Suitability for Sorghum (*Sorghum bicolor*)

Sorghum is one of the major food crop grown in Karnataka in an area of 10.47 lakh ha in Bijapur, Gulbarga, Raichur, Bidar, Belgaum, Dharwad, Bellary, Chitradurga, Mysore and Tumakuru districts. The crop requirements for growing sorghum (Table 7.2) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing sorghum was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.1.

An area of about <1 ha (<1%) is highly suitable (Class S1) for growing sorghum and are distributed in the northern part of the microwatershed. An area of about 33 ha (11%) is moderately suitable (Class S2) for growing sorghum and are distributed in the northern and southeastern part of the microwatershed. They have minor limitations of



rooting depth and texture. About 83 ha (27%) is marginally suitable (Class S3) for growing sorghum and are distributed in the central, southeastern, eastern and northern part of the microwatershed with moderate limitations of rooting depth, gravelliness, nutrient availability, calcareousness and texture. Currently not suitable (Class N1) lands occur in an area of 149 ha (50%) and are distributed in the major part of the microwatershed with moderate limitation of rooting depth.




Fig. 7.1 Land Suitability map of Sorghum

## 7.2 Land Suitability for Maize (*Zea mays*)

Maize is one of the most important food crop grown in an area of 13.37 lakh ha in almost all the districts of the State. The crop requirements for growing maize (Table 7.3) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing maize was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.2.

An area of about 15 ha (5%) is highly suitable (Class S1) for growing maize and are distributed in the northern part of the microwatershed. An area of about 18 ha (6%) is moderately suitable (Class S2) for growing maize and are distributed in the southeastern part of the microwatershed. They have minor limitation of rooting depth. About 83 ha (27%) is marginally suitable (Class S3) for growing maize and are distributed in the central, southeastern, eastern and northern part of the microwatershed with moderate limitations of rooting depth, gravelliness, nutrient availability, calcareousness and texture. Currently not suitable (Class N1) lands occur in an area of 149 ha (50%) and are

distributed in the major part of the microwatershed with moderate limitation of rooting depth.




Fig. 7.2 Land Suitability map of Maize

### 7.3 Land Suitability for Bajra (*Pennisetum glaucum*)

Bajra is one of the most important millet crop grown in an area of 2.34 lakh ha in the northern districts of Karnataka state. The crop requirements for growing bajra (Table 7.4) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing bajra was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.3.

An area of about 15 ha (5%) is highly suitable (Class S1) for growing bajra and are distributed in the northern part of the microwatershed. An area of about 18 ha (6%) is moderately suitable (Class S2) for growing bajra and are distributed in the southeastern part of the microwatershed. They have minor limitation of rooting depth. About 83 ha (27%) is marginally suitable (Class S3) for growing bajra and are distributed in the central, southeastern, eastern and northern part of the microwatershed with moderate limitations of rooting depth, nutrient availability and calcareousness. Currently not suitable (Class N1) lands occur in an area of 149 ha (50%) and are distributed in the major part of the microwatershed with moderate limitation of rooting depth.




Fig. 7.3 Land Suitability map of Bajra

#### 7.4 Land Suitability for Groundnut (*Arachis hypogaea*)

Groundnut is one of the major oilseed crop grown in an area of 6.54 lakh ha in Karnataka in most of the districts either as rainfed or irrigated crop. The crop requirements for growing groundnut (Table 7.5) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing groundnut was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.4.

An area of about 15 ha (5%) is highly suitable (Class S1) for growing groundnut and are distributed in the northern part of the microwatershed. An area of about 18 ha (6%) is moderately suitable (Class S2) for growing groundnut and are distributed in the southeastern and northern part of the microwatershed. They have minor limitations of rooting depth and texture. About 82 ha (27%) is marginally suitable (Class S3) for growing groundnut and are distributed in the central, southeastern, eastern and northern part of the microwatershed with moderate limitation of rooting depth. Currently not suitable (Class N1) lands occur in an area of 150 ha (50%) and are distributed in the major part of the microwatershed with moderate limitations of rooting depth and nutrient availability.






Fig. 7.4 Land Suitability map of Groundnut

### 7.5 Land Suitability for Sunflower (*Helianthus annuus*)

Sunflower is one of the most important oilseed crop grown in an area of 3.56 lakh ha in the State in all the districts. The crop requirements for growing sunflower (Table 7.6) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing sunflower was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.5.

An area of about 15 ha (5%) is moderately suitable (Class S2) for sunflower and are distributed in the northern part of the microwatershed. They have minor limitations of rooting depth and texture. An area of about 18 ha (6%) is marginally suitable (Class S3) for growing sunflower and are distributed in the southeastern part of the microwatershed. They have minor limitation of rooting depth. Currently not suitable (Class N1) lands occur in an area of 232 ha (77%) and are distributed in the major part of the microwatershed with moderate limitations of rooting depth and nutrient availability.




Fig. 7.5 Land Suitability map of Sunflower

## 7.6 Land Suitability for Red gram (*Cajanus Cajan*)

Redgram is one of the most important pulse crop grown in an area of 7.28 lakh ha in almost all the districts of the State. The crop requirements for growing red gram (Table 7.7) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing redgram was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.6.

An area of about 15 ha (5%) is moderately suitable (Class S2) for redgram and are distributed in the northern part of the microwatershed. They have minor limitations of rooting depth and texture. An area of about 19 ha (6%) is marginally suitable (Class S3) for growing redgram and are distributed in the southeastern and northern part of the microwatershed. They have minor limitation of rooting depth, nutrient availability and calcareousness. Currently not suitable (Class N1) lands occur in an area of 231 ha (77%) and are distributed in the major part of the microwatershed with moderate limitation of rooting depth.




Fig. 7.6 Land Suitability map of Redgram

### 7.7 Land Suitability for Bengal gram (*Cicer aerativum*)

Bengal gram is one of the most important pulse crop grown in about 9.39 lakh ha area in Bijapur, Raichur, Kalaburgi, Dharwad, Belgaum and Bellary districts. The crop requirements for growing Bengal gram (Table 7.8) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing Bengal gram was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.7.

Marginally suitable lands (Class S3) for growing bengal gram occupy an area of about 49 ha (16%) and occur in the southeastern and northern part of the microwatershed. They have moderate limitations of texture, nutrient availability, calcareousness and rooting depth. Currently not suitable (Class N1) lands occur in an area of 216 ha (72%) and are distributed in the major part of the microwatershed with severe limitations of rooting depth and texture.




Fig. 7.7 Land Suitability map of Bengal gram.

## 7.8 Land Suitability for Cotton (*Gossypium hirsutum*)

Cotton is one of the most important fibre crop grown in the State in about 8.75 lakh ha area in Raichur, Dharwad, Belgaum, Kalaburgi, Bijapur, Bidar, Bellary, Chitradurga and Chamarajnagar districts. The crop requirements for growing cotton (Table 7.9) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing cotton was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.8.

An area of about <1 ha (<1%) is moderately suitable (Class S2) for growing cotton and are distributed in the northern part of the microwatershed. They have minor limitation of rooting depth. Marginally suitable lands (Class S3) for growing cotton occupy an area of about 49 ha (16%) and occur in the southeastern and northern part of the microwatershed. They have moderate limitations of texture, nutrient availability, gravelliness, calcareousness and rooting depth. Currently not suitable (Class N1) lands occur in an area of 216 ha (72%) and are distributed in the southern major part of the microwatershed with severe limitations of rooting depth and texture.






Fig. 7.8 Land Suitability map of Cotton

## 7.9 Land Suitability for Chilli (*Capsicum annuum*)

Chilli is one of the most important spice crop grown in about 0.42 lakh ha in Karnataka State. The crop requirements for growing chilli (Table 7.10) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing chilli was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.9.

An area of about 15 ha (5%) is highly suitable (Class S1) for growing chilli and are distributed in the northern part of the microwatershed. An area of about 18 ha (6%) is moderately suitable (Class S2) for growing chilli and are distributed in the southeastern part of the microwatershed. They have minor limitations of rooting depth. About 82 ha (27%) is marginally suitable (Class S3) for growing chilli and are distributed in the central, southeastern, eastern and northern part of the microwatershed with moderate limitations of rooting depth and graveliness. Currently not suitable (Class N1) lands occur in an area of 150 ha (50%) and are distributed in the major part of the microwatershed with moderate limitations of rooting depth and nutrient availability.




Fig 7.9 Land Suitability map of Chilli

### 7.10 Land Suitability for Tomato (*Lycopersicon esculentum*)

Tomato is one of the most important vegetable crop grown in about 0.61 lakh ha covering almost all the district of the state. The crop requirements for growing tomato (Table 7.11) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing tomato was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.10.

An area of about 15 ha (5%) is highly suitable (Class S1) for growing tomato and are distributed in the northern part of the microwatershed. An area of about 18 ha (6%) is moderately suitable (Class S2) for growing tomato and are distributed in the southeastern part of the microwatershed. They have minor limitation of rooting depth. About 82 ha (27%) is marginally suitable (Class S3) for growing tomato and are distributed in the central, southeastern, eastern and northern part of the microwatershed with moderate limitations of rooting depth and gravelliness. Currently not suitable (Class N1) lands occur in an area of 150 ha (50%) and are distributed in the major part of the microwatershed with moderate limitations of rooting depth and nutrient availability.




Fig 7.10 Land Suitability map of Tomato

### 7.11 Land Suitability for Brinjal (*Solanum melongena*)

Brinjal is one of the most important vegetable crop grown in the state. The crop requirements for growing brinjal (Table 7.12) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing Brinjal was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.11.

An area of about 15 ha (5%) is highly suitable (Class S1) for growing brinjal and are distributed in the northern part of the microwatershed. An area of about 18 ha (6%) is moderately suitable (Class S2) for growing brinjal and are distributed in the southeastern part of the microwatershed. They have minor limitations of rooting depth. About 82 ha (27%) is marginally suitable (Class S3) for growing brinjal and are distributed in the central, southeastern, eastern and northern part of the microwatershed with moderate limitations of rooting depth and gravelliness. Currently not suitable (Class N1) lands occur in an area of 150 ha (50%) and are distributed in the major part of the microwatershed with moderate limitations of rooting depth and nutrient availability.




Fig 7.11 Land Suitability map of Brinjal

## 7.12 Land Suitability for Onion (*Allium cepa L.*)

Onion is one of the most important vegetable crop grown in the state. The crop requirements for growing onion (Table 7.13) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing onion was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.12.

An area of about 15 ha (5%) is highly suitable (Class S1) for growing onion and are distributed in the northern part of the microwatershed. An area of about 18 ha (6%) is moderately suitable (Class S2) for growing onion and are distributed in the southeastern part of the microwatershed. They have minor limitations of rooting depth. About 82 ha (27%) is marginally suitable (Class S3) for growing onion and are distributed in the central, southeastern, eastern and northern part of the microwatershed with moderate limitations of rooting depth and gravelliness. Currently not suitable (Class N1) lands occur in an area of 150 ha (50%) and are distributed in the major part of the microwatershed with moderate limitations of rooting depth and nutrient availability.






Fig 7.12 Land Suitability map of Onion

### 7.13 Land Suitability for Bhendi (*Abelmoschus esculentus*)

Bhendi is one of the most important vegetable crop grown in the state. The crop requirements for growing bhendi (Table 7.14) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing bhendi was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.13.

An area of about 15 ha (5%) is highly suitable (Class S1) for growing bhendi and are distributed in the northern part of the microwatershed. An area of about 18 ha (6%) is moderately suitable (Class S2) for growing bhendi and are distributed in the southeastern part of the microwatershed. They have minor limitations of rooting depth. About 82 ha (27%) is marginally suitable (Class S3) for growing bhendi and are distributed in the central, southeastern, eastern and northern part of the microwatershed with moderate limitations of rooting depth and gravelliness. Currently not suitable (Class N1) lands occur in an area of 150 ha (50%) and are distributed in the major part of the microwatershed with moderate limitations of rooting depth and nutrient availability.




Fig 7.13 Land Suitability map of Bhendi

#### 7.14 Land Suitability for Drumstick (*Moringa oleifera*)

Drumstick is one of the most important vegetable crop grown in about 2403 ha in the state. The crop requirements for growing drumstick (Table 7.15) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing drumstick was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.14.

An area of about 15 ha (5%) is moderately suitable (Class S2) for drumstick and are distributed in the northern part of the microwatershed. They have minor limitation of rooting depth. An area of about 18 ha (6%) is marginally suitable (Class S3) for growing drumstick and are distributed in the southeastern part of the microwatershed. They have minor limitation of rooting depth. About 232 ha (77%) is currently not suitable (Class N1) for growing drumstick and are distributed in the major part of the microwatershed with moderate limitations of rooting depth and nutrient availability.




Fig 7.14 Land Suitability map of Drumstick

### 7.15 Land Suitability for Mango (*Mangifera indica*)

Mango is one of the most important fruit crop grown in an area of 1.73 lakh ha in almost all the districts of the State. The crop requirements (Table 7.16) for growing mango were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing mango was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed is given in Figure 7.15.

Marginally suitable (Class S3) lands for growing mango cover an area of about 15 ha (5%) and occur in the northern part of the microwatershed. They have moderate limitations of rooting depth. Currently not suitable (Class N1) for growing mango occupy an area about 250 ha (82%) and occur in the major part of the microwatershed. They have severe limitations of rooting depth and nutrient availability.






Fig. 7.15 Land Suitability map of Mango

## 7.16 Land Suitability for Guava (*Psidium guajava*)

Guava is one of the most important fruit crop grown in an area of 0.06 lakh ha in almost all the districts of the State. The crop requirements (Table 7.17) for growing guava were matched with the soil-site characteristics (7.1) and a land suitability map for growing guava was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.16.

An area of about 15 ha (5%) is moderately suitable (Class S2) for guava and are distributed in the northern part of the microwatershed. They have minor limitation of rooting depth. An area of about 18 ha (6%) is marginally suitable (Class S3) for growing guava and are distributed in the southeastern part of the microwatershed. They have minor limitation of rooting depth. About 232 ha (77%) is currently not suitable (Class N1) for growing guava and are distributed in the major part of the microwatershed with moderate limitations of rooting depth and nutrient availability.




Fig. 7.16 Land Suitability map of Guava

### 7.17 Land Suitability for Sapota (*Manilkara zapota*)

Sapota is one of the most important fruit crop grown in an area of 29373 ha in almost all the districts of the State. The crop requirements (Table 7.18) for growing sapota were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing sapota was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.17.

An area of about 15 ha (5%) is moderately suitable (Class S2) for sapota and are distributed in the northern part of the microwatershed. They have minor limitation of rooting depth. An area of about 18 ha (6%) is marginally suitable (Class S3) for growing sapota and are distributed in the southeastern part of the microwatershed. They have minor limitation of rooting depth. About 232 ha (77%) is currently not suitable (Class N1) for growing sapota and are distributed in the major part of the microwatershed with moderate limitations of rooting depth and nutrient availability.




Fig. 7.17 Land Suitability map of Sapota

### 7.18 Land Suitability for Pomegranate (*Punica granatum*)

Pomegranate is one of the most important fruit crop commercially grown in about 18488 ha in Karnataka, mainly in Bijapur, Bagalkot, Koppal, Gadag and Chitradurga districts. The crop requirements for growing pomegranate (Table 7.19) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing pomegranate was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.18.

An area of about 15 ha (5%) is moderately suitable (Class S2) for pomegranate and are distributed in the northern part of the microwatershed. They have minor limitation of rooting depth. An area of about 18 ha (6%) is marginally suitable (Class S3) for growing pomegranate and are distributed in the southeastern part of the microwatershed. They have minor limitation of rooting depth. About 232 ha (77%) is currently not suitable (Class N1) for growing pomegranate and are distributed in the major part of the microwatershed with moderate limitations of rooting depth and nutrient availability.




Fig 7.18 Land Suitability map of Pomegranate

### 7.19 Land Suitability for Musambi (*Citrus limetta*)

Musambi is one of the important fruit crop grown in an area of 3446 ha in almost all the districts of the State. The crop requirements for growing musambi (Table 7.20) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing musambi was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.19.

An area of about 15 ha (5%) is moderately suitable (Class S2) for musambi and are distributed in the northern part of the microwatershed. They have minor limitation of rooting depth. An area of about 18 ha (6%) is marginally suitable (Class S3) for growing musambi and are distributed in the southeastern part of the microwatershed. They have minor limitation of rooting depth. About 232 ha (77%) is currently not suitable (Class N1) for growing musambi and are distributed in the major part of the microwatershed with moderate limitations of rooting depth and nutrient availability.






Fig. 7.19 Land Suitability map of Musambi

## 7.20 Land Suitability for Lime (*Citrus sp*)

Lime is one of the most important fruit crop grown in an area of 0.11 lakh ha in almost all the districts of the State. The crop requirements for growing lime (Table 7.21) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing lime was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7. 20.

An area of about 15 ha (5%) is moderately suitable (Class S2) for lime and are distributed in the northern part of the microwatershed. They have minor limitation of rooting depth. An area of about 18 ha (6%) is marginally suitable (Class S3) for growing lime and are distributed in the southeastern part of the microwatershed. They have minor limitation of rooting depth. About 232 ha (77%) is currently not suitable (Class N1) for growing lime and are distributed in the major part of the microwatershed with moderate limitations of rooting depth and nutrient availability.




Fig. 7.20 Land Suitability map of Lime

### 7.21 Land Suitability for Amla (*Phyllanthus emblica*)

Amla is one of the medicinal fruit crop grown in almost all the districts of the State. The crop requirements for growing amla (Table 7.22) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing amla was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.21.

An area of about 15 ha (5%) is highly suitable (Class S1) for growing amla and are distributed in the northern part of the microwatershed. An area of about 18 ha (6%) is moderately suitable (Class S2) for growing amla and are distributed in the southeastern part of the microwatershed. They have minor limitations of rooting depth. About 82 ha (27%) is marginally suitable (Class S3) for growing amla and are distributed in the central, southeastern, eastern and northern part of the microwatershed with moderate limitations of rooting depth and texture. Currently not suitable (Class N1) lands occur in an area of 150 ha (50%) and are distributed in the major part of the microwatershed with moderate limitations of rooting depth and nutrient availability.




Fig. 7.21 Land Suitability map of Amla

## 7.22 Land Suitability for Cashew (*Anacardium occidentale*)

Cashew is one of the most important plantation nut crop grown in an area of 0.7 lakh ha in almost all the districts of the State. The crop requirements for growing cashew (Table 7.23) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing cashew was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.22.

An area of about <1 ha (<1%) is moderately suitable (Class S2) for growing cashew and are distributed in the northern part of the microwatershed. They have minor limitation of rooting depth. Marginally (Class S3) suitable lands for growing cashew occur in an area of 15 ha (5%) and are distributed in the northern part of the microwatershed. They have moderate limitation of nutrient availability. Currently not suitable (Class N1) lands for growing cashew occur in a maximum area of 249 ha (83%) and are distributed in the major part of the microwatershed with severe limitations of rooting depth and nutrient availability.






Fig. 7.22 Land Suitability map of Cashew

### 7. 23 Land Suitability for Jackfruit (*Artocarpus heterophyllus*)

Jackfruit is one of the most important fruit crop grown in an area of 5368 ha in almost all the districts of the State. The crop requirements for growing jackfruit (Table 7.24) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing jackfruit was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.23.

An area of about 15 ha (5%) is moderately suitable (Class S2) for jackfruit and are distributed in the northern part of the microwatershed. They have minor limitation of rooting depth. An area of about 18 ha (6%) is marginally suitable (Class S3) for growing jackfruit and are distributed in the southeastern part of the microwatershed. They have minor limitation of rooting depth. About 232 ha (77%) is currently not suitable (Class N1) for growing jackfruit and are distributed in the major part of the microwatershed with moderate limitations of rooting depth and nutrient availability.




Fig. 7.23 Land Suitability map of Jackfruit

## 7.24 Land Suitability for Jamun (*Syzygium cumini*)

Jamun is an important fruit crop grown in almost all the districts of the State. The crop requirements for growing jamun (Table 25) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing jamun was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.24.

An area of about 33 ha (11%) is marginally suitable (Class S3) for growing jamun and are distributed in the northern and southeastern part of the microwatershed. They have moderate limitation of rooting depth. About 232 ha (77%) is currently not suitable (Class N1) for growing jamun and are distributed in the major part of the microwatershed with moderate limitations of rooting depth and nutrient availability.




Fig. 7.24 Land Suitability map of Jamun

### 7.25 Land Suitability for Custard Apple (*Annona reticulata*)

Custard apple is one of the most important fruit crop grown in almost all the districts of the State. The crop requirements for growing custard apple (Table 7.26) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing custard apple was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.25.

An area of about 15 ha (5%) is highly suitable (Class S1) for growing custard apple and are distributed in the northern part of the microwatershed. An area of about 18 ha (6%) is moderately suitable (Class S2) for growing custard apple and are distributed in the southeastern part of the microwatershed. They have minor limitation of rooting depth. About 82 ha (27%) is marginally suitable (Class S3) for growing custard apple and are distributed in the central, southeastern, eastern and northern part of the microwatershed with moderate limitations of rooting depth and texture. Currently not suitable (Class N1) lands occur in an area of 150 ha (50%) and are distributed in the major part of the microwatershed with moderate limitations of rooting depth and nutrient availability.




Fig. 7.25 Land Suitability map of Custard Apple

## 7.26 Land Suitability for Tamarind (*Tamarindus indica*)

Tamarind is one of the most important spice crop grown in almost all the districts of the state. The crop requirements for growing tamarind (Table 7.27) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing tamarind was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Fig. 7.26.

Marginally suitable (Class S3) lands for growing tamarind cover an area of about 15 ha (5%) and occur in the northern part of the microwatershed. They have moderate limitation of rooting depth. Currently not suitable (Class N1) lands occur an area about 250 ha (83%) and occur in the major part of the microwatershed. They have severe limitations of rooting depth and nutrient availability.






Fig. 7.26 Land Suitability map of Tamarind

### 7.27 Land Suitability for Mulberry (*Morus nigra*)

Mulberry is the important leaf crop grown for rearing of silkworms in about 1.6 lakh ha area in all the districts of the state. The crop requirements for growing mulberry (Table 7.28) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing mulberry was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.27.

An area of about 15 ha (5%) is moderately suitable (Class S2) for mulberry and are distributed in the northern part of the microwatershed. They have minor limitation of rooting depth. An area of about 18 ha (6%) is marginally suitable (Class S3) for growing mulberry and are distributed in the southeastern part of the microwatershed. They have minor limitation of rooting depth. About 232 ha (77%) is currently not suitable (Class N1) for growing mulberry and are distributed in the major part of the microwatershed with moderate limitations of rooting depth and nutrient availability.




Fig 7.27 Land Suitability map of Mulberry

## 7.28 Land Suitability for Marigold (*Tagetes sps.*)

Marigold is one of the most important flower crop grown in an area of 9108 ha in almost all the districts of the State. The crop requirements (Table 7.29) for growing marigold were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing marigold was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.28.

An area of about 15 ha (5%) is highly suitable (Class S1) for growing marigold and are distributed in the northern part of the microwatershed. An area of about 18 ha (6%) is moderately suitable (Class S2) for growing marigold and are distributed in the southeastern part of the microwatershed. They have minor limitations of rooting depth. About 82 ha (27%) is marginally suitable (Class S3) for growing marigold and are distributed in the central, southeastern, eastern and northern part of the microwatershed with moderate limitations of rooting depth and gravelliness. Currently not suitable (Class N1) lands occur in an area of 150 ha (50%) and are distributed in the major part of the microwatershed with moderate limitations of rooting depth and nutrient availability.






Fig. 7.28 Land Suitability map of Marigold

## 7.29 Land Suitability for Chrysanthemum (*Dendranthema grandiflora*)

Chrysanthemum is one of the most important flower crop grown in an area of 4978 ha in almost all the districts of the State. The crop requirements (Table 7.30) for growing chrysanthemum were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing chrysanthemum was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.29.

An area of about 15 ha (5%) is highly suitable (Class S1) for growing chrysanthemum and are distributed in the northern part of the microwatershed. An area of about 18 ha (6%) is moderately suitable (Class S2) for growing chrysanthemum and are distributed in the southeastern part of the microwatershed. They have minor limitation of rooting depth. About 82 ha (27%) is marginally suitable (Class S3) for growing chrysanthemum and are distributed in the central, southeastern, eastern and northern part of the microwatershed with moderate limitations of rooting depth and graveliness. Currently not suitable (Class N1) lands occur in an area of 150 ha (50%) and are distributed in the major part of the microwatershed with moderate limitations of rooting depth and nutrient availability.




Fig. 7.29 Land Suitability map of Chrysanthemum

**Table 7.1 Soil-Site Characteristics of Kotageri-2 Microwatershed**

Soil Map Units	Climate (P) (mm)	Growing period (Days)	Drainage Class	Soil depth (cm)	Soil texture		Gravelliness		AWC (mm/m)	Slope (%)	Erosion	pH	EC (dSm <sup>-1</sup> )	ESP (%)	CEC [Cmol (p <sup>+</sup> )kg <sup>-1</sup> ]	BS (%)
					Sur-face	Sub-surface	Surface (%)	Sub-surface (%)								
BDPiB3	866	150	WD	<25	sc	scl	<15	<15	<50	1-3	moderate	8.58	0.262	0.35	18.10	100
KKRcB2	866	150	WD	<25	sl	sl	<15	10-15	<50	1-3	moderate	5.85	0.027	1.17	2.6	60.90
BDLbC3	866	150	WD	25-50	ls	sl	<15	<15	<50	3-5	severe	6.20	0.074	0.20	4.20	93
BDLhB2	866	150	WD	25-50	scl	sl	<15	<15	<50	1-3	moderate	6.20	0.074	0.20	4.20	93
BDLhB2g1	866	150	WD	25-50	scl	sl	15-35	<15	<50	1-3	moderate	6.20	0.074	0.20	4.20	93
BDLiB2	866	150	WD	25-50	sc	sl	<15	<15	<50	1-3	moderate	6.20	0.074	0.20	4.20	93
DSBcB2	866	150	WD	25-50	sl	g c	<15	35-60	<50	3-5	moderate	5.93	0.04	0.14	3.60	73
JNKiB2	866	150	WD	50-75	sc	scl	<15	<15	51-100	1-3	moderate	8.42	0.148	0.18	14.50	100
GWDmB2	866	150	MWD	75-100	c	scl	<15	<15	101-150	1-3	moderate	9.89	0.74	17.40	8.35	100
SHTcB2	866	150	WD	75-100	sl	scl	<15	15-35	51-100	1-3	moderate	7.26	0.199	0.86	10.60	100
PGPcB2	866	150	WD	75-100	sl	sc	<15	<15	51-100	1-3	moderate	6.83	0.210	2.83	3.15	100

\*Symbols and abbreviations are according to Field Guide for LRI under Sujala-III Project, Karnataka

**Table 7.2 Land suitability criteria for Sorghum**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	26–30	30–34; 24–26	34–40; 20–24	>40; <20
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, c (red), c (black)	scl, cl	ls, sl	-
	pH	1:2.5	5.5-7.8	5.0-5.5 7.8-9.0	>9.0	-
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	10-15
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

**Table 7.3 Land suitability criteria for Maize**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	30-34	35-38 26-30	38-40 26-20	
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	Mm				
	Rainfall in growing season	Mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc	c (red), c (black)	ls, sl	-
	pH	1:2.5	5.5-7.8	5.0-5.5 7.8-9.0	>9.0	-
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	Cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	-
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

**Table 7.4 Land suitability criteria for Bajra**

Land use requirement		Rating				
Soil –site characteristics	Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)	
Climatic regime	Mean temperature in growing season	°C	28-32	33-38 24-27	39-40 20-23	<20
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm	500-750	400-500	200-400	<200
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl, scl, cl,sc,c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.8	5.0-5.5 7.8-9.0	5.5-6.0 >9.0	
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	15-35	35-60	>60	
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	1-3	3-5	5-10	>10



**Table 7.5 Land suitability criteria for Groundnut**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	24–33	22–24; 33–35	20–22; 35–40	<20; >40
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	Mm				
	Rainfall in growing season	Mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. Well drained	Poorly drained	Very Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl	sl,cl, sc	c (red), c (black), ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	Cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<35	35-60	>60	
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.6 Land suitability criteria for Sunflower**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	24–30	30–34; 20–24	34–38; 16–20	>38; <16
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	mod. Well drained	-	Poorly to very drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	cl, sc,c (red), c (black)	scl	ls, sl	-
	pH	1:2.5	6.5-7.8	7.8-8.4 5.5-6.5	8.4-9.0; 5.0-5.5	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.7 Land suitability criteria for Redgram**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	30-35(G) 20-25(AV) 15-18 (F&PS) 35-40(M)	25-30(G) 20-25 (AV) 12-15 (F&PS) 30-35(M)	20-25(G) 15-20(AV) 10-12 (F&PS) 25-30(M)	< 20 <15 <10 <25
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	Mm				
	Rainfall in growing season	Mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. Well drained	Poorly drained	Very Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, c (red)	c (black),sl, scl, cl	ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-9.0	5.0-5.5 >9.0	-
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	Cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-50	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<1.0	1.0-2.0	>2.0	
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.8 Land suitability criteria for Bengal gram**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	20–25	25–30; 15–20	30–35; 10–15	>35; <10
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. Well drained	Poorly drained	Very Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	c(black)	-	c (red), scl, cl, sc	ls, sl
	pH	1:2.5	6.0-7.8	5.0-6.0 7.8-9.0	>9.0	-
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	-
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.9 Land suitability criteria for Cotton**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	22-32	>32	<19	-
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well to moderately well	Poorly drained/Some what excessively drained	-	very poorly/excessively drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, c (red,black)	cl	scl	ls, sl
	pH	1:2.5	6.5-7.8	7.8-8.4	5.5-6.5 8.4->9.0	<5.5
	CEC	C mol (p+)Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	50-100	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	<3	3-5	-	>5

**Table 7.10 Land suitability criteria for Chilli**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	25-32	33-35 20-25	35-38 <20	>38
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc	c (black), sl	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10



**Table 7.11 Land suitability criteria for Tomato**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	25-28	29-32 20-24	15-19 33-36	<15 >36
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl, scl, cl, sc, c (red)	-	ls, c(black)	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.12 Land suitability criteria for Brinjal**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	Well drained	Moderately well drained	Poorly drained	V. Poorly drained
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class				
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl, scl, cl, sc c (red)	-	ls, c (black)	-
	pH	1:2.5	6.0-7.3	7.3-8.4 5.0-6.0	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	>60
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.13 Land suitability criteria for Onion**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	20-30	30-35	35-40	>40
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately /imperfectly	-	Poorly to V poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl,scl,cl,sc,c (red)	-	c (Black),ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<1.0	1.0-2.0	2.0-4.0	<4
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.14 Land suitability criteria for Bhendi**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	25-28	29-32 20-24	15-19 33-36	<15 >36
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Imperfectly drained	Poorly to very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl,sc, c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.15 Land suitability criteria for Drumstick**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, scl, cl, c (red)	sl, c (black)	ls	S
	pH	1:2.5	6.0-7.3	5.0-5.5 7.3-7.8	5.5-6.0 7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<35	35-60	60-80	>80
Soil toxicity	Salinity (EC saturation extract)	ds/m				
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-10	-	>10

**Table 7.16 Land suitability criteria for Mango**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-32	24-27 33-35	36-40	20-24
	Min temp. before flowering	°C	10-15	15-22	>22	-
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration	Days				
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V. Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	-	ls, sl, c (black)	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>150	100-150	75-100	<75
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10



**Table 7.17 Land suitability criteria for Guava**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-32	33-36 24-27	37-42 20-23	
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	sl	c (black), ls	-
	pH	1:2.5	6.0-7.8	5.0-6.0	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.18 Land suitability criteria for Sapota**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-32	33-36 24-27	37-42 20-23	>42 <18
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	-	Poorly to very drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	sl	ls, c (black)	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.19 Land suitability criteria for Pomegranate**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	30-34	35-38 25-29	39-40 15-24	
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl,cl, sc, c (red)	c (black),sl	ls	-
	pH	1:2.5	5.5-7.8	7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.20 Land suitability criteria for Musambi**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-30	31-35 24-27	36-40 20-23	>40 <20
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately drained	poorly	Very poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c	sl	ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.21 Land suitability criteria for Lime**

Land use requirement		Rating				
Soil –site characteristics	Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)	
Climatic regime	Mean temperature in growing season	°C	28-30	31-35 24-27	36-40 20-23	>40 <20
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately drained	poorly	Very poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c	sl	ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.22 Land suitability criteria for Amla**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. well drained	Poorly drained	V. Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	c (black)	ls, sl	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15-35	35-60	60-80	-
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10



**Table 7.23 Land suitability criteria for Cashew**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	32 to 34	28 to 32; 34 to 38	24 to 28; 38 to 40	<20; >40
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	-	sl, ls	c (black)
	pH	1:2.5	5.5-6.5	5.0-5.5 6.5-7.3	7.3-7.8	>7.8
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-10	>10	-

**Table 7.24 Land suitability criteria for Jackfruit**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. well	Poorly	V. Poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	-	sl, ls, c (black)	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	>60
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10-

**Table 7.25 Land suitability criteria for Jamun**

Land use requirement		Rating				
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well	Mod. well	Poorly	V.Poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c(red)	sl, c (black)	ls	-
	pH	1:2.5	6.0-7.8	5.0-6.0	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>150	100-150	50-100	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	>60
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

**Table 7.26 Land suitability criteria for Custard apple**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	Scl, cl, sc, c (red), c (black)	-	Sl, ls	-
	pH	1:2.5	6.0-7.3	5.5-6.0 7.3-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15-35	35-60	60-80	-
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	>5	-

**Table 7.27 Land suitability criteria for Tamarind**

Land use requirement		Rating				
Soil –site characteristics	Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)	
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod.well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl,sc, c (red)	sl, c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>150	100-150	75-100	<75
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

**Table 7.28 Land suitability criteria for Mulberry**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	24–28	22–24; 28–32	32–38; 22–18	>38; <18
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V. Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, cl, scl	c (red)	c (black), sl, ls	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.8-8.4	7.3-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	0-35	35-60	60-80	>80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10



**Table 7.29 Land suitability criteria for Marigold**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	18-23	17-15 24-35	35-40 10-14	>40 <10
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl,scl, cl, sc, c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%				
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.30 Land suitability criteria for Chrysanthemum**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	18-23	17-15 24-35	35-40 10-14	>40 <10
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl,scl, cl, sc, c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%				
Erosion hazard	Slope	%	<3	3-5	5-10	>10

### 7.30 Land Management Units (LMUs)

The 11 soil map units identified in Kotageri-2 microwatershed have been grouped into 5 Land Management Units (LMUs) for the purpose of preparing a Proposed Crop Plan. Land Management Units are grouped based on the similarities in respect of the type of soil, the depth of the soil, the surface soil texture, gravel content, AWC, slope, erosion etc. and a Land Management Units map (Fig. 7.30) has been generated. These Land Management Units are expected to behave similarly for a given level of management. The map units that have been grouped into 5 Land Management Units along with brief description of soil and site characteristics are given below.

LMU	Soil map units	Soil and site characteristics
1	127.GWDmB2	Moderately deep sodic soils (75-100cm), 1-3 % slopes, non-gravelly (<15 %), moderate erosion.
2	40.PGPcB2	Moderately deep, sandy clay soils (75-100 cm), 1-3 % slopes, non-gravelly (<15%), moderate erosion.
3	36.SHThB2	Moderately deep sandy clay loam soils (75-100 cm), 1-3% slopes, non-gravelly (<15%), moderate erosion.
4	22.JNKiB2	Moderately shallow, sandy clay loam soils (50-75 cm), 1-3 % slopes, non-gravelly (<15%), moderate erosion.
5	3.BDLbC34.BDLhB2 162.BDLhB2g15.BDLiB2 119.BDPiB3121.DSBcB2 175.KKRcB2	Very shallow to shallow sandy clay loam to sandy loam soils (<25-50 cm), 1-3 % slopes, non-gravelly to very gravelly (<15 to 60%), moderate and severe erosion.




Fig. 7.30 Land Management Units Map Kotageri-2 Microwatershed

### **7.31 Proposed Crop Plan for Kotageri-2 Microwatershed**

After assessing the land suitability for the 29 crops, the Proposed Crop Plan has been prepared for the 5 identified LMUs by considering only the highly (Class S1) and moderately (Class S2) suitable lands for each of the 29 crops. The resultant proposed crop plan is presented below in Table 7.31.

**Table 7.31 Proposed Crop Plan for Kotageri-2 Microwatershed**

LMU	Soil Map Units	Survey Number	Field Crops/ Commercial crops	Horticulture Crops (Rainfed/Irrigated )	Suitable Interventions
1	127.GWDmB2	<b>Kootagera:</b> 181,182,183,188,199	-	<b>Agri-Silvi-Pasture</b> Ber, Aonla, Acacia sp. Dhaincha, Rhodes grass, Para grass ,Bermuda grass	Application of gypsum, iron pyrites and elemental sulphur. Addition of farm yard manure, green manures and providing subsurface drainage
2	40.PGPcB2	<b>Kootagera :</b> 28	Sunflower, Sorghum, Maize, Groundnut, Red gram, Bajra	<b>Fruit crops:</b> Mango, Musambi, Sapota, Tamarind, Pomegranate, Amla, Custard apple, Guava, Jackfruit, Jamun, Lime <b>Vegetables:</b> Tomato, Onion, Bhendi, Chilli, Brinjal, Drumstick, Coriander <b>Flowers:</b> Marigold, Chrysanthemum	Application of FYM, Biofertilizers and micronutrients, drip irrigation, mulching, suitable soil and water conservation practices
3	36.SHThB2	<b>Kootagera:</b> 6,7,8,9,10,11,12,13,14,15,16,17,18,26,27,323,179,184,185,186,187,189,190,191,192,193,324,325,326,327,328,329,337,338,339,340, 557	Sorghum, Maize, Groundnut, Red gram, Bajra, Sunflower,	<b>Fruit crops:</b> Musambi, Sapota, Pomegranate, Amla, Custard apple, Guava, Jackfruit, Lime <b>Vegetables:</b> Tomato, Onion, Bhendi, Chilli, Brinjal, Drumstick, Coriander <b>Flowers:</b> Marigold, Chrysanthemum	Application of FYM, Biofertilizers and micronutrients, drip irrigation, mulching, suitable soil and water conservation practices
4	22.JNKiB2	<b>Gajarakota :</b> 849,855,921,922,923,924,935,937 <b>Kootagera:</b> 442,445,446	Maize, Sorghum Groundnut, Bajra	<b>Fruit crops:</b> Amla, Custard apple <b>Vegetables:</b> Tomato, Chilli, Brinjal, Bhendi, Onion <b>Flowers:</b> Marigold, Chrysanthemum	Application of FYM, Biofertilizers and micronutrients, drip irrigation, mulching, suitable soil and water

LMU	Soil Map Units	Survey Number	Field Crops/ Commercial crops	Horticulture Crops (Rainfed/Irrigated )	Suitable Interventions
					conservation practices
5	3.BDLbC3 4.BDLhB2 162.BDLhB2g1 5.BDLiB2 119.BDPiB3 121.DSBcB2 175.KKRcB2	<b>Fatthepura :</b> 1,2,3,4,5,11,20 <b>Gajarakota:</b> 1139,847,848 <b>Kootagera:</b> 342,343,347,348, 349,350,351,353,354, 355,356,357,358,359,360,361, 362,363,364,365,367,369,370, 371,372,373,374,376,377,378, 379,380,381,385,386,387,388, 389,390,391,393,394,395,397, 398,399,400,401,402,403,406, 407,408,409,410,411,412,413, 414,415,416,417,418,419,420, 421,422,423,424,425,426,428, 429,433,434,435,436,437,443, 444,448,449,450,452,454,455 /1,455/2,455/3,455/4,456,457, 458,459,460,461,462,463,464, 465,466,467,468,469,470,471, 472,473,474,475,476/1,476/2, 476/3,476/4,476/5,477,478,47 9,480,481,482,483,484,485,48 6,487,488,489,490,491,492,49 3,494,495,496,497,498,499,50 0,532,533,534,535,536,537,53 8,541,542,543,544,545,546,54 7,548,549,550,551,553,554,55 5, 556	-	<b>Agri-Silvi-Pasture:</b> Hybrid Napier, <i>Styloxanthes hamata</i> , <i>Styloxanthes scabra</i>	Use of short duration varieties, sowing across the slope, drip irrigation and mulching is recommended

## SOIL HEALTH MANAGEMENT

### 8.1 Soil Health

Soil health is basic to plant health and plant health is basic to human and bovine health. Soil is fundamental to crop production. Without soil, no food could be produced nor would livestock be fed on a large scale. Because it is finite and fragile, soil is a precious resource that requires special care from its users.

Soil health or the capacity of the soil to function is critical to human survival. Soil health has been defined as: “the capacity of the soil to function as a living system without adverse effect on the ecosystem”. Healthy soils maintain a diverse community of soil organisms that help to form beneficial symbiotic associations with plant roots, recycle essential plant nutrients, improve soil structure with positive repercussions for soil, water and nutrient holding capacity and ultimately improve crop production and also contribute to mitigating climate change by maintaining or increasing its carbon content.

Functional interactions of soil biota with organic and inorganic components, air and water determine a soil’s potential to store and release nutrients and water to plants and to promote and sustain plant growth. Thus, maintaining soil health is vital to crop production and conserve soil resource base for sustaining agriculture.

#### **The most important characteristics of a healthy soil are**

- Good soil tilth
- Sufficient soil depth
- Good water storage and good drainage
- Adequate supply, but not excess of nutrients
- Large population of beneficial organisms
- Small proportion of plant pathogens and insect pests
- Low weed pressure
- Free of chemicals and toxins that may harm the crop
- Resistance to degradation
- Resilience when unfavorable conditions occur

#### **Characteristics of Kotageri-2 Microwatershed**

- ❖ The soil phases identified in the microwatershed belonged to different soil series, BDP series occupies maximum area of 132 ha (44%) followed by BDL 67 ha (22%), JNK 18 ha (6%), KKR 17 ha (6%), SHT 15 ha (5%), DSB 15 ha (5%), GWD 1 ha (<1%) and PGP 5 ha (<1%).
- ❖ As per land capability classification an area of 265 ha in the microwatershed falls under arable land category (Class II, III & IV). The major limitations identified in the arable lands were soil and erosion.
- ❖ On the basis of soil reaction about 252 ha (83%) area is neutral and 13 ha (4%) is slightly alkaline in soil reaction in the microwatershed.



## **Soil Health Management**

The following actions are required to improve the current land husbandry practices that provide a sound basis for the successful adoption of sustainable crop production system.

### **Neutral soils**

Entire cultivated area of 265 ha is under neutral soils.

1. Regular addition of organic manure, green manuring, green leaf manuring, crop residue incorporation and mulching needs to be taken up to improve the soil organic matter status.
2. Application of Biofertilizers, (*Azospirillum*, *Azotobacter*, *Rhizobium*).
3. Application of 100 per cent RDF.
4. Need based micronutrient applications.

Besides the above recommendations, the best transfer of technology options are also to be adopted.

### **Soil Degradation**

Soil erosion is one of the major factors affecting the soil health in the microwatershed. Out of total 302 ha area in the microwatershed, entire cultivated area of 121 ha (40%) and 144 ha (48%) is suffering from moderate and severe erosion respectively. These areas need immediate soil and water conservation and, other land development and land husbandry practices for restoring soil health.

### **Dissemination of Information and Communication of Benefits**

Any large scale implementation of soil health management requires that supporting information is made available widely, particularly through channels familiar to farmers and extension workers. Given the very high priority attached to soil-health especially by the Central Government on issuing Soil-Health Cards to all the farmers, media outlets like Regional, State and National Newspapers, Radio and Dooradarshan programs in local languages but also modern information and communication technologies such as Cellular phones and the Internet, which can be much more effective in reaching the younger farmers.

### **Inputs for Net Planning (Saturation Plan) and Interventions needed**

Net planning (Saturation Plan) in IWMP is focusing on preparation of

1. Soil and Water Conservation Plan for each plot or farm.
2. Productivity enhancement measures/ interventions for existing crops/livestock/other farm enterprises.
3. Diversification of farming mainly with perennial horticultural crops and livestock.
4. Improving livelihood opportunities and income generating activities.

In this connection, how various outputs of Sujala-III are of use in addressing these objectives of Net Planning (Saturation Plan) are briefly presented below.


- ❖ **Soil Depth:** The depth of a soil decides the amount of moisture and nutrients it can hold, what crops can be taken up or not, depending on the rooting depth and the length of growing period available for raising any crop. Deeper the soil, better for a wide variety of crops. If sufficient depth is not available for growing deep rooted crops, either choose medium or short duration crops or deeper planting pits need to be opened and additional good quality soil brought from outside has to be filled into the planting pits.
- ❖ **Surface Soil Texture:** Lighter soil texture in the top soil means, better rain water infiltration, less run-off and soil moisture conservation, less capillary rise and less evaporation losses. Lighter surface textured soils are amenable to good soil tilth and are highly suitable for crops like groundnut, root vegetables (carrot, radish, potato etc) but not ideal for crops that need stagnant water like lowland paddy. Heavy textured soils are poor in water infiltration and percolation. They are prone for sheet erosion; such soils can be improved by sand mulching. The technology that is developed by the AICRP-Dryland Agriculture, Vijayapura, Karnataka can be adopted.
- ❖ **Gravelliness:** More gravel content is favorable for run-off harvesting but poor in soil moisture storage and nutrient availability. It is a significant parameter that decides the kind of crop to be raised.
- ❖ **Land Capability Classification:** The land capability map shows the areas suitable and not suitable for agriculture and the major constraints in each of the plot/survey number. Hence, one can decide what kind of enterprise is possible in each of these units. In general, erosion and soil are the major constraints in Kotageri-2 microwatershed.
- ❖ **Organic Carbon:** The OC content (an index of available Nitrogen) is low (<0.5%) in about 207 ha (69%), medium (0.5-0.75%) in about 58 ha (19%) and <1 ha (<1%) is high (>0.75%) in organic carbon. The areas that are medium and low in OC needs to be further improved by applying farmyard manure and rotating crops with cereals and legumes or mixed cropping.
- ❖ **Promoting Green Manuring:** Growing of green manuring crops costs Rs. 1250/ha (green manuring seeds) and about Rs. 2000/ha towards cultivation that totals to Rs. 3250/- per ha. On the other hand, application of organic manure @ 10 tons/ha costs Rs. 5000/ha. The practice needs to be continued for 2-3 years or more. Nitrogen fertilizer needs to be supplemented by 25% in addition to the recommended level in 265 ha area where OC is low and medium (<0.5-0.75%). For example, for rainfed maize, recommended level is 50 kg N per ha and an additional 12 kg /ha needs to be applied for all the crops grown in these plots.
- ❖ **Available Phosphorus:** Available phosphorus is medium (23-57 kg/ha) covering an area of 168 ha (56%) and high (>57 kg/ha) covering an area of 97 ha (32%) in the microwatershed. For all the crops 25% additional P needs to be applied where available P is medium.

- ❖ **Available Potassium:** Available potassium is medium (145-337 kg/ha) covering a maximum area of 245 ha (81%) and low (145-337 kg/ha) covering an area of 19 ha (6%) in the microwatershed. All the plots, where available potassium is medium and low, additional 25% potassium may be applied.
- ❖ **Available Sulphur:** Available sulphur is a very critical nutrient for oilseed crops, it is medium (10-20ppm) in an area of 186 ha (62%) and low (<10 ppm) covers in an area of 79 ha (26%). Medium and low areas need to be applied with magnesium sulphate or gypsum or Factamphos (p) fertilizer (13% sulphur) for 2-3 years for the deficiency to be corrected.
- ❖ **Available Boron:** An area of 219 ha (73%) is low (<0.5ppm) and about 45 ha (15%) is medium (0.5-1.0ppm) in available boron. For these low and medium areas, application of sodium borate @ 10 kg/ha as soil application or 0.2 % borax as foliar spray is recommended.
- ❖ **Available Iron:** Available iron content is sufficient (>4.5 ppm) in the entire cultivated area of the microwatershed.
- ❖ **Available Manganese:** Entire cultivated area in the microwatershed is sufficient in available manganese content.
- ❖ **Available Copper:** Entire cultivated area in the microwatershed is sufficient in available copper content.
- ❖ **Available Zinc:** Available zinc content is deficient (<0.6 ppm) covering an area of 218 ha (72%) and sufficient (>0.6 ppm) in about 47 ha (15%) in the microwatershed. Application of zinc sulphate 25 kg/ha is recommended for the deficient areas.
- ❖ **Land Suitability for various crops:** Areas that are highly, moderately and marginally suitable for growing various crops are indicated. Along with the suitability, various constraints that are limiting the productivity are also indicated. For example, in case of cotton, gravel content, rooting depth and salinity/alkalinity are the major constraints in various plots. With suitable management interventions, the productivity can be enhanced. In order to increase the water holding capacity of light textured soils, growing of green manure crops and application of organic manure is recommended.

## SOIL AND WATER CONSERVATION TREATMENT PLAN

For preparing soil and water conservation treatment plan for Kotageri-2 microwatershed, the land resource inventory database generated under Sujala-III project has been transformed as information through series of interpretative (thematic) maps using soil phase map as a base. The various thematic maps (1:7920 scale) generated were

- Soil depth
- Surface soil texture
- Available water capacity
- Soil slope
- Soil gravelliness
- Land capability
- Present land use and land cover
- Crop suitability
- Rainfall
- Hydrology
- Water Resources
- Socio-economic data
- Contour plan with existing features- network of waterways, pothissa boundaries, cut up/ minor terraces etc.
- Cadastral map (1:7920 scale)
- Satellite imagery (1:7920 scale)



Apart from these, Hand Level/ Hydro Marker/ Dumpy Level/ Total Station and Kathedars' List to be collected.

### Steps for Survey and Preparation of Treatment Plan

The boundaries of Land User Groups' and Survey No. boundaries are traced in the field.

- Naming of user groups and farmers
- Identification of arable and non arable lands
- Identification of drainage lines and gullies
- Identification of non treatable areas
- Identification of priority areas in the arable lands
- Treatment plan for arable lands
- Location of water harvesting and recharge structures

### 9.1 Treatment Plan

The treatment plan recommended for arable lands is briefly described below

#### 9.1.1 Arable Land Treatment

## A. BUNDING

Steps for Survey and Preparation of Treatment Plan		<p><b>USER GROUP-1</b></p> <p>CLASSIFICATION OF GULLIES</p> <p>ಕೊರಕಲಿನ ವರ್ಗೀಕರಣ</p>
<ul style="list-style-type: none"> <li>• Cadastral map (1:7920 scale) is enlarged to a scale of 1:2500 scale</li> <li>• Existing network of waterways, pothissa boundaries, grass belts, natural drainage lines/ watercourse, cut ups/ terraces are marked on the cadastral map to the scale</li> <li>• Drainage lines are demarcated into</li> </ul>		
Small gullies	(up to 5 ha catchment)	
Medium gullies	(5-15 ha catchment)	
Ravines	(15-25 ha catchment) and	
Halla/Nala	(more than 25ha catchment)	

### Measurement of Land Slope

Land slope is estimated or determined by the study and interpretation of contours or by measurement in the field using simple instruments like Hand Level or Hydromarker.



Vertical and Horizontal intervals between bunds as recommended by the Watershed Development Department.

Slope percentage	Vertical interval (m)	Corresponding Horizontal Distance (m)
2 - 3%	0.6	24
3 - 4%	0.9	21
4 - 5%	0.9	21
5 - 6%	1.2	21
6 - 7%	1.2	21

**Note:** (i) The above intervals are maximum.

(ii) Considering the slope class and erosion status (A1... A=0-1 % slope, 1= slight erosion) the intervals have to be decided.

**Bund length recording:** Considering the contour plan and the existing grass belts/partitions, the bunds are aligned and lengths are measured.

## Section of the Bund

Bund section is decided considering the soil texture class and gravelliness class (bg<sub>0</sub>... b=loamy sand, g<sub>0</sub> = <15% gravel). The recommended Sections for different soils are given below.

### Recommended Bund Section



Top width (m)	Base width (m)	Height (m)	Side slope (Z:1;H:V)	Cross section (sq m)	Soil Texture	Remarks
0.3	0.9	0.3	01:01	0.18	Sandy loam	Vegetative bund
0.3	1.2	0.3	1.5:1	0.225	Sandy clay	
0.3	1.2	0.5	0.9:1	0.375	Red gravelly soils	
0.3	1.2	0.6	0.75:1	0.45		
0.3	1.5	0.6	01:01	0.54	Red sandy loam	
0.3	2.1	0.6	1.5:1	0.72	Very shallow black soils	
0.45	2	0.75	01:01	0.92		
0.45	2.4	0.75	1.3:1	1.07	Shallow black soils	
0.6	3.1	0.7	1.78:1	1.29	Medium black soils	
0.5	3	0.85	1.47:1	1.49		

## Formation of Trench cum Bund

Dimensions of the Borrow Pits/Trenches to be excavated (machinery are decided considering the Bund Section).

Details of Borrow Pit dimensions are given below:


### TRENCH CUM BUND

0.45 Sq.m section

IDEAL FOR HORTICULTURE CROPS

### 'A' FRAME FOR INTERBUND MANAGEMENT



1. ಸಮಸಾತನ ಉಳುವೆ
2. ಸಮಸಾತನ ಬಿತ್ತನೆ/ನಾಟಿ

### Size of Borrow Pits/ Trench recommended for Trench cum Bund (by machinery)

Bund section	Bund length	Earth quantity	Pit				Berm (pit to pit)	Soil depth class
			L(m)	W(m)	D(m)	Quantity (m <sup>3</sup> )		
m <sup>2</sup>	M	m <sup>3</sup>					m	
0.375	6	2.25	5.85	0.85	0.45	2.24	0.15	Shallow
0.45	6	2.7	5.4	1.2	0.43	2.79	0.6	Shallow
0.45	6	2.7	5	0.85	0.65	2.76	1	Moderately Shallow
0.54	5.6	3.02	5.5	0.85	0.7	3.27	0.1	Moderately shallow
0.54	5.5	2.97	5	1.2	0.5	3	0.5	Shallow
0.72	6.2	4.46	6	1.2	0.7	5.04	0.2	Moderately shallow
0.72	5.2	3.74	5.1	0.85	0.9	3.9	0.1	Moderately deep

#### B. Water Ways

1. Existing waterways are marked on the cadastral map (1:7920 scale) and their dimensions are recorded.
2. Considering the contour plan of the MWS, additional waterways/ modernization of the existing ones can be thought of.
3. The design details are given in the Manual.

#### C. Farm Ponds

Waterways and the catchment area will give an indication on the size of the Farm Pond. Location of the pond can be decided based on the contour plan/ field condition and farmers' need/desire.

#### D. Diversion Channel

Existing EPT/ CPT are marked on the cadastral map. Looking to the need, these can be modernized or fresh diversion channel can be proposed and runoff from this can be stored in Gokatte/ Recharge ponds.

#### 9.1.2 Non-Arable Land Treatment

Depending on the gravelliness and crops preferred by the farmers, the concerned authorities can decide appropriate treatment plan. The recommended treatments may be Contour Trench, Staggered Trench, Crescent Bund, Boulder Bund or Pebble Bund.



### **9.1.3 Treatment of Natural Water Course/ Drainage Lines**

- a) The cadastral map has to be updated as regards the network of drainage lines (gullies/ *nalas*/ *hallas*) and existing structures are marked to the scale and storage capacity of the existing water bodies are documented.
- b) The drainage line will be demarcated into Upper Reach, Middle Reach and Lower Reach.
- c) Considering the Catchment, *Nala* bed and bank conditions, suitable structures are decided.
- d) Number of storage structures (Check dam/ *Nala* bund/ Percolation tank) will be decided considering the commitments and available runoff from water budgeting and quality of water in the wells and site suitability.
- e) Detailed Leveling Survey using Dumpy Level / Total Station has to be carried out to arrive at the site-specific designs as shown in the Manual.
- f) The location of ground water recharge structures are decided by examining the lineaments and fracture zones from geological maps.
- g) Rainfall intensity data of the nearest Rain Gauge Station is considered for Hydrologic Designs.
- h) Silt load to the Storage/Recharge structures is reduced by providing vegetative, boulder and earthen checks in the natural water course. Location and design details are given in the Manual.

### **9.2 Recommended Soil and Water Conservation Measures**

The appropriate conservation structures best suited for each of the land parcel/ survey number (Appendix-I) are selected based on the slope per cent, severity of erosion, amount of rainfall, land use and soil type. The different kinds of conservation structures recommended are:

1. Graded / Strengthening of Bunds
2. Trench cum Bunds (TCB)
3. Trench cum Bunds / Strengthening
4. Crescent Bunds

A map (Fig. 9.1) showing soil and water conservation plan with different kinds of structures recommended has been prepared which shows the spatial distribution and extent of area. An area about 147 ha (49%) requires trench cum bunding and 118 ha (39%) needs Graded bunding.

The conservation plan prepared may be presented to all the stakeholders including farmers and after considering their suggestions, the conservation plan for the microwatershed may be finalised in a participatory approach.




Fig. 9.1 Soil and Water Conservation Plan map of Kotageri-2 Microwatershed

### 9.3 Greening of Microwatershed

As part of the greening programme in the watersheds, it is envisaged to plant a variety of horticultural and other tree plants that are edible, economical and produce lot of biomass which helps to restore the ecological balance in the watersheds. The lands that are suitable for greening programme are non-arable lands (land capability classes V, VI VII and VIII) and also the lands that are not suitable or marginally suitable and field bunds for growing annual and perennial crops. The method of planting these trees is given below.

It is recommended to open pits during the 1<sup>st</sup> week of March along the contour and heap the dug out soil on the lower side of the slope in order to harness the flowing water and facilitate weathering of soil in the pit. Exposure of soil in the pit also prevents spread of pests and diseases due to scorching sun rays. The pits should be filled with mixture of soil and organic manure during the second week of April and keep ready with sufficiently tall seedlings produced either in poly bags or in root trainer nurseries so that planting can be done during the 2<sup>nd</sup> or 3<sup>rd</sup> week of April depending on the rainfall.

The tree species suitable for the area considering rainfall, temperature and adaptability is listed below; waterlogged areas are recommended to be planted with species like Nerale (*Syzgium cumini*) and Bamboo. Dry areas are to be planted with species like Honge, Bevu, Seetaphal etc.

<b>Dry Deciduous Species</b>			<b>Temp (°C)</b>	<b>Rainfall (mm)</b>
1.	Bevu	<i>Azadiracta indica</i>	21–32	400 –1,200
2.	Tapasi	<i>Holoptelia integrifolia</i>	20-30	500 - 1000
3.	Seetaphal	<i>Anona Squamosa</i>	20-40	400 - 1000
4.	Honge	<i>Pongamia pinnata</i>	20 -50	500– 2,500
5.	Kamara	<i>Hardwickia binata</i>	25 -35	400 - 1000
6.	Bage	<i>Albezzia lebbek</i>	20 - 45	500 - 1000
7.	Ficus	<i>Ficus bengalensis</i>	20 - 50	500–2,500
8.	Sisso	<i>Dalbargia Sissoo</i>	20 - 50	500 -2000
9.	Ailanthus	<i>Ailanthus excelsa</i>	20 - 50	500 - 1000
10.	Hale	<i>Wrightia tinctoria</i>	25 - 45	500 - 1000
11.	Uded	<i>Steriospermum chelanoides</i>	25 - 45	500 -2000
12.	Dhupa	<i>Boswella Serrata</i>	20 - 40	500 - 2000
13.	Nelli	<i>Emblia Officinalis</i>	20 - 50	500 -1500
14.	Honne	<i>Pterocarpus marsupium</i>	20 - 40	500 - 2000
<b>Moist Deciduous Species</b>			<b>Temp (°C)</b>	<b>Rainfall (mm)</b>
15.	Teak	<i>Tectona grandis</i>	20 - 50	500-5000
16.	Nandi	<i>Legarstroemia lanceolata</i>	20 - 40	500 - 4000
17.	Honne	<i>Pterocarpus marsupium</i>	20 - 40	500 - 3000
18.	Mathi	<i>Terminalia alata</i>	20 -50	500 - 2000
19.	Shivane	<i>Gmelina arborea</i>	20 -50	500 -2000
20.	Kindal	<i>T.Paniculata</i>	20 - 40	500 - 1500
21.	Beete	<i>Dalbargia latifolia</i>	20 - 40	500 - 1500
22.	Tare	<i>T. belerica</i>	20 - 40	500 - 2000
23.	Bamboo	<i>Bambusa arundinasia</i>	20 - 40	500 - 2500
24.	Bamboo	<i>Dendrocalamus strictus</i>	20 – 40	500 – 2500
25.	Muthuga	<i>Butea monosperma</i>	20 - 40	400 - 1500
26.	Hippe	<i>Madhuca latifolia</i>	20 - 40	500 - 2000
27.	Sandal	<i>Santalum album</i>	20 - 50	400 - 1000
28.	Nelli	<i>Emblia officinalis</i>	20 - 40	500 - 2000
29.	Nerale	<i>Sizyium cumini</i>	20 - 40	500 - 2000
30.	Dhaman	<i>Grevia tilifolia</i>	20 - 40	500 - 2000
31.	Kaval	<i>Careya arborea</i>	20 - 40	500 - 2000
32.	Harada	<i>Terminalia chebula</i>	20 - 40	500 - 2000



## References

1. FAO (1976) Framework for Land Evaluation, Food and Agriculture Organization, Rome. 72 pp.
2. FAO (1983) Guidelines for Land Evaluation for Rainfed Agriculture, FAO, Rome, 237 pp.
3. IARI (1971) Soil Survey Manual, All India Soil and Land Use Survey Organization, IARI, New Delhi, 121 pp.
4. Katyal, J.C. and Rattan, R.K. (2003) Secondary and Micronutrients; Research Gap and Future Needs. Fert. News 48 (4); 9-20.
5. Naidu, L.G.K., Ramamurthy, V., Challa, O., Hegde, R. and Krishnan, P. (2006) Manual Soil Site Suitability Criteria for Major Crops, NBSS Publ. No. 129, NBSS & LUP, Nagpur, 118 pp.
6. Natarajan, A. and Dipak Sarkar (2010) Field Guide for Soil Survey, National Bureau of Soil Survey and Land Use Planning (ICAR), Nagpur, India.
7. Natarajan, A., Rajendra Hegde, Raj, J.N. and Shivananda Murthy, H.G. (2015) Implementation Manual for Sujala-III Project, Watershed Development Department, Bengaluru, Karnataka.
8. Sarma, V.A.K., Krishnan, P. and Budihal, S.L. (1987) Laboratory Manual, Tech. Bull. 23, NBSS & LUP, Nagpur.
9. Sehgal, J.L. (1990) Soil Resource Mapping of Different States of India; Why and How?, National Bureau of Soil Survey and Land Use Planning, Nagpur, 49 pp.
10. Shivaprasad, C.R., R.S. Reddy, J. Sehgal and M. Velayuthum (1998) Soils of Karnataka for Optimizing Land Use, NBSS Publ. No. 47b, NBSS & LUP, Nagpur, India.
11. Soil Survey Staff (2006) Keys to Soil Taxonomy, Tenth edition, U.S. Department of Agriculture/ NRCS, Washington DC, U.S.A.
12. Soil Survey Staff (2012) Soil Survey Manual, Handbook No. 18, USDA, Washington DC, USA.



**Appendix I**  
**Kotageri-2 (1B2d) Microwatershed**  
**Soil Phase Information**

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Gajarakota	847	2.26	BDLiB2	LMU-5	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Gajarakota	848	2.32	BDLhB2	LMU-5	Shallow (25-50 cm)	Sandy clay loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Gajarakota	849	0.51	JNKiB2	LMU-4	Moderately shallow (50-75 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Ies	Graded bunding
Gajarakota	855	3.57	JNKiB2	LMU-4	Moderately shallow (50-75 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut+Redgram (Gn+Rg)	Not Available	Ies	Graded bunding
Gajarakota	921	3.88	JNKiB2	LMU-4	Moderately shallow (50-75 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut+Redgram (Gn+Rg)	Not Available	Ies	Graded bunding
Gajarakota	922	0.85	JNKiB2	LMU-4	Moderately shallow (50-75 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut (Gn)	Not Available	Ies	Graded bunding
Gajarakota	923	0.73	JNKiB2	LMU-4	Moderately shallow (50-75 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut (Gn)	Not Available	Ies	Graded bunding
Gajarakota	924	0.44	JNKiB2	LMU-4	Moderately shallow (50-75 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Ies	Graded bunding
Gajarakota	935	0.31	JNKiB2	LMU-4	Moderately shallow (50-75 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut (Gn)	Not Available	Ies	Graded bunding
Gajarakota	937	0.33	JNKiB2	LMU-4	Moderately shallow (50-75 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut+Redgram (Gn+Rg)	Not Available	Ies	Graded bunding
Gajarakota	1139	0.15	BDPiB3	LMU-5	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Rock outcrops (Rc)	Not Available	IVes	Trench cum bunding
Kootagera	1	0.3	Habitatio n	Others	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others
Kootagera	2	0.28	Habitatio n	Others	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others
Kootagera	3	0.75	Habitatio n	Others	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others
Kootagera	4	0.61	Habitatio n	Others	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others
Kootagera	5	0.7	Habitatio n	Others	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others
Kootagera	6	0.8	SHThB2	LMU-3	Moderately deep (75-100 cm)	Sandy clay loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Habitation	Not Available	Ies	Graded bunding
Kootagera	7	0.13	SHThB2	LMU-3	Moderately deep (75-100 cm)	Sandy clay loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Ies	Graded bunding
Kootagera	8	0.32	SHThB2	LMU-3	Moderately deep (75-100 cm)	Sandy clay loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Ies	Graded bunding
Kootagera	9	0.22	SHThB2	LMU-3	Moderately deep (75-100 cm)	Sandy clay loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Ies	Graded bunding
Kootagera	10	0.32	SHThB2	LMU-3	Moderately deep (75-100 cm)	Sandy clay loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Ies	Graded bunding
Kootagera	11	0.1	SHThB2	LMU-3	Moderately deep (75-100 cm)	Sandy clay loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Ies	Graded bunding



Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Kootagera	12	0.7	SHThB2	LMU-3	Moderately deep (75-100 cm)	Sandy clay loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Kootagera	13	0.52	SHThB2	LMU-3	Moderately deep (75-100 cm)	Sandy clay loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Kootagera	14	0.77	SHThB2	LMU-3	Moderately deep (75-100 cm)	Sandy clay loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Kootagera	15	0.39	SHThB2	LMU-3	Moderately deep (75-100 cm)	Sandy clay loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Kootagera	16	0.29	SHThB2	LMU-3	Moderately deep (75-100 cm)	Sandy clay loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Kootagera	17	0.57	SHThB2	LMU-3	Moderately deep (75-100 cm)	Sandy clay loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Kootagera	18	0.13	SHThB2	LMU-3	Moderately deep (75-100 cm)	Sandy clay loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Kootagera	26	0.31	SHThB2	LMU-3	Moderately deep (75-100 cm)	Sandy clay loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Kootagera	27	1.34	SHThB2	LMU-3	Moderately deep (75-100 cm)	Sandy clay loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Kootagera	28	0.3	PGPcB2	LMU-2	Moderately deep (75-100 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Trench cum bunding
Kootagera	179	0.004	SHThB2	LMU-3	Moderately deep (75-100 cm)	Sandy clay loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Kootagera	181	0.04	GWDmB 2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IVes	Graded bunding
Kootagera	182	0.11	GWDmB 2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IVes	Graded bunding
Kootagera	183	0.12	GWDmB 2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IVes	Graded bunding
Kootagera	184	0.01	SHThB2	LMU-3	Moderately deep (75-100 cm)	Sandy clay loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Kootagera	185	0.06	SHThB2	LMU-3	Moderately deep (75-100 cm)	Sandy clay loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Kootagera	186	0.07	SHThB2	LMU-3	Moderately deep (75-100 cm)	Sandy clay loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Kootagera	187	0.05	SHThB2	LMU-3	Moderately deep (75-100 cm)	Sandy clay loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Kootagera	188	0.15	GWDmB 2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IVes	Graded bunding
Kootagera	189	0.68	SHThB2	LMU-3	Moderately deep (75-100 cm)	Sandy clay loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Kootagera	190	0.1	SHThB2	LMU-3	Moderately deep (75-100 cm)	Sandy clay loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Kootagera	191	0.17	SHThB2	LMU-3	Moderately deep (75-100 cm)	Sandy clay loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Kootagera	192	0.16	SHThB2	LMU-3	Moderately deep (75-100 cm)	Sandy clay loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Kootagera	193	0.18	SHThB2	LMU-3	Moderately deep (75-100 cm)	Sandy clay loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Kootagera	199	0.06	GWDmB2	LMU-1	Moderately deep (75-100 cm)	Clay	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IVes	Graded bunding
Kootagera	323	0.1	SHThB2	LMU-3	Moderately deep (75-100 cm)	Sandy clay loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Kootagera	324	0.69	SHThB2	LMU-3	Moderately deep (75-100 cm)	Sandy clay loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Habitation	Not Available	Iles	Graded bunding
Kootagera	325	1.01	SHThB2	LMU-3	Moderately deep (75-100 cm)	Sandy clay loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Kootagera	326	1.03	SHThB2	LMU-3	Moderately deep (75-100 cm)	Sandy clay loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Kootagera	327	0.52	SHThB2	LMU-3	Moderately deep (75-100 cm)	Sandy clay loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Kootagera	328	0.23	SHThB2	LMU-3	Moderately deep (75-100 cm)	Sandy clay loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Kootagera	329	0.05	SHThB2	LMU-3	Moderately deep (75-100 cm)	Sandy clay loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Kootagera	330	0.1	Habitatio n	Other s	Others	Others	Others	Others	Others	Others	Paddy (Pd)	Not Available	Others	Others
Kootagera	331	1.06	Habitatio n	Other s	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others
Kootagera	332	1.11	Habitatio n	Other s	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others
Kootagera	333	0.3	Habitatio n	Other s	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others
Kootagera	334	0.57	Habitatio n	Other s	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others
Kootagera	335	0.68	Habitatio n	Other s	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others
Kootagera	336	0.61	Habitatio n	Other s	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others
Kootagera	337	1.85	SHThB2	LMU-3	Moderately deep (75-100 cm)	Sandy clay loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Habitation	Not Available	Iles	Graded bunding
Kootagera	338	0.0006	SHThB2	LMU-3	Moderately deep (75-100 cm)	Sandy clay loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Kootagera	339	0.02	SHThB2	LMU-3	Moderately deep (75-100 cm)	Sandy clay loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Kootagera	340	0.15	SHThB2	LMU-3	Moderately deep (75-100 cm)	Sandy clay loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Kootagera	342	0.54	BDLhB2	LMU-5	Shallow (25-50 cm)	Sandy clay loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIIes	Graded bunding
Kootagera	343	0.5	BDLhB2	LMU-5	Shallow (25-50 cm)	Sandy clay loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIIes	Graded bunding
Kootagera	344	1.51	Habitatio n	Other s	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others
Kootagera	345	0.46	Habitatio n	Other s	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others
Kootagera	346	1.27	Habitatio n	Other s	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Kootagera	347	0.33	BDLhB2	LMU-5	Shallow (25-50 cm)	Sandy clay loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IIes	Graded bunding
Kootagera	348	2.35	BDLhB2g1	LMU-5	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IIes	Graded bunding
Kootagera	349	1.08	BDLhB2	LMU-5	Shallow (25-50 cm)	Sandy clay loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Kootagera	350	2.49	BDLhB2	LMU-5	Shallow (25-50 cm)	Sandy clay loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Kootagera	351	0.4	BDLhB2	LMU-5	Shallow (25-50 cm)	Sandy clay loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Kootagera	353	0.56	KKRcB2	LMU-5	Very shallow (<25 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IVes	Graded bunding
Kootagera	354	0.49	BDLhB2	LMU-5	Shallow (25-50 cm)	Sandy clay loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IIes	Graded bunding
Kootagera	355	0.38	KKRcB2	LMU-5	Very shallow (<25 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IVes	Graded bunding
Kootagera	356	0.43	KKRcB2	LMU-5	Very shallow (<25 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IVes	Graded bunding
Kootagera	357	0.79	KKRcB2	LMU-5	Very shallow (<25 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IVes	Graded bunding
Kootagera	358	0.57	KKRcB2	LMU-5	Very shallow (<25 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IVes	Graded bunding
Kootagera	359	0.17	KKRcB2	LMU-5	Very shallow (<25 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IVes	Graded bunding
Kootagera	360	0.07	KKRcB2	LMU-5	Very shallow (<25 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IVes	Graded bunding
Kootagera	361	0.06	KKRcB2	LMU-5	Very shallow (<25 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IVes	Graded bunding
Kootagera	362	0.12	KKRcB2	LMU-5	Very shallow (<25 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IVes	Graded bunding
Kootagera	363	0.42	KKRcB2	LMU-5	Very shallow (<25 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IVes	Graded bunding
Kootagera	364	0.22	KKRcB2	LMU-5	Very shallow (<25 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IVes	Graded bunding
Kootagera	365	0.47	KKRcB2	LMU-5	Very shallow (<25 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IVes	Graded bunding
Kootagera	366	0.36	Waterbody	Others	Others	Others	Others	Others	Others	Others	Paddy (Pd)	Not Available	Others	Others
Kootagera	367	0.51	KKRcB2	LMU-5	Very shallow (<25 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IVes	Graded bunding
Kootagera	368	0.23	Waterbody	Others	Others	Others	Others	Others	Others	Others	Paddy (Pd)	Not Available	Others	Others
Kootagera	369	0.6	KKRcB2	LMU-5	Very shallow (<25 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IVes	Graded bunding
Kootagera	370	0.22	KKRcB2	LMU-5	Very shallow (<25 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IVes	Graded bunding
Kootagera	371	0.66	KKRcB2	LMU-5	Very shallow (<25 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IVes	Graded bunding

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Kootagera	372	0.23	KKRcB2	LMU-5	Very shallow (<25 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IVes	Graded bunding
Kootagera	373	0.25	KKRcB2	LMU-5	Very shallow (<25 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IVes	Graded bunding
Kootagera	374	0.04	KKRcB2	LMU-5	Very shallow (<25 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IVes	Graded bunding
Kootagera	375	0.04	Waterbody	Others	Others	Others	Others	Others	Others	Others	Paddy (Pd)	Not Available	Others	Others
Kootagera	376	0.33	KKRcB2	LMU-5	Very shallow (<25 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IVes	Graded bunding
Kootagera	377	0.06	KKRcB2	LMU-5	Very shallow (<25 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IVes	Graded bunding
Kootagera	378	0.16	KKRcB2	LMU-5	Very shallow (<25 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IVes	Graded bunding
Kootagera	379	0.54	KKRcB2	LMU-5	Very shallow (<25 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IVes	Graded bunding
Kootagera	380	0.45	KKRcB2	LMU-5	Very shallow (<25 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IVes	Graded bunding
Kootagera	381	0.48	KKRcB2	LMU-5	Very shallow (<25 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IVes	Graded bunding
Kootagera	382	0.04	Waterbody	Others	Others	Others	Others	Others	Others	Others	Paddy (Pd)	Not Available	Others	Others
Kootagera	383	0.1	Waterbody	Others	Others	Others	Others	Others	Others	Others	Paddy (Pd)	Not Available	Others	Others
Kootagera	384	0.24	Waterbody	Others	Others	Others	Others	Others	Others	Others	Paddy (Pd)	Not Available	Others	Others
Kootagera	385	0.39	KKRcB2	LMU-5	Very shallow (<25 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IVes	Graded bunding
Kootagera	386	0.72	KKRcB2	LMU-5	Very shallow (<25 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IVes	Graded bunding
Kootagera	387	0.21	KKRcB2	LMU-5	Very shallow (<25 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IVes	Graded bunding
Kootagera	388	0.39	KKRcB2	LMU-5	Very shallow (<25 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IVes	Graded bunding
Kootagera	389	0.2	KKRcB2	LMU-5	Very shallow (<25 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IVes	Graded bunding
Kootagera	390	0.25	KKRcB2	LMU-5	Very shallow (<25 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IVes	Graded bunding
Kootagera	391	0.33	KKRcB2	LMU-5	Very shallow (<25 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IVes	Graded bunding
Kootagera	392	0.21	Waterbody	Others	Others	Others	Others	Others	Others	Others	Paddy (Pd)	Not Available	Others	Others
Kootagera	393	0.11	KKRcB2	LMU-5	Very shallow (<25 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IVes	Graded bunding
Kootagera	394	0.09	KKRcB2	LMU-5	Very shallow (<25 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IVes	Graded bunding
Kootagera	395	0.09	KKRcB2	LMU-5	Very shallow (<25 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IVes	Graded bunding



Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Kootagera	396	0.21	Waterbody	Others	Others	Others	Others	Others	Others	Others	Paddy (Pd)	Not Available	Others	Others
Kootagera	397	0.22	DSBcB2	LMU-5	Shallow (25-50 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IIes	Trench cum bunding
Kootagera	398	0.41	DSBcB2	LMU-5	Shallow (25-50 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IIes	Trench cum bunding
Kootagera	399	0.14	DSBcB2	LMU-5	Shallow (25-50 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IIes	Trench cum bunding
Kootagera	400	0.15	BDPiB3	LMU-5	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Paddy (Pd)	Not Available	IVes	Trench cum bunding
Kootagera	401	0.12	BDPiB3	LMU-5	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Paddy (Pd)	Not Available	IVes	Trench cum bunding
Kootagera	402	2.82	BDPiB3	LMU-5	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Redgram (Rg)	Not Available	IVes	Trench cum bunding
Kootagera	403	3.52	DSBcB2	LMU-5	Shallow (25-50 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram (Gg)	Not Available	IIes	Trench cum bunding
Kootagera	404	6.35	Waterbody	Others	Others	Others	Others	Others	Others	Others	Waterbody	Not Available	Others	Others
Kootagera	405	5.93	Waterbody	Others	Others	Others	Others	Others	Others	Others	Waterbody	Not Available	Others	Others
Kootagera	406	0.21	KKRcB2	LMU-5	Very shallow (<25 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IVes	Graded bunding
Kootagera	407	0.38	KKRcB2	LMU-5	Very shallow (<25 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IVes	Graded bunding
Kootagera	408	0.2	KKRcB2	LMU-5	Very shallow (<25 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IVes	Graded bunding
Kootagera	409	0.16	KKRcB2	LMU-5	Very shallow (<25 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IVes	Graded bunding
Kootagera	410	0.3	KKRcB2	LMU-5	Very shallow (<25 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IVes	Graded bunding
Kootagera	411	0.42	DSBcB2	LMU-5	Shallow (25-50 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IIes	Trench cum bunding
Kootagera	412	0.44	KKRcB2	LMU-5	Very shallow (<25 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IVes	Graded bunding
Kootagera	413	0.04	KKRcB2	LMU-5	Very shallow (<25 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IVes	Graded bunding
Kootagera	414	0.44	KKRcB2	LMU-5	Very shallow (<25 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IVes	Graded bunding
Kootagera	415	0.07	KKRcB2	LMU-5	Very shallow (<25 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IVes	Graded bunding
Kootagera	416	0.52	KKRcB2	LMU-5	Very shallow (<25 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IVes	Graded bunding
Kootagera	417	0.27	KKRcB2	LMU-5	Very shallow (<25 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IVes	Graded bunding
Kootagera	418	0.32	KKRcB2	LMU-5	Very shallow (<25 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IVes	Graded bunding
Kootagera	419	0.42	KKRcB2	LMU-5	Very shallow (<25 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IVes	Graded bunding

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Kootagera	420	0.59	KKRcB2	LMU-5	Very shallow (<25 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IVes	Graded bunding
Kootagera	421	0.66	KKRcB2	LMU-5	Very shallow (<25 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IVes	Graded bunding
Kootagera	422	0.17	KKRcB2	LMU-5	Very shallow (<25 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IVes	Graded bunding
Kootagera	423	1.04	KKRcB2	LMU-5	Very shallow (<25 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IVes	Graded bunding
Kootagera	424	2.49	BDLhB2	LMU-5	Shallow (25-50 cm)	Sandy clay loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram (Gg)	Not Available	IIIes	Graded bunding
Kootagera	425	0.2	BDLhB2	LMU-5	Shallow (25-50 cm)	Sandy clay loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIIes	Graded bunding
Kootagera	426	0.2	BDLhB2	LMU-5	Shallow (25-50 cm)	Sandy clay loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIIes	Graded bunding
Kootagera	428	0.31	BDLhB2	LMU-5	Shallow (25-50 cm)	Sandy clay loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIIes	Graded bunding
Kootagera	429	1.64	BDLhB2	LMU-5	Shallow (25-50 cm)	Sandy clay loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Greengram (Rg+Gg)	Not Available	IIIes	Graded bunding
Kootagera	433	0.26	BDLhB2	LMU-5	Shallow (25-50 cm)	Sandy clay loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram (Gg)	Not Available	IIIes	Graded bunding
Kootagera	434	0.62	BDLhB2	LMU-5	Shallow (25-50 cm)	Sandy clay loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram (Gg)	Not Available	IIIes	Graded bunding
Kootagera	435	0.54	BDLhB2	LMU-5	Shallow (25-50 cm)	Sandy clay loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIIes	Graded bunding
Kootagera	436	0.48	BDLhB2	LMU-5	Shallow (25-50 cm)	Sandy clay loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram (Gg)	Not Available	IIIes	Graded bunding
Kootagera	437	0.25	BDLhB2	LMU-5	Shallow (25-50 cm)	Sandy clay loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram (Gg)	Not Available	IIIes	Graded bunding
Kootagera	442	0.18	JNKiB2	LMU-4	Moderately shallow (50-75 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram (Gg)	Not Available	Iles	Graded bunding
Kootagera	443	0.9	BDLhB2	LMU-5	Shallow (25-50 cm)	Sandy clay loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram (Gg)	Not Available	IIIes	Graded bunding
Kootagera	444	0.58	BDLhB2	LMU-5	Shallow (25-50 cm)	Sandy clay loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram (Gg)	Not Available	IIIes	Graded bunding
Kootagera	445	6.79	JNKiB2	LMU-4	Moderately shallow (50-75 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Kootagera	446	2.11	JNKiB2	LMU-4	Moderately shallow (50-75 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Kootagera	447	4.48	Waterbody	Others	Others	Others	Others	Others	Others	Others	Waterbody	Not Available	Others	Others
Kootagera	448	5.45	DSBcB2	LMU-5	Shallow (25-50 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram (Gg)	Not Available	IIIes	Trench cum bunding
Kootagera	449	3.83	DSBcB2	LMU-5	Shallow (25-50 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram (Gg)	Not Available	IIIes	Trench cum bunding
Kootagera	450	1.82	BDPiB3	LMU-5	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Rock outcrops (Rc)	Not Available	IVes	Trench cum bunding
Kootagera	452	1.15	BDPiB3	LMU-5	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Redgram (Rg)	Not Available	IVes	Trench cum bunding

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Kootagera	454	1.31	BDPiB3	LMU-5	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Scrub land (SI)	Not Available	IVes	Trench cum bunding
Kootagera	455/1	44.81	BDPiB3	LMU-5	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Rock outcrops (Rc)	Not Available	IVes	Trench cum bunding
Kootagera	455/2	0.98	BDPiB3	LMU-5	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Redgram (Rg)	Not Available	IVes	Trench cum bunding
Kootagera	455/3	1.54	BDPiB3	LMU-5	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Rock outcrops (Rc)	Not Available	IVes	Trench cum bunding
Kootagera	455/4	1.09	BDPiB3	LMU-5	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Rock outcrops (Rc)	Not Available	IVes	Trench cum bunding
Kootagera	456	0.34	BDPiB3	LMU-5	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Redgram (Rg)	Not Available	IVes	Trench cum bunding
Kootagera	457	0.23	BDPiB3	LMU-5	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Redgram (Rg)	Not Available	IVes	Trench cum bunding
Kootagera	458	0.22	BDPiB3	LMU-5	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Redgram (Rg)	Not Available	IVes	Trench cum bunding
Kootagera	459	0.91	BDPiB3	LMU-5	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Redgram (Rg)	Not Available	IVes	Trench cum bunding
Kootagera	460	1.45	BDPiB3	LMU-5	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Redgram (Rg)	Not Available	IVes	Trench cum bunding
Kootagera	461	0.5	BDPiB3	LMU-5	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Redgram (Rg)	Not Available	IVes	Trench cum bunding
Kootagera	462	1.04	BDPiB3	LMU-5	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Redgram (Rg)	Not Available	IVes	Trench cum bunding
Kootagera	463	1.95	BDPiB3	LMU-5	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Redgram (Rg)	Not Available	IVes	Trench cum bunding
Kootagera	464	3.92	BDPiB3	LMU-5	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Redgram (Rg)	Not Available	IVes	Trench cum bunding
Kootagera	465	6.09	BDPiB3	LMU-5	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Redgram (Rg)	Not Available	IVes	Trench cum bunding
Kootagera	466	1.25	BDPiB3	LMU-5	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Redgram (Rg)	Not Available	IVes	Trench cum bunding
Kootagera	467	0.41	BDPiB3	LMU-5	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Redgram (Rg)	Not Available	IVes	Trench cum bunding
Kootagera	468	0.15	BDPiB3	LMU-5	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Redgram (Rg)	Not Available	IVes	Trench cum bunding
Kootagera	469	0.3	BDPiB3	LMU-5	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Redgram (Rg)	Not Available	IVes	Trench cum bunding
Kootagera	470	0.62	BDPiB3	LMU-5	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Redgram (Rg)	Not Available	IVes	Trench cum bunding
Kootagera	471	0.76	BDPiB3	LMU-5	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Redgram (Rg)	Not Available	IVes	Trench cum bunding
Kootagera	472	1.04	BDPiB3	LMU-5	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Redgram (Rg)	Not Available	IVes	Trench cum bunding
Kootagera	473	0.26	BDPiB3	LMU-5	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Redgram (Rg)	Not Available	IVes	Trench cum bunding
Kootagera	474	0.07	BDPiB3	LMU-5	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Redgram (Rg)	Not Available	IVes	Trench cum bunding



Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Kootagera	475	1.39	BDPiB3	LMU-5	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Redgram (Rg)	Not Available	IVes	Trench cum bunding
Kootagera	476/1	44.35	BDPiB3	LMU-5	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Rock outcrops (Rc)	Not Available	IVes	Trench cum bunding
Kootagera	476/2	0.29	BDPiB3	LMU-5	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Redgram (Rg)	Not Available	IVes	Trench cum bunding
Kootagera	476/3	0.6	BDPiB3	LMU-5	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Redgram (Rg)	Not Available	IVes	Trench cum bunding
Kootagera	476/4	2.14	BDPiB3	LMU-5	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Redgram (Rg)	Not Available	IVes	Trench cum bunding
Kootagera	476/5	0.69	BDPiB3	LMU-5	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Redgram (Rg)	Not Available	IVes	Trench cum bunding
Kootagera	477	0.04	BDLhB2g 1	LMU-5	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IIIes	Graded bunding
Kootagera	478	0.53	BDLhB2g 1	LMU-5	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IIIes	Graded bunding
Kootagera	479	2.15	BDLbC3	LMU-5	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Redgram (Rg)	Not Available	IVes	Graded bunding
Kootagera	480	0.62	BDLhB2g 1	LMU-5	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IIIes	Graded bunding
Kootagera	481	0.53	BDLhB2g 1	LMU-5	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IIIes	Graded bunding
Kootagera	482	0.17	BDLhB2g 1	LMU-5	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IIIes	Graded bunding
Kootagera	483	0.37	BDLhB2g 1	LMU-5	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IIIes	Graded bunding
Kootagera	484	0.7	BDLhB2g 1	LMU-5	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIIes	Graded bunding
Kootagera	485	0.27	BDLhB2g 1	LMU-5	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIIes	Graded bunding
Kootagera	486	0.69	BDLhB2g 1	LMU-5	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIIes	Graded bunding
Kootagera	487	0.43	BDLhB2g 1	LMU-5	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIIes	Graded bunding
Kootagera	488	0.68	BDLhB2g 1	LMU-5	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIIes	Graded bunding
Kootagera	489	0.5	BDLhB2g 1	LMU-5	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IIIes	Graded bunding
Kootagera	490	1.12	BDLhB2g 1	LMU-5	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IIIes	Graded bunding
Kootagera	491	0.38	BDLbC3	LMU-5	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Paddy (Pd)	Not Available	IVes	Graded bunding
Kootagera	492	5.09	BDLbC3	LMU-5	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Redgram (Rg)	Not Available	IVes	Graded bunding
Kootagera	493	0.58	BDLhB2g 1	LMU-5	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIIes	Graded bunding
Kootagera	494	0.56	BDLbC3	LMU-5	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Redgram (Rg)	Not Available	IVes	Graded bunding

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Kootagera	495	0.51	BDLbC3	LMU-5	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Redgram (Rg)	Not Available	IVes	Graded bunding
Kootagera	496	1.79	BDLhB2g 1	LMU-5	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	IIIs	Graded bunding
Kootagera	497	1.34	BDLbC3	LMU-5	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Redgram (Rg)	Not Available	IVes	Graded bunding
Kootagera	498	0.82	BDLbC3	LMU-5	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Redgram (Rg)	Not Available	IVes	Graded bunding
Kootagera	499	0.77	BDLbC3	LMU-5	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Redgram (Rg)	Not Available	IVes	Graded bunding
Kootagera	500	1.89	BDPiB3	LMU-5	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Redgram (Rg)	Not Available	IVes	Trench cum bunding
Kootagera	532	1.3	BDLbC3	LMU-5	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Redgram+Cotton (Rg+Ct)	Not Available	IVes	Graded bunding
Kootagera	533	0.87	BDLbC3	LMU-5	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Redgram (Rg)	Not Available	IVes	Graded bunding
Kootagera	534	0.31	BDLbC3	LMU-5	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Redgram (Rg)	Not Available	IVes	Graded bunding
Kootagera	535	0.05	BDLbC3	LMU-5	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Redgram (Rg)	Not Available	IVes	Graded bunding
Kootagera	536	0.92	BDLbC3	LMU-5	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Redgram (Rg)	Not Available	IVes	Graded bunding
Kootagera	537	1.24	BDLhB2g 1	LMU-5	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIIs	Graded bunding
Kootagera	538	0.004	BDLhB2g 1	LMU-5	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIIs	Graded bunding
Kootagera	541	0.17	BDLhB2g 1	LMU-5	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIIs	Graded bunding
Kootagera	542	0.34	BDLhB2g 1	LMU-5	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIIs	Graded bunding
Kootagera	543	0.92	BDLhB2g 1	LMU-5	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIIs	Graded bunding
Kootagera	544	1.53	BDLhB2g 1	LMU-5	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIIs	Graded bunding
Kootagera	545	0.72	BDLhB2g 1	LMU-5	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIIs	Graded bunding
Kootagera	546	0.89	BDLhB2g 1	LMU-5	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	IIIs	Graded bunding
Kootagera	547	0.45	BDLhB2g 1	LMU-5	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIIs	Graded bunding
Kootagera	548	0.66	BDLhB2g 1	LMU-5	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	IIIs	Graded bunding
Kootagera	549	0.98	BDLhB2g 1	LMU-5	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIIs	Graded bunding
Kootagera	550	0.35	BDLhB2g 1	LMU-5	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIIs	Graded bunding
Kootagera	551	1.13	BDLhB2g 1	LMU-5	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIIs	Graded bunding

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Kootagera	552	0.81	Habitations	Others	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others
Kootagera	553	0.47	BDLhB2g1	LMU-5	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIIes	Graded bunding
Kootagera	554	1.03	BDLhB2g1	LMU-5	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIIes	Graded bunding
Kootagera	555	0.69	BDLhB2g1	LMU-5	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIIes	Graded bunding
Kootagera	556	1.39	BDLhB2g1	LMU-5	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	IIIes	Graded bunding
Kootagera	557	1.8	SHTbB2	LMU-3	Moderately deep (75-100 cm)	Sandy clay loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	IIes	Graded bunding
Fatthepura	1	0.2	BDPiB3	LMU-5	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Forest (Fo)	Not Available	IVes	Trench cum bunding
Fatthepura	2	0.42	BDPiB3	LMU-5	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Forest (Fo)	Not Available	IVes	Trench cum bunding
Fatthepura	3	0.68	BDPiB3	LMU-5	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Forest (Fo)	Not Available	IVes	Trench cum bunding
Fatthepura	4	0.27	BDPiB3	LMU-5	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Forest (Fo)	Not Available	IVes	Trench cum bunding
Fatthepura	5	0.002	BDPiB3	LMU-5	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Forest (Fo)	Not Available	IVes	Trench cum bunding
Fatthepura	11	0.02	BDPiB3	LMU-5	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Forest (Fo)	Not Available	IVes	Trench cum bunding
Fatthepura	20	0.09	BDPiB3	LMU-5	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Forest (Fo)	Not Available	IVes	Trench cum bunding































Village	Survey No	Soil Reaction	Salinity	Organic Carbon	Available Phosphorus	Available Potassium	Available Sulphur	Available Boron	Available Iron	Available Manganese	Available Copper	Available Zinc
Fatthepura	2	Neutral (pH 6.5 - 7.3)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	Low (<145 kg/ha)	Medium (10 - 20 ppm)	Medium (0.5 - 1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Fatthepura	3	Neutral (pH 6.5 - 7.3)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	Low (<145 kg/ha)	Medium (10 - 20 ppm)	Medium (0.5 - 1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Fatthepura	4	Neutral (pH 6.5 - 7.3)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	Low (<145 kg/ha)	Medium (10 - 20 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Fatthepura	5	Neutral (pH 6.5 - 7.3)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	Low (<145 kg/ha)	Medium (10 - 20 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Fatthepura	11	Neutral (pH 6.5 - 7.3)	Non saline (<2 dsm)	Low (< 0.5 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	Medium (10 - 20 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Fatthepura	20	Neutral (pH 6.5 - 7.3)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	Low (<145 kg/ha)	Medium (10 - 20 ppm)	Medium (0.5 - 1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)



**Appendix III**  
**Kotageri-2 (1B2d) Microwatershed**  
**Soil Suitability Information**

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry	
Gajarakota	847	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Gajarakota	848	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Gajarakota	849	N1r	S2r	S3r	S2rt	S3r	S3t	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	N1n	S3r	S3r	S2r	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Gajarakota	855	N1r	S2r	S3r	S2rt	S3r	S3t	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	N1n	S3r	S3r	S2r	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Gajarakota	921	N1r	S2r	S3r	S2rt	S3r	S3t	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	N1n	S3r	S3r	S2r	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Gajarakota	922	N1r	S2r	S3r	S2rt	S3r	S3t	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	N1n	S3r	S3r	S2r	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Gajarakota	923	N1r	S2r	S3r	S2rt	S3r	S3t	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	N1n	S3r	S3r	S2r	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Gajarakota	924	N1r	S2r	S3r	S2rt	S3r	S3t	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	N1n	S3r	S3r	S2r	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Gajarakota	935	N1r	S2r	S3r	S2rt	S3r	S3t	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	N1n	S3r	S3r	S2r	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Gajarakota	937	N1r	S2r	S3r	S2rt	S3r	S3t	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	N1n	S3r	S3r	S2r	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Gajarakota	1139	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Kootagera	1	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	
Kootagera	2	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Kootagera	3	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Kootagera	4	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Kootagera	5	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Kootagera	6	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r	
Kootagera	7	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r	
Kootagera	8	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r	
Kootagera	9	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r	
Kootagera	10	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r	
Kootagera	11	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r	
Kootagera	12	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r	
Kootagera	13	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r	

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry
Kootagera	14	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r
Kootagera	15	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r
Kootagera	16	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r
Kootagera	17	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r
Kootagera	18	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r
Kootagera	26	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r
Kootagera	27	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r
Kootagera	28	S3r	S1	S2r	S1	S2r	S2r	S3r	S2r	S3t	S2r	S2r	S1	S2r	S1	S2r	S3r	S2r	S2t	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r
Kootagera	179	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r
Kootagera	181	N1n	S3nz	N1n	S3nz	N1n	S3nz	N1n	N1n	S3nz	N1n	S3nz	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3nz	N1n	N1n	N1n	N1n
Kootagera	182	N1n	S3nz	N1n	S3nz	N1n	S3nz	N1n	N1n	S3nz	N1n	S3nz	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3nz	N1n	N1n	N1n	N1n
Kootagera	183	N1n	S3nz	N1n	S3nz	N1n	S3nz	N1n	N1n	S3nz	N1n	S3nz	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3nz	N1n	N1n	N1n	N1n
Kootagera	184	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r
Kootagera	185	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r
Kootagera	186	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r
Kootagera	187	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r
Kootagera	188	N1n	S3nz	N1n	S3nz	N1n	S3nz	N1n	N1n	S3nz	N1n	S3nz	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3nz	N1n	N1n	N1n	N1n
Kootagera	189	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r
Kootagera	190	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r
Kootagera	191	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r
Kootagera	192	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r
Kootagera	193	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r
Kootagera	199	N1n	S3nz	N1n	S3nz	N1n	S3nz	N1n	N1n	S3nz	N1n	S3nz	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3nz	N1n	N1n	N1n	N1n
Kootagera	323	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r
Kootagera	324	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r
Kootagera	325	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r
Kootagera	326	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r





Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry
Kootagera	383	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Kootagera	384	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Kootagera	385	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Kootagera	386	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Kootagera	387	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Kootagera	388	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Kootagera	389	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Kootagera	390	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Kootagera	391	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Kootagera	392	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Kootagera	393	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Kootagera	394	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Kootagera	395	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Kootagera	396	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Kootagera	397	N1r	S3rg	N1r	S3rg	N1r	S3rg	N1r	N1r	S3rt	N1r	N1r	S3r	N1r	S3r	N1r	N1r	N1r	S3r	S3rg	S3rg	S3rg	S3rg	S3rg	S3rg	N1r	S3r	S3rg	S3rg	N1r
Kootagera	398	N1r	S3rg	N1r	S3rg	N1r	S3rg	N1r	N1r	S3rt	N1r	N1r	S3r	N1r	S3r	N1r	N1r	N1r	S3r	S3rg	S3rg	S3rg	S3rg	S3rg	S3rg	N1r	S3r	S3rg	S3rg	N1r
Kootagera	399	N1r	S3rg	N1r	S3rg	N1r	S3rg	N1r	N1r	S3rt	N1r	N1r	S3r	N1r	S3r	N1r	N1r	N1r	S3r	S3rg	S3rg	S3rg	S3rg	S3rg	S3rg	N1r	S3r	S3rg	S3rg	N1r
Kootagera	400	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Kootagera	401	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Kootagera	402	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Kootagera	403	N1r	S3rg	N1r	S3rg	N1r	S3rg	N1r	N1r	S3rt	N1r	N1r	S3r	N1r	S3r	N1r	N1r	N1r	S3r	S3rg	S3rg	S3rg	S3rg	S3rg	S3rg	N1r	S3r	S3rg	S3rg	N1r
Kootagera	404	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Kootagera	405	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Kootagera	406	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Kootagera	407	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Kootagera	408	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r



Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry	
Kootagera	409	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	
Kootagera	410	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Kootagera	411	N1r	S3rg	N1r	S3rg	N1r	S3rg	N1r	N1r	S3rt	N1r	N1r	S3r	N1r	S3r	N1r	N1r	N1r	S3r	S3rg	S3rg	S3rg	S3rg	S3rg	N1r	S3r	S3rg	S3rg	N1r	N1r	
Kootagera	412	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Kootagera	413	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Kootagera	414	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Kootagera	415	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Kootagera	416	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Kootagera	417	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Kootagera	418	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Kootagera	419	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Kootagera	420	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Kootagera	421	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Kootagera	422	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Kootagera	423	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Kootagera	424	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Kootagera	425	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Kootagera	426	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Kootagera	428	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Kootagera	429	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Kootagera	433	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Kootagera	434	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Kootagera	435	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Kootagera	436	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Kootagera	437	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Kootagera	442	N1r	S2r	S3r	S2rt	S3r	S3t	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	N1n	S3r	S3r	S2r	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Kootagera	443	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry	
Kootagera	444	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Kootagera	445	N1r	S2r	S3r	S2rt	S3r	S3t	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	N1n	S3r	S3r	S2r	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Kootagera	446	N1r	S2r	S3r	S2rt	S3r	S3t	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	N1n	S3r	S3r	S2r	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Kootagera	447	Other s	Other s	Other s	Other s	Other s	Other s	Other s	Other s	Other s	Other s	Other s	Other s	Other s	Other s	Other s	Other s	Other s	Other s	Other s	Other s	Other s	Other s	Other s	Other s	Other s	Other s	Other s	Other s	Other s	
Kootagera	448	N1r	S3rg	N1r	S3rg	N1r	S3rg	N1r	N1r	S3rt	N1r	N1r	S3r	N1r	S3r	N1r	N1r	N1r	S3r	S3rg	S3rg	S3rg	S3rg	S3rg	N1r	S3r	S3rg	S3rg	N1r	N1r	
Kootagera	449	N1r	S3rg	N1r	S3rg	N1r	S3rg	N1r	N1r	S3rt	N1r	N1r	S3r	N1r	S3r	N1r	N1r	N1r	S3r	S3rg	S3rg	S3rg	S3rg	S3rg	N1r	S3r	S3rg	S3rg	N1r	N1r	
Kootagera	450	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Kootagera	452	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Kootagera	454	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Kootagera	455/ 1	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Kootagera	455/ 2	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Kootagera	455/ 3	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Kootagera	455/ 4	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Kootagera	456	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Kootagera	457	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Kootagera	458	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Kootagera	459	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Kootagera	460	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Kootagera	461	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Kootagera	462	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Kootagera	463	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Kootagera	464	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Kootagera	465	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Kootagera	466	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Kootagera	467	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Kootagera	468	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry
Kootagera	469	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Kootagera	470	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Kootagera	471	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Kootagera	472	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Kootagera	473	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Kootagera	474	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Kootagera	475	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Kootagera	476/1	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Kootagera	476/2	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Kootagera	476/3	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Kootagera	476/4	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Kootagera	476/5	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Kootagera	477	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r
Kootagera	478	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r
Kootagera	479	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r
Kootagera	480	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r
Kootagera	481	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r
Kootagera	482	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r
Kootagera	483	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r
Kootagera	484	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r
Kootagera	485	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r
Kootagera	486	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r
Kootagera	487	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r
Kootagera	488	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r
Kootagera	489	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r
Kootagera	490	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry	
Kootagera	491	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Kootagera	492	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Kootagera	493	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Kootagera	494	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Kootagera	495	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Kootagera	496	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Kootagera	497	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Kootagera	498	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Kootagera	499	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Kootagera	500	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Kootagera	532	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Kootagera	533	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Kootagera	534	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Kootagera	535	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Kootagera	536	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Kootagera	537	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Kootagera	538	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Kootagera	541	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Kootagera	542	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Kootagera	543	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Kootagera	544	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Kootagera	545	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Kootagera	546	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Kootagera	547	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Kootagera	548	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Kootagera	549	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Kootagera	550	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry
Kootagera	551	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r
Kootagera	552	Other s	Other s	Other s	Other s	Other s	Other s	Other s	Other s	Other s	Other s	Other s	Other s	Other s	Other s	Other s	Other s	Other s	Other s	Other s	Other s	Other s	Other s	Other s	Other s	Other s	Other s	Other s	Other s	Other s
Kootagera	553	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r
Kootagera	554	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r
Kootagera	555	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r
Kootagera	556	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r
Kootagera	557	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S3n	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r
Fatthepura	1	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Fatthepura	2	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Fatthepura	3	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Fatthepura	4	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Fatthepura	5	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Fatthepura	11	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Fatthepura	20	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r

# **PART-B**

**SOCIO-ECONOMIC STATUS OF FARM HOUSEHOLDS**





## CONTENTS

1.	Findings of the socio-economic survey	1-3
2.	Introduction	5
3	Methodology	7-8
4	Salient features of the survey	9-31
5	Summary	33-37



## LIST OF TABLES

1	Households sampled for socio economic survey	9
2	Population characteristics	9
3	Age wise classification of household members	9
4	Education level of household members	10
5	Occupation of household heads	10
6	Occupation of family members	11
7	Institutional participation of household members	11
8	Type of house owned by households	11
9	Durable assets owned by households	12
10	Average value of durable assets owned by households	12
11	Farm implements owned by households	12
12	Average value of farm implements	13
13	Livestock possession by households	13
14	Average labour availability	13
15	Adequacy of hired labour	14
16	Distribution of land (ha)	14
17	Average land value (Rs./ha)	14
18	Status of bore wells	14
19	Status of open wells	15
20	Source of irrigation	15
21	Depth of water(Avg in meters)	15
22	Irrigated area (ha)	15
23	Cropping pattern	15
24	Cropping intensity	16
25	Possession of bank account and savings	16
26	Borrowing status	16
27	Source of credit borrowed by households	16
28	Avg. Credit amount	17
29	Purpose of credit borrowed (institutional Source)	17
30	Purpose of credit borrowed (Private Source)	17
31	Repayment status of household (institutional Source)	17
32	Repayment status of household (Private Source)	18
33	Opinion regarding institutional sources of credit	18
34	Opinion regarding Non- institutional sources of credit	18
35.a	Cost of cultivation of Red gram	19

35.b	Cost of cultivation of paddy	20
35.c	Cost of cultivation of Green gram	21
35.d	Cost of cultivation of Groundnut	22
35.e	Cost of cultivation of Cotton	23
36	Adequacy of fodder	24
37	Annual gross income	24
38	Average annual expenditure	24
39	Horticultural species grown	24
40	Interest towards cultivation of horticulture crops	25
41	Forest species grown	25
42	Average additional investment capacity	25
43	Source of funds for additional investment	26
44	Marketing of the agricultural produce	26
45	Marketing channels used for sale of agricultural produce	26
46	Mode of transport of agricultural produce	27
47	Incidence of soil and water erosion problems	27
48	Interest shown towards soil testing	27
49	Soil and water conservation practices and structures adopted	27
50	Status of soil and water conservation structures	27
51	Agencies involved in the soil and water conservation structures	28
52	Usage pattern of fuel for domestic use	28
53	Source of drinking water	28
54	Source of light	28
55	Existence of sanitary toilet facility	29
56	Possession of public distribution system (PDS) card	29
57	Participation in NREGA programme	29
58	Adequacy of food items	29
59	Inadequacy of food items	30
60	Response on market surplus of food items	30
61	Farming constraints experienced	30

**FINDINGS OF THE SOCIO-ECONOMIC SURVEY**

- ❖ *The survey was conducted in Kotageri-2 is located at North latitude 16° 54' 2.29" and 16° 52' 44.243" and East longitude 77° 15' 21.73" and 77° 13' 47.096" covering an area of about 301.61 ha coming unde Kootagera, Gajarakota and Fatthepura Villages of Yadagiri taluk.*
- ❖ *Socio-economic analysis of Kotageri-2 micro watersheds of Shivapur sub-watershed, Yadgiri taluk & District indicated that, out of the total sample of 36 total respondents, 15 (41.67 %) were marginal, 10 (27.78%)were small, 5 (13.89 %) were Semi medium and 2 (5.56 %) were medium and 1 (2.78 %) were large farmers.*
- ❖ *The population characteristics of households indicated that, there were 108 (54.82%) men and 89 (45.18 %) were women. The average population of landless was 4.7, marginal farmers were 4.7, small farmers were 6.3, semi medium farmers were 4.5 and medium farmers were 10.*
- ❖ *Majority of the respondents (47.21%) were in the age group of 16-35 years.*
- ❖ *Education level of the sample households indicated that, there were 37.06 per cent illiterates, 67.02 per cent pre university education and 5.08 per cent attained graduation.*
- ❖ *About, 69.44 per cent of household heads practicing agriculture and 16.67 per cent of the household heads were engaged as agricultural labourers.*
- ❖ *Agriculture was the major occupation for 46.19 per cent of the household members.*
- ❖ *In the study area, 55.56 per cent of the households possess katcha house and 38.89 per cent possess pucca house.*
- ❖ *The durable assets owned by the households showed that, 58.33 per cent possess TV, 2.78 per cent possess mixer grinder, 91.67 per cent possess mobile phones and 22.22 per cent possess motor cycles.*
- ❖ *Farm implements owned by the households indicated that, 44.44 per cent of the households possess plough, 8.33 per cent possess tractor, 19.44 per cent possess bullock cart and 38.89 per cent possess sprayer.*
- ❖ *Regarding livestock possession by the households, 25.00 per cent possess local cow and 19.44 per cent possess buffalo.*
- ❖ *The average labour availability in the study area showed that, own men and women labour availability in the micro watershed was 1.56 and 2.06 each, while the hired labour (women) availability was 12.94 and the hired labour (men) availability was 8.57.*
- ❖ *Further, 11.11 per cent of the households opined that hired labour was inadequate during the agricultural season.*

- ❖ *Out of the total land holding of the sample respondents 42.05 per cent (38.73 ha) of the area is under dry condition and the remaining 56.91 per cent area is irrigated land.*
- ❖ *There were 11.00 live bore wells and 12.00 dry bore wells among the sampled households.*
- ❖ *Bore well was the major source of irrigation for 30.56 per cent of the households.*
- ❖ *The major crops grown by sample farmers are Red gram, Paddy, Green gram, Groundnut and Cotton and cropping intensity was recorded as 100.00 per cent.*
- ❖ *Out of the sample households 94.44 percent possessed bank account and 41.67 per cent of them have savings in the account.*
- ❖ *About 83.33 per cent of the respondents borrowed credit from various sources.*
- ❖ *Among the credit borrowed by households, 22.22 per cent have borrowed loan from Cooperative bank , 100 per cent have borrowed loan from Grameena Bank, and 33.33 per cent have borrowed loan from money lender.*
- ❖ *Majority of the respondents (92.86%) have borrowed loan for agriculture purpose.*
- ❖ *Regarding the opinion on institutional sources of credit, 64.29 per cent of the households opined that credit helped to perform timely agricultural operations, while, only 7.14 per cent respondents opined that loan amount was adequate to fulfil their requirement.*
- ❖ *The per hectare cost of cultivation for Red gram, Paddy, Green gram, Groundnut and Cotton was Rs.45496.44 , 93955.24, 54349.84, 45538.76 and 45070.85 with benefit cost ratio of 1:1.20, 1: 1.10, 1: 0.92, 1: 1.20 and 1:1.80 respectively.*
- ❖ *Further, 41.67 per cent of the households opined that dry fodder was adequate .*
- ❖ *The average annual gross income of the farmers was Rs. 137441.67 in micro-watershed, of which Rs. 60461.11 comes from agriculture.*
- ❖ *Sampled households have grown 10 horticulture trees and 12 forestry trees together in the fields and back yards.*
- ❖ *Households have an average investment capacity of Rs. 2777.78 for land development and Rs. 4166.67 for irrigation facility.*
- ❖ *Source of funds for additional investment is concerned, 2.78 per cent a Government subsidy for land development activities.*
- ❖ *Regarding marketing channels, 44.44 per cent of the households have sold agricultural produce to the local/village merchants, while, 55.56 per cent have sold in regulated markets.*
- ❖ *Further, 77.78 per cent of the households have used tractor for the transport of agriculture commodity.*
- ❖ *Majority of the farmers (44.44%) have experienced soil and water erosion problems in the watershed and 83.33 per cent of the households were interested towards soil testing.*

- ❖ *About, 92.89 per cent of farmers practicing summer ploughing as soil and water conservation practice.*
- ❖ *Fire was the major source of fuel for domestic use for 47.22 per cent of the households and 50.00 per cent households has LPG connection.*
- ❖ *Piped supply was the major source for drinking water for 86.11 per cent of the households.*
- ❖ *Electricity was the major source of light for 100.00 per cent of the households.*
- ❖ *In the study area, 41.67 per cent of the households possess toilet facility.*
- ❖ *Regarding possession of PDS card, 91.67 per cent of the households possessed BPL card, 2.78 per cent of the household's possessed APL card and 5.56 per cent of the household's were not having ration cards.*
- ❖ *Households opined that, the requirement of cereals (94.44%), pulses (83.33%) and oilseeds (63.89%) are adequate for consumption.*
- ❖ *Farming constraints experienced by households in the micro watersheds were lower fertility status of the soil (75.00%) wild animal menace on farm field (66.67%), frequent incidence of pest and diseases (83.33%), inadequacy of irrigation water (61.11%), high cost of fertilizers and plant protection chemicals (88.89%), high rate of interest on credit (83.33%), low price for the agricultural commodities (83.33%), lack of marketing facilities in the area (63.89%), inadequate extension services (30.56%) and lack of transport for safe transport of the agricultural produce to the market (63.89%).*





## INTRODUCTION

Soil and water are the two precious natural resources which are essential for crop production and existence of life on earth. Rainfed agriculture is under severe stress due to various constraints related to agriculture like uneven and erratic distribution of rainfall, indiscriminate use of fertilizers, chemicals and pesticides, adoption of improper land management practices, soil erosion, decline in soil fertility, decline in ground water resources leading to low crop productivity. The area under rainfed agriculture has to be managed effectively using the best available practices to enhance the production of food, fodder and fuel. This is possible if the land resources are characterized at each parcel of land through detailed land resource inventory using the best available techniques of remote sensing, GPS and GIS. The watershed development programs are aimed at the sustainable distribution of its resources and the process of creating and implementing plans, programs, and projects to sustain and enhance watershed functions that affect the plant, animal and human communities within a watershed boundary.

World Bank funded KWDP II, SUJALA III project was implemented in with Broad objective of demonstrating more effective watershed management through greater integration of programmes related to rain-fed agriculture, innovative and science based approaches and strengthen institutional capacities and If successful, it is expected that the systems and tools could be mainstreamed into the overall IWMP in the State of Karnataka and in time, throughout other IWMP operations in India. With this background the socio-economic survey has been carried out with following specific objectives:

1. To understand the demographic features of the households in the micro-watershed
2. To understand the extent of family labour available and additional employment opportunities available within the village.
3. To know the status of assets of households in the micro-watershed for suggesting possible improvements.
4. To study the cropping pattern, cropped area and productivity levels of different households in micro-watershed.
5. To determine the type and extent of livestock owned by different categories of HHs
6. Availability of fodder and level of livestock management.

### **Scope and importance of survey**

Survey helps in identification of different socio-economic and resource use-patterns of farmers at the Micro watershed. Household survey provides demographic features, labor force, and levels of education; land ownership and asset position (including livestock and other household assets) of surveyed households; and cropping patterns, input intensities, and average crop yields from farmers' fields. It also discusses crop utilization and the degree of commercialization of production in the areas; farmers' access to and utilization of credit from formal and informal sources; and the level of adoption and use of soil, water, and pest management technologies.



## **METHODOLOGY**

The description of the methods, components selected for the survey and procedures followed in conducting the baseline survey are furnished under the following heads.

### **1. Description of the study area**

Yadgir District is one of the 30 districts of Karnataka state in southern India. This district was carved out from the erstwhile Gulbarga district as the 30th district of Karnataka on 10 April 2010. Yadgir town is the administrative headquarters of the district. The district comprises of 3 taluks namely, Shahapur, Yadgiri and Shorapur (There are 16 hoblies, 117 Gram Panchayats, 4 Municipalities, 8 Towns/ Urban agglomeration and 487 inhabited & 32 un-inhabited villages The district occupies an area of 5,160.88 km<sup>2</sup>.

Yadgir district is the second smallest district in the state, area wise is very rich in cultural traditions. The vast stretch of fertile black soil of the district is known for bumper red gram and jowar crops. The district is a "Daal bowl" of the state. The district is also known for cluster of cement industries and a distinct stone popularly known as "Malakheda Stone". Two main rivers, Krishna and Bhima, and a few tributaries flow in this region. Krishna and Bhima Rivers drain the district. They constitute the two major river basins of the district. Kagna and Amarja are the two sub - basins of Bhima River, which occur within the geographical area of the district

According to the 2011 census Yadgir district has a population of 1, 172,985, roughly equal to the nation of Timor-Leste or the US state of Rhode Island. This gives it a ranking of 404th in India (out of a total of 640). The district has a population density of 224 inhabitants per square kilometre (580/sq mi). Its population growth rate over the decade 2001-2011 was 22.67%. Yadgir has a sex ratio of 984 females for every 1000 males, and a literacy rate of 52.36%.

### **2. Locale of the survey and description of the micro-watershed and**

The study was conducted in Kotageri-2 micro-watershed (Shivapur sub-watershed, Yadgiri taluk & District) is located at North latitude 16<sup>o</sup> 54' 2.29" and 16<sup>o</sup> 52' 44.243" and East longitude 77<sup>o</sup> 15' 21.73" and 77<sup>o</sup> 13' 47.096" covering an area of about 301.61 ha bounded by unde Kootagera, Gajarakota and Fatthepura Villages.

### **3. Selection of the respondents for the study**

The micro-watershed is marked with 320 square meters grids. One farmer from every alternate grid in the micro-watershed was selected for the study and interviewed for socio-economic data. Totally 36 households were interviewed for the survey.

#### **4. The parameters considered for socio-economic survey of households**

Two forms of data were collected from the micro-watershed which includes primary data from the farm households and secondary data about the villages under the micro-watershed jurisdiction.

The following parameters were considered for the primary data collection about the socio-economic data of the households, (1) Demographic information, (2) Farm and durable assets owned by households, (3) Livestock possession, (4) Labour availability, (5) Level of migration in the village, Land holding, (7) Cropping pattern, (8) Source of irrigation, (9) Borrowing status, (10) Cost of cultivation of major crops, (11) Economics of subsidiary activities, (12) Fodder availability, (13) Family annual income from different sources, (14) Horticulture and forestry species grown, (15) Additional investment capacity, (16) Marketing practices, (17) Status of soil and water conservation structure, (18) Access to basic needs and (19) Constraints and suggestion.

The following parameters were considered for the secondary data regarding the villages under the micro-watershed jurisdiction, (1) Number of villages in each micro-watershed jurisdiction, (2) Village wise number of households, (3) Geographical area of the villages, (4) Cultivable area including rainfed and irrigated, (5) Number and type of house in each village, (6) Human and livestock population, (7) Facilities in the village such as roads, transport facility for conveyance, drinking water supply, street light and (8) Community based organizations in the villages.

#### **5. Development of interview schedule and data collection**

Taking into the consideration the objectives of the survey, an interview schedule was prepared after thorough consultation with the experts in the field of social sciences. A comprehensive interview schedule covering all the major parameters for measuring the socio-economic situation was developed.

#### **6. Tools used to analyze the data**

The statistical components such as frequency and percentage were used to analyze the data.

#### **Abbreviations used in the report**

LL=Landless

MF=Marginal Farmers

SF=Small farmers

SMF=Semi medium farmers

MDF=Medium farmers

LF=Large Farmers

## FINDINGS OF THE SURVEY

This chapter deals with systematic presentation of results of the survey. Keeping in view the objectives, the salient features of the survey are presented under the following headings.

**Households sampled for socio-economic survey:** The data on households sampled for socio economic survey in Kotageri-2 Micro watershed is presented in Table 1 and it indicated that 36 farmers were sampled in Kotageri-2 micro-watershed among households surveyed 15 (41.67%) were marginal, 10 (27.78%) were small, 5 (13.89 %) were semi medium, 2 (5.56 %) were medium and 1 (2.78 %) were large farmers. 3 landless farmers were also interviewed for the survey.

**Table 1. Households sampled for socio economic survey in Kotageri-2 micro-watershed**

Sl. No.	Particulars	LL (3)		MF (15)		SF (10)		SMF (5)		MDF (2)		LF (1)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Farmers	3	8.33	15	41.7	10	27.8	5	13.9	2	5.56	1	3	36	100

**Population characteristics:** The population characteristics of households sampled for socio-economic survey in Kotageri-2 Micro watershed is presented in Table 2. The data indicated that, there were 108 (54.82%) men and 89 (45.18%) were women. The average population of landless was 4.7, marginal farmers were 4.7, small farmers were 6.3, semi medium farmers were 4.5 and medium farmers were 10.

**Table 2. Population characteristics in Kotageri-2 micro-watershed**

Sl. No.	Particulars	LL (14)		MF (71)		SF (63)		SMF (30)		MDF (9)		LF (10)		All (197)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Men	7	50	35	49	38	60	16	53.3	5	55.6	7	70	108	54.8
2	Women	7	50	36	51	25	40	14	46.7	4	44.4	3	30	89	45.2
Total		14	100	71	100	63	100	30	100	9	100	10	100	197	100
Average		4.7		4.7		6.3		6.0		4.5		10.0		5.5	

**Age wise classification of population:** The age wise classification of household members in Kotageri-2 Micro watershed is presented in Table 3. The indicated that, 35 (17.77%) of population were 0-15 years of age, 93 (47.21%) were 16-35 years of age, 49(24.87%) were 36-60 years of age and 20 (10.15 %) were above 61 years of age.

**Table 3: Age wise classification of members of the household in Kotageri-2 micro-watershed**

Sl. No.	Particulars	LL (14)		MF (71)		SF (63)		SMF (30)		MDF (9)		LF (10)		All (197)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	0-15 years of age	7	50	14	19.7	11	17.5	3	10	0	0	0	0	35	17.77
2	16-35 years of age	4	28.6	33	46.5	33	52.4	17	56.67	5	56	1	10	93	47.21
3	36-60 years of age	3	21.4	15	21.1	13	20.6	8	26.67	3	33	7	70	49	24.87
4	> 61 years	0	0	9	12.7	6	9.52	2	6.67	1	11	2	20	20	10.15
Total		14	100	71	100	63	100	30	100	9	100	10	100	197	100

**Education level of household members:** Education level of household members in Kotageri-2 Micro watershed is presented in Table 4. The results indicated that, there were 37.06 per cent of illiterates, 28.43 per cent of them had primary school education, 2.54 per cent middle school education, and 13.71 per cent high school education, 11.17 per cent of them had PUC education, 0.51 per cent of them had Diploma, 5.08 per cent attained graduation, and 0.51 them had other education.

**Table 4. Education level of members of the household in Kotageri-2 micro-watershed**

Sl. No.	Particulars	LL (14)		MF (71)		SF (63)		SMF (30)		MDF (9)		LF (10)		All (197)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Illiterate	2	14.3	23	32.4	29	46	12	40	4	44.44	3	30	73	37.1
2	Primary School	11	78.6	21	29.6	15	23.8	4	13.3	1	11.11	4	40	56	28.4
3	Middle School	0	0	2	2.82	2	3.17	1	3.33	0	0	0	0	5	2.54
4	High School	0	0	8	11.3	10	15.9	6	20	1	11.11	2	20	27	13.7
5	PUC	1	7.14	12	16.9	4	6.35	3	10	2	22.22	0	0	22	11.2
6	Diploma	0	0	0	0	0	0	0	0	1	11.11	0	0	1	0.51
7	ITI	0	0	0	0	0	0	2	6.67	0	0	0	0	2	1.02
8	Degree	0	0	4	5.63	3	4.76	2	6.67	0	0	1	10	10	5.08
9	Others	0	0	1	1.41	0	0	0	0	0	0	0	0	1	0.51
Total		14	100	71	100	63	100	30	100	9	100	10	100	197	100

**Occupation of head of households:** The data regarding the occupation of the household heads in Kotageri-2 Micro watershed is presented in Table 5. The results indicate that, 69.44 per cent of household heads were practicing agriculture, 16.67 per cent of the household heads were agricultural Labour.

**Table 5: Occupation of heads of households in Kotageri-2 micro-watershed**

Sl. No.	Particulars	LL (3)		MF (15)		SF (10)		SMF (5)		MDF (2)		LF (1)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Agriculture	0	0	11	73	8	80	5	100	0	0	1	100	25	69.44
2	Agricultural Labour	1	33	2	13	2	20	0	0	1	50	0	0	6	16.67
3	General Labour	1	33	0	0	0	0	0	0	0	0	0	0	1	2.78
4	Household industry	0	0	1	6.7	0	0	0	0	0	0	0	0	1	2.78
5	Artisans	1	33	0	0	0	0	0	0	0	0	0	0	1	2.78
6	Trade & Business	0	0	1	6.7	0	0	0	0	1	50	0	0	2	5.56
Total		3	100	15	100	10	100	5	100	2	100	1	100	36	100

**Occupation of the members of the household:** The data regarding the occupation of the household members in Kotageri-2 Micro watershed is presented in Table 6. The results indicate that, agriculture was the major occupation for 46.19 per cent of the household members, 12.18 per cent were agricultural labour, 2.54 per cent were general labour, 27.41 per cent were working in pursuing education, 7.61 per cent were involved as housewife, and 0.51 per cent were children.



**Table 6: Occupation of members of the household in Kotageri-2 micro-watershed**

Sl. No.	Particulars	LL (14)		MF (71)		SF (63)		SMF (30)		MDF (9)		LF (10)		All (197)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Agriculture	0	0	32	45.1	31	49.21	16	53.33	3	33	9	90	91	46.2
2	Agricultural Labour	1	7.14	6	8.45	11	17.46	4	13.33	2	22	0	0	24	12.2
3	General Labour	3	21.4	1	1.41	1	1.59	0	0	0	0	0	0	5	2.54
4	Household industry	0	0	1	1.41	0	0	0	0	0	0	0	0	1	0.51
5	Artisans	2	14.3	0	0	0	0	0	0	0	0	0	0	2	1.02
6	Private Service	0	0	1	1.41	0	0	1	3.33	0	0	0	0	2	1.02
7	Trade & Business	0	0	1	1.41	0	0	0	0	1	11	0	0	2	1.02
8	Student	8	57.1	24	33.8	15	23.81	5	16.67	2	22	0	0	54	27.4
9	Housewife	0	0	4	5.63	5	7.94	4	13.33	1	11	1	10	15	7.61
10	Children	0	0	1	1.41	0	0	0	0	0	0	0	0	1	0.51
Total		14	100	71	100	63	100	30	100	9	100	10	100	197	100

**Institutional Participation of household members:** The data regarding the institutional participation of the household members in Kotageri-2 Micro watershed is presented in Table 7. The results show that, out of the total family members in the households 2.54 per cent of them are participating in self help group.

**Table 7: Institutional Participation of household member in Kotageri-2 micro-watershed**

Sl.No.	Particulars	LL (14)		MF (71)		SF (63)		SMF (30)		MDF (9)		LF (10)		All (197)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Self Help Group	0	0	2	2.82	1	1.59	0	0	2	22.22	0	0	5	2.54
2	No Participation	14	100	69	97.2	62	98.4	30	100	7	77.78	10	100	192	97.5
Total		14	100	71	100	63	100	30	100	9	100	10	100	197	100

**Type of house owned:** The data regarding the type of house owned by the households in Kotageri-2 Micro watershed is presented in Table 8. The results indicate that, 5.56 percent possess thatched house, 55.56 per cent of the households possess Katcha house and 38.89 per cent possess pucca house.

**Table 8. Type of house owned by households in Kotageri-2 micro-watershed**

Sl. No.	Particulars	LL (3)		MF (15)		SF (10)		SMF (5)		MDF (2)		LF (1)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Thatched	0	0	1	6.7	1	10	0	0	0	0	0	0	2	5.56
2	Katcha	2	67	8	53	8	80	2	40	0	0	0	0	20	55.56
3	Pucca/RCC	1	33	6	40	1	10	3	60	2	100	1	100	14	38.89
Total		3	100	15	100	10	100	5	100	2	100	1	100	36	100

**Durable assets owned by the households:** The data regarding the Durable Assets owned by the households in Kotageri-2 Micro watershed is presented in Table 9. The results shows that, 58.33 per cent possess TV, 2.78 per cent possess mixer grinder, 5.56 per cent possess refrigerator, 8.33 per cent possess Bicycle, 22.22 per cent possess motor cycle, and 91.67 per cent possess mobile phones.

**Table 9. Durable assets owned by households in Kotageri-2 micro-watershed**

Sl. No.	Particulars	LL (3)		MF (15)		SF (10)		SMF (5)		MDF (2)		LF (1)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Television	1	33	9	60	5	50	3	60	2	100	1	100	21	58.33
2	Mixer/Grinder	0	0	0	0	0	0	1	20	0	0	0	0	1	2.78
3	Refrigerator	0	0	1	6.7	0	0	0	0	1	50	0	0	2	5.56
4	Bicycle	1	33	1	6.7	1	10	0	0	0	0	0	0	3	8.33
4	Motor Cycle	0	0	1	6.7	3	30	3	60	0	0	1	100	8	22.22
5	Auto	0	0	0	0	1	10	0	0	0	0	0	0	1	2.78
6	Mobile Phone	3	100	13	87	10	100	4	80	2	100	1	100	33	91.67
7	Blank	0	0	2	13	0	0	1	20	0	0	0	0	3	8.33

**Average value of durable assets:** The data regarding the average value of durable assets owned by the households in Kotageri-2 Micro watershed is presented in Table 10. The result shows that, the average value of television was Rs.4833.00, mixer grinder was Rs.2000.00, refrigerator was 8000.00, bicycle was Rs.1666.00, motor cycle was Rs. 51250.00 and mobile phone was Rs.2250.00.

**Table 10. Average value of durable assets owned in Kotageri-2 micro-watershed**

Average Value (Rs.)

Sl.No.	Particulars	LL (3)	MF (15)	SF (10)	SMF (5)	MDF (2)	LF (1)	All (36)
1	Television	7000	4222	5200	5166	4500	6000	4833
2	Mixer/Grinder	0	0	0	2000	0	0	2000
3	Refrigerator	0	8000	0	0	8000	0	8000
4	Bicycle	2000	1500	1500	0	0	0	1666
5	Motor Cycle	0	60000	51666	48333	0	50000	51250
6	Auto	0	0	80000	0	0	0	80000
7	Mobile Phone	2375	1843	2125	2062	3375	3000	2250

**Farm implements owned:** The data regarding the farm implements owned by the households in Kotageri-2 Micro watershed is presented in Table 11. About 19.44 per cent of the households possess Bullock Cart, 44.44 per cent possess plough and 33.33 per cent possess Seed/Fertilizer Drill and Sprinkler, 38.89 per cent possess Sprayer, 44.44 per cent possess Weeder, 8.33 per cent possess tractor.

**Table 11. Farm implements owned in Kotageri-2 micro-watershed**

Sl. No.	Particulars	LL (3)		MF (15)		SF (10)		SMF (5)		MDF (2)		LF (1)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Bullock Cart	0	0	2	13.3	3	30	0	0	1	50	1	100	7	19.44
2	Plough	0	0	5	33.3	7	70	2	40	1	50	1	100	16	44.44
3	Seed/Fertilizer Drill	0	0	2	13.3	6	60	2	40	1	50	1	100	12	33.33
4	Power Tiller	0	0	0	0	1	10	0	0	0	0	1	100	2	5.56
5	Tractor	0	0	0	0	1	10	1	20	0	0	1	100	3	8.33
6	Sprayer	0	0	2	13.3	6	60	4	80	1	50	1	100	14	38.89
7	Weeder	0	0	4	26.7	8	80	3	60	1	50	0	0	16	44.44
8	Maize Huller	0	0	0	0	1	10	0	0	0	0	0	0	1	2.78
9	Blank	3	100	9	60	1	10	1	20	1	50	0	0	15	41.67

**Average value of farm implements:** The data regarding the average value of farm Implements owned by the households in Kotageri-2 Micro watershed is presented in Table 12. The results show that the average value of plough was Rs.3375.00, bullock Cart was Rs.21142.00, seed/fertilizer drill was Rs.7850.00, sprayer was Rs.2835.00, weeder was Rs.160.00 and tractor was Rs. 650000.

**Table 12. Average value of farm implements in Kotageri-2 micro-watershed**

Average Value (Rs.)								
Sl. No.	Particulars	LL (3)	MF (15)	SF (10)	SMF (5)	MDF(2)	LF (1)	All (36)
1	Bullock Cart	0	20000	22666	0	16000	24000	21142
2	Plough	0	2200	4428	2250	3500	4000	3375
3	Seed/Fertilizer Drill	0	3300	12166	3250	3600	4500	7850
4	Power Tiller	0	0	100000	0	0	58000	79000
5	Tractor	0	0	400000	800000	0	750000	650000
6	Sprayer	0	2600	2766	2900	2800	3500	2835
7	Weeder	0	88	222	98	100	0	160
8	Maize Huller	0	0	65000	0	0	0	65000

**Livestock possession by the households:** The data regarding the Livestock possession by the households in Kotageri-2 Micro watershed is presented in Table 13. The indicate that, 36.11 per cent of the households possess bullocks, 25.00 per cent possess local cow, 19.44 per cent possess buffalo, 2.78 per cent possess crossbred cow, 13.89 per cent possess goat, and 22.22 per cent were poultry birds.

**Table 13. Livestock possession by households in Kotageri-2 micro-watershed**

Sl. No.	Particulars	LL (3)		MF (15)		SF (10)		SMF (5)		MDF (2)		LF (1)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Bullock	0	0	3	20	6	60	2	40	1	50	1	100	13	36.11
2	Local cow	1	33	1	6.7	6	60	0	0	0	0	1	100	9	25
3	Crossbred cow	1	33	0	0	0	0	0	0	0	0	0	0	1	2.78
4	Buffalo	0	0	1	6.7	3	30	2	40	1	50	0	0	7	19.44
5	Goat	0	0	1	6.7	2	20	1	20	1	50	0	0	5	13.89
6	Poultry birds	0	0	2	13	4	40	0	0	1	50	1	100	8	22.22
7	blank	2	67	11	73	1	10	2	40	0	0	0	0	16	44.44

**Average Labour availability:** The data regarding the average labour availability in Kotageri-2 Micro watershed is presented in Table 14. The indicated that, own labour men available in the micro watershed was 2.06, women available in the micro watershed was 1.56, hired labour (men) available was 8.57 and hired labour (women) available was 12.94.

**Table 14. Average labour availability in Kotageri-2 micro-watershed**

Sl. No.	Particulars	LL (3)	MF (15)	SF (10)	SMF (5)	MDF (2)	LF (1)	All (36)
		N	N	N	N	N	N	N
1	Hired labour Female	0	12.5	14.8	13	20	25	12.94
2	Own Labour Female	0	1.4	1.9	2.2	1.5	2	1.56
3	Own labour Male	0	1.47	2.8	2.6	2	7	2.06
4	Hired labour Male	0	7.86	10.2	9.6	15	10	8.57

**Adequacy of hired labour:** The data regarding the adequacy of hired labour in Kotageri-2 Micro watershed is presented in Table 15. The results indicate that, 88.89 per cent of the household opined that hired labour was adequate, and 11.11 per cent of the household opined that hired labour was Inadequate.

**Table 15. Adequacy of hired labour in Kotageri-2 micro-watershed**

Sl. No.	Particulars	LL (3)		MF (15)		SF (10)		SMF (5)		MDF (2)		LF (1)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Adequate	3	100	15	100	8	80	4	80	2	100	0	0	32	88.9
2	Inadequate	0	0	0	0	2	20	1	20	0	0	1	100	4	11.1

**Distribution of land (ha):** The data regarding the distribution of land (ha) in Kotageri-2 Micro watershed is presented in Table 21. The results indicate that, 16.29 ha (42.05%) of dry land and 22.04 ha (56.91 %) of irrigated land.

**Table 16. Distribution of land (ha) in Kotageri-2 micro-watershed**

Sl. No.	Particulars	MF (15)		SF (10)		SMF (5)		MDF (2)		LF (1)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Dry	6.17	85.92	7.59	66.21	2.53	30.57	0	0	0	0	16.29	42.05
2	Irrigated	0.61	8.45	3.87	33.79	5.73	69.43	5.35	100	6.48	100	22.04	56.91
3	Permanent Fallow	0.4	5.63	0	0	0	0	0	0	0	0	0.4	1.04
Total		7.18	100	11.46	100	8.26	100	5.35	100	6.48	100	38.73	100

**Average value of land (ha):** The data regarding the average land value (Rs./ha) in Kotageri-2 Micro watershed is presented in Table 17. The results show that the average value of dry land was Rs.687475.15 and the average value of irrigated land was Rs.453997.43.

**Table 17. Average value of land (ha) in Kotageri-2 micro-watershed**

Sl. No.	Particulars	LL (3)	MF (15)	SF (10)	SMF (5)	MDF (2)	LF (1)	All (36)
		N	N	N	N	N	N	N
1	Dry	0	1149967	474240	197916.7	0	0	687475.2
2	Irrigated	0	1152667	1032393	592660.6	151338.9	169812.5	453997.4
3	Permanent Fallow	0	1152667	0	0	0	0	1152667

**Status of bore wells:** The data regarding the status of bore wells in Kotageri-2 Micro watershed is presented in Table 18. The results indicate that, there were 12 De-functioning bore wells and 11 functioning bore wells among the sampled households in micro watershed.

**Table 18. Status of bore wells in Kotageri-2 micro-watershed**

Sl. No.	Particulars	LL (3)	MF (15)	SF (10)	SMF (5)	MDF (2)	LF (1)	All (36)
		N	N	N	N	N	N	N
1	De-functioning	0	0	3	2	1	6	12
2	Functioning	0	0	4	3	2	2	11

**Status of open wells:** The data regarding the status of open wells in Kotageri-2 Micro watershed is presented in Table 19. The results indicate that, there were 1 functioning open wells among the sampled households in micro watershed.

**Table 19. Status of open wells in Kotageri-2 micro-watershed**

Sl. No.	Particulars	LL (3)	MF (15)	SF (10)	SMF (5)	MDF (2)	LF (1)	All (36)
		N	N	N	N	N	N	N
1	Functioning	0	0	0	1	0	0	1

**Source of irrigation:** The data regarding the source of irrigation in Kotageri-2 Micro watershed is presented in Table 20. The results that open well were major source of irrigation for 2.78 per cent of the households and bore well for 30.56 per cent of the households.

**Table 20. Source of irrigation in Kotageri-2 micro-watershed**

Sl. No.	Particulars	LL (3)		MF (15)		SF (10)		SMF (5)		MDF (2)		LF (1)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Bore Well	0	0	0	0	4	40	3	60	2	100	2	200	11	30.56
2	Canal	0	0	1	6.67	0	0	0	0	0	0	0	0	1	2.78
3	Open Well	0	0	0	0	0	0	1	20	0	0	0	0	1	2.78

**Depth of water (Avg. In meters):** The data regarding the depth of water in Kotageri-2 Micro watershed is presented in Table 21. The results revealed that, the depth of open well was 2.54 meter and depth of bore well was 14.01 meter.

**Table 21. Depth of water (Avg. In meters) in Kotageri-2 micro-watershed**

Sl. No.	Particulars	LL (3)	MF (15)	SF (10)	SMF (5)	MDF (2)	LF (1)	All (36)
		N	N	N	N	N	N	N
1	Bore Well	0	0	17.98	32	51.82	60.96	14.01
3	Open Well	0	0	0	18.29	0	0	2.54

**Irrigated Area (ha):** The data regarding the irrigated area (ha) in Kotageri-2 Micro watershed is presented in Table 22. The results indicate that, the availability of irrigation water was used for kharif crops was 9.83 ha.

**Table 22. Irrigated Area (ha) in Kotageri-2 micro-watershed**

Sl.No.	Particulars	LL (3)	MF (15)	SF (10)	SMF (5)	MDF (2)	LF (1)	All (36)
1	Kharif	0	0.4	1.33	1.62	0	6.48	9.83

**Table 23. Cropping pattern in Kotageri-2 micro-watershed**

Sl.No.	Particulars	LL (3)	MF (15)	SF (10)	SMF (5)	MDF (2)	LF (1)	All (36)
1	Kharif - Red gram	0	5.67	8.32	5.43	0.9	2.02	22.34
2	Kharif - Groundnut	0	0	0	0	0	2.83	2.83
3	Kharif - Paddy	0	0.2	0.4	1.62	0.4	0	2.63
4	Rabi - Groundnut	0	0	0.93	0	1.62	0	2.55
5	Rabi - Jowar	0	0.51	1.7	0	0	0	2.2
6	Kharif - Cotton	0	0	0	1.21	0	0	1.21
7	Kharif - Greengram	0	0.4	0	0	0	0	0.4

**Cropping pattern:** The data regarding the cropping pattern in Kotageri-2 Micro watershed is presented in Table 23. The results indicate that, farmers have grown Red gram (22.34 ha), Groundnut (5.38 ha), Paddy (2.63 ha), Jowar (2.2 ha) Cotton (1.21 ha) and Greengram (0.40 ha).

**Cropping intensity:** The data regarding the cropping intensity in Kotageri-2 Micro watershed is presented in Table 24. The results indicate that, the cropping intensity was 100.00 per cent.

**Table 24. Cropping intensity (%) in Kotageri-2 micro-watershed**

Sl. No.	Particulars	LL (3)	MF (15)	SF (10)	SMF (5)	MDF (2)	LF (1)	All (36)
1	Cropping Intensity	0	100	100	100	100	100	100

**Possession of bank account and savings:** The data regarding the possession of bank account and saving in Kotageri-2 micro-watershed is presented in Table 25. The results indicate that, 94.44 cent of the households' posses bank account and 41.67 per cent of them have savings.

**Table 25. Possession of Bank account and savings in Kotageri-2 micro-watershed**

Sl. No.	Particulars	LL (3)		MF (15)		SF (10)		SMF (5)		MDF (2)		LF (1)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Account	3	100	13	86.67	10	100	5	100	2	100	1	100	34	94.44
2	Savings	0	0	4	26.67	5	50	3	60	2	100	1	100	15	41.67

**Borrowing status:** The data regarding the borrowing status in Kotageri-2 micro-watershed is presented in Table 26. The results indicate that, 83.33 percent of the sample farmers have borrowed credit from different sources.

**Table 26. Borrowing status in Kotageri-2 micro-watershed**

Sl. No.	Particulars	LL (3)		MF (15)		SF (10)		SMF (5)		MDF (2)		LF (1)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Credit Aailed	3	100	14	93.33	7	70	3	60	2	100	1	100	30	83.33

**Source of credit:** The data regarding the source of credit availed by households in Kotageri-2 micro-watershed is presented in Table 27. The results show that, 22.22 per cent have borrowed loan from Cooperative bank, 100 per cent have borrowed loan from Grameena Bank, and 33.33 per cent have borrowed loan from money lender.

**Table 27. Source of credit borrowed by households in Kotageri-2 micro-watershed**

Sl. No.	Particulars	LL (0)		MF (2)		SF (3)		SMF (2)		MDF (2)		LF (0)		All (9)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Cooperative Bank	0	0	1	50	1	33.3	0	0	0	0	0	0	2	22.22
2	Grameena Bank	0	0	2	100	3	100	2	100	2	100	0	0	9	100
3	Money Lender	0	0	1	50	2	66.7	0	0	0	0	0	0	3	33.33

**Avg. Credit amount:** The data regarding the avg. Credit amount in Kotageri-2 micro-watershed is presented in Table 28. The results show that, farmers have borrowed Avg. Credit of Rs.196666.67 from different sources.

**Table 28. Avg. Credit amount in Kotageri-2 micro-watershed**

Sl. No.	Particulars	LL (0)	MF (2)	SF (3)	SMF (2)	MDF (2)	LF (0)	All (9)
		N	N	N	N	N	N	N
1	Average Credit	0	96500	379333	100000	49500	0	196667

**Purpose of credit borrowed (institutional Source):** The data regarding the purpose of credit borrowed - Institutional Credit in Kotageri-2 micro-watershed is presented in Table 29. The results indicate that, 92.86 per cent of the households have borrowed loan for agriculture and income generating activities (7.14 %).

**Table 29. Purpose of credit borrowed (institutional Source) by households in Kotageri-2 micro-watershed**

Sl. No	Particulars	LL (0)		MF (2)		SF (3)		SMF (2)		MDF (2)		LF (0)		All (9)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Agriculture production	0	0	3	100	5	100	3	100	1	100	1	100	13	92.9
4	Income generating activities	0	0	0	0	0	0	0	0	0	0	0	0	1	7.14

**Purpose of credit borrowed (Private Source):** The data regarding the purpose of credit borrowed – Private Source in Kotageri-2 micro-watershed is presented in Table 30. The results indicate that, 66.67 per cent of the households have borrowed loan for agriculture and 33.33 per cent of Social functions like marriage.

**Table 30. Purpose of credit borrowed (Private Source) by households in Kotageri-2 micro-watershed**

Sl. No.	Particulars	LL (0)		MF (1)		SF (2)		SMF (0)		MDF (0)		LF (0)		All (3)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Agriculture production	0	0	1	100	1	50	0	0	0	0	0	0	2	66.67
2	Social functions like marriage	0	0	0	0	1	50	0	0	0	0	0	0	1	33.33

**Repayment status of household (institutional Source):** The data regarding the repayment status of credit borrowed from institutional Source by households in Kotageri-2 micro watershed is presented in Table 31. The results indicate that, 7.14 per cent of the households have partially paid and 92.86 per cent have unpaid.

**Table 31. Repayment status of household (institutional Source) in Kotageri-2 micro-watershed**

Sl. No.	Particulars	LL (1)		MF (3)		SF (5)		SMF (3)		MDF (1)		LF (1)		All (14)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Partially paid	0	0	1	33.3	0	0	0	0	0	0	0	0	1	7.14
2	Un paid	1	100	2	66.7	5	100	3	100	1	100	1	100	13	92.86

**Repayment status of household (Private Source):** The data regarding the repayment status of credit borrowed from private sources by households in Kotageri-2 micro



watershed is presented in Table 32. The results indicate that, 33.33 per cent of the households have partially paid and 66.7 percent have unpaid.

**Table 32. Repayment status of household (Private Source) in Kotageri-2 micro-watershed**

Sl. No.	Particulars	LL (0)		MF (1)		SF (2)		SMF (0)		MDF (0)		LF (0)		All (3)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Partially paid	0	0	0	0	1	50	0	0	0	0	0	0	1	33.3
2	Un paid	0	0	1	100	1	50	0	0	0	0	0	0	2	66.7

**Opinion regarding institutional sources of credit:** The data regarding the opinion on institutional sources of credit in Kotageri-2 micro watershed is presented in Table 33. The results indicate that, 64.29 per cent of the households opined that credit helped to perform timely agricultural operations, 28.57 per cent Higher rate of interest and 7.14 per cent Loan amount was adequate to fulfil the requirement.

**Table 33. Opinion regarding institutional sources of credit in Kotageri-2 micro-watershed**

Sl. No.	Particulars	LL (1)		MF (3)		SF (5)		SMF (3)		MDF (1)		LF (1)		All (14)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Helped to perform timely agricultural operations	0	0	2	66.7	4	80	2	67	1	100	0	0	9	64.3
2	Higher rate of interest	1	100	0	0	1	20	1	33	0	0	1	100	4	28.6
3	Loan amount was adequate to fulfil the requirement	0	0	1	33.3	0	0	0	0	0	0	0	0	1	7.14

**Opinion regarding Non- institutional sources of credit:** The data regarding the opinion on non-institutional sources of credit in Kotageri-2 micro watershed is presented in Table 34. The results indicate that, 33 per cent of the households Loan amount was adequate to fulfil the requirement and 67 per cent of the households Higher rate of interest.

**Table 34. Opinion regarding Non- institutional sources of credit in Kotageri-2 micro-watershed**

Sl. No.	Particulars	LL (0)		MF (1)		SF (2)		SMF (0)		MDF (0)		LF (0)		All (3)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Loan amount was adequate to fulfil the requirement	0	0	0	0	1	50	0	0	0	0	0	0	1	33
2	Higher rate of interest	0	0	1	100	1	50	0	0	0	0	0	0	2	67

**Cost of Cultivation of Red gram:** The data regarding the cost of cultivation (Rs/ha) of Red gram in Kotageri-2 micro watershed is presented in Table 35.a. The results indicate that, the total cost of cultivation (Rs/ha) for Red gram was Rs. 45496.44. The gross income realized by the farmers was Rs. 56182.96. The net income from Red gram cultivation was Rs.10686.52, thus the benefit cost ratio was found to be 1:1.20.

**Table 35(a). Cost of Cultivation of Red gram in Kotageri-2 micro-watershed**

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
<b>I</b>	<b>Cost A1</b>				
1	Hired Human Labour	Man days	53.4	12118.8	26.64
2	Bullock	Pairs/day	3.66	2794.04	6.14
3	Tractor	Hours	3.06	2216.02	4.87
4	Machinery	Hours	0	0	0
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	8.64	809.93	1.78
6	Seed Inter Crop	Kgs.	0	0	0
7	FYM	Quintal	2.62	6539.09	14.37
8	Fertilizer + micronutrients	Quintal	4.4	3385.73	7.44
9	Pesticides (PPC)	Kgs/liters	2.32	1551.17	3.41
10	Irrigation	Number	4.26	0	0
11	Repairs		0	713.04	1.57
12	Msc. Charges (Marketing costs etc)		0	47.83	0.11
13	Depreciation charges		0	1218.65	2.68
14	Land revenue and Taxes		0	0.72	0
<b>II</b>	<b>Cost B1</b>				
16	Interest on working capital			1474.41	3.24
17	<b>Cost B1 = (Cost A1 + sum of 15 and 16)</b>			32869.42	72.25
<b>III</b>	<b>Cost B2</b>				
18	Rental Value of Land			295.83	0.65
19	<b>Cost B2 = (Cost B1 + Rental value)</b>			33165.26	72.9
<b>IV</b>	<b>Cost C1</b>				
20	Family Human Labour		33.87	8194.31	18.01
21	<b>Cost C1 = (Cost B2 + Family Labour)</b>			41359.57	90.91
<b>V</b>	<b>Cost C2</b>				
22	Risk Premium			0.83	0
23	<b>Cost C2 = (Cost C1 + Risk Premium)</b>			41360.4	90.91
<b>VI</b>	<b>Cost C3</b>				
24	Managerial Cost			4136.04	9.09
25	<b>Cost C3 = (Cost C2 + Managerial Cost)</b>			45496.44	100
<b>VII</b>	<b>Economics of the Crop</b>				
a.	Main Product	a) Main Product (q)		11.21	50523.51
		b) Main Crop Sales Price (Rs.)			4508.7
	By Product	e) Main Product (q)		2.76	5659.44
		f) Main Crop Sales Price (Rs.)			2052.17
b.	Gross Income (Rs.)			56182.96	
c.	Net Income (Rs.)			10686.52	
d.	Cost per Quintal (Rs./q.)			4060.08	
e.	Benefit Cost Ratio (BC Ratio)			1:1.2	

**Cost of Cultivation of Paddy:** The data regarding the cost of cultivation (Rs/ha) of Paddy in Kotageri-2 micro watershed is presented in Table 35.b. The results indicate that, the total cost of cultivation (Rs/ha) for Paddy was Rs. 93955.24. The gross income realized by the farmers was Rs. 101184.24. The net income from Paddy cultivation was Rs.7228.99, thus the benefit cost ratio was found to be 1:1.10.

**Table 35(b). Cost of Cultivation of Paddy in Kotageri-2 micro-watershed**

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
<b>I</b>	<b>Cost A1</b>				
1	Hired Human Labour	Man days	104.56	21711.3	23.11
2	Bullock	Pairs/day	2.68	2519.4	2.68
3	Tractor	Hours	3.71	2815.8	3
4	Machinery	Hours	0	0	0
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	59.9	20090.36	21.38
6	Seed Inter Crop	Kgs.	0	0	0
7	FYM	Quintal	2.47	6175	6.57
8	Fertilizer + micronutrients	Quintal	10.91	7739.33	8.24
9	Pesticides (PPC)	Kgs / liters	3.09	2099.5	2.23
10	Irrigation	Number	10.5	0	0
11	Repairs		0	1333.33	1.42
12	Msc. Charges (Marketing costs etc)		0	1016.67	1.08
13	Depreciation charges		0	522.44	0.56
14	Land revenue and Taxes		0	4.12	0
<b>II</b>	<b>Cost B1</b>				
16	Interest on working capital			4335.1	4.61
17	<b>Cost B1 = (Cost A1 + sum of 15 and 16)</b>			70362.36	74.89
<b>III</b>	<b>Cost B2</b>				
18	Rental Value of Land			333.33	0.35
19	<b>Cost B2 = (Cost B1 + Rental value)</b>			70695.69	75.24
<b>IV</b>	<b>Cost C1</b>				
20	Family Human Labour		60.1	14696.5	15.64
21	<b>Cost C1 = (Cost B2 + Family Labour)</b>			85392.19	90.89
<b>V</b>	<b>Cost C2</b>				
22	Risk Premium			21.67	0.02
23	<b>Cost C2 = (Cost C1 + Risk Premium)</b>			85413.86	90.91
<b>VI</b>	<b>Cost C3</b>				
24	Managerial Cost			8541.39	9.09
25	<b>Cost C3 = (Cost C2 + Managerial Cost)</b>			93955.24	100
<b>VII</b>	<b>Economics of the Crop</b>				
a.	Main Product	a) Main Product (q)		49.61	98384.9
		b) Main Crop Sales Price (Rs.)			1983.33
	By Product	e) Main Product (q)		3.29	2799.33
		f) Main Crop Sales Price (Rs.)			850
b.	Gross Income (Rs.)			101184.24	
c.	Net Income (Rs.)			7228.99	
d.	Cost per Quintal (Rs./q.)			1894.04	
e.	Benefit Cost Ratio (BC Ratio)			1:1.1	

**Cost of Cultivation of Green gram:** The data regarding the cost of cultivation (Rs/ha) of Green gram in Kotageri-2 micro watershed is presented in Table 35.c. The results indicate, the total cost of cultivation (Rs/ha) for Green gram was Rs.54349.84. The gross income realized by the farmers was Rs. 49400.00. The net income from Green gram cultivation was Rs. -4949.84, thus the benefit cost ratio was found to be 1:0.92.

**Table 35(c). Cost of Cultivation of Green gram in Kotageri-2 micro-watershed**

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
<b>I</b>	<b>Cost A1</b>				
1	Hired Human Labour	Man days	113.62	24082.5	44.31
2	Bullock	Pairs/day	4.94	3458	6.36
3	Tractor	Hours	0	0	0
4	Machinery	Hours	0	0	0
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	9.88	691.6	1.27
6	Seed Inter Crop	Kgs.	0	0	0
7	FYM	Quintal	0	0	0
8	Fertilizer + micronutrients	Quintal	4.94	3458	6.36
9	Pesticides (PPC)	Kgs / liters	2.47	1482	2.73
10	Irrigation	Number	0	0	0
11	Repairs		0	0	0
12	Msc. Charges (Marketing costs etc)		0	0	0
13	Depreciation charges		0	0.05	0
14	Land revenue and Taxes		0	0	0
<b>II</b>	<b>Cost B1</b>				
16	Interest on working capital			675.79	1.24
17	<b>Cost B1 = (Cost A1 + sum of 15 and 16)</b>			33847.94	62.28
<b>III</b>	<b>Cost B2</b>				
18	Rental Value of Land			0	0
19	<b>Cost B2 = (Cost B1 + Rental value)</b>			33847.94	62.28
<b>IV</b>	<b>Cost C1</b>				
20	Family Human Labour		51.87	15561	28.63
21	<b>Cost C1 = (Cost B2 + Family Labour)</b>			49408.94	90.91
<b>V</b>	<b>Cost C2</b>				
22	Risk Premium			0	0
23	<b>Cost C2 = (Cost C1 + Risk Premium)</b>			49408.94	90.91
<b>VI</b>	<b>Cost C3</b>				
24	Managerial Cost			4940.89	9.09
25	<b>Cost C3 = (Cost C2 + Managerial Cost)</b>			54349.84	100
<b>VII</b>	<b>Economics of the Crop</b>				
a.	Main Product	a) Main Product (q)		9.88	49400
		b) Main Crop Sales Price (Rs.)			5000
b.	Gross Income (Rs.)				49400
c.	Net Income (Rs.)				-4949.84
d.	Cost per Quintal (Rs./q.)				5501
e.	Benefit Cost Ratio (BC Ratio)				1:0.92

**Cost of Cultivation of Groundnut:** The data regarding the cost of cultivation (Rs/ha) of Groundnut in Kotageri-2 micro watershed is presented in Table 35.d. The results indicate that, the total cost of cultivation (Rs/ha) for Groundnut was Rs. 45538.76. The gross income realized by the farmers was Rs.54829.84. The net income from Groundnut cultivation was Rs. 9291.08, thus the benefit cost ratio was found to be 1:1.20.

**Table 35(d). Cost of Cultivation of Groundnut in Kotageri-2 micro-watershed**

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
<b>I</b>	<b>Cost A1</b>				
1	Hired Human Labour	Man days	16.82	3079.99	6.76
2	Bullock	Pairs/day	3.14	3002.47	6.59
3	Tractor	Hours	4.27	3419.48	7.51
4	Machinery	Hours	0	0	0
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	120.57	12301.37	27.01
6	Seed Inter Crop	Kgs.	0	0	0
7	FYM	Quintal	1.08	2696.51	5.92
8	Fertilizer + micronutrients	Quintal	3.25	2583.39	5.67
9	Pesticides (PPC)	Kgs / liters	2.01	1053.49	2.31
10	Irrigation	Number	4.55	0	0
11	Repairs		0	2000	4.39
12	Msc. Charges (Marketing costs etc)		0	0	0
13	Depreciation charges		0	2161.99	4.75
14	Land revenue and Taxes		0	0	0
<b>II</b>	<b>Cost B1</b>				
16	Interest on working capital			2236.17	4.91
17	<b>Cost B1 = (Cost A1 + sum of 15 and 16)</b>			34534.86	75.84
<b>III</b>	<b>Cost B2</b>				
18	Rental Value of Land			316.67	0.7
19	<b>Cost B2 = (Cost B1 + Rental value)</b>			34851.53	76.53
<b>IV</b>	<b>Cost C1</b>				
20	Family Human Labour		30.36	6547.35	14.38
21	<b>Cost C1 = (Cost B2 + Family Labour)</b>			41398.87	90.91
<b>V</b>	<b>Cost C2</b>				
22	Risk Premium			0	0
23	<b>Cost C2 = (Cost C1 + Risk Premium)</b>			41398.87	90.91
<b>VI</b>	<b>Cost C3</b>				
24	Managerial Cost			4139.89	9.09
25	<b>Cost C3 = (Cost C2 + Managerial Cost)</b>			45538.76	100
<b>VII</b>	<b>Economics of the Crop</b>				
a.	Main Product	a) Main Product (q)		12.11	52058.69
		b) Main Crop Sales Price (Rs.)			4300
	By Product	e) Main Product (q)		2.08	2771.15
		f) Main Crop Sales Price (Rs.)			1333.33
b.	Gross Income (Rs.)			54829.84	
c.	Net Income (Rs.)			9291.08	
d.	Cost per Quintal (Rs./q.)			3761.46	
e.	Benefit Cost Ratio (BC Ratio)			1:1.2	

**Cost of Cultivation of Cotton:** The data regarding the cost of cultivation (Rs/ha) of Cotton in Kotageri-2 micro watershed is presented in Table 35.e. The results indicate that, the total cost of cultivation (Rs/ha) for Cotton was Rs.45070.85. The gross income realized by the farmers was Rs. 79698.67. The net income from Cotton cultivation was Rs. 34627.82, thus the benefit cost ratio was found to be 1:1.80.

**Table 35(e). Cost of Cultivation of Cotton in Kotageri-2 micro-watershed**

Sl. No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
<b>I</b>	<b>Cost A1</b>				
1	Hired Human Labour	Man days	58.46	9756.5	21.65
2	Bullock	Pairs/day	0	0	0
3	Tractor	Hours	2.47	1976	4.38
4	Machinery	Hours	0	0	0
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	2.47	3952	8.77
6	Seed Inter Crop	Kgs.	0	0	0
7	FYM	Quintal	0	0	0
8	Fertilizer + micronutrients	Quintal	2.47	1482	3.29
9	Pesticides (PPC)	Kgs / liters	1.65	823.33	1.83
10	Irrigation	Number	2.47	0	0
11	Repairs		0	3000	6.66
12	Msc. Charges (Marketing costs etc)		0	0	0
13	Depreciation charges		0	13218.45	29.33
14	Land revenue and Taxes		0	0	0
<b>II</b>	<b>Cost B1</b>				
16	Interest on working capital			750.88	1.67
17	<b>Cost B1 = (Cost A1 + sum of 15 and 16)</b>			34959.17	77.56
<b>III</b>	<b>Cost B2</b>				
18	Rental Value of Land			333.33	0.74
19	<b>Cost B2 = (Cost B1 + Rental value)</b>			35292.5	78.3
<b>IV</b>	<b>Cost C1</b>				
20	Family Human Labour		23.88	5681	12.6
21	<b>Cost C1 = (Cost B2 + Family Labour)</b>			40973.5	90.91
<b>V</b>	<b>Cost C2</b>				
22	Risk Premium			0	0
23	<b>Cost C2 = (Cost C1 + Risk Premium)</b>			40973.5	90.91
<b>VI</b>	<b>Cost C3</b>				
24	Managerial Cost			4097.35	9.09
25	<b>Cost C3 = (Cost C2 + Managerial Cost)</b>			45070.85	100
<b>VII</b>	<b>Economics of the Crop</b>				
a.	Main Product	a) Main Product (q)		18.11	79698.67
		b) Main Crop Sales Price (Rs.)			4400
b.	Gross Income (Rs.)				79698.67
c.	Net Income (Rs.)				34627.82
d.	Cost per Quintal (Rs./q.)				2488.27
e.	Benefit Cost Ratio (BC Ratio)				1:1.8

**Adequacy of fodder:** The data regarding the adequacy of fodder in Kotageri-2 Micro watershed is presented in Table 44. The results indicate that, 41.67 per cent of the households opined that dry fodder was adequate and 5.56 per cent of them opined dry fodder was inadequate.

**Table 36. Adequacy of fodder in Kotageri-2 micro-watershed**

Sl. No.	Particulars	LL (3)		MF (15)		SF (10)		SMF (5)		MDF (2)		LF (1)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Adequate-Dry Fodder	0	0	5	33.33	6	60	2	40	1	50	1	100	15	41.67
2	Inadequate-Dry Fodder	0	0	0	0	1	10	0	0	1	50	0	0	2	5.56

**Average annual gross income:** The data regarding the annual gross income in Kotageri-2 Micro watershed is presented in Table 37. The results indicate that, the farmers have annual gross income of Rs. 137441.67 in micro-watershed, of which Rs. 60461.11 is from agriculture itself.

**Table 37. Average annual gross income in Kotageri-2 micro-watershed**

Sl.No.	Particulars	LL (3)	MF (15)	SF (10)	SMF (5)	MDF (2)	LF (1)	All (36)
		Rs.	Rs.	Rs.	Rs.	Rs.	Rs.	Rs.
1	Business	0	8000	0	0	40000	0	5555.56
2	Wage	86666.7	50166.7	85500	84000	20000	75000	66736.1
3	Agriculture	0	26573.3	60920	108760	104500	416000	60461.1
4	Dairy Farm	6666.67	0	8496	864	6000	47520	4688.89
	Income(Rs.)	93333.3	84740	154916	193624	170500	538520	137442

**Average annual Expenditure:** The data regarding the average annual expenditure in Kotageri-2 Micro watershed is presented in Table 38. The results indicate that, the farmers have annual gross expenditure of Rs. 647238.10 in micro-watershed, of which Rs. 30272.22 is from agriculture itself.

**Table 38. Average annual Expenditure in Kotageri-2 micro-watershed**

Sl. No.	Particulars	LL (3)	MF (15)	SF (10)	SMF (5)	MDF (2)	LF (1)	All (36)
		Rs.	Rs.	Rs.	Rs.	Rs.	Rs.	Rs.
1	Business	0	25000	0	0	35000	0	2361.11
2	Wage	41666.7	20357.1	43000	38800	15000	52000	30583.3
3	Agriculture	0	14064.3	28950	44566.7	32000	240000	30272.2
4	Dairy Farm	1000	0	4833.33	2500	6000	2500	1138.89
	Total	42666.7	59421.4	76783.3	85866.7	88000	294500	647238

**Table 39. Horticulture species grown in Kotageri-2 micro-watershed**

Sl. No.	Particulars	LL (3)		MF (15)		SF (10)		SMF (5)		MDF (2)		LF (1)		All (36)	
		F	B	F	B	F	B	F	B	F	B	F	B	F	B
9	Custard apple	0	0	0	0	6	0	0	0	2	2	0	0	8	2

\*F= Field B=Back Yard

**Horticulture species grown:** The data regarding horticulture species grown in Kotageri-2 Micro watershed is presented in Table 39. The results indicate that, the total number of



horticultural trees grown (both field and backyard) by the sampled households were clustered apple (10).

**Interest towards cultivation of horticulture crops:** The data regarding Table (40) indicates that, 5.56 percent of the households shown interest to cultivate horticultural crops.

**Table 40. Interest towards cultivation of horticulture crops in Kotageri-2 micro-watershed**

Sl. No.	Particulars	LL (3)		MF (15)		SF (10)		SMF (5)		MDF (2)		LF (1)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Interested towards cultivation of horticulture crops	0	0	1	6.7	1	10	0	0	0	0	0	0	2	5.56

**Forest species grown:** The data regarding forest species grown in Kotageri-2 Micro watershed is presented in Table 41. The results indicate that, households have planted 33 neem trees, 2 tamarind trees and 2 acacia trees, together in both field and backyard.

**Table 41. Forest species grown in Kotageri-2 micro-watershed**

Sl.No.	Particulars	LL (3)		MF (15)		SF (10)		SMF (5)		MDF (2)		LF (1)		All (36)	
		F	B	F	B	F	B	F	B	F	B	F	B	F	B
1	Neem	0	0	17	3	5	4	2	0	1	1	0	0	25	8
2	Tamarind	0	0	0	0	0	2	0	0	0	0	0	0	0	2
3	Acacia	0	0	0	2	0	0	0	0	0	0	0	0	0	2

\*F= Field B=Back Yard

**Average additional investment capacity:** The data regarding average additional investment capacity in Kotageri-2 Micro watershed is presented in Table 42. The results indicate that, households have an average investment capacity of Rs. 2777.78 for land development, Rs. 4166.67 for creation of irrigation facility, Rs.3333.33 for adoption of improved livestock breeds, Rs.2222.22 for adoption of improved crop production activities.

**Table 42. Average additional investment capacity of households in Kotageri-2 micro-watershed**

Sl. No.	Particulars	LL (3)	MF (15)	SF (10)	SMF (5)	MDF (2)	LF (1)	All (36)
		Rs.	Rs.	Rs.	Rs.	Rs.	Rs.	Rs.
1	Land development	0	1333.33	5000	6000	0	0	2777.78
2	Irrigation facility	0	10000	0	0	0	0	4166.67
3	Improved crop production	0	1333.33	10000	0	0	0	3333.33
4	Improved livestock management	0	0	5000	0	15000	0	2222.22

**Source of funds for additional investment:** The data regarding source of funds for additional investment in Kotageri-2 Micro watershed is presented in Table 43. The results

indicate that, the sources of finance raised from bank as a Government subsidy for land development was 2.78 and 5.56 per cent, for irrigation facility.

**Table 43. Source of funds for additional investment in Kotageri-2 micro-watershed**

Sl. No	Item	Land development		Irrigation facility		Improved crop production	
		N	%	N	%	N	%
1	Government subsidy	1	2.78	2	5.56	1	2.78

**Marketing of agricultural produce:** The data regarding marketing of the agricultural produce in Kotageri-2 Micro watershed is presented in Table 44. The results indicated that, 100.00 percent of output of Cotton was sold in the market with average price of Rs. 4400.00; 50.00 percent of output of Greengram was sold in the market with average price of Rs. 5000.00; 82.09 percent of output of Groundnut was sold in the market with average price of Rs. 4300.00 and 82.61 percent of output of Jowar was sold in the market with average price of Rs. 2650.00; 64 percent of output of Paddy was sold in the market with average price of Rs. 1983; and 75 percent of output of Redgram was sold in the market with average price of Rs. 4509.

**Table 44. Marketing of agricultural produce in Kotageri-2 micro-watershed**

Sl. No	Crops	Output obtained (q)	Output retained (q)	Output sold (q)	Output sold (%)	Avg. Price obtained (Rs/q)
1	Cotton	22	0	22	100	4400
2	Greengram	4	2	2	50	5000
3	Groundnut	67	12	55	82	4300
4	Jowar	23	4	19	83	2650
5	Paddy	138	50	88	64	1983
6	Redgram	239	60	179	75	4509

**Marketing channels used for sale of agricultural produce:** The data regarding marketing channels used for sale of agricultural produce in Kotageri-2 Micro watershed is presented in Table 45. The results indicated that, 44.44 cent of the households have sold agricultural produce to the local/village merchants and 55.56 per cent of regulated market.

**Table 45. Marketing channels used for sale of agricultural produce in Kotageri-2 micro-watershed**

Sl. No.	Particulars	LL (3)		MF (15)		SF (10)		MF (5)		IDF (2)		LF (1)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Local/village Merchant	0	0	7	47	4	40	4	80	1	50	0	0	16	44.44
2	Regulated Market	0	0	7	47	6	60	3	60	2	100	2	200	20	55.56

**Mode of transport of agricultural produce:** The data regarding mode of transport of agricultural produce in Kotageri-2 Micro watershed is presented in Table 46. The results indicated that, 77.78 cent of the households have used tractor and 16.67 per cent have used Cart and 5.56 per cent have used Truck.

**Table 46. Mode of transport of agricultural produce in Kotageri-2 micro-watershed**

Sl. No.	Particulars	LL (3)		MF (15)		SF (10)		SMF (5)		MDF (2)		LF (1)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Cart	0	0	5	33	1	10	0	0	0	0	0	0	6	16.67
2	Tractor	0	0	8	53	8	80	7	140	3	150	2	200	28	77.78
3	Truck	0	0	1	6.7	1	10	0	0	0	0	0	0	2	5.56

**Incidence of soil and water erosion problems:** The data regarding incidence of incidence of soil and water erosion problems in Kotageri-2 Micro watershed is presented in Table 47. The results indicate that, 44.44 per cent of the households have experienced soil and water erosion problems.

**Table 47. Incidence of soil and water erosion problems in Kotageri-2 micro-watershed**

Sl. No.	Particulars	LL (3)		MF (15)		SF (10)		SMF (5)		MDF (2)		LF (1)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Soil and water erosion problems in the farm	0	0	8	53	7	70	1	20	0	0	0	0	16	44.44

**Interest towards soil testing:** The data regarding Interest shown towards soil testing in Kotageri-2 Micro watershed is presented in Table 48. The results indicated that, 83.33 per cent of the households were interested towards soil testing.

**Table 48. Interest regarding soil testing in Kotageri-2 micro-watershed**

Sl.No.	Particulars	LL (3)		MF (15)		SF (10)		SMF (5)		MDF (2)		LF (1)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Interest in soil test	0	0	13	87	9	90	5	100	2	100	1	100	30	83.33

**Soil and water conservation practices and structures adopted:** The data regarding soil and water conservation practices and structures adopted in Kotageri-2 Micro watershed is presented in Table 49. The results indicated that 92.89 per cent of farmers practicing summer ploughing as soil and water conservation practice.

**Table 49. Soil and water conservation practices and structures adopted in Kotageri-2 micro-watershed**

Sl No.	Particulars	LL (3)		MF (15)		SF (10)		SMF (5)		MDF (2)		LF (1)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Field Bunding	0	0	1	6.7	2	20	1	20	0	0	0	0	4	11.11
2	Farm Pond	0	0	2	13	0	0	0	0	0	0	0	0	2	5.56

**Table 50. Status of soil and water conservation structures in Kotageri-2 micro-watershed**

Sl. No	Item	Good		Slightly Damaged		Severely Damaged		Full Replacement Required	
		N	%	N	%	N	%	N	%
1	Farm Pond	2	100	0	0	0	0	0	0

**Status of soil and water conservation structures:** The data regarding status soil and water conservation structures adopted in Kotageri-2 Micro watershed is presented in

Table 50. The results indicated that, the households have adopted field bunding as a soil and water conservation structures out of which 100.00 per cent was in good condition.

**Agencies involved in the soil and water conservation structures:** The data regarding Agencies involved in the soil and water conservation structures adopted in Kotageri-2 Micro watershed is presented in Table 51. The results indicated that, 5.56 per cent of the households have adopted by their own 11.11 per cent were done by Government

**Table 51. Agencies involved in the soil and water conservation structures in Kotageri-2 micro-watershed**

Sl. No.	Particulars	LL (3)		MF (15)		SF (10)		SMF (5)		MDF (2)		LF (1)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Own	0	0	1	6.7	0	0	1	20	0	0	0	0	2	5.56
2	Govt.	0	0	2	13	2	20	0	0	0	0	0	0	4	11.11

**Usage pattern of fuel for domestic use:** The data on usage pattern of fuel for domestic use in Kotageri-2 Micro watershed is presented in Table 52. The results indicated that, firewood was the major source of fuel for domestic use for 47.22 per cent of the households followed by LPG (50.00%).

**Table 52. Usage pattern of fuel for domestic use in Kotageri-2 micro-watershed**

Sl. No.	Particulars	LL (3)		MF (15)		SF (10)		SMF (5)		MDF (2)		LF (1)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Fire Wood	1	33.3	10	66.7	4	40	2	40	0	0	0	0	17	47.22
2	LPG	2	66.7	5	33.3	6	60	3	60	1	50	1	100	18	50

**Source of drinking water:** The data on source of drinking water in Kotageri-2 Micro watershed is presented in Table 53. The results indicated that, Piped supply of water was the major source for drinking water for 86.11 per cent of the households followed by bore well water (13.89%).

**Table 53. Source of drinking water in Kotageri-2 micro-watershed**

Sl. No.	Particulars	LL (3)		MF (15)		SF (10)		SMF (5)		MDF (2)		LF (1)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Piped supply	3	100	14	93.3	8	80	4	80	2	100	0	0	31	86.11
2	Bore Well	0	0	1	6.67	2	20	1	20	0	0	1	100	5	13.89

**Source of light:** The data on source of light in Kotageri-2 Micro watershed is presented in Table 54. The results indicated that, electricity was the major source of light for 100.00 per cent of the households.

**Table 54. Source of light in Kotageri-2 micro-watershed**

Sl. No.	Particulars	LL (3)		MF (15)		SF (10)		SMF (5)		MDF (2)		LF (1)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Electricity	3	100	15	100	10	100	5	100	2	100	1	100	36	100

**Existence of sanitary toilet facility:** The data on availability of toilet facility in Kotageri-2 Micro watershed is presented in Table 55. The results indicated that, 41.67 per cent of the households possess toilets.

**Table 55. Existence of sanitary toilet facility in Kotageri-2 micro-watershed**

Sl. No.	Particulars	LL (3)		MF (15)		SF (10)		SMF (5)		MDF (2)		LF (1)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Sanitary toilet facility	2	66.7	6	40	3	30	1	20	2	100	1	100	15	41.7

**Possession of PDS card:** The data regarding possession of PDS card in Kotageri-2 Micro watershed is presented in Table 56. The results indicated that, 91.67 per cent of the households possessed BPL card, 2.78 per cent possessed APL card and 5.56 per cent do not possess PDS card.

**Table 56. Possession of PDS card in Kotageri-2 micro-watershed**

Sl. No.	Particulars	LL (3)		MF (15)		SF (10)		SMF (5)		MDF (2)		LF (1)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	APL	0	0	1	6.67	0	0	0	0	0	0	0	0	1	2.78
2	BPL	3	100	13	86.7	10	100	4	80	2	100	1	100	33	91.67
3	Not Possessed	0	0	1	6.67	0	0	1	20	0	0	0	0	2	5.56

**Participation in NREGA programme:** The data regarding Participation in NREGA programme in Kotageri-2 Micro watershed is presented in Table 57. The results indicated that, only 13.89 per cent of the households have participated in NREGA programme.

**Table 57. Participation in NREGA programme in Kotageri-2 micro-watershed**

Sl. No.	Particulars	LL (3)		MF (15)		SF (10)		SMF (5)		MDF (2)		LF (1)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Participation in NREGA programme	0	0	3	20	2	20	0	0	0	0	0	0	5	13.9

**Adequacy of food items:** The data regarding adequacy of food items in Kotageri-2 Micro watershed is presented in Table 58. The results indicated that, the extent of adequacy of food items for cereals, pulses, Oilseeds and vegetables were 94.44, 83.33, 63.89, 58.33 per cent respectively, similarly for Fruits (25.00%), milk (25.00%), Egg (25.00%), and Meat (16.67%).

**Table 58. Adequacy of food items in Kotageri-2 micro-watershed**

Sl.No.	Particulars	LL (3)		MF (15)		SF (10)		SMF (5)		MDF (2)		LF (1)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Cereals	3	100	14	93.3	10	100	4	80	2	100	1	100	34	94.44
2	Pulses	2	66.7	11	73.3	9	90	5	100	2	100	1	100	30	83.33
3	Oilseed	1	33.3	10	66.7	5	50	4	80	2	100	1	100	23	63.89
4	Vegetables	3	100	9	60	6	60	1	20	1	50	1	100	21	58.33
5	Fruits	0	0	7	46.7	0	0	0	0	1	50	1	100	9	25
6	Milk	0	0	4	26.7	3	30	1	20	1	50	0	0	9	25
7	Egg	1	33.3	4	26.7	3	30	0	0	1	50	0	0	9	25
8	Meat	0	0	4	26.7	2	20	0	0	0	0	0	0	6	16.67

**Inadequacy of food items:** The data regarding in adequacy of food items in Kotageri-2 Micro watershed is presented in Table 59. The results indicated that, the extent of in adequacy of food items for cereals, pulses, Oilseeds and vegetables were 5.56, 16.67, 36.11, 41.67 and 80.56 per cent respectively, similarly for fruits (75.00%), milk (58.33%), egg (72.22%) and meat (80.56%).

**Table 59. Inadequacy of food items in Kotageri-2 micro-watershed**

Sl. No.	Particulars	LL (3)		MF (15)		SF (10)		SMF (5)		MDF (2)		LF (1)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Cereals	0	0	1	6.67	0	0	1	20	0	0	0	0	2	5.56
2	Pulses	1	33.3	4	26.7	1	10	0	0	0	0	0	0	6	16.67
3	Oilseed	2	66.7	5	33.3	5	50	1	20	0	0	0	0	13	36.11
4	Vegetables	0	0	6	40	4	40	4	80	1	50	0	0	15	41.67
5	Fruits	3	100	8	53.3	10	100	5	100	1	50	0	0	27	75
6	Milk	2	66.7	11	73.3	3	30	4	80	1	50	0	0	21	58.33
7	Egg	2	66.7	11	73.3	6	60	5	100	1	50	1	100	26	72.22
8	Meat	3	100	11	73.3	7	70	5	100	2	100	1	100	29	80.56

**Response on market surplus of food items:** The data regarding adequacy of food items in Kotageri-2 Micro watershed is presented in Table 60. The results indicated that, the extent of market surplus adequacy of food items for milk (16.67%), egg (2.78%) and meat (2.78%).

**Table 60. Response on market surplus of food items in Kotageri-2 micro-watershed**

Sl. No.	Particulars	LL (3)		MF (15)		SF (10)		SMF (5)		MDF (2)		LF (1)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Milk	1	33.3	0	0	4	40	0	0	0	0	1	100	6	16.67
2	Egg	0	0	0	0	1	10	0	0	0	0	0	0	1	2.78
3	Meat	0	0	0	0	1	10	0	0	0	0	0	0	1	2.78

**Table 61. Farming constraints experienced in Kotageri-2 micro-watershed**

Sl. No	Particulars	MF (15)		SF (10)		SMF (5)		MDF (2)		LF (1)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Lower fertility status of the soil	14	93.33	8	80	3	60	2	100	0	0	27	75
2	Wild animal menace on farm field	8	53.33	8	80	5	100	2	100	1	100	24	66.67
3	Frequent incidence of pest and diseases	13	86.67	10	100	4	80	2	100	1	100	30	83.33
4	Inadequacy of irrigation water	10	66.67	7	70	3	60	1	50	1	100	22	61.11
5	High cost of Fertilizers and plant protection chemicals	14	93.33	10	100	5	100	2	100	1	100	32	88.89
6	High rate of interest on credit	13	86.67	9	90	5	100	2	100	1	100	30	83.33
7	Low price for the agricultural commodities	13	86.67	9	90	5	100	2	100	1	100	30	83.33
8	Lack of marketing facilities in the area	12	80	7	70	2	40	1	50	1	100	23	63.89
9	Inadequate extension services	3	20	4	40	2	40	2	100	0	0	11	30.56
10	Lack of transport for safe transport of the Agril produce to the market.	13	86.67	5	50	3	60	1	50	1	100	23	63.89

**Farming constraints:** The data regarding farming constraints experienced by households in Kotageri-2 Micro watershed is presented in Table 61. The results indicated that, lower fertility status of the soil was the constraint experienced by (75.00 %) per cent of the households, wild animal menace on farm field (66.67%), frequent incidence of pest and diseases (83.33%), inadequacy of irrigation water (61.11%), high cost of fertilizers and plant protection chemicals (88.89%), high rate of interest on credit (83.33%), low price for the agricultural commodities (83.33 %), lack of marketing facilities in the area (63.89%), inadequate extension services (30.56 %), lack of transport for safe transport of the agricultural produce to the market (63.89%).





**SUMMARY AND IMPLICATIONS**

In order to assess the socio-economic condition of the farmers in the watershed 36 households located in the micro watershed were interviewed for the survey. The study was conducted in Kotageri-2 micro-watershed (Shivapur sub-watershed, Yadgiri taluk & District) is located at North latitude 16<sup>0</sup> 54' 2.29" and 16<sup>0</sup> 52' 44.243" and East longitude 77<sup>0</sup> 15' 21.73" and 77<sup>0</sup> 13' 47.096" covering an area of about 301.61 ha bounded by unde Kootagera, Gajarakota and Fatthepura Villages.

Socio-economic analysis indicated that, out of the total sample of 36 total respondents, 15 (41.67 %) were marginal, 10 (27.78%) were small, 5 (13.89 %) were Semi medium and 2 (5.56 %) were medium and 1 (2.78 %) were large farmers. The population characteristics of households indicated that, there were 108 (54.82 %) men and 89 (45.18 %) were women. The average population of landless was 4.7, marginal farmers were 4.7, small farmers were 6.3, semi medium farmers were 4.5 and medium farmers were 10. Majority of the respondents (47.21 %) were in the age group of 16-35 years. Education level of the sample households indicated that, there were 37.06 per cent illiterates, 67.02 per cent pre university education and 5.08 per cent attained graduation. About, 69.44 per cent of household heads practicing agriculture and 16.67 per cent of the household heads were engaged as agricultural labourers. Agriculture was the major occupation for 46.19 per cent of the household members.

In the study area, 55.56 per cent of the households possess Katcha house and 38.89 per cent possess Pucca house. The durable assets owned by the households showed that, 58.33 per cent possess TV, 2.78 per cent possess mixer grinder, 91.67 per cent possess mobile phones and 22.22 per cent possess motor cycles. Farm implements owned by the households indicated that, 44.44 per cent of the households possess plough, 8.33 per cent possess tractor, 19.44 per cent possess bullock cart and 38.89 per cent possess sprayer. Regarding livestock possession by the households, 25.00 per cent possess local cow and 19.44 per cent possess buffalo.

The average labour availability in the study area showed that, own men and women labour availability in the micro watershed was 1.56 and 2.06 each, while the hired labour (women) availability was 12.94 and the hired labour (men) availability was 8.57. Further, 11.11 per cent of the households opined that hired labour was inadequate during the agricultural season.

Out of the total land holding of the sample respondents 42.05 per cent (38.73 ha) of the area is under dry condition and the remaining 56.91 per cent area is irrigated land. There were 11.00 live bore wells and 12.00 dry bore wells among the sampled households. Bore well was the major source of irrigation for 30.56 per cent of the households. The major crops grown by sample farmers are Red gram, Paddy, Green gram, Groundnut and Cotton and cropping intensity was recorded as 100.00 per cent.

Out of the sample households 94.44 percent possessed bank account and 41.67 per cent of them have savings in the account. About 83.33 per cent of the respondents borrowed credit from various sources. Among the credit borrowed by households, 22.22 per cent have borrowed loan from Cooperative bank, 100 per cent have borrowed loan from Grameena Bank, and 33.33 per cent have borrowed loan from money lender. Majority of the respondents (92.86 %) have borrowed loan for agriculture purpose. Regarding the opinion on institutional sources of credit, 64.29 per cent of the households opined that credit helped to perform timely agricultural operations, while, only 7.14 per cent respondents opined that loan amount was adequate to fulfil their requirement.

The per hectare cost of cultivation for Red gram, Paddy, Green gram, Groundnut and Cotton was Rs.45496.44, 93955.24, 54349.84, 45538.76 and 45070.85 with benefit cost ratio of 1:1.20, 1: 1.10, 1: 0.92, 1: 1.20 and 1:1.80 respectively.

Further, 41.67 per cent of the households opined that dry fodder was adequate

The average annual gross income of the farmers was Rs. 137441.67 in micro-watershed, of which Rs. 60461.11 comes from agriculture.

Sampled households have grown 10 horticulture trees and 12 forestry trees together in the fields and back yards.

Households have an average investment capacity of Rs. 2777.78 for land development and Rs. 4166.67 for irrigation facility. Source of funds for additional investment is concerned, 2.78 per cent a Government subsidy for land development activities.

Regarding marketing channels, 44.44 per cent of the households have sold agricultural produce to the local/village merchants, while, 55.56 per cent have sold in regulated markets. Further, 77.78 per cent of the households have used tractor for the transport of agriculture commodity.

Majority of the farmers (44.44 %) have experienced soil and water erosion problems in the watershed and 83.33 per cent of the households were interested towards soil testing. About, 92.89 per cent of farmers practicing summer ploughing as soil and water conservation practice.

Fire was the major source of fuel for domestic use for 47.22 per cent of the households and 50.00 per cent households has LPG connection. Piped supply was the major source for drinking water for 86.11 per cent of the households. Electricity was the major source of light for 100.00 per cent of the households. In the study area, 41.67 per cent of the households possess toilet facility. Regarding possession of PDS card, 91.67 per cent of the households possessed BPL card, 2.78 per cent of the household's possessed APL card and 5.56 per cent of the household's were not having ration

cards. Households opined that, the requirement of cereals (94.44%), pulses (83.33%) and oilseeds (63.89%) are adequate for consumption.

Farming constraints experienced by households in the micro watersheds were lower fertility status of the soil (75.00%) wild animal menace on farm field (66.67%), frequent incidence of pest and diseases (83.33%), inadequacy of irrigation water (61.11%), high cost of fertilizers and plant protection chemicals (88.89%), high rate of interest on credit (83.33%), low price for the agricultural commodities (83.33%), lack of marketing facilities in the area (63.89%), inadequate extension services (30.56%), lack of transport for safe transport of the agricultural produce to the market (63.89%).

### **Implications of the survey**

- ✓ Result indicated that, there were 37.06 per cent were illiterate hence, extension methodologies such as demonstration, street play, drama, video shows will be effective in dissemination of the technologies in the micro watershed.
- ✓ The data indicate that, 55.56 per cent of the households possess katcha house. Hence, the development department while implementing the watershed plan should focus on agriculture to enhance the productivity of major crops in the area to increase the income of the farmers.
- ✓ Results indicated that the local institutional participation of the household members in the micro watershed is minimal hence, activities like membership campaign, awareness creation about the benefits of membership in local institutions and strengths of organized groups must be conveyed.
- ✓ Majority of the households in the watershed have experience in use of mobile phones, and television hence, these mass media can be effectively utilized for transfer of technology as well as for information dissemination.
- ✓ The farm machinery/implement possession in the micro watershed was found to be minimum the reasons may lack of knowledge or lack of financial ability which can be addressed through training on use of different farm implements, providing information on different sources of finance for purchase of farm implements.
- ✓ The possession of livestock such as crossbred cow found is less hence, farmers must be made aware of the benefits of crossbred cow in increased milk production.
- ✓ The possession of livestock such as sheep, goat and poultry was found to be low hence, farmers may be informed the role of subsidiary enterprises in enhancing the income and information on financial support for subsidiary activities.
- ✓ The data indicate that, job/work was the reason for all the migrants hence, farmers may be trained on profitable agriculture or self employment such as animal husbandry, plate making, sheep rearing, goat rearing, rabbit rearing with suitable information on sources of financial support.

- ✓ The results indicate that there was a change in quality of life due to migration hence, the developmental departments should take actions to arrest migration and to improve the quality of the life in rural areas.
- ✓ Households possess 16.29ha (42.05 %) of dry land and 22.04ha (56.91 %) of irrigated land hence, the availability of the dry land agricultural technologies such as short duration crops, high yielding drought resistance crop varieties, drip irrigation technology and subsidy information will be helpful for the farmers to enhance the productivity of land and as well as farmers income.
- ✓ Few of the bore well in micro watershed found non functional hence, farmers may be trained on possibility of bore well rejuvenation.
- ✓ Bore well was major source of irrigation for 30.56 per cent of the households. hence, in order to increase the area under irrigation as well as to increase the water use efficiency farmers may trained on drip irrigation and provide the information on subsidy for drip irrigation equipment's along with the information on different agencies which provides the financial assistance for drip irrigation.
- ✓ The cropping intensity in the micro watershed was found to be (100.00 %) hence, care must be taken by the implementing agency to bring uncultivated land into cultivation through suitable measures.
- ✓ Many of the household members have borrowed loan from cooperative banks which has higher rate of interest hence, farmers may be sensitized on the different sources of credit with lesser interest rate such SHGs etc.
- ✓ The results indicated the non availability of both green and dry fodder throughout the year hence, fodder development activities can be taken up in the micro watershed.
- ✓ The average annual gross income of the households Rs.60461.11 from agriculture, Rs.5555.56 from business and Rs. 66736.11 from wages and. Agriculture was found to be the major source of income for households hence; the development activities should focus on productivity enhancement, marketing arrangements and agricultural technology dissemination to have a direct impact on the farmers.
- ✓ The cultivation of forest species is found minimal hence; information and production technology related to agro-forestry and integrated farming system.
- ✓ The data indicated that, 44.44 per cent of the households have experienced soil and water erosion problems. Hence, those farmers who reported the soil and water erosion problems may be given attention while implementation of the watershed development plan.
- ✓ The data indicated that, 83.33 per cent of the households have interest in soil testing hence, farmers must be provided with the information on various institutions which are involved in soil testing for the benefit of the farmers.

- ✓ Except summer ploughing the adoption of other soil and water conservation structures is minimum hence, the farmers in the micro watershed should be sensitized on the use of different conservation structures for soil water conservation.
- ✓ Cereals and pulses found be adequate for per cent of the households respectively hence, farm households and the farm women must be trained on importance of balanced nutrition and role of vegetable, milk, egg, meat in balanced diet.
- ✓ Lower fertility status of the soil (75.00%), wild animal menace on farm field (66.67%), frequent incidence of pest and diseases (83.33%), high cost of fertilizers and plant protection chemicals (88.89%), high rate of interest on credit (83.33%), low price for the agricultural commodities (83.33%), lack of marketing facilities in the area (63.89%), inadequate extension services (30.56%), lack of transport for safe transport of the agricultural produce to the market (63.89%) were the major farming constraints experienced hence, these constraints must be addressed immediately for the welfare of the farmers. Awareness to be created among the farmers to approach nearest KVKs/RSKs and other developmental departments for technical and for subsidized inputs and utilize the well established regulated markets, approaching the contract firms, direct markets to avoid the involvement of middlemen.