



हर कदम, हर डगर  
किसानों का हमसफर  
भारतीय कृषि अनुसंधान परिषद

*Agrisearch with a human touch*

**LAND RESOURCE INVENTORY AND SOCIO-ECONOMIC STATUS OF  
FARM HOUSEHOLDS FOR WATERSHED PLANNING AND  
DEVELOPMENT**

**BELAGIRI-1 (4D5B1H1d) MICROWATERSHED**

**Hattakuni Hobli, Yadgir Taluk and District, Karnataka**

**Karnataka Watershed Development Project – II**

**SUJALA – III**

**World Bank funded Project**



**ICAR – NATIONAL BUREAU OF SOIL SURVEY AND LAND USE PLANNING**



ICAR - NBSS & LUP



**WATERSHED DEVELOPMENT DEPARTMENT  
GOVT. OF KARNATAKA, BANGALORE**



## **About ICAR - NBSS&LUP**

The ICAR-National Bureau of Soil Survey and Land Use Planning (ICAR-NBSS&LUP), Nagpur, a premier Institute of the Indian Council of Agricultural Research (ICAR), was set up during 1976 with the objective to prepare soil resource maps at national, state and district levels and to provide research inputs in soil resource mapping and its applications, land evaluation, land use planning, land resource management, and database management using GIS for optimising land use on different kinds of soils in the country.

The Bureau has been engaged in carrying out soil resource survey, agro-ecological and soil degradation mapping at the country, state and district levels for qualitative assessment and monitoring the soil health towards viable land use planning. The research activities have resulted in identifying the soil potentials and problems, and the various applications of the soil surveys with the ultimate objective of sustainable agricultural development. The Bureau has the mandate to correlate and classify soils of the country and maintain a National Register of all the established soil series. The Institute is also imparting in-service training to staff of the soil survey agencies in the area of soil survey, land evaluation and soil survey interpretations for land use planning. The Bureau in collaboration with Panjab rao Krishi Vidyapeeth, Akola is running post-graduate teaching and research programme in land resource management, leading to M.Sc. and Ph.D. degrees.

**Citation:** Rajendra Hegde, Ramesh Kumar, S.C., B.A. Dhanorkar, S. Srinivas, M. Lalitha, K.V. Niranjana, R.S. Reddy and S.K. Singh (2019). "Land Resource Inventory and Socio-Economic Status of Farm Households for Watershed Planning and Development of Belagiri-1 (4D5B1H1d) Microwatershed, Hattakuni Hobli, Yadgir Taluk and District, Karnataka", ICAR-NBSS&LUP Sujala MWS Publ.256, ICAR – NBSS & LUP, RC, Bangalore. p.131 & 33.

### **TO OBTAIN COPIES,**

**Please write to:**

**Director, ICAR - NBSS & LUP,**

Amaravati Road, NAGPUR - 440 033, India

Phone : (0712) 2500386, 2500664, 2500545 (O)

Telefax : 0712-2522534

E-Mail : [director@nbsslup.ernet.in](mailto:director@nbsslup.ernet.in)

Website URL : [nbsslup.in](http://nbsslup.in)

Or

**Head, Regional Centre, ICAR - NBSS&LUP, Hebbal, Bangalore - 560 024**

Phone : (080) 23412242, 23510350 (O)

Telefax : 080-23510350

E-Mail : [nbssrcb@gmail.com](mailto:nbssrcb@gmail.com)

ICAR-NBSS&LUP Sujala MWS Publ.256



हर कदम, हर डगर  
किसानों का हमसफर  
भारतीय कृषि अनुसंधान परिषद

*Agri*search with a human touch

**LAND RESOURCE INVENTORY AND SOCIO-ECONOMIC  
STATUS OF FARM HOUSEHOLDS FOR WATERSHED  
PLANNING AND DEVELOPMENT**

**BELAGIRI-1 (4D5B1H1d) MICROWATERSHED**

**Hattakuni Hobli, Yadgir Taluk and District, Karnataka**

**Karnataka Watershed Development Project – II**

**Sujala-III**

**World Bank funded Project**



**ICAR – NATIONAL BUREAU OF SOIL SURVEY AND LAND USE  
PLANNING**



ICAR - NBSS & LUP



**WATERSHED DEVELOPMENT DEPARTMENT, GOVT. OF  
KARNATAKA, BANGALORE**







## PREFACE

In Karnataka, as in other Indian States, the livelihoods of rural people are intertwined with farming pursuits. The challenges in agriculture are seriously threatening the livelihood of a large number of farmers as they have been practicing farming in contextual factors beyond their control. Climatic factors are the most important ones and have become much more significant in recent times due to rapid climate changes induced by intensive anthropogenic activities affecting our ecosystem in multiple ways. Climate change has become the reality, it is happening and efforts to evolve and demonstrate climate resilient technologies have become essential. Due to the already over stressed scenario of agrarian sector, the climate change is resulting in manifold increase in the complexities, pushing the rural mass to face more and more unpredictable situations. The rising temperatures and unpredictable rainfall patterns are going to test seriously the informed decisions farmers have to make in order to survive in farming and sustain their livelihood.

It is generally recognized that impacts of climate change shall not be uniform across the globe. It is said that impact of climate change is more severe in South Asia. Based on the analysis of meteorological data, it is predicted that in India, there will be upward trend in mean temperature, downward trend in relative humidity, annual rainfall and number of wet days in a year. Also, in general, phenomena like erratic monsoon, spread of tropical diseases, rise in sea levels, changes in availability of fresh water, frequent floods, droughts, heat waves, storms and hurricanes are predicted. Each one of these adverse situations are already being experienced in various parts of India and also at the global level. Decline in agricultural productivity of small and marginal farmers becoming more vulnerable is already witnessed.

In Karnataka, more than 60 per cent of the population live in rural areas and depend on agriculture and allied activities for their livelihood. Though the state has achieved significant progress in increasing the yield of many crops, there is tremendous pressure on the land resources due to the growing and competing demands of various land uses. This is reflected in the alarming rate of land degradation observed. Already more than 50 per cent of the area is affected by various forms of degradation. If this trend continues, the sustainability of the fragile ecosystem will be badly affected. The adverse effects of change in the climatic factors are putting additional stress on the land resources and the farmers dependent on this.

The natural resources (land, water and vegetation) of the state need adequate and constant care and management, backed by site-specific technological interventions and investments particularly by the government. Detailed database pertaining to the nature of the land resources, their constraints, inherent potentials and suitability for various land

based rural enterprises, crops and other uses is a prerequisite for preparing location-specific action plans, which are in tune with the inherent capability of the resources. Any effort to evolve climate resilient technologies has to be based on the baseline scientific database. Then only one can expect effective implementation of climate resilient technologies, monitor the progress, make essential review of the strategy, and finally evaluate the effectiveness of the implemented programs. The information available at present on the land resources of the state are of general nature and useful only for general purpose planning. Since the need of the hour is to have site-specific information suitable for farm level planning and detailed characterization and delineation of the existing land resources of an area into similar management units is the only option.

ICAR-NBSS&LUP, Regional Centre, Bangalore has taken up a project sponsored by the Karnataka Watershed Development Project-II, (Sujala-III), Government of Karnataka funded by the World Bank under Component-1 Land Resource Inventory. This study was taken up to demonstrate the utility of such a database in reviewing, monitoring and evaluating all the land based watershed development programs on a scientific footing. To meet the requirements of various land use planners at grassroots level, the present study on “Land Resource Inventory and Socio-Economic Status of Farm Households for Watershed Planning and Development of Belagiri-1 Microwatershed, Yadgir Taluk and District, Karnataka” for integrated development was taken up in collaboration with the State Agricultural Universities, IISC, KRSRAC, KSNDMC as Consortia partners. The project provides detailed land resource information at cadastral level (1:7920 scale) for all the plots and socio-economic status of farm households covering thirty per cent farmers randomly selected representing landed and landless class of farmers in the microwatershed. The project report with the accompanying maps for the Microwatershed will provide required detailed database for evolving effective land use plan, alternative land use options and conservation plans for the planners, administrators, agricultural extension personnel, KVK officials, developmental departments and other land users to manage the land resources in a sustainable manner.

It is hoped that this database will be useful to the planners, administrators and developmental agencies working in the area in not only for formulating location specific developmental schemes but also for their effective monitoring at the village/watershed level.

Nagpur

Date: 12-07-2019

**S.K. SINGH**

Director, ICAR - NBSS&LUP, Nagpur

## Contributors

<b>Dr. Rajendra Hegde</b> Principal Scientist, Head & Project Leader, Sujala-III Project ICAR-NBSS&LUP, Regional Centre, Bangalore	<b>Dr. S.K.Singh</b> Director, ICAR-NBSS&LUP Coordinator, Sujala-III Project Nagpur
<b>Soil Survey, Mapping &amp; Report Preparation</b>	
Dr. B.A. Dhanorkar	Sh. R.S. Reddy
Dr. K.V. Niranjana	Sh. Venkata Giriyappa
	Mr. Somashekar T N
	Smt. Chaitra, S.P.
	Dr. Gopali bardhan
	Ms. Arpitha
	Dr. Mahendra Kumar, M.B.
<b>Field Work</b>	
Sh. C.BacheGowda	Sh. Mahesh, D.B.
Sh. Somashekar	Sh. Ashok S Sindagi
Sh. M. Jayaramaiah	Sh. Veerabhadrapa B.
Sh. Paramesha, K.	Sh. Shankarappa
Sh. B. M. Narayana Reddy	Sh. Anand
	Sh. Arun N Kambar.
	Sh Kamalesh Awate
	Sh. Sharaan Kumar Huppar
	Sh. Yogesh H.N.
	Sh. Kalaveerachari R Kammar
<b>GIS Work</b>	
Dr. S.Srinivas	Sh. A.G.Devendra Prasad
Sh. D.H.Venkatesh	Sh. Prakashanaik, M.K.
Smt.K.Sujatha	Sh. Abhijith Sastry, N.S.
Smt. K.V.Archana	Sh. Sudip Kumar Suklabaidya
Sh. N. Maddileti	Sh. Avinash, K.N.
	Sh. Amar Suputhra, S
	Sh. Deepak, M.J.
	Smt. K.Karunya Lakshmi
	Ms. Seema, K.V.
	Ms. A. Rajab Nisha

<b>Laboratory Analysis</b>	
Dr. K.M.Nair	Ms. Steffi Peter
Smt. Arti Koyal	Ms. Thara, V.R
Smt. Parvathy	Ms. Roopa, G.
	Ms. Swati, H.
	Sh. Shantaveera Swami
	Ms. Shwetha, N.K.
	Smt. Ishrat Haji
	Ms. P. Pavan Kumari
	Ms. Padmaja
	Ms. Veena, M.
<b>Socio-Economic Analysis</b>	
Dr. S.C. Ramesh Kumar	Sh. M.K. Prakashanaik
	Ms. Karuna V. Kulkarni
	Mrs. Sowmya A.N
	Sh. Vinod R
	Sh. Basavaraja
	Sh. Vijay Kumar Lamani
	Ms. Sowmya K.B
	Mrs. Prathibha, D.G
	Sh. Rajendra,D
<b>Soil &amp; Water Conservation</b>	
Sh. Sunil P. Maske	
<b>Watershed Development Department, GoK, Bangalore</b>	
Sh. Rajeev Ranjan IFS Project Director & Commissioner, WDD	Dr. A. Natarajan NRM Consultant, Sujala-III Project
Dr. S.D. Pathak IFS Executive Director & Chief Conservator of Forests, WDD	

# **PART-A**

## **LAND RESOURCE INVENTORY**



## Contents

Preface		
Contributors		
Executive Summary		
Chapter 1	Introduction	1
Chapter 2	Geographical Setting	3
2.1	Location and Extent	3
2.2	Geology	3
2.3	Physiography	4
2.4	Drainage	4
2.5	Climate	4
2.6	Natural Vegetation	6
2.7	Land Utilization	6
Chapter 3	Survey Methodology	9
3.1	Base maps	9
3.2	Image Interpretation for Physiography	9
3.3	Field Investigation	12
3.4	Soil Mapping	14
3.5	Laboratory Characterization	15
Chapter 4	The Soils	15
4.1	Soils of granite gneiss landscape	19
Chapter 5	Interpretation for Land Resource Management	35
5.1	Land Capability Classification	35
5.2	Soil Depth	37
5.3	Surface Soil Texture	38
5.4	Soil Gravelliness	39
5.5	Available Water Capacity	40
5.6	Soil Slope	41
5.7	Soil Erosion	42
Chapter 6	Fertility Status	45
6.1	Soil Reaction (pH)	45
6.2	Electrical Conductivity (EC)	45
6.3	Organic Carbon (OC)	45
6.4	Available Phosphorus	47
6.5	Available Potassium	47
6.6	Available Sulphur	47
6.7	Available Boron	48
6.8	Available Iron	48
6.9	Available Manganese	48
6.10	Available Copper	48
6.11	Available Zinc	52
Chapter 7	Land Suitability for Major Crops	53

7.1	Land suitability for Sorghum	53
7.2	Land suitability for Maize	54
7.3	Land suitability for Bajra	55
7.4	Land suitability for Groundnut	56
7.5	Land suitability for Sunflower	57
7.6	Land suitability for Redgram	58
7.7	Land suitability for Bengal gram	59
7.8	Land suitability for Cotton	60
7.9	Land suitability for Chilli	61
7.10	Land suitability for Tomato	62
7.11	Land suitability for Brinjal	63
7.12	Land suitability for Onion	64
7.13	Land suitability for Bhendi	65
7.14	Land suitability for Drumstick	66
7.15	Land suitability for Mango	67
7.16	Land suitability for Guava	68
7.17	Land suitability for Sapota	69
7.18	Land Suitability for Pomegranate	70
7.19	Land Suitability for Musambi	71
7.20	Land Suitability for Lime	72
7.21	Land Suitability for Amla	73
7.22	Land Suitability for Cashew	74
7.23	Land Suitability for Jackfruit	75
7.24	Land Suitability for Jamun	76
7.25	Land Suitability for Custard apple	77
7.26	Land Suitability for Tamarind	78
7.27	Land Suitability for Mulberry	79
7.28	Land Suitability for Marigold	80
7.29	Land Suitability for Chrysanthemum	81
7.30	Land use classes	113
7.31	Proposed Crop Plan	114
Chapter 8	Soil Health Management	117
Chapter 9	Soil and Water conservation Treatment Plan	123
9.1	Treatment Plan	123
9.2	Recommended Soil and Water Conservation measures	127
9.3	Greening of Microwatershed	128
	References	131
	Appendix I	I-X
	Appendix II	XI-XIX
	Appendix III	XIX-XXVIII



## LIST OF TABLES

2.1	Mean Monthly Rainfall, PET, 1/2 PET at Yadgir Taluk & District	5
2.2	Land Utilization in Yadgir district	6
3.1	Differentiating Characteristics used for Identifying Soil Series	14
3.2	Soil map unit description of Belagiri-1Microwatershed	15
4.1	Physical and Chemical Characteristics of Soil Series identified in Belagiri-1microwatershed	26
7.1	Soil-Site Characteristics of Belagiri-1Microwatershed	83
7.2	Land suitability criteria for Sorghum	84
7.3	Land suitability criteria for Maize	85
7.4	Land suitability criteria for Bajra	86
7.5	Land suitability criteria for Groundnut	87
7.6	Land suitability criteria for Sunflower	88
7.7	Land suitability criteria for Red gram	89
7.8	Land suitability criteria for Bengal gram	90
7.9	Land suitability criteria for Cotton	91
7.10	Land suitability criteria for Chilli	92
7.11	Land suitability criteria for Tomato	93
7.12	Land suitability criteria for Brinjal	94
7.13	Land suitability criteria for Onion	95
7.14	Land suitability criteria for Bhendi	96
7.15	Land suitability criteria for Drumstick	97
7.16	Land suitability criteria for Mango	98
7.17	Land suitability criteria for Guava	99
7.18	Land suitability criteria for Sapota	100
7.19	Land suitability criteria for Pomegranate	101
7.20	Land suitability criteria for Musambi	102
7.21	Land suitability criteria for Lime	103
7.22	Land suitability criteria for Amla	104
7.23	Land suitability criteria for Cashew	105
7.24	Land suitability criteria for Jackfruit	106
7.25	Land suitability criteria for Jamun	107
7.26	Land suitability criteria for Custard apple	108
7.27	Land suitability criteria for Tamarind	109

7.28	Land suitability criteria for Mulberry	110
7.29	Land suitability criteria for Marigold	111
7.30	Land suitability criteria for Chrysanthemum	112
7.31	Proposed Crop Plan for Belagiri-1Microwatershed	115

## LIST OF FIGURES

2.1	Location map of Belagiri-1 Microwatershed	3
2.2 a	Granite and granite gneiss rock formation	4
2.3	Rainfall distribution in Yadgir Taluk & District	5
2.4	Current Land use map of Belagiri-1Microwatershed	7
2.5 a & b	Different crops and cropping systems in Belagiri-1Microwatershed	7
3.1	Scanned and Digitized Cadastral map of Belagiri-1Microwatershed	11
3.2	Satellite image of Belagiri-1Microwatershed	11
3.3	Cadastral map overlaid on IRS PAN+LISS IV merged imagery of Belagiri-1Microwatershed	12
3.4	Location of profiles in a transect	13
3.5	Soil phase or management units of Belagiri-1Microwatershed	17
5.1	Land Capability Classification map of Belagiri-1Microwatershed	37
5.2	Soil Depth map of Belagiri-1Microwatershed	38
5.3	Surface Soil Texture map of Belagiri-1Microwatershed	39
5.4	Soil Gravelliness map of Belagiri-1Microwatershed	40
5.5	Soil Available Water Capacity map of Belagiri-1Microwatershed	41
5.6	Soil Slope map of Belagiri-1Microwatershed	42
5.7	Soil Erosion map of Belagiri-1Microwatershed	43
6.1	Soil Reaction (pH) map of Belagiri-1Microwatershed	46
6.2	Electrical Conductivity (EC) map of Belagiri-1Microwatershed	46
6.3	Soil Organic Carbon (OC) map of Belagiri-1Microwatershed	47
6.4	Soil Available Phosphorus map of Belagiri-1Microwatershed	48
6.5	Soil Available Potassium map of Belagiri-1Microwatershed	49
6.6	Soil Available Sulphur map of Belagiri-1Microwatershed	49
6.7	Soil Available Boron map of Belagiri-1Microwatershed	50
6.8	Soil Available Iron map of Belagiri-1Microwatershed	50
6.9	Soil Available Manganese map of Belagiri-1Microwatershed	51
6.10	Soil Available Copper map of Belagiri-1Microwatershed	51
6.11	Soil Available Zinc map of Belagiri-1Microwatershed	52
7.1	Land suitability for Sorghum	54
7.2	Land suitability for Maize	55
7.3	Land suitability for Bajra	56

7.4	Land suitability for Groundnut	57
7.5	Land suitability for Sunflower	58
7.6	Land suitability for Red gram	59
7.7	Land suitability for Bengal gram	60
7.8	Land suitability for Cotton	61
7.9	Land suitability for Chilli	62
7.10	Land suitability for Tomato	63
7.11	Land suitability for Brinjal	64
7.12	Land suitability for Onion	65
7.13	Land suitability for Bhendi	66
7.14	Land suitable for Drumstick	67
7.15	Land suitability for Mango	68
7.16	Land suitability for Guava	69
7.17	Land suitability for Sapota	70
7.18	Land suitability for Pomegranate	71
7.19	Land suitability for Musambi	72
7.20	Land suitability for Lime	73
7.21	Land suitability for Amla	74
7.22	Land suitability for Cashew	75
7.23	Land suitability for Jackfruit	76
7.24	Land suitability for Jamun	77
7.25	Land suitability for Custard apple	78
7.26	Land suitability for Tamarind	79
7.27	Land suitability for Mulberry	80
7.28	Land suitability for Marigold	81
7.29	Land suitability for Chrysanthemum	82
7.30	Land management unit map of Belagiri-1 Microwatershed	114
9.1	Soil and water conservation map of Belagiri-1 Microwatershed	128

## **EXECUTIVE SUMMARY**

*The land resource inventory of Belagiri-1 Microwatershed was conducted using village cadastral maps and IRS satellite imagery on 1:7920 scale. The false colour composites of IRS imagery were interpreted for physiography and the physiographic delineations were used as base for mapping soils. The soils were studied in several transects and a soil map was prepared with phases of soil series as mapping units. Random checks were made all over the area outside the transects to confirm and validate the soil map unit boundaries. The soil map shows the geographic distribution and extent, characteristics, classification, behavior and use potentials of the soils in the microwatershed.*

*The present study covers an area of 602 ha in Yadgir taluk & district, Karnataka. The climate is semiarid and categorized as drought-prone with an average annual rainfall of 866 mm, of which about 652 mm is received during south-west monsoon, 138 mm during north-east and the remaining 76 mm during the rest of the year. An area of 410 ha in the microwatershed is covered by soils, 165 ha by rock outcrops and 28 ha by others (habitation and water body). The salient findings from the land resource inventory are summarized briefly below.*

- ❖ The soils belong to 9 soil series and 15 soil phases (management units) and 5 land management units.*
- ❖ The length of crop growing period is about 120-150 days starting from 1<sup>st</sup> week of June to 4<sup>th</sup> week of October.*
- ❖ From the master soil map, several interpretative and thematic maps like land capability, soil depth, surface soil texture, soil gravelliness, available water capacity, soil slope and soil erosion were generated.*
- ❖ Soil fertility status maps for macro and micronutrients were generated based on the surface soil samples collected at every 320 m grid interval.*
- ❖ Land suitability for growing 29 major agricultural and horticultural crops was assessed and maps showing the degree of suitability along with constraints were generated.*
- ❖ Entire area in the microwatershed is suitable for agriculture.*
- ❖ About <1 per cent area is very shallow (<25 cm), 68 per cent area of the microwatershed has soils that are moderately deep to very deep (75 - >150 cm).*
- ❖ About 40 per cent area in the microwatershed has loamy soils and 28 per cent clayey soils at the surface.*
- ❖ Maximum of 65 per cent area in the microwatershed is non gravelly (<15%) and 3 per cent is gravelly (15-35%).*
- ❖ About 41 per cent area of the microwatershed is very high (>200 mm/m) in available water capacity, 25 per cent is low (51-100 mm/m) and 1 per cent area is very low (<50 mm/m).*
- ❖ About 11 per cent area in the microwatershed has nearly level (0-1% slope) lands, 56 per cent has very gently sloping (1-3% slope) lands and 1 per cent area is gently sloping (3.5% slope).*

- ❖ *An area of about 11 per cent is severely (e3) eroded, 57 per cent is moderately (e2) eroded and 11 per cent area is slightly (e1) eroded.*
- ❖ *An area of about 17 per cent area is neutral (pH 6.5-7.3) in soil reaction, 37 per cent soils is slightly to moderately alkaline (pH 7.3-8.4) and 14 per cent soils are strongly to very strongly alkaline (pH 8.4 - >9.0).*
- ❖ *The Electrical Conductivity (EC) of entire soils of the microwatershed is dominantly  $<2 \text{ dsm}^{-1}$  indicating that the soils are non-saline.*
- ❖ *About 9 per cent of the soils are low (<0.5%) in organic carbon, 55 per cent medium (0.5-0.75%) and high in 4 per cent area.*
- ❖ *4 per cent area is high (>57 kg/ha) in available phosphorus and 64 per cent area is medium (23-57 kg/ha).*
- ❖ *About 60 per cent is medium (145-337 kg/ha) in available potassium and 8 per cent is low (<145 kg/ha).*
- ❖ *Available sulphur is low (<10 ppm) in an area of about 59 per cent and medium (10 -20 ppm) in 9 per cent.*
- ❖ *Available boron is low (<0.5 ppm) in the entire area of the microwatershed.*
- ❖ *Available iron is sufficient (>4.5 ppm) in the entire area of the microwatershed.*
- ❖ *Available manganese and copper are sufficient in all the soils of the microwatershed.*
- ❖ *Available zinc is deficient (<0.6 ppm) in an area of 60 per cent and sufficient (>0.6 ppm) in 8 per cent area of the microwatershed.*
- ❖ *The land suitability for 29 major crops grown in the microwatershed were assessed and the areas that are highly suitable (S1) and moderately suitable (S2) are given below. It is however to be noted that a given soil may be suitable for various crops but what specific crop to be grown may be decided by the farmer looking to his capacity to invest on various inputs, marketing infrastructure, market price and finally the demand and supply position.*

**Land suitability for various crops in the Microwatershed**

<b>Crop</b>	<b>Suitability Area in ha (%)</b>		<b>Crop</b>	<b>Suitability Area in ha (%)</b>	
	<b>Highly suitable (S1)</b>	<b>Moderately suitable (S2)</b>		<b>Highly suitable (S1)</b>	<b>Moderately suitable (S2)</b>
<i>Sorghum</i>	249(41)	94(16)	<i>Guava</i>	-	152(25)
<i>Maize</i>	-	338(56)	<i>Sapota</i>	-	152(25)
<i>Bajra</i>	-	408(68)	<i>Pomegranate</i>	-	401(67)
<i>Groundnut</i>	-	158(26)	<i>Musambi</i>	174(29)	227(38)
<i>Sunflower</i>	132(22)	211(35)	<i>Lime</i>	174(29)	227(38)
<i>Redgram</i>	-	401(67)	<i>Amla</i>	18(3)	330(55)
<i>Bengal gram</i>	249(41)	18(3)	<i>Cashew</i>	-	18(3)
<i>Cotton</i>	132(22)	135(22)	<i>Jackfruit</i>	-	94(16)
<i>Chilli</i>	-	402(67)	<i>Jamun</i>	-	307(51)
<i>Tomato</i>	-	269(45)	<i>Custard apple</i>	322(54)	26(4)
<i>Brinjal</i>	151(25)	255(42)	<i>Tamarind</i>	-	307(51)
<i>Onion</i>	281(47)	10(2)	<i>Mulberry</i>	-	99(17)
<i>Bhendi</i>	281(47)	10(2)	<i>Marigold</i>	-	401(67)
<i>Drumstick</i>	-	406(68)	<i>Chrysanthemum</i>	-	401(67)
<i>Mango</i>	-	16(3)			

- ❖ *Apart from the individual crop suitability, a proposed crop plan has been prepared for the identified LMUs by considering only the highly and moderately suitable lands for different crops and cropping systems with food, fodder, fibre and other horticulture crops.*
- ❖ *Maintaining soil-health is vital for crop production and conserve soil and land resource base for maintaining ecological balance and to mitigate climate change. For this, several ameliorative measures have been suggested for these problematic soils like saline/alkali, highly eroded, sandy soils etc.,*
- ❖ *Soil and water conservation treatment plan has been prepared that would help in identifying the sites to be treated and also the type of structures required.*
- ❖ *As part of the greening programme, several tree species have been suggested to be planted in marginal and submarginal lands, field bunds and also in the hillocks, mounds and ridges. This would help in not only supplementing the farm income but also provide fodder and fuel and generate lot of biomass which would help in maintaining an ecological balance and also contribute to mitigating the climate change.*





## **INTRODUCTION**

Soil being a vital natural resource on whose proper use depends the life supporting systems of a country and the socioeconomic development of its people. Soils provide food, fodder, fibre and fuel for meeting the basic human and animal needs. With the ever increasing growth in human and animal population, the demand on soil for more food and fodder production is on the increase. The area available for agriculture is about 51 per cent of the total geographical area and more than 60 per cent of the people are still dependant on agriculture for their livelihood. However, the capacity of a soil to produce is limited and the limits to the production are set by its intrinsic characteristics, agro-climatic setting, and, use and management. There is, therefore, tremendous pressure on land and water resources, which is causing decline in soil-health and stagnation in productivity. The soils have been degrading at an estimated rate of one million hectares per year and ground water levels have been receding at an alarming rate resulting in decline in the ground water resource. Further, land degradation has emerged as a serious problem which has already affected about 38 lakh ha of cultivated area in the State. Soil erosion alone has degraded about 35 lakh ha. Almost all the uncultivated areas are facing various degrees of degradation, particularly soil erosion; salinity and alkalinity has emerged as a major problem in more than 3.5 lakh ha in the irrigated areas of the State. Nutrient depletion and declining factor productivity is common in both rainfed and irrigated areas. The degradation is continuing at an alarming rate and there appears to be no systematic effort among the stakeholders to contain this process. In recent times, an aberration of weather due to climate change phenomenon has added another dimension leading to unpredictable situation to be tackled by the farmers.

In this critical juncture, the challenge before us is not only to increase the productivity per unit area which is steadily declining and showing a fatigue syndrome, but also to prevent or at least reduce the severity of degradation. If the situation is not reversed at the earliest, then the sustainability of the already fragile crop production system and the overall ecosystem will be badly affected in the state. Added to this, every year there is a significant diversion of farm lands and water resources for non-agricultural purposes. Thus, developing strategies to slow down the degradation process or reclaim the soils to normal condition and ensure sustainability of production system are the major issues today. This demands a systematic appraisal of our soil and land resources with respect to their extent, geographic distribution, characteristics, behaviour and use potential, which is very important for developing an effective land use and cropping systems for augmenting agricultural production on a sustainable basis.

The soil and land resource inventories made so far in Karnataka had limited utility because the surveys were of different types, scales and intensities carried out at different times with specific objectives. Hence, there is an urgent need to generate detailed site-

specific farm level database on various land resources for all the villages/watersheds in a time bound manner that would help to protect the valuable soil and land resources and also to stabilize the farm production. Therefore, the land resource inventory required for farm level planning is the one which investigates all the parameters which are critical for productivity *viz.*, soils, site characteristics like slope, erosion, gravelliness and stoniness, climate, water, topography, geology, hydrology, vegetation, crops, land use pattern, animal population, socio-economic conditions, infrastructure, marketing facilities and various schemes and developmental works of the government etc. From the data collected at farm level, the specific problems and potentials of the area can be identified and highlighted, conservation measures required for the area can be planned on a scientific footing, suitability of the area for various uses can be worked out and finally viable and sustainable land use options suitable for each and every land holding can be prescribed.

The Land Resource Inventory is basically done for identifying potential and problem areas, developing sustainable land use plans, estimation of surface run off and water harvesting potential, preparation of soil and water conservation plans, land degradation/desertification etc. The Bureau is presently engaged in developing an LRI methodology using high resolution satellite remote sensing data and Digital Elevation Model (DEM) data to prepare Landscape Ecological Units (LEU) map representing agro-ecosystem as a whole. The LEU is preferred over landform as the base map for LRI. LEU is the assemblage of landform, slope and land use. An attempt has already been made to upscale the soil resource information from 1:250000 and 1:50000 scale to the LEU map in Goa and in some other states.

The land resource inventory aims to provide site-specific database for Belagiri-1microwatershed in Yadgir Taluk and Yadgir District, Karnataka State for the Karnataka Watershed Development Department. The database was generated by using cadastral map of the village as a base along with high resolution IRS LISS IV and Cartosat-1 merged satellite imagery. Later, an attempt will be made to uplink this LRI data generated at 1:7920 scale under Sujala-III Project to the proposed Landscape Ecological Units (LEUs) map.

The study was organized and executed by the ICAR- National Bureau of Soil Survey and Land Use Planning, Regional Centre, Bangalore under Generation of Land Resource Inventory Data Base Component-1 of the Sujala-III Project funded by the World Bank.

## GEOGRAPHICAL SETTING

### 2.1 Location and Extent

The Belagiri-1 microwatershed is located in the northern part of Karnataka in Yadgir Taluk & District, Karnataka State (Fig.2.1). It comprises parts of Sutharahosalli, Honagera and Belagera villages. It lies between  $16^{\circ} 48'$  and  $16^{\circ} 50'$  North latitudes and  $77^{\circ} 12'$  and  $77^{\circ} 14'$  East longitudes covering an area of about 602 ha. It is about 12 km southeast of Yadgir town and is surrounded by Sutharahosahalli on the north, Belagera on the northeast, east, south and southeast and Honagera village on the western side.




Fig.2.1 Location map of Belagiri-1 Microwatershed

### 2.2 Geology

Major rock formations observed in the microwatershed are granite gneiss (Figs.2.2). They are essentially pink to gray and are coarse to medium grained. They consist primarily of quartz, feldspar, biotite and hornblende. The gray granite gneisses are highly

weathered, fractured and fissured upto a depth of about 10 m. Dolerite dykes and quartz veins are common with variable width and found to occur in Belagiri-1 microwatershed.



Fig.2.2 Granite and granite gneiss rocks

### **2.3 Physiography**

Physiographically, the area has been identified as granite gneiss landscape based on geology. The area has been further subdivided into five landforms, *viz*; mounds/ridges, summits, side slopes and very gently sloping uplands, plains and valleys based on slope and its relief features. The elevation ranges from 399-431 m above MSL. The mounds and ridges are mostly covered by rock outcrops.

### **2.4 Drainage**

The area is drained by several parallel streams like Bori, Amerja and Kanga which finally join the river Bhima along its course. Though, they are not perennial, during rainy season they carry large quantities of rain water. The microwatershed has only few small tanks which are not capable of storing the water that flows during the rainy season. Due to this, the ground water recharge is very much affected. This is reflected in the failure of many bore wells in the villages. If the available rain water is properly harnessed by constructing new tanks and recharge structures at appropriate places in the villages, then the drinking and irrigation needs of the area can be easily met. The drainage network is parallel to sub parallel and dendritic.


### **2.5 Climate**

The Yadgir district lies in the northern plains of Karnataka and falls under semiarid tract of the state and is categorized as drought-prone with total annual rainfall of 866 mm (Table 2.1). Of the total rainfall, maximum of 652 mm is received during the south-west monsoon period from June to September, the north-east monsoon from

October to early December contributes about 138 mm and the remaining 76 mm during the rest of the year. The summer season starts during the middle of February and continues up to the first week of June. The period from December to the middle of February is the coldest season. December is the coldest month with mean daily maximum and minimum temperatures being 29.5<sup>0</sup>C and 10<sup>0</sup>C respectively. During peak summer, temperature shoots up to 45<sup>0</sup>C. Relative humidity varies from 26% in summer to 62% in winter. Rainfall distribution is shown in Figure 2.3. The average Potential Evapo-Transpiration (PET) is 141 mm and varies from a low of 81 mm in December to 199 mm in the month of May. The PET is always higher than precipitation in all the months except July, August and September. Generally, the Length of crop Growing Period (LGP) is 120-150 days and starts from 1<sup>st</sup> week of June to 4<sup>th</sup> week of October.

**Table 2.1 Mean Monthly Rainfall, PET, 1/2 PET at Yadgir Taluk, Yadgir District**

Sl. No.	Months	Rainfall	PET	1/2 PET
1	January	4.30	86.0	43.0
2	February	2.30	125.5	62.7
3	March	15.10	166.0	83.0
4	April	18.50	179.8	89.9
5	May	36.0	198.8	97.9
6	June	118.0	175.1	87.5
7	July	171.80	156.3	78.1
8	August	182.9	150.3	75.1
9	September	179.7	142.0	71.0
10	October	105.3	138.5	69.2
11	November	26.4	97.60	48.6
12	December	6.0	80.90	40.4
<b>Total</b>		<b>866.3</b>		



**Fig 2.3 Rainfall distribution in Yadgir Taluk, Yadgir District**

## 2.6 Natural Vegetation

The natural vegetation is sparse comprising few tree species, shrubs and herbs. The mounds, ridges and boulders occupy very sizeable area which is under thin to moderately thick forest vegetation. Still, there are some remnants of the past forest cover which can be seen in patches in some ridges and hillocks in the microwatershed.

Apart from the continuing deforestation, the presence of large population of goats, sheep and other cattle in the microwatershed is causing vegetative degradation of whatever little vegetation left in the area. The uncontrolled grazing has left no time for the regeneration of the vegetative cover. This leads to the accelerated rate of erosion on the hill slopes resulting in the formation of deep gullies in the foot slopes that eventually result in the heavy siltation of tanks and reservoirs in the microwatershed.

## 2.7 Land Utilization

About 72 per cent area (Table 2.2) in Yadgir district is cultivated at present. An area of about 2 per cent is permanently under pasture, 20 per cent under current fallows and 6 per cent under non-agricultural land, and 5 per cent under currently barren. Forests occupy an area of about 7 per cent and the tree cover is in a very poor state. Most of the mounds, ridges and bouldery areas have very poor vegetative cover. Major crops grown in the area are sorghum, maize, cotton, sunflower, groundnut, red gram and paddy. The cropping intensity is 120 per cent in the taluk. While carrying out land resource inventory, the land use/land cover particulars are collected from all the survey numbers and a current land use map of the microwatershed is prepared. The current land use map prepared shows the arable and non-arable lands, other land uses and different types of crops grown in the area. The current land use map of Belagiri-1 microwatershed is presented in Fig.2.4. The different crops and cropping systems adopted in the microwatershed is presented in the Figures 2.5 a & b.

**Table 2.2 Land Utilization in Yadgir District**

Sl. No.	Agricultural land use	Area ( ha)	Per cent
1.	Total geographical area	516088	-
2.	Total cultivated area	373617	72.4
3.	Area sown more than once	74081	14.3
4.	Cropping intensity	-	119.8
5.	Trees and grooves	737	0.14
6.	Forest	33773	6.54
7.	Cultivable wasteland	2385	0.46
8.	Permanent Pasture land	11755	2.28
9.	Barren land	27954	5.41
10.	Non- Agriculture land	29623	5.73
11.	Current Fallows	105212	20.4






Fig.2.4 Current Land Use map of Belagiri-1Microwatershed



Fig 2.5 a. Different Crops and Cropping Systems in Belagiri-1Microwatershed






Fig 2.5 b. Different Crops and Cropping Systems in Belagiri-1 Microwatershed



## SURVEY METHODOLOGY

The purpose of land resource inventory is to delineate similar areas (soil series and phases), which respond or expected to respond similarly for a given level of management. This was achieved in Belagiri-1 microwatershed by the detailed study of all the soil characteristics (depth, texture, colour, structure, consistence, coarse fragments, porosity, soil reaction, soil horizons etc.) and site characteristics (slope of the land, erosion, drainage, occurrence of rock fragments etc.) followed by grouping of similar areas based on soil-site characteristics into homogeneous (management units) units, and showing the area extent and their geographic distribution on the microwatershed cadastral map. The detailed survey at 1:7920 scale was carried out in an area of 602 ha. The methodology followed for carrying out land resource inventory was as per the guidelines given in Soil Survey Manual (IARI, 1971; Soil Survey Staff, 2006; Natarajan *et al.*, 2015) which is briefly described below.

### 3.1 Base Maps

The detailed survey of the land resources occurring in the microwatershed was carried out by using digitized cadastral map and satellite imagery as base supplied by KSRSAC. The cadastral map shows field boundaries with their survey numbers, location of tanks, streams and other permanent features of the area (Fig. 3.1). Apart from the cadastral map, remote sensing data products from Cartosat-1 and LISS IV merged at the scale of 1:7920 were used in conjunction with the cadastral map to identify the landscapes, landforms and other surface features. The imagery helped in the identification and delineation of boundaries between hills, uplands and lowlands, water bodies, forest and vegetated areas, roads, habitations and other cultural features of the area (Fig. 3.2). The cadastral map was overlaid on the satellite imagery (Fig. 3.3) that helps to identify the parcel boundaries and other permanent features. Apart from cadastral maps and images, toposheets of the area (1:50,000 scale) were also used for initial traversing, identification of geology and landforms, drainage features, present land use and also for selection of transects in the microwatershed.

### 3.2 Image Interpretation for Physiography

False Colour Composites (FCCs) of Cartosat-I and LISS-IV merged satellite data covering microwatershed area was visually interpreted using image interpretation elements and all the available collateral data with local knowledge. The delineated physiographic boundaries were transferred on to a cadastral map overlaid on satellite imagery. Physiographically, the area has been identified as granite gneiss landscape. It was divided into five landforms, *viz.*; ridges and mounds, gently and very gently sloping uplands and lowlands based on slope and image characteristics. They were further

subdivided into physiographic/image interpretation units based on image characteristics. The image interpretation legend for physiography is given below.

## **Image Interpretation Legend for Physiography**

### **G- Granite Gneiss Landscape**

G1	Hills/ Ridges/ Mounds
G11	Summits
G12	Side slopes
G121	Side slopes with dark grey tones
G2	Uplands
G21	Summits
G22	Gently sloping uplands
G221	Gently sloping uplands, yellowish green (eroded)
G222	Gently sloping uplands, yellowish white (severely eroded)
G23	Very gently sloping uplands
G231	Very gently sloping uplands, yellowish green
G232	Very gently sloping uplands, medium green and pink
G233	Very gently sloping uplands, pink and green (scrub land)
G234	Very gently sloping uplands, medium greenish grey
G235	Very gently sloping uplands, yellowish white (eroded)
G236	Very gently sloping uplands, dark green
G237	Very gently sloping uplands, medium pink (coconut garden)
G238	Very gently sloping uplands, pink and bluish white (eroded)
G24	Valleys/ lowlands
G241	Valleys, pink tones
G242	Valleys gray mixed with pink tones




Fig 3.1 Scanned and Digitized Cadastral map of Belagiri-1Microwatershed




Fig.3.2 Satellite Image of Belagiri-1Microwatershed




Fig.3.3 Cadastral map overlaid on IRS PAN+LISS IV merged imagery of Belagiri-1 Microwatershed

### 3.3 Field Investigation

The field boundaries and survey numbers given on the cadastral sheet were located on the ground by following permanent features like roads, cart tracks, *nallas*, streams, tanks etc., and wherever changes were noticed, they were incorporated on the microwatershed cadastral map. Preliminary traverse of the microwatershed was carried out with the help of cadastral map, imagery and toposheets. While traversing, landforms and physiographic units identified were checked and preliminary soil legend was prepared by studying soils at few selected places. Then, intensive traversing of each physiographic unit like hills, ridges, uplands and valleys was carried out. Based on the variability observed on the surface, transects (Fig. 3.4) were selected across the slope covering all the landform units in the microwatershed (Natarajan and Dipak Sarkar, 2010).




Fig: 3.4. Location of profiles in a transect

In the selected transect, soil profiles were located (Fig. 3.4) at closely spaced intervals to take care of any change in the land features like break in slope, erosion, gravel, stones etc. In the selected sites, soil profiles (vertical cut showing the soil layers from surface to the rock) were opened upto 200 cm or to the depth limited by rock or hard substratum and studied in detail for all their morphological and physical characteristics. The soil and site characteristics were recorded for all profile sites on a standard proforma as per the guidelines given in USDA Soil Survey Manual (Soil Survey Staff, 2012). Apart from the transect study, profiles were also studied at random, almost like in a grid pattern, outside the transect areas.

Based on the soil characteristics, the soils were grouped into different soil series. Soil series is the most homogeneous unit having similar horizons and properties and behaves similarly for a given level of management. Soil depth, texture, colour, kind of horizon and horizon sequence, calcareousness, amount and nature of gravel present, nature of substratum etc, were used as the major differentiating characteristics for identifying soil series occurring in the area. The differentiating characteristics used for identifying the soil series are given in Table 3.1. Based on the above characteristics, 9 soil series were identified in the Belagiri-1microwatershed.

**Table 3.1 Differentiating Characteristics used for identifying Soil Series**  
(Characteristics are of Series Control Section)

<b>Soils of Granite gneiss Landscape</b>							
<b>Sl. no</b>	<b>Soil Series</b>	<b>Depth (cm)</b>	<b>Colour (moist)</b>	<b>Texture</b>	<b>Gravel (%)</b>	<b>Horizon sequence</b>	<b>Calcareousness</b>
<b>Soil of Granite and Granite Gneiss Landscape</b>							
1	BDL (Badiyala)	25-50	7.5 YR 2.5/3,2.5/2,3/3 10YR 3/4,4/3	sl	<15	Ap-Bw	e
2	BLC (Balichakra)	75-100	2.5YR 5/3,2.5/4 5YR 4/3,3/3	scl	<15	Ap-Bt	-
3	KBD (Kalabelagundi)	75-100	2.5YR 4/4,3/4 5YR 4/2,4/3	gscl	35-60	Ap-Bt	-
4	HSL (Hosalli)	75-100	10YR 5/4,4/4,4/6	sc	<15	Ap-Bw	e
5	YDR (Yadgir)	100-150	10YR 4/3,4/4 2.5YR 4/3,5/3	sl	<15	Ap-AC	-
6	ANR (Anur)	100-150	10YR 4/3,4/1	c	<15	Ap-Bw	es
7	MDG (Mundargi)	100-150	10YR 4/4,3/3 7.5YR 4/4	scl	<15	Ap-Bw	-
8	MDR (Madhwara)	>150	10YR 3/1,3/2,2/1,2/2	scl	<15	Ap-Bw	e
9	BMN (Bhimanahalli)	>150	10YR 3/1	c	<15	Ap-Bss	es

### 3.4 Soil Mapping

The area under each soil series was further separated into soil phases and their boundaries delineated on the cadastral map based on the variations observed in the texture of the surface soil, slope, erosion, presence of gravel, stoniness etc. A soil phase is a subdivision of soil series based mostly on surface features that affect its use and management. The soil mapping units are shown on the map (Fig.3.5) in the form of symbols. During the survey many soil profile pits, few minipits and a few auger bores representing different landforms occurring in the microwatershed were studied. In addition to the profile study, spot observations in the form of minipits, road cuts, terrace cuts etc., were studied to validate the soil boundaries on the soil map. The soil map shows the geographic distribution of 15 mapping units representing 9 soil series occurring in the microwatershed. The soil map unit (soil legend) description is presented in Table 3.2. The soil phase map (management units) shows the distribution of 15 soil phases mapped in the microwatershed. Each mapping unit (soil phase) delineated on the map has similar soil and site characteristics. In other words, all the farms or survey numbers included in one soil phase will have similar management needs and have to be treated accordingly.

### 3.5 Land Management Units (LMU's)

The 15 soil phases identified and mapped in the microwatershed were grouped into 5 Land Management Units (LMU's) for the purpose of preparing a Proposed Crop Plan for sustained development of the microwatershed. The database (soil phases) generated under LRI was utilized for identifying Land Management Units (LMU's) based on the management needs. One or more than one soil site characteristic having influence on the management have been chosen for identification and delineation of LMUs. For Belagiri-1 microwatershed, five soil and site characteristics, namely soil depth, soil texture, slope, erosion and gravel content have been considered for defining LMUs. The Land Management Units are expected to behave similarly for a given level of management.

### 3.6 Laboratory Characterization

Soil samples for each soil series were collected from representative master profiles for laboratory characterization by following the methods outlined in the Laboratory Manual (Sarma *et al*, 1987). Surface soil samples collected from farmer's fields (57 samples) for fertility status (major and micronutrients) at 320 m grid interval in the year 2017 were analyzed in the laboratory (Katyal and Rattan, 2003). By linking the soil fertility data to the survey numbers through GIS, soil fertility maps were generated by using Kriging method for the microwatershed.

**Table 3.2 Soil map unit description of Belagiri-1Microwatershed**

Soil Map unit No.	Soil Series	Soil Phase	Mapping Unit Description	Area in ha (%)
<b>Soils of Granite Gneiss Landscape</b>				
	BDL		Badiyala soils are shallow (25-50 cm), well drained, have dark brown to very dark brown and dark yellowish brown, slightly calcareous sandy loam soils occurring on very gently to gently sloping uplands under cultivation	<b>3 (0.42)</b>
6		BDLiB3	Sandy clay surface, slope 1-3%,severe erosion	3 (0.42)
	BLC		Balichakra soils are moderately deep (75-100 cm), well drained, have reddish brown to dark reddish brown, sandy clay loam red soils occurring on very gently sloping uplands under cultivation	<b>18(3.03)</b>
37		BLCcB2	Sandy loam surface, slope 1-3%, moderate erosion	17 (2.87)
38		BLCiB2	Sandy clay surface, slope 1-3%, moderate erosion	1 (0.16)
	KBD		Kalabelagundi soils are moderately deep (75-100 cm), well drained, have reddish brown to dark reddish brown and dark reddish gray, gravelly sandy clay loam soils occurring on very gently sloping uplands under cultivation	<b>5(0.9)</b>
130		KBDhB2	Sandy clay loam surface, slope 1-3%,	0 (0.08)

			moderate erosion	
164		KBDcC2g1	Sandy loam surface, slope 3-5%, moderate erosion, gravelly (15-35%)	5 (0.82)
	HSL	Hosalli soils are moderately deep (75-100 cm), moderately well drained, have yellowish brown to dark yellowish brown, slightly calcareous sandy clay soils occurring on very gently sloping uplands under cultivation		<b>76(12.63)</b>
33		HSLiB2	Sandy clay surface, slope 1-3%, moderate erosion	71 (11.81)
126		HSLhB2	Sandy clay loam surface, slope 1-3%, moderate erosion	5 (0.82)
	YDR	Yadgir soils are deep (100-150 cm), well drained, have brown to dark yellowish brown and olive brown, sodic sandy loam soils occurring on very gently sloping uplands under cultivation		<b>58(9.69)</b>
42		YDRcB2	Sandy loam surface, slope 1-3%, moderate erosion	46 (7.71)
43		YDRiB2	Sandy clay surface, slope 1-3%, moderate erosion	12 (1.98)
	ANR	Anur soils are deep (100-150 cm), moderately well drained, have dark gray to brown, calcareous, sodic clay soils occurring on very gently sloping uplands under cultivation		<b>0.07 (0.01)</b>
167		ANRcA1	Sandy loam surface, slope 0-1%, slight erosion	0.07 (0.01)
	MDG	Mundargi soils are deep (100-150 cm), moderately well drained, have brown to dark yellowish brown, sandy clay loam soils occurring on very gently sloping uplands under cultivation		<b>16 (2.67)</b>
149		MDGhB2g1	Sandy clay loam surface, slope 1-3%, moderate erosion, gravelly (15-35%)	16 (2.67)
	MDR	Madhwara soils are very deep (>150 cm), moderately well drained, have very dark gray to very dark brown, slightly calcareous sandy clay loam soils occurring on nearly level to very gently sloping uplands under cultivation		<b>216(35.95)</b>
59		MDRcB2	Sandy loam surface, slope 1-3%, moderate erosion	33 (5.48)
60		MDRiA1	Sandy clay surface, slope 0-1%, slight erosion	66 (11.03)
132		MDRhB2	Sandy clay loam surface, slope 1-3%, moderate erosion	117 (19.44)
	BMN	Bhimanahalli soils are very deep (>150 cm), moderately well drained, have very dark gray, calcareous cracking clay black soils occurring on very gently sloping uplands under cultivation		<b>17 (2.74)</b>
62		BMNmB2	Clay surface, slope 1-3%, moderate erosion	17 (2.74)
999		Rock out crops	Rocklands, both massive and bouldery with little or no soil	165 (27.33)
1000		Others	Habitation and water body	28 (4.62)






Fig 3.5 Soil Phase or Management Units - Belagiri-1 Microwatershed



## THE SOILS

Detailed information pertaining to the nature, extent and their distribution of different kinds of soils occurring in Belagiri-1 microwatershed is provided in this chapter. The microwatershed area has been identified as granite gneiss landscape based on geology. In all, 9 soil series are identified. Soil formation is the result of the combined effect of environmental and terrain factors that are reflected in soil morphology. In the granite gneiss landscape, it is by parent material, relief and climate.

A brief description of each of the 9 soil series identified followed by 15 soil phases (management units) mapped are furnished below. The physical and chemical characteristics of soil series identified in Belagiri-1 microwatershed are given in Table 4.1 along with soil classification. The soils in any one map unit differ from place to place in their depth, texture, slope, gravelliness, erosion or any other site characteristic that affect management. The soil phase map can be used for identifying the suitability of areas for growing specific crops or for other alternative uses and also for deciding the type of conservation structures needed. The detailed information on soil and site-characteristics like soil depth, surface soil texture, slope, erosion, gravelliness, AWC, LCC etc, with respect to each of the soil phase identified is given village/survey number wise for the microwatershed in Appendix-I.

### 4.1 Soils of granite gneiss landscape

In this landscape, 9 soil series are identified and mapped. Of these, MDR series occupies maximum area of 216 ha (36%) followed by HSL 76 ha (13%), YDR 58 ha (10%), BLC 18 ha (3%), BMN 17 ha (3%), MDG 16 ha (3%), KBD 5 ha (1%), BDL 3 ha (<1%) and ANR 0.07 ha (<1%). Brief description of each series identified and number of soil phases mapped is given below.

**4.1.1 Badiyala (BDL) Series:** Badiyala soils are shallow (25-50 cm), well drained, have very dark brown, dark yellow brown and dark brown, slightly calcareous sandy loam soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation. The Badiyala series has been classified as a member of the coarse-loamy, mixed, isohyperthermic family of Fluventic Haplustepts.

The thickness of the solum ranges from 28 to 50 cm. The thickness of A horizon ranges from 4 to 12 cm. Its colour is in 10YR hue with value 3 to 4 and chroma 3 to 4. The texture is loamy sand, sandy clay loam and sandy clay. The thickness of B horizon ranges from 27 to 45 cm. Its colour is in 10 YR and 7.5 YR hue with value 2 to 4 and chroma 3 to 4. Its texture is sandy loam to sandy clay loam and is slightly calcareous. The available water capacity is very low (<50mm/m). Only one phase was identified and mapped.



Landscape and Soil Profile characteristics of Badiyala (BDL) Series

**4.1.2 Balichakra (BLC) Series:** Balichakra soils are moderately deep (75-100 cm), well drained, have dark reddish brown to reddish brown, sandy clay loam soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation. The Balichakra series has been classified as a member of the fine-loamy, mixed, isohyperthermic family of Typic Haplustalfs.

The thickness of the solum ranges from 80 to 100 cm. The thickness of A horizon ranges from 10 to 16 cm. Its colour is in hue 5 YR with value and chroma of 3 to 4. Its texture varies from sandy clay loam and sandy clay. The thickness of B horizon ranges from 70 to 88 cm. Its colour is in hue 2.5 YR and 5 YR with value 3 to 5 and chroma 3 to 4. Its texture is sandy clay loam to sandy clay and is slightly calcareous. The available water capacity is medium (101-150 mm/m). Two phases were identified and mapped.



Landscape and Soil Profile characteristics of Balichakra (BLC) Series



**4.1.3 Kalabelagundi (KBD) Series:** Kalabelagundi soils are moderately deep (75-100 cm), well drained, have dark reddish brown to dark reddish grey and reddish brown, gravelly sandy clay loam soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation. The Kalabelagundi series has been classified as a member of the loamy-skeletal, mixed, isohyperthermic family of Typic Haplustalfs.

The thickness of the solum ranges from 75 to 98 cm. The thickness of A horizon ranges from 10 to 19 cm. Its colour is in hue 5 YR and 7.5 YR with value 3 to 4 and chroma 2 to 4. Its texture varies from loamy sand to sandy loam and sandy clay loam. The thickness of B horizon ranges from 70 to 84 cm. Its colour is in hue 5 YR and 2.5YR with value 3 to 4 and chroma 2 to 4. Its texture is sandy clay loam to sandy clay with gravel content of 35-60 per cent. The available water capacity is very low (<50 mm/m). Two phases were identified and mapped.



Landscape and Soil Profile characteristics of Kalabelagundi (KBD) Series

**4.1.4 Hosalli (HSL) Series:** Hosalli soils are moderately deep (75-100 cm), moderately well drained, have dark yellowish brown to yellowish brown, slightly calcareous sandy clay soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation. The Hosalli series has been classified as a member of the fine, mixed, isohyperthermic family of Typic Haplustepts.

The thickness of the solum ranges from 76 to 100 cm. The thickness of A horizon ranges from 6 to 15 cm. Its colour is in hue 10 YR and 7.5 YR with value 3 to 5 and chroma 2 to 4. Its texture varies from loamy sand to sandy loam and sandy clay loam. The thickness of B horizon ranges from 62 to 93 cm. Its colour is in hue 10 YR with value 3 to 4 and chroma 2 to 4. Its texture varies from sandy clay loam to sandy clay and clay and is slightly calcareous. The available water capacity is medium (101-150 mm/m). Two phases were identified and mapped.



Landscape and Soil Profile characteristics of Hosalli (HSL) Series

**4.1.5 Yadgir (YDR) Series:** Yadgir soils are deep (100-150 cm), well drained, have very dark yellowish brown to light olive brown, sodic sandy loam soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation. The Yadgir series has been classified as a member of the coarse-loamy, mixed isohyperthermic family of Fluventic Haplustepts.

The thickness of the soil ranges from 105 to 145 cm. The thickness of A horizon ranges from 6 to 10 cm. Its colour is in 10 YR hue with value 4 and chroma 3. The texture is loamy sand. The thickness of subsurface horizons range from 95 to 130 cm. Its colour is in 10 YR and 2.5 Y hue with value 4 to 5 and chroma 3 to 4. Texture is loamy sand to sandy loam and sandy clay loam and are sodic soils. The available water capacity is low (51-100 mm/m). Two phases were identified and mapped.



Landscape and Soil Profile characteristics of Yadgir (YDR) Series



**4.1.6 Anur (ANR) Series:** Anur soils are deep (100-150 cm), moderately well drained, have dark gray to dark brown, calcareous sodic clay soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation. The Anur series has been classified as a member of the fine, mixed (calc), isohyperthermic family of Typic Haplustepts.

The thickness of the solum ranges from 102 to 148 cm. The thickness of A-horizon ranges from 9 to 17 cm. Its colour is in 10 YR hue with value 3 to 4 and chroma 2 to 4. The texture ranges from loamy sand to sandy clay loam and sandy clay and are calcareous. The thickness of B horizon ranges from 102 to 135 cm. Its colour is in 10 YR hue with value 3 to 5 and chroma 1 to 6. Texture is sandy clay loam to sandy clay and clay and are calcareous sodic soils. The available water capacity is very high (>200 mm/m). Only one phase was identified and mapped.



Landscape and Soil Profile characteristics of Anur (ANR) Series

**4.1.7 Mundargi (MDG) Series:** Mundargi soils are deep (100-150 cm), moderately well drained, have dark brown to dark yellowish brown, sandy clay loam soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation. The Mundargi series has been classified as a member of the fine-loamy, mixed, isohyperthermic family of Fluventic Haplustepts.

The thickness of the solum ranges from 100 to 149 cm. The thickness of A horizon ranges from 8 to 20 cm. Its colour is in 10 YR hue with value 3 and chroma 1 to 4. The texture ranges from sandy loam to sandy clay loam and sandy clay. The thickness of B horizon ranges from 105 to 140 cm. Its colour is in 10 YR hue with value 3 to 4 and chroma 2 to 4. The texture varies from sandy loam to sandy clay loam and sandy clay. The available water capacity is very high (>200 mm/m). Only one phase was identified and mapped.



Landscape and Soil Profile characteristics of Mundargi (MDG) Series

**4.1.8 Madhwara (MDR) Series:** Madhwara soils are very deep (>150 cm), well drained, have black to very dark brown and very dark gray to very dark grayish brown, slightly calcareous sandy clay loam soils. They are developed from weathered granite gneiss and occur on nearly level to very gently sloping uplands under cultivation. The Madhwara series has been classified as a member of the fine-loamy, mixed, isohyperthermic family of Fluventic Haplustepts.

The thickness of the solum is more than 150 cm. The thickness of A horizon ranges from 10 to 16 cm. Its colour is in 10 YR hue with value 3 to 5 and chroma 2 to 3. Texture varies from sandy clay and clay. The thickness of B horizon is >150 cm. Its colour is in 10 YR hue with value 3 to 5 and chroma 1 to 3. Texture varies from sandy clay loam to sandy clay and is slightly calcareous. The available water capacity is very high (>200 mm/m). Three phases were identified and mapped.



Landscape and Soil Profile characteristics of Madhwara (MDR) Series



**4.1.9 Bhimanahalli (BMN) Series:** Bhimanahalli soils are very deep (>150 cm), moderately well drained, have very dark gray calcareous cracking clay soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation. The Bhimanahalli series has been classified as a member of the fine, smectitic (calc), isohyperthermic family of Typic Haplusterts.

The thickness of the solum is more than 150 cm. The thickness of A horizon ranges from 6 to 13 cm. Its colour is in 10 YR hue with value 3 to 4 and chroma 1 to 2 with clay texture. The thickness of B horizon ranges from 163 to 176 cm. Its colour is in 10 YR hue with value 2 to 4 and chroma 1. Its texture is clay and is calcareous. The available water capacity is very high (>200 mm/m). Only one phase was identified and mapped.



Landscape and Soil Profile characteristics of Bhimanahalli (BMN) Series

**Table: 4.1 Physical and Chemical Characteristics of Soil Series identified in Belagiri-1microwatershed**

**Soil Series:** Badiyala (BDL) **Pedon:** R-5

**Location:** 16°37'10.0"N 77°20'21.5", Gudalagunta village, Balichakra hobli, Yadgir taluk and district

**Analysis at:** NBSS&LUP, Regional Centre, Bengaluru **Classification:** Coarse-loamy, mixed, isohyperthermic Fluventic Haplustepts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-12	Ap	87.13	7.04	5.83	10.03	24.32	23.61	23.51	5.67	<15	ls	6.27	2.44
12-28	Bw1	64.63	13.30	22.07	6.74	13.07	22.30	17.01	5.50	<15	scl	16.34	7.83
28-52	BC	73.11	12.02	14.87	3.93	16.03	26.89	18.41	7.86	<15	sl	12.94	5.47

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO <sub>3</sub>	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP						
	Water	CaCl <sub>2</sub>	M KCl				dS m <sup>-1</sup>	%	%	Ca	Mg					K	Na	Total	cmol kg <sup>-1</sup>	%	%
0-12	6.20	-	-	0.074	1.00	0.00	2.80	0.98	0.14	0.01	3.92	4.20	0.72	93	0.20						
12-28	9.04	-	-	0.253	0.80	3.20	-	-	0.16	0.69	-	16.90	0.77	100	4.09						
28-52	9.41	-	-	0.364	1.10	3.60	-	-	0.16	1.39	-	11.10	0.75	100	12.52						

*Contd...*

**Soil Series:** Balichakra (BLC) **Pedon:** T1/P2

**Location:** 16°33'25.0"N 77°20'52.3"E, Sowrashtralli village, Sydhapura hobli, Yadgir taluk and district

**Analysis at:** NBSS&LUP, Regional Centre, Bengaluru **Classification:** Fine-loamy, mixed, isohyperthermic Typic Haplustalfs

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-8	Ap	65.46	8.38	26.16	12.51	18.72	18.82	10.44	4.96	-	scl	15.15	8.63
8-19	Bt1	63.48	8.16	28.36	12.80	15.84	17.21	12.49	5.14	-	scl	16.45	8.81
19-40	Bt2	52.64	11.58	35.79	13.19	13.19	14.35	8.23	3.69	-	sc	21.49	10.36
40-75	BC	55.14	10.71	34.15	14.10	14.42	14.63	7.53	4.45	-	scl	17.77	8.99

Depth (cm)	pH (1:2.5)			E.C. (1:2.5) dS m <sup>-1</sup>	O.C. %	CaCO <sub>3</sub> %	Exchangeable bases					CEC	CEC/Clay	Base saturation %	ESP %
	Water	CaCl <sub>2</sub>	M KCl				Ca	Mg	K	Na	Total				
0-8	6.75	-	-	0.19	0.72	0.00	12.18	3.10	0.43	0.22	15.92	16.80	0.64	95	1.31
8-19	7.23	-	-	0.12	0.68	0.84	11.37	2.50	0.23	0.18	14.28	14.77	0.52	97	1.24
19-40	7.13	-	-	0.08	0.50	0.48	13.80	2.82	0.18	0.09	16.89	17.66	0.49	96	0.51
40-75	7.07	-	-	0.07	0.35	0.84	13.00	2.90	0.17	0.10	16.16	17.55	0.51	92	0.57

Contd...

**Soil Series:** Kalabelagundi (KBD) **Pedon:** R-13

**Location:** 16°43'78.3''n 77°13'71.4''E, Halagera village, Yadgir hobli, Yadgir taluk and district

**Analysis at:** NBSS&LUP, Regional Centre, Bengaluru **Classification:** Loamy-skeletal, mixed, isohyperthermic Typic Haplustalfs

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-11	Ap	72.35	5.19	22.46	7.19	14.29	19.01	25.28	6.58	15	scl	15.12	8.16
11-35	Bt1	73.20	5.81	20.99	13.66	18.67	16.79	17.62	6.47	20	scl	11.58	7.29
35-64	Bt2	51.68	7.30	41.03	29.41	8.00	4.86	5.62	3.78	40	sc	19.86	14.24
64-89	BC	64.35	3.51	32.15	21.84	12.03	14.87	10.23	5.38	40	scl	16.72	10.36

Depth (cm)	pH (1:2.5)			E.C. (1:2.5) dS m <sup>-1</sup>	O.C. %	CaCO <sub>3</sub> %	Exchangeable bases					CEC	CEC/Clay	Base saturation %	ESP %
	Water	CaCl <sub>2</sub>	M KCl				Ca	Mg	K	Na	Total				
0-11	7.84	-	-	0.604	0.88	0.52	8.69	2.17	0.44	0.49	11.78	11.50	0.51	100	4.27
11-35	5.57	-	-	0.181	0.68	0.00	6.40	1.63	0.18	0.14	8.36	9.10	0.43	92	1.57
35-64	7.42	-	-	0.098	0.44	1.05	15.82	2.34	0.12	0.76	19.04	19.60	0.48	97	3.90
64-89	6.66	-	-	0.165	0.56	0.65	10.45	4.00	0.09	0.43	14.97	15.10	0.47	99	2.86

Contd...

**Soil Series:** Hosalli (HSL) **Pedon:** R-3

**Location:** 16°46'60.3"N 77°05'47.6"E, Mudhanala village, Yadgir hobli, Yadgir taluk and district

**Analysis at:** NBSS&LUP, Regional Centre, Bengaluru **Classification:** Fine, mixed, isohyperthermic Typic Haplustepts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-10	Ap	88.43	5.15	6.42	5.69	6.40	36.04	27.31	12.99	-	s	7.40	2.74
10-30	Bw1	58.47	7.24	34.29	4.26	9.37	19.91	19.28	5.64	-	scl	19.07	11.57
30-50	Bw2	51.43	12.67	35.90	3.49	8.89	16.72	15.87	6.46	<15	sc	21.64	12.44
50-90	Bw3	49.89	13.64	36.47	2.43	2.96	20.61	16.17	7.72	<15	sc	21.12	12.95

Depth (cm)	pH (1:2.5)			E.C. (1:2.5) dS m <sup>-1</sup>	O.C. %	CaCO <sub>3</sub> %	Exchangeable bases					CEC	CEC/Clay	Base saturation %	ESP %
	Water	CaCl <sub>2</sub>	M KCl				Ca	Mg	K	Na	Total				
0-10	7.16	-	-	0.117	0.48	0.00	2.83	1.50	0.15	0.29	4.76	4.90	0.76	97	5.94
10-30	6.91	-	-	0.040	0.36	0.00	10.64	5.43	0.10	0.26	16.43	17.80	0.52	92	1.47
30-50	8.17	-	-	0.182	0.24	1.43	-	-	0.12	0.22	-	19.90	0.55	100	1.08
50-90	8.60	-	-	0.148	0.20	4.29	-	-	0.13	0.16	-	19.70	0.54	100	0.81

Contd...

**Soil Series:** Yadgir (YDR) **Pedon:** R-5

**Location:** 16°35'43.6"N 77°17'06.4"E, Kanikal village, Balichakra hobli, Yadgir taluk and district

**Analysis at:** NBSS&LUP, Regional Centre, Bengaluru **Classification:** Coarse-loamy, mixed, isohyperthermic Fluevntic Haplustepts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-14	Ap	86.59	8.77	4.64	23.19	26.92	14.11	15.22	7.16	-	ls	6.97	2.68
14-43	C1	73.39	11.31	15.30	6.76	20.27	24.87	15.66	5.83	-	sl	12.14	7.22
43-89	C2	80.41	3.75	15.84	8.06	13.47	36.73	15.71	6.43	-	sl	22.84	10.18
89-110	C3	63.55	5.40	31.05	8.10	23.05	19.00	9.87	3.53	15-35	scl	38.46	17.70

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO <sub>3</sub>	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP
	Water	CaCl <sub>2</sub>	M KCl				Ca	Mg	K	Na	Total				
				dS m <sup>-1</sup>	%	%	cmol kg <sup>-1</sup>					%	%		
0-14	7.25	-	-	0.114	0.56	0.00	2.29	0.86	0.07	0.03	3.25	3.40	0.73	96	0.31
14-43	9.47	-	-	0.371	0.32	1.30	14.71	4.28	0.38	1.54	20.91	12.70	0.83	165	4.86
43-89	10.30	-	-	0.820	0.16	0.52	1.70	0.98	0.15	6.62	9.45	8.61	0.54	110	30.77
89-110	10.80	-	-	1.440	0.12	0.91	1.02	2.00	0.29	14.43	17.74	16.17	0.52	110	35.688

Contd...

**Soil Series:** Anur (ANR) **Pedon:** R-15

**Location:** 16°32'45.0"N 77°23'57.4"E, Duppalli village, Sydhapura hobli, Yadgir taluk and district

**Analysis at:** NBSS&LUP, Regional Centre, Bengaluru **Classification:** Fine, mixed (calc), isohyperthermic Typic Haplustepts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-18	Ap	64.60	13.44	21.96	7.33	10.42	18.68	20.12	8.05	<15	scl	16.59	7.96
18-49	Bw1	56.66	12.19	31.15	4.73	9.80	18.66	17.02	6.45	-	scl	33.38	13.51
49-95	Bw2	39.94	17.81	42.25	3.09	3.30	15.44	10.65	7.45	<15	c	44.68	25.23
95-123	Bw3	30.65	17.58	51.77	1.50	5.57	10.18	9.65	3.75	<15	c	54.94	32.07

Depth (cm)	pH (1:2.5)			E.C. (1:2.5) dS m <sup>-1</sup>	O.C. %	CaCO <sub>3</sub> %	Exchangeable bases					CEC	CEC/Clay	Base saturation %	ESP %
	Water	CaCl <sub>2</sub>	M KCl				Ca	Mg	K	Na	Total				
0-18	10.17	-	-	0.365	0.48	6.11	-	-	0.25	3.52	-	19.90	0.91	100	7.08
18-49	10.32	-	-	1.38	0.30	6.76	-	-	0.21	16.03	-	24.60	0.79	100	26.07
49-95	10.08	-	-	2.55	0.17	6.11	-	-	0.33	21.49	-	32.60	0.77	100	26.36
95-123	9.92	-	-	2.56	0.12	7.93	-	-	0.51	26.03	-	36.00	0.70	100	28.92

Contd...

**Soil Series:** Mundargi (MDG) **Pedon:** R-2

**Location:** 16°46'82.4"N 77°04'85.2"E, Thumakura village, Yadgir hobli, Yadgir taluk and district

**Analysis at:** NBSS&LUP, Regional Centre, Bengaluru **Classification:** Fine-loamy, mixed, isohyperthermic Fluventic Haplustepts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-9	Ap	81.23	12.97	5.80	4.84	10.19	14.83	37.94	13.42	<15	ls	11.75	3.31
9-20	A2	76.82	16.19	6.98	4.96	10.12	20.75	27.53	13.46	-	ls	14.52	3.99
20-46	Bw1	42.43	17.43	40.15	2.26	5.59	11.49	14.93	8.16	-	c	34.90	21.14
46-90	Bw2	54.51	16.56	28.93	4.72	5.03	19.92	16.67	8.18	-	scl	36.73	18.88
90-110	Bw3	53.69	11.00	35.30	9.57	9.89	16.23	13.01	4.99	-	sc	38.72	20.53

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO <sub>3</sub>	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP
	Water	CaCl <sub>2</sub>	M KCl				Ca	Mg	K	Na	Total				
				dS m <sup>-1</sup>	%	%	cmol kg <sup>-1</sup>							%	%
0-9	8.2	-	-	0.399	0.44	0.78	-	-	0.16	0.38	-	4.90	0.84	100	3.08
9-20	8.44	-	-	0.075	0.29	1.82	-	-	0.05	0.35	-	4.90	0.70	100	2.88
20-46	9.39	-	-	0.451	0.32	2.73	-	-	0.12	5.22	-	20.77	0.52	100	10.06
46-90	9.75	-	-	0.616	0.24	3.25	-	-	0.12	5.72	-	16.56	0.57	100	13.82
90-110	9.72	-	-	0.725	0.24	3.64	-	-	0.14	6.84	-	19.76	0.56	100	13.836

Contd...



**Soil Series:** Madhawara (MDR) **Pedon:** T<sub>2</sub> P<sub>2</sub>

**Location:** 16°43'48.9"N 77°18'38.3"E, Yaleri village, Balichakra hobli, Yadgir taluk and district

**Analysis at:** NBSS&LUP, Regional Centre, Bengaluru **Classification:** Fine-loamy, mixed, isohyperthermic Fluventic Haplustepts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-11	Ap	58.94	20.74	20.32	5.41	7.28	13.31	20.89	12.06	-	scl	16.47	8.85
11-30	Bw1	55.52	19.32	25.16	5.00	7.19	13.12	19.69	10.52	-	scl	18.25	10.18
30-53	Bw2	53.95	19.15	26.90	4.68	7.48	12.58	19.65	9.56	-	scl	26.99	14.02
53-117	Bw3	52.68	19.51	27.81	2.84	5.47	14.72	20.82	8.83	-	scl	37.86	17.40
117-160	Bw4	49.95	17.27	32.79	2.11	5.07	14.15	20.49	8.13	-	scl	44.15	20.38

Depth (cm)	pH (1:2.5)			E.C. (1:2.5) dS m <sup>-1</sup>	O.C. %	CaCO <sub>3</sub> %	Exchangeable bases					CEC	CEC/Clay	Base saturation %	ESP %
	Water	CaCl <sub>2</sub>	M KCl				Ca	Mg	K	Na	Total				
0-11	8.31	-	-	0.33	0.46	2.76	-	-	0.45	0.47	-	20.57	1.01	100	0.90
11-30	9.25	-	-	0.20	0.31	4.20	-	-	0.19	1.40	-	23.98	0.95	100	2.34
30-53	9.78	-	-	0.40	0.19	5.76	-	-	0.16	1.53	-	24.53	0.91	100	2.49
53-117	9.94	-	-	0.88	0.23	4.80	-	-	0.18	9.09	-	24.31	0.87	100	14.96
117-160	9.98	-	-	0.93	0.15	3.00	-	-	0.24	11.09	-	28.27	0.86	100	15.69

Contd...

**Soil Series:** Bhimanahalli (BMN) **Pedon:** R-3

**Location:** 16°31'82.4"N 77°12'70.8"E, Bheemanahalli village, Sydhapura hobli, Yadgir taluk and district

**Analysis at:** NBSS&LUP, Regional Centre, Bengaluru **Classification:** Fine, smectitic (calc), isohyperthermic Typic Haplusterts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-8	Ap	20.34	19.94	59.72	2.68	5.03	3.75	5.25	3.64	-	c	50.19	33.49
8-40	Bss1	19.61	22.76	57.62	1.94	2.59	5.28	4.96	4.85	-	c	43.22	29.05
40-70	Bss2	21.25	17.65	61.10	3.02	5.26	3.91	5.48	3.58	-	c	44.30	30.25
70-120	Bss3	19.08	22.29	58.63	1.75	5.04	3.84	5.15	3.29	-	c	43.26	30.31
120-170	Bss4	11.11	20.44	68.45	2.04	1.93	1.70	2.83	2.61	-	c	51.33	33.51

Depth (cm)	pH (1:2.5)			E.C. (1:2.5) dS m <sup>-1</sup>	O.C. %	CaCO <sub>3</sub> %	Exchangeable bases					CEC	CEC/Clay	Base saturation %	ESP %
	Water	CaCl <sub>2</sub>	M KCl				Ca	Mg	K	Na	Total				
0-8	8.2	-	-	0.284	0.72	4.94	-	-	1.20	0.34	-	52.70	0.88	100	0.65
8-40	8.44	-	-	0.139	0.40	7.28	-	-	0.30	0.48	-	52.06	0.90	100	0.93
40-70	8.32	-	-	0.202	0.40	6.37	-	-	0.18	0.40	-	52.52	0.86	100	0.77
70-120	9.3	-	-	0.282	0.36	6.89	-	-	0.27	0.38	-	50.97	0.87	100	0.75
120-170	8.47	-	-	0.305	0.37	8.19	-	-	0.28	0.91	-	58.19	0.85	100	1.57

## INTERPRETATION FOR LAND RESOURCE MANAGEMENT

The most important soil and site characteristics that affect the land use and conservation needs of an area are land capability, land irrigability, soil depth, soil texture, coarse fragments, available water capacity, soil slope, soil erosion, soil reaction etc. These are interpreted from the data base generated through land resource inventory and several thematic maps are generated. These would help in identifying the areas suitable for growing crops and, soil and water conservation measures and structures needed thus helping to maintain good soil health for sustained crop production. The various interpretative and thematic maps generated are described below.

### 5.1 Land Capability Classification

Land capability classification is an interpretative grouping of soil map units (soil phases) mainly based on inherent soil characteristics, external land features and environmental factors that limit the use of land for agriculture, pasture, forestry, or other uses on a sustained basis (IARI, 1971). The land and soil characteristics used to group the land resources in an area into various land capability classes, subclasses and units are

*Soil Characteristics:* Depth, texture, gravelliness, calcareousness.

*Land characteristics:* Slope, erosion, drainage, rock outcrops.

*Climate:* Total rainfall and its distribution, and length of crop growing period.

The land capability classification system is divided into land capability classes, subclasses and units based on the level of information available. Eight land capability classes are recognized. They are

*Class I:* They are very good lands that have no limitations or very few limitations that restrict their use.

*Class II:* They are good lands that have minor limitations and require moderate conservation practices.

*Class III:* They are moderately good lands that have moderate limitations that reduce the choice of crops or that require special conservation practices.

*Class IV:* They are fairly good lands that have very severe limitations that reduce the choice of crops or that require very careful management.

*Class V:* Soils in these lands are not likely to erode, but have other limitations like wetness that are impractical to remove and as such not suitable for agriculture, but suitable for pasture or forestry with minor limitations.

*Class VI:* The lands have severe limitations that make them generally unsuitable for cultivation, but suitable for pasture or forestry with moderate limitations.

*Class VII:* The lands have very severe limitations that make them unsuitable for cultivation, but suitable for pasture or forestry with major limitations.

*Class VIII:* Soil and other miscellaneous areas (rock lands) that have very severe limitations that nearly preclude their use for any crop production, but suitable for wildlife, recreation and installation of wind mills.

The land capability subclasses are recognised based on the dominant limitations observed within a given land capability class. The subclasses are designated by adding a lower case letter like 'e', 'w', 's', or 'c' to the class numeral. The subclass "e" indicates that the main hazard is risk of erosion, "w" indicates drainage or wetness as a limitation for plant growth, "s" indicates shallow soil depth, coarse or heavy textures, calcareousness, salinity/alkalinity or gravelliness and "c" indicates limitation due to climate.

The land capability subclasses have been further subdivided into land capability units based on the kinds of limitations present in each subclass. Ten land capability units are used in grouping the soil map units. They are stony or rocky (0), erosion hazard (slope, erosion) (1), coarse texture (sand, loamy sand, sandy loam) (2), fine texture (cracking clay, silty clay) (3), slowly permeable subsoil (4), coarse underlying material (5), salinity/alkali (6), stagnation, overflow, high ground water table (7), soil depth (8) and fertility problems (9). The capability units thus identified have similar soil and land characteristics that respond similarly to a given level of management. The soils of the microwatershed have been classified upto land capability subclass level.

The 15 soil map units identified in Belagiri-1 microwatershed are grouped under 3 land capability classes and 5 land capability subclasses. An entire area of 410 ha (68%) in the microwatershed is suitable for agriculture. About 165 ha (27%) area is having rock outcrops and about 28 ha (5%) is covered by others (water body & habitation) (Fig. 5.1).

Good cultivable lands (Class II) cover an area of about 67 per cent and are distributed in the major part of the microwatershed with minor problems of soil and erosion. Moderately good cultivable lands (Class III) cover an area of about 1 per cent and are distributed in the northern part of the microwatershed with moderate problems of soil and erosion. Fairly good cultivable lands (Class IV) lands occur in <1 per area of the microwatershed, and they have severe problem of soil and erosion and distributed in the southern part of the microwatershed.




Fig. 5.1 Land Capability map of Belagiri-1 Microwatershed

## 5.2 Soil Depth

Soil depth refers to the depth of the soil occurring above the parent material or hard rock. The depth of the soil determines the effective rooting depth for plants and in accordance with soil texture, mineralogy and gravel content, the capacity of the soil column to hold water and nutrient availability. Soil depth is one of the most important soil characteristic that is used in differentiating soils into different soil series. The soil depth classes used in identifying soils in the field are very shallow (<25 cm), shallow (25-50 cm), moderately shallow (50-75 cm), moderately deep (75-100 cm), deep (100-150 cm) and very deep (>150 cm). They were used to classify the soils into different depth classes and a soil depth map was generated. The area extent and their geographical distribution in the microwatershed is given in Fig. 5.2.

Shallow (25-50 cm) soils occur in a small area of 3 ha (<1%) and are distributed in the southern part of the microwatershed. Moderately deep (75-100 cm) soils occur in an area of 100 ha (17%) and is distributed in the northern, central and southern part of the microwatershed. Deep spoils occur in an area of 74 ha (12%) and are distributed in all parts of the microwatershed. Very deep (>150 cm) soils cover a maximum area of 233 ha (39%) and are distributed in the major part of the microwatershed.

The most productive lands covering 307 ha (51%) with respect to soil rooting depth where all climatically adapted annual and perennial crops can be grown are deep to very deep (100 - >150 cm depth) soils occurring in the major part of the microwatershed.

The problem soils occupy only 3 ha (<1%) area where only short duration crops can be grown occasionally and the probability of crop failure is very high.




Fig. 5.2 Soil Depth map of Belagiri-1 Microwatershed

### 5.3 Surface Soil Texture

Texture is an expression to indicate the coarseness or fineness of the soil as determined by the relative proportion of primary particles of sand, silt and clay. It has a direct bearing on the structure, porosity, adhesion and consistence. The surface layer of a soil to a depth of about 25 cm is the layer that is most used by crops and plants. The surface soil textural class provides a guide to understanding soil-water retention and availability, nutrient holding capacity, infiltration, workability, drainage, physical and chemical behaviour, microbial activity and crop suitability. The textural classes used for LRI were used to classify and a surface soil texture map was generated. The area extent and their geographical distribution in the microwatershed is shown in Figure 5.3.

Maximum area of about 241 ha (40%) of the microwatershed has loamy soils at the surface and are distributed in all parts of the microwatershed. An area of 170 ha (28%) of the microwatershed has soils that are clayey. Both soils have high potential for soil-water retention and availability, and nutrient retention and availability, but clay soils have more problems of drainage, infiltration, workability and other physical problems.






Fig. 5.3 Surface Soil Texture map of Belagiri-1 Microwatershed

#### 5.4 Soil Graveliness

Gravel is the term used for describing coarse fragments between 2 mm and 7.5 cm diameter and stones for those between 7.5 cm and 25 cm. The presence of gravel and stones in soil reduces the volume of soil responsible for moisture and nutrient storage, drainage, infiltration and runoff, and hinders plant growth by impeding root growth and seedling emergence, intercultural operations and farm mechanization. The graveliness classes used in LRI were used to classify the soils and using these classes a graveliness map was generated. The area extent and their geographic distribution in the microwatershed is shown in Figure 5.4.

Non gravelly (<15%) soils cover maximum area of 389 ha (65%) of the microwatershed. These are the most productive soils, where all climatically adapted short and long duration crops can be grown. Gravelly (15-35%) soils occur in an area of 21 ha (3%) and distributed in the southern and northern part of the microwatershed; these lands are low in moisture holding capacity and hence growing of short duration crops is ideal with best management practice.




Fig. 5.4 Soil Gravelliness map of Belagiri-1 Microwatershed

### 5.5 Available Water Capacity

The soil available water capacity (AWC) is estimated based on the ability of the soil column to retain water between the tensions of 0.33 and 15 bar in a depth of 100 cm or the entire solum if the soil is shallower. The AWC of the soils (soil series) as estimated by considering the soil texture, mineralogy, soil depth and gravel content (Sehgal *et al.*, 1990) and accordingly the soil map units were grouped into five AWC classes *viz*, very low (<50 mm/m), low (50-100 mm/m), medium (100-150 mm/m), high (150-200 mm/m) and very high (>200 mm/m) and using these values, an AWC map was generated. The area extent and their geographic distribution of different AWC classes in the microwatershed is given in Figure 5.5.

Small area of about 8 ha (1%) in the microwatershed has soils that are very low (<50 mm/m) in available water capacity and is distributed in the southern and northern part of the microwatershed. An area of about 153 ha (25%) is low (51-100 mm/m) in available water capacity and are distributed in all parts of the microwatershed. Very high (>200 mm/m) in 249 ha (41%) and are distributed in the major part of the microwatershed.

An area of about 161 ha (27%) in the microwatershed has soils that are problematic with regard to available water capacity. Here, only short duration crops can be grown and probability of the crop failure is very high. These areas are best put to other alternative uses. An area of 249 ha (41%) are potential areas with regard to AWC where all climatically adapted annual and perennial crops can be grown.






Fig. 5.5 Soil Available Water Capacity map of Belagiri-1 Microwatershed

## 5.6 Soil Slope

Soil slope refers to the inclination of the surface of the land. It is defined by gradient, shape and length, and is an integral feature of any soil as a natural body. Slope is considered important in soil genesis, land use and land development. The length and gradient of slope influences the rate of runoff, infiltration, erosion and deposition. The soil map units were grouped into single slope class and a slope map was generated showing the area extent and their geographic distribution in the microwatershed (Fig. 5.6).

An area of about 67 ha (11%) of the microwatershed falls under nearly level (0-1% slope) and 338 ha (56%) under very gently sloping (1-3% slope) lands, thus these areas have high potential in respect of soil slopes. In these areas, all climatically adapted annual and perennial crops can be grown without much soil and water conservation and other land development measures. Gently sloping (3-5%) lands occur in 5 ha (1%) and are distributed in the northern part of the microwatershed. In these areas the soil and water conservation measures should be adopted in order to increase the productivity of soils.




Fig. 5.6 Soil Slope map of Belagiri-1 Microwatershed

## 5.7 Soil Erosion

Soil erosion refers to the wearing away of the earth's surface by the forces of water, wind and ice involving detachment and transport of soil by raindrop impact. It is used for accelerated soil erosion resulting from disturbance of the natural landscape by burning, excessive grazing and indiscriminate felling of forest trees and tillage, all usually by man. The erosion classes showing an estimate of the current erosion status as judged from field observations in the form of rills, gullies or a carpet of gravel on the surface are recorded. Four erosion classes, viz, slight erosion (e1), moderate erosion (e2), severe erosion (e3) and very severe erosion (e4) are recognized. The soil map units were grouped into different erosion classes and a soil erosion map generated. The area extent and their spatial distribution in the microwatershed is given in Figure 5.7.

Soils that are slightly eroded (e1 class) cover an area of 67 ha (11%) and are distributed in the central part of the microwatershed. Moderately eroded (e2 class) soils cover a maximum area of 341 ha (57%) and are distributed in the major part of the microwatershed. Severely eroded soils (e3 class) cover an area of 3 ha (<1%) and are distributed in the southern part of the microwatershed.

An area of about 344 ha of the microwatershed is problematic because of moderate and severe erosion. For these areas, taking up of soil and water conservation and other land development measures are needed.




Fig. 5.7 Soil Erosion map of Belagiri-1 Microwatershed



## FERTILITY STATUS

Soil fertility plays an important role in increasing crop yield. The adoption of high yielding varieties that require high amounts of nutrients has resulted in deficiency symptoms in crops and plants due to imbalanced fertilization and poor inherent fertility status as these areas are characterised by low rainfall and high temperatures. Hence, it is necessary to know the fertility (macro and micro nutrients) status of the soils of the watersheds for assessing the kind and amount of fertilizers required for each of the crop intended to be grown. For this purpose, the surface soil samples collected from the grid points (one soil sample at every 320 m interval) all over the microwatershed through land resource inventory in the year 2017 were analysed for pH, EC, organic carbon, available phosphorus and potassium, and for micronutrients like zinc, boron, copper, iron and manganese, and secondary nutrient sulphur.

Soil fertility data generated has been assessed and individual maps for all the nutrients for the microwatershed have been generated using Kriging method under GIS. The village/survey number wise fertility data for the microwatershed is given in Appendix-II.

### 6.1 Soil Reaction (pH)

The soil analysis of the Belagiri-1 microwatershed for soil reaction (pH) showed that an area of 104 ha (17%) is neutral (pH 6.5-7.3) and are distributed in the northwestern, southwestern, central and northern part of the microwatershed. Slightly alkaline (pH 7.3-7.8) occur in 119 ha (20%) area and are distributed in the central, western and northern part of the microwatershed. An area of about 105 ha (17%) is moderately alkaline (pH 7.8-8.4) and are distributed in the central, western, and southern part of the microwatershed. An area of about 54 ha (9%) is strongly alkaline (pH 8.4-9.0) and are distributed in the southern and central part of the microwatershed. Very strongly alkaline (pH >9.0) soils occur in 27 ha (5%) and is distributed in the southern part of the microwatershed. (Fig. 6.1).

### 6.2 Electrical Conductivity (EC)

The Electrical Conductivity in entire area of the microwatershed is non saline (<2 dS/m) (Fig. 6.2) and as such the soils are nonsaline.

### 6.3 Organic Carbon

The soil organic carbon content (an index of available Nitrogen) in the soils of the microwatershed is high (>0.75 %) in 24 ha (4%) and are distributed in the southeastern part of the microwatershed. Medium (0.5-0.75%) in about 331 ha (55%) and are distributed in the major part of the microwatershed, whereas low (<0.5%) in an area of about 55 ha (9%) and are distributed in the western part of the microwatershed (Fig. 6.3).






Fig.6.1 Soil Reaction (pH) map of Belagiri-1Microwatershed




Fig.6.2 Electrical Conductivity (EC) map of Belagiri-1Microwatershed




Fig.6.3 Soil Organic Carbon map of Belagiri-1 Microwatershed

#### 6.4 Available Phosphorus

Available phosphorus content is medium (23-57 kg/ha) in a maximum area of about 384 ha (64%) and occur in the major part of the microwatershed. High (>57 kg/ha) in an area of 26 ha (4%) and are distributed in the southeastern part of the microwatershed (Fig. 6.4).

#### 6.5 Available Potassium

Available potassium content is medium (145-337 kg/ha) in maximum area of about 364 ha (60%) and are distributed in the major part of the microwatershed (Fig. 6.5). Low (<145 kg/ha) in an area of 46 ha (8%) and are distributed in the western part of the microwatershed.

#### 6.6 Available Sulphur

Maximum area of about 354 ha (59%) is low (<10 ppm) in available sulphur content and are distributed in the major part of the microwatershed. Medium (10-20 ppm) in an area of about 56 ha (9%) and is distributed in the southern part of the microwatershed (Fig. 6.6).

### 6.7 Available Boron

Available boron content is low (<0.5 ppm) in the entire area of the microwatershed (Fig. 6.7).

### 6.8 Available Iron

Available iron content is sufficient (>4.5 ppm) in the entire area of the microwatershed (Fig 6.8).

### 6.9 Available Manganese

Available manganese content is sufficient (>1.0 ppm) in the entire microwatershed area (Fig 6.9).

### 6.10 Available Copper

Available copper content is sufficient (>0.2 ppm) in the entire microwatershed area (Fig 6.10).




Fig.6.4 Soil Available Phosphorus map of Belagiri-1Microwatershed






Fig.6.5 Soil Available Potassium map of Belagiri-1Microwatershed




Fig.6.6 Soil Available Sulphur map of Belagiri-1Microwatershed




Fig.6.7 Soil Available Boron map of Belagiri-1Microwatershed




Fig.6.8 Soil Available Iron map of Belagiri-1Microwatershed






Fig.6.9 Soil Available Manganese map of Belagiri-1 Microwatershed




Fig.6.10 Soil Available Copper map of Belagiri-1 Microwatershed

## 6.11 Available Zinc

Available zinc content is deficient (<0.6 ppm) in maximum area of 362 ha (60%) and is distributed in the major part of the microwatershed. About 48 ha (8%) is sufficient (>0.6 ppm) and is distributed in the northern and central part of the microwatershed (Fig 6.11).




Fig.6.11 Soil Available Zinc map of Belagiri-1Microwatershed

## LAND SUITABILITY FOR MAJOR CROPS

The soil and land resource units (soil phases) of Belagiri-1 microwatershed were assessed for their suitability for growing food, fodder, fibre and other horticulture crops by following the procedure as outlined in FAO, 1976 and 1983. Crop requirements were developed for each of the crop from the available research data and also by referring to Naidu *et. al.* (2006) and Natarajan *et. al.* (2015). The soil and land characteristics were matched with the crop requirement to arrive at the crop suitability. The soil and land characteristics table (Table 7.1) and crop requirement tables (Table 7.2 to Table 7.30) are given at the end. In FAO land suitability classification, two orders are recognized. Order S-Suitable and Order N-Not suitable. The orders have classes, subclasses and units. Order-S has three classes, Class S1-Highly Suitable, Class S2-Moderately Suitable and Class S3- Marginally Suitable. Order N has two classes, N1-Currently not Suitable and N2- Permanently not Suitable. There are no subclasses within the Class S1 as they will have very minor or no limitations for crop growth. Classes S2, S3, N1 and N2 are divided into subclasses based on the kinds of limitations encountered. The limitations that affect crop production are ‘c’ for erratic rainfall and its distribution and length of growing period (LGP), ‘e’ for erosion hazard, ‘r’ for rooting condition, ‘t’ for lighter or heavy texture, ‘g’ for gravelliness or stoniness, ‘n’ for nutrient availability, ‘l’ for topography, ‘m’ for moisture availability, ‘w’ for drainage, ‘s’ for sodium and ‘z’ for calcareousness. These limitations are indicated as lower case letters to the Class symbol. For example, moderately suitable lands with the limitations of soil depth and erosion are designated as S2re. For the microwatershed, the soil mapping units were evaluated and classified up to subclass level.

Using the above criteria, the soil map units of the microwatershed were evaluated and land suitability maps for 29 major agricultural and horticultural crops were generated. The detailed information on the kind of suitability of each of the soil phase for the crops assessed are given village/ survey number wise for the microwatershed in Appendix-III.

### 7.1 Land Suitability for Sorghum (*Sorghum bicolor*)

Sorghum is one of the major crop grown in Karnataka in an area of 10.47 lakh ha in Bijapur, Gulbarga, Raichur, Bidar, Belgaum, Dharwad, Bellary, Chitradurga, Mysore and Tumakuru districts. The crop requirements for growing sorghum (Table 7.2) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing sorghum was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.1.

Highly suitable (Class S1) lands for growing sorghum occur in a maximum area of 249 ha (41%) and are distributed in the major part of the microwatershed. An area of

about 94 ha (16%) is moderately suitable (Class S2) for growing sorghum and are distributed in the northern, central and southern part of the microwatershed. They have minor limitations of calcareousness and texture. An area of about 66 ha (11%) is marginally suitable (Class S3) for growing sorghum and is distributed in all parts of the microwatershed with moderate limitations rooting depth, gravelliness, calcareousness and texture.




Fig. 7.1 Land Suitability map of Sorghum

## 7.2 Land Suitability for Maize (*Zea mays*)

Maize is one of the most important food crop grown in an area of 13.37 lakh ha in almost all the districts of the State. The crop requirements for growing maize (Table 7.3) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing maize was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.2.

No highly suitable (Class S1) lands available for growing maize in the microwatershed. Moderately suitable (Class S2) lands occur in a maximum area of 338 ha (56%) and are distributed in the major part of the microwatershed with minor limitations of texture, drainage and calcareousness. Marginally suitable lands (Class S3) for growing maize occupy an area of 71 ha (12%) and occur in all parts of the microwatershed. They have moderate limitations of rooting depth, gravelliness, calcareousness and texture.






Fig. 7.2 Land Suitability map of Maize

### 7.3 Land Suitability for Bajra (*Pennisetum glaucum*)

Bajra is one of the most important millet crop grown in an area of 2.34 lakh ha in the northern districts of Karnataka state. The crop requirements for growing bajra (Table 7.4) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing bajra was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.3.

No Highly (Class S1) suitable lands available for growing bajra in the microwatershed. Maximum area of about 408 ha (68%) is moderately suitable (Class S2) for growing bajra and are distributed in the major part of the microwatershed. They have minor limitations of texture, gravelliness, drainage and calcareousness. Marginally suitable lands (Class S3) occupy very small area of 3 ha (<1%) and are distributed in the southern part of the microwatershed. They have moderate limitations of rooting depth and texture.




Fig. 7.3 Land Suitability map of Bajra

#### 7.4 Land Suitability for Groundnut (*Arachis hypogaea*)

Groundnut is one of the major oilseed crop grown in an area of 6.54 lakh ha in Karnataka in most of the districts either as rainfed or irrigated crop. The crop requirements for growing groundnut (Table 7.5) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing groundnut was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.4.

There are no highly suitable (Class S1) lands available for growing groundnut in the microwatershed. Moderately suitable (Class S2) lands occur in an area of 158 ha (26%) and are distributed in all parts of the microwatershed. They have minor limitations of gravelliness, calcareousness and texture. Marginally suitable lands (Class S3) for growing groundnut occupy a maximum area of about 252 ha (42%) with moderate limitations of texture, drainage and rooting depth.






Fig. 7.4 Land Suitability map of Groundnut

### 7.5 Land Suitability for Sunflower (*Helianthus annuus*)

Sunflower is one of the most important oilseed crop grown in an area of 4.1 lakh ha in the State in all the districts. The crop requirements for growing sunflower (Table 7.6) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing sunflower was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.5.

Highly suitable (Class S1) lands for growing sunflower occupy an area of 132 ha (22%) and are distributed in the northern, central, western and southern part of the microwatershed. Maximum area of about 211 ha (35%) is moderately suitable (Class S2) for sunflower and are distributed in the major part of the microwatershed. They have minor limitations of rooting depth, calcareousness and drainage. An area of about 63 ha (11%) is marginally suitable (Class S3) and is distributed in all parts of the microwatershed with moderate limitations of gravelliness, calcareousness and texture. Currently not suitable (Class N1) lands occur in a very small area of 3 ha (<1%) and is distributed in the southern part of the microwatershed with severe limitation of rooting depth.




Fig. 7.5 Land Suitability map of Sunflower

## 7.6 Land suitability criteria for Red gram (*Cajanus Cajan*)

Redgram is one of the most important pulse crop grown in an area of 7.28 lakh ha in almost all the districts of the State. The crop requirements for growing red gram (Table 7.7) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing redgram was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.6.

No highly suitable (Class S1) lands are available for growing redgram in the microwatershed. Maximum area of about 401 ha (67%) is moderately suitable (Class S2) for growing redgram and are distributed in the major part of the microwatershed. They have minor limitations of rooting depth, texture, drainage and calcareousness. Marginally suitable lands (Class S3) for growing redgram occupy an area of about 8 ha (1%) and occur in the southern and northern part of the microwatershed. They have moderate limitations of rooting depth, gravelliness and texture.




Fig. 7.6 Land Suitability map of Redgram

### 7.7 Land Suitability for Bengal gram (*Cicer aerativum*)

Bengal gram is one of the most important pulse crop grown in about 9.39 lakh ha area in Bijapur, Raichur, Kalaburgi, Dharwad, Belgaum and Bellary districts. The crop requirements for growing Bengal gram (Table 7.8) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing Bengal gram was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.7.

Highly (Class S1) suitable lands for growing Bengal gram occupy a maximum area of 249 ha (41%) and are distributed in the major part of the microwatershed. An area of about 18 ha (3%) is moderately suitable (Class S2) for growing Bengal gram and are distributed in the southern part of the microwatershed. They have minor limitations of calcareousness and texture. Marginally suitable lands (Class S3) occupy an area of about 84 ha (14%) and are distributed in the central, southern and northern part of the microwatershed. They have moderate limitations of rooting depth, gravelliness, texture and calcareousness. Currently not suitable (Class N1) lands occur in an area of 58 ha (10%) and are distributed in the central, southwestern and northern part of the microwatershed with severe limitations of rooting depth and texture.




Fig. 7.7 Land Suitability map of Bengal gram

### 7.8 Land Suitability for Cotton (*Gossypium hirsutum*)

Cotton is one of the most important fibre crop grown in the State in about 8.75 lakh ha area in Raichur, Dharwad, Belgaum, Kalaburgi, Bijapur, Bidar, Bellary, Chitradurga and Chamarajnar district. The crop requirements for growing cotton (Table 7.9) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing cotton was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.8.

Highly suitable (Class S1) lands for growing cotton occur in an area of 132 ha (22%) and are distributed in the northern, western and southern part of the microwatershed. Moderately suitable (Class S2) lands are found to occur in a maximum area of about 135 ha (22%). These soils have minor limitations of rooting depth and calcareousness. They are distributed in the major part of the microwatershed. Marginally suitable (Class S3) lands for cotton occur in an area of 84 ha (14%) with moderate limitations of rooting depth, gravelliness, texture and calcareousness and are distributed in the central and northern part the microwatershed. Currently not suitable (Class N1) lands occur in an area of 58 ha (10%) and are distributed in the central, northern and western part of the microwatershed with severe limitations of calcareousness and texture.






Fig. 7.8 Land Suitability map of Cotton

### 7.9 Land Suitability for Chilli (*Capsicum annuum*)

Chilli is one of the most important vegetable and spice crop grown in about 0.42 lakh ha in Karnataka State. The crop requirements for growing chilli (Table 7.10) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing chilli was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.9.

There are no highly (Class S1) suitable lands available for growing chilli crop in the microwatershed. Maximum area of about 402 ha (67%) is moderately suitable (Class S2) for growing chilli and are distributed in the major part of the microwatershed. They have minor limitations of texture, drainage and calcareousness. Marginally suitable lands (Class S3) occupy an area of 8 ha (1%) and are distributed in the northern and southern part of the microwatershed. They have moderate limitations of rooting depth, gravelliness and texture.




Fig 7.9 Land Suitability map of Chilli

### 7.10 Land Suitability for Tomato (*Lycopersicon esculentum*)

Tomato is one of the most important vegetable crop grown in about 0.61 lakh ha covering almost all the district of the state. The crop requirements for growing tomato (Table 7.11) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing tomato was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.10.

No highly suitable (Class S1) lands available for growing tomato in the microwatershed. Maximum area of 269 ha (45%) is moderately suitable (Class S2) and is distributed in the major part of the microwatershed with minor limitations of texture, drainage and calcareousness. An area of 141 ha (23%) is marginally suitable for tomato (Class S3) and is distributed in the northern, western, central and southern part of the microwatershed. They have moderate limitations of rooting depth, gravelliness, drainage and texture.






Fig 7.10 Land Suitability map of Tomato

### 7.11 Land Suitability for Brinjal (*Solanum melongena*)

Brinjal is one of the most important vegetable crop grown in the state. The crop requirements for growing brinjal (Table 7.12) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing brinjal was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.11.

Highly (Class S1) suitable lands for growing brinjal occur in an area of 151 ha (25%) and are distributed in the western, central and southern part of the microwatershed. Maximum area of about 255 ha (42%) is moderately suitable (Class S2) for brinjal and is distributed in the major part of the microwatershed. They have minor limitations of texture, calcareousness and gravelliness. A very small area of 3 ha (<1%) is marginally suitable and is distributed in the southern part of the microwatershed with moderate limitation of rooting depth.




Fig 7.11 Land Suitability map of Brinjal

## 7.12 Land Suitability for Onion (*Allium cepa* L.,)

Onion is one of the most important vegetable crop grown in the state. The crop requirements for growing onion (Table 7.13) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing onion was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.12.

Highly (Class S1) suitable lands for growing onion occur in a maximum area of 281 ha (47%) and are distributed in the major part of the microwatershed. Small area of about 10 ha (2%) is moderately suitable (Class S2) for onion and is distributed in the northern part of the microwatershed. They have minor limitations of gravelliness, texture and calcareousness. An area of 119 ha (20%) is marginally suitable and is distributed in all parts of the microwatershed with moderate limitations of rooting depth and texture.




Fig 7.12 Land Suitability map of Onion

### 7.13 Land Suitability for Bhendi (*Abelmoschus esculentus*)

Bhendi is one of the most important vegetable crop grown in the state. The crop requirements for growing bhendi (Table 7.14) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing bhendi was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.13.

Highly (Class S1) suitable lands for growing bhendi occur in a maximum area of 281 ha (47%) and are distributed in the major part of the microwatershed. Small area of about 10 ha (2%) is moderately suitable (Class S2) for bhendi and is distributed in the northern part of the microwatershed. They have minor limitations of texture, calcareousness and gravelliness. An area of 119 ha (20%) is marginally suitable and is distributed in all parts of the microwatershed with moderate limitations of rooting depth, and texture.




Fig 7.13 Land Suitability map of Bhendi

#### 7.14 Land Suitability for Drumstick (*Moringa oleifera*)

Drumstick is one of the most important vegetable crop grown in about 2403 ha in the state. The crop requirements for growing drumstick (Table 7.15) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing drumstick was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.14.

There are no highly (Class S1) suitable lands available for growing drumstick in the microwatershed. Maximum area of about 406 ha (68%) is moderately suitable (Class S2) for drumstick and is distributed in the major part of the microwatershed. They have minor limitations of rooting depth, texture, gravelliness, calcareousness and drainage. Currently not suitable (Class N1) lands for growing drumstick occur in a very small area of 3 ha (<1%) and are distributed in the southern part of the microwatershed. They have severe limitations of rooting depth and texture.






Fig 7.14 Land Suitability map of Drumstick

### 7.15 Land suitability for Mango (*Mangifera indica*)

Mango is one of the most important fruit crop grown in an area of 1.73 lakh ha in almost all the districts of the State. The crop requirements (Table 7.16) for growing mango were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing mango was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed is given in Figure 7.15.

There are no highly suitable (Class S1) lands available for growing mango in the microwatershed. Moderately suitable (Class S2) lands occur in an area of 16 ha (3%) and are distributed in the southern part of the microwatershed with minor limitation of rooting depth. Maximum area of 391 ha (65%) is marginally suitable (Class S3) for growing mango with moderate limitations of texture, gravelliness, calcareousness and rooting depth and are distributed in the major part of the microwatershed. Very small area of about 3 ha (<1%) is currently not suitable (Class N1) for growing mango and are distributed in the southern part of the microwatershed with severe limitation of rooting depth.




Fig. 7.15 Land Suitability map of Mango

### 7.16 Land suitability for Guava (*Psidium guajava*)

Guava is one of the most important fruit crop grown in an area of 6558 ha in almost all the districts of the State. The crop requirements (Table 7.17) for growing guava were matched with the soil-site characteristics (7.1) and a land suitability map for growing guava was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.16.

No highly (Class S1) suitable lands available for growing guava in the microwatershed. Moderately suitable (Class S2) lands occur in an area of 152 ha (25%) and are distributed in all parts of the microwatershed with minor limitations of rooting depth, texture and calcareousness. Marginally suitable (Class S3) lands cover a maximum area of about 254 ha (42%) and are distributed in the major part of the microwatershed. They have moderate limitations of texture and gravelliness. Very small area of about 3 ha (<1%) is currently not suitable (N) for growing guava and occur in the southern part of the microwatershed with severe limitations of rooting depth and texture.






Fig. 7.16 Land Suitability map of Guava

### 7.17 Land suitability for Sapota (*Manilkara zapota*)

Sapota is one of the most important fruit crop grown in an area of 29373 ha in almost all the districts of the State. The crop requirements (Table 7.18) for growing sapota were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing sapota was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.17.

No highly (Class S1) suitable lands available for growing sapota in the microwatershed. Moderately suitable (Class S2) lands occur in an area of 152 ha (25%) and are distributed in all parts of the microwatershed with minor limitations of rooting depth, texture and calcareousness. Maximum area of about 254 ha (42%) is marginally suitable (Class S3) for growing sapota and are distributed in the major part of the microwatershed. They have moderate limitations of texture and gravelliness. Very small area of 3 ha (<1%) is currently not suitable (Class N1) for growing sapota and occur in the southern part of the microwatershed with severe limitation of rooting depth.




Fig. 7.17 Land Suitability map of Sapota

### 7.18 Land Suitability for Pomegranate (*Punica granatum*)

Pomegranate is one of the most important fruit crop commercially grown in about 18488 ha in Karnataka, mainly in Bijapur, Bagalkot, Koppal, Gadag and Chitradurga districts. The crop requirements for growing pomegranate (Table 7.19) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing pomegranate was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.18.

No highly (Class S1) suitable lands available for growing pomegranate in the microwatershed. Maximum area of about 401 ha (67%) is moderately suitable (Class S2) for growing pomegranate and is distributed in the major part of the microwatershed. They have minor limitations of rooting depth, texture, calcareousness and gravelliness. Small area of 5 ha (1%) is marginally suitable for pomegranate and is distributed in the northern part of the microwatershed with moderate limitation of gravelliness. Very small area of about 3 ha (<1%) is currently not suitable (Class N1) for growing pomegranate and is distributed in the southern part of the microwatershed with severe limitation of rooting depth.




Fig 7.18 Land Suitability map of Pomegranate

### 7.19 Land Suitability for Musambi (*Citrus limetta*)

Musambi is one of the important fruit crop grown in an area of 3446 ha in almost all the districts of the State. The crop requirements for growing musambi (Table 7.20) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing musambi was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.19.

Highly suitable (Class S1) lands for growing Musambi occur in an area of 174 ha (29%) and are distributed in all parts of the microwatershed. Maximum area of about 227 ha (38%) is moderately suitable (Class S2) for growing Musambi and are distributed in the major part of the microwatershed. They have minor limitations of calcareousness and rooting depth. Small area of about 5 ha (1%) is marginally suitable and is distributed in the northern part of the microwatershed with moderate limitation of gravelliness. Currently not suitable (Class N1) lands occur in a very small area of 3 ha (<1%) and are distributed in the southern part of the microwatershed with severe limitation of rooting depth.




Fig. 7.19 Land Suitability map of Musambi

## 7.20 Land Suitability for Lime (*Citrus sp*)

Lime is one of the most important fruit crop grown in an area of 0.11 lakh ha in almost all the districts of the State. The crop requirements for growing lime (Table 7.21) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing lime was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7. 20.

Highly suitable (Class S1) lands for growing Lime occur in an area of 174 ha (29%) and are distributed in all parts of the microwatershed. Maximum area of about 227 ha (38%) is moderately suitable (Class S2) for growing lime and are distributed in the major part of the microwatershed. They have minor limitations of calcareousness and rooting depth. Small area of about 5 ha (1%) is marginally suitable and is distributed in the northern part of the microwatershed with moderate limitation of gravelliness. Currently not suitable (Class N1) lands occur in a very small area of 3 ha (<1%) and are distributed in the southern part of the microwatershed with severe limitation of rooting depth.






Fig. 7.20 Land Suitability map of Lime

### 7.21 Land Suitability for Amla (*Phyllanthus emblica*)

Amla is one of the medicinal fruit crop grown in almost all the districts of the State. The crop requirements for growing amla (Table 7.22) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing amla was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.21.

Highly suitable (Class S1) lands for growing Amla occur in an area of 18 ha (3%) and are distributed in the southern part of the microwatershed. Maximum area of about 330 ha (55%) has soils that are moderately suitable (Class S2) for growing Amla with minor limitations of gravelliness, texture and calcareousness and are distributed in the major part of the microwatershed. An area of 61 ha (10%) is marginally suitable (Class S3) for growing amla with moderate limitations of rooting depth, texture and gravelliness and is distributed in all parts of the microwatershed.




Fig. 7.21 Land Suitability map of Amla

## 7.22 Land Suitability for Cashew (*Anacardium occidentale*)

Cashew is one of the most important plantation nut crop grown in an area of 0.7 lakh ha in almost all the districts of the State. The crop requirements for growing cashew (Table 7.23) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing cashew was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.22.

An area of 18 ha (3%) is moderately suitable (Class S2) for cashew and are distributed in the southern part of the microwatershed with minor limitations of rooting depth and texture. About 63 ha (11%) area is marginally suitable for cashew and is distributed in all parts of the microwatershed with moderate limitations of gravelliness, calcareousness and texture. Maximum area of 328 ha (54%) is currently not suitable for cashew and is distributed in the major part of the microwatershed with severe limitations of rooting depth, texture and calcareousness.






Fig. 7.22 Land Suitability map of Cashew

### 7.23 Land Suitability for Jackfruit (*Artocarpus heterophyllus*)

Jackfruit is one of the most important fruit crop grown in an area of 5368 ha in almost all the districts of the State. The crop requirements for growing jackfruit (Table 7.24) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing jackfruit was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.23.

No highly suitable (Class S1) lands available for growing Jackfruit in the microwatershed. Moderately suitable (Class S2) lands occur in an area of 94 ha (16%) and are distributed in the northern, central and southern part of the microwatershed with minor limitations of rooting depth and calcareousness. Marginally suitable (Class S3) lands occupy a maximum area of about 312 ha (52%) and are distributed in the major part of the microwatershed. They have moderate limitations of texture, gravelliness and calcareousness. Very small area of about 3 ha (<1%) is currently not suitable (Class N1) and is distributed in the southern part of the microwatershed with severe limitations of rooting depth and texture.




Fig. 7.23 Land Suitability map of Jackfruit

## 7.24 Land Suitability for Jamun (*Syzygium cumini*)

Jamun is an important fruit crop grown in almost all the districts of the State. The crop requirements for growing jamun (Table 25) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing jamun was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.24.

No highly suitable (Class S1) lands available for growing Jamun in the microwatershed. Maximum area of about 307 ha (51%) is moderately suitable (Class S2) for growing Jamun and are distributed in the major part of the microwatershed. They have minor limitations of texture, rooting depth and calcareousness. An area of about 99 ha (17%) is marginally suitable (Class S3) for growing Jamun and is distributed in all parts of the microwatershed. They have moderate limitations of gravelliness, calcareousness and rooting depth. Very small area of about 3 ha (<1%) is currently not suitable (Class N1) and is distributed in the southern part of the microwatershed with severe limitations of rooting depth and texture.




Fig. 7.24 Land Suitability map of Jamun

### 7.25 Land Suitability for Custard Apple (*Annona reticulata*)

Custard apple is one of the most important fruit crop grown in almost all the districts of the State. The crop requirements for growing custard apple (Table 7.26) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing custard apple was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.25.

Highly suitable (Class S1) lands for growing custard apple occur in maximum area of 322 ha (54%) and is distributed in the major part of the microwatershed. An area of about 26 ha (4%) is moderately suitable (Class S2) for growing custard apple and is distributed in the northern and southern part of the microwatershed with minor limitations of calcareousness and gravelliness. Marginally suitable (Class S3) lands occur in an area of 61 ha (10%) and are distributed in all parts of the microwatershed with moderate limitations of texture, rooting depth and calcareousness.




Fig. 7.25 Land Suitability map of Custard Apple

### 7.26 Land Suitability for Tamarind (*Tamarindus indica*)

Tamarind is one of the most important spice crop grown in almost all the districts of the state. The crop requirements for growing tamarind (Table 7.27) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing tamarind was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.26.

No highly suitable (Class S1) lands available for growing Tamarind in the microwatershed. Maximum area of about 307 ha (51%) is moderately suitable (Class S2) for growing Tamarind and are distributed in the major part of the microwatershed. They have minor limitations of texture and rooting depth. Marginally suitable (Class S3) lands for growing Tamarind occupy an area of about 99 ha (17%) and are distributed in the northern, central and southern part of the microwatershed. They have moderate limitations of rooting depth, gravelliness and calcareousness. Very small area of about 3 ha (<1%) is currently not suitable (Class N1) for growing Tamarind and occur in the southern part of the microwatershed with severe limitations of rooting depth and texture.






Fig. 7.26 Land Suitability map of Tamarind

### 7.27 Land Suitability for Mulberry (*Morus nigra*)

Mulberry is one of the important leaf crop grown for rearing silk worms in about 1.6 lakh ha area in all the districts of the state. The crop requirements for growing mulberry (Table 7.28) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing mulberry was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.27.

No highly (Class S1) suitable lands available for growing mulberry in the microwatershed. An area of about 99 ha (17%) is moderately (Class S2) suitable for growing mulberry and are distributed in the northern, central and southern part of the microwatershed. They have moderate limitations of gravelliness, rooting depth and calcareousness. Marginally suitable (Class S3) lands occur in a maximum area of 307 ha (51%) and are distributed in the major part of the microwatershed with moderate limitations of texture, calcareousness and drainage. Currently not suitable lands (Class N1) occupy an area of about 3 ha (<1%) and distributed in the southern part of the microwatershed. They have severe limitations of rooting depth and texture.






Fig 7.27 Land Suitability map of Mulberry

### 7.28 Land suitability for Marigold (*Tagetes sps.*)

Marigold is one of the most important flower crop grown in an area of 9108 ha in almost all the districts of the State. The crop requirements (Table 7.29) for growing marigold were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing marigold was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.28.

No highly suitable (Class S1) lands available for growing Marigold in the microwatershed. Maximum area of about 401 ha (67%) is moderately suitable (Class S2) for growing Marigold and are distributed in the major part of the microwatershed. They have minor limitations of texture, drainage and calcareousness. Marginally suitable (Class S3) lands occupy an area of about 8 ha (1%) and are distributed in the northern and southern part of the microwatershed. They have moderate limitations of rooting depth, gravelliness and texture.




Fig. 7.28 Land Suitability map of Marigold

### 7.29 Land Suitability for Chrysanthemum (*Dendranthema grandiflora*)

Chrysanthemum is one of the most important flower crop grown in an area of 4978 ha in almost all the districts of the State. The crop requirements (Table 7.30) for growing chrysanthemum were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing chrysanthemum was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.29.

No highly suitable (Class S1) lands available for growing Chrysanthemum in the microwatershed. Maximum area of about 401 ha (67%) is moderately suitable (Class S2) for growing Chrysanthemum and are distributed in the major part of the microwatershed. They have minor limitations of texture, calcareousness and drainage. Marginally suitable (Class S3) lands for growing Chrysanthemum occupy an area of about 8 ha (1%) and are distributed in the northern and southern part of the microwatershed. They have moderate limitations of rooting depth, gravelliness and texture.




Fig. 7.29 Land Suitability map of Chrysanthemum

**Table 7.1 Soil-Site Characteristics of Belagiri-1Microwatershed**

Soil Map Units	Climate (P) (mm)	Growing period (Days)	Drain-age Class	Soil depth (cm)	Soil texture		Gravelliness		AWC (mm/m)	Slope (%)	Erosion	pH	EC (dSm <sup>-1</sup> )	ESP (%)	CEC [Cmol (p <sup>+</sup> )kg <sup>-1</sup> ]	BS (%)
					Sur-face	Sub-surface	Surface (%)	Sub-surface (%)								
BDLiB3	866	150	WD	25-50	sc	sl	<15	<15	<50	1-3	severe	6.20	0.07	0.20	4.20	90
BLCcB2	866	150	WD	75-100	sl	scl	<15	<15	51-100	1-3	moderate	6.75	0.19	1.31	16.80	95
BLCiB2	866	150	WD	75-100	sc	scl	<15	<15	51-100	1-3	moderate	6.75	0.19	1.31	16.80	95
KBDhB2	866	150	WD	75-100	scl	scl	<15	35-60	<50	1-3	moderate	7.84	0.60	4.27	11.50	100
KBDcC2g1	866	150	WD	75-100	sl	scl	15-35	35-60	<50	3-5	moderate	7.84	0.60	4.27	11.50	100
HSLiB2	866	150	MWD	75-100	sc	sc	<15	<15	101-150	1-3	moderate	7.16	0.11	5.94	4.90	97
HSLhB2	866	150	MWD	75-100	scl	sc	<15	<15	101-150	1-3	moderate	7.16	0.11	5.94	4.90	97
YDRcB2	866	150	WD	100-150	sl	sl	<15	<15	51-100	1-3	moderate	7.25	0.11	0.31	3.40	96
YDRiB2	866	150	WD	100-150	sc	sl	<15	<15	51-100	1-3	moderate	7.25	0.11	0.31	3.40	96
ANRcA1	866	150	MWD	100-150	sl	c	<15	<15	>200	0-1	slight	10.17	0.36	7.08	19.90	100
MDGhB2g1	866	150	WD	100-150	scl	scl	15-35	<15	>200	1-3	moderate	8.2	0.40	3.08	4.90	100
MDRcB2	866	150	WD	>150	sl	scl	<15	<15	>200	1-3	moderate	8.31	0.33	0.90	20.57	100
MDRiA1	866	150	WD	>150	sc	scl	<15	<15	>200	0-1	slight	8.31	0.33	0.90	20.57	100
MDRhB2	866	150	WD	>150	scl	scl	<15	<15	>200	1-3	moderate	8.31	0.33	0.90	20.57	100
BMNmB2	866	150	MWD	>150	c	c	<15	<15	>200	1-3	moderate	8.20	0.28	0.65	52.70	100

\*Symbols and abbreviations are according to Field Guide for LRI under Sujala-III Project, Karnataka

**Table 7.2 Land suitability criteria for Sorghum**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	26–30	30–34; 24–26	34–40; 20–24	>40; <20
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, c (red), c (black)	scl, cl	ls, sl	-
	pH	1:2.5	5.5-7.8	5.0-5.5 7.8-9.0	>9.0	-
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	10-15
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	0-3	3-5	5-10	>10



**Table 7.3 Land suitability criteria for Maize**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	30-34	35-38 26-30	38-40 26-20	
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc	c (red), c (black)	ls, sl	-
	pH	1:2.5	5.5-7.8	5.0-5.5 7.8-9.0	>9.0	-
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	-
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

**Table 7.4 Land suitability criteria for Bajra**

Land use requirement		Rating				
Soil –site characteristics	Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)	
Climatic regime	Mean temperature in growing season	°C	28-32	33-38 24-27	39-40 20-23	<20
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm	500-750	400-500	200-400	<200
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl, scl, cl,sc,c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.8	5.0-5.5 7.8-9.0	5.5-6.0 >9.0	
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	15-35	35-60	>60	
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	1-3	3-5	5-10	>10

**Table 7.5 Land suitability criteria for Groundnut**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	24–33	22–24; 33–35	20–22; 35–40	<20; >40
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. Well drained	Poorly drained	Very Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl	sl,cl, sc	c (red), c (black), ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<35	35-60	>60	
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.6 Land suitability criteria for Sunflower**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	24–30	30–34; 20–24	34–38; 16–20	>38; <16
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	mod. Well drained	-	Poorly to very drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	cl, sc,c (red), c (black)	scl	ls, sl	-
	pH	1:2.5	6.5-7.8	7.8-8.4 5.5-6.5	8.4-9.0; 5.0-5.5	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.7 Land suitability criteria for Red gram**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	30-35(G) 20-25(AV) 15-18 (F&PS) 35-40(M)	25-30(G) 20-25 (AV) 12-15 (F&PS) 30-35(M)	20-25(G) 15-20(AV) 10-12 (F&PS) 25-30(M)	< 20 <15 <10 <25
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. Well drained	Poorly drained	Very Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, c (red)	c (black),sl, scl, cl	ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-9.0	5.0-5.5 >9.0	-
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-50	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<1.0	1.0-2.0	>2.0	
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	<3	3-5	5-10	>10



**Table 7.8 Land suitability criteria for Bengal gram**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	20–25	25–30; 15–20	30–35; 10–15	>35; <10
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. Well drained	Poorly drained	Very Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	c(black)	-	c (red), scl, cl, sc	ls, sl
	pH	1:2.5	6.0-7.8	5.0-6.0 7.8-9.0	>9.0	-
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	-
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.9 Land suitability criteria for Cotton**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	22-32	>32	<19	-
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well to moderately well	Poorly drained/Some what excessively drained	-	very poorly/excessively drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, c (red,black)	cl	scl	ls, sl
	pH	1:2.5	6.5-7.8	7.8-8.4	5.5-6.5 8.4->9.0	<5.5
	CEC	C mol (p+)Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	50-100	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	<3	3-5	-	>5

**Table 7.10 Land suitability criteria for Chilli**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	25-32	33-35 20-25	35-38 <20	>38
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc	c (black), sl	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.11 Land suitability criteria for Tomato**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	25-28	29-32 20-24	15-19 33-36	<15 >36
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl, scl, cl, sc, c (red)	-	ls, c(black)	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.12 Land suitability criteria for Brinjal**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	Well drained	Moderately well drained	Poorly drained	V. Poorly drained
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class				
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl, scl, cl, sc c (red)	-	ls, c (black)	-
	pH	1:2.5	6.0-7.3	7.3-8.4 5.0-6.0	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	>60
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10



**Table 7.13 Land suitability criteria for Onion**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	20-30	30-35	35-40	>40
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately /imperfectly	-	Poorly to V poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl,scl,cl,sc,c (red)	-	c (Black),ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<1.0	1.0-2.0	2.0-4.0	<4
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.14 Land suitability criteria for Bhendi**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	25-28	29-32 20-24	15-19 33-36	<15 >36
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Imperfectly drained	Poorly to very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl,sc, c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.15 Land suitability criteria for Drumstick**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, scl, cl, c (red)	sl, c (black)	ls	s
	pH	1:2.5	6.0-7.3	5.0-5.5 7.3-7.8	5.5-6.0 7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<35	35-60	60-80	>80
Soil toxicity	Salinity (EC saturation extract)	ds/m				
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-10	-	>10

**Table 7.16 Land suitability criteria for Mango**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-32	24-27 33-35	36-40	20-24
	Min temp. before flowering	°C	10-15	15-22	>22	-
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration	Days				
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V. Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	-	ls, sl, c (black)	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>150	100-150	75-100	<75
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.17 Land suitability criteria for Guava**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-32	33-36 24-27	37-42 20-23	
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	sl	c (black), ls	-
	pH	1:2.5	6.0-7.8	5.0-6.0	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.18 Land suitability criteria for Sapota**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-32	33-36 24-27	37-42 20-23	>42 <18
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	-	Poorly to very drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	sl	ls, c (black)	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10



**Table 7.19 Land suitability criteria for Pomegranate**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	30-34	35-38 25-29	39-40 15-24	
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl,cl, sc, c (red)	c (black),sl	ls	-
	pH	1:2.5	5.5-7.8	7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.20 Land suitability criteria for Musambi**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-30	31-35 24-27	36-40 20-23	>40 <20
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately drained	poorly	Very poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c	sl	ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.21 Land suitability criteria for Lime**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-30	31-35 24-27	36-40 20-23	>40 <20
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately drained	poorly	Very poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c	sl	ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.22 Land suitability criteria for Amla**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. well drained	Poorly drained	V. Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	c (black)	ls, sl	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15-35	35-60	60-80	-
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

**Table 7.23 Land suitability criteria for Cashew**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	32 to 34	28 to 32; 34 to 38	24 to 28; 38 to 40	<20; >40
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	-	sl, ls	c (black)
	pH	1:2.5	5.5-6.5	5.0-5.5 6.5-7.3	7.3-7.8	>7.8
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-10	>10	-

**Table 7.24 Land suitability criteria for Jackfruit**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. well	Poorly	V. Poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	-	sl, ls, c (black)	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	>60
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10-



**Table 7.25 Land suitability criteria for Jamun**

Land use requirement		Rating				
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well	Mod. well	Poorly	V.Poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c(red)	sl, c (black)	ls	-
	pH	1:2.5	6.0-7.8	5.0-6.0	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>150	100-150	50-100	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	>60
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

**Table 7.26 Land suitability criteria for Custard apple**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	Scl, cl, sc, c (red), c (black)	-	Sl, ls	-
	pH	1:2.5	6.0-7.3	5.5-6.0 7.3-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15-35	35-60	60-80	-
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	>5	-

**Table 7.27 Land suitability criteria for Tamarind**

Land use requirement		Rating				
Soil –site characteristics	Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)	
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod.well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl,sc, c (red)	sl, c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>150	100-150	75-100	<75
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

**Table 7.28 Land suitability criteria for Mulberry**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	24–28	22–24; 28–32	32–38; 22–18	>38; <18
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V. Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, cl, scl	c (red)	c (black), sl, ls	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.8-8.4	7.3-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	0-35	35-60	60-80	>80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

**Table 7.29 Land suitability criteria for Marigold**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	18-23	17-15 24-35	35-40 10-14	>40 <10
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl,scl, cl, sc, c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%				
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.30 Land suitability criteria for Chrysanthemum**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	18-23	17-15 24-35	35-40 10-14	>40 <10
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl,scl, cl, sc, c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%				
Erosion hazard	Slope	%	<3	3-5	5-10	>10



### 7.30 Land Management Units (LMUs)

The 15 soil map units identified in Belagiri-1microwatershed have been grouped into 5 Land Management Units (LMU's) for the purpose of preparing a Proposed Crop Plan. Land Management Units are grouped based on the similarities in respect of the type of soil, the depth of the soil, the surface soil texture, gravel content, AWC, slope, erosion etc. and a Land Management Units map (Fig. 7.30) has been generated. These Land Management Units are expected to behave similarly for a given level of management.

The 15 map units that have been grouped into 5 Land Management Units along with brief description of soil and site characteristics are given below.

LMU NO.	Soil map units	Soil and site characteristics
1	59.MDRcB2 60.MDRiA1 132.MDRhB2 149.MDGhB2g1 42.YDRcB2 43.YDRiB2	Deep to very deep, black sandy clay loam to sandy loam, 0-3% slopes, non gravelly to gravelly, slight to moderate erosion.
2	130.KBDhB2 164.KBDcC2g1	Moderately deep, red gravelly loamy, 1-5% slopes, non gravelly to gravelly, moderate erosion.
3	33.HSLiB2 62.BMNmB2 126.HSLhB2 167.ANRcA1	Moderately deep to very deep, black calcareous clay, 0-3% slopes, non gravelly, slight to moderate erosion.
4	37.BLCcB2 38.BLCiB2	Moderately deep, red loam, 1-3%, slopes, non gravelly, moderate erosion
5	6.BDLiB3	Shallow, sandy loam, 1-3% slopes, non gravelly severe erosion.




Fig. 7.30 Land Management Units Map- Belagiri-1 Microwatershed

### 7.31 Proposed Crop Plan for Belagiri-1 Microwatershed

After assessing the land suitability for the 29 crops, the Proposed Crop Plan has been prepared for the 5 identified LMUs by considering only the highly (Class S1) and moderately (Class S2) suitable lands for each of the 29 crops. The resultant proposed crop plan is presented below in Table 7.31.

**Table 7.31 Proposed Crop Plan for Belagiri-1Microwatershed**

LMU No	Mapping Units	Survey Number	Soil Characteristics	Field Crops	Horticulture Crops	Suitable Interventions
1	59.MDRcB2 60.MDRiA1 132.MDRhB2 149.MDGhB2g1 42.YDRcB2 43.YDRiB2	<b>Belagera:</b> 5(1),5(2),6(1),6(2),7(1),7(2),7(3),8(1),8(2),9(1),9(2),10(1),10(2),11(1),11(2),12(1),12(2),13,14,15,16,17,18,19,20,21,22,23,24,25,26,27,28,29,30,31,32,33,34,35,37,38,39,40,41,42,43,44,45,46,47,48,49,50,51,52,53,54,55,56,57,58,59,60,61,62,95,96,97,98,99,100,101(3),101/1,101/2,102(1),102(2),102(3),103(1),125,127,133,134,136,137,138,139,140,154,157,158,159,161(1),164,165,250,251,252,255,256,257,258,259,260,261,289,290,292,293,295,299,300,301,302,303,304,305(1),305(2),305(3),306,307,308,309(1),309(2),310(1),310(3),311,312(1) <b>Honagera:</b> 206,208,209,210,211,226	Deep to very deep, black sandy clay loam to sandy loam, 0-3% slopes, non gravelly to gravelly, slight to moderate erosion.	Sunflower, Sorghum, Maize, Groundnut, Soybean, Safflower, Linseed, Bajra, Mulberry	<b>Fruit crops:</b> Mango, Sapota, Pomegranate, Guava, Lime, Musambi, Jamun, Jackfruit, Tamarind, Amla, Custard apple <b>Vegetables:</b> Onion, Tomato, Bhendi, Drumstick, Chilli, Coriander <b>Flowers:</b> Marigold, Chrysanthemum	Application of FYM, Biofertilizers and micronutrients, drip irrigation, Mulching, suitable soil and water conservation practices
2	130.KBDhB2 164.KBDcC2g1	<b>Sutharahosalli:</b> 93,94	Moderately deep, red gravelly loamy, 1-5% slopes, non gravelly to gravelly, moderate erosion.	Groundnut, Bajra, Horse gram, Castor, Mulberry	<b>Fruit crops:</b> Musambi, Lime, Jamun, Jackfruit Amla, Custard apple, Tamarind <b>Vegetable crops:</b> Drumstick, Curry leaves	Drip irrigation, mulching, suitable soil and water conservation practises (Crescent Bunding with Catch Pit etc)
3	33.HSLiB2 62.BMNmB2	<b>Belagera:</b> 94,103(2),104,105,106,107,108,109,110,112(2),113,1	Moderately deep to very deep,	Sunflower, Sorghum,	<b>Fruit crops:</b> Pomegranate, Lime,	Application of FYM, Biofertilizers

	126.HSLhB2 167.ANRcA1	15,116,117,118,119,120,121,122,123,124(1),124(2),126,128(1),128(2), 129,130,131,132,160 <b>Sutharahosalli:</b> 91,92,95,96,97,98,99	black calcareous clay, 0-3% slopes, non gravelly, slight to moderate erosion.	Maize, Soybean, Cotton, Bengal gram, Safflower, Linseed, Bajra	Musambi, Tamarind, Jamun, Amla, Custard apple <b>Vegetables:</b> Drumstick, Chilli, Bhendi, Cluster bean, Coriander <b>Flowers:</b> Marigold, Chrysanthemum	and micronutrients, drip irrigation, Mulching, suitable soil and water conservation practices
4	37.BLCcB2 38.BLCiB2	<b>Belagera:</b> 141,149,150,151,152,153	Moderately deep, red loam, 1-3%, slopes, non gravelly, moderate erosion	Maize, Sorghum, Sunflower, Bajra, Finger millet, Groundnut, Red gram, Cowpea, Field bean, Castor, Mulberry	<b>Fruit crops:</b> Mango, Pomegranate, Guava, Sapota, Jackfruit, Jamun, Tamarind, Lime, Musambi, Amla, Custard apple, Cashew <b>Vegetable crops:</b> Drumstick, Tomato, Bhendi, Chilli, Brinjal, Onion, Curry leaves <b>Flower crops:</b> Marigold, Chrysanthemum, Jasmine, Crossandra	Drip irrigation, mulching, suitable soil and water conservation practises (Crescent Bunding with Catch Pit etc)
5	6.BDLiB3	<b>Belagera:</b> 148	Shallow, sandy loam, 1-3% slopes, non gravelly, severe erosion.	-	<b>Agri-Silvi-Pasture:</b> Hybrid Napier, <i>Styloxanthes hamata</i> , <i>Styloxanthes scabra</i>	Use of short duration varieties, sowing across the slope, drip irrigation and mulching is recommended

## SOIL HEALTH MANAGEMENT

### 8.1 Soil Health

Soil health is basic to plant health and plant health is basic to human and bovine health. Soil is fundamental to crop production. Without soil, no food could be produced nor would livestock be fed on a large scale. Because it is finite and fragile, soil is a precious resource that requires special care from its users.

Soil health or the capacity of the soil to function is critical to human survival. Soil health has been defined as: “the capacity of the soil to function as a living system without adverse effect on the ecosystem”. Healthy soils maintain a diverse community of soil organisms that help to form beneficial symbiotic associations with plant roots, recycle essential plant nutrients, improve soil structure with positive repercussions for soil, water and nutrient holding capacity and ultimately improve crop production and also contribute to mitigating climate change by maintaining or increasing its carbon content.

Functional interactions of soil biota with organic and inorganic components, air and water determine a soil’s potential to store and release nutrients and water to plants and to promote and sustain plant growth. Thus, maintaining soil health is vital to crop production and conserve soil resource base for sustaining agriculture.

#### The most important characteristics of a healthy soil are

- Good soil tilth
- Sufficient soil depth
- Good water storage and good drainage
- Adequate supply, but not excess of nutrients
- Large population of beneficial organisms
- Small proportion of plant pathogens and insect pests
- Low weed pressure
- Free of chemicals and toxins that may harm the crop
- Resistance to degradation
- Resilience when unfavorable conditions occur

#### Characteristics of Belagiri-1 Microwatershed

- ❖ The soil phases identified in the microwatershed belonged to the soil series of MDR 216 ha (36%), HSL 76 ha (13%), YDR 58 ha (10%), BLC 18 ha (3%), BMN 17 (3%), MDG 16 ha (3%), KBD 5 ha (1%), BDL 3 ha (<1%) and ANR 0.07 ha (<1%).
- ❖ As per land capability classification entire area of the microwatershed falls under arable land category (Class II, III & IV). The major limitations identified in the arable lands were soil and erosion.
- ❖ On the basis of soil reaction, about 104 ha (17%) is neutral, 224 ha (37%) area is slightly to moderately alkaline (pH 7.3-8.4) and 81 ha (14%) is strongly to very strongly alkaline (pH 8.4 - >9.0).

## ❖ Soil Health Management

The following actions are required to improve the current land husbandry practices that provide a sound basis for the successful adoption of sustainable crop production system.

### Acid soils

1. Growing of crops suitable for a particular soil pH.
2. Amelioration of soils through the application of amendments (liming materials).

Liming materials:

1.  $\text{CaCO}_3$  (Calcium Carbonate).
2. Dolomite [ $\text{Ca Mg} (\text{CO}_3)_2$ ]
3. Quick lime (Cao)
4. Slaked lime [ $\text{Ca} (\text{OH})_2$ ]

For normal pH and pH 4.8 (35 t/ha) and pH 6.0-7.0 (4 t/ha) lime is required.

### Alkaline soils

(Slightly alkaline to very strongly alkaline soils cover about 305 ha area in the microwatershed.

1. Regular addition of organic manure, green manuring, green leaf manuring, crop residue incorporation and mulching needs to be taken up to improve the soil organic matter status.
2. Application of biofertilizers (Azospirillum, Azotobacter, Rhizobium).
3. Application of 25% extra N and P (125 % RDN&P).
4. Application of  $\text{ZnSO}_4$  – 12.5 kg/ha (once in three years).
5. Application of Boron – 5kg/ha (once in three years).

### Neutral soils

Neutral soils occur in 104 ha area in the microwatershed.

1. Regular addition of organic manure, green manuring, green leaf manuring, crop residue incorporation and mulching needs to be taken up to improve the soil organic matter status.
2. Application of biofertilizers, (Azospirillum, Azotobacter, Rhizobium).
3. Application of 100 per cent RDF.
4. Need based micronutrient applications.

Besides the above recommendations, the best transfer of technology options are also to be adopted.

### Soil Degradation

Soil erosion is one of the major factor affecting the soil health in the microwatershed. Out of total 602 ha area in the microwatershed, an area of about 67 ha (11%) is suffering from slight erosion, about 341 ha (57%) is suffering from moderate



erosion and 3 ha (<1%) from severe erosion. In areas of moderate and severe erosion immediate soil and water conservation and, other land development and land husbandry practices are required for restoring soil health.

### **Dissemination of Information and Communication of Benefits**

Any large scale implementation of soil health management requires that supporting information is made available widely, particularly through channels familiar to farmers and extension workers. Given the very high priority attached to soil-health especially by the Central Government on issuing Soil-Health Cards to all the farmers, media outlets like Regional, State and National Newspapers, Radio and Dooradarshan programs in local languages but also modern information and communication technologies such as Cellular phones and the Internet, which can be much more effective in reaching the younger farmers.

### **Inputs for Net Planning (Saturation Plan) and Interventions needed**

Net planning (Saturation Plan) in IWMP is focusing on preparation of

1. Soil and Water Conservation Plan for each plot or farm.
2. Productivity enhancement measures/ interventions for existing crops/livestock/other farm enterprises.
3. Diversification of farming mainly with perennial horticultural crops and livestock.
4. Improving livelihood opportunities and income generating activities.

In this connection, how various outputs of Sujala-III are of use in addressing these objectives of Net Planning (Saturation Plan) are briefly presented below.

- ❖ **Soil Depth:** The depth of a soil decides the amount of moisture and nutrients it can hold, what crops can be taken up or not, depending on the rooting depth and the length of growing period available for raising any crop. Deeper the soil, better for a wide variety of crops. If sufficient depth is not available for growing deep rooted crops, either choose medium or short duration crops or deeper planting pits need to be opened and additional good quality soil brought from outside has to be filled into the planting pits.
- ❖ **Surface Soil Texture:** Lighter soil texture in the top soil means, better rain water infiltration, less run-off and soil moisture conservation, less capillary rise and less evaporation losses. Lighter surface textured soils are amenable to good soil tilth and are highly suitable for crops like groundnut, root vegetables (carrot, raddish, potato etc) but not ideal for crops that need stagnant water like lowland paddy. Heavy textured soils are poor in water infiltration and percolation. They are prone for sheet erosion; such soils can be improved by sand mulching. The technology that is developed by the AICRP-Dryland Agriculture, Vijayapura, Karnataka can be adopted.

- ❖ **Gravelliness:** More gravel content is favorable for run-off harvesting but poor in soil moisture storage and nutrient availability. It is a significant parameter that decides the kind of crop to be raised.
- ❖ **Land Capability Classification:** The land capability map shows the areas suitable and not suitable for agriculture and the major constraints in each of the plot/survey number. Hence, one can decide what kind of enterprise is possible in each of these units. In general, erosion and soil are the major constraints in Belagiri-1 microwatershed.
- ❖ **Organic Carbon:** The OC content (an index of available Nitrogen) is high (>0.75%) in an area of 24 ha (4%), medium (0.5-0.75%) in 331 ha (55%) area and low (<0.5%) in 55 ha (9%). The areas that are medium and low in OC needs to be further improved by applying farmyard manure and crop rotation with cereals and legumes or mixed cropping.
- ❖ **Promoting green manuring:** Growing of green manuring crops costs Rs. 1250/ha (green manuring seeds) and about Rs. 2000/ha towards cultivation that totals to Rs. 3250/- per ha. On the other hand, application of organic manure @ 10 tons/ha costs Rs. 5000/ha. The practice needs to be continued for 2-3 years or more. Nitrogen fertilizer needs to be supplemented by 25% in addition to the recommended level in 386 ha area where OC is low and medium (<0.5 - 0.75%). For example, for rainfed maize, recommended level is 50 kg N per ha and an additional 12 kg /ha needs to be applied for all the crops grown in these plots.
- ❖ **Available Phosphorus:** Available Phosphorus is medium (23-57 kg/ha) in 384 ha (64%) of the microwatershed and high in an area of 26 ha (4%). In medium areas, for all the crops 25% additional P needs to be applied.
- ❖ **Available Potassium:** Available potassium is medium (145-337 kg/ha) in an area of 364 ha (60%) of the microwatershed and low (<145 kg/ha) in 46 ha (8%). All the plots, where available potassium is low and medium, for all the crops, additional 25% potassium may be applied.
- ❖ **Available Sulphur:** Available sulphur is a very critical nutrient for oilseed crops. It is medium in 56 ha (9%) and low in 354 ha (59%). Low and medium areas need to be applied with magnesium sulphate or gypsum or Factamphos (p) fertilizer (13% sulphur) for 2-3 years for the deficiency to be corrected.
- ❖ **Available Boron:** Entire area of 410 ha (68%) is low in available boron. Application of sodium tetra borate @ 10 kg/ha as soil application or 0.2 % borax as foliar spray is recommended.
- ❖ **Available Iron:** Entire area of the microwatershed is sufficient (>4.5 ppm) in available iron content.
- ❖ **Available Manganese:** All the soils in the microwatershed are sufficient (>1.0 ppm) in available manganese.

- ❖ **Available Copper:** All the soils in the microwatershed are sufficient ( $>0.2$  ppm) in available copper.
- ❖ **Available Zinc:** Maximum area of about 362 ha (60%) is deficient ( $<0.6$  ppm) in available zinc content. Application of zinc sulphate @25 kg/ha is recommended for these areas. About 48 ha (8%) area is sufficient ( $>0.6$  ppm).
- ❖ **Soil Alkalinity:** Maximum area of 305 ha (51%) in the microwatershed has soils that are slightly to very strongly alkaline. These areas need application of gypsum and wherever calcium is in excess, iron pyrites and element sulphur can be recommended. Management practices like treating repeatedly with good quality water to drain out the excess salts and provision of subsurface drainage and growing of salt tolerant crops like Casuarina, Acacia, Neem, Ber etc, are recommended.


**Land Suitability for various crops:** Areas that are highly, moderately and marginally suitable and not suitable for growing various crops are indicated. Along with the suitability, various constraints that are limiting the productivity are also indicated. For example, in case of cotton, gravel content, rooting depth and salinity/alkalinity are the major constraints in various plots. With suitable management interventions, the productivity can be enhanced. In order to increase the water holding capacity of light textured soils, growing of green manure crops and application of organic manure is recommended.



## SOIL AND WATER CONSERVATION TREATMENT PLAN

For preparing soil and water conservation treatment plan for Belagiri-1 microwatershed, the land resource inventory database generated under Sujala-III project has been transformed as information through series of interpretative (thematic) maps using soil phase map as a base. The various thematic maps (1:7920 scale) generated were

- Soil depth
- Surface soil texture
- Available water capacity
- Soil slope
- Soil gravelliness
- Land capability
- Present land use and land cover
- Crop suitability
- Rainfall
- Hydrology
- Water Resources
- Socio-economic data
- Contour plan with existing features- network of waterways, pottissa boundaries, cut up/ minor terraces etc.
- Cadastral map (1:7920 scale)
- Satellite imagery (1:7920 scale)



Apart from these, Hand Level/ Hydro Marker/ Dumpy Level/ Total Station and Kathedars' List to be collected.

### Steps for Survey and Preparation of Treatment Plan

The boundaries of Land User Groups' and Survey No. boundaries are traced in the field.

- Naming of user groups and farmers
- Identification of arable and non arable lands
- Identification of drainage lines and gullies
- Identification of non treatable areas
- Identification of priority areas in the arable lands
- Treatment plan for arable lands
- Location of water harvesting and recharge structures

## 9.1 Treatment Plan

The treatment plan recommended for arable lands is briefly described below


### 9.1.1 Arable Land Treatment

#### A. BUNDING

Steps for Survey and Preparation of Treatment Plan		<b>USER GROUP-1</b> <b>CLASSIFICATION OF GULLIES</b> 
<ul style="list-style-type: none"> <li>Cadastral map (1:7920 scale) is enlarged to a scale of 1:2500 scale</li> <li>Existing network of waterways, pothissa boundaries, grass belts, natural drainage lines/ watercourse, cut ups/ terraces are marked on the cadastral map to the scale</li> <li>Drainage lines are demarcated into</li> </ul>		
Small gullies	(up to 5 ha catchment)	
Medium gullies	(5-15 ha catchment)	
Ravines	(15-25 ha catchment) and	
<i>Halla/Nala</i>	(more than 25ha catchment)	

#### Measurement of Land Slope

Land slope is estimated or determined by the study and interpretation of contours or by measurement in the field using simple instruments like Hand Level or Hydromarker.



$$\text{FALL: } 1.5 - 0.6 = 0.9 \text{ m.}$$

Vertical and Horizontal intervals between bunds as recommended by the Watershed Development Department.

Slope percentage	Vertical interval (m)	Corresponding Horizontal Distance (m)
2 - 3%	0.6	24
3 - 4%	0.9	21
4 - 5%	0.9	21
5 - 6%	1.2	21
6 - 7%	1.2	21



**Note:** (i) The above intervals are maximum.

(ii) Considering the slope class and erosion status (A1... A=0-1 % slope, 1= slight erosion) the intervals have to be decided.

**Bund length recording:** Considering the contour plan and the existing grass belts/partitions, the bunds are aligned and lengths are measured.

**Section of the Bund**

Bund section is decided considering the soil texture class and gravelliness class (bg<sub>0</sub>... b=loamy sand, g<sub>0</sub> = <15% gravel). The recommended Sections for different soils are given below.

**Recommended Bund Section**

Top width (m)	Base width (m)	Height (m)	Side slope (Z:1;H:V)	Cross section (sq m)	Soil Texture	Remarks
0.3	0.9	0.3	01:01	0.18	Sandy loam	Vegetative bund
0.3	1.2	0.3	1.5:1	0.225	Sandy clay	
0.3	1.2	0.5	0.9:1	0.375	Red gravelly soils	
0.3	1.2	0.6	0.75:1	0.45		
0.3	1.5	0.6	01:01	0.54	Red sandy loam	
0.3	2.1	0.6	1.5:1	0.72	Very shallow black clayey soils	
0.45	2	0.75	01:01	0.92		
0.45	2.4	0.75	1.3:1	1.07	Shallow black clayey soils	
0.6	3.1	0.7	1.78:1	1.29	Medium black clayey soils	
0.5	3	0.85	1.47:1	1.49		

**Formation of Trench cum Bund**

Dimensions of the Borrow Pits/Trenches to be excavated (machinery are decided considering the Bund Section).

Details of Borrow Pit dimensions are given below:

**TRENCH CUM BUND**

**'A' FRAME FOR INTERBUND MANAGEMENT**

### Size of Borrow Pits/ Trench recommended for Trench cum Bund (by machinery)

Bund section	Bund length	Earth quantity	Pit				Berm (pit to pit)	Soil depth class
			L(m)	W(m)	D(m)	Quantity (m <sup>3</sup> )		
m <sup>2</sup>	m	m <sup>3</sup>					m	
0.375	6	2.25	5.85	0.85	0.45	2.24	0.15	Shallow
0.45	6	2.7	5.4	1.2	0.43	2.79	0.6	Shallow
0.45	6	2.7	5	0.85	0.65	2.76	1	Moderately Shallow
0.54	5.6	3.02	5.5	0.85	0.7	3.27	0.1	Moderately shallow
0.54	5.5	2.97	5	1.2	0.5	3	0.5	Shallow
0.72	6.2	4.46	6	1.2	0.7	5.04	0.2	Moderately shallow
0.72	5.2	3.74	5.1	0.85	0.9	3.9	0.1	Moderately deep

#### B. Water Ways

- Existing waterways are marked on the cadastral map (1:7920 scale) and their dimensions are recorded.
- Considering the contour plan of the MWS, additional waterways/ modernization of the existing ones can be thought of.
- The design details are given in the Manual.

#### C. Farm Ponds

Waterways and the catchment area will give an indication on the size of the Farm Pond. Location of the pond can be decided based on the contour plan/ field condition and farmers' need/desire.

#### D. Diversion Channel

Existing EPT/ CPT are marked on the cadastral map. Looking to the need, these can be modernized or fresh diversion channel can be proposed and runoff from this can be stored in *Gokatte*/ Recharge ponds.

#### 9.1.2 Non-Arable Land Treatment

Depending on the gravelliness and crops preferred by the farmers, the concerned authorities can decide appropriate treatment plan. The recommended treatments may be Contour Trench, Staggered Trench, Crescent Bund, Boulder Bund or Pebble Bund.

### **9.1.3 Treatment of Natural Water Course/ Drainage Lines**

- a) The cadastral map has to be updated as regards the network of drainage lines (gullies/*nalas/hallas*) and existing structures are marked to the scale and storage capacity of the existing water bodies are documented.
- b) The drainage line will be demarcated into Upper Reach, Middle Reach and Lower Reach.
- c) Considering the Catchment, *Nala* bed and bank conditions, suitable structures are decided.
- d) Number of storage structures (Check dam/*Nala* bund/ Percolation tank) will be decided considering the commitments and available runoff from water budgeting and quality of water in the wells and site suitability.
- e) Detailed Leveling Survey using Dumpy Level / Total Station has to be carried out to arrive at the site-specific designs as shown in the Manual.
- f) The location of ground water recharge structures are decided by examining the lineaments and fracture zones from geological maps.
- g) Rainfall intensity data of the nearest Rain Gauge Station is considered for Hydrologic Designs.
- h) Silt load to the Storage/Recharge Structures is reduced by providing vegetative, boulder and earthen checks in the natural water course. Location and design details are given in the Manual.

### **9.2 Recommended Soil and Water Conservation Measures**

The appropriate conservation structures best suited for each of the land parcel/ survey number (Appendix-I) are selected based on the slope per cent, severity of erosion, amount of rainfall, land use and soil type. The different kinds of conservation structures recommended are:

1. Graded / Strengthening of Bunds
2. Trench cum Bunds (TCB)
3. Trench cum Bunds / Strengthening
4. Crescent Bunds

A map (Fig. 9.1) showing soil and water conservation plan with different kinds of structures recommended has been prepared which shows the spatial distribution and extent of area. An area of about 24 ha (4%) needs Trench Cum Bunding, maximum area of about 320 ha (53%) needs Graded Bunding and 67 ha (11%) needs strengthening of existing bunds.

The conservation plan prepared may be presented to all the stakeholders including farmers and after considering their suggestions, the conservation plan for the microwatershed may be finalised in a participatory approach.




Fig. 9.1 Soil and Water Conservation Plan map of Belagiri-1 Microwatershed

### 9.3 Greening of Microwatershed

As part of the greening programme in the watersheds, it is envisaged to plant a variety of horticultural and other tree plants that are edible, economical and produce lot of biomass which helps to restore the ecological balance in the watersheds. The lands that are suitable for greening programme are non-arable lands (land capability classes V, VI VII and VIII) and also the lands that are not suitable or marginally suitable and field bunds for growing annual and perennial crops. The method of planting these trees is given below.

It is recommended to open pits during the 1<sup>st</sup> week of March along the contour and heap the dugout soil on the lower side of the slope in order to harness the flowing water and facilitate weathering of soil in the pit. Exposure of soil in the pit also prevents spread of pests and diseases due to scorching sun rays. The pits should be filled with mixture of soil and organic manure during the second week of April and keep ready with sufficiently tall seedlings produced either in poly bags or in root trainer nurseries so that planting can be done during the 2<sup>nd</sup> or 3<sup>rd</sup> week of April depending on the rainfall.

The tree species suitable for the area considering rainfall, temperature and adaptability is listed below; waterlogged areas are recommended to be planted with species like Nerale (*Syzgium cumini*) and Bamboo. Dry areas are to be planted with species like Honge, Bevu, Seetaphal etc.

<b>Dry Deciduous Species</b>			<b>Temp (°C)</b>	<b>Rainfall (mm)</b>
1.	Bevu	<i>Azadiracta indica</i>	21–32	400 –1,200
2.	Tapasi	<i>Holoptelia integrifolia</i>	20-30	500 - 1000
3.	Seetaphal	<i>Anona Squamosa</i>	20-40	400 - 1000
4.	Honge	<i>Pongamia pinnata</i>	20 -50	500– 2,500
5.	Kamara	<i>Hardwickia binata</i>	25 -35	400 - 1000
6.	Bage	<i>Albezzia lebbek</i>	20 - 45	500 - 1000
7.	Ficus	<i>Ficus bengalensis</i>	20 - 50	500–2,500
8.	Sisso	<i>Dalbargia Sissoo</i>	20 - 50	500 -2000
9.	Ailanthus	<i>Ailanthus excelsa</i>	20 - 50	500 - 1000
10.	Hale	<i>Wrightia tinctoria</i>	25 - 45	500 - 1000
11.	Uded	<i>Steriospermum chelanoides</i>	25 - 45	500 -2000
12.	Dhupa	<i>Boswella Serrata</i>	20 - 40	500 - 2000
13.	Nelli	<i>Emblca Officinalis</i>	20 - 50	500 -1500
14.	Honne	<i>Pterocarpus marsupium</i>	20 - 40	500 - 2000
<b>Moist Deciduous Species</b>			<b>Temp (°C)</b>	<b>Rainfall (mm)</b>
15.	Teak	<i>Tectona grandis</i>	20 - 50	500-5000
16.	Nandi	<i>Legarstroemia lanceolata</i>	20 - 40	500 - 4000
17.	Honne	<i>Pterocarpus marsupium</i>	20 - 40	500 - 3000
18.	Mathi	<i>Terminalia alata</i>	20 -50	500 - 2000
19.	Shivane	<i>Gmelina arborea</i>	20 -50	500 -2000
20.	Kindal	<i>T.Paniculata</i>	20 - 40	500 - 1500
21.	Beete	<i>Dalbargia latifolia</i>	20 - 40	500 - 1500
22.	Tare	<i>T. belerica</i>	20 - 40	500 - 2000
23.	Bamboo	<i>Bambusa arundinasia</i>	20 - 40	500 - 2500
24.	Bamboo	<i>Dendrocalamus strictus</i>	20 – 40	500 – 2500
25.	Muthuga	<i>Butea monosperma</i>	20 - 40	400 - 1500
26.	Hippe	<i>Madhuca latifolia</i>	20 - 40	500 - 2000
27.	Sandal	<i>Santalum album</i>	20 - 50	400 - 1000
28.	Nelli	<i>Emblca officinalis</i>	20 - 40	500 - 2000
29.	Nerale	<i>Sizyzium cumini</i>	20 - 40	500 - 2000
30.	Dhaman	<i>Grevia tilifolia</i>	20 - 40	500 - 2000
31.	Kaval	<i>Careya arborea</i>	20 - 40	500 - 2000
32.	Harada	<i>Terminalia chebula</i>	20 - 40	500 - 2000



## References

1. FAO (1976) Framework for Land Evaluation, Food and Agriculture Organization, Rome.72 pp.
2. FAO (1983) Guidelines for Land Evaluation for Rainfed Agriculture, FAO, Rome, 237 pp.
3. IARI (1971) Soil Survey Manual, All India Soil and Land Use Survey Organization, IARI, New Delhi, 121 pp.
4. Katyal, J.C. and Rattan, R.K. (2003) Secondary and Micronutrients; Research Gap and Future Needs. Fert. News 48 (4); 9-20.
5. Naidu, L.G.K., Ramamurthy, V., Challa, O., Hegde, R. and Krishnan, P. (2006) Manual Soil Site Suitability Criteria for Major Crops, NBSS Publ. No. 129, NBSS & LUP, Nagpur, 118 pp.
6. Natarajan, A. and Dipak Sarkar (2010) Field Guide for Soil Survey, National Bureau of Soil Survey and Land Use Planning (ICAR), Nagpur, India.
7. Natarajan, A., Rajendra Hegde, Raj, J.N. and Shivananda Murthy, H.G. (2015) Implementation Manual for Sujala-III Project, Watershed Development Department, Bengaluru, Karnataka.
8. Sarma, V.A.K., Krishnan, P. and Budihal, S.L. (1987) Laboratory Manual, Tech. Bull. 23, NBSS &LUP, Nagpur.
9. Sehgal, J.L. (1990) Soil Resource Mapping of Different States of India; Why and How?, National Bureau of Soil Survey and Land Use Planning, Nagpur, 49 pp.
10. Shivaprasad, C.R., R.S. Reddy, J. Sehgal and M. Velayuthum (1998) Soils of Karnataka for Optimizing Land Use, NBSS Publ. No. 47b, NBSS & LUP, Nagpur, India.
11. Soil Survey Staff (2006) Keys to Soil Taxonomy, Tenth edition, U.S. Department of Agriculture/ NRCS, Washington DC, U.S.A.
12. Soil Survey Staff (2012) Soil Survey Manual, Handbook No. 18, USDA, Washington DC, USA.





**Appendix I**  
**Belagiri1\_1H1d Microwatershed**  
**Soil Phase Information**

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	WELLS	Land Capability	Conservation Plan
Belagera	1	0.02	Habitation	Others	Others	Others	Others	Others	Others	Others	Not Available (NA)	Not Available	Others	Others
Belagera	2	0.11	Habitation	Others	Others	Others	Others	Others	Others	Others	Not Available (NA)	Not Available	Others	Others
Belagera	3	0.32	Habitation	Others	Others	Others	Others	Others	Others	Others	Not Available (NA)	Not Available	Others	Others
Belagera	4	3.98	Habitation	Others	Others	Others	Others	Others	Others	Others	Redgram (Rg)	Not Available	Others	Others
Belagera	5(1)	1.73	MDRhB2	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iie	Graded bunding
Belagera	5(2)	1.42	MDRhB2	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iie	Graded bunding
Belagera	6(1)	1.34	MDRhB2	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	1 Bore Well	Iie	Graded bunding
Belagera	6(2)	1.36	MDRhB2	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iie	Graded bunding
Belagera	7(1)	2.57	MDRhB2	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iie	Graded bunding
Belagera	7(2)	0.76	MDRhB2	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iie	Graded bunding
Belagera	7(3)	0.57	MDRhB2	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iie	Graded bunding
Belagera	8(1)	1.12	MDRhB2	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iie	Graded bunding
Belagera	8(2)	0.48	MDRhB2	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iie	Graded bunding
Belagera	9(1)	1.56	MDRhB2	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Iie	Graded bunding
Belagera	9(2)	2.53	MDRhB2	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Iie	Graded bunding
Belagera	10(1)	1.48	MDRhB2	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iie	Graded bunding
Belagera	10(2)	1.08	MDRhB2	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iie	Graded bunding
Belagera	11(1)	1.27	MDRhB2	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iie	Graded bunding
Belagera	11(2)	1.56	MDRhB2	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iie	Graded bunding
Belagera	12(1)	1.15	MDRhB2	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iie	Graded bunding
Belagera	12(2)	1.75	YDRcB2	LMU-1	Deep (100-150 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iies	Graded bunding
Belagera	13	2.93	YDRcB2	LMU-1	Deep (100-150 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iies	Graded bunding

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	WELLS	Land Capability	Conservation Plan
Belagera	14	3.82	MDRhB2	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iie	Graded bunding
Belagera	15	5.42	MDRhB2	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iie	Graded bunding
Belagera	16	0.46	MDRhB2	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iie	Graded bunding
Belagera	17	2.55	MDRhB2	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iie	Graded bunding
Belagera	18	5.37	MDRhB2	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Iie	Graded bunding
Belagera	19	0.24	MDRhB2	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iie	Graded bunding
Belagera	20	3.53	MDRhB2	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Iie	Graded bunding
Belagera	21	4.19	MDRhB2	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Iie	Graded bunding
Belagera	22	3.2	MDRhB2	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut (Gn)	Not Available	Iie	Graded bunding
Belagera	23	2.97	MDRhB2	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iie	Graded bunding
Belagera	24	0.2	MDRhB2	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iie	Graded bunding
Belagera	25	3.81	MDRhB2	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Iie	Graded bunding
Belagera	26	1.71	MDRhB2	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut (Gn)	Not Available	Iie	Graded bunding
Belagera	27	2.1	MDRhB2	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Iie	Graded bunding
Belagera	28	4.99	YDRiB2	LMU-1	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Iies	Graded bunding
Belagera	29	0.37	YDRiB2	LMU-1	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iies	Graded bunding
Belagera	30	1.33	YDRiB2	LMU-1	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Iies	Graded bunding
Belagera	31	2.59	MDRhB2	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iie	Graded bunding
Belagera	32	0.8	MDRiA1	LMU-1	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Not Available (NA)	Not Available	Iis	Graded bunding
Belagera	33	1.23	MDRhB2	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iie	Graded bunding
Belagera	34	3.11	MDRhB2	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iie	Graded bunding
Belagera	35	2.31	MDRiA1	LMU-1	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Redgram (Rg)	Not Available	Iis	Graded bunding
Belagera	36	3.05	Habitation	Others	Others	Others	Others	Others	Others	Others	Redgram (Rg)	Not Available	Others	Others
Belagera	37	1.1	MDRiA1	LMU-1	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Not Available (NA)	Not Available	Iis	Graded bunding

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	WELLS	Land Capability	Conservation Plan
Belagera	38	4.42	MDRiA1	LMU-1	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Not Available (NA)	Not Available	IIs	Graded bunding
Belagera	39	2.07	MDRiA1	LMU-1	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Not Available (NA)	Not Available	IIs	Graded bunding
Belagera	40	4.32	MDRiA1	LMU-1	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Redgram (Rg)	Not Available	IIs	Graded bunding
Belagera	41	2.39	MDRiA1	LMU-1	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Redgram (Rg)	Not Available	IIs	Graded bunding
Belagera	42	2.06	MDRiA1	LMU-1	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Redgram (Rg)	Not Available	IIs	Graded bunding
Belagera	43	4.46	MDRiA1	LMU-1	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Redgram (Rg)	Not Available	IIs	Graded bunding
Belagera	44	0.49	MDRiA1	LMU-1	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Not Available (NA)	Not Available	IIs	Graded bunding
Belagera	45	0.5	YDRcB2	LMU-1	Deep (100-150 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIs	Graded bunding
Belagera	46	0.48	YDRcB2	LMU-1	Deep (100-150 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIs	Graded bunding
Belagera	47	7.14	MDRiA1	LMU-1	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Cotton (Ct)	Not Available	IIs	Graded bunding
Belagera	48	2.91	MDRiA1	LMU-1	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Redgram (Rg)	Not Available	IIs	Graded bunding
Belagera	49	5.46	YDRiB2	LMU-1	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	IIs	Graded bunding
Belagera	50	0.08	YDRiB2	LMU-1	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	IIs	Graded bunding
Belagera	51	3	YDRcB2	LMU-1	Deep (100-150 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	IIs	Graded bunding
Belagera	52	0.48	MDRcB2	LMU-1	Very deep (>150 cm)	Sandy loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIs	Graded bunding
Belagera	53	4.6	YDRcB2	LMU-1	Deep (100-150 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIs	Graded bunding
Belagera	54	1.28	YDRcB2	LMU-1	Deep (100-150 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIs	Graded bunding
Belagera	55	0.92	YDRcB2	LMU-1	Deep (100-150 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIs	Graded bunding
Belagera	56	0.72	MDRiA1	LMU-1	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Not Available (NA)	Not Available	IIs	Graded bunding
Belagera	57	1.96	YDRcB2	LMU-1	Deep (100-150 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIs	Graded bunding
Belagera	58	3.94	MDRcB2	LMU-1	Very deep (>150 cm)	Sandy loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIs	Graded bunding
Belagera	59	7.52	YDRcB2	LMU-1	Deep (100-150 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIs	Graded bunding
Belagera	60	4.87	YDRcB2	LMU-1	Deep (100-150 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIs	Graded bunding
Belagera	61	6.74	YDRcB2	LMU-1	Deep (100-150 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy+Scrubland+Waterbody (Pd+Sl+Wb)	Not Available	IIs	Graded bunding

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	WELLS	Land Capability	Conservation Plan
Belagera	62	2.14	YDRcB2	LMU-1	Deep (100-150 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Belagera	64	0.16	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Not Available (NA)	Not Available	Ro	Ro
Belagera	72	126.01	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro (Rc)	Not Available	Ro	Ro
Belagera	90	0.37	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Not Available (NA)	Not Available	Ro	Ro
Belagera	91	6.46	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Not Available (NA)	Not Available	Ro	Ro
Belagera	92	9.96	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Not Available (NA)	Not Available	Ro	Ro
Belagera	93	7.37	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Not Available (NA)	Not Available	Ro	Ro
Belagera	94	7.22	HSLiB2	LMU-3	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut+Cotton (Gn+Ct)	Not Available	Iles	Graded bunding
Belagera	95	7.4	MDRiA1	LMU-1	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Cotton+Paddy (Ct+Pd)	Not Available	IIs	Graded bunding
Belagera	96	4.67	MDRiA1	LMU-1	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Groundnut (Gn)	Not Available	IIs	Graded bunding
Belagera	97	4.36	MDRiA1	LMU-1	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Groundnut (Gn)	Not Available	IIs	Graded bunding
Belagera	98	7.16	MDRiA1	LMU-1	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Cotton (Ct)	Not Available	IIs	Graded bunding
Belagera	99	2.97	MDRiA1	LMU-1	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Cotton (Ct)	Not Available	IIs	Graded bunding
Belagera	100	2.09	MDRiA1	LMU-1	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Groundnut (Gn)	1 Bore Well	IIs	Graded bunding
Belagera	101/1	0.76	MDRiA1	LMU-1	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Groundnut (Gn)	Not Available	IIs	Graded bunding
Belagera	101/2	0.64	MDRiA1	LMU-1	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Groundnut (Gn)	Not Available	IIs	Graded bunding
Belagera	101(3)	0.69	MDRiA1	LMU-1	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Groundnut (Gn)	Not Available	IIs	Graded bunding
Belagera	102(1)	0.73	MDRiA1	LMU-1	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Groundnut (Gn)	Not Available	IIs	Graded bunding
Belagera	102(2)	0.65	MDRiA1	LMU-1	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Groundnut (Gn)	Not Available	IIs	Graded bunding
Belagera	102(3)	0.74	MDRiA1	LMU-1	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Groundnut (Gn)	Not Available	IIs	Graded bunding
Belagera	103(1)	2.72	MDRcB2	LMU-1	Very deep (>150 cm)	Sandy loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Belagera	103(2)	1.05	HSLiB2	LMU-3	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut (Gn)	Not Available	Iles	Graded bunding
Belagera	104	1.9	HSLiB2	LMU-3	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Belagera	105	2.74	HSLiB2	LMU-3	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	1 Bore Well	Iles	Graded bunding

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	WELLS	Land Capability	Conservation Plan
Belagera	106	5.55	HSLiB2	LMU-3	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Belagera	107	2.39	HSLiB2	LMU-3	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Belagera	108	3.85	HSLiB2	LMU-3	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Iles	Graded bunding
Belagera	109	5.44	HSLiB2	LMU-3	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Fallow Land (Rg+Fl)	Not Available	Iles	Graded bunding
Belagera	110	2.34	HSLiB2	LMU-3	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Iles	Graded bunding
Belagera	111	7.74	Waterbody	Others	Others	Others	Others	Others	Others	Others	Waterbody+Fallow Land (Wb+Fl)	Not Available	Others	Others
Belagera	112(1)	0.39	Habitation	Others	Others	Others	Others	Others	Others	Others	Not Available (NA)	Not Available	Others	Others
Belagera	112(2)	0.45	BMNmB2	LMU-3	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iles	Graded bunding
Belagera	113	0.2	BMNmB2	LMU-3	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iles	Graded bunding
Belagera	114	0.25	Habitation	Others	Others	Others	Others	Others	Others	Others	Not Available (NA)	Not Available	Others	Others
Belagera	115	0.67	BMNmB2	LMU-3	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iles	Graded bunding
Belagera	116	0.42	BMNmB2	LMU-3	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iles	Graded bunding
Belagera	117	1.43	BMNmB2	LMU-3	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iles	Graded bunding
Belagera	118	0.64	BMNmB2	LMU-3	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iles	Graded bunding
Belagera	119	1.01	BMNmB2	LMU-3	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iles	Graded bunding
Belagera	120	0.78	BMNmB2	LMU-3	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iles	Graded bunding
Belagera	121	0.88	BMNmB2	LMU-3	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iles	Graded bunding
Belagera	122	0.96	BMNmB2	LMU-3	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iles	Graded bunding
Belagera	123	0.57	BMNmB2	LMU-3	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iles	Graded bunding
Belagera	124(1)	1.58	BMNmB2	LMU-3	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iles	Graded bunding
Belagera	124(2)	2.36	BMNmB2	LMU-3	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iles	Graded bunding
Belagera	125	1.97	MDGhB2g1	LMU-1	Deep (100-150 cm)	Sandy clay loam	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iles	Graded bunding
Belagera	126	2.38	BMNmB2	LMU-3	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iles	Graded bunding
Belagera	127	8.85	MDGhB2g1	LMU-1	Deep (100-150 cm)	Sandy clay loam	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	1 Bore Well	Iles	Graded bunding



Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	WELLS	Land Capability	Conservation Plan
Belagera	128(1)	2.49	HSLiB2	LMU-3	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Belagera	128(2)	0.98	HSLiB2	LMU-3	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iles	Graded bunding
Belagera	129	0.2	HSLiB2	LMU-3	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iles	Graded bunding
Belagera	130	2.26	HSLiB2	LMU-3	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Belagera	131	3.37	HSLiB2	LMU-3	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Iles	Graded bunding
Belagera	132	7.45	HSLiB2	LMU-3	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Iles	Graded bunding
Belagera	133	6.66	YDRcB2	LMU-1	Deep (100-150 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Scrubland (Sl)	Not Available	Iles	Graded bunding
Belagera	134	6.09	YDRcB2	LMU-1	Deep (100-150 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Scrubland (Sl)	Not Available	Iles	Graded bunding
Belagera	135	7.62	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Redgram (Rg)	Not Available	Ro	Ro
Belagera	136	0	MDRhB2	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Ile	Graded bunding
Belagera	137	1.17	MDRhB2	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Ile	Graded bunding
Belagera	138	7.77	MDRhB2	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Ile	Graded bunding
Belagera	139	8.09	MDRhB2	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Ile	Graded bunding
Belagera	140	3.5	MDRhB2	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	1 Bore Well	Ile	Graded bunding
Belagera	141	2.95	BLCcB2	LMU-4	Moderately deep (75-100 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Iles	TCB
Belagera	148	2.82	BDLiB3	LMU-5	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Cotton+Ro (Ct+Rc)	Not Available	Ives	Graded bunding
Belagera	149	1.02	BLCcB2	LMU-4	Moderately deep (75-100 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iles	TCB
Belagera	150	2.32	BLCcB2	LMU-4	Moderately deep (75-100 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Iles	TCB
Belagera	151	4.86	BLCcB2	LMU-4	Moderately deep (75-100 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Iles	TCB
Belagera	152	6.81	BLCcB2	LMU-4	Moderately deep (75-100 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iles	TCB
Belagera	153	3.42	BLCcB2	LMU-4	Moderately deep (75-100 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Iles	TCB
Belagera	154	1.83	MDGhB2g1	LMU-1	Deep (100-150 cm)	Sandy clay loam	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton+Fallow Land (Ct+Fl)	Not Available	Iles	Graded bunding
Belagera	157	2.63	MDGhB2g1	LMU-1	Deep (100-150 cm)	Sandy clay loam	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Iles	Graded bunding
Belagera	158	1.62	MDGhB2g1	LMU-1	Deep (100-150 cm)	Sandy clay loam	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut+Cotton (Gn+Ct)	Not Available	Iles	Graded bunding



Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	WELLS	Land Capability	Conservation Plan
Belagera	159	0.13	ADGhB2g1	LMU-1	Deep (100-150 cm)	Sandy clay loam	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton+Redgram (Ct+Rg)	Not Available	Iles	Graded bunding
Belagera	160	0.13	MNmB2	LMU-3	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Belagera	161(1)	0.44	ADGhB2g1	LMU-1	Deep (100-150 cm)	Sandy clay loam	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iles	Graded bunding
Belagera	164	0.12	ADGhB2g1	LMU-1	Deep (100-150 cm)	Sandy clay loam	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iles	Graded bunding
Belagera	165	0.57	ADGhB2g1	LMU-1	Deep (100-150 cm)	Sandy clay loam	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iles	Graded bunding
Belagera	250	0.68	MDRhB2	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Ile	Graded bunding
Belagera	251	0.4	MDRhB2	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Ile	Graded bunding
Belagera	252	1.34	ADGhB2g1	LMU-1	Deep (100-150 cm)	Sandy clay loam	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iles	Graded bunding
Belagera	253	0.34	habitation	Others	Others	Others	Others	Others	Others	Others	Not Available (NA)	Not Available	Others	Others
Belagera	254	2.96	habitation	Others	Others	Others	Others	Others	Others	Others	Not Available (NA)	Not Available	Others	Others
Belagera	255	2.45	MDRhB2	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Ile	Graded bunding
Belagera	256	0.87	MDRhB2	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Ile	Graded bunding
Belagera	257	0.43	MDRhB2	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Ile	Graded bunding
Belagera	258	0.68	MDRhB2	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Ile	Graded bunding
Belagera	259	0.39	MDRhB2	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Ile	Graded bunding
Belagera	260	0.29	MDRhB2	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Ile	Graded bunding
Belagera	261	0.65	MDRhB2	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Ile	Graded bunding
Belagera	289	0.16	MDRhB2	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Ile	Graded bunding
Belagera	290	0.49	MDRhB2	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Ile	Graded bunding
Belagera	292	0.11	YDRcB2	LMU-1	Deep (100-150 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Belagera	293	1.62	YDRcB2	LMU-1	Deep (100-150 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iles	Graded bunding
Belagera	295	1.72	MDRhB2	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Ile	Graded bunding
Belagera	299	0.32	MDRhB2	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Ile	Graded bunding
Belagera	300	0.28	MDRhB2	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Ile	Graded bunding

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	WELLS	Land Capability	Conservation Plan
Belagera	301	0.42	MDRhB2	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Ile	Graded bunding
Belagera	302	0.19	MDRhB2	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Ile	Graded bunding
Belagera	303	4.79	MDRhB2	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Ile	Graded bunding
Belagera	304	0.67	MDRhB2	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Ile	Graded bunding
Belagera	305(1)	1	MDRhB2	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Ile	Graded bunding
Belagera	305(2)	1.04	MDRhB2	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Ile	Graded bunding
Belagera	305(3)	0.71	MDRhB2	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Ile	Graded bunding
Belagera	306	0.39	MDRhB2	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Ile	Graded bunding
Belagera	307	0.55	MDRhB2	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Ile	Graded bunding
Belagera	308	0.37	MDRhB2	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Ile	Graded bunding
Belagera	309(1)	1.95	MDRhB2	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Ile	Graded bunding
Belagera	309(2)	1.17	MDRhB2	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Ile	Graded bunding
Belagera	310(1)	0.5	MDRhB2	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Ile	Graded bunding
Belagera	310(2)	1.23	Habitation	Others	Others	Others	Others	Others	Others	Others	Not Available (NA)	Not Available	Others	Others
Belagera	310(3)	0.83	MDRhB2	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Ile	Graded bunding
Belagera	311	0.63	MDRhB2	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Ile	Graded bunding
Belagera	312(1)	1.14	MDRhB2	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Ile	Graded bunding
Belagera	312(2)	1.31	Habitation	Others	Others	Others	Others	Others	Others	Others	Not Available (NA)	Not Available	Others	Others
Belagera	313(1)	0.15	Habitation	Others	Others	Others	Others	Others	Others	Others	Not Available (NA)	Not Available	Others	Others
Belagera	313(2)	0.82	Habitation	Others	Others	Others	Others	Others	Others	Others	Not Available (NA)	Not Available	Others	Others
Honagera	206	0.25	MDRcB2	LMU-1	Very deep (>150 cm)	Sandy loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut (Gn)	Not Available	Iles	Graded bunding
Honagera	208	5.53	MDRcB2	LMU-1	Very deep (>150 cm)	Sandy loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut (Gn)	Not Available	Iles	Graded bunding
Honagera	209	3.56	MDRcB2	LMU-1	Very deep (>150 cm)	Sandy loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Honagera	210	0.7	MDRcB2	LMU-1	Very deep (>150 cm)	Sandy loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar+Groundnut (Jw+Gn)	Not Available	Iles	Graded bunding

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	WELLS	Land Capability	Conservation Plan
Honagera	211	2.29	MDRcB2	LMU-1	Very deep (>150 cm)	Sandy loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut (Gn)	Not Available	Iles	Graded bunding
Honagera	226	0.16	MDRcB2	LMU-1	Very deep (>150 cm)	Sandy loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Sutharaho salli	91	0	HSLhB2	LMU-3	Moderately deep (75-100 cm)	Sandy clay loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iles	Graded bunding
Sutharaho salli	92	3.96	HSLhB2	LMU-3	Moderately deep (75-100 cm)	Sandy clay loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut (Gn)	Not Available	Iles	Graded bunding
Sutharaho salli	93	2.01	KBDcC2g1	LMU-2	Moderately deep (75-100 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Moderate	Redgram (Rg)	Not Available	IIles	TCB
Sutharaho salli	94	3.33	KBDcC2g1	LMU-2	Moderately deep (75-100 cm)	Sandy loam	Gravelly (15-35%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Moderate	Redgram (Rg)	Not Available	IIles	TCB
Sutharaho salli	95	0.25	HSLhB2	LMU-3	Moderately deep (75-100 cm)	Sandy clay loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut (Gn)	Not Available	Iles	Graded bunding
Sutharaho salli	96	0.15	HSLiB2	LMU-3	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy+Groundnut (Pd+Gn)	Not Available	Iles	Graded bunding
Sutharaho salli	97	6.82	HSLiB2	LMU-3	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy+Groundnut (Pd+Gn)	Not Available	Iles	Graded bunding
Sutharaho salli	98	4.3	HSLiB2	LMU-3	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	Iles	Graded bunding
Sutharaho salli	99	0.23	HSLiB2	LMU-3	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut (Gn)	Not Available	Iles	Graded bunding
Sutharaho salli	100	15.74	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro (Rc)	Not Available	Ro	Ro

Note:Ro- Rock outcrops





















Village	Survey Number	Soil Reaction	Salinity	Organic Carbon	Available Phosphorus	Available Potassium	Available Sulphur	Available Boron	Available Iron	Available Manganese	Available Copper	Available Zinc
Sutharah osalli	93	Neutral (pH 6.5 - 7.3)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Sutharah osalli	94	Neutral (pH 6.5 - 7.3)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Sutharah osalli	95	Neutral (pH 6.5 - 7.3)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Sutharah osalli	96	Neutral (pH 6.5 - 7.3)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Sutharah osalli	97	Neutral (pH 6.5 - 7.3)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Sutharah osalli	98	Neutral (pH 6.5 - 7.3)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Sutharah osalli	99	Neutral (pH 6.5 - 7.3)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Sutharah osalli	100	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro



**Appendix III**  
**Belagiri\_1H1d Microwatershed**  
**Soil Suitability Information**

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengalgram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry	
Belagera	1	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Belagera	2	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Belagera	3	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Belagera	4	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Belagera	5(1)	S3tz	S2tw	S3t	S1	S3t	S2z	S2t	S2z	S1	S2zw	S2tw	S2t	S3t	S1	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2tz	S1	S1	S2tw	S3tw	
Belagera	5(2)	S3tz	S2tw	S3t	S1	S3t	S2z	S2t	S2z	S1	S2zw	S2tw	S2t	S3t	S1	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2tz	S1	S1	S2tw	S3tw	
Belagera	6(1)	S3tz	S2tw	S3t	S1	S3t	S2z	S2t	S2z	S1	S2zw	S2tw	S2t	S3t	S1	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2tz	S1	S1	S2tw	S3tw	
Belagera	6(2)	S3tz	S2tw	S3t	S1	S3t	S2z	S2t	S2z	S1	S2zw	S2tw	S2t	S3t	S1	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2tz	S1	S1	S2tw	S3tw	
Belagera	7(1)	S3tz	S2tw	S3t	S1	S3t	S2z	S2t	S2z	S1	S2zw	S2tw	S2t	S3t	S1	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2tz	S1	S1	S2tw	S3tw	
Belagera	7(2)	S3tz	S2tw	S3t	S1	S3t	S2z	S2t	S2z	S1	S2zw	S2tw	S2t	S3t	S1	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2tz	S1	S1	S2tw	S3tw	
Belagera	7(3)	S3tz	S2tw	S3t	S1	S3t	S2z	S2t	S2z	S1	S2zw	S2tw	S2t	S3t	S1	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2tz	S1	S1	S2tw	S3tw	
Belagera	8(1)	S3tz	S2tw	S3t	S1	S3t	S2z	S2t	S2z	S1	S2zw	S2tw	S2t	S3t	S1	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2tz	S1	S1	S2tw	S3tw	
Belagera	8(2)	S3tz	S2tw	S3t	S1	S3t	S2z	S2t	S2z	S1	S2zw	S2tw	S2t	S3t	S1	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2tz	S1	S1	S2tw	S3tw	
Belagera	9(1)	S3tz	S2tw	S3t	S1	S3t	S2z	S2t	S2z	S1	S2zw	S2tw	S2t	S3t	S1	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2tz	S1	S1	S2tw	S3tw	
Belagera	9(2)	S3tz	S2tw	S3t	S1	S3t	S2z	S2t	S2z	S1	S2zw	S2tw	S2t	S3t	S1	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2tz	S1	S1	S2tw	S3tw	
Belagera	10(1)	S3tz	S2tw	S3t	S1	S3t	S2z	S2t	S2z	S1	S2zw	S2tw	S2t	S3t	S1	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2tz	S1	S1	S2tw	S3tw	
Belagera	10(2)	S3tz	S2tw	S3t	S1	S3t	S2z	S2t	S2z	S1	S2zw	S2tw	S2t	S3t	S1	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2tz	S1	S1	S2tw	S3tw	
Belagera	11(1)	S3tz	S2tw	S3t	S1	S3t	S2z	S2t	S2z	S1	S2zw	S2tw	S2t	S3t	S1	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2tz	S1	S1	S2tw	S3tw	
Belagera	11(2)	S3tz	S2tw	S3t	S1	S3t	S2z	S2t	S2z	S1	S2zw	S2tw	S2t	S3t	S1	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2tz	S1	S1	S2tw	S3tw	
Belagera	12(1)	S3tz	S2tw	S3t	S1	S3t	S2z	S2t	S2z	S1	S2zw	S2tw	S2t	S3t	S1	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2tz	S1	S1	S2tw	S3tw	
Belagera	12(2)	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz	
Belagera	13	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz	
Belagera	14	S3tz	S2tw	S3t	S1	S3t	S2z	S2t	S2z	S1	S2zw	S2tw	S2t	S3t	S1	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2tz	S1	S1	S2tw	S3tw	

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengalgram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry	
Belagera	15	S3tz	S2tw	S3t	S1	S3t	S2z	S2t	S2z	S1	S2zw	S2tw	S2t	S3t	S1	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2tz	S1	S1	S2tw	S3tw	
Belagera	16	S3tz	S2tw	S3t	S1	S3t	S2z	S2t	S2z	S1	S2zw	S2tw	S2t	S3t	S1	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2tz	S1	S1	S2tw	S3tw	
Belagera	17	S3tz	S2tw	S3t	S1	S3t	S2z	S2t	S2z	S1	S2zw	S2tw	S2t	S3t	S1	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2tz	S1	S1	S2tw	S3tw	
Belagera	18	S3tz	S2tw	S3t	S1	S3t	S2z	S2t	S2z	S1	S2zw	S2tw	S2t	S3t	S1	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2tz	S1	S1	S2tw	S3tw	
Belagera	19	S3tz	S2tw	S3t	S1	S3t	S2z	S2t	S2z	S1	S2zw	S2tw	S2t	S3t	S1	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2tz	S1	S1	S2tw	S3tw	
Belagera	20	S3tz	S2tw	S3t	S1	S3t	S2z	S2t	S2z	S1	S2zw	S2tw	S2t	S3t	S1	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2tz	S1	S1	S2tw	S3tw	
Belagera	21	S3tz	S2tw	S3t	S1	S3t	S2z	S2t	S2z	S1	S2zw	S2tw	S2t	S3t	S1	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2tz	S1	S1	S2tw	S3tw	
Belagera	22	S3tz	S2tw	S3t	S1	S3t	S2z	S2t	S2z	S1	S2zw	S2tw	S2t	S3t	S1	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2tz	S1	S1	S2tw	S3tw	
Belagera	23	S3tz	S2tw	S3t	S1	S3t	S2z	S2t	S2z	S1	S2zw	S2tw	S2t	S3t	S1	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2tz	S1	S1	S2tw	S3tw	
Belagera	24	S3tz	S2tw	S3t	S1	S3t	S2z	S2t	S2z	S1	S2zw	S2tw	S2t	S3t	S1	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2tz	S1	S1	S2tw	S3tw	
Belagera	25	S3tz	S2tw	S3t	S1	S3t	S2z	S2t	S2z	S1	S2zw	S2tw	S2t	S3t	S1	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2tz	S1	S1	S2tw	S3tw	
Belagera	26	S3tz	S2tw	S3t	S1	S3t	S2z	S2t	S2z	S1	S2zw	S2tw	S2t	S3t	S1	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2tz	S1	S1	S2tw	S3tw	
Belagera	27	S3tz	S2tw	S3t	S1	S3t	S2z	S2t	S2z	S1	S2zw	S2tw	S2t	S3t	S1	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2tz	S1	S1	S2tw	S3tw	
Belagera	28	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz	
Belagera	29	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz	
Belagera	30	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz	
Belagera	31	S3tz	S2tw	S3t	S1	S3t	S2z	S2t	S2z	S1	S2zw	S2tw	S2t	S3t	S1	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2tz	S1	S1	S2tw	S3tw	
Belagera	32	S3tz	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tz	S2t	S1	S2tw	S3tw	
Belagera	33	S3tz	S2tw	S3t	S1	S3t	S2z	S2t	S2z	S1	S2zw	S2tw	S2t	S3t	S1	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2tz	S1	S1	S2tw	S3tw	
Belagera	34	S3tz	S2tw	S3t	S1	S3t	S2z	S2t	S2z	S1	S2zw	S2tw	S2t	S3t	S1	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2tz	S1	S1	S2tw	S3tw	
Belagera	35	S3tz	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tz	S2t	S1	S2tw	S3tw	
Belagera	36	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Belagera	37	S3tz	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tz	S2t	S1	S2tw	S3tw	
Belagera	38	S3tz	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tz	S2t	S1	S2tw	S3tw	
Belagera	39	S3tz	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tz	S2t	S1	S2tw	S3tw	
Belagera	40	S3tz	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tz	S2t	S1	S2tw	S3tw	
Belagera	41	S3tz	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tz	S2t	S1	S2tw	S3tw	



Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengalgram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry
Belagera	42	S3tz	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tz	S2t	S1	S2tw	S3tw
Belagera	43	S3tz	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tz	S2t	S1	S2tw	S3tw
Belagera	44	S3tz	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tz	S2t	S1	S2tw	S3tw
Belagera	45	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Belagera	46	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Belagera	47	S3tz	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tz	S2t	S1	S2tw	S3tw
Belagera	48	S3tz	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tz	S2t	S1	S2tw	S3tw
Belagera	49	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Belagera	50	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Belagera	51	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Belagera	52	S3tz	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tz	S2t	S1	S2tw	S3tw
Belagera	53	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Belagera	54	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Belagera	55	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Belagera	56	S3tz	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tz	S2t	S1	S2tw	S3tw
Belagera	57	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Belagera	58	S3tz	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tz	S2t	S1	S2tw	S3tw
Belagera	59	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Belagera	60	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Belagera	61	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Belagera	62	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Belagera	64	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro
Belagera	72	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro
Belagera	90	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro
Belagera	91	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro
Belagera	92	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro
Belagera	93	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengalgram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry	
Belagera	94	S3rz	S2tz	S2rz	S2tz	S2rz	S3tz	S3rz	S2rz	S3tz	S2rz	S2rz	S2tz	S2rz	S1	N1tz	S3rz	S2rz	S2z	S1	S2z	S2z	S2z	S2z	S2rz	S2z	S2t	S1	S2rz	S2rz	
Belagera	95	S3tz	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tz	S2t	S1	S2tw	S3tw	
Belagera	96	S3tz	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tz	S2t	S1	S2tw	S3tw	
Belagera	97	S3tz	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tz	S2t	S1	S2tw	S3tw	
Belagera	98	S3tz	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tz	S2t	S1	S2tw	S3tw	
Belagera	99	S3tz	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tz	S2t	S1	S2tw	S3tw	
Belagera	100	S3tz	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tz	S2t	S1	S2tw	S3tw	
Belagera	101/1	S3tz	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tz	S2t	S1	S2tw	S3tw	
Belagera	101/2	S3tz	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tz	S2t	S1	S2tw	S3tw	
Belagera	101(3)	S3tz	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tz	S2t	S1	S2tw	S3tw	
Belagera	102(1)	S3tz	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tz	S2t	S1	S2tw	S3tw	
Belagera	102(2)	S3tz	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tz	S2t	S1	S2tw	S3tw	
Belagera	102(3)	S3tz	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tz	S2t	S1	S2tw	S3tw	
Belagera	103(1)	S3tz	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tz	S2t	S1	S2tw	S3tw	
Belagera	103(2)	S3rz	S2tz	S2rz	S2tz	S2rz	S3tz	S3rz	S2rz	S3tz	S2rz	S2rz	S2tz	S2rz	S1	N1tz	S3rz	S2rz	S2z	S1	S2z	S2z	S2z	S2z	S2rz	S2z	S2t	S1	S2rz	S2rz	
Belagera	104	S3rz	S2tz	S2rz	S2tz	S2rz	S3tz	S3rz	S2rz	S3tz	S2rz	S2rz	S2tz	S2rz	S1	N1tz	S3rz	S2rz	S2z	S1	S2z	S2z	S2z	S2z	S2rz	S2z	S2t	S1	S2rz	S2rz	
Belagera	105	S3rz	S2tz	S2rz	S2tz	S2rz	S3tz	S3rz	S2rz	S3tz	S2rz	S2rz	S2tz	S2rz	S1	N1tz	S3rz	S2rz	S2z	S1	S2z	S2z	S2z	S2z	S2rz	S2z	S2t	S1	S2rz	S2rz	
Belagera	106	S3rz	S2tz	S2rz	S2tz	S2rz	S3tz	S3rz	S2rz	S3tz	S2rz	S2rz	S2tz	S2rz	S1	N1tz	S3rz	S2rz	S2z	S1	S2z	S2z	S2z	S2z	S2rz	S2z	S2t	S1	S2rz	S2rz	
Belagera	107	S3rz	S2tz	S2rz	S2tz	S2rz	S3tz	S3rz	S2rz	S3tz	S2rz	S2rz	S2tz	S2rz	S1	N1tz	S3rz	S2rz	S2z	S1	S2z	S2z	S2z	S2z	S2rz	S2z	S2t	S1	S2rz	S2rz	
Belagera	108	S3rz	S2tz	S2rz	S2tz	S2rz	S3tz	S3rz	S2rz	S3tz	S2rz	S2rz	S2tz	S2rz	S1	N1tz	S3rz	S2rz	S2z	S1	S2z	S2z	S2z	S2z	S2rz	S2z	S2t	S1	S2rz	S2rz	
Belagera	109	S3rz	S2tz	S2rz	S2tz	S2rz	S3tz	S3rz	S2rz	S3tz	S2rz	S2rz	S2tz	S2rz	S1	N1tz	S3rz	S2rz	S2z	S1	S2z	S2z	S2z	S2z	S2rz	S2z	S2t	S1	S2rz	S2rz	
Belagera	110	S3rz	S2tz	S2rz	S2tz	S2rz	S3tz	S3rz	S2rz	S3tz	S2rz	S2rz	S2tz	S2rz	S1	N1tz	S3rz	S2rz	S2z	S1	S2z	S2z	S2z	S2z	S2rz	S2z	S2t	S1	S2rz	S2rz	
Belagera	111	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Belagera	112(1)	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Belagera	112(2)	S3t	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw	
Belagera	113	S3t	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw	
Belagera	114	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengalgram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry	
Belagera	115	S3t	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw	
Belagera	116	S3t	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw	
Belagera	117	S3t	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw	
Belagera	118	S3t	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw	
Belagera	119	S3t	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw	
Belagera	120	S3t	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw	
Belagera	121	S3t	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw	
Belagera	122	S3t	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw	
Belagera	123	S3t	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw	
Belagera	124(1)	S3t	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw	
Belagera	124(2)	S3t	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw	
Belagera	125	S2r	S2tw	S3t	S1	S3t	S1	S2t	S2z	S1	S1	S2tw	S2t	S3t	S2z	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2t	S1	S1	S2tw	S3tw	
Belagera	126	S3t	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw	
Belagera	127	S2r	S2tw	S3t	S1	S3t	S1	S2t	S2z	S1	S1	S2tw	S2t	S3t	S2z	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2t	S1	S1	S2tw	S3tw	
Belagera	128(1)	S3rz	S2tz	S2rz	S2tz	S2rz	S3tz	S3rz	S2rz	S3tz	S2rz	S2rz	S2tz	S2rz	S1	N1tz	S3rz	S2rz	S2z	S1	S2z	S2z	S2z	S2z	S2rz	S2z	S2t	S1	S2rz	S2rz	
Belagera	128(2)	S3rz	S2tz	S2rz	S2tz	S2rz	S3tz	S3rz	S2rz	S3tz	S2rz	S2rz	S2tz	S2rz	S1	N1tz	S3rz	S2rz	S2z	S1	S2z	S2z	S2z	S2z	S2rz	S2z	S2t	S1	S2rz	S2rz	
Belagera	129	S3rz	S2tz	S2rz	S2tz	S2rz	S3tz	S3rz	S2rz	S3tz	S2rz	S2rz	S2tz	S2rz	S1	N1tz	S3rz	S2rz	S2z	S1	S2z	S2z	S2z	S2z	S2rz	S2z	S2t	S1	S2rz	S2rz	
Belagera	130	S3rz	S2tz	S2rz	S2tz	S2rz	S3tz	S3rz	S2rz	S3tz	S2rz	S2rz	S2tz	S2rz	S1	N1tz	S3rz	S2rz	S2z	S1	S2z	S2z	S2z	S2z	S2rz	S2z	S2t	S1	S2rz	S2rz	
Belagera	131	S3rz	S2tz	S2rz	S2tz	S2rz	S3tz	S3rz	S2rz	S3tz	S2rz	S2rz	S2tz	S2rz	S1	N1tz	S3rz	S2rz	S2z	S1	S2z	S2z	S2z	S2z	S2rz	S2z	S2t	S1	S2rz	S2rz	
Belagera	132	S3rz	S2tz	S2rz	S2tz	S2rz	S3tz	S3rz	S2rz	S3tz	S2rz	S2rz	S2tz	S2rz	S1	N1tz	S3rz	S2rz	S2z	S1	S2z	S2z	S2z	S2z	S2rz	S2z	S2t	S1	S2rz	S2rz	
Belagera	133	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz	
Belagera	134	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz	
Belagera	135	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro
Belagera	136	S3tz	S2tw	S3t	S1	S3t	S2z	S2t	S2z	S1	S2zw	S2tw	S2t	S3t	S1	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2tz	S1	S1	S2tw	S3tw	
Belagera	137	S3tz	S2tw	S3t	S1	S3t	S2z	S2t	S2z	S1	S2zw	S2tw	S2t	S3t	S1	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2tz	S1	S1	S2tw	S3tw	
Belagera	138	S3tz	S2tw	S3t	S1	S3t	S2z	S2t	S2z	S1	S2zw	S2tw	S2t	S3t	S1	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2tz	S1	S1	S2tw	S3tw	
Belagera	139	S3tz	S2tw	S3t	S1	S3t	S2z	S2t	S2z	S1	S2zw	S2tw	S2t	S3t	S1	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2tz	S1	S1	S2tw	S3tw	

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengalgram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry	
Belagera	140	S3tz	S2tw	S3t	S1	S3t	S2z	S2t	S2z	S1	S2zw	S2tw	S2t	S3t	S1	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2tz	S1	S1	S2tw	S3tw	
Belagera	141	S3rz	S2z	S2rz	S2z	S2rt	S2rz	S3rz	S2rz	S2tz	S2rz	S2rz	S1	S2rz	S1	S2rt	S3rz	S2rz	S2tz	S1	S2tz	S2z	S2tz	S2tz	S2rz	S2z	S1	S1	S2rz	S2rz	
Belagera	148	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	
Belagera	149	S3rz	S2z	S2rz	S2z	S2rt	S2rz	S3rz	S2rz	S2tz	S2rz	S2rz	S1	S2rz	S1	S2rt	S3rz	S2rz	S2tz	S1	S2tz	S2z	S2tz	S2tz	S2rz	S2z	S1	S1	S2rz	S2rz	
Belagera	150	S3rz	S2z	S2rz	S2z	S2rt	S2rz	S3rz	S2rz	S2tz	S2rz	S2rz	S1	S2rz	S1	S2rt	S3rz	S2rz	S2tz	S1	S2tz	S2z	S2tz	S2tz	S2rz	S2z	S1	S1	S2rz	S2rz	
Belagera	151	S3rz	S2z	S2rz	S2z	S2rt	S2rz	S3rz	S2rz	S2tz	S2rz	S2rz	S1	S2rz	S1	S2rt	S3rz	S2rz	S2tz	S1	S2tz	S2z	S2tz	S2tz	S2rz	S2z	S1	S1	S2rz	S2rz	
Belagera	152	S3rz	S2z	S2rz	S2z	S2rt	S2rz	S3rz	S2rz	S2tz	S2rz	S2rz	S1	S2rz	S1	S2rt	S3rz	S2rz	S2tz	S1	S2tz	S2z	S2tz	S2tz	S2rz	S2z	S1	S1	S2rz	S2rz	
Belagera	153	S3rz	S2z	S2rz	S2z	S2rt	S2rz	S3rz	S2rz	S2tz	S2rz	S2rz	S1	S2rz	S1	S2rt	S3rz	S2rz	S2tz	S1	S2tz	S2z	S2tz	S2tz	S2rz	S2z	S1	S1	S2rz	S2rz	
Belagera	154	S2r	S2tw	S3t	S1	S3t	S1	S2t	S2z	S1	S1	S2tw	S2t	S3t	S2z	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2t	S1	S1	S2tw	S3tw	
Belagera	157	S2r	S2tw	S3t	S1	S3t	S1	S2t	S2z	S1	S1	S2tw	S2t	S3t	S2z	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2t	S1	S1	S2tw	S3tw	
Belagera	158	S2r	S2tw	S3t	S1	S3t	S1	S2t	S2z	S1	S1	S2tw	S2t	S3t	S2z	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2t	S1	S1	S2tw	S3tw	
Belagera	159	S2r	S2tw	S3t	S1	S3t	S1	S2t	S2z	S1	S1	S2tw	S2t	S3t	S2z	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2t	S1	S1	S2tw	S3tw	
Belagera	160	S3t	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw	
Belagera	161(1)	S2r	S2tw	S3t	S1	S3t	S1	S2t	S2z	S1	S1	S2tw	S2t	S3t	S2z	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2t	S1	S1	S2tw	S3tw	
Belagera	164	S2r	S2tw	S3t	S1	S3t	S1	S2t	S2z	S1	S1	S2tw	S2t	S3t	S2z	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2t	S1	S1	S2tw	S3tw	
Belagera	165	S2r	S2tw	S3t	S1	S3t	S1	S2t	S2z	S1	S1	S2tw	S2t	S3t	S2z	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2t	S1	S1	S2tw	S3tw	
Belagera	250	S3tz	S2tw	S3t	S1	S3t	S2z	S2t	S2z	S1	S2zw	S2tw	S2t	S3t	S1	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2tz	S1	S1	S2tw	S3tw	
Belagera	251	S3tz	S2tw	S3t	S1	S3t	S2z	S2t	S2z	S1	S2zw	S2tw	S2t	S3t	S1	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2tz	S1	S1	S2tw	S3tw	
Belagera	252	S2r	S2tw	S3t	S1	S3t	S1	S2t	S2z	S1	S1	S2tw	S2t	S3t	S2z	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2t	S1	S1	S2tw	S3tw	
Belagera	253	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Belagera	254	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Belagera	255	S3tz	S2tw	S3t	S1	S3t	S2z	S2t	S2z	S1	S2zw	S2tw	S2t	S3t	S1	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2tz	S1	S1	S2tw	S3tw	
Belagera	256	S3tz	S2tw	S3t	S1	S3t	S2z	S2t	S2z	S1	S2zw	S2tw	S2t	S3t	S1	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2tz	S1	S1	S2tw	S3tw	
Belagera	257	S3tz	S2tw	S3t	S1	S3t	S2z	S2t	S2z	S1	S2zw	S2tw	S2t	S3t	S1	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2tz	S1	S1	S2tw	S3tw	
Belagera	258	S3tz	S2tw	S3t	S1	S3t	S2z	S2t	S2z	S1	S2zw	S2tw	S2t	S3t	S1	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2tz	S1	S1	S2tw	S3tw	
Belagera	259	S3tz	S2tw	S3t	S1	S3t	S2z	S2t	S2z	S1	S2zw	S2tw	S2t	S3t	S1	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2tz	S1	S1	S2tw	S3tw	
Belagera	260	S3tz	S2tw	S3t	S1	S3t	S2z	S2t	S2z	S1	S2zw	S2tw	S2t	S3t	S1	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2tz	S1	S1	S2tw	S3tw	

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengalgram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry	
Belagera	261	S3tz	S2tw	S3t	S1	S3t	S2z	S2t	S2z	S1	S2zw	S2tw	S2t	S3t	S1	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2tz	S1	S1	S2tw	S3tw	
Belagera	289	S3tz	S2tw	S3t	S1	S3t	S2z	S2t	S2z	S1	S2zw	S2tw	S2t	S3t	S1	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2tz	S1	S1	S2tw	S3tw	
Belagera	290	S3tz	S2tw	S3t	S1	S3t	S2z	S2t	S2z	S1	S2zw	S2tw	S2t	S3t	S1	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2tz	S1	S1	S2tw	S3tw	
Belagera	292	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz	
Belagera	293	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz	
Belagera	295	S3tz	S2tw	S3t	S1	S3t	S2z	S2t	S2z	S1	S2zw	S2tw	S2t	S3t	S1	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2tz	S1	S1	S2tw	S3tw	
Belagera	299	S3tz	S2tw	S3t	S1	S3t	S2z	S2t	S2z	S1	S2zw	S2tw	S2t	S3t	S1	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2tz	S1	S1	S2tw	S3tw	
Belagera	300	S3tz	S2tw	S3t	S1	S3t	S2z	S2t	S2z	S1	S2zw	S2tw	S2t	S3t	S1	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2tz	S1	S1	S2tw	S3tw	
Belagera	301	S3tz	S2tw	S3t	S1	S3t	S2z	S2t	S2z	S1	S2zw	S2tw	S2t	S3t	S1	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2tz	S1	S1	S2tw	S3tw	
Belagera	302	S3tz	S2tw	S3t	S1	S3t	S2z	S2t	S2z	S1	S2zw	S2tw	S2t	S3t	S1	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2tz	S1	S1	S2tw	S3tw	
Belagera	303	S3tz	S2tw	S3t	S1	S3t	S2z	S2t	S2z	S1	S2zw	S2tw	S2t	S3t	S1	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2tz	S1	S1	S2tw	S3tw	
Belagera	304	S3tz	S2tw	S3t	S1	S3t	S2z	S2t	S2z	S1	S2zw	S2tw	S2t	S3t	S1	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2tz	S1	S1	S2tw	S3tw	
Belagera	305(1)	S3tz	S2tw	S3t	S1	S3t	S2z	S2t	S2z	S1	S2zw	S2tw	S2t	S3t	S1	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2tz	S1	S1	S2tw	S3tw	
Belagera	305(2)	S3tz	S2tw	S3t	S1	S3t	S2z	S2t	S2z	S1	S2zw	S2tw	S2t	S3t	S1	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2tz	S1	S1	S2tw	S3tw	
Belagera	305(3)	S3tz	S2tw	S3t	S1	S3t	S2z	S2t	S2z	S1	S2zw	S2tw	S2t	S3t	S1	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2tz	S1	S1	S2tw	S3tw	
Belagera	306	S3tz	S2tw	S3t	S1	S3t	S2z	S2t	S2z	S1	S2zw	S2tw	S2t	S3t	S1	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2tz	S1	S1	S2tw	S3tw	
Belagera	307	S3tz	S2tw	S3t	S1	S3t	S2z	S2t	S2z	S1	S2zw	S2tw	S2t	S3t	S1	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2tz	S1	S1	S2tw	S3tw	
Belagera	308	S3tz	S2tw	S3t	S1	S3t	S2z	S2t	S2z	S1	S2zw	S2tw	S2t	S3t	S1	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2tz	S1	S1	S2tw	S3tw	
Belagera	309(1)	S3tz	S2tw	S3t	S1	S3t	S2z	S2t	S2z	S1	S2zw	S2tw	S2t	S3t	S1	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2tz	S1	S1	S2tw	S3tw	
Belagera	309(2)	S3tz	S2tw	S3t	S1	S3t	S2z	S2t	S2z	S1	S2zw	S2tw	S2t	S3t	S1	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2tz	S1	S1	S2tw	S3tw	
Belagera	310(1)	S3tz	S2tw	S3t	S1	S3t	S2z	S2t	S2z	S1	S2zw	S2tw	S2t	S3t	S1	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2tz	S1	S1	S2tw	S3tw	
Belagera	310(2)	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Belagera	310(3)	S3tz	S2tw	S3t	S1	S3t	S2z	S2t	S2z	S1	S2zw	S2tw	S2t	S3t	S1	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2tz	S1	S1	S2tw	S3tw	
Belagera	311	S3tz	S2tw	S3t	S1	S3t	S2z	S2t	S2z	S1	S2zw	S2tw	S2t	S3t	S1	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2tz	S1	S1	S2tw	S3tw	
Belagera	312(1)	S3tz	S2tw	S3t	S1	S3t	S2z	S2t	S2z	S1	S2zw	S2tw	S2t	S3t	S1	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2tz	S1	S1	S2tw	S3tw	
Belagera	312(2)	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Belagera	313(1)	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengalgram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry	
Belagera	313(2)	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Honagera	206	S3tz	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tz	S2t	S1	S2tw	S3tw	
Honagera	208	S3tz	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tz	S2t	S1	S2tw	S3tw	
Honagera	209	S3tz	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tz	S2t	S1	S2tw	S3tw	
Honagera	210	S3tz	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tz	S2t	S1	S2tw	S3tw	
Honagera	211	S3tz	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tz	S2t	S1	S2tw	S3tw	
Honagera	226	S3tz	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tz	S2t	S1	S2tw	S3tw	
Sutharahosalli	91	S3rz	S3tz	S2rz	S2tz	S2rz	S3tz	S3rz	S2rz	S3tz	S2rz	S2rz	S2tz	S2rz	S2z	N1tz	S3rz	S2rz	S2z	S2tz	S2z	S2z	S2z	S2z	S2rz	S2z	S2tz	S2z	S2rz	S2rz	
Sutharahosalli	92	S3rz	S3tz	S2rz	S2tz	S2rz	S3tz	S3rz	S2rz	S3tz	S2rz	S2rz	S2tz	S2rz	S2z	N1tz	S3rz	S2rz	S2z	S2tz	S2z	S2z	S2z	S2z	S2rz	S2z	S2tz	S2z	S2rz	S2rz	
Sutharahosalli	93	S3rg	S3g	S3g	S3g	S3g	S3g	S3rg	S3g	S3g	S3g	S3g	S2g	S3g	S2g	S3g	S3rg	S3g	S2g	S2g	S3g	S3g	S3g	S3g	S3g	S2g	S2g	S2g	S2g	S2g	
Sutharahosalli	94	S3rg	S3g	S3g	S3g	S3g	S3g	S3rg	S3g	S3g	S3g	S3g	S2g	S3g	S2g	S3g	S3rg	S3g	S2g	S2g	S3g	S3g	S3g	S3g	S3g	S2g	S2g	S2g	S2g	S2g	
Sutharahosalli	95	S3rz	S3tz	S2rz	S2tz	S2rz	S3tz	S3rz	S2rz	S3tz	S2rz	S2rz	S2tz	S2rz	S2z	N1tz	S3rz	S2rz	S2z	S2tz	S2z	S2z	S2z	S2z	S2rz	S2z	S2tz	S2z	S2rz	S2rz	
Sutharahosalli	96	S3rz	S2tz	S2rz	S2tz	S2rz	S3tz	S3rz	S2rz	S3tz	S2rz	S2rz	S2tz	S2rz	S1	N1tz	S3rz	S2rz	S2z	S1	S2z	S2z	S2z	S2z	S2rz	S2z	S2t	S1	S2rz	S2rz	
Sutharahosalli	97	S3rz	S2tz	S2rz	S2tz	S2rz	S3tz	S3rz	S2rz	S3tz	S2rz	S2rz	S2tz	S2rz	S1	N1tz	S3rz	S2rz	S2z	S1	S2z	S2z	S2z	S2z	S2rz	S2z	S2t	S1	S2rz	S2rz	
Sutharahosalli	98	S3rz	S2tz	S2rz	S2tz	S2rz	S3tz	S3rz	S2rz	S3tz	S2rz	S2rz	S2tz	S2rz	S1	N1tz	S3rz	S2rz	S2z	S1	S2z	S2z	S2z	S2z	S2rz	S2z	S2t	S1	S2rz	S2rz	
Sutharahosalli	99	S3rz	S2tz	S2rz	S2tz	S2rz	S3tz	S3rz	S2rz	S3tz	S2rz	S2rz	S2tz	S2rz	S1	N1tz	S3rz	S2rz	S2z	S1	S2z	S2z	S2z	S2z	S2rz	S2z	S2t	S1	S2rz	S2rz	
Sutharahosalli	100	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	

Ro-Rock outcrops

# **PART-B**

**SOCIO-ECONOMIC STATUS OF FARM HOUSEHOLDS**





## CONTENTS

1.	Findings of the socio-economic survey	1-3
2.	Introduction	5
3	Methodology	6-7
4	Salient features of the survey	9-27
5	Summary	29-33



## LIST OF TABLES

1	Households sampled for socio economic survey	9
2	Population characteristics	9
3	Age wise classification of household members	9
4	Education level of household members	10
5	Occupation of household heads	10
6	Occupation of family members	10
7	Institutional participation of household members	11
8	Type of house owned by households	11
9	Durable assets owned by households	11
10	Average value of durable assets owned by households	11
11	Farm implements owned by households	12
12	Average value of farm implements	12
13	Livestock possession by households	12
14	Average labour availability	13
15	Adequacy of hired labour	13
16	Distribution of land (ha)	13
17	Average land value (Rs./ha)	13
18	Status of bore wells	13
19	Source of irrigation	14
20	Depth of water(Avg in meters)	14
21	Irrigated area (ha)	14
22	Cropping pattern	14
23	Cropping intensity	14
24	Possession of bank account and saving	15
25	Borrowing status	15
26	Source of credit	15
27	Avg. credit borrowed	15
28	Purpose of credit borrowed from institutional sources	15
29	Purpose of credit borrowed from Non-institutional sources	16
30	Repayment status of household from institutional sources	16
31	Repayment status of household from Non-institutional sources	16

32	Opinion on institutional sources of credit	16
33	Opinion on Non-institutional sources of credit	16
34.a	Cost of cultivation of Red gram	17
34. b	Cost of cultivation of Groundnut	18
34. c	Cost of cultivation of Cotton	19
34. d	Cost of cultivation of Green gram	20
34. e	Cost of cultivation of Paddy	21
35	Adequacy of fodder	22
36	Annual gross income	22
37	Average annual expenditure	22
38	Horticultural species grown	22
39	Average additional investment capacity	23
40	Source of funds for additional investment	23
41	Marketing of the agricultural produce	23
42	Marketing channels used for sale of agricultural produce	24
43	Mode of transport of agricultural produce	24
44	Incidence of soil and water erosion problems	24
45	Interest shown towards soil testing	24
46	Soil and water conservation practices and structures	24
47	Status soil and water conservation structures	25
48	Agencies involved in the soil and water conservation structures	25
49	Usage pattern of fuel for domestic use	25
50	Source of drinking water	25
51	Source of light	26
52	Existence of sanitary toilet facility	26
53	Possession of public distribution system (PDS) card	26
54	Participation in NREGA programme	26
55	Adequacy of food items	26
56	Inadequacy of food items	27
57	Farming constraints experienced	27

**FINDINGS OF THE SOCIO-ECONOMIC SURVEY**

- ❖ *The survey was conducted in Belagiri-1 is located at North latitude 16<sup>0</sup> 50' 28.182" and 16<sup>0</sup> 48' 54.3" and East longitude 77<sup>0</sup> 14' 23.172" and 77<sup>0</sup> 11' 59.467" covering an area of about 601.86 ha coming unde Belagera, Sutharahosalli and Honagera villages of Yadagiri taluk.*
- ❖ *Socio-economic analysis of Belagiri-1 micro watersheds of Belagiri sub-watershed, Yadgiri taluk & District indicated that, out of the total sample of 34 farmers were sampled in Belagiri-1 micro-watershed among households surveyed 12 (35.29%) were marginal, 11 (32.35%) were small, 6 (17.65 %) were semi medium and 2 (5.88 %) were medium farmers. 3 landless farmers were also interviewed for the survey.*
- ❖ *The population characteristics of households indicated that, there were 118 (59.00%) men and 82 (41.00 %) were women. The average population of landless was 5.7, marginal farmers were 5, small farmers were 6.9, semi medium farmers were 6 and medium farmers were 5.5.*
- ❖ *Majority of the respondents (48.50%) were in the age group of 16-35 years.*
- ❖ *Education level of the sample households indicated that, there were 16.00 per cent illiterates, 82.50 per cent pre university education and 7.50 per cent attained graduation.*
- ❖ *About, 88.24 per cent of household heads practicing agriculture and 11.76 per cent of the household heads were engaged as agricultural labourers.*
- ❖ *Agriculture was the major occupation for 46.00 per cent of the household members.*
- ❖ *In the study area, 79.41 per cent of the households possess katcha house and 17.65 per cent possess pucca house.*
- ❖ *The durable assets owned by the households showed that, 85.29 per cent possess TV, 5.88 per cent possess mixer grinder, 102.94 per cent possess mobile phones and 26.47 per cent possess motor cycles.*
- ❖ *Farm implements owned by the households indicated that, 50.00 per cent of the households possess plough, 2.94 per cent possess tractor, 5.88 per cent possess bullock cart and 26.47 per cent possess sprayer.*
- ❖ *Regarding livestock possession by the households, 20.59 per cent possess local cow and 14.71 per cent possess buffalo.*
- ❖ *The average labour availability in the study area showed that, own labour men available in the micro watershed was 2.24, women available in the micro watershed was 1.50, hired labour (men) available was 6.18 and hired labour (women) available was 11.03.*
- ❖ *Further, 44.12 per cent of the households opined that hired labour was inadequate during the agricultural season.*

- ❖ *Out of the total land holding of the sample respondents 76.04 per cent (39.75 ha) of the area is under dry condition and the remaining 23.96 per cent area is irrigated land.*
- ❖ *There were 7.00 live bore wells and 7.00 dry bore wells among the sampled households.*
- ❖ *Bore well was the major source of irrigation for 32.35 per cent of the households.*
- ❖ *The major crops grown by sample farmers are Red gram, Groundnut, Cotton, Green gram and Paddy and cropping intensity was recorded as 100.00 per cent.*
- ❖ *Out of the sample households 97.06 percent possessed bank account and 47.06 per cent of them have savings in the account.*
- ❖ *About 100.00 per cent of the respondents borrowed credit from various sources.*
- ❖ *Among the credit borrowed by households, 21.43 per cent have borrowed loan from commercial banks and 71.43 per cent from co-operative/Grameena bank.*
- ❖ *Majority of the respondents (100.00%) have borrowed loan for agriculture purpose.*
- ❖ *Regarding the opinion on institutional sources of credit, 100.00 per cent of the households opined that credit helped to perform timely agricultural operations.*
- ❖ *The per hectare cost of cultivation for Red gram, Groundnut, Cotton, Green gram and Paddy was Rs.37413.18, 62616.52, 26530.56, 28811.74 and 66062.03 with benefit cost ratio of 1:1.95, 1: 1.70, 1: 2.10, 1: 1.90 and 1:0.80 respectively.*
- ❖ *Further, 67.65 per cent of the households opined that dry fodder was adequate and 5.88 per cent of the households have opined that the green fodder was adequate.*
- ❖ *The average annual gross income of the farmers was Rs. 170735.29 in micro-watershed, of which Rs. 78852.94 comes from agriculture.*
- ❖ *Sampled households have grown 25 horticulture trees.*
- ❖ *Households have an average investment capacity of Rs. 3441.18 for land development.*
- ❖ *Source of funds for additional investment is concerned, 2.94 per cent depends on own funds and 14.71 per cent depends on bank loan for land development activities.*
- ❖ *Regarding marketing channels, 32.35 per cent of the households have sold agricultural produce to the local/village merchants, while, 61.76 per cent have sold in regulated markets.*
- ❖ *Further, 50.00 per cent of the households have used tractor for the transport of agriculture commodity.*
- ❖ *Majority of the farmers (85.29%) have experienced soil and water erosion problems in the watershed and 79.41 per cent of the households were interested towards soil testing.*
- ❖ *Fire was the major source of fuel for domestic use for 97.06 per cent of the households and 2.94 per cent households has LPG connection.*



- ❖ *Piped supply was the major source for drinking water for 91.18 per cent of the households.*
- ❖ *Electricity was the major source of light for 100.00 per cent of the households.*
- ❖ *In the study area, 47.06 per cent of the households possess toilet facility.*
- ❖ *Regarding possession of PDS card, 97.06 per cent of the households possessed BPL card, 2.94 per cent of the household's possessed APL card.*
- ❖ *Households opined that, the requirement of cereals (91.18%), pulses (70.59%) and oilseeds (61.76%) are adequate for consumption.*
- ❖ *Farming constraints experienced by households in the micro watersheds were lower fertility status of the soil (82.35%) wild animal menace on farm field (32.35%), frequent incidence of pest and diseases (70.59%), inadequacy of irrigation water (76.47%), high cost of fertilizers and plant protection chemicals (82.35%), high rate of interest on credit (82.35%), low price for the agricultural commodities (85.29%), lack of marketing facilities in the area (76.47%), inadequate extension services (23.53%), lack of transport for safe transport of the agricultural produce to the market (76.47%).*



## INTRODUCTION

Soil and water are the two precious natural resources which are essential for crop production and existence of life on earth. Rainfed agriculture is under severe stress due to various constraints related to agriculture like uneven and erratic distribution of rainfall, indiscriminate use of fertilizers, chemicals and pesticides, adoption of improper land management practices, soil erosion, decline in soil fertility, decline in ground water resources leading to low crop productivity. The area under rainfed agriculture has to be managed effectively using the best available practices to enhance the production of food, fodder and fuel. This is possible if the land resources are characterized at each parcel of land through detailed land resource inventory using the best available techniques of remote sensing, GPS and GIS. The watershed development programs are aimed at the sustainable distribution of its resources and the process of creating and implementing plans, programs, and projects to sustain and enhance watershed functions that affect the plant, animal and human communities within a watershed boundary.

World Bank funded KWDP II, SUJALA III project was implemented in with Broad objective of demonstrating more effective watershed management through greater integration of programmes related to rain-fed agriculture, innovative and science based approaches and strengthen institutional capacities and If successful, it is expected that the systems and tools could be mainstreamed into the overall IWMP in the State of Karnataka and in time, throughout other IWMP operations in India. With this background the socio-economic survey has been carried out with following specific objectives:

1. To understand the demographic features of the households in the micro-watershed
2. To understand the extent of family labour available and additional employment opportunities available within the village.
3. To know the status of assets of households in the micro-watershed for suggesting possible improvements.
4. To study the cropping pattern, cropped area and productivity levels of different households in micro-watershed.
5. To determine the type and extent of livestock owned by different categories of HHs
6. Availability of fodder and level of livestock management.

### **Scope and importance of survey**

Survey helps in identification of different socio-economic and resource use-patterns of farmers at the Micro watershed. Household survey provides demographic features, labor force, and levels of education; land ownership and asset position (including livestock and other household assets) of surveyed households; and cropping patterns, input intensities, and average crop yields from farmers' fields. It also discusses crop utilization and the degree of commercialization of production in the areas; farmers' access to and utilization of credit from formal and informal sources; and the level of adoption and use of soil, water, and pest management technologies.

## METHODOLOGY

The description of the methods, components selected for the survey and procedures followed in conducting the baseline survey are furnished under the following heads.

### 1. Description of the study area

Yadgir District is one of the 30 districts of Karnataka state in southern India. This district was carved out from the erstwhile Gulbarga district as the 30th district of Karnataka on 10 April 2010. Yadgir town is the administrative headquarters of the district. The district comprises of 3 taluks namely, Shahapur, Yadgiri and Shorapur (There are 16 hoblies, 117 Gram Panchayats, 4 Municipalities, 8 Towns/ Urban agglomeration and 487 inhabited & 32 un-inhabited villages The district occupies an area of 5,160.88 km<sup>2</sup>).

Yadgir district is the second smallest district in the state, area wise is very rich in cultural traditions. The vast stretch of fertile black soil of the district is known for bumper red gram and jowar crops. The district is a "Daal bowl" of the state. The district is also known for cluster of cement industries and a distinct stone popularly known as "Malakheda Stone". Two main rivers, Krishna and Bhima, and a few tributaries flow in this region. Krishna and Bhima Rivers drain the district. They constitute the two major river basins of the district. Kagna and Amarja are the two sub - basins of Bhima River, which occur within the geographical area of the district

According to the 2011 census Yadgir district has a population of 1, 172,985, roughly equal to the nation of Timor-Leste or the US state of Rhode Island. This gives it a ranking of 404th in India (out of a total of 640). The district has a population density of 224 inhabitants per square kilometre (580/sq mi). Its population growth rate over the decade 2001-2011 was 22.67%. Yadgir has a sex ratio of 984 females for every 1000 males, and a literacy rate of 52.36%.

### 2. Locale of the survey and description of the micro-watershed and

The study was conducted in Belagiri-1 micro-watershed (Belagiri sub-watershed, Yadgiri taluk & District) is located at North latitude 16<sup>o</sup> 50' 28.182" and 16<sup>o</sup> 48' 54.3" and East longitude 77<sup>o</sup> 14' 23.172" and 77<sup>o</sup> 11' 59.467" covering an area of about 601.86 ha bounded by unde Belagera, Sutharahosalli and Honagera Villages.

### 3. Selection of the respondents for the study

The micro-watershed is marked with 320 square meters grids. One farmer from every alternate grid in the micro-watershed was selected for the study and interviewed for socio-economic data. Totally 34 households were interviewed for the survey.

### 4. The parameters considered for socio-economic survey of households

Two forms of data were collected from the micro-watershed which includes primary data from the farm households and secondary data about the villages under the micro-watershed jurisdiction.

The following parameters were considered for the primary data collection about the socio-economic data of the households, (1) Demographic information, (2) Farm and durable assets owned by households, (3) Livestock possession, (4) Labour availability, (5) Level of migration in the village, Land holding, (7) Cropping pattern, (8) Source of irrigation, (9) Borrowing status, (10) Cost of cultivation of major crops, (11) Economics of subsidiary activities, (12) Fodder availability, (13) Family annual income from different sources, (14) Horticulture and forestry species grown, (15) Additional investment capacity, (16) Marketing practices, (17) Status of soil and water conservation structure, (18) Access to basic needs and (19) Constraints and suggestion.

The following parameters were considered for the secondary data regarding the villages under the micro-watershed jurisdiction, (1) Number of villages in each micro-watershed jurisdiction, (2) Village wise number of households, (3) Geographical area of the villages, (4) Cultivable area including rainfed and irrigated, (5) Number and type of house in each village, (6) Human and livestock population, (7) Facilities in the village such as roads, transport facility for conveyance, drinking water supply, street light and (8) Community based organizations in the villages.

## **5. Development of interview schedule and data collection**

Taking into the consideration the objectives of the survey, an interview schedule was prepared after thorough consultation with the experts in the field of social sciences. A comprehensive interview schedule covering all the major parameters for measuring the socio-economic situation was developed.

## **6. Tools used to analyze the data**

The statistical components such as frequency and percentage were used to analyze the data.

### **Abbreviations used in the report**

- LL=Landless
- MF=Marginal Farmers
- SF=Small farmers
- SMF=Semi medium farmers
- MDF=Medium farmers
- LF=Large Farmers



## FINDINGS OF THE SURVEY

This chapter deals with systematic presentation of results of the survey. Keeping in view the objectives, the salient features of the survey are presented under the following headings.

**Households sampled for socio-economic survey:** The data on households sampled for socio economic survey in Belagiri-1 Micro watershed is presented in Table 1 and it indicated that 34 farmers were sampled in Belagiri-1 micro-watershed among households surveyed 12 (35.29%) were marginal, 11 (32.35%) were small, 6 (17.65 %) were semi medium and 2 (5.88 %) were medium farmers. 3 landless farmers were also interviewed for the survey.

**Table 1. Households sampled for socio economic survey in Belagiri-1 micro-watershed**

Sl.No.	Particulars	LL (3)		MF (12)		SF (11)		SMF (6)		MDF (2)		All (34)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Farmers	3	8.82	12	35.3	11	32.4	6	17.7	2	5.88	34	100

**Population characteristics:** The population characteristics of households sampled for socio-economic survey in Belagiri-1 Micro watershed is presented in Table 2. The data indicated that, there were 118 (59.00%) men and 82 (41.00%) were women. The average population of landless was 5.7, marginal farmers were 5, small farmers were 6.9, semi medium farmers were 6 and medium farmers were 5.5.

**Table 2. Population characteristics in Belagiri-1 micro-watershed**

Sl.No.	Particulars	LL (17)		MF (60)		SF (76)		SMF (36)		MDF (11)		All (200)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Men	8	47.1	33	55	47	62	22	61.1	8	72.7	118	59
2	Women	9	52.9	27	45	29	38	14	38.9	3	27.3	82	41
Total		17	100	60	100	76	100	36	100	11	100	200	100
Average		5.7		5.0		6.9		6.0		5.5		5.9	

**Age wise classification of population:** The age wise classification of household members in Belagiri-1 Micro watershed is presented in Table 3. The indicated that, 34 (17.00%) of population were 0-15 years of age, 97 (48.50%) were 16-35 years of age, 51(25.50%) were 36-60 years of age and 18 (9.00 %) were above 61 years of age.

**Table 3: Age wise classification of members of the household in Belagiri-1 micro-watershed**

Sl.No.	Particulars	LL (17)		MF (60)		SF (76)		SMF (36)		MDF (11)		All (200)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	0-15 years of age	9	52.9	14	23.3	8	10.5	3	8.33	0	0	34	17
2	16-35 years of age	5	29.4	26	43.3	45	59.2	16	44.44	5	45	97	48.5
3	36-60 years of age	2	11.8	16	26.7	18	23.7	11	30.56	4	36	51	25.5
4	> 61 years	1	5.88	4	6.67	5	6.58	6	16.67	2	18	18	9
Total		17	100	60	100	76	100	36	100	11	100	200	100



**Education level of household members:** Education level of household members in Belagiri-1 Micro watershed is presented in Table 4. The results indicated that, there were 16.00 per cent of illiterates, 52.00 per cent of them had primary school education, 1.00 per cent middle school education, 17.50 per cent high school education, 6.00 per cent of them had PUC education, 7.50 per cent attained graduation.

**Table 4. Education level of members of the household in Belagiri-1 micro-watershed**

Sl.No.	Particulars	LL (17)		MF (60)		SF (76)		SMF (36)		MDF (11)		All (200)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Illiterate	3	17.7	8	13.3	10	13.2	9	25	2	18.18	32	16
2	Primary School	14	82.4	33	55	38	50	13	36.1	6	54.55	104	52
3	Middle School	0	0	0	0	2	2.63	0	0	0	0	2	1
4	High School	0	0	11	18.3	13	17.1	9	25	2	18.18	35	17.5
5	PUC	0	0	3	5	7	9.21	2	5.56	0	0	12	6
6	Degree	0	0	5	8.33	6	7.89	3	8.33	1	9.09	15	7.5
Total		17	100	60	100	76	100	36	100	11	100	200	100

**Occupation of head of households:** The data regarding the occupation of the household heads in Belagiri-1 Micro watershed is presented in Table 5. The results indicate that, 88.24 per cent of households heads were practicing agriculture, 11.76 per cent of the household heads were agricultural Labour.

**Table 5: Occupation of heads of households in Belagiri-1 micro-watershed**

Sl.No.	Particulars	LL (3)		MF (12)		SF (11)		SMF (6)		MDF (2)		All (34)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Agriculture	0	0	11	92	11	100	6	100	2	100	30	88.24
2	Agricultural Labour	2	67	1	8.3	1	9.09	0	0	0	0	4	11.76
3	General Labour	1	33	1	8.3	0	0	0	0	0	0	2	5.88
Total		3	100	13	100	12	100	6	100	2	100	36	100

**Occupation of the members of the household:** The data regarding the occupation of the household members in Belagiri-1 Micro watershed is presented in Table 6. The results indicate that, agriculture was the major occupation for 46.00 per cent of the household members, 3.50 per cent were agricultural labour, 3.00 per cent were general labour, 0.50 per cent were working in government sector, 25.00 per cent were working in pursuing education, 20.00 per cent were involved as housewife.

**Table 6: Occupation of members of the household in Belagiri-1 micro-watershed**

Sl.No.	Particulars	LL (17)		MF (60)		SF (76)		SMF (36)		MDF (11)		All (200)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Agriculture	0	0	28	46.7	33	43.42	22	61.11	9	82	92	46
2	Agricultural Labour	3	17.7	1	1.67	3	3.95	0	0	0	0	7	3.5
3	General Labour	2	11.8	2	3.33	2	2.63	0	0	0	0	6	3
4	Government Service	0	0	0	0	0	0	0	0	1	9.1	1	0.5
5	Private Service	0	0	2	3.33	2	2.63	0	0	0	0	4	2
6	Student	9	52.9	19	31.7	14	18.42	8	22.22	0	0	50	25
7	Housewife	3	17.7	8	13.3	22	28.95	6	16.67	1	9.1	40	20
Total		17	100	60	100	76	100	36	100	11	100	200	100

**Institutional Participation of household members:** The data regarding the institutional participation of the household members in Belagiri-1 Micro watershed is presented in Table 7. The results show that, out of the total family members in the households 1 per cent of them are participating in NGOs.

**Table 7: Institutional Participation of household member in Belagiri-1 micro-watershed**

Sl.No.	Particulars	LL (17)		MF (60)		SF (76)		SMF (36)		MDF (11)		All (200)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	No Participation	16	94	59	98.3	76	100	36	100	11	100	198	99
2	NGOs	1	5.9	1	1.67	0	0	0	0	0	0	2	1
Total		17	100	60	100	76	100	36	100	11	100	200	100

**Type of house owned:** The data regarding the type of house owned by the households in Belagiri-1 Micro watershed is presented in Table 8. The results indicate that, 2.94 percent possess thatched house, 79.41 per cent of the households possess katcha house, 17.65 per cent possess pacca house.

**Table 8. Type of house owned by households in Belagiri-1 micro-watershed**

Sl.No.	Particulars	LL (3)		MF (12)		SF (11)		SMF (6)		MDF (2)		All (34)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Thatched	1	33	0	0	0	0	0	0	0	0	1	2.94
2	Katcha	2	67	10	83	9	81.82	5	83.3	1	50	27	79.41
3	Pucca/RCC	0	0	2	17	2	18.18	1	16.7	1	50	6	17.65
Total		3	100	12	100	11	100	6	100	2	100	34	100

**Durable assets owned by the households:** The data regarding the Durable Assets owned by the households in Belagiri-1 Micro watershed is presented in Table 9. The results shows that, 85.29 per cent possess TV, 5.88 per cent possess mixer grinder, 26.47 per cent possess motor cycle, 102.94 per cent possess mobile phones.

**Table 9. Durable assets owned by households in Belagiri-1 micro-watershed**

Sl.No.	Particulars	LL (3)		MF (12)		SF (11)		SMF (6)		MDF (2)		All (34)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Television	1	33	11	92	9	81.8	6	100	2	100	29	85.29
2	Mixer/Grinder	0	0	0	0	1	9.09	1	17	0	0	2	5.88
3	Motor Cycle	0	0	3	25	3	27.3	2	33	1	50	9	26.47
4	Mobile Phone	3	100	12	100	11	100	7	117	2	100	35	102.94

**Average value of durable assets:** The data regarding the average value of durable assets owned by the households in Belagiri-1 Micro watershed is presented in Table 10. The result shows that, the average value of television was Rs.6517.00, mixer grinder was Rs.1600.00, motor cycle was Rs. 59444.00, mobile phone was Rs.2408.00.

**Table 10. Average value of durable assets owned in Belagiri-1 micro-watershed**

Average Value (Rs.)

Sl.No.	Particulars	LL (3)	MF (12)	SF (11)	SMF (6)	MDF (2)	All (34)
1	Television	7000	6727	5666	7166	7000	6517
2	Mixer/Grinder	0	0	2000	1200	0	1600
3	Motor Cycle	0	56666	61666	60000	60000	59444
4	Mobile Phone	2000	1778	3071	2000	2500	2408

**Farm implements owned:** The data regarding the farm implements owned by the households in Belagiri-1 Micro watershed is presented in Table 11. About 5.88 per cent of the households possess Bullock Cart, 50.00 per cent possess plough and 29.41 per cent possess Seed/Fertilizer Drill and Sprinkler, 26.47 per cent possess Sprayer, 29.41 per cent possess Weeder, 2.94 per cent possess tractor.

**Table 11. Farm implements owned in Belagiri-1 micro-watershed**

Sl.No.	Particulars	LL (3)		MF (12)		SF (11)		SMF (6)		MDF (2)		All (34)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Bullock Cart	0	0	0	0	1	9.09	1	16.7	0	0	2	5.88
2	Plough	0	0	7	58.3	5	45.45	3	50	2	100	17	50
3	Seed/Fertilizer Drill	0	0	4	33.3	3	27.27	1	16.7	2	100	10	29.41
4	Tractor	0	0	0	0	0	0	1	16.7	0	0	1	2.94
5	Sprayer	0	0	4	33.3	2	18.18	2	33.3	1	50	9	26.47
6	Weeder	0	0	6	50	3	27.27	1	16.7	0	0	10	29.41
7	Blank	3	100	5	41.7	6	54.55	3	50	0	0	17	50

**Average value of farm implements:** The data regarding the average value of farm Implements owned by the households in Belagiri-1 Micro watershed is presented in Table 12. The results show that the average value of plough was Rs.4688.00, bullock Cart was Rs.18500.00, seed/fertilizer drill was Rs.2477.00, sprayer and weeder was Rs.130.00 and tractor was Rs. 800000.

**Table 12. Average value of farm implements in Belagiri-1 micro-watershed**

Sl.No.	Particulars	Average Value (Rs.)					
		LL (3)	MF (12)	SF (11)	SMF (6)	MDF (2)	All (34)
1	Bullock Cart	0	0	15000	22000	0	18500
2	Plough	0	5357	4040	5000	3500	4688
3	Seed/Fertilizer Drill	0	3250	3166	3000	3750	3300
4	Tractor	0	0	0	800000	0	800000
5	Sprayer	0	2525	2350	2500	2500	2477
6	Weeder	0	132	125	150	0	130

**Livestock possession by the households:** The data regarding the Livestock possession by the households in Belagiri-1 Micro watershed is presented in Table 13. The indicate that, 52.94 per cent of the households possess bullocks, 20.59 per cent possess local cow, 14.71 per cent possess buffalo, 2.94 per cent possess sheep, 5.88 per cent were poultry birds.

**Table 13. Livestock possession by households in Belagiri-1 micro-watershed**

Sl.No.	Particulars	LL (3)		MF (12)		SF (11)		SMF (6)		MDF (2)		All (34)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Bullock	0	0	6	50	6	54.55	4	67	2	100	18	52.94
2	Local cow	0	0	1	8.3	3	27.27	2	33	1	50	7	20.59
3	Buffalo	0	0	3	25	0	0	2	33	0	0	5	14.71
4	Sheep	0	0	1	8.3	0	0	0	0	0	0	1	2.94
5	Poultry birds	0	0	2	17	0	0	0	0	0	0	2	5.88
6	blank	3	100	5	42	5	45.45	3	50	0	0	16	47.06

**Average Labour availability:** The data regarding the average labour availability in Belagiri-1 Micro watershed is presented in Table 14. The indicated that, own labour men

available in the micro watershed was 2.24, women available in the micro watershed was 1.50, hired labour (men) available was 6.18 and hired labour (women) available was 11.03.

**Table 14. Average labour availability in Belagiri-1 micro-watershed**

Sl.No.	Particulars	LL (3)	MF (12)	SF (11)	SMF (6)	MDF (2)	All (34)
		N	N	N	N	N	N
1	Hired labour Female	0	10.4	13.18	13.33	12.5	11.03
2	Own Labour Female	0	1.17	2.09	2	1	1.5
3	Own labour Male	0	1.83	2.82	2.83	3	2.24
4	Hired labour Male	0	5	7.73	8.33	7.5	6.18

**Adequacy of hired labour:** The data regarding the adequacy of hired labour in Belagiri-1 Micro watershed is presented in Table 15. The results indicate that, 55.88 per cent of the household opined that hired labour was adequate, 44.12 per cent of the household opined that hired labour was Inadequate.

**Table 15. Adequacy of hired labour in Belagiri-1 micro-watershed**

Sl.No.	Particulars	LL (3)		MF (12)		SF (11)		SMF (6)		MDF (2)		All (34)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Adequate	3	100	9	75	4	36.4	2	33.3	1	50	19	55.9
2	Inadequate	0	0	3	25	7	63.6	4	66.7	1	50	15	44.1

**Distribution of land (ha):** The data regarding the distribution of land (ha) in Belagiri-1 Micro watershed is presented in Table 16. The results indicate that, 30.23 ha (76.04%) of dry land and 9.53 ha (23.96 %) of irrigated land.

**Table 16. Distribution of land (ha) in Belagiri-1 micro-watershed**

Sl.No.	Particulars	LL (3)		MF (12)		SF (11)		SMF (6)		MDF (2)		All (34)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Dry	0	0	8.36	100	12.85	85.14	9.02	73.59	0	0	30.23	76.04
2	Irrigated	0	0	0	0	2.24	14.86	3.24	26.41	4.05	100	9.53	23.96
Total		0	100	8.36	100	15.09	100	12.26	100	4.05	100	39.75	100

**Average value of land (ha):** The data regarding the average land value (Rs./ha) in Belagiri-1 Micro watershed is presented in Table 17. The results show that the average value of dry land was Rs.459673.32 and the average value of irrigated land was Rs.503653.36.

**Table 17. Average value of land (ha) in Belagiri-1 micro-watershed**

Sl.No.	Particulars	LL (3)	MF (12)	SF (11)	SMF (6)	MDF (2)	All (34)
		N	N	N	N	N	N
1	Dry	0	741598.1	412315	265948.9	0	459673.3
2	Irrigated	0	0	891696.8	432250	345800	503653.4

**Table 18. Status of bore wells in Belagiri-1 micro-watershed**

Sl.No.	Particulars	LL (3)	MF (12)	SF (11)	SMF (6)	MDF (2)	All (34)
		N	N	N	N	N	N
1	De-functioning	0	0	1	2	4	7
2	Functioning	0	0	1	2	4	7

**Status of bore wells:** The data regarding the status of bore wells in Belagiri-1 Micro watershed is presented in Table 18. The results indicate that, there were 7 De-functioning

bore wells and 7 functioning bore wells among the sampled households in micro watershed.

**Source of irrigation:** The data regarding the source of irrigation in Belagiri-1 Micro watershed is presented in Table 19. The results show that bore well for 20.59 per cent of the households.

**Table 19. Source of irrigation in Belagiri-1 micro-watershed**

Sl.No.	Particulars	LL (3)		MF (12)		SF (11)		SMF (6)		MDF (2)		All (34)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Bore Well	0	0	0	0	1	9.09	2	33.3	4	200	7	20.59
4	Tank	0	0	0	0	4	36.36	0	0	0	0	4	11.76

**Depth of water (Avg. In meters):** The data regarding the depth of water in Belagiri-1 Micro watershed is presented in Table 20. The results revealed that, the depth of bore well was 8.38 meter.

**Table 20. Depth of water (Avg. In meters) in Belagiri-1 micro-watershed**

Sl.No.	Particulars	LL (3)	MF (12)	SF (11)	SMF (6)	MDF (2)	All (34)
		N	N	N	N	N	N
1	Bore Well	0	0	8.73	12.7	56.39	8.38

**Irrigated Area (ha):** The data regarding the irrigated area (ha) in Belagiri-1 Micro watershed is presented in Table 21. The results indicate that, the availability of irrigation water was used for kharif crops was 7.91 ha.

**Table 21. Irrigated Area (ha) in Belagiri-1 micro-watershed**

Sl.No.	Particulars	LL (3)	MF (12)	SF (11)	SMF (6)	MDF (2)	All (34)
1	Kharif	0	0	2.24	1.62	4.05	7.91
	Total	0	0	2.24	1.62	4.05	7.91

**Cropping pattern:** The data regarding the cropping pattern in Belagiri-1 Micro watershed is presented in Table 22. The results indicate that, farmers have grown Red gram (26.26 ha), Cotton (3.79 ha), Groundnut (2.54 ha), Green gram (2.02 ha), Paddy (2.02 ha) and Jowar (1.26 ha).

**Table 22. Cropping pattern in Belagiri-1 micro-watershed**

Sl.No.	Particulars	LL (3)	MF (12)	SF (11)	SMF (6)	MDF (2)	All (34)
1	Kharif - Red gram	0	5.67	9.54	9.02	2.02	26.26
2	Kharif - Cotton	0	0	1.76	0	2.02	3.79
3	Kharif - Groundnut	0	0	0.92	1.62	0	2.54
4	Kharif - Green gram	0	0.81	1.21	0	0	2.02
5	Kharif - Paddy	0	0	0.4	1.62	0	2.02
6	Kharif - Jowar	0	1.88	0	0	0	1.88
7	Rabi - Jowar	0	0	1.26	0	0	1.26

**Cropping intensity:** The data regarding the cropping intensity in Belagiri-1 Micro watershed is presented in Table 23. The results indicate that, the cropping intensity was 100.00 per cent.

**Table 23. Cropping intensity (%) in Belagiri-1 micro-watershed**

Sl.No.	Particulars	LL (3)	MF (12)	SF (11)	SMF (6)	MDF (2)	All (34)
1	Cropping Intensity	0	100	100	100	100	100

**Possession of bank account and savings:** The data regarding the possession of bank account and saving in Belagiri-1 micro-watershed is presented in Table 24. The results indicate that, 97.06 cent of the households posses bank account and 47.06 per cent of them have savings.

**Table 24. Possession of Bank account and savings in Belagiri-1 micro-watershed**

Sl.No.	Particulars	LL (3)		MF (12)		SF (11)		SMF (6)		MDF (2)		All (34)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Account	2	66.67	12	100	11	100	6	100	2	100	33	97.06
2	Savings	0	0	3	25	8	72.73	5	83.33	0	0	16	47.06

**Borrowing status:** The data regarding the borrowing status in Belagiri-1 micro-watershed is presented in Table 25. The results indicate that, 100.00 percent of the sample farmers have borrowed credit from different sources.

**Table 25. Borrowing status in Belagiri-1 micro-watershed**

Sl.No.	Particulars	LL (3)		MF (12)		SF (11)		SMF (6)		MDF (2)		All (34)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Credit Availed	3	100	12	100	11	100	6	100	2	100	34	100

**Source of credit:** The data regarding the source of credit availed by households in Belagiri-1 micro-watershed is presented in Table 26. The results show that, 21.43 per cent have borrowed loan from commercial banks and 7.14 per cent have borrowed loan from Cooperative bank, 71.43 per cent have borrowed loan from Grameena Bank, 57.14 per cent have borrowed loan from money lender.

**Table 26. Source of credit borrowed by households in Belagiri-1 micro-watershed**

Sl.No.	Particulars	LL (0)		MF (2)		SF (8)		SMF (4)		MDF (0)		All (14)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Commercial Bank	0	0	0	0	2	25	1	25	0	0	3	21.43
2	Cooperative Bank	0	0	0	0	1	12.5	0	0	0	0	1	7.143
3	Grameena Bank	0	0	2	100	5	62.5	3	75	0	0	10	71.43
4	Money Lender	0	0	2	100	3	37.5	2	50	0	0	8	57.14

**Avg. Credit amount:** The data regarding the avg. Credit amount in Belagiri-1 micro-watershed is presented in Table 27. The results show that, farmers have borrowed Avg. Credit of Rs.24000.00 from different sources.

**Table 27. Avg. Credit amount in Belagiri-1 micro-watershed**

Sl.No.	Particulars	LL (0)		MF (2)		SF (8)		SMF (4)		MDF (0)		All (14)	
		N	N	N	N	N	N	N	N	N	N		
1	Average Credit	0	12500	45800	13700	0	24000						

**Purpose of credit borrowed (institutional Source):** The results indicate (Table 28) that, 100.00 per cent of the households have borrowed loan for agriculture.

**Table 28. Purpose of credit borrowed (institutional Source) by households in Belagiri-1 micro-watershed**

SN	Particulars	LL (0)		MF (2)		SF (8)		SMF (4)		MDF (0)		All (14)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Agriculture production	0	0	2	100	8	100	4	100	0	0	14	100

**Purpose of credit borrowed (Private Source):** The data regarding the purpose of credit borrowed – Private Source in Belagiri-1 micro-watershed is presented in Table 29. The results indicate that, 100.00 per cent of the households have borrowed loan for agriculture.

**Table 29. Purpose of credit borrowed (Private Source) by households in Belagiri-1 micro-watershed**

Sl. No.	Particulars	LL (0)		MF (3)		SF (3)		SMF (2)		All (8)	
		N	%	N	%	N	%	N	%	N	%
1	Agriculture production	0	0	3	100	3	100	2	100	8	100

**Repayment status of household (institutional Source):** The results indicate (Table 30) that, 14.29 per cent of the households have partially paid, 85.71 per cent have unpaid.

**Table 30. Repayment status of household (institutional Source) in Belagiri-1 micro-watershed**

Sl.No.	Particulars	LL (0)		MF (2)		SF (8)		SMF (4)		All (14)	
		N	%	N	%	N	%	N	%	N	%
1	Partially paid	0	0	0	0	0	0	2	50	2	14.29
2	Un paid	0	0	2	100	8	100	2	50	12	85.71

**Repayment status of household (Private Source):** The data regarding the repayment status of credit borrowed from private sources by households in Belagiri-1 micro watershed is presented in Table 31. The results indicate that, 100 per cent of the households have unpaid.

**Table 31. Repayment status of household (Private Source) in Belagiri-1 micro-watershed**

Sl.No.	Particulars	LL (0)		MF (3)		SF (3)		SMF (2)		MDF (0)		All (8)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Un paid	0	0	3	100	3	100	2	100	0	0	8	100

**Opinion regarding institutional sources of credit:** The results indicate (Table 32) that, 100.00 per cent of the households opined that credit helped to perform timely agricultural operations.

**Table 32. Opinion regarding institutional sources of credit in Belagiri-1 micro-watershed**

Sl. No.	Particulars	MF (2)		SF (8)		SMF (4)		All (14)	
		N	%	N	%	N	%	N	%
1	Helped to perform timely agricultural operations	2	100	8	100	4	100	14	100

**Opinion regarding Non- institutional sources of credit:** The results indicate (Table 33) that, 100.00 per cent of the households opined that credit helped to perform timely agricultural operations.

**Table 33. Opinion regarding Non- institutional sources of credit in Belagiri-1 micro-watershed**

Sl. No.	Particulars	LL (0)		MF (3)		SF (3)		SMF (2)		MDF (0)		All (8)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Helped to perform timely agricultural operations	0	0	3	100	3	100	2	100	0	0	8	100



**Cost of Cultivation of Red gram:** The data regarding the cost of cultivation (Rs/ha) of Red gram in Belagiri-1 micro watershed is presented in Table 34.a. The results indicate that, the total cost of cultivation (Rs/ha) for Red gram was Rs. 37413.18. The gross income realized by the farmers was Rs. 72967.10. The net income from Red gram cultivation was Rs.35553.93, thus the benefit cost ratio was found to be 1:1.95.

**Table 34(a). Cost of Cultivation of Red gram in Belagiri-1 micro-watershed**

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
<b>I</b>	<b>Cost A1</b>				
1	Hired Human Labour	Man days	57.98	12500.14	33.41
2	Bullock	Pairs/day	0.82	475.35	1.27
3	Tractor	Hours	4.69	3676.23	9.83
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	10.33	883.5	2.36
7	FYM	Quintal	1.5	3131.95	8.37
8	Fertilizer + micronutrients	Quintal	3.85	3182.64	8.51
9	Pesticides (PPC)	Kgs / liters	1.98	1664.73	4.45
10	Irrigation	Number	2.67	0	0
11	Repairs		0	20	0.05
12	Msc. Charges (Marketing costs etc)		0	122.5	0.33
13	Depreciation charges		0	77.14	0.21
14	Land revenue and Taxes		0	0.82	0
<b>II</b>	<b>Cost B1</b>				
16	Interest on working capital			1063.66	2.84
17	<b>Cost B1 = (Cost A1 + sum of 15 and 16)</b>			26798.66	71.63
<b>III</b>	<b>Cost B2</b>				
18	Rental Value of Land			288.33	0.77
19	<b>Cost B2 = (Cost B1 + Rental value)</b>			27087	72.4
<b>IV</b>	<b>Cost C1</b>				
20	Family Human Labour		27.43	6923.98	18.51
21	<b>Cost C1 = (Cost B2 + Family Labour)</b>			34010.98	90.91
<b>V</b>	<b>Cost C2</b>				
22	Risk Premium			1	0
23	<b>Cost C2 = (Cost C1 + Risk Premium)</b>			34011.98	90.91
<b>VI</b>	<b>Cost C3</b>				
24	Managerial Cost			3401.2	9.09
25	<b>Cost C3 = (Cost C2 + Managerial Cost)</b>			37413.18	100
<b>VII</b>	<b>Economics of the Crop</b>				
	Main Product	a) Main Product (q)		14.86	71547.99
		b) Main Crop Sales Price (Rs.)			4815
a.	By Product	e) Main Product (q)		1.62	1419.12
		f) Main Crop Sales Price (Rs.)			875
b.	Gross Income (Rs.)			72967.1	
c.	Net Income (Rs.)			35553.93	
d.	Cost per Quintal (Rs./q.)			2517.81	
e.	Benefit Cost Ratio (BC Ratio)			1:1.95	

**Cost of Cultivation of Groundnut:** The data regarding the cost of cultivation (Rs/ha) of Groundnut in Belagiri-1 micro watershed is presented in Table 34.b. The results indicate that, the total cost of cultivation (Rs/ha) for Groundnut was Rs. 62616.52. The gross income realized by the farmers was Rs. 103715.52. The net income from Groundnut cultivation was Rs.41099.00, thus the benefit cost ratio was found to be 1:1.70.

**Table 34(b). Cost of Cultivation of Groundnut in Belagiri-1 micro-watershed**

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
<b>I</b>	<b>Cost A1</b>				
1	Hired Human Labour	Man days	60.27	13228.65	21.13
2	Bullock	Pairs/day	1.24	864.5	1.38
3	Tractor	Hours	2.79	2234.97	3.57
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	205.11	14357.56	22.93
7	FYM	Quintal	2.01	5962.14	9.52
8	Fertilizer + micronutrients	Quintal	5.5	4361.74	6.97
9	Pesticides (PPC)	Kgs / liters	2.01	1905.55	3.04
10	Irrigation	Number	6.75	0	0
11	Repairs		0	200	0.32
12	Msc. Charges (Marketing costs etc)		0	3250	5.19
13	Depreciation charges		0	180.94	0.29
14	Land revenue and Taxes		0	8.23	0.01
<b>II</b>	<b>Cost B1</b>				
16	Interest on working capital			3191.64	5.1
17	<b>Cost B1 = (Cost A1 + sum of 15 and 16)</b>			49745.9	79.45
<b>III</b>	<b>Cost B2</b>				
18	Rental Value of Land			333.33	0.53
19	<b>Cost B2 = (Cost B1 + Rental value)</b>			50079.24	79.98
<b>IV</b>	<b>Cost C1</b>				
20	Family Human Labour		25.8	6834.87	10.92
21	<b>Cost C1 = (Cost B2 + Family Labour)</b>			56914.1	90.89
<b>V</b>	<b>Cost C2</b>				
22	Risk Premium			10	0.02
23	<b>Cost C2 = (Cost C1 + Risk Premium)</b>			56924.1	90.91
<b>VI</b>	<b>Cost C3</b>				
24	Managerial Cost			5692.41	9.09
25	<b>Cost C3 = (Cost C2 + Managerial Cost)</b>			62616.52	100
<b>VII</b>	<b>Economics of the Crop</b>				
a.	Main Product	a) Main Product (q)		20.51	102553.97
		b) Main Crop Sales Price (Rs.)			5000
	By Product	e) Main Product (q)		1.16	1161.55
		f) Main Crop Sales Price (Rs.)			1000
b.	Gross Income (Rs.)				103715.52
c.	Net Income (Rs.)				41099
d.	Cost per Quintal (Rs./q.)				3052.86
e.	Benefit Cost Ratio (BC Ratio)				1:1.7

**Cost of Cultivation of Cotton:** The data regarding the cost of cultivation (Rs/ha) of Cotton in Belagiri-1 micro watershed is presented in Table 34.c. The results indicate, the total cost of cultivation (Rs/ha) for Cotton was Rs.26530.56. The gross income realized by the farmers was Rs. 56176.89. The net income from Cotton cultivation was Rs. 29646.33, thus the benefit cost ratio was found to be 1:2.10.

**Table 34(c). Cost of Cultivation of Cotton in Belagiri-1 micro-watershed**

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
<b>I</b>	<b>Cost A1</b>				
1	Hired Human Labour	Man days	42.92	7958.08	30
2	Bullock	Pairs/day	1.35	942.01	3.55
3	Tractor	Hours	1.14	908.51	3.42
4	Machinery	Hours	0	0	0
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	3.22	3866.83	14.57
6	Seed Inter Crop	Kgs.	0	0	0
7	FYM	Quintal	0	0	0
8	Fertilizer + micronutrients	Quintal	3.11	2621.46	9.88
9	Pesticides (PPC)	Kgs / liters	1.59	1592.72	6
10	Irrigation	Number	1.48	0	0
11	Repairs		0	100	0.38
12	Misc. Charges (Marketing costs etc)		0	2500	9.42
13	Depreciation charges		0	187.29	0.71
14	Land revenue and Taxes		0	4.12	0.02
<b>II</b>	<b>Cost B1</b>				
16	Interest on working capital			970.32	3.66
17	<b>Cost B1 = (Cost A1 + sum of 15 and 16)</b>			21651.35	81.61
<b>III</b>	<b>Cost B2</b>				
18	Rental Value of Land			308.33	1.16
19	<b>Cost B2 = (Cost B1 + Rental value)</b>			21959.68	82.77
<b>IV</b>	<b>Cost C1</b>				
20	Family Human Labour		8.17	2154.01	8.12
21	<b>Cost C1 = (Cost B2 + Family Labour)</b>			24113.69	90.89
<b>V</b>	<b>Cost C2</b>				
22	Risk Premium			5	0.02
23	<b>Cost C2 = (Cost C1 + Risk Premium)</b>			24118.69	90.91
<b>VI</b>	<b>Cost C3</b>				
24	Managerial Cost			2411.87	9.09
25	<b>Cost C3 = (Cost C2 + Managerial Cost)</b>			26530.56	100
<b>VII</b>	<b>Economics of the Crop</b>				
a.	Main Product	a) Main Product (q)	11.95	56176.89	
		b) Main Crop Sales Price (Rs.)		4700	
b.	Gross Income (Rs.)			56176.89	
c.	Net Income (Rs.)			29646.33	
d.	Cost per Quintal (Rs./q.)			2219.66	
e.	Benefit Cost Ratio (BC Ratio)			1:2.1	

**Cost of Cultivation of Green gram:** The data regarding the cost of cultivation (Rs/ha) of Green gram in Belagiri-1 micro watershed is presented in Table 34.d. The results indicate that, the total cost of cultivation (Rs/ha) for Green gram was Rs. 28811.74. The gross income realized by the farmers was Rs.55575.00. The net income from Green gram cultivation was Rs. 26763.26, thus the benefit cost ratio was found to be 1:1.90.

**Table 34(d). Cost of Cultivation of Green gram in Belagiri-1 micro-watershed**

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
<b>I</b>	<b>Cost A1</b>				
1	Hired Human Labour	Man days	57.22	11732.5	40.72
2	Bullock	Pairs/day	0.62	432.25	1.5
3	Tractor	Hours	3.29	2634.67	9.14
4	Machinery	Hours	0	0	0
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	7.41	518.7	1.8
6	Seed Inter Crop	Kgs.	0	0	0
7	FYM	Quintal	0.82	1235	4.29
8	Fertilizer + micronutrients	Quintal	3.09	2305.33	8
9	Pesticides (PPC)	Kgs / liters	1.44	1440.83	5
12	Msc. Charges (Marketing costs etc)		0	0	0
13	Depreciation charges		0	102.51	0.36
14	Land revenue and Taxes		0	0	0
<b>II</b>	<b>Cost B1</b>				
16	Interest on working capital			659.98	2.29
17	<b>Cost B1 = (Cost A1 + sum of 15 and 16)</b>			21061.78	73.1
<b>III</b>	<b>Cost B2</b>				
18	Rental Value of Land			283.33	0.98
19	<b>Cost B2 = (Cost B1 + Rental value)</b>			21345.11	74.08
<b>IV</b>	<b>Cost C1</b>				
20	Family Human Labour		21.41	4847.38	16.82
21	<b>Cost C1 = (Cost B2 + Family Labour)</b>			26192.49	90.91
<b>V</b>	<b>Cost C2</b>				
22	Risk Premium			0	0
23	<b>Cost C2 = (Cost C1 + Risk Premium)</b>			26192.49	90.91
<b>VI</b>	<b>Cost C3</b>				
24	Managerial Cost			2619.25	9.09
25	<b>Cost C3 = (Cost C2 + Managerial Cost)</b>			28811.74	100
<b>VII</b>	<b>Economics of the Crop</b>				
a.	Main Product	a) Main Product (q)		11.12	55575
		b) Main Crop Sales Price (Rs.)			5000
b.	Gross Income (Rs.)				55575
c.	Net Income (Rs.)				26763.26
d.	Cost per Quintal (Rs./q.)				2592.15
e.	Benefit Cost Ratio (BC Ratio)				1:1.9

**Cost of Cultivation of Paddy:** The data regarding the cost of cultivation (Rs/ha) of Paddy in Belagiri-1 micro watershed is presented in Table 34.e. The results indicate that, the total cost of cultivation (Rs/ha) for Paddy was Rs.66062.03. The gross income realized by the farmers was Rs. 52101.56. The net income from Paddy cultivation was Rs. -13960.47, thus the benefit cost ratio was found to be 1:0.80.

**Table 34(e). Cost of Cultivation of Paddy in Belagiri-1 micro-watershed**

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
<b>I</b>	<b>Cost A1</b>				
1	Hired Human Labour	Man days	87.99	20254	30.66
2	Bullock	Pairs/day	4.32	3211	4.86
3	Tractor	Hours	5.56	4569.5	6.92
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	61.75	3643.25	5.51
7	FYM	Quintal	1.24	1235	1.87
8	Fertilizer + micronutrients	Quintal	4.01	3241.88	4.91
9	Pesticides (PPC)	Kgs / liters	3.4	3396.25	5.14
10	Irrigation	Number	19.76	0	0
11	Repairs		0	200	0.3
12	Msc. Charges (Marketing costs etc)		0	1500	2.27
13	Depreciation charges		0	5153.04	7.8
14	Land revenue and Taxes		0	8.23	0.01
<b>II</b>	<b>Cost B1</b>				
16	Interest on working capital			1383.17	2.09
17	<b>Cost B1 = (Cost A1 + sum of 15 and 16)</b>			47795.31	72.35
<b>III</b>	<b>Cost B2</b>				
18	Rental Value of Land			333.33	0.5
19	<b>Cost B2 = (Cost B1 + Rental value)</b>			48128.64	72.85
<b>IV</b>	<b>Cost C1</b>				
20	Family Human Labour		49.71	11917.75	18.04
21	<b>Cost C1 = (Cost B2 + Family Labour)</b>			60046.39	90.89
<b>V</b>	<b>Cost C2</b>				
22	Risk Premium			10	0.02
23	<b>Cost C2 = (Cost C1 + Risk Premium)</b>			60056.39	90.91
<b>VI</b>	<b>Cost C3</b>				
24	Managerial Cost			6005.64	9.09
25	<b>Cost C3 = (Cost C2 + Managerial Cost)</b>			66062.03	100
<b>VII</b>	<b>Economics of the Crop</b>				
a.	Main Product	a) Main Product (q)		33.96	48396.56
		b) Main Crop Sales Price (Rs.)			1425
	By Product	e) Main Product (q)		3.71	3705
		f) Main Crop Sales Price (Rs.)			1000
b.	Gross Income (Rs.)			52101.56	
c.	Net Income (Rs.)			-13960.47	
d.	Cost per Quintal (Rs./q.)			1945.15	
e.	Benefit Cost Ratio (BC Ratio)			1:0.8	

**Adequacy of fodder:** The data regarding the adequacy of fodder in Belagiri-1 Micro watershed is presented in Table 35. The results indicate that, 67.65 per cent of the households opined that dry fodder was adequate. With respect to green fodder availability, 5.88 percent of them opined it was sufficient.

**Table 35. Adequacy of fodder in Belagiri-1 micro-watershed**

Sl.No.	Particulars	LL (3)		MF (12)		SF (11)		SMF (6)		MDF (2)		All (34)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Adequate-Dry Fodder	0	0	7	58.33	9	81.82	5	83.3	2	100	23	67.65
2	Adequate-Green Fodder	0	0	1	8.33	1	9.09	0	0	0	0	2	5.88

**Average annual gross income:** The data regarding the annual gross income in Belagiri-1 Micro watershed is presented in Table 36. The results indicate that, the farmers have annual gross income of Rs. 170735.29 in micro-watershed, of which Rs. 78852.94 is from agriculture itself.

**Table 36. Average annual gross income in Belagiri-1 micro-watershed**

Sl.No.	Particulars	LL (3)	MF (12)	SF (11)	SMF (6)	MDF (2)	All (34)
		Rs.	Rs.	Rs.	Rs.	Rs.	Rs.
1	Service/salary	0	23833.3	11818.2	0	60000	15764.7
2	Business	0	0	0	20000	0	3529.41
3	Wage	76333.3	48333.3	77818.2	81333.3	107500	69647.1
4	Agriculture	0	37533.3	87963.6	165500	135000	78852.9
5	Dairy Farm	0	3916.67	1363.64	5000	4000	2941.18
Income(Rs.)		76333.3	113617	178964	271833	306500	170735

**Average annual Expenditure:** The data regarding the average annual expenditure in Belagiri-1 Micro watershed is presented in Table 37. The results indicate that, the farmers have annual gross expenditure of Rs. 609365.15 in micro-watershed, of which Rs. 36867.65 is from agriculture itself.

**Table 37. Average annual Expenditure in Belagiri-1 micro-watershed**

Sl.No.	Particulars	LL (3)	MF (12)	SF (11)	SMF (6)	MDF (2)	All (34)
		Rs.	Rs.	Rs.	Rs.	Rs.	Rs.
1	Service/salary	0	60000	25000	0	60000	6764.71
2	Business	0	0	0	53000	0	1558.82
3	Wage	32666.7	21090.9	34500	49600	52500	30235.3
4	Agriculture	0	19545.5	42545.5	73416.7	65000	36867.7
5	Dairy Farm	0	11500	4000	3000	2000	1029.41
Total		32666.7	112136	106045	179017	179500	609365

**Horticulture species grown:** The data regarding horticulture species grown in Belagiri-1 Micro watershed is presented in Table 38. The results indicate that, the total number of horticultural trees grown (both field and backyard) by the sampled households were clustered apple (25).

**Table 38. Horticulture species grown in Belagiri-1 micro-watershed**

Sl.No.	Particulars	LL (3)		MF (12)		SF (11)		SMF (6)		MDF (2)		All (34)	
		F	B	F	B	F	B	F	B	F	B	F	B
1	Custard apple	0	0	10	0	15	0	0	0	0	0	25	0

\*F= Field B=Back Yard

**Average additional investment capacity:** The data regarding average additional investment capacity in Belagiri-1 Micro watershed is presented in Table 39. The results indicate that, households have an average investment capacity of Rs. 3441.18 for land development, Rs.294.12 for adoption of improved livestock breeds.

**Table 39. Average additional investment capacity of households in Belagiri-1 micro-watershed**

Sl.No.	Particulars	LL (3)	MF (12)	SF (11)	SMF (6)	MDF (2)	All (34)
		Rs.	Rs.	Rs.	Rs.	Rs.	Rs.
1	Land development	0	833.33	7454.55	4166.67	0	3441.18
2	Improved crop production	0	0	909.09	0	0	294.12

**Source of funds for additional investment:** The data regarding source of funds for additional investment in Belagiri-1 Micro watershed is presented in Table 40. The results indicate that, the sources of finance raised from bank as a loan and from own sources for land development were 14.71 and 2.94 per cent.

**Table 40. Source of funds for additional investment in Belagiri-1 micro-watershed**

Sl. No	Item	Land development		Improved crop production	
		N	%	N	%
1	Loan from bank	5	14.71	1	2.94

**Marketing of agricultural produce:** The data regarding marketing of the agricultural produce in Belagiri-1 Micro watershed is presented in Table 41. The results indicated that, 100.00 percent of output of Cotton was sold in the market with average price of Rs. 4700.00; 90.91 percent of output of Green gram was sold in the market with average price of Rs. 5000.00; 89.09 percent of output of Groundnut was sold in the market with average price of Rs. 5000.00 and 85.29 percent of output of Jowar was sold in the market with average price of Rs. 2350.00.

**Table 41. Marketing of agricultural produce in Belagiri-1 micro-watershed**

Sl.No	Crops	Output obtained (q)	Output retained (q)	Output sold (q)	Output sold (%)	Avg. Price obtained (Rs/q)
1	Cotton	46	0	46	100	4700
2	Green gram	22	2	20	91	5000
3	Groundnut	55	6	49	89	5000
4	Jowar	34	5	29	85	2350
5	Paddy	65	1	64	98	1425
6	Red gram	400	33	367	92	4815

**Marketing channels used for sale of agricultural produce:** The data regarding marketing channels used for sale of agricultural produce in Belagiri-1 Micro watershed is presented in Table 42. The results indicated that, 32.35 cent of the households have sold agricultural produce to the local/village merchants, 61.76 per cent of regulated market.

**Mode of transport of agricultural produce:** The data regarding mode of transport of agricultural produce in Belagiri-1 Micro watershed is presented in Table 43. The results



indicated that, 50.00 cent of the households have used tractor, 23.53 per cent have used Cart for the transport of agriculture commodity.

**Table 42. Marketing channels used for sale of agricultural produce in Belagiri-1 micro-watershed**

Sl.No.	Particulars	LL (3)		MF (12)		SF (11)		SMF (6)		MDF (2)		All (34)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Local/village Merchant	0	0	4	33	4	36.4	2	33.3	1	50	11	32.35
2	Regulated Market	0	0	8	67	8	72.7	4	66.7	1	50	21	61.76

**Table 43. Mode of transport of agricultural produce in Belagiri-1 micro-watershed**

Sl.No.	Particulars	LL (3)		MF (12)		SF (11)		SMF (6)		MDF (2)		All (34)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Cart	0	0	4	33	3	27.3	0	0	1	50	8	23.53
2	Tractor	0	0	4	33	7	63.6	5	83.3	1	50	17	50
3	Truck	0	0	4	33	2	18.2	1	16.7	0	0	7	20.59

**Incidence of soil and water erosion problems:** The data regarding incidence of incidence of soil and water erosion problems in Belagiri-1 Micro watershed is presented in Table 44. The results indicate that, 85.29 per cent of the households have experienced soil and water erosion problems.

**Table 44. Incidence of soil and water erosion problems in Belagiri-1 micro-watershed**

Sl.No.	Particulars	LL (3)		MF (12)		SF (11)		SMF (6)		MDF (2)		All (34)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Soil and water erosion problems in the farm	0	0	10	83	11	100	6	100	2	100	29	85.29

**Interest towards soil testing:** The data regarding Interest shown towards soil testing in Belagiri-1 Micro watershed is presented in Table 45. The results indicated that, 79.41 per cent of the households were interested towards soil testing.

**Table 45. Interest regarding soil testing in Belagiri-1 micro-watershed**

Sl.No.	Particulars	LL (3)		MF (12)		SF (11)		SMF (6)		MDF (2)		All (34)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Interest in soil test	0	0	10	83	11	100	4	67	2	100	27	79.41

### Soil and water conservation practices and structures adopted

The data regarding soil and water conservation practices and structures adopted in Belagiri-1 Micro watershed is presented in Table 46. The results indicated that 100 per cent of farmers practicing summer ploughing as soil and water conservation practice.

**Table 46. Soil and water conservation practices and structures adopted in Belagiri-1 micro-watershed**

Sl.No.	Particulars	LL (3)		MF (12)		SF (11)		SMF (6)		MDF (2)		All (34)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Field Bunding	0	0	1	8.3	3	27	0	0	0	0	4	11.76

**Status of soil and water conservation structures:** The data regarding status soil and water conservation structures adopted in Belagiri-1 Micro watershed is presented in Table

47. The results indicated that, the households have adopted field bunding as a soil and water conservation structures out of which 25.00 per cent was in good condition, 75.00 per cent was slightly damaged.

**Table 47. Status of soil and water conservation structures in Belagiri-1 micro-watershed**

Sl.No	Item	Good		Slightly Damaged		Severely Damaged		Full Replacement Required	
		N	%	N	%	N	%	N	%
1	Field Bunding	1	25	3	75	0	0	0	0

**Agencies involved in the soil and water conservation structures:** The data regarding Agencies involved in the soil and water conservation structures adopted in Belagiri-1 Micro watershed is presented in Table 48. The results indicated that, 11.76 per cent were done by Govt.

**Table 48. Agencies involved in the soil and water conservation structures in Belagiri-1 micro-watershed**

Sl.No.	Particulars	LL (3)		MF (12)		SF (11)		SMF (6)		MDF (2)		All (34)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Govt.	0	0	1	8.3	3	27.27	0	0	0	0	4	11.76

**Usage pattern of fuel for domestic use:** The data on usage pattern of fuel for domestic use in Belagiri-1 Micro watershed is presented in Table 49. The results indicated that, firewood was the major source of fuel for domestic use for 97.06 per cent of the households followed by LPG (2.94%).

**Table 49. Usage pattern of fuel for domestic use in Belagiri-1 micro-watershed**

Sl.No.	Particulars	LL (3)		MF (12)		SF (11)		SMF (6)		MDF (2)		All (34)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Fire Wood	3	100	11	91.7	11	100	6	100	2	100	33	97.06
2	LPG	0	0	1	8.33	0	0	0	0	0	0	1	2.94

**Source of drinking water:** The data on source of drinking water in Belagiri-1 Micro watershed is presented in Table 50. The results indicated that, piped waters supply was the major source for drinking water for 91.18 per cent of the households followed by bore well water (8.82%).

**Table 50. Source of drinking water in Belagiri-1 micro-watershed**

Sl.No.	Particulars	LL (3)		MF (12)		SF (11)		SMF (6)		MDF (2)		All (34)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Piped supply	3	100	12	100	11	100	4	66.7	1	50	31	91.18
2	Bore Well	0	0	0	0	0	0	2	33.3	1	50	3	8.82

**Source of light:** The data on source of light in Belagiri-1 Micro watershed is presented in Table 51. The results indicated that, electricity was the major source of light for 100.00 per cent of the households.

**Existence of sanitary toilet facility:** The data on availability of toilet facility in Belagiri-1 Micro watershed is presented in Table 52. The results indicated that, 47.06 per cent of the households possess toilets.

**Table 51. Source of light in Belagiri-1 micro-watershed**

Sl.No.	Particulars	LL (3)		MF (12)		SF (11)		SMF (6)		MDF (2)		All (34)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Electricity	3	100	12	100	11	100	6	100	2	100	34	100

**Table 52. Existence of sanitary toilet facility in Belagiri-1 micro-watershed**

Sl.No.	Particulars	LL (3)		MF (12)		SF (11)		SMF (6)		MDF (2)		All (34)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Sanitary toilet facility	1	33.3	12	100	1	9.09	1	17	1	50	16	47.1

**Possession of PDS card:** The data regarding possession of PDS card in Belagiri-1 Micro watershed is presented in Table 53. The results indicated that, 97.06 per cent of the households possessed BPL card, 2.94 per cent possessed APL card.

**Table 53. Possession of PDS card in Belagiri-1 micro-watershed**

Sl.No.	Particulars	LL (3)		MF (12)		SF (11)		SMF (6)		MDF (2)		All (34)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	APL	0	0	0	0	1	9.09	0	0	0	0	1	2.94
2	BPL	3	100	12	100	10	90.91	6	100	2	100	33	97.06

**Participation in NREGA programme:** The data regarding Participation in NREGA programme in Belagiri-1 Micro watershed is presented in Table 54. The results indicated that, only 5.88 percent of the households have participated in NREGA programme.

**Table 54. Participation in NREGA programme in Belagiri-1 micro-watershed**

Sl. No.	Particulars	LL (3)		MF (12)		SF (11)		SMF (6)		MDF (2)		All (34)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Participation in NREGA programme	0	0	1	8.33	1	9.09	0	0	0	0	2	5.88

**Adequacy of food items:** The data regarding adequacy of food items in Belagiri-1 Micro watershed is presented in Table 55. The results indicated that, the extent of adequacy of food items for cereals, pulses, Oilseeds and vegetables were 91.18, 70.59, 61.76, 88.24 per cent respectively, similarly for Fruits (14.71%), milk (17.65%) and Egg (17.65%).

**Table 55. Adequacy of food items in Belagiri-1 micro-watershed**

Sl.No.	Particulars	LL (3)		MF (12)		SF (11)		SMF (6)		MDF (2)		All (34)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Cereals	0	0	13	108	10	90.91	6	100	2	100	31	91.18
2	Pulses	0	0	8	66.7	9	81.82	6	100	1	50	24	70.59
3	Oilseed	0	0	6	50	8	72.73	5	83.3	2	100	21	61.76
4	Vegetables	0	0	11	91.7	11	100	6	100	2	100	30	88.24
5	Fruits	0	0	2	16.7	2	18.18	1	16.7	0	0	5	14.71
6	Milk	0	0	3	25	2	18.18	1	16.7	0	0	6	17.65
7	Egg	0	0	2	16.7	0	0	3	50	1	50	6	17.65

**Inadequacy of food items:** The data regarding in adequacy of food items in Belagiri-1 Micro watershed is presented in Table 56. The results indicated that, the extent of in adequacy of food items for cereals, pulses, Oilseeds and vegetables were 2.94, 20.59,

26.47, 2.94 and 88.24 per cent respectively, similarly for fruits (76.47%), milk (55.88%), egg (70.59%) and meat (88.24%).

**Table 56. Inadequacy of food items in Belagiri-1 micro-watershed**

Sl.No.	Particulars	LL (3)		MF (12)		SF (11)		SMF (6)		MDF (2)		All (34)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Cereals	0	0	0	0	1	9.09	0	0	0	0	1	2.94
2	Pulses	0	0	4	33.3	2	18.18	0	0	1	50	7	20.59
3	Oilseed	0	0	6	50	3	27.27	0	0	0	0	9	26.47
4	Vegetables	0	0	1	8.33	0	0	0	0	0	0	1	2.94
5	Fruits	0	0	10	83.3	9	81.82	5	83.3	2	100	26	76.47
6	Milk	0	0	7	58.3	8	72.73	3	50	1	50	19	55.88
7	Egg	0	0	9	75	11	100	3	50	1	50	24	70.59
8	Meat	0	0	11	91.7	11	100	6	100	2	100	30	88.24

**Farming constraints:** The data regarding farming constraints experienced by households in Belagiri-1 Micro watershed is presented in Table 57. The results indicated that, lower fertility status of the soil was the constraint experienced by (82.35 %) per cent of the households, wild animal menace on farm field (32.35%), frequent incidence of pest and diseases (70.59%), inadequacy of irrigation water (76.47%), high cost of fertilizers and plant protection chemicals (82.35%), high rate of interest on credit (82.35%), low price for the agricultural commodities (85.29 %), lack of marketing facilities in the area (76.47%), inadequate extension services (23.53 %), lack of transport for safe transport of the agricultural produce to the market (76.47%).

**Table 57. Farming constraints experienced in Belagiri-1 micro-watershed**

SN	Particulars	LL (3)		MF (12)		SF (11)		SMF (6)		MDF (2)		All (34)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Lower fertility status of the soil	0	0	12	100	8	72.73	6	100	2	100	28	82.35
2	Wild animal menace on farm field	0	0	5	41.67	5	45.45	1	16.67	0	0	11	32.35
3	Frequent incidence of pest and diseases	0	0	8	66.67	10	90.91	5	83.33	1	50	24	70.59
4	Inadequacy of irrigation water	0	0	10	83.33	9	81.82	5	83.33	2	100	26	76.47
5	High cost of Fertilizers and plant protection chemicals	0	0	11	91.67	9	81.82	6	100	2	100	28	82.35
6	High rate of interest on credit	0	0	11	91.67	9	81.82	6	100	2	100	28	82.35
7	Low price for the agricultural commodities	0	0	12	100	10	90.91	5	83.33	2	100	29	85.29
8	Lack of marketing facilities in the area	0	0	9	75	10	90.91	5	83.33	2	100	26	76.47
9	Inadequate extension services	0	0	3	25	3	27.27	2	33.33	0	0	8	23.53
10	Lack of transport for safe transport of the Agril produce to the market.	0	0	11	91.67	9	81.82	4	66.67	2	100	26	76.47



**SUMMARY AND IMPLICATIONS**

In order to assess the socio-economic condition of the farmers in the watershed 34 households located in the micro watershed were interviewed for the survey. The study was conducted in Belagiri-1 micro-watershed (Belagiri sub-watershed, Yadgiri taluk & District) is located at North latitude 16° 50' 28.182" and 16° 48' 54.3" and East longitude 77° 14' 23.172" and 77° 11' 59.467" covering an area of about 601.86 ha bounded by under Belagera, Sutharahosalli and Honagera villages.

Socio-economic analysis of Belagiri-1 micro watersheds of Belagiri sub-watershed, Yadgiri taluk & District indicated that, out of the total sample of 34 farmers were sampled in Belagiri-1 micro-watershed among households surveyed 12 (35.29%) were marginal, 11 (32.35%) were small, 6 (17.65 %) were semi medium and 2 (5.88 %) were medium farmers. 3 landless farmers were also interviewed for the survey. The population characteristics of households indicated that, there were 118 (59.00%) men and 82 (41.00 %) were women. The average population of landless was 5.7, marginal farmers were 5, small farmers were 6.9, semi medium farmers were 6 and medium farmers were 5.5. Majority of the respondents (48.50%) were in the age group of 16-35 years.

Education level of the sample households indicated that, there were 16.00 per cent illiterates, 82.50 per cent pre university education and 7.50 per cent attained graduation. About, 88.24 per cent of household heads practicing agriculture and 11.76 per cent of the household heads were engaged as agricultural labourers. Agriculture was the major occupation for 46.00 per cent of the household members.

In the study area, 79.41 per cent of the households possess katcha house and 17.65 per cent possess pucca house. The durable assets owned by the households showed that, 85.29 per cent possess TV, 5.88 per cent possess mixer grinder, 102.94 per cent possess mobile phones and 26.47 per cent possess motor cycles.

Farm implements owned by the households indicated that, 50.00 per cent of the households possess plough, 2.94 per cent possess tractor, 5.88 per cent possess bullock cart and 26.47 per cent possess sprayer. Regarding livestock possession by the households, 20.59 per cent possess local cow and 14.71 per cent possess buffalo.

The average labour availability in the study area showed that, own labour men available in the micro watershed was 2.24, women available in the micro watershed was 1.50, hired labour (men) available was 6.18 and hired labour (women) available was 11.03. Further, 44.12 per cent of the households opined that hired labour was inadequate during the agricultural season.

Out of the total land holding of the sample respondents 76.04 per cent (39.75 ha) of the area is under dry condition and the remaining 23.96 per cent area is irrigated land. There were 7.00 live bore wells and 7.00 dry bore wells among the sampled households.

Bore well was the major source of irrigation for 32.35 per cent of the households. The major crops grown by sample farmers are Red gram, Groundnut, Cotton, Green gram and Paddy and cropping intensity was recorded as 100.00 per cent.

Out of the sample households 97.06 percent possessed bank account and 47.06 per cent of them have savings in the account. About 100.00 per cent of the respondents borrowed credit from various sources. Among the credit borrowed by households, 21.43 per cent have borrowed loan from commercial banks and 71.43 per cent from co-operative/Grameena bank.

Majority of the respondents (100.00%) have borrowed loan for agriculture purpose. Regarding the opinion on institutional sources of credit, 100.00 per cent of the households opined that credit helped to perform timely agricultural operations. The per hectare cost of cultivation for Red gram, Groundnut, Cotton, Green gram and Paddy was Rs.37413.18, 62616.52, 26530.56, 28811.74 and 66062.03 with benefit cost ratio of 1:1.95, 1: 1.70, 1: 2.10, 1: 1.90 and 1:0.80 respectively.

Further, 67.65 per cent of the households opined that dry fodder was adequate and 5.88 per cent of the households have opined that the green fodder was adequate. The average annual gross income of the farmers was Rs. 170735.29 in micro-watershed, of which Rs. 78852.94 comes from agriculture. Sampled households have grown 25 horticulture trees.

Households have an average investment capacity of Rs. 3441.18 for land development. Source of funds for additional investment is concerned, 2.94 per cent depends on own funds and 14.71 per cent depends on bank loan for land development activities.

Regarding marketing channels, 32.35 per cent of the households have sold agricultural produce to the local/village merchants, while, 61.76 per cent have sold in regulated markets. Further, 50.00 per cent of the households have used tractor for the transport of agriculture commodity.

Majority of the farmers (85.29%) have experienced soil and water erosion problems in the watershed and 79.41 per cent of the households were interested towards soil testing. Fire was the major source of fuel for domestic use for 97.06 per cent of the households and 2.94 per cent households has LPG connection.

Piped supply was the major source for drinking water for 91.18 per cent of the households. Electricity was the major source of light for 100.00 per cent of the households. In the study area, 47.06 per cent of the households possess toilet facility. Regarding possession of PDS card, 97.06 per cent of the households possessed BPL card, 2.94 per cent of the household's possessed APL card. Households opined that, the requirement of cereals (91.18%), pulses (70.59%) and oilseeds (61.76%) are adequate for consumption.



Farming constraints experienced by households in the micro watersheds were lower fertility status of the soil (82.35%) wild animal menace on farm field (32.35%), frequent incidence of pest and diseases (70.59%), inadequacy of irrigation water (76.47%), high cost of fertilizers and plant protection chemicals (82.35%), high rate of interest on credit (82.35%), low price for the agricultural commodities (85.29%), lack of marketing facilities in the area (76.47%), inadequate extension services (23.53%), lack of transport for safe transport of the agricultural produce to the market (76.47%).

### **Implications of the survey**

- ✓ Result indicated that, there were 16.00 per cent were illiterate hence, extension methodologies such as demonstration, street play, drama, video shows will be effective in dissemination of the technologies in the micro watershed.
- ✓ The data indicate that, 79.41 per cent of the households possess katcha house. Hence, the development department while implementing the watershed plan should focus on agriculture to enhance the productivity of major crops in the area to increase the income of the farmers.
- ✓ Results indicated that the local institutional participation of the household members in the micro watershed is minimal hence, activities like membership campaign, awareness creation about the benefits of membership in local institutions and strengths of organized groups must be conveyed.
- ✓ Majority of the households in the watershed have experience in use of mobile phones, and television hence, these mass media can be effectively utilized for transfer of technology as well as for information dissemination.
- ✓ The farm machinery/implement possession in the micro watershed was found to be minimum the reasons may lack of knowledge or lack of financial ability which can be addressed through training on use of different farm implements, providing information on different sources of finance for purchase of farm implements.
- ✓ The possession of livestock such as crossbred cow found is less hence, farmers must be made aware of the benefits of crossbred cow in increased milk production.
- ✓ The possession of livestock such as sheep, goat and poultry was found to be low hence, farmers may be informed the role of subsidiary enterprises in enhancing the income and information on financial support for subsidiary activities.
- ✓ The data indicate that, job/work was the reason for all the migrants hence, farmers may be trained on profitable agriculture or self employment such as animal husbandry, plate making, sheep rearing, goat rearing, rabbit rearing with suitable information on sources of financial support.
- ✓ The results indicate that there was a change in quality of life due to migration hence, the developmental departments should take actions to arrest migration and to improve the quality of the life in rural areas.
- ✓ Households possess 30.23ha (76.04 %) of dry land and 9.53ha (23.96 %) of irrigated land hence, the availability of the dry land agricultural technologies such as short

duration crops, high yielding drought resistance crop varieties, drip irrigation technology and subsidy information will be helpful for the farmers to enhance the productivity of land and as well as farmers income.

- ✓ Few of the bore well in micro watershed found non functional hence, farmers may be trained on possibility of bore well rejuvenation.
- ✓ Bore well was major source of irrigation for 20.59 per cent of the households. hence, in order to increase the area under irrigation as well as to increase the water use efficiency farmers may trained on drip irrigation and provide the information on subsidy for drip irrigation equipment's along with the information on different agencies which provides the financial assistance for drip irrigation.
- ✓ The cropping intensity in the micro watershed was found to be (100.00 %) hence, care must be taken by the implementing agency to bring uncultivated land into cultivation through suitable measures.
- ✓ Many of the household members have borrowed loan from cooperative banks which has higher rate of interest hence, farmers may be sensitized on the different sources of credit with lesser interest rate such SHGs etc.
- ✓ The results indicated the non availability of both green and dry fodder throughout the year hence, fodder development activities can be taken up in the micro watershed.
- ✓ The average annual gross income of the households Rs.78852.94 from agriculture, Rs.3529.41 from business and Rs. 69647.06 from wages and. Agriculture was found to be the major source of income for households hence; the development activities should focus on productivity enhancement, marketing arrangements and agricultural technology dissemination to have a direct impact on the farmers.
- ✓ The cultivation of forest species is found minimal hence, information and production technology related to agro-forestry and integrated farming system.
- ✓ The data indicated that, 85.29 per cent of the households have experienced soil and water erosion problems. Hence, those farmers who reported the soil and water erosion problems may be given attention while implementation of the watershed development plan.
- ✓ The data indicated that, 79.41 per cent of the households have interest in soil testing hence, farmers must be provided with the information on various institutions which are involved in soil testing for the benefit of the farmers.
- ✓ Except summer ploughing the adoption of other soil and water conservation structures is minimum hence, the farmers in the micro watershed should be sensitized on the use of different conservation structures for soil water conservation.
- ✓ Cereals and pulses found be adequate for per cent of the households respectively hence, farm households and the farm women must be trained on importance of balanced nutrition and role of vegetable, milk, egg, meat in balanced diet.
- ✓ Lower fertility status of the soil (82.35%), wild animal menace on farm field (32.35%), frequent incidence of pest and diseases (70.59%), high cost of fertilizers

and plant protection chemicals (82.35%), high rate of interest on credit (82.35%), low price for the agricultural commodities (85.29%), lack of marketing facilities in the area (76.47%), inadequate extension services (23.53%), lack of transport for safe transport of the agricultural produce to the market (76.47%) were the major farming constraints experienced hence, these constraints must be addressed immediately for the welfare of the farmers. Awareness to be created among the farmers to approach nearest KVKs/RSKs and other developmental departments for technical and for subsidized inputs and utilize the well established regulated markets, approaching the contract firms, direct markets to avoid the involvement of middlemen.