



हर कदम, हर डगर
किसानों का हमसफर
भारतीय कृषि अनुसंधान परिषद

Agrisearch with a human touch

**LAND RESOURCE INVENTORY AND SOCIO-ECONOMIC STATUS OF
FARM HOUSEHOLDS FOR WATERSHED PLANNING AND
DEVELOPMENT**

YADGIR RF-1 (4D5B2A1a) MICROWATERSHED

Yadgir Taluk and District, Karnataka

Karnataka Watershed Development Project – II

SUJALA – III

World Bank funded Project



THE WORLD BANK



ICAR – NATIONAL BUREAU OF SOIL SURVEY AND LAND USE PLANNING



ICAR - NBSS & LUP

**WATERSHED DEVELOPMENT DEPARTMENT
GOVT. OF KARNATAKA, BANGALORE**



About ICAR - NBSS&LUP

The ICAR-National Bureau of Soil Survey and Land Use Planning (ICAR-NBSS&LUP), Nagpur, a premier Institute of the Indian Council of Agricultural Research (ICAR), was set up during 1976 with the objective to prepare soil resource maps at national, state and district levels and to provide research inputs in soil resource mapping and its applications, land evaluation, land use planning, land resource management, and database management using GIS for optimising land use on different kinds of soils in the country.

The Bureau has been engaged in carrying out soil resource survey, agro-ecological and soil degradation mapping at the country, state and district levels for qualitative assessment and monitoring the soil health towards viable land use planning. The research activities have resulted in identifying the soil potentials and problems, and the various applications of the soil surveys with the ultimate objective of sustainable agricultural development. The Bureau has the mandate to correlate and classify soils of the country and maintain a National Register of all the established soil series. The Institute is also imparting in-service training to staff of the soil survey agencies in the area of soil survey, land evaluation and soil survey interpretations for land use planning. The Bureau in collaboration with Panjabrao Krishi Vidyapeeth, Akola is running post-graduate teaching and research programme in land resource management, leading to M.Sc. and Ph.D. degrees.

Citation: Rajendra Hegde, Ramesh Kumar, S.C., K.V. Niranjana, S. Srinivas, M.Lalitha, B.A. Dhanorkar, R.S. Reddy and S.K. Singh (2019). "Land resource inventory and socio-economic status of farm households for watershed planning and development of Yadgir Rf-1 (4D5B2A1a) Microwatershed, Yadgir Taluk and District, Karnataka", ICAR-NBSS &LUP Sujala MWS Publ.414, ICAR – NBSS & LUP, RC, Bangalore. p.119 & 31.

TO OBTAIN COPIES,

Please write to:

Director, ICAR - NBSS & LUP,

Amaravati Road, NAGPUR - 440 033, India

Phone : (0712) 2500386, 2500664, 2500545 (O)

Telefax : 0712-2522534

E-Mail : director@nbsslup.ernet.in

Website URL : nbsslup.in

Or

Head, Regional Centre, ICAR - NBSS&LUP, Hebbal, Bangalore - 560 024

Phone : (080) 23412242, 23510350 (O)

Telefax : 080-23510350

E-Mail : nbssrcb@gmail.com



**LAND RESOURCE INVENTORY AND SOCIO-ECONOMIC
STATUS OF FARM HOUSEHOLDS FOR WATERSHED
PLANNING AND DEVELOPMENT**

YADGIR RF-1 (4D5B2A1a) MICROWATERSHED

Yadgir Taluk and District, Karnataka

Karnataka Watershed Development Project – II

Sujala-III

World Bank funded Project



**ICAR – NATIONAL BUREAU OF SOIL SURVEY AND LAND
USE PLANNING**



ICAR - NBSS & LUP



**WATERSHED DEVELOPMENT DEPARTMENT, GOVT. OF
KARNATAKA, BANGALORE**



PREFACE

In Karnataka, as in other Indian States, the livelihoods of rural people are intertwined with farming pursuits. The challenges in agriculture are seriously threatening the livelihood of a large number of farmers as they have been practicing farming in contextual factors beyond their control. Climatic factors are the most important ones and have become much more significant in recent times due to rapid climate changes induced by intensive anthropogenic activities affecting our ecosystem in multiple ways. Climate change has become the reality, it is happening and efforts to evolve and demonstrate climate resilient technologies have become essential. Due to the already over stressed scenario of agrarian sector, the climate change is resulting in manifold increase in the complexities, pushing the rural mass to face more and more unpredictable situations. The rising temperatures and unpredictable rainfall patterns are going to test seriously the informed decisions farmers have to make in order to survive in farming and sustain their livelihood.

It is generally recognized that impacts of climate change shall not be uniform across the globe. It is said that impact of climate change is more severe in South Asia. Based on the analysis of meteorological data, it is predicted that in India, there will be upward trend in mean temperature, downward trend in relative humidity, annual rainfall and number of wet days in a year. Also, in general, phenomena like erratic monsoon, spread of tropical diseases, rise in sea levels, changes in availability of fresh water, frequent floods, droughts, heat waves, storms and hurricanes are predicted. Each one of these adverse situations are already being experienced in various parts of India and also at the global level. Decline in agricultural productivity of small and marginal farmers becoming more vulnerable is already witnessed.

In Karnataka, more than 60 per cent of the population live in rural areas and depend on agriculture and allied activities for their livelihood. Though the state has achieved significant progress in increasing the yield of many crops, there is tremendous pressure on the land resources due to the growing and competing demands of various land uses. This is reflected in the alarming rate of land degradation observed. Already more than 50 per cent of the area is affected by various forms of degradation. If this trend continues, the sustainability of the fragile ecosystem will be badly affected. The adverse effects of change in the climatic factors are putting additional stress on the land resources and the farmers dependent on this.

The natural resources (land, water and vegetation) of the state need adequate and constant care and management, backed by site-specific technological interventions and investments particularly by the government. Detailed database pertaining to the nature of

the land resources, their constraints, inherent potentials and suitability for various land based rural enterprises, crops and other uses is a prerequisite for preparing location-specific action plans, which are in tune with the inherent capability of the resources. Any effort to evolve climate resilient technologies has to be based on the baseline scientific database. Then only one can expect effective implementation of climate resilient technologies, monitor the progress, make essential review of the strategy, and finally evaluate the effectiveness of the implemented programs. The information available at present on the land resources of the state are of general nature and useful only for general purpose planning. Since the need of the hour is to have site-specific information suitable for farm level planning and detailed characterization and delineation of the existing land resources of an area into similar management units is the only option.

ICAR-NBSS&LUP, Regional Centre, Bangalore has taken up a project sponsored by the Karnataka Watershed Development Project-II, (Sujala-III), Government of Karnataka funded by the World Bank under Component -1 Land Resource Inventory. This study was taken up to demonstrate the utility of such a database in reviewing, monitoring and evaluating all the land based watershed development programs on a scientific footing. To meet the requirements of various land use planners at grassroots level, the present study on “Land Resource Inventory and Socio-Economic Status of Farm Households for Watershed Planning and Development of for Yadgir Rf-1 microwatershed in Yadgir Taluk and District, Karnataka” for integrated development was taken up in collaboration with the State Agricultural Universities, IISC, KRSRAC, KSNDMC as Consortia partners. The project provides detailed land resource information at cadastral level (1:7920 scale) for all the plots and socio-economic status of farm households covering thirty per cent farmers randomly selected representing landed and landless class of farmers in the micro-watershed. The project report with the accompanying maps for the microwatershed will provide required detailed database for evolving effective land use plan, alternative land use options and conservation plans for the planners, administrators, agricultural extension personnel, KVK officials, developmental departments and other land users to manage the land resources in a sustainable manner.

It is hoped that this database will be useful to the planners, administrators and developmental agencies working in the area in not only for formulating location specific developmental schemes but also for their effective monitoring at the village/watershed level.

Nagpur

Date:15-10-2019

S.K. SINGH

Director, ICAR - NBSS&LUP Nagpur

Contributors

Dr. Rajendra Hegde Principal Scientist, Head & Project Leader, Sujala-III Project ICAR-NBSS&LUP, Regional Centre, Bangalore	Dr. S.K.Singh Director, ICAR-NBSS&LUP Coordinator, Sujala-III Project Nagpur
Soil Survey, Mapping & Report Preparation	
Dr. B.A. Dhanorkar	Sh. R.S. Reddy
Dr. K.V. Niranjana	Mr. Somashekar T N
	Smt. Chaitra, S.P.
	Dr. Gopali bardhan
	Dr. Mahendra Kumar, M.B.
	Ms. Arpitha, G.M.
Field Work	
Sh. C.BacheGowda	Sh. Mahesh, D.B.
Sh. Somashekar	Sh. Ashok S Sindagi
Sh. M. Jayaramaiah	Sh. Veerabhadrapa B.
Sh. Paramesha, K.	Sh. Shankarappa
Sh. B. M. Narayana Reddy	Sh. Anand
	Sh. Arun N Kambar.
	Sh Kamalesh Awate
	Sh. Sharaan Kumar Huppar
	Sh. Yogesh H.N.
	Sh. Kalaveerachari R Kammar
GIS Work	
Dr. S.Srinivas	Sh. A.G.Devendra Prasad
Sh. D.H.Venkatesh	Sh. Prakashanaik, M.K.
Smt.K.Sujatha	Sh. Abhijith Sastry, N.S.
Smt. K.V.Archana	Sh. Sudip Kumar Suklabaidya
Sh. N. Maddileti	Sh. Avinash, K.N.
	Sh. Amar Suputhra, S
	Sh. Deepak, M.J.
	Smt. K.Karunya Lakshmi
	Ms. Seema, K.V.
	Ms. A. Rajab Nisha

Laboratory Analysis	
Dr. K.M.Nair	Ms. Steffi Peter
Smt. Arti Koyal	Ms. Thara, V.R
Smt. Parvathy	Ms. Roopa, G.
	Ms. Swati, H.
	Sh. Shantaveera Swami
	Ms. Shwetha, N.K.
	Smt. Ishrat Haji
	Ms. P. Pavan Kumari
	Ms. Padmaja
	Ms. Veena, M.
Socio-Economic Analysis	
Dr. S.C. Ramesh Kumar	Sh. M.K. Prakashanaik
	Ms. Karuna V. Kulkarni
	Mrs. Sowmya A.N
	Sh. Vinod R
	Sh. Basavaraja
	Sh. Vijay Kumar Lamani
	Ms. Sowmya K.B
	Mrs. Prathibha, D.G
	Sh. Rajendra,D
Soil & Water Conservation	
Sh. Sunil P. Maske	
Watershed Development Department, GoK, Bangalore	
Sh. Rajeev Ranjan IFS Project Director & Commissioner, WDD	Dr. A. Natarajan NRM Consultant, Sujala-III Project
Dr. S.D. Pathak IFS Executive Director & Chief Conservator of Forests, WDD	

PART-A

LAND RESOURCE INVENTORY

Contents

Preface		
Contributors		
Executive Summary		
Chapter 1	Introduction	1
Chapter 2	Geographical Setting	3
2.1	Location and Extent	3
2.2	Geology	3
2.3	Physiography	4
2.4	Drainage	4
2.5	Climate	4
2.6	Natural Vegetation	6
2.7	Land Utilization	6
Chapter 3	Survey Methodology	11
3.1	Base maps	11
3.2	Image Interpretation for Physiography	11
3.3	Field Investigation	14
3.4	Soil Mapping	15
3.5	Land Management Units	15
3.6	Laboratory Characterization	16
Chapter 4	The Soils	19
4.1	Soils of granite gneiss landscape	19
Chapter 5	Interpretation for Land Resource Management	25
5.1	Land Capability Classification	25
5.2	Soil Depth	27
5.3	Surface Soil Texture	28
5.4	Soil Gravelliness	29
5.5	Available Water Capacity	30
5.6	Soil Slope	31
5.7	Soil Erosion	32
Chapter 6	Fertility Status	35
6.1	Soil Reaction (pH)	35
6.2	Electrical Conductivity (EC)	35
6.3	Organic Carbon (OC)	35
6.4	Available Phosphorus	37
6.5	Available Potassium	37
6.6	Available Sulphur	37
6.7	Available Boron	37
6.8	Available Iron	38
6.9	Available Manganese	38
6.10	Available Copper	38
6.11	Available Zinc	42

Chapter 7	Land Suitability for Major Crops	43
7.1	Land suitability for Sorghum	43
7.2	Land suitability for Maize	44
7.3	Land suitability for Bajra	45
7.4	Land suitability for Groundnut	46
7.5	Land suitability for Sunflower	47
7.6	Land suitability for Redgram	48
7.7	Land suitability for Bengal gram	49
7.8	Land suitability for Cotton	50
7.9	Land suitability for Chilli	51
7.10	Land suitability for Tomato	52
7.11	Land suitability for Brinjal	53
7.12	Land suitability for Onion	54
7.13	Land suitability for Bhendi	55
7.14	Land suitability for Drumstick	56
7.15	Land suitability for Mango	57
7.16	Land suitability for Guava	58
7.17	Land suitability for Sapota	59
7.18	Land Suitability for Pomegranate	60
7.19	Land Suitability for Musambi	61
7.20	Land Suitability for Lime	62
7.21	Land Suitability for Amla	63
7.22	Land Suitability for Cashew	64
7.23	Land Suitability for Jackfruit	65
7.24	Land Suitability for Jamun	66
7.25	Land Suitability for Custard apple	67
7.26	Land Suitability for Tamarind	68
7.27	Land Suitability for Mulberry	69
7.28	Land Suitability for Marigold	70
7.29	Land Suitability for Chrysanthemum	71
7.30	Land Management Units (LMUs)	103
7.31	Proposed Crop Plan for Yadgir Rf-1 Microwatershed	104
Chapter 8	Soil Health Management	107
Chapter 9	Soil and Water conservation Treatment Plan	111
9.1	Treatment Plan	111
9.2	Recommended Soil and Water Conservation measures	115
9.3	Greening of Microwatershed	116
	References	119
	Appendix I	I
	Appendix II	III
	Appendix III	V

LIST OF TABLES

2.1	Mean Monthly Rainfall, PET, 1/2 PET at Yadgir Taluk & District	5
2.2	Land Utilization in Yadgir district	7
3.1	Differentiating Characteristics used for Identifying Soil Series	15
3.2	Soil map unit description of Yadgir Rf-1Microwatershed	16
4.1	Physical and Chemical Characteristics of Soil Series identified in Yadgir Rf-1microwatershed	22
7.1	Soil-Site Characteristics of Yadgir Rf-1 Microwatershed	73
7.2	Crop suitability for Sorghum	74
7.3	Crop suitability for Maize	75
7.4	Crop suitability for Bajra	76
7.5	Crop suitability for Groundnut	77
7.6	Crop suitability for Sunflower	78
7.7	Crop suitability for Redgram	79
7.8	Crop suitability for Bengal gram	80
7.9	Crop suitability for Cotton	81
7.10	Crop suitability for Chilli	82
7.11	Crop suitability for Tomato	83
7.12	Crop suitability for Brinjal	84
7.13	Crop suitability for Onion	85
7.14	Crop suitability for Bhendi	86
7.15	Crop suitability for Drumstick	87
7.16	Crop suitability for Mango	88
7.17	Crop suitability for Guava	89
7.18	Crop suitability for Sapota	90
7.19	Crop suitability for Pomegranate	91
7.20	Crop suitability for Musambi	92
7.21	Crop suitability for Lime	93
7.22	Crop suitability for Amla	94
7.23	Crop suitability for Cashew	95
7.24	Crop suitability for Jackfruit	96

7.25	Crop suitability for Jamun	97
7.26	Crop suitability for Custard apple	98
7.27	Crop suitability for Tamarind	99
7.28	Crop suitability for Mulberry	100
7.29	Crop suitability for Marigold	101
7.30	Crop suitability for Chrysanthemum	102
7.31	Proposed Crop Plan for Yadgir Rf-1Microwatershed	105

LIST OF FIGURES

2.1	Location map of Yadgir Rf-1Microwatershed	3
2.2	Granite and granite gneiss rock formation	4
2.3	Rainfall distribution in Yadgir Taluk & District	5
2.4	Natural vegetation of Yadgir Rf-1Microwatershed	6
2.5	Current Land use map of Yadgir Rf-1Microwatershed	7
2.6 a	Different crops and cropping systems in Yadgir Rf-1Microwatershed	8
2.6 b	Different crops and cropping systems in Yadgir Rf-1Microwatershed	9
3.1	Scanned and Digitized Cadastral map of Yadgir Rf-1Microwatershed	12
3.2	Satellite image of Yadgir Rf-1Microwatershed	13
3.3	Cadastral map overlaid on IRS PAN+LISS IV merged imagery of Yadgir Rf-1Microwatershed	13
3.4	Location of profiles in a transect	14
3.5	Soil phase or management units of Yadgir Rf-1Microwatershed	17
5.1	Land Capability Classification map of Yadgir Rf-1Microwatershed	27
5.2	Soil Depth map of Yadgir Rf-1Microwatershed	28
5.3	Surface Soil Texture map of Yadgir Rf-1Microwatershed	29
5.4	Soil Gravelliness map of Yadgir Rf-1Microwatershed	30
5.5	Soil Available Water Capacity map Yadgir Rf-1Microwatershed	31
5.6	Soil Slope map of Yadgir Rf-1Microwatershed	32
5.7	Soil Erosion map of Yadgir Rf-1Microwatershed	33
6.1	Soil Reaction (pH) map of Yadgir Rf-1Microwatershed	36
6.2	Electrical Conductivity (EC) map of Yadgir Rf-1Microwatershed	36
6.3	Soil Organic Carbon (OC) map of Yadgir Rf-1Microwatershed	37
6.4	Soil Available Phosphorus map of Yadgir Rf-1Microwatershed	38
6.5	Soil Available Potassium map of Yadgir Rf-1Microwatershed	39
6.6	Soil Available Sulphur map of Yadgir Rf-1Microwatershed	39
6.7	Soil Available Boron map of Yadgir Rf-1Microwatershed	40
6.8	Soil Available Iron map of Yadgir Rf-1Microwatershed	40
6.9	Soil Available Manganese map of Yadgir Rf-1Microwatershed	41
6.10	Soil Available Copper map of Yadgir Rf-1Microwatershed	41
6.11	Soil Available Zinc map of Yadgir Rf-1Microwatershed	42
7.1	Land suitability for Sorghum	44

7.2	Land suitability for Maize	45
7.3	Land suitability for Bajra	46
7.4	Land suitability for Groundnut	47
7.5	Land suitability for Sunflower	48
7.6	Land suitability for Redgram	49
7.7	Land suitability for Bengal gram	50
7.8	Land suitability for Cotton	51
7.9	Land suitability for Chilli	52
7.10	Land suitability for Tomato	53
7.11	Land suitable for Brinjal	54
7.12	Land suitable for Onion	55
7.13	Land suitable for Bhendi	56
7.14	Land suitable for Drumstick	57
7.15	Land suitability for Mango	58
7.16	Land suitability for Guava	59
7.17	Land suitability for Sapota	60
7.18	Land suitability for Pomegranate	61
7.19	Land suitability for Musambi	62
7.20	Land suitability for Lime	63
7.21	Land suitability for Amla	64
7.22	Land suitability for Cashew	65
7.23	Land suitability for Jackfruit	66
7.24	Land suitability for Jamun	67
7.25	Land suitability for Custard apple	68
7.26	Land suitability for Tamarind	69
7.27	Land suitability for Mulberry	70
7.28	Land suitability for Marigold	71
7.29	Land suitability for Chrysanthemum	72
7.30	Land management units map of Yadgir Rf-1Microwatershed	103
9.1	Soil and water conservation plan map of Yadgir Rf-1Microwatershed	116

EXECUTIVE SUMMARY

The land resource inventory of Yadgir Rf-1Microwatershed was conducted using village cadastral maps and IRS satellite imagery on 1:7920 scale. The false colour composites of IRS imagery were interpreted for physiography and the physiographic delineations were used as base for mapping soils. The soils were studied in several transects and a soil map was prepared with phases of soil series as mapping units. Random checks were made all over the area outside the transects to confirm and validate the soil map unit boundaries. The soil map shows the geographic distribution and extent, characteristics, classification, behavior and use potentials of the soils in the microwatershed.

The present study covers an area of 306 ha in Yadgir taluk & district, Karnataka. The climate is semiarid and categorized as drought-prone with an average annual rainfall of 866 mm, of which about 652 mm is received during south-west monsoon, 138 mm during north-east and the remaining 76 mm during the rest of the year. An area of 141(46%) ha in the microwatershed is covered by soils and about 165 ha (54%) covered by rock outcrops. The salient findings from the land resource inventory are summarized briefly below.

- ❖ The soils belong to 3 soil series and 4 soil phases (management units) and 2 land management units.*
- ❖ The length of crop growing period is about 120-150 days starting from 1st week of June to 4th week of October.*
- ❖ From the master soil map, several interpretative and thematic maps like land capability, soil depth, surface soil texture, soil gravelliness, available water capacity, soil slope and soil erosion were generated.*
- ❖ Soil fertility status maps for macro and micronutrients were generated based on the surface soil samples collected at every 320 m grid interval.*
- ❖ Land suitability for growing 29 major agricultural and horticultural crops was assessed and maps showing the degree of suitability along with constraints were generated.*
- ❖ An area about 141 ha (46%) in the microwatershed is suitable for agriculture.*
- ❖ About 31 per cent soils are very shallow and shallow (<25 -50 cm) and 15 per cent soils deep (100-150) in the microwatershed.*
- ❖ About 31 per cent are sandy soils and 16 percent soils are loamy at the surface in the microwatershed.*
- ❖ An area of 22 per cent soils are non gravelly (<15%) and about 24 per cent soils are gravelly (15-35%) soils in the microwatershed.*
- ❖ About 16 per cent soils are low (51-100 mm/m) and 31 per cent area is very low (<50 mm/m) available water capacity.*

- ❖ *Entire cultivated area in the microwatershed has very gently sloping (1-3% slope) lands.*
- ❖ *Entire cultivated area in the microwatershed has moderately (e2) eroded lands*
- ❖ *An area of about 20 per cent soils are moderately acid (pH 5.5-6.5) about 7 per cent soils are strongly acid (pH 5.0-5.5) and about 20 per cent soils are neutral (pH 6.5-7.3)*
- ❖ *The Electrical Conductivity (EC) of the soils in the entire cultivated area of the microwatershed is dominately $<2 \text{ d}^{\text{sm-1}}$ indicating that the soils are non-saline.*
- ❖ *An area of about 39 per cent is low ($<0.5\%$) and 7 percent is medium (0.50-0.75%) in organic carbon content.*
- ❖ *An area of 46 percent is high ($>57 \text{ kg/ha}$) in available phosphorus and about <1 percent is medium (23-57 kg/ha) in available phosphorus.*
- ❖ *Entire cultivated area in the microwatershed is medium (145-337 kg/ha) in available Potassium content.*
- ❖ *An area of about 38 percent is medium (10-20ppm) and 8 percent area is low ($<10\text{ppm}$) in available sulphur.*
- ❖ *Available boron is low ($<0.5 \text{ ppm}$) in a maximum area of about 43 per cent and medium (0.5-1.0 ppm) is 3 percent area.*
- ❖ *Available iron content is sufficient ($>4.5 \text{ ppm}$) in the entire cultivated area of the microwatershed.*
- ❖ *Available manganese and copper are sufficient in all the soils of the microwatershed.*
- ❖ *Available zinc content is deficient ($<0.6 \text{ ppm}$) in the entire cultivated area of the microwatershed*
- ❖ *The land suitability for 29 major crops grown in the microwatershed were assessed and the areas that are highly suitable (S1) and moderately suitable (S2) are given below. It is however to be noted that a given soil may be suitable for various crops but what specific crop to be grown may be decided by the farmer looking to his capacity to invest on various inputs, marketing infrastructure, market price and finally the demand and supply position.*

Land suitability for various crops in the Microwatershed

Crop	Suitability Area in ha (%)		Crop	Suitability Area in ha (%)	
	Highly suitable (S1)	Moderately suitable (S2)		Highly suitable (S1)	Moderately suitable (S2)
<i>Sorghum</i>	-	-	<i>Guava</i>	-	-
<i>Maize</i>	-	-	<i>Sapota</i>	-	-
<i>Bajra</i>	-	-	<i>Pomegranate</i>	-	-
<i>Groundnut</i>	-	-	<i>Musambi</i>	-	-
<i>Sunflower</i>	-	-	<i>Lime</i>	-	-
<i>Redgram</i>	-	-	<i>Amla</i>	-	-
<i>Bengal gram</i>	-	-	<i>Cashew</i>	-	-
<i>Cotton</i>	-	-	<i>Jackfruit</i>	-	-
<i>Chilli</i>	-	-	<i>Jamun</i>	-	-
<i>Tomato</i>	-	-	<i>Custard apple</i>	-	-
<i>Brinjal</i>	-	-	<i>Tamarind</i>	-	-
<i>Onion</i>	-	-	<i>Mulberry</i>	-	-
<i>Bhendi</i>	-	-	<i>Marigold</i>	-	-
<i>Drumstick</i>	-	-	<i>Chrysanthemum</i>	-	-
<i>Mango</i>	-	-			

- ❖ *Apart from the individual crop suitability, a proposed crop plan has been prepared for the identified LMUs by considering only the highly and moderately suitable lands for different crops and cropping systems with food, fodder, fiber and horticulture crops.*
- ❖ *Maintaining soil-health is vital for crop production and conserve soil and land resource base for maintaining ecological balance and to mitigate climate change. For this, several ameliorative measures have been suggested for these problematic soils like saline/alkali, highly eroded, sandy soils etc.*
- ❖ *Soil and water conservation treatment plan has been prepared that would help in identifying the sites to be treated and also the type of structures required.*
- ❖ *As part of the greening programme, several tree species have been suggested to be planted in marginal and submarginal lands, field bunds and also in the hillocks, mounds and ridges. This would help in not only supplementing the farm income but also provide fodder and fuel to generate lot of biomass which would help in maintaining an ecological balance and also contribute to mitigating the climate change.*

INTRODUCTION

Land is a scarce resource and basic unit for any material production. It can support the needs of the growing population, provided they use the land in a rational and judicious manner. But what is happening in many areas of the state is a cause for concern to everyone involved in the management of land resources at the grassroots level. The area available for agriculture is about 51 per cent of the total geographical area and more than 60 per cent of the people are still dependant on agriculture for their livelihood. The limited land area is under severe stress and strain due to increasing population pressure and competing demands of various land uses. Due to this, every year there is significant diversion of farm lands and water resources for non-agricultural purposes. Apart from this, due to lack of interest in farmers for farming, large tracts of cultivable lands are turning into fallows in many areas and this trend is continuing at an alarming rate.

Further, land degradation has emerged as a serious problem which has already affected about 38 lakh ha of cultivated area in the state. Soil erosion alone has degraded about 35 lakh ha. Almost all the uncultivated areas are facing various degrees of degradation, particularly soil erosion. Salinity and alkalinity has emerged as a major problem in more than 3.5 lakh ha in the irrigated areas of the state. Nutrient depletion and declining factor productivity is common in both rainfed and irrigated areas. The degradation is continuing at an alarming rate and there appears to be no systematic effort among the stakeholders to contain this process. In recent times, an aberration of weather due to climate change phenomenon has added another dimension leading to unpredictable situations to be tackled by the farmers.

In this critical juncture, the challenge before us is not only to increase the productivity per unit area which is steadily declining and showing a fatigue syndrome, but also to prevent or at least reduce the severity of degradation. If the situation is not reversed at the earliest, then the sustainability of the already fragile crop production system and the overall ecosystem will be badly affected in the state. The continued neglect and unscientific use of the resources for a long time has led to the situation observed at present in the state. It is a known fact and established beyond doubt by many studies in the past that the cause for all kinds of degradation is the neglect and irrational use of the land resources. Hence, there is an urgent need to generate a detailed site-specific farm level database on various land resources for all the villages/watersheds in a time bound manner that would help to protect the valuable soil and land resources and also to stabilize the farm production.

Therefore, the land resource inventory required for farm level planning is the one which investigates not only the surface but also consider the other parameters which are critical for productivity *viz.*, soils, climate, water, minerals and rocks, topography, geology, hydrology, vegetation, crops, land use pattern, animal population, socio-

economic conditions, infrastructure, marketing facilities and various schemes and developmental works of the government etc. From the data collected at farm level, the specific problems and potentials of the area can be identified and highlighted, conservation measures required for the area can be planned on a scientific footing, suitability of the area for various uses can be worked out and finally viable and sustainable land use options suitable for each and every land holding can be prescribed.

The Land Resource Inventory is basically done for identifying the potential and problem areas, developing sustainable land use plans, estimation of surface run off and water harvesting potential, preparation of soil and water conservation plans, land degradation/desertification etc. The Bureau is presently engaged in developing an LRI methodology using high resolution satellite remote sensing data and Digital Elevation Model (DEM) data to prepare Landscape Ecological Units (LEU) map representing agro-ecosystem as a whole. The LEU is preferred over landform as the base map for LRI. LEU is the assemblage of landform, slope and land use. An attempt has already been made to upscale the soil resource information from 1:250000 and 1:50000 scale to the LEU map in Goa and other states.

The land resource inventory aims to provide site specific database for Yadgir Rf-1 microwatershed in Yadgir Taluk & District, Karnataka State for the Karnataka Watershed Development Department. The database was generated by using cadastral map of the village as a base along with high resolution IRS LISS IV and Cartosat-1 merged satellite imagery. Later, an attempt will be made to uplink this LRI data generated at 1:7920 scale under Sujala-III Project to the proposed Landscape Ecological Units (LEUs) map.

The study was organized and executed by the ICAR- National Bureau of Soil Survey and Land Use Planning, Regional Centre, Bangalore under Generation of Land Resource Inventory Data Base Component-1 of the Sujala-III Project funded by the World Bank.

GEOGRAPHICAL SETTING

2.1 Location and Extent

The Yadgir Rf-1 microwatershed is located in the northern part of Karnataka in Yadgir Taluk & District, Karnataka State (Fig.2.1). It comprises parts of Allura B, Bheemanahalli and Baggalamadu villages. It lies between $16^{\circ} 55'$ and $16^{\circ} 56'$ North latitudes and $77^{\circ} 9'$ and $77^{\circ} 11'$ East longitudes, covering an area of about 306 ha, It is on the northern side of Yadgir town and is surrounded by Allura B on the south, west and north, Bheemanahalli on the east and Baggalamadu on southern side of the microwatershed.




Fig.2.1 Location map of Yadgir Rf-1 Microwatershed

2.2 Geology

Major rock formations observed in the microwatershed are granite gneiss (Figs.2.2a). Granite gneisses are essentially pink to gray and are coarse to medium grained. They

consist primarily of quartz, feldspar, biotite and hornblende. The gray granite gneisses are highly weathered, fractured and fissured upto a depth of about 10 m. Dolerite dykes and quartz veins are common with variable width and found to occur in Yadgir Rf-1 microwatershed.



Fig.2.2a Granite and granite gneiss rocks

2.3 Physiography

Physiographically, the area has been identified as granite gneiss based on geology. The area has been further subdivided into five landforms, *viz;* mounds/ridges, summits, side slopes and very gently sloping uplands, plains and valleys based on slope and its relief features. The elevation ranges from 469-508 m above MSL. The mounds and ridges are mostly covered by rock outcrops.

2.4 Drainage

The area is drained by several parallel streams like Bori, Amerja and Kanga which finally join the river Bhima along its course. Though, they are not perennial, during rainy season they carry large quantities of rain water. The microwatershed has only few small tanks which are not capable of storing the water that flows during the rainy season. Due to this, the ground water recharge is very much affected. This is reflected in the failure of many bore wells in the villages. If the available rain water is properly harnessed by constructing new tanks and recharge structures at appropriate places in the villages, then the drinking and irrigation needs of the area can be easily met. The drainage network is parallel to sub parallel and dendritic.

2.5 Climate

The Yadgir district lies in the northern plains of Karnataka and falls under semiarid tract of the state and is categorized as drought-prone with total annual rainfall of 866 mm (Table 2.1). Of the total rainfall, maximum of 652 mm is received during the south-west monsoon period from June to September; the north-east monsoon from

October to early December contributes about 138 mm and the remaining 76 mm during the rest of the year. The summer season starts during the middle of February and continues up to the first week of June. The period from December to the middle of February is the coldest season. December is the coldest month with mean daily maximum and minimum temperatures being 29.5⁰C and 10⁰C respectively. During peak summer, temperature shoots up to 45⁰C. Relative humidity varies from 26% in summer to 62% in winter. Rainfall distribution is shown in Figure 2.3. The average Potential Evapo-Transpiration (PET) is 141 mm and varies from a low of 81 mm in December to 199 mm in the month of May. The PET is always higher than precipitation in all the months except end of June to end of September. Generally, the Length of crop Growing Period (LGP) is 120-150 days and starts from 1st week of June to 4th week of October.

Table 2.1 Mean Monthly Rainfall, PET, 1/2 PET at Yadgir Taluk, Yadgir District

Sl. No.	Months	Rainfall	PET	1/2 PET
1	January	4.30	86.0	43.0
2	February	2.30	125.5	62.7
3	March	15.10	166.0	83.0
4	April	18.50	179.8	89.9
5	May	36.0	198.8	97.9
6	June	118.0	175.1	87.5
7	July	171.80	156.3	78.1
8	August	182.9	150.3	75.1
9	September	179.7	142.0	71.0
10	October	105.3	138.5	69.2
11	November	26.4	97.60	48.6
12	December	6.0	80.90	40.4
Total		866.3		




Fig 2.3 Rainfall distribution in Yadgir Taluk, Yadgir District

2.6 Natural Vegetation

The natural vegetation is sparse comprising few tree species, shrubs and herbs. The mounds, ridges and boulders occupy very sizeable area which is under thin to moderately thick forest vegetation. Still, there are some remnants of the past forest cover which can be seen in patches in some ridges and hillocks in the microwatershed (Fig 2.4).

Apart from the continuing deforestation, the presence of large population of goats, sheep and other cattle in the microwatershed is causing vegetative degradation of whatever little vegetation left in the area. The uncontrolled grazing has left no time for the regeneration of the vegetative cover. This leads to the accelerated rate of erosion on the hill slopes resulting in the formation of deep gullies in the foot slopes that eventually result in the heavy siltation of tanks and reservoirs in the microwatershed.



Fig 2.4 Natural vegetation of Yadgir Rf-1 Microwatershed

2.7 Land Utilization

About 72 per cent area (Table 2.2) in Yadgir district is cultivated at present. An area of about 2 per cent is permanently under pasture, 20 per cent under current fallows and 6 per cent under non-agricultural land and 5 per cent under currently barren. Forests occupy an area of about 7 per cent and the tree cover is in a very poor state. Most of the mounds, ridges and bouldery areas have very poor vegetative cover. Major crops grown in the area are sorghum, maize, cotton, sunflower, groundnut, red gram, mango, pomegranate, marigold and sapota. While carrying out land resource inventory, the land use/land cover particulars are collected from all the survey numbers and a current land use map of the microwatershed is prepared. The current land use map prepared shows the arable and non-arable lands, other land uses and different types of crops grown in the area. The current land use map of Yadgir Rf-1 microwatershed is presented in Fig.2.5. The different crops and cropping systems adopted in the microwatershed are presented in Figures 2.7 a & b.

Table 2.2 Land Utilization in Yadgir District

Sl. No.	Agricultural land use	Area (ha)	Per cent
1	Total geographical area	516088	-
2	Total cultivated area	373617	72.4
3	Area sown more than once	74081	14.3
4	Cropping intensity	-	119.8
5	Trees and grooves	737	0.14
6	Forest	33773	6.54
7	Cultivable wasteland	2385	0.46
8	Permanent Pasture land	11755	2.28
9	Barren land	27954	5.41
10	Non- Agriculture land	29623	5.73
11	Current Fallows	105212	20.4




Fig.2.5 Current Land Use map of Yadgir Rf-1 Microwatershed




Fig. 2.7 a. Different Crops and Cropping Systems in Yadgir Rf-1 Microwatershed



Fig. 2.7 b. Different Crops and Cropping Systems in Yadgir Rf-1 Microwatershed

SURVEY METHODOLOGY

The purpose of land resource inventory is to delineate similar areas (soil series and phases), which respond or expected to respond similarly to a given level of management. This was achieved in Yadgir Rf-1 microwatershed by the detailed study of all the soil characteristics (depth, texture, colour, structure, consistence, coarse fragments, porosity, soil reaction, soil horizons etc.) and site characteristics (slope of the land, erosion, drainage, occurrence of rock fragments etc.) followed by grouping of similar areas based on soil-site characteristics into homogeneous (management units) units, and showing the area extent and their geographic distribution on the microwatershed cadastral map. The detailed survey at 1:7920 scale was carried out in an area of 306 ha. The methodology followed for carrying out land resource inventory was as per the guidelines given in Soil Survey Manual (IARI, 1971; Soil Survey Staff, 2006; Natarajan *et al.*, 2015) which is briefly described below.

3.1 Base Maps

The detailed survey of the land resources occurring in the microwatershed was carried out by using digitized cadastral and IRS satellite imagery map as base supplied by KSRSAC. The cadastral map shows field boundaries with their survey numbers, location of tanks, streams and other permanent features of the area (Fig. 3.1). Apart from the cadastral map, remote sensing data products from Cartosat-1 and LISS IV merged at the scale of 1:7920 were used in conjunction with the cadastral map to identify the landscapes, landforms and other surface features. The imagery helped in the identification and delineation of boundaries between hills, uplands and lowlands, water bodies, forest and vegetated areas, roads, habitations and other cultural features of the area (Fig. 3.2). The cadastral map was overlaid on the satellite imagery (Fig. 3.3) that helps to identify the parcel boundaries and other permanent features. Apart from cadastral maps and images, toposheets of the area (1:50,000 scale) were also used for initial traversing, identification of geology and landforms, drainage features, present land use and also for selection of transects in the microwatershed.

3.2 Image Interpretation for Physiography

False Colour Composites (FCCs) of Cartosat-I and LISS-IV merged satellite data covering microwatershed area was visually interpreted using image interpretation elements and all the available collateral data with local knowledge. The delineated physiographic boundaries were transferred on to a cadastral map overlaid on satellite imagery. Physiographically, the area has been identified as granite gneiss landscape. It was divided into five landforms, *viz.*; ridges and mounds, gently and very gently sloping uplands and lowlands based on slope and image characteristics. They were further subdivided into physiographic/image interpretation units based on image characteristics. The image interpretation legend for physiography is given below.

Image Interpretation Legend for Physiography

G- Granite Gneiss Landscape

G1	Hills/ Ridges/ Mounds
G11	Summits
G12	Side slopes
G121	Side slopes with dark grey tones
G2	Uplands
G21	Summits
G22	Gently sloping uplands
G221	Gently sloping uplands, yellowish green (eroded)
G222	Gently sloping uplands, yellowish white (severely eroded)
G23	Very gently sloping uplands
G231	Very gently sloping uplands, yellowish green
G232	Very gently sloping uplands, medium green and pink
G233	Very gently sloping uplands, pink and green (scrub land)
G234	Very gently sloping uplands, medium greenish grey
G235	Very gently sloping uplands, yellowish white (eroded)
G236	Very gently sloping uplands, dark green
G237	Very gently sloping uplands, medium pink (coconut garden)
G238	Very gently sloping uplands, pink and bluish white (eroded)
G24	Valleys/ lowlands
G241	Valleys, pink tones
G242	Valleys gray mixed with pink tones




Fig 3.1 Scanned and Digitized Cadastral map of Yadgir Rf-1 Microwatershed




Fig.3.2 Satellite Image of Yadgir Rf-1 Microwatershed




Fig.3.3 Cadastral map overlaid on IRS PAN+LISS IV merged imagery of Yadgir Rf-1 Microwatershed

3.3 Field Investigation

The field boundaries and survey numbers given on the cadastral sheet were located on the ground by following permanent features like roads, cart tracks, *nallas*, streams, tanks etc., and wherever changes were noticed, they were incorporated on the microwatershed cadastral map. Preliminary traverse of the microwatershed was carried out with the help of cadastral map, imagery and toposheets. While traversing, landforms and physiographic units identified were checked and preliminary soil legend was prepared by studying soils at few selected places. Then, intensive traversing of each physiographic unit like hills, ridges, uplands and valleys was carried out. Based on the variability observed on the surface, transects (Fig. 3.4) were selected across the slope covering all the landform units in the microwatershed (Natarajan and Dipak Sarkar, 2010).




Fig: 3.4. Location of profiles in a transect

In the selected transect, soil profiles were located (Fig. 3.4) at closely spaced intervals to take care of any change in the land features like break in slope, erosion, gravel, stones etc. In the selected sites, profiles (vertical cut showing the soil layers from surface to the rock) were opened upto 200 cm or to the depth limited by rock or hard substratum and studied in detail for all their morphological and physical characteristics. The soil and site characteristics were recorded for all profile sites on a standard proforma as per the guidelines given in USDA Soil Survey Manual (Soil Survey Staff, 2012). Apart from the transect study, profiles were also studied at random, almost like in a grid pattern, outside the transect areas.

Based on the soil characteristics, the soils were grouped into different soil series. Soil series is the most homogeneous unit having similar horizons and properties and behaves similarly for a given level of management. Soil depth, texture, colour, kind of horizon and horizon sequence, calcareousness, amount and nature of gravel present, nature of substratum etc, were used as the major differentiating characteristics for

identifying soil series occurring in the area. The differentiating characteristics used for identifying the soil series are given in Table 3.1. Based on the above characteristics, 3 soil series were identified in the Yadgir Rf-1 microwatershed.

Table 3.1 Differentiating Characteristics used for identifying Soil Series
(Characteristics are of Series Control Section)

Soils of Granite gneiss Landscape							
Sl. no	Soil Series	Depth (cm)	Colour (moist)	Texture	Gravel (%)	Horizon sequence	Calcareous-ness
1	KKR (Kakalawar)	<25	7.5YR 4/3 10YR 6/3	sl	10-15	Ap-AC	-
2	HTK (Hattikuni)	25-50	10YR 4/6, 4/4 7.5YR 4/4, 3/3	sl	10-25	Ap-AC	-
3	YDR (Yadgir)	100-150	10YR 4/3,4/4, 2.5YR 4/3,5/3,	sl	<15	Ap-Ac	-

3.4 Soil Mapping

The area under each soil series was further separated into soil phases and their boundaries delineated on the cadastral map based on the variations observed in the texture of the surface soil, slope, erosion, presence of gravel, stoniness etc. A soil phase is a subdivision of soil series based mostly on surface features that affect its use and management. The soil mapping units are shown on the map (Fig.3.5) in the form of symbols. During the survey many profile pits, few minipits and a few auger bores representing different landforms occurring in the microwatershed were studied. In addition to the profile study, spot observations in the form of minipits, road cuts, terrace cuts etc., were studied to validate the soil boundaries on the soil map.

The soil map shows the geographic distribution of 4 mapping units representing 3 soil series occurring in the microwatershed. The soil map unit (soil legend) description is presented in Table 3.2. The soil phase map (management units) shows the distribution of 4 soil phases mapped in the microwatershed. Each mapping unit (soil phase) delineated on the map has similar soil and site characteristics. In other words, all the farms or survey numbers included in one phase will have similar management needs and have to be treated accordingly.

3.5 Land Management Units

The 4 soil phases identified and mapped in the microwatershed were grouped into 2 Land Management Units (LMU's) for the purpose of preparing a Proposed Crop Plan for sustained development of the microwatershed. The database (soil phases) generated under LRI was utilized for identifying Land Management Units (LMU's) based on the management needs. One or more than one soil site characteristic having influence on the management have been chosen for identification and delineation of LMUs. For Yadgir Rf-1 microwatershed, five soil and site characteristics, namely soil depth, soil texture,

slope, erosion and gravel content have been considered for defining LMUs. The land use classes are expected to behave similarly for a given level of management.

3.6 Laboratory Characterization

Soil samples were collected from representative master profiles for laboratory characterization by following the methods outlined in the Laboratory Manual (Sarma *et al*, 1987). Surface soil samples collected from farmer's fields (11 samples) for fertility status (major and micronutrients) at 320 m grid interval in the year 2017 were analyzed in the laboratory (Katyal and Rattan, 2003). By linking the soil fertility data to the survey numbers through GIS, soil fertility maps were generated by using Kriging method for the microwatershed.

Table 3.2 Soil map unit description of Yadgir Rf-1 Microwatershed

*Soil map unit No.	Soil Series	Soil Phase	Mapping Unit Description	Area in ha (%)
Soils of Granite and Granite Gneiss Landscape				
	KKR	Kakalawar soils are very shallow (<25 cm), well drained, have dark brown sandy loam soils occurring on very gently sloping uplands under cultivation		60 (19.7)
153		KKRbB2g1	Loamy sand surface, slope 1-3%, moderate erosion, gravelly (15-35%)	60 (19.7)
	HTK	Hattikuni soils are shallow (25-50 cm), well drained, have dark yellowish brown sandy loam soils occurring on very gently sloping uplands under cultivation		33 (10.83)
156		HTKbB2	Loamy sand surface, slope 1-3%, moderate erosion	33 (10.83)
	YDR	Yadgir soils are deep (100-150 cm), well drained, have brown to dark yellowish brown and olive brown, sodic sandy loam soils occurring on very gently sloping uplands under cultivation		47(15.5)
42		YDRcB2	Sandy loam surface, slope 1-3%, moderate erosion	33 (10.93)
154		YDRcB2g1	Sandy loam surface, slope 1-3%, moderate erosion, gravelly (15-35%)	14 (4.57)
999		Rock outcrops	Rock lands, both massive and bouldery with little or no soil	165 (53.97)




Fig 3.5 Soil Phase or Management Units - Yadgir Rf-1 Microwatershed

THE SOILS

Detailed information pertaining to the nature, extent and their distribution of different kinds of soils occurring in Yadgir Rf-1 microwatershed is provided in this chapter. The microwatershed area has been identified as granite gneiss landscape based on geology. In all, 3 soil series are identified. Soil formation is the result of the combined effect of environmental and terrain factors that are reflected in soil morphology. In the granite gneiss landscape, it is by parent material, relief and climate.

A brief description of each of the 3 soil series identified followed by 4 soil phases (management units) mapped under each series are furnished below. The physical and chemical characteristics of soil series identified in Yadgir Rf-1 microwatershed are given in Table 4.1 along with soil classification. The soils in any one map unit differ from place to place in their depth, texture, slope, gravelliness, erosion or any other site characteristic that affect management. The soil phase map can be used for identifying the suitability of areas for growing specific crops or for other alternative uses and also for deciding the type of conservation structures needed. The detailed information on soil and site-characteristics like soil depth, surface soil texture, slope, erosion, gravelliness, AWC, LCC etc, with respect to each of the soil phase identified is given village/survey number wise for the microwatershed in Appendix-I.

4.1 Soils of granite gneiss landscape

In this landscape, 3 soil series are identified and mapped. KKR series occupies maximum area of 60 ha (20%) followed by YDR 47 ha (16%) and HTK 33 ha (11%). Brief description of each series identified and number of soil phases mapped is given below.

4.1.1 Kakalawar (KKR) Series: Kakalawar soils are very shallow (<25cm), well drained, have dark brown to light brown, sandy loam soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation. The Kakalawar series has been classified as a member of the mixed, isohyperthermic family of Lithic Ustipsamments.

The thickness of the soil is less than 25 cm. Its colour is in 10 YR and 7.5 YR hue with value 4 to 6 and chroma 3 to 4. The texture varies from loamy sand to sand. The available water capacity is very low (<50 mm/m). One phase was identified and mapped.



Landscape and Soil Profile characteristics of Kakalawar (KKR) Series

4.1.2 (HTK) Series: Hattikuni soils are shallow (25-50 cm), well drained, have dark brown to dark yellowish brown sandy loam soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation. The Hattikuni series has been classified as a member of the mixed, isohyperthermic family of Lithic Ustipsammments.

The thickness of the soil ranges from 36 to 50 cm. The thickness of A horizon ranges from 8 to 12 cm. Its colour is in 10YR and 7.5 YR hue with value 3 to 4 and chroma 4 to 6. The texture varies from loamy sand to sandy loam. The thickness of subsurface horizon ranges from 28 to 42 cm. Its colour is in 10 YR and 7.5 YR hue with value 3 to 4 and chroma 4 to 6. Its texture varies from loamy sand to sand and sandy loam. The available water capacity is very low (<50 mm/m). One phase was identified and mapped



Landscape and Soil Profile characteristics of Hattikuni (HTK) Series

4.1.3 Yadgir (YDR) Series: Yadgir soils are deep (100-150 cm), well drained, have very dark yellowish brown to light olive brown, sodic sandy loam soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation. The Yadgir series has been classified as a member of the coarse-loamy, mixed, isohyperthermic family of Fuluventic Haplustepts.

The thickness of the soil ranges from 105 to 145 cm. The thickness of A horizon ranges from 6 to 10 cm. Its colour is in 10 YR hue with value 4 and chroma 3. The texture is loamy sand. The thickness of subsurface horizons ranges from 95 to 130 cm. Its colour is in 10 YR and 2.5 Y hue with value 4 to 5 and chroma 3 to 4. Texture is sandy loam and sandy clay loam and are sodic soils. The available water capacity is low (51-100 mm/m). Two phases were identified and mapped.



Landscape and Soil Profile characteristics of Yadgir (YDR) Series

Table: 4.1 Physical and Chemical Characteristics of Soil Series identified in Yadgir Rf-1 microwatershed

Soil Series: Kakalawar (KKR), Pedon: R-7

Location: 16°50'25.9"N 77°15'97.1"E, Yampada village, Gurumitkal hobli, Yadgir taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bengaluru **Classification:** Mixed, isohyperthermic, Lithic Ustipsamments

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-22	Ap	83.81	10.37	5.82	17.31	20.65	17.91	5.67	22.27	10-20	ls	9.77	4.65

Depth (cm)	pH (1:2.5)			E.C. (1:2.5) dS m ⁻¹	O.C. %	CaCO ₃ %	Exchangeable bases					CEC	CEC/Clay	Base saturation %	ESP %
	Water	CaCl ₂	M KCl				Ca	Mg	K	Na	Total				
							cmol kg ⁻¹								
0-22	5.85	-	-	0.027	0.19	-	0.72	0.21	0.62	0.03	1.58	2.6	0.45	60.90	1.17

Contd...

Soil Series: Hattikuni (HTK), Pedon: R-7

Location: 16°50'46.5"N 77°10'16.4"E, Yaddalli village, Hattikuni hobli, Yadgir taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bengaluru **Classification:** Mixed, isohyperthermic Lithic Ustipsamments

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-12	Ap	90.89	5.62	3.49	8.50	13.46	29.86	29.55	9.51	20	s	7.73	3.16
12-22	A1	89.97	6.53	3.50	7.19	13.48	29.48	29.79	10.03	20	s	8.00	3.05
22-45	A2	87.20	6.43	6.38	11.09	14.42	31.55	7.16	22.98	40	ls	7.67	3.96

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO ₃	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP
	Water	CaCl ₂	M KCl	dS m ⁻¹	%	%	Ca	Mg	K	Na	Total			%	%
							cmol kg ⁻¹								
0-12	6.81	-	-	0.062	0.07	-	2.35	0.50	0.16	0.01	3.02	3.0	0.86	100	0.38
12.0-22	6.80	-	-	0.050	0.21	-	1.67	0.30	0.09	0.01	2.07	2.4	0.69	86.30	0.45
22-45	6.85	-	-	0.044	0.19	-	1.82	0.42	0.10	0.06	2.40	2.6	0.41	92.41	2.17

Contd...

Soil Series: Yadgir (YDR) **Pedon:** R-5

Location: 16°35'43.6"N 77°17'06.4"E, Kanikal village, Balichakra hobli, Yadgir taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bengaluru **Classification:** Coarse-loamy, mixed, isohyperthermic Fuluventic Haplustepts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-14	Ap	73.39	11.31	15.30	6.76	20.27	24.87	15.66	5.83	-	sl	12.14	7.22
14-43	A2	86.59	8.77	4.64	23.19	26.92	14.11	15.22	7.16	-	ls	6.97	2.68
43-89	Bw1	80.41	3.75	15.84	8.06	13.47	36.73	15.71	6.43	-	sl	22.84	10.18
89-110	Bw2	63.55	5.40	31.05	8.10	23.05	19.00	9.87	3.53	15-35	scl	38.46	17.70

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO ₃	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP			
	Water	CaCl ₂	M KCl				dS m ⁻¹	%	%	Ca	Mg					K	Na	Total
										cmol kg ⁻¹								
0-14	9.47	-	-	0.371	0.32	1.30	14.71	4.28	0.38	1.54	20.91	12.70	0.83	165	4.86			
14-43	7.25	-	-	0.114	0.56	0.00	2.29	0.86	0.07	0.03	3.25	3.40	0.73	96	0.31			
43-89	10.30	-	-	0.820	0.16	0.52	1.70	0.98	0.15	6.62	9.45	8.61	0.54	110	30.77			
89-110	10.80	-	-	1.440	0.12	0.91	1.02	2.00	0.29	14.43	17.74	16.17	0.52	110	35.688			

INTERPRETATION FOR LAND RESOURCE MANAGEMENT

The most important soil and site characteristics that affect the land use and conservation needs of an area are land capability, soil depth, soil texture, coarse fragments, available water capacity, soil slope, soil erosion, soil reaction etc. These are interpreted from the data base generated through land resource inventory and several thematic maps are generated. These would help in identifying the areas suitable for growing crops and, soil and water conservation measures and structures needed thus helping to maintain good soil health for sustained crop production. The various interpretative and thematic maps generated are described below.

5.1 Land Capability Classification

Land capability classification is an interpretative grouping of soil map units (soil phases) mainly based on inherent soil characteristics, external land features and environmental factors that limit the use of land for agriculture, pasture, forestry, or other uses on a sustained basis (IARI, 1971). The land and soil characteristics used to group the land resources in an area into various land capability classes, subclasses and units are

Soil Characteristics: Depth, texture, gravelliness, calcareousness.

Land characteristics: Slope, erosion, drainage, rock outcrops.

Climate: Total rainfall and its distribution, and length of crop growing period.

The Land capability classification system is divided into land capability classes, subclasses and units based on the level of information available. Eight land capability classes are recognized. They are

Class I: They are very good lands that have no limitations or very few limitations that restrict their use.

Class II: They are good lands that have minor limitations and require moderate conservation practices.

Class III: They are moderately good lands that have moderate limitations that reduce the choice of crops or that require special conservation practices.

Class IV: They are fairly good lands that have very severe limitations that reduce the choice of crops or that require very careful management.

Class V: Soils in these lands are not likely to erode, but have other limitations like wetness that are impractical to remove and as such not suitable for agriculture, but suitable for pasture or forestry with minor limitations.

Class VI: The lands have severe limitations that make them generally unsuitable for cultivation, but suitable for pasture or forestry with moderate limitations.

Class VII: The lands have very severe limitations that make them unsuitable for cultivation, but suitable for pasture or forestry with major limitations.

Class VIII: Soil and other miscellaneous areas (rock lands) that have very severe limitations that nearly preclude their use for any crop production, but suitable for wildlife, recreation and installation of wind mills.

The land capability subclasses are recognised based on the dominant limitations observed within a given land capability class. The subclasses are designated by adding a lower case letter like 'e', 'w', 's', or 'c' to the class numeral. The subclass "e" indicates that the main hazard is risk of erosion, "w" indicates drainage or wetness as a limitation for plant growth, "s" indicates shallow soil depth, coarse or heavy textures, calcareousness, salinity/alkalinity or gravelliness and "c" indicates limitation due to climate.

The land capability subclasses have been further subdivided into land capability units based on the kinds of limitations present in each subclass. Ten land capability units are used in grouping the soil map units. They are stony or rocky (0), erosion hazard (slope, erosion) (1), coarse texture (sand, loamy sand, sandy loam) (2), fine texture (cracking clay, silty clay) (3), slowly permeable subsoil (4), coarse underlying material (5), salinity/alkali (6), stagnation, overflow, high ground water table (7), soil depth (8) and fertility problems (9). The capability units thus identified have similar soil and land characteristics that respond similarly to a given level of management. The soils of the microwatershed have been classified upto land capability subclass level.

The 4 soil map units identified in the Yadgir Rf-1 microwatershed are grouped under 2 land capability classes and 2 subclasses. An area about 141 ha (46%) in the microwatershed is suitable for agriculture and about 165 ha (54%) covered by Rock outcrops in the microwatershed (Fig. 5.1).




Fig. 5.1 Land Capability map of Yadgir Rf-1 Microwatershed

Moderately good lands (Class III) cover an area of about 33 ha (11%) and are distributed in the northeastern, eastern and southeastern part of the microwatershed. They have moderate limitations of soil and erosion. Fairly good lands (Class IV) cover an area of about 108 ha (35%) and are distributed in the major part of the cultivated area. They have moderate limitations of soil and erosion.

5.2 Soil Depth

Soil depth refers to the depth of the soil occurring above the parent material or hard rock. The depth of the soil determines the effective rooting depth for plants and in accordance with soil texture, mineralogy and gravel content, the capacity of the soil column to hold water and nutrient availability. Soil depth is one of the most important soil characteristic that is used in differentiating soils into different soil series. The soil depth classes used in identifying soils in the field are very shallow (<25 cm), shallow (25-50 cm), moderately shallow (50-75 cm), moderately deep (75-100 cm), deep (100-150 cm) and very deep (>150 cm). They were used to classify the soils into different depth classes and a soil depth map was generated. The area extent and their geographical distribution in the microwatershed is given in Fig. 5.2.




Fig. 5.2 Soil Depth map of Yadgir Rf-1 Microwatershed

Very shallow (<25cm) soils cover an area of 60 ha (20%) and are distributed in the major part of the cultivated area. Shallow (25-50 cm) soils cover an area of 33 ha (11%) and are distributed in the northeastern, eastern, southeastern part of the microwatershed. Deep (100-150 cm) soils cover an area of 47 ha (16%) and are distributed in the central, eastern and northeastern part of the microwatershed.

The most productive lands of 47 ha (16%) with respect to soil rooting depth where all climatically adapted annual and perennial crops can be grown are deep (> 100 cm) soils. Shallow and very shallow soils cover about 93 ha (31%) area where short duration crops can be grown and probability of crop failure is high

5.3 Surface Soil Texture

Texture is an expression to indicate the coarseness or fineness of the soil as determined by the relative proportion of primary particles of sand, silt and clay. It has a direct bearing on the structure, porosity, adhesion and consistence. The surface layer of a soil to a depth of about 25 cm is the layer that is most used by crops and plants. The surface soil textural class provides a guide to understanding soil-water retention and availability, nutrient holding capacity, infiltration, workability, drainage, physical and chemical behaviour, microbial activity and crop suitability. The textural classes used for LRI were used to classify and a surface soil texture map was generated. The area extent and their geographical distribution in the microwatershed is shown in Figure 5.3.




Fig. 5.3 Surface Soil Texture map of Yadgir Rf-1 Microwatershed

An area of about 93 ha (31%) is sandy and is distributed in the major part of the cultivated area. An area of 47 ha (15%) has soils that are loamy at the surface and occur in the central, northeastern and eastern part of the microwatershed.

An area of 47 ha (16%) the microwatershed is most productive with respect to surface soil texture. The productive lands are loamy soils (16%) which also have high potential for soil-water retention and nutrient availability but have no drainage or other physical problems. The sandy soils (31%) are problematic but productive for root and tuber crops, but these soils have the major limitation of moisture and nutrient retention capacity, hence frequent and shallow irrigation with balanced fertilizer application is to be followed in order to get better crop yields.

5.4 Soil Gravelliness

Gravel is the term used for describing coarse fragments between 2 mm and 7.5 cm diameter and stones for those between 7.5 cm and 25 cm. The presence of gravel and stones in soil reduces the volume of soil responsible for moisture and nutrient storage, drainage, infiltration and runoff, and hinders plant growth by impeding root growth and seedling emergence, intercultural operations and farm mechanization. The gravelliness classes used in LRI were used to classify the soils and using these classes a gravelliness map was generated. The area extent and their geographic distribution in the microwatershed is shown in Figure 5.4.




Fig. 5.4 Soil Gravelliness map of Yadgir Rf-1 Microwatershed

An area of about 67 ha (22%) is non gravelly (<15%) and are distributed in the eastern, northeastern and southeastern part of the microwatershed and about 74 ha (24%) is gravelly (15-35%) soils and are distributed in the major part of the cultivated area.

The most productive soils (22%) that are non gravelly (<15%), where all climatically adapted long duration crops can be grown.

5.5 Available Water Capacity

The soil available water capacity (AWC) is estimated based on the ability of the soil column to retain water between the tensions of 0.33 and 15 bar in a depth of 100 cm or the entire solum if the soil is shallower. The AWC of the soils (soil series) as estimated by considering the soil texture, mineralogy, soil depth and gravel content (Sehgal *et al.*, 1990) and accordingly the soil map units were grouped into five AWC classes *viz*, very low (<50 mm/m), low (50-100 mm/m), medium (100-150 mm/m), high (150-200 mm/m) and very high (>200 mm/m) and using these values, an AWC map was generated. The area extent and their geographic distribution of different AWC classes in the microwatershed is given in Figure 5.5.




Fig. 5.5 Soil Available Water Capacity map of Yadgir Rf-1 Microwatershed

An area of about 47 ha (16%) and 93 ha (31%) in the microwatershed has soils that are low (51-100 mm/m) and very low (<50mm/m) in available water capacity and are distributed in the major part of the cultivated area.

An area of 140 ha (46%) in the microwatershed has problematic with regard to available water capacity. Here, only short duration crops can be grown and the probability of crop failure is very high. These areas are best put to other alternative uses.

5.6 Soil Slope

Soil slope refers to the inclination of the surface of the land. It is defined by gradient, shape and length, and is an integral feature of any soil as a natural body. Slope is considered important in soil genesis, land use and land development. The length and gradient of slope influences the rate of runoff, infiltration, erosion and deposition. The soil map units were grouped into two slope classes and a slope map was generated showing the area extent and their geographic distribution in the microwatershed (Fig. 5.6).

Entire cultivated area falls under very gently sloping (1-3% slope) lands in the microwatershed.




Fig. 5.6 Soil Slope map of Yadgir Rf-1 Microwatershed

Entire cultivated area in the microwatershed has high potential in respect of soil slopes. In these areas, all climatically adapted annual and perennial crops can be grown without much soil and water conservation and other land development measures.

5.7 Soil Erosion

Soil erosion refers to the wearing away of the earth's surface by the forces of water, wind and ice involving detachment and transport of soil by raindrop impact. It is used for accelerated soil erosion resulting from disturbance of the natural landscape by burning, excessive grazing and indiscriminate felling of forest trees and tillage, all usually by man. The erosion classes showing an estimate of the current erosion status as judged from field observations in the form of rills, gullies or a carpet of gravel on the surface are recorded. Four erosion classes, viz, slight erosion (e1), moderate erosion (e2), severe erosion (e3) and very severe erosion (e4) are recognized. The soil map units were grouped into different erosion classes and a soil erosion map generated. The area extent and their spatial distribution in the microwatershed is given in Figure 5.7.

Entire cultivated area is moderately eroded (e2 class) in the microwatershed.

Entire cultivated area in the microwatershed is problematic because of moderate erosion. For these areas, taking up soil and water conservation and other land development measures are needed.




Fig. 5.7 Soil Erosion map of Yadgir Rf-1 Microwatershed

FERTILITY STATUS

Soil fertility plays an important role in increasing crop yield. The adoption of high yielding varieties that require high amounts of nutrients has resulted in deficiency symptoms in crops and plants due to imbalanced fertilization and poor inherent fertility status as these areas are characterised by low rainfall and high temperatures. Hence, it is necessary to know the fertility (macro and micro nutrients) status of the soils of the watersheds for assessing the kind and amount of fertilizers required for each of the crop intended to be grown. For this purpose, the surface soil samples collected from the grid points (one soil sample at every 320 m interval) all over the microwatershed through land resource inventory in the year 2017 were analysed for pH, EC, organic carbon, available phosphorus and potassium, and for micronutrients like zinc, boron, copper, iron and manganese, and secondary nutrient sulphur.

Soil fertility data generated has been assessed and individual maps for all the nutrients for the microwatershed have been generated using Kriging method under GIS. The village/survey number wise fertility data for the microwatershed is given in Appendix-II.

6.1 Soil Reaction (pH)

The soil analysis of the Yadgir Rf-1 microwatershed for soil reaction (pH) showed that an area of about 61 ha (20%) is moderately acid (pH 5.5- 6.5) and are distributed in the major part of the cultivated area. About 20 ha (7%) is strongly acid (pH 5.0-5.5) and are distributed in the central and eastern part of the microwatershed. An area of about 60 ha (20%) is neutral (6.5-7.3) and are distributed in the northeastern, southern and southeastern part of the microwatershed. (fig.6.1). In all, major area of about 81 ha is under acidic and 60 ha is neutral.

6.2 Electrical Conductivity (EC)

The Electrical Conductivity of the soils in the microwatershed area is $<2 \text{ dS m}^{-1}$ (Fig 6.2) and as such the soils are non-saline.

6.3 Organic Carbon

The soil organic carbon content (an index of available Nitrogen) in the soils of the microwatershed is medium (0.5-0.75%) in about 21 ha (7%) and are distributed in the southeastern part of the microwatershed and 120 ha (39%) is low ($<0.5\%$) in organic carbon and are distributed in the major part of cultivated area (Fig. 6.3).




Fig.6.1 Soil Reaction (pH) map of Yadgir Rf-1 Microwatershed




Fig.6.2 Electrical Conductivity (EC) map of Yadgir Rf-1 Microwatershed




Fig.6.3 Soil Organic Carbon map of Yadgir Rf-1 Microwatershed

6.4 Available Phosphorus

Available phosphorus content is medium (23-57 kg/ha) covering an area of about <1 ha (<1%) and occur in the northern part of the microwatershed and high (>57 kg/ha) covering an area of about 141 ha (46%) and occur in the major part of the cultivated area (Fig. 6.4).

6.5 Available Potassium

Entire cultivated area in the microwatershed is medium (145-337 kg/ha) in available potassium content (Fig.6.5).

6.6 Available Sulphur

Available sulphur is medium (10-20 ppm) in an area of about 117 ha (38%) and occur in the major part of the cultivated area and low (<10 ppm) covering an area of about 24 ha (8%) and occur in the northeastern part of the microwatershed (Fig. 6.6).

6.7 Available Boron

Available boron content is medium (0.5-1.0 ppm) in an area of 11 ha (3%) and are distributed in the southeastern part of the microwatershed. An area of about 130 ha (43%) is low (<0.5 ppm) in available boron and are distributed in the major part of the cultivated area (Fig. 6.7).

6.8 Available Iron

Available iron content is sufficient (>4.5 ppm) in the entire cultivated area of the microwatershed (Fig 6.8).

6.9 Available Manganese

Available manganese content is sufficient (>1.0 ppm) in the entire cultivated area of the microwatershed (Fig 6.9).

6.10 Available Copper

Available copper content is sufficient (>0.2 ppm) in the entire cultivated area of the microwatershed area (Fig 6.10).




Fig.6.4 Soil Available Phosphorus map of Yadgir Rf-1 Microwatershed




Fig.6.5 Soil Available Potassium map of Yadgir Rf-1 Microwatershed




Fig.6.6 Soil Available Sulphur map of Yadgir Rf-1 Microwatershed




Fig.6.7 Soil Available Boron map of Yadgir Rf-1 Microwatershed




Fig.6.8 Soil Available Iron map of Yadgir Rf-1 Microwatershed




Fig.6.9 Soil Available Manganese map of Yadgir Rf-1 Microwatershed




Fig.6.10 Soil Available Copper map of Yadgir Rf-1 Microwatershed

6.11 Available Zinc

Available zinc content is deficient (<0.6 ppm) in the entire cultivated area of the microwatershed (Fig 6.11).




Fig.6.11 Soil Available Zinc map of Yadgir Rf-1 Microwatershed

LAND SUITABILITY FOR MAJOR CROPS

The soil and land resource units (soil phases) of Yadgir Rf-1 microwatershed were assessed for their suitability for growing food, fodder, fibre and other horticulture crops by following the procedure as outlined in FAO, 1976 and 1983. Crop requirements were developed for each of the crop from the available research data and also by referring to Naidu *et. al.* (2006) and Natarajan *et. al* (2015). The soil and land characteristics were matched with the crop requirement to arrive at the crop suitability. The soil and land characteristics (Table 7.1) and crop requirement (Table 7.2 to 7.30) are given at the end of the chapter. In FAO land suitability classification, two orders are recognized. Order S-Suitable and Order N-Not suitable. The orders have classes, subclasses and units. Order S has three classes, Class S1-Highly Suitable, Class S2-Moderately Suitable and Class S3- Marginally Suitable. Order N has two classes, N1-Currently not Suitable and N2-Permanently not Suitable. There are no subclasses within the Class S1 as they will have very minor or no limitations for crop growth. Classes S2, S3 and N1 are divided into subclasses based on the kinds of limitations encountered. The limitations that affect crop production are 'c' for erratic rainfall and its distribution and length of growing period (LGP), 'e' for erosion hazard, 'r' for rooting condition, 't' for lighter or heavy texture, 'g' for gravelliness or stoniness, 'n' for nutrient availability, 'l' for topography, 'm' for moisture availability, 'w' for drainage and 'z' for calcareousness. These limitations are indicated as lower case letters to the Class symbol. For example, moderately suitable lands with the limitations of soil depth and erosion are designated as S2re. For the microwatershed, the soil mapping units were evaluated and classified up to subclass level.

Using the above criteria, the soil map units of the microwatershed were evaluated and land suitability maps for 29 major annual and perennial crops were generated. The detailed information on the kind of suitability of each of the soil phase for the crops assessed are given village/ survey number wise for the microwatershed in Appendix-IV.

7.1 Land Suitability for Sorghum (*Sorghum bicolor*)

Sorghum is one of the major food crop grown in Karnataka in an area of 10.47 lakh ha in Bijapur, Gulbarga, Raichur, Bidar, Belgaum, Dharwad, Bellary, Chitradurga, Mysore and Tumakuru districts. The crop requirements for growing sorghum (Table 7.2) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing sorghum was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.1.

An area of about 80 ha (26%) is marginally suitable (Class S3) for growing sorghum and are distributed in the major part of the cultivated area with moderate limitations of rooting depth, nutrient availability and texture. Currently not suitable (Class

N1) lands occur in an area of 60 ha (20%) and are distributed in the central, southern and southeastern part of the microwatershed with severe limitation of rooting depth.




Fig. 7.1 Land Suitability map of Sorghum

7.2 Land Suitability for Maize (*Zea mays*)

Maize is one of the most important food crop grown in an area of 13.37 lakh ha in almost all the districts of the State. The crop requirements for growing maize (Table 7.3) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing maize was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.2.

An area of about 80 ha (26%) is marginally suitable (Class S3) for growing maize and are distributed in the major part of the cultivated area with moderate limitations of rooting depth, nutrient availability and texture. Currently not suitable (Class N1) lands occur in an area of 60 ha (20%) and are distributed in the central, southern and southeastern part of the microwatershed with severe limitation of rooting depth.




Fig. 7.2 Land Suitability map of Maize

7.3 Land Suitability for Bajra (*Pennisetum glaucum*)

Bajra is one of the most important millet crop grown in an area of 2.34 lakh ha in the northern districts of Karnataka state. The crop requirements for growing bajra (Table 7.4) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing bajra was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.3.

An area of about 80 ha (26%) is marginally suitable (Class S3) for growing bajra and are distributed in the major part of the cultivated area with moderate limitations of rooting depth, nutrient availability and texture. Currently not suitable (Class N1) lands occur in an area of 60 ha (20%) and are distributed in the central, southern and southeastern part of the microwatershed with severe limitation of rooting depth.




Fig. 7.3 Land Suitability map of Bajra

7.4 Land Suitability for Groundnut (*Arachis hypogaea*)

Groundnut is one of the major oilseed crop grown in an area of 6.54 lakh ha in Karnataka in most of the districts either as rainfed or irrigated crop. The crop requirements for growing groundnut (Table 7.5) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing groundnut was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.4.

Marginally suitable lands (Class S3) for growing groundnut occupy an area of about 33 ha (11%) and are distributed in the northeastern, eastern and southeastern part of the microwatershed. They have moderate limitation of rooting depth. About 107 ha (35%) is currently not suitable (Class N1) for growing groundnut and are distributed in the major part of the cultivated area with severe limitations of rooting depth and nutrient availability.




Fig. 7.4 Land Suitability map of Groundnut

7.5 Land Suitability for Sunflower (*Helianthus annus*)

Sunflower is one of the most important oilseed crop grown in an area of 3.56 lakh ha in the State in all the districts. The crop requirements for growing sunflower (Table 7.6) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing sunflower was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.5.

Currently not suitable (Class N1) lands for growing sunflower occur in the entire cultivated area which is about 140 ha (46%) with severe limitations of rooting depth and nutrient availability.




Fig. 7.5 Land Suitability map of Sunflower

7.6 Land Suitability for Red gram (*Cajanus Cajan*)

Redgram is one of the most important pulse crop grown in an area of 7.28 lakh ha in almost all the districts of the State. The crop requirements for growing red gram (Table 7.7) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing redgram was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.6.

An area of about 47 ha (16%) is marginally suitable (Class S3) for growing redgram and are distributed in the central, northeastern and eastern part of the microwatershed. They have moderate limitation of nutrient availability. About 93 ha (31%) is currently not suitable (Class N1) and are distributed in the major part of the cultivated area with severe limitation of rooting depth.




Fig. 7.6 Land Suitability map of Redgram

7.7 Land Suitability for Bengal gram (*Cicer arietinum*)

Bengal gram is one of the most important pulse crop grown in about 9.39 lakh ha area in Bijapur, Raichur, Kalaburgi, Dharwad, Belgaum and Bellary districts. The crop requirements for growing Bengal gram (Table 7.8) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing Bengal gram was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.7.

Currently not suitable (Class N1) lands for growing bengal gram occur in the entire cultivated area of about 141 ha (46%) with severe limitations of rooting depth and nutrient availability.




Fig. 7.7 Land Suitability map of Bengal gram.

7.8 Land Suitability for Cotton (*Gossypium hirsutum*)

Cotton is one of the most important fibre crop grown in the State in about 8.75 lakh ha area in Raichur, Dharwad, Belgaum, Kalaburgi, Bijapur, Bidar, Bellary, Chitradurga and Chamarajnagar districts. The crop requirements for growing cotton (Table 7.9) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing cotton was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.8.

Currently not suitable (Class N1) lands for growing cotton occur in the entire cultivated area of about 141 ha (46%) with severe limitations of rooting depth and texture.




Fig. 7.8 Land Suitability map of Cotton

7.9 Land Suitability for Chilli (*Capsicum annuum*)

Chilli is one of the most important spice crop grown in about 0.42 lakh ha in Karnataka State. The crop requirements for growing chilli (Table 7.10) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing chilli was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.9.

Marginally suitable lands (Class S3) for growing chilli occupy an area of about 33 ha (11%) and are distributed in the northeastern, eastern and southeastern part of the microwatershed. They have moderate limitation of rooting depth. About 107 ha (35%) is currently not suitable (Class N1) for growing chilli and are distributed in the major part of the cultivated area with severe limitations of rooting depth and nutrient availability.




Fig 7.9 Land Suitability map of Chilli

7.10 Land Suitability for Tomato (*Lycopersicon esculentum*)

Tomato is one of the most important vegetable crop grown in about 0.61 lakh ha covering almost all the district of the state. The crop requirements for growing tomato (Table 7.11) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing tomato was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.10.

Marginally suitable lands (Class S3) for growing tomato occupy an area of about 33 ha (11%) and are distributed in the northeastern, eastern and southeastern part of the microwatershed. They have moderate limitation of rooting depth. About 107 ha (35%) is currently not suitable (Class N1) for growing tomato and are distributed in the major part of the cultivated area with severe limitations of rooting depth and nutrient availability.




Fig 7.10 Land Suitability map of Tomato

7.11 Land Suitability for Brinjal (*Solanum melongena*)

Brinjal is one of the most important vegetable crop grown in the state. The crop requirements for growing brinjal (Table 7.12) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing Brinjal was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.11.

Marginally suitable lands (Class S3) for growing brinjal occupy an area of about 33 ha (11%) and are distributed in the northeastern, eastern and southeastern part of the microwatershed. They have moderate limitation of rooting depth. About 107 ha (35%) is currently not suitable (Class N1) for growing brinjal and are distributed in the major part of the cultivated area with severe limitations of rooting depth and nutrient availability.




Fig 7.11 Land Suitability map of Brinjal

7.12 Land Suitability for Onion (*Allium cepa L.*)

Onion is one of the most important vegetable crop grown in the state. The crop requirements for growing onion (Table 7.13) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing onion was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.12.

Marginally suitable lands (Class S3) for growing onion occupy an area of about 33 ha (11%) and are distributed in the northeastern, eastern and southeastern part of the microwatershed. They have moderate limitation of rooting depth. About 107 ha (35%) is currently not suitable (Class N1) for growing onion and are distributed in the major part of the cultivated area with severe limitations of rooting depth and nutrient availability.



Fig 7.12 Land Suitability map of Onion

7.13 Land Suitability for Bhendi (*Abelmoschus esculentus*)

Bhendi is one of the most important vegetable crop grown in the state. The crop requirements for growing bhendi (Table 7.14) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing bhendi was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.13.

Marginally suitable lands (Class S3) for growing bhendi occupy an area of about 33 ha (11%) and are distributed in the northeastern, eastern and southeastern part of the microwatershed. They have moderate limitation of rooting depth. About 107 ha (35%) is currently not suitable (Class N1) for growing bhendi and are distributed in the major part of the cultivated area with severe limitations of rooting depth and nutrient availability.




Fig 7.13 Land Suitability map of Bhendi

7.14 Land Suitability for Drumstick (*Moringa oleifera*)

Drumstick is one of the most important vegetable crop grown in about 2403 ha in the state. The crop requirements for growing drumstick (Table 7.15) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing drumstick was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.14.

Currently not suitable (Class N1) lands for growing drumstick occur in the entire cultivated area of about 140 ha (46%) with severe limitations of rooting depth and nutrient availability.




Fig 7.14 Land Suitability map of Drumstick

7.15 Land Suitability for Mango (*Mangifera indica*)

Mango is one of the most important fruit crop grown in an area of 1.73 lakh ha in almost all the districts of the State. The crop requirements (Table 7.16) for growing mango were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing mango was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed is given in Figure 7.15.

Currently not suitable (Class N1) lands for growing mango occur in the entire cultivated area of about 140 ha (46%) with severe limitations of rooting depth and nutrient availability.




Fig. 7.15 Land Suitability map of Mango

7.16 Land Suitability for Guava (*Psidium guajava*)

Guava is one of the most important fruit crop grown in an area of 0.06 lakh ha in almost all the districts of the State. The crop requirements (Table 7.17) for growing guava were matched with the soil-site characteristics (7.1) and a land suitability map for growing guava was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.16.

Currently not suitable (Class N1) lands for growing guava occur in the entire cultivated area of about 140 ha (46%) with severe limitations of rooting depth and nutrient availability.




Fig. 7.16 Land Suitability map of Guava

7.17 Land Suitability for Sapota (*Manilkara zapota*)

Sapota is one of the most important fruit crop grown in an area of 29373 ha in almost all the districts of the State. The crop requirements (Table 7.18) for growing sapota were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing sapota was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.17.

Currently not suitable (Class N1) lands for growing sapota occur in the entire cultivated area of about 140 ha (46%) with severe limitations of rooting depth and nutrient availability.




Fig. 7.17 Land Suitability map of Sapota

7.18 Land Suitability for Pomegranate (*Punica granatum*)

Pomegranate is one of the most important fruit crop commercially grown in about 18488 ha in Karnataka, mainly in Bijapur, Bagalkot, Koppal, Gadag and Chitradurga districts. The crop requirements for growing pomegranate (Table 7.19) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing pomegranate was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.18.

Currently not suitable (Class N1) lands for growing pomegranate occur in the entire cultivated area of about 140 ha (46%) with severe limitations of rooting depth and nutrient availability.




Fig 7.18 Land Suitability map of Pomegranate

7.19 Land Suitability for Musambi (*Citrus limetta*)

Musambi is one of the important fruit crop grown in an area of 3446 ha in almost all the districts of the State. The crop requirements for growing musambi (Table 7.20) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing musambi was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.19.

Currently not suitable (Class N1) lands for growing musambi occur in the entire cultivated area of about 140 ha (46%) with severe limitations of rooting depth and nutrient availability.




Fig. 7.19 Land Suitability map of Musambi

7.20 Land Suitability for Lime (*Citrus sp*)

Lime is one of the most important fruit crop grown in an area of 0.11 lakh ha in almost all the districts of the State. The crop requirements for growing lime (Table 7.21) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing lime was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7. 20.

Currently not suitable (Class N1) lands for growing lime occur in the entire cultivated area of about 140 ha (46%) with severe limitations of rooting depth and nutrient availability.




Fig. 7.20 Land Suitability map of Lime

7.21 Land Suitability for Amla (*Phyllanthus emblica*)

Amla is one of the medicinal fruit crop grown in almost all the districts of the State. The crop requirements for growing amla (Table 7.22) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing amla was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.21.

Marginally suitable lands (Class S3) for growing amla occupy an area of about 33 ha (11%) and are distributed in the northeastern, eastern and southeastern part of the microwatershed. They have moderate limitations of rooting depth and texture. About 107 ha (35%) is currently not suitable (Class N1) for growing amla and are distributed in the major part of the cultivated area with severe limitations of rooting depth and nutrient availability.




Fig. 7.21 Land Suitability map of Amla

7.22 Land Suitability for Cashew (*Anacardium occidentale*)

Cashew is one of the most important plantation nut crop grown in an area of 0.7 lakh ha in almost all the districts of the State. The crop requirements for growing cashew (Table 7.23) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing cashew was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.22.

Currently not suitable (Class N1) lands for growing cashew occur in the entire cultivated area of about 140 ha (46%) with severe limitations of rooting depth and nutrient availability.




Fig. 7.22 Land Suitability map of Cashew

7. 23 Land Suitability for Jackfruit (*Artocarpus heterophyllus*)

Jackfruit is one of the most important fruit crop grown in an area of 5368 ha in almost all the districts of the State. The crop requirements for growing jackfruit (Table 7.24) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing jackfruit was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.23.

Currently not suitable (Class N1) lands for growing jackfruit occur in the entire cultivated area of about 140 ha (46%) with severe limitations of rooting depth and nutrient availability.




Fig. 7.23 Land Suitability map of Jackfruit

7.24 Land Suitability for Jamun (*Syzygium cumini*)

Jamun is an important fruit crop grown in almost all the districts of the State. The crop requirements for growing jamun (Table 25) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing jamun was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.24.

Currently not suitable (Class N1) lands for growing jamun occur in the entire cultivated area of about 140 ha (46%) with severe limitations of rooting depth and nutrient availability.




Fig. 7.24 Land Suitability map of Jamun

7.25 Land Suitability for Custard Apple (*Annona reticulata*)

Custard apple is one of the most important fruit crop grown in almost all the districts of the State. The crop requirements for growing custard apple (Table 7.26) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing custard apple was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.25.

Marginally suitable lands (Class S3) for growing custard apple occupy an area of about 33 ha (11%) and are distributed in the northeastern, eastern and southeastern part of the microwatershed. They have moderate limitations of rooting depth and texture. About 107 ha (35%) is currently not suitable (Class N1) for growing custard apple and are distributed in the major cultivated area of the microwatershed with severe limitations of rooting depth and nutrient availability.




Fig. 7.25 Land Suitability map of Custard Apple

7.26 Land Suitability for Tamarind (*Tamarindus indica*)

Tamarind is one of the most important spice crop grown in almost all the districts of the state. The crop requirements for growing tamarind (Table 7.27) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing tamarind was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Fig. 7.26.

Currently not suitable (Class N1) lands for growing tamarind occur in the entire cultivated area of about 140 ha (46%) with severe limitations of rooting depth and nutrient availability.




Fig. 7.26 Land Suitability map of Tamarind

7.27 Land Suitability for Mulberry (*Morus nigra*)

Mulberry is the important leaf crop grown for rearing of silkworms in about 1.6 lakh ha area in all the districts of the state. The crop requirements for growing mulberry (Table 7.28) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing mulberry was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.27.

Currently not suitable (Class N1) lands for growing mulberry occur in the entire cultivated area of about 140 ha (46%) with severe limitations of rooting depth and nutrient availability.




Fig 7.27 Land Suitability map of Mulberry

7.28 Land Suitability for Marigold (*Tagetes sps.*)

Marigold is one of the most important flower crop grown in an area of 9108 ha in almost all the districts of the State. The crop requirements (Table 7.29) for growing marigold were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing marigold was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.28.

Marginally suitable lands (Class S3) for growing marigold occupy an area of about 33 ha (11%) and are distributed in the northeastern, eastern and southeastern part of the microwatershed. They have moderate limitation of rooting depth and texture. About 107 ha (35%) is currently not suitable (Class N1) for growing marigold and are distributed in the major part of the cultivated area with severe limitations of rooting depth and nutrient availability.




Fig. 7.28 Land Suitability map of Marigold

7.29 Land Suitability for Chrysanthemum (*Dendranthema grandiflora*)

Chrysanthemum is one of the most important flower crop grown in an area of 4978 ha in almost all the districts of the State. The crop requirements (Table 7.30) for growing chrysanthemum were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing chrysanthemum was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.29.

Marginally suitable lands (Class S3) for growing chrysanthemum occupy an area of about 33 ha (11%) and are distributed in the northeastern, eastern and southeastern part of the microwatershed. They have moderate limitation of rooting depth. About 107 ha (35%) is currently not suitable (Class N1) for growing chrysanthemum and are distributed in the major part of the cultivated area with severe limitations of rooting depth and nutrient availability.




Fig. 7.29 Land Suitability map of Chrysanthemum

Table 7.1 Soil-Site Characteristics of Yadgir Rf-1 Microwatershed

Soil Map Units	Climate (P) (mm)	Growing period (Days)	Drain-age Class	Soil depth (cm)	Soil texture		Gravelliness		AWC (mm/m)	Slope (%)	Erosion	pH	EC (dSm ⁻¹)	ESP (%)	CEC [Cmol (p ⁺)kg ⁻¹]	BS (%)
					Sur-face	Sub-surface	Surface (%)	Sub-surface (%)								
KKRbB2g1	866	150	WD	<25	ls	sl	15-35	10-15	<50	1-3	moderate	5.85	0.027	1.17	2.6	60.90
HTKbB2	866	150	WD	25-50	ls	sl	<15	10-25	<50	1-3	moderate	6.81	0.062	0.38	3.0	100
YDRcB2	866	150	WD	100-150	sl	sl	<15	<15	51-100	1-3	moderate	9.47	0.371	4.86	12.70	165
YDRcB2g1	866	150	WD	100-150	sl	sl	15-35	<15	51-100	1-3	moderate	9.47	0.371	4.86	12.70	165

*Symbols and abbreviations are according to Field Guide for LRI under Sujala-III Project, Karnataka

Table 7.2 Land suitability criteria for Sorghum

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	26–30	30–34; 24–26	34–40; 20–24	>40; <20
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, c (red), c (black)	scl, cl	ls, sl	-
	pH	1:2.5	5.5-7.8	5.0-5.5 7.8-9.0	>9.0	-
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	10-15
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

Table 7.3 Land suitability criteria for Maize

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	30-34	35-38 26-30	38-40 26-20	
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	Mm				
	Rainfall in growing season	Mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc	c (red), c (black)	ls, sl	-
	pH	1:2.5	5.5-7.8	5.0-5.5 7.8-9.0	>9.0	-
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	Cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	-
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

Table 7.4 Land suitability criteria for Bajra

Land use requirement		Rating				
Soil –site characteristics	Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)	
Climatic regime	Mean temperature in growing season	°C	28-32	33-38 24-27	39-40 20-23	<20
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm	500-750	400-500	200-400	<200
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl, scl, cl,sc,c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.8	5.0-5.5 7.8-9.0	5.5-6.0 >9.0	
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	15-35	35-60	>60	
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	1-3	3-5	5-10	>10

Table 7.5 Land suitability criteria for Groundnut

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	24–33	22–24; 33–35	20–22; 35–40	<20; >40
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	Mm				
	Rainfall in growing season	Mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. Well drained	Poorly drained	Very Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl	sl,cl, sc	c (red), c (black), ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	Cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<35	35-60	>60	
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.6 Land suitability criteria for Sunflower

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	24–30	30–34; 20–24	34–38; 16–20	>38; <16
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	mod. Well drained	-	Poorly to very drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	cl, sc,c (red), c (black)	scl	ls, sl	-
	pH	1:2.5	6.5-7.8	7.8-8.4 5.5-6.5	8.4-9.0; 5.0-5.5	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.7 Land suitability criteria for Redgram

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	30-35(G) 20-25(AV) 15-18 (F&PS) 35-40(M)	25-30(G) 20-25 (AV) 12-15 (F&PS) 30-35(M)	20-25(G) 15-20(AV) 10-12 (F&PS) 25-30(M)	< 20 <15 <10 <25
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	Mm				
	Rainfall in growing season	Mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. Well drained	Poorly drained	Very Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, c (red)	c (black),sl, scl, cl	ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-9.0	5.0-5.5 >9.0	-
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	Cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-50	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<1.0	1.0-2.0	>2.0	
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.8 Land suitability criteria for Bengal gram

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	20–25	25–30; 15–20	30–35; 10–15	>35; <10
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. Well drained	Poorly drained	Very Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	c(black)	-	c (red), scl, cl, sc	ls, sl
	pH	1:2.5	6.0-7.8	5.0-6.0 7.8-9.0	>9.0	-
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	-
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.9 Land suitability criteria for Cotton

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	22-32	>32	<19	-
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well to moderately well	Poorly drained/Some what excessively drained	-	very poorly/excessively drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, c (red,black)	cl	scl	ls, sl
	pH	1:2.5	6.5-7.8	7.8-8.4	5.5-6.5 8.4->9.0	<5.5
	CEC	C mol (p+)Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	50-100	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	<3	3-5	-	>5

Table 7.10 Land suitability criteria for Chilli

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	25-32	33-35 20-25	35-38 <20	>38
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc	c (black), sl	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.11 Land suitability criteria for Tomato

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	25-28	29-32 20-24	15-19 33-36	<15 >36
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl, scl, cl, sc, c (red)	-	ls, c(black)	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.12 Land suitability criteria for Brinjal

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	Well drained	Moderately well drained	Poorly drained	V. Poorly drained
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class				
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl, scl, cl, sc c (red)	-	ls, c (black)	-
	pH	1:2.5	6.0-7.3	7.3-8.4 5.0-6.0	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	>60
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.13 Land suitability criteria for Onion

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	20-30	30-35	35-40	>40
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately /imperfectly	-	Poorly to V poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl,scl,cl,sc,c (red)	-	c (Black),ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<1.0	1.0-2.0	2.0-4.0	<4
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.14 Land suitability criteria for Bhendi

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	25-28	29-32 20-24	15-19 33-36	<15 >36
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Imperfectly drained	Poorly to very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl,sc, c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.15 Land suitability criteria for Drumstick

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, scl, cl, c (red)	sl, c (black)	ls	S
	pH	1:2.5	6.0-7.3	5.0-5.5 7.3-7.8	5.5-6.0 7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<35	35-60	60-80	>80
Soil toxicity	Salinity (EC saturation extract)	ds/m				
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-10	-	>10

Table 7.16 Land suitability criteria for Mango

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-32	24-27 33-35	36-40	20-24
	Min temp. before flowering	°C	10-15	15-22	>22	-
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration	Days				
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V. Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	-	ls, sl, c (black)	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>150	100-150	75-100	<75
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.17 Land suitability criteria for Guava

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-32	33-36 24-27	37-42 20-23	
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	sl	c (black), ls	-
	pH	1:2.5	6.0-7.8	5.0-6.0	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.18 Land suitability criteria for Sapota

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-32	33-36 24-27	37-42 20-23	>42 <18
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	-	Poorly to very drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	sl	ls, c (black)	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.19 Land suitability criteria for Pomegranate

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	30-34	35-38 25-29	39-40 15-24	
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl,cl, sc, c (red)	c (black),sl	ls	-
	pH	1:2.5	5.5-7.8	7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.20 Land suitability criteria for Musambi

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-30	31-35 24-27	36-40 20-23	>40 <20
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately drained	poorly	Very poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c	sl	ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.21 Land suitability criteria for Lime

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-30	31-35 24-27	36-40 20-23	>40 <20
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately drained	poorly	Very poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c	sl	ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.22 Land suitability criteria for Amla

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. well drained	Poorly drained	V. Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	c (black)	ls, sl	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15-35	35-60	60-80	-
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

Table 7.23 Land suitability criteria for Cashew

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	32 to 34	28 to 32; 34 to 38	24 to 28; 38 to 40	<20; >40
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	-	sl, ls	c (black)
	pH	1:2.5	5.5-6.5	5.0-5.5 6.5-7.3	7.3-7.8	>7.8
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-10	>10	-

Table 7.24 Land suitability criteria for Jackfruit

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. well	Poorly	V. Poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	-	sl, ls, c (black)	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	>60
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10-

Table 7.25 Land suitability criteria for Jamun

Land use requirement		Rating				
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well	Mod. well	Poorly	V.Poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c(red)	sl, c (black)	ls	-
	pH	1:2.5	6.0-7.8	5.0-6.0	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>150	100-150	50-100	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	>60
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

Table 7.26 Land suitability criteria for Custard apple

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	Scl, cl, sc, c (red), c (black)	-	Sl, ls	-
	pH	1:2.5	6.0-7.3	5.5-6.0 7.3-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15-35	35-60	60-80	-
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	>5	-

Table 7.27 Land suitability criteria for Tamarind

Land use requirement		Rating				
Soil –site characteristics	Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)	
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod.well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl,sc, c (red)	sl, c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>150	100-150	75-100	<75
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

Table 7.28 Land suitability criteria for Mulberry

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	24–28	22–24; 28–32	32–38; 22–18	>38; <18
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V. Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, cl, scl	c (red)	c (black), sl, ls	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.8-8.4	7.3-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	0-35	35-60	60-80	>80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

Table 7.29 Land suitability criteria for Marigold

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	18-23	17-15 24-35	35-40 10-14	>40 <10
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl,scl, cl, sc, c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%				
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.30 Land suitability criteria for Chrysanthemum

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	18-23	17-15 24-35	35-40 10-14	>40 <10
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl,scl, cl, sc, c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%				
Erosion hazard	Slope	%	<3	3-5	5-10	>10

7.30 Land Management Units (LMUs)

The 4 soil map units identified in Yadgir Rf-1 microwatershed have been grouped into 2 Land Management Units (LMUs) for the purpose of preparing a Proposed Crop Plan. Land Management Units are grouped based on the similarities in respect of the type of soil, the depth of the soil, the surface soil texture, gravel content, AWC, slope, erosion etc. and a Land Management Units map (Fig. 7.30) has been generated. These Land Management Units are expected to behave similarly for a given level of management.




Fig. 7.30 Land Management Units Map Yadgir Rf-1 Microwatershed

The map units that have been grouped into 2 Land Management Units along with brief description of soil and site characteristics are given below.

LMU	Soil map units	Soil and site characteristics
1	42.YDRcB2 154.YDRcB2g1	Deep, sodic soils (100-150cm), 1-3 % slopes, non gravelly to gravelly (<15 - 35 %), moderate erosion.
2	153.KKRbB2g1 156.HTKbB2	Very shallow to shallow sandy loam soils (<25 to 50 cm), 1-3 % slopes, non gravelly to gravelly (<15 - 35 %), moderate erosion.

7.31 Proposed Crop Plan for Yadgir Rf-1 Microwatershed

After assessing the land suitability for the 29 crops, the Proposed Crop Plan has been prepared for the 2 identified LMUs by considering only the highly (Class S1) and moderately (Class S2) suitable lands for each of the 29 crops. The resultant proposed crop plan is presented below in Table 7.31.

There are no highly and moderately suitable lands available for growing any of the major crops grown because of sodicity and shallow rooting depth.

Table 7.31 Proposed Crop Plan for Yadgir Rf-1 Microwatershed

LMU	Soil Map Units	Survey Number	Soil and site characteristics	Field Crops/ Commercial crops	Horticulture Crops (Rainfed/Irrigated)	Suitable Interventions
1	42.YDRcB2 154.YDRcB2g1	Allura.B: 383,385(1), 385(2),386	Deep, sodic soils (100-150cm), 1-3 % slopes, non gravelly to gravelly (<15 - 35 %), moderate erosion.	-	Agri-Silvi-Pasture Ber, Aonla, <i>Acacia</i> sp. Dhaincha, Rhodes grass, Para grass ,Bermuda grass	Application of gypsum, iron pyrites and elemental sulphur. Addition of farm yard manures, green manures and providing subsurface drainage
2	153.KKRbB2g 1 156.HTKbB2	Allura.B: 380,380(1), 380(2),381,382,384 Bheemanahalli: 159, 168,169	Very shallow to shallow sandy loam soils (<25 to 50 cm), 1-3 % slopes, non gravelly to gravelly (<15 - 35 %), moderate erosion.	-	Agri-Silvi-Pasture: Hybrid Napier, <i>Styloxanthes hamata</i> , <i>Styloxanthes scabra</i>	Use of short duration varieties, sowing across the slope, drip irrigation and mulching is recommended

SOIL HEALTH MANAGEMENT

8.1 Soil Health

Soil health is basic to plant health and plant health is basic to human and bovine health. Soil is fundamental to crop production. Without soil, no food could be produced nor would livestock be fed on a large scale. Because it is finite and fragile, soil is a precious resource that requires special care from its users.

Soil health or the capacity of the soil to function is critical to human survival. Soil health has been defined as: “the capacity of the soil to function as a living system without adverse effect on the ecosystem”. Healthy soils maintain a diverse community of soil organisms that help to form beneficial symbiotic associations with plant roots, recycle essential plant nutrients, improve soil structure with positive repercussions for soil, water and nutrient holding capacity and ultimately improve crop production and also contribute to mitigating climate change by maintaining or increasing its carbon content.

Functional interactions of soil biota with organic and inorganic components, air and water determine a soil’s potential to store and release nutrients and water to plants and to promote and sustain plant growth. Thus, maintaining soil health is vital to crop production and conserve soil resource base for sustaining agriculture.

The most important characteristics of a healthy soil are

- Good soil tilth
- Sufficient soil depth
- Good water storage and good drainage
- Adequate supply, but not excess of nutrients
- Large population of beneficial organisms
- Small proportion of plant pathogens and insect pests
- Low weed pressure
- Free of chemicals and toxins that may harm the crop
- Resistance to degradation
- Resilience when unfavorable conditions occur

Characteristics of Yadgir Rf-1 Microwatershed

- ❖ The soil phases identified in the microwatershed belonged to different soil series, KKR series occupies maximum area of 60 ha (20%) followed by YDR 47 ha (16%) and HTK 33 ha (11%)
- ❖ As per land capability classification an area of 141 ha in the microwatershed falls under arable land category (Class III & IV). The major limitations identified in the arable lands were soil and erosion.
- ❖ On the basis of soil reaction an area of 61 ha (20%) is moderately acid (pH 5.5 -6.5), about 20 ha (7%) is strongly acid (pH 5.0 -5.5) and about 60 ha (20%) is neutral (pH 6.5 -7.3) in the microwatershed.

❖ Soil Health Management

The following actions are required to improve the current land husbandry practices that provide a sound basis for the successful adoption of sustainable crop production system.

Acid soils

An area of 81 ha is under acid soils

1. Growing of crops suitable for particular soil pH.
2. Ameliorating the soils through the application of amendments (liming materials).

Liming materials:

1. CaCO_3 (Calcium Carbonate).
2. Dolomite [$\text{Ca Mg}(\text{CO}_3)_2$]
3. Quick lime (Cao)
4. Slaked lime [$\text{Ca}(\text{OH})_2$]

For normal pH and pH 4.8 (35 t/ha) and pH 6.0-7.0 (4 t/ha) lime is required.

Neutral soils

About 60 ha is under neutral soils.

1. Regular addition of organic manure, green manuring, green leaf manuring, crop residue incorporation and mulching needs to be taken up to improve the soil organic matter status.
2. Application of Biofertilizers, (Azospirillum, Azotobacter, Rhizobium).
3. Application of 100 per cent RDF.
4. Need based micronutrient applications.

Besides the above recommendations, the best transfer of technology options are also to be adopted.

Soil Degradation

Soil erosion is one of the major factors affecting the soil health in the microwatershed. Out of total 306 ha area in the microwatershed, entire cultivated area of about 141 ha (46%) is suffering from moderate erosion. These areas need immediate soil and water conservation and other land development and land husbandry practices for restoring soil health.

Dissemination of Information and Communication of Benefits

Any large scale implementation of soil health management requires that supporting information is made available widely, particularly through channels familiar to farmers and extension workers. Given the very high priority attached to soil-health especially by the Central Government on issuing Soil-Health Cards to all the farmers, media outlets like Regional, State and National Newspapers, Radio and Dooradarshan programs in local languages but also modern information and communication

technologies such as Cellular phones and the Internet, which can be much more effective in reaching the younger farmers.

Inputs for Net Planning (Saturation Plan) and Interventions needed

Net planning (Saturation Plan) in IWMP is focusing on preparation of

1. Soil and Water Conservation Plan for each plot or farm.
2. Productivity enhancement measures/ interventions for existing crops/livestock/other farm enterprises.
3. Diversification of farming mainly with perennial horticultural crops and livestock.
4. Improving livelihood opportunities and income generating activities.

In this connection, how various outputs of Sujala-III are of use in addressing these objectives of Net Planning (Saturation Plan) are briefly presented below.


- ❖ **Soil Depth:** The depth of a soil decides the amount of moisture and nutrients it can hold, what crops can be taken up or not, depending on the rooting depth and the length of growing period available for raising any crop. Deeper the soil, better for a wide variety of crops. If sufficient depth is not available for growing deep rooted crops, either choose medium or short duration crops or deeper planting pits need to be opened and additional good quality soil brought from outside has to be filled into the planting pits.
- ❖ **Surface Soil Texture:** Lighter soil texture in the top soil means, better rain water infiltration, less run-off and soil moisture conservation, less capillary rise and less evaporation losses. Lighter surface textured soils are amenable to good soil tilth and are highly suitable for crops like groundnut, root vegetables (carrot, raddish, potato etc) but not ideal for crops that need stagnant water like lowland paddy. Heavy textured soils are poor in water infiltration and percolation. They are prone for sheet erosion; such soils can be improved by sand mulching. The technology that is developed by the AICRP-Dryland Agriculture, Vijayapura, Karnataka can be adopted.
- ❖ **Gravelliness:** More gravel content is favorable for run-off harvesting but poor in soil moisture storage and nutrient availability. It is a significant parameter that decides the kind of crop to be raised.
- ❖ **Land Capability Classification:** The land capability map shows the areas suitable and not suitable for agriculture and the major constraints in each of the plot/survey number. Hence, one can decide what kind of enterprise is possible in each of these units. In general, erosion and soil are the major constraints in Yadgir Rf-1 microwatershed.
- ❖ **Organic Carbon:** The OC content (an index of available Nitrogen) is medium (0.5-0.75%) is 21 ha (7%) and low (<0.5%) IN 120 ha (39%) area. The areas that are medium and low in OC needs to be further improved by applying farmyard manure and rotating crops with cereals and legumes or mixed cropping.

- ❖ **Promoting Green Manuring:** Growing of green manuring crops costs Rs. 1250/ha (green manuring seeds) and about Rs. 2000/ha towards cultivation that totals to Rs. 3250/- per ha. On the other hand, application of organic manure @ 10 tons/ha costs Rs. 5000/ha. The practice needs to be continued for 2-3 years or more. Nitrogen fertilizer needs to be supplemented by 25% in addition to the recommended level in 141 ha area where OC is low and medium (<0.5-0.75%). For example, for rainfed maize, recommended level is 50 kg N per ha and an additional 12 kg /ha needs to be applied for all the crops grown in these plots.
- ❖ **Available Phosphorus:** Available Phosphorus is medium (23-57 kg/ha) covering an area of <1 ha (<1%) and high (>57 kg/ha) in an area of 141 ha (46%) in the microwatershed. For all the crops 25% additional P needs to be applied where available P is medium.
- ❖ **Available Potassium:** Entire cultivated area in the microwatershed is medium (145-337 kg/ha) in available potassium content. All the plots, where available potassium is medium, additional 25% potassium may be applied.
- ❖ **Available Sulphur:** Available sulphur is a very critical nutrient for oilseed crops, it is medium (10-20ppm) in an area of 117 ha (38%) and low (<10 ppm) in an area of 24 ha (8%). Medium and low areas need to be applied with magnesium sulphate or gypsum or Factamphos (p) fertilizer (13% sulphur) for 2-3 years for the deficiency to be corrected.
- ❖ **Available Boron:** An area of 130 ha (43%) is low (<0.5ppm) and about 11 ha (3%) is medium (0.5-1.0ppm) in available boron. For these low and medium areas application of sodium borate @ 10 kg/ha as soil application or 0.2 % borax as foliar spray is recommended.
- ❖ **Available Iron:** Available iron content is sufficient (>4.5 ppm) in the entire cultivated area of the microwatershed.
- ❖ **Available Manganese:** Entire cultivated area in the microwatershed is sufficient in available manganese content.
- ❖ **Available Copper:** Entire cultivated area in the microwatershed is sufficient in available copper content.
- ❖ **Available Zinc:** Available zinc content is deficient (<0.6 ppm) in the entire cultivated area of the microwatershed. Application of zinc sulphate 25 kg/ha is recommended for the deficient areas.
- ❖ **Land Suitability for various crops:** Areas that are highly, moderately and marginally suitable for growing various crops are indicated. Along with the suitability, various constraints that are limiting the productivity are also indicated. For example, in case of cotton, gravel content, rooting depth and salinity/alkalinity are the major constraints in various plots. With suitable management interventions, the productivity can be enhanced. In order to increase the water holding capacity of light textured soils, growing of green manure crops and application of organic manure is recommended.

SOIL AND WATER CONSERVATION TREATMENT PLAN

For preparing soil and water conservation treatment plan for Yadgir Rf-1 microwatershed, the land resource inventory database generated under Sujala-III project has been transformed as information through series of interpretative (thematic) maps using soil phase map as a base. The various thematic maps (1:7920 scale) generated were

- Soil depth
- Surface soil texture
- Available water capacity
- Soil slope
- Soil gravelliness
- Land capability
- Present land use and land cover
- Crop suitability
- Rainfall
- Hydrology
- Water Resources
- Socio-economic data
- Contour plan with existing features- network of waterways, pothera boundaries, cut up/ minor terraces etc.
- Cadastral map (1:7920 scale)
- Satellite imagery (1:7920 scale)



Apart from these, Hand Level/ Hydro Marker/ Dumpy Level/ Total Station and Kathedars' List to be collected.

Steps for Survey and Preparation of Treatment Plan

The boundaries of Land User Groups' and Survey No. boundaries are traced in the field.

- Naming of user groups and farmers
- Identification of arable and non arable lands
- Identification of drainage lines and gullies
- Identification of non treatable areas
- Identification of priority areas in the arable lands
- Treatment plan for arable lands
- Location of water harvesting and recharge structures

9.1 Treatment Plan

The treatment plan recommended for arable lands is briefly described below

9.1.1 Arable Land Treatment


A. BUNDING

Steps for Survey and Preparation of Treatment Plan	
<ul style="list-style-type: none"> • Cadastral map (1:7920 scale) is enlarged to a scale of 1:2500 scale • Existing network of waterways, pothissa boundaries, grass belts, natural drainage lines/ watercourse, cut ups/ terraces are marked on the cadastral map to the scale • Drainage lines are demarcated into 	
Small gullies	(up to 5 ha catchment)
Medium gullies	(5-15 ha catchment)
Ravines	(15-25 ha catchment) and
Halla/Nala	(more than 25ha catchment)

USER GROUP-1

CLASSIFICATION OF GULLIES

ಕೊರಕಲಿನ ವರ್ಗೀಕರಣ



Measurement of Land Slope

Land slope is estimated or determined by the study and interpretation of contours or by measurement in the field using simple instruments like Hand Level or Hydromarker.



Vertical and Horizontal intervals between bunds as recommended by the Watershed Development Department.

Slope percentage	Vertical interval (m)	Corresponding Horizontal Distance (m)
2 - 3%	0.6	24
3 - 4%	0.9	21
4 - 5%	0.9	21
5 - 6%	1.2	21
6 - 7%	1.2	21

Note: (i) The above intervals are maximum.

(ii) Considering the slope class and erosion status (A1... A=0-1 % slope, 1= slight erosion) the intervals have to be decided.

Bund length recording: Considering the contour plan and the existing grass belts/partitions, the bunds are aligned and lengths are measured.

Section of the Bund

Bund section is decided considering the soil texture class and gravelliness class (bg₀... b=loamy sand, g₀ = <15% gravel). The recommended Sections for different soils are given below.

Recommended Bund Section



Top width (m)	Base width (m)	Height (m)	Side slope (Z:1;H:V)	Cross section (sq m)	Soil Texture	Remarks
0.3	0.9	0.3	01:01	0.18	Sandy loam	Vegetative bund
0.3	1.2	0.3	1.5:1	0.225	Sandy clay	
0.3	1.2	0.5	0.9:1	0.375	Red gravelly soils	
0.3	1.2	0.6	0.75:1	0.45		
0.3	1.5	0.6	01:01	0.54	Red sandy loam	
0.3	2.1	0.6	1.5:1	0.72	Very shallow black soils	
0.45	2	0.75	01:01	0.92		
0.45	2.4	0.75	1.3:1	1.07	Shallow black soils	
0.6	3.1	0.7	1.78:1	1.29	Medium black soils	
0.5	3	0.85	1.47:1	1.49		

Formation of Trench cum Bund

Dimensions of the Borrow Pits/Trenches to be excavated (machinery are decided considering the Bund Section).

Details of Borrow Pit dimensions are given below:


TRENCH CUM BUND

0.45 Sq.m section

IDEAL FOR HORTICULTURE CROPS

'A' FRAME FOR INTERBUND MANAGEMENT



1. ಸಮವಾತಕ ಉಳುವೆ
2. ಸಮವಾತಕ ಬಿತ್ತನೆ/ನಾಟಿ

Size of Borrow Pits/ Trench recommended for Trench cum Bund (by machinery)

Bund section	Bund length	Earth quantity	Pit				Berm (pit to pit)	Soil depth class
			L(m)	W(m)	D(m)	Quantity (m ³)		
m ²	M	m ³					m	
0.375	6	2.25	5.85	0.85	0.45	2.24	0.15	Shallow
0.45	6	2.7	5.4	1.2	0.43	2.79	0.6	Shallow
0.45	6	2.7	5	0.85	0.65	2.76	1	Moderately Shallow
0.54	5.6	3.02	5.5	0.85	0.7	3.27	0.1	Moderately shallow
0.54	5.5	2.97	5	1.2	0.5	3	0.5	Shallow
0.72	6.2	4.46	6	1.2	0.7	5.04	0.2	Moderately shallow
0.72	5.2	3.74	5.1	0.85	0.9	3.9	0.1	Moderately deep

B. Water Ways

1. Existing waterways are marked on the cadastral map (1:7920 scale) and their dimensions are recorded.
2. Considering the contour plan of the MWS, additional waterways/ modernization of the existing ones can be thought of.
3. The design details are given in the Manual.

C. Farm Ponds

Waterways and the catchment area will give an indication on the size of the Farm Pond. Location of the pond can be decided based on the contour plan/ field condition and farmers' need/desire.

D. Diversion Channel

Existing EPT/ CPT are marked on the cadastral map. Looking to the need, these can be modernized or fresh diversion channel can be proposed and runoff from this can be stored in Gokatte/ Recharge ponds.

9.1.2 Non-Arable Land Treatment

Depending on the gravelliness and crops preferred by the farmers, the concerned authorities can decide appropriate treatment plan. The recommended treatments may be Contour Trench, Staggered Trench, Crescent Bund, Boulder Bund or Pebble Bund.

9.1.3 Treatment of Natural Water Course/ Drainage Lines

- a) The cadastral map has to be updated as regards the network of drainage lines (gullies/ *nalas/ hallas*) and existing structures are marked to the scale and storage capacity of the existing water bodies are documented.
- b) The drainage line will be demarcated into Upper Reach, Middle Reach and Lower Reach.
- c) Considering the Catchment, *Nala* bed and bank conditions, suitable structures are decided.
- d) Number of storage structures (Check dam/ *Nala* bund/ Percolation tank) will be decided considering the commitments and available runoff from water budgeting and quality of water in the wells and site suitability.
- e) Detailed Leveling Survey using Dumpy Level / Total Station has to be carried out to arrive at the site-specific designs as shown in the Manual.
- f) The location of ground water recharge structures are decided by examining the lineaments and fracture zones from geological maps.
- g) Rainfall intensity data of the nearest Rain Gauge Station is considered for Hydrologic Designs.
- h) Silt load to the Storage/Recharge structures is reduced by providing vegetative, boulder and earthen checks in the natural water course. Location and design details are given in the Manual.

9.2 Recommended Soil and Water Conservation Measures

The appropriate conservation structures best suited for each of the land parcel/ survey number (Appendix-I) are selected based on the slope per cent, severity of erosion, amount of rainfall, land use and soil type. The different kinds of conservation structures recommended are:

1. Graded / Strengthening of Bunds
2. Trench cum Bunds (TCB)
3. Trench cum Bunds / Strengthening
4. Crescent Bunds

A map (Fig. 9.1) showing soil and water conservation plan with different kinds of structures recommended has been prepared which shows the spatial distribution and extent of area. Entire cultivated area of about 141 ha (46%) needs Graded bunding.

The conservation plan prepared may be presented to all the stakeholders including farmers and after considering their suggestions, the conservation plan for the microwatershed may be finalised in a participatory approach.




Fig. 9.1 Soil and Water Conservation Plan map of Yadgir Rf-1 Microwatershed

9.3 Greening of Microwatershed

As part of the greening programme in the watersheds, it is envisaged to plant a variety of horticultural and other tree plants that are edible, economical and produce lot of biomass which helps to restore the ecological balance in the watersheds. The lands that are suitable for greening programme are non-arable lands (land capability classes V, VI VII and VIII) and also the lands that are not suitable or marginally suitable and field bunds for growing annual and perennial crops. The method of planting these trees is given below.

It is recommended to open pits during the 1st week of March along the contour and heap the dug out soil on the lower side of the slope in order to harness the flowing water and facilitate weathering of soil in the pit. Exposure of soil in the pit also prevents spread of pests and diseases due to scorching sun rays. The pits should be filled with mixture of soil and organic manure during the second week of April and keep ready with sufficiently tall seedlings produced either in poly bags or in root trainer nurseries so that planting can be done during the 2nd or 3rd week of April depending on the rainfall.

The tree species suitable for the area considering rainfall, temperature and adaptability is listed below; waterlogged areas are recommended to be planted with species like Nerale (*Syzgium cumini*) and Bamboo. Dry areas are to be planted with species like Honge, Bevu, Seetaphal *etc.*

Dry Deciduous Species			Temp (°C)	Rainfall (mm)
1.	Bevu	<i>Azadiracta indica</i>	21–32	400 –1,200
2.	Tapasi	<i>Holoptelia integrifolia</i>	20-30	500 - 1000
3.	Seetaphal	<i>Anona Squamosa</i>	20-40	400 - 1000
4.	Honge	<i>Pongamia pinnata</i>	20 -50	500– 2,500
5.	Kamara	<i>Hardwickia binata</i>	25 -35	400 - 1000
6.	Bage	<i>Albezzia lebbek</i>	20 - 45	500 - 1000
7.	Ficus	<i>Ficus bengalensis</i>	20 - 50	500–2,500
8.	Sisso	<i>Dalbargia Sissoo</i>	20 - 50	500 -2000
9.	Ailanthus	<i>Ailanthus excelsa</i>	20 - 50	500 - 1000
10.	Hale	<i>Wrightia tinctoria</i>	25 - 45	500 - 1000
11.	Uded	<i>Steriospermum chelanoides</i>	25 - 45	500 -2000
12.	Dhupa	<i>Boswella Serrata</i>	20 - 40	500 - 2000
13.	Nelli	<i>Emblca Officinalis</i>	20 - 50	500 -1500
14.	Honne	<i>Pterocarpus marsupium</i>	20 - 40	500 - 2000
Moist Deciduous Species			Temp (°C)	Rainfall (mm)
15.	Teak	<i>Tectona grandis</i>	20 - 50	500-5000
16.	Nandi	<i>Legarstroemia lanceolata</i>	20 - 40	500 - 4000
17.	Honne	<i>Pterocarpus marsupium</i>	20 - 40	500 - 3000
18.	Mathi	<i>Terminalia alata</i>	20 -50	500 - 2000
19.	Shivane	<i>Gmelina arborea</i>	20 -50	500 -2000
20.	Kindal	<i>T.Paniculata</i>	20 - 40	500 - 1500
21.	Beete	<i>Dalbargia latifolia</i>	20 - 40	500 - 1500
22.	Tare	<i>T. belerica</i>	20 - 40	500 - 2000
23.	Bamboo	<i>Bambusa arundinasia</i>	20 - 40	500 - 2500
24.	Bamboo	<i>Dendrocalamus strictus</i>	20 – 40	500 – 2500
25.	Muthuga	<i>Butea monosperma</i>	20 - 40	400 - 1500
26.	Hippe	<i>Madhuca latifolia</i>	20 - 40	500 - 2000
27.	Sandal	<i>Santalum album</i>	20 - 50	400 - 1000
28.	Nelli	<i>Emblca officinalis</i>	20 - 40	500 - 2000
29.	Nerale	<i>Sizyzium cumini</i>	20 - 40	500 - 2000
30.	Dhaman	<i>Grevia tilifolia</i>	20 - 40	500 - 2000
31.	Kaval	<i>Careya arborea</i>	20 - 40	500 - 2000
32.	Harada	<i>Terminalia chebula</i>	20 - 40	500 - 2000

References

1. FAO (1976) Framework for Land Evaluation, Food and Agriculture Organization, Rome.72 pp.
2. FAO (1983) Guidelines for Land Evaluation for Rainfed Agriculture, FAO, Rome, 237 pp.
3. IARI (1971) Soil Survey Manual, All India Soil and Land Use Survey Organization, IARI, New Delhi, 121 pp.
4. Katyal, J.C. and Rattan, R.K. (2003) Secondary and Micronutrients; Research Gap and Future Needs. Fert. News 48 (4); 9-20.
5. Naidu, L.G.K., Ramamurthy, V., Challa, O., Hegde, R. and Krishnan, P. (2006) Manual Soil Site Suitability Criteria for Major Crops, NBSS Publ. No. 129, NBSS & LUP, Nagpur, 118 pp.
6. Natarajan, A. and Dipak Sarkar (2010) Field Guide for Soil Survey, National Bureau of Soil Survey and Land Use Planning (ICAR), Nagpur, India.
7. Natarajan, A., Rajendra Hegde, Raj, J.N. and Shivananda Murthy, H.G. (2015) Implementation Manual for Sujala-III Project, Watershed Development Department, Bengaluru, Karnataka.
8. Sarma, V.A.K., Krishnan, P. and Budihal, S.L. (1987) Laboratory Manual, Tech. Bull. 23, NBSS &LUP, Nagpur.
9. Sehgal, J.L. (1990) Soil Resource Mapping of Different States of India; Why and How?, National Bureau of Soil Survey and Land Use Planning, Nagpur, 49 pp.
10. Shivaprasad, C.R., R.S. Reddy, J. Sehgal and M. Velayuthum (1998) Soils of Karnataka for Optimizing Land Use, NBSS Publ. No. 47b, NBSS & LUP, Nagpur, India.
11. Soil Survey Staff (2006) Keys to Soil Taxonomy, Tenth edition, U.S. Department of Agriculture/ NRCS, Washington DC, U.S.A.
12. Soil Survey Staff (2012) Soil Survey Manual, Handbook No. 18, USDA, Washington DC, USA.

Appendix I
Yadgir Rf-1 (4D5B2A1a) Microwatershed
Soil Phase Information

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Allura .B	380(1)	0.05	HTKbB2	LMU-2	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIIes	Graded bunding
Allura .B	380(2)	0.19	HTKbB2	LMU-2	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIIes	Graded bunding
Allura .B	385(1)	1.26	YDRcB2g1	LMU-1	Deep (100-150 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IVes	Graded bunding
Allura .B	385(2)	1.92	YDRcB2	LMU-1	Deep (100-150 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram (Gg)	Not Available	IVes	Graded bunding
Allura .B	378	239.21	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro+Greengram+Redgram (Rc+Gg+Rg)	Not Available	Ro	Ro
Allura .B	380	0.95	HTKbB2	LMU-2	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIIes	Graded bunding
Allura .B	381	0.88	HTKbB2	LMU-2	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIIes	Graded bunding
Allura .B	382	4.06	HTKbB2	LMU-2	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIIes	Graded bunding
Allura .B	383	5.76	YDRcB2	LMU-1	Deep (100-150 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Greengram (Rg+Gg)	Not Available	IVes	Graded bunding
Allura .B	384	1.63	HTKbB2	LMU-2	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIIes	Graded bunding
Allura .B	386	2.83	YDRcB2	LMU-1	Deep (100-150 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IVes	Graded bunding
Baggalamadu	1	1.06	HTKbB2	LMU-2	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIIes	Graded bunding
Bheemanahalli	159	37.85	HTKbB2	LMU-2	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Fallow land (Rg+Fl)	Not Available	IIIes	Graded bunding
Bheemanahalli	168	4.33	HTKbB2	LMU-2	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIIes	Graded bunding
Bheemanahalli	169	4.18	HTKbB2	LMU-2	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIIes	Graded bunding

Appendix II

Yadgir Rf-1 (4D5B2A1a) Microwatershed

Soil Fertility Information

Village	Survey Number	Soil Reaction	Salinity	Organic Carbon	Available Phosphorus	Available Potassium	Available Sulphur	Available Boron	Available Iron	Available Manganese	Available Copper	Available Zinc
Allura .B	380(1)	Neutral (pH 6.5 - 7.3)	Non saline (<2 dsm)	Low (< 0.5 %)	High (> 57 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Allura .B	380(2)	Neutral (pH 6.5 - 7.3)	Non saline (<2 dsm)	Low (< 0.5 %)	High (> 57 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Allura .B	385(1)	Neutral (pH 6.5 - 7.3)	Non saline (<2 dsm)	Low (< 0.5 %)	High (> 57 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Allura .B	385(2)	Slightly acid (pH 6.0 - 6.5)	Non saline (<2 dsm)	Low (< 0.5 %)	High (> 57 kg/ha)	Medium (145 - 337 kg/ha)	Medium (10 - 20 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Allura .B	378	Rock outcrops	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro
Allura .B	380	Slightly alkaline (pH 7.3 - 7.8)	Non saline (<2 dsm)	Low (< 0.5 %)	High (> 57 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Allura .B	381	Slightly alkaline (pH 7.3 - 7.8)	Non saline (<2 dsm)	Low (< 0.5 %)	High (> 57 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Allura .B	382	Neutral (pH 6.5 - 7.3)	Non saline (<2 dsm)	Low (< 0.5 %)	High (> 57 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Allura .B	383	Slightly acid (pH 6.0 - 6.5)	Non saline (<2 dsm)	Low (< 0.5 %)	High (> 57 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Allura .B	384	Neutral (pH 6.5 - 7.3)	Non saline (<2 dsm)	Low (< 0.5 %)	High (> 57 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Allura .B	386	Moderately acid (pH 5.5 - 6.0)	Non saline (<2 dsm)	Low (< 0.5 %)	High (> 57 kg/ha)	Medium (145 - 337 kg/ha)	Medium (10 - 20 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Baggala madu	1	Neutral (pH 6.5 - 7.3)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	High (> 57 kg/ha)	Medium (145 - 337 kg/ha)	Medium (10 - 20 ppm)	Medium (0.5 - 1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Bheema nahalli	159	Neutral (pH 6.5 - 7.3)	Non saline (<2 dsm)	Low (< 0.5 %)	High (> 57 kg/ha)	Medium (145 - 337 kg/ha)	Medium (10 - 20 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Bheema nahalli	168	Neutral (pH 6.5 - 7.3)	Non saline (<2 dsm)	Low (< 0.5 %)	High (> 57 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Bheema nahalli	169	Neutral (pH 6.5 - 7.3)	Non saline (<2 dsm)	Low (< 0.5 %)	High (> 57 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)

Appendix III
Yadgir Rf-1 (4D5B2A1a) Microwatershed
Soil Suitability Information

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry	
Allura .B	380(1)	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1r	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Allura .B	380(2)	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1r	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Allura .B	385(1)	N1n	S3nt	N1n	S3nt	N1n	N1t	N1n	N1n	N1t	N1n	S3n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n	N1n
Allura .B	385(2)	N1n	S3nt	N1n	S3nt	N1n	N1t	N1n	N1n	N1t	N1n	S3n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n	N1n
Allura .B	378	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro
Allura .B	380	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1r	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Allura .B	381	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1r	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Allura .B	382	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1r	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Allura .B	383	N1n	S3nt	N1n	S3nt	N1n	N1t	N1n	N1n	N1t	N1n	S3n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n	N1n
Allura .B	384	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1r	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Allura .B	386	N1n	S3nt	N1n	S3nt	N1n	N1t	N1n	N1n	N1t	N1n	S3n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n	N1n
Baggalamadu	1	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1r	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Bheemanahalli	159	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1r	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Bheemanahalli	168	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1r	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Bheemanahalli	169	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1r	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	

PART-B

SOCIO-ECONOMIC STATUS OF FARM HOUSEHOLDS

CONTENTS

1.	Findings of the socio-economic survey	1-3
2.	Introduction	5
3	Methodology	7-8
4	Salient features of the survey	9-26
5	Summary	27-31

LIST OF TABLES

1	Households sampled for socio economic survey	9
2	Population characteristics	9
3	Age wise classification of household members	9
4	Education level of household members	10
5	Occupation of household heads	10
6	Occupation of family members	10
7	Institutional participation of household members	11
8	Type of house owned by households	11
9	Durable assets owned by households	11
10	Average value of durable assets owned by households	12
11	Farm implements owned by households	12
12	Average value of farm implements	12
13	Livestock possession by households	13
14	Average labour availability	13
15	Adequacy of hired labour	13
16	Distribution of land (ha)	13
17	Average land value (Rs./ha)	14
18	Cropping pattern	14
19	Cropping intensity	14
20	Possession of bank account and saving	14
21	Borrowing status	14
22	Source of credit	15
23	Avg. credit borrowed	15
24	Purpose of credit borrowed from institutional sources	15
25	Purpose of credit borrowed from Non-institutional sources	15
26	Repayment status of household from institutional sources	16
27	Repayment status of household from Non-institutional sources	16
28	Opinion on institutional sources of credit	16
29	Opinion on Non-institutional sources of credit	16
30.a	Cost of cultivation of Red gram	17

30.b	Cost of cultivation of Green gram	18
30.c	Cost of cultivation of Maize	19
31	Adequacy of fodder	20
32	Annual gross income	20
33	Average annual expenditure	20
34	Horticultural species grown	20
35	Interest to cultivate horticultural crops	21
36	Forest species grown	21
37	Average additional investment capacity	21
38	Source of funds for additional investment	21
39	Marketing of the agricultural produce	22
40	Marketing channels used for sale of agricultural produce	22
41	Mode of transport of agricultural produce	22
42	Incidence of soil and water erosion problems	23
43	Interest shown towards soil testing	23
44	Soil and water conservation practices and structures	23
45	Status soil and water conservation structures	23
46	Agencies involved in the soil and water conservation structures	24
47	Usage pattern of fuel for domestic use	24
48	Source of drinking water	24
49	Source of light	24
50	Existence of sanitary toilet facility	24
51	Possession of public distribution system (PDS) card	25
52	Participation in NREGA programme	25
53	Adequacy of food items	25
54	Inadequacy of food items	25
55	Farming constraints experienced	26

FINDINGS OF THE SOCIO-ECONOMIC SURVEY

- ❖ *The survey was conducted in Yadgir Rf-1 is located at North latitude 16° 56' 33.469" and 16° 56' 0.445" and East longitude 77° 11' 20.418" and 77° 9' 57.462" covering an area of about 142.59 ha coming unde Baggalamadu, Alur. B and Bhimanahalli villages of Yadagiri taluk.*
- ❖ *Socio-economic analysis of Yadgir Rf-1 micro watersheds of Chikka Alur sub-watershed, Yadgiri taluk & District indicated that, out of the total sample of 35 farmers were sampled in Yadgir Rf-1 micro-watershed among households surveyed 10 (28.57%) were marginal, 13 (37.14%) were small, 5 (14.29 %) were semi medium and 2 (5.71 %) were medium farmers. 5 landless farmers were also interviewed for the survey.*
- ❖ *The population characteristics of households indicated that, there were 97 (58.79%) men and 68 (41.21 %) were women. The average population of landless was 5.2, marginal farmers were 3.4, small farmers were 5.2, semi medium farmers were 5.2 and medium farmers were 6.*
- ❖ *Majority of the respondents (43.03%) were in the age group of 16-35 years.*
- ❖ *Education level of the sample households indicated that, there were 46.67 per cent illiterates, 49.08 per cent pre university education and 4.24 per cent attained graduation.*
- ❖ *About, 48.57 per cent of household heads practicing agriculture and 42.86 per cent of the household heads were engaged as agricultural labourers.*
- ❖ *Agriculture was the major occupation for 29.09 per cent of the household members.*
- ❖ *In the study area, 68.57 per cent of the households possess katcha house and 2.86 per cent possess pucca house.*
- ❖ *The durable assets owned by the households showed that, 60.00 per cent possess TV, 57.14 per cent possess mixer grinder, 97.14 per cent possess mobile phones and 14.29 per cent possess motor cycles.*
- ❖ *Farm implements owned by the households indicated that, 34.29 per cent of the households possess plough, 20.00 per cent possess bullock cart and 11.43 per cent possess sprayer.*
- ❖ *Regarding livestock possession by the households, 11.43 per cent possess local cow and 2.86 per cent possess buffalo.*
- ❖ *The average labour availability in the study area showed that, own labour men available in the micro watershed was 1.57, women available in the micro watershed was 1.30, hired labour (men) available was 7.23 and hired labour (women) available was 11.1.*
- ❖ *Out of the total land holding of the sample respondents 100.00 per cent (49.19 ha) of the area is under dry condition.*

- ❖ *The major crops grown by sample farmers are Red gram, Green gram, Maize and cropping intensity was recorded as 100.00 per cent.*
- ❖ *Out of the sample households 97.14 percent possessed bank account and 17.14 per cent of them have savings in the account.*
- ❖ *About 68.57 per cent of the respondents borrowed credit from various sources.*
- ❖ *Among the credit borrowed by households, 6.25 per cent have borrowed loan from commercial banks and 75.00 per cent from co-operative/Grameena bank.*
- ❖ *Majority of the respondents (100.00%) have borrowed loan for agriculture purpose.*
- ❖ *Regarding the opinion on institutional sources of credit, 92.86 per cent of the households opined that credit helped to perform timely agricultural operations, while, only 7.14 per cent respondents opined that loan amount was adequate to fulfil their requirement.*
- ❖ *The per hectare cost of cultivation for Red gram, Green gram and Maize was Rs.21228.66, 15877.36 and 20572.72 with benefit cost ratio of 1:1.12, 1: 1.90 and 1: 1.72 respectively.*
- ❖ *Further, 20.00 per cent of the households opined that dry fodder was adequate.*
- ❖ *The average annual gross income of the farmers was Rs. 75218.57 in micro-watershed, of which Rs. 37504.29 comes from agriculture.*
- ❖ *Sampled households have grown 10 horticulture trees and 20 forestry trees together in the fields and back yards.*
- ❖ *About 60.00 per cent of the households shown interest to cultivate horticultural crops.*
- ❖ *Households have an average investment capacity of Rs. 2200.00 for land development.*
- ❖ *Source of funds for additional investment is concerned, 54.29 per cent depends on bank loan for land development activities.*
- ❖ *Regarding marketing channels, 85.71 per cent of the households have sold agricultural produce to the local/village merchants.*
- ❖ *Further, 31.43 per cent of the households have used tractor for the transport of agriculture commodity.*
- ❖ *Majority of the farmers (54.29%) have experienced soil and water erosion problems in the watershed and 85.71 per cent of the households were interested towards soil testing.*
- ❖ *Fire was the major source of fuel for domestic use for 45.71 per cent of the households and 54.29 per cent households has LPG connection.*
- ❖ *Piped supply was the major source for drinking water for 100.00 per cent of the households.*
- ❖ *Electricity was the major source of light for 100.00 per cent of the households.*
- ❖ *In the study area, 57.14 per cent of the households possess toilet facility.*

- ❖ *Regarding possession of PDS card, 100.00 per cent of the households possessed BPL card.*
- ❖ *Households opined that, the requirement of cereals (100.00%), pulses (100.00%) and oilseeds (57.14%) are adequate for consumption.*
- ❖ *Farming constraints experienced by households in the micro watersheds were lower fertility status of the soil (85.71%) wild animal menace on farm field (82.86%), frequent incidence of pest and diseases (85.71%), inadequacy of irrigation water (85.71%), high cost of fertilizers and plant protection chemicals (80.00%), high rate of interest on credit (80.00%), low price for the agricultural commodities (82.86%), lack of marketing facilities in the area (74.29%), inadequate extension services (71.43%) and lack of transport for safe transport of the agricultural produce to the market (82.86%).*

INTRODUCTION

Soil and water are the two precious natural resources which are essential for crop production and existence of life on earth. Rainfed agriculture is under severe stress due to various constraints related to agriculture like uneven and erratic distribution of rainfall, indiscriminate use of fertilizers, chemicals and pesticides, adoption of improper land management practices, soil erosion, decline in soil fertility, decline in ground water resources leading to low crop productivity. The area under rainfed agriculture has to be managed effectively using the best available practices to enhance the production of food, fodder and fuel. This is possible if the land resources are characterized at each parcel of land through detailed land resource inventory using the best available techniques of remote sensing, GPS and GIS. The watershed development programs are aimed at the sustainable distribution of its resources and the process of creating and implementing plans, programs, and projects to sustain and enhance watershed functions that affect the plant, animal and human communities within a watershed boundary.

World Bank funded KWDP II, SUJALA III project was implemented in with Broad objective of demonstrating more effective watershed management through greater integration of programmes related to rain-fed agriculture, innovative and science based approaches and strengthen institutional capacities and If successful, it is expected that the systems and tools could be mainstreamed into the overall IWMP in the State of Karnataka and in time, throughout other IWMP operations in India. With this background the socio-economic survey has been carried out with following specific objectives:

1. To understand the demographic features of the households in the micro-watershed
2. To understand the extent of family labour available and additional employment opportunities available within the village.
3. To know the status of assets of households in the micro-watershed for suggesting possible improvements.
4. To study the cropping pattern, cropped area and productivity levels of different households in micro-watershed.
5. To determine the type and extent of livestock owned by different categories of HHs
6. Availability of fodder and level of livestock management.

Scope and importance of survey

Survey helps in identification of different socio-economic and resource use-patterns of farmers at the Micro watershed. Household survey provides demographic features, labour force, and levels of education; land ownership and asset position (including livestock and other household assets) of surveyed households; and cropping patterns, input intensities, and average crop yields from farmers' fields. It also discusses crop utilization and the degree of commercialization of production in the areas; farmers' access to and utilization of credit from formal and informal sources; and the level of adoption and use of soil, water, and pest management technologies.

METHODOLOGY

The description of the methods, components selected for the survey and procedures followed in conducting the baseline survey are furnished under the following heads.

1. Description of the study area

Yadgir District is one of the 30 districts of Karnataka state in southern India. This district was carved out from the erstwhile Gulbarga district as the 30th district of Karnataka on 10 April 2010. Yadgir town is the administrative headquarters of the district. The district comprises of 3 taluks namely, Shahapur, Yadgiri and Shorapur (There are 16 hoblies, 117 Gram Panchayats, 4 Municipalities, 8 Towns/ Urban agglomeration and 487 inhabited & 32 un-inhabited villages The district occupies an area of 5,160.88 km².

Yadgir district is the second smallest district in the state, area wise is very rich in cultural traditions. The vast stretch of fertile black soil of the district is known for bumper red gram and jowar crops. The district is a "Daal bowl" of the state. The district is also known for cluster of cement industries and a distinct stone popularly known as "Malakheda Stone". Two main rivers, Krishna and Bhima, and a few tributaries flow in this region. Krishna and Bhima Rivers drain the district. They constitute the two major river basins of the district. Kagna and Amarja are the two sub - basins of Bhima River, which occur within the geographical area of the district

According to the 2011 census Yadgir district has a population of 1, 172,985, roughly equal to the nation of Timor-Leste or the US state of Rhode Island. This gives it a ranking of 404th in India (out of a total of 640). The district has a population density of 224 inhabitants per square kilometre (580/sq mi). Its population growth rate over the decade 2001-2011 was 22.67%. Yadgir has a sex ratio of 984 females for every 1000 males, and a literacy rate of 52.36%.

2. Locale of the survey and description of the micro-watershed and

The study was conducted in Yadgir Rf-1 micro-watershed (Chikka Alur sub-watershed, Yadgiri taluk & District) is located at North latitude 16^o 56' 33.469" and 16^o 56' 0.445" and East longitude 77^o 11' 20.418" and 77^o 9' 57.462" covering an area of about 142.59 ha bounded by unde Baggalamadu, Alur. B and Bhimanahalli Villages.

3. Selection of the respondents for the study

The micro-watershed is marked with 320 square meters grids. One farmer from every alternate grid in the micro-watershed was selected for the study and interviewed for socio-economic data. Totally 35 households were interviewed for the survey.

4. The parameters considered for socio-economic survey of households

Two forms of data were collected from the micro-watershed which includes primary data from the farm households and secondary data about the villages under the micro-watershed jurisdiction.

The following parameters were considered for the primary data collection about the socio-economic data of the households, (1) Demographic information, (2) Farm and durable assets owned by households, (3) Livestock possession, (4) Labour availability, (5) Level of migration in the village, Land holding, (7) Cropping pattern, (8) Source of irrigation, (9) Borrowing status, (10) Cost of cultivation of major crops, (11) Economics of subsidiary activities, (12) Fodder availability, (13) Family annual income from different sources, (14) Horticulture and forestry species grown, (15) Additional investment capacity, (16) Marketing practices, (17) Status of soil and water conservation structure, (18) Access to basic needs and (19) Constraints and suggestion.

The following parameters were considered for the secondary data regarding the villages under the micro-watershed jurisdiction, (1) Number of villages in each micro-watershed jurisdiction, (2) Village wise number of households, (3) Geographical area of the villages, (4) Cultivable area including rainfed and irrigated, (5) Number and type of house in each village, (6) Human and livestock population, (7) Facilities in the village such as roads, transport facility for conveyance, drinking water supply, street light and (8) Community based organizations in the villages.

5. Development of interview schedule and data collection

Taking into the consideration the objectives of the survey, an interview schedule was prepared after thorough consultation with the experts in the field of social sciences. A comprehensive interview schedule covering all the major parameters for measuring the socio-economic situation was developed.

6. Tools used to analyze the data

The statistical components such as frequency and percentage were used to analyze the data.

Abbreviations used in the report

LL=Landless

MF=Marginal Farmers

SF=Small farmers

SMF=Semi medium farmers

MDF=Medium farmers

LF=Large Farmers

FINDINGS OF THE SURVEY

This chapter deals with systematic presentation of results of the survey. Keeping in view the objectives, the salient features of the survey are presented under the following headings.

Households sampled for socio-economic survey: The data on households sampled for socio economic survey in Yadgir Rf-1 Micro watershed is presented in Table 1 and it indicated that 35 farmers were sampled in Yadgir Rf-1 micro-watershed among households surveyed 10 (28.57%) were marginal, 13 (37.14%) were small, 5 (14.29 %) were semi medium and 2 (5.71 %) were medium farmers. 5 landless farmers were also interviewed for the survey.

Table 1. Households sampled for socio economic survey in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (5)		MF (10)		SF (13)		SMF (5)		MDF (2)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Farmers	5	14.3	10	28.6	13	37.1	5	14.3	2	5.71	35	100

Population characteristics: The population characteristics of households sampled for socio-economic survey in Yadgir Rf-1 Micro watershed is presented in Table 2. The data indicated that, there were 97 (58.79%) men and 68 (41.21%) were women. The average population of landless was 5.2, marginal farmers were 3.4, small farmers were 5.2, semi medium farmers were 5.2 and medium farmers were 6.

Table 2. Population characteristics in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (26)		MF (34)		SF (67)		SMF (26)		MDF (12)		All (165)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Men	13	50	21	62	39	58	16	61.5	8	66.7	97	58.8
2	Women	13	50	13	38	28	42	10	38.5	4	33.3	68	41.2
Total		26	100	34	100	67	100	26	100	12	100	165	100
Average		5.2		3.4		5.2		5.2		6.0		4.7	

Age wise classification of population: The age wise classification of household members in Yadgir Rf-1 Micro watershed is presented in Table 3. The indicated that, 29 (17.58%) of population were 0-15 years of age, 71 (43.03%) were 16-35 years of age, 54(32.73%) were 36-60 years of age and 11 (6.67 %) were above 61 years of age.

Table 3: Age wise classification of members of the household in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (26)		MF (34)		SF (67)		SMF (26)		MDF (12)		All (165)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	0-15 years of age	3	11.5	7	20.6	13	19.4	3	11.54	3	25	29	17.58
2	16-35 years of age	14	53.9	10	29.4	29	43.3	15	57.69	3	25	71	43.03
3	36-60 years of age	7	26.9	16	47.1	22	32.8	6	23.08	3	25	54	32.73
4	> 61 years	2	7.69	1	2.94	3	4.48	2	7.69	3	25	11	6.67
Total		26	100	34	100	67	100	26	100	12	100	165	100

Education level of household members: Education level of household members in Yadgir Rf-1 Micro watershed is presented in Table 4. The results indicated that, there were 46.67 per cent of illiterates, 16.97 per cent of them had primary school education, 7.27 per cent middle school education, 16.36 per cent high school education, 4.24 per cent of them had PUC education, 4.24 per cent attained graduation and 1.82 them had other education.

Table 4. Education level of members of the household in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (26)		MF (34)		SF (67)		SMF (26)		MDF (12)		All (165)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Illiterate	10	38.5	19	55.9	29	43.3	14	53.9	5	41.67	77	46.7
2	Primary School	2	7.69	5	14.7	13	19.4	5	19.2	3	25	28	17
3	Middle School	4	15.4	1	2.94	6	8.96	1	3.85	0	0	12	7.27
4	High School	7	26.9	4	11.8	11	16.4	3	11.5	2	16.67	27	16.4
5	PUC	2	7.69	1	2.94	3	4.48	1	3.85	0	0	7	4.24
6	ITI	1	3.85	0	0	2	2.99	0	0	0	0	3	1.82
7	Degree	0	0	2	5.88	3	4.48	1	3.85	1	8.33	7	4.24
8	Masters	0	0	0	0	0	0	1	3.85	0	0	1	0.61
9	Others	0	0	2	5.88	0	0	0	0	1	8.33	3	1.82
Total		26	100	34	100	67	100	26	100	12	100	165	100

Occupation of head of households: The data regarding the occupation of the household heads in Yadgir Rf-1 Micro watershed is presented in Table 5. The results indicate that, 48.57 per cent of households heads were practicing agriculture, 42.86 per cent of the household heads were agricultural Labour and housewife (5.71%).

Table 5: Occupation of heads of households in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (5)		MF (10)		SF (13)		SMF (5)		MDF (2)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Agriculture	0	0	5	50	9	69.23	1	20	2	100	17	48.57
2	Agricultural Labour	3	60	5	50	4	30.77	3	60	0	0	15	42.86
3	General Labour	1	20	0	0	0	0	0	0	0	0	1	2.86
4	Housewife	1	20	0	0	0	0	1	20	0	0	2	5.71
Total		5	100	10	100	13	100	5	100	2	100	35	100

Table 6: Occupation of members of the household in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (26)		MF (34)		SF (67)		SMF (26)		MDF (12)		All (165)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Agriculture	0	0	11	32.4	26	38.81	4	15.38	7	58	48	29.1
2	Agricultural Labour	8	30.8	14	41.2	12	17.91	6	23.08	2	17	42	25.5
3	General Labour	10	38.5	0	0	4	5.97	3	11.54	0	0	17	10.3
4	Private Service	0	0	1	2.94	7	10.45	5	19.23	0	0	13	7.88
5	Housewife	3	11.5	0	0	0	0	1	3.85	0	0	4	2.42
6	Children	0	0	2	5.88	0	0	0	0	1	8.3	3	1.82
Total		26	100	34	100	67	100	26	100	12	100	165	100

Occupation of the members of the household: The data regarding the occupation of the household members in Yadgir Rf-1 Micro watershed is presented in Table 6. The results indicate that, agriculture was the major occupation for 29.09 per cent of the household members, 25.45 per cent were agricultural labour, 10.30 per cent were general labour, 23.03 per cent were working in pursuing education, 2.42 per cent were involved as housewife and 1.82 per cent were children.

Institutional Participation of household members: The data regarding the institutional participation of the household members in Yadgir Rf-1 Micro watershed is presented in Table 7. The results show that, out of the total family members in the households, 0.61 per cent of them were participating in gram panchayat, 2.42 per cent of them were participating in Self Help Group.

Table 7: Institutional Participation of household member in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (26)		MF (34)		SF (67)		SMF (26)		MDF (12)		All (165)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Gram Panchayat	0	0	0	0	1	1.49	0	0	0	0	1	0.61
2	Self Help Group	0	0	3	8.82	1	1.49	0	0	0	0	4	2.42
3	NGOs	0	0	0	0	0	0	0	0	1	8.33	1	0.61
4	No Participation	26	100	31	91.2	65	97	26	100	11	91.67	159	96.4
Total		26	100	34	100	67	100	26	100	12	100	165	100

Type of house owned: The data regarding the type of house owned by the households in Yadgir Rf-1 Micro watershed is presented in Table 8. The results indicate that, 28.57 percent possess thatched house, 68.57 per cent of the households possess katcha house, 2.86 per cent possess pucca house and 0.00 percent possess semi pucca house.

Table 8. Type of house owned by households in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (5)		MF (10)		SF (13)		SMF (5)		MDF (2)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Thatched	0	0	6	60	4	30.77	0	0	0	0	10	28.57
2	Katcha	5	100	3	30	9	69.23	5	100	2	100	24	68.57
3	Pucca/RCC	0	0	1	10	0	0	0	0	0	0	1	2.86
Total		5	100	10	100	13	100	5	100	2	100	35	100

Table 9. Durable assets owned by households in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (5)		MF (10)		SF (13)		SMF (5)		MDF (2)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Television	5	100	2	20	7	53.9	5	100	2	100	21	60
2	Mixer/Grinder	5	100	2	20	7	53.9	4	80	2	100	20	57.14
3	Bicycle	4	80	2	20	6	46.2	1	20	0	0	13	37.14
4	Motor Cycle	1	20	1	10	3	23.1	0	0	0	0	5	14.29
5	Mobile Phone	5	100	9	90	13	100	5	100	2	100	34	97.14
6	Blank	0	0	1	10	0	0	0	0	0	0	1	2.86

Durable assets owned by the households: The data regarding the Durable Assets owned by the households in Yadgir Rf-1 Micro watershed is presented in Table 9. The results shows that, 60.00 per cent possess TV, 57.14 per cent possess mixer grinder, 37.14 per cent possess Bicycle, 14.29 per cent possess motor cycle, 97.14 per cent possess mobile phones.

Average value of durable assets: The data regarding the average value of durable assets owned by the households in Yadgir Rf-1 Micro watershed is presented in Table 10. The result shows that, the average value of television was Rs.2904.00, mixer grinder was Rs.1500.00, bicycle was Rs.1000.00, motor cycle was Rs. 33000.00, mobile phone was Rs.1390.00.

Table 10. Average value of durable assets owned in Yadgir Rf-1 micro-watershed

Average Value (Rs.)							
Sl.No.	Particulars	LL (5)	MF (10)	SF (13)	SMF (5)	MDF (2)	All (35)
1	Television	2000	2000	2142	4200	5500	2904
2	Mixer/Grinder	1000	1000	1571	2000	2000	1500
3	Bicycle	1000	1000	1000	1000	0	1000
4	Motor Cycle	35000	30000	33333	0	0	33000
5	Mobile Phone	1183	1827	1054	1625	3000	1390

Farm implements owned: The data regarding the farm implements owned by the households in Yadgir Rf-1 Micro watershed is presented in Table 11. About 20.00 per cent of the households possess Bullock Cart, 34.29 per cent possess plough, 11.43 per cent possess Sprayer, 57.14 per cent possess Weeder.

Table 11. Farm implements owned in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (5)		MF (10)		SF (13)		SMF (5)		MDF (2)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Bullock Cart	0	0	1	10	6	46.15	0	0	0	0	7	20
2	Plough	0	0	3	30	8	61.54	0	0	1	50	12	34.29
3	Sprayer	0	0	1	10	3	23.08	0	0	0	0	4	11.43
4	Weeder	5	100	5	50	8	61.54	1	20	1	50	20	57.14
5	Chaff Cutter	0	0	0	0	6	46.15	0	0	0	0	6	17.14
6	Blank	0	0	5	50	5	38.46	4	80	1	50	15	42.86

Table 12. Average value of farm implements in Yadgir Rf-1 micro-watershed

Average Value (Rs.)							
Sl.No.	Particulars	LL (5)	MF (10)	SF (13)	SMF (5)	MDF (2)	All (35)
1	Bullock Cart	0	20000	20000	0	0	20000
2	Plough	0	1166	708	0	2500	854
3	Sprayer	0	4000	4000	0	0	4000
4	Weeder	23	46	35	25	100	35
5	Chaff Cutter	0	0	4000	0	0	4000

Average value of farm implements: The data regarding the average value of farm Implements owned by the households in Yadgir Rf-1 Micro watershed is presented in

Table 12. The results show that the average value of plough was Rs.854.00, bullock Cart was Rs.20000.00, seed/fertilizer drill was Rs.4000.00, sprayer and weeder was Rs.35.00.

Livestock possession by the households: The data regarding the Livestock possession by the households in Yadgir Rf-1 Micro watershed is presented in Table 13. The results indicate that, 34.29 per cent of the households possess bullocks, 11.43 per cent possess local cow, 2.86 per cent possess buffalo.

Table 13. Livestock possession by households in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (5)		MF (10)		SF (13)		SMF (5)		MDF (2)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Bullock	0	0	3	30	8	61.54	0	0	1	50	12	34.29
2	Local cow	0	0	2	20	2	15.38	0	0	0	0	4	11.43
3	Buffalo	0	0	0	0	1	7.69	0	0	0	0	1	2.86
4	blank	5	100	6	60	3	23.08	5	100	1	50	20	57.14

Average Labour availability: The data regarding the average labour availability in Yadgir Rf-1 Micro watershed is presented in Table 14. The indicated that, own labour men available in the micro watershed was 1.57, women available in the micro watershed was 1.30, hired labour (men) available was 7.23 and hired labour (women) available was 11.1.

Table 14. Average labour availability in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (5)	MF (10)	SF (13)	SMF (5)	MDF (2)	All (35)
		N	N	N	N	N	N
1	Hired labour Female	0	9.2	12.08	13.2	9	11.1
2	Own Labour Female	0	1.2	1.38	1	2	1.3
3	Own labour Male	0	1.2	1.69	1.6	2.5	1.57
4	Hired labour Male	0	5.3	7.92	9.8	6	7.23

Adequacy of hired labour: The data regarding the adequacy of hired labour in Yadgir Rf-1 Micro watershed is presented in Table 15. The results indicate that, 85.71 per cent of the household opined that hired labour was adequate.

Table 15. Adequacy of hired labour in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (5)		MF (10)		SF (13)		SMF (5)		MDF (2)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Adequate	0	0	10	100	13	100	5	100	2	100	30	85.7

Distribution of land (ha): The data regarding the distribution of land (ha) in Yadgir Rf-1 Micro watershed is presented in Table 16. The results indicate that, 49.19 ha (100.00%) of dry land.

Table 16. Distribution of land (ha) in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (5)		MF (10)		SF (13)		SMF (5)		MDF (2)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Dry	0	0	7.98	100	19.22	100	13.08	100	8.9	100	49.19	100
	Total	0	100	7.98	100	19.22	100	13.08	100	8.9	100	49.19	100

Average value of land (ha): The data regarding the average land value (Rs./ha) in Yadgir Rf-1 Micro watershed is presented in Table 17. The results show that the average value of dry land was Rs.317031.43.

Table 17. Average value of land (ha) in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (5)	MF (10)	SF (13)	SMF (5)	MDF (2)	All (35)
		N	N	N	N	N	N
1	Dry	0	600912.3	327600	206406.7	202090.9	317031.4

Cropping pattern: The data regarding the cropping pattern in Yadgir Rf-1 Micro watershed is presented in Table 18. The results indicate that, farmers have grown Red gram (32.53 ha), Maize (7.69 ha), Red gram (7.37 ha) and Green gram (1.62 ha).

Table 18. Cropping pattern in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (5)	MF (10)	SF (13)	SMF (5)	MDF (2)	All (35)
1	Kharif - Red gram	0	4.75	8.62	10.25	8.91	32.53
2	Rabi - Maize	0	3.24	1.62	2.83	0	7.69
3	Rabi - Red gram	0	0	7.37	0	0	7.37
4	Kharif - Green gram	0	0	1.62	0	0	1.62
Total		0	7.99	19.23	13.08	8.91	49.21

Cropping intensity: The data regarding the cropping intensity in Yadgir Rf-1 Micro watershed is presented in Table 19. The results indicate that, the cropping intensity was 100.00 per cent.

Table 19. Cropping intensity (%) in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (5)	MF (10)	SF (13)	SMF (5)	MDF (2)	All (35)
1	Cropping Intensity	0	100	100	100	100	100

Possession of bank account and savings: The data regarding the possession of bank account and saving in Yadgir Rf-1 micro-watershed is presented in Table 20. The results indicate that, 97.14 cent of the households posses bank account and 17.14 per cent of them have savings.

Table 20. Possession of Bank account and savings in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (5)		MF (10)		SF (13)		SMF (5)		MDF (2)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Account	5	100	10	100	13	100	4	80	2	100	34	97.14
2	Savings	0	0	1	10	4	30.77	1	20	0	0	6	17.14

Borrowing status: The data regarding the borrowing status in Yadgir Rf-1 micro-watershed is presented in Table 21. The results indicate that, 68.57 percent of the sample farmers have borrowed credit from different sources.

Table 21. Borrowing status in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (5)		MF (10)		SF (13)		SMF (5)		MDF (2)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Credit Availed	5	100	6	60	7	53.9	4	80	2	100	24	68.57

Source of credit: The data regarding the source of credit availed by households in Yadgir Rf-1 micro-watershed is presented in Table 22. The results shows that, 6.25 per cent have

borrowed loan from commercial banks and 6.25 per cent have borrowed loan from Cooperative bank, 75.00 per cent have borrowed loan from Grameena Bank, 6.25 per cent have borrowed loan from money lender, 25.00 per cent have borrowed loan from SHGs/CBOs.

Table 22. Source of credit borrowed by households in Yadgir Rf-1 micro-watershed

Sl. No.	Particulars	LL (5)		MF (4)		SF (6)		SMF (1)		MDF (0)		All (16)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Commercial Bank	0	0	1	25	0	0	0	0	0	0	1	6.25
2	Cooperative Bank	0	0	0	0	0	0	1	100	0	0	1	6.25
3	Grameena Bank	0	0	1	25	9	150	1	100	0	0	12	75
4	Money Lender	1	20	0	0	0	0	0	0	0	0	1	6.25
5	SHGs/CBOs	0	0	3	75	1	16.7	0	0	0	0	4	25

Avg. Credit amount: The data regarding the avg. Credit amount in Yadgir Rf-1 micro-watershed is presented in Table 23. The results show that, farmers have borrowed Avg. Credit of Rs.54687.50 from different sources.

Table 23. Avg. Credit amount in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (5)	MF (4)	SF (6)	SMF (1)	MDF (0)	All (16)
		N	N	N	N	N	N
1	Average Credit	6000	35000	83333.3	105000	0	54687.5

Purpose of credit borrowed (institutional Source): The data regarding the purpose of credit borrowed - Institutional Credit in Yadgir Rf-1 micro-watershed is presented in Table 24. The results indicate that, 100.00 per cent of the households have borrowed loan for agriculture.

Table 24. Purpose of credit borrowed (institutional Source) by households in Yadgir Rf-1 micro-watershed

SN	Particulars	LL (0)		MF (2)		SF (9)		SMF (2)		MDF (1)		All (14)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Agriculture production	0	0	2	100	9	100	2	100	1	100	14	100

Purpose of credit borrowed (Private Source): The data regarding the purpose of credit borrowed – Private Source in Yadgir Rf-1 micro-watershed is presented in Table 25. The results indicate that, 80.00 per cent of the households have borrowed loan for agriculture.

Table 25. Purpose of credit borrowed (Private Source) by households in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (1)		MF (3)		SF (1)		SMF (0)		MDF (0)		All (5)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Agriculture production	0	0	3	100	1	100	0	0	0	0	4	80
2	Social functions like marriage	1	100	0	0	0	0	0	0	0	0	1	20

Repayment status of household (institutional Source): The data regarding the repayment status of credit borrowed from institutional Source by households in Yadgir Rf-1 micro watershed is presented in Table 26. The results indicate that, 7.14 per cent of the households have partially paid, 78.57 per cent have unpaid and 14.29 percent have fully paid.

Table 26. Repayment status of household (institutional Source) in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (0)		MF (2)		SF (9)		SMF (2)		MDF (1)		All (14)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Partially paid	0	0	1	50	0	0	0	0	0	0	1	7.14
2	Un paid	0	0	0	0	8	88.9	2	100	1	100	11	78.57
3	Fully paid	0	0	1	50	1	11.1	0	0	0	0	2	14.29

Repayment status of household (Private Source): The data regarding the repayment status of credit borrowed from private sources by households in Yadgir Rf-1 micro watershed is presented in Table 27. The results indicate that, 100.00 per cent of the households have partially paid.

Table 27. Repayment status of household (Private Source) in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (1)		MF (3)		SF (1)		SMF (0)		MDF (0)		All (5)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Partially paid	1	100	3	100	1	100	0	0	0	0	5	100

Opinion regarding institutional sources of credit: The data regarding the opinion on institutional sources of credit in Yadgir Rf-1 micro watershed is presented in Table 28. The results indicate that, 92.86 per cent of the households opined that credit helped to perform timely agricultural operations, 7.14 per cent Loan amount was adequate to fulfil the requirement.

Table 28. Opinion regarding institutional sources of credit in Yadgir Rf-1 micro-watershed

Sl. No.	Particulars	MF (2)		SF (9)		SMF (2)		MDF (1)		All (14)	
		N	%	N	%	N	%	N	%	N	%
1	Helped to perform timely agricultural operations	1	50	9	100	2	100	1	100	13	92.9
2	Loan amount was adequate to fulfil the requirement	1	50	0	0	0	0	0	0	1	7.14

Opinion regarding Non- institutional sources of credit: The data regarding the opinion on non-institutional sources of credit in Yadgir Rf-1 micro watershed is presented in Table 29. The results indicate that, 92.86 per cent of the households opined that credit helped to perform timely agricultural operations, 7.14 per cent Loan amount was adequate to fulfil the requirement.

Table 29. Opinion regarding Non- institutional sources of credit in Yadgir Rf-1 micro-watershed

Sl. No.	Particulars	LL (1)		MF (3)		SF (1)		SMF (0)		MDF (0)		All (5)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Helped to perform timely agricultural operations	0	0	1	33	0	0	0	0	0	0	1	20
2	Loan amount was adequate to fulfil the requirement	1	100	2	67	1	100	0	0	0	0	4	80

Cost of Cultivation of Red gram: The data regarding the cost of cultivation (Rs/ha) of Red gram in Yadgir Rf-1 micro watershed is presented in Table 30.a. The results indicate that, the total cost of cultivation (Rs/ha) for Red gram was Rs. 21228.66. The gross income realized by the farmers was Rs. 23771.05. The net income from Red gram cultivation was Rs.2542.39, thus the benefit cost ratio was found to be 1:1.12.

Table 30(a). Cost of Cultivation of Red gram in Yadgir Rf-1 micro-watershed

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
I	Cost A1				
1	Hired Human Labour	Man days	27.53	5349.02	25.2
2	Bullock	Pairs/day	1.46	1254.78	5.91
3	Tractor	Hours	1.81	1660.76	7.82
4	Machinery	Hours	0	0	0
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	12.98	1145.26	5.39
6	Seed Inter Crop	Kgs.	0	0	0
7	FYM	Quintal	1.16	2904.07	13.68
8	Fertilizer + micronutrients	Quintal	1.93	2288.21	10.78
9	Pesticides (PPC)	Kgs /liters	0.89	781.2	3.68
10	Irrigation	Number	0	0	0
11	Repairs		0	0	0
12	Msc. Charges (Marketing costs etc)		0	0	0
13	Depreciation charges		0	114.39	0.54
14	Land revenue and Taxes		0	0	0
II	Cost B1				
16	Interest on working capital			854.25	4.02
17	Cost B1 = (Cost A1 + sum of 15 and 16)			16351.93	77.03
III	Cost B2				
18	Rental Value of Land			283.33	1.33
19	Cost B2 = (Cost B1 + Rental value)			16635.26	78.36
IV	Cost C1				
20	Family Human Labour		11.18	2663.52	12.55
21	Cost C1 = (Cost B2 + Family Labour)			19298.79	90.91
V	Cost C2				
22	Risk Premium			0	0
23	Cost C2 = (Cost C1 + Risk Premium)			19298.79	90.91
VI	Cost C3				
24	Managerial Cost			1929.88	9.09
25	Cost C3 = (Cost C2 + Managerial Cost)			21228.66	100
VII	Economics of the Crop				
a.	Main Product	a) Main Product (q)		4.74	23771.05
		b) Main Crop Sales Price (Rs.)			5012.5
b.	Gross Income (Rs.)			23771.05	
c.	Net Income (Rs.)			2542.39	
d.	Cost per Quintal (Rs./q.)			4476.4	
e.	Benefit Cost Ratio (BC Ratio)			1:1.12	

Cost of Cultivation of Green gram: The data regarding the cost of cultivation (Rs/ha) of Green gram in Yadgir Rf-1 micro watershed is presented in Table 30.b. The results indicate that, the total cost of cultivation (Rs/ha) for Green gram was Rs. 15877.36. The gross income realized by the farmers was Rs. 29640.00. The net income from Green gram cultivation was Rs.13762.64, thus the benefit cost ratio was found to be 1:1.90.

Table 30(b). Cost of Cultivation of Green gram in Yadgir Rf-1 micro-watershed

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
I	Cost A1				
1	Hired Human Labour	Man days	9.88	1667.25	10.5
2	Bullock	Pairs/day	1.85	1852.5	11.67
3	Tractor	Hours	0.62	617.5	3.89
4	Machinery	Hours	0	0	0
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	9.26	741	4.67
6	Seed Inter Crop	Kgs.	0	0	0
7	FYM	Quintal	1.24	1852.5	11.67
8	Fertilizer + micronutrients	Quintal	1.24	1482	9.33
9	Pesticides (PPC)	Kgs / liters	0.62	617.5	3.89
10	Irrigation	Number	0	0	0
11	Repairs		0	0	0
13	Depreciation charges		0	33.35	0.21
14	Land revenue and Taxes		0	0	0
II	Cost B1				
16	Interest on working capital			563.16	3.55
17	Cost B1 = (Cost A1 + sum of 15 and 16)			9426.76	59.37
III	Cost B2				
18	Rental Value of Land			283.33	1.78
19	Cost B2 = (Cost B1 + Rental value)			9710.09	61.16
IV	Cost C1				
20	Family Human Labour		20.38	4723.88	29.75
21	Cost C1 = (Cost B2 + Family Labour)			14433.96	90.91
V	Cost C2				
22	Risk Premium			0	0
23	Cost C2 = (Cost C1 + Risk Premium)			14433.96	90.91
VI	Cost C3				
24	Managerial Cost			1443.4	9.09
25	Cost C3 = (Cost C2 + Managerial Cost)			15877.36	100
VII	Economics of the Crop				
a.	Main Product	a) Main Product (q)		7.41	29640
		b) Main Crop Sales Price (Rs.)			4000
b.	Gross Income (Rs.)			29640	
c.	Net Income (Rs.)			13762.64	
d.	Cost per Quintal (Rs./q.)			2142.69	
e.	Benefit Cost Ratio (BC Ratio)			1:1.9	

Cost of Cultivation of Maize: The data regarding the cost of cultivation (Rs/ha) of Maize in Yadgir Rf-1 micro watershed is presented in Table 30.c. The results indicate, the total cost of cultivation (Rs/ha) for Maize was Rs.20572.72. The gross income realized by the farmers was Rs. 35037.66. The net income from Maize cultivation was Rs. 14464.93, thus the benefit cost ratio was found to be 1:1.72.

Table 30(c). Cost of Cultivation of Maize in Yadgir Rf-1 micro-watershed

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
I	Cost A1				
1	Hired Human Labour	Man days	22.71	4801.5	23.34
2	Bullock	Pairs/day	1.8	1079.74	5.25
3	Tractor	Hours	2.61	1958.36	9.52
4	Machinery	Hours	0	0	0
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	10.39	805.93	3.92
6	Seed Inter Crop	Kgs.	0	0	0
7	FYM	Quintal	0.84	2095.09	10.18
8	Fertilizer + micronutrients	Quintal	2.91	3493.29	16.98
9	Pesticides (PPC)	Kgs / liters	0	0	0
10	Irrigation	Number	0	0	0
13	Depreciation charges		0	75.48	0.37
14	Land revenue and Taxes		0	0	0
II	Cost B1				
16	Interest on working capital			767.32	3.73
17	Cost B1 = (Cost A1 + sum of 15 and 16)			15076.7	73.28
III	Cost B2				
18	Rental Value of Land			283.33	1.38
19	Cost B2 = (Cost B1 + Rental value)			15360.03	74.66
IV	Cost C1				
20	Family Human Labour		12.72	3342.44	16.25
21	Cost C1 = (Cost B2 + Family Labour)			18702.47	90.91
V	Cost C2				
22	Risk Premium			0	0
23	Cost C2 = (Cost C1 + Risk Premium)			18702.47	90.91
VI	Cost C3				
24	Managerial Cost			1870.25	9.09
25	Cost C3 = (Cost C2 + Managerial Cost)			20572.72	100
VII	Economics of the Crop				
a.	Main Product	a) Main Product (q)		14.41	34305.83
		b) Main Crop Sales Price (Rs.)			2380
	By Product	e) Main Product (q)		1.08	731.83
		f) Main Crop Sales Price (Rs.)			680
b.	Gross Income (Rs.)			35037.66	
c.	Net Income (Rs.)			14464.93	
d.	Cost per Quintal (Rs./q.)			1427.25	
e.	Benefit Cost Ratio (BC Ratio)			1:1.72	

Adequacy of fodder: The data regarding the adequacy of fodder in Yadgir Rf-1 Micro watershed is presented in Table 31. The results indicate that, 20.00 per cent of the households opined that dry fodder was adequate and 2.86 per cent of them opined dry fodder was inadequate.

Table 31. Adequacy of fodder in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (5)		MF (10)		SF (13)		SMF (5)		MDF (2)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Adequate-Dry Fodder	0	0	3	30	3	23.08	0	0	1	50	7	20
2	Inadequate-Dry Fodder	0	0	0	0	1	7.69	0	0	0	0	1	2.86

Average annual gross income: The data regarding the annual gross income in Yadgir Rf-1 Micro watershed is presented in Table 32. The results indicate that, the farmers have annual gross income of Rs. 75218.57 in micro-watershed, of which Rs. 37504.29 is from agriculture itself.

Table 32. Average annual gross income in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (5)	MF (10)	SF (13)	SMF (5)	MDF (2)	All (35)
		Rs.	Rs.	Rs.	Rs.	Rs.	Rs.
1	Service/salary	0	0	21538.5	32000	0	12571.4
2	Wage	76000	14400	20384.6	18200	0	25142.9
3	Agriculture	20000	20170	46111.5	47300	87500	37504.3
Income(Rs.)		96000	34570	88034.6	97500	87500	75218.6

Average annual Expenditure: The data regarding the average annual expenditure in Yadgir Rf-1 Micro watershed is presented in Table 33. The results indicate that, the farmers have annual gross expenditure of Rs. 262414.76 in micro-watershed, of which Rs. 21171.43 is from agriculture itself.

Table 33. Average annual Expenditure in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (5)	MF (10)	SF (13)	SMF (5)	MDF (2)	All (35)
		Rs.	Rs.	Rs.	Rs.	Rs.	Rs.
1	Service/salary	0	0	35000	12000	0	3685.71
2	Wage	45500	2888.89	3818.18	4500	0	7657.14
3	Agriculture	52000	12100	26307.7	29800	38500	21171.4
Total		97500	14988.9	65125.9	46300	38500	262415

Table 34. Horticulture species grown in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (5)		MF (10)		SF (13)		SMF (5)		MDF (2)		All (35)	
		F	B	F	B	F	B	F	B	F	B	F	B
1	Coconut	0	0	0	0	10	0	0	0	0	0	10	0

*F= Field B=Back Yard

Horticulture species grown: The data regarding horticulture species grown in Yadgir Rf-1 Micro watershed is presented in Table 34. The results indicate that, the total number

of horticultural trees grown (both field and backyard) by the sampled households were coconut (10).

Interest towards cultivation of horticulture crops: The data regarding Table (35) indicates that, 60.00 per cent of the households shown interest to cultivate horticultural crops.

Table 35. Interest towards cultivation of horticulture crops in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (5)		MF (10)		SF (13)		SMF (5)		MDF (2)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Interested towards cultivation of horticulture crops	0	0	9	90	10	77	0	0	2	100	21	60

Forest species grown: The data regarding forest species grown in Yadgir Rf-1 Micro watershed is presented in Table 36. The results indicate that, households have planted 18 neem trees, 2 tamarind trees together in both field and backyard.

Table 36. Forest species grown in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (5)		MF (10)		SF (13)		SMF (5)		MDF (2)		All (35)	
		F	B	F	B	F	B	F	B	F	B	F	B
1	Neem	0	0	2	0	15	1	0	0	0	0	17	1
2	Tamarind	0	0	1	0	1	0	0	0	0	0	2	0

*F= Field B=Back Yard

Average additional investment capacity: The data regarding average additional investment capacity in Yadgir Rf-1 Micro watershed is presented in Table 37. The results indicate that, households have an average investment capacity of Rs. 2200.00 for land development, Rs.3828.57 for adoption of improved livestock breeds, Rs.457.14 for adoption of improved crop production activities.

Table 37. Average additional investment capacity of households in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (5)	MF (10)	SF (13)	SMF (5)	MDF (2)	All (35)
		Rs.	Rs.	Rs.	Rs.	Rs.	Rs.
1	Land development	0	2100	1923.08	3600	6500	2200
2	Improved crop production	0	3100	3307.69	6200	14500	3828.57
3	Improved livestock management	0	600	615.38	0	1000	457.14

Table 38. Source of funds for additional investment in Yadgir Rf-1 micro-watershed

Sl.No	Item	Land development		Irrigation facility		Improved crop production		Improved livestock management	
		N	%	N	%	N	%	N	%
1	Own funds	19	54.29	0	0	19	54.29	7	20

Source of funds for additional investment: The data regarding source of funds for additional investment in Yadgir Rf-1 Micro watershed is presented in Table 38. The results indicate that, the sources of finance raised from own sources for land development was 54.29.

Marketing of agricultural produce: The data regarding marketing of the agricultural produce in Yadgir Rf-1 Micro watershed is presented in Table 39. The results indicated that, 58.33 percent of output of Green gram was sold in the market with average price of Rs. 4000.00; 90.29 percent of output of Maize was sold in the market with average price of Rs. 2380.00; 86.11 percent of output of Red gram was sold in the market with average price of Rs. 5012.50.

Table 39. Marketing of agricultural produce in Yadgir Rf-1 micro-watershed

Sl. No	Crops	Output obtained (q)	Output retained (q)	Output sold (q)	Output sold (%)	Avg. Price obtained (Rs/q)
1	Green gram	12	5	7	58	4000
2	Maize	103	10	93	90	2380
3	Red gram	180	25	155	86	5013

Marketing channels used for sale of agricultural produce: The data regarding marketing channels used for sale of agricultural produce in Yadgir Rf-1 Micro watershed is presented in Table 40. The results indicated that, 85.71 cent of the households have sold agricultural produce to the local/village merchants.

Table 40. Marketing channels used for sale of agricultural produce in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (5)		MF (10)		SF (13)		SMF (5)		MDF (2)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Local/village Merchant	0	0	10	100	13	100	5	100	2	100	30	85.71

Mode of transport of agricultural produce: The data regarding mode of transport of agricultural produce in Yadgir Rf-1 Micro watershed is presented in Table 41. The results indicated that, 31.43 cent of the households have used tractor, 54.29 per cent have used Cart for the transport of agriculture commodity.

Table 41. Mode of transport of agricultural produce in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (5)		MF (10)		SF (13)		SMF (5)		MDF (2)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Cart	0	0	6	60	8	61.5	4	80	1	50	19	54.29
2	Tractor	0	0	4	40	5	38.5	1	20	1	50	11	31.43

Incidence of soil and water erosion problems: The data regarding incidence of incidence of soil and water erosion problems in Yadgir Rf-1 Micro watershed is presented

in Table 42. The results indicate that, 54.29 per cent of the households have experienced soil and water erosion problems.

Table 42. Incidence of soil and water erosion problems in Yadgir Rf-1 micro-watershed

Sl. No.	Particulars	LL (5)		MF (10)		SF (13)		SMF (5)		MDF (2)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Soil and water erosion problems in the farm	0	0	6	60	7	53.9	4	80	2	100	19	54.29

Interest towards soil testing: The data regarding Interest shown towards soil testing in Yadgir Rf-1 Micro watershed is presented in Table 43. The results indicated that, 85.71 per cent of the households were interested towards soil testing.

Table 43. Interest regarding soil testing in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (5)		MF (10)		SF (13)		SMF (5)		MDF (2)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Interest in soil test	0	0	10	100	13	100	5	100	2	100	30	85.71

Soil and water conservation practices and structures adopted: The data regarding soil and water conservation practices and structures adopted in Yadgir Rf-1 Micro watershed is presented in Table 44. The results indicated that 100 per cent of farmers practicing summer ploughing as soil and water conservation practice.

Table 44. Soil and water conservation practices and structures adopted in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (5)		MF (10)		SF (13)		SMF (5)		MDF (2)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Field Bunding	0	0	1	10	2	15	0	0	0	0	3	8.57
2	Farm Pond	0	0	0	0	1	7.7	0	0	0	0	1	2.86

Status of soil and water conservation structures: The data regarding status soil and water conservation structures adopted in Yadgir Rf-1 Micro watershed is presented in Table 45. The results indicated that, the households have adopted field bunding as a soil and water conservation structures out of which 100.00 per cent was in good condition.

Table 45. Status of soil and water conservation structures in Yadgir Rf-1 micro-watershed

Sl. No	Item	Good		Slightly Damaged		Severely Damaged	
		N	%	N	%	N	%
1	Farm Pond	1	100	0	0	0	0

Agencies involved in the soil and water conservation structures: The data regarding Agencies involved in the soil and water conservation structures adopted in Yadgir Rf-1 Micro watershed is presented in Table 46. The results indicated that, 2.86 per cent of the households have adopted by their own, 8.57 per cent were done by Govt.

Table 46. Agencies involved in the soil and water conservation structures in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (5)		MF (10)		SF (13)		SMF (5)		MDF (2)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Own	0	0	1	10	0	0	0	0	0	0	1	2.86
2	Govt.	0	0	0	0	3	23.08	0	0	0	0	3	8.57

Usage pattern of fuel for domestic use: The data on usage pattern of fuel for domestic use in Yadgir Rf-1 Micro watershed is presented in Table 47. The results indicated that, firewood was the major source of fuel for domestic use for 45.71 per cent of the households followed by LPG (54.29%).

Table 47. Usage pattern of fuel for domestic use in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (5)		MF (10)		SF (13)		SMF (5)		MDF (2)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Fire Wood	4	80	5	50	5	38.5	0	0	2	100	16	45.71
2	LPG	1	20	5	50	8	61.5	5	100	0	0	19	54.29

Source of drinking water: The data on source of drinking water in Yadgir Rf-1 Micro watershed is presented in Table 48. The results indicated that, piped waters supply (100.00 %).

Table 48. Source of drinking water in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (5)		MF (10)		SF (13)		SMF (5)		MDF (2)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Piped supply	5	100	10	100	13	100	5	100	2	100	35	100

Source of light: The data on source of light in Yadgir Rf-1 Micro watershed is presented in Table 49. The results indicated that, electricity was the major source of light for 100.00 per cent of the households.

Table 49. Source of light in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (5)		MF (10)		SF (13)		SMF (5)		MDF (2)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Electricity	5	100	10	100	13	100	5	100	2	100	35	100

Existence of sanitary toilet facility: The data on availability of toilet facility in Yadgir Rf-1 Micro watershed is presented in Table 50. The results indicated that, 57.14 per cent of the households possess toilets.

Table 50. Existence of sanitary toilet facility in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (5)		MF (10)		SF (13)		SMF (5)		MDF (2)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Sanitary toilet facility	5	100	7	70	3	23.08	2	40	3	150	20	57.1

Possession of PDS card: The data regarding possession of PDS card in Yadgir Rf-1 Micro watershed is presented in Table 51. The results indicated that, 100.00 per cent of the households possessed BPL card.

Table 51. Possession of PDS card in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (5)		MF (10)		SF (13)		SMF (5)		MDF (2)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	BPL	5	100	10	100	13	100	5	100	2	100	35	100

Participation in NREGA programme: The data regarding Participation in NREGA programme in Yadgir Rf-1 Micro watershed is presented in Table 52. The results indicated that, only 48.57 percent of the households have participated in NREGA programme.

Table 52. Participation in NREGA programme in Yadgir Rf-1 micro-watershed

Sl. No.	Particulars	LL (5)		MF (10)		SF (13)		SMF (5)		MDF (2)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Participation in NREGA programme	0	0	5	50	10	76.9	3	60	0	0	17	48.6

Adequacy of food items: The data regarding adequacy of food items in Yadgir Rf-1 Micro watershed is presented in Table 53. The results indicated that, the extent of adequacy of food items for cereals, pulses, Oilseeds and vegetables were 100.00, 100.00, 57.14, 40.00 per cent respectively, similarly for Fruits (42.86%), milk (42.86%).

Table 53. Adequacy of food items in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (5)		MF (10)		SF (13)		SMF (5)		MDF (2)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Cereals	5	100	10	100	13	100	5	100	2	100	35	100
2	Pulses	5	100	10	100	13	100	5	100	2	100	35	100
3	Oilseed	0	0	6	60	8	61.54	4	80	2	100	20	57.14
4	Vegetables	0	0	3	30	5	38.46	4	80	2	100	14	40
5	Fruits	5	100	4	40	5	38.46	1	20	0	0	15	42.86
6	Milk	5	100	4	40	5	38.46	1	20	0	0	15	42.86

Table 54. Inadequacy of food items in Yadgir Rf-1 micro-watershed

Sl.No.	Particulars	LL (5)		MF (10)		SF (13)		SMF (5)		MDF (2)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Oilseed	5	100	4	40	5	38.46	1	20	0	0	15	42.86
2	Vegetables	5	100	7	70	8	61.54	1	20	0	0	21	60
3	Fruits	0	0	6	60	8	61.54	4	80	2	100	20	57.14
4	Milk	0	0	6	60	8	61.54	4	80	2	100	20	57.14
5	Egg	5	100	10	100	13	100	5	100	2	100	35	100
6	Meat	5	100	10	100	13	100	5	100	2	100	35	100

Inadequacy of food items: The data regarding in adequacy of food items in Yadgir Rf-1 Micro watershed is presented in Table 54. The results indicated that, the extent of in

adequacy of food items for Oilseeds and vegetables were 42.86, 60.00 and 100.00 per cent respectively, similarly for fruits (57.14%), milk (57.14%), egg (100.00%) and meat (100.00%).

Farming constraints: The data regarding farming constraints experienced by households in Yadgir Rf-1 Micro watershed is presented in Table 55. The results indicated that, lower fertility status of the soil was the constraint experienced by (85.71 %) per cent of the households, wild animal menace on farm field (82.86%), frequent incidence of pest and diseases (85.71%), inadequacy of irrigation water (85.71%), high cost of fertilizers and plant protection chemicals (80.00%), high rate of interest on credit (80.00%), low price for the agricultural commodities (82.86 %), lack of marketing facilities in the area (74.29%), inadequate extension services (71.43 %) and lack of transport for safe transport of the agricultural produce to the market (82.86%).

Table 55. Farming constraints experienced in Yadgir Rf-1 micro-watershed

SN	Particulars	MF (10)		SF (13)		SMF (5)		MDF (2)		All (35)	
		N	%	N	%	N	%	N	%	N	%
1	Lower fertility status of the soil	10	100	13	100	5	100	2	100	30	85.71
2	Wild animal menace on farm field	10	100	13	100	5	100	1	50	29	82.86
3	Frequent incidence of pest and diseases	10	100	13	100	5	100	2	100	30	85.71
4	Inadequacy of irrigation water	10	100	13	100	5	100	2	100	30	85.71
5	High cost of Fertilizers and plant protection chemicals	10	100	13	100	4	80	1	50	28	80
6	High rate of interest on credit	10	100	13	100	3	60	2	100	28	80
7	Low price for the agricultural commodities	10	100	13	100	4	80	2	100	29	82.86
8	Lack of marketing facilities in the area	9	90	13	100	3	60	1	50	26	74.29
9	Inadequate extension services	9	90	11	84.62	4	80	1	50	25	71.43
10	Lack of transport for safe transport of the Agril produce to the market.	10	100	13	100	4	80	2	100	29	82.86

SUMMARY AND IMPLICATIONS

In order to assess the socio-economic condition of the farmers in the watershed 35 households located in the micro watershed were interviewed for the survey. The study was conducted in Yadgir Rf-1 micro-watershed (Chikka Alur sub-watershed, Yadgiri taluk & District) is located at North latitude $16^{\circ} 56' 33.469''$ and $16^{\circ} 56' 0.445''$ and East longitude $77^{\circ} 11' 20.418''$ and $77^{\circ} 9' 57.462''$ covering an area of about 142.59 ha bounded by unde Baggalamadu, Alur. B and Bhimanahalli Villages.

Socio-economic analysis of Yadgir Rf-1 micro watersheds of Chikka Alur sub-watershed, Yadgiri taluk & District indicated that, out of the total sample of 35 farmers were sampled in Yadgir Rf-1 micro-watershed among households surveyed 10 (28.57%) were marginal, 13 (37.14%) were small, 5 (14.29 %) were semi medium and 2 (5.71 %) were medium farmers. 5 landless farmers were also interviewed for the survey. The population characteristics of households indicated that, there were 97 (58.79%) men and 68 (41.21 %) were women. The average population of landless was 5.2, marginal farmers were 3.4, small farmers were 5.2, semi medium farmers were 5.2 and medium farmers were 6. Majority of the respondents (43.03%) were in the age group of 16-35 years.

Education level of the sample households indicated that, there were 46.67 per cent illiterates, 49.08 per cent pre university education and 4.24 per cent attained graduation. About, 48.57 per cent of household heads practicing agriculture and 42.86 per cent of the household heads were engaged as agricultural labourers. Agriculture was the major occupation for 29.09 per cent of the household members.

In the study area, 68.57 per cent of the households possess katcha house and 2.86 per cent possess pucca house. The durable assets owned by the households showed that, 60.00 per cent possess TV, 57.14 per cent possess mixer grinder, 97.14 per cent possess mobile phones and 14.29 per cent possess motor cycles.

Farm implements owned by the households indicated that, 34.29 per cent of the households possess plough, 20.00 per cent possess bullock cart and 11.43 per cent possess sprayer. Regarding livestock possession by the households, 11.43 per cent possess local cow and 2.86 per cent possess buffalo.

The average labour availability in the study area showed that, own labour men available in the micro watershed was 1.57, women available in the micro watershed was 1.30, hired labour (men) available was 7.23 and hired labour (women) available was 11.1.

Out of the total land holding of the sample respondents 100.00 per cent (49.19 ha) of the area is under dry condition. The major crops grown by sample farmers are Red gram, Green gram, Maize and cropping intensity was recorded as 100.00 per cent.

Out of the sample households 97.14 percent possessed bank account and 17.14 per cent of them have savings in the account. About 68.57 per cent of the respondents borrowed credit from various sources. Among the credit borrowed by households, 6.25 per cent have borrowed loan from commercial banks and 75.00 per cent from co-operative/Grameena bank. Majority of the respondents (100.00%) have borrowed loan for agriculture purpose.

Regarding the opinion on institutional sources of credit, 92.86 per cent of the households opined that credit helped to perform timely agricultural operations, while, only 7.14 per cent respondents opined that loan amount was adequate to fulfil their requirement. The per hectare cost of cultivation for Red gram, Green gram and Maize was Rs.21228.66 , 15877.36 and 20572.72 with benefit cost ratio of 1:1.12, 1: 1.90 and 1: 1.72 respectively.

Further, 20.00 per cent of the households opined that dry fodder was adequate. The average annual gross income of the farmers was Rs. 75218.57 in micro-watershed, of which Rs. 37504.29 comes from agriculture. Sampled households have grown 10 horticulture trees and 20 forestry trees together in the fields and back yards. About 60.00 per cent of the households shown interest to cultivate horticultural crops.

Households have an average investment capacity of Rs. 2200.00 for land development. Source of funds for additional investment is concerned, 54.29 per cent depends on bank loan for land development activities.

Regarding marketing channels, 85.71 per cent of the households have sold agricultural produce to the local/village merchants. Further, 31.43 per cent of the households have used tractor for the transport of agriculture commodity. Majority of the farmers (54.29%) have experienced soil and water erosion problems in the watershed and 85.71 per cent of the households were interested towards soil testing.

Fire was the major source of fuel for domestic use for 45.71 per cent of the households and 54.29 per cent households has LPG connection. Piped supply was the major source for drinking water for 100.00 per cent of the households. Electricity was the major source of light for 100.00 per cent of the households.

In the study area, 57.14 per cent of the households possess toilet facility. Regarding possession of PDS card, 100.00 per cent of the households possessed BPL card. Households opined that, the requirement of cereals (100.00%), pulses (100.00%) and oilseeds (57.14%) are adequate for consumption.

Farming constraints experienced by households in the micro watersheds were lower fertility status of the soil (85.71%) wild animal menace on farm field (82.86%), frequent incidence of pest and diseases (85.71%), inadequacy of irrigation water (85.71%), high cost of fertilizers and plant protection chemicals (80.00%), high rate of

interest on credit (80.00%), low price for the agricultural commodities (82.86%), lack of marketing facilities in the area (74.29%), inadequate extension services (71.43%), lack of transport for safe transport of the agricultural produce to the market (82.86%).

Implications of the survey

- ✓ Result indicated that, there were 46.67 per cent were illiterate hence, extension methodologies such as demonstration, street play, drama, video shows will be effective in dissemination of the technologies in the micro watershed.
- ✓ The data indicate that, 68.57 per cent of the households possess katcha house. Hence, the development department while implementing the watershed plan should focus on agriculture to enhance the productivity of major crops in the area to increase the income of the farmers.
- ✓ Results indicated that the local institutional participation of the household members in the micro watershed is minimal hence, activities like membership campaign, awareness creation about the benefits of membership in local institutions and strengths of organized groups must be conveyed.
- ✓ Majority of the households in the watershed have experience in use of mobile phones, and television hence, these mass media can be effectively utilized for transfer of technology as well as for information dissemination.
- ✓ The farm machinery/implement possession in the micro watershed was found to be minimum the reasons may lack of knowledge or lack of financial ability which can be addressed through training on use of different farm implements, providing information on different sources of finance for purchase of farm implements.
- ✓ The possession of livestock such as crossbred cow found is less hence, farmers must be made aware of the benefits of crossbred cow in increased milk production.
- ✓ The possession of livestock such as sheep, goat and poultry was found to be low hence, farmers may be informed the role of subsidiary enterprises in enhancing the income and information on financial support for subsidiary activities.
- ✓ The data indicate that, job/work was the reason for all the migrants hence, farmers may be trained on profitable agriculture or self employment such as animal husbandry, plate making, sheep rearing, goat rearing, rabbit rearing with suitable information on sources of financial support.
- ✓ The results indicate that there was a change in quality of life due to migration hence, the developmental departments should take actions to arrest migration and to improve the quality of the life in rural areas.
- ✓ Households possess 49.19ha (100.00 %) of dry land of irrigated land hence, the availability of the dry land agricultural technologies such as short duration crops, high yielding drought resistance crop varieties, drip irrigation technology and subsidy information will be helpful for the farmers to enhance the productivity of land and as well as farmers income.

- ✓ Few of the bore well in micro watershed found non functional hence, farmers may be trained on possibility of bore well rejuvenation.
- ✓ Bore well was major source of irrigation for 0.00 per cent of the households. Hence, in order to increase the area under irrigation as well as to increase the water use efficiency farmers may trained on drip irrigation and provide the information on subsidy for drip irrigation equipment's along with the information on different agencies which provides the financial assistance for drip irrigation.
- ✓ The cropping intensity in the micro watershed was found to be (100.00 %) hence, care must be taken by the implementing agency to bring uncultivated land into cultivation through suitable measures.
- ✓ Many of the household members have borrowed loan from cooperative banks which has higher rate of interest hence, farmers may be sensitized on the different sources of credit with lesser interest rate such SHGs etc.
- ✓ The results indicated the non availability of both green and dry fodder throughout the year hence, fodder development activities can be taken up in the micro watershed.
- ✓ The average annual gross income of the households Rs.37504.29 from agriculture and Rs. 25142.86 from wages. Agriculture was found to be the major source of income for households hence; the development activities should focus on productivity enhancement, marketing arrangements and agricultural technology dissemination to have a direct impact on the farmers.
- ✓ The cultivation of forest species is found minimal hence, information and production technology related to agro-forestry and integrated farming system.
- ✓ The data indicated that, 54.29 per cent of the households have experienced soil and water erosion problems. Hence, those farmers who reported the soil and water erosion problems may be given attention while implementation of the watershed development plan.
- ✓ The data indicated that, 85.71 per cent of the households have interest in soil testing hence, farmers must be provided with the information on various institutions which are involved in soil testing for the benefit of the farmers.
- ✓ Except summer ploughing the adoption of other soil and water conservation structures is minimum hence, the farmers in the micro watershed should be sensitized on the use of different conservation structures for soil water conservation.
- ✓ Cereals and pulses found be adequate for per cent of the households respectively hence, farm households and the farm women must be trained on importance of balanced nutrition and role of vegetable, milk, egg, meat in balanced diet.
- ✓ Lower fertility status of the soil (85.71%), wild animal menace on farm field (82.86%), frequent incidence of pest and diseases (85.71%), high cost of fertilizers and plant protection chemicals (80.00%), high rate of interest on credit (80.00%), low price for the agricultural commodities (82.86%), lack of marketing facilities in the area (74.29%), inadequate extension services (71.43%), lack of transport for safe

transport of the agricultural produce to the market (82.86%) were the major farming constraints experienced hence, these constraints must be addressed immediately for the welfare of the farmers. Awareness to be created among the farmers to approach nearest KVKs/RSKs and other developmental departments for technical and for subsidized inputs and utilize the well established regulated markets, approaching the contract firms, direct markets to avoid the involvement of middlemen.