



हर कदम, हर डगर
किसानों का हमसफर
भारतीय कृषि अनुसंधान परिषद

AgriSearch with a human touch

**LAND RESOURCE INVENTORY AND SOCIO-ECONOMIC STATUS OF
FARM HOUSEHOLDS FOR WATERSHED PLANNING AND
DEVELOPMENT**

KESALAPURA-1 (4D4A2Q1e) MICRO WATERSHED

Alavandi Hobli, Koppal Taluk and District, Karnataka

Karnataka Watershed Development Project – II

SUJALA – III

World Bank funded Project



ICAR – NATIONAL BUREAU OF SOIL SURVEY AND LAND USE PLANNING



ICAR - NBSS & LUP



**WATERSHED DEVELOPMENT DEPARTMENT
GOVT. OF KARNATAKA, BANGALORE**



About ICAR - NBSS&LUP

The ICAR-National Bureau of Soil Survey and Land Use Planning (ICAR-NBSS&LUP), Nagpur, a premier Institute of the Indian Council of Agricultural Research (ICAR), was set up during 1976 with the objective to prepare soil resource maps at national, state and district levels and to provide research inputs in soil resource mapping and its applications, land evaluation, land use planning, land resource management, and database management using GIS for optimising land use on different kinds of soils in the country.

The Bureau has been engaged in carrying out soil resource survey, agro-ecological and soil degradation mapping at the country, state and district levels for qualitative assessment and monitoring the soil health towards viable land use planning. The research activities have resulted in identifying the soil potentials and problems, and the various applications of the soil surveys with the ultimate objective of sustainable agricultural development. The Bureau has the mandate to correlate and classify soils of the country and maintain a National Register of all the established soil series. The Institute is also imparting in-service training to staff of the soil survey agencies in the area of soil survey, land evaluation and soil survey interpretations for land use planning. The Bureau in collaboration with Panjabrao Krishi Vidyapeeth, Akola is running post-graduate teaching and research programme in land resource management, leading to M.Sc. and Ph.D. degrees.

Citation: Rajendra Hegde, Ramesh Kumar, S.C., K.V. Niranjana, S. Srinivas, M.Lalitha, B.A. Dhanorkar, R.S. Reddy and S.K. Singh (2019). "Land Resource Inventory and Socio-Economic Status of Farm Households for Watershed Planning and Development of Kesalapura-1 (4D4A2Q1e) Microwatershed, Alavandi Hobli, Koppal Taluk and District, Karnataka", ICAR-NBSS&LUP Sujala MWS Publ.361, ICAR – NBSS & LUP, RC, Bangalore. p.143 & 39.

TO OBTAIN COPIES,

Please write to:

Director, ICAR - NBSS & LUP,

Amaravati Road, NAGPUR - 440 033, India

Phone : (0712) 2500386, 2500664, 2500545 (O)

Telefax : 0712-2522534

E-Mail : director@nbsslup.ernet.in

Website URL : nbsslup.in

Or

Head, Regional Centre, ICAR - NBSS&LUP, Hebbal, Bangalore - 560 024

Phone : (080) 23412242, 23510350 (O)

Telefax : 080-23510350

E-Mail : nbssrcb@gmail.com

ICAR-NBSS&LUP Sujala MWS Publ.361



हर कदम, हर डगर
किसानों का हमसफर
भारतीय कृषि अनुसंधान परिषद

*Agri*search with a human touch

**LAND RESOURCE INVENTORY AND SOCIO-ECONOMIC
STATUS OF FARM HOUSEHOLDS FOR WATERSHED
PLANNING AND DEVELOPMENT**

KESALAPURA-1 (4D4A2Q1e) MICRO WATERSHED

Alavandi Hobli, Koppal Taluk and District, Karnataka

Karnataka Watershed Development Project – II

Sujala-III

World Bank funded Project



**ICAR – NATIONAL BUREAU OF SOIL SURVEY AND LAND USE
PLANNING**



ICAR - NBSS & LUP



**WATERSHED DEVELOPMENT DEPARTMENT, GOVT. OF
KARNATAKA, BANGALORE**



PREFACE

In Karnataka, as in other Indian States, the livelihoods of rural people are intertwined with farming pursuits. The challenges in agriculture are seriously threatening the livelihood of a large number of farmers as they have been practicing farming in contextual factors beyond their control. Climatic factors are the most important ones and have become much more significant in recent times due to rapid climate changes induced by intensive anthropogenic activities affecting our ecosystem in multiple ways. Climate change has become the reality, it is happening and efforts to evolve and demonstrate climate resilient technologies have become essential. Due to the already over stressed scenario of agrarian sector, the climate change is resulting in manifold increase in the complexities, pushing the rural mass to face more and more unpredictable situations. The rising temperatures and unpredictable rainfall patterns are going to test seriously the informed decisions farmers have to make in order to survive in farming and sustain their livelihood.

It is generally recognized that impacts of climate change shall not be uniform across the globe. It is said that impact of climate change is more severe in South Asia. Based on the analysis of meteorological data, it is predicted that in India, there will be upward trend in mean temperature, downward trend in relative humidity, annual rainfall and number of wet days in a year. Also, in general, phenomena like erratic monsoon, spread of tropical diseases, rise in sea levels, changes in availability of fresh water, frequent floods, droughts, heat waves, storms and hurricanes are predicted. Each one of these adverse situations are already being experienced in various parts of India and also at the global level. Decline in agricultural productivity of small and marginal farmers becoming more vulnerable is already witnessed.

In Karnataka, more than 60 per cent of the population live in rural areas and depend on agriculture and allied activities for their livelihood. Though the state has achieved significant progress in increasing the yield of many crops, there is tremendous pressure on the land resources due to the growing and competing demands of various land uses. This is reflected in the alarming rate of land degradation observed. Already more than 50 per cent of the area is affected by various forms of degradation. If this trend continues, the sustainability of the fragile ecosystem will be badly affected. The adverse effects of change in the climatic factors are putting additional stress on the land resources and the farmers dependent on this.

The natural resources (land, water and vegetation) of the state need adequate and constant care and management, backed by site-specific technological interventions and investments particularly by the government. Detailed database pertaining to the nature of

the land resources, their constraints, inherent potentials and suitability for various land based rural enterprises, crops and other uses is a prerequisite for preparing location-specific action plans, which are in tune with the inherent capability of the resources. Any effort to evolve climate resilient technologies has to be based on the baseline scientific database. Then only one can expect effective implementation of climate resilient technologies, monitor the progress, make essential review of the strategy, and finally evaluate the effectiveness of the implemented programs. The information available at present on the land resources of the state are of general nature and useful only for general purpose planning. Since the need of the hour is to have site-specific information suitable for farm level planning and detailed characterization and delineation of the existing land resources of an area into similar management units is the only option.

ICAR-NBSS&LUP, Regional Centre, Bangalore has taken up a project sponsored by the Karnataka Watershed Development Project-II, (Sujala-III), Government of Karnataka funded by the World Bank under Component -1 Land Resource Inventory. This study was taken up to demonstrate the utility of such a database in reviewing, monitoring and evaluating all the land based watershed development programs on a scientific footing. To meet the requirements of various land use planners at grassroots level, the present study on “Land Resource Inventory and Socio-Economic Status of Farm Households for Watershed Planning and Development of Kesalapura-1 microwatershed in Koppal Taluk, Koppal District, Karnataka” for integrated development was taken up in collaboration with the State Agricultural Universities, IISC, KRSRAC, KSNDMC as Consortia partners. The project provides detailed land resource information at cadastral level (1:7920 scale) for all the plots and socio-economic status of farm households covering thirty per cent farmers randomly selected representing landed and landless class of farmers in the micro-watershed. The project report with the accompanying maps for the microwatershed will provide required detailed database for evolving effective land use plan, alternative land use options and conservation plans for the planners, administrators, agricultural extension personnel, KVK officials, developmental departments and other land users to manage the land resources in a sustainable manner.

It is hoped that this database will be useful to the planners, administrators and developmental agencies working in the area in not only for formulating location specific developmental schemes but also for their effective monitoring at the village/watershed level.

Nagpur
Date:13-09-2019

S.K. SINGH
Director, ICAR - NBSS&LUP Nagpur

Contributors

Dr. Rajendra Hegde Principal Scientist, Head & Project Leader, Sujala-III Project ICAR-NBSS&LUP, Regional Centre, Bangalore	Dr. S.K.Singh Director, ICAR-NBSS&LUP Coordinator, Sujala-III Project Nagpur
Soil Survey, Mapping & Report Preparation	
Dr. K.V. Niranjana	Sh. R.S. Reddy
Dr. B.A. Dhanorkar	Sh. Venkata Giriappa
	Dr. Gopali Bardhan
	Smt. Chaitra, S.P.
	Dr. Gayathri, B.
	Dr. Savitha, H.R.
	Sh. Nagendra, B.R.
	Sh. Somashekar T.N
	Ms. Arpitha, G.M.
Field Work	
Sh. C. Bache Gowda	Sh. Mayur Patil
Sh. Somashekar	Sh. Arun Kumar, S.
Sh. M. Jayaramaiah	Sh. Sunil Raj
	Sh. Yogesh Kumar, B.
	Sh. Vikas, N.K.
	Sh. Arun Kumar, S.G.
	Sh. Umesh Jadiyahappa Madolli
	Sh. Praveen Kumar P. Achalkar
	Sh. Veerabhadraswamy
	Sh. Vinay
	Sh. Shankarappa, K.
	Sh. Lankesh, R.S.
	Sh. Appanna B. Hattigoudar
	Sh. Maharudra
GIS Work	
Dr. S.Srinivas	Sh. A.G. Devendra Prasad
Sh. D. H.Venkatesh	Sh. Abhijith Sastry, N.S.
Smt. K.Sujatha	Sh. Nagendra Babu Kolukondu
Smt. K. V. Archana	Sh. Avinash
Sh. N. Maddileti	Sh. Amar Suputhra, S.
	Sh. Deepak M.J.
	Sh. Madappaswamy
	Smt. K. Karunya Lakshmi
	Ms. Seema, K.V.
	Ms. Ramireddy Lakshmi Silpa
	Ms. Bhanu Rekha, T.
	Ms. Rajata Bhat
	Ms. Shruthi
	Ms. Suman, S.

Laboratory Analysis	
Dr. M. Lalitha	Ms. Thara, V.R.
Smt. Arti Koyal	Ms. Roopa, G.
Smt. Parvathy, S.	Ms. Vindhya, N.G.
	Ms. Shwetha N.K.
	Ms. Pavana Kumari, P.
	Ms. Leelavathy, K.U.
	Ms. Rashmi, N.
	Ms. Padmaja, S.
	Ms. Veena, M.
	Ms. Chaithrashree B
	Ms. Shwetha N
Socio-economic Analysis	
Dr. Ramesh Kumar, S.C.	Sh. Prakashanaik, M.K.
	Ms. Shraddha Hegde
	Sh. Vinod R
	Sh. Basavaraj
	Ms. Sowmya K.B
	Mrs. Prathibha, D.G
	Sh. Rajendra,D
Soil & Water Conservation	
Sh. Sunil P. Maske	
Watershed Development Department, GOK, Bangalore	
Sh. Rajeev Ranjan IFS Project Director & Commissioner, WDD	Dr. A. Natarajan NRM Consultant, Sujala-III Project
Dr. S.D. Pathak IFS Executive Director & Chief Conservator of Forests, WDD	

PART-A

LAND RESOURCE INVENTORY

Contents

Preface		
Contributors		
Executive Summary		
Chapter 1	Introduction	1
Chapter 2	Geographical Setting	3
2.1	Location and Extent	3
2.2	Geology	3
2.3	Physiography	4
2.4	Drainage	5
2.5	Climate	5
2.6	Natural Vegetation	6
2.7	Land Utilization	7
Chapter 3	Survey Methodology	11
3.1	Base maps	11
3.2	Image Interpretation for Physiography	12
3.3	Field Investigation	14
3.4	Soil mapping	16
3.5	Laboratory Characterization	16
3.6	Land Management Units	19
Chapter 4	The Soils	23
4.1	Soils of Granite gneiss	23
4.2	Alluvial Landscape	31
Chapter 5	Interpretation for Land Resource Management	43
5.1	Land Capability Classification	43
5.2	Soil Depth	45
5.3	Surface Soil Texture	46
5.4	Soil Gravelliness	47
5.5	Available Water Capacity	48
5.6	Soil Slope	49
5.7	Soil Erosion	50
Chapter 6	Fertility Status	53
6.1	Soil Reaction (pH)	53
6.2	Electrical Conductivity (EC)	53
6.3	Organic Carbon (OC)	53
6.4	Available Phosphorus	56
6.5	Available Potassium	56
6.6	Available Sulphur	56
6.7	Available Boron	56
6.8	Available Iron	56
6.9	Available Manganese	56
6.10	Available Copper	56
6.11	Available Zinc	60
Chapter 7	Land Suitability for Major Crops	61

7.1	Land suitability for Sorghum	61
7.2	Land suitability for Maize	62
7.3	Land suitability for Bajra	63
7.4	Land suitability for Groundnut	64
7.5	Land suitability for Sunflower	65
7.6	Land suitability for Red gram	66
7.7	Land suitability for Bengalgram	67
7.8	Land suitability for Cotton	68
7.9	Land suitability for Chilli	69
7.10	Land suitability for Tomato	70
7.11	Land suitability for Brinjal	71
7.12	Land suitability for onion	72
7.13	Land suitability for Bhindi	73
7.14	Land suitability for Drumstick	74
7.15	Land suitability for Mango	75
7.16	Land suitability for Guava	76
7.17	Land suitability for Sapota	77
7.18	Land suitability for Pomegranate	78
7.19	Land suitability for Musambi	79
7.20	Land suitability for Lime	80
7.21	Land suitability for Amla	81
7.22	Land suitability for Cashew	82
7.23	Land suitability for Jackfruit	83
7.24	Land Suitability for Jamun	84
7.25	Land Suitability for Custard apple	85
7.26	Land Suitability for Tamarind	86
7.27	Land Suitability for Mulberry	87
7.28	Land Suitability for Marigold	88
7.29	Land suitability for Chrysanthemum	89
7.30	Land suitability for Jasmine	90
7.31	Land suitability for Crossandra	91
7.32	Land management units	125
7.33	Proposed Crop Plan	126
Chapter 8	Soil Health Management	129
Chapter 9	Soil and Water conservation Treatment Plan	135
9.1	Treatment Plan	136
9.2	Recommended Soil and Water Conservation measures	139
9.3	Greening of microwatershed	140
	References	143
	Appendix I	I-V
	Appendix II	VI-X
	Appendix III	XI-XIV

LIST OF TABLES

2.1	Mean Monthly Rainfall, PET, 1/2 PET at Koppal Taluk and District	5
2.2	Land Utilization in Koppal District	7
3.1	Differentiating Characteristics used for Identifying Soil Series	15
3.2	Soil map unit description of Kesalapura-1 microwatershed	17
4.1	Physical and chemical characteristics of soil series identified in Kesalapura-1 microwatershed	32
7.1	Soil-Site Characteristics of Kesalapura-1 microwatershed	93
7.2	Land suitability for Sorghum	94
7.3	Land suitability for Maize	95
7.4	Land suitability for Bajra	96
7.5	Land suitability for Groundnut	97
7.6	Land suitability for Sunflower	98
7.7	Land suitability for Red gram	99
7.8	Land suitability for Bengalgram	100
7.9	Land suitability for Cotton	101
7.10	Land suitability for Chilli	102
7.11	Land suitability for Tomato	103
7.12	Land suitability for Brinjal	104
7.13	Land suitability for onion	105
7.14	Land suitability for Bhindi	106
7.15	Land suitability for Drumstick	107
7.16	Land suitability for Mango	108
7.17	Land suitability for Guava	109
7.18	Land suitability for Sapota	110
7.19	Land suitability for Pomegranate	111
7.20	Land suitability for Musambi	112
7.21	Land suitability for Lime	113
7.22	Land suitability for Amla	114
7.23	Land suitability for Cashew	115
7.24	Land suitability for Jackfruit	116

7.25	Land Suitability for Jamun	117
7.26	Land Suitability for Custard apple	118
7.27	Land Suitability for Tamarind	119
7.28	Land Suitability for Mulberry	120
7.29	Land Suitability for Marigold	121
7.30	Land suitability for Chrysanthemum	122
7.31	Land suitability for Jasmine	123
7.32	Land suitability for Crossandra	124
7.33	Proposed Crop Plan for Kesalapura-1 microwatershed	127

LIST OF FIGURES

2.1	Location map of Kesalapura-1 microwatershed	3
2.2a	Granite and granite gneiss rocks	4
2.2b	Alluvial rocks	4
2.3	Rainfall distribution in Koppal Taluk, Koppal District	6
2.4	Natural vegetation of Kesalapura-1 microwatershed	6
2.5a	Different crops and cropping systems in Kesalapura-1 microwatershed	8
2.5b	Different crops and cropping systems in Kesalapura-1 microwatershed	9
2.6	Current Land use – Kesalapura-1 microwatershed	10
2.7	Location of Wells- Kesalapura-1 microwatershed	10
3.1	Scanned and Digitized Cadastral map of Kesalapura-1 microwatershed	13
3.2	Satellite image of Kesalapura-1 microwatershed	13
3.3	Cadastral map overlaid on IRS PAN+LISS IV merged imagery of Kesalapura-1 microwatershed	14
3.4	Location of profiles in a transect	15
3.5	Soil phase or management units of Kesalapura-1 microwatershed	21
5.1	Land Capability Classification of Kesalapura-1 microwatershed	44
5.2	Soil Depth map of Kesalapura-1 microwatershed	46
5.3	Surface Soil Texture map of Kesalapura-1 microwatershed	47
5.4	Soil Gravelliness map of Kesalapura-1 microwatershed	48
5.5	Soil Available Water Capacity map of Kesalapura-1 microwatershed	49
5.6	Soil Slope map of Kesalapura-1 microwatershed	50
5.7	Soil Erosion map of Kesalapura-1 microwatershed	51
6.1	Soil Reaction (pH) map of Timmapur-2Microwatershed	54
6.2	Electrical Conductivity (EC) map of Kesalapura-1 microwatershed	54
6.3	Soil Organic Carbon (OC) map of Kesalapura-1 microwatershed	55
6.4	Soil Available Phosphorus map of Kesalapura-1 microwatershed	55
6.5	Soil Available Potassium map of Kesalapura-1 microwatershed	57
6.6	Soil Available Sulphur map of Kesalapura-1 microwatershed	57
6.7	Soil Available Boron map of Kesalapura-1 microwatershed	58
6.8	Soil Available Iron map of Kesalapura-1 microwatershed	58
6.9	Soil Available Manganese map of Kesalapura-1 microwatershed	59
6.10	Soil Available Copper map of Kesalapura-1 microwatershed	59

6.11	Soil Available Zinc map of Kesalapura-1 microwatershed	60
7.1	Land suitability for Sorghum	62
7.2	Land suitability for Maize	63
7.3	Land suitability for Bajra	64
7.4	Land suitability for Groundnut	65
7.5	Land suitability for Sunflower	66
7.6	Land suitability for Red gram	67
7.7	Land suitability for Bengalgram	68
7.8	Land suitability for Cotton	69
7.9	Land suitability for Chilli	70
7.10	Land suitability for Tomato	71
7.11	Land suitability for Brinjal	72
7.12	Land suitability for onion	73
7.13	Land suitability for Bhindi	74
7.14	Land suitability for Drumstick	75
7.15	Land suitability for Mango	76
7.16	Land suitability for Guava	77
7.17	Land suitability for Sapota	78
7.18	Land suitability for Pomegranate	79
7.19	Land suitability for Musambi	80
7.20	Land suitability for Lime	81
7.21	Land suitability for Amla	82
7.22	Land suitability for Cashew	83
7.23	Land suitability for Jackfruit	84
7.24	Land Suitability for Jamun	85
7.25	Land Suitability for Custard apple	86
7.26	Land Suitability for Tamarind	87
7.27	Land Suitability for Mulberry	88
7.28	Land Suitability for Marigold	89
7.29	Land suitability for Chrysanthemum	90
7.30	Land suitability for Jasmine	91
7.31	Land suitability for Crossandra	92
7.32	Land Management Units of Kesalapura-1 microwatershed	126
9.2	Soil and water conservation map of Kesalapura-1 microwatershed	140

EXECUTIVE SUMMARY

The land resource inventory of Kosalapura-1 microwatershed was conducted using village cadastral maps and IRS satellite imagery on 1:7920 scale. The false colour composites of IRS imagery were interpreted for physiography and these physiographic delineations were used as base for mapping soils. The soils were studied in several transects and a soil map was prepared with phases of soil series as mapping units. Random checks were made all over the area outside the transects to confirm and validate the soil map unit boundaries. The soil map shows the geographic distribution and extent, characteristics, classification, behavior and use potentials of the soils in the microwatershed.

The present study covers an area of 575 ha in Koppal taluk and district, Karnataka. The climate is semiarid and categorized as drought - prone with an average annual rainfall of 662 mm, of which about 424 mm is received during south-west monsoon, 161 mm during north-east and the remaining 77 mm during the rest of the year. An area of about 96 per cent is covered by soils and 4 per cent by habitation and water bodies, settlements and others. The salient findings from the land resource inventory are summarized briefly below.

- ❖ The soils belong to 12 soil series and 18 soil phases (management units) and 5 land management units.*
- ❖ The length of crop growing period is <90 days and starts from 2nd week of August to 2nd week of November.*
- ❖ From the master soil map, several interpretative and thematic maps like land capability, soil depth, surface soil texture, soil gravelliness, available water capacity, soil slope and soil erosion were generated.*
- ❖ Soil fertility status maps for macro and micronutrients were generated based on the surface soil samples collected at every 320 m grid interval.*
- ❖ Land suitability for growing 31 major agricultural and horticultural crops were assessed and maps showing the degree of suitability along with constraints were generated.*
- ❖ Entire area is suitable for agriculture.*
- ❖ About 15 per cent of the soils are moderately shallow (50-75 cm), 43 per cent of the soils are moderately deep (75-100 cm), 21 per cent area has deep (100-150 cm) and 17 per cent area has very deep (>150 cm) soils.*
- ❖ An area of about 55 per cent has loamy soils and 41 per cent has clayey soils at the surface.*
- ❖ About 42 per cent of the area has non-gravelly (<15%) soils, 44 per cent gravelly (15-35% gravel) and 10 per cent very gravelly (35-60%) soils.*
- ❖ About 24 per cent are very low (<50 mm/m), 20 per cent low (51-100 mm/m), 36 per cent medium (101-150 mm/m), 15 per cent high (151-200 mm/m) and 2 per cent very high (>200 mm/m) in available water capacity.*

- ❖ *About 1 per cent area has nearly level (0-1%) and 95 per cent area has very gently sloping (1-3%) lands.*
- ❖ *An area of about 37 per cent has soils that are slightly eroded (e1) and 59 per cent moderately eroded (e2) lands.*
- ❖ *An area of about 61 per cent are moderately alkaline (pH pH 7.8-8.4), 29 per cent are strongly alkaline (pH 8.4-9.0) and 5 per cent are very strongly alkaline (pH >9.0) in soil reaction.*
- ❖ *The Electrical Conductivity (EC) of the soils is $<2 \text{ dS m}^{-1}$ and as such the soils are non-saline.*
- ❖ *Organic carbon is low ($<0.5\%$) in 18 per cent, medium (0.5-0.75%) in 70 per cent and high ($>0.75\%$) in 8 per cent area of the soils.*
- ❖ *Available phosphorus is medium (23-57 kg/ha) in the entire area of the microwatershed.*
- ❖ *About 96 per cent are medium (145-337 kg/ha) and <1 per cent soils are high (>337 kg/ha) in available potassium content.*
- ❖ *Available sulphur is low (<10 ppm) in 94 per cent and medium (10-20 ppm) in 2 per cent in the microwatershed.*
- ❖ *Available boron is low (0.5 ppm) in about 2 per cent, medium (0.5-1.0 ppm) in 82 per cent and high (>1.0 ppm) in 12 per cent area.*
- ❖ *Available iron is deficient (<4.5 ppm) in the entire area of the microwatershed.*
- ❖ *Available zinc is deficient (<0.6 ppm) in 6 per cent and sufficient (>0.6 ppm) in about 90 per cent area.*
- ❖ *Available manganese and copper are sufficient in all the soils.*
- ❖ *The land suitability for 31 major agricultural and horticultural crops grown in the microwatershed were assessed and the areas that are highly suitable (S1) and moderately suitable (S2) are given below. It is however to be noted that a given soil may be suitable for various crops but what specific crop to be grown may be decided by the farmer looking to his capacity to invest on various inputs, marketing infrastructure, market price and finally the demand and supply position.*

Land suitability for various crops in the microwatershed

Crop	Suitability Area in ha (%)		Crop	Suitability Area in ha (%)	
	Highly suitable (S1)	Moderately suitable (S2)		Highly suitable (S1)	Moderately suitable (S2)
<i>Sorghum</i>	192 (33)	227 (39)	<i>Sapota</i>	123 (21)	224 (39)
<i>Maize</i>	85 (15)	333 (58)	<i>Pomegranate</i>	123 (21)	282 (49)
<i>Bajra</i>	266 (46)	226 (39)	<i>Musambi</i>	132 (23)	273 (47)
<i>Groundnut</i>	38 (7)	457 (80)	<i>Lime</i>	132 (23)	320 (56)
<i>Sunflower</i>	132 (23)	273 (47)	<i>Amla</i>	300 (52)	252 (44)
<i>Red gram</i>	123 (21)	234 (41)	<i>Cashew</i>	99 (17)	286 (50)
<i>Bengalgram</i>	10 (2)	435 (76)	<i>Jackfruit</i>	123 (21)	272 (47)
<i>Cotton</i>	96 (17)	323 (56)	<i>Jamun</i>	86 (15)	356 (62)
<i>Chilli</i>	219 (38)	153 (27)	<i>Custard apple</i>	300 (52)	252 (44)
<i>Tomato</i>	219 (38)	153 (27)	<i>Tamarind</i>	86 (15)	119 (21)
<i>Brinjal</i>	204 (36)	212 (37)	<i>Mulberry</i>	147 (26)	307 (53)
<i>Onion</i>	61 (11)	298 (52)	<i>Marigold</i>	123 (21)	296 (52)
<i>Bhendi</i>	61 (11)	355 (62)	<i>Chrysanthemum</i>	123 (21)	296 (52)
<i>Drumstick</i>	147 (26)	221 (39)	<i>Jasmine</i>	123 (21)	239 (42)
<i>Mango</i>	86 (15)	62 (11)	<i>Crossandra</i>	123 (21)	239 (42)
<i>Guava</i>	74 (13)	320 (56)			

- ❖ *Apart from the individual crop suitability, a proposed crop plan has been prepared for the 5 identified LMUs by considering only the highly and moderately suitable lands for different crops and cropping systems with food, fodder, fibre and other horticulture crops that helps in maintaining productivity and ecological balance in the microwatershed.*
- ❖ *Maintaining soil-health is vital for crop production and conserve soil and land resource base for maintaining ecological balance and to mitigate climate change. For this, several ameliorative measures have been suggested for these problematic soils like saline/alkali, highly eroded, sandy soils etc.*
- ❖ *Soil and water conservation treatment plan has been prepared that would help in identifying the sites to be treated and also the type of structures required.*
- ❖ *As part of the greening programme, several tree species have been suggested to be planted in marginal and submarginal lands, field bunds and also in the hillocks, mounds and ridges. That would help in supplementing the farm income, provide fodder and fuel, and generate lot of biomass which in turn would help in maintaining the ecological balance and contribute to mitigating the climate change.*

INTRODUCTION

Land is a scarce resource and basic unit for any material production. It can support the needs of the growing population, provided they use the land in a rational and judicious manner. But what is happening in many areas of the state is a cause for concern to everyone involved in the management of land resources at the grassroots level. The area available for agriculture is about 51 per cent of the total area and more than 60 per cent of the people are still dependant on agriculture for their livelihood. The limited land area is under severe stress and strain due to increasing population pressure and competing demands of various land uses. Due to this, every year there is significant diversion of farm lands and water resources for non-agricultural purposes. Apart from this, due to lack of interest in farmers for farming, large tracts of cultivable lands are turning into fallows in many areas and this trend is continuing at an alarming rate.

Further, land degradation has emerged as a serious problem which has already affected about 38 lakh ha of cultivated area in the state. Soil erosion alone has degraded about 35 lakh ha. Almost all the uncultivated areas are facing various degrees of degradation, particularly soil erosion. Salinity and alkalinity has emerged as a major problem in more than 3.5 lakh ha in the irrigated areas of the state. Nutrient depletion and declining factor productivity is common in both rainfed and irrigated areas. The degradation is continuing at an alarming rate and there appears to be no systematic effort among the stakeholders to contain this process. In recent times, an aberration of weather due to climate change phenomenon has added another dimension leading to unpredictable situations to be tackled by the farmers.

In this critical juncture, the challenge before us is not only to increase the productivity per unit area which is steadily declining and showing a fatigue syndrome, but also to prevent or at least reduce the severity of degradation. If the situation is not reversed at the earliest, then the sustainability of the already fragile crop production system and the overall ecosystem will be badly affected in the state.

The continued neglect and unscientific use of the resources for a long time has led to the situation observed at present in the state. It is a known fact and established beyond doubt by many studies in the past that the cause for all kinds of degradation is the neglect and irrational use of the land resources. Hence, there is urgent need to generate a detailed site-specific farm level database on various land resources for all the villages/watersheds in a time bound manner that would help to protect the valuable soil and land resources and also to stabilize the farm production.

Therefore, the land resource inventory required for farm level planning is the one which investigates not only the surface but also consider the other parameters which are critical for productivity *viz.*, soils, climate, water, minerals and rocks, topography, geology, hydrology, vegetation, crops, land use pattern, animal population, socio-economic conditions, infrastructure, marketing facilities and various schemes and

developmental works of the government etc. From the data collected at farm level, the specific problems and potentials of the area can be identified and highlighted, conservation measures required for the area can be planned on a scientific footing, suitability of the area for various uses can be worked out and finally viable and sustainable land use options suitable for each and every land holding can be prescribed.

The Land Resource Inventory is basically done for identifying potential and problem areas, developing sustainable land use plans, estimation of surface run off and water harvesting potential, preparation of soil and water conservation plans, land degradation/desertification etc. The Bureau is presently engaged in developing an LRI methodology using high resolution satellite remote sensing data and Digital Elevation Model (DEM) data to prepare Landscape Ecological Units (LEU) map representing agro-ecosystem as a whole. The LEU is preferred over landform as the base map for LRI. LEU is the assemblage of landform, slope and land use. An attempt was made to upscale the soil resource information from 1:250000 and 1:50000 scale to the LEU map in Goa and other states.

The land resource inventory aims to provide site-specific database for Kesalapura-1 Microwatershed in Koppal Taluk and District, Karnataka State for the Karnataka Watershed Development Department. The database was generated by using cadastral map of the village as a base along with high resolution IRS LISS IV and Cartosat-1 merged satellite imagery. Later, an attempt will be made to uplink this LRI data generated at 1:7920 scale under Sujala-III Project to the proposed Landscape Ecological Units (LEUs) map.

The study was organized and executed by the ICAR- National Bureau of Soil Survey and Land Use Planning, Regional Centre, Bangalore under Generation of Land Resource Inventory Data Base Component-1 of the Sujala-III Project funded by the World Bank.

GEOGRAPHICAL SETTING

2.1 Location and Extent

The Kesalapura-1 Microwatershed is located in the central part of northern Karnataka in Koppal Taluk and District, Karnataka State (Fig. 2.1). It comprises parts of Kasalapura, Halavagali and Hesaru Villages. It lies between $15^{\circ}28'$ – $15^{\circ}10'$ North latitudes and $76^{\circ}56'$ – $76^{\circ}57'$ East longitudes and covers an area of 575 ha. It is about 32 km from Koppal town. It is surrounded by Kasalapura village on the south, Halavagali village on the east, Hyderanagara village on the north and Hesaru village on the western side.




Fig. 2.1 Location map of Kesalapura-1 Microwatershed

2.2 Geology

Major rock formations observed in the microwatershed are granite gneiss and alluvium (Figs. 2.2a and b). Granite gneisses are essentially pink to gray and are coarse to medium grained. They consist primarily of quartz, feldspar, biotite and hornblende. The gray granite gneisses are highly weathered, fractured and fissured upto a depth of about

10 m. Dolerite dykes and quartz veins are common with variable width and found to occur in the village. The thickness of the alluvium generally is limited to less than a meter, except in river valleys where it is very deep extending to tens of meters. Such soils are transported and represent palaeo black soils originally formed at higher elevation, but now occupying river valleys.



Fig. 2.2 a Granite and granite gneiss rocks



Fig. 2.2 b Alluvium

2.3 Physiography

Physiographically, the area has been identified as granite gneiss and alluvial landscapes based on geology. The microwatershed area has been further divided into summits, very gently sloping uplands and nearly level plains based on slope and its relief features. The elevation ranges from 502 to 516 m in the gently sloping uplands.

2.4 Drainage

The area is drained by several small seasonal streams that join Hire *halla* and Chenna *halla* along its course. Though, the streams are not perennial, during rainy season they carry large quantities of rain water. The microwatershed has only few small tanks which are not able to store the water flowing during the rainy season. Due to this, the ground water recharge is very much affected in the villages. This is reflected in the failure of many bore wells in the villages. If the available rain water is properly harnessed by constructing tanks and recharge structures at appropriate places in the villages, then the drinking and irrigation needs of the area can be easily met. The drainage network is dendritic to sub parallel.

2.5 Climate

The district falls under semiarid tract of the state and is categorized as drought-prone with an average annual rainfall of 662 mm (Table 2.1). Maximum of 424 mm precipitation takes place during the south-west monsoon period from June to September, north-east monsoon contributes about 161 mm and prevails from October to early December and the remaining 77 mm takes place during the rest of the year. The winter season is from December to February. During April and May, the temperatures reach up to 45 °C and in December and January, the temperatures will go down to 16 °C. Rainfall distribution is shown in Figure 2.3. The average Potential Evapo Transpiration (PET) is 145 mm and varies from a low of 101 mm in December and 193 mm in the month of May. The PET is always higher than precipitation in all the months except in the month of September. Generally, the Length of crop Growing Period (LGP) is <90 days and starts from 2nd week of August to 2nd week of November.

Table 2.1 Mean Monthly Rainfall, PET, 1/2 PET at Koppal Taluk and District

Sl. No.	Months	Rainfall	PET	1/2 PET
1	January	1.60	116.70	58.35
2	February	1.50	129.20	64.60
3	March	14.10	169.80	84.90
4	April	18.10	180.60	90.30
5	May	41.60	193.50	96.75
6	June	85.80	167.90	83.95
7	July	72.10	156.20	78.10
8	August	110.50	152.50	76.25
9	September	155.60	138.50	69.25
10	October	116.30	122.30	61.15
11	November	36.00	106.40	53.20
12	December	9.10	101.00	50.50
	TOTAL	662.30	144.55	




Fig. 2.3 Rainfall distribution in Koppal Taluk and District

2.6 Natural Vegetation

The natural vegetation is sparse comprising few tree species, shrubs and herbs. The mounds, ridges and boulders occupy sizeable areas which are under thin to moderately thick forest vegetation. Still, there are some remnants of the past forest cover which can be seen in patches in some ridges and hillocks in the microwatershed (Fig 2.4).

Apart from the continuing deforestation, the presence of large population of goats, sheep and other cattle in the microwatershed is causing vegetative degradation of whatever little vegetation left in the area. The uncontrolled grazing has left no time for the regeneration of the vegetative cover. This leads to the accelerated rate of erosion on the hill slopes, resulting in the formation of deep gullies in the foot slopes and eventually resulting in the heavy siltation of few tanks and reservoirs in the microwatershed.



Fig 2.4 Natural vegetation of Kesalapura-1 Microwatershed

2.7 Land Utilization

About 91 per cent area (Table 2.2) in Koppal district is cultivated at present and about 16 per cent of the area is sown more than once. The cropping intensity is 118 per cent. An area of about 3 per cent is currently barren. Forests occupy a small area of about 5 per cent and the tree cover is in a very poor state. Most of the mounds, ridges and bouldery areas have very poor vegetative cover. Major crops grown in the area are sorghum, maize, bajra, cotton, safflower, sunflower, red gram, horse gram, onion, mulberry, pomegranate, sugarcane, bengalgram and groundnut (Fig 2.5). While carrying out land resource inventory, the land use/land cover particulars are collected from all the survey numbers and a current land use map of the microwatershed is prepared. The current land use map prepared shows the arable and non-arable lands, other land uses and different types of crops grown in the area. The current land use map of Kesalapura-1 Microwatershed is presented in Fig. 2.6. Simultaneously, enumeration of existing wells (bore wells and open wells) and other soil and water conservation structures in the microwatershed is made and their location in different survey numbers is marked on the cadastral map. Map showing the location of wells and other water bodies and conservation structures in Kesalapura-1 Microwatershed is given Fig. 2.7.

Table 2.2 Land Utilization in Koppal District

Sl. no.	Agricultural land use	Area (ha)	Per cent
1	Total geographical area	552495	
2	Total cultivated area	500542	90.6
3	Area sown more than once	92696	16.8
4	Trees and groves	210	0.04
5	Cropping intensity	-	118
6	Forest	29451	5.33
7	Cultivable wasteland	2568	0.46
8	Permanent Pasture land	14675	2.66
9	Barren land	16627	3.01
10	Non agricultural land	40591	7.35
11	Current fallow	19660	3.56



Fig. 2.5 (a) Different crops and cropping systems in Kesalapura-1 Microwatershed



Fig. 2.5 (b) Different crops and cropping systems in Kesalapura-1 Microwatershed




Fig. 2.6 Current Land Use – Kesalapura-1 Microwatershed




Fig. 2.7 Location of wells and conservation structures-Kesalapura-1 Microwatershed

SURVEY METHODOLOGY

The purpose of land resource inventory is to delineate similar areas (soil series and phases), which respond or expected to respond similarly for a given level of management. This was achieved in Kesalapura-1 Microwatershed by the detailed study of all the soil characteristics (depth, texture, colour, structure, consistence, coarse fragments, porosity, soil reaction, soil horizons etc.) and site characteristics (slope, erosion, drainage, occurrence of rock fragments etc.) followed by grouping of similar areas based on soil-site characteristics into homogeneous (management units) units and showing their extent and geographic distribution on the microwatershed cadastral map. The detailed soil survey at 1:7920 scale was carried out in 575 ha area. The methodology followed for carrying out land resource inventory was as per the guidelines given in Soil Survey Manual (IARI, 1971; Soil Survey Staff, 2006; Natarajan *et al.*, 2015) which is briefly described below.

3.1 Base Maps

The detailed survey of the land resources occurring in the microwatershed was carried out by using digitized cadastral map and satellite imagery as base supplied by the KRSRAC. The cadastral map shows field boundaries with their survey numbers, location of tanks, streams and other permanent features of the area (Fig. 3.1). Apart from the cadastral map, remote sensing data products from Cartosat-1 and LISS IV merged at the scale of 1:7920 were used in conjunction with the cadastral map to identify the geology, landscapes, landforms and other surface features. The imagery helped in the identification and delineation of boundaries between hills, uplands and lowlands, water bodies, forest and vegetated areas, roads, habitations and other cultural features of the area (Fig. 3.2). The cadastral map was overlaid on the satellite imagery (Fig. 3.3) that helps to identify the parcel boundaries and other permanent features. Apart from cadastral maps and images, toposheets of the area (1:50,000 scale) were used for initial traversing, identification of geology, landscapes and landforms, drainage features, present land use and also for selection of transects in the microwatershed.

3.2 Image Interpretation for Physiography

False Colour Composites (FCC) of Cartosat-I and LISS-IV merged satellite data covering the microwatershed area was visually interpreted using image interpretation elements and all the available collateral data with local knowledge. The delineated physiographic boundaries were transferred on to a cadastral map overlaid on satellite imagery. Physiographically, the area has been identified as granite gneiss and alluvial landscapes and is divided into landforms such as ridges, mounds and uplands based on slope. They were further subdivided into physiographic/ image interpretation units based on image characteristics. The image interpretation legend for physiography is given below.

Image Interpretation Legend for Physiography

G- Granite gneiss landscape

G1	Hills/ Ridges/ Mounds
G11	Summits
G12	Side slopes
G121	Side slopes with dark grey tones
G2	Uplands
G21	Summits
G22	Gently sloping uplands
G221	Gently sloping uplands, yellowish green (eroded)
G222	Gently sloping uplands, yellowish white (severely eroded)
G23	Very gently sloping uplands
G231	Very gently sloping uplands, yellowish green
G232	Very gently sloping uplands, medium green and pink
G233	Very gently sloping uplands, pink and green (scrub land)
G234	Very gently sloping uplands, medium greenish grey
G235	Very gently sloping uplands, yellowish white (eroded)
G236	Very gently sloping uplands, dark green
G237	Very gently sloping uplands, medium pink (coconut garden)
G238	Very gently sloping uplands, pink and bluish white (eroded)

DSe Alluvial landscape

Dse 1 Summit

- Dse 11 Nearly level Summit with dark grey tone
- Dse 12 Nearly level Summit with medium grey tone
- Dse 13 Nearly level Summit with whitish grey tone
- Dse 14 Nearly level Summit with whitish tone (Calcareousness)
- Dse 15 Nearly level Summit with pinkish grey tone
- Dse 16 Nearly level Summit with medium pink tone
- Dse 17 Nearly level Summit with bluish white tone
- Dse 18 Nearly level Summit with greenish grey tone

Dse 2 Very gently sloping

- Dse 21 Very gently sloping, whitish tone
- Dse 22 Very gently sloping, greyish pink tone
- Dse 23 Very gently sloping, whitish grey tone
- Dse 24 Very gently sloping, medium grey tone
- Dse 25 Very gently sloping, medium pink tone
- Dse 26 Very gently sloping, dark grey tone
- Dse 27 Very gently sloping, bluish grey tone
- Dse 28 Very gently sloping, greenish grey tone
- Dse 29 Very gently sloping, Pinkish grey

Dsa 25 – Nearly Level Lands

- Dsa 251- Nearly level, Grayish green tone
- Dsa 252- Nearly level, Bluish grey tone
- Dsa 253- Nearly level, Light green tone
- Dsa 254- Nearly level, Medium green tone
- Dsa 255- Nearly level, Greenish pink tone
- Dsa 256- Nearly level, Whitish green
- Dsa 257- Nearly level, Pink tone
- Dsa 258- Nearly level, Whitish grey tone
- Dsa 259- Nearly level, Grayish Pink




Fig. 3.1 Scanned and Digitized Cadastral map of Kesalapura-1 Microwatershed




Fig. 3.2 Satellite Image of Kesalapura-1 Microwatershed




Fig. 3.3 Cadastral map overlaid on IRS PAN+LISS IV merged imagery of Kesalapura-1 Microwatershed

3.3 Field Investigation

The field boundaries and survey numbers given on the cadastral sheet were located on the ground by following permanent features like roads, cart tracks, *nallas*, streams, tanks etc., and wherever changes were noticed, they were incorporated on the microwatershed cadastral map. Preliminary traverse of the microwatershed was carried out with the help of cadastral map, imagery and toposheets. While traversing, landforms and physiographic units identified were checked and preliminary soil legend was prepared by studying soils at few selected places. Then, intensive traversing of each physiographic unit like hills, ridges, uplands and plains was carried out. Based on the variability observed on the surface, transects (Fig. 3.4) were selected across the slope covering all the landform units in the microwatershed (Natarajan and Dipak Sarkar, 2010).




Fig. 3.4 Location of profiles in a transect

In the selected transect, soil profiles (Fig. 3.4) were located at closely spaced intervals to take care of any change in the land features like break in slope, erosion, gravel, stones etc. In the selected sites, soil profiles (vertical cut showing the soil layers from surface to the rock) were opened upto 200 cm or to the depth limited by rock or hard substratum and studied in detail for all their morphological and physical characteristics. The soil and site characteristics were recorded for all profile sites on a standard proforma as per the guidelines given in USDA Soil Survey Manual (Soil Survey Staff, 2012). Apart from the transect study, profiles were also studied at random, almost like in a grid pattern, outside the transect areas to validate the soil map unit boundaries.

Based on the soil characteristics, the soils were grouped into different soil series. Soil series is the most homogeneous unit having similar horizons and properties and behaves similarly for a given level of management. Soil depth, texture, colour, kind of horizon and horizon sequence, amount and nature of gravel present, calcareousness, nature of substratum etc, were used as the major differentiating characteristics for identifying soil series occurring in the area. The differentiating characteristics used for identifying the soil series are given in Table 3.1. Based on the above characteristics, 12 soil series were identified in Kesalapura-1 Microwatershed.

**Table 3.1 Differentiating Characteristics used for identifying Soil Series
(Characteristics are of Series Control Section)**

Soils of Granite gneiss Landscape							
Sl. No.	Soil Series	Depth (cm)	Colour (moist)	Texture	Gravel (%)	Horizon sequence	Calcareousness
1	Mukhadahalli (MKH)	50-75	5YR3/3,3/4,4/3, 5/4,6/6 2.5YR3/4	gsc	>35	Ap-Bt-Cr	
2	Lakkur	50-75	2.5YR 2.5/3, 2.5/4,	gsc	40-60	Ap-Bt-Bc-Cr	

	(LKR)		3/4, 3/6				
3	Hooradhahalli (HDH)	75-100	2.5YR2.5/4,3/4, 3/6	gsc-gc	>35	Ap-Bt-Cr	
4	Bisarahalli (BSR)	75-100	5 YR 3/3, 3/4	gsc	15-35	Ap-Bt-Cr	
5	Chikkamegheri (CKM)	75-100	2.5YR2.5/3,3/4, 3/6	sc	-	Ap-Bt-Cr	
6	Huliyapura (HLP)	75-100	7.5YR3/3,4/6 10YR4/6	scl	-	Ap-Bw-C	-
7	Kumchahalli (KMH)	100-150	2.5YR3/4, 3/6	sc	<15	Bt-Cr	
8	Balapur (BPR)	100-150	2.5YR2.5/4,3/4	gsc-gc	>35	Ap-Bt-Cr	-
9	Giddadapalya (GDP)	100-150	2.5YR3/4, 3/6	gsc-gc	30-60 after 60 cm	Ap-Bt-Cr	-
10	Ranatur (RTR)	>150	2.5YR2.5/3,2.5/4, 3/3,4/6	c	-	Ap-Bt	
11	Kavalakkeri (KLR)	>150	10YR2/1,3/1, 3/2 7.5YR2.5/1,3/2	sc	-	Ap-Bw	e-es
Soils of Alluvial Landscape							
12	Budagumpa (BGP)	>150	7.5YR3/2,5/1 10YR4/1,4/4	c	<15	Ap-Bw	es

3.4 Soil Mapping

The area under each soil series was further separated into 18 soil phases and their boundaries delineated on the cadastral map based on the variations observed in the texture of the surface soil, slope, erosion, presence of gravel, stoniness etc. A soil phase is a subdivision of soil series based mostly on surface features that affect its use and management.

The soil mapping units are shown on the soil map (Fig. 3.5) in the form of symbols. During the survey many soil profile pits, few minipits and a few auger bores representing different landforms occurring in the microwatershed were studied. In addition to the profile study, spot observations in the form of minipits, road cuts, terrace cuts etc., were studied to validate the soil boundaries on the soil map. The soil map shows the geographic distribution of 18 mapping units representing 12 soil series occurring in the microwatershed. The soil map unit (soil legend) description is presented in Table 3.2. The soil phase map (management units) shows the distribution of 18 soil phases mapped in the microwatershed. Each mapping unit (soil phase) delineated on the map has similar soil and site characteristics. In other words, all the farms or survey numbers included in one phase will have similar management needs and have to be treated accordingly.

3.5 Laboratory Characterization

Soil samples for each soil series were collected from representative master profiles for laboratory characterization by following the methods outlined in the

Laboratory Manual (Sarma *et al.*, 1987). Surface soil samples collected in the year 2018 from Kesalapura-1 farmer's fields for fertility status (major and micronutrients) at 320 m grid interval were analyzed in the laboratory (Katyala and Rattan, 2003). By linking the soil fertility data to the survey numbers through GIS, soil fertility maps were generated using Kriging method for the microwatershed.

Table 3.2 Soil map unit description of Kesalapura-1 Microwatershed

Soil map unit No*	Soil Series	Soil Phase Symbol	Mapping Unit Description	Area in ha (%)
Soils of Granite and Granite gneiss landscape				
	MKH		Mukhadahalli soils are moderately shallow (50-75 cm), well drained, have dark brown to reddish brown gravelly sandy clay soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation	32 (10.74)
75		MKHcB1g1	Sandy loam surface, slope 1-3%, slight erosion, gravelly (15-35%)	44 (7.63)
77		MKHcB2g1	Sandy loam surface, slope 1-3%, moderate erosion, gravelly (15-35%)	18 (3.11)
	LKR		Lakkur soils are moderately shallow (50-75cm), well drained, have reddish brown to dark red gravelly sandy clay red soils. They have developed from granite gneiss and occur on nearly level to very gently and gently sloping uplands under cultivation	26 (4.58)
452		LKRhB2g1	Sandy clay loam surface, slope 1-3%, moderate erosion, gravelly (15-35%)	26 (4.58)
	HDH		Hooradhahalli soils are moderately deep (75-100 cm), well drained, have red to dark red and reddish brown gravelly sandy clay to clay soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation	47 (8.23)
123		HDHhB2g1	Sandy clay loam surface, slope 1-3%, moderate erosion, gravelly (15-35%)	47 (8.23)
	BSR		Bisarahalli soils are moderately deep (75-100 cm), well drained, have dark reddish brown gravelly sandy clay red soils. They have developed from weathered granite gneiss and occur on very gently sloping uplands under cultivation	52 (9.0)
162		BSRhB2g1	Sandy clay loam surface, slope 1-3%, moderate erosion, gravelly (15-35%)	37 (6.39)
167		BSRiB2	Sandy clay surface, slope 1-3%, moderate erosion	15 (2.61)
	CKM		Chikkamegheri soils are moderately deep (75-100 cm), well drained, have dark brown to dark reddish brown and red sandy clay soils. They have developed from weathered granite gneiss and occur on nearly level to	140 (24.19)

Soil map unit No*	Soil Series	Soil Phase Symbol	Mapping Unit Description	Area in ha (%)
			very gently sloping uplands	
173		CKMcB2g2	Sandy loam surface, slope 1-3%, moderate erosion, very gravelly (35-60%)	48 (8.31)
174		CKMhB1g1	Sandy clay loam surface, slope 1-3%, slight erosion, gravelly (15-35%)	47 (8.13)
178		CKMiB1	Sandy clay surface, slope 1-3%, slight erosion	45 (7.75)
	HLP		Huliyapura soils are moderately deep (75-100 cm), well drained, have dark- strong brown to dark yellowish brown sandy clay loam soils. They have developed from weathered granite gneiss and occur on very gently sloping low lands under cultivation	9 (1.6)
436		HLPcB2g1	Sandy loam surface, slope 1-3%, moderate erosion, gravelly (15-35%)	9 (1.6)
	KMH		Kumchahalli soils are deep (100-150cm), well drained, have dark reddish brown to dark red sandy clay soils. They have developed from granite gneiss and occur on nearly level to very gently sloping uplands under cultivation	62 (10.67)
198		KMHhB1g1	Sandy clay loam surface, slope 1-3%, slight erosion, gravelly (15-35%)	25 (4.31)
201		KMHhB2	Sandy clay surface, slope 1-3%, moderate erosion	25 (4.31)
	BPR		Balapur soils are deep (100-150 cm), well drained, have dark reddish brown to dark red gravelly sandy clay to clay soils. These soils are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation	12 (2.16)
232		BPRhB2g2	Sandy clay loam surface, slope 1-3%, moderate erosion, very gravelly (35-60%)	12 (2.16)
	GDP		Giddadapalya soils are deep (100-150 cm), well drained, have dark reddish brown to dark red gravelly sandy clay to clay soils. They are developed from weathered granite gneiss and occur on very gently sloping uplands under cultivation	48 (8.41)
269		GDPiB2	Sandy clay surface, slope 1-3%, moderate erosion	48 (8.41)
	RTR		Ranatur soils are very deep (> 150 cm), well drained, have dark reddish brown to dark red clayey soils. They have developed from weathered granite gneiss and occur on very gently sloping uplands under cultivation	38 (6.58)
288		RTRiB2	Sandy clay surface, slope 1-3%, moderate erosion	38 (6.58)
	KLR		Kavalakkeri soils are very deep (>150 cm), moderately well drained, black to very dark brown calcareous	10 (1.71)

Soil map unit No*	Soil Series	Soil Phase Symbol	Mapping Unit Description	Area in ha (%)
			cracking sandy clay soils. They have developed from weathered granite gneiss and occur on nearly level to very gently sloping lowlands under cultivation	
473		KLRmA1	Clay surface, slope 0-1%, slight erosion	4 (0.67)
476		KLRhB2	Sandy loam surface, slope 1-3%, moderate erosion	6 (1.04)
Soils of Alluvial Landscape				
	BGP		Budagumpa soils are very deep (>150 cm), well drained, black calcareous sodic clay soils. They have developed from alluvium and occur on very gently sloping plains under cultivation	48 (8.28)
396		BGPmB1	Clay surface, slope 1-3%, slight erosion	48 (8.28)
1000	Others		Habitation and waterbody	22 (3.86)

*Soil map unit numbers are continuous for the taluk, not the microwatersheds

3.6 Land Management Units (LMU's)

The 18 soil phases identified and mapped in the microwatershed were regrouped into 5 Land Management Units (LMU's) for the purpose of preparing a Proposed Crop Plan for sustained development of the microwatershed. The database (soil phases) generated under LRI was utilized for identifying Land Management Units (LMU's) based on the management needs. One or more than one soil site characteristic having influence on the management have been chosen for identification and delineation of LMUs. For Kesalapura-1 Microwatershed, five soil and site characteristics, namely soil depth, soil texture, slope, erosion and gravel content have been considered for defining LMUs. The Land Management Units are expected to behave similarly for a given level of management.




Fig 3.4 Soil Phase or Management Units-Kesalapura-1 Microwatershed

THE SOILS

Detailed information pertaining to the nature, extent and distribution of different kinds of soils occurring in Kesalapura-1 Microwatershed is provided in this chapter. The microwatershed area has been identified as granite gneiss and alluvial landscapes based on geology. In all, 12 soil series are identified. Soil formation is the result of the combined effect of environmental and terrain factors that are reflected in soil morphology. The soil formation is dominantly influenced by the parent material, climate, time and relief.

A brief description of each of the 12 soil series identified followed by 18 soil phases (management units) mapped (Fig. 3.4) are furnished below. The physical and chemical characteristics of soil series identified in Kesalapura-1 Microwatershed are given in Table 4.1. The soils in any one map unit differ from place to place in their depth, texture, slope, gravelliness, erosion or any other site characteristic that affect management. The soil phase map can be used for identifying the suitability of areas for growing specific crops or for other alternative uses and also for deciding the type of conservation structures needed. The detailed information on soil and site-characteristics like soil depth, surface soil texture, slope, erosion, gravelliness, AWC, LCC etc, with respect to each of the soil phase identified is given village/survey number wise for the microwatershed in Appendix-I.

4.1 Soils of granite and granite gneiss landscape

In this landscape, 11 soil series are identified and mapped. Of these, Chikkamegheri (CKM) series occupies maximum area of 140 (24%), Kumchahalli (KMH) 62 ha (11%), Bisarahalli (BSR) 52 ha (9%), Giddadapalya (GDP) 48 ha (8%), Hooradhahalli (HDH) 47 ha (8%), Ranatur (RTR) 38 ha (7%), Mukhadahalli (MKH) 32 ha (11%), Lakkur (LKR) 26 ha (5%), Balapur (BPR) 12 ha (2%), Kavalakkeri (KLR) 10 ha (2%), and Huliypura (HLP) occupy an area of about 9 ha (2%) in the microwatershed. The brief description of each soil series along with the soil phases identified and mapped is given below.

4.1.1 Mukhadahalli (MKH) Series: Mukhadahalli soils are moderately shallow (50-75 cm), well drained, have dark brown to reddish brown, gravelly sandy clay soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands. The Mukhadahalli series has been classified as a member of the clayey-skeletal, mixed, isohyperthermic family of Typic Haplustalfs.

The thickness of the solum ranges from 51 to 72 cm. The thickness of A horizon ranges from 12 to 17 cm. Its colour is in 5 YR and 7.5 YR hue with value 3 to 4 and chroma 2 to 4. The texture varies from loamy sand to sandy loam with 20 to 45 per cent gravel. The thickness of B horizon ranges from 40 to 68 cm. Its colour is in 2.5 YR and 5

YR hue with value and chroma 3 to 6. Texture is sandy clay loam to sandy clay with 35 to 50 per cent gravel. The available water capacity is low (50-100 mm/m). Two phases were identified and mapped.



Landscape and soil profile characteristics of Mukhadahalli (MKH) Series

4.1.2 Lakkur (LKR) Series: Lakkur soils are moderately shallow (50-75cm), well drained, have reddish brown to dark red gravelly sandy clay red soils. They have developed from weathered granite gneiss and occur on nearly level to very gently and gently sloping uplands. The Lakkur series has been classified as a member of the clayey-skeletal, mixed, isohyperthermic family of Typic Rhodustalfs.

The thickness of the solum ranges from 51 to 74 cm. The thickness of A horizon ranges from 12 to 18 cm. Its colour is in 5YR and 2.5 YR hue with value 3 to 4 and chroma 4 to 6. The texture varies from loamy sand to sandy clay loam with 15 to 50 per cent gravel. The thickness of B horizon ranges from 39 to 58 cm. Its colour is in 2.5 YR hue with value 3 to 4 and chroma 4 to 6. Texture is sandy clay with 40 to 60 per cent gravel. The available water capacity is low (50-100 mm/m). Only one phase was identified and mapped.



Landscape and soil profile characteristics of Lakkur (LKR) Series

4.1.3 Hooradhahalli (HDH) Series: Hooradhahalli soils are moderately deep (75-100 cm), well drained, have red to dark red and reddish brown gravelly sandy clay to clay soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands. The Hooradhahalli series has been classified as a member of the clayey-skeletal, mixed, isohyperthermic family of Rhodic Paleustalfs.

The thickness of the solum ranges from 76 to 100 cm. The thickness of A horizon ranges from 11 to 19 cm. Its colour is in 5 YR and 2.5 YR hue with value 3 to 4 and chroma 3 to 6. The texture varies from loamy sand to sandy clay with 15 to 50 per cent gravel. The thickness of B horizon varies from 65 to 83 cm. Its colour is in 2.5 YR hue with value 2.5 to 3 and chroma 4 to 6. Texture is sandy clay to clay with 35 to 50 per cent gravel. The available water capacity is low (50-100 mm/m). Only one phase was identified and mapped.



Landscape and soil profile characteristics of Hooradhahalli (HDH) Series

4.1.4 Bisarahalli (BSR) Series: Bisarahalli soils are moderately deep (75-100 cm), well drained, have dark reddish brown gravelly sandy clay red soils. They have developed from weathered granite gneiss and occur on very gently sloping uplands under cultivation. The Bisarahalli series has been classified as a member of the fine, mixed isohyperthermic family of Typic Paleustalfs.

The thickness of the solum ranges from 75 to 98 cm. The thickness of A horizon ranges from 17 to 25 cm. Its colour is in 5 YR hue with value 3 to 4 and chroma 3 to 6. The texture ranges from sandy clay loam to sandy clay with 15 to 35 per cent gravel. The thickness of B horizon ranges from 61 to 79 cm. Its colour is in 5 YR hue with value 3 and chroma 3 to 4. Its texture is gravelly sandy clay with gravel content of 15-35 per cent. The available water capacity is low (50-100 mm/m). Two phases were identified and mapped.



Landscape and soil profile characteristics of Bisarahalli (BSR) Series

4.1.5 Chikkamegheri (CKM) Series: Chikkamegheri soils are moderately deep (75-100 cm), well drained, have dark brown to dark reddish brown and red sandy clay soils. They have developed from weathered granite gneiss and occur on nearly level to very gently sloping uplands. The Chikkamegheri series has been classified as a member of the fine, mixed, isohyperthermic family of Rhodic Paleustalfs.

The thickness of the solum ranges from 76 to 100 cm. The thickness of A horizon ranges from 11 to 24 cm. Its colour is in 7.5 YR, 5YR and 2.5 YR hue with value 2 to 4 and chroma 3 to 6. The texture varies from sandy clay loam to sandy clay with 10 to 15 per cent gravel. The thickness of B horizon ranges from 65 to 86 cm. Its colour is in 2.5 YR hue with value 2.5 to 3 and chroma 3 to 6. Its texture is dominantly sandy clay to clay. The available water capacity is medium (100-150 mm/m). Three phases were identified and mapped.



Landscape and soil profile characteristics of Chikkamegheri (CKM) Series

4.1.6 Huliyapura (HLP) Series: Huliyapura soils are moderately deep (75-100 cm), well drained, have dark- strong brown to dark yellowish brown sandy clay loam soils. They have developed from weathered granite gneiss and occur on very gently sloping low lands under cultivation. The Huliyapura series has been classified as a member of fine-loamy, mixed, isohyperthermic family of Typic Haplustepts.

The thickness of the solum ranges from 75 to 98 cm. The thickness of A-horizon ranges from 18 to 22 cm. Its colour is in 5 YR and 10 YR hue with value 3 to 4 and chroma 4. The texture is sandy clay loam. The thickness of B-horizon ranges from 56 to 75 cm. Its colour is in 5 YR, 7.5 YR and 10 YR hue with value 3 to 4 and chroma 2 to 6. Its texture is sandy clay. The available water capacity is low (50-100 mm/m). Only one phase was identified and mapped.



Landscape and soil profile characteristics of Huliyapura (HLP) Series.

4.1.7 Kumchahalli (KMH) Series: Kumchahalli soils are deep (100-150 cm), well drained, have dark reddish brown to dark red sandy clay soils. They have developed from weathered granite gneiss and occur on nearly level to very gently sloping uplands. The Kumchahalli series has been classified as a member of the fine, mixed, isohyperthermic family of Typic Rhodustalfs.

The thickness of the solum ranges from 102 to 150 cm. The thickness of A horizon ranges from 11 to 23 cm. Its colour is in 5 YR and 2.5 YR hue with value 2.5 to 3 and chroma 3 to 6. The texture is dominantly sandy clay. The thickness of B horizon ranges from 95 to 132 cm. Its colour is in 2.5 YR hue with value 3 and chroma 4 to 6. Its texture is dominantly sandy clay loam to sandy clay. The available water capacity is high (150-200 mm/m). Two phases were identified and mapped.



Landscape and soil profile characteristics of Kumchahalli (KMH) Series

4.1.8 Balapur (BPR) Series: Balapur soils are deep (100-150 cm), well drained, have dark reddish brown to dark red gravelly sandy clay to clay soils. These soils are developed from weathered granite gneiss and occur on very gently to gently sloping uplands. The Balapur series has been classified as a member of the clayey-skeletal, mixed, isohyperthermic family of Typic Rhodustalfs.

The thickness of the solum ranges from 102 to 147 cm. The thickness of A horizon ranges from 12 to 17 cm. Its colour is in 5 YR and 2.5 YR hue with value and chroma 3 to 4. The texture ranges from loamy sand to sandy clay with 15 to 50 per cent gravel. The thickness of B horizon ranges from 90 to 132 cm. Its colour is in 2.5 YR hue with value 2.5 to 3 and chroma 4 to 6. Texture is sandy clay to clay with 35 to 50 per cent gravel. The available water capacity is medium (100-150 mm/m). Only one phase was identified and mapped.



Landscape and soil profile characteristics of Balapur (BPR) Series

4.1.9 Giddadapalya (GDP) Series: Giddadapalya soils are deep (100-150 cm), well drained, have dark reddish brown to dark red gravelly sandy clay to clay soils. They are developed from weathered granite gneiss and occur on very gently sloping uplands under cultivation. The Giddadapalya series has been classified as a member of the fine, mixed, isohyperthermic family of Rhodic Paleustalfs.

The thickness of the solum ranges from 106 to 145 cm. The thickness of A-horizon ranges from 12 to 13 cm. Its colour is in 5 YR hue with value and chroma 3 to 4. The texture ranges from sandy loam with 10 to 15 per cent gravel. The thickness of B-horizon ranges from 106 to 123 cm. Its colour is in 2.5 YR hue with value 3 to 4 and chroma 3 to 6. Texture is sandy clay to clay with 35 to 75 per cent gravel. The available water capacity is low (51-100 mm/m). Only one phase was identified and mapped.



Landscape and soil profile characteristics of Giddadapalya (GDP) Series.

4.1.10 Ranatur (RTR) Series: Ranatur soils are very deep (> 150 cm), well drained, have dark reddish brown to dark red clayey soils. They have developed from weathered granite gneiss and occur on very gently sloping uplands. The Ranatur series has been classified as a member of the fine, mixed, isohyperthermic family of Rhodic Paleustalfs.

The thickness of the solum is more than 150 cm. The thickness of A horizon ranges from 8 to 14 cm. Its colour is in 5 YR and 2.5 YR hue with value 2.5 to 4 and chroma 3 to 6. The texture varies from sandy loam to sand clay. The thickness of B horizon is more than 150 cm. Its colour is in 2.5 YR hue with value 2.5 to 3 and chroma 3 to 6. Its texture is clay. The available water capacity is high (150-200 mm/m). Only one phase was identified and mapped.



Landscape and soil profile characteristics of Ranatur (RTR) Series

4.1.11 Kavalakkeri (KLR) Series: Kavalakkeri soils are very deep (>150 cm), moderately well drained, black to very dark brown calcareous cracking sandy clay soils. They have developed from granite gneiss and occur on nearly level to very gently sloping lowlands under cultivation. The Kavalakkeri series has been classified as a member of the fine, mixed, (calc) isohyperthermic family of Fluventic Haplustepts.

The thickness of the solum is more than 150 cm. The thickness of A horizon ranges from 18 to 29 cm. Its colour is in 7.5 and 10YR hue with value 3 to 4 and chroma 2 to 4. The texture is sandy clay. The thickness of B horizon ranges from 131-155 cm. Its colour is in 7.5YR and 10 YR hue with value 2 to 4 and chroma 1 to 4. Its texture is sandy clay to clay. The available water capacity is very high (>200 mm/). Two phases were identified and mapped.



Landscape and soil profile characteristics of Kavalakkeri (KLR) Series

4.2 Soils of Alluvial landscape

In this landscape, only one soil series has been identified and mapped. Budagumpa (BGP) series occupies an area of 48 ha (8%) in the microwatershed. The brief description of soil series along with the soil phases identified and mapped is given below.

4.2.1 Budagumpa (BGP) Series: Budagumpa soils are very deep (>150 cm), well drained, black calcareous sodic clay soils. They have developed from alluvium and occur on very gently sloping plains under cultivation. The Budagumpa series has been classified as a member of the fine, mixed, (calc) isohyperthermic family of Typic Haplustepts.

The thickness of the solum is more than 150 cm. The thickness of A horizon ranges from 16 to 26 cm. Its colour is in 7.5 YR and 10 YR hue with value 2 to 3 and chroma 2 to 4. The texture varies from sandy clay to clay with 5 to 10 per cent gravel. The thickness of B horizon ranges from 130 to 160 cm. Its colour is in 10 YR and 7.5 YR hue with value 3 to 5 and chroma 1 to 4. Its texture is clay with gravel content of <15 per cent. These soils are calcareous that increase with depth. The available water capacity is very high (>200 mm/m). Only one phase was identified and mapped.



Landscape and soil Profile Characteristics of Budagumpa (BGP) Series

Table: 4.1 Physical and Chemical Characteristics of Soil Series identified in Kesalapura-1 Microwatershed

Series Name: Mukahadahalli (MKH), **Pedon:** R-11

Location: 15°22'05.4"N, 76°04'10.3"E, Halageri village, Koppal Taluk and District

Analysis at: NBSS&LUP, Regional Centre, Bangalore. **Classification:** Clayey-skeletal, mixed, isohyperthermic Typic Haplustalfs

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-19	Ap	65.71	8.83	25.46	9.27	9.06	14.42	21.52	11.43	70	scl	16.54	8.60
19-32	Bt	55.89	11.13	32.98	6.47	9.18	11.89	19.19	9.18	50	scl	19.24	12.78
32-58	Bt	47.95	10.41	41.63	17.52	3.78	9.13	9.55	7.97	50	sc	24.03	16.02

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO ₃	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP
	Water	CaCl ₂	M KCl				Ca	Mg	K	Na	Total				
				dS m ⁻¹	%	%	cmol kg ⁻¹					%	%		
0-19	7.38	-	-	0.09	0.2	0.00	8.97	4.32	0.26	0.22	13.77	14.84	0.58	93	1.49
19-32	7.5	-	-	0.106	0.41	0.00	15.98	3.27	0.16	0.50	19.91	20.88	0.63	95	2.38
32-58	7.46	-	-	0.173	0.49	0.00	19.71	4.53	0.23	1.32	25.79	25.76	0.62	100	5.11

Contd...

Soil Series: Lakkur (LKR), **Pedon:** RM-8.

Location: 15°04'26.3"N, 75°37'84.1"E, (4D4A3I1f), Belhatti village, Shirahatti taluk, Gadag district

Analysis at: NBSS&LUP, Regional Centre, Bengaluru **Classification:** Clayey-skeletal, mixed, isohyperthermic Typic Rhodustalfs

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-21	Ap	74.00	8.34	17.66	9.62	11.57	15.76	23.13	13.92	20	sl	-	-
21-35	Bt	54.37	10.48	35.14	16.33	8.64	9.69	11.59	8.11	40	sc	-	-
35-56	Bc	48.37	13.46	38.17	10.96	7.69	9.17	11.28	9.27	60	sc	-	-

Depth (cm)	pH (1:2.5)			E.C. (1:2.5) dS m ⁻¹	O.C. %	CaCO ₃ %	Exchangeable bases					CEC	CEC/Clay	Base saturation %	ESP %
	Water	CaCl ₂	M KCl				Ca	Mg	K	Na	Total				
0-21	8.18	-	-	0.30	0.56	0.94	-	-	0.31	0.55	0.86	12.19	0.69	100.00	4.51
21-35	8.17	-	-	0.30	0.52	1.29	-	-	0.19	0.84	1.03	22.18	0.63	100.00	3.79
35-56	7.95	-	-	0.46	0.48	1.99	-	-	0.24	0.58	0.82	22.94	0.60	100.00	2.53

Contd...

Soil Series: Hooradhahalli (HDH), **Pedon:** RM-69

Location: 13°24'31"N, 76°33'41"E, (4D3D8G2d), Hesarahalli village, Chikkanayakanahalli taluk, Tumukura district

Analysis at: NBSS&LUP, Regional Centre, Bengaluru **Classification:** Clayey-skeletal, mixed, isohyperthermic Rhodic Paleustalfs

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-18	Ap	72.56	15.17	12.27	4.57	8.33	17.38	23.88	18.39	35	sl	-	-
18-33	Bt1	56.29	10.75	32.96	7.88	10.24	13.41	14.43	10.34	55	scl	-	-
33-58	Bt2	46.66	10.79	42.55	10.79	9.87	8.43	9.04	8.53	55	sc	-	-
58-90	Bt3	43.09	13.63	43.27	9.90	8.25	7.32	8.76	8.87	45	c	-	-

Depth (cm)	pH (1:2.5)			E.C. (1:2.5) dS m ⁻¹	O.C. %	CaCO ₃ %	Exchangeable bases					CEC	CEC/Clay	Base saturation %	ESP %
	Water	CaCl ₂	M KCl				Ca	Mg	K	Na	Total				
0-18	6.54	-	-	0.07	0.60	0.00	2.68	1.38	0.44	0.42	4.91	5.84	0.48	84.07	7.11
18-33	5.90	-	-	0.07	0.52	0.00	3.99	1.27	0.09	0.37	5.71	8.61	0.26	66.32	4.29
33-58	6.16	-	-	0.07	0.44	0.00	4.92	1.67	0.08	0.55	7.22	10.00	0.24	72.23	5.50
58-90	6.39	-	-	0.06	0.40	0.00	4.30	2.02	0.08	0.46	6.87	9.21	0.21	74.61	5.05

Contd...

Series Name: Bissarahalli (BSR) **Pedon:** R-9

Location: 15°25'21.0"N, 76°11'42.0"E Hatti village, Koppal Taluk and District

Analysis at: NBSS&LUP, Regional Centre, Bangalore. **Classification:** Fine, mixed, isohyperthermic Typic Paleustalfs

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-14	Ap	70.11	9.29	20.60	22.31	15.97	11.98	9.83	10.03	20	scl	13.22	7.81
14-57	Bt1	47.27	7.52	45.20	27.04	8.28	4.61	2.10	5.24	25	sc	16.39	13.31
57-80	Bt2	41.93	8.67	49.40	21.95	6.83	4.76	4.66	3.73	30	c	21.41	15.41
80-99	Bt3	49.02	9.87	41.11	19.90	10.78	6.84	6.42	5.08	40	sc	21.82	14.24

Depth (cm)	pH (1:2.5)			E.C. (1:2.5) dS m ⁻¹	O.C. %	CaCO ₃ %	Exchangeable bases					CEC	CEC/Clay	Base saturation %	ESP %
	Water	CaCl ₂	M KCl				Ca	Mg	K	Na	Total				
0-14	6.59	-	-	0.12	0.73	-	4.47	1.77	0.06	0.53	6.82	8.80	0.43	77.55	6.00
14-57	7.02	-	-	0.04	0.48	-	5.85	2.31	0.06	0.20	8.43	14.70	0.33	57.32	1.36
57-80	7.00	-	-	0.05	0.28	-	11.74	2.26	0.08	0.22	14.31	15.60	0.32	91.73	1.44
80-99	6.90	-	-	0.06	0.18	-	13.70	2.16	0.08	0.14	16.08	16.50	0.40	97.44	0.83

Contd...

Series Name: Chikkamegheri (CKM), **Pedon:** RM-2

Location: 15°21'40"N, 76°16'43"E, Gudanhalli village, Koppal Taluk and District

Analysis at: NBSS&LUP, Regional Centre, Bangalore. **Classification:** Fine, mixed, isohyperthermic Rhodic Paleustalfs

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-10	Ap	66.80	5.51	27.69	10.14	10.04	20.29	14.75	11.58	-	scl	20.59	7.15
10-25	Bt1	39.52	7.17	53.32	8.75	9.59	7.27	8.43	5.48	-	c	26.96	13.99
25-38	Bt2	42.00	7.16	50.84	13.16	8.74	6.42	8.53	5.16	-	c	26.51	13.42
38-55	Bt3	41.77	10.31	47.92	15.19	8.54	6.33	7.38	4.32	10	c	25.28	14.10
55-70	Bt4	44.03	8.96	47.01	15.72	9.22	6.92	6.81	5.35	20	c	24.30	14.35
70-90	Bt5	56.02	8.46	35.52	11.41	17.07	12.36	10.26	4.92	25	sc	20.59	13.06

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO ₃	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP
	Water	CaCl ₂	M KCl				Ca	Mg	K	Na	Total				
				dS m ⁻¹	%	%	cmol kg ⁻¹					%	%		
0-10	7.99	-	-	0.326	0.83	4.44	9.35	4.76	0.28	0.54	14.93	12.50	0.45	119	1.73
10-25	7.36	-	-	0.345	0.99	2.40	10.37	4.84	0.10	1.18	16.48	17.60	0.33	94	2.67
25-38	6.69	-	-	0.477	0.79	0.00	10.25	4.20	0.09	1.61	16.15	16.10	0.32	100	4.00
38-55	6.45	-	-	0.548	0.63	0.00	9.43	2.86	0.10	1.52	13.91	14.80	0.31	94	4.11
55-70	6.35	-	-	0.532	0.71	0.00	9.59	2.79	0.11	1.66	14.16	14.60	0.31	97	4.56
70-90	6.44	-	-	0.613	0.27	0.00	9.58	3.10	0.19	1.87	14.74	14.70	0.41	100	5.08

Contd...

Series Name: Kumchahalli (KMH), **Pedon:** RM-9

Location: 15°20'05"N, 76°13'21"E, Basapura village, Koppal Taluk and District

Analysis at: NBSS&LUP, Regional Centre, Bangalore. **Classification:** Fine, mixed, isohyperthermic Typic Rhodustalfs

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-13	Ap	51.76	9.05	39.19	7.99	8.84	13.42	14.38	7.14	-	sc	20.08	13.69
13-27	A21	53.50	8.12	38.38	7.00	11.05	15.21	14.33	5.91	-	sc	17.05	12.32
27-43	A22	63.60	5.01	31.40	3.85	11.56	24.52	18.52	5.14	-	scl	11.76	9.09
43-64	Bt1	48.74	5.91	45.35	8.87	9.31	12.49	12.27	5.81	10	sc	16.68	13.35
64-84	Bt2	45.13	8.90	45.97	9.86	7.12	10.95	10.62	6.57	20	sc	17.45	13.42
84-114	BC	65.04	6.94	28.02	10.49	16.21	17.80	13.88	6.67	40	scl	13.20	9.75

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO ₃	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP			
	Water	CaCl ₂	M KCl				dS m ⁻¹	%	%	Ca	Mg					K	Na	Total
0-13	7.2	-	-	0.193	0.81	3.00	9.69	3.93	1.41	0.08	15.10	15.07	0.38	100	0.54			
13-27	7.13	-	-	0.161	0.7	3.00	8.69	3.57	1.29	0.16	13.70	13.75	0.36	100	1.14			
27-43	7.31	-	-	0.096	0.89	2.64	5.19	2.36	1.07	0.24	8.86	9.46	0.30	94	2.51			
43-64	7.65	-	-	0.089	1.16	2.52	8.25	2.88	0.72	0.35	12.20	12.65	0.28	96	2.79			
64-84	7.98	-	-	0.1	0.38	3.12	10.49	2.88	0.26	0.41	14.04	14.63	0.32	96	2.78			
84-114	8.23	-	-	0.121	0.58	2.88	8.02	1.87	0.09	0.43	10.41	10.67	0.38	98	4.02			

Contd...

Soil Series: Balapur (BPR), **Pedon:** RM-78

Location: 13°26'39"N, 76°35'03"E, (4D3D8G2c), Kasaba, Chikkanayakanahalli taluk, Tumakuru district

Analysis at: NBSS&LUP, Regional Centre, Bengaluru **Classification:** Clayey-skeletal, mixed, isohyperthermic Typic Rhodustalfs

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-12	Ap	65.66	18.66	15.68	4.14	6.16	13.33	21.82	20.20	-	sl	-	-
12-34	Bt1	61.91	11.52	26.57	2.36	6.78	12.53	21.36	18.89	-	scl	-	-
34-60	Bt2	51.81	11.24	36.94	4.66	5.70	12.23	15.96	13.26	30	sc	-	-
60-84	Bt3	46.61	9.02	44.37	14.70	6.88	7.51	8.97	8.55	55	sc	-	-
84-112	Bt4	48.75	12.92	38.33	15.73	8.13	6.87	8.23	9.79	60	sc	-	-
112-127	Bc	50.98	24.74	24.28	5.25	4.63	5.15	10.92	25.03	50	scl	-	-

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO ₃	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP
	Water	CaCl ₂	M KCl				Ca	Mg	K	Na	Total				
				dS m ⁻¹	%	%	cmol kg ⁻¹					%	%		
0-12	6.64	-	-	0.03	0.56	0.00	1.90	1.32	0.21	0.03	3.46	5.45	0.35	63.48	0.51
12-34	6.99	-	-	0.02	0.48	0.00	3.66	1.90	0.07	0.08	5.70	7.82	0.29	72.93	0.96
34-60	7.29	-	-	0.02	0.40	0.00	5.13	2.08	0.11	0.20	7.52	11.19	0.30	67.18	1.75
60-84	7.50	-	-	0.02	0.32	0.00	5.83	6.36	0.13	0.23	12.55	12.38	0.28	101.43	1.83
84-112	7.54	-	-	0.02	0.24	0.00	6.02	6.59	0.11	0.25	12.96	12.77	0.33	101.49	1.97
112-127	7.90	-	-	0.02	0.20	0.00	8.04	3.62	0.07	0.32	12.04	12.47	0.51	96.56	2.55

Contd...

Series Name: Giddadapalya (GDP), **Pedon:** R-8

Location: 15°25'26"N, 76°10'59"E, Kalakeri village, Koppal Taluk and District

Analysis at: NBSS&LUP, Regional Centre, Bangalore. **Classification:** Fine, mixed, isohyperthermic Rhodic Paleustalfs

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-16	Ap	74.95	9.24	15.81	18.43	18.94	13.85	14.97	8.76	-	sl	11.88	5.09
16-43	Bt1	41.69	13.89	44.42	9.84	10.90	7.41	7.62	5.93	-	c	23.13	14.53
43-61	Bt2	47.67	6.13	46.19	21.14	10.15	5.29	6.45	4.65	-	sc	21.60	11.87
61-83	Bt3	52.52	7.10	40.38	24.42	10.59	5.66	7.55	4.30	40	sc	19.51	11.35
83-119	Bt4	43.76	11.59	44.65	20.15	7.56	5.77	5.46	4.83	60	c	20.80	12.06
119-139	Bt5	54.93	9.84	35.23	29.70	10.49	5.50	5.92	3.32	50	sc	15.24	11.97

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO ₃	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP			
	Water	CaCl ₂	M KCl				dS m ⁻¹	%	%	Ca	Mg					K	Na	Total
0-16	7.88	-	-	0.103	0.79	-	5.98	1.35	0.05	0.22	7.60	7.8	0.49	97	2.87			
16-43	7.81	-	-	0.117	0.66	-	13.99	1.97	0.08	0.46	16.50	16.9	0.38	98	2.74			
43-61	7.74	-	-	0.132	0.51	-	12.70	2.18	0.08	0.69	15.64	15.9	0.34	98	4.36			
61-83	7.72	-	-	0.142	0.39	-	11.46	2.22	0.08	0.66	14.41	14.6	0.36	99	4.53			
83-119	7.58	-	-	0.115	0.22	-	11.30	2.70	0.09	0.73	14.82	15.3	0.34	97	4.79			
119-139	7.50	-	-	0.113	0.22	-	10.03	2.19	0.07	0.65	12.95	13.2	0.37	98	4.89			

Contd...

Soil Series: Ranatur (RTR), **Pedon:** RM-87

Location: 13°21'49.0"N, 76°38'06"E, (4B3D4L2a), J C Pura village, Chikkanayakanahalli taluk, Tumakuru district

Analysis at: NBSS&LUP, Regional Centre, Bengaluru **Classification:** Fine, mixed, isohyperthermic Rhodic Paleustalfs

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-17	Ap	84.16	9.46	6.38	2.22	18.57	26.14	24.32	12.92	-	ls	-	-
17-47	Bt1	51.14	8.30	40.56	1.66	13.49	14.52	13.59	7.88	-	sc	-	-
47-89	Bt2	51.99	11.01	37.00	1.94	13.99	15.32	13.18	7.56	-	sc	-	-
89-123	Bt3	51.58	9.07	39.35	3.47	14.50	14.61	11.64	7.35	-	sc	-	-
123-152	Bt4	47.89	8.88	43.23	2.27	12.36	14.21	11.12	7.93	-	sc	-	-
152-198	Bt5	43.37	13.17	43.45	2.48	9.83	13.25	10.87	6.94	-	c	-	-

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO ₃	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP
	Water	CaCl ₂	M KCl				Ca	Mg	K	Na	Total				
				dS m ⁻¹	%	%	cmol kg ⁻¹					%	%		
0-17	5.08	-	-	0.03	0.52	0.00	3.68	0.72	0.06	0.19	4.65	9.21	1.44	50.50	2.06
17-47	6.28	-	-	0.03	0.48	0.00	3.93	0.72	0.08	0.07	4.80	7.92	0.20	60.59	0.94
47-89	6.42	-	-	0.03	0.40	0.00	4.40	0.74	0.08	0.06	5.28	7.52	0.20	70.15	0.79
89-123	6.50	-	-	0.02	0.32	0.00	4.44	0.76	0.09	0.07	5.36	7.82	0.20	68.58	0.93
123-152	6.52	-	-	0.02	0.28	0.00	4.40	0.71	0.09	0.07	5.26	8.22	0.19	64.00	0.81
152-198	7.09	-	-	0.02	0.24	0.00	6.10	0.98	0.10	0.20	7.38	9.60	0.22	76.89	2.09

Contd...

Series Name: Kavalakeri (KLR), **Pedon :** R-5

Location: 15°27'55.2"N, 76°15'48.0" E Kenchanadoni village, Koppal Taluk and District

Analysis at: NBSS&LUP, Regional Centre, Bangalore. **Classification:** Fine, mixed, (calc) isohyperthermic Typic Haplustepts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-21	Ap	41.67	28.70	29.62	6.62	10.58	5.70	8.00	10.76	-	cl	22.02	15.06
21-40	Bw1	32.23	29.16	38.61	3.76	4.03	3.04	8.24	13.16	-	cl	26.28	19.49
40-70	Bw2	37.41	26.13	36.46	7.52	6.25	4.62	8.61	10.42	-	cl	26.65	18.87
70-106	Bw3	46.43	18.15	35.42	13.93	14.29	5.98	5.98	6.25	-	sc	22.83	17.66
106-137	Bw4	55.64	12.91	31.45	10.59	8.16	12.67	11.46	12.76	-	scl	24.04	12.85
137-162	Bw5	47.16	16.68	36.16	2.88	4.80	5.68	17.12	16.68	-	sc	30.46	16.24

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO ₃	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP
	Water	CaCl ₂	M KCl				Ca	Mg	K	Na	Total				
				dS m ⁻¹	%	%	cmol kg ⁻¹					%	%		
0-21	7.11	-	-	0.33	0.82	8.84	-	-	0.10	0.67	-	19.50	0.66	100.00	3.42
21-40	7.50	-	-	0.32	0.40	6.63	-	-	0.15	0.99	-	23.20	0.60	100.00	4.26
40-70	7.68	-	-	0.33	0.34	8.19	-	-	0.09	1.18	-	21.90	0.60	100.00	5.38
70-106	7.82	-	-	0.23	0.42	6.50	-	-	0.07	1.36	-	21.80	0.62	100.00	6.23
106-137	7.86	-	-	0.23	0.32	3.57	-	-	0.08	0.95	-	17.30	0.55	100.00	5.47
137-162	7.75	-	-	0.31	0.38	3.90	-	-	0.09	1.01	-	22.10	0.61	100.00	4.55

Contd...

Series Name: Budagumpa (BGP), **Pedon:** R-21

Location: 15°23'45"N, 76°08'52"E Neregalla village, Koppal Taluk and District

Analysis at: NBSS&LUP, Regional Centre, Bangalore. **Classification:** Fine, mixed, (calc) isohyperthermic Typic Haplustepts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-16	Ap	58.30	18.10	23.60	6.34	11.75	11.66	17.44	11.10	-	scl	18.24	10.29
16-38	Bw1	44.26	18.39	37.36	4.71	9.79	9.32	12.24	8.19	-	cl	32.99	18.12
38-68	Bw2	37.84	24.91	37.25	3.66	7.51	8.45	10.89	7.32	-	cl	39.50	22.32
68-83	Bw3	19.17	19.89	60.93	0.87	3.47	3.85	6.07	4.91	-	c	47.27	28.52
83-107	Bw4	14.76	23.22	62.02	0.63	2.41	3.25	4.61	3.87	-	c	46.10	29.36
107-131	Bw5	11.86	17.75	70.39	0.85	2.73	2.45	3.20	2.64	-	c	50.52	28.09
131-160	Bw6	14.48	18.21	67.31	2.23	2.50	2.59	3.84	3.31	-	c	59.14	28.35

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO ₃	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP			
	Water	CaCl ₂	M KCl				dS m ⁻¹	%	%	Ca	Mg					K	Na	Total
										cmol kg ⁻¹								
0-16	9.20	-	-	0.27	0.51	6.24	-	-	0.42	3.11	-	19.60	0.83	100.00	3.84			
16-38	9.29	-	-	0.88	0.35	5.98	-	-	0.17	9.36	-	28.40	0.76	100.00	15.38			
38-68	8.95	-	-	2.37	0.31	4.81	-	-	0.31	24.10	-	34.90	0.94	100.00	42.65			
68-83	8.65	-	-	4.28	0.33	4.42	-	-	0.39	27.95	-	45.10	0.74	100.00	25.94			
83-107	8.10	-	-	9.50	0.30	3.38	-	-	0.44	31.29	-	44.10	0.71	100.00	12.82			
107-131	8.16	-	-	9.32	0.22	2.73	-	-	0.63	37.86	-	47.20	0.67	100.00	20.37			
131-160	8.49	-	-	5.29	0.19	3.51	-	-	0.60	34.82	-	43.70	0.65	100.00	48.66			

INTERPRETATION FOR LAND RESOURCE MANAGEMENT

The most important soil and site characteristics that affect the land use and conservation needs of an area are land capability, land irrigability, soil depth, soil texture, coarse fragments, available water capacity, soil slope, soil erosion, soil reaction etc. These are interpreted from the data base generated through land resource inventory and several thematic maps are generated. These would help in identifying the areas suitable for growing crops and, soil and water conservation measures and structures needed thus helping to maintain good soil health for sustained crop production. The various thematic maps generated are described below.

5.1 Land Capability Classification

Land capability classification is an interpretative grouping of soil map units (soil phases) mainly based on inherent soil characteristics, external land features and environmental factors that limit the use of land for agriculture, pasture, forestry or other uses on a sustained basis (IARI, 1971). The land and soil characteristics used to group the land resources in an area into various land capability classes, subclasses and units are

Soil characteristics: Soil depth, soil texture, coarse fragments, soil reaction, available water capacity, calcareousness, salinity/alkali *etc.*

Land characteristics: Slope, erosion, drainage, rock outcrops.

Climate: Total rainfall and its distribution, and length of crop growing period.

The land capability classification system is divided into land capability classes, subclasses and units based on the level of information available. Eight land capability classes are recognized. They are

Class I: They are very good lands that have no limitations or very few limitations that restrict their use.

Class II: They are good lands that have minor limitations and require moderate conservation practices.

Class III: They are moderately good lands that have severe limitations that reduce the choice of crops or that require special conservation practices.

Class IV: They are fairly good lands that have very severe limitations that reduce the choice of crops or that require very careful management.

Class V: Soils in these lands are not likely to erode, but have other limitations like wetness that are impractical to remove and as such not suitable for agriculture, but suitable for pasture or forestry with minor limitations.

Class VI: The lands have severe limitations that make them generally unsuitable for cultivation, but suitable for pasture or forestry with moderate limitations.

Class VII: The lands have very severe limitations that make them unsuitable for cultivation, but suitable for pasture or forestry with major limitations.

Class VIII: Soil and other miscellaneous areas (rock lands) that have very severe limitations that nearly preclude their use for any crop production, but suitable for wildlife, recreation and installation of wind mills.

The land capability subclasses are recognised based on the dominant limitations observed within a given land capability class. The subclasses are designated by adding a lower case letter like ‘e’, ‘w’, ‘s’, or ‘c’ to the class numeral. The subclass “e” indicates that the main hazard is risk of erosion, “w” indicates drainage or wetness as a limitation for plant growth, “s” indicates shallow soil depth, coarse or heavy textures, calcareousness, salinity/alkalinity or gravelliness and “c” indicates limitation due to climate.

The land capability subclasses have been further subdivided into land capability units based on the kinds of limitations present in each subclass. Ten land capability units are used in grouping the soil map units. They are stony or rocky (0), erosion hazard (slope, erosion) (1), coarse texture (sand, loamy sand, sandy loam) (2), fine texture (cracking clay, silty clay) (3), slowly permeable subsoil (4), coarse underlying material (5), salinity/alkali (6), stagnation, overflow, high ground water table (7), soil depth (8) and fertility problems (9). The capability units thus identified have similar soil and land characteristics that respond similarly to a given level of management. The soils of the microwatershed have been classified upto land capability subclass level.

The 18 soil map units identified in the Kesalapura-1 Microwatershed are grouped under two land capability classes and seven land capability subclasses (Fig. 5.1).




Fig. 5.1 Land Capability map of Kesalapura-1 Microwatershed

Entire cultivated area of the microwatershed is suitable for agriculture. Maximum area of 404 ha (70%) are good lands (Class II) that have minor limitations and require moderate conservation practices and are distributed in the major part of the microwatershed. Moderately good lands (Class III) cover an area of 149 ha (26%) and are distributed in the northwestern, eastern and central part of the microwatershed with moderate problems of soil that require special conservation practices. The other miscellaneous areas cover about 4 per cent is habitations and water bodies.

5.2 Soil Depth

Soil depth refers to the depth of the soil occurring above the parent material or hard rock. The depth of the soil determines the effective rooting depth for plants and in accordance with soil texture, mineralogy and gravel content, the capacity of the soil column to hold water and nutrient availability. Soil depth is one of the most important soil characteristic that is used in differentiating soils into different soil series. The soil depth classes used in identifying soils in the field are very shallow (<25 cm), shallow (25-50 cm), moderately shallow (50-75 cm), moderately deep (75-100 cm), deep (100-150 cm) and very deep (>150 cm). They were used to classify the soils into different depth classes and a soil depth map was generated (Fig. 5.2).

An area of 88 ha (15%) is moderately shallow (50-75 cm) and are distributed in the northwestern, southeastern and central part of the microwatershed. Maximum area of about 247 ha (43%) is moderately deep (75-100 cm) and are distributed in the major part of the microwatershed. An area of 122 ha (21%) is deep (100-150 cm) and are distributed in the central, eastern and southern part of the microwatershed. Very deep (>150 cm) soils occupy an area of 95 ha (17%) and are distributed in the northern and western part of the microwatershed.

The most problem lands with an area of about 88 ha (15%) having moderately shallow (50-75 cm) rooting depth. They are suitable for growing short duration agricultural crops but well suited for pasture, forestry or other recreational purposes. The most productive lands cover a maximum area about 217 ha (38%) where all climatically adapted long duration crops be grown.




Fig. 5.2 Soil Depth map of Kesalapura-1 Microwatershed

5.3 Surface Soil Texture

Texture is an expression to indicate the coarseness or fineness of the soil as determined by the relative proportion of primary particles of sand, silt and clay. It has a direct bearing on the structure, porosity, adhesion and consistence. The surface layer of a soil to a depth of about 25 cm is the layer that is most used by crops and plants. The surface soil textural class provides a guide to understanding soil-water retention and availability, nutrient holding capacity, infiltration, workability, drainage, physical and chemical behaviour, microbial activity and crop suitability. The textural classes used for LRI were used to classify and a surface soil texture map was generated (fig. 5.3). The area extent and their spatial distribution in the microwatershed is shown in figure 5.3.

Maximum area of 319 ha (55%) has loamy soils at the surface and are distributed in the major part of the microwatershed. An area of 233 ha (41%) has clayey soils at the surface and are distributed in the northern and western of the microwatershed (Fig. 5.3).

The most productive lands 233 ha (41%) with respect to surface soil texture are the clayey soils that have high potential for soil-water retention and availability, and nutrient retention and availability, but have more problems of drainage, infiltration, workability and other physical problems. The other most productive lands 319 ha (55%) are loamy soils which also have high potential for AWC, nutrient availability but have no drainage or other physical problems compared to loamy soils.




Fig. 5.3 Surface Soil Texture map of Kosalapura-1 Microwatershed

5.4 Soil Gravelliness

Gravel is the term used for describing coarse fragments between 2 mm and 7.5 cm diameter and stones for those between 7.5 cm and 25 cm. The presence of gravel and stones in soil reduces the volume of soil responsible for moisture and nutrient storage, drainage, infiltration and runoff, and hinders plant growth by impeding root growth and seedling emergence, intercultural operations and farm mechanization. The gravelliness classes used in LRI were used to classify the soils and using these classes a gravelliness map was generated. The area extent and their spatial distribution in the microwatershed is given in figure 5.4.

The soils that are non-gravelly (<15% gravel) cover a maximum area of 240 ha (42%) and are distributed in the northern, central, western and eastern part of the microwatershed. An area of 253 ha (44%) is covered by gravelly (15-35% gravel) soils and are distributed in the major part of the microwatershed. An area of 60 ha (10%) is very gravelly (35-60%) and are distributed in the southern, eastern and central part of the microwatershed (Fig. 5.4).

The most productive lands with respect to gravelliness are found to be 42%. They are non-gravelly with less than 15 per cent gravel and have potential for growing both annual and perennial crops. The problem soils that are gravelly (15-35%) to extremely gravelly (60-80%) cover 313 ha (54%) where only short or medium duration crops can be grown.




Fig. 5.4 Soil Gravelliness map of Kesimalapura-1 Microwatershed

5.5 Available Water Capacity

The soil available water capacity (AWC) is estimated based on the ability of the soil column to retain water between the tensions of 0.33 and 15 bar in a depth of 100 cm or the entire solum if the soil is shallower. The AWC of the soils (soil series) as estimated by considering the soil texture, mineralogy, soil depth and gravel content (Sehgal *et al.*, 1990) and accordingly the soil map units were grouped into five AWC classes *viz.*, very low (<50 mm/m), low (50-100 mm/m), medium (100-150 mm/m), high (150-200 mm/m) and very high (>200 mm/m) and using these values, an AWC map was generated (Fig. 5.5), showing the area extent and their spatial distribution in the microwatershed.

An area of about 135 ha (24%) are very low (<50 mm/m) in available water capacity and are distributed in the northern, northwestern, southeastern and central part of the microwatershed. An area of about 112 ha (20%) has soils that are low (51-100 mm/m) in available water capacity and are distributed in the eastern, northeastern, central and northwestern part of the microwatershed. Soils with medium (101-150 mm/m) available water capacity occupy a maximum area of 210 ha (36%) and are distributed in the major part of the microwatershed. An area of about 95 ha (17%) is high (151-200 mm/m) to very high (>200 mm/m) in available water capacity and are distributed in the northern and western part of the microwatershed.

An area of about 135 ha (24%) in the microwatershed has soils that are problematic with regard to available water capacity. Here, only short duration crops can be grown and the probability of crop failure is very high. These areas are best put to other alternative uses. The potential soils with respect to AWC cover about 95 ha (17%) that

have high to very high AWC, where all climatically adapted long duration crops can be grown.




Fig. 5.5 Soil Available Water Capacity map of Kosalapura-1 Microwatershed

5.6 Soil Slope

Soil slope refers to the inclination of the surface of the land. It is defined by gradient, shape and length, and is an integral feature of any soil as a natural body. Slope is considered important in soil genesis, land use and land development. The length and gradient of slope influences the rate of runoff, infiltration, erosion and deposition. The soil map units were grouped into four slope classes and a slope map was generated showing the area extent and their geographic distribution of different slope classes in the microwatershed (Fig. 5.6).

An area of 4 ha (1%) is nearly level (0-1%) and are distributed in the northern part of the microwatershed. Major area of about 549 ha (95%) falls under very gently sloping (1-3% slope) lands and are distributed in all parts of the microwatershed. In all these lands, all climatically adapted annual and perennial crops can be grown without much soil and water conservation and other land development measures.




Fig. 5.6 Soil Slope map of Kesalapura-1 Microwatershed

5.7 Soil Erosion

Soil erosion refers to the wearing away of the earth’s surface by the forces of water, wind and ice involving detachment and transport of soil by raindrop impact. It is used for accelerated soil erosion resulting from disturbance of the natural landscape by burning, excessive grazing and indiscriminate felling of forest trees and tillage, all usually by man. The erosion classes showing an estimate of the current erosion status as judged from field observations in the form of rills, gullies or a carpet of gravel on the surface are recorded. Four erosion classes, viz, slight erosion (e1), moderate erosion (e2), severe erosion (e3) and very severe erosion (e4) are recognized. The soil map units were grouped into different erosion classes and a soil erosion map generated. The area extent and their spatial distribution in the microwatershed is given in Figure 5.7.

Soils that are slightly eroded (e1 Class) occupy an area of about 211 ha (37%) and are distributed in the northern, northwestern, eastern, southern and western part of the microwatershed. Moderately eroded (e2 Class) soils cover an area of 341 ha (59%) and are distributed in the major part of the microwatershed.

An area of about 341 ha (59%) in the microwatershed is problematic because of moderate erosion. These areas need soil and water conservation and other land development measures for restoring the soil health.




Fig. 5.7 Soil Erosion map of Kesalapura-1 Microwatershed

FERTILITY STATUS

Soil fertility plays an important role in increasing crop yield. The adoption of high yielding varieties that require high amounts of nutrients has resulted in deficiency symptoms in crops and plants due to imbalanced fertilization and poor inherent fertility status, as these areas are characterised by low rainfall and high temperatures. Hence, it is necessary to know the fertility (macro and micro nutrients) status of the soils of the watersheds for assessing the kind and amount of fertilizers required for each of the crop intended to be grown. For this purpose, the surface soil samples collected from the grid points (one soil sample at every 320 m grid interval) all over the microwatershed through land resource inventory in the year 2018 were analysed for pH, EC, organic carbon, available phosphorus and potassium and for micronutrients like zinc, boron, copper, iron and manganese, and secondary nutrient sulphur.

Soil fertility data generated has been assessed and individual maps for all the nutrients for the microwatershed have been generated by using the Kriging method under GIS. The village/survey number wise fertility data for the microwatershed is given in Appendix-II.

6.1 Soil Reaction (pH)

The soil analysis of the Kesalapura-1 Microwatershed for soil reaction (pH) showed that a maximum area of 353 ha (61%) is moderately alkaline (pH 7.8-8.4) and are distributed in the major part of the microwatershed. An area of 168 ha (29%) is strongly alkaline (pH 8.4-9.0) and are distributed in the western and northern part of the microwatershed. An area of about 31 ha (5%) is very strongly alkaline (pH >9.0) and are distributed in the northern and northwestern part of the microwatershed. Thus, all soils in the microwatershed are alkaline covering 552 ha.

6.2 Electrical Conductivity (EC)

The Electrical Conductivity of the soils is $<2 \text{ dS m}^{-1}$ in the entire microwatershed (Fig. 6.2) area and as such the soils are nonsaline.

6.3 Organic Carbon

The soil organic carbon content (an index of available Nitrogen) in the soils of the microwatershed is low ($<0.5\%$) covering an area of 103 ha (18%) and is distributed in the southern part of the microwatershed. Maximum area of 403 ha (70%) is medium (0.5-0.75%) and are distributed in the major part of the microwatershed. An area of 46 ha (8%) is high ($>0.75\%$) and are distributed in the northeastern part of the microwatershed (Fig. 6.3).




Fig. 6.1 Soil Reaction (pH) map of Kesalapura-1 Microwatershed




Fig. 6.2 Electrical Conductivity (EC) map of Kesalapura-1 Microwatershed




Fig. 6.3 Soil Organic Carbon map of Kesalapura-1 Microwatershed




Fig. 6.4 Soil Available Phosphorus map of Kesalapura-1 Microwatershed

6.4 Available Phosphorus

Major cultivated area of about 552 ha (96%) is medium (23-57 kg/ha) and is distributed in the all parts of the microwatershed (Fig. 6.4).

6.5 Available Potassium

Medium (145-337 kg/ha) in available potassium content occupy a maximum area of 549 ha (96%) and are distributed in the major part of the microwatershed. An area of about 3 ha (1%) is high (>337 kg/ha) and are distributed in the northern part of the microwatershed (Fig. 6.5).

6.6 Available Sulphur

Soils that are low in available sulphur (<10 ppm) in a maximum area of 543 ha (94%) and is distributed in the major part of the microwatershed. An area of 10 ha (2%) is medium (10-20 ppm) and is distributed in the southwestern part of the microwatershed (Fig. 6.6).

6.7 Available Boron

Available boron content is low (<0.5 ppm) in an area of 12 ha (2%) and are distributed in the southern part of the microwatershed. Maximum area of about 473 ha (82%) is medium (0.5-1.0 ppm) in available boron and are distributed in the major part of the microwatershed. An area of 67 ha (12%) is high (>1.0 ppm) and is distributed in the western part of the microwatershed (Fig. 6.7).

6.8 Available Iron

Entire area of about 552 ha (96%) is deficient (<4.5 ppm) in the microwatershed (Fig. 6.8).

6.9 Available Manganese

Available manganese content is sufficient (>1.0 ppm) in the entire microwatershed area (Fig. 6.9).

6.10 Available Copper

Available copper content is sufficient (>0.2 ppm) in the entire microwatershed area (Fig. 6.10).




Fig. 6.5 Soil Available Potassium map of Kosalapura-1 Microwatershed




Fig. 6.6 Soil Available Sulphur map of Kosalapura-1 Microwatershed




Fig. 6.7 Soil Available Boron map of Kesalapura-1 Microwatershed




Fig. 6.8 Soil Available Iron map of Kesalapura-1 Microwatershed




Fig. 6.9 Soil Available Manganese map of Kesalapura-1 Microwatershed




Fig. 6.10 Soil Available Copper map of Kesalapura-1 Microwatershed

6.11 Available Zinc

Available zinc content is deficient (<0.6 ppm) in an area of 33 ha (6%) and are distributed in the southern part of the microwatershed. Maximum area of 520 ha (90%) is sufficient (>0.6 ppm) and are distributed in the major part of the microwatershed (Fig. 6.11).




Fig. 6.11 Soil Available Zinc map of Kesalapura-1 Microwatershed

LAND SUITABILITY FOR MAJOR CROPS

The soil and land resource units (soil phases) of Kesalapura-1 Microwatershed were assessed for their suitability for growing food, fodder, fibre and other horticulture crops by following the procedure as outlined in FAO, 1976 and 1983. Crop requirements were developed for each of the crop from the available research data and also by referring to Naidu *et. al.* (2006) and Natarajan *et. al.* (2015). The crop requirements (Table 7.2 to 7.33) were matched with the soil and land characteristics (Table 7.1) to arrive at the crop suitability. The criteria tables are given at the end of the Chapter. In FAO land suitability classification, two orders are recognized. Order S-Suitable and Order N- Not suitable. The orders have classes, subclasses and units. Order-S has three classes, Class S1-Highly Suitable, Class S2-Moderately Suitable and Class S3-Marginally Suitable. Order N has two Classes, N1-Currently not Suitable and N2-Permanently not Suitable. There are no subclasses within the Class S1 as they will have very minor or no limitations for crop growth. Classes S2, S3, N1 and N2 are divided into subclasses based on the kinds of limitations encountered. The limitations that affect crop production are ‘c’ for erratic rainfall and its distribution and length of growing period (LGP), ‘e’ for erosion hazard, ‘r’ for rooting condition, ‘t’ for lighter or heavy texture, ‘g’ for gravelliness or stoniness, ‘n’ for nutrient availability, ‘l’ for topography, ‘m’ for moisture availability, ‘z’ for calcareousness ‘s’ for sodium and ‘w’ for drainage. These limitations are indicated as lower case letters to the class symbol. For example, moderately suitable lands with the limitations of soil depth and erosion are designated as S2re. For the microwatershed, the soil mapping units were evaluated and classified up to subclass level.

Using the above criteria, the soil map units of the microwatershed were evaluated and land suitability maps for 31 major agricultural and horticultural crops were generated. The detailed information on the kind of suitability of each of the soil phase for the crops assessed are given village/ survey number wise for the microwatershed in Appendix-III.

7.1 Land Suitability for Sorghum (*Sorghum bicolor*)

Sorghum is one of the major food crop grown in Karnataka in an area of 10.47 lakh ha in Bijapur, Gulbarga, Raichur, Bidar, Belgaum, Dharwad, Bellary, Chitradurga, Mysore and Chamarajnar districts. The crop requirements for growing sorghum (Table 7.2) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing sorghum was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure. 7.1.

An area of 192 ha (33%) is highly suitable (Class S1) lands for growing sorghum and are distributed in the northern, western, central and eastern part of the microwatershed. Maximum area of 227 ha (39%) is moderately suitable (Class S2) and

are distributed in the major part of the microwatershed. They have minor limitations of gravelliness, calcareousness, drainage and rooting condition. An area of about 133 ha (23%) is marginally suitable (Class S3) for growing sorghum and are distributed in the northern, southern, central and southeastern part of the microwatershed with moderate limitations of gravelliness and rooting condition.




Fig. 7.1 Land Suitability map of Sorghum

7.2 Land Suitability for Maize (*Zea mays*)

Maize is one of the most important food crop grown in an area of 13.37 lakh ha in almost all the districts of the State. The crop requirements for growing maize (Table 7.3) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing maize was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.2.

An area of 85 ha (15%) is highly suitable (Class S1) for growing maize and are distributed in the eastern and central part of the microwatershed. Maximum area of 333 ha (58%) is moderately suitable (Class S2) for growing maize and are distributed in the major part of the microwatershed with minor limitations of calcareousness, gravelliness, rooting condition, drainage and texture. Marginally suitable (Class S3) lands cover an area of 133 ha (23%) and are distributed in the northern, southern, southeastern and central part of the microwatershed. They have moderate limitations of gravelliness, rooting condition and gravelliness.




Fig. 7.2 Land Suitability map of Maize

7.3 Land Suitability for Bajra (*Pennisetum glaucum*)

Bajra is one of the major food crop grown in an area of 2.34 lakh ha in the northern districts of the Karnataka State. The crop requirements for growing bajra (Table 7.4) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing bajra was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.3.

Maximum area of 266 ha (46%) is highly suitable (Class S1) for growing bajra and are distributed in the major part of the microwatershed. An area of 226 ha (39%) is moderately suitable (Class S2) and are distributed in the major part of the microwatershed with minor limitations of texture, gravelliness, rooting condition, drainage and calcareousness. Marginally suitable (Class S3) lands cover an area of 60 ha (10%) and are distributed in the southern, eastern and central part of the microwatershed. They have moderate limitation of gravelliness.




Fig. 7.3 Land Suitability map of Bajra

7.4 Land Suitability for Groundnut (*Arachis hypogaea*)

Groundnut is one of the major oilseed crop grown in an area of 6.54 lakh ha in Karnataka in most of the districts either as rainfed or irrigated crop. The crop requirements for growing groundnut (Table 7.5) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing groundnut was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.4.

An area of 38 ha (7%) is highly suitable (Class S1) for growing groundnut and are distributed in the western part of the microwatershed. Maximum area of 457 ha (80%) is moderately suitable (Class S2) and are distributed in the major part of the microwatershed. They have minor limitations of rooting condition, texture, drainage and graveliness. An area of 58 ha (10%) is marginally suitable (Class S3) and are distributed in the northern and western part of the microwatershed. They have moderate limitations of calcareousness and texture.




Fig. 7.4 Land Suitability map of Groundnut

7.5 Land Suitability for Sunflower (*Helianthus annus*)

Sunflower is one of the most important oilseed crop grown in an area of 3.56 lakh ha in the State in all the districts. The crop requirements for growing sunflower (Table 7.6) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing sunflower was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.5.

An area of 132 ha (23%) is highly suitable (Class S1) for growing sunflower and are distributed in the northern, western, central and eastern part of the microwatershed. Maximum area of 273 ha (47%) is moderately suitable (Class S2) and are distributed in the major part of the microwatershed. They have minor limitations of calcareousness, gravelliness, drainage and rooting condition. An area of 147 ha (26%) is marginally suitable (Class S3) for growing sunflower and are distributed in the northern, eastern and northwestern part of the microwatershed with moderate limitations of gravelliness and rooting condition.




Fig. 7.5 Land Suitability map of Sunflower

7.6 Land Suitability for Red gram (*Cajanus cajan*)

Redgram is one of the most important pulse crop grown in an area of 7.28 lakh ha in almost all the districts of the State. The crop requirements for growing redgram (Table 7.7) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing redgram was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.6.

An area of 123 ha (21%) is highly suitable (Class S1) for growing red gram and are distributed in the central, eastern and western part of the microwatershed. Moderately suitable (Class S2) lands occupy a maximum area of 234 ha (41%) and are distributed in the major part of the microwatershed with minor limitations of gravelliness, texture, rooting condition, drainage and calcareousness. Marginally suitable (Class S3) lands cover an area of 195 ha (34%) and are distributed in the northern, northwestern, southern, central and eastern part of the microwatershed. They have moderate limitations of gravelliness and rooting condition.




Fig. 7.6 Land Suitability map of Redgram

7.7 Land Suitability for Bengalgram (*Cicer arietinum*)

Bengalgram is one of the major pulse crop grown in an area of 9.39 lakh ha in northern Karnataka in Bijapur, Gulbarga, Raichur, Bidar, Belgaum, Dharwad and Bellary districts. The crop requirements for growing Bengalgram (Table 7.8) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing Bengalgram was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.7.

An area of 10 ha (2%) is highly suitable (Class S1) for growing bengalgram and are distributed in the northern part of the microwatershed. Moderately suitable lands (Class S2) occupy a maximum area of 435 ha (76%) and are distributed in the major part of the microwatershed with minor limitations of gravelliness, calcareousness, texture, drainage and rooting condition. Marginally suitable (Class S3) lands cover an area of 107 ha (19%) and are distributed in the northern, eastern, southern and central part of the microwatershed. They have moderate limitation of gravelliness.




Fig. 7.7 Land Suitability map of Bengalgram

7.8 Land Suitability for Cotton (*Gossypium hirsutum*)

Cotton is one of the most important fibre crop grown in the State in about 8.75 lakh ha area in Raichur, Dharwad, Belgaum, Gulbarga, Bijapur, Bidar, Bellary, Chitradurga and Chamarajnagar districts. The crop requirements for growing cotton (Table 7.9) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing cotton was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.8.

An area of 96 ha (17%) is highly suitable (Class S1) for growing cotton and are distributed in the northern, western, central and eastern part of the microwatershed. Moderately suitable (Class S2) lands occupy a maximum area of 323 ha (56%) and are distributed in the major part of the microwatershed. They have minor limitations of rooting condition, texture, calcareousness, drainage and gravelliness. Marginally suitable (Class S3) lands cover an area of 134 ha (23%) and are distributed in the northern, central, southern and eastern part of the microwatershed. They have moderate limitations of gravelliness and rooting condition.




Fig. 7.8 Land Suitability map of Cotton

7.9 Land Suitability for Chilli (*Capsicum annum L*)

Chilli is one of the most important commercial spice crop grown in an area of 0.89 lakh ha in all the districts of Karnataka State. The crop requirements for growing chilli (Table 7.10) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing chilli was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.9.

Maximum area of 219 ha (38%) is highly (Class S1) suitable for growing chilli and are distributed in the major part of the microwatershed. An area of 153 ha (27%) is moderately suitable (Class S2) for growing chilli and are distributed in the northern, northwestern, central and southern part of the microwatershed. They have minor limitations of rooting condition, gravelliness, texture, drainage and calcareousness. An area of 181 ha (32%) is marginally suitable (Class S3) and are distributed in the northern, western, southern, central and southeastern part of the microwatershed. They have moderate limitations of texture, rooting condition, gravelliness and calcareousness.




Fig. 7.9 Land Suitability map of Chilli

7.10 Land Suitability for Tomato (*Solanum lycopersicum*)

Tomato is one of the most important vegetable crop grown in an area of 0.65 lakh ha in almost all the districts of the State. The crop requirements (Table 7.11) for growing tomato were matched with the soil-site characteristics (Table 7.11) and a land suitability map for growing tomato was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.10.

Maximum area of 219 ha (38%) is highly (Class S1) suitable lands for growing tomato and are distributed in the major part of the microwatershed. An area of 153 ha (27%) is moderately suitable (Class S2) and are distributed in the northern, northwestern, central and southern part of the microwatershed. They have minor limitations of rooting condition, gravelliness, texture, drainage and calcareousness. Marginally suitable (Class S3) lands occupy an area of 181 ha (32%) and are distributed in the northern, central, western, southern and southeastern part of the microwatershed with moderate limitations of gravelliness, rooting condition, texture and calcareousness.




Fig. 7.10 Land Suitability map of Tomato

7.11 Land Suitability for Brinjal (*Solanum melongena*)

Brinjal is one of the most important vegetable crop grown in all the districts. The crop requirements for growing brinjal (Table 7.12) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing brinjal was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.11.

An area of 204 ha (36%) is highly suitable (Class S1) for growing brinjal and are distributed in the northern, central, southern and northeastern part of the microwatershed. Maximum area of about 212 ha (37%) is moderately suitable (Class S2) for growing brinjal and are distributed in the major part of the microwatershed with minor limitations of texture, rooting condition, calcareousness, drainage and gravelliness. Marginally suitable lands (Class S3) occur in an area of 135 ha (24%) and are distributed in the northern, northwestern, southeastern and central part of the microwatershed with moderate limitation of gravelliness.




Fig. 7.11 Land Suitability map of Brinjal

7.12 Land Suitability for Onion (*Allium cepa*)

Onion is one of the most important vegetable crop grown in Raichur, Dharwad, Belgaum, Gulbarga, Bijapur, Bidar, Bellary, Chitradurga and Tumakuru districts. The crop requirements for growing onion (Table 7.13) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing Onion was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.12.

An area of 61 ha (11%) is highly (Class S1) suitable for growing onion and are distributed in the central and southern part of the microwatershed. Maximum area of 298 ha (52%) is moderately suitable (Class S2) and are distributed in the major part of the microwatershed. They have minor limitations of gravelliness, drainage and texture. Marginally suitable lands (Class S3) occupy an area of 192 ha (34%) and are distributed in the northern, southwestern, central and southeastern part of the microwatershed with moderate limitations of gravelliness, calcareousness and texture.




Fig. 7.12 Land Suitability map of Onion

7.13 Land Suitability for Bhendi (*Abelmoschus esculentus*)

Bhendi is one of the most important vegetable crop grown in all the districts. The crop requirements for growing bhendi (Table 7.14) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing bhendi was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.13.

An area of 61 ha (11%) is highly suitable (Class S1) for growing bhendi and are distributed in the central and southern part of the microwatershed. Maximum area of about 355 ha (62%) is moderately suitable (Class S2) and are distributed in the major part of the microwatershed with minor limitations of texture, calcareousness, gravelliness, drainage and rooting condition. An area of 135 ha (24%) is marginally suitable (Class S3) for growing bhendi and are distributed in the northern, central and southeastern part of the microwatershed with moderate limitation of gravelliness.




Fig. 7.13 Land Suitability map of Bhendi

7.14 Land Suitability for Drumstick (*Moringa oleifera*)

Drumstick is one of the most important vegetable crop grown in 2403 ha area in the State. The crop requirements for growing drumstick (Table 7.15) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing drumstick was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.14.

An area of 147 ha (26%) is highly suitable (Class S1) for growing drumstick and are distributed in the western, central, eastern and southern part of the microwatershed. Maximum area of 221 ha (39%) is moderately suitable (Class S2) and are distributed in the major part of the microwatershed. They have minor limitations of gravelliness, texture, rooting condition, drainage and calcareousness. An area of 183 ha (32%) is marginally suitable (Class S3) for growing drumstick and are distributed in the northern, central, southern and southeastern part of the microwatershed. They have moderate limitations of gravelliness and rooting condition.




Fig. 7.14 Land Suitability map of Drumstick

7.15 Land Suitability for Mango (*Mangifera indica*)

Mango is one of the most important fruit crop grown in about 1.73 lakh ha in almost all the districts of the State. The crop requirements (Table 7.16) for growing mango were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing mango was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.15.

An area of 86 ha (15%) is highly (Class S1) suitable for growing mango and are distributed in the western, eastern and central part of the microwatershed. An area of 62 ha (11%) is moderately suitable (Class S2) and are distributed in the southern and central part of the microwatershed. They have minor limitations rooting condition and gravelliness. Marginally suitable (Class S3) lands cover a maximum area of 316 ha (55%) and are distributed in the major part of the microwatershed. They have moderate limitations of texture, gravelliness, calcareousness and rooting condition. An area of 88 ha (15%) is currently not suitable (Class N1) for growing mango and occur in the northwestern, central and southeastern part of the microwatershed with severe limitations of gravelliness and rooting condition.




Fig. 7.15 Land Suitability map of Mango

7.16 Land suitability for Guava (*Psidium guajava*)

Guava is one of the most important fruit crop grown in an area of about 0.64 lakh ha in almost all the districts of the State. The crop requirements (Table 7.17) for growing guava were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing guava was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.16.

An area of 74 ha (13%) is highly (Class S1) for growing guava and are distributed in the western, eastern and central part of the microwatershed. Moderately (Class S2) suitable lands occupy in a maximum area of 320 ha (56%) and are distributed in the major part of the microwatershed. Marginally suitable (Class S3) lands cover an area of 157 ha (27%) and are distributed in the northern, western, central and southeastern part of the microwatershed. They have moderate limitations of gravelliness, texture, rooting condition and drainage.




Fig. 7.16 Land Suitability map of Guava

7.17 Land Suitability for Sapota (*Manilkara zapota*)

Sapota is one of the most important fruit crop grown in an area of about 29373 ha in almost all the districts of the State. The crop requirements (Table 7.18) for growing sapota were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing sapota was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.17.

An area of 123 ha (21%) is highly (Class S1) suitable for growing sapota and are distributed in the western, central and eastern part of the microwatershed. Moderately (Class S2) suitable lands occupy a maximum area of 224 ha (39%) and are distributed in the major part of the microwatershed with minor limitations of gravelliness, rooting condition and drainage. Marginally suitable (Class S3) lands cover an area of 205 ha (36%) and are distributed in the northern, central, western, southern and southeastern part of the microwatershed. They have moderate limitations of gravelliness, rooting condition, texture and calcareousness.




Fig. 7.17 Land Suitability map of Sapota

7.18 Land Suitability for Pomegranate (*Punica granatum*)

Pomegranate is one of the commercially grown fruit crop in about 18488 ha in Karnataka mainly in Bijapur, Bagalkot, Koppal, Gadag and Chitradurga districts. The crop requirements for growing pomegranate (Table 7.19) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing pomegranate was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.18.

An area of 123 ha (21%) is highly suitable (Class S1) for growing pomegranate and are distributed in the western, central and eastern part of the microwatershed. Moderately suitable (Class S2) lands occupy a maximum area of 282 ha (49%) and are distributed in the major part of the microwatershed. They have minor limitations of texture, rooting condition, gravelliness, drainage and calcareousness. An area of 148 ha (26%) is marginally suitable (Class S3) for growing pomegranate and are distributed in the northwestern, central, southern and southeastern part of the microwatershed. They have moderate limitations of gravelliness and rooting condition.




Fig. 7.18 Land Suitability map of Pomegranate

7.19 Land Suitability for Musambi (*Citrus limetta*)

Musambi is one of the most important fruit crop grown in almost all the districts of the State. The crop requirements for growing musambi (Table 7.20) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing musambi was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.19.

An area of 132 ha (23%) is highly suitable (Class S1) for growing musambi and are distributed in the northern, western, central and eastern part of the microwatershed. Maximum area of 273 ha (47%) is moderately suitable (Class S2) and are distributed in the major part of the microwatershed. They have minor limitations of calcareousness, rooting condition, drainage and gravelliness. Marginally suitable (Class S3) lands occur in an area of 148 ha (26%) and are distributed in the northwestern, central, southern and southeastern part of the microwatershed with moderate limitations of gravelliness and rooting condition.




Fig. 7.19 Land Suitability map of Musambi

7.20 Land Suitability for Lime (*Citrus sp*)

Lime is one of the most important fruit crop grown in an area of 0.11 lakh ha in almost all the districts of the State. The crop requirements for growing lime (Table 7.21) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing lime was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.20.

An area of 132 ha (23%) is highly suitable (Class S1) for growing lime and are distributed in the northern, western, central and eastern part of the microwatershed. Maximum area of 320 ha (56%) is moderately suitable (Class S2) and are distributed in the major part of the microwatershed. They have minor limitations of calcareousness, rooting condition, drainage and gravelliness. Marginally suitable (Class S3) lands occur in an area of 100 ha (17%) and distributed in the northwestern, central and southeastern part of the microwatershed with moderate limitations of gravelliness and rooting condition.




Fig. 7.20 Land Suitability map of Lime

7.21 Land Suitability for Amla (*Phyllanthus emblica*)

Amla is one of the most important medicinal crop grown in 151 ha area and distributed in almost all the districts of the State. The crop requirements for growing amla (Table 7.22) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing amla was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.21.

Maximum area of 300 ha (52%) is highly suitable (Class S1) for growing amla and are distributed in the major part of the microwatershed. An area of 252 ha (44%) has soils that are moderately suitable (Class S2) and are distributed in the northern, central, western, northwestern, southern and southeastern part of the microwatershed. They have minor limitations of rooting condition, gravelliness, texture, drainage and calcareousness.




Fig. 7.21 Land Suitability map of Amla

7.22 Land Suitability for Cashew (*Anacardium occidentale*)

Cashew is one of the most important nut crop grown in an area of 1.24 lakh ha in almost all the districts of the State. The crop requirements for growing cashew (Table 7.23) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing cashew was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.22.

An area of 99 ha (17%) is highly (Class S1) suitable for growing cashew and are distributed in the western, central and southern part of the microwatershed. Moderately (Class S2) suitable lands occur in a maximum area of 286 ha (50%) and are distributed in the major part of the microwatershed with minor limitations of texture, gravelliness and rooting condition. Marginally suitable (Class S3) lands occur in an area of 100 ha (17%) for growing cashew and are distributed in the northwestern, central and southeastern part of the microwatershed with moderate limitations of rooting condition and gravelliness. An area of about 67 ha (12%) is currently not suitable (Class N1) for growing cashew and are distributed in the northern, western and southern part of the microwatershed with severe limitations of texture, drainage and calcareousness.




Fig. 7.22 Land Suitability map of Cashew

7.23 Land Suitability for Jackfruit (*Artocarpus heterophyllus*)

Jackfruit is one of the most important fruit crop grown in 5368 ha in all the districts of the State. The crop requirements for growing jackfruit (Table 7.24) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing jackfruit was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in figure 7.23.

An area of 123 ha (21%) is highly (Class S1) for growing jackfruit and are distributed in the western, central and eastern part of the microwatershed. Moderately (Class S2) suitable lands occupy a maximum area of 272 ha (47%) and are distributed in the major part of the microwatershed. They have minor limitations of gravelliness, rooting condition and drainage. Marginally suitable (Class S3) lands cover an area of 157 ha (27%) and are distributed in the northern, northwestern, western, central and southeastern part of the microwatershed. They have moderate limitations of gravelliness, rooting condition, texture and calcareousness.




Fig. 7.23 Land Suitability map of Jackfruit

7.24 Land Suitability for Jamun (*Syzygium cumini*)

Jamun is an important fruit crop grown in almost all the districts of the State. The crop requirements for growing jamun (Table 7.25) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing jamun was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.24.

An area of 86 ha (15%) is highly suitable (Class S1) for growing jamun and are distributed in the western, central and eastern part of the microwatershed. Maximum area of 356 ha (62%) is moderately suitable (Class S2) and occur in the major part of the microwatershed. They have minor limitations of rooting condition, texture, gravelliness and calcareousness. Marginally suitable (Class S3) lands cover an area of 109 ha (19%) and are distributed in the northwestern, central and southeastern part of the microwatershed with moderate limitations of rooting condition, drainage and gravelliness.




Fig. 7.24 Land Suitability map of Jamun

7.25 Land Suitability for Custard Apple (*Annona reticulata*)

Custard apple is one of the most important fruit crop grown in almost all the districts of the State. The crop requirements for growing custard apple (Table 7.26) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing custard apple was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.25.

Maximum area of 300 ha (52%) is highly (Class S1) suitable for growing custard apple and are distributed in the major part of the microwatershed. An area of 252 ha (44%) is moderately suitable (Class S2) and are distributed in the northern, western, central, southern and southeastern part of the microwatershed. They have minor limitations of gravelliness, rooting condition, drainage and calcareousness.




Fig. 7.25 Land Suitability map of Custard Apple

7.26 Land Suitability for Tamarind (*Tamarindus indica*)

Tamarind is one of the most important spice crop grown in 14897 ha in all the districts of the State. The crop requirements for growing tamarind (Table 7.27) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing tamarind was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.26.

An area of 86 ha (15%) is highly (Class S1) suitable for growing tamarind and are distributed in the western, central and eastern part of the microwatershed. An area of 119 ha (21%) is moderately suitable (Class S2) and occur in the northern, central, southern and western part of the microwatershed. They have minor limitations of rooting condition, texture and calcareousness. Maximum area of 259 ha (45%) is marginally suitable (Class S3) and occur in major part of the microwatershed with moderate limitations of gravelliness, drainage and rooting condition. An area of 88 ha (15%) is currently not suitable (Class N1) and are distributed in the northwestern, central and southeastern part of the microwatershed with severe limitations of rooting condition and gravelliness.




Fig. 7.26 Land Suitability map of Tamarind

7.27 Land Suitability for Mulberry (*Morus nigra*)

Mulberry is the most important leaf crop grown for rearing silkworms in about 1.66 lakh ha in all the districts of the State. The crop requirements for growing mulberry (Table 7.28) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing mulberry was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.27.

An area of 147 ha (26%) is highly suitable (Class S1) for growing mulberry and are distributed in the western, central, eastern and southern part of the microwatershed. Moderately suitable (Class S2) lands occupy a maximum area of 307 ha (53%) and are distributed in the major part of the microwatershed. They have minor limitations of gravelliness, calcareousness, rooting condition, drainage and texture. Marginally suitable (Class S3) lands cover an area of 90 ha (17%) and are distributed in the northern, northwestern and southeastern part of the microwatershed. They have moderate limitations of texture, rooting condition, gravelliness and calcareousness.




Fig. 7.27 Land Suitability map of Mulberry

7.28 Land Suitability for Marigold (*Tagetes erecta*)

Marigold is one of the most important flower crop grown in an area of 1858 ha in almost all the districts of the State. The crop requirements for growing marigold (Table 7.29) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing marigold was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.28.

An area of 123 ha (21%) is highly suitable (Class S1) for growing marigold and are distributed in the western, central and eastern part of the microwatershed. Maximum area of 296 ha (52%) is moderately suitable (Class S2) and are distributed in the major part of the microwatershed. They have minor limitations of texture, gravelliness, rooting condition, drainage and calcareousness. An area of 133 ha (23%) is marginally suitable (Class S3) and are distributed in the northern, southern, central and southeastern part of the microwatershed. They have moderate limitations of gravelliness and rooting condition.




Fig. 7.28 Land Suitability map of Marigold

7.29 Land Suitability for Chrysanthemum (*Chrysanthemum indicum*)

Chrysanthemum is one of the most important flower crop grown in an area of 4978 ha in almost all the districts of the State. The crop requirements for growing chrysanthemum (Table 7.30) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing chrysanthemum was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed is given in Figure 7.29.

An area of 123 ha (21%) is highly suitable (Class S1) lands for growing chrysanthemum and are distributed in the western, central and eastern part of the microwatershed. Maximum area of 296 ha (52%) is moderately suitable (Class S2) for and are distributed in the major part of the microwatershed. They have minor limitations of calcareousness, gravelliness, rooting condition, drainage and texture. An area of 133 ha (23%) is marginally suitable (Class S3) and occur in the northern, southern, central and southeastern part of the microwatershed. They have moderate limitations of gravelliness and rooting condition.




Fig. 7.29 Land Suitability map of Chrysanthemum

7. 30 Land Suitability for Jasmine (*Jasminum sp.*)

Jasmine is one of the most important flower crop grown in an area of 6146 ha in almost all the districts of the State. The crop requirements (Table 7.31) for growing jasmine were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing jasmine was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.30.

An area of 123 ha (21%) is highly suitable lands (Class S1) for growing jasmine and are distributed in the western, central and eastern part of the microwatershed. Maximum area of 239 ha (42%) is moderately suitable (Class S2) and are distributed in the major part of the microwatershed. They have minor limitations of rooting condition, gravelliness, drainage and calcareousness. An area of 190 ha (33%) is marginally suitable (Class S3) for growing jasmine and are distributed in the northern, central, western, southern and eastern part of the microwatershed. They have moderate limitations of gravelliness, rooting condition, texture and calcareousness.




Fig. 7.30 Land Suitability map of Jasmine

7.31 Land Suitability for Crossandra (*Crossandra in fundibuliformis*)

Crossandra is one of the most important flower crop grown in an area of 6146 ha in almost all the districts of the State. The crop requirements (Table 7.32) for growing crossandra were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing crossandra was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.31.

An area of 123 ha (21%) is highly suitable lands (Class S1) lands for growing crossandra and are distributed in the western, central and eastern part of the microwatershed. Maximum area of 239 ha (42%) is moderately suitable (Class S2) for growing crossandra and occur in the major part of the microwatershed. They have minor limitations of gravelliness, drainage, rooting condition and texture. An area of 190 ha (33%) is marginally suitable (Class S3) and are distributed in the northern, central, western, southern and eastern part of the microwatershed. They have moderate limitations of gravelliness, rooting condition, texture and calcareousness.




Fig. 7.31 Land Suitability map of Crossandra

Table 7.1 Soil-Site Characteristics of Kesalapura-1 Microwatershed

Soil Map Units	Climate (P) (mm)	Growing period (Days)	Drainage Class	Soil depth (cm)	Soil texture		Gravelliness		AWC (mm/m)	Slope (%)	Erosion	pH	EC	ESP	CEC [Cmol (p ⁺) kg ⁻¹]	BS (%)
					Surf-ace	Sub-surface	Surf-ace	Sub-surface								
MKHcB1g1	662	90	WD	50-75	sl	gsc	15-35	>35	50-100	1-3	Slight	7.38	0.09	1.49	14.84	93
MKHcB2g1	662	90	WD	50-75	sl	gsc	15-35	>35	50-100	1-3	Moderate	7.38	0.09	1.49	14.84	93
LKRhB2g1	662	90	WD	50-75	scl	gsc	15-35	35-60	50-100	1-3	Moderate	8.18	0.30	4.51	12.19	100
HDHhB2g1	662	90	WD	75-100	scl	gsc-gc	35-60	>35	50-100	1-3	Moderate	6.54	0.07	7.11	5.84	84.07
BSRhB2g1	662	90	WD	75-100	scl	gsc	15-35	15-35	50-100	1-3	Moderate	6.59	0.12	6.0	8.80	77.55
BSRiB2	662	90	WD	75-100	sc	gsc	-	15-35	50-100	1-3	Moderate	6.59	0.12	6.0	8.80	77.55
CKMcB2g2	662	90	WD	75-100	sl	sc	35-60	-	100-150	1-3	Moderate	7.99	0.32	1.73	12.50	119
CKMhB1g1	662	90	WD	75-100	scl	sc	15-35	-	100-150	1-3	Moderate	7.99	0.32	1.73	12.50	119
CKMiB1	662	90	WD	75-100	sc	sc	-	-	100-150	1-3	Slight	7.99	0.32	1.73	12.50	119
HLPcB2g1	662	90	WD	75-100	sl	scl	15-35	-	50-100	1-3	Moderate	-	-	-	-	-
KMHhB1g1	662	90	WD	100-150	scl	sc	15-35	<15	150-200	1-3	Slight	7.2	0.19	0.54	15.07	100
KMHiB2	662	90	WD	100-150	sc	sc	-	<15	150-200	1-3	Moderate	7.2	0.19	0.54	15.07	100
BPRhB2g2	662	90	WD	100-150	scl	gsc-gc	35-60	>35	100-150	1-3	Moderate	6.64	0.03	0.51	5.45	63.45
GDPiB2	662	90	WD	100-150	sc	gsc-gc	-	>35	51-100	1-3	Moderate	7.88	0.10	2.87	7.8	97
RTRiB2	662	90	WD	>150	sc	c	-	-	150-200	1-3	Moderate	5.08	0.03	2.06	9.21	50.50
KLRmA1	662	90	MWD	>150	c	sc	-	-	>200	0-1	Slight	7.11	0.33	3.42	19.50	100
KLRhB2	662	90	MWD	>150	scl	sc	-	-	>200	1-3	Moderate	7.11	0.33	3.42	19.50	100
BGPmB1	662	90	MWD	>150	c	c	-	<15	>200	1-3	Slight	9.20	0.27	3.84	19.60	100

*Symbols and abbreviations are according to Field Guide for LRI under Sujala-III Project, Karnataka

Table 7.2 Land suitability criteria for Sorghum

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime 1	Mean temperature in growing season	°C	26–30	30–34; 24–26	34–40; 20–24	>40; <20
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristics					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, c (red), c (black)	scl, cl	ls, sl	-
	pH	1:2.5	5.5-7.8	5.0-5.5 7.8-9.0	>9.0	-
	CEC	C mol (p+)/K g				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	10-15
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

Table 7.3 Land suitability criteria for Maize

Land use requirement		Rating				
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	30-34	35-38 26-30	38-40 26-20	
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc	c (red), c (black)	ls, sl	-
	pH	1:2.5	5.5-7.8	5.0-5.5 7.8-9.0	>9.0	-
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	-
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

Table 7.4 Land suitability criteria for Bajra

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-32	33-38 24-27	39-40 20-23	<20
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm	500-750	400-500	200-400	<200
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl, scl, cl,sc,c (red)	C (black)	ls	-
	pH	1:2.5	6.0-7.8	5.0-5.5 7.8-9.0	5.5-6.0 >9.0	
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	15-35	35-60	>60	
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	1-3	3-5	5-10	>10

Table 7.5 Land suitability criteria for Groundnut

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	24–33	22–24; 33–35	20–22; 35–40	<20; >40
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. Well drained	Poorly drained	Very Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl	sl,cl, sc	c (red), c (black), ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<35	35-60	>60	
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.6 Land suitability criteria for Sunflower

Land use requirement			Rating			
Soil –site characteristics	Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)	
Climatic regime	Mean temperature in growing season	°C	24–30	30–34; 20–24	34–38; 16–20	>38; <16
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	mod. Well drained	-	Poorly to very drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	cl, sc,c (red), c (black)	scl	ls, sl	-
	pH	1:2.5	6.5-7.8	7.8-8.4 5.5-6.5	8.4-9.0; 5.0-5.5	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.7 Land suitability criteria for Red gram

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	30-35(G) 20-25(AV) 15-18 (F&PS) 35-40(M)	25-30(G) 20-25 (AV) 12-15 (F&PS) 30-35(M)	20-25(G) 15-20(AV) 10-12 (F&PS) 25-30(M)	< 20 <15 <10 <25
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. Well drained	Poorly drained	Very Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, c (red)	c (black),sl, scl, cl	ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-9.0	5.0-5.5 >9.0	-
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-50	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<1.0	1.0-2.0	>2.0	
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.8 Land suitability criteria for Bengal gram

Land use requirement			Rating			
Soil –site characteristics	Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)	
Climatic regime	Mean temperature in growing season	°C	20–25	25–30; 15–20	30–35; 10–15	>35; <10
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. Well drained	Poorly drained	Very Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	C (black)	-	c (red), scl, cl, sc	ls, sl
	pH	1:2.5	6.0-7.8	5.0-6.0 7.8-9.0	>9.0	-
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	-
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.9 Land suitability criteria for Cotton

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	22-32	>32	<19	-
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well to moderately well	Poorly drained/Some what excessively drained	-	very poorly/excessively drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, c (red,black)	cl	scl	ls, sl
	pH	1:2.5	6.5-7.8	7.8-8.4	5.5-6.5 8.4->9.0	<5.5
	CEC	C mol (p+)Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	50-100	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	<3	3-5	-	>5

Table 7.10 Land suitability criteria for Chilli

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	25-32	33-35 20-25	35-38 <20	>38
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc	c (black), sl	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.11 Land suitability criteria for Tomato

Land use requirement		Rating				
Soil –site characteristics	Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)	
Climatic regime	Mean temperature in growing season	°C	25-28	29-32 20-24	15-19 33-36	<15 >36
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl, scl, cl, sc, c (red)	-	ls, c(black)	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.12 Land suitability criteria for Brinjal

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	Well drained	Moderately well drained	Poorly drained	V. Poorly drained
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class				
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl, scl, cl, sc c (red)	-	ls, c (black)	-
	pH	1:2.5	6.0-7.3	7.3-8.4 5.0-6.0	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	>60
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.13 Land suitability criteria for Onion

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	20-30	30-35	35-40	>40
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately /imperfectly	-	Poorly to V poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl,scl,cl,sc,c (red)	-	c (Black),ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<1.0	1.0-2.0	2.0-4.0	<4
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.14 Land suitability criteria for Bhendi

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	25-28	29-32 20-24	15-19 33-36	<15 >36
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Imperfectly drained	Poorly to very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl,sc, c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.15 Land suitability criteria for Drumstick

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, scl, cl, c (red)	sl, c (black)	ls	s
	pH	1:2.5	6.0-7.3	5.0-5.5 7.3-7.8	5.5-6.0 7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<35	35-60	60-80	>80
Soil toxicity	Salinity (EC saturation extract)	dS/m				
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-10	-	>10

Table 7.16 Land suitability criteria for Mango

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-32	24-27 33-35	36-40	20-24
	Min temp. before flowering	°C	10-15	15-22	>22	-
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration	Days				
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	-	ls, sl, c (black)	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>150	100-150	75-100	<75
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.17 Land suitability criteria for Guava

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-32	33-36 24-27	37-42 20-23	
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	sl	c (black), ls	-
	pH	1:2.5	6.0-7.8	5.0-6.0	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.18 Land suitability criteria for Sapota

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-32	33-36 24-27	37-42 20-23	>42 <18
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	-	Poorly to very drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	sl	ls, c (black)	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.19 Land suitability criteria for Pomegranate

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	30-34	35-38 25-29	39-40 15-24	
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl,cl, sc, c (red)	c (black),sl	ls	-
	pH	1:2.5	5.5-7.8	7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.20 Land suitability criteria for Musambi

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-30	31-35 24-27	36-40 20-23	>40 <20
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately drained	poorly	Very poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c	sl	ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.21 Land suitability criteria for Lime

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-30	31-35 24-27	36-40 20-23	>40 <20
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately drained	poorly	Very poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c	sl	ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.22 Land suitability criteria for Amla

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. well drained	Poorly drained	V. Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	c (black)	ls, sl	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15-35	35-60	60-80	-
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

Table 7.23 Land suitability criteria for Cashew

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	32 to 34	28 to 32; 34 to 38	24 to 28; 38 to 40	<20; >40
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	-	sl, ls	c (black)
	pH	1:2.5	5.5-6.5	5.0-5.5 6.5-7.3	7.3-7.8	>7.8
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-10	>10	-

Table 7.24 Land suitability criteria for Jackfruit

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. well	Poorly	V. Poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	-	sl, ls, c (black)	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	>60
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10-

Table 7.25 Land suitability criteria for Jamun

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well	Mod. well	Poorly	V.Poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c(red)	sl, c (black)	ls	-
	pH	1:2.5	6.0-7.8	5.0-6.0	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>150	100-150	50-100	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	>60
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

Table 7.26 Land suitability criteria for Custard apple

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	Scl, cl, sc, c (red), c (black)	-	Sl, ls	-
	pH	1:2.5	6.0-7.3	5.5-6.0 7.3-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15-35	35-60	60-80	-
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	>5	-

Table 7.27 Land suitability criteria for Tamarind

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod.well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl,sc, c (red)	sl, c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>150	100-150	75-100	<75
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

Table 7.28 Land suitability criteria for Mulberry

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	24–28	22–24; 28–32	32–38; 22–18	>38; <18
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V. Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, cl, scl	c (red)	c (black), sl, ls	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.8-8.4	7.3-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	0-35	35-60	60-80	>80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

Note: Suitability evaluation only for Mulberry leaf not for Silk worm rearing

Table 7.29 Land suitability criteria for Marigold

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	18-23	17-15 24-35	35-40 10-14	>40 <10
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl,scl, cl, sc, c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%				
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.30 Land suitability criteria for Chrysanthemum

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	18-23	17-15 24-35	35-40 10-14	>40 <10
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl,scl, cl, sc, c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%				
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.31 Land suitability criteria for Jasmine (irrigated)

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	18-23	17-15 24-35	35-40 10-14	-
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	sl	ls, c (black)	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%				
Erosion hazard	Slope	%	<3	3-5	5-10	>10

7.32 Land suitability criteria for Crossandra

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	-	Poorly to very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c(red)	sl,	c (black),ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	dS/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%				
Erosion hazard	Slope	%	<3	3-5	5-10	>10

7.32 Land Management Units (LMUs)

The 18 soil map units identified in Kesalapura-1 Microwatershed have been grouped into 5 Land Management Units (LMUs) for the purpose of preparing a Proposed Crop Plan. Land Management Units are grouped based on the similarities in respect of the type of soil, the depth of the soil, the surface soil texture, gravel content, AWC, slope, erosion etc. and a Land Management Units map (Fig. 7.32) has been generated. These Land Management Units are expected to behave similarly for a given level of management.

The map units that have been grouped into five Land Management Units along with brief description of soil and site characteristics are given below.

LMU	Soil map unit number	Mapping unit	Soil and site characteristics
1	436, 473,476	HLPcB2g1, KLRmA1, KLRhB2	Moderately deep to very deep sandy clay lowland soils
2	396	BGPmB1	Very deep, black calcareous clay soils
3	162, 167, 173, 174, 178, 198, 201, 288	BSRhB2g1, BSRiB2, CKMcB2g2, CKMhB1g1, CKMiB1, KMhB1g1, KMHiB2, RTRiB2	Moderately deep to very deep, red sandy clay to clay soils
4	123, 232,269	HDHhB2g1, BPRhB2g2, GDPiB2	Moderately deep to deep, red gravelly sandy clay to clay soils
5	75, 77, 452	MKHcB1g1, MKHcB2g1, LKRhB2g1	Moderately shallow, red gravelly loamy soils




Fig 7.32 Land Management Units map of Kesalapura-1 Microwatershed

7.33 Proposed Crop Plan for Kesalapura-1 Microwatershed

After assessing the land suitability for the 31 crops, the proposed crop plan has been prepared for the 5 identified LMUs by considering only the highly (Class S1) and moderately (Class S2) suitable lands for each of the 31 crops. The resultant proposed crop plan is presented in Table 7.33.

Table 7.33 Proposed Crop Plan for Kesalapura-1 Microwatershed

LMU	Soil Map Units	Survey Number	Soil characters	Field Crops	Horticulture Crops	Suitable Interventions
LMU 1 19 ha (3%)	436.HLPcB2g1 473.KLRmA1 476.KLRhB2	Kasalapura: 16	Moderately deep to very deep sandy clay lowland soils	Maize, sorghum, bajra, cotton	Fruit crops: Custard Apple, Amla, Musambi, Lime Vegetable crops: Brinjal, Tomato, Chillies, Drumstick, Coriander Flower crops: Marigold, Jasmine Chrysanthemum,	Providing proper drainage, addition of organic manures, green leaf manuring, suitable conservation practises
LMU 2 48 ha (8%)	396.BGPmB1	Hesrur : xx	Very deep, black calcareous clay soils	Maize, Sorghum, Sunflower, Cotton, Bengal gram, Safflower, Linseed, Bajra , Soyabean	Fruit crops: Mango, Sapota, Pomegranate, Jamun, Lime, Musambi, Tamarind, Amla, Custard apple Vegetables: Drumstick, Chilli, Coriander, Tomato, Bhendi Flowers: Marigold, Chrysanthemum, Crossandra, Jasmine	Application of FYM, Biofertilizers and micronutrients, drip irrigation, mulching, suitable soil and water conservation practises
LMU 3 290 ha (50%)	162.BSRhB2g1 167.BSRiB2 173.CKMcb2g2 174.CKMhB1g1 178.CKMiB1 198.KMHhB1g1 201.KMHhB2 288.RTRiB2	Hesrur : xx Kasalapura: 10,11,12,19,23,24,25,26,27,28,29,30, 31,45,46,47,48,58	Moderately deep to very deep, red sandy clay to clay soils	Maize, Sorghum, Sunflower, Bajra, Finger millet, Groundnut, Red gram, Cowpea, Field bean, Castor, Mulberry	Fruit crops: Pomegranate, Guava, Sapota, Jackfruit, Tamarind, Lime, Musambi, Amla, Custard apple Vegetable crops: Drumstick, Tomato, Bhendi, Chilli, Brinjal, Onion, Curry leaves Flower crops: Marigold, Chrysanthemum, Jasmine, Crossandra	Drip irrigation, mulching, suitable soil and water conservation practises (Crescent Bunding with Catch Pit etc)
LMU 4 108 ha (19%)	123.HDHhB2g1 232.BPRhB2g2 269.GDPIB2	Kasalapura: 7,8,9,13,14,15,17,20,21,22,33,34,36,40,41,42,43,44,49	Moderately deep to deep, red gravelly sandy clay to clay soils	Groundnut, Bajra, Horse gram, Castor, Mulberry	Fruit crops: Musambi, Lime, Jamun, Jackfruit Amla, Custard apple, Tamarind Vegetable crops: Drumstick, Curry leaves	Drip irrigation, mulching, suitable soil and water conservation practises (Crescent Bunding with Catch Pit etc)
LMU 5 88 ha (15%)	75.MKHcB1g1 77.MKHcB2g1 452.LKRhB2g1	Halavagali : 53 Kasalapura: 32,37,38,39,50,51,52,53,54,55,56,57	Moderately shallow, red gravelly loamy soils	Groundnut, Bajra, Castor	Fruit crops: Lime, Musambi, Amla, Cashew, Custard apple,	Drip irrigation, mulching, suitable soil and water conservation practises (Crescent Bunding with Catch Pit etc)

SOIL HEALTH MANAGEMENT

8.1 Soil Health

Soil health is basic to plant health and plant health is basic to human and bovine health. Soil is fundamental to crop production. Without soil, no food could be produced nor would livestock be fed on a large scale. Because it is finite and fragile, soil is a precious resource that requires special care from its users.

Soil health or the capacity of the soil to function is critical to human survival. Soil health has been defined as: “the capacity of the soil to function as a living system without adverse effect on the ecosystem”. Healthy soils maintain a diverse community of soil organisms that help to form beneficial symbiotic associations with plant roots, recycle essential plant nutrients, improve soil structure with positive repercussions for soil, water and nutrient holding capacity and ultimately improve crop production and also contribute to mitigating climate change by maintaining or increasing its carbon content.

Functional interactions of soil biota with organic and inorganic components, air and water determine a soil’s potential to store and release nutrients, and water to plants and to promote and sustain plant growth. Thus, maintaining soil health is vital to crop production and conserve soil resource base for sustaining agriculture.

The most important characteristics of a healthy soil are

- Good soil tilth
- Sufficient soil depth
- Good water storage and good drainage
- Adequate supply, but not excess of nutrients
- Large population of beneficial organisms
- Small proportion of plant pathogens and insect pests
- Low weed pressure
- Free of chemicals and toxins that may harm the crop
- Resistance to degradation
- Resilience when unfavourable conditions occur

Characteristics of Kesalapura-1 Microwatershed

- ❖ The soil phases with sizeable area identified in the microwatershed belonged to the soil series of Chikkamegheri (CKM) 140 (24%), Kumchahalli (KMH) 62 ha (11%), Bisarahalli (BSR) 52 ha (9%), Giddadapalya (GDP) 48 ha (8%), Budagumpa (BGP) 48 ha (8%), Hooradhahalli (HDH) 47 ha (8%), Ranatur (RTR) 38 ha (7%), Mukhadahalli (MKH) 32 ha (11%), Lakkur (LKR) 26 ha (5%), Balapur (BPR) 12 ha (2%), Kavalakkeri (KLR) 10 ha (2%) and Huliyaपुरa (HLP) occupy an area of about 9 ha (2%) in the microwatershed.

- ❖ As per land capability classification, entire area in the microwatershed falls under arable land category (Class II & III). The major limitations identified in the arable lands were soil, wetness and erosion.
- ❖ On the basis of soil reaction, a major area of 353 ha (61%) is moderately alkaline (pH 7.8-8.4), 168 ha (29%) is strongly alkaline (pH 8.4-9.0) and about 31 ha (5%) is very strongly alkaline (pH >9.0) in the microwatershed. Entire area in the microwatershed is alkaline in reaction.

Soil Health Management

The following actions are required to improve the current land husbandry practices that provide a sound basis for the successful adoption of sustainable crop production system.

Alkaline soils

Moderately to very strongly alkaline soils cover an entire cultivated area of 552 ha.

1. Regular addition of organic manure, green manuring, green leaf manuring, crop residue incorporation and mulching needs to be taken up to improve the soil organic matter status.
2. Application of biofertilizers (Azospirillum, Azotobacter, Rhizobium).
3. Application of 25% extra N and P (125 % RDN&P).
4. Application of ZnSO₄ – 12.5 kg/ha (once in three years).
5. Application of Boron – 5 kg/ha (once in three years).

Soil Degradation

Soil erosion is one of the major factor affecting the soil health in the microwatershed. Out of total 575 ha area in the microwatershed, an area of about 211 ha (37%) is suffering from slight erosion and 341 ha (59%) is suffering from moderate erosion. The areas suffering from moderate erosion need immediate soil and water conservation and, other land development and land husbandry practices for restoring soil health.

Dissemination of Information and Communication of Benefits

Any large scale implementation of soil health management requires that supporting information is made available widely, particularly through channels familiar to farmers and extension workers. Given the very high priority attached to soil health especially by the Central Government on issuing Soil-Health Cards to all the farmers, media outlets like Regional, State and National Newspapers, Radio and Dooradarshan programs in local languages but also modern information and communication technologies such as Cellular phones and the Internet, which can be much more effective in reaching the younger farmers.

Inputs for Net Planning (Saturation Plan) and Interventions needed

Net planning (Saturation plan) in IWMP is focusing on preparation of

1. Soil and Water Conservation Treatment Plans for each plot or farm.
2. Productivity enhancement measures/ interventions for existing crops/livestock/other farm enterprises.
3. Diversification of farming mainly with perennial horticultural crops and livestock.
4. Improving livelihood opportunities and income generating activities.

In this connection, how various outputs of Sujala-III are of use in addressing these objectives of Net Planning are briefly presented below.

- ❖ **Soil Depth:** The depth of a soil decides the amount of moisture and nutrients it can hold, what crops can be taken up or not, depending on the rooting depth and the length of growing period available for raising any crop. Deeper the soil, better for a wide variety of crops. If sufficient depth is not available for growing deep rooted crops, either choose medium or short duration crops or deeper planting pits need to be opened and additional good quality soil brought from outside has to be filled into the planting pits.
- ❖ **Surface Soil Texture:** Lighter soil texture in the top soil means, better rain water infiltration, less run-off and soil moisture conservation, less capillary rise and less evaporation losses. Lighter surface textured soils are amenable to good soil tilth and are highly suitable for crops like groundnut, root vegetables (carrot, raddish, potato etc) but not ideal for crops that need stagnant water like lowland paddy. Heavy textured soils are poor in water infiltration and percolation. They are prone for sheet erosion; such soils can be improved by sand mulching. The technology that is developed by the AICRP-Dryland Agriculture, Vijayapura, Karnataka can be adopted.
- ❖ **Gravelliness:** More gravel content is favorable for run-off harvesting but poor in soil moisture storage and nutrient availability. It is a significant parameter that decides the kind of crop to be raised.
- ❖ **Land Capability Classification:** The land capability map shows the areas suitable and not suitable for agriculture and the major constraints in each of the plot/survey number. Hence, one can decide what kind of enterprise is possible in each of these units. In general, erosion and soil are the major constraints in Kesalapura-1 Microwatershed.
- ❖ **Organic Carbon:** The OC content (an index of available Nitrogen) is low (<0.5%) in 103 ha (18%), medium (0.5-0.75%) in 403 ha (70%) and high (>0.75%) in 46 ha (8%). The areas that are low and medium in OC needs to be further improved by applying farmyard manure and rotating crops with cereals and legumes or mixed cropping.
- ❖ **Promoting green manuring:** Growing of green manuring crops costs Rs. 1250/ha (green manuring seeds) and about Rs. 2000/ha towards cultivation that totals to Rs. 3250/- per ha. On the other hand, application of organic manure @ 10 tons/ha costs

Rs. 5000/ha. The practice needs to be continued for 2-3 years or more. Nitrogen fertilizer needs to be supplemented by 25% in addition to the recommended level in the area where OC is low and medium. For example, for rainfed maize, recommended level is 50 kg N per ha and an additional 12 kg /ha needs to be applied for all the crops grown in these plots.


- ❖ **Available Phosphorus:** Entire area of about 552 ha (96%) is medium (23-57 kg/ha) in available phosphorus. Hence for all the crops, 25% additional P-needs to be applied where it is medium.
- ❖ **Available Potassium:** Available potassium is medium (145-337 kg/ha) in 549 ha (96%) and high (>337 kg/ha) in 3 (1%) in the microwatershed. Additional 25% potassium needs to be applied in areas where it is medium.
- ❖ **Available Sulphur:** Available sulphur is a very critical nutrient for oilseed crops. Available sulphur is low (<10 ppm) in 543 ha (94%) and medium (10-20 ppm) in 10 ha (2%) in the microwatershed. These areas need to be applied with magnesium sulphate or gypsum or Factamphos (p) fertilizer (13% sulphur) for 2-3 years for the deficiency to be corrected.
- ❖ **Available Boron:** An area of about 12 ha (2%) is low (<0.5 ppm) and 473 ha (82%) is medium (0.5-1.0 ppm) in available boron content. The areas that are low and medium need to be applied with sodium borate @ 10 kg/ha as soil application or 0.2% borax as foliar spray to correct the deficiency. It is high (>1.0 ppm) in 67 ha (12%) in the microwatershed.
- ❖ **Available Iron:** Entire area of 552 ha (96%) is deficient (<4.5 ppm) in available iron in the microwatershed. To manage iron deficiency, iron sulphate@25 kg/ha needs to be applied for 2-3 years in the deficient areas.
- ❖ **Available Manganese:** Entire area in the microwatershed is sufficient (>1.0 ppm) in available manganese.
- ❖ **Available Copper:** Entire area is sufficient (>0.2 ppm) in available copper in the microwatershed.
- ❖ **Available Zinc:** An area of 33 ha (8%) is deficient (<0.6 ppm) in available zinc content in the microwatershed. Application of zinc sulphate @ 25 kg/ha is to be followed in areas that are deficient in available zinc. It is sufficient (>0.6 ppm) in an area of 520 ha (90%) in the microwatershed.
- ❖ **Soil Alkalinity:** Entire area of the microwatershed has 552 ha (96%) soils that are moderately to very strongly alkaline. These areas need application of gypsum and wherever calcium is in excess, iron pyrites and element sulphur can be recommended. Management practices like treating repeatedly with good quality water to drain out the excess salts and provision of subsurface drainage and growing of salt tolerant crops like Casuarina, Acasia, Neem, Ber etc, are recommended.

- ❖ **Land suitability for various crops:** Areas that are highly, moderately and marginally suitable and not suitable for growing various crops are indicated. Along with the suitability, various constraints that are limiting the productivity are also indicated. For example, in case of cotton, gravel content, rooting depth and salinity/alkalinity are the major constraints in various plots. With suitable management interventions, the productivity can be enhanced. In order to increase water holding capacity of light textured soils, growing of green manure crops and application of organic manure is recommended.

SOIL AND WATER CONSERVATION TREATMENT PLAN

For preparing soil and water conservation treatment plan for Kesalapura-1 Microwatershed, the land resource inventory database generated under Sujala-III project has been transformed as information through series of interpretative (thematic) maps using soil phase map as a base. The various thematic maps (1:7920 scale) generated were

- Soil depth
 - Surface soil texture
 - Available water capacity
 - Soil slope
 - Soil gravelliness
 - Land capability
 - Present land use and land cover
 - Crop suitability maps
 - Rainfall map
 - Hydrology
 - Water Resources
 - Socio-economic data
 - Contour plan with existing features- network of waterways, pottissa boundaries, cut up/ minor terraces etc.
 - Cadastral map (1:7920 scale)
 - Satellite imagery (1:7920 scale)
- Apart from these, Hand Level/ Hydro Marker/ Dumpy Level/ Total Station and Kathedars' List to be collected.



Steps for Survey and Preparation of Treatment Plan

The boundaries of Land User Groups' and Survey No. boundaries are traced in the field.

- Naming of user groups and farmers
- Identification of arable and non arable lands
- Identification of drainage lines and gullies
- Identification of non treatable areas
- Identification of priority areas in the arable lands
- Treatment plan for arable lands
- Location of water harvesting and recharge structures

9.1 Treatment Plan

The treatment plan recommended for arable lands is briefly described below.


9.1.1 Arable Land Treatment

A. BUNDING

Steps for Survey and Preparation of Treatment Plan		USER GROUP-1
Cadastral map (1:7920 scale) is enlarged to a scale of 1:2500 scale		
Existing network of waterways, pottissa boundaries, grass belts, natural drainage lines/ watercourse, cut ups/ terraces are marked on the cadastral map to the scale		
Drainage lines are demarcated into		
Small gullies	(up to 5 ha catchment)	<p>CLASSIFICATION OF GULLIES</p> <p>ಕೊಳೆಕಾಲಿನ ವರ್ಗೀಕರಣ</p> <p>UPPER REACH: 15 Ha.</p> <p>MIDDLE REACH: 15+10=25 ಹೆ.</p> <p>LOWER REACH: 25 ಹೆಕ್ಟೇರ್ ಗಿಂತ ಅಧಿಕ</p> <p>POINT OF CONCENTRATION</p>
Medium gullies	(5-15 ha catchment)	
Ravines	(15-25 ha catchment) and	
Halla/Nala	(more than 25ha catchment)	

Measurement of Land Slope

Land slope is estimated or determined by the study and interpretation of contours or by measurement in the field using simple instruments like Hand level or Hydromarker.



Vertical and Horizontal intervals between bunds as recommended by the Watershed Development Department.

Slope percentage	Vertical interval (m)	Corresponding Horizontal Distance (m)
2 - 3%	0.6	24
3 - 4%	0.9	21
4 - 5%	0.9	21
5 - 6%	1.2	21
6 - 7%	1.2	21

Note: i) The above intervals are maximum.

(ii) Considering the slope class and erosion status (A1... A= 0-1% slope, 1= slight erosion) the intervals have to be decided.

Bund length recording: Considering the contour plan and the existing grass belts/partitions, the bunds are aligned and lengths are measured.

Section of the Bund

Bund section is decided considering the soil texture class and gravelliness class ($bg_0 \dots b$ = loamy sand, $g_0 = <15\%$ gravel). The recommended sections for different soils are given below.

Recommended Bund Section



Top width (m)	Base width (m)	Height (m)	Side slope (Z:1;H:V)	Cross section (sq m)	Soil Texture	Remarks
0.3	0.9	0.3	01:01	0.18	Sandy loam	Vegetative bund
0.3	1.2	0.3	1.5:1	0.225	Sandy clay	
0.3	1.2	0.5	0.9:1	0.375	Red gravelly soils	
0.3	1.2	0.6	0.75:1	0.45		
0.3	1.5	0.6	01:01	0.54	Red sandy loam	
0.3	2.1	0.6	1.5:1	0.72	Very shallow black clayey soils	
0.45	2	0.75	01:01	0.92		
0.45	2.4	0.75	1.3:1	1.07	Shallow black clayey soils	
0.6	3.1	0.7	1.78:1	1.29	Medium black clayey soils	
0.5	3	0.85	1.47:1	1.49		

Formation of Trench cum Bund

Dimensions of the Borrow Pits/ Trenches to be excavated (machinery are decided considering the Bund Section).


Details of Borrow Pit dimensions are given below

TRENCH CUM BUND

IDEAL FOR HORTICULTURE CROPS

'A' FRAME FOR INTERBUND MANAGEMENT



1. ಸಮಸಾಕಳ ಉಳುವು
2. ಸಮಸಾಕಳ ಬಿತ್ತನೆ/ನಾಟ

ಸಮಸಾಕಳ ರೀತಿ

Size of Borrow Pits/ Trench recommended for Trench cum Bund (by machinery)

Bund section	Bund length	Earth quantity	Pit				Berm (pit to pit)	Soil depth Class
			L(m)	W(m)	D(m)	QUANTITY (m ³)		
m ²	m	m ³	L(m)	W(m)	D(m)	QUANTITY (m ³)	m	
0.375	6	2.25	5.85	0.85	0.45	2.24	0.15	Shallow
0.45	6	2.7	5.4	1.2	0.43	2.79	0.6	Shallow
0.45	6	2.7	5	0.85	0.65	2.76	1	Moderately Shallow
0.54	5.6	3.02	5.5	0.85	0.7	3.27	0.1	Moderately shallow
0.54	5.5	2.97	5	1.2	0.5	3	0.5	Shallow
0.72	6.2	4.46	6	1.2	0.7	5.04	0.2	Moderately shallow
0.72	5.2	3.74	5.1	0.85	0.9	3.9	0.1	Moderately deep

B. Waterways

- a) Existing waterways are marked on the cadastral map (1:7920 scale) and their dimensions are recorded.
- b) Considering the contour plan of the MWS, additional waterways/ modernization of the existing ones can be thought of.
- c) The design details are given in the Manual.

C. Farm Ponds

Waterways and the catchment area will give an indication on the size of the Farm Pond. Location of the pond can be decided based on the contour plan/ field condition and farmers' need/desire.

D. Diversion channel

Existing EPT/ CPT are marked on the cadastral map. Looking to the need, these can be modernized or fresh diversion channel can be proposed and runoff from this can be stored in *Gokatte/* Recharge ponds.

9.1.2 Non-Arable Land Treatment

Depending on the gravelliness and crops preferred by the farmers, the concerned authorities can decide appropriate treatment plan. The recommended treatments may be Contour Trench, Staggered Trench, Crescent Bund, Boulder Bund or Pebble Bund.

9.1.3 Treatment of Natural Water Course/ Drainage Lines

- a) The cadastral map has to be updated as regards the network of drainage lines (gullies/ *nalas/hallas*) and existing structures are marked to the scale and storage capacity of the existing water bodies are documented.
- b) The drainage line will be demarcated into Upper Reach, Middle Reach and Lower Reach.
- c) Considering the Catchment, *Nala* bed and bank conditions, suitable structures are decided.
- d) Number of storage structures (Check dam/ *Nala* bund/ Percolation tank) will be decided considering the commitments and available runoff in water budgeting and quality of water in the wells and site suitability.
- e) Detailed Leveling Survey using Dumpy Level / Total Station has to be carried out to arrive at the site-specific designs as shown in the Manual.
- f) The location of ground water recharge structures are decided by examining the lineaments and fracture zones from geological maps.
- g) Rainfall intensity data of the nearest Rain Gauge Station is considered for Hydrologic Designs.
- h) Silt load to the Storage/Recharge Structures is reduced by providing vegetative, boulder and earthen checks in the natural water course. Location and design details are given in the Manual.

9.2 Recommended Soil and Water Conservation Measures

The appropriate conservation structures best suited for each of the land parcel/ survey number (Appendix-I) are selected based on the slope per cent, severity of erosion, amount of rainfall, land use and soil type. The different kinds of conservation structures recommended are

1. Graded / Strengthening of Bunds
2. Trench cum Bunds (TCB)
3. Trench cum Bunds / Strengthening
4. Crescent Bunds

A map (Fig. 9.1) showing soil and water conservation plan with different kinds of structures recommended has been prepared which shows the spatial distribution and extent of area. An area of about 486 ha (85%) requires Trench cum Bunding, about 63 ha (11%) area requires Graded Bunding and 4ha (1%) requires Strengthening of existing bunds in the microwatershed. The conservation plan prepared may be presented to all the stakeholders including farmers and after including their suggestions, the conservation plan for the microwatershed may be finalised in a participatory approach.




Fig. 9.1 Soil and Water Conservation Plan map of Kesalapura-1 Microwatershed

Greening of Microwatershed

As part of the greening programme in the watersheds, it is envisaged to plant a variety of horticultural and other tree plants that are edible, economical and produce lot of biomass which helps to restore the ecological balance in the watersheds. The lands that are suitable for greening programme are non-arable lands (land capability classes V, VI VII and VIII) and also the lands that are not suitable or marginally suitable for growing annual and perennial crops. The method of planting these trees is given below.

It is recommended to open the pits during the 1st week of March along the contour and heap the dug out soil on the lower side of the slope in order to harness the flowing water and facilitate weathering of soil in the pit. Exposure of soil in the pit also prevents spread of pests and diseases due to scorching sun rays. The pits should be filled with mixture of soil and organic manure during the second week of April and keep ready with sufficiently tall seedlings produced either in poly bags or in root trainer nurseries so that planting can be done during the 2nd or 3rd week of April depending on the rainfall.

The tree species suitable for the area considering rainfall, temperature and adaptability is listed below; waterlogged areas are recommended to be planted with species like Neral (*Syzizium cumini*) and Bamboo. Dry areas are to be planted with species like Honge, Bevu, Seetaphal etc.

Dry Deciduous Species			Temp (°C)	Rainfall (mm)
1.	Bevu	<i>Azadiracta indica</i>	21–32	400 –1,200
2.	Tapasi	<i>Holoptelia integrifolia</i>	20-30	500 - 1000
3.	Seetaphal	<i>Anona Squamosa</i>	20-40	400 - 1000
4.	Honge	<i>Pongamia pinnata</i>	20 -50	500–2,500
5.	Kamara	<i>Hardwickia binata</i>	25 -35	400 - 1000
6.	Bage	<i>Albezzia lebbek</i>	20 - 45	500 - 1000
7.	Ficus	<i>Ficus bengalensis</i>	20 - 50	500–2,500
8.	Sisso	<i>Dalbargia Sissoo</i>	20 - 50	500 -2000
9.	Ailanthus	<i>Ailanthus excelsa</i>	20 - 50	500 - 1000
10.	Hale	<i>Wrightia tinctoria</i>	25 - 45	500 - 1000
11.	Uded	<i>Steriospermum chelanoides</i>	25 - 45	500 -2000
12.	Dhupa	<i>Boswella Serrata</i>	20 - 40	500 - 2000
13.	Nelli	<i>Emblica Officinalis</i>	20 - 50	500 -1500
14.	Honne	<i>Pterocarpus marsupium</i>	20 - 40	500 - 2000
Moist Deciduous Species			Temp (°C)	Rainfall (mm)
15.	Teak	<i>Tectona grandis</i>	20 - 50	500-5000
16.	Nandi	<i>Legarstroemia lanceolata</i>	20 - 40	500 - 4000
17.	Honne	<i>Pterocarpus marsupium</i>	20 - 40	500 - 3000
18.	Mathi	<i>Terminalia alata</i>	20 -50	500 - 2000
19.	Shivane	<i>Gmelina arborea</i>	20 -50	500 -2000
20.	Kindal	<i>T.Paniculata</i>	20 - 40	500 - 1500
21.	Beete	<i>Dalbargia latifolia</i>	20 - 40	500 - 1500
22.	Tare	<i>T. belerica</i>	20 - 40	500 - 2000
23.	Bamboo	<i>Bambusa arundinasia</i>	20 - 40	500 - 2500
24.	Bamboo	<i>Dendrocalamus strictus</i>	20 – 40	500 – 2500
25.	Muthuga	<i>Butea monosperma</i>	20 - 40	400 - 1500
26.	Hippe	<i>Madhuca latifolia</i>	20 - 40	500 - 2000
27.	Sandal	<i>Santalum album</i>	20 - 50	400 - 1000
28.	Nelli	<i>Emblica officinalis</i>	20 - 40	500 - 2000
29.	Nerale	<i>Sizyium cumini</i>	20 - 40	500 - 2000
30.	Dhaman	<i>Grevia tilifolia</i>	20 - 40	500 - 2000
31.	Kaval	<i>Careya arborea</i>	20 - 40	500 - 2000
32.	Harada	<i>Terminalia chebula</i>	20 - 40	500 - 2000

References

1. FAO (1976) Framework for Land Evaluation, Food and Agriculture Organization, Rome.72 pp.
2. FAO (1983) Guidelines for Land Evaluation for Rainfed Agriculture, FAO, Rome, 237 pp.
3. IARI (1971) Soil Survey Manual, All India Soil and Land Use Survey Organization, IARI, New Delhi, 121 pp.
4. Katyal, J.C. and Rattan, R.K. (2003) Secondary and Micronutrients; Research Gap and future needs. Fert. News 48 (4); 9-20.
5. Naidu, L.G.K., Ramamurthy, V., Challa, O., Hegde, R. and Krishnan, P. (2006) Manual Soil Site Suitability Criteria for Major Crops, NBSS Publ. No. 129, NBSS &LUP, Nagpur, 118 pp.
6. Natarajan, A. and Dipak Sarkar (2010) Field Guide for Soil Survey, National Bureau of Soil Survey and Land Use Planning (ICAR), Nagpur, India.
7. Natarajan, A., Rajendra Hegde, Raj, J.N. and Shivananda Murthy, H.G. (2015) Implementation Manual for Sujala-III Project, Watershed Development Department, Bengaluru, Karnataka.
8. Sarma, V.A.K., Krishnan, P. and Budihal, S.L. (1987) Laboratory Manual, Tech. Bull. 23, NBSS &LUP, Nagpur.
9. Sehgal, J.L. (1990) Soil Resource Mapping of Different States of India; Why and How?, National Bureau of Soil Survey and Land Use Planning, Nagpur, 49 pp.
10. Shivaprasad, C.R., R.S. Reddy, J. Sehgal and M. Velayuthum (1998) Soils of Karnataka for Optimizing Land Use, NBSS Publ. No. 47b, NBSS & LUP, Nagpur, India.
11. Soil Survey Staff (2006) Keys to Soil Taxonomy, Tenth edition, U.S. Department of Agriculture/ NRCS, Washington DC, U.S.A.
12. Soil Survey Staff (2012) Soil Survey Manual, Handbook No. 18, USDA, Washington DC, USA.

Appendix I
Keslapur-1 (2Q2e) Microwatershed
Soil Phase Information

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Soil Gravelliness	Surface Soil Texture	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Kasalapura	7	3.32	GDPiB2	LMU-4	Deep (100-150 cm)	Non gravelly (<15%)	Sandy clay	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIes	Trench cum bunding
Kasalapura	8	5.77	GDPiB2	LMU-4	Deep (100-150 cm)	Non gravelly (<15%)	Sandy clay	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Sugarcane (Sc)	Not Available	IIes	Trench cum bunding
Kasalapura	9	4.74	GDPiB2	LMU-4	Deep (100-150 cm)	Non gravelly (<15%)	Sandy clay	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Sugarcane+Maize (Sc+Mz)	1 Borewell	IIes	Trench cum bunding
Kasalapura	10	3.06	KMHiB2	LMU-3	Deep (100-150 cm)	Non gravelly (<15%)	Sandy clay	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Sugarcane+Maize (Sc+Mz)	Not Available	Ile	Trench cum bunding
Kasalapura	11	2.5	KMHiB2	LMU-3	Deep (100-150 cm)	Non gravelly (<15%)	Sandy clay	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Sugarcane+Maize (Sc+Mz)	Not Available	Ile	Trench cum bunding
Kasalapura	12	5.28	KMHiB2	LMU-3	Deep (100-150 cm)	Non gravelly (<15%)	Sandy clay	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Maize+Sugarcane (Mz+Sc)	Not Available	Ile	Trench cum bunding
Kasalapura	13	5.04	HDHhB2g1	LMU-4	Moderately deep (75-100 cm)	Gravelly (15-35%)	Sandy clay loam	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Sugarcane+Maize+Current fallow (Sc+Mz+Cf)	Not Available	Iles	Trench cum bunding
Kasalapura	14	6.12	HDHhB2g1	LMU-4	Moderately deep (75-100 cm)	Gravelly (15-35%)	Sandy clay loam	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Sugarcane (Sc)	Not Available	Iles	Trench cum bunding
Kasalapura	15	4.48	HDHhB2g1	LMU-4	Moderately deep (75-100 cm)	Gravelly (15-35%)	Sandy clay loam	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Maize+Cotton (Mz+Ct)	1 Borewell	Iles	Trench cum bunding
Kasalapura	16	5.93	KLRhB2	LMU-1	Very deep (>150 cm)	Non gravelly (<15%)	Sandy clay loam	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow+Maize (Cf+Mz)	Not Available	Ilew	Graded bunding
Kasalapura	17	6.63	HDHhB2g1	LMU-4	Moderately deep (75-100 cm)	Gravelly (15-35%)	Sandy clay loam	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow+Maize+Bajra (Cf+Mz+Bj)	2 Borewell	Iles	Trench cum bunding
Kasalapura	19	11.15	CKMiB1	LMU-3	Moderately deep (75-100 cm)	Non gravelly (<15%)	Sandy clay	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Slight	Sugarcane (Sc)	1 Borewell	IIs	Trench cum bunding
Kasalapura	20	5.53	HDHhB2g1	LMU-4	Moderately deep (75-100 cm)	Gravelly (15-35%)	Sandy clay loam	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Sugarcane+Maize (Sc+Mz)	1 Borewell	Iles	Trench cum bunding
Kasalapura	21	5.16	HDHhB2g1	LMU-4	Moderately deep (75-100 cm)	Gravelly (15-35%)	Sandy clay loam	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Sugarcane (Sc)	Not Available	Iles	Trench cum bunding
Kasalapura	22	6.47	HDHhB2g1	LMU-4	Moderately deep (75-100 cm)	Gravelly (15-35%)	Sandy clay loam	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Sugarcane (Sc)	Not Available	Iles	Trench cum bunding
Kasalapura	23	6.53	KMHiB2	LMU-3	Deep (100-150 cm)	Non gravelly (<15%)	Sandy clay	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Sugarcane+Cotton+Current fallow (Sc+Ct+Cf)	Not Available	Ile	Trench cum bunding
Kasalapura	24	8.06	KMHiB2	LMU-3	Deep (100-150 cm)	Non gravelly (<15%)	Sandy clay	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Sugarcane+Maize (Sc+Mz)	Not Available	Ile	Trench cum bunding
Kasalapura	25	7.13	CKMiB1	LMU-3	Moderately deep (75-100 cm)	Non gravelly (<15%)	Sandy clay	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Slight	Sugarcane+Maize (Sc+Mz)	1 Farm Pond	IIs	Trench cum bunding
Kasalapura	26	7.85	CKMiB1	LMU-3	Moderately deep (75-100 cm)	Non gravelly (<15%)	Sandy clay	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Slight	Sugarcane+Maize+Current fallow (Sc+Mz+Cf)	2 Borewell	IIs	Trench cum bunding
Kasalapura	27	4.78	CKMiB1	LMU-3	Moderately deep (75-100 cm)	Non gravelly (<15%)	Sandy clay	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Slight	Bajra+Jowar+Maize (Bj+Jw+Mz)	Not Available	IIs	Trench cum bunding

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Soil Gravelliness	Surface Soil Texture	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Kasalapura	28	0.11	CKMiB1	LMU-3	Moderately deep (75-100 cm)	Non gravelly (<15%)	Sandy clay	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Slight	Current fallow (Cf)	Not Available	IIs	Trench cum bunding
Kasalapura	29	0.54	CKMiB1	LMU-3	Moderately deep (75-100 cm)	Non gravelly (<15%)	Sandy clay	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Slight	Current fallow (Cf)	Not Available	IIs	Trench cum bunding
Kasalapura	30	2.97	CKMiB1	LMU-3	Moderately deep (75-100 cm)	Non gravelly (<15%)	Sandy clay	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Slight	Sugarcane+Maize (Sc+Mz)	1 Borewell	IIs	Trench cum bunding
Kasalapura	31	6.34	CKMiB1	LMU-3	Moderately deep (75-100 cm)	Non gravelly (<15%)	Sandy clay	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Slight	Maize+Bajra+Current fallow (Mz+Bj+Cf)	2 Borewell	IIs	Trench cum bunding
Kasalapura	32	6.22	MKHcB2g1	LMU-5	Moderately shallow (50-75 cm)	Gravelly (15-35%)	Sandy loam	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow+Habitation (Cf+Hb)	Not Available	IIIs	Trench cum bunding
Kasalapura	33	5.16	GDPiB2	LMU-4	Deep (100-150 cm)	Non gravelly (<15%)	Sandy clay	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Maize+Sugarcane+Cotton (Mz+Sc+Ct)	1 Borewell	IIIs	Trench cum bunding
Kasalapura	34	5.93	GDPiB2	LMU-4	Deep (100-150 cm)	Non gravelly (<15%)	Sandy clay	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Maize+Sugarcane+Cotton (Mz+Sc+Ct)	2 Borewell	IIIs	Trench cum bunding
Kasalapura	35	5.07	Habitation	Others	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others
Kasalapura	36	3.75	GDPiB2	LMU-4	Deep (100-150 cm)	Non gravelly (<15%)	Sandy clay	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Maize+Drumstick (Mz+Ds)	Not Available	IIIs	Trench cum bunding
Kasalapura	37	4.22	MKHcB2g1	LMU-5	Moderately shallow (50-75 cm)	Gravelly (15-35%)	Sandy loam	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Sugarcane+Maize (Sc+Mz)	Not Available	IIIs	Trench cum bunding
Kasalapura	38	1.89	MKHcB2g1	LMU-5	Moderately shallow (50-75 cm)	Gravelly (15-35%)	Sandy loam	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Sugarcane+Maize (Sc+Mz)	1 Borewell	IIIs	Trench cum bunding
Kasalapura	39	4.96	MKHcB2g1	LMU-5	Moderately shallow (50-75 cm)	Gravelly (15-35%)	Sandy loam	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	1 Borewell	IIIs	Trench cum bunding
Kasalapura	40	3.95	GDPiB2	LMU-4	Deep (100-150 cm)	Non gravelly (<15%)	Sandy clay	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Bajra (Bj)	Not Available	IIIs	Trench cum bunding
Kasalapura	41	0.04	GDPiB2	LMU-4	Deep (100-150 cm)	Non gravelly (<15%)	Sandy clay	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIIs	Trench cum bunding
Kasalapura	42	0.48	BPRhB2g2	LMU-4	Deep (100-150 cm)	Very gravelly (35-60%)	Sandy clay loam	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIIs	Trench cum bunding
Kasalapura	43	0.96	BPRhB2g2	LMU-4	Deep (100-150 cm)	Very gravelly (35-60%)	Sandy clay loam	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIIs	Trench cum bunding
Kasalapura	44	3.94	BPRhB2g2	LMU-4	Deep (100-150 cm)	Very gravelly (35-60%)	Sandy clay loam	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Maize+Bajra (Mz+Bj)	Not Available	IIIs	Trench cum bunding
Kasalapura	45	5.62	KMHiB2	LMU-3	Deep (100-150 cm)	Non gravelly (<15%)	Sandy clay	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow (Cf)	Not Available	Ile	Trench cum bunding
Kasalapura	46	2.45	KMHiB2	LMU-3	Deep (100-150 cm)	Non gravelly (<15%)	Sandy clay	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow (Cf)	Not Available	Ile	Trench cum bunding
Kasalapura	47	2.38	KMHiB2	LMU-3	Deep (100-150 cm)	Non gravelly (<15%)	Sandy clay	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Bajra (Bj)	1 Borewell	Ile	Trench cum bunding
Kasalapura	48	4.39	KMHiB2	LMU-3	Deep (100-150 cm)	Non gravelly (<15%)	Sandy clay	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Current fallow (Cf)	Not Available	Ile	Trench cum bunding
Kasalapura	49	2.08	BPRhB2g2	LMU-4	Deep (100-150 cm)	Very gravelly (35-60%)	Sandy clay loam	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Bajra (Bj)	Not Available	IIIs	Trench cum bunding
Kasalapura	50	1.33	LKRhB2g1	LMU-5	Moderately shallow (50-75 cm)	Gravelly (15-35%)	Sandy clay loam	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Maize (Mz)	Not Available	IIIs	Trench cum bunding
Kasalapura	51	1.29	LKRhB2g1	LMU-5	Moderately shallow (50-75 cm)	Gravelly (15-35%)	Sandy clay loam	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Sugarcane (Sc)	Not Available	IIIs	Trench cum bunding

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Soil Gravelliness	Surface Soil Texture	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Kasalapura	52	0.99	LKRhB2g1	LMU-5	Moderately shallow (50-75 cm)	Gravelly (15-35%)	Sandy clay loam	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Sugarcane (Sc)	Not Available	IIIes	Trench cum bunding
Kasalapura	53	1.27	LKRhB2g1	LMU-5	Moderately shallow (50-75 cm)	Gravelly (15-35%)	Sandy clay loam	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Sugarcane (Sc)	Not Available	IIIes	Trench cum bunding
Kasalapura	54	0.87	LKRhB2g1	LMU-5	Moderately shallow (50-75 cm)	Gravelly (15-35%)	Sandy clay loam	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Sugarcane (Sc)	Not Available	IIIes	Trench cum bunding
Kasalapura	55	0.06	LKRhB2g1	LMU-5	Moderately shallow (50-75 cm)	Gravelly (15-35%)	Sandy clay loam	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIIes	Trench cum bunding
Kasalapura	56	0.27	LKRhB2g1	LMU-5	Moderately shallow (50-75 cm)	Gravelly (15-35%)	Sandy clay loam	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Sugarcane (Sc)	Not Available	IIIes	Trench cum bunding
Kasalapura	57	7.5	LKRhB2g1	LMU-5	Moderately shallow (50-75 cm)	Gravelly (15-35%)	Sandy clay loam	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Maize+Mulberry+Cotton (Mz+Mu+Ct)	1 Borewell	IIIes	Trench cum bunding
Kasalapura	58	35.64	KMHhB1g1	LMU-3	Deep (100-150 cm)	Gravelly (15-35%)	Sandy clay loam	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Slight	Maize+Bajra+Sugarcane (Mz+Bj+Sc)	2 Borewell	IIs	Trench cum bunding
Halavagali	53	51.72	LKRhB2g1	LMU-5	Moderately shallow (50-75 cm)	Gravelly (15-35%)	Sandy clay loam	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Waterbody	Not Available	IIIes	Trench cum bunding
Hesaru	38	1.02	BGPmB1	LMU-2	Very deep (>150 cm)	Non gravelly (<15%)	Clay	High (151-200 mm/m)	Very gently sloping (1-3%)	Slight	Not Available (NA)	Not Available	IIs	Graded bunding
Hesaru	39	2.91	BGPmB1	LMU-2	Very deep (>150 cm)	Non gravelly (<15%)	Clay	High (151-200 mm/m)	Very gently sloping (1-3%)	Slight	Not Available (NA)	Not Available	IIs	Graded bunding
Hesaru	40	2.04	BGPmB1	LMU-2	Very deep (>150 cm)	Non gravelly (<15%)	Clay	High (151-200 mm/m)	Very gently sloping (1-3%)	Slight	Not Available (NA)	Not Available	IIs	Graded bunding
Hesaru	41	0.36	BGPmB1	LMU-2	Very deep (>150 cm)	Non gravelly (<15%)	Clay	High (151-200 mm/m)	Very gently sloping (1-3%)	Slight	Not Available (NA)	Not Available	IIs	Graded bunding
Hesaru	42	4.26	RTRiB2	LMU-3	Very deep (>150 cm)	Non gravelly (<15%)	Sandy clay	High (151-200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIE	Trench cum bunding
Hesaru	86	1.98	RTRiB2	LMU-3	Very deep (>150 cm)	Non gravelly (<15%)	Sandy clay	High (151-200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIE	Trench cum bunding
Hesaru	88	2.93	MKHcB1g1	LMU-5	Moderately shallow (50-75 cm)	Gravelly (15-35%)	Sandy loam	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Slight	Not Available (NA)	Not Available	IIIIs	Trench cum bunding
Hesaru	89	1.62	MKHcB1g1	LMU-5	Moderately shallow (50-75 cm)	Gravelly (15-35%)	Sandy loam	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Slight	Not Available (NA)	Not Available	IIIIs	Trench cum bunding
Hesaru	90	2.95	BSRhB2g1	LMU-3	Moderately deep (75-100 cm)	Gravelly (15-35%)	Sandy clay loam	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIEs	Trench cum bunding
Hesaru	91	2.52	MKHcB1g1	LMU-5	Moderately shallow (50-75 cm)	Gravelly (15-35%)	Sandy loam	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Slight	Not Available (NA)	Not Available	IIIIs	Trench cum bunding
Hesaru	92	2.37	MKHcB1g1	LMU-5	Moderately shallow (50-75 cm)	Gravelly (15-35%)	Sandy loam	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Slight	Not Available (NA)	Not Available	IIIIs	Trench cum bunding
Hesaru	94	0.07	BSRhB2g1	LMU-3	Moderately deep (75-100 cm)	Gravelly (15-35%)	Sandy clay loam	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIEs	Trench cum bunding
Hesaru	95	3.91	BSRhB2g1	LMU-3	Moderately deep (75-100 cm)	Gravelly (15-35%)	Sandy clay loam	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIEs	Trench cum bunding
Hesaru	96	11.32	MKHcB1g1	LMU-5	Moderately shallow (50-75 cm)	Gravelly (15-35%)	Sandy loam	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Slight	Not Available (NA)	Not Available	IIIIs	Trench cum bunding
Hesaru	97	1.89	MKHcB1g1	LMU-5	Moderately shallow (50-75 cm)	Gravelly (15-35%)	Sandy loam	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Slight	Not Available (NA)	Not Available	IIIIs	Trench cum bunding
Hesaru	98	2.56	KLRmA1	LMU-1	Very deep (>150 cm)	Non gravelly (<15%)	Clay	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Not Available (NA)	Not Available	IIw	Graded bunding

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Soil Gravelliness	Surface Soil Texture	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Hesaru	99	3.64	HDHhB2g1	LMU-4	Moderately deep (75-100 cm)	Gravelly (15-35%)	Sandy clay loam	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iies	Trench cum bunding
Hesaru	100	10.04	MKHcB1g1	LMU-5	Moderately shallow (50-75 cm)	Gravelly (15-35%)	Sandy loam	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Slight	Not Available (NA)	Not Available	IIIs	Trench cum bunding
Hesaru	101	9.78	MKHcB1g1	LMU-5	Moderately shallow (50-75 cm)	Gravelly (15-35%)	Sandy loam	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Slight	Not Available (NA)	Not Available	IIIs	Trench cum bunding
Hesaru	102	10.62	MKHcB1g1	LMU-5	Moderately shallow (50-75 cm)	Gravelly (15-35%)	Sandy loam	Very Low (<50 mm/m)	Very gently sloping (1-3%)	Slight	Not Available (NA)	Not Available	IIIs	Trench cum bunding
Hesaru	103	12.05	RTRiB2	LMU-3	Very deep (>150 cm)	Non gravelly (<15%)	Sandy clay	High (151-200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIE	Trench cum bunding
Hesaru	104	9.21	RTRiB2	LMU-3	Very deep (>150 cm)	Non gravelly (<15%)	Sandy clay	High (151-200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIE	Trench cum bunding
Hesaru	105	5.94	BSRhB2g1	LMU-3	Moderately deep (75-100 cm)	Gravelly (15-35%)	Sandy clay loam	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iies	Trench cum bunding
Hesaru	106	9.25	BSRhB2g1	LMU-3	Moderately deep (75-100 cm)	Gravelly (15-35%)	Sandy clay loam	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iies	Trench cum bunding
Hesaru	107	7.65	BSRhB2g1	LMU-3	Moderately deep (75-100 cm)	Gravelly (15-35%)	Sandy clay loam	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iies	Trench cum bunding
Hesaru	108	6.96	BSRhB2g1	LMU-3	Moderately deep (75-100 cm)	Gravelly (15-35%)	Sandy clay loam	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iies	Trench cum bunding
Hesaru	109	8.68	CKMcB2g2	LMU-3	Moderately deep (75-100 cm)	Very gravelly (35-60%)	Sandy loam	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iies	Trench cum bunding
Hesaru	110	3.56	RTRiB2	LMU-3	Very deep (>150 cm)	Non gravelly (<15%)	Sandy clay	High (151-200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIE	Trench cum bunding
Hesaru	111	8.19	CKMcB2g2	LMU-3	Moderately deep (75-100 cm)	Very gravelly (35-60%)	Sandy loam	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iies	Trench cum bunding
Hesaru	112	10.08	CKMcB2g2	LMU-3	Moderately deep (75-100 cm)	Very gravelly (35-60%)	Sandy loam	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iies	Trench cum bunding
Hesaru	113	9.2	CKMhB1g1	LMU-3	Moderately deep (75-100 cm)	Gravelly (15-35%)	Sandy clay loam	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Slight	Not Available (NA)	Not Available	IIs	Trench cum bunding
Hesaru	114	7.31	CKMhB1g1	LMU-3	Moderately deep (75-100 cm)	Gravelly (15-35%)	Sandy clay loam	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Slight	Not Available (NA)	Not Available	IIs	Trench cum bunding
Hesaru	115	6.02	CKMhB1g1	LMU-3	Moderately deep (75-100 cm)	Gravelly (15-35%)	Sandy clay loam	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Slight	Not Available (NA)	Not Available	IIs	Trench cum bunding
Hesaru	116	4.4	CKMhB1g1	LMU-3	Moderately deep (75-100 cm)	Gravelly (15-35%)	Sandy clay loam	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Slight	Not Available (NA)	Not Available	IIs	Trench cum bunding
Hesaru	117	7.67	CKMhB1g1	LMU-3	Moderately deep (75-100 cm)	Gravelly (15-35%)	Sandy clay loam	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Slight	Not Available (NA)	Not Available	IIs	Trench cum bunding
Hesaru	118	7.42	CKMhB1g1	LMU-3	Moderately deep (75-100 cm)	Gravelly (15-35%)	Sandy clay loam	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Slight	Not Available (NA)	Not Available	IIs	Trench cum bunding
Hesaru	119	7.76	CKMhB1g1	LMU-3	Moderately deep (75-100 cm)	Gravelly (15-35%)	Sandy clay loam	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Slight	Not Available (NA)	Not Available	IIs	Trench cum bunding
Hesaru	120	6.97	CKMhB1g1	LMU-3	Moderately deep (75-100 cm)	Gravelly (15-35%)	Sandy clay loam	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Slight	Not Available (NA)	Not Available	IIs	Trench cum bunding
Hesaru	121	7.12	BGPmB1	LMU-2	Very deep (>150 cm)	Non gravelly (<15%)	Clay	High (151-200 mm/m)	Very gently sloping (1-3%)	Slight	Not Available (NA)	Not Available	IIs	Graded bunding
Hesaru	122	9.62	BGPmB1	LMU-2	Very deep (>150 cm)	Non gravelly (<15%)	Clay	High (151-200 mm/m)	Very gently sloping (1-3%)	Slight	Not Available (NA)	Not Available	IIs	Graded bunding

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Soil Gravelliness	Surface Soil Texture	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Hesaru	123	3.02	CKMcB2g2	LMU-3	Moderately deep (75-100 cm)	Very gravelly (35-60%)	Sandy loam	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIs	Trench cum bunding
Hesaru	124/A	10.37	CKMcB2g2	LMU-3	Moderately deep (75-100 cm)	Very gravelly (35-60%)	Sandy loam	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIs	Trench cum bunding
Hesaru	125	8.78	BGPmB1	LMU-2	Very deep (>150 cm)	Non gravelly (<15%)	Clay	High (151-200 mm/m)	Very gently sloping (1-3%)	Slight	Not Available (NA)	Not Available	IIs	Graded bunding
Hesaru	126	10.21	CKMcB2g2	LMU-3	Moderately deep (75-100 cm)	Very gravelly (35-60%)	Sandy loam	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIs	Trench cum bunding
Hesaru	127	0.55	CKMcB2g2	LMU-3	Moderately deep (75-100 cm)	Very gravelly (35-60%)	Sandy loam	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIs	Trench cum bunding
Hesaru	142	1.37	BGPmB1	LMU-2	Very deep (>150 cm)	Non gravelly (<15%)	Clay	High (151-200 mm/m)	Very gently sloping (1-3%)	Slight	Not Available (NA)	Not Available	IIs	Graded bunding
Hesaru	143/A	6.44	BGPmB1	LMU-2	Very deep (>150 cm)	Non gravelly (<15%)	Clay	High (151-200 mm/m)	Very gently sloping (1-3%)	Slight	Not Available (NA)	Not Available	IIs	Graded bunding
Hesaru	143/B	0.13	BGPmB1	LMU-2	Very deep (>150 cm)	Non gravelly (<15%)	Clay	High (151-200 mm/m)	Very gently sloping (1-3%)	Slight	Not Available (NA)	Not Available	IIs	Graded bunding

Appendix III
Keslapur-1 (2Q2e) Microwatershed
Soil Suitability Information

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Jasmine	Bhendi	Brinjal	Crossandra	Drumstick	Mulberry	Onion	
Kasalapura	7	S1	S1	S1	S1	S2t	S1	S1	S1	S2t	S1	S1	S1	S1	S1	S2t	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S2tg	S2g	S1	S1	S1	S2tg	
Kasalapura	8	S1	S1	S1	S1	S2t	S1	S1	S1	S2t	S1	S1	S1	S1	S1	S2t	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S2tg	S2g	S1	S1	S1	S2tg	
Kasalapura	9	S1	S1	S1	S1	S2t	S1	S1	S1	S2t	S1	S1	S1	S1	S1	S2t	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S2tg	S2g	S1	S1	S1	S2tg	
Kasalapura	10	S2r	S1	S1	S1	S1	S2t	S2r	S1	S2t	S1	S1	S1	S1	S1	S1	S2r	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	
Kasalapura	11	S2r	S1	S1	S1	S1	S2t	S2r	S1	S2t	S1	S1	S1	S1	S1	S1	S2r	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	
Kasalapura	12	S2r	S1	S1	S1	S1	S2t	S2r	S1	S2t	S1	S1	S1	S1	S1	S1	S2r	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	
Kasalapura	13	S3rg	S3g	S2rg	S3g	S2rg	S3rg	S3rg	S2rg	S3g	S3rg	S3g	S2rg	S2rg	S2rg	S2rg	S2rg	S2rg	S2g	S3g	S3g	S3g	S3g	S2rg	S2g	S3g	S3g	S3g	S3g	S3g	S3g	S2g	S3g
Kasalapura	14	S3rg	S3g	S2rg	S3g	S2rg	S3rg	S3rg	S2rg	S3g	S3rg	S3g	S2rg	S2rg	S2rg	S2rg	S2rg	S2rg	S2g	S3g	S3g	S3g	S3g	S2rg	S2g	S3g	S3g	S3g	S3g	S3g	S3g	S2g	S3g
Kasalapura	15	S3rg	S3g	S2rg	S3g	S2rg	S3rg	S3rg	S2rg	S3g	S3rg	S3g	S2rg	S2rg	S2rg	S2rg	S2rg	S2rg	S2g	S3g	S3g	S3g	S3g	S2rg	S2g	S3g	S3g	S3g	S3g	S3g	S3g	S2g	S3g
Kasalapura	16	S3tz	S2tz	S3tz	S1	S3tz	S1	S2tz	S1	S1	S1	S2tz	S1	S3tz	S1	N1t	S2tz	S1	S3t	S2tz	S2tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S3tz	S3tz	
Kasalapura	17	S3rg	S3g	S2rg	S3g	S2rg	S3rg	S3rg	S2rg	S3g	S3rg	S3g	S2rg	S2rg	S2rg	S2rg	S2rg	S2rg	S2g	S3g	S3g	S3g	S3g	S2rg	S2g	S3g	S3g	S3g	S3g	S3g	S3g	S2g	S3g
Kasalapura	19	S3r	S2t	S2rg	S1	S2rt	S2r	S3r	S2r	S2t	S2r	S2r	S1	S2r	S1	S2rt	S2r	S2r	S2t	S1	S1	S2t	S2t	S2r	S1	S2t	S2t	S1	S2t	S2r	S2r	S2t	
Kasalapura	20	S3rg	S3g	S2rg	S3g	S2rg	S3rg	S3rg	S2rg	S3g	S3rg	S3g	S2rg	S2rg	S2rg	S2rg	S2rg	S2rg	S2g	S3g	S3g	S3g	S3g	S2rg	S2g	S3g	S3g	S3g	S3g	S3g	S3g	S2g	S3g
Kasalapura	21	S3rg	S3g	S2rg	S3g	S2rg	S3rg	S3rg	S2rg	S3g	S3rg	S3g	S2rg	S2rg	S2rg	S2rg	S2rg	S2rg	S2g	S3g	S3g	S3g	S3g	S2rg	S2g	S3g	S3g	S3g	S3g	S3g	S3g	S2g	S3g
Kasalapura	22	S3rg	S3g	S2rg	S3g	S2rg	S3rg	S3rg	S2rg	S3g	S3rg	S3g	S2rg	S2rg	S2rg	S2rg	S2rg	S2rg	S2g	S3g	S3g	S3g	S3g	S2rg	S2g	S3g	S3g	S3g	S3g	S3g	S3g	S2g	S3g
Kasalapura	23	S2r	S1	S1	S1	S1	S2t	S2r	S1	S2t	S1	S1	S1	S1	S1	S1	S2r	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1
Kasalapura	24	S2r	S1	S1	S1	S1	S2t	S2r	S1	S2t	S1	S1	S1	S1	S1	S1	S2r	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1
Kasalapura	25	S3r	S2t	S2rg	S1	S2rt	S2r	S3r	S2r	S2t	S2r	S2r	S1	S2r	S1	S2rt	S2r	S2r	S2t	S1	S1	S2t	S2t	S2r	S1	S2t	S2t	S1	S2t	S2r	S2r	S2t	
Kasalapura	26	S3r	S2t	S2rg	S1	S2rt	S2r	S3r	S2r	S2t	S2r	S2r	S1	S2r	S1	S2rt	S2r	S2r	S2t	S1	S1	S2t	S2t	S2r	S1	S2t	S2t	S1	S2t	S2r	S2r	S2t	
Kasalapura	27	S3r	S2t	S2rg	S1	S2rt	S2r	S3r	S2r	S2t	S2r	S2r	S1	S2r	S1	S2rt	S2r	S2r	S2t	S1	S1	S2t	S2t	S2r	S1	S2t	S2t	S1	S2t	S2r	S2r	S2t	
Kasalapura	28	S3r	S2t	S2rg	S1	S2rt	S2r	S3r	S2r	S2t	S2r	S2r	S1	S2r	S1	S2rt	S2r	S2r	S2t	S1	S1	S2t	S2t	S2r	S1	S2t	S2t	S1	S2t	S2r	S2r	S2t	
Kasalapura	29	S3r	S2t	S2rg	S1	S2rt	S2r	S3r	S2r	S2t	S2r	S2r	S1	S2r	S1	S2rt	S2r	S2r	S2t	S1	S1	S2t	S2t	S2r	S1	S2t	S2t	S1	S2t	S2r	S2r	S2t	
Kasalapura	30	S3r	S2t	S2rg	S1	S2rt	S2r	S3r	S2r	S2t	S2r	S2r	S1	S2r	S1	S2rt	S2r	S2r	S2t	S1	S1	S2t	S2t	S2r	S1	S2t	S2t	S1	S2t	S2r	S2r	S2t	
Kasalapura	31	S3r	S2t	S2rg	S1	S2rt	S2r	S3r	S2r	S2t	S2r	S2r	S1	S2r	S1	S2rt	S2r	S2r	S2t	S1	S1	S2t	S2t	S2r	S1	S2t	S2t	S1	S2t	S2r	S2r	S2t	

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Jasmine	Bhendi	Brinjal	Crossandra	Drumstick	Mulberry	Onion	
Kasalapura	32	N1r	S2rg	S3rg	S2rg	S3rg	S2rg	N1r	S3rg	S2rt	S3rg	S3rg	S2rg	S3rg	S2rg	S3rg	S3rg	S3rg	S2rg	S2rg	S2rg	S2rg	S2rg	S3rg	S2rg	S2rg	S3g	S3g	S2rg	S3rg	S3rg	S3g	
Kasalapura	33	S1	S1	S1	S1	S2t	S1	S1	S1	S2t	S1	S1	S1	S1	S1	S2t	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S1	S2tg	S2g	S1	S1	S1	S2tg
Kasalapura	34	S1	S1	S1	S1	S2t	S1	S1	S1	S2t	S1	S1	S1	S1	S1	S2t	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S1	S2tg	S2g	S1	S1	S1	S2tg
Kasalapura	35	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Kasalapura	36	S1	S1	S1	S1	S2t	S1	S1	S1	S2t	S1	S1	S1	S1	S1	S2t	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S1	S2tg	S2g	S1	S1	S1	S2tg
Kasalapura	37	N1r	S2rg	S3rg	S2rg	S3rg	S2rg	N1r	S3rg	S2rt	S3rg	S3rg	S2rg	S3rg	S2rg	S3rg	S3rg	S3rg	S2rg	S2rg	S2rg	S2rg	S2rg	S3rg	S2rg	S2rg	S3g	S3g	S2rg	S3rg	S3rg	S3g	
Kasalapura	38	N1r	S2rg	S3rg	S2rg	S3rg	S2rg	N1r	S3rg	S2rt	S3rg	S3rg	S2rg	S3rg	S2rg	S3rg	S3rg	S3rg	S2rg	S2rg	S2rg	S2rg	S2rg	S3rg	S2rg	S2rg	S3g	S3g	S2rg	S3rg	S3rg	S3g	
Kasalapura	39	N1r	S2rg	S3rg	S2rg	S3rg	S2rg	N1r	S3rg	S2rt	S3rg	S3rg	S2rg	S3rg	S2rg	S3rg	S3rg	S3rg	S2rg	S2rg	S2rg	S2rg	S2rg	S3rg	S2rg	S2rg	S3g	S3g	S2rg	S3rg	S3rg	S3g	
Kasalapura	40	S1	S1	S1	S1	S2t	S1	S1	S1	S2t	S1	S1	S1	S1	S1	S2t	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S1	S2tg	S2g	S1	S1	S1	S2tg
Kasalapura	41	S1	S1	S1	S1	S2t	S1	S1	S1	S2t	S1	S1	S1	S1	S1	S2t	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S1	S2tg	S2g	S1	S1	S1	S2tg
Kasalapura	42	S3rg	S3g	S3g	S3g	S3g	S3g	S3rg	S3g	S3g	S3g	S3g	S2g	S3g	S2g	S3g	S3g	S3g	S2g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S2g	S2g	S3g	S2g	S2g	S2g
Kasalapura	43	S3rg	S3g	S3g	S3g	S3g	S3g	S3rg	S3g	S3g	S3g	S3g	S2g	S3g	S2g	S3g	S3g	S3g	S2g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S2g	S2g	S3g	S2g	S2g	S2g
Kasalapura	44	S3rg	S3g	S3g	S3g	S3g	S3g	S3rg	S3g	S3g	S3g	S3g	S2g	S3g	S2g	S3g	S3g	S3g	S2g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S2g	S2g	S3g	S2g	S2g	S2g
Kasalapura	45	S2r	S1	S1	S1	S1	S2t	S2r	S1	S2t	S1	S1	S1	S1	S1	S1	S2r	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1
Kasalapura	46	S2r	S1	S1	S1	S1	S2t	S2r	S1	S2t	S1	S1	S1	S1	S1	S1	S2r	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1
Kasalapura	47	S2r	S1	S1	S1	S1	S2t	S2r	S1	S2t	S1	S1	S1	S1	S1	S1	S2r	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1
Kasalapura	48	S2r	S1	S1	S1	S1	S2t	S2r	S1	S2t	S1	S1	S1	S1	S1	S1	S2r	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1
Kasalapura	49	S3rg	S3g	S3g	S3g	S3g	S3g	S3rg	S3g	S3g	S3g	S3g	S2g	S3g	S2g	S3g	S3g	S3g	S2g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S3g	S2g	S2g	S3g	S2g	S2g	S2g
Kasalapura	50	N1rg	S3rg	S3rg	S3rg	S3rg	S3rg	N1rg	S3rg	S2rt	S3rg	S3rg	S2rg	S3rg	S2rg	S3rg	S3rg	S3rg	S2rg	S3rg	S3rg	S3rg	S3rg	S3rg	S3rg	S2rg	S3rg	S3g	S3g	S3rg	S3rg	S3rg	S3g
Kasalapura	51	N1rg	S3rg	S3rg	S3rg	S3rg	S3rg	N1rg	S3rg	S2rt	S3rg	S3rg	S2rg	S3rg	S2rg	S3rg	S3rg	S3rg	S2rg	S3rg	S3rg	S3rg	S3rg	S3rg	S3rg	S2rg	S3rg	S3g	S3g	S3rg	S3rg	S3rg	S3g
Kasalapura	52	N1rg	S3rg	S3rg	S3rg	S3rg	S3rg	N1rg	S3rg	S2rt	S3rg	S3rg	S2rg	S3rg	S2rg	S3rg	S3rg	S3rg	S2rg	S3rg	S3rg	S3rg	S3rg	S3rg	S3rg	S2rg	S3rg	S3g	S3g	S3rg	S3rg	S3rg	S3g
Kasalapura	53	N1rg	S3rg	S3rg	S3rg	S3rg	S3rg	N1rg	S3rg	S2rt	S3rg	S3rg	S2rg	S3rg	S2rg	S3rg	S3rg	S3rg	S2rg	S3rg	S3rg	S3rg	S3rg	S3rg	S3rg	S2rg	S3rg	S3g	S3g	S3rg	S3rg	S3rg	S3g
Kasalapura	54	N1rg	S3rg	S3rg	S3rg	S3rg	S3rg	N1rg	S3rg	S2rt	S3rg	S3rg	S2rg	S3rg	S2rg	S3rg	S3rg	S3rg	S2rg	S3rg	S3rg	S3rg	S3rg	S3rg	S3rg	S2rg	S3rg	S3g	S3g	S3rg	S3rg	S3rg	S3g
Kasalapura	55	N1rg	S3rg	S3rg	S3rg	S3rg	S3rg	N1rg	S3rg	S2rt	S3rg	S3rg	S2rg	S3rg	S2rg	S3rg	S3rg	S3rg	S2rg	S3rg	S3rg	S3rg	S3rg	S3rg	S3rg	S2rg	S3rg	S3g	S3g	S3rg	S3rg	S3rg	S3g
Kasalapura	56	N1rg	S3rg	S3rg	S3rg	S3rg	S3rg	N1rg	S3rg	S2rt	S3rg	S3rg	S2rg	S3rg	S2rg	S3rg	S3rg	S3rg	S2rg	S3rg	S3rg	S3rg	S3rg	S3rg	S3rg	S2rg	S3rg	S3g	S3g	S3rg	S3rg	S3rg	S3g
Kasalapura	57	N1rg	S3rg	S3rg	S3rg	S3rg	S3rg	N1rg	S3rg	S2rt	S3rg	S3rg	S2rg	S3rg	S2rg	S3rg	S3rg	S3rg	S2rg	S3rg	S3rg	S3rg	S3rg	S3rg	S3rg	S2rg	S3rg	S3g	S3g	S3rg	S3rg	S3rg	S3g
Kasalapura	58	S2rg	S2g	S2g	S2g	S2g	S2t	S2rg	S2g	S2gt	S2g	S2g	S1	S2g	S1	S1	S2rg	S2g	S2t	S2g	S2g	S2g	S2g	S2g	S2g	S2g	S1	S1	S2g	S1	S1	S1	

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Jasmine	Bhendi	Brinjal	Crossandra	Drumstick	Mulberry	Onion	
Halavagali	53	N1rg	S3rg	S3rg	S3rg	S3rg	S3rg	N1rg	S3rg	S2rt	S3rg	S3rg	S2rg	S3rg	S2rg	S3rg	S3rg	S3rg	S2rg	S3rg	S3rg	S3rg	S3rg	S3rg	S2rg	S3rg	S3g	S3g	S3rg	S3rg	S3rg	S3g	
Hesaru	38	S3tz	S2tz	S3tz	S2z	S3tz	S2z	S2tz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2tz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	
Hesaru	39	S3tz	S2tz	S3tz	S2z	S3tz	S2z	S2tz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2tz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	
Hesaru	40	S3tz	S2tz	S3tz	S2z	S3tz	S2z	S2tz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2tz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	
Hesaru	41	S3tz	S2tz	S3tz	S2z	S3tz	S2z	S2tz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2tz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	
Hesaru	42	S1	S2t	S1	S1	S1	S1	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S2t	S2t	S1	S1	S1	S2t
Hesaru	86	S1	S2t	S1	S1	S1	S1	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S2t	S2t	S1	S1	S1	S2t
Hesaru	88	N1r	S2rg	S3rg	S2rg	S3rg	S2rg	N1r	S3rg	S2rt	S3rg	S3rg	S2rg	S3rg	S2rg	S3rg	S3rg	S3rg	S2rg	S2rg	S2rg	S2rg	S2rg	S3rg	S2rg	S2rg	S3g	S3g	S2rg	S3rg	S3rg	S3g	
Hesaru	89	N1r	S2rg	S3rg	S2rg	S3rg	S2rg	N1r	S3rg	S2rt	S3rg	S3rg	S2rg	S3rg	S2rg	S3rg	S3rg	S3rg	S2rg	S2rg	S2rg	S2rg	S2rg	S2rg	S3rg	S2rg	S2rg	S3g	S3g	S2rg	S3rg	S3rg	S3g
Hesaru	90	S3r	S2tg	S2rg	S2g	S2rt	S2rg	S3r	S2rg	S2gt	S2r	S2gt	S1	S2rg	S1	S2rt	S2rg	S2rg	S2t	S1	S1	S2tg	S2tg	S2rg	S1	S2tg	S2t	S1	S2gt	S2r	S2r	S2t	
Hesaru	91	N1r	S2rg	S3rg	S2rg	S3rg	S2rg	N1r	S3rg	S2rt	S3rg	S3rg	S2rg	S3rg	S2rg	S3rg	S3rg	S3rg	S2rg	S2rg	S2rg	S2rg	S2rg	S3rg	S2rg	S2rg	S3g	S3g	S2rg	S3rg	S3rg	S3g	
Hesaru	92	N1r	S2rg	S3rg	S2rg	S3rg	S2rg	N1r	S3rg	S2rt	S3rg	S3rg	S2rg	S3rg	S2rg	S3rg	S3rg	S3rg	S2rg	S2rg	S2rg	S2rg	S2rg	S3rg	S2rg	S2rg	S3g	S3g	S2rg	S3rg	S3rg	S3g	
Hesaru	94	S3r	S2tg	S2rg	S2g	S2rt	S2rg	S3r	S2rg	S2gt	S2r	S2gt	S1	S2rg	S1	S2rt	S2rg	S2rg	S2t	S1	S1	S2tg	S2tg	S2rg	S1	S2tg	S2t	S1	S2gt	S2r	S2r	S2t	
Hesaru	95	S3r	S2tg	S2rg	S2g	S2rt	S2rg	S3r	S2rg	S2gt	S2r	S2gt	S1	S2rg	S1	S2rt	S2rg	S2rg	S2t	S1	S1	S2tg	S2tg	S2rg	S1	S2tg	S2t	S1	S2gt	S2r	S2r	S2t	
Hesaru	96	N1r	S2rg	S3rg	S2rg	S3rg	S2rg	N1r	S3rg	S2rt	S3rg	S3rg	S2rg	S3rg	S2rg	S3rg	S3rg	S3rg	S2rg	S2rg	S2rg	S2rg	S2rg	S3rg	S2rg	S2rg	S3g	S3g	S2rg	S3rg	S3rg	S3g	
Hesaru	97	N1r	S2rg	S3rg	S2rg	S3rg	S2rg	N1r	S3rg	S2rt	S3rg	S3rg	S2rg	S3rg	S2rg	S3rg	S3rg	S3rg	S2rg	S2rg	S2rg	S2rg	S2rg	S3rg	S2rg	S2rg	S3g	S3g	S2rg	S3rg	S3rg	S3g	
Hesaru	98	S3tz	S2tz	S3tz	S1	S3tz	S1	S2tz	S1	S1	S1	S2tz	S1	S3tz	S1	N1t	S2tz	S1	S3t	S2tz	S2tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S3tz	S3tz	
Hesaru	99	S3rg	S3g	S2rg	S3g	S2rg	S3rg	S3rg	S2rg	S3g	S3rg	S3g	S2rg	S2rg	S2rg	S2rg	S2rg	S2rg	S2g	S3g	S3g	S3g	S3g	S2rg	S2g	S3g	S3g	S3g	S3g	S3g	S3g	S2g	S3g
Hesaru	100	N1r	S2rg	S3rg	S2rg	S3rg	S2rg	N1r	S3rg	S2rt	S3rg	S3rg	S2rg	S3rg	S2rg	S3rg	S3rg	S3rg	S2rg	S2rg	S2rg	S2rg	S2rg	S3rg	S2rg	S2rg	S3g	S3g	S2rg	S3rg	S3rg	S3g	
Hesaru	101	N1r	S2rg	S3rg	S2rg	S3rg	S2rg	N1r	S3rg	S2rt	S3rg	S3rg	S2rg	S3rg	S2rg	S3rg	S3rg	S3rg	S2rg	S2rg	S2rg	S2rg	S2rg	S3rg	S2rg	S2rg	S3g	S3g	S2rg	S3rg	S3rg	S3g	
Hesaru	102	N1r	S2rg	S3rg	S2rg	S3rg	S2rg	N1r	S3rg	S2rt	S3rg	S3rg	S2rg	S3rg	S2rg	S3rg	S3rg	S3rg	S2rg	S2rg	S2rg	S2rg	S2rg	S3rg	S2rg	S2rg	S3g	S3g	S2rg	S3rg	S3rg	S3g	
Hesaru	103	S1	S2t	S1	S1	S1	S1	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S2t	S2t	S1	S1	S1	S2t
Hesaru	104	S1	S2t	S1	S1	S1	S1	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S2t	S2t	S1	S1	S1	S2t
Hesaru	105	S3r	S2tg	S2rg	S2g	S2rt	S2rg	S3r	S2rg	S2gt	S2r	S2gt	S1	S2rg	S1	S2rt	S2rg	S2rg	S2t	S1	S1	S2tg	S2tg	S2rg	S1	S2tg	S2t	S1	S2gt	S2r	S2r	S2t	
Hesaru	106	S3r	S2tg	S2rg	S2g	S2rt	S2rg	S3r	S2rg	S2gt	S2r	S2gt	S1	S2rg	S1	S2rt	S2rg	S2rg	S2t	S1	S1	S2tg	S2tg	S2rg	S1	S2tg	S2t	S1	S2gt	S2r	S2r	S2t	
Hesaru	107	S3r	S2tg	S2rg	S2g	S2rt	S2rg	S3r	S2rg	S2gt	S2r	S2gt	S1	S2rg	S1	S2rt	S2rg	S2rg	S2t	S1	S1	S2tg	S2tg	S2rg	S1	S2tg	S2t	S1	S2gt	S2r	S2r	S2t	
Hesaru	108	S3r	S2tg	S2rg	S2g	S2rt	S2rg	S3r	S2rg	S2gt	S2r	S2gt	S1	S2rg	S1	S2rt	S2rg	S2rg	S2t	S1	S1	S2tg	S2tg	S2rg	S1	S2tg	S2t	S1	S2gt	S2r	S2r	S2t	

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Jasmine	Bhendi	Brinjal	Crossandra	Drumstick	Mulberry	Onion
Hesaru	109	S3rg	S3g	S3rg	S3g	S2rt	S3g	S3rg	S2rg	S3g	S2rg	S3g	S2g	S2rg	S2g	S2rt	S2rg	S3rg	S2g	S3g	S3g	S3g	S3g	S3rg	S3g	S3g	S2tg	S2g	S3g	S3g	S2g	S2tg
Hesaru	110	S1	S2t	S1	S1	S1	S1	S1	S1	S2t	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S2t	S2t	S1	S1	S1	S2t
Hesaru	111	S3rg	S3g	S3rg	S3g	S2rt	S3g	S3rg	S2rg	S3g	S2rg	S3g	S2g	S2rg	S2g	S2rt	S2rg	S3rg	S2g	S3g	S3g	S3g	S3g	S3rg	S3g	S3g	S2tg	S2g	S3g	S3g	S2g	S2tg
Hesaru	112	S3rg	S3g	S3rg	S3g	S2rt	S3g	S3rg	S2rg	S3g	S2rg	S3g	S2g	S2rg	S2g	S2rt	S2rg	S3rg	S2g	S3g	S3g	S3g	S3g	S3rg	S3g	S3g	S2tg	S2g	S3g	S3g	S2g	S2tg
Hesaru	113	S3rg	S2tg	S2rg	S2g	S2rt	S2rg	S3rg	S2rg	S2gt	S2r	S2gt	S1	S2rg	S1	S2rt	S2rg	S2rg	S2t	S2g	S2g	S2tg	S2tg	S2rg	S1	S2tg	S2t	S1	S2gt	S2r	S2r	S2t
Hesaru	114	S3rg	S2tg	S2rg	S2g	S2rt	S2rg	S3rg	S2rg	S2gt	S2r	S2gt	S1	S2rg	S1	S2rt	S2rg	S2rg	S2t	S2g	S2g	S2tg	S2tg	S2rg	S1	S2tg	S2t	S1	S2gt	S2r	S2r	S2t
Hesaru	115	S3rg	S2tg	S2rg	S2g	S2rt	S2rg	S3rg	S2rg	S2gt	S2r	S2gt	S1	S2rg	S1	S2rt	S2rg	S2rg	S2t	S2g	S2g	S2tg	S2tg	S2rg	S1	S2tg	S2t	S1	S2gt	S2r	S2r	S2t
Hesaru	116	S3rg	S2tg	S2rg	S2g	S2rt	S2rg	S3rg	S2rg	S2gt	S2r	S2gt	S1	S2rg	S1	S2rt	S2rg	S2rg	S2t	S2g	S2g	S2tg	S2tg	S2rg	S1	S2tg	S2t	S1	S2gt	S2r	S2r	S2t
Hesaru	117	S3rg	S2tg	S2rg	S2g	S2rt	S2rg	S3rg	S2rg	S2gt	S2r	S2gt	S1	S2rg	S1	S2rt	S2rg	S2rg	S2t	S2g	S2g	S2tg	S2tg	S2rg	S1	S2tg	S2t	S1	S2gt	S2r	S2r	S2t
Hesaru	118	S3rg	S2tg	S2rg	S2g	S2rt	S2rg	S3rg	S2rg	S2gt	S2r	S2gt	S1	S2rg	S1	S2rt	S2rg	S2rg	S2t	S2g	S2g	S2tg	S2tg	S2rg	S1	S2tg	S2t	S1	S2gt	S2r	S2r	S2t
Hesaru	119	S3rg	S2tg	S2rg	S2g	S2rt	S2rg	S3rg	S2rg	S2gt	S2r	S2gt	S1	S2rg	S1	S2rt	S2rg	S2rg	S2t	S2g	S2g	S2tg	S2tg	S2rg	S1	S2tg	S2t	S1	S2gt	S2r	S2r	S2t
Hesaru	120	S3rg	S2tg	S2rg	S2g	S2rt	S2rg	S3rg	S2rg	S2gt	S2r	S2gt	S1	S2rg	S1	S2rt	S2rg	S2rg	S2t	S2g	S2g	S2tg	S2tg	S2rg	S1	S2tg	S2t	S1	S2gt	S2r	S2r	S2t
Hesaru	121	S3tz	S2tz	S3tz	S2z	S3tz	S2z	S2tz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2tz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz
Hesaru	122	S3tz	S2tz	S3tz	S2z	S3tz	S2z	S2tz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2tz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz
Hesaru	123	S3rg	S3g	S3rg	S3g	S2rt	S3g	S3rg	S2rg	S3g	S2rg	S3g	S2g	S2rg	S2g	S2rt	S2rg	S3rg	S2g	S3g	S3g	S3g	S3g	S3rg	S3g	S3g	S2tg	S2g	S3g	S3g	S2g	S2tg
Hesaru	124/ A	S3rg	S3g	S3rg	S3g	S2rt	S3g	S3rg	S2rg	S3g	S2rg	S3g	S2g	S2rg	S2g	S2rt	S2rg	S3rg	S2g	S3g	S3g	S3g	S3g	S3rg	S3g	S3g	S2tg	S2g	S3g	S3g	S2g	S2tg
Hesaru	125	S3tz	S2tz	S3tz	S2z	S3tz	S2z	S2tz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2tz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz
Hesaru	126	S3rg	S3g	S3rg	S3g	S2rt	S3g	S3rg	S2rg	S3g	S2rg	S3g	S2g	S2rg	S2g	S2rt	S2rg	S3rg	S2g	S3g	S3g	S3g	S3g	S3rg	S3g	S3g	S2tg	S2g	S3g	S3g	S2g	S2tg
Hesaru	127	S3rg	S3g	S3rg	S3g	S2rt	S3g	S3rg	S2rg	S3g	S2rg	S3g	S2g	S2rg	S2g	S2rt	S2rg	S3rg	S2g	S3g	S3g	S3g	S3g	S3rg	S3g	S3g	S2tg	S2g	S3g	S3g	S2g	S2tg
Hesaru	142	S3tz	S2tz	S3tz	S2z	S3tz	S2z	S2tz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2tz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz
Hesaru	143/ A	S3tz	S2tz	S3tz	S2z	S3tz	S2z	S2tz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2tz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz
Hesaru	143/ B	S3tz	S2tz	S3tz	S2z	S3tz	S2z	S2tz	S2z	S2z	S2z	S2tz	S2tz	S3tz	S2z	N1tz	S2tz	S2z	S3tz	S3tz	S3tz	S2tz	S2tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz	S2tz	S2tz	S3tz

PART-B

SOCIO-ECONOMIC STATUS OF FARM HOUSEHOLDS

CONTENTS

1.	Salient findings of the survey	1-5
2.	Introduction	7
3	Methodology	9
4	Salient features of the survey	11-34
5	Summary	35-39

LIST OF TABLES

1	Households sampled for socio economic survey	11
2	Population characteristics	11
3	Age wise classification of household members	11
4	Education level of household members	12
5	Occupation of household heads	12
6	Occupation of family members	12
7	Institutional participation of household members	13
8	Type of house owned by households	13
9	Durable assets owned by households	13
10	Average value of durable assets owned by households	14
11	Farm implements owned by households	14
12	Average value of farm implements owned by households	15
13	Livestock possession by households	15
14	Average labour availability	15
15	Adequacy of hired labour	16
16	Distribution of land (ha)	16
17	Average land value (Rs./ha)	16
18	Status of bore wells	17
19	Source of irrigation	17
20	Depth of water	17
21	Irrigated area (ha)	17
22	Cropping pattern	18
23	Cropping intensity	18
24	Possession of Bank account	18
25	Borrowing status	18
26	Cost of cultivation of Sugarcane	19
27	Cost of cultivation of Groundnut	20
28	Cost of cultivation of Maize	21
29	Cost of cultivation of Bajra	22
30	Cost of cultivation of Jowar	23
31	Cost of cultivation of Chilly	24

32	Cost of cultivation of Bengalgram	25
33	Cost of cultivation of Sunflower	26
34	Cost of cultivation of Mulberry	27
35	Cost of cultivation of Cotton	28
36	Adequacy of fodder	29
37	Average annual gross income	29
38	Average annual expenditure	29
39	Horticulture species grown	30
40	Forest species grown	30
41	Marketing of the agricultural produce	30
42	Marketing channels used for sale of agricultural produce	31
43	Mode of transport of agricultural produce	31
44	Incidence of soil and water erosion problems	31
45	Interest towards soil testing	32
46	Usage pattern of fuel for domestic use	32
47	Source of drinking water	32
48	Source of light	32
49	Existence of sanitary toilet facility	32
50	Possession of public distribution system(PDS) card	33
51	Participation in NREGA programme	33
52	Adequacy of food items	33
53	Response on inadequacy of food items	34
54	Farming constraints experienced	34

SALIENT FINDINGS OF THE SURVEY

- ❖ *The data indicated that there were 82 (50.31%) men and 81 (49.69%) women among the sampled households.*
- ❖ *The average family size of landless farmers' was 4.67, marginal farmers' was 5.71, small farmers' was 6.29, semi medium farmers' was 4.40, medium farmers' was 4 and large farmers' was 5.*
- ❖ *The data indicated that, 30 (18.40%) people were in 0-15 years of age, 71 (43.56%) were in 16-35 years of age, 44 (26.99%) were in 36-60 years of age and 18 (11.04%) were above 61 years of age.*
- ❖ *The results indicated that Kesalapura-1 had 20.86 per cent illiterates, 36.81 per cent of them had primary school education, 9.82 per cent of them had middle school education, 20.25 per cent of them had high school education, 4.91 per cent of them had PUC education, 0.61 per cent of them did ITI, 4.29 per cent of them had degree education and 0.61 per cent did Maters.*
- ❖ *The results indicate that, 84.38 per cent of households were practicing agriculture, 12.50 per cent of the households were agricultural labourers and 3.13 per cent were in government service.*
- ❖ *The results indicate that agriculture was the major occupation for 62.58 per cent of the household members, 14.72 per cent were agricultural laborers, 0.61 per cent was in government service, 20.25 per cent were students, 1.23 per cent were children and 0.61 per cent were housewives.*
- ❖ *The results show that 99.39 per cent of the population in the micro watershed has not participated in any local institutions and only 0.61 per cent have participated in raitha sangha.*
- ❖ *The results indicate that 3.13 per cent of the households possess thatched house, 71.88 per cent of the households possess Katcha house and 25 per cent of the households possess pucca/RCC house.*
- ❖ *The results show that 65.63 per cent of the households possess TV, 75 per cent of the households possess Mixer grinder, 12.50 per cent of the households possess bicycle, 65.63 per cent of the households possess motor cycle, 6.25 per cent possess refrigerator, another 6.25 per cent possess car/four wheel and 93.75 per cent of the households possess mobile phones.*
- ❖ *The results show that the average value of television was Rs. 2595, mixer grinder was Rs.1362, bicycle was Rs. 12850, motor cycle was Rs.33272, refrigerator was Rs.13500, car/wheeler was Rs.150000 and mobile phone was Rs.1264.*
- ❖ *About 6.25 per cent of the households possess bullock cart, 21.88 per cent of the households possess plough, 3.13 per cent of them possess seed/fertilizer drill, 9.38 per cent of them possess power tiller, 37.50 per cent of them possess tractor, 37.50 per cent of them possess sprayer, 9.38 per cent of them possess sprinkler, 12.50 per*

cent possess harvester, 50 per cent of them possess chaff cutter, 18.75 per cent of them possess earth remover/duster and 71.88 per cent of them possess weeder.

- ❖ The results show that the average value of bullock cart was Rs.16500, the average value of plough was Rs.823, seed/fertilizer drill was Rs.25000, irrigation pump was Rs.21250, power tiller was Rs.15000, tractor was Rs.296000, the average value of sprayer was Rs.1900, weeder was Rs.24, harvester was Rs.13333, earth remover/duster was Rs.19500 and chaff cutter was Rs.817.
- ❖ The results indicate that, 21.88 per cent of the households possess bullocks, crossbred cow and buffalo. About 43.75 per cent of the households possess local cow and 18.75 per cent of the households possess poultry birds.
- ❖ The results indicate that, average own labour men available in the micro watershed was 2.07, average own labour (women) available was 1.93, average hired labour (men) available was 7.79 and average hired labour (women) available was 7.36.
- ❖ The results indicate that, 87.50 per cent of the households opined that the hired labour was adequate.
- ❖ The results indicate that, households of the Kesalapura-1 micro-watershed possess 24.74 ha (45.42%) of dry land and 29.73 ha (54.58%) of irrigated land. Marginal farmers possess 3.04 ha (70.92%) of dry land and 1.25 ha (29.08%). Small farmers possess 4.3 ha (57.97%) of dry land and 3.12 ha (42.03%) of irrigated land. Semi medium farmers possess 2.83 ha (17.02%) of dry land and 13.81 per cent (82.98%) of irrigated land. Medium farmers possess 1.62 ha (12.29%) dry land and 11.56 (87.71%) of irrigated land. Large farmers possess 12.95 ha (100%) of dry land.
- ❖ The results indicate that, the average value of dry land was Rs. 187078 and average value of irrigated land was Rs. 416878. In case of marginal famers, the average land value was Rs. 526232 for dry land and Rs. 1443506 for irrigated land. In case of small famers, the average land value was Rs. 146525 for dry land and was Rs. 737792 for irrigated land. In case of semi medium famers, the average land value was Rs. 211714 for dry land and Rs. 419748 for irrigated land. In case of medium famers, the average land value was Rs. 370500 for dry land and Rs. 216211 for irrigated land. In case of large farmers, the average land value for dry land was Rs. 92625.
- ❖ The results indicate that, there were 20 functioning bore wells in the micro watershed.
- ❖ The results indicate that, bore well was the major irrigation source in the micro water shed for 62.50 per cent of the farmers.
- ❖ The results indicate that, the depth of bore well was found to be 44.10 meters.
- ❖ The results indicate that, marginal, small, semi medium and medium farmers had irrigated area of 1.21 ha, 7.65 ha, 12.01 ha and 11.48 ha respectively.

- ❖ *The results indicate that, farmers have grown sugarcane (15.91 ha), groundnut (4.45 ha), maize (7.03 ha), sunflower (2.86 ha), pearl millet (5.19 ha), bengalgram (2.05 ha), chilly (1.21 ha), cotton (0.4 ha), mulberry (0.03 ha) and jowar (0.4 ha).*
- ❖ *Marginal farmers have grown sugarcane, maize, sunflower, pearl millet, bengalgram, jowar and mulberry. Small farmers have grown sugarcane, maize, groundnut, sunflower, pearl millet, groundnut and cotton. Semi medium farmers have grown sugarcane, maize, bajra, pearl millet, bengalgram and chilly. Medium farmers have grown sugarcane, maize, groundnut, sunflower and chilly. Large farmers have grown bajra only.*
- ❖ *The results indicate that, the cropping intensity in Kesalapura-1 micro-watershed was found to be 72.08 per cent. In case of marginal farmers it was 92.48 per cent, small farmers it was 94.18 per cent, in case of semi medium farmers it was 74.11, medium farmers it was 62.52 per cent and in case of large farmers it was 50 per cent.*
- ❖ *The results indicate that, 46.88 per cent of the households have bank account.*
- ❖ *The results indicate that, 46.88 per cent of the households have availed credit from different sources.*
- ❖ *The results indicate that, the total cost of cultivation for sugarcane was Rs. 60569.99. The gross income realized by the farmers was Rs. 525781.85. The net income from sugarcane cultivation was Rs. 465211.86. Thus the benefit cost ratio was found to be 1:8.68.*
- ❖ *The total cost of cultivation for groundnut was Rs. 42208.67. The gross income realized by the farmers was Rs. 102827.47. The net income from groundnut cultivation was Rs. 60618.81. Thus the benefit cost ratio was found to be 1:2.44.*
- ❖ *The total cost of cultivation for maize was Rs. 43481.62. The gross income realized by the farmers was Rs. 54611.46. The net income from maize cultivation was Rs. 11129.85. Thus the benefit cost ratio was found to be 1:1.26.*
- ❖ *The total cost of cultivation for bajra was Rs. 19721.25. The gross income realized by the farmers was Rs. 22622.75. The net income from bajra cultivation was Rs. 2901.51. Thus the benefit cost ratio was found to be 1:1.15.*
- ❖ *The total cost of cultivation for Jowar was Rs. 61598.57. The gross income realized by the farmers was Rs. 55328. The net income from Jowar cultivation was Rs. - 6270.57. Thus the benefit cost ratio was found to be 1:0.9.*
- ❖ *The total cost of cultivation for chilly was Rs. 223901.85. The gross income realized by the farmers was Rs. 322643.75. The net income from chilly cultivation was Rs. 98741.90. Thus the benefit cost ratio was found to be 1:1.44.*
- ❖ *The total cost of cultivation for Bengalgram was Rs. 52147.87. The gross income realized by the farmers was Rs. 97556.48. The net income from Bengalgram cultivation was Rs. 45408.62. Thus the benefit cost ratio was found to be 1:1.87.*

- ❖ *The total cost of cultivation for Sunflower was Rs. 32268.06. The gross income realized by the farmers was Rs. 64321.25. The net income from Sunflower cultivation was Rs. 32053.20. Thus the benefit cost ratio was found to be 1:1.99.*
- ❖ *The total cost of cultivation for Mulberry was Rs. 849677.77. The gross income realized by the farmers was Rs. 2161250.05. The net income from Mulberry cultivation was Rs. 1311572.28. Thus the benefit cost ratio was found to be 1:2.54.*
- ❖ *the total cost of cultivation for cotton was Rs. 48650.76. The gross income realized by the farmers was Rs. 118560. The net income from cotton cultivation was Rs. 69909.24. Thus the benefit cost ratio was found to be 1:2.44.*
- ❖ *The results indicate that, 65.63 per cent of the households opined that dry fodder was adequate and 40.63 per cent opined that green fodder was adequate.*
- ❖ *The results indicate that the average annual gross income was Rs. 37667 for landless farmers, for marginal farmers it was Rs. 181857, for small farmers it was Rs. 118516, for semi medium farmers it was Rs. 225550, for medium farmers it was Rs. 355000 and for large farmers it was Rs.120000.*
- ❖ *The results indicate that the average annual expenditure is Rs. 33462. For landless households it was Rs. 6889, for marginal farmers it was Rs. 68184, for small farmers it was Rs. 16335, for semi medium farmers it was Rs. 16350, for medium farmers it was Rs. 58750 and for large farmers it was Rs.60000.*
- ❖ *The results indicate that, households have planted 69 coconut trees, 2 lemon trees, 2 lime trees and 7 mango trees in the field and 8 coconut trees in the backyard.*
- ❖ *The results indicate that, households have planted 75 neem tree, 1 banyan tree and 29 teak trees in their fields.*
- ❖ *The results indicated that, cotton, maize, mulberry, sugarcane and sunflower were sold to the extent of 100 per cent. Bajra was sold to the extent of 86.67 per cent, Bengal gram to the extent of 54.55 per cent, chilly to the extent of 94.29 per cent, groundnut to the extent of 52.38 per cent and jowar was sold to the extent of 68.75 per cent.*
- ❖ *The results indicated that, about 18.75 per cent of the famers have sold their produce in regulated markets, 53.13 per cent of the farmers have sold to local/village merchants and another 53.13 per cent of the farmers have sold to agents/traders.*
- ❖ *The results indicated that, 87.50 per cent of the households have used tractor, 25 per cent of the households have used cart, 9.38 per cent of the households have used truck and 3.13 per cent have carried head loads as a mode of transportation for their agricultural produce.*
- ❖ *The results indicated that, 46.88 per cent of the households have experienced soil and water erosion problems in the farm i.e., 71.43 per cent of the marginal farmers, 28.5 per cent of the small farmers, 40 per cent of semi medium farmers and 75 per cent of medium farmers and 100 per cent of large farmers have experienced soil and water erosion problems.*

- ❖ *The results indicated that, 50 per cent have shown interest in soil test.*
- ❖ *The results indicated that, 53.13 per cent of the households used firewood and 46.88 per cent have used LPG as a source of fuel.*
- ❖ *The results indicated that, bore well was the major source of drinking water for 34.38 per cent of the households and piped supply was the source of drinking water for 59.38 per cent of the households in the micro watershed.*
- ❖ *Electricity was the major source of light for 100 per cent of the households in micro watershed.*
- ❖ *The results indicated that, 43.75 per cent of the households possess sanitary toilet.*
- ❖ *The results indicated that, 90.63 per cent of the sampled households possessed BPL card and 9.38 per cent of the households did not possess any PDS cards.*
- ❖ *The results indicated that, 46.88 per cent of the households participated in NREGA programme.*
- ❖ *The results indicated that, cereals were adequate for 100 per cent of the households, pulses were adequate for 62.50 per cent, oilseeds were adequate for 68.75 per cent, vegetables were adequate for 62.50 per cent, fruits were adequate for 6.25 per cent, milk was adequate for 90.63 per cent, eggs were adequate for 56.25 per cent and meat were adequate for 46.88 per cent.*
- ❖ *The results indicated that, pulses were inadequate for 37.50 per cent of the households, oilseeds were inadequate for 31.25 per cent, vegetables were inadequate for 37.50 per cent, fruits were inadequate for 53.13 per cent, eggs were inadequate for 28.13 per cent, milk was inadequate for 9.38 per cent of the households and meat was inadequate for 37.50 per cent of the households.*
- ❖ *The results indicated that, lower fertility status of the soil was the constraint experienced by 50 per cent of the households, wild animal menace on farm field (59.38%), frequent incidence of pest and diseases (71.88%), inadequacy of irrigation water (31.25%), high cost of fertilizers and plant protection chemicals (84.38%), high rate of interest on credit (71.88%), low price for the agricultural commodities (43.75%), lack of marketing facilities in the area (65.63%), lack of transport for safe transport of the agricultural produce to the market (43.75%), inadequate extension services (9.38%), less rainfall (53.13%) and source of agri-technology information (15.63%).*

INTRODUCTION

Soil and water are the two precious natural resources which are essential for crop production and existence of life on earth. Rainfed agriculture is under severe stress due to various constraints related to agriculture like uneven and erratic distribution of rainfall, indiscriminate use of fertilizers, chemicals and pesticides, adoption of improper land management practices, soil erosion, decline in soil fertility, decline in ground water resources leading to low crop productivity. The area under rainfed agriculture has to be managed effectively using the best available practices to enhance the production of food, fodder and fuel. This is possible if the land resources are characterized at each parcel of land through detailed land resource inventory using the best available techniques of remote sensing, GPS and GIS. The watershed development programs are aimed at the sustainable distribution of its resources and the process of creating and implementing plans, programs, and projects to sustain and enhance watershed functions that affect the plant, animal and human communities within a watershed boundary.

World Bank funded KWDP II, SUJALA III project was implemented in with Broad objective of demonstrating more effective watershed management through greater integration of programmes related to rain-fed agriculture, innovative and science based approaches and strengthen institutional capacities and If successful, it is expected that the systems and tools could be mainstreamed into the overall IWMP in the State of Karnataka and in time, throughout other IWMP operations in India. With this background the socio-economic survey has been carried out with following specific objectives:

1. To understand the demographic features of the households in the micro-watershed
2. To understand the extent of family labour available and additional employment opportunities available within the village.
3. To know the status of assets of households in the micro-watershed for suggesting possible improvements.
4. To study the cropping pattern, cropped area and productivity levels of different households in micro-watershed.
5. To determine the type and extent of livestock owned by different categories of HHs
6. Availability of fodder and level of livestock management.

Scope and importance of survey

Survey helps in identification of different socio-economic and resource use-patterns of farmers at the Micro watershed. Household survey provides demographic features, labour force, and levels of education; land ownership and asset position (including livestock and other household assets) of surveyed households; and cropping patterns, input intensities, and average crop yields from farmers' fields. It also discusses crop utilization and the degree of commercialization of production in the areas; farmers' access to and utilization of credit from formal and informal sources; and the level of adoption and use of soil, water, and pest management technologies.

METHODOLOGY

The description of the methods, components selected for the survey and procedures followed in conducting the baseline survey are furnished under the following heads.

Description of the study area

Koppal district is an administrative district in the state of Karnataka in India. In the past Koppal was referred to as 'Kopana Nagara'. Koppal, now a district headquarters is ancient Kopana a major holy place of the Jainas. The district occupies an area of 7,190 km² and has a population of 1,196,089, which 16.58% were urban as of 2001. The Koppal district was formed after split of Raichur district.

Geographers are very particular about the physiography or relief of a region. It plays a very important role in the spatial analysis of agricultural situation of the study area. The undulating topography with black cotton soil shrips, cut across by numerous nalas or streams is the major characteristic feature of the study region. Three physiographic divisions have made considering the local conditions of landforms and crops grown in the district. On the basis of physiography, Koppal district can be divided into three major divisions.³ They are (a) Koppal & Yelburga plateau, (b) Maidan division, (c) Tungabhadra valley. The district is part of Krishna basin the main streams draining the area are Maskinala, Ilkal-nadi and Hirenala. These are Ephemeral in nature, these come under Tungabhadra sub-basin. The drainage exhibit dentritic to subdentritic with drainage density varies from 1.4 to 7.0kms/sq.km.

According to the 2011 census Koppal district has a population of 1,391,292, roughly equal to the nation of Swaziland or the US state of Hawaii. This gives it a ranking of 350th in India (out of a total of 640). The district has a population density of 250 inhabitants per square kilometre (650/sq mi). Its population growth rate over the decade 2001-2011 was 16.32%. Koppal has a sex ratio of 983 females for every 1000 males, and a literacy rate of 67.28%.

Description of the micro watershed

Kesalapura-1 micro-watershed (Bettageri sub-watershed, Koppal Taluk and District) is located at North latitude 15⁰10'16.428'' to 15⁰8'51.552'' and East longitude 75⁰57'42.996'' to 75⁰57'0.618'' E covering an area of 304.53 ha and spread across Kesalapura and Halavagali villages.

Methodology followed in assessing socio-economic status of households

In order to assess the socio-economic condition of the farmers in the watershed a comprehensive questionnaire was prepared. Major components such as demographic conditions, migration details, food consumption and family expenditure pattern, material possession, land holding, land use management, cropping pattern, cost of cultivation of crops, livestock management. The statistical components such as frequency and percentage were used to analyze the data. About 32 households located in the micro watershed were interviewed for the survey.

SALIENT FEATURES OF THE SURVEY

Households sampled for socio-economic survey: The data on households sampled for socio economic survey in Kesalapura-1 micro-watershed is presented in Table 1 and it indicated that 32 farmers were sampled in Kesalapura-1 micro-watershed among them 3 (9.38%) were landless, 7 (21.88%) were marginal farmers, 7 (21.88%) were small farmers, 10 (31.25%) were semi medium farmers, 4 (12.50%) were medium farmers and 1 (3.13%) was large farmer.

Table 1: Households sampled for socio economic survey in Kesalapura-1 micro-watershed

Sl.No.	Particulars	LL (3)		MF (7)		SF (7)		SMF (10)		MDF (4)		LF (1)		All (32)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Farmers	3	9.38	7	21.88	7	21.88	10	31.25	4	12.50	1	3.13	32	100.00

Population characteristics: The population characteristics of households sampled for socio-economic survey in Kesalapura-1 micro-watershed is presented in Table 2. The data indicated that there were 82 (50.31%) men and 81 (49.69%) women among the sampled households. The average family size of landless farmers' was 4.67, marginal farmers' was 5.71, small farmers' was 6.29, semi medium farmers' was 4.40, medium farmers' was 4 and large farmers' was 5.

Table 2: Population characteristics of Kesalapura-1 micro-watershed

Sl. No.	Particulars	LL (14)		MF (40)		SF (44)		SMF (44)		MDF (16)		LF (5)		All (163)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Male	6	42.86	19	47.50	22	50.00	22	50.00	11	68.75	2	40.00	82	50.31
2	Female	8	57.14	21	52.50	22	50.00	22	50.00	5	31.25	3	60.00	81	49.69
Total		14	100.00	40	100.00	44	100.00	44	100.00	16	100.00	5	100.00	163	100.00
Average		4.67		5.71		6.29		4.40		4.00		5.00		5.09	

Age wise classification of population: The age wise classification of household members in Kesalapura-1 micro-watershed is presented in Table 3. The data indicated that, 30 (18.40%) people were in 0-15 years of age, 71 (43.56%) were in 16-35 years of age, 44 (26.99%) were in 36-60 years of age and 18 (11.04%) were above 61 years of age.

Table 3: Age wise classification of household members in Kesalapura-1 micro-watershed

Sl. No.	Particulars	LL (14)		MF (40)		SF (44)		SMF(44)		MDF(16)		LF (5)		All (163)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	0-15 years of age	4	28.57	8	20.00	10	22.73	5	11.36	1	6.25	2	40.00	30	18.40
2	16-35 years of age	5	35.71	21	52.50	19	43.18	20	45.45	5	31.25	1	20.00	71	43.56
3	36-60 years of age	4	28.57	9	22.50	10	22.73	12	27.27	8	50.00	1	20.00	44	26.99
4	> 61 years	1	7.14	2	5.00	5	11.36	7	15.91	2	12.50	1	20.00	18	11.04
Total		14	100.00	40	100.00	44	100.00	44	100.00	16	100.00	5	100.00	163	100.00

Education level of household members: Education level of household members in Kesalapura-1 micro-watershed is presented in Table 4. The results indicated that

Kesalapura-1 had 20.86 per cent illiterates, 36.81 per cent of them had primary school education, 9.82 per cent of them had middle school education, 20.25 per cent of them had high school education, 4.91 per cent of them had PUC education, 0.61 per cent of them did ITI, 4.29 per cent of them had degree education and 0.61 per cent did Masters.

Table 4. Education level of household members in Kesalapura-1 micro-watershed

Sl. No.	Particulars	LL (14)		MF (40)		SF (44)		SMF (44)		MDF (16)		LF (5)		All (163)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Illiterate	3	21.43	9	22.50	15	34.09	4	9.09	2	12.50	1	20.00	34	20.86
2	Primary School	3	21.43	9	22.50	16	36.36	23	52.27	7	43.75	2	40.00	60	36.81
3	Middle School	4	28.57	5	12.50	3	6.82	2	4.55	2	12.50	0	0.00	16	9.82
4	High School	3	21.43	12	30.00	6	13.64	10	22.73	1	6.25	1	20.00	33	20.25
5	PUC	1	7.14	1	2.50	2	4.55	2	4.55	2	12.50	0	0.00	8	4.91
6	ITI	0	0.00	1	2.50	0	0.00	0	0.00	0	0.00	0	0.00	1	0.61
7	Degree	0	0.00	1	2.50	1	2.27	3	6.82	2	12.50	0	0.00	7	4.29
8	Masters	0	0.00	0	0.00	0	0.00	0	0.00	0	0.00	1	20.00	1	0.61
9	Others	0	0.00	2	5.00	1	2.27	0	0.00	0	0.00	0	0.00	3	1.84
Total		14	100.00	40	100.00	44	100.00	44	100.00	16	100.00	5	100.00	163	100.00

Occupation of household heads: The data regarding the occupation of the household heads in Kesalapura-1 micro-watershed is presented in Table 5. The results indicate that, 84.38 per cent of households were practicing agriculture, 12.50 per cent of the households were agricultural labourers and 3.13 per cent were in government service.

Table 5: Occupation of household heads in Kesalapura-1 micro-watershed

Sl. No.	Particulars	LL (3)		MF (7)		SF (7)		SMF(10)		MDF(4)		LF (1)		All (32)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Agriculture	0	0.00	6	85.71	6	85.71	10	100.00	4	100.00	1	100.00	27	84.38
2	Agricultural Labour	3	100.00	1	14.29	0	0.00	0	0.00	0	0.00	0	0.00	4	12.50
3	Government Service	0	0.00	1	14.29	0	0.00	0	0.00	0	0.00	0	0.00	1	3.13
Total		3	100.00	8	100.00	6	100.00	10	100.00	4	100.00	1	100.00	32	100.00

Table 6: Occupation of family members in Kesalapura-1 micro-watershed

Sl. No.	Particulars	LL (14)		MF (40)		SF (44)		SMF(44)		MDF(16)		LF (5)		All (163)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Agriculture	0	0.00	27	67.50	22	50.00	36	81.82	14	87.50	3	60.00	102	62.58
2	Agricultural Labour	10	71.43	1	2.50	10	22.73	3	6.82	0	0.00	0	0.00	24	14.72
3	Government Service	0	0.00	1	2.50	0	0.00	0	0.00	0	0.00	0	0.00	1	0.61
4	Student	4	28.57	10	25.00	10	22.73	5	11.36	2	12.50	2	40.00	33	20.25
5	Housewife	0	0.00	0	0.00	1	2.27	0	0.00	0	0.00	0	0.00	1	0.61
6	Children	0	0.00	1	2.50	1	2.27	0	0.00	0	0.00	0	0.00	2	1.23
Total		14	100.00	40	100.00	44	100.00	44	100.00	16	100.00	5	100.00	163	100.00

Occupation of the household members: The data regarding the occupation of the household members in Kesalapura-1 micro-watershed is presented in Table 6. The results indicate that agriculture was the major occupation for 62.58 per cent of the household members, 14.72 per cent were agricultural laborers, 0.61 per cent was in government

service, 20.25 per cent were students, 1.23 per cent were children and 0.61 per cent were housewives.

Institutional participation of the household members: The data regarding the institutional participation of the household members in Kesalapura-1 micro-watershed is presented in Table 7. The results show that 99.39 per cent of the population in the micro watershed has not participated in any local institutions and only 0.61 per cent have participated in raitha sangha.

Table 7. Institutional Participation of household members in Kesalapura-1 micro-watershed

Sl. No.	Particulars	LL (14)		MF (40)		SF (44)		SMF (44)		MDF (16)		LF (5)		All (163)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Raitha Sangha	0	0.00	0	0.00	0	0.00	1	2.27	0	0.00	0	0.00	1	0.61
2	No Participation	14	100.00	40	100.00	44	100.00	43	97.73	16	100.00	5	100.00	162	99.39
Total		14	100.00	40	100.00	44	100.00	44	100.00	16	100.00	5	100.00	163	100.00

Type of house owned: The data regarding the type of house owned by the households in Kesalapura-1 micro-watershed is presented in Table 8. The results indicate that 3.13 per cent of the households possess thatched house, 71.88 per cent of the households possess Katcha house and 25 per cent of the households possess pucca/RCC house.

Table 8. Type of house owned by households in Kesalapura-1 micro-watershed

Sl. No.	Particulars	LL (3)		MF (7)		SF (7)		SMF (10)		MDF (4)		LF (1)		All (32)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Thatched	0	0.00	1	14.29	0	0.00	0	0.00	0	0.00	0	0.00	1	3.13
2	Katcha	2	66.67	5	71.43	6	85.71	6	60.00	3	75.00	1	100.00	23	71.88
3	Pucca/RCC	1	33.33	1	14.29	1	14.29	4	40.00	1	25.00	0	0.00	8	25.00
Total		3	100.00	7	100.00	7	100.00	10	100.00	4	100.00	1	100.00	32	100.00

Durable Assets owned by the households: The data regarding the Durable Assets owned by the households in Kesalapura-1 micro-watershed is presented in Table 9. The results show that 65.63 per cent of the households possess TV, 75 per cent of the households possess Mixer grinder, 12.50 per cent of the households possess bicycle, 65.63 per cent of the households possess motor cycle, 6.25 per cent possess refrigerator, another 6.25 per cent possess car/four wheel and 93.75 per cent of the households possess mobile phones.

Table 9. Durable Assets owned by households in Kesalapura-1 micro-watershed

Sl. No.	Particulars	LL (3)		MF (7)		SF (7)		SMF (10)		MDF (4)		LF (1)		All (32)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Television	1	33.33	3	42.86	5	71.43	8	80.00	3	75.00	1	100.00	21	65.63
2	Mixer/Grinder	3	100.00	5	71.43	5	71.43	7	70.00	3	75.00	1	100.00	24	75.00
3	Refrigerator	0	0.00	0	0.00	0	0.00	1	10.00	1	25.00	0	0.00	2	6.25
4	Bicycle	1	33.33	1	14.29	0	0.00	1	10.00	1	25.00	0	0.00	4	12.50
5	Motor Cycle	1	33.33	5	71.43	4	57.14	7	70.00	3	75.00	1	100.00	21	65.63
6	Car/Four Wheeler	0	0.00	1	14.29	0	0.00	0	0.00	1	25.00	0	0.00	2	6.25
7	Mobile Phone	3	100.00	7	100.00	6	85.71	9	90.00	4	100.00	1	100.00	30	93.75

Average value of durable assets: The data regarding the average value of durable assets owned by the households in Kesalapura-1 micro-watershed is presented in Table 10. The results show that the average value of television was Rs. 2595, mixer grinder was Rs.1362, bicycle was Rs. 12850, motor cycle was Rs.33272, refrigerator was Rs.13500, car/wheeler was Rs.150000 and mobile phone was Rs.1264.

Table 10. Average value of durable assets owned by households in Kesalapura-1 micro-watershed
Average value (Rs.)

Sl.No.	Particulars	LL (3)	MF (7)	SF (7)	SMF (10)	MDF (4)	LF (1)	All (32)
1	Television	3,000	2,500	2,400	2,500	3,333	2,000	2,595
2	Mixer/Grinder	1,333	1,480	1,060	1,400	1,733	1,000	1,362
3	Refrigerator	0	0	0	12,000	15,000	0	13,500
4	Bicycle	25,000	700	0	700	25,000	0	12,850
5	Motor Cycle	20,000	35,000	36,250	32,125	35,000	30,000	33,272
6	Car/Four Wheeler	0	3,00,000	0	0	2,00,000	0	2,50,000
7	Mobile Phone	1,375	1,272	1,300	1,214	1,416	750	1,264

Farm Implements owned: The data regarding the farm implements owned by the households in Kesalapura-1 micro-watershed is presented in Table 11. About 6.25 per cent of the households possess bullock cart, 21.88 per cent of the households possess plough, 3.13 per cent of them possess seed/fertilizer drill, 9.38 per cent of them possess power tiller, 37.50 per cent of them possess tractor, 37.50 per cent of them possess sprayer, 9.38 per cent of them possess sprinkler, 12.50 per cent possess harvester, 50 per cent of them possess chaff cutter, 18.75 per cent of them possess earth remover/duster and 71.88 per cent of them possess weeder.

Table 11. Farm Implements owned by households in Kesalapura-1 micro-watershed

Sl. No.	Particulars	LL (3)		MF (7)		SF (7)		SMF (10)		MDF (4)		LF (1)		All (32)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Bullock Cart	0	0.00	0	0.00	0	0.00	2	20.00	0	0.00	0	0.00	2	6.25
2	Plough	0	0.00	2	28.57	1	14.29	4	40.00	0	0.00	0	0.00	7	21.88
3	Seed/Fertilizer Drill	0	0.00	0	0.00	0	0.00	1	10.00	0	0.00	0	0.00	1	3.13
4	Irrigation Pump	0	0.00	0	0.00	0	0.00	3	30.00	1	25.00	0	0.00	4	12.50
5	Power Tiller	0	0.00	1	14.29	0	0.00	2	20.00	0	0.00	0	0.00	3	9.38
6	Tractor	1	33.33	2	28.57	2	28.57	4	40.00	3	75.00	0	0.00	12	37.50
7	Sprayer	0	0.00	2	28.57	1	14.29	6	60.00	2	50.00	1	100.00	12	37.50
8	Sprinkler	0	0.00	1	14.29	1	14.29	0	0.00	1	25.00	0	0.00	3	9.38
9	Weeder	3	100.00	5	71.43	3	42.86	9	90.00	2	50.00	1	100.00	23	71.88
10	Harvester	0	0.00	0	0.00	1	14.29	2	20.00	1	25.00	0	0.00	4	12.50
11	Chaff Cutter	2	66.67	4	57.14	2	28.57	5	50.00	2	50.00	1	100.00	16	50.00
12	Earth remover/Duster	0	0.00	1	14.29	1	14.29	3	30.00	1	25.00	0	0.00	6	18.75
13	Blank	0	0.00	2	28.57	3	42.86	1	10.00	0	0.00	0	0.00	6	18.75

Average value of farm implements: The data regarding the average value of farm Implements owned by the households in Kesalapura-1 micro-watershed is presented in Table 12. The results show that the average value of bullock cart was Rs.16500, the average value of plough was Rs.823, seed/fertilizer drill was Rs.25000, irrigation pump

was Rs.21250, power tiller was Rs.15000, tractor was Rs.296000, the average value of sprayer was Rs.1900, weeder was Rs.24, harvester was Rs.13333, earth remover/duster was Rs.19500 and chaff cutter was Rs.817.

Table 12. Average value of farm implements owned by households in Kesalapura-1 micro-watershed

Sl.No.	Particulars	Average Value (Rs.)						
		LL (3)	MF (7)	SF (7)	SMF (10)	MDF (4)	LF (1)	All (32)
1	Bullock Cart	0	0	0	16500	0	0	16500
2	Plough	0	500	200	2250	0	0	823
3	Seed/Fertilizer Drill	0	0	0	25000	0	0	25000
4	Irrigation Pump	0	0	0	20000	25000	0	21250
5	Power Tiller	0	20000	0	12500	0	0	15000
6	Tractor	200000	151000	250000	362500	366666	0	296000
7	Sprayer	0	2000	4000	2071	1250	1000	1900
8	Weeder	22	26	25	26	18	18	24
9	Harvester	0	0	30000	6250	25000	0	13333
10	Chaff Cutter	500	850	750	1000	750	500	817
11	Earth remover/Duster	0	18000	20000	18000	25000	0	19500

Livestock possession by the households: The data regarding the Livestock possession by the households in Kesalapura-1 micro-watershed is presented in Table 13. The results indicate that, 21.88 per cent of the households possess bullocks, crossbred cow and buffalo. About 43.75 per cent of the households possess local cow and 18.75 per cent of the households possess poultry birds.

Table 13. Livestock possession by households in Kesalapura-1 micro-watershed

Sl. No.	Particulars	LL (3)		MF (7)		SF (7)		SMF (10)		MDF (4)		LF (1)		All (32)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Bullock	1	33.33	3	42.86	1	14.29	1	10.00	1	25.00	0	0.00	7	21.88
2	Local cow	1	33.33	3	42.86	4	57.14	4	40.00	1	25.00	1	100.00	14	43.75
3	Crossbred cow	0	0.00	1	14.29	1	14.29	4	40.00	1	25.00	0	0.00	7	21.88
4	Buffalo	0	0.00	2	28.57	3	42.86	2	20.00	0	0.00	0	0.00	7	21.88
5	Poultry birds	2	66.67	2	28.57	0	0.00	1	10.00	0	0.00	1	100.00	6	18.75
6	blank	0	0.00	3	42.86	1	14.29	1	10.00	3	75.00	0	0.00	8	25.00

Average Labour availability: The data regarding the average labour availability in Kesalapura-1 micro-watershed is presented in Table 14. The results indicate that, average own labour men available in the micro watershed was 2.07, average own labour (women) available was 1.93, average hired labour (men) available was 7.79 and average hired labour (women) available was 7.36.

Table 14. Average Labour availability in Kesalapura-1 micro-watershed

Sl.No.	Particulars	LL (3)	MF (7)	SF (7)	SMF (10)	MDF (4)	LF (1)	All (32)
		N	N	N	N	N	N	N
1	Hired labour Female	0.00	5.57	6.43	11.22	4.00	5.00	7.36
2	Hired labour Male	0.00	5.86	7.14	11.78	4.00	5.00	7.79
3	Own Labour Female	0.00	2.00	2.43	1.78	1.25	2.00	1.93
4	Own labour Male	0.00	2.14	2.14	1.89	2.50	1.00	2.07

Adequacy of Hired Labour: The data regarding the adequacy of hired labour in Kesalapura-1 micro-watershed is presented in Table 15. The results indicate that, 87.50 per cent of the households opined that the hired labour was adequate.

Table 15. Adequacy of Hired Labour in Kesalapura-1 micro-watershed

Sl.No.	Particulars	LL (3)		MF (7)		SF (7)		SMF (10)		MDF (4)		LF (1)		All (32)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Adequate	0	0.00	7	100.00	7	100.00	9	90.00	4	100.00	1	100.00	28	87.50

Distribution of land (ha): The data regarding the distribution of land (ha) in Kesalapura-1 micro-watershed is presented in Table 16. The results indicate that, households of the Kesalapura-1 micro-watershed possess 24.74 ha (45.42%) of dry land and 29.73 ha (54.58%) of irrigated land. Marginal farmers possess 3.04 ha (70.92%) of dry land and 1.25 ha (29.08%). Small farmers possess 4.3 ha (57.97%) of dry land and 3.12 ha (42.03%) of irrigated land. Semi medium farmers possess 2.83 ha (17.02%) of dry land and 13.81 per cent (82.98%) of irrigated land. Medium farmers possess 1.62 ha (12.29%) dry land and 11.56 (87.71%) of irrigated land. Large farmers possess 12.95 ha (100%) of dry land.

Table 16. Distribution of land (Ha) in Kesalapura-1 micro-watershed

Sl. No.	Particulars	LL (3)		MF (7)		SF (7)		SMF (10)		MDF (4)		LF (1)		All (32)	
		ha	%	ha	%	ha	%	ha	%	ha	%	ha	%	ha	%
1	Dry	0	0	3.04	70.92	4.3	57.97	2.83	17.02	1.62	12.29	12.95	100	24.74	45.42
2	Irrigated	0	0	1.25	29.08	3.12	42.03	13.81	82.98	11.56	87.71	0	0	29.73	54.58
Total		0	100	4.29	100	7.41	100	16.65	100	13.18	100	12.95	100	54.47	100

Average land value (Rs./ha): The data regarding the average land value (Rs./ha) in Kesalapura-1 micro-watershed is presented in Table 17. The results indicate that, the average value of dry land was Rs. 187078 and average value of irrigated land was Rs. 416878. In case of marginal famers, the average land value was Rs. 526232 for dry land and Rs. 1443506 for irrigated land. In case of small famers, the average land value was Rs. 146525 for dry land and was Rs. 737792 for irrigated land. In case of semi medium famers, the average land value was Rs. 211714 for dry land and Rs. 419748 for irrigated land. In case of medium famers, the average land value was Rs. 370500 for dry land and Rs. 216211 for irrigated land. In case of large farmers, the average land value for dry land was Rs. 92625.

Table 17. Average land value (Rs./ha) in Kesalapura-1 micro-watershed

Sl.No.	Particulars	LL (3)	MF (7)	SF (7)	SMF (10)	MDF (4)	LF (1)	All (32)
1	Dry	0	526232	146525	211714	370500	92625	187078
2	Irrigated	0	1443506	737792	419748	216211	0	416878

Status of bore wells: The data regarding the status of bore wells in Kesalapura-1 micro-watershed is presented in Table 18. The results indicate that, there were 20 functioning bore wells in the micro watershed.

Table 18. Status of bore wells in Kesalapura-1 micro-watershed

Sl. No.	Particulars	LL (3)	MF (7)	SF (7)	SMF (10)	MDF (4)	LF (1)	All (32)
		N	N	N	N	N	N	N
1	Functioning	0	3	4	9	4	0	20

Source of irrigation: The data regarding the source of irrigation in Kesalapura-1 micro-watershed is presented in Table 19. The results indicate that, bore well was the major irrigation source in the micro water shed for 62.50 per cent of the farmers.

Table 19. Source of irrigation in Kesalapura-1 micro-watershed

Sl.No.	Particulars	LL (3)		MF (7)		SF (7)		SMF (10)		MDF (4)		LF (1)		All (32)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Bore Well	0	0.00	3	42.86	4	57.14	9	90.00	4	100.00	0	0.00	20	62.50

Depth of water (Avg in meters): The data regarding the depth of water in Kesalapura-1 micro-watershed is presented in Table 20. The results indicate that, the depth of bore well was found to be 44.10 meters.

Table 20. Depth of water (Avg in meters) in Kesalapura-1 micro-watershed

Sl.No.	Particulars	LL (3)	MF (7)	SF (7)	SMF (10)	MDF (4)	LF (1)	All (32)
1	Bore Well	0.00	23.95	44.85	67.97	62.48	0.00	44.10

Irrigated Area (ha): The data regarding the irrigated area (ha) in Kesalapura-1 micro-watershed is presented in Table 21. The results indicate that, marginal, small, semi medium and medium farmers had irrigated area of 1.21 ha, 7.65 ha, 12.01 ha and 11.48 ha respectively.

Table 21. Irrigated Area (ha) in Kesalapura-1 micro-watershed

Sl.No.	Particulars	LL (3)	MF (7)	SF (7)	SMF (10)	MDF (4)	LF (1)	All (32)
1	Kharif	0.00	1.21	2.79	11.20	11.48	0.00	26.69
2	Perennial Crops	0.00	0.00	3.64	0.00	0.00	0.00	3.64
3	Rabi	0.00	0.00	1.21	0.00	0.00	0.00	1.21
4	Summer	0.00	0.00	0.00	0.81	0.00	0.00	0.81
Total		0.00	1.21	7.65	12.01	11.48	0.00	32.36

Cropping pattern: The data regarding the cropping pattern in Kesalapura-1 micro-watershed is presented in Table 22. The results indicate that, farmers have grown sugarcane (15.91 ha), groundnut (4.45 ha), maize (7.03 ha), sunflower (2.86 ha), pearl millet (5.19 ha), bengalgram (2.05 ha), chilly (1.21 ha), cotton (0.4 ha), mulberry (0.03 ha) and jowar (0.4 ha). Marginal farmers have grown sugarcane, maize, sunflower, pearl millet, bengalgram, jowar and mulberry. Small farmers have grown sugarcane, maize, groundnut, sunflower, pearl millet, groundnut and cotton. Semi medium farmers have grown sugarcane, maize, bajra, pearl millet, bengalgram and chilly. Medium farmers have grown sugarcane, maize, groundnut, sunflower and chilly. Large farmers have grown bajra only.

Table 22. Cropping pattern in Kesalapura-1 micro-watershed (Area in ha)

Sl. No.	Particulars	LL (3)	MF (7)	SF (7)	SMF (10)	MDF (4)	LF (1)	All (32)
1	Kharif - Sugarcane	0	0.81	1.35	7.94	5.81	0	15.91
2	Kharif - Maize	0	1.36	1.21	1.62	2.83	0	7.03
3	Kharif - Bajra	0	0	0	2.93	0	3.24	6.17
4	Kharif - Groundnut	0	0	0.81	0	2.43	0	3.24
5	Kharif - Sunflower	0	0.81	0.83	0	1.21	0	2.86
6	Kharif - Pearl millet [bajra]	0	0	2.68	0	0	0	2.68
7	Kharif - Pearl millet (Sajje)	0	0.87	0	1.64	0	0	2.51
8	Rabi - Bengal gram	0	0.4	0	1.64	0	0	2.05
9	Kharif - Chilly	0	0	0	0.4	0.81	0	1.21
10	Rabi - Groundnut	0	0	1.21	0	0	0	1.21
11	Kharif - Cotton	0	0	0.4	0	0	0	0.4
12	Kharif - Jowar	0	0.4	0	0	0	0	0.4
13	Kharif - Mulberry	0	0.03	0	0	0	0	0.03
	Total	0	4.69	8.51	16.18	13.1	3.24	45.72

Cropping intensity: The data regarding the cropping intensity in Kesalapura-1 micro-watershed is presented in Table 23. The results indicate that, the cropping intensity in Kesalapura-1 micro-watershed was found to be 72.08 per cent. In case of marginal farmers it was 92.48 per cent, small farmers it was 94.18 per cent, in case of semi medium farmers it was 74.11, medium farmers it was 62.52 per cent and in case of large farmers it was 50 per cent.

Table 23. Cropping intensity (%) in Kesalapura-1 micro-watershed

Sl.No.	Particulars	LL (3)	MF (7)	SF(7)	SMF (10)	MDF (4)	LF (1)	All (32)
1	Cropping Intensity	0.00	91.48	94.18	74.11	62.52	50.00	72.08

Possession of Bank account and savings: The data regarding the cropping intensity in Kesalapura-1 micro-watershed is presented in Table 24. The results indicate that, 46.88 per cent of the households have bank account.

Table 24. Possession of Bank account and savings in Kesalapura-1 micro-watershed

Sl. No.	Particulars	LL (3)		MF (7)		SF (7)		SMF (10)		MDF (4)		LF (1)		All (32)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Account	0	0.00	5	71.43	2	28.57	5	50.00	3	75.00	0	0.00	15	46.88
2	Savings	0	0.00	0	0.00	0	0.00	0	0.00	0	0.00	0	0.00	0	0.00

Borrowing status: The data regarding the cropping intensity in Kesalapura-1 micro-watershed is presented in Table 25. The results indicate that, 46.88 per cent of the households have availed credit from different sources.

Table 25. Borrowing status in Kesalapura-1 micro-watershed

Sl.No.	Particulars	LL (3)		MF (7)		SF (7)		SMF (10)		MDF (4)		LF (1)		All (32)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Credit Availed	0	0.00	5	71.43	2	28.57	5	50.00	3	75.00	0	0.00	15	46.88

Cost of Cultivation of sugarcane: The data regarding the cost of cultivation of sugarcane in Kesalapura-1 micro-watershed is presented in Table 26. The results indicate that, the total cost of cultivation for sugarcane was Rs. 60569.99. The gross income realized by the farmers was Rs. 525781.85. The net income from sugarcane cultivation was Rs. 465211.86. Thus the benefit cost ratio was found to be 1:8.68.

Table 26. Cost of Cultivation of Sugarcane in Kesalapura-1 micro-watershed

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
I	Cost A1				
1	Hired Human Labour	Man days	62.99	11731.21	19.37
2	Bullock	Pairs/day	0.58	290.83	0.48
3	Tractor	Hours	5.12	3292.12	5.44
4	Machinery	Hours	0.27	214.07	0.35
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	2332.20	15797.11	26.08
6	Seed Inter Crop	Kgs.	0.00	0.00	0.00
7	FYM	Quintal	27.49	4281.70	7.07
8	Fertilizer + micronutrients	Quintal	6.81	5932.48	9.79
9	Pesticides (PPC)	Kgs / liters	0.60	482.28	0.80
10	Irrigation	Number	11.82	0.00	0.00
11	Repairs		0.00	0.00	0.00
12	Msc. Charges (Marketing costs etc)		0.00	0.00	0.00
13	Depreciation charges		0.00	2820.47	4.66
14	Land revenue and Taxes		0.00	3.16	0.01
II	Cost B1				
16	Interest on working capital			3179.30	5.25
17	Cost B1 = (Cost A1 + sum of 15 and 16)			48024.72	79.29
III	Cost B2				
18	Rental Value of Land			425.00	0.70
19	Cost B2 = (Cost B1 + Rental value)			48449.72	79.99
IV	Cost C1				
20	Family Human Labour		31.52	6613.32	10.92
21	Cost C1 = (Cost B2 + Family Labour)			55063.04	90.91
V	Cost C2				
22	Risk Premium			0.58	0.00
23	Cost C2 = (Cost C1 + Risk Premium)			55063.63	90.91
VI	Cost C3				
24	Managerial Cost			5506.36	9.09
25	Cost C3 = (Cost C2 + Managerial Cost)			60569.99	100.00
VII	Economics of the Crop				
a.	Main Product	a) Main Product (q)	806.83	525781.85	
		b) Main Crop Sales Price (Rs.)		651.67	
b.	Gross Income (Rs.)			525781.85	
c.	Net Income (Rs.)			465211.86	
d.	Cost per Quintal (Rs./q.)			75.07	
e.	Benefit Cost Ratio (BC Ratio)			1:8.68	

Cost of Cultivation of Groundnut: The data regarding the cost of cultivation of groundnut in Kesalapura-1 micro-watershed is presented in Table 27. The results indicate that, the total cost of cultivation for groundnut was Rs. 42208.67. The gross income realized by the farmers was Rs. 102827.47. The net income from groundnut cultivation was Rs. 60618.81. Thus the benefit cost ratio was found to be 1:2.44.

Table 27. Cost of Cultivation of Groundnut in Kesalapura-1 micro-watershed

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
I	Cost A1				
1	Hired Human Labour	Man days	22.02	4003.46	9.48
2	Bullock	Pairs/day	0.00	0.00	0.00
3	Tractor	Hours	3.09	2295.04	5.44
4	Machinery	Hours	0.82	658.67	1.56
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	94.68	11238.50	26.63
6	Seed Inter Crop	Kgs.	0.00	0.00	0.00
7	FYM	Quintal	12.97	3025.75	7.17
8	Fertilizer + micronutrients	Quintal	6.38	6298.50	14.92
9	Pesticides (PPC)	Kgs / liters	0.00	0.00	0.00
10	Irrigation	Number	4.94	0.00	0.00
11	Repairs		0.00	0.00	0.00
12	Msc. Charges (Marketing costs etc)		0.00	0.00	0.00
13	Depreciation charges		0.00	2063.69	4.89
14	Land revenue and Taxes		0.00	3.57	0.01
II	Cost B1				
16	Interest on working capital			2467.61	5.85
17	Cost B1 = (Cost A1 + sum of 15 and 16)			32054.78	75.94
III	Cost B2				
18	Rental Value of Land			377.78	0.90
19	Cost B2 = (Cost B1 + Rental value)			32432.56	76.84
IV	Cost C1				
20	Family Human Labour		31.08	5938.29	14.07
21	Cost C1 = (Cost B2 + Family Labour)			38370.85	90.91
V	Cost C2				
22	Risk Premium			0.67	0.00
23	Cost C2 = (Cost C1 + Risk Premium)			38371.52	90.91
VI	Cost C3				
24	Managerial Cost			3837.15	9.09
25	Cost C3 = (Cost C2 + Managerial Cost)			42208.67	100.00
VII	Economics of the Crop				
a.	Main Product	a) Main Product (q)	27.79	102813.75	
		b) Main Crop Sales Price (Rs.)		3700.00	
	By Product	e) Main Product (q)	0.41	13.72	
		f) Main Crop Sales Price (Rs.)		33.33	
b.	Gross Income (Rs.)			102827.47	
c.	Net Income (Rs.)			60618.81	
d.	Cost per Quintal (Rs./q.)			1518.98	
e.	Benefit Cost Ratio (BC Ratio)			1:2.44	

Cost of Cultivation of Maize: The data regarding the cost of cultivation of maize in Kesalapura-1 micro-watershed is presented in Table 28. The results indicate that, the total cost of cultivation for maize was Rs. 43481.62. The gross income realized by the farmers was Rs. 54611.46. The net income from maize cultivation was Rs. 11129.85. Thus the benefit cost ratio was found to be 1:1.26.

Table 28. Cost of Cultivation of Maize in Kesalapura-1 micro-watershed

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
I	Cost A1				
1	Hired Human Labour	Man days	31.52	5328.92	12.26
2	Bullock	Pairs/day	1.50	791.26	1.82
3	Tractor	Hours	2.05	1150.82	2.65
4	Machinery	Hours	0.31	282.29	0.65
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	22.47	4102.21	9.43
6	Seed Inter Crop	Kgs.	0.00	0.00	0.00
7	FYM	Quintal	15.62	11913.38	27.40
8	Fertilizer + micronutrients	Quintal	5.54	4602.85	10.59
9	Pesticides (PPC)	Kgs / liters	0.81	978.02	2.25
10	Irrigation	Number	3.29	0.00	0.00
11	Repairs		0.00	0.00	0.00
12	Msc. Charges (Marketing costs etc)		0.00	0.00	0.00
13	Depreciation charges		0.00	1572.49	3.62
14	Land revenue and Taxes		0.00	3.60	0.01
II	Cost B1				
16	Interest on working capital			2591.64	5.96
17	Cost B1 = (Cost A1 + sum of 15 and 16)			33317.48	76.62
III	Cost B2				
18	Rental Value of Land			354.17	0.81
19	Cost B2 = (Cost B1 + Rental value)			33671.65	77.44
IV	Cost C1				
20	Family Human Labour		27.82	5856.59	13.47
21	Cost C1 = (Cost B2 + Family Labour)			39528.24	90.91
V	Cost C2				
22	Risk Premium			0.50	0.00
23	Cost C2 = (Cost C1 + Risk Premium)			39528.74	90.91
VI	Cost C3				
24	Managerial Cost			3952.87	9.09
25	Cost C3 = (Cost C2 + Managerial Cost)			43481.62	100.00
VII	Economics of the Crop				
a.	Main Product	a) Main Product (q)	44.54	53168.67	
		b) Main Crop Sales Price (Rs.)		1193.75	
	By Product	e) Main Product (q)	2.73	1442.79	
		f) Main Crop Sales Price (Rs.)		528.75	
b.	Gross Income (Rs.)			54611.46	
c.	Net Income (Rs.)			11129.85	
d.	Cost per Quintal (Rs./q.)			976.25	
e.	Benefit Cost Ratio (BC Ratio)			1:1.26	

Cost of Cultivation of Bajra: The data regarding the cost of cultivation of bajra in Kesalapura-1 micro-watershed is presented in Table 29. The results indicate that, the total cost of cultivation for bajra was Rs. 19721.25. The gross income realized by the farmers was Rs. 22622.75. The net income from bajra cultivation was Rs. 2901.51. Thus the benefit cost ratio was found to be 1:1.15.

Table 29. Cost of Cultivation of Bajra in Kesalapura-1 micro-watershed

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
I	Cost A1				
1	Hired Human Labour	Man days	21.47	3794.60	19.24
2	Bullock	Pairs/day	0.25	158.42	0.80
3	Tractor	Hours	3.63	2383.82	12.09
4	Machinery	Hours	0.31	259.52	1.32
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	20.06	2547.17	12.92
6	Seed Inter Crop	Kgs.	0.00	0.00	0.00
7	FYM	Quintal	6.55	1870.40	9.48
8	Fertilizer + micronutrients	Quintal	2.48	2447.00	12.41
9	Pesticides (PPC)	Kgs / liters	0.58	582.55	2.95
10	Irrigation	Number	9.13	0.00	0.00
11	Repairs		0.00	0.00	0.00
12	Msc. Charges (Marketing costs etc)		0.00	0.00	0.00
13	Depreciation charges		0.00	789.97	4.01
14	Land revenue and Taxes		0.00	3.41	0.02
II	Cost B1				
16	Interest on working capital			893.77	4.53
17	Cost B1 = (Cost A1 + sum of 15 and 16)			15730.64	79.76
III	Cost B2				
18	Rental Value of Land			361.90	1.84
19	Cost B2 = (Cost B1 + Rental value)			16092.54	81.60
IV	Cost C1				
20	Family Human Labour		9.21	1834.86	9.30
21	Cost C1 = (Cost B2 + Family Labour)			17927.41	90.90
V	Cost C2				
22	Risk Premium			1.00	0.01
23	Cost C2 = (Cost C1 + Risk Premium)			17928.41	90.91
VI	Cost C3				
24	Managerial Cost			1792.84	9.09
25	Cost C3 = (Cost C2 + Managerial Cost)			19721.25	100.00
VII	Economics of the Crop				
a.	Main Product	a) Main Product (q)		18.95	22603.73
		b) Main Crop Sales Price (Rs.)			1192.86
	By Product	e) Main Product (q)		0.17	19.02
		f) Main Crop Sales Price (Rs.)			114.29
b.	Gross Income (Rs.)			22622.75	
c.	Net Income (Rs.)			2901.51	
d.	Cost per Quintal (Rs./q.)			1040.74	
e.	Benefit Cost Ratio (BC Ratio)			1:1.15	

Cost of Cultivation of Jowar: The data regarding the cost of cultivation of Jowar in Kesalapura-1 micro-watershed is presented in Table 30. The results indicate that, the total cost of cultivation for Jowar was Rs. 61598.57. The gross income realized by the farmers was Rs. 55328. The net income from Jowar cultivation was Rs. -6270.57. Thus the benefit cost ratio was found to be 1:0.9.

Table 30. Cost of Cultivation of Jowar in Kesalapura-1 micro-watershed

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
I	Cost A1				
1	Hired Human Labour	Man days	49.40	8398.00	13.63
2	Bullock	Pairs/day	2.47	1729.00	2.81
3	Tractor	Hours	4.94	4446.00	7.22
4	Machinery	Hours	0.00	0.00	0.00
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	17.29	1037.40	1.68
6	Seed Inter Crop	Kgs.	0.00	0.00	0.00
7	FYM	Quintal	4.94	5928.00	9.62
8	Fertilizer + micronutrients	Quintal	7.41	5112.90	8.30
9	Pesticides (PPC)	Kgs /liters	2.47	1976.00	3.21
10	Irrigation	Number	24.70	0.00	0.00
11	Repairs		0.00	0.00	0.00
12	Msc. Charges (Marketing costs etc)		0.00	0.00	0.00
13	Depreciation charges		0.00	16949.14	27.52
14	Land revenue and Taxes		0.00	3.29	0.01
II	Cost B1				
16	Interest on working capital			1686.64	2.74
17	Cost B1 = (Cost A1 + sum of 15 and 16)			47266.37	76.73
III	Cost B2				
18	Rental Value of Land			333.33	0.54
19	Cost B2 = (Cost B1 + Rental value)			47599.70	77.27
IV	Cost C1				
20	Family Human Labour		39.52	8398.00	13.63
21	Cost C1 = (Cost B2 + Family Labour)			55997.70	90.91
V	Cost C2				
22	Risk Premium			1.00	0.00
23	Cost C2 = (Cost C1 + Risk Premium)			55998.70	90.91
VI	Cost C3				
24	Managerial Cost			5599.87	9.09
25	Cost C3 = (Cost C2 + Managerial Cost)			61598.57	100.00
VII	Economics of the Crop				
a.	Main Product	a) Main Product (q)		39.52	55328.00
		b) Main Crop Sales Price (Rs.)			1400.00
b.	Gross Income (Rs.)			55328.00	
c.	Net Income (Rs.)			-6270.57	
d.	Cost per Quintal (Rs./q.)			1558.67	
e.	Benefit Cost Ratio (BC Ratio)			1:0.9	

Cost of cultivation of chilly: The data regarding the cost of cultivation of chilly in Kesalapura-1 micro-watershed is presented in Table 31. The results indicate that, the total cost of cultivation for chilly was Rs. 223901.85. The gross income realized by the farmers was Rs. 322643.75. The net income from chilly cultivation was Rs. 98741.90. Thus the benefit cost ratio was found to be 1:1.44.

Table 31. Cost of Cultivation of chilly in Kesalapura-1 micro-watershed

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
I	Cost A1				
1	Hired Human Labour	Man days	86.45	19389.50	8.66
2	Bullock	Pairs/day	1.24	0.00	0.00
3	Tractor	Hours	4.94	3458.00	1.54
4	Machinery	Hours	2.47	4446.00	1.99
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	311.22	115472.50	51.57
6	Seed Inter Crop	Kgs.	0.00	0.00	0.00
7	FYM	Quintal	4.94	9880.00	4.41
8	Fertilizer + micronutrients	Quintal	8.65	6928.35	3.09
9	Pesticides (PPC)	Kgs / liters	2.47	1976.00	0.88
10	Irrigation	Number	24.70	0.00	0.00
11	Repairs		0.00	0.00	0.00
12	Msc. Charges (Marketing costs etc)		0.00	0.00	0.00
13	Depreciation charges		0.00	3745.14	1.67
14	Land revenue and Taxes		0.00	3.71	0.00
II	Cost B1				
16	Interest on working capital			16110.94	7.20
17	Cost B1 = (Cost A1 + sum of 15 and 16)			181410.13	81.02
III	Cost B2				
18	Rental Value of Land			400.00	0.18
19	Cost B2 = (Cost B1 + Rental value)			181810.13	81.20
IV	Cost C1				
20	Family Human Labour		59.28	21736.00	9.71
21	Cost C1 = (Cost B2 + Family Labour)			203546.13	90.91
V	Cost C2				
22	Risk Premium			1.00	0.00
23	Cost C2 = (Cost C1 + Risk Premium)			203547.13	90.91
VI	Cost C3				
24	Managerial Cost			20354.71	9.09
25	Cost C3 = (Cost C2 + Managerial Cost)			223901.85	100.00
VII	Economics of the Crop				
a.	Main Product	a) Main Product (q)	117.33	322643.75	
		b) Main Crop Sales Price (Rs.)		2750.00	
b.	Gross Income (Rs.)			322643.75	
c.	Net Income (Rs.)			98741.90	
d.	Cost per Quintal (Rs./q.)			1908.39	
e.	Benefit Cost Ratio (BC Ratio)			1:1.44	

Cost of cultivation of Bengalgram: The data regarding the cost of cultivation of Bengalgram in Kesalapura-1 micro-watershed is presented in Table 32. The results indicate that, the total cost of cultivation for Bengalgram was Rs. 52147.87. The gross income realized by the farmers was Rs. 97556.48. The net income from Bengalgram cultivation was Rs. 45408.62. Thus the benefit cost ratio was found to be 1:1.87.

Table 32. Cost of Cultivation of Bengalgram in Kesalapura-1 micro-watershed

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
I	Cost A1				
1	Hired Human Labour	Man days	34.43	5288.60	10.14
2	Bullock	Pairs/day	3.71	2593.50	4.97
3	Tractor	Hours	3.69	2953.05	5.66
4	Machinery	Hours	0.00	0.00	0.00
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	52.44	8817.78	16.91
6	Seed Inter Crop	Kgs.	0.00	0.00	0.00
7	FYM	Quintal	0.00	0.00	0.00
8	Fertilizer + micronutrients	Quintal	7.37	7223.84	13.85
9	Pesticides (PPC)	Kgs / liters	1.54	1231.35	2.36
10	Irrigation	Number	15.39	0.00	0.00
11	Repairs		0.00	0.00	0.00
12	Msc. Charges (Marketing costs etc)		0.00	0.00	0.00
13	Depreciation charges		0.00	11215.91	21.51
14	Land revenue and Taxes		0.00	2.88	0.01
II	Cost B1				
16	Interest on working capital			2072.94	3.98
17	Cost B1 = (Cost A1 + sum of 15 and 16)			41399.84	79.39
III	Cost B2				
18	Rental Value of Land			333.33	0.64
19	Cost B2 = (Cost B1 + Rental value)			41733.17	80.03
IV	Cost C1				
20	Family Human Labour		26.76	5672.48	10.88
21	Cost C1 = (Cost B2 + Family Labour)			47405.65	90.91
V	Cost C2				
22	Risk Premium			1.50	0.00
23	Cost C2 = (Cost C1 + Risk Premium)			47407.15	90.91
VI	Cost C3				
24	Managerial Cost			4740.72	9.09
25	Cost C3 = (Cost C2 + Managerial Cost)			52147.87	100.00
VII	Economics of the Crop				
a.	Main Product	a) Main Product (q)	21.21	97556.48	
		b) Main Crop Sales Price (Rs.)		4600.00	
b.	Gross Income (Rs.)			97556.48	
c.	Net Income (Rs.)			45408.62	
d.	Cost per Quintal (Rs./q.)			2458.89	
e.	Benefit Cost Ratio (BC Ratio)			1:1.87	

Cost of cultivation of Sunflower: The data regarding the cost of cultivation of Sunflower in Kesalapura-1 micro-watershed is presented in Table 33. The results indicate that, the total cost of cultivation for Sunflower was Rs. 32268.06. The gross income realized by the farmers was Rs. 64321.25. The net income from Sunflower cultivation was Rs. 32053.20. Thus the benefit cost ratio was found to be 1:1.99.

Table 33. Cost of Cultivation of Sunflower in Kesalapura-1 micro-watershed

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
I	Cost A1				
1	Hired Human Labour	Man days	34.35	6312.16	19.56
2	Bullock	Pairs/day	0.00	0.00	0.00
3	Tractor	Hours	3.81	2469.00	7.65
4	Machinery	Hours	0.00	0.00	0.00
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	8.19	3302.61	10.23
6	Seed Inter Crop	Kgs.	0.00	0.00	0.00
7	FYM	Quintal	13.55	3167.83	9.82
8	Fertilizer + micronutrients	Quintal	4.47	3664.50	11.36
9	Pesticides (PPC)	Kgs / liters	1.20	959.22	2.97
10	Irrigation	Number	9.59	0.00	0.00
11	Repairs		0.00	0.00	0.00
12	Msc. Charges (Marketing costs etc)		0.00	0.00	0.00
13	Depreciation charges		0.00	2855.89	8.85
14	Land revenue and Taxes		0.00	2.74	0.01
II	Cost B1				
16	Interest on working capital			1331.38	4.13
17	Cost B1 = (Cost A1 + sum of 15 and 16)			24065.34	74.58
III	Cost B2				
18	Rental Value of Land			400.00	1.24
19	Cost B2 = (Cost B1 + Rental value)			24465.34	75.82
IV	Cost C1				
20	Family Human Labour		21.79	4868.59	15.09
21	Cost C1 = (Cost B2 + Family Labour)			29333.93	90.91
V	Cost C2				
22	Risk Premium			0.67	0.00
23	Cost C2 = (Cost C1 + Risk Premium)			29334.60	90.91
VI	Cost C3				
24	Managerial Cost			2933.46	9.09
25	Cost C3 = (Cost C2 + Managerial Cost)			32268.06	100.00
VII	Economics of the Crop				
a.	Main Product	a) Main Product (q)		18.92	64321.25
		b) Main Crop Sales Price (Rs.)			3400.00
b.	Gross Income (Rs.)			64321.25	
c.	Net Income (Rs.)			32053.20	
d.	Cost per Quintal (Rs./q.)			1705.68	
e.	Benefit Cost Ratio (BC Ratio)			1:1.99	

Cost of cultivation of Mulberry: The data regarding the cost of cultivation of Mulberry in Kesalapura-1 micro-watershed is presented in Table 34. The results indicate that, the total cost of cultivation for Mulberry was Rs. 849677.77. The gross income realized by the farmers was Rs. 2161250.05. The net income from Mulberry cultivation was Rs. 1311572.28. Thus the benefit cost ratio was found to be 1:2.54.

Table 34. Cost of Cultivation of Mulberry in Kesalapura-1 micro-watershed

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
I	Cost A1				
1	Hired Human Labour	Man days	1358.50	213037.50	25.07
2	Bullock	Pairs/day	123.50	74100.00	8.72
3	Tractor	Hours	0.00	0.00	0.00
4	Machinery	Hours	0.00	0.00	0.00
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	61.75	111150.00	13.08
6	Seed Inter Crop	Kgs.	0.00	0.00	0.00
7	FYM	Quintal	926.25	111150.00	13.08
8	Fertilizer + micronutrients	Quintal	61.75	74100.00	8.72
9	Pesticides (PPC)	Kgs / liters	0.00	0.00	0.00
10	Irrigation	Number	92.63	0.00	0.00
11	Repairs		0.00	0.00	0.00
12	Msc. Charges (Marketing costs etc)		0.00	0.00	0.00
13	Depreciation charges		0.00	1636.38	0.19
14	Land revenue and Taxes		0.00	4.94	0.00
II	Cost B1				
16	Interest on working capital			35568.00	4.19
17	Cost B1 = (Cost A1 + sum of 15 and 16)			620746.83	73.06
III	Cost B2				
18	Rental Value of Land			400.00	0.05
19	Cost B2 = (Cost B1 + Rental value)			621146.83	73.10
IV	Cost C1				
20	Family Human Labour		741.00	151287.50	17.81
21	Cost C1 = (Cost B2 + Family Labour)			772434.33	90.91
V	Cost C2				
22	Risk Premium			0.00	0.00
23	Cost C2 = (Cost C1 + Risk Premium)			772434.33	90.91
VI	Cost C3				
24	Managerial Cost			77243.43	9.09
25	Cost C3 = (Cost C2 + Managerial Cost)			849677.77	100.00
VII	Economics of the Crop				
a.	Main Product	a) Main Product (q)		3087.50	2161250.05
		b) Main Crop Sales Price (Rs.)			700.00
b.	Gross Income (Rs.)				2161250.05
c.	Net Income (Rs.)				1311572.28
d.	Cost per Quintal (Rs./q.)				275.20
e.	Benefit Cost Ratio (BC Ratio)				1:2.54

Cost of cultivation of cotton: The data regarding the cost of cultivation of cotton in Kesalapura-1 micro-watershed is presented in Table 35. The results indicate that, the total cost of cultivation for cotton was Rs. 48650.76. The gross income realized by the farmers was Rs. 118560. The net income from cotton cultivation was Rs. 69909.24. Thus the benefit cost ratio was found to be 1:2.44.

Table 35. Cost of Cultivation of cotton in Kesalapura-1 micro-watershed

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
I	Cost A1				
1	Hired Human Labour	Man days	46.93	7286.50	14.98
2	Bullock	Pairs/day	2.47	1235.00	2.54
3	Tractor	Hours	2.47	1482.00	3.05
4	Machinery	Hours	2.47	1976.00	4.06
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	7.41	2371.20	4.87
6	Seed Inter Crop	Kgs.	0.00	0.00	0.00
7	FYM	Quintal	2.47	2470.00	5.08
8	Fertilizer + micronutrients	Quintal	4.94	3211.00	6.60
9	Pesticides (PPC)	Kgs / liters	2.47	3458.00	7.11
10	Irrigation	Number	0.00	0.00	0.00
11	Repairs		0.00	0.00	0.00
12	Msc. Charges (Marketing costs etc)		0.00	0.00	0.00
13	Depreciation charges		0.00	0.00	0.00
14	Land revenue and Taxes		0.00	2.47	0.01
II	Cost B1				
16	Interest on working capital			1381.46	2.84
17	Cost B1 = (Cost A1 + sum of 15 and 16)			24873.63	51.13
III	Cost B2				
18	Rental Value of Land			333.33	0.69
19	Cost B2 = (Cost B1 + Rental value)			25206.97	51.81
IV	Cost C1				
20	Family Human Labour		93.86	19019.00	39.09
21	Cost C1 = (Cost B2 + Family Labour)			44225.97	90.90
V	Cost C2				
22	Risk Premium			2.00	0.00
23	Cost C2 = (Cost C1 + Risk Premium)			44227.97	90.91
VI	Cost C3				
24	Managerial Cost			4422.80	9.09
25	Cost C3 = (Cost C2 + Managerial Cost)			48650.76	100.00
VII	Economics of the Crop				
a.	Main Product	a) Main Product (q)	24.70	118560.00	
		b) Main Crop Sales Price (Rs.)		4800.00	
b.	Gross Income (Rs.)			118560.00	
c.	Net Income (Rs.)			69909.24	
d.	Cost per Quintal (Rs./q.)			1969.67	
e.	Benefit Cost Ratio (BC Ratio)			1:2.44	

Adequacy of fodder: The data regarding the adequacy of fodder in Kesalapura-1 micro-watershed is presented in Table 36. The results indicate that, 65.63 per cent of the households opined that dry fodder was adequate and 40.63 per cent opined that green fodder was adequate.

Table 36. Adequacy of fodder in Kesalapura-1 micro-watershed

Sl. No.	Particulars	LL (3)		MF (7)		SF (7)		SMF (10)		MDF (4)		LF (1)		All (32)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Adequate-Dry Fodder	0	0.00	4	57.14	7	100.00	8	80.00	1	25.00	1	100.00	21	65.63
2	Adequate-Green Fodder	0	0.00	1	14.29	5	71.43	6	60.00	0	0.00	1	100.00	13	40.63

Average annual gross income: The data regarding the average annual gross income in Kesalapura-1 micro-watershed is presented in Table 37. The results indicate that the average annual gross income was Rs. 37667 for landless farmers, for marginal farmers it was Rs. 181857, for small farmers it was Rs. 118516, for semi medium farmers it was Rs. 225550, for medium farmers it was Rs. 355000 and for large farmers it was Rs.120000.

Table 37. Average annual gross income in Kesalapura-1 micro-watershed

(Avg value in Rs.)

Sl.No.	Particulars	LL (3)	MF (7)	SF (7)	SMF (10)	MDF (4)	LF (1)	All (32)
1	Service/salary	0	14286	0	0	0	0	3125
2	Business	0	57143	0	0	0	0	12500
3	Wage	35000	10000	27857	8000	0	20000	14688
4	Agriculture	0	98286	86129	214950	355000	100000	155013
5	Farm income	0	714	0	0	0	0	156
6	Dairy Farm	2667	1429	4530	2600	0	0	2366
	Income(Rs.)	37667	181857	118516	225550	355000	120000	187847

Average annual expenditure: The data regarding the average annual expenditure in Kesalapura-1 micro-watershed is presented in Table 38. The results indicate that the average annual expenditure is Rs. 33462. For landless households it was Rs. 6889, for marginal farmers it was Rs. 68184, for small farmers it was Rs. 16335, for semi medium farmers it was Rs. 16350, for medium farmers it was Rs. 58750 and for large farmers it was Rs.60000.

Table 38. Average annual expenditure in Kesalapura-1 micro-watershed

(Avg value in Rs.)

Sl.No.	Particulars	LL (3)	MF (7)	SF (7)	SMF (10)	MDF (4)	LF (1)	All (32)
1	Service/salary	0	50000	0	0	0	0	1563
2	Business	0	360000	0	0	0	0	11250
3	Wage	16667	0	41250	25000	0	10000	8594
4	Agriculture	0	59286	61429	132000	235000	50000	98594
5	Farm income	0	3000	0	0	0	0	94
6	Dairy Farm	4000	5000	11667	6500	0	0	1781
	Total	20667	477286	114345	163500	235000	60000	1070798
	Average	6889	68184	16335	16350	58750	60000	33462

Horticulture species grown: The data regarding horticulture species grown in Kesalapura-1 micro-watershed is presented in Table 39. The results indicate that, households have planted 69 coconut trees, 2 lemon trees, 2 lime trees and 7 mango trees in the field and 8 coconut trees in the backyard.

Table 39. Horticulture species grown in Kesalapura-1 micro-watershed

Sl.No.	Particulars	LL (3)		MF (7)		SF (7)		SMF (10)		MDF (4)		LF (1)		All (32)	
		F	B	F	B	F	B	F	B	F	B	F	B	F	B
1	Coconut	0	0	4	1	37	1	24	6	4	0	0	0	69	8
2	Lemon	0	0	0	0	2	0	0	0	0	0	0	0	2	0
3	lime	0	0	0	0	0	0	2	0	0	0	0	0	2	0
4	Mango	0	0	1	0	3	0	3	0	0	0	0	0	7	0

*F= Field B=Back Yard

Forest species grown: The data regarding forest species grown in Kesalapura-1 micro-watershed is presented in Table 40. The results indicate that, households have planted 75 neem tree, 1 banyan tree and 29 teak trees in their fields.

Table 40. Forest species grown in Kesalapura-1 micro-watershed

Sl.No.	Particulars	LL (3)		MF (7)		SF (7)		SMF (10)		MDF (4)		LF (1)		All (32)	
		F	B	F	B	F	B	F	B	F	B	F	B	F	B
1	Teak	0	0	0	0	2	0	27	0	0	0	0	0	29	0
2	Neem	0	0	3	1	13	5	34	3	21	0	4	0	75	9
3	Banyan	0	0	0	0	0	0	0	0	1	0	0	0	1	0

*F= Field B=Back Yard

Marketing of the agricultural produce: The data regarding marketing of the agricultural produce in Kesalapura-1 micro-watershed is presented in Table 41. The results indicated that, cotton, maize, mulberry, sugarcane and sunflower were sold to the extent of 100 per cent. Bajra was sold to the extent of 86.67 per cent, Bengal gram to the extent of 54.55 per cent, chilly to the extent of 94.29 per cent, groundnut to the extent of 52.38 per cent and jowar was sold to the extent of 68.75 per cent.

Table 41. Marketing of the agricultural produce in Kesalapura-1 micro-watershed

Sl.No	Crops	Output obtained (q)	Output retained (q)	Output sold (q)	Output sold (%)	Avg. Price obtained (Rs/q)
1	Bajra	240	32	208	86.67	1192.86
2	Bengalgram	33	15	18	54.55	4600.0
3	Chilly	175	10	165	94.29	2750.0
4	Cotton	10	0	10	100.00	4800.0
5	Groundnut	105	50	55	52.38	3700.0
6	Jowar	16	5	11	68.75	1400.0
7	Maize	361	0	361	100.00	1193.75
8	Mulberry	100	0	100	100.00	700.0
9	Sugarcane	10461	0	10461	100.00	651.67
10	Sunflower	58	0	58	100.00	3400.0

Marketing Channels used for sale of agricultural produce: The data regarding marketing channels used for sale of agricultural produce in Kesalapura-1 micro-watershed is presented in Table 42. The results indicated that, about 18.75 per cent of the famers have sold their produce in regulated markets, 53.13 per cent of the farmers have sold to local/village merchants and another 53.13 per cent of the farmers have sold to agents/traders.

Table 42. Marketing Channels used for sale of agricultural produce in Kesalapura-1 micro-watershed

Sl.No.	Particulars	LL (3)		MF (7)		SF (7)		SMF (10)		MDF (4)		LF (1)		All (32)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Agent/Traders	0	0.00	3	42.86	7	100.00	6	60.00	1	25.00	0	0.00	17	53.13
2	Local/village Merchant	0	0.00	4	57.14	0	0.00	5	50.00	7	175.00	1	100.00	17	53.13
3	Regulated Market	0	0.00	2	28.57	2	28.57	2	20.00	0	0.00	0	0.00	6	18.75

Mode of transport of agricultural produce: The data regarding mode of transport of agricultural produce in Kesalapura-1 micro-watershed is presented in Table 43. The results indicated that, 87.50 per cent of the households have used tractor, 25 per cent of the households have used cart, 9.38 per cent of the households have used truck and 3.13 per cent have carried head loads as a mode of transportation for their agricultural produce.

Table 43. Mode of transport of agricultural produce in Kesalapura-1 micro-watershed

Sl.No.	Particulars	LL (3)		MF (7)		SF (7)		SMF (10)		MDF (4)		LF (1)		All (32)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Head Load	0	0.00	0	0.00	1	14.29	0	0.00	0	0.00	0	0.00	1	3.13
2	Cart	0	0.00	3	42.86	4	57.14	1	10.00	0	0.00	0	0.00	8	25.00
3	Tractor	0	0.00	6	85.71	3	42.86	11	110.00	7	175.00	1	100.00	28	87.50
4	Truck	0	0.00	0	0.00	1	14.29	1	10.00	1	25.00	0	0.00	3	9.38

Incidence of soil and water erosion problems: The data regarding incidence of soil and water erosion problems in Kesalapura-1 micro-watershed is presented in Table 44. The results indicated that, 46.88 per cent of the households have experienced soil and water erosion problems in the farm i.e., 71.43 per cent of the marginal farmers, 28.5 per cent of the small farmers, 40 per cent of semi medium farmers and 75 per cent of medium farmers and 100 per cent of large farmers have experienced soil and water erosion problems.

Table 44. Incidence of soil and water erosion problems in Kesalapura-1 micro-watershed

Sl. No.	Particulars	LL (3)		MF (7)		SF (7)		SMF (10)		MDF (4)		LF (1)		All (32)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Soil and water erosion problems in the farm	0	0.00	5	71.43	2	28.5	4	40.00	3	75.00	1	100	15	46.88

Interest shown towards soil testing: The data regarding incidence of soil and water erosion problems in Kesalapura-1 micro-watershed is presented in Table 45. The results indicated that, 50 per cent have shown interest in soil test.

Table 45. Interest shown towards soil testing in Kesalapura-1 micro-watershed

Sl. No.	Particulars	LL (3)		MF (7)		SF (7)		SMF (10)		MDF (4)		LF (1)		All (32)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Interest in soil test	0	0.00	5	71.43	2	28.57	5	50.00	3	75.00	1	100.00	16	50.00

Usage pattern of fuel for domestic use: The data regarding usage pattern of fuel for domestic use in Kesalapura-1 micro-watershed is presented in Table 46. The results indicated that, 53.13 per cent of the households used firewood and 46.88 per cent have used LPG as a source of fuel.

Table 46. Usage pattern of fuel for domestic use in Kesalapura-1 micro-watershed

Sl. No.	Particulars	LL (3)		MF (7)		SF (7)		SMF (10)		MDF (4)		LF (1)		All (32)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Fire Wood	3	100.00	4	57.14	4	57.14	4	40.00	2	50.00	0	0.00	17	53.13
2	LPG	0	0.00	3	42.86	3	42.86	6	60.00	2	50.00	1	100.00	15	46.88

Source of drinking water: The data regarding source of drinking water in Kesalapura-1 micro-watershed is presented in Table 47. The results indicated that, bore well was the major source of drinking water for 34.38 per cent of the households and piped supply was the source of drinking water for 59.38 per cent of the households in the micro watershed.

Table 47. Source of drinking water in Kesalapura-1 micro-watershed

Sl.No.	Particulars	LL (3)		MF (7)		SF (7)		SMF (10)		MDF (4)		LF (1)		All (32)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Piped supply	2	66.67	6	85.71	2	28.57	5	50.00	3	75.00	1	100.00	19	59.38
2	Bore Well	0	0.00	1	14.29	6	85.71	4	40.00	0	0.00	0	0.00	11	34.38

Source of light: The data regarding source of light in Kesalapura-1 micro-watershed is presented in Table 48. The results indicated that, Electricity was the major source of light for 100 per cent of the households in micro watershed.

Table 48. Source of light in Kesalapura-1 micro-watershed

Sl.No.	Particulars	LL (3)		MF (7)		SF (7)		SMF (10)		MDF (4)		LF (1)		All (32)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Electricity	3	100.00	7	100.00	7	100.00	10	100.00	4	100.00	1	100.00	32	100.00

Table 49. Existence of Sanitary toilet facility in Kesalapura-1 micro-watershed

Sl.No.	Particulars	LL (3)		MF (7)		SF (7)		SMF (10)		MDF (4)		LF (1)		All (32)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Sanitary toilet facility	1	33.33	3	42.86	3	42.86	2	20.00	4	100.00	1	100.00	14	43.75

Existence of Sanitary toilet facility: The data regarding existence of sanitary toilet facility in Kesalapura-1 micro-watershed is presented in Table 49. The results indicated that, 43.75 per cent of the households possess sanitary toilet.

Possession of PDS card: The data regarding possession of PDS card in Kesalapura-1 micro-watershed is presented in Table 50. The results indicated that, 90.63 per cent of the sampled households possessed BPL card and 9.38 per cent of the households did not possess any PDS cards.

Table 50. Possession of PDS card in Kesalapura-1 micro-watershed

Sl.No.	Particulars	LL (3)		MF (7)		SF (7)		SMF (10)		MDF (4)		LF (1)		All (32)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	BPL	3	100	6	85.71	7	100	9	90	3	75	1	100	29	90.63
2	Not Possessed	0	0	1	14.29	0	0	1	10	1	25	0	0	3	9.38

Participation in NREGA program: The data regarding participation in NREGA programme in Kesalapura-1 micro-watershed is presented in Table 51. The results indicated that, 46.88 per cent of the households participated in NREGA programme.

Table 51. Participation in NREGA programme in Kesalapura-1 micro-watershed

Sl. No.	Particulars	LL (3)		MF (7)		SF (7)		SMF (10)		MDF (4)		LF (1)		All (32)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Participation in NREGA programme	3	100.00	2	28.57	2	28.57	5	50.00	2	50.00	1	100.00	15	46.88

Adequacy of food items: The data regarding adequacy of food items in Kesalapura-1 micro-watershed is presented in Table 52. The results indicated that, cereals were adequate for 100 per cent of the households, pulses were adequate for 62.50 per cent, oilseeds were adequate for 68.75 per cent, vegetables were adequate for 62.50 per cent, fruits were adequate for 6.25 per cent, milk was adequate for 90.63 per cent, eggs were adequate for 56.25 per cent and meat were adequate for 46.88 per cent.

Table 52. Adequacy of food items in Kesalapura-1 micro-watershed

Sl.No.	Particulars	LL (3)		MF (7)		SF (7)		SMF (10)		MDF (4)		LF (1)		All (32)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Cereals	3	100.00	7	100.00	7	100.00	10	100.00	4	100.00	1	100.00	32	100.00
2	Pulses	3	100.00	5	71.43	2	28.57	5	50.00	4	100.00	1	100.00	20	62.50
3	Oilseed	2	66.67	5	71.43	6	85.71	8	80.00	1	25.00	0	0.00	22	68.75
4	Vegetables	0	0.00	3	42.86	6	85.71	7	70.00	3	75.00	1	100.00	20	62.50
5	Fruits	1	33.33	1	14.29	0	0.00	0	0.00	0	0.00	0	0.00	2	6.25
6	Milk	2	66.67	5	71.43	7	100.00	10	100.00	4	100.00	1	100.00	29	90.63
7	Egg	0	0.00	6	85.71	6	85.71	4	40.00	2	50.00	0	0.00	18	56.25
8	Meat	1	33.33	3	42.86	5	71.43	4	40.00	2	50.00	0	0.00	15	46.88

Response on Inadequacy of food items: The data regarding inadequacy of food items in Kesalapura-1 micro-watershed is presented in Table 53. The results indicated that, pulses were inadequate for 37.50 per cent of the households, oilseeds were inadequate for 31.25

per cent, vegetables were inadequate for 37.50 per cent, fruits were inadequate for 53.13 per cent, eggs were inadequate for 28.13 per cent, milk was inadequate for 9.38 per cent of the households and meat was inadequate for 37.50 per cent of the households.

Table 53. Response on Inadequacy of food items in Kesalapura-1 micro-watershed

Sl.No.	Particulars	LL (3)		MF (7)		SF (7)		SMF (10)		MDF (4)		LF (1)		All (32)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Pulses	0	0.00	2	28.57	5	71.43	5	50.00	0	0.00	0	0.00	12	37.50
2	Oilseed	1	33.33	2	28.57	1	14.29	2	20.00	3	75.00	1	100.00	10	31.25
3	Vegetables	3	100.00	4	57.14	1	14.29	3	30.00	1	25.00	0	0.00	12	37.50
4	Fruits	2	66.67	4	57.14	2	28.57	5	50.00	3	75.00	1	100.00	17	53.13
5	Milk	1	33.33	2	28.57	0	0.00	0	0.00	0	0.00	0	0.00	3	9.38
6	Egg	3	100.00	0	0.00	0	0.00	4	40.00	1	25.00	1	100.00	9	28.13
7	Meat	2	66.67	3	42.86	1	14.29	4	40.00	1	25.00	1	100.00	12	37.50

Farming constraints: The data regarding farming constraints experienced by households in Kesalapura-1 micro-watershed is presented in Table 54. The results indicated that, lower fertility status of the soil was the constraint experienced by 50 per cent of the households, wild animal menace on farm field (59.38%), frequent incidence of pest and diseases (71.88%), inadequacy of irrigation water (31.25%), high cost of fertilizers and plant protection chemicals (84.38%), high rate of interest on credit (71.88%), low price for the agricultural commodities (43.75%), lack of marketing facilities in the area (65.63%), lack of transport for safe transport of the agricultural produce to the market (43.75%), inadequate extension services (9.38%), less rainfall (53.13%) and source of agri-technology information (15.63%).

Table 54. Farming constraints Experienced in Kesalapura-1 micro-watershed

Sl. No.	Particulars	MF (7)		SF (7)		SMF (10)		MDF (4)		LF (1)		All (32)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Lower fertility status of the soil	5	71.43	2	28.57	5	50	3	75	1	100	16	50
2	Wild animal menace on farm field	4	57.14	5	71.43	6	60	3	75	1	100	19	59.38
3	Frequent incidence of pest and diseases	6	85.71	4	57.14	8	80	4	100	1	100	23	71.88
4	Inadequacy of irrigation water	1	14.29	1	14.29	4	40	3	75	1	100	10	31.25
5	High cost of Fertilizers and plant protection chemicals	6	85.71	5	71.43	10	100	5	125	1	100	27	84.38
6	High rate of interest on credit	6	85.71	3	42.86	9	90	4	100	1	100	23	71.88
7	Low price for the agricultural commodities	5	71.43	1	14.29	4	40	3	75	1	100	14	43.75
8	Lack of marketing facilities in the area	6	85.71	1	14.29	10	100	3	75	1	100	21	65.63
9	Inadequate extension services	1	14.29	0	0	1	10	0	0	1	100	3	9.38
10	Lack of transport for safe transport of the Agril produce to the market.	2	28.57	2	28.57	6	60	3	75	1	100	14	43.75
11	Less rainfall	2	28.57	9	128.57	5	50	1	25	0	0	17	53.13
12	Source of Agri-technology information(Newspaper/TV/Mobile)	0	0	1	14.29	3	30	1	25	0	0	5	15.63

SUMMARY

In order to assess the socio-economic condition of the farmers in the watershed a comprehensive questionnaire was prepared. Major components such as demographic conditions, migration details, food consumption and family expenditure pattern, material possession, land holding, land use management, cropping pattern, cost of cultivation of crops, livestock management. The statistical components such as frequency and percentage were used to analyze the data. About 32 households located in the micro watershed were interviewed for the survey.

The data indicated that there were 82 (50.31%) men and 81 (49.69%) women among the sampled households. The average family size of landless farmers' was 4.67, marginal farmers' was 5.71, small farmers' was 6.29, semi medium farmers' was 4.40, medium farmers' was 4 and large farmers' was 5.

The data indicated that, 30 (18.40%) people were in 0-15 years of age, 71 (43.56%) were in 16-35 years of age, 44 (26.99%) were in 36-60 years of age and 18 (11.04%) were above 61 years of age.

The results indicated that Kesalapura-1 had 20.86 per cent illiterates, 36.81 per cent of them had primary school education, 9.82 per cent of them had middle school education, 20.25 per cent of them had high school education, 4.91 per cent of them had PUC education, 0.61 per cent of them did ITI, 4.29 per cent of them had degree education and 0.61 per cent did Maters.

The results indicate that, 84.38 per cent of households were practicing agriculture, 12.50 per cent of the households were agricultural labourers and 3.13 per cent were in government service. The results indicate that agriculture was the major occupation for 62.58 per cent of the household members, 14.72 per cent were agricultural laborers, 0.61 per cent was in government service, 20.25 per cent were students, 1.23 per cent were children and 0.61 per cent were housewives.

The results show that 99.39 per cent of the population in the micro watershed has not participated in any local institutions and only 0.61 per cent have participated in raitha sangha.

The results indicate that 3.13 per cent of the households possess thatched house, 71.88 per cent of the households possess Katcha house and 25 per cent of the households possess pucca/RCC house.

The results show that 65.63 per cent of the households possess TV, 75 per cent of the households possess Mixer grinder, 12.50 per cent of the households possess bicycle, 65.63 per cent of the households possess motor cycle, 6.25 per cent possess refrigerator, another 6.25 per cent possess car/four wheel and 93.75 per cent of the households possess mobile phones. The results show that the average value of television was Rs. 2595, mixer

grinder was Rs.1362, bicycle was Rs. 12850, motor cycle was Rs.33272, refrigerator was Rs.13500, car/wheeler was Rs.150000 and mobile phone was Rs.1264.

About 6.25 per cent of the households possess bullock cart, 21.88 per cent of the households possess plough, 3.13 per cent of them possess seed/fertilizer drill, 9.38 per cent of them possess power tiller, 37.50 per cent of them possess tractor, 37.50 per cent of them possess sprayer, 9.38 per cent of them possess sprinkler, 12.50 per cent possess harvester, 50 per cent of them possess chaff cutter, 18.75 per cent of them possess earth remover/duster and 71.88 per cent of them possess weeder. The results show that the average value of bullock cart was Rs.16500, the average value of plough was Rs.823, seed/fertilizer drill was Rs.25000, irrigation pump was Rs.21250, power tiller was Rs.15000, tractor was Rs.296000, the average value of sprayer was Rs.1900, weeder was Rs.24, harvester was Rs.13333, earth remover/duster was Rs.19500 and chaff cutter was Rs.817.

The results indicate that, 21.88 per cent of the households possess bullocks, crossbred cow and buffalo. About 43.75 per cent of the households possess local cow and 18.75 per cent of the households possess poultry birds.

The results indicate that, average own labour men available in the micro watershed was 2.07, average own labour (women) available was 1.93, average hired labour (men) available was 7.79 and average hired labour (women) available was 7.36. The results indicate that, 87.50 per cent of the households opined that the hired labour was adequate.

The results indicate that, households of the Kesalapura-1 micro-watershed possess 24.74 ha (45.42%) of dry land and 29.73 ha (54.58%) of irrigated land. Marginal farmers possess 3.04 ha (70.92%) of dry land and 1.25 ha (29.08%). Small farmers possess 4.3 ha (57.97%) of dry land and 3.12 ha (42.03%) of irrigated land. Semi medium farmers possess 2.83 ha (17.02%) of dry land and 13.81 per cent (82.98%) of irrigated land. Medium farmers possess 1.62 ha (12.29%) dry land and 11.56 (87.71%) of irrigated land. Large farmers possess 12.95 ha (100%) of dry land.

The results indicate that, the average value of dry land was Rs. 187078 and average value of irrigated land was Rs. 416878. In case of marginal famers, the average land value was Rs. 526232 for dry land and Rs. 1443506 for irrigated land. In case of small famers, the average land value was Rs. 146525 for dry land and was Rs. 737792 for irrigated land. In case of semi medium famers, the average land value was Rs. 211714 for dry land and Rs. 419748 for irrigated land. In case of medium famers, the average land value was Rs. 370500 for dry land and Rs. 216211 for irrigated land. In case of large farmers, the average land value for dry land was Rs. 92625.

The results indicate that, there were 20 functioning bore wells in the micro watershed. The results indicate that, bore well was the major irrigation source in the

micro water shed for 62.50 per cent of the farmers. The results indicate that, the depth of bore well was found to be 44.10 meters.

The results indicate that, marginal, small, semi medium and medium farmers had irrigated area of 1.21 ha, 7.65 ha, 12.01 ha and 11.48 ha respectively. The results indicate that, farmers have grown sugarcane (15.91 ha), groundnut (4.45 ha), maize (7.03 ha), sunflower (2.86 ha), pearl millet (5.19 ha), bengalgram (2.05 ha), chilly (1.21 ha), cotton (0.4 ha), mulberry (0.03 ha) and jowar (0.4 ha).

Marginal farmers have grown sugarcane, maize, sunflower, pearl millet, bengalgram, jowar and mulberry. Small farmers have grown sugarcane, maize, groundnut, sunflower, pearl millet, groundnut and cotton. Semi medium farmers have grown sugarcane, maize, bajra, pearl millet, bengalgram and chilly. Medium farmers have grown sugarcane, maize, groundnut, sunflower and chilly. Large farmers have grown bajra only.

The results indicate that, the cropping intensity in Kesalapura-1 micro-watershed was found to be 72.08 per cent. In case of marginal farmers it was 92.48 per cent, small farmers it was 94.18 per cent, in case of semi medium farmers it was 74.11, medium farmers it was 62.52 per cent and in case of large farmers it was 50 per cent.

The results indicate that, 46.88 per cent of the households have bank account. The results indicate that, 46.88 per cent of the households have availed credit from different sources.

The results indicate that, the total cost of cultivation for sugarcane was Rs. 60569.99. The gross income realized by the farmers was Rs. 525781.85. The net income from sugarcane cultivation was Rs. 465211.86. Thus the benefit cost ratio was found to be 1:8.68. The total cost of cultivation for groundnut was Rs. 42208.67. The gross income realized by the farmers was Rs. 102827.47. The net income from groundnut cultivation was Rs. 60618.81. Thus the benefit cost ratio was found to be 1:2.44. The total cost of cultivation for maize was Rs. 43481.62. The gross income realized by the farmers was Rs. 54611.46. The net income from maize cultivation was Rs. 11129.85. Thus the benefit cost ratio was found to be 1:1.26. The total cost of cultivation for bajra was Rs. 19721.25. The gross income realized by the farmers was Rs. 22622.75. The net income from bajra cultivation was Rs. 2901.51. Thus the benefit cost ratio was found to be 1:1.15. The total cost of cultivation for Jowar was Rs. 61598.57. The gross income realized by the farmers was Rs. 55328. The net income from Jowar cultivation was Rs. -6270.57. Thus the benefit cost ratio was found to be 1:0.9. The total cost of cultivation for chilly was Rs. 223901.85. The gross income realized by the farmers was Rs. 322643.75. The net income from chilly cultivation was Rs. 98741.90. Thus the benefit cost ratio was found to be 1:1.44. The total cost of cultivation for Bengalgram was Rs. 52147.87. The gross income realized by the farmers was Rs. 97556.48. The net income from Bengalgram cultivation

was Rs. 45408.62. Thus the benefit cost ratio was found to be 1:1.87. The total cost of cultivation for Sunflower was Rs. 32268.06. The gross income realized by the farmers was Rs. 64321.25. The net income from Sunflower cultivation was Rs. 32053.20. Thus the benefit cost ratio was found to be 1:1.99. The total cost of cultivation for Mulberry was Rs. 849677.77. The gross income realized by the farmers was Rs. 2161250.05. The net income from Mulberry cultivation was Rs. 1311572.28. Thus the benefit cost ratio was found to be 1:2.54. The total cost of cultivation for cotton was Rs. 48650.76. The gross income realized by the farmers was Rs. 118560. The net income from cotton cultivation was Rs. 69909.24. Thus the benefit cost ratio was found to be 1:2.44.

The results indicate that, 65.63 per cent of the households opined that dry fodder was adequate and 40.63 per cent opined that green fodder was adequate.

The results indicate that the average annual gross income was Rs. 37667 for landless farmers, for marginal farmers it was Rs. 181857, for small farmers it was Rs. 118516, for semi medium farmers it was Rs. 225550, for medium farmers it was Rs. 355000 and for large farmers it was Rs.120000.

The results indicate that the average annual expenditure is Rs. 33462. For landless households it was Rs. 6889, for marginal farmers it was Rs. 68184, for small farmers it was Rs. 16335, for semi medium farmers it was Rs. 16350, for medium farmers it was Rs. 58750 and for large farmers it was Rs.60000.

The results indicate that, households have planted 69 coconut trees, 2 lemon trees, 2 lime trees and 7 mango trees in the field and 8 coconut trees in the backyard. The results indicate that, households have planted 75 neem tree, 1 banyan tree and 29 teak trees in their fields.

The results indicated that, cotton, maize, mulberry, sugarcane and sunflower were sold to the extent of 100 per cent. Bajra was sold to the extent of 86.67 per cent, Bengal gram to the extent of 54.55 per cent, chilly to the extent of 94.29 per cent, groundnut to the extent of 52.38 per cent and jowar was sold to the extent of 68.75 per cent.

The results indicated that, about 18.75 per cent of the famers have sold their produce in regulated markets, 53.13 per cent of the farmers have sold to local/village merchants and another 53.13 per cent of the farmers have sold to agents/traders.

The results indicated that, 87.50 per cent of the households have used tractor, 25 per cent of the households have used cart, 9.38 per cent of the households have used truck and 3.13 per cent have carried head loads as a mode of transportation for their agricultural produce.

The results indicated that, 46.88 per cent of the households have experienced soil and water erosion problems in the farm i.e., 71.43 per cent of the marginal farmers, 28.5 per cent of the small farmers, 40 per cent of semi medium farmers and 75 per cent of

medium farmers and 100 per cent of large farmers have experienced soil and water erosion problems. The results indicated that, 50 per cent have shown interest in soil test.

The results indicated that, 53.13 per cent of the households used firewood and 46.88 per cent have used LPG as a source of fuel. The results indicated that, bore well was the major source of drinking water for 34.38 per cent of the households and piped supply was the source of drinking water for 59.38 per cent of the households in the micro watershed.

Electricity was the major source of light for 100 per cent of the households in micro watershed. The results indicated that, 43.75 per cent of the households possess sanitary toilet. The results indicated that, 90.63 per cent of the sampled households possessed BPL card and 9.38 per cent of the households did not possess any PDS cards. The results indicated that, 46.88 per cent of the households participated in NREGA programme.

The results indicated that, cereals were adequate for 100 per cent of the households, pulses were adequate for 62.50 per cent, oilseeds were adequate for 68.75 per cent, vegetables were adequate for 62.50 per cent, fruits were adequate for 6.25 per cent, milk was adequate for 90.63 per cent, eggs were adequate for 56.25 per cent and meat were adequate for 46.88 per cent.

The results indicated that, pulses were inadequate for 37.50 per cent of the households, oilseeds were inadequate for 31.25 per cent, vegetables were inadequate for 37.50 per cent, fruits were inadequate for 53.13 per cent, eggs were inadequate for 28.13 per cent, milk was inadequate for 9.38 per cent of the households and meat was inadequate for 37.50 per cent of the households.

The results indicated that, lower fertility status of the soil was the constraint experienced by 50 per cent of the households, wild animal menace on farm field (59.38%), frequent incidence of pest and diseases (71.88%), inadequacy of irrigation water (31.25%), high cost of fertilizers and plant protection chemicals (84.38%), high rate of interest on credit (71.88%), low price for the agricultural commodities (43.75%), lack of marketing facilities in the area (65.63%), lack of transport for safe transport of the agricultural produce to the market (43.75%), inadequate extension services (9.38%), less rainfall (53.13%) and source of agri-technology information (15.63%).