



हर कदम, हर डगर
किसानों का हमसफर
भारतीय कृषि अनुसंधान परिषद

AgriSearch with a human touch

**LAND RESOURCE INVENTORY AND SOCIO-ECONOMIC STATUS OF
FARM HOUSEHOLDS FOR WATERSHED PLANNING AND
DEVELOPMENT**

SUDAPUR-2 (4D5B4J2a) MICROWATERSHED

Gurumitkal Hobli, Yadgir Taluk and District, Karnataka

Karnataka Watershed Development Project – II

SUJALA – III

World Bank funded Project



ICAR – NATIONAL BUREAU OF SOIL SURVEY AND LAND USE PLANNING



ICAR - NBSS & LUP



**WATERSHED DEVELOPMENT DEPARTMENT
GOVT. OF KARNATAKA, BANGALORE**



About ICAR - NBSS&LUP

The ICAR-National Bureau of Soil Survey and Land Use Planning (ICAR-NBSS&LUP), Nagpur, a premier Institute of the Indian Council of Agricultural Research (ICAR), was set up during 1976 with the objective to prepare soil resource maps at national, state and district levels and to provide research inputs in soil resource mapping and its applications, land evaluation, land use planning, land resource management, and database management using GIS for optimising land use on different kinds of soils in the country.

The Bureau has been engaged in carrying out soil resource survey, agro-ecological and soil degradation mapping at the country, state and district levels for qualitative assessment and monitoring the soil health towards viable land use planning. The research activities have resulted in identifying the soil potentials and problems, and the various applications of the soil surveys with the ultimate objective of sustainable agricultural development. The Bureau has the mandate to correlate and classify soils of the country and maintain a National Register of all the established soil series. The Institute is also imparting in-service training to staff of the soil survey agencies in the area of soil survey, land evaluation and soil survey interpretations for land use planning. The Bureau in collaboration with Panjabrao Krishi Vidyapeeth, Akola is running post-graduate teaching and research programme in land resource management, leading to M.Sc. and Ph.D. degrees.

Citation: Rajendra Hegde, Ramesh Kumar, S.C., B.A. Dhanorkar, S. Srinivas, M.Lalitha, K.V. Niranjana, R.S. Reddy and S.K. Singh (2019). "Land Resource Inventory and Socio-Economic Status of Farm Households for Watershed Planning and Development of Sudapur 2 (4D5B4J2a) Microwatershed, Gurmitkal Hobli, Yadgir Taluk and District, Karnataka", ICAR-NBSS&LUP Sujala MWS Publ.350, ICAR – NBSS & LUP, RC, Bangalore. p.129 & 31.

TO OBTAIN COPIES,

Please write to:

Director, ICAR - NBSS & LUP,

Amaravati Road, NAGPUR - 440 033, India

Phone : (0712) 2500386, 2500664, 2500545 (O)

Telefax : 0712-2522534

E-Mail : director@nbsslup.ernet.in

Website URL : nbsslup.in

Or

Head, Regional Centre, ICAR - NBSS&LUP, Hebbal, Bangalore - 560 024

Phone : (080) 23412242, 23510350 (O)

Telefax : 080-23510350

E-Mail : nbssrcb@gmail.com

ICAR-NBSS&LUP Sujala MWS Publ.350



हर कदम, हर डगर
किसानों का हमसफर
भारतीय कृषि अनुसंधान परिषद

*Agri*search with a human touch

LAND RESOURCE INVENTORY AND SOCIO-ECONOMIC STATUS OF FARM HOUSEHOLDS FOR WATERSHED PLANNING AND DEVELOPMENT

SUDAPUR-2 (4D5B4J2a) MICROWATERSHED

Gurumitkal Hobli, Yadgir Taluk and District, Karnataka


Karnataka Watershed Development Project – II

Sujala-III

World Bank funded Project



ICAR – NATIONAL BUREAU OF SOIL SURVEY AND LAND USE
PLANNING



WATERSHED DEVELOPMENT DEPARTMENT, GOVT. OF
KARNATAKA, BANGALORE



PREFACE

In Karnataka, as in other Indian States, the livelihoods of rural people are intertwined with farming pursuits. The challenges in agriculture are seriously threatening the livelihood of a large number of farmers as they have been practicing farming in contextual factors beyond their control. Climatic factors are the most important ones and have become much more significant in recent times due to rapid climate changes induced by intensive anthropogenic activities affecting our ecosystem in multiple ways. Climate change has become the reality, it is happening and efforts to evolve and demonstrate climate resilient technologies have become essential. Due to the already over stressed scenario of agrarian sector, the climate change is resulting in manifold increase in the complexities, pushing the rural mass to face more and more unpredictable situations. The rising temperatures and unpredictable rainfall patterns are going to test seriously the informed decisions farmers have to make in order to survive in farming and sustain their livelihood.

It is generally recognized that impacts of climate change shall not be uniform across the globe. It is said that impact of climate change is more severe in South Asia. Based on the analysis of meteorological data, it is predicted that in India, there will be upward trend in mean temperature, downward trend in relative humidity, annual rainfall and number of wet days in a year. Also, in general, phenomena like erratic monsoon, spread of tropical diseases, rise in sea levels, changes in availability of fresh water, frequent floods, droughts, heat waves, storms and hurricanes are predicted. Each one of these adverse situations are already being experienced in various parts of India and also at the global level. Decline in agricultural productivity of small and marginal farmers becoming more vulnerable is already witnessed.

In Karnataka, more than 60 per cent of the population live in rural areas and depend on agriculture and allied activities for their livelihood. Though the state has achieved significant progress in increasing the yield of many crops, there is tremendous pressure on the land resources due to the growing and competing demands of various land uses. This is reflected in the alarming rate of land degradation observed. Already more than 50 per cent of the area is affected by various forms of degradation. If this trend continues, the sustainability of the fragile ecosystem will be badly affected. The adverse effects of change in the climatic factors are putting additional stress on the land resources and the farmers dependent on this.

The natural resources (land, water and vegetation) of the state need adequate and constant care and management, backed by site-specific technological interventions and investments particularly by the government. Detailed database pertaining to the nature of the land resources, their constraints, inherent potentials and suitability for various land

based rural enterprises, crops and other uses is a prerequisite for preparing location-specific action plans, which are in tune with the inherent capability of the resources. Any effort to evolve climate resilient technologies has to be based on the baseline scientific database. Then only one can expect effective implementation of climate resilient technologies, monitor the progress, make essential review of the strategy, and finally evaluate the effectiveness of the implemented programs. The information available at present on the land resources of the state are of general nature and useful only for general purpose planning. Since the need of the hour is to have site-specific information suitable for farm level planning and detailed characterization and delineation of the existing land resources of an area into similar management units is the only option.

ICAR-NBSS&LUP, Regional Centre, Bangalore has taken up a project sponsored by the Karnataka Watershed Development Project-II, (Sujala-III), Government of Karnataka funded by the World Bank under Component-1 Land Resource Inventory. This study was taken up to demonstrate the utility of such a database in reviewing, monitoring and evaluating all the land based watershed development programs on a scientific footing. To meet the requirements of various land use planners at grassroots level, the present study on “Land Resource Inventory and Socio-Economic Status of Farm Households for Watershed Planning and Development of Sudapur-2 Microwatershed, Yadgir Taluk and District, Karnataka” for integrated development was taken up in collaboration with the State Agricultural Universities, IISC, KRSRAC, KSNDMC as Consortia partners. The project provides detailed land resource information at cadastral level (1:7920 scale) for all the plots and socio-economic status of farm households covering thirty per cent farmers randomly selected representing landed and landless class of farmers in the microwatershed. The project report with the accompanying maps for the Microwatershed will provide required detailed database for evolving effective land use plan, alternative land use options and conservation plans for the planners, administrators, agricultural extension personnel, KVK officials, developmental departments and other land users to manage the land resources in a sustainable manner.

It is hoped that this database will be useful to the planners, administrators and developmental agencies working in the area in not only for formulating location specific developmental schemes but also for their effective monitoring at the village/watershed level.

Nagpur

Date: 06-09-2019

S.K. SINGH

Director, ICAR - NBSS&LUP, Nagpur

Contributors

Dr. Rajendra Hegde Principal Scientist, Head & Project Leader, Sujala-III Project ICAR-NBSS&LUP, Regional Centre, Bangalore	Dr. S.K.Singh Director, ICAR-NBSS&LUP Coordinator, Sujala-III Project Nagpur
Soil Survey, Mapping & Report Preparation	
Dr. B.A. Dhanorkar	Sh. R.S. Reddy
Dr. K.V. Niranjana	Sh. Venkata Giriappa
	Mr. Somashekar T N
	Smt. Chaitra, S.P.
	Dr. Gopali bardhan
	Ms. Arpitha
	Dr. Mahendra Kumar, M.B.
Field Work	
Sh. C.BacheGowda	Sh. Mahesh, D.B.
Sh. Somashekar	Sh. Ashok S Sindagi
Sh. M. Jayaramaiah	Sh. Veerabhadrapa B.
Sh. Paramesha, K.	Sh. Shankarappa
Sh. B. M. Narayana Reddy	Sh. Anand
	Sh. Arun N Kambar.
	Sh Kamalesh Awate
	Sh. Sharaan Kumar Huppar
	Sh. Yogesh H.N.
	Sh. Kalaveerachari R Kammar
GIS Work	
Dr. S.Srinivas	Sh. A.G.Devendra Prasad
Sh. D.H.Venkatesh	Sh. Prakashanaik, M.K.
Smt.K.Sujatha	Sh. Abhijith Sastry, N.S.
Smt. K.V.Archana	Sh. Sudip Kumar Suklabaidya
Sh. N. Maddileti	Sh. Avinash, K.N.
	Sh. Amar Suputhra, S
	Sh. Deepak, M.J.
	Smt. K.Karunya Lakshmi
	Ms. Seema, K.V.
	Ms. A. Rajab Nisha

Laboratory Analysis	
Dr. K.M.Nair	Ms. Steffi Peter
Smt. Arti Koyal	Ms. Thara, V.R
Smt. Parvathy	Ms. Roopa, G.
	Ms. Swati, H.
	Sh. Shantaveera Swami
	Ms. Shwetha, N.K.
	Smt. Ishrat Haji
	Ms. P. Pavan Kumari
	Ms. Padmaja
	Ms. Veena, M.
Socio-Economic Analysis	
Dr. S.C. Ramesh Kumar	Sh. M.K. Prakashanaik
	Ms. Karuna V. Kulkarni
	Mrs. Sowmya A.N
	Sh. Vinod R
	Sh. Basavaraja
	Sh. Vijay Kumar Lamani
	Ms. Sowmya K.B
	Mrs. Prathibha, D.G
	Sh. Rajendra,D
Soil & Water Conservation	
Sh. Sunil P. Maske	
Watershed Development Department, GoK, Bangalore	
Sh. Rajeev Ranjan IFS Project Director & Commissioner, WDD	Dr. A. Natarajan NRM Consultant, Sujala-III Project
Dr. S.D. Pathak IFS Executive Director & Chief Conservator of Forests, WDD	

PART-A

LAND RESOURCE INVENTORY

Contents

Preface		
Contributors		
Executive Summary		
Chapter 1	Introduction	1
Chapter 2	Geographical Setting	3
2.1	Location and Extent	3
2.2	Geology	3
2.3	Physiography	4
2.4	Drainage	4
2.5	Climate	4
2.6	Natural Vegetation	6
2.7	Land Utilization	6
Chapter 3	Survey Methodology	11
3.1	Base maps	11
3.2	Image Interpretation for Physiography	11
3.3	Field Investigation	14
3.4	Soil Mapping	16
3.5	Land Management Units	16
3.5	Laboratory Characterization	17
Chapter 4	The Soils	21
4.1	Soils of granite gneiss landscape	21
Chapter 5	Interpretation for Land Resource Management	33
5.1	Land Capability Classification	33
5.2	Soil Depth	35
5.3	Surface Soil Texture	36
5.4	Soil Gravelliness	37
5.5	Available Water Capacity	38
5.6	Soil Slope	39
5.7	Soil Erosion	40
Chapter 6	Fertility Status	43
6.1	Soil Reaction (pH)	43
6.2	Electrical Conductivity (EC)	43
6.3	Organic Carbon (OC)	43
6.4	Available Phosphorus	45
6.5	Available Potassium	45
6.6	Available Sulphur	45
6.7	Available Boron	46
6.8	Available Iron	46
6.9	Available Manganese	46
6.10	Available Copper	46
6.11	Available Zinc	50

Chapter 7	Land Suitability for Major Crops	51
7.1	Land suitability for Sorghum	51
7.2	Land suitability for Maize	52
7.3	Land suitability for Bajra	53
7.4	Land suitability for Groundnut	54
7.5	Land suitability for Sunflower	55
7.6	Land suitability for Redgram	56
7.7	Land suitability for Bengal gram	57
7.8	Land suitability for Cotton	58
7.9	Land suitability for Chilli	59
7.10	Land suitability for Tomato	60
7.11	Land suitability for Brinjal	61
7.12	Land suitability for Onion	62
7.13	Land suitability for Bhendi	63
7.14	Land suitability for Drumstick	64
7.15	Land suitability for Mango	65
7.16	Land suitability for Guava	66
7.17	Land suitability for Sapota	67
7.18	Land Suitability for Pomegranate	68
7.19	Land Suitability for Musambi	69
7.20	Land Suitability for Lime	70
7.21	Land Suitability for Amla	71
7.22	Land Suitability for Cashew	72
7.23	Land Suitability for Jackfruit	73
7.24	Land Suitability for Jamun	74
7.25	Land Suitability for Custard apple	75
7.26	Land Suitability for Tamarind	76
7.27	Land Suitability for Mulberry	77
7.28	Land Suitability for Marigold	78
7.29	Land Suitability for Chrysanthemum	79
7.30	Land management units	80
7.31	Proposed Crop Plan	112
Chapter 8	Soil Health Management	115
Chapter 9	Soil and Water conservation Treatment Plan	121
9.1	Treatment Plan	122
9.2	Recommended Soil and Water Conservation measures	125
9.3	Greening of Microwatershed	126
	References	129
	Appendix I	I-VIII
	Appendix II	IX-XVI
	Appendix III	XVII-XXIV

LIST OF TABLES

2.1	Mean Monthly Rainfall, PET, 1/2 PET at Yadgir Taluk & District	5
2.2	Land Utilization in Yadgir district	7
3.1	Differentiating Characteristics used for Identifying Soil Series	16
3.2	Soil map unit description of Sudapur-2 Microwatershed	17
4.1	Physical and Chemical Characteristics of Soil Series identified in Sudapur-2 microwatershed	26
7.1	Soil-Site Characteristics of Sudapur-2 Microwatershed	81
7.2	Land suitability criteria for Sorghum	82
7.3	Land suitability criteria for Maize	83
7.4	Land suitability criteria for Bajra	84
7.5	Land suitability criteria for Groundnut	85
7.6	Land suitability criteria for Sunflower	86
7.7	Land suitability criteria for Redgram	87
7.8	Land suitability criteria for Bengal gram	88
7.9	Land suitability criteria for Cotton	89
7.10	Land suitability criteria for Chilli	90
7.11	Land suitability criteria for Tomato	91
7.12	Land suitability criteria for Brinjal	92
7.13	Land suitability criteria for Onion	93
7.14	Land suitability criteria for Bhendi	94
7.15	Land suitability criteria for Drumstick	95
7.16	Land suitability criteria for Mango	96
7.17	Land suitability criteria for Guava	97
7.18	Land suitability criteria for Sapota	98
7.19	Land suitability criteria for Pomegranate	99
7.20	Land suitability criteria for Musambi	100
7.21	Land suitability criteria for Lime	101
7.22	Land suitability criteria for Amla	102
7.23	Land suitability criteria for Cashew	103
7.24	Land suitability criteria for Jackfruit	104

7.25	Land suitability criteria for Jamun	105
7.26	Land suitability criteria for Custard apple	106
7.27	Land suitability criteria for Tamarind	107
7.28	Land suitability criteria for Mulberry	108
7.29	Land suitability criteria for Marigold	109
7.30	Land suitability criteria for Chrysanthemum	110
7.31	Proposed Crop Plan for Sudapur-2 Microwatershed	113

LIST OF FIGURES

2.1	Location map of Sudapur-2 Microwatershed	3
2.2 a	Granite and granite gneiss rock formation	4
2.3	Rainfall distribution in Yadgir Taluk & District	5
2.4	Natural vegetation of Sudapur-2 Microwatershed	6
2.5	Current Land use map of Sudapur-2 Microwatershed	7
2.6 a & b	Different crops and cropping systems in Sudapur-2 Microwatershed	8
2.7	Location of Wells in Sudapur-2 microwatershed	9
3.1	Scanned and Digitized Cadastral map of Sudapur-2 Microwatershed	13
3.2	Satellite image of Sudapur-2 Microwatershed	13
3.3	Cadastral map overlaid on IRS PAN+LISS IV merged imagery of Sudapur-2 Microwatershed	14
3.4	Location of profiles in a transect	15
3.5	Soil phase or management units of Sudapur-2 Microwatershed	19
5.1	Land Capability Classification map of Sudapur-2 Microwatershed	35
5.2	Soil Depth map of Sudapur-2 Microwatershed	36
5.3	Surface Soil Texture map of Sudapur-2 Microwatershed	37
5.4	Soil Gravelliness map of Sudapur-2 Microwatershed	38
5.5	Soil Available Water Capacity map of Sudapur-2 Microwatershed	39
5.6	Soil Slope map of Sudapur-2 Microwatershed	40
5.7	Soil Erosion map of Sudapur-2 Microwatershed	41
6.1	Soil Reaction (pH) map of Sudapur-2 Microwatershed	44
6.2	Electrical Conductivity (EC) map of Sudapur-2 Microwatershed	44
6.3	Soil Organic Carbon (OC) map of Sudapur-2 Microwatershed	45
6.4	Soil Available Phosphorus map of Sudapur-2 Microwatershed	46
6.5	Soil Available Potassium map of Sudapur-2 Microwatershed	47
6.6	Soil Available Sulphur map of Sudapur-2 Microwatershed	47
6.7	Soil Available Boron map of Sudapur-2 Microwatershed	48
6.8	Soil Available Iron map of Sudapur-2 Microwatershed	48
6.9	Soil Available Manganese map of Sudapur-2 Microwatershed	49
6.10	Soil Available Copper map of Sudapur-2 Microwatershed	49
6.11	Soil Available Zinc map of Sudapur-2 Microwatershed	50
7.1	Land suitability for Sorghum	52

7.2	Land suitability for Maize	53
7.3	Land suitability for Bajra	54
7.4	Land suitability for Groundnut	55
7.5	Land suitability for Sunflower	56
7.6	Land suitability for Redgram	57
7.7	Land suitability for Bengal gram	58
7.8	Land suitability for Cotton	59
7.9	Land suitability for Chilli	60
7.10	Land suitability for Tomato	61
7.11	Land suitability for Brinjal	62
7.12	Land suitability for Onion	63
7.13	Land suitability for Bhendi	64
7.14	Land suitable for Drumstick	65
7.15	Land suitability for Mango	66
7.16	Land suitability for Guava	67
7.17	Land suitability for Sapota	68
7.18	Land suitability for Pomegranate	69
7.19	Land suitability for Musambi	70
7.20	Land suitability for Lime	71
7.21	Land suitability for Amla	72
7.22	Land suitability for Cashew	73
7.23	Land suitability for Jackfruit	74
7.24	Land suitability for Jamun	75
7.25	Land suitability for Custard apple	76
7.26	Land suitability for Tamarind	77
7.27	Land suitability for Mulberry	78
7.28	Land suitability for Marigold	79
7.29	Land suitability for Chrysanthemum	80
7.30	Land management map of Sudapur-2 Microwatershed	112
9.1	Soil and water conservation map of Sudapur-2 Microwatershed	126

EXECUTIVE SUMMARY

The land resource inventory of Sudapur-2 Microwatershed was conducted using village cadastral maps and IRS satellite imagery on 1:7920 scale. The false colour composites of IRS imagery were interpreted for physiography and the physiographic delineations were used as base for mapping soils. The soils were studied in several transects and a soil map was prepared with phases of soil series as mapping units. Random checks were made all over the area outside the transects to confirm and validate the soil map unit boundaries. The soil map shows the geographic distribution and extent, characteristics, classification, behavior and use potentials of the soils in the microwatershed.

The present study covers an area of 589 ha in Yadgir taluk & district, Karnataka. The climate is semiarid and categorized as drought-prone with an average annual rainfall of 866 mm, of which about 652 mm is received during south-west monsoon, 138 mm during north-east and the remaining 76 mm during the rest of the year. An area of 480 ha in the microwatershed is covered by soils, 108 ha by rock outcrops and 1 ha by others (habitation and water body). The salient findings from the land resource inventory are summarized briefly below.

- ❖ The soils belong to 6 soil series and 8 soil phases (management units) and 4 land management units.*
- ❖ The length of crop growing period is about 120-150 days starting from 1st week of June to 4th week of October.*
- ❖ From the master soil map, several interpretative and thematic maps like land capability, soil depth, surface soil texture, soil gravelliness, available water capacity, soil slope and soil erosion were generated.*
- ❖ Soil fertility status maps for macro and micronutrients were generated based on the surface soil samples collected at every 320 m grid interval.*
- ❖ Land suitability for growing 29 major agricultural and horticultural crops was assessed and maps showing the degree of suitability along with constraints were generated.*
- ❖ About 480 ha area in the microwatershed is suitable for agriculture.*
- ❖ About 13 per cent area is shallow (25-50 cm), 2 per cent area is moderately shallow (50-75 cm), 19 per cent area is moderately deep (75-100 cm), 10 per cent area of the microwatershed has soils that are deep (100-150 cm) and 39 per cent area of the microwatershed has soils that are very deep (>150 cm).*
- ❖ About 13 per cent loamy soils and 68 per cent clayey soils at the surface.*
- ❖ About 48 per cent area in the microwatershed is gravelly (15-35%) and 34 per cent is non gravelly (<15%).*

- ❖ *About 13 per cent is very low (<50 mm/m) in available water capacity, 20 per cent is low (51-100 mm/m) and 48 per cent is very high (>200 mm/m).*
- ❖ *About 62 per cent area in the microwatershed has very gently sloping (1-3% slope) and 19 per cent has nearly level (0-1%) lands.*
- ❖ *An area of about 19 per cent is slightly eroded (e1) and 62 per cent is moderately (e2) eroded.*
- ❖ *An area of about 44 per cent is slightly acid (pH 6.0-6.5) in soil reaction and 37 per cent area is neutral (pH 6.5-7.3).*
- ❖ *The Electrical Conductivity (EC) of the entire soils of the microwatershed is dominantly <2 dsm⁻¹ indicating that the soils are non-saline.*
- ❖ *Available organic carbon content is high (>0.75%) in the entire microwatershed area.*
- ❖ *An area of about 18 per cent is medium (23-57 kg/ha) in available phosphorus, 52 per cent area is low (<23 kg/ha) and 11 per cent is high (>57 kg/ha) in the microwatershed.*
- ❖ *Available potassium content is medium (145-337 kg/ha) in the entire area of the microwatershed.*
- ❖ *Available sulphur is medium (10-20 ppm) in an area of 22 per cent and low in an area of 59 per cent of the microwatershed.*
- ❖ *Available boron is low (<0.5 ppm) in an area of about 80 per cent and medium (0.5-1.0 ppm) in 2 per cent area of the microwatershed.*
- ❖ *Available iron is sufficient (>4.5 ppm) in the entire cultivated area of the microwatershed.*
- ❖ *Available manganese and copper are sufficient in all the cultivated soils of the microwatershed.*
- ❖ *Available zinc is deficient (<0.6 ppm) in an area of 44 per cent and sufficient (>0.6 ppm) in 38 per cent of area of the microwatershed.*
- ❖ *The land suitability for 29 major crops grown in the microwatershed were assessed and the areas that are highly suitable (S1) and moderately suitable (S2) are given below. It is however to be noted that a given soil may be suitable for various crops but what specific crop to be grown may be decided by the farmer looking to his capacity to invest on various inputs, marketing infrastructure, market price and finally the demand and supply position.*

Land suitability for various crops in the Microwatershed

Crop	Suitability Area in ha (%)		Crop	Suitability Area in ha (%)	
	Highly suitable (S1)	Moderately suitable (S2)		Highly suitable (S1)	Moderately suitable (S2)
<i>Sorghum</i>	-	404 (69)	<i>Guava</i>	-	110 (19)
<i>Maize</i>	33 (6)	371 (63)	<i>Sapota</i>	-	110 (19)
<i>Bajra</i>	110 (19)	294 (50)	<i>Pomegranate</i>	-	394 (67)
<i>Groundnut</i>	110 (19)	10(2)	<i>Musambi</i>	-	394 (67)
<i>Sunflower</i>	-	-	<i>Lime</i>	-	394 (67)
<i>Redgram</i>	-	394 (67)	<i>Amla</i>	110 (19)	10 (2)
<i>Bengal gram</i>	-	284 (48)	<i>Cashew</i>	-	110 (19)
<i>Cotton</i>	-	284 (48)	<i>Jackfruit</i>	-	110 (19)
<i>Chilli</i>	33 (6)	371 (63)	<i>Jamun</i>	-	-
<i>Tomato</i>	33 (6)	87 (15)	<i>Custard apple</i>	110 (19)	294 (50)
<i>Brinjal</i>	33 (6)	87 (15)	<i>Tamarind</i>	-	-
<i>Onion</i>	33 (6)	87 (15)	<i>Mulberry</i>	-	110 (19)
<i>Bhendi</i>	33 (6)	371 (63)	<i>Marigold</i>	33 (6)	371 (63)
<i>Drumstick</i>	-	110 (19)	<i>Chrysanthemum</i>	33 (6)	371 (63)
<i>Mango</i>	-	-			

- ❖ *Apart from the individual crop suitability, a proposed crop plan has been prepared for the identified LMUs by considering only the highly and moderately suitable lands for different crops and cropping systems with food, fodder, fibre and other horticulture crops.*
- ❖ *Maintaining soil-health is vital for crop production and conserve soil and land resource base for maintaining ecological balance and to mitigate climate change. For this, several ameliorative measures have been suggested for these problematic soils like saline/alkali, highly eroded, sandy soils etc.,*
- ❖ *Soil and water conservation treatment plan has been prepared that would help in identifying the sites to be treated and also the type of structures required.*
- ❖ *As part of the greening programme, several tree species have been suggested to be planted in marginal and submarginal lands, field bunds and also in the hillocks, mounds and ridges. This would help in not only supplementing the farm income but also provide fodder and fuel and generate lot of biomass which would help in maintaining an ecological balance and also contribute to mitigating the climate change.*

INTRODUCTION

Soil being a vital natural resource on whose proper use depends the life supporting systems of a country and the socioeconomic development of its people. Soils provide food, fodder, fibre and fuel for meeting the basic human and animal needs. With the ever increasing growth in human and animal population, the demand on soil for more food and fodder production is on the increase. The area available for agriculture is about 51 per cent of the total geographical area and more than 60 per cent of the people are still dependant on agriculture for their livelihood. However, the capacity of a soil to produce is limited and the limits to the production are set by its intrinsic characteristics, agro-climatic setting, and, use and management. There is, therefore, tremendous pressure on land and water resources, which is causing decline in soil-health and stagnation in productivity. The soils have been degrading at an estimated rate of one million hectares per year and ground water levels have been receding at an alarming rate resulting in decline in the ground water resource. Further, land degradation has emerged as a serious problem which has already affected about 38 lakh ha of cultivated area in the State. Soil erosion alone has degraded about 35 lakh ha. Almost all the uncultivated areas are facing various degrees of degradation, particularly soil erosion; salinity and alkalinity has emerged as a major problem in more than 3.5 lakh ha in the irrigated areas of the State. Nutrient depletion and declining factor productivity is common in both rainfed and irrigated areas. The degradation is continuing at an alarming rate and there appears to be no systematic effort among the stakeholders to contain this process. In recent times, an aberration of weather due to climate change phenomenon has added another dimension leading to unpredictable situation to be tackled by the farmers.

In this critical juncture, the challenge before us is not only to increase the productivity per unit area which is steadily declining and showing a fatigue syndrome, but also to prevent or at least reduce the severity of degradation. If the situation is not reversed at the earliest, then the sustainability of the already fragile crop production system and the overall ecosystem will be badly affected in the state. Added to this, every year there is a significant diversion of farm lands and water resources for non-agricultural purposes. Thus, developing strategies to slow down the degradation process or reclaim the soils to normal condition and ensure sustainability of production system are the major issues today. This demands a systematic appraisal of our soil and land resources with respect to their extent, geographic distribution, characteristics, behaviour and use potential, which is very important for developing an effective land use and cropping systems for augmenting agricultural production on a sustainable basis.

The soil and land resource inventories made so far in Karnataka had limited utility because the surveys were of different types, scales and intensities carried out at different times with specific objectives. Hence, there is an urgent need to generate detailed site-

specific farm level database on various land resources for all the villages/watersheds in a time bound manner that would help to protect the valuable soil and land resources and also to stabilize the farm production. Therefore, the land resource inventory required for farm level planning is the one which investigates all the parameters which are critical for productivity *viz.*, soils, site characteristics like slope, erosion, gravelliness and stoniness, climate, water, topography, geology, hydrology, vegetation, crops, land use pattern, animal population, socio-economic conditions, infrastructure, marketing facilities and various schemes and developmental works of the government etc. From the data collected at farm level, the specific problems and potentials of the area can be identified and highlighted, conservation measures required for the area can be planned on a scientific footing, suitability of the area for various uses can be worked out and finally viable and sustainable land use options suitable for each and every land holding can be prescribed.

The Land Resource Inventory is basically done for identifying potential and problem areas, developing sustainable land use plans, estimation of surface run off and water harvesting potential, preparation of soil and water conservation plans, land degradation/desertification etc. The Bureau is presently engaged in developing an LRI methodology using high resolution satellite remote sensing data and Digital Elevation Model (DEM) data to prepare Landscape Ecological Units (LEU) map representing agro-ecosystem as a whole. The LEU is preferred over landform as the base map for LRI. LEU is the assemblage of landform, slope and land use. An attempt has already been made to upscale the soil resource information from 1:250000 and 1:50000 scale to the LEU map in Goa and in some other states.

The land resource inventory aims to provide site-specific database for Sudapur-2 microwatershed in Yadgir Taluk and Yadgir District, Karnataka State for the Karnataka Watershed Development Department. The database was generated by using cadastral map of the village as a base along with high resolution IRS LISS IV and Cartosat-1 merged satellite imagery. Later, an attempt will be made to uplink this LRI data generated at 1:7920 scale under Sujala-III Project to the proposed Landscape Ecological Units (LEUs) map.

The study was organized and executed by the ICAR- National Bureau of Soil Survey and Land Use Planning, Regional Centre, Bangalore under Generation of Land Resource Inventory Data Base Component-1 of the Sujala-III Project funded by the World Bank.

GEOGRAPHICAL SETTING

2.1 Location and Extent

The Sudapur-2 microwatershed is located in the northern part of Karnataka in Yadgir Taluk & District, Karnataka State (Fig.2.1). It comprises parts of Siddhapur.B, Gajarakota and Mitathapadamapalli villages. It lies between $16^{\circ} 56'$ and $16^{\circ} 54'$ North latitudes and $77^{\circ} 18'$ and $77^{\circ} 20'$ East longitudes covering an area of about 589 ha. It is about 42 km southeast of Yadgir town and is surrounded by Siddhapur.B on the north and northeast, Gajarakota on the northwest, southwest, western, central, southern and Mitathapadamapalli village on the southeast and eastern side.




Fig.2.1 Location map of Sudapur-2 Microwatershed

2.2 Geology

Major rock formations observed in the microwatershed are granite gneiss (Figs.2.2). They are essentially pink to gray and are coarse to medium grained. They consist

primarily of quartz, feldspar, biotite and hornblende. The gray granite gneisses are highly weathered, fractured and fissured upto a depth of about 10 m. Dolerite dykes and quartz veins are common with variable width and found to occur in Sudapur-2 microwatershed.



Fig.2.2 Granite and granite gneiss rocks

2.3 Physiography

Physiographically, the area has been identified as granite gneiss landscape based on geology. The area has been further subdivided into five landforms, *viz*; mounds/ridges, summits, side slopes and very gently sloping uplands, plains and valleys based on slope and its relief features. The elevation ranges from 537-603 m above MSL. The mounds and ridges are mostly covered by rock outcrops.

2.4 Drainage

The area is drained by several parallel streams like Bori, Amerja and Kanga which finally join the river Bhima along its course. Though, they are not perennial, during rainy season they carry large quantities of rain water. The microwatershed has only few small tanks which are not capable of storing the water that flows during the rainy season. Due to this, the ground water recharge is very much affected. This is reflected in the failure of many bore wells in the villages. If the available rain water is properly harnessed by constructing new tanks and recharge structures at appropriate places in the villages, then the drinking and irrigation needs of the area can be easily met. The drainage network is parallel to sub parallel and dendritic.

2.5 Climate

The Yadgir district lies in the northern plains of Karnataka and falls under semiarid tract of the state and is categorized as drought- prone with total annual rainfall of 866 mm (Table 2.1). Of the total rainfall, maximum of 652 mm is received during the south–west monsoon period from June to September, the north-east monsoon from

October to early December contributes about 138 mm and the remaining 76 mm during the rest of the year. The summer season starts during the middle of February and continues up to the first week of June. The period from December to the middle of February is the coldest season. December is the coldest month with mean daily maximum and minimum temperatures being 29.5⁰C and 10⁰C respectively. During peak summer, temperature shoots up to 45⁰C. Relative humidity varies from 26% in summer to 62% in winter. Rainfall distribution is shown in Figure 2.3. The average Potential Evapo-Transpiration (PET) is 141 mm and varies from a low of 81 mm in December to 199 mm in the month of May. The PET is always higher than precipitation in all the months except July, August and September. Generally, the Length of crop Growing Period (LGP) is 120-150 days and starts from 1st week of June to 4th week of October.

Table 2.1 Mean Monthly Rainfall, PET, 1/2 PET at Yadgir Taluk, Yadgir District

Sl. No.	Months	Rainfall	PET	1/2 PET
1	January	4.30	86.0	43.0
2	February	2.30	125.5	62.7
3	March	15.10	166.0	83.0
4	April	18.50	179.8	89.9
5	May	36.0	198.8	97.9
6	June	118.0	175.1	87.5
7	July	171.80	156.3	78.1
8	August	182.9	150.3	75.1
9	September	179.7	142.0	71.0
10	October	105.3	138.5	69.2
11	November	26.4	97.60	48.6
12	December	6.0	80.90	40.4
Total		866.3		




Fig 2.3 Rainfall distribution in Yadgir Taluk, Yadgir District

2.6 Natural Vegetation

The natural vegetation is sparse comprising few tree species, shrubs and herbs. The mounds, ridges and boulders occupy very sizeable area which is under thin to moderately thick forest vegetation. Still, there are some remnants of the past forest cover which can be seen in patches in some ridges and hillocks in the microwatershed (Fig 2.4).

Apart from the continuing deforestation, the presence of large population of goats, sheep and other cattle in the microwatershed is causing vegetative degradation of whatever little vegetation left in the area. The uncontrolled grazing has left no time for the regeneration of the vegetative cover. This leads to the accelerated rate of erosion on the hill slopes resulting in the formation of deep gullies in the foot slopes that eventually result in the heavy siltation of tanks and reservoirs in the microwatershed.



Fig 2.4 Natural vegetation of Sudapur-2 microwatershed

2.7 Land Utilization

About 72 per cent area (Table 2.2) in Yadgir district is cultivated at present. An area of about 2 per cent is permanently under pasture, 20 per cent under current fallows and 6 per cent under non-agricultural land, and 5 per cent under currently barren. Forests occupy an area of about 7 per cent and the tree cover is in a very poor state. Most of the mounds, ridges and bouldery areas have very poor vegetative cover. Major crops grown in the area are sorghum, maize, cotton, sunflower, groundnut, red gram and paddy. The cropping intensity is 120 per cent in the taluk. While carrying out land resource inventory, the land use/land cover particulars are collected from all the survey numbers and a current land use map of the microwatershed is prepared. The current land use map prepared shows the arable and non-arable lands, other land uses and different types of crops grown in the area. The current land use map of Sudapur-2 microwatershed is presented in Fig.2.5.

The different crops and cropping systems adopted in the microwatershed is presented in the Figures 2.6. The location of wells in Sudapur-2 microwatershed is given in Fig.2.7.

Table 2.2 Land Utilization in Yadgir District

Sl. No.	Agricultural land use	Area (ha)	Per cent
1.	Total geographical area	516088	-
2.	Total cultivated area	373617	72.4
3.	Area sown more than once	74081	14.3
4.	Cropping intensity	-	119.8
5.	Trees and grooves	737	0.14
6.	Forest	33773	6.54
7.	Cultivable wasteland	2385	0.46
8.	Permanent Pasture land	11755	2.28
9.	Barren land	27954	5.41
10.	Non- Agriculture land	29623	5.73
11.	Current Fallows	105212	20.4




Fig.2.5 Current Land Use map of Sudapur-2 Microwatershed





Fig 2.6 Different Crops and Cropping Systems in Sudapur-2 Microwatershed




Fig 2.7 Location of wells in Sudapur-2 Microwatershed

SURVEY METHODOLOGY

The purpose of land resource inventory is to delineate similar areas (soil series and phases), which respond or expected to respond similarly for a given level of management. This was achieved in Sudapur-2 microwatershed by the detailed study of all the soil characteristics (depth, texture, colour, structure, consistence, coarse fragments, porosity, soil reaction, soil horizons etc.) and site characteristics (slope of the land, erosion, drainage, occurrence of rock fragments etc.) followed by grouping of similar areas based on soil-site characteristics into homogeneous (management units) units, and showing the area extent and their geographic distribution on the microwatershed cadastral map. The detailed survey at 1:7920 scale was carried out in an area of 589 ha. The methodology followed for carrying out land resource inventory was as per the guidelines given in Soil Survey Manual (IARI, 1971; Soil Survey Staff, 2006; Natarajan *et al.*, 2015) which is briefly described below.

3.1 Base Maps

The detailed survey of the land resources occurring in the microwatershed was carried out by using digitized cadastral map and satellite imagery as base supplied by KRSRSAC. The cadastral map shows field boundaries with their survey numbers, location of tanks, streams and other permanent features of the area (Fig. 3.1). Apart from the cadastral map, remote sensing data products from Cartosat-1 and LISS IV merged at the scale of 1:7920 were used in conjunction with the cadastral map to identify the landscapes, landforms and other surface features. The imagery helped in the identification and delineation of boundaries between hills, uplands and lowlands, water bodies, forest and vegetated areas, roads, habitations and other cultural features of the area (Fig. 3.2). The cadastral map was overlaid on the satellite imagery (Fig. 3.3) that helps to identify the parcel boundaries and other permanent features. Apart from cadastral maps and images, toposheets of the area (1:50,000 scale) were also used for initial traversing, identification of geology and landforms, drainage features, present land use and also for selection of transects in the microwatershed.

3.2 Image Interpretation for Physiography

False Colour Composites (FCCs) of Cartosat-I and LISS-IV merged satellite data covering microwatershed area was visually interpreted using image interpretation elements and all the available collateral data with local knowledge. The delineated physiographic boundaries were transferred on to a cadastral map overlaid on satellite imagery. Physiographically, the area has been identified as granite gneiss landscape. It was divided into five landforms, *viz*; ridges and mounds, gently and very gently sloping uplands and lowlands based on slope and image characteristics. They were further

subdivided into physiographic/image interpretation units based on image characteristics. The image interpretation legend for physiography is given below.

Image Interpretation Legend for Physiography

G- Granite Gneiss Landscape

G1	Hills/ Ridges/ Mounds
G11	Summits
G12	Side slopes
G121	Side slopes with dark grey tones
G2	Uplands
G21	Summits
G22	Gently sloping uplands
G221	Gently sloping uplands, yellowish green (eroded)
G222	Gently sloping uplands, yellowish white (severely eroded)
G23	Very gently sloping uplands
G231	Very gently sloping uplands, yellowish green
G232	Very gently sloping uplands, medium green and pink
G233	Very gently sloping uplands, pink and green (scrub land)
G234	Very gently sloping uplands, medium greenish grey
G235	Very gently sloping uplands, yellowish white (eroded)
G236	Very gently sloping uplands, dark green
G237	Very gently sloping uplands, medium pink (coconut garden)
G238	Very gently sloping uplands, pink and bluish white (eroded)
G24	Valleys/ lowlands
G241	Valleys, pink tones
G242	Valleys gray mixed with pink tones




Fig 3.1 Scanned and Digitized Cadastral map of Sudapur-2 Microwatershed




Fig.3.2 Satellite Image of Sudapur-2 Microwatershed




Fig.3.3 Cadastral map overlaid on IRS PAN+LISS IV merged imagery of Sudapur-2 Microwatershed

3.3 Field Investigation

The field boundaries and survey numbers given on the cadastral sheet were located on the ground by following permanent features like roads, cart tracks, *nallas*, streams, tanks etc., and wherever changes were noticed, they were incorporated on the microwatershed cadastral map. Preliminary traverse of the microwatershed was carried out with the help of cadastral map, imagery and toposheets. While traversing, landforms and physiographic units identified were checked and preliminary soil legend was prepared by studying soils at few selected places. Then, intensive traversing of each physiographic unit like hills, ridges, uplands and valleys was carried out. Based on the variability observed on the surface, transects (Fig. 3.4) were selected across the slope covering all the landform units in the microwatershed (Natarajan and Dipak Sarkar, 2010).




Fig: 3.4. Location of profiles in a transect

In the selected transect, soil profiles were located (Fig. 3.4) at closely spaced intervals to take care of any change in the land features like break in slope, erosion, gravel, stones etc. In the selected sites, soil profiles (vertical cut showing the soil layers from surface to the rock) were opened upto 200 cm or to the depth limited by rock or hard substratum and studied in detail for all their morphological and physical characteristics. The soil and site characteristics were recorded for all profile sites on a standard proforma as per the guidelines given in USDA Soil Survey Manual (Soil Survey Staff, 2012). Apart from the transect study, profiles were also studied at random, almost like in a grid pattern, outside the transect areas.

Based on the soil characteristics, the soils were grouped into different soil series. Soil series is the most homogeneous unit having similar horizons and properties and behaves similarly for a given level of management. Soil depth, texture, colour, kind of horizon and horizon sequence, calcareousness, amount and nature of gravel present, nature of substratum etc, were used as the major differentiating characteristics for identifying soil series occurring in the area. The differentiating characteristics used for identifying the soil series are given in Table 3.1. Based on the above characteristics, 6 soil series were identified in the Sudapur-2 microwatershed.

Table 3.1 Differentiating Characteristics used for identifying Soil Series
(Characteristics are of Series Control Section)

Soils of Granite gneiss Landscape							
Sl. no	Soil Series	Depth (cm)	Colour (moist)	Texture	Gravel (%)	Horizon sequence	Calcareousness
1	BDL (Badiyala)	25-50	7.5 YR 2.5/3,2.5/2,3/3 10YR 3/4,4/3	sl	-	Ap-Bw	e
2	VNK (Vanakanahalli)	25-50	2.5 YR 3/4	sc	-	Ap-Bt-Cr	-
3	JNK (Jinkera)	50-75	10YR5/3,3/2 7.5YR3/4	scl	-	Ap-Bw	e
4	BLC (Balichakra)	75-100	2.5YR5/3,2.5/4 5YR4/3,3/3	sc	-	Ap-Bt	es
5	NGP (Nagalapur)	100-150	10YR3/2,3/1,2/1	c	-	Ap-Bss	es
6	BMN (Bhimanahalli)	>150	10YR 3/1	c	-	Ap-Bss	es

3.4 Soil Mapping

The area under each soil series was further separated into soil phases and their boundaries delineated on the cadastral map based on the variations observed in the texture of the surface soil, slope, erosion, presence of gravel, stoniness etc. A soil phase is a subdivision of soil series based mostly on surface features that affect its use and management. The soil mapping units are shown on the map (Fig.3.5) in the form of symbols. During the survey many soil profile pits, few minipits and a few auger bores representing different landforms occurring in the microwatershed were studied. In addition to the profile study, spot observations in the form of minipits, road cuts, terrace cuts etc., were studied to validate the soil boundaries on the soil map. The soil map shows the geographic distribution of 8 mapping units representing 6 soil series occurring in the microwatershed. The soil map unit (soil legend) description is presented in Table 3.2. The soil phase map (management units) shows the distribution of 8 soil phases mapped in the microwatershed. Each mapping unit (soil phase) delineated on the map has similar soil and site characteristics. In other words, all the farms or survey numbers included in one soil phase will have similar management needs and have to be treated accordingly.

3.5 Land Management Units (LMU's)

The 8 soil phases identified and mapped in the microwatershed were grouped into 4 Land Management Units (LMU's) for the purpose of preparing a Proposed Crop Plan for sustained development of the microwatershed. The database (soil phases) generated under LRI was utilized for identifying Land Management Units (LMU's) based on the management needs. One or more than one soil site characteristic having influence on the management have been chosen for identification and delineation of LMUs. For Sudapur-2 microwatershed, five soil and site characteristics, namely soil depth, soil texture, slope,

erosion and gravel content have been considered for defining LMUs. The Land Management Units are expected to behave similarly for a given level of management.

3.6 Laboratory Characterization

Soil samples for each soil series were collected from representative master profiles for laboratory characterization by following the methods outlined in the Laboratory Manual (Sarma *et al*, 1987). Surface soil samples collected from farmer's fields (57 samples) for fertility status (major and micronutrients) at 320 m grid interval in the year 2017 were analyzed in the laboratory (Katyal and Rattan, 2003). By linking the soil fertility data to the survey numbers through GIS, soil fertility maps were generated by using Kriging method for the microwatershed.

Table 3.2 Soil map unit description of Sudapur-2 Microwatershed

*Soil map unit No.	Soil Series	Soil Phase	Mapping Unit Description	Area in ha (%)
Soils of Granite and Granite Gneiss Landscape				
	BDL		Badiyala soils are shallow (25-50 cm), well drained, have dark brown to very dark brown and dark yellowish brown, slightly calcareous sandy loam soils occurring on very gently to gently sloping uplands under cultivation	42 (7.14)
5		BDLiB2	Sandy clay surface, slope 1-3%, moderate erosion	42 (7.14)
	VNK		Vanakanahalli soils are shallow (25-50 cm), well drained, have dark reddish brown, sandy clay red soils occurring on very gently to moderately sloping uplands under cultivation	34 (5.72)
109		VNKmB2g1	Clay surface, slope 1-3%, moderate erosion, gravelly (15-35%)	34 (5.72)
	JNK		Jinkera soils are moderately shallow (50-75 cm), well drained, have dark brown to very dark grayish brown, slightly calcareous sandy clay loam soils occurring on very gently sloping uplands under cultivation	10 (1.75)
152		JNKmB2	Clay surface, slope 1-3%, moderate erosion	10 (1.75)
	BLC		Balichakra soils are moderately deep (75-100 cm), well drained, have reddish brown to dark reddish brown, slightly calcareous sandy clay loam red soils occurring on very gently sloping uplands under cultivation	110 (18.69)
155		BLCcB2g1	Sandy loam surface, slope 1-3%, moderate erosion, gravelly (15-35%)	77 (13.01)
38		BLCiB2	Sandy clay surface, slope 1-3%, moderate erosion	33 (5.68)
	NGP		Nagalapur soils are deep (100-150 cm), moderately well drained, have very dark gray to very dark grayish brown, black calcareous cracking clay soils occurring on very gently sloping uplands under cultivation	57 (9.61)
146		NGPmB2g1	Clay surface, slope 1-3%, moderate erosion, gravelly (15-35%)	57 (9.61)
	BMN		Bhimanahalli soils are very deep (>150 cm), moderately well drained, have very dark gray, calcareous cracking clay black soils occurring on very gently sloping uplands under	227 (38.5)

*Soil map unit No.	Soil Series	Soil Phase	Mapping Unit Description	Area in ha (%)
		cultivation		
159		BMNmA1	Clay surface, slope 0-1%, slight erosion	113 (19.2)
63		BMNmB2g1	Clay surface, slope 1-3%, moderate erosion, gravelly (15-35%)	114 (19.3)
999		Rock outcrops	Rock lands, both massive and bouldery with little or no soil	108 (18.34)
1000		Others	Water body	1 (0.25)




Fig 3.5 Soil Phase or Management Units - Sudapur-2 Microwatershed

THE SOILS

Detailed information pertaining to the nature, extent and distribution of different kinds of soils occurring in Sudapur-2 microwatershed is provided in this chapter. The microwatershed area has been identified as granite gneiss landscape based on geology. In all, 6 soil series are identified. Soil formation is the result of the combined effect of environmental and terrain factors that are reflected in soil morphology. In the granite gneiss landscape, it is by parent material, relief and climate.

A brief description of each of the 6 soil series identified followed by 8 soil phases (management units) mapped are furnished below. The physical and chemical characteristics of soil series identified in Sudapur-2 microwatershed are given in Table 4.1 along with soil classification. The soils in any one map unit differ from place to place in their depth, texture, slope, gravelliness, erosion or any other site characteristic that affect management. The soil phase map can be used for identifying the suitability of areas for growing specific crops or for other alternative uses and also for deciding the type of conservation structures needed. The detailed information on soil and site-characteristics like soil depth, surface soil texture, slope, erosion, gravelliness, AWC, LCC etc, with respect to each of the soil phase identified is given village/survey number wise for the microwatershed in Appendix-I.

4.1 Soils of granite gneiss landscape

In this landscape, 6 soil series are identified and mapped. Of these, BMN series occupies a maximum area of 227 ha (39%) followed by BLC 110 ha (19%), NGP 57 ha (10%), BDL 42 ha (7%), VNK 34 ha (6%) and JNK 10 ha (2%). Brief description of each series identified and number of soil phases mapped is given below.

4.1.1 Badiyala (BDL) Series: Badiyala soils are shallow (25-50 cm), well drained, have very dark brown, dark yellow brown and dark brown, slightly calcareous sandy loam soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation. The Badiyala series has been classified as a member of the coarse-loamy, mixed, isohyperthermic family of Fluventic Haplustepts

The thickness of the solum ranges from 28 to 50 cm. The thickness of A horizon ranges from 4 to 12 cm. Its colour is in 10YR hue with value 3 to 4 and chroma 3 to 4. The texture is loamy sand, sandy clay loam and sandy clay. The thickness of B horizon ranges from 27 to 45 cm. Its colour is in 10 YR and 7.5 YR hue with value 2 to 4 and chroma 3 to 4. Its texture is sandy loam to sandy clay loam and is slightly calcareous. The available water capacity is very low (<50mm/m).



Landscape and Soil Profile characteristics of Badiyala (BDL) Series

4.1.2 Vanakanahalli (VNK) Series: Vanakanahalli soils are shallow (25-50 cm), well drained, have dark reddish brown sandy clay red soils. They have developed from weathered granite gneiss and occur on very gently sloping uplands under cultivation. The Vanakanahalli series has been classified as a member of the fine, mixed isohyperthermic family of Typic (Paralithic) Haplustalfs.

The thickness of the solum ranges from 25 to 49 cm. The thickness of A horizon ranges from 7 to 16 cm. Its colour is in 2.5 YR and 5 YR with value 3 and chroma 2 to 4. The texture is sandy loam to sandy clay loam and sandy clay. The thickness of B horizon ranges from 20 to 40 cm. Its colour is in 2.5 YR and 5 YR hue with value 3 to 4 and chroma 3 to 4. Its texture is sandy clay. The available water capacity is very low (<50 mm/m).



Landscape and Soil Profile characteristics of Vanakanahalli (VNK) Series

4.1.3 Jinkera (JNK) Series: Jinkera soils are moderately shallow (50-75 cm), well drained, have very dark gray to very dark grayish brown and dark brown, slightly calcareous sandy clay loam soils. They are developed from weathered granite gneiss and occur on very gently sloping uplands under cultivation. The Jinkera series has been classified as a member of the fine-loamy, mixed, isohyperthermic family of Typic Haplustepts.

The thickness of the solum ranges from 51-75 cm. Thickness of A horizon ranges from 6 to 11 cm. Its colour is in hue 10 YR and 7.5 YR with value and chroma of 3 to 4. The texture varies from sandy loam to sandy clay. The thickness of B horizon ranges from 53 to 66 cm. Its colour is in 10 YR and 7.5 YR hue with value and chroma of 2 to 4. The texture varies from sandy clay loam to sandy clay and is slightly calcareous. The available water capacity is low (51-100 mm/m). Only one phase was identified and mapped.



Landscape and Soil Profile characteristics of Jinkera (JNK) Series

4.1.4 Balichakra (BLC) Series: Balichakra soils are moderately deep (75-100 cm), well drained, dark reddish brown to reddish brown, slightly calcareous sandy clay loam soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation. The Balichakra series has been classified as a member of the fine-loamy, mixed, isohyperthermic family of Typic Haplustalfs.

The thickness of the solum ranges from 80 to 100 cm. The thickness of A horizon ranges from 10 to 16 cm. Its colour is in hue 5 YR with value and chroma of 3 to 4. Its texture varies from sandy clay loam and sandy clay. The thickness of B horizon ranges from 70 to 88 cm. Its colour is in hue 2.5 YR and 5 YR with value 3 to 5 and chroma 3 to 4. Its texture is sandy clay loam to sandy clay and is slightly calcareous. The available water capacity is medium (101-150 mm/m).



Landscape and Soil Profile characteristics of Balichakra (BLC) Series

4.1.5 Naglapur (NGP) Series: Naglapur soils are deep (100-150 cm), moderately well drained, have black to very dark grayish brown, calcareous cracking clay soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation. The Naglapur series has been classified as a member of the very fine, smectitic (calcareous), isohyperthermic family of Typic Haplusterts.

The thickness of the solum ranges from 110 to 150 cm. The thickness of A horizon ranges from 6 to 25 cm. Its colour is in 10 YR hue with value 3 to 5 and chroma 1 to 3. The texture varies from sandy loam to sandy clay and clay. The thickness of B horizon ranges from 110 to 141 cm. Its colour is in 10 YR hue with value 2 to 3 and chroma 1 to 2. Texture is clay and is calcareous. The available water capacity is very high (>200 mm/m).



Landscape and Soil Profile characteristics of Naglapur (NGP) Series

4.1.6 Bhimanahalli (BMN) Series: Bhimanahalli soils are very deep (>150 cm), moderately well drained, very dark gray calcareous cracking clay soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation. The Bhimanahalli series has been classified as a member of the fine, smectitic (calcareous), isohyperthermic family of Typic Haplusterts.

The thickness of the solum is more than 150 cm. The thickness of A horizon ranges from 6 to 13 cm. Its colour is in 10 YR hue with value 3 to 4 and chroma 1 to 2 with clay texture. The thickness of B horizon ranges from 163 to 176 cm. Its colour is in 10 YR hue with value 2 to 4 and chroma 1. Its texture is clay and is calcareous. The available water capacity is very high (>200 mm/m).



Landscape and Soil Profile characteristics of Bhimanahalli (BMN) Series

Table: 4.1 Physical and Chemical Characteristics of Soil Series identified in Sudapur-2 microwatershed

Soil Series: Badiyala (BDL) **Pedon:** R-5

Location: 16°37'10.0"N 77°20'21.5", Gudalagunta village, Balichakra hobli, Yadgir taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bengaluru **Classification:** Coarse-loamy, mixed, isohyperthermic Fluventic Haplustepts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-12	Ap	87.13	7.04	5.83	10.03	24.32	23.61	23.51	5.67	<15	ls	6.27	2.44
12-28	Bw1	64.63	13.30	22.07	6.74	13.07	22.30	17.01	5.50	<15	scl	16.34	7.83
28-52	BC	73.11	12.02	14.87	3.93	16.03	26.89	18.41	7.86	<15	sl	12.94	5.47

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO ₃	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP
	Water	CaCl ₂	M KCl				Ca	Mg	K	Na	Total				
	dS m ⁻¹			%	%	cmol kg ⁻¹					%	%			
0-12	6.20	-	-	0.074	1.00	0.00	2.80	0.98	0.14	0.01	3.92	4.20	0.72	93	0.20
12-28	9.04	-	-	0.253	0.80	3.20	-	-	0.16	0.69	-	16.90	0.77	100	4.09
28-52	9.41	-	-	0.364	1.10	3.60	-	-	0.16	1.39	-	11.10	0.75	100	12.52

Contd...

Soil Series: Vanakanahalli (VNK) **Pedon:** R-15

Location: 16°43'49.5"N 77°17'17.9"E, Yaleri village, Balichakra hobli, Yadgiri taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bengaluru **Classification:** clayey, mixed isohyperthermic (Paralithic) Haplustalfs

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-18	Ap	82.61	8.09	9.30	6.77	8.59	21.13	34.58	11.53	-	ls	8.85	3.53
18-50	Bt	54.51	8.73	36.77	4.93	6.18	14.15	20.75	8.49	-	sc	18.88	11.63

Depth (cm)	pH (1:2.5)			E.C. (1:2.5) dS m ⁻¹	O.C. %	CaCO ₃ %	Exchangeable bases					CEC	CEC/Clay	Base saturation %	ESP %
	Water	CaCl ₂	M KCl				Ca	Mg	K	Na	Total				
0-18	5.37	-	-	0.11	0.60	0.00	2.96	1.45	0.13	0.14	4.68	6.27	0.67	75	2.22
18-50	4.71	-	-	0.05	0.81	0.00	5.56	2.24	0.10	0.05	7.95	13.31	0.36	60	0.38

Contd...

Soil Series: Jinkera (JNK) **Pedon:** R-1

Location: 16°45'13.5"N 77°10'59.8"E, Varkanahalli village, Yadgir hobli, Yadgir taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bengaluru **Classification:** Fine-loamy, mixed, isohyperthermic Typic Haplustepts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-15	Ap	66.84	13.62	19.54	12.15	21.22	11.23	12.56	9.68	10	sl	14.42	7.70
15-38	Bw1	59.08	12.11	28.81	12.53	12.42	17.85	8.77	7.52	20	scl	18.21	12.23
38-50	Bw2	68.21	11.68	20.11	17.90	21.81	10.60	10.80	7.10	10	scl	14.54	8.96

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO ₃	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP
	Water	CaCl ₂	M KCl				Ca	Mg	K	Na	Total				
				dS m ⁻¹	%	%	cmol kg ⁻¹					%	%		
0-15	8.42	-	-	0.148	0.70	0.65	-	-	0.15	0.03	-	14.50	0.74	100	0.18
15-38	8.38	-	-	0.226	0.31	2.21	-	-	0.09	0.23	-	21.70	0.75	100	1.05
38-50	8.40	-	-	0.195	0.25	1.17	-	-	0.07	0.19	-	15.90	0.79	100	1.23

Contd...

Soil Series: Balichakra (BLC) **Pedon:** T1/P2

Location: 16°33'25.0"N 77°20'52.3"E, Sowrashtralli village, Sydhapura hobli, Yadgir taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bengaluru **Classification:** Fine-loamy, mixed, isohyperthermic Typic Haplustalfs

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-8	Ap	65.46	8.38	26.16	12.51	18.72	18.82	10.44	4.96	-	scl	15.15	8.63
8-19	BA	63.48	8.16	28.36	12.80	15.84	17.21	12.49	5.14	-	scl	16.45	8.81
19-40	Bt	52.64	11.58	35.79	13.19	13.19	14.35	8.23	3.69	-	sc	21.49	10.36
40-75	BC	55.14	10.71	34.15	14.10	14.42	14.63	7.53	4.45	-	scl	17.77	8.99

Depth (cm)	pH (1:2.5)			E.C. (1:2.5) dS m ⁻¹	O.C. %	CaCO ₃ %	Exchangeable bases					CEC	CEC/Clay	Base saturation %	ESP %
	Water	CaCl ₂	M KCl				Ca	Mg	K	Na	Total				
0-8	6.75	-	-	0.19	0.72	0.00	12.18	3.10	0.43	0.22	15.92	16.80	0.64	95	1.31
8-19	7.23	-	-	0.12	0.68	0.84	11.37	2.50	0.23	0.18	14.28	14.77	0.52	97	1.24
19-40	7.13	-	-	0.08	0.50	0.48	13.80	2.82	0.18	0.09	16.89	17.66	0.49	96	0.51
40-75	7.07	-	-	0.07	0.35	0.84	13.00	2.90	0.17	0.10	16.16	17.55	0.51	92	0.57

Contd...

Soil Series: Naglapur (NGP) **Pedon:** R-8

Location: 16°52'84.1"N 77°22'99.4"E, Gurumitkal village, Gurumitkal hobli, Yadgir taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bengaluru **Classification:** Very fine, smectitic (calcareous), isohyperthermic Typic Haplusterts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-10	Ap	7.53	19.88	72.59	1.00	0.78	0.89	2.10	2.77	-	c	44.31	32.79
10-35	Bss1	6.55	18.76	74.68	0.80	0.92	0.80	1.72	2.30	-	c	43.09	31.62
35-60	Bss2	6.58	21.05	72.37	0.69	0.46	1.04	1.50	2.89	-	c	46.52	32.52
60-102	Bss3	7.48	19.74	72.78	1.61	1.38	0.69	1.61	2.19	-	c	51.12	35.62

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO ₃	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP			
	Water	CaCl ₂	M KCl				dS m ⁻¹	%	%	Ca	Mg					K	Na	Total
										cmol kg ⁻¹								
0-10	7.42	-	-	0.24	0.84	1.30	-	-	0.84	0.15	-	67.10	0.92	100	0.22			
10-35	8.52	-	-	0.291	0.64	2.86	-	-	0.17	0.29	-	65.20	0.87	100	0.45			
35-60	7.89	-	-	0.134	0.62	4.55	-	-	0.15	0.20	-	65.00	0.90	100	0.30			
60-102	8.68	-	-	0.213	0.54	8.32	-	-	0.17	0.15	-	64.10	0.88	100	0.24			

Contd...

Soil Series: Bhimanahalli (BMN) **Pedon:** R-3

Location: 16°31'82.4"N 77°12'70.8"E, Bheemanahalli village, Sydhapura hobli, Yadgir taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bengaluru **Classification:** Fine, smectitic (calcareous), isohyperthermic Typic Haplusterts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-8	Ap	20.34	19.94	59.72	2.68	5.03	3.75	5.25	3.64	-	c	50.19	33.49
8-40	Bss1	19.61	22.76	57.62	1.94	2.59	5.28	4.96	4.85	-	c	43.22	29.05
40-70	Bss2	21.25	17.65	61.10	3.02	5.26	3.91	5.48	3.58	-	c	44.30	30.25
70-120	Bss3	19.08	22.29	58.63	1.75	5.04	3.84	5.15	3.29	-	c	43.26	30.31
120-170	Bss4	11.11	20.44	68.45	2.04	1.93	1.70	2.83	2.61	-	c	51.33	33.51

Depth (cm)	pH (1:2.5)			E.C. (1:2.5) dS m ⁻¹	O.C. %	CaCO ₃ %	Exchangeable bases					CEC	CEC/Clay	Base saturation %	ESP %
	Water	CaCl ₂	M KCl				Ca	Mg	K	Na	Total				
0-8	8.2	-	-	0.284	0.72	4.94	-	-	1.20	0.34	-	52.70	0.88	100	0.65
8-40	8.44	-	-	0.139	0.40	7.28	-	-	0.30	0.48	-	52.06	0.90	100	0.93
40-70	8.32	-	-	0.202	0.40	6.37	-	-	0.18	0.40	-	52.52	0.86	100	0.77
70-120	9.3	-	-	0.282	0.36	6.89	-	-	0.27	0.38	-	50.97	0.87	100	0.75
120-170	8.47	-	-	0.305	0.37	8.19	-	-	0.28	0.91	-	58.19	0.85	100	1.57

INTERPRETATION FOR LAND RESOURCE MANAGEMENT

The most important soil and site characteristics that affect the land use and conservation needs of an area are land capability, land irrigability, soil depth, soil texture, coarse fragments, available water capacity, soil slope, soil erosion, soil reaction etc. These are interpreted from the data base generated through land resource inventory and several thematic maps are generated. These would help in identifying the areas suitable for growing crops and, soil and water conservation measures and structures needed thus helping to maintain good soil health for sustained crop production. The various interpretative and thematic maps generated are described below.

5.1 Land Capability Classification

Land capability classification is an interpretative grouping of soil map units (soil phases) mainly based on inherent soil characteristics, external land features and environmental factors that limit the use of land for agriculture, pasture, forestry, or other uses on a sustained basis (IARI, 1971). The land and soil characteristics used to group the land resources in an area into various land capability classes, subclasses and units are

Soil Characteristics: Depth, texture, gravelliness, calcareousness.

Land characteristics: Slope, erosion, drainage, rock outcrops.

Climate: Total rainfall and its distribution, and length of crop growing period.

The land capability classification system is divided into land capability classes, subclasses and units based on the level of information available. Eight land capability classes are recognized. They are

Class I: They are very good lands that have no limitations or very few limitations that restrict their use.

Class II: They are good lands that have minor limitations and require moderate conservation practices.

Class III: They are moderately good lands that have moderate limitations that reduce the choice of crops or that require special conservation practices.

Class IV: They are fairly good lands that have very severe limitations that reduce the choice of crops or that require very careful management.

Class V: Soils in these lands are not likely to erode, but have other limitations like wetness that are impractical to remove and as such not suitable for agriculture, but suitable for pasture or forestry with minor limitations.

Class VI: The lands have severe limitations that make them generally unsuitable for cultivation, but suitable for pasture or forestry with moderate limitations.

Class VII: The lands have very severe limitations that make them unsuitable for cultivation, but suitable for pasture or forestry with major limitations.

Class VIII: Soil and other miscellaneous areas (rock lands) that have very severe limitations that nearly preclude their use for any crop production, but suitable for wildlife, recreation and installation of wind mills.

The land capability subclasses are recognised based on the dominant limitations observed within a given land capability class. The subclasses are designated by adding a lower case letter like 'e', 'w', 's', or 'c' to the class numeral. The subclass "e" indicates that the main hazard is risk of erosion, "w" indicates drainage or wetness as a limitation for plant growth, "s" indicates shallow soil depth, coarse or heavy textures, calcareousness, salinity/alkalinity or gravelliness and "c" indicates limitation due to climate.

The land capability subclasses have been further subdivided into land capability units based on the kinds of limitations present in each subclass. Ten land capability units are used in grouping the soil map units. They are stony or rocky (0), erosion hazard (slope, erosion) (1), coarse texture (sand, loamy sand, sandy loam) (2), fine texture (cracking clay, silty clay) (3), slowly permeable subsoil (4), coarse underlying material (5), salinity/alkali (6), stagnation, overflow, high ground water table (7), soil depth (8) and fertility problems (9). The capability units thus identified have similar soil and land characteristics that respond similarly to a given level of management. The soils of the microwatershed have been classified upto land capability subclass level.

The 8 soil map units identified in Sudapur-2 microwatershed are grouped under 2 land capability classes and 3 land capability subclasses. An area of 480 ha (81%) in the microwatershed is suitable for agriculture. About 108 ha (18%) is under rock outcrops and 1 ha (<1%) is covered by others (water body & habitation) (Fig. 5.1).

Good lands (Class II) cover an area of about 69 per cent and are distributed in the major part of the microwatershed with minor problems of soil and erosion. Moderately good lands (Class III) cover an area of about 13 per cent and are distributed in the northeastern and northern part of the microwatershed with moderate problems of soil and erosion.




Fig. 5.1 Land Capability map of Sudapur-2 Microwatershed

5.2 Soil Depth

Soil depth refers to the depth of the soil occurring above the parent material or hard rock. The depth of the soil determines the effective rooting depth for plants and in accordance with soil texture, mineralogy and gravel content, the capacity of the soil column to hold water and nutrient availability. Soil depth is one of the most important soil characteristic that is used in differentiating soils into different soil series. The soil depth classes used in identifying soils in the field are very shallow (<25 cm), shallow (25-50 cm), moderately shallow (50-75 cm), moderately deep (75-100 cm), deep (100-150 cm) and very deep (>150 cm). They were used to classify the soils into different depth classes and a soil depth map was generated. The area extent and their geographical distribution in the microwatershed is given in Fig. 5.2.

Shallow (25-50 cm) soils occur in an area of 76 ha (13%) and are distributed in the northeastern and northern part of the microwatershed. Moderately shallow (50-75 cm) soils occur in an area of 10 ha (2%) and are distributed in the western part of the microwatershed. Moderately deep (75-100 cm) soils occur in an area of 110 ha (19%) and are distributed in the eastern, central, northeastern and northern part of the microwatershed. Deep (100-150 cm) soils occur in an area of 57 ha (10%) and are distributed in the eastern, southeastern and southern part of the microwatershed. Very deep (>150 cm) soils occur in an area of 227 ha (39%) and are distributed in the western, southwestern, northwestern and southern part of the microwatershed. An area of 394 ha (68%) soils in the microwatershed are potential where all climatically adapted long

duration crops can be grown. The problem soils occupy an area of 76 ha (13%) where only short duration crops can be grown occasionally and the probability of crop failure is very high.




Fig. 5.2 Soil Depth map of Sudapur-2 Microwatershed

5.3 Surface Soil Texture

Texture is an expression to indicate the coarseness or fineness of the soil as determined by the relative proportion of primary particles of sand, silt and clay. It has a direct bearing on the structure, porosity, adhesion and consistence. The surface layer of a soil to a depth of about 25 cm is the layer that is most used by crops and plants. The surface soil textural class provides a guide to understanding soil-water retention and availability, nutrient holding capacity, infiltration, workability, drainage, physical and chemical behaviour, microbial activity and crop suitability. The textural classes used for LRI were used to classify and a surface soil texture map was generated. The area extent and their geographical distribution in the microwatershed is shown in Figure 5.3.

An area of 77 ha (13%) of the microwatershed has soils that are loamy and are distributed in the northern and northeastern part. An area of 404 ha (68%) of the microwatershed has soils that are clayey and are distributed in the major part of the microwatershed. Both clayey and loamy soils have high potential for soil-water retention and availability, and nutrient retention and availability, but clay soils have more problems of drainage, infiltration, workability and other physical problems.




Fig. 5.3 Surface Soil Texture map of Sudapur-2 Microwatershed

5.4 Soil Gravelliness

Gravel is the term used for describing coarse fragments between 2 mm and 7.5 cm diameter and stones for those between 7.5 cm and 25 cm. The presence of gravel and stones in soil reduces the volume of soil responsible for moisture and nutrient storage, drainage, infiltration and runoff, and hinders plant growth by impeding root growth and seedling emergence, intercultural operations and farm mechanization. The gravelliness classes used in LRI were used to classify the soils and using these classes a gravelliness map was generated. The area extent and their geographic distribution in the microwatershed is shown in Figure 5.4.

Non gravelly (<15%) soils cover an area of 199 ha (34%) and are distributed in all parts of the microwatershed. These are the most productive soils, where all climatically adapted short and long duration crops can be grown. Gravelly (15-35%) soils occur in a maximum area of 281 ha (48%) and distributed in the northern, northeastern, southern, western and southwestern part of the microwatershed. These lands are low in moisture holding capacity and hence growing of short duration crops is ideal with best management practice.




Fig. 5.4 Soil Gravelliness map of Sudapur-2 Microwatershed

5.5 Available Water Capacity

The soil available water capacity (AWC) is estimated based on the ability of the soil column to retain water between the tensions of 0.33 and 15 bar in a depth of 100 cm or the entire solum if the soil is shallower. The AWC of the soils (soil series) as estimated by considering the soil texture, mineralogy, soil depth and gravel content (Sehgal *et al.*, 1990) and accordingly the soil map units were grouped into five AWC classes *viz*, very low (<50 mm/m), low (50-100 mm/m), medium (100-150 mm/m), high (150-200 mm/m) and very high (>200 mm/m) and using these values, an AWC map was generated. The area extent and their geographic distribution of different AWC classes in the microwatershed is given in Figure 5.5.

An area of about 76 ha (13%) in the microwatershed has soils that are very low (<50 mm/m) in available water capacity and is distributed in the northern and northeastern part of the microwatershed. An area of 120 ha (20%) is low (51-100 mm/m) and distributed in the northeastern, eastern, central, northern and western part. An area of about 284 ha (48%) is very high (>200 mm/m) in available water capacity and are distributed in the central, northwestern, western, southern, southeastern, southwestern and eastern part of the microwatershed.

An area of about 196 ha (33%) in the microwatershed has soils that are problematic with regard to available water capacity. Here, only short duration crops can be grown and probability of the crop failure is very high. These areas are best put to other

alternative uses. Potential soils cover about 48 per cent area where all climatically adapted long duration crops can be grown.




Fig. 5.5 Soil Available Water Capacity map of Sudapur-2 Microwatershed

5.6 Soil Slope

Soil slope refers to the inclination of the surface of the land. It is defined by gradient, shape and length, and is an integral feature of any soil as a natural body. Slope is considered important in soil genesis, land use and land development. The length and gradient of slope influences the rate of runoff, infiltration, erosion and deposition. The soil map units were grouped into single slope class and a slope map was generated showing the area extent and their geographic distribution in the microwatershed (Fig. 5.6).

An area of about 113 ha (19%) is nearly level (0-1%) lands and 367 ha (62%) of the microwatershed falls under very gently sloping (1-3% slope) lands. Thus these areas have high potential in respect of soil slopes. In these areas, all climatically adapted annual and perennial crops can be grown without much soil and water conservation and other land development measures.




Fig. 5.6 Soil Slope map of Sudapur-2 Microwatershed

5.7 Soil Erosion

Soil erosion refers to the wearing away of the earth's surface by the forces of water, wind and ice involving detachment and transport of soil by raindrop impact. It is used for accelerated soil erosion resulting from disturbance of the natural landscape by burning, excessive grazing and indiscriminate felling of forest trees and tillage, all usually by man. The erosion classes showing an estimate of the current erosion status as judged from field observations in the form of rills, gullies or a carpet of gravel on the surface are recorded. Four erosion classes, viz, slight erosion (e1), moderate erosion (e2), severe erosion (e3) and very severe erosion (e4) are recognized. The soil map units were grouped into different erosion classes and a soil erosion map generated. The area extent and their spatial distribution in the microwatershed is given in Figure 5.7.

Soils that are slightly eroded (e1 class) cover an area of 113 ha (19%) and are distributed in the western, southwestern and southern part of the microwatershed. Moderately eroded soils (e2 class) cover an area of 367 ha (62%) and are distributed in the major part of the microwatershed.

Maximum area of 367 ha (62%) in the microwatershed is problematic because of moderate erosion. For these areas, taking up of soil and water conservation and other land development measures are needed.




Fig. 5.7 Soil Erosion map of Sudapur-2 Microwatershed

FERTILITY STATUS

Soil fertility plays an important role in increasing crop yield. The adoption of high yielding varieties that require high amounts of nutrients has resulted in deficiency symptoms in crops and plants due to imbalanced fertilization and poor inherent fertility status as these areas are characterised by low rainfall and high temperatures. Hence, it is necessary to know the fertility (macro and micro nutrients) status of the soils of the watersheds for assessing the kind and amount of fertilizers required for each of the crop intended to be grown. For this purpose, the surface soil samples collected from the grid points (one soil sample at every 320 m interval) all over the microwatershed through land resource inventory in the year 2017 were analysed for pH, EC, organic carbon, available phosphorus and potassium, and for micronutrients like zinc, boron, copper, iron and manganese, and secondary nutrient sulphur.

Soil fertility data generated has been assessed and individual maps for all the nutrients for the microwatershed have been generated using Kriging method under GIS. The village/survey number wise fertility data for the microwatershed is given in Appendix-II.

6.1 Soil Reaction (pH)

The soil analysis of the Sudapur-2 microwatershed for soil reaction (pH) showed that an area of 261 ha (44%) is slightly acid (pH 6.0-6.5) and are distributed in the major part of the microwatershed. An area of 219 ha (37%) is neutral (pH 6.5-7.3) and are distributed in the southwestern, northwestern, central, western and southern part of the microwatershed (Fig. 6.1).

6.2 Electrical Conductivity (EC)

The Electrical Conductivity in entire area of the microwatershed is <2 dS/m (Fig. 6.2) and as such the soils are non-saline.

6.3 Organic Carbon

The soil organic carbon content (an index of available Nitrogen) is high (>0.75 %) in the entire area of the microwatershed (Fig. 6.3).




Fig.6.1 Soil Reaction (pH) map of Sudapur-2 Microwatershed




Fig.6.2 Electrical Conductivity (EC) map of Sudapur-2 Microwatershed




Fig.6.3 Soil Organic Carbon map of Sudapur-2 Microwatershed

6.4 Available Phosphorus

Available phosphorus content is low (<23 kg/ha) in a maximum area of about 306 ha (52%) and occur in the major part of the microwatershed. Medium (23-57 kg/ha) in an area of 108 ha (18%) and occur in the eastern, central, northern and northeastern part of the microwatershed. high (>57 kg/ha) in an area of 66 ha (11%) and occur in the northern and northeastern part of the microwatershed (Fig. 6.4).

6.5 Available Potassium

Available potassium content is medium (145-337 kg/ha) in the entire area of the microwatershed (Fig. 6.5).

6.6 Available Sulphur

Available sulphur content is medium (10-20 ppm) in an area of 130 ha (22%) and are distributed in the northern, northeastern and eastern part of the microwatershed. Low in a maximum area of about 350 ha (59%) and are distributed in the major part of the microwatershed (Fig. 6.6).

6.7 Available Boron

Available boron content is low (<0.5 ppm) in a maximum area of about 470 ha (80%) and are distributed in the all parts of the microwatershed. Medium (0.5-1.0ppm) in a small area of 9 ha (2%) and are distributed in the southern part of the microwatershed (Fig. 6.7).

6.8 Available Iron

Available iron content is sufficient (>4.5 ppm) in the entire area of the microwatershed (Fig 6.8).

6.9 Available Manganese

Available manganese content is sufficient (>1.0 ppm) in the entire microwatershed area (Fig 6.9).

6.10 Available Copper

Available copper content is sufficient (>0.2 ppm) in the entire microwatershed area (Fig 6.10).




Fig.6.4 Soil Available Phosphorus map of Sudapur-2 Microwatershed




Fig.6.5 Soil Available Potassium map of Sudapur-2 Microwatershed




Fig.6.6 Soil Available Sulphur map of Sudapur-2 Microwatershed




Fig.6.7 Soil Available Boron map of Sudapur-2 Microwatershed




Fig.6.8 Soil Available Iron map of Sudapur-2 Microwatershed




Fig.6.9 Soil Available Manganese map of Sudapur-2 Microwatershed




Fig.6.10 Soil Available Copper map of Sudapur-2 Microwatershed

6.11 Available Zinc

Available zinc content is deficient (<0.6 ppm) in an area of 258 ha (44%) and is distributed in the western, northwestern, central, southern, southwestern and southeastern part of the microwatershed. An area of about 222 ha (38%) is sufficient (>0.6 ppm) and is distributed in southwestern, northern, eastern, northwestern, central and northeastern part of the microwatershed (Fig 6.11).




Fig.6.11 Soil Available Zinc map of Sudapur-2 Microwatershed

LAND SUITABILITY FOR MAJOR CROPS

The soil and land resource units (soil phases) of Sudapur-2 microwatershed were assessed for their suitability for growing food, fodder, fibre and other horticulture crops by following the procedure as outlined in FAO, 1976 and 1983. Crop requirements were developed for each of the crop from the available research data and also by referring to Naidu *et. al.* (2006) and Natarajan *et. al* (2015). The soil and land characteristics (Table 7.1) were matched with the crop requirement (Tables 7.2 to 7.30) to arrive at the crop suitability. The soil and land characteristics table and crop requirement tables are given at the end of the chapter. In FAO land suitability classification, two orders are recognized. Order S-Suitable and Order N-Not suitable. The orders have classes, subclasses and units. Order-S has three classes, Class S1-Highly Suitable, Class S2-Moderately Suitable and Class S3- Marginally Suitable. Order N has two classes, N1-Currently not Suitable and N2- Permanently not Suitable. There are no subclasses within the Class S1 as they will have very minor or no limitations for crop growth. Classes S2, S3, N1 and N2 are divided into subclasses based on the kinds of limitations encountered. The limitations that affect crop production are ‘c’ for erratic rainfall and its distribution and length of growing period (LGP), ‘e’ for erosion hazard, ‘r’ for rooting condition, ‘t’ for lighter or heavy texture, ‘g’ for gravelliness or stoniness, ‘n’ for nutrient availability, ‘l’ for topography, ‘m’ for moisture availability, ‘w’ for drainage, ‘s’ for sodium and ‘z’ for calcareousness. These limitations are indicated as lower case letters to the Class symbol. For example, moderately suitable lands with the limitations of soil depth and erosion are designated as S2re. For the microwatershed, the soil mapping units were evaluated and classified up to subclass level.

Using the above criteria, the soil map units of the microwatershed were evaluated and land suitability maps for 29 major agricultural and horticultural crops were generated. The detailed information on the kind of suitability of each of the soil phase for the crops assessed are given village/ survey number wise for the microwatershed in Appendix-III.

7.1 Land Suitability for Sorghum (*Sorghum bicolor*)

Sorghum is one of the major food crop grown in Karnataka in an area of 10.47 lakh ha in Bijapur, Gulbarga, Raichur, Bidar, Belgaum, Dharwad, Bellary, Chitradurga, Mysore and Tumakuru districts. The crop requirements for growing sorghum (Table 7.2) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing sorghum was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.1.

Highly suitable (Class S1) lands for growing sorghum not occur in the microwatershed. Maximum area of about 404 ha (69%) is moderately suitable (Class S2)

for growing sorghum and are distributed in the major part of the microwatershed. They have minor limitations of rooting depth, calcareousness, texture and gravelliness. An area of about 76 ha (13%) is marginally suitable (Class S3) for growing sorghum and is distributed in the northern and northeastern part of the microwatershed with moderate limitations of rooting depth and texture.




Fig. 7.1 Land Suitability map of Sorghum

7.2 Land Suitability for Maize (*Zea mays*)

Maize is one of the most important food crop grown in an area of 13.37 lakh ha in almost all the districts of the State. The crop requirements for growing maize (Table 7.3) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing maize was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.2.

Highly suitable (Class S1) lands available for growing maize occur in an area of 33 ha (6%) and are distributed in the northern part of the microwatershed. Moderately suitable (Class S2) lands occur in an area of 371 ha (63%) and are distributed in the major part of the microwatershed with minor limitations of gravelliness, rooting depth, calcareousness and texture. Marginally suitable lands (Class S3) for growing maize occupy an area of 76 ha (13%) and occur in the northern and northeastern part of the microwatershed. They have moderate limitations of rooting depth and texture.




Fig. 7.2 Land Suitability map of Maize

7.3 Land Suitability for Bajra (*Pennisetum glaucum*)

Bajra is one of the most important millet crop grown in an area of 2.34 lakh ha in the northern districts of Karnataka state. The crop requirements for growing bajra (Table 7.4) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing bajra was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.3.

Highly suitable (Class S1) lands available for growing bajra occur in an area of 110 ha (19%) and are distributed in the northern, northeastern, central and eastern part of the microwatershed. Moderately suitable (Class S2) lands occur in an area of 294 ha (50%) and are distributed in the major part of the microwatershed with minor limitations of rooting depth, calcareousness and texture. Marginally suitable lands (Class S3) for growing bajra occupy an area of 76 ha (13%) and occur in the northern and northeastern part of the microwatershed. They have moderate limitation of rooting depth.




Fig. 7.3 Land Suitability map of Bajra

7.4 Land Suitability for Groundnut (*Arachis hypogaea*)

Groundnut is one of the major oilseed crop grown in an area of 6.54 lakh ha in Karnataka in most of the districts either as rainfed or irrigated crop. The crop requirements for growing groundnut (Table 7.5) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing groundnut was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.4.

Highly suitable (Class S1) lands available for growing groundnut occur in an area of 110 ha (19%) and are distributed in the northern, northeastern, central and eastern part of the microwatershed. Moderately suitable (Class S2) lands occur in an area of 10 ha (2%) and are distributed in the western part of the microwatershed with minor limitation of rooting depth. Marginally suitable lands (Class S3) for growing groundnut occupy an area of about 360 ha (61%) with moderate limitations of texture, calcareousness and rooting depth and are distributed in the major part of the microwatershed.




Fig. 7.4 Land Suitability map of Groundnut

7.5 Land Suitability for Sunflower (*Helianthus annus*)

Sunflower is one of the most important oilseed crop grown in an area of 4.1 lakh ha in the State in all the districts. The crop requirements for growing sunflower (Table 7.6) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing sunflower was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.5.

There are no highly suitable (Class S1) lands available for growing sunflower in the microwatershed. An area of about 394 ha (67%) is marginally suitable (Class S3) and is distributed in the major part of the microwatershed with moderate limitations of rooting depth, calcareousness and texture. Currently not suitable (Class N1) lands occur in an area of 76 ha (13%) and are distributed in the northern and northeastern part of the microwatershed with severe limitation of rooting depth.




Fig. 7.5 Land Suitability map of Sunflower

7.6 Land Suitability for Red gram (*Cajanus Cajan*)

Redgram is one of the most important pulse crop grown in an area of 7.28 lakh ha in almost all the districts of the State. The crop requirements for growing red gram (Table 7.7) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing redgram was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.6.

Highly suitable (Class S1) lands for growing red gram are not available in the microwatershed. An area of about 394 ha (67%) is moderately suitable (Class S2) and is distributed in the major part of the microwatershed with minor limitations of rooting depth, calcareousness and texture. An area of about 10 ha (2%) is marginally suitable (Class S3) and is distributed in the western part of the microwatershed with moderate limitation of rooting depth. Currently not suitable (Class N1) lands occur in an area of 76 ha (13%) and are distributed in the northern and northeastern part of the microwatershed with severe limitation of rooting depth.




Fig. 7.6 Land Suitability map of Redgram

7.7 Land Suitability for Bengal gram (*Cicer aerativum*)

Bengal gram is one of the most important pulse crop grown in about 9.39 lakh ha area in Bijapur, Raichur, Kalaburgi, Dharwad, Belgaum and Bellary districts. The crop requirements for growing Bengal gram (Table 7.8) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing Bengal gram was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.7.

No highly suitable (Class S1) lands available for growing bengal gram in the microwatershed. An area of about 284 ha (48%) is moderately suitable (Class S2) and is distributed in the major part of the microwatershed with minor limitation of calcareousness. Marginally suitable lands (Class S3) occupy an area of about 154 ha (26%) and are distributed in the western, eastern, northern, central and northeastern part of the microwatershed. They have moderate limitations of rooting depth and texture. Currently not suitable (Class N1) lands occur in an area of 42 ha (7%) and are distributed in the northern and northeastern part of the microwatershed with severe limitation of texture.




Fig. 7.7 Land Suitability map of Bengal gram

7.8 Land Suitability for Cotton (*Gossypium hirsutum*)

Cotton is one of the most important fibre crop grown in the State in about 8.75 lakh ha area in Raichur, Dharwad, Belgaum, Kalaburgi, Bijapur, Bidar, Bellary, Chitradurga and Chamarajnagar districts. The crop requirements for growing cotton (Table 7.9) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing cotton was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.8.

No highly suitable (Class S1) lands available for growing cotton in the microwatershed. An area of about 284 ha (48%) is moderately suitable (Class S2) and is distributed in the major part of the microwatershed with minor limitation of calcareousness. Marginally suitable lands (Class S3) occupy an area of about 154 ha (26%) and are distributed in the western, eastern, northern, central and northeastern part of the microwatershed. They have moderate limitations of rooting depth and texture. Currently not suitable (Class N1) lands occur in an area of 42 ha (7%) and are distributed in the northern and northeastern part of the microwatershed with severe limitation of texture.




Fig. 7.8 Land Suitability map of Cotton

7.9 Land Suitability for Chilli (*Capsicum annuum*)

Chilli is one of the most important spice crop grown in about 0.42 lakh ha in Karnataka State. The crop requirements for growing chilli (Table 7.10) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing chilli was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.9.

Highly suitable (Class S1) lands available for growing chilli occur in an area of 33 ha (6%) and are distributed in the northern part of the microwatershed. Moderately suitable (Class S2) lands occur in an area of 371 ha (63%) and are distributed in the major part of the microwatershed with minor limitations of gravelliness, rooting depth, calcareousness and texture. Marginally suitable lands (Class S3) for growing chilli occupy an area of 76 ha (13%) and occur in the northern and northeastern part of the microwatershed. They have moderate limitation of rooting depth.




Fig 7.9 Land Suitability map of Chilli

7.10 Land Suitability for Tomato (*Lycopersicon esculentum*)

Tomato is one of the most important vegetable crop grown in about 0.61 lakh ha covering almost all the district of the state. The crop requirements for growing tomato (Table 7.11) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing tomato was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.10.

Highly suitable (Class S1) lands available for growing tomato occur in an area of 33 ha (6%) and are distributed in the northern part of the microwatershed. Moderately suitable (Class S2) lands occur in an area of 87 ha (15%) and are distributed in the western, central northeastern and eastern part of the microwatershed with minor limitations of gravelliness and rooting depth. Marginally suitable lands (Class S3) for growing tomato occupy an area of 360 ha (61%) and occur in the major part of the microwatershed. They have moderate limitations of rooting depth and texture.




Fig 7.10 Land Suitability map of Tomato

7.11 Land Suitability for Brinjal (*Solanum melongena*)

Brinjal is one of the most important vegetable crop grown in the state. The crop requirements for growing brinjal (Table 7.12) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing brinjal was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.11.

Highly suitable (Class S1) lands for growing brinjal occur in an area of 33 ha (6%) and are distributed in the northern part of the microwatershed. Moderately suitable (Class S2) lands occur in an area of 87 ha (15%) and are distributed in the western, central northeastern and eastern part of the microwatershed with minor limitations of gravelliness and rooting depth. Marginally suitable lands (Class S3) for growing brinjal occupy an area of 360 ha (61%) and occur in the major part of the microwatershed. They have moderate limitations of rooting depth and texture.




Fig 7.11 Land Suitability map of Brinjal

7.12 Land Suitability for Onion (*Allium cepa* L.)

Onion is one of the most important vegetable crop grown in the state. The crop requirements for growing onion (Table 7.13) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing onion was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.12.

Highly suitable (Class S1) lands for growing onion occur in an area of 33 ha (6%) and are distributed in the northern part of the microwatershed. Moderately suitable (Class S2) lands occur in an area of 87 ha (15%) and are distributed in the western, central northeastern and eastern part of the microwatershed with minor limitations of gravelliness and rooting depth. Marginally suitable lands (Class S3) for growing onion occupy an area of 360 ha (61%) and occur in the major part of the microwatershed. They have moderate limitations of rooting depth, calcareousness and texture.




Fig 7.12 Land Suitability map of Onion

7.13 Land Suitability for Bhendi (*Abelmoschus esculentus*)

Bhendi is one of the most important vegetable crop grown in the state. The crop requirements for growing bhendi (Table 7.14) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing bhendi was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.13.

Highly suitable (Class S1) lands for growing bhendi occur in an area of 33 ha (6%) and are distributed in the northern part of the microwatershed. Moderately suitable (Class S2) lands occur in an area of 371 ha (63%) and are distributed in the major part of the microwatershed with minor limitations of gravelliness, rooting depth, calcareousness and texture. Marginally suitable lands (Class S3) for growing bhendi occupy an area of 76 ha (13%) and occur in the northern and northeastern part of the microwatershed. They have moderate limitation of rooting depth.




Fig 7.13 Land Suitability map of Bhendi

7.14 Land Suitability for Drumstick (*Moringa oleifera*)

Drumstick is one of the most important vegetable crop grown in about 2403 ha in the state. The crop requirements for growing drumstick (Table 7.15) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing drumstick was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.14.

No highly suitable (Class S1) lands available for growing drumstick in the microwatershed. Moderately suitable (Class S2) lands occur in an area of 110 ha (19%) and are distributed in the central, northern, northeastern and eastern part of the microwatershed with minor limitation of rooting depth. Marginally suitable lands (Class S3) for growing drumstick occupy an area of 294 ha (50%) and occur in the major part of the microwatershed. They have moderate limitations of rooting depth and calcareousness. Currently not suitable (Class N1) lands occur in an area of 76 ha (13%) and are distributed in the northern and northeastern part of the microwatershed with severe limitation of rooting depth.




Fig 7.14 Land Suitability map of Drumstick

7.15 Land Suitability for Mango (*Mangifera indica*)

Mango is one of the most important fruit crop grown in an area of 1.73 lakh ha in almost all the districts of the State. The crop requirements (Table 7.16) for growing mango were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing mango was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed is given in Figure 7.15.

Marginally suitable lands (Class S3) for growing mango occupy an area of 394 ha (67%) and occur in the major part of the microwatershed. They have moderate limitations of rooting depth and texture. Currently not suitable (Class N1) lands occur in an area of 76 ha (13%) and are distributed in the northern, western and northeastern part of the microwatershed with severe limitation of rooting depth.




Fig. 7.15 Land Suitability map of Mango

7.16 Land Suitability for Guava (*Psidium guajava*)

Guava is one of the most important fruit crop grown in an area of 6558 ha in almost all the districts of the State. The crop requirements (Table 7.17) for growing guava were matched with the soil-site characteristics (7.1) and a land suitability map for growing guava was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.16.

No highly suitable (Class S1) lands available for growing guava in the microwatershed. Moderately suitable (Class S2) lands occur in an area of 110 ha (19%) and are distributed in the central, northern, northeastern and eastern part of the microwatershed with minor limitation of rooting depth. Marginally suitable lands (Class S3) for growing guava occupy an area of 294 ha (50%) and occur in the major part of the microwatershed. They have moderate limitations of rooting depth, texture and calcareousness. Currently not suitable (Class N1) lands occur in an area of 76 ha (13%) and are distributed in the northern and northeastern part of the microwatershed with severe limitation of rooting depth.




Fig. 7.16 Land Suitability map of Guava

7.17 Land Suitability for Sapota (*Manilkara zapota*)

Sapota is one of the most important fruit crop grown in an area of 29373 ha in almost all the districts of the State. The crop requirements (Table 7.18) for growing sapota were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing sapota was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.17.

No highly suitable (Class S1) lands available for growing sapota in the microwatershed. Moderately suitable (Class S2) lands occur in an area of 110 ha (19%) and are distributed in the central, northern, northeastern and eastern part of the microwatershed with minor limitation of rooting depth. Marginally suitable lands (Class S3) for growing sapota occupy an area of 294 ha (50%) and occur in the major part of the microwatershed. They have moderate limitations of rooting depth and. Currently not suitable (Class N1) lands occur in an area of 76 ha (13%) and are distributed in the northern and northeastern part of the microwatershed with severe limitation of rooting depth.




Fig. 7.17 Land Suitability map of Sapota

7.18 Land Suitability for Pomegranate (*Punica granatum*)

Pomegranate is one of the most important fruit crop commercially grown in about 18488 ha in Karnataka, mainly in Bijapur, Bagalkot, Koppal, Gadag and Chitradurga districts. The crop requirements for growing pomegranate (Table 7.19) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing pomegranate was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.18.

There are no highly (Class S1) suitable lands available for growing pomegranate in the microwatershed. An area of about 394 ha (67%) is moderately suitable (Class S2) and is distributed in the major part of the microwatershed with minor limitations of rooting depth, calcareousness and texture. An area of about 10 ha (2%) is marginally suitable (Class S3) and is distributed in the western part of the microwatershed with moderate limitation of rooting depth. Currently not suitable (Class N1) lands occur in an area of 76 ha (13%) and are distributed in the northern and northeastern part of the microwatershed with severe limitation of rooting depth.




Fig 7.18 Land Suitability map of Pomegranate

7.19 Land Suitability for Musambi (*Citrus limetta*)

Musambi is one of the important fruit crop grown in an area of 3446 ha in almost all the districts of the State. The crop requirements for growing musambi (Table 7.20) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing musambi was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.19.

There are no highly (Class S1) suitable lands available for growing musambi in the microwatershed. An area of about 394 ha (67%) is moderately suitable (Class S2) and is distributed in the major part of the microwatershed with minor limitations of rooting depth and calcareousness. An area of about 10 ha (2%) is marginally suitable (Class S3) and is distributed in the western part of the microwatershed with moderate limitation of rooting depth. Currently not suitable (Class N1) lands occur in an area of 76 ha (13%) and are distributed in the northern and northeastern part of the microwatershed with severe limitation of rooting depth.




Fig. 7.19 Land Suitability map of Musambi

7.20 Land Suitability for Lime (*Citrus sp*)

Lime is one of the most important fruit crop grown in an area of 0.11 lakh ha in almost all the districts of the State. The crop requirements for growing lime (Table 7.21) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing lime was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7. 20.

There are no highly (Class S1) suitable lands available for growing lime in the microwatershed. An area of about 394 ha (67%) is moderately suitable (Class S2) and is distributed in the major part of the microwatershed with minor limitations of rooting depth and calcareousness. An area of about 10 ha (2%) is marginally suitable (Class S3) and is distributed in the western part of the microwatershed with moderate limitation of rooting depth. Currently not suitable (Class N1) lands occur in an area of 76 ha (13%) and are distributed in the northern and northeastern part of the microwatershed with severe limitation of rooting depth.




Fig. 7.20 Land Suitability map of Lime

7.21 Land Suitability for Amla (*Phyllanthus emblica*)

Amla is one of the medicinal fruit crop grown in almost all the districts of the State. The crop requirements for growing amla (Table 7.22) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing amla was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.21.

Highly suitable (Class S1) lands for growing amla occur in an area of 110 ha (19%) and are distributed in the northern, northeastern, central and eastern part of the microwatershed. Moderately suitable (Class S2) lands occur in an area of 10 ha (2%) and are distributed in the western part of the microwatershed with minor limitation of rooting depth. Marginally suitable lands (Class S3) for growing amla occupy an area of 360 ha (61%) and occur in the major part of the microwatershed. They have moderate limitations of rooting depth, calcareousness and texture.




Fig. 7.21 Land Suitability map of Amla

7.22 Land Suitability for Cashew (*Anacardium occidentale*)

Cashew is one of the most important plantation nut crop grown in an area of 0.7 lakh ha in almost all the districts of the State. The crop requirements for growing cashew (Table 7.23) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing cashew was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.22.

No highly suitable (Class S1) lands available for growing cashew in the microwatershed. Moderately suitable (Class S2) lands occur in an area of 110 ha (19%) and are distributed in the central, northern, northeastern and eastern part of the microwatershed with minor limitations of rooting depth and nutrient availability. Currently not suitable (Class N1) lands occur in an area of 370 ha (63%) and are distributed in the major part of the microwatershed with severe limitations of rooting depth, texture and nutrient availability.




Fig. 7.22 Land Suitability map of Cashew

7. 23 Land Suitability for Jackfruit (*Artocarpus heterophyllus*)

Jackfruit is one of the most important fruit crop grown in an area of 5368 ha in almost all the districts of the State. The crop requirements for growing jackfruit (Table 7.24) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing jackfruit was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.23.

No highly suitable (Class S1) lands available for growing jackfruit in the microwatershed. Moderately suitable (Class S2) lands occur in an area of 110 ha (19%) and are distributed in the central, northern, northeastern and eastern part of the microwatershed with minor limitations of rooting depth. Marginally suitable lands (Class S3) for growing jackfruit occupy an area of 294 ha (50%) and occur in the major part of the microwatershed. They have moderate limitations of rooting depth, texture and calcareousness. Currently not suitable (Class N1) lands occur in an area of 76 ha (13%) and are distributed in the northern and northeastern part of the microwatershed with severe limitation of rooting depth.




Fig. 7.23 Land Suitability map of Jackfruit

7.24 Land Suitability for Jamun (*Syzygium cumini*)

Jamun is an important fruit crop grown in almost all the districts of the State. The crop requirements for growing jamun (Table 25) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing jamun was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.24.

Marginally suitable lands (Class S3) for growing jamun occupy an area of 404 ha (69%) and occur in the major part of the microwatershed. They have moderate limitations of rooting depth and calcareousness. Currently not suitable (Class N1) lands occur in an area of 76 ha (13%) and are distributed in the northern and northeastern part of the microwatershed with severe limitation of rooting depth.




Fig. 7.24 Land Suitability map of Jamun

7.25 Land Suitability for Custard Apple (*Annona reticulata*)

Custard apple is one of the most important fruit crop grown in almost all the districts of the State. The crop requirements for growing custard apple (Table 7.26) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing custard apple was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.25.

Highly (Class S1) suitable lands for growing custard apple occupy an area of 110 ha (19%) and are distributed in the central, northern, northeastern and eastern part of the microwatershed. Moderately suitable (Class S2) lands occur in an area of 294 ha (50%) and are distributed in the major part of the microwatershed with minor limitations of rooting depth and calcareousness. Marginally suitable lands (Class S3) for growing custard apple occupy an area of 76 ha (13%) and occur in the northern and northeastern part of the microwatershed. They have moderate limitations of rooting depth and texture.




Fig. 7.25 Land Suitability map of Custard Apple

7.26 Land Suitability for Tamarind (*Tamarindus indica*)

Tamarind is one of the most important spice crop grown in almost all the districts of the state. The crop requirements for growing tamarind (Table 7.27) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing tamarind was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.26.

Marginally suitable lands (Class S3) for growing tamarind occupy an area of 394 ha (69%) and occur in the major part of the microwatershed. They have moderate limitations of rooting depth and calcareousness. Currently not suitable (Class N1) lands occur in an area of 86 ha (15%) and are distributed in the northern, western and northeastern part of the microwatershed with severe limitation of rooting depth.




Fig. 7.26 Land Suitability map of Tamarind

7.27 Land Suitability for Mulberry (*Morus nigra*)

Mulberry is one of the important leaf crop grown for rearing silk worms in about 1.6 lakh ha area in all the districts of the state. The crop requirements for growing mulberry (Table 7.28) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing mulberry was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.27.

No highly suitable (Class S1) lands available for growing mulberry in the microwatershed. Moderately suitable (Class S2) lands occur in an area of 110 ha (19%) and are distributed in the central, northern, northeastern and eastern part of the microwatershed with minor limitations of rooting depth. Marginally suitable lands (Class S3) for growing mulberry occupy an area of 294 ha (50%) and occur in the major part of the microwatershed. They have moderate limitations of rooting depth, texture and calcareousness. Currently not suitable (Class N1) lands occur in an area of 76 ha (13%) and are distributed in the northern and northeastern part of the microwatershed with severe limitation of rooting depth.




Fig 7.27 Land Suitability map of Mulberry

7.28 Land Suitability for Marigold (*Tagetes sps.*)

Marigold is one of the most important flower crop grown in an area of 9108 ha in almost all the districts of the State. The crop requirements (Table 7.29) for growing marigold were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing marigold was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.28.

Highly suitable (Class S1) lands available for growing marigold occur in an area of 33 ha (6%) and are distributed in the northern part of the microwatershed. Moderately suitable (Class S2) lands occur in an area of 371 ha (63%) and are distributed in the major part of the microwatershed with minor limitations of gravelliness, rooting depth, calcareousness and texture. Marginally suitable lands (Class S3) for growing marigold occupy an area of 76 ha (13%) and occur in the northern and northeastern part of the microwatershed. They have moderate limitation of rooting depth.




Fig. 7.28 Land Suitability map of Marigold

7.29 Land Suitability for Chrysanthemum (*Dendranthema grandiflora*)

Chrysanthemum is one of the most important flower crop grown in an area of 4978 ha in almost all the districts of the State. The crop requirements (Table 7.30) for growing chrysanthemum were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing chrysanthemum was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.29.

Highly suitable (Class S1) lands available for growing chrysanthemum occur in an area of 33 ha (6%) and are distributed in the northern part of the microwatershed. Moderately suitable (Class S2) lands occur in an area of 371 ha (63%) and are distributed in the major part of the microwatershed with minor limitations of gravelliness, rooting depth, calcareousness and texture. Marginally suitable lands (Class S3) for growing chrysanthemum occupy an area of 76 ha (13%) and occur in the northern and northeastern part of the microwatershed. They have moderate limitation of rooting depth.




Fig. 7.29 Land Suitability map of Chrysanthemum

Table 7.1 Soil-Site Characteristics of Sudapur-2 Microwatershed

Soil Map Units	Climate (P) (mm)	Growing period (Days)	Drain-age Class	Soil depth (cm)	Soil texture		Gravelliness		AWC (mm/m)	Slope (%)	Erosion	pH	EC (dSm ₁ ⁻¹)	ESP (%)	CEC [Cmol (p ⁺)kg ₁ ⁻¹]	BS (%)
					Sur-face	Sub-surface	Surface (%)	Sub-surface (%)								
BDLiB2	866	150	WD	25-50	sc	sl	<15	<15	<50	1-3	moderate	6.20	0.074	0.20	4.20	93
VNKmB2g1	866	150	WD	25-50	c	sc	15-35	<15	<50	1-3	moderate	5.37	0.11	2.22	6.27	75
JNKmB2	866	150	WD	50-75	c	scl	<15	<15	51-100	1-3	moderate	8.42	0.148	0.18	14.50	100
BLCcB2g1	866	150	WD	75-100	sl	scl	15-35	<15	51-100	1-3	moderate	6.75	0.19	1.31	16.80	95
BLCiB2	866	150	WD	75-100	sc	scl	<15	<15	51-100	1-3	moderate	6.75	0.19	1.31	16.80	95
NGPmB2g1	866	150	MW	100-150	c	c	15-35	<15	>200	1-3	moderate	7.42	0.24	0.22	67.10	100
BMNmA1	866	150	MW	>150	c	c	<15	<15	>200	0-1	slight	8.2	0.28	0.365	52.70	100
BMNmB2g1	866	150	MW	>150	c	c	15-35	<15	>200	1-3	moderate	8.2	0.28	0.365	52.70	100

*Symbols and abbreviations are according to Field Guide for LRI under Sujala-III Project, Karnataka

Table 7.2 Land suitability criteria for Sorghum

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	26–30	30–34; 24–26	34–40; 20–24	>40; <20
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, c (red), c (black)	scl, cl	ls, sl	-
	pH	1:2.5	5.5-7.8	5.0-5.5 7.8-9.0	>9.0	-
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	10-15
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

Table 7.3 Land suitability criteria for Maize

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	30-34	35-38 26-30	38-40 26-20	
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc	c (red), c (black)	ls, sl	-
	pH	1:2.5	5.5-7.8	5.0-5.5 7.8-9.0	>9.0	-
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	-
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

Table 7.4 Land suitability criteria for Bajra

Land use requirement		Rating				
Soil –site characteristics	Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)	
Climatic regime	Mean temperature in growing season	°C	28-32	33-38 24-27	39-40 20-23	<20
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm	500-750	400-500	200-400	<200
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl, scl, cl,sc,c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.8	5.0-5.5 7.8-9.0	5.5-6.0 >9.0	
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	15-35	35-60	>60	
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	1-3	3-5	5-10	>10

Table 7.5 Land suitability criteria for Groundnut

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	24–33	22–24; 33–35	20–22; 35–40	<20; >40
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. Well drained	Poorly drained	Very Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl	sl,cl, sc	c (red), c (black), ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<35	35-60	>60	
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.6 Land suitability criteria for Sunflower

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	24–30	30–34; 20–24	34–38; 16–20	>38; <16
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	mod. Well drained	-	Poorly to very drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	cl, sc,c (red), c (black)	scl	ls, sl	-
	pH	1:2.5	6.5-7.8	7.8-8.4 5.5-6.5	8.4-9.0; 5.0-5.5	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.7 Land suitability criteria for Redgram

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	30-35(G) 20-25(AV) 15-18 (F&PS) 35-40(M)	25-30(G) 20-25 (AV) 12-15 (F&PS) 30-35(M)	20-25(G) 15-20(AV) 10-12 (F&PS) 25-30(M)	< 20 <15 <10 <25
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. Well drained	Poorly drained	Very Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, c (red)	c (black),sl, scl, cl	ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-9.0	5.0-5.5 >9.0	-
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-50	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<1.0	1.0-2.0	>2.0	
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.8 Land suitability criteria for Bengal gram

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	20–25	25–30; 15–20	30–35; 10–15	>35; <10
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. Well drained	Poorly drained	Very Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	c(black)	-	c (red), scl, cl, sc	ls, sl
	pH	1:2.5	6.0-7.8	5.0-6.0 7.8-9.0	>9.0	-
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	-
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.9 Land suitability criteria for Cotton

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	22-32	>32	<19	-
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well to moderately well	Poorly drained/Some what excessively drained	-	very poorly/excessively drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, c (red,black)	cl	scl	ls, sl
	pH	1:2.5	6.5-7.8	7.8-8.4	5.5-6.5 8.4->9.0	<5.5
	CEC	C mol (p+)Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	50-100	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	<3	3-5	-	>5

Table 7.10 Land suitability criteria for Chilli

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	25-32	33-35 20-25	35-38 <20	>38
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc	c (black), sl	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.11 Land suitability criteria for Tomato

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	25-28	29-32 20-24	15-19 33-36	<15 >36
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl, scl, cl, sc, c (red)	-	ls, c(black)	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.12 Land suitability criteria for Brinjal

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	Well drained	Moderately well drained	Poorly drained	V. Poorly drained
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class				
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl, scl, cl, sc c (red)	-	ls, c (black)	-
	pH	1:2.5	6.0-7.3	7.3-8.4 5.0-6.0	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	>60
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.13 Land suitability criteria for Onion

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	20-30	30-35	35-40	>40
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately /imperfectly	-	Poorly to V poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl,scl,cl,sc,c (red)	-	c (Black),ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<1.0	1.0-2.0	2.0-4.0	<4
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.14 Land suitability criteria for Bhendi

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	25-28	29-32 20-24	15-19 33-36	<15 >36
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Imperfectly drained	Poorly to very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl,sc, c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.15 Land suitability criteria for Drumstick

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, scl, cl, c (red)	sl, c (black)	ls	s
	pH	1:2.5	6.0-7.3	5.0-5.5 7.3-7.8	5.5-6.0 7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<35	35-60	60-80	>80
Soil toxicity	Salinity (EC saturation extract)	ds/m				
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-10	-	>10

Table 7.16 Land suitability criteria for Mango

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-32	24-27 33-35	36-40	20-24
	Min temp. before flowering	°C	10-15	15-22	>22	-
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration	Days				
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V. Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	-	ls, sl, c (black)	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>150	100-150	75-100	<75
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.17 Land suitability criteria for Guava

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-32	33-36 24-27	37-42 20-23	
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	sl	c (black), ls	-
	pH	1:2.5	6.0-7.8	5.0-6.0	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.18 Land suitability criteria for Sapota

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-32	33-36 24-27	37-42 20-23	>42 <18
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	-	Poorly to very drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	sl	ls, c (black)	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.19 Land suitability criteria for Pomegranate

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	30-34	35-38 25-29	39-40 15-24	
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl,cl, sc, c (red)	c (black),sl	ls	-
	pH	1:2.5	5.5-7.8	7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.20 Land suitability criteria for Musambi

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-30	31-35 24-27	36-40 20-23	>40 <20
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately drained	poorly	Very poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c	sl	ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.21 Land suitability criteria for Lime

Land use requirement		Rating				
Soil –site characteristics	Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)	
Climatic regime	Mean temperature in growing season	°C	28-30	31-35 24-27	36-40 20-23	>40 <20
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately drained	poorly	Very poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c	sl	ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.22 Land suitability criteria for Amla

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. well drained	Poorly drained	V. Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	c (black)	ls, sl	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15-35	35-60	60-80	-
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

Table 7.23 Land suitability criteria for Cashew

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	32 to 34	28 to 32; 34 to 38	24 to 28; 38 to 40	<20; >40
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	-	sl, ls	c (black)
	pH	1:2.5	5.5-6.5	5.0-5.5 6.5-7.3	7.3-7.8	>7.8
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-10	>10	-

Table 7.24 Land suitability criteria for Jackfruit

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. well	Poorly	V. Poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	-	sl, ls, c (black)	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	>60
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10-

Table 7.25 Land suitability criteria for Jamun

Land use requirement		Rating				
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well	Mod. well	Poorly	V.Poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c(red)	sl, c (black)	ls	-
	pH	1:2.5	6.0-7.8	5.0-6.0	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>150	100-150	50-100	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	>60
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

Table 7.26 Land suitability criteria for Custard apple

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	Scl, cl, sc, c (red), c (black)	-	Sl, ls	-
	pH	1:2.5	6.0-7.3	5.5-6.0 7.3-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15-35	35-60	60-80	-
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	>5	-

Table 7.27 Land suitability criteria for Tamarind

Land use requirement		Rating				
Soil –site characteristics	Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)	
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod.well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl,sc, c (red)	sl, c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>150	100-150	75-100	<75
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

Table 7.28 Land suitability criteria for Mulberry

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	24–28	22–24; 28–32	32–38; 22–18	>38; <18
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V. Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, cl, scl	c (red)	c (black), sl, ls	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.8-8.4	7.3-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	0-35	35-60	60-80	>80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

Table 7.29 Land suitability criteria for Marigold

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	18-23	17-15 24-35	35-40 10-14	>40 <10
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl,scl, cl, sc, c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%				
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.30 Land suitability criteria for Chrysanthemum

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	18-23	17-15 24-35	35-40 10-14	>40 <10
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl,scl, cl, sc, c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%				
Erosion hazard	Slope	%	<3	3-5	5-10	>10

7.30 Land Management Units (LMUs)

The 8 soil map units identified in Sudapur-2 microwatershed have been grouped into 4 Land Management Units (LMU's) for the purpose of preparing a Proposed Crop Plan. Land Management Units are grouped based on the similarities in respect of the type of soil, the depth of the soil, the surface soil texture, gravel content, AWC, slope, erosion etc. and a Land Management Units map (Fig. 7.30) has been generated. These Land Management Units are expected to behave similarly for a given level of management.

The 8 map units that have been grouped into 4 Land Management Units along with brief description of soil and site characteristics are given below.

LMU	Soil map units	Soil and site characteristics
1	159.BMNmA1 63.BMNmB2g1 146.NGPmB2g1	Deep to very deep (100 to >150 cm), black calcareous clay soils, 0-3% slopes, non gravelly to gravelly (<15-35%), slight to moderate erosion.
2	155.BLCcB2g1 38.BLCiB2	Moderately deep (75 to 100 cm), sandy clay soils, 1-3% slopes, non gravelly to gravelly (<15-35%), moderate erosion.
3	152.JNKmB2	Moderately shallow (50 -75 cm), sandy clay loamy soils, 1-3% slopes, non gravelly (<15%), moderate erosion.
4	5.BDLiB2 109.VNKmB2g1	Shallow (25-50 cm), sandy loam to sandy clay soils, 1-3% slopes, non gravelly to gravelly (<15-35%), moderate erosion.




Fig. 7.30 Land Management Units Map- Sudapur-2 Microwatershed

7.31 Proposed Crop Plan for Sudapur-2 Microwatershed

After assessing the land suitability for the 29 crops, the Proposed Crop Plan has been prepared for the 4 identified LMUs by considering only the highly (Class S1) and moderately (Class S2) suitable lands for each of the 29 crops. The resultant proposed crop plan is presented below in Table 7.31.

Table 7.31 Proposed Crop Plan for Sudapur-2 Microwatershed

LMU	Soil Map Units	Survey Number	Field Crops/ Commercial crops	Horticulture Crops (Rainfed/Irrigated)	Suitable Interventions
1	159.BMNmA1 63.BMNmB2g1 146.NGPmB2g1 (Deep to very deep, black calcareous clay soils)	Gajarakota: 254,256,257,283,289,290,291,292,293,294,295,296,297,298,299,300,301,302,303,304,306,307,308,309,310,311,312,313,338,351,352,353,354,355,356,359,360,361,362,364,365,366,367,368,369,370,371,375,376, 377 Mitathapadamapalli: 188,189,190,191,192/1,192/2,193,194,195,196/1,196/2,196/3, 197,255, 256,257,258,259,260,262/14,262/20,262/21, 265, 269/2,270 Yadhalapura : 26	Maize, Sorghum, Sunflower, Cotton, Red gram, Bengalgram, Bajra	Fruit crops: Lime, Musambi, Custard apple, Pomegranate Vegetables: Chilli, Bhendi Flowers: Marigold, Chrysanthemum	Application of FYM, Bio-fertilizers and micronutrients, drip irrigation, mulching, suitable soil and water conservation practices
2	155.BLCcB2g1 38.BLCiB2 (Moderately deep sandy clay loam soils)	Mitathapadamapalli : 261,263,264 Siddhapura.B : 42,44/1,46,51,52,53/1,53/2,54,55,57,58,59,60,61,90,91,92, 94/1,94/2, 94/3,95,96, 97,98,99	Sunflower, Sorghum, Maize, Groundnut, Red gram, Bajra	Fruit crops: Musambi, Sapota, Pomegranate, Amla, Custard apple, Guava, Jackfruit, Lime Vegetables: Tomato, Onion, Bhendi, Chilli, Brinjal, Drumstick, Coriander Flowers: Marigold, Chrysanthemum	Application of FYM, Bio-fertilizers and micronutrients, drip irrigation, mulching, suitable soil and water conservation practices
3	152.JNKmB2 (Moderately shallow, sandy clay loam soils)	Gajarakota : 252,255,374,383	Maize, Sorghum Groundnut, Bajra	Fruit crops: Amla, Custard apple Vegetables: Tomato, Chilli, Brinjal, Bhendi, Onion Flowers: Marigold, Chrysanthemum	Application of FYM, Bio-fertilizers and micronutrients, drip irrigation, mulching, suitable soil and water conservation practices
4	5.BDLiB2 109.VNKmB2g1 (Shallow sandy loam to sandy clay soils)	Siddhapura.B : 14/1,14/2,14/3,15,16,17,18/1,37,44/2,45,47,48,62,63,64,65,66,67,68, 69,70,71,72,73,74/1,74/2,74/3,75,76,77,78,79,80,81,82,83,84,85,86,87,88,89,107/1, 107/2,108/3,109	-	Agri-Silvi-Pasture: Hybrid Napier, <i>Styloxanthes hamata</i> , <i>Styloxanthes scabra</i>	Use of short duration varieties, sowing across the slope, drip irrigation and mulching is recommended

SOIL HEALTH MANAGEMENT

8.1 Soil Health

Soil health is basic to plant health and plant health is basic to human and bovine health. Soil is fundamental to crop production. Without soil, no food could be produced nor would livestock be fed on a large scale. Because it is finite and fragile, soil is a precious resource that requires special care from its users.

Soil health or the capacity of the soil to function is critical to human survival. Soil health has been defined as: “the capacity of the soil to function as a living system without adverse effect on the ecosystem”. Healthy soils maintain a diverse community of soil organisms that help to form beneficial symbiotic associations with plant roots, recycle essential plant nutrients, improve soil structure with positive repercussions for soil, water and nutrient holding capacity and ultimately improve crop production and also contribute to mitigating climate change by maintaining or increasing its carbon content.

Functional interactions of soil biota with organic and inorganic components, air and water determine a soil’s potential to store and release nutrients and water to plants and to promote and sustain plant growth. Thus, maintaining soil health is vital to crop production and conserve soil resource base for sustaining agriculture.

The most important characteristics of a healthy soil are

- Good soil tilth
- Sufficient soil depth
- Good water storage and good drainage
- Adequate supply, but not excess of nutrients
- Large population of beneficial organisms
- Small proportion of plant pathogens and insect pests
- Low weed pressure
- Free of chemicals and toxins that may harm the crop
- Resistance to degradation
- Resilience when unfavorable conditions occur

Characteristics of Sudapur-2 Microwatershed

- ❖ The soil phases identified in the microwatershed belonged to the soil series of BMN series occupying a maximum area of 227 ha (39%) followed by BLC 110 ha (19%), NGP 57 ha (10%), BDL 42 ha (7%), VNK 34 ha (6%) and JNK 10 ha (2%).
- ❖ As per land capability classification entire area of the microwatershed falls under arable land category (Class II & III). The major limitations identified in the arable lands were soil and erosion.
- ❖ On the basis of soil reaction, about 261 ha (44%) area is slightly acid and 219 ha (37%) is neutral in soil reaction.

❖ **Soil Health Management**

The following actions are required to improve the current land husbandry practices that provide a sound basis for the successful adoption of sustainable crop production system.

Acid soils

Acid soils cover about 261 ha area in the microwatershed.

1. Growing of crops suitable for a particular soil pH.
2. Amelioration of soils through the application of amendments (liming materials).

Liming materials:

1. CaCO_3 (Calcium Carbonate).
2. Dolomite [$\text{Ca Mg} (\text{CO}_3)_2$]
3. Quick lime (Cao)
4. Slaked lime [$\text{Ca} (\text{OH})_2$]

For normal pH and pH 4.8 (35 t/ha) and pH 6.0-7.0 (4 t/ha) lime is required.

Alkaline soils

Alkaline soils are not occurring in the microwatershed.

1. Regular addition of organic manure, green manuring, green leaf manuring, crop residue incorporation and mulching needs to be taken up to improve the soil organic matter status.
2. Application of biofertilizers (Azospirillum, Azotobacter, Rhizobium).
3. Application of 25% extra N and P (125 % RDN&P).
4. Application of ZnSO_4 – 12.5 kg/ha (once in three years).
5. Application of Boron – 5kg/ha (once in three years).

Neutral soils

Neutral soils occur in 219 ha area in the microwatershed.

1. Regular addition of organic manure, green manuring, green leaf manuring, crop residue incorporation and mulching needs to be taken up to improve the soil organic matter status.
2. Application of biofertilizers, (Azospirillum, Azotobacter, Rhizobium).
3. Application of 100 per cent RDF.
4. Need based micronutrient applications.

Besides the above recommendations, the best transfer of technology options are also to be adopted.

Soil Degradation

Soil erosion is one of the major factor affecting the soil health in the microwatershed. Out of total 589 ha area in the microwatershed, about 113 ha (19%) is

suffering from slight erosion and 367 ha (62%) is suffering from moderate erosion. In areas of moderate erosion immediate soil and water conservation and, other land development and land husbandry practices are required for restoring soil health.

Dissemination of Information and Communication of Benefits

Any large scale implementation of soil health management requires that supporting information is made available widely, particularly through channels familiar to farmers and extension workers. Given the very high priority attached to soil-health especially by the Central Government on issuing Soil-Health Cards to all the farmers, media outlets like Regional, State and National Newspapers, Radio and Dooradarshan programs in local languages but also modern information and communication technologies such as Cellular phones and the Internet, which can be much more effective in reaching the younger farmers.

Inputs for Net Planning (Saturation Plan) and Interventions needed

Net planning (Saturation Plan) in IWMP is focusing on preparation of

1. Soil and Water Conservation Plan for each plot or farm.
2. Productivity enhancement measures/ interventions for existing crops/livestock/other farm enterprises.
3. Diversification of farming mainly with perennial horticultural crops and livestock.
4. Improving livelihood opportunities and income generating activities.

In this connection, how various outputs of Sujala-III are of use in addressing these objectives of Net Planning (Saturation Plan) are briefly presented below.

- ❖ **Soil Depth:** The depth of a soil decides the amount of moisture and nutrients it can hold, what crops can be taken up or not, depending on the rooting depth and the length of growing period available for raising any crop. Deeper the soil, better for a wide variety of crops. If sufficient depth is not available for growing deep rooted crops, either choose medium or short duration crops or deeper planting pits need to be opened and additional good quality soil brought from outside has to be filled into the planting pits.
- ❖ **Surface Soil Texture:** Lighter soil texture in the top soil means, better rain water infiltration, less run-off and soil moisture conservation, less capillary rise and less evaporation losses. Lighter surface textured soils are amenable to good soil tilth and are highly suitable for crops like groundnut, root vegetables (carrot, raddish, potato etc) but not ideal for crops that need stagnant water like lowland paddy. Heavy textured soils are poor in water infiltration and percolation. They are prone for sheet erosion; such soils can be improved by sand mulching. The technology that is developed by the AICRP-Dryland Agriculture, Vijayapura, Karnataka can be adopted.

- ❖ **Gravelliness:** More gravel content is favorable for run-off harvesting but poor in soil moisture storage and nutrient availability. It is a significant parameter that decides the kind of crop to be raised.
- ❖ **Land Capability Classification:** The land capability map shows the areas suitable and not suitable for agriculture and the major constraints in each of the plot/survey number. Hence, one can decide what kind of enterprise is possible in each of these units. In general, erosion and soil are the major constraints in Sudapur-2 microwatershed.
- ❖ **Organic Carbon:** The OC content (an index of available Nitrogen) is high (>0.75%) in the entire area of the microwatershed.
- ❖ **Promoting green manuring:** Growing of green manuring crops costs Rs. 1250/ha (green manuring seeds) and about Rs. 2000/ha towards cultivation that totals to Rs. 3250/- per ha. On the other hand, application of organic manure @ 10 tons/ha costs Rs. 5000/ha. The practice needs to be continued for 2-3 years or more.
- ❖ **Available Phosphorus:** Available Phosphorus is medium (23-57 kg/ha) in an area 108 ha (18%) and low (<23 kg/ha) in an area of 306 ha (52%) of the microwatershed. In medium and low areas, for all the crops 25% additional P needs to be applied.
- ❖ **Available Potassium:** Available potassium is medium (145-337 kg/ha) in the entire area of the microwatershed. All the plots, where available potassium is low and medium, for all the crops, additional 25% potassium may be applied.
- ❖ **Available Sulphur:** Available sulphur is a very critical nutrient for oilseed crops. It is medium in an area of 130 ha (22%) and low in an area of 350 ha (59%) of the microwatershed. Medium and low areas need to be applied with magnesium sulphate or gypsum or Factamphos (p) fertilizer (13% sulphur) for 2-3 years for the deficiency to be corrected.
- ❖ **Available Boron:** An area of 470 ha (80%) is low (<0.5 ppm) and 9 ha (2%) is medium (0.5-1.0 ppm) in available boron. Application of sodium tetra borate @ 10 kg/ha as soil application or 0.2 % borax as foliar spray is recommended for low and medium areas.
- ❖ **Available Iron:** Entire area of the microwatershed is sufficient (>4.5 ppm) in available iron content.
- ❖ **Available Manganese:** All the soils in the microwatershed are sufficient (>1.0 ppm) in available manganese.
- ❖ **Available Copper:** All the soils in the microwatershed are sufficient (>0.2 ppm) in available copper.
- ❖ **Available Zinc:** An area of about 258 ha (44%) is deficient (<0.6 ppm) and an area of 222 ha (38%) is sufficient in available zinc content. Application of zinc sulphate @25 kg/ha is recommended for deficient areas.
- ❖ **Soil Alkalinity:** Alkaline soils are not occurring in the microwatershed. Alkaline soils area need application of gypsum and wherever calcium is in excess, iron pyrites and


element sulphur can be recommended. Management practices like treating repeatedly with good quality water to drain out the excess salts and provision of subsurface drainage and growing of salt tolerant crops like Casuarina, Acacia, Neem, Ber etc, are recommended.

Land Suitability for various crops: Areas that are highly, moderately and marginally suitable and not suitable for growing various crops are indicated. Along with the suitability, various constraints that are limiting the productivity are also indicated. For example, in case of cotton, gravel content, rooting depth and salinity/alkalinity are the major constraints in various plots. With suitable management interventions, the productivity can be enhanced. In order to increase the water holding capacity of light textured soils, growing of green manure crops and application of organic manure is recommended.

SOIL AND WATER CONSERVATION TREATMENT PLAN

For preparing soil and water conservation treatment plan for Sudapur-2 microwatershed, the land resource inventory database generated under Sujala-III project has been transformed as information through series of interpretative (thematic) maps using soil phase map as a base. The various thematic maps (1:7920 scale) generated were

- Soil depth
- Surface soil texture
- Available water capacity
- Soil slope
- Soil gravelliness
- Land capability
- Present land use and land cover
- Crop suitability
- Rainfall
- Hydrology
- Water Resources
- Socio-economic data
- Contour plan with existing features- network of waterways, pothissa boundaries, cut up/ minor terraces etc.
- Cadastral map (1:7920 scale)
- Satellite imagery (1:7920 scale)



Apart from these, Hand Level/ Hydro Marker/ Dumpy Level/ Total Station and Kathedars' List to be collected.

Steps for Survey and Preparation of Treatment Plan

The boundaries of Land User Groups' and Survey No. boundaries are traced in the field.

- Naming of user groups and farmers
- Identification of arable and non arable lands
- Identification of drainage lines and gullies
- Identification of non treatable areas
- Identification of priority areas in the arable lands
- Treatment plan for arable lands
- Location of water harvesting and recharge structures

9.1 Treatment Plan

The treatment plan recommended for arable lands is briefly described below


9.1.1 Arable Land Treatment

A. BUNDING

Steps for Survey and Preparation of Treatment Plan		USER GROUP-1 CLASSIFICATION OF GULLIES
<ul style="list-style-type: none"> Cadastral map (1:7920 scale) is enlarged to a scale of 1:2500 scale Existing network of waterways, pottissa boundaries, grass belts, natural drainage lines/ watercourse, cut ups/ terraces are marked on the cadastral map to the scale Drainage lines are demarcated into 		
Small gullies	(up to 5 ha catchment)	
Medium gullies	(5-15 ha catchment)	
Ravines	(15-25 ha catchment) and	
<i>Halla/Nala</i>	(more than 25ha catchment)	

Measurement of Land Slope

Land slope is estimated or determined by the study and interpretation of contours or by measurement in the field using simple instruments like Hand Level or Hydromarker.



$$\text{FALL: } 1.5 - 0.6 = 0.9 \text{ m.}$$

Vertical and Horizontal intervals between bunds as recommended by the Watershed Development Department.

Slope percentage	Vertical interval (m)	Corresponding Horizontal Distance (m)
2 - 3%	0.6	24
3 - 4%	0.9	21
4 - 5%	0.9	21
5 - 6%	1.2	21
6 - 7%	1.2	21

Note: (i) The above intervals are maximum.

(ii) Considering the slope class and erosion status (A1... A=0-1 % slope, 1= slight erosion) the intervals have to be decided.

Bund length recording: Considering the contour plan and the existing grass belts/partitions, the bunds are aligned and lengths are measured.

Section of the Bund

Bund section is decided considering the soil texture class and gravelliness class (bg₀... b=loamy sand, g₀ = <15% gravel). The recommended Sections for different soils are given below.

Recommended Bund Section

Top width (m)	Base width (m)	Height (m)	Side slope (Z:1;H:V)	Cross section (sq m)	Soil Texture	Remarks
0.3	0.9	0.3	01:01	0.18	Sandy loam	Vegetative bund
0.3	1.2	0.3	1.5:1	0.225	Sandy clay	
0.3	1.2	0.5	0.9:1	0.375	Red gravelly soils	
0.3	1.2	0.6	0.75:1	0.45		
0.3	1.5	0.6	01:01	0.54	Red sandy loam	
0.3	2.1	0.6	1.5:1	0.72	Very shallow black clayey soils	
0.45	2	0.75	01:01	0.92		
0.45	2.4	0.75	1.3:1	1.07	Shallow black clayey soils	
0.6	3.1	0.7	1.78:1	1.29	Medium black clayey soils	
0.5	3	0.85	1.47:1	1.49		

Formation of Trench cum Bund

Dimensions of the Borrow Pits/Trenches to be excavated (machinery are decided considering the Bund Section).

Details of Borrow Pit dimensions are given below:

TRENCH CUM BUND

'A' FRAME FOR INTERBUND MANAGEMENT

1. ಸುಮಾರು 1.2 ಮೀಟರ್ ಉದ್ದವು
2. ಸುಮಾರು 1.2 ಮೀಟರ್ ವಿಸ್ತಾರ/ನಾಟಿ

ಇಳಿಬಂಡು

ಸುಮಾರು 1.2 ಮೀಟರ್

Size of Borrow Pits/ Trench recommended for Trench cum Bund (by machinery)

Bund section	Bund length	Earth quantity	Pit				Berm (pit to pit)	Soil depth class
			L(m)	W(m)	D(m)	Quantity (m ³)		
m ²	m	m ³					m	
0.375	6	2.25	5.85	0.85	0.45	2.24	0.15	Shallow
0.45	6	2.7	5.4	1.2	0.43	2.79	0.6	Shallow
0.45	6	2.7	5	0.85	0.65	2.76	1	Moderately Shallow
0.54	5.6	3.02	5.5	0.85	0.7	3.27	0.1	Moderately shallow
0.54	5.5	2.97	5	1.2	0.5	3	0.5	Shallow
0.72	6.2	4.46	6	1.2	0.7	5.04	0.2	Moderately shallow
0.72	5.2	3.74	5.1	0.85	0.9	3.9	0.1	Moderately deep

B. Water Ways

- Existing waterways are marked on the cadastral map (1:7920 scale) and their dimensions are recorded.
- Considering the contour plan of the MWS, additional waterways/ modernization of the existing ones can be thought of.
- The design details are given in the Manual.

C. Farm Ponds

Waterways and the catchment area will give an indication on the size of the Farm Pond. Location of the pond can be decided based on the contour plan/ field condition and farmers' need/desire.

D. Diversion Channel

Existing EPT/ CPT are marked on the cadastral map. Looking to the need, these can be modernized or fresh diversion channel can be proposed and runoff from this can be stored in *Gokatte*/ Recharge ponds.

9.1.2 Non-Arable Land Treatment

Depending on the gravelliness and crops preferred by the farmers, the concerned authorities can decide appropriate treatment plan. The recommended treatments may be Contour Trench, Staggered Trench, Crescent Bund, Boulder Bund or Pebble Bund.

9.1.3 Treatment of Natural Water Course/ Drainage Lines

- a) The cadastral map has to be updated as regards the network of drainage lines (gullies/*nalas/hallas*) and existing structures are marked to the scale and storage capacity of the existing water bodies are documented.
- b) The drainage line will be demarcated into Upper Reach, Middle Reach and Lower Reach.
- c) Considering the Catchment, *Nala* bed and bank conditions, suitable structures are decided.
- d) Number of storage structures (Check dam/*Nala* bund/ Percolation tank) will be decided considering the commitments and available runoff from water budgeting and quality of water in the wells and site suitability.
- e) Detailed Leveling Survey using Dumpy Level / Total Station has to be carried out to arrive at the site-specific designs as shown in the Manual.
- f) The location of ground water recharge structures are decided by examining the lineaments and fracture zones from geological maps.
- g) Rainfall intensity data of the nearest Rain Gauge Station is considered for Hydrologic Designs.
- h) Silt load to the Storage/Recharge Structures is reduced by providing vegetative, boulder and earthen checks in the natural water course. Location and design details are given in the Manual.

9.2 Recommended Soil and Water Conservation Measures

The appropriate conservation structures best suited for each of the land parcel/ survey number (Appendix-I) are selected based on the slope per cent, severity of erosion, amount of rainfall, land use and soil type. The different kinds of conservation structures recommended are:

1. Graded / Strengthening of Bunds
2. Trench cum Bunds (TCB)
3. Trench cum Bunds / Strengthening
4. Crescent Bunds

A map (Fig. 9.1) showing soil and water conservation plan with different kinds of structures recommended has been prepared which shows the spatial distribution and extent of area. Small area of about 144 ha (24%) needs Trench Cum Bunding. An area of about 223 ha (38%) needs Graded Bunding and an area of about 113 ha (19%) needs strengthening of existing bunds.

The conservation plan prepared may be presented to all the stakeholders including farmers and after considering their suggestions, the conservation plan for the microwatershed may be finalised in a participatory approach.




Fig. 9.1 Soil and Water Conservation Plan map of Sudapur-2 Microwatershed

9.3 Greening of Microwatershed

As part of the greening programme in the watersheds, it is envisaged to plant a variety of horticultural and other tree plants that are edible, economical and produce lot of biomass which helps to restore the ecological balance in the watersheds. The lands that are suitable for greening programme are non-arable lands (land capability classes V, VI VII and VIII) and also the lands that are not suitable or marginally suitable and field bunds for growing annual and perennial crops. The method of planting these trees is given below.

It is recommended to open pits during the 1st week of March along the contour and heap the dugout soil on the lower side of the slope in order to harness the flowing water and facilitate weathering of soil in the pit. Exposure of soil in the pit also prevents spread of pests and diseases due to scorching sun rays. The pits should be filled with mixture of soil and organic manure during the second week of April and keep ready with sufficiently tall seedlings produced either in poly bags or in root trainer nurseries so that planting can be done during the 2nd or 3rd week of April depending on the rainfall.

The tree species suitable for the area considering rainfall, temperature and adaptability is listed below; waterlogged areas are recommended to be planted with species like Nerale (*Syzgium cumini*) and Bamboo. Dry areas are to be planted with species like Honge, Bevu, Seetaphal etc.

Dry Deciduous Species			Temp (°C)	Rainfall (mm)
1.	Bevu	<i>Azadiracta indica</i>	21–32	400 –1,200
2.	Tapasi	<i>Holoptelia integrifolia</i>	20-30	500 - 1000
3.	Seetaphal	<i>Anona Squamosa</i>	20-40	400 - 1000
4.	Honge	<i>Pongamia pinnata</i>	20 -50	500– 2,500
5.	Kamara	<i>Hardwickia binata</i>	25 -35	400 - 1000
6.	Bage	<i>Albezzia lebbek</i>	20 - 45	500 - 1000
7.	Ficus	<i>Ficus bengalensis</i>	20 - 50	500–2,500
8.	Sisso	<i>Dalbargia Sissoo</i>	20 - 50	500 -2000
9.	Ailanthus	<i>Ailanthus excelsa</i>	20 - 50	500 - 1000
10.	Hale	<i>Wrightia tinctoria</i>	25 - 45	500 - 1000
11.	Uded	<i>Steriospermum chelanoides</i>	25 - 45	500 -2000
12.	Dhupa	<i>Boswella Serrata</i>	20 - 40	500 - 2000
13.	Nelli	<i>Emblca Officinalis</i>	20 - 50	500 -1500
14.	Honne	<i>Pterocarpus marsupium</i>	20 - 40	500 - 2000
Moist Deciduous Species			Temp (°C)	Rainfall (mm)
15.	Teak	<i>Tectona grandis</i>	20 - 50	500-5000
16.	Nandi	<i>Legarstroemia lanceolata</i>	20 - 40	500 - 4000
17.	Honne	<i>Pterocarpus marsupium</i>	20 - 40	500 - 3000
18.	Mathi	<i>Terminalia alata</i>	20 -50	500 - 2000
19.	Shivane	<i>Gmelina arborea</i>	20 -50	500 -2000
20.	Kindal	<i>T.Paniculata</i>	20 - 40	500 - 1500
21.	Beete	<i>Dalbargia latifolia</i>	20 - 40	500 - 1500
22.	Tare	<i>T. belerica</i>	20 - 40	500 - 2000
23.	Bamboo	<i>Bambusa arundinasia</i>	20 - 40	500 - 2500
24.	Bamboo	<i>Dendrocalamus strictus</i>	20 – 40	500 – 2500
25.	Muthuga	<i>Butea monosperma</i>	20 - 40	400 - 1500
26.	Hippe	<i>Madhuca latifolia</i>	20 - 40	500 - 2000
27.	Sandal	<i>Santalum album</i>	20 - 50	400 - 1000
28.	Nelli	<i>Emblca officinalis</i>	20 - 40	500 - 2000
29.	Nerale	<i>Sizyium cumini</i>	20 - 40	500 - 2000
30.	Dhaman	<i>Grevia tilifolia</i>	20 - 40	500 - 2000
31.	Kaval	<i>Careya arborea</i>	20 - 40	500 - 2000
32.	Harada	<i>Terminalia chebula</i>	20 - 40	500 - 2000

References

1. FAO (1976) Framework for Land Evaluation, Food and Agriculture Organization, Rome. 72 pp.
2. FAO (1983) Guidelines for Land Evaluation for Rainfed Agriculture, FAO, Rome, 237 pp.
3. IARI (1971) Soil Survey Manual, All India Soil and Land Use Survey Organization, IARI, New Delhi, 121 pp.
4. Katyal, J.C. and Rattan, R.K. (2003) Secondary and Micronutrients; Research Gap and Future Needs. Fert. News 48 (4); 9-20.
5. Naidu, L.G.K., Ramamurthy, V., Challa, O., Hegde, R. and Krishnan, P. (2006) Manual Soil Site Suitability Criteria for Major Crops, NBSS Publ. No. 129, NBSS & LUP, Nagpur, 118 pp.
6. Natarajan, A. and Dipak Sarkar (2010) Field Guide for Soil Survey, National Bureau of Soil Survey and Land Use Planning (ICAR), Nagpur, India.
7. Natarajan, A., Rajendra Hegde, Raj, J.N. and Shivananda Murthy, H.G. (2015) Implementation Manual for Sujala-III Project, Watershed Development Department, Bengaluru, Karnataka.
8. Sarma, V.A.K., Krishnan, P. and Budihal, S.L. (1987) Laboratory Manual, Tech. Bull. 23, NBSS & LUP, Nagpur.
9. Sehgal, J.L. (1990) Soil Resource Mapping of Different States of India; Why and How?, National Bureau of Soil Survey and Land Use Planning, Nagpur, 49 pp.
10. Shivaprasad, C.R., R.S. Reddy, J. Sehgal and M. Velayuthum (1998) Soils of Karnataka for Optimizing Land Use, NBSS Publ. No. 47b, NBSS & LUP, Nagpur, India.
11. Soil Survey Staff (2006) Keys to Soil Taxonomy, Tenth edition, U.S. Department of Agriculture/ NRCS, Washington DC, U.S.A.
12. Soil Survey Staff (2012) Soil Survey Manual, Handbook No. 18, USDA, Washington DC, USA.

Appendix I
Sudapur-2 (4J2a) Microwatershed
Soil Phase Information

Village	Survey NO	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Gajarakota	252	6.91	JNkMB2	LMU-3	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar+Redgram (Jw+Rg)	Not Available	Iles	Graded bunding
Gajarakota	254	5.99	BMNmA1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Jowar+Redgram (Jw+Rg)	Not Available	IIs	Strengthening of existing bunds
Gajarakota	255	0.75	JNkMB2	LMU-3	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar+Redgram (Jw+Rg)	Not Available	Iles	Graded bunding
Gajarakota	256	1.28	BMNmA1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Redgram (Rg)	Not Available	IIs	Strengthening of existing bunds
Gajarakota	257	0.05	BMNmA1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Redgram (Rg)	Not Available	IIs	Strengthening of existing bunds
Gajarakota	283	0	BMNmA1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Jowar (Jw)	Not Available	IIs	Strengthening of existing bunds
Gajarakota	289	1.04	BMNmA1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Cotton (Ct)	Not Available	IIs	Strengthening of existing bunds
Gajarakota	290	5.39	BMNmA1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Jowar+Redgram (Jw+Rg)	Not Available	IIs	Strengthening of existing bunds
Gajarakota	291	2.31	BMNmA1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Redgram (Rg)	Not Available	IIs	Strengthening of existing bunds
Gajarakota	292	6.94	BMNmB2g1	LMU-1	Very deep (>150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Gajarakota	293	4.62	BMNmB2g1	LMU-1	Very deep (>150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton+Redgram (Ct+Rg)	Not Available	Iles	Graded bunding
Gajarakota	294	5.63	BMNmB2g1	LMU-1	Very deep (>150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar+Redgram (Jw+Rg)	Not Available	Iles	Graded bunding
Gajarakota	295	4.71	BMNmA1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Redgram (Rg)	Not Available	IIs	Strengthening of existing bunds
Gajarakota	296	7.72	BMNmA1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Cotton+Redgram (Ct+Rg)	Not Available	IIs	Strengthening of existing bunds
Gajarakota	297	5.64	BMNmA1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Redgram (Rg)	Not Available	IIs	Strengthening of existing bunds
Gajarakota	298	5.16	BMNmA1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Jowar (Jw)	Not Available	IIs	Strengthening of existing bunds
Gajarakota	299	4.86	BMNmA1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Jowar (Jw)	Not Available	IIs	Strengthening of existing bunds
Gajarakota	300	5.58	BMNmB2g1	LMU-1	Very deep (>150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	Iles	Graded bunding
Gajarakota	301	4.14	BMNmA1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Redgram (Rg)	Not Available	IIs	Strengthening of existing bunds
Gajarakota	302	4.41	BMNmA1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Jowar+Redgram (Jw+Rg)	Not Available	IIs	Strengthening of existing bunds
Gajarakota	303	4.08	BMNmA1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Redgram (Rg)	Not Available	IIs	Strengthening of existing bunds
Gajarakota	304	7.18	BMNmA1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Cotton+Redgram (Ct+Rg)	Not Available	IIs	Strengthening of existing bunds

Village	Survey NO	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Gajarakota	305	6.85	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Jowar (Jw)	Not Available	Ro	Ro
Gajarakota	306	7.13	BMNmA1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Jowar+Redgram (Jw+Rg)	Not Available	IIs	Strengthening of existing bunds
Gajarakota	307	8.31	BMNmA1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Cotton+Jowar+Redgram (Ct+Jw+Rg)	Not Available	IIs	Strengthening of existing bunds
Gajarakota	308	9.22	BMNmA1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Jowar+Redgram (Jw+Rg)	Not Available	IIs	Strengthening of existing bunds
Gajarakota	309	2.91	BMNmA1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Cotton+Jowar+Redgram (Ct+Jw+Rg)	Not Available	IIs	Strengthening of existing bunds
Gajarakota	310	4.18	BMNmA1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Jowar+Redgram (Jw+Rg)	Not Available	IIs	Strengthening of existing bunds
Gajarakota	311	1.91	BMNmA1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Jowar+Redgram (Jw+Rg)	Not Available	IIs	Strengthening of existing bunds
Gajarakota	312	4.02	BMNmA1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Jowar+Redgram (Jw+Rg)	Not Available	IIs	Strengthening of existing bunds
Gajarakota	313	0	BMNmA1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Cotton+Redgram (Ct+Rg)	Not Available	IIs	Strengthening of existing bunds
Gajarakota	338	0	BMNmB2g1	LMU-1	Very deep (>150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar+Redgram (Jw+Rg)	Not Available	IIs	Graded bunding
Gajarakota	351	0.33	BMNmB2g1	LMU-1	Very deep (>150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	IIs	Graded bunding
Gajarakota	352	0.02	BMNmB2g1	LMU-1	Very deep (>150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIs	Graded bunding
Gajarakota	353	2.13	BMNmB2g1	LMU-1	Very deep (>150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Scrub land (Rg+Sl)	Not Available	IIs	Graded bunding
Gajarakota	354	5.66	BMNmB2g1	LMU-1	Very deep (>150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIs	Graded bunding
Gajarakota	355	4.12	BMNmB2g1	LMU-1	Very deep (>150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar+Redgram (Jw+Rg)	Not Available	IIs	Graded bunding
Gajarakota	356	2.91	BMNmB2g1	LMU-1	Very deep (>150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton+Redgram (Ct+Rg)	Not Available	IIs	Graded bunding
Gajarakota	359	4.53	BMNmB2g1	LMU-1	Very deep (>150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar+Redgram (Jw+Rg)	Not Available	IIs	Graded bunding
Gajarakota	360	5.24	BMNmB2g1	LMU-1	Very deep (>150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar+Redgram (Jw+Rg)	Not Available	IIs	Graded bunding
Gajarakota	361	4.81	BMNmB2g1	LMU-1	Very deep (>150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar+Redgram (Jw+Rg)	Not Available	IIs	Graded bunding
Gajarakota	362	0.94	BMNmB2g1	LMU-1	Very deep (>150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIs	Graded bunding
Gajarakota	363	8.08	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Redgram+Scrub land (Rg+Sl)	Not Available	Ro	Ro
Gajarakota	364	6.83	BMNmB2g1	LMU-1	Very deep (>150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar+Redgram (Jw+Rg)	Not Available	IIs	Graded bunding
Gajarakota	365	8.1	BMNmB2g1	LMU-1	Very deep (>150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar+Redgram (Jw+Rg)	Not Available	IIs	Graded bunding
Gajarakota	366	6.72	BMNmB2g1	LMU-1	Very deep (>150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton+Redgram (Ct+Rg)	Not Available	IIs	Graded bunding

Village	Survey NO	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Gajarakota	367	5.8	BMNmB2g1	LMU-1	Very deep (>150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton+Redgram (Ct+Rg)	Not Available	Iles	Graded bunding
Gajarakota	368	6.5	BMNmB2g1	LMU-1	Very deep (>150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton+Redgram (Ct+Rg)	Not Available	Iles	Graded bunding
Gajarakota	369	8.52	BMNmB2g1	LMU-1	Very deep (>150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Gajarakota	370	2.77	BMNmB2g1	LMU-1	Very deep (>150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Gajarakota	371	2.07	BMNmB2g1	LMU-1	Very deep (>150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar+Redgram (Jw+Rg)	Not Available	Iles	Graded bunding
Gajarakota	374	2.2	JNKmB2	LMU-3	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar+Redgram (Jw+Rg)	Not Available	Iles	Graded bunding
Gajarakota	375	1.02	BMNmB2g1	LMU-1	Very deep (>150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Gajarakota	376	5.58	BMNmA1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Jowar+Redgram (Jw+Rg)	Not Available	IIs	Strengthening of existing bunds
Gajarakota	377	5.71	BMNmA1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Redgram (Rg)	Not Available	IIs	Strengthening of existing bunds
Gajarakota	383	0.06	JNKmB2	LMU-3	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Mitathapad amapalli	188	0.28	BMNmB2g1	LMU-1	Very deep (>150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar+Redgram (Jw+Rg)	Not Available	Iles	Graded bunding
Mitathapad amapalli	189	3.53	BMNmA1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Redgram (Rg)	Not Available	IIs	Strengthening of existing bunds
Mitathapad amapalli	190	3.36	BMNmA1	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Not Available (NA)	Not Available	IIs	Strengthening of existing bunds
Mitathapad amapalli	191	4.26	NGPmB2g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	Iles	Graded bunding
Mitathapad amapalli	192/1	2.19	NGPmB2g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Mitathapad amapalli	192/2	1.7	NGPmB2g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iles	Graded bunding
Mitathapad amapalli	193	1.52	NGPmB2g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	Iles	Graded bunding
Mitathapad amapalli	194	3.97	NGPmB2g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Mitathapad amapalli	195	1.72	NGPmB2g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Mitathapad amapalli	196/1	0.1	NGPmB2g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iles	Graded bunding
Mitathapad amapalli	196/2	0.74	NGPmB2g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iles	Graded bunding
Mitathapad amapalli	196/3	1.46	NGPmB2g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	Iles	Graded bunding
Mitathapad amapalli	197	2.17	NGPmB2g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Bengalgram+Redgram (Bg+Rg)	Not Available	Iles	Graded bunding
Mitathapad amapalli	255	0.59	NGPmB2g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding

Village	Survey NO	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Mitathapad amapalli	256	4.08	NGPmB2g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Mitathapad amapalli	257	3.24	NGPmB2g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Mitathapad amapalli	258	0.95	NGPmB2g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Mitathapad amapalli	259	0.32	NGPmB2g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar+Redgram (Jw+Rg)	Not Available	Iles	Graded bunding
Mitathapad amapalli	260	4.99	NGPmB2g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar+Redgram (Jw+Rg)	Not Available	Iles	Graded bunding
Mitathapad amapalli	261	5.22	BLCcB2g1	LMU-2	Moderately deep (75-100 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	Iles	Trench cum bunding
Mitathapad amapalli	262/1	1.63	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Not Available (NA)	Not Available	Ro	Ro
Mitathapad amapalli	262/10	2.23	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Redgram (Rg)	Not Available	Ro	Ro
Mitathapad amapalli	262/11	2.55	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Scrub land (Sl)	Not Available	Ro	Ro
Mitathapad amapalli	262/12	2.22	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Scrub land (Sl)	Not Available	Ro	Ro
Mitathapad amapalli	262/13	2.33	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Scrub land (Sl)	Not Available	Ro	Ro
Mitathapad amapalli	262/14	2.27	NGPmB2g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Mitathapad amapalli	262/15	2.09	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Scrub land (Sl)	Not Available	Ro	Ro
Mitathapad amapalli	262/16	2.34	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Redgram (Rg)	Not Available	Ro	Ro
Mitathapad amapalli	262/17	2.74	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Scrub land (Sl)	Not Available	Ro	Ro
Mitathapad amapalli	262/18	0.79	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Not Available (NA)	Not Available	Ro	Ro
Mitathapad amapalli	262/19	2.05	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Redgram+Scrub land (Rg+Sl)	Not Available	Ro	Ro
Mitathapad amapalli	262/20	2.88	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Redgram (Rg)	Not Available	Ro	Ro
Mitathapad amapalli	262/20	3.24	NGPmB2g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Scrub land (Sl)	Not Available	Iles	Graded bunding
Mitathapad amapalli	262/21	0.27	NGPmB2g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iles	Graded bunding
Mitathapad amapalli	262/3	2.29	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Redgram (Rg)	Not Available	Ro	Ro
Mitathapad amapalli	262/4	2.67	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Jowar (Jw)	Not Available	Ro	Ro
Mitathapad amapalli	262/5	2.39	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Scrub land (Sl)	Not Available	Ro	Ro
Mitathapad amapalli	262/6	2.23	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Scrub land (Sl)	Not Available	Ro	Ro

Village	Survey NO	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Mitathapad amapalli	262/7	2.44	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Redgram (Rg)	Not Available	Ro	Ro
Mitathapad amapalli	262/8	2.64	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Redgram (Rg)	Not Available	Ro	Ro
Mitathapad amapalli	262/9	2.45	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Redgram (Rg)	Not Available	Ro	Ro
Mitathapad amapalli	263	3.96	BLCcB2g1	LMU-2	Moderately deep (75-100 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Trench cum bunding
Mitathapad amapalli	264	2.76	BLCcB2g1	LMU-2	Moderately deep (75-100 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	Iles	Trench cum bunding
Mitathapad amapalli	265	2.89	NGPmB2g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	Iles	Graded bunding
Mitathapad amapalli	269/2	2.11	NGPmB2g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	Iles	Graded bunding
Mitathapad amapalli	270	8.17	NGPmB2g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar+Redgram (Jw+Rg)	Not Available	Iles	Graded bunding
Siddhapura .B	14/1	1.53	BDLiB2	LMU-4	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram+Redgram (Gg+Rg)	Not Available	IIles	Graded bunding
Siddhapura .B	14/2	1.06	BDLiB2	LMU-4	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram (Gg)	Not Available	IIles	Graded bunding
Siddhapura .B	14/3	0.82	BDLiB2	LMU-4	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIles	Graded bunding
Siddhapura .B	15	2.75	BDLiB2	LMU-4	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIles	Graded bunding
Siddhapura .B	16	1.56	BDLiB2	LMU-4	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIles	Graded bunding
Siddhapura .B	17	0.73	BDLiB2	LMU-4	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIles	Graded bunding
Siddhapura .B	18/1	0.03	BDLiB2	LMU-4	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIles	Graded bunding
Siddhapura .B	37	0.81	VNKmB2g1	LMU-4	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIles	Trench cum bunding
Siddhapura .B	42	0.4	BLCcB2g1	LMU-2	Moderately deep (75-100 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Trench cum bunding
Siddhapura .B	44/1	0.42	BLCcB2g1	LMU-2	Moderately deep (75-100 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Trench cum bunding
Siddhapura .B	44/2	0.65	VNKmB2g1	LMU-4	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIles	Trench cum bunding
Siddhapura .B	45	4.51	VNKmB2g1	LMU-4	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIles	Trench cum bunding
Siddhapura .B	46	0.59	BLCcB2g1	LMU-2	Moderately deep (75-100 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iles	Trench cum bunding
Siddhapura .B	47	1.03	VNKmB2g1	LMU-4	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIles	Trench cum bunding
Siddhapura .B	48	3.31	VNKmB2g1	LMU-4	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram+Redgram (Gg+Rg)	Not Available	IIles	Trench cum bunding
Siddhapura .B	50	37.6	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Forest (Fo)	Not Available	Ro	Ro

Village	Survey NO	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Siddhapura .B	51	3.48	BLCcB2g1	LMU-2	Moderately deep (75-100 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram+Redgram (Gg+Rg)	Not Available	Iles	Trench cum bunding
Siddhapura .B	52	4.74	BLCcB2g1	LMU-2	Moderately deep (75-100 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram+Redgram (Gg+Rg)	Not Available	Iles	Trench cum bunding
Siddhapura .B	53/1	2.86	BLCcB2g1	LMU-2	Moderately deep (75-100 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Trench cum bunding
Siddhapura .B	53/2	0.93	BLCcB2g1	LMU-2	Moderately deep (75-100 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Trench cum bunding
Siddhapura .B	54	4.42	BLCcB2g1	LMU-2	Moderately deep (75-100 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram+Redgram+Scrub land (Gg+Rg+Sl)	Not Available	Iles	Trench cum bunding
Siddhapura .B	55	6.29	BLCcB2g1	LMU-2	Moderately deep (75-100 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Trench cum bunding
Siddhapura .B	56	47.48	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Forest (Fo)	Not Available	Ro	Ro
Siddhapura .B	57	3.37	BLCcB2g1	LMU-2	Moderately deep (75-100 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Trench cum bunding
Siddhapura .B	58	0.79	BLCcB2g1	LMU-2	Moderately deep (75-100 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iles	Trench cum bunding
Siddhapura .B	59	6.2	BLCcB2g1	LMU-2	Moderately deep (75-100 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram+Paddy+Redgram (Gg+Pd+Rg)	1 Bore well	Iles	Trench cum bunding
Siddhapura .B	60	4.96	BLCcB2g1	LMU-2	Moderately deep (75-100 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram+Redgram (Gg+Rg)	Not Available	Iles	Trench cum bunding
Siddhapura .B	61	3.53	BLCcB2g1	LMU-2	Moderately deep (75-100 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Trench cum bunding
Siddhapura .B	62	3.63	VNKmB2g1	LMU-4	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram+Redgram (Gg+Rg)	Not Available	IIles	Trench cum bunding
Siddhapura .B	63	2.31	VNKmB2g1	LMU-4	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIles	Trench cum bunding
Siddhapura .B	64	5.08	VNKmB2g1	LMU-4	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram+Redgram (Gg+Rg)	Not Available	IIles	Trench cum bunding
Siddhapura .B	65	1.06	VNKmB2g1	LMU-4	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram+Redgram (Gg+Rg)	Not Available	IIles	Trench cum bunding
Siddhapura .B	66	1.06	VNKmB2g1	LMU-4	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIles	Trench cum bunding
Siddhapura .B	67	3.05	VNKmB2g1	LMU-4	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram+Redgram (Gg+Rg)	Not Available	IIles	Trench cum bunding
Siddhapura .B	68	0.5	VNKmB2g1	LMU-4	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIles	Trench cum bunding
Siddhapura .B	69	0.43	BDLiB2	LMU-4	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IIles	Graded bunding
Siddhapura .B	70	0.59	BDLiB2	LMU-4	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IIles	Graded bunding
Siddhapura .B	71	0.17	BDLiB2	LMU-4	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IIles	Graded bunding
Siddhapura .B	72	0.51	BDLiB2	LMU-4	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IIles	Graded bunding

Village	Survey NO	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Siddhapura .B	73	0.45	BDLiB2	LMU-4	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IIes	Graded bunding
Siddhapura .B	74/1	1	VNKmB2g1	LMU-4	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Trench cum bunding
Siddhapura .B	74/2	1.42	VNKmB2g1	LMU-4	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IIes	Trench cum bunding
Siddhapura .B	74/3	0.9	VNKmB2g1	LMU-4	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IIes	Trench cum bunding
Siddhapura .B	75	0.2	VNKmB2g1	LMU-4	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Trench cum bunding
Siddhapura .B	76	4.15	VNKmB2g1	LMU-4	Shallow (25-50 cm)	Clay	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Trench cum bunding
Siddhapura .B	77	0.58	BDLiB2	LMU-4	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIes	Graded bunding
Siddhapura .B	78	0.44	BDLiB2	LMU-4	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Fallow land (Fl)	Not Available	IIes	Graded bunding
Siddhapura .B	79	2.98	BDLiB2	LMU-4	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram+Redgram (Gg+Rg)	Not Available	IIes	Graded bunding
Siddhapura .B	80	0.82	BDLiB2	LMU-4	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Siddhapura .B	81	0.76	BDLiB2	LMU-4	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Siddhapura .B	82	1.16	BDLiB2	LMU-4	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IIes	Graded bunding
Siddhapura .B	83	2.72	BDLiB2	LMU-4	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Siddhapura .B	84	0.11	BDLiB2	LMU-4	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIes	Graded bunding
Siddhapura .B	85	0.38	BDLiB2	LMU-4	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IIes	Graded bunding
Siddhapura .B	86	0.89	BDLiB2	LMU-4	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IIes	Graded bunding
Siddhapura .B	87	0.42	BDLiB2	LMU-4	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IIes	Graded bunding
Siddhapura .B	88	0.98	BDLiB2	LMU-4	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IIes	Graded bunding
Siddhapura .B	89	0.33	BDLiB2	LMU-4	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IIes	Graded bunding
Siddhapura .B	90	0.75	BLCcB2g1	LMU-2	Moderately deep (75-100 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iies	Trench cum bunding
Siddhapura .B	91	0.71	BLCcB2g1	LMU-2	Moderately deep (75-100 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iies	Trench cum bunding
Siddhapura .B	92	1.19	BLCcB2g1	LMU-2	Moderately deep (75-100 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iies	Trench cum bunding
Siddhapura .B	93	2.92	Waterbody	Others	Others	Others	Others	Others	Others	Others	Waterbody	Not Available	Others	Others
Siddhapura .B	94/1	2.32	BLCcB2g1	LMU-2	Moderately deep (75-100 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram (Gg)	Not Available	Iies	Trench cum bunding

Village	Survey NO	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Siddhapura .B	94/2	0.25	BLCcB2g1	LMU-2	Moderately deep (75-100 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iles	Trench cum bunding
Siddhapura .B	94/3	0.34	BLCcB2g1	LMU-2	Moderately deep (75-100 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Trench cum bunding
Siddhapura .B	95	0.59	BLCcB2g1	LMU-2	Moderately deep (75-100 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Trench cum bunding
Siddhapura .B	96	2.13	BLCcB2g1	LMU-2	Moderately deep (75-100 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram (Gg)	1 Bore well	Iles	Trench cum bunding
Siddhapura .B	97	2.57	BLCiB2	LMU-2	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Trench cum bunding
Siddhapura .B	98	0.85	BLCiB2	LMU-2	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Trench cum bunding
Siddhapura .B	99	2.87	BLCiB2	LMU-2	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Trench cum bunding
Siddhapura .B	100	2.32	BLCiB2	LMU-2	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Trench cum bunding
Siddhapura .B	101	6.29	BLCiB2	LMU-2	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram+Redgram (Gg+Rg)	Not Available	Iles	Trench cum bunding
Siddhapura .B	102	4.57	BLCiB2	LMU-2	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Bengalgram+Redgram (Bg+Rg)	Not Available	Iles	Trench cum bunding
Siddhapura .B	103	2.09	BLCiB2	LMU-2	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Trench cum bunding
Siddhapura .B	104	3.45	BLCiB2	LMU-2	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Bengalgram+Redgram (Bg+Rg)	Not Available	Iles	Trench cum bunding
Siddhapura .B	105	4.63	BLCiB2	LMU-2	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Bengalgram+Redgram (Bg+Rg)	Not Available	Iles	Trench cum bunding
Siddhapura .B	106	0.5	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Redgram (Rg)	Not Available	Ro	Ro
Siddhapura .B	107/1	1.65	BDLiB2	LMU-4	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram+Redgram (Gg+Rg)	Not Available	IIles	Graded bunding
Siddhapura .B	107/2	1.46	BDLiB2	LMU-4	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIles	Graded bunding
Siddhapura .B	108/3	1.11	BDLiB2	LMU-4	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram+Redgram (Gg+Rg)	Not Available	IIles	Graded bunding
Siddhapura .B	109	0.03	BDLiB2	LMU-4	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram+Redgram (Gg+Rg)	Not Available	IIles	Graded bunding
Siddhapura .B	127	1.66	BLCiB2	LMU-2	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Trench cum bunding
Yadhalapura	26	1.56	NGPmB2g1	LMU-1	Deep (100-150 cm)	Clay	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	Iles	Graded bunding

Ro- Rock outcrops

Village	Survey No	Soil Reaction	Salinity	Organic Carbon	Available Phosphorus	Available Potassium	Available Sulphur	Available Boron	Available Iron	Available Manganese	Available Copper	Available Zinc
Mitathapada mapalli	257	Slightly acid (pH 6.0 - 6.5)	Non saline (<2 dsm)	High (> 0.75 %)	Low (< 23 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Mitathapada mapalli	258	Slightly acid (pH 6.0 - 6.5)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Mitathapada mapalli	259	Slightly acid (pH 6.0 - 6.5)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	Medium (10 - 20 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)
Mitathapada mapalli	260	Slightly acid (pH 6.0 - 6.5)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Mitathapada mapalli	261	Slightly acid (pH 6.0 - 6.5)	Non saline (<2 dsm)	High (> 0.75 %)	Low (< 23 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)
Mitathapada mapalli	262/1	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro
Mitathapada mapalli	262/10	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro
Mitathapada mapalli	262/11	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro
Mitathapada mapalli	262/12	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro
Mitathapada mapalli	262/13	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro
Mitathapada mapalli	262/14	Slightly acid (pH 6.0 - 6.5)	Non saline (<2 dsm)	High (> 0.75 %)	Low (< 23 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Mitathapada mapalli	262/15	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro
Mitathapada mapalli	262/16	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro
Mitathapada mapalli	262/17	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro
Mitathapada mapalli	262/18	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro
Mitathapada mapalli	262/19	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro
Mitathapada mapalli	262/2	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro
Mitathapada mapalli	262/20	Slightly acid (pH 6.0 - 6.5)	Non saline (<2 dsm)	High (> 0.75 %)	Low (< 23 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Mitathapada mapalli	262/21	Slightly acid (pH 6.0 - 6.5)	Non saline (<2 dsm)	High (> 0.75 %)	Low (< 23 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Mitathapada mapalli	262/3	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro
Mitathapada mapalli	262/4	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro
Mitathapada mapalli	262/5	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro
Mitathapada mapalli	262/6	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro
Mitathapada mapalli	262/7	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro

Village	Survey No	Soil Reaction	Salinity	Organic Carbon	Available Phosphorus	Available Potassium	Available Sulphur	Available Boron	Available Iron	Available Manganese	Available Copper	Available Zinc
Siddhapura .B	96	Slightly acid (pH 6.0 – 6.5)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 – 57 kg/ha)	Medium (145 – 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)
Siddhapura .B	97	Slightly acid (pH 6.0 – 6.5)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 – 57 kg/ha)	Medium (145 – 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)
Siddhapura .B	98	Slightly acid (pH 6.0 – 6.5)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 – 57 kg/ha)	Medium (145 – 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)
Siddhapura .B	99	Slightly acid (pH 6.0 – 6.5)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 – 57 kg/ha)	Medium (145 – 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)
Siddhapura .B	100	Slightly acid (pH 6.0 – 6.5)	Non saline (<2 dsm)	High (> 0.75 %)	Low (< 23 kg/ha)	Medium (145 – 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)
Siddhapura .B	101	Slightly acid (pH 6.0 – 6.5)	Non saline (<2 dsm)	High (> 0.75 %)	Low (< 23 kg/ha)	Medium (145 – 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)
Siddhapura .B	102	Neutral (pH 6.5 – 7.3)	Non saline (<2 dsm)	High (> 0.75 %)	Low (< 23 kg/ha)	Medium (145 – 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)
Siddhapura .B	103	Slightly acid (pH 6.0 – 6.5)	Non saline (<2 dsm)	High (> 0.75 %)	Low (< 23 kg/ha)	Medium (145 – 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)
Siddhapura .B	104	Slightly acid (pH 6.0 – 6.5)	Non saline (<2 dsm)	High (> 0.75 %)	Low (< 23 kg/ha)	Medium (145 – 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)
Siddhapura .B	105	Neutral (pH 6.5 – 7.3)	Non saline (<2 dsm)	High (> 0.75 %)	Low (< 23 kg/ha)	Medium (145 – 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)
Siddhapura .B	106	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro
Siddhapura .B	107/1	Slightly acid (pH 6.0 – 6.5)	Non saline (<2 dsm)	High (> 0.75 %)	High (> 57 kg/ha)	Medium (145 – 337 kg/ha)	Medium (10 – 20 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)
Siddhapura .B	107/2	Slightly acid (pH 6.0 – 6.5)	Non saline (<2 dsm)	High (> 0.75 %)	High (> 57 kg/ha)	Medium (145 – 337 kg/ha)	Medium (10 – 20 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)
Siddhapura .B	108/3	Slightly acid (pH 6.0 – 6.5)	Non saline (<2 dsm)	High (> 0.75 %)	High (> 57 kg/ha)	Medium (145 – 337 kg/ha)	Medium (10 – 20 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)
Siddhapura .B	109	Slightly acid (pH 6.0 – 6.5)	Non saline (<2 dsm)	High (> 0.75 %)	High (> 57 kg/ha)	Medium (145 – 337 kg/ha)	Medium (10 – 20 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)
Siddhapura .B	127	Slightly acid (pH 6.0 – 6.5)	Non saline (<2 dsm)	High (> 0.75 %)	Low (< 23 kg/ha)	Medium (145 – 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)
Yadhalapura	26	Slightly acid (pH 6.0 – 6.5)	Non saline (<2 dsm)	High (> 0.75 %)	Low (< 23 kg/ha)	Medium (145 – 337 kg/ha)	Low (<10 ppm)	Medium (0.5 – 1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)

Appendix III
Sudapur-2 (4J2a) Microwatershed
Soil Suitability Information

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry
Gajarakota	252	N1r	S2r	S3r	S2rt	S3r	S3t	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	N1n	S3r	S3r	S2r	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r
Gajarakota	254	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz
Gajarakota	255	N1r	S2r	S3r	S2rt	S3r	S3t	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	N1n	S3r	S3r	S2r	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r
Gajarakota	256	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz
Gajarakota	257	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz
Gajarakota	283	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz
Gajarakota	289	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz
Gajarakota	290	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz
Gajarakota	291	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz
Gajarakota	292	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz
Gajarakota	293	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz
Gajarakota	294	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz
Gajarakota	295	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz
Gajarakota	296	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz
Gajarakota	297	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz
Gajarakota	298	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz
Gajarakota	299	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz
Gajarakota	300	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz
Gajarakota	301	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz
Gajarakota	302	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz
Gajarakota	303	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz
Gajarakota	304	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz
Gajarakota	305	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro
Gajarakota	306	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry	
Gajarakota	307	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz	
Gajarakota	308	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz	
Gajarakota	309	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz	
Gajarakota	310	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz	
Gajarakota	311	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz	
Gajarakota	312	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz	
Gajarakota	313	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz	
Gajarakota	338	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz	
Gajarakota	351	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz	
Gajarakota	352	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz	
Gajarakota	353	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz	
Gajarakota	354	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz	
Gajarakota	355	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz	
Gajarakota	356	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz	
Gajarakota	359	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz	
Gajarakota	360	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz	
Gajarakota	361	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz	
Gajarakota	362	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz	
Gajarakota	363	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro
Gajarakota	364	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz	
Gajarakota	365	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz	
Gajarakota	366	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz	
Gajarakota	367	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz	
Gajarakota	368	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz	
Gajarakota	369	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz	
Gajarakota	370	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz	

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry
Gajarakota	371	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz
Gajarakota	374	N1r	S2r	S3r	S2rt	S3r	S3t	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	N1n	S3r	S3r	S2r	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r
Gajarakota	375	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz
Gajarakota	376	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz
Gajarakota	377	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz
Gajarakota	383	N1r	S2r	S3r	S2rt	S3r	S3t	N1r	S3r	S3t	S3r	S3r	S2r	S3r	S2r	N1n	S3r	S3r	S2r	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r
Mitathapadam apalli	188	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz
Mitathapadam apalli	189	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz
Mitathapadam apalli	190	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3tz	S3t	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz
Mitathapadam apalli	191	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3t	S3tz	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz
Mitathapadam apalli	192	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3t	S3tz	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz
Mitathapadam apalli	192 /1	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3t	S3tz	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz
Mitathapadam apalli	192 /2	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3t	S3tz	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz
Mitathapadam apalli	193	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3t	S3tz	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz
Mitathapadam apalli	194	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3t	S3tz	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz
Mitathapadam apalli	195	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3t	S3tz	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz
Mitathapadam apalli	196 /1	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3t	S3tz	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz
Mitathapadam apalli	196 /2	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3t	S3tz	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz
Mitathapadam apalli	196 /3	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3t	S3tz	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz
Mitathapadam apalli	197	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3t	S3tz	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz
Mitathapadam apalli	255	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3t	S3tz	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz
Mitathapadam apalli	256	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3t	S3tz	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz
Mitathapadam apalli	257	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3t	S3tz	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz
Mitathapadam apalli	258	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3t	S3tz	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry	
apalli																															
Mitathapadam apalli	259	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3t	S3tz	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz	
Mitathapadam apalli	260	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3t	S3tz	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz	
Mitathapadam apalli	261	S3r	S2g	S2r	S2gt	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S2rn	S3r	S2r	S1	S2g	S2g	S2g	S2g	S2g	S2r	S1	S2g	S2g	S2r	S2r	
Mitathapadam apalli	262 /1	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	
Mitathapadam apalli	262 /10	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	
Mitathapadam apalli	262 /11	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	
Mitathapadam apalli	262 /12	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	
Mitathapadam apalli	262 /13	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	
Mitathapadam apalli	262 /14	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3t	S3tz	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz	
Mitathapadam apalli	262 /15	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	
Mitathapadam apalli	262 /16	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	
Mitathapadam apalli	262 /17	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	
Mitathapadam apalli	262 /18	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	
Mitathapadam apalli	262 /19	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	
Mitathapadam apalli	262 /2	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	
Mitathapadam apalli	262 /20	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3t	S3tz	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz	
Mitathapadam apalli	262 /21	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3t	S3tz	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz	
Mitathapadam apalli	262 /3	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	
Mitathapadam apalli	262 /4	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	
Mitathapadam apalli	262 /5	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	
Mitathapadam	262	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry	
apalli	/6																														
Mitathapadam apalli	262/7	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro
Mitathapadam apalli	262/8	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro
Mitathapadam apalli	262/9	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro
Mitathapadam apalli	263	S3r	S2g	S2r	S2gt	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S2rn	S3r	S2r	S1	S2g	S2g	S2g	S2g	S2g	S2r	S1	S2g	S2g	S2r	S2r	S2r
Mitathapadam apalli	264	S3r	S2g	S2r	S2gt	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S2rn	S3r	S2r	S1	S2g	S2g	S2g	S2g	S2g	S2r	S1	S2g	S2g	S2r	S2r	S2r
Mitathapadam apalli	265	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3t	S3tz	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz	S3tz
Mitathapadam apalli	269/2	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3t	S3tz	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz	S3tz
Mitathapadam apalli	270	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3t	S3tz	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz	S3tz
Siddhapura .B	14/1	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	S3r	N1r	N1r
Siddhapura .B	14/2	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	S3r	N1r	N1r
Siddhapura .B	14/3	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	S3r	N1r	N1r
Siddhapura .B	15	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	S3r	N1r	N1r
Siddhapura .B	16	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	S3r	N1r	N1r
Siddhapura .B	17	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	S3r	N1r	N1r
Siddhapura .B	18/1	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	S3r	N1r	N1r
Siddhapura .B	37	N1r	S3r	N1r	S3r	N1r	S3r	N1r	N1r	S3rt	N1r	N1r	S3r	N1r	S3r	N1r	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	S3r	N1r	N1r
Siddhapura .B	42	S3r	S2g	S2r	S2gt	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S2rn	S3r	S2r	S1	S2g	S2g	S2g	S2g	S2g	S2r	S1	S2g	S2g	S2r	S2r	S2r
Siddhapura .B	44/1	S3r	S2g	S2r	S2gt	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S2rn	S3r	S2r	S1	S2g	S2g	S2g	S2g	S2g	S2r	S1	S2g	S2g	S2r	S2r	S2r
Siddhapura .B	44/2	N1r	S3r	N1r	S3r	N1r	S3r	N1r	N1r	S3rt	N1r	N1r	S3r	N1r	S3r	N1r	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	S3r	N1r	N1r
Siddhapura .B	45	N1r	S3r	N1r	S3r	N1r	S3r	N1r	N1r	S3rt	N1r	N1r	S3r	N1r	S3r	N1r	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	S3r	N1r	N1r
Siddhapura .B	46	S3r	S2g	S2r	S2gt	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S2rn	S3r	S2r	S1	S2g	S2g	S2g	S2g	S2g	S2r	S1	S2g	S2g	S2r	S2r	S2r
Siddhapura .B	47	N1r	S3r	N1r	S3r	N1r	S3r	N1r	N1r	S3rt	N1r	N1r	S3r	N1r	S3r	N1r	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	S3r	N1r	N1r

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry	
Siddhapura .B	48	N1r	S3r	N1r	S3r	N1r	S3r	N1r	N1r	S3rt	N1r	N1r	S3r	N1r	S3r	N1r	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Siddhapura .B	50	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro
Siddhapura .B	51	S3r	S2g	S2r	S2gt	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S2rn	S3r	S2r	S1	S2g	S2g	S2g	S2g	S2g	S2r	S1	S2g	S2g	S2r	S2r	
Siddhapura .B	52	S3r	S2g	S2r	S2gt	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S2rn	S3r	S2r	S1	S2g	S2g	S2g	S2g	S2g	S2r	S1	S2g	S2g	S2r	S2r	
Siddhapura .B	53/1	S3r	S2g	S2r	S2gt	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S2rn	S3r	S2r	S1	S2g	S2g	S2g	S2g	S2g	S2r	S1	S2g	S2g	S2r	S2r	
Siddhapura .B	53/2	S3r	S2g	S2r	S2gt	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S2rn	S3r	S2r	S1	S2g	S2g	S2g	S2g	S2g	S2r	S1	S2g	S2g	S2r	S2r	
Siddhapura .B	54	S3r	S2g	S2r	S2gt	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S2rn	S3r	S2r	S1	S2g	S2g	S2g	S2g	S2g	S2r	S1	S2g	S2g	S2r	S2r	
Siddhapura .B	55	S3r	S2g	S2r	S2gt	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S2rn	S3r	S2r	S1	S2g	S2g	S2g	S2g	S2g	S2r	S1	S2g	S2g	S2r	S2r	
Siddhapura .B	56	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro
Siddhapura .B	57	S3r	S2g	S2r	S2gt	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S2rn	S3r	S2r	S1	S2g	S2g	S2g	S2g	S2g	S2r	S1	S2g	S2g	S2r	S2r	
Siddhapura .B	58	S3r	S2g	S2r	S2gt	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S2rn	S3r	S2r	S1	S2g	S2g	S2g	S2g	S2g	S2r	S1	S2g	S2g	S2r	S2r	
Siddhapura .B	59	S3r	S2g	S2r	S2gt	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S2rn	S3r	S2r	S1	S2g	S2g	S2g	S2g	S2g	S2r	S1	S2g	S2g	S2r	S2r	
Siddhapura .B	60	S3r	S2g	S2r	S2gt	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S2rn	S3r	S2r	S1	S2g	S2g	S2g	S2g	S2g	S2r	S1	S2g	S2g	S2r	S2r	
Siddhapura .B	61	S3r	S2g	S2r	S2gt	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S2rn	S3r	S2r	S1	S2g	S2g	S2g	S2g	S2g	S2r	S1	S2g	S2g	S2r	S2r	
Siddhapura .B	62	N1r	S3r	N1r	S3r	N1r	S3r	N1r	N1r	S3rt	N1r	N1r	S3r	N1r	S3r	N1r	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Siddhapura .B	63	N1r	S3r	N1r	S3r	N1r	S3r	N1r	N1r	S3rt	N1r	N1r	S3r	N1r	S3r	N1r	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Siddhapura .B	64	N1r	S3r	N1r	S3r	N1r	S3r	N1r	N1r	S3rt	N1r	N1r	S3r	N1r	S3r	N1r	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Siddhapura .B	65	N1r	S3r	N1r	S3r	N1r	S3r	N1r	N1r	S3rt	N1r	N1r	S3r	N1r	S3r	N1r	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Siddhapura .B	66	N1r	S3r	N1r	S3r	N1r	S3r	N1r	N1r	S3rt	N1r	N1r	S3r	N1r	S3r	N1r	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Siddhapura .B	67	N1r	S3r	N1r	S3r	N1r	S3r	N1r	N1r	S3rt	N1r	N1r	S3r	N1r	S3r	N1r	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Siddhapura .B	68	N1r	S3r	N1r	S3r	N1r	S3r	N1r	N1r	S3rt	N1r	N1r	S3r	N1r	S3r	N1r	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Siddhapura .B	69	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Siddhapura .B	70	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Siddhapura .B	71	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Siddhapura .B	72	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Siddhapura .B	73	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry	
Siddhapura .B	74/1	N1r	S3r	N1r	S3r	N1r	S3r	N1r	N1r	S3rt	N1r	N1r	S3r	N1r	S3r	N1r	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Siddhapura .B	74/2	N1r	S3r	N1r	S3r	N1r	S3r	N1r	N1r	S3rt	N1r	N1r	S3r	N1r	S3r	N1r	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Siddhapura .B	74/3	N1r	S3r	N1r	S3r	N1r	S3r	N1r	N1r	S3rt	N1r	N1r	S3r	N1r	S3r	N1r	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Siddhapura .B	75	N1r	S3r	N1r	S3r	N1r	S3r	N1r	N1r	S3rt	N1r	N1r	S3r	N1r	S3r	N1r	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Siddhapura .B	76	N1r	S3r	N1r	S3r	N1r	S3r	N1r	N1r	S3rt	N1r	N1r	S3r	N1r	S3r	N1r	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Siddhapura .B	77	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Siddhapura .B	78	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Siddhapura .B	79	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Siddhapura .B	80	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Siddhapura .B	81	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Siddhapura .B	82	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Siddhapura .B	83	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Siddhapura .B	84	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Siddhapura .B	85	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Siddhapura .B	86	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Siddhapura .B	87	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Siddhapura .B	88	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Siddhapura .B	89	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Siddhapura .B	90	S3r	S2g	S2r	S2gt	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S2rn	S3r	S2r	S1	S2g	S2g	S2g	S2g	S2g	S2r	S1	S2g	S2g	S2r	S2r	
Siddhapura .B	91	S3r	S2g	S2r	S2gt	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S2rn	S3r	S2r	S1	S2g	S2g	S2g	S2g	S2g	S2r	S1	S2g	S2g	S2r	S2r	
Siddhapura .B	92	S3r	S2g	S2r	S2gt	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S2rn	S3r	S2r	S1	S2g	S2g	S2g	S2g	S2g	S2r	S1	S2g	S2g	S2r	S2r	
Siddhapura .B	93	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs
Siddhapura .B	94/1	S3r	S2g	S2r	S2gt	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S2rn	S3r	S2r	S1	S2g	S2g	S2g	S2g	S2g	S2r	S1	S2g	S2g	S2r	S2r	
Siddhapura .B	94/2	S3r	S2g	S2r	S2gt	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S2rn	S3r	S2r	S1	S2g	S2g	S2g	S2g	S2g	S2r	S1	S2g	S2g	S2r	S2r	
Siddhapura .B	94/3	S3r	S2g	S2r	S2gt	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S2rn	S3r	S2r	S1	S2g	S2g	S2g	S2g	S2g	S2r	S1	S2g	S2g	S2r	S2r	

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry	
Siddhapura .B	95	S3r	S2g	S2r	S2gt	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S2rn	S3r	S2r	S1	S2g	S2g	S2g	S2g	S2g	S2r	S1	S2g	S2g	S2r	S2r	
Siddhapura .B	96	S3r	S2g	S2r	S2gt	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S2rn	S3r	S2r	S1	S2g	S2g	S2g	S2g	S2g	S2r	S1	S2g	S2g	S2r	S2r	
Siddhapura .B	97	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S2rn	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r	
Siddhapura .B	98	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S2rn	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r	
Siddhapura .B	99	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S2rn	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r	
Siddhapura .B	100	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S2rn	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r	
Siddhapura .B	101	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S2rn	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r	
Siddhapura .B	102	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S2rn	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r	
Siddhapura .B	103	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S2rn	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r	
Siddhapura .B	104	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S2rn	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r	
Siddhapura .B	105	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S2rn	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r	
Siddhapura .B	106	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro
Siddhapura .B	107 /1	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Siddhapura .B	107 /2	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Siddhapura .B	108 /3	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Siddhapura .B	109	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1n	N1r	N1r	S3r	S3r	S3r	S3r	S3r	S3r	N1r	S3r	S3r	S3r	N1r	N1r	
Siddhapura .B	127	S3r	S1	S2r	S2t	S2r	S3t	S3r	S2r	S3t	S2rt	S2rt	S1	S2r	S1	S2rn	S3r	S2r	S1	S1	S1	S1	S1	S1	S2r	S1	S1	S1	S2r	S2r	
Yadhalapura	26	S3t	S2tz	S3t	S2z	S3tz	S2z	S3z	S2z	S2z	S2z	S2tz	S3z	S3tz	S2z	N1t	S3z	S2z	S3t	S3tz	S2tz	S3t	S2tz	S2tz	S2tz	S2tz	S3t	S2tz	S3z	S3tz	

PART-B

SOCIO-ECONOMIC STATUS OF FARM HOUSEHOLDS

CONTENTS

1.	Salient findings of the survey	1-4
2.	Introduction	5
3.	Methodology	7
4.	Salient features of the survey	9-25
5.	Summary	27-31

LIST OF TABLES

1	Households sampled for socio economic survey	9
2	Population characteristics	9
3	Age wise classification of household members	9
4	Education level of household members	10
5	Occupation of household heads	10
6	Occupation of family members	10
7	Institutional participation of household members	11
8	Type of house owned by households	11
9	Durable assets owned by households	11
10	Average value of durable assets owned by households	12
11	Farm implements owned by households	12
12	Average value of farm implements	12
13	Livestock possession by households	13
14	Average labour availability	13
15	Adequacy of hired labour	14
16	Distribution of land (ha)	14
17	Average land value (Rs./ha)	14
18	Source of irrigation	15
19	Irrigated area (ha)	15
20	Cropping pattern	15
21	Cropping intensity	15
22	Cost of cultivation of Cotton	16
23	Cost of cultivation of Green gram	17
24	Cost of cultivation of red gram	18
25	Cost of cultivation of paddy	19
26	Adequacy of fodder	20
27	Annual gross income	20
28	Average annual expenditure	20
29	Horticulture species grown	21
30	Forest species grown	21
31	Average additional investment capacity	21

32	Source of additional investment	22
33	Marketing of the agricultural produce	22
34	Marketing channels used for sale of agricultural produce	22
35	Mode of transport of agricultural produce	23
36	Incidence of soil and water erosion problems	23
37	Interest shown towards soil testing	23
38	Usage pattern of fuel for domestic use	23
39	Source of drinking water	23
40	Source of light	24
41	Existence of sanitary toilet facility	24
42	Possession of public distribution system (PDS) card	24
43	Participation in NREGA programme	24
44	Adequacy of food items	25
45	Inadequacy of food items	25
46	Farming constraints experienced	25

SALIENT FINDINGS OF THE SURVEY

- ❖ *The data on households sampled for socio economic survey indicated that 35 farmers were sampled in Sudapur-2 micro-watershed among them 1 (2.86 %) were landless, 19 (54.29 %) were marginal farmers, 7 (20 %) were small farmers and semi medium farmers and 1 (2.86 %) were medium farmers.*
- ❖ *The data indicated that there were 95 (51.91 %) men and 88 (48.09 %) women among the sampled households. The average family size of landless farmers' was 5, marginal farmers' was 5.57, small farmers' was 4.14, semi medium farmers' was 5.57 and medium farmers' was 4.*
- ❖ *The data indicated that, 39 (21.31 %) people were in 0-15 years of age, 83 (45.36 %) were in 16-35 years of age, 40 (21.86 %) were in 36-60 years of age and 21 (11.48 %) were above 61 years of age.*
- ❖ *The results indicated that Sudapur-2 had 55.19 per cent illiterates, 1.09 per cent Functional Literate, 14.75 per cent of them had primary school, 1.64 per cent of them had middle school, 12.02 per cent of them had high school education, 4.37 per cent of them had PUC, 1.24 per cent of them had Diploma, 0.62 per cent of them had ITI and 5.46 per cent of them had Degree education.*
- ❖ *The results indicate that, 68.57 per cent of household heads were practicing agriculture, 25.71 per cent of the household heads were agricultural labourers, 5.71 per cent of the household heads were in Private Service and 2.86 per cent of the household heads were Housewives.*
- ❖ *The results indicate that agriculture was the major occupation for 46.99 per cent of the household members, 10.93 per cent were agricultural labourers, 0.55 per cent were General Labour, 1.09 per cent were Government Service, 3.28 per cent were Private Service, 0.55 per cent were Trade & Business, 19.67 per cent were Student, 9.84 per cent were Housewife and 7.10 per cent were children.*
- ❖ *The results show that, 100 per cent of the population in the micro watershed has not participated in any institutions.*
- ❖ *The results indicate that 11.43 per cent of the households possess thatched house, 77.14 per cent of the households possess katcha house and 11.43 per cent of them possess pucca/RCC house.*
- ❖ *The results show that 88.57 per cent of the households possess TV, 20 per cent of the households possess mixer/grinder, 5.71 per cent of the households possess Refrigerator, 14.29 per cent of the households possess Bicycle, 40 per cent of the households possess Motor Cycle and 100 per cent of the households possess mobile phones.*
- ❖ *The results show that the average value of television was Rs. 9,000, mixer/grinder was Rs. 2,000, Refrigerator was Rs. 8,500, Bicycle was Rs. 2,000, motor cycle was Rs. 55,500 and mobile phone was Rs. 2,630.*
- ❖ *About 14.29 per cent of the households possess bullock cart, 40 per cent of them possess plough, 22.86 per cent of them possess seed/fertilizer drill, 2.86 per cent of*

- them possess Harvester, 17.14 per cent of them possess Sprayer and 34.29 per cent of them possess Weeder.
- ❖ The results show that the average value of bullock cart was Rs. 28,000, plough was Rs. 1,500, seed/fertilizer drill was Rs. 1,687, Harvester was Rs. 100, sprayer was Rs. 3,750 and weeder was Rs. 30.
 - ❖ The results indicate that, 42.86 per cent of the households possess bullocks, 20 per cent of the households possess local cow and 5.71 per cent of the households possess Buffalo, Goat and Poultry birds.
 - ❖ The results indicate that, average own labour men available in the micro watershed was 1.62, average own labour (women) available was 1.68, average hired labour (men) available was 11.32 and average hired labour (women) available was 8.09.
 - ❖ The results indicate that, 94.29 per cent of the households opined that the hired labour was adequate.
 - ❖ The results indicate that, households of the Sudapur-2 micro-watershed possess 32.73 ha (82.87 %) of dry land, 5.59 ha (14.16 %) of irrigated land and 1.17 ha (2.97 %) of Permanent Fallow land. Marginal farmers possess 11.80 ha (96.68 %) of dry land and 0.40 ha (3.32 %) of irrigated land. Small farmers possess 9.07 ha (100 %) of dry land. Semi medium farmers possess 8.09 ha (60.94 %) of dry land and 5.19 ha (39.06 %) of irrigated land. Medium farmers possess 3.76 ha (76.23 %) of dry land and 1.17 ha (23.77 %) of Permanent Fallow land.
 - ❖ The results indicate that, the average value of dry land was Rs. 540,608.39, the average value of irrigated land was Rs. 714,905.94 and the average value of permanent fallow land was Rs. 1,703,448.22. In case of marginal farmers, the average land value was Rs. 948,696.85 for dry land and the average value of irrigated land was Rs. 1,976,000. In case of small farmers, the average land value was Rs. 396,787.15 for dry land. In case of semi medium farmers, the average land value was Rs. 247,000 for dry land and the average value of irrigated land was Rs. 616,536.66. In case of medium farmers, the average land value was Rs. 239,032.26 for dry land and the average value of permanent fallow land was Rs. 1,703,448.22.
 - ❖ The results indicate that, Bore Well was the major irrigation source in the micro watershed for 14.29 per cent of the farmers.
 - ❖ The results indicate that, marginal and semi medium farmers had an irrigated area of 0.40 ha and 5.19 ha.
 - ❖ The results indicate that, farmers have grown Green gram (2.02 ha), cotton (2.86 ha), paddy (4.30 ha), Red gram (26.56 ha) and Sorghum (1.62 ha). Marginal farmers have grown sorghum, red gram and cotton. Small farmers have grown red gram, cotton and green gram. Semi medium farmers have grown cotton, red gram and paddy. Medium farmers have grown red gram and paddy.
 - ❖ The results indicate that, the cropping intensity in Sudapur-2 micro-watershed was found to be 89.51 per cent.
 - ❖ The results indicate that, the total cost of cultivation for Cotton was Rs. 35806.62. The gross income realized by the farmers was Rs. 38739.82. The net income from

Cotton cultivation was Rs. 2933.20. Thus the benefit cost ratio was found to be 1: 1.08.

- ❖ The results indicate that, the total cost of cultivation for green gram was Rs. 18525.98. The gross income realized by the farmers was Rs. 49400. The net income from green gram cultivation was Rs. 30874.02. Thus the benefit cost ratio was found to be 1: 2.67.*
- ❖ The results indicate that, the total cost of cultivation for Red gram was Rs. 32506.88. The gross income realized by the farmers was Rs. 62432.47. The net income from Red gram cultivation was Rs. 29925.59. Thus the benefit cost ratio was found to be 1: 1.92.*
- ❖ The results indicate that, the total cost of cultivation for Paddy was Rs. 127772.06. The gross income realized by the farmers was Rs. 189158.35. The net income from Paddy cultivation was Rs. 61386.29. Thus the benefit cost ratio was found to be 1: 1.48.*
- ❖ The results indicate that, 57.14 per cent of the households opined that dry fodder was adequate and 54.29 per cent of the households opined that green fodder was adequate.*
- ❖ The results indicate that the annual gross income was Rs. 85,000 for landless farmers, for marginal farmers it was Rs. 121,564.21, for small farmers it was Rs. 126,557.14, semi medium farmers it was Rs. 233,428.57 and medium farmers it was Rs. 80,800.*
- ❖ The results indicate that the average annual expenditure is Rs. 19,493.59. For landless households it was Rs. 55,000, for marginal farmers it was Rs. 9,988.19, for small farmers it was Rs. 23,000, for semi medium farmers it was Rs. 29,500 and medium farmers it was Rs. 70,000.*
- ❖ The results indicate that, sampled households have grown 4 Custard apple trees in their field also 1 Coconut and 13 Custard apple trees in backyard.*
- ❖ The results indicate that, households have planted 4 Teak, 39 neem and 4 tamarind in their field and also 2 Tamarind, 2 acacia trees and 11 neem in backyard.*
- ❖ The results indicated that, households have an average investment capacity of Rs. 3,257.14 for land development Rs. 4,285.71 for Irrigation facility, Rs. 2,600 for improved crop production and Rs. 1,714.29 for improved livestock management.*
- ❖ The results indicated that Government subsidy was the source of additional investment for 5.71 per cent for irrigation facility, Loan from bank was the source of additional investment for 8.57 per cent for land development and 5.71 per cent for improved crop production and improved livestock management, Own funds was the source of additional investment for 5.71 per cent for land development, improved crop production and improved livestock management, soft loan was the source of additional investment for 17.14 per cent for land development, 2.86 per cent for irrigation facility and improved livestock management, 11.43 per cent for improved crop production.*

- ❖ *The results indicated that, cotton was sold to the extent of 100 per cent, Green gram was sold to the extent of 95.0 per cent, Paddy was sold to the extent of 96.92 per cent, Red gram was sold to the extent of 85.98 per cent and Sorghum to the extent of 88.24 per cent.*
- ❖ *The results indicated that, about 100 per cent of the farmers sold their produce to local/village merchants and 5.71 per cent of the farmers sold their produce to Regulated Market.*
- ❖ *The results indicated that, 100 per cent of the households have used Tractor as a mode of transportation.*
- ❖ *The results indicated that, 34.29 per cent of the households have experienced soil and water erosion problems in the farm.*
- ❖ *The results indicated that, 97.14 per cent have shown interest in soil test.*
- ❖ *The results indicated that, 85.71 per cent of the households used firewood, 5.71 per cent of the households used Kerosene and 14.29 per cent of the households used LPG as a source of fuel.*
- ❖ *The results indicated that, piped supply was the major source of drinking water for 94.29 per cent and 5.71 per cent of the households used bore well in the micro watershed.*
- ❖ *The results indicated that, Electricity was the major source of light for 100 per cent of the households in micro watershed.*
- ❖ *The results indicated that, 48.57 per cent of the households possess sanitary toilet facility.*
- ❖ *The results indicated that, 100 per cent of the sampled households possessed BPL cards.*
- ❖ *The results indicated that, 88.57 per cent of the households participated in NREGA programme.*
- ❖ *The results indicated that, cereals were adequate for 100 per cent of the households, pulses were adequate for 97.14 per cent of the households, Fruits were adequate for 20 per cent, vegetables were adequate for 91.43 per cent, milk, Egg and meat were adequate for 100 per cent.*
- ❖ *The results indicated that, pulses were inadequate for 2.86 per cent of the households, oilseeds were inadequate for 100 per cent, vegetables were inadequate for 11.43 per cent and fruits were inadequate for 80 per cent of the households.*
- ❖ *The results indicated that, lower fertility status of the soil was the constraint experienced by 100 per cent of the households,, wild animal menace on farm field, Inadequacy of irrigation water, High cost of Fertilizers and plant protection chemicals and Frequent incidence of pest and diseases was the constraint experienced by 97.14 per cent of the households, High rate of interest on credit (94.29 %) and low price for the agricultural commodities (5.71 %).*

INTRODUCTION

Soil and water are the two precious natural resources which are essential for crop production and existence of life on earth. Rainfed agriculture is under severe stress due to various constraints related to agriculture like uneven and erratic distribution of rainfall, indiscriminate use of fertilizers, chemicals and pesticides, adoption of improper land management practices, soil erosion, decline in soil fertility, decline in ground water resources leading to low crop productivity. The area under rainfed agriculture has to be managed effectively using the best available practices to enhance the production of food, fodder and fuel. This is possible if the land resources are characterized at each parcel of land through detailed land resource inventory using the best available techniques of remote sensing, GPS and GIS. The watershed development programs are aimed at the sustainable distribution of its resources and the process of creating and implementing plans, programs, and projects to sustain and enhance watershed functions that affect the plant, animal and human communities within a watershed boundary.

World Bank funded KWDP II, SUJALA III project was implemented in with Broad objective of demonstrating more effective watershed management through greater integration of programmes related to rain-fed agriculture, innovative and science based approaches and strengthen institutional capacities and If successful, it is expected that the systems and tools could be mainstreamed into the overall IWMP in the State of Karnataka and in time, throughout other IWMP operations in India. With this background the socio-economic survey has been carried out with following specific objectives:

1. To understand the demographic features of the households in the micro-watershed
2. To understand the extent of family labour available and additional employment opportunities available within the village.
3. To know the status of assets of households in the micro-watershed for suggesting possible improvements.
4. To study the cropping pattern, cropped area and productivity levels of different households in micro-watershed.
5. To determine the type and extent of livestock owned by different categories of HHs
6. Availability of fodder and level of livestock management.

Scope and importance of survey

Survey helps in identification of different socio-economic and resource use-patterns of farmers at the Micro watershed. Household survey provides demographic features, labour force, and levels of education; land ownership and asset position (including livestock and other household assets) of surveyed households; and cropping patterns, input intensities, and average crop yields from farmers' fields. It also discusses crop utilization and the degree of commercialization of production in the areas; farmers' access to and utilization of credit from formal and informal sources; and the level of adoption and use of soil, water, and pest management technologies.

METHODOLOGY

The description of the methods, components selected for the survey and procedures followed in conducting the baseline survey are furnished under the following heads.

Description of the study area

Yadgir District is one of the 30 districts of Karnataka state in southern India. This district was carved out from the erstwhile Gulbarga district as the 30th district of Karnataka on 10 April 2010. Yadgir town is the administrative headquarters of the district. The district comprises of 3 taluks namely, Shahapur, Yadgiri and Shorapur (There are 16 hoblies, 117 Gram Panchayats, 4 Municipalities, 8 Towns/ Urban agglomeration and 487 inhabited & 32 un-inhabited villages The district occupies an area of 5,160.88 km².

Yadgir district is the second smallest district in the state, area wise is very rich in cultural traditions. The vast stretch of fertile black soil of the district is known for bumper red gram and jowar crops. The district is a "Daal bowl" of the state. The district is also known for cluster of cement industries and a distinct stone popularly known as "Malakheda Stone". Two main rivers, Krishna and Bhima, and a few tributaries flow in this region. Krishna and Bhima Rivers drain the district. They constitute the two major river basins of the district. Kagna and Amarja are the two sub - basins of Bhima River, which occur within the geographical area of the district

According to the 2011 census Yadgir district has a population of 1, 172,985, roughly equal to the nation of Timor-Leste or the US state of Rhode Island. This gives it a ranking of 404th in India (out of a total of 640). The district has a population density of 224 inhabitants per square kilometre (580/sq mi). Its population growth rate over the decade 2001-2011 was 22.67%. Yadgir has a sex ratio of 984 females for every 1000 males, and a literacy rate of 52.36%.

Description of the micro watershed

Sudapur-2 micro-watershed in Duganur sub-watershed (Yadgir taluk and district) is located in between 16^o55'36.491'' to 16^o53'33.602'' North latitudes and 77^o19'58.212'' to 77^o18'4.961'' East longitudes, covering an area of about 589.01 ha, bounded by Gajarkota, Mitathapadamapalli, Yadhlapura and Siddapura.B villages.

Methodology followed in assessing socio-economic status of households

In order to assess the socio-economic condition of the farmers in the watershed a comprehensive questionnaire was prepared. Major components such as demographic conditions, migration details, food consumption and family expenditure pattern, material possession, land holding, land use management, cropping pattern, cost of cultivation of crops, livestock management. The statistical components such as frequency and percentage were used to analyze the data. About 35 households located in the micro-watershed were interviewed for the survey.

SALIENT FEATURES OF THE SURVEY

Households sampled for socio-economic survey: The data on households sampled for socio economic survey in Sudapur-2 micro-watershed is presented in Table 1 and it indicated that 35 farmers were sampled in Sudapur-2 micro-watershed among them 1 (2.86 %) were landless, 19 (54.29 %) were marginal farmers, 7 (20 %) were small farmers and semi medium farmers and 1 (2.86 %) were medium farmers.

Table 1: Households sampled for socio economic survey in Sudapur-2 micro-watershed

Sl.No.	Particulars	LL (1)		MF (19)		SF (7)		SMF (7)		MDF (1)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Farmers	1	2.86	19	54.29	7	20	7	20	1	2.86	35	100

Population characteristics: The population characteristics of households sampled for socio-economic survey in Sudapur-2 micro-watershed is presented in Table 2. The data indicated that there were 95 (51.91 %) men and 88 (48.09 %) women among the sampled households. The average family size of landless farmers' was 5, marginal farmers' was 5.57, small farmers' was 4.14, semi medium farmers' was 5.57 and medium farmers' was 4.

Table 2: Population characteristics of Sudapur-2 micro-watershed

Sl.No.	Particulars	LL (5)		MF (106)		SF (29)		SMF (39)		MDF (4)		All (183)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Men	3	60	54	50.94	15	51.72	20	51.28	3	75	95	51.91
2	Women	2	40	52	49.06	14	48.28	19	48.72	1	25	88	48.09
Total		5	100	106	100	29	100	39	100	4	100	183	100
Average		5		5.57		4.14		5.57		4		5.22	

Age wise classification of population: The age wise classification of household members in Sudapur-2 micro-watershed is presented in Table 3. The data indicated that, 39 (21.31 %) people were in 0-15 years of age, 83 (45.36 %) were in 16-35 years of age, 40 (21.86 %) were in 36-60 years of age and 21 (11.48 %) were above 61 years of age.

Table 3: Age wise classification of household members in Sudapur-2 micro-watershed

Sl. No.	Particulars	LL (5)		MF (106)		SF (29)		SMF (39)		MDF (4)		All (183)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	0-15 years of age	1	20	25	23.58	10	34.48	3	7.69	0	0	39	21.31
2	16-35 years of age	2	40	49	46.23	13	44.83	17	43.59	2	50	83	45.36
3	36-60 years of age	2	40	18	16.98	4	13.79	15	38.46	1	25	40	21.86
4	> 61 years	0	0	14	13.21	2	6.90	4	10.26	1	25	21	11.48
Total		5	100	106	100	29	100	39	100	4	100	183	100

Education level of household members: Education level of household members in Sudapur-2 micro-watershed is presented in Table 4. The results indicated that Sudapur-2

had 55.19 per cent illiterates, 1.09 per cent Functional Literate, 14.75 per cent of them had primary school, 1.64 per cent of them had middle school, 12.02 per cent of them had high school education, 4.37 per cent of them had PUC, 1.24 per cent of them had Diploma, 0.62 per cent of them had ITI and 5.46 per cent of them had Degree education.

Table 4. Education level of household members in Sudapur-2 micro-watershed

Sl. No.	Particulars	LL (5)		MF (106)		SF (29)		SMF (39)		MDF (4)		All (183)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Illiterate	4	80	55	51.89	16	55.17	23	58.97	3	75	101	55.19
2	Functional Literate	0	0	1	0.94	0	0	1	2.56	0	0	2	1.09
3	Primary School	0	0	19	17.92	5	17.24	3	7.69	0	0	27	14.75
4	Middle School	0	0	3	2.83	0	0	0	0	0	0	3	1.64
5	High School	0	0	15	14.15	1	3.45	6	15.38	0	0	22	12.02
6	PUC	0	0	4	3.77	1	3.45	3	7.69	0	0	8	4.37
7	Degree	0	0	4	3.77	3	10.34	2	5.13	1	25	10	5.46
8	Others	1	20	5	4.72	3	10.34	1	2.56	0	0	10	5.46
Total		5	100	106	100	29	100	39	100	4	100	183	100

Occupation of household heads: The data regarding the occupation of the household heads in Sudapur-2 micro-watershed is presented in Table 5. The results indicate that, 68.57 per cent of household heads were practicing agriculture, 25.71 per cent of the household heads were agricultural labourers, 5.71 per cent of the household heads were in Private Service and 2.86 per cent of the household heads were Housewives.

Table 5: Occupation of household heads in Sudapur-2 micro-watershed

Sl. No.	Particulars	LL (1)		MF (19)		SF (7)		SMF (7)		MDF (1)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Agriculture	0	0	13	68.42	5	71.43	6	85.71	0	0	24	68.57
2	Agricultural Labour	1	100	6	31.58	0	0	1	14.29	1	100	9	25.71
3	Private Service	0	0	0	0	2	28.57	0	0	0	0	2	5.71
4	Housewife	0	0	1	5.26	0	0	0	0	0	0	1	2.86
Total		1	100	20	100	7	100	7	100	1	100	36	100

Table 6: Occupation of family members in Sudapur-2 micro-watershed

Sl. No.	Particulars	LL (5)		MF (106)		SF (29)		SMF (39)		MDF (4)		All (183)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Agriculture	0	0	47	44.34	15	51.72	23	58.97	1	25	86	46.99
2	Agricultural Labour	4	80	12	11.32	0	0	2	5.13	2	50	20	10.93
3	General Labour	0	0	0	0	0	0	1	2.56	0	0	1	0.55
4	Government Service	0	0	1	0.94	0	0	1	2.56	0	0	2	1.09
5	Private Service	0	0	2	1.89	3	10.34	1	2.56	0	0	6	3.28
6	Trade & Business	0	0	0	0	0	0	1	2.56	0	0	1	0.55
7	Student	0	0	24	22.64	7	24.14	4	10.26	1	25	36	19.67
8	Housewife	0	0	13	12.26	0	0	5	12.82	0	0	18	9.84
9	Children	1	20	7	6.60	4	13.79	1	2.56	0	0	13	7.10
Total		5	100	106	100	29	100	39	100	4	100	183	100

Occupation of the household members: The data regarding the occupation of the household members in Sudapur-2 micro-watershed is presented in Table 6. The results indicate that agriculture was the major occupation for 46.99 per cent of the household members, 10.93 per cent were agricultural labourers, 0.55 per cent were General Labour, 1.09 per cent were Government Service, 3.28 per cent were Private Service, 0.55 per cent were Trade & Business, 19.67 per cent were Student, 9.84 per cent were Housewife and 7.10 per cent were children.

Institutional participation of the household members: The data regarding the institutional participation of the household members in Sudapur-2 micro-watershed is presented in Table 7. The results show that, 100 per cent of the population in the micro watershed has not participated in any institutions.

Table 7. Institutional Participation of household members in Sudapur-2 micro-watershed

Sl.No.	Particulars	LL (5)		MF (106)		SF (29)		SMF (39)		MDF (4)		All (183)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	No Participation	5	100	106	100	29	100	39	100	4	100	183	100
	Total	5	100	106	100	29	100	39	100	4	100	183	100

Type of house owned: The data regarding the type of house owned by the households in Sudapur-2 micro-watershed is presented in Table 8. The results indicate that 11.43 per cent of the households possess thatched house, 77.14 per cent of the households possess katcha house and 11.43 per cent of them possess pucca/RCC house.

Table 8. Type of house owned by households in Sudapur-2 micro-watershed

Sl.No.	Particulars	LL (1)		MF (19)		SF (7)		SMF (7)		MDF (1)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Thatched	0	0	4	21.05	0	0	0	0	0	0	4	11.43
2	Katcha	1	100	13	68.42	5	71.43	7	100	1	100	27	77.14
3	Pucca/RCC	0	0	2	10.53	2	28.57	0	0	0	0	4	11.43
	Total	1	100	19	100	7	100	7	100	1	100	35	100

Table 9. Durable Assets owned by households in Sudapur-2 micro-watershed

Sl.No.	Particulars	LL (1)		MF (19)		SF (7)		SMF (7)		MDF (1)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Television	1	100	16	84.21	7	100	6	85.71	1	100	31	88.57
2	Mixer/Grinder	0	0	3	15.79	4	57.14	0	0	0	0	7	20
3	Refrigerator	0	0	0	0	1	14.29	1	14.29	0	0	2	5.71
4	Bicycle	0	0	1	5.26	4	57.14	0	0	0	0	5	14.29
5	Motor Cycle	0	0	8	42.11	4	57.14	2	28.57	0	0	14	40
6	Mobile Phone	1	100	19	100	7	100	7	100	1	100	35	100

Durable Assets owned by the households: The data regarding the Durable Assets owned by the households in Sudapur-2 micro-watershed is presented in Table 9. The results show that 88.57 per cent of the households possess TV, 20 per cent of the

households possess mixer/grinder, 5.71 per cent of the households possess Refrigerator, 14.29 per cent of the households possess Bicycle, 40 per cent of the households possess Motor Cycle and 100 per cent of the households possess mobile phones.

Average value of durable assets: The data regarding the average value of durable assets owned by the households in Sudapur-2 micro-watershed is presented in Table 10. The results show that the average value of television was Rs. 9,000, mixer/grinder was Rs. 2,000, Refrigerator was Rs. 8,500, Bicycle was Rs. 2,000, motor cycle was Rs. 55,500 and mobile phone was Rs. 2,630.

Table 10. Average value of durable assets owned by households in Sudapur-2 micro-watershed Average value (Rs.)

Sl.No.	Particulars	LL (1)	MF (19)	SF (7)	SMF (7)	MDF (1)	All (35)
1	Television	9,000	9,000	9,000	9,000	9,000	9,000
2	Mixer/Grinder	0	2,000	2,000	0	0	2,000
3	Refrigerator	0	0	8,000	9,000	0	8,500
4	Bicycle	0	2,000	2,000	0	0	2,000
5	Motor Cycle	0	54,875	58,250	52,500	0	55,500
6	Mobile Phone	2,000	2,400	2,818	3,250	2,000	2,630

Farm Implements owned: The data regarding the farm implements owned by the households in Sudapur-2 micro-watershed is presented in Table 11. About 14.29 per cent of the households possess bullock cart, 40 per cent of them possess plough, 22.86 per cent of them possess seed/fertilizer drill, 2.86 per cent of them possess Harvester, 17.14 per cent of them possess Sprayer and 34.29 per cent of them possess Weeder.

Table 11. Farm Implements owned by households in Sudapur-2 micro-watershed

Sl.No.	Particulars	LL (1)		MF (19)		SF (7)		SMF (7)		MDF (1)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Bullock Cart	0	0	3	15.79	2	28.57	0	0	0	0	5	14.29
2	Plough	0	0	10	52.63	2	28.57	2	28.57	0	0	14	40
3	Seed/Fertilizer Drill	0	0	6	31.58	1	14.29	1	14.29	0	0	8	22.86
4	Sprayer	0	0	3	15.79	1	14.29	2	28.57	0	0	6	17.14
5	Weeder	0	0	7	36.84	4	57.14	1	14.29	0	0	12	34.29
6	Harvester	0	0	1	5.26	0	0	0	0	0	0	1	2.86
7	Blank	1	100	8	42.11	3	42.86	4	57.14	1	100	17	48.57

Table 12. Average value of farm implements owned by households in Sudapur-2 micro-watershed Average Value (Rs.)

Sl.No.	Particulars	LL (1)	MF (19)	SF (7)	SMF (7)	MDF (1)	All (35)
1	Bullock Cart	0	20,000	40,000	0	0	28,000
2	Plough	0	1,500	1,500	1,500	0	1,500
3	Seed/Fertilizer Drill	0	1,666	2,000	1,500	0	1,687
4	Sprayer	0	1,833	3,500	2,250	0	2,250
5	Weeder	0	52	60	100	0	56
6	Harvester	0	100	0	0	0	100

Average value of farm implements: The data regarding the average value of farm Implements owned by the households in Sudapur-2 micro-watershed is presented in Table 12. The results show that the average value of bullock cart was Rs. 28,000, plough was Rs. 1,500, seed/fertilizer drill was Rs. 1,687, Harvester was Rs. 100, sprayer was Rs. 3,750 and weeder was Rs. 30.

Livestock possession by the households: The data regarding the Livestock possession by the households in Sudapur-2 micro-watershed is presented in Table 13. The results indicate that, 42.86 per cent of the households possess bullocks, 20 per cent of the households possess local cow and 5.71 per cent of the households possess Buffalo, Goat and Poultry birds.

Table 13. Livestock possession by households in Sudapur-2 micro-watershed

Sl.No.	Particulars	LL (1)		MF (19)		SF (7)		SMF (7)		MDF (1)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Bullock	0	0	10	52.63	2	28.57	3	42.86	0	0	15	42.86
2	Local cow	0	0	4	21.05	1	14.29	2	28.57	0	0	7	20
3	Buffalo	0	0	1	5.26	0	0	1	14.29	0	0	2	5.71
4	Goat	0	0	1	5.26	0	0	1	14.29	0	0	2	5.71
5	Poultry birds	0	0	1	5.26	0	0	1	14.29	0	0	2	5.71
6	blank	1	100	7	36.84	5	71.43	2	28.57	1	100	16	45.71

Average Labour availability: The data regarding the average labour availability in Sudapur-2 micro-watershed is presented in Table 14. The results indicate that, average own labour men available in the micro watershed was 1.62, average own labour (women) available was 1.68, average hired labour (men) available was 11.32 and average hired labour (women) available was 8.09.

In case of marginal farmers, average own labour men available was 1.74, average own labour (women) was 1.79, average hired labour (men) was 9.74 and average hired labour (women) available was 6.58. In case of small farmers, average own labour men available was 0.71, average own labour (women) was 1.14, average hired labour (men) was 17.14 and average hired labour (women) available was 12.14. In case of semi medium farmers, average own labour men available was 2.29, average own labour (women) was 2, average hired labour (men) was 8.57 and average hired labour (women) available was 6.43. In case of medium farmers, average own labour men available was 1, average own labour (women) was 1, average hired labour (men) was 20 and average hired labour (women) available was 20.

Table 14. Average Labour availability in Sudapur-2 micro-watershed

Sl.No.	Particulars	LL (1)	MF (19)	SF (7)	SMF (7)	MDF (1)	All (35)
		N	N	N	N	N	N
1	Hired labour Female	0	6.58	12.14	6.43	20	8.09
2	Own Labour Female	0	1.79	1.14	2	1	1.68
3	Own labour Male	0	1.74	0.71	2.29	1	1.62
4	Hired labour Male	0	9.74	17.14	8.57	20	11.32

Adequacy of Hired Labour: The data regarding the adequacy of hired labour in Sudapur-2 micro-watershed is presented in Table 15. The results indicate that, 94.29 per cent of the households opined that the hired labour was adequate.

Table 15. Adequacy of Hired Labour in Sudapur-2 micro-watershed

Sl.No.	Particulars	LL (1)		MF (19)		SF (7)		SMF (7)		MDF (1)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Adequate	0	0	18	94.74	7	100	7	100	1	100	33	94.29

Distribution of land (ha): The data regarding the distribution of land (ha) in Sudapur-2 micro-watershed is presented in Table 16. The results indicate that, households of the Sudapur-2 micro-watershed possess 32.73 ha (82.87 %) of dry land, 5.59 ha (14.16 %) of irrigated land and 1.17 ha (2.97 %) of Permanent Fallow land. Marginal farmers possess 11.80 ha (96.68 %) of dry land and 0.40 ha (3.32 %) of irrigated land. Small farmers possess 9.07 ha (100 %) of dry land. Semi medium farmers possess 8.09 ha (60.94 %) of dry land and 5.19 ha (39.06 %) of irrigated land. Medium farmers possess 3.76 ha (76.23 %) of dry land and 1.17 ha (23.77 %) of Permanent Fallow land.

Table 16. Distribution of land (Ha) in Sudapur-2 micro-watershed

Sl. No.	Particulars	MF (19)		SF (7)		SMF (7)		MDF (1)		All (35)	
		ha	%	ha	%	ha	%	ha	%	ha	%
1	Dry	11.80	96.68	9.07	100	8.09	60.94	3.76	76.23	32.73	82.87
2	Irrigated	0.40	3.32	0	0	5.19	39.06	0	0	5.59	14.16
3	Permanent Fallow	0	0	0	0	0	0	1.17	23.77	1.17	2.97
Total		12.21	100	9.07	100	13.28	100	4.94	100	39.49	100

Average land value (Rs./ha): The data regarding the average land value (Rs./ha) in Sudapur-2 micro-watershed is presented in Table 17. The results indicate that, the average value of dry land was Rs. 540,608.39, the average value of irrigated land was Rs. 714,905.94 and the average value of permanent fallow land was Rs. 1,703,448.22. In case of marginal famers, the average land value was Rs. 948,696.85 for dry land and the average value of irrigated land was Rs. 1,976,000. In case of small famers, the average land value was Rs. 396,787.15 for dry land. In case of semi medium famers, the average land value was Rs. 247,000 for dry land and the average value of irrigated land was Rs. 616,536.66. In case of medium farmers, the average land value was Rs. 239,032.26 for dry land and the average value of permanent fallow land was Rs. 1,703,448.22.

Table 17. Average land value (Rs./ha) in Sudapur-2 micro-watershed

Sl. No.	Particulars	MF (19)	SF (7)	SMF (7)	MDF (1)	All (35)
		N	N	N	N	N
1	Dry	948,696.85	396,787.15	247,000	239,032.26	540,608.39
2	Irrigated	1,976,000	0	616,536.66	0	714,905.94
3	Permanent Fallow	0	0	0	1,703,448.22	1,703,448.22

Source of irrigation: The data regarding the source of irrigation in Sudapur-2 micro-watershed is presented in Table 18. The results indicate that, Bore Well was the major irrigation source in the micro water shed for 14.29 per cent of the farmers.

Table 18. Source of irrigation in Sudapur-2 micro-watershed

Sl.No.	Particulars	LL (1)		MF (19)		SF (7)		SMF (7)		MDF (1)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Bore Well	0	0	1	5.26	0	0	4	57.14	0	0	5	14.29

Irrigated Area (ha): The data regarding the irrigated area (ha) in Sudapur-2 micro-watershed is presented in Table 19. The results indicate that, marginal and semi medium farmers had an irrigated area of 0.40 ha and 5.19 ha.

Table 19. Irrigated Area (ha) in Sudapur-2 micro-watershed

Sl.No.	Particulars	LL (1)	MF (19)	SF (7)	SMF (7)	MDF (1)	All (35)
1	Kharif	0	0.40	0	5.19	0	5.60

Cropping pattern: The data regarding the cropping pattern in Sudapur-2 micro-watershed is presented in Table 20. The results indicate that, farmers have grown Green gram (2.02 ha), cotton (2.86 ha), paddy (4.30 ha), Red gram (26.56 ha) and Sorghum (1.62 ha). Marginal farmers have grown sorghum, red gram and cotton. Small farmers have grown red gram, cotton and green gram. Semi medium farmers have grown cotton, red gram and paddy. Medium farmers have grown red gram and paddy.

Table 20. Cropping pattern in Sudapur-2 micro-watershed (Area in ha)

Sl.No.	Particulars	LL (1)	MF (19)	SF (7)	SMF (7)	MDF (1)	All(35)
1	Kharif - Cotton	0	0.43	1.21	1.21	0	2.86
2	Kharif - Greengram	0	0	2.02	0	0	2.02
3	Kharif - Paddy	0	0	0	4.18	0.12	4.30
4	Kharif - Red gram (togari)	0	10.16	6.88	7.89	1.62	26.56
5	Kharif - Sorghum	0	1.62	0	0	0	1.62
Total		0	12.21	10.12	13.29	1.74	37.36

Cropping intensity: The data regarding the cropping intensity in Sudapur-2 micro-watershed is presented in Table 21. The results indicate that, the cropping intensity in Sudapur-2 micro-watershed was found to be 89.51 per cent.

Table 21. Cropping intensity (%) in Sudapur-2 micro-watershed

Sl.No.	Particulars	LL (1)	MF (19)	SF (7)	SMF (7)	MDF (1)	All (35)
1	Cropping Intensity	0	100	83.84	84.54	100	89.51

Cost of cultivation of Cotton: The data regarding the cost of cultivation of Cotton in Sudapur-2 micro-watershed is presented in Table 22. The results indicate that, the total cost of cultivation for Cotton was Rs. 35806.62. The gross income realized by the farmers was Rs. 38739.82. The net income from Cotton cultivation was Rs. 2933.20. Thus the benefit cost ratio was found to be 1: 1.08.

Table 22. Cost of Cultivation of Cotton in Sudapur-2 micro-watershed

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
I	Cost A1				
1	Hired Human Labour	Man days	51.31	9806.06	27.39
2	Bullock	Pairs/day	3.98	2386.11	6.66
3	Tractor	Hours	3.25	2597.38	7.25
4	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	4.57	4343.73	12.13
5	FYM	Quintal	3.15	630.70	1.76
6	Fertilizer + micronutrients	Quintal	5.85	4990.33	13.94
7	Pesticides (PPC)	Kgs / liters	1.33	1325.62	3.70
8	Depreciation charges		0	25.41	0.07
9	Land revenue and Taxes		0	3.29	0.01
II	Cost B1				
10	Interest on working capital			1354.97	3.78
11	Cost B1 = (Cost A1 + sum of 15 and 16)			27463.61	76.70
III	Cost B2				
12	Rental Value of Land			333.33	0.93
13	Cost B2 = (Cost B1 + Rental value)			27796.94	77.63
IV	Cost C1				
14	Family Human Labour		20.80	4753.53	13.28
15	Cost C1 = (Cost B2 + Family Labour)			32550.47	90.91
V	Cost C2				
16	Risk Premium			1	0
17	Cost C2 = (Cost C1 + Risk Premium)			32551.47	90.91
VI	Cost C3				
18	Managerial Cost			3255.15	9.09
19	Cost C3 = (Cost C2 + Managerial Cost)			35806.62	100
VII	Economics of the Crop				
a.	Main Product	a) Main Product (q)		7.50	38739.82
		b) Main Crop Sales Price (Rs.)			5166.67
b.	Gross Income (Rs.)			38739.82	
c.	Net Income (Rs.)			2933.20	
d.	Cost per Quintal (Rs./q.)			4775.47	
e.	Benefit Cost Ratio (BC Ratio)			1:1.08	

Cost of cultivation of Green gram: The data regarding the cost of cultivation of green gram in Sudapur-2 micro-watershed is presented in Table 23. The results indicate that, the total cost of cultivation for green gram was Rs. 18525.98. The gross income realized by the farmers was Rs. 49400. The net income from green gram cultivation was Rs. 30874.02. Thus the benefit cost ratio was found to be 1: 2.67.

Table 23. Cost of Cultivation of Green gram in Sudapur-2 micro-watershed

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
I	Cost A1				
1	Hired Human Labour	Man days	37.46	6545.50	35.33
2	Bullock	Pairs/day	2.88	1729	9.33
3	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	7.41	889.20	4.80
4	FYM	Quintal	2.06	411.67	2.22
5	Fertilizer + micronutrients	Quintal	2.88	2276.52	12.29
6	Pesticides (PPC)	Kgs / liters	1.03	1029.17	5.56
7	Depreciation charges		0	723.72	3.91
8	Land revenue and Taxes		0	3.29	0.02
II	Cost B1				
9	Interest on working capital			552.91	2.98
10	Cost B1 = (Cost A1 + sum of 15 and 16)			14160.97	76.44
III	Cost B2				
11	Rental Value of Land			333.33	1.80
12	Cost B2 = (Cost B1 + Rental value)			14494.30	78.24
IV	Cost C1				
13	Family Human Labour		10.70	2346.50	12.67
14	Cost C1 = (Cost B2 + Family Labour)			16840.80	90.90
V	Cost C2				
15	Risk Premium			1	0.01
16	Cost C2 = (Cost C1 + Risk Premium)			16841.80	90.91
VI	Cost C3				
17	Managerial Cost			1684.18	9.09
18	Cost C3 = (Cost C2 + Managerial Cost)			18525.98	100
VII	Economics of the Crop				
a.	Main Product	a) Main Product (q)		9.88	49400
		b) Main Crop Sales Price (Rs.)			5000
b.	Gross Income (Rs.)			49400	
c.	Net Income (Rs.)			30874.02	
d.	Cost per Quintal (Rs./q.)			1875.10	
e.	Benefit Cost Ratio (BC Ratio)			1:2.67	

Cost of cultivation of Red gram: The data regarding the cost of cultivation of Red gram in Sudapur-2 micro-watershed is presented in Table 24. The results indicate that, the total cost of cultivation for Red gram was Rs. 32506.88. The gross income realized by the farmers was Rs. 62432.47. The net income from Red gram cultivation was Rs. 29925.59. Thus the benefit cost ratio was found to be 1: 1.92.

Table 24. Cost of Cultivation of Red gram in Sudapur-2 micro-watershed

Sl. No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
I	Cost A1				
1	Hired Human Labour	Man days	40.84	7260.94	22.34
2	Bullock	Pairs/day	5.08	3166.36	9.74
3	Tractor	Hours	4.85	3913.45	12.04
4	Machinery	Hours	0.40	329.33	1.01
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	8.75	1056.27	3.25
6	Seed Inter Crop	Kgs.	0	0	0
7	FYM	Quintal	2.96	591.23	1.82
8	Fertilizer + micronutrients	Quintal	4.50	3827.28	11.77
9	Pesticides (PPC)	Kgs /liters	1.42	1419.85	4.37
10	Irrigation	Number	4.94	0	0
11	Repairs		0	0	0
12	Msc. Charges (Marketing costs etc)		0	0	0
13	Depreciation charges		0	175.70	0.54
14	Land revenue and Taxes		0	3.29	0.01
II	Cost B1				
16	Interest on working capital			827.48	2.55
17	Cost B1 = (Cost A1 + sum of 15 and 16)			22571.20	69.44
III	Cost B2				
18	Rental Value of Land			333.33	1.03
19	Cost B2 = (Cost B1 + Rental value)			22904.53	70.46
IV	Cost C1				
20	Family Human Labour		28.69	6646.18	20.45
21	Cost C1 = (Cost B2 + Family Labour)			29550.71	90.91
V	Cost C2				
22	Risk Premium			1	0
23	Cost C2 = (Cost C1 + Risk Premium)			29551.71	90.91
VI	Cost C3				
24	Managerial Cost			2955.17	9.09
25	Cost C3 = (Cost C2 + Managerial Cost)			32506.88	100
VII	Economics of the Crop				
a.	Main Product	a) Main Product (q)		11.55	51130.07
		b) Main Crop Sales Price (Rs.)			4425.93
	By Product	e) Main Product (q)		8.16	11302.40
		f) Main Crop Sales Price (Rs.)			1385.19
b.	Gross Income (Rs.)			62432.47	
c.	Net Income (Rs.)			29925.59	
d.	Cost per Quintal (Rs./q.)			2813.86	
e.	Benefit Cost Ratio (BC Ratio)			1:1.92	

Cost of Cultivation of Paddy: The data regarding the cost of cultivation of Paddy in Sudapur-2 micro-watershed is presented in Table 25. The results indicate that, the total cost of cultivation for Paddy was Rs. 127772.06. The gross income realized by the farmers was Rs. 189158.35. The net income from Paddy cultivation was Rs. 61386.29. Thus the benefit cost ratio was found to be 1: 1.48.

Table 25. Cost of Cultivation of Paddy in Sudapur-2 micro-watershed

Sl.No	Particulars	Units	Phy Units	Value (Rs.)	% to C3
I	Cost A1				
1	Hired Human Labour	Man days	145.17	30692.91	24.02
2	Bullock	Pairs/day	5.04	3025.75	2.37
3	Tractor	Hours	5.98	4781.29	3.74
4	Machinery	Hours	0	0	0
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	107.23	45614.09	35.70
6	Seed Inter Crop	Kgs.	0	0	0
7	FYM	Quintal	1.44	288.17	0.23
8	Fertilizer + micronutrients	Quintal	13.41	10573.86	8.28
9	Pesticides (PPC)	Kgs /liters	2.60	2604.54	2.04
10	Irrigation	Number	0	0	0
11	Repairs		0	0	0
12	Msc. Charges (Marketing costs etc)		0	0	0
13	Depreciation charges		0	17.03	0.01
14	Land revenue and Taxes		0	3.29	0
II	Cost B1				
16	Interest on working capital			7089.80	5.55
17	Cost B1 = (Cost A1 + sum of 15 and 16)			104690.72	81.94
III	Cost B2				
18	Rental Value of Land			333.33	0.26
19	Cost B2 = (Cost B1 + Rental value)			105024.05	82.20
IV	Cost C1				
20	Family Human Labour		39.70	11131.37	8.71
21	Cost C1 = (Cost B2 + Family Labour)			116155.42	90.91
V	Cost C2				
22	Risk Premium			1	0
23	Cost C2 = (Cost C1 + Risk Premium)			116156.42	90.91
VI	Cost C3				
24	Managerial Cost			11615.64	9.09
25	Cost C3 = (Cost C2 + Managerial Cost)			127772.06	100
VII	Economics of the Crop				
a.	Main Product	a) Main Product (q)	105.45	168713.89	
		b) Main Crop Sales Price (Rs.)		1600	
	By Product	e) Main Product (q)	13.63	20444.46	
		f) Main Crop Sales Price (Rs.)		1500	
b.	Gross Income (Rs.)			189158.35	
c.	Net Income (Rs.)			61386.29	
d.	Cost per Quintal (Rs./q.)			1211.73	
e.	Benefit Cost Ratio (BC Ratio)			1:1.48	

Adequacy of fodder: The data regarding the adequacy of fodder in Sudapur-2 micro-watershed is presented in Table 26. The results indicate that, 57.14 per cent of the households opined that dry fodder was adequate and 54.29 per cent of the households opined that green fodder was adequate.

Table 26. Adequacy of fodder in Sudapur-2 micro-watershed

Sl.No.	Particulars	LL (1)		MF (19)		SF (7)		SMF(7)		MDF(1)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Adequate-Dry Fodder	0	0	13	68.42	2	28.57	5	71.43	0	0	20	57.14
2	Adequate-Green Fodder	0	0	13	68.42	2	28.57	4	57.14	0	0	19	54.29

Annual gross income: The data regarding the annual gross income in Sudapur-2 micro-watershed is presented in Table 27. The results indicate that the annual gross income was Rs. 85,000 for landless farmers, for marginal farmers it was Rs. 121,564.21, for small farmers it was Rs. 126,557.14, semi medium farmers it was Rs. 233,428.57 and medium farmers it was Rs. 80,800.

Table 27. Annual gross income in Sudapur-2 micro-watershed

(Avg. value in Rs.)

Sl.No.	Particulars	LL (1)	MF (19)	SF (7)	SMF (7)	MDF (1)	All (35)
		Rs.	Rs.	Rs.	Rs.	Rs.	Rs.
1	Service/salary	0	15,789.47	35,714.29	0	0	15,714.29
2	Business	0	0	0	11,428.57	0	2,285.71
3	Wage	85,000	56,842.11	23,571.43	14,285.71	0	40,857.14
4	Agriculture	0	48,442.11	67,271.43	194,228.57	80,800	80,905.71
5	Dairy Farm	0	490.53	0	4,914.29	0	1,249.14
6	Goat Farming	0	0	0	8,571.43	0	1,714.29
	Income(Rs.)	85,000	121,564.21	126,557.14	233,428.57	80,800	142,726.29

Average annual expenditure: The data regarding the average annual expenditure in Sudapur-2 micro-watershed is presented in Table 28. The results indicate that the average annual expenditure is Rs. 19,493.59. For landless households it was Rs. 55,000, for marginal farmers it was Rs. 9,988.19, for small farmers it was Rs. 23,000, for semi medium farmers it was Rs. 29,500 and medium farmers it was Rs. 70,000.

Table 28. Average annual expenditure in Sudapur-2 micro-watershed

(Avg. value in Rs.)

Sl.No.	Particulars	LL (1)	MF (19)	SF (7)	SMF (7)	MDF (1)	All (35)
		Rs.	Rs.	Rs.	Rs.	Rs.	Rs.
1	Service/salary	0	112,500	90,000	0	0	11,571.43
2	Business	0	0	0	45,000	0	1,285.71
3	Wage	55,000	44,117.65	27,000	22,500	0	27,371.43
4	Agriculture	0	30,157.89	44,000	110,000	70,000	49,171.43
5	Dairy Farm	0	3,000	0	9,000	0	685.71
6	Goat Farming	0	0	0	20,000	0	571.43
	Total	55,000	189,775.54	161,000	206,500	70,000	682,275.54
	Average	55,000	9,988.19	23,000	29,500	70,000	19,493.59

Horticulture species grown: The data regarding horticulture species grown in Sudapur-2 micro-watershed is presented in Table 29. The results indicate that, sampled households have grown 4 Custard apple trees in their field also 1 Coconut and 13 Custard apple trees in backyard.

Table 29. Horticulture species grown in Sudapur-2 micro-watershed

Sl. No.	Particulars	LL (1)		MF (19)		SF (7)		SMF (7)		MDF (1)		All (35)	
		F	B	F	B	F	B	F	B	F	B	F	B
1	Coconut	0	0	0	0	0	0	0	1	0	0	0	1
2	Custard apple	0	0	0	2	2	8	2	3	0	0	4	13

*F= Field B=Back Yard

Forest species grown: The data regarding forest species grown in Sudapur-2 micro-watershed is presented in Table 30. The results indicate that, households have planted 4 Teak, 39 neem and 4 tamarind in their field and also 2 Tamarind, 2 acacia trees and 11 neem in backyard.

Table 30: Forest species grown in Sudapur-2 micro-watershed

Sl.No.	Particulars	LL (1)		MF (19)		SF (7)		SMF (7)		MDF (1)		All (35)	
		F	B	F	B	F	B	F	B	F	B	F	B
1	Neem	0	0	16	4	14	1	6	6	3	0	39	11
2	Tamarind	0	0	2	1	1	1	1	0	0	0	4	2
3	Acacia	0	0	0	0	0	2	0	0	0	0	0	2

*F= Field B=Back Yard

Average Additional investment capacity: The data regarding average additional investment capacity in Sudapur-2 micro-watershed is presented in Table 31. The results indicated that, households have an average investment capacity of Rs. 3,257.14 for land development Rs. 4,285.71 for Irrigation facility, Rs. 2,600 for improved crop production and Rs. 1,714.29 for improved livestock management.

Table 31: Average Additional investment capacity in Sudapur-2 micro-watershed

Sl.No.	Particulars	MF (19)	SF (7)	SMF (7)	MDF (1)	All (35)
		Rs.	Rs.	Rs.	Rs.	Rs.
1	Land development	3,105.26	2,857.14	4,428.57	4,000	3,257.14
2	Irrigation facility	5,263.16	7,142.86	0	0	4,285.71
3	Improved crop production	421.05	2,857.14	8,428.57	4,000	2,600
4	Improved livestock management	421.05	0	7,428.57	0	1,714.29

Source of additional investment: The data regarding source of funds for additional investment in Sudapur-2 micro-watershed is presented in Table 32. The results indicated that Government subsidy was the source of additional investment for 5.71 per cent for irrigation facility, Loan from bank was the source of additional investment for 8.57 per cent for land development and 5.71 per cent for improved crop production and improved livestock management, Own funds was the source of additional investment for 5.71 per cent for land development, improved crop production and improved livestock

management, soft loan was the source of additional investment for 17.14 per cent for land development, 2.86 per cent for irrigation facility and improved livestock management, 11.43 per cent for improved crop production.

Table 32: Source of funds for additional investment capacity in Sudapur-2 micro-watershed

Sl. No	Item	Land development		Irrigation facility		Improved crop production		Improved livestock management	
		N	%	N	%	N	%	N	%
1	Government subsidy	0	0.0	2	5.71	0	0.0	0	0.0
2	Loan from bank	3	8.57	0	0.0	2	5.71	2	5.71
3	Own funds	2	5.71	0	0.0	2	5.71	2	5.71
4	Soft loan	6	17.14	1	2.86	4	11.43	1	2.86

Marketing of the agricultural produce: The data regarding marketing of the agricultural produce in Sudapur-2 micro-watershed is presented in Table 33. The results indicated that, cotton was sold to the extent of 100 per cent, Greengram was sold to the extent of 95.0 per cent, Paddy was sold to the extent of 96.92 per cent, Redgram was sold to the extent of 85.98 per cent and Sorghum to the extent of 88.24 per cent.

Table 33. Marketing of the agricultural produce in Sudapur-2 micro-watershed

Sl.No	Crops	Output obtained (q)	Output retained (q)	Output sold (q)	Output sold (%)	Avg. Price obtained (Rs/q)
1	Cotton	20.0	0.0	20.0	100.0	5166.67
2	Greengram	20.0	1.0	19.0	95.0	5000.0
3	Paddy	325.0	10.0	315.0	96.92	1600.0
4	Redgram	264.0	37.0	227.0	85.98	4596.15
5	Sorghum	17.0	2.0	15.0	88.24	2350.0

Marketing Channels used for sale of agricultural produce: The data regarding marketing channels used for sale of agricultural produce in Sudapur-2 micro-watershed is presented in Table 34. The results indicated that, about 100 per cent of the farmers sold their produce to local/village merchants and 5.71 per cent of the farmers sold their produce to Regulated Market.

Table 34. Marketing Channels used for sale of agricultural produce in Sudapur-2 micro-watershed

Sl.No.	Particulars	LL (1)		MF (19)		SF (7)		SMF (7)		MDF (1)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Local/village Merchant	0	0	18	94.74	8	114.29	8	114.29	1	100	35	100
2	Regulated Market	0	0	1	5.26	1	14.29	0	0	0	0	2	5.71

Mode of transport of agricultural produce: The data regarding mode of transport of agricultural produce in Sudapur-2 micro-watershed is presented in Table 35. The results indicated that, 100 per cent of the households have used Tractor as a mode of transportation.

Table 35. Mode of transport of agricultural produce in Sudapur-2 micro-watershed

Sl.No.	Particulars	LL (1)		MF (19)		SF (7)		SMF (7)		MDF (1)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Tractor	0	0	19	100	7	100	7	100	1	100	35	100

Incidence of soil and water erosion problems: The data regarding incidence of soil and water erosion problems in Sudapur-2 micro-watershed is presented in Table 36. The results indicated that, 34.29 per cent of the households have experienced soil and water erosion problems in the farm.

Table 36. Incidence of soil and water erosion problems in Sudapur-2 micro-watershed

Sl.No.	Particulars	LL (1)		MF (19)		SF (7)		SMF (7)		MDF (1)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Soil and water erosion problems in the farm	0	0	8	42.11	1	14.29	2	28.57	1	100	12	34.29

Interest shown towards soil testing: The data regarding Interest shown towards soil testing in Sudapur-2 micro-watershed is presented in Table 37. The results indicated that, 97.14 per cent have shown interest in soil test.

Table 37. Interest shown towards soil testing in Sudapur-2 micro-watershed

Sl.No.	Particulars	LL (1)		MF (19)		SF (7)		SMF (7)		MDF (1)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Interest in soil test	0	0	19	100	7	100	7	100	1	100	34	97.14

Usage pattern of fuel for domestic use: The data regarding usage pattern of fuel for domestic use in Sudapur-2 micro-watershed is presented in Table 38. The results indicated that, 85.71 per cent of the households used firewood, 5.71 per cent of the households used Kerosene and 14.29 per cent of the households used LPG as a source of fuel.

Table 38. Usage pattern of fuel for domestic use in Sudapur-2 micro-watershed

Sl.No.	Particulars	LL (1)		MF (19)		SF (7)		SMF (7)		MDF (1)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Fire Wood	1	100	19	100	4	57.14	6	85.71	0	0	30	85.71
2	LPG	0	0	0	0	3	42.86	1	14.29	1	100	5	14.29

Table 39. Source of drinking water in Sudapur-2 micro-watershed

Sl.No.	Particulars	LL (1)		MF (19)		SF (7)		SMF (7)		MDF (1)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Piped supply	1	100	18	94.74	6	85.71	7	100	1	100	33	94.29
2	Bore Well	0	0	1	5.26	1	14.29	0	0	0	0	2	5.71

Source of drinking water: The data regarding source of drinking water in Sudapur-2 micro-watershed is presented in Table 39. The results indicated that, piped supply was the

major source of drinking water for 94.29 per cent and 5.71 per cent of the households used bore well in the micro watershed.

Source of light: The data regarding source of light in Sudapur-2 micro-watershed is presented in Table 40. The results indicated that, Electricity was the major source of light for 100 per cent of the households in micro watershed.

Table 40. Source of light in Sudapur-2 micro-watershed

Sl.No.	Particulars	LL (1)		MF (19)		SF (7)		SMF (7)		MDF (1)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Electricity	1	100	19	100	7	100	7	100	1	100	35	100

Existence of Sanitary toilet facility: The data regarding existence of sanitary toilet facility in Sudapur-2 micro-watershed is presented in Table 41. The results indicated that, 48.57 per cent of the households possess sanitary toilet facility.

Table 41. Existence of Sanitary toilet facility in Sudapur-2 micro-watershed

Sl.No.	Particulars	LL (1)		MF (19)		SF (7)		SMF (7)		MDF (1)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Sanitary toilet facility	1	100	7	36.84	4	57.14	4	57.14	1	100	17	48.57

Possession of PDS card: The data regarding possession of PDS card in Sudapur-2 micro-watershed is presented in Table 42. The results indicated that, 100 per cent of the sampled households possessed BPL cards.

Table 42. Possession of PDS card in Sudapur-2 micro-watershed

Sl.No.	Particulars	LL (1)		MF (19)		SF (7)		SMF (7)		MDF (1)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	BPL	1	100	19	100	7	100	7	100	1	100	35	100

Participation in NREGA program: The data regarding participation in NREGA programme in Sudapur-2 micro-watershed is presented in Table 43. The results indicated that, 88.57 per cent of the households participated in NREGA programme.

Table 43. Participation in NREGA programme in Sudapur-2 micro-watershed

Sl.No.	Particulars	LL (1)		MF (19)		SF (7)		SMF (7)		MDF (1)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Participation in NREGA programme	1	100	18	94.74	4	57.14	7	100	1	100	31	88.57

Adequacy of food items: The data regarding adequacy of food items in Sudapur-2 micro-watershed is presented in Table 44. The results indicated that, cereals were adequate for 100 per cent of the households, pulses were adequate for 97.14 per cent of the households, Fruits were adequate for 20 per cent, vegetables were adequate for 91.43 per cent, milk, Egg and meat were adequate for 100 per cent.

Table 44. Adequacy of food items in Sudapur-2 micro-watershed

Sl.No.	Particulars	LL (1)		MF (19)		SF (7)		SMF (7)		MDF (1)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Cereals	1	100	19	100	7	100	7	100	1	100	35	100
2	Pulses	1	100	19	100	6	85.71	7	100	1	100	34	97.14
3	Vegetables	1	100	17	89.47	7	100	6	85.71	1	100	32	91.43
4	Fruits	0	0	3	15.79	1	14.29	2	28.57	1	100	7	20
5	Milk	1	100	19	100	7	100	7	100	1	100	35	100
6	Egg	1	100	19	100	7	100	7	100	1	100	35	100
7	Meat	1	100	19	100	7	100	7	100	1	100	35	100

Response on Inadequacy of food items: The data regarding inadequacy of food items in Sudapur-2 micro-watershed is presented in Table 45. The results indicated that, pulses were inadequate for 2.86 per cent of the households, oilseeds were inadequate for 100 per cent, vegetables were inadequate for 11.43 per cent and fruits were inadequate for 80 per cent of the households.

Table 45. Response on Inadequacy of food items in Sudapur-2 micro-watershed

Sl.No.	Particulars	LL (1)		MF (19)		SF (7)		SMF (7)		MDF (1)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Pulses	0	0	0	0	1	14.29	0	0	0	0	1	2.86
2	Oilseed	1	100	19	100	7	100	7	100	1	100	35	100
3	Vegetables	0	0	2	10.53	1	14.29	1	14.29	0	0	4	11.43
4	Fruits	1	100	16	84.21	6	85.71	5	71.43	0	0	28	80

Farming constraints: The data regarding farming constraints experienced by households in Sudapur-2 micro-watershed is presented in Table 46. The results indicated that, lower fertility status of the soil was the constraint experienced by 100 per cent of the households,, wild animal menace on farm field, Inadequacy of irrigation water, High cost of Fertilizers and plant protection chemicals and Frequent incidence of pest and diseases was the constraint experienced by 97.14 per cent of the households, High rate of interest on credit (94.29 %) and low price for the agricultural commodities (5.71 %).

Table 46. Farming constraints Experienced in Sudapur-2 micro-watershed

Sl. No.	Particulars	MF (19)		SF (7)		SMF (7)		MDF (1)		All (35)	
		N	%	N	%	N	%	N	%	N	%
1	Lower fertility status of the soil	20	105.26	7	100	7	100	1	100	35	100
2	Wild animal menace on farm field	19	100	7	100	7	100	1	100	34	97.14
3	Frequent incidence of pest and diseases	19	100	7	100	7	100	1	100	34	97.14
4	Inadequacy of irrigation water	19	100	7	100	7	100	1	100	34	97.14
5	High cost of Fertilizers and plant protection chemicals	19	100	7	100	7	100	1	100	34	97.14
6	High rate of interest on credit	18	94.74	7	100	7	100	1	100	33	94.29
7	Low price for the agricultural commodities	1	5.26	0	0	1	14.29	0	0	2	5.71

SUMMARY

In order to assess the socio-economic condition of the farmers in the watershed a comprehensive questionnaire was prepared. Major components such as demographic conditions, migration details, food consumption and family expenditure pattern, material possession, land holding, land use management, cropping pattern, cost of cultivation of crops, livestock management. The statistical components such as frequency and percentage were used to analyze the data. About 35 households located in the micro watershed were interviewed for the survey.

The data on households sampled for socio economic survey indicated that 35 farmers were sampled in Sudapur-2 micro-watershed among them 1 (2.86 %) were landless, 19 (54.29 %) were marginal farmers, 7 (20 %) were small farmers and semi medium farmers and 1 (2.86 %) were medium farmers.

The data indicated that there were 95 (51.91 %) men and 88 (48.09 %) women among the sampled households. The average family size of landless farmers' was 5, marginal farmers' was 5.57, small farmers' was 4.14, semi medium farmers' was 5.57 and medium farmers' was 4.

The data indicated that, 39 (21.31 %) people were in 0-15 years of age, 83 (45.36 %) were in 16-35 years of age, 40 (21.86 %) were in 36-60 years of age and 21 (11.48 %) were above 61 years of age.

The results indicated that Sudapur-2 had 55.19 per cent illiterates, 1.09 per cent Functional Literate, 14.75 per cent of them had primary school, 1.64 per cent of them had middle school, 12.02 per cent of them had high school education, 4.37 per cent of them had PUC, 1.24 per cent of them had Diploma, 0.62 per cent of them had ITI and 5.46 per cent of them had Degree education.

The results indicate that, 68.57 per cent of household heads were practicing agriculture, 25.71 per cent of the household heads were agricultural labourers, 5.71 per cent of the household heads were in Private Service and 2.86 per cent of the household heads were Housewives.

The results indicate that agriculture was the major occupation for 46.99 per cent of the household members, 10.93 per cent were agricultural labourers, 0.55 per cent were General Labour, 1.09 per cent were Government Service, 3.28 per cent were Private Service, 0.55 per cent were Trade & Business, 19.67 per cent were Student, 9.84 per cent were Housewife and 7.10 per cent were children.

The results show that, 100 per cent of the population in the micro watershed has not participated in any institutions. The results indicate that 11.43 per cent of the

households possess thatched house, 77.14 per cent of the households possess katcha house and 11.43 per cent of them possess pucca/RCC house.

The results show that 88.57 per cent of the households possess TV, 20 per cent of the households possess mixer/grinder, 5.71 per cent of the households possess Refrigerator, 14.29 per cent of the households possess Bicycle, 40 per cent of the households possess Motor Cycle and 100 per cent of the households possess mobile phones.

The results show that the average value of television was Rs. 9,000, mixer/grinder was Rs. 2,000, Refrigerator was Rs. 8,500, Bicycle was Rs. 2,000, motor cycle was Rs. 55,500 and mobile phone was Rs. 2,630.

About 14.29 per cent of the households possess bullock cart, 40 per cent of them possess plough, 22.86 per cent of them possess seed/fertilizer drill, 2.86 per cent of them possess Harvester, 17.14 per cent of them possess Sprayer and 34.29 per cent of them possess Weeder.

The results show that the average value of bullock cart was Rs. 28,000, plough was Rs. 1,500, seed/fertilizer drill was Rs. 1,687, Harvester was Rs. 100, sprayer was Rs. 3,750 and weeder was Rs. 30.

The results indicate that, 42.86 per cent of the households possess bullocks, 20 per cent of the households possess local cow and 5.71 per cent of the households possess Buffalo, Goat and Poultry birds.

The results indicate that, average own labour men available in the micro watershed was 1.62, average own labour (women) available was 1.68, average hired labour (men) available was 11.32 and average hired labour (women) available was 8.09.

The results indicate that, 94.29 per cent of the households opined that the hired labour was adequate. The results indicate that, households of the Sudapur-2 micro-watershed possess 32.73 ha (82.87 %) of dry land, 5.59 ha (14.16 %) of irrigated land and 1.17 ha (2.97 %) of Permanent Fallow land. Marginal farmers possess 11.80 ha (96.68 %) of dry land and 0.40 ha (3.32 %) of irrigated land. Small farmers possess 9.07 ha (100 %) of dry land. Semi medium farmers possess 8.09 ha (60.94 %) of dry land and 5.19 ha (39.06 %) of irrigated land. Medium farmers possess 3.76 ha (76.23 %) of dry land and 1.17 ha (23.77 %) of Permanent Fallow land .

The results indicate that, the average value of dry land was Rs. 540,608.39, the average value of irrigated land was Rs. 714,905.94 and the average value of permanent fallow land was Rs. 1,703,448.22. In case of marginal famers, the average land value was Rs. 948,696.85 for dry land and the average value of irrigated land was Rs. 1,976,000. In case of small famers, the average land value was Rs. 396,787.15 for dry land. In case of semi medium famers, the average land value was Rs. 247,000 for dry land and the

average value of irrigated land was Rs. 616,536.66. In case of medium farmers, the average land value was Rs. 239,032.26 for dry land and the average value of permanent fallow land was Rs. 1,703,448.22.

The results indicate that, Bore Well was the major irrigation source in the micro water shed for 14.29 per cent of the farmers. The results indicate that, marginal and semi medium farmers had an irrigated area of 0.40 ha and 5.19 ha.

The results indicate that, farmers have grown Green gram (2.02 ha), cotton (2.86 ha), paddy (4.30 ha), Red gram (26.56 ha) and Sorghum (1.62 ha). Marginal farmers have grown sorghum, red gram and cotton. Small farmers have grown red gram, cotton and green gram. Semi medium farmers have grown cotton, red gram and paddy. Medium farmers have grown red gram and paddy.

The results indicate that, the cropping intensity in Sudapur-2 micro-watershed was found to be 89.51 per cent. The results indicate that, the total cost of cultivation for Cotton was Rs. 35806.62. The gross income realized by the farmers was Rs. 38739.82. The net income from Cotton cultivation was Rs. 2933.20. Thus the benefit cost ratio was found to be 1: 1.08.

The results indicate that, the total cost of cultivation for green gram was Rs. 18525.98. The gross income realized by the farmers was Rs. 49400. The net income from green gram cultivation was Rs. 30874.02. Thus the benefit cost ratio was found to be 1: 2.67.

The results indicate that, the total cost of cultivation for Red gram was Rs. 32506.88. The gross income realized by the farmers was Rs. 62432.47. The net income from Red gram cultivation was Rs. 29925.59. Thus the benefit cost ratio was found to be 1: 1.92.

The results indicate that, the total cost of cultivation for Paddy was Rs. 127772.06. The gross income realized by the farmers was Rs. 189158.35. The net income from Paddy cultivation was Rs. 61386.29. Thus the benefit cost ratio was found to be 1: 1.48.

The results indicate that, 57.14 per cent of the households opined that dry fodder was adequate and 54.29 per cent of the households opined that green fodder was adequate. The results indicate that the annual gross income was Rs. 85,000 for landless farmers, for marginal farmers it was Rs. 121,564.21, for small farmers it was Rs. 126,557.14, semi medium farmers it was Rs. 233,428.57 and medium farmers it was Rs. 80,800.

The results indicate that the average annual expenditure is Rs. 19,493.59. For landless households it was Rs. 55,000, for marginal farmers it was Rs. 9,988.19, for small farmers it was Rs. 23,000, for semi medium farmers it was Rs. 29,500 and medium farmers it was Rs. 70,000.

The results indicate that, sampled households have grown 4 Custard apple trees in their field also 1 Coconut and 13 Custard apple trees in backyard. The results indicate that, households have planted 4 Teak, 39 neem and 4 tamarind in their field and also 2 Tamarind, 2 acacia trees and 11 neem in backyard.

The results indicated that, households have an average investment capacity of Rs. 3,257.14 for land development Rs. 4,285.71 for Irrigation facility, Rs. 2,600 for improved crop production and Rs. 1,714.29 for improved livestock management.

The results indicated that Government subsidy was the source of additional investment for 5.71 per cent for irrigation facility, Loan from bank was the source of additional investment for 8.57 per cent for land development and 5.71 per cent for improved crop production and improved livestock management, Own funds was the source of additional investment for 5.71 per cent for land development, improved crop production and improved livestock management, soft loan was the source of additional investment for 17.14 per cent for land development, 2.86 per cent for irrigation facility and improved livestock management, 11.43 per cent for improved crop production.

The results indicated that, cotton was sold to the extent of 100 per cent, Greengram was sold to the extent of 95.0 per cent, Paddy was sold to the extent of 96.92 per cent, Red gram was sold to the extent of 85.98 per cent and Sorghum to the extent of 88.24 per cent.

The results indicated that, about 100 per cent of the farmers sold their produce to local/village merchants and 5.71 per cent of the farmers sold their produce to Regulated Market. The results indicated that, 100 per cent of the households have used Tractor as a mode of transportation.

The results indicated that, 34.29 per cent of the households have experienced soil and water erosion problems in the farm. The results indicated that, 97.14 per cent have shown interest in soil test. The results indicated that, 85.71 per cent of the households used firewood, 5.71 per cent of the households used Kerosene and 14.29 per cent of the households used LPG as a source of fuel.

The results indicated that, piped supply was the major source of drinking water for 94.29 per cent and 5.71 per cent of the households used bore well in the micro watershed. The results indicated that, Electricity was the major source of light for 100 per cent of the households in micro watershed.

The results indicated that, 48.57 per cent of the households possess sanitary toilet facility. The results indicated that, 100 per cent of the sampled households possessed BPL cards. The results indicated that, 88.57 per cent of the households participated in NREGA programme.

The results indicated that, cereals were adequate for 100 per cent of the households, pulses were adequate for 97.14 per cent of the households, Fruits were adequate for 20 per cent, vegetables were adequate for 91.43 per cent, milk, Egg and meat were adequate for 100 per cent.

The results indicated that, pulses were inadequate for 2.86 per cent of the households, oilseeds were inadequate for 100 per cent, vegetables were inadequate for 11.43 per cent and fruits were inadequate for 80 per cent of the households.

The results indicated that, lower fertility status of the soil was the constraint experienced by 100 per cent of the households,, wild animal menace on farm field, Inadequacy of irrigation water, High cost of Fertilizers and plant protection chemicals and Frequent incidence of pest and diseases was the constraint experienced by 97.14 per cent of the households, High rate of interest on credit (94.29 %) and low price for the agricultural commodities (5.71 %).