



हर कदम, हर डगर
किसानों का हमसफर
भारतीय कृषि अनुसंधान परिषद

Agrisearch with a human touch

**LAND RESOURCE INVENTORY AND SOCIO-ECONOMIC STATUS OF
FARM HOUSEHOLDS FOR WATERSHED PLANNING AND
DEVELOPMENT**

BELAGUNDA-3 (4D5B1N1d) MICROWATERSHED

Yadgir Taluk and District, Karnataka

Karnataka Watershed Development Project – II

SUJALA – III

World Bank funded Project



THE WORLD BANK

**ICAR – NATIONAL BUREAU OF SOIL SURVEY AND LAND USE
PLANNING**

**WATERSHED DEVELOPMENT DEPARTMENT
GOVT. OF KARNATAKA, BANGALORE**



ICAR - NBSS & LUP



About ICAR - NBSS&LUP

The ICAR-National Bureau of Soil Survey and Land Use Planning (ICAR-NBSS&LUP), Nagpur, a premier Institute of the Indian Council of Agricultural Research (ICAR), was set up during 1976 with the objective to prepare soil resource maps at national, state and district levels and to provide research inputs in soil resource mapping and its applications, land evaluation, land use planning, land resource management, and database management using GIS for optimising land use on different kinds of soils in the country.

The Bureau has been engaged in carrying out soil resource survey, agro-ecological and soil degradation mapping at the country, state and district levels for qualitative assessment and monitoring the soil health towards viable land use planning. The research activities have resulted in identifying the soil potentials and problems, and the various applications of the soil surveys with the ultimate objective of sustainable agricultural development. The Bureau has the mandate to correlate and classify soils of the country and maintain a National Register of all the established soil series. The Institute is also imparting in-service training to staff of the soil survey agencies in the area of soil survey, land evaluation and soil survey interpretations for land use planning. The Bureau in collaboration with Panjabrao Krishi Vidyapeeth, Akola is running post-graduate teaching and research programme in land resource management, leading to M.Sc. and Ph.D. degrees.

Citation: Rajendra Hegde, Ramesh Kumar, S.C., B.A. Dhanorkar, S. Srinivas, M. Lalitha, K.V. Niranjana, R.S. Reddy and S.K. Singh (2019). "Land resource inventory and socio-economic status of farm households for watershed planning and development of Belagunda-3 (4D5B1N1d) Microwatershed, Yadgir Taluk and District, Karnataka", ICAR-NBSS&LUP Sujala MWS Publ.396, ICAR – NBSS LUP, RC, Bangalore. p.121 & 33.

TO OBTAIN COPIES,

Please write to:

Director, ICAR - NBSS & LUP,

Amaravati Road, NAGPUR - 440 033, India

Phone : (0712) 2500386, 2500664, 2500545 (O)

Telefax : 0712-2522534

E-Mail : director@nbsslup.ernet.in

Website URL : nbsslup.in

Or

Head, Regional Centre, ICAR - NBSS&LUP, Hebbal, Bangalore - 560 024

Phone : (080) 23412242, 23510350 (O)

Telefax : 080-23510350

E-Mail : nbssrcb@gmail.com



**LAND RESOURCE INVENTORY AND SOCIO-ECONOMIC
STATUS OF FARM HOUSEHOLDS FOR WATERSHED
PLANNING AND DEVELOPMENT**

BELAGUNDA-3 (4D5B1N1d) MICROWATERSHED

Yadgir Taluk and District, Karnataka

Karnataka Watershed Development Project – II

Sujala-III

World Bank funded Project



**ICAR – NATIONAL BUREAU OF SOIL SURVEY AND LAND
USE PLANNING**



ICAR - NBSS & LUP



**WATERSHED DEVELOPMENT DEPARTMENT, GOVT. OF
KARNATAKA, BANGALORE**



PREFACE

In Karnataka, as in other Indian States, the livelihoods of rural people are intertwined with farming pursuits. The challenges in agriculture are seriously threatening the livelihood of a large number of farmers as they have been practicing farming in contextual factors beyond their control. Climatic factors are the most important ones and have become much more significant in recent times due to rapid climate changes induced by intensive anthropogenic activities affecting our ecosystem in multiple ways. Climate change has become the reality, it is happening and efforts to evolve and demonstrate climate resilient technologies have become essential. Due to the already over stressed scenario of agrarian sector, the climate change is resulting in manifold increase in the complexities, pushing the rural mass to face more and more unpredictable situations. The rising temperatures and unpredictable rainfall patterns are going to test seriously the informed decisions farmers have to make in order to survive in farming and sustain their livelihood.

It is generally recognized that impacts of climate change shall not be uniform across the globe. It is said that impact of climate change is more severe in South Asia. Based on the analysis of meteorological data, it is predicted that in India, there will be upward trend in mean temperature, downward trend in relative humidity, annual rainfall and number of wet days in a year. Also, in general, phenomena like erratic monsoon, spread of tropical diseases, rise in sea levels, changes in availability of fresh water, frequent floods, droughts, heat waves, storms and hurricanes are predicted. Each one of these adverse situations are already being experienced in various parts of India and also at the global level. Decline in agricultural productivity of small and marginal farmers becoming more vulnerable is already witnessed.

In Karnataka, more than 60 per cent of the population live in rural areas and depend on agriculture and allied activities for their livelihood. Though the state has achieved significant progress in increasing the yield of many crops, there is tremendous pressure on the land resources due to the growing and competing demands of various land uses. This is reflected in the alarming rate of land degradation observed. Already more than 50 per cent of the area is affected by various forms of degradation. If this trend continues, the sustainability of the fragile ecosystem will be badly affected. The adverse effects of change in the climatic factors are putting additional stress on the land resources and the farmers dependent on this.

The natural resources (land, water and vegetation) of the state need adequate and constant care and management, backed by site-specific technological interventions and investments particularly by the government. Detailed database pertaining to the nature of the land resources, their constraints, inherent potentials and suitability for various land

based rural enterprises, crops and other uses is a prerequisite for preparing location-specific action plans, which are in tune with the inherent capability of the resources. Any effort to evolve climate resilient technologies has to be based on the baseline scientific database. Then only one can expect effective implementation of climate resilient technologies, monitor the progress, make essential review of the strategy, and finally evaluate the effectiveness of the implemented programs. The information available at present on the land resources of the state are of general nature and useful only for general purpose planning. Since the need of the hour is to have site-specific information suitable for farm level planning and detailed characterization and delineation of the existing land resources of an area into similar management units is the only option.

ICAR-NBSS&LUP, Regional Centre, Bangalore has taken up a project sponsored by the Karnataka Watershed Development Project-II, (Sujala-III), Government of Karnataka funded by the World Bank under Component -1 Land Resource Inventory. This study was taken up to demonstrate the utility of such a database in reviewing, monitoring and evaluating all the land based watershed development programs on a scientific footing. To meet the requirements of various land use planners at grassroots level, the present study on “Land Resource Inventory and Socio-Economic Status of Farm Households for Watershed Planning and Development of for Belagunda-3 microwatershed in Yadgir Taluk and District, Karnataka” for integrated development was taken up in collaboration with the State Agricultural Universities, IISC, KRSAC, KSNDMC as Consortia partners. The project provides detailed land resource information at cadastral level (1:7920 scale) for all the plots and socio-economic status of farm households covering thirty per cent farmers randomly selected representing landed and landless class of farmers in the micro-watershed. The project report with the accompanying maps for the microwatershed will provide required detailed database for evolving effective land use plan, alternative land use options and conservation plans for the planners, administrators, agricultural extension personnel, KVK officials, developmental departments and other land users to manage the land resources in a sustainable manner.

It is hoped that this database will be useful to the planners, administrators and developmental agencies working in the area in not only for formulating location specific developmental schemes but also for their effective monitoring at the village/watershed level.

Nagpur

Date:05-10-2019

S.K. SINGH

Director, ICAR - NBSS&LUP Nagpur

Contributors

Dr. Rajendra Hegde Principal Scientist, Head & Project Leader, Sujala-III Project ICAR-NBSS&LUP, Regional Centre, Bangalore	Dr. S.K.Singh Director, ICAR-NBSS&LUP Coordinator, Sujala-III Project Nagpur
Soil Survey, Mapping & Report Preparation	
Dr. B.A. Dhanorkar	Sh. R.S. Reddy
Dr. K.V. Niranjana	Mr. Somashekar T N
	Smt. Chaitra, S.P.
	Dr. Gopali bardhan
	Ms. Arpitha
	Dr. Mahendra Kumar, M.B.
Field Work	
Sh. C.BacheGowda	Sh. Mahesh, D.B.
Sh. Somashekar	Sh. Ashok S Sindagi
Sh. M. Jayaramaiah	Sh. Veerabhadrapa B.
Sh. Paramesha, K.	Sh. Kailas
Sh. B. M. Narayana Reddy	Sh. Anand
	Sh. Arun N Kambar.
	Sh Kamalesh Awate
	Sh. Sharaan Kumar Huppar
	Sh. Yogesh H.N.
	Sh. Kalaveerachari R Kammar
GIS Work	
Dr. S.Srinivas	Sh. A.G.Devendra Prasad
Sh. D.H.Venkatesh	Sh. Prakashanaik, M.K.
Smt.K.Sujatha	Sh. Abhijith Sastry, N.S.
Smt. K.V.Archana	Sh. Sudip Kumar Suklabaidya
Sh. N. Maddileti	Sh. Avinash, K.N.
	Sh. Amar Suputhra, S
	Sh. Deepak, M.J.
	Smt. K.Karunya Lakshmi
	Ms. Seema, K.V.
	Ms. A. Rajab Nisha

Laboratory Analysis	
Dr. K.M.Nair	Ms. Steffi Peter
Smt. Arti Koyal	Ms. Thara, V.R
Smt. Parvathy	Ms. Roopa, G.
	Ms. Swati, H.
	Sh. Shantaveera Swami
	Ms. Shwetha, N.K.
	Smt. Ishrat Haji
	Ms. P. Pavan Kumari
	Ms. Padmaja
	Ms. Veena, M.
Socio-Economic Analysis	
Dr. S.C. Ramesh Kumar	Sh. M.K. Prakashanaik,
	Ms. Karuna V. Kulkarni,
	Mrs. Sowmya A.N,
	Sh. Vinod R,
	Sh. Basavaraja,
	Sh. Vijay Kumar Lamani,
	Ms. Sowmya K.B.,
	Mrs. Prathibha, D.G,
	Sh. Rajendra,D,
Soil & Water Conservation	
Sh. Sunil P. Maske	
Watershed Development Department, GoK, Bangalore	
Sh. Rajeev Ranjan IFS Project Director & Commissioner, WDD	Dr. A. Natarajan NRM Consultant, Sujala-III Project
Dr. S.D. Pathak IFS Executive Director & Chief Conservator of Forests, WDD	

PART-A

LAND RESOURCE INVENTORY

Contents

Preface		
Contributors		
Executive Summary		
Chapter 1	Introduction	1
Chapter 2	Geographical Setting	3
2.1	Location and Extent	3
2.2	Geology	3
2.3	Physiography	4
2.4	Drainage	4
2.5	Climate	4
2.6	Natural Vegetation	6
2.7	Land Utilization	6
Chapter 3	Survey Methodology	9
3.1	Base maps	9
3.2	Image Interpretation for Physiography	9
3.3	Field Investigation	12
3.4	Soil Mapping	13
3.4	Land Management Units	13
3.5	Laboratory Characterization	14
Chapter 4	The Soils	19
4.1	Soils of granite gneiss landscape	19
Chapter 5	Interpretation for Land Resource Management	27
5.1	Land Capability Classification	27
5.2	Soil Depth	29
5.3	Surface Soil Texture	30
5.4	Soil Gravelliness	31
5.5	Available Water Capacity	32
5.6	Soil Slope	33
5.7	Soil Erosion	34
Chapter 6	Fertility Status	37
6.1	Soil Reaction (pH)	37
6.2	Electrical Conductivity (EC)	37
6.3	Organic Carbon (OC)	37
6.4	Available Phosphorus	39
6.5	Available Potassium	39
6.6	Available Sulphur	39
6.7	Available Boron	40
6.8	Available Iron	40
6.9	Available Manganese	40
6.10	Available Copper	40
6.11	Available Zinc	44

Chapter 7	Land Suitability for Major Crops	45
7.1	Land suitability for Sorghum	45
7.2	Land suitability for Maize	46
7.3	Land suitability for Bajra	47
7.4	Land suitability for Groundnut	48
7.5	Land suitability for Sunflower	49
7.6	Land suitability for Redgram	50
7.7	Land suitability for Bengal gram	51
7.8	Land suitability for Cotton	52
7.9	Land suitability for Chilli	53
7.10	Land suitability for Tomato	54
7.11	Land suitability for Brinjal	55
7.12	Land suitability for Onion	56
7.13	Land suitability for Bhendi	57
7.14	Land suitability for Drumstick	58
7.15	Land suitability for Mango	59
7.16	Land suitability for Guava	60
7.17	Land suitability for Sapota	61
7.18	Land Suitability for Pomegranate	62
7.19	Land Suitability for Musambi	63
7.20	Land Suitability for Lime	64
7.21	Land Suitability for Amla	65
7.22	Land Suitability for Cashew	66
7.23	Land Suitability for Jackfruit	67
7.24	Land Suitability for Jamun	68
7.25	Land Suitability for Custard apple	69
7.26	Land Suitability for Tamarind	70
7.27	Land Suitability for Mulberry	71
7.28	Land Suitability for Marigold	72
7.29	Land Suitability for Chrysanthemum	73
7.30	Land Management Units	105
7.31	Proposed Crop Plan	105
Chapter 8	Soil Health Management	107
Chapter 9	Soil and Water conservation Treatment Plan	113
9.1	Treatment Plan	114
9.2	Recommended Soil and Water Conservation measures	117
9.3	Greening of Microwatershed	118
	References	121
	Appendix I	I-III
	Appendix II	V-VII
	Appendix III	IX-XI

LIST OF TABLES

2.1	Mean Monthly Rainfall, PET, 1/2 PET at Yadgir Taluk & District	5
2.2	Land Utilization in Yadgir district	7
3.1	Differentiating Characteristics used for Identifying Soil Series	13
3.2	Soil map unit description of Belagunda 3 Microwatershed	14
4.1	Physical and Chemical Characteristics of Soil Series identified in Belagunda 3 microwatershed	23
7.1	Soil-Site Characteristics of Belagunda 3 Microwatershed	75
7.2	Land suitability criteria for Sorghum	76
7.3	Land suitability criteria for Maize	77
7.4	Land suitability criteria for Bajra	78
7.5	Land suitability criteria for Groundnut	79
7.6	Land suitability criteria for Sunflower	80
7.7	Land suitability criteria for Redgram	81
7.8	Land suitability criteria for Bengal gram	82
7.9	Land suitability criteria for Cotton	83
7.10	Land suitability criteria for Chilli	84
7.11	Land suitability criteria for Tomato	85
7.12	Land suitability criteria for Brinjal	86
7.13	Land suitability criteria for Onion	87
7.14	Land suitability criteria for Bhendi	88
7.15	Land suitability criteria for Drumstick	89
7.16	Land suitability criteria for Mango	90
7.17	Land suitability criteria for Guava	91
7.18	Land suitability criteria for Sapota	92
7.19	Land suitability criteria for Pomegranate	93
7.20	Land suitability criteria for Musambi	94
7.21	Land suitability criteria for Lime	95
7.22	Land suitability criteria for Amla	96
7.23	Land suitability criteria for Cashew	97
7.24	Land suitability criteria for Jackfruit	98
7.25	Land suitability criteria for Jamun	99
7.26	Land suitability criteria for Custard apple	100
7.27	Land suitability criteria for Tamarind	101

7.28	Land suitability criteria for Mulberry	102
7.29	Land suitability criteria for Marigold	103
7.30	Land suitability criteria for Chrysanthemum	104
7.31	Proposed Crop Plan for Belagunda 3 Microwatershed	106

LIST OF FIGURES

2.1	Location map of Belagunda 3 Microwatershed	3
2.2 a	Granite and granite gneiss rock formation	4
2.3	Rainfall distribution in Yadgir Taluk & District	5
2.4	Natural vegetation of Belagunda-3 microwatershed	6
2.5	Current Land use map of Belagunda 3 Microwatershed	7
2.6 a & b	Different crops and cropping systems in Belagunda 3 Microwatershed	7
3.1	Scanned and Digitized Cadastral map of Belagunda 3 Microwatershed	10
3.2	Satellite image of Belagunda 3 Microwatershed	11
3.3	Cadastral map overlaid on IRS PAN+LISS IV merged imagery of Belagunda 3 Microwatershed	11
3.4	Location of profiles in a transect	12
3.5	Soil phase or management units of Belagunda 3 Microwatershed	17
5.1	Land Capability Classification map of Belagunda 3 Microwatershed	29
5.2	Soil Depth map of Belagunda 3 Microwatershed	30
5.3	Surface Soil Texture map of Belagunda 3 Microwatershed	31
5.4	Soil Gravelliness map of Belagunda 3 Microwatershed	32
5.5	Soil Available Water Capacity map of Belagunda 3 Microwatershed	33
5.6	Soil Slope map of Belagunda 3 Microwatershed	34
5.7	Soil Erosion map of Belagunda 3 Microwatershed	35
6.1	Soil Reaction (pH) map of Belagunda 3 Microwatershed	38
6.2	Electrical Conductivity (EC) map of Belagunda 3 Microwatershed	38
6.3	Soil Organic Carbon (OC) map of Belagunda 3 Microwatershed	39
6.4	Soil Available Phosphorus map of Belagunda 3 Microwatershed	40
6.5	Soil Available Potassium map of Belagunda 3 Microwatershed	41
6.6	Soil Available Sulphur map of Belagunda 3 Microwatershed	41
6.7	Soil Available Boron map of Belagunda 3 Microwatershed	42
6.8	Soil Available Iron map of Belagunda 3 Microwatershed	42
6.9	Soil Available Manganese map of Belagunda 3 Microwatershed	43
6.10	Soil Available Copper map of Belagunda 3 Microwatershed	43
6.11	Soil Available Zinc map of Belagunda 3 Microwatershed	44
7.1	Land suitability for Sorghum	46
7.2	Land suitability for Maize	47

7.3	Land suitability for Bajra	48
7.4	Land suitability for Groundnut	49
7.5	Land suitability for Sunflower	50
7.6	Land suitability for Redgram	51
7.7	Land suitability for Bengal gram	52
7.8	Land suitability for Cotton	53
7.9	Land suitability for Chilli	54
7.10	Land suitability for Tomato	55
7.11	Land suitability for Brinjal	56
7.12	Land suitability for Onion	57
7.13	Land suitability for Bhendi	58
7.14	Land suitable for Drumstick	59
7.15	Land suitability for Mango	60
7.16	Land suitability for Guava	61
7.17	Land suitability for Sapota	62
7.18	Land suitability for Pomegranate	63
7.19	Land suitability for Musambi	64
7.20	Land suitability for Lime	65
7.21	Land suitability for Amla	66
7.22	Land suitability for Cashew	67
7.23	Land suitability for Jackfruit	68
7.24	Land suitability for Jamun	69
7.25	Land suitability for Custard apple	70
7.26	Land suitability for Tamarind	71
7.27	Land suitability for Mulberry	72
7.28	Land suitability for Marigold	73
7.29	Land suitability for Chrysanthemum	74
7.30	Land management units map of Belagunda 3 Microwatershed	105
9.1	Soil and water conservation map of Belagunda 3 Microwatershed	118

EXECUTIVE SUMMARY

The land resource inventory of Belagunda-3 Microwatershed was conducted using village cadastral maps and IRS satellite imagery on 1:7920 scale. The false colour composites of IRS imagery were interpreted for physiography and the physiographic delineations were used as base for mapping soils. The soils were studied in several transects and a soil map was prepared with phases of soil series as mapping units. Random checks were made all over the area outside the transects to confirm and validate the soil map unit boundaries. The soil map shows the geographic distribution and extent, characteristics, classification, behavior and use potentials of the soils in the microwatershed.

The present study covers an area of 485 ha in Yadgir taluk & district, Karnataka. The climate is semiarid and categorized as drought-prone with an average annual rainfall of 866 mm, of which about 652 mm is received during south-west monsoon, 138 mm during north-east and the remaining 76 mm during the rest of the year. An area of 415 ha in the microwatershed is covered by soils and 71 ha by others (habitation and water body). The salient findings from the land resource inventory are summarized briefly below.

- ❖ The soils belong to 5 soil series and 5 soil phases (management units) and 2 land management units.*
- ❖ The length of crop growing period is about 120-150 days starting from 1st week of June to 4th week of October.*
- ❖ From the master soil map, several interpretative and thematic maps like land capability, soil depth, surface soil texture, soil gravelliness, available water capacity, soil slope and soil erosion were generated.*
- ❖ Soil fertility status maps for macro and micronutrients were generated based on the surface soil samples collected at every 320 m grid interval.*
- ❖ Land suitability for growing 29 major agricultural and horticultural crops was assessed and maps showing the degree of suitability along with constraints were generated.*
- ❖ Entire area in the microwatershed is suitable for agriculture.*
- ❖ About 19 per cent area are deep (100 - 150 cm) and 67 per cent area of the microwatershed has soils that are very deep (>150 cm).*
- ❖ About 5 per cent area in the microwatershed has loamy and 80 per cent clayey soils at the surface.*
- ❖ Entire area of the microwatershed is non gravelly (<15%).*
- ❖ About 80 per cent area of the microwatershed is very high (>200 mm/m) in available water capacity and 5 per cent area medium (101-150 mm/m) in available water capacity.*

- ❖ *Maximum of 50 per cent area in the microwatershed is nearly level (0-1%) and 35 per cent area is very gently sloping (1-3%) lands.*
- ❖ *An area of about 69 per cent in the microwatershed is slightly (e1) eroded and 17 per cent area is moderately (e2) eroded lands.*
- ❖ *An area of about 23 per cent is moderately alkaline (pH 7.8-8.4), 58 per cent is strongly alkaline (pH 8.4- 9.0) and 4 per cent is very strongly alkaline (pH > 9.0) in reaction*
- ❖ *The electrical conductivity is non saline (<2 dsm⁻¹) in a maximum area of about 71 per cent, 8 per cent is low (2 – 4 dsm⁻¹) and 6 per cent is medium (4 – 8 dsm⁻¹) in the microwatershed.*
- ❖ *33 per cent area of the microwatershed is low (<0.5%) and 52 per cent is medium (0.5 – 0.75 %) in organic carbon.*
- ❖ *26 per cent area is high (>57 kg/ha) in available phosphorus, 28 per area is medium (23-57 kg/ha) and 31per area is low (<23 kg/ha).*
- ❖ *About 60 per cent is medium (145-337 kg/ha) in available potassium and 25 per cent is high (>337 kg/ha).*
- ❖ *Available sulphur is low (<10 ppm) in an area of about <1 per cent, medium (10 - 20 ppm) in 6 per cent and 79 per cent is high (>20 ppm).*
- ❖ *About <1 per cent area is low (<0.5 ppm) in available boron, 60 per cent is medium (0.5-1.0 ppm) and 26 per cent is high (>1.0 ppm).*
- ❖ *About 28 per cent area is deficient (<4.5 ppm) in available iron and 57 per cent is sufficient (>4.5 ppm).*
- ❖ *Available manganese and copper are sufficient in all the soils of the microwatershed.*
- ❖ *Available zinc is deficient (<0.6 ppm) in all the soils of the microwatershed.*
- ❖ *The land suitability for 29 major crops grown in the microwatershed were assessed and the areas that are highly suitable (S1) and moderately suitable (S2) are given below. It is however to be noted that a given soil may be suitable for various crops but what specific crop to be grown may be decided by the farmer looking to his capacity to invest on various inputs, marketing infrastructure, market price and finally the demand and supply position.*

Land suitability for various crops in the Microwatershed

Crop	Suitability Area in ha (%)		Crop	Suitability Area in ha (%)	
	Highly suitable (S1)	Moderately suitable (S2)		Highly suitable (S1)	Moderately suitable (S2)
<i>Sorghum</i>	-	269(55)	<i>Guava</i>	25(5)	-
<i>Maize</i>	25(5)	244(50)	<i>Sapota</i>	25(5)	-
<i>Bajra</i>	25(5)	244(50)	<i>Pomegranate</i>	25(5)	-
<i>Groundnut</i>	25(5)	244(50)	<i>Musambi</i>	25(5)	-
<i>Sunflower</i>	-	25(5)	<i>Lime</i>	25(5)	-
<i>Redgram</i>	-	269(55)	<i>Amla</i>	25(5)	-
<i>Bengal gram</i>	-	-	<i>Cashew</i>	-	25(5)
<i>Cotton</i>	-	-	<i>Jackfruit</i>	25(5)	-
<i>Chilli</i>	25(5)	-	<i>Jamun</i>	25(5)	-
<i>Tomato</i>	25(5)	-	<i>Custard apple</i>	25(5)	-
<i>Brinjal</i>	25(5)	-	<i>Tamarind</i>	25(5)	-
<i>Onion</i>	25(5)	-	<i>Mulberry</i>	25(5)	-
<i>Bhendi</i>	25(5)	-	<i>Marigold</i>	25(5)	-
<i>Drumstick</i>	25(5)	-	<i>Chrysanthemum</i>	25(5)	-
<i>Mango</i>	25(5)	-			

- ❖ *Apart from the individual crop suitability, a proposed crop plan has been prepared for the identified LMUs by considering only the highly and moderately suitable lands for different crops and cropping systems with food, fodder, fibre and other horticulture crops.*
- ❖ *Maintaining soil-health is vital for crop production and conserve soil and land resource base for maintaining ecological balance and to mitigate climate change. For this, several ameliorative measures have been suggested for these problematic soils like saline/alkali, highly eroded, sandy soils etc.,*
- ❖ *Soil and water conservation treatment plan has been prepared that would help in identifying the sites to be treated and also the type of structures required.*
- ❖ *As part of the greening programme, several tree species have been suggested to be planted in marginal and submarginal lands, field bunds and also in the hillocks, mounds and ridges. This would help in not only supplementing the farm income but also provide fodder and fuel and generate lot of biomass which would help in maintaining an ecological balance and also contribute to mitigating the climate change.*

INTRODUCTION

Land is a scarce resource and basic unit for any material production. It can support the needs of the growing population, provided they use the land in a rational and judicious manner. But what is happening in many areas of the state is a cause for concern to everyone involved in the management of land resources at the grassroots level. The area available for agriculture is about 51 per cent of the total geographical area and more than 60 per cent of the people are still dependant on agriculture for their livelihood. The limited land area is under severe stress and strain due to increasing population pressure and competing demands of various land uses. Due to this, every year there is significant diversion of farm lands and water resources for non-agricultural purposes. Apart from this, due to lack of interest in farmers for farming, large tracts of cultivable lands are turning into fallows in many areas and this trend is continuing at an alarming rate.

Further, land degradation has emerged as a serious problem which has already affected about 38 lakh ha of cultivated area in the state. Soil erosion alone has degraded about 35 lakh ha. Almost all the uncultivated areas are facing various degrees of degradation, particularly soil erosion. Salinity and alkalinity has emerged as a major problem in more than 3.5 lakh ha in the irrigated areas of the state. Nutrient depletion and declining factor productivity is common in both rainfed and irrigated areas. The degradation is continuing at an alarming rate and there appears to be no systematic effort among the stakeholders to contain this process. In recent times, an aberration of weather due to climate change phenomenon has added another dimension leading to unpredictable situations to be tackled by the farmers.

In this critical juncture, the challenge before us is not only to increase the productivity per unit area which is steadily declining and showing a fatigue syndrome, but also to prevent or at least reduce the severity of degradation. If the situation is not reversed at the earliest, then the sustainability of the already fragile crop production system and the overall ecosystem will be badly affected in the state. The continued neglect and unscientific use of the resources for a long time has led to the situation observed at present in the state. It is a known fact and established beyond doubt by many studies in the past that the cause for all kinds of degradation is the neglect and irrational use of the land resources. Hence, there is an urgent need to generate a detailed site-specific farm level database on various land resources for all the villages/watersheds in a time bound manner that would help to protect the valuable soil and land resources and also to stabilize the farm production.

Therefore, the land resource inventory required for farm level planning is the one which investigates not only the surface but also consider the other parameters which are critical for productivity *viz.*, soils, climate, water, minerals and rocks, topography, geology, hydrology, vegetation, crops, land use pattern, animal population, socio-

economic conditions, infrastructure, marketing facilities and various schemes and developmental works of the government etc. From the data collected at farm level, the specific problems and potentials of the area can be identified and highlighted, conservation measures required for the area can be planned on a scientific footing, suitability of the area for various uses can be worked out and finally viable and sustainable land use options suitable for each and every land holding can be prescribed.

The Land Resource Inventory is basically done for identifying the potential and problem areas, developing sustainable land use plans, estimation of surface run off and water harvesting potential, preparation of soil and water conservation plans, land degradation/desertification etc. The Bureau is presently engaged in developing an LRI methodology using high resolution satellite remote sensing data and Digital Elevation Model (DEM) data to prepare Landscape Ecological Units (LEU) map representing agro-ecosystem as a whole. The LEU is preferred over landform as the base map for LRI. LEU is the assemblage of landform, slope and land use. An attempt has already been made to upscale the soil resource information from 1:250000 and 1:50000 scale to the LEU map in Goa and other states.

The land resource inventory aims to provide site-specific database for Belagunda-3 microwatershed in Yadgir Taluk & District, Karnataka State for the Karnataka Watershed Development Department. The database was generated by using cadastral map of the village as a base along with high resolution IRS LISS IV and Cartosat-1 merged satellite imagery. Later, an attempt will be made to uplink this LRI data generated at 1:7920 scale under Sujala-III Project to the proposed Landscape Ecological Units (LEUs) map.

The study was organized and executed by the ICAR- National Bureau of Soil Survey and Land Use Planning, Regional Centre, Bangalore under Generation of Land Resource Inventory Data Base Component-1 of the Sujala-III Project funded by the World Bank.

GEOGRAPHICAL SETTING

2.1 Location and Extent

The Belagunda-3 microwatershed is located in the northern part of Karnataka in Yadgir Taluk and District, Karnataka State (Fig.2.1). It comprises parts of Belagundi & Anura. B villages. It lies between $16^{\circ} 32'$ – $16^{\circ} 33'$ north latitudes and $77^{\circ} 10'$ – $77^{\circ} 12'$ east longitudes, covering an area of about 484.87 ha. It is about 28 km southeast of Yadgir town and is surrounded by Belagundi on the southeast, Munagala on the south and Anura. B on the north and western side of the microwatershed.




Fig.2.1 Location map of Belagunda-3 microwatershed

2.2 Geology

Major rock formations observed in the microwatershed are granite gneiss (Figs.2.2). Granite gneisses are essentially pink to gray and are coarse to medium grained. They consist primarily of quartz, feldspar, biotite and hornblende. The gray granite gneisses are highly weathered, fractured and fissured upto a depth of about 10 m. Dolerite dykes and

quartz veins are common with variable width and found to occur in Belagunda-3 microwatershed.



Fig.2.2 Granite and granite gneiss rocks formation

2.3 Physiography

Physiographically, the area has been identified as granite gneiss landscape. The area has been further subdivided into five landforms, *viz;* mounds/ridges, summits, side slopes and very gently sloping uplands, plains and valleys based on slope and its relief features. The elevation ranges from 352-361 m above MSL. The mounds and ridges are mostly covered by rock outcrops.

2.4 Drainage

The area is drained by several parallel streams like Bori, Amerja and Kanga which finally join the river Bhima along its course. Though, they are not perennial, during rainy season they carry large quantities of rain water. The microwatershed has only few small tanks which are not capable of storing the water that flows during the rainy season. Due to this, the ground water recharge is very much affected. This is reflected in the failure of many bore wells in the villages. If the available rain water is properly harnessed by constructing new tanks and recharge structures at appropriate places in the villages, then the drinking and irrigation needs of the area can be easily met. The drainage network is parallel to sub parallel and dendritic.

2.5 Climate

The Yadgir district lies in the northern plains of Karnataka and falls under semiarid tract of the state and is categorized as drought- prone with total annual rainfall of 866 mm (Table 2.1). Of the total rainfall, maximum of 652 mm is received during the south–west monsoon period from June to September, the north-east monsoon from October to early December contributes about 138 mm and the remaining 76 mm during

the rest of the year. The summer season starts during the middle of February and continues up to the first week of June. The period from December to the middle of February is the coldest season. December is the coldest month with mean daily maximum and minimum temperatures being 29.5⁰C and 10⁰C respectively. During peak summer, temperature shoots up to 45⁰C. Relative humidity varies from 26% in summer to 62% in winter. Rainfall distribution is shown in Figure 2.3. The average Potential Evapo-Transpiration (PET) is 141 mm and varies from a low of 81 mm in December to 199 mm in the month of May. The PET is always higher than precipitation in all the months except end of June to end of September. Generally, the Length of crop Growing Period (LGP) is 120-150 days and starts from 1st week of June to 4th week of October.

Table 2.1 Mean Monthly Rainfall, PET, 1/2 PET at Yadgir Taluk, Yadgir District

Sl. No.	Months	Rainfall	PET	1/2 PET
1	January	4.30	86.0	43.0
2	February	2.30	125.5	62.7
3	March	15.10	166.0	83.0
4	April	18.50	179.8	89.9
5	May	36.0	198.8	97.9
6	June	118.0	175.1	87.5
7	July	171.80	156.3	78.1
8	August	182.9	150.3	75.1
9	September	179.7	142.0	71.0
10	October	105.3	138.5	69.2
11	November	26.4	97.60	48.6
12	December	6.0	80.90	40.4
Total		866.3		




Fig 2.3 Rainfall distribution in Yadgir Taluk and District

2.6 Natural Vegetation

The natural vegetation is sparse comprising few tree species, shrubs and herbs. The mounds, ridges and boulders occupy very sizeable area which is under thin to moderately thick forest vegetation. Still, there are some remnants of the past forest cover which can be seen in patches in some ridges and hillocks in the microwatershed (Fig 2.4).

Apart from the continuing deforestation, the presence of large population of goats, sheep and other cattle in the microwatershed is causing vegetative degradation of whatever little vegetation left in the area. The uncontrolled grazing has left no time for the regeneration of the vegetative cover. This leads to the accelerated rate of erosion on the hill slopes resulting in the formation of deep gullies in the foot slopes that eventually result in the heavy siltation of tanks and reservoirs in the microwatershed.



Fig 2.4 Natural vegetation of Belagunda-3 microwatershed

2.7 Land Utilization

About 72 per cent area (Table 2.2) in Yadgir district is cultivated at present. An area of about 2 per cent is permanently under pasture, 20 per cent under current fallows and 6 per cent under non-agricultural land and 5 per cent under currently barren. Forests occupy an area of about 7 per cent and the tree cover is in a very poor state. Most of the mounds, ridges and bouldery areas have very poor vegetative cover. Major crops grown in the area are sorghum, maize, cotton, sunflower, groundnut, red gram, mango, pomegranate, marigold and sapota. While carrying out land resource inventory, the land use/land cover particulars are collected from all the survey numbers and a current land use map of the microwatershed is prepared. The current land use map prepared shows the arable and non-arable lands, other land uses and different types of crops grown in the area. The current land use map of Belagunda-3 microwatershed is presented in Fig.2.5. The different crops and cropping systems adopted in the microwatershed is presented in Figures 2.6.

Table 2.2 Land Utilization in Yadgir District

Sl. No.	Agricultural land use	Area (ha)	Per cent
1	Total geographical area	516088	-
2	Total cultivated area	373617	72.4
3	Area sown more than once	74081	14.3
4	Cropping intensity	-	119.8
5	Trees and grooves	737	0.14
6	Forest	33773	6.54
7	Cultivable wasteland	2385	0.46
8	Permanent Pasture land	11755	2.28
9	Barren land	27954	5.41
10	Non- Agriculture land	29623	5.73
11	Current Fallows	105212	20.4





Fig.2.5 Current Land Use map of Belagunda-3 microwatershed



Fig. 2.6. Different Crops and Cropping Systems in Belagunda-3 microwatershed



Sunflower




Groundnut



Mango



Pomegranate



Cotton



Sorghum

Fig. 2.6. Different Crops and Cropping Systems in Belagunda-3 microwatershed

SURVEY METHODOLOGY

The purpose of land resource inventory is to delineate similar areas (soil series and phases), which respond or expected to respond similarly to a given level of management. This was achieved in Belagunda-3 microwatershed by the detailed study of all the soil characteristics (depth, texture, colour, structure, consistence, coarse fragments, porosity, soil reaction, soil horizons etc.) and site characteristics (slope of the land, erosion, drainage, occurrence of rock fragments etc.) followed by grouping of similar areas based on soil-site characteristics into homogeneous (management units) units, and showing the area extent and their geographic distribution on the microwatershed cadastral map. The detailed survey at 1:7920 scale was carried out in an area of 489 ha. The methodology followed for carrying out land resource inventory was as per the guidelines given in Soil Survey Manual (IARI, 1971; Soil Survey Staff, 2006; Natarajan *et al.*, 2015) which is briefly described below.

3.1 Base Maps

The detailed survey of the land resources occurring in the microwatershed was carried out by using digitized cadastral map and IRS satellite imagery as base supplied by KRSAC. The cadastral map shows field boundaries with their survey numbers, location of tanks, streams and other permanent features of the area (Fig. 3.1). Apart from the cadastral map, remote sensing data products from Cartosat-1 and LISS IV merged at the scale of 1:7920 were used in conjunction with the cadastral map to identify the landscapes, landforms and other surface features. The imagery helped in the identification and delineation of boundaries between hills, uplands and lowlands, water bodies, forest and vegetated areas, roads, habitations and other cultural features of the area (Fig. 3.2). The cadastral map was overlaid on the satellite imagery (Fig. 3.3) that helps to identify the parcel boundaries and other permanent features. Apart from cadastral maps and images, toposheets of the area (1:50,000 scale) were also used for initial traversing, identification of geology and landforms, drainage features, present land use and also for selection of transects in the microwatershed.

3.2 Image Interpretation for Physiography

False Colour Composites (FCCs) of Cartosat-I and LISS-IV merged satellite data covering microwatershed area was visually interpreted using image interpretation elements and all the available collateral data with local knowledge. The delineated physiographic boundaries were transferred on to a cadastral map overlaid on satellite imagery. Physiographically, the area has been identified as granite gneiss landscape. It was divided into five landforms, *viz.*; ridges and mounds, gently and very gently sloping uplands and lowlands based on slope and image characteristics. They were further subdivided into physiographic/image interpretation units based on image characteristics. The image interpretation legend for physiography is given below.

Image Interpretation Legend for Physiography

G- Granite Gneiss Landscape

G1	Hills/ Ridges/ Mounds
G11	Summits
G12	Side slopes
G121	Side slopes with dark grey tones
G2	Uplands
G21	Summits
G22	Gently sloping uplands
G221	Gently sloping uplands, yellowish green (eroded)
G222	Gently sloping uplands, yellowish white (severely eroded)
G23	Very gently sloping uplands
G231	Very gently sloping uplands, yellowish green
G232	Very gently sloping uplands, medium green and pink
G233	Very gently sloping uplands, pink and green (scrub land)
G234	Very gently sloping uplands, medium greenish grey
G235	Very gently sloping uplands, yellowish white (eroded)
G236	Very gently sloping uplands, dark green
G237	Very gently sloping uplands, medium pink (coconut garden)
G238	Very gently sloping uplands, pink and bluish white (eroded)
G24	Valleys/ lowlands
G241	Valleys, pink tones
G242	Valleys gray mixed with pink tones




Fig 3.1 Scanned and Digitized Cadastral map of Belagunda-3 microwatershed




Fig.3.2 Satellite Image of Belagunda-3 microwatershed




Fig.3.3 Cadastral map overlaid on IRS PAN+LISS IV merged imagery of Belagunda-3 microwatershed

3.3 Field Investigation

The field boundaries and survey numbers given on the cadastral sheet were located on the ground by following permanent features like roads, cart tracks, *nallas*, streams, tanks etc., and wherever changes were noticed, they were incorporated on the microwatershed cadastral map. Preliminary traverse of the microwatershed was carried out with the help of cadastral map, imagery and toposheets. While traversing, landforms and physiographic units identified were checked and preliminary soil legend was prepared by studying soils at few selected places. Then, intensive traversing of each physiographic unit like hills, ridges, uplands and valleys was carried out. Based on the variability observed on the surface, transects (Fig. 3.4) were selected across the slope covering all the landform units in the microwatershed (Natarajan and Dipak Sarkar, 2010).




Fig: 3.4. Location of profiles in a transect

In the selected transect, soil profiles were located (Fig. 3.4) at closely spaced intervals to take care of any change in the land features like break in slope, erosion, gravel, stones etc. In the selected sites, profiles (vertical cut showing the soil layers from surface to the rock) were opened upto 200 cm or to the depth limited by rock or hard substratum and studied in detail for all their morphological and physical characteristics. The soil and site characteristics were recorded for all profile sites on a standard proforma as per the guidelines given in USDA Soil Survey Manual (Soil Survey Staff, 2012). Apart from the transect study, profiles were also studied at random, almost like in a grid pattern, outside the transect areas.

Based on the soil characteristics, the soils were grouped into different soil series. Soil series is the most homogeneous unit having similar horizons and properties and behaves similarly for a given level of management. Soil depth, texture, colour, kind of horizon and horizon sequence, calcareousness, amount and nature of gravel present, nature of substratum *etc*, were used as the major differentiating characteristics for

identifying soil series occurring in the area. The differentiating characteristics used for identifying the soil series are given in Table 3.1. Based on the above characteristics, 5 soil series were identified in the Belagunda-3 microwatershed.

Table 3.1 Differentiating Characteristics used for identifying soil Series
(Characteristics are of Series Control Section)

Soils of Granite gneiss Landscape							
Sl. no	Soil Series	Depth (cm)	Colour (moist)	Texture	Gravel (%)	Horizon sequence	Calcareousness
1	BMD (Bommaraladoddi)	>150	5YR 3/3, 4/1, 4/3, 4/6	scl	-	Ap-Bt	e
2	MDR (Madhwara)	>150	10YR 3/1, 3/2, 2/1, 2/2	scl	-	Ap-Bw	e
3	ANR (Anur)	100-150	10YR 4/3,4/1	sc-c	-	Ap-Bw	es
4	VKS (Vankasambar)	100-150	10YR 5/3, 4/2, 2/1, 2/2, 3/2, 4/3	scl	-	Ap-Bw	es
5	TMK (Thumakur)	>150	10YR 3/1, 3/2, 3/3, 4/3	c	-	Ap-Bw	e

3.4 Soil Mapping

The area under each soil series was further separated into soil phases and their boundaries delineated on the cadastral map based on the variations observed in the texture of the surface soil, slope, erosion, presence of gravel, stoniness etc. A soil phase is a subdivision of soil series based mostly on surface features that affect its use and management. The soil mapping units are shown on the map (Fig.3.5) in the form of symbols. During the survey many profile pits, few minipits and a few auger bores representing different landforms occurring in the microwatershed were studied. In addition to the profile study, spot observations in the form of minipits, road cuts, terrace cuts etc., were studied to validate the soil boundaries on the soil map.

The soil map shows the geographic distribution of 5 mapping units representing 5 soil series occurring in the microwatershed. The soil map unit (soil legend) description is presented in Table 3.2. The soil phase map (management units) shows the distribution of 5 soil phases mapped in the microwatershed. Each mapping unit (soil phase) delineated on the map has similar soil and site characteristics. In other words, all the farms or survey numbers included in one phase will have similar management needs and have to be treated accordingly.

3.5 Land Management Units

The 5 soil phases identified and mapped in the microwatershed were grouped into 2 Land Management Units (LMU's) for the purpose of preparing a Proposed Crop Plan for sustained development of the microwatershed. The database (soil phases) generated under LRI was utilized for identifying Land Management Units (LMU's) based on the

management needs. One or more than one soil site characteristic having influence on the management have been chosen for identification and delineation of LMUs. For Belagunda-3 microwatershed, five soil and site characteristics, namely soil depth, soil texture, slope, erosion and gravel content have been considered for defining LMUs. The Land Management Units are expected to behave similarly for a given level of management.

3.6 Laboratory Characterization

Soil samples were collected from representative master profiles for laboratory characterization by following the methods outlined in the Laboratory Manual (Sarma *et al*, 1987). Surface soil samples collected from farmer's fields (47 samples) for fertility status (major and micronutrients) at 320 m grid interval in the year 2018 were analyzed in the laboratory (Katyral and Rattan, 2003). By linking the soil fertility data to the survey numbers through GIS, soil fertility maps were generated by using Kriging method for the microwatershed.

Table 3.2 Soil map unit description of Belagunda-3 microwatershed

*Soil map unit No.	Soil Series	Soil Phase	Mapping Unit Description	Area in ha(%)
Soils of Granite and Granite Gneiss Landscape				
	BMD		Bomraddoddi soils are very deep (>150 cm), well drained, have dark gray, reddish brown to dark reddish brown and yellowish red sandy clay loam soils occurring on very gently sloping uplands under cultivation	25 (5.09)
64		BMDcB2	Sandy loam surface, slope 1-3%, moderate erosion	25 (5.09)
	MDR		Madhwara soils are very deep (>150 cm), moderately well drained, have very dark gray to very dark brown, slightly calcareous sandy clay loam soils occurring on nearly level to very gently sloping uplands under cultivation	244 (50.26)
60		MDRiA1	Sandy clay surface, slope 0-1%, slight erosion	244 (50.26)
	ANR		Anur soils are deep (100-150 cm), moderately well drained, have dark gray to brown, calcareous sodic cracking clay soils occurring on very gently sloping uplands under cultivation	1 (0.16)
55		ANRiB2	Sandy clay surface, slope 1-3%, moderate erosion	1 (0.16)
	VKS		Vankasambar soils are deep (100-150 cm), moderately well drained, very dark brown to brown, sodic calcareous sandy clay loam soils occurring on very gently to gently sloping lowlands under cultivation	89 (18.39)
100		VKSmB1	Clay surface, slope 1-3%, slight erosion	89 (18.39)

*Soil map unit No.	Soil Series	Soil Phase	Mapping Unit Description	Area in ha(%)
	TMK		Thumakur soils are very deep (>150 cm), moderately well drained, have brown to very dark grayish brown, sodic clay black soils occurring on nearly level to very gently sloping lowlands under cultivation	56 (11.54)
104		TMKiB2	Sandy clay surface, slope 1-3%, moderate erosion	56 (11.54)
1000		Others	Habitation and water body	71 (14.55)

* Soil map unit numbers are continuous for the taluk, not for the microwatershed




Fig 3.5 Soil phase or Management Units - Belagunda-3 microwatershed

THE SOILS

Detailed information pertaining to the nature, extent and their distribution of different kinds of soils occurring in Belagunda-3 microwatershed is provided in this chapter. The microwatershed area has been identified as granite gneiss landscape based on geology. In all, 5 soil series are identified. Soil formation is the result of the combined effect of environmental and terrain factors that are reflected in soil morphology. In the granite gneiss landscape, it is by parent material, relief and climate.

A brief description of each of the 5 soil series identified followed by 5 soil phases (management units) mapped under each series are furnished below. The physical and chemical characteristics of soil series identified in Belagunda-3 microwatershed are given in Table 4.1 along with soil classification. The soils in any one map unit differ from place to place in their depth, texture, slope, gravelliness, erosion or any other site characteristic that affect management. The soil phase map can be used for identifying the suitability of areas for growing specific crops or for other alternative uses and also for deciding the type of conservation structures needed. The detailed information on soil and site-characteristics like soil depth, surface soil texture, slope, erosion, gravelliness, AWC, LCC etc, with respect to each of the soil phase identified is given village/survey number wise for the microwatershed in Appendix-I.

4.1 Soils of granite gneiss landscape

In this landscape, 5 soil series are identified and mapped. Of these, MDR series occupies maximum area of 244 ha (50%) followed by VKS 89 ha (18%), TMK 56 ha (12%), BMD 25 ha (5%) and ANR 1 ha (<1%). Brief description of each series identified and number of soil phases mapped is given below.

4.1.1 Bomraldoddi (BMD) Series: Bomraldoddi soils are very deep (>150 cm), well drained, have dark reddish brown to dark grey, reddish brown, dark brown and yellowish red, slightly calcareous sandy clay loam soils. They are developed from weathered granite gneiss and occur on nearly level to very gently sloping uplands under cultivation. The Bommaraladoddi series has been classified as a member of the fine-loamy, mixed, isohyperthermic family of Rhodic Paleustalfs.

The thickness of the solum is more than 150 cm. The thickness of A horizon ranges from 11 to 17 cm. Its colour is in 5 YR hue with value 3 to 4 and chroma 1 to 5. Texture varies from sandy loam to sandy clay loam. The thickness of B horizon is more than 150 cm. Its colour is in hue 5 YR with value 4 and chroma 1 to 6. Texture varies from sandy clay loam to sandy clay and is slightly calcareous. The available water capacity is high (151-200 mm/m). Only one phase was identified and mapped.

4.1.2 Madhwara (MDR) Series: Madhwara soils are very deep (>150 cm), well drained, have black to very dark brown and very dark gray to very dark grayish brown, slightly calcareous sandy clay loam soils. They are developed from weathered granite gneiss and occur on nearly level to very gently sloping uplands under cultivation. The Madhwara series has been classified as a member of the fine-loamy, mixed, isohyperthermic family of Fluventic Haplustepts.

The thickness of the solum is more than 150 cm. The thickness of A horizon ranges from 10 to 16 cm. Its colour is in 10 YR hue with value 3 to 5 and chroma 2 to 3. Texture varies from sandy clay and clay. The thickness of B horizon is >150 cm. Its colour is in 10 YR hue with value 3 to 5 and chroma 1 to 3. Texture varies from sandy clay loam to sandy clay and is slightly calcareous. The available water capacity is very high (>200 mm/m). Only one phase was identified and mapped.



Landscape and Soil Profile characteristics of Madhwara (MDR) Series

4.1.3 Anur (ANR) Series: Anur soils are deep (100-150 cm), moderately well drained, have dark gray to dark brown, calcareous sodic, cracking clay soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation. The Anur series has been classified as a member of the fine, mixed (calcareous), isohyperthermic family of Typic Haplustepts.

The thickness of the solum ranges from 102 to 148 cm. The thickness of A-horizon ranges from 9 to 17 cm. Its colour is in 10 YR hue with value 3 to 4 and chroma 2 to 4. The texture ranges from loamy sand to sandy clay loam and sandy clay and are calcareous. The thickness of B horizon ranges from 102 to 135 cm. Its colour is in 10 YR hue with value 3 to 5 and chroma 1 to 6. Texture is sandy clay loam to sandy clay and clay and is calcareous sodic soils. The available water capacity is very high (>200 mm/m). Only one phase was identified and mapped.



Landscape and Soil Profile characteristics of Anur (ANR) Series

4.1.4 Vankasambar (VKS) Series: Vankasambar soils are deep (100-150 cm), well drained, very dark brown to brown, sodic calcareous sandy clay loam soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping lowlands under cultivation. The Vankasambar series has been classified as a member of the fine-loamy, mixed (calcareous), isohyperthermic family of **Fulventic** Haplustepts.

The thickness of the solum ranges from 120 to 150 cm. The thickness of A horizon ranges from 9 to 22 cm. Its colour is in 10 YR hue with value 4 to 5 and chroma 2 to 5. The texture varies from loamy sand, sandy clay loam and sandy clay. The thickness of B horizon ranges from 102 to 138 cm. Its colour is in 10 YR hue with value 2 to 5 and chroma 2 to 4. Texture is sandy clay loam to sandy clay and is calcareous sodic soils. The available water capacity is very high (>200 mm/m). Only one phase was identified and mapped.



Landscape and Soil Profile characteristics of Vankasambaar (VKS) Series

4.1.5 Thumakur (TMK) Series: Thumakur soils are very deep (>150 cm), moderately well drained, have very dark gray to dark brown, slightly calcareous sodic clay soils. They are developed from weathered granite gneiss and occur on nearly level to very gently sloping low lands under cultivation. The Thumakur series has been classified as a member of the fine, mixed, isohyperthermic family of Typic Haplustepts.

The thickness of the solum ranges from 150-200 cm. The thickness of A horizon ranges from 7 to 14 cm. Its colour is in 10 YR hue with value 3 to 5 and chroma 1 to 3. Texture varies from sandy loam to sandy clay and clay. The thickness of B horizon is >150 cm. Its colour is in 10 YR hue with value 3 to 5 and chroma 1 to 3. Texture varies from sandy clay to clay and is slightly calcareous sodic soils. The available water capacity is very high (>200 mm/m). Only one phase was identified and mapped.



Landscape and Soil Profile characteristics of Thumakur (TMK) Series

Table 4.1 Physical and Chemical characteristics of soil series identified in Belagunda-3 microwatershed

Soil Series: Madhawara (MDR) **Pedon:** T₂ P₂

Location: 16°43'48.9"N 77°18'38.3"E, Yaleri village, Balichakra hobli, Yadgir taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bengaluru **Classification:** Fine-loamy, mixed, isohyperthermic Fluventic Haplustepts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-11	Ap	58.94	20.74	20.32	5.41	7.28	13.31	20.89	12.06	-	scl	16.47	8.85
11-30	Bw1	55.52	19.32	25.16	5.00	7.19	13.12	19.69	10.52	-	scl	18.25	10.18
30-58	Bw2	53.95	19.15	26.90	4.68	7.48	12.58	19.65	9.56	-	scl	26.99	14.02
58-117	Bw3	52.68	19.51	27.81	2.84	5.47	14.72	20.82	8.83	-	scl	37.86	17.40
117-160	Bw4	49.95	17.27	32.79	2.11	5.07	14.15	20.49	8.13	-	scl	44.15	20.38

Depth (cm)	pH (1:2.5)			E.C. (1:2.5) dS m ⁻¹	O.C. %	CaCO ₃ %	Exchangeable bases					CEC	CEC/Clay	Base saturation %	ESP %
	Water	CaCl ₂	M KCl				Ca	Mg	K	Na	Total				
0-11	8.31	-	-	0.33	0.46	2.76	-	-	0.45	0.47	-	20.57	1.01	100	0.90
11-30	9.25	-	-	0.20	0.31	4.20	-	-	0.19	1.40	-	23.98	0.95	100	2.34
30-58	9.78	-	-	0.40	0.19	5.76	-	-	0.16	1.53	-	24.53	0.91	100	2.49
58-117	9.94	-	-	0.88	0.23	4.80	-	-	0.18	9.09	-	24.31	0.87	100	14.96
117-160	9.98	-	-	0.93	0.15	3.00	-	-	0.24	11.09	-	28.27	0.86	100	15.69

Contd...

Soil Series: Anur (ANR) **Pedon:** R-15

Location: 16°32'45.0"N 77°23'57.4"E, Duppalli village, Sydhapura hobli, Yadgir taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bengaluru **Classification:** Fine, mixed (calcareous), isohyperthermic Typic Haplustepts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-18	Ap	64.60	13.44	21.96	7.33	10.42	18.68	20.12	8.05	<15	scl	16.59	7.96
18-49	Bw1	56.66	12.19	31.15	4.73	9.80	18.66	17.02	6.45	-	scl	33.38	13.51
49-95	Bw2	39.94	17.81	42.25	3.09	3.30	15.44	10.65	7.45	<15	c	44.68	25.23
95-123	Bw3	30.65	17.58	51.77	1.50	5.57	10.18	9.65	3.75	<15	c	54.94	32.07

Depth (cm)	pH (1:2.5)			E.C. (1:2.5) dS m ⁻¹	O.C. %	CaCO ₃ %	Exchangeable bases					CEC	CEC/Clay	Base saturation %	ESP %
	Water	CaCl ₂	M KCl				Ca	Mg	K	Na	Total				
0-18	10.17	-	-	0.365	0.48	6.11	-	-	0.25	3.52	-	19.90	0.91	100	7.08
18-49	10.32	-	-	1.38	0.30	6.76	-	-	0.21	16.03	-	24.60	0.79	100	26.07
49-95	10.08	-	-	2.55	0.17	6.11	-	-	0.33	21.49	-	32.60	0.77	100	26.36
95-123	9.92	-	-	2.56	0.12	7.93	-	-	0.51	26.03	-	36.00	0.70	100	28.92

Contd...

Soil Series: Vankasambar (VKS) **Pedon:** R-11

Location: 16°34'49.4"N 77°22'46.5"E, Baddepalli village, Sydhapura hobli, Yadgir taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bengaluru **Classification:** Fine-loamy, mixed, calcareous, isohyperthermic Fulventic Haplustepts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-14	Ap	61.32	10.31	28.37	7.14	12.07	16.04	19.03	7.05	-	scl	20.65	11.25
14-37	Bw1	62.63	8.72	28.65	9.88	14.50	16.19	15.57	6.49	-	scl	24.37	11.33
37-80	Bw2	61.43	9.14	29.43	4.84	15.45	18.01	16.73	6.40	-	scl	41.96	13.39
80-108	Bw3	55.39	11.75	32.86	4.06	5.99	23.87	15.39	6.08	-	scl	45.20	15.45

Depth (cm)	pH (1:2.5)			E.C. (1:2.5) dS m ⁻¹	O.C. %	CaCO ₃ %	Exchangeable bases					CEC	CEC/Clay	Base saturation %	ESP %
	Water	CaCl ₂	M KCl				Ca	Mg	K	Na	Total				
0-14	9.1	-	-	0.586	0.96	5.72	-	-	0.54	1.74	-	17.57	0.62	100	3.97
14-37	10.35	-	-	0.595	0.52	7.80	-	-	0.50	4.24	-	16.65	0.58	100	10.19
37-80	10.39	-	-	2.14	0.28	12.35	-	-	0.64	15.89	-	13.45	0.46	100	47.24
80-108	11.15	-	-	3	0.32	11.70	-	-	0.74	20.69	-	22.58	0.69	100	36.656

Contd...

Soil Series: Thumakuru (TMK) **Pedon:** R-10

Location: 16°38'01.3"N 77°16'49.8"E, Kilankera village, Balichakra hobli, Yadgir taluk and district

Analysis at: NBSS&LUP, Regional Centre, Bengaluru **Classification:** Fine, mixed, isohyperthermic Typic Haplustepts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-12	Ap	62.92	15.76	21.32	5.56	9.37	21.83	18.33	7.83	-	scl	17.98	6.60
12-29	Bw1	45.91	18.53	35.56	6.08	8.18	15.41	11.43	4.82	-	sc	33.40	11.79
29-74	Bw2	48.47	16.24	35.29	5.93	9.84	16.40	11.75	4.55	-	sc	28.66	11.19
74-132	Bw3	38.25	20.59	41.16	3.21	8.23	14.64	8.97	3.21	-	c	38.85	14.72
132-158	Bw4	36.87	19.99	43.14	3.54	7.61	13.08	8.57	4.07	-	c	44.36	15.75

Depth (cm)	pH (1:2.5)			E.C. (1:2.5) dS m ⁻¹	O.C. %	CaCO ₃ %	Exchangeable bases					CEC	CEC/Clay	Base saturation %	ESP %
	Water	CaCl ₂	M KCl				Ca	Mg	K	Na	Total				
0-12	9.60	-	-	0.35	0.48	1.44	-	-	0.23	3.62	-	21.83	1.02	100	6.63
12-29	9.72	-	-	1.27	0.50	1.44	-	-	0.59	20.88	-	30.50	0.86	100	27.39
29-74	9.16	-	-	3.44	0.31	3.72	-	-	0.38	25.84	-	28.68	0.81	100	36.04
74-132	9.33	-	-	2.52	0.23	4.92	-	-	0.82	20.25	-	34.99	0.85	100	23.148
132-158	9.23	-	-	2.07	0.31	3.48	-	-	0.70	21.03	-	34.24	0.79	100	24.564

INTERPRETATION FOR LAND RESOURCE MANAGEMENT

The most important soil and site characteristics that affect the land use and conservation needs of an area are land capability, soil depth, soil texture, coarse fragments, available water capacity, soil slope, soil erosion, soil reaction etc. These are interpreted from the data base generated through land resource inventory and several thematic maps are generated. These would help in identifying the areas suitable for growing crops and, soil and water conservation measures and structures needed thus helping to maintain good soil health for sustained crop production. The various interpretative and thematic maps generated are described below.

5.1 Land Capability Classification

Land capability classification is an interpretative grouping of soil map units (soil phases) mainly based on inherent soil characteristics, external land features and environmental factors that limit the use of land for agriculture, pasture, forestry, or other uses on a sustained basis (IARI, 1971). The land and soil characteristics used to group the land resources in an area into various land capability classes, subclasses and units are

Soil Characteristics: Depth, texture, gravelliness, calcareousness.

Land characteristics: Slope, erosion, drainage, rock outcrops.

Climate: Total rainfall and its distribution, and length of crop growing period.

The Land capability classification system is divided into land capability classes, subclasses and units based on the level of information available. Eight land capability classes are recognized. They are

Class I: They are very good lands that have no limitations or very few limitations that restrict their use.

Class II: They are good lands that have minor limitations and require moderate conservation practices.

Class III: They are moderately good lands that have moderate limitations that reduce the choice of crops or that require special conservation practices.

Class IV: They are fairly good lands that have very severe limitations that reduce the choice of crops or that require very careful management.

Class V: Soils in these lands are not likely to erode, but have other limitations like wetness that are impractical to remove and as such not suitable for agriculture, but suitable for pasture or forestry with minor limitations.

Class VI: The lands have severe limitations that make them generally unsuitable for cultivation, but suitable for pasture or forestry with moderate limitations.

Class VII: The lands have very severe limitations that make them unsuitable for cultivation, but suitable for pasture or forestry with major limitations.

Class VIII: Soil and other miscellaneous areas (rock lands) that have very severe limitations that nearly preclude their use for any crop production, but suitable for wildlife, recreation and installation of wind mills.

The land capability subclasses are recognised based on the dominant limitations observed within a given land capability class. The subclasses are designated by adding a lower case letter like 'e', 'w', 's', or 'c' to the class numeral. The subclass "e" indicates that the main hazard is risk of erosion, "w" indicates drainage or wetness as a limitation for plant growth, "s" indicates shallow soil depth, coarse or heavy textures, calcareousness, salinity/alkalinity or gravelliness and "c" indicates limitation due to climate.

The land capability subclasses have been further subdivided into land capability units based on the kinds of limitations present in each subclass. Ten land capability units are used in grouping the soil map units. They are stony or rocky (0), erosion hazard (slope, erosion) (1), coarse texture (sand, loamy sand, sandy loam) (2), fine texture (cracking clay, silty clay) (3), slowly permeable subsoil (4), coarse underlying material (5), salinity/alkali (6), stagnation, overflow, high ground water table (7), soil depth (8) and fertility problems (9). The capability units thus identified have similar soil and land characteristics that respond similarly to a given level of management. The soils of the microwatershed have been classified upto land capability subclass level.

The 5 soil map units identified in the Belagunda-3 microwatershed are grouped under 2 land capability classes and 4 subclasses. Entire area in the microwatershed is suitable for agriculture (Fig. 5.1).

Good lands (Class II) cover a maximum area of about 55 per cent and are distributed in the major part of the microwatershed with minor problems of soil and erosion. Fairly good (Class IV) cover an area of about 30 per cent and is distributed in the western and northwestern part of the microwatershed with very severe problems of soil and erosion.




Fig. 5.1 Land Capability Classification map of Belagunda-3 microwatershed

5.2 Soil Depth

Soil depth refers to the depth of the soil occurring above the parent material or hard rock. The depth of the soil determines the effective rooting depth for plants and in accordance with soil texture, mineralogy and gravel content, the capacity of the soil column to hold water and nutrient availability. Soil depth is one of the most important soil characteristic that is used in differentiating soils into different soil series. The soil depth classes used in identifying soils in the field are very shallow (<25 cm), shallow (25-50 cm), moderately shallow (50-75 cm), moderately deep (75-100 cm), deep (100-150 cm) and very deep (>150 cm). They were used to classify the soils into different depth classes and a soil depth map was generated. The area extent and their geographical distribution in the microwatershed is given in Fig. 5.2.

Deep to very deep (100 to >150 cm) soils occupy entire area of the microwatershed.

The most productive lands with respect to soil rooting depth where all climatically adapted annual and perennial crops can be grown is deep (100 - >150 cm depth) soils of the microwatershed.




Fig. 5.2 Soil depth map of Belagunda-3 microwatershed

5.3 Surface Soil Texture

Texture is an expression to indicate the coarseness or fineness of the soil as determined by the relative proportion of primary particles of sand, silt and clay. It has a direct bearing on the structure, porosity, adhesion and consistence. The surface layer of a soil to a depth of about 25 cm is the layer that is most used by crops and plants. The surface soil textural class provides a guide to understanding soil-water retention and availability, nutrient holding capacity, infiltration, workability, drainage, physical and chemical behaviour, microbial activity and crop suitability. The textural classes used for LRI were used to classify and a surface soil texture map was generated. The area extent and their geographical distribution in the microwatershed is shown in Figure 5.3.

Maximum area of about 389 ha (80%) of the microwatershed has clayey soils at the surface and are distributed in the major part of the microwatershed. An area of 25 ha (5%) has soils that are loamy and are distributed in the northwestern part of the microwatershed. Clayey and loamy soils have high potential for soil-water retention and availability, and nutrient retention and availability, but clay soils have more problems of drainage, infiltration, work ability and other physical problems.




Fig. 5.3 Surface soil texture map of Belagunda-3 microwatershed

5.4 Soil Gravelliness

Gravel is the term used for describing coarse fragments between 2 mm and 7.5 cm diameter and stones for those between 7.5 cm and 25 cm. The presence of gravel and stones in soil reduces the volume of soil responsible for moisture and nutrient storage, drainage, infiltration and runoff, and hinders plant growth by impeding root growth and seedling emergence, intercultural operations and farm mechanization. The gravelliness classes used in LRI were used to classify the soils and using these classes a gravelliness map was generated. The area extent and their geographic distribution in the microwatershed is shown in Figure 5.4.

Entire area of the microwatershed has non gravelly (<15%) soils.

The most productive soils that are non gravelly (<15%), where all climatically adapted long duration crops can be grown.




Fig. 5.4 Soil gravelliness map of Belagunda-3 microwatershed

5.5 Available Water Capacity

The soil available water capacity (AWC) is estimated based on the ability of the soil column to retain water between the tensions of 0.33 and 15 bar in a depth of 100 cm or the entire solum if the soil is shallower. The AWC of the soils (soil series) as estimated by considering the soil texture, mineralogy, soil depth and gravel content (Sehgal *et al.*, 1990) and accordingly the soil map units were grouped into five AWC classes *viz.*, very low (<50 mm/m), low (50-100 mm/m), medium (100-150 mm/m), high (150-200 mm/m) and very high (>200 mm/m) and using these values, an AWC map was generated. The area extent and their geographic distribution of different AWC classes in the microwatershed is given in Figure 5.5.

Maximum area of about 390 ha (80%) in the microwatershed have soils that are very high (>200 mm/m) in available water capacity and are distributed in the major part of the microwatershed. An area of about 25 ha (5%) is medium (101 - 150 mm/m) in available water capacity and are distributed in the northwestern part of the microwatershed.

An area of 25 ha (5%) in the microwatershed has soils that are somewhat problematic with regard to available water capacity. Here, only medium or short duration crops can be grown and the probability of crop failure is very high. These areas are best put to other alternative uses. Maximum area of 390 ha (80%) are potential areas with regard to AWC where all climatically adapted annual and perennial crops can be grown.




Fig. 5.5 Soil available water capacity map of Belagunda-3 microwatershed

5.6 Soil Slope

Soil slope refers to the inclination of the surface of the land. It is defined by gradient, shape and length, and is an integral feature of any soil as a natural body. Slope is considered important in soil genesis, land use and land development. The length and gradient of slope influences the rate of runoff, infiltration, erosion and deposition. The soil map units were grouped into two slope classes and a slope map was generated showing the area extent and their geographic distribution in the microwatershed (Fig. 5.6).

Maximum area of about 244 ha (50%) in the microwatershed falls under nearly level (0-1% slope) lands and are distributed in the major parts of the microwatershed. An area of 171 ha (35%) falls under very gently sloping (1-3%) lands and are distributed in the western and northwestern part of the microwatershed.

Entire area in the microwatershed has soils that have high potential in respect of soil slopes. In these areas, all climatically adapted annual and perennial crops can be grown without much soil and water conservation and other land development measures.




Fig. 5.6 Soil slope map of Belagunda-3 microwatershed

5.7 Soil Erosion

Soil erosion refers to the wearing away of the earth's surface by the forces of water, wind and ice involving detachment and transport of soil by raindrop impact. It is used for accelerated soil erosion resulting from disturbance of the natural landscape by burning, excessive grazing and indiscriminate felling of forest trees and tillage, all usually by man. The erosion classes showing an estimate of the current erosion status as judged from field observations in the form of rills, gullies or a carpet of gravel on the surface are recorded. Four erosion classes, viz, slight erosion (e1), moderate erosion (e2), severe erosion (e3) and very severe erosion (e4) are recognized. The soil map units were grouped into different erosion classes and a soil erosion map generated. The area extent and their spatial distribution in the microwatershed is given in Figure 5.7.

Maximum area of about 333 ha (69%) in the microwatershed falls under slightly eroded (e1 class) lands and are distributed in the maximum part of the microwatershed. An area of about 81 ha (17%) in the microwatershed falls under moderately eroded (e2 class) lands and are distributed in the northwestern part of the microwatershed.

An area of 81 ha in the microwatershed has soils that are problematic because of moderate erosion. For these areas, taking up of soil and water conservation and other land development measures are needed.




Fig. 5.7 Soil erosion map of Belagunda-3 microwatershed

FERTILITY STATUS

Soil fertility plays an important role in increasing crop yield. The adoption of high yielding varieties that require high amounts of nutrients has resulted in deficiency symptoms in crops and plants due to imbalanced fertilization and poor inherent fertility status as these areas are characterised by low rainfall and high temperatures. Hence, it is necessary to know the fertility (macro and micro nutrients) status of the soils of the watersheds for assessing the kind and amount of fertilizers required for each of the crop intended to be grown. For this purpose, the surface soil samples collected from the grid points (one soil sample at every 320 m interval) all over the microwatershed through land resource inventory in the year 2018 were analysed for pH, EC, organic carbon, available phosphorus and potassium, and for micronutrients like zinc, boron, copper, iron manganese and secondary nutrient sulphur.

Soil fertility data generated has been assessed and individual maps for all the nutrients for the microwatershed have been generated using Kriging method under GIS. The village/survey number wise fertility data for the microwatershed is given in Appendix-II.

6.1 Soil Reaction (pH)

The soil analysis of the Belagunda-3 microwatershed for soil reaction (pH) showed that an area of about 111 ha (23%) is moderately alkaline (pH 7.8-8.4) and distributed in the eastern and northwestern part of the microwatershed. Maximum area of about 281 ha (58%) is strongly alkaline (pH 8.4- 9.0) and are distributed in the major part of the microwatershed. An area of about 22 ha (4%) is very strongly alkaline (pH >9.0) and are distributed in the southern part of the microwatershed (Fig. 6.1). Entire area in the microwatershed is alkaline in reaction.

6.2 Electrical Conductivity (EC)

The electrical conductivity is non saline ($<2 \text{ dsm}^{-1}$) in a maximum area of about 347 ha (71%) and are distributed in the major part of the microwatershed. An area of 39 ha (8%) is low ($2-4 \text{ dsm}^{-1}$) and are distributed in the southeastern part of the microwatershed. An area of 29 ha (6%) is medium ($4 - 8 \text{ dsm}^{-1}$) and are distributed in the southeastern part of the microwatershed.

6.3 Organic Carbon

Organic carbon content is low ($<0.5 \%$) in an area of about 162 ha (33%) and are distributed in the northwestern, western, southwestern, southern, southeastern and northeastern part of the microwatershed. Maximum area of 252 ha (52%) is medium ($0.5 - 0.75 \%$) and are distributed in the major part of the microwatershed (Fig. 6.3).




Fig.6.1 Soil reaction (pH) map of Belagunda-3 microwatershed




Fig.6.2 Electrical conductivity (EC) map of Belagunda-3 microwatershed




Fig.6.3 Soil organic carbon map of Belagunda-3 microwatershed

6.4 Available Phosphorus

Available phosphorus content is low (<23 kg/ha) in a maximum area of about 152 ha (31%) and are distributed in the major part of the microwatershed. Medium (23-57 kg/ha) in an area of about 134 ha (28%) and are distributed in the central, western, northwestern and southern part of the microwatershed. High (>57 kg/ha) in an area of 128 ha (26%) and are distributed in the northern and southwestern part of the microwatershed (Fig. 6.4).

6.5 Available Potassium

Available potassium content is medium (145-337 kg/ha) in a maximum area of about 291 ha (60%) and are distributed in the major part of the microwatershed. High (>337 kg/ha) in an area of 123 ha (25%) and is distributed in the northern, central and southern part of the microwatershed (Fig. 6.5).

6.6 Available Sulphur

An area of about 2 ha (<1%) is low (<10 ppm) in available sulphur content and are distributed in the northwestern part of the microwatershed. Medium (10 - 20 ppm) in an area of about 27 ha (6%) and is distributed in the northwestern part of the microwatershed. High (> 20 ppm) in a maximum area of about 385 ha (79%) and is distributed in the major part of the microwatershed (Fig. 6.6).

6.7 Available Boron

Available boron content is low (<0.5 ppm) in an area of about 0.3 ha and are distributed in the northern part of the microwatershed. Maximum area of 290 ha (60%) is medium (0.5-1.0 ppm) in available boron content and are distributed in the major part of the microwatershed. An area of 124 ha (26%) is high (>1.0 ppm) in available boron content and are distributed in the eastern and southeastern part of the microwatershed (Fig. 6.7).

6.8 Available Iron

Available iron content is deficient in an area of 138 ha (28%) (<4.5 ppm) and are distributed in the eastern, northeastern, southeastern and northwestern part of the microwatershed. Sufficient in 277 ha (57%) (>4.5 ppm) and is distributed in the major part of the microwatershed (Fig 6.8).

6.9 Available Manganese

Available manganese content is sufficient (>1.0 ppm) in the entire microwatershed area (Fig 6.9).

6.10 Available Copper

Available copper content is sufficient (>0.2 ppm) in the entire microwatershed area (Fig 6.10).




Fig.6.4 Soil available phosphorus map of Belagunda-3 microwatershed




Fig.6.5 Soil available potassium map of Belagunda-3 microwatershed




Fig.6.6 Soil available sulphur map of Belagunda-3 microwatershed




Fig.6.7 Soil available boron map of Belagunda-3 microwatershed




Fig.6.8 Soil available iron map of Belagunda-3 microwatershed




Fig.6.9 Soil available manganese map of Belagunda-3 microwatershed




Fig.6.10 Soil available copper map of Belagunda-3 microwatershed

6.11 Available Zinc

Available zinc content is deficient (<0.6 ppm) in the entire microwatershed area (Fig 6.11).




Fig.6.11 Soil available zinc map of Belagunda-3 microwatershed

LAND SUITABILITY FOR MAJOR CROPS

The soil and land resource units (soil phases) of Belagunda-3 microwatershed were assessed for their suitability for growing food, fodder, fibre and other horticulture crops by following the procedure as outlined in FAO, 1976 and 1983. Crop requirements were developed for each of the crop from the available research data and also by referring to Naidu *et. al.* (2006) and Natarajan *et. al.* (2015). The crop requirements (Table 7.2 to 7.30) were matched with the soil and land characteristics (Table 7.1) to arrive at the crop suitability. The criteria tables are given at the end of the chapter. In FAO land suitability classification, two orders are recognized. Order S-Suitable and Order N-Not suitable. The orders have classes, subclasses and units. Order-S has three classes, Class S1-Highly Suitable, Class S2-Moderately Suitable and Class S3- Marginally Suitable. Order N has two classes, N1-Currently not Suitable and N2- Permanently not Suitable. There are no subclasses within the Class S1 as they will have very minor or no limitation for crop growth. Classes S2 and S3 are divided into subclasses based on the kinds of limitations encountered. The limitations that affect crop production are ‘c’ for erratic rainfall and its distribution and length of growing period (LGP), ‘e’ for erosion hazard, ‘r’ for rooting condition, ‘t’ for lighter or heavy texture, ‘g’ for gravelliness or stoniness, ‘n’ for nutrient availability, ‘l’ for topography, ‘m’ for moisture availability, ‘w’ for drainage and ‘z’ for calcareousness. These limitations are indicated as lower case letters to the class symbol. For example, moderately suitable lands with the limitations of soil depth and erosion are designated as S2re. For the microwatershed, the soil mapping units were evaluated and classified up to subclass level.

Using the above criteria, the soil map units of the microwatershed were evaluated and land suitability maps for 29 major agricultural and horticultural crops were generated. The detailed information on the kind of suitability of each of the soil phase for the crops assessed are given village/ survey number wise for the microwatershed in Appendix-III.

7.1 Land Suitability for Sorghum (*Sorghum bicolor*)

Sorghum is one of the major food crop grown in Karnataka in an area of 10.47 lakh ha in Bijapur, Gulbarga, Raichur, Bidar, Belgaum, Dharwad, Bellary, Chitradurga, Mysore and Tumakuru districts. The crop requirements for growing sorghum (Table 7.2) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing sorghum was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.1.

There are no highly suitable (Class S1) lands available for growing sorghum in the microwatershed. Maximum area of about 269 ha (55%) is moderately suitable (Class S2) for growing sorghum and are distributed in the major part of the microwatershed. They have minor limitations of texture and nutrient availability. An area of about 146 ha (30%)

is marginally suitable (Class S3) for growing sorghum and is distributed in the western and northeastern part of the microwatershed with moderate limitation of nutrient availability.




Fig. 7.1 Land suitability map of Sorghum

7.2 Land Suitability for Maize (*Zea mays*)

Maize is one of the most important food crop grown in an area of 13.37 lakh ha in almost all the districts of the State. The crop requirements for growing maize (Table 7.3) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing maize was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.2.

Highly suitable (Class S1) lands for growing maize occur in an area of 25 ha (5%) and are distributed in the northwestern part of the microwatershed. Maximum area of about 244 ha (50%) is moderately suitable (Class S2) for growing maize and are distributed in the major part of the microwatershed. They have minor limitation of nutrient availability. An area of about 146 ha (30%) is marginally suitable (Class S3) for growing maize and is distributed in the western and northwestern part of the microwatershed with moderate limitation nutrient availability.




Fig. 7.2 Land suitability map of Maize

7.3 Land Suitability for Bajra (*Pennisetum glaucum*)

Bajra is one of the most important millet crop grown in an area of 2.34 lakh ha in the northern districts of Karnataka state. The crop requirements for growing bajra (Table 7.4) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing bajra was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.3.

Highly suitable (Class S1) lands for growing bajra occur in an area of 25 ha (5%) and are distributed in the northwestern part of the microwatershed. Maximum area of about 244 ha (50%) is moderately suitable (Class S2) for growing bajra and are distributed in the major part of the microwatershed. They have minor limitation of nutrient availability. An area of about 146 ha (30%) is marginally suitable (Class S3) for growing bajra and is distributed in the western and northwestern part of the microwatershed with moderate limitation nutrient availability.




Fig. 7.3 Land suitability map of Bajra

7.4 Land Suitability for Groundnut (*Arachis hypogaea*)

Groundnut is one of the major oilseed crop grown in an area of 6.54 lakh ha in Karnataka in most of the districts either as rainfed or irrigated crop. The crop requirements for growing groundnut (Table 7.5) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing groundnut was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.4.

Highly suitable (Class S1) lands for growing groundnut occur in an area of 25 ha (5%) and are distributed in the northwestern part of the microwatershed. Maximum area of about 244 ha (50%) is marginally suitable (Class S3) for growing groundnut and are distributed in the major part of the microwatershed. They have moderate limitation of nutrient availability. An area of about 146 ha (30%) is not suitable (Class N1) for growing groundnut and is distributed in the western and northwestern part of the microwatershed with severe limitation nutrient availability.




Fig. 7.4 Land suitability map of Groundnut

7.5 Land Suitability for Sunflower (*Helianthus annus*)

Sunflower is one of the most important oilseed crop grown in an area of 3.56 lakh ha in the State in all the districts. The crop requirements for growing sunflower (Table 7.6) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing sunflower was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.5.

There are no highly suitable (Class S1) lands available for growing sunflower in the microwatershed. An area of about 25 ha (5%) is moderately suitable (Class S2) for growing sunflower and are distributed in the northwestern part of the microwatershed. They have minor limitation of texture. Maximum area of about 244 ha (50%) is marginally suitable (Class S3) for growing sunflower and is distributed in the major part of the microwatershed with moderate limitation of nutrient availability. An area of about 146 ha (30%) is not suitable (Class N1) for growing sunflower and is distributed in the western and northwestern part of the microwatershed with severe limitation nutrient availability.




Fig. 7.5 Land suitability map of Sunflower

7.6 Land Suitability for Red gram (*Cajanus Cajan*)

Red gram is one of the most important pulse crop grown in an area of 7.28 lakh ha in almost all the districts of the State. The crop requirements for growing red gram (Table 7.7) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing red gram was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.6.

There are no highly suitable (Class S1) lands available for growing red gram in the microwatershed. Maximum area of about 269 ha (55%) is moderately suitable (Class S2) for growing red gram and are distributed in the major part of the microwatershed. They have minor limitations of texture and nutrient availability. An area of about 146 ha (30%) is marginally suitable (Class S3) for growing red gram and is distributed in the western and northeastern part of the microwatershed with moderate limitation of nutrient availability.




Fig. 7.6 Land suitability map of Red gram

7.7 Land Suitability for Bengal gram (*Cicer aerativum*)

Bengal gram one of is the most important pulse crop grown in about 9.39 lakh ha area in Bijapur, Raichur, Kalaburgi, Dharwad, Belgaum and Bellary districts. The crop requirements for growing bengal gram (Table 7.8) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing bengal gram was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.7.

No highly and moderately suitable (Class S1 and S2) lands are available for growing bengal gram in the microwatershed. Marginally suitable lands (Class S3) for growing bengal gram occupy entire area of the microwatershed. They have moderate limitations of texture and nutrient availability.




Fig. 7.7 Land suitability map of Bengal gram

7.8 Land Suitability for Cotton (*Gossypium hirsutum*)

Cotton is one of the most important fibre crop grown in the State in about 8.75 lakh ha area in Raichur, Dharwad, Belgaum, Kalaburgi, Bijapur, Bidar, Bellary, Chitradurga and Chamarajnagar districts. The crop requirements for growing cotton (Table 7.9) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing cotton was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.8.

No highly and moderately suitable (Class S1 and S2) lands are available for growing cotton in the microwatershed. Marginally suitable lands (Class S3) for growing cotton occupy an entire cultivated area of the microwatershed. They have moderate limitations of texture and nutrient availability.




Fig. 7.8 Land suitability map of Cotton

7.9 Land Suitability for Chilli (*Capsicum annuum*)

Chilli is one of the most important vegetable and spice crop grown in about 0.42 lakh ha in Karnataka state. The crop requirements for growing chilli (Table 7.10) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing chilli was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.9.

Highly suitable (Class S1) lands for growing chilli occur in an area of 25 ha (5%) and are distributed in the northwestern part of the microwatershed. Maximum area of about 244 ha (50%) is marginally suitable (Class S3) for growing chilli and are distributed in the major part of the microwatershed. They have moderate limitation of nutrient availability. An area of about 146 ha (30%) is not suitable (Class N1) for growing chilli and is distributed in the western and northwestern part of the microwatershed with severe limitation of nutrient availability.




Fig 7.9 Land suitability map of Chilli

7.10 Land Suitability for Tomato (*Lycopersicon esculentum*)

Tomato is one of the most important vegetable crop grown in about 0.61 lakh ha covering almost all the district of the state. The crop requirements for growing tomato (Table 7.11) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing tomato was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.10.

Highly suitable (Class S1) lands for growing tomato occur in an area of 25 ha (5%) and are distributed in the northwestern part of the microwatershed. Maximum area of about 244 ha (50%) is marginally suitable (Class S3) for growing tomato and are distributed in the major part of the microwatershed. They have moderate limitation of nutrient availability. An area of about 146 ha (30%) is not suitable (Class N1) for growing tomato and is distributed in the western and northwestern part of the microwatershed with severe limitation of nutrient availability.




Fig 7.10 Land suitability map of Tomato

7.11 Land Suitability for Brinjal (*Solanum melongena*)

Brinjal is one of the most important vegetable crop grown in the state. The crop requirements for growing brinjal (Table 7.12) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing brinjal was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.11.

Highly suitable (Class S1) lands for growing brinjal occur in an area of 25 ha (5%) and are distributed in the northwestern part of the microwatershed. Maximum area of about 244 ha (50%) is marginally suitable (Class S3) for growing brinjal and are distributed in the major part of the microwatershed. They have moderate limitation of nutrient availability. An area of about 146 ha (30%) is not suitable (Class N1) for growing brinjal and is distributed in the western and northwestern part of the microwatershed with severe limitation of nutrient availability.




Fig 7.11 Land suitability map of Brinjal

7.12 Land Suitability for Onion (*Allium cepa L.*)

Onion is one of the most important vegetable crop grown in the state. The crop requirements for growing onion (Table 7.13) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing onion was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.12.

Highly suitable (Class S1) lands for growing onion occur in an area of 25 ha (5%) and are distributed in the northwestern part of the microwatershed. Maximum area of about 390 ha (80%) is not suitable (Class N1) for growing onion and are distributed in the major part of the microwatershed. They have severe limitation of nutrient availability.




Fig 7.12 Land suitability map of Onion

7.13 Land Suitability for Bhendi (*Abelmoschus esculentus*)

Bhendi is one of the most important vegetable crop grown in the state. The crop requirements for growing bhendi (Table 7.14) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing bhendi was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.13.

Highly suitable (Class S1) lands for growing bhendi occur in an area of 25 ha (5%) and are distributed in the northwestern part of the microwatershed. Maximum area of about 244 ha (50%) is marginally suitable (Class S3) for growing bhendi and are distributed in the major part of the microwatershed. They have moderate limitation of nutrient availability. An area of about 146 ha (30%) is not suitable (Class N1) for growing bhendi and is distributed in the western and northwestern part of the microwatershed with severe limitation of nutrient availability.




Fig 7.13 Land suitability map of Bhendi

7.14 Land Suitability for Drumstick (*Moringa oleifera*)

Drumstick is one of the most important vegetable crop grown in about 2403 ha in the state. The crop requirements for growing drumstick (Table 7.15) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing drumstick was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.14.

Highly suitable (Class S1) lands for growing drumstick occur in an area of 25 ha (5%) and are distributed in the northwestern part of the microwatershed. Maximum area of about 390 ha (80%) is not suitable (Class N1) for growing drumstick and are distributed in the major part of the microwatershed. They have severe limitation of nutrient availability.




Fig 7.14 Land suitability map of Drumstick

7.15 Land Suitability for Mango (*Mangifera indica*)

Mango is one of the most important fruit crop grown in an area of 1.73 lakh ha in almost all the districts of the state. The crop requirements (Table 7.16) for growing mango were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing mango was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed is given in Figure 7.15.

Highly suitable (Class S1) lands for growing mango occur in an area of 25 ha (5%) and are distributed in the northwestern part of the microwatershed. Maximum area of about 244 ha (50%) is marginally suitable (Class S3) for growing mango and are distributed in the major part of the microwatershed. They have moderate limitation of nutrient availability. An area of about 146 ha (30%) is not suitable (Class N1) for growing mango and is distributed in the western and northwestern part of the microwatershed with severe limitation of nutrient availability.




Fig. 7.15 Land suitability map of Mango

7.16 Land Suitability for Guava (*Psidium guajava*)

Guava is one of the most important fruit crop grown in an area of 0.06 lakh ha in almost all the districts of the State. The crop requirements (Table 7.17) for growing guava were matched with the soil-site characteristics (7.1) and a land suitability map for growing guava was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.16.

Highly suitable (Class S1) lands for growing guava occur in an area of 25 ha (5%) and are distributed in the northwestern part of the microwatershed. Maximum area of about 390 ha (80%) is not suitable (Class N1) for growing guava and are distributed in the major part of the microwatershed. They have severe limitation of nutrient availability.




Fig. 7.16 Land suitability map of Guava

7.17 Land Suitability for Sapota (*Manilkara zapota*)

Sapota is one of the most important fruit crop grown in an area of 29373 ha in almost all the districts of the State. The crop requirements (Table 7.18) for growing sapota were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing sapota was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.17.

Highly suitable (Class S1) lands for growing sapota occur in an area of 25 ha (5%) and are distributed in the northwestern part of the microwatershed. Maximum area of about 244 ha (50%) is marginally suitable (Class S3) for growing sapota and are distributed in the major part of the microwatershed. They have moderate limitation of nutrient availability. An area of about 146 ha (30%) is not suitable (Class N1) for growing sapota and is distributed in the western and northwestern part of the microwatershed with severe limitation of nutrient availability.




Fig. 7.17 Land suitability map of Sapota

7.18 Land Suitability for Pomegranate (*Punica granatum*)

Pomegranate is one of the most important fruit crop commercially grown in about 18488 ha in Karnataka, mainly in Bijapur, Bagalkot, Koppal, Gadag and Chitradurga districts. The crop requirements for growing pomegranate (Table 7.19) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing pomegranate was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.18.

Highly suitable (Class S1) lands for growing pomegranate occur in an area of 25 ha (5%) and are distributed in the northwestern part of the microwatershed. Maximum area of about 244 ha (50%) is marginally suitable (Class S3) for growing pomegranate and are distributed in the major part of the microwatershed. They have moderate limitation of nutrient availability. An area of about 146 ha (30%) is not suitable (Class N1) for growing pomegranate and is distributed in the western and northwestern part of the microwatershed with severe limitation of nutrient availability.




Fig 7.18 Land suitability map of Pomegranate

7.19 Land Suitability for Musambi (*Citrus limetta*)

Musambi is one of the important fruit crop grown in an area of 3446 ha in almost all the districts of the State. The crop requirements for growing musambi (Table 7.20) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing musambi was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.19.

Highly suitable (Class S1) lands for growing musambi occur in an area of 25 ha (5%) and are distributed in the northwestern part of the microwatershed. Maximum area of about 244 ha (50%) is marginally suitable (Class S3) for growing musambi and are distributed in the major part of the microwatershed. They have moderate limitation of nutrient availability. An area of about 146 ha (30%) is not suitable (Class N1) for growing musambi and is distributed in the western and northwestern part of the microwatershed with severe limitation of nutrient availability.




Fig. 7.19 Land suitability map of Musambi

7.20 Land Suitability for Lime (*Citrus sp*)

Lime is one of the most important fruit crop grown in an area of 0.11 lakh ha in almost all the districts of the State. The crop requirements for growing lime (Table 7.21) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing lime was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7. 20.

Highly suitable (Class S1) lands for growing lime occur in an area of 25 ha (5%) and are distributed in the northwestern part of the microwatershed. Maximum area of about 244 ha (50%) is marginally suitable (Class S3) for growing lime and are distributed in the major part of the microwatershed. They have moderate limitation of nutrient availability. An area of about 146 ha (30%) is not suitable (Class N1) for growing lime and is distributed in the western and northwestern part of the microwatershed with severe limitation of nutrient availability.




Fig. 7.20 Land suitability map of Lime

7.21 Land Suitability for Amla (*Phyllanthus emblica*)

Amla is one of the medicinal fruit crop grown in almost all the districts of the State. The crop requirements for growing amla (Table 7.22) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing amla was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.21.

Highly suitable (Class S1) lands for growing amla occur in an area of 25 ha (5%) and are distributed in the northwestern part of the microwatershed. Maximum area of about 390 ha (80%) is not suitable (Class N1) for growing amla and are distributed in the major part of the microwatershed. They have severe limitation of nutrient availability.




Fig. 7.21 Land suitability map of Amla

7.22 Land Suitability for Cashew (*Anacardium occidentale*)

Cashew is one of the most important plantation nut crop grown in an area of 0.7 lakh ha in almost all the districts of the state. The crop requirements for growing cashew (Table 7.23) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing cashew was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.22.

There are no highly suitable (Class S1) lands available for growing cashew in the microwatershed. An area of about 25 ha (5%) is moderately suitable (Class S2) for growing cashew and are distributed in the northwestern part of the microwatershed. They have minor limitation of nutrient availability. Maximum area of about 390 ha (80%) is not suitable (Class N1) for growing cashew and is distributed in the major part of the microwatershed with severe limitations nutrient availability and texture.




Fig. 7.22 Land suitability map of Cashew

7. 23 Land Suitability for Jackfruit (*Artocarpus heterophyllus*)

Jackfruit is one of the most important fruit crop grown in an area of 5368 ha in almost all the districts of the State. The crop requirements for growing jackfruit (Table 7.24) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing jackfruit was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.23.

Highly suitable (Class S1) lands for growing jackfruit occur in an area of 25 ha (5%) and are distributed in the northwestern part of the microwatershed. Maximum area of about 390 ha (80%) is not suitable (Class N1) for growing jackfruit and are distributed in the major part of the microwatershed. They have severe limitation of nutrient availability.




Fig. 7.23 Land suitability map of Jackfruit

7.24 Land Suitability for Jamun (*Syzygium cumini*)

Jamun is one of the important fruit crop grown in almost all the districts of the State. The crop requirements for growing jamun (Table 25) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing jamun was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.24.

Highly suitable (Class S1) lands for growing jamun occur in an area of 25 ha (5%) and are distributed in the northwestern part of the microwatershed. Maximum area of about 390 ha (80%) is not suitable (Class N1) for growing jamun and are distributed in the major part of the microwatershed. They have severe limitation of nutrient availability.




Fig. 7.24 Land suitability map of Jamun

7.25 Land Suitability for Custard Apple (*Annona reticulata*)

Custard apple is one of the most important fruit crop grown in almost all the districts of the State. The crop requirements for growing custard apple (Table 7.26) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing custard apple was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.25.

Highly suitable (Class S1) lands for growing custard apple occur in an area of 25 ha (5%) and are distributed in the northwestern part of the microwatershed. Maximum area of about 244 ha (50%) is marginally suitable (Class S3) for growing custard apple and are distributed in the major part of the microwatershed. They have moderate limitation of nutrient availability. An area of about 146 ha (30%) is not suitable (Class N1) for growing custard apple and is distributed in the western and northwestern part of the microwatershed with severe limitation of nutrient availability.




Fig. 7.25 Land suitability map of Custard Apple

7.26 Land Suitability for Tamarind (*Tamarindus indica*)

Tamarind is one of the most important spice crop grown in almost all the districts of the state. The crop requirements for growing tamarind (Table 7.27) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing tamarind was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Fig. 7.26.

Highly suitable (Class S1) lands for growing tamarind occur in an area of 25 ha (5%) and are distributed in the northwestern part of the microwatershed. Maximum area of about 390 ha (80%) is not suitable (Class N1) for growing tamarind and are distributed in the major part of the microwatershed. They have severe limitation of nutrient availability.




Fig. 7.26 Land suitability map of Tamarind

7.27 Land Suitability for Mulberry (*Morus nigra*)

Mulberry is one of the important leaf crop grown for rearing silkworms in about 1.6 lakh ha area in all the districts of the state. The crop requirements for growing mulberry (Table 7.28) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing mulberry was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.27.

Highly suitable (Class S1) lands for growing mulberry occur in an area of 25 ha (5%) and are distributed in the northwestern part of the microwatershed. Maximum area of about 390 ha (80%) is not suitable (Class N1) for growing mulberry and are distributed in the major part of the microwatershed. They have severe limitation of nutrient availability.




Fig 7.27 Land suitability map of Mulberry

7.28 Land Suitability for Marigold (*Tagetes sps.*)

Marigold is one of the most important flower crop grown in an area of 9108 ha in almost all the districts of the State. The crop requirements (Table 7.29) for growing marigold were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing marigold was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.28.

Highly suitable (Class S1) lands for growing marigold occur in an area of 25 ha (5%) and are distributed in the northwestern part of the microwatershed. Maximum area of about 244 ha (50%) is marginally suitable (Class S3) for growing marigold and are distributed in the major part of the microwatershed. They have moderate limitation of nutrient availability. An area of about 146 ha (30%) is not suitable (Class N1) for growing marigold and is distributed in the western and northwestern part of the microwatershed with severe limitation of nutrient availability.




Fig. 7.28 Land suitability map of Marigold

7.29 Land Suitability for Chrysanthemum (*Dendranthema grandiflora*)

Chrysanthemum is one of the most important flower crop grown in an area of 4978 ha in almost all the districts of the State. The crop requirements (Table 7.30) for growing chrysanthemum were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing chrysanthemum was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.29.

Highly suitable (Class S1) lands for growing chrysanthemum occur in an area of 25 ha (5%) and are distributed in the northwestern part of the microwatershed. Maximum area of about 244 ha (50%) is marginally suitable (Class S3) for growing chrysanthemum and are distributed in the major part of the microwatershed. They have moderate limitation of nutrient availability. An area of about 146 ha (30%) is not suitable (Class N1) for growing chrysanthemum and is distributed in the western and northwestern part of the microwatershed with severe limitation of nutrient availability.




Fig. 7.29 Land suitability map of Chrysanthemum

Table 7.1 Soil-Site Characteristics of Belagunda-3 Microwatershed

Soil Map Units	Climate (P) (mm)	Growing period (Days)	Drain-age Class	Soil depth (cm)	Soil texture		Gravelliness		AWC (mm/m)	Slope (%)	Erosion	pH	EC (dSm ⁻¹)	ESP (%)	CEC [Cmol (p ⁺)kg ⁻¹]	BS (%)
					Sur-face	Sub-surface	Surface (%)	Sub-surface (%)								
TMKiB2	866	150	MW	>150	sc	c	<15	<15	>200	1-3	Moderate	9.60	0.35	6.63	21.83	100
ANRiB2	866	150	MW	100-150	sc	c	<15	<15	>200	1-3	Moderate	10.17	0.365	7.08	19.90	100
VKSmB1	866	150	W	100-150	c	scl	<15	<15	>200	1-3	Moderate	9.1	0.586	3.97	17.57	100
BMDcB2	866	150	W	>150	sl	scl	<15	<15	151-200	1-3	Moderate	-	-	-	-	-
MDRiA1	866	150	W	>150	sc	scl	<15	<15	>200	0-1	Slight	8.31	0.33	0.90	20.57	100

*Symbols and abbreviations are according to Field Guide for LRI under Sujala-III Project, Karnataka

Table 7.2 Land suitability criteria for Sorghum

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	26–30	30–34; 24–26	34–40; 20–24	>40; <20
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, c (red), c (black)	scl, cl	ls, sl	-
	pH	1:2.5	5.5-7.8	5.0-5.5 7.8-9.0	>9.0	-
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	10-15
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

Table 7.3 Land suitability criteria for Maize

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	30-34	35-38 26-30	38-40 26-20	
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc	c (red), c (black)	ls, sl	-
	pH	1:2.5	5.5-7.8	5.0-5.5 7.8-9.0	>9.0	-
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	-
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

Table 7.4 Land suitability criteria for Bajra

Land use requirement		Rating				
Soil –site characteristics	Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)	
Climatic regime	Mean temperature in growing season	°C	28-32	33-38 24-27	39-40 20-23	<20
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm	500-750	400-500	200-400	<200
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl, scl, cl,sc,c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.8	5.0-5.5 7.8-9.0	5.5-6.0 >9.0	
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	15-35	35-60	>60	
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	1-3	3-5	5-10	>10

Table 7.5 Land suitability criteria for Groundnut

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	24–33	22–24; 33–35	20–22; 35–40	<20; >40
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. Well drained	Poorly drained	Very Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl	sl,cl, sc	c (red), c (black), ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<35	35-60	>60	
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.6 Land suitability criteria for Sunflower

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	24–30	30–34; 20–24	34–38; 16–20	>38; <16
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	mod. Well drained	-	Poorly to very drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	cl, sc,c (red), c (black)	scl	ls, sl	-
	pH	1:2.5	6.5-7.8	7.8-8.4 5.5-6.5	8.4-9.0; 5.0-5.5	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.7 Land suitability criteria for Redgram

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	30-35(G) 20-25(AV) 15-18 (F&PS) 35-40(M)	25-30(G) 20-25 (AV) 12-15 (F&PS) 30-35(M)	20-25(G) 15-20(AV) 10-12 (F&PS) 25-30(M)	< 20 <15 <10 <25
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. Well drained	Poorly drained	Very Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, c (red)	c (black),sl, scl, cl	ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-9.0	5.0-5.5 >9.0	-
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-50	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<1.0	1.0-2.0	>2.0	
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.8 Land suitability criteria for Bengal gram

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	20–25	25–30; 15–20	30–35; 10–15	>35; <10
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. Well drained	Poorly drained	Very Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	c(black)	-	c (red), scl, cl, sc	ls, sl
	pH	1:2.5	6.0-7.8	5.0-6.0 7.8-9.0	>9.0	-
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	-
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.9 Land suitability criteria for Cotton

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	22-32	>32	<19	-
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well to moderately well	Poorly drained/Some what excessively drained	-	very poorly/excessively drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, c (red,black)	cl	scl	ls, sl
	pH	1:2.5	6.5-7.8	7.8-8.4	5.5-6.5 8.4->9.0	<5.5
	CEC	C mol (p+)Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	50-100	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	<3	3-5	-	>5

Table 7.10 Land suitability criteria for Chilli

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	25-32	33-35 20-25	35-38 <20	>38
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc	c (black), sl	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.11 Land suitability criteria for Tomato

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	25-28	29-32 20-24	15-19 33-36	<15 >36
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl, scl, cl, sc, c (red)	-	ls, c(black)	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.12 Land suitability criteria for Brinjal

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	Well drained	Moderately well drained	Poorly drained	V. Poorly drained
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class				
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl, scl, cl, sc c (red)	-	ls, c (black)	-
	pH	1:2.5	6.0-7.3	7.3-8.4 5.0-6.0	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	>60
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.13 Land suitability criteria for Onion

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	20-30	30-35	35-40	>40
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately /imperfectly	-	Poorly to V poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl,scl,cl,sc,c (red)	-	c (Black),ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<1.0	1.0-2.0	2.0-4.0	<4
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.14 Land suitability criteria for Bhendi

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	25-28	29-32 20-24	15-19 33-36	<15 >36
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Imperfectly drained	Poorly to very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl,sc, c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.15 Land suitability criteria for Drumstick

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, scl, cl, c (red)	sl, c (black)	ls	s
	pH	1:2.5	6.0-7.3	5.0-5.5 7.3-7.8	5.5-6.0 7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<35	35-60	60-80	>80
Soil toxicity	Salinity (EC saturation extract)	ds/m				
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-10	-	>10

Table 7.16 Land suitability criteria for Mango

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-32	24-27 33-35	36-40	20-24
	Min temp. before flowering	°C	10-15	15-22	>22	-
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration	Days				
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V. Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	-	ls, sl, c (black)	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>150	100-150	75-100	<75
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.17 Land suitability criteria for Guava

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-32	33-36 24-27	37-42 20-23	
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	sl	c (black), ls	-
	pH	1:2.5	6.0-7.8	5.0-6.0	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.18 Land suitability criteria for Sapota

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-32	33-36 24-27	37-42 20-23	>42 <18
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	-	Poorly to very drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	sl	ls, c (black)	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.19 Land suitability criteria for Pomegranate

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	30-34	35-38 25-29	39-40 15-24	
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl,cl, sc, c (red)	c (black),sl	ls	-
	pH	1:2.5	5.5-7.8	7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.20 Land suitability criteria for Musambi

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-30	31-35 24-27	36-40 20-23	>40 <20
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately drained	poorly	Very poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c	sl	ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.21 Land suitability criteria for Lime

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-30	31-35 24-27	36-40 20-23	>40 <20
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately drained	poorly	Very poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c	sl	ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.22 Land suitability criteria for Amla

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. well drained	Poorly drained	V. Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	c (black)	ls, sl	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15-35	35-60	60-80	-
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

Table 7.23 Land suitability criteria for Cashew

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	32 to 34	28 to 32; 34 to 38	24 to 28; 38 to 40	<20; >40
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	-	sl, ls	c (black)
	pH	1:2.5	5.5-6.5	5.0-5.5 6.5-7.3	7.3-7.8	>7.8
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-10	>10	-

Table 7.24 Land suitability criteria for Jackfruit

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. well	Poorly	V. Poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	-	sl, ls, c (black)	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	>60
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10-

Table 7.25 Land suitability criteria for Jamun

Land use requirement		Rating				
Soil –site characteristics	Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)	
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well	Mod. well	Poorly	V.Poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c(red)	sl, c (black)	ls	-
	pH	1:2.5	6.0-7.8	5.0-6.0	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>150	100-150	50-100	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	>60
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

Table 7.26 Land suitability criteria for Custard apple

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	Scl, cl, sc, c (red), c (black)	-	Sl, ls	-
	pH	1:2.5	6.0-7.3	5.5-6.0 7.3-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15-35	35-60	60-80	-
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	>5	-

Table 7.27 Land suitability criteria for Tamarind

Land use requirement		Rating				
Soil –site characteristics	Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)	
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod.well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl,sc, c (red)	sl, c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>150	100-150	75-100	<75
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

Table 7.28 Land suitability criteria for Mulberry

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	24–28	22–24; 28–32	32–38; 22–18	>38; <18
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V. Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, cl, scl	c (red)	c (black), sl, ls	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.8-8.4	7.3-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO ₃ in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	0-35	35-60	60-80	>80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

Table 7.29 Land suitability criteria for Marigold

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	18-23	17-15 24-35	35-40 10-14	>40 <10
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl,scl, cl, sc, c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%				
Erosion hazard	Slope	%	<3	3-5	5-10	>10

Table 7.30 Land suitability criteria for Chrysanthemum

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	18-23	17-15 24-35	35-40 10-14	>40 <10
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl,scl, cl, sc, c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%				
Erosion hazard	Slope	%	<3	3-5	5-10	>10

7.30 Land Management Units (LMUs)

The 5 soil map units identified in Belagunda-3 microwatershed have been grouped into 2 Land Management Units (LMU's) for the purpose of preparing a Proposed Crop Plan. Land Management Units are grouped based on the similarities in respect of the type of soil, the depth of the soil, the surface soil texture, gravel content, AWC, slope, erosion etc. and a Land Management Units map (Fig. 7.30) has been generated. These Land Management Units are expected to behave similarly for a given level of management.

The 5 map units that have been grouped into 2 Land Management Units along with brief description of soil and site characteristics are given below.

LMU	Soil map units	Soil and site characteristics
1	104.TMKiB2 55.ANRiB2 100.VKSmB1	Deep to very deep (100 to >150cm), 1-3 % slopes, non-gravelly (<15%), slight to moderate erosion
2	64.BMDcB2 60.MDRiA1	Very deep (>150cm), 0-3 % slopes, non-gravelly (<15%), slight to moderate erosion




Fig. 7.3 Land Management Units Map-Belagunda-3 microwatershed

7.31 Proposed crop plan for Belagunda-3 microwatershed

After assessing the land suitability for the 29 crops, the Proposed Crop Plan has been prepared for the 2 identified LMUs by considering only the highly (Class S1) and moderately (Class S2) suitable lands for each of the 29 crops. The resultant proposed crop plan is presented below in Table 7.31.

Table 7.31 Proposed crop plan for Belagunda-3 microwatershed

LMU	Soil Map Units	Survey Number	Soil Characteristics	Field Crops/ Commercial crops	Horticulture Crops (Rainfed/Irrigated)	Suitable Interventions
1	104.TMKiB2 55.ANRiB2 100.VKSmB1 (Deep to very deep, sodic soils)	Anura.B: 74,129,131,132,133,134,135,136,137,138,139,140,141,148,149,150,151,152,153,154,155,156,157,158,164,165,166,167,168,169,170,171,172,173,174,175,176, 186,187,188	Deep to very deep (100 to >150cm), sodic soils 1-3 % slopes, non-gravelly (<15%), slight to moderate erosion	-	Agri-Silvi-Pasture Ber, Aonla, Acacia sp. Dhaincha, Rhodes grass, Para grass, Bermuda grass	Application of gypsum, iron pyrites and elemental sulphur. Addition of farm yard manures, green manures and providing subsurface drainage
2	64.BMDcB2 60.MDRiA1 (Very deep, sandy clay loam soils)	Anura.B: 98,99,100,101,102,103,104,105,106,107,108,109,110,111,112,125/1,125/2,126,127/1,127/2,128/2,130,142,143,144,145,146,147,159,160,161,162,163,177,178,179/1,179/2,179/3,180/1, 182 Belagundi: 1,178,179,180,182,183, 184,185,186,187,188,189, 190, 191,192	Very deep (>150cm), 0-3 % slopes, non-gravelly (<15%), slight to moderate erosion	Sunflower, Sorghum, Maize, Groundnut, Red gram, Bajra	Fruit crops: Mango, Musambi, Sapota, Tamarind, Pomegranate, Amla, Custard apple, Guava, Jackfruit, Jamun, Lime Vegetables: Tomato, Onion, Bhendi, Chilli, Brinjal, Drumstick, Coriander Flowers: Marigold, Chrysanthemum	Application of FYM, biofertilizers and micronutrients, drip irrigation, mulching, suitable soil and water conservation practices

SOIL HEALTH MANAGEMENT

8.1 Soil Health

Soil health is basic to plant health and plant health is basic to human and bovine health. Soil is fundamental to crop production. Without soil, no food could be produced nor would livestock be fed on a large scale. Because it is finite and fragile, soil is a precious resource that requires special care from its users.

Soil health or the capacity of the soil to function is critical to human survival. Soil health has been defined as: “the capacity of the soil to function as a living system without adverse effect on the ecosystem”. Healthy soils maintain a diverse community of soil organisms that help to form beneficial symbiotic associations with plant roots, recycle essential plant nutrients, improve soil structure with positive repercussions for soil, water and nutrient holding capacity and ultimately improve crop production and also contribute to mitigating climate change by maintaining or increasing its carbon content.

Functional interactions of soil biota with organic and inorganic components, air and water determine a soil’s potential to store and release nutrients and water to plants and to promote and sustain plant growth. Thus, maintaining soil health is vital to crop production and conserve soil resource base for sustaining agriculture.

The most important characteristics of a healthy soil are

- Good soil tilth
- Sufficient soil depth
- Good water storage and good drainage
- Adequate supply, but not excess of nutrients
- Large population of beneficial organisms
- Small proportion of plant pathogens and insect pests
- Low weed pressure
- Free of chemicals and toxins that may harm the crop
- Resistance to degradation
- Resilience when unfavorable conditions occur

Characteristics of Belagunda-3 microwatershed

- ❖ The soil phases identified in the microwatershed belonged to the soil series of MDR 244 ha (50%), VKS 89 ha (18%), TMK 56 ha (12%), BMD 25 ha (5%) and ANR 1 ha (<1%).
- ❖ As per land capability classification entire area of the microwatershed falls under arable land category (Class II & IV). The major limitations identified in the arable lands were soil and erosion.
- ❖ On the basis of soil reaction, an area of about 111 ha (23%) is moderately alkaline (pH 7.8- 8.4), 281 ha(58%) is strongly alkaline (pH 8.4- 9.0) and 22 ha(4%) is very strongly alkaline(pH > 9.0).

Soil Health Management

The following actions are required to improve the current land husbandry practices that provide a sound basis for the successful adoption of sustainable crop production system.

Acid soils

No acid soils found in this microwatershed.

1. Growing of crops suitable for particular soil pH.
2. Ameliorating the soils through the application of amendments (liming materials).

Liming materials:

1. CaCO_3 (Calcium Carbonate).
2. Dolomite [$\text{Ca Mg} (\text{CO}_3)_2$]
3. Quick lime (Cao)
4. Slaked lime [$\text{Ca} (\text{OH})_2$]

For normal pH and pH 4.8 (35 t/ha) and pH 6.0-7.0 (4 t/ha) lime is required.

Alkaline soils

Entire area in the microwatershed is alkaline in reaction.

1. Regular addition of organic manure, green manuring, green leaf manuring, crop residue incorporation and mulching needs to be taken up to improve the soil organic matter status.
2. Application of biofertilizers (Azospirillum, Azotobacter, Rhizobium).
3. Application of 25% extra N and P (125 % RDN&P).
4. Application of ZnSO_4 – 12.5 kg/ha (once in three years).
5. Application of Boron – 5kg/ha (once in three years).

Neutral soils

No neutral soil found in this microwatershed.

1. Regular addition of organic manure, green manuring, green leaf manuring, crop residue incorporation and mulching needs to be taken up to improve the soil organic matter status.
2. Application of biofertilizers, (Azospirillum, Azotobacter, Rhizobium).
3. Application of 100 per cent RDF.
4. Need based micronutrient applications.

Besides the above recommendations, the best transfer of technology options are also to be adopted.

Soil Degradation

Soil erosion is one of the major factor affecting the soil health in the microwatershed. Out of total 485 ha area in the microwatershed, an area of about 81 ha is suffering from moderate erosion and 333 ha is suffering from slight erosion. These areas

need immediate soil and water conservation and other land development and land husbandry practices for restoring soil health.

Dissemination of Information and Communication of Benefits

Any large scale implementation of soil health management requires that supporting information is made available widely, particularly through channels familiar to farmers and extension workers. Given the very high priority attached to soil-health especially by the Central Government on issuing Soil-Health Cards to all the farmers, media outlets like Regional, State and National Newspapers, Radio and Dooradarshan programs in local languages but also modern information and communication technologies such as Cellular phones and the Internet, which can be much more effective in reaching the younger farmers.

Inputs for Net Planning (Saturation Plan) and Interventions needed

Net planning (Saturation Plan) in IWMP is focusing on preparation of

1. Soil and Water Conservation Plan for each plot or farm.
2. Productivity enhancement measures/ interventions for existing crops/livestock/other farm enterprises.
3. Diversification of farming mainly with perennial horticultural crops and livestock.
4. Improving livelihood opportunities and income generating activities.

In this connection, how various outputs of Sujala-III are of use in addressing these objectives of Net Planning (Saturation Plan) are briefly presented below.

- ❖ **Soil Depth:** The depth of a soil decides the amount of moisture and nutrients it can hold, what crops can be taken up or not, depending on the rooting depth and the length of growing period available for raising any crop. Deeper the soil, better for a wide variety of crops. If sufficient depth is not available for growing deep rooted crops, either choose medium or short duration crops or deeper planting pits need to be opened and additional good quality soil brought from outside has to be filled into the planting pits.
- ❖ **Surface Soil Texture:** Lighter soil texture in the top soil means, better rain water infiltration, less run-off and soil moisture conservation, less capillary rise and less evaporation losses. Lighter surface textured soils are amenable to good soil tilth and are highly suitable for crops like groundnut, root vegetables (carrot, raddish, potato etc) but not ideal for crops that need stagnant water like lowland paddy. Heavy textured soils are poor in water infiltration and percolation. They are prone for sheet erosion; such soils can be improved by sand mulching. The technology that is developed by the AICRP-Dry land Agriculture, Vijayapura, Karnataka can be adopted.


- ❖ **Gravelliness:** More gravel content is favorable for run-off harvesting but poor in soil moisture storage and nutrient availability. It is a significant parameter that decides the kind of crop to be raised.
- ❖ **Land Capability Classification:** The land capability map shows the areas suitable and not suitable for agriculture and the major constraints in each of the plot/survey number. Hence, one can decide what kind of enterprise is possible in each of these units. In general, erosion and soil are the major constraints in Belagunda-3 microwatershed.
- ❖ **Organic Carbon:** The OC content (an index of available Nitrogen) is low (<0.5) in an area of 162 ha (33%) and medium (0.5 – 0.75 %) in an area of 252 ha (52%) of the microwatershed. The areas that are low and medium in OC needs to be further improved by applying farmyard manure and rotating crops with cereals and legumes or mixed cropping.
- ❖ **Promoting Green Manuring:** Growing of green manuring crops costs Rs. 1250/ha (green manuring seeds) and about Rs. 2000/ha towards cultivation that totals to Rs. 3250/- per ha. On the other hand, application of organic manure @ 10 tons/ha costs Rs. 5000/ha. The practice needs to be continued for 2-3 years or more. Nitrogen fertilizer needs to be supplemented by 25% in addition to the recommended level. For example, for rainfed maize, recommended level is 50 kg N per ha and an additional 12 kg /ha needs to be applied for all the crops grown in these plots.
- ❖ **Available Phosphorus:** Available phosphorus is low (<23 kg/ha) in an area of 152 ha (31%), medium (23-57 kg/ha) in 134 ha (28%) area and high (>57 kg/ha) in an area of 128 ha (26%) of the microwatershed. In medium and low areas, for all the crops 25% additional P needs to be applied.
- ❖ **Available Potassium:** Available potassium is medium (145-337 kg/ha) in an area of 291 ha (60%) of the microwatershed and high (>337 kg/ha) in 123 ha (25%). In medium areas, for all the crops 25% additional potassium needs to be applied.
- ❖ **Available Sulphur:** Available sulphur is a very critical nutrient for oilseed crops. It is low(<10 ppm) in 2 ha (70%), medium (10 - 20 ppm) in 27 ha (6%) and high (>20 ppm) in 385 ha (79%). Low and medium areas need to be applied with magnesium sulphate or gypsum or Factamphos (p) fertilizer (13% sulphur) for 2-3 years for the deficiency to be corrected.
- ❖ **Available Boron:** An area of 290 ha (60%) is medium (0.5 – 1.0ppm), 0.3 ha is low (<0.5 ppm) and 124 ha (26%) is high (>1.0 ppm) in the microwatershed. For medium and low areas, application of sodium borate @ 10 kg/ha as soil application or 0.2 % borax as foliar spray is recommended.
- ❖ **Available Iron:** An area of 138 ha (28%) is deficient (<4.5 ppm) and maximum area of 277 ha (57%) sufficient (>4.5 ppm) in available iron. For the deficient areas, iron sulphate @ 25 kg/ha need to be applied for 2-3 years.

- ❖ **Available Manganese:** All the soils in the microwatershed are sufficient (>1.0 ppm) in available manganese.
- ❖ **Available Copper:** All the soils in the microwatershed are sufficient (>0.2 ppm) in available copper.
- ❖ **Available Zinc:** All the soils in the microwatershed are deficient (<0.6 ppm) in available zinc content. Application of zinc sulphate @25 kg/ha is recommended for these areas.
- ❖ **Soil Alkalinity:** Entire area in the microwatershed is alkaline in reaction. These areas need application of gypsum and wherever calcium is in excess, iron pyrites and element sulphur can be recommended. Management practices like treating repeatedly with good quality water to drain out the excess salts and provision of subsurface drainage and growing of salt tolerant crops like Casuarina, Acacia, Neem, Ber etc, are recommended.
- ❖ **Land Suitability for Various Crops:** Areas that are highly, moderately and marginally suitable for growing various crops are indicated. Along with the suitability, various constraints that are limiting the productivity are also indicated. For example, in case of cotton, rooting depth, texture and calcareousness are the major constraints in various plots. With suitable management interventions, the productivity can be enhanced. In order to increase the water holding capacity of light textured soils, growing of green manure crops and application of organic manure is recommended.

SOIL AND WATER CONSERVATION TREATMENT PLAN

For preparing soil and water conservation treatment plan for Belagunda-3 microwatershed, the land resource inventory database generated under Sujala-III project has been transformed as information through series of interpretative (thematic) maps using soil phase map as a base. The various thematic maps (1:7920 scale) generated were

- Soil depth
- Surface soil texture
- Available water capacity
- Soil slope
- Soil gravelliness
- Land capability
- Present land use and land cover
- Crop suitability
- Rainfall
- Hydrology
- Water Resources
- Socio-economic data
- Contour plan with existing features- network of waterways, pottissa boundaries, cut up/ minor terraces etc.
- Cadastral map (1:7920 scale)
- Satellite imagery (1:7920 scale)



Apart from these, Hand Level/ Hydro Marker/ Dumpy Level/ Total Station and Kathedars' List to be collected.

Steps for Survey and Preparation of Treatment Plan

The boundaries of Land User Groups' and Survey No. boundaries are traced in the field.

- Naming of user groups and farmers
- Identification of arable and non arable lands
- Identification of drainage lines and gullies
- Identification of non treatable areas
- Identification of priority areas in the arable lands
- Treatment plan for arable lands
- Location of water harvesting and recharge structures

9.1 Treatment Plan

The treatment plan recommended for arable lands is briefly described below

9.1.1 Arable Land Treatment

A. BUNDING

Steps for Survey and Preparation of Treatment Plan	
<ul style="list-style-type: none"> • Cadastral map (1:7920 scale) is enlarged to a scale of 1:2500 scale • Existing network of waterways, pottissa boundaries, grass belts, natural drainage lines/ watercourse, cut ups/ terraces are marked on the cadastral map to the scale • Drainage lines are demarcated into 	
Small gullies	(up to 5 ha catchment)
Medium gullies	(5-15 ha catchment)
Ravines	(15-25 ha catchment) and
Halla/Nala	(more than 25ha catchment)

USER GROUP-1

CLASSIFICATION OF GULLIES

ಕೊರಕಾಲಿನ ವರ್ಗೀಕರಣ

UPPER REACH: 15 Ha.

MIDDLE REACH: 15 + 10 = 25 ಹೆ.

LOWER REACH: 25 ಹೆಕ್ಟಾರ್ ನಿಂತ ಅಧಿಕ

POINT OF CONCENTRATION

Measurement of Land Slope

Land slope is estimated or determined by the study and interpretation of contours or by measurement in the field using simple instruments like Hand Level or Hydromarker.



Vertical and Horizontal intervals between bunds as recommended by the Watershed Development Department.

Slope percentage	Vertical interval (m)	Corresponding Horizontal Distance (m)
2 - 3%	0.6	24
3 - 4%	0.9	21
4 - 5%	0.9	21
5 - 6%	1.2	21
6 - 7%	1.2	21

Note: (i) The above intervals are maximum.

(ii) Considering the slope class and erosion status (A1... A=0-1 % slope, 1= slight erosion) the intervals have to be decided.

Bund length recording: Considering the contour plan and the existing grass belts/partitions, the bunds are aligned and lengths are measured.

Section of the Bund

Bund section is decided considering the soil texture class and gravelliness class (bg₀... b=loamy sand, g₀ = <15% gravel). The recommended Sections for different soils are given below.

Recommended Bund Section

Top width (m)	Base width (m)	Height (m)	Side slope (Z:1;H:V)	Cross section (sq m)	Soil Texture	Remarks
0.3	0.9	0.3	01:01	0.18	Sandy loam	Vegetative bund
0.3	1.2	0.3	1.5:1	0.225	Sandy clay	
0.3	1.2	0.5	0.9:1	0.375	Red gravelly soils	
0.3	1.2	0.6	0.75:1	0.45		
0.3	1.5	0.6	01:01	0.54	Red sandy loam	
0.3	2.1	0.6	1.5:1	0.72	Very shallow black soils	
0.45	2	0.75	01:01	0.92		
0.45	2.4	0.75	1.3:1	1.07	Shallow black soils	
0.6	3.1	0.7	1.78:1	1.29	Medium black soils	
0.5	3	0.85	1.47:1	1.49		

Formation of Trench cum Bund

Dimensions of the Borrow Pits/Trenches to be excavated (machinery are decided considering the Bund Section).

Details of Borrow Pit dimensions are given below:

TRENCH CUM BUND

WATER STORAGE AREA

0.45 Sq.m section

IDEAL FOR HORTICULTURE CROPS

'A' FRAME FOR INTERBUND MANAGEMENT

1. ಸಮಸಾತಳ ಉಳಂಪು

2. ಸಮಸಾತಳ ಬಿತ್ತನೆ/ನಾಟಿ

Size of Borrow Pits/ Trench recommended for Trench cum Bund (by machinery)

Bund section	Bund length	Earth quantity	Pit				Berm (pit to pit)	Soil depth class
			L(m)	W(m)	D(m)	Quantity (m ³)		
m ²	m	m ³					m	
0.375	6	2.25	5.85	0.85	0.45	2.24	0.15	Shallow
0.45	6	2.7	5.4	1.2	0.43	2.79	0.6	Shallow
0.45	6	2.7	5	0.85	0.65	2.76	1	Moderately Shallow
0.54	5.6	3.02	5.5	0.85	0.7	3.27	0.1	Moderately shallow
0.54	5.5	2.97	5	1.2	0.5	3	0.5	Shallow
0.72	6.2	4.46	6	1.2	0.7	5.04	0.2	Moderately shallow
0.72	5.2	3.74	5.1	0.85	0.9	3.9	0.1	Moderately deep

B. Water Ways

1. Existing waterways are marked on the cadastral map (1:7920 scale) and their dimensions are recorded.
2. Considering the contour plan of the MWS, additional waterways/ modernization of the existing ones can be thought of.
3. The design details are given in the Manual.

C. Farm Ponds

Waterways and the catchment area will give an indication on the size of the Farm Pond. Location of the pond can be decided based on the contour plan/ field condition and farmers' need/desire.

D. Diversion Channel

Existing EPT/ CPT are marked on the cadastral map. Looking to the need, these can be modernized or fresh diversion channel can be proposed and runoff from this can be stored in Gokatte/ Recharge ponds.

9.1.2 Non-Arable Land Treatment

Depending on the gravelliness and crops preferred by the farmers, the concerned authorities can decide appropriate treatment plan. The recommended treatments may be Contour Trench, Staggered Trench, Crescent Bund, Boulder Bund or Pebble Bund.

9.1.3 Treatment of Natural Water Course/ Drainage Lines

- a) The cadastral map has to be updated as regards the network of drainage lines (gullies/ *nalas*/ *hallas*) and existing structures are marked to the scale and storage capacity of the existing water bodies are documented.
- b) The drainage line will be demarcated into Upper Reach, Middle Reach and Lower Reach.
- c) Considering the Catchment, *Nala* bed and bank conditions, suitable structures are decided.
- d) Number of storage structures (Check dam/ *Nala* bund/ Percolation tank) will be decided considering the commitments and available runoff from water budgeting and quality of water in the wells and site suitability.
- e) Detailed Leveling Survey using Dumpy Level / Total Station has to be carried out to arrive at the site-specific designs as shown in the Manual.
- f) The location of ground water recharge structures are decided by examining the lineaments and fracture zones from geological maps.
- g) Rainfall intensity data of the nearest Rain Gauge Station is considered for Hydrologic Designs.
- h) Silt load to the Storage/Recharge Structures is reduced by providing vegetative, boulder and earthen checks in the natural water course. Location and design details are given in the Manual.

9.2 Recommended Soil and Water Conservation Measures

The appropriate conservation structures best suited for each of the land parcel/ survey number (Appendix-I) are selected based on the slope per cent, severity of erosion, amount of rainfall, land use and soil type. The different kinds of conservation structures recommended are:

1. Graded / Strengthening of Bunds
2. Trench cum Bunds (TCB)
3. Trench cum Bunds / Strengthening
4. Crescent Bunds

A map (Fig. 9.1) showing soil and water conservation plan with different kinds of structures recommended has been prepared which shows the spatial distribution and extent of area. An area of about 25 ha (5%) needs trench cum bunding, 244 ha (50%) needs strengthening of existing bunds and an area of about 146 ha (30%) needs graded bunding.

The conservation plan prepared may be presented to all the stakeholders including farmers and after considering their suggestions, the conservation plan for the microwatershed may be finalised in a participatory approach.




Fig. 9.1 Soil and water conservation plan map of Belagunda-3 microwatershed

9.3 Greening of microwatershed

As part of the greening programme in the watersheds, it is envisaged to plant a variety of horticultural and other tree plants that are edible, economical and produce lot of biomass which helps to restore the ecological balance in the watersheds. The lands that are suitable for greening programme are non-arable lands (land capability classes V, VI VII and VIII) and also the lands that are not suitable or marginally suitable and field bunds for growing annual and perennial crops. The method of planting these trees is given below.

It is recommended to open pits during the 1st week of March along the contour and heap the dugout soil on the lower side of the slope in order to harness the flowing water and facilitate weathering of soil in the pit. Exposure of soil in the pit also prevents spread of pests and diseases due to scorching sun rays. The pits should be filled with mixture of soil and organic manure during the second week of April and keep ready with sufficiently tall seedlings produced either in poly bags or in root trainer nurseries so that planting can be done during the 2nd or 3rd week of April depending on the rainfall.

The tree species suitable for the area considering rainfall, temperature and adaptability is listed below; waterlogged areas are recommended to be planted with species like Nerale (*Sizyzium cumini*) and Bamboo. Dry areas are to be planted with species like Honge, Bevu, Seetaphal etc.

Dry deciduous species			Temp (°C)	Rainfall (mm)
1.	Bevu	<i>Azadiracta indica</i>	21–32	400 –1,200
2.	Tapasi	<i>Holoptelia integrifolia</i>	20-30	500 - 1000
3.	Seetaphal	<i>Anona Squamosa</i>	20-40	400 - 1000
4.	Honge	<i>Pongamia pinnata</i>	20 -50	500– 2,500
5.	Kamara	<i>Hardwickia binata</i>	25 -35	400 - 1000
6.	Bage	<i>Albezzia lebbek</i>	20 - 45	500 - 1000
7.	Ficus	<i>Ficus bengalensis</i>	20 - 50	500–2,500
8.	Sisso	<i>Dalbargia Sissoo</i>	20 - 50	500 -2000
9.	Ailanthus	<i>Ailanthus excelsa</i>	20 - 50	500 - 1000
10.	Hale	<i>Wrightia tinctoria</i>	25 - 45	500 - 1000
11.	Uded	<i>Steriospermum chelanoides</i>	25 - 45	500 -2000
12.	Dhupa	<i>Boswellia Serrata</i>	20 - 40	500 - 2000
13.	Nelli	<i>Emblica Officinalis</i>	20 - 50	500 -1500
14.	Honne	<i>Pterocarpus marsupium</i>	20 - 40	500 - 2000
Moist Deciduous Species			Temp (°C)	Rainfall (mm)
15.	Teak	<i>Tectona grandis</i>	20 - 50	500-5000
16.	Nandi	<i>Legarstroemia lanceolata</i>	20 - 40	500 - 4000
17.	Honne	<i>Pterocarpus marsupium</i>	20 - 40	500 - 3000
18.	Mathi	<i>Terminalia alata</i>	20 -50	500 - 2000
19.	Shivane	<i>Gmelina arborea</i>	20 -50	500 -2000
20.	Kindal	<i>T.Paniculata</i>	20 - 40	500 - 1500
21.	Beete	<i>Dalbargia latifolia</i>	20 - 40	500 - 1500
22.	Tare	<i>T. belerica</i>	20 - 40	500 - 2000
23.	Bamboo	<i>Bambusa arundinasia</i>	20 - 40	500 - 2500
24.	Bamboo	<i>Dendrocalamus strictus</i>	20 – 40	500 – 2500
25.	Muthuga	<i>Butea monosperma</i>	20 - 40	400 - 1500
26.	Hippe	<i>Madhuca latifolia</i>	20 - 40	500 - 2000
27.	Sandal	<i>Santalum album</i>	20 - 50	400 - 1000
28.	Nelli	<i>Emblica officinalis</i>	20 - 40	500 - 2000
29.	Nerale	<i>Sizyzium cumini</i>	20 - 40	500 - 2000
30.	Dhaman	<i>Grevia tilifolia</i>	20 - 40	500 - 2000
31.	Kaval	<i>Careya arborea</i>	20 - 40	500 - 2000
32.	Harada	<i>Terminalia chebula</i>	20 - 40	500 - 2000

References

1. FAO (1976) Framework for Land Evaluation, Food and Agriculture Organization, Rome.72 pp.
2. FAO (1983) Guidelines for Land Evaluation for Rainfed Agriculture, FAO, Rome, 237 pp.
3. IARI (1971) Soil Survey Manual, All India Soil and Land Use Survey Organization, IARI, New Delhi, 121 pp.
4. Katyal, J.C. and Rattan, R.K. (2003) Secondary and Micronutrients; Research Gap and Future Needs. Fert. News 48 (4); 9-20.
5. Naidu, L.G.K., Ramamurthy, V., Challa, O., Hegde, R. and Krishnan, P. (2006) Manual Soil Site Suitability Criteria for Major Crops, NBSS Publ. No. 129, NBSS & LUP, Nagpur, 118 pp.
6. Natarajan, A. and Dipak Sarkar (2010) Field Guide for Soil Survey, National Bureau of Soil Survey and Land Use Planning (ICAR), Nagpur, India.
7. Natarajan, A., Rajendra Hegde, Raj, J.N. and Shivananda Murthy, H.G. (2015) Implementation Manual for Sujala-III Project, Watershed Development Department, Bengaluru, Karnataka.
8. Sarma, V.A.K., Krishnan, P. and Budihal, S.L. (1987) Laboratory Manual, Tech. Bull. 23, NBSS &LUP, Nagpur.
9. Sehgal, J.L. (1990) Soil Resource Mapping of Different States of India; Why and How?, National Bureau of Soil Survey and Land Use Planning, Nagpur, 49 pp.
10. Shivaprasad, C.R., R.S. Reddy, J. Sehgal and M. Velayuthum (1998) Soils of Karnataka for Optimizing Land Use, NBSS Publ. No. 47b, NBSS & LUP, Nagpur, India.
11. Soil Survey Staff (2006) Keys to Soil Taxonomy, Tenth edition, U.S. Department of Agriculture/ NRCS, Washington DC, U.S.A.
12. Soil Survey Staff (2012) Soil Survey Manual, Handbook No. 18, USDA, Washington DC, USA.

Appendix I
Belagunda-3 (4D5B1N1d) Microwatershed
Soil Phase Information

Village	Survey Number	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Machanura	RIVER_BHIMA	3.29	Waterbody	Others	Others	Others	Others	Others	Others	Others	Waterbody	Not Available	Others	Others
Bilhara	RIVER_BHIMA	0.69	Waterbody	Others	Others	Others	Others	Others	Others	Others	Waterbody	Not Available	Others	Others
Belagundi	1	0.15	MDRiA1	LMU-2	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Paddy (Pd)	Not Available	IIs	Graded bunding
Belagundi	178	5.53	MDRiA1	LMU-2	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Paddy (Pd)	Not Available	IIs	Graded bunding
Belagundi	179	1.91	MDRiA1	LMU-2	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Paddy (Pd)	Not Available	IIs	Graded bunding
Belagundi	180	0.64	MDRiA1	LMU-2	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Paddy (Pd)	Not Available	IIs	Graded bunding
Belagundi	182	1.95	MDRiA1	LMU-2	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Paddy (Pd)	Not Available	IIs	Graded bunding
Belagundi	183	6.63	MDRiA1	LMU-2	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Paddy (Pd)	Not Available	IIs	Graded bunding
Belagundi	184	4.34	MDRiA1	LMU-2	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Paddy (Pd)	Not Available	IIs	Graded bunding
Belagundi	185	4.88	MDRiA1	LMU-2	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Paddy (Pd)	Not Available	IIs	Graded bunding
Belagundi	186	5.67	MDRiA1	LMU-2	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Paddy (Pd)	Not Available	IIs	Graded bunding
Belagundi	187	3.43	MDRiA1	LMU-2	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Paddy (Pd)	Not Available	IIs	Graded bunding
Belagundi	188	6.56	MDRiA1	LMU-2	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Paddy (Pd)	Not Available	IIs	Graded bunding
Belagundi	189	4.15	MDRiA1	LMU-2	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Paddy (Pd)	Not Available	IIs	Graded bunding
Belagundi	190	6.49	MDRiA1	LMU-2	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Paddy (Pd)	Not Available	IIs	Graded bunding
Belagundi	191	7.8	MDRiA1	LMU-2	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Paddy (Pd)	Not Available	IIs	Graded bunding
Belagundi	192	5.44	MDRiA1	LMU-2	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Paddy (Pd)	Not Available	IIs	Graded bunding
Anura B	74	0.73	ANRiB2	LMU-1	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IVes	Graded bunding
Anura B	98	0.57	MDRiA1	LMU-2	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Cotton (Ct)	Not Available	IIs	Graded bunding
Anura B	99	7.11	MDRiA1	LMU-2	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Cotton (Ct)	Not Available	IIs	Graded bunding
Anura B	100	4.08	MDRiA1	LMU-2	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Scrub land (SI)	Not Available	IIs	Graded bunding
Anura B	101	0.45	MDRiA1	LMU-2	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Scrub land (SI)	Not Available	IIs	Graded bunding

Village	Survey Number	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Anura B	102	3.35	MDRiA1	LMU-2	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Redgram (Rg)	Not Available	IIs	Graded bunding
Anura B	103	6.54	MDRiA1	LMU-2	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Paddy (Pd)	Not Available	IIs	Graded bunding
Anura B	104	4.85	MDRiA1	LMU-2	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Paddy (Pd)	Not Available	IIs	Graded bunding
Anura B	105	3.18	MDRiA1	LMU-2	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Paddy (Pd)	Not Available	IIs	Graded bunding
Anura B	106	6.73	MDRiA1	LMU-2	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Paddy (Pd)	Not Available	IIs	Graded bunding
Anura B	107	7.7	MDRiA1	LMU-2	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Paddy (Pd)	Not Available	IIs	Graded bunding
Anura B	108	4.75	MDRiA1	LMU-2	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Paddy (Pd)	Not Available	IIs	Graded bunding
Anura B	109	5.01	MDRiA1	LMU-2	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Paddy (Pd)	Not Available	IIs	Graded bunding
Anura B	110	4.13	MDRiA1	LMU-2	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Paddy (Pd)	Not Available	IIs	Graded bunding
Anura B	111	2.21	MDRiA1	LMU-2	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Paddy (Pd)	Not Available	IIs	Graded bunding
Anura B	112	3.22	MDRiA1	LMU-2	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Paddy (Pd)	Not Available	IIs	Graded bunding
Anura B	125/1	2.05	BMDcB2	LMU-2	Very deep (>150 cm)	Sandy loam	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	IIs	TCB
Anura B	125/2	0.45	BMDcB2	LMU-2	Very deep (>150 cm)	Sandy loam	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	IIs	TCB
Anura B	126	1.26	BMDcB2	LMU-2	Very deep (>150 cm)	Sandy loam	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	IIs	TCB
Anura B	127/1	0.93	BMDcB2	LMU-2	Very deep (>150 cm)	Sandy loam	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	IIs	TCB
Anura B	127/2	0.52	BMDcB2	LMU-2	Very deep (>150 cm)	Sandy loam	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	IIs	TCB
Anura B	128/2	0.18	BMDcB2	LMU-2	Very deep (>150 cm)	Sandy loam	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIs	TCB
Anura B	129	1.72	VKSmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Paddy (Pd)	Not Available	IVs	Graded bunding
Anura B	130	4.61	MDRiA1	LMU-2	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Paddy (Pd)	Not Available	IIs	Graded bunding
Anura B	131	4.35	VKSmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Paddy (Pd)	Not Available	IVs	Graded bunding
Anura B	132	4.16	VKSmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Paddy (Pd)	Not Available	IVs	Graded bunding
Anura B	133	2.39	TMKiB2	LMU-1	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	IVs	Graded bunding
Anura B	134	4.71	VKSmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Cotton (Ct)	Not Available	IVs	Graded bunding
Anura B	135	3.15	VKSmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Cotton (Ct)	Not Available	IVs	Graded bunding

Village	Survey Number	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Anura B	136	2.94	VKSmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Cotton (Ct)	Not Available	IVs	Graded bunding
Anura B	137	3.57	VKSmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Cotton (Ct)	Not Available	IVs	Graded bunding
Anura B	138	1.53	VKSmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Paddy (Pd)	Not Available	IVs	Graded bunding
Anura B	139	0.82	VKSmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Paddy (Pd)	Not Available	IVs	Graded bunding
Anura B	140	4.53	VKSmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Paddy (Pd)	Not Available	IVs	Graded bunding
Anura B	141	3.59	VKSmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Cotton (Ct)	Not Available	IVs	Graded bunding
Anura B	142	2.43	MDRiA1	LMU-2	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Paddy (Pd)	Not Available	IIs	Graded bunding
Anura B	143	5.61	MDRiA1	LMU-2	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Paddy (Pd)	Not Available	IIs	Graded bunding
Anura B	144	4.32	MDRiA1	LMU-2	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Cotton (Ct)	Not Available	IIs	Graded bunding
Anura B	145	5.08	MDRiA1	LMU-2	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Cotton (Ct)	Not Available	IIs	Graded bunding
Anura B	146	6.59	MDRiA1	LMU-2	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Cotton (Ct)	Not Available	IIs	Graded bunding
Anura B	147	4.77	MDRiA1	LMU-2	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Paddy (Pd)	Not Available	IIs	Graded bunding
Anura B	148	4.06	VKSmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Paddy (Pd)	Not Available	IVs	Graded bunding
Anura B	149	3.73	VKSmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Paddy (Pd)	Not Available	IVs	Graded bunding
Anura B	150	5.75	VKSmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Paddy (Pd)	Not Available	IVs	Graded bunding
Anura B	151	3.07	VKSmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Paddy (Pd)	Not Available	IVs	Graded bunding
Anura B	152	1.86	VKSmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Paddy (Pd)	Not Available	IVs	Graded bunding
Anura B	153	3.45	TMKiB2	LMU-1	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	IVes	Graded bunding
Anura B	154	1.44	VKSmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Paddy (Pd)	Not Available	IVs	Graded bunding
Anura B	155	1.63	TMKiB2	LMU-1	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	IVes	Graded bunding
Anura B	156	1.05	TMKiB2	LMU-1	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	IVes	Graded bunding
Anura B	157	4.45	VKSmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Paddy (Pd)	Not Available	IVs	Graded bunding
Anura B	158	7.35	VKSmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Paddy (Pd)	Not Available	IVs	Graded bunding
Anura B	159	4.25	BMDcB2	LMU-2	Very deep (>150 cm)	Sandy loam	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton+Jowar (Ct+Jw)	Not Available	IIs	TCB

Village	Survey Number	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Anura B	160	1.48	BMDcB2	LMU-2	Very deep (>150 cm)	Sandy loam	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	Iles	TCB
Anura B	161	0.11	BMDcB2	LMU-2	Very deep (>150 cm)	Sandy loam	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iles	TCB
Anura B	162	4.66	BMDcB2	LMU-2	Very deep (>150 cm)	Sandy loam	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Iles	TCB
Anura B	163	4.29	BMDcB2	LMU-2	Very deep (>150 cm)	Sandy loam	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Iles	TCB
Anura B	164	4.45	TMKiB2	LMU-1	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	IVes	Graded bunding
Anura B	165	6.94	TMKiB2	LMU-1	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IVes	Graded bunding
Anura B	166	4.32	VKSmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Paddy (Pd)	Not Available	IVs	Graded bunding
Anura B	167	4.59	VKSmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Paddy (Pd)	Not Available	IVs	Graded bunding
Anura B	168	3.52	VKSmB1	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Slight	Paddy (Pd)	Not Available	IVs	Graded bunding
Anura B	169	3.93	TMKiB2	LMU-1	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IVes	Graded bunding
Anura B	170	3.2	TMKiB2	LMU-1	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IVes	Graded bunding
Anura B	171	8.1	TMKiB2	LMU-1	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	IVes	Graded bunding
Anura B	172	3.49	TMKiB2	LMU-1	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	IVes	Graded bunding
Anura B	173	3.1	TMKiB2	LMU-1	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	IVes	Graded bunding
Anura B	174	1.6	TMKiB2	LMU-1	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	IVes	Graded bunding
Anura B	175	2.29	TMKiB2	LMU-1	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	IVes	Graded bunding
Anura B	176	2.77	TMKiB2	LMU-1	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	IVes	Graded bunding
Anura B	177	4.38	BMDcB2	LMU-2	Very deep (>150 cm)	Sandy loam	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	Iles	TCB
Anura B	178	0.19	BMDcB2	LMU-2	Very deep (>150 cm)	Sandy loam	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iles	TCB
Anura B	179/1	1.18	BMDcB2	LMU-2	Very deep (>150 cm)	Sandy loam	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	Iles	TCB
Anura B	179/2	0.35	BMDcB2	LMU-2	Very deep (>150 cm)	Sandy loam	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iles	TCB
Anura B	179/3	1.66	BMDcB2	LMU-2	Very deep (>150 cm)	Sandy loam	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	Iles	TCB
Anura B	180/1	1.15	BMDcB2	LMU-2	Very deep (>150 cm)	Sandy loam	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Iles	TCB
Anura B	181	0.12	Habitatio n	Others	Others	Others	Others	Others	Others	Others	Not Available (NA)	Not Available	Others	Others

Village	Survey Number	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Anura B	182	0.02	BMDcB2	LMU-2	Very deep (>150 cm)	Sandy loam	Non gravelly (<15%)	Medium (101-150 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	IIs	TCB
Anura B	183	0.14	Habitatio n	Others	Others	Others	Others	Others	Others	Others	Not Available (NA)	Not Available	Others	Others
Anura B	186	0.51	TMKiB2	LMU-1	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar+Paddy (Jw+Pd)	Not Available	IVes	Graded bunding
Anura B	187	0.45	TMKiB2	LMU-1	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IVes	Graded bunding
Anura B	188	1.24	TMKiB2	LMU-1	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	IVes	Graded bunding
NA	NA	79.32	MDRiA1	LMU-2	Very deep (>150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Not Available (NA)	Not Available	IIs	Graded bunding

Appendix II
Belagunda-3 (4D5B1N1d) Microwatershed
Soil Fertility Information

Village	Survey Number	Soil Reaction	Salinity	Organic Carbon	Available Phosphorus	Available Potassium	Available Sulphur	Available Boron	Available Iron	Available Manganese	Available Copper	Available Zinc
Machanura	RIVER_BHIMA	Soil Reaction	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Bilhara	RIVER_BHIMA	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Belagundi	1	Others	Medium (4 – 8 dsm)	Low (< 0.5 %)	Low (< 23 kg/ha)	High (> 337 kg/ha)	High (> 20 ppm)	High (> 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Belagundi	178	Moderately alkaline (pH 7.8 – 8.4)	Low (2 – 4 dsm)	Medium (0.5 – 0.75 %)	Low (< 23 kg/ha)	Medium (145 – 337 kg/ha)	High (> 20 ppm)	High (> 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Belagundi	179	Moderately alkaline (pH 7.8 – 8.4)	Low (2 – 4 dsm)	Medium (0.5 – 0.75 %)	Low (< 23 kg/ha)	Medium (145 – 337 kg/ha)	High (> 20 ppm)	High (> 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Belagundi	180	Moderately alkaline (pH 7.8 – 8.4)	Low (2 – 4 dsm)	Low (< 0.5 %)	Low (< 23 kg/ha)	Medium (145 – 337 kg/ha)	High (> 20 ppm)	High (> 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Belagundi	182	Moderately alkaline (pH 7.8 – 8.4)	Medium (4 – 8 dsm)	Low (< 0.5 %)	Low (< 23 kg/ha)	Medium (145 – 337 kg/ha)	High (> 20 ppm)	High (> 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Belagundi	183	Moderately alkaline (pH 7.8 – 8.4)	Medium (4 – 8 dsm)	Low (< 0.5 %)	Low (< 23 kg/ha)	Medium (145 – 337 kg/ha)	High (> 20 ppm)	High (> 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Belagundi	184	Moderately alkaline (pH 7.8 – 8.4)	Medium (4 – 8 dsm)	Low (< 0.5 %)	Low (< 23 kg/ha)	Medium (145 – 337 kg/ha)	High (> 20 ppm)	High (> 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Belagundi	185	Moderately alkaline (pH 7.8 – 8.4)	Medium (4 – 8 dsm)	Low (< 0.5 %)	Low (< 23 kg/ha)	Medium (145 – 337 kg/ha)	High (> 20 ppm)	High (> 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Belagundi	186	Moderately alkaline (pH 7.8 – 8.4)	Low (2 – 4 dsm)	Medium (0.5 – 0.75 %)	Low (< 23 kg/ha)	Medium (145 – 337 kg/ha)	High (> 20 ppm)	High (> 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Belagundi	187	Moderately alkaline (pH 7.8 – 8.4)	Low (2 – 4 dsm)	Medium (0.5 – 0.75 %)	Low (< 23 kg/ha)	Medium (145 – 337 kg/ha)	High (> 20 ppm)	High (> 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Belagundi	188	Moderately alkaline (pH 7.8 – 8.4)	Low (2 – 4 dsm)	Medium (0.5 – 0.75 %)	Low (< 23 kg/ha)	Medium (145 – 337 kg/ha)	High (> 20 ppm)	High (> 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Belagundi	189	Moderately alkaline (pH 7.8 – 8.4)	Medium (4 – 8 dsm)	Medium (0.5 – 0.75 %)	Low (< 23 kg/ha)	Medium (145 – 337 kg/ha)	High (> 20 ppm)	High (> 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Belagundi	190	Moderately alkaline (pH 7.8 – 8.4)	Low (2 – 4 dsm)	Medium (0.5 – 0.75 %)	Low (< 23 kg/ha)	Medium (145 – 337 kg/ha)	High (> 20 ppm)	High (> 1.0 ppm)	Sufficient (> 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)

Village	Survey Number	Soil Reaction	Salinity	Organic Carbon	Available Phosphorus	Available Potassium	Available Sulphur	Available Boron	Available Iron	Available Manganese	Available Copper	Available Zinc
Belagundi	191	Strongly alkaline (pH 8.4 - 9.0)	Medium (4 - 8 dsm)	Medium (0.5 - 0.75 %)	Low (< 23 kg/ha)	High (> 337 kg/ha)	High (> 20 ppm)	High (> 1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Belagundi	192	Strongly alkaline (pH 8.4 - 9.0)	Low (2 - 4 dsm)	Medium (0.5 - 0.75 %)	Low (< 23 kg/ha)	Medium (145 - 337 kg/ha)	High (> 20 ppm)	High (> 1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Anura B	74	Strongly alkaline (pH 8.4 - 9.0)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Low (< 23 kg/ha)	Medium (145 - 337 kg/ha)	High (> 20 ppm)	Medium (0.5 - 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Anura B	98	Strongly alkaline (pH 8.4 - 9.0)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Low (< 23 kg/ha)	Medium (145 - 337 kg/ha)	High (> 20 ppm)	Medium (0.5 - 1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Anura B	99	Strongly alkaline (pH 8.4 - 9.0)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	High (> 20 ppm)	Medium (0.5 - 1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Anura B	100	Strongly alkaline (pH 8.4 - 9.0)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	High (> 337 kg/ha)	High (> 20 ppm)	Medium (0.5 - 1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Anura B	101	Strongly alkaline (pH 8.4 - 9.0)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	High (> 337 kg/ha)	High (> 20 ppm)	Medium (0.5 - 1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Anura B	102	Strongly alkaline (pH 8.4 - 9.0)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	High (> 337 kg/ha)	High (> 20 ppm)	Medium (0.5 - 1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Anura B	103	Strongly alkaline (pH 8.4 - 9.0)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	High (> 337 kg/ha)	High (> 20 ppm)	Medium (0.5 - 1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Anura B	104	Strongly alkaline (pH 8.4 - 9.0)	Non saline (<2 dsm)	Low (< 0.5 %)	Medium (23 - 57 kg/ha)	High (> 337 kg/ha)	High (> 20 ppm)	Medium (0.5 - 1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Anura B	105	Very strongly alkaline (pH > 9.0)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	High (> 337 kg/ha)	High (> 20 ppm)	Medium (0.5 - 1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Anura B	106	Strongly alkaline (pH 8.4 - 9.0)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	High (> 337 kg/ha)	High (> 20 ppm)	Medium (0.5 - 1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Anura B	107	Strongly alkaline (pH 8.4 - 9.0)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	High (> 337 kg/ha)	High (> 20 ppm)	Medium (0.5 - 1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Anura B	108	Strongly alkaline (pH 8.4 - 9.0)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	High (> 337 kg/ha)	High (> 20 ppm)	Medium (0.5 - 1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Anura B	109	Strongly alkaline (pH 8.4 - 9.0)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	High (> 57 kg/ha)	High (> 337 kg/ha)	High (> 20 ppm)	Medium (0.5 - 1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Anura B	110	Strongly alkaline (pH 8.4 - 9.0)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	High (> 57 kg/ha)	High (> 337 kg/ha)	High (> 20 ppm)	Medium (0.5 - 1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)

Village	Survey Number	Soil Reaction	Salinity	Organic Carbon	Available Phosphorus	Available Potassium	Available Sulphur	Available Boron	Available Iron	Available Manganese	Available Copper	Available Zinc
		(pH 7.8 - 8.4)										
Anura B	181	Moderately alkaline (pH 7.8 - 8.4)	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Anura B	182	Others	Non saline (<2 dsm)	Low (< 0.5 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	Medium (0.5 - 1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Anura B	183	Moderately alkaline (pH 7.8 - 8.4)	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Anura B	186	Others	Non saline (<2 dsm)	Low (< 0.5 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	Medium (10 - 20 ppm)	Medium (0.5 - 1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Anura B	187	Moderately alkaline (pH 7.8 - 8.4)	Non saline (<2 dsm)	Low (< 0.5 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	Medium (10 - 20 ppm)	Medium (0.5 - 1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Anura B	188	Strongly alkaline (pH 8.4 - 9.0)	Non saline (<2 dsm)	Low (< 0.5 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	Medium (10 - 20 ppm)	Medium (0.5 - 1.0 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
NA	NA	Strongly alkaline (pH 8.4 - 9.0)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Low (< 23 kg/ha)	Medium (145 - 337 kg/ha)	High (> 20 ppm)	Medium (0.5 - 1.0 ppm)	Deficient (< 4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)

Appendix III
Belagunda-3 (4D5B1N1d) Microwatershed
Soil Suitability Information

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry	
Machanura	RIVER_BHIMA	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs
Bilhara	RIVER_BHIMA	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs
Belagundi	1	S3n	S2n	S3n	S2tn	N1n	S3tn	N1n	S3n	S3t	S3n	S2tn	N1n	N1n	S3n	N1n	N1n	S3n	S3n	N1n	S3n	S3n	S3n	S3n	S3n	S2n	S3n	S3n	N1n	N1n	
Belagundi	178	S3n	S2n	S3n	S2tn	N1n	S3tn	N1n	S3n	S3t	S3n	S2tn	N1n	N1n	S3n	N1n	N1n	S3n	S3n	N1n	S3n	S3n	S3n	S3n	S3n	S2n	S3n	S3n	N1n	N1n	
Belagundi	179	S3n	S2n	S3n	S2tn	N1n	S3tn	N1n	S3n	S3t	S3n	S2tn	N1n	N1n	S3n	N1n	N1n	S3n	S3n	N1n	S3n	S3n	S3n	S3n	S3n	S2n	S3n	S3n	N1n	N1n	
Belagundi	180	S3n	S2n	S3n	S2tn	N1n	S3tn	N1n	S3n	S3t	S3n	S2tn	N1n	N1n	S3n	N1n	N1n	S3n	S3n	N1n	S3n	S3n	S3n	S3n	S3n	S2n	S3n	S3n	N1n	N1n	
Belagundi	182	S3n	S2n	S3n	S2tn	N1n	S3tn	N1n	S3n	S3t	S3n	S2tn	N1n	N1n	S3n	N1n	N1n	S3n	S3n	N1n	S3n	S3n	S3n	S3n	S3n	S2n	S3n	S3n	N1n	N1n	
Belagundi	183	S3n	S2n	S3n	S2tn	N1n	S3tn	N1n	S3n	S3t	S3n	S2tn	N1n	N1n	S3n	N1n	N1n	S3n	S3n	N1n	S3n	S3n	S3n	S3n	S3n	S2n	S3n	S3n	N1n	N1n	
Belagundi	184	S3n	S2n	S3n	S2tn	N1n	S3tn	N1n	S3n	S3t	S3n	S2tn	N1n	N1n	S3n	N1n	N1n	S3n	S3n	N1n	S3n	S3n	S3n	S3n	S3n	S2n	S3n	S3n	N1n	N1n	
Belagundi	185	S3n	S2n	S3n	S2tn	N1n	S3tn	N1n	S3n	S3t	S3n	S2tn	N1n	N1n	S3n	N1n	N1n	S3n	S3n	N1n	S3n	S3n	S3n	S3n	S3n	S2n	S3n	S3n	N1n	N1n	
Belagundi	186	S3n	S2n	S3n	S2tn	N1n	S3tn	N1n	S3n	S3t	S3n	S2tn	N1n	N1n	S3n	N1n	N1n	S3n	S3n	N1n	S3n	S3n	S3n	S3n	S3n	S2n	S3n	S3n	N1n	N1n	
Belagundi	187	S3n	S2n	S3n	S2tn	N1n	S3tn	N1n	S3n	S3t	S3n	S2tn	N1n	N1n	S3n	N1n	N1n	S3n	S3n	N1n	S3n	S3n	S3n	S3n	S3n	S2n	S3n	S3n	N1n	N1n	
Belagundi	188	S3n	S2n	S3n	S2tn	N1n	S3tn	N1n	S3n	S3t	S3n	S2tn	N1n	N1n	S3n	N1n	N1n	S3n	S3n	N1n	S3n	S3n	S3n	S3n	S3n	S2n	S3n	S3n	N1n	N1n	
Belagundi	189	S3n	S2n	S3n	S2tn	N1n	S3tn	N1n	S3n	S3t	S3n	S2tn	N1n	N1n	S3n	N1n	N1n	S3n	S3n	N1n	S3n	S3n	S3n	S3n	S3n	S2n	S3n	S3n	N1n	N1n	
Belagundi	190	S3n	S2n	S3n	S2tn	N1n	S3tn	N1n	S3n	S3t	S3n	S2tn	N1n	N1n	S3n	N1n	N1n	S3n	S3n	N1n	S3n	S3n	S3n	S3n	S3n	S2n	S3n	S3n	N1n	N1n	
Belagundi	191	S3n	S2n	S3n	S2tn	N1n	S3tn	N1n	S3n	S3t	S3n	S2tn	N1n	N1n	S3n	N1n	N1n	S3n	S3n	N1n	S3n	S3n	S3n	S3n	S3n	S2n	S3n	S3n	N1n	N1n	
Belagundi	192	S3n	S2n	S3n	S2tn	N1n	S3tn	N1n	S3n	S3t	S3n	S2tn	N1n	N1n	S3n	N1n	N1n	S3n	S3n	N1n	S3n	S3n	S3n	S3n	S3n	S2n	S3n	S3n	N1n	N1n	
Anura B	74	N1n	S3n	N1n	S3n	N1n	S3n	N1n	N1n	S3n	N1n	S3n	N1n	N1n	N1n	N1tn	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n
Anura B	98	S3n	S2n	S3n	S2tn	N1n	S3tn	N1n	S3n	S3t	S3n	S2tn	N1n	N1n	S3n	N1n	N1n	S3n	S3n	N1n	S3n	S3n	S3n	S3n	S3n	S2n	S3n	S3n	N1n	N1n	
Anura B	99	S3n	S2n	S3n	S2tn	N1n	S3tn	N1n	S3n	S3t	S3n	S2tn	N1n	N1n	S3n	N1n	N1n	S3n	S3n	N1n	S3n	S3n	S3n	S3n	S3n	S2n	S3n	S3n	N1n	N1n	
Anura B	100	S3n	S2n	S3n	S2tn	N1n	S3tn	N1n	S3n	S3t	S3n	S2tn	N1n	N1n	S3n	N1n	N1n	S3n	S3n	N1n	S3n	S3n	S3n	S3n	S3n	S2n	S3n	S3n	N1n	N1n	
Anura B	101	S3n	S2n	S3n	S2tn	N1n	S3tn	N1n	S3n	S3t	S3n	S2tn	N1n	N1n	S3n	N1n	N1n	S3n	S3n	N1n	S3n	S3n	S3n	S3n	S3n	S2n	S3n	S3n	N1n	N1n	
Anura B	102	S3n	S2n	S3n	S2tn	N1n	S3tn	N1n	S3n	S3t	S3n	S2tn	N1n	N1n	S3n	N1n	N1n	S3n	S3n	N1n	S3n	S3n	S3n	S3n	S3n	S2n	S3n	S3n	N1n	N1n	
Anura B	103	S3n	S2n	S3n	S2tn	N1n	S3tn	N1n	S3n	S3t	S3n	S2tn	N1n	N1n	S3n	N1n	N1n	S3n	S3n	N1n	S3n	S3n	S3n	S3n	S3n	S2n	S3n	S3n	N1n	N1n	

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Margold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry
Anura B	104	S3n	S2n	S3n	S2tn	N1n	S3tn	N1n	S3n	S3t	S3n	S2tn	N1n	N1n	S3n	N1n	N1n	S3n	S3n	N1n	S3n	S3n	S3n	S3n	S3n	S2n	S3n	S3n	N1n	N1n
Anura B	105	S3n	S2n	S3n	S2tn	N1n	S3tn	N1n	S3n	S3t	S3n	S2tn	N1n	N1n	S3n	N1n	N1n	S3n	S3n	N1n	S3n	S3n	S3n	S3n	S3n	S2n	S3n	S3n	N1n	N1n
Anura B	106	S3n	S2n	S3n	S2tn	N1n	S3tn	N1n	S3n	S3t	S3n	S2tn	N1n	N1n	S3n	N1n	N1n	S3n	S3n	N1n	S3n	S3n	S3n	S3n	S3n	S2n	S3n	S3n	N1n	N1n
Anura B	107	S3n	S2n	S3n	S2tn	N1n	S3tn	N1n	S3n	S3t	S3n	S2tn	N1n	N1n	S3n	N1n	N1n	S3n	S3n	N1n	S3n	S3n	S3n	S3n	S3n	S2n	S3n	S3n	N1n	N1n
Anura B	108	S3n	S2n	S3n	S2tn	N1n	S3tn	N1n	S3n	S3t	S3n	S2tn	N1n	N1n	S3n	N1n	N1n	S3n	S3n	N1n	S3n	S3n	S3n	S3n	S3n	S2n	S3n	S3n	N1n	N1n
Anura B	109	S3n	S2n	S3n	S2tn	N1n	S3tn	N1n	S3n	S3t	S3n	S2tn	N1n	N1n	S3n	N1n	N1n	S3n	S3n	N1n	S3n	S3n	S3n	S3n	S3n	S2n	S3n	S3n	N1n	N1n
Anura B	110	S3n	S2n	S3n	S2tn	N1n	S3tn	N1n	S3n	S3t	S3n	S2tn	N1n	N1n	S3n	N1n	N1n	S3n	S3n	N1n	S3n	S3n	S3n	S3n	S3n	S2n	S3n	S3n	N1n	N1n
Anura B	111	S3n	S2n	S3n	S2tn	N1n	S3tn	N1n	S3n	S3t	S3n	S2tn	N1n	N1n	S3n	N1n	N1n	S3n	S3n	N1n	S3n	S3n	S3n	S3n	S3n	S2n	S3n	S3n	N1n	N1n
Anura B	112	S3n	S2n	S3n	S2tn	N1n	S3tn	N1n	S3n	S3t	S3n	S2tn	N1n	N1n	S3n	N1n	N1n	S3n	S3n	N1n	S3n	S3n	S3n	S3n	S3n	S2n	S3n	S3n	N1n	N1n
Anura B	125/1	S1	S1	S1	S2t	S1	S3t	S1	S1	S3t	S2t	S2t	S1	S1	S1	S2n	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1
Anura B	125/2	S1	S1	S1	S2t	S1	S3t	S1	S1	S3t	S2t	S2t	S1	S1	S1	S2n	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1
Anura B	126	S1	S1	S1	S2t	S1	S3t	S1	S1	S3t	S2t	S2t	S1	S1	S1	S2n	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1
Anura B	127/1	S1	S1	S1	S2t	S1	S3t	S1	S1	S3t	S2t	S2t	S1	S1	S1	S2n	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1
Anura B	127/2	S1	S1	S1	S2t	S1	S3t	S1	S1	S3t	S2t	S2t	S1	S1	S1	S2n	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1
Anura B	128/2	S1	S1	S1	S2t	S1	S3t	S1	S1	S3t	S2t	S2t	S1	S1	S1	S2n	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1
Anura B	129	N1n	S3n	N1n	S3n	N1n	S3nt	N1n	N1n	S3nt	N1n	S3n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n
Anura B	130	S3n	S2n	S3n	S2tn	N1n	S3tn	N1n	S3n	S3t	S3n	S2tn	N1n	N1n	S3n	N1n	N1n	S3n	S3n	N1n	S3n	S3n	S3n	S3n	S3n	S2n	S3n	S3n	N1n	N1n
Anura B	131	N1n	S3n	N1n	S3n	N1n	S3nt	N1n	N1n	S3nt	N1n	S3n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n
Anura B	132	N1n	S3n	N1n	S3n	N1n	S3nt	N1n	N1n	S3nt	N1n	S3n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n
Anura B	133	N1n	S3n	N1n	S3n	N1n	S3n	N1n	N1n	S3n	N1n	S3n	N1n	N1n	N1n	N1tn	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n
Anura B	134	N1n	S3n	N1n	S3n	N1n	S3nt	N1n	N1n	S3nt	N1n	S3n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n
Anura B	135	N1n	S3n	N1n	S3n	N1n	S3nt	N1n	N1n	S3nt	N1n	S3n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n
Anura B	136	N1n	S3n	N1n	S3n	N1n	S3nt	N1n	N1n	S3nt	N1n	S3n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n
Anura B	137	N1n	S3n	N1n	S3n	N1n	S3nt	N1n	N1n	S3nt	N1n	S3n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n
Anura B	138	N1n	S3n	N1n	S3n	N1n	S3nt	N1n	N1n	S3nt	N1n	S3n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n
Anura B	139	N1n	S3n	N1n	S3n	N1n	S3nt	N1n	N1n	S3nt	N1n	S3n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n
Anura B	140	N1n	S3n	N1n	S3n	N1n	S3nt	N1n	N1n	S3nt	N1n	S3n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Margold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry	
Anura B	141	N1n	S3n	N1n	S3n	N1n	S3nt	N1n	N1n	S3nt	N1n	S3n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n	
Anura B	142	S3n	S2n	S3n	S2tn	N1n	S3tn	N1n	S3n	S3t	S3n	S2tn	N1n	N1n	S3n	N1n	N1n	S3n	S3n	N1n	S3n	S3n	S3n	S3n	S3n	S2n	S3n	S3n	N1n	N1n	
Anura B	143	S3n	S2n	S3n	S2tn	N1n	S3tn	N1n	S3n	S3t	S3n	S2tn	N1n	N1n	S3n	N1n	N1n	S3n	S3n	N1n	S3n	S3n	S3n	S3n	S3n	S2n	S3n	S3n	N1n	N1n	
Anura B	144	S3n	S2n	S3n	S2tn	N1n	S3tn	N1n	S3n	S3t	S3n	S2tn	N1n	N1n	S3n	N1n	N1n	S3n	S3n	N1n	S3n	S3n	S3n	S3n	S3n	S2n	S3n	S3n	N1n	N1n	
Anura B	145	S3n	S2n	S3n	S2tn	N1n	S3tn	N1n	S3n	S3t	S3n	S2tn	N1n	N1n	S3n	N1n	N1n	S3n	S3n	N1n	S3n	S3n	S3n	S3n	S3n	S2n	S3n	S3n	N1n	N1n	
Anura B	146	S3n	S2n	S3n	S2tn	N1n	S3tn	N1n	S3n	S3t	S3n	S2tn	N1n	N1n	S3n	N1n	N1n	S3n	S3n	N1n	S3n	S3n	S3n	S3n	S3n	S2n	S3n	S3n	N1n	N1n	
Anura B	147	S3n	S2n	S3n	S2tn	N1n	S3tn	N1n	S3n	S3t	S3n	S2tn	N1n	N1n	S3n	N1n	N1n	S3n	S3n	N1n	S3n	S3n	S3n	S3n	S3n	S2n	S3n	S3n	N1n	N1n	
Anura B	148	N1n	S3n	N1n	S3n	N1n	S3nt	N1n	N1n	S3nt	N1n	S3n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n
Anura B	149	N1n	S3n	N1n	S3n	N1n	S3nt	N1n	N1n	S3nt	N1n	S3n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n
Anura B	150	N1n	S3n	N1n	S3n	N1n	S3nt	N1n	N1n	S3nt	N1n	S3n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n
Anura B	151	N1n	S3n	N1n	S3n	N1n	S3nt	N1n	N1n	S3nt	N1n	S3n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n
Anura B	152	N1n	S3n	N1n	S3n	N1n	S3nt	N1n	N1n	S3nt	N1n	S3n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n
Anura B	153	N1n	S3n	N1n	S3n	N1n	S3n	N1n	N1n	S3n	N1n	S3n	N1n	N1n	N1n	N1tn	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n
Anura B	154	N1n	S3n	N1n	S3n	N1n	S3nt	N1n	N1n	S3nt	N1n	S3n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n
Anura B	155	N1n	S3n	N1n	S3n	N1n	S3n	N1n	N1n	S3n	N1n	S3n	N1n	N1n	N1n	N1tn	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n
Anura B	156	N1n	S3n	N1n	S3n	N1n	S3n	N1n	N1n	S3n	N1n	S3n	N1n	N1n	N1n	N1tn	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n
Anura B	157	N1n	S3n	N1n	S3n	N1n	S3nt	N1n	N1n	S3nt	N1n	S3n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n
Anura B	158	N1n	S3n	N1n	S3n	N1n	S3nt	N1n	N1n	S3nt	N1n	S3n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n
Anura B	159	S1	S1	S1	S2t	S1	S3t	S1	S1	S3t	S2t	S2t	S1	S1	S1	S2n	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1
Anura B	160	S1	S1	S1	S2t	S1	S3t	S1	S1	S3t	S2t	S2t	S1	S1	S1	S2n	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1
Anura B	161	S1	S1	S1	S2t	S1	S3t	S1	S1	S3t	S2t	S2t	S1	S1	S1	S2n	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1
Anura B	162	S1	S1	S1	S2t	S1	S3t	S1	S1	S3t	S2t	S2t	S1	S1	S1	S2n	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1
Anura B	163	S1	S1	S1	S2t	S1	S3t	S1	S1	S3t	S2t	S2t	S1	S1	S1	S2n	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1
Anura B	164	N1n	S3n	N1n	S3n	N1n	S3n	N1n	N1n	S3n	N1n	S3n	N1n	N1n	N1n	N1tn	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n
Anura B	165	N1n	S3n	N1n	S3n	N1n	S3n	N1n	N1n	S3n	N1n	S3n	N1n	N1n	N1n	N1tn	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n
Anura B	166	N1n	S3n	N1n	S3n	N1n	S3nt	N1n	N1n	S3nt	N1n	S3n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n
Anura B	167	N1n	S3n	N1n	S3n	N1n	S3nt	N1n	N1n	S3nt	N1n	S3n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry	
Anura B	168	N1n	S3n	N1n	S3n	N1n	S3nt	N1n	N1n	S3nt	N1n	S3n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n	
Anura B	169	N1n	S3n	N1n	S3n	N1n	S3n	N1n	N1n	S3n	N1n	S3n	N1n	N1n	N1n	N1tn	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n
Anura B	170	N1n	S3n	N1n	S3n	N1n	S3n	N1n	N1n	S3n	N1n	S3n	N1n	N1n	N1n	N1tn	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n
Anura B	171	N1n	S3n	N1n	S3n	N1n	S3n	N1n	N1n	S3n	N1n	S3n	N1n	N1n	N1n	N1tn	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n
Anura B	172	N1n	S3n	N1n	S3n	N1n	S3n	N1n	N1n	S3n	N1n	S3n	N1n	N1n	N1n	N1tn	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n
Anura B	173	N1n	S3n	N1n	S3n	N1n	S3n	N1n	N1n	S3n	N1n	S3n	N1n	N1n	N1n	N1tn	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n
Anura B	174	N1n	S3n	N1n	S3n	N1n	S3n	N1n	N1n	S3n	N1n	S3n	N1n	N1n	N1n	N1tn	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n
Anura B	175	N1n	S3n	N1n	S3n	N1n	S3n	N1n	N1n	S3n	N1n	S3n	N1n	N1n	N1n	N1tn	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n
Anura B	176	N1n	S3n	N1n	S3n	N1n	S3n	N1n	N1n	S3n	N1n	S3n	N1n	N1n	N1n	N1tn	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n
Anura B	177	S1	S1	S1	S2t	S1	S3t	S1	S1	S3t	S2t	S2t	S1	S1	S1	S2n	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1
Anura B	178	S1	S1	S1	S2t	S1	S3t	S1	S1	S3t	S2t	S2t	S1	S1	S1	S2n	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1
Anura B	179/1	S1	S1	S1	S2t	S1	S3t	S1	S1	S3t	S2t	S2t	S1	S1	S1	S2n	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1
Anura B	179/2	S1	S1	S1	S2t	S1	S3t	S1	S1	S3t	S2t	S2t	S1	S1	S1	S2n	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1
Anura B	179/3	S1	S1	S1	S2t	S1	S3t	S1	S1	S3t	S2t	S2t	S1	S1	S1	S2n	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1
Anura B	180/1	S1	S1	S1	S2t	S1	S3t	S1	S1	S3t	S2t	S2t	S1	S1	S1	S2n	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1
Anura B	181	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Anura B	182	S1	S1	S1	S2t	S1	S3t	S1	S1	S3t	S2t	S2t	S1	S1	S1	S2n	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1	S1
Anura B	183	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs
Anura B	186	N1n	S3n	N1n	S3n	N1n	S3n	N1n	N1n	S3n	N1n	S3n	N1n	N1n	N1n	N1tn	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n
Anura B	187	N1n	S3n	N1n	S3n	N1n	S3n	N1n	N1n	S3n	N1n	S3n	N1n	N1n	N1n	N1tn	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n
Anura B	188	N1n	S3n	N1n	S3n	N1n	S3n	N1n	N1n	S3n	N1n	S3n	N1n	N1n	N1n	N1tn	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	N1n	S3n	N1n	N1n	N1n	N1n
NA	NA	S3n	S2n	S3n	S2tn	N1n	S3tn	N1n	S3n	S3t	S3n	S2tn	N1n	N1n	S3n	N1n	N1n	S3n	S3n	N1n	S3n	S3n	S3n	S3n	S3n	S2n	S3n	S3n	N1n	N1n	

PART-B

SOCIO-ECONOMIC STATUS OF FARM HOUSEHOLDS

CONTENTS

1.	Findings of the socio-economic survey	1-3
2.	Introduction	5-6
3	Chapter II- Methodology	7-8
4	Salient features of the survey	9-27
5	Summary	29-33

LIST OF TABLES

1	Households sampled for socio economic survey	9
2	Population characteristics	9
3	Age wise classification of household members	9
4	Education level of household members	10
5	Occupation of household heads	10
6	Occupation of family members	10
7	Institutional participation of household members	11
8	Type of house owned by households	11
9	Durable assets owned by households	11
10	Average value of durable assets owned by households	11
11	Farm implements owned by households	12
12	Average value of farm implements	12
13	Livestock possession by households	13
14	Average labour availability	13
15	Adequacy of hired labour	13
16	Distribution of land (ha)	13
17	Average land value (Rs./ha)	14
18	Cropping pattern	14
19	Cropping intensity	14
20	Possession of bank account and saving	14
21	Borrowing status	14
22	Source of credit	15
23	Avg. Credit amount	15
24	Purpose of credit borrowed - Institutional Credit	15
25	Purpose of credit borrowed – Private Credit	15
26	Repayment status of credit borrowed from institutional Source	16
27	Repayment status of credit borrowed from private Source	16
28.a	Cost of cultivation of Red gram	17
28.b	Cost of cultivation of Groundnut	18
28.c	Cost of cultivation of Jowar	19

28.d	Cost of cultivation of Paddy	20
28.e	Cost of cultivation of Maize	21
29	Adequacy of fodder	22
30	Annual gross income	22
31	Average annual expenditure	22
32	Horticultural species grown	22
33	Forest species grown	23
34	Average additional investment capacity	23
35	Source of funds for additional investment	23
36	Marketing of the agricultural produce	24
37	Marketing channels used for sale of agricultural produce	24
38	Mode of transport of agricultural produce	24
39	Incidence of soil and water erosion problems	24
40	Interest shown towards soil testing	25
41	Usage pattern of fuel for domestic use	25
42	Source of drinking water	25
43	Source of light	25
44	Existence of sanitary toilet facility	25
45	Possession of public distribution system (PDS) card	26
46	Participation in NREGA programme	26
47	Adequacy of food items	26
48	Inadequacy of food items	26
49	Farming constraints experienced	27

FINDINGS OF THE SOCIO-ECONOMIC SURVEY

- ❖ *The survey was conducted in Belagunda-3 is located at North latitude 16^o 34' 11.759" and 16^o 32'43.028" and East longitude 77^o 12' 8.355" and 77^o 10' 11.177" covering an area of about 474.74 ha coming under under Anura. B and Belagundi villages of Yadagiri taluk.*
- ❖ *Socio-economic analysis of Belagunda-3 micro watersheds of Belagunda sub-watershed, Yadgiri taluk & District indicated that, out of the total sample of 35 farmers were sampled in Belagunda-3 micro-watershed among households surveyed 8 (22.86%) were marginal, 19(54.29%) were small, 3 (8.57 %) were semi medium and 2 (5.71 %) were medium farmers. 3 landless farmers were also interviewed for the survey.*
- ❖ *The population characteristics of households indicated that, there were 91 (54.49%) men and 76 (45.51 %) were women. The average population of landless was 4.3, marginal farmers were 5.4, small farmers were 4.5, semi medium farmers were 6 and medium farmers were 3.5.*
- ❖ *Majority of the respondents (44.31%) were in the age group of 16-35 years.*
- ❖ *Education level of the sample households indicated that, there were 53.29 per cent illiterates, 42.52 per cent pre university education and 4.79 per cent attained graduation.*
- ❖ *About, 88.57 per cent of household heads practicing agriculture and 5.71 per cent of the household heads were engaged as agricultural labourers.*
- ❖ *Agriculture was the major occupation for 64.67 per cent of the household members.*
- ❖ *In the study area, 82.86 per cent of the households possess katcha house and 5.71 per cent possess pucca house.*
- ❖ *The durable assets owned by the households showed that, 77.14 per cent possess TV, 40.00 per cent possess mixer grinder, 94.29 per cent possess mobile phones and 31.43 per cent possess motor cycles.*
- ❖ *Farm implements owned by the households indicated that, 25.71 per cent of the households possess plough, 2.86 per cent possess tractor, 2.86 per cent possess bullock cart and 5.71 per cent possess sprayer.*
- ❖ *Regarding livestock possession by the households, 14.29 per cent possess local cow and 2.86 per cent possess buffalo.*
- ❖ *The average labour availability in the study area showed that, own labour men available in the micro watershed was 1.85, women available in the micro watershed was 1.18, hired labour (men) available was 13.09 and hired labour (women) available was 9.71.*

- ❖ *Out of the total land holding of the sample respondents 71.75 per cent (43.07 ha) of the area is under dry condition and the remaining 22.97 per cent area is irrigated land.*
- ❖ *The major crops grown by sample farmers are Red gram, Groundnut, Jowar, Paddy and Maize and cropping intensity was recorded as 100.00 per cent.*
- ❖ *Out of the sample households 85.71 percent possessed bank account and 54.29 per cent of them have savings in the account.*
- ❖ *About 100.00 per cent of the respondents borrowed credit from various sources.*
- ❖ *Among the credit borrowed by households, 55.00 per cent have borrowed loan from commercial banks and 15.00 per cent from co-operative/Grameena bank.*
- ❖ *Majority of the respondents (100.00%) have borrowed loan for agriculture purpose.*
- ❖ *The per hectare cost of cultivation for Red gram, Groundnut, Jowar, Paddy and Maize was Rs.29784.68, 37139.36, 37139.36, 36528.36 and 22730.15 with benefit cost ratio of 1:1.70, 1: 1.00, 1: 1.00, 1: 0.70 and 1:1.40 respectively.*
- ❖ *Further, 25.71 per cent of the households opined that dry fodder was adequate.*
- ❖ *The average annual gross income of the farmers was Rs. 97142.86 in micro-watershed, of which Rs. 50114.29 comes from agriculture.*
- ❖ *Sampled households have grown 1 horticulture trees and 51 forestry trees together in the fields and back yards.*
- ❖ *Households have an average investment capacity of Rs. 2628.57 for land development and Rs. 16428.57 for irrigation facility.*
- ❖ *Source of funds for additional investment is concerned, 2.86 per cent depends on own funds for land development activities.*
- ❖ *Regarding marketing channels, 91.43 per cent of the households have sold agricultural produce to the local/village merchants.*
- ❖ *Further, 91.43 per cent of the households have used tractor for the transport of agriculture commodity.*
- ❖ *Majority of the farmers (28.57%) have experienced soil and water erosion problems in the watershed and 94.29 per cent of the households were interested towards soil testing.*
- ❖ *Fire was the major source of fuel for domestic use for 65.71 per cent of the households and 37.14 per cent households has LPG connection.*
- ❖ *Piped supply was the major source for drinking water for 100.00 per cent of the households.*
- ❖ *Electricity was the major source of light for 97.14 per cent of the households.*
- ❖ *In the study area, 51.43 per cent of the households possess toilet facility.*
- ❖ *Regarding possession of PDS card, 100.00 per cent of the households possessed BPL card.*

- ❖ *Households opined that, the requirement of cereals (100.00%), pulses (97.14%) and oilseeds (40.00%) are adequate for consumption.*
- ❖ *Farming constraints experienced by households in the micro watersheds were lower fertility status of the soil (97.14%) wild animal menace on farm field (91.43%), frequent incidence of pest and diseases (91.43%), inadequacy of irrigation water (80.00%), high cost of fertilizers and plant protection chemicals (91.43%), high rate of interest on credit (91.43%), low price for the agricultural commodities (60.00%), lack of marketing facilities in the area (60.00%), inadequate extension services (54.29%), lack of transport for safe transport of the agricultural produce to the market (60.00%) and Less rainfall (2.86%).*

INTRODUCTION

Soil and water are the two precious natural resources which are essential for crop production and existence of life on earth. Rainfed agriculture is under severe stress due to various constraints related to agriculture like uneven and erratic distribution of rainfall, indiscriminate use of fertilizers, chemicals and pesticides, adoption of improper land management practices, soil erosion, decline in soil fertility, decline in ground water resources leading to low crop productivity. The area under rainfed agriculture has to be managed effectively using the best available practices to enhance the production of food, fodder and fuel. This is possible if the land resources are characterized at each parcel of land through detailed land resource inventory using the best available techniques of remote sensing, GPS and GIS. The watershed development programs are aimed at the sustainable distribution of its resources and the process of creating and implementing plans, programs, and projects to sustain and enhance watershed functions that affect the plant, animal and human communities within a watershed boundary.

World Bank funded KWDP II, SUJALA III project was implemented in with Broad objective of demonstrating more effective watershed management through greater integration of programmes related to rain-fed agriculture, innovative and science based approaches and strengthen institutional capacities and If successful, it is expected that the systems and tools could be mainstreamed into the overall IWMP in the State of Karnataka and in time, throughout other IWMP operations in India. With this background the socio-economic survey has been carried out with following specific objectives:

1. To understand the demographic features of the households in the micro-watershed
2. To understand the extent of family labour available and additional employment opportunities available within the village.
3. To know the status of assets of households in the micro-watershed for suggesting possible improvements.
4. To study the cropping pattern, cropped area and productivity levels of different households in micro-watershed.
5. To determine the type and extent of livestock owned by different categories of HHs
6. Availability of fodder and level of livestock management.

Scope and importance of survey

Survey helps in identification of different socio-economic and resource use-patterns of farmers at the Micro watershed. Household survey provides demographic features, labour force, and levels of education; land ownership and asset position (including livestock and other household assets) of surveyed households; and cropping patterns, input intensities, and average crop yields from farmers' fields. It

also discusses crop utilization and the degree of commercialization of production in the areas; farmers' access to and utilization of credit from formal and informal sources; and the level of adoption and use of soil, water, and pest management technologies.

METHODOLOGY

The description of the methods, components selected for the survey and procedures followed in conducting the baseline survey are furnished under the following heads.

1. Description of the study area

Yadgir District is one of the 30 districts of Karnataka state in southern India. This district was carved out from the erstwhile Gulbarga district as the 30th district of Karnataka on 10 April 2010. Yadgir town is the administrative headquarters of the district. The district comprises of 3 taluks namely, Shahapur, Yadgiri and Shorapur (There are 16 hoblies, 117 Gram Panchayats, 4 Municipalities, 8 Towns/ Urban agglomeration and 487 inhabited & 32 un-inhabited villages The district occupies an area of 5,160.88 km².

Yadgir district is the second smallest district in the state, area wise is very rich in cultural traditions. The vast stretch of fertile black soil of the district is known for bumper red gram and jowar crops. The district is a "Daal bowl" of the state. The district is also known for cluster of cement industries and a distinct stone popularly known as "Malakheda Stone". Two main rivers, Krishna and Bhima, and a few tributaries flow in this region. Krishna and Bhima Rivers drain the district. They constitute the two major river basins of the district. Kagna and Amarja are the two sub-basins of Bhima River, which occur within the geographical area of the district

According to the 2011 census Yadgir district has a population of 1, 172,985, roughly equal to the nation of Timor-Leste or the US state of Rhode Island. This gives it a ranking of 404th in India (out of a total of 640). The district has a population density of 224 inhabitants per square kilometre (580/sq mi). Its population growth rate over the decade 2001-2011 was 22.67%. Yadgir has a sex ratio of 984 females for every 1000 males, and a literacy rate of 52.36%.

2. Locale of the survey and description of the micro-watershed and

The study was conducted in Belagunda-3 micro-watershed (Belagunda sub-watershed, Yadgiri taluk & District) is located at North latitude 16⁰ 34' 11.759" and 16⁰ 32' 43.028" and East longitude 77⁰ 12' 8.355" and 77⁰ 10' 11.177" covering an area of about 474.74 ha bounded by under Anura. B and Belagundi Villages.

3. Selection of the respondents for the study

The micro-watershed is marked with 320 square meters grids. One farmer from every alternate grid in the micro-watershed was selected for the study and interviewed for socio-economic data. Totally 35 households were interviewed for the survey.

4. The parameters considered for socio-economic survey of households

Two forms of data were collected from the micro-watershed which includes primary data from the farm households and secondary data about the villages under the micro-watershed jurisdiction.

The following parameters were considered for the primary data collection about the socio-economic data of the households, (1) Demographic information, (2) Farm and durable assets owned by households, (3) Livestock possession, (4) Labour availability, (5) Level of migration in the village, Land holding, (7) Cropping pattern, (8) Source of irrigation, (9) Borrowing status, (10) Cost of cultivation of major crops, (11) Economics of subsidiary activities, (12) Fodder availability, (13) Family annual income from different sources, (14) Horticulture and forestry species grown, (15) Additional investment capacity, (16) Marketing practices, (17) Status of soil and water conservation structure, (18) Access to basic needs and (19) Constraints and suggestion.

The following parameters were considered for the secondary data regarding the villages under the micro-watershed jurisdiction, (1) Number of villages in each micro-watershed jurisdiction, (2) Village wise number of households, (3) Geographical area of the villages, (4) Cultivable area including rainfed and irrigated, (5) Number and type of house in each village, (6) Human and livestock population, (7) Facilities in the village such as roads, transport facility for conveyance, drinking water supply, street light and (8) Community based organizations in the villages.

5. Development of interview schedule and data collection

Taking into the consideration the objectives of the survey, an interview schedule was prepared after thorough consultation with the experts in the field of social sciences. A comprehensive interview schedule covering all the major parameters for measuring the socio-economic situation was developed.

6. Tools used to analyze the data

The statistical components such as frequency and percentage were used to analyze the data.

Abbreviations used in the report

LL=Landless

MF=Marginal Farmers

SF=Small farmers

SMF=Semi medium farmers

MDF=Medium farmers

LF=Large Farmers

FINDINGS OF THE SURVEY

This chapter deals with systematic presentation of results of the survey. Keeping in view the objectives, the salient features of the survey are presented under the following headings.

Households sampled for socio-economic survey: The data on households sampled for socio economic survey in Belagunda-3 Micro watershed is presented in Table 1 and it indicated that 35 farmers were sampled in Belagunda-3 micro-watershed among households surveyed 8 (22.86%) were marginal, 19(54.29%) were small, 3 (8.57 %) were semi medium and 2 (5.71 %) were medium farmers. 3 landless farmers were also interviewed for the survey.

Table 1. Households sampled for socio economic survey in Belagunda-3 micro-watershed

Sl.No.	Particulars	LL (3)		MF (8)		SF (19)		SMF (3)		MDF (2)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Farmers	3	8.57	8	22.9	19	54.3	3	8.57	2	5.71	35	100

Population characteristics: The population characteristics of households sampled for socio-economic survey in Belagunda-3 Micro watershed is presented in Table 2. The data indicated that, there were 91 (54.49%) men and 76 (45.51%) were women. The average population of landless was 4.3, marginal farmers were 5.4, small farmers were 4.5, semi medium farmers were 6 and medium farmers were 3.5.

Table 2. Population characteristics in Belagunda-3 micro-watershed

Sl. No.	Particulars	LL (13)		MF (43)		SF (86)		SMF (18)		MDF (7)		All (167)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Men	3	23.1	21	49	51	59	12	66.7	4	57.1	91	54.5
2	Women	10	76.9	22	51	35	41	6	33.3	3	42.9	76	45.5
Total		13	100	43	100	86	100	18	100	7	100	167	100
Average		4.3		5.4		4.5		6.0		3.5		4.8	

Age wise classification of population: The age wise classification of household members in Belagunda-3 Micro watershed is presented in Table 3. The indicated that, 35 (20.96%) of population were 0-15 years of age, 74 (44.31%) were 16-35 years of age, 44(26.35%) were 36-60 years of age and 14 (8.38 %) were above 61 years of age.

Table 3: Age wise classification of members of the household in Belagunda-3 micro-watershed

Sl. No.	Particulars	LL (13)		MF (43)		SF (86)		SMF (18)		MDF(7)		All (167)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	0-15 years of age	4	30.8	10	23.3	17	19.8	4	22.22	0	0	35	20.96
2	16-35 years of age	7	53.9	17	39.5	39	45.4	9	50	2	29	74	44.31
3	36-60 years of age	2	15.4	13	30.2	22	25.6	4	22.22	3	43	44	26.35
4	> 61 years	0	0	3	6.98	8	9.3	1	5.56	2	29	14	8.38
Total		13	100	43	100	86	100	18	100	7	100	167	100

Education level of household members: Education level of household members in Belagunda-3 Micro watershed is presented in Table 4. The results indicated that, there were 53.29 per cent of illiterates, 30.54 per cent of them had primary school education, 8.38 per cent high school education, 1.80 per cent of them had PUC education, 4.79 per cent attained graduation and 1.20 them had other education.

Table 4. Education level of members of the household in Belagunda-3 micro-watershed

Sl. No.	Particulars	LL (13)		MF (43)		SF (86)		SMF (18)		MDF (7)		All (167)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Primary School	6	46.2	13	30.2	26	30.2	5	27.8	1	14.29	51	30.5
2	High School	1	7.69	2	4.65	10	11.6	1	5.56	0	0	14	8.38
3	PUC	0	0	2	4.65	0	0	1	5.56	0	0	3	1.8
4	Degree	0	0	2	4.65	2	2.33	3	16.7	1	14.29	8	4.79
5	Others	1	7.69	1	2.33	0	0	0	0	0	0	2	1.2
Total		13	100	43	100	86	100	18	100	7	100	167	100

Occupation of head of households: The data regarding the occupation of the household heads in Belagunda-3 Micro watershed is presented in Table 5. The results indicate that, 88.57 per cent of households heads were practicing agriculture, 5.71 per cent of the household heads were agricultural Labour and housewife (2.86%).

Table 5: Occupation of heads of households in Belagunda-3 micro-watershed

Sl. No.	Particulars	LL (3)		MF (8)		SF (19)		SMF (3)		MDF (2)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Agriculture	1	33	8	100	17	89.47	3	100	2	100	31	88.57
2	Agricultural Labour	2	67	0	0	0	0	0	0	0	0	2	5.71
3	Housewife	0	0	0	0	1	5.26	0	0	0	0	1	2.86
Total		3	100	8	100	18	100	3	100	2	100	34	100

Occupation of the members of the household: The data regarding the occupation of the household members in Belagunda-3 Micro watershed is presented in Table 6. The results indicate that, agriculture was the major occupation for 64.67 per cent of the household members, 5.39 per cent were agricultural labour, 25.15 per cent were working in pursuing education, 2.99 per cent were involved as housewife and 1.20 per cent were childrens.

Table 6: Occupation of members of the household in Belagunda-3 micro-watershed

Sl. No.	Particulars	LL (13)		MF (43)		SF (86)		SMF (18)		MDF (7)		All (167)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Agriculture	4	30.8	26	60.5	59	68.6	12	66.67	7	100	108	64.7
2	Agricultural Labour	5	38.5	0	0	3	3.49	1	5.56	0	0	9	5.39
3	Private Service	0	0	1	2.33	0	0	0	0	0	0	1	0.6
4	Student	3	23.1	12	27.9	22	25.58	5	27.78	0	0	42	25.2
5	Housewife	0	0	3	6.98	2	2.33	0	0	0	0	5	2.99
6	Children	1	7.69	1	2.33	0	0	0	0	0	0	2	1.2
Total		13	100	43	100	86	100	18	100	7	100	167	100

Institutional Participation of household members: The data regarding the institutional participation of the household members in Belagunda-3 Micro watershed is presented in Table 7. The results show that, out of the total family members in the households 100 per cent of them were not participating in any of the institutions.

Table 7: Institutional Participation of household member in Belagunda-3 micro-watershed

Sl.No.	Particulars	LL (13)		MF (43)		SF (86)		SMF (18)		MDF (7)		All (167)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	No Participation	13	100	43	100	86	100	18	100	7	100	167	100
	Total	13	100	43	100	86	100	18	100	7	100	167	100

Type of house owned: The data regarding the type of house owned by the households in Belagunda-3 Micro watershed is presented in Table 8. The results indicate that, 14.29 percent possess thatched house, 82.86 per cent of the households possess katcha house and 5.71 per cent possess pacca house.

Table 8. Type of house owned by households in Belagunda-3 micro-watershed

Sl.No.	Particulars	LL (3)		MF (8)		SF (19)		SMF (3)		MDF (2)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Thatched	1	33	1	13	3	15.79	0	0	0	0	5	14.29
2	Katcha	2	67	7	88	16	84.21	2	66.7	2	100	29	82.86
3	Pucca/RCC	0	0	1	13	0	0	1	33.3	0	0	2	5.71
	Total	3	100	9	100	19	100	3	100	2	100	36	100

Durable assets owned by the households: The data regarding the Durable Assets owned by the households in Belagunda-3 Micro watershed is presented in Table 9. The results shows that, 77.14 per cent possess TV, 40.00 per cent possess mixer grinder, 22.86 per cent possess Bicycle, 31.43 per cent possess motor cycle and 94.29 per cent possess mobile phones.

Table 9. Durable assets owned by households in Belagunda-3 micro-watershed

Sl.No.	Particulars	LL (3)		MF (8)		SF (19)		SMF (3)		MDF (2)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Television	2	67	5	63	14	73.7	3	100	3	150	27	77.14
2	Mixer/Grinder	1	33	2	25	10	52.6	0	0	1	50	14	40
3	Bicycle	0	0	1	13	7	36.8	0	0	0	0	8	22.86
4	Motor Cycle	0	0	2	25	8	42.1	1	33	0	0	11	31.43
5	Mobile Phone	3	100	6	75	19	100	3	100	2	100	33	94.29
6	Blank	0	0	1	13	0	0	0	0	0	0	1	2.86

Table 10. Average value of durable assets owned in Belagunda-3 micro-watershed

Sl.No.	Particulars	LL (3)		MF (8)		SF (19)		SMF (3)		MDF (2)		All (35)	
		Average Value (Rs.)		Average Value (Rs.)		Average Value (Rs.)		Average Value (Rs.)		Average Value (Rs.)		Average Value (Rs.)	
1	Television	4500		2600		2285		2833		4666		2833	
2	Mixer/Grinder	2000		1000		1000		0		2000		1142	
3	Bicycle	0		1000		1000		0		0		1000	
4	Motor Cycle	0		35000		33500		35000		0		33909	
5	Mobile Phone	3000		900		1225		2000		2000		1356	

Average value of durable assets: The data regarding the average value of durable assets owned by the households in Belagunda-3 Micro watershed is presented in Table 10. The result shows that, the average value of television was Rs.2833.00, mixer grinder was Rs.1142.00, bicycle was Rs.1000.00, motor cycle was Rs. 33909.00 and mobile phone was Rs.1356.00.

Farm implements owned: The data regarding the farm implements owned by the households in Belagunda-3 Micro watershed is presented in Table 11. About 2.86 per cent of the households possess Bullock Cart, 25.71 per cent possess plough and 8.57 per cent possess Seed/Fertilizer Drill and Sprinkler, 5.71 per cent possess Sprayer, 68.57 per cent possess Weeder and 2.86 per cent possess tractor.

Table 11. Farm implements owned in Belagunda-3 micro-watershed

Sl.No.	Particulars	LL (3)		MF (8)		SF (19)		SMF (3)		MDF (2)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Bullock Cart	0	0	1	12.5	0	0	0	0	0	0	1	2.86
2	Plough	0	0	3	37.5	4	21.05	2	66.7	0	0	9	25.71
3	Seed/Fertilizer Drill	0	0	1	12.5	1	5.26	1	33.3	0	0	3	8.57
4	Tractor	0	0	0	0	1	5.26	0	0	0	0	1	2.86
5	Sprayer	0	0	0	0	1	5.26	1	33.3	0	0	2	5.71
6	Chaff Cutter	0	0	0	0	1	5.26	0	0	0	0	1	2.86
7	Blank	1	33	1	12.5	6	31.58	1	33.3	2	100	11	31.43

Average value of farm implements: The data regarding the average value of farm Implements owned by the households in Belagunda-3 Micro watershed is presented in Table 12. The results show that the average value of plough was Rs.1433.00, bullock Cart was Rs.25000.00, seed/fertilizer drill was Rs.3400.00, sprayer and weeder was Rs.79.00 and tractor Rs. 400000.

Table 12. Average value of farm implements in Belagunda-3 micro-watershed

Sl.No.	Particulars	Average Value (Rs.)					
		LL (3)	MF (8)	SF (19)	SMF (3)	MDF(2)	All (35)
1	Bullock Cart	0	25000	0	0	0	25000
2	Plough	0	2500	900	2500	0	1433
3	Seed/Fertilizer Drill	0	3500	30000	3000	0	12166
4	Tractor	0	0	400000	0	0	400000
5	Sprayer	0	0	4000	2800	0	3400
6	Weeder	100	62	82	100	0	79
7	Harvester	0	0	50000	0	0	50000
8	Thresher	0	0	48000	0	0	48000
9	Chaff Cutter	0	0	2000	0	0	2000

Livestock possession by the households: The data regarding the Livestock possession by the households in Belagunda-3 Micro watershed is presented in Table 13. The indicate that, 25.71 per cent of the households possess bullocks, 14.29 per cent possess local cow, 2.86 per cent possess buffalo, 2.86 per cent possess sheep, 2.86 per cent possess goat and 2.86 per cent were poultry birds.

Table 13. Livestock possession by households in Belagunda-3 micro-watershed

Sl.No.	Particulars	LL (3)		MF (8)		SF (19)		SMF (3)		MDF (2)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Bullock	0	0	3	38	4	21.05	2	67	0	0	9	25.71
2	Local cow	0	0	1	13	3	15.79	1	33	0	0	5	14.29
3	Buffalo	0	0	1	13	0	0	0	0	0	0	1	2.86
4	Sheep	0	0	0	0	1	5.26	0	0	0	0	1	2.86
5	Goat	0	0	1	13	0	0	0	0	0	0	1	2.86
6	Poultry birds	0	0	0	0	1	5.26	0	0	0	0	1	2.86
7	blank	3	100	4	50	12	63.16	0	0	2	100	21	60

Average Labour availability: The data regarding the average labour availability in Belagunda-3 Micro watershed is presented in Table 14. The indicated that, own labour men available in the micro watershed was 1.85, women available in the micro watershed was 1.18, hired labour (men) available was 13.09 and hired labour (women) available was 9.71.

Table 14. Average labour availability in Belagunda-3 micro-watershed

Sl.No.	Particulars	LL (3)	MF (8)	SF (19)	SMF (3)	MDF (2)	All (35)
		N	N	N	N	N	N
1	Hired labour Female	5	7.5	12.11	6.67	5	9.71
2	Own Labour Female	1	1	1.32	1	1	1.18
3	Own labour Male	1	2.5	1.63	2.67	1	1.85
4	Hired labour Male	7.5	10.6	15.53	11.67	7.5	13.09

Adequacy of hired labour: The data regarding the adequacy of hired labour in Belagunda-3 Micro watershed is presented in Table 15. The results indicate that, 97.14 per cent of the household opined that hired labour was adequate.

Table 15. Adequacy of hired labour in Belagunda-3 micro-watershed

Sl.No.	Particulars	LL (3)		MF (8)		SF (19)		SMF (3)		MDF (2)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Adequate	2	67	8	100	19	100	3	100	2	100	34	97.1

Distribution of land (ha): The data regarding the distribution of land (ha) in Belagunda-3 Micro watershed is presented in Table 16. The results indicate that, 31.86 ha (71.75%) of dry land and 10.20 ha (22.97 %) of irrigated land.

Table 16. Distribution of land (ha) in Belagunda-3 micro-watershed

Sl. No.	Particulars	LL (0)		MF (2)		SF (4)		SMF (0)		MDF (0)		All (6)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Dry	0	0	6.54	100	25.32	96.16	0	0	0	0	31.86	71.75
2	Irrigated	0	0	0	0	0	0	4.46	100	5.73	100	10.2	22.97
3	Permanent Fallow	0	0	0	0	1.01	3.84	0	0	0	0	1.01	2.28
	Total	0	100	6.54	100	26.33	100	4.46	100	5.73	100	43.07	100

Average value of land (ha): The data regarding the average land value (Rs./ha) in Belagunda-3 Micro watershed is presented in Table 17. The results show that the average value of dry land was Rs.432948.05, and the average value of irrigated land was Rs.392063.49.

Table 17. Average value of land (ha) in Belagunda-3 micro-watershed

Sl. No.	Particulars	LL (3)		MF (8)		SF (19)		SMF (3)		MDF (2)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Dry	0		634313.1		380941.4		0		0		432948.1	
2	Irrigated	0		0		0		537443.3		278899.1		392063.5	
3	Permanent Fallow	0		0		691600		0		0		691600	

Cropping pattern: The data regarding the cropping pattern in Belagunda-3 Micro watershed is presented in Table 18. The results indicate that, farmers have grown Cotton (13.59 ha), Red gram (8.97 ha), Paddy (8.77 ha), Maize (5.77 ha), Groundnut (3.99 ha) and Jowar (2.35 ha).

Table 18. Cropping pattern in Belagunda-3 micro-watershed

Sl.No.	Particulars	LL (3)	MF (8)	SF (19)	SMF (3)	MDF (2)	All (35)
1	Kharif - Cotton	0	3.91	9.68	0	0	13.59
2	Kharif - Red gram (togari)	0	1.72	7.24	0	0	8.97
3	Kharif - Paddy	1.34	0	0	1.69	5.74	8.77
4	Kharif - Maize	0	0.91	4.86	0	0	5.77
5	Kharif - Groundnut	0	0	1.21	2.77	0	3.99
6	Kharif - Jowar	0	0	2.35	0	0	2.35
Total	0	1.34	6.54	25.34	4.47	5.74	43.43

Cropping intensity: The data regarding the cropping intensity in Belagunda-3 Micro watershed is presented in Table 19. The results indicate that, the cropping intensity was 100.00 per cent.

Table 19. Cropping intensity (%) in Belagunda-3 micro-watershed

Sl.No.	Particulars	LL (3)	MF (8)	SF (19)	SMF (3)	MDF (2)	All (35)
1	Cropping Intensity	100	100	100	100	100	100

Possession of bank account and savings: The data regarding the possession of bank account and saving in Belagunda-3 micro-watershed is presented in Table 20. The results indicate that, 85.71 cent of the households posses bank account and 54.29 per cent of them have savings.

Table 20. Possession of Bank account and savings in Belagunda-3 micro-watershed

Sl.No.	Particulars	LL (3)		MF (8)		SF (19)		SMF (3)		MDF (2)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Account	2	66.67	7	87.5	17	89.47	2	66.67	2	100	30	85.71
2	Savings	0	0	5	62.5	12	63.16	1	33.33	1	50	19	54.29

Borrowing status: The data regarding the borrowing status in Belagunda-3 micro-watershed is presented in Table 21. The results indicate that, 100.00 percent of the sample farmers have borrowed credit from different sources.

Table 21. Borrowing status in Belagunda-3 micro-watershed

Sl.No.	Particulars	LL (3)		MF (8)		SF (19)		SMF (3)		MDF (2)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Credit Availed	3	100	8	100	19	100	3	100	2	100	35	100

Source of credit: The data regarding the source of credit availed by households in Belagunda-3 micro-watershed is presented in Table 22. The results shows that, 55.00 per cent have borrowed loan from commercial banks and 10.00 per cent have borrowed loan from Cooperative bank and 20.00 per cent have borrowed loan from Friends/Relatives, 15.00 per cent have borrowed loan from Grameena Bank, 45.00 per cent have borrowed loan from money lender and 15.00 per cent have borrowed loan from SHGs/CBOs.

Table 22. Source of credit borrowed by households in Belagunda-3 micro-watershed

Sl.No.	Particulars	LL (1)		MF (5)		SF (10)		SMF (2)		MDF (2)		All (20)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Commercial Bank	0	0	0	0	8	80	2	100	1	50	11	55
2	Cooperative Bank	0	0	1	12.5	1	10	0	0	0	0	2	10
3	Friends/Relatives	0	0	1	12.5	3	30	0	0	0	0	4	20
4	Grameena Bank	1	100	0	0	1	10	0	0	1	50	3	15
5	Money Lender	1	100	4	50	4	40	0	0	0	0	9	45
6	SHGs/CBOs	0	0	0	0	3	30	0	0	0	0	3	15

Avg. Credit amount: The data regarding the avg. Credit amount in Belagunda-3 micro-watershed is presented in Table 23. The results show that, farmers have borrowed Avg. Credit of Rs.35333.33 from different sources.

Table 23. Avg. Credit amount in Belagunda-3 micro-watershed

Sl.No.	Particulars	LL (1)	MF (5)	SF (10)	SMF (2)	MDF (2)	All (20)
		N	N	N	N	N	N
1	Average Credit	10000	12000	15000	25000	150000	35333.3

Purpose of credit borrowed (institutional Source): The data regarding the purpose of credit borrowed - Institutional Credit in Belagunda-3 micro-watershed is presented in Table 24. The results indicate that, 100.00 per cent of the households have borrowed loan for agriculture.

Table 24. Purpose of credit borrowed (institutional Source) by households in Belagunda-3 micro-watershed

SN	Particulars	LL (1)		MF (1)		SF (10)		SMF (2)		MDF (2)		All (16)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Agriculture production	1	100	1	100	10	100	2	100	2	100	16	100

Purpose of credit borrowed (Private Source): The data regarding the purpose of credit borrowed – Private Source in Belagunda-3 micro-watershed is presented in Table 25. The results indicate that, 62.50 per cent of the households have borrowed loan for agriculture production.

Table 25. Purpose of credit borrowed (Private Source) by households in Belagunda-3 micro-watershed

Sl.No.	Particulars	LL (1)		MF (5)		SF (10)		SMF (0)		MDF (0)		All (16)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Agriculture production	0	0	3	60	7	70	0	0	0	0	10	62.5

Repayment status of household (institutional Source): The data regarding the repayment status of credit borrowed from institutional Source by households in Belagunda-3 micro watershed is presented in Table 26. The results indicate that, 100.00 per cent have unpaid.

Table 26. Repayment status of household (institutional Source) in Belagunda-3 micro-watershed

Sl.No.	Particulars	LL (1)		MF (1)		SF (10)		SMF (2)		MDF (2)		All (16)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Un paid	1	100	1	100	10	100	2	100	2	100	16	100

Repayment status of household (Private Source): The data regarding the repayment status of credit borrowed from private sources by households in Belagunda-3 micro watershed is presented in Table 27. The results indicate that, 100 per cent has unpaid.

Table 27. Repayment status of household (Private Source) in Belagunda-3 micro-watershed

Sl.No.	Particulars	LL (1)		MF (5)		SF (10)		SMF (0)		MDF (0)		All (16)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Un paid	1	100	5	100	10	100	0	0	0	0	16	100

Cost of Cultivation of Red gram: The data regarding the cost of cultivation (Rs/ha) of Red gram in Belagunda-3 micro watershed is presented in Table 28.a. The results indicate that, the total cost of cultivation (Rs/ha) for Red gram was Rs. 29784.68. The gross income realized by the farmers was Rs. 50223.15. The net income from Red gram cultivation was Rs.20438.47, thus the benefit cost ratio was found to be 1:1.70.

Table 28(a). Cost of Cultivation of Red gram in Belagunda-3 micro-watershed

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
I	Cost A1				
1	Hired Human Labour	Man days	31.51	5416.36	18.19
2	Bullock	Pairs/day	3.91	3517.21	11.81
3	Tractor	Hours	2.73	2064.33	6.93
4	Machinery	Hours	0.61	368.21	1.24
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	10.85	900.34	3.02
6	FYM	Quintal	1.72	3476.17	11.67
7	Fertilizer + micronutrients	Quintal	3.46	2923.31	9.81
8	Pesticides (PPC)	Kgs / liters	1.46	3511.72	11.79
9	Depreciation charges		0	120.54	0.4
II	Cost B1				
10	Interest on working capital			1298.58	4.36
11	Cost B1 = (Cost A1 + sum of 15 and 16)			23596.76	79.22
III	Cost B2				
12	Rental Value of Land			166.67	0.56
13	Cost B2 = (Cost B1 + Rental value)			23763.43	79.78
IV	Cost C1				
14	Family Human Labour		13.65	3303.56	11.09
15	Cost C1 = (Cost B2 + Family Labour)			27066.99	90.88
V	Cost C2				
16	Risk Premium			10	0.03
17	Cost C2 = (Cost C1 + Risk Premium)			27076.99	90.91
VI	Cost C3				
18	Managerial Cost			2707.7	9.09
19	Cost C3 = (Cost C2 + Managerial Cost)			29784.68	100
VII	Economics of the Crop				
		a) Main Product (q)	9.77	50223.15	
a.	Main Product	b) Main Crop Sales Price (Rs.)		5142.86	
b.	Gross Income (Rs.)			50223.15	
c.	Net Income (Rs.)			20438.47	
d.	Cost per Quintal (Rs./q.)			3049.96	
e.	Benefit Cost Ratio (BC Ratio)			1:1.7	

Cost of Cultivation of Groundnut: The data regarding the cost of cultivation (Rs/ha) of Groundnut in Belagunda-3 micro watershed is presented in Table 28.b. The results indicate that, the total cost of cultivation (Rs/ha) for Groundnut was Rs. 37139.36. The gross income realized by the farmers was Rs. 36695.38. The net income from Groundnut cultivation was Rs.-443.98, thus the benefit cost ratio was found to be 1:1.00.

Table 28(b). Cost of Cultivation of Groundnut in Belagunda-3 micro-watershed

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
I	Cost A1				
1	Hired Human Labour	Man days	23.13	4432.75	11.94
2	Bullock	Pairs/day	6.75	5971.44	16.08
3	Tractor	Hours	2.31	1951.22	5.25
4	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	95.46	7636.41	20.56
5	FYM	Quintal	1.39	3091.47	8.32
6	Fertilizer + micronutrients	Quintal	3.27	2814.57	7.58
7	Pesticides (PPC)	Kgs / liters	1.28	1001.46	2.7
8	Irrigation	Number	2.29	0	0
9	Depreciation charges		0	58.04	0.16
II	Cost B1				
10	Interest on working capital			1746.47	4.7
11	Cost B1 = (Cost A1 + sum of 15 and 16)			28703.85	77.29
III	Cost B2				
12	Rental Value of Land			111.11	0.3
13	Cost B2 = (Cost B1 + Rental value)			28814.96	77.59
IV	Cost C1				
14	Family Human Labour		18.94	4938.09	13.3
15	Cost C1 = (Cost B2 + Family Labour)			33753.05	90.88
V	Cost C2				
16	Risk Premium			10	0.03
17	Cost C2 = (Cost C1 + Risk Premium)			33763.05	90.91
VI	Cost C3				
18	Managerial Cost			3376.31	9.09
19	Cost C3 = (Cost C2 + Managerial Cost)			37139.36	100
VII	Economics of the Crop				
a.	Main Product	a) Main Product (q)		7.98	36695.38
		b) Main Crop Sales Price (Rs.)			4600
b.	Gross Income (Rs.)			36695.38	
c.	Net Income (Rs.)			-443.98	
d.	Cost per Quintal (Rs./q.)			4655.66	
e.	Benefit Cost Ratio (BC Ratio)			1:1	

Cost of Cultivation of Jowar: The data regarding the cost of cultivation (Rs/ha) of Jowar in Belagunda-3 micro watershed is presented in Table 28.c. The results indicate, the total cost of cultivation (Rs/ha) for Jowar was Rs.37139.36. The gross income realized by the farmers was Rs. 36695.38. The net income from Jowar cultivation was Rs. -443.98, thus the benefit cost ratio was found to be 1:1.00.

Table 28(c). Cost of Cultivation of Jowar in Belagunda-3 micro-watershed

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
I	Cost A1				
1	Hired Human Labour	Man days	23.13	4432.75	11.94
2	Bullock	Pairs/day	6.75	5971.44	16.08
3	Tractor	Hours	2.31	1951.22	5.25
4	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	95.46	7636.41	20.56
5	FYM	Quintal	1.39	3091.47	8.32
6	Fertilizer + micronutrients	Quintal	3.27	2814.57	7.58
7	Pesticides (PPC)	Kgs / liters	1.28	1001.46	2.7
8	Irrigation	Number	2.29	0	0
9	Depreciation charges		0	58.04	0.16
II	Cost B1				
16	Interest on working capital			1746.47	4.7
17	Cost B1 = (Cost A1 + sum of 15 and 16)			28703.85	77.29
III	Cost B2				
18	Rental Value of Land			111.11	0.3
19	Cost B2 = (Cost B1 + Rental value)			28814.96	77.59
IV	Cost C1				
20	Family Human Labour		18.94	4938.09	13.3
21	Cost C1 = (Cost B2 + Family Labour)			33753.05	90.88
V	Cost C2				
22	Risk Premium			10	0.03
23	Cost C2 = (Cost C1 + Risk Premium)			33763.05	90.91
VI	Cost C3				
24	Managerial Cost			3376.31	9.09
25	Cost C3 = (Cost C2 + Managerial Cost)			37139.36	100
VII	Economics of the Crop				
a.	Main Product	a) Main Product (q)		7.98	36695.38
		b) Main Crop Sales Price (Rs.)			4600
b.	Gross Income (Rs.)			36695.38	
c.	Net Income (Rs.)			-443.98	
d.	Cost per Quintal (Rs./q.)			4655.66	
e.	Benefit Cost Ratio (BC Ratio)			1:1	

Cost of Cultivation of Paddy: The data regarding the cost of cultivation (Rs/ha) of Paddy in Belagunda-3 micro watershed is presented in Table 28.d. The results indicate that, the total cost of cultivation (Rs/ha) for Paddy was Rs. 36528.36. The gross income realized by the farmers was Rs.25200.94. The net income from Paddy cultivation was Rs. -11327.41, thus the benefit cost ratio was found to be 1:0.70.

Table 28(d). Cost of Cultivation of Paddy in Belagunda-3 micro-watershed

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
I	Cost A1				
1	Hired Human Labour	Man days	70.35	12151	33.26
2	Bullock	Pairs/day	1.62	1299.58	3.56
3	Tractor	Hours	1.84	1780.62	4.87
4	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	71.92	5753.26	15.75
5	FYM	Quintal	1.54	3858.69	10.56
6	Fertilizer + micronutrients	Quintal	4.47	3737.15	10.23
7	Pesticides (PPC)	Kgs / liters	0.66	661.07	1.81
8	Irrigation	Number	1.97	0	0
9	Depreciation charges		0	0.01	0
II	Cost B1				
10	Interest on working capital			1682.42	4.61
11	Cost B1 = (Cost A1 + sum of 15 and 16)			30923.8	84.66
III	Cost B2				
12	Rental Value of Land			125	0.34
13	Cost B2 = (Cost B1 + Rental value)			31048.8	85
IV	Cost C1				
14	Family Human Labour		8.87	2148.8	5.88
15	Cost C1 = (Cost B2 + Family Labour)			33197.6	90.88
V	Cost C2				
16	Risk Premium			10	0.03
17	Cost C2 = (Cost C1 + Risk Premium)			33207.6	90.91
VI	Cost C3				
18	Managerial Cost			3320.76	9.09
19	Cost C3 = (Cost C2 + Managerial Cost)			36528.36	100
VII	Economics of the Crop				
a.	Main Product	a) Main Product (q)		24.59	25200.94
		b) Main Crop Sales Price (Rs.)			1025
b.	Gross Income (Rs.)			25200.94	
c.	Net Income (Rs.)			-11327.41	
d.	Cost per Quintal (Rs./q.)			1485.72	
e.	Benefit Cost Ratio (BC Ratio)			1:0.7	

Cost of Cultivation of Maize: The data regarding the cost of cultivation (Rs/ha) of Maize in Belagunda-3 micro watershed is presented in Table 28.e. The results indicate that, the total cost of cultivation (Rs/ha) for Maize was Rs.22730.15. The gross income realized by the farmers was Rs. 31521.66. The net income from Maize cultivation was Rs. 8791.51, thus the benefit cost ratio was found to be 1:1.40.

Table 28(e). Cost of Cultivation of Maize in Belagunda-3 micro-watershed

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
I	Cost A1				
1	Hired Human Labour	Man days	24.68	5329.03	23.44
2	Bullock	Pairs/day	3.64	2355.93	10.36
3	Tractor	Hours	2.56	2110.65	9.29
4	Machinery	Hours	0.15	92.63	0.41
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	8.22	543.31	2.39
6	FYM	Quintal	0.62	926.25	4.07
7	Fertilizer + micronutrients	Quintal	3.41	3504.31	15.42
8	Pesticides (PPC)	Kgs / liters	0.74	507.72	2.23
9	Depreciation charges		0	1559.12	6.86
II	Cost B1				
10	Interest on working capital			658.99	2.9
11	Cost B1 = (Cost A1 + sum of 15 and 16)			17587.95	77.38
III	Cost B2				
12	Rental Value of Land			166.67	0.73
13	Cost B2 = (Cost B1 + Rental value)			17754.61	78.11
IV	Cost C1				
14	Family Human Labour		11.17	2899.16	12.75
15	Cost C1 = (Cost B2 + Family Labour)			20653.78	90.87
V	Cost C2				
16	Risk Premium			10	0.04
17	Cost C2 = (Cost C1 + Risk Premium)			20663.78	90.91
VI	Cost C3				
18	Managerial Cost			2066.38	9.09
19	Cost C3 = (Cost C2 + Managerial Cost)			22730.15	100
VII	Economics of the Crop				
a.	Main Product	a) Main Product (q)		13.71	31521.66
		b) Main Crop Sales Price (Rs.)			2300
b.	Gross Income (Rs.)			31521.66	
c.	Net Income (Rs.)			8791.51	
d.	Cost per Quintal (Rs./q.)			1658.52	
e.	Benefit Cost Ratio (BC Ratio)			1:1.4	

Adequacy of fodder: The data regarding the adequacy of fodder in Belagunda-3 Micro watershed is presented in Table 29. The results indicate that, 25.71 per cent of the households opined that dry fodder was adequate.

Table 29. Adequacy of fodder in Belagunda-3 micro-watershed

Sl.No.	Particulars	LL (3)		MF (8)		SF (19)		SMF (3)		MDF (2)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Adequate-Dry Fodder	0	0	4	50	3	15.79	2	66.7	0	0	9	25.71

Average annual gross income: The data regarding the annual gross income in Belagunda-3 Micro watershed is presented in Table 30. The results indicate that, the farmers have annual gross income of Rs. 97142.86 in micro-watershed, of which Rs. 50114.29 is from agriculture itself.

Table 30. Average annual gross income in Belagunda-3 micro-watershed

Sl.No.	Particulars	LL (3)	MF (8)	SF (19)	SMF (3)	MDF (2)	All (35)
		Rs.	Rs.	Rs.	Rs.	Rs.	Rs.
1	Service/salary	0	17500	0	0	0	4000
2	Wage	81000	63375	35315.8	13333.3	10000	42314.3
3	Agriculture	14666.7	40375	55568.4	59066.7	77000	50114.3
4	Goat Farming	0	0	1315.79	0	0	714.29
Income(Rs.)		95666.7	121250	92200	72400	87000	97142.9

Average annual Expenditure: The data regarding the average annual expenditure in Belagunda-3 Micro watershed is presented in Table 31. The results indicate that, the farmers have annual gross expenditure of Rs. 261711.95 in micro-watershed, of which Rs. 24928.57 is from agriculture itself.

Table 31. Average annual Expenditure in Belagunda-3 micro-watershed

Sl.No.	Particulars	LL (3)	MF (8)	SF (19)	SMF (3)	MDF (2)	All (35)
		Rs.	Rs.	Rs.	Rs.	Rs.	Rs.
1	Service/salary	0	60000	0	0	0	1714.29
2	Wage	26666.7	25750	14117.7	5000	5000	15457.1
3	Agriculture	25000	19125	30552.6	13000	37500	24928.6
Total		51666.7	104875	44670.3	18000	42500	261712

Horticulture species grown: The data regarding horticulture species grown in Belagunda-3 Micro watershed is presented in Table 32. The results indicate that, the total number of horticultural trees grown (both field and backyard) by the sampled households was Mango (1).

Table 32. Horticulture species grown in Belagunda-3 micro-watershed

Sl.No.	Particulars	LL (3)		MF (8)		SF (19)		SMF (3)		MDF (2)		All (35)	
		F	B	F	B	F	B	F	B	F	B	F	B
1	Mango	0	0	0	0	1	0	0	0	0	0	1	0

*F= Field B=Back Yard

Forest species grown: The data regarding forest species grown in Belagunda-3 Micro watershed is presented in Table 33. The results indicate that, households have

planted 37 neem trees, 4 tamarind trees and 10 banyan trees together in both field and backyard.

Table 33. Forest species grown in Belagunda-3 micro-watershed

Sl.No.	Particulars	LL (3)		MF (8)		SF (19)		SMF (3)		MDF (2)		All (35)	
		F	B	F	B	F	B	F	B	F	B	F	B
1	Neem	0	0	6	4	21	3	3	0	0	0	30	7
2	Tamarind	0	0	1	0	3	0	0	0	0	0	4	0
3	Banyan	0	0	3	0	7	0	0	0	0	0	10	0

*F= Field B=Back Yard

Average additional investment capacity: The data regarding average additional investment capacity in Belagunda-3 Micro watershed is presented in Table 34. The results indicate that, households have an average investment capacity of Rs. 2628.57 for land development, Rs. 16428.57 for creation of irrigation facility, Rs.1714.29 for adoption of improved livestock breeds and Rs.114.29 adoption of improved crop production activities.

Table 34. Average additional investment capacity of households in Belagunda-3 micro-watershed

Sl. No	Particulars	LL (3)	MF (8)	SF (19)	SMF (3)	MDF (2)	All (35)
		Rs.	Rs.	Rs.	Rs.	Rs.	Rs.
1	Land development	2000	5125	1000	3333.33	8000	2628.57
2	Irrigation facility	0	37500	14473.7	0	0	16428.6
3	Improved crop production	1666.67	1125	947.37	2666.67	10000	1714.29
4	Improved livestock management	0	0	0	1333.33	0	114.29

Source of funds for additional investment: The data regarding source of funds for additional investment in Belagunda-3 Micro watershed is presented in Table 35. The results indicate that, the sources of finance raised from bank as a loan and from own sources for land development was 2.86 per cent, for irrigation facility was 8.57, the sources of finance raised from Own funds for land development was 28.57 per cent, for irrigation facility was 8.57.

Table 35. Source of funds for additional investment in Belagunda-3 micro-watershed

Sl. No	Item	Land development		Irrigation facility		Improved crop production		Improved livestock management	
		N	%	N	%	N	%	N	%
1	Government subsidy	0	0	3	8.57	0	0	0	0
2	Loan from bank	1	2.86	3	8.57	0	0	0	0
3	Own funds	10	28.57	0	0	10	28.57	1	2.86

Marketing of agricultural produce: The data regarding marketing of the agricultural produce in Belagunda-3 Micro watershed is presented in Table 36. The results indicated that, 100.00 per cent of output of Cotton was sold in the market with

average price of Rs. 4704.17; 81.25 per cent of output of Groundnut was sold in the market with average price of Rs. 4600.00; 94.74 per cent of output of Jowar was sold in the market with average price of Rs. 2150.00; 75.31 per cent of output of Maize was sold in the market with average price of Rs. 2300.00 and 88.10 per cent of output of Paddy was sold in the market with average price of Rs. 1025.00.

Table 36. Marketing of agricultural produce in Belagunda-3 micro-watershed

Sl.No	Crops	Output obtained (q)	Output retained (q)	Output sold (q)	Output sold (%)	Avg. Price obtained (Rs/q)
1	Cotton	122	0	122	100	4704
2	Groundnut	32	6	26	81	4600
3	Jowar	19	1	18	95	2150
4	Maize	81	20	61	75	2300
5	Paddy	210	25	185	88	1025
6	Red gram	87	10	77	89	5143

Marketing channels used for sale of agricultural produce: The data regarding marketing channels used for sale of agricultural produce in Belagunda-3 Micro watershed is presented in Table 37. The results indicated that, 91.43 cent of the households have sold agricultural produce to the local/village merchants.

Table 37. Marketing channels used for sale of agricultural produce in Belagunda-3 micro-watershed

Sl.No.	Particulars	LL (3)		MF (8)		SF (19)		SMF (3)		MDF (2)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Local/village Merchant	1	33.33	8	100	18	94.7	3	100	2	100	32	91.43

Mode of transport of agricultural produce: The data regarding mode of transport of agricultural produce in Belagunda-3 Micro watershed is presented in Table 38. The results indicated that, 91.43 cent of the households have used tractor for the transport of agriculture commodity.

Table 38. Mode of transport of agricultural produce in Belagunda-3 micro-watershed

Sl.No.	Particulars	LL (3)		MF (8)		SF (19)		SMF (3)		MDF (2)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Tractor	1	33.33	8	100	18	94.7	3	100	2	100	32	91.43

Incidence of soil and water erosion problems: The data regarding incidence of incidence of soil and water erosion problems in Belagunda-3 Micro watershed is presented in Table 39. The results indicate that, 28.57 per cent of the households have experienced soil and water erosion problems.

Table 39. Incidence of soil and water erosion problems in Belagunda-3 micro-watershed

Sl. No.	Particulars	LL (3)		MF (8)		SF (19)		SMF (3)		MDF (2)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Soil and water erosion problems in the farm	1	33	3	38	3	15.8	1	33	2	100	10	28.57

Table 40. Interest regarding soil testing in Belagunda-3 micro-watershed

Sl.No.	Particulars	LL (3)		MF (8)		SF (19)		SMF (3)		MDF (2)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Interest in soil test	1	33.33	8	100	19	100	3	100	2	100	33	94.29

Interest towards soil testing: The data regarding Interest shown towards soil testing in Belagunda-3 Micro watershed is presented in Table 40. The results indicated that, 94.29 per cent of the households were interested towards soil testing.

Usage pattern of fuel for domestic use: The data on usage pattern of fuel for domestic use in Belagunda-3 Micro watershed is presented in Table 41. The results indicated that, firewood was the major source of fuel for domestic use for 65.71 per cent of the households followed by LPG (37.14%), Kerosene (2.86 %).

Table 41. Usage pattern of fuel for domestic use in Belagunda-3 micro-watershed

Sl.No.	Particulars	LL (3)		MF (8)		SF (19)		SMF (3)		MDF (2)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Fire Wood	2	66.7	7	87.5	11	57.9	1	33.3	2	100	23	65.71
2	Kerosene	0	0	1	12.5	1	5.26	0	0	0	0	2	5.71
3	LPG	1	33.3	1	12.5	9	47.4	2	66.7	0	0	13	37.14

Source of drinking water: The data on source of drinking water in Belagunda-3 Micro watershed is presented in Table 42. The results indicated that, piped waters supply was the major source for drinking water for 100 per cent of the households.

Table 42. Source of drinking water in Belagunda-3 micro-watershed

Sl.No.	Particulars	LL (3)		MF (8)		SF (19)		SMF (3)		MDF (2)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Piped supply	3	100	8	100	19	100	3	100	2	100	35	100

Source of light: The data on source of light in Belagunda-3 Micro watershed is presented in Table 43. The results indicated that, electricity was the major source of light for 97.14 per cent of the households followed by kerosene lamp (2.86 %).

Table 43. Source of light in Belagunda-3 micro-watershed

Sl.No.	Particulars	LL (3)		MF (8)		SF (19)		SMF (3)		MDF (2)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Kerosene Lamp	1	33.3	0	0	0	0	0	0	0	0	1	2.86
2	Electricity	2	66.7	8	100	19	100	3	100	2	100	34	97.14

Existence of sanitary toilet facility: The data on availability of toilet facility in Belagunda-3 Micro watershed is presented in Table 44. The results indicated that, 51.43 per cent of the households possess toilets.

Table 44. Existence of sanitary toilet facility in Belagunda-3 micro-watershed

Sl.No.	Particulars	LL (3)		MF (8)		SF (19)		SMF (3)		MDF (2)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Sanitary toilet facility	2	66.7	5	63	8	42.11	2	67	1	50	18	51.4

Possession of PDS card: The data regarding possession of PDS card in Belagunda-3 Micro watershed is presented in Table 45. The results indicated that, 100.00 per cent of the households possessed BPL card.

Table 45. Possession of PDS card in Belagunda-3 micro-watershed

Sl.No.	Particulars	LL (3)		MF (8)		SF (19)		SMF (3)		MDF (2)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	BPL	3	100	8	100	19	100	3	100	2	100	35	100

Participation in NREGA programme: The data regarding Participation in NREGA programme in Belagunda-3 Micro watershed is presented in Table 46. The results indicated that, only 48.57 per cent of the households have participated in NREGA programme.

Table 46. Participation in NREGA programme in Belagunda-3 micro-watershed

Sl.No.	Particulars	LL (3)		MF (8)		SF (19)		SMF (3)		MDF (2)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Participation in NREGA programme	0	0	3	37.5	13	68.4	1	33.3	0	0	17	48.6

Adequacy of food items: The data regarding adequacy of food items in Belagunda-3 Micro watershed is presented in Table 47. The results indicated that, the extent of adequacy of food items for cereals, pulses, Oilseeds and vegetables were 100.00, 97.14, 40.00, 28.57 per cent respectively, similarly for Fruits (45.71%) and milk (42.86%).

Table 47. Adequacy of food items in Belagunda-3 micro-watershed

Sl.No.	Particulars	LL (3)		MF (8)		SF (19)		SMF (3)		MDF (2)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Cereals	2	66.7	8	100	19	100	4	133	2	100	35	100
2	Pulses	2	66.7	8	100	19	100	3	100	2	100	34	97.14
3	Oilseed	1	33.3	4	50	4	21.05	3	100	2	100	14	40
4	Vegetables	1	33.3	2	25	3	15.79	2	66.7	2	100	10	28.57
5	Fruits	1	33.3	4	50	11	57.89	0	0	0	0	16	45.71
6	Milk	1	33.3	3	37.5	11	57.89	0	0	0	0	15	42.86

Table 48. Inadequacy of food items in Belagunda-3 micro-watershed

Sl.No.	Particulars	LL (3)		MF (8)		SF (19)		SMF (3)		MDF (2)		All (35)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Cereals	1	33.3	0	0	0	0	0	0	0	0	1	2.86
2	Pulses	1	33.3	0	0	0	0	0	0	0	0	1	2.86
3	Oilseed	2	66.7	4	50	15	78.95	0	0	0	0	21	60
4	Vegetables	2	66.7	6	75	16	84.21	1	33.3	0	0	25	71.43
5	Fruits	1	33.3	3	37.5	5	26.32	1	33.3	0	0	10	28.57
6	Milk	2	66.7	5	62.5	8	42.11	3	100	2	100	20	57.14
7	Egg	2	66.7	6	75	16	84.21	1	33.3	0	0	25	71.43
8	Meat	2	66.7	6	75	16	84.21	1	33.3	0	0	25	71.43

Inadequacy of food items: The data regarding in adequacy of food items in Belagunda-3 Micro watershed is presented in Table 48. The results indicated that, the

extent of inadequacy of food items for cereals, pulses, Oilseeds and vegetables were 2.86, 2.86, 60.00, 71.43 and 71.43 per cent respectively, similarly for fruits (28.57%), milk (57.14%), egg (71.43%) and meat (71.43%).

Farming constraints: The data regarding farming constraints experienced by households in Belagunda-3 Micro watershed is presented in Table 49. The results indicated that, lower fertility status of the soil was the constraint experienced by (97.14 %) per cent of the households, wild animal menace on farm field (91.43%), frequent incidence of pest and diseases (91.43%), inadequacy of irrigation water (80.00%), high cost of fertilizers and plant protection chemicals (91.43%), high rate of interest on credit (91.43%), low price for the agricultural commodities (60.00 %), lack of marketing facilities in the area (60.00%), inadequate extension services (54.29 %), lack of transport for safe transport of the agricultural produce to the market (60.00%) and less rainfall (2.86%).

Table 49. Farming constraints experienced in Belagunda-3 micro-watershed

SN	Particulars	MF (8)		SF (19)		SMF (3)		MDF (2)		All (35)	
		N	%	N	%	N	%	N	%	N	%
1	Lower fertility status of the soil	8	100	20	105.26	3	100	2	100	34	97.14
2	Wild animal menace on farm field	8	100	18	94.74	3	100	2	100	32	91.43
3	Frequent incidence of pest and diseases	8	100	18	94.74	3	100	2	100	32	91.43
4	Inadequacy of irrigation water	7	87.5	15	78.95	3	100	2	100	28	80
5	High cost of Fertilizers and plant protection chemicals	8	100	18	94.74	3	100	2	100	32	91.43
6	High rate of interest on credit	8	100	18	94.74	3	100	2	100	32	91.43
7	Low price for the agricultural commodities	6	75	14	73.68	1	33.33	0	0	21	60
8	Lack of marketing facilities in the area	6	75	14	73.68	1	33.33	0	0	21	60
9	Inadequate extension services	4	50	14	73.68	1	33.33	0	0	19	54.29
10	Lack of transport for safe transport of the Agril produce to the market.	5	62.5	15	78.95	1	33.33	0	0	21	60
11	Less rainfall	1	12.5	0	0	0	0	0	0	1	2.86

SUMMARY AND IMPLICATIONS

In order to assess the socio-economic condition of the farmers in the watershed 35 households located in the micro watershed were interviewed for the survey. The study was conducted in Belagunda-3 micro-watershed (Belagunda sub-watershed, Yadgiri taluk & District) is located at North latitude 16° 34' 11.759" and 16° 32' 43.028" and East longitude 77° 12' 8.355" and 77° 10' 11.177" covering an area of about 474.74 ha bounded by under under Anura. B and Belagundi Villages.

Socio-economic analysis of Belagunda-3 micro watersheds of Belagunda sub-watershed, Yadgiri taluk & District indicated that, out of the total sample of 35 farmers were sampled in Belagunda-3 micro-watershed among households surveyed 8 (22.86%) were marginal, 19(54.29%) were small, 3 (8.57 %) were semi medium and 2 (5.71 %) were medium farmers. 3 landless farmers were also interviewed for the survey. The population characteristics of households indicated that, there were 91 (54.49%) men and 76 (45.51 %) were women. The average population of landless was 4.3, marginal farmers were 5.4, small farmers were 4.5, semi medium farmers were 6 and medium farmers were 3.5.

Majority of the respondents (44.31%) were in the age group of 16-35 years. Education level of the sample households indicated that, there were 53.29 per cent illiterates, 42.52 per cent pre university education and 4.79 per cent attained graduation. About, 88.57 per cent of household heads practicing agriculture and 5.71 per cent of the household heads were engaged as agricultural labourers. Agriculture was the major occupation for 64.67 per cent of the household members.

In the study area, 82.86 per cent of the households possess katcha house and 5.71 per cent possess pucca house. The durable assets owned by the households showed that, 77.14 per cent possess TV, 40.00 per cent possess mixer grinder, 94.29 per cent possess mobile phones and 31.43 per cent possess motor cycles.

Farm implements owned by the households indicated that, 25.71 per cent of the households possess plough, 2.86 per cent possess tractor, 2.86 per cent possess bullock cart and 5.71 per cent possess sprayer. Regarding livestock possession by the households, 14.29 per cent possess local cow and 2.86 per cent possess buffalo.

The average labour availability in the study area showed that, own labour men available in the micro watershed was 1.85, women available in the micro watershed was 1.18, hired labour (men) available was 13.09 and hired labour (women) available was 9.71. Out of the total land holding of the sample respondents 71.75 per cent (43.07 ha) of the area is under dry condition and the remaining 22.97 per cent area is irrigated land.

The major crops grown by sample farmers are Red gram, Groundnut, Jowar, Paddy and Maize and cropping intensity was recorded as 100.00 per cent. Out of the sample households 85.71 percent possessed bank account and 54.29 per cent of them have savings in the account. About 100.00 per cent of the respondents borrowed credit from various sources.

Among the credit borrowed by households, 55.00 per cent have borrowed loan from commercial banks and 15.00 per cent from co-operative/Grameena bank. Majority of the respondents (100.00%) have borrowed loan for agriculture purpose. The per hectare cost of cultivation for Red gram, Groundnut, Jowar, Paddy and Maize was Rs.29784.68 , 37139.36, 37139.36, 36528.36 and 22730.15 with benefit cost ratio of 1:1.70, 1: 1.00, 1: 1.00, 1: 0.70 and 1:1.40 respectively.

Further, 25.71 per cent of the households opined that dry fodder was adequate and 0.00 per cent of the households have opined that the green fodder was adequate. The average annual gross income of the farmers was Rs. 97142.86 in micro-watershed, of which Rs. 50114.29 comes from agriculture. Sampled households have grown 1 horticulture trees and 51 forestry trees together in the fields and back yards.

Households have an average investment capacity of Rs. 2628.57 for land development and Rs. 16428.57 for irrigation facility. Source of funds for additional investment is concerned, 2.86 per cent depends on own funds for land development activities. Regarding marketing channels, 91.43 per cent of the households have sold agricultural produce to the local/village merchants. Further, 91.43 per cent of the households have used tractor for the transport of agriculture commodity.

Majority of the farmers (28.57%) have experienced soil and water erosion problems in the watershed and 94.29 per cent of the households were interested towards soil testing. Fire was the major source of fuel for domestic use for 65.71 per cent of the households and 37.14 per cent households has LPG connection. Piped supply was the major source for drinking water for 100.00 per cent of the households.

Electricity was the major source of light for 97.14 per cent of the households. In the study area, 51.43 per cent of the households possess toilet facility. Regarding possession of PDS card, 100.00 per cent of the households possessed BPL card. Households opined that, the requirement of cereals (100.00%), pulses (97.14%) and oilseeds (40.00%) are adequate for consumption.

Farming constraints experienced by households in the micro watersheds were lower fertility status of the soil (97.14%) wild animal menace on farm field (91.43%), frequent incidence of pest and diseases (91.43%), inadequacy of irrigation water (80.00%), high cost of fertilizers and plant protection chemicals (91.43%), high rate of interest on credit (91.43%), low price for the agricultural commodities (60.00%),

lack of marketing facilities in the area (60.00%), inadequate extension services (54.29%), lack of transport for safe transport of the agricultural produce to the market (60.00%) and Less rainfall (2.86%).

Implications of the survey

- ✓ Result indicated that, there were 53.29 per cent were illiterate hence, extension methodologies such as demonstration, street play, drama, video shows will be effective in dissemination of the technologies in the micro watershed.
- ✓ The data indicate that, 82.86 per cent of the households possess katcha house. Hence, the development department while implementing the watershed plan should focus on agriculture to enhance the productivity of major crops in the area to increase the income of the farmers.
- ✓ Results indicated that the local institutional participation of the household members in the micro watershed is minimal hence, activities like membership campaign, awareness creation about the benefits of membership in local institutions and strengths of organized groups must be conveyed.
- ✓ Majority of the households in the watershed have experience in use of mobile phones, and television hence, these mass media can be effectively utilized for transfer of technology as well as for information dissemination.
- ✓ The farm machinery/implement possession in the micro watershed was found to be minimum the reasons may lack of knowledge or lack of financial ability which can be addressed through training on use of different farm implements, providing information on different sources of finance for purchase of farm implements.
- ✓ The possession of livestock such as crossbred cow found is less hence, farmers must be made aware of the benefits of crossbred cow in increased milk production.
- ✓ The possession of livestock such as sheep, goat and poultry was found to be low hence, farmers may be informed the role of subsidiary enterprises in enhancing the income and information on financial support for subsidiary activities.
- ✓ The data indicate that, job/work was the reason for all the migrants hence, farmers may be trained on profitable agriculture or self employment such as animal husbandry, plate making, sheep rearing, goat rearing, rabbit rearing with suitable information on sources of financial support.
- ✓ The results indicate that there was a change in quality of life due to migration hence, the developmental departments should take actions to arrest migration and to improve the quality of the life in rural areas.
- ✓ Households possess 31.86ha (71.75 %) of dry land and 10.20ha (22.97 %) of irrigated land hence, the availability of the dryland agricultural technologies such as short duration crops, high yielding drought resistance crop varieties, drip

irrigation technology and subsidy information will be helpful for the farmers to enhance the productivity of land and as well as farmers income.

- ✓ Few of the bore well in micro watershed found non functional hence, farmers may be trained on possibility of bore well rejuvenation.
- ✓ In order to increase the area under irrigation as well as to increase the water use efficiency farmers may train on drip irrigation and provides the information on subsidy for drip irrigation equipment's along with the information on different agencies which provides the financial assistance for drip irrigation.
- ✓ The cropping intensity in the micro watershed was found to be (100.00 %) hence, care must be taken by the implementing agency to bring uncultivated land into cultivation through suitable measures.
- ✓ Many of the household members have borrowed loan from cooperative banks which has higher rate of interest hence, farmers may be sensitized on the different sources of credit with lesser interest rate such SHGs etc.
- ✓ The results indicated the non availability of both green and dry fodder throughout the year hence, fodder development activities can be taken up in the micro watershed.
- ✓ The average annual gross income of the households Rs.50114.29 from agriculture and Rs. 42314.29 from wages and. Agriculture was found to be the major source of income for households hence; the development activities should focus on productivity enhancement, marketing arrangements and agricultural technology dissemination to have a direct impact on the farmers.
- ✓ The cultivation of forest species is found minimal hence, information and production technology related to agro-forestry and integrated farming system.
- ✓ The data indicated that, 28.57 per cent of the households have experienced soil and water erosion problems. Hence, those farmers who reported the soil and water erosion problems may be given attention while implementation of the watershed development plan.
- ✓ The data indicated that, 94.29 per cent of the households have interest in soil testing hence, farmers must be provided with the information on various institutions which are involved in soil testing for the benefit of the farmers.
- ✓ Except summer ploughing the adoption of other soil and water conservation structures is minimum hence, the farmers in the micro watershed should be sensitized on the use of different conservation structures for soil water conservation.
- ✓ Cereals and pulses found be adequate for per cent of the households respectively hence, farm households and the farm women must be trained on importance of balanced nutrition and role of vegetable, milk, egg, meat in balanced diet.
- ✓ Lower fertility status of the soil (97.14%), wild animal menace on farm field (91.43%), frequent incidence of pest and diseases (91.43%), high cost of

fertilizers and plant protection chemicals (91.43%), high rate of interest on credit (91.43%), low price for the agricultural commodities (60.00%), lack of marketing facilities in the area (60.00%), inadequate extension services (54.29%), lack of transport for safe transport of the agricultural produce to the market (60.00%) were the major farming constraints experienced hence, these constraints must be addressed immediately for the welfare of the farmers. Awareness to be created among the farmers to approach nearest KVKs/RSKs and other developmental departments for technical and for subsidized inputs and utilize the well established regulated markets, approaching the contract firms, direct markets to avoid the involvement of middlemen.