



हर कदम, हर डगर  
किसानों का हमसफर  
भारतीय कृषि अनुसंधान परिषद

*AgriSearch with a human touch*

**LAND RESOURCE INVENTORY AND SOCIO-ECONOMIC STATUS OF  
FARM HOUSEHOLDS FOR WATERSHED PLANNING AND  
DEVELOPMENT**

**MUNDRAGI -3 (4D5B1H2c) MICROWATERSHED**

**Yadgir Hobli, Yadgir Taluk and District, Karnataka**

**Karnataka Watershed Development Project – II**

**SUJALA – III**

**World Bank funded Project**



**ICAR – NATIONAL BUREAU OF SOIL SURVEY AND LAND USE PLANNING**



ICAR - NBSS & LUP



**WATERSHED DEVELOPMENT DEPARTMENT  
GOVT. OF KARNATAKA, BANGALORE**



## **About ICAR - NBSS&LUP**

The ICAR-National Bureau of Soil Survey and Land Use Planning (ICAR-NBSS&LUP), Nagpur, a premier Institute of the Indian Council of Agricultural Research (ICAR), was set up during 1976 with the objective to prepare soil resource maps at national, state and district levels and to provide research inputs in soil resource mapping and its applications, land evaluation, land use planning, land resource management, and database management using GIS for optimising land use on different kinds of soils in the country.

The Bureau has been engaged in carrying out soil resource survey, agro-ecological and soil degradation mapping at the country, state and district levels for qualitative assessment and monitoring the soil health towards viable land use planning. The research activities have resulted in identifying the soil potentials and problems, and the various applications of the soil surveys with the ultimate objective of sustainable agricultural development. The Bureau has the mandate to correlate and classify soils of the country and maintain a National Register of all the established soil series. The Institute is also imparting in-service training to staff of the soil survey agencies in the area of soil survey, land evaluation and soil survey interpretations for land use planning. The Bureau in collaboration with Panjabrao Krishi Vidyapeeth, Akola is running post-graduate teaching and research programme in land resource management, leading to M.Sc. and Ph.D. degrees.

**Citation:** Rajendra Hegde, Ramesh Kumar, S.C., B.A. Dhanorkar, S. Srinivas, M. Lalitha, K.V. Niranjana, R.S. Reddy and S.K. Singh (2019). "Land Resource Inventory and Socio-Economic Status of Farm Households for Watershed Planning and Development of Mundragi-3 (4D5B1H2d) Microwatershed, Yadgir Hobli, Yadgir Taluk and District, Karnataka", ICAR-NBSS&LUP Sujala MWS Publ.276, ICAR – NBSS & LUP, RC, Bangalore. p.131 & 26.

### **TO OBTAIN COPIES,**

**Please write to:**

**Director, ICAR - NBSS & LUP,**

Amaravati Road, NAGPUR - 440 033, India

Phone : (0712) 2500386, 2500664, 2500545 (O)

Telefax : 0712-2522534

E-Mail : [director@nbsslup.ernet.in](mailto:director@nbsslup.ernet.in)

Website URL : [nbsslup.in](http://nbsslup.in)

Or

**Head, Regional Centre, ICAR - NBSS&LUP, Hebbal, Bangalore - 560 024**

Phone : (080) 23412242, 23510350 (O)

Telefax : 080-23510350

E-Mail : [nbssrcb@gmail.com](mailto:nbssrcb@gmail.com)

ICAR-NBSS&LUP Sujala MWS Publ.276



**LAND RESOURCE INVENTORY AND SOCIO-ECONOMIC  
STATUS OF FARM HOUSEHOLDS FOR WATERSHED  
PLANNING AND DEVELOPMENT**

**MUNDRAGI -3 (4D5B1H2c) MICROWATERSHED**

**Yadgir Hobli, Yadgir Taluk and District, Karnataka**

**Karnataka Watershed Development Project – II  
Sujala-III**

**World Bank funded Project**



**ICAR – NATIONAL BUREAU OF SOIL SURVEY AND LAND  
USE PLANNING**



ICAR - NBSS & LUP



**WATERSHED DEVELOPMENT DEPARTMENT, GOVT. OF  
KARNATAKA, BANGALORE**





## PREFACE

In Karnataka, as in other Indian States, the livelihoods of rural people are intertwined with farming pursuits. The challenges in agriculture are seriously threatening the livelihood of a large number of farmers as they have been practicing farming in contextual factors beyond their control. Climatic factors are the most important ones and have become much more significant in recent times due to rapid climate changes induced by intensive anthropogenic activities affecting our ecosystem in multiple ways. Climate change has become the reality, it is happening and efforts to evolve and demonstrate climate resilient technologies have become essential. Due to the already over stressed scenario of agrarian sector, the climate change is resulting in manifold increase in the complexities, pushing the rural mass to face more and more unpredictable situations. The rising temperatures and unpredictable rainfall patterns are going to test seriously the informed decisions farmers have to make in order to survive in farming and sustain their livelihood.

It is generally recognized that impacts of climate change shall not be uniform across the globe. It is said that impact of climate change is more severe in South Asia. Based on the analysis of meteorological data, it is predicted that in India, there will be upward trend in mean temperature, downward trend in relative humidity, annual rainfall and number of wet days in a year. Also, in general, phenomena like erratic monsoon, spread of tropical diseases, rise in sea levels, changes in availability of fresh water, frequent floods, droughts, heat waves, storms and hurricanes are predicted. Each one of these adverse situations are already being experienced in various parts of India and also at the global level. Decline in agricultural productivity of small and marginal farmers becoming more vulnerable is already witnessed.

In Karnataka, more than 60 per cent of the population live in rural areas and depend on agriculture and allied activities for their livelihood. Though the state has achieved significant progress in increasing the yield of many crops, there is tremendous pressure on the land resources due to the growing and competing demands of various land uses. This is reflected in the alarming rate of land degradation observed. Already more than 50 per cent of the area is affected by various forms of degradation. If this trend continues, the sustainability of the fragile ecosystem will be badly affected. The adverse effects of change in the climatic factors are putting additional stress on the land resources and the farmers dependent on this.

The natural resources (land, water and vegetation) of the state need adequate and constant care and management, backed by site-specific technological interventions and investments particularly by the government. Detailed database pertaining to the nature of the land resources, their constraints, inherent potentials and suitability for various land based rural enterprises, crops and other uses is a prerequisite for preparing location-specific action plans, which are in tune with the inherent capability of the resources. Any

effort to evolve climate resilient technologies has to be based on the baseline scientific database. Then only one can expect effective implementation of climate resilient technologies, monitor the progress, make essential review of the strategy, and finally evaluate the effectiveness of the implemented programs. The information available at present on the land resources of the state are of general nature and useful only for general purpose planning. Since the need of the hour is to have site-specific information suitable for farm level planning and detailed characterization and delineation of the existing land resources of an area into similar management units is the only option.

ICAR-NBSS&LUP, Regional Centre, Bangalore has taken up a project sponsored by the Karnataka Watershed Development Project-II, (Sujala-III), Government of Karnataka funded by the World Bank under Component-1 Land Resource Inventory. This study was taken up to demonstrate the utility of such a database in reviewing, monitoring and evaluating all the land based watershed development programs on a scientific footing. To meet the requirements of various land use planners at grassroots level, the present study on “Land Resource Inventory and Socio-Economic Status of Farm Households for Watershed Planning and Development of Mundragi-3 Microwatershed, Yadgir Taluk and District, Karnataka” for integrated development was taken up in collaboration with the State Agricultural Universities, IISC, KSRSAC, KSNDMC as Consortia partners. The project provides detailed land resource information at cadastral level (1:7920 scale) for all the plots and socio-economic status of farm households covering thirty per cent farmers randomly selected representing landed and landless class of farmers in the microwatershed. The project report with the accompanying maps for the Microwatershed will provide required detailed database for evolving effective land use plan, alternative land use options and conservation plans for the planners, administrators, agricultural extension personnel, KVK officials, developmental departments and other land users to manage the land resources in a sustainable manner.

It is hoped that this database will be useful to the planners, administrators and developmental agencies working in the area in not only for formulating location specific developmental schemes but also for their effective monitoring at the village/watershed level.

Nagpur

Date: 25-07-2019

**S.K. SINGH**

Director, ICAR - NBSS&LUP, Nagpur

## Contributors

<b>Dr. Rajendra Hegde</b> Principal Scientist, Head & Project Leader, Sujala-III Project ICAR-NBSS&LUP, Regional Centre, Bangalore	<b>Dr. S.K.Singh</b> Director, ICAR-NBSS&LUP Coordinator, Sujala-III Project Nagpur
<b>Soil Survey, Mapping &amp; Report Preparation</b>	
Dr. B.A. Dhanorkar	Sh. R.S. Reddy
Dr. K.V. Niranjana	Mr. Somashekar T N
	Smt. Chaitra, S.P.
	Dr. Gopali bardhan
	Ms. Arpitha
	Dr. Mahendra Kumar, M.B.
<b>Field Work</b>	
Sh. C.BacheGowda	Sh. Mahesh, D.B.
Sh. Somashekar	Sh. Ashok S Sindagi
Sh. M. Jayaramaiah	Sh. Veerabhadrapa B.
Sh. Paramesha, K.	Sh. Shankarappa
Sh. B. M. Narayana Reddy	Sh. Anand
	Sh. Arun N Kambar.
	Sh Kamalesh Awate
	Sh. Sharaan Kumar Huppar
	Sh. Yogesh H.N.
	Sh. Kalaveerachari R Kammar
<b>GIS Work</b>	
Dr. S.Srinivas	Sh. A.G.Devendra Prasad
Sh. D.H.Venkatesh	Sh. Prakashanaik, M.K.
Smt.K.Sujatha	Sh. Abhijith Sastry, N.S.
Smt. K.V.Archana	Sh. Sudip Kumar Suklabaidya
Sh. N. Maddileti	Sh. Avinash, K.N.
	Sh. Amar Suputhra, S
	Sh. Deepak, M.J.
	Smt. K.Karunya Lakshmi
	Ms. Seema, K.V.
	Ms. A. Rajab Nisha

<b>Laboratory Analysis</b>	
Dr. K.M.Nair	Ms. Steffi Peter
Smt. Arti Koyal	Ms. Thara, V.R
Smt. Parvathy	Ms. Roopa, G.
	Ms. Swati, H.
	Sh. Shantaveera Swami
	Ms. Shwetha, N.K.
	Smt. Ishrat Haji
	Ms. P. Pavan Kumari
	Ms. Padmaja
	Ms. Veena, M.
<b>Socio-Economic Analysis</b>	
Dr. S.C. Ramesh Kumar	Sh. M.K. Prakashanaik
	Ms. Karuna V. Kulkarni
	Mrs. Sowmya A.N
	Sh. Vinod R
	Sh. Basavaraja
	Sh. Vijay Kumar Lamani
	Ms. Sowmya K.B
	Mrs. Prathibha, D.G
	Sh. Rajendra,D
<b>Soil &amp; Water Conservation</b>	
Sh. Sunil P. Maske	
<b>Watershed Development Department, GoK, Bangalore</b>	
Sh. Rajeev Ranjan IFS Project Director & Commissioner, WDD	Dr. A. Natarajan NRM Consultant, Sujala-III Project
Dr. S.D. Pathak IFS Executive Director & Chief Conservator of Forests, WDD	



# **PART-A**

## **LAND RESOURCE INVENTORY**



## Contents

Preface		
Contributors		
Executive Summary		
Chapter 1	Introduction	1
Chapter 2	Geographical Setting	3
2.1	Location and Extent	3
2.2	Geology	4
2.3	Physiography	4
2.4	Drainage	4
2.5	Climate	5
2.6	Natural Vegetation	6
2.7	Land Utilization	6
Chapter 3	Survey Methodology	11
3.1	Base maps	11
3.2	Image Interpretation for Physiography	11
3.3	Field Investigation	14
3.4	Soil Mapping	15
3.5	Land Management Units	16
3.6	Laboratory Characterization	16
Chapter 4	The Soils	21
4.1	Soils of granite gneiss landscape	21
Chapter 5	Interpretation for Land Resource Management	35
5.1	Land Capability Classification	35
5.2	Soil Depth	37
5.3	Surface Soil Texture	38
5.4	Soil Gravelliness	39
5.5	Available Water Capacity	40
5.6	Soil Slope	41
5.7	Soil Erosion	42
Chapter 6	Fertility Status	45
6.1	Soil Reaction (pH)	45
6.2	Electrical Conductivity (EC)	45
6.3	Organic Carbon (OC)	45
6.4	Available Phosphorus	47
6.5	Available Potassium	47
6.6	Available Sulphur	47
6.7	Available Boron	47
6.8	Available Iron	48
6.9	Available Manganese	48
6.10	Available Copper	48
6.11	Available Zinc	48

Chapter 7	Land Suitability for Major Crops	53
7.1	Land suitability for Sorghum	53
7.2	Land suitability for Maize	54
7.3	Land suitability for Bajra	55
7.4	Land suitability for Groundnut	56
7.5	Land suitability for Sunflower	57
7.6	Land suitability for Redgram	58
7.7	Land suitability for Bengal gram	59
7.8	Land suitability for Cotton	60
7.9	Land suitability for Chilli	61
7.10	Land suitability for Tomato	62
7.11	Land suitability for Brinjal	63
7.12	Land suitability for Onion	64
7.13	Land suitability for Bhendi	65
7.14	Land suitability for Drumstick	66
7.15	Land suitability for Mango	67
7.16	Land suitability for Guava	68
7.17	Land suitability for Sapota	69
7.18	Land Suitability for Pomegranate	70
7.19	Land Suitability for Musambi	71
7.20	Land Suitability for Lime	72
7.21	Land Suitability for Amla	73
7.22	Land Suitability for Cashew	74
7.23	Land Suitability for Jackfruit	75
7.24	Land Suitability for Jamun	76
7.25	Land Suitability for Custard apple	77
7.26	Land Suitability for Tamarind	78
7.27	Land Suitability for Mulberry	79
7.28	Land Suitability for Marigold	80
7.29	Land Suitability for Chrysanthemum	81
7.30	Land Management Units	113
7.31	Proposed Crop Plan	114
Chapter 8	Soil Health Management	117
Chapter 9	Soil and Water conservation Treatment Plan	123
9.1	Treatment Plan	124
9.2	Recommended Soil and Water Conservation measures	127
9.3	Greening of Microwatershed	128
	References	131
	Appendix I	I-IV
	Appendix II	V-XII
	Appendix III	XIII-XVIII

## LIST OF TABLES

2.1	Mean Monthly Rainfall, PET, 1/2 PET at Yadgir Taluk & District	5
2.2	Land Utilization in Yadgir taluk	7
3.1	Differentiating Characteristics used for Identifying Soil Series	15
3.2	Soil map unit description of Mundragi-3 Microwatershed	16
4.1	Physical and Chemical Characteristics of Soil Series identified in Mundragi-3 microwatershed	27
7.1	Soil-Site Characteristics of Mundragi-3 Microwatershed	83
7.2	Land suitability for Sorghum	84
7.3	Land suitability for Maize	85
7.4	Land suitability for Bajra	86
7.5	Land suitability for Groundnut	87
7.6	Land suitability for Sunflower	88
7.7	Land suitability for Redgram	89
7.8	Land suitability for Bengal gram	90
7.9	Land suitability for Cotton	91
7.10	Land suitability for Chilli	92
7.11	Land suitability for Tomato	93
7.12	Land suitability for Brinjal	94
7.13	Land suitability for Onion	95
7.14	Land suitability for Bhendi	96
7.15	Land suitability for Drumstick	97
7.16	Land suitability for Mango	98
7.17	Land suitability for Guava	99
7.18	Land suitability for Sapota	100
7.19	Land suitability for Pomegranate	101
7.20	Land suitability for Musambi	102
7.21	Land suitability for Lime	103
7.22	Land suitability for Amla	104
7.23	Land suitability for Cashew	105
7.24	Land suitability for Jackfruit	106

7.25	Land suitability for Jamun	107
7.26	Land suitability for Custard apple	108
7.27	Land suitability for Tamarind	109
7.28	Land suitability for Mulberry	110
7.29	Land suitability for Marigold	111
7.30	Land suitability for Chrysanthemum	112
7.31	Proposed Crop Plan for Mundragi-3 Microwatershed	115

## LIST OF FIGURES

2.1	Location map of Mundragi-3 Microwatershed	3
2.2	Granite and granite gneiss rock formation	4
2.3	Rainfall distribution in Yadgir Taluk & District	5
2.4	Natural vegetation of Mundragi-3 Microwatershed	6
2.5	Current Land use map of Mundragi-3 Microwatershed	7
2.6 a	Major crops and cropping systems in Mundragi-3 Microwatershed	8
2.7	Location of wells - Mundragi-3 Microwatershed	9
3.1	Scanned and Digitized Cadastral map of Mundragi-3 Microwatershed	12
3.2	Satellite image of Mundragi-3 Microwatershed	13
3.3	Cadastral map overlaid on IRS PAN+LISS IV merged imagery of Mundragi-3 Microwatershed	13
3.4	Location of profiles in a transect	14
3.5	Soil phase or management units of Mundragi-3 Microwatershed	19
5.1	Land Capability Classification map of Mundragi-3 Microwatershed	37
5.2	Soil Depth map of Mundragi-3 Microwatershed	38
5.3	Surface Soil Texture map of Mundragi-3 Microwatershed	39
5.4	Soil Gravelliness map of Mundragi-3 Microwatershed	40
5.5	Soil Available Water Capacity map of Mundragi-3 Microwatershed	41
5.6	Soil Slope map of Mundragi-3 Microwatershed	42
5.7	Soil Erosion map of Mundragi-3 Microwatershed	43
6.1	Soil Reaction (pH) map of Mundragi-3 Microwatershed	46
6.2	Electrical Conductivity (EC) map of Mundragi-3 Microwatershed	46
6.3	Soil Organic Carbon (OC) map of Mundragi-3 Microwatershed	47
6.4	Soil Available Phosphorus map of Mundragi-3 Microwatershed	48
6.5	Soil Available Potassium map of Mundragi-3 Microwatershed	49
6.6	Soil Available Sulphur map of Mundragi-3 Microwatershed	49
6.7	Soil Available Boron map of Mundragi-3 Microwatershed	50
6.8	Soil Available Iron map of Mundragi-3 Microwatershed	50
6.9	Soil Available Manganese map of Mundragi-3 Microwatershed	51
6.10	Soil Available Copper map of Mundragi-3 Microwatershed	51
6.11	Soil Available Zinc map of Mundragi-3 Microwatershed	52
7.1	Land suitability for Sorghum	54

7.2	Land suitability for Maize	55
7.3	Land suitability for Bajra	56
7.4	Land suitability for Groundnut	57
7.5	Land suitability for Sunflower	58
7.6	Land suitability for Redgram	59
7.7	Land suitability for Bengal gram	60
7.8	Land suitability for Cotton	61
7.9	Land suitability for Chilli	62
7.10	Land suitability for Tomato	63
7.11	Land suitability for Brinjal	64
7.12	Land suitability for Onion	65
7.13	Land suitability for Bhendi	66
7.14	Land suitable for Drumstick	67
7.15	Land suitability for Mango	68
7.16	Land suitability for Guava	69
7.17	Land suitability for Sapota	70
7.18	Land suitability for Pomegranate	71
7.19	Land suitability for Musambi	72
7.20	Land suitability for Lime	73
7.21	Land suitability for Amla	74
7.22	Land suitability for Cashew	75
7.23	Land suitability for Jackfruit	76
7.24	Land suitability for Jamun	77
7.25	Land suitability for Custard apple	78
7.26	Land suitability for Tamarind	79
7.27	Land suitability for Mulberry	80
7.28	Land suitability for Marigold	81
7.29	Land suitability for Chrysanthemum	82
7.30	Land Management Units map of Mundragi-3 Microwatershed	113
9.1	Soil and water conservation map of Mundragi-3 Microwatershed	128



## **EXECUTIVE SUMMARY**

*The land resource inventory of Mundragi-3 Microwatershed was conducted using village cadastral maps and IRS satellite imagery on 1:7920 scale. The false colour composites of IRS imagery were interpreted for physiography and the physiographic delineations were used as base for mapping soils. The soils were studied in several transects and a soil map was prepared with phases of soil series as mapping units. Random checks were made all over the area outside the transects to confirm and validate the soil map unit boundaries. The soil map shows the geographic distribution and extent, characteristics, classification, behavior and use potentials of the soils in the microwatershed.*

*The present study covers an area of 598 ha in Yadgir taluk & district, Karnataka. The climate is semiarid and categorized as drought-prone with an average annual rainfall of 866 mm, of which about 652 mm is received during south-west monsoon, 138 mm during north-east and the remaining 76 mm during the rest of the year. An area of 453 ha in the microwatershed is covered by soils, 25 ha by mining/industrial area, 74 ha by rock outcrops and about 46 ha by others (habitation and water bodies). The salient findings from the land resource inventory are summarized briefly below.*

- ❖ *The soils belong to 8 soil series and 10 soil phases (management units) and 4 land management units.*
- ❖ *The length of crop growing period is about 120-150 days starting from 1<sup>st</sup> week of June to 4<sup>th</sup> week of October.*
- ❖ *From the master soil map, several interpretative and thematic maps like land capability, soil depth, surface soil texture, soil gravelliness, available water capacity, soil slope and soil erosion were generated.*
- ❖ *Soil fertility status maps for macro and micronutrients were generated based on the surface soil samples collected at every 320 m grid interval.*
- ❖ *Land suitability for growing 29 major agricultural and horticultural crops was assessed and maps showing the degree of suitability along with constraints were generated.*
- ❖ *Entire area in the microwatershed is suitable for agriculture.*
- ❖ *About 5 per cent area of the microwatershed has soils that are moderately deep to very deep (75 - >150 cm) and 71 per cent soils are very shallow to moderately shallow (0-75 cm).*
- ❖ *About 52 per cent area in the microwatershed has loamy soils and 24 per cent clayey soils.*
- ❖ *About 53 per cent area is non gravelly (<15%) and 23 per cent is gravelly (15-35%).*

- ❖ *About 4 per cent area of the microwatershed is very high (>200 mm/m) in available water capacity, 3 per cent area low (51-100 mm/m) and 69 per cent area very low (<50 mm/m) in available water capacity.*
- ❖ *An area of about <1 per cent soils are nearly level (0-1%) and 75% area of microwatershed has very gently sloping (1-3% slope) lands.*
- ❖ *An area of about <1 per cent is slightly (e1) eroded, 75 per cent area is moderately (e2) eroded.*
- ❖ *An area of about 32 per cent soils are neutral (pH 6.5-7.3) in soil reaction and 43 per cent soils are slightly alkaline (pH 7.3-8.4).*
- ❖ *The Electrical Conductivity (EC) of the soils in the entire area of the microwatershed is dominantly <2 dsm<sup>-1</sup> indicating that the soils are non-saline.*
- ❖ *About 76 per cent medium (0.5-0.75%) in organic carbon content and <1 per cent high (>0.75).*
- ❖ *About 26 per cent area is low (<23kg/ha) and 49 per area is medium (23-57 kg/ha) in available phosphorus.*
- ❖ *About an area of 9 per cent is low (145 kg/ha) is low and 67 per cent medium (145-337 kg/ha) in available potassium.*
- ❖ *Available sulphur is low (<10 ppm) in an area of about 9 per cent and medium (10 -20 ppm) in 66 per cent of the microwatershed.*
- ❖ *Available boron is low (<0.5 ppm) in an entire area of the microwatershed.*
- ❖ *Available iron is sufficient (>4.5 ppm) in the whole area of the microwatershed.*
- ❖ *Available manganese and copper are sufficient in all the soils of the microwatershed.*
- ❖ *Available zinc is deficient (<0.6 ppm) in 53 per cent and sufficient (>0.6 ppm) in 23 per cent of the microwatershed.*
- ❖ *The land suitability for 29 major crops grown in the microwatershed were assessed and the areas that are highly suitable (S1) and moderately suitable (S2) are given below. It is however to be noted that a given soil may be suitable for various crops but what specific crop to be grown may be decided by the farmer looking to his capacity to invest on various inputs, marketing infrastructure, market price and finally the demand and supply position.*

**Land suitability for various crops in the Microwatershed**

<b>Crop</b>	<b>Suitability Area in ha (%)</b>		<b>Crop</b>	<b>Suitability Area in ha (%)</b>	
	<b>Highly suitable (S1)</b>	<b>Moderately suitable (S2)</b>		<b>Highly suitable (S1)</b>	<b>Moderately suitable (S2)</b>
<i>Sorghum</i>	3(<1)	40(7)	<i>Guava</i>	-	3(<1)
<i>Maize</i>	-	43(7)	<i>Sapota</i>	-	3(<1)
<i>Bajra</i>	-	43(7)	<i>Pomegranate</i>	-	28(5)
<i>Groundnut</i>	-	3 (<1)	<i>Musambi</i>	22(4)	6(1)
<i>Sunflower</i>	-	28(5)	<i>Lime</i>	22(4)	6(1)
<i>Redgram</i>	-	27(5)	<i>Amla</i>	3 (<1)	40(7)
<i>Bengal gram</i>	24(4)	15(3)	<i>Cashew</i>	-	-
<i>Cotton</i>	24(4)	15(3)	<i>Jackfruit</i>	-	3(<1)
<i>Chilli</i>	-	21(3)	<i>Jamun</i>	-	25(4)
<i>Tomato</i>	-	18(3)	<i>Custard apple</i>	27(5)	15(3)
<i>Brinjal</i>	3(<1)	40(7)	<i>Tamarind</i>	-	25(4)
<i>Onion</i>	3(<1)	18(3)	<i>Mulberry</i>	-	3(<1)
<i>Bhendi</i>	5(<1)	37(7)	<i>Marigold</i>	-	42(7)
<i>Drumstick</i>	-	27(5)	<i>Chrysanthemum</i>	-	42(7)
<i>Mango</i>	-	-			

- ❖ *Apart from the individual crop suitability, a proposed crop plan has been prepared for the identified LUCs by considering only the highly and moderately suitable lands for different crops and cropping systems with food, fodder, fibre and horticulture crops.*
- ❖ *Maintaining soil-health is vital to crop production and conserve soil and land resource base for maintaining ecological balance and to mitigate climate change. For this, several ameliorative measures have been suggested to these problematic soils like saline/alkali, highly eroded, sandy soils etc.,*
- ❖ *Soil and water conservation treatment plan has been prepared that would help in identifying the sites to be treated and also the type of structures required.*
- ❖ *As part of the greening programme, several tree species have been suggested to be planted in marginal and sub marginal lands, field bunds and also in the hillocks, mounds and ridges. This would help in not only supplementing the farm income but also provide fodder and fuel and generate lot of biomass which would help in maintaining an ecological balance and also contribute to mitigating the climate change.*



## **INTRODUCTION**

Land is a scarce resource and basic unit for any material production. It can support the needs of the growing population, provided they use the land in a rational and judicious manner. But what is happening in many areas of the state is a cause for concern to everyone involved in the management of land resources at the grassroots level. The area available for agriculture is about 51 per cent of the total geographical area and more than 60 per cent of the people are still dependant on agriculture for their livelihood. The limited land area is under severe stress and strain due to increasing population pressure and competing demands of various land uses. Due to this, every year there is significant diversion of farm lands and water resources for non-agricultural purposes. Apart from this, due to lack of interest in farmers for farming, large tracts of cultivable lands are turning into fallows in many areas and this trend is continuing at an alarming rate.

Further, land degradation has emerged as a serious problem which has already affected about 38 lakh ha of cultivated area in the state. Soil erosion alone has degraded about 35 lakh ha. Almost all the uncultivated areas are facing various degrees of degradation, particularly soil erosion. Salinity and alkalinity has emerged as a major problem in more than 3.5 lakh ha in the irrigated areas of the state. Nutrient depletion and declining factor productivity is common in both rainfed and irrigated areas. The degradation is continuing at an alarming rate and there appears to be no systematic effort among the stakeholders to contain this process. In recent times, an aberration of weather due to climate change phenomenon has added another dimension leading to unpredictable situations to be tackled by the farmers.

In this critical juncture, the challenge before us is not only to increase the productivity per unit area which is steadily declining and showing a fatigue syndrome, but also to prevent or at least reduce the severity of degradation. If the situation is not reversed at the earliest, then the sustainability of the already fragile crop production system and the overall ecosystem will be badly affected in the state. The continued neglect and unscientific use of the resources for a long time has led to the situation observed at present in the state. It is a known fact and established beyond doubt by many studies in the past that the cause for all kinds of degradation is the neglect and irrational use of the land resources. Hence, there is an urgent need to generate a detailed site-specific farm level database on various land resources for all the villages/watersheds in a time bound manner that would help to protect the valuable soil and land resources and also to stabilize the farm production.

Therefore, the land resource inventory required for farm level planning is the one which investigates not only the surface but also consider the other parameters which are critical for productivity *viz.*, soils, climate, water, minerals and rocks, topography, geology, hydrology, vegetation, crops, land use pattern, animal population, socio-

economic conditions, infrastructure, marketing facilities and various schemes and developmental works of the government etc. From the data collected at farm level, the specific problems and potentials of the area can be identified and highlighted, conservation measures required for the area can be planned on a scientific footing, suitability of the area for various uses can be worked out and finally viable and sustainable land use options suitable for each and every land holding can be prescribed.

The Land Resource Inventory is basically done for identifying the potential and problem areas, developing sustainable land use plans, estimation of surface run off and water harvesting potential, preparation of soil and water conservation plans, land degradation/desertification etc. The Bureau is presently engaged in developing an LRI methodology using high resolution satellite remote sensing data and Digital Elevation Model (DEM) data to prepare Landscape Ecological Units (LEU) map representing agro-ecosystem as a whole. The LEU is preferred over landform as the base map for LRI. LEU is the assemblage of landform, slope and land use. An attempt has already been made to upscale the soil resource information from 1:250000 and 1:50000 scale to the LEU map in Goa and other states.

The land resource inventory aims to provide site-specific database for Mundragi-3 microwatershed in Yadgir Taluk & District, Karnataka State for the Karnataka Watershed Development Department. The database was generated by using cadastral map of the village as a base along with high resolution IRS LISS IV and Cartosat-1 merged satellite imagery. Later, an attempt will be made to uplink this LRI data generated at 1:7920 scale under Sujala-III Project to the proposed Landscape Ecological Units (LEUs) map.

The study was organized and executed by the ICAR- National Bureau of Soil Survey and Land Use Planning, Regional Centre, Bangalore under Generation of Land Resource Inventory Data Base Component-1 of the Sujala-III Project funded by the World Bank.

## GEOGRAPHICAL SETTING

### 2.1 Location and Extent

The Mundragi-3 microwatershed is located in the northern part of Karnataka in Yadgir Taluk & District, Karnataka State (Fig.2.1). It comprises parts of Yadhagiri.b, Mundaragi & Varkanahalli Villages. It lies between  $16^{\circ} 45'$  -  $16^{\circ} 46'$  North latitudes and  $77^{\circ} 9'$  -  $77^{\circ} 12'$  East longitudes covering an area of about 598.36 ha. It is about 5 km northwest of Yadgir town and is surrounded by Mundaragi on the north and east, Yadhagiri.B on the south and southwest and Varkanahalli village on the southern side.




Fig.2.1 Location map of Mundragi-3 Microwatershed

### 2.2 Geology

Major rock formation observed in the microwatershed are granite gneiss (Fig.2.2). Granite gneisses are essentially pink to gray and are coarse to medium grained. They consist primarily of quartz, feldspar, biotite and hornblende. The gray granite gneisses are highly weathered, fractured and fissured up to a depth of about 10 m. Dolerite dykes and

quartz veins are common with variable width and found to occur in Mundragi-3 microwatershed.



Fig.2.2 Granite and granite gneiss rocks

### **2.3 Physiography**

Physiographically, the area has been identified as granite and gneiss landscape based on geology. The area has been further subdivided into five landforms, *viz*.; mounds/ridges, summits, side slopes and very gently sloping uplands, plains and valleys based on slope and its relief features. The elevation ranges from 378-406 m above MSL. The mounds and ridges are mostly covered by rock outcrops.

### **2.4 Drainage**

The area is drained by several parallel streams like Bori, Amerja and Kanga which finally join the river Bhima along its course. Though, they are not perennial, during rainy season they carry large quantities of rain water. The microwatershed has only few small tanks which are not capable of storing the water that flows during the rainy season. Due to this, the ground water recharge is very much affected. This is reflected in the failure of many bore wells in the villages. If the available rain water is properly harnessed by constructing new tanks and recharge structures at appropriate places in the villages, then the drinking and irrigation needs of the area can be easily met. The drainage network is parallel to sub parallel and dendritic.

### **2.5 Climate**


The Yadgir district lies in the northern plains of Karnataka and falls under semiarid tract of the state and is categorized as drought-prone with total annual rainfall of 866 mm (Table 2.1). Of the total rainfall, maximum of 652 mm is received during the south-west monsoon period from June to September, the north-east monsoon from October to early December contributes about 138 mm and the remaining 76 mm during



the rest of the year. The summer season starts during the middle of February and continues up to the first week of June. The period from December to the middle of February is the coldest season. December is the coldest month with mean daily maximum and minimum temperatures being 29.5<sup>0</sup>C and 10<sup>0</sup>C respectively. During peak summer, temperature shoots up to 45<sup>0</sup>C. Relative humidity varies from 26% in summer to 62% in winter. Rainfall distribution is shown in Figure 2.3. The average Potential Evapo-Transpiration (PET) is 141 mm and varies from a low of 81 mm in December to 199 mm in the month of May. The PET is always higher than precipitation in all the months except end of June to end of September. Generally, the Length of crop Growing Period (LGP) is 120-150 days and starts from 1<sup>st</sup> week of June to 4<sup>th</sup> week of October.

**Table 2.1 Mean Monthly Rainfall, PET, 1/2 PET at Yadgir Taluk, Yadgir District**

Sl. No.	Months	Rainfall	PET	1/2 PET
1	January	4.30	86.0	43.0
2	February	2.30	125.5	62.7
3	March	15.10	166.0	83.0
4	April	18.50	179.8	89.9
5	May	36.0	198.8	97.9
6	June	118.0	175.1	87.5
7	July	171.80	156.3	78.1
8	August	182.9	150.3	75.1
9	September	179.7	142.0	71.0
10	October	105.3	138.5	69.2
11	November	26.4	97.60	48.6
12	December	6.0	80.90	40.4
<b>Total</b>		<b>866.3</b>		



**Fig 2.3 Rainfall distribution in Yadgir Taluk, Yadgir District**

## 2.6 Natural Vegetation

The natural vegetation is sparse comprising few tree species, shrubs and herbs. The mounds, ridges and boulders occupy very sizeable area which is under thin to moderately thick forest vegetation. Still, there are some remnants of the past forest cover which can be seen in patches in some ridges and hillocks in the microwatershed (Fig 2.4).

Apart from the continuing deforestation, the presence of large population of goats, sheep and other cattle in the microwatershed is causing vegetative degradation of whatever little vegetation left in the area. The uncontrolled grazing has left no time for the regeneration of the vegetative cover. This leads to the accelerated rate of erosion on the hill slopes resulting in the formation of deep gullies in the foot slopes that eventually result in the heavy siltation of tanks and reservoirs in the microwatershed.




Fig 2.4 Natural vegetation of Mundragi-3 Microwatershed

## 2.7 Land Utilization


About 72 per cent area (Table 2.2) in Yadgir district is cultivated at present. An area of about 2 per cent is permanently under pasture, 20 per cent under current fallows and 6 per cent under non-agricultural land and 5 per cent under currently barren. Forests occupy an area of about 7 per cent and the tree cover is in a very poor state. Most of the mounds, ridges and bouldery areas have very poor vegetative cover. Major crops grown in the area are paddy, cotton, groundnut and red gram. While carrying out land resource inventory, the land use/land cover particulars are collected from all the survey numbers and a current land use map of the microwatershed is prepared. The current land use map prepared shows the arable and non-arable lands, other land uses and different types of crops grown in the area. The current land use map of Mundragi-3 microwatershed is presented in Fig.2.5. The different crops and cropping systems adopted in the microwatershed is presented in the Figures 2.6. Map showing the location of wells in the Mundragi-3 microwatershed is given in fig 2.7.

**Table 2.2 Land Utilization in Yadgir District**

Sl. No.	Agricultural land use	Area ( ha)	Per cent
1	Total geographical area	516088	-
2	Total cultivated area	373617	72.4
3	Area sown more than once	74081	14.3
4	Cropping intensity	-	119.8
5	Trees and grooves	737	0.14
6	Forest	33773	6.54
7	Cultivable wasteland	2385	0.46
8	Permanent Pasture land	11755	2.28
9	Barren land	27954	5.41
10	Non- Agriculture land	29623	5.73
11	Current Fallows	105212	20.4



**Fig.2.5 Current Land Use map of Mundragi-3 Microwatershed**



Groundnut



Cotton



Redgram



Paddy

Fig. 2.6 . Different Crops and Cropping Systems in Mundragi-3 Microwatershed




Fig 2.7 Location of wells - Mundragi-3 Microwatershed



## SURVEY METHODOLOGY

The purpose of land resource inventory is to delineate similar areas (soil series and phases), which respond or expected to respond similarly to a given level of management. This was achieved in Mundragi-3 microwatershed by the detailed study of all the soil characteristics (depth, texture, colour, structure, consistence, coarse fragments, porosity, soil reaction, soil horizons etc.) and site characteristics (slope of the land, erosion, drainage, occurrence of rock fragments etc.) followed by grouping of similar areas based on soil-site characteristics into homogeneous (management units) units, and showing the area extent and their geographic distribution on the microwatershed cadastral map. The detailed survey at 1:7920 scale was carried out in an area of 598 ha. The methodology followed for carrying out land resource inventory was as per the guidelines given in Soil Survey Manual (IARI, 1971; Soil Survey Staff, 2006; Natarajan *et al.*, 2015) which is briefly described below.

### 3.1 Base Maps

The detailed survey of the land resources occurring in the microwatershed was carried out by using digitized cadastral map and IRS satellite imagery as base supplied by KRSRSAC. The cadastral map shows field boundaries with their survey numbers, location of tanks, streams and other permanent features of the area (Fig. 3.1). Apart from the cadastral map, remote sensing data products from Cartosat-1 and LISS IV merged at the scale of 1:7920 were used in conjunction with the cadastral map to identify the landscapes, landforms and other surface features. The imagery helped in the identification and delineation of boundaries between hills, uplands and lowlands, water bodies, forest and vegetated areas, roads, habitations and other cultural features of the area (Fig. 3.2). The cadastral map was overlaid on the satellite imagery (Fig. 3.3) that helps to identify the parcel boundaries and other permanent features. Apart from cadastral maps and images, toposheets of the area (1:50,000 scale) were also used for initial traversing, identification of geology and landforms, drainage features, present land use and also for selection of transects in the microwatershed.

### 3.2 Image Interpretation for Physiography

False Colour Composites (FCCs) of Cartosat-I and LISS-IV merged satellite data covering microwatershed area was visually interpreted using image interpretation elements and all the available collateral data with local knowledge. The delineated physiographic boundaries were transferred on to a cadastral map overlaid on satellite imagery. Physiographically, the area has been identified as granite and granite gneiss landscape. It was divided into five landforms, *viz*; ridges and mounds, gently and very gently sloping uplands and lowlands based on slope and image characteristics. They were

further subdivided into physiographic/image interpretation units based on image characteristics. The image interpretation legend for physiography is given below.

### Image Interpretation Legend for Physiography

#### G- Granite Gneiss Landscape

- G1 Hills/ Ridges/ Mounds
  - G11 Summits
  - G12 Side slopes
    - G121 Side slopes with dark grey tones
- G2 Uplands
  - G21 Summits
  - G22 Gently sloping uplands
    - G221 Gently sloping uplands, yellowish green (eroded)
    - G222 Gently sloping uplands, yellowish white (severely eroded)
  - G23 Very gently sloping uplands
    - G231 Very gently sloping uplands, yellowish green
    - G232 Very gently sloping uplands, medium green and pink
    - G233 Very gently sloping uplands, pink and green (scrub land)
    - G234 Very gently sloping uplands, medium greenish grey
    - G235 Very gently sloping uplands, yellowish white (eroded)
    - G236 Very gently sloping uplands, dark green
    - G237 Very gently sloping uplands, medium pink (coconut garden)
    - G238 Very gently sloping uplands, pink and bluish white (eroded)
  - G24 Valleys/ lowlands
    - G241 Valleys, pink tones
    - G242 Valleys gray mixed with pink tones




Fig 3.1 Scanned and Digitized Cadastral map of Mundragi-3 Microwatershed






Fig.3.2 Satellite Image of Mundragi-3 Microwatershed




Fig.3.3 Cadastral map overlaid on IRS PAN+LISS IV merged imagery of Mundragi-3 Microwatershed

### 3.3 Field Investigation

The field boundaries and survey numbers given on the cadastral sheet were located on the ground by following permanent features like roads, cart tracks, *nallas*, streams, tanks etc., and wherever changes were noticed, they were incorporated on the microwatershed cadastral map. Preliminary traverse of the microwatershed was carried out with the help of cadastral map, imagery and toposheets. While traversing, landforms and physiographic units identified were checked and preliminary soil legend was prepared by studying soils at few selected places. Then, intensive traversing of each physiographic unit like hills, ridges, uplands and valleys was carried out. Based on the variability observed on the surface, transects (Fig. 3.4) were selected across the slope covering all the landform units in the microwatershed (Natarajan and Dipak Sarkar, 2010).




Fig: 3.4. Location of profiles in a transect

In the selected transect, soil profiles were located (Fig. 3.4) at closely spaced intervals to take care of any change in the land features like break in slope, erosion, gravel, stones etc. In the selected sites, profiles (vertical cut showing the soil layers from surface to the rock) were opened up to 200 cm or to the depth limited by rock or hard substratum and studied in detail for all their morphological and physical characteristics. The soil and site characteristics were recorded for all profile sites on a standard proforma as per the guidelines given in USDA Soil Survey Manual (Soil Survey Staff, 2012). Apart from the transect study, profiles were also studied at random, almost like in a grid pattern, outside the transect areas.

Based on the soil characteristics, the soils were grouped into different soil series. Soil series is the most homogeneous unit having similar horizons and properties and behaves similarly for a given level of management. Soil depth, texture, colour, kind of horizon and horizon sequence, calcareousness, amount and nature of gravel present, nature of substratum etc, were used as the major differentiating characteristics for identifying soil series occurring in the area. The differentiating characteristics used for

identifying the soil series are given in Table 3.1. Based on the above characteristics, 08 soil series were identified in the Mundragi-3 microwatershed.

**Table 3.1 Differentiating Characteristics used for identifying Soil Series**  
(Characteristics are of Series Control Section)

Soils of Granite gneiss Landscape							
Sl. no	Soil Series	Depth (cm)	Colour (moist)	Texture	Gravel (%)	Horizon sequence	Calcareous-ness
1	BDL (Badiyala)	25-50	7.5YR 2.5/3,2.5/2,3/3 10YR 3/4,4/3	sl	-	Ap-Bw	e
2	SBR (Sambra)	50-75	10YR 7/1 7.5YR 7/4	ls-s	-	Ap-AC	-
3	JNK (Jinkera)	50-75	10YR5/3,3/2 7.5YR3/4	scl	-	Ap-Bw	e
4	HSL (Hosalli)	75-100	10YR 5/4,4/4,4/6	sc	-	Ap-Bw	e
5	NGP (Nagalapur)	100-150	10YR3/2,3/1,2/1	c	-	Ap-Bss	es
6	BDP (Baddeppalli)	<25	7.5YR 3/2,3/4 5YR 3/4	scl	-	Ap-Ac	es
7	HTK (Hattikuni)	25-50	10YR4/6,4/4 7.5YR3/4,3/3	sl	10-25	Ap-Ac	-
8	TMK (Thumakur)	>150	10YR 3/1,3/2,3/3,4/3	c	-	Ap-Bw	e

### 3.4 Soil Mapping

The area under each soil series was further separated into soil phases and their boundaries delineated on the cadastral map based on the variations observed in the texture of the surface soil, slope, erosion, presence of gravel, stoniness etc. A soil phase is a subdivision of soil series based mostly on surface features that affect its use and management. The soil mapping units are shown on the map (Fig.3.5) in the form of symbols. During the survey many profile pits, few minipits and a few auger bores representing different landforms occurring in the microwatershed were studied. In addition to the profile study, spot observations in the form of minipits, road cuts, terrace cuts etc., were studied to validate the soil boundaries on the soil map.

The soil map shows the geographic distribution of 10 mapping units representing 8 soil series occurring in the microwatershed. The soil map unit (soil legend) description is presented in Table 3.2. The soil phase map (management units) shows the distribution of 10 soil phases mapped in the microwatershed. Each mapping unit (soil phase) delineated on the map has similar soil and site characteristics. In other words, all the farms or survey numbers included in one phase will have similar management needs and have to be treated accordingly.

### 3.5 Land Management Units

The 10 soil phases identified and mapped in the microwatershed were grouped into 4 Land Management Units (LMU's) for the purpose of preparing a Proposed Crop Plan for sustained development of the microwatershed. The database (soil phases) generated under LRI was utilized for identifying Land Management Units (LMU's) based on the management needs. One or more than one soil site characteristic having influence on the management have been chosen for identification and delineation of LMUs. For Mundragi-3 microwatershed, five soil and site characteristics, namely soil depth, soil texture, slope, erosion and gravel content have been considered for defining LMUs. The Land Management Units are expected to behave similarly for a given level of management.

### 3.6 Laboratory Characterization

Soil samples were collected from representative master profiles for laboratory characterization by following the methods outlined in the Laboratory Manual (Sarma *et al*, 1987). Surface soil samples collected from farmer's fields (58 samples) for fertility status (major and micronutrients) at 320 m grid interval in the year 2018 were analyzed in the laboratory (Katyal and Rattan, 2003). By linking the soil fertility data to the survey numbers through GIS, soil fertility maps were generated by using Kriging method for the microwatershed.

**Table 3.2 Soil map unit description of Mundragi-3 Microwatershed**

*Soil map unit No.	Soil Series	Soil Phase	Mapping Unit Description	Area in ha (%)
<b>Soils of Granite and Granite Gneiss Landscape</b>				
	BDL		Badiyala soils are shallow (25-50 cm), well drained, have dark brown to very dark brown and dark yellowish brown, slightly calcareous, sandy loam soils occurring on very gently to gently sloping uplands under cultivation	<b>245(40.9)</b>
5		BDLiB2	Sandy clay surface, slope 1-3, moderate erosion	109(18.26)
162		BDLhB2g1	Sandy clay loam surface, slope 1-3%, moderate erosion, gravelly (15-35%)	136(22.71)
	SBR		Sambara soils are moderately shallow (50-75 cm), somewhat excessively drained, have light gray to pink, loamy sand soils occurring on very gently to gently sloping uplands under cultivation	<b>78 (13.02)</b>
11		SBRcB2	Sandy loam surface, slope 1-3%, moderate erosion	78 (13.02)
	JNK		Jinkera soils are moderately shallow (50-75 cm), well	<b>15(2.56)</b>

		drained, have dark brown to very dark grayish brown, slightly calcareous, sandy clay loam soils occurring on very gently sloping uplands under cultivation		
20		JNKcB2	Sandy loam surface, slope 1-3%, moderate erosion	5 (0.81)
22		JNKiB2	Sandy clay surface, slope 1-3, moderate erosion	10 (1.75)
	HSL	Hosalli soils are moderately deep (75-100 cm), moderately well drained, have yellowish brown to dark yellowish brown, slightly calcareous, sandy clay soils occurring on very gently sloping uplands under cultivation		<b>3 (0.49)</b>
32		HSLcB2	Sandy loam surface, slope 1-3%, moderate erosion	3 (0.49)
	NGP	Nagalapur soils are deep (100-150 cm), moderately well drained, have very dark gray to very dark grayish brown, black calcareous, cracking clay soils occurring on very gently sloping uplands under cultivation		<b>22 (3.65)</b>
49		NGPmB2	Clay surface, slope 1-3%, moderate erosion	22 (3.65)
	BDP	Baddeppalli soils are very shallow (<25 cm), well drained, have dark brown to dark reddish brown, calcareous, sandy clay loam soils occurring on very gently sloping uplands under cultivation		<b>9 (1.54)</b>
118		BDPcB2	Sandy loam surface, slope 1-3%, moderate erosion	9 (1.54)
	HTK	Hattikuni soils are shallow (25-50 cm), well drained, have dark yellowish brown sandy loam soils occurring on very gently sloping uplands under cultivation		<b>78 (13.06)</b>
165		HTKcB2	Sandy loam surface, slope 1-3%, moderate erosion	78 (13.06)
	TMK	Thumakur soils are very deep (>150 cm), moderately well drained, have brown to very dark grayish brown, slightly calcareous, sodic, clay black soils occurring on nearly level to very gently sloping lowlands under cultivation		<b>3 (0.43)</b>
103		TMKhA1	Sandy clay loam surface, slope 0-1%, slight erosion	3 (0.43)
994		Mining/Industrial area		<b>25 (4.24)</b>
999	Rock outcrops	Rock lands, both massive and boulder with little or no soil		<b>74 (12.44)</b>
1000	Others	Habitation and Water body		<b>46 (7.61)</b>






Fig 3.5 Soil Phase or Management Units - Mundragi-3 Microwatershed





## THE SOILS

Detailed information pertaining to the nature, extent and their distribution of different kinds of soils occurring in Mundragi-3 microwatershed is provided in this chapter. The microwatershed area has been identified as granite and gneiss landscape based on geology. In all, 8 soil series are identified. Soil formation is the result of the combined effect of environmental and terrain factors that are reflected in soil morphology. In the granite gneiss landscape, it is by parent material, relief and climate.

A brief description of each of the 8 soil series identified followed by 10 soil phases (management units) mapped under each series are furnished below. The physical and chemical characteristics of soil series identified in Mundragi-3 microwatershed are given in Table 4.1 along with soil classification. The soils in any one map unit differ from place to place in their depth, texture, slope, gravelliness, erosion or any other site characteristic that affect management. The soil phase map can be used for identifying the suitability of areas for growing specific crops or for other alternative uses and also for deciding the type of conservation structures needed. The detailed information on soil and site-characteristics like soil depth, surface soil texture, slope, erosion, gravelliness, AWC, LCC etc, with respect to each of the soil phase identified is given village/survey number wise for the microwatershed in Appendix-I.

### 4.1 Soils of granite gneiss landscape

In this landscape, 8 soil series are identified and mapped. Of these, BDL series occupies maximum area of 245 ha (41%) followed by HTK 78 ha (13%), SBR 78 ha (13%) and NGP 22 ha (4%). The other series occupy minor area in the microwatershed. Brief description of each series identified and number of soil phases mapped is given below.

**4.1.1 Badiyala (BDL) Series:** Badiyala soils are shallow (25-50 cm), well drained, have very dark brown, dark yellow brown and dark brown, slightly calcareous, sandy loam soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation. The Badiyala series has been classified as a member of the coarse-loamy, mixed, isohyperthermic family of Fluventic Haplustepts.

The thickness of the solum ranges from 28 to 50 cm. The thickness of A horizon ranges from 4 to 12 cm. Its colour is in 10YR hue with value 3 to 4 and chroma 3 to 4. The texture is loamy sand, sandy clay loam and sandy clay. The thickness of B horizon ranges from 27 to 45 cm. Its colour is in 10 YR and 7.5 YR hue with value 2 to 4 and chroma 3 to 4. Its texture is sandy loam to sandy clay loam and is slightly calcareous. The available water capacity is very low (<50mm/m). Two phases were identified and mapped.



Landscape and Soil Profile characteristics of Badiyala (BDL) Series

**4.1.2 Sambra (SBR) Series:** Sambra soils are moderately shallow (50-75 cm), somewhat excessively drained, have light grey to reddish yellow, loamy sand soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation. The Sambra series has been classified as a member of the mixed, isohyperthermic family of Typic Ustipsamments

The thickness of the soil ranges from 52-75 cm. Thickness of A horizon ranges from 8 to 23 cm. Its colour is in hue 10 YR and 7.5 YR with value 3 and chroma 1 to 4. The texture varies from loamy sand to sandy loam. The thickness of subsurface horizons ranges from 41 to 66 cm. Its colour is in 10 YR and 7.5 YR hue with value 3 to 5 and chroma 1 to 4. The texture is loamy sand. The available water capacity is very low (<50 mm/m). Only one phase was identified and mapped.



Landscape and Soil Profile characteristics of Sambra (SBR) Series

**4.1.3 Jinkera (JNK) Series:** Jinkera soils are moderately shallow (50-75 cm), well drained, have very dark gray to very dark grayish brown and dark brown, slightly calcareous, sandy clay loam soils. They are developed from weathered granite gneiss and occur on very gently sloping uplands under cultivation. The Jinkera series has been classified as a member of the fine-loamy, mixed, isohyperthermic family of Typic Haplustepts.

The thickness of the solum ranges from 51-75 cm. Thickness of A horizon ranges from 6 to 11 cm. Its colour is in hue 10 YR and 7.5 YR with value and chroma of 3 to 4. The texture varies from sandy loam to sandy clay. The thickness of B horizon ranges from 53 to 66 cm. Its colour is in 10 YR and 7.5 YR hue with value and chroma of 2 to 4. The texture varies from sandy clay loam to sandy clay and is slightly calcareous. The available water capacity is low (51-100 mm/m). Two phases were identified and mapped.



Landscape and Soil Profile characteristics of Jinkera (JNK) Series

**4.1.4 Hosalli (HSL) Series:** Hosalli soils are moderately deep (75-100 cm), moderately well drained, have dark yellowish brown to yellowish brown, slightly calcareous, sandy clay soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation. The Hosalli series has been classified as a member of the fine, mixed, isohyperthermic family of Typic Haplustepts.

The thickness of the solum ranges from 76 to 100 cm. The thickness of A horizon ranges from 6 to 15 cm. Its colour is in hue 10 YR and 7.5 YR with value 3 to 5 and chroma 2 to 4. Its texture varies from loamy sand to sandy loam and sandy clay loam. The thickness of B horizon ranges from 62 to 93 cm. Its colour is in hue 10 YR with value 3 to 4 and chroma 2 to 4. Its texture varies from sandy clay loam to sandy clay and clay and is slightly calcareous. The available water capacity is medium (101-150 mm/m). Only one phase was identified and mapped.



Landscape and Soil Profile characteristics of Hosalli (HSL) Series

**4.1.5 Naglapur (NGP) Series:** Naglapur soils are deep (100-150 cm), moderately well drained, have black to very dark grayish brown, calcareous cracking clay soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation. The Naglapur series has been classified as a member of the very fine, smectitic (calcareous), isohyperthermic family of Typic Haplusterts.

The thickness of the solum ranges from 110 to 150 cm. The thickness of A horizon ranges from 6 to 25 cm. Its colour is in 10 YR hue with value 3 to 5 and chroma 1 to 3. The texture varies from sandy loam to sandy clay and clay. The thickness of B horizon ranges from 110 to 141 cm. Its colour is in 10 YR hue with value 2 to 3 and chroma 1 to 2. Texture is clay and is calcareous. The available water capacity is very high (>200 mm/m). Only one phase was identified and mapped.



Landscape and Soil Profile characteristics of Naglapur (NGP) Series

**4.1.6 Baddeppalli (BDP) Series:** Baddeppalli soils are very shallow (<25cm), well drained, have dark brown to dark reddish brown, calcareous, sandy clay loam soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation. The Baddeppalli series has been classified as a member of the loamy, mixed (calcareous), isohyperthermic family of Lithic Ustorthents.

The thickness of the soil is less than 25 cm. Its colour is in 7.5 YR and 5 YR hue with value 3 and chroma 2 to 4. The texture varies from sandy clay loam to sandy clay and is calcareous. The available water capacity is very low (<50 mm/m). Only one phase was identified and mapped.



Landscape and Soil Profile characteristics of Baddeppalli (BDP) Series

**4.1.7 Hattikuni soils** are shallow (25-50 cm), well drained, have dark brown to dark yellowish brown sandy loam soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation. The Hattikuni series has been classified as a member of the mixed, isohyperthermic family of Lithic Ustipsamments.

The thickness of the soil ranges from 36 to 50 cm. The thickness of A horizon ranges from 8 to 12 cm. Its colour is in 10YR and 7.5 YR hue with value 3 to 4 and chroma 4 to 6. The texture varies from loamy sand to sandy loam. The thickness of subsurface horizon ranges from 28 to 42 cm. Its colour is in 10 YR and 7.5 YR hue with value 3 to 4 and chroma 4 to 6. Its texture varies from loamy sand to sand and sandy loam. The available water capacity is very low (<50 mm/m). Only one phase was identified and mapped.



Landscape and Soil Profile characteristics of Hattikuni (HTK) Series

**4.1.8 Thumakur (TMK) Series:** Thumakur soils are very deep (>150 cm), moderately well drained, have very dark gray to dark brown, slightly calcareous, sodic clay soils. They are developed from weathered granite gneiss and occur on nearly level to very gently sloping low lands under cultivation. The Thumakur series has been classified as a member of the fine, mixed, isohyperthermic family of Typic Haplustepts.

The thickness of the solum ranges from 150-200 cm. The thickness of A horizon ranges from 7 to 14 cm. Its colour is in 10 YR hue with value 3 to 5 and chroma 1 to 3. Texture varies from sandy loam to sandy clay and clay. The thickness of B horizon is >150 cm. Its colour is in 10 YR hue with value 3 to 5 and chroma 1 to 3. Texture varies from sandy clay to clay and is slightly calcareous, sodic soils. The available water capacity is very high (>200 mm/m). Only one phase was identified and mapped.



Landscape and Soil Profile characteristics of Thumakur (TMK) Series

**Table: 4.1 Physical and Chemical Characteristics of Soil Series identified in Mundragi-3 microwatershed**

**Soil Series:** Badiyala (BDL) **Pedon:** R-5

**Location:** 16<sup>0</sup>37'10.0"N 77<sup>0</sup>20'21.5", Gudalagunta village, Balichakra hobli, Yadgir taluk and district

**Analysis at:** NBSS&LUP, Regional Centre, Bengaluru **Classification:** Coarse-loamy, mixed, isohyperthermic, Fluventic Haplustepts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-12	Ap	87.13	7.04	5.83	10.03	24.32	23.61	23.51	5.67	<15	ls	6.27	2.44
12-28	Bw1	64.63	13.30	22.07	6.74	13.07	22.30	17.01	5.50	<15	scl	16.34	7.83
28-52	BC	73.11	12.02	14.87	3.93	16.03	26.89	18.41	7.86	<15	sl	12.94	5.47

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO <sub>3</sub>	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP
	Water	CaCl <sub>2</sub>	M KCl				Ca	Mg	K	Na	Total				
				dS m <sup>-1</sup>	%	%	cmol kg <sup>-1</sup>					%	%		
0-12	6.20	-	-	0.074	1.00	0.00	2.80	0.98	0.14	0.01	3.92	4.20	0.72	93	0.20
12-28	9.04	-	-	0.253	0.80	3.20	-	-	0.16	0.69	-	16.90	0.77	100	4.09
28-52	9.41	-	-	0.364	1.10	3.60	-	-	0.16	1.39	-	11.10	0.75	100	12.52

*Contd...*

**Soil Series:** Sambara (SBR) **Pedon:** R-10

**Location:** 16<sup>0</sup>42'04.5"N 77<sup>0</sup>14'35.3"E, Jinatera village, Balichakra hobli, Yadgir taluk and district

**Analysis at:** NBSS&LUP, Regional Centre, Bengaluru **Classification:** Mixed, isohyperthermic Typic Ustipsamments

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-9	Ap	81.90	8.22	9.88	23.76	14.05	23.76	10.62	9.71	-	ls	9.45	2.69
9-17	C1	84.08	6.59	9.33	21.30	20.69	17.65	17.65	6.80	-	ls	7.84	2.65
17-60	C2	86.86	6.17	6.98	11.53	21.54	25.08	23.46	5.26	-	ls	5.48	2.62
60-78	C3	87.27	6.92	5.81	15.05	20.91	26.36	19.29	5.66	-	ls	5.19	2.81

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO <sub>3</sub>	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP			
	Water	CaCl <sub>2</sub>	M KCl				dS m <sup>-1</sup>	%	%	Ca	Mg					K	Na	Total
										cmol kg <sup>-1</sup>								
0-9	8.24	-	-	0.145	0.61	0.91	-	-	0.12	0.09	-	7.50	0.76	100	1.15			
9-17	8.21	-	-	0.068	0.57	0.39	-	-	0.06	0.12	-	6.70	0.72	100	1.82			
17-60	8.47	-	-	0.080	0.38	0.48	-	-	0.03	0.17	-	2.70	0.39	100	6.34			
60-78	8.50	-	-	0.081	0.30	0.52	-	-	0.03	0.17	-	2.70	0.46	100	6.43			

Contd...



**Soil Series:** Jinkera (JNK) **Pedon:** R-1

**Location:** 16°45'13.5"N 77°10'59.8"E, Varkanahalli village, Yadgir hobli, Yadgir taluk and district

**Analysis at:** NBSS&LUP, Regional Centre, Bengaluru **Classification:** Fine-loamy, mixed, isohyperthermic Typic Haplustepts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-15	Ap	66.84	13.62	19.54	12.15	21.22	11.23	12.56	9.68	10	sl	14.42	7.70
15-38	Bw1	59.08	12.11	28.81	12.53	12.42	17.85	8.77	7.52	20	scl	18.21	12.23
38-50	Bw2	68.21	11.68	20.11	17.90	21.81	10.60	10.80	7.10	10	scl	14.54	8.96

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO <sub>3</sub>	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP
	Water	CaCl <sub>2</sub>	M KCl				Ca	Mg	K	Na	Total				
				dS m <sup>-1</sup>	%	%	cmol kg <sup>-1</sup>						%	%	
0-15	8.42	-	-	0.148	0.70	0.65	-	-	0.15	0.03	-	14.50	0.74	100	0.18
15-38	8.38	-	-	0.226	0.31	2.21	-	-	0.09	0.23	-	21.70	0.75	100	1.05
38-50	8.40	-	-	0.195	0.25	1.17	-	-	0.07	0.19	-	15.90	0.79	100	1.23

*Contd...*

**Soil Series:** Hosalli (HSL) **Pedon:** R-3

**Location:** 16°46'60.3"N 77°05'47.6"E, Mudhanala village, Yadgir hobli, Yadgir taluk and district

**Analysis at:** NBSS&LUP, Regional Centre, Bengaluru **Classification:** Fine, mixed, isohyperthermic Typic Haplustepts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-10	Ap	88.43	5.15	6.42	5.69	6.40	36.04	27.31	12.99	-	s	7.40	2.74
10-30	Bw1	58.47	7.24	34.29	4.26	9.37	19.91	19.28	5.64	-	scl	19.07	11.57
30-50	Bw2	51.43	12.67	35.90	3.49	8.89	16.72	15.87	6.46	<15	sc	21.64	12.44
50-90	Bw3	49.89	13.64	36.47	2.43	2.96	20.61	16.17	7.72	<15	sc	21.12	12.95

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO <sub>3</sub>	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP			
	Water	CaCl <sub>2</sub>	M KCl				dS m <sup>-1</sup>	%	%	Ca	Mg					K	Na	Total
										cmol kg <sup>-1</sup>						%	%	
0-10	7.16	-	-	0.117	0.48	0.00	2.83	1.50	0.15	0.29	4.76	4.90	0.76	97	5.94			
10-30	6.91	-	-	0.040	0.36	0.00	10.64	5.43	0.10	0.26	16.43	17.80	0.52	92	1.47			
30-50	8.17	-	-	0.182	0.24	1.43	-	-	0.12	0.22	-	19.90	0.55	100	1.08			
50-90	8.60	-	-	0.148	0.20	4.29	-	-	0.13	0.16	-	19.70	0.54	100	0.81			

Contd...

**Soil Series:** Naglapur (NGP) **Pedon:** R-8

**Location:** 16°52'84.1"N 77°22'99.4"E, Gurumitkal village, Gurumitkal hobli, Yadgir taluk and district

**Analysis at:** NBSS&LUP, Regional Centre, Bengaluru **Classification:** Very fine, smectitic (calcareous), isohyperthermic Typic Haplusterts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-10	Ap	7.53	19.88	72.59	1.00	0.78	0.89	2.10	2.77	-	c	44.31	32.79
10-35	Bss1	6.55	18.76	74.68	0.80	0.92	0.80	1.72	2.30	-	c	43.09	31.62
35-60	Bss2	6.58	21.05	72.37	0.69	0.46	1.04	1.50	2.89	-	c	46.52	32.52
60-102	Bss3	7.48	19.74	72.78	1.61	1.38	0.69	1.61	2.19	-	c	51.12	35.62

Depth (cm)	pH (1:2.5)			E.C. (1:2.5) dS m <sup>-1</sup>	O.C. %	CaCO <sub>3</sub> %	Exchangeable bases					CEC	CEC/Clay	Base saturation %	ESP %
	Water	CaCl <sub>2</sub>	M KCl				Ca	Mg	K	Na	Total				
0-10	7.42	-	-	0.24	0.84	1.30	-	-	0.84	0.15	-	67.10	0.92	100	0.22
10-35	8.52	-	-	0.291	0.64	2.86	-	-	0.17	0.29	-	65.20	0.87	100	0.45
35-60	7.89	-	-	0.134	0.62	4.55	-	-	0.15	0.20	-	65.00	0.90	100	0.30
60-102	8.68	-	-	0.213	0.54	8.32	-	-	0.17	0.15	-	64.10	0.88	100	0.24

Contd...

**Soil Series:** Baddeppalli (BDP) **Pedon:** R-11

**Location:** 16°43'84.4"N 77°14'06.4"E, Halagera village, Yadgir hobli, Yadgir taluk and district

**Analysis at:** NBSS&LUP, Regional Centre, Bengaluru **Classification:** Loamy, mixed (calcareous), isohyperthermic, Lithic Ustorthents

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-16	Ap	58.67	17.02	24.31	19.03	13.74	9.62	10.57	5.71	<15	scl	16.19	8.18

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO <sub>3</sub>	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP
	Water	CaCl <sub>2</sub>	M KCl				Ca	Mg	K	Na	Total				
				dS m <sup>-1</sup>	%	%	cmol kg <sup>-1</sup>					%	%		
0-16	8.58	-	-	0.262	1.60	7.67	-	-	0.24	0.06	-	18.10	0.74	100	0.35

*Contd...*

**Soil Series:** Hattikuni (HTK), Pedon: R-7

**Location:** 16°50'46.5"N 77°10'16.4"E, Yaddalli village, Hattikuni hobli, Yadgir taluk and district

**Analysis at:** NBSS&LUP, Regional Centre, Bengaluru **Classification:** Mixed, isohyperthermic, Lithic Ustipsamments

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-12	Ap	90.89	5.62	3.49	8.50	13.46	29.86	29.55	9.51	20	s	7.73	3.16
12-22	A1	89.97	6.53	3.50	7.19	13.48	29.48	29.79	10.03	20	s	8.00	3.05
22-45	A2	87.20	6.43	6.38	11.09	14.42	31.55	7.16	22.98	40	ls	7.67	3.96

Depth (cm)	pH (1:2.5)			E.C. (1:2.5) dS m <sup>-1</sup>	O.C. %	CaCO <sub>3</sub> %	Exchangeable bases					CEC	CEC/Clay	Base saturation %	ESP %
	Water	CaCl <sub>2</sub>	M KCl				Ca	Mg	K	Na	Total				
0-12	6.81	-	-	0.062	0.07	-	2.35	0.50	0.16	0.01	3.02	3.0	0.86	100	0.38
12-22	6.80	-	-	0.050	0.21	-	1.67	0.30	0.09	0.01	2.07	2.4	0.69	86.30	0.45
22-45	6.85	-	-	0.044	0.19	-	1.82	0.42	0.10	0.06	2.40	2.6	0.41	92.41	2.17

*Contd...*

**Soil Series:** Thumakuru (TMK) **Pedon:** R-10

**Location:** 16°38'01.3"N 77°16'49.8"E, Kilankera village, Balichakra hobli, Yadgir taluk and district

**Analysis at:** NBSS&LUP, Regional Centre, Bengaluru **Classification:** Fine, mixed, isohyperthermic Typic Haplustepts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-12	Ap	62.92	15.76	21.32	5.56	9.37	21.83	18.33	7.83	-	scl	17.98	6.60
12-29	Bw1	45.91	18.53	35.56	6.08	8.18	15.41	11.43	4.82	-	sc	33.40	11.79
29-74	Bw2	48.47	16.24	35.29	5.93	9.84	16.40	11.75	4.55	-	sc	28.66	11.19
74-132	Bw3	38.25	20.59	41.16	3.21	8.23	14.64	8.97	3.21	-	c	38.85	14.72
132-158	Bw4	36.87	19.99	43.14	3.54	7.61	13.08	8.57	4.07	-	c	44.36	15.75

Depth (cm)	pH (1:2.5)			E.C. (1:2.5) dS m <sup>-1</sup>	O.C. %	CaCO <sub>3</sub> %	Exchangeable bases					CEC	CEC/Clay	Base saturation %	ESP %
	Water	CaCl <sub>2</sub>	M KCl				Ca	Mg	K	Na	Total				
0-12	9.60	-	-	0.35	0.48	1.44	-	-	0.23	3.62	-	21.83	1.02	100	6.63
12-29	9.72	-	-	1.27	0.50	1.44	-	-	0.59	20.88	-	30.50	0.86	100	27.39
29-74	9.16	-	-	3.44	0.31	3.72	-	-	0.38	25.84	-	28.68	0.81	100	36.04
74-132	9.33	-	-	2.52	0.23	4.92	-	-	0.82	20.25	-	34.99	0.85	100	23.148
132-158	9.23	-	-	2.07	0.31	3.48	-	-	0.70	21.03	-	34.24	0.79	100	24.564

## INTERPRETATION FOR LAND RESOURCE MANAGEMENT

The most important soil and site characteristics that affect the land use and conservation needs of an area are land capability, soil depth, soil texture, coarse fragments, available water capacity, soil slope, soil erosion, soil reaction etc. These are interpreted from the data base generated through land resource inventory and several thematic maps are generated. These would help in identifying the areas suitable for growing crops, soil and water conservation measures and structures needed thus helping to maintain good soil health for sustained crop production. The various interpretative and thematic maps generated are described below.

### 5.1 Land Capability Classification

Land capability classification is an interpretative grouping of soil map units (soil phases) mainly based on inherent soil characteristics, external land features and environmental factors that limit the use of land for agriculture, pasture, forestry, or other uses on a sustained basis (IARI, 1971). The land and soil characteristics used to group the land resources in an area into various land capability classes, subclasses and units are

*Soil Characteristics:* Depth, texture, gravelliness, calcareousness.

*Land characteristics:* Slope, erosion, drainage, rock outcrops.

*Climate:* Total rainfall and its distribution, and length of crop growing period.

The Land capability classification system is divided into land capability classes, subclasses and units based on the level of information available. Eight land capability classes are recognized. They are

*Class I:* They are very good lands that have no limitations or very few limitations that restrict their use.

*Class II:* They are good lands that have minor limitations and require moderate conservation practices.

*Class III:* They are moderately good lands that have moderate limitations that reduce the choice of crops or that require special conservation practices.

*Class IV:* They are fairly good lands that have very severe limitations that reduce the choice of crops or that require very careful management.

*Class V:* Soils in these lands are not likely to erode, but have other limitations like wetness that are impractical to remove and as such not suitable for agriculture, but suitable for pasture or forestry with minor limitations.

*Class VI:* The lands have severe limitations that make them generally unsuitable for cultivation, but suitable for pasture or forestry with moderate limitations.

*Class VII:* The lands have very severe limitations that make them unsuitable for cultivation, but suitable for pasture or forestry with major limitations.

*Class VIII:* Soil and other miscellaneous areas (rock lands) that have very severe limitations that nearly preclude their use for any crop production, but suitable for wildlife, recreation and installation of wind mills.

The land capability subclasses are recognised based on the dominant limitations observed within a given land capability class. The subclasses are designated by adding a lower case letter like 'e', 'w', 's', or 'c' to the class numeral. The subclass "e" indicates that the main hazard is risk of erosion, "w" indicates drainage or wetness as a limitation for plant growth, "s" indicates shallow soil depth, coarse or heavy textures, calcareousness, salinity/alkalinity or gravelliness and "c" indicates limitation due to climate.

The land capability subclasses have been further subdivided into land capability units based on the kinds of limitations present in each subclass. Ten land capability units are used in grouping the soil map units. They are stony or rocky (0), erosion hazard (slope, erosion) (1), coarse texture (sand, loamy sand, sandy loam) (2), fine texture (cracking clay, silty clay) (3), slowly permeable subsoil (4), coarse underlying material (5), salinity/alkali (6), stagnation, overflow, high ground water table (7), soil depth (8) and fertility problems (9). The capability units thus identified have similar soil and land characteristics that respond similarly to a given level of management. The soils of the microwatershed have been classified up to 1 and capability subclass level.

The 10 soil map units identified in the Mundragi-3 microwatershed are grouped under 3 land capability classes and 4 subclasses. Entire area in the microwatershed is suitable for agriculture (Fig. 5.1).

Good cultivable lands (Class II) cover an area of about 20 per cent and are distributed in the northern, southern, southwestern and northeastern part of the microwatershed with minor problems of soil and erosion. Moderately good cultivable lands (Class III) cover an area of about 54 per cent and are distributed in the major part of the microwatershed with moderate problems of soil and erosion. Fairly good cultivation lands (Class IV) cover an area of about 2 per cent and are distributed in the eastern part of the microwatershed with very severe problems of soil and erosion.






Fig. 5.1 Land Capability map of Mundragi-3 Microwatershed

## 5.2 Soil Depth

Soil depth refers to the depth of the soil occurring above the parent material or hard rock. The depth of the soil determines the effective rooting depth for plants and in accordance with soil texture, mineralogy and gravel content, the capacity of the soil column to hold water and nutrient availability. Soil depth is one of the most important soil characteristic that is used in differentiating soils into different soil series. The soil depth classes used in identifying soils in the field are very shallow (<25 cm), shallow (25-50 cm), moderately shallow (50-75 cm), moderately deep (75-100 cm), deep (100-150 cm) and very deep (>150 cm). They were used to classify the soils into different depth classes and a soil depth map was generated. The area extent and their geographical distribution in the microwatershed is given in Fig. 5.2.

Very shallow (<25 cm) soils occur in an area of 9 ha (2%) and are distributed in the eastern part of the microwatershed. Shallow (25-50 cm) soils occur in a maximum area of 323 ha (54%) and are distributed in the major part of the microwatershed. Moderately shallow (50-75 cm) and moderately deep (75-100 cm) soils occupy an area of about 93 ha (16%) and 3 ha (<1%) respectively of the microwatershed and are distributed in the northeastern, western and southern part of the microwatershed. Deep (100-150 cm) soils cover an area of 22 ha (4%) and are distributed in the northern and eastern part of the microwatershed. Very deep (>150 cm) soils occur in a small area of 3 ha (<1%) and are distributed in the northern part of the microwatershed.




Fig. 5.2 Soil Depth map of Mundragi-3 Microwatershed

The most productive lands cover an area of 25 ha (4%) with respect to soil rooting depth where all climatically adapted annual and perennial crops can be grown are deep to very deep (100->150 cm depth) soils occurring in the northern and eastern part of the microwatershed. The problematic soils cover about 56 per cent area where the soils are shallow and very shallow which are suitable for short duration crops and probability of crop failure is high.

### 5.3 Surface Soil Texture

Texture is an expression to indicate the coarseness or fineness of the soil as determined by the relative proportion of primary particles of sand, silt and clay. It has a direct bearing on the structure, porosity, adhesion and consistence. The surface layer of a soil to a depth of about 25 cm is the layer that is most used by crops and plants. The surface soil textural class provides a guide to understanding soil-water retention and availability, nutrient holding capacity, infiltration, workability, drainage, physical and chemical behaviour, microbial activity and crop suitability. The textural classes used for LRI were used to classify and a surface soil texture map was generated. The area extent and their geographical distribution in the microwatershed is shown in Figure 5.3.

An area of about 311 ha (52%) is loamy and are distributed in the major part of the microwatershed. An area of 142 ha (24%) has soils that are clayey at the surface and occur in the central, northern, eastern and northeastern part of the microwatershed.

Entire area has most productive lands with respect to surface soil texture. The clayey soils (24%) have high potential for soil-water retention and availability, and nutrient retention and availability, but have more problems of drainage, infiltration, workability and other physical problems. The other productive lands are loamy soils (52%) which also have high potential for soil-water retention and nutrient availability but have no drainage or other physical problems.




Fig. 5.3 Surface Soil Texture map of Mundragi-3 Microwatershed

#### 5.4 Soil Gravelliness

Gravel is the term used for describing coarse fragments between 2 mm and 7.5 cm diameter and stones for those between 7.5 cm and 25 cm. The presence of gravel and stones in soil reduces the volume of soil responsible for moisture and nutrient storage, drainage, infiltration and runoff, and hinders plant growth by impeding root growth and seedling emergence, intercultural operations and farm mechanization. The gravelliness classes used in LRI were used to classify the soils and using these classes a gravelliness map was generated. The area extent and their geographic distribution in the microwatershed is shown in Figure 5.4.

Non gravelly (<15%) soils cover a maximum area of about 317 ha (53%) and are distributed in the major part of the microwatershed. An area of about 136 ha (23%) is gravelly (15-35%) and are distributed in the central, northern and western part of the microwatershed.




Fig. 5.4 Soil Gravelliness map of Mundragi-3 Microwatershed

The problem soils (23%) that are gravelly (15-35%), where only short or medium duration crops can be grown. The most productive soils (53%) that are non gravelly (<15%) where, all climatically adapted long duration crops can be grown.

### 5.5 Available Water Capacity

The soil available water capacity (AWC) is estimated based on the ability of the soil column to retain water between the tensions of 0.33 and 15 bar in a depth of 100 cm or the entire solum if the soil is shallower. The AWC of the soils (soil series) as estimated by considering the soil texture, mineralogy, soil depth and gravel content (Sehgal *et al.*, 1990) and accordingly the soil map units were grouped into five AWC classes *viz*, very low (<50 mm/m), low (50-100 mm/m), medium (100-150 mm/m), high (150-200 mm/m) and very high (>200 mm/m) and using these values, an AWC map was generated. The area extent and their geographic distribution of different AWC classes in the microwatershed is given in Figure 5.5.

A major area of about 410 ha (69%) in the microwatershed has soils that are very low (<50 mm/m) in available water capacity and are distributed in the major part of the microwatershed. An area of about 18 ha (3%) in the microwatershed has soils that are low (51-100 mm/m) in available water capacity and are distributed in the northern, southern and northeastern part of the microwatershed. Very high (>200 mm/m) in 24 ha (4%) and are distributed in the northern and central part of the microwatershed.




Fig. 5.5 Soil Available Water Capacity map of Mundragi-3 Microwatershed

About 428 ha (72%) area in the microwatershed has soils that are problematic with regard to available water capacity. Here, only short duration crops can be grown and the probability of crop failure is very high. These areas are best put to other alternative uses. An area of 24 ha (4%) are potential with regard to AWC where all climatically adapted annual and perennial crops can be grown.

### 5.6 Soil Slope

Soil slope refers to the inclination of the surface of the land. It is defined by gradient, shape and length, and is an integral feature of any soil as a natural body. Slope is considered important in soil genesis, land use and land development. The length and gradient of slope influences the rate of runoff, infiltration, erosion and deposition. The soil map units were grouped into two slope classes and a slope map was generated showing the area extent and their geographic distribution in the microwatershed (Fig. 5.6).

An area of about 3 ha (<1%) falls under nearly level (0-1%) and are distributed in the northern part of the microwatershed. An area of about 450 ha (75%) falls under very gently sloping (1-3% slope) lands and are distributed in all parts of the microwatershed.

In these areas, all climatically adapted annual and perennial crops can be grown without much soil and water conservation and other land development measures.




Fig. 5.6 Soil Slope map of Mundragi-3 Microwatershed

## 5.7 Soil Erosion

Soil erosion refers to the wearing away of the earth's surface by the forces of water, wind and ice involving detachment and transport of soil by raindrop impact. It is used for accelerated soil erosion resulting from disturbance of the natural landscape by burning, excessive grazing and indiscriminate felling of forest trees and tillage, all usually by man. The erosion classes showing an estimate of the current erosion status as judged from field observations in the form of rills, gullies or a carpet of gravel on the surface are recorded. Four erosion classes, viz, slight erosion (e1), moderate erosion (e2), severe erosion (e3) and very severe erosion (e4) are recognized. The soil map units were grouped into different erosion classes and a soil erosion map generated. The area extent and their spatial distribution in the microwatershed is given in Figure 5.7.

Soils that are slightly eroded (e1 class) cover an area of 3 ha (<1%) and are distributed in the northern part of the microwatershed. Moderately eroded (e2 class) soils cover an area of 450 ha (75%) and are distributed in all parts of the microwatershed.

Almost entire area in the microwatershed is problematic because of moderate erosion. For these areas, taking up soil and water conservation and other land development measures are needed.




Fig. 5.7 Soil Erosion map of Mundragi-3 Microwatershed





## **FERTILITY STATUS**

Soil fertility plays an important role in increasing crop yield. The adoption of high yielding varieties that require high amounts of nutrients has resulted in deficiency symptoms in crops and plants due to imbalanced fertilization and poor inherent fertility status as these areas are characterised by low rainfall and high temperatures. Hence, it is necessary to know the fertility (macro and micro nutrients) status of the soils of the watersheds for assessing the kind and amount of fertilizers required for each of the crop intended to be grown. For this purpose, the surface soil samples collected from the grid points (one soil sample at every 320 m interval) all over the microwatershed through land resource inventory in the year 2018 were analysed for pH, EC, organic carbon, available phosphorus and potassium, and for micronutrients like zinc, boron, copper, iron and manganese, and secondary nutrient sulphur.

Soil fertility data generated has been assessed and individual maps for all the nutrients for the microwatershed have been generated using Kriging method under GIS. The village/survey number wise fertility data for the microwatershed is given in Appendix-II.

### **6.1 Soil Reaction (pH)**

The soil analysis of the Mundragi-3 microwatershed for soil reaction (pH) showed that an area of about 193 ha (32%) is neutral (pH 6.5-7.3) and are distributed in the northern, western, northwestern, eastern and northeastern part of the microwatershed. A maximum area of about 260 ha (43%) is slightly alkaline (pH 7.3-7.8) and are distributed in the major part of the microwatershed. In all, major area of about 193 ha is neutral and 260 ha is alkaline.

### **6.2 Electrical Conductivity (EC)**

The Electrical Conductivity of the soils of the entire microwatershed area is  $<2 \text{ dS m}^{-1}$  (Fig 6.2) and as such the soils are non-saline.

### **6.3 Organic Carbon**

The soil organic carbon content (an index of available Nitrogen) in the soils of the microwatershed is high ( $>0.75$ ) in an area of about less than 1 ha ( $<1\%$ ) and are distributed in the southeastern part of the microwatershed and medium (0.5-0.75%) covering a maximum area of about 453 ha (76%) and are distributed in the major part of the microwatershed (Fig. 6.3).




Fig.6.1 Soil Reaction (pH) map of Mundragi-3 Microwatershed




Fig.6.2 Electrical Conductivity (EC) map of Mundragi-3 Microwatershed




Fig.6.3 Soil Organic Carbon map of Mundragi-3 Microwatershed

#### 6.4 Available Phosphorus

Available phosphorus content is low (<23 kg/ha) in an area of 158 ha (26%) and are distributed in the northern, southern, western, northwestern and southwestern part of the microwatershed. Medium (23-57 kg/ha) in available phosphorus occur in a maximum area of about 295 ha (49%) and are distributed in the major part of the microwatershed (Fig. 6.4).

#### 6.5 Available Potassium

Available potassium content is low (<145 kg/ha) in an area of about 52 ha (9%) and are distributed in the northeastern and southeastern part of the microwatershed. Medium (145-337 kg/ha) in available potassium occur in a maximum area of about 401 ha (67%) and are distributed in all parts of the microwatershed (Fig. 6.5).

#### 6.6 Available Sulphur

An area of about 57 ha (9%) is low (<10 ppm) in available sulphur content and are distributed in the northern, northeastern and northwestern part of the microwatershed and medium (10-20 ppm) in a maximum area of about 396 ha (66%) and are distributed in the major part of the microwatershed (Fig. 6.6).

#### 6.7 Available Boron

Available boron content is low (<0.5 ppm) in the entire microwatershed (Fig. 6.7).

### 6.8 Available Iron

Available iron content is sufficient (>4.5 ppm) in the entire microwatershed (Fig 6.8).

### 6.9 Available Manganese

Available manganese content is sufficient (>1.0 ppm) in the entire microwatershed area (Fig 6.9).

### 6.10 Available Copper

Available copper content is sufficient (>0.2 ppm) in the entire microwatershed area (Fig 6.10).

### 6.11 Available Zinc

Available zinc content is deficient (<0.6 ppm) in the maximum area of about 318 ha (53%) and are distributed in the major part of the microwatershed and sufficient (>0.6 ppm) in an area of about 135 ha (23%) and are distributed in the southern and southeastern part of the microwatershed (Fig 6.11).




Fig.6.4 Soil Available Phosphorus map of Mundragi-3 Microwatershed




Fig.6.5 Soil Available Potassium map of Mundragi-3 Microwatershed




Fig.6.6 Soil Available Sulphur map of Mundragi-3 Microwatershed




Fig.6.7 Soil Available Boron map of Mundragi-3 Microwatershed




Fig.6.8 Soil Available Iron map of Mundragi-3 Microwatershed




Fig.6.9 Soil Available Manganese map of Mundragi-3 Microwatershed




Fig.6.10 Soil Available Copper map of Mundragi-3 Microwatershed




Fig.6.11 Soil Available Zinc map of Mundragi-3 Microwatershed



## LAND SUITABILITY FOR MAJOR CROPS

The soil and land resource units (soil phases) of Mundragi-3 microwatershed were assessed for their suitability for growing food, fodder, fibre and other horticulture crops by following the procedure as outlined in FAO, 1976 and 1983. Crop requirements were developed for each of the crop from the available research data and also by referring to Naidu *et. al.* (2006) and Natarajan *et. al* (2015). The soil and land characteristics were matched with the crop requirement to arrive at the crop suitability. The soil and land characteristics table (Table 7.1) and crop requirement tables (Tables 7.2 to Tables 7.30) are given at the end. In FAO land suitability classification, two orders are recognized. Order S-Suitable and Order N-Not suitable. The orders have classes, subclasses and units. Order-S has three classes, Class S1-Highly Suitable, Class S2-Moderately Suitable and Class S3- Marginally Suitable. Order N has two classes, N1-Currently not Suitable and N2- Permanently not Suitable. There are no subclasses within the Class S1 as they will have very minor or no limitations for crop growth. Classes S2, S3 and N1 are divided into subclasses based on the kinds of limitations encountered. The limitations that affect crop production are 'c' for erratic rainfall and its distribution and length of growing period (LGP), 'e' for erosion hazard, 'r' for rooting condition, 't' for lighter or heavy texture, 'g' for gravelliness or stoniness, 'n' for nutrient availability, 'l' for topography, 'm' for moisture availability, 'w' for drainage and 'z' for calcareousness. These limitations are indicated as lower case letters to the Class symbol. For example, moderately suitable lands with the limitations of soil depth and erosion are designated as S2re. For the microwatershed, the soil mapping units were evaluated and classified up to subclass level.

Using the above criteria, the soil map units of the microwatershed were evaluated and land suitability maps for 29 major annual and perennial crops were generated. The detailed information on the kind of suitability of each of the soil phase for the crops assessed are given village/ survey number wise for the microwatershed in Appendix-III.

### 7.1 Land Suitability for Sorghum (*Sorghum bicolor*)

Sorghum is one of the major crop grown in Karnataka in an area of 10.47 lakh ha in Bijapur, Gulbarga, Raichur, Bidar, Belgaum, Dharwad, Bellary, Chitradurga, Mysore and Tumakuru districts. The crop requirements for growing sorghum (Table 7.2) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing sorghum was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.1.

A small area of about 3 ha (<1%) is highly suitable (Class S1) for growing sorghum and are distributed in the northern part of the microwatershed with no limitations. An area of about 40 ha (7%) is moderately suitable (Class S2) for growing sorghum and are distributed in the northern, eastern, northwestern and northeastern part

of the microwatershed. They have minor limitations of texture, drainage, calcareousness and rooting depth. Maximum area of about 401 ha (67%) is marginally suitable (Class S3) for growing sorghum and are distributed in the major part of the microwatershed with moderate limitations rooting depth and texture. An area of about 9 ha (2%) is currently not suitable (Class N1) and are distributed in the eastern part of the microwatershed with severe limitation of rooting depth.




Fig. 7.1 Land Suitability map of Sorghum

## 7.2 Land Suitability for Maize (*Zea mays*)

Maize is one of the most important food crop grown in an area of 13.37 lakh ha in almost all the districts of the State. The crop requirements for growing maize (Table 7.3) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing maize was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.2.

An area of about 43 ha (7%) is moderately suitable (Class S2) for growing maize and occur in the northern, southern, northwestern and northeastern part of the microwatershed. They have minor limitations of texture, calcareousness and drainage. Marginally suitable lands (Class S3) for growing maize occupy a maximum area of about 401 ha (67%) and occur in all parts of the microwatershed. They have moderate limitations of rooting depth and texture. An area of about 9 ha (2%) is currently not

suitable (Class N1) and are distributed in the eastern part of the microwatershed with severe limitation of rooting depth.




Fig. 7.2 Land Suitability map of Maize

### 7.3 Land Suitability for Bajra (*Pennisetum glaucum*)

Bajra is one of the most important millet crop grown in an area of 2.34 lakh ha in the northern districts of Karnataka state. The crop requirements for growing bajra (Table 7.4) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing bajra was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.3.

An area of about 43 ha (7%) is moderately suitable (Class S2) for growing bajra and occur in the northern, southern, northwestern and northeastern part of the microwatershed. They have minor limitations of texture, calcareousness and drainage. Marginally suitable lands (Class S3) for growing bajra occupy a maximum area of about 401 ha (67%) and occur in all parts of the microwatershed. They have moderate limitations of rooting depth and texture. An area of about 9 ha (2%) is currently not suitable (Class N1) and are distributed in the eastern part of the microwatershed with severe limitation of rooting depth.




Fig. 7.3 Land Suitability map of Bajra

#### 7.4 Land Suitability for Groundnut (*Arachis hypogaea*)

Groundnut is one of the major oilseed crop grown in an area of 6.54 lakh ha in Karnataka in most of the districts either as rainfed or irrigated crop. The crop requirements for growing groundnut (Table 7.5) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing groundnut was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.4.

A small area of about 3 ha (<1%) is moderately suitable (Class S2) for groundnut and are distributed in the northwestern part of the microwatershed. It has minor limitation of calcareousness. Marginally suitable lands (Class S3) for growing groundnut occupy a maximum area of about 440 ha (74%) and are distributed in all parts of the microwatershed. They have moderate limitations of texture, drainage and rooting depth. An area of about 9 ha (2%) is currently not suitable (Class N1) and are distributed in the eastern part of the microwatershed with severe limitation of rooting depth.




Fig. 7.4 Land Suitability map of Groundnut

## 7.5 Land Suitability for Sunflower (*Helianthus annus*)

Sunflower is one of the most important oilseed crop grown in an area of 3.56 lakh ha in the State in all the districts. The crop requirements for growing sunflower (Table 7.6) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing sunflower was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.5.

An area of about 28 ha (5%) is moderately suitable (Class S2) for growing sunflower and are distributed in the northern, northwestern and eastern part of the microwatershed. It has minor limitations of rooting depth, drainage and calcareousness. Marginally suitable (Class S3) lands for sunflower are found to occur in an area of about 93 ha (16%) with moderate limitations of rooting depth and texture and are distributed in the northeastern, southern and western part of the microwatershed. A maximum area of about 333 ha (56%) is currently not suitable (Class N1) and are distributed in the major part of the microwatershed with severe limitation of rooting depth.




Fig. 7.5 Land Suitability map of Sunflower

## 7.6 Land Suitability for Red gram (*Cajanus Cajan*)

Redgram is one of the most important pulse crop grown in an area of 7.28 lakh ha in almost all the districts of the State. The crop requirements for growing red gram (Table 7.7) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing redgram was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.6.

An area of about 27 ha (5%) is moderately suitable (Class S2) for growing redgram and are distributed in the northern, northwestern and eastern part of the microwatershed. They have minor limitations of rooting depth, calcareousness, texture and drainage. Marginally suitable lands (Class S3) for growing redgram occupy a maximum area of about 338 ha (57%) and occur in the major part of the microwatershed. They have moderate limitations of rooting depth and texture. An area of about 87 ha (15%) is currently not suitable (Class N1) and are distributed in the central, eastern, western and northwestern part of the microwatershed with severe limitation of rooting depth.




Fig. 7.6 Land Suitability map of Redgram

### 7.7 Land Suitability for Bengal gram (*Cicer aerativum*)

Bengal gram is one of the most important pulse crop grown in about 9.39 lakh ha area in Bijapur, Raichur, Kalaburgi, Dharwad, Belgaum and Bellary districts. The crop requirements for growing Bengal gram (Table 7.8) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing Bengal gram was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.7.

Highly (Class S1) suitable lands for growing Bengal gram occur in an area of about 24 ha (4%) and are distributed in the northern and eastern part of the microwatershed. An area of about 15 ha (3%) is moderately suitable (Class S2) for growing Bengal gram and are distributed in the southern and northeastern part of the microwatershed. It has minor limitation of rooting depth. Marginally suitable lands (Class S3) occupy a maximum area of about 248 ha (41%) and are distributed in the major part of the microwatershed. They have moderate limitations of rooting depth, texture and calcareousness. Currently not suitable (Class N1) lands occur in an area of 165 ha (28%) and are distributed in the central, northern, southern, eastern, western and northwestern part of the microwatershed with severe limitations of texture and rooting depth.




Fig. 7.7 Land Suitability map of Bengal gram.

### 7.8 Land Suitability for Cotton (*Gossypium hirsutum*)

Cotton is one of the most important fibre crop grown in the State in about 8.75 lakh ha area in Raichur, Dharwad, Belgaum, Kalaburgi, Bijapur, Bidar, Bellary, Chitradurga and Chamarajnagar districts. The crop requirements for growing cotton (Table 7.9) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing cotton was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.8.

Highly (Class S1) suitable lands for growing cotton occur in an area of about 24 ha (4%) and are distributed in the northern and eastern part of the microwatershed. An area of about 15 ha (3%) is moderately suitable (Class S2) for growing cotton and are distributed in the southern and northeastern part of the microwatershed. It has minor limitation of rooting depth. Marginally suitable lands (Class S3) occupy a maximum area of about 248 ha (41%) and are distributed in the major part of the microwatershed. They have moderate limitations of rooting depth, texture and calcareousness. Currently not suitable (Class N1) lands occur in an area of 165 ha (28%) and are distributed in the central, northern, southern, eastern, western and northwestern part of the microwatershed with severe limitations of texture and rooting depth.






Fig. 7.8 Land Suitability map of Cotton

### 7.9 Land Suitability for Chilli (*Capsicum annuum*)

Chilli is one of the most important spice crop grown in about 0.42 lakh ha in Karnataka state. The crop requirements for growing chilli (Table 7.10) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing chilli was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.9.

An area of about 21 ha (3%) is moderately suitable (Class S2) for growing chilli and are distributed in the northwestern and northeastern part of the microwatershed. They have minor limitations of texture, drainage, calcareousness and rooting depth. Marginally suitable lands (Class S3) occupy a major area of about 423 ha (71%) and are distributed in the major part of the microwatershed. They have moderate limitations of rooting depth, drainage and texture. An area of about 9 ha (2%) is currently not suitable (Class N1) and are distributed in the eastern part of the microwatershed with severe limitation of rooting depth.




Fig 7.9 Land Suitability map of Chilli

### 7.10 Land Suitability for Tomato (*Lycopersicon esculentum*)

Tomato is one of the most important vegetable crop grown in about 0.61 lakh ha covering almost all the district of the state. The crop requirements for growing tomato (Table 7.11) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing tomato was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.10.

An area of about 18 ha (3%) is moderately suitable (Class S2) for growing tomato and are distributed in the northwestern, northeastern and southeastern part of the microwatershed. They have minor limitations of calcareousness and rooting depth. Marginally suitable lands (Class S3) occupy a major area of about 425 ha (71%) and are distributed in all parts of the microwatershed. They have moderate limitations of rooting depth, drainage and texture. An area of about 9 ha (2%) is currently not suitable (Class N1) and are distributed in the eastern part of the microwatershed with severe limitation of rooting depth.




Fig 7.10 Land Suitability map of Tomato

### 7.11 Land Suitability for Brinjal (*Solanum melongena*)

Brinjal is one of the most important vegetable crop grown in the state. The crop requirements for growing Brinjal (Table 7.12) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing Brinjal was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.11.

Highly (Class S1) suitable lands for growing brinjal occur in a small area of about 3 ha (<1%) and are distributed in the northern part of the microwatershed. An area of about 40 ha (7%) is moderately suitable (Class S2) for brinjal and is distributed in the northern, eastern, northwestern, northeastern and southeastern part of the microwatershed. They have minor limitations of texture and rooting depth. A maximum area of 401 ha (67%) is marginally suitable (Class S3) and is distributed in the major part of the microwatershed with moderate limitations of rooting depth and texture. An area of about 9 ha (2%) is currently not suitable (Class N1) and are distributed in the eastern part of the microwatershed with severe limitation of rooting depth.




Fig 7.11 Land Suitability map of Brinjal

## 7.12 Land Suitability for Onion (*Allium cepa L.*)

Onion is one of the most important vegetable crop grown in the state. The crop requirements for growing onion (Table 7.13) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing onion was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.12.

Highly (Class S1) suitable lands for growing onion occur in a small area of 3 ha (<1%) and are distributed in the northwestern part of the microwatershed. An area of about 18 ha (3%) is moderately suitable (Class S2) for onion and is distributed in the northwestern, northeastern and southeastern part of the microwatershed. They have minor limitations of texture and rooting depth. A maximum area of 423 ha (71%) is marginally suitable (Class S3) and is distributed in all parts of the microwatershed with moderate limitations of rooting depth and texture. An area of about 9 ha (2%) is currently not suitable (Class N1) and are distributed in the eastern part of the microwatershed with severe limitation of rooting depth.




Fig 7.12 Land Suitability map of Onion

### 7.13 Land Suitability for Bhendi (*Abelmoschus esculentus*)

Bhendi is one of the most important vegetable crop grown in the state. The crop requirements for growing bhendi (Table 7.14) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing bhendi was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.13.

Highly (Class S1) suitable lands for growing bhendi occur in a small area of 5 ha (<1%) and are distributed in the northwestern part of the microwatershed. An area of about 37 ha (7%) is moderately suitable (Class S2) for bhendi and is distributed in the northern, northeastern, eastern and southeastern part of the microwatershed. They have minor limitations of texture and rooting depth. A maximum area of about 401 ha (67%) is marginally suitable (Class S3) and is distributed in the major part of the microwatershed with moderate limitations of rooting depth and texture. An area of about 9 ha (2%) is currently not suitable (Class N1) and are distributed in the eastern part of the microwatershed with severe limitation of rooting depth.




Fig 7.13 Land Suitability map of Bhendi

#### 7.14 Land Suitability for Drumstick (*Moringa oleifera*)

Drumstick is one of the most important vegetable crop grown in about 2403 ha in the state. The crop requirements for growing drumstick (Table 7.15) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing drumstick was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.14.

An area of about 27 ha (5%) is moderately suitable (Class S2) for growing drumstick and are distributed in the northern, northwestern and eastern part of the microwatershed. They have minor limitations of rooting depth, texture, calcareousness and drainage. An area of about 93 ha (16%) is marginally suitable (Class S3) for growing drumstick and are distributed in the southern, northeastern, western and southeastern part of the microwatershed. They have moderate limitations of rooting depth and texture. A maximum area of about 332 ha (56%) is currently not suitable (Class N1) for growing drumstick and are distributed in the major part of the microwatershed. They have severe limitations of rooting depth and texture.




Fig 7.14 Land Suitability map of Drumstick

### 7.15 Land suitability for Mango (*Mangifera indica*)

Mango is one of the most important fruit crop grown in an area of 1.73 lakh ha in almost all the districts of the State. The crop requirements (Table 7.16) for growing mango were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing mango was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed is given in Figure 7.15.

An area of 28 ha (5%) is marginally suitable (Class S3) for growing mango with moderate limitations of calcareousness, drainage, texture and rooting depth and are distributed in the northern, eastern and northwestern part of the microwatershed. A maximum area of about 426 ha (71%) is currently not suitable (Class N1) for growing mango and occur in all parts of the microwatershed with severe limitation of rooting depth.




Fig. 7.15 Land Suitability map of Mango

### 7.16 Land suitability for Guava (*Psidium guajava*)

Guava is one of the most important fruit crop grown in an area of 0.06 lakh ha in almost all the districts of the State. The crop requirements (Table 7.17) for growing guava were matched with the soil-site characteristics (7.1) and a land suitability map for growing guava was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.16.

An area of about 3 ha (<1%) is moderately suitable (Class S2) for guava and are distributed in the northwestern part of the microwatershed. It has minor limitations of rooting depth and calcareousness. Marginally suitable (Class S3) lands cover an area of about 118 ha (20%) and are distributed in the northern, southern, eastern, western, northeastern and southeastern part of the microwatershed. They have moderate limitations of rooting depth, texture and drainage. A maximum area of about 332 ha (56%) is currently not suitable (Class N1) for growing guava and occur in the major part of the microwatershed with severe limitations of rooting depth and texture.






Fig. 7.16 Land Suitability map of Guava

### 7.17 Land suitability for Sapota (*Manilkara zapota*)

Sapota is one of the most important fruit crop grown in an area of 29373 ha in almost all the districts of the State. The crop requirements (Table 7.18) for growing sapota were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing sapota was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.17.

An area of about 3 ha (<1%) is moderately suitable (Class S2) for sapota and are distributed in the northwestern part of the microwatershed. It has minor limitations of rooting depth and calcareousness. Marginally suitable (Class S3) lands cover an area of about 118 ha (20%) and are distributed in the northern, southern, eastern, western, northeastern and southeastern part of the microwatershed. They have moderate limitations of rooting depth, texture and drainage. A maximum area of about 333 ha (56%) is currently not suitable (Class N1) for growing sapota and occur in the major part of the microwatershed with severe limitation of rooting depth.




Fig. 7.17 Land Suitability map of Sapota

### 7.18 Land Suitability for Pomegranate (*Punica granatum*)

Pomegranate is one of the most important fruit crop commercially grown in about 18488 ha in Karnataka, mainly in Bijapur, Bagalkot, Koppal, Gadag and Chitradurga districts. The crop requirements for growing pomegranate (Table 7.19) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing pomegranate was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.18.

An area of 28 ha (5%) is moderately suitable (Class S2) for growing pomegranate with minor limitations of calcareousness, drainage, texture and rooting depth and are distributed in the northern, eastern and northwestern part of the microwatershed. An area of about 93 ha (16%) is marginally suitable (Class S3) for growing pomegranate and are distributed in the northeastern, southeastern, southern and western part of the microwatershed. They have moderate limitations of rooting depth and texture. A maximum area of about 333 ha (56%) is currently not suitable (Class N1) for growing pomegranate and occur in the major part of the microwatershed with severe limitation of rooting depth.




Fig 7.18 Land Suitability map of Pomegranate

### 7.19 Land Suitability for Musambi (*Citrus limetta*)

Musambi is one of the important fruit crop grown in an area of 3446 ha in almost all the districts of the State. The crop requirements for growing musambi (Table 7.20) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing musambi was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.19.

Highly (Class S1) suitable lands for growing musambi occur in an area of 22 ha (4%) and are distributed in the northern and eastern part of the microwatershed. An area of about 6 ha (1%) is moderately suitable (Class S2) for growing musambi and are distributed in the northern and northwestern part of the microwatershed. They have minor limitations of drainage, rooting depth and calcareousness. Marginally suitable (Class S3) lands occupy an area of about 93 ha (16%) and are distributed in the northeastern, southern, southeastern and western part of the microwatershed. They have moderate limitations of rooting depth and texture. A maximum area of about 333 ha (56%) is currently not suitable (Class N1) for growing musambi and occur in the major part of the microwatershed with severe limitation of rooting depth.




Fig. 7.19 Land Suitability map of Musambi

## 7.20 Land Suitability for Lime (*Citrus sp*)

Lime is one of the most important fruit crop grown in an area of 0.11 lakh ha in almost all the districts of the State. The crop requirements for growing lime (Table 7.21) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing lime was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7. 20.

Highly (Class S1) suitable lands for growing lime occur in an area of 22 ha (4%) and are distributed in the northern and eastern part of the microwatershed. An area of about 6 ha (1%) is moderately suitable (Class S2) for growing lime and are distributed in the northern and northwestern part of the microwatershed. They have minor limitations of drainage, rooting depth and calcareousness. Marginally suitable (Class S3) lands occupy an area of about 93 ha (16%) and are distributed in the northeastern, southern, southeastern and western part of the microwatershed. They have moderate limitations of rooting depth and texture. A maximum area of about 333 ha (56%) is currently not suitable (Class N1) for growing lime and occur in the major part of the microwatershed with severe limitation of rooting depth.




Fig. 7.20 Land Suitability map of Lime

### 7.21 Land Suitability for Amla (*Phyllanthus emblica*)

Amla is one of the medicinal fruit crop grown in almost all the districts of the State. The crop requirements for growing amla (Table 7.22) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing amla was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.21.

A small area of about 3 ha (<1%) is highly suitable (Class S1) for growing amla and is distributed in the northern part of the microwatershed. An area of about 40 ha (7%) has soils that are moderately suitable (Class S2) for growing amla with minor limitations of texture, calcareousness and rooting depth and are distributed in the northern, northeastern, northwestern and southeastern part of the microwatershed. A maximum area of 401 ha (67%) is marginally suitable (Class S3) with moderate limitations of rooting depth and texture and are distributed in the major part of the microwatershed. An area of about 9 ha (2%) is currently not suitable (Class N1) and are distributed in the eastern part of the microwatershed with severe limitation of rooting depth.




Fig. 7.21 Land Suitability map of Amla

## 7.22 Land Suitability for Cashew (*Anacardium occidentale*)

Cashew is one of the most important plantation nut crop grown in an area of 0.7 lakh ha in almost all the districts of the State. The crop requirements for growing cashew (Table 7.23) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing cashew was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.22.

The marginally suitable (Class S3) lands cover an area of about 78 ha (13%) and occur in the southern and western part of the microwatershed. It has moderate limitations of rooting depth and texture. A maximum area of about 375 ha (63%) is currently not suitable (Class N1) and are distributed in all parts of the microwatershed with severe limitations of rooting depth, texture and calcareousness.




Fig. 7.22 Land Suitability map of Cashew

### 7. 23 Land Suitability for Jackfruit (*Artocarpus heterophyllus*)

Jackfruit is one of the most important fruit crop grown in an area of 5368 ha in almost all the districts of the State. The crop requirements for growing jackfruit (Table 7.24) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing jackfruit was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.23.

A small area of about 3 ha (<1%) has soils that are moderately suitable (Class S2) for growing jackfruit with minor limitations of calcareousness and rooting depth and are distributed in the northwestern part of the microwatershed. Marginally suitable (Class S3) lands for growing jackfruit occupy an area of about 118 ha (20%) and are distributed in the northern, northeastern, southern, southeastern and western part of the microwatershed. They have moderate limitations of rooting depth, texture and drainage. A maximum area of about 332 ha (56%) is currently not suitable (Class N1) and are distributed in the major part of the microwatershed with severe limitations of rooting depth and texture.




Fig. 7.23 Land Suitability map of Jackfruit

## 7.24 Land Suitability for Jamun (*Syzygium cumini*)

Jamun is an important fruit crop grown in almost all the districts of the State. The crop requirements for growing jamun (Table 7.25) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing jamun was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.24.

An area of about 25 ha (4%) has soils that are moderately suitable (Class S2) for growing jamun with minor limitations of texture and drainage and are distributed in the northern and eastern part of the microwatershed. Marginally suitable (Class S3) lands for growing jamun occupy an area of about 96 ha (16%) and are distributed in the northern, northeastern, southern, southeastern and western part of the microwatershed. They have moderate limitations of rooting depth, texture and drainage. A maximum area of about 332 ha (56%) is currently not suitable (Class N1) and are distributed in the major part of the microwatershed with severe limitations of rooting depth and texture.






Fig. 7.24 Land Suitability map of Jamun

## 7.25 Land Suitability for Custard Apple (*Annona reticulata*)

Custard apple is one of the most important fruit crop grown in almost all the districts of the State. The crop requirements for growing custard apple (Table 7.26) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing custard apple was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.25.

An area of 27 ha (5%) is highly suitable (Class S1) for growing custard apple and are distributed in the northern, northwestern and eastern part of the microwatershed. An area of about 15 ha (3%) has soils that are moderately suitable (Class S2) for growing custard apple with minor limitation of rooting depth and are distributed in the northeastern and southeastern part of the microwatershed. A maximum area of about 401 ha (67%) is marginally suitable (Class S3) for growing custard apple and are distributed in the major part of the microwatershed with moderate limitations of rooting depth and texture. An area of about 9 ha (2%) is currently not suitable (Class N1) and are distributed in the eastern part of the microwatershed with severe limitation of rooting depth.




Fig. 7.25 Land Suitability map of Custard Apple

## 7.26 Land Suitability for Tamarind (*Tamarindus indica*)

Tamarind is one of the most important spice crop grown in almost all the districts of the state. The crop requirements for growing tamarind (Table 7.27) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing tamarind was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Fig. 7.26.

An area of about 25 ha (4%) is moderately suitable (Class S2) for growing tamarind and are distributed in the northern and eastern part of the microwatershed. They have minor limitations of texture and drainage. Marginally suitable (Class S3) lands for growing tamarind occupy a small area of about 3 ha (<1%) and are distributed in the northwestern part of the microwatershed. It has moderate limitations of rooting depth and calcareousness. A maximum area of about 426 ha (71%) is currently not suitable (Class N1) for growing tamarind and occur in all parts of the microwatershed with severe limitations of rooting depth and texture.




Fig. 7.26 Land Suitability map of Tamarind

### 7.27 Land Suitability for Mulberry (*Morus nigra*)

Mulberry is an important leaf crop grown for rearing silkworms in about 1.6 lakh ha area in all the districts of the state. The crop requirements for growing mulberry (Table 7.28) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing mulberry was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.27.

A small area of about 3 ha (<1%) has soils that are moderately suitable (Class S2) for growing mulberry with minor limitations of calcareousness and rooting depth and are distributed in the northwestern part of the microwatershed. Marginally suitable (Class S3) lands for growing mulberry occupy an area of about 117 ha (20%) and are distributed in the northern, northeastern, southern, southeastern and western part of the microwatershed. They have moderate limitations of rooting depth, texture and drainage. A maximum area of about 332 ha (56%) is currently not suitable (Class N1) and are distributed in the major part of the microwatershed with severe limitations of rooting depth and texture.




Fig 7.27 Land Suitability map of Mulberry

### 7.28 Land suitability for Marigold (*Tagetes sps.*)

Marigold is one of the most important flower crop grown in an area of 9108 ha in almost all the districts of the State. The crop requirements (Table 7.29) for growing marigold were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing marigold was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.28.

An area of about 42 ha (7%) is moderately suitable (Class S2) for growing marigold and are distributed in the northern, northeastern, northwestern, eastern and southeastern part of the microwatershed. They have minor limitations of texture, drainage, rooting depth and calcareousness. Marginally suitable (Class S3) lands for growing marigold occupy a maximum area of about 401 ha (67%) and are distributed in the major part of the microwatershed. They have moderate limitations of texture and rooting depth. An area of about 9 ha (2%) is currently not suitable (Class N1) and are distributed in the eastern part of the microwatershed with severe limitation of rooting depth.




Fig. 7.28 Land Suitability map of Marigold

### 7.29 Land Suitability for Chrysanthemum (*Dendranthema grandiflora*)

Chrysanthemum is one of the most important flower crop grown in an area of 4978 ha in almost all the districts of the State. The crop requirements (Table 7.30) for growing chrysanthemum were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing chrysanthemum was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.29.

An area of about 42 ha (7%) is moderately suitable (Class S2) for growing chrysanthemum and are distributed in the northern, northeastern, northwestern, eastern and southeastern part of the microwatershed. They have minor limitations of texture, drainage, rooting depth and calcareousness. Marginally suitable (Class S3) lands for growing chrysanthemum occupy a maximum area of about 401 ha (67%) and are distributed in the major part of the microwatershed. They have moderate limitations of texture and rooting depth. An area of about 9 ha (2%) is currently not suitable (Class N1) and are distributed in the eastern part of the microwatershed with severe limitation of rooting depth.




Fig. 7.29 Land Suitability map of Chrysanthemum

**Table 7.1 Soil-Site Characteristics of Mundragi-3 Microwatershed**

Soil Map Units	Climate (P) (mm)	Growing period (Days)	Drainage Class	Soil depth (cm)	Soil texture		Gravelliness		AWC (mm/m)	Slope (%)	Erosion	pH	EC (dSm <sup>-1</sup> )	ESP (%)	CEC [Cmol (p <sup>+</sup> )kg <sup>-1</sup> ]	BS (%)
					Sur-face	Sub-surface	Surface (%)	Sub-surface (%)								
BDLiB2	866	150	WD	25-50	sc	sl	<15	<15	<50	1-3	moderate	6.20	0.074	0.20	4.20	93
BDLhB2g1	866	150	WD	25-50	scl	sl	15-35	<15	<50	1-3	moderate	6.20	0.074	0.20	4.20	93
SBRcB2	866	150	Sed	50-75	sl	ls	<15	<15	<50	1-3	moderate	8.24	0.145	1.15	7.50	100
JNKcB2	866	150	WD	50-75	sl	scl	<15	<15	51-100	1-3	moderate	8.42	0.148	0.18	14.50	100
JNKiB2	866	150	WD	50-75	sc	scl	<15	<15	51-100	1-3	moderate	8.42	0.148	0.18	14.50	100
HSLcB2	866	150	MW	75-100	sl	sc	<15	<15	101-150	1-3	moderate	7.16	0.117	5.94	4.90	97
NGPmB2	866	150	MW	100-150	c	c	<15	<15	>200	1-3	moderate	7.42	0.24	0.22	67.10	100
BDPcB2	866	150	WD	<25	sl	scl	<15	<15	<50	1-3	moderate	8.58	0.262	0.35	18.10	100
HTKcB2	866	150	WD	25-50	sl	sl	<15	10-25	<50	1-3	moderate	6.81	0.062	0.38	3	101
TMKhA1	866	150	MW	>150	scl	c	<15	<15	>200	0-1	slight	9.60	0.35	6.63	21.83	100

\*Symbols and abbreviations are according to Field Guide for LRI under Sujala-III Project, Karnataka

**Table 7.2 Land suitability criteria for Sorghum**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	26–30	30–34; 24–26	34–40; 20–24	>40; <20
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, c (red), c (black)	scl, cl	ls, sl	-
	pH	1:2.5	5.5-7.8	5.0-5.5 7.8-9.0	>9.0	-
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	10-15
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	0-3	3-5	5-10	>10



**Table 7.3 Land suitability criteria for Maize**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	30-34	35-38 26-30	38-40 26-20	
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc	c (red), c (black)	ls, sl	-
	pH	1:2.5	5.5-7.8	5.0-5.5 7.8-9.0	>9.0	-
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	-
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

**Table 7.4 Land suitability criteria for Bajra**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-32	33-38 24-27	39-40 20-23	<20
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm	500-750	400-500	200-400	<200
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl, scl, cl,sc,c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.8	5.0-5.5 7.8-9.0	5.5-6.0 >9.0	
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	15-35	35-60	>60	
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	1-3	3-5	5-10	>10

**Table 7.5 Land suitability criteria for Groundnut**

Land use requirement			Rating			
Soil –site characteristics	Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)	
Climatic regime	Mean temperature in growing season	°C	24–33	22–24; 33–35	20–22; 35–40	<20; >40
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. Well drained	Poorly drained	Very Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl	sl,cl, sc	c (red), c (black), ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<35	35-60	>60	
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.6 Land suitability criteria for Sunflower**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	24–30	30–34; 20–24	34–38; 16–20	>38; <16
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	mod. Well drained	-	Poorly to very drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	cl, sc,c (red), c (black)	scl	ls, sl	-
	pH	1:2.5	6.5-7.8	7.8-8.4 5.5-6.5	8.4-9.0; 5.0-5.5	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.7 Land suitability criteria for Redgram**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	30-35(G) 20-25(AV) 15-18 (F&PS) 35-40(M)	25-30(G) 20-25 (AV) 12-15 (F&PS) 30-35(M)	20-25(G) 15-20(AV) 10-12 (F&PS) 25-30(M)	< 20 <15 <10 <25
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. Well drained	Poorly drained	Very Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, c (red)	c (black),sl, scl, cl	ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-9.0	5.0-5.5 >9.0	-
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-50	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<1.0	1.0-2.0	>2.0	
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.8 Land suitability criteria for Bengal gram**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	20–25	25–30; 15–20	30–35; 10–15	>35; <10
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. Well drained	Poorly drained	Very Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	c(black)	-	c (red), scl, cl, sc	ls, sl
	pH	1:2.5	6.0-7.8	5.0-6.0 7.8-9.0	>9.0	-
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	-
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.9 Land suitability criteria for Cotton**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	22-32	>32	<19	-
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well to moderately well	Poorly drained/Some what excessively drained	-	very poorly/excessively drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, c (red,black)	cl	scl	ls, sl
	pH	1:2.5	6.5-7.8	7.8-8.4	5.5-6.5 8.4->9.0	<5.5
	CEC	C mol (p+)Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	50-100	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	<3	3-5	-	>5

**Table 7.10 Land suitability criteria for Chilli**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	25-32	33-35 20-25	35-38 <20	>38
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc	c (black), sl	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10



**Table 7.11 Land suitability criteria for Tomato**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	25-28	29-32 20-24	15-19 33-36	<15 >36
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl, scl, cl, sc, c (red)	-	ls, c(black)	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.12 Land suitability criteria for Brinjal**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	Well drained	Moderately well drained	Poorly drained	V. Poorly drained
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class				
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl, scl, cl, sc c (red)	-	ls, c (black)	-
	pH	1:2.5	6.0-7.3	7.3-8.4 5.0-6.0	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	>60
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.13 Land suitability criteria for Onion**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	20-30	30-35	35-40	>40
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately /imperfectly	-	Poorly to V poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl,scl,cl,sc,c (red)	-	c (Black),ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<1.0	1.0-2.0	2.0-4.0	<4
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.14 Land suitability criteria for Bhendi**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	25-28	29-32 20-24	15-19 33-36	<15 >36
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Imperfectly drained	Poorly to very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl,sc, c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.15 Land suitability criteria for Drumstick**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, scl, cl, c (red)	sl, c (black)	ls	s
	pH	1:2.5	6.0-7.3	5.0-5.5 7.3-7.8	5.5-6.0 7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<35	35-60	60-80	>80
Soil toxicity	Salinity (EC saturation extract)	ds/m				
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-10	-	>10

**Table 7.16 Land suitability criteria for Mango**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-32	24-27 33-35	36-40	20-24
	Min temp. before flowering	°C	10-15	15-22	>22	-
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration	Days				
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V. Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	-	ls, sl, c (black)	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>150	100-150	75-100	<75
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.17 Land suitability criteria for Guava**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-32	33-36 24-27	37-42 20-23	
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	sl	c (black), ls	-
	pH	1:2.5	6.0-7.8	5.0-6.0	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.18 Land suitability criteria for Sapota**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-32	33-36 24-27	37-42 20-23	>42 <18
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	-	Poorly to very drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	sl	ls, c (black)	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10



**Table 7.19 Land suitability criteria for Pomegranate**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	30-34	35-38 25-29	39-40 15-24	
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl,cl, sc, c (red)	c (black),sl	ls	-
	pH	1:2.5	5.5-7.8	7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.20 Land suitability criteria for Musambi**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-30	31-35 24-27	36-40 20-23	>40 <20
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately drained	poorly	Very poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c	sl	ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.21 Land suitability criteria for Lime**

Land use requirement		Rating				
Soil –site characteristics	Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)	
Climatic regime	Mean temperature in growing season	°C	28-30	31-35 24-27	36-40 20-23	>40 <20
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately drained	poorly	Very poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c	sl	ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.22 Land suitability criteria for Amla**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. well drained	Poorly drained	V. Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	c (black)	ls, sl	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15-35	35-60	60-80	-
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

**Table 7.23 Land suitability criteria for Cashew**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	32 to 34	28 to 32; 34 to 38	24 to 28; 38 to 40	<20; >40
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	-	sl, ls	c (black)
	pH	1:2.5	5.5-6.5	5.0-5.5 6.5-7.3	7.3-7.8	>7.8
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-10	>10	-

**Table 7.24 Land suitability criteria for Jackfruit**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. well	Poorly	V. Poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	-	sl, ls, c (black)	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	>60
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10-

**Table 7.25 Land suitability criteria for Jamun**

Land use requirement		Rating				
Soil –site characteristics	Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)	
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well	Mod. well	Poorly	V.Poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c(red)	sl, c (black)	ls	-
	pH	1:2.5	6.0-7.8	5.0-6.0	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>150	100-150	50-100	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	>60
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

**Table 7.26 Land suitability criteria for Custard apple**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	Scl, cl, sc, c (red), c (black)	-	Sl, ls	-
	pH	1:2.5	6.0-7.3	5.5-6.0 7.3-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15-35	35-60	60-80	-
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	>5	-



**Table 7.27 Land suitability criteria for Tamarind**

Land use requirement		Rating				
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod.well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl,sc, c (red)	sl, c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>150	100-150	75-100	<75
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

**Table 7.28 Land suitability criteria for Mulberry**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	24–28	22–24; 28–32	32–38; 22–18	>38; <18
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V. Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, cl, scl	c (red)	c (black), sl, ls	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.8-8.4	7.3-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	0-35	35-60	60-80	>80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

**Table 7.29 Land suitability criteria for Marigold**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	18-23	17-15 24-35	35-40 10-14	>40 <10
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl,scl, cl, sc, c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%				
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.30 Land suitability criteria for Chrysanthemum**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	18-23	17-15 24-35	35-40 10-14	>40 <10
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl,scl, cl, sc, c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%				
Erosion hazard	Slope	%	<3	3-5	5-10	>10

### 7.30 Land Management Units (LMUs)

The 10 soil map units identified in Mundragi-3 microwatershed have been grouped into 4 Land Management Units (LMU's) for the purpose of preparing a Proposed Crop Plan. Land Management Units are grouped based on the similarities in respect of the type of soil, the depth of the soil, the surface soil texture, gravel content, AWC, slope, erosion etc. and a Land Management Units map (Fig. 7.30) has been generated. These Land Management Units are expected to behave similarly for a given level of management.

The map units that have been grouped into 4 Land Management Units along with brief description of soil and site characteristics are given below.

LUC No.	Soil map units	Soil and site characteristics
1	103.TMKhA1	Very deep (> 150 cm), lowland clay soils, 0-1 % slopes, non-gravelly (<15%), slight erosion.
2	32.HSLcB2 49.NGPmB2	Moderately deep to deep (75 to 150 cm), black clay soils, 1-3 % slopes, non-gravelly (<15%), moderate erosion.
3	11.SBRcB2 20.JNKcB2 22.JNKiB2	Moderately shallow (50 to 75 cm), sandy clay loam to loamy sand soils, 1-3% slopes, non-gravelly (<15%), moderate erosion.
4	5.BDLiB2 118.BDPcB2 162.BDLhB2g1 165.HTKcB2	Shallow (25 to 50 cm), sandy loam soils, 1-3 % slopes, gravelly to non-gravelly (0 to 35%), moderate erosion.




Fig. 7.30 Land Management Units Map- Mundragi-3 Microwatershed

### **7.31 Proposed Crop Plan for Mundragi-3 Microwatershed**

After assessing the land suitability for the 29 crops, the Proposed Crop Plan has been prepared for the 4 identified LMUs by considering only highly (Class S1) and moderately (Class S2) suitable lands for each of the 29 crops. The resultant proposed crop plan is presented below in Table 7.31.

**Table 7.31 Proposed Crop Plan for Mundragi-3 Microwatershed**

<b>LMU</b>	<b>Soil Map Units</b>	<b>Survey Number</b>	<b>Field Crops/ Commercial crops</b>	<b>Horticulture Crops (Rainfed/Irrigated )</b>	<b>Suitable Interventions</b>
1	103.TMKhA1 (Very deep, lowland clay soils)	<b>Mundaragi :17</b>	Sorghum, maize, Bajra	<b>Fruit crops:</b> Custard Apple, Amla, Ber, Aonla <b>Vegetable crops:</b> Brinjal, Tomato, Chillies, Drumstick, Coriander	Providing proper drainage, addition of organic manures, green leaf manuring, suitable conservation practices
2	32.HSLcB2 49.NGPmB2 (Moderately deep to deep black clay soils)	<b>Mundaragi:</b> 14,15,16,23,24,516,517,5 18/1,522/1,610/2,613/1,613/2,614/1,6 15/1, 615/2	Sunflower, Sorghum, Maize, Soybean, Cotton, Bengal gram, Safflower, Linseed, Bajra	<b>Fruit crops:</b> Pomegranate, Lime, Musambi, Tamarind, Jamun, Amla, Custard apple <b>Vegetables:</b> Drumstick, Chilli, Bhendi, Cluster bean, Coriander <b>Flowers:</b> Marigold, Chrysanthemum	Application of FYM, Biofertilizers and micronutrients, drip irrigation, Mulching, suitable soil and water conservation practices
3	11.SBRcB2 20.JNKcB2 22.JNKiB2 (Moderately shallow, sandy clay loam to loamy sand soils)	<b>Mundaragi :</b> 454,455,456,508 <b>Varkanahalli :</b> 183 <b>Yadagiri.B:</b> 30,31,33,34,35,36,39, 40,41,43,44,45,46,47,48,49,52	Sorghum, Bajra, Groundnut	<b>Fruit crops:</b> Amla, Custard apple <b>Vegetables:</b> Coriander, Bhendi <b>Flowers:</b> Marigold, Jasmine Chrysanthemum	Application of FYM, Biofertilizers and micronutrients, drip irrigation, mulching, suitable soil and water conservation practices
4	5.BDLiB2 118.BDPcB2 162.BDLhB2g1 165.HTKcB2 (Shallow, sandy loam soils)	<b>Mundaragi:</b> 13,450,451,484,486,487, 488,489,491,492,505,509,510,511,512 ,513,514,515,523,524,525,526,527,52 8,529,530,531,532,533,534,535,536,5 37,538,539,540,543/1,543/2,543/3,544 ,545,546,547,548,549,550,553/1,553/2 ,557,565,572,584,585,586,587,588,58 9,590/1,590/2,591,592,593,594/1,594/ 2,596,597,599,600,601,602,603,604,6 05,607,608,610/1,611,612,613/3,614/2	-	<b>Agri-Silvi-Pasture:</b> Hybrid Napier, <i>Styloxanthes hamata</i> , <i>Styloxanthes scabra</i>	Use of short duration varieties, sowing across the slope, drip irrigation and mulching is recommended





## **SOIL HEALTH MANAGEMENT**

### **8.1 Soil Health**

Soil health is basic to plant health and plant health is basic to human and bovine health. Soil is fundamental to crop production. Without soil, no food could be produced nor would livestock be fed on a large scale. Because it is finite and fragile, soil is a precious resource that requires special care from its users.

Soil health or the capacity of the soil to function is critical to human survival. Soil health has been defined as: “the capacity of the soil to function as a living system without adverse effect on the ecosystem”. Healthy soils maintain a diverse community of soil organisms that help to form beneficial symbiotic associations with plant roots, recycle essential plant nutrients, improve soil structure with positive repercussions for soil, water and nutrient holding capacity and ultimately improve crop production and also contribute to mitigating climate change by maintaining or increasing its carbon content.

Functional interactions of soil biota with organic and inorganic components, air and water determine a soil’s potential to store and release nutrients and water to plants and to promote and sustain plant growth. Thus, maintaining soil health is vital to crop production and conserve soil resource base for sustaining agriculture.

#### **The most important characteristics of a healthy soil are**

- Good soil tilth
- Sufficient soil depth
- Good water storage and good drainage
- Adequate supply, but not excess of nutrients
- Large population of beneficial organisms
- Small proportion of plant pathogens and insect pests
- Low weed pressure
- Free of chemicals and toxins that may harm the crop
- Resistance to degradation
- Resilience when unfavorable conditions occur

#### **Characteristics of Mundragi-3 Microwatershed**

- ❖ The soil phases identified in the microwatershed belonged to the soil series of BDL 245 ha (41%), SBR 78 ha (13%), JNK 15 ha (3%), HSL 3 ha (<1%), NGP 22 ha (4%), BDP 9 ha (2%), HTK 78 ha (13%) and TMK 3 ha (<1%).
- ❖ As per land capability classification entire area of the microwatershed falls under arable land category (Class II, III & IV). The major limitations identified in the arable lands were soil erosion and soil limitation.
- ❖ On the basis of soil reaction, 193 ha (32%) is neutral (pH 6.5 -7.3) and 260 ha (44%) area is slightly alkaline (pH 7.3-7.8).

## **Soil Health Management**

The following actions are required to improve the current land husbandry practices that provide a sound basis for the successful adoption of sustainable crop production system.

### **Acid soils**

1. Growing of crops suitable for particular soil pH.
2. Ameliorating the soils through the application of amendments (liming materials).

Liming materials:

1.  $\text{CaCO}_3$  (Calcium Carbonate).
2. Dolomite [ $\text{Ca Mg} (\text{CO}_3)_2$ ]
3. Quick lime (CaO)
4. Slaked lime [ $\text{Ca} (\text{OH})_2$ ]

For normal pH and pH 4.8 (35 t/ha) and pH 6.0-7.0 (4 t/ha) lime is required.

### **Alkaline soils**

Slightly alkaline soils cover about 260 ha area.

1. Regular addition of organic manure, green manuring, green leaf manuring, crop residue incorporation and mulching needs to be taken up to improve the soil organic matter status.
2. Application of biofertilizers (Azospirillum, Azotobacter, Rhizobium).
3. Application of 25% extra N and P (125 % RDN&P).
4. Application of  $\text{ZnSO}_4$  – 12.5 kg/ha (once in three years).
5. Application of Boron – 5kg/ha (once in three years).

### **Neutral soils**

Neutral soils cover about 193 ha area in the microwatershed.

1. Regular addition of organic manure, green manuring, green leaf manuring, crop residue incorporation and mulching needs to be taken up to improve the soil organic matter status.
2. Application of biofertilizers, (Azospirillum, Azotobacter, Rhizobium).
3. Application of 100 per cent RDF.
4. Need based micronutrient applications.

Besides the above recommendations, the best transfer of technology options are also to be adopted.

### **Soil Degradation**

Soil erosion is one of the major factor affecting the soil health in the microwatershed. Out of total 598 ha area in the microwatershed, an area of about 450 ha is suffering from moderate erosion. These areas need immediate soil and water

conservation and, other land development and land husbandry practices for restoring soil health.

### **Dissemination of Information and Communication of Benefits**

Any large scale implementation of soil health management requires that supporting information is made available widely, particularly through channels familiar to farmers and extension workers. Given the very high priority attached to soil-health especially by the Central Government on issuing Soil-Health Cards to all the farmers, media outlets like Regional, State and National Newspapers, Radio and Dooradarshan programs in local languages but also modern information and communication technologies such as Cellular phones and the Internet, which can be much more effective in reaching the younger farmers.

### **Inputs for Net Planning (Saturation Plan) and Interventions needed**

Net planning (Saturation Plan) in IWMP is focusing on preparation of

1. Soil and Water Conservation Plan for each plot or farm.
2. Productivity enhancement measures/ interventions for existing crops/livestock/other farm enterprises.
3. Diversification of farming mainly with perennial horticultural crops and livestock.
4. Improving livelihood opportunities and income generating activities.

In this connection, how various outputs of Sujala-III are of use in addressing these objectives of Net Planning (Saturation Plan) are briefly presented below.

- ❖ **Soil Depth:** The depth of a soil decides the amount of moisture and nutrients it can hold, what crops can be taken up or not, depending on the rooting depth and the length of growing period available for raising any crop. Deeper the soil, better for a wide variety of crops. If sufficient depth is not available for growing deep rooted crops, either choose medium or short duration crops or deeper planting pits need to be opened and additional good quality soil brought from outside has to be filled into the planting pits.
- ❖ **Surface Soil Texture:** Lighter soil texture in the top soil means, better rain water infiltration, less run-off and soil moisture conservation, less capillary rise and less evaporation losses. Lighter surface textured soils are amenable to good soil tilth and are highly suitable for crops like groundnut, root vegetables (carrot, raddish, potato etc) but not ideal for crops that need stagnant water like lowland paddy. Heavy textured soils are poor in water infiltration and percolation. They are prone for sheet erosion; such soils can be improved by sand mulching. The technology that is developed by the AICRP-Dryland Agriculture, Vijayapura, Karnataka can be adopted.
- ❖ **Gravelliness:** More gravel content is favorable for run-off harvesting but poor in soil moisture storage and nutrient availability. It is a significant parameter that decides the kind of crop to be raised.

- ❖ **Land Capability Classification:** The land capability map shows the areas suitable and not suitable for agriculture and the major constraints in each of the plot/survey number. Hence, one can decide what kind of enterprise is possible in each of these units. In general, erosion, wetness and soil are the major constraints in Mundragi-3 microwatershed.
- ❖ **Organic Carbon:** The OC content (an index of available Nitrogen) is medium (0.5-0.75%) in 453 ha (76%) and high (>0.75%) in less than 1 ha (<1%). The areas that are medium in OC needs to be further improved by applying farmyard manure and rotating crops with cereals and legumes or mixed cropping.
- ❖ **Promoting green manuring:** Growing of green manuring crops costs Rs. 1250/ha (green manuring seeds) and about Rs. 2000/ha towards cultivation that totals to Rs. 3250/- per ha. On the other hand, application of organic manure @ 10 tons/ha costs Rs. 5000/ha. The practice needs to be continued for 2-3 years or more. Nitrogen fertilizer needs to be supplemented by 25% in addition to the recommended level in 453 ha area where OC is medium (0.5-0.75%). For example, for rainfed maize, recommended level is 50 kg N per ha and an additional 12 kg /ha needs to be applied for all the crops grown in these plots.
- ❖ **Available Phosphorus:** Available Phosphorus is low (<23 kg/ha) in an area of 158 ha (26%) and medium (23-57 kg/ha) in 295 ha (49%) of the microwatershed. For all the crops 25% additional P needs to be applied where available P is low and medium.
- ❖ **Available Potassium:** Available potassium is low (<145 kg/ha) in an area of about 52 ha (9%) and medium (145-337 kg/ha) in an area of 401 ha (67%) of the microwatershed. All the plots, where available potassium is low and medium, for all the crops, additional 25 % potassium may be applied.
- ❖ **Available Sulphur:** Available sulphur is a very critical nutrient for oilseed crops. It is medium in 396 ha (66%) and low in 57 ha (9%). Low and medium areas need to be applied with magnesium sulphate or gypsum or Factamphos (p) fertilizer (13% sulphur) for 2-3 years for the deficiency to be corrected.
- ❖ **Available Boron:** Entire area of 453 ha (76%) is low (<0.5 ppm) in available boron content. For these areas, application of sodium borate @ 10 kg/ha as soil application or 0.2 % borax as foliar spray is recommended.
- ❖ **Available Iron:** Entire area of 453 ha (76%) is sufficient in available iron in the microwatershed. For the deficient areas, iron sulphate @ 25 kg/ha need to be applied for 2-3 years.
- ❖ **Available Zinc:** An area of 318 ha (53%) in the microwatershed is deficient (<0.6 ppm) and 135 ha (23%) is sufficient (>0.6 ppm) in available zinc content. Application of zinc sulphate @25 kg/ha is recommended for the deficient areas.
- ❖ **Soil Alkalinity:** The microwatershed has 260 ha (44%) area under slightly alkaline. These areas need application of gypsum and wherever calcium is in excess, iron pyrites and element sulphur can be recommended. Management practices like treating

repeatedly with good quality water to drain out the excess salts and provision of subsurface drainage and growing of salt tolerant crops like Casuarina, Acacia, Neem, Ber etc, are recommended.


- ❖ **Land Suitability for various crops:** Areas that are highly, moderately and marginally suitable for growing various crops are indicated. Along with the suitability, various constraints that are limiting the productivity are also indicated. For example, in case of cotton, gravel content, rooting depth and salinity/alkalinity are the major constraints in various plots. With suitable management interventions, the productivity can be enhanced. In order to increase the water holding capacity of light textured soils, growing of green manure crops and application of organic manure is recommended.



## SOIL AND WATER CONSERVATION TREATMENT PLAN

For preparing soil and water conservation treatment plan for Mundragi-3 microwatershed, the land resource inventory database generated under Sujala-III project has been transformed as information through series of interpretative (thematic) maps using soil phase map as a base. The various thematic maps (1:7920 scale) generated were

- Soil depth
- Surface soil texture
- Available water capacity
- Soil slope
- Soil gravelliness
- Land capability
- Present land use and land cover
- Crop suitability
- Rainfall
- Hydrology
- Water Resources
- Socio-economic data
- Contour plan with existing features- network of waterways, pothissa boundaries, cut up/ minor terraces etc.
- Cadastral map (1:7920 scale)
- Satellite imagery (1:7920 scale)



Apart from these, Hand Level/ Hydro Marker/ Dumpy Level/ Total Station and Kathedars' List to be collected.

### Steps for Survey and Preparation of Treatment Plan

The boundaries of Land User Groups' and Survey No. boundaries are traced in the field.

- Naming of user groups and farmers
- Identification of arable and non arable lands
- Identification of drainage lines and gullies
- Identification of non treatable areas
- Identification of priority areas in the arable lands
- Treatment plan for arable lands
- Location of water harvesting and recharge structures

## 9.1 Treatment Plan

The treatment plan recommended for arable lands is briefly described below

### 9.1.1 Arable Land Treatment

#### A. BUNDING

Steps for Survey and Preparation of Treatment Plan		<p><b>USER GROUP-1</b></p> <p>CLASSIFICATION OF GULLIES</p> <p>ಕೊರಕಾಲಿನ ವರ್ಗೀಕರಣ</p>
<ul style="list-style-type: none"> <li>Cadastral map (1:7920 scale) is enlarged to a scale of 1:2500 scale</li> <li>Existing network of waterways, pothissa boundaries, grass belts, natural drainage lines/ watercourse, cut ups/ terraces are marked on the cadastral map to the scale</li> <li>Drainage lines are demarcated into</li> </ul>		
Small gullies	(up to 5 ha catchment)	
Medium gullies	(5-15 ha catchment)	
Ravines	(15-25 ha catchment) and	
<i>Halla/Nala</i>	(more than 25ha catchment)	

#### Measurement of Land Slope

Land slope is estimated or determined by the study and interpretation of contours or by measurement in the field using simple instruments like Hand Level or Hydromarker.



Vertical and Horizontal intervals between bunds as recommended by the Watershed Development Department.

Slope percentage	Vertical interval (m)	Corresponding Horizontal Distance (m)
2 - 3%	0.6	24
3 - 4%	0.9	21
4 - 5%	0.9	21
5 - 6%	1.2	21
6 - 7%	1.2	21

**Note:** (i) The above intervals are maximum.

(ii) Considering the slope class and erosion status (A1... A=0-1 % slope, 1= slight erosion) the intervals have to be decided.



**Bund length recording:** Considering the contour plan and the existing grass belts/partitions, the bunds are aligned and lengths are measured.

**Section of the Bund**

Bund section is decided considering the soil texture class and gravelliness class (bg<sub>0</sub>... b=loamy sand, g<sub>0</sub> = <15% gravel). The recommended Sections for different soils are given below.

**Recommended Bund Section**

Top width (m)	Base width (m)	Height (m)	Side slope (Z:1;H:V)	Cross section (sq m)	Soil Texture	Remarks
0.3	0.9	0.3	01:01	0.18	Sandy loam	Vegetative bund
0.3	1.2	0.3	1.5:1	0.225	Sandy clay	
0.3	1.2	0.5	0.9:1	0.375	Red gravelly soils	
0.3	1.2	0.6	0.75:1	0.45		
0.3	1.5	0.6	01:01	0.54	Red sandy loam	
0.3	2.1	0.6	1.5:1	0.72	Very shallow black soils	
0.45	2	0.75	01:01	0.92		
0.45	2.4	0.75	1.3:1	1.07	Shallow black soils	
0.6	3.1	0.7	1.78:1	1.29	Medium black soils	
0.5	3	0.85	1.47:1	1.49		

**Formation of Trench cum Bund**

Dimensions of the Borrow Pits/Trenches to be excavated (machinery are decided considering the Bund Section).

Details of Borrow Pit dimensions are given below:

**TRENCH CUM BUND**

**'A' FRAME FOR INTERBUND MANAGEMENT**

**Size of Borrow Pits/ Trench recommended for Trench cum Bund (by machinery)**

Bund section	Bund length	Earth quantity	Pit				Berm (pit to pit)	Soil depth class
			L(m)	W(m)	D(m)	Quantity (m <sup>3</sup> )		
m <sup>2</sup>	m	m <sup>3</sup>					m	
0.375	6	2.25	5.85	0.85	0.45	2.24	0.15	Shallow
0.45	6	2.7	5.4	1.2	0.43	2.79	0.6	Shallow
0.45	6	2.7	5	0.85	0.65	2.76	1	Moderately Shallow
0.54	5.6	3.02	5.5	0.85	0.7	3.27	0.1	Moderately shallow
0.54	5.5	2.97	5	1.2	0.5	3	0.5	Shallow
0.72	6.2	4.46	6	1.2	0.7	5.04	0.2	Moderately shallow
0.72	5.2	3.74	5.1	0.85	0.9	3.9	0.1	Moderately deep

**B. Water Ways**

1. Existing waterways are marked on the cadastral map (1:792 scale) and their dimensions are recorded.
2. Considering the contour plan of the MWS, additional waterways/ modernization of the existing ones can be thought of.
3. The design details are given in the Manual.

**C. Farm Ponds**

Waterways and the catchment area will give an indication on the size of the Farm Pond. Location of the pond can be decided based on the contour plan/ field condition and farmers' need/desire.

**D. Diversion Channel**

Existing EPT/ CPT are marked on the cadastral map. Looking to the need, these can be modernized or fresh diversion channel can be proposed and runoff from this can be stored in Gokatte/ Recharge ponds.

**9.1.2 Non-Arable Land Treatment**

Depending on the gravelliness and crops preferred by the farmers, the concerned authorities can decide appropriate treatment plan. The recommended treatments may be Contour Trench, Staggered Trench, Crescent Bund, Boulder Bund or Pebble Bund.

### 9.1.3 Treatment of Natural Water Course/ Drainage Lines

- a) The cadastral map has to be updated as regards the network of drainage lines (gullies/ *nalas*/ *hallas*) and existing structures are marked to the scale and storage capacity of the existing water bodies are documented.
- b) The drainage line will be demarcated into Upper Reach, Middle Reach and Lower Reach.
- c) Considering the Catchment, *Nala* bed and bank conditions, suitable structures are decided.
- d) Number of storage structures (Check dam/ *Nala* bund/ Percolation tank) will be decided considering the commitments and available runoff from water budgeting and quality of water in the wells and site suitability.
- e) Detailed Levelling Survey using Dumpy Level / Total Station has to be carried out to arrive at the site-specific designs as shown in the Manual.
- f) The location of ground water recharge structures are decided by examining the lineaments and fracture zones from geological maps.
- g) Rainfall intensity data of the nearest Rain Gauge Station is considered for Hydrologic Designs.
- h) Silt load to the Storage/Recharge structures is reduced by providing vegetative, boulder and earthen checks in the natural water course. Location and design details are given in the Manual.

### 9.2 Recommended Soil and Water Conservation Measures

The appropriate conservation structures best suited for each of the land parcel/ survey number (Appendix-I) are selected based on the slope per cent, severity of erosion, amount of rainfall, land use and soil type. The different kinds of conservation structures recommended are:

1. Graded / Strengthening of Bunds
2. Trench cum Bunds (TCB)
3. Trench cum Bunds / Strengthening
4. Crescent Bunds

A map (Fig. 9.1) showing soil and water conservation plan with different kinds of structures recommended has been prepared which shows the spatial distribution and extent of area. An area of about 9 ha (2%) needs Trench cum bunding, 441 ha (74%) needs Graded Bunding and 3 ha (<1%) needs strengthening of existing bunds.

The conservation plan prepared may be presented to all the stakeholders including farmers and after considering their suggestions, the conservation plan for the microwatershed may be finalised in a participatory approach.




Fig. 9.1 Soil and Water Conservation Plan map of Mundragi-3 Microwatershed

### 9.3 Greening of Microwatershed

As part of the greening programme in the watersheds, it is envisaged to plant a variety of horticultural and other tree plants that are edible, economical and produce lot of biomass which helps to restore the ecological balance in the watersheds. The lands that are suitable for greening programme are non-arable lands (land capability classes V, VI VII and VIII) and also the lands that are not suitable or marginally suitable and field bunds for growing annual and perennial crops. The method of planting these trees is given below.

It is recommended to open pits during the 1<sup>st</sup> week of March along the contour and heap the dug-out soil on the lower side of the slope in order to harness the flowing water and facilitate weathering of soil in the pit. Exposure of soil in the pit also prevents spread of pests and diseases due to scorching sun rays. The pits should be filled with mixture of soil and organic manure during the second week of April and keep ready with sufficiently tall seedlings produced either in poly bags or in root trainer nurseries so that planting can be done during the 2<sup>nd</sup> or 3<sup>rd</sup> week of April depending on the rainfall.

The tree species suitable for the area considering rainfall, temperature and adaptability is listed below; waterlogged areas are recommended to be planted with species like Nerale (*Syzyzium cumini*) and Bamboo. Dry areas are to be planted with species like Honge, Bevu, Seetaphal etc.

<b>Dry Deciduous Species</b>			<b>Temp (°C)</b>	<b>Rainfall (mm)</b>
1.	Bevu	<i>Azadiracta indica</i>	21–32	400 –1,200
2.	Tapasi	<i>Holoptelia integrifolia</i>	20-30	500 - 1000
3.	Seetaphal	<i>Anona Squamosa</i>	20-40	400 - 1000
4.	Honge	<i>Pongamia pinnata</i>	20 -50	500– 2,500
5.	Kamara	<i>Hardwickia binata</i>	25 -35	400 - 1000
6.	Bage	<i>Albezzia lebbek</i>	20 - 45	500 - 1000
7.	Ficus	<i>Ficus bengalensis</i>	20 - 50	500–2,500
8.	Sisso	<i>Dalbargia Sissoo</i>	20 - 50	500 -2000
9.	Ailanthus	<i>Ailanthus excelsa</i>	20 - 50	500 - 1000
10.	Hale	<i>Wrightia tinctoria</i>	25 - 45	500 - 1000
11.	Uded	<i>Steriospermum chelanoides</i>	25 - 45	500 -2000
12.	Dhupa	<i>Boswella Serrata</i>	20 - 40	500 - 2000
13.	Nelli	<i>Embllica Officinalis</i>	20 - 50	500 -1500
14.	Honne	<i>Pterocarpus marsupium</i>	20 - 40	500 - 2000
<b>Moist Deciduous Species</b>			<b>Temp (°C)</b>	<b>Rainfall (mm)</b>
15.	Teak	<i>Tectona grandis</i>	20 - 50	500-5000
16.	Nandi	<i>Legarstroemia lanceolata</i>	20 - 40	500 - 4000
17.	Honne	<i>Pterocarpus marsupium</i>	20 - 40	500 - 3000
18.	Mathi	<i>Terminalia alata</i>	20 -50	500 - 2000
19.	Shivane	<i>Gmelina arborea</i>	20 -50	500 -2000
20.	Kindal	<i>T.Paniculata</i>	20 - 40	500 - 1500
21.	Beete	<i>Dalbargia latifolia</i>	20 - 40	500 - 1500
22.	Tare	<i>T. belerica</i>	20 - 40	500 - 2000
23.	Bamboo	<i>Bambusa arundinasia</i>	20 - 40	500 - 2500
24.	Bamboo	<i>Dendrocalamus strictus</i>	20 – 40	500 – 2500
25.	Muthuga	<i>Butea monosperma</i>	20 - 40	400 - 1500
26.	Hippe	<i>Madhuca latifolia</i>	20 - 40	500 - 2000
27.	Sandal	<i>Santalum album</i>	20 - 50	400 - 1000
28.	Nelli	<i>Embllica officinalis</i>	20 - 40	500 - 2000
29.	Nerale	<i>Sizyzium cumini</i>	20 - 40	500 - 2000
30.	Dhaman	<i>Grevia tilifolia</i>	20 - 40	500 - 2000
31.	Kaval	<i>Careya arborea</i>	20 - 40	500 - 2000
32.	Harada	<i>Terminalia chebula</i>	20 - 40	500 - 2000



## References

1. FAO (1976) Framework for Land Evaluation, Food and Agriculture Organization, Rome.72 pp.
2. FAO (1983) Guidelines for Land Evaluation for Rainfed Agriculture, FAO, Rome, 237 pp.
3. IARI (1971) Soil Survey Manual, All India Soil and Land Use Survey Organization, IARI, New Delhi, 121 pp.
4. Katyal, J.C. and Rattan, R.K. (2003) Secondary and Micronutrients; Research Gap and Future Needs. Fert. News 48 (4); 9-20.
5. Naidu, L.G.K., Ramamurthy, V., Challa, O., Hegde, R. and Krishnan, P. (2006) Manual Soil Site Suitability Criteria for Major Crops, NBSS Publ. No. 129, NBSS & LUP, Nagpur, 118 pp.
6. Natarajan, A. and Dipak Sarkar (2010) Field Guide for Soil Survey, National Bureau of Soil Survey and Land Use Planning (ICAR), Nagpur, India.
7. Natarajan, A., Rajendra Hegde, Raj, J.N. and Shivananda Murthy, H.G. (2015) Implementation Manual for Sujala-III Project, Watershed Development Department, Bengaluru, Karnataka.
8. Sarma, V.A.K., Krishnan, P. and Budihal, S.L. (1987) Laboratory Manual, Tech. Bull. 23, NBSS &LUP, Nagpur.
9. Sehgal, J.L. (1990) Soil Resource Mapping of Different States of India; Why and How?, National Bureau of Soil Survey and Land Use Planning, Nagpur, 49 pp.
10. Shivaprasad, C.R., R.S. Reddy, J. Sehgal and M. Velayuthum (1998) Soils of Karnataka for Optimizing Land Use, NBSS Publ. No. 47b, NBSS & LUP, Nagpur, India.
11. Soil Survey Staff (2006) Keys to Soil Taxonomy, Tenth edition, U.S. Department of Agriculture/ NRCS, Washington DC, U.S.A.
12. Soil Survey Staff (2012) Soil Survey Manual, Handbook No. 18, USDA, Washington DC, USA.





**Appendix I**  
**Mundragi-3 (1H2c) Microwatershed**  
**Soil Phase Information**

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Varkanahalli	182/1	0.65	RO	RO	RO	RO	RO	RO	RO	RO	Greengram (Gg)	Not Available	RO	RO
Varkanahalli	183	2.79	JNKcB2	LMU-3	Moderately shallow (50-75 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram+Groundnut (Gg+Gn)	Not Available	Iles	Graded bunding
Mundaragi	13	0.19	BDLhB2g1	LMU-4	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Paddy+Scrub land (Rg+pd+Sl)	Not Available	IIles	Graded bunding
Mundaragi	14	0.88	NGPmB2	LMU-2	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iles	Graded bunding
Mundaragi	15	0.05	NGPmB2	LMU-2	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Mundaragi	16	0.61	NGPmB2	LMU-2	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Mundaragi	17	2.35	TMKhA1	LMU-1	Very deep (>150 cm)	Sandy clay loam	Non gravelly (<15%)	Very high (>200 mm/m)	Nearly level (0-1%)	Slight	Habitation	Not Available	IIws	Graded bunding
Mundaragi	23	0.68	HSLcB2	LMU-2	Moderately deep (75-100 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Iles	Graded bunding
Mundaragi	24	0.09	HSLcB2	LMU-2	Moderately deep (75-100 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Scrub land (Sl)	Not Available	Iles	Graded bunding
Mundaragi	416	15.64	RO	RO	RO	RO	RO	RO	RO	RO	RO	Not Available	RO	RO
Mundaragi	450	1.38	BDPcB2	LMU-4	Very shallow (<25 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	RO	Not Available	IVes	TCB
Mundaragi	451	5.43	BDPcB2	LMU-4	Very shallow (<25 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IVes	TCB
Mundaragi	452	8.01	RO	RO	RO	RO	RO	RO	RO	RO	Redgram (Rg)	Not Available	RO	RO
Mundaragi	453	6.61	RO	RO	RO	RO	RO	RO	RO	RO	RO	Not Available	RO	RO
Mundaragi	454	4.68	JNKiB2	LMU-3	Moderately shallow (50-75 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Mundaragi	455	2.14	JNKiB2	LMU-3	Moderately shallow (50-75 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Mundaragi	456	1.85	JNKiB2	LMU-3	Moderately shallow (50-75 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Mundaragi	484	1.2	BDLiB2	LMU-4	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIles	Graded bunding
Mundaragi	486	0.28	BDLiB2	LMU-4	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut (Gn)	Not Available	IIles	Graded bunding
Mundaragi	487	0.21	BDLiB2	LMU-4	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIles	Graded bunding
Mundaragi	488	1.08	BDLiB2	LMU-4	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIles	Graded bunding
Mundaragi	489	5.55	BDLiB2	LMU-4	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut (Gn)	Not Available	IIles	Graded bunding

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Mundaragi	490	6.71	RO	RO	RO	RO	RO	RO	RO	RO	Groundnut+Paddy (Gn+Pd)	Not Available	RO	RO
Mundaragi	491	9.07	BDLiB2	LMU-4	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut (Gn)	Not Available	IIes	Graded bunding
Mundaragi	492	4.67	BDLiB2	LMU-4	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	IIes	Graded bunding
Mundaragi	505	0.1	BDLiB2	LMU-4	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Mundaragi	508	1.71	JNKC2	LMU-3	Moderately shallow (50-75 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Cotton (Rg+Ct)	Not Available	Iies	Graded bunding
Mundaragi	509	2.35	BDLiB2	LMU-4	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Cotton (Rg+Ct)	Not Available	IIes	Graded bunding
Mundaragi	510	7.73	BDLiB2	LMU-4	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Mundaragi	511	5.22	BDLiB2	LMU-4	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Mundaragi	512	6.07	BDLiB2	LMU-4	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Groundnut (Rg+Gn)	Not Available	IIes	Graded bunding
Mundaragi	513	7.12	BDLiB2	LMU-4	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Groundnut (Rg+Gn)	Not Available	IIes	Graded bunding
Mundaragi	514	7.01	BDLiB2	LMU-4	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Mundaragi	515	4.86	BDLiB2	LMU-4	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	IIes	Graded bunding
Mundaragi	516	6.64	NGPmB2	LMU-2	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Cotton (Rg+Ct)	Not Available	Iies	Graded bunding
Mundaragi	517	1.26	NGPmB2	LMU-2	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iies	Graded bunding
Mundaragi	518/1	0.15	NGPmB2	LMU-2	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Scrub land (Sl)	Not Available	Iies	Graded bunding
Mundaragi	521/1	0.4	Habitation	Others	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others
Mundaragi	521/2	0.48	Habitation	Others	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others
Mundaragi	522/1	0.71	NGPmB2	LMU-2	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Habitation	Not Available	Iies	Graded bunding
Mundaragi	522/2	0.36	Habitation	Others	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others
Mundaragi	522/3	0.34	Habitation	Others	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others
Mundaragi	522/4	0.53	Habitation	Others	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others
Mundaragi	523	2.41	BDLhB2g1	LMU-4	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Scrub land (Sl)	Not Available	IIes	Graded bunding
Mundaragi	524	1.95	BDLhB2g1	LMU-4	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Scrub land (Sl)	Not Available	IIes	Graded bunding
Mundaragi	525	3.04	BDLhB2g1	LMU-4	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IIes	Graded bunding

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Graveliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Mundaragi	526	1.8	BDLhB2g1	LMU-4	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Mundaragi	527	4.83	BDLhB2g1	LMU-4	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Mundaragi	528	4.82	BDLhB2g1	LMU-4	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	IIes	Graded bunding
Mundaragi	529	7.49	BDLhB2g1	LMU-4	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	IIes	Graded bunding
Mundaragi	530	0.4	BDLhB2g1	LMU-4	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IIes	Graded bunding
Mundaragi	531	0.33	BDLiB2	LMU-4	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IIes	Graded bunding
Mundaragi	532	0.18	BDLhB2g1	LMU-4	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIes	Graded bunding
Mundaragi	533	0.37	BDLiB2	LMU-4	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIes	Graded bunding
Mundaragi	534	7.61	BDLiB2	LMU-4	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut (Gn)	Not Available	IIes	Graded bunding
Mundaragi	535	3.59	BDLiB2	LMU-4	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut (Gn)	Not Available	IIes	Graded bunding
Mundaragi	536	4.31	BDLiB2	LMU-4	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut (Gn)	Not Available	IIes	Graded bunding
Mundaragi	537	4	BDLiB2	LMU-4	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Mundaragi	538	3.8	BDLiB2	LMU-4	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut+Cotton (Gn+Ct)	Not Available	IIes	Graded bunding
Mundaragi	539	5.26	BDLiB2	LMU-4	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Groundnut (Rg+Gn)	Not Available	IIes	Graded bunding
Mundaragi	540	5.33	BDLiB2	LMU-4	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut (Gn)	Not Available	IIes	Graded bunding
Mundaragi	541	6.28	Waterbody	Others	Others	Others	Others	Others	Others	Others	Waterbody	Not Available	Others	Others
Mundaragi	542	5.07	Waterbody	Others	Others	Others	Others	Others	Others	Others	Scrub land (SI)	Not Available	Others	Others
Mundaragi	543/1	1.57	BDLiB2	LMU-4	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Scrub land (SI)	Not Available	IIes	Graded bunding
Mundaragi	543/2	0.09	BDLiB2	LMU-4	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIes	Graded bunding
Mundaragi	543/3	1.14	BDLiB2	LMU-4	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Waterbody	Not Available	IIes	Graded bunding
Mundaragi	544	3.98	BDLhB2g1	LMU-4	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	IIes	Graded bunding
Mundaragi	545	5.17	BDLhB2g1	LMU-4	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Cotton (Rg+Ct)	Not Available	IIes	Graded bunding
Mundaragi	546	4.87	BDLhB2g1	LMU-4	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	IIes	Graded bunding
Mundaragi	547	8.09	BDLhB2g1	LMU-4	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Graveliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Mundaragi	548	4.85	BDLhB2g1	LMU-4	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Mundaragi	549	3.56	BDLhB2g1	LMU-4	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Mundaragi	550	4.14	HTKcB2	LMU-4	Shallow (25-50 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIes	Graded bunding
Mundaragi	553/1	4.11	BDLiB2	LMU-4	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Mundaragi	553/2	0.9	BDLiB2	LMU-4	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	IIes	Graded bunding
Mundaragi	557	7.93	BDLiB2	LMU-4	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Scrub land (Rg+Sl)	Not Available	IIes	Graded bunding
Mundaragi	565	3.62	HTKcB2	LMU-4	Shallow (25-50 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Scrub land (Sl)	Not Available	IIes	Graded bunding
Mundaragi	572	5.21	HTKcB2	LMU-4	Shallow (25-50 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	RO	Not Available	IIes	Graded bunding
Mundaragi	584	7.59	BDLhB2g1	LMU-4	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Mundaragi	585	6.12	BDLhB2g1	LMU-4	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Mundaragi	586	4.71	HTKcB2	LMU-4	Shallow (25-50 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Mundaragi	587	4.94	HTKcB2	LMU-4	Shallow (25-50 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Mundaragi	588	5.89	HTKcB2	LMU-4	Shallow (25-50 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	RO	Not Available	IIes	Graded bunding
Mundaragi	589	6.93	HTKcB2	LMU-4	Shallow (25-50 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton+Rockout crops (Ct+Rc)	Not Available	IIes	Graded bunding
Mundaragi	590/1	0.84	HTKcB2	LMU-4	Shallow (25-50 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton+Rockout crops (Ct+Rc)	Not Available	IIes	Graded bunding
Mundaragi	590/2	5.1	HTKcB2	LMU-4	Shallow (25-50 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton+Rockout crops (Ct+Rc)	Not Available	IIes	Graded bunding
Mundaragi	591	5.47	HTKcB2	LMU-4	Shallow (25-50 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Scrub land (Sl)	Not Available	IIes	Graded bunding
Mundaragi	592	3.47	HTKcB2	LMU-4	Shallow (25-50 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Scrub land (Sl)	Not Available	IIes	Graded bunding
Mundaragi	593	1.46	HTKcB2	LMU-4	Shallow (25-50 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Scrub land (Sl)	Not Available	IIes	Graded bunding
Mundaragi	594/1	4.35	HTKcB2	LMU-4	Shallow (25-50 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Scrub land (Sl)	Not Available	IIes	Graded bunding
Mundaragi	594/2	1.44	HTKcB2	LMU-4	Shallow (25-50 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Scrub land (Sl)	Not Available	IIes	Graded bunding
Mundaragi	596	4.06	HTKcB2	LMU-4	Shallow (25-50 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Scrub land (Sl)	Not Available	IIes	Graded bunding
Mundaragi	597	4.02	HTKcB2	LMU-4	Shallow (25-50 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Scrub land (Sl)	Not Available	IIes	Graded bunding
Mundaragi	598	7.22	Habitation	Others	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Mundaragi	599	2.89	HTKcB2	LMU-4	Shallow (25-50 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Scrub land (SI)	Not Available	IIes	Graded bunding
Mundaragi	600	5.99	HTKcB2	LMU-4	Shallow (25-50 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Scrub land (SI)	Not Available	IIes	Graded bunding
Mundaragi	601	8.91	BDLhB2g1	LMU-4	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Scrub land (SI)	Not Available	IIes	Graded bunding
Mundaragi	602	4.87	HTKcB2	LMU-4	Shallow (25-50 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Mundaragi	603	7.33	HTKcB2	LMU-4	Shallow (25-50 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Mundaragi	604	5.72	HTKcB2	LMU-4	Shallow (25-50 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Mundaragi	605	7.96	BDLhB2g1	LMU-4	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Scrub land (SI)	Not Available	IIes	Graded bunding
Mundaragi	606	8.19	Habitation	Others	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others
Mundaragi	607	6.19	BDLhB2g1	LMU-4	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Scrub land (SI)	Not Available	IIes	Graded bunding
Mundaragi	608	7.43	BDLhB2g1	LMU-4	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Scrub land+Habitation (SI+Hb)	Not Available	IIes	Graded bunding
Mundaragi	609	5.1	Habitation	Others	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others
Mundaragi	610/1	8.32	BDLhB2g1	LMU-4	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Scrub land+Fallow land (SI+FI)	Not Available	IIes	Graded bunding
Mundaragi	610/2	0.79	NGPmB2	LMU-2	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iies	Graded bunding
Mundaragi	611	7.18	BDLhB2g1	LMU-4	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Scrub land+Fallow land+Redgram (SI+FI+Rg)	Not Available	IIes	Graded bunding
Mundaragi	612	2.6	BDLhB2g1	LMU-4	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Scrubland+Groundnut (SI+Gn)	Not Available	IIes	Graded bunding
Mundaragi	613/1	1.59	NGPmB2	LMU-2	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Habitation	Not Available	Iies	Graded bunding
Mundaragi	613/2	0.39	NGPmB2	LMU-2	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Habitation	Not Available	Iies	Graded bunding
Mundaragi	613/3	1.89	BDLhB2g1	LMU-4	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Fallow land (FI)	Not Available	IIes	Graded bunding
Mundaragi	614/1	1.92	NGPmB2	LMU-2	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iies	Graded bunding
Mundaragi	614/2	0.04	BDLhB2g1	LMU-4	Shallow (25-50 cm)	Sandy clay loam	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Mundaragi	615/1	0.67	NGPmB2	LMU-2	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iies	Graded bunding
Mundaragi	615/2	0.54	NGPmB2	LMU-2	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iies	Graded bunding
Mundaragi	617	0.28	Habitation	Others	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Yadhagiri .B	30	0.41	SBRcB2	LMU-3	Moderately shallow (50-75 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Fallow land (Fl)	Not Available	Iles	Graded bunding
Yadhagiri .B	31	1.41	SBRcB2	LMU-3	Moderately shallow (50-75 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Fallow land (Fl)	Not Available	Iles	Graded bunding
Yadhagiri .B	33	7.37	SBRcB2	LMU-3	Moderately shallow (50-75 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Fallow land (Fl)	Not Available	Iles	Graded bunding
Yadhagiri .B	34	0.19	SBRcB2	LMU-3	Moderately shallow (50-75 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Fallow land (Fl)	Not Available	Iles	Graded bunding
Yadhagiri .B	35	0.15	SBRcB2	LMU-3	Moderately shallow (50-75 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Fallow land (Fl)	Not Available	Iles	Graded bunding
Yadhagiri .B	36	7.37	SBRcB2	LMU-3	Moderately shallow (50-75 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Fallow land (Fl)	Not Available	Iles	Graded bunding
Yadhagiri .B	37	4.88	MI	MI	MI	MI	MI	MI	MI	MI	Not Available (NA)	Not Available	MI	MI
Yadhagiri .B	38	8.65	MI	MI	MI	MI	MI	MI	MI	MI	RO	Not Available	MI	MI
Yadhagiri .B	39	0.12	SBRcB2	LMU-3	Moderately shallow (50-75 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	RO	Not Available	Iles	Graded bunding
Yadhagiri .B	40	5.24	SBRcB2	LMU-3	Moderately shallow (50-75 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Fallow land (Fl)	Not Available	Iles	Graded bunding
Yadhagiri .B	41	6.03	SBRcB2	LMU-3	Moderately shallow (50-75 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Fallow land (Fl)	Not Available	Iles	Graded bunding
Yadhagiri .B	42	8.08	RO	RO	RO	RO	RO	RO	RO	RO	RO	Not Available	RO	RO
Yadhagiri .B	43	9.31	SBRcB2	LMU-3	Moderately shallow (50-75 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Iles	Graded bunding
Yadhagiri .B	44	8.44	SBRcB2	LMU-3	Moderately shallow (50-75 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Iles	Graded bunding
Yadhagiri .B	45	7.8	SBRcB2	LMU-3	Moderately shallow (50-75 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram+Cotton (Gg+Ct)	Not Available	Iles	Graded bunding
Yadhagiri .B	46	6.79	SBRcB2	LMU-3	Moderately shallow (50-75 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram+Cotton (Gg+Ct)	Not Available	Iles	Graded bunding
Yadhagiri .B	47	7.28	SBRcB2	LMU-3	Moderately shallow (50-75 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Iles	Graded bunding
Yadhagiri .B	48	4.59	SBRcB2	LMU-3	Moderately shallow (50-75 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Iles	Graded bunding
Yadhagiri .B	49	1.21	SBRcB2	LMU-3	Moderately shallow (50-75 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Yadhagiri .B	52	6.1	SBRcB2	LMU-3	Moderately shallow (50-75 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Cotton (Rg+Ct)	1 Borewell	Iles	Graded bunding
Yadhagiri .B	53	4.39	RO	RO	RO	RO	RO	RO	RO	RO	Redgram+Fallow land (Rg+Fl)	Not Available	RO	RO
Yadhagiri .B	73	29.55	RO	RO	RO	RO	RO	RO	RO	RO	RO	Not Available	RO	RO

MI- Mining/ industrial

**Appendix II**  
**Mundragi-3 Microwatershed**  
**Soil Fertility Information**

.Village	Survey No	Soil Reaction	Salinity	Organic Carbon	Available Phosphorus	Available Potassium	Available Sulphur	Available Boron	Available Iron	Available Manganese	Available Copper	Available Zinc
Varkanahalli	182/1	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro
Varkanahalli	183	Slightly alkaline (pH 7.3 - 7.8)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	Medium (10 - 20 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)
Mundaragi	13	Neutral (pH 6.5 - 7.3)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Mundaragi	14	Neutral (pH 6.5 - 7.3)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Mundaragi	15	Neutral (pH 6.5 - 7.3)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Mundaragi	16	Neutral (pH 6.5 - 7.3)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Mundaragi	17	Neutral (pH 6.5 - 7.3)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Low (< 23 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Mundaragi	23	Neutral (pH 6.5 - 7.3)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Low (< 23 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Mundaragi	24	Neutral (pH 6.5 - 7.3)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Low (< 23 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Mundaragi	416	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro
Mundaragi	450	Neutral (pH 6.5 - 7.3)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Mundaragi	451	Neutral (pH 6.5 - 7.3)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	Medium (10 - 20 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Mundaragi	452	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro
Mundaragi	453	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro
Mundaragi	454	Neutral (pH 6.5 - 7.3)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	Low (<145 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Mundaragi	455	Neutral (pH 6.5 - 7.3)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Mundaragi	456	Neutral (pH 6.5 - 7.3)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	Low (<145 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Mundaragi	484	Neutral (pH 6.5 - 7.3)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Mundaragi	486	Neutral (pH 6.5 - 7.3)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Mundaragi	487	Neutral (pH 6.5 - 7.3)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Mundaragi	488	Neutral (pH 6.5 - 7.3)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	Low (<145 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Mundaragi	489	Neutral (pH 6.5 - 7.3)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	Low (<145 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Mundaragi	490	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro











.Village	Survey No	Soil Reaction	Salinity	Organic Carbon	Available Phosphorus	Available Potassium	Available Sulphur	Available Boron	Available Iron	Available Manganese	Available Copper	Available Zinc
Yadhagiri .B	36	Slightly alkaline (pH 7.3 - 7.8)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Low (< 23 kg/ha)	Medium (145 - 337 kg/ha)	Medium (10 - 20 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Yadhagiri .B	37	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI
Yadhagiri .B	38	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI
Yadhagiri .B	39	Slightly alkaline (pH 7.3 - 7.8)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Low (< 23 kg/ha)	Medium (145 - 337 kg/ha)	Medium (10 - 20 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Yadhagiri .B	40	Slightly alkaline (pH 7.3 - 7.8)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Low (< 23 kg/ha)	Medium (145 - 337 kg/ha)	Medium (10 - 20 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Yadhagiri .B	41	Slightly alkaline (pH 7.3 - 7.8)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Low (< 23 kg/ha)	Medium (145 - 337 kg/ha)	Medium (10 - 20 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Yadhagiri .B	42	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro
Yadhagiri .B	43	Slightly alkaline (pH 7.3 - 7.8)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Low (< 23 kg/ha)	Medium (145 - 337 kg/ha)	Medium (10 - 20 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Yadhagiri .B	44	Slightly alkaline (pH 7.3 - 7.8)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	Medium (10 - 20 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)
Yadhagiri .B	45	Slightly alkaline (pH 7.3 - 7.8)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	Medium (10 - 20 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)
Yadhagiri .B	46	Slightly alkaline (pH 7.3 - 7.8)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	Medium (10 - 20 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)
Yadhagiri .B	47	Slightly alkaline (pH 7.3 - 7.8)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	Medium (10 - 20 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)
Yadhagiri .B	48	Slightly alkaline (pH 7.3 - 7.8)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	Medium (10 - 20 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)
Yadhagiri .B	49	Slightly alkaline (pH 7.3 - 7.8)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	Medium (10 - 20 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)
Yadhagiri .B	52	Slightly alkaline (pH 7.3 - 7.8)	Non saline (<2 dsm)	Medium (0.5 - 0.75 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	Medium (10 - 20 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)
Yadhagiri .B	53	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro
Yadhagiri .B	73	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro

**Appendix III**  
**Mundragi-3 Microwatershed**  
**Soil Suitability Information**

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry	
Varkanahalli	182/1	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	
Varkanahalli	183	N1r	S2t	S3r	S2r	S3r	S2r	N1r	S3r	S2r	S3r	S3r	S2r	S3r	S2r	N1rt	S3r	S3r	S3t	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Mundaragi	13	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	
Mundaragi	14	S3t	S2t	S3t	S2w	S3t	S1	S2t	S1	S1	S2w	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S3tw	S3tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw	
Mundaragi	15	S3t	S2t	S3t	S2w	S3t	S1	S2t	S1	S1	S2w	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S3tw	S3tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw	
Mundaragi	16	S3t	S2t	S3t	S2w	S3t	S1	S2t	S1	S1	S2w	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S3tw	S3tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw	
Mundaragi	17	S3tw	S2tw	S3tw	S1	S3tw	S1	S2tw	S2zw	S1	S2rw	S2tw	S1	S3tw	S1	S1tz	S2tw	S2zw	S3tw	S2t	S2tw	S3tw	S2tw	S2tw	S2tw	S2t	S1	S1	S2tw	S3tw	
Mundaragi	23	S3rz	S2tz	S2rz	S2tz	S2rz	S3tz	S3rz	S2rz	S3tz	S2rz	S2rz	S2tz	S2rz	S1	N1tz	S3rz	S2rz	S2z	S1	S2z	S2z	S2z	S2z	S2rz	S2z	S2t	S1	S2rz	S2rz	
Mundaragi	24	S3rz	S2tz	S2rz	S2tz	S2rz	S3tz	S3rz	S2rz	S3tz	S2rz	S2rz	S2tz	S2rz	S1	N1tz	S3rz	S2rz	S2z	S1	S2z	S2z	S2z	S2z	S2rz	S2z	S2t	S1	S2rz	S2rz	
Mundaragi	416	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	
Mundaragi	450	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Mundaragi	451	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Mundaragi	452	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	
Mundaragi	453	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	
Mundaragi	454	N1r	S2t	S3r	S2r	S3r	S2r	N1r	S3r	S2r	S3r	S3r	S2r	S3r	S2r	N1rt	S3r	S3r	S3t	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Mundaragi	455	N1r	S2t	S3r	S2r	S3r	S2r	N1r	S3r	S2r	S3r	S3r	S2r	S3r	S2r	N1rt	S3r	S3r	S3t	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Mundaragi	456	N1r	S2t	S3r	S2r	S3r	S2r	N1r	S3r	S2r	S3r	S3r	S2r	S3r	S2r	N1rt	S3r	S3r	S3t	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Mundaragi	484	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	
Mundaragi	486	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	
Mundaragi	487	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	
Mundaragi	488	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	
Mundaragi	489	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	
Mundaragi	490	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	
Mundaragi	491	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry	
Mundaragi	492	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	
Mundaragi	505	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	
Mundaragi	508	N1r	S2t	S3r	S2r	S3r	S2r	N1r	S3r	S2r	S3r	S3r	S2r	S3r	S2r	N1rt	S3r	S3r	S3t	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Mundaragi	509	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	
Mundaragi	510	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	
Mundaragi	511	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	
Mundaragi	512	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	
Mundaragi	513	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	
Mundaragi	514	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	
Mundaragi	515	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	
Mundaragi	516	S3t	S2t	S3t	S2w	S3t	S1	S2t	S1	S1	S2w	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S3tw	S3tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw	
Mundaragi	517	S3t	S2t	S3t	S2w	S3t	S1	S2t	S1	S1	S2w	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S3tw	S3tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw	
Mundaragi	518/1	S3t	S2t	S3t	S2w	S3t	S1	S2t	S1	S1	S2w	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S3tw	S3tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw	
Mundaragi	521/1	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Mundaragi	521/2	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Mundaragi	522/1	S3t	S2t	S3t	S2w	S3t	S1	S2t	S1	S1	S2w	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S3tw	S3tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw	
Mundaragi	522/2	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Mundaragi	522/3	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Mundaragi	522/4	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Mundaragi	523	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	
Mundaragi	524	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	
Mundaragi	525	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	
Mundaragi	526	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	
Mundaragi	527	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	
Mundaragi	528	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	
Mundaragi	529	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry	
Mundaragi	530	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	
Mundaragi	531	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	
Mundaragi	532	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	
Mundaragi	533	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	
Mundaragi	534	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	
Mundaragi	535	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	
Mundaragi	536	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	
Mundaragi	537	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	
Mundaragi	538	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	
Mundaragi	539	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	
Mundaragi	540	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	
Mundaragi	541	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Mundaragi	542	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Mundaragi	543/1	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	
Mundaragi	543/2	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	
Mundaragi	543/3	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	
Mundaragi	544	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	
Mundaragi	545	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	
Mundaragi	546	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	
Mundaragi	547	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	
Mundaragi	548	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	
Mundaragi	549	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	
Mundaragi	550	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Mundaragi	553/1	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	
Mundaragi	553/2	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	
Mundaragi	557	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry	
Mundaragi	565	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Mundaragi	572	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Mundaragi	584	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	
Mundaragi	585	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	
Mundaragi	586	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Mundaragi	587	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Mundaragi	588	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Mundaragi	589	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Mundaragi	590/1	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Mundaragi	590/2	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Mundaragi	591	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Mundaragi	592	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Mundaragi	593	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Mundaragi	594/1	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Mundaragi	594/2	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Mundaragi	596	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Mundaragi	597	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Mundaragi	598	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Mundaragi	599	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Mundaragi	600	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Mundaragi	601	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	
Mundaragi	602	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Mundaragi	603	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Mundaragi	604	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Mundaragi	605	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	
Mundaragi	606	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others



Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry	
Mundaragi	607	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	
Mundaragi	608	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	
Mundaragi	609	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Mundaragi	610/1	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	
Mundaragi	610/2	S3t	S2t	S3t	S2w	S3t	S1	S2t	S1	S1	S2w	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S3tw	S3tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw	
Mundaragi	611	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	
Mundaragi	612	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	
Mundaragi	613/1	S3t	S2t	S3t	S2w	S3t	S1	S2t	S1	S1	S2w	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S3tw	S3tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw	
Mundaragi	613/2	S3t	S2t	S3t	S2w	S3t	S1	S2t	S1	S1	S2w	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S3tw	S3tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw	
Mundaragi	613/3	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	
Mundaragi	614/1	S3t	S2t	S3t	S2w	S3t	S1	S2t	S1	S1	S2w	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S3tw	S3tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw	
Mundaragi	614/2	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	
Mundaragi	615/1	S3t	S2t	S3t	S2w	S3t	S1	S2t	S1	S1	S2w	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S3tw	S3tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw	
Mundaragi	615/2	S3t	S2t	S3t	S2w	S3t	S1	S2t	S1	S1	S2w	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S3tw	S3tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw	
Mundaragi	617	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Yadhagiri .B	30	N1r	S3t	S3rt	S3t	S3rt	N1t	N1r	S3rt	N1t	S3t	S3t	S3t	S3rt	S3t	S3rt	S3rt	S3rt	S3t	S3t	S3t	S3t	S3t	S3t	S3rt	S3t	S3t	S3t	S3t	S3t	
Yadhagiri .B	31	N1r	S3t	S3rt	S3t	S3rt	N1t	N1r	S3rt	N1t	S3t	S3t	S3t	S3rt	S3t	S3rt	S3rt	S3rt	S3t	S3t	S3t	S3t	S3t	S3t	S3rt	S3t	S3t	S3t	S3t	S3t	S3t
Yadhagiri .B	33	N1r	S3t	S3rt	S3t	S3rt	N1t	N1r	S3rt	N1t	S3t	S3t	S3t	S3rt	S3t	S3rt	S3rt	S3rt	S3t	S3t	S3t	S3t	S3t	S3t	S3rt	S3t	S3t	S3t	S3t	S3t	S3t
Yadhagiri .B	34	N1r	S3t	S3rt	S3t	S3rt	N1t	N1r	S3rt	N1t	S3t	S3t	S3t	S3rt	S3t	S3rt	S3rt	S3rt	S3t	S3t	S3t	S3t	S3t	S3t	S3rt	S3t	S3t	S3t	S3t	S3t	S3t
Yadhagiri .B	35	N1r	S3t	S3rt	S3t	S3rt	N1t	N1r	S3rt	N1t	S3t	S3t	S3t	S3rt	S3t	S3rt	S3rt	S3rt	S3t	S3t	S3t	S3t	S3t	S3t	S3rt	S3t	S3t	S3t	S3t	S3t	S3t
Yadhagiri .B	36	N1r	S3t	S3rt	S3t	S3rt	N1t	N1r	S3rt	N1t	S3t	S3t	S3t	S3rt	S3t	S3rt	S3rt	S3rt	S3t	S3t	S3t	S3t	S3t	S3t	S3rt	S3t	S3t	S3t	S3t	S3t	S3t
Yadhagiri .B	37	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI
Yadhagiri .B	38	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI	MI
Yadhagiri .B	39	N1r	S3t	S3rt	S3t	S3rt	N1t	N1r	S3rt	N1t	S3t	S3t	S3t	S3rt	S3t	S3rt	S3rt	S3rt	S3t	S3t	S3t	S3t	S3t	S3t	S3rt	S3t	S3t	S3t	S3t	S3t	S3t
Yadhagiri .B	40	N1r	S3t	S3rt	S3t	S3rt	N1t	N1r	S3rt	N1t	S3t	S3t	S3t	S3rt	S3t	S3rt	S3rt	S3rt	S3t	S3t	S3t	S3t	S3t	S3t	S3rt	S3t	S3t	S3t	S3t	S3t	S3t
Yadhagiri .B	41	N1r	S3t	S3rt	S3t	S3rt	N1t	N1r	S3rt	N1t	S3t	S3t	S3t	S3rt	S3t	S3rt	S3rt	S3rt	S3t	S3t	S3t	S3t	S3t	S3t	S3rt	S3t	S3t	S3t	S3t	S3t	S3t

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry	
Yadhagiri .B	42	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro
Yadhagiri .B	43	N1r	S3t	S3rt	S3t	S3rt	N1t	N1r	S3rt	N1t	S3t	S3t	S3t	S3rt	S3t	S3rt	S3rt	S3rt	S3t	S3t	S3t	S3t	S3t	S3t	S3rt	S3t	S3t	S3t	S3t	S3t	S3t
Yadhagiri .B	44	N1r	S3t	S3rt	S3t	S3rt	N1t	N1r	S3rt	N1t	S3t	S3t	S3t	S3rt	S3t	S3rt	S3rt	S3rt	S3t	S3t	S3t	S3t	S3t	S3t	S3rt	S3t	S3t	S3t	S3t	S3t	S3t
Yadhagiri .B	45	N1r	S3t	S3rt	S3t	S3rt	N1t	N1r	S3rt	N1t	S3t	S3t	S3t	S3rt	S3t	S3rt	S3rt	S3rt	S3t	S3t	S3t	S3t	S3t	S3t	S3rt	S3t	S3t	S3t	S3t	S3t	S3t
Yadhagiri .B	46	N1r	S3t	S3rt	S3t	S3rt	N1t	N1r	S3rt	N1t	S3t	S3t	S3t	S3rt	S3t	S3rt	S3rt	S3rt	S3t	S3t	S3t	S3t	S3t	S3t	S3rt	S3t	S3t	S3t	S3t	S3t	S3t
Yadhagiri .B	47	N1r	S3t	S3rt	S3t	S3rt	N1t	N1r	S3rt	N1t	S3t	S3t	S3t	S3rt	S3t	S3rt	S3rt	S3rt	S3t	S3t	S3t	S3t	S3t	S3t	S3rt	S3t	S3t	S3t	S3t	S3t	S3t
Yadhagiri .B	48	N1r	S3t	S3rt	S3t	S3rt	N1t	N1r	S3rt	N1t	S3t	S3t	S3t	S3rt	S3t	S3rt	S3rt	S3rt	S3t	S3t	S3t	S3t	S3t	S3t	S3rt	S3t	S3t	S3t	S3t	S3t	S3t
Yadhagiri .B	49	N1r	S3t	S3rt	S3t	S3rt	N1t	N1r	S3rt	N1t	S3t	S3t	S3t	S3rt	S3t	S3rt	S3rt	S3rt	S3t	S3t	S3t	S3t	S3t	S3t	S3rt	S3t	S3t	S3t	S3t	S3t	S3t
Yadhagiri .B	52	N1r	S3t	S3rt	S3t	S3rt	N1t	N1r	S3rt	N1t	S3t	S3t	S3t	S3rt	S3t	S3rt	S3rt	S3rt	S3t	S3t	S3t	S3t	S3t	S3t	S3rt	S3t	S3t	S3t	S3t	S3t	S3t
Yadhagiri .B	53	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro
Yadhagiri .B	73	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro

Ro-Rock outcrops

# **PART-B**

**SOCIO-ECONOMIC STATUS OF FARM HOUSEHOLDS**



## CONTENTS

1.	Findings of the socio-economic survey	1-2
2.	Introduction	3
3	Methodology	4-5
4	Salient features of the survey	7-22
5	Summary	23-26



## LIST OF TABLES

1	Households sampled for socio economic survey	7
2	Population characteristics	7
3	Age wise classification of household members	7
4	Education level of household members	8
5	Occupation of household heads	8
6	Occupation of family members	8
7	Institutional participation of household members	9
8	Type of house owned by households	9
9	Durable assets owned by households	9
10	Average value of durable assets owned by households	9
11	Farm implements owned by households	10
12	Average value of farm implements	10
13	Livestock possession by households	10
14	Average labour availability	10
15	Adequacy of hired labour	11
16	Distribution of land (ha)	11
17	Average land value (Rs./ha)	11
18	Status of bore wells	11
19	Source of irrigation	11
20	Depth of water(Avg in meters)	12
21	Irrigated area (ha)	12
22	Cropping pattern	12
23	Cropping intensity	12
24.a	Cost of cultivation of Red gram	13
24.b	Cost of cultivation of Cotton	14
24.c	Cost of cultivation of Groundnut	15
24.d	Cost of cultivation of Sorghum	16
24.e	Cost of cultivation of Green gram	17
25	Adequacy of fodder	18
26	Annual gross income	18
27	Average annual expenditure	18

28	Horticultural species grown	18
29	Forest species grown	19
30	Average additional investment capacity	19
31	Source of funds for additional investment	19
32	Marketing of the agricultural produce	19
33	Marketing channels used for sale of agricultural produce	20
34	Mode of transport of agricultural produce	20
35	Incidence of soil and water erosion problems	20
36	Interest shown towards soil testing	20
37	Usage pattern of fuel for domestic use	20
38	Source of drinking water	21
39	Source of light	21
40	Existence of sanitary toilet facility	21
41	Possession of public distribution system (PDS) card	21
42	Adequacy of food items	21
43	Inadequacy of food items	22
44	Farming constraints experienced	22



**FINDINGS OF THE SOCIO-ECONOMIC SURVEY**

- ❖ *The survey was conducted in Mundragi-3 is located at North latitude 16° 46' 39.624" and 16° 45' 25.814" and East longitude 77° 12' 9.516" and 77° 9' 33.277" covering an area of about 598.12 ha coming unde Mundaragi and Yadhagiri B villages of Yadagiri taluk.*
- ❖ *Socio-economic analysis of Mundragi-3 micro watersheds of Belagiri sub-watershed, Yadgiri taluk & District indicated that, out of the total sample of 36 total respondents, 18 (50.00 %) were marginal, 10 (27.78%)were small, 2 (5.56 %) were Semi medium and 2 (5.56 %) were medium farmers.*
- ❖ *The population characteristics of households indicated that, there were 78 (57.35%) men and 58 (42.65 %) were women.*
- ❖ *Majority of the respondents (47.06%) were in the age group of 16-35 years.*
- ❖ *Education level of the sample households indicated that, there were 66.91 per cent illiterates, 29.41 per cent pre university education and 3.68 per cent attained graduation.*
- ❖ *About, 61.11 per cent of household heads practicing agriculture and 36.11 per cent of the household heads were engaged as agricultural labourers.*
- ❖ *Agriculture was the major occupation for 37.50 per cent of the household members.*
- ❖ *In the study area, 91.67 per cent of the households possess katcha house and 2.78 per cent possess pucca house.*
- ❖ *The durable assets owned by the households showed that, 88.89 per cent possess TV, 8.33 per cent possess mixer grinder, 86.11 per cent possess mobile phones and 2.78 per cent possess motor cycles.*
- ❖ *Farm implements owned by the households indicated that, 22.22 per cent of the households possess plough and 5.56 per cent possess bullock cart.*
- ❖ *Regarding livestock possession by the households, 16.67 per cent possess local cow and 2.78 per cent possess buffalo.*
- ❖ *The average labour availability in the study area showed that, own labour men available in the micro watershed was 1.97, women available in the micro watershed was 1.41, hired labour (men) available was 7.62 and hired labour (women) available was 6.79.*
- ❖ *Out of the total land holding of the sample respondents 56.78 per cent (26.96 ha) of the area is under dry condition and the remaining 43.22 per cent area is irrigated land.*
- ❖ *There were 11.00 live bore wells among the sampled households.*
- ❖ *Bore/open well was the major source of irrigation for 30.56 per cent of the households.*
- ❖ *The major crops grown by sample farmers are Red gram, Cotton, Groundnut, Sorghum and Green gram and cropping intensity was recorded as 100.00 per cent.*

- ❖ *The per hectare cost of cultivation for Red gram, Cotton, Groundnut, Sorghum and Green gram was Rs.37056.78, 53774.67, 76635.07, 39040.33 and 108930.93 with benefit cost ratio of 1:1.90, 1: 1.60, 1: 1.10, 1: 2.10 and 1:1.02 respectively.*
- ❖ *Further, 33.33 per cent of the households opined that dry fodder was adequate and 16.67 per cent of the households have opined that the green fodder was adequate.*
- ❖ *The average annual gross income of the farmers was Rs. 114444.44 in micro-watershed, of which Rs. 57250.00 comes from agriculture.*
- ❖ *Sampled households have grown 8 horticulture trees and 78 forestry trees together in the fields and back yards.*
- ❖ *Households have an average investment capacity of Rs. 2555.56 for land development and Rs. 833.33 for irrigation facility.*
- ❖ *Source of funds for additional investment is concerned, 8.33 per cent depends on bank loan for land development activities.*
- ❖ *Regarding marketing channels, 83.33 per cent of the households have sold agricultural produce to the local/village merchants, while, 2.78 per cent have sold in regulated markets.*
- ❖ *Further, 86.11 per cent of the households have used tractor for the transport of agriculture commodity.*
- ❖ *Majority of the farmers (86.11%) have experienced soil and water erosion problems in the watershed and 80.56 per cent of the households were interested towards soil testing.*
- ❖ *Fire was the major source of fuel for domestic use for 100.00 per cent of the households.*
- ❖ *Piped supply was the major source for drinking water for 97.22 per cent of the households.*
- ❖ *Electricity was the major source of light for 100.00 per cent of the households.*
- ❖ *In the study area, 77.78 per cent of the households possess toilet facility.*
- ❖ *Regarding possession of PDS card, 100.00 per cent of the households possessed BPL card.*
- ❖ *Households opined that, the requirement of cereals (100.00%), pulses (83.33%) and oilseeds (2.78%) are adequate for consumption.*
- ❖ *Farming constraints experienced by households in the micro watersheds were lower fertility status of the soil (88.89%) wild animal menace on farm field (8.33%), frequent incidence of pest and diseases (77.78%), inadequacy of irrigation water (8.33%), high cost of fertilizers and plant protection chemicals (77.78%), high rate of interest on credit (5.56%), low price for the agricultural commodities (80.56%), lack of marketing facilities in the area (75.00%), inadequate extension services (8.33%), lack of transport for safe transport of the agricultural produce to the market(80.56%).*

## INTRODUCTION

Soil and water are the two precious natural resources which are essential for crop production and existence of life on earth. Rainfed agriculture is under severe stress due to various constraints related to agriculture like uneven and erratic distribution of rainfall, indiscriminate use of fertilizers, chemicals and pesticides, adoption of improper land management practices, soil erosion, decline in soil fertility, decline in ground water resources leading to low crop productivity. The area under rainfed agriculture has to be managed effectively using the best available practices to enhance the production of food, fodder and fuel. This is possible if the land resources are characterized at each parcel of land through detailed land resource inventory using the best available techniques of remote sensing, GPS and GIS. The watershed development programs are aimed at the sustainable distribution of its resources and the process of creating and implementing plans, programs, and projects to sustain and enhance watershed functions that affect the plant, animal and human communities within a watershed boundary.

World Bank funded KWDP II, SUJALA III project was implemented in with Broad objective of demonstrating more effective watershed management through greater integration of programmes related to rain-fed agriculture, innovative and science based approaches and strengthen institutional capacities and If successful, it is expected that the systems and tools could be mainstreamed into the overall IWMP in the State of Karnataka and in time, throughout other IWMP operations in India. With this background the socio-economic survey has been carried out with following specific objectives:

1. To understand the demographic features of the households in the micro-watershed
2. To understand the extent of family labour available and additional employment opportunities available within the village.
3. To know the status of assets of households in the micro-watershed for suggesting possible improvements.
4. To study the cropping pattern, cropped area and productivity levels of different households in micro-watershed.
5. To determine the type and extent of livestock owned by different categories of HHs
6. Availability of fodder and level of livestock management.

### **Scope and importance of survey**

Survey helps in identification of different socio-economic and resource use-patterns of farmers at the Micro watershed. Household survey provides demographic features, labor force, and levels of education; land ownership and asset position (including livestock and other household assets) of surveyed households; and cropping patterns, input intensities, and average crop yields from farmers' fields. It also discusses crop utilization and the degree of commercialization of production in the areas; farmers' access to and utilization of credit from formal and informal sources; and the level of adoption and use of soil, water, and pest management technologies.

## METHODOLOGY

The description of the methods, components selected for the survey and procedures followed in conducting the baseline survey are furnished under the following heads.

### 1. Description of the study area

Yadgir District is one of the 30 districts of Karnataka state in southern India. This district was carved out from the erstwhile Gulbarga district as the 30th district of Karnataka on 10 April 2010. Yadgir town is the administrative headquarters of the district. The district comprises of 3 taluks namely, Shahapur, Yadgiri and Shorapur (There are 16 hoblies, 117 Gram Panchayats, 4 Municipalities, 8 Towns/ Urban agglomeration and 487 inhabited & 32 un-inhabited villages The district occupies an area of 5,160.88 km<sup>2</sup>).

Yadgir district is the second smallest district in the state, area wise is very rich in cultural traditions. The vast stretch of fertile black soil of the district is known for bumper red gram and jowar crops. The district is a "Daal bowl" of the state. The district is also known for cluster of cement industries and a distinct stone popularly known as "Malakheda Stone". Two main rivers, Krishna and Bhima, and a few tributaries flow in this region. Krishna and Bhima Rivers drain the district. They constitute the two major river basins of the district. Kagna and Amarja are the two sub - basins of Bhima River, which occur within the geographical area of the district

According to the 2011 census Yadgir district has a population of 1, 172,985, roughly equal to the nation of Timor-Leste or the US state of Rhode Island. This gives it a ranking of 404th in India (out of a total of 640). The district has a population density of 224 inhabitants per square kilometre (580/sq mi). Its population growth rate over the decade 2001-2011 was 22.67%. Yadgir has a sex ratio of 984 females for every 1000 males, and a literacy rate of 52.36%.

### 2. Locale of the survey and description of the micro-watershed and

The study was conducted in Mundragi-3 micro-watershed (Belagiri sub-watershed, Yadgiri taluk & District) is located at North latitude 16<sup>o</sup> 46' 39.624" and 16<sup>o</sup> 45' 25.814" and East longitude 77<sup>o</sup> 12' 9.516" and 77<sup>o</sup> 9' 33.277" covering an area of about 598.12 ha bounded by unde Mundaragi and Yadhagiri B Villages.

### 3. Selection of the respondents for the study

The micro-watershed is marked with 320 square meters grids. One farmer from every alternate grid in the micro-watershed was selected for the study and interviewed for socio-economic data. Totally 36 households were interviewed for the survey.

### 4. The parameters considered for socio-economic survey of households

Two forms of data were collected from the micro-watershed which includes primary data from the farm households and secondary data about the villages under the micro-watershed jurisdiction.

The following parameters were considered for the primary data collection about the socio-economic data of the households, (1) Demographic information, (2) Farm and durable assets owned by households, (3) Livestock possession, (4) Labour availability, (5) Level of migration in the village, Land holding, (7) Cropping pattern, (8) Source of irrigation, (9) Borrowing status, (10) Cost of cultivation of major crops, (11) Economics of subsidiary activities, (12) Fodder availability, (13) Family annual income from different sources, (14) Horticulture and forestry species grown, (15) Additional investment capacity, (16) Marketing practices, (17) Status of soil and water conservation structure, (18) Access to basic needs and (19) Constraints and suggestion.

The following parameters were considered for the secondary data regarding the villages under the micro-watershed jurisdiction, (1) Number of villages in each micro-watershed jurisdiction, (2) Village wise number of households, (3) Geographical area of the villages, (4) Cultivable area including rain fed and irrigated, (5) Number and type of house in each village, (6) Human and livestock population, (7) Facilities in the village such as roads, transport facility for conveyance, drinking water supply, street light and (8) Community based organizations in the villages.

## **5. Development of interview schedule and data collection**

Taking into the consideration the objectives of the survey, an interview schedule was prepared after thorough consultation with the experts in the field of social sciences. A comprehensive interview schedule covering all the major parameters for measuring the socio-economic situation was developed.

## **6. Tools used to analyze the data**

The statistical components such as frequency and percentage were used to analyze the data.

### **Abbreviations used in the report**

LL=Landless

MF=Marginal Farmers

SF=Small farmers

SMF=Semi medium farmers

MDF=Medium farmers

LF=Large Farmers



## FINDINGS OF THE SURVEY

This chapter deals with systematic presentation of results of the survey. Keeping in view the objectives, the salient features of the survey are presented under the following headings.

**Households sampled for socio-economic survey:** The data on households sampled for socio economic survey in Mundragi-3 Micro watershed is presented in Table 1 and it indicated that 36 farmers were sampled in Mundragi-3 micro-watershed among households surveyed 18 (50.00%) were marginal, 10 (27.78%) were small, 2 (5.56 %) were semi medium, 2 (5.56 %) were medium farmers. 4 landless farmers were also interviewed for the survey.

**Table 1. Households sampled for socio economic survey in Mundragi-3 micro-watershed**

Sl.No.	Particulars	LL (4)		MF (18)		SF (10)		SMF (2)		MDF (2)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Farmers	4	11.1	18	50	10	27.8	2	5.56	2	5.56	36	100

**Population characteristics:** The population characteristics of households sampled for socio-economic survey in Mundragi-3 Micro watershed is presented in Table 2. The data indicated that, there were 78 (57.35%) men and 58 (42.65%) were women.

**Table 2. Population characteristics in Mundragi-3 micro-watershed**

Sl.No.	Particulars	LL (12)		MF (67)		SF (34)		SMF (14)		MDF (9)		All (136)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Men	8	66.7	39	58	16	47	8	57.1	7	77.8	78	57.4
2	Women	4	33.3	28	42	18	53	6	42.9	2	22.2	58	42.7
Total		12	100	67	100	34	100	14	100	9	100	136	100
Average		3.0		3.7		3.4		7.0		4.5		3.8	

**Age wise classification of population:** The age wise classification of household members in Mundragi-3 Micro watershed is presented in Table 3. The indicated that, 20 (14.71%) of population were 0-15 years of age, 64 (47.06%) were 16-35 years of age, 40(29.41%) were 36-60 years of age and 12 (8.82 %) were above 61 years of age.

**Table 3: Age wise classification of members of the household in Mundragi-3 micro-watershed**

Sl.No.	Particulars	LL (12)		MF (67)		SF (34)		SMF (14)		MDF (9)		All (136)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	0-15 years of age	3	25	13	19.4	4	11.8	0	0	0	0	20	14.71
2	16-35 years of age	6	50	28	41.8	15	44.1	10	71.43	5	56	64	47.06
3	36-60 years of age	3	25	20	29.9	13	38.2	1	7.14	3	33	40	29.41
4	> 61 years	0	0	6	8.96	2	5.88	3	21.43	1	11	12	8.82
Total		12	100	67	100	34	100	14	100	9	100	136	100

**Education level of household members:** Education level of household members in Mundragi-3 Micro watershed is presented in Table 4. The results indicated that, there were 66.91 per cent of illiterates, 15.44 per cent of them had primary school education, 2.21 per

cent middle school education, 8.82 per cent high school education, 1.47 per cent of them had PUC education, 0.74 per cent of them had ITI, 3.68 per cent attained graduation and 0.74 them had other education.

**Table 4. Education level of members of the household in Mundragi-3 micro-watershed**

Sl.No.	Particulars	LL (12)		MF (67)		SF (34)		SMF (14)		MDF (9)		All (136)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Illiterate	9	75	37	55.2	26	76.5	14	100	5	55.56	91	66.9
2	Primary School	3	25	12	17.9	5	14.7	0	0	1	11.11	21	15.4
3	Middle School	0	0	3	4.48	0	0	0	0	0	0	3	2.21
4	High School	0	0	8	11.9	3	8.82	0	0	1	11.11	12	8.82
5	PUC	0	0	2	2.99	0	0	0	0	0	0	2	1.47
6	ITI	0	0	0	0	0	0	0	0	1	11.11	1	0.74
7	Degree	0	0	4	5.97	0	0	0	0	1	11.11	5	3.68
8	Others	0	0	1	1.49	0	0	0	0	0	0	1	0.74
Total		12	100	67	100	34	100	14	100	9	100	136	100

**Occupation of head of households:** The data regarding the occupation of the household heads in Mundragi-3 Micro watershed is presented in Table 5. The results indicate that, 61.11 per cent of households heads were practicing agriculture, 36.11 per cent of the household heads were agricultural Labour and trade and business (2.78%).

**Table 5: Occupation of heads of households in Mundragi-3 micro-watershed**

Sl.No.	Particulars	LL (4)		MF (18)		SF (10)		SMF (2)		MDF (2)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Agriculture	0	0	12	67	6	60	2	100	2	100	22	61.11
2	Agricultural Labour	4	100	5	28	4	40	0	0	0	0	13	36.11
3	Trade & Business	0	0	1	5.6	0	0	0	0	0	0	1	2.78
Total		4	100	18	100	10	100	2	100	2	100	36	100

**Occupation of the members of the household:** The data regarding the occupation of the household members in Mundragi-3 Micro watershed is presented in Table 6. The results indicate that, agriculture was the major occupation for 37.50 per cent of the household members, 24.26 per cent were agricultural labour, 0.74 per cent were general labour, government sector and trade and business, 15.44 per cent were working in pursuing education, 16.91 per cent were involved as housewife and 3.68 per cent were children's.

**Table 6: Occupation of members of the household in Mundragi-3 micro-watershed**

Sl.No.	Particulars	LL (12)		MF (67)		SF (34)		SMF (14)		MDF (9)		All (136)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Agriculture	0	0	24	35.8	13	38.24	7	50	7	78	51	37.5
2	Agricultural Labour	7	58.3	15	22.4	9	26.47	2	14.29	0	0	33	24.3
3	General Labour	0	0	1	1.49	0	0	0	0	0	0	1	0.74
4	Government Service	0	0	1	1.49	0	0	0	0	0	0	1	0.74
5	Trade & Business	0	0	1	1.49	0	0	0	0	0	0	1	0.74
6	Student	3	25	14	20.9	4	11.76	0	0	0	0	21	15.4
7	Housewife	2	16.7	8	11.9	7	20.59	4	28.57	2	22	23	16.9
8	Children	0	0	3	4.48	1	2.94	1	7.14	0	0	5	3.68
Total		12	100	67	100	34	100	14	100	9	100	136	100



**Institutional Participation of household members:** The data regarding the institutional participation of the household members in Mundragi-3 Micro watershed is presented in Table 7. The results show that, out of the total family members in the households 100 per cent were not participating in any of the institutions.

**Table 7: Institutional Participation of household member in Mundragi-3 micro-watershed**

Sl.No.	Particulars	LL (12)		MF (67)		SF (34)		SMF (14)		MDF (9)		All (136)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	No Participation	12	100	67	100	34	100	14	100	9	100	136	100
	Total	12	100	67	100	34	100	14	100	9	100	136	100

**Type of house owned:** The data regarding the type of house owned by the households in Mundragi-3 Micro watershed is presented in Table 8. The results indicate that, 5.56 percent possess thatched house, 91.67 per cent of the households possess katcha house and 2.78 per cent possess pucca house.

**Table 8. Type of house owned by households in Mundragi-3 micro-watershed**

Sl.No.	Particulars	LL (4)		MF (18)		SF (10)		SMF (2)		MDF (2)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Thatched	1	25	0	0	1	10	0	0	0	0	2	5.56
2	Katcha	3	75	18	100	8	80	2	100	2	100	33	91.67
3	Pucca/RCC	0	0	0	0	1	10	0	0	0	0	1	2.78
	Total	4	100	18	100	10	100	2	100	2	100	36	100

**Durable assets owned by the households:** The data regarding the Durable Assets owned by the households in Mundragi-3 Micro watershed is presented in Table 9. The results shows that, 88.89 per cent possess TV, 8.33 per cent possess mixer grinder, 2.78 per cent possess motor cycle, 86.11 per cent possess mobile phones.

**Table 9. Durable assets owned by households in Mundragi-3 micro-watershed**

Sl.No.	Particulars	LL (4)		MF (18)		SF (10)		SMF (2)		MDF (2)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Television	4	100	16	89	8	80	2	100	2	100	32	88.89
2	Mixer/Grinder	0	0	1	5.6	0	0	1	50	1	50	3	8.33
3	Motor Cycle	0	0	1	5.6	0	0	0	0	0	0	1	2.78
4	Mobile Phone	4	100	15	83	8	80	2	100	2	100	31	86.11

**Average value of durable assets:** The data regarding the average value of durable assets owned by the households in Mundragi-3 Micro watershed is presented in Table 10. The result shows that, the average value of television was Rs.9000.00, mixer grinder was Rs.2666.00, motor cycle was Rs. 35000.00 and mobile phone was Rs.2934.00.

**Table 10. Average value of durable assets owned in Mundragi-3 micro-watershed**

Sl.No.	Particulars	LL (4)	MF (18)	SF (10)	SMF (2)	MDF (2)	All (36)
1	Television	9000	9062	8875	9000	9000	9000
2	Mixer/Grinder	0	2000	0	2000	4000	2666
3	Motor Cycle	0	35000	0	0	0	35000
4	Mobile Phone	3500	2863	2727	3200	3000	2934

**Farm implements owned:** The data regarding the farm implements owned by the households in Mundragi-3 Micro watershed is presented in Table 11. About 5.56 per cent of the households possess Bullock Cart, 22.22 per cent possess plough and 13.89 per cent possess Weeder.

**Table 11. Farm implements owned in Mundragi-3 micro-watershed**

Sl.No.	Particulars	LL (4)		MF (18)		SF (10)		SMF (2)		MDF (2)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Bullock Cart	0	0	1	5.56	1	10	0	0	0	0	2	5.56
2	Plough	0	0	3	16.7	2	20	1	50	2	100	8	22.22
3	Weeder	0	0	1	5.56	1	10	1	50	2	100	5	13.89

**Average value of farm implements:** The data regarding the average value of farm Implements owned by the households in Mundragi-3 Micro watershed is presented in Table 12. The results show that the average value of plough was Rs.1400.00, bullock Cart was Rs.17000.00 and weeder was Rs.50.00.

**Table 12. Average value of farm implements in Mundragi-3 micro-watershed**

Sl.No.	Particulars	Average Value (Rs.)					
		LL (4)	MF (18)	SF (10)	SMF (2)	MDF (2)	All (36)
1	Bullock Cart	0	10000	24000	0	0	17000
2	Plough	0	1466	1600	1200	1200	1400
3	Weeder	0	50	50	50	50	50

**Livestock possession by the households:** The data regarding the Livestock possession by the households in Mundragi-3 Micro watershed is presented in Table 13. The indicate that, 22.22 per cent of the households possess bullocks, 16.67 per cent possess local cow, 2.78 per cent possess buffalo, 5.56 per cent possess crossbred cow and 2.78 per cent possess goat.

**Table 13. Livestock possession by households in Mundragi-3 micro-watershed**

Sl.No.	Particulars	LL (4)		MF (18)		SF (10)		SMF (2)		MDF (2)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Bullock	0	0	4	22	2	20	1	50	1	50	8	22.22
2	Local cow	0	0	4	22	2	20	0	0	0	0	6	16.67
3	Crossbred cow	0	0	1	5.6	0	0	0	0	1	50	2	5.56
4	Buffalo	0	0	1	5.6	0	0	0	0	0	0	1	2.78
5	Goat	0	0	1	5.6	0	0	0	0	0	0	1	2.78

**Average Labour availability:** The data regarding the average labour availability in Mundragi-3 Micro watershed is presented in Table 14. The indicated that, own labour men available in the micro watershed was 1.97, women available in the micro watershed was 1.41, hired labour (men) available was 7.62 and hired labour (women) available was 6.79.

**Table 14. Average labour availability in Mundragi-3 micro-watershed**

Sl.No.	Particulars	LL (4)	MF (18)	SF (10)	SMF (2)	MDF (2)	All (36)
1	Hired labour Female	0	5.06	4.8	6.67	32.5	6.59
2	Own Labour Female	1	1.39	1.3	2	1.5	1.41
3	Own labour Male	2	1.94	1.7	2.67	2.5	1.97
4	Hired labour Male	0	6.11	5.8	10	32.5	7.62

**Adequacy of hired labour:** The data regarding the adequacy of hired labour in Mundragi-3 Micro watershed is presented in Table 15. The results indicate that, 97.22 per cent of the household opined that hired labour was adequate.

**Table 15. Adequacy of hired labour in Mundragi-3 micro-watershed**

Sl.No.	Particulars	LL (4)		MF (18)		SF (10)		SMF (2)		MDF (2)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Adequate	1	25	19	106	10	100	3	150	2	100	35	97.2

**Distribution of land (ha):** The data regarding the distribution of land (ha) in Mundragi-3 Micro watershed is presented in Table 16. The results indicate that, 15.31 ha (56.78%) of dry land and 11.65 ha (43.22 %) of irrigated land.

**Table 16. Distribution of land (ha) in Mundragi-3 micro-watershed**

Sl.No.	Particulars	LL (4)		MF (18)		SF (10)		SMF (2)		MDF (2)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Dry	0	0	7.6	98.38	4.99	48.43	2.71	69	0	0	15.31	56.78
2	Irrigated	0	0	0.13	1.62	5.31	51.57	1.22	31	4.99	100	11.65	43.22
Total		0	100	7.73	100	10.3	100	3.93	100	4.99	100	26.96	100

**Average value of land (ha):** The data regarding the average land value (Rs./ha) in Mundragi-3 Micro watershed is presented in Table 17. The results show that the average value of dry land was Rs.334383.93 and the average value of irrigated land was Rs.307999.31.

**Table 17. Average value of land (ha) in Mundragi-3 micro-watershed**

Sl.No.	Particulars	LL (4)	MF (18)	SF (10)	SMF (2)	MDF (2)	All (36)
1	Dry	0	383842.5	300486.6	258059.7	0	334383.9
2	Irrigated	0	717096.8	319802	328239.2	280226.9	307999.3

**Status of bore wells:** The data regarding the status of bore wells in Mundragi-3 Micro watershed is presented in Table 18. The results indicate that, there were 11 functioning bore wells among the sampled households in micro watershed.

**Table 18. Status of bore wells in Mundragi-3 micro-watershed**

Sl.No.	Particulars	LL (4)	MF (18)	SF (10)	SMF (2)	MDF (2)	All (36)
1	Functioning	0	2	6	1	2	11

**Source of irrigation:** The data regarding the source of irrigation in Mundragi-3 Micro watershed is presented in Table 19. The results that open well were major source of irrigation for 0.00 per cent of the households and bore well for 30.56 per cent of the households.

**Table 19. Source of irrigation in Mundragi-3 micro-watershed**

Sl.No.	Particulars	LL (4)		MF (18)		SF (10)		SMF (2)		MDF (2)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Bore Well	0	0	2	11.1	6	60	1	50	2	100	11	30.56

**Depth of water (Avg. In meters):** The data regarding the depth of water in Mundragi-3 Micro watershed is presented in Table 20. The results revealed that, the depth of bore well was 32.60 meter.

**Table 20. Depth of water (Avg. In meters) in Mundragi-3 micro-watershed**

Sl.No.	Particulars	LL (4)	MF (18)	SF (10)	SMF (2)	MDF (2)	All (36)
1	Bore Well	0	11.85	64.01	53.34	106.68	32.6

**Irrigated Area (ha):** The data regarding the irrigated area (ha) in Mundragi-3 Micro watershed is presented in Table 21. The results indicate that, the availability of irrigation water was used for kharif crops was 10.68 ha.

**Table 21. Irrigated Area (ha) in Mundragi-3 micro-watershed**

Sl.No.	Particulars	LL (4)	MF (18)	SF (10)	SMF (2)	MDF (2)	All (36)
1	Kharif	0	0.06	4.4	1.22	5	10.68
	Total	0	0.06	4.4	1.22	5	10.68

**Cropping pattern:** The data regarding the cropping pattern in Mundragi-3 Micro watershed is presented in Table 22. The results indicate that, farmers have grown groundnut (12.35 ha), red gram (7.89 ha), sorghum (1.74 ha), cotton (1.37 ha), green gram (0.64 ha) and paddy (0.06 ha).

**Table 22. Cropping pattern in Mundragi-3 micro-watershed**

Sl.No.	Particulars	LL (4)	MF (18)	SF (10)	SMF (2)	MDF (2)	All (36)
1	Kharif - Groundnut	0	3.39	5.66	1.22	2.08	12.35
2	Kharif - Red gram	0	1.75	3.43	2.71	0	7.89
3	Kharif - Sorghum	0	1.74	0	0	0	1.74
4	Kharif - Cotton	0	0.15	1.21	0	0	1.37
5	Kharif - Green gram	0	0.64	0	0	0	0.64
6	Kharif - Paddy	0	0.06	0	0	0	0.06
	Total	0	7.73	10.31	3.93	2.08	24.05

**Cropping intensity:** The data regarding the cropping intensity in Mundragi-3 Micro watershed is presented in Table 23. The results indicate that, the cropping intensity was 100.00 per cent.

**Table 23. Cropping intensity (%) in Mundragi-3 micro-watershed**

Sl.No.	Particulars	LL (4)	MF (18)	SF (10)	SMF (2)	MDF (2)	All (36)
1	Cropping Intensity	0	100	100	100	100	100

**Cost of Cultivation of Red gram:** The data regarding the cost of cultivation (Rs/ha) of Red gram in Mundragi-3 micro watershed is presented in Table 24.a. The results indicate that, the total cost of cultivation (Rs/ha) for Red gram was Rs. 37056.78. The gross income realized by the farmers was Rs. 69847.73. The net income from Red gram cultivation was Rs.32790.96, thus the benefit cost ratio was found to be 1:1.90.

**Table 24(a). Cost of Cultivation of Red gram in Mundragi-3 micro-watershed**

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
<b>I</b>	<b>Cost A1</b>				
1	Hired Human Labour	Man days	44.74	8947.25	24.14
2	Bullock	Pairs/day	1.36	813.22	2.19
3	Tractor	Hours	5.75	4602.6	12.42
4	Machinery	Hours	0	0	0
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	9.39	1304.4	3.52
7	FYM	Quintal	2.52	503.89	1.36
8	Fertilizer + micronutrients	Quintal	6.32	4936.75	13.32
9	Pesticides (PPC)	Kgs/liters	1.95	1951.47	5.27
10	Irrigation	Number	0	0	0
11	Repairs		0	0	0
12	Msc. Charges (Marketing costs etc)		0	0	0
13	Depreciation charges		0	0.03	0
14	Land revenue and Taxes		0	3.29	0.01
<b>II</b>	<b>Cost B1</b>				
16	Interest on working capital			1043.7	2.82
17	<b>Cost B1 = (Cost A1 + sum of 15 and 16)</b>			24106.61	65.05
<b>III</b>	<b>Cost B2</b>				
18	Rental Value of Land			333.33	0.9
19	<b>Cost B2 = (Cost B1 + Rental value)</b>			24439.95	65.95
<b>IV</b>	<b>Cost C1</b>				
20	Family Human Labour		39.58	9247.03	24.95
21	<b>Cost C1 = (Cost B2 + Family Labour)</b>			33686.98	90.91
<b>V</b>	<b>Cost C2</b>				
22	Risk Premium			1	0
23	<b>Cost C2 = (Cost C1 + Risk Premium)</b>			33687.98	90.91
<b>VI</b>	<b>Cost C3</b>				
24	Managerial Cost			3368.8	9.09
25	<b>Cost C3 = (Cost C2 + Managerial Cost)</b>			37056.78	100
<b>VII</b>	<b>Economics of the Crop</b>				
a.	Main Product	a) Main Product (q)		13.51	66725.77
		b) Main Crop Sales Price (Rs.)			4937.5
	By Product	e) Main Product (q)		7.14	3121.96
		f) Main Crop Sales Price (Rs.)			437.5
b.	Gross Income (Rs.)			69847.73	
c.	Net Income (Rs.)			32790.96	
d.	Cost per Quintal (Rs./q.)			2742.09	
e.	Benefit Cost Ratio (BC Ratio)			1:1.9	

**Cost of Cultivation of Cotton:** The data regarding the cost of cultivation (Rs/ha) of Cotton in Mundragi-3 micro watershed is presented in Table 24.b. The results indicate that, the total cost of cultivation (Rs/ha) for Cotton was Rs. 53774.67. The gross income realized by the farmers was Rs. 85041.67. The net income from Cotton cultivation was Rs.31267.00, thus the benefit cost ratio was found to be 1:1.60.

**Table 24(b). Cost of Cultivation of Cotton in Mundragi-3 micro-watershed**

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C <sub>3</sub>
<b>I</b>	<b>Cost A1</b>				
1	Hired Human Labour	Man days	74.85	15009.76	27.91
2	Bullock	Pairs/day	5.77	3498.81	6.51
3	Tractor	Hours	0.34	274.44	0.51
4	Machinery	Hours	0	0	0
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	6.87	6528.35	12.14
6	Seed Inter Crop	Kgs.	0	0	0
7	FYM	Quintal	1.51	301.89	0.56
8	Fertilizer + micronutrients	Quintal	11.64	9143.31	17
9	Pesticides (PPC)	Kgs / liters	2.56	2555.46	4.75
10	Irrigation	Number	0	0	0
11	Repairs		0	0	0
12	Msc. Charges (Marketing costs etc)		0	0	0
13	Depreciation charges		0	89.65	0.17
14	Land revenue and Taxes		0	3.29	0.01
<b>II</b>	<b>Cost B1</b>				
16	Interest on working capital			2223.6	4.14
17	<b>Cost B1 = (Cost A1 + sum of 15 and 16)</b>			39628.56	73.69
<b>III</b>	<b>Cost B2</b>				
18	Rental Value of Land			222.22	0.41
19	<b>Cost B2 = (Cost B1 + Rental value)</b>			39850.78	74.11
<b>IV</b>	<b>Cost C1</b>				
20	Family Human Labour		39.85	9034.28	16.8
21	<b>Cost C1 = (Cost B2 + Family Labour)</b>			48885.06	90.91
<b>V</b>	<b>Cost C2</b>				
22	Risk Premium			1	0
23	<b>Cost C2 = (Cost C1 + Risk Premium)</b>			48886.06	90.91
<b>VI</b>	<b>Cost C3</b>				
24	Managerial Cost			4888.61	9.09
25	<b>Cost C3 = (Cost C2 + Managerial Cost)</b>			53774.67	100
<b>VII</b>	<b>Economics of the Crop</b>				
a.	Main Product	a) Main Product (q)		17.01	85041.67
		b) Main Crop Sales Price (Rs.)			5000
b.	Gross Income (Rs.)			85041.67	
c.	Net Income (Rs.)			31267	
d.	Cost per Quintal (Rs./q.)			3161.67	
e.	Benefit Cost Ratio (BC Ratio)			1:1.6	

**Cost of Cultivation of Groundnut:** The data regarding the cost of cultivation (Rs/ha) of Groundnut in Mundragi-3 micro watershed is presented in Table 24.c. The results indicate, the total cost of cultivation (Rs/ha) for Groundnut was Rs.76635.07. The gross income realized by the farmers was Rs. 85112.24. The net income from Groundnut cultivation was Rs. 8477.18, thus the benefit cost ratio was found to be 1:1.10.

**Table 24(c). Cost of Cultivation of Groundnut in Mundragi-3 micro-watershed**

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
<b>I</b>	<b>Cost A1</b>				
1	Hired Human Labour	Man days	71.82	14050.15	18.33
2	Bullock	Pairs/day	6.45	3894.59	5.08
3	Tractor	Hours	3.22	2577.87	3.36
4	Machinery	Hours	0.21	171.55	0.22
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	91.46	16421.36	21.43
7	FYM	Quintal	4.82	963.63	1.26
8	Fertilizer + micronutrients	Quintal	9.93	8495.47	11.09
9	Pesticides (PPC)	Kgs/liters	3.54	3543.83	4.62
10	Irrigation	Number	0	0	0
11	Repairs		0	0	0
12	Msc. Charges (Marketing costs etc)		0	0	0
13	Depreciation charges		0	60.88	0.08
14	Land revenue and Taxes		0	3.29	0
<b>II</b>	<b>Cost B1</b>				
16	Interest on working capital			3531.03	4.61
17	<b>Cost B1 = (Cost A1 + sum of 15 and 16)</b>			53713.66	70.09
<b>III</b>	<b>Cost B2</b>				
18	Rental Value of Land			291.67	0.38
19	<b>Cost B2 = (Cost B1 + Rental value)</b>			54005.33	70.47
<b>IV</b>	<b>Cost C1</b>				
20	Family Human Labour		66.21	15661.91	20.44
21	<b>Cost C1 = (Cost B2 + Family Labour)</b>			69667.24	90.91
<b>V</b>	<b>Cost C2</b>				
22	Risk Premium			1	0
23	<b>Cost C2 = (Cost C1 + Risk Premium)</b>			69668.24	90.91
<b>VI</b>	<b>Cost C3</b>				
24	Managerial Cost			6966.82	9.09
25	<b>Cost C3 = (Cost C2 + Managerial Cost)</b>			76635.07	100
<b>VII</b>	<b>Economics of the Crop</b>				
a.	Main Product	a) Main Product (q)		16.93	74608.08
		b) Main Crop Sales Price (Rs.)			4406.44
	By Product	e) Main Product (q)		10.84	10504.17
		f) Main Crop Sales Price (Rs.)			969.19
b.	Gross Income (Rs.)			85112.24	
c.	Net Income (Rs.)			8477.18	
d.	Cost per Quintal (Rs./q.)			4526.15	
e.	Benefit Cost Ratio (BC Ratio)			1:1.1	

**Cost of Cultivation of Sorghum:** The data regarding the cost of cultivation (Rs/ha) of Sorghum in Mundragi-3 micro watershed is presented in Table 24.d. The results indicate that, the total cost of cultivation (Rs/ha) for Sorghum was Rs. 39040.33. The gross income realized by the farmers was Rs.83200.00. The net income from Sorghum cultivation was Rs. 44159.67, thus the benefit cost ratio was found to be 1:2.10.

**Table 24(d). Cost of Cultivation of Sorghum in Mundragi-3 micro-watershed**

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
<b>I</b>	<b>Cost A1</b>				
1	Hired Human Labour	Man days	46.53	8358.41	21.41
2	Bullock	Pairs/day	0	0	0
3	Tractor	Hours	5.17	4136.36	10.6
4	Machinery	Hours	0	0	0
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	80.95	11286.36	28.91
7	FYM	Quintal	2.3	460.91	1.18
8	Fertilizer + micronutrients	Quintal	3.43	2707.55	6.94
9	Pesticides (PPC)	Kgs / liters	1.12	1122.73	2.88
10	Irrigation	Number	0	0	0
11	Repairs		0	0	0
12	Msc. Charges (Marketing costs etc)		0	0	0
13	Depreciation charges		0	0.02	0
14	Land revenue and Taxes		0	3.29	0.01
<b>II</b>	<b>Cost B1</b>				
16	Interest on working capital			1869.43	4.79
17	<b>Cost B1 = (Cost A1 + sum of 15 and 16)</b>			29945.06	76.7
<b>III</b>	<b>Cost B2</b>				
18	Rental Value of Land			333.33	0.85
19	<b>Cost B2 = (Cost B1 + Rental value)</b>			30278.39	77.56
<b>IV</b>	<b>Cost C1</b>				
20	Family Human Labour		21.98	5211.82	13.35
21	<b>Cost C1 = (Cost B2 + Family Labour)</b>			35490.21	90.91
<b>V</b>	<b>Cost C2</b>				
22	Risk Premium			1	0
23	<b>Cost C2 = (Cost C1 + Risk Premium)</b>			35491.21	90.91
<b>VI</b>	<b>Cost C3</b>				
24	Managerial Cost			3549.12	9.09
25	<b>Cost C3 = (Cost C2 + Managerial Cost)</b>			39040.33	100
<b>VII</b>	<b>Economics of the Crop</b>				
a.	Main Product	a) Main Product (q)	9.28	37109.09	
		b) Main Crop Sales Price (Rs.)		4000	
	By Product	e) Main Product (q)	23.05	46090.91	
		f) Main Crop Sales Price (Rs.)		2000	
b.	Gross Income (Rs.)			83200	
c.	Net Income (Rs.)			44159.67	
d.	Cost per Quintal (Rs./q.)			4208.17	
e.	Benefit Cost Ratio (BC Ratio)			1:2.1	



**Cost of Cultivation of Green gram:** The data regarding the cost of cultivation (Rs/ha) of Green gram in Mundragi-3 micro watershed is presented in Table 24.e. The results indicate that, the total cost of cultivation (Rs/ha) for Green gram was Rs.108930.93. The gross income realized by the farmers was Rs. 113854.67. The net income from Green gram cultivation was Rs. 4923.74, thus the benefit cost ratio was found to be 1:1.02.

**Table 24(e). Cost of Cultivation of Green gram in Mundragi-3 micro-watershed**

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
<b>I</b>	<b>Cost A1</b>				
1	Hired Human Labour	Man days	138.61	28175.3	25.87
2	Bullock	Pairs/day	0	0	0
3	Tractor	Hours	22.77	18216.75	16.72
4	Machinery	Hours	0	0	0
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	14.52	2032.48	1.87
6	Seed Inter Crop	Kgs.	0	0	0
7	FYM	Quintal	10.94	2187.6	2.01
8	Fertilizer + micronutrients	Quintal	22.57	19666.53	18.05
9	Pesticides (PPC)	Kgs / liters	5.47	5469	5.02
10	Irrigation	Number	0	0	0
11	Repairs		0	0	0
12	Msc. Charges (Marketing costs etc)		0	0	0
13	Depreciation charges		0	0.11	0
14	Land revenue and Taxes		0	3.29	0
<b>II</b>	<b>Cost B1</b>				
16	Interest on working capital			3522.79	3.23
17	<b>Cost B1 = (Cost A1 + sum of 15 and 16)</b>			79273.86	72.77
<b>III</b>	<b>Cost B2</b>				
18	Rental Value of Land			333.33	0.31
19	<b>Cost B2 = (Cost B1 + Rental value)</b>			79607.19	73.08
<b>IV</b>	<b>Cost C1</b>				
20	Family Human Labour		85.71	19419.93	17.83
21	<b>Cost C1 = (Cost B2 + Family Labour)</b>			99027.12	90.91
<b>V</b>	<b>Cost C2</b>				
22	Risk Premium			1	0
23	<b>Cost C2 = (Cost C1 + Risk Premium)</b>			99028.12	90.91
<b>VI</b>	<b>Cost C3</b>				
24	Managerial Cost			9902.81	9.09
25	<b>Cost C3 = (Cost C2 + Managerial Cost)</b>			108930.93	100
<b>VII</b>	<b>Economics of the Crop</b>				
a.	Main Product	a) Main Product (q)	22.77	113854.67	
		b) Main Crop Sales Price (Rs.)		5000	
		f) Main Crop Sales Price (Rs.)		375	
b.	Gross Income (Rs.)			113854.67	
c.	Net Income (Rs.)			4923.74	
d.	Cost per Quintal (Rs./q.)			4783.77	
e.	Benefit Cost Ratio (BC Ratio)			1:1.02	

**Adequacy of fodder:** The data regarding the adequacy of fodder in Mundragi-3 Micro watershed is presented in Table 25. The results indicate that, 33.33 per cent of the households opined that dry fodder was adequate and 16.67 percent of them opined it was sufficient.

**Table 25. Adequacy of fodder in Mundragi-3 micro-watershed**

Sl.No.	Particulars	LL (4)		MF (18)		SF (10)		SMF (2)		MDF (2)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Adequate-Dry Fodder	0	0	6	33.33	5	50	1	50	0	0	12	33.33
2	Adequate-Green Fodder	0	0	3	16.67	3	30	0	0	0	0	6	16.67

**Average annual gross income:** The data regarding the annual gross income in Mundragi-3 Micro watershed is presented in Table 26. The results indicate that, the farmers have annual gross income of Rs. 114444.44 in micro-watershed, of which Rs. 57250.00 is from agriculture itself.

**Table 26. Average annual gross income in Mundragi-3 micro-watershed**

Sl.No.	Particulars	LL (4)	MF (18)	SF (10)	SMF (2)	MDF (2)	All (36)
1	Wage	63500	42222.2	64000	100000	40000	53722.2
2	Agriculture	0	50833.3	59200	152000	125000	57250
3	Dairy Farm	0	3333.33	0	0	0	1666.67
4	Goat Farming	0	3611.11	0	0	0	1805.56
	Income(Rs.)	63500	100000	123200	252000	165000	114444

**Average annual Expenditure:** The data regarding the average annual expenditure in Mundragi-3 Micro watershed is presented in Table 27. The results indicate that, the farmers have annual gross expenditure of Rs. 466352.94 in micro-watershed, of which Rs. 29027.78 is from agriculture itself.

**Table 27. Average annual Expenditure in Mundragi-3 micro-watershed**

Sl.No.	Particulars	LL (4)	MF (18)	SF (10)	SMF (2)	MDF (2)	All (36)
1	Wage	32000	29600	38500	85000	16500	29861.1
2	Agriculture	0	25352.9	30900	92500	60000	29027.8
3	Dairy Farm	0	35000	0	0	0	972.22
4	Goat Farming	0	21000	0	0	0	1166.67
	Total	32000	110953	69400	177500	76500	466353

**Horticulture species grown:** The data regarding horticulture species grown in Mundragi-3 Micro watershed is presented in Table 28. The results indicate that, the total number of horticultural trees grown (both field and backyard) by the sampled households were Jack fruit (2).

**Table 28. Horticulture species grown in Mundragi-3 micro-watershed**

Sl.No.	Particulars	LL (4)		MF (18)		SF (10)		SMF (2)		MDF (2)		All (36)	
		F	B	F	B	F	B	F	B	F	B	F	B
1	Jack fruit	0	0	0	0	0	0	0	0	2	0	2	0

\*F= Field B=Back Yard

**Forest species grown:** The data regarding forest species grown in Mundragi-3 Micro watershed is presented in Table 29. The results indicate that, households have planted 75 neem trees, 1 tamarind trees and 2 banyan trees together in both field and backyard.

**Table 29. Forest species grown in Mundragi-3 micro-watershed**

Sl.No.	Particulars	LL (4)		MF (18)		SF (10)		SMF (2)		MDF (2)		All (36)	
		F	B	F	B	F	B	F	B	F	B	F	B
1	Neem	0	0	15	9	21	3	12	0	15	0	63	12
2	Tamarind	0	0	1	0	0	0	0	0	0	0	1	0
3	Banyan	0	0	0	0	2	0	0	0	0	0	2	0

\*F= Field B=Back Yard

**Average additional investment capacity:** The data regarding average additional investment capacity in Mundragi-3 Micro watershed is presented in Table 30. The results indicate that, households have an average investment capacity of Rs. 2555.56 for land development, Rs. 833.33 for creation of irrigation facility and Rs.277.78 for adoption of improved crop production activities.

**Table 30. Average additional investment capacity of households in Mundragi-3 micro-watershed**

Sl.No.	Particulars	LL (4)	MF (18)	SF (10)	SMF (2)	MDF (2)	All (36)
1	Land development	0	3444.44	0	0	15000	2555.56
2	Irrigation facility	0	0	0	0	15000	833.33
3	Improved crop production	0	277.78	0	0	2500	277.78

**Source of funds for additional investment:** The data regarding source of funds for additional investment in Mundragi-3 Micro watershed is presented in Table 31. The results indicate that, the sources of finance raised from bank as a loan and from own sources for land development were 8.33 per cent and for improved crop production was 5.56 per cent.

**Table 31. Source of funds for additional investment in Mundragi-3 micro-watershed**

Sl.No	Item	Land development		Improved crop production	
		N	%	N	%
1	Loan from bank	3	8.33	2	5.56

**Marketing of agricultural produce:** The data regarding marketing of the agricultural produce in Mundragi-3 Micro watershed is presented in Table 32. The results indicated that, 100.00 percent of output of cotton was sold in the market; 91.07 percent of output of green gram was sold in the market; 92.11 percent of output of groundnut was sold in the market; 66.67 percent of output of paddy was sold in the market; 74 percent of output of red gram was sold in the market and 50 percent of output of sorghum was sold in the market

**Table 32. Marketing of agricultural produce in Mundragi-3 micro-watershed**

Sl.No	Crops	Output obtained (q)	Output retained (q)	Output sold (q)	Output sold (%)	Avg. Price obtained (Rs/q)
1	Cotton	45	0	45	100	7500
2	Green gram	504	45	459	91	5000
3	Groundnut	152	12	140	92	4406
4	Paddy	15	5	10	67	1200
5	Red gram	96	25	71	74	4938
6	Sorghum	16	8	8	50	4000

**Marketing channels used for sale of agricultural produce:** The data regarding marketing channels used for sale of agricultural produce in Mundragi-3 Micro watershed is

presented in Table 33. The results indicated that, 83.33 cent of the households have sold agricultural produce to the local/village merchants and 2.78 per cent of regulated market.

**Table 33. Marketing channels used for sale of agricultural produce in Mundragi-3 micro-watershed**

Sl.No	Particulars	LL (4)		MF (18)		SF (10)		SMF (2)		MDF (2)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Local/village Merchant	0	0	18	100	8	80	2	100	2	100	30	83.33
2	Regulated Market	0	0	0	0	1	10	0	0	0	0	1	2.78

**Mode of transport of agricultural produce:** The data regarding mode of transport of agricultural produce in Mundragi-3 Micro watershed is presented in Table 34. The results indicated that, 86.11 cent of the households have used tractor.

**Table 34. Mode of transport of agricultural produce in Mundragi-3 micro-watershed**

Sl.No.	Particulars	LL (4)		MF (18)		SF (10)		SMF (2)		MDF (2)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Tractor	0	0	18	100	9	90	2	100	2	100	31	86.11

**Incidence of soil and water erosion problems:** The data regarding incidence of incidence of soil and water erosion problems in Mundragi-3 Micro watershed is presented in Table 35. The results indicate that, 86.11 per cent of the households have experienced soil and water erosion problems.

**Table 35. Incidence of soil and water erosion problems in Mundragi-3 micro-watershed**

Sl.No.	Particulars	LL (4)		MF (18)		SF (10)		SMF (2)		MDF (2)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Soil and water erosion problems in the farm	0	0	17	94	10	100	2	100	2	100	31	86.11

**Interest towards soil testing:** The data regarding Interest shown towards soil testing in Mundragi-3 Micro watershed is presented in Table 36. The results indicated that, 80.56 per cent of the households were interested towards soil testing.

**Table 36. Interest regarding soil testing in Mundragi-3 micro-watershed**

Sl.No.	Particulars	LL (4)		MF (18)		SF (10)		SMF (2)		MDF (2)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Interest in soil test	0	0	16	89	9	90	2	100	2	100	29	80.56

**Usage pattern of fuel for domestic use:** The data on usage pattern of fuel for domestic use in Mundragi-3 Micro watershed is presented in Table 37. The results indicated that, firewood was the major source of fuel for domestic use for 100.00 per cent of the households.

**Table 37. Usage pattern of fuel for domestic use in Mundragi-3 micro-watershed**

Sl.No.	Particulars	LL (4)		MF (18)		SF (10)		SMF (2)		MDF (2)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Fire Wood	4	100	18	100	10	100	2	100	2	100	36	100

**Source of drinking water:** The data on source of drinking water in Mundragi-3 Micro watershed is presented in Table 38. The results indicated that, piped supply of water was the major source for drinking water for 97.22 per cent of the households.

**Table 38. Source of drinking water in Mundragi-3 micro-watershed**

Sl.No.	Particulars	LL (4)		MF (18)		SF (10)		SMF (2)		MDF (2)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Piped supply	4	100	18	100	9	90	2	100	2	100	35	97.22

**Source of light:** The data on source of light in Mundragi-3 Micro watershed is presented in Table 39. The results indicated that, electricity was the major source of light for 100.00 per cent of the households.

**Table 39. Source of light in Mundragi-3 micro-watershed**

Sl.No.	Particulars	LL (4)		MF (18)		SF (10)		SMF (2)		MDF (2)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Electricity	4	100	18	100	10	100	2	100	2	100	36	100

**Existence of sanitary toilet facility:** The data on availability of toilet facility in Mundragi-3 Micro watershed is presented in Table 40. The results indicated that, 77.78 per cent of the households possess toilets.

**Table 40. Existence of sanitary toilet facility in Mundragi-3 micro-watershed**

Sl.No.	Particulars	LL (4)		MF (18)		SF (10)		SMF (2)		MDF (2)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Sanitary toilet facility	1	25	16	89	8	80	2	100	1	50	28	77.8

**Possession of PDS card:** The data regarding possession of PDS card in Mundragi-3 Micro watershed is presented in Table 41. The results indicated that, 100.00 per cent of the households possessed BPL card.

**Table 41. Possession of PDS card in Mundragi-3 micro-watershed**

Sl.No.	Particulars	LL (4)		MF (18)		SF (10)		SMF (2)		MDF (2)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	BPL	4	100	18	100	10	100	2	100	2	100	36	100

**Adequacy of food items:** The data regarding adequacy of food items in Mundragi-3 Micro watershed is presented in Table 42. The results indicated that, the extent of adequacy of food items for cereals, pulses, Oilseeds and vegetables were 100.00, 83.33, 2.78, 88.89 per cent respectively, similarly for Fruits (11.11%), milk (97.22%), Egg (100.00%) and Meat (97.22%).

**Table 42. Adequacy of food items in Mundragi-3 micro-watershed**

Sl.No.	Particulars	LL (4)		MF (18)		SF (10)		SMF (2)		MDF (2)		LF (0)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Cereals	4	100	18	100	10	100	2	100	2	100	0	0	36	100
2	Pulses	4	100	14	77.8	8	80	2	100	2	100	0	0	30	83.33
3	Oilseed	0	0	0	0	1	10	0	0	0	0	0	0	1	2.78
4	Vegetables	4	100	16	88.9	9	90	2	100	1	50	0	0	32	88.89
5	Fruits	0	0	1	5.56	3	30	0	0	0	0	0	0	4	11.11
6	Milk	4	100	17	94.4	10	100	2	100	2	100	0	0	35	97.22
7	Egg	4	100	18	100	10	100	2	100	2	100	0	0	36	100
8	Meat	4	100	18	100	9	90	2	100	2	100	0	0	35	97.22

**Inadequacy of food items:** The data regarding in adequacy of food items in Mundragi-3 Micro watershed is presented in Table 43. The results indicated that, the extent of in adequacy of food items for pulses, Oilseeds and vegetables were 16.67, 94.44 and 11.11 per cent respectively, similarly for fruits (86.11%) and milk (2.78%).

**Table 43. Inadequacy of food items in Mundragi-3 micro-watershed**

Sl.No.	Particulars	LL (4)		MF (18)		SF (10)		SMF (2)		MDF (2)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Pulses	0	0	4	22.2	2	20	0	0	0	0	6	16.67
2	Oilseed	4	100	17	94.4	9	90	2	100	2	100	34	94.44
3	Vegetables	0	0	2	11.1	1	10	0	0	1	50	4	11.11
4	Fruits	4	100	16	88.9	7	70	2	100	2	100	31	86.11
5	Milk	0	0	1	5.56	0	0	0	0	0	0	1	2.78

**Farming constraints:** The data regarding farming constraints experienced by households in Mundragi-3 Micro watershed is presented in Table 44. The results indicated that, lower fertility status of the soil was the constraint experienced by (88.89 %) per cent of the households, wild animal menace on farm field (8.33%), frequent incidence of pest and diseases (77.78%), inadequacy of irrigation water (8.33%), high cost of fertilizers and plant protection chemicals (77.78%), high rate of interest on credit (5.56%), low price for the agricultural commodities (80.56 %), lack of marketing facilities in the area (75.00%), inadequate extension services (8.33 %), lack of transport for safe transport of the agricultural produce to the market (80.56%).

**Table 44a. Farming constraints experienced in Mundragi-3 micro-watershed**

SN	Particulars	LL (4)		MF (18)		SF (10)		SMF (2)		MDF (2)		All (36)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Lower fertility status of the soil	0	0	18	100	10	100	2	100	2	100	32	88.89
2	Wild animal menace on farm field	0	0	0	0	3	30	0	0	0	0	3	8.33
3	Frequent incidence of pest and diseases	0	0	17	94.44	7	70	2	100	2	100	28	77.78
4	Inadequacy of irrigation water	0	0	1	5.56	2	20	0	0	0	0	3	8.33
5	High cost of Fertilizers and plant protection chemicals	0	0	15	83.33	10	100	2	100	1	50	28	77.78
6	High rate of interest on credit	0	0	2	11.11	0	0	0	0	0	0	2	5.56
7	Low price for the agricultural commodities	0	0	16	88.89	9	90	2	100	2	100	29	80.56
8	Lack of marketing facilities in the area	0	0	16	88.89	7	70	2	100	2	100	27	75
9	Inadequate extension services	0	0	2	11.11	1	10	0	0	0	0	3	8.33
10	Lack of transport for safe transport of the Agril produce to the market.	0	0	16	88.89	9	90	2	100	2	100	29	80.56

**SUMMARY AND IMPLICATIONS**

In order to assess the socio-economic condition of the farmers in the watershed 36 households located in the micro watershed were interviewed for the survey. The study was conducted in Mundragi-3 micro-watershed (Belagiri sub-watershed, Yadgiri taluk & District) is located at North latitude 16<sup>0</sup> 46' 39.624" and 16<sup>0</sup> 45' 25.814" and East longitude 77<sup>0</sup> 12' 9.516" and 77<sup>0</sup> 9' 33.277" covering an area of about 598.12 ha bounded by unde Mundaragi and Yadhagiri B Villages.

Socio-economic analysis indicated that, out of the total sample of 36 respondents, - 18 (50.00%) were marginal, 10(27.78%) were small and 2 (5.56%) were semi medium, 2 (5.56%) were medium farmers. The population characteristics of households indicated that, there were 78 (57.35%) men and 58 (42.65%) were women. Majority of the respondents (47.06%) were in the age group of 35-60 years. Education level of the sample households indicated that, majority there were 66.91 per cent illiterates and only 3.68 per cent attained graduation. About, 61.11 per cent of household heads practicing agriculture and 36.11 per cent of the household heads were engaged as agricultural labourers. Agriculture was the major occupation for 37.50 per cent of the household members.

In the study area, 91.67 per cent of the households possess katcha house and 2.78 per cent possess pucca house. The durable assets owned by the households showed that, 88.89 per cent possess TV, 8.33 per cent possess mixer grinder and 86.11 per cent possess mobile phones. Farm implements owned by the households indicated that, 22.22 per cent of the households possess plough. Regarding livestock possession by the households, 16.67 per cent possess local cow and 2.78 per cent possess buffalo respectively.

The average labour availability in the study area showed that, own labour men available in the micro watershed was 1.97, women available in the micro watershed was 1.41, hired labour (men) available was 7.62 and hired labour (women) available was 6.79.

Out of the total land holding of the sample respondents (26.96 ha), 56.78 per cent of the area is under dry condition and the remaining 43.22 per cent area is irrigated land. Bore well was the major source of irrigation for 30.56 per cent of the households. The major crops grown by sample farmers are Red gram, Cotton, Groundnut, Sorghum and Green gram and cropping intensity was recorded as 100.00 per cent.

The per hectare cost of cultivation for Red gram, Cotton, Groundnut, Sorghum and Green gram was Rs.37056.78 , 53774.67, 76635.07, 39040.33 and 108930.93 with benefit cost ratio of 1:1.90, 1: 1.60, 1: 1.10, 1: 2.10 and 1:1.02 respectively.

Further, 33.33 per cent of the households opined that dry fodder was adequate and 16.67 per cent of the households have opined that the green fodder was adequate.

The average annual gross income of the farmers was Rs. 114444.44 in micro-watershed, of which Rs. 57250.00 comes from agriculture.

The total number of horticultural trees grown (both field and backyard) by the sampled households were Jack fruit (2) and forest species have planted 75 neem trees, 1 tamarind trees and 2 banyan trees together in both field and backyard.

Households have an average investment capacity of Rs. 2555.56 for land development, Rs. 833.33 for creation of irrigation facility and Rs.277.78 for adoption of improved crop production activities. Source of funds raised from bank as a loan and from own sources for land development were 8.33 per cent and for improved crop production was 5.56 per cent.

Regarding marketing channels, 83.33 per cent of the households have sold agricultural produce to the local/village merchants, while, 2.78 per cent have sold by Agents/Traders. Further, 86.11 per cent of the households have used tractor for the transport of agriculture commodity.

Majority of the farmers (86.11 %) have experienced soil and water erosion problems in the watershed and 80.56 per cent of the households were interested towards soil testing.

Firewood connection was the major source of fuel for domestic use for 100.00 per cent of the households. Piped supply was the major source for drinking water for 97.22 per cent of the households. Electricity was the major source of light for 100.00 per cent of the households. In the study area, 77.78 per cent of the households possess toilet facility. Regarding possession of PDS card, 100.00 per cent of the households possessed BPL card and 0.00 per cent do not possess PDS card. Cereals (100.00%), pulses (83.33%), oilseeds (2.78%) were adequate for consumption.

Farming constraints experienced by households in the micro watersheds were lower fertility status of the soil (88.89%) wild animal menace on farm field (8.33%), frequent incidence of pest and diseases (77.78%), inadequacy of irrigation water (8.33%), high cost of fertilizers and plant protection chemicals (77.78%), high rate of interest on credit (5.56%), low price for the agricultural commodities (80.56%), lack of marketing facilities in the area (75.00%), inadequate extension services (8.33%), lack of transport for safe transport of the agricultural produce to the market (80.56%).

#### **Implications of the survey**

- ✓ Result indicated that, there were 66.91 per cent were illiterate hence, extension methodologies such as demonstration, street play, drama, video shows will be effective in dissemination of the technologies in the micro watershed.
- ✓ The data indicate that, 91.67 per cent of the households possess katcha house. Hence, the development department while implementing the watershed plan should focus on agriculture to enhance the productivity of major crops in the area to increase the income of the farmers.
- ✓ Results indicated that the local institutional participation of the household members in the micro watershed is minimal hence, activities like membership campaign, awareness



creation about the benefits of membership in local institutions and strengths of organized groups must be conveyed.

- ✓ Majority of the households in the watershed have experience in use of mobile phones, and television hence, these mass media can be effectively utilized for transfer of technology as well as for information dissemination.
- ✓ The farm machinery/implement possession in the micro watershed was found to be minimum the reasons may lack of knowledge or lack of financial ability which can be addressed through training on use of different farm implements, providing information on different sources of finance for purchase of farm implements.
- ✓ The possession of livestock such as crossbred cow found is less hence, farmers must be made aware of the benefits of crossbred cow in increased milk production.
- ✓ The possession of livestock such as sheep, goat and poultry was found to be low hence, farmers may be informed the role of subsidiary enterprises in enhancing the income and information on financial support for subsidiary activities.
- ✓ Households possess 15.31ha (56.78 %) of dry land and 11.65ha (43.22 %) of irrigated land hence, the availability of the dry land agricultural technologies such as short duration crops, high yielding drought resistance crop varieties, drip irrigation technology and subsidy information will be helpful for the farmers to enhance the productivity of land and as well as farmers income.
- ✓ Few of the bore well in micro watershed found non functional hence, farmers may be trained on possibility of bore well rejuvenation.
- ✓ Bore well was major source of irrigation for 30.56 per cent of the households. hence, in order to increase the area under irrigation as well as to increase the water use efficiency farmers may trained on drip irrigation and provide the information on subsidy for drip irrigation equipment's along with the information on different agencies which provides the financial assistance for drip irrigation.
- ✓ The total number of horticultural trees grown (both field and backyard) by the sampled households were Jack fruit (2) and forest species have planted 75 neem trees, 1 tamarind trees and 2 banyan trees together in both field and backyard. Hence, production technologies related to these crops can be made available to the farmers for better adoption.
- ✓ The cropping intensity in the micro watershed was found to be (100.00 %) hence, care must be taken by the implementing agency to bring uncultivated land into cultivation through suitable measures.
- ✓ Many of the household members have borrowed loan from cooperative banks which has higher rate of interest hence, farmers may be sensitized on the different sources of credit with lesser interest rate such SHGs etc.
- ✓ The results indicated the non availability of both green and dry fodder throughout the year hence, fodder development activities can be taken up in the micro watershed.

- ✓ The average annual gross income of the households Rs.57250.00 from agriculture and Rs. 53722.22 from wages and. Agriculture was found to be the major source of income for households hence; the development activities should focus on productivity enhancement, marketing arrangements and agricultural technology dissemination to have a direct impact on the farmers.
- ✓ The cultivation of forest species is found minimal hence, information and production technology related to agro-forestry and integrated farming system.
- ✓ The data indicated that, 86.11 per cent of the households have experienced soil and water erosion problems. Hence, those farmers who reported the soil and water erosion problems may be given attention while implementation of the watershed development plan.
- ✓ The data indicated that, 80.56 per cent of the households have interest in soil testing hence, farmers must be provided with the information on various institutions which are involved in soil testing for the benefit of the farmers.
- ✓ Except summer ploughing the adoption of other soil and water conservation structures is minimum hence, the farmers in the micro watershed should be sensitized on the use of different conservation structures for soil water conservation.
- ✓ Cereals and pulses found be adequate for per cent of the households respectively hence, farm households and the farm women must be trained on importance of balanced nutrition and role of vegetable, milk, egg, meat in balanced diet.
- ✓ Lower fertility status of the soil (88.89%), wild animal menace on farm field (8.33%), frequent incidence of pest and diseases (77.78%), high cost of fertilizers and plant protection chemicals (77.78%), high rate of interest on credit (5.56%), low price for the agricultural commodities (80.56%), lack of marketing facilities in the area (75.00%), inadequate extension services (8.33%), lack of transport for safe transport of the agricultural produce to the market (80.56%) were the major farming constraints experienced hence, these constraints must be addressed immediately for the welfare of the farmers. Awareness to be created among the farmers to approach nearest KVKs/RSKs and other developmental departments for technical and for subsidized inputs and utilize the well established regulated markets, approaching the contract firms, direct markets to avoid the involvement of middlemen.