



हर कदम, हर डगर  
किसानों का हमसफर  
भारतीय कृषि अनुसंधान परिषद

*AgriSearch with a human touch*

**LAND RESOURCE INVENTORY AND SOCIO-ECONOMIC STATUS OF  
FARM HOUSEHOLDS FOR WATERSHED PLANNING AND  
DEVELOPMENT**

**HIMALAPUR (4D2D6A2a) MICROWATERSHED**

**Gurumitkal Hobli, Yadgir Taluk and District, Karnataka**

**Karnataka Watershed Development Project – II**

**SUJALA – III**

**World Bank funded Project**



**ICAR – NATIONAL BUREAU OF SOIL SURVEY AND LAND USE PLANNING**



ICAR - NBSS & LUP



**WATERSHED DEVELOPMENT DEPARTMENT  
GOVT. OF KARNATAKA, BANGALORE**



## **About ICAR - NBSS&LUP**

The ICAR-National Bureau of Soil Survey and Land Use Planning (ICAR-NBSS&LUP), Nagpur, a premier Institute of the Indian Council of Agricultural Research (ICAR), was set up during 1976 with the objective to prepare soil resource maps at national, state and district levels and to provide research inputs in soil resource mapping and its applications, land evaluation, land use planning, land resource management, and database management using GIS for optimising land use on different kinds of soils in the country.

The Bureau has been engaged in carrying out soil resource survey, agro-ecological and soil degradation mapping at the country, state and district levels for qualitative assessment and monitoring the soil health towards viable land use planning. The research activities have resulted in identifying the soil potentials and problems, and the various applications of the soil surveys with the ultimate objective of sustainable agricultural development. The Bureau has the mandate to correlate and classify soils of the country and maintain a National Register of all the established soil series. The Institute is also imparting in-service training to staff of the soil survey agencies in the area of soil survey, land evaluation and soil survey interpretations for land use planning. The Bureau in collaboration with Panjabrao Krishi Vidyapeeth, Akola is running post-graduate teaching and research programme in land resource management, leading to M.Sc. and Ph.D. degrees.

**Citation:** Rajendra Hegde, Ramesh kumar, S.C., B.A. Dhanorkar, S. Srinivas, M.Lalitha, K.V. Niranjana, R.S. Reddy and S.K. Singh (2019). "Land Resource Inventory and Socio-Economic Status of Farm Households for Watershed Planning and Development of Himalapur (4D2D6A2a) Microwatershed, Gurumitkal Hobli, Yadgir Taluk and District, Karnataka", ICAR-NBSS&LUP Sujala MWS Publ.303, ICAR – NBSS & LUP, RC, Bangalore. p.123 & 26.

### **TO OBTAIN COPIES,**

**Please write to:**

**Director, ICAR - NBSS & LUP,**

Amaravati Road, NAGPUR - 440 033, India

Phone : (0712) 2500386, 2500664, 2500545 (O)

Telefax : 0712-2522534

E-Mail : director@nbsslup.ernet.in

Website URL : nbsslup.in

Or

**Head, Regional Centre, ICAR - NBSS&LUP, Hebbal, Bangalore - 560 024**

Phone : (080) 23412242, 23510350 (O)

Telefax : 080-23510350

E-Mail : nbssrcb@gmail.com

ICAR-NBSS&LUP Sujala MWS Publ.303



हर कदम, हर डगर  
किसानों का हमसफर  
भारतीय कृषि अनुसंधान परिषद

*Agr*search with a human touch

**LAND RESOURCE INVENTORY AND SOCIO-ECONOMIC  
STATUS OF FARM HOUSEHOLDS FOR WATERSHED  
PLANNING AND DEVELOPMENT**

**HIMALAPUR (4D2D6A2a) MICROWATERSHED**

**Gurumitkal Hobli, Yadgir Taluk and District, Karnataka**

**Karnataka Watershed Development Project – II**

**Sujala-III**

**World Bank funded Project**



**ICAR – NATIONAL BUREAU OF SOIL SURVEY AND LAND USE  
PLANNING**



ICAR - NBSS & LUP



**WATERSHED DEVELOPMENT DEPARTMENT, GOVT. OF  
KARNATAKA, BANGALORE**





## PREFACE

In Karnataka, as in other Indian States, the livelihoods of rural people are intertwined with farming pursuits. The challenges in agriculture are seriously threatening the livelihood of a large number of farmers as they have been practicing farming in contextual factors beyond their control. Climatic factors are the most important ones and have become much more significant in recent times due to rapid climate changes induced by intensive anthropogenic activities affecting our ecosystem in multiple ways. Climate change has become the reality, it is happening and efforts to evolve and demonstrate climate resilient technologies have become essential. Due to the already over stressed scenario of agrarian sector, the climate change is resulting in manifold increase in the complexities, pushing the rural mass to face more and more unpredictable situations. The rising temperatures and unpredictable rainfall patterns are going to test seriously the informed decisions farmers have to make in order to survive in farming and sustain their livelihood.

It is generally recognized that impacts of climate change shall not be uniform across the globe. It is said that impact of climate change is more severe in South Asia. Based on the analysis of meteorological data, it is predicted that in India, there will be upward trend in mean temperature, downward trend in relative humidity, annual rainfall and number of wet days in a year. Also, in general, phenomena like erratic monsoon, spread of tropical diseases, rise in sea levels, changes in availability of fresh water, frequent floods, droughts, heat waves, storms and hurricanes are predicted. Each one of these adverse situations are already being experienced in various parts of India and also at the global level. Decline in agricultural productivity of small and marginal farmers becoming more vulnerable is already witnessed.

In Karnataka, more than 60 per cent of the population live in rural areas and depend on agriculture and allied activities for their livelihood. Though the state has achieved significant progress in increasing the yield of many crops, there is tremendous pressure on the land resources due to the growing and competing demands of various land uses. This is reflected in the alarming rate of land degradation observed. Already more than 50 per cent of the area is affected by various forms of degradation. If this trend continues, the sustainability of the fragile ecosystem will be badly affected. The adverse effects of change in the climatic factors are putting additional stress on the land resources and the farmers dependent on this.

The natural resources (land, water and vegetation) of the state need adequate and constant care and management, backed by site-specific technological interventions and investments particularly by the government. Detailed database pertaining to the nature of the land resources, their constraints, inherent potentials and suitability for various land

based rural enterprises, crops and other uses is a prerequisite for preparing location-specific action plans, which are in tune with the inherent capability of the resources. Any effort to evolve climate resilient technologies has to be based on the baseline scientific database. Then only one can expect effective implementation of climate resilient technologies, monitor the progress, make essential review of the strategy, and finally evaluate the effectiveness of the implemented programs. The information available at present on the land resources of the state are of general nature and useful only for general purpose planning. Since the need of the hour is to have site-specific information suitable for farm level planning and detailed characterization and delineation of the existing land resources of an area into similar management units is the only option.

ICAR-NBSS&LUP, Regional Centre, Bangalore has taken up a project sponsored by the Karnataka Watershed Development Project-II, (Sujala-III), Government of Karnataka funded by the World Bank under Component-1 Land Resource Inventory. This study was taken up to demonstrate the utility of such a database in reviewing, monitoring and evaluating all the land based watershed development programs on a scientific footing. To meet the requirements of various land use planners at grassroots level, the present study on “Land Resource Inventory and Socio-Economic Status of Farm Households for Watershed Planning and Development of Honageri-2Microwatershed, Yadgir Taluk and District, Karnataka” for integrated development was taken up in collaboration with the State Agricultural Universities, IISC, KRSRAC, KSNDMC as Consortia partners. The project provides detailed land resource information at cadastral level (1:7920 scale) for all the plots and socio-economic status of farm households covering thirty per cent farmers randomly selected representing landed and landless class of farmers in the microwatershed. The project report with the accompanying maps for the Microwatershed will provide required detailed database for evolving effective land use plan, alternative land use options and conservation plans for the planners, administrators, agricultural extension personnel, KVK officials, developmental departments and other land users to manage the land resources in a sustainable manner.

It is hoped that this database will be useful to the planners, administrators and developmental agencies working in the area in not only for formulating location specific developmental schemes but also for their effective monitoring at the village/watershed level.

Nagpur  
Date: 09-08-2019

**S.K. SINGH**  
Director, ICAR - NBSS&LUP Nagpur

## Contributors

<b>Dr. Rajendra Hegde</b> Principal Scientist, Head & Project Leader, Sujala-III Project ICAR-NBSS&LUP, Regional Centre, Bangalore	<b>Dr. S.K.Singh</b> Director, ICAR-NBSS&LUP Coordinator, Sujala-III Project Nagpur
<b>Soil Survey, Mapping &amp; Report Preparation</b>	
Dr. B.A. Dhanorkar	Sh. R.S. Reddy
Dr. K.V. Niranjana	Sh. Somashekar T N
	Smt. Chaitra, S.P.
	Dr. Gopali bardhan
	Ms. Arpitha G.M
	Dr. Mahendra kumar M.B
<b>Field Work</b>	
Sh. C.BacheGowda	Sh. Mahesh, D.B.
Sh. Somashekar	Sh. Ashok S Sindagi
Sh. M. Jayaramaiah	Sh. Veerabhadrapa B.
Sh. Paramesha, K.	Sh. Shankarappa
Sh. B. M. Narayana Reddy	Sh. Anand
	Sh. Arun N Kambar.
	Sh Kamalesh Awate
	Sh. Sharaan Kumar Huppar
	Sh. Yogesh H.N.
	Sh. Kalaveerachari R Kammar
<b>GIS Work</b>	
Dr. S.Srinivas	Sh. A.G.Devendra Prasad
Sh. D.H.Venkatesh	Sh. Prakashanaik, M.K.
Smt.K.Sujatha	Sh. Abhijith Sastry, N.S.
Smt. K.V.Archana	Sh. Sudip Kumar Suklabaidya
Sh. N. Maddileti	Sh. Avinash, K.N.
	Sh. Amar Suputhra, S
	Sh. Deepak, M.J.
	Smt. K.Karunya Lakshmi
	Ms. Seema, K.V.
	Ms. A. Rajab Nisha

<b>Laboratory Analysis</b>	
Dr. K.M.Nair	Ms. Steffi Peter
Smt. Arti Koyal	Ms. Thara, V.R
Smt. Parvathy	Ms. Roopa, G.
	Ms. Swati, H.
	Sh. Shantaveera Swami
	Ms. Shwetha, N.K.
	Smt. Ishrat Haji
	Ms. P. Pavan Kumari
	Ms. Padmaja
	Ms. Veena, M.
<b>Socio-Economic Analysis</b>	
Dr. S.C. Ramesh Kumar	Sh. M.K. Prakashanaik
	Ms. Karuna V. Kulkarni
	Mrs. Sowmya A.N
	Sh. Vinod R
	Sh. Basavaraja
	Sh. Vijay Kumar Lamani
	Ms. Sowmya K.B
	Mrs. Prathibha, D.G
	Sh. Rajendra,D
<b>Soil &amp; Water Conservation</b>	
Sh. Sunil P. Maske	
<b>Watershed Development Department, GoK, Bangalore</b>	
Sh. Rajeev Ranjan IFS Project Director & Commissioner, WDD	Dr. A. Natarajan NRM Consultant, Sujala-III Project
Dr. S.D. Pathak IFS Executive Director & Chief Conservator of Forests, WDD	



# **PART-A**

## **LAND RESOURCE INVENTORY**



## Contents

Preface		
Contributors		
Executive Summary		
Chapter 1	Introduction	1
Chapter 2	Geographical Setting	3
2.1	Location and Extent	3
2.2	Geology	3
2.3	Physiography	4
2.4	Drainage	4
2.5	Climate	4
2.6	Natural Vegetation	6
2.7	Land Utilization	6
Chapter 3	Survey Methodology	10
3.1	Base maps	10
3.2	Image Interpretation for Physiography	10
3.3	Field Investigation	13
3.4	Soil Mapping	14
3.5	Land Management Units	14
3.6	Laboratory Characterization	15
Chapter 4	The Soils	19
4.1	Soils of granite gneiss landscape	19
Chapter 5	Interpretation for Land Resource Management	29
5.1	Land Capability Classification	29
5.2	Soil Depth	31
5.3	Surface Soil Texture	32
5.4	Soil Gravelliness	33
5.5	Available Water Capacity	34
5.6	Soil Slope	35
5.7	Soil Erosion	36
Chapter 6	Fertility Status	39
6.1	Soil Reaction (pH)	39
6.2	Electrical Conductivity (EC)	39
6.3	Organic Carbon (OC)	39
6.4	Available Phosphorus	41
6.5	Available Potassium	41
6.6	Available Sulphur	41
6.7	Available Boron	42
6.8	Available Iron	42
6.9	Available Manganese	42
6.10	Available Copper	42
6.11	Available Zinc	46

Chapter 7	Land Suitability for Major Crops	47
7.1	Land suitability for Sorghum	47
7.2	Land suitability for Maize	48
7.3	Land suitability for Bajra	49
7.4	Land suitability for Groundnut	50
7.5	Land suitability for Sunflower	51
7.6	Land suitability for Redgram	52
7.7	Land suitability for Bengal gram	53
7.8	Land suitability for Cotton	54
7.9	Land suitability for Chilli	55
7.10	Land suitability for Tomato	56
7.11	Land suitability for Brinjal	57
7.12	Land suitability for Onion	58
7.13	Land suitability for Bhendi	59
7.14	Land suitability for Drumstick	60
7.15	Land suitability for Mango	61
7.16	Land suitability for Guava	62
7.17	Land suitability for Sapota	63
7.18	Land Suitability for Pomegranate	64
7.19	Land Suitability for Musambi	65
7.20	Land Suitability for Lime	66
7.21	Land Suitability for Amla	67
7.22	Land Suitability for Cashew	68
7.23	Land Suitability for Jackfruit	69
7.24	Land Suitability for Jamun	70
7.25	Land Suitability for Custard apple	71
7.26	Land Suitability for Tamarind	72
7.27	Land Suitability for Mulberry	73
7.28	Land Suitability for Marigold	74
7.29	Land Suitability for Chrysanthemum	75
7.30	Land Management Units (LMUs)	107
7.31	Proposed Crop Plan for Himalapur Microwatershed	108
Chapter 8	Soil Health Management	111
Chapter 9	Soil and Water conservation Treatment Plan	115
9.1	Treatment Plan	116
9.2	Recommended Soil and Water Conservation measures	119
9.3	Greening of Microwatershed	120
	References	123
	Appendix I	I-XIV
	Appendix II	XV-XXVIII
	Appendix III	XXIX-XL

## LIST OF TABLES

2.1	Mean Monthly Rainfall, PET, 1/2 PET at Yadgir Taluk & District	5
2.2	Land Utilization in Yadgir district	7
3.1	Differentiating Characteristics used for Identifying Soil Series	14
3.2	Soil map unit description of Himalapur Microwatershed	15
4.1	Physical and Chemical Characteristics of Soil Series identified in Himalapur microwatershed	23
7.1	Soil-Site Characteristics of Himalapur Microwatershed	77
7.2	Land suitability for Sorghum	78
7.3	Land suitability for Maize	79
7.4	Land suitability for Bajra	80
7.5	Land suitability for Groundnut	81
7.6	Land suitability for Sunflower	82
7.7	Land suitability for Redgram	83
7.8	Land suitability for Bengal gram	84
7.9	Land suitability for Cotton	85
7.10	Land suitability for Chilli	86
7.11	Land suitability for Tomato	87
7.12	Land suitability for Brinjal	88
7.13	Land suitability for Onion	89
7.14	Land suitability for Bhendi	90
7.15	Land suitability for Drumstick	91
7.16	Land suitability for Mango	92
7.17	Land suitability for Guava	93
7.18	Land suitability for Sapota	94
7.19	Land suitability for Pomegranate	95
7.20	Land suitability for Musambi	96
7.21	Land suitability for Lime	97
7.22	Land suitability for Amla	98
7.23	Land suitability for Cashew	99
7.24	Land suitability for Jackfruit	100
7.25	Land suitability for Jamun	101
7.26	Land suitability for Custard apple	102
7.27	Land suitability for Tamarind	103

7.28	Land suitability for Mulberry	104
7.29	Land suitability for Marigold	105
7.30	Land suitability for Chrysanthemum	106
7.31	Proposed Crop Plan for Himalapur Microwatershed	109

## LIST OF FIGURES

2.1	Location map of Himalapur Microwatershed	3
2.2	Granite and granite gneiss rock formation	4
2.3	Rainfall distribution in Yadgir Taluk & District	5
2.4	Natural vegetation of Himalapur Microwatershed	6
2.5	Current Land use map of Himalapur Microwatershed	7
2.6	Location of wells map of Himalapur Microwatershed.	8
2.7 a	Different crops and cropping systems in Himalapur Microwatershed	8
2.7 b	Different crops and cropping systems in Himalapur Microwatershed	9
3.1	Scanned and Digitized Cadastral map of Himalapur Microwatershed	11
3.2	Satellite image of Himalapur Microwatershed	12
3.3	Cadastral map overlaid on IRS PAN+LISS IV merged imagery of Himalapur Microwatershed	12
3.4	Location of profiles in a transect	13
3.5	Soil phase or management units of Himalapur Microwatershed	17
5.1	Land Capability Classification map of Himalapur Microwatershed	31
5.2	Soil Depth map of Himalapur Microwatershed	32
5.3	Surface Soil Texture map of Himalapur Microwatershed	33
5.4	Soil Gravelliness map of Himalapur Microwatershed	34
5.5	Soil Available Water Capacity map Himalapur Microwatershed	35
5.6	Soil Slope map of Himalapur Microwatershed	36
5.7	Soil Erosion map of Himalapur Microwatershed	37
6.1	Soil Reaction (pH) map of Himalapur Microwatershed	40
6.2	Electrical Conductivity (EC) map of Himalapur Microwatershed	40
6.3	Soil Organic Carbon (OC) map of Himalapur Microwatershed	41
6.4	Soil Available Phosphorus map of Himalapur Microwatershed	41
6.5	Soil Available Potassium map of Himalapur Microwatershed	42
6.6	Soil Available Sulphur map of Himalapur Microwatershed	43
6.7	Soil Available Boron map of Himalapur Microwatershed	43
6.8	Soil Available Iron map of Himalapur Microwatershed	44
6.9	Soil Available Manganese map of Himalapur Microwatershed	44
6.10	Soil Available Copper map of Himalapur Microwatershed	45
6.11	Soil Available Zinc map of Himalapur Microwatershed	45
7.1	Land suitability for Sorghum	48

7.2	Land suitability for Maize	49
7.3	Land suitability for Bajra	50
7.4	Land suitability for Groundnut	51
7.5	Land suitability for Sunflower	52
7.6	Land suitability for Redgram	53
7.7	Land suitability for Bengal gram	54
7.8	Land suitability for Cotton	55
7.9	Land suitability for Chilli	56
7.10	Land suitability for Tomato	57
7.11	Land suitable for Brinjal	58
7.12	Land suitable for Onion	59
7.13	Land suitable for Bhendi	60
7.14	Land suitable for Drumstick	61
7.15	Land suitability for Mango	62
7.16	Land suitability for Guava	63
7.17	Land suitability for Sapota	64
7.18	Land suitability for Pomegranate	65
7.19	Land suitability for Musambi	66
7.20	Land suitability for Lime	67
7.21	Land suitability for Amla	68
7.22	Land suitability for Cashew	69
7.23	Land suitability for Jackfruit	70
7.24	Land suitability for Jamun	71
7.25	Land suitability for Custard apple	72
7.26	Land suitability for Tamarind	73
7.27	Land suitability for Mulberry	74
7.28	Land suitability for Marigold	75
7.29	Land suitability for Chrysanthemum	76
7.30	Land management units map of Himalapur Microwatershed	107
9.1	Soil and water conservation plan map of Himalapur Microwatershed	120



## **EXECUTIVE SUMMARY**

*The land resource inventory of Himalapur Microwatershed was conducted using village cadastral maps and IRS satellite imagery on 1:7920 scale. The false colour composites of IRS imagery were interpreted for physiography and the physiographic delineations were used as base for mapping soils. The soils were studied in several transects and a soil map was prepared with phases of soil series as mapping units. Random checks were made all over the area outside the transects to confirm and validate the soil map unit boundaries. The soil map shows the geographic distribution and extent, characteristics, classification, behavior and use potentials of the soils in the microwatershed.*

*The present study covers an area of 498 ha in Yadgir taluk & district, Karnataka. The climate is semiarid and categorized as drought-prone with an average annual rainfall of 866 mm, of which about 652 mm is received during south-west monsoon, 138 mm during north-east and the remaining 76 mm during the rest of the year. An area of 330 ha in the microwatershed is covered by soils, about 152 ha covered by forest and 17 ha by others (Habitation and water bodies). The salient findings from the land resource inventory are summarized briefly below.*

- ❖ The soils belong to 5 soil series and 7 soil phases (management units) and 3 land management units.*
- ❖ The length of crop growing period is about 120-150 days starting from 1<sup>st</sup> week of June to 4<sup>th</sup> week of October.*
- ❖ From the master soil map, several interpretative and thematic maps like land capability, soil depth, surface soil texture, soil gravelliness, available water capacity, soil slope and soil erosion were generated.*
- ❖ Soil fertility status maps for macro and micronutrients were generated based on the surface soil samples collected at every 320 m grid interval.*
- ❖ Land suitability for growing 29 major agricultural and horticultural crops was assessed and maps showing the degree of suitability along with constraints were generated.*
- ❖ An area about 330 ha (66%) in the microwatershed is suitable for agriculture.*
- ❖ About 27 per cent area of the microwatershed has soils that are deep to very deep (100 - >150 cm) whereas 5 per cent soils are moderately shallow (50 -75 cm), 34 per cent soils are very shallow to shallow (<25 -50 cm).*
- ❖ About 8 per cent area in the microwatershed has sandy soils and 57 per cent clayey soils at the surface.*
- ❖ Entire cultivated area in the microwatershed has non gravelly (<15%) soils.*
- ❖ About 27 per cent area of the microwatershed is very high (>200 mm/m) in available water capacity, 5 per cent soils are low (51-100 mm/m) and 34 per cent area very low (<50 mm/m) available water capacity.*

- ❖ *About 58 per cent area of the microwatershed has very gently sloping (1-3% slope) land and 8 per cent is gently sloping (3-5%) soils.*
- ❖ *An area of about 33 per cent area is moderately (e2) eroded and 33 percent soils are severely eroded (e3).*
- ❖ *An area of about 46 per cent soils are neutral (pH 6.5-7.3) in soil reaction, about 20 per cent soil are slightly alkaline (pH 7.3-7.8) soils.*
- ❖ *The Electrical Conductivity (EC) of the soils in the entire cultivated area of the microwatershed is dominated by  $<2 \text{ ds}^{\text{m}^{-1}}$  indicating that the soils are non-saline.*
- ❖ *Entire cultivated area of the microwatershed is high ( $>0.75$ ) in organic carbon content.*
- ❖ *About 19 percent is medium (23-57 kg/ha) in available phosphorus and 48 percent soils are low ( $<23 \text{ kg/ha}$ ) in available phosphorus.*
- ❖ *About 6 percent of the soils are low ( $<145 \text{ kg/ha}$ ) in available potassium, 36 percent soils are medium (145-337kg/ha) and 23 percent area high ( $>337 \text{ kg/ha}$ ) in available potassium.*
- ❖ *An area of about 32 percent is medium (10-20ppm) and 34 per cent area is low ( $<10 \text{ ppm}$ ) in available sulphur*
- ❖ *Available boron is low ( $<0.5 \text{ ppm}$ ) in an area of about 65 per cent and medium (0.5-1.0 ppm) of about  $<1$  per cent. Soils.*
- ❖ *Available iron content is sufficient ( $>4.5 \text{ ppm}$ ) in the entire cultivated area of the microwatershed.*
- ❖ *Available manganese and copper are sufficient in all the soils of the microwatershed.*
- ❖ *Available zinc is deficient ( $<0.6 \text{ ppm}$ ) which covers an area of 49 percent and sufficient ( $>0.6 \text{ ppm}$ ) covers an area of 17 per cent in the microwatershed.*
- ❖ *The land suitability for 29 major crops grown in the microwatershed were assessed and the areas that are highly suitable (S1) and moderately suitable (S2) are given below. It is however to be noted that a given soil may be suitable for various crops but what specific crop to be grown may be decided by the farmer looking to his capacity to invest on various inputs, marketing infrastructure, market price and finally the demand and supply position.*

**Land suitability for various crops in the Microwatershed**

<b>Crop</b>	<b>Suitability Area in ha (%)</b>		<b>Crop</b>	<b>Suitability Area in ha (%)</b>	
	<b>Highly suitable (S1)</b>	<b>Moderately suitable (S2)</b>		<b>Highly suitable (S1)</b>	<b>Moderately suitable (S2)</b>
<i>Sorghum</i>	60(12)	99(20)	<i>Sapota</i>	-	-)
<i>Maize</i>	-	159(32)	<i>Pomegranate</i>	-	134(27)
<i>Bajra</i>	-	159(32)	<i>Musambi</i>	134(27)	-
<i>Groundnut</i>	-	-	<i>Lime</i>	134(27)	-
<i>Sunflower</i>	60(12)	74(15)	<i>Amla</i>	-	159(32)
<i>Redgram</i>	-	134(27)	<i>Cashew</i>	-	-
<i>Bengal gram</i>	134(27)	25(5)	<i>Jackfruit</i>	-	-
<i>Cotton</i>	134(27)	25(5)	<i>Jamun</i>	-	134(27)
<i>Chilli</i>	-	85(17)	<i>Custard apple</i>	134(27)	25(5)
<i>Tomato</i>	-	85(17)	<i>Tamarind</i>	-	134(27)
<i>Drumstick</i>	-	134(27)	<i>Mulberry</i>	-	-
<i>Mango</i>	-	-	<i>Marigold</i>	-	159(32)
<i>Guava</i>	-	-	<i>Chrysanthemum</i>	-	159(32)
<i>Brinjal</i>	-	159(32)	<i>Bhendi</i>	-	159(32)
<i>Onion</i>	-	25(5)			

- ❖ *Apart from the individual crop suitability, a proposed crop plan has been prepared for the identified LMUs by considering only the highly and moderately suitable lands for different crops and cropping systems with food, fodder, fiber and horticulture crops.*
- ❖ *Maintaining soil-health is vital to crop production and conserve soil and land resource base for maintaining ecological balance and to mitigate climate change. For this, several ameliorative measures have been suggested to these problematic soils like saline/alkali, highly eroded, sandy soils etc.*
- ❖ *Soil and water conservation treatment plan has been prepared that would help in identifying the sites to be treated and also the type of structures required.*
- ❖ *As part of the greening programme, several tree species have been suggested to be planted in marginal and submarginal lands, field bunds and also in the hillocks, mounds and ridges. This would help in not only supplementing the farm income but also provide fodder and fuel to generate lot of biomass which would help in maintaining an ecological balance and also contribute to mitigating the climate change.*



## **INTRODUCTION**

Land is a scarce resource and basic unit for any material production. It can support the needs of the growing population, provided they use the land in a rational and judicious manner. But what is happening in many areas of the state is a cause for concern to everyone involved in the management of land resources at the grassroots level. The area available for agriculture is about 51 per cent of the total geographical area and more than 60 per cent of the people are still dependant on agriculture for their livelihood. The limited land area is under severe stress and strain due to increasing population pressure and competing demands of various land uses. Due to this, every year there is significant diversion of farm lands and water resources for non-agricultural purposes. Apart from this, due to lack of interest in farmers for farming, large tracts of cultivable lands are turning into fallows in many areas and this trend is continuing at an alarming rate.

Further, land degradation has emerged as a serious problem which has already affected about 38 lakh ha of cultivated area in the state. Soil erosion alone has degraded about 35 lakh ha. Almost all the uncultivated areas are facing various degrees of degradation, particularly soil erosion. Salinity and alkalinity has emerged as a major problem in more than 3.5 lakh ha in the irrigated areas of the state. Nutrient depletion and declining factor productivity is common in both rainfed and irrigated areas. The degradation is continuing at an alarming rate and there appears to be no systematic effort among the stakeholders to contain this process. In recent times, an aberration of weather due to climate change phenomenon has added another dimension leading to unpredictable situations to be tackled by the farmers.

In this critical juncture, the challenge before us is not only to increase the productivity per unit area which is steadily declining and showing a fatigue syndrome, but also to prevent or at least reduce the severity of degradation. If the situation is not reversed at the earliest, then the sustainability of the already fragile crop production system and the overall ecosystem will be badly affected in the state. The continued neglect and unscientific use of the resources for a long time has led to the situation observed at present in the state. It is a known fact and established beyond doubt by many studies in the past that the cause for all kinds of degradation is the neglect and irrational use of the land resources. Hence, there is an urgent need to generate a detailed site-specific farm level database on various land resources for all the villages/watersheds in a time bound manner that would help to protect the valuable soil and land resources and also to stabilize the farm production.

Therefore, the land resource inventory required for farm level planning is the one which investigates not only the surface but also consider the other parameters which are critical for productivity *viz.*, soils, climate, water, minerals and rocks, topography, geology, hydrology, vegetation, crops, land use pattern, animal population, socio-

economic conditions, infrastructure, marketing facilities and various schemes and developmental works of the government etc. From the data collected at farm level, the specific problems and potentials of the area can be identified and highlighted, conservation measures required for the area can be planned on a scientific footing, suitability of the area for various uses can be worked out and finally viable and sustainable land use options suitable for each and every land holding can be prescribed.

The Land Resource Inventory is basically done for identifying the potential and problem areas, developing sustainable land use plans, estimation of surface run off and water harvesting potential, preparation of soil and water conservation plans, land degradation/desertification etc. The Bureau is presently engaged in developing an LRI methodology using high resolution satellite remote sensing data and Digital Elevation Model (DEM) data to prepare Landscape Ecological Units (LEU) map representing agro-ecosystem as a whole. The LEU is preferred over landform as the base map for LRI. LEU is the assemblage of landform, slope and land use. An attempt has already been made to upscale the soil resource information from 1:250000 and 1:50000 scale to the LEU map in Goa and other states.

The land resource inventory aims to provide site specific database for Himalapur microwatershed in Yadgir Taluk & District, Karnataka State for the Karnataka Watershed Development Department. The database was generated by using cadastral map of the village as a base along with high resolution IRS LISS IV and Cartosat-1 merged satellite imagery. Later, an attempt will be made to uplink this LRI data generated at 1:7920 scale under Sujala-III Project to the proposed Landscape Ecological Units (LEUs) map.

The study was organized and executed by the ICAR- National Bureau of Soil Survey and Land Use Planning, Regional Centre, Bangalore under Generation of Land Resource Inventory Data Base Component-1 of the Sujala-III Project funded by the World Bank.

## GEOGRAPHICAL SETTING

### 2.1 Location and Extent

The Himalapur microwatershed is located in the northern part of Karnataka in Yadgir Taluk & District, Karnataka State (Fig.2.1). It comprises parts of Chapetla, Boodhura and Himalapur village. It lies between 16<sup>0</sup> 50' and 16<sup>0</sup> 52' North latitudes and 77<sup>0</sup> 19' and 77<sup>0</sup> 20' East longitudes, covering an area of about 498 ha, It is about 30 km north of Yadgir town and is surrounded by Chapetla on the north, Boodhura on east, Himalapur on southern side of the microwatershed.




Fig.2.1 Location map of Himalapur Microwatershed

### 2.2 Geology

Major rock formations observed in the microwatershed are granite gneiss (Figs.2.2). Granite gneisses are essentially pink to gray and are coarse to medium grained. They

consist primarily of quartz, feldspar, biotite and hornblende. The gray granite gneisses are highly weathered, fractured and fissured upto a depth of about 10 m. Dolerite dykes and quartz veins are common with variable width and found to occur in Himalapur microwatershed.



Fig.2.2 Granite and granite gneiss rocks

### **2.3 Physiography**

Physiographically, the area has been identified as granite gneiss based on geology. The area has been further subdivided into five landforms, viz; mounds/ridges, summits, side slopes and very gently sloping uplands, plains and valleys based on slope and its relief features. The elevation ranges from 541-629 m above MSL. The mounds and ridges are mostly covered by rock outcrops.

### **2.4 Drainage**

The area is drained by several parallel streams like Bori, Amerja and Kanga which finally join the river Bhima along its course. Though, they are not perennial, during rainy season they carry large quantities of rain water. The microwatershed has only few small tanks which are not capable of storing the water that flows during the rainy season. Due to this, the ground water recharge is very much affected. This is reflected in the failure of many bore wells in the villages. If the available rain water is properly harnessed by constructing new tanks and recharge structures at appropriate places in the villages, then the drinking and irrigation needs of the area can be easily met. The drainage network is parallel to sub parallel and dendritic.

### **2.5 Climate**


The Yadgir district lies in the northern plains of Karnataka and falls under semiarid tract of the state and is categorized as drought- prone with total annual rainfall of 866 mm (Table 2.1). Of the total rainfall, maximum of 652 mm is received during the



south–west monsoon period from June to September, the north-east monsoon from October to early December contributes about 138 mm and the remaining 76 mm during the rest of the year. The summer season starts during the middle of February and continues up to the first week of June. The period from December to the middle of February is the coldest season. December is the coldest month with mean daily maximum and minimum temperatures being 29.5<sup>0</sup>C and 10<sup>0</sup>C respectively. During peak summer, temperature shoots up to 45<sup>0</sup>C. Relative humidity varies from 26% in summer to 62% in winter. Rainfall distribution is shown in Figure 2.3. The average Potential Evapo-Transpiration (PET) is 141 mm and varies from a low of 81 mm in December to 199 mm in the month of May. The PET is always higher than precipitation in all the months except end of June to end of September. Generally, the Length of crop Growing Period (LGP) is 120-150 days and starts from 1<sup>st</sup> week of June to 4<sup>th</sup> week of October.

**Table 2.1 Mean Monthly Rainfall, PET, 1/2 PET at Yadgir Taluk, Yadgir District**

Sl. No.	Months	Rainfall	PET	1/2 PET
1	January	4.30	86.0	43.0
2	February	2.30	125.5	62.7
3	March	15.10	166.0	83.0
4	April	18.50	179.8	89.9
5	May	36.0	198.8	97.9
6	June	118.0	175.1	87.5
7	July	171.80	156.3	78.1
8	August	182.9	150.3	75.1
9	September	179.7	142.0	71.0
10	October	105.3	138.5	69.2
11	November	26.4	97.60	48.6
12	December	6.0	80.90	40.4
<b>Total</b>		<b>866.3</b>		



**Fig 2.3 Rainfall distribution in Yadgir Taluk, Yadgir District**

## 2.6 Natural Vegetation

The natural vegetation is sparse comprising few tree species, shrubs and herbs. The mounds, ridges and boulders occupy very sizeable area which is under thin to moderately thick forest vegetation. Still, there are some remnants of the past forest cover which can be seen in patches in some ridges and hillocks in the microwatershed (Fig 2.4).

Apart from the continuing deforestation, the presence of large population of goats, sheep and other cattle in the microwatershed is causing vegetative degradation of whatever little vegetation left in the area. The uncontrolled grazing has left no time for the regeneration of the vegetative cover. This leads to the accelerated rate of erosion on the hill slopes resulting in the formation of deep gullies in the foot slopes that eventually result in the heavy siltation of tanks and reservoirs in the microwatershed.




Fig 2.4 Natural vegetation of Himalapur Microwatershed

## 2.7 Land Utilization

About 72 per cent area (Table 2.2) in Yadgir district is cultivated at present. An area of about 2 per cent is permanently under pasture, 20 per cent under current fallows and 6 per cent under non-agricultural land and 5 per cent under currently barren. Forests occupy an area of about 7 per cent and the tree cover is in a very poor state. Most of the mounds, ridges and bouldery areas have very poor vegetative cover. Major crops grown in the area are sorghum, maize, cotton, sunflower, groundnut, red gram, mango, pomegranate, marigold and sapota. While carrying out land resource inventory, the land use/land cover particulars are collected from all the survey numbers and a current land use map of the microwatershed is prepared. The current land use map prepared shows the arable and non-arable lands, other land uses and different types of crops grown in the area. The current land use map of Himalapur microwatershed is presented in Fig.2.5. The location of wells in the Himalapur microwatershed is shown in fig. 2.6. The different crops and cropping systems adopted in the microwatershed are presented in Figures 2.7 a & b.

**Table 2.2 Land Utilization in Yadgir District**

Sl. No.	Agricultural land use	Area ( ha)	Per cent
1	Total geographical area	516088	-
2	Total cultivated area	373617	72.4
3	Area sown more than once	74081	14.3
4	Cropping intensity	-	119.8
5	Trees and grooves	737	0.14
6	Forest	33773	6.54
7	Cultivable wasteland	2385	0.46
8	Permanent Pasture land	11755	2.28
9	Barren land	27954	5.41
10	Non- Agriculture land	29623	5.73
11	Current Fallows	105212	20.4



**Fig.2.5 Current Land Use map of Himalapur Microwatershed**




Fig.2.6 Location of wells map of Himalapur Microwatershed.




Fig. 2.7 a. Different Crops and Cropping Systems in Himalapur Microwatershed



Sunflower




Groundnut



Mango



Pomegranate



Cotton



Sorghum

Fig. 2.7 b. Different Crops and Cropping Systems in Himalapur Microwatershed

## SURVEY METHODOLOGY

The purpose of land resource inventory is to delineate similar areas (soil series and phases), which respond or expected to respond similarly to a given level of management. This was achieved in Himalapur microwatershed by the detailed study of all the soil characteristics (depth, texture, colour, structure, consistence, coarse fragments, porosity, soil reaction, soil horizons etc.) and site characteristics (slope of the land, erosion, drainage, occurrence of rock fragments etc.) followed by grouping of similar areas based on soil-site characteristics into homogeneous (management units) units, and showing the area extent and their geographic distribution on the microwatershed cadastral map. The detailed survey at 1:7920 scale was carried out in an area of 498 ha. The methodology followed for carrying out land resource inventory was as per the guidelines given in Soil Survey Manual (IARI, 1971; Soil Survey Staff, 2006; Natarajan *et al.*, 2015) which is briefly described below.

### 3.1 Base Maps

The detailed survey of the land resources occurring in the microwatershed was carried out by using digitized cadastral and IRS satellite imagery map as base supplied by KRSRAC. The cadastral map shows field boundaries with their survey numbers, location of tanks, streams and other permanent features of the area (Fig. 3.1). Apart from the cadastral map, remote sensing data products from Cartosat-1 and LISS IV merged at the scale of 1:7920 were used in conjunction with the cadastral map to identify the landscapes, landforms and other surface features. The imagery helped in the identification and delineation of boundaries between hills, uplands and lowlands, water bodies, forest and vegetated areas, roads, habitations and other cultural features of the area (Fig. 3.2). The cadastral map was overlaid on the satellite imagery (Fig. 3.3) that helps to identify the parcel boundaries and other permanent features. Apart from cadastral maps and images, toposheets of the area (1:50,000 scale) were also used for initial traversing, identification of geology and landforms, drainage features, present land use and also for selection of transects in the microwatershed.

### 3.2 Image Interpretation for Physiography

False Colour Composites (FCCs) of Cartosat-I and LISS-IV merged satellite data covering microwatershed area was visually interpreted using image interpretation elements and all the available collateral data with local knowledge. The delineated physiographic boundaries were transferred on to a cadastral map overlaid on satellite imagery. Physiographically, the area has been identified as granite and granite gneiss landscape. It was divided into five landforms, *viz*; ridges and mounds, gently and very gently sloping uplands and lowlands based on slope and image characteristics. They were further subdivided into physiographic/image interpretation units based on image characteristics. The image interpretation legend for physiography is given below.

## Image Interpretation Legend for Physiography

### G- Granite Gneiss Landscape

G1	Hills/ Ridges/ Mounds
G11	Summits
G12	Side slopes
G121	Side slopes with dark grey tones
G2	Uplands
G21	Summits
G22	Gently sloping uplands
G221	Gently sloping uplands, yellowish green (eroded)
G222	Gently sloping uplands, yellowish white (severely eroded)
G23	Very gently sloping uplands
G231	Very gently sloping uplands, yellowish green
G232	Very gently sloping uplands, medium green and pink
G233	Very gently sloping uplands, pink and green (scrub land)
G234	Very gently sloping uplands, medium greenish grey
G235	Very gently sloping uplands, yellowish white (eroded)
G236	Very gently sloping uplands, dark green
G237	Very gently sloping uplands, medium pink (coconut garden)
G238	Very gently sloping uplands, pink and bluish white (eroded)
G24	Valleys/ lowlands
G241	Valleys, pink tones
G242	Valleys gray mixed with pink tones




Fig 3.1 Scanned and Digitized Cadastral map of Himalapur Microwatershed




Fig.3.2 Satellite Image of Himalapur Microwatershed




Fig.3.3 Cadastral map overlaid on IRS PAN+LISS IV merged imagery of Himalapur Microwatershed



### 3.3 Field Investigation

The field boundaries and survey numbers given on the cadastral sheet were located on the ground by following permanent features like roads, cart tracks, *nallas*, streams, tanks etc., and wherever changes were noticed, they were incorporated on the microwatershed cadastral map. Preliminary traverse of the microwatershed was carried out with the help of cadastral map, imagery and toposheets. While traversing, landforms and physiographic units identified were checked and preliminary soil legend was prepared by studying soils at few selected places. Then, intensive traversing of each physiographic unit like hills, ridges, uplands and valleys was carried out. Based on the variability observed on the surface, transects (Fig. 3.4) were selected across the slope covering all the landform units in the microwatershed (Natarajan and Dipak Sarkar, 2010).




Fig: 3.4. Location of profiles in a transect

In the selected transect, soil profiles were located (Fig. 3.4) at closely spaced intervals to take care of any change in the land features like break in slope, erosion, gravel, stones etc. In the selected sites, profiles (vertical cut showing the soil layers from surface to the rock) were opened upto 200 cm or to the depth limited by rock or hard substratum and studied in detail for all their morphological and physical characteristics. The soil and site characteristics were recorded for all profile sites on a standard proforma as per the guidelines given in USDA Soil Survey Manual (Soil Survey Staff, 2012). Apart from the transect study, profiles were also studied at random, almost like in a grid pattern, outside the transect areas.

Based on the soil characteristics, the soils were grouped into different soil series. Soil series is the most homogeneous unit having similar horizons and properties and behaves similarly for a given level of management. Soil depth, texture, colour, kind of horizon and horizon sequence, calcareousness, amount and nature of gravel present, nature of substratum etc, were used as the major differentiating characteristics for

identifying soil series occurring in the area. The differentiating characteristics used for identifying the soil series are given in Table 3.1. Based on the above characteristics, 5 soil series were identified in the Himalapur microwatershed.

**Table 3.1 Differentiating Characteristics used for identifying Soil Series**  
(Characteristics are of Series Control Section)

<b>Soils of Granite gneiss Landscape</b>							
<b>Sl. no</b>	<b>Soil Series</b>	<b>Depth (cm)</b>	<b>Colour (moist)</b>	<b>Texture</b>	<b>Gravel (%)</b>	<b>Horizon sequence</b>	<b>Calcareousness</b>
<b>Soil of Granite and Granite Gneiss Landscape</b>							
1	BDP (Baddeppalli)	<25	7.5YR 3/2,3/4 5YR 3/4	scl	<15	Ap-AC	es
2	BDL (Badiyala)	25-50	7.5YR2.5/3,2.5/2,3/3 10YR 3/4,4/3	sl	<15	Ap-Bw	e
3	JNK (Jinkera)	50-75	10YR 3/1,3/2 7.5YR 3/4	scl	<15	Ap-Bw	e
4	NGP (Naglapur)	100-150	10YR 3/2,3/1,2/1	c	<15	Ap-Bss	es
5	BMN (Bhimanahalli)	>150	10YR 3/1	c	<15	Ap-Bss	es

### 3.4 Soil Mapping

The area under each soil series was further separated into soil phases and their boundaries delineated on the cadastral map based on the variations observed in the texture of the surface soil, slope, erosion, presence of gravel, stoniness etc. A soil phase is a subdivision of soil series based mostly on surface features that affect its use and management. The soil mapping units are shown on the map (Fig.3.5) in the form of symbols. During the survey about many profile pits, few minipits and a few auger bores representing different landforms occurring in the microwatershed were studied. In addition to the profile study, spot observations in the form of minipits, road cuts, terrace cuts etc., were studied to validate the soil boundaries on the soil map.

The soil map shows the geographic distribution of 7 mapping units representing 5 soil series occurring in the microwatershed. The soil map unit (soil legend) description is presented in Table 3.2. The soil phase map (management units) shows the distribution of 7 soil phases mapped in the microwatershed. Each mapping unit (soil phase) delineated on the map has similar soil and site characteristics. In other words, all the farms or survey numbers included in one phase will have similar management needs and have to be treated accordingly.

### 3.5 Land Management Units

The 7 soil phases identified and mapped in the microwatershed were grouped into 3 Land Management Units (LMU's) for the purpose of preparing a Proposed Crop Plan for sustained development of the microwatershed. The database (soil phases) generated under LRI was utilized for identifying Land Management Units (LMU's) based on the

management needs. One or more than one soil site characteristic having influence on the management have been chosen for identification and delineation of LMUs. For Himalapur microwatershed, five soil and site characteristics, namely soil depth, soil texture, slope, erosion and gravel content have been considered for defining LMUs. The land use classes are expected to behave similarly for a given level of management.

### 3.6 Laboratory Characterization

Soil samples were collected from representative master profiles for laboratory characterization by following the methods outlined in the Laboratory Manual (Sarma *et al*, 1987). Surface soil samples collected from farmer's fields (50 samples) for fertility status (major and micronutrients) at 320 m grid interval in the year 2017 were analyzed in the laboratory (Katyal and Rattan, 2003). By linking the soil fertility data to the survey numbers through GIS, soil fertility maps were generated by using Kriging method for the microwatershed.

**Table 3.2 Soil map unit description of Himalapur Microwatershed**

<b>**Soil map unit No.</b>	<b>Soil Series</b>	<b>Soil Phase</b>	<b>Mapping Unit Description</b>	<b>Area in ha (%)</b>
<b>Soils of Granite and Granite Gneiss Landscape</b>				
	BDP		Baddeppalli soils are very shallow (<25 cm), well drained, have dark brown to dark reddish brown, calcareous sandy clay loam soils occurring on very gently sloping uplands under cultivation	<b>94 (18.78)</b>
119		BDPiB3	Sandy clay surface, slope 1-3%, severe erosion	94 (18.78)
	BDL		Badiyala soils are shallow (25-50 cm), well drained, have dark brown to very dark brown and dark yellowish brown, slightly calcareous sandy loam soils occurring on very gently to gently sloping uplands under cultivation	<b>77 (15.37)</b>
3		BDLbC3	Loamy sand surface, slope 3-5%, severe erosion	42 (8.37)
5		BDLiB2	Sandy clay surface, slope 1-3%, moderate erosion	7 (1.42)
6		BDLiB3	Sandy clay surface, slope 1-3%, severe erosion	28 (5.58)
	JNK		Jinkera soils are moderately shallow (50-75 cm), well drained, have dark brown to very dark grayish brown, slightly calcareous sandy clay loam soils occurring on very gently sloping uplands under cultivation	<b>25 (5.03)</b>
152		JNKmB2	Clay surface, slope 1-3%, moderate erosion	25 (5.03)
	NGP		Nagalapur soils are deep (100-150 cm), moderately well drained, have very dark gray to very dark grayish brown, black calcareous cracking clay soils occurring on very gently sloping uplands under cultivation	<b>74 (14.92)</b>

49		NGPmB2	Clay surface, slope 1-3%, moderate erosion	74 (14.92)
	BMN	Bhimanahalli soils are very deep (>150 cm), moderately well drained, have very dark gray, calcareous cracking clay black soils occurring on very gently sloping uplands under cultivation		<b>60 (12.0)</b>
62		BMNmB2	Clay surface, slope 1-3%, moderate erosion	60 (12.0)
900		Forest		<b>152 (30.47)</b>
1000	Others	Habitation and Water body		<b>17 (3.43)</b>




Fig 3.5 Soil Phase or Management Units - Himalapur Microwatershed



## THE SOILS

Detailed information pertaining to the nature, extent and their distribution of different kinds of soils occurring in Himalapur microwatershed is provided in this chapter. The microwatershed area has been identified as granite gneiss landscape based on geology. In all, 5 soil series are identified. Soil formation is the result of the combined effect of environmental and terrain factors that are reflected in soil morphology. In the granite gneiss landscape, it is by parent material, relief and climate.

A brief description of each of the 5 soil series identified followed by 7 soil phases (management units) mapped under each series are furnished below. The physical and chemical characteristics of soil series identified in Himalapur microwatershed are given in Table 4.1 along with soil classification. The soils in any one map unit differ from place to place in their depth, texture, slope, gravelliness, erosion or any other site characteristic that affect management. The soil phase map can be used for identifying the suitability of areas for growing specific crops or for other alternative uses and also for deciding the type of conservation structures needed. The detailed information on soil and site-characteristics like soil depth, surface soil texture, slope, erosion, gravelliness, AWC, LCC etc, with respect to each of the soil phase identified is given village/survey number wise for the microwatershed in Appendix-I.

### 4.1 Soils of granite gneiss landscape

In this landscape, 5 soil series are identified and mapped. Of these, BDP series occupies maximum area of 94 ha (19%) followed by BDL 77 ha (15%), NGP 74 ha (15%), BMN 60 ha (12%), JNK 25 ha (5%), Brief description of each series identified and number of soil phases mapped is given below.

**4.1.1 Baddeppalli (BDP) Series:** Baddeppalli soils are very shallow (<25cm), well drained, have dark brown to dark reddish brown, calcareous sandy clay loam soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation. The Baddeppalli series has been classified as a member of the loamy, mixed (calcareous), isohyperthermic family of Lithic Ustorthents.

The thickness of the soil is less than 25 cm. Its colour is in 7.5 YR and 5 YR hue with value 3 and chroma 2 to 4. The texture varies from sandy clay loam to sandy clay and is calcareous. The available water capacity is very low (<50 mm/m). One phase was identified and mapped.



Landscape and Soil Profile characteristics of Baddeppalli (BDP) Series

**4.1.2 Badiyala (BDL) Series:** Badiyala soils are shallow (25-50 cm), well drained, have very dark brown, dark yellow brown and dark brown, slightly calcareous sandy loam soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation. The Badiyala series has been classified as a member of the coarse-loamy, mixed, isohyperthermic family of Fluventic Haplustepts

The thickness of the solum ranges from 28 to 50 cm. The thickness of A horizon ranges from 4 to 12 cm. Its colour is in 10YR hue with value 3 to 4 and chroma 3 to 4. The texture is loamy sand, sandy clay loam and sandy clay. The thickness of B horizon ranges from 27 to 45 cm. Its colour is in 10 YR and 7.5 YR hue with value 2 to 4 and chroma 3 to 4. Its texture is sandy loam to sandy clay loam and is slightly calcareous. The available water capacity is very low (<50mm/m). Three phases were indentified and mapped.



Landscape and Soil Profile characteristics of Badiyala (BDL) Series



**4.1.3 Jinkera (JNK) Series:** Jinkera soils are moderately shallow (50-75 cm), well drained, have very dark gray to very dark grayish brown and dark brown, slightly calcareous sandy clay loam soils. They are developed from weathered granite gneiss and occur on very gently sloping uplands under cultivation. The Jinkera series has been classified as a member of the fine-loamy, mixed, isohyperthermic family of Typic Haplustepts.

The thickness of the solum ranges from 51-75 cm. Thickness of A horizon ranges from 6 to 11 cm. Its colour is in hue 10 YR and 7.5 YR with value and chroma of 3 to 4. The texture varies from sandy loam to sandy clay. The thickness of B horizon ranges from 53 to 66 cm. Its colour is in 10 YR and 7.5 YR hue with value and chroma of 2 to 4. The texture varies from sandy clay loam to sandy clay and is slightly calcareous. The available water capacity is low (51-100 mm/m). One phase was indentified and mapped.



Landscape and Soil Profile characteristics of Jinkera (JNK) Series

**4.1.4 Naglapur (NGP) Series:** Naglapur soils are deep (100-150 cm), moderately well drained, have black to very dark grayish brown, calcareous cracking clay soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation. The Naglapur series has been classified as a member of the very fine, smectitic (calcareous), isohyperthermic family of Typic Haplusterts.

The thickness of the solum ranges from 110 to 150 cm. The thickness of A horizon ranges from 6 to 25 cm. Its colour is in 10 YR hue with value 3 to 5 and chroma 1 to 3. The texture varies from sandy loam to sandy clay and clay. The thickness of B horizon ranges from 110 to 141 cm. Its colour is in 10 YR hue with value 2 to 3 and chroma 1 to 2. Texture is clay and is calcareous. The available water capacity is very high (>200 mm/m). One phase was indentified and mapped.



Landscape and Soil Profile characteristics of Naglapur (NGP) Series

**4.1.5 Bhimanahalli (BMN) Series:** Bhimanahalli soils are very deep (>150 cm), moderately well drained, very dark gray calcareous cracking clay soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation. The Bhimanahalli series has been classified as a member of the fine, smectitic (calcareous), isohyperthermic family of Typic Haplusterts.

The thickness of the solum is more than 150 cm. The thickness of A horizon ranges from 6 to 13 cm. Its colour is in 10 YR hue with value 3 to 4 and chroma 1 to 2 with clay texture. The thickness of B horizon ranges from 163 to 176 cm. Its colour is in 10 YR hue with value 2 to 4 and chroma 1. Its texture is clay and is calcareous. The available water capacity is very high (>200 mm/m). One phase was identified and mapped.



Landscape and Soil Profile characteristics of Bhimanahalli (BMN) Series

**Table: 4.1 Physical and Chemical Characteristics of Soil Series identified in Himalapur microwatershed**

**Soil Series:** Baddeppalli (BDP) **Pedon:** R-11

**Location:** 16°43'84.4"N 77°14'06.4"E, Halagera village, Yadgir hobli, Yadgir taluk and district

**Analysis at:** NBSS&LUP, Regional Centre, Bengaluru **Classification:** Loamy, mixed (calcareous), isohyperthermic, Lithic Ustorthents

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-16	Ap	58.67	17.02	24.31	19.03	13.74	9.62	10.57	5.71	<15	scl	16.19	8.18

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO <sub>3</sub>	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP
	Water	CaCl <sub>2</sub>	M KCl				Ca	Mg	K	Na	Total				
				dS m <sup>-1</sup>	%	%	cmol kg <sup>-1</sup>					%	%		
0-16	8.58	-	-	0.262	1.60	7.67	-	-	0.24	0.06	-	18.10	0.74	100	0.35

*Contd...*

**Soil Series:** Badiyala (BDL) **Pedon:** R-5

**Location:** 16°37'10.0"N 77°20'21.5", Gudalagunta village, Balichakra hobli, Yadgir taluk and district

**Analysis at:** NBSS&LUP, Regional Centre, Bengaluru **Classification:** Coarse-loamy, mixed, isohyperthermic, Fluventic Haplustepts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-12	Ap	87.13	7.04	5.83	10.03	24.32	23.61	23.51	5.67	<15	ls	6.27	2.44
12-28	Bw	64.63	13.30	22.07	6.74	13.07	22.30	17.01	5.50	<15	scl	16.34	7.83
28-50	BC	73.11	12.02	14.87	3.93	16.03	26.89	18.41	7.86	<15	sl	12.94	5.47

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO <sub>3</sub>	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP
	Water	CaCl <sub>2</sub>	M KCl				Ca	Mg	K	Na	Total				
				dS m <sup>-1</sup>	%	%	cmol kg <sup>-1</sup>							%	%
0-12	6.20	-	-	0.074	1.00	0.00	2.80	0.98	0.14	0.01	3.92	4.20	0.72	93	0.20
12-28	9.04	-	-	0.253	0.80	3.20	-	-	0.16	0.69	-	16.90	0.77	100	4.09
28-50	9.41	-	-	0.364	1.10	3.60	-	-	0.16	1.39	-	11.10	0.75	100	12.52

*Contd...*

**Soil Series:** Jinkera (JNK) **Pedon:** R-1

**Location:** 16°45'13.5"N 77°10'59.8"E, Varkanahalli village, Yadgir hobli, Yadgir taluk and district

**Analysis at:** NBSS&LUP, Regional Centre, Bengaluru **Classification:** Fine-loamy, mixed, isohyperthermic Typic Haplustepts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-15	Ap	66.84	13.62	19.54	12.15	21.22	11.23	12.56	9.68	10	sl	14.42	7.70
15-38	Bw1	59.08	12.11	28.81	12.53	12.42	17.85	8.77	7.52	20	scl	18.21	12.23
38-52	Bw2	68.21	11.68	20.11	17.90	21.81	10.60	10.80	7.10	10	scl	14.54	8.96

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO <sub>3</sub>	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP
	Water	CaCl <sub>2</sub>	M KCl				Ca	Mg	K	Na	Total				
				dS m <sup>-1</sup>	%	%	cmol kg <sup>-1</sup>							%	%
0-15	8.42	-	-	0.148	0.70	0.65	-	-	0.15	0.03	-	14.50	0.74	100	0.18
15-38	8.38	-	-	0.226	0.31	2.21	-	-	0.09	0.23	-	21.70	0.75	100	1.05
38-52	8.40	-	-	0.195	0.25	1.17	-	-	0.07	0.19	-	15.90	0.79	100	1.23

*Contd...*

**Soil Series:** Naglapur (NGP) **Pedon:** R-8

**Location:** 16°52'84.1"N 77°22'99.4"E, Gurumitkal village, Gurumitkal hobli, Yadgir taluk and district

**Analysis at:** NBSS&LUP, Regional Centre, Bengaluru **Classification:** Very fine, smectitic (calcareous), isohyperthermic Typic Haplusterts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-10	Ap	7.53	19.88	72.59	1.00	0.78	0.89	2.10	2.77	-	c	44.31	32.79
10-35	Bss1	6.55	18.76	74.68	0.80	0.92	0.80	1.72	2.30	-	c	43.09	31.62
35-60	Bss2	6.58	21.05	72.37	0.69	0.46	1.04	1.50	2.89	-	c	46.52	32.52
60-102	Bss3	7.48	19.74	72.78	1.61	1.38	0.69	1.61	2.19	-	c	51.12	35.62

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO <sub>3</sub>	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP
	Water	CaCl <sub>2</sub>	M KCl				Ca	Mg	K	Na	Total				
				dS m <sup>-1</sup>	%	%	cmol kg <sup>-1</sup>					%	%		
0-10	7.42	-	-	0.24	0.84	1.30	-	-	0.84	0.15	-	67.10	0.92	100	0.22
10-35	8.52	-	-	0.291	0.64	2.86	-	-	0.17	0.29	-	65.20	0.87	100	0.45
35-60	7.89	-	-	0.134	0.62	4.55	-	-	0.15	0.20	-	65.00	0.90	100	0.30
60-102	8.68	-	-	0.213	0.54	8.32	-	-	0.17	0.15	-	64.10	0.88	100	0.24

Contd...

**Soil Series:** Bhimanahalli (BMN) **Pedon:** R-3

**Location:** 16°31'82.4"N 77°12'70.8"E, Bheemanahalli village, Sydhapura hobli, Yadgir taluk and district

**Analysis at:** NBSS&LUP, Regional Centre, Bengaluru **Classification:** Fine, smectitic (calcareous), isohyperthermic Typic Haplusterts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-8	Ap	20.34	19.94	59.72	2.68	5.03	3.75	5.25	3.64	-	c	50.19	33.49
8-40	Bss1	19.61	22.76	57.62	1.94	2.59	5.28	4.96	4.85	-	c	43.22	29.05
40-70	Bss2	21.25	17.65	61.10	3.02	5.26	3.91	5.48	3.58	-	c	44.30	30.25
70-120	Bss3	19.08	22.29	58.63	1.75	5.04	3.84	5.15	3.29	-	c	43.26	30.31
120-170	Bss4	11.11	20.44	68.45	2.04	1.93	1.70	2.83	2.61	-	c	51.33	33.51

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO <sub>3</sub>	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP			
	Water	CaCl <sub>2</sub>	M KCl				dS m <sup>-1</sup>	%	%	Ca	Mg					K	Na	Total
										cmol kg <sup>-1</sup>						%	%	
0-8	8.2	-	-	0.284	0.72	4.94	-	-	1.20	0.34	-	52.70	0.88	100	0.65			
8-40	8.44	-	-	0.139	0.40	7.28	-	-	0.30	0.48	-	52.06	0.90	100	0.93			
40-70	8.32	-	-	0.202	0.40	6.37	-	-	0.18	0.40	-	52.52	0.86	100	0.77			
70-120	9.3	-	-	0.282	0.36	6.89	-	-	0.27	0.38	-	50.97	0.87	100	0.75			
120-170	8.47	-	-	0.305	0.37	8.19	-	-	0.28	0.91	-	58.19	0.85	100	1.57			





## INTERPRETATION FOR LAND RESOURCE MANAGEMENT

The most important soil and site characteristics that affect the land use and conservation needs of an area are land capability, soil depth, soil texture, coarse fragments, available water capacity, soil slope, soil erosion, soil reaction etc. These are interpreted from the data base generated through land resource inventory and several thematic maps are generated. These would help in identifying the areas suitable for growing crops and, soil and water conservation measures and structures needed thus helping to maintain good soil health for sustained crop production. The various interpretative and thematic maps generated are described below.

### 5.1 Land Capability Classification

Land capability classification is an interpretative grouping of soil map units (soil phases) mainly based on inherent soil characteristics, external land features and environmental factors that limit the use of land for agriculture, pasture, forestry, or other uses on a sustained basis (IARI, 1971). The land and soil characteristics used to group the land resources in an area into various land capability classes, subclasses and units are

*Soil Characteristics:* Depth, texture, gravelliness, calcareousness.

*Land characteristics:* Slope, erosion, drainage, rock outcrops.

*Climate:* Total rainfall and its distribution, and length of crop growing period.

The Land capability classification system is divided into land capability classes, subclasses and units based on the level of information available. Eight land capability classes are recognized. They are

*Class I:* They are very good lands that have no limitations or very few limitations that restrict their use.

*Class II:* They are good lands that have minor limitations and require moderate conservation practices.

*Class III:* They are moderately good lands that have moderate limitations that reduce the choice of crops or that require special conservation practices.

*Class IV:* They are fairly good lands that have very severe limitations that reduce the choice of crops or that require very careful management.

*Class V:* Soils in these lands are not likely to erode, but have other limitations like wetness that are impractical to remove and as such not suitable for agriculture, but suitable for pasture or forestry with minor limitations.

*Class VI:* The lands have severe limitations that make them generally unsuitable for cultivation, but suitable for pasture or forestry with moderate limitations.

*Class VII:* The lands have very severe limitations that make them unsuitable for cultivation, but suitable for pasture or forestry with major limitations.

*Class VIII:* Soil and other miscellaneous areas (rock lands) that have very severe limitations that nearly preclude their use for any crop production, but suitable for wildlife, recreation and installation of wind mills.

The land capability subclasses are recognised based on the dominant limitations observed within a given land capability class. The subclasses are designated by adding a lower case letter like 'e', 'w', 's', or 'c' to the class numeral. The subclass "e" indicates that the main hazard is risk of erosion, "w" indicates drainage or wetness as a limitation for plant growth, "s" indicates shallow soil depth, coarse or heavy textures, calcareousness, salinity/alkalinity or gravelliness and "c" indicates limitation due to climate.

The land capability subclasses have been further subdivided into land capability units based on the kinds of limitations present in each subclass. Ten land capability units are used in grouping the soil map units. They are stony or rocky (0), erosion hazard (slope, erosion) (1), coarse texture (sand, loamy sand, sandy loam) (2), fine texture (cracking clay, silty clay) (3), slowly permeable subsoil (4), coarse underlying material (5), salinity/alkali (6), stagnation, overflow, high ground water table (7), soil depth (8) and fertility problems (9). The capability units thus identified have similar soil and land characteristics that respond similarly to a given level of management. The soils of the microwatershed have been classified upto land capability subclass level.

The 7 soil map units identified in the Himalapur microwatershed are grouped under 3 land capability classes and 3 subclasses. An area about 330 ha (66%) in the microwatershed is suitable for agriculture (Fig. 5.1). About 152 ha covered by forest and others (Habitation & water bodies) cover an area of about 17 ha in the microwatershed.




Fig. 5.1 Land Capability map of Himalapur Microwatershed

Good cultivable lands (Class II) cover an area of 159 ha (32%) and are distributed in the central, eastern, western and northern part of the microwatershed. They have minor limitations of soil and erosion. Moderately cultivable lands (Class III) cover an area of about 7 ha (1%) and are distributed in the eastern part of the microwatershed. They have moderate limitations of soil and erosion. Fairly cultivable lands (Class IV) cover about 163 ha (33%) and are distributed in the major part of the microwatershed. They have severe limitations of soil and erosion.

## 5.2 Soil Depth

Soil depth refers to the depth of the soil occurring above the parent material or hard rock. The depth of the soil determines the effective rooting depth for plants and in accordance with soil texture, mineralogy and gravel content, the capacity of the soil column to hold water and nutrient availability. Soil depth is one of the most important soil characteristic that is used in differentiating soils into different soil series. The soil depth classes used in identifying soils in the field are very shallow (<25 cm), shallow (25-50 cm), moderately shallow (50-75 cm), moderately deep (75-100 cm), deep (100-150 cm) and very deep (>150 cm). They were used to classify the soils into different depth classes and a soil depth map was generated. The area extent and their geographical distribution in the microwatershed is given in Fig. 5.2.




Fig. 5.2 Soil Depth map of Himalapur Microwatershed

Very shallow (<25 cm) soils cover an area of 94 ha (19%) and are distributed in the northern and southern part of the microwatershed. Shallow (25-50 cm) soils cover an area of 77 ha (15%) and are distributed in the northern, southern and eastern part of the microwatershed. Moderately shallow (50-75 cm) soils cover an area of 25 ha (5%) and are distributed in the northern part of the microwatershed. Deep (100-150 cm) soils cover an area of 74 ha (15%) and are distributed in the central, eastern and western part of the microwatershed. Very deep (>150 cm) soils cover an area of 60 ha (12%) and are distributed in the western and eastern part of the microwatershed.

The most productive lands 134 ha (27%) with respect to soil rooting depth where all climatically adapted annual and perennial crops can be grown are deep to very deep (100 to >150 cm depth) soils occurring in the major part of the microwatershed.

### 5.3 Surface Soil Texture

Texture is an expression to indicate the coarseness or fineness of the soil as determined by the relative proportion of primary particles of sand, silt and clay. It has a direct bearing on the structure, porosity, adhesion and consistence. The surface layer of a soil to a depth of about 25 cm is the layer that is most used by crops and plants. The surface soil textural class provides a guide to understanding soil-water retention and availability, nutrient holding capacity, infiltration, workability, drainage, physical and chemical behaviour, microbial activity and crop suitability. The textural classes used for LRI were used to classify and a surface soil texture map was generated. The area extent and their geographical distribution in the microwatershed is shown in Figure 5.3.




Fig. 5.3 Surface Soil Texture map of Himalapur Microwatershed

An area of about 42 ha (8%) area is sandy and is distributed in the southern part of the microwatershed. An area of 287 ha (57%) has soils that are clayey at the surface and occur in the major part of the microwatershed.

Major area of (57%) the microwatershed has most productive with respect to surface soil texture. The clayey soils (57%) have high potential for soil-water retention and availability, and nutrient retention and availability, but have more problems of drainage, infiltration, workability and other physical problems. The other productive lands are loamy soils (8%) which also have high potential for soil-water retention and nutrient availability but have no drainage or other physical problems.

#### 5.4 Soil Gravelliness

Gravel is the term used for describing coarse fragments between 2 mm and 7.5 cm diameter and stones for those between 7.5 cm and 25 cm. The presence of gravel and stones in soil reduces the volume of soil responsible for moisture and nutrient storage, drainage, infiltration and runoff, and hinders plant growth by impeding root growth and seedling emergence, intercultural operations and farm mechanization. The gravelliness classes used in LRI were used to classify the soils and using these classes a gravelliness map was generated. The area extent and their geographic distribution in the microwatershed is shown in Figure 5.4.




Fig. 5.4 Soil Gravelliness map of Himalapur Microwatershed

Entire cultivated area in the microwatershed has non gravelly (<15%) soils. These are the most productive soils, where all climatically adapted long duration crops can be grown.

### 5.5 Available Water Capacity

The soil available water capacity (AWC) is estimated based on the ability of the soil column to retain water between the tensions of 0.33 and 15 bar in a depth of 100 cm or the entire solum if the soil is shallower. The AWC of the soils (soil series) as estimated by considering the soil texture, mineralogy, soil depth and gravel content (Sehgal *et al.*, 1990) and accordingly the soil map units were grouped into five AWC classes *viz*, very low (<50 mm/m), low (50-100 mm/m), medium (100-150 mm/m), high (150-200 mm/m) and very high (>200 mm/m) and using these values, an AWC map was generated. The area extent and their geographic distribution of different AWC classes in the microwatershed is given in Figure 5.5.




Fig. 5.5 Soil Available Water Capacity map of Himalapur Microwatershed

An area of about 25 ha (5%) and 170 ha (34%) in the microwatershed has soils that are low (51-100 mm/m) and very low (<50mm/m) available water capacity and are distributed in the major part of the microwatershed. Very high (>200 mm/m) in an area of 134 ha (27%) and are distributed in the central, western and eastern part of the microwatershed.

About 195 ha (39%) area in the microwatershed has soils that are problematic with regard to available water capacity. Here, only short duration crops can be grown and the probability of crop failure is very high. These areas are best put to other alternative uses. An area of 134 ha (27%) are potential areas with regard to AWC where all climatically adapted annual and perennial crops can be grown.

## 5.6 Soil Slope

Soil slope refers to the inclination of the surface of the land. It is defined by gradient, shape and length, and is an integral feature of any soil as a natural body. Slope is considered important in soil genesis, land use and land development. The length and gradient of slope influences the rate of runoff, infiltration, erosion and deposition. The soil map units were grouped into two slope classes and a slope map was generated showing the area extent and their geographic distribution in the microwatershed (Fig. 5.6).

Maximum area of about 288 ha (58%) falls under very gently sloping (1-3% slope) lands and are distributed in the major part of the microwatershed. An area of about

42 ha (8%) falls under gently sloping (3-5% slope) lands and is distributed in the southern part of the microwatershed.




Fig. 5.6 Soil Slope map of Himalapur Microwatershed

An area of about 288 ha (58%) in the microwatershed, have high potential in respect of soil slopes. In these areas, all climatically adapted annual and perennial crops can be grown without much soil and water conservation and other land development measures.

### 5.7 Soil Erosion

Soil erosion refers to the wearing away of the earth's surface by the forces of water, wind and ice involving detachment and transport of soil by raindrop impact. It is used for accelerated soil erosion resulting from disturbance of the natural landscape by burning, excessive grazing and indiscriminate felling of forest trees and tillage, all usually by man. The erosion classes showing an estimate of the current erosion status as judged from field observations in the form of rills, gullies or a carpet of gravel on the surface are recorded. Four erosion classes, viz, slight erosion (e1), moderate erosion (e2), severe erosion (e3) and very severe erosion (e4) are recognized. The soil map units were grouped into different erosion classes and a soil erosion map generated. The area extent and their spatial distribution in the microwatershed is given in Figure 5.7.

Soils that are moderately eroded (e2 class) cover an area of 166 ha (33%) and are distributed in the central, western, eastern and northern part of the microwatershed. Soils



that are severely eroded (e3 class) cover an area of 163 ha (33%) and are distributed in the northern, eastern and southern part of the microwatershed.

An area of 329 ha (66%) in the microwatershed is problematic because of moderate to severe erosion. For these areas, taking up soil and water conservation and other land development measures are needed.




Fig. 5.7 Soil Erosion map of Himalapur Microwatershed



## **FERTILITY STATUS**

Soil fertility plays an important role in increasing crop yield. The adoption of high yielding varieties that require high amounts of nutrients has resulted in deficiency symptoms in crops and plants due to imbalanced fertilization and poor inherent fertility status as these areas are characterised by low rainfall and high temperatures. Hence, it is necessary to know the fertility (macro and micro nutrients) status of the soils of the watersheds for assessing the kind and amount of fertilizers required for each of the crop intended to be grown. For this purpose, the surface soil samples collected from the grid points (one soil sample at every 320 m interval) all over the microwatershed through land resource inventory in the year 2017 were analysed for pH, EC, organic carbon, available phosphorus and potassium, and for micronutrients like zinc, boron, copper, iron and manganese, and secondary nutrient sulphur.

Soil fertility data generated has been assessed and individual maps for all the nutrients for the microwatershed have been generated using Kriging method under GIS. The village/survey number wise fertility data for the microwatershed is given in Appendix-II.

### **6.1 Soil Reaction (pH)**

The soil analysis of the Himalapur microwatershed for soil reaction (pH) showed that an area of about 228 ha (46%) is neutral (pH 6.5-7.3) and are distributed in the major part of the microwatershed. About 101 ha (20%) is slightly alkaline (pH 7.3-7.8) and are distributed in the eastern and northern part of the microwatershed.(Fig.6.1). In all, major area of about 101 ha is alkaline, 228 ha is under neutral soils.

### **6.2 Electrical Conductivity (EC)**

The Electrical Conductivity of the soils in the microwatershed area is  $<2 \text{ dS m}^{-1}$  (Fig 6.2) and as such the soils are non-saline.

### **6.3 Organic Carbon**

The soil organic carbon content (an index of available Nitrogen) is high ( $>0.75\%$ ) in the entire cultivated area of the microwatershed (Fig. 6.3).




Fig.6.1 Soil Reaction (pH) map of Himalapur Microwatershed




Fig.6.2 Electrical Conductivity (EC) map of Himalapur Microwatershed




Fig.6.3 Soil Organic Carbon map of Himalapur Microwatershed

#### 6.4 Available Phosphorus

Available phosphorus content is low (<23 kg/ha) which covers an area of about 237 ha (48%) and occur in the major part of the microwatershed. Available phosphorus content is medium (23-57 kg/ha) which covers an area of about 92 ha (19%) and occur in southern, central and northern part of the microwatershed. (Fig. 6.4).

#### 6.5 Available Potassium

Available potassium content is low (<145kg/ha) in an area of 32 ha (6%) and occur in the southern part of the microwatershed. Available potassium content is medium (145-337 kg/ha) in an area of 180 ha (36%) and occur in the major part of the microwatershed. High (>337kg/ha) which covers an area of about 117 ha (23%) and occur in the central, eastern and northern part of the microwatershed (Fig. 6.5).

#### 6.6 Available Sulphur

Available sulphur is low (<10 ppm) which covers a maximum area of 169 ha (34%) in the microwatershed and are distributed in the major part of the microwatershed. Available sulphur is medium (10-20 ppm) in an area of 160 ha (32%) and are distributed in the southern, western and northern part of the microwatershed (Fig. 6.6).

## 6.7 Available Boron

Available boron content is medium (0.5-1.0 ppm) in an area of 4 ha (<1%) and are distributed in the northern part of the microwatershed. Maximum area of about 326 ha (65%) is low (<0.5 ppm) in available boron and are distributed in the major part of the microwatershed (Fig. 6.7).

## 6.8 Available Iron

Available iron content is sufficient (>4.5 ppm) in the entire area of the microwatershed. (Fig 6.8).

## 6.9 Available Manganese

Available manganese content is sufficient (>1.0 ppm) in the entire cultivated area of the microwatershed (Fig 6.9).

## 6.10 Available Copper

Available copper content is sufficient (>0.2 ppm) in the entire cultivated area of the microwatershed area (Fig 6.10).




Fig.6.4 Soil Available Phosphorus map of Himalapur Microwatershed




Fig.6.5 Soil Available Potassium map of Himalapur Microwatershed




Fig.6.6 Soil Available Sulphur map of Himalapur Microwatershed




Fig.6.7 Soil Available Boron map of Himalapur Microwatershed




Fig.6.8 Soil Available Iron map of Himalapur Microwatershed






Fig.6.9 Soil Available Manganese map of Himalapur Microwatershed




Fig.6.10 Soil Available Copper map of Himalapur Microwatershed

## 6.11 Available Zinc

Available zinc content is deficient (<0.6 ppm) which covers a maximum area of 244 ha (49%) and are distributed in major part of the microwatershed and sufficient (>0.6 ppm) which cover an area of 86 ha (17%) and are distributed in the southern, eastern and northern part of the microwatershed (Fig 6.11).




Fig.6.11 Soil Available Zinc map of Himalapur Microwatershed

## LAND SUITABILITY FOR MAJOR CROPS

The soil and land resource units (soil phases) of Himalapur microwatershed were assessed for their suitability for growing food, fodder, fibre and other horticulture crops by following the procedure as outlined in FAO, 1976 and 1983. Crop requirements were developed for each of the crop from the available research data and also by referring to Naidu *et. al.* (2006) and Natarajan *et. al* (2015). The soil and land characteristics were matched with the crop requirement to arrive at the crop suitability. The soil and land characteristics (Table 7.1) and crop requirement (Table 7.2 to 7.30) are given at the end of the chapter. In FAO land suitability classification, two orders are recognized. Order S-Suitable and Order N-Not suitable. The orders have classes, subclasses and units. Order-S has three classes, Class S1-Highly Suitable, Class S2-Moderately Suitable and Class S3- Marginally Suitable. Order N has two classes, N1-Currently not Suitable and N2-Permanently not Suitable. There are no subclasses within the Class S1 as they will have very minor or no limitations for crop growth. Classes S2, S3 and N1 are divided into subclasses based on the kinds of limitations encountered. The limitations that affect crop production are ‘c’ for erratic rainfall and its distribution and length of growing period (LGP), ‘e’ for erosion hazard, ‘r’ for rooting condition, ‘t’ for lighter or heavy texture, ‘g’ for gravelliness or stoniness, ‘n’ for nutrient availability, ‘l’ for topography, ‘m’ for moisture availability, ‘w’ for drainage and ‘z’ for calcareousness. These limitations are indicated as lower case letters to the Class symbol. For example, moderately suitable lands with the limitations of soil depth and erosion are designated as S2re. For the microwatershed, the soil mapping units were evaluated and classified up to subclass level.

Using the above criteria, the soil map units of the microwatershed were evaluated and land suitability maps for 29 major annual and perennial crops were generated. The detailed information on the kind of suitability of each of the soil phase for the crops assessed are given village/ survey number wise for the microwatershed in Appendix-IV.

### 7.1 Land Suitability for Sorghum (*Sorghum bicolor*)

Sorghum is one of the major crop grown in Karnataka in an area of 10.47 lakh ha in Bijapur, Gulbarga, Raichur, Bidar, Belgaum, Dharwad, Bellary, Chitradurga, Mysore and Tumakuru districts. The crop requirements for growing sorghum (Table 7.2) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing sorghum was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.1.

An area of about 60 ha (12%) is highly suitable (Class S1) for growing sorghum and are distributed in the eastern and western part of the microwatershed. An area of about 99 ha (20%) is moderately suitable (Class S2) for growing sorghum and are distributed in the central, western, eastern and northern part of the microwatershed. They have minor

limitations of rooting depth and drainage. An area of about 77 ha (15%) is marginally suitable (Class S3) for growing sorghum and are distributed in the southern, eastern and northern part of the microwatershed with moderate limitation of rooting depth. Currently not suitable (Class N1) lands occur in an area of 94 ha (19%) and are distributed in the southern and northern part of the microwatershed with severe limitation of rooting depth.




Fig. 7.1 Land Suitability map of Sorghum

## 7.2 Land Suitability for Maize (*Zea mays*)

Maize is one of the most important food crop grown in an area of 13.37 lakh ha in almost all the districts of the State. The crop requirements for growing maize (Table 7.3) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing maize was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.2.

Moderately suitable (Class S2) lands for growing maize cover an area of about 159 ha (32%) and occur in the major part of the microwatershed. They have minor limitation of texture. An area of about 77 ha (15%) is marginally suitable (Class S3) for growing maize and are distributed in the southern, eastern and northern part of the microwatershed with moderate limitationS of rooting depth and texture. Currently not suitable (Class N1) lands occur in an area of 94 ha (19%) and are distributed in the southern and northern part of the microwatershed with severe limitation of rooting depth.




Fig. 7.2 Land Suitability map of Maize

### 7.3 Land Suitability for Bajra (*Pennisetum glaucum*)

Bajra is one of the most important millet crop grown in an area of 2.34 lakh ha in the northern districts of Karnataka state. The crop requirements for growing bajra (Table 7.4) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing bajra was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.3.

Moderately suitable (Class S2) lands for growing bajra cover an area of about 159 ha (32%) and occur in the major part of the microwatershed. They have minor limitations of texture, rooting depth and drainage. An area of about 77 ha (15%) is marginally suitable (Class S3) for growing bajra and are distributed in the southern, eastern and northern part of the microwatershed with moderate limitations of rooting depth and texture. Currently not suitable (Class N1) lands occur in an area of 94 ha (19%) and are distributed in the southern and northern part of the microwatershed with severe limitation of rooting depth.




Fig. 7.3 Land Suitability map of Bajra

#### 7.4 Land Suitability for Groundnut (*Arachis hypogaea*)

Groundnut is one of the major oilseed crop grown in an area of 6.54 lakh ha in Karnataka in most of the districts either as rainfed or irrigated crop. The crop requirements for growing groundnut (Table 7.5) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing groundnut was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.4.

Marginally suitable lands (Class S3) for growing groundnut occupy maximum area of about 236 ha (47%) and are distributed in the major part of the microwatershed. They have moderate limitations of texture, rooting depth and drainage. Currently not suitable (Class N1) lands occur in an area of 94 ha (19%) and are distributed in the southern and northern part of the microwatershed with severe limitation of rooting depth.




Fig. 7.4 Land Suitability map of Groundnut

### 7.5 Land Suitability for Sunflower (*Helianthus annus*)

Sunflower is one of the most important oilseed crop grown in an area of 3.56 lakh ha in the State in all the districts. The crop requirements for growing sunflower (Table 7.6) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing sunflower was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.5.

An area of about 60 ha (12%) is highly suitable (Class S1) for growing sunflower and are distributed in the eastern and western part of the microwatershed. An area of about 74 ha (15%) is moderately suitable (Class S2) for sunflower and are distributed in the central, eastern and western part of the microwatershed. They have minor limitation of drainage. Marginally suitable lands (Class S3) for growing sunflower occupy an area of about 25 ha (5%) and is distributed in the northern part of the microwatershed. They have moderate limitation of rooting depth. Currently not suitable (Class N1) lands occur in an area of 170 ha (34%) and are distributed in the major part of the microwatershed with severe limitation of rooting depth.




Fig. 7.5 Land Suitability map of Sunflower

## 7.6 Land suitability criteria for Red gram (*Cajanus Cajan*)

Redgram is one of the most important pulse crop grown in an area of 7.28 lakh ha in almost all the districts of the State. The crop requirements for growing red gram (Table 7.7) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing redgram was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.6.

Moderately suitable (Class S2) lands for growing redgram cover an area of about 134 ha (27%) and occur in the eastern, central and western part of the microwatershed. They have minor limitations of texture and drainage. Marginally suitable lands (Class S3) for growing redgram occupy an area about 102 ha (20%) and occur in the eastern, northern and southern part of the microwatershed. They have moderate limitations of rooting depth and texture. Currently not suitable (Class N1) lands occur in an area of 94 ha (19%) and are distributed in the southern and northern part of the microwatershed with severe limitation of rooting depth






Fig. 7.6 Land Suitability map of Redgram

### 7.7 Land Suitability for Bengal gram (*Cicer aerativum*)

Bengal gram is one of the most important pulse crop grown in about 9.39 lakh ha area in Bijapur, Raichur, Kalaburgi, Dharwad, Belgaum and Bellary districts. The crop requirements for growing Bengal gram (Table 7.8) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing Bengal gram was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.7.

Highly (Class S1) suitable lands for growing bengal gram occur in an area of 134 ha (27%) and are distributed in the central, western and eastern part of the microwatershed. An area of about 25 ha (5%) is moderately suitable (Class S2) for growing bengal gram and are distributed in the northern part of the microwatershed. They have minor limitation of rooting depth. Marginally suitable lands (Class S3) for growing bengalgram occupy an area of about 77 ha (15%) and occur in the southern, eastern and northern part of the microwatershed. They have moderate limitation of rooting depth. Currently not suitable (Class N1) lands occur in an area of 94 ha (19%) and are distributed in the southern and northern part of the microwatershed with severe limitation of rooting depth.




Fig. 7.7 Land Suitability map of Bengal gram.

### 7.8 Land Suitability for Cotton (*Gossypium hirsutum*)

Cotton is one of the most important fibre crop grown in the State in about 8.75 lakh ha area in Raichur, Dharwad, Belgaum, Kalaburgi, Bijapur, Bidar, Bellary, Chitradurga and Chamarajnagar districts. The crop requirements for growing cotton (Table 7.9) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing cotton was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.8.

Highly (Class S1) suitable lands for growing cotton occur in an area of 134 ha (27%) and are distributed in the central, eastern and western part of the microwatershed. An area of about 25 ha (5%) is moderately suitable (Class S2) for growing cotton and are distributed in the northern part of the microwatershed. They have minor limitation of rooting depth. Marginally suitable lands (Class S3) for growing cotton occupy an area of about 77 ha (15%) and occur in the southern, northern and eastern part of the microwatershed. They have moderate limitation of rooting depth. Currently not suitable (Class N1) lands occur in an area of 94 ha (19%) and are distributed in the southern and northern part of the microwatershed with severe limitation of rooting depth.




Fig. 7.8 Land Suitability map of Cotton

### 7.9 Land Suitability for Chilli (*Capsicum annuum*)

Chilli is one of the most important spice crop grown in about 0.42 lakh ha in Karnataka State. The crop requirements for growing chilli (Table 7.10) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing chilli was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.9.

An area of about 85 ha (17%) is moderately suitable (Class S2) for growing chilli and are distributed in the northern, eastern and western part of the microwatershed. They have minor limitations of texture, drainage and rooting depth. Marginally suitable lands (Class S3) occupy an area of about 151 ha (30%) and are distributed in the central, northern, western, southern and eastern part of the microwatershed. They have moderate limitations of texture, rooting depth and drainage. Currently not suitable (Class N1) lands occur in an area of 94 ha (19%) and are distributed in the southern and northern part of the microwatershed with severe limitation of rooting depth.




Fig 7.9 Land Suitability map of Chilli

### 7.10 Land Suitability for Tomato (*Lycopersicon esculentum*)

Tomato is one of the most important vegetable crop grown in about 0.61 lakh ha covering almost all the district of the state. The crop requirements for growing tomato (Table 7.11) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing tomato was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.10.

An area of about 85 ha (17%) is moderately suitable (Class S2) for growing tomato and are distributed in the northern, eastern and western part of the microwatershed. They have minor limitations of texture, drainage and rooting depth. Marginally suitable lands (Class S3) occupy an area of about 151 ha (30%) and are distributed in the major part of the microwatershed. They have moderate limitations of texture, rooting depth and drainage. Currently not suitable (Class N1) lands occur in an area of 94 ha (19%) and are distributed in the southern and northern part of the microwatershed with severe limitation of rooting depth.




Fig 7.10 Land Suitability map of Tomato

### 7.11 Land Suitability for Brinjal (*Solanum melongena*)

Brinjal is one of the most important vegetable crop grown in the state. The crop requirements for growing brinjal (Table 7.12) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing Brinjal was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.11.

Moderately suitable (Class S2) lands for growing brinjal cover an area of about 159 ha (32%) and occur in the major part of the microwatershed. They have minor limitations of texture and rooting depth. An area of about 77 ha (15%) is marginally suitable (Class S3) for growing brinjal and are distributed in the southern, eastern and northern part of the microwatershed with moderate limitation of rooting depth. Currently not suitable (Class N1) lands occur in an area of 94 ha (19%) and are distributed in the southern and northern part of the microwatershed with severe limitation of rooting depth.




Fig 7.11 Land Suitability map of Brinjal

## 7.12 Land Suitability for Onion (*Allium cepa L.*)

Onion is one of the most important vegetable crop grown in the state. The crop requirements for growing onion (Table 7.13) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing onion was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.12.

Moderately suitable lands (Class S2) for growing onion occupy an area of about 25 ha (5%) and are distributed in the northern part of the microwatershed. They have minor limitation of rooting depth. Marginally (Class S3) suitable lands for growing onion occur in an area of 211 ha (42%) and are distributed in the major part of the microwatershed. Currently not suitable (Class N1) lands occur in an area of 94 ha (19%) and are distributed in the southern and northern part of the microwatershed with severe limitation of rooting depth.




Fig 7.12 Land Suitability map of Onion

### 7.13 Land Suitability for Bhendi (*Abelmoschus esculentus*)

Bhendi is one of the most important vegetable crop grown in the state. The crop requirements for growing bhendi (Table 7.14) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing bhendi was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.13.

Moderately suitable (Class S2) lands for growing bhendi cover an area of about 159 ha (32%) and occur in the major part of the microwatershed. They have minor limitation of texture and rooting depth. An area of about 77 ha (15%) is marginally suitable (Class S3) for growing bhendi and are distributed in the southern, eastern and northern part of the microwatershed with moderate limitation of rooting depth. Currently not suitable (class N1) lands occur in an area of 94 ha (19%) and are distributed in the southern and northern part of the microwatershed with severe limitation of rooting depth.




Fig 7.13 Land Suitability map of Bhendi

#### 7.14 Land Suitability for Drumstick (*Moringa oleifera*)

Drumstick is one of the most important vegetable crop grown in about 2403 ha in the state. The crop requirements for growing drumstick (Table 7.15) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing drumstick was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.14.

Moderately suitable (Class S2) lands for growing drumstick cover an area of about 134 ha (27%) and occur in the central, eastern and western part of the microwatershed. They have minor limitations of texture and drainage. An area of about 25 ha (5%) is marginally suitable (Class S3) for growing drumstick and are distributed in the northern part of the microwatershed. They have moderate limitation of rooting depth. Currently not suitable (Class N1) lands occur in an area of 171 ha (34%) and are distributed in the major part of the microwatershed with severe limitations of texture and rooting depth.






Fig 7.14 Land Suitability map of Drumstick

### 7.15 Land suitability for Mango (*Mangifera indica*)

Mango is one of the most important fruit crop grown in an area of 1.73 lakh ha in almost all the districts of the State. The crop requirements (Table 7.16) for growing mango were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing mango was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed is given in Figure 7.15.

Marginally suitable (Class S3) lands for growing mango cover an area of about 134 ha (27%) and occur in the central, eastern and western part of the microwatershed. They have moderate limitation of texture. Currently not suitable (Class N1) for growing mango occupy an area about 195 ha (39%) and occur in the major part of the microwatershed. They have severe limitation of rooting depth.




Fig. 7.15 Land Suitability map of Mango

### 7.16 Land suitability for Guava (*Psidium guajava*)

Guava is one of the most important fruit crop grown in an area of 0.06 lakh ha in almost all the districts of the State. The crop requirements (Table 7.17) for growing guava were matched with the soil-site characteristics (7.1) and a land suitability map for growing guava was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.16.

Marginally suitable (Class S3) lands for growing guava cover an area of about 159 ha (32%) and occur in the central, western, eastern and northern part of the microwatershed. They have moderate limitations of texture and rooting depth. Currently not suitable (Class N1) lands occur in an area of 171 ha (34%) and are distributed in the major part of the microwatershed with severe limitations of texture and rooting depth.




Fig. 7.16 Land Suitability map of Guava

### 7.17 Land suitability for Sapota (*Manilkara zapota*)

Sapota is one of the most important fruit crop grown in an area of 29373 ha in almost all the districts of the State. The crop requirements (Table 7.18) for growing sapota were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing sapota was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.17.

Marginally suitable (Class S3) lands for growing sapota cover an area of about 159 ha (32%) and occur in the central, western, and northern part of the microwatershed. They have minor limitations of texture and rooting depth. Currently not suitable (Class N1) lands occur in an area of 170 ha (34%) and are distributed in the major part of the microwatershed with severe limitation of rooting depth.




Fig. 7.17 Land Suitability map of Sapota

### 7.18 Land Suitability for Pomegranate (*Punica granatum*)

Pomegranate is one of the most important fruit crop commercially grown in about 18488 ha in Karnataka, mainly in Bijapur, Bagalkot, Koppal, Gadag and Chitradurga districts. The crop requirements for growing pomegranate (Table 7.19) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing pomegranate was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.18.

Moderately suitable (Class S2) lands for growing pomegranate cover an area of about 134 ha (27%) and occur in the central, eastern and western part of the microwatershed. They have minor limitation of texture. An area of about 25 ha (5%) is marginally suitable (Class S3) for growing pomegranate and are distributed in the northern part of the microwatershed. They have moderate limitation of rooting depth. Currently not suitable (Class N1) lands occur in an area of 170 ha (34%) and are distributed in the major part of the microwatershed with severe limitation of rooting depth.




Fig 7.18 Land Suitability map of Pomegranate

### 7.19 Land Suitability for Musambi (*Citrus limetta*)

Musambi is one of the important fruit crop grown in an area of 3446 ha in almost all the districts of the State. The crop requirements for growing musambi (Table 7.20) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing musambi was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.19.

Highly suitable (Class S1) lands for growing musambi cover an area of about 134 ha (27%) and occur in the central, eastern and western part of the microwatershed. An area of about 25 ha (5%) is marginally suitable (Class S3) for growing musambi and are distributed in the northern part of the microwatershed. They have moderate limitations of rooting depth. Currently not suitable (Class N1) lands occur in an area of 170 ha (34%) and are distributed in the major part of the microwatershed with severe limitation of rooting depth.




Fig. 7.19 Land Suitability map of Musambi

## 7.20 Land Suitability for Lime (*Citrus sp*)

Lime is one of the most important fruit crop grown in an area of 0.11 lakh ha in almost all the districts of the State. The crop requirements for growing lime (Table 7.21) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing lime was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7. 20.

Highly suitable (Class S1) lands for growing lime cover an area of about 134 ha (27%) and occur in the central, eastern and western part of the microwatershed. An area of about 25 ha (5%) is marginally suitable (Class S3) for growing lime and are distributed in the northern part of the microwatershed. They have moderate limitation of rooting depth. Currently not suitable (Class N1) lands occur in an area of 170 ha (34%) and are distributed in the southern, northern and eastern part of the microwatershed with severe limitations of rooting depth.




Fig. 7.20 Land Suitability map of Lime

### 7.21 Land Suitability for Amla (*Phyllanthus emblica*)

Amla is one of the medicinal, fruit crop grown in almost all the districts of the State. The crop requirements for growing amla (Table 7.22) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing amla was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.21.

Moderately suitable (Class S2) lands for growing amla cover an area of about 159 ha (32%) and occur in the major part of the microwatershed. They have minor limitations of texture and rooting depth. An area of about 77 ha (15%) is marginally suitable (Class S3) for growing amla and are distributed in the southern, eastern and northern part of the microwatershed with moderate limitations of rooting depth and texture. Currently not suitable (Class N1) lands occur in an area of 94 ha (19%) and are distributed in the southern and northern part of the microwatershed with severe limitation of rooting depth.




Fig. 7.21 Land Suitability map of Amla

## 7.22 Land Suitability for Cashew (*Anacardium occidentale*)

Cashew is one of the most important plantation nut crop grown in an area of 0.7 lakh ha in almost all the districts of the State. The crop requirements for growing cashew (Table 7.23) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing cashew was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.22.

Currently not suitable (Class N1) lands for growing cashew occur a maximum area of 330 ha (66%) and are distributed in the entire cultivated area of the microwatershed with severe limitations of rooting depth and texture.






Fig. 7.22 Land Suitability map of Cashew

### 7. 23 Land Suitability for Jackfruit (*Artocarpus heterophyllus*)

Jackfruit is one of the most important fruit crop grown in an area of 5368 ha in almost all the districts of the State. The crop requirements for growing jackfruit (Table 7.24) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing jackfruit was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.23.

Marginally suitable (Class S3) lands for growing jackfruit cover an area of about 159 ha (32%) and occur in the central, western, eastern and northern part of the microwatershed. They have moderate limitations of texture and rooting depth. Currently not suitable (Class N1) lands occur in an area of 171 ha (34%) and are distributed in the major part of the microwatershed with severe limitations of texture and rooting depth.




Fig. 7.23 Land Suitability map of Jackfruit

#### 7.24 Land Suitability for Jamun (*Syzygium cumini*)

Jamun is an important fruit crop grown in almost all the districts of the State. The crop requirements for growing jamun (Table 25) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing jamun was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.24.

Moderately suitable (Class S2) lands for growing jamun cover an area of about 134 ha (27%) and occur in the central, eastern and western part of the microwatershed. They have minor limitation of texture. An area of about 25 ha (5%) is marginally suitable (Class S3) for growing jamun and are distributed in the northern part of the microwatershed. They have moderate limitation of rooting depth. Currently not suitable (Class N1) lands occur in an area of 171 ha (34%) and are distributed in the major part of the microwatershed with severe limitations of texture and rooting depth.




Fig. 7.24 Land Suitability map of Jamun

### 7.25 Land Suitability for Custard Apple (*Annona reticulata*)

Custard apple is one of the most important fruit crop grown in almost all the districts of the State. The crop requirements for growing custard apple (Table 7.26) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing custard apple was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.25.

Highly (Class S1) suitable lands for growing custard apple occur in an area of 134 ha (27%) and are distributed in the major part of the microwatershed. An area of about 25 ha (5%) is moderately suitable (Class S2) for growing custard apple and are distributed in the northern part of the microwatershed. They have minor limitation of rooting depth. Marginally suitable lands (Class S3) for growing custard apple occupy an area of about 77 ha (15%) and occur in the southern, northern and eastern part of the microwatershed. They have moderate limitation of rooting depth. Currently not suitable (Class N1) lands occur in an area of 94 ha (19%) and are distributed in the southern and northern part of the microwatershed with severe limitation of rooting depth.




Fig. 7.25 Land Suitability map of Custard Apple

## 7.26 Land Suitability for Tamarind (*Tamarindus indica*)

Tamarind is one of the most important spice crop grown in almost all the districts of the state. The crop requirements for growing tamarind (Table 7.27) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing tamarind was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Fig. 7.26.

Moderately (Class S2) suitable lands for growing tamarind occur in an area of 134 ha (27%) and are distributed in the central, eastern and western part of the microwatershed. And they have minor limitation of texture. Currently not suitable lands (Class N1) for occur in an area about 196 ha (39%) and occur in the northern, eastern and southern part of the microwatershed. They have severe limitations of rooting depth and texture.




Fig. 7.26 Land Suitability map of Tamarind

### 7.27 Land Suitability for Mulberry (*Morus nigra*)

Mulberry is the important crop grown for rearing of silkworms in about 1.6 lakh ha area in all the districts of the state. The crop requirements for growing mulberry (Table 7.28) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing mulberry was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.27.

Marginally suitable (Class S3) lands for growing mulberry cover an area of about 159 ha (32%) and occur in the central, western, eastern and northern part of the microwatershed. They have moderate limitations of texture, drainage and rooting depth. Currently not suitable (Class N1) lands occur in an area of 171 ha (34%) and are distributed in the major part of the microwatershed with severe limitations of texture and rooting depth.




Fig 7.27 Land Suitability map of Mulberry

## 7.28 Land suitability for Marigold (*Tagetes sps.*)

Marigold is one of the most important flower crop grown in an area of 9108 ha in almost all the districts of the State. The crop requirements (Table 7.29) for growing marigold were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing marigold was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.28.

Moderately suitable (Class S2) lands for growing marigold cover an area of about 159 ha (32%) and occur in the major part of the microwatershed. They have minor limitations of texture, rooting depth and drainage. An area of about 77 ha (15%) is marginally suitable (Class S3) for growing marigold and are distributed in the southern, eastern and northern part of the microwatershed with moderate limitations of rooting depth and texture. Currently not suitable (Class N1) lands occur in an area of 94 ha (19%) and are distributed in the southern and northern part of the microwatershed with severe limitation of rooting depth.




Fig. 7.28 Land Suitability map of Marigold

### 7.29 Land Suitability for Chrysanthemum (*Dendranthema grandiflora*)

Chrysanthemum is one of the most important flower crop grown in an area of 4978 ha in almost all the districts of the State. The crop requirements (Table 7.30) for growing chrysanthemum were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing chrysanthemum was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.29.

Moderately suitable (Class S2) lands for growing chrysanthemum cover an area of about 159 ha (32%) and occur in the major part of the microwatershed. They have minor limitations of texture, rooting depth and drainage. An area of about 77 ha (15%) is marginally suitable (Class S3) for growing chrysanthemum and are distributed in the southern, eastern and northern part of the microwatershed with moderate limitations of rooting depth and texture. Currently not suitable (Class N1) lands occur in an area of 94 ha (19%) and are distributed in the southern and northern part of the microwatershed with severe limitation of rooting depth.




Fig. 7.29 Land Suitability map of Chrysanthemum



**Table 7.1 Soil-Site Characteristics of Himalapur Microwatershed**

Soil Map Units	Climate (P) (mm)	Growing period (Days)	Drain-age Class	Soil depth (cm)	Soil texture		Gravelliness		AWC (mm/m)	Slope (%)	Erosion	pH	EC (dSm <sup>-1</sup> )	ESP (%)	CEC [Cmol (p <sup>+</sup> )kg <sup>-1</sup> ]	BS (%)
					Sur-face	Sub-surface	Surface (%)	Sub-surface (%)								
BDPiB3	866	150	WD	<25	sc	scl	<15	<15	<50	1-3	moderate	8.58	0.262	0.35	18.10	100
BDLbC3	866	150	WD	25-50	ls	sl	<15	<15	<50	3-5	severe	6.20	0.074	0.20	4.20	93
BDLiB2	866	150	WD	25-50	sc	sl	<15	<15	<50	1-3	moderate	6.20	0.074	0.20	4.20	93
BDLiB3	866	150	WD	25-50	sc	sl	<15	<15	<50	1-3	moderate	6.20	0.074	0.20	4.20	93
JNKmB2	866	150	WD	50-75	c	scl	<15	<15	51-100	1-3	moderate	8.42	0.148	0.18	14.50	100
NGPmB2	866	150	MWD	100-150	c	c	<15	<15	>200	1-3	moderate	7.42	0.24	0.22	67.10	100
BMNmB2	866	150	MWD	>150	c	c	<15	<15	>200	1-3	moderate	8.20	0.284	0.65	52.70	100

\*Symbols and abbreviations are according to Field Guide for LRI under Sujala-III Project, Karnataka

**Table 7.2 Land suitability criteria for Sorghum**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	26–30	30–34; 24–26	34–40; 20–24	>40; <20
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, c (red), c (black)	scl, cl	ls, sl	-
	pH	1:2.5	5.5-7.8	5.0-5.5 7.8-9.0	>9.0	-
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	10-15
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

**Table 7.3 Land suitability criteria for Maize**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	30-34	35-38 26-30	38-40 26-20	
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc	c (red), c (black)	ls, sl	-
	pH	1:2.5	5.5-7.8	5.0-5.5 7.8-9.0	>9.0	-
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	-
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

**Table 7.4 Land suitability criteria for Bajra**

Land use requirement		Rating				
Soil –site characteristics	Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)	
Climatic regime	Mean temperature in growing season	°C	28-32	33-38 24-27	39-40 20-23	<20
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm	500-750	400-500	200-400	<200
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl, scl, cl,sc,c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.8	5.0-5.5 7.8-9.0	5.5-6.0 >9.0	
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	15-35	35-60	>60	
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	1-3	3-5	5-10	>10

**Table 7.5 Land suitability criteria for Groundnut**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	24–33	22–24; 33–35	20–22; 35–40	<20; >40
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. Well drained	Poorly drained	Very Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl	sl,cl, sc	c (red), c (black), ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<35	35-60	>60	
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.6 Land suitability criteria for Sunflower**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	24–30	30–34; 20–24	34–38; 16–20	>38; <16
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	mod. Well drained	-	Poorly to very drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	cl, sc,c (red), c (black)	scl	ls, sl	-
	pH	1:2.5	6.5-7.8	7.8-8.4 5.5-6.5	8.4-9.0; 5.0-5.5	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.7 Land suitability criteria for Redgram**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	30-35(G) 20-25(AV) 15-18 (F&PS) 35-40(M)	25-30(G) 20-25 (AV) 12-15 (F&PS) 30-35(M)	20-25(G) 15-20(AV) 10-12 (F&PS) 25-30(M)	< 20 <15 <10 <25
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. Well drained	Poorly drained	Very Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, c (red)	c (black),sl, scl, cl	ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-9.0	5.0-5.5 >9.0	-
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-50	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<1.0	1.0-2.0	>2.0	
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.8 Land suitability criteria for Bengal gram**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	20–25	25–30; 15–20	30–35; 10–15	>35; <10
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. Well drained	Poorly drained	Very Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	c(black)	-	c (red), scl, cl, sc	ls, sl
	pH	1:2.5	6.0-7.8	5.0-6.0 7.8-9.0	>9.0	-
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	-
Erosion hazard	Slope	%	<3	3-5	5-10	>10



**Table 7.9 Land suitability criteria for Cotton**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	22-32	>32	<19	-
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well to moderately well	Poorly drained/Some what excessively drained	-	very poorly/excessively drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, c (red,black)	cl	scl	ls, sl
	pH	1:2.5	6.5-7.8	7.8-8.4	5.5-6.5 8.4->9.0	<5.5
	CEC	C mol (p+)Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	50-100	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	<3	3-5	-	>5

**Table 7.10 Land suitability criteria for Chilli**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	25-32	33-35 20-25	35-38 <20	>38
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc	c (black), sl	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.11 Land suitability criteria for Tomato**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	25-28	29-32 20-24	15-19 33-36	<15 >36
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl, scl, cl, sc, c (red)	-	ls, c(black)	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.12 Land suitability criteria for Brinjal**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	Well drained	Moderately well drained	Poorly drained	V. Poorly drained
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class				
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl, scl, cl, sc c (red)	-	ls, c (black)	-
	pH	1:2.5	6.0-7.3	7.3-8.4 5.0-6.0	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	>60
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.13 Land suitability criteria for Onion**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	20-30	30-35	35-40	>40
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately /imperfectly	-	Poorly to V poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl,scl,cl,sc,c (red)	-	c (Black),ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<1.0	1.0-2.0	2.0-4.0	<4
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.14 Land suitability criteria for Bhendi**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	25-28	29-32 20-24	15-19 33-36	<15 >36
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Imperfectly drained	Poorly to very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl,sc, c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.15 Land suitability criteria for Drumstick**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, scl, cl, c (red)	sl, c (black)	ls	s
	pH	1:2.5	6.0-7.3	5.0-5.5 7.3-7.8	5.5-6.0 7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<35	35-60	60-80	>80
Soil toxicity	Salinity (EC saturation extract)	ds/m				
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-10	-	>10

**Table 7.16 Land suitability criteria for Mango**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-32	24-27 33-35	36-40	20-24
	Min temp. before flowering	°C	10-15	15-22	>22	-
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration	Days				
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V. Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	-	ls, sl, c (black)	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>150	100-150	75-100	<75
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10



**Table 7.17 Land suitability criteria for Guava**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-32	33-36 24-27	37-42 20-23	
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	sl	c (black), ls	-
	pH	1:2.5	6.0-7.8	5.0-6.0	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.18 Land suitability criteria for Sapota**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-32	33-36 24-27	37-42 20-23	>42 <18
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	-	Poorly to very drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	sl	ls, c (black)	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.19 Land suitability criteria for Pomegranate**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	30-34	35-38 25-29	39-40 15-24	
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl,cl, sc, c (red)	c (black),sl	ls	-
	pH	1:2.5	5.5-7.8	7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.20 Land suitability criteria for Musambi**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-30	31-35 24-27	36-40 20-23	>40 <20
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately drained	poorly	Very poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c	sl	ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.21 Land suitability criteria for Lime**

Land use requirement		Rating				
Soil –site characteristics	Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)	
Climatic regime	Mean temperature in growing season	°C	28-30	31-35 24-27	36-40 20-23	>40 <20
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately drained	poorly	Very poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c	sl	ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.22 Land suitability criteria for Amla**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. well drained	Poorly drained	V. Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	c (black)	ls, sl	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15-35	35-60	60-80	-
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

**Table 7.23 Land suitability criteria for Cashew**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	32 to 34	28 to 32; 34 to 38	24 to 28; 38 to 40	<20; >40
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	-	sl, ls	c (black)
	pH	1:2.5	5.5-6.5	5.0-5.5 6.5-7.3	7.3-7.8	>7.8
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-10	>10	-

**Table 7.24 Land suitability criteria for Jackfruit**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. well	Poorly	V. Poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	-	sl, ls, c (black)	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	>60
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10-



**Table 7.25 Land suitability criteria for Jamun**

Land use requirement		Rating				
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well	Mod. well	Poorly	V.Poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c(red)	sl, c (black)	ls	-
	pH	1:2.5	6.0-7.8	5.0-6.0	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>150	100-150	50-100	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	>60
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

**Table 7.26 Land suitability criteria for Custard apple**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	Scl, cl, sc, c (red), c (black)	-	Sl, ls	-
	pH	1:2.5	6.0-7.3	5.5-6.0 7.3-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15-35	35-60	60-80	-
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	>5	-

**Table 7.27 Land suitability criteria for Tamarind**

Land use requirement		Rating				
Soil –site characteristics	Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)	
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod.well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl,sc, c (red)	sl, c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>150	100-150	75-100	<75
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

**Table 7.28 Land suitability criteria for Mulberry**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	24–28	22–24; 28–32	32–38; 22–18	>38; <18
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V. Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, cl, scl	c (red)	c (black), sl, ls	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.8-8.4	7.3-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	0-35	35-60	60-80	>80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

**Table 7.29 Land suitability criteria for Marigold**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	18-23	17-15 24-35	35-40 10-14	>40 <10
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl,scl, cl, sc, c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%				
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.30 Land suitability criteria for Chrysanthemum**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	18-23	17-15 24-35	35-40 10-14	>40 <10
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl,scl, cl, sc, c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%				
Erosion hazard	Slope	%	<3	3-5	5-10	>10

### 7.30 Land Management Units (LMUs)

The 7 soil map units identified in Himalapur microwatershed have been grouped into 3 Land Management Units (LMUs) for the purpose of preparing a Proposed Crop Plan. Land Management Units are grouped based on the similarities in respect of the type of soil, the depth of the soil, the surface soil texture, gravel content, AWC, slope, erosion etc. and a Land Management Units map (Fig. 7.30) has been generated. These Land Management Units are expected to behave similarly for a given level of management.

The map units that have been grouped into 3 Land Management Units along with brief description of soil and site characteristics are given below.

LUC	Soil map units	Soil and site characteristics
1	62.BMNmB2 49.NGPmB2	Deep to very deep, black calcareous clay soils (100 - >150cm), 1-3 % slopes, non-gravelly (<15 %), moderate erosion.
2	152.JNKmB2	Moderately shallow, sandy, clay loam soils (50-75cm), 1-3 % slopes, non-gravelly (<15%), moderate erosion.
3	3.BDLbC3 5.BDLiB2 6.BDLiB3 119.BDPiB3	Very shallow to shallow, sandy clay loam to sandy loam soils (<25 to 50cm) 1- 5% slopes, non-gravelly (<15%), moderate to severe erosion.




Fig. 7.30 Land Management Units Map- Himalapur Microwatershed

### **7.31 Proposed Crop Plan for Himalapur Microwatershed**

After assessing the land suitability for the 29 crops, the Proposed Crop Plan has been prepared for the 3 identified LMUs by considering only the highly (Class S1) and moderately (Class S2) suitable lands for each of the 29 crops. The resultant proposed crop plan is presented below in Table 7.31.



**Table 7.31 Proposed Crop Plan for Himalapur Microwatershed**

LMU	Soil Map Units	Survey Number	Soil and site characteristics	Field Crops/ Commercial crops	Horticulture Crops (Rainfed/Irrigated )	Suitable Interventions
1	62.BMNmB2 49.NGPmB2	<b>Chapetla :</b> 107,108,109 <b>Himalapura:</b> 11,18,19,20,22,23,24,25,26,27,34,35,36,37,38,39,40,41,42,43,44,45,46,47,49,74,281,283,284,304,305,306,307,308,312,313,315,316,317,318,319,320,321,322,323,324,325,326,327,328,33,330,332,333,334,335,336,337,338,339,340,341,342,343,372,373,385,386,387,388,389,390,391, 392, 393,395	Deep to very deep, black, calcareous clay soils (100 >150cm), 1-3 % slopes, non-gravelly (<15 %), moderate erosion.	Maize, Sorghum, Sunflower, Cotton, Red gram, Bengalgram, Bajra	<b>Fruit crops:</b> Lime, Musambi, Custard apple, Pomegranate <b>Vegetables:</b> Chilli, Bhendi <b>Flowers:</b> Marigold ,Chrysanthemum	Application of FYM, Biofertilizers and micronutrients, drip irrigation, mulching, suitable soil and water conservation practices
2	152.JNKmB2	<b>Chapetla :</b> 18,85,86,87,89,149,150, 151,152,153	Moderately shallow, sandy clay loam soils (50-75cm), 1-3% slopes, nongravelly(<15%), moderate erosion.	Maize, Sorghum, Bajra	<b>Fruit crops:</b> Amla, Custard apple <b>Vegetables:</b> Tomato, Chilli, Brinjal, Bhendi, Onion <b>Flowers:</b> Marigold, Chrysanthemum	Application of FYM, Biofertilizers and micronutrients, drip irrigation, mulching, suitable soil and water conservation practices
3	3.BDLbC3 5.BDLiB2 6.BDLiB3 119.BDPiB3	<b>Chapetla :</b> 90,91,98,99,100,101,102, 103,104,105, 106 <b>Himalapura:</b> 2,3,4,5,6,7,12,13,14,16,17,21,31,32,48,51,52,53,54,55,56,57,58,59,60,61,62,63,65,75,77,79,80,81,82,83,84,85,86,87,88,90,91,92,93,94,95,96,97,98,99,100,101,126,127,132,133,134,135,136,140,141,148,149,150,151,152,153,154,155,156,157,180,181,182,183,226,227,228,229,230,231,232,235,236,237,238,239,240,241,242,243,244,245,290,292,293,294,295,296,297,298,299,300,301,302,303,329,331,332,333,334,335,336,337,338,339,340,341,342,343,372,373,385,386,387,388,389,390,391,392,393,395	Shallow to very shallow, sandy clay loam to sandy loam soils (25-50 to <25cm) 1- 5% slopes, non-gravelly (<15%) moderate to severe erosion.	-	<b>Agri-Silvi-Pastur:</b> , <i>Styloxanthes hamata</i> , <i>Styloxanthes scabra</i>	Use of short duration varieties, sowing across the slope, drip irrigation and mulching is recommended



## SOIL HEALTH MANAGEMENT

### 8.1 Soil Health

Soil health is basic to plant health and plant health is basic to human and bovine health. Soil is fundamental to crop production. Without soil, no food could be produced nor would livestock be fed on a large scale. Because it is finite and fragile, soil is a precious resource that requires special care from its users.

Soil health or the capacity of the soil to function is critical to human survival. Soil health has been defined as: “the capacity of the soil to function as a living system without adverse effect on the ecosystem”. Healthy soils maintain a diverse community of soil organisms that help to form beneficial symbiotic associations with plant roots, recycle essential plant nutrients, improve soil structure with positive repercussions for soil, water and nutrient holding capacity and ultimately improve crop production and also contribute to mitigating climate change by maintaining or increasing its carbon content.

Functional interactions of soil biota with organic and inorganic components, air and water determine a soil’s potential to store and release nutrients and water to plants and to promote and sustain plant growth. Thus, maintaining soil health is vital to crop production and conserve soil resource base for sustaining agriculture.

#### **The most important characteristics of a healthy soil are**

- Good soil tilth
- Sufficient soil depth
- Good water storage and good drainage
- Adequate supply, but not excess of nutrients
- Large population of beneficial organisms
- Small proportion of plant pathogens and insect pests
- Low weed pressure
- Free of chemicals and toxins that may harm the crop
- Resistance to degradation
- Resilience when unfavorable conditions occur

#### **Characteristics of Himalapur Microwatershed**

- ❖ The soil phases identified in the microwatershed belonged to the soil series, of these, BDP series occupies maximum area of 94 ha (19%) followed by BMN 77 ha (15%), NGP 74 ha (15%), BMN 60 ha (12%), JNK 25 ha (5%).
- ❖ As per land capability classification an area of 330 ha in the microwatershed falls under arable land category (Class II, III & IV). The major limitations identified in the arable lands were soil and erosion.

- ❖ On the basis of soil reaction an area of 228 ha (46%) is neutral (pH 6.5 -7.3), about 101 ha (20%) is slightly alkaline (pH 7.3-7.8) in the microwatershed.

### **Soil Health Management**

The following actions are required to improve the current land husbandry practices that provide a sound basis for the successful adoption of sustainable crop production system.

#### **Alkaline soils**

Slightly alkaline soils cover an area about 101 ha in the microwatershed

1. Regular addition of organic manure, green manuring, green leaf manuring, crop residue incorporation and mulching needs to be taken up to improve the soil organic matter status.
2. Application of Biofertilizers (Azospirillum, Azotobacter, Rhizobium).
3. Application of 25% extra N and P (125 % RDN&P).
4. Application of ZnSO<sub>4</sub> – 12.5 kg/ha (once in three years).
5. Application of Boron – 5kg/ha (once in three years).

#### **Neutral soils**

About 228 ha is under neutral soils.

1. Regular addition of organic manure, green manuring, green leaf manuring, crop residue incorporation and mulching needs to be taken up to improve the soil organic matter status.
2. Application of Biofertilizers, (Azospirillum, Azotobacter, Rhizobium).
3. Application of 100 per cent RDF.
4. Need based micronutrient applications.

Besides the above recommendations, the best transfer of technology options are also to be adopted.

### **Soil Degradation**

Soil erosion is one of the major factors affecting the soil health in the microwatershed. Out of total 498 ha area in the microwatershed, about 329 ha is suffering from moderate to erosion. These areas need immediate soil and water conservation and, other land development and land husbandry practices for restoring soil health.

### **Dissemination of Information and Communication of Benefits**

Any large scale implementation of soil health management requires that supporting information is made available widely, particularly through channels familiar to farmers and extension workers. Given the very high priority attached to soil-health especially by the Central Government on issuing Soil-Health Cards to all the farmers, media outlets like Regional, State and National Newspapers, Radio and Dooradarshan programs in local languages but also modern information and communication

technologies such as Cellular phones and the Internet, which can be much more effective in reaching the younger farmers.

### **Inputs for Net Planning (Saturation Plan) and Interventions needed**

Net planning (Saturation Plan) in IWMP is focusing on preparation of

1. Soil and Water Conservation Plan for each plot or farm.
2. Productivity enhancement measures/ interventions for existing crops/livestock/other farm enterprises.
3. Diversification of farming mainly with perennial horticultural crops and livestock.
4. Improving livelihood opportunities and income generating activities.

In this connection, how various outputs of Sujala-III are of use in addressing these objectives of Net Planning (Saturation Plan) are briefly presented below.

- ❖ **Soil Depth:** The depth of a soil decides the amount of moisture and nutrients it can hold, what crops can be taken up or not, depending on the rooting depth and the length of growing period available for raising any crop. Deeper the soil, better for a wide variety of crops. If sufficient depth is not available for growing deep rooted crops, either choose medium or short duration crops or deeper planting pits need to be opened and additional good quality soil brought from outside has to be filled into the planting pits.
- ❖ **Surface Soil Texture:** Lighter soil texture in the top soil means, better rain water infiltration, less run-off and soil moisture conservation, less capillary rise and less evaporation losses. Lighter surface textured soils are amenable to good soil tilth and are highly suitable for crops like groundnut, root vegetables (carrot, radish, potato etc) but not ideal for crops that need stagnant water like lowland paddy. Heavy textured soils are poor in water infiltration and percolation. They are prone for sheet erosion; such soils can be improved by sand mulching. The technology that is developed by the AICRP-Dryland Agriculture, Vijayapura, Karnataka can be adopted.
- ❖ **Gravelliness:** More gravel content is favorable for run-off harvesting but poor in soil moisture storage and nutrient availability. It is a significant parameter that decides the kind of crop to be raised.
- ❖ **Land Capability Classification:** The land capability map shows the areas suitable and not suitable for agriculture and the major constraints in each of the plot/survey number. Hence, one can decide what kind of enterprise is possible in each of these units. In general, erosion and soil are the major constraints in Himalapur microwatershed.
- ❖ **Organic Carbon:** The OC content (an index of available Nitrogen) is High (>0.75%) in the entire microwatershed area.
- ❖ **Available Phosphorus:** Available Phosphorus is low (<23 kg/ha) in an area of 237 ha (48%), medium (23-57 kg/ha) in an area of 92 ha (19%) of the microwatershed. For


all the crops 25% additional P needs to be applied where available P is low and medium.

- ❖ **Available Potassium:** Available potassium is low (<145 kg/ha) in an area of 32 ha (6%), medium (145-337 kg/ha) in an area of 180 ha (36%) and high (>337 kg/ha) area of 117 ha (23%) the microwatershed. All the plots, where available potassium is low and medium, for all the crops, additional 25% potassium may be applied.
- ❖ **Available Sulphur:** Available sulphur is a very critical nutrient for oilseed crops, it is medium in 160 ha (32%) and low in 169 ha (34%). Low and medium areas need to be applied with magnesium sulphate or gypsum or Factamphos (p) fertilizer (13% sulphur) for 2-3 years for the deficiency to be corrected.
- ❖ **Available Boron:** An area of 326 ha (65%) is low and 4 ha (<1%) is medium in available boron. For these areas, application of sodium borate @ 10 kg/ha as soil application or 0.2 % borax as foliar spray is recommended.
- ❖ **Available Iron:** Available iron is sufficient (>4.5ppm) in the entire cultivated area of the microwatershed.
- ❖ **Available Manganese:** Entire cultivated area of the microwatershed is sufficient in available manganese content.
- ❖ **Available Copper:** Entire cultivated area of the microwatershed is sufficient in available copper content.
- ❖ **Available Zinc:** Available zinc content is deficient (<0.6 ppm) in a maximum area of 244 ha (49%) and sufficient (>0.6 ppm) cover an area of 86 ha (17%) of the microwatershed. Application of zinc sulphate 25 kg/ha is recommended for the deficient areas.
- ❖ **Land Suitability for various crops:** Areas that are highly, moderately and marginally suitable for growing various crops are indicated. Along with the suitability, various constraints that are limiting the productivity are also indicated. For example, in case of cotton, gravel content, rooting depth and salinity/alkalinity are the major constraints in various plots. With suitable management interventions, the productivity can be enhanced. In order to increase the water holding capacity of light textured soils, growing of green manure crops and application of organic manure is recommended.

## SOIL AND WATER CONSERVATION TREATMENT PLAN

For preparing soil and water conservation treatment plan for Himalapur microwatershed, the land resource inventory database generated under Sujala-III project has been transformed as information through series of interpretative (thematic) maps using soil phase map as a base. The various thematic maps (1:7920 scale) generated were

- Soil depth
- Surface soil texture
- Available water capacity
- Soil slope
- Soil gravelliness
- Land capability
- Present land use and land cover
- Crop suitability
- Rainfall
- Hydrology
- Water Resources
- Socio-economic data
- Contour plan with existing features- network of waterways, pothissa boundaries, cut up/ minor terraces etc.
- Cadastral map (1:7920 scale)
- Satellite imagery (1:7920 scale)



Apart from these, Hand Level/ Hydro Marker/ Dumpy Level/ Total Station and Kathedars' List to be collected.

### Steps for Survey and Preparation of Treatment Plan

The boundaries of Land User Groups' and Survey No. boundaries are traced in the field.

- Naming of user groups and farmers
- Identification of arable and non arable lands
- Identification of drainage lines and gullies
- Identification of non treatable areas
- Identification of priority areas in the arable lands
- Treatment plan for arable lands
- Location of water harvesting and recharge structures

## 9.1 Treatment Plan

The treatment plan recommended for arable lands is briefly described below

### 9.1.1 Arable Land Treatment

#### A. BUNDING

Steps for Survey and Preparation of Treatment Plan		<p><b>USER GROUP-1</b></p> <p><b>CLASSIFICATION OF GULLIES</b></p> <p>ಕೊರಕಲಿನ ವರ್ಗೀಕರಣ</p>
<ul style="list-style-type: none"> <li>• Cadastral map (1:7920 scale) is enlarged to a scale of 1:2500 scale</li> <li>• Existing network of waterways, pothissa boundaries, grass belts, natural drainage lines/ watercourse, cut ups/ terraces are marked on the cadastral map to the scale</li> <li>• Drainage lines are demarcated into</li> </ul>		
Small gullies	(up to 5 ha catchment)	
Medium gullies	(5-15 ha catchment)	
Ravines	(15-25 ha catchment) and	
<i>Halla/Nala</i>	(more than 25ha catchment)	

#### Measurement of Land Slope

Land slope is estimated or determined by the study and interpretation of contours or by measurement in the field using simple instruments like Hand Level or Hydromarker.



Vertical and Horizontal intervals between bunds as recommended by the Watershed Development Department.

Slope percentage	Vertical interval (m)	Corresponding Horizontal Distance (m)
2 - 3%	0.6	24
3 - 4%	0.9	21
4 - 5%	0.9	21
5 - 6%	1.2	21
6 - 7%	1.2	21



**Note:** (i) The above intervals are maximum.

(ii) Considering the slope class and erosion status (A1... A=0-1 % slope, 1= slight erosion) the intervals have to be decided.

**Bund length recording:** Considering the contour plan and the existing grass belts/partitions, the bunds are aligned and lengths are measured.

### Section of the Bund

Bund section is decided considering the soil texture class and gravelliness class (bg<sub>0</sub>... b=loamy sand, g<sub>0</sub> = <15% gravel). The recommended Sections for different soils are given below.

**Recommended Bund Section**



Top width (m)	Base width (m)	Height (m)	Side slope (Z:1;H:V)	Cross section (sq m)	Soil Texture	Remarks
0.3	0.9	0.3	01:01	0.18	Sandy loam	Vegetative bund
0.3	1.2	0.3	1.5:1	0.225	Sandy clay	
0.3	1.2	0.5	0.9:1	0.375	Red gravelly soils	
0.3	1.2	0.6	0.75:1	0.45		
0.3	1.5	0.6	01:01	0.54	Red sandy loam	
0.3	2.1	0.6	1.5:1	0.72	Very shallow black soils	
0.45	2	0.75	01:01	0.92		
0.45	2.4	0.75	1.3:1	1.07	Shallow black soils	
0.6	3.1	0.7	1.78:1	1.29	Medium black soils	
0.5	3	0.85	1.47:1	1.49		

### Formation of Trench cum Bund

Dimensions of the Borrow Pits/Trenches to be excavated (machinery are decided considering the Bund Section).


Details of Borrow Pit dimensions are given below:

#### TRENCH CUM BUND

IDEAL FOR HORTICULTURE CROPS

#### 'A' FRAME FOR INTERBUND MANAGEMENT



1. ಸಮವಾತಳ ಉಳಿವೆ
2. ಸಮವಾತಳ ಬಿತ್ತನೆ/ನಾಟಿ

### Size of Borrow Pits/ Trench recommended for Trench cum Bund (by machinery)

Bund section	Bund length	Earth quantity	Pit				Berm (pit to pit)	Soil depth class
			L(m)	W(m)	D(m)	Quantity (m <sup>3</sup> )		
m <sup>2</sup>	m	m <sup>3</sup>					m	
0.375	6	2.25	5.85	0.85	0.45	2.24	0.15	Shallow
0.45	6	2.7	5.4	1.2	0.43	2.79	0.6	Shallow
0.45	6	2.7	5	0.85	0.65	2.76	1	Moderately Shallow
0.54	5.6	3.02	5.5	0.85	0.7	3.27	0.1	Moderately shallow
0.54	5.5	2.97	5	1.2	0.5	3	0.5	Shallow
0.72	6.2	4.46	6	1.2	0.7	5.04	0.2	Moderately shallow
0.72	5.2	3.74	5.1	0.85	0.9	3.9	0.1	Moderately deep

#### B. Water Ways

1. Existing waterways are marked on the cadastral map (1:7920 scale) and their dimensions are recorded.
2. Considering the contour plan of the MWS, additional waterways/ modernization of the existing ones can be thought of.
3. The design details are given in the Manual.

#### C. Farm Ponds

Waterways and the catchment area will give an indication on the size of the Farm Pond. Location of the pond can be decided based on the contour plan/ field condition and farmers' need/desire.

#### D. Diversion Channel

Existing EPT/ CPT are marked on the cadastral map. Looking to the need, these can be modernized or fresh diversion channel can be proposed and runoff from this can be stored in Gokatte/ Recharge ponds.

#### 9.1.2 Non-Arable Land Treatment

Depending on the gravelliness and crops preferred by the farmers, the concerned authorities can decide appropriate treatment plan. The recommended treatments may be Contour Trench, Staggered Trench, Crescent Bund, Boulder Bund or Pebble Bund.

### **9.1.3 Treatment of Natural Water Course/ Drainage Lines**

- a) The cadastral map has to be updated as regards the network of drainage lines (gullies/ *nalas*/ *hallas*) and existing structures are marked to the scale and storage capacity of the existing water bodies are documented.
- b) The drainage line will be demarcated into Upper Reach, Middle Reach and Lower Reach.
- c) Considering the Catchment, *Nala* bed and bank conditions, suitable structures are decided.
- d) Number of storage structures (Check dam/ *Nala* bund/ Percolation tank) will be decided considering the commitments and available runoff from water budgeting and quality of water in the wells and site suitability.
- e) Detailed Leveling Survey using Dumpy Level / Total Station has to be carried out to arrive at the site-specific designs as shown in the Manual.
- f) The location of ground water recharge structures are decided by examining the lineaments and fracture zones from geological maps.
- g) Rainfall intensity data of the nearest Rain Gauge Station is considered for Hydrologic Designs.
- h) Silt load to the Storage/Recharge structures is reduced by providing vegetative, boulder and earthen checks in the natural water course. Location and design details are given in the Manual.

### **9.2 Recommended Soil and Water Conservation Measures**

The appropriate conservation structures best suited for each of the land parcel/ survey number (Appendix-I) are selected based on the slope per cent, severity of erosion, amount of rainfall, land use and soil type. The different kinds of conservation structures recommended are:

1. Graded / Strengthening of Bunds
2. Trench cum Bunds (TCB)
3. Trench cum Bunds / Strengthening
4. Crescent Bunds

A map (Fig. 9.1) showing soil and water conservation plan with different kinds of structures recommended has been prepared which shows the spatial distribution and extent of area. An area about 94 ha (19%) requires trench cum bunding, and 236 ha (47%) needs Graded Bunding.

The conservation plan prepared may be presented to all the stakeholders including farmers and after considering their suggestions, the conservation plan for the microwatershed may be finalised in a participatory approach.




Fig. 9.1 Soil and Water Conservation Plan map of Himalapur Microwatershed

### 9.3 Greening of Microwatershed

As part of the greening programme in the watersheds, it is envisaged to plant a variety of horticultural and other tree plants that are edible, economical and produce lot of biomass which helps to restore the ecological balance in the watersheds. The lands that are suitable for greening programme are non-arable lands (land capability classes V, VI VII and VIII) and also the lands that are not suitable or marginally suitable and field bunds for growing annual and perennial crops. The method of planting these trees is given below.

It is recommended to open pits during the 1<sup>st</sup> week of March along the contour and heap the dug out soil on the lower side of the slope in order to harness the flowing water and facilitate weathering of soil in the pit. Exposure of soil in the pit also prevents spread of pests and diseases due to scorching sun rays. The pits should be filled with mixture of soil and organic manure during the second week of April and keep ready with sufficiently tall seedlings produced either in poly bags or in root trainer nurseries so that planting can be done during the 2<sup>nd</sup> or 3<sup>rd</sup> week of April depending on the rainfall.

The tree species suitable for the area considering rainfall, temperature and adaptability is listed below; waterlogged areas are recommended to be planted with species like Nerale (*Syzyzium cumini*) and Bamboo. Dry areas are to be planted with species like Honge, Bevu, Seetaphal etc.

<b>Dry Deciduous Species</b>			<b>Temp (°C)</b>	<b>Rainfall (mm)</b>
1.	Bevu	<i>Azadiracta indica</i>	21–32	400 –1,200
2.	Tapasi	<i>Holoptelia integrifolia</i>	20-30	500 - 1000
3.	Seetaphal	<i>Anona Squamosa</i>	20-40	400 - 1000
4.	Honge	<i>Pongamia pinnata</i>	20 -50	500– 2,500
5.	Kamara	<i>Hardwickia binata</i>	25 -35	400 - 1000
6.	Bage	<i>Albezzia lebbek</i>	20 - 45	500 - 1000
7.	Ficus	<i>Ficus bengalensis</i>	20 - 50	500–2,500
8.	Sisso	<i>Dalbargia Sissoo</i>	20 - 50	500 -2000
9.	Ailanthus	<i>Ailanthus excelsa</i>	20 - 50	500 - 1000
10.	Hale	<i>Wrightia tinctoria</i>	25 - 45	500 - 1000
11.	Uded	<i>Steriospermum chelanoides</i>	25 - 45	500 -2000
12.	Dhupa	<i>Boswella Serrata</i>	20 - 40	500 - 2000
13.	Nelli	<i>Emblia Officinalis</i>	20 - 50	500 -1500
14.	Honne	<i>Pterocarpus marsupium</i>	20 - 40	500 - 2000
<b>Moist Deciduous Species</b>			<b>Temp (°C)</b>	<b>Rainfall (mm)</b>
15.	Teak	<i>Tectona grandis</i>	20 - 50	500-5000
16.	Nandi	<i>Legarstroemia lanceolata</i>	20 - 40	500 - 4000
17.	Honne	<i>Pterocarpus marsupium</i>	20 - 40	500 - 3000
18.	Mathi	<i>Terminalia alata</i>	20 -50	500 - 2000
19.	Shivane	<i>Gmelina arborea</i>	20 -50	500 -2000
20.	Kindal	<i>T.Paniculata</i>	20 - 40	500 - 1500
21.	Beete	<i>Dalbargia latifolia</i>	20 - 40	500 - 1500
22.	Tare	<i>T. belerica</i>	20 - 40	500 - 2000
23.	Bamboo	<i>Bambusa arundinasia</i>	20 - 40	500 - 2500
24.	Bamboo	<i>Dendrocalamus strictus</i>	20 – 40	500 – 2500
25.	Muthuga	<i>Butea monosperma</i>	20 - 40	400 - 1500
26.	Hippe	<i>Madhuca latifolia</i>	20 - 40	500 - 2000
27.	Sandal	<i>Santalum album</i>	20 - 50	400 - 1000
28.	Nelli	<i>Emblia officinalis</i>	20 - 40	500 - 2000
29.	Nerale	<i>Sizyium cumini</i>	20 - 40	500 - 2000
30.	Dhaman	<i>Grevia tilifolia</i>	20 - 40	500 - 2000
31.	Kaval	<i>Careya arborea</i>	20 - 40	500 - 2000
32.	Harada	<i>Terminalia chebula</i>	20 - 40	500 - 2000



## References

1. FAO (1976) Framework for Land Evaluation, Food and Agriculture Organization, Rome.72 pp.
2. FAO (1983) Guidelines for Land Evaluation for Rainfed Agriculture, FAO, Rome, 237 pp.
3. IARI (1971) Soil Survey Manual, All India Soil and Land Use Survey Organization, IARI, New Delhi, 121 pp.
4. Katyal, J.C. and Rattan, R.K. (2003) Secondary and Micronutrients; Research Gap and Future Needs. Fert. News 48 (4); 9-20.
5. Naidu, L.G.K., Ramamurthy, V., Challa, O., Hegde, R. and Krishnan, P. (2006) Manual Soil Site Suitability Criteria for Major Crops, NBSS Publ. No. 129, NBSS & LUP, Nagpur, 118 pp.
6. Natarajan, A. and Dipak Sarkar (2010) Field Guide for Soil Survey, National Bureau of Soil Survey and Land Use Planning (ICAR), Nagpur, India.
7. Natarajan, A., Rajendra Hegde, Raj, J.N. and Shivananda Murthy, H.G. (2015) Implementation Manual for Sujala-III Project, Watershed Development Department, Bengaluru, Karnataka.
8. Sarma, V.A.K., Krishnan, P. and Budihal, S.L. (1987) Laboratory Manual, Tech. Bull. 23, NBSS &LUP, Nagpur.
9. Sehgal, J.L. (1990) Soil Resource Mapping of Different States of India; Why and How?, National Bureau of Soil Survey and Land Use Planning, Nagpur, 49 pp.
10. Shivaprasad, C.R., R.S. Reddy, J. Sehgal and M. Velayuthum (1998) Soils of Karnataka for Optimizing Land Use, NBSS Publ. No. 47b, NBSS & LUP, Nagpur, India.
11. Soil Survey Staff (2006) Keys to Soil Taxonomy, Tenth edition, U.S. Department of Agriculture/ NRCS, Washington DC, U.S.A.
12. Soil Survey Staff (2012) Soil Survey Manual, Handbook No. 18, USDA, Washington DC, USA.





**Appendix-I**  
**Himalapur (6A2a) Microwatershed**  
**Soil Phase Information**

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Boodhura	138	0.2	BDLiB2	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Rock outcrops	Not Available	IIes	Graded bunding
Boodhura	139	0.59	BDLiB2	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Cotton (Rg+Ct)	Not Available	IIes	Graded bunding
Boodhura	140	2.33	BDLiB2	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Cotton (Rg+Ct)	Not Available	IIes	Graded bunding
Chapetla	18	1.71	JNKmB2	LMU-2	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Chapetla	85	1.84	JNKmB2	LMU-2	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Chapetla	86	0.18	JNKmB2	LMU-2	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Chapetla	87	6.03	JNKmB2	LMU-2	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Chapetla	89	3.17	JNKmB2	LMU-2	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Chapetla	90	5.44	BDPiB3	LMU-3	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Redgram (Rg)	Not Available	IVes	Trench cum bunding
Chapetla	91	0.7	BDPiB3	LMU-3	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Redgram (Rg)	Not Available	IVes	Trench cum bunding
Chapetla	98	4.88	BDLiB3	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Redgram (Rg)	Not Available	IVes	Graded bunding
Chapetla	99	4.21	BDLiB3	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Redgram (Rg)	Not Available	IVes	Graded bunding
Chapetla	100	4.42	BDPiB3	LMU-3	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Habitation	Not Available	IVes	Trench cum bunding
Chapetla	101	3.83	BDPiB3	LMU-3	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Redgram (Rg)	Not Available	IVes	Trench cum bunding
Chapetla	102	6.83	BDPiB3	LMU-3	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Jowar+Redgram (Jw+Rg)	Not Available	IVes	Trench cum bunding
Chapetla	103	5.44	BDPiB3	LMU-3	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Redgram (Rg)	Not Available	IVes	Trench cum bunding
Chapetla	104	3.63	BDPiB3	LMU-3	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Redgram (Rg)	Not Available	IVes	Trench cum bunding
Chapetla	105	7.11	BDPiB3	LMU-3	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Redgram (Rg)	Not Available	IVes	Trench cum bunding
Chapetla	106	5.51	BDPiB3	LMU-3	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Redgram (Rg)	Not Available	IVes	Trench cum bunding
Chapetla	107	1.06	BMNmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Chapetla	108	1.2	BMNmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Chapetla	109	0.08	BMNmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Chapetla	149	1.19	JNKmB2	LMU-2	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iles	Graded bunding
Chapetla	150	3.61	JNKmB2	LMU-2	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Chapetla	151	5.72	JNKmB2	LMU-2	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Chapetla	152	3.28	JNKmB2	LMU-2	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Chapetla	153	2.59	JNKmB2	LMU-2	Moderately shallow (50-75 cm)	Clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Himalapura	1	0.5	Habitation	Others	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others
Himalapura	2	2.09	BDLiB3	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Jowar+Redgram+Habitation (Jw+Rg+Hb)	Not Available	IVes	Graded bunding
Himalapura	3	0.09	BDLiB3	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Not Available (NA)	Not Available	IVes	Graded bunding
Himalapura	4	0.07	BDLiB3	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Not Available (NA)	Not Available	IVes	Graded bunding
Himalapura	5	0.56	BDLiB3	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Jowar (Jw)	Not Available	IVes	Graded bunding
Himalapura	6	0.26	BDLiB3	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Jowar (Jw)	Not Available	IVes	Graded bunding
Himalapura	7	1.95	BDLiB3	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Redgram (Rg)	Not Available	IVes	Graded bunding
Himalapura	11	0.02	BMNmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton+Paddy+Redgram (Ct+Pd+Rg)	Not Available	Iles	Graded bunding
Himalapura	12	0.7	BDLiB3	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Cotton (Ct)	Not Available	IVes	Graded bunding
Himalapura	13	4.03	BDLiB3	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Groundnut+Redgram (Gn+Rg)	Not Available	IVes	Graded bunding
Himalapura	14	0.08	BDLiB3	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Habitation	Not Available	IVes	Graded bunding
Himalapura	15	0.14	Habitation	Others	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others
Himalapura	16	2.95	BDLiB3	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Cotton (Ct)	Not Available	IVes	Graded bunding
Himalapura	17	2.47	BDLiB3	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Redgram (Rg)	Not Available	IVes	Graded bunding
Himalapura	18	1.69	BMNmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Habitation (Rg+Hb)	1 Borewell	Iles	Graded bunding
Himalapura	19	1.48	BMNmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Iles	Graded bunding
Himalapura	20	1.79	BMNmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut+Redgram (Gn+Rg)	Not Available	Iles	Graded bunding
Himalapura	21	1.09	BDLiB3	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Groundnut (Gn)	1 Borewell	IVes	Graded bunding
Himalapura	22	3.72	BMNmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Iles	Graded bunding

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Himalapura	23	4.03	NGPmB2	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy+Cotton (Pd+Ct)	Not Available	Iles	Graded bunding
Himalapura	24	1.72	BMNmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton+Jowar+Paddy (Ct+Jw+Pd)	Not Available	Iles	Graded bunding
Himalapura	25	1.41	BMNmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar+Cotton (Jw+Ct)	Not Available	Iles	Graded bunding
Himalapura	26	1.13	BMNmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Iles	Graded bunding
Himalapura	27	2.1	BMNmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton+Jowar+Paddy (Ct+Jw+Pd)	Not Available	Iles	Graded bunding
Himalapura	31	2.28	BDLiB2	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Rock outcrops (Rg+Rc)	Not Available	IIles	Graded bunding
Himalapura	32	2.26	BDLiB2	LMU-3	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIles	Graded bunding
Himalapura	33	0.89	NGPmB2	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Himalapura	34	1.63	NGPmB2	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Himalapura	35	1.2	NGPmB2	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Iles	Graded bunding
Himalapura	36	3.85	NGPmB2	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton+Paddy+Redgram (Ct+Pd+Rg)	Not Available	Iles	Graded bunding
Himalapura	37	2.38	NGPmB2	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Himalapura	38	1.75	BMNmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Himalapura	39	1.17	NGPmB2	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Himalapura	40	2.33	NGPmB2	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar+Redgram (Jw+Rg)	Not Available	Iles	Graded bunding
Himalapura	41	0.47	NGPmB2	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Himalapura	42	0.25	NGPmB2	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Himalapura	43	0.29	NGPmB2	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Himalapura	44	0.63	NGPmB2	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Himalapura	45	0.96	NGPmB2	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Himalapura	46	0.96	NGPmB2	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Himalapura	47	2.56	NGPmB2	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Himalapura	48	1.09	BDLbc3	LMU-3	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Redgram (Rg)	Not Available	IVes	Graded bunding
Himalapura	49	2	NGPmB2	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy+Cotton (Pd+Ct)	Not Available	Iles	Graded bunding

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Himalapura	50	3.12	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Not Available (NA)	Not Available	Forest	Forest
Himalapura	51	3.45	BDLbC3	LMU-3	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Paddy+Cotton (Pd+Ct)	Not Available	IVes	Graded bunding
Himalapura	52	0.86	BDLbC3	LMU-3	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Redgram (Rg)	Not Available	IVes	Graded bunding
Himalapura	53	1	BDLbC3	LMU-3	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Paddy (Pd)	Not Available	IVes	Graded bunding
Himalapura	54	0.99	BDLbC3	LMU-3	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Cotton+Redgram (Ct+Rg)	Not Available	IVes	Graded bunding
Himalapura	55	0.62	BDLbC3	LMU-3	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Paddy (Pd)	Not Available	IVes	Graded bunding
Himalapura	56	1.57	BDLbC3	LMU-3	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Paddy (Pd)	Not Available	IVes	Graded bunding
Himalapura	57	0.57	BDLbC3	LMU-3	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Not Available (NA)	Not Available	IVes	Graded bunding
Himalapura	58	0.74	BDLbC3	LMU-3	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Redgram (Rg)	Not Available	IVes	Graded bunding
Himalapura	59	1.47	BDLbC3	LMU-3	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Not Available (NA)	Not Available	IVes	Graded bunding
Himalapura	60	0.97	BDLbC3	LMU-3	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Not Available (NA)	Not Available	IVes	Graded bunding
Himalapura	61	0.75	BDLbC3	LMU-3	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Paddy (Pd)	Not Available	IVes	Graded bunding
Himalapura	62	0.58	BDLbC3	LMU-3	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Paddy (Pd)	Not Available	IVes	Graded bunding
Himalapura	63	0.72	BDLbC3	LMU-3	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Paddy (Pd)	Not Available	IVes	Graded bunding
Himalapura	64	1.1	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Not Available (NA)	Not Available	Forest	Forest
Himalapura	65	2.45	BDLbC3	LMU-3	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Paddy (Pd)	Not Available	IVes	Graded bunding
Himalapura	66	1.34	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Not Available (NA)	Not Available	Forest	Forest
Himalapura	67	3.22	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Rock outcrops	Not Available	Forest	Forest
Himalapura	68	0.83	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Rock outcrops	Not Available	Forest	Forest
Himalapura	69	1.22	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Rock outcrops	Not Available	Forest	Forest
Himalapura	70	1.16	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Rock outcrops	Not Available	Forest	Forest
Himalapura	71	1.56	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Rock outcrops	Not Available	Forest	Forest
Himalapura	73	1.05	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Not Available (NA)	Not Available	Forest	Forest
Himalapura	75	1.91	BDPiB3	LMU-3	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Not Available (NA)	Not Available	IVes	Trench cum bunding

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Himalapura	76	1.35	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Redgram (Rg)	Not Available	Forest	Forest
Himalapura	77	1.41	BDPiB3	LMU-3	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Redgram (Rg)	Not Available	IVes	Trench cum bunding
Himalapura	78	0.47	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Paddy (Pd)	Not Available	Forest	Forest
Himalapura	79	0.17	BDLbc3	LMU-3	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Not Available (NA)	Not Available	IVes	Graded bunding
Himalapura	80	0.15	BDLbc3	LMU-3	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Paddy (Pd)	Not Available	IVes	Graded bunding
Himalapura	81	0.15	BDLbc3	LMU-3	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Paddy (Pd)	Not Available	IVes	Graded bunding
Himalapura	82	0.27	BDLbc3	LMU-3	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Paddy (Pd)	Not Available	IVes	Graded bunding
Himalapura	83	0.22	BDLbc3	LMU-3	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Paddy (Pd)	Not Available	IVes	Graded bunding
Himalapura	84	0.11	BDLbc3	LMU-3	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Paddy (Pd)	Not Available	IVes	Graded bunding
Himalapura	85	0.11	BDLbc3	LMU-3	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Paddy (Pd)	Not Available	IVes	Graded bunding
Himalapura	86	0.19	BDLbc3	LMU-3	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Paddy (Pd)	Not Available	IVes	Graded bunding
Himalapura	87	0.29	BDLbc3	LMU-3	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Paddy (Pd)	Not Available	IVes	Graded bunding
Himalapura	88	0.26	BDLbc3	LMU-3	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Paddy (Pd)	Not Available	IVes	Graded bunding
Himalapura	89	2.41	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Rock outcrops	Not Available	Forest	Forest
Himalapura	90	0.35	BDLbc3	LMU-3	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Paddy (Pd)	Not Available	IVes	Graded bunding
Himalapura	91	0.19	BDLbc3	LMU-3	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Paddy (Pd)	Not Available	IVes	Graded bunding
Himalapura	92	0.23	BDLbc3	LMU-3	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Not Available (NA)	Not Available	IVes	Graded bunding
Himalapura	93	0.13	BDLbc3	LMU-3	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Paddy (Pd)	Not Available	IVes	Graded bunding
Himalapura	94	0.81	BDLbc3	LMU-3	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Paddy (Pd)	Not Available	IVes	Graded bunding
Himalapura	95	0.22	BDLbc3	LMU-3	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Paddy (Pd)	Not Available	IVes	Graded bunding
Himalapura	96	0.13	BDLbc3	LMU-3	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Paddy (Pd)	Not Available	IVes	Graded bunding
Himalapura	97	0.27	BDLbc3	LMU-3	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Paddy (Pd)	Not Available	IVes	Graded bunding
Himalapura	98	0.21	BDLbc3	LMU-3	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Paddy (Pd)	Not Available	IVes	Graded bunding
Himalapura	99	0.3	BDLbc3	LMU-3	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Paddy (Pd)	Not Available	IVes	Graded bunding

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Himalapura	100	0.49	BDLbc3	LMU-3	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Paddy (Pd)	Not Available	Ives	Graded bunding
Himalapura	101	0.42	BDLbc3	LMU-3	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Paddy (Pd)	Not Available	Ives	Graded bunding
Himalapura	102	2.99	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Not Available (NA)	Not Available	Forest	Forest
Himalapura	103	0.14	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Not Available (NA)	Not Available	Forest	Forest
Himalapura	104	0.29	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Not Available (NA)	Not Available	Forest	Forest
Himalapura	105	0.33	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Not Available (NA)	Not Available	Forest	Forest
Himalapura	106	0.29	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Not Available (NA)	Not Available	Forest	Forest
Himalapura	107	0.14	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Not Available (NA)	Not Available	Forest	Forest
Himalapura	108	0.13	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Not Available (NA)	Not Available	Forest	Forest
Himalapura	109	0.35	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Not Available (NA)	Not Available	Forest	Forest
Himalapura	110	0.15	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Not Available (NA)	Not Available	Forest	Forest
Himalapura	111	0.26	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Not Available (NA)	Not Available	Forest	Forest
Himalapura	112	0.1	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Not Available (NA)	Not Available	Forest	Forest
Himalapura	113	0.14	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Not Available (NA)	Not Available	Forest	Forest
Himalapura	114	0.16	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Waterbody	Not Available	Forest	Forest
Himalapura	115	0.25	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Waterbody	Not Available	Forest	Forest
Himalapura	116	0.2	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Waterbody	Not Available	Forest	Forest
Himalapura	117	0.05	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Not Available (NA)	Not Available	Forest	Forest
Himalapura	118	0.16	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Not Available (NA)	Not Available	Forest	Forest
Himalapura	119	0.2	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Not Available (NA)	Not Available	Forest	Forest
Himalapura	120	0.06	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Not Available (NA)	Not Available	Forest	Forest
Himalapura	121	2.45	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Waterbody	Not Available	Forest	Forest
Himalapura	122	0.18	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Not Available (NA)	Not Available	Forest	Forest
Himalapura	123	0.08	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Not Available (NA)	Not Available	Forest	Forest

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Himalapura	124	0.13	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Not Available (NA)	Not Available	Forest	Forest
Himalapura	125	0.12	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Not Available (NA)	Not Available	Forest	Forest
Himalapura	126	1.39	BDPiB3	LMU-3	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Not Available (NA)	Not Available	IVes	Trench cum bunding
Himalapura	127	1.56	BDPiB3	LMU-3	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Not Available (NA)	Not Available	IVes	Trench cum bunding
Himalapura	132	0.49	BDPiB3	LMU-3	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Not Available (NA)	Not Available	IVes	Trench cum bunding
Himalapura	133	0.24	BDPiB3	LMU-3	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Not Available (NA)	Not Available	IVes	Trench cum bunding
Himalapura	134	0.12	BDPiB3	LMU-3	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Not Available (NA)	Not Available	IVes	Trench cum bunding
Himalapura	135	0.01	BDPiB3	LMU-3	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Redgram (Rg)	Not Available	IVes	Trench cum bunding
Himalapura	136	1.02	BDPiB3	LMU-3	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Not Available (NA)	Not Available	IVes	Trench cum bunding
Himalapura	137	0.24	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Not Available (NA)	Not Available	Forest	Forest
Himalapura	138	0.14	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Not Available (NA)	Not Available	Forest	Forest
Himalapura	139	0.22	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Not Available (NA)	Not Available	Forest	Forest
Himalapura	140	0.88	BDPiB3	LMU-3	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Not Available (NA)	Not Available	IVes	Trench cum bunding
Himalapura	141	0.12	BDPiB3	LMU-3	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Not Available (NA)	Not Available	IVes	Trench cum bunding
Himalapura	148	0.03	BDPiB3	LMU-3	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Redgram (Rg)	Not Available	IVes	Trench cum bunding
Himalapura	149	1.32	BDPiB3	LMU-3	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Redgram (Rg)	Not Available	IVes	Trench cum bunding
Himalapura	150	0.79	BDPiB3	LMU-3	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Waterbody	Not Available	IVes	Trench cum bunding
Himalapura	151	2.99	BDPiB3	LMU-3	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Redgram (Rg)	Not Available	IVes	Trench cum bunding
Himalapura	152	0.47	BDPiB3	LMU-3	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Redgram (Rg)	Not Available	IVes	Trench cum bunding
Himalapura	153	0.31	BDPiB3	LMU-3	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Not Available (NA)	Not Available	IVes	Trench cum bunding
Himalapura	154	0.06	BDPiB3	LMU-3	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Not Available (NA)	Not Available	IVes	Trench cum bunding
Himalapura	155	1.79	BDPiB3	LMU-3	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Groundnut (Gn)	Not Available	IVes	Trench cum bunding
Himalapura	156	0.31	BDPiB3	LMU-3	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Redgram (Rg)	Not Available	IVes	Trench cum bunding
Himalapura	157	0.01	BDPiB3	LMU-3	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Not Available (NA)	Not Available	IVes	Trench cum bunding

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Himalapura	180	0.04	BDPiB3	LMU-3	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Paddy (Pd)	Not Available	IVes	Trench cum bunding
Himalapura	181	0.52	BDPiB3	LMU-3	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Not Available (NA)	Not Available	IVes	Trench cum bunding
Himalapura	182	0.27	BDPiB3	LMU-3	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Redgram (Rg)	Not Available	IVes	Trench cum bunding
Himalapura	183	0.15	BDPiB3	LMU-3	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Redgram+Habitation (Rg+Hb)	Not Available	IVes	Trench cum bunding
Himalapura	221	0.12	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Redgram+Rock outcrops (Rg+Rc)	Not Available	Forest	Forest
Himalapura	226	1.21	BDLbC3	LMU-3	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Redgram (Rg)	Not Available	IVes	Graded bunding
Himalapura	227	0.68	BDPiB3	LMU-3	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Redgram (Rg)	Not Available	IVes	Trench cum bunding
Himalapura	228	1.68	BDPiB3	LMU-3	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Redgram (Rg)	Not Available	IVes	Trench cum bunding
Himalapura	229	0.37	BDPiB3	LMU-3	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Redgram (Rg)	Not Available	IVes	Trench cum bunding
Himalapura	230	0.33	BDPiB3	LMU-3	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Redgram (Rg)	Not Available	IVes	Trench cum bunding
Himalapura	231	2.38	BDLbC3	LMU-3	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Redgram (Rg)	Not Available	IVes	Graded bunding
Himalapura	232	1.48	BDLbC3	LMU-3	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Very sloping (3-5%)	Severe	Redgram+Rock outcrops (Rg+Rc)	Not Available	IVes	Graded bunding
Himalapura	233	0.11	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Not Available (NA)	Not Available	Forest	Forest
Himalapura	234	0.16	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Not Available (NA)	Not Available	Forest	Forest
Himalapura	235	0.8	BDPiB3	LMU-3	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Redgram+Rock outcrops (Rg+Rc)	Not Available	IVes	Trench cum bunding
Himalapura	236	0.43	BDPiB3	LMU-3	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Redgram+Rock outcrops (Rg+Rc)	Not Available	IVes	Trench cum bunding
Himalapura	237	0.26	BDPiB3	LMU-3	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Redgram+Rock outcrops (Rg+Rc)	Not Available	IVes	Trench cum bunding
Himalapura	238	0.47	BDPiB3	LMU-3	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Redgram+Rock outcrops (Rg+Rc)	Not Available	IVes	Trench cum bunding
Himalapura	239	1.27	BDPiB3	LMU-3	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Redgram (Rg)	Not Available	IVes	Trench cum bunding
Himalapura	240	1.27	BDPiB3	LMU-3	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Jowar+Paddy+Ground nut (Jw+Pd+Gn)	Not Available	IVes	Trench cum bunding
Himalapura	241	1.69	BDPiB3	LMU-3	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Jowar+Paddy+Ground nut (Jw+Pd+Gn)	Not Available	IVes	Trench cum bunding
Himalapura	242	3.64	BDPiB3	LMU-3	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Not Available (NA)	Not Available	IVes	Trench cum bunding
Himalapura	243	2.17	BDPiB3	LMU-3	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Not Available (NA)	Not Available	IVes	Trench cum bunding
Himalapura	244	0.3	BDPiB3	LMU-3	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Waterbody	Not Available	IVes	Trench cum bunding



Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Himalapura	245	0.47	BDPiB3	LMU-3	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Waterbody	Not Available	Ives	Trench cum bunding
Himalapura	246	0.12	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Waterbody	Not Available	Forest	Forest
Himalapura	247	0.18	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Waterbody	Not Available	Forest	Forest
Himalapura	248	0.67	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Not Available (NA)	Not Available	Forest	Forest
Himalapura	249	2.44	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Rock outcrops	Not Available	Forest	Forest
Himalapura	250	1.62	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Rock outcrops	Not Available	Forest	Forest
Himalapura	251	1.38	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Rock outcrops	Not Available	Forest	Forest
Himalapura	252	1.46	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Rock outcrops	Not Available	Forest	Forest
Himalapura	253	1.02	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Rock outcrops	Not Available	Forest	Forest
Himalapura	254	2.88	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Rock outcrops	Not Available	Forest	Forest
Himalapura	255	2.57	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Rock outcrops	Not Available	Forest	Forest
Himalapura	256	0.26	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Rock outcrops	Not Available	Forest	Forest
Himalapura	257	0.45	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Rock outcrops	Not Available	Forest	Forest
Himalapura	258	0.22	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Rock outcrops	Not Available	Forest	Forest
Himalapura	259	1.2	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Rock outcrops	Not Available	Forest	Forest
Himalapura	260	0.27	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Rock outcrops	Not Available	Forest	Forest
Himalapura	261	0.15	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Rock outcrops	Not Available	Forest	Forest
Himalapura	262	35.65	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Rock outcrops	Not Available	Forest	Forest
Himalapura	263	4.12	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Rock outcrops	Not Available	Forest	Forest
Himalapura	264	2	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Rock outcrops	Not Available	Forest	Forest
Himalapura	265	1.17	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Rock outcrops	Not Available	Forest	Forest
Himalapura	274	2.4	NGPmB2	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton+Redgram (Ct+Rg)	Not Available	Iles	Graded bunding
Himalapura	275	1.09	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Not Available (NA)	Not Available	Forest	Forest
Himalapura	276	0.56	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Not Available (NA)	Not Available	Forest	Forest

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Himalapura	277	1.11	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Not Available (NA)	Not Available	Forest	Forest
Himalapura	278	4.85	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Rock outcrops	Not Available	Forest	Forest
Himalapura	279	1.1	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Rock outcrops	Not Available	Forest	Forest
Himalapura	280	3.64	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Rock outcrops	Not Available	Forest	Forest
Himalapura	281	1.83	NGPmB2	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Rock outcrops	Not Available	Iles	Graded bunding
Himalapura	282	2.01	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Rock outcrops	Not Available	Forest	Forest
Himalapura	283	4.63	NGPmB2	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iles	Graded bunding
Himalapura	284	1.52	NGPmB2	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Rock outcrops	Not Available	Iles	Graded bunding
Himalapura	285	2.57	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Not Available (NA)	Not Available	Forest	Forest
Himalapura	286	4.4	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Scrub land (SI)	Not Available	Forest	Forest
Himalapura	287	1.32	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Rock outcrops	Not Available	Forest	Forest
Himalapura	288	2.53	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Rock outcrops	Not Available	Forest	Forest
Himalapura	289	0.35	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Not Available (NA)	Not Available	Forest	Forest
Himalapura	290	1.76	BDLbc3	LMU-3	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Not Available (NA)	Not Available	IVes	Graded bunding
Himalapura	291	10.32	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Rock outcrops	Not Available	Forest	Forest
Himalapura	292	0.18	BDLbc3	LMU-3	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Waterbody	Not Available	IVes	Graded bunding
Himalapura	293	0.21	BDLbc3	LMU-3	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Waterbody	Not Available	IVes	Graded bunding
Himalapura	294	0.17	BDLbc3	LMU-3	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Waterbody	Not Available	IVes	Graded bunding
Himalapura	295	0.16	BDLbc3	LMU-3	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Not Available (NA)	Not Available	IVes	Graded bunding
Himalapura	296	0.2	BDLbc3	LMU-3	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Waterbody	Not Available	IVes	Graded bunding
Himalapura	297	0.38	BDLbc3	LMU-3	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Waterbody	Not Available	IVes	Graded bunding
Himalapura	298	2.18	BDLbc3	LMU-3	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Redgram (Rg)	Not Available	IVes	Graded bunding
Himalapura	299	1.28	BDLbc3	LMU-3	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Not Available (NA)	Not Available	IVes	Graded bunding
Himalapura	300	0.67	BDLbc3	LMU-3	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Not Available (NA)	Not Available	IVes	Graded bunding

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Himalapura	301	0.85	BDLbC3	LMU-3	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Waterbody	Not Available	IVes	Graded bunding
Himalapura	302	0.79	BDLbC3	LMU-3	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Not Available (NA)	Not Available	IVes	Graded bunding
Himalapura	303	2.25	BDLbC3	LMU-3	Shallow (25-50 cm)	Loamy sand	Non gravelly (<15%)	Very low (<50 mm/m)	Gently sloping (3-5%)	Severe	Not Available (NA)	Not Available	IVes	Graded bunding
Himalapura	304	0.96	NGPmB2	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iles	Graded bunding
Himalapura	305	2.9	NGPmB2	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Himalapura	306	1.47	NGPmB2	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Himalapura	307	5.66	NGPmB2	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton+Redgram+Scrub land (Ct+Rg+Sl)	Not Available	Iles	Graded bunding
Himalapura	308	4.09	NGPmB2	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Himalapura	309	0.94	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Not Available (NA)	Not Available	Forest	Forest
Himalapura	310	0.59	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Waterbody	Not Available	Forest	Forest
Himalapura	311	3.02	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Not Available (NA)	Not Available	Forest	Forest
Himalapura	312	0.71	NGPmB2	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iles	Graded bunding
Himalapura	313	1.01	NGPmB2	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iles	Graded bunding
Himalapura	314	2.49	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Not Available (NA)	Not Available	Forest	Forest
Himalapura	315	0.34	NGPmB2	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iles	Graded bunding
Himalapura	316	1.2	NGPmB2	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iles	Graded bunding
Himalapura	317	1.7	NGPmB2	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Paddy (Rg+Pd)	Not Available	Iles	Graded bunding
Himalapura	318	0.76	NGPmB2	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Himalapura	319	0.92	NGPmB2	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Himalapura	320	0.42	NGPmB2	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iles	Graded bunding
Himalapura	321	1.27	BMNmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton+Habitation (Ct+Hb)	Not Available	Iles	Graded bunding
Himalapura	322	1.8	BMNmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar+Habitation (Jw+Hb)	Not Available	Iles	Graded bunding
Himalapura	323	0.32	BMNmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Himalapura	324	0.45	BMNmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iles	Graded bunding

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Himalapura	325	2.47	NGPmB2	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Himalapura	326	0.37	BMNmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton+Habitation (Ct+Hb)	Not Available	Iles	Graded bunding
Himalapura	327	0.37	BMNmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Iles	Graded bunding
Himalapura	328	0.9	BMNmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton+Habitation (Ct+Hb)	Not Available	Iles	Graded bunding
Himalapura	329	1.02	BDPiB3	LMU-3	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Jowar (Jw)	Not Available	IVes	Trench cum bunding
Himalapura	330	1.55	BMNmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	Iles	Graded bunding
Himalapura	331	0.66	BDPiB3	LMU-3	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Jowar (Jw)	Not Available	IVes	Trench cum bunding
Himalapura	332	1.08	BMNmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar+Cotton (Jw+Ct)	Not Available	Iles	Graded bunding
Himalapura	333	1.5	BMNmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Himalapura	334	3.06	BMNmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton+Redgram (Ct+Rg)	Not Available	Iles	Graded bunding
Himalapura	335	1.13	BMNmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Himalapura	336	1.29	BMNmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	Iles	Graded bunding
Himalapura	337	1.75	BMNmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Himalapura	338	1.29	BMNmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	Iles	Graded bunding
Himalapura	339	1.35	BMNmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	Iles	Graded bunding
Himalapura	340	3.38	NGPmB2	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Iles	Graded bunding
Himalapura	341	3.4	NGPmB2	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iles	Graded bunding
Himalapura	342	3.49	NGPmB2	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton+Redgram+Scrub land (Ct+Rg+Sl)	Not Available	Iles	Graded bunding
Himalapura	343	4.18	NGPmB2	LMU-1	Deep (100-150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton+Redgram (Ct+Rg)	Not Available	Iles	Graded bunding
Himalapura	344	1.32	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Redgram (Rg)	Not Available	Forest	Forest
Himalapura	345	3.54	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Redgram (Rg)	Not Available	Forest	Forest
Himalapura	346	2.1	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Not Available (NA)	Not Available	Forest	Forest
Himalapura	347	2.09	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Not Available (NA)	Not Available	Forest	Forest
Himalapura	348	2.26	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Rock outcrops	Not Available	Forest	Forest

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Himalapura	349	2.45	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Redgram+Rock outcrops+Jowar (Rg+Rc+Jw)	Not Available	Forest	Forest
Himalapura	350	2.69	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Not Available (NA)	Not Available	Forest	Forest
Himalapura	351	1.63	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Redgram (Rg)	Not Available	Forest	Forest
Himalapura	360	0.01	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Not Available (NA)	Not Available	Forest	Forest
Himalapura	371	0.19	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Not Available (NA)	Not Available	Forest	Forest
Himalapura	372	0.91	BMNmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar+Cotton (Jw+Ct)	Not Available	Iles	Graded bunding
Himalapura	373	2.7	BMNmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar+Cotton (Jw+Ct)	Not Available	Iles	Graded bunding
Himalapura	385	0	BMNmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Iles	Graded bunding
Himalapura	386	0.82	BMNmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Himalapura	387	2.46	BMNmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar+Redgram (Jw+Rg)	Not Available	Iles	Graded bunding
Himalapura	388	1.19	BMNmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Iles	Graded bunding
Himalapura	389	1.07	BMNmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	Iles	Graded bunding
Himalapura	390	1.98	BMNmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar+Redgram (Jw+Rg)	Not Available	Iles	Graded bunding
Himalapura	391	2.09	BMNmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar+Cotton (Jw+Ct)	Not Available	Iles	Graded bunding
Himalapura	392	1.67	BMNmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton+Redgram (Ct+Rg)	Not Available	Iles	Graded bunding
Himalapura	393	1.16	BMNmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Himalapura	394	3.24	BDPiB3	LMU-3	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Jowar+Cotton (Jw+Ct)	Not Available	IVes	Trench cum bunding
Himalapura	395	4.13	BMNmB2	LMU-1	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar+Redgram (Jw+Rg)	Not Available	Iles	Graded bunding
Himalapura	396	0.58	BDPiB3	LMU-3	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Cotton (Ct)	Not Available	IVes	Trench cum bunding
Himalapura	397	1.74	BDPiB3	LMU-3	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Cotton (Ct)	Not Available	IVes	Trench cum bunding
Himalapura	398	2.39	BDPiB3	LMU-3	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Cotton (Ct)	Not Available	IVes	Trench cum bunding
Himalapura	399	2.54	BDPiB3	LMU-3	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Jowar+Cotton (Jw+Ct)	Not Available	IVes	Trench cum bunding
Himalapura	400	1.26	BDPiB3	LMU-3	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Jowar+Cotton (Jw+Ct)	Not Available	IVes	Trench cum bunding
Himalapura	401	0.25	BDPiB3	LMU-3	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Redgram (Rg)	Not Available	IVes	Trench cum bunding

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Himalapura	402	0.72	BDPiB3	LMU-3	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Redgram (Rg)	Not Available	IVes	Trench cum bunding
Himalapura	403	1.33	BDPiB3	LMU-3	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Habitation	Not Available	IVes	Trench cum bunding
Himalapura	404	0.12	Habitation	Others	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others
Himalapura	405	0.62	Habitation	Others	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others
Himalapura	406	0.44	BDPiB3	LMU-3	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Habitation	Not Available	IVes	Trench cum bunding
Himalapura	407	0.81	Habitation	Others	Others	Others	Others	Others	Others	Others	Not Available (NA)	Not Available	Others	Others
Himalapura	408	1.31	Habitation	Others	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others
Himalapura	409	1.14	BDPiB3	LMU-3	Very shallow (<25 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Severe	Jowar+Habitation (Jw+Hb)	Not Available	IVes	Trench cum bunding
Himalapura	410	0.37	Habitation	Others	Others	Others	Others	Others	Others	Others	Not Available (NA)	Not Available	Others	Others













Village	Survey No	Soil Reaction	Salinity	Organic Carbon	Available Phosphorus	Available Potassium	Available Sulphur	Available Boron	Available Iron	Available Manganese	Available Copper	Available Zinc
Himalapura	104	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest
Himalapura	105	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest
Himalapura	106	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest
Himalapura	107	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest
Himalapura	108	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest
Himalapura	109	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest
Himalapura	110	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest
Himalapura	111	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest
Himalapura	112	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest
Himalapura	113	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest
Himalapura	114	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest
Himalapura	115	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest
Himalapura	116	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest
Himalapura	117	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest
Himalapura	118	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest
Himalapura	119	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest
Himalapura	120	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest
Himalapura	121	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest
Himalapura	122	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest
Himalapura	123	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest
Himalapura	124	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest
Himalapura	125	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest
Himalapura	126	Neutral (pH 6.5 - 7.3)	Non saline (<2 dsm)	High (> 0.75 %)	Low (< 23 kg/ha)	Low (<145 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)
Himalapura	127	Neutral (pH 6.5 - 7.3)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 - 57 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)
Himalapura	132	Neutral (pH 6.5 - 7.3)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 - 57 kg/ha)	Low (<145 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)
Himalapura	133	Neutral (pH 6.5 - 7.3)	Non saline (<2 dsm)	High (> 0.75 %)	Low (< 23 kg/ha)	Low (<145 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)
Himalapura	134	Neutral (pH 6.5 - 7.3)	Non saline (<2 dsm)	High (> 0.75 %)	Low (< 23 kg/ha)	Low (<145 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)
Himalapura	135	Neutral (pH 6.5 - 7.3)	Non saline (<2 dsm)	High (> 0.75 %)	Low (< 23 kg/ha)	Low (<145 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)
Himalapura	136	Neutral (pH 6.5 - 7.3)	Non saline (<2 dsm)	High (> 0.75 %)	Low (< 23 kg/ha)	Low (<145 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)





Village	Survey No	Soil Reaction	Salinity	Organic Carbon	Available Phosphorus	Available Potassium	Available Sulphur	Available Boron	Available Iron	Available Manganese	Available Copper	Available Zinc
Himalapura	258	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest
Himalapura	259	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest
Himalapura	260	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest
Himalapura	261	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest
Himalapura	262	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest
Himalapura	263	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest
Himalapura	264	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest
Himalapura	265	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest
Himalapura	274	Neutral (pH 6.5 - 7.3)	Non saline (<2 dsm)	High (> 0.75 %)	Low (< 23 kg/ha)	Medium (145 - 337 kg/ha)	Medium (10 - 20 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Himalapura	275	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest
Himalapura	276	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest
Himalapura	277	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest
Himalapura	278	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest
Himalapura	279	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest
Himalapura	280	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest
Himalapura	281	Neutral (pH 6.5 - 7.3)	Non saline (<2 dsm)	High (> 0.75 %)	Low (< 23 kg/ha)	Medium (145 - 337 kg/ha)	Medium (10 - 20 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Himalapura	282	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest
Himalapura	283	Neutral (pH 6.5 - 7.3)	Non saline (<2 dsm)	High (> 0.75 %)	Low (< 23 kg/ha)	Medium (145 - 337 kg/ha)	Medium (10 - 20 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Himalapura	284	Neutral (pH 6.5 - 7.3)	Non saline (<2 dsm)	High (> 0.75 %)	Low (< 23 kg/ha)	Medium (145 - 337 kg/ha)	Medium (10 - 20 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Himalapura	285	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest
Himalapura	286	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest
Himalapura	287	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest
Himalapura	288	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest
Himalapura	289	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest
Himalapura	290	Neutral (pH 6.5 - 7.3)	Non saline (<2 dsm)	High (> 0.75 %)	Low (< 23 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)
Himalapura	291	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest
Himalapura	292	Neutral (pH 6.5 - 7.3)	Non saline (<2 dsm)	High (> 0.75 %)	Low (< 23 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)
Himalapura	293	Neutral (pH 6.5 - 7.3)	Non saline (<2 dsm)	High (> 0.75 %)	Low (< 23 kg/ha)	Medium (145 - 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Sufficient (> 0.6 ppm)









Village	Survey No	Soil Reaction	Salinity	Organic Carbon	Available Phosphorus	Available Potassium	Available Sulphur	Available Boron	Available Iron	Available Manganese	Available Copper	Available Zinc
Himalapura	398	Neutral (pH 6.5 - 7.3)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 - 57 kg/ha)	High (> 337 kg/ha)	Low (<10 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Himalapura	399	Neutral (pH 6.5 - 7.3)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 - 57 kg/ha)	High (> 337 kg/ha)	Medium (10 - 20 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Himalapura	400	Neutral (pH 6.5 - 7.3)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 - 57 kg/ha)	High (> 337 kg/ha)	Medium (10 - 20 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Himalapura	401	Neutral (pH 6.5 - 7.3)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 - 57 kg/ha)	High (> 337 kg/ha)	Medium (10 - 20 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Himalapura	402	Neutral (pH 6.5 - 7.3)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 - 57 kg/ha)	High (> 337 kg/ha)	Medium (10 - 20 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Himalapura	403	Neutral (pH 6.5 - 7.3)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 - 57 kg/ha)	High (> 337 kg/ha)	Medium (10 - 20 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Himalapura	404	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Himalapura	405	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Himalapura	406	Neutral (pH 6.5 - 7.3)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 - 57 kg/ha)	High (> 337 kg/ha)	Medium (10 - 20 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Himalapura	407	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Himalapura	408	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Himalapura	409	Neutral (pH 6.5 - 7.3)	Non saline (<2 dsm)	High (> 0.75 %)	Medium (23 - 57 kg/ha)	High (> 337 kg/ha)	Medium (10 - 20 ppm)	Low (< 0.5 ppm)	Sufficient (>4.5 ppm)	Sufficient (> 1.0 ppm)	Sufficient (> 0.2 ppm)	Deficient (< 0.6 ppm)
Himalapura	410	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others



**Appendix III**  
**Himalapur (6A2a) Microwatershed**  
**Soil Suitability Information**

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brijjal	Bhendi	Drumstick	Mulberry	
Boodhura	138	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	
Boodhura	139	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	
Boodhura	140	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	
Chapetla	18	N1r	S2t	S3r	S2r	S3r	S2r	N1r	S3r	S2r	S3r	S3r	S2r	S3r	S2r	N1rt	S3r	S3r	S3t	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Chapetla	85	N1r	S2t	S3r	S2r	S3r	S2r	N1r	S3r	S2r	S3r	S3r	S2r	S3r	S2r	N1rt	S3r	S3r	S3t	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Chapetla	86	N1r	S2t	S3r	S2r	S3r	S2r	N1r	S3r	S2r	S3r	S3r	S2r	S3r	S2r	N1rt	S3r	S3r	S3t	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Chapetla	87	N1r	S2t	S3r	S2r	S3r	S2r	N1r	S3r	S2r	S3r	S3r	S2r	S3r	S2r	N1rt	S3r	S3r	S3t	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Chapetla	89	N1r	S2t	S3r	S2r	S3r	S2r	N1r	S3r	S2r	S3r	S3r	S2r	S3r	S2r	N1rt	S3r	S3r	S3t	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Chapetla	90	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Chapetla	91	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Chapetla	98	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	
Chapetla	99	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	
Chapetla	100	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Chapetla	101	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Chapetla	102	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Chapetla	103	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Chapetla	104	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Chapetla	105	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Chapetla	106	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Chapetla	107	S3t	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw	
Chapetla	108	S3t	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw	
Chapetla	109	S3t	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw	
Chapetla	149	N1r	S2t	S3r	S2r	S3r	S2r	N1r	S3r	S2r	S3r	S3r	S2r	S3r	S2r	N1rt	S3r	S3r	S3t	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Chapetla	150	N1r	S2t	S3r	S2r	S3r	S2r	N1r	S3r	S2r	S3r	S3r	S2r	S3r	S2r	N1rt	S3r	S3r	S3t	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Chapetla	151	N1r	S2t	S3r	S2r	S3r	S2r	N1r	S3r	S2r	S3r	S3r	S2r	S3r	S2r	N1rt	S3r	S3r	S3t	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry
Chapetla	152	N1r	S2t	S3r	S2r	S3r	S2r	N1r	S3r	S2r	S3r	S3r	S2r	S3r	S2r	N1rt	S3r	S3r	S3t	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r
Chapetla	153	N1r	S2t	S3r	S2r	S3r	S2r	N1r	S3r	S2r	S3r	S3r	S2r	S3r	S2r	N1rt	S3r	S3r	S3t	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r
Himalapura	1	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs
Himalapura	2	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt
Himalapura	3	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt
Himalapura	4	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt
Himalapura	5	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt
Himalapura	6	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt
Himalapura	7	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt
Himalapura	11	S3t	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw
Himalapura	12	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt
Himalapura	13	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt
Himalapura	14	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt
Himalapura	15	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs	Othe rs
Himalapura	16	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt
Himalapura	17	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt
Himalapura	18	S3t	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw
Himalapura	19	S3t	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw
Himalapura	20	S3t	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw
Himalapura	21	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt
Himalapura	22	S3t	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw
Himalapura	23	S3t	S2t	S3t	S2w	S3t	S1	S2t	S1	S1	S2w	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S3tw	S3tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw
Himalapura	24	S3t	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw
Himalapura	25	S3t	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw
Himalapura	26	S3t	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw
Himalapura	27	S3t	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw
Himalapura	31	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry
Himalapura	32	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt
Himalapura	33	S3t	S2t	S3t	S2w	S3t	S1	S2t	S1	S1	S2w	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S3tw	S3tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw
Himalapura	34	S3t	S2t	S3t	S2w	S3t	S1	S2t	S1	S1	S2w	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S3tw	S3tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw
Himalapura	35	S3t	S2t	S3t	S2w	S3t	S1	S2t	S1	S1	S2w	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S3tw	S3tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw
Himalapura	36	S3t	S2t	S3t	S2w	S3t	S1	S2t	S1	S1	S2w	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S3tw	S3tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw
Himalapura	37	S3t	S2t	S3t	S2w	S3t	S1	S2t	S1	S1	S2w	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S3tw	S3tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw
Himalapura	38	S3t	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw
Himalapura	39	S3t	S2t	S3t	S2w	S3t	S1	S2t	S1	S1	S2w	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S3tw	S3tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw
Himalapura	40	S3t	S2t	S3t	S2w	S3t	S1	S2t	S1	S1	S2w	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S3tw	S3tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw
Himalapura	41	S3t	S2t	S3t	S2w	S3t	S1	S2t	S1	S1	S2w	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S3tw	S3tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw
Himalapura	42	S3t	S2t	S3t	S2w	S3t	S1	S2t	S1	S1	S2w	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S3tw	S3tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw
Himalapura	43	S3t	S2t	S3t	S2w	S3t	S1	S2t	S1	S1	S2w	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S3tw	S3tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw
Himalapura	44	S3t	S2t	S3t	S2w	S3t	S1	S2t	S1	S1	S2w	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S3tw	S3tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw
Himalapura	45	S3t	S2t	S3t	S2w	S3t	S1	S2t	S1	S1	S2w	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S3tw	S3tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw
Himalapura	46	S3t	S2t	S3t	S2w	S3t	S1	S2t	S1	S1	S2w	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S3tw	S3tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw
Himalapura	47	S3t	S2t	S3t	S2w	S3t	S1	S2t	S1	S1	S2w	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S3tw	S3tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw
Himalapura	48	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt
Himalapura	49	S3t	S2t	S3t	S2w	S3t	S1	S2t	S1	S1	S2w	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S3tw	S3tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw
Himalapura	50	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st	Fore st
Himalapura	51	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt
Himalapura	52	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt
Himalapura	53	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt
Himalapura	54	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt
Himalapura	55	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt
Himalapura	56	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt
Himalapura	57	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt
Himalapura	58	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt









Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry	
Himalapura	152	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	
Himalapura	153	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Himalapura	154	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Himalapura	155	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Himalapura	156	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Himalapura	157	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Himalapura	180	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Himalapura	181	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Himalapura	182	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Himalapura	183	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Himalapura	221	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest
Himalapura	226	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	
Himalapura	227	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Himalapura	228	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Himalapura	229	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Himalapura	230	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Himalapura	231	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	
Himalapura	232	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	
Himalapura	233	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest
Himalapura	234	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest
Himalapura	235	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Himalapura	236	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Himalapura	237	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Himalapura	238	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Himalapura	239	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Himalapura	240	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Himalapura	241	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r





Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry	
Himalapura	304	S3t	S2t	S3t	S2w	S3t	S1	S2t	S1	S1	S2w	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S3tw	S3tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw	
Himalapura	305	S3t	S2t	S3t	S2w	S3t	S1	S2t	S1	S1	S2w	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S3tw	S3tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw	
Himalapura	306	S3t	S2t	S3t	S2w	S3t	S1	S2t	S1	S1	S2w	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S3tw	S3tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw	
Himalapura	307	S3t	S2t	S3t	S2w	S3t	S1	S2t	S1	S1	S2w	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S3tw	S3tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw	
Himalapura	308	S3t	S2t	S3t	S2w	S3t	S1	S2t	S1	S1	S2w	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S3tw	S3tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw	
Himalapura	309	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest
Himalapura	310	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest
Himalapura	311	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest
Himalapura	312	S3t	S2t	S3t	S2w	S3t	S1	S2t	S1	S1	S2w	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S3tw	S3tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw	
Himalapura	313	S3t	S2t	S3t	S2w	S3t	S1	S2t	S1	S1	S2w	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S3tw	S3tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw	
Himalapura	314	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest	Forest
Himalapura	315	S3t	S2t	S3t	S2w	S3t	S1	S2t	S1	S1	S2w	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S3tw	S3tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw	
Himalapura	316	S3t	S2t	S3t	S2w	S3t	S1	S2t	S1	S1	S2w	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S3tw	S3tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw	
Himalapura	317	S3t	S2t	S3t	S2w	S3t	S1	S2t	S1	S1	S2w	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S3tw	S3tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw	
Himalapura	318	S3t	S2t	S3t	S2w	S3t	S1	S2t	S1	S1	S2w	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S3tw	S3tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw	
Himalapura	319	S3t	S2t	S3t	S2w	S3t	S1	S2t	S1	S1	S2w	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S3tw	S3tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw	
Himalapura	320	S3t	S2t	S3t	S2w	S3t	S1	S2t	S1	S1	S2w	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S3tw	S3tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw	
Himalapura	321	S3t	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw	
Himalapura	322	S3t	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw	
Himalapura	323	S3t	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw	
Himalapura	324	S3t	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw	
Himalapura	325	S3t	S2t	S3t	S2w	S3t	S1	S2t	S1	S1	S2w	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S3tw	S3tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw	
Himalapura	326	S3t	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw	
Himalapura	327	S3t	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw	
Himalapura	328	S3t	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw	
Himalapura	329	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Himalapura	330	S3t	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw	



Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry	
Himalapura	387	S3t	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw	
Himalapura	388	S3t	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw	
Himalapura	389	S3t	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw	
Himalapura	390	S3t	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw	
Himalapura	391	S3t	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw	
Himalapura	392	S3t	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw	
Himalapura	393	S3t	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw	
Himalapura	394	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Himalapura	395	S3t	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S3t	S2tw	S2tw	S2tw	S2tw	S2t	S2tw	S2t	S2t	S2tw	S3tw	
Himalapura	396	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Himalapura	397	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Himalapura	398	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Himalapura	399	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Himalapura	400	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Himalapura	401	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Himalapura	402	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Himalapura	403	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Himalapura	404	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Himalapura	405	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Himalapura	406	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Himalapura	407	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Himalapura	408	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Himalapura	409	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Himalapura	410	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others



# **PART-B**

**SOCIO-ECONOMIC STATUS OF FARM HOUSEHOLDS**



## CONTENTS

1.	Findings of the socio-economic survey	1-2
2.	Introduction	3
3	Methodology	5-6
4	Salient features of the survey	7-21
5	Summary	23-26



## LIST OF TABLES

1	Households sampled for socio economic survey	7
2	Population characteristics	7
3	Age wise classification of household members	7
4	Education level of household members	8
5	Occupation of household heads	8
6	Occupation of family members	8
7	Institutional participation of household members	9
8	Type of house owned by households	9
9	Durable assets owned by households	9
10	Average value of durable assets owned by households	9
11	Farm implements owned by households	10
12	Average value of farm implements	10
13	Livestock possession by households	10
14	Average labour availability	10
15	Adequacy of hired labour	11
16	Distribution of land (ha)	11
17	Average land value (Rs./ha)	11
18	Status of bore wells	11
19	Source of irrigation	11
20	Depth of water(Avg in meters)	12
21	Irrigated area (ha)	12
22	Cropping pattern	12
23	Cropping intensity	12
24.a	Cost of cultivation of Red gram	13
24.b	Cost of cultivation of Cotton	14
24.c	Cost of cultivation of Groundnut	15
24.d	Cost of cultivation of Bengal Gram	16
25	Adequacy of fodder	17
26	Annual gross income	17
27	Average annual expenditure	17
28	Horticultural species grown	17

29	Forest species grown	18
30	Marketing of the agricultural produce	18
31	Marketing channels used for sale of agricultural produce	18
32	Mode of transport of agricultural produce	19
33	Incidence of soil and water erosion problems	19
34	Interest shown towards soil testing	19
35	Usage pattern of fuel for domestic use	19
36	Source of drinking water	19
37	Source of light	20
38	Existence of sanitary toilet facility	20
39	Possession of public distribution system (PDS) card	20
40	Adequacy of food items	20
41	Inadequacy of food items	21
42	Farming constraints experienced	21

**FINDINGS OF THE SOCIO-ECONOMIC SURVEY**

- ❖ *The survey was conducted in Himalapur is located at North latitude 16<sup>o</sup> 52' 34.382" and 16<sup>o</sup> 50' 38.077" and East longitude 77<sup>o</sup> 20' 51.651" and 77<sup>o</sup> 19' 20.923" covering an area of about 498.01 ha coming under Himalapura and Chapetla Villages of Yadagiri taluk.*
- ❖ *Socio-economic analysis of Himalapur micro watersheds of Mokadampur sub-watershed, Yadgiri taluk & District indicated that, out of the total sample of 34 farmers were sampled in Himalapur micro-watershed among households surveyed 25 (73.53%) were marginal, 4 (11.76%) were small and 2 (5.88 %) were semi medium farmers. 3 landless farmers were also interviewed for the survey.*
- ❖ *The population characteristics of households indicated that, there were 83 (60.58%) men and 54 (39.42 %) were women. The average population of landless was 3.7, marginal farmers were 3.9, small farmers were 4.3 and semi medium farmers were 6.*
- ❖ *Majority of the respondents (42.34%) were in the age group of 16-35 years.*
- ❖ *Education level of the sample households indicated that, there were 49.64 per cent illiterates, 0.73 percent were functional literates, 47.45 per cent pre university education and 3.65 per cent attained graduation.*
- ❖ *About, 76.47 per cent of household heads practicing agriculture and 5.88 per cent of the household heads were engaged as agricultural labourers.*
- ❖ *Agriculture was the major occupation for 35.04 per cent of the household members.*
- ❖ *In the study area, 94.12 per cent of the households possess katcha house and 2.94 per cent possess pucca house.*
- ❖ *The durable assets owned by the households showed that, 100.00 per cent possess TV.*
- ❖ *Farm implements owned by the households indicated that, 8.82 per cent of the households possess plough.*
- ❖ *The average labour availability in the study area showed that, own labour men available in the micro watershed was 1.53, women available in the micro watershed was 1.38, hired labour (men) available was 8.91 and hired labour (women) available was 7.34.*
- ❖ *Out of the total land holding of the sample respondents 94.95 per cent (24.04 ha) of the area is under dry condition and the remaining 5.05 per cent area is irrigated land.*
- ❖ *There were 1.00 live bore wells among the sampled households.*
- ❖ *Bore well was the major source of irrigation for 2.94 per cent of the households.*

- ❖ *The major crops grown by sample farmers are Red gram, Cotton, Groundnut, Bengal gram and cropping intensity was recorded as 100.00 per cent.*
- ❖ *The per hectare cost of cultivation for Red gram, Cotton, Groundnut and Bengal gram was Rs.36642.31, 78378.30, 30333.70 and 116785.65 with benefit cost ratio of 1:3.50, 1: 1.10, 1: 3.30 and 1: 0.99 respectively.*
- ❖ *Further, 2.94 per cent of the households opined that dry fodder was adequate and 2.94 per cent of the households have opined that the green fodder was adequate.*
- ❖ *The average annual gross income of the farmers was Rs. 90000.00 in micro-watershed, of which Rs. 39794.12 comes from agriculture.*
- ❖ *Sampled households have grown 1 horticulture trees and 61 forestry trees together in the fields and back yards.*
- ❖ *Regarding marketing channels, 91.18 per cent of the households have sold agricultural produce to the local/village merchants.*
- ❖ *Further, 91.18 per cent of the households have used tractor for the transport of agriculture commodity.*
- ❖ *Majority of the farmers (91.18%) have experienced soil and water erosion problems in the watershed and 91.18 per cent of the households were interested towards soil testing.*
- ❖ *Fire was the major source of fuel for domestic use for 100.00 per cent of the households.*
- ❖ *Piped supply was the major source for drinking water for 100.00 per cent of the households.*
- ❖ *Electricity was the major source of light for 100.00 per cent of the households.*
- ❖ *In the study area, 100.00 per cent of the households possess toilet facility.*
- ❖ *Regarding possession of PDS card, 100.00 per cent of the households possessed BPL card.*
- ❖ *Households opined that, the requirement of cereals (100.00%), pulses (94.12%) are adequate for consumption.*
- ❖ *Farming constraints experienced by households in the micro watersheds were lower fertility status of the soil (91.18%) wild animal menace on farm field (8.82%), frequent incidence of pest and diseases (88.24%), inadequacy of irrigation water (2.94%), high cost of fertilizers and plant protection chemicals (88.24%), high rate of interest on credit (14.71%), low price for the agricultural commodities (76.47%), lack of marketing facilities in the area (79.41%), inadequate extension services (8.82%), lack of transport for safe transport of the agricultural produce to the market (64.71%).*



## **INTRODUCTION**

Soil and water are the two precious natural resources which are essential for crop production and existence of life on earth. Rainfed agriculture is under severe stress due to various constraints related to agriculture like uneven and erratic distribution of rainfall, indiscriminate use of fertilizers, chemicals and pesticides, adoption of improper land management practices, soil erosion, decline in soil fertility, decline in ground water resources leading to low crop productivity. The area under rainfed agriculture has to be managed effectively using the best available practices to enhance the production of food, fodder and fuel. This is possible if the land resources are characterized at each parcel of land through detailed land resource inventory using the best available techniques of remote sensing, GPS and GIS. The watershed development programs are aimed at the sustainable distribution of its resources and the process of creating and implementing plans, programs, and projects to sustain and enhance watershed functions that affect the plant, animal and human communities within a watershed boundary.

World Bank funded KWDP II, SUJALA III project was implemented in with Broad objective of demonstrating more effective watershed management through greater integration of programmes related to rain-fed agriculture, innovative and science based approaches and strengthen institutional capacities and If successful, it is expected that the systems and tools could be mainstreamed into the overall IWMP in the State of Karnataka and in time, throughout other IWMP operations in India. With this background the socio-economic survey has been carried out with following specific objectives:

1. To understand the demographic features of the households in the micro-watershed
2. To understand the extent of family labour available and additional employment opportunities available within the village.
3. To know the status of assets of households in the micro-watershed for suggesting possible improvements.
4. To study the cropping pattern, cropped area and productivity levels of different households in micro-watershed.
5. To determine the type and extent of livestock owned by different categories of HHs
6. Availability of fodder and level of livestock management.

### **Scope and importance of survey**

Survey helps in identification of different socio-economic and resource use-patterns of farmers at the Micro watershed. Household survey provides demographic features, labor force, and levels of education; land ownership and asset position (including livestock and other household assets) of surveyed households; and cropping patterns, input intensities, and average crop yields from farmers' fields. It also discusses crop utilization and the degree of commercialization of production in the areas; farmers' access to and utilization of credit from formal and informal sources; and the level of adoption and use of soil, water, and pest management technologies.



## METHODOLOGY

The description of the methods, components selected for the survey and procedures followed in conducting the baseline survey are furnished under the following heads.

### **1. Description of the study area**

Yadgir District is one of the 30 districts of Karnataka state in southern India. This district was carved out from the erstwhile Gulbarga district as the 30th district of Karnataka on 10 April 2010. Yadgir town is the administrative headquarters of the district. The district comprises of 3 taluks namely, Shahapur, Yadgiri and Shorapur (There are 16 hoblies, 117 Gram Panchayats, 4 Municipalities, 8 Towns/ Urban agglomeration and 487 inhabited & 32 un-inhabited villages The district occupies an area of 5,160.88 km<sup>2</sup>.

Yadgir district is the second smallest district in the state, area wise is very rich in cultural traditions. The vast stretch of fertile black soil of the district is known for bumper red gram and jowar crops. The district is a "Daal bowl" of the state. The district is also known for cluster of cement industries and a distinct stone popularly known as "Malakheda Stone". Two main rivers, Krishna and Bhima, and a few tributaries flow in this region. Krishna and Bhima Rivers drain the district. They constitute the two major river basins of the district. Kagna and Amarja are the two sub - basins of Bhima River, which occur within the geographical area of the district

According to the 2011 census Yadgir district has a population of 1, 172,985, roughly equal to the nation of Timor-Leste or the US state of Rhode Island. This gives it a ranking of 404th in India (out of a total of 640). The district has a population density of 224 inhabitants per square kilometre (580/sq mi). Its population growth rate over the decade 2001-2011 was 22.67%. Yadgir has a sex ratio of 984 females for every 1000 males, and a literacy rate of 52.36%.

### **2. Locale of the survey and description of the micro-watershed and**

The study was conducted in Himalapur micro-watershed (Mokadampur sub-watershed, Yadgiri taluk & District) is located at North latitude 16<sup>o</sup> 52' 34.382" and 16<sup>o</sup> 50' 38.077" and East longitude 77<sup>o</sup> 20' 51.651" and 77<sup>o</sup> 19' 20.923" covering an area of about 498.01 ha bounded by under Himalapura and Chapetla Villages.

### **3. Selection of the respondents for the study**

The micro-watershed is marked with 320 square meters grids. One farmer from every alternate grid in the micro-watershed was selected for the study and interviewed for socio-economic data. Totally 34 households were interviewed for the survey.

#### **4. The parameters considered for socio-economic survey of households**

Two forms of data were collected from the micro-watershed which includes primary data from the farm households and secondary data about the villages under the micro-watershed jurisdiction.

The following parameters were considered for the primary data collection about the socio-economic data of the households, (1) Demographic information, (2) Farm and durable assets owned by households, (3) Livestock possession, (4) Labour availability, (5) Level of migration in the village, Land holding, (7) Cropping pattern, (8) Source of irrigation, (9) Borrowing status, (10) Cost of cultivation of major crops, (11) Economics of subsidiary activities, (12) Fodder availability, (13) Family annual income from different sources, (14) Horticulture and forestry species grown, (15) Additional investment capacity, (16) Marketing practices, (17) Status of soil and water conservation structure, (18) Access to basic needs and (19) Constraints and suggestion.

The following parameters were considered for the secondary data regarding the villages under the micro-watershed jurisdiction, (1) Number of villages in each micro-watershed jurisdiction, (2) Village wise number of households, (3) Geographical area of the villages, (4) Cultivable area including rainfed and irrigated, (5) Number and type of house in each village, (6) Human and livestock population, (7) Facilities in the village such as roads, transport facility for conveyance, drinking water supply, street light and (8) Community based organizations in the villages.

#### **5. Development of interview schedule and data collection**

Taking into the consideration the objectives of the survey, an interview schedule was prepared after thorough consultation with the experts in the field of social sciences. A comprehensive interview schedule covering all the major parameters for measuring the socio-economic situation was developed.

#### **6. Tools used to analyze the data**

The statistical components such as frequency and percentage were used to analyze the data.

#### **Abbreviations used in the report**

LL=Landless

MF=Marginal Farmers

SF=Small farmers

SMF=Semi medium farmers

MDF=Medium farmers

LF=Large Farmers

## FINDINGS OF THE SURVEY

This chapter deals with systematic presentation of results of the survey. Keeping in view the objectives, the salient features of the survey are presented under the following headings.

**Households sampled for socio-economic survey:** The data on households sampled for socio economic survey in Himalapur Micro watershed is presented in Table 1 and it indicated that 34 farmers were sampled in Himalapur micro-watershed among households surveyed 25 (73.53%) were marginal, 4 (11.76%) were small and 2 (5.88 %) were semi medium farmers. 3 landless farmers were also interviewed for the survey.

**Table 1. Households sampled for socio economic survey in Himalapur micro-watershed**

Sl.No.	Particulars	LL (3)		MF (25)		SF (4)		SMF (2)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Farmers	3	8.82	25	73.5	4	11.8	2	5.88	34	100

**Population characteristics:** The population characteristics of households sampled for socio-economic survey in Himalapur Micro watershed is presented in Table 2. The data indicated that, there were 83 (60.58%) men and 54 (39.42%) were women. The average population of landless was 3.7, marginal farmers were 3.9, small farmers were 4.3 and semi medium farmers were 6.

**Table 2. Population characteristics in Himalapur micro-watershed**

Sl.No.	Particulars	LL (11)		MF (97)		SF (17)		SMF (12)		All (137)	
		N	%	N	%	N	%	N	%	N	%
1	Men	6	54.6	59	61	11	65	7	58.3	83	60.6
2	Women	5	45.5	38	39	6	35	5	41.7	54	39.4
Total		11	100	97	100	17	100	12	100	137	100
Average		3.7		3.9		4.3		6.0		4.0	

**Age wise classification of population:** The age wise classification of household members in Himalapur Micro watershed is presented in Table 3. The indicated that, 31 (22.63%) of population were 0-15 years of age, 58 (42.34%) were 16-35 years of age, 44(32.12%) were 36-60 years of age and 4 (2.92 %) were above 61 years of age.

**Table 3: Age wise classification of members of the household in Himalapur micro-watershed**

Sl.No.	Particulars	LL (11)		MF (97)		SF (17)		SMF (12)		All (137)	
		N	%	N	%	N	%	N	%	N	%
1	0-15 years of age	4	36.4	25	25.8	1	5.88	1	8.33	31	22.63
2	16-35 years of age	4	36.4	41	42.3	9	52.9	4	33.33	58	42.34
3	36-60 years of age	3	27.3	28	28.9	7	41.2	6	50	44	32.12
4	> 61 years	0	0	3	3.09	0	0	1	8.33	4	2.92
Total		11	100	97	100	17	100	12	100	137	100

**Education level of household members:** Education level of household members in Himalapur Micro watershed is presented in Table 4. The results indicated that, there were 49.64 per cent of illiterates, 0.73 per cent of functional literate, 23.36 per cent of them had primary school education, 2.92 per cent middle school education, 10.95 per cent high school education, 5.11 per cent of them had PUC education, 0.73 per cent of them had Diploma, 3.65 per cent attained graduation and 2.92 them had other education.

**Table 4. Education level of members of the household in Himalapur micro-watershed**

Sl.No.	Particulars	LL (11)		MF (97)		SF (17)		SMF (12)		All (137)	
		N	%	N	%	N	%	N	%	N	%
1	Illiterate	5	45.5	47	48.5	8	47.1	8	66.7	68	49.6
2	Functional Literate	0	0	1	1.03	0	0	0	0	1	0.73
3	Primary School	0	0	26	26.8	4	23.5	2	16.7	32	23.4
4	Middle School	1	9.09	2	2.06	0	0	1	8.33	4	2.92
5	High School	4	36.4	9	9.28	2	11.8	0	0	15	11
6	PUC	0	0	6	6.19	1	5.88	0	0	7	5.11
7	Diploma	0	0	1	1.03	0	0	0	0	1	0.73
8	Others	0	0	3	3.09	0	0	1	8.33	4	2.92
Total		11	100	97	100	17	100	12	100	137	100

**Occupation of head of households:** The data regarding the occupation of the household heads in Himalapur Micro watershed is presented in Table 5. The results indicate that, 76.47 per cent of households heads were practicing agriculture, 5.88 per cent of the household heads were agricultural Labour and housewife (14.71%).

**Table 5: Occupation of heads of households in Himalapur micro-watershed**

Sl.No.	Particulars	LL (3)		MF (25)		SF (4)		SMF (2)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Agriculture	0	0	21	84	3	75	2	100	26	76.47
2	Agricultural Labour	2	67	0	0	0	0	0	0	2	5.88
3	Housewife	1	33	3	12	1	25	0	0	5	14.71
Total		3	100	24	100	4	100	2	100	33	100

**Table 6: Occupation of members of the household in Himalapur micro-watershed**

Sl.No.	Particulars	LL (11)		MF (97)		SF (17)		SMF (12)		All (137)	
		N	%	N	%	N	%	N	%	N	%
1	Agriculture	0	0	34	35.1	8	47.06	6	50	48	35
2	Agricultural Labour	4	36.4	0	0	0	0	0	0	4	2.92
3	Student	4	36.4	31	32	2	11.76	0	0	37	27
4	Others	0	0	2	2.06	0	0	0	0	2	1.46
5	Housewife	3	27.3	27	27.8	7	41.18	4	33.33	41	29.9
6	Children	0	0	3	3.09	0	0	2	16.67	5	3.65
Total		11	100	97	100	17	100	12	100	137	100

**Occupation of the members of the household:** The data regarding the occupation of the household members in Himalapur Micro watershed is presented in Table 6. The results indicate that, agriculture was the major occupation for 35.04 per cent of the household

members, 2.92 per cent were agricultural labour, 27.01 per cent were working in pursuing education, 29.93 per cent were involved as housewife and 3.65 per cent were children.

**Institutional Participation of household members:** The data regarding the institutional participation of the household members in Himalapur Micro watershed is presented in Table 7. The results show that, out of the total family members in the households 100 per cent of them were not participating in any of the institutions.

**Table 7: Institutional Participation of household member in Himalapur micro-watershed**

Sl.No.	Particulars	LL (11)		MF (97)		SF (17)		SMF (12)		All (137)	
		N	%	N	%	N	%	N	%	N	%
1	No Participation	11	100	97	100	17	100	12	100	137	100
Total		11	100	97	100	17	100	12	100	137	100

**Type of house owned:** The data regarding the type of house owned by the households in Himalapur Micro watershed is presented in Table 8. The results indicate that, 2.94 percent possess thatched house, 94.12 per cent of the households possess katcha house, 2.94 per cent possess pacca house.

**Table 8. Type of house owned by households in Himalapur micro-watershed**

Sl.No.	Particulars	LL (3)		MF (25)		SF (4)		SMF (2)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Thatched	0	0	1	4	0	0	0	0	1	2.94
2	Katcha	3	100	23	92	4	100	2	100	32	94.12
3	Pucca/RCC	0	0	1	4	0	0	0	0	1	2.94
Total		3	100	25	100	4	100	2	100	34	100

**Durable assets owned by the households:** The data regarding the Durable Assets owned by the households in Himalapur Micro watershed is presented in Table 9. The results shows that, 100.00 per cent possess TV and Mobile phone.

**Table 9. Durable assets owned by households in Himalapur micro-watershed**

Sl.No.	Particulars	LL (3)		MF (25)		SF (4)		SMF (2)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Television	3	100	25	100	4	100	2	100	34	100
2	Mobile Phone	3	100	25	100	4	100	2	100	34	100

**Average value of durable assets:** The data regarding the average value of durable assets owned by the households in Himalapur Micro watershed is presented in Table 10. The result shows that, the average value of television was Rs.9000.00 and mobile phone was Rs.2982.00.

**Table 10. Average value of durable assets owned in Himalapur micro-watershed**

Sl.No.	Particulars	Average Value (Rs.)									
		LL (3)		MF (25)		SF (4)		SMF (2)		All (34)	
1	Television	9000		9000		9000		9000		9000	
2	Mobile Phone	3000		2947		3375		2666		2982	

**Farm implements owned:** The data regarding the farm implements owned by the households in Himalapur Micro watershed is presented in Table 11. About 0.00 per cent of the households possess Bullock Cart, 8.82 per cent possess plough and 17.65 per cent possess Weeder.

**Table 11. Farm implements owned in Himalapur micro-watershed**

Sl.No.	Particulars	LL (3)		MF (25)		SF (4)		SMF (2)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Plough	0	0	2	8	0	0	1	50	3	8.82
2	Weeder	0	0	3	12	2	50	1	50	6	17.65
3	Blank	3	100	20	80	2	50	1	50	26	76.47

**Average value of farm implements:** The data regarding the average value of farm Implements owned by the households in Himalapur Micro watershed is presented in Table 12. The results show that the average value of plough was Rs.1200.00 and weeder was Rs.768.00.

**Table 12. Average value of farm implements in Himalapur micro-watershed**

Average Value (Rs.)

Sl.No.	Particulars	LL (3)	MF (25)	SF (4)	SMF (2)	All (34)
1	Plough	0	1200	0	1200	1200
2	Weeder	0	50	2025	50	768

**Livestock possession by the households:** The data regarding the Livestock possession by the households in Himalapur Micro watershed is presented in Table 13. The results indicate that, 14.71 per cent of the households possess bullocks.

**Table 13. Livestock possession by households in Himalapur micro-watershed**

Sl.No.	Particulars	LL (3)		MF (25)		SF (4)		SMF (2)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Bullock	0	0	2	8	2	50	1	50	5	14.71
2	blank	3	100	23	92	2	50	1	50	29	85.29

**Average Labour availability:** The data regarding the average labour availability in Himalapur Micro watershed is presented in Table 14. The indicated that, own labour men available in the micro watershed was 1.53, women available in the micro watershed was 1.38, hired labour (men) available was 8.91 and hired labour (women) available was 7.34.

**Table 14. Average labour availability in Himalapur micro-watershed**

Sl.No.	Particulars	LL (3)	MF (25)	SF (4)	SMF (2)	All (34)
		N	N	N	N	N
1	Hired labour Female	0	5.96	11.25	17.5	7.34
2	Own Labour Female	0	1.23	1.75	2.5	1.38
3	Own labour Male	0	1.35	2	3	1.53
4	Hired labour Male	0	7.31	15	17.5	8.91

**Adequacy of hired labour:** The data regarding the adequacy of hired labour in Himalapur Micro watershed is presented in Table 15. The results indicate that, 94.12 per cent of the household opined that hired labour was adequate.



**Table 15. Adequacy of hired labour in Himalapur micro-watershed**

Sl.No.	Particulars	LL (3)		MF (25)		SF (4)		SMF (2)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Adequate	0	0	26	104	4	100	2	100	32	94.1

**Distribution of land (ha):** The data regarding the distribution of land (ha) in Himalapur Micro watershed is presented in Table 16. The results indicate that, 22.82 ha (94.95%) of dry land and 1.21 ha (5.05 %) of irrigated land.

**Table 16. Distribution of land (ha) in Himalapur micro-watershed**

Sl.No.	Particulars	LL (3)		MF (25)		SF (4)		SMF (2)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Dry	0	0	14.7	100	5.65	100	2.44	66.78	22.82	94.95
2	Irrigated	0	0	0	0	0	0	1.21	33.22	1.21	5.05
Total		0	100	14.7	100	5.65	100	3.65	100	24.04	100

**Average value of land (ha):** The data regarding the average land value (Rs./ha) in Himalapur Micro watershed is presented in Table 17. The results show that the average value of dry land was Rs.540298.81 and the average value of irrigated land was Rs.247000.00.

**Table 17. Average value of land (ha) in Himalapur micro-watershed**

Sl.No.	Particulars	LL (3)	MF (25)	SF (4)	SMF (2)	All (34)
		N	N	N	N	N
1	Dry	0	674160.7	300788	286733	540298.8
2	Irrigated	0	0	0	247000	247000

**Status of bore wells:** The data regarding the status of bore wells in Himalapur Micro watershed is presented in Table 18. The results indicate that, there was 1 functioning bore wells among the sampled households in micro watershed.

**Table 18. Status of bore wells in Himalapur micro-watershed**

Sl.No.	Particulars	LL (3)	MF (25)	SF (4)	SMF (2)	All (34)
		N	N	N	N	N
1	De-functioning	0	0	0	0	0
2	Functioning	0	0	0	1	1

**Source of irrigation:** The data regarding the source of irrigation in Himalapur Micro watershed is presented in Table 19. The results that bore well were major source of irrigation for 2.94 per cent of the households.

**Table 19. Source of irrigation in Himalapur micro-watershed**

Sl.No.	Particulars	LL (3)		MF (25)		SF (4)		SMF (2)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Bore Well	0	0	0	0	0	0	1	50	1	2.94

**Depth of water (Avg. In meters):** The data regarding the depth of water in Himalapur Micro watershed is presented in Table 20. The results revealed that, the depth of bore well was 3.14 meter.

**Table 20. Depth of water (Avg. In meters) in Himalapur micro-watershed**

Sl.No.	Particulars	LL (3)	MF (25)	SF (4)	SMF (2)	All (34)
		N	N	N	N	N
1	Bore Well	0	0	0	53.34	3.14

**Irrigated Area (ha):** The data regarding the irrigated area (ha) in Himalapur Micro watershed is presented in Table 21. The results indicate that, the availability of irrigation water was used for kharif crops was 1.21 ha.

**Table 21. Irrigated Area (ha) in Himalapur micro-watershed**

Sl.No.	Particulars	LL (3)	MF (25)	SF (4)	SMF (2)	All (34)
1	Kharif	0	0	0	1.21	1.21
Total		0	0	0	1.21	1.21

**Cropping pattern:** The data regarding the cropping pattern in Himalapur Micro watershed is presented in Table 22. The results indicate that, farmers have grown Red gram (9.83 ha), Cotton (8.51 ha), Bengal gram (3.13 ha) and Groundnut (2.56 ha).

**Table 22. Cropping pattern in Himalapur micro-watershed**

Sl.No.	Particulars	LL (3)	MF (25)	SF (4)	SMF (2)	All (34)
1	Kharif - Red gram (togari)	0	7.28	2.55	0	9.83
2	Kharif - Cotton	0	6.06	0	2.44	8.51
3	Kharif - Bengal gram	0	1.37	1.76	0	3.13
4	Kharif - Groundnut	0	0	1.34	1.21	2.56
Total		0	14.72	5.65	3.66	24.02

**Cropping intensity:** The data regarding the cropping intensity in Himalapur Micro watershed is presented in Table 23. The results indicate that, the cropping intensity was 100.00 per cent.

**Table 23. Cropping intensity (%) in Himalapur micro-watershed**

Sl.No.	Particulars	LL (3)	MF (25)	SF (4)	SMF (2)	All (34)
1	Cropping Intensity	0	100	100	100	100

**Cost of Cultivation of Red gram:** The data regarding the cost of cultivation (Rs/ha) of Red gram in Himalapur micro watershed is presented in Table 24.a. The results indicate that, the total cost of cultivation (Rs/ha) for Red gram was Rs. 36642.31. The gross income realized by the farmers was Rs. 128502.20. The net income from Red gram cultivation was Rs.91859.89, thus the benefit cost ratio was found to be 1:3.50.

**Table 24(a). Cost of Cultivation of Red gram in Himalapur micro-watershed**

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
<b>I</b>	<b>Cost A1</b>				
1	Hired Human Labour	Man days	62.01	11575.58	31.59
2	Bullock	Pairs/day	0.87	519.3	1.42
3	Tractor	Hours	5.74	4589.18	12.52
4	Machinery	Hours	1.4	1121.38	3.06
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	6.16	765.4	2.09
6	Seed Inter Crop	Kgs.	0	0	0
7	FYM	Quintal	3.08	1118.38	3.05
8	Fertilizer + micronutrients	Quintal	4.08	3148.66	8.59
9	Pesticides (PPC)	Kgs / liters	1.54	1538.59	4.2
10	Irrigation	Number	0	0	0
13	Depreciation charges		0	9.82	0.03
14	Land revenue and Taxes		0	3.29	0.01
<b>II</b>	<b>Cost B1</b>				
16	Interest on working capital			788.64	2.15
17	<b>Cost B1 = (Cost A1 + sum of 15 and 16)</b>			25178.23	68.71
<b>III</b>	<b>Cost B2</b>				
18	Rental Value of Land			333.33	0.91
19	<b>Cost B2 = (Cost B1 + Rental value)</b>			25511.56	69.62
<b>IV</b>	<b>Cost C1</b>				
20	Family Human Labour		33.65	7798.63	21.28
21	<b>Cost C1 = (Cost B2 + Family Labour)</b>			33310.19	90.91
<b>V</b>	<b>Cost C2</b>				
22	Risk Premium			1	0
23	<b>Cost C2 = (Cost C1 + Risk Premium)</b>			33311.19	90.91
<b>VI</b>	<b>Cost C3</b>				
24	Managerial Cost			3331.12	9.09
25	<b>Cost C3 = (Cost C2 + Managerial Cost)</b>			36642.31	100
<b>VII</b>	<b>Economics of the Crop</b>				
a.	Main Product	a) Main Product (q)		25.45	113532.22
		b) Main Crop Sales Price (Rs.)			4461.54
	By Product	e) Main Product (q)		14.42	14969.98
		f) Main Crop Sales Price (Rs.)			1038.46
b.	Gross Income (Rs.)			128502.2	
c.	Net Income (Rs.)			91859.89	
d.	Cost per Quintal (Rs./q.)			1439.95	
e.	Benefit Cost Ratio (BC Ratio)			1:3.5	

**Cost of Cultivation of Cotton;** The data regarding the cost of cultivation (Rs/ha) of Cotton in Himalapur micro watershed is presented in Table 24.b. The results indicate that, the total cost of cultivation (Rs/ha) for Cotton was Rs. 78378.30. The gross income realized by the farmers was Rs. 85727.69. The net income from Cotton cultivation was Rs.7349.38, thus the benefit cost ratio was found to be 1:1.10.

**Table 24(b). Cost of Cultivation of Cotton in Himalapur micro-watershed**

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
<b>I</b>	<b>Cost A1</b>				
1	Hired Human Labour	Man days	110.78	21332.59	27.22
2	Bullock	Pairs/day	0	0	0
3	Tractor	Hours	10.21	8171.58	10.43
4	Machinery	Hours	0	0	0
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	7.57	7195.96	9.18
6	Seed Inter Crop	Kgs.	0	0	0
7	FYM	Quintal	6.86	1372.68	1.75
8	Fertilizer + micronutrients	Quintal	17.44	13774	17.57
9	Pesticides (PPC)	Kgs / liters	3.43	3431.71	4.38
10	Irrigation	Number	0	0	0
11	Repairs		0	0	0
12	Msc. Charges (Marketing costs etc)		0	0	0
13	Depreciation charges		0	43.65	0.06
14	Land revenue and Taxes		0	3.29	0
<b>II</b>	<b>Cost B1</b>				
16	Interest on working capital			3093.04	3.95
17	<b>Cost B1 = (Cost A1 + sum of 15 and 16)</b>			58418.51	74.53
<b>III</b>	<b>Cost B2</b>				
18	Rental Value of Land			277.78	0.35
19	<b>Cost B2 = (Cost B1 + Rental value)</b>			58696.28	74.89
<b>IV</b>	<b>Cost C1</b>				
20	Family Human Labour		55.45	12555.72	16.02
21	<b>Cost C1 = (Cost B2 + Family Labour)</b>			71252	90.91
<b>V</b>	<b>Cost C2</b>				
22	Risk Premium			1	0
23	<b>Cost C2 = (Cost C1 + Risk Premium)</b>			71253	90.91
<b>VI</b>	<b>Cost C3</b>				
24	Managerial Cost			7125.3	9.09
25	<b>Cost C3 = (Cost C2 + Managerial Cost)</b>			78378.3	100
<b>VII</b>	<b>Economics of the Crop</b>				
a.	Main Product	a) Main Product (q)		18.7	85727.69
		b) Main Crop Sales Price (Rs.)			4583.33
b.	Gross Income (Rs.)				85727.69
c.	Net Income (Rs.)				7349.38
d.	Cost per Quintal (Rs./q.)				4190.41
e.	Benefit Cost Ratio (BC Ratio)				1:1.1

**Cost of Cultivation of Groundnut:** The data regarding the cost of cultivation (Rs/ha) of Groundnut in Himalapur micro watershed is presented in Table 24.c. The results indicate, the total cost of cultivation (Rs/ha) for Groundnut was Rs.30333.70. The gross income realized by the farmers was Rs. 100684.74. The net income from Groundnut cultivation was Rs. 70351.04, thus the benefit cost ratio was found to be 1:3.30.

**Table 24(c). Cost of Cultivation of Groundnut in Himalapur micro-watershed**

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
<b>I</b>	<b>Cost A1</b>				
1	Hired Human Labour	Man days	35.91	7068.52	23.3
2	Bullock	Pairs/day	2.88	1729	5.7
3	Tractor	Hours	5.21	4166.27	13.73
4	Machinery	Hours	0.82	658.67	2.17
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	2.06	370.5	1.22
6	Seed Inter Crop	Kgs.	0	0	0
7	FYM	Quintal	1.57	313.46	1.03
8	Fertilizer + micronutrients	Quintal	3.88	3064.09	10.1
9	Pesticides (PPC)	Kgs / liters	1.16	1155.64	3.81
10	Irrigation	Number	0	0	0
13	Depreciation charges		0	11.45	0.04
14	Land revenue and Taxes		0	3.29	0.01
<b>II</b>	<b>Cost B1</b>				
16	Interest on working capital			588.56	1.94
17	<b>Cost B1 = (Cost A1 + sum of 15 and 16)</b>			19129.44	63.06
<b>III</b>	<b>Cost B2</b>				
18	Rental Value of Land			333.33	1.1
19	<b>Cost B2 = (Cost B1 + Rental value)</b>			19462.78	64.16
<b>IV</b>	<b>Cost C1</b>				
20	Family Human Labour		34.96	8112.31	26.74
21	<b>Cost C1 = (Cost B2 + Family Labour)</b>			27575.09	90.91
<b>V</b>	<b>Cost C2</b>				
22	Risk Premium			1	0
23	<b>Cost C2 = (Cost C1 + Risk Premium)</b>			27576.09	90.91
<b>VI</b>	<b>Cost C3</b>				
24	Managerial Cost			2757.61	9.09
25	<b>Cost C3 = (Cost C2 + Managerial Cost)</b>			30333.7	100
<b>VII</b>	<b>Economics of the Crop</b>				
a.	Main Product	a) Main Product (q)		15.67	62692.37
		b) Main Crop Sales Price (Rs.)			4000
	By Product	e) Main Product (q)		19	37992.37
		f) Main Crop Sales Price (Rs.)			2000
b.	Gross Income (Rs.)			100684.74	
c.	Net Income (Rs.)			70351.04	
d.	Cost per Quintal (Rs./q.)			1935.4	
e.	Benefit Cost Ratio (BC Ratio)			1:3.3	

**Cost of Cultivation of Bengal gram:** The data regarding the cost of cultivation (Rs/ha) of Bengal gram in Himalapur micro watershed is presented in Table 24.d. The results indicate that, the total cost of cultivation (Rs/ha) for Bengal gram was Rs. 116785.65. The gross income realized by the farmers was Rs.115490.62. The net income from Bengal gram cultivation was Rs. -1295.04, thus the benefit cost ratio was found to be 1:0.99.

**Table 24(d). Cost of Cultivation of Bengal gram in Himalapur micro-watershed**

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
<b>I</b>	<b>Cost A1</b>				
1	Hired Human Labour	Man days	227.02	43145.26	36.94
2	Bullock	Pairs/day	0	0	0
3	Tractor	Hours	10.43	8340.87	7.14
4	Machinery	Hours	0	0	0
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	16.32	2340.44	2
6	Seed Inter Crop	Kgs.	0	0	0
7	FYM	Quintal	8.16	1632.76	1.4
8	Fertilizer + micronutrients	Quintal	16.61	13123.08	11.24
9	Pesticides (PPC)	Kgs / liters	7.17	7169.39	6.14
10	Irrigation	Number	0	0	0
11	Repairs		0	0	0
12	Msc. Charges (Marketing costs etc)		0	0	0
13	Depreciation charges		0	0.14	0
14	Land revenue and Taxes		0	3.29	0
<b>II</b>	<b>Cost B1</b>				
16	Interest on working capital			2912	2.49
17	<b>Cost B1 = (Cost A1 + sum of 15 and 16)</b>			78667.24	67.36
<b>III</b>	<b>Cost B2</b>				
18	Rental Value of Land			333.33	0.29
19	<b>Cost B2 = (Cost B1 + Rental value)</b>			79000.57	67.65
<b>IV</b>	<b>Cost C1</b>				
20	Family Human Labour		119.23	27167.2	23.26
21	<b>Cost C1 = (Cost B2 + Family Labour)</b>			106167.78	90.91
<b>V</b>	<b>Cost C2</b>				
22	Risk Premium			1	0
23	<b>Cost C2 = (Cost C1 + Risk Premium)</b>			106168.78	90.91
<b>VI</b>	<b>Cost C3</b>				
24	Managerial Cost			10616.88	9.09
25	<b>Cost C3 = (Cost C2 + Managerial Cost)</b>			116785.65	100
<b>VII</b>	<b>Economics of the Crop</b>				
a.	Main Product	a) Main Product (q)	22	115490.62	
		b) Main Crop Sales Price (Rs.)		5250	
b.	Gross Income (Rs.)			115490.62	
c.	Net Income (Rs.)			-1295.04	
d.	Cost per Quintal (Rs./q.)			5308.87	
e.	Benefit Cost Ratio (BC Ratio)			1:0.99	

**Adequacy of fodder:** The data regarding the adequacy of fodder in Himalapur Micro watershed is presented in Table 25. The results indicate that, 2.94 per cent of the households opined that dry fodder was adequate. With respect to green fodder availability, 2.94 percent of them opined it was sufficient.

**Table 25. Adequacy of fodder in Himalapur micro-watershed**

Sl.No.	Particulars	LL (3)		MF (25)		SF (4)		SMF (2)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Adequate-Dry Fodder	0	0	0	0	1	25	0	0	1	2.94
2	Adequate-Green Fodder	0	0	0	0	1	25	0	0	1	2.94

**Average annual gross income:** The data regarding the annual gross income in Himalapur Micro watershed is presented in Table 26. The results indicate that, the farmers have annual gross income of Rs. 90000.00 in micro-watershed, of which Rs. 39794.12 is from agriculture itself.

**Table 26. Average annual gross income in Himalapur micro-watershed**

Sl.No.	Particulars	LL (3)	MF (25)	SF (4)	SMF (2)	All (34)
		Rs.	Rs.	Rs.	Rs.	Rs.
1	Wage	54000	47800	57500	60000	50205.9
2	Agriculture	0	35760	72750	84000	39794.1
Income(Rs.)		54000	83560	130250	144000	90000

**Average annual Expenditure:** The data regarding the average annual expenditure in Himalapur Micro watershed is presented in Table 27. The results indicate that, the farmers have annual gross expenditure of Rs. 176943.33 in micro-watershed, of which Rs. 16235.29 is from agriculture itself.

**Table 27. Average annual Expenditure in Himalapur micro-watershed**

Sl.No.	Particulars	LL (3)	MF (25)	SF (4)	SMF (2)	All (34)
		Rs.	Rs.	Rs.	Rs.	Rs.
1	Wage	33333.3	20600	25750	24500	22558.8
2	Agriculture	0	15760	22000	35000	16235.3
Total		33333.3	36360	47750	59500	176943

**Horticulture species grown:** The data regarding horticulture species grown in Himalapur Micro watershed is presented in Table 28. The results indicate that, the total number of horticultural trees grown (both field and backyard) by the sampled households was Mango (1).

**Table 28. Horticulture species grown in Himalapur micro-watershed**

Sl.No.	Particulars	LL (3)		MF (25)		SF (4)		SMF (2)		All (34)	
		F	B	F	B	F	B	F	B	F	B
1	Mango	0	0	1	0	0	0	0	0	1	0

\*F= Field B=Back Yard

**Forest species grown:** The data regarding forest species grown in Himalapur Micro watershed is presented in Table 29. The results indicate that, households have planted 58 neem trees and 3 banyan trees together in both field and backyard.

**Table 29. Forest species grown in Himalapur micro-watershed**

Sl.No.	Particulars	LL (3)		MF (25)		SF (4)		SMF (2)		All (34)	
		F	B	F	B	F	B	F	B	F	B
1	Neem	0	0	49	0	7	0	2	0	58	0
2	Banyan	0	0	2	0	1	0	0	0	3	0

\*F= Field B=Back Yard

**Marketing of agricultural produce:** The data regarding marketing of the agricultural produce in Himalapur Micro watershed is presented in Table 30. The results indicated that, 74.36 percent of output of Bengal gram was sold in the market with average price of Rs. 5250.00; 100.00 percent of output of Cotton was sold in the market with average price of Rs. 4583.33; 75.00 percent of output of Groundnut was sold in the market with average price of Rs. 4000.00 and 76.56 percent of output of Red gram was sold in the market with average price of Rs. 4461.54.

**Table 30. Marketing of agricultural produce in Himalapur micro-watershed**

Sl. No	Crops	Output obtained (q)	Output retained (q)	Output sold (q)	Output sold (%)	Avg. Price obtained (Rs/q)
1	Bengal gram (Kadale)	39	10	29	74	5250
2	Cotton	103	0	103	100	4583
3	Groundnut	40	10	30	75	4000
4	Red gram	192	45	147	77	4462

**Marketing channels used for sale of agricultural produce:** The data regarding marketing channels used for sale of agricultural produce in Himalapur Micro watershed is presented in Table 31. The results indicated that, 91.18 cent of the households have sold agricultural produce to the local/village merchants.

**Table 31. Marketing channels used for sale of agricultural produce in Himalapur micro-watershed**

Sl.No.	Particulars	LL (3)		MF (25)		SF (4)		SMF (2)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Local/village Merchant	0	0	25	100	4	100	2	100	31	91.18

**Mode of transport of agricultural produce:** The data regarding mode of transport of agricultural produce in Himalapur Micro watershed is presented in Table 32. The results indicated that, 91.18 cent of the households have used tractor for the transport of agriculture commodity.



**Table 32. Mode of transport of agricultural produce in Himalapur micro-watershed**

Sl.No.	Particulars	LL (3)		MF (25)		SF (4)		SMF (2)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Tractor	0	0	25	100	4	100	2	100	31	91.18

**Incidence of soil and water erosion problems:** The data regarding incidence of incidence of soil and water erosion problems in Himalapur Micro watershed is presented in Table 33. The results indicate that, 91.18 per cent of the households have experienced soil and water erosion problems.

**Table 33. Incidence of soil and water erosion problems in Himalapur micro-watershed**

Sl.No.	Particulars	LL (3)		MF (25)		SF (4)		SMF (2)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Soil and water erosion problems in the farm	0	0	25	100	4	100	2	100	31	91.18

**Interest towards soil testing:** The data regarding Interest shown towards soil testing in Himalapur Micro watershed is presented in Table 34. The results indicated that, 91.18 per cent of the households were interested towards soil testing.

**Table 34. Interest regarding soil testing in Himalapur micro-watershed**

Sl.No.	Particulars	LL (3)		MF (25)		SF (4)		SMF (2)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Interest in soil test	0	0	25	100	4	100	2	100	31	91.18

**Usage pattern of fuel for domestic use:** The data on usage pattern of fuel for domestic use in Himalapur Micro watershed is presented in Table 35. The results indicated that, firewood was the major source of fuel for domestic use for 100.00.

**Table 35. Usage pattern of fuel for domestic use in Himalapur micro-watershed**

Sl.No.	Particulars	LL (3)		MF (25)		SF (4)		SMF (2)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Fire Wood	3	100	25	100	4	100	2	100	34	100

**Source of drinking water:** The data on source of drinking water in Himalapur Micro watershed is presented in Table 36. The results indicated that, piped waters supply was the major source for drinking water for 100 per cent of the households.

**Table 36. Source of drinking water in Himalapur micro-watershed**

Sl.No.	Particulars	LL (3)		MF (25)		SF (4)		SMF (2)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Piped supply	3	100	25	100	4	100	2	100	34	100

**Source of light:** The data on source of light in Himalapur Micro watershed is presented in Table 37. The results indicated that, electricity was the major source of light for 100.00 per cent of the households.

**Table 37. Source of light in Himalapur micro-watershed**

Sl.No.	Particulars	LL (3)		MF (25)		SF (4)		SMF (2)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Electricity	3	100	25	100	4	100	2	100	34	100

**Existence of sanitary toilet facility:** The data on availability of toilet facility in Himalapur Micro watershed is presented in Table 38. The results indicated that, 100.00 per cent of the households possess toilets.

**Table 38. Existence of sanitary toilet facility in Himalapur micro-watershed**

Sl.No.	Particulars	LL (3)		MF (25)		SF (4)		SMF (2)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Sanitary toilet facility	3	100	25	100	4	100	2	100	34	100

**Possession of PDS card:** The data regarding possession of PDS card in Himalapur Micro watershed is presented in Table 39. The results indicated that, 100.00 per cent of the households possessed BPL card.

**Table 39. Possession of PDS card in Himalapur micro-watershed**

Sl.No.	Particulars	LL (3)		MF (25)		SF (4)		SMF (2)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	BPL	3	100	25	100	4	100	2	100	34	100

**Adequacy of food items:** The data regarding adequacy of food items in Himalapur Micro watershed is presented in Table 40. The results indicated that, the extent of adequacy of food items for cereals, pulses, Oilseeds and vegetables were 100.00, 94.12, 0.00, 100.00 per cent respectively, similarly for Fruits (2.94%), milk (100.00%), Egg (97.06%) and Meat (100.00%).

**Table 40. Adequacy of food items in Himalapur micro-watershed**

Sl.No.	Particulars	LL (3)		MF (25)		SF (4)		SMF (2)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Cereals	3	100	25	100	4	100	2	100	34	100
2	Pulses	3	100	23	92	4	100	2	100	32	94.12
3	Vegetables	2	66.7	26	104	4	100	2	100	34	100
4	Fruits	0	0	1	4	0	0	0	0	1	2.94
5	Milk	3	100	25	100	4	100	2	100	34	100
6	Egg	3	100	24	96	4	100	2	100	33	97.06
7	Meat	3	100	25	100	4	100	2	100	34	100

**Inadequacy of food items:** The data regarding in adequacy of food items in Himalapur Micro watershed is presented in Table 41. The results indicated that, the extent of in adequacy of food items for pulses, Oilseeds and vegetables were 5.88, 94.12, 2.94 per cent respectively, similarly for fruits (97.06%) and egg (2.94%).

**Table 41. Inadequacy of food items in Himalapur micro-watershed**

Sl.No.	Particulars	LL (3)		MF (25)		SF (4)		SMF (2)		All (34)	
		N	%	N	%	N	%	N	%	N	%
1	Pulses	0	0	2	8	0	0	0	0	2	5.88
2	Oilseed	2	66.7	24	96	4	100	2	100	32	94.12
3	Vegetables	1	33.3	0	0	0	0	0	0	1	2.94
4	Fruits	3	100	24	96	4	100	2	100	33	97.06
5	Egg	0	0	1	4	0	0	0	0	1	2.94

**Farming constraints:** The data regarding farming constraints experienced by households in Himalapur Micro watershed is presented in Table 42. The results indicated that, lower fertility status of the soil was the constraint experienced by (91.18 %) per cent of the households, wild animal menace on farm field (8.82%), frequent incidence of pest and diseases (88.24%), inadequacy of irrigation water (2.94%), high cost of fertilizers and plant protection chemicals (88.24%), high rate of interest on credit (14.71%), low price for the agricultural commodities (76.47 %), lack of marketing facilities in the area (79.41%), inadequate extension services (8.82 %) and lack of transport for safe transport of the agricultural produce to the market (64.71%).

**Table 42. Farming constraints experienced in Himalapur micro-watershed**

SN	Particulars	MF (25)		SF (4)		SMF (2)		All (34)	
		N	%	N	%	N	%	N	%
1	Lower fertility status of the soil	25	100	4	100	2	100	31	91.18
2	Wild animal menace on farm field	3	12	0	0	0	0	3	8.82
3	Frequent incidence of pest and diseases	24	96	4	100	2	100	30	88.24
4	Inadequacy of irrigation water	1	4	0	0	0	0	1	2.94
5	High cost of Fertilizers and plant protection chemicals	24	96	4	100	2	100	30	88.24
6	High rate of interest on credit	3	12	1	25	1	50	5	14.71
7	Low price for the agricultural commodities	21	84	3	75	2	100	26	76.47
8	Lack of marketing facilities in the area	24	96	2	50	1	50	27	79.41
9	Inadequate extension services	3	12	0	0	0	0	3	8.82
10	Lack of transport for safe transport of the Agril produce to the market.	17	68	3	75	2	100	22	64.71



**SUMMARY AND IMPLICATIONS**

In order to assess the socio-economic condition of the farmers in the watershed 34 households located in the micro watershed were interviewed for the survey. The study was conducted in Himalapur micro-watershed (Mokadampur sub-watershed, Yadgiri taluk & District) is located at North latitude  $16^{\circ} 52' 34.382''$  and  $16^{\circ} 50' 38.077''$  and East longitude  $77^{\circ} 20' 51.651''$  and  $77^{\circ} 19' 20.923''$  covering an area of about 498.01 ha bounded by under Himalapura and Chapetla Villages.

Socio-economic analysis of Himalapur micro watersheds of Mokadampur sub-watershed, Yadgiri taluk & District indicated that, out of the total sample of 34 farmers were sampled in Himalapur micro-watershed among households surveyed 25 (73.53%) were marginal, 4 (11.76%) were small and 2 (5.88 %) were semi medium farmers. 3 landless farmers were also interviewed for the survey. The population characteristics of households indicated that, there were 83 (60.58%) men and 54 (39.42 %) were women. The average population of landless was 3.7, marginal farmers were 3.9, small farmers were 4.3 and semi medium farmers were 6.

Majority of the respondents (42.34%) were in the age group of 16-35 years. Education level of the sample households indicated that, there were 49.64 per cent illiterates, 0.73 percent were functional literates, 47.45 per cent pre university education and 3.65 per cent attained graduation. About, 76.47 per cent of household heads practicing agriculture and 5.88 per cent of the household heads were engaged as agricultural labourers. Agriculture was the major occupation for 35.04 per cent of the household members.

In the study area, 94.12 per cent of the households possess katcha house and 2.94 per cent possess pucca house. The durable assets owned by the households showed that, 100.00 per cent possess TV. Farm implements owned by the households indicated that, 8.82 per cent of the households possess plough. The average labour availability in the study area showed that, own labour men available in the micro watershed was 1.53, women available in the micro watershed was 1.38, hired labour (men) available was 8.91 and hired labour (women) available was 7.34.

Out of the total land holding of the sample respondents 94.95 per cent (24.04 ha) of the area is under dry condition and the remaining 5.05 per cent area is irrigated land. There were 1.00 live bore wells among the sampled households. Bore well was the major source of irrigation for 2.94 per cent of the households. The major crops grown by sample farmers are Red gram, Cotton, Groundnut, Bengal gram and cropping intensity was recorded as 100.00 per cent.

The per hectare cost of cultivation for Red gram, Cotton, Groundnut, Bengal gram and 0 was Rs.36642.31 , 78378.30, 30333.70, 116785.65 with benefit cost ratio of 1:3.50, 1: 1.10, 1: 3.30, 1: 0.99 respectively. Further, 2.94 per cent of the households opined that dry fodder was adequate and 2.94 per cent of the households have opined that the green fodder was adequate.

The average annual gross income of the farmers was Rs. 90000.00 in micro-watershed, of which Rs. 39794.12 comes from agriculture. Sampled households have grown 1 horticulture trees and 61 forestry trees together in the fields and back yards. Regarding marketing channels, 91.18 per cent of the households have sold agricultural produce to the local/village merchants. Further, 91.18 per cent of the households have used tractor for the transport of agriculture commodity.

Majority of the farmers (91.18%) have experienced soil and water erosion problems in the watershed and 91.18 per cent of the households were interested towards soil testing. Fire was the major source of fuel for domestic use for 100.00 per cent of the households. Piped supply was the major source for drinking water for 100.00 per cent of the households. Electricity was the major source of light for 100.00 per cent of the households.

In the study area, 100.00 per cent of the households possess toilet facility. Regarding possession of PDS card, 100.00 per cent of the households possessed BPL card. Households opined that, the requirement of cereals (100.00%), pulses (94.12%) are adequate for consumption.

Farming constraints experienced by households in the micro watersheds were lower fertility status of the soil (91.18%) wild animal menace on farm field (8.82%), frequent incidence of pest and diseases (88.24%), inadequacy of irrigation water (2.94%), high cost of fertilizers and plant protection chemicals (88.24%), high rate of interest on credit (14.71%), low price for the agricultural commodities (76.47%), lack of marketing facilities in the area (79.41%), inadequate extension services (8.82%), lack of transport for safe transport of the agricultural produce to the market (64.71%).

### **Implications of the survey**

- ✓ Result indicated that, there were 49.64 per cent were illiterate hence, extension methodologies such as demonstration, street play, drama, video shows will be effective in dissemination of the technologies in the micro watershed.
- ✓ The data indicate that, 94.12 per cent of the households possess katcha house. Hence, the development department while implementing the watershed plan should focus on agriculture to enhance the productivity of major crops in the area to increase the income of the farmers.

- ✓ Results indicated that the local institutional participation of the household members in the micro watershed is minimal hence, activities like membership campaign, awareness creation about the benefits of membership in local institutions and strengths of organized groups must be conveyed.
- ✓ Majority of the households in the watershed have experience in use of mobile phones, and television hence, these mass media can be effectively utilized for transfer of technology as well as for information dissemination.
- ✓ The farm machinery/implement possession in the micro watershed was found to be minimum the reasons may lack of knowledge or lack of financial ability which can be addressed through training on use of different farm implements, providing information on different sources of finance for purchase of farm implements.
- ✓ The possession of livestock such as crossbred cow found is less hence, farmers must be made aware of the benefits of crossbred cow in increased milk production.
- ✓ The possession of livestock such as sheep, goat and poultry was found to be low hence, farmers may be informed the role of subsidiary enterprises in enhancing the income and information on financial support for subsidiary activities.
- ✓ The data indicate that, job/work was the reason for all the migrants hence, farmers may be trained on profitable agriculture or self employment such as animal husbandry, plate making, sheep rearing, goat rearing, rabbit rearing with suitable information on sources of financial support.
- ✓ The results indicate that there was a change in quality of life due to migration hence, the developmental departments should take actions to arrest migration and to improve the quality of the life in rural areas.
- ✓ Households possess 22.82ha (94.95 %) of dry land and 1.21ha (5.05 %) of irrigated land hence, the availability of the dry land agricultural technologies such as short duration crops, high yielding drought resistance crop varieties, drip irrigation technology and subsidy information will be helpful for the farmers to enhance the productivity of land and as well as farmers income.
- ✓ Few of the bore well in micro watershed found non functional hence, farmers may be trained on possibility of bore well rejuvenation.
- ✓ Bore well was major source of irrigation for 2.94 per cent of the households. Hence, in order to increase the area under irrigation as well as to increase the water use efficiency farmers may trained on drip irrigation and provide the information on subsidy for drip irrigation equipment's along with the information on different agencies which provides the financial assistance for drip irrigation.
- ✓ The cropping intensity in the micro watershed was found to be (100.00 %) hence, care must be taken by the implementing agency to bring uncultivated land into cultivation through suitable measures.

- ✓ Many of the household members have borrowed loan from cooperative banks which has higher rate of interest hence, farmers may be sensitized on the different sources of credit with lesser interest rate such SHGs etc.
- ✓ The results indicated the non availability of both green and dry fodder throughout the year hence, fodder development activities can be taken up in the micro watershed.
- ✓ The average annual gross income of the households Rs.39794.12 from agriculture and Rs. 50205.88 from wages. Agriculture was found to be the major source of income for households hence; the development activities should focus on productivity enhancement, marketing arrangements and agricultural technology dissemination to have a direct impact on the farmers.
- ✓ The cultivation of forest species is found minimal hence, information and production technology related to agro-forestry and integrated farming system.
- ✓ The data indicated that, 91.18 per cent of the households have experienced soil and water erosion problems. Hence, those farmers who reported the soil and water erosion problems may be given attention while implementation of the watershed development plan.
- ✓ The data indicated that, 91.18 per cent of the households have interest in soil testing hence, farmers must be provided with the information on various institutions which are involved in soil testing for the benefit of the farmers.
- ✓ Except summer ploughing the adoption of other soil and water conservation structures is minimum hence, the farmers in the micro watershed should be sensitized on the use of different conservation structures for soil water conservation.
- ✓ Cereals and pulses found be adequate for per cent of the households respectively hence, farm households and the farm women must be trained on importance of balanced nutrition and role of vegetable, milk, egg, meat in balanced diet.
- ✓ Lower fertility status of the soil (91.18%), wild animal menace on farm field (8.82%), frequent incidence of pest and diseases (88.24%), high cost of fertilizers and plant protection chemicals (88.24%), high rate of interest on credit (14.71%), low price for the agricultural commodities (76.47%), lack of marketing facilities in the area (79.41%), inadequate extension services (8.82%), lack of transport for safe transport of the agricultural produce to the market (64.71%) were the major farming constraints experienced hence, these constraints must be addressed immediately for the welfare of the farmers. Awareness to be created among the farmers to approach nearest KVKs/RSKs and other developmental departments for technical and for subsidized inputs and utilize the well established regulated markets, approaching the contract firms, direct markets to avoid the involvement of middlemen.