



हर कदम, हर डगर  
किसानों का हमसफर  
भारतीय कृषि अनुसंधान परिषद

*AgriSearch with a human touch*

**LAND RESOURCE INVENTORY AND SOCIO-ECONOMIC STATUS OF  
FARM HOUSEHOLDS FOR WATERSHED PLANNING AND  
DEVELOPMENT**

**RAISABAD HOSALLI-1 (4D5B1J1c) MICROWATERSHED**

**Yadgir & Hattakuni Hobli, Yadgir Taluk and District, Karnataka**

**Karnataka Watershed Development Project – II**

**SUJALA – III**

**World Bank funded Project**



**ICAR – NATIONAL BUREAU OF SOIL SURVEY AND LAND USE PLANNING**



ICAR - NBSS & LUP



**WATERSHED DEVELOPMENT DEPARTMENT  
GOVT. OF KARNATAKA, BANGALORE**



## **About ICAR - NBSS&LUP**

The ICAR-National Bureau of Soil Survey and Land Use Planning (ICAR-NBSS&LUP), Nagpur, a premier Institute of the Indian Council of Agricultural Research (ICAR), was set up during 1976 with the objective to prepare soil resource maps at national, state and district levels and to provide research inputs in soil resource mapping and its applications, land evaluation, land use planning, land resource management, and database management using GIS for optimising land use on different kinds of soils in the country.

The Bureau has been engaged in carrying out soil resource survey, agro-ecological and soil degradation mapping at the country, state and district levels for qualitative assessment and monitoring the soil health towards viable land use planning. The research activities have resulted in identifying the soil potentials and problems, and the various applications of the soil surveys with the ultimate objective of sustainable agricultural development. The Bureau has the mandate to correlate and classify soils of the country and maintain a National Register of all the established soil series. The Institute is also imparting in-service training to staff of the soil survey agencies in the area of soil survey, land evaluation and soil survey interpretations for land use planning. The Bureau in collaboration with Panjabrao Krishi Vidyapeeth, Akola is running post-graduate teaching and research programme in land resource management, leading to M.Sc. and Ph.D. degrees.

**Citation:** Rajendra Hegde, Ramesh kumar, S.C., B.A. Dhanorkar, S. Srinivas, M.Lalitha, K.V. Niranjana, R.S. Reddy and S.K. Singh (2019). "Land Resource Inventory and Socio-Economic Status of Farm Households for Watershed Planning and Development of Raisabad Hosalli-1 (4D5B1J1c) Microwatershed, Yadgir & Hattakuni Hobli, Yadgir Taluk and District, Karnataka", ICAR-NBSS&LUP Sujala MWS Publ.277, ICAR – NBSS & LUP, RC, Bangalore. p.143 & 30.

### **TO OBTAIN COPIES,**

**Please write to:**

**Director, ICAR - NBSS & LUP,**

Amaravati Road, NAGPUR - 440 033, India

Phone : (0712) 2500386, 2500664, 2500545 (O)

Telefax : 0712-2522534

E-Mail : [director@nbsslup.ernet.in](mailto:director@nbsslup.ernet.in)

Website URL : [nbsslup.in](http://nbsslup.in)

Or

**Head, Regional Centre, ICAR - NBSS&LUP, Hebbal, Bangalore - 560 024**

Phone : (080) 23412242, 23510350 (O)

Telefax : 080-23510350

E-Mail : [nbssrcb@gmail.com](mailto:nbssrcb@gmail.com)

ICAR-NBSS&LUP Sujala MWS Publ.277



हर कदम, हर डगर  
किसानों का हमसफर  
भारतीय कृषि अनुसंधान परिषद

*Agri search with a human touch*

**LAND RESOURCE INVENTORY AND SOCIO-ECONOMIC  
STATUS OF FARM HOUSEHOLDS FOR WATERSHED  
PLANNING AND DEVELOPMENT**

**RAISABAD HOSALLI-1 (4D5B1J1c) MICROWATERSHED**

**Yadgir & Hattakuni Hobli, Yadgir Taluk and District, Karnataka**

**Karnataka Watershed Development Project – II**

**Sujala-III**

**World Bank funded Project**



**ICAR – NATIONAL BUREAU OF SOIL SURVEY AND LAND  
USE PLANNING**



ICAR - NBSS & LUP



**WATERSHED DEVELOPMENT DEPARTMENT, GOVT. OF  
KARNATAKA, BANGALORE**





## PREFACE

In Karnataka, as in other Indian States, the livelihoods of rural people are intertwined with farming pursuits. The challenges in agriculture are seriously threatening the livelihood of a large number of farmers as they have been practicing farming in contextual factors beyond their control. Climatic factors are the most important ones and have become much more significant in recent times due to rapid climate changes induced by intensive anthropogenic activities affecting our ecosystem in multiple ways. Climate change has become the reality, it is happening and efforts to evolve and demonstrate climate resilient technologies have become essential. Due to the already over stressed scenario of agrarian sector, the climate change is resulting in manifold increase in the complexities, pushing the rural mass to face more and more unpredictable situations. The rising temperatures and unpredictable rainfall patterns are going to test seriously the informed decisions farmers have to make in order to survive in farming and sustain their livelihood.

It is generally recognized that impacts of climate change shall not be uniform across the globe. It is said that impact of climate change is more severe in South Asia. Based on the analysis of meteorological data, it is predicted that in India, there will be upward trend in mean temperature, downward trend in relative humidity, annual rainfall and number of wet days in a year. Also, in general, phenomena like erratic monsoon, spread of tropical diseases, rise in sea levels, changes in availability of fresh water, frequent floods, droughts, heat waves, storms and hurricanes are predicted. Each one of these adverse situations are already being experienced in various parts of India and also at the global level. Decline in agricultural productivity of small and marginal farmers becoming more vulnerable is already witnessed.

In Karnataka, more than 60 per cent of the population live in rural areas and depend on agriculture and allied activities for their livelihood. Though the state has achieved significant progress in increasing the yield of many crops, there is tremendous pressure on the land resources due to the growing and competing demands of various land uses. This is reflected in the alarming rate of land degradation observed. Already more than 50 per cent of the area is affected by various forms of degradation. If this trend continues, the sustainability of the fragile ecosystem will be badly affected. The adverse effects of change in the climatic factors are putting additional stress on the land resources and the farmers dependent on this.

The natural resources (land, water and vegetation) of the state need adequate and constant care and management, backed by site-specific technological interventions and investments particularly by the government. Detailed database pertaining to the nature of the land resources, their constraints, inherent potentials and suitability for various land

based rural enterprises, crops and other uses is a prerequisite for preparing location-specific action plans, which are in tune with the inherent capability of the resources. Any effort to evolve climate resilient technologies has to be based on the baseline scientific database. Then only one can expect effective implementation of climate resilient technologies, monitor the progress, make essential review of the strategy, and finally evaluate the effectiveness of the implemented programs. The information available at present on the land resources of the state are of general nature and useful only for general purpose planning. Since the need of the hour is to have site-specific information suitable for farm level planning and detailed characterization and delineation of the existing land resources of an area into similar management units is the only option.

ICAR-NBSS&LUP, Regional Centre, Bangalore has taken up a project sponsored by the Karnataka Watershed Development Project-II, (Sujala-III), Government of Karnataka funded by the World Bank under Component-1 Land Resource Inventory. This study was taken up to demonstrate the utility of such a database in reviewing, monitoring and evaluating all the land based watershed development programs on a scientific footing. To meet the requirements of various land use planners at grassroots level, the present study on “Land Resource Inventory and Socio-Economic Status of Farm Households for Watershed Planning and Development of Raisabad Hosalli-1 Microwatershed, Yadgir Taluk and District, Karnataka” for integrated development was taken up in collaboration with the State Agricultural Universities, IISC, KRSRAC, KSNDMC as Consortia partners. The project provides detailed land resource information at cadastral level (1:7920 scale) for all the plots and socio-economic status of farm households covering thirty per cent farmers randomly selected representing landed and landless class of farmers in the micro-watershed. The project report with the accompanying maps for the Microwatershed will provide required detailed database for evolving effective land use plan, alternative land use options and conservation plans for the planners, administrators, agricultural extension personnel, KVK officials, developmental departments and other land users to manage the land resources in a sustainable manner.

It is hoped that this database will be useful to the planners, administrators and developmental agencies working in the area in not only for formulating location specific developmental schemes but also for their effective monitoring at the village/watershed level.

Nagpur

Date: 25-07-2019

**S.K. SINGH**

Director, ICAR NBSS&LUP, Nagpur

## Contributors

<b>Dr. Rajendra Hegde</b> Principal Scientist, Head & Project Leader, Sujala-III Project ICAR-NBSS&LUP, Regional Centre, Bangalore	<b>Dr. S.K.Singh</b> Director, ICAR-NBSS&LUP Coordinator, Sujala-III Project Nagpur
<b>Soil Survey, Mapping &amp; Report Preparation</b>	
Dr. B.A. Dhanorkar	Sh. R.S. Reddy
Dr. K.V. Niranjana	Mr. Somashekar T N
	Smt. Chaitra, S.P.
	Dr. Gopali bardhan
	Dr. Mahendra Kumar, M.B.
	Ms. Arpitha, G.M.
<b>Field Work</b>	
Sh. C.BacheGowda	Sh. Mahesh, D.B.
Sh. Somashekar	Sh. Ashok S Sindagi
Sh. M. Jayaramaiah	Sh. Veerabhadrapa B.
Sh. Paramesha, K.	Sh. Shankarappa
Sh. B. M. Narayana Reddy	Sh. Anand
	Sh. Arun N Kambar.
	Sh Kamalesh Awate
	Sh. Sharaan Kumar Huppar
	Sh. Yogesh H.N.
	Sh. Kalaveerachari R Kammar
<b>GIS Work</b>	
Dr. S.Srinivas	Sh. A.G.Devendra Prasad
Sh. D.H.Venkatesh	Sh. Prakashanaik, M.K.
Smt.K.Sujatha	Sh. Abhijith Sastry, N.S.
Smt. K.V.Archana	Sh. Sudip Kumar Suklabaidya
Sh. N. Maddileti	Sh. Avinash, K.N.
	Sh. Amar Suputhra, S
	Sh. Deepak, M.J.
	Smt. K.Karunya Lakshmi
	Ms. Seema, K.V.
	Ms. A. Rajab Nisha

<b>Laboratory Analysis</b>	
Dr. K.M.Nair	Ms. Steffi Peter
Smt. Arti Koyal	Ms. Thara, V.R
Smt. Parvathy	Ms. Roopa, G.
	Ms. Swati, H.
	Sh. Shantaveera Swami
	Ms. Shwetha, N.K.
	Smt. Ishrat Haji
	Ms. P. Pavan Kumari
	Ms. Padmaja
	Ms. Veena, M.
<b>Socio-Economic Analysis</b>	
Dr. S.C. Ramesh Kumar	Sh.M.K. Prakashanaik
	Mrs. Sowmya A N
	Ms. Karuna V Kulkarni
	Sh. Vijay Kumar Lamani
	Sh. Basavaraja
	Sh. Vinodh R
	Ms. Sowmya K.B
	Mrs. Prathibha, D.G
	Sh. Rajendra,D
<b>Soil &amp; Water Conservation</b>	
Sh. Sunil P. Maske	
<b>Watershed Development Department, GoK, Bangalore</b>	
Sh. Rajeev Ranjan IFS Project Director & Commissioner, WDD	Dr. A. Natarajan NRM Consultant, Sujala-III Project
Dr. S.D. Pathak IFS Executive Director & Chief Conservator of Forests, WDD	



# **PART-A**

## **LAND RESOURCE INVENTORY**



## Contents

Preface		
Contributors		
Executive Summary		
Chapter 1	Introduction	1
Chapter 2	Geographical Setting	3
2.1	Location and Extent	3
2.2	Geology	3
2.3	Physiography	4
2.4	Drainage	4
2.5	Climate	4
2.6	Natural Vegetation	6
2.7	Land Utilization	6
Chapter 3	Survey Methodology	11
3.1	Base maps	11
3.2	Image Interpretation for Physiography	11
3.3	Field Investigation	14
3.4	Soil Mapping	16
3.5	Land Management Units	17
3.6	Laboratory Characterization	17
Chapter 4	The Soils	23
4.1	Soils of granite gneiss landscape	23
Chapter 5	Interpretation for Land Resource Management	47
5.1	Land Capability Classification	47
5.2	Soil Depth	49
5.3	Surface Soil Texture	50
5.4	Soil Gravelliness	51
5.5	Available Water Capacity	52
5.6	Soil Slope	53
5.7	Soil Erosion	54
Chapter 6	Fertility Status	57
6.1	Soil Reaction (pH)	57
6.2	Electrical Conductivity (EC)	57
6.3	Organic Carbon (OC)	57
6.4	Available Phosphorus	59
6.5	Available Potassium	59
6.6	Available Sulphur	59
6.7	Available Boron	60
6.8	Available Iron	60
6.9	Available Manganese	60
6.10	Available Copper	60
6.11	Available Zinc	64

Chapter 7	Land Suitability for Major Crops	65
7.1	Land suitability for Sorghum	65
7.2	Land suitability for Maize	66
7.3	Land suitability for Bajra	67
7.4	Land suitability for Groundnut	68
7.5	Land suitability for Sunflower	69
7.6	Land suitability for Red gram	70
7.7	Land suitability for Bengal gram	71
7.8	Land suitability for Cotton	72
7.9	Land suitability for Chilli	73
7.10	Land suitability for Tomato	74
7.11	Land suitability for Brinjal	75
7.12	Land suitability for Onion	76
7.13	Land suitability for Bhendi	77
7.14	Land suitability for Drumstick	78
7.15	Land suitability for Mango	79
7.16	Land suitability for Guava	80
7.17	Land suitability for Sapota	81
7.18	Land Suitability for Pomegranate	82
7.19	Land Suitability for Musambi	83
7.20	Land Suitability for Lime	84
7.21	Land Suitability for Amla	85
7.22	Land Suitability for Cashew	86
7.23	Land Suitability for Jackfruit	87
7.24	Land Suitability for Jamun	88
7.25	Land Suitability for Custard apple	89
7.26	Land Suitability for Tamarind	90
7.27	Land Suitability for Mulberry	91
7.28	Land Suitability for Marigold	92
7.29	Land Suitability for Chrysanthemum	93
7.30	Land management units	125
7.31	Proposed Crop Plan	126
Chapter 8	Soil Health Management	129
Chapter 9	Soil and Water conservation Treatment Plan	135
9.1	Treatment Plan	136
9.2	Recommended Soil and Water Conservation measures	139
9.3	Greening of Microwatershed	140
	References	143
	Appendix I	I-XII
	Appendix II	XIII-XXVI
	Appendix III	XXV-XXXVII

## LIST OF TABLES

2.1	Mean Monthly Rainfall, PET, 1/2 PET at Yadgir Taluk & District	5
2.2	Land Utilization in Yadgir district	7
3.1	Differentiating Characteristics used for Identifying soil Series	16
3.2	Soil map unit description of Raisabad Hosalli-1 microwatershed	17
4.1	Physical and chemical characteristics of soil series identified in Raisabad Hosalli-1 microwatershed	33
7.1	Soil-Site Characteristics of Raisabad Hosalli-1 microwatershed	95
7.2	Land suitability for Sorghum	96
7.3	Land suitability for Maize	97
7.4	Land suitability for Bajra	98
7.5	Land suitability for Groundnut	99
7.6	Land suitability for Sunflower	100
7.7	Land suitability for Red gram	101
7.8	Land suitability for Bengal gram	102
7.9	Land suitability for Cotton	103
7.10	Land suitability for Chilli	104
7.11	Land suitability for Tomato	105
7.12	Land suitability for Brinjal	106
7.13	Land suitability for Onion	107
7.14	Land suitability for Bhendi	108
7.15	Land suitability for Drumstick	109
7.16	Land suitability for Mango	110
7.17	Land suitability for Guava	111
7.18	Land suitability for Sapota	112
7.19	Land suitability for Pomegranate	113
7.20	Land suitability for Musambi	114
7.21	Land suitability for Lime	115
7.22	Land suitability for Amla	116
7.23	Land suitability for Cashew	117
7.24	Land suitability for Jackfruit	118

7.25	Land suitability for Jamun	119
7.26	Land suitability for Custard apple	120
7.27	Land suitability for Tamarind	121
7.28	Land suitability for Mulberry	122
7.29	Land suitability for Marigold	123
7.30	Land suitability for Chrysanthemum	124
7.31	Proposed Crop Plan for Raisabad Hosalli-1 Microwatershed	127

## LIST OF FIGURES

2.1	Location map of Raisabad Hosalli-1 microwatershed	3
2.2	Granite and granite gneiss rock formation	4
2.3	Rainfall distribution in Yadgir Taluk and District	5
2.4	Natural vegetation of Raisabad Hosalli-1 microwatershed	6
2.5	Current land use map of Raisabad Hosalli-1 microwatershed	7
2.6 a	Different crops and cropping systems in Raisabad Hosalli-1 microwatershed	8
2.7	Location of wells- Raisabad Hosalli-1 microwatershed	9
3.1	Scanned and Digitized Cadastral map of Raisabad Hosalli-1 microwatershed	13
3.2	Satellite image of Raisabad Hosalli-1 microwatershed	13
3.3	Cadastral map overlaid on IRS PAN+LISS IV merged imagery of Raisabad Hosalli-1 microwatershed	14
3.4	Location of profiles in a transect	15
3.5	Soil phase or Management Units of Raisabad Hosalli-1 microwatershed	21
5.1	Land Capability Classification map of Raisabad Hosalli-1 microwatershed	49
5.2	Soil depth map of Raisabad Hosalli-1 microwatershed	50
5.3	Surface soil texture map of Raisabad Hosalli-1 microwatershed	51
5.4	Soil gravelliness map of Raisabad Hosalli-1 microwatershed	52
5.5	Soil available water capacity map Raisabad Hosalli-1 microwatershed	53
5.6	Soil slope map of Raisabad Hosalli-1 microwatershed	54
5.7	Soil erosion map of Raisabad Hosalli-1 microwatershed	55
6.1	Soil reaction (pH) map of Raisabad Hosalli-1 microwatershed	58
6.2	Electrical conductivity (EC) map of Raisabad Hosalli-1 microwatershed	58
6.3	Soil organic carbon (OC) map of Raisabad Hosalli-1 microwatershed	59
6.4	Soil available phosphorus map of Raisabad Hosalli-1 microwatershed	60
6.5	Soil available potassium map of Raisabad Hosalli-1 microwatershed	61
6.6	Soil available sulphur map of Raisabad Hosalli-1 microwatershed	61
6.7	Soil available boron map of Raisabad Hosalli-1 microwatershed	62
6.8	Soil available iron map of Raisabad Hosalli-1 microwatershed	62
6.9	Soil available manganese map of Raisabad Hosalli-1 microwatershed	63

6.10	Soil available copper map of Raisabad Hosalli-1 microwatershed	63
6.11	Soil available zinc map of Raisabad Hosalli-1 microwatershed	64
7.1	Land suitability for Sorghum	66
7.2	Land suitability for Maize	67
7.3	Land suitability for Bajra	68
7.4	Land suitability for Groundnut	69
7.5	Land suitability for Sunflower	70
7.6	Land suitability for Red gram	71
7.7	Land suitability for Bengal gram	72
7.8	Land suitability for Cotton	73
7.9	Land suitability for Chilli	74
7.10	Land suitability for Tomato	75
7.11	Land suitable for Brinjal	76
7.12	Land suitable for Onion	77
7.13	Land suitable for Bhendi	78
7.14	Land suitable for Drumstick	79
7.15	Land suitability for Mango	80
7.16	Land suitability for Guava	81
7.17	Land suitability for Sapota	82
7.18	Land suitability for Pomegranate	83
7.19	Land suitability for Musambi	84
7.20	Land suitability for Lime	85
7.21	Land suitability for Amla	86
7.22	Land suitability for Cashew	87
7.23	Land suitability for Jackfruit	88
7.24	Land suitability for Jamun	89
7.25	Land suitability for Custard apple	90
7.26	Land suitability for Tamarind	91
7.27	Land suitability for Mulberry	92
7.28	Land suitability for Marigold	93
7.29	Land suitability for Chrysanthemum	94
7.30	Land management units map of Raisabad Hosalli-1 microwatershed	126
9.1	Soil and water conservation plan map of Raisabad Hosalli-1 microwatershed	140



## **EXECUTIVE SUMMARY**

*The land resource inventory of Raisabad Hosalli-I Microwatershed was conducted using village cadastral maps and IRS satellite imagery on 1:7920 scale. The false colour composites of IRS imagery were interpreted for physiography and the physiographic delineations were used as base for mapping soils. The soils were studied in several transects and a soil map was prepared with phases of soil series as mapping units. Random checks were made all over the area outside the transects to confirm and validate the soil map unit boundaries. The soil map shows the geographic distribution and extent, characteristics, classification, behavior and use potentials of the soils in the microwatershed.*

*The present study covers an area of 887 ha in Yadgir taluk & district, Karnataka. The climate is semiarid and categorized as drought-prone with an average annual rainfall of 866 mm, of which about 652 mm is received during south-west monsoon, 138 mm during north-east and the remaining 76 mm during the rest of the year. An area of 817 ha in the microwatershed is covered by soils, 6 ha by rock outcrops and about 64 ha by others (habitation and water bodies). The salient findings from the land resource inventory are summarized briefly below.*

- ❖ The soils belong to 14 soil series and 22 soil phases (management units) and 9 land management units.*
- ❖ The length of crop growing period is about 120-150 days starting from 1<sup>st</sup> week of June to 4<sup>th</sup> week of October.*
- ❖ From the master soil map, several interpretative and thematic maps like land capability, soil depth, surface soil texture, soil gravelliness, available water capacity, soil slope and soil erosion were generated.*
- ❖ Soil fertility status maps for macro and micronutrients were generated based on the surface soil samples collected at every 320 m grid interval.*
- ❖ Land suitability for growing 29 major agricultural and horticultural crops was assessed and maps showing the degree of suitability along with constraints were generated.*
- ❖ Entire area in the microwatershed is suitable for agriculture.*
- ❖ About 67 per cent area of the microwatershed has soils that are moderately deep to very deep (75 - >150 cm) and 25 per cent soils are very shallow to moderately shallow (<25 - 75 cm).*
- ❖ About 3 per cent area in the microwatershed has sandy soils, 52 per cent soils are loamy and 37 per cent clayey soils at the surface.*
- ❖ About 64 per cent of the microwatershed area is non gravelly (<15%) and 28 per cent of the microwatershed area is gravelly (15 – 35%).*

- ❖ *About 25 per cent area of the microwatershed is very high (>200 mm/m) in available water capacity, 39 per cent is low (51-100 mm/m) and 28 per cent is very low (<50 mm/m).*
- ❖ *About <1 per cent area in the microwatershed has nearly level (0-1% slope) lands, 92 per cent has very gently sloping (1-3% slope) lands and <1 per cent area is gently sloping (3.5% slope).*
- ❖ *An area of about <1 is severely (e3) eroded, 92 per cent is moderately (e2) eroded and <1 per cent area is slightly (e1) eroded.*
- ❖ *An area of about 4 per cent soils are slightly acid (pH 6.0-6.5) in soil reaction, an area of 51 per cent is neutral (pH 6.5-7.3), 21 per cent soils are slightly alkaline (pH 7.3-7.8), 15 per cent soils are moderately alkaline (pH 7.8-8.4) and <1 per cent is strongly alkaline (pH 8.4-9.0).*
- ❖ *The Electrical Conductivity (EC) of the soils in the entire area of the microwatershed is dominantly <2 dsm<sup>-1</sup> indicating that the soils are non-saline.*
- ❖ *Available organic carbon is high (>0.75) in an area of 24 per cent and medium (0.5-0.75%) in an area of 68 per cent.*
- ❖ *About 20 per cent is low (<23 kg/ha) in available phosphorus, 70 per cent is medium (23-57 kg/ha) and high (>57 kg/ha) in an area of one per cent.*
- ❖ *About 8 per cent is low (145 kg/ha) in available potassium, 82 per cent is medium (145-337 kg/ha) and 2 per cent is high (>337 kg/ha).*
- ❖ *Available sulphur is low (<10 ppm) in an area of about 14 per cent and medium (10-20 ppm) in 78 per cent.*
- ❖ *Available boron content is low (<0.5 ppm) in an area of about 63 per cent and medium (0.5-1.0 ppm) in 29 per cent.*
- ❖ *Available iron is sufficient (>4.5 ppm) in the entire area of the microwatershed.*
- ❖ *Available manganese and copper are sufficient in all the soils of the microwatershed.*
- ❖ *Available zinc is deficient <0.6 ppm) in the entire area of the microwatershed.*
- ❖ *The land suitability for 29 major crops grown in the microwatershed were assessed and the areas that are highly suitable (S1) and moderately suitable (S2) are given below. It is however to be noted that a given soil may be suitable for various crops but what specific crop to be grown may be decided by the farmer looking to his capacity to invest on various inputs, marketing infrastructure, market price and finally the demand and supply position.*

**Land suitability for various crops in the Microwatershed**

<b>Crop</b>	<b>Suitability Area in ha (%)</b>		<b>Crop</b>	<b>Suitability Area in ha (%)</b>	
	<b>Highly suitable (S1)</b>	<b>Moderately suitable (S2)</b>		<b>Highly suitable (S1)</b>	<b>Moderately suitable (S2)</b>
<i>Sorghum</i>	243(27)	106(12)	<i>Guava</i>	-	223(25)
<i>Maize</i>	<1 (<1)	348(39)	<i>Sapota</i>	-	223(25)
<i>Bajra</i>	<1 (<1)	674(76)	<i>Pomegranate</i>	-	465(52)
<i>Groundnut</i>	-	367(41)	<i>Musambi</i>	303(34)	162(18)
<i>Sunflower</i>	225(25)	33(4)	<i>Lime</i>	303(34)	162(18)
<i>Red gram</i>	-	407(46)	<i>Amla</i>	74(8)	392(44)
<i>Bengal gram</i>	243(27)	106(12)	<i>Cashew</i>	-	15(2)
<i>Cotton</i>	219(25)	127(14)	<i>Jackfruit</i>	-	15(2)
<i>Chilli</i>	-	556(63)	<i>Jamun</i>	-	450(51)
<i>Tomato</i>	<1 (<1)	408(46)	<i>Custard apple</i>	192(22)	274(31)
<i>Brinjal</i>	217(24)	456(51)	<i>Tamarind</i>	-	450(51)
<i>Onion</i>	89(13)	216(32)	<i>Mulberry</i>	-	133(15)
<i>Bhendi</i>	217(24)	456(51)	<i>Marigold</i>	-	557(63)
<i>Drumstick</i>	-	584(66)	<i>Chrysanthemum</i>	-	557(63)
<i>Mango</i>	-	160(18)			

- ❖ *Apart from the individual crop suitability, a proposed crop plan has been prepared for the identified LMUs by considering only the highly and moderately suitable lands for different crops and cropping systems with food, fodder, fibre and horticulture crops.*
- ❖ *Maintaining soil-health is vital to crop production and conserve soil and land resource base for maintaining ecological balance and to mitigate climate change. For this, several ameliorative measures have been suggested to these problematic soils like saline/alkali, highly eroded, sandy soils etc.,*
- ❖ *Soil and water conservation treatment plan has been prepared that would help in identifying the sites to be treated and also the type of structures required.*
- ❖ *As part of the greening programme, several tree species have been suggested to be planted in marginal and sub marginal lands, field bunds and also in the hillocks, mounds and ridges. This would help in not only supplementing the farm income but also provide fodder and fuel and generate lot of biomass which would help in maintaining an ecological balance and also contribute to mitigating the climate change.*



## **INTRODUCTION**

Soil being a vital natural resource on whose proper use depends the life supporting systems of a country and the socioeconomic development of its people. Soils provide food, fodder, fibre and fuel for meeting the basic human and animal needs. With the ever increasing growth in human and animal population, the demand on soil for more food and fodder production is on the increase. The area available for agriculture is about 51 per cent of the total geographical area and more than 60 per cent of the people are still dependant on agriculture for their livelihood. However, the capacity of a soil to produce is limited and the limits to the production are set by its intrinsic characteristics, agro-climatic setting, and, use and management. There is, therefore, tremendous pressure on land and water resources, which is causing decline in soil-health and stagnation in productivity. The soils have been degrading at an estimated rate of one million hectares per year and ground water levels have been receding at an alarming rate resulting in decline in the ground water resource. Further, land degradation has emerged as a serious problem which has already affected about 38 lakh ha of cultivated area in the State. Soil erosion alone has degraded about 35 lakh ha. Almost all the uncultivated areas are facing various degrees of degradation, particularly soil erosion; salinity and alkalinity has emerged as a major problem in more than 3.5 lakh ha in the irrigated areas of the State. Nutrient depletion and declining factor productivity is common in both rainfed and irrigated areas. The degradation is continuing at an alarming rate and there appears to be no systematic effort among the stakeholders to contain this process. In recent times, an aberration of weather due to climate change phenomenon has added another dimension leading to unpredictable situation to be tackled by the farmers.

In this critical juncture, the challenge before us is not only to increase the productivity per unit area which is steadily declining and showing a fatigue syndrome, but also to prevent or at least reduce the severity of degradation. If the situation is not reversed at the earliest, then the sustainability of the already fragile crop production system and the overall ecosystem will be badly affected in the state. Added to this, every year there is a significant diversion of farm lands and water resources for non-agricultural purposes. Thus, developing strategies to slow down the degradation process or reclaim the soils to normal condition and ensure sustainability of production system are the major issues today. This demands a systematic appraisal of our soil and land resources with respect to their extent, geographic distribution, characteristics, behaviour and uses potential, which is very important for developing an effective land use and cropping systems for augmenting agricultural production on a sustainable basis.

The soil and land resource inventories made so far in Karnataka had limited utility because the surveys were of different types, scales and intensities carried out at different times with specific objectives. Hence, there is an urgent need to generate detailed site-

specific farm level database on various land resources for all the villages/watersheds in a time bound manner that would help to protect the valuable soil and land resources and also to stabilize the farm production. Therefore, the land resource inventory required for farm level planning is the one which investigates all the parameters which are critical for productivity *viz.*, soils, site characteristics like slope, erosion, gravelliness and stoniness, climate, water, topography, geology, hydrology, vegetation, crops, land use pattern, animal population, socio-economic conditions, infrastructure, marketing facilities and various schemes and developmental works of the government etc. From the data collected at farm level, the specific problems and potentials of the area can be identified and highlighted, conservation measures required for the area can be planned on a scientific footing, suitability of the area for various uses can be worked out and finally viable and sustainable land use options suitable for each and every land holding can be prescribed.

The Land Resource Inventory is basically done for identifying potential and problem areas, developing sustainable land use plans, estimation of surface run off and water harvesting potential, preparation of soil and water conservation plans, land degradation/desertification etc. The Bureau is presently engaged in developing an LRI methodology using high resolution satellite remote sensing data and Digital Elevation Model (DEM) data to prepare Landscape Ecological Units (LEU) map representing agro-ecosystem as a whole. The LEU is preferred over landform as the base map for LRI. LEU is the assemblage of landform, slope and land use. An attempt has already been made to upscale the soil resource information from 1:250000 and 1:50000 scale to the LEU map in Goa and in some other states.

The land resource inventory aims to provide site-specific database for Raisabad Hosalli-1 microwatershed in Yadgir Taluk and Yadgir District, Karnataka State for the Karnataka Watershed Development Department. The database was generated by using cadastral map of the village as a base along with high resolution IRS LISS IV and Cartosat-1 merged satellite imagery. Later, an attempt will be made to uplink this LRI data generated at 1:7920 scale under Sujala-III Project to the proposed Landscape Ecological Units (LEUs) map.

The study was organized and executed by the ICAR- National Bureau of Soil Survey and Land Use Planning, Regional Centre, Bangalore under Generation of Land Resource Inventory Data Base Component-1 of the Sujala-III Project funded by the World Bank.

## GEOGRAPHICAL SETTING

### 2.1 Location and Extent

The Raisabad Hosalli-1 microwatershed is located in the northern part of Karnataka in Yadgir Taluk & District, Karnataka State (Fig.2.1). It comprises parts of Mylapura, Rasabadha Hosalli, Yaleri, Naglapura, Panchasheelanagar and Jinatera villages. It lies between  $16^{\circ}42'$  –  $16^{\circ}43'$  North latitudes and  $77^{\circ}13'$  –  $77^{\circ}16'$  East longitudes, covering an area of about 887 ha. It is about 19 km southeast of Yadgir town and is surrounded by Mylapura on the north, Rasabadha Hosalli on the northeast and southeast, Yaleri village on the east, Naglapura village on south, Panchasheelanagar on the east, Halagera village on the west and Jinatera village on the southwestern side.




Fig.2.1 Location map of Raisabad Hosalli-1 Microwatershed

### 2.2 Geology

Major rock formation observed in the microwatershed are granite gneiss (Figs.2.2). They are essentially pink to gray and are coarse to medium grained. They consist

primarily of quartz, feldspar, biotite and hornblende. The gray granite gneisses are highly weathered, fractured and fissured upto a depth of about 10 m. Dolerite dykes and quartz veins are common with variable width and found to occur in Raisabad Hosalli-1 microwatershed.



Fig.2.2 Granite and granite gneiss rocks

### 2.3 Physiography

Physiographically, the area has been identified as granite gneiss landscape based on geology. The area has been further subdivided into five landforms, viz; mounds/ridges, summits, side slopes and very gently sloping uplands, plains and valleys based on slope and its relief features. The elevation ranges from 390-415 m above MSL. The mounds and ridges are mostly covered by rock outcrops.

### 2.4 Drainage

The area is drained by several parallel streams like Bori, Amerja and Kanga which finally join the river Bhima along its course. Though, they are not perennial, during rainy season they carry large quantities of rain water. The microwatershed has only few small tanks which are not capable of storing the water that flows during the rainy season. Due to this, the ground water recharge is very much affected. This is reflected in the failure of many bore wells in the villages. If the available rain water is properly harnessed by constructing new tanks and recharge structures at appropriate places in the villages, then the drinking and irrigation needs of the area can be easily met. The drainage network is parallel to sub parallel and dendritic.

### 2.5 Climate


The Yadgir district lies in the northern plains of Karnataka and falls under semiarid tract of the state and is categorized as drought- prone with total annual rainfall of 866 mm (Table 2.1). Of the total rainfall, maximum of 652 mm is received during the south-west monsoon period from June to September, the north-east monsoon from October to early December contributes about 138 mm and the remaining 76 mm during



the rest of the year. The summer season starts during the middle of February and continues up to the first week of June. The period from December to the middle of February is the coldest season. December is the coldest month with mean daily maximum and minimum temperatures being 29.5<sup>0</sup>C and 10<sup>0</sup>C respectively. During peak summer, temperature shoots up to 45<sup>0</sup>C. Relative humidity varies from 26% in summer to 62% in winter. Rainfall distribution is shown in Figure 2.3. The average Potential Evapo-Transpiration (PET) is 141 mm and varies from a low of 81 mm in December to 199 mm in the month of May. The PET is always higher than precipitation in all the months except July, August and September. Generally, the Length of crop Growing Period (LGP) is 120-150 days and starts from 1<sup>st</sup> week of June to 4<sup>th</sup> week of October.

**Table 2.1 Mean Monthly Rainfall, PET, 1/2 PET at Yadgir Taluk, Yadgir District**

Sl. No.	Months	Rainfall	PET	1/2 PET
1	January	4.30	86.0	43.0
2	February	2.30	125.5	62.7
3	March	15.10	166.0	83.0
4	April	18.50	179.8	89.9
5	May	36.0	198.8	97.9
6	June	118.0	175.1	87.5
7	July	171.80	156.3	78.1
8	August	182.9	150.3	75.1
9	September	179.7	142.0	71.0
10	October	105.3	138.5	69.2
11	November	26.4	97.60	48.6
12	December	6.0	80.90	40.4
<b>Total</b>		<b>866.3</b>		



**Fig 2.3 Rainfall distribution in Yadgir Taluk, Yadgir District**

## 2.6 Natural Vegetation

The natural vegetation is sparse comprising few tree species, shrubs and herbs. The mounds, ridges and boulders occupy very sizeable area which is under thin to moderately thick forest vegetation. Still, there are some remnants of the past forest cover which can be seen in patches in some ridges and hillocks in the microwatershed.

Apart from the continuing deforestation, the presence of large population of goats, sheep and other cattle in the microwatershed is causing vegetative degradation of whatever little vegetation left in the area. The uncontrolled grazing has left no time for the regeneration of the vegetative cover. This leads to the accelerated rate of erosion on the hill slopes resulting in the formation of deep gullies in the foot slopes that eventually result in the heavy siltation of tanks and reservoirs in the microwatershed.



Fig 2.4 Natural vegetation of Raisabad Hosalli-1 microwatershed


## 2.7 Land Utilization

About 72 per cent area (Table 2.2) in Yadgir district is cultivated at present. An area of about 2 per cent is permanently under pasture, 20 per cent under current fallows and 6 per cent under non-agricultural land, and 5 per cent under currently barren. Forests occupy an area of about 7 per cent and the tree cover is in a very poor state. Most of the mounds, ridges and bouldery areas have very poor vegetative cover. Major crops grown in the area are sorghum, maize, cotton, sunflower, groundnut, red gram and paddy. The cropping intensity is 120 per cent in the taluk. While carrying out land resource inventory, the land use/land cover particulars are collected from all the survey numbers and a current land use map of the microwatershed is prepared. The current land use map prepared shows the arable and non-arable lands, other land uses and different types of crops grown in the area. The current land use map of Raisabad Hosalli-1 microwatershed is presented in Fig.2.5. The different crops and cropping systems adopted in the microwatershed is

presented in the Figures 2.6 a & b. The occurrence and distribution of wells and bore wells in Raisabad Hosalli-1 microwatershed is show in figure 2.7.

**Table 2.2 Land Utilization in Yadgir District**

Sl. No.	Agricultural land use	Area ( ha)	Per cent
1.	Total geographical area	516088	-
2.	Total cultivated area	373617	72.4
3.	Area sown more than once	74081	14.3
4.	Cropping intensity	-	119.8
5.	Trees and grooves	737	0.14
6.	Forest	33773	6.54
7.	Cultivable wasteland	2385	0.46
8.	Permanent Pasture land	11755	2.28
9.	Barren land	27954	5.41
10.	Non- Agriculture land	29623	5.73
11.	Current Fallows	105212	20.4



**Fig.2.5 Current Land Use map of Raisabad Hosalli-1 Microwatershed**




Fig 2.6 a. Different Crops and Cropping Systems in Raisabad Hosalli-1 Microwatershed




Fig 2.7 Location of wells-Raisabad Hosalli-1 microwatershed



## SURVEY METHODOLOGY

The purpose of land resource inventory is to delineate similar areas (soil series and phases), which respond or expected to respond similarly for a given level of management. This was achieved in Raisabad Hosalli-1 microwatershed by the detailed study of all the soil characteristics (depth, texture, colour, structure, consistence, coarse fragments, porosity, soil reaction, soil horizons etc.) and site characteristics (slope of the land, erosion, drainage, occurrence of rock fragments etc.) followed by grouping of similar areas based on soil-site characteristics into homogeneous (management units) units, and showing the area extent and their geographic distribution on the microwatershed cadastral map. The detailed survey at 1:7920 scale was carried out in an area of 887 ha. The methodology followed for carrying out land resource inventory was as per the guidelines given in Soil Survey Manual (IARI, 1971; Soil Survey Staff, 2006; Natarajan *et al.*, 2015) which is briefly described below.

### 3.1 Base Maps

The detailed survey of the land resources occurring in the microwatershed was carried out by using digitized cadastral map and IRS satellite imagery as base supplied by KRSAC. The cadastral map shows field boundaries with their survey numbers, location of tanks, streams and other permanent features of the area (Fig. 3.1). Apart from the cadastral map, remote sensing data products from Cartosat-1 and LISS IV merged at the scale of 1:7920 were used in conjunction with the cadastral map to identify the landscapes, landforms and other surface features. The imagery helped in the identification and delineation of boundaries between hills, uplands and lowlands, water bodies, forest and vegetated areas, roads, habitations and other cultural features of the area (Fig. 3.2). The cadastral map was overlaid on the satellite imagery (Fig. 3.3) that helps to identify the parcel boundaries and other permanent features. Apart from cadastral maps and images, toposheets of the area (1:50,000 scale) were also used for initial traversing, identification of geology and landforms, drainage features, present land use and also for selection of transects in the microwatershed.

### 3.2 Image Interpretation for Physiography

False Colour Composites (FCCs) of Cartosat-I and LISS-IV merged satellite data covering microwatershed area was visually interpreted using image interpretation elements and all the available collateral data with local knowledge. The delineated physiographic boundaries were transferred on to a cadastral map overlaid on satellite imagery. Physiographically, the area has been identified as granite gneiss landscape. It was divided into five landforms, *viz*; ridges and mounds, gently and very gently sloping uplands and lowlands based on slope. They were further subdivided into

physiographic/image interpretation units based on image characteristics. The image interpretation legend for physiography is given below.

## **Image Interpretation Legend for Physiography**

### **G- Granite Gneiss Landscape**

G1	Hills/ Ridges/ Mounds
G11	Summits
G12	Side slopes
G121	Side slopes with dark grey tones
G2	Uplands
G21	Summits
G22	Gently sloping uplands
G221	Gently sloping uplands, yellowish green (eroded)
G222	Gently sloping uplands, yellowish white (severely eroded)
G23	Very gently sloping uplands
G231	Very gently sloping uplands, yellowish green
G232	Very gently sloping uplands, medium green and pink
G233	Very gently sloping uplands, pink and green (scrub land)
G234	Very gently sloping uplands, medium greenish grey
G235	Very gently sloping uplands, yellowish white (eroded)
G236	Very gently sloping uplands, dark green
G237	Very gently sloping uplands, medium pink (coconut garden)
G238	Very gently sloping uplands, pink and bluish white (eroded)
G24	Valleys/ lowlands
G241	Valleys, pink tones
G242	Valleys gray mixed with pink tones






Fig 3.1 Scanned and Digitized Cadastral map of Raisabad Hosalli-1 Microwatershed




Fig.3.2 Satellite Image of Raisabad Hosalli-1 Microwatershed




Fig.3.3 Cadastral map overlaid on IRS PAN+LISS IV merged imagery of Raisabad Hosalli-1 Microwatershed

### 3.3 Field Investigation

The field boundaries and survey numbers given on the cadastral sheet were located on the ground by following permanent features like roads, cart tracks, *nallas*, streams, tanks etc., and wherever changes were noticed, they were incorporated on the microwatershed cadastral map. Preliminary traverse of the microwatershed was carried out with the help of cadastral map, imagery and toposheets. While traversing, landforms and physiographic units identified were checked and preliminary soil legend was prepared by studying soils at few selected places. Then, intensive traversing of each physiographic unit like hills, ridges, uplands and valleys was carried out. Based on the variability observed on the surface, transects (Fig. 3.4) were selected across the slope covering all the landform units in the microwatershed (Natarajan and Dipak Sarkar, 2010).




Fig: 3.4. Location of profiles in a transect

In the selected transect, soil profiles were located (Fig. 3.4) at closely spaced intervals to take care of any change in the land features like break in slope, erosion, gravel, stones etc. In the selected sites, soil profiles (vertical cut showing the soil layers from surface to the rock) were opened upto 200 cm or to the depth limited by rock or hard substratum and studied in detail for all their morphological and physical characteristics. The soil and site characteristics were recorded for all profile sites on a standard proforma as per the guidelines given in USDA Soil Survey Manual (Soil Survey Staff, 2012). Apart from the transect study, profiles were also studied at random, almost like in a grid pattern, outside the transect areas.

Based on the soil characteristics, the soils were grouped into different soil series. Soil series is the most homogeneous unit having similar horizons and properties and behaves similarly for a given level of management. Soil depth, texture, colour, kind of horizon and horizon sequence, calcareousness, amount and nature of gravel present, nature of substratum etc, were used as the major differentiating characteristics for identifying soil series occurring in the area. The differentiating characteristics used for identifying the soil series are given in Table 3.1. Based on the above characteristics, 14 soil series were identified in the Raisabad Hosalli-1 microwatershed.

**Table 3.1 Differentiating Characteristics used for identifying Soil Series**  
(Characteristics are of Series Control Section)

<b>Soils of Granite gneiss Landscape</b>							
<b>Sl. no</b>	<b>Soil Series</b>	<b>Depth (cm)</b>	<b>Colour (moist)</b>	<b>Texture</b>	<b>Gravel (%)</b>	<b>Horizon sequence</b>	<b>Calcareousness</b>
1	BDL (Badiyala)	25-50	7.5 YR 2.5/3, 2.5/2, 3/3 10YR 3/4, 4/3	sl	-	Ap-Bw	e
2	HLG (Halagera)	50-75	10YR 3/2, 4/4 7.5YR 4/3, 4/2	scl	-	Ap-Bw	es
3	JNK (Jinkera)	50-75	10YR 3/1,3/2 7.5YR 3/4	scl	-	Ap-Bw	e
4	YLR (Yalleri)	50-75	2.5YR 3/4, 4/4, 5YR 3/4, 7.5YR 4/4	c	15-35	Ap-Bt	-
5	BLC (Balichakra)	75-100	2.5YR 5/3, 2.5/4, 5YR 4/3, 3/3	scl	-	Ap-Bt	-
6	PGP (Poglapur)	75-100	5YR 4/6, 3/3, 7.5YR 4/4	sc	-	Ap-Bt	-
7	YDR (Yadgir)	100-150	10YR4/3,4/4 2.5Y4/3,5/3	sl	-	Ap-Ac	-
8	MDG (Mundargi)	100-150	10YR 4/4,3/3 7.5YR 4/4	scl	-	Ap-Bw	-
9	BDP (Baddepalli)	<25	7.5YR 3/2,3/4 5YR 3/4	scl	-	Ap-Ac	es
10	KBD (Kalabelagundi )	75-100	2.5Y4/4,3/4 5YR 4/2, 4/3	g scl	35-60	Ap-Bt	-
11	HTK (Hattikuni)	25-50	10YR 4/6,4/4, 7.5YR 4/4,3/3	sl	10-25	Ap-Ac	-
12	KDH (Kadechoor)	75-100	10YR 3/2	sc	-	Ap-Bw	e
13	SGR (Sangawar)	>150	10YR 3/1,4/1	c	-	Ap-Bss	es
14	KDP (Kondapur)	>150	7.5YR 5/6, 10YR 4/2,4/4, 5/3	s	-	A-C	-

### 3.4 Soil Mapping

The area under each soil series was further separated into soil phases and their boundaries delineated on the cadastral map based on the variations observed in the texture of the surface soil, slope, erosion, presence of gravel, stoniness etc. A soil phase is a subdivision of soil series based mostly on surface features that affect its use and management. The soil mapping units are shown on the map (Fig.3.5) in the form of symbols. During the survey many soil profile pits, few minipits and a few auger bores representing different landforms occurring in the microwatershed were studied. In

addition to the profile study, spot observations in the form of minipits, road cuts, terrace cuts etc., were studied to validate the soil boundaries on the soil map.

The soil map shows the geographic distribution of 22 mapping units representing 14 soil series occurring in the microwatershed. The soil map unit (soil legend) description is presented in Table 3.2. The soil phase map (management units) shows the distribution of 22 soil phases mapped in the microwatershed. Each mapping unit (soil phase) delineated on the map has similar soil and site characteristics. In other words, all the farms or survey numbers included in one soil phase will have similar management needs and have to be treated accordingly.

### 3.5 Land Management Units (LMU's)

The 22 soil phases identified and mapped in the microwatershed were grouped into 9 Land Management Units (LMU's) for the purpose of preparing a Proposed Crop Plan for sustained development of the microwatershed. The database (soil phases) generated under LRI was utilized for identifying Land Management Units (LMU's) based on the management needs. One or more than one soil site characteristic having influence on the management have been chosen for identification and delineation of LMUs. For Raisabad Hosalli-1 microwatershed, five soil and site characteristics, namely soil depth, soil texture, slope, erosion and gravel content have been considered for defining LMUs. The Land Management Units are expected to behave similarly for a given level of management.

### 3.6 Laboratory Characterization

Soil samples for each soil series were collected from representative master profiles for laboratory characterization by following the methods outlined in the Laboratory Manual (Sarma *et al*, 1987). Surface soil samples collected from farmer's fields (86 samples) for fertility status (major and micronutrients) at 320 m grid interval in the year 2018 were analyzed in the laboratory (Katyal and Rattan, 2003). By linking the soil fertility data to the survey numbers through GIS, soil fertility maps were generated by using Kriging method for the microwatershed.

**Table 3.2 Soil map unit description of Raisabad Hosalli-1 Microwatershed**

Soil Map unit No.	Soil Series	Soil Phase	Mapping Unit Description	Area in ha (%)
	BDL		Badiyala soils are shallow (25-50 cm), well drained, have dark brown to very dark brown and dark yellowish brown, slightly calcareous sandy loam soils occurring on very gently to gently sloping uplands under cultivation	78 (8.81)
		BDLiB2	Sandy clay surface, slope 1-3%, moderate erosion	78 (8.81)
	HLG		Halagera soils are moderately shallow (50-75 cm), well drained, have very dark grayish brown to dark yellowish brown, calcareous sandy clay loam soils occurring on	13 (1.41)

		very gently sloping uplands under cultivation		
16		HLGcB2	Sandy loam surface, slope 1-3%, moderate erosion	4 (0.61)
	JNK	Jinkera soils are moderately shallow (50-75 cm), well drained, have dark brown to very dark grayish brown, slightly calcareous sandy clay loam soils occurring on very gently sloping uplands under cultivation		<b>77 (8.68)</b>
20		JNKcB2	Sandy loam surface, slope 1-3%, moderate erosion	25 (2.8)
23		JNKiB2g1	Sandy clay surface, slope 1-3%, moderate erosion, gravelly (15-35%)	52 (5.88)
	YLR	Yalleri soils are moderately shallow (50-75 cm), well drained, have brown to reddish brown and dark reddish brown, clay red soils occurring on very gently to gently sloping uplands under cultivation		<b>1 (0.06)</b>
30		YLRcC3	Sandy loam surface, slope 3-5%, severe erosion	1 (0.06)
	BLC	Balichakra soils are moderately deep (75-100 cm), well drained, have reddish brown to dark reddish brown, sandy clay loam red soils occurring on very gently sloping uplands under cultivation		<b>15 (1.65)</b>
38		BLCiB2	Sandy clay surface, slope 1-3, moderate erosion	15 (1.65)
	PGP	Poglapur soils are moderately deep (75-100 cm), well drained, have dark brown to dark reddish brown and yellowish red sandy clay soils occurring on very gently sloping uplands under cultivation		<b>0 (0.01)</b>
40		PGPcB2	Sandy loam surface, slope 1-3%, moderate erosion	0 (0.01)
	YDR	Yadgir soils are deep (100-150 cm), well drained, have brown to dark yellowish brown and olive brown, sodic sandy loam soils occurring on very gently sloping uplands under cultivation		<b>207 (23.41)</b>
42		YDRcB2	Sandy loam surface, slope 1-3%, moderate erosion	106 (12.01)
154		YDRcB2g1	Sandy loam surface, slope 1-3%, moderate erosion, gravelly (15-35%)	101 (11.4)
-	MDG	Mundargi soils are deep (100-150 cm), moderately well drained, have brown to dark yellowish brown, sandy clay loam soils occurring on very gently sloping uplands under cultivation		<b>160 (18.1)</b>
58		MDGiB2	Sandy clay surface, slope 1-3, moderate erosion	95 (10.75)
149		MDGhB2g1	Sandy clay loam surface, slope 1-3%, moderate erosion, gravelly (15-35%)	65 (7.35)
	BDP	Baddeppalli soils are very shallow (<25 cm), well drained, have dark brown to dark reddish brown, calcareous sandy clay loam soils occurring on very gently sloping uplands under cultivation		<b>22(2.46)</b>

118		BDPcB2	Sandy loam surface, slope 1-3%, moderate erosion	18 (2.06)
120		BDPhB2	Sandy clay loam surface, slope 1-3%, moderate erosion	4 (0.4)
	KBD	Kalabelagundi soils are moderately deep (75-100 cm), well drained, have reddish brown to dark reddish brown and dark reddish gray, gravelly sandy clay loam soils occurring on very gently sloping uplands under cultivation		<b>118(13.3)</b>
130		KBDhB2	Sandy clay loam surface, slope 1-3%, moderate erosion	52 (5.86)
131		KBDiB2	Sandy clay surface, slope 1-3, moderate erosion	66 (7.44)
	HTK	Hattikuni soils are shallow (25-50 cm), well drained, have dark yellowish brown sandy loam soils occurring on very gently sloping uplands under cultivation		<b>31 (3.47)</b>
161		HTKbB2g1	Loamy sand surface, slope 1-3%, moderate erosion, gravelly (15-35%)	31 (3.47)
	KDH	Kadechoor soils are moderately deep (75-100 cm), moderately well drained, have very dark grayish brown to dark brown, slightly calcareous sandy clay soils occurring on very gently to gently sloping lowlands under cultivation		<b>23.36(5.63)</b>
99		KDHcB2	Sandy loam surface, slope 1-3%, moderate erosion	23 (2.59)
157		KDHiA1	Sandy clay surface, slope 0-1%, slight erosion	0.36 (0.04)
	SGR	Sangwar soils are very deep (>150 cm), moderately well drained, have dark gray to very dark gray, sodic, calcareous cracking clay black soils occurring on very gently sloping lowlands under cultivation		<b>59(6.644)</b>
106		SGRmB2	Clay surface, slope 1-3%, moderate erosion	18 (2.05)
141		SGRcB2	Sandy loam surface, slope 1-3%, moderate erosion	41 (4.59)
143		SGRiB2	Sandy clay surface, slope 1-3, moderate erosion	0.03(0.004)
	KDP	Kondapur soils are very deep (>150 cm), somewhat excessively drained, have strong brown to dark grayish brown and brown sandy soils occurring on very gently to gently sloping lowlands under cultivation.		<b>13 (1.46)</b>
180		KDPhB2	Sandy clay loam surface, slope 1-3%, moderate erosion	13 (1.46)
999	Rock outcrops	Rock lands, both massive and bouldery with little or no soil		<b>6 (0.68)</b>
1000		Habitation and Water body		<b>64 (7.24)</b>








Fig 3.5 Soil Phase or Management Units - Raisabad Hosalli-1



## THE SOILS

Detailed information pertaining to the nature, extent and their distribution of different kinds of soils occurring in Raisabad Hosalli-1 microwatershed is provided in this chapter. The microwatershed area has been identified as granite gneiss landscape based on geology. In all, 9 soil series are identified. Soil formation is the result of the combined effect of environmental and terrain factors that are reflected in soil morphology. In the granite gneiss landscape, it is by parent material, relief and climate.

A brief description of each of the 14 soil series identified followed by 22 soil phases (management units) mapped are furnished below. The physical and chemical characteristics of soil series identified in Raisabad Hosalli-1 microwatershed are given in Table 4.1 along with soil classification. The soils in any one map unit differ from place to place in their depth, texture, slope, gravelliness, erosion or any other site characteristic that affect management. The soil phase map can be used for identifying the suitability of areas for growing specific crops or for other alternative uses and also for deciding the type of conservation structures needed. The detailed information on soil and site-characteristics like soil depth, surface soil texture, slope, erosion, gravelliness, AWC, LCC etc, with respect to each of the soil phase identified is given village/survey number wise for the microwatershed in Appendix-I.

### 4.1 Soils of granite gneiss landscape

In this landscape, 14 soil series are identified and mapped. Of these, YDR series occupies maximum area of 207 ha (23%) followed by MDG 80 ha (18%), KBD 118 ha (13%), BDL 78 ha (9%), JNK 77 ha (9%), SGR 59 ha (6%), HTK 31 ha (3%), KDH 23 ha (3%), BDP 22 ha (2%), BLC 15 ha (2%), HLG 13 ha (1%), KDP 13 ha (1%) YLR 1 ha (<1%), PGP 0 ha (<1%). The rock outcrops and others (habitation and water body) occupy an area of 6 ha (<1%) and 64 ha (7%) respectively. Brief description of each series identified and number of soil phases mapped is given below.

**4.1.1 Badiyala (BDL) Series:** Badiyala soils are shallow (25-50 cm), well drained, have very dark brown, dark yellow brown and dark brown, slightly calcareous sandy loam soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation. The Badiyala series has been classified as a member of the coarse-loamy, mixed, isohyperthermic family of Fluventic Haplustepts.

The thickness of the solum ranges from 28 to 50 cm. The thickness of A horizon ranges from 4 to 12 cm. Its colour is in 10YR hue with value 3 to 4 and chroma 3 to 4. The texture is loamy sand, sandy clay loam and sandy clay. The thickness of B horizon ranges from 27 to 45 cm. Its colour is in 10 YR and 7.5 YR hue with value 2 to 4 and chroma 3 to 4. Its texture is sandy loam to sandy clay loam and is slightly calcareous. The available water capacity is very low (<50mm/m). Only one phase was identified and mapped.



Landscape and Soil Profile characteristics of Badiyala (BDL) Series

**4.1.2 Halagera (HLG) Series:** Halagera soils are moderately shallow (50-75 cm), well drained, have very dark grayish brown to dark yellowish brown, calcareous sandy clay loam soils. They are developed from weathered granite gneiss and occur on very gently sloping uplands under cultivation. The Halagera series has been classified as a member of the fine-loamy, mixed (calcareous), isohyperthermic family of Typic Haplusteps.

The thickness of the solum ranges from 51 to 75 cm. The thickness of A horizon ranges from 9 to 15 cm. Its colour is in 7.5 YR and 10 YR hue with value 3 to 4 and chroma 2 to 4. The texture is loamy sand to sandy clay loam. The thickness of B horizon ranges from 44 to 66 cm. Its colour is in 10 YR and 7.5 YR hue with value 3 to 4 and chroma 2 to 3. Its texture varies from sandy clay loam to sandy clay and is calcareous. The available water capacity is low (51-100 mm/m). Only one phase was identified and mapped.



Landscape and Soil Profile characteristics of Vanakanahalli (VNK) Series

**4.1.3 Jinkera (JNK) Series:** Jinkera soils are moderately shallow (50-75 cm), well drained, have very dark gray to very dark grayish brown and dark brown, slightly calcareous sandy clay loam soils. They are developed from weathered granite gneiss and occur on very gently sloping uplands under cultivation. The Jinkera series has been classified as a member of the fine-loamy, mixed, isohyperthermic family of Typic Haplustepts.

The thickness of the solum ranges from 51-75 cm. Thickness of A horizon ranges from 6 to 11 cm. Its colour is in hue 10 YR and 7.5 YR with value and chroma of 3 to 4. The texture varies from sandy loam to sandy clay. The thickness of B horizon ranges from 53 to 66 cm. Its colour is in 10 YR and 7.5 YR hue with value and chroma of 2 to 4. The texture varies from sandy clay loam to sandy clay and is slightly calcareous. The available water capacity is low (51-100 mm/m). Two phases were identified and mapped.



Landscape and Soil Profile characteristics of Jinkera (JNK) Series

**4.1.4 Yalleri (YLR) Series:** Yalleri soils are moderately shallow (50-75 cm), well drained, have very dark reddish brown to dark brown, gravelly clay soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation. The Yalleri series has been classified as a member of the fine, mixed, isohyperthermic family of Typic Paleustalfs.

The thickness of the solum ranges from 50 to 74 cm. The thickness of A horizon ranges from 10 to 13 cm. Its colour is in 7.5 YR and 5 YR hue with value and chroma 2 to 4. The texture is sandy loam, loamy sand, and sandy clay loam. The thickness of B horizon ranges from 45 to 64 cm. Its colour is in 10 YR, 7.5 YR and 5 YR hue with value 2 to 4 and chroma 2 to 4. Its texture is clay with gravel content of 15-35 per cent. The available water capacity is low (51-100 mm/m). Only one phase was identified and mapped.



Landscape and Soil Profile characteristics of Yalleri (YLR) Series

**4.2.5 Balichakra (BLC) Series:** Balichakra soils are moderately deep (75-100 cm), well drained, dark reddish brown to reddish brown, sandy clay loam soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation. The Balichakra series has been classified as a member of the fine-loamy, mixed, isohyperthermic family of Typic Haplustalfs.

The thickness of the solum ranges from 80 to 100 cm. The thickness of A horizon ranges from 10 to 16 cm. Its colour is in hue 5 YR with value and chroma of 3 to 4. Its texture varies from sandy clay loam and sandy clay. The thickness of B horizon ranges from 70 to 88 cm. Its colour is in hue 2.5 YR and 5 YR with value 3 to 5 and chroma 3 to 4. Its texture is sandy clay loam to sandy clay and is slightly calcareous. The available water capacity is medium (101-150 mm/m). Only one phase was identified and mapped.



Landscape and Soil Profile characteristics of Balichakra (BLC) Series

**4.2.6 Poglapur (PGP) Series:** Poglapur soils are moderately deep (75-100 cm), well drained, have dark brown to dark reddish brown and yellowish red sandy clay red soils. They have developed from granite gneiss and occur on very gently sloping uplands under cultivation. The Poglapur series has been classified as a member of the fine, mixed, isohyperthermic family of Rhodic Paleustalfs.

The thickness of the solum ranges from 78 to 100 cm. The thickness of A horizon ranges from 8 to 17 cm. Its colour is in 7.5 YR hue with value 3 and chroma 3 to 4. Its texture varies from loamy sand to sandy clay loam and sandy clay. The thickness of B horizon ranges from 65 to 92 cm. Its colour is in 2.5 YR, 5 YR and 7.5 YR hue with value 2 to 4 and chroma 2 to 4. Its texture is sandy clay and clay. The available water capacity is medium (101-150 mm/m). Only one phase was identified and mapped.



Landscape and Soil Profile characteristics of Poglapur (PGP) Series

**4.1.7 Yadgir (YDR) Series:** Yadgir soils are deep (100-150 cm), well drained, have very dark yellowish brown to light olive brown, sodic sandy loam soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation. The Yadgir series has been classified as a member of the coarse-loamy, isohyperthermic family of Fluventic Haplustepts.

The thickness of the soil ranges from 105 to 145 cm. The thickness of A horizon ranges from 6 to 10 cm. Its colour is in 10 YR hue with value 4 and chroma 3. The texture is loamy sand. The thickness of subsurface horizons ranges from 95 to 130 cm. Its colour is in 10 YR and 2.5 Y hue with value 4 to 5 and chroma 3 to 4. Textures is loamy sand to sandy loam and sandy clay loam and are sodic soils. The available water capacity is low (51-100 mm/m). Two phases were identified and mapped.



Landscape and Soil Profile characteristics of Yadgir (YDR) Series

**4.1.8 Mundargi (MDG) Series:** Mundargi soils are deep (100-150 cm), moderately well drained, dark brown to dark yellowish brown, sandy clay loam soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation. The Mundargi series has been classified as a member of the fine-loamy, mixed, isohyperthermic family of Fluventic Haplustepts.

The thickness of the solum ranges from 100 to 149 cm. The thickness of A horizon ranges from 8 to 20 cm. Its colour is in 10 YR hue with value 3 and chroma 1 to 4. The texture ranges from sandy loam to sandy clay loam and sandy clay. The thickness of B horizon ranges from 105 to 140 cm. Its colour is in 10 YR hue with value 3 to 4 and chroma 2 to 4. The texture varies from sandy loam to sandy clay loam and sandy clay. The available water capacity is very high (>200 mm/m). Two phases were identified and mapped.



Landscape and Soil Profile characteristics of Mundargi (MDG) Series



**4.1.9 Baddeppalli (BDP) Series:** Baddeppalli soils are very shallow (<25cm), well drained, have dark brown to dark reddish brown, calcareous sandy clay loam soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation. The Baddeppalli series has been classified as a member of the loamy, mixed (calcareous), isohyperthermic family of Lithic Ustorthents.

The thickness of the soil is less than 25 cm. Its colour is in 7.5 YR and 5 YR hue with value 3 and chroma 2 to 4. The texture varies from sandy clay loam to sandy clay and is calcareous. The available water capacity is very low (<50 mm/m). Two phases were identified and mapped.



Landscape and Soil Profile characteristics of Baddeppalli (BDP) Series

**4.2.10 Kalabelagundi (KBD) Series:** Kalabelagundi soils are moderately deep (75-100 cm), well drained, have dark reddish brown to dark reddish grey and reddish brown, gravelly sandy clay loam soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation. The Kalabelagundi series has been classified as a member of the loamy-skeletal, mixed, isohyperthermic family of Typic Haplustalfs.

The thickness of the solum ranges from 75 to 98 cm. The thickness of A horizon ranges from 10 to 19 cm. Its colour is in hue 5 YR and 7.5 YR with value 3 to 4 and chroma 2 to 4. Its texture varies from loamy sand to sandy loam and sandy clay loam. The thickness of B horizon ranges from 70 to 84 cm. Its colour is in hue 5 YR and 2.5YR with value 3 to 4 and chroma 2 to 4. Its texture is sandy clay loam to sandy clay with gravel content of 35-60 per cent. The available water capacity is very low (<50 mm/m). Two phases were identified and mapped.



Landscape and Soil Profile characteristics of Kalabelagundi (KBD) Series

**4.1.11 Hattikuni (HTK) Series:** Hattikuni soils are shallow (25-50 cm), well drained, have dark brown to dark yellowish brown sandy loam soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping uplands under cultivation. The Hattikuni series has been classified as a member of the mixed, isohyperthermic family of Lithic Ustipsamments.

The thickness of the soil ranges from 36 to 50 cm. The thickness of A horizon ranges from 8 to 12 cm. Its colour is in 10YR and 7.5 YR hue with value 3 to 4 and chroma 4 to 6. The texture varies from loamy sand to sandy loam. The thickness of subsurface horizon ranges from 28 to 42 cm. Its colour is in 10 YR and 7.5 YR hue with value 3 to 4 and chroma 4 to 6. Its texture varies from loamy sand to sand and sandy loam. The available water capacity is very low (<50 mm/m). Only one phase was identified and mapped.



Landscape and Soil Profile characteristics of Hattikuni (HTK) Series

**4.2.12 Kadechoor (KDH) Series:** Kadechoor soils are moderately deep (75-100 cm), moderately well drained, have very dark grayish brown to dark brown, slightly calcareous sandy clay soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping lowlands under cultivation. The Kadechoor series has been classified as a member of the fine, mixed, isohyperthermic family of Typic Haplustepts.

The thickness of the solum ranges from 75 to 98 cm. The thickness of A horizon ranges from 6 to 11 cm. Its colour is in hue 10 YR with value 3 to 4 and chroma 2. Its texture varies from sandy loam to sandy clay. The thickness of B horizon ranges from 73 to 90 cm. Its colour is in hue 10 YR with value 3 to 4 and chroma 2 to 3. The texture is sandy clay loam to sandy clay and is slightly calcareous. The available water capacity is low (51-100 mm/m). Two phases were identified and mapped.



Landscape and Soil Profile characteristics of Kadechoor (KDH) Series

**4.1.9 Sangwar (SGR) Series:** Sangwar soils are very deep (>150 cm), moderately well drained, have very dark gray to dark gray, sodic calcareous cracking clay soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping lowlands under cultivation. The Sangwar series has been classified as a member of the fine, smectic (calcareous), isohyperthermic family of Sodic Haplusterts.

The thickness of the solum is more than 150 cm. The thickness of A horizon ranges from 9 to 20 cm. Its colour is in 10 YR hue with value 3 to 5 and chroma 1 to 2 with sandy clay loam to sandy clay and clay texture. The thickness of B horizon ranges from 157 to 174 cm. Its colour is in 10 YR hue with value 3 to 4 and chroma 1 to 2. Its texture varies from sandy clay to clay and is calcareous. They are sodic with ESP ranging from 29 - 65%. The available water capacity is very high (>200 mm/m). Three phases were identified and mapped.



Landscape and Soil Profile characteristics of Sangwar (SGR) Series

**4.1.6 Kondapur (KDP) Series:** Kondapur soils are very deep (>150 cm), somewhat excessively drained, have strong brown to dark grayish brown and brown sandy soils. They are developed from weathered granite gneiss and occur on very gently to gently sloping lowlands under cultivation. The Kondapur series has been classified as a member of the mixed, isohyperthermic family of Typic Ustipsamments.

The thickness of the solum is more than 150 cm. The thickness of A horizon ranges from 9 to 18 cm. Its colour is in 10 YR hue with value 3 and chroma 2 to 3 with clay texture. The thickness of B horizon ranges from 159 to 162 cm. Its colour is in 10 YR and 7.5YR hue with value 4 to 5 and chroma 2 to 6. Its texture varies from sand to loamy sand and sandy loam and is stratified. The available water capacity is low (51-100 mm/m). Only one phase was identified and mapped.



Landscape and Soil Profile characteristics of Kondapur (KDP) Series

**Table: 4.1 Physical and chemical characteristics of soil series identified in Raisabad Hosalli-1 microwatershed**

**Soil Series:** Badiyala (BDL) **Pedon:** R-5

**Location:** 16°37'10.0"N 77°20'21.5", Gudalagunta village, Balichakra hobli, Yadgir taluk and district

**Analysis at:** NBSS&LUP, Regional Centre, Bengaluru **Classification:** Coarse-loamy, mixed, isohyperthermic Fluventic Haplustepts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-12	Ap	87.13	7.04	5.83	10.03	24.32	23.61	23.51	5.67	<15	ls	6.27	2.44
12-28	Bw1	64.63	13.30	22.07	6.74	13.07	22.30	17.01	5.50	<15	scl	16.34	7.83
28-52	BC	73.11	12.02	14.87	3.93	16.03	26.89	18.41	7.86	<15	sl	12.94	5.47

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO <sub>3</sub>	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP			
	Water	CaCl <sub>2</sub>	M KCl				dS m <sup>-1</sup>	%	%	Ca	Mg					K	Na	Total
										cmol kg <sup>-1</sup>						%	%	
0-12	6.20	-	-	0.074	1.00	0.00	2.80	0.98	0.14	0.01	3.92	4.20	0.72	93	0.20			
12-28	9.04	-	-	0.253	0.80	3.20	-	-	0.16	0.69	-	16.90	0.77	100	4.09			
28-52	9.41	-	-	0.364	1.10	3.60	-	-	0.16	1.39	-	11.10	0.75	100	12.52			

*Contd...*

**Soil Series:** Halagera (HLG) **Pedon:** R-4

**Location:** 16°44'29.3"N 77°13'56.3"E, Halagera village, Yadgir hobli, Yadgir taluk and district

**Analysis at:** NBSS&LUP, Regional Centre, Bengaluru **Classification:** Fine-loamy, mixed (calcareous), isohyperthermic Typic Haplustepts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-8	Ap	81.02	8.42	10.56	10.41	24.08	18.98	19.08	8.47	<15	ls	9.10	4.79
8-22	Bw1	61.00	11.50	27.50	8.29	9.35	21.89	14.35	7.12	<15	scl	16.91	12.28
22-53	Bw2	61.41	13.80	24.79	15.98	15.67	12.62	11.78	5.36	15-35	scl	17.08	11.26

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO <sub>3</sub>	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP
	Water	CaCl <sub>2</sub>	M KCl				Ca	Mg	K	Na	Total				
				dS m <sup>-1</sup>	%	%	cmol kg <sup>-1</sup>					%	%		
0-8	8.49	-	-	0.185	0.30	2.99	-	-	0.24	0.06	-	8.80	0.83	100	0.69
8-22	8.57	-	-	0.116	0.45	4.03	-	-	0.11	0.02	-	19.50	0.71	100	0.12
22-53	8.70	-	-	0.113	0.27	7.67	-	-	0.11	0.05	-	15.50	0.63	100	0.33

*Contd...*

**Soil Series:** Jinkera (JNK) **Pedon:** R-1

**Location:** 16°45'13.5"N 77°10'59.8"E, Varkanahalli village, Yadgir hobli, Yadgir taluk and district

**Analysis at:** NBSS&LUP, Regional Centre, Bengaluru **Classification:** Fine-loamy, mixed, isohyperthermic Typic Haplustepts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-15	Ap	66.84	13.62	19.54	12.15	21.22	11.23	12.56	9.68	10	sl	14.42	7.70
15-38	Bw1	59.08	12.11	28.81	12.53	12.42	17.85	8.77	7.52	20	scl	18.21	12.23
38-50	Bw2	68.21	11.68	20.11	17.90	21.81	10.60	10.80	7.10	10	scl	14.54	8.96

Depth (cm)	pH (1:2.5)			E.C. (1:2.5) dS m <sup>-1</sup>	O.C. %	CaCO <sub>3</sub> %	Exchangeable bases					CEC	CEC/Clay	Base saturation %	ESP %
	Water	CaCl <sub>2</sub>	M KCl				Ca	Mg	K	Na	Total				
0-15	8.42	-	-	0.148	0.70	0.65	-	-	0.15	0.03	-	14.50	0.74	100	0.18
15-38	8.38	-	-	0.226	0.31	2.21	-	-	0.09	0.23	-	21.70	0.75	100	1.05
38-50	8.40	-	-	0.195	0.25	1.17	-	-	0.07	0.19	-	15.90	0.79	100	1.23

*Contd...*

**Soil Series:** Yalleri (YLR) **Pedon:** R-16

**Location:** 16°32'54.3"N 77°22'71.2"E, Duppalli village, Sydhapura hobli, Yadgir taluk and district

**Analysis at:** NBSS&LUP, Regional Centre, Bengaluru **Classification:** Fine, mixed, isohyperthermic Typic Paleustalfs

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-5	Ap	81.69	5.44	12.87	6.10	8.65	33.88	21.57	11.50	-	sl	8.60	3.37
5-34	Bt1	38.78	6.73	54.49	3.38	9.91	12.42	8.93	4.14	-	c	25.33	15.82
34-75	Bt2	40.35	2.90	56.75	12.91	6.83	10.30	7.48	2.82	35-60	c	24.49	16.20

Depth (cm)	pH (1:2.5)			E.C. (1:2.5) dS m <sup>-1</sup>	O.C. %	CaCO <sub>3</sub> %	Exchangeable bases					CEC	CEC/Clay	Base saturation %	ESP %
	Water	CaCl <sub>2</sub>	M KCl				Ca	Mg	K	Na	Total				
0-5	6.91	-	-	0.069	0.70	0.00	5.29	1.37	0.28	0.03	6.96	6.90	0.54	100	0.45
5-34	7.05	-	-	0.053	0.62	0.00	16.43	3.89	0.26	0.09	20.67	21.60	0.40	96	0.42
34-75	7.25	-	-	0.058	0.59	0.00	15.22	3.46	0.25	0.14	19.06	19.90	0.35	96	0.69

*Contd...*



**Soil Series:** Balichakra (BLC) **Pedon:** T1/P2

**Location:** 16°33'25.0"N 77°20'52.3"E, Sowrashtralli village, Sydhapura hobli, Yadgir taluk and district

**Analysis at:** NBSS&LUP, Regional Centre, Bengaluru **Classification:** Fine-loamy, mixed, isohyperthermic Typic Haplustalfs

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-8	Ap	65.46	8.38	26.16	12.51	18.72	18.82	10.44	4.96	-	scl	15.15	8.63
8-19	Bt1	63.48	8.16	28.36	12.80	15.84	17.21	12.49	5.14	-	scl	16.45	8.81
19-40	Bt2	52.64	11.58	35.79	13.19	13.19	14.35	8.23	3.69	-	sc	21.49	10.36
40-75	BC	55.14	10.71	34.15	14.10	14.42	14.63	7.53	4.45	-	scl	17.77	8.99

Depth (cm)	pH (1:2.5)			E.C. (1:2.5) dS m <sup>-1</sup>	O.C. %	CaCO <sub>3</sub> %	Exchangeable bases					CEC	CEC/Clay	Base saturation %	ESP %
	Water	CaCl <sub>2</sub>	M KCl				Ca	Mg	K	Na	Total				
0-8	6.75	-	-	0.19	0.72	0.00	12.18	3.10	0.43	0.22	15.92	16.80	0.64	95	1.31
8-19	7.23	-	-	0.12	0.68	0.84	11.37	2.50	0.23	0.18	14.28	14.77	0.52	97	1.24
19-40	7.13	-	-	0.08	0.50	0.48	13.80	2.82	0.18	0.09	16.89	17.66	0.49	96	0.51
40-75	7.07	-	-	0.07	0.35	0.84	13.00	2.90	0.17	0.10	16.16	17.55	0.51	92	0.57

Contd...

**Soil Series:** Poglapur (PGP) **Pedon:** R-6

**Location:** 16°34'45.2"N 77°10'96.4"E, Anura B village, Sydhapura hobli, Yadgir taluk and district

**Analysis at:** NBSS&LUP, Regional Centre, Bengaluru **Classification:** Fine, mixed, isohyperthermic Rhodic Paleustalfs

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-15	Ap	91.81	4.70	3.49	17.80	30.23	15.57	20.93	7.28	-	s	4.94	2.29
15-50	Bt1	46.83	4.99	48.17	11.92	16.22	8.59	6.77	3.33	10	sc	24.59	17.37
50-90	Bt2	45.81	4.73	49.46	17.10	14.09	6.45	5.16	3.01	15	sc	24.44	16.57
90-125	Bt3	58.92	5.86	35.22	28.51	10.45	10.98	5.49	3.48	15	sc	21.73	10.30

Depth (cm)	pH (1:2.5)			E.C. (1:2.5) dS m <sup>-1</sup>	O.C. %	CaCO <sub>3</sub> %	Exchangeable bases					CEC	CEC/Clay	Base saturation %	ESP %
	Water	CaCl <sub>2</sub>	M KCl				Ca	Mg	K	Na	Total				
0-15	6.83	-	-	0.210	0.76	0.00	1.79	0.88	0.41	0.09	3.16	3.15	0.90	100	2.83
15-50	6.20	-	-	0.105	0.48	0.00	12.27	4.45	0.30	0.39	17.40	17.54	0.36	99	2.22
50-90	6.23	-	-	0.080	0.40	0.00	11.51	3.92	0.28	0.37	16.09	17.33	0.35	93	2.16
90-125	6.49	-	-	0.068	0.20	0.00	11.19	3.62	0.27	0.40	15.49	17.43	0.49	89	2.29

Contd...

**Soil Series:** Yadgir (YDR) **Pedon:** R-5

**Location:** 16°35'43.6"N 77°17'06.4"E, Kanikal village, Balichakra hobli, Yadgir taluk and district

**Analysis at:** NBSS&LUP, Regional Centre, Bengaluru **Classification:** Coarse-loamy, mixed, isohyperthermic Fluventic Haplustepts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-14	Ap	86.59	8.77	4.64	23.19	26.92	14.11	15.22	7.16	-	ls	6.97	2.68
14-43	C1	73.39	11.31	15.30	6.76	20.27	24.87	15.66	5.83	-	sl	12.14	7.22
43-89	C2	80.41	3.75	15.84	8.06	13.47	36.73	15.71	6.43	-	sl	22.84	10.18
89-110	C3	63.55	5.40	31.05	8.10	23.05	19.00	9.87	3.53	15-35	scl	38.46	17.70

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO <sub>3</sub>	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP
	Water	CaCl <sub>2</sub>	M KCl				Ca	Mg	K	Na	Total				
				dS m <sup>-1</sup>	%	%	cmol kg <sup>-1</sup>								
0-14	7.25	-	-	0.114	0.56	0.00	2.29	0.86	0.07	0.03	3.25	3.40	0.73	96	0.31
14-43	9.47	-	-	0.371	0.32	1.30	14.71	4.28	0.38	1.54	20.91	12.70	0.83	165	4.86
43-89	10.30	-	-	0.820	0.16	0.52	1.70	0.98	0.15	6.62	9.45	8.61	0.54	110	30.77
89-110	10.80	-	-	1.440	0.12	0.91	1.02	2.00	0.29	14.43	17.74	16.17	0.52	110	35.688

Contd...

**Soil Series:** Mundargi (MDG) **Pedon:** R-2

**Location:** 16°46'82.4"N 77°04'85.2"E, Thumakura village, Yadgir hobli, Yadgir taluk and district

**Analysis at:** NBSS&LUP, Regional Centre, Bengaluru **Classification:** Fine-loamy, mixed, isohyperthermic Fluventic Haplustepts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-9	Ap	81.23	12.97	5.80	4.84	10.19	14.83	37.94	13.42	<15	ls	11.75	3.31
9-20	A2	76.82	16.19	6.98	4.96	10.12	20.75	27.53	13.46	-	ls	14.52	3.99
20-46	Bw1	42.43	17.43	40.15	2.26	5.59	11.49	14.93	8.16	-	c	34.90	21.14
46-90	Bw2	54.51	16.56	28.93	4.72	5.03	19.92	16.67	8.18	-	scl	36.73	18.88
90-110	Bw3	53.69	11.00	35.30	9.57	9.89	16.23	13.01	4.99	-	sc	38.72	20.53

Depth (cm)	pH (1:2.5)			E.C. (1:2.5) dS m <sup>-1</sup>	O.C. %	CaCO <sub>3</sub> %	Exchangeable bases					CEC	CEC/Clay	Base saturation %	ESP %
	Water	CaCl <sub>2</sub>	M KCl				Ca	Mg	K	Na	Total				
0-9	8.2	-	-	0.399	0.44	0.78	-	-	0.16	0.38	-	4.90	0.84	100	3.08
9-20	8.44	-	-	0.075	0.29	1.82	-	-	0.05	0.35	-	4.90	0.70	100	2.88
20-46	9.39	-	-	0.451	0.32	2.73	-	-	0.12	5.22	-	20.77	0.52	100	10.06
46-90	9.75	-	-	0.616	0.24	3.25	-	-	0.12	5.72	-	16.56	0.57	100	13.82
90-110	9.72	-	-	0.725	0.24	3.64	-	-	0.14	6.84	-	19.76	0.56	100	13.83

Contd...

**Soil Series:** Baddeppalli (BDP) **Pedon:** R-11

**Location:** 16°43'84.4"N 77°14'06.4"E, Halagera village, Yadgir hobli, Yadgir taluk and district

**Analysis at:** NBSS&LUP, Regional Centre, Bengaluru **Classification:** Loamy, mixed (calcareous), isohyperthermic Lithic Ustorthents

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-16	Ap	58.67	17.02	24.31	19.03	13.74	9.62	10.57	5.71	<15	scl	16.19	8.18

Depth (cm)	pH (1:2.5)			E.C. (1:2.5)	O.C.	CaCO <sub>3</sub>	Exchangeable bases					CEC	CEC/Clay	Base saturation	ESP
	Water	CaCl <sub>2</sub>	M KCl				Ca	Mg	K	Na	Total				
				dS m <sup>-1</sup>	%	%	cmol kg <sup>-1</sup>						%	%	
0-16	8.58	-	-	0.262	1.60	7.67	-	-	0.24	0.06	-	18.10	0.74	100	0.35

*Contd...*

**Soil Series:** Kalabelagundi (KBD) **Pedon:** R-13

**Location:** 16°43'78.3"n 77°13'71.4"E, Halagera village, Yadgir hobli, Yadgir taluk and district

**Analysis at:** NBSS&LUP, Regional Centre, Bengaluru **Classification:** Loamy-skeletal, mixed, isohyperthermic Typic Haplustalfs

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-11	Ap	72.35	5.19	22.46	7.19	14.29	19.01	25.28	6.58	15	scl	15.12	8.16
11-35	Bt1	73.20	5.81	20.99	13.66	18.67	16.79	17.62	6.47	20	scl	11.58	7.29
35-64	Bt2	51.68	7.30	41.03	29.41	8.00	4.86	5.62	3.78	40	sc	19.86	14.24
64-89	BC	64.35	3.51	32.15	21.84	12.03	14.87	10.23	5.38	40	scl	16.72	10.36

Depth (cm)	pH (1:2.5)			E.C. (1:2.5) dS m <sup>-1</sup>	O.C. %	CaCO <sub>3</sub> %	Exchangeable bases					CEC	CEC/Clay	Base saturation %	ESP %
	Water	CaCl <sub>2</sub>	M KCl				Ca	Mg	K	Na	Total				
0-11	7.84	-	-	0.604	0.88	0.52	8.69	2.17	0.44	0.49	11.78	11.50	0.51	100	4.27
11-35	5.57	-	-	0.181	0.68	0.00	6.40	1.63	0.18	0.14	8.36	9.10	0.43	92	1.57
35-64	7.42	-	-	0.098	0.44	1.05	15.82	2.34	0.12	0.76	19.04	19.60	0.48	97	3.90
64-89	6.66	-	-	0.165	0.56	0.65	10.45	4.00	0.09	0.43	14.97	15.10	0.47	99	2.86

Contd...

**Soil Series:** Hattikuni (HTK), Pedon: R-7

**Location:** 16°50'46.5"N 77°10'16.4"E, Yaddalli village, Hattikuni hobli, Yadgir taluk and district

**Analysis at:** NBSS&LUP, Regional Centre, Bengaluru **Classification:** Mixed, isohyperthermic, Lithic Ustipsamments

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-12	Ap	90.89	5.62	3.49	8.50	13.46	29.86	29.55	9.51	20	s	7.73	3.16
12-22	A1	89.97	6.53	3.50	7.19	13.48	29.48	29.79	10.03	20	s	8.00	3.05
22-45	A2	87.20	6.43	6.38	11.09	14.42	31.55	7.16	22.98	40	ls	7.67	3.96

Depth (cm)	pH (1:2.5)			E.C. (1:2.5) dS m <sup>-1</sup>	O.C. %	CaCO <sub>3</sub> %	Exchangeable bases					CEC	CEC/Clay	Base saturation %	ESP %
	Water	CaCl <sub>2</sub>	M KCl				Ca	Mg	K	Na	Total				
0-12	6.81	-	-	0.062	0.07	-	2.35	0.50	0.16	0.01	3.02	3.0	0.86	100	0.38
12.0-22	6.80	-	-	0.050	0.21	-	1.67	0.30	0.09	0.01	2.07	2.4	0.69	86.30	0.45
22-45	6.85	-	-	0.044	0.19	-	1.82	0.42	0.10	0.06	2.40	2.6	0.41	92.41	2.17

*Contd...*

**Soil Series:** Kadechoor (KDH) Pedon: T1/P3

**Location:** 16°31'15.0"N 77°20'52.2"E, Kadechoora village, Sydhapura hobli, Yadgir taluk and district

**Analysis at:** NBSS&LUP, Regional Centre, Bengaluru **Classification:** Fine, mixed, isohyperthermic Typic Haplustepts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-18	Ap	75.81	4.05	20.14	7.09	16.85	24.77	19.10	8.01	-	scl	13.70	6.92
18-40	Bw1	57.82	7.95	34.23	2.38	13.52	21.68	14.97	5.27	-	scl	22.10	13.10
40-78	Bw2	50.54	10.54	38.92	1.99	4.51	24.19	12.91	6.95	<15	sc	24.00	14.54

Depth (cm)	pH (1:2.5)			E.C. (1:2.5) dS m <sup>-1</sup>	O.C. %	CaCO <sub>3</sub> %	Exchangeable bases					CEC	CEC/Clay	Base saturation %	ESP %
	Water	CaCl <sub>2</sub>	M KCl				Ca	Mg	K	Na	Total				
0-18	8.22	-	-	0.198	0.84	0.91	-	-	0.41	0.33	-	12.26	0.61	100	2.71
18-40	8.71	-	-	0.163	0.64	1.56	-	-	0.18	0.26	-	20.31	0.59	100	1.27
40-78	8.92	-	-	0.17	0.40	2.90	-	-	0.16	0.37	-	21.41	0.55	100	1.71

*Contd...*



**Soil Series:** Sangwar (SGR) Pedon: R-4

**Location:** 16°32'25.9"N 77°12'52.6"E, Bheemanahalli village, Sydhapura hobli, Yadgir taluka and district

**Analysis at:** NBSS&LUP, Regional Centre, Bengaluru **Classification:** Fine, smectic (calcareous), isohyperthermic Sodic Haplusterts

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-8	Ap	37.30	18.18	44.52	4.91	6.76	12.10	4.80	8.72	-	c	32.36	23.18
8-30	BA	42.04	17.77	40.19	8.28	16.34	7.42	6.13	3.87	-	c	29.89	20.87
30-70	Bss1	33.77	18.63	47.60	5.45	11.66	6.21	6.75	3.70	-	c	37.04	26.13
70-100	Bss2	26.95	18.65	54.40	5.39	9.79	4.95	4.07	2.75	-	c	43.07	32.05
100-150	Bss3	14.35	17.32	68.33	2.69	4.15	2.35	2.69	2.47	-	c	55.74	38.19

Depth (cm)	pH (1:2.5)			E.C. (1:2.5) dS m <sup>-1</sup>	O.C. %	CaCO <sub>3</sub> %	Exchangeable bases					CEC	CEC/Clay	Base saturation %	ESP %
	Water	CaCl <sub>2</sub>	M KCl				Ca	Mg	K	Na	Total				
0-8	8.3	-	-	6.49	1.48	6.69	-	-	1.32	10.09	-	34.77	0.78	100	11.61
8-30	9.09	-	-	2.54	0.64	6.76	-	-	0.75	10.00	-	33.76	0.84	100	11.85
30-70	9.23	-	-	2.6	0.28	6.63	-	-	0.42	11.55	-	38.98	0.82	100	11.86
70-100	9.39	-	-	3.01	0.36	6.89	-	-	0.73	27.73	-	42.46	0.78	100	26.132
100-150	9.28	-	-	4.0	0.24	7.15	-	-	0.80	27.78	-	47.67	0.70	100	23.308

Contd...

**Soil Series:** Kondapura (KDP) **Pedon:** R-2

**Location:** 16°42'03.6"N 77°17'20.7"E, Yaleri village, Balichakra hobli, Yadgir taluk and district

**Analysis at:** NBSS&LUP, Regional Centre, Bengaluru **Classification:** Mixed, isohyperthermic Typic Ustipsamments

Depth (cm)	Horizon	Size class and particle diameter (mm)								Coarse fragments w/w (%)	Texture Class (USDA)	% Moisture	
		Total			Sand							1/3 Bar	15 Bar
		Sand (2.0-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2.0-1.0)	Coarse (1.0-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)				
0-10	Ap	91.15	7.70	1.16	24.04	23.94	23.74	12.58	6.84	20	s	4.59	1.61
10-30	C1	92.15	2.64	5.21	21.45	23.06	26.08	10.47	11.08	20	s	4.43	1.63
30-59	C2	86.75	5.69	7.56	17.49	23.96	27.00	12.94	5.36	20	ls	6.60	2.20
59-118	C3	94.00	2.55	3.45	23.60	27.20	26.70	13.50	3.00	20	s	3.15	0.92
118-157	C4	89.34	7.77	2.89	23.84	24.55	24.65	11.47	4.83	10	s	6.07	1.44

Depth (cm)	pH (1:2.5)			E.C. (1:2.5) dS m <sup>-1</sup>	O.C. %	CaCO <sub>3</sub> %	Exchangeable bases					CEC	CEC/Clay	Base saturation %	ESP %
	Water	CaCl <sub>2</sub>	M KCl				Ca	Mg	K	Na	Total				
0-10	6.55	-	-	0.07	0.48	0.00	1.12	0.31	0.08	0.03	1.54	2.53	2.19	61	1.37
10-30	6.66	-	-	0.03	0.28	0.00	1.38	0.44	0.06	0.05	1.93	2.86	0.55	67	1.90
30-59	6.85	-	-	0.03	0.15	0.00	1.87	0.66	0.06	0.11	2.69	3.85	0.51	70	2.94
59-118	7.06	-	-	0.03	0.11	0.00	0.93	0.31	0.03	0.06	1.33	2.03	0.59	66	3.10
118-157	7.15	-	-	0.03	0.03	0.00	1.51	0.66	0.03	0.08	2.29	2.68	0.92	85	2.93

## INTERPRETATION FOR LAND RESOURCE MANAGEMENT

The most important soil and site characteristics that affect the land use and conservation needs of an area are land capability, land irrigability, soil depth, soil texture, coarse fragments, available water capacity, soil slope, soil erosion, soil reaction etc. These are interpreted from the data base generated through land resource inventory and several thematic maps are generated. These would help in identifying the areas suitable for growing crops and, soil and water conservation measures and structures needed thus helping to maintain good soil health for sustained crop production. The various interpretative and thematic maps generated are described below.

### 5.1 Land Capability Classification

Land capability classification is an interpretative grouping of soil map units (soil phases) mainly based on inherent soil characteristics, external land features and environmental factors that limit the use of land for agriculture, pasture, forestry, or other uses on a sustained basis (IARI, 1971). The land and soil characteristics used to group the land resources in an area into various land capability classes, subclasses and units are

*Soil Characteristics:* Depth, texture, gravelliness, calcareousness.

*Land characteristics:* Slope, erosion, drainage, rock outcrops.

*Climate:* Total rainfall and its distribution, and length of crop growing period.

The land capability classification system is divided into land capability classes, subclasses and units based on the level of information available. Eight land capability classes are recognized. They are

*Class I:* They are very good lands that have no limitations or very few limitations that restrict their use.

*Class II:* They are good lands that have minor limitations and require moderate conservation practices.

*Class III:* They are moderately good lands that have moderate limitations that reduce the choice of crops or that require special conservation practices.

*Class IV:* They are fairly good lands that have very severe limitations that reduce the choice of crops or that require very careful management.

*Class V:* Soils in these lands are not likely to erode, but have other limitations like wetness that are impractical to remove and as such not suitable for agriculture, but suitable for pasture or forestry with minor limitations.

*Class VI:* The lands have severe limitations that make them generally unsuitable for cultivation, but suitable for pasture or forestry with moderate limitations.

*Class VII:* The lands have very severe limitations that make them unsuitable for cultivation, but suitable for pasture or forestry with major limitations.

*Class VIII:* Soil and other miscellaneous areas (rock lands) that have very severe limitations that nearly preclude their use for any crop production, but suitable for wildlife, recreation and installation of wind mills.

The land capability subclasses are recognised based on the dominant limitations observed within a given land capability class. The subclasses are designated by adding a lower case letter like 'e', 'w', 's', or 'c' to the class numeral. The subclass "e" indicates that the main hazard is risk of erosion, "w" indicates drainage or wetness as a limitation for plant growth, "s" indicates shallow soil depth, coarse or heavy textures, calcareousness, salinity/alkalinity or gravelliness and "c" indicates limitation due to climate.

The land capability subclasses have been further subdivided into land capability units based on the kinds of limitations present in each subclass. Ten land capability units are used in grouping the soil map units. They are stony or rocky (0), erosion hazard (slope, erosion) (1), coarse texture (sand, loamy sand, sandy loam) (2), fine texture (cracking clay, silty clay) (3), slowly permeable subsoil (4), coarse underlying material (5), salinity/alkali (6), stagnation, overflow, high ground water table (7), soil depth (8) and fertility problems (9). The capability units thus identified have similar soil and land characteristics that respond similarly to a given level of management. The soils of the microwatershed have been classified upto land capability subclass level.

The 22 soil map units identified in Raisabad Hosalli-1 microwatershed are grouped under 3 land capability classes and 5 land capability subclasses. An entire area of 817 ha (92%) in the microwatershed is suitable for agriculture. About 6 ha (<1%) area is having rock outcrops and about 64 ha (7%) is covered by others (water body & habitation) (Fig. 5.1).

Good cultivable lands (Class II) cover in an area of about 63 per cent and are distributed in the major part of the microwatershed with minor problems of soil, drainage and erosion. Moderately good cultivable lands (Class III) cover an area of about 26 per cent and are distributed in all parts of the microwatershed except north, central and southeast. They have moderate problems of soil and erosion. Fairly good cultivable lands (Class IV) cover an area of about 4 per cent and are distributed in the western, central and northeastern part of the microwatershed with very severe problems of soil and erosion.




Fig. 5.1 Land Capability map of Raisabad Hosalli-1 Microwatershed

## 5.2 Soil Depth

Soil depth refers to the depth of the soil occurring above the parent material or hard rock. The depth of the soil determines the effective rooting depth for plants and in accordance with soil texture, mineralogy and gravel content, the capacity of the soil column to hold water and nutrient availability. Soil depth is one of the most important soil characteristic that is used in differentiating soils into different soil series. The soil depth classes used in identifying soils in the field are very shallow (<25 cm), shallow (25-50 cm), moderately shallow (50-75 cm), moderately deep (75-100 cm), deep (100-150 cm) and very deep (>150 cm). They were used to classify the soils into different depth classes and a soil depth map was generated. The area extent and their geographical distribution in the microwatershed is given in Fig. 5.2.

Very shallow (<25 cm) soils occur in an area of 22 ha (2%) and are distributed in the northwestern and northeastern part of the microwatershed. Shallow (25-50 cm) soils occur in an area of 109 ha (12%) and are distributed in the northeastern, eastern and southern part of the microwatershed. Moderately shallow (50-75 cm) soils occur in an area of 90 ha (10%) and are distributed in the northwestern, western, southwestern and northeastern part of the microwatershed. Moderately deep (75-100 cm) soils occur in an area of 156 ha (18%) and are distributed in the northern, northwestern, western and southwestern part of the microwatershed. Deep (100-150 cm) soils cover an area of 381 ha (43%) and are distributed in the major part of the microwatershed. Very deep (>150

cm) soils cover an area of 59 ha (7%) and are distributed in the western and northern part of the microwatershed.

The most productive lands covering 440 ha (50%) with respect to soil rooting depth where all climatically adapted annual and perennial crops can be grown are deep to very deep (100 - >150 cm depth) soils. The problem soils occupy an area of 131 ha (15%) where only short duration crops can be grown occasionally and the probability of crop failure is very high.




Fig. 5.2 Soil Depth map of Raisabad Hosalli-1 Microwatershed

### 5.3 Surface Soil Texture

Texture is an expression to indicate the coarseness or fineness of the soil as determined by the relative proportion of primary particles of sand, silt and clay. It has a direct bearing on the structure, porosity, adhesion and consistence. The surface layer of a soil to a depth of about 25 cm is the layer that is most used by crops and plants. The surface soil textural class provides a guide to understanding soil-water retention and availability, nutrient holding capacity, infiltration, workability, drainage, physical and chemical behaviour, microbial activity and crop suitability. The textural classes used for LRI were used to classify and a surface soil texture map was generated. The area extent and their geographical distribution in the microwatershed is shown in Figure 5.3.

An area of about 31 ha (3%) is sandy at the surface and are distributed in the eastern part of the microwatershed. Maximum area of about 462 ha (52%) of the microwatershed has loamy soils at the surface and are distributed in the major part of the

microwatershed. An area of 325 ha (37%) of the microwatershed has soils that are clayey and are distributed in the southern, southwestern, northern and northeastern part of the microwatershed. Loamy and clayey soils have high potential for soil-water retention and availability, and nutrient retention and availability, but clay soils have more problems of drainage, infiltration, workability and other physical problems. Sandy soils are problematic where tuber crops can be grown and require frequent irrigation and poor nutrient status.




Fig. 5.3 Surface Soil Texture map of Raisabad Hosalli-1 Microwatershed

#### 5.4 Soil Gravelliness

Gravel is the term used for describing coarse fragments between 2 mm and 7.5 cm diameter and stones for those between 7.5 cm and 25 cm. The presence of gravel and stones in soil reduces the volume of soil responsible for moisture and nutrient storage, drainage, infiltration and runoff, and hinders plant growth by impeding root growth and seedling emergence, intercultural operations and farm mechanization. The gravelliness classes used in LRI were used to classify the soils and using these classes a gravelliness map was generated. The area extent and their geographic distribution in the microwatershed is shown in Figure 5.4.

Non gravelly (<15%) soils cover a maximum area of 567 ha (64%) and distributed in the major part of the microwatershed. These are the most productive soils, where all climatically adapted short and long duration crops can be grown. Gravelly (15-35%) soils occur in an area of 249 ha (28%) and distributed in the central, northern, northeastern,

eastern and southeastern part of the microwatershed; these lands are low in moisture holding capacity and hence growing of short duration crops is ideal with best management practice.




Fig. 5.4 Soil Gravelliness map of Raisabad Hosalli-1 Microwatershed

### 5.5 Available Water Capacity

The soil available water capacity (AWC) is estimated based on the ability of the soil column to retain water between the tensions of 0.33 and 15 bar in a depth of 100 cm or the entire solum if the soil is shallower. The AWC of the soils (soil series) as estimated by considering the soil texture, mineralogy, soil depth and gravel content (Sehgal *et al.*, 1990) and accordingly the soil map units were grouped into five AWC classes *viz*, very low (<50 mm/m), low (50-100 mm/m), medium (100-150 mm/m), high (150-200 mm/m) and very high (>200 mm/m) and using these values, an AWC map was generated. The area extent and their geographic distribution of different AWC classes in the microwatershed is given in Figure 5.5.

An area of about 249 ha (28%) in the microwatershed has soils that are very low (<50 mm/m) in available water capacity and is distributed in all parts of the microwatershed except north, central and southeast. Maximum area of about 349 ha (39%) is low (51-100 mm/m) in available water capacity and are distributed in the major part of the microwatershed. Very high (>200 mm/m) in 219 ha (25%) and are distributed in the western, northern, northeastern and southeastern part of the microwatershed.



Maximum area of about 598 ha (67%) in the microwatershed has soils that are problematic with regard to available water capacity. Here, only short duration crops can be grown and probability of the crop failure is very high. These areas are best put to other alternative uses. An area of 219 ha (25%) are potential with regard to AWC where all climatically adapted annual and perennial crops can be grown.




Fig. 5.5 Soil Available Water Capacity map of Raisabad Hosalli-1 Microwatershed

### 5.6 Soil Slope

Soil slope refers to the inclination of the surface of the land. It is defined by gradient, shape and length, and is an integral feature of any soil as a natural body. Slope is considered important in soil genesis, land use and land development. The length and gradient of slope influences the rate of runoff, infiltration, erosion and deposition. The soil map units were grouped into single slope class and a slope map was generated showing the area extent and their geographic distribution in the microwatershed (Fig. 5.6).

An area of about <1 ha (<1%) of the microwatershed falls under nearly level (0-1% slope) and 816 ha (92%) under very gently sloping (1-3% slope) lands, thus these areas have high potential in respect of soil slopes. In these areas, all climatically adapted annual and perennial crops can be grown without much soil and water conservation and other land development measures.




Fig. 5.6 Soil Slope map of Raisabad Hosalli-1 Microwatershed

## 5.7 Soil Erosion

Soil erosion refers to the wearing away of the earth's surface by the forces of water, wind and ice involving detachment and transport of soil by raindrop impact. It is used for accelerated soil erosion resulting from disturbance of the natural landscape by burning, excessive grazing and indiscriminate felling of forest trees and tillage, all usually by man. The erosion classes showing an estimate of the current erosion status as judged from field observations in the form of rills, gullies or a carpet of gravel on the surface are recorded. Four erosion classes, viz, slight erosion (e1), moderate erosion (e2), severe erosion (e3) and very severe erosion (e4) are recognized. The soil map units were grouped into different erosion classes and a soil erosion map generated. The area extent and their spatial distribution in the microwatershed is given in Figure 5.7.

Soils that are slightly eroded (e1 class) cover an area of <1 ha (<1%) and are distributed in the western part of the microwatershed. Moderately eroded (e2 class) soils cover a maximum area of 816 ha (92%) and are distributed in the major part of the microwatershed. Severely eroded soils (e3 class) cover an area of 1 ha (<1%) and are distributed in the southwestern part of the microwatershed.

An area of about 817 ha of the microwatershed is problematic because of moderate and severe erosion. For these areas, taking up of soil and water conservation and other land development measures are needed.




Fig. 5.7 Soil Erosion map of Raisabad Hosalli-1 Microwatershed



## **FERTILITY STATUS**

Soil fertility plays an important role in increasing crop yield. The adoption of high yielding varieties that require high amounts of nutrients has resulted in deficiency symptoms in crops and plants due to imbalanced fertilization and poor inherent fertility status as these areas are characterised by low rainfall and high temperatures. Hence, it is necessary to know the fertility (macro and micro nutrients) status of the soils of the watersheds for assessing the kind and amount of fertilizers required for each of the crop intended to be grown. For this purpose, the surface soil samples collected from the grid points (one soil sample at every 320 m interval) all over the microwatershed through land resource inventory in the year 2018 were analysed for pH, EC, organic carbon, available phosphorus and potassium, and for micronutrients like zinc, boron, copper, iron and manganese, and secondary nutrient sulphur.

Soil fertility data generated has been assessed and individual maps for all the nutrients for the microwatershed have been generated using Kriging method under GIS. The village/survey number wise fertility data for the microwatershed is given in Appendix-II.

### **6.1 Soil Reaction (pH)**

The soil analysis of the Raisabad Hosalli-1 microwatershed for soil reaction (pH) showed that an area of 34 ha (4%) is slightly acid (pH 6.0-6.5) and are distributed in the northeastern part of the microwatershed. Maximum area of 453 ha (51%) is neutral (pH 6.5-7.3) and are distributed in the major part of the microwatershed. Slightly alkaline (pH 7.3-7.8) occur in an area of 188 ha (21%) and are distributed in the southwestern, central and northern part of the microwatershed. Moderately alkaline (pH 7.8-8.4) occur in an area of 135 ha (15%) and are distributed in the western and northwestern part of the microwatershed. Strongly alkaline (pH 8.4-9.0) occur in an area of 7 ha (<1%) and are distributed in the western part of the microwatershed (Fig. 6.1). Thus major soils (453 ha) in the microwatershed are under neutral followed by alkaline soils (330 ha) and 34 ha under slightly acid.

### **6.2 Electrical Conductivity (EC)**

The Electrical Conductivity in entire area of the microwatershed is non saline (<2 dS/m) (Fig. 6.2) and as such the soils are nonsaline.

### **6.3 Organic Carbon**

The soils organic carbon content (an index of available Nitrogen) of the microwatershed is high (>0.75) in an area of about 212 ha (24%) and are distributed in the western and northeastern part of the microwatershed. Medium (0.5-0.75%) covering a maximum area of about 605 ha (68%) and are distributed in the major part of the microwatershed (Fig.6.3).




Fig.6.1 Soil Reaction (pH) map of Raisabad Hosalli-1 Microwatershed




Fig.6.2 Electrical Conductivity (EC) map of Raisabad Hosalli-1 Microwatershed




Fig.6.3 Soil Organic Carbon map of Raisabad Hosalli-1 Microwatershed

#### 6.4 Available Phosphorus

Available phosphorus content is low (<23 kg/ha) in an area of about 179 ha (20%) and distributed in the eastern and western part of the microwatershed. The available phosphorus content is medium (23-57 kg/ha) in a maximum area of about 625 ha (70%) and occur in the major part of the microwatershed and available phosphorus content is high (>57 kg/ha) in an area of 13 ha (1%) and distributed in the northern and northeastern part of the microwatershed (Fig. 6.4).

#### 6.5 Available Potassium

Available potassium content is high (>337 kg/ha) in an area of about 16 ha (2%) and are distributed in the central part of the microwatershed. Available potassium content is medium (145-337 kg/ha) in an area of about 726 ha (82%) and are distributed in the major part of the microwatershed (Fig. 6.5). Low (<145 kg/ha) in available potassium content occur in an area of 75 ha (8%) and are distributed in the northeastern, southern and southwestern part of the microwatershed.

#### 6.6 Available Sulphur

An area of about 123 ha (14%) is low (<10 ppm) in available sulphur content and are distributed in the eastern and northeastern part of the microwatershed. Medium (10-20 ppm) in a maximum area of about 693 ha (78%) and is distributed in the major part of the microwatershed (Fig. 6.6).

### 6.7 Available Boron

Available boron content is medium (0.5-1.0 ppm) in an area of 256 ha (29%) and are distributed in the western, southern, southwestern and northeastern part of the microwatershed. Maximum area of about 561 ha (63%) is low (<0.5 ppm) in available boron and are distributed in the major part of the microwatershed (Fig. 6.7).

### 6.8 Available Iron

Available iron content is sufficient (>4.5 ppm) in the entire area of the microwatershed (Fig 6.8).

### 6.9 Available Manganese

Available manganese content is sufficient (>1.0 ppm) in the entire area of the microwatershed (Fig 6.9).

### 6.10 Available Copper

Available copper content is sufficient (>0.2 ppm) in the entire area of the microwatershed (Fig 6.10).




Fig.6.4 Soil Available Phosphorus map of Raisabad Hosalli-1 Microwatershed






Fig.6.5 Soil Available Potassium map of Raisabad Hosalli-1 Microwatershed




Fig.6.6 Soil Available Sulphur map of Raisabad Hosalli-1 Microwatershed




Fig.6.7 Soil Available Boron map of Raisabad Hosalli-1 Microwatershed




Fig.6.8 Soil Available Iron map of Raisabad Hosalli-1 Microwatershed




Fig.6.9 Soil Available Manganese map of Raisabad Hosalli-1 Microwatershed




Fig.6.10 Soil Available Copper map of Raisabad Hosalli-1 Microwatershed

### 6.11 Available Zinc

Available Zinc content is deficient (<0.6 ppm) in the entire area of the microwatershed (Fig.6.11).




Fig.6.11 Soil Available Zinc map of Raisabad Hosalli-1 Microwatershed

## LAND SUITABILITY FOR MAJOR CROPS

The soil and land resource units (soil phases) of Raisabad Hosalli-1 microwatershed were assessed for their suitability for growing food, fodder, fibre and other horticulture crops by following the procedure as outlined in FAO, 1976 and 1983. Crop requirements were developed for each of the crop from the available research data and also by referring to Naidu *et. al.* (2006) and Natarajan *et. al* (2015). The soil and land characteristics were matched with the crop requirement to arrive at the crop suitability. The soil and land characteristics table (table 7.1) and crop requirement tables (tables 7.2 to 7.30) are given at the end. In FAO land suitability classification, two orders are recognized. Order S-Suitable and Order N-Not suitable. The orders have classes, subclasses and units. Order-S has three classes, Class S1-Highly Suitable, Class S2-Moderately Suitable and Class S3- Marginally Suitable. Order N has two classes, N1- Currently not Suitable and N2- Permanently not Suitable. There are no subclasses within the Class S1 as they will have very minor or no limitations for crop growth. Classes S2, S3, N1 and N2 are divided into subclasses based on the kinds of limitations encountered. The limitations that affect crop production are ‘c’ for erratic rainfall and its distribution and length of growing period (LGP), ‘e’ for erosion hazard, ‘r’ for rooting condition, ‘t’ for lighter or heavy texture, ‘g’ for gravelliness or stoniness, ‘n’ for nutrient availability, ‘l’ for topography, ‘m’ for moisture availability, ‘w’ for drainage, ‘s’ for sodium and ‘z’ for calcareousness. These limitations are indicated as lower case letters to the Class symbol. For example, moderately suitable lands with the limitations of soil depth and erosion are designated as S2re. For the microwatershed, the soil mapping units were evaluated and classified up to subclass level.

Using the above criteria, the soil map units of the microwatershed were evaluated and land suitability maps for 29 major agricultural and horticultural crops were generated. The detailed information on the kind of suitability of each of the soil phase for the crops assessed are given village/ survey number wise for the microwatershed in Appendix-III.

### 7.1 Land Suitability for Sorghum (*Sorghum bicolor*)

Sorghum is one of the major crop grown in Karnataka in an area of 10.47 lakh ha in Bijapur, Gulbarga, Raichur, Bidar, Belgaum, Dharwad, Bellary, Chitradurga, Mysore and Tumakuru districts. The crop requirements for growing sorghum (Table 7.2) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing sorghum was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.1.

Highly suitable (Class S1) lands for growing sorghum occur in an area of 243 ha (27%) and are distributed in the western, northern, northeastern and southeastern part of the microwatershed. An area of about 106 ha (12%) is moderately suitable (Class S2) for

growing sorghum and are distributed in the southwestern, northwestern, northern and northeastern part of the microwatershed. They have minor limitations of rooting depth, topography, calcareousness and gravelliness. An area of about 435 ha (49%) is marginally suitable (Class S3) for growing sorghum and is distributed in the major part of the microwatershed with moderate limitations rooting depth, texture, calcareousness and gravelliness. An area of about 35 ha (4%) is currently not suitable (Class N1) and are distributed in the northeastern part of the microwatershed with severe limitations of rooting depth and texture.




Fig. 7.1 Land Suitability map of Sorghum

## 7.2 Land Suitability for Maize (*Zea mays*)

Maize is one of the most important food crop grown in an area of 13.37 lakh ha in almost all the districts of the State. The crop requirements for growing maize (Table 7.3) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing maize was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.2.

Highly suitable (Class S1) lands for growing maize occur in an area of <1 ha (<1%) and are distributed in the northeastern part of the microwatershed. Moderately suitable (Class S2) lands occur in an area of 348 ha (39%) and are distributed in all parts of the microwatershed except central, east and south. They have minor limitations of rooting depth, topography, calcareousness, texture, gravelliness and drainage. Marginally suitable

lands (Class S3) for growing maize occupy a maximum area of 435 ha (49%) and occur in the major part of the microwatershed. They have moderate limitations of rooting depth, texture, calcareousness and gravelliness. An area of about 35 ha (4%) is currently not suitable (Class N1) and are distributed in the northeastern part of the microwatershed with severe limitations of rooting depth and texture.




Fig. 7.2 Land Suitability map of Maize

### 7.3 Land Suitability for Bajra (*Pennisetum glaucum*)

Bajra is one of the most important millet crop grown in an area of 2.34 lakh ha in the northern districts of Karnataka state. The crop requirements for growing bajra (Table 7.4) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing bajra was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.3.

Highly suitable (Class S1) lands for growing bajra occur in an area of <1 ha (<1%) and are distributed in the northeastern part of the microwatershed. Maximum area of about 674 ha (76%) is moderately suitable (Class S2) for growing bajra and are distributed in the major part of the microwatershed. They have minor limitations of texture, rooting depth, gravelliness, topography and calcareousness. Marginally suitable lands (Class S3) occupy an area of 122 ha (14%) and are distributed in the southern, eastern and northeastern part of the microwatershed. They have moderate limitations of rooting depth and texture. An area of about 22 ha (2%) is currently not suitable (Class

N1) and are distributed in the northeastern part of the microwatershed with severe limitation of rooting depth.




Fig. 7.3 Land Suitability map of Bajra

#### 7.4 Land Suitability for Groundnut (*Arachis hypogaea*)

Groundnut is one of the major oilseed crop grown in an area of 6.54 lakh ha in Karnataka in most of the districts either as rainfed or irrigated crop. The crop requirements for growing groundnut (Table 7.5) were matched with the soil-site characteristics (Table 7.1) of the soils of the microwatershed and a land suitability map for growing groundnut was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed are given in Figure 7.4.

There are no highly suitable (Class S1) lands available for growing groundnut in the microwatershed. Moderately suitable (Class S2) lands occur in an area of 367 ha (41%) and are distributed in all parts of the microwatershed. They have minor limitations of rooting depth, texture, topography, gravelliness and calcareousness. Marginally suitable lands (Class S3) for growing groundnut occupy a maximum area of about 429 ha (48%) with moderate limitations of texture, drainage and rooting depth. They are distributed in the major part of the microwatershed except central and southeast. An area of about 22 ha (2%) is currently not suitable (Class N1) and are distributed in the northeastern part of the microwatershed with severe limitation of rooting depth.






Fig. 7.4 Land Suitability map of Groundnut

### 7.5 Land Suitability for Sunflower (*Helianthus annus*)

Sunflower is one of the most important oilseed crop grown in an area of 4.1 lakh ha in the State in all the districts. The crop requirements for growing sunflower (Table 7.6) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing sunflower was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.5.

Highly suitable (Class S1) lands for growing sunflower occupy an area of 225 ha (25%) and are distributed in the western, northern, northeastern and southwestern part of the microwatershed. An area of about 33 ha (4%) is moderately suitable (Class S2) for sunflower and is distributed in the northern part of the microwatershed. They have minor limitations of rooting depth, drainage and calcareousness. An area of about 417 ha (47%) is marginally suitable (Class S3) and is distributed in all parts of the microwatershed except north. They have moderate limitations of rooting depth, texture, topography and calcareousness. Currently not suitable (Class N1) lands occur in an area of 144 ha (16%) and are distributed in the southern, eastern and northeastern part of the microwatershed with severe limitations of rooting depth and texture.




Fig. 7.5 Land Suitability map of Sunflower

### 7.6 Land suitability criteria for Red gram (*Cajanus Cajan*)

Redgram is one of the most important pulse crop grown in an area of 7.28 lakh ha in almost all the districts of the State. The crop requirements for growing red gram (Table 7.7) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing redgram was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.6.

No highly suitable (Class S1) lands are available for growing redgram in the microwatershed. An area of about 407 ha (46%) is moderately suitable (Class S2) for growing redgram and are distributed in all parts of the microwatershed except southwest and northwest. They have minor limitations of rooting depth, texture, drainage, gravelliness and calcareousness. Marginally suitable lands (Class S3) for growing redgram occupy an area of about 346 ha (39%) and occur in all parts part of the microwatershed except central and southeast. They have moderate limitations of rooting depth, gravelliness, texture, topography and drainage. Currently not suitable (Class N1) lands occur in an area of 66 ha (7%) and are distributed in the eastern and northeastern part of the microwatershed with severe limitations of rooting depth and texture.




Fig. 7.6 Land Suitability map of Redgram

### 7.7 Land Suitability for Bengal gram (*Cicer aerativum*)

Bengal gram is one of the most important pulse crop grown in about 9.39 lakh ha area in Bijapur, Raichur, Kalaburgi, Dharwad, Belgaum and Bellary districts. The crop requirements for growing Bengal gram (Table 7.8) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing Bengal gram was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.7.

Highly suitable (Class S1) lands for growing bengalgram occupy an area of 243 ha (27%) and are distributed in the western, northern, northeastern and southwestern part of the microwatershed. An area of about 106 ha (12%) is moderately suitable (Class S2) for growing bengalgram and are distributed in the southwestern, northwestern, northern and northeastern part of the microwatershed. They have minor limitations of rooting depth, texture, calcareousness, topography and gravelliness. Marginally suitable lands (Class S3) for growing bengalgram occupy an area of about 196 ha (22%) and occur in the southern, southwestern, northwestern and northeastern part of the microwatershed. They have moderate limitations of rooting depth and gravelliness. Currently not suitable (Class N1) lands occur in an area of 274 ha (31%) and are distributed in the central, southern, southeastern, eastern and northeastern part of the microwatershed with severe limitations of texture, rooting depth and calcareousness.




Fig. 7.7 Land Suitability map of Bengal gram

### 7.8 Land Suitability for Cotton (*Gossypium hirsutum*)

Cotton is one of the most important fibre crop grown in the State in about 8.75 lakh ha area in Raichur, Dharwad, Belgaum, Kalaburgi, Bijapur, Bidar, Bellary, Chitradurga and Chamarajnagar districts. The crop requirements for growing cotton (Table 7.9) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing cotton was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.8.

Highly suitable (Class S1) lands for growing cotton occur in an area of 219 ha (25%) and are distributed in the western, northern, northeastern and southwestern part of the microwatershed. Moderately suitable (Class S2) lands are found to occur in an area of 127 ha (14%). These soils have minor limitations of rooting depth, calcareousness and gravelliness. They are distributed in the southwestern, northwestern, northern and northeastern part of the microwatershed. Marginally suitable (Class S3) lands for cotton occur in an area of 197 ha (22%) with moderate limitations of rooting depth, topography and gravelliness. They are distributed in the southern, southwestern, northwestern and northeastern part the microwatershed. Currently not suitable (Class N1) lands occur in an area of 274 ha (31%) and are distributed in the central, southern, southeastern, eastern and northeastern part of the microwatershed with severe limitations of texture, rooting depth and calcareousness.




Fig. 7.8 Land Suitability map of Cotton

### 7.9 Land Suitability for Chilli (*Capsicum annuum*)

Chilli is one of the most important spice crop grown in about 0.42 lakh ha in Karnataka State. The crop requirements for growing chilli (Table 7.10) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing chilli was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.9.

There are no highly (Class S1) suitable lands available for growing chilli crop in the microwatershed. Maximum area of about 556 ha (63%) is moderately suitable (Class S2) for growing chilli and are distributed in the major part of the microwatershed. They have minor limitations of texture, drainage, calcareousness, gravelliness, topography and rooting depth. Marginally suitable lands (Class S3) occupy an area of 227 ha (26%) and are distributed in the southern, southwestern, northwestern, northeastern and eastern part of the microwatershed. They have moderate limitations of rooting depth, gravelliness and texture. Currently not suitable (Class N1) lands occur in an area of 35 ha (4%) and are distributed in the northeastern part of the microwatershed with severe limitations of texture and rooting depth.




Fig 7.9 Land Suitability map of Chilli

### 7.10 Land Suitability for Tomato (*Lycopersicon esculentum*)

Tomato is one of the most important vegetable crop grown in about 0.61 lakh ha covering almost all the district of the state. The crop requirements for growing tomato (Table 7.11) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing tomato was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.10.

Highly suitable (Class S1) lands for growing tomato occur in an area of <1 ha (<1%) and are distributed in the northeastern part of the microwatershed. An area of 408 ha (46%) is moderately suitable (Class S2) and is distributed in all parts of the microwatershed except western. They have minor limitations of texture, drainage, calcareousness, gravelliness, topography and rooting depth. An area of 374 ha (42%) is marginally suitable for tomato (Class S3) and is distributed in all parts of the microwatershed except central, southeast and northeast. They have moderate limitations of rooting depth and texture, drainage and gravelliness. Currently not suitable (Class N1) lands occur in an area of 35 ha (4%) and are distributed in the northeastern part of the microwatershed with severe limitations of texture and rooting depth.




Fig 7.10 Land Suitability map of Tomato

### 7.11 Land Suitability for Brinjal (*Solanum melongena*)

Brinjal is one of the most important vegetable crop grown in the state. The crop requirements for growing brinjal (Table 7.12) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing brinjal was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.11.

Highly suitable (Class S1) lands for growing brinjal occur in an area of 217 ha (24%) and are distributed in the western, northern, northeastern and southwestern part of the microwatershed. Maximum area of about 456 ha (51%) is moderately suitable (Class S2) for brinjal and is distributed in the major part of the microwatershed with minor limitations of texture, gravelliness and rooting depth. An area of 109 ha (12%) is marginally suitable (Class S3) and is distributed in the southern, eastern and northeastern part of the microwatershed. They have moderate limitations of rooting depth and texture. Currently not suitable (Class N1) lands occur in an area of 35 ha (4%) and are distributed in the northeastern part of the microwatershed with severe limitations of texture and rooting depth.




Fig 7.11 Land Suitability map of Brinjal

## 7.12 Land Suitability for Onion (*Allium cepa L.*)

Onion is one of the most important vegetable crop grown in the state. The crop requirements for growing onion (Table 7.13) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing onion was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.12.

Highly (Class S1) suitable lands for growing onion occur in an area of 89 ha (13%) and are distributed in the northern, eastern and southeastern part of the microwatershed. An area of about 216 ha (32%) is moderately suitable (Class S2) for onion and is distributed in all parts of the microwatershed except north. They have minor limitations of texture and rooting depth. An area of 200 ha (30%) is marginally suitable (Class S3) and is distributed in all parts of the microwatershed except north and central. They have moderate limitations of rooting depth and texture.






Fig 7.12 Land Suitability map of Onion

### 7.13 Land Suitability for Bhendi (*Abelmoschus esculentus*)

Bhendi is one of the most important vegetable crop grown in the state. The crop requirements for growing bhendi (Table 7.14) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing bhendi was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.13.

Highly suitable (Class S1) lands for growing bhendi occur in an area of 217 ha (24%) and are distributed in the western, northern, northeastern and southwestern part of the microwatershed. An area of about 456 ha (51%) is moderately suitable (Class S2) for bhendi and is distributed in the major part of the microwatershed. They have minor limitations of texture, gravelliness and rooting depth. An area of 109 ha (12%) is marginally suitable (Class S3) and is distributed in the southern, eastern and northeastern part of the microwatershed. They have moderate limitation of rooting depth. Currently not suitable (Class N1) lands occur in an area of 35 ha (4%) and are distributed in the northeastern part of the microwatershed with severe limitations of texture and rooting depth.




Fig 7.13 Land Suitability map of Bhendi

#### 7.14 Land Suitability for Drumstick (*Moringa oleifera*)

Drumstick is one of the most important vegetable crop grown in about 2403 ha in the state. The crop requirements for growing drumstick (Table 7.15) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing drumstick was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.14.

There are no highly (Class S1) suitable lands available for growing drumstick in the microwatershed. An area of about 584 ha (66%) is moderately suitable (Class S2) for drumstick and is distributed in the major part of the microwatershed. They have minor limitations of rooting depth, texture, calcareousness, gravelliness and drainage. Marginally suitable lands (Class S3) for growing drumstick occupy an area of about 91 ha (10%) and occur in the southwestern, northwestern and northeastern part of the microwatershed. They have moderate limitations of rooting depth, topography and gravelliness. Currently not suitable (Class N1) lands for growing drumstick occur in an area of 144 ha (16%) and are distributed in the southern, eastern and northeastern part of the microwatershed. They have severe limitations of rooting depth and texture.




Fig 7.14 Land Suitability map of Drumstick

### 7.15 Land suitability for Mango (*Mangifera indica*)

Mango is one of the most important fruit crop grown in an area of 1.73 lakh ha in almost all the districts of the State. The crop requirements (Table 7.16) for growing mango were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing mango was generated. The area extent and their geographic distribution of different suitability subclasses in the microwatershed is given in Figure 7.15.

There are no highly suitable (Class S1) lands available for growing mango in the microwatershed. An area of 160 ha (18%) is moderately suitable (Class S2) for growing mango with minor limitation of rooting depth. They are distributed in the northern, northeastern and southeastern part of the microwatershed. An area of 423 ha (48%) is marginally suitable (Class S3) for growing mango with moderate limitations of texture, calcareousness, drainage, gravelliness and rooting depth. They are distributed in all parts of the microwatershed. An area of about 235 ha (26%) is currently not suitable (Class N1) for growing mango and are distributed in the northeastern, northwestern, southeastern, southwestern and eastern part of the microwatershed. They have severe limitations of rooting depth, texture, topography and calcareousness.




Fig. 7.15 Land Suitability map of Mango

### 7.16 Land suitability for Guava (*Psidium guajava*)

Guava is one of the most important fruit crop grown in an area of 6558 ha in almost all the districts of the State. The crop requirements (Table 7.17) for growing guava were matched with the soil-site characteristics (7.1) and a land suitability map for growing guava was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.16.

No highly (Class S1) suitable lands available for growing guava in the microwatershed. Moderately suitable (Class S2) lands occur in an area of 223 ha (25%) and are distributed in the central, northern, southern, southeastern and northeastern part of the microwatershed with minor limitations of rooting depth, calcareousness and texture. Marginally suitable (Class S3) lands cover an area of about 451 ha (51%) and are distributed in the major part of the microwatershed. They have moderate limitations of rooting depth, texture, calcareousness, topography, gravelliness and drainage. An area of about 144 ha (16%) is currently not suitable (Class N1) for growing guava and occur in the southern, southeastern and northeastern part of the microwatershed. They have severe limitations of rooting depth and texture.




Fig. 7.16 Land Suitability map of Guava

### 7.17 Land suitability for Sapota (*Manilkara zapota*)

Sapota is one of the most important fruit crop grown in an area of 29373 ha in almost all the districts of the State. The crop requirements (Table 7.18) for growing sapota were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing sapota was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.17.

No highly (Class S1) suitable lands available for growing sapota in the microwatershed. Moderately suitable (Class S2) lands occur in an area of 223 ha (25%) and are distributed in the central, northern, southern, southeastern and northeastern part of the microwatershed with minor limitations of rooting depth, calcareousness and texture. Marginally suitable (Class S3) lands cover an area of about 451 ha (51%) and are distributed in the major part of the microwatershed. They have moderate limitations of rooting depth, texture, calcareousness, topography, gravelliness and drainage. An area of about 144 ha (16%) is currently not suitable (Class N1) for growing sapota and occur in the southern, southeastern and northeastern part of the microwatershed. They have severe limitations of rooting depth and texture.




Fig. 7.17 Land Suitability map of Sapota

### 7.18 Land Suitability for Pomegranate (*Punica granatum*)

Pomegranate is one of the most important fruit crop commercially grown in about 18488 ha in Karnataka, mainly in Bijapur, Bagalkot, Koppal, Gadag and Chitradurga districts. The crop requirements for growing pomegranate (Table 7.19) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing pomegranate was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.18.

No highly (Class S1) suitable lands available for growing pomegranate in the microwatershed. Moderately suitable (Class S2) lands occur in an area of 465 ha (52%) and are distributed in the major part of the microwatershed with minor limitations of rooting depth, texture, calcareousness, gravelliness and drainage. An area of about 209 ha (23%) is marginally suitable (Class S3) for growing pomegranate and are distributed in the southwestern, northwestern and northeastern part of the microwatershed. They have moderate limitations of rooting depth, gravelliness, calcareousness and topography. An area of about 144 ha (16%) is currently not suitable (Class N1) for growing pomegranate and occur in the southern, southeastern and northeastern part of the microwatershed. They have severe limitations of rooting depth and texture.




Fig 7.18 Land Suitability map of Pomegranate

### 7.19 Land Suitability for Musambi (*Citrus limetta*)

Musambi is one of the important fruit crop grown in an area of 3446 ha in almost all the districts of the State. The crop requirements for growing musambi (Table 7.20) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing musambi was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.19.

Highly suitable (Class S1) lands for growing musambi occur in an area of 303 ha (34%) and are distributed in the central southern, southeastern eastern, northern and northeastern part of the microwatershed. An area of about 162 ha (18%) is moderately suitable (Class S2) for growing musambi and are distributed in the northern, western and northeastern part of the microwatershed. They have minor limitations of drainage and rooting depth and calcareousness. An area of about 209 ha (23%) is marginally suitable (Class S3) and is distributed in the southwestern, northwestern and northeastern part of the microwatershed. They have moderate limitations of rooting depth, calcareousness, gravelliness and topography. An area of about 144 ha (16%) is currently not suitable (Class N1) for growing musambi and occur in the southern, southeastern, eastern and northeastern part of the microwatershed. They have severe limitations of rooting depth and texture.




Fig. 7.19 Land Suitability map of Musambi

## 7.20 Land Suitability for Lime (*Citrus sp*)

Lime is one of the most important fruit crop grown in an area of 0.11 lakh ha in almost all the districts of the State. The crop requirements for growing lime (Table 7.21) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing lime was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7. 20.

Highly suitable (Class S1) lands for growing lime occur in an area of 303 ha (34%) and are distributed in the central, southern, southeastern, eastern, northern and northeastern part of the microwatershed. An area of about 162 ha (18%) is moderately suitable (Class S2) for growing lime and are distributed in the northern, western and northeastern part of the microwatershed. They have minor limitations of drainage, rooting depth and calcareousness. An area of about 209 ha (23%) is marginally suitable (Class S3) and is distributed in the southwestern, northwestern and northeastern part of the microwatershed. They have moderate limitations of rooting depth, calcareousness, gravelliness and topography. An area of about 144 ha (16%) is currently not suitable (Class N1) for growing lime and occur in the southern, southeastern, eastern and northeastern part of the microwatershed. They have severe limitations of rooting depth and texture.






Fig. 7.20 Land Suitability map of Lime

### 7.21 Land Suitability for Amla (*Phyllanthus emblica*)

Amla is one of the medicinal fruit crop grown in almost all the districts of the State. The crop requirements for growing amla (Table 7.22) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing amla was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.21.

Highly (Class S1) suitable lands for growing amla occur in an area of 74 ha (8%) and are distributed in the western and northern part of the microwatershed. An area of about 392 ha (44%) is moderately suitable (Class S2) for amla and is distributed in all parts of the microwatershed except central, south and east. They have minor limitations of texture, calcareousness, gravelliness, topography, drainage and rooting depth. An area of 317 ha (36%) is marginally suitable (Class S3) and is distributed in the central, southern, southeastern, eastern and northeastern part of the microwatershed. They have moderate limitations of rooting depth, calcareousness and texture. An area of about 35 ha (4%) is currently not suitable (Class N1) for growing lime and occur in the northeastern part of the microwatershed. They have severe limitations of rooting depth and texture.




Fig. 7.21 Land Suitability map of Amla

## 7.22 Land Suitability for Cashew (*Anacardium occidentale*)

Cashew is one of the most important plantation nut crop grown in an area of 0.7 lakh ha in almost all the districts of the State. The crop requirements for growing cashew (Table 7.23) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing cashew was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.22.

No highly (Class S1) suitable lands available for growing cashew in the microwatershed. An area of 15 ha (2%) is moderately suitable (Class S2) for cashew and are distributed in the northern and northeastern part of the microwatershed with minor limitations of rooting depth and texture. About 327 ha (37%) area is marginally suitable (Class S3) for cashew and is distributed in all parts of the microwatershed except north and northeast. They have moderate limitations of texture, calcareousness, gravelliness, topography and rooting depth. Maximum area of 476 ha (54%) is currently not suitable (Class N1) for cashew and is distributed in the major part of the microwatershed with severe limitations of rooting depth, texture, calcareousness and drainage.




Fig. 7.22 Land Suitability map of Cashew

### 7. 23 Land Suitability for Jackfruit (*Artocarpus heterophyllus*)

Jackfruit is one of the most important fruit crop grown in an area of 5368 ha in almost all the districts of the State. The crop requirements for growing jackfruit (Table 7.24) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing jackfruit was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.23.

No highly (Class S1) suitable lands available for growing jackfruit in the microwatershed. Moderately suitable (Class S2) lands occur in an area of 15 ha (2%) and are distributed in the northern and northeastern part of the microwatershed with minor limitations of rooting and calcareousness. An area of about 659 ha (74%) is marginally suitable (Class S3) for growing jackfruit and are distributed in all parts of the microwatershed. They have moderate limitations of rooting depth, texture, graveliness, calcareousness, topography and drainage. An area of about 144 ha (16%) is currently not suitable (Class N1) for growing jackfruit and occur in the southern, southeastern, eastern and northeastern part of the microwatershed. They have severe limitations of rooting depth and texture.




Fig. 7.23 Land Suitability map of Jackfruit

#### 7.24 Land Suitability for Jamun (*Syzygium cumini*)

Jamun is an important fruit crop grown in almost all the districts of the State. The crop requirements for growing jamun (Table 25) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing jamun was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.24.

No highly (Class S1) suitable lands available for growing jamun in the microwatershed. Moderately suitable (Class S2) lands occur in an area of 450 ha (51%) and are distributed in the major part of the microwatershed with minor limitations of rooting depth, texture, calcareousness and drainage. An area of about 223 ha (25%) is marginally suitable (Class S3) for growing jamun and are distributed in the southwestern, northwestern and northeastern part of the microwatershed. They have moderate limitations of rooting depth, gravelliness, calcareousness and topography. An area of about 144 ha (16%) is currently not suitable (Class N1) for growing jamun and occur in the southern and northeastern part of the microwatershed. They have severe limitations of rooting depth and texture.




Fig. 7.24 Land Suitability map of Jamun

### 7.25 Land Suitability for Custard Apple (*Annona reticulata*)

Custard apple is one of the most important fruit crop grown in almost all the districts of the State. The crop requirements for growing custard apple (Table 7.26) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing custard apple was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.25.

Highly suitable (Class S1) lands for growing custard apple occur in an area of 192 ha (22%) and are distributed in the western, northern and northeastern and part of the microwatershed. An area of about 274 ha (31%) is moderately suitable (Class S2) for growing custard apple and are distributed in the southwestern, northwestern, northern, northeastern and southeastern part of the microwatershed. They have minor limitations of rooting depth, topography, calcareousness and graveliness. An area of about 330 ha (37%) is marginally suitable (Class S3) for growing custard apple and is distributed in the central, southern, southeastern, eastern and northeastern part of the microwatershed with moderate limitations rooting depth, texture and calcareousness. An area of about 22 ha (2%) is currently not suitable (Class N1) and are distributed in the northeastern part of the microwatershed with severe limitation of rooting depth.




Fig. 7.25 Land Suitability map of Custard Apple

## 7.26 Land Suitability for Tamarind (*Tamarindus indica*)

Tamarind is one of the most important spice crop grown in almost all the districts of the state. The crop requirements for growing tamarind (Table 7.27) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing tamarind was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are shown in Figure 7.26.

No highly suitable (Class S1) lands available for growing tamarind in the microwatershed. An area of about 450 ha (51%) is moderately suitable (Class S2) for growing tamarind and are distributed in the major part of the microwatershed. They have minor limitations of texture, rooting depth and drainage. Marginally suitable (Class S3) lands for growing tamarind occupy an area of about 133 ha (15%) and are distributed in the northern, northwestern, western, southwestern and northeastern part of the microwatershed. They have moderate limitations of rooting depth, gravelliness and calcareousness. An area of about 235 ha (26%) is currently not suitable (Class N1) for growing tamarind and occur in the northwestern, southwestern, southern, eastern and northeastern part of the microwatershed. They have severe limitations of rooting depth, calcareousness, topography and texture.




Fig. 7.26 Land Suitability map of Tamarind

### 7.27 Land Suitability for Mulberry (*Morus nigra*)

Mulberry is one of the important leaf crop grown for rearing silk worms in about 1.6 lakh ha area in all the districts of the state. The crop requirements for growing mulberry (Table 7.28) were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing mulberry was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed is given in Figure 7.27.

No highly (Class S1) suitable lands available for growing mulberry in the microwatershed. Moderately suitable (Class S2) lands occur in an area of 133 ha (15%) and are distributed in the northern, northwestern and southwestern part of the microwatershed with minor limitations of calcareousness, gravelliness and rooting depth. An area of about 557 ha (61%) is marginally suitable (Class S3) for growing mulberry and are distributed in the major part of the microwatershed. They have moderate limitations of rooting depth, drainage, topography, calcareousness and texture. Currently not suitable (Class N1) lands for growing mulberry occur in an area of 148 ha (16%) and are distributed in the southern, eastern and northeastern part of the microwatershed. They have severe limitations of rooting depth and texture.




Fig 7.27 Land Suitability map of Mulberry

### 7.28 Land suitability for Marigold (*Tagetes sps.*)

Marigold is one of the most important flower crop grown in an area of 9108 ha in almost all the districts of the State. The crop requirements (Table 7.29) for growing marigold were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing marigold was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.28.

No highly (Class S1) suitable lands available for growing marigold in the microwatershed. Moderately suitable (Class S2) lands occur in an area of 557 ha (63%) and are distributed in the major part of the microwatershed with minor limitations of calcareousness, drainage, texture, gravelliness, topography and rooting depth. An area of about 240 ha (27%) is marginally suitable (Class S3) for growing marigold and are distributed in all parts of the microwatershed except north, southeast and central. They have moderate limitations of rooting depth, gravelliness and texture. Currently not suitable (Class N1) lands for growing marigold occur in an area of 22 ha (2%) and are distributed in the northeastern part of the microwatershed. They have severe limitation of rooting depth.






Fig. 7.28 Land Suitability map of Marigold

### 7.29 Land Suitability for Chrysanthemum (*Dendranthema grandiflora*)

Chrysanthemum is one of the most important flower crop grown in an area of 4978 ha in almost all the districts of the State. The crop requirements (Table 7.30) for growing chrysanthemum were matched with the soil-site characteristics (Table 7.1) and a land suitability map for growing chrysanthemum was generated. The area extent and their geographical distribution of different suitability subclasses in the microwatershed are given in Figure 7.29.

No highly (Class S1) suitable lands available for growing chrysanthemum in the microwatershed. Moderately suitable (Class S2) lands occur in an area of 557 ha (63%) and are distributed in the major part of the microwatershed with minor limitations of calcareousness, drainage, texture, gravelliness, topography and rooting depth. An area of about 240 ha (27%) is marginally suitable (Class S3) for growing chrysanthemum and are distributed in all parts of the microwatershed except north, southeast and central. They have moderate limitations of rooting depth, gravelliness and texture. Currently not suitable (Class N1) lands for growing chrysanthemum occur in an area of 22 ha (2%) and are distributed in the northeastern part of the microwatershed. They have severe limitation of rooting depth.




Fig. 7.29 Land Suitability map of Chrysanthemum

**Table 7.1 Soil-site characteristics of Raisabad Hosalli-1 microwatershed**

Soil Map Units	Climate (P) (mm)	Growing period (Days)	Drainage Class	Soil depth (cm)	Soil texture		Gravelliness		AWC (mm/m)	Slope (%)	Erosion	pH	EC (dSm <sup>-1</sup> )	ESP (%)	CEC [Cmol (p <sup>+</sup> )kg <sup>-1</sup> ]	BS (%)
					Surface	Sub-surface	Surface (%)	Sub-surface (%)								
BDLiB2	866	150	W	25-50	sc	sl	<15	-	<50	1-3	Moderate	6.20	0.074	0.20	4.20	93
HLGcB2	866	150	W	50-75	sl	scl	<15	-	51-100	1-3	Moderate	8.49	0.185	0.69	8.80	100
JNKcB2	866	150	W	50-75	sl	scl	<15	-	51-100	1-3	Moderate	8.42	0.148	0.18	14.50	100
JNKiB2g1	866	150	W	50-75	sc	scl	15-35	-	51-100	3-5	Moderate	8.42	0.148	0.18	14.50	100
YLRcC3	866	150	W	50-75	sl	c	<15	15-35	51-100	1-3	Severe	6.91	0.069	0.45	6.90	100
BLCiB2	866	150	W	75-100	sc	scl	<15	-	51-100	1-3	Moderate	6.75	0.19	1.31	16.80	95
PGPcB2	866	150	W	75-100	sl	sc	<15	-	51-100	1-3	Moderate	6.83	0.210	2.83	3.15	100
YDRcB2	866	150	W	100-150	sl	sl	<15	-	51-100	1-3	Moderate	7.25	0.114	0.31	3.40	96
YDRcB2g1	866	150	W	100-150	sl	sl	15-35	-	51-100	1-3	Moderate	7.25	0.114	0.31	3.40	96
MDGiB2	866	150	W	100-150	sc	scl	<15	-	>200	1-3	Moderate	8.20	0.399	3.08	4.90	100
MDGhB2g1	866	150	W	100-150	scl	scl	15-35	-	>200	1-3	Moderate	8.20	0.399	3.08	4.90	100
BDPcB2	866	150	W	<25	scl	scl	<15	-	<50	1-3	Moderate	8.58	0.262	0.35	18.10	100
BDPhB2	866	150	W	<25	scl	scl	<15	-	<50	1-3	Moderate	8.58	0.262	0.35	18.10	100
KBDhB2	866	150	W	75-100	scl	g scl	<15	35-60	<50	1-3	Moderate	7.84	0.604	4.27	11.50	100
KBDiB2	866	150	W	75-100	sc	g scl	<15	35-60	<50	1-3	Moderate	7.84	0.604	4.27	11.50	100
HTKbB2g1	866	150	W	25-50	sl	sl	15-35	10-25	<50	1-3	Moderate	6.81	0.062	0.38	3.00	100
KDHcB2	866	150	MW	75-100	sl	sc	<15	-	101-150	1-3	Moderate	8.22	0.198	2.71	12.26	100
KDHiA1	866	150	MW	75-100	sc	sc	<15	-	101-150	0-1	Slight	8.22	0.198	2.71	12.26	100
SGRmB2	866	150	MW	>150	c	c	<15	-	>200	1-3	Moderate	8.3	1.48	11.61	34.77	100
SGRcB2	866	150	MW	>150	sl	c	<15	-	>200	1-3	Moderate	8.3	1.48	11.61	34.77	100
SGRiB2	866	150	MW	>150	sc	c	<15	-	>200	1-3	Moderate	8.3	1.48	11.61	34.77	100
KDPhB2	866	150	SED	>150	scl	s	<15	-	51-100	1-3	Moderate	6.55	0.07	1.37	2.53	61

\*Symbols and abbreviations are according to Field Guide for LRI under Sujala-III Project, Karnataka

**Table 7.2 Land suitability criteria for Sorghum**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	26–30	30–34; 24–26	34–40; 20–24	>40; <20
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, c (red), c (black)	scl, cl	ls, sl	-
	pH	1:2.5	5.5-7.8	5.0-5.5 7.8-9.0	>9.0	-
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	10-15
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

**Table 7.3 Land suitability criteria for Maize**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	30-34	35-38 26-30	38-40 26-20	
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc	c (red), c (black)	ls, sl	-
	pH	1:2.5	5.5-7.8	5.0-5.5 7.8-9.0	>9.0	-
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	-
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

**Table 7.4 Land suitability criteria for Bajra**

Land use requirement		Rating				
Soil –site characteristics	Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)	
Climatic regime	Mean temperature in growing season	°C	28-32	33-38 24-27	39-40 20-23	<20
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm	500-750	400-500	200-400	<200
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl, scl, cl,sc,c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.8	5.0-5.5 7.8-9.0	5.5-6.0 >9.0	
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	15-35	35-60	>60	
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	1-3	3-5	5-10	>10

**Table 7.5 Land suitability criteria for Groundnut**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	24–33	22–24; 33–35	20–22; 35–40	<20; >40
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. Well drained	Poorly drained	Very Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl	sl,cl, sc	c (red), c (black), ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<35	35-60	>60	
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.6 Land suitability criteria for Sunflower**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	24–30	30–34; 20–24	34–38; 16–20	>38; <16
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	mod. Well drained	-	Poorly to very drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	cl, sc,c (red), c (black)	scl	ls, sl	-
	pH	1:2.5	6.5-7.8	7.8-8.4 5.5-6.5	8.4-9.0; 5.0-5.5	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10



**Table 7.7 Land suitability criteria for Redgram**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	30-35(G) 20-25(AV) 15-18 (F&PS) 35-40(M)	25-30(G) 20-25 (AV) 12-15 (F&PS) 30-35(M)	20-25(G) 15-20(AV) 10-12 (F&PS) 25-30(M)	< 20 <15 <10 <25
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. Well drained	Poorly drained	Very Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, c (red)	c (black),sl, scl, cl	ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-9.0	5.0-5.5 >9.0	-
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-50	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<1.0	1.0-2.0	>2.0	
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.8 Land suitability criteria for Bengal gram**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	20–25	25–30; 15–20	30–35; 10–15	>35; <10
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. Well drained	Poorly drained	Very Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	c(black)	-	c (red), scl, cl, sc	ls, sl
	pH	1:2.5	6.0-7.8	5.0-6.0 7.8-9.0	>9.0	-
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	-
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.9 Land suitability criteria for Cotton**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	22-32	>32	<19	-
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well to moderately well	Poorly drained/Some what excessively drained	-	very poorly/excessively drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, c (red,black)	cl	scl	ls, sl
	pH	1:2.5	6.5-7.8	7.8-8.4	5.5-6.5 8.4->9.0	<5.5
	CEC	C mol (p+)Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	50-100	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	5-10	10-15	>15	
Erosion hazard	Slope	%	<3	3-5	-	>5

**Table 7.10 Land suitability criteria for Chilli**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	25-32	33-35 20-25	35-38 <20	>38
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc	c (black), sl	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.11 Land suitability criteria for Tomato**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	25-28	29-32 20-24	15-19 33-36	<15 >36
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl, scl, cl, sc, c (red)	-	ls, c(black)	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.12 Land suitability criteria for Brinjal**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	Well drained	Moderately well drained	Poorly drained	V. Poorly drained
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class				
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl, scl, cl, sc c (red)	-	ls, c (black)	-
	pH	1:2.5	6.0-7.3	7.3-8.4 5.0-6.0	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	>60
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.13 Land suitability criteria for Onion**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	20-30	30-35	35-40	>40
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately /imperfectly	-	Poorly to V poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl,scl,cl,sc,c (red)	-	c (Black),ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<1.0	1.0-2.0	2.0-4.0	<4
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.14 Land suitability criteria for Bhendi**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	25-28	29-32 20-24	15-19 33-36	<15 >36
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Imperfectly drained	Poorly to very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl,sc, c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10



**Table 7.15 Land suitability criteria for Drumstick**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, scl, cl, c (red)	sl, c (black)	ls	s
	pH	1:2.5	6.0-7.3	5.0-5.5 7.3-7.8	5.5-6.0 7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<35	35-60	60-80	>80
Soil toxicity	Salinity (EC saturation extract)	ds/m				
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-10	-	>10

**Table 7.16 Land suitability criteria for Mango**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-32	24-27 33-35	36-40	20-24
	Min temp. before flowering	°C	10-15	15-22	>22	-
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration	Days				
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V. Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	-	ls, sl, c (black)	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>150	100-150	75-100	<75
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.17 Land suitability criteria for Guava**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-32	33-36 24-27	37-42 20-23	
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	sl	c (black), ls	-
	pH	1:2.5	6.0-7.8	5.0-6.0	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.18 Land suitability criteria for Sapota**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-32	33-36 24-27	37-42 20-23	>42 <18
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	-	Poorly to very drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	sl	ls, c (black)	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.19 Land suitability criteria for Pomegranate**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	30-34	35-38 25-29	39-40 15-24	
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl,cl, sc, c (red)	c (black),sl	ls	-
	pH	1:2.5	5.5-7.8	7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.20 Land suitability criteria for Musambi**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	28-30	31-35 24-27	36-40 20-23	>40 <20
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately drained	poorly	Very poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c	sl	ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.21 Land suitability criteria for Lime**

Land use requirement		Rating				
Soil –site characteristics	Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)	
Climatic regime	Mean temperature in growing season	°C	28-30	31-35 24-27	36-40 20-23	>40 <20
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately drained	poorly	Very poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c	sl	ls	-
	pH	1:2.5	6.0-7.8	5.5-6.0 7.8-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.22 Land suitability criteria for Amla**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. well drained	Poorly drained	V. Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	c (black)	ls, sl	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15-35	35-60	60-80	-
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10



**Table 7.23 Land suitability criteria for Cashew**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	32 to 34	28 to 32; 34 to 38	24 to 28; 38 to 40	<20; >40
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	moderately well drained	Poorly drained	Very poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	-	sl, ls	c (black)
	pH	1:2.5	5.5-6.5	5.0-5.5 6.5-7.3	7.3-7.8	>7.8
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	<3	3-10	>10	-

**Table 7.24 Land suitability criteria for Jackfruit**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. well	Poorly	V. Poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c (red)	-	sl, ls, c (black)	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	>60
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10-

**Table 7.25 Land suitability criteria for Jamun**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well	Mod. well	Poorly	V.Poorly
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl, sc, c(red)	sl, c (black)	ls	-
	pH	1:2.5	6.0-7.8	5.0-6.0	7.8-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>150	100-150	50-100	<50
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	>60
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

**Table 7.26 Land suitability criteria for Custard apple**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod. well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	Scl, cl, sc, c (red), c (black)	-	Sl, ls	-
	pH	1:2.5	6.0-7.3	5.5-6.0 7.3-8.4	5.0-5.5 8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15-35	35-60	60-80	-
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	>5	-

**Table 7.27 Land suitability criteria for Tamarind**

Land use requirement		Rating				
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C				
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Mod.well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	scl, cl,sc, c (red)	sl, c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-7.8	7.8-8.4	>8.4
	CEC	C mol (p+)/ Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>150	100-150	75-100	<75
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

**Table 7.28 Land suitability criteria for Mulberry**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	24–28	22–24; 28–32	32–38; 22–18	>38; <18
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V. Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sc, cl, scl	c (red)	c (black), sl, ls	-
	pH	1:2.5	5.5-7.3	5.0-5.5 7.8-8.4	7.3-8.4	>8.4
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO <sub>3</sub> in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>100	75-100	50-75	<50
	Stoniness	%				
	Coarse fragments	Vol %	0-35	35-60	60-80	>80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2	2-4	4-8	>8
	Sodicity (ESP)	%	<5	5-10	10-15	>15
Erosion hazard	Slope	%	0-3	3-5	5-10	>10

**Table 7.29 Land suitability criteria for Marigold**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	18-23	17-15 24-35	35-40 10-14	>40 <10
	Mean max. temp. in growing season	°C				
	Mean min. tempt. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl,scl, cl, sc, c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%				
Erosion hazard	Slope	%	<3	3-5	5-10	>10

**Table 7.30 Land suitability criteria for Chrysanthemum**

Land use requirement			Rating			
Soil –site characteristics		Unit	Highly suitable (S1)	Moderately suitable (S2)	Marginally suitable (S3)	Not suitable (N1)
Climatic regime	Mean temperature in growing season	°C	18-23	17-15 24-35	35-40 10-14	>40 <10
	Mean max. temp. in growing season	°C				
	Mean min. temp. in growing season	°C				
	Mean RH in growing season	%				
	Total rainfall	mm				
	Rainfall in growing season	mm				
Land quality	Soil-site characteristic					
Moisture availability	Length of growing period for short duration	Days				
	Length of growing period for long duration					
	AWC	mm/m				
Oxygen availability to roots	Soil drainage	Class	Well drained	Moderately well drained	Poorly drained	V.Poorly drained
	Water logging in growing season	Days				
Nutrient availability	Texture	Class	sl,scl, cl, sc, c (red)	c (black)	ls	-
	pH	1:2.5	6.0-7.3	5.0-6.0 7.3-8.4	8.4-9.0	>9.0
	CEC	C mol (p+)/Kg				
	BS	%				
	CaCO3 in root zone	%		<5	5-10	>10
	OC	%				
Rooting conditions	Effective soil depth	cm	>75	50-75	25-50	<25
	Stoniness	%				
	Coarse fragments	Vol %	<15	15-35	35-60	60-80
Soil toxicity	Salinity (EC saturation extract)	ds/m	<2.0	2-4	4-8	>8.0
	Sodicity (ESP)	%				
Erosion hazard	Slope	%	<3	3-5	5-10	>10



### 7.30 Land Management Units (LMUs)

The 22 soil map units identified in Raisabad Hosalli-1 microwatershed have been grouped into 9 Land Management Units (LMU's) for the purpose of preparing a Proposed Crop Plan. Land Management Units are grouped based on the similarities in respect of the type of soil, the depth of the soil, the surface soil texture, gravel content, AWC, slope, erosion etc. And a Land Management Units map (Fig. 7.30) has been generated. These Land Management Units are expected to behave similarly for a given level of management.

The 22 map units that have been grouped into 9 Land Management Units along with brief description of soil and site characteristics are given below.

LMU	Soil map units	Soil and site characteristics
1	180.KDPhB2	Very deep, lowland sandy soils, 1-3% slopes, non gravelly, moderate erosion
2	42.YDRcB2 154.YDRcB2g1	Deep, sandy loam soils, 1-3% slopes, non gravelly to gravelly, moderate erosion
3	99.KDHcB2 106.SGRmB2 141.SGRcB2 143.SGRiB2 157.KDHiA1	Moderately deep to very deep, lowland clay soils, 0-3% slopes, non gravelly, moderate erosion
4	130.KBDhB2 131.KBDiB2	Moderately deep, red gravelly loam soils, 1-3% slopes, non gravelly, moderate erosion
5	58.MDGiB2 149.MDGhB2g1	Deep, black loamy soils, 1-3% slopes, non gravelly to gravelly, moderate erosion
6	38.BLCiB2 40.PGPcB2	Moderately deep to deep, red sandy clay to sandy clay loam soils, 1-3% slopes, non gravelly, moderate erosion
7	30.YLRcC3	Moderately shallow, red clay soils, 3-5% slopes, non gravelly, severe erosion
8	16.HLGcB2 20.JNKcB2 23.JNKiB2g1	Moderately shallow, sandy clay loam soils, 1-3% slopes, non gravelly to gravelly, moderate erosion
9	5.BDLiB2 118.BDPcB2 120.BDPhB2 161.HTKbB2g1	Shallow to very shallow, sandy loam soils, 1-3% slopes, non gravelly to gravelly, moderate erosion




Fig. 7.30 Land Management Units Map- Raisabad Hosalli-1 Microwatershed

### 7.31 Proposed Crop Plan for Raisabad Hosalli-1 Microwatershed

After assessing the land suitability for the 29 crops, the Proposed Crop Plan has been prepared for the 9 identified LMUs by considering only the highly (Class S1) and moderately (Class S2) suitable lands for each of the 29 crops. The resultant proposed crop plan is presented below in Table 7.31.

**Table 7.31 Proposed Crop Plan for Raisabad Hosalli-1 Microwatershed**

<b>LMU</b>	<b>Mapping Units</b>	<b>Survey Number</b>	<b>Soil Characteristics</b>	<b>Field Crops</b>	<b>Horticulture Crops</b>	<b>Suitable Interventions</b>
1	180.KDPhB2	<b>Panchasheelanagara:</b> 47,48	Very deep, lowland sandy soils), 1-3% slopes, non gravelly, moderate erosion	-	<b>Agri-Silvi-Pasture:</b> Hybrid Napier, <i>Styloxanthes hamata</i> , <i>Styloxanthes scabra</i>	Use of short duration varieties, sowing across the slope, and mulching is recommended
2	42.YDRcB2 154.YDRcB2g1	<b>Naglapura :</b> 196,205 <b>Risabadha Hosalli:</b> 74,75,76,77, 78,79,80,83,84,85,86,87,88,89,103, 104,105,106,107,108,109,110,111,112,113,114,115,116,117,118,119,120,121,122,123,124,125,126,127,128,129,130,131,132,133,134,135,136,137,138,139,140,141,142,143,144,145,146,147,148,149,150,151,152,153,154,155,156,157/1,157/2,158,159,160,161,162,163,164,165,166,167,168,169,170,171,172,173,174,175,176,177,178,179	Deep, sandy loam soils, 1-3% slopes, non gravelly to gravelly, moderate erosion	Sorghum, Groundnut, Red gram, Horse gram	<b>Fruit crops:</b> Sapota, Pomegranate, Guava, Lime, Musambi, Jamun, Tamarind, Amla, Custard apple <b>Vegetables:</b> Onion, Tomato, Bhendi, Drumstick, Chilli, Coriander <b>Flowers:</b> Marigold, Chrysanthemum	Application of FYM, biofertilizers and micronutrients, drip irrigation, mulching, suitable soil and water conservation practices
3	99.KDHcB2 106.SGRmB2 141.SGRcB2 143.SGRiB2 157.KDHiA1	<b>Halagera:</b> 185,192,193,194,195/1,195/2,198,199,200,202,223,410 <b>Mylapura:</b> 51,53,57,58,59,60,61,62,63,64	Moderately deep to very deep, lowland clay soils, 0-3% slopes, non gravelly, moderate erosion	Sorghum, maize, Bajra	<b>Fruit crops:</b> Custard Apple, Amla <b>Flower crops:</b> Marigold, Chrysanthemum, Jasmine	Providing proper drainage, addition of organic manures, green leaf manuring, suitable conservation practices
4	130.KBDhB2 131.KBDiB2	<b>Halagera:</b> 156,157,159/2,159/3,160,166,167,168,169,170,171,187,188,189,190,191,196,197,204,205,206,207,208,209,210,211,212,213,214,215 <b>Risabadha Hosalli:</b> 15,181/A,183	Moderately deep, red gravelly loamy soils, 1-3% slopes, non gravelly, moderate erosion	Groundnut, Bajra, Horse gram, Castor, Mulberry	<b>Fruit crops:</b> Musambi, Lime, Jackfruit, Amla, Custard apple, Tamarind <b>Vegetable crops:</b> Drumstick, Curry leaves	Drip irrigation, mulching, suitable soil and water conservation practices (Crescent Bunding with Catch Pit etc)
5	58.MDGiB2 149.MDGH2g1	<b>Halagera :</b> 159/1 <b>Mylapura:</b> 123,124,125,126,127,128,131	Deep, black loamy soils, 1-3% slopes, non gravelly to	Sunflower, Sorghum, Soybean,	<b>Fruit crops:</b> Pomegranate, Lime, Musambi, Tamarind, Jamun, Amla, Custard apple	Application of FYM, biofertilizers and micronutrients, drip

		<b>Risabadha Hosalli:</b> 10,11,12,13,14,16,17,18,19,20,21,22,23,24,25,27,28,29,30,31,32,33,34,35,4,46/4,47,48,55,56,57,59,60,64,7,82	gravelly, moderate erosion	Cotton, Bengal gram, Maize, Safflower, Linseed, Bajra	<b>Vegetables:</b> Drumstick, Chilli, Bhendi, Cluster bean, Coriander <b>Flowers:</b> Marigold, Chrysanthemum	irrigation, mulching, suitable soil and water conservation practices
6	38.BLCiB2 40.PGPcB2	<b>Mylapura:</b> 52/1,52/2,54,55,56 <b>Panchasheelanagara:</b> 46	Moderately deep to deep, red sandy clay to sandy clay loam soils, 1-3% slopes, non gravelly, moderate erosion	Maize, Sorghum, Sunflower, Bajra, Finger millet, Castor, Cowpea, Groundnut, Red gram, Field bean, Mulberry	<b>Fruit crops:</b> Mango, Pomegranate, Guava, Sapota, Jackfruit, Jamun, Tamarind, Lime, Musambi, Amla, Custard apple, Cashew <b>Vegetable crops:</b> Drumstick, Tomato, Bhendi, Chilli, Brinjal, Onion, Curry leaves <b>Flower crops:</b> Marigold, Chrysanthemum, Jasmine, Crossandra	Drip irrigation, mulching, suitable soil and water conservation practices (Crescent Bunding with Catch Pit etc)
7	30.YLRcC3	<b>Jinatera :</b> 253,256,258	Moderately shallow, red clay soils, 3-5% slopes, non gravelly, severe erosion	Maize, Groundnut, Bajra, Red gram	<b>Fruit crops:</b> Amla, Custard apple <b>Vegetables:</b> Tomato, Chilli, Brinjal <b>Flowers:</b> Marigold Chrysanthemum	Drip irrigation, mulching, suitable soil and water conservation practices (Crescent Bunding with Catch Pit etc)
8	16.HLGcB2 20.JNKcB2 23.JNKiB2g1	<b>Halagera:</b> 158,164,165,186,203,216, 217,218,219,221,222,224 <b>Risabadha Hosalli:</b> 45,46/1,46/10,46/5,46/6,46/7,46/8,46/9,49,50,51,52,53,54,65	Moderately shallow, sandy clay loam soils, 1-3% slopes, non gravelly to gravelly, moderate erosion	Sorghum, Bajra, Coriander	<b>Fruit crops:</b> Amla, Custard apple <b>Vegetables:</b> Coriander, Bhendi <b>Flowers:</b> Marigold, Jasmine Chrysanthemum	Application of FYM, Biofertilizers and micronutrients, drip irrigation, mulching, suitable soil and water conservation practices
9	5.BDLiB2 118.BDPcB2 120.BDPbB2 161.HTKbB2g1	<b>Naglapura :</b> 195,206,207 <b>Risabadha Hosalli :</b> 26,58,66,67,68,69,71,72,73,90,91,92,93,94,95,96,97,98,99,100,101,102 <b>Yaleri:</b> 691,693,694,695,696,697,698,699,702	Shallow to very shallow, sandy loam soils, 1-3% slopes, non gravelly to gravelly, moderate erosion	-	<b>Agri-Silvi-Pasture:</b> Custard apple, <i>Styloxanthes hamata</i> , Glyricidia, <i>Styloxanthes scabra</i>	Use of short duration varieties, sowing across the slope and split application of nitrogen fertilizers

## SOIL HEALTH MANAGEMENT

### 8.1 Soil Health

Soil health is basic to plant health and plant health is basic to human and bovine health. Soil is fundamental to crop production. Without soil, no food could be produced nor would livestock be fed on a large scale. Because it is finite and fragile, soil is a precious resource that requires special care from its users.

Soil health or the capacity of the soil to function is critical to human survival. Soil health has been defined as: “the capacity of the soil to function as a living system without adverse effect on the ecosystem”. Healthy soils maintain a diverse community of soil organisms that help to form beneficial symbiotic associations with plant roots, recycle essential plant nutrients, improve soil structure with positive repercussions for soil, water and nutrient holding capacity and ultimately improve crop production and also contribute to mitigating climate change by maintaining or increasing its carbon content.

Functional interactions of soil biota with organic and inorganic components, air and water determine a soil’s potential to store and release nutrients and water to plants and to promote and sustain plant growth. Thus, maintaining soil health is vital to crop production and conserve soil resource base for sustaining agriculture.

#### **The most important characteristics of a healthy soil are**

- Good soil tilth
- Sufficient soil depth
- Good water storage and good drainage
- Adequate supply, but not excess of nutrients
- Large population of beneficial organisms
- Small proportion of plant pathogens and insect pests
- Low weed pressure
- Free of chemicals and toxins that may harm the crop
- Resistance to degradation
- Resilience when unfavorable conditions occur

#### **Characteristics of Raisabad Hosalli-1 Microwatershed**

- ❖ The soil phases identified in the microwatershed belonged to the soil series of YDR series occupying maximum area of 207 ha (23%) followed by MDG 80 ha (18%), KBD 118 ha (13%), BDL 78 ha (9%), JNK 77 ha (9%), SGR 59 ha (6%), HTK 31 ha (3%), KDH 23 ha (3%), BDP 22 ha (2%), BLC 15 ha (2%), HLG 13 ha (1%), KDP 13 ha (1%) YLR 1 ha (<1%), PGP 0 ha (<1%).
- ❖ As per land capability classification, entire area of the microwatershed falls under arable land category (Class II, III & IV). The major limitations identified in the arable lands were soil, erosion and drainage.

- ❖ On the basis of soil reaction, about 34 ha (4%) is slightly acid (pH 6.0-6.5), 453 ha (51%) is neutral (pH 6.5-7.3), 188 ha (21 %) is slightly alkaline (pH 7.3-7.8), moderately alkaline 135 ha (15%) and strongly alkaline 7 ha (<1%).

#### ❖ **Soil Health Management**

The following actions are required to improve the current land husbandry practices that provide a sound basis for the successful adoption of sustainable crop production system.

#### **Acid soils**

Slightly acidic soils cover about 34 ha area in the microwatershed.

1. Growing of crops suitable for a particular soil pH.
2. Amelioration of soils through the application of amendments (liming materials).

Liming materials:

1. CaCO<sub>3</sub> (Calcium Carbonate).
2. Dolomite [Ca Mg (CO<sub>3</sub>)<sub>2</sub>]
3. Quick lime (CaO)
4. Slaked lime [Ca (OH)<sub>2</sub>]

For normal pH and pH 4.8 (35 t/ha) and pH 6.0-7.0 (4 t/ha) lime is required.

#### **Alkaline soils**

Slightly alkaline to strongly alkaline soils cover about 330 ha area in the microwatershed.

1. Regular addition of organic manure, green manuring, green leaf manuring, crop residue incorporation and mulching needs to be taken up to improve the soil organic matter status.
2. Application of biofertilizers (Azospirillum, Azotobacter, Rhizobium).
3. Application of 25% extra N and P (125 % RDN&P).
4. Application of ZnSO<sub>4</sub> – 12.5 kg/ha (once in three years).
5. Application of Boron – 5kg/ha (once in three years).

#### **Neutral soils**

Neutral soils occur in 453ha area in the microwatershed.

1. Regular addition of organic manure, green manuring, green leaf manuring, crop residue incorporation and mulching needs to be taken up to improve the soil organic matter status.
2. Application of biofertilizers, (Azospirillum, Azotobacter, Rhizobium).
3. Application of 100 per cent RDF.
4. Need based micronutrient applications.

Besides the above recommendations, the best transfer of technology options are also to be adopted.

## **Soil Degradation**

Soil erosion is one of the major factor affecting the soil health in the microwatershed. Out of total 887 ha area in the microwatershed, an area of about <1 ha (<1%) is suffering from slight erosion, about 816 ha (92%) is suffering from moderate erosion and one ha (<1%) from severe erosion. In areas of moderate and severe erosion immediate soil and water conservation and, other land development and land husbandry practices are required for restoring soil health.

## **Dissemination of Information and Communication of Benefits**

Any large scale implementation of soil health management requires that supporting information is made available widely, particularly through channels familiar to farmers and extension workers. Given the very high priority attached to soil-health especially by the Central Government on issuing Soil-Health Cards to all the farmers, media outlets like Regional, State and National Newspapers, Radio and Dooradarshan programs in local languages but also modern information and communication technologies such as Cellular phones and the Internet, which can be much more effective in reaching the younger farmers.

## **Inputs for Net Planning (Saturation Plan) and Interventions needed**

Net planning (Saturation Plan) in IWMP is focusing on preparation of

1. Soil and Water Conservation Plan for each plot or farm.
2. Productivity enhancement measures/ interventions for existing crops/livestock/other farm enterprises.
3. Diversification of farming mainly with perennial horticultural crops and livestock.
4. Improving livelihood opportunities and income generating activities.

In this connection, how various outputs of Sujala-III are of use in addressing these objectives of Net Planning (Saturation Plan) are briefly presented below.

- ❖ **Soil Depth:** The depth of a soil decides the amount of moisture and nutrients it can hold, what crops can be taken up or not, depending on the rooting depth and the length of growing period available for raising any crop. Deeper the soil, better for a wide variety of crops. If sufficient depth is not available for growing deep rooted crops, either choose medium or short duration crops or deeper planting pits need to be opened and additional good quality soil brought from outside has to be filled into the planting pits.
- ❖ **Surface Soil Texture:** Lighter soil texture in the top soil means, better rain water infiltration, less run-off and soil moisture conservation, less capillary rise and less evaporation losses. Lighter surface textured soils are amenable to good soil tilth and are highly suitable for crops like groundnut, root vegetables (carrot, raddish, potato etc) but not ideal for crops that need stagnant water like lowland paddy. Heavy textured soils are poor in water infiltration and percolation. They are prone for sheet erosion; such soils can

be improved by sand mulching. The technology that is developed by the AICRP-Dryland Agriculture, Vijayapura, Karnataka can be adopted.

- ❖ **Gravelliness:** More gravel content is favorable for run-off harvesting but poor in soil moisture storage and nutrient availability. It is a significant parameter that decides the kind of crop to be raised.
- ❖ **Land Capability Classification:** The land capability map shows the areas suitable and not suitable for agriculture and the major constraints in each of the plot/survey number. Hence, one can decide what kind of enterprise is possible in each of these units. In general, erosion and soil are the major constraints in Raisabad Hosalli-1 microwatershed.
- ❖ **Organic Carbon:** The OC content (an index of available Nitrogen) is medium (0.5-0.75) in 605 ha (68%) and high (>0.75%) in 212 (24%). Area under medium in OC needs to be further improved by applying farmyard manure and rotating crops with cereals and legumes or mixed cropping.
- ❖ **Promoting Green Manuring:** Growing of green manuring crops costs Rs. 1250/ha (green manuring seeds) and about Rs. 2000/ha towards cultivation that totals to Rs. 3250/- per ha. On the other hand, application of organic manure @ 10 tons/ha costs Rs. 5000/ha. The practice needs to be continued for 2-3 years or more. Nitrogen fertilizer needs to be supplemented by 25% in addition to the recommended level in 605 ha area where OC is medium (<0.5 - 0.75%). For example, for rainfed maize, recommended level is 50 kg N per ha and an additional 12 kg /ha needs to be applied for all the crops grown in these plots.
- ❖ **Available Phosphorus:** Available phosphorus is low (<23 kg/ha) in 179 ha (20%) of the microwatershed, medium (23-57 kg/ha) in an area of 625 ha (70%) and high (23-57 kg/ha) in an area of 13 ha (1%) In low and medium areas, for all the crops 25% additional P needs to be applied.
- ❖ **Available Potassium:** Available potassium is high (>337 kg/ha) in an area of 16 ha (2%) of the microwatershed, medium (145-337 kg/ha) in an area of 726 ha (82%) of the microwatershed and low (<145 kg/ha) in an area of 75 ha (8%). All the plots, where available potassium is low and medium, for all the crops, additional 25% potassium may be applied.
- ❖ **Available Sulphur:** Available sulphur is a very critical nutrient for oilseed crops. It is low in an area of 123 ha (14%) and medium in an area of 693 ha (78%). Low and medium areas need to be applied with magnesium sulphate or gypsum or factomphos (p) fertilizer (13% sulphur) for 2-3 years for the deficiency to be corrected.
- ❖ **Available Boron:** An area of 561 ha (63%) is low (<0.5 ppm) and an area of about 256 ha (29%) is medium (0.5-1.0 ppm). For these areas, application of sodium borate @ 10 kg/ha as soil application or 0.2 % borax as foliar spray is recommended.
- ❖ **Available Iron:** Entire area of the microwatershed is sufficient (>4.5 ppm) in available iron content.



- ❖ **Available Manganese:** All the soils in the microwatershed are sufficient (>1.0 ppm) in available manganese.
- ❖ **Available Copper:** All the soils in the microwatershed are sufficient (>0.2 ppm) in available copper.
- ❖ **Available Zinc:** Entire area of the microwatershed is deficient (<0.6 ppm) in available zinc content. Application of zinc sulphate @ 25 kg/ha is recommended for these areas. About 72 ha (11%) area is sufficient (>0.6 ppm).
- ❖ **Soil Alkalinity:** An area of 330 ha (37%) in the microwatershed has soils that are slightly alkaline to strongly alkaline. These areas need application of gypsum and wherever calcium is in excess, iron pyrites and element sulphur can be recommended. Management practices like treating repeatedly with good quality water to drain out the excess salts and provision of subsurface drainage and growing of salt tolerant crops like Casuarina, Acacia, Neem, Ber etc, are recommended.


**Land Suitability for Various Crops:** Areas that are highly, moderately and marginally suitable and not suitable for growing various crops are indicated. Along with the suitability, various constraints that are limiting the productivity are also indicated. For example, in case of cotton, gravel content, rooting depth and salinity/alkalinity are the major constraints in various plots. With suitable management interventions, the productivity can be enhanced. In order to increase the water holding capacity of light textured soils, growing of green manure crops and application of organic manure is recommended.



## SOIL AND WATER CONSERVATION TREATMENT PLAN

For preparing soil and water conservation treatment plan for Raisabad Hosalli-1 microwatershed, the land resource inventory database generated under Sujala-III project has been transformed as information through series of interpretative (thematic) maps using soil phase map as a base. The various thematic maps (1:7920 scale) generated were

- Soil depth
- Surface soil texture
- Available water capacity
- Soil slope
- Soil gravelliness
- Land capability
- Present land use and land cover
- Crop suitability
- Rainfall
- Hydrology
- Water Resources
- Socio-economic data
- Contour plan with existing features- network of waterways, pottissa boundaries, cut up/ minor terraces etc.
- Cadastral map (1:7920 scale)
- Satellite imagery (1:7920 scale)



Apart from these, Hand Level/ Hydro Marker/ Dumpy Level/ Total Station and Kathedars' List to be collected.

### Steps for Survey and Preparation of Treatment Plan

The boundaries of Land User Groups' and Survey No. boundaries are traced in the field.

- Naming of user groups and farmers
- Identification of arable and non arable lands
- Identification of drainage lines and gullies
- Identification of non treatable areas
- Identification of priority areas in the arable lands
- Treatment plan for arable lands
- Location of water harvesting and recharge structures

## 9.1 Treatment Plan

The treatment plan recommended for arable lands is briefly described below


### 9.1.1 Arable Land Treatment

#### A. BUNDING

Steps for Survey and Preparation of Treatment Plan		<b>USER GROUP-1</b>  <b>CLASSIFICATION OF GULLIES</b>  
<ul style="list-style-type: none"> <li>• Cadastral map (1:7920 scale) is enlarged to a scale of 1:2500 scale</li> <li>• Existing network of waterways, pothissa boundaries, grass belts, natural drainage lines/ watercourse, cut ups/ terraces are marked on the cadastral map to the scale</li> <li>• Drainage lines are demarcated into</li> </ul>		
Small gullies	(up to 5 ha catchment)	
Medium gullies	(5-15 ha catchment)	
Ravines	(15-25 ha catchment) and	
<i>Halla/Nala</i>	(more than 25ha catchment)	

#### Measurement of Land Slope

Land slope is estimated or determined by the study and interpretation of contours or by measurement in the field using simple instruments like Hand Level or Hydromarker.



$$\text{FALL: } 1.5 - 0.6 = 0.9 \text{ m.}$$

Vertical and Horizontal intervals between bunds as recommended by the Watershed Development Department.

Slope percentage	Vertical interval (m)	Corresponding Horizontal Distance (m)
2 - 3%	0.6	24
3 - 4%	0.9	21
4 - 5%	0.9	21
5 - 6%	1.2	21
6 - 7%	1.2	21

**Note:** (i) The above intervals are maximum.

(ii) Considering the slope class and erosion status (A1... A=0-1 % slope, 1= slight erosion) the intervals have to be decided.

**Bund length recording:** Considering the contour plan and the existing grass belts/partitions, the bunds are aligned and lengths are measured.

### Section of the Bund

Bund section is decided considering the soil texture class and gravelliness class (bg<sub>0</sub>... b=loamy sand, g<sub>0</sub> = <15% gravel). The recommended Sections for different soils are given below.

**Recommended Bund Section**

Top width (m)	Base width (m)	Height (m)	Side slope (Z:1;H:V)	Cross section (sq m)	Soil Texture	Remarks
0.3	0.9	0.3	01:01	0.18	Sandy loam	Vegetative bund
0.3	1.2	0.3	1.5:1	0.225	Sandy clay	
0.3	1.2	0.5	0.9:1	0.375	Red gravelly soils	
0.3	1.2	0.6	0.75:1	0.45		
0.3	1.5	0.6	01:01	0.54	Red sandy loam	
0.3	2.1	0.6	1.5:1	0.72	Very shallow black clayey soils	
0.45	2	0.75	01:01	0.92		
0.45	2.4	0.75	1.3:1	1.07	Shallow black clayey soils	
0.6	3.1	0.7	1.78:1	1.29	Medium black clayey soils	
0.5	3	0.85	1.47:1	1.49		

### Formation of Trench cum Bund

Dimensions of the Borrow Pits/Trenches to be excavated (machinery are decided considering the Bund Section).

**Details of Borrow Pit dimensions are given below:**

**TRENCH CUM BUND**

WATER STORAGE AREA

0.45 Sq.m section

IDEAL FOR HORTICULTURE CROPS

**'A' FRAME FOR INTERBUND MANAGEMENT**

1. ಸಮವಾತಳ ಉಳಿಸುವೆ

2. ಸಮವಾತಳ ಬಿತ್ತನೆ/ನಾಟಿ

### Size of Borrow Pits/ Trench recommended for Trench cum Bund (by machinery)

Bund section	Bund length	Earth quantity	Pit				Berm (pit to pit)	Soil depth class
			L(m)	W(m)	D(m)	Quantity (m <sup>3</sup> )		
m <sup>2</sup>	m	m <sup>3</sup>					m	
0.375	6	2.25	5.85	0.85	0.45	2.24	0.15	Shallow
0.45	6	2.7	5.4	1.2	0.43	2.79	0.6	Shallow
0.45	6	2.7	5	0.85	0.65	2.76	1	Moderately shallow
0.54	5.6	3.02	5.5	0.85	0.7	3.27	0.1	Moderately shallow
0.54	5.5	2.97	5	1.2	0.5	3	0.5	Shallow
0.72	6.2	4.46	6	1.2	0.7	5.04	0.2	Moderately shallow
0.72	5.2	3.74	5.1	0.85	0.9	3.9	0.1	Moderately deep

#### B. Water Ways

- Existing waterways are marked on the cadastral map (1:7920 scale) and their dimensions are recorded.
- Considering the contour plan of the MWS, additional waterways/ modernization of the existing ones can be thought of.
- The design details are given in the Manual.

#### C. Farm Ponds

Waterways and the catchment area will give an indication on the size of the Farm Pond. Location of the pond can be decided based on the contour plan/ field condition and farmers' need/desire.

#### D. Diversion Channel

Existing EPT/ CPT are marked on the cadastral map. Looking to the need, these can be modernized or fresh diversion channel can be proposed and runoff from this can be stored in *Gokatte*/ Recharge ponds.

#### 9.1.2 Non-Arable Land Treatment

Depending on the gravelliness and crops preferred by the farmers, the concerned authorities can decide appropriate treatment plan. The recommended treatments may be Contour Trench, Staggered Trench, Crescent Bund, Boulder Bund or Pebble Bund.

### **9.1.3 Treatment of Natural Water Course/ Drainage Lines**

- a) The cadastral map has to be updated as regards the network of drainage lines (gullies/*nalas/hallas*) and existing structures are marked to the scale and storage capacity of the existing water bodies are documented.
- b) The drainage line will be demarcated into Upper Reach, Middle Reach and Lower Reach.
- c) Considering the Catchment, *Nala* bed and bank conditions, suitable structures are decided.
- d) Number of storage structures (Check dam/*Nala* bund/ Percolation tank) will be decided considering the commitments and available runoff from water budgeting and quality of water in the wells and site suitability.
- e) Detailed Leveling Survey using Dumpy Level / Total Station has to be carried out to arrive at the site-specific designs as shown in the Manual.
- f) The location of ground water recharge structures are decided by examining the lineaments and fracture zones from geological maps.
- g) Rainfall intensity data of the nearest Rain Gauge Station is considered for Hydrologic Designs.
- h) Silt load to the Storage/Recharge Structures is reduced by providing vegetative, boulder and earthen checks in the natural water course. Location and design details are given in the Manual.

### **9.2 Recommended Soil and Water Conservation Measures**

The appropriate conservation structures best suited for each of the land parcel/ survey number (Appendix-I) are selected based on the slope per cent, severity of erosion, amount of rainfall, land use and soil type. The different kinds of conservation structures recommended are:

1. Graded / Strengthening of Bunds
2. Trench cum Bunds (TCB)
3. Trench cum Bunds / Strengthening
4. Crescent Bunds

A map (Fig. 9.1) showing soil and water conservation plan with different kinds of structures recommended has been prepared which shows the spatial distribution and extent of area. An area of about 155 ha (17%) needs Trench Cum Bunding, maximum area of about 648 ha (73%) needs Graded Bunding and an area of 13 ha (2%) needs Strengthening of existing bunds. The conservation plan prepared may be presented to all the stakeholders including farmers and after considering their suggestions, the conservation plan for the microwatershed may be finalised in a participatory approach.




Fig. 9.1 Soil and Water Conservation Plan map of Raisabad Hosalli-1 Microwatershed

### 9.3 Greening of Microwatershed

As part of the greening programme in the watersheds, it is envisaged to plant a variety of horticultural and other tree plants that are edible, economical and produce lot of biomass which helps to restore the ecological balance in the watersheds. The lands that are suitable for greening programme are non-arable lands (land capability classes V, VI VII and VIII) and also the lands that are not suitable or marginally suitable and field bunds for growing annual and perennial crops. The method of planting these trees is given below.

It is recommended to open pits during the 1<sup>st</sup> week of March along the contour and heap the dugout soil on the lower side of the slope in order to harness the flowing water and facilitate weathering of soil in the pit. Exposure of soil in the pit also prevents spread of pests and diseases due to scorching sun rays. The pits should be filled with mixture of soil and organic manure during the second week of April and keep ready with sufficiently tall seedlings produced either in poly bags or in root trainer nurseries so that planting can be done during the 2<sup>nd</sup> or 3<sup>rd</sup> week of April depending on the rainfall.

The tree species suitable for the area considering rainfall, temperature and adaptability is listed below; waterlogged areas are recommended to be planted with species like Nerale (*Syzgium cumini*) and Bamboo. Dry areas are to be planted with species like Honge, Bevu, Seetaphal *etc.*



<b>Dry Deciduous Species</b>			<b>Temp (°C)</b>	<b>Rainfall (mm)</b>
1.	Bevu	<i>Azadiracta indica</i>	21–32	400 –1,200
2.	Tapasi	<i>Holoptelia integrifolia</i>	20-30	500 - 1000
3.	Seetaphal	<i>Anona Squamosa</i>	20-40	400 - 1000
4.	Honge	<i>Pongamia pinnata</i>	20 -50	500– 2,500
5.	Kamara	<i>Hardwickia binata</i>	25 -35	400 - 1000
6.	Bage	<i>Albezzia lebbek</i>	20 - 45	500 - 1000
7.	Ficus	<i>Ficus bengalensis</i>	20 - 50	500–2,500
8.	Sisso	<i>Dalbargia Sissoo</i>	20 - 50	500 -2000
9.	Ailanthus	<i>Ailanthus excelsa</i>	20 - 50	500 - 1000
10.	Hale	<i>Wrightia tinctoria</i>	25 - 45	500 - 1000
11.	Uded	<i>Steriospermum chelanoides</i>	25 - 45	500 -2000
12.	Dhupa	<i>Boswella Serrata</i>	20 - 40	500 - 2000
13.	Nelli	<i>Embllica Officinalis</i>	20 - 50	500 -1500
14.	Honne	<i>Pterocarpus marsupium</i>	20 - 40	500 - 2000
<b>Moist Deciduous Species</b>			<b>Temp (°C)</b>	<b>Rainfall (mm)</b>
15.	Teak	<i>Tectona grandis</i>	20 - 50	500-5000
16.	Nandi	<i>Legarstroemia lanceolata</i>	20 - 40	500 - 4000
17.	Honne	<i>Pterocarpus marsupium</i>	20 - 40	500 - 3000
18.	Mathi	<i>Terminalia alata</i>	20 -50	500 - 2000
19.	Shivane	<i>Gmelina arborea</i>	20 -50	500 -2000
20.	Kindal	<i>T.Paniculata</i>	20 - 40	500 - 1500
21.	Beete	<i>Dalbargia latifolia</i>	20 - 40	500 - 1500
22.	Tare	<i>T. belerica</i>	20 - 40	500 - 2000
23.	Bamboo	<i>Bambusa arundinasia</i>	20 - 40	500 - 2500
24.	Bamboo	<i>Dendrocalamus strictus</i>	20 – 40	500 – 2500
25.	Muthuga	<i>Butea monosperma</i>	20 - 40	400 - 1500
26.	Hippe	<i>Madhuca latifolia</i>	20 - 40	500 - 2000
27.	Sandal	<i>Santalum album</i>	20 - 50	400 - 1000
28.	Nelli	<i>Embllica officinalis</i>	20 - 40	500 - 2000
29.	Nerale	<i>Sizygium cumini</i>	20 - 40	500 - 2000
30.	Dhaman	<i>Grevia tilifolia</i>	20 - 40	500 - 2000
31.	Kaval	<i>Careya arborea</i>	20 - 40	500 - 2000
32.	Harada	<i>Terminalia chebula</i>	20 - 40	500 - 2000



## References

1. FAO (1976) Framework for Land Evaluation, Food and Agriculture Organization, Rome. 72 pp.
2. FAO (1983) Guidelines for Land Evaluation for Rainfed Agriculture, FAO, Rome, 237 pp.
3. IARI (1971) Soil Survey Manual, All India Soil and Land Use Survey Organization, IARI, New Delhi, 121 pp.
4. Katyal, J.C. and Rattan, R.K. (2003) Secondary and Micronutrients; Research Gap and Future Needs. Fert. News 48 (4); 9-20.
5. Naidu, L.G.K., Ramamurthy, V., Challa, O., Hegde, R. and Krishnan, P. (2006) Manual Soil Site Suitability Criteria for Major Crops, NBSS Publ. No. 129, NBSS & LUP, Nagpur, 118 pp.
6. Natarajan, A. and Dipak Sarkar (2010) Field Guide for Soil Survey, National Bureau of Soil Survey and Land Use Planning (ICAR), Nagpur, India.
7. Natarajan, A., Rajendra Hegde, Raj, J.N. and Shivananda Murthy, H.G. (2015) Implementation Manual for Sujala-III Project, Watershed Development Department, Bengaluru, Karnataka.
8. Sarma, V.A.K., Krishnan, P. and Budihal, S.L. (1987) Laboratory Manual, Tech. Bull. 23, NBSS & LUP, Nagpur.
9. Sehgal, J.L. (1990) Soil Resource Mapping of Different States of India; Why and How?, National Bureau of Soil Survey and Land Use Planning, Nagpur, 49 pp.
10. Shivaprasad, C.R., R.S. Reddy, J. Sehgal and M. Velayuthum (1998) Soils of Karnataka for Optimizing Land Use, NBSS Publ. No. 47b, NBSS & LUP, Nagpur, India.
11. Soil Survey Staff (2006) Keys to Soil Taxonomy, Tenth edition, U.S. Department of Agriculture/ NRCS, Washington DC, U.S.A.
12. Soil Survey Staff (2012) Soil Survey Manual, Handbook No. 18, USDA, Washington DC, USA.



**Appendix I**  
**Raisabad Hosalli (1J1c) Microwatershed**  
**Soil Phase Information**

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Gravelliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Halagera	156	0.11	KBDhB2	LMU-4	Moderately deep (75-100 cm)	Sandy clay loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Trench cum bunding
Halagera	157	2.96	KBDhB2	LMU-4	Moderately deep (75-100 cm)	Sandy clay loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Trench cum bunding
Halagera	158	0.21	HLGcB2	LMU-8	Moderately shallow (50-75 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Halagera	159/1	0.94	MDGiB2	LMU-5	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Halagera	159/2	0.73	KBDhB2	LMU-4	Moderately deep (75-100 cm)	Sandy clay loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Trench cum bunding
Halagera	159/3	1.64	KBDhB2	LMU-4	Moderately deep (75-100 cm)	Sandy clay loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Trench cum bunding
Halagera	160	0.25	KBDhB2	LMU-4	Moderately deep (75-100 cm)	Sandy clay loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Trench cum bunding
Halagera	164	1.44	HLGcB2	LMU-8	Moderately shallow (50-75 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Halagera	165	2.22	HLGcB2	LMU-8	Moderately shallow (50-75 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Halagera	166	2.81	KBDhB2	LMU-4	Moderately deep (75-100 cm)	Sandy clay loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Trench cum bunding
Halagera	167	2.29	KBDhB2	LMU-4	Moderately deep (75-100 cm)	Sandy clay loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Trench cum bunding
Halagera	168	6.16	KBDhB2	LMU-4	Moderately deep (75-100 cm)	Sandy clay loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Trench cum bunding
Halagera	169	4.97	KBDhB2	LMU-4	Moderately deep (75-100 cm)	Sandy clay loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Trench cum bunding
Halagera	170	3.58	KBDhB2	LMU-4	Moderately deep (75-100 cm)	Sandy clay loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	IIes	Trench cum bunding
Halagera	171	1.19	KBDhB2	LMU-4	Moderately deep (75-100 cm)	Sandy clay loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut (Gn)	Not Available	IIes	Trench cum bunding
Halagera	185	1.11	SGRcB2	LMU-3	Very deep (>150 cm)	Sandy loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IIws	Graded bunding
Halagera	186	2.05	HLGcB2	LMU-8	Moderately shallow (50-75 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Graded bunding
Halagera	187	2.36	KBDhB2	LMU-4	Moderately deep (75-100 cm)	Sandy clay loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	IIes	Trench cum bunding
Halagera	188	6.04	KBDhB2	LMU-4	Moderately deep (75-100 cm)	Sandy clay loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	IIes	Trench cum bunding
Halagera	189	3.52	KBDhB2	LMU-4	Moderately deep (75-100 cm)	Sandy clay loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Jowar (Jw)	Not Available	IIes	Trench cum bunding
Halagera	190	5.73	KBDhB2	LMU-4	Moderately deep (75-100 cm)	Sandy clay loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Trench cum bunding
Halagera	191	3.98	KBDhB2	LMU-4	Moderately deep (75-100 cm)	Sandy clay loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Trench cum bunding

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Graveliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Halagera	192	5.38	SGRcB2	LMU-3	Very deep (>150 cm)	Sandy loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIws	Graded bunding
Halagera	193	8.6	SGRcB2	LMU-3	Very deep (>150 cm)	Sandy loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	1 Borewell	IIws	Graded bunding
Halagera	194	7.19	SGRcB2	LMU-3	Very deep (>150 cm)	Sandy loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIws	Graded bunding
Halagera	195/1	5.87	SGRcB2	LMU-3	Very deep (>150 cm)	Sandy loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIws	Graded bunding
Halagera	195/2	1.95	SGRcB2	LMU-3	Very deep (>150 cm)	Sandy loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIws	Graded bunding
Halagera	196	6.86	KBDiB2	LMU-4	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	IIIs	Trench cum bunding
Halagera	197	8.02	KBDiB2	LMU-4	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IIIs	Trench cum bunding
Halagera	198	6.77	SGRcB2	LMU-3	Very deep (>150 cm)	Sandy loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IIws	Graded bunding
Halagera	199	25.51	KDHcB2	LMU-3	Moderately deep (75-100 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut+Redgram (Gn+Rg)	Not Available	IIws	Graded bunding
Halagera	200	4.52	SGRcB2	LMU-3	Very deep (>150 cm)	Sandy loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Scrub land (Sl)	Not Available	IIws	Graded bunding
Halagera	201	19.64	Waterbody	Others	Others	Others	Others	Others	Others	Others	Waterbody	Not Available	Others	Others
Halagera	202	4.68	KDHcB2	LMU-3	Moderately deep (75-100 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIws	Graded bunding
Halagera	203	9.28	JNKC2	LMU-8	Moderately shallow (50-75 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut+Fallow land (Gn+Fl)	Not Available	IIes	Graded bunding
Halagera	204	1.61	KBDiB2	LMU-4	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	IIIs	Trench cum bunding
Halagera	205	5.61	KBDiB2	LMU-4	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIIs	Trench cum bunding
Halagera	206	8.04	KBDiB2	LMU-4	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	IIIs	Trench cum bunding
Halagera	207	4.72	KBDiB2	LMU-4	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	IIIs	Trench cum bunding
Halagera	208	3.33	KBDiB2	LMU-4	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	IIIs	Trench cum bunding
Halagera	209	2.76	KBDiB2	LMU-4	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	IIIs	Trench cum bunding
Halagera	210	5.69	KBDiB2	LMU-4	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	IIIs	Trench cum bunding
Halagera	211	0.09	KBDiB2	LMU-4	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut+Redgram (Gn+Rg)	Not Available	IIIs	Trench cum bunding
Halagera	212	0.16	KBDiB2	LMU-4	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut+Redgram (Gn+Rg)	Not Available	IIIs	Trench cum bunding
Halagera	213	7.75	KBDiB2	LMU-4	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut+Redgram (Gn+Rg)	Not Available	IIIs	Trench cum bunding
Halagera	214	5	KBDiB2	LMU-4	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIIs	Trench cum bunding

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Graveliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Halagera	215	3.41	KBDiB2	LMU-4	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIes	Trench cum bunding
Halagera	216	3	JNKcB2	LMU-8	Moderately shallow (50-75 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iies	Graded bunding
Halagera	217	5.19	JNKcB2	LMU-8	Moderately shallow (50-75 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iies	Graded bunding
Halagera	218	3.96	JNKcB2	LMU-8	Moderately shallow (50-75 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iies	Graded bunding
Halagera	219	0.31	JNKcB2	LMU-8	Moderately shallow (50-75 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iies	Graded bunding
Halagera	221	0.08	JNKcB2	LMU-8	Moderately shallow (50-75 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iies	Graded bunding
Halagera	222	4.87	JNKcB2	LMU-8	Moderately shallow (50-75 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Fallow land (Fl)	Not Available	Iies	Graded bunding
Halagera	223	0.91	KDHcB2	LMU-3	Moderately deep (75-100 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIws	Graded bunding
Halagera	224	0.03	JNKcB2	LMU-8	Moderately shallow (50-75 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iies	Graded bunding
Halagera	383	0.01	Waterbody	Others	Others	Others	Others	Others	Others	Others	Paddy (Pd)	Not Available	Others	Others
Halagera	384	0	Waterbody	Others	Others	Others	Others	Others	Others	Others	Paddy (Pd)	Not Available	Others	Others
Halagera	386	0.67	Waterbody	Others	Others	Others	Others	Others	Others	Others	Paddy (Pd)	Not Available	Others	Others
Halagera	410	0	SGRcB2	LMU-3	Very deep (>150 cm)	Sandy loam	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Scrub land (Sl)	Not Available	IIws	Graded bunding
Jinatera	253	0.39	YLRcC3	LMU-7	Moderately shallow (50-75 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Gently sloping (3-5%)	Severe	Groundnut (Gn)	Not Available	IIies	Trench cum bunding
Jinatera	256	0.18	YLRcC3	LMU-7	Moderately shallow (50-75 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Gently sloping (3-5%)	Severe	Groundnut (Gn)	Not Available	IIies	Trench cum bunding
Jinatera	258	0	YLRcC3	LMU-7	Moderately shallow (50-75 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Gently sloping (3-5%)	Severe	Groundnut (Gn)	Not Available	IIies	Trench cum bunding
Mylapura	51	0.34	SGRmB2	LMU-3	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Waterbody	Not Available	IIws	Graded bunding
Mylapura	52/1	1.19	BLCiB2	LMU-6	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iies	Trench cum bunding
Mylapura	52/2	2.31	BLCiB2	LMU-6	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iies	Trench cum bunding
Mylapura	53	3.43	SGRmB2	LMU-3	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram (Gg)	Not Available	IIws	Graded bunding
Mylapura	54	3.8	BLCiB2	LMU-6	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Iies	Trench cum bunding
Mylapura	55	5.11	BLCiB2	LMU-6	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram (Gg)	Not Available	Iies	Trench cum bunding
Mylapura	56	4.28	BLCiB2	LMU-6	Moderately deep (75-100 cm)	Sandy clay	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram (Gg)	Not Available	Iies	Trench cum bunding
Mylapura	57	1.04	SGRmB2	LMU-3	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram (Gg)	Not Available	IIws	Graded bunding

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Graveliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Mylapura	58	3.38	SGRmB2	LMU-3	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram (Gg)	Not Available	IIws	Graded bunding
Mylapura	59	1.3	SGRmB2	LMU-3	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram (Gg)	Not Available	IIws	Graded bunding
Mylapura	60	0.37	SGRmB2	LMU-3	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram (Gg)	Not Available	IIws	Graded bunding
Mylapura	61	3.09	SGRmB2	LMU-3	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram (Gg)	Not Available	IIws	Graded bunding
Mylapura	62	3.81	SGRmB2	LMU-3	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram (Gg)	Not Available	IIws	Graded bunding
Mylapura	63	0.54	SGRmB2	LMU-3	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram (Gg)	Not Available	IIws	Graded bunding
Mylapura	64	1.83	SGRmB2	LMU-3	Very deep (>150 cm)	Clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram (Gg)	Not Available	IIws	Graded bunding
Mylapura	123	0.93	MDGhB2g1	LMU-5	Deep (100-150 cm)	Sandy clay loam	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IIes	Graded bunding
Mylapura	124	0.75	MDGhB2g1	LMU-5	Deep (100-150 cm)	Sandy clay loam	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IIes	Graded bunding
Mylapura	125	1.59	MDGhB2g1	LMU-5	Deep (100-150 cm)	Sandy clay loam	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IIes	Graded bunding
Mylapura	126	1.39	MDGhB2g1	LMU-5	Deep (100-150 cm)	Sandy clay loam	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IIes	Graded bunding
Mylapura	127	0.7	MDGhB2g1	LMU-5	Deep (100-150 cm)	Sandy clay loam	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IIes	Graded bunding
Mylapura	128	0.18	MDGhB2g1	LMU-5	Deep (100-150 cm)	Sandy clay loam	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IIes	Graded bunding
Mylapura	131	0.23	MDGhB2g1	LMU-5	Deep (100-150 cm)	Sandy clay loam	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IIes	Graded bunding
Naglapura	195	1.37	BDLiB2	LMU-9	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut (Gn)	Not Available	IIes	Graded bunding
Naglapura	196	2.46	YDRcB2	LMU-2	Deep (100-150 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut (Gn)	Not Available	IIes	Graded bunding
Naglapura	205	1.1	YDRcB2	LMU-2	Deep (100-150 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut (Gn)	Not Available	IIes	Graded bunding
Naglapura	206	5.65	BDLiB2	LMU-9	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Groundnut (Gn)	Not Available	IIes	Graded bunding
Naglapura	207	4.54	BDLiB2	LMU-9	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton+Groundnut (Ct+Gn)	Not Available	IIes	Graded bunding
Panchashe elanagara	46	0	PGPcB2	LMU-6	Moderately deep (75-100 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Scrub land (Sl)	Not Available	IIes	Trench cum bunding
Panchashe elanagara	47	5.62	KDPhB2	LMU-1	Deep (100-150 cm)	Sandy clay loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram+Cow pea (Gg+Cp)	Not Available	IVs	Graded bunds
Panchashe elanagara	48	4.36	KDPhB2	LMU-1	Deep (100-150 cm)	Sandy clay loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Fallow land (Fl)	Not Available	IVs	Graded bunds
Panchashe elanagara	49	1.57	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Not Available	Ro	Ro
Risabadha Hosalli	1	1.99	Habitation	Others	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others



Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Graveliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Risabadha Hosalli	2	0.82	Waterbody	Others	Others	Others	Others	Others	Others	Others	Waterbody	Not Available	Others	Others
Risabadha Hosalli	3	2.91	Waterbody	Others	Others	Others	Others	Others	Others	Others	Waterbody	Not Available	Others	Others
Risabadha Hosalli	4	5.14	MDGhB2g1	LMU-5	Deep (100-150 cm)	Sandy clay loam	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Waterbody	Not Available	Iles	Graded bunding
Risabadha Hosalli	5	0.56	Waterbody	Others	Others	Others	Others	Others	Others	Others	Paddy (Pd)	Not Available	Others	Others
Risabadha Hosalli	6	0.72	Waterbody	Others	Others	Others	Others	Others	Others	Others	Not Available (NA)	Not Available	Others	Others
Risabadha Hosalli	7	0.93	MDGiB2	LMU-5	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Risabadha Hosalli	8	0.18	Waterbody	Others	Others	Others	Others	Others	Others	Others	Not Available (NA)	Not Available	Others	Others
Risabadha Hosalli	9	0.67	Habitation	Others	Others	Others	Others	Others	Others	Others	Not Available (NA)	Not Available	Others	Others
Risabadha Hosalli	10	0.91	MDGiB2	LMU-5	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Risabadha Hosalli	11	4.96	MDGiB2	LMU-5	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Risabadha Hosalli	12	3.96	MDGiB2	LMU-5	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Risabadha Hosalli	13	5.03	MDGiB2	LMU-5	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Paddy (Rg+Pd)	Not Available	Iles	Graded bunding
Risabadha Hosalli	14	5.97	MDGiB2	LMU-5	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Iles	Graded bunding
Risabadha Hosalli	15	3.11	KBDhB2	LMU-4	Moderately deep (75-100 cm)	Sandy clay loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	IIles	Trench cum bunding
Risabadha Hosalli	16	5.48	MDGiB2	LMU-5	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Risabadha Hosalli	17	4.42	MDGiB2	LMU-5	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Risabadha Hosalli	18	3.3	MDGiB2	LMU-5	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Risabadha Hosalli	19	4.54	MDGiB2	LMU-5	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Risabadha Hosalli	20	3.17	MDGiB2	LMU-5	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Risabadha Hosalli	21	3.46	MDGiB2	LMU-5	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Risabadha Hosalli	22	4.76	MDGhB2g1	LMU-5	Deep (100-150 cm)	Sandy clay loam	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Risabadha Hosalli	23	8.39	MDGhB2g1	LMU-5	Deep (100-150 cm)	Sandy clay loam	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Paddy (Rg+Pd)	Not Available	Iles	Graded bunding
Risabadha Hosalli	24	5.6	MDGhB2g1	LMU-5	Deep (100-150 cm)	Sandy clay loam	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Risabadha Hosalli	25	5.97	MDGhB2g1	LMU-5	Deep (100-150 cm)	Sandy clay loam	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Graveliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Risabadha Hosalli	26	0.09	BDLiB2	LMU-9	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton+Not Available (Rg)	Not Available	Iles	Graded bunding
Risabadha Hosalli	27	0.27	MDGhB2g1	LMU-5	Deep (100-150 cm)	Sandy clay loam	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Risabadha Hosalli	28	4.35	MDGhB2g1	LMU-5	Deep (100-150 cm)	Sandy clay loam	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Risabadha Hosalli	29	1.52	MDGhB2g1	LMU-5	Deep (100-150 cm)	Sandy clay loam	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Risabadha Hosalli	30	4.66	MDGhB2g1	LMU-5	Deep (100-150 cm)	Sandy clay loam	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Risabadha Hosalli	31	4.06	MDGhB2g1	LMU-5	Deep (100-150 cm)	Sandy clay loam	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram+Paddy (Rg+Pd)	Not Available	Iles	Graded bunding
Risabadha Hosalli	32	6.9	MDGhB2g1	LMU-5	Deep (100-150 cm)	Sandy clay loam	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Risabadha Hosalli	33	5.65	MDGhB2g1	LMU-5	Deep (100-150 cm)	Sandy clay loam	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Risabadha Hosalli	34	1.36	MDGhB2g1	LMU-5	Deep (100-150 cm)	Sandy clay loam	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Risabadha Hosalli	35	0.69	MDGhB2g1	LMU-5	Deep (100-150 cm)	Sandy clay loam	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Risabadha Hosalli	45	1.02	JNKiB2g1	LMU-8	Moderately shallow (50-75 cm)	Sandy clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram (Gg)	Not Available	Iles	Graded bunding
Risabadha Hosalli	46/1	2.53	JNKiB2g1	LMU-8	Moderately shallow (50-75 cm)	Sandy clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Risabadha Hosalli	46/10	0.68	JNKiB2g1	LMU-8	Moderately shallow (50-75 cm)	Sandy clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Risabadha Hosalli	46/4	0.07	MDGiB2	LMU-5	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Risabadha Hosalli	46/5	0.18	JNKiB2g1	LMU-8	Moderately shallow (50-75 cm)	Sandy clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Risabadha Hosalli	46/6	0.2	JNKiB2g1	LMU-8	Moderately shallow (50-75 cm)	Sandy clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Risabadha Hosalli	46/7	0.37	JNKiB2g1	LMU-8	Moderately shallow (50-75 cm)	Sandy clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Risabadha Hosalli	46/8	0.86	JNKiB2g1	LMU-8	Moderately shallow (50-75 cm)	Sandy clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Risabadha Hosalli	46/9	0.58	JNKiB2g1	LMU-8	Moderately shallow (50-75 cm)	Sandy clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Risabadha Hosalli	47	0.59	MDGiB2	LMU-5	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Risabadha Hosalli	48	7.22	MDGiB2	LMU-5	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram (Gg)	Not Available	Iles	Graded bunding
Risabadha Hosalli	49	7.68	JNKiB2g1	LMU-8	Moderately shallow (50-75 cm)	Sandy clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Iles	Graded bunding
Risabadha Hosalli	50	4.11	JNKiB2g1	LMU-8	Moderately shallow (50-75 cm)	Sandy clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram (Gg)	Not Available	Iles	Graded bunding
Risabadha Hosalli	51	10.18	JNKiB2g1	LMU-8	Moderately shallow (50-75 cm)	Sandy clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Iles	Graded bunding

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Graveliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Risabadha Hosalli	52	2.72	JNKiB2g1	LMU-8	Moderately shallow (50-75 cm)	Sandy clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Iles	Graded bunding
Risabadha Hosalli	53	6.84	JNKiB2g1	LMU-8	Moderately shallow (50-75 cm)	Sandy clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton+Redgram (Ct+Rg)	Not Available	Iles	Graded bunding
Risabadha Hosalli	54	7.21	JNKiB2g1	LMU-8	Moderately shallow (50-75 cm)	Sandy clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Risabadha Hosalli	55	8.36	MDGiB2	LMU-5	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton+Paddy+Redgram (Ct+Pd+Rg)	Not Available	Iles	Graded bunding
Risabadha Hosalli	56	7.89	MDGiB2	LMU-5	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Risabadha Hosalli	57	4.61	MDGiB2	LMU-5	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Risabadha Hosalli	58	6.43	BDLiB2	LMU-9	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIles	Graded bunding
Risabadha Hosalli	59	7.85	MDGiB2	LMU-5	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Risabadha Hosalli	60	9.29	MDGiB2	LMU-5	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton+Redgram (Ct+Rg)	Not Available	Iles	Graded bunding
Risabadha Hosalli	61	4.32	Waterbody	Others	Others	Others	Others	Others	Others	Others	Not Available (NA)	Not Available	Others	Others
Risabadha Hosalli	62	1.35	Waterbody	Others	Others	Others	Others	Others	Others	Others	Waterbody	Not Available	Others	Others
Risabadha Hosalli	63	4.76	Waterbody	Others	Others	Others	Others	Others	Others	Others	Waterbody	Not Available	Others	Others
Risabadha Hosalli	64	8.56	MDGiB2	LMU-5	Deep (100-150 cm)	Sandy clay	Non gravelly (<15%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Risabadha Hosalli	65	3.56	JNKiB2g1	LMU-8	Moderately shallow (50-75 cm)	Sandy clay	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Risabadha Hosalli	66	6.56	HTKbB2g1	LMU-9	Shallow (25-50 cm)	Loamy sand	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IIles	Graded bunding
Risabadha Hosalli	67	7.23	HTKbB2g1	LMU-9	Shallow (25-50 cm)	Loamy sand	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	IIles	Graded bunding
Risabadha Hosalli	68	1.84	HTKbB2g1	LMU-9	Shallow (25-50 cm)	Loamy sand	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton+Ragi+Habitation (Ct+Rg+Hb)	Not Available	IIles	Graded bunding
Risabadha Hosalli	69	0.01	HTKbB2g1	LMU-9	Shallow (25-50 cm)	Loamy sand	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram (Gg)	Not Available	IIles	Graded bunding
Risabadha Hosalli	71	0.51	HTKbB2g1	LMU-9	Shallow (25-50 cm)	Loamy sand	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIles	Graded bunding
Risabadha Hosalli	72	2.92	HTKbB2g1	LMU-9	Shallow (25-50 cm)	Loamy sand	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton+Groundnut (Ct+Gn)	Not Available	IIles	Graded bunding
Risabadha Hosalli	73	8.28	HTKbB2g1	LMU-9	Shallow (25-50 cm)	Loamy sand	Gravelly (15-35%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	IIles	Graded bunding
Risabadha Hosalli	74	7.13	YDRcB2g1	LMU-2	Deep (100-150 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Iles	Graded bunding
Risabadha Hosalli	75	9.36	YDRcB2g1	LMU-2	Deep (100-150 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Iles	Graded bunding
Risabadha Hosalli	76	2.76	YDRcB2g1	LMU-2	Deep (100-150 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Iles	Graded bunding

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Graveliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Risabadha Hosalli	77	4.98	YDRcB2g1	LMU-2	Deep (100-150 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram (Gg)	Not Available	Iles	Graded bunding
Risabadha Hosalli	78	5.63	YDRcB2g1	LMU-2	Deep (100-150 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Banana (Ba)	Not Available	Iles	Graded bunding
Risabadha Hosalli	79	4.11	YDRcB2	LMU-2	Deep (100-150 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Risabadha Hosalli	80	0.67	YDRcB2	LMU-2	Deep (100-150 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram (Gg)	Not Available	Iles	Graded bunding
Risabadha Hosalli	82	0.19	MDGhB2g1	LMU-5	Deep (100-150 cm)	Sandy clay loam	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram (Gg)	Not Available	Iles	Graded bunding
Risabadha Hosalli	83	7.39	YDRcB2	LMU-2	Deep (100-150 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Risabadha Hosalli	84	8.6	YDRcB2	LMU-2	Deep (100-150 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram (Gg)	Not Available	Iles	Graded bunding
Risabadha Hosalli	85	7.05	YDRcB2	LMU-2	Deep (100-150 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Iles	Graded bunding
Risabadha Hosalli	86	6.32	YDRcB2	LMU-2	Deep (100-150 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Risabadha Hosalli	87	5.26	YDRcB2	LMU-2	Deep (100-150 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram (Gg)	Not Available	Iles	Graded bunding
Risabadha Hosalli	88	3.55	YDRcB2	LMU-2	Deep (100-150 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Iles	Graded bunding
Risabadha Hosalli	89	5.72	YDRcB2	LMU-2	Deep (100-150 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Iles	Graded bunding
Risabadha Hosalli	90	7	BDLiB2	LMU-9	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIles	Graded bunding
Risabadha Hosalli	91	1.58	BDLiB2	LMU-9	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIles	Graded bunding
Risabadha Hosalli	92	4.61	BDLiB2	LMU-9	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IIles	Graded bunding
Risabadha Hosalli	93	3.82	BDLiB2	LMU-9	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IIles	Graded bunding
Risabadha Hosalli	94	6.15	BDLiB2	LMU-9	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	IIles	Graded bunding
Risabadha Hosalli	95	7.03	BDLiB2	LMU-9	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy+Cotton (Pd+Ct)	Not Available	IIles	Graded bunding
Risabadha Hosalli	96	5.22	BDLiB2	LMU-9	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IIles	Graded bunding
Risabadha Hosalli	97	6.46	BDLiB2	LMU-9	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIles	Graded bunding
Risabadha Hosalli	98	0.96	BDLiB2	LMU-9	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IIles	Graded bunding
Risabadha Hosalli	99	0.9	BDLiB2	LMU-9	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IIles	Graded bunding
Risabadha Hosalli	100	0.32	BDLiB2	LMU-9	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	IIles	Graded bunding
Risabadha Hosalli	101	9.37	BDLiB2	LMU-9	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IIles	Graded bunding

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Graveliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Risabadha Hosalli	102	7.63	BDLiB2	LMU-9	Shallow (25-50 cm)	Sandy clay	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton+Greengram (Ct+Gg)	Not Available	Iles	Graded bunding
Risabadha Hosalli	103	8.68	YDRcB2	LMU-2	Deep (100-150 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram (Gg)	Not Available	Iles	Graded bunding
Risabadha Hosalli	104	7.85	YDRcB2	LMU-2	Deep (100-150 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Risabadha Hosalli	105	3.07	YDRcB2	LMU-2	Deep (100-150 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Risabadha Hosalli	106	0.66	YDRcB2	LMU-2	Deep (100-150 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Risabadha Hosalli	107	0.65	YDRcB2	LMU-2	Deep (100-150 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Risabadha Hosalli	108	0.84	YDRcB2	LMU-2	Deep (100-150 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Risabadha Hosalli	109	1.83	YDRcB2	LMU-2	Deep (100-150 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Risabadha Hosalli	110	2.68	YDRcB2	LMU-2	Deep (100-150 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Risabadha Hosalli	111	8.79	YDRcB2	LMU-2	Deep (100-150 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton+Ragi (Ct+Rg)	Not Available	Iles	Graded bunding
Risabadha Hosalli	112	1.04	YDRcB2	LMU-2	Deep (100-150 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Iles	Graded bunding
Risabadha Hosalli	113	4.12	YDRcB2	LMU-2	Deep (100-150 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Risabadha Hosalli	114	4.28	YDRcB2	LMU-2	Deep (100-150 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Risabadha Hosalli	115	6.41	YDRcB2	LMU-2	Deep (100-150 cm)	Sandy loam	Non gravelly (<15%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Iles	Graded bunding
Risabadha Hosalli	116	6.37	YDRcB2g1	LMU-2	Deep (100-150 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Risabadha Hosalli	117	0.76	YDRcB2g1	LMU-2	Deep (100-150 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Risabadha Hosalli	118	0.4	YDRcB2g1	LMU-2	Deep (100-150 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Risabadha Hosalli	119	1.1	YDRcB2g1	LMU-2	Deep (100-150 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Risabadha Hosalli	120	0.51	YDRcB2g1	LMU-2	Deep (100-150 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iles	Graded bunding
Risabadha Hosalli	121	0.81	YDRcB2g1	LMU-2	Deep (100-150 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Risabadha Hosalli	122	1.18	YDRcB2g1	LMU-2	Deep (100-150 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Risabadha Hosalli	123	1.22	YDRcB2g1	LMU-2	Deep (100-150 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Risabadha Hosalli	124	0.87	YDRcB2g1	LMU-2	Deep (100-150 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Risabadha Hosalli	125	0.89	YDRcB2g1	LMU-2	Deep (100-150 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Graveliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Risabadha Hosalli	126	1.34	YDRcB2g1	LMU-2	Deep (100-150 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Risabadha Hosalli	127	6.79	YDRcB2g1	LMU-2	Deep (100-150 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Greengram (Gg)	Not Available	Iles	Graded bunding
Risabadha Hosalli	128	5.17	YDRcB2g1	LMU-2	Deep (100-150 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Iles	Graded bunding
Risabadha Hosalli	129	0.39	YDRcB2g1	LMU-2	Deep (100-150 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Risabadha Hosalli	130	0.3	YDRcB2g1	LMU-2	Deep (100-150 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Risabadha Hosalli	131	0.66	YDRcB2g1	LMU-2	Deep (100-150 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Risabadha Hosalli	132	0.5	YDRcB2g1	LMU-2	Deep (100-150 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Risabadha Hosalli	133	0.62	YDRcB2g1	LMU-2	Deep (100-150 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Risabadha Hosalli	134	0.68	YDRcB2g1	LMU-2	Deep (100-150 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Risabadha Hosalli	135	1.14	YDRcB2g1	LMU-2	Deep (100-150 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Risabadha Hosalli	136	1.12	YDRcB2g1	LMU-2	Deep (100-150 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Risabadha Hosalli	137	0.7	YDRcB2g1	LMU-2	Deep (100-150 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Risabadha Hosalli	138	0.87	YDRcB2g1	LMU-2	Deep (100-150 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Risabadha Hosalli	139	0.75	YDRcB2g1	LMU-2	Deep (100-150 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Risabadha Hosalli	140	0.88	YDRcB2g1	LMU-2	Deep (100-150 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Risabadha Hosalli	141	0.88	YDRcB2g1	LMU-2	Deep (100-150 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Risabadha Hosalli	142	0.89	YDRcB2g1	LMU-2	Deep (100-150 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Risabadha Hosalli	143	0.65	YDRcB2g1	LMU-2	Deep (100-150 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Risabadha Hosalli	144	3.71	YDRcB2g1	LMU-2	Deep (100-150 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Risabadha Hosalli	145	0.41	YDRcB2g1	LMU-2	Deep (100-150 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Risabadha Hosalli	146	0.28	YDRcB2g1	LMU-2	Deep (100-150 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Risabadha Hosalli	147	1.14	YDRcB2g1	LMU-2	Deep (100-150 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Risabadha Hosalli	148	1.23	YDRcB2g1	LMU-2	Deep (100-150 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Risabadha Hosalli	149	0.6	YDRcB2g1	LMU-2	Deep (100-150 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Graveliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Risabadha Hosalli	150	0.74	YDRcB2g1	LMU-2	Deep (100-150 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Risabadha Hosalli	151	0.46	YDRcB2g1	LMU-2	Deep (100-150 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Risabadha Hosalli	152	2.45	YDRcB2g1	LMU-2	Deep (100-150 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Risabadha Hosalli	153	0.61	YDRcB2g1	LMU-2	Deep (100-150 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Risabadha Hosalli	154	4.79	MDGhB2g1	LMU-5	Deep (100-150 cm)	Sandy clay loam	Gravelly (15-35%)	Very high (>200 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Risabadha Hosalli	155	3.93	YDRcB2g1	LMU-2	Deep (100-150 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	Iles	Graded bunding
Risabadha Hosalli	156	0.42	YDRcB2g1	LMU-2	Deep (100-150 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Risabadha Hosalli	157/1	1.23	YDRcB2g1	LMU-2	Deep (100-150 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Risabadha Hosalli	157/2	1.25	YDRcB2g1	LMU-2	Deep (100-150 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Risabadha Hosalli	158	0.71	YDRcB2g1	LMU-2	Deep (100-150 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Risabadha Hosalli	159	0.6	YDRcB2g1	LMU-2	Deep (100-150 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Risabadha Hosalli	160	0.28	YDRcB2g1	LMU-2	Deep (100-150 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Risabadha Hosalli	161	0.19	YDRcB2g1	LMU-2	Deep (100-150 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Risabadha Hosalli	162	0.62	YDRcB2g1	LMU-2	Deep (100-150 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Risabadha Hosalli	163	1.1	YDRcB2g1	LMU-2	Deep (100-150 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Risabadha Hosalli	164	0.39	YDRcB2g1	LMU-2	Deep (100-150 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Risabadha Hosalli	165	0.21	YDRcB2g1	LMU-2	Deep (100-150 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Risabadha Hosalli	166	0.35	YDRcB2g1	LMU-2	Deep (100-150 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Risabadha Hosalli	167	0.98	YDRcB2g1	LMU-2	Deep (100-150 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Risabadha Hosalli	168	0.67	YDRcB2g1	LMU-2	Deep (100-150 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Risabadha Hosalli	169	3.53	YDRcB2g1	LMU-2	Deep (100-150 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	Iles	Graded bunding
Risabadha Hosalli	170	0.55	YDRcB2g1	LMU-2	Deep (100-150 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Risabadha Hosalli	171	0.62	YDRcB2g1	LMU-2	Deep (100-150 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Risabadha Hosalli	172	0.5	YDRcB2g1	LMU-2	Deep (100-150 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding

Village	Survey No	Area (ha)	Soil Phase	LMU	Soil Depth	Surface Soil Texture	Soil Graveliness	Available Water Capacity	Slope	Soil Erosion	Current Land Use	Wells	Land Capability	Conservation Plan
Risabadha Hosalli	173	0.4	YDRcB2g1	LMU-2	Deep (100-150 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Risabadha Hosalli	174	0.71	YDRcB2g1	LMU-2	Deep (100-150 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Risabadha Hosalli	175	0.6	YDRcB2g1	LMU-2	Deep (100-150 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Risabadha Hosalli	176	0.57	YDRcB2g1	LMU-2	Deep (100-150 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iles	Graded bunding
Risabadha Hosalli	177	0.39	YDRcB2g1	LMU-2	Deep (100-150 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Not Available (NA)	Not Available	Iles	Graded bunding
Risabadha Hosalli	178	1.01	YDRcB2g1	LMU-2	Deep (100-150 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Risabadha Hosalli	179	0.35	YDRcB2g1	LMU-2	Deep (100-150 cm)	Sandy loam	Gravelly (15-35%)	Low (51-100 mm/m)	Very gently sloping (1-3%)	Moderate	Paddy (Pd)	Not Available	Iles	Graded bunding
Risabadha Hosalli	180	0.8	Habitation	Others	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others
Risabadha Hosalli	181/A	4.52	KBDhB2	LMU-4	Moderately deep (75-100 cm)	Sandy clay loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	IIles	Trench cum bunding
Risabadha Hosalli	181/B	0.46	Habitation	Others	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others
Risabadha Hosalli	182	2	Habitation	Others	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others
Risabadha Hosalli	183	2.82	KBDhB2	LMU-4	Moderately deep (75-100 cm)	Sandy clay loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Cotton (Ct)	Not Available	IIles	Trench cum bunding
Risabadha Hosalli	184	1.9	Habitation	Others	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others
Risabadha Hosalli	185	4.22	Habitation	Others	Others	Others	Others	Others	Others	Others	Habitation	Not Available	Others	Others
Yaleri	691	2.16	BDPhB2	LMU-9	Very shallow (<25 cm)	Sandy clay loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IVes	Trench cum bunding
Yaleri	693	0.31	BDPcB2	LMU-9	Very shallow (<25 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IVes	Trench cum bunding
Yaleri	694	6.65	BDPcB2	LMU-9	Very shallow (<25 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IVes	Trench cum bunding
Yaleri	695	2.52	BDPcB2	LMU-9	Very shallow (<25 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IVes	Trench cum bunding
Yaleri	696	1.77	BDPcB2	LMU-9	Very shallow (<25 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IVes	Trench cum bunding
Yaleri	697	3.56	BDPcB2	LMU-9	Very shallow (<25 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IVes	Trench cum bunding
Yaleri	698	6.17	BDPcB2	LMU-9	Very shallow (<25 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IVes	Trench cum bunding
Yaleri	699	0.88	BDPcB2	LMU-9	Very shallow (<25 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IVes	Trench cum bunding
Yaleri	702	1.39	BDPcB2	LMU-9	Very shallow (<25 cm)	Sandy loam	Non gravelly (<15%)	Very low (<50 mm/m)	Very gently sloping (1-3%)	Moderate	Redgram (Rg)	Not Available	IVes	Trench cum bunding





























**Appendix III**  
**Raisabad Hosalli1 (1J1c) Microwatershed**  
**Soil Suitability Information**

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthe mum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry	
Halagera	156	S3rg	S3g	S3g	S3g	S3g	S3g	S3rg	S3g	S3g	S3g	S3g	S2g	S3g	S2g	S3g	S3rg	S3g	S2g	S2g	S3g	S3g	S3g	S3g	S3g	S2g	S2g	S2g	S2g	S2g	
Halagera	157	S3rg	S3g	S3g	S3g	S3g	S3g	S3rg	S3g	S3g	S3g	S3g	S2g	S3g	S2g	S3g	S3rg	S3g	S2g	S2g	S3g	S3g	S3g	S3g	S3g	S2g	S2g	S2g	S2g	S2g	
Halagera	158	N1rz	S2rz	S3rz	S2rz	S3rz	S2rz	N1rz	S3rz	S2rz	S3rz	S3rz	S2rz	S3rz	S2rz	N1tz	S3rz	S3rz	S2rz	S2r	S2rz	S2rz	S2rz	S2rz	S3rz	S2rz	S2r	S2r	S3rz	S3rz	
Halagera	159/1	S2r	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S1	S2tw	S2tw	S2tw	S2tw	S2t	S2t	S1	S1	S2tw	S3tw	
Halagera	159/2	S3rg	S3g	S3g	S3g	S3g	S3g	S3rg	S3g	S3g	S3g	S3g	S2g	S3g	S2g	S3g	S3rg	S3g	S2g	S2g	S3g	S3g	S3g	S3g	S3g	S2g	S2g	S2g	S2g	S2g	
Halagera	159/3	S3rg	S3g	S3g	S3g	S3g	S3g	S3rg	S3g	S3g	S3g	S3g	S2g	S3g	S2g	S3g	S3rg	S3g	S2g	S2g	S3g	S3g	S3g	S3g	S3g	S2g	S2g	S2g	S2g	S2g	
Halagera	160	S3rg	S3g	S3g	S3g	S3g	S3g	S3rg	S3g	S3g	S3g	S3g	S2g	S3g	S2g	S3g	S3rg	S3g	S2g	S2g	S3g	S3g	S3g	S3g	S3g	S2g	S2g	S2g	S2g	S2g	
Halagera	164	N1rz	S2rz	S3rz	S2rz	S3rz	S2rz	N1rz	S3rz	S2rz	S3rz	S3rz	S2rz	S3rz	S2rz	N1tz	S3rz	S3rz	S2rz	S2r	S2rz	S2rz	S2rz	S2rz	S3rz	S2rz	S2r	S2r	S3rz	S3rz	
Halagera	165	N1rz	S2rz	S3rz	S2rz	S3rz	S2rz	N1rz	S3rz	S2rz	S3rz	S3rz	S2rz	S3rz	S2rz	N1tz	S3rz	S3rz	S2rz	S2r	S2rz	S2rz	S2rz	S2rz	S3rz	S2rz	S2r	S2r	S3rz	S3rz	
Halagera	166	S3rg	S3g	S3g	S3g	S3g	S3g	S3rg	S3g	S3g	S3g	S3g	S2g	S3g	S2g	S3g	S3rg	S3g	S2g	S2g	S3g	S3g	S3g	S3g	S3g	S2g	S2g	S2g	S2g	S2g	
Halagera	167	S3rg	S3g	S3g	S3g	S3g	S3g	S3rg	S3g	S3g	S3g	S3g	S2g	S3g	S2g	S3g	S3rg	S3g	S2g	S2g	S3g	S3g	S3g	S3g	S3g	S2g	S2g	S2g	S2g	S2g	
Halagera	168	S3rg	S3g	S3g	S3g	S3g	S3g	S3rg	S3g	S3g	S3g	S3g	S2g	S3g	S2g	S3g	S3rg	S3g	S2g	S2g	S3g	S3g	S3g	S3g	S3g	S2g	S2g	S2g	S2g	S2g	
Halagera	169	S3rg	S3g	S3g	S3g	S3g	S3g	S3rg	S3g	S3g	S3g	S3g	S2g	S3g	S2g	S3g	S3rg	S3g	S2g	S2g	S3g	S3g	S3g	S3g	S3g	S2g	S2g	S2g	S2g	S2g	
Halagera	170	S3rg	S3g	S3g	S3g	S3g	S3g	S3rg	S3g	S3g	S3g	S3g	S2g	S3g	S2g	S3g	S3rg	S3g	S2g	S2g	S3g	S3g	S3g	S3g	S3g	S2g	S2g	S2g	S2g	S2g	
Halagera	171	S3rg	S3g	S3g	S3g	S3g	S3g	S3rg	S3g	S3g	S3g	S3g	S2g	S3g	S2g	S3g	S3rg	S3g	S2g	S2g	S3g	S3g	S3g	S3g	S3g	S2g	S2g	S2g	S2g	S2g	
Halagera	185	S3tw	S2tw	S3tw	S1	S3tw	S1	S2tw	S2w	S1	S1	S3tw	S1	S3tw	S1	N1tw	S2tw	S2w	S3tw	S2t	S2tw	S3tw	S2tw	S2tw	S2tw	S2t	S2t	S2t	S2t	S2tw	S3tw
Halagera	186	N1rz	S2rz	S3rz	S2rz	S3rz	S2rz	N1rz	S3rz	S2rz	S3rz	S3rz	S2rz	S3rz	S2rz	N1tz	S3rz	S3rz	S2rz	S2r	S2rz	S2rz	S2rz	S2rz	S3rz	S2rz	S2r	S2r	S3rz	S3rz	
Halagera	187	S3rg	S3g	S3g	S3g	S3g	S3g	S3rg	S3g	S3g	S3g	S3g	S2g	S3g	S2g	S3g	S3rg	S3g	S2g	S2g	S3g	S3g	S3g	S3g	S3g	S2g	S2g	S2g	S2g	S2g	
Halagera	188	S3rg	S3g	S3g	S3g	S3g	S3g	S3rg	S3g	S3g	S3g	S3g	S2g	S3g	S2g	S3g	S3rg	S3g	S2g	S2g	S3g	S3g	S3g	S3g	S3g	S2g	S2g	S2g	S2g	S2g	
Halagera	189	S3rg	S3g	S3g	S3g	S3g	S3g	S3rg	S3g	S3g	S3g	S3g	S2g	S3g	S2g	S3g	S3rg	S3g	S2g	S2g	S3g	S3g	S3g	S3g	S3g	S2g	S2g	S2g	S2g	S2g	
Halagera	190	S3rg	S3g	S3g	S3g	S3g	S3g	S3rg	S3g	S3g	S3g	S3g	S2g	S3g	S2g	S3g	S3rg	S3g	S2g	S2g	S3g	S3g	S3g	S3g	S3g	S2g	S2g	S2g	S2g	S2g	
Halagera	191	S3rg	S3g	S3g	S3g	S3g	S3g	S3rg	S3g	S3g	S3g	S3g	S2g	S3g	S2g	S3g	S3rg	S3g	S2g	S2g	S3g	S3g	S3g	S3g	S3g	S2g	S2g	S2g	S2g	S2g	
Halagera	192	S3tw	S2tw	S3tw	S1	S3tw	S1	S2tw	S2w	S1	S1	S3tw	S1	S3tw	S1	N1tw	S2tw	S2w	S3tw	S2t	S2tw	S3tw	S2tw	S2tw	S2tw	S2t	S2t	S2t	S2t	S2tw	S3tw
Halagera	193	S3tw	S2tw	S3tw	S1	S3tw	S1	S2tw	S2w	S1	S1	S3tw	S1	S3tw	S1	N1tw	S2tw	S2w	S3tw	S2t	S2tw	S3tw	S2tw	S2tw	S2tw	S2t	S2t	S2t	S2t	S2tw	S3tw
Halagera	194	S3tw	S2tw	S3tw	S1	S3tw	S1	S2tw	S2w	S1	S1	S3tw	S1	S3tw	S1	N1tw	S2tw	S2w	S3tw	S2t	S2tw	S3tw	S2tw	S2tw	S2tw	S2t	S2t	S2t	S2t	S2tw	S3tw

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry	
Halagera	195/1	S3tw	S2tw	S3tw	S1	S3tw	S1	S2tw	S2w	S1	S1	S3tw	S1	S3tw	S1	N1tw	S2tw	S2w	S3tw	S2t	S2tw	S3tw	S2tw	S2tw	S2t	S2t	S2t	S2t	S2tw	S3tw	
Halagera	195/2	S3tw	S2tw	S3tw	S1	S3tw	S1	S2tw	S2w	S1	S1	S3tw	S1	S3tw	S1	N1tw	S2tw	S2w	S3tw	S2t	S2tw	S3tw	S2tw	S2tw	S2t	S2t	S2t	S2t	S2tw	S3tw	
Halagera	196	S3rg	S3g	S3g	S3g	S3g	S3g	S3rg	S3g	S3g	S3g	S3g	S2g	S3g	S2g	S3g	S3rg	S3g	S2g	S2g	S3g	S3g	S3g	S3g	S3g	S2g	S2g	S2g	S2g	S2g	
Halagera	197	S3rg	S3g	S3g	S3g	S3g	S3g	S3rg	S3g	S3g	S3g	S3g	S2g	S3g	S2g	S3g	S3rg	S3g	S2g	S2g	S3g	S3g	S3g	S3g	S3g	S2g	S2g	S2g	S2g	S2g	
Halagera	198	S3tw	S2tw	S3tw	S1	S3tw	S1	S2tw	S2w	S1	S1	S3tw	S1	S3tw	S1	N1tw	S2tw	S2w	S3tw	S2t	S2tw	S3tw	S2tw	S2tw	S2tw	S2t	S2t	S2t	S2tw	S3tw	
Halagera	199	S3tw	S2tw	S3tw	S1	S3tw	S2r	S2tw	S2w	S1	S1	S2tw	S2tw	S3tw	S1	N1tz	S2tw	S2w	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2tw	S2t	S1	S1	S2tw	S3tw	
Halagera	200	S3tw	S2tw	S3tw	S1	S3tw	S1	S2tw	S2w	S1	S1	S3tw	S1	S3tw	S1	N1tw	S2tw	S2w	S3tw	S2t	S2tw	S3tw	S2tw	S2tw	S2tw	S2t	S2t	S2t	S2tw	S3tw	
Halagera	201	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Halagera	202	S3tw	S2tw	S3tw	S1	S3tw	S2r	S2tw	S2w	S1	S1	S2tw	S2tw	S3tw	S1	N1tz	S2tw	S2w	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2tw	S2t	S1	S1	S2tw	S3tw	
Halagera	203	N1r	S2t	S3r	S2r	S3r	S2r	N1r	S3r	S2r	S3r	S3r	S2r	S3r	S2r	N1rt	S3r	S3r	S3t	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Halagera	204	S3rg	S3g	S3g	S3g	S3g	S3g	S3rg	S3g	S3g	S3g	S3g	S2g	S3g	S2g	S3g	S3rg	S3g	S2g	S2g	S3g	S3g	S3g	S3g	S3g	S2g	S2g	S2g	S2g	S2g	
Halagera	205	S3rg	S3g	S3g	S3g	S3g	S3g	S3rg	S3g	S3g	S3g	S3g	S2g	S3g	S2g	S3g	S3rg	S3g	S2g	S2g	S3g	S3g	S3g	S3g	S3g	S2g	S2g	S2g	S2g	S2g	
Halagera	206	S3rg	S3g	S3g	S3g	S3g	S3g	S3rg	S3g	S3g	S3g	S3g	S2g	S3g	S2g	S3g	S3rg	S3g	S2g	S2g	S3g	S3g	S3g	S3g	S3g	S2g	S2g	S2g	S2g	S2g	
Halagera	207	S3rg	S3g	S3g	S3g	S3g	S3g	S3rg	S3g	S3g	S3g	S3g	S2g	S3g	S2g	S3g	S3rg	S3g	S2g	S2g	S3g	S3g	S3g	S3g	S3g	S2g	S2g	S2g	S2g	S2g	
Halagera	208	S3rg	S3g	S3g	S3g	S3g	S3g	S3rg	S3g	S3g	S3g	S3g	S2g	S3g	S2g	S3g	S3rg	S3g	S2g	S2g	S3g	S3g	S3g	S3g	S3g	S2g	S2g	S2g	S2g	S2g	
Halagera	209	S3rg	S3g	S3g	S3g	S3g	S3g	S3rg	S3g	S3g	S3g	S3g	S2g	S3g	S2g	S3g	S3rg	S3g	S2g	S2g	S3g	S3g	S3g	S3g	S3g	S2g	S2g	S2g	S2g	S2g	
Halagera	210	S3rg	S3g	S3g	S3g	S3g	S3g	S3rg	S3g	S3g	S3g	S3g	S2g	S3g	S2g	S3g	S3rg	S3g	S2g	S2g	S3g	S3g	S3g	S3g	S3g	S2g	S2g	S2g	S2g	S2g	
Halagera	211	S3rg	S3g	S3g	S3g	S3g	S3g	S3rg	S3g	S3g	S3g	S3g	S2g	S3g	S2g	S3g	S3rg	S3g	S2g	S2g	S3g	S3g	S3g	S3g	S3g	S2g	S2g	S2g	S2g	S2g	
Halagera	212	S3rg	S3g	S3g	S3g	S3g	S3g	S3rg	S3g	S3g	S3g	S3g	S2g	S3g	S2g	S3g	S3rg	S3g	S2g	S2g	S3g	S3g	S3g	S3g	S3g	S2g	S2g	S2g	S2g	S2g	
Halagera	213	S3rg	S3g	S3g	S3g	S3g	S3g	S3rg	S3g	S3g	S3g	S3g	S2g	S3g	S2g	S3g	S3rg	S3g	S2g	S2g	S3g	S3g	S3g	S3g	S3g	S2g	S2g	S2g	S2g	S2g	
Halagera	214	S3rg	S3g	S3g	S3g	S3g	S3g	S3rg	S3g	S3g	S3g	S3g	S2g	S3g	S2g	S3g	S3rg	S3g	S2g	S2g	S3g	S3g	S3g	S3g	S3g	S2g	S2g	S2g	S2g	S2g	
Halagera	215	S3rg	S3g	S3g	S3g	S3g	S3g	S3rg	S3g	S3g	S3g	S3g	S2g	S3g	S2g	S3g	S3rg	S3g	S2g	S2g	S3g	S3g	S3g	S3g	S3g	S2g	S2g	S2g	S2g	S2g	
Halagera	216	N1r	S2t	S3r	S2r	S3r	S2r	N1r	S3r	S2r	S3r	S3r	S2r	S3r	S2r	N1rt	S3r	S3r	S3t	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Halagera	217	N1r	S2t	S3r	S2r	S3r	S2r	N1r	S3r	S2r	S3r	S3r	S2r	S3r	S2r	N1rt	S3r	S3r	S3t	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Halagera	218	N1r	S2t	S3r	S2r	S3r	S2r	N1r	S3r	S2r	S3r	S3r	S2r	S3r	S2r	N1rt	S3r	S3r	S3t	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Halagera	219	N1r	S2t	S3r	S2r	S3r	S2r	N1r	S3r	S2r	S3r	S3r	S2r	S3r	S2r	N1rt	S3r	S3r	S3t	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	
Halagera	221	N1r	S2t	S3r	S2r	S3r	S2r	N1r	S3r	S2r	S3r	S3r	S2r	S3r	S2r	N1rt	S3r	S3r	S3t	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S3r	S3r	

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry	
Halagera	222	N1r	S2t	S3r	S2r	S3r	S2r	N1r	S3r	S2r	S3r	S3r	S2r	S3r	S2r	N1rt	S3r	S3r	S3t	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S2r	S3r	S3r
Halagera	223	S3tw	S2tw	S3tw	S1	S3tw	S2r	S2tw	S2w	S1	S1	S2tw	S2tw	S3tw	S1	N1tz	S2tw	S2w	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2tw	S2t	S1	S1	S2tw	S3tw	
Halagera	224	N1r	S2t	S3r	S2r	S3r	S2r	N1r	S3r	S2r	S3r	S3r	S2r	S3r	S2r	N1rt	S3r	S3r	S3t	S2r	S2r	S2r	S2r	S2r	S3r	S2r	S2r	S2r	S2r	S3r	S3r
Halagera	383	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Halagera	384	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Halagera	386	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Halagera	410	S3tw	S2tw	S3tw	S1	S3tw	S1	S2tw	S2w	S1	S1	S3tw	S1	S3tw	S1	N1tw	S2tw	S2w	S3tw	S2t	S2tw	S3tw	S2tw	S2tw	S2tw	S2t	S2t	S2t	S2tw	S3tw	
Jinatera	253	N1rl	S2rl	S3rl	S2rl	S3rl	S3rl	N1rl	S3rl	S2rl	S3rl	S3rl	S2rl	S3rl	S2rl	S3rl	S3rl	S3rl	S2rl	S2r	S2rl	S2rl	S2rl	S2rl	S3rl	S2rl	S2r	S2r	S3rl	S3rl	
Jinatera	256	N1rl	S2rl	S3rl	S2rl	S3rl	S3rl	N1rl	S3rl	S2rl	S3rl	S3rl	S2rl	S3rl	S2rl	S3rl	S3rl	S3rl	S2rl	S2r	S2rl	S2rl	S2rl	S2rl	S3rl	S2rl	S2r	S2r	S3rl	S3rl	
Jinatera	258	N1rl	S2rl	S3rl	S2rl	S3rl	S3rl	N1rl	S3rl	S2rl	S3rl	S3rl	S2rl	S3rl	S2rl	S3rl	S3rl	S3rl	S2rl	S2r	S2rl	S2rl	S2rl	S2rl	S3rl	S2rl	S2r	S2r	S3rl	S3rl	
Mylapura	51	S3tw	S2tw	S3tw	S1	S3tw	S1	S2tw	S2w	S1	S2w	S3tw	S1	S3tw	S1	N1tw	S2tw	S2w	S3tw	S3t	S2tw	S3tw	S2tw	S2tw	S2tw	S2t	S1	S1	S2tw	S3tw	
Mylapura	52/1	S3rz	S2z	S2rz	S2z	S2rt	S2rz	S3rz	S2rz	S2tz	S2rz	S2rz	S1	S2rz	S1	S2rt	S3rz	S2rz	S2tz	S1	S2tz	S2z	S2tz	S2tz	S2rz	S2z	S1	S1	S2rz	S2rz	
Mylapura	52/2	S3rz	S2z	S2rz	S2z	S2rt	S2rz	S3rz	S2rz	S2tz	S2rz	S2rz	S1	S2rz	S1	S2rt	S3rz	S2rz	S2tz	S1	S2tz	S2z	S2tz	S2tz	S2rz	S2z	S1	S1	S2rz	S2rz	
Mylapura	53	S3tw	S2tw	S3tw	S1	S3tw	S1	S2tw	S2w	S1	S2w	S3tw	S1	S3tw	S1	N1tw	S2tw	S2w	S3tw	S3t	S2tw	S3tw	S2tw	S2tw	S2tw	S2t	S1	S1	S2tw	S3tw	
Mylapura	54	S3rz	S2z	S2rz	S2z	S2rt	S2rz	S3rz	S2rz	S2tz	S2rz	S2rz	S1	S2rz	S1	S2rt	S3rz	S2rz	S2tz	S1	S2tz	S2z	S2tz	S2tz	S2rz	S2z	S1	S1	S2rz	S2rz	
Mylapura	55	S3rz	S2z	S2rz	S2z	S2rt	S2rz	S3rz	S2rz	S2tz	S2rz	S2rz	S1	S2rz	S1	S2rt	S3rz	S2rz	S2tz	S1	S2tz	S2z	S2tz	S2tz	S2rz	S2z	S1	S1	S2rz	S2rz	
Mylapura	56	S3rz	S2z	S2rz	S2z	S2rt	S2rz	S3rz	S2rz	S2tz	S2rz	S2rz	S1	S2rz	S1	S2rt	S3rz	S2rz	S2tz	S1	S2tz	S2z	S2tz	S2tz	S2rz	S2z	S1	S1	S2rz	S2rz	
Mylapura	57	S3tw	S2tw	S3tw	S1	S3tw	S1	S2tw	S2w	S1	S2w	S3tw	S1	S3tw	S1	N1tw	S2tw	S2w	S3tw	S3t	S2tw	S3tw	S2tw	S2tw	S2tw	S2t	S1	S1	S2tw	S3tw	
Mylapura	58	S3tw	S2tw	S3tw	S1	S3tw	S1	S2tw	S2w	S1	S2w	S3tw	S1	S3tw	S1	N1tw	S2tw	S2w	S3tw	S3t	S2tw	S3tw	S2tw	S2tw	S2tw	S2t	S1	S1	S2tw	S3tw	
Mylapura	59	S3tw	S2tw	S3tw	S1	S3tw	S1	S2tw	S2w	S1	S2w	S3tw	S1	S3tw	S1	N1tw	S2tw	S2w	S3tw	S3t	S2tw	S3tw	S2tw	S2tw	S2tw	S2t	S1	S1	S2tw	S3tw	
Mylapura	60	S3tw	S2tw	S3tw	S1	S3tw	S1	S2tw	S2w	S1	S2w	S3tw	S1	S3tw	S1	N1tw	S2tw	S2w	S3tw	S3t	S2tw	S3tw	S2tw	S2tw	S2tw	S2t	S1	S1	S2tw	S3tw	
Mylapura	61	S3tw	S2tw	S3tw	S1	S3tw	S1	S2tw	S2w	S1	S2w	S3tw	S1	S3tw	S1	N1tw	S2tw	S2w	S3tw	S3t	S2tw	S3tw	S2tw	S2tw	S2tw	S2t	S1	S1	S2tw	S3tw	
Mylapura	62	S3tw	S2tw	S3tw	S1	S3tw	S1	S2tw	S2w	S1	S2w	S3tw	S1	S3tw	S1	N1tw	S2tw	S2w	S3tw	S3t	S2tw	S3tw	S2tw	S2tw	S2tw	S2t	S1	S1	S2tw	S3tw	
Mylapura	63	S3tw	S2tw	S3tw	S1	S3tw	S1	S2tw	S2w	S1	S2w	S3tw	S1	S3tw	S1	N1tw	S2tw	S2w	S3tw	S3t	S2tw	S3tw	S2tw	S2tw	S2tw	S2t	S1	S1	S2tw	S3tw	
Mylapura	64	S3tw	S2tw	S3tw	S1	S3tw	S1	S2tw	S2w	S1	S2w	S3tw	S1	S3tw	S1	N1tw	S2tw	S2w	S3tw	S3t	S2tw	S3tw	S2tw	S2tw	S2tw	S2t	S1	S1	S2tw	S3tw	
Mylapura	123	S2r	S2tw	S3t	S1	S3t	S1	S2t	S2z	S1	S1	S2tw	S2t	S3t	S2z	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2t	S1	S1	S2tw	S3tw	
Mylapura	124	S2r	S2tw	S3t	S1	S3t	S1	S2t	S2z	S1	S1	S2tw	S2t	S3t	S2z	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2t	S1	S1	S2tw	S3tw	

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry	
Mylapura	125	S2r	S2tw	S3t	S1	S3t	S1	S2t	S2z	S1	S1	S2tw	S2t	S3t	S2z	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2t	S1	S1	S2tw	S3tw	
Mylapura	126	S2r	S2tw	S3t	S1	S3t	S1	S2t	S2z	S1	S1	S2tw	S2t	S3t	S2z	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2t	S1	S1	S2tw	S3tw	
Mylapura	127	S2r	S2tw	S3t	S1	S3t	S1	S2t	S2z	S1	S1	S2tw	S2t	S3t	S2z	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2t	S1	S1	S2tw	S3tw	
Mylapura	128	S2r	S2tw	S3t	S1	S3t	S1	S2t	S2z	S1	S1	S2tw	S2t	S3t	S2z	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2t	S1	S1	S2tw	S3tw	
Mylapura	131	S2r	S2tw	S3t	S1	S3t	S1	S2t	S2z	S1	S1	S2tw	S2t	S3t	S2z	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2t	S1	S1	S2tw	S3tw	
Naglapura	195	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	
Naglapura	196	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz	
Naglapura	205	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz	
Naglapura	206	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	
Naglapura	207	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	
Panchashee lanagara	46	S3r	S1	S2r	S1	S2rt	S2r	S3r	S2r	S2t	S2r	S2r	S1	S2r	S1	S2rt	S3r	S2r	S2t	S2t	S2t	S1	S2t	S2t	S2r	S1	S1	S2t	S2r	S2r	
Panchashee lanagara	47	N1t	N1t	N1t	N1t	N1t	N1t	N1t	N1t	N1t	N1t	N1t	N1t	N1t	S3	N1t	N1t	N1t	S2t	S3t	N1t	N1t	S3t	S3t	N1t	S3t	N1t	N1t	N1t	N1t	
Panchashee lanagara	48	N1t	N1t	N1t	N1t	N1t	N1t	N1t	N1t	N1t	N1t	N1t	N1t	N1t	S3	N1t	N1t	N1t	S2t	S3t	N1t	N1t	S3t	S3t	N1t	S3t	N1t	N1t	N1t	N1t	
Panchashee lanagara	49	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	Ro	
Risabadha Hosalli	1	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	
Risabadha Hosalli	2	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Risabadha Hosalli	3	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Risabadha Hosalli	4	S2r	S2tw	S3t	S1	S3t	S1	S2t	S2z	S1	S1	S2tw	S2t	S3t	S2z	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2t	S1	S1	S2tw	S3tw	
Risabadha Hosalli	5	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Risabadha Hosalli	6	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Risabadha Hosalli	7	S2r	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S1	S2tw	S2tw	S2tw	S2tw	S2t	S2t	S1	S1	S2tw	S3tw	
Risabadha Hosalli	8	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Risabadha Hosalli	9	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Risabadha Hosalli	10	S2r	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S1	S2tw	S2tw	S2tw	S2tw	S2t	S2t	S1	S1	S2tw	S3tw	



Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry
Risabadha Hosalli	11	S2r	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S1	S2tw	S2tw	S2tw	S2tw	S2t	S2t	S1	S1	S2tw	S3tw
Risabadha Hosalli	12	S2r	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S1	S2tw	S2tw	S2tw	S2tw	S2t	S2t	S1	S1	S2tw	S3tw
Risabadha Hosalli	13	S2r	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S1	S2tw	S2tw	S2tw	S2tw	S2t	S2t	S1	S1	S2tw	S3tw
Risabadha Hosalli	14	S2r	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S1	S2tw	S2tw	S2tw	S2tw	S2t	S2t	S1	S1	S2tw	S3tw
Risabadha Hosalli	15	S3rg	S3g	S3g	S3g	S3g	S3g	S3rg	S3g	S3g	S3g	S3g	S2g	S3g	S2g	S3g	S3rg	S3g	S2g	S2g	S3g	S3g	S3g	S3g	S3g	S2g	S2g	S2g	S2g	S2g
Risabadha Hosalli	16	S2r	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S1	S2tw	S2tw	S2tw	S2tw	S2t	S2t	S1	S1	S2tw	S3tw
Risabadha Hosalli	17	S2r	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S1	S2tw	S2tw	S2tw	S2tw	S2t	S2t	S1	S1	S2tw	S3tw
Risabadha Hosalli	18	S2r	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S1	S2tw	S2tw	S2tw	S2tw	S2t	S2t	S1	S1	S2tw	S3tw
Risabadha Hosalli	19	S2r	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S1	S2tw	S2tw	S2tw	S2tw	S2t	S2t	S1	S1	S2tw	S3tw
Risabadha Hosalli	20	S2r	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S1	S2tw	S2tw	S2tw	S2tw	S2t	S2t	S1	S1	S2tw	S3tw
Risabadha Hosalli	21	S2r	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S1	S2tw	S2tw	S2tw	S2tw	S2t	S2t	S1	S1	S2tw	S3tw
Risabadha Hosalli	22	S2r	S2tw	S3t	S1	S3t	S1	S2t	S2z	S1	S1	S2tw	S2t	S3t	S2z	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2t	S1	S1	S2tw	S3tw
Risabadha Hosalli	23	S2r	S2tw	S3t	S1	S3t	S1	S2t	S2z	S1	S1	S2tw	S2t	S3t	S2z	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2t	S1	S1	S2tw	S3tw
Risabadha Hosalli	24	S2r	S2tw	S3t	S1	S3t	S1	S2t	S2z	S1	S1	S2tw	S2t	S3t	S2z	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2t	S1	S1	S2tw	S3tw
Risabadha Hosalli	25	S2r	S2tw	S3t	S1	S3t	S1	S2t	S2z	S1	S1	S2tw	S2t	S3t	S2z	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2t	S1	S1	S2tw	S3tw
Risabadha Hosalli	26	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt
Risabadha Hosalli	27	S2r	S2tw	S3t	S1	S3t	S1	S2t	S2z	S1	S1	S2tw	S2t	S3t	S2z	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2t	S1	S1	S2tw	S3tw
Risabadha Hosalli	28	S2r	S2tw	S3t	S1	S3t	S1	S2t	S2z	S1	S1	S2tw	S2t	S3t	S2z	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2t	S1	S1	S2tw	S3tw
Risabadha Hosalli	29	S2r	S2tw	S3t	S1	S3t	S1	S2t	S2z	S1	S1	S2tw	S2t	S3t	S2z	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2t	S1	S1	S2tw	S3tw
Risabadha Hosalli	30	S2r	S2tw	S3t	S1	S3t	S1	S2t	S2z	S1	S1	S2tw	S2t	S3t	S2z	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2t	S1	S1	S2tw	S3tw
Risabadha Hosalli	31	S2r	S2tw	S3t	S1	S3t	S1	S2t	S2z	S1	S1	S2tw	S2t	S3t	S2z	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2t	S1	S1	S2tw	S3tw
Risabadha Hosalli	32	S2r	S2tw	S3t	S1	S3t	S1	S2t	S2z	S1	S1	S2tw	S2t	S3t	S2z	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2t	S1	S1	S2tw	S3tw

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry
Risabadha Hosalli	33	S2r	S2tw	S3t	S1	S3t	S1	S2t	S2z	S1	S1	S2tw	S2t	S3t	S2z	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2t	S1	S1	S2tw	S3tw
Risabadha Hosalli	34	S2r	S2tw	S3t	S1	S3t	S1	S2t	S2z	S1	S1	S2tw	S2t	S3t	S2z	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2t	S1	S1	S2tw	S3tw
Risabadha Hosalli	35	S2r	S2tw	S3t	S1	S3t	S1	S2t	S2z	S1	S1	S2tw	S2t	S3t	S2z	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2t	S1	S1	S2tw	S3tw
Risabadha Hosalli	45	N1r	S2tg	S3r	S2rg	S3r	S2rg	N1r	S3r	S2rg	S3r	S3r	S2r	S3r	S2r	N1rt	S3r	S3r	S3t	S2r	S2rg	S2rg	S2rg	S2rg	S3r	S2r	S2r	S2r	S3r	S3r
Risabadha Hosalli	46/1	N1r	S2tg	S3r	S2rg	S3r	S2rg	N1r	S3r	S2rg	S3r	S3r	S2r	S3r	S2r	N1rt	S3r	S3r	S3t	S2r	S2rg	S2rg	S2rg	S2rg	S3r	S2r	S2r	S2r	S3r	S3r
Risabadha Hosalli	46/10	N1r	S2tg	S3r	S2rg	S3r	S2rg	N1r	S3r	S2rg	S3r	S3r	S2r	S3r	S2r	N1rt	S3r	S3r	S3t	S2r	S2rg	S2rg	S2rg	S2rg	S3r	S2r	S2r	S2r	S3r	S3r
Risabadha Hosalli	46/4	S2r	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S1	S2tw	S2tw	S2tw	S2tw	S2t	S2t	S1	S1	S2tw	S3tw
Risabadha Hosalli	46/5	N1r	S2tg	S3r	S2rg	S3r	S2rg	N1r	S3r	S2rg	S3r	S3r	S2r	S3r	S2r	N1rt	S3r	S3r	S3t	S2r	S2rg	S2rg	S2rg	S2rg	S3r	S2r	S2r	S2r	S3r	S3r
Risabadha Hosalli	46/6	N1r	S2tg	S3r	S2rg	S3r	S2rg	N1r	S3r	S2rg	S3r	S3r	S2r	S3r	S2r	N1rt	S3r	S3r	S3t	S2r	S2rg	S2rg	S2rg	S2rg	S3r	S2r	S2r	S2r	S3r	S3r
Risabadha Hosalli	46/7	N1r	S2tg	S3r	S2rg	S3r	S2rg	N1r	S3r	S2rg	S3r	S3r	S2r	S3r	S2r	N1rt	S3r	S3r	S3t	S2r	S2rg	S2rg	S2rg	S2rg	S3r	S2r	S2r	S2r	S3r	S3r
Risabadha Hosalli	46/8	N1r	S2tg	S3r	S2rg	S3r	S2rg	N1r	S3r	S2rg	S3r	S3r	S2r	S3r	S2r	N1rt	S3r	S3r	S3t	S2r	S2rg	S2rg	S2rg	S2rg	S3r	S2r	S2r	S2r	S3r	S3r
Risabadha Hosalli	46/9	N1r	S2tg	S3r	S2rg	S3r	S2rg	N1r	S3r	S2rg	S3r	S3r	S2r	S3r	S2r	N1rt	S3r	S3r	S3t	S2r	S2rg	S2rg	S2rg	S2rg	S3r	S2r	S2r	S2r	S3r	S3r
Risabadha Hosalli	47	S2r	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S1	S2tw	S2tw	S2tw	S2tw	S2t	S2t	S1	S1	S2tw	S3tw
Risabadha Hosalli	48	S2r	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S1	S2tw	S2tw	S2tw	S2tw	S2t	S2t	S1	S1	S2tw	S3tw
Risabadha Hosalli	49	N1r	S2tg	S3r	S2rg	S3r	S2rg	N1r	S3r	S2rg	S3r	S3r	S2r	S3r	S2r	N1rt	S3r	S3r	S3t	S2r	S2rg	S2rg	S2rg	S2rg	S3r	S2r	S2r	S2r	S3r	S3r
Risabadha Hosalli	50	N1r	S2tg	S3r	S2rg	S3r	S2rg	N1r	S3r	S2rg	S3r	S3r	S2r	S3r	S2r	N1rt	S3r	S3r	S3t	S2r	S2rg	S2rg	S2rg	S2rg	S3r	S2r	S2r	S2r	S3r	S3r
Risabadha Hosalli	51	N1r	S2tg	S3r	S2rg	S3r	S2rg	N1r	S3r	S2rg	S3r	S3r	S2r	S3r	S2r	N1rt	S3r	S3r	S3t	S2r	S2rg	S2rg	S2rg	S2rg	S3r	S2r	S2r	S2r	S3r	S3r
Risabadha Hosalli	52	N1r	S2tg	S3r	S2rg	S3r	S2rg	N1r	S3r	S2rg	S3r	S3r	S2r	S3r	S2r	N1rt	S3r	S3r	S3t	S2r	S2rg	S2rg	S2rg	S2rg	S3r	S2r	S2r	S2r	S3r	S3r
Risabadha Hosalli	53	N1r	S2tg	S3r	S2rg	S3r	S2rg	N1r	S3r	S2rg	S3r	S3r	S2r	S3r	S2r	N1rt	S3r	S3r	S3t	S2r	S2rg	S2rg	S2rg	S2rg	S3r	S2r	S2r	S2r	S3r	S3r
Risabadha Hosalli	54	N1r	S2tg	S3r	S2rg	S3r	S2rg	N1r	S3r	S2rg	S3r	S3r	S2r	S3r	S2r	N1rt	S3r	S3r	S3t	S2r	S2rg	S2rg	S2rg	S2rg	S3r	S2r	S2r	S2r	S3r	S3r
Risabadha Hosalli	55	S2r	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S1	S2tw	S2tw	S2tw	S2tw	S2t	S2t	S1	S1	S2tw	S3tw
Risabadha Hosalli	56	S2r	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S1	S2tw	S2tw	S2tw	S2tw	S2t	S2t	S1	S1	S2tw	S3tw

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry	
Risabadha Hosalli	57	S2r	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S1	S2tw	S2tw	S2tw	S2tw	S2t	S2t	S1	S1	S2tw	S3tw	
Risabadha Hosalli	58	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt	
Risabadha Hosalli	59	S2r	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S1	S2tw	S2tw	S2tw	S2tw	S2t	S2t	S1	S1	S2tw	S3tw	
Risabadha Hosalli	60	S2r	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S1	S2tw	S2tw	S2tw	S2tw	S2t	S2t	S1	S1	S2tw	S3tw	
Risabadha Hosalli	61	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Risabadha Hosalli	62	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Risabadha Hosalli	63	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Risabadha Hosalli	64	S2r	S2t	S3t	S1	S3t	S1	S2t	S1	S1	S1	S2tw	S2t	S3t	S1	N1t	S2t	S1	S3tw	S1	S2tw	S2tw	S2tw	S2tw	S2t	S2t	S1	S1	S2tw	S3tw	
Risabadha Hosalli	65	N1r	S2tg	S3r	S2rg	S3r	S2rg	N1r	S3r	S2rg	S3r	S3r	S2r	S3r	S2r	N1rt	S3r	S3r	S3t	S2r	S2rg	S2rg	S2rg	S2rg	S3r	S2r	S2r	S2r	S3r	S3r	
Risabadha Hosalli	66	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Risabadha Hosalli	67	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Risabadha Hosalli	68	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Risabadha Hosalli	69	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Risabadha Hosalli	71	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Risabadha Hosalli	72	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Risabadha Hosalli	73	N1r	S3rt	N1r	S3rt	N1r	N1t	N1r	N1r	N1t	N1r	N1r	S3rt	N1r	S3rt	N1rt	N1r	N1r	S3rt	S3r	S3rt	S3rt	S3r	S3r	N1r	S3rt	S3rt	S3r	N1r	N1r	
Risabadha Hosalli	74	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz	
Risabadha Hosalli	75	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz	
Risabadha Hosalli	76	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz	
Risabadha Hosalli	77	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz	
Risabadha Hosalli	78	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz	
Risabadha Hosalli	79	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz	

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry
Risabadha Hosalli	80	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Risabadha Hosalli	82	S2r	S2tw	S3t	S1	S3t	S1	S2t	S2z	S1	S1	S2tw	S2t	S3t	S2z	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2t	S1	S1	S2tw	S3tw
Risabadha Hosalli	83	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Risabadha Hosalli	84	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Risabadha Hosalli	85	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Risabadha Hosalli	86	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Risabadha Hosalli	87	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Risabadha Hosalli	88	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Risabadha Hosalli	89	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Risabadha Hosalli	90	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt
Risabadha Hosalli	91	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt
Risabadha Hosalli	92	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt
Risabadha Hosalli	93	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt
Risabadha Hosalli	94	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt
Risabadha Hosalli	95	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt
Risabadha Hosalli	96	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt
Risabadha Hosalli	97	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt
Risabadha Hosalli	98	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt
Risabadha Hosalli	99	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt
Risabadha Hosalli	100	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt
Risabadha Hosalli	101	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt
Risabadha Hosalli	102	N1r	S3rt	N1r	S3r	N1rt	S3r	N1rt	N1r	S3r	N1r	S3rt	S3rt	N1rt	S3r	N1rt	N1rt	N1r	S3rt	S3r	S3rt	S3rt	S3rt	S3rt	N1r	S3rt	S3r	S3r	N1rt	N1rt

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry
Risabadha Hosalli	103	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Risabadha Hosalli	104	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Risabadha Hosalli	105	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Risabadha Hosalli	106	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Risabadha Hosalli	107	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Risabadha Hosalli	108	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Risabadha Hosalli	109	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Risabadha Hosalli	110	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Risabadha Hosalli	111	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Risabadha Hosalli	112	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Risabadha Hosalli	113	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Risabadha Hosalli	114	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Risabadha Hosalli	115	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Risabadha Hosalli	116	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Risabadha Hosalli	117	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Risabadha Hosalli	118	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Risabadha Hosalli	119	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Risabadha Hosalli	120	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Risabadha Hosalli	121	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Risabadha Hosalli	122	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Risabadha Hosalli	123	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Risabadha Hosalli	124	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry
Risabadha Hosalli	125	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Risabadha Hosalli	126	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Risabadha Hosalli	127	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Risabadha Hosalli	128	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Risabadha Hosalli	129	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Risabadha Hosalli	130	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Risabadha Hosalli	131	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Risabadha Hosalli	132	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Risabadha Hosalli	133	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Risabadha Hosalli	134	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Risabadha Hosalli	135	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Risabadha Hosalli	136	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Risabadha Hosalli	137	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Risabadha Hosalli	138	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Risabadha Hosalli	139	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Risabadha Hosalli	140	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Risabadha Hosalli	141	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Risabadha Hosalli	142	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Risabadha Hosalli	143	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Risabadha Hosalli	144	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Risabadha Hosalli	145	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Risabadha Hosalli	146	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry
Risabadha Hosalli	147	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Risabadha Hosalli	148	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Risabadha Hosalli	149	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Risabadha Hosalli	150	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Risabadha Hosalli	151	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Risabadha Hosalli	152	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Risabadha Hosalli	153	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Risabadha Hosalli	154	S2r	S2tw	S3t	S1	S3t	S1	S2t	S2z	S1	S1	S2tw	S2t	S3t	S2z	N1t	S2t	S2z	S3tw	S1	S2tw	S3tw	S2tw	S2tw	S2t	S2t	S1	S1	S2tw	S3tw
Risabadha Hosalli	155	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Risabadha Hosalli	156	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Risabadha Hosalli	157/1	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Risabadha Hosalli	157/2	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Risabadha Hosalli	158	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Risabadha Hosalli	159	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Risabadha Hosalli	160	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Risabadha Hosalli	161	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Risabadha Hosalli	162	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Risabadha Hosalli	163	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Risabadha Hosalli	164	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Risabadha Hosalli	165	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Risabadha Hosalli	166	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz
Risabadha Hosalli	167	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz

Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry	
Risabadha Hosalli	168	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz	
Risabadha Hosalli	169	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz	
Risabadha Hosalli	170	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz	
Risabadha Hosalli	171	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz	
Risabadha Hosalli	172	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz	
Risabadha Hosalli	173	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz	
Risabadha Hosalli	174	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz	
Risabadha Hosalli	175	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz	
Risabadha Hosalli	176	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz	
Risabadha Hosalli	177	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz	
Risabadha Hosalli	178	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz	
Risabadha Hosalli	179	S3tz	S3tz	S2tz	S3tz	S2tz	N1tz	S2rt	S1	N1tz	S3tz	S2tz	S3tz	S3tz	S3tz	S3tz	S2rz	S1	S2tz	S1	S2tz	S2z	S2z	S2z	S2tz	S2z	S2t	S2t	S2tz	S3tz	
Risabadha Hosalli	180	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Risabadha Hosalli	181/A	S3rg	S3g	S3g	S3g	S3g	S3g	S3rg	S3g	S3g	S3g	S3g	S2g	S3g	S2g	S3g	S3rg	S3g	S2g	S2g	S3g	S3g	S3g	S3g	S3g	S2g	S2g	S2g	S2g	S2g	
Risabadha Hosalli	181/B	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Risabadha Hosalli	182	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Risabadha Hosalli	183	S3rg	S3g	S3g	S3g	S3g	S3g	S3rg	S3g	S3g	S3g	S3g	S2g	S3g	S2g	S3g	S3rg	S3g	S2g	S2g	S3g	S3g	S3g	S3g	S3g	S2g	S2g	S2g	S2g	S2g	
Risabadha Hosalli	184	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Risabadha Hosalli	185	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others	Others
Yaleri	691	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	
Yaleri	693	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Yaleri	694	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r



Village	Survey Number	Mango	Maize	Sapota	Sorghum	Guava	Cotton	Tamarind	Lime	Bengal gram	Sunflower	Red gram	Amla	Jackfruit	Custard-apple	Cashew	Jamun	Musambi	Groundnut	Onion	Chilly	Tomato	Marigold	Chrysanthemum	Pomegranate	Bajra	Brinjal	Bhendi	Drumstick	Mulberry
Yaleri	695	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Yaleri	696	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Yaleri	697	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Yaleri	698	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Yaleri	699	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r
Yaleri	702	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r	N1r



# **PART-B**

**SOCIO-ECONOMIC STATUS OF FARM HOUSEHOLDS**



## CONTENTS

1.	Salient findings of the survey	1-4
2.	Introduction	5
3	Methodology	7
4	Salient features of the survey	9-26
5	Summary	27-30



## LIST OF TABLES

1	Households sampled for socio economic survey	9
2	Population characteristics	9
3	Age wise classification of household members	9
4	Education level of household members	10
5	Occupation of household heads	10
6	Occupation of family members	10
7	Institutional participation of household members	11
8	Type of house owned by households	11
9	Durable assets owned by households	11
10	Average value of durable assets owned by households	12
11	Farm implements owned by households	12
12	Average value of farm implements	12
13	Livestock possession by households	13
14	Average labour availability	13
15	Adequacy of hired labour	14
16	Distribution of land (ha)	14
17	Average land value	15
18	Status of bore wells	15
19	Source of irrigation	15
20	Depth of water	15
21	Irrigated area	15
22	Cropping pattern	16
23	Cropping intensity	16
24	Cost of cultivation of Cotton	17
25	Cost of cultivation of Green gram	18
26	Cost of cultivation of Red gram	19
27	Cost of cultivation of Groundnut	20
28	Adequacy of fodder	21
29	Annual gross income	21
30	Average annual expenditure	21
31	Horticulture species grown	22

32	Forest species grown	22
33	Average additional investment capacity	22
34	Source of additional investment	22
35	Marketing of the agricultural produce	23
36	Marketing channels used for sale of agricultural produce	23
37	Mode of transport of agricultural produce	23
38	Incidence of soil and water erosion problems	23
39	Interest towards soil testing	24
40	Usage pattern of fuel for domestic use	24
41	Source of drinking water	24
42	Source of light	24
43	Existence of sanitary toilet facility	24
44	Possession of public distribution system(PDS) card	25
45	Participation in NREGA programme	25
46	Adequacy of food items	25
47	Response on inadequacy of food items	25
48	Farming constraints experienced	26



**SALIENT FINDINGS OF THE SURVEY**

- ❖ *The data indicated that there were 125 (63.45%) men and 72 (36.55%) women among the sampled households.*
- ❖ *The average family size of landless farmers' was 3.5, marginal farmers' was 4.5, small farmers' was 6, semi medium farmers' was 5.1, medium farmers were 4.3 and large farmers was 3.*
- ❖ *The data indicated that, 37 (18.78%) people were in 0-15 years of age, 75 (38.07%) were in 16-35 years of age, 69 (35.03%) were in 36-60 years of age and 16 (8.12%) were above 61 years of age.*
- ❖ *The results indicated that Raisabad Hosalli-1 had 59.90 per cent illiterates, 1.02 per cent of them functional literate, 15.74 per cent of them had primary school, 3.55 per cent of them had Middle school and PUC education, 11.17 per cent of them had high school and 3.05 per cent of them had degree education.*
- ❖ *The results indicate that, 85 per cent of household heads were practicing agriculture, 10 per cent of the household heads were agricultural labourers and 5 cent of the household heads was housewives.*
- ❖ *The results indicate that agriculture was the major occupation for 58.88 per cent of the household members, 6.09 per cent were agricultural labourers, 1.52 per cent were private service, 22.34 per cent student, 9.14 per cent were housewives and 2.03 per cent were children.*
- ❖ *The results show that, 100 per cent of the population in the micro watershed has not participated in any local institutions.*
- ❖ *The results indicate that 7.5 per cent of the households possess thatched, 67.5 per cent of the households possess katcha house and 25 per cent of the households possess pucca/RCC house.*
- ❖ *The results show that 45 per cent of the households possess TV, 2.50 per cent of the households possess radio and landline phone, 7.50 per cent of the households possess mixer/grinder, 5 per cent of the households possess bicycle, 20 per cent of the households possess motor cycle and 95 per cent of the households possess mobile phones.*
- ❖ *The results show that the average value of television was Rs. 6,393, mixer/grinder was Rs. 1,642, motor cycle was Rs. 56,333, auto was Rs. 160,000 and mobile phone was Rs. 2,474.*
- ❖ *About 10 per cent each of the households possess bullock cart, 30 per cent each of the households possess plough, 7.50 per cent of the households possess sprayer and sprinkler and 65 per cent of the households possess weeder.*
- ❖ *The results show that the average value of bullock cart was Rs. 16,500, plough was Rs. 1,575, sprayer was Rs. 2,166, sprinkler was Rs. 2,333 and the average value of weeder was Rs. 57.*

- ❖ *The results indicate that, 35 per cent of the households possess bullocks, 22.5 per cent of the households possess local cow, 2.5 per cent of the households possess crossbreed cow, 17.5 per cent of the households possess buffalo and 7.50 per cent of the households possess goat.*
- ❖ *The results indicate that, average own labour men available in the micro watershed was 2.18, average own labour (women) available was 1.45, average hired labour (men) available was 10 and average hired labour (women) available was 9.05.*
- ❖ *The results indicate that, 95 per cent of the households opined that the hired labour was adequate.*
- ❖ *The results indicate that, households of the Raisabad Hosalli-1 micro-watershed possess 23.63 ha (46.73%) of dry land, 25.80 ha (51.03%) of irrigated land and 1.13 ha (2.24%) of permanent fallow land. Marginal farmers possess 5.4 ha (73.64%) of dry land, 1.7 ha (23.06%) of irrigated land and 0.2 ha (3.29%) of permanent fallow land. Small farmers possess 5.16 ha (49.3%) of dry land and 5.29 ha (50.6%) of irrigated land. Semi medium farmers possess 6.16 ha (38.1%) of dry land and 9.11 ha (56.3%) of irrigated land and 0.89 ha (5.51 %) of permanent fallow land. Medium farmers possess 6.88 ha (62.96%) of dry land and 4.05 ha (37.04%) of irrigated land. Large farmers possess 5.67 ha (100%) of irrigated land.*
- ❖ *The results indicate that, the average value of dry land was Rs. 340,587.53, the average value of irrigated land was Rs. 604,328.73 and the average value of permanent fallow land was Rs. 573,392.85. In case of marginal famers, the average land value was Rs. 690,715.90 for dry land, the average land value was Rs. 1,587,857.12 and the average land value was Rs.1,646,666.60. In case of small famers, the average land value was Rs. 407,142.86 for dry land and the average land value was Rs. 832,159.27 for irrigated land. In case of semi medium famers, the average land value was Rs. 227,051.87 for dry land, the average land value was Rs. 636,711.11 for irrigated land and the average land value was Rs. 280,681.82 for permanent fallow land. In case of medium famers, the average land value was Rs. 116,235.29 for dry land and the average land value was Rs. 469,300 for irrigated land. In case of large famers, the average land value was Rs. 141,142.86 for irrigated land.*
- ❖ *The results indicate that, there were 20 functioning bore wells in the micro watershed.*
- ❖ *The results indicate that, bore well was the major irrigation source in the micro water shed for 50 per cent of the farmers.*
- ❖ *The results indicate that, the depth of bore well was found to be 52.96 meters.*
- ❖ *The results indicate that, marginal, small, semi medium, medium and large farmers had an irrigated area of 1.70 ha, 5.29 ha, 9.19 ha, 4.05 ha and 5.67 ha respectively.*
- ❖ *The results indicate that, farmers have grown cotton (20.11%), green gram (15.35 ha), groundnut (3.62 ha) and red gram (4.20 ha). Marginal farmers have grown*

cotton, green gram and red gram. Small farmers have grown cotton, groundnut and red gram. Semi medium farmers have grown cotton, green gram and groundnut. Medium farmers have grown cotton, green gram and red gram. Large farmers have grown cotton.

- ❖ The results indicate that, the cropping intensity in Raisabad Hosalli-1 micro-watershed was found to be 96.78 per cent.
- ❖ The results indicate that, the total cost of cultivation for Cotton was Rs. 37968.84. The gross income realized by the farmers was Rs. 124061.50. The net income from Cotton cultivation was Rs. 86092.66. Thus the benefit cost ratio was found to be 1:3.27.
- ❖ The total cost of cultivation for green gram was Rs. 27795.50. The gross income realized by the farmers was Rs. 54874.85. The net income from green gram cultivation was Rs. 27079.35. Thus the benefit cost ratio was found to be 1:1.97.
- ❖ The total cost of cultivation for Red gram was Rs. 65271.49. The gross income realized by the farmers was Rs. 149815.42. The net income from Red gram cultivation was Rs. 84543.93. Thus the benefit cost ratio was found to be 1:2.3.
- ❖ The total cost of cultivation for groundnut was Rs. 31661.34. The gross income realized by the farmers was Rs. 72163.14. The net income from groundnut cultivation was Rs. 40501.81. Thus the benefit cost ratio was found to be 1:2.28.
- ❖ The results indicate that, 47.50 per cent of the households opined that dry fodder and green fodder was adequate.
- ❖ The results indicate that the annual gross income was Rs. 50,000 for landless farmers, for marginal farmers it was Rs. 124,764.29, for small farmers it was Rs. 139,500, semi medium farmers it was Rs. 192,400, for medium farmers it was Rs. 376,500 and large farmers it was Rs. 636,000.
- ❖ The results indicate that the average annual expenditure is Rs. 37,494.69. For landless farmers it was Rs. 37,500, marginal farmers it was Rs. 16,490.78, for small farmers it was Rs. 11,750, for semi medium farmers it was Rs. 25,241.67, for medium farmers it was Rs. 120,666.67 and for large farmers it was Rs. 462,000.
- ❖ The results indicate that, households have planted 31 mango and 1 sapota trees in their field.
- ❖ The results indicate that, households have planted 83 neem, 6 tamarind and 1 banyan trees in their field and also 22 neem trees in their backyard.
- ❖ The results indicated that, households have an average investment capacity of Rs. 200 for land development.
- ❖ The results indicated that loan from bank was the source of additional investment for 2.44 per cent for land development. Own funds was the source of additional investment for 7.32 per cent for land development.

- ❖ *The results indicated that, cotton was sold to the extent of 99.54 per cent, green gram and red gram was sold to the extent of 93.75 per cent and groundnut was sold to the extent of 98.25 per cent.*
- ❖ *The results indicated that, about 90 per cent of the farmers sold their produce to local/village merchant and 7.50 per cent of the farmers sold their produce to regulated market.*
- ❖ *The results indicated that, 97.5 per cent of the households have used tractor as a mode of transportation.*
- ❖ *The results indicated that, 50 per cent of the households have experienced soil and water erosion problems in the farm.*
- ❖ *The results indicated that, 80 per cent have shown interest in soil test.*
- ❖ *The results indicated that, 90 per cent of the households used fire wood and 10 per cent of the households used LPG as a source of fuel.*
- ❖ *The results indicated that, piped supply was the major source of drinking water for 92.5 per cent and 7.50 per cent of the households used bore well in the micro watershed.*
- ❖ *Electricity was the major source of light for 100 per cent of the households in micro watershed.*
- ❖ *The results indicated that, 57.5 per cent of the households possess sanitary toilet facility.*
- ❖ *The results indicated that, 100 per cent of the sampled households possessed BPL cards.*
- ❖ *The results indicated that, 87.5 per cent of the households participated in NREGA programme.*
- ❖ *The results indicated that, cereals milk and egg were adequate for 100 per cent of the households, pulses were adequate for 80 per cent, oilseed were adequate for 5 per cent, vegetables were adequate for 85 per cent, fruits were adequate for 22.50 per cent and meat were adequate for 90 per cent.*
- ❖ *The results indicated that, pulse were inadequate for 20 per cent of the households, oilseed were inadequate for 95 per cent, vegetables were inadequate for 15 per cent, fruits were inadequate for 77.50 per cent and meat were inadequate for 10 per cent of the households.*
- ❖ *The results indicated that, lower fertility status of the soil in the area was the constraint experienced by 82.5 per cent of the households, wild animal menace on farm field (22.5%), frequent incidence of pest and diseases (87.5%), Inadequacy of irrigation water (12.5%), high cost of fertilizer and plant protection chemicals (85%), high rate of interest on credit (27.5%), low price for the agricultural commodities (65%), lack of marketing facilities in the area (25%), inadequate extension service (2.5%), Lack of transport for safe transport of the Agril produce to the market (12.5%), less rainfall and Source of Agri-technology information (10%).*

## **INTRODUCTION**

Soil and water are the two precious natural resources which are essential for crop production and existence of life on earth. Rainfed agriculture is under severe stress due to various constraints related to agriculture like uneven and erratic distribution of rainfall, indiscriminate use of fertilizers, chemicals and pesticides, adoption of improper land management practices, soil erosion, decline in soil fertility, decline in ground water resources leading to low crop productivity. The area under rainfed agriculture has to be managed effectively using the best available practices to enhance the production of food, fodder and fuel. This is possible if the land resources are characterized at each parcel of land through detailed land resource inventory using the best available techniques of remote sensing, GPS and GIS. The watershed development programs are aimed at the sustainable distribution of its resources and the process of creating and implementing plans, programs, and projects to sustain and enhance watershed functions that affect the plant, animal and human communities within a watershed boundary.

World Bank funded KWDP II, SUJALA III project was implemented in with Broad objective of demonstrating more effective watershed management through greater integration of programmes related to rain-fed agriculture, innovative and science based approaches and strengthen institutional capacities and If successful, it is expected that the systems and tools could be mainstreamed into the overall IWMP in the State of Karnataka and in time, throughout other IWMP operations in India. With this background the socio-economic survey has been carried out with following specific objectives:

1. To understand the demographic features of the households in the micro-watershed
2. To understand the extent of family labour available and additional employment opportunities available within the village.
3. To know the status of assets of households in the micro-watershed for suggesting possible improvements.
4. To study the cropping pattern, cropped area and productivity levels of different households in micro-watershed.
5. To determine the type and extent of livestock owned by different categories of HHs
6. Availability of fodder and level of livestock management.

### **Scope and importance of survey**

Survey helps in identification of different socio-economic and resource use-patterns of farmers at the Micro watershed. Household survey provides demographic features, labour force, and levels of education; land ownership and asset position (including livestock and other household assets) of surveyed households; and cropping patterns, input intensities, and average crop yields from farmers' fields. It also discusses crop utilization and the degree of commercialization of production in the areas; farmers' access to and utilization of credit from formal and informal sources; and the level of adoption and use of soil, water, and pest management technologies.



## **METHODOLOGY**

The description of the methods, components selected for the survey and procedures followed in conducting the baseline survey are furnished under the following heads.

### **Description of the study area**

Yadgir District is one of the 30 districts of Karnataka state in southern India. This district was carved out from the erstwhile Gulbarga district as the 30th district of Karnataka on 10 April 2010. Yadgir town is the administrative headquarters of the district. The district comprises of 3 taluks namely, Shahapur, Yadgiri and Shorapur (There are 16 hoblies, 117 Gram Panchayats, 4 Municipalities, 8 Towns/ Urban agglomeration and 487 inhabited & 32 un-inhabited villages The district occupies an area of 5,160.88 km<sup>2</sup>.

Yadgir district is the second smallest district in the state, area wise is very rich in cultural traditions. The vast stretch of fertile black soil of the district is known for bumper red gram and jowar crops. The district is a "Daal bowl" of the state. The district is also known for cluster of cement industries and a distinct stone popularly known as "Malakheda Stone". Two main rivers, Krishna and Bhima, and a few tributaries flow in this region. Krishna and Bhima Rivers drain the district. They constitute the two major river basins of the district. Kagna and Amarja are the two sub - basins of Bhima River, which occur within the geographical area of the district

According to the 2011 census Yadgir district has a population of 1, 172,985, roughly equal to the nation of Timor-Leste or the US state of Rhode Island. This gives it a ranking of 404th in India (out of a total of 640). The district has a population density of 224 inhabitants per square kilometre (580/sq mi). Its population growth rate over the decade 2001-2011 was 22.67%. Yadgir has a sex ratio of 984 females for every 1000 males, and a literacy rate of 52.36%.

### **Description of the micro watershed**

Raisabad Hosalli-1 micro-watershed in Haligeri sub-watershed (Yadgir taluk and district) is located in between 16<sup>o</sup>44'20.9" to 16<sup>o</sup> 42'24.692" North latitudes 77<sup>o</sup> 16'41.097" to 77<sup>o</sup>15'54.469" East longitudes, covering an area of about 886.45 ha, bounded by Panchasheelanagara, Mylapura, Yaleri and Raisabadha hosalli villages.

### **Methodology followed in assessing socio-economic status of households**

In order to assess the socio-economic condition of the farmers in the watershed a comprehensive questionnaire was prepared. Major components such as demographic conditions, migration details, food consumption and family expenditure pattern, material possession, land holding, land use management, cropping pattern, cost of cultivation of crops, livestock management. The statistical components such as frequency and percentage were used to analyze the data. About 40 households located in the micro-watershed were interviewed for the survey.





## SALIENT FEATURES OF THE SURVEY

**Households sampled for socio-economic survey:** The data on households sampled for socio economic survey in Raisabad Hosalli-1 micro-watershed is presented in Table 1 and it indicated that 40 farmers were sampled in Raisabad Hosalli-1 micro-watershed among them 2 (5%) were landless, 14 (35%) were marginal, 10 (25%) were small and semi medium farmers, medium farmers were 3 (7.5%) and large farmers were 1 (2.5%).

**Table 1: Households sampled for socio economic survey in Raisabad Hosalli-1 micro-watershed**

Sl.No.	Particulars	LL (2)		MF (14)		SF (10)		SMF (10)		MDF (3)		LF (1)		All (40)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Farmers	2	5	14	35	10	25	10	25	3	7.50	1	2.50	40	100

**Population characteristics:** The population characteristics of households sampled for socio-economic survey in Raisabad Hosalli-1 micro-watershed is presented in Table 2. The data indicated that there were 125 (63.45%) men and 72 (36.55%) women among the sampled households. The average family size of landless farmers' was 3.5, marginal farmers' was 4.5, small farmers' was 6, semi medium farmers' was 5.1, medium farmers was 4.3 and large farmers was 3.

**Table 2: Population characteristics of Raisabad Hosalli-1 micro-watershed**

Sl.No.	Particulars	LL (7)		MF (63)		SF (60)		SMF (51)		MDF (13)		LF (3)		All (197)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Men	4	57.14	43	68.25	35	58.33	35	68.63	7	53.85	1	33.33	125	63.45
2	Women	3	42.86	20	31.75	25	41.67	16	31.37	6	46.15	2	66.67	72	36.55
	Total	7	100	63	100	60	100	51	100	13	100	3	100	197	100
	Average		3.5		4.5		6		5.1		4.3		3		4.9

**Age wise classification of population:** The age wise classification of household members in Raisabad Hosalli-1 micro-watershed is presented in Table 3. The data indicated that, 37 (18.78%) people were in 0-15 years of age, 75 (38.07%) were in 16-35 years of age, 69 (35.03%) were in 36-60 years of age and 16 (8.12%) were above 61 years of age.

**Table 3: Age wise classification of household members in Raisabad Hosalli-1 micro-watershed**

Sl. No.	Particulars	LL (7)		MF (63)		SF (60)		SMF (51)		MDF (13)		LF (3)		All (197)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	0-15 years of age	0	0	15	23.81	14	23.33	5	9.80	2	15.38	1	33.33	37	18.78
2	16-35 years of age	5	71.43	20	31.75	22	36.67	24	47.06	4	30.77	0	0	75	38.07
3	36-60 years of age	2	28.57	24	38.10	17	28.33	18	35.29	6	46.15	2	66.67	69	35.03
4	> 61 years	0	0	4	6.35	7	11.67	4	7.84	1	7.69	0	0	16	8.12
	Total	7	100	63	100	60	100	51	100	13	100	3	100	197	100

**Education level of household members:** Education level of household members in Raisabad Hosalli-1 micro-watershed is presented in Table 4. The results indicated that Raisabad Hosalli-1 had 59.90 per cent illiterates, 1.02 per cent of them functional literate, 15.74 per cent of them had primary school, 3.55 per cent of them had Middle school and PUC education, 11.17 per cent of them had high school and 3.05 per cent of them had degree education.

**Table 4. Education level of household members in Raisabad Hosalli-1 micro-watershed**

Sl. No.	Particulars	LL (7)		MF (63)		SF (60)		SMF(51)		MDF(13)		LF (3)		All (197)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Illiterate	6	85.71	34	53.97	36	60	35	68.63	6	46.15	1	33.33	118	59.90
2	Functional Literate	1	14.29	0	0	0	0	1	1.96	0	0	0	0	2	1.02
3	Primary School	0	0	16	25.40	10	16.67	2	3.92	2	15.38	1	33.33	31	15.74
4	Middle School	0	0	3	4.76	2	3.33	1	1.96	1	7.69	0	0	7	3.55
5	High School	0	0	6	9.52	5	8.33	7	13.73	3	23.08	1	33.33	22	11.17
6	PUC	0	0	2	3.17	2	3.33	3	5.88	0	0	0	0	7	3.55
7	Degree	0	0	2	3.17	2	3.33	1	1.96	1	7.69	0	0	6	3.05
Total		7	100	63	100	60	100	51	100	13	100	3	100	197	100

**Occupation of household heads:** The data regarding the occupation of the household heads in Raisabad Hosalli-1 micro-watershed is presented in Table 5. The results indicate that, 85 per cent of household heads were practicing agriculture, 10 per cent of the household heads were agricultural labourers and 5 cent of the household heads was housewives.

**Table 5: Occupation of household heads in Raisabad Hosalli-1 micro-watershed**

Sl.No.	Particulars	LL (2)		MF (14)		SF (10)		SMF (10)		MDF (3)		LF (1)		All (40)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Agriculture	0	0	13	92.86	8	80	9	90	3	100	1	100	34	85
2	Agricultural Labour	2	100	1	7.14	1	10	0	0	0	0	0	0	4	10
3	Housewife	0	0	0	0	1	10	1	10	0	0	0	0	2	5
Total		2	100	14	100	10	100	10	100	3	100	1	100	40	100

**Table 6: Occupation of family members in Raisabad Hosalli-1 micro-watershed**

Sl. No.	Particulars	LL(7)		MF (63)		SF (60)		SMF(51)		MDF(13)		LF (3)		All (197)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Agriculture	0	0	36	57.14	36	60	38	74.51	5	38.46	1	33.33	116	58.88
2	Agricultural Labour	7	100	2	3.17	2	3.33	1	1.96	0	0	0	0	12	6.09
3	Private Service	0	0	0	0	1	1.67	1	1.96	1	7.69	0	0	3	1.52
4	Student	0	0	19	30.16	14	23.33	7	13.73	3	23.08	1	33.33	44	22.34
5	Housewife	0	0	6	9.52	4	6.67	3	5.88	4	30.77	1	33.33	18	9.14
6	Children	0	0	0	0	3	5	1	1.96	0	0	0	0	4	2.03
Total		7	100	63	100	60	100	51	100	13	100	3	100	197	100

**Occupation of the household members:** The data regarding the occupation of the household members in Raisabad Hosalli-1 micro-watershed is presented in Table 6. The results indicate that agriculture was the major occupation for 58.88 per cent of the

household members, 6.09 per cent were agricultural labourers, 1.52 per cent were private service, 22.34 per cent student, 9.14 per cent were housewives and 2.03 per cent were children.

**Institutional participation of the household members:** The data regarding the institutional participation of the household members in Raisabad Hosalli-1 micro-watershed is presented in Table 7. The results show that, 100 per cent of the population in the micro watershed has not participated in any local institutions. .

**Table 7. Institutional Participation of household members in Raisabad Hosalli-1 micro-watershed**

Sl.No.	Particulars	LL (7)		MF (63)		SF (60)		SMF (51)		MDF (13)		LF (3)		All (197)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	No Participation	7	100	63	100	60	100	51	100	13	100	3	100	197	100
	Total	7	100	63	100	60	100	51	100	13	100	3	100	197	100

**Type of house owned:** The data regarding the type of house owned by the households in Raisabad Hosalli-1 micro-watershed is presented in Table 8. The results indicate that 7.5 per cent of the households possess thatched, 67.5 per cent of the households possess katcha house and 25 per cent of the households possess pucca/RCC house.

**Table 8. Type of house owned by households in Raisabad Hosalli-1 micro-watershed**

Sl.No.	Particulars	LL (2)		MF (14)		SF (10)		SMF (10)		MDF (3)		LF (1)		All (40)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Thatched	1	50	2	14.29	0	0	0	0	0	0	0	0	3	7.50
2	Katcha	1	50	8	57.14	10	100	6	60	1	33.33	1	100	27	67.50
3	Pucca/RCC	0	0	4	28.57	0	0	4	40	2	66.67	0	0	10	25
	Total	2	100	14	100	10	100	10	100	3	100	1	100	40	100

**Durable Assets owned by the households:** The data regarding the Durable Assets owned by the households in Raisabad Hosalli-1 micro-watershed is presented in Table 9. The results show that 45 per cent of the households possess TV, 2.50 per cent of the households possess radio and landline phone, 7.50 per cent of the households possess mixer/grinder, 5 per cent of the households possess bicycle, 20 per cent of the households possess motor cycle and 95 per cent of the households possess mobile phones.

**Table 9. Durable Assets owned by households in Raisabad Hosalli-1 micro-watershed**

Sl.No.	Particulars	LL (2)		MF (14)		SF (10)		SMF (10)		MDF (3)		LF (1)		All (40)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Radio	0	0	0	0	0	0	1	10	0	0	0	0	1	2.50
2	Television	0	0	6	42.86	6	60	3	30	2	66.67	1	100	18	45
3	Mixer/Grinder	0	0	0	0	2	20	1	10	0	0	0	0	3	7.50
4	Bicycle	0	0	0	0	0	0	2	20	0	0	0	0	2	5
5	Motor Cycle	0	0	2	14.29	3	30	2	20	1	33.33	0	0	8	20
6	Landline Phone	0	0	1	7.14	0	0	0	0	0	0	0	0	1	2.50
7	Mobile Phone	2	100	13	92.86	9	90	10	100	3	100	1	100	38	95

**Average value of durable assets:** The data regarding the average value of durable assets owned by the households in Raisabad Hosalli-1 micro-watershed is presented in Table 10. The results show that the average value of radio was Rs. 1,000, television was Rs. 9,111, mixer/grinder was Rs. 2,000, bicycle was Rs. 2,000, motor cycle was Rs. 52,500, landline was Rs. 2,000 and mobile phone was Rs. 2,692.

**Table 10. Average value of durable assets owned by households in Raisabad Hosalli-1 micro-watershed** Average value (Rs.)

Sl.No.	Particulars	LL (2)	MF (14)	SF (10)	SMF (10)	MDF (3)	LF (1)	All (40)
1	Radio	0	0	0	1,000	0	0	1,000
2	Television	0	9,000	9,166	9,333	9,000	9,000	9,111
3	Mixer/Grinder	0	0	2,000	2,000	0	0	2,000
4	Bicycle	0	0	0	2,000	0	0	2,000
5	Motor Cycle	0	65,000	41,666	50,000	65,000	0	52,500
6	Landline Phone	0	2,000	0	0	0	0	2,000
7	Mobile Phone	3,000	2,575	2,137	3,538	2,666	2,000	2,692

**Farm Implements owned:** The data regarding the farm implements owned by the households in Raisabad Hosalli-1 micro-watershed is presented in Table 11. About 10 per cent each of the households possess bullock cart, 30 per cent each of the households possess plough, 7.50 per cent of the households possess sprayer and sprinkler and 65 per cent of the households possess weeder.

**Table 11. Farm Implements owned by households in Raisabad Hosalli-1 micro-watershed**

Sl.No.	Particulars	LL (2)		MF (14)		SF (10)		SMF (10)		MDF (3)		LF (1)		All (40)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Bullock Cart	0	0	0	0	2	20	2	20	0	0	0	0	4	10
2	Plough	0	0	2	14.29	4	40	5	50	1	33.33	0	0	12	30
3	Sprayer	0	0	1	7.14	2	20	0	0	0	0	0	0	3	7.50
4	Sprinkler	0	0	2	14.29	0	0	1	10	0	0	0	0	3	7.50
5	Weeder	1	50	10	71.43	6	60	5	50	3	100	1	100	26	65
6	Blank	1	50	2	14.29	0	0	1	10	0	0	0	0	4	10

**Average value of farm implements:** The data regarding the average value of farm Implements owned by the households in Raisabad Hosalli-1 micro-watershed is presented in Table 12. The results show that the average value of bullock cart was Rs. 16,500, plough was Rs. 1,575, sprayer was Rs. 2,166, sprinkler was Rs.2,333 and the average value of weeder was Rs. 57.

**Table 12. Average value of farm implements owned by households in Raisabad Hosalli-1 micro-watershed** Average Value (Rs.)

Sl.No.	Particulars	LL (2)	MF (14)	SF (10)	SMF(10)	MDF (3)	LF (1)	All (40)
1	BullockCart	0	0	17,500	15,500	0	0	16,500
2	Plough	0	1,500	1,750	1,480	1,500	0	1,575
3	Sprayer	0	2,500	2,000	0	0	0	2,166
4	Sprinkler	0	2,500	0	2,000	0	0	2,333
5	Weeder	50	60	56	50	57	66	57

**Livestock possession by the households:** The data regarding the Livestock possession by the households in Raisabad Hosalli-1 micro-watershed is presented in Table 13. The results indicate that, 35 per cent of the households possess bullocks, 22.5 per cent of the households possess local cow, 2.5 per cent of the households possess crossbreed cow, 17.5 per cent of the households possess buffalo and 7.50 per cent of the households possess goat.

**Table 13. Livestock possession by households in Raisabad Hosalli-1 micro-watershed**

Sl.No.	Particulars	LL (2)		MF (14)		SF (10)		SMF (10)		MDF (3)		LF (1)		All (40)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Bullock	1	50	5	35.71	4	40	3	30	0	0	1	100	14	35
2	Local cow	0	0	3	21.43	2	20	2	20	1	33.33	1	100	9	22.50
3	Crossbred cow	0	0	1	7.14	0	0	0	0	0	0	0	0	1	2.50
4	Buffalo	0	0	3	21.43	0	0	4	40	0	0	0	0	7	17.50
5	Goat	0	0	2	14.29	0	0	1	10	0	0	0	0	3	7.50
6	blank	1	50	7	50	6	60	4	40	2	66.67	0	0	20	50

**Average Labour availability:** The data regarding the average labour availability in Raisabad Hosalli-1 micro-watershed is presented in Table 14. The results indicate that, average own labour men available in the micro watershed was 2.18, average own labour (women) available was 1.45, average hired labour (men) available was 10 and average hired labour (women) available was 9.05.

In case of marginal farmers, average own labour men available was 1.93, average own labour (women) was 1.29, average hired labour (men) was 6.79 and average hired labour (women) available was 5.64. In case of small farmers, average own labour men available was 2.30, average own labour (women) was 1.5, average hired labour (men) was 8.5 and average hired labour (women) available was 7.50. In case of semi medium farmers, average own labour men available was 2.7, average own labour (women) was 1.6, average hired labour (men) was 9.5 and average hired labour (women) available was 8.5. In case of medium farmers, average own labour men available and average own labour (women) was 1.67, average hired labour (men) and average hired labour (women) available was 23.33. In case of large farmers, average own labour men available and average own labour (women) was 1, average hired labour (men) and average hired labour (women) available was 35.

**Table 14. Average Labour availability in Raisabad Hosalli-1 micro-watershed**

Sl.No.	Particulars	LL (2)	MF (14)	SF (10)	SMF (10)	MDF (3)	LF (1)	All (40)
1	Hired labour Female	0	5.64	7.50	8.50	23.33	35	9.05
2	Own Labour Female	0	1.29	1.50	1.60	1.67	1	1.45
3	Own labour Male	0	1.93	2.30	2.70	1.67	1	2.18
4	Hired labour Male	0	6.79	8.50	9.50	23.33	35	10

**Adequacy of Hired Labour:** The data regarding the adequacy of hired labour in Raisabad Hosalli-1 micro-watershed is presented in Table 15. The results indicate that, 95 per cent of the households opined that the hired labour was adequate.

**Table 15. Adequacy of Hired Labour in Raisabad Hosalli-1 micro-watershed**

Sl.No.	Particulars	LL (2)		MF (14)		SF (10)		SMF (10)		MDF (3)		LF (1)		All (40)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Adequate	0	0	14	100	10	100	10	100	3	100	1	100	38	95

**Distribution of land (ha):** The data regarding the distribution of land (ha) in Raisabad Hosalli-1 micro-watershed is presented in Table 16. The results indicate that, households of the Raisabad Hosalli-1 micro-watershed possess 23.63 ha (46.73%) of dry land, 25.80 ha (51.03%) of irrigated land and 1.13 ha (2.24%) of permanent fallow land. Marginal farmers possess 5.4 ha (73.64%) of dry land, 1.7 ha (23.06%) of irrigated land and 0.2 ha (3.29%) of permanent fallow land. Small farmers possess 5.16 ha (49.3%) of dry land and 5.29 ha (50.6%) of irrigated land. Semi medium farmers possess 6.16 ha (38.1%) of dry land and 9.11 ha (56.3%) of irrigated land and 0.89 ha (5.51 %) of permanent fallow land. Medium farmers possess 6.88 ha (62.96%) of dry land and 4.05 ha (37.04%) of irrigated land. Large farmers possess 5.67 ha (100%) of irrigated land.

**Table 16. Distribution of land (Ha) in Raisabad Hosalli-1 micro-watershed**

Sl. No.	Particulars	LL (2)		MF (14)		SF (10)		SMF (10)		MDF (3)		LF (1)		All (40)	
		ha	%	ha	%	ha	%	ha	%	ha	%	ha	%	ha	%
1	Dry	0	0	5.4	73.64	5.16	49.3	6.16	38.1	6.88	62.96	0	0	23.63	46.73
2	Irrigated	0	0	1.7	23.06	5.29	50.6	9.11	56.3	4.05	37.04	5.67	100	25.80	51.03
3	Permanent Fallow	0	0	0.2	3.29	0	0	0.89	5.51	0	0	0	0	1.13	2.24
Total		0	100	7.3	100	10.4	100	16.1	100	10.9	100	5.67	100	50.56	100

**Average land value (Rs./ha):** The data regarding the average land value (Rs./ha) in Raisabad Hosalli-1 micro-watershed is presented in Table 17. The results indicate that, the average value of dry land was Rs. 340,587.53, the average value of irrigated land was Rs. 604,328.73 and the average value of permanent fallow land was Rs. 573,392.85.

In case of marginal famers, the average land value was Rs. 690,715.90 for dry land, the average land value was Rs. 1,587,857.12 and the average land value was Rs.1,646,666.60. In case of small famers, the average land value was Rs. 407,142.86 for dry land and the average land value was Rs. 832,159.27 for irrigated land. In case of semi medium famers, the average land value was Rs. 227,051.87 for dry land, the average land value was Rs. 636,711.11 for irrigated land and the average land value was Rs. 280,681.82 for permanent fallow land. In case of medium famers, the average land value was Rs. 116,235.29 for dry land and the average land value was Rs. 469,300 for irrigated

land. In case of large famers, the average land value was Rs. 141,142.86 for irrigated land.

**Table 17. Average land value (Rs./ha) in Raisabad Hosalli-1 micro-watershed**

Sl. No.	Particulars	MF (14)	SF (10)	SMF (10)	MDF (3)	LF (1)	All (40)
1	Dry	690,715.90	407,142.86	227,051.87	116,235.29	0	340,587.53
2	Irrigated	1,587,857.12	832,159.27	636,711.11	469,300	141,142.86	604,328.73
3	Permanent Fallow	1,646,666.60	0	280,681.82	0	0	573,392.85

**Status of bore wells:** The data regarding the status of bore wells in Raisabad Hosalli-1 micro-watershed is presented in Table 18. The results indicate that, there were 20 functioning bore wells in the micro watershed.

**Table 18. Status of bore wells in Raisabad Hosalli-1 micro-watershed**

Sl.No.	Particulars	LL (2)	MF (14)	SF (10)	SMF (10)	MDF (3)	LF (1)	All (40)
1	Functioning	0	4	6	7	2	1	20

**Source of irrigation:** The data regarding the source of irrigation in Raisabad Hosalli-1 micro-watershed is presented in Table 19. The results indicate that, bore well was the major irrigation source in the micro water shed for 50 per cent of the farmers.

**Table 19. Source of irrigation in Raisabad Hosalli-1 micro-watershed**

Sl.No.	Particulars	LL (2)		MF (14)		SF (10)		SMF (10)		MDF (3)		LF (1)		All (40)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Bore Well	0	0	4	28.57	6	60	7	70	2	66.67	1	100	20	50

**Depth of Water (Avg. in meters):** The data regarding the depth of water in Raisabad Hosalli-1 micro-watershed is presented in Table 20. The results indicate that, the depth of bore well was found to be 52.96 meters.

**Table 20. Depth of water (Avg in meters) in Raisabad Hosalli-1 micro-watershed**

Sl.No.	Particulars	LL (2)	MF (14)	SF (10)	SMF (10)	MDF (3)	LF (1)	All (40)
1	Bore Well	0	30.48	62.48	74.68	71.12	106.68	52.96

**Irrigated Area (ha):** The data regarding the irrigated area (ha) in Raisabad Hosalli-1 micro-watershed is presented in Table 21. The results indicate that, marginal, small, semi medium, medium and large farmers had an irrigated area of 1.70 ha, 5.29 ha, 9.19 ha, 4.05 ha and 5.67 ha respectively.

**Table 21. Irrigated Area (ha) in Raisabad Hosalli-1micro-watershed**

Sl.No.	Particulars	LL (2)	MF (14)	SF (10)	SMF (10)	MDF (3)	LF (1)	All (40)
1	Kharif	0	1.70	5.29	9.19	4.05	5.67	25.89
	Total	0	1.70	5.29	9.19	4.05	5.67	25.89

**Cropping pattern:** The data regarding the cropping pattern in Raisabad Hosalli-1 micro-watershed is presented in Table 22. The results indicate that, farmers have grown cotton (20.11%), green gram (15.35 ha), groundnut (3.62 ha) and red gram (4.20 ha). Marginal farmers have grown cotton, green gram and red gram. Small farmers have grown cotton, groundnut and red gram. Semi medium farmers have grown cotton, green gram and groundnut. Medium farmers have grown cotton, green gram and red gram. Large farmers have grown cotton.

**Table 22. Cropping pattern in Raisabad Hosalli-1 micro-watershed** (Area in ha)

Sl.No.	Particulars	LL (2)	MF (14)	SF (10)	SMF (10)	MDF (3)	LF (1)	All (40)
1	Kharif - Cotton	0	2.66	4.02	5.75	4.86	2.83	20.11
2	Kharif - Greengram	0	3.51	0	8.60	3.24	0	15.35
3	Kharif - Groundnut	0	0	2.60	1.01	0	0	3.62
4	Kharif - Red gram	0	0.96	1.21	0	2.02	0	4.20
Total		0	7.13	10.45	15.36	10.12	5.67	48.72

**Cropping intensity:** The data regarding the cropping intensity in Raisabad Hosalli-1 micro-watershed is presented in Table 23. The results indicate that, the cropping intensity in Raisabad Hosalli-1 micro-watershed was found to be 96.78 per cent.

**Table 23. Cropping intensity (%) in Raisabad Hosalli-1 micro-watershed**

Sl.No.	Particulars	LL (2)	MF (14)	SF (10)	SMF (10)	MDF (3)	LF (1)	All (40)
1	Cropping Intensity	0	100	100	94.99	92.59	100	96.78



**Cost of cultivation of Cotton:** The data regarding the cost of cultivation of Cotton in Raisabad Hosalli-1 micro-watershed is presented in Table 24. The results indicate that, the total cost of cultivation for Cotton was Rs. 37968.84. The gross income realized by the farmers was Rs. 124061.50. The net income from Cotton cultivation was Rs. 86092.66. Thus the benefit cost ratio was found to be 1:3.27.

**Table 24. Cost of Cultivation of Cotton in Raisabad Hosalli-1 micro-watershed**

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
<b>I</b>	<b>Cost A1</b>				
1	Hired Human Labour	Man days	50.42	7926.97	20.88
2	Bullock	Pairs/day	4.18	2506.72	6.60
3	Tractor	Hours	2.41	1931.94	5.09
4	Machinery	Hours	0	0	0
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	5.01	4652.52	12.25
6	Seed Inter Crop	Kgs.	0	0	0
7	FYM	Quintal	3.01	602.48	1.59
8	Fertilizer + micronutrients	Quintal	7.33	5896.73	15.53
9	Pesticides (PPC)	Kgs / liters	1.67	1673.76	4.41
10	Irrigation	Number	1.55	0	0
11	Repairs		0	0	0
12	Msc. Charges (Marketing costs etc)		0	0	0
13	Depreciation charges		0	18.39	0.05
14	Land revenue and Taxes		0	3.29	0.01
<b>II</b>	<b>Cost B1</b>				
16	Interest on working capital			1539.18	4.05
17	<b>Cost B1 = (Cost A1 + sum of 15 and 16)</b>			26751.98	70.46
<b>III</b>	<b>Cost B2</b>				
18	Rental Value of Land			380.95	1
19	<b>Cost B2 = (Cost B1 + Rental value)</b>			27132.93	71.46
<b>IV</b>	<b>Cost C1</b>				
20	Family Human Labour		30.75	7383.20	19.45
21	<b>Cost C1 = (Cost B2 + Family Labour)</b>			34516.13	90.91
<b>V</b>	<b>Cost C2</b>				
22	Risk Premium			1	0
23	<b>Cost C2 = (Cost C1 + Risk Premium)</b>			34517.13	90.91
<b>VI</b>	<b>Cost C3</b>				
24	Managerial Cost			3451.71	9.09
25	<b>Cost C3 = (Cost C2 + Managerial Cost)</b>			37968.84	100
<b>VII</b>	<b>Economics of the Crop</b>				
a.	Main Product	a) Main Product (q)	24.74	124061.50	
		b) Main Crop Sales Price (Rs.)		5014.29	
b.	Gross Income (Rs.)			124061.50	
c.	Net Income (Rs.)			86092.66	
d.	Cost per Quintal (Rs./q.)			1534.61	
e.	Benefit Cost Ratio (BC Ratio)			1:3.27	

**Cost of Cultivation of Green gram:** The data regarding the cost of cultivation of green gram in Raisabad Hosalli-1 micro-watershed is presented in Table 25. The results indicate that, the total cost of cultivation for green gram was Rs. 27795.50. The gross income realized by the farmers was Rs. 54874.85. The net income from green gram cultivation was Rs. 27079.35. Thus the benefit cost ratio was found to be 1:1.97.

**Table 25. Cost of Cultivation of green gram in Raisabad Hosalli-1 micro-watershed**

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
<b>I</b>	<b>Cost A1</b>				
1	Hired Human Labour	Man days	49.96	9165.70	32.98
2	Bullock	Pairs/day	1.88	1125.53	4.05
3	Tractor	Hours	1.85	1423.88	5.12
4	Machinery	Hours	0.27	186.20	0.67
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	10.25	1379.03	4.96
7	FYM	Quintal	2.48	861.62	3.10
8	Fertilizer + micronutrients	Quintal	5.24	4717.14	16.97
9	Pesticides (PPC)	Kgs / liters	1.21	1212.61	4.36
10	Irrigation	Number	2.26	0	0
11	Repairs		0	0	0
12	Msc. Charges (Marketing costs etc)		0	0	0
13	Depreciation charges		0	57.58	0.21
14	Land revenue and Taxes		0	3.29	0.01
<b>II</b>	<b>Cost B1</b>				
16	Interest on working capital			980.57	3.53
17	<b>Cost B1 = (Cost A1 + sum of 15 and 16)</b>			21113.16	75.96
<b>III</b>	<b>Cost B2</b>				
18	Rental Value of Land			333.33	1.20
19	<b>Cost B2 = (Cost B1 + Rental value)</b>			21446.49	77.16
<b>IV</b>	<b>Cost C1</b>				
20	Family Human Labour		14.65	3821.15	13.75
21	<b>Cost C1 = (Cost B2 + Family Labour)</b>			25267.64	90.91
<b>V</b>	<b>Cost C2</b>				
22	Risk Premium			1	0
23	<b>Cost C2 = (Cost C1 + Risk Premium)</b>			25268.64	90.91
<b>VI</b>	<b>Cost C3</b>				
24	Managerial Cost			2526.86	9.09
25	<b>Cost C3 = (Cost C2 + Managerial Cost)</b>			27795.50	100
<b>VII</b>	<b>Economics of the Crop</b>				
a.	Main Product	a) Main Product (q)		11.96	54869
		b) Main Crop Sales Price (Rs.)			4588.46
	By Product	e) Main Product (q)		0.08	5.85
		f) Main Crop Sales Price (Rs.)			76.92
b.	Gross Income (Rs.)			54874.85	
c.	Net Income (Rs.)			27079.35	
d.	Cost per Quintal (Rs./q.)			2324.42	
e.	Benefit Cost Ratio (BC Ratio)			1:1.97	

**Cost of cultivation of Red gram:** The data regarding the cost of cultivation of Red gram in Raisabad Hosalli-1 micro-watershed is presented in Table 26. The results indicate that, the total cost of cultivation for Red gram was Rs. 65271.49. The gross income realized by the farmers was Rs. 149815.42. The net income from Red gram cultivation was Rs. 84543.93. Thus the benefit cost ratio was found to be 1:2.3.

**Table 26. Cost of Cultivation of Red gram in Raisabad Hosalli-1 micro-watershed**

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
<b>I</b>	<b>Cost A1</b>				
1	Hired Human Labour	Man days	47.91	7797.09	11.95
2	Bullock	Pairs/day	4.94	2964	4.54
3	Tractor	Hours	4.98	3816.89	5.85
4	Machinery	Hours	0	0	0
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	12.13	1484.79	2.27
7	FYM	Quintal	5.71	1142.81	1.75
8	Fertilizer + micronutrients	Quintal	14.02	11555.29	17.70
9	Pesticides (PPC)	Kgs / liters	3.98	3979.02	6.10
10	Irrigation	Number	4.94	0	0
11	Repairs		0	0	0
12	Msc. Charges (Marketing costs etc)		0	0	0
13	Depreciation charges		0	34.03	0.05
14	Land revenue and Taxes		0	3.29	0.01
<b>II</b>	<b>Cost B1</b>				
16	Interest on working capital			2179.55	3.34
17	<b>Cost B1 = (Cost A1 + sum of 15 and 16)</b>			34956.77	53.56
<b>III</b>	<b>Cost B2</b>				
18	Rental Value of Land			266.67	0.41
19	<b>Cost B2 = (Cost B1 + Rental value)</b>			35223.44	53.96
<b>IV</b>	<b>Cost C1</b>				
20	Family Human Labour		97.56	24113.28	36.94
21	<b>Cost C1 = (Cost B2 + Family Labour)</b>			59336.72	90.91
<b>V</b>	<b>Cost C2</b>				
22	Risk Premium			1	0
23	<b>Cost C2 = (Cost C1 + Risk Premium)</b>			59337.72	90.91
<b>VI</b>	<b>Cost C3</b>				
24	Managerial Cost			5933.77	9.09
25	<b>Cost C3 = (Cost C2 + Managerial Cost)</b>			65271.49	100
<b>VII</b>	<b>Economics of the Crop</b>				
a.	Main Product	a) Main Product (q)		31.37	147444.22
		b) Main Crop Sales Price (Rs.)			4700
	By Product	e) Main Product (q)		5.93	2371.20
		f) Main Crop Sales Price (Rs.)			400
b.	Gross Income (Rs.)			149815.42	
c.	Net Income (Rs.)			84543.93	
d.	Cost per Quintal (Rs./q.)			2080.62	
e.	Benefit Cost Ratio (BC Ratio)			1:2.3	

**Cost of cultivation of Groundnut:** The data regarding the cost of cultivation of groundnut in Raisabad Hosalli-1 micro-watershed is presented in Table 27. The results indicate that, the total cost of cultivation for groundnut was Rs. 31661.34. The gross income realized by the farmers was Rs. 72163.14. The net income from groundnut cultivation was Rs. 40501.81. Thus the benefit cost ratio was found to be 1:2.28.

**Table 27. Cost of Cultivation of groundnut in Raisabad Hosalli-1 micro-watershed**

Sl.No	Particulars	Units	Phy Units	Value(Rs.)	% to C3
<b>I</b>	<b>Cost A1</b>				
1	Hired Human Labour	Man days	52.21	10532.57	33.27
2	Bullock	Pairs/day	0.91	547.20	1.73
3	Tractor	Hours	2.84	2036.47	6.43
4	Machinery	Hours	2.57	1898.09	5.99
5	Seed Main Crop (Establishment and Maintenance)	Kgs (Rs.)	11.89	1783.73	5.63
6	Seed Inter Crop	Kgs.	0	0	0
7	FYM	Quintal	7.10	1420.71	4.49
8	Fertilizer + micronutrients	Quintal	4.39	3888.77	12.28
9	Pesticides (PPC)	Kgs / liters	0.84	841.58	2.66
10	Irrigation	Number	0	0	0
11	Repairs		0	0	0
12	Msc. Charges (Marketing costs etc)		0	0	0
13	Depreciation charges		0	98.82	0.31
14	Land revenue and Taxes		0	3.29	0.01
<b>II</b>	<b>Cost B1</b>				
16	Interest on working capital			952.29	3.01
17	<b>Cost B1 = (Cost A1 + sum of 15 and 16)</b>			24003.51	75.81
<b>III</b>	<b>Cost B2</b>				
18	Rental Value of Land			333.33	1.05
19	<b>Cost B2 = (Cost B1 + Rental value)</b>			24336.84	76.87
<b>IV</b>	<b>Cost C1</b>				
20	Family Human Labour		17.23	4445.19	14.04
21	<b>Cost C1 = (Cost B2 + Family Labour)</b>			28782.03	90.91
<b>V</b>	<b>Cost C2</b>				
22	Risk Premium			1	0
23	<b>Cost C2 = (Cost C1 + Risk Premium)</b>			28783.03	90.91
<b>VI</b>	<b>Cost C3</b>				
24	Managerial Cost			2878.30	9.09
25	<b>Cost C3 = (Cost C2 + Managerial Cost)</b>			31661.34	100
<b>VII</b>	<b>Economics of the Crop</b>				
a.	Main Product	a) Main Product (q)		15.46	72163.14
		b) Main Crop Sales Price (Rs.)			4666.67
b.	Gross Income (Rs.)				72163.14
c.	Net Income (Rs.)				40501.81
d.	Cost per Quintal (Rs./q.)				2047.48
e.	Benefit Cost Ratio (BC Ratio)				1:2.28

**Adequacy of fodder:** The data regarding the adequacy of fodder in Raisabad Hosalli-1 micro-watershed is presented in Table 28. The results indicate that, 47.50 per cent of the households opined that dry fodder and green fodder was adequate.

**Table 28. Adequacy of fodder in Raisabad Hosalli-1 micro-watershed**

Sl. No.	Particulars	LL (2)		MF (14)		SF (10)		SMF (10)		MDF (3)		LF (1)		All (40)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Adequate-Dry Fodder	0	0	7	50	4	40	6	60	1	33.33	1	100	19	47.50
2	Adequate-Green Fodder	0	0	7	50	4	40	6	60	1	33.33	1	100	19	47.50

**Annual gross income:** The data regarding the annual gross income in Raisabad Hosalli-1 micro-watershed is presented in Table 29. The results indicate that the annual gross income was Rs. 50,000 for landless farmers, for marginal farmers it was Rs. 124,764.29, for small farmers it was Rs. 139,500, semi medium farmers it was Rs. 192,400, for medium farmers it was Rs. 376,500 and large farmers it was Rs. 636,000.

**Table 29. Annual gross income in Raisabad Hosalli-1 micro-watershed**

(Avg value in Rs.)

Sl.No.	Particulars	LL (2)	MF (14)	SF (10)	SMF(10)	MDF (3)	LF (1)	All (40)
1	Service/salary	0	0	0	20,000	33,333.33	0	7,500
2	Wage	50,000	44,285.71	34,000	51,500	20,000	20,000	41,375
3	Agriculture	0	46,785.71	99,500	118,940	319,166.67	616,000	110,322.50
4	Dairy Farm	0	3,121.43	6,000	1,960	4,000	0	3,382.50
5	Goat Farming	0	30,571.43	0	0	0	0	10,700
	Income(Rs.)	50,000	124,764.29	139,500	192,400	376,500	636,000	173,280

**Average annual expenditure:** The data regarding the average annual expenditure in Raisabad Hosalli-1 micro-watershed is presented in Table 30. The results indicate that the average annual expenditure is Rs. 37,494.69. For landless farmers it was Rs. 37,500, marginal farmers it was Rs. 16,490.78, for small farmers it was Rs. 11,750, for semi medium farmers it was Rs. 25,241.67, for medium farmers it was Rs. 120,666.67 and for large farmers it was Rs. 462,000.

**Table 30. Average annual expenditure in Raisabad Hosalli-1 micro-watershed**

(Avg value in Rs.)

Sl.No.	Particulars	LL (2)	MF (14)	SF (10)	SMF (10)	MDF (3)	LF (1)	All (40)
1	Service/salary	0	0	0	140,000	75,000	0	5,375
2	Wage	75,000	27,928.57	21,500	36,666.67	16,000	12,000	26,375
3	Agriculture	0	33,442.31	66,000	70,500	265,000	450,000	76,118.75
4	Dairy Farm	0	13,500	30,000	5,250	6,000	0	1,837.50
5	Goat Farming	0	156,000	0	0	0	0	7,800
	Total	75,000	230,870.8	117,500	252,416.6	362,000	462,000	1,499,787.5
	Average	37,500	16,490.78	11,750	25,241.67	120,666.6	462,000	37,494.69

**Horticulture species grown:** The data regarding horticulture species grown in Raisabad Hosalli-1 micro-watershed is presented in Table 31. The results indicate that, households have planted 31 mango and 1 sapota trees in their field.

**Table 31: Horticulture species grown in Raisabad Hosalli-1 micro-watershed**

Sl.No.	Particulars	LL (2)		MF (14)		SF (10)		SMF (10)		MDF (3)		LF (1)		All (40)	
		F	B	F	B	F	B	F	B	F	B	F	B	F	B
1	Mango	0	0	12	0	6	0	11	0	1	0	1	0	31	0
2	Sapota	0	0	1	0	0	0	0	0	0	0	0	0	1	0

**Forest species grown:** The data regarding forest species grown in Raisabad Hosalli-1 micro-watershed is presented in Table 32. The results indicate that, households have planted 83 neem, 6 tamarind and 1 banyan trees in their field and also 22 neem trees in their backyard.

**Table 32: Forest species grown in Raisabad Hosalli-1 micro-watershed**

Sl.No.	Particulars	LL (2)		MF (14)		SF (10)		SMF (10)		MDF (3)		LF (1)		All (40)	
		F	B	F	B	F	B	F	B	F	B	F	B	F	B
1	Neem	0	0	26	0	20	16	26	6	6	0	5	0	83	22
2	Tamarind	0	0	3	0	0	0	1	0	0	0	2	0	6	0
3	Banyan	0	0	1	0	0	0	0	0	0	0	0	0	1	0

\*F= Field B=Back Yard

**Average Additional investment capacity:** The data regarding average additional investment capacity in Raisabad Hosalli-1 micro-watershed is presented in Table 33. The results indicated that, households have an average investment capacity of Rs. 200 for land development.

**Table 33: Average additional investment capacity in Raisabad Hosalli-1 micro-watershed**

Sl.No.	Particulars	LL (2)	MF (14)	SF (10)	SMF (10)	MDF (3)	LF (1)	All (40)
1	Land development	0	71.43	200	500	0	0	200

**Source of additional investment:** The data regarding source of funds for additional investment in Raisabad Hosalli-1 micro-watershed is presented in Table 34. The results indicated that loan from bank was the source of additional investment for 2.44 per cent for land development. Own funds was the source of additional investment for 7.32 per cent for land development.

**Table 34: Source of funds for additional investment capacity in Raisabad Hosalli-1 micro –watershed**

Sl.No	Item	Land development	
		N	%
1	Loan from bank	1	2.44
2	Own funds	3	7.32

**Marketing of the agricultural produce:** The data regarding marketing of the agricultural produce in Raisabad Hosalli-1 micro-watershed is presented in Table 35. The results indicated that, cotton was sold to the extent of 99.54 per cent, green gram and red gram was sold to the extent of 93.75 per cent and groundnut was sold to the extent of 98.25 per cent.

**Table 35. Marketing of the agricultural produce in Raisabad Hosalli-1 micro-watershed**

Sl.No	Crops	Output obtained (q)	Output retained (q)	Output sold (q)	Output sold (%)	Avg. Price obtained (Rs/q)
1	Cotton	439.0	2.0	437.0	99.54	5014.29
2	Greengram	224.0	14.0	210.0	93.75	4588.46
3	Groundnut	57.0	1.0	56.0	98.25	4666.67
4	Redgram	80.0	5.0	75.0	93.75	4700.0

**Marketing Channels used for sale of agricultural produce:** The data regarding marketing channels used for sale of agricultural produce in Raisabad Hosalli-1 micro-watershed is presented in Table 36. The results indicated that, about 90 per cent of the farmers sold their produce to local/village merchant and 7.50 per cent of the farmers sold their produce to regulated market.

**Table 36. Marketing Channels used for sale of agricultural produce in Raisabad Hosalli-1 micro-watershed**

Sl. No.	Particulars	MF (14)		SF (10)		SMF (10)		MDF (3)		LF (1)		All (40)	
		N	%	N	%	N	%	N	%	N	%	N	%
1	Local/village Merchant	13	92.86	8	80	9	90	4	133.33	2	200	36	90
2	Regulated Market	0	0	2	20	1	10	0	0	0	0	3	7.50

**Mode of transport of agricultural produce:** The data regarding mode of transport of agricultural produce in Raisabad Hosalli-1 micro-watershed is presented in Table 37. The results indicated that, 97.5 per cent of the households have used tractor as a mode of transportation.

**Table 37. Mode of transport of agricultural produce in Raisabad Hosalli-1 micro-watershed**

Sl.No.	Particulars	LL (2)		MF (14)		SF (10)		SMF (10)		MDF (3)		LF (1)		All (40)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Tractor	0	0	13	92.86	10	100	10	100	4	133.33	2	200	39	97.50

**Incidence of soil and water erosion problems:** The data regarding incidence of soil and water erosion problems in Raisabad Hosalli-1 micro-watershed is presented in Table 38. The results indicated that, 50 per cent of the households have experienced soil and water erosion problems in the farm.

**Table 38. Incidence of soil and water erosion problems in Raisabad Hosalli-1 micro-watershed**

Sl.No.	Particulars	LL(2)		MF(14)		SF(10)		SMF(10)		MDF(3)		LF(1)		All(40)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Soil and water erosion problems in the farm	0	0	9	64.29	5	50	4	40	1	33.33	1	100	20	50

**Interest shown towards soil testing:** The data regarding Interest shown towards soil testing in Raisabad Hosalli-1 micro-watershed is presented in Table 39. The results indicated that, 80 per cent have shown interest in soil test.

**Table 39. Interest shown towards soil testing in Raisabad Hosalli-1 micro-watershed**

Sl.No.	Particulars	LL (2)		MF (14)		SF (10)		SMF (10)		MDF (3)		LF (1)		All (40)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Interest in soil test	0	0	12	85.71	7	70	9	90	3	100	1	100	32	80

**Usage pattern of fuel for domestic use:** The data regarding usage pattern of fuel for domestic use in Raisabad Hosalli-1 micro-watershed is presented in Table 40. The results indicated that, 90 per cent of the households used fire wood and 10 per cent of the households used LPG as a source of fuel.

**Table 40. Usage pattern of fuel for domestic use in Raisabad Hosalli-1 micro-watershed**

Sl.No.	Particulars	LL (2)		MF (14)		SF (10)		SMF (10)		MDF (3)		LF (1)		All (40)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Fire Wood	2	100	11	78.57	10	100	9	90	3	100	1	100	36	90
2	LPG	0	0	3	21.43	0	0	1	10	0	0	0	0	4	10

**Source of drinking water:** The data regarding source of drinking water in Raisabad Hosalli-1 micro-watershed is presented in Table 41. The results indicated that, piped supply was the major source of drinking water for 92.5 per cent and 7.50 per cent of the households used bore well in the micro watershed.

**Table 41. Source of drinking water in Raisabad Hosalli-1 micro-watershed**

Sl.No.	Particulars	LL (2)		MF (14)		SF (10)		SMF (10)		MDF (3)		LF (1)		All (40)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Piped supply	2	100	13	92.86	8	80	10	100	3	100	1	100	37	92.50
2	Bore Well	0	0	1	7.14	2	20	0	0	0	0	0	0	3	7.50

**Source of light:** The data regarding source of light in Raisabad Hosalli-1 micro-watershed is presented in Table 42. The results indicated that, Electricity was the major source of light for 100 per cent of the households in micro watershed.

**Table 42. Source of light in Raisabad Hosalli-1 micro-watershed**

Sl.No.	Particulars	LL (2)		MF (14)		SF (10)		SMF (10)		MDF (3)		LF (1)		All (40)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Electricity	2	100	14	100	10	100	10	100	3	100	1	100	40	100

**Table 43. Existence of Sanitary toilet facility in Raisabad Hosalli-1 micro-watershed**

Sl.No.	Particulars	LL (2)		MF (14)		SF (10)		SMF (10)		MDF (3)		LF (1)		All (40)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Sanitary toilet facility	1	50	7	50	7	70	5	50	2	66.67	1	100	23	57.50

**Existence of Sanitary toilet facility:** The data regarding existence of sanitary toilet facility in Raisabad Hosalli-1 micro-watershed is presented in Table 43. The results indicated that, 57.5 per cent of the households possess sanitary toilet facility.



**Possession of PDS card:** The data regarding possession of PDS card in Raisabad Hosalli-1 micro-watershed is presented in Table 44. The results indicated that, 100 per cent of the sampled households possessed BPL cards.

**Table 44. Possession of PDS card in Raisabad Hosalli-1 micro-watershed**

Sl.No.	Particulars	LL (2)		MF (14)		SF (10)		SMF (10)		MDF (3)		LF (1)		All (40)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	BPL	2	100	14	100	10	100	10	100	3	100	1	100	40	100

**Participation in NREGA program:** The data regarding participation in NREGA programme in Raisabad Hosalli-1 micro-watershed is presented in Table 45. The results indicated that, 87.5 per cent of the households participated in NREGA programme.

**Table 45. Participation in NREGA programme in Raisabad Hosalli-1 micro-watershed**

Sl.No.	Particulars	LL(2)		MF (14)		SF(10)		SMF(10)		MDF(3)		LF(1)		All (40)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Participation in NREGA programme	2	100	11	78.57	10	100	8	80	3	100	1	100	35	87.50

**Adequacy of food items:** The data regarding adequacy of food items in Raisabad Hosalli-1 micro-watershed is presented in Table 46. The results indicated that, cereals milk and egg were adequate for 100 per cent of the households, pulses were adequate for 80 per cent, oilseed were adequate for 5 per cent, vegetables were adequate for 85 per cent, fruits were adequate for 22.50 per cent and meat were adequate for 90 per cent.

**Table 46. Adequacy of food items in Raisabad Hosalli-1 micro-watershed**

Sl.No.	Particulars	LL (2)		MF (14)		SF (10)		SMF (10)		MDF (3)		LF (1)		All (40)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Cereals	2	100	14	100	10	100	10	100	3	100	1	100	40	100
2	Pulses	2	100	10	71.43	7	70	10	100	2	66.67	1	100	32	80
3	Oilseed	1	50	0	0	0	0	1	10	0	0	0	0	2	5
4	Vegetables	1	50	14	100	7	70	8	80	3	100	1	100	34	85
5	Fruits	0	0	3	21.43	3	30	3	30	0	0	0	0	9	22.50
6	Milk	2	100	14	100	10	100	10	100	3	100	1	100	40	100
7	Egg	2	100	14	100	10	100	10	100	3	100	1	100	40	100
8	Meat	2	100	13	92.86	8	80	9	90	3	100	1	100	36	90

**Table 47. Response on Inadequacy of food items in Raisabad Hosalli-1 micro-watershed**

Sl.No.	Particulars	LL (2)		MF (14)		SF (10)		SMF (10)		MDF (3)		LF (1)		All (40)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Pulses	0	0	4	28.57	3	30	0	0	1	33.33	0	0	8	20
2	Oilseed	1	50	14	100	10	100	9	90	3	100	1	100	38	95
3	Vegetables	1	50	0	0	3	30	2	20	0	0	0	0	6	15
4	Fruits	2	100	11	78.57	7	70	7	70	3	100	1	100	31	77.50
5	Meat	0	0	1	7.14	2	20	1	10	0	0	0	0	4	10

**Response on Inadequacy of food items:** The data regarding inadequacy of food items in Raisabad Hosalli-1 micro-watershed is presented in Table 47. The results indicated that, pulse were inadequate for 20 per cent of the households, oilseed were inadequate for 95 per cent, vegetables were inadequate for 15 per cent, fruits were inadequate for 77.50 per cent and meat were inadequate for 10 per cent of the households.

**Farming constraints:** The data regarding farming constraints experienced by households in Raisabad Hosalli-1 micro-watershed is presented in Table 48. The results indicated that, lower fertility status of the soil in the area was the constraint experienced by 82.5 per cent of the households, wild animal menace on farm field (22.5%), frequent incidence of pest and diseases (87.5%), Inadequacy of irrigation water (12.5%), high cost of fertilizer and plant protection chemicals (85%), high rate of interest on credit (27.5%), low price for the agricultural commodities (65%), lack of marketing facilities in the area (25%), inadequate extension service (2.5%), Lack of transport for safe transport of the Agril produce to the market (12.5%), less rainfall and Source of Agri-technology information (10%).

**Table 48. Farming constraints Experienced in Raisabad Hosalli-1 micro-watershed**

Sl. No.	Particulars	LL(2)		MF (14)		SF(10)		SMF(10)		MDF(3)		LF(1)		All (40)	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%
1	Lower fertility status of the soil	0	0	13	92.86	7	70	9	90	3	100	1	100	33	82.50
2	Wild animal menace on farm field	0	0	4	28.57	4	40	1	10	0	0	0	0	9	22.50
3	Frequent incidence of pest and diseases	0	0	13	92.86	9	90	9	90	3	100	1	100	35	87.50
4	Inadequacy of irrigation water	0	0	4	28.57	0	0	1	10	0	0	0	0	5	12.50
5	High cost of Fertilizers and plant protection chemicals	0	0	12	85.71	9	90	9	90	3	100	1	100	34	85
6	High rate of interest on credit	0	0	3	21.43	4	40	4	40	0	0	0	0	11	27.50
7	Low price for the agricultural commodities	0	0	11	78.57	5	50	6	60	3	100	1	100	26	65
8	Lack of marketing facilities in the area	0	0	4	28.57	4	40	2	20	0	0	0	0	10	25
9	Inadequate extension services	0	0	1	7.14	0	0	0	0	0	0	0	0	1	2.50
10	Lack of transport for safe transport of the Agril produce to the market.	0	0	1	7.14	3	30	1	10	0	0	0	0	5	12.50
11	Less rainfall	0	0	0	0	3	30	1	10	0	0	0	0	4	10
12	Source of Agri-technology information	0	0	0	0	3	30	1	10	0	0	0	0	4	10

**SUMMARY**

In order to assess the socio-economic condition of the farmers in the watershed a comprehensive questionnaire was prepared. Major components such as demographic conditions, migration details, food consumption and family expenditure pattern, material possession, land holding, land use management, cropping pattern, cost of cultivation of crops, livestock management. The statistical components such as frequency and percentage were used to analyze the data. About 40 households located in the micro watershed were interviewed for the survey.

The data indicated that there were 125 (63.45%) men and 72 (36.55%) women among the sampled households. The average family size of landless farmers' was 3.5, marginal farmers' was 4.5, small farmers' was 6, semi medium farmers' was 5.1, medium farmers was 4.3 and large farmers was 3. The data indicated that, 37 (18.78%) people were in 0-15 years of age, 75 (38.07%) were in 16-35 years of age, 69 (35.03%) were in 36-60 years of age and 16 (8.12%) were above 61 years of age.

The results indicated that Raisabad Hosalli-1 had 59.90 per cent illiterates, 1.02 per cent of them functional literate, 15.74 per cent of them had primary school, 3.55 per cent of them had Middle school and PUC education, 11.17 per cent of them had high school and 3.05 per cent of them had degree education.

The results indicate that, 85 per cent of household heads were practicing agriculture, 10 per cent of the household heads were agricultural labourers and 5 cent of the household heads was housewives. The results indicate that agriculture was the major occupation for 58.88 per cent of the household members, 6.09 per cent were agricultural labourers, 1.52 per cent were private service, 22.34 per cent student, 9.14 per cent were housewives and 2.03 per cent were children.

The results show that, 100 per cent of the population in the micro watershed has not participated in any local institutions. The results indicate that 7.5 per cent of the households possess thatched, 67.5 per cent of the households possess katcha house and 25 per cent of the households possess pucca/RCC house.

The results show that 45 per cent of the households possess TV, 2.50 per cent of the households possess radio and landline phone, 7.50 per cent of the households possess mixer/grinder, 5 per cent of the households possess bicycle, 20 per cent of the households possess motor cycle and 95 per cent of the households possess mobile phones. The results show that the average value of television was Rs. 6,393, mixer/grinder was Rs. 1,642, motor cycle was Rs. 56,333, auto was Rs. 160,000 and mobile phone was Rs. 2,474.

About 10 per cent each of the households possess bullock cart, 30 per cent each of the households possess plough, 7.50 per cent of the households possess sprayer and

sprinkler and 65 per cent of the households possess weeder. The results show that the average value of bullock cart was Rs. 16,500, plough was Rs. 1,575, sprayer was Rs. 2,166, sprinkler was Rs.2,333 and the average value of weeder was Rs. 57.

The results indicate that, 35 per cent of the households possess bullocks, 22.5 per cent of the households possess local cow, 2.5 per cent of the households possess crossbreed cow, 17.5 per cent of the households possess buffalo and 7.50 per cent of the households possess goat.

The results indicate that, average own labour men available in the micro watershed was 2.18, average own labour (women) available was 1.45, average hired labour (men) available was 10 and average hired labour (women) available was 9.05. The results indicate that, 95 per cent of the households opined that the hired labour was adequate.

The results indicate that, households of the Raisabad Hosalli-1 micro-watershed possess 23.63 ha (46.73%) of dry land, 25.80 ha (51.03%) of irrigated land and 1.13 ha (2.24%) of permanent fallow land. Marginal farmers possess 5.4 ha (73.64%) of dry land, 1.7 ha (23.06%) of irrigated land and 0.2 ha (3.29%) of permanent fallow land. Small farmers possess 5.16 ha (49.3%) of dry land and 5.29 ha (50.6%) of irrigated land. Semi medium farmers possess 6.16 ha (38.1%) of dry land and 9.11 ha (56.3%) of irrigated land and 0.89 ha (5.51 %) of permanent fallow land. Medium farmers possess 6.88 ha (62.96%) of dry land and 4.05 ha (37.04%) of irrigated land. Large farmers possess 5.67 ha (100%) of irrigated land.

The results indicate that, the average value of dry land was Rs. 340,587.53, the average value of irrigated land was Rs. 604,328.73 and the average value of permanent fallow land was Rs. 573,392.85. In case of marginal famers, the average land value was Rs. 690,715.90 for dry land, the average land value was Rs. 1,587,857.12 and the average land value was Rs.1,646,666.60. In case of small famers, the average land value was Rs. 407,142.86 for dry land and the average land value was Rs. 832,159.27 for irrigated land. In case of semi medium famers, the average land value was Rs. 227,051.87 for dry land, the average land value was Rs. 636,711.11 for irrigated land and the average land value was Rs. 280,681.82 for permanent fallow land. In case of medium famers, the average land value was Rs. 116,235.29 for dry land and the average land value was Rs. 469,300 for irrigated land. In case of large famers, the average land value was Rs. 141,142.86 for irrigated land.

The results indicate that, there were 20 functioning bore wells in the micro watershed. The results indicate that, bore well was the major irrigation source in the micro water shed for 50 per cent of the farmers. The results indicate that, the depth of bore well was found to be 52.96 meters.

The results indicate that, marginal, small, semi medium, medium and large farmers had an irrigated area of 1.70 ha, 5.29 ha, 9.19 ha, 4.05 ha and 5.67 ha respectively. The results indicate that, farmers have grown cotton (20.11%), green gram (15.35 ha), groundnut (3.62 ha) and red gram (4.20 ha). Marginal farmers have grown cotton, green gram and red gram. Small farmers have grown cotton, groundnut and red gram. Semi medium farmers have grown cotton, green gram and groundnut. Medium farmers have grown cotton, green gram and red gram. Large farmers have grown cotton. The results indicate that, the cropping intensity in Raisabad Hosalli-1 micro-watershed was found to be 96.78 per cent.

The results indicate that, the total cost of cultivation for Cotton was Rs. 37968.84. The gross income realized by the farmers was Rs. 124061.50. The net income from Cotton cultivation was Rs. 86092.66. Thus the benefit cost ratio was found to be 1:3.27. The total cost of cultivation for green gram was Rs. 27795.50. The gross income realized by the farmers was Rs. 54874.85. The net income from green gram cultivation was Rs. 27079.35. Thus the benefit cost ratio was found to be 1:1.97. The total cost of cultivation for Red gram was Rs. 65271.49. The gross income realized by the farmers was Rs. 149815.42. The net income from Red gram cultivation was Rs. 84543.93. Thus the benefit cost ratio was found to be 1:2.3. The total cost of cultivation for groundnut was Rs. 31661.34. The gross income realized by the farmers was Rs. 72163.14. The net income from groundnut cultivation was Rs. 40501.81. Thus the benefit cost ratio was found to be 1:2.28.

The results indicate that, 47.50 per cent of the households opined that dry fodder and green fodder was adequate.

The results indicate that the annual gross income was Rs. 50,000 for landless farmers, for marginal farmers it was Rs. 124,764.29, for small farmers it was Rs. 139,500, semi medium farmers it was Rs. 192,400, for medium farmers it was Rs. 376,500 and large farmers it was Rs. 636,000. The results indicate that the average annual expenditure is Rs. 37,494.69. For landless farmers it was Rs. 37,500, marginal farmers it was Rs. 16,490.78, for small farmers it was Rs. 11,750, for semi medium farmers it was Rs. 25,241.67, for medium farmers it was Rs. 120,666.67 and for large farmers it was Rs. 462,000.

The results indicate that, households have planted 31 mango and 1 sapota trees in their field. The results indicate that, households have planted 83 neem, 6 tamarind and 1 banyan trees in their field and also 22 neem trees in their backyard.

The results indicated that, households have an average investment capacity of Rs. 200 for land development. The results indicated that loan from bank was the source of additional investment for 2.44 per cent for land development. Own funds was the source of additional investment for 7.32 per cent for land development.

The results indicated that, cotton was sold to the extent of 99.54 per cent, green gram and red gram was sold to the extent of 93.75 per cent and groundnut was sold to the extent of 98.25 per cent.

The results indicated that, about 90 per cent of the farmers sold their produce to local/village merchant and 7.50 per cent of the farmers sold their produce to regulated market. The results indicated that, 97.5 per cent of the households have used tractor as a mode of transportation.

The results indicated that, 50 per cent of the households have experienced soil and water erosion problems in the farm. The results indicated that, 80 per cent have shown interest in soil test.

The results indicated that, 90 per cent of the households used fire wood and 10 per cent of the households used LPG as a source of fuel. The results indicated that, piped supply was the major source of drinking water for 92.5 per cent and 7.50 per cent of the households used bore well in the micro watershed.

Electricity was the major source of light for 100 per cent of the households in micro watershed. The results indicated that, 57.5 per cent of the households possess sanitary toilet facility. The results indicated that, 100 per cent of the sampled households possessed BPL cards. The results indicated that, 87.5 per cent of the households participated in NREGA programme.

The results indicated that, cereals milk and egg were adequate for 100 per cent of the households, pulses were adequate for 80 per cent, oilseed were adequate for 5 per cent, vegetables were adequate for 85 per cent, fruits were adequate for 22.50 per cent and meat were adequate for 90 per cent. The results indicated that, pulse were inadequate for 20 per cent of the households, oilseed were inadequate for 95 per cent, vegetables were inadequate for 15 per cent, fruits were inadequate for 77.50 per cent and meat were inadequate for 10 per cent of the households.

The results indicated that, lower fertility status of the soil in the area was the constraint experienced by 82.5 per cent of the households, wild animal menace on farm field (22.5%), frequent incidence of pest and diseases (87.5%), Inadequacy of irrigation water (12.5%), high cost of fertilizer and plant protection chemicals (85%), high rate of interest on credit (27.5%), low price for the agricultural commodities (65%), lack of marketing facilities in the area (25%), inadequate extension service (2.5%), Lack of transport for safe transport of the Agril produce to the market (12.5%), less rainfall and Source of Agri-technology information (10%).